

**W L Willowood**

# TEBUCON 3.6SC

For control of specified diseases on various crops.

**ACTIVE INGREDIENT:**

Tebuconazole, alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol ..... 38.7%

**OTHER INGREDIENTS:** ..... 61.3%

**TOTAL:** ..... 100.0%

Contains 3.6 pounds tebuconazole per gallon.

**STOP - Read the label before use  
KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See inside label booklet for First Aid, Precautionary Statements and Directions for Use including Storage and Disposal instructions.

EPA Reg. No. 87290-13

EPA Est. No. 70989-AR-001

**Manufactured For:**  
Willowood, LLC  
1600 NW Garden Valley Blvd. #120  
Roseburg, OR 97471

**NET CONTENTS: 2.5 Gallons**

### FIRST AID

<b>If swallowed:</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time or your poison control center at 1-800-222-1222.

**NOTE TO PHYSICIAN:** No specific antidote. Treat symptomatically. The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing vapor or spray mist.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

##### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY REQUIREMENTS

##### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

**Ground Water Advisory:** Tebuconazole is known to leach through soil into ground under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground-water contamination.

**Surface Water Advisory:** Willowood Tebucon 3.6SC may contaminate water through drift of spray in wind. Willowood Tebucon 3.6SC has a high potential for runoff for several months or more after application. Poorly draining soils and soils with shallow water tables are more prone to runoff that contains this product. A level, well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall-runoff. Runoff of Willowood Tebucon 3.6SC will be reduced by avoiding applications when rainfall is forecasted within 48 hours.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI). The REI for each crop is listed in the application directions associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride or viton
- Shoes plus socks

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store above 28°F or agitate before use.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### PRODUCT INFORMATION

Read the entire Directions for Use and Conditions of Sale before using this product.

**Spray Volume:** Willowood Tebucon 3.6SC may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment. Check equipment calibration frequently. Complete coverage and uniform application are essential for the most effective results, especially when lower spray volumes are applied. If necessary, increase the spray volume per acre for complete crop coverage.

**Chemigation:** Apply Willowood Tebucon 3.6SC through irrigation equipment only to crops and diseases for which the chemigation use is specified. Apply Willowood Tebucon 3.6SC only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move; or drip (trickle) irrigation systems. Do not apply Willowood Tebucon 3.6SC through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either

automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. Do not apply when wind speed favors drift beyond the area intended for treatment.

Maintain continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. Pesticide may be applied continuously for the duration of the water application.

**Mixing:** Add labeled amount of Willowood Tebucon 3.6SC into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the Willowood Tebucon 3.6SC should be thoroughly dispersed prior to the addition of other materials. Do not tank mix with products containing a prohibition against tank mixing. Follow the most restrictive labeling requirements of any tank mix product.

**Compatibility:** To determine the compatibility of Willowood Tebucon 3.6SC with other products, the following procedure should be followed: Pour the recommended proportions of the products into a suitable container of water, mix thoroughly and allow to stand at least five (5) minutes. If the combination remains mixed or can be re-mixed readily, the mixture is considered physically compatible. For further information, contact your local Willowood USA, LLC representative.

**Resistance Management Statement**

Willowood Tebucon 3.6SC is a Group 3 fungicide which exhibits no known cross-resistance to other fungicide groups. However, fungal pathogens are known to develop resistance to products with the same mode of action when used repeatedly. Any fungal population may contain or develop individuals that are resistant to Willowood Tebucon 3.6SC and other Group 3 fungicides. If Group 3 fungicides are used repeatedly in the same field or in successive years as the primary method of control for targeted diseases, the resistant isolates may eventually dominate the fungal population. Because resistance development cannot be predicted, the use of Willowood Tebucon 3.6SC should conform to resistance management strategies established for the crop and use area. Such strategies may include rotation and/or tank mixing with products having different modes of action or limiting the total number of applications per season. Contact your local extension specialist, certified crop advisor, and/or manufacturer for fungicide resistance management and/or integrated disease management recommendations for specific crops and resistant disease populations. Willowood USA, LLC encourages responsible resistance management to ensure effective long-term control of the fungal diseases on this label.

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Asparagus	Rust ( <i>Puccinia</i> spp.)	4 to 6 fl. oz. per acre
	<b>Notes:</b> Apply Willowood Tebucon 3.6SC as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of Willowood Tebucon 3.6SC per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days of harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).	
<b>General Comments:</b> Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating Willowood Tebucon 3.6SC with other DMI fungicides may lead to resistance.		
<b>Restricted-entry interval (REI)</b> = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Barley	Rust ( <i>Puccinia</i> spp.)	4 fl. oz. per acre
	Head blight ( <i>Fusarium</i> spp.) – Suppression	
<p><b>Notes:</b> Apply Willowood Tebucon 3.6SC in a minimum of 10 gallons of spray solution per acre by ground or in a minimum of 5 gallons of spray solution per acre by air. A maximum of 4 fl. oz. of Willowood Tebucon 3.6SC may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Grazing livestock or feeding of green forage is permitted 6 or more days after the last application of Willowood Tebucon 3.6SC. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.</p> <p><b>Application timing directions:</b></p> <p><b>Rusts:</b> Apply Willowood Tebucon 3.6SC at the earliest sign of rust pustules on foliage.</p> <p><b>Fusarium head blight:</b> Optimal timing of Willowood Tebucon 3.6SC for Fusarium head blight suppression is when main stem heads have fully emerged (Feekes 10.5) on 50% of the plants.</p>		
<p><b>General Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Beans (fresh & dry except succulent shelled)	Rust ( <i>Uromyces appendiculatus</i> )	4 to 6 fl. oz. per acre
	<p><b>Notes:</b> Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: Willowood Tebucon 3.6SC may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6SC per acre per crop season. Beans, dry: Willowood Tebucon 3.6SC may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of Willowood Tebucon 3.6SC per acre per crop season.</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust ( <i>Puccinia</i> spp.) Northern leaf blight ( <i>Helminthosporium turcicum</i> ) Southern leaf blight ( <i>Helminthosporium maydis</i> ) Northern leaf spot ( <i>Helminthosporium carbonum</i> ) Gray leaf spot ( <i>Cercospora zea-maydis</i> )	4 to 6 fl. oz. per acre
	<p><b>Notes:</b> Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of Willowood Tebucon 3.6SC may be applied per acre per crop season. Sweet corn: Willowood Tebucon 3.6SC may be applied up to 7 days before the harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: Willowood Tebucon 3.6SC may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restricted-entry interval (REI) for sweet corn = 19 hours.</b></p> <p><b>Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.</b></p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Cotton	Southwestern cotton rust ( <i>Puccinia cacabata</i> )	6 to 8 fl. oz. per acre
	<p><b>Notes:</b> Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. Willowood Tebucon 3.6SC may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6SC per acre per crop season.</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (includes hyotan, cucuzza, hechima and Chinese okra) <i>Momordica</i> spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternut squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew ( <i>Sphaerotheca fuliginea</i> / <i>Podosphaera xanthii</i> ) ( <i>Erysiphe cichoracearum</i> )  Gummy stem blight - suppression ( <i>Didymella bryoniae</i> ) (watermelon, squash, pumpkin, and melons only)	4 to 6 fl. oz. per acre  8 fl. oz. per acre
<p><b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. Willowood Tebucon 3.6SC may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6SC per acre per crop season.</p>		
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		
<p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		



APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot ( <i>Sclerotium cepivorum</i> )	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust ( <i>Puccinia allii</i> , <i>Puccinia porri</i> ) Purple blotch ( <i>Alternaria porri</i> )	4 to 6 fl. oz. per acre
<p><b>White rot:</b> For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz Willowood Tebucon 3.6SC per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p><b>Rust:</b> For the control of rust make foliar applications at the rate of 4 to 6 fl. oz Willowood Tebucon 3.6SC per acre per application. Repeat at an interval of 10 to 14 days. Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p><b>Notes:</b> Do not apply more than 32.5 fl. oz. Willowood Tebucon 3.6SC per acre per season if an in-furrow treatment is made. If Willowood Tebucon 3.6SC is not applied as an in-furrow treatment then do not apply more than 12 fl oz. Willowood Tebucon 3.6SC per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).</p>		
<p><b>General Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Fruiting Vegetables Group Eggplant Groundcherry Pepino Pepper Tomatillo Tomato	Early blight ( <i>Alternaria solani</i> )	8 fl. oz. per acre
	<p><b>Notes:</b> Apply Willowood Tebucon 3.6SC as a foliar spray using an interval of 7 days. Do not apply more than 48 fl. oz. of Willowood Tebucon 3.6SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).</p>	

**General Comments:** For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted-entry interval (REI) = 12 hours.**

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Grasses Grown For Seed	Rusts ( <i>Puccinia</i> spp.)	4 to 8 fl. oz. per acre
	Apply the specified rate of Willowood Tebucon 3.6SC as soon as weather conditions are favorable for rust development or when first rust pustules are present. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.	
	Powdery mildew	4 to 8 fl. oz. per acre
Apply specified rate of Willowood Tebucon 3.6SC when powdery mildew first appears on the leaves. Repeat applications at 14- to 16-day intervals. Under heavy disease pressure use 6 to 8 fl oz/A and shorter spray intervals.		

**General Comments:** Apply the specified rate in a minimum of 20 gallons of water per acre with ground sprayers or in a minimum of 10 gallons of water per acre with aircraft. Thorough coverage is important for optimum disease control.

For optimum benefit, the lowest specified rate of a spray surfactant should be tank mixed with Willowood Tebucon 3.6SC.

A maximum of 16 fluid ounces (1 pint) may be applied per acre per crop season. Willowood Tebucon 3.6SC may be applied up to 4 days before harvest. Chaff, screenings and straw from treated areas may be used for feed purposes; however, do not forage, cut green crop, or use seed for feed purposes. Regrowth may be grazed starting 17 days after last application.

**Restricted-entry interval (REI) = 12 hours.**

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Green onion Leek Spring onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot ( <i>Sclerotium cepivorum</i> ) suppression only Rust ( <i>Puccinia allii</i> , <i>Puccinia porri</i> ) Purple blotch ( <i>Alternaria porri</i> )	4 to 6 fl. oz per acre
For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply Willowood Tebucon 3.6SC in a protective spray schedule or when weather conditions are favorable for rust development.		
<b>Notes:</b> Do not apply more than 24 fl. oz. Willowood Tebucon 3.6SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).		

**General Comments:** For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted-entry interval (REI) = 12 hours.**

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Hops	Powdery mildew ( <i>Sphaerotheca humuli</i> / <i>Sphaerotheca macularis</i> )	4 to 8 fl. oz. per acre
	<b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. Willowood Tebucon 3.6SC may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of Willowood Tebucon 3.6SC per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.	

**General Comments:** For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restricted-entry interval (REI) = 12 hours.**

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuna Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot ( <i>Cercospora brassicicola</i> )	3 to 4 fl. oz. per acre
	Powdery mildew ( <i>Erysiphe cruciferarum</i> ) Alternaria leaf spot ( <i>Alternaria brassicicola</i> )	
<b>Notes:</b> Do not apply more than 16 fl. oz. Willowood Tebucon 3.6SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days). Do not apply more often than once every 10 days.		

**General Comments:** For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).

**Restriction:** Application to turnip greens is limited to East of the Rockies.

**Restricted-entry interval (REI) = 12 hours.**

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Garden beet roots and tops (leaves)	Cercospora leaf spot ( <i>Cercospora beticola</i> )	3 to 7.2 fl. oz. per acre
	<b>Notes:</b> Make applications on a 14 day interval. Do not apply more than 28.8 fl. oz. Willowood Tebucon 3.6SC per acre per season. Do not apply within 7 days of harvest (PHI = 7 days).	
<b>General Comments:</b> For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest recommended rate of a spray surfactant may be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
<b>Restricted-entry interval (REI)</b> = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Lychee	Anthrachnose ( <i>Colletotrichum gloeosporioides</i> )	4 to 6 fl. oz. per acre
	<b>Notes:</b> Begin first application of Willowood Tebucon 3.6SC as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specified dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of Willowood Tebucon 3.6SC per acre per season. Willowood Tebucon 3.6SC can be applied up to and including the day of harvest (PHI = 0 days).	
<b>General Comments:</b> For optimum disease control, the lowest labeled rate of a non-ionic spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
<b>Restricted-entry interval (REI)</b> = 2 days.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Okra	Cercospora leaf spot ( <i>Cercospora</i> spp.)	4 to 6 fl. oz. per acre
	<b>Notes:</b> Apply specific dosage of Willowood Tebucon 3.6SC in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of Willowood Tebucon 3.6SC per acre per season.	
<b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
<b>Restricted-entry interval (REI)</b> = 12 hours.		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Peanut	<b>SOILBORNE:</b> Sclerotium stem and pod rot (white mold, southern blight, southern stem rot) Rhizoctonia limb rot Rhizoctonia pod rot (Virginia and North Carolina only)  <b>FOLIAR:</b> Early leaf spot Late Leaf spot Leaf rust Web blotch ( <i>Phoma</i> ) Pepper spot ( <i>Leptoshaerulina</i> )	7.2 fl. oz. per acre
	<b>FOUR-APPLICATION SPRAY PROGRAM:</b> Apply the specified rate in a preventive spray schedule. See table below for proper timing of applications. Applications of chlorothalonil should be made prior to and following applications of Willowood Tebucon 3.6SC to discourage development of resistant strains of fungi. For optimum control of foliar diseases such as leaf rust, web blotch, and pepper spot, the lowest label specified rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC.  <b>LEAF SPOT ADVISORY SCHEDULE:</b> For control of soilborne diseases in an advisory schedule, apply Willowood Tebucon 3.6SC in the first advisory spray in July and continue Willowood Tebucon 3.6SC applications at 14-day intervals. Applications after August 15 should be tank mixed with chlorothalonil for resistance management purposes.	
<p><b>GENERAL DIRECTIONS:</b> For optimum control of the specified soilborne diseases, four consecutive applications of Willowood Tebucon 3.6SC must be made at 14-day intervals.</p> <p>A maximum of 28.8 fluid ounces of Willowood Tebucon 3.6SC may be applied per crop season. Willowood Tebucon 3.6SC may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas.</p> <p>Willowood Tebucon 3.6SC is a sterol demethylation inhibitor (DMI) fungicide. Chlorothalonil may be tank mixed at the rate of 12 ounces of active ingredient with Willowood Tebucon 3.6SC as a leaf spot resistance management strategy. A spray surfactant is not necessary when Willowood Tebucon 3.6SC is tank mixed with chlorothalonil. Mixing or alternating Willowood Tebucon 3.6SC with other DMI fungicides may lead to resistance.</p> <p>Willowood Tebucon 3.6SC must be carried by rainfall or irrigation into the root and pod zone for control of root and pod rots caused by <i>Sclerotium rolfsii</i> and <i>Rhizoctonia solani</i>. Drought conditions will decrease the effectiveness of Willowood Tebucon 3.6SC against the root and pod rots.</p> <p>Use Willowood Tebucon 3.6SC in conjunction with cultural practices that are known to reduce the severity of soilborne diseases, such as proper crop rotation practices.</p> <p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		
<b>Timing of Willowood Tebucon 3.6SC Application for Optimum Control of White Mold and Rhizoctonia Limb and Pod Rot</b>		
<b>Spray Program</b>	<b>Willowood Tebucon 3.6SC Application No.</b>	<b>Chlorothalonil Application No.</b>
7 Applications	3,4,5 and 6	1,2 and 7

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Pecan	Brown leaf spot ( <i>Sirosporium diffusum</i> ) Downy spot ( <i>Mycosphaerella caryigena</i> ) Liver spot ( <i>Gnomonia caryae</i> ) Scab ( <i>Cladosporium caryigenum</i> ) Vein spot ( <i>Gnomonia nerviseda</i> ) Zonate leaf spot ( <i>Grovesinia pyramidalis</i> )	4 to 8 fl. oz. per acre
<p><b>Notes:</b> Apply Willowood Tebucon 3.6SC in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. Willowood Tebucon 3.6SC should be applied at 4 fl. oz. per acre in a tank-mix with the recommended rate of Super-Tin® in cover sprays. Follow label directions for the use of SuperTin. Do not add a surfactant to the spray solution when tank-mixing Willowood Tebucon 3.6SC with SuperTin. Apply Willowood Tebucon 3.6SC in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of Willowood Tebucon 3.6SC to full-size mature trees, and 4 to 6 fl. oz. per acre of Willowood Tebucon 3.6SC to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of Willowood Tebucon 3.6SC may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.</p>		
<p><b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p>		
<p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		

APPLICATION DIRECTIONS		
CROP	DISEASE(S)	RATE OF WILLOWOOD TEBUCON 3.6SC
Soybean	Rust ( <i>Phakopsora pachyrhizi</i> ) Powdery Mildew ( <i>Microsphaera diffusa</i> )	3 to 4 fl. oz. per acre

**Use Directions:** Apply Willowood Tebucon 3.6SC as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use the higher rates and shorter spray intervals when disease pressure is severe. The lowest labeled rate of spray surfactant must be tank-mixed with Willowood Tebucon 3.6SC. Apply Willowood Tebucon 3.6SC in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.

**Restrictions:** Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply more than 12 fl. oz/a per use season.

**Restricted-entry interval (REI) = 12 hours.**

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Sunflower	Rust ( <i>Puccinia helianthi</i> )	4 to 6 fl. oz. per acre
	<b>Notes:</b> Apply specific dosage of Willowood Tebucon 3.6SC at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Application may be repeated at 14 days if necessary to maintain control of the disease. Apply specified dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of Willowood Tebucon 3.6SC per acre per season or within 50 days of harvest.	
<b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Contact your state Extension Service or Willowood USA, LLC representative for a list of approved surfactants. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
<b>Restricted-entry interval (REI) = 12 hours.</b>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot ( <i>Cercospora brassicicola</i> )	4 to 7.2 fl. oz. per acre
	<b>Notes:</b> Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-day intervals. Willowood Tebucon 3.6SC may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of Willowood Tebucon 3.6SC per acre per crop season.	
<b>General Comments:</b> For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).		
<b>Restricted-entry interval (REI) = 12 hours.</b>		

APPLICATION DIRECTIONS		
CROP	DISEASE	RATE OF WILLOWOOD TEBUCON 3.6SC

Wheat	Rusts leaf, stem, and stripe ( <i>Puccinia</i> spp.) Head blight or scab ( <i>Fusarium</i> spp.) - Suppression	4 fl. oz. per acre
	<p><b>Notes:</b> Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. A maximum of 4 fl. oz. of Willowood Tebucon 3.6SC may be applied per acre per crop season. Do not apply within 30 days of harvest. Straw may be fed or used for bedding. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with Willowood Tebucon 3.6SC. Apply Willowood Tebucon 3.6SC in a minimum of 10 gallons of spray solution per acre by ground, or in a minimum of 5 gallons of spray solution per acre by air.</p> <p><b>Application timing directions:</b> Rusts: Apply Willowood Tebucon 3.6SC at the earliest sign of rust pustules on foliage. Fusarium head blight: Optimal timing of Willowood Tebucon 3.6SC for Fusarium head blight suppression is the beginning of flowering on main stem heads (Feekes 10.51).</p>	
<p><b>General Comments:</b> For optimum disease control, the lowest specified rate of a spray surfactant should be tank-mixed with Willowood Tebucon 3.6SC. Willowood Tebucon 3.6SC must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, Willowood Tebucon 3.6SC will be resistant to weathering. Willowood Tebucon 3.6SC is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p><b>Restricted-entry interval (REI) = 12 hours.</b></p>		

<b>SEED TREATMENT - Corn</b> (Sweet Corn, Field Corn, Field Corn Grown For Seed, and Popcorn)		
For control of soilborne and seedborne Fusarium and soilborne and seedborne head smut.		
<b>SEED LABELING:</b> To meet U.S. Federal Seed Act requirements, all seed treated with Willowood Tebucon 3.6SC must be labeled:		
<b>TREATED SEED. DO NOT USE FOR FOOD, FEED OR OIL PURPOSES. Treated with Tebuconazole.</b>		
<b>USE PRECAUTION:</b> When using formulations that do not contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.		
DISEASE	RATE Fl Oz/CWT	DIRECTIONS FOR USE
<b>Soilborne and Seedborne</b> Fusarium	0.071	Apply as a seed treatment using standard slurry or mist-type seed treatment equipment. Uniform application of seed is necessary to ensure seed safety and best disease protection. Seed should be sound and well cured prior to treatment. Product should be diluted with sufficient water to ensure complete seed coverage. Consult a seed treatment specialist regarding slurry rates recommended for the crop to be treated with Willowood Tebucon 3.6SC. The length of control will vary depending on the rate used.
<b>Soilborne and Seedborne</b> Head smut ( <i>Sphacelotheca reilana</i> )	0.27 – 0.54	

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.**

Apply only during alternate years in fields adjacent to aquatic areas listed above.

Do not apply by ground or air within 100 feet of aquatic areas listed above.

Do not cultivate within 10 feet of an aquatic area to allow growth of a vegetative filter strip.

**Spray Drift Management:** For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used, and must not exceed 75% of the wing span or rotor diameter.



Use the largest droplet size consistent with pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure. Apply in a minimum of 5 gallons of spray solution per acre by aircraft spray equipment.

Spray should be released at the lowest possible height consistent with good pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

#### **ROTATIONAL CROPS**

Treated areas may be replanted with any crop specified on this label as soon as practical after last application. Any crop not specified on this label may be planted into treated areas 120 days after last application.

#### **CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Willowood USA, LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Willowood USA, LLC and Seller harmless for any claims relating to such factors.

Willowood USA, LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Willowood USA, LLC, and Buyer and User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WILLOWOOD USA, LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent consistent with applicable law, neither Willowood USA, LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF WILLOWOOD USA, LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF WILLOWOOD USA, LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Willowood USA, LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Willowood USA, LLC.

EPA 20110304