

7 Literaturverzeichnis

- Agee, C.S. and Hill, N.S. (1994): Ergovaline variability in *Acremonium*-infected tall fescue due to environment and plant genotype. *Crop Sci.* 34, 221-226.
- Amalric, C., Sallanon, H., Monnet, F., Hitmi, A., Coudret, A. (1999): Gas exchange and chlorophyll fluorescence in symbiotic and non-symbiotic ryegrass under water stress. *Photosynthetica* 37, 107-112.
- Analytis, S., Kranz, J. und Stumpf, A. (1971): Eine Methode zur Berechnung der Blattfläche. *Angew. Botanik XLV*, 111-114.
- Anonym (1988): Richtlinien für die Durchführung von landwirtschaftlichen Wertprüfungen und Sortenversuchen. Bundessortenamt. Verlag Alfred Strothe. Pp. 173.
- Anonym (2000): Beschreibende Sortenlist 2000. Rasengräser. Bundessortenamt. Verlag Alfred Strothe. Pp. 173.
- Arachevaleta, M., Bacon, C.W., Hoveland, C.S. and Radcliffe D.E. (1989): Effect of the tall fescue endophyte on plant response to environmental stress. *Agron.J.* 81, 83-90.
- Arachevaleta, M., Bacon, C.W., Plattner, R.D., Hoveland, C.S. and Radcliffe D.E. (1992): Accumulation of ergopeptide alkaloids in symbiotic tall fescue grown under deficits of soil water and nitrogen fertiliser. *Appl. Environ. Microbiol.* 58, 857-861.
- Azevedo M. D. and Welty R. E. (1995): A study on the fungal endophyte *Acremonium coenophialum* in the roots of tall fescue seedlings. *Mycologia*, 87 (3), 289-297.
- Bacon, C.W., Porter, J.K., Robbins, J.D. and Luttrell, E.S. (1977): *Epichloe typhina* from tall fescue grasses. *Appl. Environ. Microbiol.* 35, 576-581.
- Bacon, C.W. and Siegel, M.R. (1988): Endophyte parasitism of tall fescue. *J. Prod. Agric.* 1, 45-55.
- Baldwin, J. C., Karthikeyan, A.S. and Raghothama, K. G. (2001): LEPS2, a phosphorus starvation-induced novel acid phosphatase from tomato. *Plant Physiology* 125, 728-737.
- Balfourier, F. and Charmet, G. (1991): Spaced plant evaluation of Mediterranean germplast collection of perennial ryegrass. *Euphytica* 57, 57-66.
- Ball, O.J.P., Prestidge, R.A. and Sprosen, J.M. (1995): Interrelationships between *Acremonium lolii*, peramine, and Lolitrem B in perennial ryegrass. *Appl. Environ. Microbiol.* 61, 1527-1533.
- Barker, D.J., Hume, D.E. and Quigley, P.E. (1997): Negligible physiological responses to water deficit in endophyte-infected and uninfected perennial ryegrass. In: *Neotyphodium/Grass Interactions*. C.W. Bacon and N.S. Hill (Ed.). Plenum Press. New York and London, 137-140.
- Belesky, D.P., Evans J.J. and Wilkinson, S.R. (1985): Amino acid composition of tall fescue seed produced from fungal endophyte (*Acremonium coenophialum*)-free and infected plants. *Agron. J.* 77, 796-798.
- Belesky, D.P., Devine, O.J., Pallas, J.E.jr. and Stringer, W.C. (1987): Photosynthetic activity of tall fescue as influenced by a fungal endophyte. *Photosynthetica* 21, 82.

- Belesky, D.P., Stringer, W.C. and Hill, N.S. (1989): Influence of endophyte and water regime upon tall fescue accessions. I. Growth characteristics. Ann. Bot. 63, 495-503.
- Belesky, D.P. and Fedders, J.M. (1995): Tall fescue development in response to *Acremonium coenophialum* and soil acidity. Crop Sci. 35, 529-533.
- Belesky, D.P. and Fedders, J.M. (1996): Does endophyte influence regrowth of tall fescue? Ann. Bot.. 78 (1996), 499-505.
- Bouton, J.H., Gates, R.N., Belesky, D.P. and Owsley, M. (1993): Yield and persistence of tall fescue in the south-eastern coastal plain after removal of its endophyte. Agron. J. 85, 52-55.
- Buck, G.W., West, C.P. and Elbersen, H.W. (1997): Endophyte effect on drought tolerance in diverse *Festuca* species. In: *Neotyphodium/Grass Interactions*. C.W. Bacon and N.S. Hill (Ed.). Plenum Press. New York and London, pp. 141-144.
- Bumerl, J. and Mika, V. (1991): Infection of meadow fescue and *Festuca arundinacea* by the fungus *Acremonium coenophialum*: The effect on seed yield and quality. Scientific studies 1991, Oseva, Breeding Institute for Fodder Plants, Troubsko, 187-191.
- Bunyard, B. and McInnis, T., Jr. (1990): Evidence for elevated phytohormone levels in endophyte infected tall fescue. Int. Symp. on *Acremonium/Grass Interactions*, Abstracts. S.S. Quinsenberry and R.E. Joost (Ed.). New Orleans, 185-188.
- Burhan, W. (1984): Development of *Acremonium coenophialum* and accumulation of N-acetyl loline and N-formyl loline in tall fescue (*Festuca arundinacea* Schreb.). MT. University of Kentucky. Lexington. US. Pp. 64.
- Bush, L. P., Fannin, F. F., Siegel, M. R., Dahlman, D.L. and Burton, H. R. (1993): Chemistry, occurrence and biological effects of saturated pyrrolizidine alkaloids associated with endophyte grass interactions. Agriculture Ecosystems & Environment 44, 81-102.
- Cabral, D., Novas, M.V. and White, J.F. (2000): *Neotyphodium* endophytes: incidence, diversity, and hosts in South America. In: 4th International *Neotyphodium/Grass Interactions* Symposium. Book of Abstract. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp 34.
- Cagaš, B. (1991): Genetic resources of tall fescue (*Festuca arundinacea* Schreb.) and *Acremonium coenophialum*. Report of a working group on forages, 4th Meeting, Budapest. Pp. 76-81.
- Cagaš, B. (1999): The endophytes (*Neotyphodium* spp.) in ecotypes of *Lolium perenne* and the characteristics of their localities. Grassland Ecology V. Proceedings of the 5th Ecological Conference. 23 - 25 November 1999, Banska Bystrica, Slovakia, 180-184.
- Cagaš, B. (2000): The endophytes *Neotyphodium* spp., their incidence in the Czech republic, problems, questions, work with E+ plants. COST 828 "Endophyte Meeting". Institut für Acker- und Pflanzenbau Halle, Germany, 6./7. April 2000.
- Camacho, R. and Malavolta, E. (2000): Effect of phosphorus supply on growth and acid phosphatase activity of grain sorghum hybrids. Phyton – International Journal of Experimental Botany 67, 13-20.

Carrow, R.N. (1996): Drought avoidance characteristics of diverse tall fescue cultivars. *Crop Sci.* 36, 371-377.

Charmet, G., Balfourier, F. and Bion, A. (1990): Agronomic evaluation of a collection of French perennial ryegrass populations. *Genetic Resources and Crop Evolution* 40, 77-89.

Cheplick, G.P., Clay, K. and Marks, S. (1989): Interactions between infection by endophytic fungi and nutrient limitation in the grasses *Lolium perenne* and *Festuca arundinacea*. *New Phytol.* 111, 89-97.

Christensen, M. J., Ball, O. J. P., Bennett, R. J. and Schardl, C. L. (1997): Fungal and host genotype effects on compatibility and vascular colonization by *Epichloe festucae*. *Mycological Research* 101, 493-501.

Christensen, M.J., Easton, H.S., Simpson, W.R. and Tapper, B.A. (1998): Occurrence of the fungal endophyte *Neotyphodium coenophialum* in leaf blades of tall fescue and implications for stock health. *New Zealand Journal of Agricultural Research* 41, 595-602.

Clay, K. (1987): Effects of fungal endophytes on the seed and seedling biology of *Lolium perenne* and *Festuca arundinacea*. *Oecologia* 73, 358-362.

Clement, S. L., Wilson, A. D., Lester, D. G. and Davitt, C. M. (1997): Fungal endophytes of wild barley and their effects on *Diuraphis noxia* population development. *Entomologia Experimentalis et Aplicata* 82, 275-281.

Cook, K. (1987): Distribution of *Acremonium coenophialum* in developing seedlings and inflorescences of *Festuca arundinacea*. M.S. Thesis, Oregon State Univ., Corvallis. 62 pp.

Cook R., Lewis C. G. and Mizen K. A. (1991): Effects on plant-parasitic nematodes of infection of perennial ryegrass, *Lolium perenne*, by the endophytic fungus, *Acremonium lolii*. *Crop Protection* 91, 403-407.

Daniels, L.B. (1989): Historical perspective of fescue toxicosis in Arkansas. In: Proceedings of the 1989 Arkansas Fescue Toxicosis Conference. C.P. West (Ed.). US. Arkansas Agricultural Experiment Station. Pp. 1-6.

Dapprich, P. (1996): Untersuchungen zur morphologischen und molekularbiologischen Charakterisierung der Gräserendophyten *Acremonium lolii* und *Acremonium uncinatum* und deren Auswirkungen auf ihren Wirt. Dissertation. Paderborn, Germany. Pp 150.

Dapprich, P., Paul, V.H. and Krohn, K. (1996): Incidence of *Acremonium* endophytes in selected German pastures and the contents of alkaloids in *Lolium perenne*. *Bulletin OILB/srop* 19, 103-114.

Da Silva, E.A.A., Pinho E.V.D., Vieira M.D.G.C., de Carvalho M.L.M. and Machado I.D. (2000): Alteration of the izoenzyme patterns in corn seeds infected by fungi. *Pesquisa Agropecuaria Brasileira* 35, 1725-1732.

De Batista, J.P., Bouton, J.H., Bacon, C.W. and Siegel, M.R. (1990a): Rhizome and herbage production of endophyte-removed tall fescue clones and populations. *Agron. J.* 82, 651-654.

De Battista, J.P., Bacon, C.W., Severson, R., Plattner, R.D. and Bouton, J.H. (1990b): Indole acetic acid production by the fungal endophyte of tall fescue. *Agron. J.* 82, 878-880.

Dörffling, K. (1982): Das Hormonsystem der Pflanze. Georg Thieme Verlag Stuttgart New York. Pp. 303.

Dymock, J.J., Rowan, D.D. and McGee, I.R. (1989): Effects of endophyte-produced mycotoxins on Argentine stem weevil and the cutworm *Graphania mutans*. In: Proc. 5th Australasian Conference of Grassland Invertebrate Ecology. Melbourne. Australia. Pp. 35-43.

Easton, H.S. (1993): Will endophyte strain affect variety performance. In: Proc. Sec. Int. Symp. on *Acremonium*/Grass Interactions. Hume D.E., Latch, G. C. M. and Easton, H. S. (Ed.). Palmerston North. New Zealand. Pp. 195-199.

Eerens, J.P.J., Easton, H.S., Lukas, R.J., White, J.G.H. and Miller, K.B. (1997): Influence of the ryegrass endophyte on pasture production and composition in a cool-moist environment. *Neotyphodium*/Grass Interactions. C.W. Bacon and N.S. Hill (Ed.). Plenum Press. New York and London, 157-160.

Eerens, J.P.J., Lucas, R.J., Easton, H.S., White, J.G.H. (1998): Influence of the ryegrass endophyte (*Neotyphodium lolii*) in a cool-moist environment II. New Zealand Journal of Agricultural Research 41, 191-199.

Eggestein, S. (1997): Untersuchungen zum Vorkommen von Endophyten der Gattung *Acremonium* in *Festuca*-Arten Europas sowie zu Symbioseeffekten bei *Festuca pratensis*. Diss., Halle, 115S.

Elbersen, H.W., Buck, G.W., West, C.P. and Joost, R.E. (1994): Water loss from tall fescue leaves is decreased by endophyte. Arkansas Farm Research 43, 8-9.

Elbersen, H.W. and West, C.P. (1996): Growth and water relations of field-grown tall fescue as influenced by drought and endophyte. Grass and Forage Science 51, 333-342.

Elmi, A.A., West, C.P. and Turner, K.E. (1989): *Acremonium* enhances osmotic adjustment in tall fescue. Arkansas Farm Research 38, 7.

Elmi, A.A., West, C.P., Turner, K.E. and Oosterhuis, D.M. (1990): *Acremonium coenophialum* effects on tall fescue water relations. Proc. Int. Symp. on *Acremonium*/Grass Interactions. S.S. Quinsenberry and R.E. Joost (Ed.). New Orleans, 137-140.

Elmi, A.A. and West, C.P. (1995): Endophyte infection effects on stomatal conductance, osmotic adjustment and drought recovery of tall fescue. New Phytol. 131, 61-67.

Emile, J.C., Ravault, J.P., Chabosseau, J.M. and Ghesquiere, M. (1997): Effect of endophyte-infected tall fescue hay on dairy heifers under European intensive management. XVIII International Grassland Congress, Canada.

Endo, T. (1981): Developmental modification and hybridisation of allelic acid phosphatase isoenzymes in homo- and heterozygotes for the Acp-1 locus in rice. Biochemical Genetics 19, 373-384.

Fannin, F.F., Bush, L.P., Siegel, M.R. and Rowan, D.D. (1990): Analysis of peramin in fungal endophyte infected grasses by reversed-phase thin-layer chromatography. J. Chromatogr. 503, 288-292.

Feuerstein, U. (2000): Utilisation of endophyte-mediated advantages in plant breeding. COST 828 "Seed Science in the field of genetically controlled stress physiology" Endophyte Meeting. April 6./7., Halle. Germany.

Fink-Gremmels, J. and Bull, S. (2000): Prevalence of ryegrass staggers syndrome in horses in the Netherlands. In: 4th International *Neotyphodium*/Grass Interactions Symposium. Book of Abstracts. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp 72.

Fletcher, L.R. and Harvey, I.C. (1981): An association of a *Lolium* endophyte with ryegrass staggers. N.Z. Vet. J. 29, 185-186.

Freeman, E.M. (1903): The seed-fungus of *Lolium temulentum* L. the darnel. Philos. Trans. R. Soc. London Ser. B 196, 1-27.

Gallagher, R.T., Hawkes, A.D., Steyn, P.S. and Vleggaar, R. (1984): Termogenic neurotoxins from perennial ryegrass causing ryegrass staggers disorder of livestock: structure elucidation of lolitrem. Br. J. Chem. Soc., Chem. Commun. Pp, 614-616.

Gallagher, R.T., Smith, G.S. and Sprosen, J.M. (1987): The distribution and accumulation of Lolitrem B neurotoxin in developing perennial ryegrass plants. Proc. 4th Animal Science Congress of the Asian-Australasian Assoc. of Animal Production Societies. Hamilton. New Zealand. Pp 404.

Gams, W., Petrini O. and Schmidt D. (1990): *Acremonium uncinatum*, a new endophyte in *Festuca pratensis*. Mycotaxon 15, 311-318.

Garner, G.B., Rottinghaus, G.E., Cornell, C.M. and Testereci, H. (1993): Chemistry of compounds associated with endophyte/grass interaction: ergovaline and ergopeptine-related alkaloids. Agriculture, Ecosystems and Environment 44, 65-80.

Gaume A., Machler F., De Leon C., Narro L., and Frossard, E (2001): Low-P tolerance by maize (*Zea mays* L.) genotypes: Significance of root growth, and organic acids and acid phosphatase root exudation. Plant and Soil 228, 253-264.

Glenn, A.E., Bacon, C.W., Price, R.P. and Hanlin, R.T. (1996): Molecular phylogeny of *Acremonium* and its Taxonomic implications. Mycologia 88, 369-383.

Goodin, J.R. (1972): Chemical regulation of growth in leaves and tillers. In: The Biology and Utilisation of Grasses. V.B. Younger and C.M. McKell (Ed.). Academic Press. New York and London. Pp. 135-146.

Gottlieb, L.D. (1982): Conservation and duplication of isozymes in plants. Science 216, 373-380.

Guillaumin, J.J., Frain, M., Pichon, N. and Ravel, C. (2000): Survey of the fungal endophytes in wild grass species of the Auvergne Region (Central France). In: 4th International *Neotyphodium*/Grass Interactions Symposium. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp 85-92.

Guy, P.L. (1992): Incidence of *Acremonium lolii* and lack of correlation with barley yellow dwarf viruses in tasmanian perennial ryegrass pastures. Plant Pathology 41, 29-34.

Gwinn, K.D. and Bernard, E.C. (1993): Interactions of endophyte-infected grasses with the nematodes *Meloidogyne marylandi* and *Pratylenchus scribneri*. In: Proceedings of the Second International Symposium on *Acremonium*/Grass Interactions. D.E. Hume, G.C.M. Latch and H.S. Easton (Ed.) Palmerston North. New Zealand. Pp. 156-160.

Hahn, H. (1996): Untersuchungen zur elektrophoretischen Charakterisierung ausgewählter Sorten und Zuchtstämme von *Festuca pratensis* Huds. und *Festulolium*-Bastarden. Dissertation. Halle/Saale. Germany. S. 139.

Hesse U. and Latch G.C.M. (1999): Influence of *Neotyphodium lolii* and barley yellow dwarf virus, individually and combined, on the growth of *Lolium perenne*. Australasian Plant Pathology 28, 240-247.

Hiatt E.E. and Hill N.S. (1997): *Neotyphodium coenophialum* mycelial protein and herbage mass effects on ergot alkaloid concentration in tall fescue. Journal of Chemical Ecology 23, 2721-2736.

Hill, N.S., Stringer, W.C., Rottinghaus, G.E., Belesky, D.P., Parrott, W.A. and Pope, D.D. (1990): Growth, morphological, and chemical component responses of tall fescue to *Acremonium coenophialum*. Crop Sci. 30, 156-161.

Hill, N.S., Pachon, J.G. and Bacon, C.W. (1996): *Acremonium coenophialum*-mediated short- and long-term drought acclimation in tall fescue. Crop Sci. 36, 665-672.

Hill, N.S., Beleski, D.P. and Stringer W.C. (1998): Encroachment of endophyte-infected on endophyte-free tall fescue. Annals of Botany 81, 483-488.

Holder, T.L., West, C.P., Turner, K.E., McConnell, M.E. and Piper, E.L. (1994): Incidence and viability of *Acremonium* endophytes in tall fescue and meadow fescue plant introductions. Crop Sci. Madison, Wis. : Crop Science of America, 34, 252-254.

Hoveland, C.S. (1993): Importance and economic significance of the *Acremonium*-endophytes to performance of animals and grass plants. Agriculture Ecosystems & Environment 44, 3-12.

Huizing, H.J., Molen W., Kloek W. and Nijs A. P. M (1991): Detection of lolines in endophyte-containing meadow fescue in the Netherlands and the effect of elevated temperature on induction of lolines in endophyte-infected perennial ryegrass. Blackwell Scientific Publications 46, 441-445.

Hume, D.E., Popay, A.J. and Barker, D.J. (1993): Effect of *Acremonium* endophyte on growth of ryegrass and tall fescue under varying levels of soil moisture and Argentine stem weevil attack. Proc. Sec. Int. Symp.on *Acremonium*/Grass Interactions. Hume D.E., Latch, G. C. M. and Easton, H. S. (Ed.). New Zealand: Palmerston North pp.161.

Joost, R.E. (1995): *Acremonium* in fescue and ryegrass: Boon or bane? A review. J. Anim. Sci. 73, 881-888.

Justus, M., Witte, L. and Hartmann, T. (1997): Levels and tissue distribution of loline alkaloids in endophyte-infected *Festuca pratensis*. Phytochemistry 44, 51-57.

Kennedy, C.W. and Bush, L.P. (1983): Effect of environmental and management factors on the accumulation of N-acetyl and N-formyl loline alkaloids in tall fescue. Crop Science 23, 547-552.

Keogh, R.G. and Lawrence, T. (1987): Influence of *Acremonium lolii* presence on emergence and growth of ryegrass seedlings. NZ. J. Agric. Res. 30, 507-510.

- Keogh, R.G., Tapper, B.A. and Fletcher, R.H. (1991): Distributions of the fungal endophyte *Acremonium lolii*, and of the alkaloids Lolitrem B and peramine, within perennial ryegrass. NZ. J. Agric. Research. 35. 35-38.
- Kimmons, C.A., Gwinn, K.D. and Bernard E.C. (1990): Nematode reproduction on endophyte-free tall fescue. Plant Dis. 74, 757-761.
- Kirpatrick T.L., Barham J.D., and Batteman R.J. (1990): Host status for *Meloidogyne graminis* of tall fescue clones with and without the endophyte *Acremonium coenophialum*. In: Proceedings of the International Symposium on *Acremonium*/Grass Interactions, S. S. Quisenberry and R. E. Joost (Ed.) USA: New Orleans, 154-156.
- Koga, H., Christensen, M.J. and Bennett, R.J. (1993a): Incompatibility of some grass-*Acremonium* endophyte associations. Mycol. Res. 97, 1237-1244.
- Koga, H., Christensen, M.J. and Bennett, R.J. (1993b): Grass/*Acremonium* associations at the cellular level. Proc. Sec. Int. Symp.on *Acremonium*/Grass Interactions. Hume D.E., Latch, G. C. M. and Easton, H. S. (Ed.). New Zealand: Palmerston North, 52.
- Lane, G.A., Tapper,B.A., Davies, E., Hume, D.E., Latch, G.C.M., Barker, D.J., Easton, H.S. and Rolston, M.P. (1997): Effect of growth conditions on alkaloid concentrations in perennial ryegrass naturally infected with endophyte. *Neotyphodium*/Grass Interactions. C.W. Bacon and N.S. Hill (Ed.). Plenum Press. New York and London, 179-182.
- Latch, G.C.M. (1983): Incidence and control of ryegrass endophytes in New Zealand. Proceedings of Forage and Turfgrass Endophyte Workshop. Oregon (USA): Oregon State University Extention Service, 29-34.
- Latch, G.C.M. (1993): Physiological interactions of endophytic fungi and their hosts. Biotic stress tolerance imparted to grasses by endophytes. Agriculture, Ecosystems and Environment 44, 143-156.
- Latch, G.C.M. and Christensen, M.J. (1982): Ryegrass endophyte, incidence, and control. NZ. J.Agric.Res. 25, 443-448.
- Latch, G.C.M., Christensen, M.J. and Samuels, G.J. (1984): Five endophytes of *Lolium* and *Festuca* in New Zealand. Mycotorax 20, 535-550.
- Latch, G.C.M., Hunt, W.F. and Musgrave, D.R. (1985): Endophytic fungi affect growth of perennial ryegrass. NZ. J. Agric.Res. 28, 165-168.
- Latch, G.C.M., Potter L. R. and Tyler B. F. (1987): Incidence of endophytes in seeds from collections of *Lolium* and *Festuca* species. Annals of Applied Biology 111, 59-64.
- LeBars, J. and LeBars, P. (1996): Recent acute and subacute mycotoxicoses recognised in France. Vet. Research 27, 383-394.
- Leuchtmann, A. (1992): Systematics, Distribution, and Host Specificity of Grass Endophytes. Natural Toxins 1, 150-162.
- Leuchtmann, A. and Clay, K. (1993): Nonreciprocal compatibility between *Epichloe typhina* and four host grasses. Mycologia Bronx, The New York Botanical Garden. 85, 157-163.

Leuchtmann, A., Schardl, C.L. and Siegel, M.R. (1994): Sexual compatibility and taxonomy of a new species of *Epichloe* symbiotic with fine fescue. *Mycologia* 86, 802-812.

Levitt, J.: Responses of plants to environmental stresses. Vol. II. Water, radiation, salt, and other stresses. Academic Press. New York (1980).

Lewis, G.C. (1990): Herbage yield of nine genotypes of perennial ryegrass with and without infection by ryegrass endophyte. *Tests of Agrochemicals and Cultivars* 11, *Ann. Appl. Biol.* (Suppl.) 116: 108-109.

Lewis, G.C. (1993): Incidence of infection of grasses by endophytic fungi in the UK, and effects of infection on animal health, pest and disease damage and plant growth. *Proceedings of the Conference on Harmful and Beneficial Microorganisms in Pastures. Turf and Grassland, Padernborn*, pp. 161-167.

Lewis, G.C. (2000): Effect of drought stress on genotypes of *Lolium perenne* and other grass species with and without *Neotyphodium/Epichloe* infection. In: *4th International Neotyphodium/Grass Interactions Symposium*. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp. 201-206.

Lewis, G.C. and Clements, R.O. (1986): A survey of ryegrass endophyte (*Acremonium loliae*) in the U.K. and its apparent ineffectuality on a seed pest. *Journal of Agricultural Science, Cambridge*, 107, 633-638.

Lewis, G.C. and Clements, R.O. (1990): Effect of *Acremonium lolii* on herbage yield of *Lolium perenne* at three sites in the United Kingdom. *Proc. Int. Symp. on Acremonium /Grass Interactions*. S. S. Quinsenberry and R. E. Joost (Ed.). USA: New Orleans, 160.

Lewis, G.C., Bakken, A.K., MacDuff, J.H. and Raistrick, N. (1996): Effect of infection by the endophytic fungus *Acremonium lolii* on growth and nitrogen uptake by perennial ryegrass (*Lolium perenne*) in flowing solution culture. *Ann. Appl. Biol.* 129, 451-460.

Lewis, G.C., Ravel, C., Naffaa, W., Astier, C. and Charmet, G. (1997): Occurrence of *Acremonium* endophytes in wild populations of *Lolium* spp. in European countries and a relationship between level of infection and climate in France. *Ann. Appl. Biol.* 130, 227-238.

Lyons, P.C., Plattner, R.D. and Bacon, C.W. (1986): Occurrence of peptide and clavine ergot alkaloids in tall fescue grass. *Science (Washington DC)*. 232, 487-489.

Lyons, P.C., Evans, J.J. and Bacon C. W. (1990): Effects of the fungal endophyte *Acremonium coenophyllum* on nitrogen accumulation and metabolism in tall fescue. *Plant Physiol.* 92, 726-732.

Malinowski, D., Leuchtmann, A., Schmidt, D. and Nosberger, J. (1997a): Growth and water status in meadow fescue is affected by *Neotyphodium* and *Phialophora* endophyte species. *Agron. J.* 89, 673-678.

Malinowski, D., Leuchtmann, A., Schmidt, D. and Nosberger, J. (1997b): Symbiosis with *Neotyphodium uncinatum* endophyte may increase the competitive ability of meadow fescue. *Agron. J.* 89, 833-839.

Malinowski D.P., Beleski D.P. and Fedders J. M. (1999): Endophyte infection may affect the competitive ability of tall fescue grown with red clover. *Journal of agronomy and Crop Science* 183, 91-101.

Marks, S. and Clay, K. (1990): Effects of CO₂ enrichment, nutrient addition, and fungal endophyte-infection on the growth of two grasses. *Oecologia* 84, 207-214.

Marks, S. and Clay, K. (1996): Physiological responses of *Festuca arundinacea* to fungal endophyte infection. *New Phytol.* 133, 727-733.

Marschall, D., Tunali, B. and Nelson, L.R. (1999): Occurrence of fungal endophytes in species of wild *Triticum*. *Crop Science* 39, 1507-1512.

Mika, V. and Bumerl, J. (1991): Occurrence of endophytic fungi in varieties of *Festuca pratensis*. Int. Symp. "Progress in grass seed production". Roznov pod Radhostem. 14-16.Mai. Pp.12-16.

Moon, C.D., Miles, C.O. and Schardl, C.L. (2000): The evolutionary origins of *Neotyphodium* endophytes from grasses indigenous to the southern hemisphere. In: 4th International *Neotyphodium*/Grass Interactions Symposium. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. pp. 289-294.

Morgan-Jones, G. and Gams, W. (1982): Notes on Hyphomycetes. XLI. An endophyte of *Festuca arundinacea* and the anamorph of *Epichloe typhina*, new taxa in one of two new sections of *Acremonium*. *Mycotaxon* 15, 311-318.

Morgan-Jones, G., White, J.F., Jr. and Piontelli, E.L. (1990): Endophyte-host associations in forage grasses. XIII. *Acremonium chilense*, an undescribed endophyte occurring in *Dactylis glomerata* in Chile. *Mycotaxon* 39, 441-454.

Mortimer, P.H. and DiMenna, M.E. (1983): Ryegrass staggers: further substantiation of *Acremonium* endophyte aetiology and the discovery of weevil resistance of ryegrass pastures infected with *Lolium* endophyte. *Proc. NZ. Grassl. Assoc.* 44, 240-243.

Moy, M., Belanger F., Duncan R., Freehof A., Leary C., Meyer W., Sullivan R. and White J. F. (2000): Identification of epiphyllous mycelial nets on leaves of grasses infected by clavicipitaceous endophytes. *Symbiosis* 28, 291-302.

Mühle, E. (1971): Krankheiten und Schädlinge der Futtergräser. S. Hirzel Verlag Leipzig. S 147-152.

Nan, Z.B. and Li, C.J. (2000): *Neotyphodium* in native grasses in China and observations on endophyte/host interactions. In: 4th International *Neotyphodium*/Grass Interactions Symposium. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp 41-50.

Neill, J.C. (1940): The endophyte of rye-grass (*Lolium perenne*). *New Zealand J. Sci. Technol.* 2, 280A-291A.

Nelson, L.R., Marshall, D. and Tunali, B. (1993): Exploration for fungal endophyte in *Lolium* and other grass species of central Turkey. In: Proceedings of the 2. International Symposium on *Acremonium*/Grass Interactions. D.E. Hume, G.C.M Latch and H.S. Easton (Ed.). Palmerston-North. New Zealand. 11-13.

Nestler, A. (1898): Über einen in der Frucht von *Lolium temulentum* L. vorkommenden Pilz. Berlin: Berichte der Deutschen Botanischen Gesellschaft 16, 207-214.

Neubauer, H. (1902): Ueber die von A. Vogl entdeckte Pilzschicht in Loliumfrüchten. *Zentralblatt Bakteriologie* 9, 652-653.

Oldenburg, E. (1994): Occurrence of fungal endophytes in *Lolium perenne*. Bulletin of the International Conference on Harmful and Beneficial Microorganisms in Grassland, Pasture and Turf. 17, 99-104.

Oldenburg, E. (1997a): Endophytische Pilze in Gräsern - Auswirkungen auf die Pflanzenqualität und die Leistung von Weidetieren. In: Bericht über die 41.Jahrestagung der Gesellschaft für Pflanzenbauwissenschaften, AG Grünland und Futterbau. vom 28. - 30. 8. in Aulendorf, 69-74.

Oldenburg, E. (1997b): Endophytic fungi and alkaloid production in perennial ryegrass in Germany. Grass and Forage Science. 52, 425-431.

Oliveira, J.A. and Castro, P. (1997): Incidence and viability of *Acremonium* endophytes in tall fescue accessions from North Spain. Genetic Resources and Crop Evolution 44, 519-522.

Oliveira, J.A., Rottinghaus G.E., Collar J. and Castro P. (1997): Perennial ryegrass endophyte in Galicia, Northwest Spain. Journal of Agricultural Science. 129, 173-177.

Pedersen, J.F., Rodriguez-Kabana R. and Shelby R.A. (1988): Ryegrass cultivars and endophyte in tall fescue affect nematodes in grass and succeeding soybean. Agron J. 80, 811-814.

Petrini, O. (1991): Fungal endophytes of tree leaves. In: Microbial Ecology of Leaves. J.A. Andrews and S.S. Hirano (Ed.). New York. Springer-Verlag, 179-197.

Petrini, O., Sieber, T.N., Toti, L. and Viret, O. (1992): Ecology, metabolite production, and substrate utilization in endophytic fungi. Natural Toxins 1, 185-196.

Pfannmöller, M., Eggstein, S. and Schöberlein, W. (1994): Endophytes in European varieties of festuca species. IOBC/WPRS Bulletin OILB/SROP 17, 105-109.

Philipson, M.N. (1991): Septal and lateral pores in the fungal endophytes of two pasture grasses. National research Council of Canada. 69, 2740-2743.

Pinkerton, B.W., Rice, J.S. and Undersander, D.J. (1990): Germination in *Festuca arundinacea* as affected by the fungal endophyte, *Acremonium coenophialum*. Proc. Int. Symp. on *Acremonium*/Grass Interactions. S.S. Quinsenberry and R.E. Joost (Ed.). New Orleans, 176.

Porter, J.K., Bacon, C.W., Plattner, R.D. and Arrendale, R.F. (1987): Ergot peptide alkaloid spectra of *Claviceps*-infected tall fescue, wheat, and barley. J. Agric. Food Chem. 35, 359-361.

Prestidge, R.A., Pottinger R.P. and Barker G.M. (1982): An association of *Lolium* endophyte with ryegrass resistance to Argentine stem weevil. Proc. N. Z. Weed Pest Control Conf. 35, 119-122.

Ravel, C., Charmet G. and Balfourier, F. (1995): Influence of the fungal endophyte *Acremonium lolii* on agronomic traits of perennial ryegrass in France. Gras and Forage Science 50, 75-80.

Ravel, C., Astier, C., Balfourier, F. and Charmet, G. (1997): *Acremonium* Endophytes of perennial ryegrass (*Lolium perenne*) in France. Session 13- Constraints on Forage and Grassland Production, 13-17.

Ravel, C., Balfourier, F. and Guillaumin, J.J. (1999): Enhancement of yield and persistence of perennial ryegrass inoculated with one endophyte isolate in France. Agronomie 19, 635-644.

Read, J.C. and Camp, B.J. (1986): The effect of the fungal endophyte *Acremonium coenophialum* in tall fescue on animal performance, toxicity, and stand maintenance. *Agron. J.* 78, 848-850.

Reed, K.F.M., Leonforte, A., Cunningham, P.J., Walsh, P.J., Walsh, J.R., Allen, D.I., Johnstone, G.R. and Kearney, G. (2000): Incidence of ryegrass endophyte (*Neotyphodium lolii*) and diversity of associated alkaloid concentrations among naturalised populations of perennial ryegrass (*Lolium perenne* L.). *Australian Journal of Agricultural Research* 51, 569-578.

Reynolds, J.H., Walker, C.L. and Sams, C.L. (1997): Photosynthesis and chlorophyll fluorescence in tall fescue. *Neotyphodium/Grass Interactions*. C.W. Bacon and N.S. Hill (Ed.). Plenum Press. New York and London, 199-200.

Rice, J.S., Pinkerton, B.W. and Undersander, D.J. (1990): Imbibition of Tall Fescue as affected by *Acremonium coenophialum*. *Proc. Int. Symp. on Acremonium/Grass Interactions*. S.S. Quinsenberry and R.E. Joost (Ed.). New Orleans, 185-188.

Rice, J.S., Ferguson, N.H., Pinkerton, B.W. and Stringer, W.C. (1994): Alteration of phenotypic variances by the endophyte of tall fescue. *Crop Sci.* 34, 1487-1489.

Richardson, M.D., Bacon, C.W. and Hoveland, C.S. (1990): The effect of endophyte removal on gas exchange in tall fescue. *Proc. Int. Symp. on Acremonium/Grass Interactions*. S.S. Quinsenberry and R.E. Joost (Ed.). New Orleans, 189-193.

Richardson, M.D., Chapman, G.W., Jr., Hoveland, C.S. and Bacon, C.W. (1992): Sugar alcohols in endophyte-infected tall fescue under drought. *Crop Sci.* 32, 1060-1061.

Richardson, M.D., Hoveland, C.S. and Bacon, C.W. (1993): Photosynthesis and stomatal conductance of symbiotic and nonsymbiotic tall fescue. *Crop Sci.* 33, 145-149.

Rowan, D.D., Hund, M.B. and Gaynor, D.L. (1986): Peramine, a novel insect feeding deterrent from ryegrass infected with the endophyte *Acremonium loliæ*. *Journal of the Chemical Society-Chemical Communications* 12, 935-936.

Rowan, D.D. (1993): Lolitrem, peramine and paxilline: mycotoxins of the ryegrass/endophyte interaction. *Agriculture, Ecosystems and Environment* 44, 103-122.

Saha, D.C., Johnson-Cicalese, J.M., Van Heemstra, M.I., and Funk, C.R. (1987): Occurrence and significance of endophytic fungi in the fine fescues. *Plant Disease* 71: 1021-1024.

Sako, N. and Stahmann, M. A. (1972): Multiple molecular forms of enzymes in barley leaves infected with *Erysiphe graminis* f.sp. *hordei*. *Physiol. Plant Pathol.* 2: 217-226

Sampson, K. (1933): The systemic infection of grasses by *Epichloe typhina* (Pers.) Tul. *Trans. Br. Mycol. Soc.* 18, 30-47.

Saikkonen, K., Ahlholm J., Helander, M., Lehtimaki, S. and Niemelainen, O. (2000): Endophytic fungi in wild and cultivated grasses in Finland. *Ecography* 23, 360-366.

Scandalios, J.G. (1974): Isozymes in development and differentiation. *Ann. Rev. Plant Physiol.* 25, 225-258.

Schardl, C.L., Leuchtmann, A., Tsai, H.F., Collet, M.A., Watt, D.M. and Scott, D.B. (1994): Origin of a fungal symbiont of perennial ryegrass by interspecific hybridization of a mutualist with the ryegrass choke pathogen, *Epichloe typhina*. Genetics. Baltimore, Md. Genetics Society of America. 136, 1307-1317.

Schardl, C.L., Leuchtmann, A., Chung, K.R., Penny, D. and Siegel, M.R. (1997): Coevolution by common descent of fungal symbionts (*Epichloe* spp.) and grass hosts. Molecular Biology and Evolution 14, 133-143.

Schardl, C.L. and Wilkinson, H.H. (2000): Hybridization and Cospeciation Hypotheses for the evolution of grass endophytes. In: Microbial Endophytes. Ed: C.W. Bacon and J.F. White. Marcel-Dekker-Inc. New York 2000.

Schilling, G. (1987): Pflanzenernährung und Düngung. Teil 1: Pflanzenernährung. VEB Deutscher Landwirtschaftsverlag DDR. Berlin. 240 S.

Schöberlein, W. (1970): Untersuchungen an mehrjährigen Gramineen über die Abhängigkeit der generativen Triebbildung vom Entwicklungsstand der Herbsttriebe und Möglichkeiten seiner Beeinflussung im Hinblick auf höhere Saatguterträge. Diss. Halle. 170 S.

Schöberlein, W. und Pfannmöller, M. (1996): Endophytische Pilze in mehrjährigen Nutzgräsern - Probleme oder Vorteile. In: „Bericht über die 47. Arbeitstagung der Vereinigung österreichischer Pflanzenzüchter“ vom 26.-28. 11. in Gumpenstein, Verl. und Druck der BA für alpenländische Landw. Gumpenstein, S. 185-191.

Schöberlein, W., Pfannmöller, M., Eggestein, S. and Szabová (1998): Effects of endophyte-infected *Festuca pratensis* plots on population density of nematode species in the soil. In: Breeding for a Multifunctional Agriculture. B. Boller and F.J. Stadelmann (Ed.). Swiss Federal Research Station for Agroecology and Agriculture. Zürich-Reckenholz. Pp. 218-220.

Siegel, M.R., Latch, G.C.M. and Johnson, M.C. (1985): Acremonium fungal endophytes of tall fescue and perennial ryegrass: significance and control. Plant Disease 69, 179-183.

Siegel, M.R., Latch, G.C.M. and Johnson, M.C. (1987): Fungal endophytes of grasses. Ann. Rev. Phytopathol. 25, 293-315.

Siegel, M.R., Latch, G.C.M., Bush, L.P., Fannin, F.F., Rowan, D.D., Tapper, B.A., Bacon, C.W. and Johnson, M.C. (1990): Fungal endophyte-infected grasses: alkaloid accumulation and aphid response. J-Chem-Ecol. New York: Plenum Press 16.,3301-3315.

Siegel, M.R., Schardl, C.L. and Phillips, T.D. (1995): Incidence and compatibility of nonclavicipitaceous fungal endophytes in *Festuca* and *Lolium* grass species. Mycologia 87, 196-202.

Smith, G.S., Johnston, C.M. and Cornforth, I.S. (1983): Comparison of nutrient solutions for growth of plants in sand culture. New Phytol. (1983) 94, 537-548.

Schmidt, D. (1991): Les endophytes de la féstue des prés. Revue Suisse Agric. 23, 369-375.

Schmidt, D. (1994): Du nouveau sue les endophytes de la féstue des prés. Revue Suisse Agric. 26, 57-63.

Shelby, R.A. and Dalrymple, L.W. (1987): Incidence and distribution of the tall fescue endophyte in the United States. *Plant Disease* 71: 783-786.

Stumpe, H., Garz, J. und Schliephake, W. (1995): Das Versuchsfeld der Landwirtschaftlichen Fakultät in Halle und seine Dauerversuche. Mitteilungen der Deutschen Bodenkundlichen Gesellschaft, Exkursionsführer zur Jahrestagung 1995 in Halle. Herausgeber: P. Hugenroth. 77, ISSN 0343-107X, S. 375-382.

Sun, S., Clarke, B.B., and Huff, D.R. (1993): Enhanced performance and new sources of *Acremonium typhinum* in fine fescues. Proc. Sec. Int. Symp.on *Acremonium*/Grass Interactions. Hume D.E., Latch, G. C. M. and Easton, H. S. (Ed.). New Zealand: Palmerston North, 23-26.

Tapper, B.A., Rowan, D.D. and Latch, G.C.M. (1989): Detection and measurement of the alkaloid peramine in endophyte-infected grasses. *J. Chromatogr.* 463: 133-138.

Thom, E.R., Clark, D.A. and Waugh, C.D. (1999): Growth, persistence, and alkaloid levels of endophyte-infected and endophyte-free ryegrass pastures grazed by dairy cows in northern New Zealand. *New Zealand Journal of Agricultural Research* 42, 241-253.

Tyler, B.F., Chorlton, K.H. and Thomas I.D. (1981): Extending genetic resources. Annual Report for Welsh Plant Breeding Station for 1980, pp. 44-46.

Van-Essen, G.J., Blom, M. and Frink-Gremmels, J. (1995): Rye-grass intoxication in horses. *Tijdschrift voor diergeneeskunde* 120, 710-711.

Van-Santen, E. and Collins, D.J. (1991): Genetic diversity in tall fescue ecotypes from Alabama. 1. Maturity, morphological traits, and disease reaction. *Plant Breeding* 107, 210-217.

Vogl, A.E. (1898): Mehl und die anderen Mehlprodukte der Cerealien und Leguminosen. *Nahrungsm. Unters. Hyg. Warenk.* 12, 25-29.

Wäli, P., Saikkonen, K., Helander, M., Lehtimäki, S. and Lehtonen, P. (2000): Seed transmitted endophytic fungi in wild grass populations in Finland. In: 4th International *Neotyphodium*/Grass Interactions Symposium. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp 93-96.

Warnstorff, K. und Dörfel, H. (1998): Ein Programm zur Kontingenztafel- und Kontrastanalyse. *Zeitschrift für Argainformatik*. 2, 38-42.

Wendel, J.F. and Weeden, N.F. (1989): Visualization and interpretation of plant isozymes. *Isozymes in Plant Biology*. Ed.: D.E. Soltis & P.S. Soltis. Dioscorides, Portland, Oregon.

West, C.P., Izekor, E., Oosterhuis, D.M. and Robbins, R.T. (1988): The effect of *Acremonium coenophialum* on the growth and nematode infestation of tall fescue. *Plant Soil* 112, 3-6.

West, C.P., Izekor, E., Elmi, A., Robbins, R.T. and Turner, K.E. (1989): Endophyte effects on drought tolerance, nematode infestation and persistence of tall fescue. In: Proceedings of the Arkansas Fescue Toxicosis Conference. C.P. West (Ed.). Arkansas Agricultural Experiment Station. US. Pp. 23-27.

West, C.P., Izekor, E., Turner, K.E. and Elmi, A. (1993): Endophyte effects on growth and persistence of tall fescue along a water-supply gradient. *Agron. J.* 85, 264-270.

- Wheatley, W.M. (2000): Endophytes as a marketing tool to increase the sail of perennial ryegrass cultivars in south-eastern Australia. In: 4th International *Neotyphodium*/Grass Interactions Symposium. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp169-174.
- White, J.F., Jr. (1988): Endophyte-host associations in forage grasses. XI A proposal concerning origin and evolution. *Mycologia* 80, 442-446.
- White, J.F. and Cole, T.G. (1986): Endophyte-host associations in forage grasses. V. Occurrence of fungal endophytes in certain species of *Bromus* and *Poa*. *Mycologia* 78, 846-850.
- White, J.F., Cole, G.T. and Morgan-Jones, G. (1987): Endophyte-host associations in forage grasses. VI. A new species of *Acremonium* isolated from *Festuca arizonica*. *Mycologia* 79, 148-152.
- White, J.F. and Morgan-Jones, G. (1987a): Endophyte-host associations in forage grasses. VII. *Acremonium chisosum*, a new species isolated from *Stipa eminens* in Texas. *Mycotaxon* 28, 179-189.
- White, J.F. and Morgan-Jones, G. (1987b): Endophyte-host associations in forage grasses. X. Cultural studies on some species of *Acremonium* Sect. *Albo-Lanosa*, including a new species, *A. starrii*. *Mycotaxon* 30, 87-95.
- White, J.F., Jr., Meyer, W., Sullivan, R., Moy, M., Balady, G. and Cabral, D. (2000): Evolution of *Epichloe/Neotyphodium* endophytes. In: 4th International *Neotyphodium*/Grass Interactions Symposium. V.H. Paul and P.D. Dapprich (Ed.) Soest, Germany. Pp 17-30.
- White, R.H., Engelke, M.C., Morton, S.J., Johnson-Cicalese, J.M. and Ruemmele, B.A. (1992): *Acremonium* endophyte effects on tall fescue drought tolerance. *Crop Sci.* 32, 1392-1396.
- Widdup, K.H. and Ryan, D.L. (1992): Forage potential of wild populations of perennial ryegrass collected from southern New Zealand farms. In: Proceedings of the New Zealand Grassland Association 54, 161-165.
- Wortha, H.-P. (1989): A new statistical method for mixed variates analyses in epidemiological research. *Biometrical Journal*. 31, 941-955.
- Yates, S.G., Plattner, R.D. and Garner, G.B. (1985): Detection of ergopeptine alkaloids in endophyte-infected, toxic Ky-31 tall fescue by mass spectrometry. *J. Agric. Food Chem.* 33, 719-722.
- Yue, Q., Logendra, S., Freehoff, A. and Richardson, M.D. (1997): Alkaloids of turf-type fine fescue (*Festuca* sp.). *Neotyphodium*/Grass Interactions. C.W. Bacon and N.S. Hill (Ed.). Plenum Press. New York and London, 285-287.
- Zabalgogeazcoa, I., Vazquez de Aldana, B. R., Garcia Criado, B. and Garcia Ciudad, A. (1997): The infection of *Festuca rubra* by the fungal endophyte *Epichloe festucae* in Mediterranean permanent grasslands. *Grass and Forage Science* 54, 91-95.
- Zadoks; J.C., Chang, T.T. and Konzak, C.F. (1974): A decimal code for the growth stages of cereals. *Weed Research - Oxford* 14, 415-121.