



**Bayer CropScience**

**Bayer CropScience LP  
P.O. Box 12014  
2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709  
1-866-99BAYER (1-866-992-2937)**

**PROLINE® 480 SC Fungicide**

**EPA Reg. No. 264-825**

**For Use On: Bushberry subgroup; Low growing berry subgroup, except strawberry; Cucurbit vegetables**

*This supplemental label expires on 11/12/2016 and must not be used or distributed after this date.*

**Supplemental Label**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Read this label and the product package label before using this product. This Supplemental Label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the product label for PROLINE® 480 SC Fungicide attached to the container.

**USE DIRECTIONS FOR SPECIFIC CROPS**

<b>APPLICATION DIRECTIONS</b>		
<b>CROP</b>	<b>DISEASES CONTROLLED</b>	<b>RATE OF PROLINE 480 SC FUNGICIDE</b>
<b>Bushberry subgroup:</b> Aronia berry; blueberry (highbush and lowbush); Chilean guava; highbush cranberry; currant (black, buffalo, and red); elderberry; European barberry; gooseberry; edible honeysuckle; huckleberry; jostaberry; juneberry (Saskatoon berry); lingonberry; native currant; salal; sea buckthorn; and cultivars, varieties, and/or hybrids of these.	Septoria leaf spot <i>(Septoria spp.)</i>	5.7 fl oz. per acre
	Monilinia blight <i>(Monilinia vaccinii-corymbosi)</i>	
	Valdensinia leaf spot <i>(Valdensinia heterodoxa)</i>	
	Leaf rust <i>(Thekopsora minima)</i>	
	Anthracnose <i>(Colletotrichum gloeosporioides)</i>	
	Botrytis blight <i>(Botrytis cinerea)</i>	
	Alternaria fruit rot <i>(Alternaria spp.)</i>	
	White pine blister rust <i>(Cronartium ribicola)</i>	
PROLINE 480 SC FUNGICIDE may be applied by either ground or chemigation application equipment. Apply PROLINE 480 SC FUNGICIDE at the first sign of disease.		

*Proline is a Registered trademark of Bayer.*

SAL 12/18/13 - PRL

**Other Requirements:** Apply up to two (2) applications of PROLINE 480 SC per year. Repeat applications as needed using a 7- to 10-day spray interval if conditions remain favorable for continued or increasing disease development.

A maximum of 11.4 fl oz. of PROLINE 480 SC may be applied per year. Do not apply within 7 days of harvest.

**APPLICATION DIRECTIONS**

CROP	DISEASES CONTROLLED	RATE OF PROLINE 480 SC FUNGICIDE
<p><b>Low growing berry subgroup, except strawberry:</b>            Bearberry; bilberry; cloudberry; cranberry; muntries; partridgeberry; and cultivars, varieties, and/or hybrids of these</p>	<p><b>Fruit rot:</b>  <i>Coleophoma empetri</i>  <i>Glomerella cingulata</i>  <i>Phyllosticta vaccinii</i>  <i>Physalospora vaccinii</i>  <i>Allantophomopsis lycopodina</i>  <i>Allantophomopsis cytispora</i>  <i>Fusicoccum putrefaciens</i>  <i>Penicillium</i> spp.  <i>Phomopsis vaccinii</i>  <i>Colletotrichum acutatum</i>  <i>Botrytis</i> spp.  <i>Monilinia</i> spp.</p> <p>Valdensinia leaf spot            (<i>Valdensinia heterodoxa</i>)</p>	<p>5.0 fl oz. per acre</p>
<p>PROLINE 480 SC FUNGICIDE may be applied by either ground or chemigation application equipment. For best control of fruit rots begin applications at early bloom. Make a second application of PROLINE 480 SC or another approved fungicide 7-10 days later.</p>		

**Other Requirements:** Apply up to two (2) applications of PROLINE 480 SC per year. Repeat applications as needed using a 7- to 10-day spray interval if conditions remain favorable for continued or increasing disease development.

A maximum of 10.0 fl oz. of PROLINE 480 SC may be applied per year. Do not apply within 45 days of harvest.

APPLICATION DIRECTIONS		
CROP	DISEASES CONTROLLED	RATE OF PROLINE 480 SC FUNGICIDE
<p><b>Cucurbit vegetables:</b>            Chayote (fruit); Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; edible gourd (includes hyotan, cucuzza, hechima, Chinese okra);  <i>Momordica</i> spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (includes cantaloupe); pumpkin; squash (summer and winter, includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon</p>	Fusarium wilt Fusarium blight <i>(Fusarium oxysporum)</i> <i>(Fusarium spp.)</i>	5.7 fl oz. per acre (soil)
	Gummy stem blight <i>(Didymella spp.)</i>  Southern blight <i>(Sclerotium roflsii)</i>  Powdery mildew <i>(Sphaerotheca fuliginea / Podosphaera xanthii)</i> <i>(Erysiphe cichoracearum)</i>	5.7 fl oz. per acre (foliar)
<p>PROLINE 480 SC FUNGICIDE may be applied by either ground or chemigation application equipment (including drip irrigation). Do not use in water used for hand transplanting. Not for use in greenhouse/transplant house.</p>		
<p><b>Other Requirements:</b> Apply up to one (1) soil application and two (2) foliar applications of PROLINE 480 SC per year. Repeat applications as needed using a 5- to 10-day spray interval if conditions remain favorable for continued or increasing disease development.</p> <p>A maximum of 17.1 fl oz. of PROLINE 480 SC may be applied per year. Do not apply within 7 days of harvest.</p>		

**For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
 For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)**

*As with any crop-protection product, always read and follow label directions.  
 For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

*Proline is a Registered trademark of Bayer.*

SAL 12/18/13 - PRL