

SECTION 24(c) SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

Roper DF Rainshield[®]

EPA Reg. No. 34704-1063

EPA SLN No. WA-130003

For Disease Control in Small-Seeded Vegetable Seed Crops

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2017.

ATTENTION

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.

• Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Directions for Use

Begin applications of Roper DF Rainshield fungicide when disease first appears or is reported in the area. Apply Roper DF Rainshield fungicide at the rate of 2 pounds per acre in thorough coverage sprays. Repeat applications on a 7 to 10 day application schedule until the threat of disease is over.

Use sufficient spray volume to achieve thorough coverage. Refer to container label for use directions and restrictions for ground or aerial applications.

Roper DF Rainshield fungicide may be applied to the following small-seeded vegetable seed crops for control of listed diseases:

Seed Crop	Diseases Controlled
Allium spp. (Leek, bunching and dry bulb onions)	Botrytis leaf blight (Botrytis spp.)
	Downy mildew (Peronospora destructor)
	Neck rot (<i>Botrytis</i> spp.)
	Purple blotch (Alternaria porri)
Arugula	Alternaria leaf spot (Alternaria sp.)
Beet	Alternaria leaf spot (Alternaria sp.)
	Cercospora leaf spot (Cercospora beticola)
	Downy mildew (Peronospora farinosa)
	Rust (Uromyces betae)
Carrot	Leaf blight (Alternaria dauci)
	Leaf spot (Cercospora carotae)
	Rust (Uromyces betae)
Coriander	Leaf spot (Alternaria spp.)
Brassica and Raphanus spp.:	Black spot (Alternaria alternata and A.
Broccoli, broccoli raab, Brussels sprout, cabbage,	brassicola)
cauliflower, Chinese cabbage, Chinese mustard,	Cercospora leaf spot (Cercospora brasiciccola)
collard, kale, kohlrabi, mustard, radish (other than	Downy mildew (Peronospora parasitica)
diakon), rape, rutabaga, spinach mustard, turnip	
Dill	Alternaria leaf spot (Alternaria sp.)
Endive	Downy mildew (Bremia lactucae)
	Leaf spot (Cercospora cichoria)

Lettuce	Anthracnose (<i>Microdochium panattoniamnum</i>) Cercospora leaf spot (<i>Cercospora longissima</i>) Downy mildew (<i>Bremia lactucae</i>)
Parsley	Alternaria leaf spot (Alternaria sp.)
Parsnip	Alternaria leaf spot (Alternaria sp.)
Spinach	Downy mildew (<i>Peronospora effusa</i> and <i>P. farinosa</i>) Leaf spot (<i>Alternaria</i> spp.) Leaf spot (<i>Cercospora beticola</i>) Leaf spot (<i>Cladosporium variable</i>) Leaf spot (<i>Stemphylium botryosum</i>)
Swiss Chard	Cercospora leaf spot (Cercospora beticola) Rust (Uromyces batae)

Restrictions / Precautions:

- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.
- Do not apply more than 24 lb (18.0 lb active ingredient) per acre per crop. Do not make more than 12 applications per crop.
- For use under this SLN label do not apply this product through any type of irrigation system.
- All seed screenings from small-seeded vegetable seed crops shall be disposed of in such a way that
 they cannot be distributed or used for human food or animal feed. The seed conditioner shall keep
 records of screening disposal for three years from the date of disposal and shall furnish the records to
 the director immediately upon request. Conditioner disposal records shall consist of documentation of
 on-farm disposal, disposal at a controlled dumpsite, incinerator, composter or other equivalent
 disposal site and shall include the lot numbers, amount of material disposed of, the grower(s), and the
 date of disposal.
- No portion of the small-seeded vegetable seed crop plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
- Seed from small-seeded vegetable seed crops shall bear a tag or container label which forbids use of the seed for human consumption or animal
- Seed from small-seeded vegetable seed crops may not be distributed for human consumption or animal feed.
- This pesticide is toxic to aquatic organisms. Roper DF Rainshield should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the Washington State Department of Agriculture Endangered Species web site at http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the Washington State Department of Agriculture Waste Pesticide Program web site at http://agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in the state of Washington.

Rainshield is a trademark of Dow AgroSciences LLC Roper is a trademark of Loveland Products, Inc.

> Section 24(c) Registrant: Loveland Products, Inc. P.O Box 1286 Greeley, Colorado 80632-1286