

# QUICK IDENTIFICATION COMMON WILDFLOWERS



**CENTRAL VICTORIA  
& THE HIGHLANDS**



---

# QUICK IDENTIFICATION LOCAL WILDFLOWERS

---

By David & Debbie Hibbert

This draft eBook is a growing educational database of some of the more common wildflowers found in Central Victoria. It is not a comprehensive resource. The book is based almost entirely on images taken by people with local links or visitors to our district. The database grows when we receive photographic submissions. If you can help build this educational resource, please contact us.

Plant count: **1** **5** **2** and growing.



## Contributors:

Ross & Nancy Coller, Geoff Edwards, Debbie Hibbert, Denis Martin,  
Kelly Petersen, Robert (Bob) Tate, Ron Turner, Arnis Heislars,  
Kathie Maynes, Jill Miles.

Contributing Photographers:  
Denis Martin, David Hibbert

Project commenced 1st April 2013  
Reached 100 wildflowers in September 2021

**An Artworkz Free Publication**





# FORWARD



By Robert (Bob) Tate


Nature study was an integral part of study in Primary Schools up until the mid 1900s. In those days nature was much closer to us than now, with the bush not far away and country students having easy access to nature on the way to and from school, and of course at home.

Unfortunately in this day and age, more emphasis is placed on environmental issues, though to appreciate this we still need to know just what is in the environment.

That is where I hope this eBook will help in raising awareness natural flora which can be seen more easily. It can help students in identifying the various species shown and thereby help them gain an interest which could eventually be in birds, animals, insects, plants, fungi, rocks or many other aspects of nature.


Bob Tate. B.V.Sc. L.D.A.

February 2013.





# COLOURS




Daisy

Guinea-flower

Lily

Orchid

Pea




Daisy

Lily



Pea

Orchid

Violet



Orchid



OTHER



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Buttercup



Also known as the common buttercup and yarraalgamba, it is found in eastern Australia from Queensland to South Australia and Tasmania. It grows to 60 cm high and prefers bushland, woodlands and open forests, but can be found up to 400 m. Flowers appear from September to April with a width to 40 mm.

Family: Ranunculaceae  
Genus: *Ranunculus*  
Species: *Ranunculus lappaceus*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Creeping woodsorrel




Also known as sleeping beauty, it is a ground cover introduced from Europe and found in all states of Australia. It grows to 20 cm and prefers bushland, woodlands, forests and is often considered a weed in parks and gardens. Its tiny flowers appear in Spring. The leaves are edible in small amounts.

Family: Oxalidaceae  
Genus: *Oxalis*  
Species: *Oxalis corniculata*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Great mullein



A biannual native of Europe, Africa and Asia and introduced to all states of Australia (except the NT) where it is a weed. It grows to 2.4 m with hairy leaves and yellow flowers which appear from November with a width to 50 mm. It is often on roadsides and has uses in traditional medicines. Leaves and stem are covered in tiny white hairs.

Family: Scrophulariaceae  
Genus: *Verbascum*  
Species: *Verbascum thapsus*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Hop goodenia



An erect native shrub endemic to eastern Australia from Queensland to SA and Tasmania. It grows to 2 m, though can appear more prostrate. It prefers moist soils in woodlands and forests. Flowers appear from September to February with a width to 20 mm. Leaves are toothed, egg shaped to elliptic, often sticky and may look varnished.

Family: Goodeniaceae  
Genus: *Goodenia*  
Species: *Goodenia ovata*



# WILDFLOWER



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Hop trefoil



Also known as field clover and low hop clover, it is a native clover of Africa, Europe and Western Asia. It was introduced to Australia where it is now naturalised in all states (except the NT). It grows to 30 cm high and flowers appear from Sept. to Dec with a width of 20 mm and it prefers grasslands, bushland, and woodlands. It is a valuable pastoral herb.

Family: Fabaceae  
Genus: *Trifolium*  
Species: *Trifolium campestre*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Ivy goodenia



A native flowering plant found in eastern Australia from Queensland to Victoria. It grows to 80 cm, though can appear more prostrate. It prefers drier soils in woodlands and forests. It flowers in Spring and Summer and has a flower width to 22 mm. Leaves are green on top and often covered with tiny hairs (tomentose) underneath.

Family: Goodeniaceae  
Genus: *Goodenia*  
Species: *G. hederacea subsp. hederacea*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Musk monkeyflower



Also known as eyebright and common musk. It is found in south east Australia from NSW to SA and Tasmania. It grows to a height of 40 cm and has a yellow flower that appears from December to February with a width to 18 mm. It shaded areas in bushland, and forests. It was once grown for its fragrance until its smell changed worldwide circa 1913.

Family: Phrymaceae  
Genus: *Erythranthe*  
Species: *Erythranthe moschata*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Shrubby St. Johns wort



Also known as sweet amber or tutsan, it is an introduced perennial weed with poisonous berries found in south eastern Australia from NSW to SA and Tasmania. It grows to a height of 70 cm, has a flower width of 25 mm and prefers wet areas in bushland, grasslands, woodlands and forests.

Family: Hypericaceae  
Genus: *Hypericum*  
Species: *Hypericum androsaemum*



# WILDFLOWER



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Small St. John's wort



Also known as grassy St. Johns wort, it is a native to Australia. It is found in all states of Australia, usually grows to around 20 cm though can reach 70 cm in some instances. Flowers appear from November to January on solitary stems with a width to 13 mm that. It prefers drier countryside, bushlands, woodlands and forests.

Family: Hypericaceae  
Genus: *Hypericum*  
Species: *Hypericum gramineum*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## St. Johns wort



An introduced perennial plant considered a noxious weed and found in all states of Australia. It grows to a height of 30 cm, has a flower width of 25 mm which appears from November to February, and prefers bushland, grasslands, woodlands and forests. It is a medicinal herb with antidepressant and anti-inflammatory properties and is cultivated commercially.

Family: Hypericaceae  
Genus: *Hypericum*  
Species: *Hypericum perforatum*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Tiny star



Also known as grass-star, star lily, African potato and yellow stars, it is a small flowering plant found in all states of Australia (except Queensland and the NT). It grows to 25 cm with flower width to 25 mm. It prefers dryer soils in bushland, woodlands and open forests and flowers from July to November with early Nancy and hardenbergia.

Family: Hypoxidaceae  
Genus: *Hypoxis*  
Species: *Hypoxis glabella*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Twiggy mullein



A biannual native of Europe, Africa and Asia introduced to all states of Australia (except the NT) where it is a weed. It grows to 2.4 m with dark green leaves and yellow flowers which appear from October to April with a width to 50 mm. Often found on roadsides, it is highly regarded in traditional medicine. Unlike *Verbascum thapsus*, it is mostly hairless.

Family: Scrophulariaceae  
Genus: *Verbascum*  
Species: *Verbascum virgatum*



# WILDFLOWER



## Wild radish



Also known as white charlock or joined charlock, it is native to Asia, Europe and Africa and introduced to all states of Australia. It grows to 75 cm, has a flower width to 25 mm and disturbed soils along roadsides. White to pale yellow flowers appear from January to March. In Australia it is an invasive weed. It is a popular species for western honey bees.

Family: Brassicaceae  
Genus: *Raphanus*  
Species: *Raphanus raphanistrum*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Yellow woodsorrel



Also known as upright yellow sorrel, it's a ground cover introduced from America and Europe and introduced to all states of Australia. It grows in clumps to 30 cm, has a flower width to 32 mm and prefers bushland, woodlands and grasslands. Edible flowers appear from late July to November. Its edible leaf is heart-shaped and citrus flavoured.

Family: Oxalidaceae  
Genus: *Oxalis*  
Species: *Oxalis stricta*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# DAISY



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Black eyes



Also known as the European umbrella milkwort, it is native to Europe and introduced to all states of Australia (except the NT). It grows to 50 cm prefers bushlands and woodlands and is rarely unbranched. Pale yellow flowers appear on long stems from Nov to Jan with a width to 20 mm. It is very similar to *Tolpis umbellate* which may be in the same patches.

Family: Asteraceae  
Genus: *Tolpis*  
Species: *Tolpis barbata*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Button everlasting



A native perennial herb also found in all states of Australia (except WA and the NT). It grows to 50 cm on a stem, prefers clay loamy soils in bushland and woodlands. Yellow flowers (rarely white) appear from September to November with a width to 25 mm. It was previously known as *Helichrysum scorpioides*.

Family: Asteraceae  
Genus: *Coronidium*  
Species: *Coronidium scorpioides*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Cape weed



Also known as cape dandelion, it is a weed in the sunflower family introduced from South Africa and found in all states of Australia. It grows to 40 cm and prefers countryside, bushland and woodlands. It has a bright yellow flower with a black centre that appears in spring with a width to 7 cm.

Family: Asteraceae  
Genus: *Arctotheca*  
Species: *Arctotheca calendula*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Cascade everlasting



An aromatic shrub in the daisy family found in south eastern Australia from NSW to Victoria and Tasmania. It grows to 2 m and has tiny cream flowers that appear in summer as a clustered flowerhead. It prefers bushland and forests in subalpine and sheltered alpine areas, often amongst granite rocks. Leaves grow to 10 mm and 1.5 mm wide.

Family: Asteraceae  
Genus: *Gnaphalieveae*  
Species: *Ozothamnus secundiflorus*



# DAISY



VIC N.S.W QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED

## Clustered everlasting



A perennial herb ground cover that grows to 1 m and prefers drier rocky soils in open bushland and woodlands. It has a yellow clustered flower head width of 100 mm heads, with individual flowers in the head growing to a width of 5 mm. It flowers from September to February, though flowers quickly dry out in hot periods.

Family: Asteraceae  
Genus: *Chrysocephalum*  
Species: *Chrysocephalum semipapposum*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common Billy buttons



Also known as woollyheads, it is native to Australia and are found in all states of Australia (except the Northern Territory). It grows to 70 cm and prefers grasslands, scrublands and eucalypt forests. They grow in soils including sands, gravels, clays and loams, and flower in Spring.

Family: Asteraceae  
Genus: *Craspedia*  
Species: *Craspedia variabilis*



VIC N.S.W QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED

## Common cats ears




Also known as false dandelion, it was introduced to all states of Australia. Forked flower stems grows to 70 cm and flowers appear from Sep -Feb with a width to 5 cm. Hairy leaves grow to 20 cm. It prefers bushlands, woodlands, grasslands and countryside. All parts of the plant exudes a milky sap and are edible. Roots can be roasted into a coffee substitute.

Family: Asteraceae  
Genus: *Hypochaeris*  
Species: *Hypochaeris radicata*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common everlasting



Also known as yellow buttons, it is a perennial herb ground cover that grows to 50 cm and prefers drier rocky soils in bushland, woodlands and grasslands. Yellow flower heads appear from September to December with a width to 15 mm. Flowers quickly dry out in hot periods. Silvery grey hairy leaves clasp tight to the stem..

Family: Asteraceae  
Genus: *Chrysocephalum*  
Species: *Chrysocephalum apiculatum*



# DAISY



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common sow-thistle



Also known as the milky tassel, it is an introduced weed in the dandelion tribe, native to Europe and Asia. It is found in all states of Australia, grows to 100 cm and prefers full sun in parks and gardens as well as bushland and woodlands. It flowers from late spring to late summer. Young leaves are edible raw or cooked like spinach. Young roots are edible.

Family: Asteraceae  
Genus: *Sonchus*  
Species: *Sonchus oleraceus*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Dandelion



Also known as the common dandelion, it is a tap-rooted perennial that grows to 70 cm and a flower width of up to 5 cm. Leaves in their rosette grow to 40 cm. It prefers bushlands, woodlands, forests, grasslands and countryside and some of the plant is edible or used for medical purposes. It is commonly found in urban lawns.

Family: Asteraceae  
Genus: *Taraxacum*  
Species: *Taraxacum officinale*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Golden everlasting




Also known as the strawflower, it is an Australian native found in all states of Australia. It grows to 1 m, though is shorter in higher altitudes. It prefers most habitats and flowers from September to May. It is a source of food for moths and butterflies and also attracts hoverflies, native bees, grasshoppers and small beetles.

Family: Asteraceae  
Genus: *Xerochrysum*  
Species: *Xerochrysum bracteatum*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Scaly buttons




A native herb found in south east Australia from QLD to SA and Tasmania. It prefers bushland, woodlands and drier forests. Yellow flowers appear from October to December on a flower stem to 30 cm with a flower width to 12 mm. Stems are motley and surprisingly rigid and strong.

Family: Asteraceae  
Genus: *Leptorhynchos*  
Species: *Leptorhynchos squamatus*



# DAISY



## Yam daisy



A native perennial herb also called murrnong, found in all states of Australia (except the Northern Territory). It grows to 30 cm, has a tufted rosette and prefers bushland, woodlands and drier forests and flowers from late Spring to late Summer. It has an edible tuber like a carrot and was a preferred food for early Aboriginals found in abundance.

Family: Asteraceae  
Genus: *Microseris*  
Species: *Microseris lanceolata*

VIC

N.S.W

QLD

S.A.

W.A.

N.T.

TAS

NATIVE

WEED

INTRODUCED

NATURALISED

# GUINEA-FLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Erect guinea-flower



One of numerous shrubs in the *Hibbertia* genus. It is a small native shrub endemic to eastern Australia from QLD to SA and Tasmania. It grows to 60 cm with a flower width of 50 mm, which appears from mid spring to mid summer. It prefers rocky soils in bushland, woodlands and forests and also appears in the Grampians.

Family: Dilleniaceae  
Genus: *Hibbertia*  
Species: *Hibbertia riparia*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Euroa guinea-flower



A native Australian shrub found in Central Victoria. It is a low growing ground shrub that reaches a height of up to 20 cm. It prefers bushlands and woodlands and drier forests. There are around 150 species of *Hibbertia* that occur in Australia. Flowers have a width to 15 mm and appear from August to November.

Family: Dilleniaceae  
Genus: *Hibbertia*  
Species: *Hibbertia humifusa ssp. erigens*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Hoary guinea-flower



One of numerous shrubs in the *Hibbertia* genus. It is a small native shrub endemic to south eastern Australia from Queensland to Victoria and Tasmania. It grows to 60 cm with a flower width to 50 mm which appears from September to December. It prefers rocky soils in bushland, woodlands and forests. It is often found near red box and red stringybark trees.

Family: Dilleniaceae  
Genus: *Hibbertia*  
Species: *Hibbertia obtusifolia*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Silky guinea-flower




One of numerous shrubs in the *Hibbertia* genus. It is a medium sized native shrub endemic to south eastern Australia from Queensland to South Australia and Tasmania. It grows to 1.5 m with a flower width of 40 mm, which appears from autumn to winter. It prefers rocky hillsides and shallow soils on ridges in bushland, woodlands and open forests.

Family: Dilleniaceae  
Genus: *Hibbertia*  
Species: *Hibbertia sericea*



# LILY



VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED

## Bulbine lily



Also known as the golden lily or yellow onion Weed, it is found in eastern Australia from Qld to SA and Tasmania. It grows to 75 cm with flower width to 25 mm. It prefers dryer soils in bushland, woodlands and open forests and flowers from late August to November with early Nancy and hardenbergia. Tubers were eaten raw or roasted by Aboriginals.

Family: *Asphodelaceae*  
Genus: *Bulbine*  
Species: *Bulbine bulbosa*



VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED

## Yellow-rush lily



Similar in colour to the bulbine lily, it is a native perennial lily found in all states of Australia and Tasmania. It grows to a height 75 cm and yellow flowers appear in September to November with a flower width to 13 mm. It prefers bushland, woodlands and drier forests and is found on sandy loams. It has six stamen and the tepals twist after flowering.

Family: *Asphodelaceae*  
Genus: *Tricoryne*  
Species: *Tricoryne elatior*



# ORCHID



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Leopard orchid



A native perennial orchid found in south eastern Australia from New South Wales to Victoria and Tasmania. It grows to a height 50 cm with two or three grass-like leaves and up to ten flowers each with a flower width to 25 mm. It prefers well drained soils in bushland, woodlands and forests.

Family: Orchidaceae  
Genus: *Diuris*  
Species: *Diuris pardina*



SITE GPS

VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Northern golden moths



A native endemic to Victoria and first described in 2006. It grows to a height of 35 cm, and has a flower width to 22 mm. It prefers drier grasslands, bushland, woodlands and forests and flowers from September to November. It is listed in the Victorian *Flora and Fauna Guarantee Act 1988* as endangered—because of farm clearing.

Family: Orchidaceae  
Genus: *Diuris*  
Species: *Diuris protena*



VIC N.S.W. QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED

## Tiger orchid




Also known as the hornet orchid, it is a native perennial tuberous orchid found in eastern Australia from Queensland to South Australia and Tasmania. It grows to a height 50 cm, has a flower width to 35 mm and flowers appear in Spring. It prefers bushland, woodlands and forests and is often found with the leopard orchid.

Family: Orchidaceae  
Genus: *Diuris*  
Species: *Diuris sulphurea*



SITE GPS

VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Wallflower orchid



Also known as the Eastern wallflower orchid, it is a native perennial tuberous orchid similar to the common donkey orchid and found in eastern Australia from Victoria to South Australia. It grows to a height 50 cm, has a flower width to 35 mm and flowers appear in Spring. It prefers bushland, woodlands and forests.

Family: Orchidaceae  
Genus: *Diuris*  
Species: *Diuris orientis*



# P E A



## Bird's-foot trefoil




A perennial also known as eggs and bacon, it is a native of Eurasia and North Africa. It is found all states of Australia (except WA & NT). It grows to 30 cm and prefers grasslands with moist soils. Yellow flowers appear from December to February with a width to 13 mm. It is widely used as livestock feed. It is regarded an invasive species in Australia.

Family: Fabaceae  
Genus: *Lotus*  
Species: *Lotus corniculatus*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Clover tree



Found in south eastern Australia from Queensland to South Australia and Tasmania. It grows to 4 m and prefers drier sclerophyll bushland, woodlands and forests. Yellow flowers appear in racemes from September to November with a width to 16 mm. Leaves are trifoliate.

Family: Fabaceae  
Genus: *Goodia*  
Species: *Goodia lotifolia*

SITE GPS

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Common aotus



Also known as the golden pea, it is a native shrub found in all states of Australia (except the NT and WA). It is a hardy evergreen, grows to 15 m and prefers wetter bushlands, woodlands and forests. It flowers from mid spring to summer and flowers have a width to 25 mm. Leaves vary from egg-shape to long and narrow and may roll under on the edges.

Family: Fabaceae  
Genus: *Aotus*  
Species: *Aotus ericoides*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Creeping bossiaea



Found in eastern Australia from Queensland to South Australia and Tasmania. It is a hardy prostrate creeping evergreen prefers eucalypt bushland, woodlands and forests. Flowers appear from September to November with a width to 22 mm. Leaves are ovate to oblong and grow alternately. They are dark on top and paler below.

Family: Fabaceae  
Genus: *Bossiaea*  
Species: *Bossiaea prostrata*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# P E A



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Gorse bitter pea



A small to medium sized hardy evergreen shrub found in all states of Australia (except the Northern Territory). It grows to 2 m and prefers poorer soils in dry bushlands, woodlands and forests. It has a yellow flower which appears from late winter to spring and has a width of up to 7 mm. Its small oval leaves terminate in a prickly point.

Family: Fabaceae  
Genus: *Daviesia*  
Species: *D. ulicifolia*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Grey parrot-pea




A hardy native pea bush found in all states of Australia (except QLD and the NT). It grows to 1.5 m has a yellow-orange flower with a width to 11 mm. It prefers drier woodlands and bushland and flowers from September to December and is drought, snow and frost tolerant. Phyllodes grow to 20 x 0.5 mm and are dark green.

Family: Fabaceae  
Genus: *Dillwynia*  
Species: *Dillwynia cinerascens*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Handsome-flat pea



A native pea bush found in eastern Australia from QLD to Victoria and Tasmania. It grows to 2 m and has a yellow-orange flower that appears from September to December with a width to 20 mm. It prefers woodlands, bushland and forests and is drought, snow and frost tolerant. Leaves are dark green on top, light green underneath and to 5 cm long.

Family: Fabaceae  
Genus: *Platylobium*  
Species: *Platylobium formosum*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Hop bitter pea



A common shrub endemic to south eastern Australia from Queensland to SA and Tasmania. It is a hardy evergreen that grows to 3 m and prefers drier rocky sclerophyll bushlands, woodlands and forests. It has clusters of flowers which appear in late spring with a width to 15 mm.

Family: Fabaceae  
Genus: *Daviesia*  
Species: *Daviesia latifolia*



# P E A



## Large-leaf bush-pea



Also known as the large-leaf bitter-pea, it is an erect shrub found in south eastern Australia from QLD to SA and Tasmania. It grows to 3 m and prefers bushland and woodlands. Yellow flowers appear in clusters of 5–11 from Sep to Nov with a width to 15 mm. Leaves are wedge to egg shaped with a point and egg-shaped pods are 5–7 mm long.

Family: Fabaceae  
Genus: *Pultenaea*  
Species: *Pultenaea daphnoides*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Mountain mirbelia



An erect spreading shrub also known as sandstone bushpea found in eastern Australia from NSW to Victoria and Tasmania. It prefers well drained soils in drier rocky areas in woodlands and bushland. It flowers in spring with a flower width to 13 mm and small hairy reddish pods. It was first described by botanist Ferdinand von Mueller.

Family: Fabaceae  
Genus: *Mirbelia*  
Species: *Mirbelia oxylobioides*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Narrow-leaf bitter pea



Found in south eastern Australia from NSW to SA. It is a broom-like shrub with multiple stems which grows to 3 m and prefers drier sclerophyll bushland, woodlands and forests. Yellow and maroon flowers appear on racemes with 5–10 flowers from Aug to Dec with a width to 12 mm. Yellowish-green phyllodes grow to 10 cm long by 6 mm wide.

Family: Fabaceae  
Genus: *Daviesia*  
Species: *Daviesia leptophylla*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Showy parrot pea



Found in south eastern Australia from Queensland to South Australia and Tasmania. It is a hardy evergreen, grows to 1 m and prefers drier sclerophyll bushland, woodlands and forests. It flowers in Spring with a flower width to 15 mm. Young flowers may show different colours to older leaves.



Family: Fabaceae  
Genus: *Dillwynia*  
Species: *D. sericea*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# P E A



## Small-leaf parrot-pea



Found in south eastern Australia from Queensland to Victoria. It is a hardy evergreen, grows to 1.5 m and prefers drier bushlands, woodlands and forests. It flowers in late winter to spring, followed by seed pods up to 5 mm long with smooth seeds.

Family: Fabaceae  
Genus: *Dillwynia*  
Species: *Dillwynia phyllicoides*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Tall bitter pea (Alpine)



A relatively rare native shrub found only in Victoria. It grows to 2 m and prefers rocky sub-alpine or protected alpine eucalypt forests with rainfall exceeding 1000 mm annually. Small yellow flowers appear in abundance in the summer months and have a flower width to 7 mm. Leaves grow to 25 cm.

Family: Fabaceae  
Genus: *Daviesia*  
Species: *Daviesia laxiflora*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Biddie-widgee



Also known as biddy-biddy, bidi-bidi or piri-piri-bur, it is found in all states of Australia (except the Northern Territory). It grows to a height of 40 cm and a head width of 30 mm. A bur is left once the flower falls, which are known for sticking to socks when walking through a patch where they are growing. It flowers from September to March.

Family: Rosaceae  
Genus: *Acaena*  
Species: *Acaena novae-zelandiae*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Cranberry heath




Also known as native cranberry, it is found in south eastern Australia from NSW to SA and Tasmania. It grows to 60 cm and prefers drier bushland and woodlands. Its red tubular flowers grow to 2 cm long and appear from February to June. Green berries appear after pollination, which later turn red and are then edible and juicy.

Family: Ericaceae  
Genus: *Astroloma*  
Species: *Astroloma humifusum*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Lake Mountain Grevillea



A relatively rare native shrub endemic to the eastern Victorian ranges and found on Lake Mountain. It grows to 2 m and prefers rocky subalpine snowgum forests. It has a red flower that appears in early to mid summer. It was previously known as the Lake Mountain form of *Grevillea victoriae*.

Family: Proteaceae  
Genus: *Grevillea*  
Species: *Grevillea monslacana*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Mountain grevillea



Also known as cat's claw and alpine grevillea, it is a native shrub found in eastern Australia from NSW to Victoria. It grows to 2 m and prefers drier bushland, woodlands and forests. The curled flower varies in colour, but often appears in a bright red and yellow form. It is not restricted to alpine areas, often preferring lower areas.

Family: Proteaceae  
Genus: *Grevillea*  
Species: *Grevillea alpina*



# WILDFLOWER



## Red bartsia



Also known as the sticky bartsia and red tarweed, it is an erect perennial herb introduced to Australia from Europe and is considered a weed. It is found in all states or Australia and grows to 30 cm. It has a tubular flower and prefers moist soils. It flowers from spring to early summer.

Family: *Orobanchaceae*  
Genus: *Parentucellia*  
Species: *Parentucellia latifolia*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Scarlet pimpernel



Also known as red chickweed and poor man's barometer, it is found in all states of Australia (except the NT). It grows to 10 cm, has an orange, red or blue flower, with a width to 6 mm. It prefers bushland, woodlands and forests. It flowers from September to April and is often considered a weed. Flowers close when barometric pressure drops sharply.

Family: *Primulaceae*  
Genus: *Anagallis*  
Species: *Anagallis arvensis*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Running postman



Also known as scarlet runner, it is an Australian native found in all states of Australia. It grows close to the ground and spreads for up to 1.5 m width. It prefers drier soils and even sandy areas found in bushland, woodlands, forests and coastal areas.

Family: *Fabaceae*  
Genus: *Kennedia*  
Species: *Kennedia prostrata*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



## Austral bugle



A flowering shrub that grows from a loose leaf rosette and is found in all states of Australia (except the NT). It grows to 40 cm, flowers from September to November and flowers have a width to 20 mm. It prefers bushland, woodlands and forests. The shrubs only last for 2–3 years each.

Family: Lamiaceae (mint family)  
Genus: *Ajuga*  
Species: *Ajuga australis*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Bindweed



Also known as great bindweed, giant bindweed and hedge bindweed, it is native to Europe and has been naturalised into south eastern Australia from NSW to South Australia and Tasmania. It grows to 2 m and its pink or white flowers have a width to 9 cm. It prefers moist soils such as those in lower lying areas and is often seen in parks and gardens.

Family: Convolvulaceae  
Genus: *Calystegia*  
Species: *Calystegia silvatica*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Blue pin cushion



A common perennial herb found in all states of Australia. It grows to a height 50 cm and has a flower width of 20 mm. It prefers bushland, woodlands and open forests. It flowers from September to January. It was first collected during the 1801-02 voyage of HMS Investigator under Matthew Flinders.

Family: Goodeniaceae  
Genus: *Brunonia*  
Species: *Brunonia australis*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Blue stars



Also known as blue squill and mudrurt, it is a tuberous perennial herb found in southern Australia from New South Wales to South Australia, Western Australia and Tasmania. It grows to 15 cm with grass-like basal leaves and a flower width to 20 mm. Flowers appear in spring. It prefers bushland, woodlands and forests and has edible tubers.

Family: Asphodelaceae  
Genus: *Chamaescilla*  
Species: *Chamaescilla corymbosa*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Brazilian vervain



Also known as Brazilian verbena, it is native to South America and found in eastern Australia from QLD to Victoria. It grows to 1 m and has a purple flowerhead which is hairy and has a width to 12 cm. It prefers bushland and woodlands and flowers from April to June. The stem is squarish and usually glabrous (not hairy) below the flowerhead.

Family: Verbenaceae  
Genus: *Verbena*  
Species: *Verbena brasiliensis*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Chicory



A woody perennial also known as blue daisy and coffeeweed. It is native to the Old World and introduced to all of Australia (except the NT). It prefers roadsides in drier areas. It grows to 1.5 m and blue to lavender flowers with a width to 3.5 cm appear from December to March. The entire plant is edible. Ground roots are a coffee substitute.

Family: Gentianaceae  
Genus: *Centaurium*  
Species: *Centaurium erythraea*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common centaury




Also known as European Centaury, it is native to Europe and a weed introduced to Australia, where it is a weed found in all states of Australia (except the NT). It prefers moist soils in forests and bushland and woodlands. It grows to 60 cm, flower has five petals (sometimes six) which appear from October to April with a width to 13 mm.

Family: Gentianaceae  
Genus: *Centaurium*  
Species: *Centaurium erythraea*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common fumitory



Also known as earth smoke and drug fumitory, it is an erect introduced member of the poppy family found in eastern Australia from Queensland to South Australia and Tasmania. It grows to 50 cm but as a climber it will climb to 1.5 m. Pink flowers appear from September to December with a length to 10 mm. It prefers damper environments.

Family: Papaveraceae  
Genus: *Fumaria*  
Species: *Fumaria officinalis*



# WILDFLOWER



## Common heath



A shrub native to south eastern Australia and is found from NSW to SA and Tasmania. It grows to 1 m and prefers bushland, woodlands, forests and alpine areas up to 1200 m. It flowers from April to October, attracts honeyeaters and is found in white, pink, scarlet, crimson and double colours. It's the State Emblem of Victoria.

Family: Ericaceae  
Genus: *Epacris*  
Species: *Epacris impressa*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Common selfheal



Also known as blue curls, woundwort and heal-all, it is found in all states of Australia (except the NT). It grows to 80 cm, though normally to 30 cm and has a composite club-like flower head with a width to 30 mm. It prefers bushland, woodlands and forests and flowers from Nov-Mar, leaves and flowers are edible. It is an important herbal medicine.

Family: Lamiaceae (mint family)  
Genus: *Prunella*  
Species: *Prunella vulgaris*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Common stalk's bill



A herbaceous hairy sticky annual native to the Mediterranean Basin and introduced to Australia where it is found in all States. It grows to 40 cm and has a flower width of 13 mm. It prefers grasslands. The entire plant is edible, with leaves having a parsley flavour. The seeds can twist as they dry and the plant is a noxious weed.

Family: Geraniaceae  
Genus: *Erodium*  
Species: *Erodium cicutarium*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Fairy aprons



A perennial bladderwort with numerous underground trailing stems with bladders found in all states of Australia (except the NT). It grows to 50 cm, has a flower width of 10 mm and prefers wetter shady areas around rocks. It flowers from August to April.

Family: Lentibulariaceae  
Genus: *Utricularia*  
Species: *Utricularia dichotoma*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



## Finger flower



A common flowering small upright shrub found in eastern Australia from Queensland to South Australia. It grows to 50 cm and prefers drier rocky or sandy soils in bushland and woodlands. Flowers appear from October to January with a width to 40 mm. Pedals may also appear wider and even egg shaped.

Family: Pittosporaceae  
Genus: *Cheiranthra*  
Species: *Cheiranthra linearis*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Grass triggerplant




A perennial herb endemic to Australia and found in all states of Australia. It grows to 100 cm and prefers areas of drier soil found in bushland, woodlands and sclerophyll forests. It is known for its pollination trigger mechanism used to spread pollen by striking insects. Pink to whitish flowers with a width to 17 mm appear from September to January.

Family: Stylidiaceae  
Genus: *Stylidium*  
Species: *Stylidium graminifolium*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Hairy pink



A meadow wildflower native to Europe and introduced into Australia where it is now found in all states (except the NT). It grows to 60 cm and develops clusters of flowers with a flower width to 10 mm from September to December. It prefers bushland, woodlands, open fields and grasslands. It is considered a weed in Australia.

Family: Caryophyllaceae  
Genus: *Petrorhagia*  
Species: *Petrorhagia dubia*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Love creeper



A slender climber found in all states of Australia (except the NT). It climbs to 2 m, has a flower width of 15 mm and prefers bushland, woodlands and forests. Pale blue to purple flowers appear in sprays from July to December. Leaves are to 50 mm long and 5 mm wide.

Family: Polygalaceae  
Genus: *Comesperma*  
Species: *Comesperma volubile*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Many-stemmed bluebell



Similar to the tall bluebell, it is a perennial plant found in all states of Australia (except the NT and Queensland). It grows to 70 cm and prefers bushland, woodlands and dry open forests. Flowers appear from September to February with a width to 37 mm though normally to 28 mm. It has multiple flower stems per plant.

Family: Campanulaceae  
Genus: *Wahlenbergia*  
Species: *Wahlenbergia multicaulis*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Native flax



Also known as Australian flax, it is a short-lived perennial flowering herb that is native to Australia and found in all states of Australia (except the NT). It grows to 1.2 m, flowers in mid Spring, has a flower width to 20 mm and prefers eucalypt bushland, woodlands and forests. Aboriginals used the stem for strong fibre and ate the small seed.

Family: Linaceae  
Genus: *Linum*  
Species: *Linum marginale*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Native raspberry



A native also known as the mountain raspberry. It is found in eastern Australia from Queensland to South Australia and Tasmania. The thorny bush grows to a height of 2 m and looks like a small version of the blackberry bush. It prefers higher altitudes. It has edible berries when ripe.

Family: Rosaceae  
Genus: *Rubus*  
Species: *Rubus parvifolius*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Onion grass



Also known as guildford grass, it is a herbaceous perennial naturalised in all states of Australia (except the NT). It grows to 20 cm and prefers bushland, woodlands, and open forests. It has tough thin grass-like leaves and a pink flower from August to November with a width to 32 mm. The corm is edible as an onion replacement and is eaten by cockatoos.

Family: Iridaceae  
Genus: *Romulea*  
Species: *Romulea rosea*



# WILDFLOWER



## Pennyroyal



A creeping or erect member of the mint family also known as mosquito plant and pudding grass, it is an introduced found in all states of Australia (except the NT). It prefers bushland and woodlands, grows to 40 cm with a flower width to 17 mm and flowers from January to March. It is poisonous and toxic to the liver.

Family: Lamiaceae (mint family)  
Genus: *Mentha*  
Species: *Mentha pulegium*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Pink bells



An upright climbing shrub found in all states of Australia (except the Northern Territory). It grows to 1 m and prefers moist to moderately dry soils and semi-shaded areas in bushland, woodlands and forests. Flowers appear from August to January width to 28 mm when fully open. It was first described by botanist John Lindley in 1838.

Family: Elaeocarpaceae  
Genus: *Tetradlea*  
Species: *Tetradlea ciliata*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Purple loosestrife




A flowering plant native to south eastern Australia as well as parts of Europe, Asia and Africa. It grows to 2 m, has purple flowers in summer to autumn and prefers wetter areas beside creeks and rivers or beside bodies of water. It has edible leaves, flowers and tubers, though they should all be eaten in small quantities.

Family: Lythraceae  
Genus: *Lythrum*  
Species: *L. salicaria*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Purple woodsorrel



A purple ground cover introduced from Africa and found in all states of Australia (except the NT). It grows to 30 cm and prefers dry to moderately moist soils in bushland, woodlands and forests. It has purple, pink, mauve to white flowers ranging in width from 10 mm to 80 mm from August to November. It is considered a weed in Australia.

Family: Oxalidaceae  
Genus: *Oxalis*  
Species: *Oxalis purpurea*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



## Rock isotome



A small herbaceous perennial plant also known as the showy isotome, it is found in south eastern Australia from Queensland to Victoria. It grows to 50 cm and has a flower width to 20 mm. It prefers rocky areas in bushland, woodlands and drier forests. It flowers from Spring to late Summer, can be highly toxic, and should not be touched.

Family: Campanulaceae  
Genus: *Isotoma*  
Species: *I. axillaris*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Round-leaved cranes-bill



Also known as the roundleaf geranium, it is an annual herb found in Victoria and Tasmania. It grows prostrate to 30 cm and flowers appear from August to December with a width to 13 mm. It prefers bushland and woodlands. Leaves are round-shaped and may turn yellow to orange-reddish when old.

Family: Geraniaceae  
Genus: *Geranium*  
Species: *Geranium rotundifolium*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Salvation Jane



Also known Patterson's curse, it is an introduced invasive poisonous annual found in all states of Australia. It grows to 1 m and prefers bushland, farmlands and roadsides. Flowers appear from October to January with a width to 20 mm. It can be eaten by cattle and sheep but not horses or animals without ruminant digestive systems. It irritates human skin.

Family: Boraginaceae  
Genus: *Echium*  
Species: *Echium plantagineum*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Slender speedwell



A native perennial groundcover herb found in south eastern Australia from NSW to SA and Tasmania. It prefers bushland and forests. Pinkish to mauve flower appears from October to December with a width to 26 mm. Stems grow to 70 cm. It spreads by rhizomes. Narrow lance shaped opposed leaves grow to 3 cm.

Family: Plantaginaceae  
Genus: *Veronica*  
Species: *Veronica gracilis*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Soft cranes-bill

Also known as cinquefoil geranium and native carrot, it is a native found in eastern Australia from Queensland to SA and Tasmania. It grows to 50 cm, flowers from August to February with a width to 15 mm and prefers bushland, woodlands and forests. Leaves can be kidney-shaped and may turn yellow to orange-reddish. Its corm is edible.



Family: Geraniaceae  
Genus: *Geranium*  
Species: *Geranium potentilloides*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Tall bluebell

Also known as the Australian bluebell or Austral bluebell, it is a perennial plant found in all states of Australia (except the Northern Territory). It grows to 70 cm and prefers bushland, woodlands and dry open forests. It flowers in spring and early summer and flowers have a width to 37 mm though normally to 28 mm.



Family: Campanulaceae  
Genus: *Wahlenbergia*  
Species: *Wahlenbergia stricta*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Tall sundew

Found in all states of Australia. It is a tall growing perennial carnivorous plant that grows to 50 cm. It prefers bushland, woodlands and drier forests and has pink to white flowers with a width to 18 mm from August to November. To survive in poor soils it traps insects and digests their nutrients and minerals instead of relying overly on the soil.



Family: Droseraceae  
Genus: *Drosera*  
Species: *Drosera peltata* ssp: *auriculata*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## True forget-me-not

An introduced species native to Europe and often called scorpion grass in the Northern Hemisphere, it is naturalised in all states of Australia and is considered a weed. It grows to 45 cm with a flower width of 11 mm. It prefers moist soils in forests. The similar species *Myosotis alpestris* is the State Flower of Alaska and Datsland, Sweden.



Family: Boraginaceae  
Genus: *Myosotis*  
Species: *Myosotis scorpioides*



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED

## Water speedwell



Also known as blue-water speedwell, brook pimpernel and sessile water-speedwell, it is found in eastern Australia from Queensland to Victoria and Tasmania. It grows to 30 cm high and prefers forests where it grows best in moist soils. A pinkish to mauve flower appears in Spring with a width to 12 mm. It's rarely found in forest environments.

Family: Plantaginaceae  
Genus: *Veronica*  
Species: *Veronica anagallis-aquatica*



VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED

## Wavy lobelia



First described in 1770 by Joseph Banks on Captain Cooks first voyage of discover, it is found in south eastern Australia from Queensland to Victoria. It grows to 40 cm, has a flower width of 20 mm and prefers eucalypt bushland, woodlands and forests. It flowers from November to March.

Family: Campanulaceae  
Genus: *Lobelia*  
Species: *Lobelia dentata*



# DAISY



## Blue bottle daisy



Also known as common lagenophora, it is a small plant found in eastern Australia from QLD to SA and Tasmania. It grows to 25 cm, has a pink, mauve, purplish or white flower with a width to 14 mm that appear from October to February and prefers shady areas in bushland, forests and woodlands.

Family: Asteraceae  
Genus: *Lagenophora*  
Species: *Lagenifera stipitata*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Purple burr daisy



Also known as bindi-eye, it is a purple prostrate native daisy found in all states of Australia (except WA and Tasmania). It grows to 60 cm, has a flower width to 50 mm and prefers eucalypt bushland and woodlands and at elevations to 1600 m. It flowers from October to January.

Family: Asteraceae  
Genus: *Calotis*  
Species: *Calotis cuneifolia*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# LILY



VIC N.S.W. QLD S.A. W.A. N.T. TAS


NATIVE WEED INTRODUCED NATURALISED

## Black-anther lily



A native found in south eastern Australia from New South Wales to South Australia and Tasmania. It grows to 80 cm and prefers bushland, woodlands, drier forests and coastal areas. It flowers in early Spring to mid summer and its berries are considered as being edible.

Family: Xanthorrhoeaceae  
Genus: *Dianella*  
Species: *Dianella admixta*



VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED

## Blue flax-lily



Also known as the blueberry lily, it is an Australian native perennial herb found in south eastern Australia from Queensland to Victoria and Tasmania. It grows to 70 cm, flowers from October to January and has a flower width of up to 15 mm. It has medium sized violet berries which grow up to 20 mm are not edible.

Family: Asphodelaceae  
Genus: *Dianella*  
Species: *Dianella caerulea*



VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED

## Chocolate lily



A native found in eastern Australia from New South Wales to South Australia and Tasmania. It grows to 1 m and prefers bushland, woodlands, drier forests and coastal areas. It flowers from September to November with a width to 30 mm and has a chocolate scent when crushed. The edible tubers were eaten by early Aborigines.

Family: Asparagaceae  
Genus: *Dichopogon*  
Species: *D. strictus*



VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED

## Common fringe-lily



A perennial herb endemic to Australia and found in all states of Australia. It grows to 60 mm in height and has a flower width to 35 mm. It prefers grasslands, bushland, woodlands and forests. It flowers from September to April and each flower only lives for one day.

Family: Asparagaceae  
Genus: *Thysanotus*  
Species: *Thysanotus tuberosus*



# LILY



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Nodding blue lily



An Australian native also known as the blind lily, found in all states of Australia (except Western Australia and Northern Territory). It grows to 1.5 m and prefers bushland, woodlands and forests. It has a flower width to 30 mm which appears in Spring. It is a favourite hardy native plant grown in urban parks and gardens.

Family: Asphodelaceae  
Genus: *Stypandra*  
Species: *Stypandra glauca*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Nodding chocolate lily




A native found in all states of Australia (except the Northern Territory and Tasmania). It grows to 30 cm and prefers bushland, woodlands, drier forests and coastal areas. It flowers in early Spring with a flower width to 17 mm. The edible juicy tubers were eaten by early Aboriginals. Recently reclassified, it is similar to the chocolate lily.

Family: Asparagaceae  
Genus: *Arthropodium*  
Species: *Arthropodium fimbriatum*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Pale vanilla lily



A pale pinkish herbaceous perennial Australian native that found in all states of Australia (except Western Australia and Northern Territory). It grows to 1 m and prefers bushland, woodlands, forests and Alpine areas. It grows best in moderately moist soils, has a strong vanilla fragrance on warmer days and has an edible tuber.

Family: Asparagaceae  
Genus: *Arthropodium*  
Species: *Arthropodium milleflorum*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Small vanilla lily




An Australian native found in south eastern Australia from Queensland to SA and Tasmania. It has grass-like leaves and grows to 35 cm. It prefers bushland, woodlands and forests. It has a flower width to 12 mm which appears from November to December. It has edible roots.

Family: Asparagaceae  
Genus: *Arthropodium*  
Species: *Arthropodium minus*



# LILY



## Tasman flax-lily



Also known as the Tasmanian Flax-lily, it is an Australian native perennial herb found in south eastern Australia from New South Wales to Victoria and Tasmania. It grows to 1.5 m, flowers in late spring and summer and has a flower width of up to 15 mm. It has medium sized violet berries which grow up to 20 mm are not edible.

Family: *Asphodelaceae*  
Genus: *Dianella*  
Species: *Dianella tasmanica*

VIC

N.S.W

QLD

S.A.

W.A.

N.T.

TAS

NATIVE

WEED

INTRODUCED

NATURALISED



# PEA



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Austral indigo



Found in all states of Australia. It is a member of the pea family that flowers from July to September. It grows to 1.3 m. It prefers semi shade in bushland, woodlands and forests. Aborigines used the flowers to create purple dye and they would stun fish with finely crushed roots put into water.

Family: Fabaceae  
Genus: *Indigofera*  
Species: *Indigofera australis*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common vetch



A member of the pea family also known as vetch, garden vetch and tare and introduced from the Northern Hemisphere to all states of Australia. It is a tall growing plant that grows to 50 cm. It prefers drier bushland and woodlands and flowers from August to December with a width to 23 mm. Seedpods appear soon after with a length to 7 cm.

Family: Fabaceae  
Genus: *Vicia*  
Species: *Vicia sativa*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Hardenbergia



An evergreen climber in the pea family also known as sarsaparilla, happy wanderer and purple coral pea, it is found in all states of Australia. It grows to 6 m and prefers most habitats except extreme climates. It flowers early from July to December with bulbine lily and early Nancy and is a good garden plant. Seeds can be dispersed by ants.

Family: Fabaceae  
Genus: *Hardenbergia*  
Species: *Hardenbergia violacea*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Hovea lanceolata



A native shrub from the pea family that it is found in eastern Australia from Queensland to Victoria. It grows to 2 m and prefers rocky well drained areas and is often found in ridges. Leaves are lanceolate in shape and can be up to 7 cm in length. It flowers in late winter to early spring with flowers to 12 mm in length.

Family: Fabaceae  
Genus: *Hovea*  
Species: *Hovea lanceolata*



# PEA



## Smooth darling pea




A small perennial native bush in the found in eastern Australia from Queensland to Victoria. It grows to 1 m and prefers bushland, woodlands and drier forests. It produces pink to purple pea-like flowers with a width to 20 mm. Flower spikes can grow to 15 cm and appear from late winter to spring.

Family: Fabaceae  
Genus: *Swainsona*  
Species: *Swainsona galegifolia*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Twining glycine



Also known as the love creeper and glycine pea, it is an Australian native found in south eastern Australia. It is a crawler that grows along the ground and has a flower width of 8 mm. It prefers bushland and forests, often found in shaded subalpine altitudes. It flowers all year, but most commonly in late winter and spring.

Family: Fabaceae  
Genus: *Glycine*  
Species: *Glycine clandestina*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# ORCHID



## Blue fingers orchid



Also known as the eastern tiny blue China orchid, it is found in southern Australia from New South Wales to Victoria and Tasmania. It grows to 70 cm and has a flower width to 30 mm. It prefers well drained soil in bushland, woodlands and forests.

Family: Orchidaceae  
Genus: *Caladenia*  
Species: *Caladenia caerulea*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Blotched hyacinth orchid




A leafless orchid found in south eastern Australia from Queensland to South Australia, it grows to a height 100 cm and a flower width to 25 mm. It prefers shaded areas in forests, bushland and woodlands. For most of the year they remain below ground, flowering from November to March.

Family: Orchidaceae  
Genus: *Dipodium*  
Species: *Dipodium punctatum*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Common bird orchid




Also known as the large bird orchid, it is found in south eastern Australia from NSW to SA and Tasmania. It has large dark green leaves to 50 cm long and prefers shaded areas in woodlands, bushland and forests and grows to 8 cm. Flowers appear from September to February with a width to 4 cm and 6–10 calli (fleshy column-like lumps).

Family: Orchidaceae  
Genus: *Chiloglottis*  
Species: *Chiloglottis valida*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Mountain sun orchid



A herbaceous orchid native to Australia, they are endemic to south-eastern Australia and found from New South Wales to Victoria. It grows to 32 cm and prefers grassy areas of alpine, subalpine and forest environments. It has a purplish flower which forms from September to January


Family: Orchidaceae  
Genus: *Thelymitra*  
Species: *Thelymitra alpina*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# ORCHID



SITE GPS  
SITE GPS

VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Plain-lip spider orchid



A native ground orchid with a single leaf that is found in eastern Australia from NSW to SA. It grows to 35 cm in small colonies and prefers bushland, woodlands and forests where it grows best in dry to moist soils. It flowers from mid September to October and has an edible tuber that was eaten by early Aboriginals.

Family: Orchidaceae  
Genus: *Caladenia*  
Species: *Caladenia clavigera*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Purple beard orchid



An Australian native found in all states of Australia (except the NT). It grows to 45 cm and prefers bushland and woodlands. It has a single dark green leaf which is developed by flowering time, a small tuber and up to nine flowers with reddish to purplish stripes. Flowers appear from August to November with a width to 22 mm and length to 27 mm.

Family: Orchidaceae  
Genus: *Calochilus*  
Species: *Calochilus robertsonii*



VIC N.S.W QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED

## Salmon sun orchid



A native found in south eastern Australia from NSW to SA and Tasmania. It grows to 30 cm and prefers bushland and woodlands in dry to moderately moist soils. Flower stems have up to five pale red to salmon flowers which open on sunny days and appear from September to November with a width to 15 mm. *Rubra* is Latin for 'red'.

Family: Orchidaceae  
Genus: *Thelymitra*  
Species: *Thelymitra rubra*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Slender sun orchid



Native to Australia and New Zealand, it is found in all states of Australia (except WA and NT). It grows to 40 cm and prefers bushland, woodlands and forests where it grows best in dry to moderately moist soils. It is one of the most widespread of all native orchids. Flowers appears from August to January with a width to 20 mm.

Family: Orchidaceae  
Genus: *Thelymitra*  
Species: *Thelymitra pauciflora*



# ORCHID



## Spotted sun orchid



A purple to mauve native found in all states of Australia (except WA and the NT). It grows to 40 cm with up to 10 flowers which appear September to December with a width to 12 mm. Flowers can differ in colour greatly. It prefers bushland and woodlands in dry to moderately moist soils. It has small spots.

Family: Orchidaceae  
Genus: *Thelymitra*  
Species: *Thelymitra ixioides*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Summer fingers



A pinkish native herb endemic to southern Australia from Victoria to South Australia and Tasmania. It has a single stem which grows to 27 cm high. It prefers bushland, woodlands and forests where it grows best in dry to moderately moist soils. It has an edible tuber and was a source of food for early Aboriginals.

Family: Orchidaceae  
Genus: *Caladenia*  
Species: *Caladenia vulgaris*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Waxlip orchid



A purple native found in all states of Australia (except Western Australia and Northern Territory) and is most widespread in NSW. It grows to 40 cm and prefers bushland, woodlands and forests where it grows best in dry to moderately moist soils. It has an edible tuber and was a source of food for early Aboriginals.

Family: Orchidaceae  
Genus: *Caladenia*  
Species: *Caladenia major*

VIC N.S.W QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# VIOLET



## Common dog violet


A violet native to Eurasia and Africa and introduced to eastern Australia from New South Wales to South Australia and Tasmania. It grows to 15 cm, has a flower width of 40 mm and prefers damp shaded areas in bushland, woodlands and forests. Flowers appear from November to December. Dark green leaves are heart shaped.



Family: *Violaceae*  
 Genus: *Viola*  
 Species: *Viola riviniana*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Ivy-leaf violet

A native violet found in south eastern Australia from NSW to SA and Tasmania. It grows to 12 cm and prefers bushland, woodlands, and forests. Flowers to 10 mm wide appear from July to December. It's likely the species explorer Sir Thomas Mitchell witnessed beside a pond in 1836 at Violet Town and named *Violet Ponds*.



Family: *Violaceae*  
 Genus: *Viola*  
 Species: *Viola hederacea*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Native violet

A native violet found in eastern Australia from New South Wales to South Australia and Tasmania. It grows to 20 cm, has a flower width of 18 mm and prefers damp shaded areas in bushland, woodlands and forests. Flowers appear from later Winter to Spring. Unlike *v. hederacea*, it is often grown in parks and gardens.



Family: *Violaceae*  
 Genus: *Viola*  
 Species: *Viola banksii*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Sweet Violet

A violet also known as wood violet, English violet and garden violet. It is native to Eurasia and Africa and introduced to all states of Australia (except the NT). It grows to 15 cm, has a flower width of 40 mm and prefers edges of bushland, forests and gardens. Flowers appear from July to September. Leaves are reported as edible.



Family: *Violaceae*  
 Genus: *Viola*  
 Species: *Viola odorata*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



## Mountain correa



*Var. latrobeana* is tree endemic to NSW and Victoria. It grows to 3 m and prefers partially shaded areas in dense scrub in bushland and forests. It is frost and snow tolerant, has egg shaped leaves and green to reddish or mauve flowers from late Autumn to Winter. Flowers grow to 50 mm long and attract honeyeaters.

Family: Rutaceae  
Genus: *Correa*  
Species: *Correa lawrenceana*  
Var. *latrobeana*

VIC

N.S.W.

QLD

S.A.

W.A.

N.T.

TAS

NATIVE

WEED

INTRODUCED

NATURALISED



# ORCHIDS



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common onion orchid



Found in all states of Australia (except the NT). It grows a flowering stem to 60 cm and prefers bushland, woodlands and forests. It flowers from October to January. The flowering section can be up to 13 cm long and can contain between 10 and 100 yellowish-green flowers with a length to 9 mm and width to 4 mm.

Family: Orchidaceae  
Genus: *Microtis*  
Species: *Microtis unifolia*



VIC N.S.W QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED

## Greenhood orchid




Found in all states of Australia (except the NT). It grows to 40 cm in clumps and flowers from late winter to early spring. It prefers bushland, woodlands and forests. The hooded flowers are pollinated by insects attracted to the orchid by nectar excreted from glands at the base of the flower.

Family: Orchidaceae  
Genus: *Pterostylis*  
Species: *Pterostylis concinna*



VIC N.S.W QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED

## Maroonhood



A terrestrial perennial herbaceous native found in NSW and QLD but recently identified at Marysville, Victoria. It grows to 35 cm and prefers moist shady areas in forests amongst leaf litter. Flowers appear from July to November, have a maroon colour over the top of the hood and to 25 mm wide.

Family: Orchidaceae  
Genus: *Pterostylis*  
Species: *Pterostylis pedunculata*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Nodding greenhood




Also known as the parrot's beak orchid, it is a native found in south eastern Australia from Queensland to South Australia and Tasmania. It grows to 30 cm with a single green hooded flower appearing between July and October. The flower droops which gives rise to its name.

Family: Orchidaceae  
Genus: *Pterostylis*  
Species: *Pterostylis nutans*



# ORCHIDS



SITE GPS

SITE GPS

VIC

N.S.W

QLD

S.A.

W.A.

N.T.

TAS

NATIVE

WEED

INTRODUCED

NATURALISED

## Small flowering leafy greenhood



A terrestrial perennial herbaceous native endemic to Victoria. It grows to 25 cm and prefers bushland and woodlands and semi dry soils. It flowers from August to October and flowers are pollinated by insects attracted to the orchid by nectar excreted from glands at the flower's base.

Family: Orchidaceae

Genus: *Pterostylis*

Species: *Pterostylis diminuta*



# WILDFLOWER



SITE GPS

VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Alpine westringia



A small shrub endemic to Victoria. It grows to 2 m and prefers bushland and forests. White flowers appear from September to November with a width to 8 mm and flower heads to 35 mm. It has narrow green leaves that are glossy on top and pale and hairy underside. It was first described by Ferdinand von Mueller in 1855.

Family: Lamiaceae (mint family)  
Genus: *Westringia*  
Species: *Westringia senifolia*



VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Black nightshade



A large-leaved plant also known as blackberry nightshade, it is native to Eurasia and introduced to all states of Australia. It grows to 120 cm and prefers damper areas in woodlands, bushland. White flowers appear from Dec to April with black berries following. It can be confused with deadly nightshade so take extreme caution.

Family: Solanaceae  
Genus: *Solanum*  
Species: *Solanum nigrum*



VIC N.S.W QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED

## Blackberry



A perennial climber found in south eastern Australia from Queensland to Victoria. It was introduced in 1851 by Baron von Mueller and is Australia's most invasive weed, being classified as a *Weed of National Significance* in 1999. It has a thorny vine-like cane and a fruit width of up to 2 cm. The berries are edible when ripe.

Family: Rosaceae  
Genus: *Rubus*  
Species: *Rubus fruticosus*



SITE GPS

VIC N.S.W QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Blue pigroot



Also known as the striped rush-leaf, it is native to the USA and Mexico and introduced to all states of Australia (except the NT). It grows to 20 cm and prefers bushland, woodlands, open forests. It has a distinctive colour band at the centre of the flower and can bloom in modified colours. White flowers appear in October to December with a width to 20 mm.

Family: Iridaceae  
Genus: *Sisyrinchium*  
Species: *Sisyrinchium micranthum*



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Chickweed



Also known as common chickweed, it is a small plant native to Europe and naturalised around the world. It is found growing in shade and often in lawns in all states of Australia. White flowers appear in spring and grow to a width of 8 mm. All above-ground parts of the plant are edible and it is grown as a crop in agriculture.

Family: Caryophyllaceae  
Genus: *Stellaria*  
Species: *Stellaria media*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common beard-heath



A member of a genus containing over 150 species of shrub and small trees. It is a native found in eastern Australia from Queensland to South Australia and Tasmania. It grows to 50 cm and prefers bushland, woodlands and open forests. White flowers appear in Spring with a width to 10 mm.

Family: Ericaceae  
Genus: *Leucopogon*  
Species: *Leucopogon virgatus*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Common rice-flower



A native also known as the drawf riceflower. It is found in eastern Australia from NSW to SA and Tasmania. It grows to 50 cm and prefers bushland, woodlands, open forests. It has a composite flowerhead with a width to 60 cm. The flowerhead has 12–52 individual white flowers with a width to 8 mm. Flowers appear from later winter to spring.

Family: Thymelaeaceae  
Genus: *Pimelea*  
Species: *Pimelea humilis*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Creamy candles



A native perennial herb also known as creamy stackhousia, It is found in all states of Australia (except the Northern Territory). It grows to 80 cm on an erect stem and prefers bushland, woodlands and forests. White to cream or yellowish flowers appear on a spike from August to January and appear a little like a candle.

Family: Celastraceae  
Genus: *Stackhousia*  
Species: *Stackhousia monogyna*



# WILDFLOWER



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Derwent speedwell



A native flowering plant found in south eastern Australia from QLD to SA and Tasmania. It grows to 1.5 m and prefers wetter areas in bushland and forests or beside waterways. Racemes of white, lilac or pale blue flowers appears from December to February with an individual width to 19 mm. Stems remain for up to two years. Leaves are long and serrated.

Family: Plantaginaceae  
Genus: *Veronica*  
Species: *Veronica derwentiana*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Dutch clover



Also known as white clover or ladino, its a herbaceous perennial native to Europe and introduced to all states of Australia. It prefers open grasslands, bushland and lawns and gardens. Edible cream flowers appear from September to February and leaves are also edible and best in salads, steamed or boiled. Once pollinated, flowers bottom drop (left).

Family: Fabaceae  
Genus: *Trifolium*  
Species: *Trifolium repens*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Dwarf boronia



A native ground cover also known as short boronia. It is found in southern Australia from NSW to South Australia and Tasmania. It is a groundcover that grows to a height 30 cm and has a flower width of 23 mm. It prefers woodlands, bushland and forests. White to pinkish-white flowers appear from October to December with a width to 8 mm.

Family: Rutaceae  
Genus: *Cyanothamnus*  
Species: *Cyanothamnus nanus*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Early Nancy



A native perennial herb found in all states of Australia (except the NT). It grows to 30 cm and prefers bushland, woodlands, grasslands and forest areas. Flowers appear August to October (one of the first with Bulbine Lily and hardenbergia) and grow to 21 mm width. Cream flowers are either male or female, with up to eight percent being cosexual.

Family: Colchicaceae  
Genus: *Wurmbea*  
Species: *Wurmbea dioica*



# WILDFLOWER



## Hairy bittercress



An invasive native of Eurasia also known as wild mustard and introduced to all states of Australia (except the NT). It grows to 30 cm and prefers moist soils in bushland, woodlands, open forests and urban gardens/lawns where it is considered a weed. It is an annual or biannual plant. Flower width to 7 mm and its above-ground parts are edible.

Family: Brassicaceae  
Genus: *Cardamine*  
Species: *Cardamine hirsuta*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Hairy tare





Also known as hairy vetch and tiny vetch, it is a small annual herb native to Europe and Western Asia and introduced to all states of Australia (except the NT). It grows to 90 cm and prefers bushland, woodlands and open forests. White flowers appear in a raceme of up to 8 (usually on one side) from September to November with a width to just 3 mm.

Family: Fabaceae  
Genus: *Vicia*  
Species: *Vicia hirsuta*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Lawn daisy



Also known as the common daisy, it is native to Europe but found in south eastern Australia from NSW to SA and Tasmania. It grows to 45 cm and flowers appear from September to March with a width to 35 cm. It prefers gardens and lawns. It is considered an invasive weed in Australia, though is often found in gardens for its flowers.

Family: Asteraceae  
Genus: *Bellis*  
Species: *Bellis perennis*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Milkmaids



A native perennial herb found in south eastern Australia from Queensland to South Australia and Tasmania. It grows to 60 cm high with each flower of each cluster head having a width to 27 mm. Flowers appear from September to November. It prefers bushland, woodlands and open drier forests. The edible tubers were eaten by early Aboriginals.

Family: Colchicaceae  
Genus: *Burchardia*  
Species: *Burchardia umbellata*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



## Prickly starwort



Also known as starwort, stitchwort and chickweed, it is a small herb found in south eastern Australia from Queensland to South Australia and Tasmania. It is a groundcover that grows to a height 30 cm and has a flower width of 23 mm. It prefers bushland and forests and the bush is prickly to touch.

Family: Caryophyllaceae  
Genus: *Stellaria*  
Species: *Stellaria pungens*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Spoon leaved sundew





Found in all states of Australia (except SA, WA and the NT), its a short carnivorous perennial plant. It prefers woodlands, bushland and open forests. It traps prey and digests their nutrients and minerals to survive in poor soils. White flowers appear in August to December with a width to 20 mm. Spoon-shaped leaves grow to 17 mm and turn from red to green.

Family: Droseraceae  
Genus: *Drosera*  
Species: *Drosera spatulata*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Onion weed



Also known as the three-corned leek and wild garlic, it was introduced from Europe and Africa and is found in all states of Australia. It has white flowers, grows to a height of 60 cm and prefers grasslands, bushland, woodlands and forests where it grows best in moist soils. The green stem is distinctly triangular in shape and all parts of this plant are edible.

Family: Amaryllidaceae  
Genus: *Allium*  
Species: *Allium triquetrum*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



## Oxeye daisy



Also known as the dog daisy, ox-eye daisy and marguerite, it native to Europe and found in south eastern Australia from QLD to SA and Tasmania. It grows to 80 cm and prefers bushland, woodlands and open forests. Flowers appear from November to February with a width to 7.5 cm and leaves can exceed 15 cm at base and decrease in side up stem.

Family: Asteraceae  
Genus: *Leucanthemum*  
Species: *Leucanthemum vulgare*

VIC N.S.W. QLD S.A. W.A. N.T. TAS

NATIVE WEED INTRODUCED NATURALISED



# WILDFLOWER



## White everlasting



Also known as the a fringed everlasting or the strawflower, it is a daisy native found in south eastern states of Australia from New South Wales to South Australia. It grows to 30 cm and prefers bushland, woodlands, open forests and even urban gardens where it has been successfully cultivated.

Family: Asteraceae  
Genus: *Chrysocephalum*  
Species: *Chrysocephalum baxteri*

VIC

N.S.W

QLD

S.A.

W.A.

N.T.

TAS

NATIVE

WEED

INTRODUCED

NATURALISED



# ORCHID



VIC N.S.W. QLD S.A. W.A. N.T. TAS




NATIVE WEED INTRODUCED NATURALISED


## White fingers

Also known as pink fingers, it is a native perennial herb found in eastern Australia from Queensland to SA and Tasmania. It grows to 20 cm on a single hairy stem and prefers grasslands, bushland, woodlands and forests as well as some coastal areas and near granite outcrops. Flowers appear from September to January with a width to 23 mm.

Family: Orchidaceae  
Genus: *Caladenia*  
Species: *Caladenia carnea*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Potato orchid

Also known as the potatoes and cinnamon bells, it is leafless native to Australia and found in all states of Australia (except Northern Territory). It grows to 60 cm and prefers bushland, woodlands, and forests with moist soils where bacterial and fungal agents can flourish. It flowers in later Spring to early Summer.

Family: Orchidaceae  
Genus: *Gastrodia*  
Species: *Gastrodia sesamoides*



VIC N.S.W. QLD S.A. W.A. N.T. TAS



NATIVE WEED INTRODUCED NATURALISED

## Parsons bands orchid

An single leaf orchid also known as the leafless parsons bands, it is found in all eastern Australia from Queensland to South Australia and Tasmania. It has white to pinkish flowers, which appear from December to June. It grows to a height of 25 cm and prefers grasslands, bushland, woodlands, forests and coastal areas.

Family: Orchidaceae  
Genus: *Eriochilus*  
Species: *Eriochilus cucullatus*





AUSTRALIAN  
**HABITATS**

**ALPINE**



**Areas of high altitude and given to snow in winter months**

**BUSHLAND**




**Bushlands are areas where the flora and fauna is predominately indigenous**

**FARMLAND**



**Paddocks used by farmers for agriculture and livestock**

**FOREST**



**Tree based habitat with an upper canopy height of more than 20 metres**



AUSTRALIAN  
**HABITATS**

**GRASSLAND**



**Grassland based habitat with few low shrubs and no tall trees**

**OPEN BUSHLAND**



**Open bushlands are areas where the flora and fauna is predominately indigenous and open**

**PARKLAND**



**Urban parkland area**

**RAINFOREST**



**A forest with a heavy annual rainfall**




AUSTRALIAN  
**HABITATS**

**SCRUB**



**Thick low growth habitat where vegetation ranges from 0.5 to 4 metres**

**URBAN**




**An area dominated by human population (characteristically a town or city)**

**WETLAND / MARSH / SWAMP**



**A low lying area of uncultivated land where water naturally collects**

**WOODLAND**



**Woodlands are tree based where the upper canopy height ranges from 8-20 metres**



# BASIC GLOSSARY

- .ssp** (*acronym*) Sub Species.
- ANNUAL** (*noun*) A plant that completes its entire life cycle within a single year. Different annual plants begin and end their cycles at different times of the year.
- ANTHROPOD** (*noun*) Any invertebrate animal having an exoskeleton, external segmented body and jointed appendages, such as ants, flies, beetles, cicadas, crabs, centipedes and dragonflies.
- BERRY** (*noun*) A small roundish juicy fruit that does not have a stone.
- CANE** (*noun*) The hollow jointed part of a tall grass, bamboo or sugar cane.
- CHITIN** (*noun*) A chemical compound that is a derivative of glucose and found in the cells walls of fungi as well as in other forms such as the exoskeletons of anthropoids such as cicadas.
- COMPOSITE FLOWER** (*noun*) A flower comprising of hundreds or even thousands of small flowers clustered together to form one large flower.
- CONIFER** (*noun*) A tree where the seeds are located within a cone.
- COROLLA** (*noun*) Whorl of petals of a flower.
- DECIDUOUS** (*adjective*) A tree that sheds all of its leaves each year.
- ENDEMIC** (*adjective*) prevalent or regularly found among a people or in a district; confined to a particular area (*biology*).
- ESCARPMENT** (*noun*) A long, steep slope, esp. one at the edge of a plateau or separating areas of land at different heights.
- FLORET** (*noun*) One of the small flowers that make up a composite flower head.
- FRUIT** (*noun*) The fleshy part of a plant that is eaten as food. Unlike vegetables, all fruits contain seeds. Fruits generally have some form of outer skin to protect them such as an apple skin.
- FUNGI** (*noun*) Any member of the group of eukaryotic organisms, including moulds yeasts, that have fruits such as mushrooms. Fungi all have chitin in their cell walls.
- HERB** (*noun*) Any seed bearing plant that has flowers or leaves that





*Artworkz*

*Serving the community*