

## TURF AND FOULING ALGAE: II. MAT & TURF SPECIES

### What are they?

Some marine algae exist as low “mats” or “turfs” (also called “turving” or “turf-like” algae by some workers). As the names imply, “mats” are flat and spreading, “turfs” are more upright like miniature mown lawns or grazed pastures, however, both terms are pretty subjective. Some mats heavily grazed by molluscs may be so low that they slip into the category of “crusts”.

Generally, the term “turf” has been used to describe:

- **communities** — mainly intertidal some running into shallow water — consisting of mixed species, all with a low profile.
- or
- **a single species** — its natural form, or a shape modified by heavy grazing.

Some workers expand their definition to include all those algae smaller than the brown algal canopy of a community. These algae should probably better be described as *understorey* species.

### Mat & turf species described in this key

Edgar, G & Shepherd, S A (2013) in *The Ecology of Australian temperate reefs*, CSIRO, described turf algae as thread-like, or leafy species in early stages of growth, ~ 10-20 mm tall. The key below uses this definition where possible, but strays to describing larger leafy and

filamentous plants that bloom seasonally on reefs and form dense, low mats a few cm high.

### Where are they found?

Some turfs may suddenly appear seasonally, often as a response to increased water temperature, light and dissolved nutrients. Others occur as permanent zones in the intertidal distributed according to a balance between resistance to radiation/drying and the distribution of animal grazers. They can be used to define zones on reefs, related to tidal levels.

### Limitations

Only macrophytes, those with some features that can be seen by the unaided eye, are considered below. Unfortunately, microscopic investigation will also be needed for definitive identifications.

### Images used below

Unless acknowledged otherwise, all images come from pressed specimens or the extensive slide collection of the algal unit, State Herbarium of S Australia, collections generated by Professor Womersley and his workers over some 60 years. Those with dark backgrounds have been taken using phase contrast or interference microscopy to highlight transparent structures. Other images may be stained dark blue.

### Scale

The coin used as a scale is 24 mm or almost 1” across

- 1a. plants forming a broken band of growth a few mm high on rock just above high tide (the supra-littoral zone), kept wet by wave splash, crumbling when dry, heavily grazed by small blue snails at lower levels ..... marine lichens (combination of fungus and microscopic alga) ..... 2.
- 1b. not as above ..... 3.
- 2a. dry plants orange-yellow. Fig. 1. .... *Caloplaca* spp
- 2b. dry plants black. Fig. 2. .... *Lichina* spp
- 3a. plants bright green ..... 4.
- 3b. plants dark red, red, amber or brown ..... 10.
- 4a. plants leafy or with narrow ribbon-like branches, bright yellow-green and with crinkled edges. Figs 3-7. .... *Ulva* (Sea lettuce) (including species with hollow parts, once placed into *Enteromorpha*) See **Southern Australian Groups at a glance: *Ulva***
- 4b. plants thread-like, yellowish, dark green or blue-green ..... 5.



Fig. 1: *Caloplaca* on boulders at Petrel Cove, SA



Fig. 2: *Lichina pygmaea*, Yorke Peninsula, SA being grazed by blue, littorinid snails (about 4 mm tall)



Fig. 3: *Ulva* mats exposed at low tide on mud flats of the Port River, S Australia, during December, plants attached to shell fragments



Fig. 4: *Ulva* (*Enteromorpha*) *compressa* mats at the edge of a flat reef, exposed at low tide during December, Rapid Bay, S Australia

Fig. 5: *Ulva* (*Enteromorpha*) *linza*, plants with broad blades



Fig. 6: *Ulva*, closely cropped on a boulder, with dead barnacles, Port River, S Australia



5a. threads consist of box-shaped cells in rows and columns; branches may be hollow. Figs 7-12. .... *Ulva* (species with hollow parts, once placed into *Enteromorpha*)

See **Southern Australian Groups at a glance: *Ulva***

5b. threads of elongate cells in single lines. .... 6.

6a. threads branched ..... 7.  
6b. threads unbranched ..... 9.

7a. branching regular, or irregular, **cross walls** occur near branches ..... 8.

7b. branching irregular, side branches **without** a cross wall, contents of parent cell continuous with side cell; plants in mats or cushions. Figs 13-15. .... *Cladophoropsis* 3 spp, 1 rare, 1 sp forms floating balls

8a. branching regular, cells usually elongate, hairs **absent**, rhizoids if present **not** tapering. Figs 16-18. .... *Cladophora*

See **Southern Australian Groups at a glance: *Cladophora* species groups I-III**

8b. branching irregular, cells irregularly shaped, often **bulging**, hairs may be **present**, rhizoids tapering to a fine point. Figs 19, 20. (next page). .... *Wittrockiella salina*




Fig. 7-9: *Ulva* (*Enteromorpha*) *ralfsii*, plants  
7. coarse hair-like appearance  
8. surface view of cells  
9. the hollow core seen after focusing through the surface




Fig. 10: *Ulva* (*Enteromorpha*) *clathrata*



Fig. 11: *Ulva* (*Enteromorpha*) *paradoxa*




Fig. 12: *Ulva* (*Enteromorpha*) *paradoxa*



Fig. 13: *Cladophoropsis herpestica*, from Elliston, S Australia, in shaded parts of the lower intertidal



Fig. 14: *Cladophoropsis membranacea*, forming low cushions at Cape Lannes, S Australia




Fig. 15: *Cladophoropsis herpestica*, side branches (*sb*) lack cross walls at their bases, and rhizoids (*rh*) arise from the bases of axial cells



Fig. 16: *Cladophora coelothrix*




Fig. 17: *Cladophora coelothrix*, branching pattern




Fig. 18 *Cladophora subsimplex* branching pattern of horizontal and upright branches

9a. threads coarse (> 0.7mm wide) plants often loose lying. Figs 21-23.

..... *Chaetomorpha* 8 spp

See **Southern Australian Groups at a glance: *Chaetomorpha***,

9b. threads fine (< 0.6 mm wide), forming tangled mats or loose-lying strands. Figs 24-29.

..... *Rhizoclonium* 4 spp

See **Southern Australian Groups at a glance: *Rhizoclonium***



Fig. 19: *Wittrockiella salina*




Fig. 20: *Wittrockiella salina*, irregularly shaped cells



Fig. 21: *Chaetomorpha linum*, detail of a thread



Fig. 22: *Chaetomorpha linum*



Fig. 23: *Chaetomorpha valida*



Fig. 24: *Rhizoclonium riparium*



Fig. 25: *Rhizoclonium tortuosum*, mixed with sea grass blades from intertidal mud flats




Fig. 26: *Rhizoclonium tortuosum*, detail of threads



Fig. 27: *Rhizoclonium curvatum*




Fig. 28: *Rhizoclonium curvatum*, detail of threads in arcs between rhizoids




Fig. 29: *Rhizoclonium tortuosum*, detail of thread, net-like chloroplasts

- 10a. plants pink, red to red-brown ..... 11.
- 10b. plants brownish ..... 14.

- 11a. plants pink to red, some bleaching to white, all *limey, brittle and jointed* (articulated); at outer edges of reefs. Figs 30-36. .... coralline red algae  
See **Pictured keys of common southern Australian marine plants: coralline red algae**

- 11b. hair-like or gristly, *not* limey; ... 12.

- 12a. plants hair-like, of dark red threads; <1mm wide; cells in a single line, with girdles and coatings of microscopic cells; may be attached to other turf plants. Figs 37-41.  
..... *Ceramium* spp, *Centroceros*  
See **Pictured key to southern Australian algae: filamentous red algae Part IV**

- 12b. plants red, some bleached to yellow, branches >1 mm wide, many cells wide ..... 13.




Fig. 30: *Cheilosporum sagittatum*, turf, shallow water, Aldinga reef edge, SA



Fig. 31: *Cheilosporum sagittatum* arrow-shaped sections of jointed branches



Fig. 32: bleaching coralline turf with brown algae, reef edge, Second Valley, SA



Fig. 33: *Corallina officinalis*




Fig. 34: *Corallina officinalis*, detail of jointed branches



Fig. 35: *Haliptilon roseum*, Stokes Bay, SA. Photo: D Muirhead



Fig. 36: *Haliptilon roseum*, detail of branches



Fig. 37: *Centroceros clavulatum*




Fig. 38: *Centroceros clavulatum*, spiny tips




Fig. 39: *Ceramium cliftonianum* forms entangled masses



Fig. 40: *Ceramium pusillum*, branch tips, cell girdles showing beneath coatings of additional (corticing) cells

Fig. 41: *Ceramium cliftonianum* detail of thread-cells with girdles of smaller cells



13a. plants stubby or in tangled masses;  
cross sections show a core of thick-  
walled rounded cells. Figs 42-46.

..... *Gelidium crinale*, *G. pusillum*,  
*Capreolia implexa*

See **Southern Australian species of  
Gelidiaceae at a glance**

13b. plants hair-like; cross sections show  
a core of well-spaced, many-armed  
cells. Figs 47-49.

.....*Gigartina brachiata*, *G. densa*

See **Southern Australian species of  
Gigartinaceae at a glance**



Fig. 42: *Gelidium pusillum*, with thin, gristly  
branches



Fig. 43: *Gelidium crinale*, forming hair-like  
turfs bleaching at the tips



Fig. 44: band or zone of red algal turf, mainly  
*Gelidium*, exposed between waves on  
granite, West I., SA



Fig. 45 *Capreolia implexa*,  
**often mistaken for *Gelidium pusillum***




Fig. 46: *Gelidium crinale*, cross section



Fig. 47: *Gigartina brachiata*, tangled masses, often  
with faint white bands on branches



Fig. 48: *Gigartina densa*, growing in stubby,  
gristly clumps at the lower edges of  
reefs in rough waters




Fig. 49: *Gigartina brachiata*, cross section


14a. plants fan-shaped, fans ~ 20-40 mm  
across, lying flat on reefs. Figs 50-52.

*Lobophora* spp, *Padina* spp.

See **Southern Australian species of  
Dictyotaceae at a glance**

14b. plants thread-like. Figs 53-55.

.....See **Turf and fouling algae I:  
Ectocarpaceae**



Figs 50, 51: *Lobophora variegata*, two views of flat masses (arrowed) at the reef edge, Cape Jervis SA



Figs 52: *Padina elegans*, fan-shaped blades, edges in-rolled, lie flat on reefs





Fig. 53: *Hincksia sordida*, Whyalla, SA



Fig. 54: *Feldmannia*, (stained blue) Whyalla, SA



Fig. 55: *Ectocarpus*, with spore sacs

## EXAMPLES OF TURF HABITATS IN SOUTH AUSTRALIA



Closely cropped turf of mixed species, greens (mainly *Ulva*) and reds (*Gelidium*) at the edge of the reef, Port Willunga



Red algal turf above green, at the edge of the reef, Port Willunga



Underwater view reef edge, Aldinga, turf of mixed red species and an emergent *Laurencia* plant (arrowed)



Seastar, *Petricia vernicina*, on turf including fan-shape *Lobophora*, characteristically fluorescing blue-green underwater, also sponges and pink, encrusting coralline red algae, 5m deep, Port Noarlunga



Slanting view of coralline turf (uppermost) and basal leaves of *Sargassum* on a granite boulder, exposed between waves, West Island



*Ulva* mats exposed at low tide, Robe reef platform

Broken red algal turf (arrowed) (*Gelidium*) bleached yellow, above the bead-like, brown canopy species, *Hormosira*, Slipway reef, Robe

