

Summary published in Mycotaxon 103: 197–200, 2008

Additions to the knowledge of lignocellulolytic basidiomycetes in forests from Santa Catarina, Southern Brazil

ELISANDRO RICARDO DRECHSLER-SANTOS¹

CLAUDIA GROPOSO² & CLARICE LOGUERCIO-LEITE³

¹ drechslersantos@yahoo.com.br

Dept. Micologia, Universidade Federal de Pernambuco, Recife, Pernambuco, Brazil

²Petrobrás – CENPES, Rio de Janeiro, Rio de Janeiro, Brazil

³Universidade Federal de Santa Catarina, Florianópolis, Santa Catarina, Brazil

Abstract — An updated checklist of the lignocellulolytic basidiomycetes in Santa Catarina State with 110 species distributed in 19 families and six orders (*Agaricales*, *Hymenochaetales*, *Polyporales*, *Russulales*, *Auriculariales*, *Tremellales*) is presented. The complete checklist is available on <http://www.mycotaxon.com/resources/weblists.html> and as Electronic Guide BASC (Basidiomycetes of Santa Catarina) as well on <http://www.cienciasbiologicas.ufsc.br/bot/micologia/index.htm>.

Introduction

Basidiomycota comprises almost 30,000 species with *Basidiomycetes* representing approximately 70% (more of the 20,000 spp.) of the phylum (Kirk et al. 2001, David 2002). The basidiomycetes include wood decomposing organisms in most ecosystems causing white and brown rots in tropical areas where they are very abundant (Nobles 1971, Dix & Webster 1995, Deacon 1997, Anagnost 1998, Highley & Dashek 1998, Bononi & Grandi 1999). In Southern Brazil, the Santa Catarina State (26°–30° S lat, 48°30'–54° W long) presents two major vegetation types: the Atlantic Rain Forest and the Atlantic Semi-deciduous Forest (Morellato & Haddad 2000). In these ecosystems, Polypore fungi were first studied by Europeans Mycologists already in the 19th century (Loguercio-Leite 1990). The contributions by Brazilian researches from the Mycology Laboratory (BOT/CCB/UFSC) in Santa Catarina State has been taking place since 1990, with several works based on collections from the Atlantic Forest (Loguercio-Leite & Wright 1991ab, Loguercio-Leite 1992, 1993, 1994; Loguercio-Leite & Wright 1995, Gerber 1996, Gerber & Loguercio-Leite 1997, Loguercio-Leite & Gerber 1997, Loguercio-Leite & Wright 1998, Loguercio-Leite et al. 1998, Gerber et al. 1999, Neves & Loguercio-Leite 1999, Gerber & Loguercio-Leite 2000, Gonçalves 2001, Loguercio-Leite et al. 2001, 2002; Groposo & Loguercio-Leite 2005). The purpose of this study was to estimate the species diversity of basidiomycetes in Santa Catarina State.

Material and methods

This study is based on the collections deposited in the herbarium FLOR since 1983 (Holmgren et al. 1990). Basidiomes were studied using macroscopic (shape, size, colors, types of hymenophore) and microscopic (presence/absence of structures, dimensions, basidiospores) characters (Ryvarden 1991). Measurements were made from slide preparations stained with 1% aqueous phloxine and 5% KOH. Melzer's reagent was used to define wall chemical characteristics. Specimens were identified to species using specialized references and comparing collections with BAFC, ICN and PACA Herbaria collections. Nomenclature, taxonomy and author citation followed Kirk et al. 2001 and databases: Centraalbureau voor Schimmelcultures – CBS (<http://www.cbs.knaw.nl/databases/>) and Index Fungorum – IFS (<http://www.indexfungorum.org/Names/Names.asp>). For geographic references consult the map of Santa Catarina State in Groposo & Loguercio-Leite (2005).

Results and discussion

As a contribution to the general checklist of the lignocellulolytic fungi (*Basidiomycetes*) in Santa Catarina State a total of 110 species is presented. The taxa are distributed in 19 families and six orders. *Polyporales* is the most representative order with 74 species in 12 families, followed by *Hymenochaetales* with 25 species in two families. The high diversity of *Polyporales* found in the present study agrees with the results of other inventories of the basidiomycetes on subtropical Southern Brazil (Groposo & Loguercio-Leite 2005) and also in tropical Northeastern Brazil (Gibertoni et al. 2004). Consequently, *Polyporaceae* is the most representative family with 45 species, followed by *Hymenochaetaceae* with 23 species. *Phellinus* s. l. is the genus with the highest number of species with 16 taxa. Most of the reported species showed a tropical distribution: 47 neotropical (almost 42.7%) and 42 pantropical species (38.2%). Only 8 and 13 species are considered widely distributed (7.3%) and cosmopolitan (11.8%), respectively. Up to now, these 110 species added here to the original list of lignocellulolytic fungi (Groposo & Loguercio-Leite 2005) increases to 157 species the total diversity of these organisms in Santa Catarina State, Brazil. The results revealed a high mycodiversity in Santa Catarina State, which corresponds to the knowledge of the basidiomycetes in the Atlantic Forest of Southern Brazil. From the total 157 species, 75.8% (119 species) are tropical (pantropical and neotropical). Besides the 4 species known only from Brazil [*Amauroderma corneri* Gulaid & Ryvarden, *Antrodiella multipileata* C.L. Leite & J.E. Wright, *Henningsia brasiliensis* (Speg.) Speg., *Skeletocutis roseolus* (Rick ex Theiss.) Rajchenb.] cited earlier by Groposo & Loguercio-

Leite (2005), we report ten additional species endemic to this country: *Phellinus bambusarum* (Rick) M.J. Larsen, *Ceriporiopsis cystidiata* C.L. Leite et al., *Rigidoporus amazonicus* Ryvarden, *Diplomitoporus dilutabilis* C.L. Leite & J.E. Wright, *Rubroporus carneoporis* C.L. Leite et al., *Tyromyces crassisporus* C.L. Leite & J.E. Wright, *Tyromyces hypocitrinus* (Berk.) Ryvarden, *Wrightoporia porilacerata* C.L. Leite et al., *Pyrofomes fulvoumbrinus* (Bres.) A. David & Rajchenb., and *Protomerulius substuppeus* (Berk. & Cooke) Ryvarden. Of these, six are known only to Santa Catarina State. Only eight of the 157 taxa cause brown rot [*Fistulina hepatica* (Schaeff.) With., *Fomitopsis feei* (Fr.) Kreisel, *Gloeophyllum striatum* (Sw.) Murrill, *Antrodia albida* (Fr.) Donk, *Laetiporus sulphureus* (Bull.) Murrill, *Amylosporus bracei* (Murrill) A. David & Rajchenb., *Wrightoporia avellanea* (Bres.) Pouzar, and *Wrightoporia porilacerata*]. Consequently, 93-95% of the lignocellulolytic fungi from Santa Catarina caused white rots, confirming the trends given for tropical and subtropical areas by several authors (Nobles 1971, Gilbertson 1980, Nakasone 1996).

Checklist for the species from Santa Catarina State – Part II:

AGARICALES Clem.

Fistulinaceae Lotsy

Fistulina hepatica (Schaeff.) With.

Bot. Arr. Brit. Pl., Edn 2 2: 405. 1792.

BASIONYM: *Boletus hepaticus* Schaeff., 1777.

DISTRIBUTION: cosmopolitan.

Ref.: Gilbertson & Ryvarden 1986; Gonçalves & Loguerio-Leite 2001.

HYMENOPCHAETALES Imazeki & Toki

Hymenochaetaceae Imazeki & Tori

Auricularia luteoumbrina (Romell) D.A. Reid

Kew Bull. 17: 279. 1963

BASIONYM: *Phaeoporus luteoumbrinus* Romell, 1901

DISTRIBUTION: neotropical.

Ref.: Reid 1963.

Hydnochaete peroxydata (Berk. ex Cooke) Dennis

Kew Bull., Addit. Ser. 3: 105. 1970

BASIONYM: *Hydnum peroxydatum* Berk. ex Cooke, 1891

DISTRIBUTION: neotropical.

Ref.: Ryvarden 1982.

Hymenochaete anomala Burt

Ann. Mo. Bot. Gdn 5: 358. 1918

DISTRIBUTION: neotropical.

Ref.: Job 1990, Parmasto 2001.

Hymenochaete damicornis (Link) Lév.

Annls Sci. Nat., Bot., sér. 3 5: 151. 1846

BASIONYM: *Stereum damicornis* Link

DISTRIBUTION: pantropical.

Ref.: Job 1990, Parmasto 2001.

Hymenochaete rhabarberina (Berk.) Cooke

Grevillea 8(48): 148. 1880

BASIONYM: *Corticium rhabarbarinum* Berk.

DISTRIBUTION: pantropical.

Ref.: Parmasto 2001.

Inonotus patouillardii (Rick) Imazeki

Bulletin of the Tokyo Sci. Mus. 6: 105. 1943

BASIONYM: *Polystictus patouillardii* Rick

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980, Ryvarden 2004.

Phellinus allardii (Bres.) S. Ahmad

Basidiomyc. W. Pakist.: 57. 1972

BASIONYM: *Fomes allardii* Bres.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980, Gerber & Loguerio-Leite 1997.

Phellinus apiahyalus (Speg.) Rajchenb. & J.E. Wright

Mycologia 79(2): 251. 1987

BASIONYM: *Fomes apiahyalus* Speg.

DISTRIBUTION: neotropical.

Ref.: Loguerio-Leite & Wright 1995, Neves & Loguerio-Leite 1999.

Phellinus bambusarum (Rick) M.J. Larsen

A survey of the world taxa. (Oslo) 3: 40. 1990

BASIONYM: *Poria bambusarum* Rick

DISTRIBUTION: neotropical (Known only from Southern Brazil).

Ref.: Rajchenberg 1987, Larsen & Cobb-Poule 1990.

Phellinus callimorphus (Lév.) Ryvarden

Prelim. Polyp. Fl. E. Afr. (Oslo): 145. 1980

BASIONYM: *Polyporus callimorphus* Lév.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980, Corner 1991, Loguerio-Leite & Wright 1995.

Phellinus cesatii (Bres.) Ryvarden

Norw. Jl. Bot. 19: 234. 1972

BASIONYM: *Poria cesatii* Bres.

DISTRIBUTION: neotropical.

Ref.: Ryvarden & Johansen 1980, Loguerio-Leite & Wright 1995.

Phellinus ferreus (Pers.) Bourdot & Galzin

Hyménomyc. de France (Sceaux): 627. 1928

BASIONYM: *Polyporus ferreus* Pers.

DISTRIBUTION: widely distributed.

Ref.: Gilbertson 1979, Ryvarden & Johansen 1980, Loguerio-Leite & Wright 1995, Ryvarden 2004.

Phellinus flavomarginatus (Murrill) Ryvarden
 Norw. Jl. Bot. 19: 234. 1972
 BASIONYM: *Fomitiporia flavomarginata* Murrill
 DISTRIBUTION: neotropical.
 Ref.: Loguerio-Leite & Wright 1995.

Phellinus glaucescens (Petch) Ryvarden
 Norw. Jl. Bot. 19: 234. 1972
 BASIONYM: *Poria glaucescens* Petch
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Larson & Cobb-poule 1990, Corner 1991.

Phellinus linteus (Berk. & M.A. Curtis) Teng
 Chung-Kuo Ti Chen-chun: 762. 1963
 BASIONYM: *Polyporus linteus* Berk. & M.A. Curtis
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Dai & Xu 1998, Ryvarden 2004.

Phellinus melanodermus (Pat.) M. Fidalgo
 Mem. N. Y. bot. Gdn. 17: 135. 1968
 BASIONYM: *Xanthochrous melanodermus* Pat.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Corner 1991, Gonçalves & Loguerio-Leite 2001.

Phellinus pullus (Mont. & Berk.) Ryvarden
 Norw. Jl. Bot. 19: 235. 1972
 BASIONYM: *Polyporus pullus* Mont. & Berk.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Larsen & Cobb-Poule 1990.

Phellinus punctatiformis (Murrill) Ryvarden
 Norw. Jl. Bot. 19: 235. 1972
 BASIONYM: *Fomitiporia punctatiformis* Murrill
 DISTRIBUTION: neotropical.
 Ref.: Ryvarden & Johansen 1980, Loguerio-Leite & Wright 1995, Ryvarden 2004.

Phellinus punctatus (Fr.) Pilát
 Atl. Champ. Europ. 3: 530. 1942
 BASIONYM: *Polyporus punctatus* Fr.
 DISTRIBUTION: widely distributed.
 Ref.: Nobles 1948, Gilbertson 1979, Ryvarden & Johansen 1980, Loguerio-Leite & Wright 1995, Ryvarden 2004.

Phellinus rhabarbarinus (Berk.) G. Cunn.
 Bull. N.Z. Dept. Sci. Industr. Res., Pl. Dis. Div. 164: 229. 1965
 BASIONYM: *Polyporus rhabarbarinus* Berk.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Corner 1991, Gerber & Loguerio-Leite 1997, Ryvarden 2004.

Phellinus sarcites (Fr.) Ryvarden
 Norw. Jl. Bot. 19: 235. 1972

BASIONYM: *Polyporus sarcites* Fr.

DISTRIBUTION: neotropical.

Ref.: Ryvarden & Johansen 1980, Ryvarden 2004.

Phellinus undulatus (Murrill) Ryvarden

Norw. Jl. Bot. 19: 235. 1972

BASIONYM: *Fomitiporia undulata* Murrill

DISTRIBUTION: neotropical.

Ref.: Ryvarden & Johansen 1980, David & Rajchenberg 1985, Loguerio-Leite & Wright 1995, Ryvarden 2004.

Phylloporia chrysites (Berk.) Ryvarden

Norw. Jl. Bot. 19: 235. 1972

BASIONYM: *Polyporus chrysites* Berk.

DISTRIBUTION: pantropical.

Ref.: Gilbertson & Ryvarden 1987, Ryvarden & Johansen 1980, Corner 1991.

Schizophoraceae Jülich

Schizophora flavigena (Berk & M.A. Curtis ex Cooke) Ryvarden

Mycotaxon 23: 186. 1985

BASIONYM: *Poria flavigena* Berk. & M.A. Curtis ex Cooke

DISTRIBUTION: cosmopolitan.

Ref.: Gilbertson & Hemmes 1997.

Schizophora paradoxula (Schrad.) Donk

Persoonia 5(1): 76. 1967.

BASIONYM: *Hydnellum paradoxum* Schrad.

DISTRIBUTION: cosmopolitan.

Ref.: Ryvarden & Johansen 1980, Gilbertson & Blackwell 1985, Nakasone 1990.

POLYPORALES Gäm.

Boreostereaceae Jülich

Mycobionia flava (Sw.) Fr.

Bull. Soc. Mycol. Fr. 10: 77. 1894

BASIONYM: *Peziza flava* Sw.

DISTRIBUTION: neotropical.

Ref.: Reid 1976.

Fomitopsidaceae Jülich

Fomitopsis feiei (Fr.) Kreisel

Ciencias Biológicas, Cuba 16: 83. 1971

BASIONYM: *Polyporus feiei* Fr.

DISTRIBUTION: neotropical.

Ref.: Ryvarden & Johansen 1980, Carranza-Morse & Gilbertson 1986, Corner 1989.

Ganodermataceae Donk

Amauroderma omphalodes (Berk.) Torrend

Broteria, Bot. 18: 131. 1920

BASIONYM: *Polyporus omphalodes* Berk.

DISTRIBUTION: neotropical.
Ref.: Furtado 1981, Ryvarden 1984.

Ganoderma annulare (Lloyd) Boedijn
Bull. Jard. Bot. Buitenzorg 16: 391. 1940
BASIONYM: *Fomes annularis* Lloyd
DISTRIBUTION: cosmopolitan.
Ref.: Gerber & Loguercio-Leite 1997.

Ganoderma lucidum (Curtis) P. Karst.
Revue Mycol. (Toulouse) 3(9): 17. 1881
BASIONYM: *Boletus lucidus* Curtis
DISTRIBUTION: widely distributed.
Ref.: Nobles 1948, Steyaert 1975, Bazzalo & Wright 1982.

Ganoderma oerstedii (Fr.) Torrend
Bull. Torrey Bot. Club 29: 606. 1902
BASIONYM: *Polyporus oerstedii* Fr.
DISTRIBUTION: neotropical.
Ref.: Bazzalo & Wright 1982, Ryvarden 2004.

Ganoderma multiplicatum (Mont.) Pat.
Bull. Soc. Mycol. Fr. 5(2,3): 74. 1889
BASIONYM: *Polyporus multiplicatus* Mont.
DISTRIBUTION: neotropical.
Ref.: Ryvarden 2004.

Humphreya coffeata (Berk.) Steyaert
Persoonia 7(1): 102. 1972
BASIONYM: *Polyporus coffeatus* Berk.
DISTRIBUTION: neotropical.
Ref.: Steyaert 1972, Ryvarden 1976.

Gloeophyllaceae Jülich

Gloeophyllum striatum (Sw.) Murrill
Torrey. Bot. Club 32(7): 370. 1905
BASIONYM: *Agaricus striatus* Sw.
DISTRIBUTION: pantropical.
Ref.: Ryvarden & Johansen 1980, Corner 1987.

Grammotheleaceae Jülich

Grammothele subargentea (Speg.) Rajchenb.
Mycotaxon 17: 280. 1983
BASIONYM: *Poria subargentea* Speg.
DISTRIBUTION: pantropical.
Ref.: Rajchenberg 1984.

Hapalopilaceae Jülich

Ceriporia xylostromatoides (Berk.) Ryvarden
Prelim. Polyp. Fl. E. Afr. (Oslo): 276. 1980
BASIONYM: *Polyoporus xylostromatoides* Berk.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980, David & Rajchenberg 1985, Gilbertson & Blackwell 1985.

Ceriporiopsis cystidiata C.L. Leite et al.

Mycotaxon 79: 286. 2001

DISTRIBUTION: neotropical (Known only from Southern Brazil).

Ref.: Loguerio-Leite et al. 2001.

Ceriporiopsis pannocincta (Romell) Gilb. & Ryvarden

Mycotaxon 22(2): 364. 1985

BASIONYM: *Polyporus pannocinctus* Romell

DISTRIBUTION: cosmopolitan.

Ref.: Gilbertson & Ryvarden 1986.

Meripilaceae Jülich

Antrodia albida (Fr.) Donk

Persoonia 4(3): 339. 1966

BASIONYM: *Daedalea albida* Fr.

DISTRIBUTION: widely distributed.

Ref.: Ryvarden & Johansen 1980, Gilbertson & Ryvarden 1986.

Hydnopolyporus fimbriatus (Fr.) D. A. Reid

Persoonia 2(2): 151. 1962

BASIONYM: *Polyporus fimbriatus* Fr.

DISTRIBUTION: neotropical.

Ref.: Reid 1962, Fidalgo 1963, Stalpers 1978.

Physisporinus sanguinolentus (Alb. & Schwein.) Pilát

Atlas Champ. Eur., Polypor., B3: 248. 1940

BASIONYM: *Boletus sanguinolentus* Alb & Schwein.

DISTRIBUTION: widely distributed.

Ref.: Gilbertson & Ryvarden 1987, Gerber & Loguerio-Leite 2000.

Rigidoporus amazonicus Ryvarden

Mycotaxon 28(2): 537. 1987

DISTRIBUTION: neotropical (Known only from Brazil).

Ref.: Ryvarden 1987, Gerber & Loguerio-Leite 2000.

Rigidoporus lineatus (Pers.) Ryvarden

Norw. J. Bot. 19: 236. 1972

BASIONYM: *Polyporus lineatus* Pers.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980.

Meruliaceae P. Karst.

Merulius tremellosus Schrad.

Spicil. Fl. Germ.: 139. 1794

DISTRIBUTION: pantropical.

Ref.: Eriksson & Ryvarden 1976, Ginn 1976.

Gloeoporus thelephoroides (Hook.) G. Cunn.

Bull. N.Z. Dept. Sci. Industr. Res., Pl. Dis. Div. 164: 111. 1965

BASIONYM: *Boletus thelephoroides* Hook.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980, Gilbertson & Ryvarden 1986, Corner 1989.

***Podoscyphaceae* Reid**

Cymatoderma caperatum (Berk. & Mont.) D.A. Reid

Kew. Bull. 10: 635. 1956

BASIONYM: *Thelephora caperata* Berk. & Mont.

DISTRIBUTION: neotropical.

Ref.: Reid 1965.

***Polyporaceae* Fr. ex Corda.**

Coriolopsis caperata (Berk.) Murrill

N. Amer. Fl. (New York) 9(2): 77. 1908

BASIONYM: *Polyporus caperatus* Berk.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980.

Coriolopsis polyzona (Pers.) Ryvarden

Norw. Jl. Bot. 19: 230. 1972

BASIONYM: *Polyporus polyzonus* Pers.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980.

Coriolopsis rigida (Berk. & Mont.) Murrill

N. Amer. Fl. (New York) 9(2): 75. 1908

BASIONYM: *Trametes rigida* Berk. & Mont.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980.

Datronia scutellata (Schwein.) Gilb. & Ryvarden

Mycotaxon 22(2): 364. 1985

BASIONYM: *Polyporus scutellatus* Schwein.

DISTRIBUTION: widely distributed.

Ref.: Gilbertson & Ryvarden 1986.

Dichomitus anoectoporus (Berk. & M.A. Curtis) Ryvarden

Mycotaxon 20(2): 331. 1984

BASIONYM: *Polyporus anoectoporus* Berk. & M.A. Curtis

DISTRIBUTION: neotropical.

Ref.: Ryvarden 1984.

Dichomitus citricremeus Masuka & Ryvarden

Mycol. Res. 103(9): 1128. 1999

DISTRIBUTION: pantropical.

Ref.: Masuka & Ryvarden 1999, Gonçalves & Loguerio-Leite 2001.

Diplomitoporus dilutabilis C.L. Leite & J.E. Wright

Mycotaxon 68: 48. 1998

DISTRIBUTION: neotropical (Known only from Southern Brazil).

Ref.: Loguerio-Leite & Wright 1998.

- Fomes fasciatus* (Sw.) Cooke
 Grevillea 14(69): 21. 1885
 BASIONYM: *Boletus fasciatus* Sw.
 DISTRIBUTION: neotropical.
 Ref.: Gilbertson & Ryvarden 1986.
- Fomitella supina* (Sw.) Murrill
 Bull. Torrey bot. Club 32(7): 365. 1905
 BASIONYM: *Boletus supinus* Sw.
 DISTRIBUTION: pantropical.
 Ref.: Gerber 1996.
- Hexagonia hydnoides* (Sw.) M. Fidalgo
 Mem. N. Y. bot. Gdn. 17: 64. 1968
 BASIONYM: *Boletus hydnoides* Sw.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980.
- Hexagonia tenuis* J.M. Hook
 Epicrisis Systematis Mycologici: 498. 1838
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Loguercio-Leite & Wright 1991.
- Laetiporus sulphureus* (Bull.) Murrill
 Annls mycol. 18(1/3): 51. 1920
 BASIONYM: *Boletus sulphureus* Bull.
 DISTRIBUTION: widely distributed.
 Ref.: Ryvarden & Johansen 1980, Corner 1984.
- Lentinus badius* (Berk.) Berk.
 J. Bot. 6: 491. 1847
 BASIONYM: *Panus badius* Berk.
 DISTRIBUTION: pantropical.
 Ref.: Pavlich 1976, Corner 1981.
- Lenzites betulina* (L.) Fr.
 Epicr. Syst. Mycol. (Upsaliae): 405. 1838
 BASIONYM: *Agaricus betulinus* L.
 DISTRIBUTION: cosmopolitan.
 Ref.: Gilbertson & Ryvarden 1987.
- Megasporoporia cavernulosa* (Berk.) Ryvarden
 Mycotaxon 16(1): 174. 1982
 BASIONYM: *Polyporus cavernulosus* Berk.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden et al. 1982.
- Megasporoporia setulosa* (Henn.) Rajchenb.
 Mycotaxon 16(1): 180. 1982
 BASIONYM: *Poria setulosa* Henn.
 DISTRIBUTION: neotropical.
 Ref.: Ryvarden et al. 1982.

- Microporellus dealbatus* (Berk. & M.A. Curtis) Murrill
 Bull. Torrey bot. Club 32(9): 483. 1905
 BASIONYM: *Polyporus dealbatus* Berk. & M.A. Curtis
 DISTRIBUTION: neotropical.
 Ref.: David & Rajchenberg 1985.
- Nigroporus vinosus* (Berk.) Murrill
 Bull. Torrey bot. Club 32(7): 361. 1905
 BASIONYM: *Polyporus vinosus* Berk.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Corner 1987.
- Pachykytospora alabamae* (Berk. & Cooke) Ryvarden
 Norw. Jl. Bot. 19: 233. (1972)
 BASIONYM: *Polyporus alabamae* Berk. & Cooke
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Gilbertson & Blackwell 1985.
- Pachykytospora papyracea* (Schwein.) Ryvarden
 Norw. Jl. Bot. 19: 233. 1972
 BASIONYM: *Boletus papyraceus* Schwein.
 DISTRIBUTION: cosmopolitan.
 Ref.: Ryvarden & Johansen 1980, Gerber & Loguerio-Leite 1997.
- Perenniporia martii* (Berk.) Ryvarden
 Norw. Jl. Bot. 19: 143. 1972
 BASIONYM: *Polyporus martius* Berk.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Gerber et al. 1999, Decock & Herrera-Figueroa 2000.
- Perenniporia medulla-panis* (Jacq.) Donk
 Persoonia 5(1): 76. 1967
 BASIONYM: *Boletus medulla-panis* Jacq.
 DISTRIBUTION: cosmopolitan.
 Ref.: Ryvarden & Johansen 1980, Gerber et al. 1999, Decock & Stalpers 2006.
- Perenniporia ohiensis* (Berk.) Ryvarden
 Norw. Jl. Bot. 19: 143. 1972
 BASIONYM: *Trametes ohiensis* Berk.
 DISTRIBUTION: neotropical.
 Ref.: Gerber et al. 1999.
- Perenniporia piperis* (Rick) Rajchenb.
 Nordic Jl. Bot. 7(5): 555. 1987
 BASIONYM: *Fomes piperis* Rick
 DISTRIBUTION: neotropical.
 Ref.: Gerber et al. 1999.
- Perenniporia stipitata* Ryvarden
 Mycotaxon 28(2): 535. 1987
 DISTRIBUTION: neotropical.
 Ref.: Decock & Ryvarden 1998, Gerber et al. 1999.

- Perenniporia tephropora* (Mont.) Ryvarden
 Norw. Jl. Bot. 19: 233. 1972
 BASIONYM: *Polyporus tephroporus* Mont.
 DISTRIBUTION: widely distributed.
 Ref.: Gilbertson & Ryvarden 1987, Ryvarden & Johansen 1980.
- Polyporus blanchetianus* Berk. & Mont.
 Annls Sci. nat., Bot., III 11:238. 1849
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980.
- Polyporus guianensis* Mont.
 Annls Sc. Nat., Bot., sér. 2 13(1): 201. 1840
 DISTRIBUTION: neotropical.
 Ref.: Ryvarden & Johansen 1980, Corner 1983.
- Polyporus infernalis* Berk.
 J. Bot., London 2: 637. 1843
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980.
- Polyporus leprieurii* Mont.
 Annls Sci. Nat., Bot., sér. 2 13: 203. 1840
 DISTRIBUTION: neotropical.
 Ref.: Ryvarden & Johansen 1980, Nuñez & Ryvarden 1995.
- Polyporus tricholoma* Mont.
 Annls Sci. Nat., Bot., sér. 2 8: 365. 1837
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Nuñez & Ryvarden 1995.
- Polyporus virgatus* Berk. & M.A. Curtis
 J. Linn. Soc., Bot. 10: 304, 1868
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Nuñez & Ryvarden 1995.
- Pseudofavolus cucullatus* (Mont.) Pat.
 Essai taxonomique: 81. 1900
 BASIONYM: *Favolus cucullatus* Mont.
 DISTRIBUTION: pantropical.
 Ref.: Ryvarden & Johansen 1980, Nuñez & Ryvarden 1995.
- Pyrofomes fulvoumbrinus* (Bres.) A. David & Rajchenb.
 Mycotaxon 22(2): 313. 1985
 BASIONYM: *Fomes fulvoumbrinus* Bres.
 DISTRIBUTION: neotropical (Known only from Southern Brazil).
 Ref.: David & Rajchenberg 1985.
- Rubroporus carneoporis* C.L. Leite et al.
 Mycotaxon 83: 223-227. 2002
 DISTRIBUTION: neotropical (Known only from Southern Brazil).
 Ref.: Loguerio-Leite et al. 2002.
- Skeletocutis diluta* (Rajchenb.) A. David & Rajchenb.

Mycotaxon 45: 144. 1992
BASIONYM: *Skeletocutis nivea* var. *diluta* Rajchenb.
DISTRIBUTION: pantropical.
Ref.: David & Rajchenberg 1992.

Stiptophyllum erubescens (Berk.) Ryvarden
Norw. Jl. Bot. 20: 4. 1973
BASIONYM: *Daedalea erubescens* Berk.
DISTRIBUTION: neotropical.
Ref.: Ryvarden 1973.

Tinctoporellus epimiltinus (Berk. & Broome) Ryvarden
Trans. Br. Mycol. Soc. 73(1): 18. 1979
BASIONYM: *Polyporus epimiltinus* Berk. & Broome
DISTRIBUTION: pantropical.
Ref.: Ryvarden & Johansen 1980.

Trametes cubensis (Mont.) Sacc.
Syll. Fung. (Abellini) 9: 198. 1891
BASIONYM: *Polyporus cubensis* Mont.
DISTRIBUTION: neotropical.
Ref.: Ryvarden 1982.

Trametes elegans (Spreng.) Fr.
Epicr. Syst. Mycol. (Upsaliae): 492. 1838
BASIONYM: *Daedalea elegans* Spreng.
DISTRIBUTION: pantropical.
Ref.: Ryvarden & Johansen 1980.

Trametes pavonia (Hook.) Ryvarden
Norw. Jl. Bot. 19: 236. 1972
BASIONYM: *Boletus pavonius* Hook.
DISTRIBUTION: neotropical.
Ref.: Gilbertson & Ryvarden 1987.

Trametes socotrana Cooke
Grevillea 11(no. 57) 39. 1882
DISTRIBUTION: pantropical.
Ref.: Ryvarden & Johansen 1980.

Tyromyces crassisporus C.L. Leite & J.E. Wright
Mycotaxon 41(1): 169. 1991
DISTRIBUTION: neotropical (Known only from Southern Brazil).
Ref.: Loguerio-Leite & Wright 1991.

Tyromyces hypocitrinus (Berk.) Ryvarden
Mycotaxon 20(2): 344. 1984
BASIONYM: *Polyporus hypocitrinus* Berk.
DISTRIBUTION: neotropical (Known only from Brazil).
Ref.: Ryvarden 1984.

Tyromyces tephrus (Pat.) Ryvarden
Occas. Pap. Farlow Herbarium of Cryptogamic Botany 18: 35. 1983

BASIONYM: *Poria tephra* Pat.

DISTRIBUTION: neotropical.

Ref.: Lowe 1966.

***Sistotrema*ceae** Jülich

Trechispora farinacea (Pers.) Liberta

Taxon 15(8): 318. 1966

BASIONYM: *Hydnium farinaceum* Pers.

DISTRIBUTION: cosmopolitan.

Ref.: Larsson 1995, Gilbertson & Hemmes 1997.

Trechispora mollusca (Pers.) Liberta

Can. J. Bot. 51(10): 1878. 1974

BASIONYM: *Boletus molluscus* Pers.

DISTRIBUTION: cosmopolitan.

Ref.: Ryvarden & Johansen 1980.

Steccherinaceae Parmasto

Flaviporus liebmannii (Fr.) Ginns

Can. J. Bot. 58(14): 1584. 1980

BASIONYM: *Polyporus liebmannii* Fr.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980.

Flaviporus subhydrophilus (Speg.) Rajchenb. & J.E. Wright

Mycologia 79(2): 259. 1987

BASIONYM: *Polystictus subhydrophilus* Speg.

DISTRIBUTION: neotropical.

Ref.: Gerber & Loguercio-Leite 1997.

Junguhnia polycystidifera (Rick) Rajchenb.

Nordic Jl. Bot. 7(5): 566. 1987

BASIONYM: *Poria polycystidifera* Rick

DISTRIBUTION: neotropical.

Ref.: Rajchenberg 1987.

Junguhnia undigera (Berk. & M.A. Curtis) Ryvarden

Mycotaxon 20(2): 359. 1984

BASIONYM: *Polyporus undiger* Berk. & M.A. Curtis

DISTRIBUTION: neotropical.

Ref.: Ryvarden 1984.

Steccherinum ochraceum (Pers.) Gray

Nat. Arr. Brit. Pl. (London) 1: 651. 1821

BASIONYM: *Hydnium ochraceum* Pers.

DISTRIBUTION: cosmopolitan.

Ref.: Gilbertson & Blackwell 1987, Ellis & Ellis 1990.

Steccherinum reniforme (Berk. & M.A. Curtis) Banker

Mem. Torrey. Bot. Club 12: 127. 1906

BASIONYM: *Hydnium reniforme* Berk. & M.A. Curtis

DISTRIBUTION: neotropical.
Ref.: Hjortstam & Larsson 1995.

RUSSULALES Kreisel ex. P.M. Kirk, P.F. Cannon & J.C. David

Bondarzewiaceae Kotl. & Pouzar

Amylosporus bracei (Murrill) A. David & Rajchenb.

Mycotaxon 22(2): 288. 1985

BASIONYM: *Poria bracei* Murrill

DISTRIBUTION: neotropical.

Ref.: David & Rajchenberg 1985.

Stecchericium seriatum (Lloyd) Maas Geest.

Proc. Kon. Ned. Akad. Wetensch., 2 Sectie 69: 325. 1966

BASIONYM: *Hydnus seriatum* Lloyd

DISTRIBUTION: neotropical.

Ref.: Burdsall & Nakasone 1983.

Wrightoporia avellanea (Bres.) Pouzar

Ceská Mykol. 20: 173. 1966

BASIONYM: *Poria avellanea* Bres.

DISTRIBUTION: pantropical.

Ref.: Ryvarden & Johansen 1980.

Wrightoporia porilacerata C.L. Leite et al.

Mycotaxon 67: 252. 1998

DISTRIBUTION: neotropical (Known only from Southern Brazil).

Ref.: Loguercio-Leite et al. 1998.

AURICULARIALES J. Schröt

Auriculariaceae Fr.

Auricularia delicata (Fr.) Henn.

Bot. Jb. 17: 492. 1893

BASIONYM: *Laschia delicata* Fr.

DISTRIBUTION: pantropical.

Ref.: Lowy 1952.

Auricularia fuscosuccinea (Mont.) Henn.

Bot. Jb. 17: 19. 1893

BASIONYM: *Hirneola fuscosuccinea* Mont.

DISTRIBUTION: pantropical.

Ref.: Lowy 1952.

Auricularia polytricha (Mont.) Sacc.

Atti Inst. Veneto Sci. Lett., ed Arti, Série 6 3: 722. 1885

BASIONYM: *Exidia polytricha* Mont.

DISTRIBUTION: cosmopolitan.

Ref.: Lowy 1952.

TREMELLALES Fr.

Aporpiaceae Bondartsev & Bondartseva

Elmerina caryaee (Schwein.) D.A. Reid

Persoonia 14(4): 471. 1992

BASIONYM: *Polyporus caryaee* Schwein.

DISTRIBUTION: cosmopolitan.

Ref.: Reid 1992.

Elmerina dimidiata (A. David) D.A. Reid 1992

Persoonia 14(4): 472. 1992

BASIONYM: *Aporpium dimidiatum* A. David

DISTRIBUTION: neotropical.

Ref.: David 1974.

Exidiaceae R. T. Moore

Protomerulius substuppeus (Berk. & Cooke) Ryvarden

Syn. Fung. (Oslo) 5: 212. 1991

BASIONYM: *Polyporus substuppeus* Berk. & Cooke

DISTRIBUTION: neotropical (Known only from Brazil).

Ref.: Rajchenberg 1987.

Acknowledgements

The work was supported by a Masters Scholarship from CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior/Brazil) to the first author. Many thanks are due to Dr. Leif Ryvarden, Dr. Mario Rajchenberg, and Dr. Orlando Popoff for expert review of the manuscript. Dr. Marcela Cáceres is warmly acknowledged for her assistance concerning the English language.

Literature cited

- Adaskaveg JE, Blanchette RA, Gilbertson RL. 1991. Decay of date palm wood by white-rot and brown-rot fungi. Canadian Journal of Botany 69: 615-629.
- Anagnos SE. 1998. Light Microscopic Diagnosis of Wood Decay. IAWA Journal 19(2): 141-167.
- Bazzalo ME, Wright JE. 1982. Survey of the Argentine species of the *Ganoderma lucidum* complex. Mycotaxon 16(1): 293-325.
- Bononi VLR, Grandi RAP. 1999. Zigomicetos, Basidiomicetos e Deuteromicetos. São Paulo: Instituto Botânica.
- Burdall HHJr, Nakasone KK. 1983. Species of effused *Aphyllophorales* (*Basidiomycotina*) from the Southeastern United States. Mycotaxon 17: 253-268.
- Carranza-Morse J, Gilbertson RL. 1986. Taxonomy of the *Fomitopsis rosea* complex (*Aphyllophorales*, *Polyporaceae*). Mycotaxon 25(2): 469-486.
- Corner EJH. 1981. The agaric genera *Lenticus*, *Panus* and *Pleurotus* with particular reference to Malaysian species. Beihefte zur Nova Hedwigia 69: 1-169.
- Corner EJH. 1983. Ad Polyporaceae I, *Amauroderma* and *Ganoderma*. Beihefte zur Nova Hedwigia 75: 1-182.
- Corner EJH. 1984. Ad Polyporaceas II, *Polyporus*, *Mycobonia*, and *Echonochaeete*, and III, *Piptoporus*, *Buglossoporus*, *Laetiporus*, *Meripilus*, and *Bondarzewia*. Beihefte zur Nova Hedwigia 78: 1-222.
- Corner EJH. 1987. Ad Polyporaceaeas IV, the genera *Daedalea*, *Flabellophora*, *Flavodon*, *Gloeophyllum*, *Heteroporus*, *Irpea*, *Lenzites*, *Microporellus*, *Nigrofomes*, *Nigroporus*,

- Oxyporus*, *Paratrichaptum*, *Rigidoporus*, *Scenidium*, *Trichaptum*, *Vanderbylia*, and *Steccherinum*. Beihefte zur Nova Hedwigia 86: 1-265.
- Corner EJH. 1989. Ad Polyporaceae VI, the genus *Trametes*. Beihefte zur Nova Hedwigia 97: 1-197.
- Corner EJH. 1991. Ad Polyporaceae VII, the xanthochroic polypores. Beihefte zur Nova Hedwigia 101: 1-175.
- Dai YC, Xu MQ. 1998. Studies on the medicinal polypore, *Phellinus baumii*, and its kin, *P. linteus*. Mycotaxon 67: 191-200.
- David A. 1974. *Aporpium dimidiatum*, nouvelle tremellale porée. Buletin Society Mycology 90(3): 179-185.
- David JC. 2002. A Preliminary Catalogue of the Names of Fungi above the Rank of Order. Constancea 83.
- David A, Rajchenberg M. 1985. Pore fungi from French Antilles and Guiana. Mycotaxon 22(2): 285-325.
- David A, Rajchenberg M. 1992. M. West African Polypores: New species and combinations. Mycotaxon 45: 131-148.
- Deacon JW. 1997. Modern Mycology. United Kingdom, Blackwell Science, 3rd edition.
- Decock C, Herrera Figueroa S. 2000. Studies in *Perenniporia*. *Navisporus ortizii*, a synonym of *Perenniporia martius*, and a note on *Navisporus* and *Perenniporia* in Cuba. Cryptogamie Mycologie 21(3): 153-162.
- Decock C, Ryvarden L. 1998. *Microporellus straminellus* comb. nov. and a note on *Perenniporia stipata*. Cryptogamie Mycologie 19(3): 171-179.
- Decock C, Stalpers JA. 2006. Studies in *Perenniporia*: *Polyporus unitus*, *Boletus medulla-panis*, the nomenclature of *Perenniporia*, *Poria* and a note on European *Perenniporia* with a resupinate basidiome. Taxon 55(3): 759-778.
- Dix NJ, Webster J. 1995. Fungal Ecology. Ed. Chapman & Hall, First edition: 161-163.
- Ejechi BO, Obuekwe CO, Ogbimi AO. 1996. Microchemical Studies of Wood Degradation by Brown Rot and White Rot Fungi in two Tropical Timbers. International Biodeterioration & Biodegradation: 119-122.
- Ellis MB, Ellis JP. 1990. Fungi Without Gills (*Hymenomycetes* and *Gasteromycetes*). An Identification Handbook. Chapman and Hall, 1st Ed. 329p.
- Eriksson J, Ryvarden L. 1976. The *Corticiaceae* of North Europe. *Hypodermella* – *Mycoacia*. Synopsis Fungorum 4, Oslo, Fungiflora.
- Fidalgo O. 1963. Studies on the type species of *Hydnopolytoporus*. Mycologia 55: 713-727.
- Furtado JS. 1981. Taxonomy of *Amauroderma* (Basidiomycetes, Polyporaceae). Memoirs of the New York Botanical Garden 34: 1-109.
- Gerber AL. 1996. Fungos Xilófilos poróides (*Aphylophorales*) on the Lagoa da Conceição, Ilha de Santa Catarina, SC, Brasil. Insula 25: 3 – 68.
- Gerber AL, Loguerico-Leite C. 1997. New records of polypores (*Aphylophorales*) from southern Brazil. Mycotaxon 62: 305-318.
- Gerber AL, Loguerico-Leite C. 2000. Polyporoid wood-rotting fungi (*Basidiomycetes*) II – New records from southern Brazil. Mycotaxon 76: 175-185.
- Gerber AL, Neves MA, Loguerico-Leite C. 1999. Some species of *Perenniporia* Murrill (*Poriales*, *Basidiomycotina*) from Southern Brazil. Revista Brasileira de Botânica 22(2): 185-193.
- Gibertoni TB, Ryvarden L, Cavalcanti MAQ. 2004. Studies in neotropical polypores 18 New species from Brazil. Synopsis Fungorum 18: 44-56.
- Gilbertson RL. 1979. The genus *Phellinus* (*Aphylophorales*: *Hymenochaetaceae*) in Western North America. Mycotaxon 9(1): 51-60.
- Gilbertson RL. 1980. Wood-rotting fungi of North América. Mycologia 72(1): 1-49.
- Gilbertson RL, Blackwell M. 1985. Notes on Wood-rotting on junipers in the Gulf Coast region. Mycotaxon 24: 325-348.

- Gilbertson RL, Blackwell M. 1987. Notes on Wood-Rotting Fungi on Junipers in the Gulf Coast Region, II. *Mycotaxon* 28(2): 369-402.
- Gilbertson RL, Hemmes DE. 1997. Notes on fungi on Hawaiian tree ferns. *Mycotaxon* 62: 465-487.
- Gilbertson RL, Ryvarden L. 1986. North American Polypores. *Synopsis Fungorum* 1. Oslo, Fungiflora, 433p.
- Gilbertson RL, Ryvarden L. 1987. North American Polypores. *Synopsis Fungorum* 2. Oslo, Fungiflora, 423p.
- Ginns J. 1976. *Merulius* s.s. and s.l., taxonomic disposition and identification of species. *Canadian Journal of Botany* 54: 100-167.
- Gonçalves GVC, Loguercio-Leite C. 2001. Biodiversidade de Fungos Poróides Xilófilos (Basidiomycetes), na Unidade de Conservação Ambiental Desterro (UCAD), Ilha de Santa Catarina, SC, Brasil. *Insula* 30: 1-19.
- Groposo C, Loguercio-leite C. 2005. Contribution to the lignocellulolytic fungi (*Basidiomycetes*) of the Atlantic Rain Forest in Southern Brazil. *Mycotaxon* 92: 103-106.
- Highley TL, Dashek WV. 1998. Biotechnology in the Study of Brown- and White-rot Decay. *Forest products Biotechnology*: 15-35.
- Hjortstam K, Larsson KH. 1995. Annotated check-list to genera and species of corticioid fungi (*Aphyllophorales*, *Basidiomycotina*) with special regards to tropical and subtropical areas. *Windahlia* 21: 1-75.
- Holmgren PK, Holmgren NH, Barnett LC. 1990. Index herbariorum: Part I: Herbaria of the World. 86th ed. Bronx, New York Botanical Garden.
- Job DJ. 1990. Le genre *Hymenochaete* dans les zones tempérées de l'hémisphère sud. *Mycologia Helvetica* 4: 1-151.
- Kirk PM, Cannon PF, David JC, Stalpers JA. 2001. Dictionary of the Fungi. 9th Ed, CAB International.
- Larsson KH. 1995. Taxonomy of *Trechispora farinacea* and proposed synonyms I. Species with a grandinoid or hydnoid hymenophore. *Symbolae Botanicae Upsalienses* 30(3): 101-118.
- Larsen MJ, Cobb-Poule LA. 1990. *Phellinus* (*Hymenochaetaceae*): a Survey of the World Taxa., *Synopsis Fungorum* 3. Oslo, Fungiflora: 206p.
- Loguercio-Leite C. 1990. Revisão histórica sobre fungos poliporóides (*Aphyllophorales*) xilófilos de Santa Catarina. *Insula* 20: 3-10.
- Loguercio-Leite C. 1992. El género *Polyporus* (*Polyporaceae*) en la Isla de Santa Catarina, Santa Catarina, Brasil. *Boletín de la Sociedad Argentina* 28(1-4): p. 27-36.
- Loguercio-Leite C. 1993. *Polyporaceae* II: *Trametes* Fr. Na Ilha de Santa Catarina, SC, Brasil. *Insula* 22: 3-20.
- Loguercio-Leite C. 1994. *Polyporaceae* na Ilha de Santa Catarina III: o gênero *Hexagonia* Fr. *Insula* 23: 3-14.
- Loguercio-Leite C, Gerber AL. 1997. Non-pileate polypores on Santa Catarina Island, SC, Brazil. *Mycotaxon* 64: 285-301.
- Loguercio-Leite C, Gerber AL, Ryvarden L. 1998. *Wrigthoporia porilacerata*, a new species of pore fungi from southern Brazil. *Mycotaxon* 67: 251-255.
- Loguercio-Leite C, Gonçalves GV, Ryvarden L. 2001. Studies in Neotropical polypores 13. *Ceriporiopsis cystidiata* sp. nov. *Mycotaxon* 79: 285-288.
- Loguercio-Leite C, Ryvarden L, Groposo C. 2002. Studies in neotropical polypores 16. *Rubroporus carneoporis* genus & species nova. *Mycotaxon* 83: 223-227.
- Loguercio-Leite C, Wright JE. 1991a. Contribution to a biogeographical study of the austroamerican xilophilous polypores (*Aphyllophorales*) from Santa Catarina Island, SC, Brazil. *Mycotaxon* 41(1): 161-166.
- Loguercio-Leite C, Wright JE. 1991b. New South American Pileate Polypores (*Polyporaceae*) from Santa Catarina Island, SC, Brazil. *Mycotaxon* 41(1): 167-172.
- Loguercio-Leite C, Wright JE. 1995. The Genus *Phellinus* (*Hymenochaetaceae*) on the Island of Santa Catarina, Brazil. *Mycotaxon* 54: 361-388.

- Loguercio-Leite C, Wright JE. 1998. *Diplomitoporus dilutabilis* a new species of *Polyporaceae* (*Aphyllophorales*) from Santa Catarina Island, Brazil. Mycotaxon 68: 47-51.
- Lowe JL. 1966. *Polyporaceae* of North America – the genus *Poria*. State University College of Forestry at Syracuse University of Technology 90: 1-183.
- Lowy B. 1952. The genus *Auricularia*. Mycologia 44: 656-692.
- Masuka AJ, Ryvarden L. 1999. *Dichomitus* in Africa. Mycological Research 103: 1126-1130.
- Morellato LPC, Haddad CFB. 2000. Introduction: the Brazilian Atlantic Forest. Biotropica 32(4b): 786-792.
- Nakasone KK. 1990. Cultural studies and identification of wood-inhabiting *Corticiaceae* and selected *Hymenomycetes* from North America. Mycological Memoir 15: 1-412.
- Nakasone KK. 1996. Diversity of lignicolous *Basidiomycetes*. In: McMiinn JW, Crossley DA. Biodiversity and Coarse Windy Debris in Southern Forests, Proceedings of the Workshop on coarse woody debris in Southern forests: effects on Biodiversity. Department of Agriculture, Forest Service, Southern Research Station, Athens. 146p.
- Nobles MK. 1948. Identification of cultures of wood-rotting fungi. Canadian Journal of Research 26: 281-431.
- Nobles MK. 1971. Cultural characters as a guide to the taxonomy of the *Polyporaceae*. In: Petersen R. Evolution in the higher *Basidiomycetes*. University of Tennessee Press, Knoxville: 169-196.
- Neves MA, Loguercio-Leite C. 1999. Cultural characteristics and taxonomy of some polypores (*Aphyllophorales*) from Santa Catarina Island, SC, Brazil. Mycotaxon 70: 193-202.
- Nuñez M, Ryvarden L. 1995. *Polyporus* (*Basidiomycotina*) and related genera. Synopsis Fungorum 10. Oslo, Fungiflora, 85p.
- Parmasto E. 2001. New Data on Rare Species of *Hydnochaete* and *Hymenochaete* (*Hymenochaetales*) Mycotaxon 91:137-163.
- Pavlich M. 1976. *Ascomycetes* y *Basidiomycetes* del Perú. I Memorias del Museo de Historia Natural "Javier Prado" 17.
- Rajchenberg M. 1984. Basidiomicetos xilófilos de la region Mesopotámica, República Argentina, V. Poliporos resupinados. Revista de Investigación Agropecuaria, INTA 19(1): 1-105.
- Rajchenberg M. 1987. Type studies of *Polyporaceae* (*Aphyllophorales*) described by J. Rick. Nordic Journal of Botany 7(5): 533-568.
- Reid DA. 1962. Notes on Fungi which have been referred to the *Thelephoraceae* sensu lato. Persoonia 2(2): 109-170.
- Reid DA. 1963. New or interesting records of Australasian *Basidiomycetes* V – *Aphyllophorales*. Kew Bulletin 17(2): 267-308.
- Reid DA. 1965. A monograph of the stipitate steroid fungi. Beihefte zur Nova Hedwigia 18: 1-483.
- Reid DA. 1976. Notes on Polypores. 2. Memoirs of the New York Botanical Garden 26(1): 179-198.
- Reid DA. 1992. The genus *Elmerina* (*Tremellales*), with accounts of two species from Queensland, Australia. Persoonia 14: 465-474.
- Ryvarden L. 1973. New Genera in the *Polyporaceae*. Norwegian Journal of Botany, 20(1): 1-5.
- Ryvarden L. 1976. Type studies in the *Polyporaceae* 7. Species described by J.M. Berkeley from 1836-1843. Kew Bulletin 31: 81-103.
- Ryvarden L. 1982. The genus *Hydnochaete* (*Hymenochaetaceae*). Mycotaxon 15: 425-447.
- Ryvarden L. 1982. Type-studies in the *Polyporaceae* 11. Species described by J.F. Montagne. Either alone or with other authors. Nordic Journal of Botany 2: 75-82.
- Ryvarden, L. 1984. Type studies in the *Polyporaceae* 16. Species described by J.M. Berkeley either alone or with other mycologists from 1856 to 1886. Mycotaxon 20(2): 329-363.
- Ryvarden L. 1987. New and noteworthy polypores from tropical America. Mycotaxon 28(2): 525-541.

- Ryvarden L. 1991. Genera of Polypores. Nomenclature and taxonomy. Synopsis Fungorum 5. Oslo, Fungiflora, 226p.
- Ryvarden L. 2004. Neotropical polypores Part 1. Synopsis Fungorum 19. Oslo, Fungiflora, 229p.
- Ryvarden L, Johansen I. 1980. A preliminary polypore flora of East Africa. Oslo, Fungiflora, 630p.
- Ryvarden L, Wright JE, Rajchenberg M. 1982. *Megasporoporia*, a new genus of resupinate polypores. Mycotaxon 16(1): 172-182.
- Singer R. 1975. The Agaricales in Modern Taxonomy. Chicago, 3rd Ed.
- Stalpers JA. 1978. Identification of wood-inhabiting *Aphylophorales* in pure culture. Studies in Mycology 16: 1-248.
- Steyaert RL. 1972. Species of *Ganoderma* and related genera mainly of the Bagor and Leiden Herbaria. Persoonia 7(1): 55-118.
- Steyaert RL. 1975. The concept and circumscription of *Ganoderma tornatum* (Pers.) Bres. Transactions of the British Mycological Society 65(3): 451 – 567.