

Invasive Shot-Hole Borer and Fusarium Dieback Field Guide

Identifying Polyphagous and Kuroshio Shot-Hole Borer in California

Background

A. Beetle-Fungal Complex

Adult female burrowing into wood (A1); colonies of the beetles symbiotic fungi recovered in the lab (A2). The invasive shot-hole borers (ISHB), *Euwallacea* spp., are invasive beetles that vector the plant disease fusarium dieback (FD). “ISHB” refers collectively to the polyphagous and Kuroshio shot-hole borers. Over 260 plant species have been attacked by ISHB and FD. The disease disrupts the flow of water and nutrients in susceptible hosts, which can kill individual branches or, in severe cases, the entire tree. It is caused by the fungi that the beetle uses as a food source: PSHB (*Fusarium euwallaceae*, *Graphium euwallaceae*, and *Paracremonium pembeum*), KSHB (*Fusarium kuroshium*, and *Graphium kuroshium*).

AKIF ESKALEN, University of California, Riverside; JOHN KABASHIMA, University of California Cooperative Extension; MONICA DIMSON, University of California Cooperative Extension; and SHANNON LYNCH, University of California, Santa Cruz.






B. Beetle Biology and Identification




ISHB bore tunnels (galleries) into host trees where they lay their eggs and grow the fungi. The two beetle species are physically identical. At 1.8 to 2.5 mm long, ISHB adult beetles are smaller than a sesame seed. The adult females (B1) are larger than the adult males (B2) and are also darker (B3). Most of the beetle's life cycle, from larva to adult (B4), is spent in the galleries. Mature siblings mate with each other so females are already pregnant when they leave to start their own galleries.

C. Signs and Symptoms



Entry holes are round and about 0.85 mm wide, the size of a ball-point pen tip (C1). The abdomen of the female beetle may be seen sticking out of the hole (C2). Tree symptoms are unique to each host species. Around the entry hole, look for dark, wet staining that sometimes dries to white or yellow (C3), thick gumming (C4), powdery white exudate (C5), or frass (C6), which resembles sawdust.



Symptoms of infection by FD pathogens include brown to black discoloration on wood beneath the bark. Scrape away bark around the entry or exit hole to reveal dark staining surrounding the gallery (C7, C8). Cross-sections of cut branches show the extent of infection (C9). Branch dieback is the result of advanced infection by ISHB's associated fungi. It may begin on a few branches (C10) and can eventually kill entire trees (C11).


D. Look-Alike Pests

Other pests may cause damage similar to that of ISHB-FD. Clues that may indicate a pest other than ISHB include

- staining, gumming, or exudate but no entry hole
- entry holes with an irregular shape (not round)
- entry holes larger or smaller than a ball-point pen tip
- the tree is not a known host of ISHB-FD (e.g., pine or most eucalyptus species)

Visit the University of California Statewide Integrated Pest Management Program website, ipm.ucanr.edu, to learn more about these pests.

Look-Alike Pests That May Have an Entry Hole




Foamy bark canker caused by *Geosmithia* sp. #41

Spread by the western oak bark beetle (*Pseudopityophthorus pubipennis*). Hosts: Coast live oak; stressed or dying trees. Look for beetles (D1) 1.7 to 2.3 mm long; reddish frass (D2), reddish sap, wet discoloration, and/or foamy liquid (D3) (a sign of infection) from an entry hole (1 mm) that is smaller than that of ISHB; dead tissue around entry hole beneath bark (D4).




Fruit tree shot-hole borer (*Scolytus rugulosus*) (D5)

Hosts: Fruit trees in *Prunus* genus, English laurel. Look for entry holes (2 mm) oozing sap or frass; the holes are larger than those of ISHB, with slightly rougher edges (D6.). Exit holes are sap free.



Western sycamore borer, *Synanthedon resplendens*

Hosts: Sycamore, oak, and ceanothus. Look for larvae 25 to 38 mm long (D7); roughened bark (D8); reddish sawdust-like frass and/or pupal cases (D9) in bark crevices or on ground; bleeding.




Lesser ambrosia beetle, *Xyleborinus saxeseni*

Hosts: Stressed and dying trees. Look for beetles 2 to 2.4 mm long (D10); entry holes (1 mm) smaller than those of ISHB (D11); reddish frass and/or sap; wet discoloration and/or dead tissue around entry hole and beneath bark (D12).



Oak ambrosia beetles, *Monarthrum dentiger*, *M. scutellare* (d)

Hosts: Oak species, tanoak, California buckeye. Look for slightly larger beetles (D13) (*M. scutellare*, 3.5 to 4.1 mm long; *M. dentiger*, 1.9 to 2.4 mm long) and entry holes (1 to 1.5 mm diameter) with bleeding, frothing, bubbling, or white boring dust (D14) that is tan when oxidized. Often attack stressed trees.



Look-Alike Pests That Lack an Entry Hole

Avocado trunk canker caused by *Phytophthora menzei* (D15, D16). Avocado branch canker and dieback caused by *Botryosphaeria* spp. and *Phomopsis* sp. Bacterial canker caused by *Xanthomonas campestris* (D17, D18). Black streak disease caused by *Botryosphaeria* spp. (D19–D22).

E. Reproductive Hosts

Reproductive host species support beetle reproduction and the growth and development of the symbiotic fungi. Each species is affected differently. Trees may be more susceptible if they are already under stress due to other pests, diseases, or environmental conditions or are in close proximity to an existing infestation. This list of species is not meant to be used as a do-not-plant list. However, as known hosts of ISHB-FD, species on this list should be closely monitored for potential infestation. Visit the Invasive Shot-Hole Borers website, www.pshb.org, for updates.

Acacia spp.

Acer buergerianum

Acer macrophyllum

Acer negundo

Acer palmatum

Acer paxii

Aesculus californica

Ailanthus altissima

Albizia julibrissin

Alectryon excelsus

Alnus rhombifolia

Archontophoenix

cunninghamiana

Baccharis salicifolia

Bauhinia variegata

Brachychiton populneus

Camellia semiserrata

Castanospermum australe

Cercidium (= *Parkinsonia*)
floridum

Cercidium (= *Parkinsonia*)
sonorae

Cocculus laurifolius

Cupaniopsis anacardioides

Erythrina coralloides

Erythrina falcata

Eucalyptus ficifolia

Fagus crenata

Ficus altissima

Ficus carica

Gleditsia triacanthos

Harpullia pendula

Howea forsteriana

Ilex cornuta

Koelreuteria bipinnata

Liquidambar styraciflua

Magnolia grandiflora

Parkinsonia aculeata

Persea americana

Platanus mexicana

Platanus racemosa

Platanus x acerifolia

Populus fremontii

Populus nigra

Populus trichocarpa

Prosopis articulata

Quercus agrifolia

Quercus chrysolepis

Quercus engelmannii

Quercus lobata

Quercus robur

Quercus suber

Ricinus communis

Salix babylonica

Salix gooddingii

Salix laevigata

Salix lasiolepis

Tamarix ramosissima

Wisteria floribunda

Xylosma avilae

Native Host Species

1. Big Leaf Maple



Acer macrophyllum

Native reproductive host

Signs/Symptoms: Staining

Native Host Species

2. Box Elder



Acer negundo

Native reproductive host

Signs/Symptoms: Staining,
bleeding, frass

Native Host Species

3. California Sycamore



Platanus racemosa

Native reproductive host

Signs/Symptoms: Staining

Native Host Species

4. Red Willow



Salix laevigata

Native reproductive host

Signs/Symptoms: Staining,
frass

Native Host Species

5. Godding's Black Willow



Salix gooddingii

Signs/Symptoms: Staining

Native Host Species

6. Fremont Cottonwood



Populus fremontii

Native reproductive host

Signs/Symptoms: Staining

Native Host Species

7. White Alder




Alnus rhombifolia

Native reproductive host

Signs/Symptoms: Staining

Native Host Species

8. Coast Live Oak



Quercus agrifolia

Native reproductive host

Signs/Symptoms: Staining

Native Host Species

9. Engelmann Oak



Quercus engelmannii
Native reproductive host
Signs/Symptoms: Staining

Native Host Species

10. Valley Oak



Quercus lobata

Native reproductive host

Signs/Symptoms: Staining

Native Host Species

11. Mesquite



Prosopis articulata

Native reproductive host

Signs/Symptoms: Staining

Agricultural Host Species

12. Avocado



Persea americana

Reproductive host

Signs/Symptoms: Sugary
exudate, staining

Non-Native Host Species

13. Castor Bean



Ricinus communis

Invasive reproductive host

Signs/Symptoms: Staining

Non-Native Host Species

14. Tree of Heaven



Ailanthus altissima

Invasive reproductive host

Signs/Symptoms: Staining

Non-Native Host Species

15. Acacia



Acacia spp.

Reproductive host

Signs/Symptoms: Gumming,
staining

Non-Native Host Species

16. Silk Tree/Mimosa



Albizia julibrissin

Reproductive host

Signs/Symptoms: Staining,
gumming

Non-Native Host Species

17. Coral Tree



Erythrina coralloides
Reproductive host
Signs/Symptoms: Staining

Non-Native Host Species

18. Palo Verde



Parkinsonia aculeata

Reproductive host

Signs/Symptoms: Staining,
frass

Non-Native Host Species

19. Moreton Bay Chestnut



Castanospermum australe

Reproductive host

Signs/Symptoms: Staining,
gumming

Non-Native Host Species

20. Chinese Flame/Goldenrain



Koelreuteria bipinnata, *K. paniculata*
FD-susceptible hosts
Signs/Symptoms: Gumming,
staining

Non-Native Host Species

21. Japanese Maple



Acer palmatum

Reproductive host

Signs/Symptoms: Staining

Non-Native Host Species

22. Trident Maple



Acer buergerianum

Reproductive host

Signs/Symptoms: Staining

Non-Native Host Species

23. English Oak



Quercus robur

Reproductive host

Signs/Symptoms: Staining

Non-Native Host Species

24. Cork Oak



Quercus suber

Reproductive host

Signs/Symptoms: Staining

Non-Native Host Species

25. Weeping Willow



Salix babylonica

Reproductive host

Signs/Symptoms: Staining,
gumming, frass

Non-Native Host Species

26. Shiny Xylosma



Xylosma avilae

Reproductive host

Signs/Symptoms: Staining

Non-Native Host Species


27. American Sweetgum



Liquidambar styraciflua
Reproductive host
Signs/Symptoms: Staining

Non-Native Host Species

28. Kruuajong



Brachychiton populneus

Reproductive host

Signs/Symptoms: Gumming

Non-Native Host Species

29. London Plane



Platanus x acerifolia

Reproductive host

Signs/Symptoms: Staining

Non-Native Host Species

30. Kentia Palm



Howea forsteriana

Reproductive host

Signs/Symptoms: Gumming,
frass

Non-Native Host Species

31. Camellia



Camellia semiserrata
Reproductive host
Signs/Symptoms: Staining

References

- Eskalen, A., et al. 2013. Host range of *Fusarium dieback* and its ambrosia beetle (Coleoptera: Scolytinae) vector in Southern California. *Plant Disease* 97(7): 938–951.
- Lynch, S., et al. 2016. Identification, pathogenicity, and abundance of *Paracremonium* sp. nov. and *Graphium euwallaceae* sp. nov. *Mycologia* 108(2): 313–329.
- O'Donnell, K., et al. 2016. Invasive Asian *Fusarium*. *Phytoparasitica* 44(4): 435–442.
- Stouthamer, R. et al. 2017. Tracing the origin of a cryptic invader. *Agricultural and Forest Entomology* 19(4): 366–375.

Photo Credits

The photographs in this publication are by Akif Eskalen, John Kabashima, or Monica Dimson except for the following. B1 and B2: Gevork Arakelian, Los Angeles County Department of Agriculture. B4: Mike Lewis, University of California, Riverside. D1, D2, D5, D7, and D13: Jack Kelly Clark, University of California Agriculture and Natural Resources. D10: Christoph Benisch, www.kerbtier.de. D14: Pavel Svihra, University of California. 11 Mesquite trunk: Tim Thibault. Leaf identification photos for 1, 2, 3, 6, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30: [SelectTree, selectree.calpoly.edu](http://selectree.calpoly.edu). Leaf identification photos 4, 7, 13, and 31: Keir Morse.

For Further Information

For additional photos and the full host list, as well as the most recent list of reproductive hosts, information, research, and news, see the University of California Agriculture and Natural Resources Invasive Shot Hole Borers website, www.pshb.org.

To order or obtain ANR publications and other products, visit the ANR Communication Services online catalog at <http://anrcatalog.ucanr.edu/> or phone 1-800-994-8849. You can also place orders by mail or request a printed catalog of our products from

University of California
Agriculture and Natural Resources
Communication Services
2801 Second Street
Davis, CA 95618

Telephone 1-800-994-8849
E-mail: anrcatalog@ucanr.edu

©2018 The Regents of the University of California. This work is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/> or send a letter to Creative Commons, PO Box 1866, Mountain View, CA 94042, USA.

Publication 8590

ISBN-13: 978-1-62711-020-4

The University of California, Division of Agriculture and Natural Resources (UC ANR) prohibits discrimination against or harassment of any person in any of its programs or activities on the basis of race, color, national origin, religion, sex, gender, gender expression, gender identity, pregnancy (which includes pregnancy, childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), genetic information (including family medical history), ancestry, marital status, age, sexual orientation, citizenship, status as a protected veteran or service in the uniformed services (as defined by the Uniformed Services Employment and Reemployment Rights Act of 1994 [USERRA]), as well as state military and naval service.

UC ANR policy prohibits retaliation against any employee or person in any of its programs or activities for bringing a complaint of discrimination or harassment. UC ANR policy also prohibits retaliation against a person who assists someone with a complaint of discrimination or harassment, or participates in any manner in an investigation or resolution of a complaint of discrimination or harassment. Retaliation includes threats, intimidation, reprisals, and/or adverse actions related to any of its programs or activities.

UC ANR is an Equal Opportunity/Affirmative Action Employer. All qualified applicants will receive consideration for employment and/or participation in any of its programs or activities without regard to race, color, religion, sex, national origin, disability, age or protected veteran status.

University policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's equal employment opportunity policies may be directed to: John Sims, Affirmative Action Contact and Title IX Officer, University of California, Agriculture and Natural Resources, 2801 Second Street, Davis, CA 95618, (530) 750-1397. Email: jsims@ucanr.edu. Website: http://ucanr.edu/sites/anrstaff/Diversity/Affirmative_Action/.

An electronic copy of this publication can be found at the ANR Communication Services catalog website, <http://anrcatalog.ucanr.edu/>.



This publication has been anonymously peer reviewed for technical accuracy by University of California scientists and other qualified professionals. This review process was managed by ANR Associate Editor for Pest Management—Agricultural Andrew Sutherland.

web-3/18-SB/CR