

NOVA ADDENDA

AD

MYCOLOGIAM VENETAM

AUCTORE

CAROLO SPEGAZZINI

Hymenomyceteæ Fr.

1. **Clavaria pistillaris** L., Saccardo Myc. Ven. Sp. p. 67.
Hab. In sylvula comitis Giustiniani prope Conegliano, aut. 1879.

Obs. Formam legi abnormem, subpyriformem, apice umbilicatam, longitudinaliter plicato-rugosam, cinereo-rubescens, alt. 14 cent., diam. 6 cent. An varietas?

Gasteromyceteæ Fr.

2. **Geaster mammosus** Fr., Vitt. Monog. Lycop. p. 165, tab. I. fig. IX.

Hab. In sylva alpina Cansiglio inter acus putrescentes Coniferarum, Aug. 1878.

Obs. In eadem sylva legi etiam *Geasterem fornictum* (Huds) Fr., *G. hygrometricum* Pers., et *Lycoperdon pyriformem* Schäff. ad caudices putrescentes Fagorum.

Hypodermeæ dBry.

3. **Ustilago receptaculorum** (DC) Tul. Sacc. Myc. Ven. Sp. p. 73.

Hab. In receptaculis florum Scorzonerae hispanicae in sylvula comitis Giustiniani, prope Conegliano, vere 1878 & 1879, rarissime.

4. **Polycystis? italica** S. & S. in litt. ad Schröt. Endogena; externe nunquam manifesta; sori fere totam seminis massam occupant, et pulvere compactinsculo, atro-brunneo obriunt: sporæ 3–6 conglobato–connatæ; una lateralis, magna, sphærica, hyalino–verruculosa, atro–fuliginea, 25–30 mm. diam., ceteræ (2–5) subglobosæ, glomerulatæ, 5–10 diam., pallide fuscae, leves, episporio crassiusculo donatae.

Hab. Intra semina Castaneæ vescæ, Conegliano 9 Novembri 1878.

Obs. An vera Ustilaginea sit, ignota germinatione, dubitandum.

5. **Uromyces Chenopodii** Thüm. in schedis. — Spieg. *Decades myc. Ital.* n. 30. (sub nom. *U. gigantei* Sp.)

Hab. In caulibus Schoberiæ maritimæ in Campo Martis, Venetiis, Martio 1879.

6. **Æcidium Compositarum** Mart. Fl. Erl. p. 314. Cook. Brit. Fung. II, p. 542 = f. *Hieracii murorum*.

Hab. In ejusdem foliis vivis in alpinis Cadubriæ a Zoldo, aest. 1877.

7. **Melampsora Betulæ** (KL) Dmz., Sacc. Myc. Ven. Sp. p. 77.

Hab. Ad folia viva Betulæ albæ a Belluno, Oct. 1878, rarissime.

Phycomycetæ dBy.

8. **Peronospora Cydarissiæ** PBy. Thuenen Myc. Univ. n. 45. Erb. Critt. Ital. Ser. II, n. 692.

Hab. In foliis Euphorbiæ Cyparissiæ a Belluno, Octobri 1878.

9. **Pilobolus Kleinii** v. Tiegh. Struct. Spor. Pilob. in Boll. d. I. Soc. Bot. de France, 1875.

Hab. In stercore humano putrescente, Conegliano, vere 1879 rarius.

10. **Cystopus candidus** (Pers.) dBy, Sacc. Mycol. Ven. p. 89. — forma *Sisymbrii officinalis* & forma *Arabidis hirsutæ*.

Hab. in eorumdem folijs a Conegliano, majo 1879.

11. **Cystopus cubicus** (Str.) dBy, Sacc. Mycol. Ven. p. 89. — forma *Tragopogonis pratensis*.

Hab. in ejusdem caulinibus, foliis floribusque a *Conegliano*, Majo 1878.

Ascomyceteæ dBy.

PYRENOMYCETEÆ Fr.

12. **Fracchiæa heterogenea** Sacc. Myc. Ven. Specimen p. 115. et Mycoh. Ven. n. 88.

Hab. In sarmenis putrescentibus Rubi fruticosi a Susegana, Majo, & Alni viridis a Belluno, Oct. 1878.

Obs. Ascii paraphysibus longioribus, crassiusculis, guttulatis, hyalinis obvallati, quæ sine jodi ope difficilime conspicuntur.

13. **Ceratostoma Venetum** Speg. Perithecia sub-carbonacea, majuscula, $\frac{1}{4}$ - $\frac{1}{3}$ -mill. diam., sparsa, superficialia, subglobosa in ostiole crassum fragile apicem versus molliusculum, desinentia; contextu peritheciorum parenchymatico-celluloso, ostioli autem prosenchymatico-fibroso; ascii (cito diffluentes) . . . : sporidia subnavicularia, inferne inaequilateraliter sub-mucronata, fuliginea, opaca, 6-9 \approx 5.

Hab. Ad putamina Coryli Avellanae putrescentia, Conegliano, Febr. 1879, rarissime.

14. **Amphisphæria culmicola** Sacc. Myc. Ven. Spec. p. 113. Fungi Ital. n. 444.

***A. melanomoides**.

Hab. In radicibus et culmis Arundinis Donacis putrescentibus Conegliano, Mart. 1879.

Obs. A typo recedit peritheciis globosis, superficiibus, papillulatis, dense gregariis; notæ fructificationis perfectissime tamen quadrant cum typo.

15. **Sphærella Coneglanensis** Sp. Perithecia primo subepidermica, dein erumpentia atque subsuperficialia, 120-150 micr. diam., laxe gregaria; contextus inferne subcarbonaceus, superne parenchymatico-polyedricus, fuligineo-ater. Ascii cylindracei vel cylindraceo-clavati,

anticee crassiuscule tunicati, 40–45 \approx 5, octospori, apaphysati. Sporidia didyma, loculo superiore majore, hyalina, 7–8 \approx $2\frac{1}{3}$ –3.

Hab. Ad folia emortua dejecta *Galyanthi floridi*, Mart. 1879, *Conegliano*.

16. **Sphærella nerviseda** Sp. Perithecia innata, sphæroidea, ostiolo perforato papillato donata, 100–120 micr. diam.; contextus minute parenchymaticus, ater. Ascii clavati, vel cylindraceo-clavati, apaphysati, 35–45 \approx 7–7 $\frac{1}{2}$, octospori. Sporidia oblongo-cylindracea, didyma quandoque curvula, ad septum vix constricta, hyalina, 15–20 \approx $2\frac{1}{2}$ –3, disticha vel oblique monosticha.

Hab. In nervis foliorum putrescentium, quae tornoso-reticulata evadunt, *Orobi verni*, *Conegliano*, vere 1879.

Obs. Cum *Sph. nemorosa* S. & S. in eadem planta vigente non comparanda.

17. **Sphærella Epicymatia** (Wallr.) Fckl. Symb. myc. p. 118. Karsten. Myc. Fenn. II. p. 96.

Hab. In thallo emortuo putrescente *Parmelia caperatae* a Bresséo (*Colli Euganei*) Febr., a Belluno et in *Sylva Cansiglio*, Aug. et Oct. 1878.

18. **Sphærella Lachesis** Sacc. Fung. Ven. Ser. II. p. 302. Spieg. Decades myc. Ital. n. 11.

Hab. In foliis emortuis putrescentibus *Hellebori viridis*, *Conegliano*, vere 1878.

19. **Sphærella prominula** Spieg. n. s. Peritheciis globulosis, primo tectis, dein subsuperficialibus, minutis, 100–120 micr. d., laxe gregariis v. sparsis, epiphyllis, atris; ascis cylindraceo-sinuositis crassiuscule tunicatis, mollibus, apaphysatis, octosporis, 70–75 \approx 8–10; sporidiis valde irregularibus, elliptico-oblongis, subimæquilateralibus, 1-septatis, ad septum valde constrictis, minute 2–8-guttulatis, hyalinis, 15–16 \approx 7–8.

Hab. ad folia emortua *Pteridis aquilinæ* a *Conegliano*, Majo 1878.

Obs. A *Sphærella Pteridis*, *S. aquilina*, *Didymella lophospora* etc. satis superque distincta.

20. **Gnomonia Veneta** Speg. Perithecia globulosa, laxe gregaria, primo parenchymati innata, tecta, dein emergentia, sicca cupulato-collapsa, $\frac{2}{10}$ - $\frac{3}{10}$ mill. diam., atra, rostro elongato, recto vel curvulo, crassiusculo, perithecium æquante vel paululum excedente, concolore instruita. Asci fusoideo-clavati, apice rotundati, crassiuscule tunicati, 45-50 \approx 7-8, octospori, aparaphysati. Sporidia elongato-fusoidea, quandoque sub-clavulata, utrinque obtusiuscula, excentrica 1-septata, h. e. loculo inferiore bis longiore quam superiore, hyalina, minute pluriguttulata, utrinque appendicibus rostriformibus, acutiusculis auctis, 15 \approx $3\frac{1}{2}$ -4,

Hab. Ad folia dejecta *Ostryæ carpinifoliae*, *Colfosco*, Martio 1879.

21. **Gnomonia vulgaris** Ces. & DNtrs. Karsten, *Myc. Fenn.* II. p. 124. Sacc. *Michelia* II. 245.

Hab. Ad folia dejecta putrescentia *Coryli Avelanæ*, *Conegliano*, vere 1879.

22. **Diaporthe** (Tetrast.) **Mahoniæ** Speg. Stroma ramos late ambiens per lignum immutatum tortuose et profunde excurrens, nigro-limitatum; peritheciis discretis, globosis, $\frac{1}{3}$ - $\frac{1}{4}$ mill. diam., ligno decorticato vel corticato totaliter vel semiimmersis, ostiolis brevibus, cylindraceo-conoideis, obtusis, exertis. Asci oblongo-fusoidei, 65-75 \approx 10, octospori, aparaphysati. Sporidia disticha vel oblique monosticha, elliptico-elongata, 12-14 \approx 4- $4\frac{1}{2}$, utrinque rotundata, medio constricto-1-septata, 2-6-guttulata, hyalina.

Hab. In ramulis emortuis *Mahoniæ Aquifolii*, *Conegliano*, 7 Mart. 1879.

23. **Diaporthe** (Euporthe) **Americana** Sp. Stroma hinc inde ramos ambiens, per lignum immutatum excurrens, nigro-limitatum; peritheciis 3-7 in eadem macula stromatica laxe-gregarie insidentibus, ligno infossis, $\frac{1}{6}$ - $\frac{1}{8}$ mill. diam., atris, ostiolo plus vel minus longo recto vel curvulo, cylindraceo, corticem perforante ornatis; contextu tenuissime membranaceo-parenchymatico, fuligineo. Asci oblongo-fusoidei, aparaphysati, 55-

60 = 7-8, granuloso-oloso fæcti. Sporidia immatura, non visa.

Hab. In ramulis emortuis Magnoliæ grandifloræ, *Susegana*, Oct. 1877.

24. **Diaporthe (E.) muralis** Sp. Stroma caulem late ambiens; peritheciis globosis, atris, ligno immutato immersis, discretis vel rarius subaggregatis, $\frac{1}{5}$ - $\frac{1}{6}$ mill. diam., ostiolo cylindraceo-conoideo, plus minusve exerto, nunc recto nunc obliquo ornatis. Ascii oblongo-fusoidei, 40-50 = 8-10, 8-spori, apophysati. Sporidia disticha vel oblique monosticha oblongo-ellipsoidæ, medio 1-septata atque constricta, utrinque obtusiuscule acuminata, 15 = 4 guttulata, hyalina.

Hab. In caulis emortuis Sedi albi, Patavii, Febr. (immatura), et S. reflexi, *Belluno* Oct. 1878.

25. **Diaporthe (Euporthe) occulta** (Fckl.) Nke, Fckl. Symb. 210.

Hab. ad strobilos putridos Cupressi semper-virentis a *Susegana*, Majo 1879, rarissime.

26. **Amphisphæria fallax** DNtrs Sfer. ital. p. 69 tab. 72. Rabh. F. E. n. 1935.

Hab. in cortice duriore Quercus pedunculatæ, in collibus di *Conegliano*, rarissime, Majo 1879.

27. **Leptosphæria nectrioides** Spig. Perithecia subcorticalia, $\frac{2}{10}$ - $\frac{3}{10}$ mill. diam., superne cortici tenaciter adhaerentia, atque eum circum circa paululum nigriscentia, sordide flavo-fuscescentia, ostiolo atro subcylindraceo epidermidem perforante donata; contextus molliusculus, tenui-membranaceus melleo-fuscidulus, ostiolum versus ater ac subcarbonaceus. Ascii fusoideo-clavati, 90-100 = 12-18 p. spor., apice rotundati, inferne in pedicellum crassiusculum, 15-20 = 5, elongati, paraphysibus paulo longioribus, filiformibus, guttulatis obvallati; sporidia disticha vel tristicha, fusoideo-elongata, 8-septata, ad septa constricta, curvula, loculo tertio superiore tumidiore, utrinque acutiuscule rotundata, fuligineo-olivacea, 35 = 6.

Hab. In sarmentis putridis Clematidis Vitalbæ in Monte *Cerva* prope *Belluno*, Oct. 1878.

28. **Leptosphaeria arvensis** Sp. Perithecia innata, subsphaeroidea, 100–150 micr. diam., parallele disposita, atra, ostiolo minuto, papillato, parum exerto, contextu fuligineo dense parenchymatico. Ascii cylindracei vel cylindraceo-clavati, 90–120 \approx 7–11, paraphysibus cerebrinatis, mucosis-obvallati, octospori. Sporidia elongato-fusoidea, disticha vel oblique monosticha, utrinque acutiuscula, 3–5-septata, loculis internis 1-guttulatis, flavidofuliginea, 30–35 \approx 5–6 $\frac{1}{2}$.

Hab. In caulis emortuis putrescentibus *Equiseti arvensis*, a *Conegliano*, Decemb. 1878.

29. **Leptosphaeria puccinoides** Speg. n. s. Peritheciis 2–3 gregariis v. solitariis, parenchymati immersis, macula fusca linearis ex innumeris hyphis mycelicis conflata insidentibus, nudoque oculo *Puccinia mæmulanti*-bus; ascis cylindraceo-clavatis, breve stipitatis, 60–65 \approx 7–8, aparaphysatis octosporis; sporidiis cylindraceo-oblongis, 5-septatis, loculo secundo superiore protuberante, ad septa vix constrictis, 18–20 \approx 3–4, pluriguttulatis, hyalinis.

Hab. ad folia dejecta *Caricis paludosaæ* a *Conegliano*, vere 1879.

30. **Sporormia ambigua** Niessl Die Arten Pyr. Spororm. in *Esterreich. Bot. Zeitsch.* p. 97, H. 2, Febr. 1874.

Hab. In fimo vaccino putrescente in montanis prope *Belluno*, Oct. 1878.

31. **Sporormia leptosphaerioides** Speg. Perithecia minuta, sparsa, subsphaeroidea, 120–140 mm. diam., ostiolo impresso perforata, contextu membranaceo parenchymatico-polyedrico atro-fuligineo. Ascii cylindracei, superne rotundati, p. sp. 100–110 \approx 10–15, inferne in pedicellum, 30–35 \approx 5, elongati, paraphysibus simplicibus filiformibus obvallati. Sporidia cylindraceo-fusoidea, tetramera, recta vel curvula, loculis duobus mediis cylindraceis, duobus extimis sub-conoideis, muco deficientia, atque loculis numquam secedentibus donata, 30–35 \approx 5–5 $\frac{1}{2}$, obscure fuligineo-olivacea.

Hab. In fimo ovinio vetusto putrescente in pratis circa *Conegliano*, aest. 1878.

32. **Melanomma canescens** Sp. Perithecia globosa vel subsphæroidea, minuscula $\frac{1}{4}$ - $\frac{1}{3}$ mill. diam., atra, sub-carbonacea, ostiolo papillato perforato donata, pilis sor-dide hyalinis, continuis vel 1-2-septatis, 100-200 \approx 5-7, undique laxe vestita (etiam in senectute?), ut oculo nudo cinereo-hirtella videntur. Asci cylindraceo-clavati, paraphysibus obvallati, octospori, 70-90 \approx 15-18. Sporidia disticha vel oblique monosticha, triseptata, ad septa non vel vix constricta, elliptica vel ovoideo-elliptica, quandoque subinaequilateralia, loculis 1-gut-tulatis, 20-25 \approx 10, primo hyalina, dein dilute fuligineo-virescentia.

Hab. Ad asseres abietinos putrescentes, Conegliano 21 Decembre 1878.

33. **Melanomma alpinum** Sp. Perithecia dense gregaria circulariter pustulatim erumpentia (more *Botryosphaeriae Bérengerianæ f. Rhamni Frangulæ*), minuta, $\frac{1}{3}$ - $\frac{1}{4}$ mill. diam., ovoidea vel subsphæroidea, papillulosa, atra, ostiolo minuto perforata. Asci cylindracei, vel leniter cylindraceo-clavati, 75-90 \approx 8-9, octospori, paraphysibus longioribus obvallati. Sporidia disticha vel sæpius oblique monosticha, primo hyalina 2-magniguttulata, dein 4-guttulata, postremo 1-septato-constricta, pallide olivaceo-fuliginea, 18 \approx 4.

Hab. Ad ramulos emortuos *Rhamni saxatilis*, *Rh. Frangulæ*, & *Hippophaëos rhamnoidis* in alpinis prope Belluno, Oct. 1878.

Obs. Species habitu *Botryosphaeriae Bérengerianæ* si-millima, sed notis fructificationis valde recedere videntur.

34. **Kalmusia dealbata** Sacc. Fung. Ven. ser. V. p. 179. Mycoth. Ven. n. 677.

Hab. Conegliano, in truncо decorticato adhuc vivo *Lyriodendri Tulipiferæ*, vere 1879.

35. **Aglaospora profusa** (Fr.) DNtrs., Sacc. Myc. Ven. Sp. p. 125. Mycoth. Ven. n. 98-650.

Hab. Vulgatissima in omnibus ramis truncisque emor-tuis *Robiniae pseud-acaciæ*, Conegliano, peren-nis, 1875-79.

Obs. Vere anni præteriti apud Coneglanum observavi

ad folia *Vitis viniferae* pelliculam tomentosulam atram nudo oculo *Capnodio* vel *Fumagini* simillimam; microscopio autem observata ab innumeris sporidiis *Aglaosporae profuse* composita videbatur. Haec sporidia, a palis Robiniaceis, quibus *Vitis maritata*, exsiliebant atque illapsa super folia, humiditatis foliorum causa, germinare incipiebant. Sporidia germinantia, ex uno vel pluribus loculis hypham, 100–150 \approx 3, emittebant hyalino-fuscidulam, quæ sub-dichotomice dividebatur atque apicibus conidia hyalina, elliptica vel ovoidea, minuta 4–5 \approx 1–2 $\frac{1}{2}$, gerebat. Quandoque invicem duo ramuli, ex eodem sporidiorum loculo orti, vertice conjugabantur, et corpusculum cylindrico-oblongum sinuoso-intestiniforme, 30–35 \approx 6–8, quandoque summitate sub-spiraliter revolutum efformabant.

SACCARDOELLA Speg. (nov. gen.) *Char.* Perithecia majuscula totaliter immersa, ostiolo papillato donata, carbonacea: asci cylindracei,—octospori, paraphysati; sporidia fusoideo-elongata, 20–30—septata, utrinque setigera.

36. **Saccardoella Montellica** Speg. Perithecia majuscula, $\frac{1}{2}$ – $\frac{3}{4}$ mill. diam., cortice immersa, ovato-sphaeroidea, superne in ostiolum acutiusculum vix exertum, perforatum desinentia, atro-carbonacea. Asci cylindracei, longissimi, 400–450 \approx 14–15, apice crasse tunicati, inferne in pedicellum breve ac robustum attenuati, paraphysibus filiformibus guttulatis, obvallati. Sporidia prægrandia, fusoideo-elongata 100–115 \approx 12, 20–30-septata, ad septa leniter constricta, loculis 3–5guttatis, utrinque acutiuscula, ac setula rigidula, 10–15 \approx 1 ornata, hyalina vel fuscidula.

Hab. Ad corticem vetustum *quercum* in sylva *Montello*, 3 Jan. 1879, rarissime.

37. **Pleospora Venturia** Sp. Perithecia superficialia, minutissima, 120–140 mmm. diam., sparsa, hypophylla, pilis rigidulis, atris, unicellulatoribus, 40–80 \approx 5–7, laxe undique vestita, contextu tenui-membranaceo parenchymatico, fuligineo-atro. Asci pauci, 5–7 in eod. perith., cylindraceo-clavati, crasse tunicati, 4–8spori, apa-

raphysati, 80–90 = 30–35. Sporidia disticha, majuscula, elliptica, utrinque obtusa, 3–5 horizontaliter, 2–4 verticaliter septata, ad septum medium constricta, primo flavida multiguttulata, dein brunneo-fuliginea 30–40 = 15.

Hab. Ad folia arida *Dryadis octopetalæ* in alveo *Plavis*, a Belluno, Oct. 1878.

38. **Pleospora malacospora** Speg. Perithecia innata; lenticulari-subsphaeroidea, $\frac{1}{5}\frac{1}{6}$ mmm. diam., ostiolo papillato protuberante exerto, contextus densissime parenchymatico fuligineo-atro. Ascii cylindraceo-clavati, apice crassissime tunicati, pauci, paraphysati(?), 100 = 16–18. Sporidia disticha, ellipsoidea irregulariter murali-septata, ad septa constrictiuscula, flavescentia, 22–24 = 10–11.

Hab. In charta stercorata putri in umbrosis prope Conegliano, vere 1879.

39. **Pleospora hyalospora** Speg. Perithecia hinc inde sparsa maculis albidis incidentia, subepidermica, lenticulari-sphaeroidea, $\frac{1}{3}\text{--}\frac{1}{2}$ mmm. diam., sicca cupulata, ostiolo minuto papillato, epidermidem perforante, donata. Ascii cylindraceo-clavati, crasse tunicati sed tenuissimi et fere diffluentes, paraphysisibus duplo longioribus obvallati. Sporidia elliptica, 3–horizontaliter, 1–verticaliter septata, ad septa, speciatim medio, constricta, perfecte hyalina, granuloso-nubilosa quandoque farcta, 24 = 12.

Hab. In sarmentis emortuis *Vitis viniferae* a Belluno, Oct. 1878.

Obs. Ob sporidia perfecte hyalina probabiliter ad novum genus trahenda species.

40. **Pleospora hispida** Niessl Notiz. ueb. n. k. Pyr. p. 193, tab. IV, f. 17.

Hab. In caulis emortuis putrescentibus *Lithospermum officinalis* a Conegliano, vere 1877.

41. **Pleospora Salsolæ** Fuck. Symb. myc. p. 131–Fung. Rh. n. 814. Speg. Dec. It. myc. n. 21.

Hab. In caulis emortuis putrescentibus *Salsolæ Kali* in Campo Martio, Venetiis, 25 Martio 1879.

42. **Rhaphidophora herpotricha** (Fr.) C. DNtrs Sche-me Sfer. 233. *Sph. herp.* Fr. DNtrs Mier. it. Dec. IX. n. 9. Perithecia sparsa vel laxe gregaria, in matrice saepius stromatico obscurata insidentia, majuscula $\frac{1}{2}$ – $\frac{3}{4}$ mill. diam., olivaceo-brunnecentia, ostiolo promi-nulo acutiusculo ornata, atque hyphis fuligineis, ana-stomosantibus undique tomentoso-vestita, contextu pa-rencymatico fuligineo densissimo. Ascii cylindrici, 140–160 \approx 10–12, paraphysati, 8–spori; sporidia bacilla-ria, 140–160 \approx 2–3, pluriguttata, hyalina.

Hab. In culmis putrescentibus *Triticum vulgaris*, Conegliano, Martio 1879.

43. **Pseudovalsa longipes** (Tul.) Sacc. Fung. Ven. Ser. IV. p. 20. *Melanconis longipes* Tul. Cke Hnd. 820.

Hab. Ad ramulos dejectos *Quercus sessiliflora*, Co-negliano, aut. 1877–78.

Obs. Ascii clavati, 8–spori, apice crassae tunicati; sporidia elongata curvula, 4–5–septata, non constricta fusca, 35–45 \approx 4–5.

HYPocreaceæ DNtrs.

44. **Nectriella villosula** Speg. Perithecia superficialia, sparsa minutissima, $\frac{1}{8}$ – $\frac{1}{6}$ mill. diam., flavescentia vel flavo-rufescens, villosa-tomentella, subsphaeroidea, ostiolo papillato ornata, contextus densissime et minute parenchymaticus, flavidus-dilute-ochraceus. Ascii cylindra-ceo-fusoidei, superne coarctati atque truncati, 45–50 \approx 4–5, paraphysati, octospori. Sporidia disticha, rarius monosticha, continua quandoque 1–3-guttulato-pseu-doseptata, hyalina, elliptica vel elliptico-elongata, 8–10 \approx $2\frac{1}{2}$ –3.

Hab. In ramulis emortuis Rosæ alpinæ in Monte Cerva prope Belluno, Oct. 1878, rarissime.

Obs. Ob sporidia quandoque 1–2–3–guttulato-pseu-doseptata ad *Calonectriam* vergit.

45. **Nectria Urceolus** Sp. Decades mycologicæ Italicae N. 16. Perithecia urceoliformia, gregaria, carneo-fla-

vescentia, minuta, superne applanata, ostiolo fuscidulo perforata, contextu tenuissime membranaceo, polyedrico-parenchymatico, dilute rosco-flavescens. Ascii cylindraceo-fusoidei, apice coaretati, praeципue in junioribus, atque tunica crassa bisveolata donati, 4-6-8-spori, 40-45 \approx 4-5, aparaphysati. Sporidia disticha rarius oblique monosticha, cylindracea, utrinque obtusiuscula, medio 1-septata et constricta, nubilosо-hyalina, 16-18 \approx 2-3.

Hab. In ramulis emortuis putrescentibus Rosæ Galliae in Monte *Cerva* prope *Belluno*, Oct. 1878.

46. **Nectria charticola** (Fuck.) Sacc. *Michelia* III, p. 289.
Fuck. *Symb. myc.* p. 176.

Hab. Conegliano, in charta putrescente stercoreata in umbrosis, Febr. 1879.

Obs. A *N. Pezicula* Sp. *Mich. II*, p. 232 facillime dignoscenda.

47. **Calonectria macrospora** S. & S. *Michelia* II, 251.
Fungi ital. N. 197.

* **C. Tarvisina** Speg. Perithecia flavo-rubentia, subglobosa, $1\frac{1}{4}$ - $1\frac{1}{3}$ mill. diam., superne intensius colorata, ostiolo subumbilicato perforata, non crasse tunicata; contextu densiusculo, flavo-rubescens, parenchymatico. Ascii clavati, paraphysisibus cerebrimis obvallati, 150-180 \approx 20. Sporidia fusoidea, quandoque curvula, utrinque acutiuscula rotundata, 7-septata, grosse 6-8-guttulata, ad septa constricta, 60 \approx 10-12, hyalina, senectute tamen pallide fuscescentia.

Hab. In ramulo emortuo putrescente Gleditschiaæ Triacanthi, a *Treviso*, 22 Novemb. 1878, rarissime.

Obs. A typo *C. macrosporæ* recedit peritheciis minoribus, superne non truncatis, minus crasse tunicatis, colore, habitatione etc.

48. **Aponectria inaurata** (B. & Br.) Sacc. *Michelia* III, p. 296. Cook. *Hndb.* p. 783.

Hab. In ramulis emortuis Ostryæ & Frangulæ in montanis prope *Belluno*, Oct. 1878.

Obs. I. Perithecia minuta, globosa, aurantiaco-cinnabarina, cæspitosa erumpentia. In eodem perithecio adsunt ascii spermatiophori, 75-80 \approx 10, aparaphysati,

50-100 sporidia minuta ovoidea vel elliptica, 3-1, hyalina foventia; ascii autem veri rariores, 8-spori, 70-80 = 7-8; sporidia elliptica hyalina spurie 1-septata, crassiuscule tunicata, utrinque obtusiuscula atque mucrone concolore acutiusculo aucta, 14-16 = 3-3 $\frac{1}{2}$.

Obs. II. Diu confusi cum *Chilonectria Cucurbitula*, in eodem ramulo concrecente, sed sedulo observata perithecia valde diversa videntur, nam minora, numquam cupulata, aurantiaco-cinnabarina nec rubella viridi-aureo tecta. An status *Chilonectriæ*? Etiam in *Pleonectria Lamyi* perithecia primo cupulata, rubella vi-ridi-aureo tecta, dein rufo-rubescentia etc. observavi.

DOThIDEACEÆ Nke.

49. **Dothidea insculpta** Wallr. Fuek. Symb. Myc. p. 223. Saccardo, Fung. Ven. p. V. p. 183. Mycoth. Ven. n. 641. Fungi italici N. 211.

Hab. In sarmenis emortuis putrescentibus Clematis Vitalbae, Conegliano, Belluno etc. autumno & vere 1876-79.

Obs. Omnia specimina quæ ipse legi a Conegliano & Belluno (nam species vulgatissima) valde differunt a descriptione Fuckelii et tabulis Saccardoi. Ascii 70-100 = 15-18, crassiuscule tunicati; sporidia bilocularia, loculo supero tumidiora, utrinque rotundata 20-25 = 6-8, diu hyalina sed tandem viridulo-fuliginea. Notis citatis exemplaria mea melius accedunt ad formam *Dothideæ Sambuci*, quam *D. insculpte*. An hæc postrema varietas *D. Sambuci*?

LOPHIOTOMEÆ Sacc.

50. **Lophiotrema? affine** Speg. Perithecia minutissima, 100-150 micr. diam., subepidermica, laxe gregaria, ostiolo compresso, tantum exerto, donata, contextu distinctissime oblongo—parenchymatico, atro. Ascii cylindracei, vel cylindraceo-clavati, apice obtusiusculi atque crassiuscule tunicati, 80-10,12, octospori. Sporidia pri-

mo 1-septata pluriguttulata utrinque acuta ac curvula, dein dilute fuscidula, 3-septata, ad septa constricta, recta, utrinque rotundata, 18 \approx 6 atque cauda, 3-4 \approx 1-1 $\frac{1}{2}$, parvula, rostriformi, acuta donata.

Hab. In ramulis emortuis Ampelopsis hederaeæ a Conegliano, aut. 1878.

51. **Lophiotrema culmifragum** Sp. Perithecia subglobosa, minuta, immersa, $\frac{1}{4}$ - $\frac{1}{3}$ mill. diam., parallele disposita, ostiolo tantum superficiem attingente, circum circa substratum parum nigrificantia. Asci cylindraceo-clavati, 80-90 \approx 8-10, paraphysibus filiformibus obvallati. Sporidia disticha primo constricto-1-septata, 4-guttulata, deinde 3-septato-constricta, curvula 4-6-guttulata, hyalina, 20-25 \approx 4-5.

Hab. In culmis putrescentibus Arundinis Donacis a Conegliano, Aug. 1878.

Obs. Cum *L. semilibero* & *L. Arundinis* in eadem matrice cohabitantibus non comparandum.

52. **Lophiotrema littorale** Speg. Perithecia laxe gregaria, $\frac{1}{4}$ - $\frac{1}{3}$ mill. diam., subsphaeroidea, ostiolo papillato compresso donata, basi tantum (rarius totaliter) ligno insculpta. Asci cylindraceo-clavati, breve-stipitati, crasse tunicati, 140-150 \approx 20-22, octospori, paraphysibus creberrimis ramulosis, guttulatis obvallati. Sporidia disticha, 5-septata, ad septa constricta, loculo tertio superiore protuberante, loculis grosse 1-2-guttulatis, hyalina, 35-40 \approx 12-14.

Hab. In ligno decorticato Salicis fragilis in insula dicta Lido apud Venetias, 25 Martio 1879.

Ob. Iodi ope asci et sporidia flavescunt; substantia antem, qua sporidia in ascis obvolvuntur, intense rubro-brunnescit et subopaca fit.

53. **Lophiostoma macrosporum** Speg. Perithecia $\frac{1}{2}$ -1 mill. diam., subsphaeroidea, basi ligno insculpta, ostiolo compresso acutiusculo donata, solitaria vel 3-5 laxe gregaria, carbonacea. Asci cylindraceo-clavati, crasse tunicati, breve pedicellati 300-350 \approx 40-50, octospori, paraphysati. Sporidia oblongo-fusiformia, utrinque acutiuscula atque hyalino-apiculato-caudata, primo hyali-

na, strato crasso obvoluta, medio 1-septato-constricta, 24–32 guttulis 1–3-seriatis donata, dein sub-fuliginea, loculo superiore intensius colorato, 12–16-pseudo-septata ibique paulum constricta 95–100 \approx 20.

Hab. In cortice vetusto dejecto *Quercus pseudorubra* in *Sylvula Giustinian* prope *Conegliano*, vere 1879, socio *L. vicino* & *L. Montellico*, cum quibus facillime confunditur, iis vero rarius reperitur.

54. **Lophiostoma collinum** Speg. Perithecia parenchymate foliorum immersa, laxissime gregaria, parallela, ostiolo compresso acutiusculo exerto, subglobosa, atra, 180–200 micr. diam. Ascii cylindraceo-clavati, superne rotundati, inferne in pedicellum brevem ac nodulosum desinentes, 150 \approx 20, paraphysibus longioribus filiformibus obvallati. Sporidia disticha, utrinque obtusiuscula, 5–7-septata, loculo superiore parum protuberante, ad septa leniter constricta, loculis raro 1–2-guttulatis, amœne flavescentia, 35–45 \approx 10–13.

Hab. Ad basim foliorum *Caricis piluliferae* a *Conegliano*, Martio, & C. patulæ a *Belluno*, Octob. 1878.

55. **Lophiostoma oreophilum** Speg. Perithecia globosa profunde in ligno immersa, ostiolo compresso apice superficiem attingente, circum circa matricem cyaneo-virescentem (fere ut in *Chrysosplenio æruginoso*) efficientia. Ascii cylindraceo-clavati, breve stipitati, 150 \approx 16–20, octospori, paraphysibus ramulosis, guttulatis obvallati. Sporidia disticha, elliptico-oblonga, utrinque acutiuscula, 5-septata, ad septa leniter medio evidentius, constricta, loculis 4 internis 1-guttulatis, amœne fuligineis, 2 extimis parvulis hyalinis, 30–35 \approx 8–10.

Hab. In ramulis emortuis putrescentibus *Corni sanguineæ*, in montanis prope *Belluno*, Oct. 1878.

Obs. Species profunditate peritheciorum, coloratione substrati (an semper?), apicibus sporidiorum hyalinis prædistincta! A *Loph. viridario* Cke diversa.

56. **Lophiostoma dumeti** Sacc. Fung. Ven. Ser. III. p. 3. Ejusd. Fung. it. N. 224.

Hab. In ramulis emortuis Fraxini excelsioris in montanis prope *Belluno*, Oct. 1878.

Obs. Ob sporidia perfecte cylindrica, parum quandoque curvata, ad septa nunquam constricta aliquantulum a typo recedit. In agro Bellunensi legi etiam; *L. macrostomoides* DNtrs in cortice populino, & *L. Nucula* in Salicino. A *Conegliano* autem in radicibus Tuniciæ *Saxifragæ* & cortice *Bignoniae Catalpæ* inveni *L. Anaxæum* Sacc.

HYSERIACEÆ Cda.

57. **Lophodermium xylomoides** Chev. Fuck. Symb. myc. p. 256. Sacc. Myc. Ven. S. p. 157. Speg. Decad. myc. Ital. N. 8.

Hab. Ad folia dejecta Crataegi Oxyacanthæ prope *Conegliano*, Martio 1879, rarissime.

58. **Lophodermium maculare** DNtrs., Fuck. Symb. myc. p. 251. Speg. Decad. myc. Ital. N. 9.

Hab. Ad folia dejecta Quercus pubescens, Rhamni saxatilis, Vaccinii Vitis-Ideæ in Monte *Cerva*, *Belluno*, Oct. 1879.

59. **Hysterographium hiascens** Rhem., Ascomyceten N. 314. Speg. Decad. myc. It. n. 7.

Hab. In cortice vetusto Quercum in *Sylva Montello* & *Giustinian*, sociis *Hysterio pulicari* & *vulgari*.

Obs. I. Singulare! Species vulgatissima sed ob suam faciem *H. pulicari* similem nondum detecta in Europa; distinguitur tamen facile etiam nudo oculo, peritheciis utrinque acutioribus, quandoque basi coaretatis, rima minus lata, atque defectu striarum longitudinalium. Specimina mea perfecte convenient cum Rhemianis tum notis externis tum fructificationis internis.

Obs. II. Legi quoque a *Conegliano* *Aylographum vagum* in foliis dejectis *Ilicis Aquifolii* vere, 1879.

ELAPHOMYCETEÆ *Tul.*

30. **Elaphomyces citrinus** Vitt. Monogr. Tub. p. 65, tab. IV, fig. XVI. Ejusd. Lycop. p. 214. tab. III. fig. XI. *Speg.* Decades myc. It. n. 5.

Hab. In terra calcareo-sabulosa inter radices Quercuum sub muscis in sylvula Comitis *Giustinian* prope *Conegliano*, vere 1879, rarius.

Obs. Odor debilis gratus. Fungus magnitudine pisi vel nucis Avellanæ; cortex levis brunneo-olivascens, thallo persistente floccoso late effuso amoene citrino-luteo obductus; peridium albido vel albido-cinereum crassiusculum; gleba atro-virescens; sporidia muriculata(?), 2-3guttulis magnis concentricis donata, fumose virescentibus, diam. 8-10 micr.

61. **Elaphomyces mutabilis** Vitt. *Vitt.* Monogr. Tub. p. 65, tab. IV, fig. XIV.; Ejusd. Lycop. p. 213.

***E. immutabilis** *Speg.* (Dec. mycol. Ital. n. 6 sub. *E. mutabili.*). Odor debilis, gratus. Fungus magnitudine pisi vel nucis avellanæ; cortex levis griseo-caerulescens, mollis, thallo persistente floccoso-fibroso albido late effuso obductus; peridium albidum; gleba virescens; sporidia muriculata(?), concentrice pluriguttulata, fuligineo-olivacea diam. 7-15 micromill.

Hab. Ad radices Quercuum sub humo non profunde delitescens in sylvula comitis *Giustinian* prope *Conegliano*, vere 1879; rarus.

Obs. A typo recedit peridio sub sectione non cya-nescente, floccis thallinis candidis nec cinereo-argentatis, odore etc.

62. **Elaphomyces Morettii** Vitt. Monogr. Tub. p. 71; tab. IV, fig. XVII. Ejusd. Lycop. p. 224.

Hab. In terra calcareo-sabulosa profunde delitescens in sylvula comitis *Giustinian*, vere 1879, rarissime.

Obs. Odor incertus fungino-aromaticus. Fungus magnitudine nucis avellanæ vel juglandis; cortex minute et obtuse verruculosus ater, fragilis; thallo non viso;

peridium albido-cinereum, crassum; gleba atro-virescens; sporidia sphæroidea, 2-4 concentrica guttulata, 25-30, flavo-brunnea, primo in ascis subglobosis, 8-sporis inclusa.

63. **Elaphomyces granulatus** Fries.

Hab. Inter radices quercum in sylvula *Giustinian* prope *Conegliano*, rarissime, vere 1878-79.

Obs. Odor debilissimus. Fungus magnitudine nucis avellanæ vel juglandis; cortex flavo-nigrescens, ex apice verrucarum flavidæ et basi nigrescente; verrucæ majusculæ; flocci thallini flavidæ-virescentes, fibrosi, rari atque apice ramulosi; peridium similare rubro-fuscescens; gleba coffeeicolor, primo flavo-albida variegata; sporidia subatra, levia, 15-25 diam.

TUBERACEÆ *Vitt.*

64. **Tuber brumale** Vitt. Monogr. Tub. p. 37. Sacc. Myc. Ven. Sp. p. 155. Speg. Decades myc. It. n. 1.

Hab. Circa radices Populorum ad ripas sabulosas torrentis *Monticano*, *Conegliano*, Dec. 1878. rarius.

65. **Tuber rufum** Pico Vitt. Monogr. Tub. p. 48. Sacc. Myc. Ven. Sp. p. 155.

Hab. Inter radices Populorum ad ripas sabulosas torrentis *Monticano* prope *Conegliano*, Dec. 1878, rarissime.

DISCOMYCETEÆ *Fr.*

66. **Pezicula Resinæ** Fuck. Symb. myc. p. 279.

Hab. In resina subputrescente Pini Strobi a *Conegliano*, Aprili 1879.

Obs. Ascis bene evolutos frustra quæsivi.

67. **Trochila Populorum** Dmz. Sacc. Michelia IV. p. 419. Speg. Decad. myc. It. n. 18.

f. Salicis Capreæ.

Hab. In ejusdem foliis dejectis putrescentibus, socia *Linospora Capreæ*, a *Conegliano*. Vere 1879.

Obs. Obturaculo vel apice ascorum jodi ope intense cærulescit.

68. **Stictis æcidiiformis** Speg. Ascomata æcidiiformia, 3–12 dense gregarie-stipata, vix $\frac{1}{10}$ – $\frac{1}{5}$ mill. diam. ceraceo-subdiaphana, tenaciussula, sordide luteolo-albida, semiimmersa, margine libero, stellatim laci-niato-lacero; contextus excipuli parenchymaticus, mi-nutus, e cellulis 2–4 micrm. diam. conflatus, dilutissime fuscidulus. Asci cylindraceo-clavati, brevissime stipitati, octospori, paraphysibus apice toruloso-clavulatis obval-lati, 65–85 \approx 8–9. Sporidia disticha vel subtristicha, cylindraceo-fusoidea, utrinque obtusiuscula, 7–10-sep-tata, ad septa plus minusve constricta, 30–35 \approx 4–4 $\frac{1}{2}$, hyalina.

Hab. Parasitica in stromatibus *Diatrypellæ verruci-formis* & *Diaporthis Mahoniæ*, in ramulis emortuis de-jectis *Mahoniæ Aquifolii*, Conegliano 7 April. 1879.

Obs. Apex ascorum et paraphysum jodi ope dilute cærulescit.

69. **Dermatea stegioides** Speg. n. s. Ascomatibus pri-mo subcutaneis, clausis, dein epidermide operculatim circumscissa sed permanente subvelatis, disciformibus, extus cinnamomeis, granuloso-tomentellis, disco cinereo v. subroseo, concavo, prominulo, $\frac{1}{2}$ –1 mill. d.; ascis cylindraceis paraphysibus filiformibus obvallatis, 35–40 \approx 2–3, octosporis; sporidiis distichis, cylindraceis v. cla-vulatis, non v. tenuissime 1-septatis, ad septum non constrictis, hyalinis, 5–6 \approx 1–1 $\frac{1}{2}$.

Hab. Ad ramulos emortuos *Quercus sessili-floræ a Colfosco*, Majo 1879, rarissime.

70. **Schyzoxylon Berkeleyanum** (Dub. & Lév.) Fuck. Symb. myc. p. 251. *Stictis decipiens* Karsten. Myc. Fenn. I. p. 240.

Hab. In sarmentis putrescentibus *Vitis viniferæ a Belluno*, Oct. 1878. & *Rosæ caninæ a Coneglia-no*, Mart. 1879.

Obs. Asci cylindracei 230–260 \approx 10–12, paraphy-sibus filiformibus longioribus obvallati; sporidia ascorum

longitudine, septulata, in articulos, 6–10 ≈ 2, secedentia, hyalina.

71. **Pyrenopeziza lichenicola** Speg. Ascomata laxe gregaria, minutissima, vix $\frac{1}{10}$ – $\frac{1}{8}$ mill. diam., atra epithecio dilutiore, sicca subsphaeroidea, humida patellaria: contextu excipuli densissime parenchymatico atro. Ascii clavato-elliptici, erasse tunicati, apophysati (?), octospori, iodi ope apice intense cærulecentes, 30–35 ≈ 12–15. Sporidia didyma, primo muco obvoluta, dein libera, loculo superiore crassiore, hyalina, ad septum constricta 6–8 ≈ 3–4.

Hab. In thallo emortuo *Parmelia caperatae* in Monte *Cerva* prope *Belluno*, Oct. 1878.

72. **Niptera cinerea** (Batsch) Fuck., Sacc. Michelia IV. p. 424.

***Niptera exsiliens** Speg. Ascomata sessilia vel brevissime pedicellata, solitaria vel gregaria, hinc inde per epidermidem ramulorum erumpentia, extus atrocærulea, epithecio pallidiore griseo, linea candida marginato, magnitudine valde ludentia, $\frac{1}{4}$ –1 mill. diam., Contextus parenchymaticus, marginem versus prosenchymaticus, densissimus, membranaceus. Ascii cylindraceo-subclavati, apice obtusiusculi, in pedicellum crassum desinentes, 55–65 ≈ 5–7, paraphysibus guttulatis crassiusculis, obvallati, octospori. Sporidia clavulata, medio uni-septata, ad septum non vel vix constricta, hyalina 10–15 ≈ 2–3.

Hab. In ramulis emortuis dejectis Quercuum, quandoque parasitica in *Colpomate quercino*, Conegliano, vere 1879.

73. **Ciboria firma** (Pers.) Fuck. Sacc. Michelia IV, p. 428. Karsten, Mycol. Fenn. I. p. 108.

Hab. In ramulis emortuis Quercum ad ripas Monticani, Conegliano, Novemb. 1878, rarissime.

Obs. Ascorum obturaculum iodi ope intense cærulescit.

74. **Lachnella corticalis** Fr., Karsten Myc. Fen. I. p. 159 (sub Helotio).

Hab. Ad corticem Quercum vetustum in M. *Cerva* prope *Belluno*, et a Conegliano, per annum, 1877–79.

Obs. Species facillime dignoscenda ob amoenitatem colorationis; epithecio roseo, purpureo, vel cinnabarinoc-aurantiaco; excipulo vivide cæruleo, vel cinereo-cæruleo; magn. $\frac{1}{2}$ a $1\frac{1}{2}$ mill. diam.

75. **Dasyphypha bicolor** (Bull.) Fckl. Symb. Mycol. 305.

Hab. Ad ramulos emortuos *Quercus* pedunculatae, a *Conegliano*, 8 April. 1879, rarius.

Obs. I. Species distinctissima et pulchella. Cupula sicca subsphæroideo-contracta, humida planiuscula, epithecio amœne luteo-aurantiaco vel cinnabariuo, extus candida vel dilutissime rosea, pilosa, stipite nullo vel brevissimo suffulta. Asci cylindracei, breve stipitati, octospori, $50 \approx 3\frac{1}{2}-4\frac{1}{2}$, obturaculo minutissimo jodo cærulecente, paraphysibus longioribus oblongo-lanciformibus crassis obvallati, superne acutissimi, $60 \approx 5-5\frac{1}{2}$. Sporidia elongato-fusoidea, vel ovoideo-elongata, hyalina, disticha vel oblique monosticha, $6-8 \approx 1\frac{1}{2}$. Cellulæ excipuli dilutissime fuscidulæ, ellipticæ $8 \approx 5$; pili cylindracei subechinulati, hinc illinc septato-constricti, 100-200 ≈ 5 .

76. **Helotium collinum** Speg. n. s. Ascomatibus laxe gregariis primo subglobosis dein explanato-patellaribus, sessilibus, extus glabris, sordide virescentibus, disco amœne griseo-cæruleo, subhygrophano, diam. $\frac{1}{4}-\frac{1}{2}$ mill.; ascis cylindraceis, a paraphysatis, 45-50 $\approx 4-5$, obturaculo vel apice jodi ope dilute cærulecente; sporidiis distichis cylindraceo-fusoideis, subclavulatisque, quandoque utrinque uniguttulatis, hyalinis, $10 \approx 1-1\frac{1}{2}$.

Hab. In pagina inferiore foliorum *Viburni Lantanae* a *Conegliano*, Mayo 1879.

77. **Calloria? pusilla** Speg. Ascomata sparsa, ephiphylla, primo epidermide tecta, dein erumpentia eaque fissâ cincta, subsuperficialia, minutissima, 120-150 micr. diam., pallide roseo-aurantiaca, sicca cupulato-concava, humida subhemisphaerica, prominula; contextus excipuli parenchymaticus, irregulariter polyedricus, tenui-membranaceus, fuscidulus. Asci cylindraceo-clavati, apice rotundati, atque tunica crassa donati, $60 \approx 8$, octospori, paraphysibus filiformibus apice leniter incrassatis ac un-

cinato-curvatis obvallati. Sporidia disticha vel oblique monosticha, elliptica $10 \approx 4$, hyalina.

Hab. In foliis dejectis putrescentibus Corni sanguineæ, a Conegliano, Apr. 1879.

Obs. Sæpius ascomata in maculis latis, vagis non circumscriptis incident.

78. **Ascobolus stictoideus** Speg. Ascomata laxissime gregaria vel sparsa, minuta, vix $\frac{1}{3}-\frac{1}{2}$ mill. diam., externe aquose albida, epithecio dilute olivaceo ex ascis exilientibus nigro-punctato, totaliter in fimo immersa, non vel labiis tantum exertis, crassiusculis atque laceris. Ascii pauci 5-10, magni 150-160 \approx 40, clavato-saccati, breviter ac crasse stipitati, superne crassiuscule tunicati, octospori, paraphysibus septulatis apice curvulis obvallati. Sporidia disticha vel irregulariter polysticha, elliptica, 25 \approx 14, primo hyalina, levia, dein amœna violacea, minute ac densissime verruculosa, postremo roseo-fuliginea.

Hab. In fimo canino putrescente, Conegliano April. 1879.

Obs. Tunica ascorum jodi ope obsolete cærulescit.

79. **Ryparobius saccoboloides** Speg. Ascomata dense gregaria, minutissima, 80-100 mil. diam., difficillime nudo oculo perspicienda, sordide roseo-fuscescentia, epithecio obscuriore virescente, primo globosa, hemisphærica, subsuperficialia; contextus excipuli minutissime ac densissime parenchymaticus, dilute roseo-fuscidulus. Ascii ovoidei vel elliptico-clavati, superne obtusi, inferne in pedicellum crassum ac nodulosum desinentes, 40-50 \approx 20-25, 60-120-sori, aparaphysati. Sporidia in glomerulo 35-40 \approx 15-20 elliptico, tenaciter muco conexa, etiam extra ascos, elliptica vel ovoidea, utrinque obtusiuscula, continua, hyalina, luce refracta dilute chlorina, crassiuscule tunicata, 5-6 \approx $2\frac{1}{2}-3$.

Hab. In fimo canino vetusto putrescente, a Conegliano, Apr. 1879.

Obs. Iodi ope ascii et sporidia rubro-fuliginea evadunt, stipites et bases ascorum autem ita colorantur ut opaci fiant.

Fungi imperfecti.

80. **Ramularia microspora** Thm. in Oest. bot. Zeit. 1877. p. 272. Mycoth. Univ. N. 982.
Hab. In hypophyllo *Teucrii Chamædryos*, *a Conegliano*, aest. 1877, rarissime.
81. **Ramularia Weigeliæ** Speg. Maculis subnullis, vagis. Acervulis hypharum hypophyllis super vel circa perithecia cujusdam *Phyllostictæ vegetantibus*, dense gregariis. Hyphis hyalinis, non vel rarius 1-2-septatis, apice nodulosis, 30-40 \approx 2-3; conidia hyalina magnitudine et forma ludentia, 5-15 \approx 3-4, subrotunda vel oblongo-cylindracea, quandoque continua, quandoque uniseptata, basi subapiculata.
Hab. In pagina inferiore foliorum *Weigeliæ roseæ a Conegliano*, aut. 1878.
82. **Ramularia Bellunensis** Speg. Maculis sordide cinerascentibus, late caules et folia ambientibus; hyphis subtortuosis, non vel rarius 1-2-septatis, apice incrassatis, hyalinis, 40-50 \approx 5; conidiis ellipticis vel elliptico-cylindraceis, utrinque obtusiusculis, medio uniseptatis ad septum non vel vix constrictis, hyalinis, 25-30 \times 5-6.
Hab. Ad folia & caules *Chrysanthemi Parthenii a Belluno*, Oct. 1878, vulgatissima.
83. **Cylindrosporium Valerianæ** Speg. Maculis magnis elongatis vel subrotundis griseis; hyphis brevibus, gracilibus, 50 \approx 3-4, hyalinis, nubiloso-granulosis, non vel 1-2-septatis, vertice irregularibus. Conidiis cylindraceis vel elliptico-elongatis, quandoque simplicibus, sed saepius 1-3-septatis, hyalino-nubilosis 15-50 \approx 5-8.
Hab. In hypophyllo *Valerianæ officinalis*, & dioicæ circa *Conegliano*, aut. 1877-78.
84. **Cercospora Bolleana** (Thm.) Speg. *Septosporium Bolleanum* Thuemen, in Oest. bot. Zeitschr. 1877. p. 12; & Contrib. micolog. del Litorale p. 10.
Hab. Ad folia viva vel languida *Fici Caricæ*, vulgatissima *a Belluno*, Oct. 1878.
85. **Cercospora Mercurialis** Pass. in Rabh. F. Eur.

Maculis argenteo-albidis, rotundis; cæspitibus hypophyllis gregariis centrum macularum sæpius occupantibus; hyphis dilute fuligineis, continuis, contortis, nodulosis, brevibus, 20–40 \approx 5–6; conidiis cylindraceo-bacillaribus, inferne attenuatis, atque acutiuscule rotundatis, antice crassiusculis et obtusis, 2–7-septatis, crasse tunicatis, hyalinis, 40–70 \approx 4–6, rectis vel curvulis.

Hab. Ad folia viva vel languida *Mercurialis perennis* in collibus *di Conegliano*, rarius, aut. 1877.

86. **Cladosporium aecidiicola** Thüm. Mycoh. Univ. n. 373.

Hab. In *Æcidio Euphorbiæ* parasitans ad folia *Euphorbiæ palustris* *a Conegliano*, Majo 1879, rassisime.

87. **Geotrichum candidum** Lk. Nees r. Esemb. Syst. d. Pilz. I. p. 29.

Hab. In ossibus putrescentibus *a Conegliano*, aut. 1876.

Obs. Conidia cylindracea utrinque truncata 5–10 \approx 4 hyalina.

88. **Graphium aspergilloides** Speg. Stipites cylindracei, longiusculi, 150–200 \approx 20–30, atro-fuliginei, compositi, apice ex hyphis undique divergentibus subgloboso-capitati; hyphæ fasciculatæ, fuligineæ, fertiles in apicem nodulosum, denticulatum desinentes; conidia e denticulis hypharum oriunda, elliptico-ovoidea vel oblongo-elliptica, 6–10 \approx 3–4, dilute fuliginea.

Hab. In pagina inferiore foliorum putrescentium *Sorbi Torminalis* *a Conegliano*, vere 1878 & 1879, rarius.

89. **Isaria stilbiformis** Speg. Gregaria vel solitaria; stipites flavescentes, 3–7 mill. alti, compositi ex hyphis fasciculatis, subcontinuis, flavidо-hyalinis, tenaciter coailitis, sursum apice divergentibus et capitulum albidum-tomentosulum subsphæroideum vel pyriforme efformantibus; apice hypharum perfecte hyalino, clavulato, vertice acutissimo atque sporidium cylindraceum utrinque acutiusculum, 8–11 \approx 2–2½, hyalinum gerente.

Hab. Ad corpora putrescentia cuiusdam minutæ spe-

- ciei *Pentatomorum* in carpineto prope *Belluno*, Oct. 1878, rarissime.
90. **Isaria arachnophila** Ditm. Deutsch. Crypt. Flor. Heft 4. tabula 55.
Hab. Ad corpora emortua *Epeiræ tubulosæ, conicæ, Drassi atri & Mygales cæmentariæ, a Belluno*, Oct. 1878, sporadice non rarius.
91. **Isaria farinosa** Tul. Fuck. Symb. myc. p. 186.
Hab. Ad corpora emortua chrysalidum *Lepido-pterorum nocturnorum a Conegliano & Belluno*. Vul-gatissima aut. 1878.
92. **Arthrinium curvatum** Kze & Schm. Fuck. Symb. myc. p. 357. Speg. Decad. myc. It. n. 28.
Hab. Ad folia emortua subputrescentia *Caricis paludosæ a Coll' Albrigo prope Conegliano*, Mar-tio 1879.
93. **Melanconium betulinum** Schm. & Kze. Fuck. Symb. myc. p. 190.
Hab. In ramis emortuis *Betulæ albæ a Belluno*, Oct. 1878.
Obs. Conidia ovata, crassiusculæ tunicata, granulosofarcta, fuliginea, $15-18 \approx 6-7\frac{1}{2}$.
94. **Gloeosporium** (Eugloespor.) **Venetum** Speg. Maculæ marginales, amorphæ, maximæ, vel centrales rotundæ parvulæ, melleæ vel ochraceæ linea fusco-purpurascente cinctæ. Acervuli minimi, prominuli, gregarii vel solitarii, atri. Conidia elliptico-cylindracea, $7-8 \approx 2-2\frac{1}{2}$, granulosa et guttulata, hyalina.
Hab. Ad folia viva vel languida *Rubi Chamæmorii a Belluno & a Conegliano*, Aut. 1877-78.
95. **Æcidiolum minutellum** Speg. Acervuli primo sub-epidermici, minimi, $\frac{1}{10}-\frac{1}{8}$ mill. diam., gregarii, dein erumpentes, læte aurantiaci. Conidia catenulata, elliptica, vel ovoidea, rarius, subsphærica, hyalina $3-4 \approx 1-2$, sterigmatibus crassiusculis, $45-50 \approx 1-2\frac{1}{2}$, subclavulatis, quandoque 1-2-septulatis suffulta.
Hab. In foliis *Scorzoneræ hispanicæ* socia *Puccinia Compositarum*, cuius videtur status microco-nidicus.

96. **Sporidesmium echinulatum** Speg. Acervuli superficiales, rotundi, atri, minuti, gregarii, quandoque plures confluentes; conidia elliptica, 30–32 ≈ 15–16, 3-septata, ad septa plus minusve constricta, septulo uno alterove longitudinali, ubique laxe et minute verruculosa fuligineo-olivacea, loculis quandoque duobus extimis pallidioribus, pedicellis hyalinis, 5–8 ≈ 2–3, suffultis.
Hab. Ad chartam putrem stercoratam in locis umbrosis prope *Conegliano*, aut. 1876–78.
97. **Hormococcus hysteroides** Speg. n. sp. Glomerulus ligno dealbato innatis, oblongo-hysteriformibus, laxe gregariis atris, parallelis, long. 1–1 $\frac{1}{2}$ mill., crass. $\frac{1}{4}$ – $\frac{1}{3}$ mill.; sporophoris dense fasciculatis, filiformibus, ramulosis, septulatis, 40–50 ≈ 2–3, fuscidulis, apice spermatia catenulata, cylindracea, utrinque truncata, 3–8 ≈ 2–2 $\frac{1}{2}$, fuscidula gerentibus.
Hab. In sarmenis putridis Rubi fruticosi a *Conegliano*, Majo 1879.
98. **Myrothecium inundatum** Tode, Fckl. Symb. p. 364.
Hab. In Cortinariis putrescentibus a Belluno, Oct. 1879.
99. **Antromyces Copridis** Fres. Beitr. p. 37. t. IV. fig. 40–45.
Hab. In terra humosa pingui (ad chrysalides?) in castaneto prope *Conegliano*, aut. 1876.
100. **Pestalozzia longiseta** Speg. Acervuli punctiformes, lenticulari–subsphaeroidei, primo tecti dein erumpentes, atri, 100–150 micrm. diam., macula rufa nigro-cincta insidentes. Conidia elliptico-fusoidea, utrinque acuminata, 20 ≈ 8, 4-septata, ad septa non vel vix constricta, loculis tribus internis fuligineo-atris, quandoque 2 superioribus intensius coloratis, duobus autem extimis parvulis atque hyalinis, superiore tribus rostellis longissimis, 30–40 ≈ 1, ornato, inferiore pedicellum, 5–15 ≈ 1 gerentibus.
Hab. Ad folia viva vel languida Rubi cæsii a *Susegana*, Novemb. 1876.
101. **Pestalozzia funerea** Dmz. Sacc. Myc. Ven. Sp. p. 198. Sequentes formas vel varietates ipse legi:

- a) *typica*. Conidia 4-septata, loculis tribus interioribus atro-olivaceis, $30-35 \approx 7-8$; rostell. 3, 15 ≈ 1 ; stip. 5-10 ≈ 1 .
- Hab.* In foliis vivis vel emortuis *Pteridis aquilinæ*, *Photiniæ serrulatae*, *Eucalypti Steweniani*, *Rubi fruticosi*, *Evonymi chinensis*, *Thujæ & Cupressi* etc.
- b) *crassipes*. Conidia oblongo-fusoidea, 4-septata, 3 loculis internis atro-olivaceis, $20-30 \approx 7$; rost. 3, 10-12 ≈ 1 ; stip. 5-10 $\approx 3-3\frac{1}{2}$.
- Hab.* In ramulis emortuis *Lonicerae Caprifolii a Conegliano*, aut. 1876.
- c) *multiseta*. Conidia elliptico-elongata, utrinque acuta, 3 loculis internis atro-olivaceis, $25-30 \approx 9-10$; rost. 5-10, 7-10 ≈ 1 ; stip. 5-7 ≈ 1 .
- Hab.* In foliis dejectis *Iridis Germanicæ*, *Conegliano*, vere 1877.
- d) *discolor*. Conidia oblongo-fusoidea, 4-septata, 3 loculis internis olivaceis, loculo medio intensius colorato, $25-30 \approx 7-8$; rost. 4-5, 15-20 ≈ 1 ; stip. 8-10 ≈ 1 .
- Hab.* In foliis emortuis *Buxi sempervirentis*, & *Serofulariæ caninæ a Tarzo prope Conegliano*, aut. 1872.
02. **Pestalozzia strobilicola** Speg. Acervuli primo lenticales tecti, dein conoidei, erumpentes, $\frac{1}{4}-\frac{1}{3}$ mill. diam., atri, gregarii, rarius sparsi; conidia elliptico-fusoidea, utrinque acuminata, 3-septata, $25 \approx 9$, loculis 2 internis atro-fuligineis, 2 extimis autem hyalinis, superiore 3-4 rostellis, $22-30 \approx 1$ ornato, inferiore pedicellum, 5-10 ≈ 1 , gerente.
- Hab.* Ad strobilos putridos *Pini sylvestris*, a *Susegana*, 17 Novemb. 1876.
03. **Pestalozzia versicolor** Speg. Acervuli globoso-lenticulares, primo subepidermici, dein erumpentes, et circum circa ex conidiis exslientibus matricem atro-inquinati; conidia fusoidea 4-septata, $30 \approx 10$, loculis 2 extimis hyalinis, 2 internis superioribus atro-olivaceis, tertio inferiore amœne flavo-virescente, 3-guttulatis; loculo supremo conoideo, 3-4 rostella,

25—30 = 1, gerente, infimo inflatulo quandoque 1—guttulato in pedicellum, 5—8 = 1, desinente.

Hab. Ad folia dejecta putrescentia *Nerii Oleandri*, *a Conegliano*, Novemb. 1876.

104. **P. monochæta** Dmz. Sacc. Myc. Spec. p. 198.

Hab. Ad folia dejecta putrescentia *Quercuum*, *a Conegliano*, aut. 1876—78.

***P. Saccardoi** Speg. Acervuli parenchymate immersi, punctiformes, densiuscule gregarii; conidia 4—septata, 20 = 5, loculis 3 internis atro—opacis; 2 extimis hyalinis; superiore acute conoideo, majusculo rostellum e basi curvato uncinato hyal. 10—15 = 1½, gerente, inferiore in pedicellum, 30—40 = 2, desinente.

Hab. Ad folia dejecta putrescentia *Quercus*, *a Conegliano & in Silva Montello*, aut. 1876-77.

105. **Pestalozzia macrospora** Ces. Sacc. Fung. Ven. Series. V. p. 198.

Hab. In frondibus emortuis *Pteridis aquilinæ* *a Conegliano*, aut. 1877-78, vulgata.

***P. macropoda** Speg. Acervuli epiphylli primo tecti dein erumpentes, conoidei, minutissimi, cinereo—fuliginei; conidia fusiformia, 35—38 = 7, 4—septata, ad septa leniter constricta, loculis 3 interioribus olivaceis, 2 extimis hyalinis, superiore rostellum unicum obliquum vel rectum, 15 = 1—1½, apice abrupte ac brevissime recurvato—uncinatum gerente, inferiore in pedicellum, 45—50 = 1, desinente.

Hab. In frondibus emortuis *Pteridis aquilinæ* *a Conegliano*, vere 1877, rarissime.

106. **Pestalozza lignicola** Cooke; an *P. truncatula* Fuck.?

Hab. In ramlis emortuis *Rosæ caninæ*, & in trabe abietina putrescente *a Conegliano*, aut. 1877-78.

Obs. Conidia 3—septata, loculis 2 interioribus primo flavidis, dein atro—olivaceis, 2 extimis hyalinis, 12—18 = 4—7, loculo supremo 2 rostellis, 20—25 = 1 ornato, inferiore in pedicellum, 20—30 = 1, desinente.

107. **Dinemasporium rhodophæum** Sp. Cupulæ parenchymate immersæ, disco superficiem non vel vix (hu-

miditate) excedente, numquam tectæ, laxe gregariæ, subparallelæ, macula nulla vel subnulla insidentes, discoidæ sed ob pressionem nervorum folii sub-quadratæ, minutæ $\frac{1}{4}$ — $\frac{1}{3}$ mill. diam. Excipuli contextus parenchymaticus, fuligineus, margine aterrimo, nitente, atque pilis brevissimis, 20—40 \approx 5, concoloribus, unicellularibus, laxe ornato; disco amoëne roseo vel roseo-aurantiaco. Spermatia elliptica, recta, biguttulata, 9 \approx 3, hyalina, utrinque setulâ rigidulâ subæquilongâ, rectâ aucta.

Hab. In foliis emortuis diu intemperiis expositis *Phragmitis communis* in insula dicta *Lido* apud Venetias, Martio 1879.

108. **Neottiospora coprophila** Speg. Perithecia minutissima, 80-70, globosa, astoma(?), irregulariter dehiscentia, contextu densiusculo, parenchymatico-indistincto; spermatia oblongo-fusoidea, 25 \approx 3, utrinque acutiuscula, nubilosso-granulosso-farcta, hyalina sessilia, apice tribus rostellis exilissimis, 20 \approx 1, ornata.

Hab. In fimo ovino vetusto in pratis circa *Conegliano*, socia *Delitschia Winteri*, Decemb. 1878.

109. **Phoma (Diaporthe) Volkameriæ** Speg. Peritheciis astomis(?) epidermide nigrificata, atque tumefacta, nitente, tectis, globoso-lenticularibus, 150-200; spermatiis hyalinis, ellipticis, utrinque attenuatis obtusiusculis, non vel 1—2—guttulatis, 7—11 \approx 3—5.

Hab. In ramulis emortuis putrescentibus Volkameriæ fragrantis, a *Conegliano*, Aug. 1876.

110. **Phoma Weigeliæ** Speg. Peritheciis lenticularibus 110-120, diam. sub epidermide nidulantibus, carbonaceis, ostiolo papillato-conoideo acutiusculo donatis. Spermatiis minutis, hyalino-chlorinis, 3—5 \approx 1—2 ellipticis vel ovoideis, obtusis, non guttulatis.

Hab. In ramulis emortuis Weigeliæ roseæ a *Conegliano*, aut. 1876.

111. **Phoma (Diaporthe?) Rosmarini** Speg. Peritheciis superficialibus, ligno circumnigrificato decorticato adnatis, lenticulari-hemisphaericis, atris rugulosis, 2—3 stromatice aggregatis, 200—250 micr. diam. carbona-

ceis. Spermatiis ovoideo-fusoideis, inferne acutiusculis, antice obtusis, hyalinis, granuloso-farctis, minute 2-guttulatis, 6-10 \approx 2-3.

Hab. In ramulis emortuis Rosmarini officinalis a Conegliano, Dec. 1876.

112. **Phoma (Diaporthe) Lagerstrœmiæ** Speg. Peritheciis ligno stromatico nigrificato immersis, 110-150 micr. diam., ostiolo pertuso, papillato donatis. Spermatiis ovoideo-ellipticis, utrinque obtusiusculis, hyalinis, 2-guttulatis, 7-10 \approx 2-3.

Hab. In ramulis emortuis Lagerstrœmiæ indicæ a Conegliano, Dec. 1876.

113. **Phoma (Diaporthe) Anaxæa** Speg. Stromate ramosos corticatos vel decorticatos late ambiente atque lignum atro-nitens efficiente. Peritheciis ligno immersis, atris, 150-180 micr. diam.; spermatiis minutissimis hyalino-chlorinosis, subellipticis, utrinque obtusiusculis, 2-3 \approx 1-1 $\frac{1}{2}$.

Hab. In ramulis emortuis Onobrychis sativæ in alveo *Plavis* (Anaxi), Aug. 1876.

114. **Phoma (Diaporthe) Nerii** Speg. Stromate late ambiente, lignum extus nigrificante. Peritheciis minutis ligno corticato vel decorticato immersis, atris. Spermatiis ellipticis vel oblongo-ellipticis, 6-9 \approx 3-4, 2-guttulatis, hyalinis.

Hab. In ramulis emortuis Nerii Oleandri a Conegliano, vere 1878.

115. **Phoma endogena** Speg. Peritheciis in membrana cotyledones obvolvente nidulantibus, griseis, quandoque subtomentellis, lenticulari-sphæroideis, tenuemembranaceis, contextu parenchymatico-celluloso, 150-200. Spermatiis hyalinis, granuloso-nubilosso-farctis, ovoideis vel ellipticis, 6-8 \approx 2-3, eguttulatis.

Hab. In seminibus maturis Castaneæ vescae corice integro vestitis, adhuc vegetis (socia quandoque *Mycogone cervina* Lk, teste ipso Saccardo) a Conegliano, aut. 1877.

116. **Phoma montana** Speg. Peritheciis subepidermicis, erumpentibus, membranaceo-carbonaceis, 110-150,

sublenticularibus. Spermatiis fusoideis utrinque acuminatis, nubiloso-granulosis, non vel 2-guttulatis, $6-10 \approx 1\frac{1}{2}-2$.

Hab. In ramis emortuis *Saturejæ montanæ a Susegana*, Martio 1877.

117. **Phyllosticta Cratægi** Speg. Maculis subrotundis, griseis; peritheciis parenchymate innatis, minutissimis, 60-80 diam., contextu laxe parenchymatico-atro; spermatiis hyalinis, minutissimis, ellipticis vel ellipsoideis, biguttulatis, $2\frac{1}{2}-3 \approx 1-1\frac{1}{2}$.

Hab. In foliis *Cratægi Oxyacanthæ a Belluno*, Oct. 1878, rarissime.

118. **Phyllosticta prunicola** Sacc. Michel. 156.

Hab. In foliis *Pruni domesticæ a Belluno*, Oct. 1878.

119. **Phyllosticta aquatica** Speg. Peritheciis subepidermicis, lenticularibus, 100-110, ostiolo latiusculo, 10, mer. d. pertuso; contextu laxissime et tenuissime parenchymatico-membranaceo, olivaceo. Spermatiis ellipticis utrinque obtusiusculis, quandoque inæquilate-ralibus, hyalinis, 4-5 \approx 2.

Hab. Ad folia putrida *Lemnæ minoris prope Conegliano*, aut. 1878.

120. **Ascochyta aquatica** Speg. Peritheciis minutis, 90-100, subepidermicis, contextu parenchymatico, densiusculo, olivaceo, ostiolo parvulo pertuso. Spermatiis oblongo-ellipticis, medio 1-septatis, ad septum non vel vix constrictis, dilute chlorinis, 7-9 \approx 2-3.

Hab. Ad folia putrida *Lemnæ minoris prope Conegliano*, aest. 1879.

121. **Ascochyta chlorospora** Speg. Peritheciis subepidermicis maculis griseis, rotundis innatis, lenticularibus, minutis, 80-90, ostiolo latiusculo pertusis; spermatiis ellipticis, medio 1-septato-constrictis, dilute chlorinis, 10-12 $\approx 3\frac{1}{2}$.

Hab. In foliis languidis *Pruni domesticæ, a Belluno*, Oct. 1878.

122. **Hendersonia charticola** Speg. Peritheciis innatis, lenticularibus vel subglobosis, primo tectis dein erum-

pentibus, 120—150, contextu membranaceo—parenchymatico denso, fuligineo, subopaco; stylosporis oblongo—ellipticis, quandoque clavulatis, 4—septatis, ad septa leniter constrictis, septo longitudinali quandoque donatis, 40 ≈ 15—16, primo fumose hyalinis, dein subfuscouscentibus.

Hab. Ad chartam stercoratam putrescentem a Conegliano, in locis umbrosis aut. 1878.

123. **Hendersonia martialis** Spieg. Peritheciis lenticulari—conoideis, minutis, 120—150 micr., sparsis, subepidermicis, ostiolo papilliformi vix exerto, contextu parenchymatico—celluloso, fusco—olivaceo; conidiis oblongo—ellipticis utrinque obtusiusculis, 2—3septatis, ad septa plus minusve constrictis, pallide olivaceis, 12—18 ≈ 5—6.

Hab. Ad caules emortuos Staticis Limonii in Campo Martis, Venetiis, 25 Mart. 1879.

124. **Diplodia Macluræ** Spieg. Peritheciis nunc sparsis, nunc in acervulos aggregatis, primo epidermide tectis, dein erumpentibus atque ab ea cinctis; stylosporis ovoideis, 23—25 ≈ 8—11, in sterigmatibus parvulis, 10 ≈ 3, acrogenis, primo continuis, guttulatis, luteolis, dein medio 1—septatis non vel vix constrictis, obscure fuligineis.

Hab. In ramulis emortuis Macluræ aurantiae a Susegana, Sept. 1876.

125. **Diplodia Iulibrissin** Spieg. Peritheciis primo subepidermide nidulantibus, dein erumpentibus, nigricantibus, subglobosis, papillatis. Stylosporis, 20—25 ≈ 10—12, dense fuligineis, medio 1—septato—constrictis.

Hab. In ramulis emortuis Albizziæ Iulibrissin a Conegliano, aut. 1876—78.

126. **Diplodia Ailanthina** Spieg. Peritheciis subglobosis, olivaceo—fuscis, papillatis, 200—250, subepidermicis, vel ligno decorticato parallele dispositis, ostiolo perforatis. Stylosporis fuligineis, 20 ≈ 7, 1—septatis ad septum non vel vix constrictis.

Hab. In ramulis emortuis *Ailanthi glandulosæ*, *a Conegliano*, aut. 1876-78.

127. **Diplodia Sophoræ** Speg. Peritheciis subepidermicis globosis, nigricantibus, ostiolo minuto pertusis, rugulosis. Stylosporis oblongo-ellipticis vel ovoideis, 1-septato-constrictis, fuligineis, 20-22 ≈ 10-12.

Hab. In ramulis emortuis *Sophoræ japonicae* *a Conegliano et Susegana*, Oct. 1876-77.

128. **Diplodia Photiniæ** Speg. Peritheciis hemisphæricis, subepidermicis, carbonaceis. Stylosporis ochraceo-fuligineis non vel 1-septatis, crassiusculæ et obscurius tunicatis, quandoque 1-2guttulatis, 22 ≈ 12-14, ad septum sæpius non constrictis.

Hab. In foliis dejectis, putrescentibus *Photiniæ serrulatae*, *a Conegliano*, Oct. 1877.

129. **Diplodia Althææ** Speg. Peritheciis superficialibus, hemisphærico-conoideis, ostiolo papillato acutiusculo donatis, 250-300. Stylosporis fuligineis, ellipticis, utrinque obtusis, 1-septatis, non vel vix constrictis, 16-20 ≈ 8-9.

Hab. In caulibus emortuis putrescentibus *Althææ roseæ* *a Conegliano*, Oct. 1877.

130. **Diplodia Lagerstroemiæ** Speg. Peritheciis atris, papillulatis, sublenticularibus, erumpentibus, 200-250, ostiolo late biante pertusis; stylosporis opace olivaceo-fuligineis, 1-septatis ad septum non vel vix constrictis, utrinque obtusiusculis, 18-20 ≈ 8-10.

Hab. In ramulis emortuis dejectis *Lagerstroemiacæ* *a Conegliano*, Aug. 1878.

131. **Diplodia Calycanthi** (Schw?) Speg. *Sphaeria Calycanthi* Schw?) Peritheciis subepidermicis, erumpentibus, atris, lenticularibus, rugulosis, 200-300. Stylosporis primo luteolis dein olivaceo-fuscis, 1-septato-constrictis, ovoideis vel ellipticis, 19-22 ≈ 9-10.

Hab. In ramulis emortuis *Calycanthi floridi* *a Conegliano*, aest. 1877.

132. **Diplodia Taxi** (Sow.) Fuck. Symb. myc. p. 394.

Hab. Ad folia emortua *Taxi baccatæ* *a Susegana*, aut. 1876-78.

133. **Diplodia Aesculi** Lév — Fuck. Symb. myc. p. 394.
Hab. Ad ramulos emortuos Aesculi Hippocastani *a Conegliano*, vere 1879.
134. **Diplodia Catalpæ** Speg. Peritheciis subepidermicis, tectis, dein erumpentibus, globosis, atris, papillulatis. Stylosporis oblongo—ellipsoideis, medio 1—septatis, non constrictis, fuligineo—olivaceis, 20—22 ≈ 8—11.
Hab. In ramulis emortuis Bignoniæ Catalpæ *a Conegliano*, aut. 1877.
135. **Diplodia Nerii** Speg. Peritheciis lenticularibus, subepidermicis, dein erumpentibus, atris, 200—250. Stylosporis ovoideis vel ellipticis, 1—septato—constrictis, olivaceo—fuligineis, 18—22 ≈ 8—10.
Hab. In ramulis et petiolis foliorum Nerii Oleandri, *a Conegliano*, aut. 1876.
136. **Diplodia Oleandri** Speg. Peritheciis laxe gregariis, circinatim dispositis (an semper?), globosis, tectis, prominulis, ostiolo papillato exerto donatis, 200—300. Stylosporis ellipticis vel ovoideis, fuligineo—olivaceis, subopacis, 25—30 ≈ 10—14, continuis, sterigmatibus hyalinis, 8—10 ≈ 5—6, suffultis.
Hab. Ad folia dejecta Nerii Oleandri *a Conegliano*, vere 1879.
137. **Septoria Ornithogali** Pass. in Thuem. Mycoth. Univ. n. 496. *S. Ornithogali* Oudm. Aanw. V. de Flr. Myc. v. Nederland n. 6 pag. 6. *S. Ornithogali* Speg. Decades Myc. Ital. n. 20.
Hab. In apicibus foliorum Ornithogali umbellati, *a Conegliano*, vere 1879.
138. **Septoria littoralis** Speg. Peritheciis 5-8 gregariis primo tectis dein erumpentibus, subsuperficialibus, globulosis, atris, ostiolo papillato, sicco umbilicato, donatis; contextu membranaceo, densissime parenchymatico, fuligineo—atro. Spermatiis cylindraceis, rectis vel curvulis apice rotundatis, basi subtruncatis, 50—65 ≈ 3 $\frac{1}{2}$ —4, 3—septatis, pluri—guttulatis, hyalinis in sterigmatibus, ampulluliformibus (ut hyphæ Chalaræ Ampullulæ), guttulatis, 10 ≈ 4—5, hyalinis.
Hab. In pagina interna vaginalium Phragmitis

communis in insula *Lido* dicta, apud Venetias, Martio 1879.

39. **Sacidium Venetum** Speg. Peritheciis sparsis, superficialibus subglobosis, 200—250 mic., atro—cærulecentibus, nitidulis, astomis, contextu atro—fuligineo, punctulato; spermatiis ellipticis, crassiuscule tunicatis utrinque rotundatis, 12—15 = 8—9, chlorino—flavidis, eguttulatis.

Hab. Ad folia viva *Salicis* purpureæ, a *Conegliano*, aut. 1877, rarissime.

Obs. Contextu peritheciorum cum *S. Spegazziniano* perfecte congruit. Ego iterum *S. Spegazzinianum* legi in caulis emortuis *Urticæ* dioicæ a *Conegliano*, æst. 1877.
