

List of the more uncommon fungi observed during the Foray:—

- | | |
|------------------------------------|----------------------------------|
| Amanita spissa Fr. | Boletus flavidus Fr. |
| Tricholoma sejunctum Sow. | tenuipes Cke. |
| flavobrunneum Fr. | bovinus Linn. |
| albobrunneum Pers. | variegatus Sow. |
| imbricatum Fr. | calopus Fr. |
| acerbum Bull. | candicans Fr. |
| Collybia nitellina Fr. | aereus Bull. |
| ambusta Fr. | purpureus Fr. |
| Mycena aurantiomarginata Fr. | duriusculus Schulz. |
| Iris Berk. | Polyporus Schweinizii Fr. |
| capillaris Schum. | intybaseus Fr. |
| Entoloma lividum Bull. | spongia Fr. |
| Leptonia formosa Fr. | amorphus Fr. |
| Pholiota radicata Bull. | spumeus Fr. |
| Stropharia squamosa Pers. var. | Fomes applanatus Wallr. |
| aurantiaca Cke. | Poria umbrina Fr. |
| Cortinarius (Myxaciium) collinitus | Hydnum Queletii Fr. |
| Fr. | cyathiforme Schaeff. |
| (Dermocybe)lepidopus Cke. | Irpex obliquus Fr. |
| miltinus Fr. | Radulum orbiculare Fr. |
| raphanoides | Grandinia granulosa Fr. |
| Pers. | Peniophora gigantea Mass. |
| (Hygrocybe) castaneus Fr. | Coniophora sulphurea Mass. |
| Gomphidius roseus Fr. | Sparassis crispa Fr. |
| gracilis Berk. & Br. | Calocera stricta Fr. |
| Hygrophorus cossus Fr. | Tremella intumescens Eng. Bot. |
| miniatus Fr. | Tremellodon gelatinosum Pers. |
| Lactarius insulsus Fr. | Rhizopogon rubescens Tul. |
| pallidus Fr. | Uredo Mülleri Schröt. |
| theiogalus Fr. | Phragmidium albidum Kühn. |
| fuliginosus Fr. | Orbilia luteorubella Karst. |
| serifluus Fr. | Dasyscypha calycina Fckl. |
| Russula caerulea Fr. | Ciboria echinophila Sacc. |
| lactea Fr. var. incarnata | Nectria inaurata B. & Br. |
| Quél. | Poronia punctata Fr. |
| azurea Bres. | Quaternaria Persoonii Tul. |
| consobrina Fr. var. inter- | conidial form. |
| media Cke. | Schirrhia rimosa Fckl., Pycnidia |
| chamaeleontina Fr. | on Arundo phragmitis. |
| Lentinus cochleatus Fr. | |