

1 **Microbial Life in a Liquid Asphalt Desert**

2
3 Dirk Schulze-Makuch¹, Shirin Haque², Marina Resendes de Sousa Antonio¹, Denzil Ali²,
4 Riad Hosein³, Young C. Song⁴, Jinshu Yang⁴, Elena Zaikova⁴, Denise M. Beckles³,
5 Edward Guinan⁵, Harry J. Lehto⁶ and Steven J. Hallam^{4,7}

6
7 ¹ School of Earth & Environmental Sciences, Washington State University, USA

8 ² Department of Physics, University of the West Indies, Trinidad & Tobago

9 ³ Department of Chemistry, University of the West Indies, Trinidad & Tobago

10 ⁴ Department of Microbiology & Immunology, University of British Columbia, Canada

11 ⁵ Department of Physics & Astronomy, Villanova University, USA

12 ⁶ Department of Physics & Astronomy, Tuorla Observatory, University of Turku, Finland

13 ⁷ Graduate Program in Bioinformatics, University of British Columbia, Canada

14
15 **Running Title:** Microbial Life in a Liquid Asphalt Desert

16
17 **Correspondence to be addressed to:** Dirk Schulze-Makuch, SEES, Webster Hall 1148,

18 Washington State University, tel: 1-509-335-1180, fax: 1-509-335-3700, email:

19 dirksm@wsu.edu or Steven Hallam, Department of Microbiology & Immunology,

20 University of British Columbia, Canada, email: shallam@interchange.ubc.ca

21
22 **Keywords:** hydrocarbons, extreme environments, water activity, Titan, ecosystem

1 **Abstract**

2 **An active microbiota, reaching up to 10^7 cells/g, was found to inhabit a naturally**
3 **occurring asphalt lake characterized by low water activity and elevated**
4 **temperature. Geochemical and molecular taxonomic approaches revealed novel and**
5 **deeply branching microbial assemblages mediating anaerobic hydrocarbon**
6 **degradation, metal respiration and C1 utilization pathways. These results open a**
7 **window into the origin and adaptive evolution of microbial life within recalcitrant**
8 **hydrocarbon matrices, and establish the site as a useful analog for the liquid**
9 **hydrocarbon environments on Saturn's moon Titan.**

10

11

12

13

14

15

16

17

18

19

20

1 **Introduction**

2
3 Pitch Lake in Trinidad & Tobago is one of three extant natural asphalt lakes on
4 Earth (Peckham, 1895). Located on the southwest peninsula of Trinidad near the town of
5 La Brea and covering an area of approximately 46 hectares, Pitch Lake is recharged by
6 pitch seepage, a form of petroleum consisting of mostly asphaltenes, from the
7 surrounding oil-rich region. During upward seepage, hydrocarbons mix with mud and
8 gases under high pressure and the lighter portion evaporates or is volatilized, producing
9 liquid asphalt. Pitch Lake has both active liquid areas and inactive areas that have
10 solidified due to surface cooling analogous to undersea pillow lavas. Long a source of
11 asphalt for road paving projects, Pitch Lake has more recently become the subject of
12 intensive field studies focusing on the biogenic potential of liquid hydrocarbon
13 environments.

14 The liquid hydrocarbon environment at Pitch Lake is of interest based on several
15 grounds. First, biodegraded oils including tar sands, shales and natural asphalts represent
16 a significant proportion of remaining global petroleum supplies, but microbial controls
17 and metabolic processes underlying hydrocarbon degradation and recovery within these
18 systems remain obscure. Second, Saturn's moon Titan exhibits many of the same
19 environmental conditions as Pitch Lake and various authors have speculated on the
20 possibility of life within the liquid hydrocarbons of Titan (e.g., Abbas and Schulze-
21 Makuch, 2002; Schulze-Makuch and Grinspoon, 2005; McKay and Smith, 2005; Baross
22 et al. 2007). Thus, Pitch Lake is an appropriate analogue site to study the adaptation
23 capabilities of life in a water-poor hydrocarbon matrix.

24
25 **Material and Methods**

26
27 We collected six samples from Pitch Lake (A-F) for geochemical analysis, from
28 which four (PL-A through D) were also used for biological characterization (Fig. 1A and
29 Table 1). Comparative reference samples were also collected from an adjacent oil well
30 (OP) and a nearby terrestrial mud volcano (Fig. 1D) known as Devil's Woodyard (MV).

31
32 *Field Activities*

33 Samples were collected from the asphalt lake with latex sampling gloves in sterile
34 sampling bottles and described *in situ*. Water activity measurements were taken in the
35 field with a Decagon p_{a_w} -kit water activity meter on select samples. Gas bubbles were
36 identified and sampled in the field with a syringe, which was purged of ambient air and
37 flushed with nitrogen prior sampling. Methane gas was also measured *in situ* emanating
38 from the gas bubbles with an open-path, hand-held laser system for the detection of
39 methane gas (Van Well et al., 2005).

40
41 *Gas analyses*

42 Syringes with gas samples were shipped on ice to Dr. Sherwood Lollar's laboratory at the
43 University of Toronto. The composition of inorganic gases (Helium, Hydrogen, Oxygen,
44 Nitrogen and Carbon Dioxide) were analyzed by injection of 300 μ l of sample in a Varian
45 3800 gas chromatograph equipped with a micro-Thermo Conductivity Detector (m-
46 TCD). The gases were separated through a 0.32 mm internal diameter, 30 m long

1 Molecular Sieve column with an argon carrier gas flow of 1 ml per minute. The column
2 oven temperature program started at 30°C for 7 minutes then the column was heated to
3 90°C with a ramp of 10 degrees per minutes, and then heated to 200°C at 30 degrees per
4 minutes and hold at this temperature for 20 minutes.

5 The composition of organic gases (hydrocarbons from methane to pentane) was
6 analyzed by injection of 300 µl of sample in a Varian 3400 gas chromatograph equipped
7 with a Flame Ionization Detector (FID). The gases were separated through a 0.52 mm
8 internal diameter, 30 m long GSQ column with an helium carrier gas flow of 7 ml per
9 minute. The column oven temperature program started at 50°C for 5 minutes then the
10 column was heated to 200°C with a ramp of 25 degrees per minutes and hold at this
11 temperature for 5 minutes. Gas compositions were calculated using certified standards
12 injected the same day. All samples run in duplicate and mean values reported.
13 Uncertainties in reported concentrations are within ±5%.

14 Carbon isotope analyses of organic gases were conducted on a GC-C-Isotope
15 Ratio Mass Spectrometry (IRMS) system composed of a Varian 3400 capillary gas
16 chromatogram, and an oxidation oven at 980°C interfaced directly to a Finnigan 252 gas
17 source mass spectrometer. Carbon dioxide, methane, ethane, propane, butanes and
18 pentanes were analyzed on a 60 m GSQ column (ID 0.32 mm). The temperature program
19 started at 35°C hold 6 minutes, ramped to 110°C at 3°C per minute and ramped to 220°C
20 at 5°C per minute, hold 5 minutes. All samples were run in duplicate and mean values
21 were reported. Total uncertainty, incorporating both accuracy and reproducibility was
22 ±0.5‰ with respect to v-PDB standard.

23 24 *Bulk Isotope Analyses*

25 Bulk isotope results were obtained using IRMS. Samples for carbon and nitrogen isotopic
26 analysis were converted to CO₂ with an elemental analyzer (ECS 4010, Costech
27 Analytical, Valencia, CA). These two gases were separated with a 3m GC column and
28 analyzed with a continuous flow isotope ratio mass spectrometer (Delta PlusXP,
29 Thermofinnigan, Bremen, Germany) at the Laboratory of Biotechnology and Bioanalysis
30 at Washington State University. In addition, select samples were analyzed at
31 IsoAnalytical in Crewe, U.K., also using EA-IRMS. The samples were first converted to
32 pure N₂ and CO₂ to permit analysis by IRMS. The samples were placed in clean tin
33 capsules and loaded into an automatic sampler. They were then dropped into a
34 combustion furnace held at 1000°C where they were combusted in the presence of an
35 excess of oxygen. The tin capsules flash combust causing the temperature in the vicinity
36 of the sample to rise to ca. 1700°C. The gaseous products of combustion were swept in a
37 helium stream over a Cr₂O₃ combustion catalyst, CuO wires to oxidize hydrocarbons and
38 silver wool to remove sulfur and halides. The resultant gases (N₂, NO_x, H₂O, O₂, and
39 CO₂) were then swept through a reduction stage of pure copper wires held at 600°C. This
40 removed any remaining oxygen and converted NO_x gases to N₂. Water was removed
41 using a magnesium perchlorate trap, while removal of CO₂ was available using a
42 selectable Carbosorb™ trap. Nitrogen and carbon dioxide were separated by a packed
43 column gas chromatograph held at an isothermal temperature. The resultant
44 chromatographic peaks sequentially entered the ion source of the IRMS where they were
45 ionized and accelerated. Gas species of different mass were separated in a magnetic field
46 and simultaneously measured by a Faraday cup universal collector array. For N₂, masses

1 28, 29 and 30 were monitored and for CO₂, masses 44, 45 and 46. The results for carbon-
2 13 that run at both labs were statistically the same.

3 For the determination of sulphur-34 the samples were first converted to pure SO₂
4 to permit analysis by IRMS. Samples were placed in clean tin capsules and loaded into an
5 automatic sampler. They were then dropped into a combustion furnace held at 1080°C
6 where they were combusted in the presence of an excess of oxygen. The tin capsules
7 flash combust causing the temperature in the vicinity of the sample to rise to ca. 1700°C.
8 The gaseous products of combustion were then swept in a helium stream through tungstic
9 oxide and zirconium oxide combustion catalysts and then reduced over high purity
10 copper wires. Water was removed using a Nafion™ membrane, permeable to only water.
11 Sulphur dioxide was separated by a packed column gas chromatograph held at an
12 isothermal temperature. The resultant SO₂ chromatographic peak entered the ion source
13 of the IRMS where it was ionized and accelerated. Gas species of different mass were
14 separated in a magnetic field and simultaneously measured by a Faraday cup universal
15 collector array. For SO₂, masses 64, 65 and 66 were monitored.

16 17 *Amino Acid Analyses*

18 The samples for amino acid analysis samples were dissolved with dimethyl sulfoxide
19 (DMSO) and analyzed with the Mars Organic Analyzer (MOA) housed at the University
20 of Berkeley, CA. MOA is a microfabricated capillary electrophoresis (CE) instrument
21 for sensitive amino acid analysis (Skelley et al., 2005). The microdevice consists of a
22 four-wafer sandwich combining glass CE separation channels, microfabricated pneumatic
23 membrane valves and pumps, and a nanoliter fluidic network.

24 25 *Phospholipid Fatty-acid (PFLA) Analysis*

26 PLFA analysis was conducted using a modified Bligh and Dyer method (White et al.,
27 1979). A one-phase chloroform-methanol buffer was used as an extractant. Lipids were
28 recovered, dissolved in chloroform, and fractionated on silicic acid columns into polar-,
29 neutral-, and glycol-lipid fractions. The PLFA was recovered as methyl esters in hexane
30 by transesterfying the polar fraction with mild alkali solution. PLFA were then analyzed
31 by gas chromatography with peaks being confirmed via electron impact mass
32 spectrometry.

33 34 *DNA Extraction*

35 Samples were ground to a fine powder under liquid nitrogen using a porcelain mortar and
36 pestle. Genomic DNA was extracted from ~10 g of powdered asphalt using the
37 PowerMax Soil DNA isolation kit according to the manufacturer's instructions (Mo Bio,
38 Carlsbad, CA). Dilute samples were washed 3X in 2 ml TE buffer (pH 8.0) and
39 concentrated down to ~200 µl using Amicon Ultra centrifugal filter devices (Millipore,
40 Bellerica, MA). DNA quality was determined by running 5µl of each sample on 1%
41 agarose gels in 1X TBE at 15 volts for 12 hours. DNA samples were subsequently used
42 for quantification of bacterial and archaeal taxonomic and functional gene markers as
43 well as clone library production.

44 45 *qPCR of Small Subunit Ribosomal RNA (SSU rRNA) Genes*

46 Total bacterial and archaeal SSU rRNA gene copy number were determined by

1 qPCR using either bacterial-specific (27F, 5'-AGAGTTTGATCCTGGCTCAG) or
2 archaeal-specific (20F, 5' - TTCCGGTTGATCCYGCCRG) forward primers coupled to
3 a universal reverse primer (DW519R, 5'-GNTTACCGCGGCKGCTG) as described
4 previously (Zaikova et al., 2010). Standards used for total bacterial and archaeal
5 quantification were derived from SSU rRNA gene clone libraries. A 10-fold dilution
6 series for each standard ranging from 3×10^1 to 3×10^7 copies per μl for both bacteria and
7 archaea was used in real time analysis. For the purposes of this study the limit of
8 detection was established based on a comparison of dissociation curves and Ct thresholds
9 for all sample replicates to be $\sim 2 \times 10^2$ copies/ml.

10 *Quantitative Polymerase Chain Reaction (qPCR) of Functional Gene Markers*

11 Functional markers for *tdo*, *alkB*, *mcrA* and *pmoA* were quantified using Bio-Dechlor
12 CENSUS real-time PCR available through Microbial Insights
13 (<http://www.microbe.com/>). Real-time PCR was performed on each sample using
14 oligonucleotides that were designed to target functional genes (such as toluene
15 dioxygenase *tdo* (Baldwin et al., 2003)) on an ABI 7000 Sequence Detection System
16 (Applied Biosystems, Foster City, CA). PCR mixtures contained 1 X Cloned Pfu Buffer
17 (Stratagene, LaJolla, CA), 0.2 mM of each of the four deoxynucleoside triphosphates,
18 SYBR green (diluted 1:30,000; Molecular Probes, Eugene, OR), and 1 U of Pfu Turbo
19 HotStart DNA polymerase (Stratagene, LaJolla, CA). Annealing temperatures, primer
20 concentrations, and MgCl_2 concentrations varied dependent of the primer set used. A
21 calibration curve was obtained by using a serial dilution of a known concentration of
22 positive control DNA. The C_T values that were obtained from each sample were then
23 compared with the standard curve to determine the original sample DNA concentration.
24

25 *PCR Amplification of SSU rRNA genes*

26 SSU rRNA gene sequences were amplified with the primers B27F (5'-
27 AGAGTTTGATCCTGGCTCAG) and U1492R (5'- GGTTACCTTAGTTACGACTT)
28 for bacteria and A20F (5'-TTCCGGTTGATCCYGCCRG) and A958R (5'-
29 YCCGGCGTTGAMTCCAATT) for archaea using the following PCR profile: 3 min at
30 94°C followed by 30 cycles of 95°C for 20 seconds, 55°C for 20 seconds, 72°C for 1
31 minute and a final extension of 10 minutes at 72°C. Each 50 μl reaction contained 1 μl of
32 template DNA, 1.25 μl each 10 μM forward and reverse primer, 2.5 U Hercules II
33 proofreading polymerase (Stratagene, La Jolla, CA), 5 μl 10 mM deoxynucleotides, 41.5
34 μl 1x buffer containing 2 mM MgCl_2 .
35

36 *PCR Amplification of methyl coenzyme M reductase subunit A (mcrA) genes*

37 With the exception of PL-C and PL-D *mcrA* gene sequences were amplified from PL, OP
38 and MV samples with the primers ME1 (5'- GCMATGCARATHGGWATGTC) and
39 ME2 (5'-TCATKGCRTAGTTDGGRTAGT) (Hales et al., 1996) using the PCR profile:
40 3 minutes at 94°C followed by 35 cycles of 94°C for 45 seconds, 50°C for 45 seconds,
41 72°C for 1.5 minutes and a final extension of 5 minutes at 72°C. Each 50 μl reaction
42 contained 1 μl of template DNA, 1.25 μl each 10 μM forward and reverse primer, 1 μl
43 Herculase II fusion DNA polymerase (Stratagene, La Jolla, CA), 5 μl 10 mM
44 deoxynucleotides, 10 μl 5x Herculase II reaction buffer, and 30.5 μl of distilled water. In
45 the case of PL-C and PL-D we used the Epicentre FailSafe PCR PreMix Selection Kit
46

1 according to the manufacturer's specifications. Using the same profile describe above
2 FailSafe amplification yielded sufficient PL-D product for downstream cloning.
3 However, PL-C remained refractory to amplification and was therefore not included in
4 the study.

6 *Clone Library Construction and Screening*

7 Amplicons were visualized on 1% agarose gels in 1X TBE and purified using the
8 Qiaquick Gel Extraction Kit (Qiagen, Valencia, CA) according to the manufacturer's
9 instructions. Samples were eluted in 30 µl of sterile filtered water (pH 7.5) and
10 concentrated down to ~6 µl using a SpeedVac system. In preparation for Topo-TA
11 cloning, 3' A-overhangs were added by incubating the purified DNA sample for 15
12 minutes at 72 °C in the presence of 1 U of BioShop Taq (Canada) and 2.5 mM dNTPs in
13 2 µl 1X PCR buffer containing 2.5 mM MgCl₂. From this reaction, 4 µl was used in a TA
14 cloning reaction (pCR4-TOPO TA cloning kit for sequencing and transformed into
15 chemically competent Mach-1-T1R cells according to the manufacturer's instructions
16 (Invitrogen, Carlsbad, CA). Transformants were picked into 96-well plates containing
17 180 µl Luria Broth (LB), supplemented with 50 µg/ml kanamycin and 10% glycerol, and
18 grown overnight at 37°C prior to storage at -80°C. Inserts were amplified directly from
19 glycerol stocks with M13F (5'- GTAAAACGACGGCCAG) and M13R (5' -
20 CAGGAAACAGCTATGAC) primers using the SSU rRNA gene PCR protocol with an
21 additional 10 minutes at 94°C added to the very beginning of the PCR profile to ensure
22 adequate cell lysis.

23 One 96-well plate per sample was selected for restriction fragment length
24 polymorphism (RFLP) analysis *i.e.* DNA fingerprinting, using the common 4-base cutter
25 Rsa I (Invitrogen, Carlsbad, CA) Each digestion reaction consisted of 2 µl of 10x React 1
26 buffer, 5 µl of M13 amplified PCR product, 13 µl of sterile filtered water and 5 U of Rsa
27 I. Reactions were incubated for 2 hours at 37°C followed by Rsa I inactivation at 65°C
28 for 10 minutes. Restriction patterns were visualized by running 5 µl of each RSA I
29 digestion mixture on a 2% agarose gel (20 cm in width) in 1X TBE for 90 minutes at 120
30 V. SSU rRNA and *mcrA* gene clones exhibiting unique restriction patterns were selected
31 for Sanger sequencing through the McGill University and Genome Quebec Innovation
32 Centre (Montreal, Quebec, Canada). A total of 492 archaeal and 496 bacterial SSU rRNA
33 gene and 406 *mcrA* clones were screened, providing 65, 72 and 74 unique restriction
34 patterns respectively, for downstream sequencing and taxonomic identification. Sequence
35 data was collected on an ABI Prism 3100 DNA sequencer (Applied Biosystems Inc,
36 Foster, CA) using Big DyeTM chemistry (PE Biosystems, Foster, CA) according to
37 manufacturer's instructions. Plasmids were bidirectionally sequenced with M13F and
38 M13R primers. Sequences were edited manually from traces using Sequencher software
39 V4.1.2 (Gene Codes Corporation, Ann Arbor, MI).

41 *Phylogenetic Analysis of Taxonomic and Functional Gene Markers*

42 Prior to phylogenetic analysis all SSU rRNA gene sequences were screened with
43 Chimera Check (DeSantis et al., 2006) using the Bellerophon (Huber et al., 2004) server
44 (<http://foo.maths.uq.edu.au/~huber/bellerophon.pl>). All sequences were analyzed using
45 the ARB software package (Ludwig et al., 2004). Sequences were imported into the full-
46 length SSU SILVA 92 database (Pruesse et al., 2004) (<http://www.arb-silva.de>) and

1 aligned to their closest relatives. For phylogenetic analysis, PL, OP, MV and selected
2 reference sequences were exported from ARB and taxonomic assignments were made
3 using the Greengenes (DeSantis et al., 2006) taxonomic hierarchy supplemented with
4 environmental reference sequences. In the case of *mcrA*, nucleotide sequences were
5 conceptually translated using the Transeq server
6 (<http://www.ebi.ac.uk/emboss/transeq>). Selected reference proteins were downloaded
7 from GenBank (<http://ncbi.nlm.nih.gov/genbank>). Both SSU rRNA gene and *mcrA*
8 sequences were realigned using the multi-sequence alignment tool MUSCLE (Edgar,
9 2004). The resulting alignments were imported into MacClade (Maddison and Maddison,
10 2003) for manual refinement.

11 SSU rRNA gene trees were inferred with PHYML (Guindon et al., 2005) using an
12 HKY + 4 Γ + I model of nucleotide evolution where the α parameter of the Γ distribution,
13 the proportion of invariable sites, and the transition/transversion ratio were estimated for
14 each dataset. The conceptually translated *mcrA* tree was inferred with PHYML using a
15 JTT + 4 Γ + I model of amino acid substitution. In all cases the confidence of each node
16 was determined by assembling a consensus tree of 100 bootstrap replicates. PHYML
17 trees were imported into the interactive Tree of Life (iTOL) server
18 (<http://www.itol.embl.de>) to convert them into radial format. Dot plots of taxon
19 abundance were generated with a custom perl script (bubble.pl).

21 *UniFrac Analysis*

22 UniFrac (<http://bmf2.colorado.edu/unifrac/index.psp>) is a web application supporting
23 multivariate statistical methods to evaluate microbial diversity based on phylogenetic
24 information (Lozupone and Knight, 2005; Lozupone et al., 2006). Distance matrices,
25 corresponding to independent alignments of archaeal or bacterial SSU rRNA gene
26 sequences were created using the Dnadist program with F84 nucleotide substitution
27 model implemented in PHYLIP (Felsenstein, 2005). Distance matrices were converted to
28 neighbor joining (NJ) trees using the Neighbor program implemented in PHYLIP (16) to
29 generate single rooted phylogenetic trees in Newick format that were in turn used as input
30 files for UniFrac analysis. Hierarchical cluster trees were constructed on the basis of the
31 distance matrix calculated by the UniFrac algorithm, and scatter-plots of principal
32 coordinates, *i.e.* ordinations, were generated by selecting the ScatterPlot and Assign
33 series options. In both forms of analyses the abundance weight of each sequence was not
34 used.

36 *Hierarchical Cluster Analysis*

37 SSU rRNA gene sequences from PL, OP MV and environmental reference studies were
38 queried against the Greengenes database. The output from the nucleotide BLAST search
39 and a modified table of the Greengenes taxonomy hierarchy was used as input files for a
40 custom perl script (tax_sum.pl) to generate a taxonomy abundance table. The values in
41 all the entries were corrected by using the following formula $A(i,k) = \ln([a(i,k) + 1]$,
42 where each '*i*th' row represents a different taxonomic group, while each '*k*th' column
43 represents a different environment. The corrected values were imported into the R
44 statistics software package (<http://www.r-project.org/>) to generate a custom heatmap.
45 Distance similarities between compositional profiles from each environment were
46 calculated using the Manhattan distance formula.

Results

Sample temperatures collected from active areas ranged between 32°C and 56°C, significantly above ambient air temperatures (Table 1), which ranged between 20°C and 25°C. Gases diffusing through the liquid asphalt form large bubbles that burst upon reaching a critical size or pressure (Fig. 1B and C). Direct measurement of such bubbles revealed gas content dominated by 72.9±3.8% methane, 1.7±0.9% ethane, 1.0±0.2% propane and trace amounts of butane, with 25.8±5.9% carbon dioxide and 1.3±0.8% nitrogen gas. Helium, hydrogen, and oxygen concentrations were below detection limits (<0.01 %).

Mean stable carbon, nitrogen and sulfur isotope ratios of Pitch Lake (PL) and Oil Well (OP) samples were more similar to one another than to the Mud Volcano (MV) sample, consistent with common or related source points (Table 2). While the ¹³C isotope fractionation for bulk carbon in PL samples was -28.1±0.4‰, carbon stable-isotope ratios for the lighter alkanes: methane, ethane, propane and butane, were -49.0±1.4 ‰, -34.1±0.2 ‰, -30.9±1.8 ‰ and -29.8±0.8 ‰, respectively, indicating potential microbial fractionation effects. However, water activity in these samples ranged between 0.49 and 0.65, at or below the reported threshold to life on Earth (Grant, 2004; Tosca et al., 2008). To resolve this seeming contradiction, we quantified the presence of amino acids such as Val (3.7±0.8 nM), Ala/Ser (13±6 nM), Gly (3±2 nM), Glu (1.3±0.6 nM), and Asp (2.3±0.7 nM) and confirmed biological activity on the basis of phospholipid fatty acid (PLFA), and taxonomic and functional gene markers. It remains to be determined whether microbial life is constrained to water or brine filled inclusions manifesting higher moisture content than the surrounding asphalt matrix, as seen in the permanent ice covers of frozen lakes and glaciers in McMurdo Dry Valleys, Antarctica (Priscu et al., 1998; Mikucki et al., 2009).

PLFA analysis was used to determine viable microbial biomass. Monoenoic and normal saturated PLFA were recovered, indicating active microbial populations ranging between 10⁶ to 10⁷ cells/g in PL-A through D (Table 3). Monoenoic PLFA occur in Gram-negative bacteria, typically Proteobacteria, while normal saturated PLFA are more widely distributed in bacteria and eukaryotes. No terminally branched saturated, branched monoenoic, mid-chain branched saturated, nor diagnostic eukaryotic PLFA were recovered. However, in samples PL-A through C several common biomarkers for Gram-negative bacteria including cy17:0 and cy19:0 were identified (Table 3).

Taxonomic structure was evaluated on the basis of bacterial and archaeal small subunit ribosomal RNA (SSU rRNA) gene profiles recovered from samples PL-A through D, OP and MV. Multivariate statistical approaches were used to compare profile relationships between samples. Common patterns were observed for both bacterial (Fig. 2A) and archaeal (Fig. 2B) domains. PL-B and OP were more similar to one another than other PL samples consistent with stable isotope analysis (Table 2). The deeper active PL-A and PL-D clustered together whereas PL-C and MV showed more variation in taxonomic structure. Consistent with PLFA results, total bacterial SSU rRNA gene copy numbers in PL samples ranged between 10⁶ to 10⁷ copies/g (Fig. 2A and Table 4).

1 Archaeal SSU rRNA copy numbers were several orders of magnitude lower, ranging
2 between 10^4 to 10^5 copies/g (Fig 2B and Table 4).

3 Active PL-A and D samples were further analyzed for a subset of functional gene
4 markers associated with hydrocarbon metabolism. PL-A contained 5.82×10^6 copies/g of
5 toluene dioxygenase (*tdo*) and 3.69×10^4 copies/g of alkane monooxygenase (*alkB*). In
6 contrast, *tdo* and *alkB* copy numbers in PL-D were below the detection limit. The
7 abundance of one carbon pathway (C1) groups based on quantification of methyl
8 coenzyme M reductase alpha (*mcrA*) and particulate methane monooxygenase alpha
9 (*pmoA*) subunit copy number did not vary considerably between sampling intervals,
10 reaching 8.51×10^5 and 1.78×10^3 copies/g in PL-A and 3.69×10^5 and 4.99×10^2
11 copies/g in PL-D respectively.

12 Within the archaeal domain, numerous SSU rRNA gene sequences affiliated with
13 anaerobic methane oxidizing archaea ANME-1a, as well as novel and deeply branching
14 lineages proximal to this group, e.g. Tar ARC I (Fig. 3 and 4) were recovered from PL,
15 OP and MV. SSU rRNA gene sequences affiliated with classical methanogens were not
16 recovered in PL, although they were identified in OP and MV (Fig. 3 and 4). Consistent
17 with these results, the majority of sequences recovered from *mcrA* clone libraries derived
18 from PL samples were affiliated with novel and deeply branching lineages proximal to
19 previously described ANME-1 subgroups (Hallam et al., 2003) (Fig. A1 of Appendix).
20 Sequences affiliated with thermoacidophilic and metal-respiring *Thermoplasmatales*
21 dominated all PL libraries and collectively defined three novel lineages, e.g. Tar ARC II,
22 Tar ARC III and Tar ARC IV (Fig. 3 and 4). Additional minority sequences affiliated
23 with uncultivated lineages from South African gold mine (SAGMEG) (Fig. 3 and 4) and
24 marine sedimentary environments (marine benthic group or MBG) were recovered from
25 MV and PL-C (Fig. 4 and A2 in the Appendix).

26 Within the bacterial domain we identified a number of SSU rRNA gene sequences
27 closely affiliated with chemolithoautotrophic sulfur-oxidizing *Thiotricales* and
28 *Campylobacteriales* (Fig. 4 and A3 in Appendix). Additional sequences affiliated with
29 sulfur-reducing *Defferibacteriales* recovered exclusively from PL-B, and sulfate-reducing
30 *Thermodesulfobacteriales*, *Nitrospirales*, *Desulfurimonadales* and *Desulfobacteriales*
31 were also identified, consistent with an active sulfur cycle (Fig 4, and A3 and A4 in
32 Appendix). However, distribution patterns of sulfur metabolizing bacteria were extremely
33 variable with minimal overlap between PL, OP and MV samples. More compositional
34 similarity was observed for sequences affiliated with higher alkane, aromatic and heavy
35 oil degradation, including *Pseudomonadales*, *Oceanospirillales*, and *Burkholderiales*
36 (Fig. 4 and A3 in Appendix). Additional sequences affiliated with desiccation resistant
37 *Acidobacteriales* and *Rhodospirillales* were recovered exclusively from inactive PL-C
38 (Fig. 4 and A4 in Appendix).

39 To place PL, OP, and MV microbiota into a broader ecological and
40 biogeochemical context we compared microbial compositional profiles between
41 hydrocarbon environments including tar pits (Kim and Crowley, 2007), methane seeps
42 (Orphan et al., 2001; Lloyd et al., 2006; Knittel et al., 2005), hydrothermal vents (Teske
43 et al., 2002; Brazelton et al., 2006), mud volcanoes (Alain et al., 2006; Lösekann et al.,
44 2007; Kormas et al., 2008), and petroleum reservoirs (Orphan et al, 2000; Pham et al,
45 2009; Fig. 4). PL, OP and MV clustered together to the exclusion of other environments
46 (Fig. 4). The archaeal component of PL was extremely divergent, dominated by novel

1 and deeply branching ANME-1 and Tar ARC lineages and reduced representation or
2 absence of classical methanogens, halophiles, and marine Group I crenarchaea. The
3 bacterial component of PL profiles was intermediate to biodegraded oil systems, methane
4 seeps and marine mud volcanoes with reduced representation of sulfate-reducing
5 δ -proteobacteria and conspicuous absence of most candidate divisions (Fig. 4). These
6 observations are broadly consistent with diverse syntrophic partnerships and electron
7 transfer processes associated with archaeal energy metabolism *e.g.* anaerobic methane
8 oxidation versus methanogenesis, in PL and environs.

10 Discussion

11
12 Pitch Lake provides a contemporary terrestrial example of an active microbial
13 community adapted for persistence and growth within a recalcitrant hydrocarbon matrix.
14 Understanding the molecular nature of these adaptations with respect to nutritional,
15 energetic and detoxification services could have important technological implications for
16 recovery and remediation efforts in biodegraded hydrocarbon reservoirs including heavy
17 oil, natural bitumen, and oil shale deposits (Head et al., 2003; Aitken et al., 2004). While
18 anaerobic hydrocarbon degradation under methanogenic conditions has been reported in
19 formation waters of deep subsurface petroleum reservoirs (Jones et al., 2008; Dolfing et
20 al., 2008), microbial community structure, carbon isotope ratios and water activity
21 measurements of Pitch Lake samples suggest variations on this theme. It will be of
22 particular interest to elucidate the biogeochemical roles of ANME-1 and Tar ARC
23 lineages in the degradation process.

24 From an astrobiological perspective, Pitch Lake represents a unique opportunity
25 to evaluate the critical limits for life in the universe (Schulze-Makuch and Irwin, 2006;
26 Shapiro and Schulze-Makuch, 2009). In particular, Pitch Lake may serve as a useful
27 analog for evaluating the potential for life in liquid hydrocarbon lakes discovered by the
28 Cassini Titan Radar Mapper on Saturn's largest moon, Titan (Elachi et al., 2006; Stofan
29 et al., 2007). Spectroscopic results also indicate the presence of methane rain on Titan
30 (Lunine et al., 1983, Lorenz, 2000) and imply a methane cycle on Titan analogous to the
31 hydrological cycle on Earth. A hydrocarbon solvent such as methane and ethane on
32 Titan's surface may improve the chances for the origin of life on Titan, based on
33 extensive experiences with organic synthesis reactions that have shown that the presence
34 of water greatly diminishes the chance of constructing nucleic acids (Schulze-Makuch
35 and Irwin, 2008). Organic reactivity in hydrocarbon solvents is no less versatile than in
36 water, and many enzymes derived from microorganisms are believed to catalyze
37 reactions by having an active site that is hydrophobic (Benner et al., 2004). Furthermore,
38 Baross et al (2007) recently suggested that the environment of Titan meets the absolute
39 requirements for life, which include thermodynamic disequilibrium, abundant carbon
40 containing molecules and heteroatoms, and a fluid environment – further concluding that
41 “this makes inescapable the conclusion that if life is an intrinsic property of chemical
42 reactivity, life should exist on Titan.” Aside from the very cold surface conditions on
43 Titan, the environmental conditions within Pitch Lake are one of the closest analog
44 environments we can find on our planet, and the discovery of a broad spectrum of
45 microbiota enhances the possibility for life in Titan's hydrocarbon lakes. In fact,
46 cryovolcanism has been inferred to exist on Titan (Sotin et al., 2005; Lopes et al., 2007)

1 and a heating of some of Titan's hydrocarbon reservoirs from below is a distinct
2 possibility further enhancing the possibility of at least pre-biotic organic reactions.

3 The absolute requirement of water for life is of much astrobiological interest and
4 should be a topic of further inquiries. In regard to Pitch Lake further research is
5 encouraged to elucidate on the adaptation mechanisms that life in Pitch Lake might
6 employ to thrive in the water-poor hydrocarbon matrix. The recent findings that *E. coli*
7 cells are able to generate up to 70 % of their intracellular water during metabolism
8 (Kreuzer-Martin et al., 2005) and that the fungus *Fusarium alkanophilum* is capable of
9 thriving in a hydrocarbon environment extracting water from light hydrocarbons
10 (Marcano et al., 2002), may point to exciting new directions of how life can adapt to life
11 in a hydrocarbon matrix with little or no liquid water.

12 13 **Conclusions**

14
15 We describe a unique, endemic microbial community in Pitch Lake, a natural asphalt lake
16 in Trinidad. The microbial organisms detected are archaea and bacteria, and thrive in a
17 hydrocarbon matrix with low water activity and toxic compounds. The environment is in
18 many aspects analogous to the surface of Titan, on which hydrocarbon lakes have been
19 detected under a thick atmosphere dominated by nitrogen and methane. Our research as
20 described above is a starting point on investigating what life's principle constraints are in
21 a hydrocarbon matrix and whether the hydrocarbon lakes on Titan could possibly contain
22 life.

23 24 **Acknowledgements**

25
26 This work was supported by a grant from the Government of Trinidad & Tobago
27 (GROTT), the Natural Sciences and Engineering Research Council (NSERC) of Canada
28 (328256-07 and STPSC 356988), Canada Foundation for Innovation (CFI 17444) and the
29 Canadian Institute for Advanced Research (CIFAR). We thank Dr. Tom Kieft for his help
30 in designing the gas sampling procedure, Dr. Barbara Sherwood Lollar for conducting the
31 gas analyses, and Drs. Alison Skelley and Christine Jayarajah for conducting amino acid
32 analyses with the Mars Organic Analyzer (MOA). We also thank Rainer Strzoda from the
33 Siemens AG to let us test his hand-held laser system for the field detection of methane
34 gas. Archaeal and bacterial SSU rRNA and *mcrA* gene sequences were deposited in
35 GenBank under the accession numbers GU120478-GU120542, GU120543-GU120617
36 and GU447199-GU447228, respectively.

1 **References**

2
3 Abbas, O. and Schulze-Makuch, D. (2002) Acetylene-based pathways for prebiotic
4 evolution on Titan. 2nd European Workshop on Exo-Astrobiology (EANA/ESA), Graz,
5 Austria, pp. 349-352.

6
7 Aitken, C.M., Jones, D.M., and Larter, S.R. (2004) Anaerobic hydrocarbon
8 biodegradation in deep subsurface oil reservoirs. *Nature* 431, 291-294.

9
10 Alain K.H.T., Musat F, Elvert M, Treude, T. and Kruger, M. (2006) Microbiological
11 investigation of methane- and hydrocarbon-discharging mud volcanoes in the Carpathian
12 Mountains, Romania. *Environmental Microbiology* 8, 574-590.

13
14 Baldwin, B.R., Nakatsu, C.H., and Nies, L. (2003) Detection and enumeration of
15 aromatic oxygenase genes by multiplex and real-time PCR. *Applied and Environmental*
16 *Microbiology* 69, 3350-3358.

17
18 Baross, J.A., Benner, S.A., Cody, G.D., Copley, S.D., Pace, N.R., Scott, J.H., Shapiro, R.,
19 Sogin, M.L., Stein, J.L., Summons, R., and Szostak, J.W. (2007) The Limits of Organic
20 Life in Planetary Systems. National Academies Press, Washington, D.C.

21
22 Benner, S.A., Ricardo, A., and Carrigan, M.A. (2004) Is there a common chemical model
23 for life in the universe? *Curr. Opin. Chem. Biol.* 8, 672-689.

24
25 Brazelton, W.J., Schrenk, M.O., Kelley, D.S., and Baross, J.A. (2006) Methane- and
26 sulfur-metabolizing microbial communities dominate the Lost City hydrothermal field
27 ecosystem. *Appl Environ Microbiol* 72, 6257-6270.

28
29 DeSantis, T.Z., Hugenholtz, P., Larsen, N., Rojas, M., Brodie, E.L., Keller, K., Huber,
30 T., Dalevi, D., Hu, P., and Andersen, G.L. (2006) Greengenes, a chimera-checked 16S
31 rRNA gene database and workbench compatible with ARB. *Appl Environ Microbiol* 72,
32 5069-5072.

33
34 Dolfing, J., Larter, S.R., and Head, I.M. (2008) Thermodynamic constraints on
35 methanogenic crude oil biodegradation. *ISME J* 2, 442-452.

36
37 Edgar, R.C. (2004) MUSCLE: multiple sequence alignment with high accuracy and high
38 throughput. *Nucleic Acids Res* 32, 1792-1797.

39
40 Elachi, C., Wall, S., Janssen, M., Stofan, E., Lopes, R., Kirk, R., Lorenz, R., Lunine, J.,
41 Paganelli, F., Soderblom, L., Wood, C., Wye, L., Zebker, H., Anderson, Y., Ostro, S.,
42 Allison, M., Boehmer, R., Callahan, P., Encrenaz, P., Flamini, E., Francescetti, G., Gim,
43 Y., Hamilton, G., Hensley, S., Johnson, W., Kelleher, K., Muhleman, D., Picardi, G.,
44 Posa, F., Roth, L., Seu, R., Shaffer, S., Stiles, B., Vetrilla, S., and West, R. (2006) Titan
45 Radar Mapper observations from Cassini's T3 fly-by. *Nature* 441, 709-713.

46

1 Felsenstein, J. (2005) PHYLIP (Phylogeny Inference Package) Distributed by the author,
2 Department of Genome Sciences, University of Washington, Seattle, WA, USA.
3
4 Grant, W.D. (2004) Life at low water activity. *Philos Trans R Soc Lond B Biol Sci* 359,
5 1249-1266; discussion 1266-1267.
6
7 Guindon, S., Lethiec, F., Duroux, P., and Gascuel, O. (2005) PHYML Online-a web
8 server for fast maximum likelihood-based phylogenetic inference. *Nucleic Acids Res* 33
9 (Web Server issue), W557-559.
10
11 Hales, B.A., Edwards, C., Ritchie, D.A., Hall, G., Pickup, R.W., and Saunders, J.R.
12 (1996) Isolation and identification of methanogen-specific DNA from blanket bog peat
13 by PCR amplification and sequence analysis. *Appl Environ Microbiol* 62, 668-675.
14
15 Hallam, S.J., Girguis, P.R., Preston, C.M., Richardson, P.M., and DeLong, E.F. (2003)
16 Identification of methyl coenzyme M reductase A (mcrA) genes associated with
17 methane-oxidizing archaea, *Appl Environ Microbiol* 69, 5483-5491.
18
19 Head, I.M., Jones, D.M., and Larter, S.R. (2003) Biological activity in the deep
20 subsurface and the origin of heavy oil. *Nature* 426, 344-352.
21
22 Huber, T., Faulkner, G., and Hugenholtz, P. (2004) Bellerophon: a program to detect
23 chimeric sequences in multiple sequence alignments. *Bioinformatics* 20, 2317-2319.
24
25 Jones, D.M., Head, I.M., Gray, D., Adams, J.J., Rowan, A.K., Aitken, C.M., Bennett, B.,
26 Huang, H., Brown, A., Bowler, B.F.J., Oldenburg, T., Erdmann, M., and Larter, S.R.
27 (2008) Crude-oil biodegradation via methanogenesis in subsurface petroleum reservoirs.
28 *Nature* 451, 176-180.
29
30 Kim, J.S. and Crowley, D.E. (2007) Microbial diversity in natural asphalts of the Rancho
31 La Brea Tar Pits. *Appl Environ Microbiol* 73, 4579-4591.
32
33 Knittel, K., Losekann, T., Boetius, A., Kort, R., and Amann, R. (2005) Diversity and
34 distribution of methanotrophic archaea at cold seeps. *Applied and Environmental*
35 *Microbiology* 71, 467-479.
36
37 Kormas, K.A., Meziti, A., Dahlmann, A., De Lange, G.J., and Lykousis, V. (2008)
38 Characterization of methanogenic and prokaryotic assemblages based on mcrA and 16S
39 rRNA gene diversity in sediments of the Kazan mud volcano (Mediterranean Sea).
40 *Geobiology* 6, 450-460.
41
42 Kreuzer-Martin, H.W., Ehleringer, J.R., and Hegg, E.L. (2005) Oxygen isotopes indicate
43 most intracellular water in long-phase *Escherichia coli* is derived from metabolism.
44 *PNAS (USA)* 102, 17337-17341.
45

- 1 Lloyd, K.G., Lapham, L., and Teske, A. (2006) An anaerobic methane-oxidizing
2 community of ANME-1b archaea in hypersaline Gulf of Mexico sediments. *Appl Environ*
3 *Microbiol* 72, 7218-7230.
4
- 5 Lopes, R.M.C., Mitchell, K.L., Stofan, E.R., Lunine, J.I., Lorenz, R., Paganelli, F., Kirk,
6 R.L., Wood, C.A., Wall, S.D., Robshaw, L.E., Fortes, A.D., Neish, C.D. Radebaugh, J.,
7 Reffet, E., Ostro, S.J., Elachi, C., Allison, M.D., Anderson, Y., Boehmer, R., Boubin, G.,
8 Callahan, P., Encrenaz, P., Flamini, E., Francescetti, G., Gim, Y., Hamilton, G., Hensley,
9 S., Janssen, M.A., Johnson, W.T.K., Kelleher, K., Muhleman, D.O., Ori, G., Orosei, R.,
10 Picardi, G., Posa, F., Roth, L.E., Seu, R., Shaffer, S., Soderblom, L.A., Stiles, B.,
11 Vetrella, S., West, R.D., Wye, L. and Zebker, H.A. (2007) Cryovolcanic features on
12 Titan's surface as revealed by the Cassini Titan Radar Mapper. *Icarus* 186, 395-412.
13
- 14 Lorenz, R.D. (2000) Post-Cassini exploration of Titan: science rationale and mission
15 concepts. *JBIS* 53, 218-234.
16
- 17 Lösekann, T., Knittel, K., Nadalig, T., Fuchs, B., Niemann, H., Boetius, A., and Amann,
18 R. (2007) Diversity and abundance of aerobic and anaerobic methane oxidizers at the
19 Haakon Mosby mud volcano, Barents Sea. *Applied and Environmental Microbiology* 73,
20 3348-3362.
21
- 22 Lozupone, C., Hamady, M., & Knight, R. (2006) UniFrac-an online tool for comparing
23 microbial community diversity in a phylogenetic context. *BMC Bioinformatics* 7, 371.
24
- 25 Lozupone, C. and Knight, R. (2005) UniFrac: a new phylogenetic method for comparing
26 microbial communities. *Appl Environ Microbiol* 71, 8228-8235.
27
- 28 Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., and 27 other authors (2004)
29 ARB: a software environment for sequence data. *Nucleic Acids Research* 32, 1363-1371.
30
- 31 Lunine, J.I., Stevenson, D.J., and Yung, Y.L. (1983) Ethane ocean on Titan. *Science* 222,
32 1229-1230.
33
- 34 Maddison, D.R. and Maddison, W.P. (2003) *MacClade 4*. Sinauer Associates, Inc.,
35 Sunderland.
36
- 37 Marcano, V., Benitez, P., and Palacios-Pru, E. (2002) Growth of lower eukaryote in non-
38 aromatic hydrocarbon media \geq C-12 and its exobiological significance. *Planet. Space Sci*
39 50, 693-709.
40
- 41 McKay, C.P. and Smith, H.D. (2005) Possibilities for methanogenic life in liquid
42 methane on the surface of Titan. *Icarus* 178, 274-276.
43
- 44 Mikucki, J.A., Pearson, A., Johnston, D.T., Turchyn, A.V., Farquhar, J., Schrag, D.P.,
45 Anbar, A.D., Priscu, J.C., and Lee, P.A. (2009) A contemporary microbially maintained
46 subglacial ferrous "ocean". *Science* 324, 397-400.

1
2 Orphan, V.J., Taylor, L.T., Hafenbradl, D., and DeLong, E.F. (2000) Culture-dependent
3 and culture-independent characterization of microbial assemblages associated with high-
4 temperature petroleum reservoirs. *Applied and Environmental Microbiology* 66, 700-711.
5
6 Orphan, V.J., Hinrichs, K.-U., Ussler, W., Paull, C.K., Taylor, L.T., Sylva, S.P., Hayes,
7 J.M., and DeLong, E.F. (2001) Comparative analysis of methane-oxidizing archaea and
8 sulfate-reducing bacteria in anoxic marine sediments. *Applied and Environmental*
9 *Microbiology* 67, 1922-1934.
10
11 Peckham, S.F. (1895) VII. - On the Pitch Lake of Trinidad. *Geological Magazine*
12 (Decade IV) 2, 416-425.
13
14 Pham, V.D., Hnatow, L.L., Zhang, S., Fallon, R.D., Jackson, S.C., Tomb, J.-F., DeLong,
15 E.F., and Keeler, S.J. (2009) Characterizing microbial diversity in production water from
16 an Alaskan mesothermic petroleum reservoir with two independent molecular methods.
17 *Environmental Microbiology* 11, 176-187.
18
19 Priscu, J.C., Fritsen, C.H., Adams, E.E., Giovannoni, S.J., Paerl, H.W., McKay, C.P.,
20 Doran, P.T., Gordon, D.A., Lanoil, B.D., and Pinckney, J.L. (1998) Perennial Antarctic
21 lake ice: an oasis for life in a polar desert. *Science* 280, 2095-2098.
22
23 Pruesse, E.C., Quast, C., Knittel, K., Fuchs, B.M, Ludwig, W., Peplies, J. and Glöckner,
24 F.O. (2007) SILVA: a comprehensive online resource for quality checked and aligned
25 ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res*, 35, 7188-7196.
26
27 Schulze-Makuch, D. and Irwin, L.N. (2006) The prospect of alien life in exotic forms on
28 other worlds. *Naturwissenschaften* 93, 155-172.
29
30 Schulze-Makuch, D. and Irwin, L. (2008) Life in the Universe: Expectations and
31 Constraints. 2nd edition, Springer, Berlin.
32
33 Shapiro, R.S. and Schulze-Makuch, D. (2009) The search for alien life in our solar
34 system: strategies and priorities. *Astrobiology* 9, 335-343.
35
36 Skelley, A.M. et al (2005) Development and evaluation of a microdevice for amino acid
37 biomarker detection and analysis on Mars. *PNAS* 102, 1041-1046.
38
39 Sotin, C., Jaumann, R., Buratti, B.J., Brown, R.H., Clark, R.N., Soderblom, L.A., Baines,
40 K.H., Bellucci, G., Bibring, J.-P., Capaccioni, F., Cerroni, P., Combes, M., Coradini, A.,
41 Cruikshank, D.P., Drossart, P., Formisano, V., Langevin, Y., Matson, D.L., McCord,
42 T.B., Nelson, R.M., Nicholson, P.D., Sicardy, B., LeMouelic, S., Rodriguez, S., Stephan,
43 K., and Scholz, C.K. (2005) Release of volatiles from a possible cryovolcano from near-
44 infrared imaging of Titan. *Nature* 435, 786-789.
45

1 Stofan, E.R., Elachi, C., Lunine, J.I., Lorenz, R.D., Stiles, B., and 34 co-authors (2007)
2 The lakes of Titan. *Nature* 445, 61-64.
3
4 Teske, A., Hinrichs, K.-U., Edgcomb, V., de Vera Gomez, A., Kysela, D., Sylva, S.P.,
5 Sogin, M.L., and Jannasch, H.W. (2002) Microbial diversity of hydrothermal sediments
6 in the Guaymas Basin: evidence for anaerobic methanotrophic communities. *Appl*
7 *Environ Microbiol* 68 , 1994-2007.

8 Tosca, N.J., Knoll, A.H., and McLennan, S.M. (2008) Water activity and the challenge
9 for life on early Mars. *Science* 320, 1204-1207.

10 Van Well, B., Murray, S., Hodgkinson, J., Pride, R., Strzoda, R., Gibson, G., and Padgett,
11 M. (2005) An open-path, hand-held laser system for the detection of methane gas. *J. Opt.*
12 *A: Pure Appl. Opt.* 7, S420-S424.

13 White, D.C., Davis, W.M., Nickels, J.S., King, J.D., and Bobbie, R.J. (1979)
14 Determination of the sedimentary microbial biomass by extractable lipid phosphate.
15 *Oecologia* 40, 51-62.
16
17 Zaikova, E., Walsh, D.A., Stilwell, C.P., Mohn, W.W., Tortell, P.D., and Hallam, S.J.
18 (2010) Microbial community dynamics in a seasonally anoxic fjord: Saanich Inlet, British
19 Columbia. *Environ Microbiol* 12, 172-191.

20

21

22

23

24

25

26

27

28

29

1 **Table 1.** Sampling information

Sample	Latitude (N) WGS84	Longitude (W) WGS84	Temperature (°C)	Depth (cm)	Description
PL-A	10° 14' 4.9"	61° 37' 36.9"	35.6	24	active bubbling low viscosity
PL-B	10° 13' 59.4"	61° 37' 46.1"	55.9	10	semi-liquid high viscosity
PL-C	10° 14' 59.4"	61° 37' 46.1"	44.2	10	inactive hard
PL-D	10° 14' 4.5"	61° 37' 37.1"	31.6	84	active bubbling low viscosity
PL-E	10° 13' 59"	61° 37' 46.0"	37.0	5	active low viscosity
PL-F	10° 14' 0.9"	61° 37' 46.1"	36.8	5	active low viscosity
OP	10° 13' 13.2"	61° 34' 13.3"	ND	0	seeping oil well
MV	10° 15' 51.3"	61° 18' 18.6"	27.7	ND	mud volcano

2 * ND = not determined

1 **Table 2.** Stable isotope ratios

	Stable Isotope Ratio		
Samples	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	$\delta^{34}\text{S}$
PL	-28.1 ‰ ± 0.4 ‰	3.1 ‰ ± 0.1 ‰	10.1 ‰ ± 0.1 ‰
OP	-28.5 ‰ ± 0.1 ‰	3.2 ‰ ± 0.2 ‰	5.6 ‰ ± 0.1 ‰
MV	-15.6 ‰ ± 0.1 ‰	4.6 ‰ ± 0.1 ‰	-4.6 ‰ ± 1.2 ‰

2 Note: Bulk carbon and nitrogen isotopic results reported in per mill relative to VPDB
 3 (Vienna Peedee belemnite) and air, and the sulfur isotope values are in per mill relative to
 4 the V-CDT (Vienna - Canyon Diablo Troilite) isotope ratio standard using NBS-127
 5 (Barium Sulphate) as the calibration material. Error limit denotes one standard deviation.

1 **Table 3.** PLFA Analysis


	Sample			
	PL-A	PL-B	PL-C	PL-D
Biomass	5.64 x 10 ⁶	7.54 x 10 ⁶	2 x 10 ⁷	3.73 x 10 ⁶
% Mono-saturated	18.64	21.36	43.86	0
Mono-Biomarkers	18:1w7c	18:1w7c cy19:0	16:1w7c 16:1w5c cy17:0 18:1w7c cy19:0	none
% Normal Saturated	81.36	78.64	56.14	100
Nsats Biomarkers	16:0	16:0 18:0	16:0 18:0	16:0

1 **Table 4.** Archaeal and bacterial SSU rRNA gene abundance (copies/g)

Sample	Total Archaea	Total Bacteria
PL-A	$2.67 \times 10^4 \pm 6.10 \times 10^3$	$1.42 \times 10^7 \pm 1.34 \times 10^6$
PL-B	$3.19 \times 10^4 \pm 2.64 \times 10^3$	$2.13 \times 10^6 \pm 3.10 \times 10^5$
PL-C	$3.27 \times 10^4 \pm 6.03 \times 10^3$	$2.46 \times 10^7 \pm 1.25 \times 10^6$
PL-D	$2.33 \times 10^5 \pm 1.62 \times 10^4$	$1.82 \times 10^7 \pm 1.12 \times 10^6$
OP	$1.65 \times 10^3 \pm 4.57 \times 10^2$	$1.94 \times 10^5 \pm 6.06 \times 10^4$
MV	$1.11 \times 10^4 \pm 2.48 \times 10^3$	$4.28 \times 10^6 \pm 2.66 \times 10^5$

1 Figures

Figure 1




2

3

4 **Fig 1.** (A) Aerial photograph of Pitch Lake. Sampling sites (PL-A through F) marked by
5 black circles (see Table S1 for sampling parameters). Samples OP and MV fall outside
6 the boundaries of this view. (B and C) Bubbles of varying sizes form on the surface of the
7 liquid asphalt under varying viscosities. (D) Terrestrial mud volcano at Devil's
8 Woodyard (MV). Scale bars indicate approximately 5 cm.


Figure 2



1

2 **Fig 2.** Principle coordinate analysis of (A) bacterial and (B) archaeal SSU rRNA gene
3 sequence profiles based on UniFrac distance matrix. The circumference of each circle in
4 the ordination corresponds to the square root of SSU rRNA gene copies/g (see scale bar)
5 measured for PL, OP and MV samples.

Figure 3




1

2 **Fig 3.** Distance tree of euryarchaeal SSU rRNA gene sequences recovered from PL, OP
3 and MV samples. Bootstrap values (%) are based on 100 replicates using the maximum
4 likelihood method and are shown for branches with greater than 50% support. The scale
5 bar represents 0.1 substitutions per site. Sequence distribution across samples is
6 represented by a series of closed circles whose circumference indicates the percentage of
7 identified RFLP patterns falling within a particular taxonomic group. The number of
8 SSU rRNA clones screened per samples is PL-A = 56(14), PL-B = 84(7), PL-C = 92(19),
9 PL-D = 89(15), OP = 93(10), MV = 78(16). Numbers in parenthesis indicate unique

- 1 RFLP patterns associated with each sample library. Sequences that belong to a specific
- 2 subgroup were highlighted as shown on the legend.

Figure 4




- 3
- 4

5 **Fig 4.** Heatmap visualization of archaeal and bacterial composition profiles based on
 6 SSU rRNA gene sequences recovered from PL, OP and MV samples and related
 7 hydrocarbon environments including Rancho La Brea Tar Pit (LB), petroleum reservoirs
 8 in Alaska (AK) and the California Coast (CA), whale fall in Monterey Cayon (MC),
 9 methane seeps including Gulf of Mexico (GoM), Eel River Basin (ER), and Hydrate
 10 Ridge (HR), hydrothermal vents including Lost City (LC) and Guaymas Basin (GB), the
 11 terrestrial mud volcano Paclele Mici (PM), and the marine mud volcanoes Haakon
 12 Mosby (HMMV), and Kazan (KMV). Color intensity in each box corresponds to the
 13 natural logarithm of total SSU rRNA gene sequences affiliated with identified taxonomic
 14 groups. Heatmap rows are organized according to the topology of the maximum
 15 likelihood tree. The histogram depicts the number of boxes representing the corrected
 16 values.


17

18 **Appendix/Supplementary Material**



1
2
3
4
5
6
7
8
9
10
11
12
13
14


Fig S1. Distance tree of *mcrA* amino acid sequences recovered from PL, OP and MV samples rooted with *Methanopyrus kandleri*. Bootstrap values (%) are based on 100 replicates using the maximum likelihood method and are shown for branches with greater than 50% support. The scale bar represents 0.1 substitutions per site. Sequence distribution across samples is represented by a series of closed circles whose circumference indicates the percentage of identified RFLP patterns falling within a particular *mcrA* lineage. The number of *mcrA* clones screened per samples is PL-A = 81(15), PL-B = 82(14), PL-D = 79(17), OP = 85(11) and MV = 79(17). Numbers in parenthesis indicate unique RFLP patterns associated with each sample library. Sequences that belong to specific *mcrA* subgroup were highlighted as shown on the legend.





1
2 **Fig S2.** Distance tree of crenarchaeal SSU rRNA gene sequences recovered from PL, OP
3 and MV samples. Bootstrap values (%) are based on 100 replicates using the maximum
4 likelihood method and are shown for branches with greater than 50% support. The scale
5 bar represents 0.08 substitutions per site. Sequence distribution across samples is
6 represented by a series of closed circles whose circumference indicates the percentage of
7 identified RFLP patterns falling within a particular taxonomic group. The number of
8 SSU rRNA clones screened per samples is PL-A = 56(14), PL-B = 84(7), PL-C = 92(19),
9 PL-D = 89(15), OP = 93(10), MV = 78(16). Numbers in parenthesis indicate unique

- 1 RFLP patterns associated with each sample library. Sequences that belong to specific
- 2 subgroups of interest are highlighted in the legend.

1



2
 3 **Fig S3.** Distance tree of proteobacterial SSU rRNA gene sequences recovered from PL,
 4 OP and MV samples. Bootstrap values (%) are based on 100 replicates using the
 5 maximum likelihood method and are shown for branches with greater than 50% support.
 6 The scale bar represents 0.2 substitutions per site. Sequence distribution across samples is
 7 represented by a series of closed circles whose circumference indicates the percentage of
 8 identified RFLP patterns falling within a particular taxonomic group. The number of
 9 SSU rRNA clones screened per samples is PL-A = 89(13), PL-B = 86(12), PL-C =
 10 84(12), PL-D = 79(10), OP = 79(16), MV = 79(14). Numbers in parenthesis indicate
 11 unique RFLP patterns associated with each sample library. Sequences that belong to
 12 specific subgroups of interest are highlighted in the legend.
 13



1
 2 **Fig S4.** Distance tree of non-proteobacterial SSU rRNA gene sequences recovered from
 3 PL, OP and MV samples. Bootstrap values (%) are based on 100 replicates using the
 4 maximum likelihood method and are shown for branches with greater than 50% support.
 5 The scale bar represents 0.3 substitutions per site. Sequence distribution across samples is
 6 represented by a series of closed circles whose circumference indicates the percentage of
 7 identified RFLP patterns falling within a particular taxonomic group. The number of
 8 SSU rRNA clones screened per samples is PL-A = 89(13), PL-B = 86(12), PL-C =
 9 84(12), PL-D = 79(10), OP = 79(16), MV = 79(14). Numbers in parenthesis indicate
 10 unique RFLP patterns associated with each sample library.
 11