

Mefentrifluconazole	Group	3	Fungicide
Pyraclostrobin	Group	11	Fungicide

## Supplemental Label

# Veltyma™

## Fungicide

**For disease control in grasses, grass grown for seed, non-grass forages, oilseeds, sugar beet, and sugarcane**

This supplemental label expires October 30, 2023, and must not be used or distributed after this date.

### Active Ingredients\*:

mefentrifluconazole: 2-[4-(4-chlorophenoxy)-2-(trifluoromethyl)phenyl]-1-(1H-1,2,4-triazole-1-yl)propan-2-ol .....	17.56%
pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester).....	17.56%
<b>Other Ingredients**:</b> .....	64.88%
<b>Total:</b> .....	100.00%

\* Veltyma™ fungicide contains 1.67 lbs mefentrifluconazole and 1.67 lbs pyraclostrobin per gallon.

**EPA Reg. No. 7969-409**  
**CAUTION/PRECAUCION**

### Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Veltyma™ fungicide** container label, EPA Reg. No. 7969-409, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Veltyma** before applying.
- Use of **Veltyma** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Veltyma**.

### Application Instructions

Refer to the **Veltyma** container label for additional instructions and restrictions.

## Crop Rotation Restrictions

**Rotational Crops:** Please see the following table for crop rotational restrictions.

Crop or Crop Group	Rotation Interval
Brassicas including broccoli, cauliflower and head cabbages (crop group 5-16) Bulb vegetables including bulb onion, spring onion and garlic (crop group 3-07) Cereals, including wheat, oats, barley, triticale, rye, rice and corn (crop group 15 and crop group 16) Cucurbits including cucumber, squash and melons (crop group 9) Fruiting vegetables including tomato, eggplant, peppers (crop group 8-10) Fresh herbs Grass and non-grass animal feeds including alfalfa and clover (crop group 17 and crop group 18) Leafy vegetables including lettuces, spinach and leafy cabbages (crop group 4-16) Legume vegetables including soybean (crop group 6 and crop group 7) Low-growing berries including strawberries (crop group 13-07G) Oilseeds including cotton, sunflower and canola (crop group 20) Peanut Root and tuber vegetables including carrot, potato, beets and sugar beet (crop group 1 and crop group 2) Stalk, stem and leaf petiole vegetables including celery and asparagus (crop group 22) Sugarcane Any other crop labeled for direct application of a product containing mefenftrifluconazole	May be planted immediately following the last application
Other food and feed crops, not listed above	May not be planted in rotation

## Detailed Use Directions

### Labeled Crops

Foliar

- When conditions favor disease or disease pressure is high:
  - Use the shorter specified interval.
  - For a rate range, use the higher specified rate.
- **DO NOT** make more than two (2) sequential applications of **Veltyma™ fungicide** before alternating to a labeled **non-Group 3** or **non-Group 11** fungicide, unless otherwise specified in the following crop-specific use sections.

#### Use Rate Conversion

fl ozs product/A	lb mefentrifluconazole/A	lb pyraclostrobin/A
5	0.07	0.07
7	0.09	0.09
8	0.10	0.10
10	0.13	0.13
13.5	0.18	0.18
20	0.26	0.26
30	0.39	0.39
40.5	0.53	0.53

**State-specific Restrictions** - Not for sale, distribution, or use in **Nassau and Suffolk counties in New York State.**

## Grasses

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Anthracnose <i>Colletotrichum</i> spp.	7 to 10	30	0
	Leaf Spot/Blight <i>Bipolaris</i> spp., <i>Cochliobolus</i> spp., <i>Drechslera</i> spp., <i>Stagonospora</i> spp.			
	Net Blotch <i>Drechslera</i> spp., <i>Helminthosporium</i> spp.			
	Powdery Mildew <i>Erysiphe</i> spp.			
	Purple Leaf Spot <i>Stagonospora</i> spp.			
	Rust <i>Puccinia</i> spp., <i>Uromyces dactylidis</i>			
	Septoria Leaf Spot <i>Septoria</i> spp.			
<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply at 14-day to 21-day intervals.</li> </ul>				

### Use Restrictions

- **DO NOT** apply more than 10 fl ozs (0.13 lb mefentrifluconazole, 0.13 lb pyraclostrobin) per acre per application.
- **DO NOT** make more than 3 applications at 10 fl ozs per acre per year.
- **DO NOT** apply more than 30 fl ozs (0.39 lb mefentrifluconazole, 0.39 lb pyraclostrobin) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.73 lb ai/acre/year of pyraclostrobin-containing products.

**Detailed Grasses (crop group 17) Crop List** - any grass, Gramineae family (either green or cured) that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage, except sugarcane and those included in the cereal grains group.

## Grass Grown for Seed

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Powdery mildew <i>Erysiphe</i> spp.	7 to 10	20	0
	Rust <i>Puccinia</i> spp.			
<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply at 14-day to 21-day intervals.</li> </ul>				

### Use Restrictions

- **DO NOT** apply more than 10 fl ozs (0.13 lb mefentrifluconazole, 0.13 lb pyraclostrobin) per acre per application.
- **DO NOT** make more than 2 applications at 10 fl ozs per acre per year.
- **DO NOT** apply more than 20 fl ozs (0.26 lb mefentrifluconazole, 0.26 lb pyraclostrobin) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.73 lb ai/acre/year of pyraclostrobin-containing products.

**Detailed Grass Grown for Seed (crop group 17) Crop List** - any grass, Gramineae family (either green or cured) that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage, except sugarcane and those included in the cereal grains group.

## Non-grass Forages

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Anthrachnose <i>Colletotrichum trifolii</i>	7 to 10	30	14
	Common leaf spot <i>Pseudopeziza medicaginis</i>			
	Downy mildew <i>Peronospora trifoliorum</i>			
	Leaf spot <i>Leptosphaerulina briosiani</i>			
	Powdery mildew <i>Erysiphe</i> spp.			
	Rhizoctonia blight/black patch <i>Rhizoctonia</i> spp.			
	Rust <i>Phakopsora</i> spp., <i>Puccinia</i> spp., <i>Uromyces</i> spp.			
	Spring black stem and leaf spot <i>Ascochyta medicaginicola</i>			
	Stagonospora leaf spot <i>Stagonospora meliloti</i>			
	Stemphylium leaf spot <i>Stemphylium</i> spp.			
	Summer black stem and leaf spot <i>Cercospora medicaginis</i>			
	Yellow leaf blotch <i>Leptotrichia medicaginis</i>			
	<p>• <b>Spray Interval</b> - Apply at 14-day to 21-day intervals.</p>			

### Use Restrictions

- **DO NOT** apply more than 10 fl ozs (0.13 lb mefentrifluconazole, 0.13 lb pyraclostrobin) per acre per application.
- **DO NOT** make more than 3 applications at 10 fl ozs per acre per year.
- **DO NOT** apply more than 30 fl ozs (0.39 lb mefentrifluconazole, 0.39 lb pyraclostrobin) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.39 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.73 lb ai/acre/year of pyraclostrobin-containing products.
- **DO NOT** apply within 14 days of grazing or harvest for forage or hay.
- **DO NOT** use on rangeland.

**Detailed Non-grass Forages (crop group 18) Crop List** - alfalfa; bean, velvet; clover (*Trifolium* spp., *Melilotus* spp.); kudzu; lespedeza; lupin; sainfoin; trefoil; vetch; vetch, crown; vetch, milk.

## Oilseeds

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Alternaria leaf spot <i>Alternaria</i> spp.			
	Blackleg <i>Leptosphaeria maculans</i>			
	Blackspot <i>Alternaria</i> spp.			
	Cercospora leaf spot <i>Cercospora</i> spp.			
	Pasmo <i>Septoria linicola</i>	7 to 10	20	21
	Powdery mildew <i>Erysiphe</i> spp.			
	Rust <i>Puccinia</i> spp., <i>Uromyces</i> spp.			
	Septoria leaf spot <i>Septoria</i> spp.			
	White rust <i>Albugo tragopogonis</i>			
<p>• <b>Spray Interval</b></p> <ul style="list-style-type: none"> <li>- Apply at 7-day to 14-day intervals.</li> <li>- Blackleg: Apply at 2-leaf to 4-leaf stage.</li> <li>- Blackspot: Apply at early pod development.</li> </ul>				

### Use Restrictions

- **DO NOT** apply more than 10 fl ozs (0.13 lb mefentrifluconazole, 0.13 lb pyraclostrobin) per acre per application.
- **DO NOT** make more than 2 applications at 10 fl oz per acre per year.
- **DO NOT** apply more than 20 fl ozs (0.26 lb mefentrifluconazole, 0.26 lb pyraclostrobin) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.73 lb ai/acre/year of pyraclostrobin-containing products.

**Detailed Oilseeds (crop group 20B) Crop List** - calendula; castor oil plant; Chinese tallowtree; euphorbia; evening primrose; jojoba; niger seed; rose hip; safflower; stokes aster; sunflower; tallowwood; tea oil plant; vernonia; cultivars, varieties, and/or hybrids of these.

## Sugar Beet

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Foliar	Cercospora leaf spot <i>Cercospora beticola</i>	7 to 10	20	7
	Powdery mildew <i>Erysiphe betae</i>			
<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply at 14-day intervals.</li> <li>• Aids in the control of Rhizoctonia stem canker and crown rot.</li> </ul>				

### Use Restrictions

- **DO NOT** apply more than 10 fl ozs (0.13 lb mefentrifluconazole, 0.13 lb pyraclostrobin) per acre per application.
- **DO NOT** make more than 2 applications per acre per year.
- **DO NOT** apply more than 20 fl ozs (0.26 lb mefentrifluconazole, 0.26 lb pyraclostrobin) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.78 lb ai/acre/year of pyraclostrobin-containing products.



# Sugarcane

Disease Controlled		Rate per Acre (fl ozs product) per application	Rate per Acre (fl ozs product) maximum per year	PHI (days)
Infurrow	Rhizoctonia seed and seedling rot <i>Rhizoctonia solani</i>	7 to 10	20	14
	Sugarcane pineapple disease <i>Ceratocystis paradoxa</i> * Fusarium seed rot, Seedling blight <i>Fusarium</i> spp.			
Foliar	Brown rust <i>Puccinia melanocephala</i>			
	Orange rust <i>Puccinia kuehnii</i>			
<ul style="list-style-type: none"> <li>• <b>Spray Interval</b> - Apply at 14-day to 28-day intervals.</li> </ul>				

## \* Suppression only

### Use Restrictions

- **DO NOT** apply more than 10 fl ozs (0.13 lb mefentrifluconazole, 0.13 lb pyraclostrobin) per acre per application.
- **DO NOT** make more than 2 applications at 10 fl ozs per acre per year.
- **DO NOT** apply more than 20 fl ozs (0.26 lb mefentrifluconazole, 0.26 lb pyraclostrobin) per acre per year.
- **DO NOT** apply more than a cumulative total of 0.26 lb ai/acre/year of mefentrifluconazole-containing products.
- **DO NOT** apply more than a cumulative total of 0.78 lb ai/acre/year of pyraclostrobin-containing products.
- **DO NOT apply this product to sugarcane through any type of irrigation system.**
- **DO NOT feed treated sugarcane commodities to livestock.**
- **Aerial Application - DO NOT** use less than 5 gallons of spray solution per acre. **Veltyma™ fungicide** can be applied by ground or air.

## SUGARCANE ONLY Infurrow and Soil-directed Banded Rates

Rate per 1000 Row Feet (fl ozs product)	Product Rate (fl ozs/A) Row width (feet)	
	5 feet	6 feet
0.80	7.0	
0.92	8.0	
0.96		7.0
1.03	9.0	
1.10		8.0
1.15	10.0	
1.24		9.0
1.38		10.0

## Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

1108

*Veltyma is a trademark of BASF.*

© 2020 BASF Corporation  
All rights reserved.

007969-00409.20201120.NVA 2020-04-554-0234  
Based on: NVA 2020-04-554-0147

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709



We create chemistry