

Evidence and Rational Based Research on Chinese Drugs

Bearbeitet von
Hildebert Wagner, Gudrun Ulrich-Merzenich

1. Auflage 2013. Buch. XII, 525 S. Hardcover

ISBN 978 3 7091 0441 5

Format (B x L): 15,5 x 23,5 cm

Gewicht: 967 g

[Weitere Fachgebiete > Medizin > Sonstige Medizinische Fachgebiete > Pharmakologie, Toxikologie](#)

Zu [Inhaltsverzeichnis](#)

schnell und portofrei erhältlich bei

The logo for beck-shop.de features the text 'beck-shop.de' in a bold, red, sans-serif font. Above the 'i' in 'shop' are three red dots of varying sizes, arranged in a slight arc. Below the main text, the words 'DIE FACHBUCHHANDLUNG' are written in a smaller, red, all-caps, sans-serif font.

beck-shop.de
DIE FACHBUCHHANDLUNG

Die Online-Fachbuchhandlung beck-shop.de ist spezialisiert auf Fachbücher, insbesondere Recht, Steuern und Wirtschaft. Im Sortiment finden Sie alle Medien (Bücher, Zeitschriften, CDs, eBooks, etc.) aller Verlage. Ergänzt wird das Programm durch Services wie Neuerscheinungsdienst oder Zusammenstellungen von Büchern zu Sonderpreisen. Der Shop führt mehr als 8 Millionen Produkte.

Chapter 2

DNA-Based Authentication of TCM-Plants: Current Progress and Future Perspectives

G. Heubl

2.1 Introduction

Traditional Chinese Medicine (TCM), with its long history, is deeply rooted in the Chinese culture and represents one of the oldest forms of medical therapy in the world. TCM has been used for thousands of years in China for health maintenance, disease prevention, and used to a lesser extent for the application of a variety of clinical therapies. Today, the global use of TCM is rapidly increasing and over 130 countries in the world are using Chinese Herbal Medicine (Hsiao 2007). The majority of drugs harvested and processed in China are of plant origin. A highly developed industry delivers medical plant preparations to the local Chinese customers and exports plant-based drugs to Asia and developing markets in Europe, Canada, and the USA.

Increasing interest by multinational pharmaceutical companies in herbal-based medicine is contributing to the significant economic growth of the global TCM market. It is estimated that 70–80 % of people worldwide rely chiefly on traditional herbal remedies as their primary form of health care (Farnsworth and Soejarto 1991; Pei 2001). Herbal medicine is becoming more main-stream and fashionable in richer countries. Therefore the TCM market has been growing at 10–20 % annually in Europe and North America over recent years (Hamilton 2003; Ten Kate and Laird 1999). In addition, there are many related botanical products sold as health foods, food supplements, herbal teas, and for various other purposes related to health and personal care. In terms of the number of species, the use of plants as medicine represents by far the most important application area of natural resources. Plants provide the predominant ingredients of medicine in most medical traditional

G. Heubl (✉)

Department Biologie I – Systematische Botanik, LMU München, Menzingerstr 67,
80638 München, Germany
e-mail: heubl@lrz.uni-muenchen.de

systems of healing and have been the source of inspiration for several new drug search endeavors of major pharmaceutical companies (Schippmann et al. 2002).

2.2 Biodiversity and Medicinal Plant Resources in China

China is one of the world's megabiodiverse countries with 32,308 species of seed plants belonging to 363 families and 3,427 genera (Ding 2002). The country spans a huge geographical area, is characterized by enormous variations in geographical, climatological, and topographical features and covers five climatic zones (cold-temperate, temperate, warm-temperate, subtropical, and tropical). The geology and geography is very complex, comprising the highest mountain ranges on Earth (the Himalayas), vaste plateaus such as Qinghai-Xizang (Tibetan) or Pamir, and arid basins such as Tarim, which contains the largest desert in China (Taklamakan). Main rivers of Asia such as Mekong, Brahmaputra, Yangtze, and Yellow River originate in Qinghai-Xizang Plateau (Lopez-Pujol and Zhao 2004). All features contribute to the enormous diversity of biomes (from rainforests to deserts) found in China, as well as to the enormous species diversity including many medicinal plants (Lopez-Pujol et al. 2006).

There is no reliable data for the total number of medicinal plants on earth and available information for countries and regions vary greatly. Estimates for the number of plant species used medicinally include 35,000–70,000 worldwide (Farnsworth and Soejarto 1991; Schippmann et al. 2002), 7,295–11,146 in China (Pei-Gen 2007; Hamilton 2004; Huang et al. 2002a, b). According to statistics there are 12,807 kinds of remedies in Chinese medicine. Of these, 11,146 (80 %) are plant based, 1,581 are animal based, and 80 are mineral based (Gao et al. 2002). Regarding the diversity of Chinese medicinal plants, only 500–600 species are commonly used in Traditional Chinese Medicine and prescribed by Chinese medical practitioners, 1,430 in Mongolian Medicine, and 1,106–3,600 in Tibetan Medicine (Pei 2001, 2002b). In the 2005 edition of the Chinese Pharmacopoeia 1,146 monographs and 538 herbal drugs have been included. The improved version of the People's Republic of China Pharmacopoeia 2010 contains 1,174 monographs (in total 4,567 monographs) and enumerates more than 4,600 varieties of species. In Europe and USA 75–150 herbal drugs are mostly used by TCM practitioners (Table 2.1).210648_Wagner

The Chinese medicinal industry represents a significant portion of the pharmaceutical industry in China. There are 1,200 Chinese medicinal, industrial enterprises that manufacture approximately 8,000 Chinese herbal medicine products. The total annual sales of functional food, TCM preparations, medicinal plant extracts, and other processed materials exceed US \$ 40 billion (Liu et al. 2011). Today, it is a big challenge to protect the medicinal plant resources from overexploitation and habitat destruction (Srivastava et al. 1996). For example, 70–80 % of natural materials were collected from the wilderness to meet the annual demand of approximate one million tons (Balunas and Kinghorn 2005). The annual sales of these natural resources have

Table 2.1 Diversity and inventory of the Chinese flora with data on medicinal plants

Resource	Total number of species	Medicinal plant species number
• Algae	5,000 ^a /8,997 ^d /12,500 ^e	115 ^a
• Mosses	2,200 ^{a,c,d,e,f}	43 ^a
• Ferns	2,300 ^e /2,600 ^{a,b,d,f}	456 ^a
• Seed plants	25,000 ^a /30,000 ^{b,c,d}	10,188 ^a
Gymnosperms	192 ^c /200 ^a /250 ^b /270 ^d	
Angiosperms	25,000 ^f /30,000 ^{c,d} /31,500 ^g	
Total plants	34,692–47,570	10,802
• Lichens	2,000 ^c	52 ^a
• Fungi	8,000 ^{f,d}	292 ^a

^aGao et al. (2002)^bLopez-Pujol and Zhao (2004)^cXue (1997)^dSEPA (1998)^eLi (2003)^fNEPA (1994)^gFlora of China Project Missouri Botanical Garden (http://www.mobot.org/press/Assets/FP/flora_china.asp). Estimated numbers of other medicinal resources in China: Medicinal animals 1,581; medicinal mine 80 (Huang et al. 2002a, b)

increased to more than 100 times in relation to the levels of 1980. Many of these natural resources are derived from species that are threatened or have become rare or endangered by large-scale exploitation (Nalawade et al. 2003; Cole et al. 2007). With the rapid increase of consumer demand for crude drugs and natural health products, many medicinal plant species are threatened with extinction due to overexploitation and habitat destruction. To ensure the sustainable use of medicinal plant resources, a conservation framework consisting of conservation strategies, cultivation practices, and various technologies has been developed. Conservation strategies include establishing in situ and ex situ conservation centers, setting up government policies and regulations, establishing methods for resource surveying and trade monitoring as well as establishing and enforcing Good Agricultural Practices (Liu et al. 2011).

About 140 new drugs have originated directly or indirectly from Chinese medicinal plants by means of modern scientific methods, confirming that these plants are an important resource (Liu and Yaniv 2005; Lee et al. 2005). The first compound derived from Chinese herbal remedies to enter the Western pharmacopoeia was Ephedrine, an amphetamine-like stimulant isolated from the Chinese medicinal herb “Ma-Huang” (*Ephedra sinica*). The next significant pharmaceutical breakthrough derived from Chinese medicine was the isolation of Artemisinin from “Qing-Hao” (*Artemisia annua*). Researchers found that Artemisinin was beneficial for fever and killed even chloroquine-resistant strains of *Plasmodium*, the parasite that causes malaria. Recent work suggests that Artemisinin may also have anticancer properties, a hypothesis which was also established for *Coix lacryma-jobi* var. *ma-yuen* the Job’s Tear. Modern pharmacological studies have demonstrated that the Coix “seed” (a bony utricle of the single female flower enclosing the fruit) and its preparation-KLT (Kanglaite Injection) possess extensive pharmacological

activities (Li 2001; Lu et al. 2008). Active components have been isolated and defined in many other Chinese herbs, for example, anthraquinone glycosides in rhubarb (*Rheum officinale*), gingerols in ginger (*Zingiber officinale*), ginkgolides in Ginkgo (*Ginkgo biloba*) or berberine, an antibacterial component from Chinese goldthread (*Coptis chinensis*) (Wang et al. 2007a).

2.3 Adulterants, Substitutes, and Confused Species

Although many Chinese medicinal plants have been used successfully in China for a long period of time they have never been subject to the stringent quality standards and regulations for pharmaceutical products in Europe or USA. To make traditional and innovative plant products acceptable for modern medicine, it is necessary to have reliable botanical, phytochemical and molecular identification tools for the identification of medicinal plant species (Liang et al. 2004).

A severe problem of the global TCM market is that many erroneous substitutes and adulterants of Chinese medicinal plants are traded due to their lower costs or due to the misidentification of species with similar morphological features.

In recent times the use and demand of herbal preparations has been growing in Western countries where medicinal plants have gained popularity and attention among physicians and patients. Among the alternative medicines TCM is becoming one of the most widely used therapies throughout the world (Normile 2003). One of the reasons for the increasing interest in herbal medicines is the belief that, being natural and traditionally used, they are hence safe and do not possess the potential for negative side effects. This coupled with lower costs compared to conventional medications is the major attraction of these treatments. Nevertheless, their natural origin is not a guarantee of safety. In literature, many reports point out the risks associated with the use of herbal products (Chan 1997; Ernst 2004; Ng et al. 2009). The quality of a herbal drug can be negatively affected by the use of inherent toxic herbs, by fraudulent action due to substitution or adulteration, by contamination, by misidentification, by confusion of species, and by inappropriate labeling. There are several incidents of Chinese herbs which document that adulterants or substitutes caused serious intoxications and even deaths (But 1994; Graham-Brown 1992; Ng et al. 2009; Chan 2003; Mazzanti et al. 2008; Zhao et al. 2007a; Chan and Critchley 1996; Gertner et al. 1995; But et al. 1996; Chen et al. 2002; McIntyre 1998; Yang and Chen 1998).

A case of encephalopathy and neuropathy was reported following ingestion of a decoction supposedly prepared from “Long Dan Cao” (*Gentiana rigescens* radix). Investigation showed that the toxicity was in fact due to adulteration of the herb with the roots of *Podophyllum emodi* or *P. hexandrum* which contain the neurotoxin podophyllotoxin in high concentration (But 1994; Ng et al. 2009). Several cases of renal damage attributed to “Fang-Ji” (*Stephania tetrandra*) in a weight-loss preparation were actually caused by “Guang Fang Ji” (*Aristolochia fangchi*) that contains the highly toxic aristolochic acid, a known nephrotoxin which causes renal failure

and urothelial carcinoma. The confusion in the latter case has obviously arisen from the similar Chinese names (Fugh-Berman 2000). Several other poisoning cases are documented and resulted from erroneous substitution of herbs.

One further example of species confusion, which had serious consequences, involves the common name “Bai Mao Teng.” This name has been used for *Solanum lyratum* and *Aristolochia mollissima*. Apparently, these herbs belong to two different families but look similar. While *S. lyratum* is not harmful, *A. mollissima* contains the toxic aristolochic acid that can cause kidney failure and cancer of the urinary tract (Zhao et al. 2006).

Another incident refers to commercial “Mu Tong,” a Chinese diuretic drug, which is associated with five species: *Clematis armandii* and *Clematis montana* (“Chuan Mu Tong”), *Akebia quinata* (“Wu Ye Mutang”), *A. trifoliata* (“San Ye Mu Tong”), *Aristolochia manshuriensis* (“Guan Mu Tong”), and *A. moupinensis*/*A. kaempferi* (“Huai Mu Tong”). This has led to a serious confusion with the consequence that substitution with *Aristolochia manshuriensis* caused renal failure (Zhu 2002; Cheung et al. 2006; Debelle et al. 2008).

Very often the high price of some herbs is an incentive for criminals to make quick profit by manufacturing counterfeit products. The more precious and rare the crude drug is, the more likely the counterfeit products will find their way into the market. A classic example is *Cordyceps sinensis* (recently renamed as *Ophiocordyceps sinensis*), a fungus parasitizing the larva of some species of insects and the dead caterpillar (Dong and Yao 2010). This scarce drug is found only in fairly inaccessible alpine regions on the Tibetan Plateau and is one of the best-known traditional Chinese medicinal products, with great benefits to human health and with a huge economic value. There are many substitutes of so-called *Cordyceps* that are traded worldwide, such as *Cordyceps militaris* (the most commonly used substitute), *C. martialis*, *C. hawkesii*, *C. liangshanensis*, *C. barnesii*, *C. cicadicola*, *C. gracilis*, *C. ramosa*, *C. ophioglossoides*, and *C. gunnii*. In addition, there are counterfeit plant products of the fungus and mimics such as *Stachys geobombycis*, *Stachys sieboldii*, and *Lycopus lucidus* that have been found on the market. Consequently there is a serious problem for authentication and quality control of *Cordyceps* (Li et al. 2006).

A review published by Zhao et al. (2006) enumerates further examples and summarizes the facts and reasons for confusion in the current Hong Kong medicinal herb market.

2.4 Authentication of Herbal Material on the DNA Level

The authentication of Chinese medicinal plants, depending on the correct identification of species, is an essential prerequisite to ensure safety, herbal drug quality, and therapeutic efficacy (Zhao et al. 2006). Identification of herbal materials, which commonly consist of dried or processed parts, is generally difficult because many useful diagnostic characters are lost during the drying process. This is particularly

true when one herb has more than one common name, or where one common name is used for more than one herb (Zhao and Li 2004). In practice, the identification of medicinal plants relies mainly on morphological, anatomical, and phytochemical characters. Many pharmacopoeias refer to macroscopic and microscopic evaluation (morphology, histology) and chemical profiling (TLC-, HPLC-, GC-fingerprinting) for quality control and standardization of raw and processed herbs (Chan 2003; Siow et al. 2005; Wagner et al. 2011). However, chemical variability within the plant material often hinders the confirmation of its botanical identity as the chemical composition is affected by growth and storage conditions as well as by the harvesting process. Otherwise microscopic examination of drugs requires botanical expertise for the unequivocal authentication, as related species often possess similar features.

With the improvements in molecular biotechnology and plant genetics in the past decades, genetic tools are considered to provide more reliability for authentication of herbal materials at the DNA level (Kumar et al. 2009). Thus in the meanwhile, various DNA-based molecular marker techniques are applied in many fields and their application is remarkably increasing for species characterization in medicinal plants (Shaw et al. 2002, 2009; Joshi et al. 2004; Zhang et al. 2007; Sucher and Carles 2008). This is especially useful in case of those species that are frequently substituted or adulterated with other species or in case of varieties that are morphologically and/or phytochemically almost indistinguishable.

Benefiting in the first place from PCR techniques, DNA markers have become a powerful tool for identification and authentication of plant, animal, fungal, and bacterial species (Yip et al. 2007; Kaplan et al. 2004; Pereira et al. 2008; Hao et al. 2010). Contrary to chemical fingerprinting which is strongly influenced by the age of the sample, physiological conditions, environmental factors, cultivation area, harvesting period, drying, and storage conditions, DNA is an extremely stable macromolecule that is not affected by external factors and therefore can be recovered from fresh, dried, and even processed biological material. Additionally the marker molecules are not tissue specific and thus can be detected at all stages of organism development. Moreover, only a small amount of a sample is sufficient for analysis.

This review provides an overview of DNA-based technologies and most commonly used assays with an emphasis on those that are based on DNA hybridization, restriction enzymes, random PCR amplifications, species-specific PCR primers, and DNA sequencing. A critical evaluation of all methods is presented focusing on their discriminatory power, sensitivity, reproducibility, user-friendliness, and costs (see Table 2.3).

The previously described incidents highlight the importance of a correct botanical classification and the need of an adequate knowledge of morphological characteristics of herbal drugs in order to perform a proper identification. As more and more people worldwide use Chinese herbs, authentication becomes an increasing problem because adulterated and substituted Chinese medicinal materials are widely common in the market (Mills and Bone 2005).

Therefore authentication of Chinese medicinal materials is the key for safety, appropriate use and maximum therapeutic potency, minimization of trading fraud,

and last but not least for the increase of consumer confidence in Chinese medicine. Identification at species level is required for quality assurance, which includes both identifying the crude plant product and evaluating its pharmaceutical quality (Wagner et al. 2011). Thus, authentication is a fundamental step for the successful, reliable clinical application and for accurate experimental studies of TCM plants. This will ensure the safe and effective use of Chinese medicinal herbs throughout the world.

2.4.1 Types of DNA Markers Used in Plant Genome Analysis

There are various types of DNA-based molecular techniques that are used to evaluate DNA polymorphism in order to authenticate plant taxa (Sucher and Carles 2008; Shaw et al. 2009; Yip et al. 2007; Kaplan et al. 2004; Pereira et al. 2008; Heubl 2010). These are hybridization-based methods, polymerase chain reaction (PCR)-based methods, and sequencing-based methods. In recent times the use of multilocus sequence analysis (MLSA), which is commonly used in phylogenetic studies, has proven its discriminatory power. Additionally DNA microarrays that contain thousands of probes are a promising new development for sensitive and high-throughput taxon identification (Trau et al. 2002; Schena et al. 1998).

2.4.1.1 RAPD (Randomly Amplified Polymorphic DNA)

The RAPD technology utilizes short synthetic oligonucleotides (10 bp long) of random sequences as primers to generate a high number of anonymous DNA fragments via PCR reaction. The large number of amplification products is generally separated on agarose gels and stained with ethidium bromide or SyBRgreen.

Using an appropriate annealing temperature in the PCR cycle, oligonucleotide primers bind to several priming sites on the complementary sequences in the template genomic DNA. If these priming sites are within an amplifiable distance discrete DNA fragments are generated. Nucleotide variation between different sets of template DNAs will result in the presence or absence of bands because of changes in the priming sites. Polymorphism of amplified fragments is caused by (1) base substitutions or deletions in the priming sites, (2) insertions that render priming sites too distant to support amplification, or (3) insertions or deletions that change the size of the amplified fragment (Weising et al. 2005).

Because of the simplicity (no prior sequence information is necessary), low costs, efficiency in developing a large number of DNA markers in a short time, and requirement for less sophisticated equipment, RAPDs have found a wide range of applications. Although the RAPD method is easy to perform, the issue of reproducibility has been an important concern. In fact, the RAPD reaction is far more sensitive than conventional PCR because of the length of a single and arbitrary primer, which is used to amplify anonymous regions of a given genome. Special

care is needed for keeping out contaminant DNA (from infections and parasites) in the material to avoid misleading patterns. RAPDs are inherited as dominant-recessive characters which mean that homozygotes and heterozygotes cannot be distinguished.

RAPD markers have found a wide range of applications in the authentication of medicinal plants. The technique has been applied in many plant groups like *Glycyrrhiza* (Yamasaki et al. 1994), *Atractylodes* (Kohjyouma et al. 1997; Chen et al. 2001), *Astragalus* (Cheng et al. 2000), *Amomum* (Wang et al. 2000), *Scutellaria* (Hosokawa et al. 2000), *Panax* (Cheung et al. 1994; Shim et al. 2003; Cui et al. 2003; Lim et al. 2007), *Aconitum* (Cole and Kuchenreuther 2001), *Ginkgo* (Fan et al. 2004), *Anectochilus* (Cheng et al. 1998), *Lycium* (Zhang et al. 2001a), *Angelica* (Watanabe et al. 1998), *Bupleurum* (Liang et al. 2002), *Dendrobium* (Zhang et al. 2001b), *Magnolia* (Guo et al. 2001), *Asarum* (Huang et al. 1998), *Apocynum* (Lu et al. 2010a), *Trollius* (Li et al. 2010a), *Phyllanthus* (Dnyaneshwar et al. 2006), *Indigofera* (Zhang et al. 1997), *Coptis* (Cheng et al. 1997), *Codonopsis* (Zhang et al. 1999), *Taraxacum* (Cao et al. 1996a), *Elephantopus* (Cao et al. 1996b), and *Rehmannia* (Cheng et al. 2002).

2.4.1.2 AP-PCR (Arbitrary Polymerase Chain Reaction)

AP-PCR (or Arbitrarily Chosen Primers ACP-PCR) is a special variation of RAPD which is using single primers approximately 10–50 bp in length (Welsh and McClelland 1990). In AP-PCR the amplification follows three steps. In the first two cycles annealing is under nonstringent conditions. Higher primer concentrations are used in the first cycle. Often primers of variable length are used and products are mostly analyzed on polyacrylamide gels. AP-PCR has been applied to various groups for identification of species and analysis of genetic variation (Munthali et al. 1992; Kersten et al. 2007). Similar to RAPDs, reproducibility can be a problem for fingerprints generated by a single primer, because small changes in annealing conditions can affect banding pattern.

2.4.1.3 DAF (DNA Amplification Fingerprinting)

DNA Amplification Fingerprinting (DAF) is a variant of the RAPD technique and was developed by Caetano-Anollés et al. (1991a). For PCR amplification, very short oligonucleotide primers of arbitrary sequence (mostly five nucleotides) are used to amplify short fragments of genomic DNA resulting in a very complex banding pattern (Caetano-Anollés et al. 1991a, b). DAF uses low stringency amplification conditions so that primers can anneal arbitrarily at multiple sites on each template DNA strand and initiate DNA synthesis. The method needs careful optimization of parameters and only two temperature cycles are required. DAF products are routinely separated by polyacrylamide gels and detected by silver staining (Chawla 2002). Another approach called ASAP (arbitrary signatures from

amplification profiles) was developed to enhance the level of informativeness of DAF reactions by using primers which contain both a 5' mini-hairpin sequence and a short "core" arbitrary 3' sequence. These arbitrary mini-hairpin primers increase detection of polymorphic DNA and direct the controlled amplification of small template molecules, thereby generating "sequence signatures" from PCR-amplified fragments (Caetano-Anollés and Gresshoff 1996).

2.4.1.4 ISSR (Inter Simple Sequence Repeat)

In higher plants, Inter Simple Sequence Repeat or ISSR markers are frequently applied because they are known to be abundant, very reproducible, highly polymorphic and easy to use (Zietkiewicz et al. 1994; Borner et al. 2002). ISSR, also known as anchored simple sequence repeat (ASSR), has been used in genetic fingerprinting, gene tagging, phylogenetic analysis, species and cultivar identification, and assessment of hybridization (Kurane et al. 2009).

The ISSR technique is nearly identical to RAPD except that ISSR primers are designed from microsatellite regions and are longer (approximately 14 bp or more) than RAPD primers. Microsatellites are very short stretches of DNA that are "hypervariable," expressed as different variants within populations and among different species. ISSR uses the presence of Simple Sequence Repeats (SSRs) which are characterized by mono-, di-, or trinucleotide repeats (e.g., AA... or AG... CAG...) that have 4–10 repeating units side-by-side. These SSRs are ubiquitous, abundant, and highly polymorphic. The primers used can be 5' or 3' anchored by 1–3 selective nucleotides to prevent internal priming and to amplify only a subset of the targeted inter-repeat regions. ISSR markers access variations in the numerous microsatellite regions dispersed throughout the genome. The PCR products are mostly separated on agarose gels and stained with ethidium bromide. Alternatively amplified DNA fragments can also be screened and detected using capillary electrophoresis which significantly increases the amount of information compared to the traditional agarose gel electrophoresis. Since ISSRs are dominant markers, the amplified fragments are scored as diallelic. Presence (0/1) of loci can be used for genetic similarity or cluster analysis. Changes in the amplified products can arise through structural changes in the region (insertions or deletions) or the loss of primer binding sites.

ISSR markers overcome the weakness of low reproducibility of RAPDs, the high costs of AFLPs, the complexity of SSRs and thus this technique is less time consuming, more cost-efficient, use of radioactivity is not required, no prior sequence information is necessary, and it shows high polymorphism.

ISSRs have been used for screening genetic diversity and authentication of *Dendrobium* (Shen et al. 2006), *Cistanche* (Shi et al. 2009), *Fritillaria* (Li et al. 2009a), *Salvia* (Song et al. 2010), *Rehmannia* (Wang et al. 2005), *Vitex* (Hu et al. 2007), *Cannabis* (Kojoma et al. 2002), *Rhodiola* (Xia et al. 2007), *Cymbidium* (Wang et al. 2009), *Ammopiptanthus* (Ge et al. 2005), *Swertia* (Joshi and Dhawan 2007), *Glycyrrhiza* (Yao et al. 2008), and *Houttuynia* (Wu et al. 2005).

2.4.1.5 AFLP (Amplified Fragment Length Polymorphism)

Amplified fragment length polymorphism (AFLP) originally developed by Zabeau and Vos (1993) is a powerful tool for DNA fingerprinting of organismal genomes and it combines the use of RFLP and PCR techniques. This multilocus approach needs no prior sequence information, it is highly reproducible, with the ability to screen a large number of loci (ca. 50–100 fragments per reaction) for polymorphisms. It is a very useful technique for DNA fingerprinting, especially when very little information on the genome of the plant under study is available (Mueller and Wolfenbarger 1999; Blears et al. 1998).

The procedure of this technique (Vos et al. 1995) is a multistep process (1) Digestion of total genomic DNA with two restriction enzymes. One restriction enzyme is a frequent cutter (four-base recognition site, e.g., MseI), the second restriction enzyme is a rare cutter (six-base recognition site, e.g., EcoRI). (2) Adapters specific to the restriction sites are ligated to the fragment ends which serve as binding sites for selective primers in PCR amplification. (3) A first PCR (preselective amplification) is performed. The PCR primers consist of a core sequence (part of the adapter), a restriction enzyme specific sequence, and 1–5 selective nucleotides (the higher the number of selective nucleotides, the lower the number of bands obtained per profile). (4) Preselective amplification products undergo another PCR run, and again a subset of those fragments is selected. Usually, for the second selective amplification, two extra nucleotides are added to the primers. (5) Amplified fragments, labeled with fluorescent or radioactive tags, are separated on acrylamide gels or with automated genetic analyzers and dominant markers are scored as the presence/absence of loci (Weising et al. 2005).

The advantages of AFLPs lie in their high genomic abundance, considerable reproducibility, in their generation of many informative bands per reaction, in their wide range of applications, and in the fact that no sequence data are required for primer construction. Compared with the widely used RFLP, AFLP is faster, less labor intensive, and provides more information.

Disadvantages include the need for purified, high molecular weight DNA, the dominance of alleles, and the possible nonhomology of comigrating fragments belonging to different loci. Because of the highly informative fingerprinting profiles, which are usually obtained as results, AFLPs can be applied in studies involving genetic identity, parentage, identification of clones and cultivars, and in phylogenetic studies of closely related species.

AFLP analyses have been used in *Panax* (Choi et al. 2008; Ha et al. 2002), *Actaea* (Zerega et al. 2002), *Plectranthus* (Passinho-Soares et al. 2006), *Caladium* (Loh et al. 1999), *Cannabis* (Datwyler and Weiblen 2006), and *Rehmannia* (Qi et al. 2008).

2.4.1.6 RAMPO (Randomly Amplified Microsatellite Polymorphism)

This method, also termed RAHM (random amplified hybridization microsatellite) or RAMS (randomly amplified microsatellites), combines arbitrarily primed PCR (RAPD) with microsatellite hybridization to produce polymorphic genetic fingerprints (Weising et al. 2005; Weising and Kahl 1998). No prior sequence information is needed. Genomic DNA is first amplified with a single arbitrary 10-mer primer (as in RAPDs) or microsatellite-complementary 15- to 16-mer PCR primer. After electrophoretic separation and staining of the PCR products, the gel is either dried or blotted onto a nylon membrane and subsequently hybridized to a radiolabeled mono-, di-, tri-, or tetranucleotide repeat probe such as [GT]₈, [GA]₈, or [CAA]₅. Subsequent autoradiography reveals reproducible, probe-dependent fingerprints that are completely different from the ethidium bromide staining patterns and that are polymorphic at an intraspecific level. An advantage of the RAMPO technique is the low complexity of banding patterns, which is considerably facilitating the detection of species-specific bands. RAMPO bands appear to be less sensitive to misinterpretation than RAPD bands, because not only the size but also the hybridization signal intensity of two bands (i.e., the presence and copy number of a certain microsatellite) are criteria for homology. The method is mainly used for identification and discrimination of genotypes within and among populations, cultivates, and germplasm e.g., in *Ficus* (Chatti et al. 2007) and *Phoenix dactylifera* (Soumaya et al. 2008). RAMPOs share their fate with other marker technologies that are partly or totally based on blot hybridization. These methods are barely used anymore, because more convenient marker systems are available for most purposes.

2.4.1.7 RFLP (Restriction Fragment Length Polymorphism)

RFLP analysis was one of the first techniques to be widely used for detecting variations at the DNA level. The principle of this method is based on the comparison of banding patterns from DNA sequences digested with specific restriction enzymes (e.g., HaeIII, EcoRI, BamHI). Restriction enzymes are endonucleases produced by bacteria with the function to cut specific DNA sequence motifs of invading DNA molecules. Each enzyme has a specific, typically palindromic recognition sequence. Consequently it recognizes and cuts DNA in a predictable way, resulting in a reproducible set of DNA fragments of different lengths. If two organisms (strains, individuals, or species) differ in the distance between sites of cleavage of a particular restriction endonuclease, then also the length of the fragments produced differs. Consequently RFLP is a result of (a) point mutation creating or destroying a restriction site and (b) insertion/deletions altering the size of a given restriction fragment. These differences in fragment lengths can be detected by gel electrophoresis, hybridization, and visualization. To detect specific fragments, the DNA restriction fragment profile of the agarose gel is transferred to

a nitrocellulose or nylon membrane. Afterwards a single-stranded DNA probe is conveniently labeled, using any standard method (e.g., a radioisotope or digoxigenin) and hybridized with the target DNA, which is stuck to the membrane. Polymorphisms are detected by the presence or absence of bands. RFLP is a robust methodology; the markers are relatively polymorphic, codominantly inherited, and highly reproducible. The method also provides opportunity to simultaneously screen numerous samples. The technique is time consuming, costly, labor intensive and requires a large quantity of good quality or undegraded DNA (Weising et al. 2005).

RFLP combined with DNA hybridization has mainly been used for phylogenetic studies in the past e.g., in *Lupinus* (Yamazaki et al. 1993), *Hedysarum* (Trifi-Farah and Marrakchi 2001), *Triticum* (Mori et al. 1997), *Musa* (Gawel et al. 1992), and for detection of *Dendrobium* (Li et al. 2005) and *Fritillaria* (Tsoi et al. 2003).

2.4.1.8 Microsatellites or SSR (Simple Sequence Repeats)

Microsatellites also known as simple sequence repeats (SSRs), short tandem repeats (STRs), or simple sequence length polymorphisms (SSLPs) are the smallest class of simple repetitive DNA sequences (Litt and Luty 1989; Gupta et al. 1996). Based on tandem repeats of short (2–6 bp) DNA sequences, these markers are highly polymorphic due to variation in the number of repeat units and dispersed throughout most eukaryotic genomes. The large number of alleles and the high levels of variability among closely related organisms made PCR-amplified microsatellites the marker system of choice for a wide variety of applications like population genetic studies, genome mapping, and marker-assisted breeding (Valdes et al. 1993; Akkaya et al. 1995; Schuler et al. 1996). It is meanwhile proven that the predominant mutation mechanism in microsatellite tracts is slipped-strand mispairing (Levinson and Gutman 1987). The repeat length at specific SSR loci is easily assayed by PCR using primers specific to conserved regions flanking the repeat. PCR fragments are usually separated on polyacrylamide gels or capillary sequencers in combination with fluorescent detection systems. The reason for the wide usage of nuclear microsatellites is their high abundance, enormous extent of allelic diversity, and suitability for automatization. Meanwhile many SSR primers deduced from flanking sequences of known microsatellites are deposited in DNA databases.

In addition to nuclear microsatellites, chloroplast microsatellites are also particularly effective markers for analysis of the genetic diversity (Clark et al. 2000) and phylogeography of plant populations (Chen et al. 2011), studying mating systems, gene flow via both pollen and seeds, detection of hybridization, and introgression (Agarwal 2008). One limitation of the approach is the need of sequence data for primer construction. Primer sequences flanking chloroplast microsatellites are usually inferred from fully or partially sequenced chloroplast genomes. In general, these primer pairs produce polymorphic PCR fragments from the species of origin and their close relatives, but transportability to more distant taxa is limited. Attempts to design universal primers to amplify chloroplast microsatellites have

resulted in a set of consensus chloroplast microsatellite primers (ccmp1–ccmp10) that aims at amplifying cpSSR regions in the chloroplast genome of dicotyledonous angiosperms (Weising and Gardner 1999). Most of the primer pairs derived from A or T mononucleotide repeats ($n = 10$) identified in the tobacco chloroplast genome, were functional as genetic markers in the Actinidiaceae, Brassicaceae, and Solanaceae (Chung and Staub 2003). Universal primers for the amplification of chloroplast microsatellites in grasses (Poaceae) have also been developed (Provan et al. 2004).

A major limitation of SSRs are the time and high development costs required to isolate and characterize each locus when preexisting DNA sequence is not available. Typically, this process requires the construction and screening of a genomic library of size-selected DNA fragments with SSR-specific probes, followed by DNA sequencing of isolated positive clones, PCR primer synthesis, and testing. With the availability of large numbers of ESTs and other DNA sequence data, development of EST-based SSR markers is less time consuming and expensive. The methodology and applications of nuclear microsatellite markers in plants and other organisms has been subjected to numerous reviews, including Goldstein and Schlötterer (1999), Zane et al. (2002), Li et al. (2002), Varshney et al. (2005), and Squirrell et al. (2003).

Microsatellites have been applied in *Panax* (Kim et al. 2007a; Jo et al. 2009), *Acanthopanax* (Kim and Chung 2007), *Dendrobium* (Fan et al. 2009), *Cymbopogon* (Kumar et al. 2007), *Bupleurum* (Chun et al. 2009), and *Schisandra* (Boqian et al. 2009).

2.4.1.9 SAMPL (Selective Amplification of Microsatellite Polymorphic Loci)

The SAMPL technique was introduced by Morgante and Vogel (1994). It combines the high and controllable multiplexing rate of the AFLP technique with the high levels of microsatellite polymorphism by using AFLP-type primers together with compound microsatellite primers (Weising et al. 2005).

SAMPLs differ from AFLPs by using primers with compound microsatellite motifs in combination with oligonucleotides complementary to the end-ligated adapters for the selective amplification step (Paglia et al. 1998). In brief, genomic DNA is digested with restriction enzymes (commonly *EcoRI* and *MseI*), and the resulting fragments are ligated to adapters. Afterwards a preamplification reaction for all ligated DNA fragments is carried out with primers annealing to the adapters. These preamplified products are then used as templates for a selective SAMPL-polymerase chain reaction (PCR) reaction that uses the adapter-primer (*EcoRI* oligo-1) in combination with an end-labeled microsatellite-based 15-mer oligonucleotide (Karp and Edwards 1997) to amplify a group of fragments from those fragments that were restricted, ligated, and preamplified. A disadvantage of multilocus SSR profiling is the capture of only some of the polymorphism associated with microsatellites due to the prevalence of dominant markers and difficulty in identifying allelic fragments in complex DNA fingerprints. This

multiplexing genome profiling technique has not been used adequately in plant genomics, although a few reports have already documented its potential for detecting polymorphisms (Molina and Kahl 2002). This method was used for analysis of genetic diversity in *Cicer* (Winter et al. 1999), *Lactuca* (Witsenboer et al. 1997), and *Tribulus* (Sarwat et al. 2008).

2.4.1.10 DAMD (Directed Amplification of Minisatellite-Region DNA)

DAMD is a DNA fingerprinting method based on amplification of the regions rich in minisatellites at relatively high stringencies by using previously found VNTR core sequences as primers (Heath et al. 1993; Somers and Demmon 2002). Minisatellites also known as variable number of tandem repeats (VNTR) or hyper-variable repeats (HVR) are similar to microsatellites (SSR) except that the tandem repeat DNA sequences are longer and generally consist of 10–60-bp motifs. Extreme variations in the tandem repeat copy number of minisatellite loci are responsible for the polymorphism observed. By using the VNTR core sequences as primers, the directed amplification of minisatellite-region DNA (DAMD) with PCR is capable of producing RAPD-like results for the identification of species (Silva et al. 2001). They are also used to generate highly variable probes for DNA fingerprinting. This method is more reproducible than RAPD due to the longer primers used.

Recently, DAMD-PCR has been applied successfully for genotyping of wheat cultivars and rice species (Zhou et al. 1997). The method has been used for authentication of *Panax* (Ha et al. 2002), *Capsicum* (Ince et al. 2009), *Salvia* (Karaca et al. 2008), and *Morus* (Bhattacharya et al. 2005).

2.4.1.11 SNP (Single Nucleotide Polymorphism)

Single nucleotide polymorphisms (SNPs) are widely observed between individuals, ecotypes, and species, serving as efficient molecular markers particularly in genetic analysis and breeding programs, also including ecological and evolutionary studies.

SNPs are single-base pair positions in the genomes of two (or more) individuals, at which different sequence alternatives (alleles) exist. Polymorphisms result from point mutations (either transition or transversion events) causing single base-pair differences between DNA sequences. According to most recent estimates, one SNP occurs every 100–300 bp (or every 1,000 bp) in any genome (Kwok 2001). SNPs are codominant, single-locus, biallelic markers and they are the most abundant molecular markers known so far. The major SNP genotyping techniques fall into at least six groups (1) direct sequencing, (2) restriction enzyme digestion (cleaved amplified polymorphic sequences = CAPS), (3) allele-specific PCR, (4) allele-specific primer extension, (5) allele-specific oligonucleotide hybridization, and (6) allele-specific oligonucleotide ligation (Weising et al. 2005).

Several molecular markers such as restriction fragment length polymorphism (RFLP), cleaved amplified polymorphic sequence (CAPS), amplification refractory mutation system (ARMS), and single-strand confirmation polymorphism (SSCP) are based on SNPs.

Many biotechnology companies are marketing DNA microarrays that can test a sample DNA for thousands of SNP sequences.

SNPs have been applied in authentication of *Perilla* varieties (Luo et al. 2006a), *Dendrobium officinale* (Ding et al. 2008), *Panax* cultivars (Wang et al. 2010a), *Boehmeria* varieties (Li et al. 2010b).

2.4.1.12 ARMS (Amplification Refractory Mutation System)

ARMS, also known as allele-specific polymerase chain reaction (AS-PCR) is a simple, timesaving, and effective method for detecting any mutations involving single base changes (SNPs) or small deletions. It has become a standard technique that allows the discrimination of alleles (Newton et al. 1989). The basis of ARMS is that oligonucleotides with a mismatched 3'-residue will not function as primers in the PCR. ARMS allows amplification of test DNA only when the target allele is contained within the sample and it does not amplify the nontarget allele. Following an ARMS reaction, the presence or absence of a PCR product is diagnostic for the presence or absence of the target allele. A main advantage of ARMS is that the amplification step and the authentication step are combined, in a way that the presence or absence of a PCR product is diagnostic for the presence or absence of the target allele. The method provides a quick screening assay that does not require any form of labeling as the amplified products are visualized simply by agarose gel electrophoresis and ethidium bromide staining. Multiplex ARMS or MARMS are a similar approach but there are several primer combinations to be optimized simultaneously, which increases the complexity of the procedure. The ARMS technique has been applied in authentication of *Alisma* (Li et al. 2007), *Panax* (Zhu et al. 2004; Diao et al. 2009), *Rheum* (Yang et al. 2004), *Dendrobium* (Ding et al. 2008; Qian et al. 2008), and *Curcuma* (Sasaki et al. 2002).

2.4.1.13 CAPS or PCR-RFLP (Cleaved Amplified Polymorphic Sequence)

CAPS, originally named PCR-RFLP, is a combination of PCR of target DNA and subsequent digestion with a restriction enzyme (Maeda et al. 1990; Lum et al. 2005). CAPS markers are generated in two steps. In the first step of a standard CAPS experiment, a defined sequence is amplified using specific 20–25 bp primers. In the second step, the PCR-product is digested with a restriction enzyme usually with a four-base recognition specificity. The digested fragments are separated on agarose gels and stained with ethidium bromide. To identify suitable combinations of amplicons and restriction enzymes, a wide range of PCR primer pairs and restriction enzymes need to be screened during the initial phase of a CAPS

project, using a small set of templates. Combinations that reveal informative polymorphisms are then applied to the full set of organisms under investigation. However, the ability of CAPS to detect DNA polymorphism is not as high as SSRs or AFLPs because nucleotide changes affecting restriction sites are essential for the detection of DNA polymorphism by CAPS. Furthermore, the development of CAPS markers is only possible where mutations disrupt or create a restriction enzyme recognition site. Advantages of CAPS include the involvement of PCR requiring only low quantities of template DNA, the codominance of alleles, and the high reproducibility. The results can be easily scored and interpreted. Compared to RFLPs, CAPS analysis does not include the laborious and technically demanding steps of Southern blot hybridization and radioactive detection procedures. However, in comparison with RFLP analysis, CAPS polymorphisms are more difficult to find because of the limited size of the amplified fragments. Furthermore, sequence data are needed to design PCR primers.

PCR-RFLP has been used for authentication of *Alisma* (Li et al. 2007), *Angelica* (Watanabe et al. 1998), *Sinopodophyllum* and *Dysosma* (Gong et al. 2006), *Ephedra* (Guo et al. 2006), *Fritillaria* (Wang et al. 2005, 2007b), *Artemisia* (Lee et al. 2009), *Panax* (Diao et al. 2009; Do et al. 2001; Lu et al. 2010b; Um et al. 2001), *Actinidia* (Zhao et al. 2007b), *Atractylodes* (Mizukami et al. 2000), *Glehnia* (Mizukami et al. 1993a), *Astragalus* (Lu et al. 2009), *Dendrobium* (Zhang et al. 2005a), *Duboisia* (Mizukami et al. 1993b), and *Codonopsis* (Fu et al. 1999).

2.4.1.14 SCAR (Sequence Characterized Amplified Region)

In 1993, Paran and Michelmore introduced a new type of RAPD-derived molecular marker, which circumvented several of the drawbacks inherent to RAPDs. A SCAR marker can be used to rapidly amplify a diagnostic nucleic acid from herbal materials using a pair of specific oligonucleotide primers designed from polymorphic RAPD (Semagn et al. 2006; McDermott et al. 1994) or ISSR (Albani et al. 2004) fragments.

Polymorphic fragments from RAPDs or ISSR are selected among amplified fingerprints. After cloning and sequencing for the selected polymorphic regions, pairs of internal primers are designed to amplify a unique and specific sequence designed as a SCAR marker. SCARs are advantageous over RAPD markers as they detect only a single locus, their amplification is less sensitive to reaction conditions, and they can potentially be converted into codominant markers (Paran et al. 1991). Prior sequence information (i.e., sequencing the polymorphic fragments) is required for designing the primers flanking the polymorphic region. As PCR inhibitory effects of ingredients can lead to false negative results, amplification of a control fragment using the same DNA template should be performed to ensure that the quality of sample DNA is suitable for PCR. The concept of generating locus-specific SCARs from anonymous PCR fragments is not restricted to RAPDs but was applied to other multilocus marker techniques such as ISSR or AFLP (Shan et al. 1999; Xu et al. 2001).

The SCAR technique has been used for authentication of *Panax* (Wang et al. 2001; Choi et al. 2008) and for discrimination of species of *Artemisia* (Lee et al. 2006a), *Phyllanthus* (Dnyaneshwar et al. 2006; Theerakulpisut et al. 2008), *Pueraria* (Devaiah and Venkatasubramanian 2008a), *Sinocalycanthus* (Ye et al. 2006), *Embelia* (Devaiah and Venkatasubramanian 2008b), and *Lycium* (Sze et al. 2008).

2.4.1.15 SSCP (Single-Strand Confirmation Polymorphism)

SSCP is a powerful mutation detection system. The principle of this technique is that under neutral condition, the single-stranded DNA (ssDNA) folds into a tertiary structure. Differences in DNA sequences (often a single base pair) alter the single stranded DNA in the tertiary conformation (by differential folding), which in turn affect the mobility of the ssDNA in a gel. Based on their mobility differences, SNPs can be detected (Orita et al. 1989).

F-SSCP is an adapted version of SSCP analysis involving amplification of the target sequence using fluorescent primers (Makino et al. 1992).

The method is not frequently applied for authentication e.g., in *Boesenbergia* (Techaprasan et al. 2008).

2.4.1.16 MSAP (Methylation-Sensitive Amplified Polymorphism)

This tool is a modification of the AFLP technique and was developed for monitoring the state of genomic DNA methylation. Genomic DNA is double-digested with one of the methylation-sensitive enzymes *HpaII* or *MspI* and then with the methylation-insensitive *EcoRI*. The resulting fragments are ligated with the corresponding double-stranded adapters, a first preselective amplification is carried out followed by a selective amplification step.

MSAP was first developed to determine DNA methylation events in dimorphic fungi (Reyna-Lopez et al. 1997) and later adapted for the detection of cytosine methylation in the rice genome (Xiong et al. 1999), pepper (Portis et al. 2004), apple (Xu et al. 2000), and Siberian ginseng (Chakrabarty et al. 2003).

2.4.1.17 LAMP (Loop-Mediated Isothermal Amplification)

Loop-mediated isothermal amplification (LAMP), which amplifies target nucleic acids with high specificity, efficiency, and rapidity under isothermal conditions was developed by Notomi et al. (2000). This method relies on auto-cycling strand displacement DNA synthesis performed by a DNA polymerase with high strand displacement activity. A specially designed set of two inner and two outer primers is used. An inner primer containing sequences of the sense and antisense strands of the target DNA initiates LAMP. The following strand displacement DNA synthesis primed by an outer primer releases a single-stranded DNA. This serves as template

for DNA synthesis primed by the second inner and outer primers that hybridize to the other end of the target, which produces a stem–loop DNA structure. In subsequent LAMP cycles, one inner primer hybridizes to the loop on the product and initiates displacement DNA synthesis, yielding the original stem–loop DNA and a new stem–loop DNA with a stem twice as long. The cycling reaction continues with accumulation of 10^9 copies of target in less than an hour. The final products are stem–loop DNAs with several inverted repeats of the target and cauliflower-like structures with multiple loops formed by annealing between alternately inverted repeats of the target in the same strand. Because LAMP recognizes the target by six distinct sequences initially and by four distinct sequences afterwards, it is expected to amplify the target sequence with high selectivity (Nagamine et al. 2001, 2002).

This technique was applied to detect *Panax ginseng* (Sasaki et al. 2008), the botanical source of Ginseng (Ginseng Radix), and to distinguish this species from *Panax japonicus*. It was also used for the detection of *Lophophora williamsii* (Sasaki et al. 2009) and *Curcuma longa* (Sasaki and Nagumo 2007).

2.4.1.18 SDA (Subtracted Diversity Array)

PCR-based plant identification techniques are often limited by their low throughput, whereas hybridization-based microarray technology represents a rapid and high-throughput tool for genotype identification. Using an innovative technique, a “Subtracted Diversity Array” (SDA) was constructed from a pooled genomic DNA library of 49 angiosperm species, from which pooled non-angiosperm genomic DNA was subtracted (Jayasinghe et al. 2007). The subtraction was carried out using the Clontech PCR-Select cDNA Subtraction Kit. This new SDA method was shown to be superior to conventional molecular identification methods in terms of accuracy, sensitivity, and efficiency, as well as capacity for high-throughput and broad application. The SDA technique was validated for potential genotyping use. The results indicated a successful subtraction of non-angiosperm DNA. This study demonstrates the potential of establishing a highly informative, reliable, and high-throughput microarray-based technique for the novel application of sequence-independent genotyping of major angiosperm clades.

Niu et al. (2011) showed that SDAs technique is suitable to differentiate two ginseng species, *Panax ginseng* and *Panax quinquefolius*, that are frequently mixed for adulteration. Further, SDA was sensitive enough to detect a deliberate adulteration of 10 % *P. quinquefolius* in *P. ginseng*. Thirty-nine species-specific features including 30 *P. ginseng* specific and nine *P. quinquefolius* specific were obtained. This resulted in a feature polymorphism rate of 10 % from the 376 features used for fingerprinting the two ginseng species. The functional characterization of 14 *Panax* species-specific features by sequencing revealed one putative ATP synthase, six putative uncharacterized proteins, and two retroelements to be different in these two species.

2.4.1.19 MLPA (Multiplexed Ligase-Dependent Probe Amplification)

An assay well suited to medicinal plant species identification is the Multiplexed Ligase-dependent Probe Amplification (MLPA) assay (Barthelson 2009; Shen and Wu 2009). MLPA is a semi-quantitative PCR-based technique initially developed by Schouten et al. (2002). It uses the sensitivity of the polymerase chain reaction, but increases the specificity by including a key ligation step for those MLPA probes that hybridize to a DNA sequence. Several key features distinguish this technique from other PCR-based techniques. First, the amplification is ligation dependent as amplification of non-ligated oligoes does not take place. Since ligation occurs at high temperatures, specificity is further ensured. Second, the amplification is highly multiplexed, allowing the detection of up to 50 targets in one single tube assay. This further allowed mutation testing at the exonic and sub-exonic level for a single gene or multiple genes in one assay, thus becoming a cost-effective medium-throughput test. Third, although this is not a real-time PCR because the primers are in excess of templates and the amplification is in linear range, the amount of amplicons generated at the end is in proportion to the templates (ligation products). Last and more importantly, a common PCR primer is used for the amplification of all target sequences, which is a key feature to ensure the relative quantification of each target with respect to a control sample. Due to its low costs, excellent sensitivity, reliability, and ease of development and implementation, the MLPA technique has become a very popular research and diagnostic tool.

2.4.1.20 Real-Time PCR

Real-time quantitative PCR allows the sensitive, specific, and reproducible quantitation of nucleic acids. Since its introduction, real-time quantitative PCR has revolutionized the field of molecular diagnostics and has become the main technical platform for nucleic acid detection in research and development, as well as in routine diagnostics (Klein 2002; Arya et al. 2005). The intention of real-time PCR is the detection of a specific DNA sequence in a sample by measuring the accumulation of amplified products during the PCR using fluorescent technology. Consequently it allows the researcher to better determine the amount of starting DNA in the sample before the amplification process. Present day real-time methods generally involve fluorescence labeling to show the amount of DNA present at each cycle of PCR. In a few cases real-time PCR has recently been used in identification of Chinese medicinal plants (Matsuyama and Nishi 2011; Slanc et al. 2006).

DNA sequence analysis of rDNA internal transcribed spacer (ITS) and fluorescence melting curve analysis using real-time PCR were applied for authentication of the traditional Chinese medicinal plant *Cimicifuga foetida* from four substitutes: *C. heracleifolia*, *C. dahurica*, *C. acerina*, and *C. simplex*. According to the melting temperature—which is a function of the GC/AT ratio, length, and nucleotide sequences of the amplified product—*C. foetida* was differentiated from all other

species (Ying et al. 2009). Based on real-time PCR technology there are further studies for authentication in the genera *Euphorbia* (Xue et al. 2008a), *Gentiana* (Xue et al. 2008b), and *Drynaria* (Xue and Xue 2008) species from adulterants.

The ongoing development of quantitative DNA-based methods using real-time PCR could enable a quantitative analysis of species composition in mixed plant materials and products in the future (Table 2.2).

2.5 DNA Sequencing Analysis

DNA sequencing is the process of determining the precise order of the nucleotides or bases (A,T,G,C) in a particular DNA molecule. The most common approach used for DNA sequencing is the dideoxy or Sanger method which was developed in the mid 1970s. It mimics the basic process used to copy DNA in a cell during chromosomal replication, except that the procedure is done in a tube or microtiter plate using a minimal set of components. Normally the length of the “sequence read” can vary from about 50 to more than 1,000 bases. If the region to be sequenced exceeds the length of a typical sequencing read internal primers have to be used to generate overlapping in order to reconstruct the complete sequence of a longer DNA region. Most large-scale DNA sequencing facilities use fluorescent dyes (rather than a radioactivity isotope) to label and detect the four bases, and capillary electrophoresis to separate DNA molecules on the basis of size. Because the end of each terminated molecule contains a dye-labeled base, the sequence of the strand complementary to the template can be determined. Modern Capillary Sequencer can run plates of 96 samples in a couple of hours and can produce read lengths of more than 700 bases. Using software provided by the manufacturers the signal of the dyes is determined for each position so that the proper base can be identified. The order of the bases is displayed in a chromatogram or trace file.

In the last decade, instrumentation for DNA sequencing has improved dramatically in terms of increasing read length and accuracy, high throughput, and decreasing costs. Meanwhile there are several next-generation DNA sequencing platforms, such as Roche’s (454) pyrosequencing system, Illumina’s Solexa Genome Sequencer or Applied Biosystem’s SOLiD Genome Sequencer.

Currently DNA sequencing is applied in various fields as in analysis of phylogenetic relationship, population genetics, systematics, and evolution (Baldwin et al. 1995). DNA polymorphisms are revealed by determining the nucleotide sequence in a defined region of the genome and aligning the sequence with homologous regions of related organisms (Alvarez and Wendel 2003).

By choosing appropriate regions of the nuclear, plastidal, or mitochondrial genome this approach provides a highly reproducible analysis at various taxonomic ranks to differentiate TCM plants from its substitutes or adulterants.

In order to ensure a correct species identification based on DNA sequence data it is necessary to have herbarium specimen for verification or a reliable database that guarantees that the reference specimen was correctly identified by a taxonomic

Table 2.2 Comparison of DNA profiling techniques for herbal authentication and quality control

	Importance for authentication	Reproducibility	Quantity of DNA required	Level of polymorphism	Locus specificity	Technical demand	Sequence information required	Automation	Running costs	Development costs
RAPD	+	Low	Low	Medium	No	Low	No	Yes	Low	Low/medium
ISSR	++	Medium/high	Low	Medium	No	Low/medium	No	Yes	Low/medium	Low
AFLP	++	High/medium	Medium	Medium	No	Medium	No	Yes	Medium	Low
RFLP	+	High	High	Medium	Yes	High	Yes	No	High	Medium/high
ARMIS	++	High	High	Low	Yes	Low	Yes	Yes	Low	Medium
SSR	+	High/medium	Low	High/medium	No	Low/medium	Yes/no	Yes	Medium	High
CAPS	++	High	Low	Low/medium	Yes	Low/high	Yes	Yes	Low	Medium
SCAR	++	High	Low	Low/medium	Yes	Low/medium	Yes	Yes	Low	Medium
RAMPO	+	Medium	Low	Medium	Yes	High	No	Yes	Medium	Medium
CAPS	++	High	Low	Low/medium	Yes	Low	No	Difficult	Medium	Medium
SSCP	+	Medium	Low	Low	Yes	Medium/high	Yes	No	Low/medium	High
LAMP	+	High	Low	Low	Yes	Medium	Yes	Yes	Low	Medium
SNP	+	High	Low	High	Yes	Medium	Yes	Yes	Low	High
Sequencing	+++	High	Low	Low/medium	Yes	Medium	Yes	Yes	Medium	Medium
DNA-barcoding										
Microarray	+++	High	Low	High	Yes	High	Yes	Yes	High	High

The table is based on the most frequent molecular tools and does not represent all DNA-based molecular techniques. Strength and weakness of DNA methods are also summarized by Weising et al. (2005), Yip et al. (2007), Pereira et al. (2008), Hao et al. (2010), Mondini et al. (2009), and Agarwal (2008)

expert. Additionally the sequence should be obtained in independent studies including related taxa. A common way to assign a particular sequence to a taxon is to perform a BLAST search (Basic Local Alignment Search Tool) in the databases of NCBI (GenBank), BOLD, MMDBD. However, care must be taken when assigning the questioned sequence to the species with the highest similarity, because several gaps and false sequences are known to be present in these databases (Heubl 2010).

There are many studies concerning the application of DNA sequence-based markers to differentiate medicinal taxa used in TCM from its substitutes or adulterants.

Sequencing analyses based on nuclear ITS have been applied to *Panax* (Ngan et al. 1999; Kim et al. 2007b), *Asarum* (Kelly 1998; Liu et al. 2005; Yamaji et al. 2007), *Astragalus* (Dong et al. 2003; Yip and Kwan 2006), *Dendrobium* (Xu et al. 2006; Ding et al. 2002; Lau et al. 2001; Zhang et al. 2003), *Fritillaria* (Wang et al. 2005), *Leonurus* (Yang et al. 2006), *Perilla* (Luo et al. 2006b), *Phyllanthus* (Lee et al. 2006b), *Rehmannia* (Albach et al. 2007), *Salvia* (Wang et al. 2005), *Swertia* (Xue et al. 2006a), *Plantago* (Sahin et al. 2007), *Bupleurum* (Yang et al. 2007), and *Euphorbia* (Xue et al. 2006b).

Another frequently used marker is the nuclear 5S rDNA intergenic spacer used for authentication of *Adenophora* (Zhao et al. 2003a), *Aconitum* (Carles et al. 2005), *Angelica* (Zhao et al. 2003b), *Astragalus* (Dong et al. 2003; Ma et al. 2000), *Curcuma* (Xia et al. 2005), *Epimedium* (Sun et al. 2004), *Fritillaria* (Cai et al. 1999), *Crocus* (Ma et al. 2001), *Ligularia* (Zhang et al. 2005b), *Pueraria* (Sun et al. 2007), *Saussurea* (Chen et al. 2008).

From nuclear DNA also 18S rDNA has been tested in *Dioscorea* (Liu et al. 2001a), *Pinellia* (Liu et al. 2001b), and *Panax* (Zhu et al. 2003).

From chloroplast DNA a couple of markers including genes, intergenic spacers, or introns are applied. The *atpB-rbcL* region was used for differentiation of *Phyllanthus* (Lee et al. 2006b), *trnC-trnD* in *Panax* (Lee and Wen 2004), *trnL-F* in *Pueraria* (Albach et al. 2007), *Rheum* (Yang et al. 2001) and *Ephedra* (Long et al. 2004), *rpl16* in *Swertia* (Xue et al. 2006b), *rpl16-rpl14* spacer in *Scutellaria* (Hosokawa et al. 2005), *atpF-atpA* in *Angelica* (Hosokawa et al. 2006), *trnD-trnT* in *Diosma* (Gong et al. 2006), *trnK* in *Actinidia* (Zhao et al. 2007c), *Attractylodes* (Mizukami et al. 2000) and *Curcuma* (Sasaki et al. 2002), *matK* in *Agastache* (Luo et al. 2002), *Panax* (Komatsu et al. 2001), *rbcL* in *Dryopteris* (Zhao et al. 2007c), *Cnidium* (Kondo et al. 1996), or *Pinellia* (Lin et al. 2006).

Recently sequencing analysis has been applied to proof the possibility of DNA-based authentication of plant extracts. Internal transcribed spacer (ITS) was successfully amplified from different extracts types from *Echinacea* species and *Matricaria chamomilla* (Novak et al. 2007).

2.6 DNA Barcoding

DNA barcoding, a term first created by Hebert et al. (2003a) is a novel molecular and bioinformatical tool designed to provide rapid, accurate, automatable, and cost-effective identification of species. Contrary to other molecular methods, it can be used on a large scale and with high reliability. For DNA barcoding, the unique nucleotide sequence patterns of small DNA fragments (400–800 bp) are used as specific reference collections to identify specimens and to discover cryptic taxa (Vijayan and Tsou 2010).

DNA barcoding uses a short genetic marker from a standard locus (alternatively from nuclear, mitochondrial, or plastidial DNA) of an organism. An ideal and successful DNA barcode marker should be suitable for a wide range of taxa (breadth of taxonomic application), routinely retrievable with a universal primer pair, be short enough to be accessible to bidirectional sequencing, and provide a unique sequence for maximal discrimination among species which means high variation between species but conserved within the species, so that the intra-specific variation will be insignificant (CBOL 2009). Additionally these markers should be flanked by evolutionary conserved regions so that universal primers can be used and they should be free of insertions or deletions to be easily alignable (Shneyer 2009) (Fig. 2.1).

2.6.1 Barcoding and Herbal Monographs

DNA authentication of medicinal species used in TCM is of major importance because there are many cases in the Chinese Pharmacopoeia where several species are listed under one common name or listed as synonyms, subspecies, or varieties in herbal monographs.

A prime example is the genus *Dendrobium* (Orchidaceae) which is represented in the Flora of China by 78 species (14 endemic). Overlapping distribution ranges coupled with high morphological variation makes proper species identification difficult. Based on processing methods, Herba Dendrobii is classified into fresh *Dendrobium*, “Fengdou Shihu,” and “Huangcao Shihu,” the latter being the predominant form of Herba Dendrobii on Chinese herbal medicine market.

As source material for Herba Dendrobii the Pharmacopoeia of the People’s Republic of China (2010 edition) lists four species, *D. catenatum* Lindl. (syn. *D. officinale* K. Kimura and Migo), *D. nobile* Lindl., *D. chrysotoxum* Lindl., and *D. fimbriatum* Hook. as the authorized plant sources. However in the Chinese Materia Medica Dictionary (2006 edition), 16 further species are enumerated as medicinal plants: *D. aduncum* Wall. ex Lindl., *D. bellatulum* Rolfe, *D. chrysanthum* Wall. ex Lindl., *D. chryseum* Rolfe, *D. crepidatum* Lindl. ex Paxt, *D. cucullatum* R. Br., *D. densiflorum* Lindl. ex Wall., *D. devonianum* Paxt, *D. hancokii* Rolfe, *D. henryi* Schltr., *D. hercoglossum* Rchb. fil., *D. linawianum* Rchb. fil., *D. loddigesii* Rolfe, *D. lohohense* Tang et Wang, *D. longicornum* Lindl., and *D. moniliforme* (L.) Sw. Further *Dendrobium* species that are not mentioned in either of these references are

Table 2.3 Plant DNA barcoding markers tested for their suitability and recommendations for barcoding in land plants and Chinese medicinal plants (Ycf5 is also known as ccsA)

DNA segment tested for suitability	Proposed/recommended DNA markers	References
nrITS, atpB-rbcL, psbM-trnD, trnC-ycf6, trnH-psbA, trnL-F, trnK-rps16, trnV-atpE rpl36-rps8, ycf6-psbM	nrITS + trnH-psbA	Kress et al. (2005)
ITS1, accD, ndhJ, matK, trnH-psbA, rbcL, rpoB, rpoC1, ycf5	rbcL + trnH-psbA	Kress and Erickson (2007)
atpF-atpH, atpH-atpI, rps15-ycf1, ndhG-ndhI, psbK-psbI, petA-psbJ, trnH-psbA	atpF-atpH + psbK-psbI	Lee et al. (2007)
rpoC1, rpoB, matK, trnH-psbA, nrITS, trnL-F	rpoC1 + rpoB + matK or rpoC1 + matK + trnH-psbA	Chase et al. (2007)
nrITS, accD, ndhJ, matK, trnH-psbA, rpoB, rpoC1, ycf5	nrITS	Sass et al. (2007)
accD, matK, trnH-psbA, rbcL, rpoB, rpoC1, UPA	matK + trnH-psbA	Newmaster et al. (2008)
TrnL (UAA) intron	trnL (UAA) intron	Taberlet et al. (2007)
matK, trnH-psbA, psbK-psbI, atpF-atpH	matK or matK + trnH-psbA + psbK-psbI	Lahaye et al. (2008a)
accD, matK, trnH-psbA, rbcL, rpoB, rpoC1, ycf5, ndhJ	matK or matK + trnH-psbA	Lahaye et al. (2008b)
Cox1, 23S rDNA, rpoB, rpoC1, rbcL, matK, trnH-psbA, atpF-atpH, psbK-psbI	matK and psbK-psbI + trnH-psbA + atpF-atpH	Fazekas et al. (2008)
trnH-psbA, rbcL, rpoC, CO1, rpoB, matK	trnH-psbA + rbcL or rpoB or rpoC	Erickson et al. (2008)
atpF-atpH, rpoB, rpoC1, rbcL, matK, psbK-psbI, trnH-psbA	rbcL + matK	Hollingsworth et al. (2009)
accD, matK, ndhA, ndhJ, ndhK, rpl22, rpoB, rpoC1, rpoC2, ycf2, ycf5, ycf9	rpoC1 + rpoB + matK	Ford et al. (2009)
atpF-atpH, matK, rbcL, rpoB, rpoC1, psbK-psbI, trnH-psbA	rbcL + matK	CBOL-PWG (2009)
psbA-trnH, matK, rbcL, rpoC1, ycf5, ITS2, and ITS	ITS2 + psbA-trnH	Chen et al. (2010)

Proposed markers are based on universality, success of PCR amplification, interspecific sequence variability, and species discriminatory power/identification capability. Several proposed combinations of loci have been summarized by Pennisi (2007), Shneyer (2009), and Vijayan and Tsou (2010). The combination *rbcL* + *trnL-F* (intron and IGS) has been proposed as a two-locus barcode for ferns (De Groot et al. 2011). The best performing single loci for mosses were the *rbcL* and *rpoC1* coding regions (Liu et al. 2010b)

additionally processed. A survey of the medicinal *Dendrobium* species revealed that in total 27 other *Dendrobium* species are also used clinically as substitutes for “Huangcao Shihu” at various places in China, a situation which may cause inconsistent therapeutic effects or even endanger safety of consumers (Li et al. 1986, 1991; Ma et al. 1995). The stems of *Dendrobium* species, have very similar morphological and anatomical characteristics, and the traditional authentication




Fig. 2.1 Major steps in the DNA barcoding process. These procedures include tissue sampling, DNA extraction, polymerase chain reaction and marker amplification, PCR product check, and sequencing. The end product of this procedure will be a unique, species-specific “barcode” sequence of the marker (gene, spacer, intron) that can be used for species identification and submitted to a DNA database (e.g., GenBank-NCBI, Bold). The original specimen is kept as a voucher. All collateral information (identification, collection data, etc.) are stored along with the DNA barcode sequence

of different “Huangcao Shihu” samples is therefore far from reliable. In addition, the chemical constituents of many *Dendrobium* species are still unknown, as proper chemical analysis methods have not been developed. However, the determination of the botanical origins of different “Huangcao Shihu” samples and their quality control through morphological and chemical studies is fraught with difficulty. Consequently DNA barcoding may offer an alternative method for the identification of the used *Dendrobium* species for herbal medicines (Xu et al. 2006; Takamiya et al. 2011).

A further remarkable example where multiple species are used in TCM is the genus *Fritillaria* (Liliaceae) which includes 24 species (15 endemic) in China. Bulbs of various *Fritillaria* species (Bulbus Fritillariae) are among the most popular herbal medicines in China and have been used as antitussive and expectorant herbs. According to the Chinese Pharmacopoeia, herbal “Beimu” is derived from the bulbs of nine *Fritillaria* species. These include *Fritillaria thunbergii* Miq. (“Zhebeimu”), *F. cirrhosa* D. Don. (Chuanbeimu), *F. unibracteata* Hiao et Hsia, *F. przewalskii* Maxim ex Batal, *F. delavayi* Franch, *F. ussuriensis* Maxim. (Pingbeimu), *F. walujewii*, *F. pallidiflora* Schrenk (“Yibeimu”), and *F. hupehensis* Hsiao et K.C (“Hubeibeimu”).

Furthermore other *Fritillaria* species (e.g., *Fritillaria anhuiensis* S.C. Chen et S.F. Yin, *Fritillaria puziensis* G.D. Yu et G.Y. Chen) are often used in different regions in China as substitutes. These distinctions can be very confusing for consumers. As there are no specific microscopic characteristics and the pattern and concentration of chemical components are often unstable, for quality control and standardization of *Fritillaria* derived herbal medicine, exact species identification using DNA barcoding is the only reliable method (Wang et al. 2007b).

Another example where the Chinese Pharmacopoeia allows some interchangeable use of herbals is the genus *Epimedium* (Berberidaceae) which is one of the most popular herbal drug (“Yinyanhua”) comprising 41 species (40 endemic) in China. Herba Epimedii is prepared from aerial parts of five *Epimedium* species listed in the Chinese Pharmacopoeia: *E. brevicornum* Maxim., *E. koreanum* Nakai, *E. sagittatum* (Sieb. et Zucc.) Maxim., *E. pubescens* Maxim., and *E. wushanense* T. S. Ying. In common with many traditional medicinal plants, a major problem with *Epimedium* is the absence of a rigorous method to authenticate species. Taxonomists have variably reported numbers ranging from 20 to 50 species (Sun et al. 2005). Traditional herbalists do not differentiate among *Epimedium* species, but rather use a mixture of species together as *Herba Epimedii*. These species differ significantly in concentrations of major and minor constituents (Wu et al. 2003). The extensive variation in morphological traits make the species difficult to classify taxonomically and there is some confusion due to lack of scientific research.

Considering the examples mentioned above there are numerous monographs in the Chinese Pharmacopoeia where multiple species (further examples *Dioscorea* and *Uncaria* each with five species listed) are a legitimate source for the preparation of herbal medicines. As these species have a different composition of bioactive compounds and are quantitatively and qualitatively not equivalent, the therapeutic efficacy of a TCM medication can vary to a high extent. For species-rich genera lacking morphological characters or taxa of controversial botany, DNA barcoding as a new tool is a prerequisite to clearly distinguish the authentic herb from the substitute/adulterant. If the barcode DNA-analysis of all frequently used

Chinese drugs becomes available in the near future, the application of this method and further DNA fingerprinting techniques should be included in monographs to optimize the quality and safety proof of the drugs (Fig. 2.2).

2.6.2 Standard Markers for Barcoding

In certain animal groups the mitochondrial cytochrome *c* oxidase subunit 1 (CO1, *cox1*) gene sequence (ca. 650 bp in length) is currently used as a universal DNA barcode (Hebert et al. 2003b; Waugh 2007). It has been introduced as standard marker as it can be easily sequenced and provides greater than 97 % species-level specificity in different animal phyla, e.g., birds (Hebert et al. 2004), mammals (Luo et al. 2011), fishes (Ward et al. 2009), amphibians (Vences et al. 2005), and various groups of arthropods (Virgilio et al. 2010). Based on this mitochondrial




Fig. 2.2 DNA sequence BLAST search against the databases of GenBank (NCBI) and BOLD using the barcoding markers *ITS* and *rbcL + matK* sequences of *Artemisia annua* as a query. NCBI (<http://www.ncbi.nlm.nih.gov>) offers the BLAST search tool to perform fast searching with rigorous statistical methods for judging the significance of matches. The Barcode of Life Data Systems (BOLD; <http://www.boldsystems.org/views/login.php>) and its Identification System (IDS) for *rbcL* and/or *matK* is an identification tool for plant species. The ID engine uses all sequences uploaded to BOLD to locate the closest match. It returns a species-level identification when possible

gene, DNA barcoding is progressing rapidly in many groups of animals, which can be reviewed online via the Canadian Barcode of Life (<http://www.bolnet.ca>) and the Consortium for the Barcode of Life (CBOL; <http://www.barcoding.si.edu>). Other alternative mitochondrial regions such as cytochrome b, 12S, 18S (SSU), 28S (LSU) rRNA, and the nuclear ITS2 region have been proposed (but not established) for animal identification (Wong et al. 2001; Chen et al. 2010; Ferri et al. 2009; Yao et al. 2010).

However, the cytochrome c oxidase marker, which is widely used in animal barcoding, seems promising only for algae (Robba et al. 2006; Lane et al. 2007) but it is not suitable for higher plants because of a much slower substitution rate in *cox1* which is two- to threefold lower than in chloroplast genes and 10- to 20-fold lower than in nuclear genes (Wolfe et al. 1987; Drouin et al. 2008). Furthermore the mitochondrial genome in plants is characterized by the occurrence of large structural rearrangements and often nonfunctional copies of mitochondrial genes (pseudogenes) which cause erroneous results. Consequently in plants appropriate barcoding regions have been analyzed, tested, and selected for discriminating

among the 260,000–422,000 species of seed plants (Govaerts 2001; Thorne 2002), but until now no agreement exists which marker is the most promising, though most researchers agree that more than one region is necessary. Consequently a multilocus approach based on the plastid genome with a focus on coding and noncoding regions (introns or spacers) is currently the most effective strategy for species identification in plants (Chase et al. 2005, 2007; Kress et al. 2005; Newmaster et al. 2006, 2008; Cowan et al. 2006; Ford et al. 2009; Kress and Erickson 2007; Lahaye et al. 2008a, b; Fazekas et al. 2008).

In order to promote the use of DNA barcoding for all eukaryotic organisms, the Consortium for the Barcode of Life (CBOL, <http://barcoding.si.edu>) was established in 2004 at the National Museum of Natural History in Washington, which currently includes more than 120 organizations from 45 nations.

In 2005 the Plant Working Group (PWG CBOL) reported that five very promising regions were chosen for further barcoding studies. These regions were *matK* (encodes a maturase and is located within the *trnK* gene), *rpoB* (RNA polymerase subunit), *rpoC1* (RNA polymerase subunit), *accD* (subunit of acetyl-CoA carboxylase), and *ccsA* (previously known as *ycf5*; the gene encoding a protein involved in cytochrome *c* biosynthesis).

In a subsequent work in 2007, additional candidate plastid DNA regions were tested in different groups (gymnosperms, angiosperms, ferns, equisetes, and mosses).

Based on assessments of recoverability, sequence quality, and levels of species discrimination, a core two-locus combination of *rbcL* and *matK* as the plant barcode was recommended. This combination was shown to successfully discriminate among 907 samples from 550 species at the species level with a probability of 72 % (CBOL Plant Working Group 2009). The group admits that the two-locus barcode is far away from perfection due to the limited identification rate, and thus further research for other appropriate candidates is necessary. Additional combinations of noncoding and coding plastid regions have been tested for barcoding purposes (Fazekas et al. 2008; Pennisi 2007; Ford et al. 2009).

Other combinations involving three plastid regions have also been proposed by various working groups that include the Royal Botanical Gardens, Kew, UK <http://www.rbgekew.org.uk/barcoding> (Chase et al. 2007).

In view of the available markers as a core barcode to identify land plants a combination of two loci from plastidal DNA, *rbcL* + *matK* was adopted as “the plant barcode” by the Executive Committee of the Consortium for the Barcoding of Life (CBOL Plant Working Group 2009) after much deliberation (e.g., Kress et al. 2005; Hollingsworth et al. 2009). They agreed that *rbcL* and *matK* are approved and the most suitable barcode regions for land plants. Based on these markers, the BOLD Identification System (BOLD-IDS) provides a species identification tool that accepts DNA sequences with a minimum sequence length of 500 bp from these two barcode region and returns a taxonomic assignment at the species level (a list of the nearest matches) when possible. The group admits that the two-locus barcode is far away from perfection due to the low identification rate and that the search is not over.

Additionally the plastidal *psbA-trnH* intergenic spacer and internal transcribed spacer region (ITS1-5.8S rRNA-ITS2) of nuclear ribosomal DNA have been tested as supplementary barcodes. Kress et al. (2005) suggested that these two noncoding regions might have potential as universal plant barcodes. Generally the ITS, which is part of the ribosomal operon, is organized in large blocks (tandem arrays of nearly 50–100 copies) in the chromosome of the nuclear organizer region. It is one of the most commonly sequenced DNA regions (ca 700 bp) used in plant phylogenetic studies at the generic and species level. The advantage of these spacers lies in their high variability. They are adjacent to the conserved 5.8S rRNA region and flanked by conserved 18S and 26S rRNA genes, which facilitate the development of primers. Meanwhile more than 50,000 plant ITS sequences have been deposited in GenBank (Hajibabaei et al. 2007). As nuclear ITS is arranged in multiple copies, paralogs may occur due to hybridization and subsequent incomplete concerted evolution. As a consequence in some groups the presence of several functional copies was detected (Alvarez and Wendel 2003). Not only the sequence is an effective barcode but also the secondary structure of the ITS2 region could provide useful information for species identification and has potential as a molecular morphological characteristic (Yao et al. 2010).

The *trnH-psbA* spacer, though short in length (ca. 450 base pairs), is one of the most variable plastid regions in angiosperms and can be easily amplified across a broad range of land plants (Shaw et al. 2007). In general intergenic spacers and introns tend to be more variable than genes and are therefore considered as better identifiers.

Recently, Chen et al. (2010) compared seven candidate DNA barcodes from Chinese medicinal plant species. This study including a total of 4,800 species from 753 distinct genera revealed that the Internal Transcribed Spacer 2 (ITS2) of nuclear rDNA and the *psbA-trnH* locus are promising universal barcodes for plant identification. A test of the discrimination ability of ITS2 showed that the rate of successful identification at species level was 92.7 % for ITS2 and 72.8 % for *psbA-trnH*.

Shaw et al. (2007) compared chloroplast genomes of Solanaceae, Fabaceae, and Poaceae and showed that no less than nine intergenic spacers, which were practically not used in molecular phylogenetic studies (*rpl32-trnL*, *trnQ-rps16*, *trnV-ndhC*, *ndhF-rpl32*, *psbD-trnT*, *psbJ-petA*, *rps16-trnK*, *atpI-atpH*, *petL-psbE*), were far more variable than the most variable ones of those spacers used earlier. Probably these regions have potential as DNA barcoding markers, if not universal, then suitable for studies on higher taxonomic level.

Moreover, further DNA barcode markers have been used to identify medicinal plants from different plant families and genera (Chen et al. 2010; Song et al. 2009; Luo et al. 2010; Pang et al. 2010; Gao et al. 2010a; Sun et al. 2010; He et al. 2010) including also medicinal Pteridophytes (Ma et al. 2010) (Table 2.3).

In 2008 several DNA barcoding projects have been initiated in China when the country became a central node of the International Barcode of Life project (*iBOL*). The Barcoding Chinese Plants Project is closely associated with the Germplasm Bank of Wild Species (GBOWS, Kunming Institute) which was founded to

safeguard and barcode about 6,000 species of vascular plants of the flora of China. Based on a nationwide seed collecting program a large data set, involving 6,286 individuals representing 1,757 species in 141 genera of 75 families of seed plants, was assembled by the China Plant BOL Group. Comprehensive analysis of the dataset was made to assess the universality, sequence quality, and discriminatory power of the chosen barcoding markers (Li et al. 2011a).

Meanwhile an integrated DNA barcode database (MMDBD) is under construction (Lou et al. 2010). This platform contains ca. 1,300 species of Chinese medicinal plants listed in the Pharmacopoeia of the People's Republic of China and it provides information on storage, retrieval, comparison, and analysis of DNA sequences, for distinguishing medicinal materials from their common substitutes and adulterants (see <http://www.cuhk.edu.hk/icm/mmdbd.htm>).

In combination with phylogenetic analyses, in recent years barcoding activities have been intensified in many genera including Chinese medicinal plants. Barcoding projects based on nuclear and/or plastidal markers has been applied in different plant families e.g., Rosaceae (Pang et al. 2011), Caprifoliaceae (Liu et al. 2010a), Asteraceae (Gao et al. 2010b), Loranthaceae (Li et al. 2009b), Euphorbiaceae (Pang et al. 2010), Fabaceae (Gao et al. 2010b), Polygonaceae (Song et al. 2009), Rutaceae (Luo et al. 2010), Myristicaceae (Newmaster et al. 2008), Lemnaceae (Wang et al. 2010b), Lamiaceae (Han et al. 2009), as well as in the genera *Panax* (Zuo et al. 2011), *Dendrobium* (Asahina et al. 2010), *Lonicera* (Sun et al. 2010), *Paris* (Zhu et al. 2010), *Aconitum* (He et al. 2010), *Taxillus* (Li et al. 2010c), *Amomum* (Zhen-Yan and Ling 2010), *Astragalus* (Guo et al. 2010), *Paeonia* (Zhang et al. 2009), in Cycadales (Sass et al. 2007), and also in medicinal Pteridophytes (Ma et al. 2010).

Molecular authentication of herbal medicinal materials has increased enormously in the last decade and its advantages are undisputed. This new tool has been included for the first time in the *Pharmacopoeia of the People's Republic of China* and its online Supplementary Note 2 as a standard method for the identification of some traditional Chinese medicines (Li et al. 2011b).

Once fully developed, DNA barcoding has the potential to completely revolutionize our knowledge of plant diversity. In the future scientists will be able to quickly identify known species and retrieve information about them. Compiling a public library of sequences linked to vouchered specimens and their scientific binomial names will make barcoding a practical and efficient tool for the identification of species, including all Chinese medicinal plants (Fig. 2.3).

2.7 DNA Microarrays (DNA Chip Technology)

The DNA chip technology developed by Fodor et al. (1991) enables the production of a "biochip" designed to identify fluorescent-labeled DNA or RNA fragments through their hybridization to oligonucleotide probes. DNA microarrays are a high-throughput technology for simultaneous analysis of multiple genes in many taxa or

Home
Search
BLAST
Data Submission
Help & Information

BLAST

Please input DNA raw sequence

```
actatttggcttacacagttctcttaaaaaatattttatagtttgggtcgattcgcggtttctctcttgg
atccatattcatttatattataggtttgtatattctattccaaaatttttatgaagtttgattccaatt
caatttcaaaccaaaatataaaaaatgcattttgcttatttattacttggataaaaaagaanaataa
tgcctttttatgttgaggtaaaaatagataaactagatagatataatagtagggggcgatgtag
ccaagtggatcaagcgagtgattgtgaat:
```

Step 1

rbcL sequence unknown taxon

Query Bar: score=100 100>=score>50 50>=score>20 20>=score>10 score<=10

Hit bar: score=100 100>=score>50 50>=score>20 20>=score>10 score<=10

gnl|mmdbd100000186|Chloroplast:ribulose-1,5-bisphosphate carboxylase/oxygenase smi-
subunit|Artemisia annua|Compositae|HERBA ARTEMISIAE ANNUAE(526bp)

Score(bit):1011.49 **Taxon identification** e-value:0
Identity:510 Query Strand:1
Alignment: Hsp Strand:1

Query:1 ATCCGCAACGGTTGGGTTCCCTTTGGAAATTTGAAGTGGAAACACGGGTTTGTCTACCGTGAGAACCACA 70
Hit:1 ATCCGCAACGGTTGGGTTCCCTTTGGAAATTTGAAGTGGAAACACGGGTTTGTCTACCGTGAGAACCACA 70
Query:71 GGTCAACAGGGTACTATGATGGAAGAAGTGGACAATGTGGAAGTTGCCCATGTTGGGTGCACCTGACTC 140

Step 2

Home
Search
BLAST
Data Submission
Help & Information

Search:

Species Information

Chinese word's stroke number Search (种属的简体中文笔画数)

-	1	2	3	4
5	6	7	8	9
10	11	12	13	14
15	16	17	18	19
20	21			

Step 3

-help | -Results | |Species| |Sequences(try to click sequence line to get detailed information)

Artemisia annua(黄花蒿)

List of available DNA sequences

ITS

18S

rbcL

psbA-trnH

seqID	DNA Region	Species Name	variant Name	TCM Name
100000058	Internal transcribed spacer	Artemisia annua	-	HERBA ARTEMISIAE ANNUAE
100000060	18S ribosomal RNA gene	Artemisia annua	-	HERBA ARTEMISIAE ANNUAE
100000186	Chloroplast ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit	Artemisia annua	-	HERBA ARTEMISIAE ANNUAE
100000254	Chloroplast psbA-trnH intergenic spacer	Artemisia annua	-	HERBA ARTEMISIAE ANNUAE

Fig. 2.3 (continued)

Medicinal Material Information

Material Name: HERBA ARTEMISIAE ANNUAE (青蒿)
Species: Artemisia annua - (黄花蒿)
Family: Compositae (菊科) **Type:** Plant
Medical Part: Dried aerial part (干燥地上部分)
Pharmacopoeia information: Pharmacopoeia of PRC
CITES (Convention on International Trade in Endangered Species):
Adulterant:

Sequence Information **Similar Sequences:** No Similar Sequences Found

Clone Number:
Voucher Number:
 >nmdbd10000186|Artemisia annua|Chloroplast ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit
 ATCCGCAACGGTTGGGTTCCCTGTTTGGAAATTTGAAGTGGAAACACGGGTTTGTCTACCGTGAGAACCACAGGTACCAGG
 GTACTATGATGGAGAAACATGGACAATGTGGAAATGCCCCATGTTGGGTGCACCTGACTCAGCCCAAGTGTGAAGGAGC
 TCCAGGAGGCAAGAAGGAGTACCCCAACGCATTATCCGTATTATCGGATTCGACACAGCTGCGTCAAGTTCAATGCATC
 AGTTTCATCGCACACAAGCCACAGGCTACTAAGCAAGAAATATGTTGCATGTACCCCTTTTGGGCACCTTACTATGTCT
 ACTGCTGTAATGTTTAGTGTGTGAAATATGCTCAAATAAGTGGCATATATTTGACTAGTGTATTTAAGAGTTAATTA
 CTTGAAAGTGTATCATACATGCCATTTTGTGTTATTGTGTGTTGAACATATTTGTACTTATTGTGTGAATGAACCTT
 CAAATTTGATATTATCGCTATTTTGTGGAAAAAAAAAAAAAAAAA

Sequence information

Reference Information
 Tkach, NV et al. Parallel evolutionary patterns in multiple lineages of arctic Artemisia L. (Asteraceae). **Evolution; international journal of organic evolution** 62 (1), 184-98 (2008) NCBI PMID: [17976192](https://pubmed.ncbi.nlm.nih.gov/17976192/)
Abstract

Taxon information

Early observers of plant evolution in the Arctic have noted a floristic similarity with temperate alpine regions and a predominance of high ploidy levels. The aim of our study was to survey these and other traits in multiple closely related but independently evolved lineages of Artemisia. Our phylogenetic study was based on 133 taxa using 3'-ETS and ITS, and on data on morphology, karyology, distribution, and ecological preferences. We compared Arctic lineages with sister groups and tested whether patterns were significantly different. We found: (1) Artemisia has independently adapted to Arctic habitats 13-18 times; (2) There were no ecological preferences of putative progenitors that might determine the colonization success in the Arctic, although most sister groups were centered in steppe habitats; (3) Plant height was distinctly reduced in Arctic lineages; (4) Arctic lineages contained no more polyploids than their respective sister groups or taxa from other habitats; (5) Enlarged flower heads have evolved repeatedly, probably for better pollinator attraction. This strategy could be a substitute for polyploidy, which is typical in other Arctic taxa. Stronger pollinator attraction should result in better outcrossing and higher heterozygosity in the offspring, which is among the main effects of polyploidy.

Fig. 2.3 Steps in the identification process using the platform MIMDB (Medicinal Materials DNA Barcode Database—<http://137.189.42.34/mherbsdb/index.php>). Search and result page with sequence similarity BLAST search using the *rbcL* sequence of *Artemisia annua* for query (steps 1–3). Information page for medicinal materials including herb name, species name, family name, medical part, pharmacopoeia information, status in CITES, adulterant, DNA sequence (with voucher), and key reference (step 4)

samples (Fodor et al. 1993; Gershon 2002). To apply this technique for identification and authentication of herbal material, it is necessary to identify a distinct DNA sequence that is unique to each species (Preeti et al. 2006a, b). The DNA sequence information is then used to synthesize a corresponding probe on a chip. These probes are capable of detecting complementary target DNA sequences if present in the test sample being analyzed. These immobilized DNA fragments are arranged in

a regular pattern on a microarray by fixation on glass slides, silicon, or nylon membranes (Gebauer 2004).

DNA extracted from the target sample and labeled with a specific fluorescent molecule is then hybridized to the microarray DNA. A positive hybridization result is detected by the intensity of the fluorescence, which reflects the stability of the hybridization between the oligonucleotide probe and the target sequence, and it is visualized with fluorescence scanning or imaging equipment. Each hybridization surface may contain a very large number of unique oligonucleotide probes (up to 400,000), permitting several thousand individual nucleotide positions to be characterized at the same time. The DNA microarray field is a combination of several technologies, including automated DNA sequencing, DNA amplification by PCR, oligonucleotide synthesis, nucleic acid labeling chemistries, and bioinformatics.

Recently this technique has been applied for the identification of various species of *Fritillaria* (Tsoi et al. 2003), *Dendrobium* (Li et al. 2005; Zhang et al. 2003), and *Bupleurum* (Lin et al. 2008). Previously, the nucleotide sequences of the nuclear 18S rRNA gene of 13 *Panax* taxa were determined. On the basis of the nucleotide differences, a DNA microarray (PNX array) was developed for the identification of various *Panax* drugs (Zhu et al. 2008). A silicon-based DNA microarray was designed and fabricated for the identification of toxic traditional Chinese medicinal plants. Species-specific oligonucleotide probes were derived from the 5S ribosomal RNA gene of *Aconitum carmichaeli*, *A. kusnezoffii*, *Alocasia macrorrhiza*, *Croton tiglium*, *Datura innoxia*, *D. metel*, *D. tatula*, *Dysosma pleiantha*, *Dy. versipellis*, *Euphorbia kansui*, *Hyoscyamus niger*, *Pinellia cordata*, *P. pedatisecta*, *P. ternata*, *Rhododendron molle*, *Strychnos nux-vomica*, *Typhonium divaricatum*, and *T. giganteum* (Carles et al. 2005).

The analyses demonstrated that DNA microarray-based technology can provide a rapid, high throughput tool for correct botanical identification, for authentication of crude plant materials, standardization, and for quality control being used for hundreds of samples simultaneously (Debouck and Goodfellow 1999).

This application of DNA microarrays will not only benefit the herbal drug industry but can also facilitate the identification of herbal products by regulatory authorities (Fig. 2.4).

2.8 Limitations of Genetic Markers in Herbal Drug Technology

Molecular authentication methods in comparison to macroscopic, microscopic, and phytochemical analyses have several advantages, which make them suitable for the identification of plants used in TCM. The DNA-based techniques are not affected by environmental factors, independent from the physical form of the plant material, and require only a low amount of material. Although DNA analysis is currently considered to be cutting-edge technology, it has certain limitations:




Fig. 2.4 Workflow overview with major steps of an automated and high throughput DNA microarray platform for species identification

- The applicability of a DNA-based method depends generally on the quality and quantity of the DNA. When plant drugs are processed under extreme conditions DNA degradation (fragmentation) is observed, making PCR amplification impossible. Drug processes involving mechanical stress, high temperature, pH variations, enzymatic activities, and fermentations affect the primary structure of DNA and cause often hydrolysis, oxidation, and deamination of the DNA. Plant-derived polyphenolics are also capable of oxidative DNA damage particularly in the presence of transition metal ions (Bauer et al. 2003; Malik et al. 2003; Novak et al. 2007).
- The isolation of high quality genomic DNA being essential for molecular applications becomes difficult for many medicinally used plants due to the presence of secondary compounds which serve mainly for chemical defense against herbivores. High concentrations of these secondary compounds (polysaccharides, tannins, essential oils, polyphenolics, alkaloids, etc.) often influence DNA extraction, PCR reaction, or restriction digestion. In tissues of medicinal plants, secondary compounds generally get accumulated and the problem becomes increasingly severe as the material gets older. Polysaccharide contaminations are particularly problematic as they can inhibit the activity of many commonly used enzymes, such as polymerases, ligases, and restriction endonucleases. Polyphenol contaminations of DNA make it resistant to restriction enzymes and interact irreversibly with proteins and nucleic acids. Choosing

the most suitable DNA extraction procedure may help to eliminate the PCR inhibitors (Friar 2005; Pirttilä et al. 2001; Fleischmann and Heubl 2010).

- Sometimes plant materials are contaminated with symbiotic or pathogenic microbes. Especially endophytic fungi often occur as symbionts living within the tissues of their angiosperm hosts. DNA isolation techniques for obtaining genomic plant DNA do not discriminate between plant and fungal DNA and PCR primers with broad applicability also amplify DNA contaminants. Fingerprinting using RAPD analyses are particularly vulnerable to these fungal contaminants because of the short length of their primers. Although the amount of endofungal DNA is presumed small as compared to the host genomic DNA, researchers need to consider its potential presence. Problems with endophytic fungi can be eliminated with plant-specific primer design for nuclear markers (e.g., ITS) (Saar et al. 2001).
- The orthology of characters is one of the fundamental and implicit assumptions in the use of DNA sequence data to reconstruct phylogeny or to establish barcodes for species. However, some studies revealed the presence of some degree of intra-individual variations among the copies of ITS1 and ITS2 sequences of the nuclear ribosomal cistron (Feliner and Rosselló 2007). Various reasons such as recent hybridization, lineage sorting, recombination among copies, high mutation rate, and pseudogene formation (nonfunctional paralogous) are considered to be the reasons for such variations (Song et al. 2008). Additionally nonfunctional copies of plastidal or mitochondrial pseudogenes, which have been sometimes transferred to the nucleus, have been detected in various eukaryotic organisms, too. For DNA barcoding as a practical molecular method to identify species only orthologous DNA sequences can be used. Consequently cloning of PCR products is sometimes inevitable (Bailey et al. 2003; Buckler et al. 1997).

2.9 Perspectives for Authentication Using DNA Barcoding

The principle of DNA barcoding is that a standardized, fast-evolving short segment of the genome from chloroplast, mitochondrial, or nuclear DNA can be used to quickly and easily identify any organism. For the process of identification and authentication, different aspects of classical taxonomy, laboratory practice, and data management must be considered.

The first step in authentication of medicinal plants is sampling of the correct taxon and determination using identification keys as provided in the Flora of China. The designated species should match the description of the “type” specimen in all relevant diagnostic features. Confirmation of the determined taxon by a plant taxonomist/systematist or expert in the field of pharmacognosy is favorable. The researcher should also give detailed information for all specimens that are used for barcoding, such as latin taxon name, collector, collection date and number, locality, geospatial coordinates (GPS data), as well as information on infraspecific variants

(variety, chemotype, hybrid) in case of cultivated plant material. These data combined with high resolution images should be deposited in public databases (e.g., BOLD, MMDBD). Special emphasis should be placed on unambiguous association of DNA barcodes with individual specimens and corresponding data records. Thus the specimens that are used for DNA barcoding need to be preserved individually, each with a unique identifying number, or sample ID. Consequently deposition of voucher specimens in local herbaria or natural history collections is important for comparison and verification (see sampling instructions BOLD/CCDB).

For barcoding studies it is also essential to reconstruct molecular phylogenies (using software e.g., PAUP, MrBayes, etc.) including a maximum of species of a taxonomic group. There are many genera of medicinal plants in China which exhibit an extraordinary diversity as *Astragalus* (401 species), *Aconitum* (211 species), *Artemisia* (186 species), *Berberis* (215 species), *Clematis* (147 species), *Polygonum* (113 species), or *Salix* (275 species) and phylogenetic information is limited or missing. Consequently in these genera misidentification is very common. In this respect the knowledge of closely related taxa and sister species relationships, the primary candidates that show low interspecific divergence and often share haplotypes, is of main interest. It is conspicuous that DNA barcoding studies generally neglect or underestimate intraspecific genetic variability and variation in the barcode region is not considered (Moritz and Cicero 2004). Thus phylogenetic reconstructions using comprehensively sampled groups are fundamental to explore intraspecific variation and interspecific divergence. For this purpose adequate sampling of several individuals across the geographic range of a species is required. There is also need to examine groups in more detail with frequent hybridization, recent radiations, introgressions, or incomplete lineage sorting (Li et al. 2011b; China Plant BOL Group).

Concerning laboratory practice and improvements, a cost-effective long-term storage for plant tissue samples (e.g., fresh tissue storage in liquid nitrogen or silica gel desiccation) has to be tested and used for DNA preservation. Furthermore DNA extraction from (old) herbarium specimens, increase of sequencing success, particularly from samples containing long mononucleotide repeats, is required. With application of new polymerases with higher fidelity and processivity or using mixtures that include repair enzymes amplification of degraded DNA can be overcome (Mitchell et al. 2005; Hajibabaei et al. 2005). Additionally development and optimization of DNA barcoding primers for a certain taxonomic group or highly degraded DNAs will greatly improve PCR success and multiplex PCR reactions that can routinely amplify barcode markers will significantly reduce laboratory costs. To add more reliability in the identification of species and to complement the barcoding regions more suitable, nuclear barcode loci (low-copy nuclear regions as ADH, waxy, leafy) must be developed. Future barcoding activities should be focused on the discriminating power of standard markers (*rbcL*, *matK*, ITS, *trnH-psbA*) in species-rich angiosperm families as Asteraceae, Fabaceae, Rosaceae, Ranunculaceae, Lamiaceae, and Apiaceae that contain many important Chinese medicinal plants. Next-generation sequencing technologies will

provide exciting new opportunities for barcoding of multiple samples or even entire plastid genomes that can be used as the ‘next-generation plant barcode’ (Chen et al. 2010; Sucher et al. 2012; Hollingsworth et al. 2011).

In the field of data management, sophisticated tools are required to automatically query segments of individual sequences, to check for different affinities or incongruence within markers, to decode the peaks in over-laid traces of sequencing chromatograms, to check for editing errors or pseudogenes, and to detect microinversions. Furthermore there is demand for improved bioinformatic search routines because sequence identification methods that use local pairwise alignments (e.g., BLAST) are unable to accurately differentiate between highly similar sequences and are not designed to cope with hierarchic phylogenetic relationships or within taxon variability (Little 2011).

Comprehensive databases containing voucher specimens, macro and microscopic data, chemical profiling, and DNA barcoding information would clearly be beneficial for the authentication of medicinal plants and for providing consumers with a safe product. The molecular detection technologies therefore undoubtedly contribute to the research and development of herbal drugs.

2.10 Guidelines for Authentication of Chinese Medicinal Plants

- For ensuring the safety, efficacy, and quality of traditional Chinese herbal medicines classical botanical methodologies have to be applied including collection and conservation of the botanical material (herbarization of plants and silica gel desiccation of leaves for DNA extraction). It is important to supply representative portions of the plant for correct identification, particularly flowering parts, fruits, and seeds. Large specimens should be dissected and mounted to show the main distinguishing characteristics.
- Additionally proper *documentation* is required: who collected the plant, date of collection, collection site (preferably by GPS coordinates), a unique collection number or code, and habitat information (surrounding vegetation, soil, exposition, and landscape). Supplementary photographs of the growing plant in its habitat should be provided. It is advisable to collect several sets of the same specimen so that, following correct identification, one can retain a specimen for later reference. The plant should be determined by an expert (taxonomist) and given a legitimate *Latin binominal name* (including genus, species, subspecies/variety, and author) according to the guidelines and rules of the International *Code of Botanical Nomenclature* (ICBN) (McNeill et al. 2006).
- Online *taxonomic sources* such as the Flora of China Checklist, The Plant List (TPL), The International Plant Names Index (IPNI), World Checklist of selected plant families or Medicinal Plant Names Index (MPNI) offer access to accepted names. If available, the local and/or English common names and relevant information, such as the cultivar name, ecotype, chemotype should also be provided.

- From the collected and properly dried plant material (one or several representative individuals) a *voucher specimen* should be prepared by mounting the plant(s) on a herbarium sheet in a format suitable for conserving the morphological key characters (e.g., flowers, fruits) of the specimen. This voucher which serves as a permanent record and reference of an individual plant species in time and space must be labeled with scientific name and detailed collection data (as mentioned above) and deposited in a registered public herbarium, museum, or repository of a certified research institute. It is appropriate to prepare two or more duplicate voucher specimens (“back ups”) that can be sent to taxonomic experts (plant taxonomists) anywhere in the world for confirmation of the species identification. Detailed information on procedures to properly collect, press, and prepare voucher specimens are available (Hildreth et al. 2007; Smillie and Khan 2010; Eisenmann et al. 2012).
- Seeds or other propagation materials should be specified, and all necessary information relating to the identity, quality, and performance of their products, as well as their breeding history should be provided. For the identification of plants grown from seed material it is essential to prepare a voucher specimen. Due to the need for herbal drugs of consistent quality and reliable supply, methods for commercial field cultivation and post-harvest processing should be developed to guarantee high standards with regard to the required bioactive constituents. To achieve genetically improved or new cultivars breeding programs should be developed.
- For *chemical standardization* it is useful to collect bulk material from the population of the reference sample to reduce potential collecting errors. These plants should be harvested during the appropriate season or time period to ensure the best possible quality of source materials. Collection practices should ensure the long-term survival of wild populations and their associated habitats. Medicinal plants/herbal drugs from species that are listed as endangered (CITES, Convention on International Trade in Endangered Species of Wild Fauna and Flora) must not be collected unless the relevant competent authority has given its authorization.
- Using DNA information for species identification, DNA sequences have to be generated to detect differences in nucleotide positions that are a prerequisite for sequence similarity search in databases. This function allows the user to conduct homology searches between the sequence of interest and the data in the public sequence databases (GenBank; MMDBD; BOLD). To make identification reliable, it is necessary to have complete information of all species of a plant genus. This can only be achieved by conducting phylogenetic reconstructions (using a combination of universal DNA barcoding markers of the nuclear and chloroplast genome) including a comprehensive sampling of species from genera of frequently used TCM plants. Only this molecular phylogenetic approach provides evidence for the identification of genetically closely related species or cryptic taxa. All analyses should include voucher numbers of samples, primer sequences, and PCR conditions for generating barcode sequences. All sequences

should be deposited in *DNA databases (GenBank; MMDBD; BOLD)* for future data retrieval, similarity search, and species identification.

- The focus of DNA barcoding analyses should be on these medicinal plant species which are worldwide and the most frequently used official Chinese plant drugs. Special attention should be paid to those medicinal plants that are frequently substituted or adulterated with other species that are morphologically and/or phytochemically indistinguishable. Based on available sequence information (barcodes), other diagnostic molecular identification tools (PCR-RFLP, ARMS, SCAR, etc.) with high discrimination power should be developed for easy, cheap, and reliable identification of species, varieties, and cultivars.
- To examine correlations between *DNA barcoding* and *chemical profiling*, chromatographic techniques (e.g., TLC, HPLC, GC as well as IR, MS, and NMR spectrometry) should be applied in an integrated strategy using the same particular plant source.
- In view of the amount of DNA barcodes deposited meanwhile in databases, it is highly recommended that molecular authentication should be implemented as a standard method for identification of Chinese herbal medicinal materials in future editions of the Pharmacopoeia of the People's Republic of China.
- Beyond that *interdisciplinary workshops/conferences* are necessary to exchange ideas, to highlight perspectives of future research, to stimulate new work in the field of TCM, and to enhance the dialogue between scientists from plant taxonomy, molecular biology, agriculture, phytochemistry, pharmacognosy, and medicine. In addition a *global information network on TCM research* is required which integrates available databases. A modernization of Traditional Chinese Medicine has become necessary and urgent to increase the worldwide acceptance.

2.11 Future Developments

New innovative automated assays and specific tools for DNA analysis are emerging and will contribute to the next generation of technologies. These are minisequencing (Pastinen et al. 1996; Cai et al. 2000), nanoscale DNA sequencing (Pastinen et al. 1996), Microsphere-based suspension arrays (Lowe 2000), and next-Generation Sequencing (Mardis 2008; Lerner and Fleischer 2010). Further extremely promising developments are the nanopore technology for identification of DNA bases with very high confidence and the arrayed primer extension reaction (APEX) which is an enzymatic genotyping method to analyze hundreds to thousands of variations in the genome simultaneously in a single multiplexed reaction (Fortina et al. 2005; Pirrung et al. 2000). Another upcoming method for large-scale multiplex analysis of nucleic acid sequences is the oligonucleotide ligation assay (OLA), which can be applied for the detection of known single nucleotide polymorphisms (SNPs) and allelic discrimination in highly polymorphic genes (Kurg et al. 2000). These novel approaches to DNA sequencing promise complete genomic analysis and have a

high multiplexing capacity and great potentials for genotyping and future taxon identification (Grossmann et al. 1994).

New high-throughput sequencing (HTS) technologies enable application of new molecular approaches. The recent introduction of massive parallel sequencing technology producing millions of DNA sequence reads (in total 0.5–60 giga base pairs) in a single run has revolutionized genomic research in biology and medicine. These so-called next-generation sequencing platforms, such as Roche/454, Illumina/Solexa, Helicos, and ABI/Solid system can sequence DNA faster and at much lower costs in comparison to the conventional 96-capillary system of Sanger sequencing.

Next-generation sequencing techniques (NGS) can be used to address new and long-standing questions previously inhibited by technological and financial limitations (Kircher and Kelsko 2010). Although no “third-generation” platform has been made commercially available yet, several companies have prototype technologies in active development.

References

- Agarwal M (2008) Advances in molecular marker techniques and their applications in plant sciences. *Plant Cell Rep* 27:617–631
- Akkaya MS, Shoemaker RC, Specht JE, Bhagwat AA, Cregan PB (1995) Integration of simple sequence repeat and markers into a soybean linkage map. *Crop Sci* 35:1439–1445
- Albach DC, Li HQ, Zhao N, Jensen SR (2007) Molecular systematics and phytochemistry of *Rehmannia* (Scrophulariaceae). *Biochem Syst Ecol* 35:293–300
- Albani MC, Battey NH, Wilkinson MJ (2004) The development of ISSR derived SCAR markers around the seasonal flowering locus (SFL) in *Fragaria vesca*. *Theor Appl Genet* 109:571–579
- Álvarez I, Wendel JF (2003) Ribosomal ITS sequences and plant phylogenetic inference. *Mol Phylogenet Evol* 29:417–434
- Arya M, Shergill IS, Williamson M, Gommersall L, Arya N, Patel HR (2005) Basic principles of real-time quantitative PCR. *Expert Rev Mol Diagn* 5(2):209–219
- Asahina H, Shinozaki J, Masuda K et al (2010) Identification of medicinal *Dendrobium* species by phylogenetic analyses using matK and rbcL sequences. *J Nat Med* 64(2):133–138
- Bailey CD, Carr TG, Harris SA, Hughes CE (2003) Characterization of angiosperm nrDNA polymorphism, paralogy, and pseudogenes. *Mol Phylogenet Evol* 29:435–455
- Baldwin BG, Sanderson MJ, Porter JM, Wojciechowski MF, Campbell CS, Donoghue MJ (1995) The ITS region of nuclear ribosomal DNA: a valuable source of evidence on angiosperm phylogeny. *Ann Missouri Bot Gard* 82:247–277
- Balunas MJ, Kinghorn AD (2005) Drug discovery from medicinal plants [J]. *Life Sci* 78 (5):431–441
- Barthelson RA (2009) Protocols for in vitro cultures and secondary metabolite analysis of aromatic and medicinal plants. Identification of medicinal plants and plant sequences: multiplexed MLPA assay. *Methods Mol Biol* 547(3):277–288
- Bauer T, Weller P, Hammes WP, Hertel C (2003) The effect of processing parameters on DNA degradation in food. *Eur Food Res Technol* 217:338–343
- Bhattacharya E, Dandin SB et al (2005) Single primer amplification reaction methods reveal exotic and indigenous mulberry varieties are similarly diverse. *J Biosci* 30:669–677
- Bleas M, De Grandis S, Lee H, Trevors J (1998) Amplified fragment length polymorphism (AFLP): a review of the procedure and its applications. *J Ind Microbiol Biotechnol* 21:99–114

- Boqian Y, Jing W, Guopei C, Ting W (2009) Isolation and characterization of polymorphic microsatellite loci in a traditional Chinese medicinal plant, *Schisandra sphenanthera*. *Cons Genetics* 10:615–617
- Borne B, Goraguer F, Joly G, Branchard M (2002) Genetic diversity in European and Argentinian cultivated potatoes (*Solanum tuberosum* subsp. *tuberosum*) detected by inter-simple sequence repeats (ISSRS). *Genome* 45:481–484
- Bornet B, Branchard M (2004) Use of ISSR fingerprints to detect microsatellites and genetic diversity in several related Brassica taxa and *Arabidopsis thaliana*. *Hereditas* 140:245–248
- Buckler ES, Ippolito SA, Holtsford TP (1997) The evolution of ribosomal DNA: divergent paralogues and phylogenetic implications. *Genetics* 145:111–125
- But PPH (1994) Herbal poisoning caused by adulterants or erroneous substitutes. *J Trop Med Hyg* 97:371–374
- But PP, Tomlinson B, Cheung KO, Yong SP, Szeto ML, Lee CK (1996) Adulterants of herbal products can cause poisoning. *Br Med J* 313:117
- Caetano-Anollés G, Gresshoff PM (1996) Generation of sequence signatures from DNA amplification fingerprints with mini-hairpin and microsatellite primers. *Biotechniques* 20:1044–1056
- Caetano-Anollés G, Bassam BJ, Gresshoff PM (1991a) DNA amplification fingerprinting: a strategy for genome analysis. *Plant Mol Biol Rep* 9:294–307
- Caetano-Anollés G, Bassam BJ, Gresshoff PM (1991b) DNA amplification using very short arbitrary oligonucleotide primers. *Biotechnology* 9:553–557
- Cai ZH, Li P, Dong TT, Tsim KW (1999) Molecular diversity of 5S-rRNA spacer domain in *Fritillaria* species revealed by PCR analysis. *Planta Med* 65:360–364
- Cai H, White PS, Torney D et al (2000) Flow cytometry-based minisequencing: a new platform for high-throughput single-nucleotide polymorphism scoring. *Genomics* 66:135–143
- Cao H, But PPH, Shaw PC (1996a) Authentication of the Chinese drug “ku-di-dan” (herba elephantopi) and its substitutes using random-primed polymerase chain reaction (PCR). *Acta Pharm Sin* 31:543–553
- Cao H, But PPH, Shaw PC (1996b) Identification of herba taraxaci and its adulterants in Hong Kong market by DNA fingerprinting with random primed PCR. *China J Chin Mater Med* 22:197–200
- Carles M, Cheung MK, Moganti S, Dong TT, Tsim KW, Ip NY et al (2005) A DNA microarray for the authentication of toxic traditional Chinese medicinal plants. *Planta Med* 71:580–584
- CBOL Plant Working Group (2009) A DNA barcode for land plants. *Proc Natl Acad Sci* 106 (31):12794–12797
- Chakrabarty D, Yu KW, Paek KY (2003) Detection of DNA methylation changes during somatic embryogenesis of Siberian ginseng (*Eleutherococcus senticosus*). *Plant Sci* 165:61–68
- Chan TYK (1997) Monitoring the safety of herbal medicine. *Drug Saf* 17:209–215
- Chan K (2003) Some aspects of toxic contaminants in herbal medicines. *Chemosphere* 52:1361–1371
- Chan TY, Critchley JA (1996) Usage and adverse effects of Chinese herbal medicines. *Hum Exp Toxicol* 15:5–12
- Chan TYK, Chan JCN, Tomlinson B, Critchley JA (1993) Chinese herbal medicines revisited: a Hong Kong perspective. *Lancet* 342:18–25
- Chase MW, Salamin N, Wilkinson M, Dunwell JM, Kesanakurthi RO et al (2005) Land plants and DNA barcodes: short-term and long-term goals. *Philos Trans R Soc Lond B* 360:1889–1895
- Chase MW, Cowan RS, Hollingsworth PM, van den Berg C, Madrinan S et al (2007) A proposal for a standardised protocol to barcode all land plants. *Taxon* 56:295–299
- Chatti K, Saddoud O, Salhi Hannachi A, Mars M, Marrakchi M, Trifi M (2007) Analysis of genetic diversity and relationships in a Tunisian Fig (*Ficus carica*) germplasm collection by random amplified microsatellite polymorphisms. *J Integr Plant Biol* 49:386–391
- Chawla HS (2002) Introduction to plant biotechnology, 2nd rev. edn. Science Publishers, Enfield, NH

- Chen KT, Su YC, Lin JG, Hsin LH, Su YP, Su CH, Li SY, Cheng JH, Mao SJ (2001) Identification of *Atractylodes* plants in Chinese herbs and formulations by random amplified polymorphic DNA. *Acta Pharm Sin* 22:493–497
- Chen J, Chen L, An Z, Shi S, Zhan Y (2002) Non-technical causes of fakes existing in Chinese medicinal material markets. *Zhongyaocai* 25:516–519
- Chen F, Chan HYE et al (2008) Authentication of *Saussurea lappa* and endangered medicinal material, by ITS DNA and 5S rRNA sequencing. *Planta Med* 74:889–892
- Chen S, Yao H, Han J, Liu C, Song J, Shi L, Zhu Y, Ma X, Gao T, Pang X, Luo K, Li Y, Li X, Jia X, Lin Y, Leon C (2010) Validation of the ITS2 region as a novel DNA barcode for identifying medicinal plant species. *PLOS ONE* 5(1):e86137
- Cheng KT, Chang HC, Su CH, Hsu FL (1997) Identification of dried rhizomes of *Coptis* species using random amplified polymorphic DNA. *Bot Bull Acad Sin* 38:241–244
- Chen JH, Zhou Y, Yang YP (2011) Isolation and characterization of 20 new microsatellite loci in *Coriaria nepalensis* (Coriariaceae). *Am J Bot* 98(6):e141–e143
- Cheng KT, Fu LC et al (1998) Identification of *Anoectochilus formosanus* and *Anoectochilus koshunensis* species with RAPD markers. *Planta Med* 64:46–49
- Cheng KT, Su B, Chen CT, Lin CC (2000) RAPD analysis of *Astragalus* medicines marketed in Taiwan. *Am J Chin Med* 28:273–278
- Cheng JL, Huang LQ, Shao AJ, Lin SF (2002) RAPD analysis on different varieties of *Rehmannia glutinosa*. *Zhongguo Zhongyao Zazhi* 27:505–508
- Cheung KS, Kwan HS, But PPH, Shaw PC (1994) Pharmacognostical identification of American and Oriental ginseng roots by genomic fingerprinting using arbitrarily primed polymerase chain reaction (AC-PCR). *J Ethnopharmacol* 42:67–69
- Cheung TP, Xue C, Leung K et al (2006) Aristolochic acids detected in some raw Chinese medicinal herbs and manufactured herbal products – a consequence of inappropriate nomenclature and imprecise labelling? *Clin Toxicol* 44:371
- Chiou SJ, Yen JH et al (2007) Authentication of medicinal herbs using PCR-Amplified ITS2 with specific primers. *Planta Med* 73:1421–1426
- Choi YE, Ahn CH, Kim BB, Yoon ES (2008) Development of species specific AFLP-derived SCAR marker for authentication of *Panax japonicus* C. A. MEYER. *Biol Pharm Bull* 31:135–138
- Chun S, Jian-He W, Shi-Lin C, Huai-Qiong C, Cheng-Min Y (2009) Development of genomic SSR and potential EST-SSR markers in *Bupleurum chinense* DC. *Afr J Biotechnol* 8:6233–6240
- Chung SM, Staub JE (2003) The development and evaluation of consensus chloroplast primer pairs that possess highly variable sequence regions in a diverse array of plant taxa. *Theor Appl Genet* 107:757–776
- Clark CM, Wentworth TM, Malley DMO (2000) Genetic discontinuity revealed by chloroplast microsatellites in eastern North American *Abies* (Pinaceae). *Am J Bot* 87:774–778
- Cole CT, Kuchenreuther MA (2001) Molecular markers reveal little genetic differentiation among *Aconitum noveboracense* and *A. columbianum* (Ranunculaceae) populations. *Am J Bot* 88:337–347
- Cole IB, Saxena PK, Murch SJ (2007) Medicinal biotechnology in the genus *Scutellaria*. *In Vitro Cell Dev Biol Plant* 43(4):318–327
- Cowan RS, Chase MW, Kress WJ, Savolainen V (2006) 300,000 species to identify: problems, progress, and prospects in DNA barcoding of land plants. *Taxon* 55:611–616
- Cui XM, Lo CK, Yip KL, Dong TT, Tsim KW (2003) Authentication of *Panax notoginseng* by 5S-rRNA spacer domain and random amplified polymorphic DNA (RAPD) analysis. *Planta Med* 69:584–586
- Dasmahapatra KK, Mallet J (2006) DNA barcodes: recent successes and future prospects. *Heredity* 97:254–255
- Datwyler SL, Weiblen GD (2006) Genetic variation in hemp and marijuana (*Cannabis sativa* L.) according to amplified fragment length polymorphisms. *J Forensic Sci* 51:371–375

- De Groot GA, During HJ, Maas JW, Schneider H, Vogel JC et al (2011) Use of *rbcl* and *trnL-F* as a two-locus DNA barcode for identification of NW European ferns: an ecological perspective. *PLoS One* 6(1):e16371
- Debelle FD, Vanherweghem JL, Nortier JL (2008) Aristolochic acid nephropathy: a worldwide problem. *Kidney Int* 74:158–169
- Debouck C, Goodfellow PN (1999) DNA microarrays in drug discovery and development. *Nat Genet* 21:48–50
- Desmarais E, Lanneluc I, Lagnel J (1998) Direct amplification of length polymorphisms (DALP), or how to get and characterize new genetic markers in many species. *Nucleic Acids Res* 26:1458–1465
- Devaiah KM, Venkatasubramanian P (2008a) Development of SCAR marker for authentication of *Pueraria tuberosa* (Roxb. ex. Willd.) DC. *Curr Sci* 94:1306–1308
- Devaiah KM, Venkatasubramanian P (2008b) Genetic characterization and authentication of *Embelia ribes* using RAPD-PCR and SCAR marker. *Planta Med* 74:194–196
- Diao Y, Lin XM et al (2009) Authentication of *Panax ginseng* from its adulterants by PCR-RFLP and ARMS. *Planta Med* 75:557–560
- Dictionary of Chinese Materia Medica (2006), 2nd edn. Shanghai Scientific & Technical, Shanghai
- Ding T (2002) The botany database: seed plants of China (SPCH), and its applications. Geoscience and Remote Sensing Symposium, 2002. IGARSS '02. 2002 I.E. International
- Ding X, Xu L, Wang Z, Zhou K, Xu H, Wang Y (2002) Authentication of stems of *Dendrobium officinale* by rDNA ITS region sequences. *Planta Med* 68:191–192
- Ding G, Zhang DZ et al (2008) SNP, ARMS and SSH authentication of medicinal *Dendrobium officinale* Kimura et Migo and application for identification of Fengdou drugs. *Biol Pharm Bull* 31:553–557
- Dnyaneshwar W, Preeti C, Kalpana J, Bhushan P (2006) Development and application of RAPD-SCAR marker for identification of *Phyllanthus emblica* L. *Biol Pharm Bull* 29:2313–2316
- Do KR, Hwang WJ, Lyu YS, An NH, Kim HM (2001) Molecular authentication of *Panax ginseng* species by RAPD analysis and PCR-RFLP. *Biol Pharm Bull* 24:872–875
- Dong CH, Yao YJ (2010) On the reliability of fungal materials used in studies on *Ophiocordyceps sinensis*. *J Ind Microbiol Biotechnol*. doi:10.1007
- Dong TT, Ma XQ, Clarke C, Song ZH, Ji ZN, Lo CK et al (2003) Phylogeny of *Astragalus* in China: molecular evidence from the DNA sequences of 5S rRNA spacer, ITS, and 18S rRNA. *J Agric Food Chem* 51:6709–6714
- Drouin G, Daoud H, Xia J (2008) Relative rates of synonymous substitutions in the mitochondrial, chloroplast and nuclear genomes of seed plants. *Mol Phylogenet Evol* 49(3):827–831
- Ernst E (2004) Risks of herbal medicinal products. *Pharmacoepidem. Drug Saf.* 13(11):767–71
- Eisenmann SW, Struwe L, Tucker AO (2012) Voucher specimens are essential for documenting source material used in medicinal plant investigations. *J Med Active Plants* 1(1):30–43
- Erickson DL, Spouge J, Resch A, Weigt LA, Kress WJ (2008) DNA barcoding in land plants: developing standards to quantify and maximize success. *Taxon* 57:1304–1316
- Fan XX, Shen L, Zhang X, Chen XY, Fu CX (2004) Assessing genetic diversity of *Ginkgo biloba* L. (*Ginkgoaceae*) populations from China by RAPD markers. *Biochem Genet* 42:269–278
- Fan WJ, Luo YM et al (2009) Development of microsatellite markers in *Dendrobium fimbriatum* Hook, an endangered Chinese endemic herb. *Mol Ecol Res* 9:373–375
- Farnsworth NR, Soejarto DD (1991) Global importance of medicinal plants. In: Akerele O, Heywood V, Syngé H (eds) *The conservation of medicinal plants*. Cambridge University Press, Cambridge, pp 25–51
- Fazekas AJ et al (2008) Multiple multilocus DNA barcodes from the plastid genome discriminate plant species equally well. *PLoS One* 3(7):e2802
- Feliner GN, Rosselló JA (2007) Better the devil you know? Guidelines for insightful utilization of nrDNA ITS in species level evolutionary studies in plants. *Mol Phylogenet Evol* 44:911–919

- Ferri G, Alù M, Corradini B, Licata M, Beduschi G (2009) Species identification through DNA “barcodes”. *Genet Test Mol Biomarkers* 13:421–426
- Fodor SP, Read JL, Pirrung MC, Stryer L, Lu AT, Solas D (1991) Light-directed, spatially addressable parallel chemical synthesis. *Science* 251:767–773
- Fleischmann A, Heubl G (2009) Overcoming DNA extraction problems from carnivorous plants. *Anal Jardín Bot Madrid* 66(2):209–215
- Fodor SP, Rava RP, Huang XC, Pease AC, Holmes CP, Adams CL (1993) Multiplexed biochemical assays with biological chips. *Nature* 364:555–556
- Ford C, Ayres KL, Toomey N, Haider N et al (2009) Selection of candidate coding DNA barcoding regions for use on land plants. *Bot J Linn Soc* 159(1):1–11
- Fortina P, Kricka LJ, Surrey S, Grodzinski P (2005) Nanobiotechnology: the promise and reality of new approaches to molecular recognition. *Trends Biotechnol* 23:168–173
- Friar EA (2005) Isolation of DNA from plants with large amounts of secondary metabolites. *Methods Enzymol* 395:3–14
- Fu RZ, Wang J, Zhang YB, Wang ZT, But PP, Li N, Shaw PC (1999) Differentiation of medicinal Codonopsis species from adulterants by polymerase chain reaction-restriction fragment length polymorphism. *Planta Med* 65:648–650
- Fugh-Berman A (2000) Herb–drug interactions. *Lancet* 355:134–138
- Gao WY, Jia W, Duan HQ, Huang L, Xiao X, Xiao P, Paek KY (2002) Good agriculture practice (GAP) and sustainable resource utilization of Chinese materia medica. *Plant Biotechnol* 4(3):103–107
- Gao T, Yao H, Song JY, Liu C, Zhu YJ, Ma XY, Pang XH, Xu HX, Chen SL (2010a) Identification of medicinal plants in the family Fabaceae using a potential DNA barcode ITS2. *J Ethnopharmacol* 130:116–121
- Gao T, Yao H, Song J, Zhu Y, Liu C, Chen S (2010b) Evaluating the feasibility of using candidate DNA barcodes in discriminating species of the large Asteraceae family. *BMC Evol Biol* 10:324
- Gawel NJ, Jarret RL, Whittmore AP (1992) Restriction fragment length polymorphism (RFLP)-based phylogenetic analysis of *Musa*. *Theor Appl Genet* 84:286–290
- Ge XJ, Yu Y, Yuan YM, Huang HW, Cheng YAN (2005) Genetic diversity and geographic differentiation in endangered *Ammopiptanthus* (Leguminosae) populations in desert regions of Northwest China as revealed by ISSR analysis. *Ann Bot* 95:843–851
- Gebauer M (2004) Microarray applications: emerging technologies and perspectives. *Drug Discov Today* 9:915–917
- Gershon D (2002) Microarray technology: an array of opportunities. *Nature* 416:885–891
- Gertner E, Marshall PS, Filandrinos D, Potek AS, Smith TM (1995) Complications resulting from the use of Chinese herbal medications containing undeclared prescription drugs. *Arthritis Rheum* 38:614–617
- Goldstein DB, Schlötterer C (1999) Microsatellites: evolution and applications. Oxford University Press, Oxford, UK
- Gong W, Fu CX, Luo YP, Qiu YX (2006) Molecular identification of *Sinopodophyllum hexandrum* and *Dysosma* species using cpDNA sequences and PCR-RFLP markers. *Planta Med* 72:650–652
- Govaerts R (2001) How many species of seed plants are there? *Taxon* 50:1085–1090
- Graham-Brown R (1992) Toxicity of Chinese herbal remedies. *Lancet* 340:673–674
- Grossmann PD, Bloch W, Brinson E et al (1994) High-density multiplex detection of nucleic acid sequences: oligonucleotide ligation assay and sequence-coded separation. *Nucleic Acids Res* 22:4527–4534
- Guo BL, Wu M, Si JP, Li JS, Xiao PG (2001) Research on DNA molecular marker of *Magnolia officinalis* Rehd. et Wils. RAPD study on certified species. *Yao Xue Xue Bao* 36:386–389
- Guo Y, Tsuruga A, Yamaguchi S, Oba K, Iwai K, Sekita S et al (2006) Sequence analysis of chloroplast chlB gene of medicinal *Ephedra* species and its application to authentication of *Ephedra* herb. *Biol Pharm Bull* 29:1207–1211

- Guo HY, Wang WW, Yang N et al (2010) DNA barcoding provides distinction between *Radix Astragali* and its adulterants. *Sci China Life Sci* 53(8):992–999
- Gupta PK, Balyan HS, Sharma PC, Ramesh B (1996) Microsatellites in plants: a new class of molecular markers. *Curr Sci* 70:45–54
- Gupta AR, Patra RC, Das DK, Gupta PK, Swarup D, Saini M (2001a) Sequence characterization and polymerase chain reaction-restriction fragment length polymorphism of the mitochondrial DNA 12S rRNA gene provides a method for species identification of Indian deer. *Mitochondrial DNA* 19:394–400
- Gupta PK, Roy JK, Prasad M (2001b) Single nucleotide polymorphisms: a new paradigm for molecular marker technology and DNA polymorphism detection with emphasis on their use in plants. *Curr Sci* 80(4):524–535
- Ha WY, Yau FC, But PP, Wang J, Shaw PC (2001) Direct amplification of length polymorphism analysis differentiates *Panax ginseng* from *P. quinquefolius*. *Planta Med* 67:587–589
- Ha WY, Shaw PC, Liu J, Yau FC, Wang J (2002) Authentication of *Panax ginseng* and *Panax quinquefolius* using amplified fragment length polymorphism (AFLP) and directed amplification of minisatellite region DNA (DAMD). *J Agric Food Chem* 50:1871–1875
- Hajibabaei M, deWaard JR, Ivanova NV, Ratnasingham S, Dooh RT, Kirk SL, Mackie PM, Hebert PDN (2005) Critical factors for assembling a high volume of DNA barcodes. *Philos Trans R Soc B* 360(1462):1959–1967
- Hajibabaei M, Singer GA, Hebert PDN, Hickey DA (2007) DNA barcoding: how it complements taxonomy, molecular phylogenetics and population genetics. *Trends Genet* 23(4):167–172
- Hamilton AC (2003) Medicinal plants and conservation: issues and approaches. Paper presented to International Plants Conservation Unit World Wildlife Foundation, UK
- Hamilton AC (2004) Medicinal plants, conservation and livelihoods. *Biodivers Conserv* 13:1477–1517
- Han J, Shi L, Song J et al (2009) Testing potential DNA barcoding regions in the Labiatae medicinal plants. *Planta Med* 75:407
- Hao DC, Chen SL, Xia PG, Peng Y (2010) Authentication of medicinal plants by DNA-based markers and genomics. *Chin Herb Med* 2(4):250–261
- Hayashi K (1992) PCR-SSCP—rapid and easy detection of DNA sequence changes. *Hum Cell* 5:180–184
- Hayden MJ, Nguyen TM, Waterman A, Chalmers KJ (2008) Multiplex-ready PCR: a new method for multiplexed SSR and SNP genotyping. *BMC Genomics* 9:80
- He J, Wong KL, Shaw PC, Wang H, Li DZ (2010) Identification of the medicinal plants in *Aconitum L.* by DNA barcoding technique. *Plant Med* 76:1622–1628
- Heath DD, Iwama GK, Devlin RH (1993) PCR primed with VNTR core sequences yields species specific patterns and hypervariable probes. *Nucleic Acids Res* 21:5782–5785
- Hebert PDN, Cywinska A, Ball SL, DeWaard JR (2003a) Biological identification through DNA barcodes. *Proc R Soc Lond B* 270:313–321
- Hebert PDN, Ratnasingham S, DeWaard JR (2003b) Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proc R Soc Lond B* 270(suppl):96–99
- Hebert PDN, Stoeckle MY, Zemlak TS, Francis CM (2004) Identification of birds through DNA barcodes. *PLoS Biol* 2(10):e312
- Heubl G (2010) New aspects of DNA-based authentication of Chinese medicinal plants by molecular biological techniques. *Planta Med* 76(17):1963–1975
- Hildreth J, Hrabeta-Robinson E, Applequist W, Betz J, Miller J (2007) Standard operating procedure for the collection and preparation of voucher plant specimens for use in the nutraceutical industry. *Anal Bioanal Chem* 389:13–17
- Hollingsworth ML, Clark A, Forrest LL, Richardson JE, Pennington RT, Long D, Cowan R, Chase MW, Gaudeul M, Hollingsworth PM (2009) Selecting barcoding loci for plants: evaluation of seven candidate loci with species level sampling in three divergent groups of land plants. *Mol Ecol Resour* 9:439–457

- Hollingsworth PM, Graham SW, Little DP (2011) Choosing and using a plant DNA barcode. *PLoS One* 6(5):e19254
- Hosokawa K, Minami M, Kawahara K, Nakamura I, Shibata T (2000) Discrimination among three species of medicinal *Scutellaria* plants using RAPD markers. *Planta Med* 66:270–272
- Hosokawa K, Minami M, Nakamura I, Hishida A, Shibata T (2005) The sequences of the plastid gene *rpl16* and the *rpl16-rpl14* spacer region allow discrimination among six species of *Scutellaria*. *J Ethnopharmacol* 99:105–108
- Hosokawa K, Hishida A, Nakamura I, Shibata T (2006) The sequences of the spacer region between the *atpF* and *atpA* genes in the plastid genome allows discrimination among three varieties of medicinal *Angelica*. *Planta Med* 72:570–571
- Hsiao JIH (2007) Patent protection for Chinese herbal medicine product invention in Taiwan. *J World Intell Prop* 10(1):1–21
- Hu Y, Zhang Q, Xin H, Qin LP, Lu BR, Rahman K et al (2007) Association between chemical and genetic variation of *Vitex rotundifolia* populations from different locations in China: its implication for quality control of medicinal plants. *Biomed Chromatogr* 21:967–975
- Huang L, Wang M, Zhou C, Li N, He X, Yang B (1998) Problems and solutions in the use of RAPD to the identification of the Chinese drugs “*xi-xin*” (*Herba asari*) and its substitutes. *Yao Xue Xue Bao* 33:778–784
- Huang LQ, Cui GH, Chen ML, Feng CQ, Yang B, Wang YY, Dai RW (2002a) Study on complex system of Chinese material medica GAP fulfilling-the situation, problems, and of Chinese material medica germplasm. *China J Chin Mater Med* 27:481–483
- Huang LQ, Cui GH, Dai RW (2002b) Study on complex system of Chinese Material Medica GAP fulfilling. *China J Chin Mater Med* 27:1–3
- ICBN, International Code of Botanical Nomenclature (2006) 2006 Electronic version of the original English text. Prepared and edited by McNeill J, Barrie FR, Burdet HM, Demoulin V, Hawksworth DL, Marhold K, Nicolson DH, Prado J, Silva PC, Skog JE, Wiersema JH, and Turland NJ. Adopted by the Seventeenth International Botanical Congress, Vienna, Austria, July 2005. Last updated 09.03.2007
- Ince AG, Karaca AGM et al (2009) Development and utilization of diagnostic DAMD-PCR markers for *Capsicum* accessions. *Genet Res Crop Evol* 56:211–221
- Jayasinghe R, Kong S, Coram TE, Kaganovitch J, Xue CC, Li CG, Pang EC (2007) Construction and validation of a prototype microarray for efficient and high-throughput genotyping of angiosperms. *Plant Biotechnol J* 5(2):282–289
- Jo BH, Suh DS et al (2009) Characterization of polymorphic microsatellite loci in cultivated and wild *Panax ginseng*. *Genes Genomics* 31:119–127
- Jones CJ, Edwards KJ, Castaglione S, Winfield MO, Sala F, et al (1993) *Theor Appl Genet* 85:985–993
- Joshi P, Dhawan V (2007) Assessment of genetic fidelity of micropropagated *Swertia chirayita* plantlets by ISSR marker assay. *Biol Plantarum* 51(1):22–26
- Joshi K, Chavan OP, Warude D (2004) Molecular markers in herbal drug technology. *Curr Sci* 87(2):159–165
- Kaplan J, Chavan P, Warude D, Patwardhan B (2004) Molecular markers in herbal drug technology. *Curr Sci* 87:159–165
- Karaca M, Ince AG et al (2008) PCR-RFLP and DAMD-PCR genotyping for *Salvia* species. *J Sci Food Agric* 88:2508–2516
- Karp A, Edwards KJ (1997) DNA markers, a global overview. In: Caetano-Anollés G, Gresshoff PM (eds) *DNA markers, protocols, applications, and overviews*. Wiley-Liss, New York
- Kelly LM (1998) Phylogenetic relationships in *Asarum* (Aristolochiaceae) based on morphology and ITS sequences. *Am J Bot* 85:1454–1467
- Kersten T, Daniel C, König GM, Knöß W (2007) The potential of PCR-related methods to identify medicinal plants in herbal medicinal products. *Planta Med* 73:256
- Kim J, Chung KW (2007) Isolation of new microsatellite-containing sequences in *Acanthopanax senticosus*. *J Plant Biol* 50:557–561

- Kim J, Jo BH, Lee KL, Yoon ES, Ryu GH, Chung KW (2007a) Identification of new microsatellite markers in *Panax ginseng*. *Mol Cells* 24:60–68
- Kim OT, Bang K, Dong S, Lee JW, Kim YC, Shin YS, Hyun DY, Lee SS, Cha SW, Seong NS (2007b) Molecular authentication of ginseng cultivars by comparison of internal transcribed spacer and 5.8S rDNA sequences. *Plant Biotechnol Rep* 1:163–167
- Kircher M, Kelsko J (2010) High-throughput DNA sequencing-concepts and limitations. *Bioessays* 32(6):524–536
- Klein D (2002) Quantification using real-time PCR technology: applications and limitations. *Trends Mol Med* 8(6):257–260
- Kohjyouma M, Nakajima S, Namera A, Shimizu R, Mizukami H, Kohda H (1997) Random amplified polymorphic DNA analysis and variation of essential oil components of *Atractylodes* plants. *Biol Pharm Bull* 20:502–506
- Kojoma M, Lida O, Makino Y, Sekita S, Satake M (2002) DNA fingerprinting of *Cannabis sativa* using inter-simple sequence repeat (ISSR) amplification. *Planta Med* 68:60–63
- Komatsu KS, Zhu S et al (2001) Phylogenetic analysis based on 18S rRNA gene and matK gene sequences of *Panax vietnamensis* and five related species. *Planta Med* 67:461–465
- Kondo K, Terabayashi S et al (1996) Phylogenetic relationship of medicinally important *Cnidium officinale* and Japanese *Apiaceae* based on rbcL sequences. *J Plant Res* 109:21–27
- Korea Herbal Pharmacopoeia (2002) Ministry of Health Family Welfare of South Korea, Seoul, Korea
- Kress WJ, Erickson DL (2007) A two-locus global DNA barcode for land plants: the coding rbcL gene complements the non-coding trnH-psbA spacer region. *PLoS One* 2(6):e508
- Kress WJ, Wurdack KJ, Zimmer EA, Weigt LA, Janzen DH (2005) Use of DNA barcodes to identify flowering plants. *Proc Natl Acad Sci USA* 102:8369–8374
- Kumar J, Verma V et al (2007) Development of simple sequence repeat markers in *Cymbopogon* species. *Planta Med* 73:262–266
- Kumar P, Gupta VK, Misra AKL, Modi DR, Pandey BK (2009) Potential of molecular markers in plant biotechnology. *Plant Omics J* 2(4):141–162
- Kurane J, Shinde V, Harsulkar A (2009) Application of ISSR marker in pharmacognosy: current update. *Phcog Rev* 3(6):216–228
- Kurg A, Tönisson N, Georgiou I, Shumaker J, Tollett J, Metspalu A (2000) Arrayed primer extension: solid-phase four-color DNA resequencing and mutation detection technology. *Genet Test* 4:1–7
- Kwok PY (2001) Methods for genotyping single nucleotide polymorphisms. *Annu Rev Genom Human Genet* 2:235–258
- Lahaye R, van der Bank M, Bogarin D, Warner J, Pupulin F, Gigot G, Maurin O, Duthoit S, Barraclough TG, Savolainen V (2008a) DNA barcoding the floras of biodiversity hotspots. *Proc Natl Acad Sci USA* 105:2923–2928
- Lahaye R, Savolainen V, Duthoit S, Maurin O, van der Bank M (2008) A test of psbK-psbI and atpF-atpH as potential plant DNA barcodes using the flora of the Kruger National Park as a model system (South Africa). *Nature Proc* hdl:10101/npre.1896.1
- Lane CE, Lindstrom SC, Saunders GW (2007) A molecular assessment of Northeast Pacific *Alaria* species (*Laminariales*, *Phaeophyceae*) with reference to the utility of DNA barcoding. *Mol Phylogenet Evol* 44(2):634–648
- Lau DT, Shaw PC, Wang J, But PP (2001) Authentication of medicinal *Dendrobium* species by the internal transcribed spacer of ribosomal DNA. *Planta Med* 67:456–460
- Lee C, Wen J (2004) Phylogeny of *Panax* using chloroplast trnC-trnD intergenic region and the utility of trnC-trnD in interspecific studies of plants. *Mol Phylogenet Evol* 31:894–903
- Lee HH, Itokawa H, Kozuka M (2005) Asian herbal products: the basis for development of high-quality dietary supplements and new medicines. In: Shi J, Ho CT, Shahidi F (eds) *Asian functional foods*. CRC, Taylor & Francis Group, Boca Raton, pp 21–72

- Lee MY, Doh EJ, Park CH, Kim YH, Kim ES, Ko BS et al (2006a) Development of SCAR marker for discrimination of *Artemisia princeps* and *A. argyi* from other *Artemisia* herbs. *Biol Pharm Bull* 29:629–633
- Lee SK, Li PT, Lau DT, Yung PP, Kong RY, Fong WF (2006b) Phylogeny of medicinal *Phyllanthus* species in China based on nuclear ITS and chloroplast *atpB-rbcL* sequences and multiplex PCR detection assay analysis. *Planta Med* 72:721–726
- Lee HL, Yi DK, Kim JS, Kim KJ (2007) Development of plant DNA barcoding markers from the variable noncoding regions of chloroplast genome. Abstract presented at the second international barcode of life conference. Academia Sinica, Taipei, Taiwan. 18–20 September 2007
- Lee JH, Lee JW, Sung JS, Bang KH, Moon SG (2009) Molecular authentication of 21 Korean *Artemisia* species (Compositae) by polymerase chain reaction-restriction fragment length polymorphism based on *trnL-F* region of chloroplast DNA. *Biol Pharm Bull* 32(11):1912–1916
- Lerner HRL, Fleischer RC (2010) Prospects for the use of next-generation sequencing methods in ornithology. *BioOne* 27(1):4–15
- Levinson G, Gutman GA (1987) Slipped-strand mispairing: a major mechanism for DNA sequence evolution. *Mol Biol Evol* 4:203–221
- Li DP (2001) Progress on mechanism of KLT injection antitumour effect. *Traditi Chin Drug Res Clin Pharmacol* 2:122–124
- Li ZY (2003) Advances in plant resource conservation in China: sustainable development of plant resources in China. *Acta Bot Sin* 45:124–130
- Li Y, Ding WL (2010) Genetic diversity assessment of *Trollius* accessions in China by RAPD markers. *Biochem Genet* 48:34–43
- Li MF, Xu GJ, Xu LS, Jin RL (1986) Survey and identification of commercial example of *Shihu* (*Dendrobium*) (I). *Chin Trad Herbal Drugs* 17:33–37
- Li MF, Xu GJ, Xu LS, Jin RL (1991) Survey and identification of commercial example of *Shihu* (*Dendrobium*) (II). *Chin Trad Herbal Drugs* 22:173–178
- Li YC, Korol AB, Fahima T, Beiles A, Nevo E (2002) Microsatellites: genomic distribution, putative functions and mutational mechanisms: a review. *Mol Ecol* 11(12):2453–2465
- Li T, Wang J, Lu Z (2005) Accurate identification of closely related *Dendrobium* species with multiple species-specific gDNA probes. *J Biochem Biophys Methods* 62:111–123
- Li SP, Yang FQ, Tsim WK (2006) Quality control of *Cordyceps sinensis*, a valued traditional Chinese medicine. *J Pharm Biomed Anal* 41:1571–1584
- Li X, Ding X, Chu B, Ding G, Gu S, Qian L et al (2007) Molecular authentication of *Alisma orientale* by PCR-RFLP and ARMS. *Planta Med* 73:67–70
- Li K, Wu W, Zheng Y, Dai Y, Xiang L, Liao K (2009a) Genetic diversity of *Fritillaria* from Sichuan province based on ISSR. *Zhongguo Zhongyao Zazhi* 34:2149–2154
- Li YH, Ruan JL, Chen SL, Lu D, Zhu KX, Zhao MH, Pei HH (2009b) Study on medicinal plants of Loranthaceae resources in China. *World Sci Technol Mod Tradit Chin Med* 5:665–669
- Li CI, Chiou SJ, Tong TS, Lee CY, Lee LT, Cheng CM (2010a) Development and validation of molecular markers for characterization of *Boehmeria nivea* var. *nivea* and *Boehmeria nivea* var. *tenacissima*. *Chin Med* 5:40
- Li YH, Ruan JL, Chen SL et al (2010b) Authentication of *Taxillus chinensis* using DNA barcoding technique. *J Med Plant Res* 4(24):2706–2709
- Li CI, Shu-Jiau Chiou SJ, Tong TS, Lee CY, Lee LT, Cheng CM (2010c) Development and validation of molecular markers for characterization of *Boehmeria nivea* var. *nivea* and *Boehmeria nivea* var. *tenacissima*. *Chin Med* 5:40
- Li DZ, Liu JQ, Chen ZD, Wang H, Ge XJ, Zhou SL, Gao LM, Fu CX, Chen SL (2011a) Plant DNA barcoding in China. *J Systemat Evol* 49(3):165–168
- Li DZ, Gao LM, Li HT, Wang H, Liu JQ, Chen ZD, Zhou SL, Chen SL, Yang YB, Fu CX, Zeng CX, Yan HF, Zhu YJ, Sun YS, Chen SY, Zhao L, Wang K, Yang T, Daun GW (2011b) Comparative analysis of a large dataset indicates that internal transcribed spacer (ITS) should be incorporated into the core barcode for seed plants. *Proc Natl Acad Sci USA* 108:19641–19646

- Liang ZT, Qin MJ, Wang ZT, Huang Y, Wang NH (2002) Identification of *Bupleurum* L. plants by RAPD technology. *Zhongcaoyao* 33:1117–1119
- Liang YZ, Kieb PS, Chan K (2004) Quality control of herbal medicines. *J Chromatogr B* 812:53–70
- Lim W, Mudge KW et al (2007) Utilization of RAPD markers to assess genetic diversity of wild populations of North American ginseng (*Panax quinquefolium*). *Planta Med* 73:71–76
- Lin J, Zhou X, Gao S, Wu W, Liu X, Sun X et al (2006) Authentication of *Pinellia ternata* and its adulterants based on PCR with specific primers. *Planta Med* 72:844–847
- Lin WY, Chen LR, Lin TY (2008) Rapid authentication of *Bupleurum* species using an array of immobilized sequence-specific oligonucleotide probes. *Planta Med* 74:464–469
- Litt M, Luty JA (1989) A hypervariable microsatellite revealed by in vitro amplification of a dinucleotide repeat within the cardiac muscle actin gene. *Am J Hum Genet* 44:397–401
- Little DP (2011) DNA barcode sequence identification incorporating taxonomic hierarchy and within taxon variability. *PLoS One* 6(8):e20552
- Liu CX, Yaniv Z (2005) Research and development of new drugs originating from Chinese plants. In: Yaniv Z, Bachrach U (eds) *Handbook of medicinal plants*. Food Products–Haworth, pp 61–96
- Liu Y, Cao H, Wang X (2001a) Application of gene technology in quality control of Chinese drugs (I) – identification of *Pinellia ternata* species from Yuncheng, Shandong using DNA sequencing. *Chin J Pharm Anal* 21:423–426
- Liu Y, He B, Cao H (2001b) Application of gene technology in quality control of Chinese drugs (II) – identification of Chinese yam (*Dioscorea polystachia* rhizome) using DNA sequencing. *Chin J Tradit Herb Drugs* 32:113–117
- Liu CS, Bai GB, Yan YN (2005) Studies on the botanical sources and DNA molecular identification of *Herba Asari* based on ITS sequence. *Zhongguo Zhongyao Zazhi* 30:329–332
- Liu Y, Yan HF, Cao T, Ge XJ (2010a) Evaluation of 10 plant barcodes in Bryophyta (Mosses). *J Systemat Evol* 48(1):36–46
- Liu Z, Chen K, Luo K, Pan H, Chen S (2010b) DNA barcoding in medicinal plants Caprifoliaceae. *Zhongguo Zhongyao Zazhi* 35(19):2527–2532
- Liu C, Yu H, Chen L (2011) Framework for sustainable use of medicinal plants in China. *Plant Divers Resour* 33(1):65–68
- Loh JP, Kiew R, Kee A, Gan LH, Gan YY (1999) Amplified fragment length polymorphism (AFLP) provides molecular markers for the identification of *Caladium bicolor* cultivars. *Ann Bot (Lond)* 84:155–161
- Long C, Kakiuchi N, Takahashi A, Komatsu K, Cai S, Mikage M (2004) Phylogenetic analysis of the DNA sequence of the non-coding region of nuclear ribosomal DNA and chloroplast of *Ephedra* plants in China. *Planta Med* 70:1080–1084
- Lopez-Pujol J, Zhao AM (2004) China: a rich flora needed of urgent conservation. *Orsis* 19:49–89
- Lopez-Pujol J, Zhang FM, Song GE (2006) Plant biodiversity in China: richly varied, endangered, and in need of conservation. *Biodivers Conserv* 15:3983–4026
- Lou SK, Wong KL, Li M et al (2010) An integrated web medicinal materials DNA database: MMDBD (Medicinal Materials DNA Barcode Database). *BMC Genomics* 11:402
- Lowe CR (2000) Nanobiotechnology: the fabrication and applications of chemical and biological nanostructures. *Curr Opin Struct Biol* 10:428–434
- Lu Y, Li CS, Dong Q (2008) Chinese herb related molecules of cancer-cell-apoptosis: a minireview of progress between Kanglaite injection and related genes. *J Exp Clin Cancer Res* 27:31
- Lu KT, Lee HC, Liu FS et al (2009) Discriminating *Astragali Radix* from *Hedysarum Radix* in Chinese medicine preparations using nested PCR and DNA sequencing methods. *J Food Drug Anal* 17:380–385
- Lu C, Zhang W, Peng X, Gu G, Chen M, Tang Z (2010a) Development of randomly amplified polymorphic DNA-sequence characterized amplified region marker for identification of *Apocynum venetum* LINN. from *A. pictum* SCHRENK. *Biol Pharm Bull* 33:522–526

- Lu KT, Lee HC, Liu FS et al (2010b) Identification of Ginseng Radix in Chinese medicine preparations by nested PCR-DNA sequencing method and nested PCR-restriction fragment length polymorphism. *J Food Drug Anal* 18:58–63
- Lum MR, Potter E, Dang T, Heber D, Hardy M, Hirsch AM (2005) Identification of botanicals and potential contaminants through RFLP and sequencing. *Planta Med* 71:841–846
- Luo JP, Cao H, Liu YP (2002) DNA sequencing and molecular identification of Patchouli and its substitute wrinkled gianthyssop. *Yao Xue Xue Bao* 37:739–742
- Luo YM, Zhang WM, Ding XY, Shen J, Bao SL, Chu BH et al (2006a) SNP marker and allele-specific diagnostic PCR for authenticating herbs of Perilla. *Acta Pharm Sin* 41:840–845
- Luo YM, Zhang WM, Ding XY, Shen J, Bao SL, Chu BH, Mao SG (2006b) SNP marker and allele-specific diagnostic PCR for authenticating herbs of Perilla. *Yao Xue Xue Bao* 41(9):840–845
- Luo K, Chen SL, Chen KL, Song JY, Yao H, Ma XY, Zhu YJ, Pang XH, Yu H, Li XW, Liu Z (2010) Assessment of candidate plant DNA barcodes using the Rutaceae family. *Sci China Life Sci* 53:701–708
- Luo A, Zhang A, Ho SY, Xu W, Zhang Y, Shi W, Cameron SL, Zhu C (2011) Potential efficacy of mitochondrial genes for animal DNA barcoding: a case study using eutherian mammals. *BMC Genomics* 12:84
- Ma GX, Xu GJ, Xu LS, Li MF (1995) Survey and identification of commercial example of Shihu (*Dendrobium*) (III). *Chin Trad Herbal Drugs* 26:370–372
- Ma XQ, Duan JA, Zhu DY, Dong TT, Tsim KW (2000) Species identification of Radix Astragali (Huangqi) by DNA sequence of its 5S-rRNA spacer domain. *Phytochemistry* 54:363–368
- Ma XQ, Zhu DY, Li SP, Dong TT, Tsim KW (2001) Authentic identification of stigma Croci (stigma of *Crocus sativus*) from its adulterants by molecular genetic analysis. *Planta Med* 67:183–186
- Ma YS, Yu H, Li YY, Yan H, Cheng X (2008) a study of genetic structure of *Stephania yunnanensis* (Menispermaceae) by DALP. *Biochem Genet* 46:227–240
- Ma XY, Xie CX, Liu C et al (2010) Species identification of medicinal pteridophytes by a DNA barcode marker, the chloroplast psbA-trnH intergenic region. *Biol Pharm Bull* 33(11):1919–1924
- Maeda M, Uryu N, Murayama N, Ishii H, Ota M, Tsuji K, Inoko H (1990) A simple and rapid method for HLA-DP genotyping by digestion of PCR-amplified DNA with allele specific restriction endonucleases. *Hum Immunol* 27:111–121
- Makino R, Yazyu H, Kishimoto Y, Sekiya T, Hayashi K (1992) F-SSCP (fluorescence-based polymerase chain reaction-singlestrand conformation polymorphism (PCR-SSCP)) analysis. *PCR Methods Appl* 2:10–13
- Malik A, Azam S, Hadi N, Hadi SM (2003) DNA degradation by water extract of green tea in the presence of copper ions: implications for anticancer properties. *Phytother Res* 17(4):358–363
- Mardis ER (2008) The impact of next-generation sequencing technology on genetics. *Trends Genet* 24(3):133–141
- Matsuyama S, Nishi K (2011) Genus identification of toxic plant by real-time PCR. *Int J Legal Med* 125(2):211–217
- Mazzanti G, Battinelli L, Daniele C, Costantini S, Ciaralli L, Evandri MG (2008) Purity control of some Chinese crude herbal drugs marketed in Italy. *Food Chem Toxicol* 46:3043–3047
- McDermott JM, Brandle U, Dutly F, Haemmerli UA, Keller S, Muller KE, Wolf MS (1994) Genetic variation in powdery mildew of barley: development of RAPD, SCAR and VNTR markers. *Phytopathology* 4:1316–1321
- McIntyre M (1998) Chinese herbs: risk, side effects, and poisoning: the case for objective reporting and analysis reveals serious misrepresentation. *J Altern Complement Med* 4(1):15–16
- Mills S, Bone K (2005) *The essential guide to herbal safety*. Churchill Livingstone, London
- Mitchell D, Willerslev E, Hansen A (2005) Damage and repair of ancient DNA. *Mutat Res* 571:265–276

- Mizukami H, Ohbayashi K, Kitamura Y, Ikenaga T (1993a) Restriction fragment length polymorphisms (RFLPs) of medicinal plants and crude drugs. I. RFLP probes allow clear identification of *Duboisia* interspecific hybrid genotypes in both fresh and dried tissues. *Biol Pharm Bull* 16:388–390
- Mizukami H, Ohbayashi K, Umetsu K, Hiraoka N (1993b) Restriction fragment length polymorphism of medicinal plants and crude drugs. II. Analysis of *Glehnia littoralis* of different geographical origin. *Biol Pharm Bull* 16:611–612
- Mizukami H, Okabe Y, Kohda H, Hiraoka N (2000) Identification of the crude drug *Atractylodes rhizome* (Byaku-jutsu) and *Atractylodes lancea* rhizome (So-jutsu) using chloroplast trnK sequence as a molecular marker. *Biol Pharm Bull* 23:589–594
- Molina C, Kahl G (2002) Genomics of two banana pathogens, genetic diversity, diagnostics, and phylogeny of *Mycosphaerella fijiensis* and *M. musicola*. In: Jain SM (ed) *Banana improvement, cellular and molecular biology, and induced mutations*. FAO/IAEA, Vienna
- Mondini L, Noorani A, Pagnotta MA (2009) Assessing plant genetic diversity by molecular tools. *Diversity* 1:19–35
- Morgante M, Vogel J (1994) Compound microsatellite primers for the detection of genetic polymorphisms. US Patent Application 08/326456
- Mori N, Moriguchi T, Nakamura C (1997) RFLP analysis of nuclear DNA for study of phylogeny and domestication of tetraploid wheat. *Genes Genet Syst* 72:153–161
- Moritz C, Cicero C (2004) DNA barcoding: promise and pitfalls. *PLoS Biol* 2(10):e354
- Mueller UG, Wolfenbarger L (1999) AFLP genotyping and fingerprinting. *Trends Ecol Evol* 14:389–394
- Munthali M, Ford-Lloyd BV, Newbury HJ (1992) The random amplification of polymorphic DNA for fingerprinting plants. *PCR Methods Appl* 1:274–276
- Nagamine K, Watanabe K, Ohtsuka K, Hase T, Notomi T (2001) Loop-mediated isothermal amplification reaction using a non-denatured template. *Clin Chem* 47:1742–1743
- Nagamine K, Hase T, Notomi T (2002) Accelerated reaction by loop-mediated isothermal amplification using loop primers. *Mol Cell Probes* 16:223–229
- Nakai R, Shoyama Y, Shiraishi S (1996) Genetic characterization of *Epimedium* species using random amplified polymorphic DNA (RAPD) and PCR-restriction fragment length polymorphism (RFLP) diagnosis. *Biol Pharm Bull* 19(1):67–70
- Nalawade SM, Sagare AP, Lee CY et al (2003) Studies on tissue culture of Chinese medicinal plant resources in Taiwan and their sustainable utilization. *Bot Bull Acad Sin* 44:79–98
- NEPA National Environmental Protection Agency (1994) China biodiversity conservation action plan. China Environmental Science Press, Beijing
- Newmaster SG, Fazekas AJ, Ragupathy S (2006) DNA barcoding in the land plants: evaluation of rbcL in a multigene tiered approach. *Can J Bot* 84:335–341
- Newmaster SG, Fazekas AJ, Steeves RAD, Janovec J (2008) Testing candidate plant barcode regions with species of recent origin in the Myristicaceae. *Mol Ecol Notes* 8:480–490
- Newton CR, Graham A, Heptinstall LE, Powell SJ, Summers C, Kalsheker N et al (1989) Analysis of any point mutation in DNA. The amplification refractory mutation system (ARMS). *Nucleic Acids Res* 17:2503–2516
- Ng KY, Cheng CI, Xu HX (2009) Safety issues of Chinese medicine: a review of intoxication cases in Hong Kong. *Chin Herb Med* 1(1):29–39
- Ngan F, Shaw P, But P, Wang J (1999) Molecular authentication of *Panax* species. *Phytochemistry* 50:787–791
- Niu L, Mantri N, Li CG, Xue C, Wohlmuth H, Pang EC (2011) Detection of *Panax quinquefolius* in *Panax ginseng* using subtracted diversity array. *J Sci Food Agric* 18. doi:10.1002/jfsa.4319
- Normile D (2003) The new face of traditional Chinese medicine. *Science* 299:188–190
- Notomi T, Okayama H, Masubuchi H, Yonekawa T, Watanabe K, Amino N, Hase T (2000) Loop-mediated isothermal amplification of DNA. *Nucleic Acids Res* 28(12):E63
- Novak J, Grausgruber-Gröger S, Luka B (2007) DNA-based authentication of plant extracts. *Food Res Int* 40(3):388–392

- Orita M, Iwahana H, Kanazawa H, Hayashi K, Sekiya T (1989) Detection of polymorphisms of human DNA by gel electrophoresis as single-strand conformation polymorphism. *Proc Natl Acad Sci USA* 86:2766–2770
- Paglia GE, Olivieri AM, Morgante M (1998) Towards second-generation STS (sequence-tagged sites) linkage maps in conifers, a genetic map of Norway spruce (*Picea abies* K.). *Mol Genet* 258:466–478
- Pang XH, Song JY, Zhu YJ, Xie CX, Chen SL (2010) Using DNA barcoding to identify species within Euphorbiaceae. *Planta Med* 76:1784–1786
- Pang X, Songa J, Zhua Y, Xub H, Huang L, Chena S (2011) Applying plant DNA barcodes for Rosaceae species identification. *Cladistics* 27:165–170
- Paran I, Michelmore RW (1993) Development of reliable PCR based markers linked to downy mildew resistance genes in lettuce. *Theor Appl Genet* 85:985–993
- Paran I, Kesseli R, Michelmore R (1991) Identification of restriction fragment-length-polymorphism and random amplified polymorphic DNA markers linked to downy mildew resistance genes in lettuce, using near isogenic lines. *Genome* 34:1021–1027
- Passinho-Soares H, Felix D, Kaplan MA, Margis-Pinheiro M, Margis R (2006) Authentication of medicinal plant botanical identity by amplified fragmented length polymorphism dominant DNA marker: inferences from the *Plectranthus* genus. *Planta Med* 72:929–931
- Pastinen T, Partanen J, Syvanen AC (1996) Multiplex, fluorescent, solid-phase minisequencing for efficient screening of DNA sequence variation. *Clin Chem* 42:1391–1397
- Pastinen T, Raitio M, Lindroos K et al (2000) A system for specific, high-throughput genotyping by allele-specific primer extension on microarrays. *Genome Res* 10:1031–1042
- Pei S (2001) Ethnobotanical approaches of traditional medicine studies: some experiences from Asia. *Pharmaceutical Botany* 39:74–79
- Pei S (2002a) A brief review of ethnobotany and its curriculum development in China. In: Shinwari ZK, Hamilton A, Khan AA (eds) In: *Proceedings of a Workshop on Curriculum Development in Applied Ethnobotany*, Nathiagali, 2–4 May 2002. WWF-Pakistan, Lahore, Pakistan
- Pei S (2002b) Ethnobotany and modernisation of Traditional Chinese Medicine. Paper at a Workshop on Wise Practices and Experiential Learning in the Conservation and Management of Himalayan Medicinal Plants, Kathmandu, Nepal, 15–20 December 2002, supported by the Ministry of Forest and Soil Conservation, Nepal, the WWF-Nepal Program, MMAPPA and PPI
- Pei-Gen X (2007) Ethnopharmacological investigation of Chinese medicinal plants. In: Chadwick DJ, Marsh J (eds) *Ethnobotany and search for new drugs*. Ciba Foundation Symposium 185 – Ethnobotany and the Search for New Drugs
- Pennisi E (2007) Wanted: a barcode for plants. *Science* 318:190–191
- Pereira F, Carneiro J, Amorim A (2008) Identification of species with DNA-based technology: current progress and challenges. *Recent Pat DNA Gene Seq* 2:187–200
- Pharmacopoeia of the People's Republic of China (2005) State pharmacopoeia committee. Chemical Industry Publisher, Beijing, China
- Pharmacopoeia of the People's Republic of China (2010) 2010 edn. China Medical Science and Technology Press, Beijing
- Pirrung MC, Connors RV, Odenbaugh AL, Montague-Smith MP, Walcott NG, Tollett JJ (2000) The arrayed primer extension method for DNA microchip analysis. Molecular computation of satisfaction problems. *J Am Chem Soc* 122:1873–1882
- Pirttilä AM, Hirsikorpi M, Kämäräinen T, Jaakola L, Hohtola A (2001) DNA isolation methods for medicinal and aromatic plants. *Plant Mol Biol Rep* 19:273a–f
- Portis E, Acquadro A, Comino C, Lanteri S (2004) Analysis of DNA methylation during germination of pepper (*Capsicum annum* L.) seeds using methylation-sensitive amplification polymorphism (MSAP). *Plant Sci* 166:169–178
- Preeti C, Kalpana J, Bhushan P (2006) DNA microarrays in herbal drug research. Advance Access Publication
- Preeti C, Kalpana J, Bhushan P (2006b) DNA microarrays in herbal drug research. *Evid Based Complement Altern Med* 3(4):447–457

- Provan J, Biss PM, McMee D, Mathews S (2004) Universal primers for the amplification of chloroplast microsatellites in grasses (Poaceae). *Mol Ecol Notes: Primer Note*
- Qi JJ, Li XE et al (2008) Genetic relationships among *Rehmannia glutinosa* cultivars and varieties. *Planta Med* 74:1846–1852
- Qian L, Ding G et al (2008) Molecular authentication of *Dendrobium loddigesii* Rolfe by amplification refractory mutation system (ARMS). *Planta Med* 74:470–473
- Qin J, Leung FC, Fung Y, Zhu D, Lin B (2005) Rapid authentication of ginseng species using microchip electrophoresis with laser-induced fluorescence detection. *Anal Bioanal Chem* 381:812–819
- Reyna-Lopez GE, Simpson J, Ruiz-Herrera J (1997) Differences in DNA methylation patterns are detectable during the dimorphic transition of fungi by amplification of restriction polymorphism. *Mol Gen Genet* 253:703–710
- Robba L, Russell SJ, Barker GL, Brodie J (2006) Assessing the use of the mitochondrial *cox1* marker for use in DNA barcoding of Red algae (Rhodophyta). *Am J Bot* 93(8):1101–1108
- Saar DE, Polans NO, Sørensen PD, Duvall MR (2001) Angiosperm DNA contamination by endophytic fungi: detection and methods of avoidance. *Plant Mol Biol Rep* 19:249–260
- Sahin FP, Yamashita H et al (2007) DNA authentication of *Plantago* herb based on nucleotide sequences of 18S-28S rRNA internal transcribed Spacer region. *Biol Pharm Bull* 30:1265–1270
- Sarwat M, Das S et al (2008) Analysis of genetic diversity through AFLP, SAMPL, ISSR and RAPD markers in *Tribulus terrestris*, a medicinal herb. *Plant Cell Rep* 27:519–528
- Sasaki Y, Nagumo S (2007) Rapid identification of *Curcuma longa* and *C. aromatica* by LAMP. *Biol Pharm Bull* 30:2229–2230
- Sasaki Y, Fushimi H et al (2002) Sequence analysis of Chinese and Japanese *Curcuma* drugs on the 18S rRNA gene and *trnK* gene and the application of amplification-refractory mutation system analysis for their authentication. *Biol Pharm Bull* 25:1593–1599
- Sasaki Y, Komatsu K, Nagumo S (2008) Rapid detection of *Panax ginseng* by loop-mediated isothermal amplification and its application to authentication of Ginseng. *Biol Pharm Bull* 31(9):1806–1808
- Sasaki Y, Fujimoto T, Aragane M, Yasuda I, Nagumo S (2009) Rapid and sensitive detection of *Lophophora williamsii* by loop-mediated isothermal amplification. *Biol Pharm Bull* 32(5):887–891
- Sass C, Little DP, Stevenson DW, Specht CD (2007) DNA barcoding in the Cycadales: testing the potential of proposed barcoding markers for species identification of Cycads. *PLoS One* 2:e1154
- Schadt EE, Turner S, Kasarskis A (2010) A window into third-generation sequencing. *Hum Mol Genet* 19:R227–R240
- Schena M, Heller RA et al (1998) Microarrays: biotechnology's discovery platform for functional genomics. *Trends Biotechnol* 16(7):301–306
- Schippmann U, Leaman DJ, Cunningham AB (2002) Impact of cultivation and gathering of medicinal plants on biodiversity: global trends and issues. Inter-Department Working Group on Biology Diversity for Food and Agriculture, Food and Agricultural Organisation of the United Nations, Rome, Italy
- Schouten JP, McElgunn CJ, Waaijer R, Zwijnenburg D, Diepvens F, Pals G (2002) Relative quantification of 40 nucleic acid sequences by multiplex ligation-dependent probe amplification. *Nucleic Acids Res* 30(12):e57
- Schuler GD, Boguski MS, Stewart EA et al (1996) A gene map of the human genome. *Science* 274:540–546
- Semagn K, Bjørnstad A, Ndjiondjop MN (2006) An overview of molecular marker methods for plants. *Afr J Biotechnol* 5:2540–2568
- SEPA State Environmental Protection Administration (1998) China's biodiversity: a country study. China Environmental Science Press, Beijing

- Shan X, Blake TK, Talbert LE (1999) Conversion of AFLP markers to sequence-specific PCR markers in barley and wheat. *Theor Appl Genet* 98:1072–1078
- Shaw PC, Ngan FN, But PPH, Wang J (2002) Molecular markers in Chinese medicinal materials. In: Shaw PC, But PPH (eds) *Authentication of Chinese medicinal material by DNA technology*. World Scientific Publishing, Singapore
- Shaw J, Lickey EB, Schilling EE, Small RL (2007) Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: the tortoise and the hare III. *Am J Bot* 94(3):275–288
- Shaw PC, Wong KKL, Chan AWK, Wong WC, But PPH (2009) Patent applications for using DNA technologies to authenticate medicinal herbal material. *J Chin Med* 4:1–11
- Shen Y, Wu BL (2009) Designing a simple multiplex ligation-dependent probe amplification (MLPA) assay for rapid detection of copy number variants in the genome. *J Genet Genomics* 36(4):257–265
- Shen J, Ding XY, Ding G, Liu DY, Tang F, He J (2006) Studies on population difference of *Dendrobium officinale* II establishment and optimization of the method of ISSR fingerprinting marker. *Zhongguo Zhongyao Zazhi* 31:291–294
- Shi HM, Wang J, Wang MY, Tu PF, Li XB (2009) Identification of *Cistanche* species by chemical and inter-simple sequence repeat fingerprinting. *Biol Pharm Bull* 32:142–146
- Shim YH, Choi JH, Park CD et al (2003) Molecular differentiation of *Panax* species by RAPD analysis. *Arch Pharm Res* 26:601–605
- Shneyer VS (2009) DNA Barcoding is a new approach in comparative genomics of plants. *Russ J Genet* 45(11):1267–1278
- Silva LM, Montes de Oca H, Diniz CR, Fortes-Dias CL (2001) Fingerprinting of cell lines by directed amplification of minisatellite-region DNA (DAMD). *Braz J Med Biol Res* 34:1405–1410
- Siow YL, Gong Y, Au-Yeung KK, Woo CW, Choy PC (2005) Emerging issues in traditional Chinese medicine. *Can J Physiol Pharmacol* 83:321–334
- Slanc P, Ravnikar M, Strukelj B (2006) Identification of individual herbal drugs in tea mixtures using restriction analysis of ITS DNA and real-time PCR. *Pharmazie* 61(11):912–915
- Smillie TJ, Khan IA (2010) A comprehensive approach to identifying and authenticating botanical products. *Clin Pharmacol Ther* 87:175–186
- Society of Japanese Pharmacopoeia (2001) *Japanese pharmacopoeia* (English version), 14th edn. Junkudo Book Shop, Tokyo
- Somers DJ, Demmon G (2002) Identification of repetitive, genome-specific probes in crucifer oilseed species. *Genome* 45:485–492
- Song H, Buhay JE, Whiting MF, Crandall KA (2008) Many species in one: DNA barcoding overestimates the number of species when nuclear mitochondrial pseudogenes are coamplified. *Proc Natl Acad Sci USA* 105(36):13486–13491
- Song JY, Yao H, Li Y, Li XW, Lin YL, Liu C, Han JP, Xie CX, Chen SL (2009) Authentication of the family Polygonaceae in Chinese pharmacopoeia by DNA barcoding technique. *J Ethnopharmacol* 124:434–439
- Song Z, Li X, Wang H, Wang J (2010) Genetic diversity and population structure of *Salvia miltiorrhiza* Bge in China revealed by ISSR and SRAP. *Genetica* 138:241–249
- Soumaya R, Dakhlaoui-Dkhil S, Salem AOM, Zehdi-Azouzi S, Rhouma A, Marrakchi M, Trifi M (2008) Genetic diversity and phylogenetic relationships in date-palms (*Phoenix dactylifera* L.) as assessed by random amplified microsatellite polymorphism markers (RAMPOs). *Scientia Hort* 117:53–57
- Squirrel J, Hollingsworth PM, Woodhead M, Russell J, Lowe AJ, Gibby M, Powell W (2003) How much effort is required to isolate nuclear microsatellites from plants? *Mol Ecol* 12:1339–1348
- Srivastava J, Lambert J, Vietmeyer N (1996) *Medicinal plants: an expanding role in development* [M]. World Bank Publications, Washington, DC

- Sucher JN, Carles MC (2008) Genome-based approaches to the authentication of medicinal plants. *Planta Med* 74:603–623
- Sucher NJ, Hennel JR, Carles MC (2012) Plant DNA fingerprinting and barcoding. *Methods Mol Biol* 862:13–22
- Sun Y, Fung KP, Leung PC, Shi D, Shaw PC (2004) Characterization of medicinal *Epimedium* species by 5S rRNA gene spacer sequencing. *Planta Med* 70:287–288
- Sun Y, Fung KP, Leung PC, Shaw PC (2005) A phylogenetic analysis of *Epimedium* (Berberidaceae) based on nuclear ribosomal DNA sequences. *Mol Phylogenet Evol* 35(1):287–291
- Sun Y, Shaw PC, Fung KP (2007) Molecular authentication of *Radix Puerariae Lobatae* and *Radix Puerariae Thomsonii* by ITS and 5S rRNA spacer sequencing. *Biol Pharm Bull* 30:173–175
- Sun ZY, Gao T, Yao H, Shi LC, Zhu YJ, Chen SL (2010) Identification of *Lonicera japonica* and its related species using the DNA barcoding method. *Plant Med*. doi:10.1055/s-0030-1250324
- Sze SCW, Song JX, Wong RNS, Feng YB, Ng TB, Tong Y, Zhang KYB (2008) Application of SCAR (sequence characterized amplified region) analysis to authenticate *Lycium barbarum* (wolfberry) and its adulterants. *Biotech Applied Biochem* 51:15–21
- Taberlet P et al (2007) Power and limitations of the chloroplast *trnL*(UAA) intron for plant DNA barcoding. *Nucleic Acids Res* 35:e14
- Takamiya T, Wongsawad P, Tajima N, Shioda N, Lu JF, Wen CL, Wu JB, Handa T, Iijima H, Kitanaka S, Yukawa T (2011) Identification of *Dendrobium* species used for herbal medicines based on ribosomal DNA internal transcribed spacer sequence. *Biol Pharm Bull* 34(5):779–82
- Techaprasan J, Klinbunga S, Jenjittiku T (2008) Genetic relationships and species authentication of *Boesenbergia* (Zingiberaceae) in Thailand based on AFLP and SSCP analyses. *Biochem Syst Ecol* 36(5–6):408–441
- Ten Kate K, Laird S (1999) The commercial use of biodiversity: access to genetic resources and benefit-sharing. Commission of the European Communities and Earthscan, London, p 398
- Theerakulpisut P, Kanawapee N, Maensiri D, Bunnag S, Chantaranothai P (2008) Development of species-specific SCAR markers for identification of three medicinal species of *Phyllanthus*. *J Syst Evol* 46:614–621
- Thorne RF (2002) How many species of seed plants are there? *Taxon* 51:511–522
- Trau D, Lee TM, Lao AI, Lenigk R, Hsing IM, Ip NY et al (2002) Genotyping on a complementary metal oxide semiconductor silicon polymerase chain reaction chip with integrated DNA microarray. *Anal Chem* 74:3168–3173
- Trifi-Farah N, Marrakchi M (2001) *Hedysarum* phylogeny mediated by RFLP analysis of nuclear ribosomal DNA. *Gen Res Crop Evol* 48:339–345
- Tsoi PY, Wu HS, Wong MS, Chen SL, Fong WF, Xiao PG et al (2003) Genotyping and species identification of *Fritillaria* by DNA chip technology. *Acta Pharm Sin* 4:185–190
- Um JY, Chung HS, Kim MS, Na HJ, Kwon HJ, Kim JJ, Lee KM, Lee SJ, Lim JP, Do KR, Hwang WJ, Lyu YS, An NH, Kim HM (2001) Molecular authentication of *Panax ginseng* species by RAPD analysis and PCR-RFLP. *Biol Pharm Bull* 24:872–875
- Valdes AM, Slatkin M, Freimer NB (1993) Allele frequencies at microsatellite loci – the stepwise mutation model revisited. *Genetics* 133:737–749
- Varshney RV, Graner A, Sorrells ME (2005) Genic microsatellite markers in plants: features and applications. *Trends Biotechnol* 23:28–55
- Vences M, Thomas M, Bonett RM, Vieites DR (2005) Deciphering amphibian diversity through DNA barcoding: chances and challenges. *Philos Trans R Soc London Ser B* 360:1859–1868
- Vijayan K, Tsou CH (2010) DNA barcoding in plants: taxonomy in a new perspective. *Curr Sci* 99 (11):1513–1541
- Virgilio M, Backeljau T, Nevado Band De Meyer M (2010) Comparative performances of DNA barcoding across insect orders. *BMC Bioinformatics* 11:206
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23:4407–4414

- Wagner H, Bauer R, Melchart D, Xiao PG, Staudinger A (2011) Chromatographic fingerprint analysis of herbal medicines: thin-layer and high performance liquid chromatography of Chinese drugs, Vol I and Vol II, 2nd edn. Springer, Wien
- Wang H, Wang Q (2005) Analysis of rDNA ITS sequences of *Radix et Rhizoma Salviae miltiorrhizae* and plants of *Salvia L.* *Chin Tradit Herb Drugs* 36:1381–1385
- Wang P, Huang F, Zhou L, Cao L, Liang S, Xu H, Liu J (2000) Analysis of *Amomun villosum* species and some adulterants of Zingiberaceae by RAPD. *Zhong Yao Cai* 23:71–74
- Wang J, Ha WY, Ngan FN, But PP, Shaw PC (2001) Application of sequence characterized amplified region (SCAR) analysis to authenticate *Panax* species and their adulterants. *Planta Med* 67:781–783
- Wang CZ, Li P, Ding JY, Jin GQ, Yuan CS (2005) Identification of *Fritillaria pallidiflora* using diagnostic PCR and PCR-RFLP based on nuclear ribosomal DNA internal transcribed spacer sequences. *Planta Med* 71:384–386
- Wang CZ, Li P, Ding JY et al (2007a) Simultaneous identification of *Bulbus Fritillariae cirrhosae* using PCR-RFLP analysis. *Phytomedicine* 14:628–632
- Wang MW, Hao XJ, Chen KX (2007b) Biological screening of natural products and drug innovation in China. *Philos Trans R Soc Lond B Biol Sci* 362(1482):1093–1105
- Wang Y, Li XE, Li XD, Qi JJ, Sun P, Zhou LL (2008) Analysis of genetic diversity of wild *Rehmannia glutinosa* by using RAPD and ISSR markers. *Zhongguo Zhongyao Zazhi* 33(22):2591–2595
- Wang HZ, Wu ZX, Lu JJ et al (2009) Molecular diversity and relationships among *Cymbidium goeringii* cultivars based on inter-simple sequence repeat (ISSR) markers. *Genetica* 136 (3):391–399
- Wang HT, Sun H, Kwon WS, Jin H, Yang DC (2010a) A PCR-based SNP marker for specific authentication of Korean ginseng (*Panax ginseng*) cultivar “Chunpoong”. *Mol Biol Rep* 37:1053–1057
- Wang W, Wu Y, Yan Y, Ermakova M, Kerstetter R et al (2010b) DNA barcoding of the Lemnaceae, a family of aquatic monocots. *BMC Plant Biol* 10(1):205
- Ward RD, Hanner R, Hebert PDN (2009) The campaign to DNA barcode all fishes, FISH-BOL. *J Fish Biol* 74:329–356
- Watanabe A, Araki S, Kobari S, Sudo H, Tsuchida T, Uno T et al (1998) In vitro propagation, restriction fragment length polymorphism, and random amplified polymorphic DNA analyses of *Angelica* plants. *Plant Cell Rep* 18:187–192
- Waugh J (2007) DNA barcoding in animal species: progress, potential and pitfalls. *Bioessays* 29 (2):188–197
- Weising K, Gardner RC (1999) A set of conserved PCR primers for the analysis of simple sequence repeat polymorphisms in chloroplast genomes of dicotyledonous angiosperms. *Genome* 42:9–11
- Weising K, Kahl G (1998) Hybridization-based microsatellite fingerprinting of plants. In: Caetano-Anolles G, Gresshoff PM (eds) *DNA markers: protocols, applications, and overviews*. Wiley, New York, pp 238–243
- Weising K, Bybom H, Wolff K, Kahl G (2005) *DNA fingerprinting in plants. Principles, methods, and application*, 2nd edn. Taylor and Francis, Boca Raton
- Welsh J, McClelland M (1990) Fingerprinting genomes using PCR with arbitrary primers. *Nucleic Acids Res* 18:7213–7218
- Winter E, Pfaff T, Udupa SM, Hiittel B, Sharma PC, Sahi S, Arreguin-Espinoza R, Weigand F, Muehlbauer FJ, Kahl G (1999) Characterization and mapping of sequence-tagged microsatellite sites in the chickpea (*Cicer arietinum L.*) genome. *Mol Gen Genet* 262:90–101
- Witsenboer H, Vogel J, Michelmore RW (1997) Identification, genetic localization, and allelic diversity of selectively amplified polymorphic loci in lettuce and wild relatives (*Lactuca* spp.). *Genome* 40:923–936
- Wolfe KH, Li WH, Sharp PM (1987) Rates of nucleotide substitution vary greatly among plant mitochondrial, chloroplast, and nuclear DNAs. *Proc Natl Acad Sci USA* 84(24):9054–9058

- Wong KL, Wang J, But PPH, Shaw PC (2001) Application of cytochrome b DNA sequences for the authentication of endangered snake species. *Forensic Sci Int* 139:49–55
- Wu H, Lien EJ, Lien LL (2003) Chemical and pharmacological investigations of *Epimedium* species: a survey. *Prog Drug Res* 60:1–57
- Wu W, Zheng YL, Chen L, Wei YM, Yang RW, Yan ZH (2005) Evaluation of genetic relationships in the genus *Houttuynia* Thunb. in China based on RAPD and ISSR markers. *Biochem Syst Ecol* 33:1141–1157
- Xia Q, Zhao KJ, Huang ZG, Zhang P, Dong TT, Li SP et al (2005) Molecular genetic and chemical assessment of *Rhizoma Curcumae* in China. *J Agric Food Chem* 53:6019–6026
- Xia T, Chen S, Chen S, Zhang D, Zhang D, Gao Q et al (2007) ISSR analysis of genetic diversity of the Qinghai-Tibet plateau endemic *Rhodiola chrysanthemifolia* (Crassulaceae). *Biochem Syst Ecol* 35:209–214
- Xiao PG (1994) Ethnopharmacological investigation of Chinese medicinal plants. *Ciba Found Symp* 185:169–173
- Xiao PG, Yong P (1998) Ethnopharmacology and research on medicinal plants in China. In: Prendergast HDV, Etkin NL, Harris DR, Houghton PJ (eds) *Plants for food and medicine. Proceedings from a Joint Conference of the Society for Economic Botany and the International Society for Ethnopharmacology*, London, Royal Botanic Gardens, Kew, UK, pp 31–39
- Xiong LZ, Xu CG, Saghai-Marouf MA, Zhang QF (1999) Pattern of cytosine methylation in an elite rice hybrid and its potential lines, detected by a methylationsensitive amplification polymorphism technique. *Mol Gen Genet* 261:439–446
- Xu M, Li X, Korban SS (2000) AFLP-Based detection of DNA methylation. *Plant Mol Biol Rep* 18:361–368
- Xu M, Huaracha E, Korban SS (2001) Development of sequence-characterized amplified regions (SCARs) from amplified fragment length polymorphism (AFLP) markers tightly linked to the Vf gene in apple. *Genome* 44:63–70
- Xu H, Wang Z, Ding X, Zhou K, Xu L (2006) Differentiation of *Dendrobium* species used as “Huangcao Shihu” by rDNA ITS sequence analysis. *Planta Med* 72:89–92
- Xue D (1997) China’s national report on implementation of the Convention on Biological Diversity. China Environmental Science Press, Beijing
- Xue CY, Xue HG (2008) Application of real-time scorpion PCR for authentication and quantification of the traditional Chinese medicinal plant *Drynaria fortunei*. *Planta Med* 74:e765–e770
- Xue CY, Li DZ, Lu JM, Yang JB, Liu JQ (2006a) Molecular authentication of the traditional Tibetan medicinal plant *Swertia mussotii*. *Planta Med* 72:1223–1226
- Xue HG, Zhou SD, He XJ, Yu Y (2006b) Molecular authentication of the traditional Chinese medicinal plant *Euphorbia pekinensis*. *Planta Med* 73:91–93
- Xue CY, Li DZ, Wang QZ (2008a) Identification and quantification of the traditional Chinese medicinal plant *Gentiana macrophylla* using Taqman real-time PCR. *Planta Med* 74 (15):1842–1845
- Xue HG, Wang H, Li DZ, Xue CY, Wang QZ (2008b) Differentiation of the traditional Chinese medicinal plants *Euphorbia humifusa* and *E. maculata* from adulterants by TaqMan real-time polymerase chain reaction. *Planta Med* 74:302–304
- Yamaji H, Fukuda T, Yokoyama J, Pak JH, Zhou C, Yang C et al (2007) Reticulate evolution and phylogeography in *Asarum* sect. *asiasarum* (Aristolochiaceae) documented in internal transcribed spacer sequences (ITS) of nuclear ribosomal DNA. *Mol Phylogenet Evol* 44:863–884
- Yamasaki M, Sato A, Shimomura K, Saito K, Murakoshi I (1994) Genetic relationships among *Glycyrrhiza* plants determined by RADP and RFLP analyses. *Biol Pharm Bull* 17:1529–1531
- Yamazaki M, Sato A, Saito K, Murakoshi I (1993) Molecular phylogeny based on RFLP and its relation with alkaloid patterns in *Lupinus* plants. *Biol Pharm Bull* 16:1182–1184
- Yang HY, Chen CF (1998) Pharmacology and toxicology of herbal medicine: subacute toxicity of commonly used Chinese drugs. *J Toxicol Sci* 23(Suppl 2):229–233
- Yang ZY, Ling Z (2010) Screening potential DNA barcode regions in *Amomum* (Zingiberaceae). *Acta Bot Yunnan* 32(5):393–400

- Yang M, Zhang D, Liu J, Zheng J (2001) A molecular marker that is specific to medicinal rhubarb based on chloroplast trnL/trnF sequences. *Planta Med* 67:784–786
- Yang DY, Hirotoishi F, Shao-Qing C, Katsuko K (2004) Polymerase chain reaction–restriction fragment length polymorphism (PCR-RFLP) and amplification refractory mutation system (ARMS) analyses of medicinally used rheum species and their application for identification of Rhei rhizoma. *Biol Pharm Bull* 27:661–669
- Yang ZY, Chao Z, Huo KK, Wu BY, Pan SL (2006) Nuclear ribosomal DNA internal transcribed spacer 1 sequences of 4 *Leonurus* species. *Nan Fang Yi Ke Da Xue Xue Bao* 26:1593–1595
- Yang ZY, Chao Z et al (2007) ITS sequence analysis used for molecular identification of the *Bupleurum* species from northwestern China. *Phytomedicine* 14:416–422
- Yao H, Zhao Z, Chen DF, Chen JK, Zhou TS (2008) ISSR primer screening and preliminary evaluation of genetic diversity in wild populations of *Glycyrrhiza uralensis*. *Biol Plantarum* 52(1):117–120
- Yao H, Song J, Liu C, Luo K, Han J et al (2010) Use of ITS2 region as the universal DNA barcode for plants and animals. *PLoS One* 5(10):e13102
- Ye Q, Qiu YX, Quo YQ, Chen JX, Yang SZ, Zhao MS, Fu CX (2006) Species-specific SCAR markers for authentication of *Sinocalycanthus chinensis*. *J Zhejiang Univ Sci B* 7:868–872
- Ying XC, Zhu D, Wang Q, Zhong WQ (2009) Application of Light Cycler polymerase chain reaction and melting curve analysis to the authentication of the traditional Chinese medicinal plant *Cimicifuga foetida*. *Planta Med* 75(8):873–874
- Yip PY, Kwan HS (2006) Molecular identification of *Astragalus membranaceus* at the species and locality levels. *J Ethnopharmacol* 106:222–229
- Yip PY, Chau CF, Mak CY, Kwan HS (2007) DNA methods for identification of Chinese medicinal materials. *J Chin Med* 2:1–19
- Zabeau M, Vos P (1993) Selective restriction fragment amplification: a general method for DNA fingerprinting. European Patent Application number 92402629.7. Publication number 0534858A1
- Zane L, Bargelloni L, Patarnello T (2002) Strategies for microsatellite isolation: a review. *Mol Ecol* 11:1–16
- Zerega NJC, Mori S, Lindqvist C, Zheng Q, Motley TJ (2002) Using amplified fragment length polymorphisms (AFLP) to identify black cohosh (*Actaea racemosa*). *Econ Bot* 56:154–164
- Zhang R, Zhang B, Ye H (1997) Identification of the Chinese herbs of Indigoferae L. by RAPD analysis. *Hongguo Zhong Yao Za Zhi* 22:72–73
- Zhang YB, Ngan FN, Wang ZT, Wang J, But PPH, Shaw PC (1999) Differentiation of *Codonopsis pilosula* using random amplified polymorphic DNA. *Planta Med* 65:57–60
- Zhang KYB, Leung HW et al (2001a) Differentiation of *Lycium barbarum* from its related *Lycium* species using random amplified polymorphic DNA. *Planta Med* 67:379–381
- Zhang M, Huang HR, Liao SM, Gao JY (2001b) Cluster analysis of *Dendrobium* by RAPD and design of specific primer for *Dendrobium candidum*. *Zhongguo Zhong Yao Za Zhi* 26:442–447
- Zhang YB, Wang J, Wang ZT, But PP, Shaw PC (2003) DNA microarray for identification of the herb of *Dendrobium* species from Chinese medicinal formulations. *Planta Med* 69:1172–1174
- Zhang M, Zhang DZ, Xu XH, Zhang T, Wang ZT (2005a) 5S rRNA gene spacer sequences from *Ligularia* medicinal plants and the identification of HPAs-containing species. *Chin J Nat Med* 3:38–40
- Zhang T, Xu LS, Wang ZT, Zhou KY, Zhang N, Shi YF (2005b) Molecular identification of medicinal plants: *Dendrobium chrysanthum*, *Dendrobium fimbriatum* and their morphologically allied species by PCR-RFLP analyses. *Yao Xue Xue Bao* 40:728–733
- Zhang YB, Shaw PC, Sze CW, Wang ZT, Tong Y (2007) Molecular authentication of Chinese herbal materials. *J Food Drug Anal* 15:1–9
- Zhang JM, Wang JX, Xia T, Zhou SL (2009) DNA barcoding: species delimitation in tree peonies. *Sci China C Life Sci* 52(6):568–578
- Zhao ZZ, Li TYS (2004) Hong Kong commonly confused Chinese medicines. Chinese Medicine Merchants Association, Hong Kong

- Zhao KJ, Dong TT, Cui XM, Tu PF, Tsim KW (2003a) Genetic distinction of radix adenophorae from its adulterants by the DNA sequence of 5S-rRNA spacer domains. *Am J Chin Med* 31:919–926
- Zhao KJ, Dong TT, Tu PF, Song ZH, Lo CK, Tsim KW (2003b) Molecular genetic and chemical assessment of radix *Angelica* (*Danggui*) in China. *J Agric Food Chem* 51:2576–2583
- Zhao Z, Hu Y, Liang Z, Yuen J, Jiang Z, Leung KSY (2006) Authentication is fundamental for standardization of Chinese medicines. *Planta Med* 72:865–874
- Zhao YP, Qiu YX et al (2007a) Authentication of *Actinidia macrosperma* using PCR-RFLP based on *trnK* sequences. *Bot Studies* 48:239–242
- Zhao ZL, Leng CH et al (2007b) Identification of *Dryopteris crassirhizoma* and the adulterant species based on cpDNA *rbcL* and translated amino acid sequences. *Planta Med* 73:1230–1233
- Zhao ZZ, Yuen JPS, Wu J, Yu T, Huang W, Zhao ZZ, Li TYS (2007c) Easily confused Chinese medicines in Hong Kong. *Ann Acad Med* 35(11):764–769
- Zhen-Yan Y, Ling Z (2010) Screening potential DNA barcode regions in *Amomum* (*Zingiberaceae*). *Acta Bot Yunnan* 32(5):393–400
- Zhou Z, Bebeli PJ et al (1997) Direct amplification of minisatellite-region DNA with VNTR core sequences in the genus *Oryza*. *Theor Appl Genet* 95:942–949
- Zhu YP (2002) Toxicity of the Chinese herb mu tong (*Aristolochia manshuriensis*). What history tells us. *Adverse Drug React Toxicol Rev* 21(4):171–177
- Zhu S, Fushimi H, Cai S, Komatsu K (2003) Phylogenetic relationship in the genus *Panax*: inferred from chloroplast *trnK* gene and nuclear 18S rRNA gene sequences. *Planta Med* 69:647–653
- Zhu S, Fushimi H, Cai S, Komatsu K (2004) Species identification from Ginseng drugs by multiplex amplification refractory mutation system (MARMS). *Planta Med* 70:189–192
- Zhu S, Hirotooshi Fushimi H, Komatsu K (2008) Development of a DNA microarray for authentication of ginseng drugs based on 18S rRNA gene sequence. *J Agric Food Chem* 56(11):3953–3959
- Zhu YJ, Si C, Hui Y et al (2010) DNA barcoding the medicinal plants of the genus *Paris*. *Yaoxue Xuebao* 45(3):376–382
- Zietkiewicz E, Rafalski A, Labuda D (1994) Genome fingerprinting by simple sequence repeat (SSR)-anchored polymerase chain reaction amplification. *Genomics* 20(2):176–183
- Zuo YJ, Chen ZJ, Kondo K et al (2011) DNA barcoding of *Panax* species. *Planta Med* 77(2):182–187