


**ICONES SELECTÆ
HYMENOMYCETUM
NON DUM DELINEATORUM.**

VOLUMEN SECUNDUM.

V.





CORTINARIUS Fr.

Velum araneosum, a pilei cuticula discretum, superficiale. Hymenophorum cum stipite contiguum. Lamellæ persistentes, aridæ, decolorantes, sporis tarde secedentibus pulverulentæ. Trama fibrillosa. Sporæ globosæ vel oblongæ, passim apiculatæ, supra chartam albam subochraceæ.

Vastum genus, peculiarem exhibens typum et a perito illico recognitum, etiamsi — complures colligens series — levibus notis videatur limitatum.

I. PHLEGMACIUM. Pileus æqualiter carnosus, viscosus. Stipes firmus, siccus.

CCCLXXXII. CORTINARIUS TRIUMPHANS Fr.

Epicr. ed. I p. 256, Monogr. II p. 4.

Diagn. Phlegmacium pileo carnoso, convexo-explanato, lævi, luteo, squamis obscurioribus, adpressis, secedentibus primum maculato; stipite solido, clavato, annulis pluribus concentricis squamosis fulvescentibus peronato; lamellis emarginatis, confertis, integerrimis, e pallido argillaceis.

In nemoribus subhumidis graminosis Sueciæ, Fenniæ, Angliæ.

Descr. Amplus, speciosus, stipitis indumento columnam triumphalem quasi imitans. Stipes firmus, a basi ovato-bulbosa attenuatus, 3—5 unc. longus, $\frac{1}{2}$ —1 unc. crassus, striatus, albo-lutescens, squamis subfulvis in plures circulos dispositis (sed facile deciduis) ornatus; cortina supera, interdum subannulata. Pileus haud admodum crassus, obtusus, regularis, 3—6 unc. latus, udus viscidus, crustallino-

flavus vel ochraceus, siccus luteus, disco squamis minutis adpressis maculæformibus variegatus vel omnino nudus; margo lævis, haud infractus. Caro pilei stipitisque compacta, alba. Lamellæ 3 lin. circiter latæ, denticulo decurrente et in zona contigua farinacea alba desinente, ad latera dense venosæ, planæ, albidae vel dilutissime in cæsium vergentes, demum argillaceo-subcinnamomeæ.

Exstat simul — in betuletis aridis — *forma minor* inferne tantum squamoso-zonata, distinctius annulata, pileo aureo, lamellis primitus cæsio-albis.

Tab. 141 f. 1 *C. triumphans* formam majorem typicam illustrat.

CCCLXXXIII. CORTINARIUS CLARICOLOR. Fr.

Epicr. I p. 257, Monogr. II p. 5.

Diagn. Phlegmacium pileo compacto, explanato, glabro, luteo, sicco nitido, saepè squamoso-diffracto; stipite solido, attenuato, primo floccoso-squamoso veloque albis; lamellis subadnexis, confertis, serratis, e cæsio-albido pallidis.

In betuletis apricis ericotosis Smolandiaæ et Hallandiaæ non raro.

Descr. Admodum versiformis, sed semper durus et compactus. Stipes nunc curtus, bulbosus, nunc elongatus, conico-attenuatus vel cylindricus, usque ad cortinam — supra quam albo-farinaceus — floccis squamosis concentricis albis vel tomento laxo vestitus; in adultis vero cortina et dein squamæ quoque disparent. Pileus totus carnosus, convexo-

explanatus demumque depresso, 3—5 unc. latus, primitus undique, mox margine tantum villo sericeo superficiali velatus (ad analogiam *Ag. caperati*), dein omnino glaber, lævis sed saepius diffracto-squamosus. Cara alba. Lamellæ nunc emarginatæ, fere liberæ, nunc adnatæ, primitus albidae vel cæsio-albæ, dein argillaceæ, acie inæquali.

Inter *C. triumphans* et *C. turmalem* medius, statura compacta, pileo rimoso-diffracto, stipite albo-squamoso cet. satis insignis.

Tab. 141 f. 2 *C. claricoloris* refert imaginem.

CCCLXXXIV. CORTINARIUS CRASSUS Fr.

Epicr. ed. I p. 257, Monogr. II p. 6.

Diagn. Phlegmacium [?] pileo crasse carnosus, plano depresso, unicolor, disco glabro, ambitu in fibrillas innatas soluto; stipite obeso, fibrilloso, albo, apice albo-farinaceo; lamellis rotundatis, confertis, integerrimis, e pallido argillaceis.

Loci graminosis humidiusculis in par. Femsjö Smolandiae rara haec atque abnormis species hactenus tantummodo lecta.

Descr. Valde notabilis, curtus, robustus, carne tamen molli, olida, pallescente. Stipes solidus, valde carnosus, 2—3 unc. longus, unc. crassus, nunc valde bulbosus, nunc aequalis, basi saepe albo-tomentosus, compressus succum

aqueum fundens. Cortina tenuissima, fugacissima. Pileus quam maxime carnosus, 3—5 unc. latus, sordide luteus, opacus, disco tantum primitus viscidulus, ambitu innato-fibrillosus. Lamellae pallidæ, dein argillaceæ [in fig. justo obscuriores].

Rectius forsitan ad Inolomata (juxta *C. opinum*) transferatur, sed manifesta, ut videtur, affinitas cum *C. balteato* heic inserendum suadet.

Tab. 142 f. 1 *C. crassum* sistit.

CCCLXXXV. CORTINARIUS BALTEATUS Fr.

Epicr. ed. I p. 257, Monogr. II p. 7.

Diagn. Phlegmacium pileo compacto, applanato, viscido, mox sicco, in floccos innatos soluto, margine subcærnescente, sericeo, inflexo; stipite obeso, solido, apice velutino cortinatoque albo; lamellis emarginatis, decurrentibus, confertis, integerrimis, e pallido subalutaceis.

In silvis, præcipue pinetis, Sueciæ admodum rarus, sed pluribus locis plures observatus.

Descr. Valde mutabilis et varians, ut diversas formas communis ægre comprehendat descriptio. Stipes admodum compactus, nunc aequalis, nunc ovato-bulbosus, 1 $\frac{1}{2}$ —2 unc. longus, unc. et ultra crassus, supra cortinam fibrillosam subtiliter velutinus sordidus, infra fibrillis adpressis ferrugina-scentibus longitudinaliter striatus vel reticulatus, in alia forma tomentosus, albus. Pileus applanato-disciformis, obtusissi-

mus, vulgo disco depresso et saepe inæqualis, 3 unc. et ultra latus, primitus viscidus (siccus nitens), mox vero siccus, cute innato-floccosa et marginem versus fibrillosa nitida, disco sordide fulvo-fuligineo vel spadiceo, zona marginali pulchre violacea vel lilacina. In alia forma pileus villosus vel tomentoso-rugosus, subferrugineus, margine violaceo-tomentoso; in alia nil violacei adest (etiam in exempl. vetustis saepe disparet), margo testaceo-spadiceus primitusque villo sericeo velatus. Caro compacta, alba. Lamellæ pulchre emarginatae, primo pallidæ, dein subalutaceæ [in iconis ex. perfecto color solito obscurior] vel rotundatae, argillaceæ — vel subdecurrentes, valde angustæ, albidae.

Tab. 142 f. 2 *C. balteati* exhibet figuræ.

CCCLXXXVI. CORTINARIUS SEBACEUS Fr.

Epicr. ed. I p. 258, Monogr. II p. 7.

Diagn. Phlegmacium pileo carnosus, infracto-expanso, subrepando, glabro sed primitus micore pruinoso albido obducto, viscidulo, pallescente; cortina tenella, fugaci, margini pilei tantum adhaerente; stipite solido, valido, aequali, albo-pallente; lamellis emarginatis, haud confertis, argillaceis.

In silvis mixtis, præcipue pinetis, regionum campestrium. In Gallia QUÉLET.

Descr. Stipes compactus, numquam bulbosus, paullum fibrillosum, 3—5 unc. et ultra longus, 1 $\frac{1}{2}$ —1 unc. crassus.

Pileus aequaliter carnosus, 2—5 unc. latus, e convexo planiusculus, unicolor, colore sebi pallescens. Caro alba, sapor mitis. Lamellæ demum latæ (4 lin. circiter), venoso-connexæ, argillaceæ vel pallide cinnamomeæ (numquam cæsiæ), ad latera pallidiores. Sporæ dilute cinnamomeæ.

Species habitu sat ignobilis ideoque saepe neglecta, sed semel rite distincta semper recognitu facilis.

Tab. 143 f. 1 *C. sebacei* exemplar majus adumbrat.

CCCLXXXVII. CORTINARIUS VARIICOLOR (Pers.)

Pers. Syn. p. 280, Fries Epicr. ed. II p. 328.

Diagn. Phlegmacium pileo compacto, convexo-expanso, viscido, discoideo, margine subtomentoso-violaceo; stipite duro, crasso, bulboso, villoso-flocculoso, cæruleo-albicante; lamellis decurrenti-emarginatis, subarcuatis, confertis, e cærulescenti-argillaceo cinnamomeis vel fuligineis.

In silvis; typica forma in pinastretis ericetosis hinc inde, vulgo gregarius.

Descr. Durus, compactus, mitis. Stipes solidus, 3 unc. et ultra longus, ultra unc. crassus, bulbosus, immarginatus. Pileus carnosus, obtusus, 4—6 unc. latus, primo spadiceus,

dein fulvo-rubiginosus, disco obscuriore, margine (sæpe repando) tomentoso violaceo, rarius totus violaceus. Caro tam stipitis quam pilei e violaceo albescens, crassa, a stipite in pileum solito magis diffusa. Lamellæ tenues, $\frac{1}{2}$ unc. latæ, acie crenulatae.

Tab. 144 f. 1 formam *C. variicoloris* luxuriantem (*herculeanum Monogr. II p. 307) reddit. Lamellarum color pristina tollit dubia, quin PERSOONII denominatio merito huic speciei sit tribuenda.

CCCLXXXVIII. CORTINARIUS SPADICEUS (Batsch?)

Batsch f. 16, Fr. Monogr. II p. 14 et 307, Epicr. ed. II p. 339.

Diagn. Phlegmacium pileo carnosu, æquali, convexo-plano, laevi, viscoso, pellicula separabili, fuligineo-spadiceo; stipite fibrilloso-striato, pallescente, cortina supera; lamellis subemarginatis, vix confertis, cyaneis, dein cinnamomeo-brunneis.

Locus humidis pinetorum rario.

Descr. Stipes solidus, sursum attenuatus vel leviter clavato-bulbosus vel cylindricus, basi spongiosus et lutescens, 3 unc. circ. longus, $\frac{1}{2}$ unc. crassus, apice nudus; cortina

supera, fibrillosa, pendula, sporis ferruginea. Pileus obtusus, regularis, glaber, 2—3 unc. latus, margine demum striatulus. Caro albo-lutescens. Lamellæ 3 lin. latæ, haud ventricosæ, ex albo-cyaneo obscure cinnamomeo-brunneæ.

Colore tam pilei quam lamellarum obscuro formas *C. elatioris* admodum refert.

Tab. 144 f. 2 *C. spadicei* icon.

CCCLXXXIX. CORTINARIUS PERCOMIS Fr.

Epicr. ed. I p. 260, Monogr. II p. 12.

Diagn. Phlegmacium pileo carnosu, convexo-plano, æquali, glaberrimo, viscoso; stipite solido, clavato-attenuato, fibrilloso, supra cortinam superam pruinoso, intus sulphureo; lamellis late emarginatis, subconfertis, sulphureis, demum decoloratis.

In silvis acifoliis Sueciæ mediæ frequens; etiam in Vestrogothia, sed in Suecia australi nondum observatus.

Descr. Elegans species, recognitu facilis. Stipes valde compactus et firmus, 3 unc. et ultra longus, $\frac{1}{2}$ — $\frac{3}{4}$ unc. crassus, sulphureus, fibrillis adpressis obscurioribus subpe-

ronatus. Pileus valde carnosus, obtusus, 2—3 unc. latus, viscidus (acubus pini cet. adglutinatis), siccus pelliculosus, gilvus. Caro tam pilei quam stipitis sulphurea. Lamellæ 4 lin. usque latæ, primo sulphureæ, dein fulventes. — Seiore autumno (Oct.) legitur varietas pallidior, pileo alutaceo, stipite breviore, lamellis minus confertis.

Tab. 143 f. 2 formas elatiores *C. percomis* exhibet.

CCCXC. CORTINARIUS RAPACEUS Fr.

Epicr. ed. I p. 263, Monogr. II p. 16.

Diagn. Phlegmacium pileo compacto, aequali, convexo-plano, laevi, glabro, viscidulo; stipite farcto, curto, bulbo depresso marginato; lamellis rotundatis, confertis, integerimis, ex albo argillaceis.

In silvis abiegnis montanis Uplandiae, Vestrogothiae cet. passim.

Descr. Parvus, compactus. Stipes marginato-bulbosus, bulbo vulgo depresso, raro ovato, uncialis vel paullum ultra, $\frac{1}{2}$ unc. crassus, a bulbo leviter attenuatus, intus medullatus, glaber, apice vix pubescens, albus vel lutescens. Pileus

carnosus, obtusus, viscidulus, mox siccus et opacus, $1\frac{1}{2}$ —2 unc. latus, argillaceo-albus, non expallens, margine regulariter inflexo. Caro alba. Lamellae rotundatae absque denticulo, admodum tenues.

C. multififormi forte nimis affinis ejusque quasi forma curta, stipite medullato lamellisque integerrimis recedens.

Tab. 145 f. 1 forma major *C. rapacei*, ad iconem a LINDGREN missam delineata, a typica satis recedens.

CCCXCI. CORTINARIUS TALUS Fr.

Epicr. ed. I p. 263, Monogr. II p. 17.

Diagn. Phlegmacium pileo carnosus, convexo-plano, aequali, laevi, glabro, viscido; stipite solido, aequali, cylindrico, glabello, basi marginato-bulboso, una cum carne palecente; lamellis emarginatis, subconfertis, ochraceo-stramineis (vix decolorantibus).

In silvis mixtis pinetisque hinc inde, nullibi copiose.

Descr. Species pulchella, *C. turbinato* affinis et coloribus potissimum distincta. Stipes 3 unc. longus, $\frac{1}{2}$ unc.

crassus, bulbo parvo subrotundo. Pileus sat tenuis, 2—3 unc. latus, lutescenti-sordidus, expallens, margine subolivaceo lutescente. Caro aquosa, sordide albopallida, maculis hyalinis guttata et linea cornea sub lamellis variegata. Lamellae stramineae vel ochraceo-pallidæ.

Tab. 145 f. 2 *C. Talum*, Upsaliæ lectum, adumbrat.

CCCXCI. CORTINARIUS PANSA Fr.

Epicr. ed. I p. 264, Monogr. II p. 18.

Diagn. Phlegmacium pileo compacto, infracto, repando, glabro, squamis innatis guttato, nitido, carne alba; stipite obeso, aequali, basi marginato cortinaque inflexa luteis; lamellis rotundatis, subconfertis, integerrimis, cyaneis.

In silvis, præcipue fagetis, suis locis abunde.

Descr. Stipes curtus, validus, unciam et ultra crassus, raro (etiam senilis) longior, basi terræ immersus et marginatus sed non bulbosus, fibrillosus. Pileus valde carnosus, obtusissimus, semper undulato-repandus, junior infractus, dein dilatatus, 3 unc. circ. latus, vulgo squamulis guttatus et subinde floccis adspersus, laete aurantio-fulvus (non ex-

pallens decolorans), viscidus, siccus nitens, disco vulgo obscurior; junior circa marginem flavo-tomentosus. Caro admodum dura, crassa, persistenter alba. Lamellæ tenues, cyaneæ [ex tab. interdum senio decoloratae].

Species haud dubie distincta, *C. vario* forte proxima; a varietate subsimili *C. glaucopodis* caute separanda.

Tab. 145 f. 3 sistit *C. Pansa* formam, in Vestrogothia a LINDGREN depictam.

CCCXCIII. CORTINARIUS SCAURUS Fr.

Monogr. II. p. 25, Agaricus S. M. I. p. 223.

Diagn. Phlegmacium pileo tenuiter carnoso, æquali, glabro, tigrino-maculato, viscoso, expallente, margine tenui demum striatulo; stipite solido, a basi spongiosa marginato-bulbosa attenuato, striato, expallente; lamellis attenuato-adnatis, angustis, tenuibus, confertis, purpurascenti-olivaceis.

Locis humidis silvarum coniferarum, præcipue in montanis, præcox.

Descr. Stipes pro ratione gracilis, basi ultra unciam, apice 3—4 lin. crassus, mollis et spongiosus (præcipue bulbosus, qui vero passim evanescit), fibrillosus, e cœruleo vel

olivaceo albescens (raro tantum — in exoletis — lutescens). Cortina fibrillosa, subvirens. Pileus e convexo planus, senio etiam depresso, 2—3 (raro 4) unc. latus, colore peculiariter fulvo-fuligineo, sicci magis fulvente. Caro tenuis, aquosa, insipida. Lamellæ normaliter purpurascenti-olivaceæ, sed variant pure olivaceæ et rarissime fuligineæ.

Tab. 146 f. 1 *C. scauri* exhibet imaginem.

CCCXCIV. CORTINARIUS CUMATILIS Fr.

Epicr. ed. I p. 269, Monogr. II p. 27.

Diagn. Phlegmacium pileo convexo, obtuso, lœvi, glutine cœruleo violaceo-tineto; stipite solido, firmo, subbulboso, apice subcortinato carneque compacta candidis, velo universalis ad basin abrupto volvaceo; lamellis adnexis, confertis, serratis, ex albo argillaceis.

In dumetis mixtis (Fagi, Coryli) atque fagetis umbrosis raro, serotinus.

Descr. Speciosus, nobilis, alii vix affinis. Stipes 2—3 unc. longus, $\frac{1}{2}$ unc. crassus, sed varius et sœpe curvatus, omnino candidus, apice modo cortinatus sed non pulverulentus; velo universalis, pelliculam pilei præbente, ad basin abrupto, cui, membranae instar separabilis adglutinatae pileo

concoloris, est adnatum. Pileus non depresso, sœpe irregularis, 3 unc. latus, viscoso-pelliculosus, amoenissime violaceus vel purpureo-violascens. Caro tam pilei quam stipitis dura, candida, immutabilis. Lamellæ attenuato-adnexæ, fere liberæ, angustæ, vix 2 lin. latæ, denticulo decurrente, apice acutæ. — Variat nunc solitarius, major, compactus, nunc subcaespitosus, stipitibus æqualibus, adscendentibus.

Tab. 146 f. 2 fungum refert *C. cumatili* saltem valde similem, sed carne molli et recentem. Upsaliæ semel lectum [et jussu Patris defuncti h. I. non omittendum: editores].

CCCXCV. CORTINARIUS INTENTUS Fr.

Epicr. ed. I p. 272, Monogr. II p. 33.

Diagn. Phlegmacium pileo carnoso, convexo-plano, molli, lœvi, viscido; stipite cavo, rigido-fragili, obsolete fibrilloso, flavescente, apice nudo; lamellis adnatis, confertissimis, angustis, lœte croceis.

In pinetis rario.

Descr. Stipes mollis, intus araneoso-faretus, apice definite cavus, 3 unc. longus, 3—4 lin. crassus, æqualis, cylindricus vel sursum leviter attenuatus, fibrillosus, flavo-pallens, fere albidus. Pileus sat crassus, valde mollis, obtusus, 2—3 unc. latus, regularis, plus minus viscidus, lœvis et contiguus (numquam granuloso-ruptus), non hygrophila-

nus, siccus sericeo-lævigatus, e varia statione vel aetate testaceo-lutescens, ochraceo-luteus vel alutaceus. Caro mollis, alba. Lamellæ adnatæ, sœpe dente decurrentes, planæ, pertenues, 1—2 lin. latæ, lœte crocatæ (colore Croci stigmatum siccatorum) sive fere igneæ, quibus semel observatis illico agnoscitur. Sapor mitis. Sporæ ferrugineo-fulvæ.

Tab. 147 f. 1 *C. intentum* spectat. Icon. dirigente cl. LINDBLAD confecta, lamellis justo latioribus e. s. p. a genuino admodum discrepat.

CCCXCVI. CORTINARIUS OLIVASCENS (Batsch?)

Batsch f. 185, Fr. Epicr. ed. I p. 273, Monogr. II p. 35.

Diagn. Phlegmacium pileo subcarnoso, convexo-expanso, glabro, viscidulo, expallente, margine striatulo; stipite fareto, attenuato, tenui, argenteo-pallescens; lamellis adfixis, subconfertis, tenuibus, ex olivello-argillaceo cinnamomeis.

In silvis abiegnis montanis, locis humidis sphagnosis sparsus, rarus.

Descr. Stipes a basi aequaliter attenuatus, fibrillosus, 3 unc. circiter longus. Pileus convexo-planus, obtusus, $1\frac{1}{2}$

—2 unc. latus, subfuligineus, expallens, fere hygrophanus. Caro tenuis, dilutior nec alba. Lamellæ planæ, integerimæ, in optima forma olivellæ, in alia argillaceæ, demum subcinnamomeæ. Sapor acris.

Species in hac grege heteroclitæ, habitu *C. anomali*, affinitate forte *C. plumboso* proxima.

Tab. 147 f. 2 *C. olivascentis* adumbratio (H. v. Post).

CCCXCVII. CORTINARIUS SUBSIMILIS (Pers.)

Agaricus Pers. Syn. p. 284; Monogr. II p. 307, Epicr. ed. II p. 353.

Diagn. Phlegmacium pileo carnosulo, e convexo expanso, lœvi, viscoso, brunneo-fuligineo, carne alba; stipite solidō, aequali, fibrilloso, fulvello; lamellis adnatis, confertis, e fuligineo subcinnamomeis.

In silvis subhumidis raro.

Descr. Stipes 2—3 unc. longus, $\frac{1}{2}$ unc. crassus, fulvellus, basi ebulbis, albicans; cortina manifesta. Pileus

obtusus, 2—3 unc. latus, marginem versus tenuis. Lamellæ adnatae, 1—2 lin. latæ, haud ventricosæ.

Tab. 147 f. 3 ad *C. subsimilem* ducenda.

II. MYXACIUM. Pileus carnosus, subtenuis, viscosus. Stipes viscosus, vix bulbosus.

CCCXCVIII. CORTINARIUS MUCIFLUUS Fr.

Epicr. ed. I p. 274, Monogr. II p. 37.

Diagn. Myxacium pileo subcarnoso, campanulato-expanso, glutine hyalino secedente oblitito, margine striato; stipite deorsum attenuato, molli, e velo fugaci floccoso-squamoso viscido; lamellis adnatis, distinctis, ex argillaceo aquose cinnamomeis.

In pinetis arenosis vel ericetosis copiose.

Descr. Stipes spongiosus, albus vel rarius cœrulescens, 3—5 unc. longus, $\frac{1}{2}$ — $\frac{3}{4}$ unc. latus. Pileus sat tenuis,

margine membranaceo striato, 2—4 unc. latus, adultus vulgo revolutus, depressus et repandus, livido-argillaceus, siccus alutaceus opacus, glutine tenui hyalino diffluente nec pellucidam crassam persistentem læte coloratam sistente. Odor dulcis. — *C. collinito* admodum affinis.

Tab. 148 f. 1 *C. mucifluum* depingit.

CCCIC. CORTINARIUS ELATIOR Fr.

Epicr. ed. I p. 274, Monogr. II p. 37; Agaricus elatus Pers. Syn. p. 332, S. M. I p. 248 (non Batsch).

Diagn. Myxarium pileo e cylindrico expanso, viscido, disco carnosulo, lævi, ceterum membranaceo, plicato-rugoso; stipite elongato, molli, valido, utrinque attenuato, e velo diffracto squamoso; lamellis adnatis, perlatis, venoso-con-nexis rugosisque, brunneo-ferrugineis.

In pinetis abundans species; etiam in fagetis, betuletis vulgaris.

Descr. Stipes solidus, 5—8 unc. et ultra longus, $\frac{1}{2}$ —1, immo 2 unc. crassus, longitudinaliter fibrosus, squamis diruptis albidis subviscosis vestitus, raro violascens. Cor-

tina viscosa, fugax. Pileus e cylindrico vel bullato campanulatus, dein explanatus subreflexus, 3—4 unc. latus, fragilis, leviter viscosus, humidus vulgo e spadiceo-livido alutaceus, siccus sordide ochraceus; ceterum colore varius: albidus, alutaceo-fuscescens, spadiceus, violaceo-brunneus, niger, circa marginem albicans, griseus margine violaceo. Lamellæ primo 3 lin. latae, dein latiores (unc. usque!), subdistantes, semper obscure, rarius violaceo-brunneæ.

Tab. 149 f. 1 *C. elatiorem optimum representat.*

CCCC. CORTINARIUS EMUNCTUS Fr.

Epicr. ed. I p. 275, Monogr. II p. 39.

Diagn. Myxarium pileo leviter carnoso, campanulato-con-vexo, lævi, glabro, viscoso; stipite farcto, molli, attenuato, glutinoso, dilute violaceo-griseo, apice nudo, albo; lamellis adnatis, subdistantibus, latis, ex argillaceo aquose cinnamomeis.

In pinetis. Juxta Upsaliam unico loco quotannis non raro.

Descr. Stipes farctus, demum cavus, 2—3 unc. et ultra longus, 3—4 lin. crassus, non raro gracilescens, 4—5-uncialis et flexuosus, extus cartilagineus, rigidus sed undique glutinoso-velatus, cortina annulari fibrillosa. Pileus obsolete gibbus,

disciformis, 1—2 unc. latus, colore peculiariter lurido, junioris griseo-violaceo, adulti in disco lutescente, demum toto griseo-alutaceo; caro tenuis, mollis, alba, demum leviter lutescens. Lamellæ subdecurrentes, minus confertæ, 3—4 lin. latae, postice connexæ, acie integerrima concolore. — Delicata species, colore inter affines inconsuetu insignis.

Tab. 148 f. 2 in *C. emunctum* ad unguem quadrat.

CCCCI. CORTINARIUS LIQUIDUS Fr.

Epicr. ed. I p. 257, Monogr. II p. 39.

Diagn. Myxarium pileo carnosulo, convexo-plano, lævi, glutinoso, sicco sericeo albicante; stipite fistuloso, attenuato, glutinoso, albo; lamellis adnato-decurrentibus, distantibus, serratis, subargillaceis.

Locis paludosis exsiccatis, inter muscos et folia coacer-vata putrida (*Fagi, Salicis auritæ* cet.).

Descr. Distinctissimus. Stipes elongatus, 3 unc. cir-citer longus, gracilis, saepe decumbens, a basi incrassata albovilloso sursum valde attenuatus, lævis, glaber, nunc rigido-fragilis, nunc mollis; cortina fibrilloso-viscida. Pileus

e convexo planus, umbone obtuso vel obsoleto, circa quem demum depresso, 1— $1\frac{1}{2}$ unc. latus, fragilis; vegetus valde glutinosus muco defluente, pallide flavus; caro aquosa, non scissilis. — In forma rigida pileus omnino exstrius, in molli, minore discus modo carnosus, pileus ceterum tenuissimus, remote striatus et margine crenulatus. Lamellæ 2—3 lin. latae, planæ, nunc distinctæ, nunc venoso-connexæ et ad latera venosæ.

Tab. 149 f. 2 *C. liquidum* illustrat.

CCCCII. CORTINARIUS SALOR Fr.

Epicr. ed. I p. 276, Monogr. II p. 40.

Diagn. Myxarium pileo conico-campanulato, viscoso, dein expanso, disco compacto gibboso, margine tenui innato-fibrilloso violaceo; stipite solido, bulboso, conico-attenuato, glabro, velo glutinoso cæruleo ad apicem tecto; lamellis adnatis, distantibus, integerrimis, griseo-argillaceis, margine violaceis.

In nemoribus frondosis (*Coryli* cet.), serotinus.

Descr. Valde nobilis, firmus. Stipes $1\frac{1}{2}$ —3 unc. longus, basi 1, apice $\frac{1}{2}$, unc. crassus, deorsim sensim elongatus — et quo magis elongatus, eo diminutior bulbus —, albus, sed junior a velo mere viscido cærulescens, vetustus expallens, mollior; cortina subapicalis, fugax. Pileus ju-

nior obtuse conicus vel parabolicus, mox campanulatus demumque explanatus, umbone lato ob discum compacte carnosum, ratione stipitis parvus, $\frac{1}{2}$ —2 unc. circ. latus, griseus, circa marginem inflexum læte violascens, demum vero concolor, tenuiter viscosus, exsiccatus marginem versus fibrillosus. Caro contigua, alba, lutescens [vel dilute cærulescens]. Lamellæ 2—3 lin. latae, primo dilute griseæ acie violacea cæsiave, adultæ griseo-argillaceæ vel demum cinnamomeæ.

Tab. 150 f. 1 est *C. Salor*, a def. LINDGREN in Vestrogothia pictus.

CCCCIII. CORTINARIUS NÆVOSUS Fr.

Monogr. II p. 41, Epicr. ed. II p. 357.

Diagn. Myxarium pileo carnoso, convexo-plano, subdepresso, helvolo-luteo, squamis maculæformibus minutis confertis ferrugineo-brunneis tigrino; stipite solido, firmo, glabro, cæsio-pallido, glutine hyalino-lutescente collinito; lamellis emarginatis, latis, confertis, e cyaneo argillaceis.

Hactenus tantum in lucis montis Kinnekulle Sueciæ a def. LINDGREN lectus; ipse numquam vidi.

Descr. Stipes primitus curtus, ventricosus, dein 2—3 unc. longus, unciam crassus, sursum attenuatus, apice cortina e sporis cinnamomea annulatus. Pileus obtusus, 2 unc. latus, carne pallida. Lamellæ pulchre emarginatæ, latae (3 lin. et ultra).

Tab. 150 f. 2 LINDGREN reddit icones *C. nævosi*.

CCCCIV. CORTINARIUS EPIPOLEUS Fr.

Epicr. ed. I p. 277, Monogr. II p. 42.

Diagn. Myxarium pileo tenui, convexo-plano, gibbo, laevi, cano-micante, disco carnoso stipiteque farcto, bulboso-clavato, obsolete violaceo viscosis; lamellis decurrentibus, apice repandis, cinnamomeis.

In silvis Sueciæ mediæ rario, insignis.

Descr. Stipes elasticus, intus spongiosus albidus, demum pertusus fragilis, 3 unc. longus, basi ultra $\frac{1}{2}$ unc.,

apice 3—4 lin. crassus; cortina fugax. Pileus disco excepto tenuis, submembranaceus (inter pelliculosos et scissiles ambiguus), 2—3 unc. latus, lente viscidus, glaber quidem sed rore peculiaris cano innato quasi sericeus, opacus. Lamellæ 3—4 lin. latæ, postice latiusculæ, constanter cinnamomeæ.

Tab. 150 f. 3 *C. epipoleum* imitatur.

1. *CORTINARIUS (PHLEGMIACUM) TRIUMPHANS*. FR. 2. *CORTINARIUS (PHLEGM.) CLARICOLOR* FR.



1. *CORTINARIUS (PHLEGMACIUM) CRASSUS. FR.* 2. *CORTINARIUS (PHLEGMACIUM) BALTEATUS. FR.*

M. A. Lindblad direx

Central-Tryckeriet Stockholm.

P. Åkerblad et Lindgren dell.

1. *CORTINARIUS (PHLEGMACIUM) SEBACEUS* FR. 2. *CORTINARIUS (PHLEGMACIUM) PERCOMIS*. FR.



1. *CORTINARIUS (PHLEGMACIUM) VARITCOLOR (PERS.)* 2. *CORTINARIUS (PHLEG.) SPADICEUS (BATSCH?)*

E.Fries direx

Central-Tryckeriet Stockholm.

P Åkerlund dell.



1. *CORTINARIUS (PHLEGMACIUM) RAPACEUS*. FR. 2. *CORTINARIUS TALUS*. FR.

E. Fries direx

3. *CORTINARIUS (PHLEGMACIUM) PANSA*. FR.

Lindgren, P Åkerblad et E Pettersson dell.

Central-Tryckeriet Stockholm.



1. *CORTINARIUS (PHLEGMACIUM) SCAURUS*. FR. 2. *CORTINARIUS (PHLEGM.) CUMATILIS*. FR.

E Fries direx

Central-Tryckeriet Stockholm

P Åkerlund et E Pettersson dell



1. *CORTINARIUS (PHLEGMACIUM) INTENTUS*. FR. 2. *CORTINARIUS (PHLEGMI.) OLIVASCENS.* (BATSCH?)

E. Fries direx

3. *CORTINARIUS (PHLEGMACIUM) SUBSIMILIS.* (PERS.)

P Åkerblad v Post et E Pettersson dell



1. *CORTINARIUS (MYXACIUM) MUCIFLUUS*. FR. 2. *CORTINARIUS (MYXACIUM) EMUNCTUS*. FR.



1. *CORTINARIUS (MYXACIUM) ELATIOR. FR.* 2. *CORTINARIUS (MYXACIUM) LIQUIDUS. FR.*

E.Fries direx

Central-Tryckeriet Stockholm.

E.Pettersson v Positum



1. CORTINARIUS (MYXACIUM) SALOR. FR. 2. CORTINARIUS (MYXACIUM) NAEVOSUS. FR.

E Fries direx

3. CORTINARIUS (MYXACIUM) EPIPOLEUS. FR.

Lindgren et E Pettersson dell.