

Identification and nomenclature of the genus Penicillium

Visagie, C.M.; Houbraken, J.; Frisvad, Jens Christian; Hong, S. B.; Klaassen, C.H.W.; Perrone, G.; Seifert, K.A.; Varga, J.; Yaguchi, T.; Samson, R.A.

Published in:
Studies in Mycology

Link to article, DOI:
[10.1016/j.simyco.2014.09.001](https://doi.org/10.1016/j.simyco.2014.09.001)

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Visagie, C. M., Houbraken, J., Frisvad, J. C., Hong, S. B., Klaassen, C. H. W., Perrone, G., ... Samson, R. A. (2014). Identification and nomenclature of the genus Penicillium. *Studies in Mycology*, 78, 343-371. DOI: [10.1016/j.simyco.2014.09.001](https://doi.org/10.1016/j.simyco.2014.09.001)

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Identification and nomenclature of the genus *Penicillium*

C.M. Visagie¹, J. Houbraken^{1*}, J.C. Frisvad^{2*}, S.-B. Hong³, C.H.W. Klaassen⁴, G. Perrone⁵, K.A. Seifert⁶, J. Varga⁷, T. Yaguchi⁸, and R.A. Samson¹

¹CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, NL-3584 CT Utrecht, The Netherlands; ²Department of Systems Biology, Building 221, Technical University of Denmark, DK-2800 Kgs. Lyngby, Denmark; ³Korean Agricultural Culture Collection, National Academy of Agricultural Science, RDA, Suwon, Korea; ⁴Medical Microbiology & Infectious Diseases, C70 Canisius Wilhelmina Hospital, 532 SZ Nijmegen, The Netherlands; ⁵Institute of Sciences of Food Production, National Research Council, Via Amendola 122/O, 70126 Bari, Italy; ⁶Biodiversity (Mycology), Agriculture and Agri-Food Canada, Ottawa, ON K1A0C6, Canada; ⁷Department of Microbiology, Faculty of Science and Informatics, University of Szeged, H-6726 Szeged, Közép fasor 52, Hungary; ⁸Medical Mycology Research Center, Chiba University, 1-8-1 Inohana, Chuo-ku, Chiba 260-8673, Japan

*Correspondence: J. Houbraken, j.houbraken@cbs.knaw.nl; J.C. Frisvad, jcf@bio.dtu.dk

Abstract: *Penicillium* is a diverse genus occurring worldwide and its species play important roles as decomposers of organic materials and cause destructive rots in the food industry where they produce a wide range of mycotoxins. Other species are considered enzyme factories or are common indoor air allergens. Although DNA sequences are essential for robust identification of *Penicillium* species, there is currently no comprehensive, verified reference database for the genus. To coincide with the move to one fungus one name in the International Code of Nomenclature for algae, fungi and plants, the generic concept of *Penicillium* was re-defined to accommodate species from other genera, such as *Chromocleista*, *Eladia*, *Eupenicillium*, *Torulomyces* and *Thysanophora*, which together comprise a large monophyletic clade. As a result of this, and the many new species described in recent years, it was necessary to update the list of accepted species in *Penicillium*. The genus currently contains 354 accepted species, including new combinations for *Aspergillus crystallinus*, *A. malodoratus* and *A. paradoxus*, which belong to *Penicillium* section *Paradoxa*. To add to the taxonomic value of the list, we also provide information on each accepted species MycoBank number, living ex-type strains and provide GenBank accession numbers to ITS, β-tubulin, calmodulin and *RPB2* sequences, thereby supplying a verified set of sequences for each species of the genus. In addition to the nomenclatural list, we recommend a standard working method for species descriptions and identifications to be adopted by laboratories working on this genus.

Key words: Aspergillaceae, Fungal identification, phylogeny, media, nomenclature.

Taxonomic novelties: New combinations: *Penicillium crystallinum* (Kwon-Chung & Fennell) Samson, Houbraken, Visagie & Frisvad, *Penicillium malodoratum* (Kwon-Chung & Fennell) Samson, Houbraken, Visagie & Frisvad, *Penicillium paradoxum* (Fennell & Raper) Samson, Houbraken, Visagie & Frisvad.

Published online 22 September 2014; <http://dx.doi.org/10.1016/j.simyco.2014.09.001>. Hard copy: June 2014.

INTRODUCTION

Penicillium is well known and one of the most common fungi occurring in a diverse range of habitats, from soil to vegetation to air, indoor environments and various food products. It has a worldwide distribution and a large economic impact on human life. Its main function in nature is the decomposition of organic materials, where species cause devastating rots as pre- and postharvest pathogens on food crops (Frisvad & Samson 2004, Pitt & Hocking 2009, Samson *et al.* 2010), as well as producing a diverse range of mycotoxins (Frisvad *et al.* 2004). Some species also have positive impacts, with the food industry exploiting some species for the production of speciality cheeses, such as Camembert or Roquefort (Thom 1906, Nelson 1970, Karahadian *et al.* 1985, Giraud *et al.* 2010) and fermented sausages (López-Díaz *et al.* 2001, Ludemann *et al.* 2010). Their degradative ability has resulted in species being screened for the production of novel enzymes (Raper & Thom 1949, Li *et al.* 2007, Adsul *et al.* 2007, Terrasan *et al.* 2010). Its biggest impact and claim to fame is the production of penicillin, which revolutionised medical approaches to treating bacterial diseases (Fleming 1929, Chain *et al.* 1940, Abraham *et al.* 1941, Thom 1945). Many other extrolites have since been discovered that are used for a wide range of applications (Frisvad *et al.* 2004). Pitt (1979) considered

it axiomatic that *Penicillium* or one of its products has affected every modern human.

It is now more than 200 years since Link (1809) introduced the generic name *Penicillium*, meaning ‘brush’, and described the three species *P. candidum*, *P. glaucum* and the generic type *P. expansum*. Since then, more than 1000 names were introduced in the genus. Many of these names are not recognisable today because descriptions were incomplete by modern criteria. Some names were published invalidly, or are now considered synonyms of other species. Thom (1930) revised all species described until 1930 and accepted 300 species. In later studies, Raper & Thom (1949) accepted 137 species, Pitt (1979) accepted 150 species, and Ramírez (1982) accepted 252 species (numbers include species described in *Eupenicillium*). At that time, a morphological species concept was used for *Penicillium* classification and identification, with DNA sequencing starting to be used during the 1990’s. DNA sequencing created the threat that old names previously considered of uncertain application, because their ex-type cultures were no longer morphologically representative, could replace more commonly used but younger names. As such, the List of “Names in Current Use” (NCU) for the family Trichocomaceae (Pitt & Samson 1993) accepted 223 species and disregarded all other names as if not published. This list was updated by Pitt *et al.* (2000) who

accepted 225 species. Species names not accepted on these lists were not to be disregarded permanently, as stated by Pitt et al. (2000), because they were not formally rejected under the nomenclatural code and could still be reintroduced in a revised taxonomy. In fact, this became common practice as many old species were shown to be distinct and were reintroduced (Peterson et al. 2005, Serra et al. 2008, Houbraken et al. 2011a,b, Houbraken et al. 2012a, Visagie et al. 2013).

The abandonment of article 59 in the new International Code of Nomenclature for algae, fungi and plants (ICN) (McNeill et al. 2012) resulted in single name nomenclature for fungi. In anticipation of this change, Houbraken & Samson (2011) redefined the genera in the family Trichocomaceae based on a four gene phylogeny. They segregated the Trichocomaceae into three families, namely the Aspergillaceae (*Aspergillus*, *Hamigera*, *Leiothecium*, *Monascus*, *Penicilliopsis*, *Penicillium*, *Phialomyces*, *Sclerocheista*, *Warcupiella*, *Xeromyces*), Thermoascaceae (*Byssochlamys*/*Paecilomyces*, *Thermoascus*) and the Trichocomaceae (*Rasamonia*, *Sagenomella*, *Talaromyces*, *Thermoascaceae*, *Trichocoma*). *Penicillium* subgenus *Biverticillium* and *Talaromyces* were shown to form a monophyletic clade distinct from the other subgenera of *Penicillium*, with these names recombined as necessary into *Talaromyces* (Samson et al. 2011). The remaining *Penicillium* species formed a monophyletic clade together with species classified in *Eupenicillium*, *Eladia*, *Hemicarpenteles*, *Torulomyces*, *Thysanophora* and *Chromocleista*. These generic names were synonymised with *Penicillium*, while their species were given *Penicillium* names (Houbraken & Samson 2011). The remaining three *Aspergillus* species, *A. paradoxus* (\equiv *Hemicarpenteles paradoxus*), *A. malodoratus* and *A. crystallinus*, phylogenetically belonging in *Penicillium*, are transferred to *Penicillium* below in the Taxonomy section. To accommodate the morphological variation, the generic diagnosis of *Penicillium* was amended in Houbraken & Samson (2011). Most importantly, in comparison with the prevailing generic concept (Raper & Thom 1949, Pitt 1979), it now excludes the acerose phialides and usually symmetrically branched conidiophores of species now included in *Talaromyces*, and was expanded to include the conidiophores with solitary phialides of species in section *Torulomyces*, and the darkly pigmented stipes that formerly characterised the genus *Thysanophora*, which show secondary growth by means of the proliferation of an apical penicillus. For infrageneric classification, the genus was divided into two subgenera, *Aspergilloides* and *Penicillium*, and 25 sections.

Thom (1954) attempted to explicitly define species concepts used for *Penicillium*. He was the pioneer of standardised working techniques and emphasised that *Penicillium* taxonomy demands a consistent, logical approach. He demonstrated these tendencies himself by taking into account infraspecies variation when delineating species. To minimise infraspecies variation, the importance of standardised working techniques were again emphasised by Pitt (1979), Samson & Pitt (1985), Okuda (1994) and Okuda et al. (2000). Although this was relatively effective when dealing with freshly isolated or wild-type strains, comparing strains using only morphology requires experience and nuance, because of the degeneration of characters in old reference material and the large number of species in the genus. New techniques incorporated into taxonomic studies resulted in the physiological species concept (Ciegler & Pitt 1970, Pitt 1973, Frisvad 1981, Frisvad & Filtenborg 1983, Cruickshank & Pitt 1987a,b, El-Banna et al. 1987, Frisvad & Filtenborg 1989, Paterson et al. 1989), phylogenetic species

concept, including Genealogical Concordance Phylogenetic Species Recognition (GCPSR) (LoBuglio et al. 1993, Berbee et al. 1995, Boysen et al. 1996, Geiser et al. 1998, Skouboe et al. 1999, O'Donnell et al. 1998, Peterson 2000a, Taylor et al. 2000) and eventually led to the combined approach using morphological, extrolite and genetic data in a polyphasic species concept (Christensen et al. 2000, Frisvad & Samson 2004). In the modern taxonomy, however, sequence data and GCPSR carries more weight than morphology or extrolite data.

Even though *Penicillium* species are very common and the taxonomic structure of the genus is well defined, species identification is still problematic. Problems include an out-dated accepted species list and a lack of a verified, complete sequence database. The aim of this paper is to address these issues. In honour of Charles Thom and Kenneth B. Raper, the fathers of *Penicillium* taxonomy, and all the other mycologists who have contributed in the last 200 years towards our current understanding of the genus, we offer this contribution towards a practical, stable, consistent, logical approach to *Penicillium* taxonomy, identification and nomenclature.

RECOMMENDED METHODS FOR IDENTIFICATION AND CHARACTERISATION OF *PENICILLIUM*

Morphological species recognition

Morphology in the past has been central to the taxonomy of *Penicillium* and along with multigene phylogenetics and extrolite profiling comprises the polyphasic species concept adopted for *Penicillium*. Morphology is the physical architecture through which an organism functions in and adapts to its environment, but some aspects may vary or be induced by specific cues in the immediate environment. As a result, strains characterised in one laboratory might look different when grown in another because of subtle differences in nutrients, temperature, lighting or humidity. This sometimes makes comparisons between different studies very difficult. These effects can be minimised using strictly standardised working techniques for medium preparation, inoculation technique and incubation conditions (Samson & Pitt 1985, Okuda 1994, Okuda et al. 2000). We recommend the following standardised methods for laboratories identifying and describing *Penicillium* species (summarised in Fig. 1).

Macromorphology

Colony characters and diameters on specific media are important features for species identification. Czapek Yeast Autolysate agar (CYA) and Malt Extract agar (MEA, Oxoid) is recommended as standard media for *Penicillium*. Even though malt extract from Oxoid is recommended, many laboratories prefer Difco. It should be noted that two different MEA formulations are widely used in modern taxonomic studies. Blakeslee's MEA was historically widely used, but lately CBS switched to a different formulation (given in Table 1). Both are suitable for characterisation, but studies should state which MEA (Oxoid or Difco) and formulation was used. The following alternative media can be used for observing additional taxonomic characters: Czapek's agar (CZ), Yeast Extract Sucrose agar (YES), Oatmeal agar (OA), Creatine Sucrose agar (CREA), Dichloran 18 % Glycerol agar (DG18), Blakeslee's MEA and CYA with 5 % NaCl (CYAS). CZ was used

Table 1. Media and stock solutions used for morphological characterisation.**Czapek stock solution (100 ml) (Pitt 1979)**

NaNO ₃	30 g
KCl	5 g
MgSO ₄ ·7H ₂ O	5 g
FeSO ₄ ·7H ₂ O	0.1 g
dH ₂ O	100 ml

*Store at 4–10 °C

Trace elements stock solution (100 ml)

CuSO ₄ ·5H ₂ O	0.5 g
ZnSO ₄ ·7H ₂ O	0.1 g
dH ₂ O	100 ml

*Store at 4–10 °C

Blakeslee's Malt extract agar (MEA) (Blakeslee 1915)

Malt extract (Oxoid)	20 g
Peptone (Oxoid)	1 g
Glucose	20 g
Trace elements	1 ml
Agar	20 g
dH ₂ O	1000 ml

*Mix well and autoclave at 121 °C for 15 min. pH 5.3 ± 0.2.

Creatine sucrose agar (CREA) (Frisvad 1981)

Sucrose	30 g
Creatine·H ₂ O	3 g
K ₃ PO ₄ ·7H ₂ O	1.6 g
MgSO ₄ ·7H ₂ O	0.5 g
KCl	0.5 g
FeSO ₄ ·7H ₂ O	0.01 g
Trace elements stock solution	1 ml
Bromocresol purple	0.05 g
Agar	20 g
dH ₂ O	1000 ml

*Mix well and autoclave at 121 °C for 15 min. pH 8.0 ± 0.2.

Czapek's agar (CZ) (Raper & Thom 1949)

Czapek concentrate	10 ml
Sucrose	30 g
Trace elements stock solution	1 ml
Agar	20 g
dH ₂ O	1000 ml

*Mix well and autoclave at 121 °C for 15 min.

Czapek Yeast Autolysate agar (CYA) (Pitt 1979)

Czapek concentrate	10 ml
Sucrose	30 g
Yeast extract (Difco)	5 g
K ₂ HPO ₄	1 g
Trace elements stock solution	1 ml
Agar	20 g
dH ₂ O	1000 ml

*Mix well and autoclave at 121 °C for 15 min. pH 6.2 ± 0.2.

Table 1. (Continued).**Czapek Yeast Autolysate agar with 5 % NaCl (CYAS)**

Czapek concentrate	10 ml
Sucrose	30 g
Yeast extract (Difco)	5 g
K ₂ HPO ₄	1 g
Trace elements stock solution	1 ml
NaCl	50 g
Agar	20 g
dH ₂ O	1000 ml

*Mix well and autoclave at 121 °C for 15 min. pH 6.2 ± 0.2.

Dichloran 18 % Glycerol agar (DG18) (Hocking & Pitt 1980)

Dichloran-Glycerol-agar-base (Oxoid)	31.5 g
Glycerol (anhydrous)	220 g
Trace elements stock solution	1 ml
Chloramphenicol	0.05 g
Agar	20 g
dH ₂ O	1000 ml

*Mix well and autoclave at 121 °C for 15 min. After autoclaving, add 0.05 chlortetracycline. pH 5.6 ± 0.2.

Malt Extract agar (MEA) (Samson et al. 2010)

Malt extract (Oxoid CM0059)	50 g
Trace elements stock solution	1 ml
dH ₂ O	1000 ml

*Mix well and autoclave at 115 °C for 10 min. pH 5.4 ± 0.2.

Oatmeal agar (OA) (Samson et al. 2010)

Oatmeal / flakes	30 g
Trace elements stock solution	1 ml
Agar	20 g
dH ₂ O	1000 ml

*First autoclave flakes (121 °C for 15 min) in 1000 ml dH₂O. Squeeze mixture through cheese cloth and use flow through, topping up to 1000 ml with dH₂O with 20 g agar. Autoclave at 121 °C for 15 min. pH 6.5 ± 0.2.**Yeast extract sucrose agar (YES)** (Frisvad 1981)

Yeast extract (Difco)	20 g
Sucrose	150 g
MgSO ₄ ·7H ₂ O	0.5 g
Trace elements stock solution	1 ml
Agar	20 g
dH ₂ O	885 ml

*Mix well and autoclave at 121 °C for 15 min. pH 6.5 ± 0.2.

in the taxonomic treatments of Raper & Thom (1949) and Ramírez (1982) and is chemically well defined. However, it is not widely used for *Penicillium* studies. YES is the recommended medium for extrolite profiling of species. Sexual reproduction most commonly occurs when strains are grown on OA and thus often provide valuable taxonomic information. OA should be prepared using organic uncooked flakes and not the unsuitable quick cook oats ("3 minute oats") or prefabricated OA formulations available. Acid production is observed by the colour reaction in CREA (from purple to yellow) and is often useful for distinguishing between closely related species. In some species acid production is followed by base production and is most often observed from the colony reverse. DG18 and CYAS often provide useful information with regards to growth rates on low water activity media. For consistent conidial colours, the addition of zinc-sulphate and copper-sulphate as trace elements (1 g ZnSO₄.7H₂O and 0.5 g CuSO₄.5H₂O in 100 ml distilled water) is of utmost importance because these metals vary widely in water sources in different locations and are critical for pigment production. For inorganic chemicals, analytical grade should be used. Experience has shown that the brand of agar used influences colony appearance. As such, it is important to test the agar for consistent character development and note the brand in species descriptions. Even though we do not recommend a brand here as standard, after extensive comparisons CBS opted to use So-BI-Gel agar (Bie & Berntsen, BBB 100030) for medium preparations. Medium formulations are given in Table 1.

Media are prepared in 90 mm Petri-dishes with a volume of 20 ml (Okuda et al. 2000). Glass Petri dishes were traditionally considered best for character observation. However, today it is not feasible using them, with polystyrene Petri dishes recommended. The Petri dishes should preferably be vented which allows for a bit of air exchange. Inoculations are made from spore suspensions in a semi-solid agar solution containing 0.2 % agar and 0.05 % Tween80 (Pitt 1979). In some laboratories, spore suspensions are made in a 30 % glycerol, 0.05 % agar and 0.05 % Tween80 solution. Variation between these two inoculation suspension solutions has not been shown. We recommend using a micropipette for inoculating the spore suspension in three-point fashion (0.5–1 µl per spot). They should not be wrapped with Parafilm, which by restricting air exchange often inhibits growth and sporulation (Okuda et al. 2000). When using walk-in incubators, laboratory rules often require plates to be placed in plastic containers or plastic bags. In this case, the boxes or bags should allow for enough aeration and care should be taken not to have too many plates in one box or in one incubator. For standard bench-top incubators, plates do not need to be incubated in boxes or bags, unless there is a strong air-current in the incubator. All media are incubated at the standard temperature of 25 °C for 7 d, with additional CYA plates at 30 and 37 °C that are useful to distinguish between species. It is crucial that temperatures are carefully checked as small differences have a large impact on colony growth. No effect on colonies has been shown for incubating Petri dishes upside down or vice versa.

After 7 d, colony diameters are measured at the widest part of the colony. Important characters used for describing *Penicillium* include colony texture, degree of sporulation, the colour of conidia, the abundance, texture and colour of mycelia, the presence and colours of soluble pigments and exudates, colony reverse colours, and degree of growth and acid production (in some species acid production followed by base production) on CREA. The use of a standard colour chart is recommended for

descriptions. Although many are available, the most commonly used colour chart in *Penicillium* is the Methuen Handbook of Color (Kornerup & Wanscher 1967), which although long out of print is available in many libraries. It is worth noting that colour names used by Raper & Thom (1949) were based on Ridgeway (1912) and are also still in use. When colour charts are not available, it is recommended to publish full color photoplates to accompany descriptions of new species. Some species produce specialised structures such as cleistothecia or sclerotia, mostly after longer incubation times, especially on OA. We thus recommend OA plates be incubated for prolonged periods to have a chance of observing these structures. For definitions and explanations of terms used in descriptions of *Penicillium* colonies, readers are referred to Frisvad & Samson (2004).

Micromorphology

Conidiophore and cleistothecium (when produced) characters of *Penicillium* are of great taxonomic importance. Conidiophore branching patterns (Fig. 2) were traditionally used in the classification of *Penicillium* (Thom 1930, Raper & Thom 1949, Pitt 1979). Although these branching patterns do not correspond perfectly with the sections currently accepted for *Penicillium*, characterising them accurately is still considered important. The conidiophores range from being simple (solitary phialides, Fig. 2A) to very complex patterns with multiple levels of branching resulting in overall symmetrical or asymmetrical patterns. Monoverticillate conidiophores (Fig. 2B) have a terminal whorl of phialides and in some species, the terminal cell of the conidiophore is slightly swollen or vesiculate; such species could be confused with diminutive *Aspergillus* conidiophores, but they have septa in the stipes unlike species of the latter genus. Divaricate conidiophores, previously also referred to as irregular (Pitt 1979, Fig. 2C), are best described as having a simple to complex branching pattern with numerous subterminal branches formed, but where conidiophore parts are divergent. Biverticillate conidiophores (Fig. 2D, E) have a whorl of three or more metulae between the end of the stipe and the phialides; the metulae may be of unequal or equal length, vary in their degree of divergence, are usually more or less cylindrical but can also be clavate or slightly vesiculate. Terverticillate conidiophores (Fig. 2F) have another level of branching between the stipe and the metulae, often just a continuation of the stipe axis and one side branch, sometimes a true whorl of three or more branches. Quaterverticillate conidiophores (Fig. 2G) are produced by only a few species, and have one extra level of branching beyond the terverticillate pattern. Terverticillate and quaterverticillate conidiophores tend to be conspicuously asymmetrical. In colonies of many species, especially as cultures begin to degenerate, there may be more than one branching pattern or intermediate forms, and it can be challenging to decide which pattern is typical or most developed.

Other important microscopic characters include the wall texture/ornamentation of stipes and conidia, as well as dimensions, ornamentation and sometimes colours of all elements of the conidiophore (Fig. 1). In our experience, wall textures are very sensitive to minute differences in media composition and aeration. For best observation of conidial ornamentation, differential interference contrast (= Nomarski) is recommended if possible; ornamentation is sometimes most conspicuously visible in air pockets in the preparation.

For microscopic observations, slides are prepared from 7–10 d old MEA colonies. Remove material from the zones where adjacent colonies are closest or growing together, from




Fig. 1. Flow diagram summarising recommended methods for the identification and characterisation of *Penicillium*. Frisvad & Thrane (1987, 1993), Smedsgaard (1997) and Klitgaard et al. (2014), refer to methods described for detecting extrolites in fungi. Lund (1995) introduced the Ehrlich reaction that tests for production of indole metabolites.




Fig. 2. Conidiophore branching patterns observed in *Penicillium*. A. Conidiophores with solitary phialides. B. Monoverticillate. C. Divaricate. D, E. Biverticillate. F. Terverticillate. G. Quaterverticillate, terms used for describing parts of conidiophores are given. Scale bar = 10 µm.

the part of the colony where the conidial colour is just starting to develop. For observing conidia, material removed from colony centers generally gives more uniform results. Lactic acid (60 %) is the most commonly used mounting fluid, although other solutions such as Shear's solution or lactic acid with cotton blue (Frisvad & Samson 2004, Samson et al. 2010) can also be used. We do not recommend lactophenol because of the corrosiveness and toxicity of phenol. Because of the abundant hydrophobic conidia produced by most species, drops of 70 % ethanol are commonly used to wash away excess conidia and to prevent air being "trapped" when mounted in lactic acid. For photographing conidiophores, we often wash away the spores two or three

times. Some species have dense colonies and then it is necessary to tease apart the conidiophores with very fine needles under the dissecting microscope.

DNA barcoding for identification and multilocus sequence typing for phylogenetic species recognition

Sequence markers

During the 1990's, DNA sequencing became one of the most powerful tools for taxonomists, because it created the opportunity for inferring relationships between species without the need for

standardising culturing regimes and eliminated problems related to deteriorated cultures. It also created the opportunity for sequence based identifications. DNA barcoding was launched to make species identification of any eukaryotic organism possible for anybody, by using a standardised short DNA sequence and a curated reference database linked to authoritatively identified vouchers (Blaxter 2003, Tautz *et al.* 2003, Hebert *et al.* 2003, Blaxter *et al.* 2005, DeSalle *et al.* 2005, Rathnasingham & Hebert 2007, Seifert *et al.* 2007, Min & Hickey 2007a,b, Schoch *et al.* 2012). Only recently however, was the internal transcribed spacer rDNA area (ITS) accepted as the official barcode for fungi (Schoch *et al.* 2012). ITS is the most widely sequenced marker for fungi, and universal primers are available (Schoch *et al.* 2012). In *Penicillium*, it works well for placing strains into a species complex or one of the 25 sections, and sometimes provides a species identification (Visagie *et al.* 2014a). Unfortunately, for *Penicillium* and many other genera of ascomycetes, the ITS is not variable enough for distinguishing all closely related species (Skouboe *et al.* 1999, Seifert *et al.* 2007, Schoch *et al.* 2012). The open source sequence repository GenBank contains a large proportion of incorrectly identified sequences, making identifications of *Penicillium* using BLAST very tricky for inexperienced workers. This particular problem is addressed in a number of publications (Köljalg *et al.* 2005, Santamaría *et al.* 2012, Köljalg *et al.* 2013, Schoch *et al.* 2014). For *Penicillium*, the International Commission of *Penicillium* and *Aspergillus* (ICPA), in conjunction with the publication of an updated accepted species list presented below, decided to include GenBank accession numbers to reference barcode sequences for each species when available.

Because of the limitations associated with ITS as a species marker in *Penicillium*, a secondary barcode or identification marker is often needed for identifying isolates to species level. The requirements for a secondary identification marker are clear. It should be easy to amplify, distinguish among closely related species and most importantly, the reference data set should be complete, meaning that there should be representative sequences for all species. It would be an added bonus if this marker is useful for phylogenetic studies, as it will by default become the gene most widely sequenced in future. Based on these criteria, we propose the use of β -tubulin (*BenA*) as the best option for a secondary identification marker for *Penicillium*. *BenA* does, however, have problems associated with it. Although not influencing BLAST identifications, alignments across a diverse genus like *Penicillium* is difficult and often contains a large proportion ambiguously aligned sites, which can make phylogenies difficult. Also, there is evidence for the amplification of *BenA* paralogous genes in *Aspergillus* (Peterson 2008, Hubka & Kolarik 2012) and *Talaromyces* (Peterson & Jurjević 2013) and it can thus be assumed that the same might be happening in some *Penicillium* species, although this has not been shown. Other possible secondary marker options include calmodulin (*CaM*) or the RNA polymerase II second largest subunit (*RPB2*) genes. Both these genes have similar discriminatory power as *BenA*. *RPB2* has the added advantage of lacking introns in the amplicon, allowing robust and easy alignments when used for phylogenies, but it is sometimes difficult to amplify and the database is incomplete. Similarly, we lack a complete *CaM* database. Thus, for routine identifications *BenA* is currently recommended, while for the description of new species, we suggest the use of ITS, *BenA*, *CaM* and *RPB2* among the markers for multilocus

sequence typing and GCPSR. We consider it good practise to include at least ITS and *BenA* sequences when describing new species, to allow others to more easily recognise the new species.

BenA can successfully be used for accurately identifying *Penicillium* species. However, as is the case for genes other than *BenA*, care should be taken in specific groups or situations. Infraspecies variation in *BenA* occurs in some *Penicillium* species as is observed in phylogenies published for *Penicillium* (Frisvad & Samson 2004, Barreto *et al.* 2011, Peterson *et al.* 2011, Houbraken *et al.* 2011b,c, Rivera & Seifert 2011, Rivera *et al.* 2012, Houbraken *et al.* 2012a, Visagie *et al.* 2013, 2014a, 2014b). This variation must be considered for identification purposes and especially when considering whether a strain might represent a new species. This means that in addition to the reference sequences of ex-type cultures sanctioned by ICPA, additional reference sequences are necessary to document sequence variation that differ from the ex-type. ICPA is currently working on populating such a validated database to capture infraspecies variation, but for the time being critical phylogenetic revisions of different sections should be referred to for reliable data. Alternatively, combining ITS, *BenA*, *CaM* and *RPB2* from a suspected new species with sequences of the same markers from related species will aid in deciding whether a species is new or not, using GCPSR as explained in detail by Taylor *et al.* (2000). This is in fact common practise in most studies describing and characterising *Penicillium* species.

β -tubulin as a secondary marker in practise

As discussed above, *BenA* works well for species identifications in *Penicillium*. *Penicillium chrysogenum* and *P. allii-sativi* is one example where identification based on *BenA* should be made with care (Houbraken *et al.* 2012a). This is a consequence of the variation observed among strains of *P. chrysogenum*, which results in *P. allii-sativi* forming a clade within the *P. chrysogenum* clade in *BenA* species trees. Even though *CaM* distinguishes *P. chrysogenum* from *P. allii-sativi*, it does not distinguish *P. chrysogenum* and *P. rubens*. As a result, molecular identifications should be made with great care in section *Chrysogena*, with the revision by Houbraken *et al.* (2012a) used as guide. *Penicillium kongii* was recently introduced as a close relative of *P. brevicompactum* (Wang & Wang 2013). Even though this species formed a coherent clade distinct from the *P. brevicompactum* ex-type strain, on examining additional strains previously identified as *P. brevicompactum*, the *P. kongii* clade is resolved within the *P. brevicompactum* clade. This clade thus requires more work to resolve the species from the complex. A similar situation is observed for *P. desertorum* and the newly described *P. glycyrrhizacola* (Chen *et al.* 2013), where more strains and additional genes included in the phylogeny will help to resolve species. Another species that cannot be identified using molecular data is *P. camemberti*, *P. caseiffulvum* and *P. commune*, which shares identical *BenA* and other gene sequences (Giraud *et al.* 2010). Although these could be considered taxonomic synonyms, their different applications and importance in the cheese industry suggest that it is of more value to keep them as separate species. Morphological identification is necessary in this case, with *P. camemberti* having white conidia, *P. caseiffulvum* having an orange reverse on YES and *P. commune* producing green conidia (Frisvad & Samson 2004).

Table 2. Primers used for amplification and sequencing.

Locus	Primer name	Direction	Primer sequence (5'-3')	Reference
Internal Transcribed Spacer (ITS)	ITS1	Forward	TCC GTA GGT GAA CCT GCG G	White <i>et al.</i> 1990
	ITS4	Reverse	TCC TCC GCT TAT TGA TAT GC	White <i>et al.</i> 1990
	V9G	Forward	TTA CGT CCC TGC CCT TTG TA	de Hoog & Gerrits van den Ende 1998
	LS266	Reverse	GCA TTC CCA AAC AAC TCG ACT C	Masclaux <i>et al.</i> 1995
β -tubulin (<i>BenA</i>)	Bt _{2a}	Forward	GGT AAC CAA ATC GGT GCT GCT TTC	Glass & Donaldson 1995
	Bt _{2b}	Reverse	ACC CTC AGT GTA GTG ACC CTT GGC	Glass & Donaldson 1995
Calmodulin (CaM)	CMD5	Forward	CCG AGT ACA AGG ARG CCT TC	Hong <i>et al.</i> 2006
	CMD6	Reverse	CCG ATR GAG GTC ATR ACG TGG	Hong <i>et al.</i> 2006
	CF1	Forward	GCC GAC TCT TTG ACY GAR GAR	Peterson <i>et al.</i> 2005
	CF4	Reverse	TTT YTG CAT CAT RAG YTG GAC	Peterson <i>et al.</i> 2005
RNA polymerase II second largest subunit (<i>RPB2</i>)	5F	Forward	GAY GAY MGW GAT CAY TTY GG	Liu <i>et al.</i> 1999
	7CR	Reverse	CCC ATR GCT TGY TTR CCC AT	Liu <i>et al.</i> 1999
	5Feur	Forward	GAY GAY CGK GAY CAY TTC GG	Houbraken <i>et al.</i> 2012b
	7CReur	Reverse	CCC ATR GCY TGY TTR CCC AT	Houbraken <i>et al.</i> 2012b

Amplification and identification

Primers used for amplification of the ITS, *BenA*, CaM and *RPB2* genes are included in Table 2 and amplification profiles are given in Table 3. A standard thermal cycle with an annealing temperature of 55 °C is generally used. Sometimes amplification success is low for CaM, especially in sections *Canescensia* and *Ramosa*. In this case, dropping the annealing temperature to 52 °C gives good results. The *RPB2* amplification is more complicated. We recommend using a touch-up PCR (50–52–55 °C) with primer pair 5Feur and 7CReur for best amplification. When amplification is problematic, the alternative touch-up PCR (48–50–52 °C) profile can be used and/or the alternative primer pair 5F and 7CR.

After sequences are obtained, there is a number of ways to use them to identify the strain. The most widely used method is the BLAST search on NCBI. Using this method, one is able to search all sequences in the database, but as noted above, there are many unidentified and misidentified sequences in the database. For ITS the RefSeq data set, accessible from the NCBI homepage (<http://www.ncbi.nlm.nih.gov/refseq/>), is now available and can be used to query ITS sequences against a verified ITS database (Schoch *et al.* 2014). Unfortunately, the RefSeq database does not cover alternative genes. For this, we suggest setting up Local BLAST files and using the verified *BenA* database provided here.

Extrolite data

Extrolites are produced by the mycelium and sporulating structures of *Penicillium* species, and exudates, diffusible pigments, and reverse colours are also mixtures of secondary metabolites. Studies of extrolite profiles were very useful for unravelling some morphological species concepts into biologically meaningful segregate species before DNA sequencing provided similar possibilities. As an example, the *P. aurantiogriseum* complex, critical contaminants of grain, was divided first into species using extrolite profiles by Frisvad & Filtenborg (1983, 1989), delimitations that were subsequently supported by *BenA* sequencing (Seifert & Louis-Seize 2000).

Identification of unknown strains using extrolites is possible for well-equipped chemical laboratories. The best way of using extrolites as identification aids is to extract and then separate them by HPLC and then partially or fully identify as many of the secondary metabolites as possible, generally using mass spectroscopy based technology (Frisvad *et al.* 2008). The media used for identification, especially CYA and YES agars, are optimal for production of most major diagnostic extrolites in *Penicillium* after incubation for 7 d at 25 °C in darkness. Agar plugs are extracted with a mixture of dichloromethane, ethyl-acetate and methanol. The metabolites extracted can then be analysed using advanced separation and detection techniques,

Table 3. Thermal cycle programs used for amplification.

Gene	Profile type	Initial denaturing	Cycles	Denaturing	Annealing	Elongation	Final elongation	Rest period
General ITS, <i>BenA</i> , CaM	standard	94 °C, 5 min	35	94 °C, 45 s	55 °C, 45 s	72 °C, 60 s	72 °C, 7 min	10 °C, ∞
General alternative	standard	94 °C, 5 min	35	94 °C, 45 s	52 °C, 45 s	72 °C, 60 s	72 °C, 7 min	10 °C, ∞
<i>RPB2</i>	touch-up	94 °C, 5 min	5	94 °C, 45 s	50 °C, 45 s	72 °C, 60 s		
			5	94 °C, 45 s	52 °C, 45 s	72 °C, 60 s		
			30	94 °C, 45 s	55 °C, 45 s	72 °C, 60 s	72 °C, 7 min	10 °C, ∞
<i>RPB2</i> alt.	touch-up	94 °C, 5 min	5	94 °C, 45 s	48 °C, 45 s	72 °C, 60 s		
			5	94 °C, 45 s	50 °C, 45 s	72 °C, 60 s		
			30	94 °C, 45 s	52 °C, 45 s	72 °C, 60 s	72 °C, 7 min	10 °C, ∞

for example ultra high performance liquid chromatography with diode array detection and high resolution mass spectrometric detection (UHPLC-DAD-HRMS) (Kildgaard *et al.* 2014, Klitgaard *et al.* 2014). However, simpler HPLC-diode array detection methods can also be used (Frisvad & Thrane 1987, 1993).

Laboratories with less sophisticated equipment can perform valuable confirmatory tests. Although Thin Layer Chromatography (TLC) is no longer considered state of the art for chemical research, it is still a useful technique for detecting coloured or uncoloured extrolites that can be used to confirm the identification of a *Penicillium* strain. For example, *P. brevicompactum* consistently produces large amounts of the colourless mycophenolic acid, and this metabolite will make a green colour reaction with ferric chloride (Clutterbuck & Raistrick 1933).

Sometimes only a few extrolites are needed to confirm the identity of a *Penicillium* isolate. If an isolate produces griseofulvin and roquefortine C, it can only be *P. coprophilum*, *P. griseofulvum* or *P. sclerotigenum* and if the isolate also produces cyclopiazonic acid, it can only be *P. griseofulvum* (Frisvad & Samson 2004).

Identification based purely on secondary metabolites is not yet possible for all species of *Penicillium*, but future databases will be better developed and an optimal battery of media for secondary metabolite production may allow this method to be used for identification of all species in the future.

Misidentifications in *Penicillium*

A few examples demonstrate why *Penicillium* identification is difficult or how misleading results might be obtained in practise using morphology or sequencing. For example, a fungus producing penicillones A and B and chloctanspirones A, B and terrestrols was identified as *P. terrestre* (patent strain CCTCC M 204077, strain unavailable) (Liu *et al.* 2005). The name *P. terrestre* was used by Raper & Thom (1949), but has since been considered a synonym of *P. crustosum*. However, additional secondary metabolites produced by the isolate, such as sorbicillin and trichodimerol, are never seen in *P. crustosum* (Frisvad *et al.* 2004), suggesting the strain was *P. chrysogenum* or *P. rubens*. Liu *et al.* (2005) did not describe how their isolate was identified. Another strain of *P. chrysogenum* or *P. rubens*, SD-118 (HQ652873), was misidentified as a *P. commune* (Shang *et al.* 2012, Zhao *et al.* 2012), based on "100 % sequence identity" with *P. commune* (FJ499541). The production of chrysogine, sorbicillins and meleagrin indicated that the strain, SD-118, was in fact *P. chrysogenum*. We propose that the use of the reference sequence databases provided here, will minimise these type of misidentifications.

Matrix-assisted laser desorption/ionization time-of-flight mass spectrometry fingerprinting (MALDI-TOF MS)

Another detection technique receiving a lot of attention, especially from the medical field, is matrix-assisted laser desorption/ionization time-of-flight mass spectrometry fingerprinting (MALDI-TOF MS). It was successfully applied to the identification of bacteria (Hettick *et al.* 2006, Siegrist *et al.* 2007) and yeast species (Amiri-Eliasi & Fenselau 2001, Kolecka *et al.* 2013). Only a limited number of studies included or focused on *Penicillium* (Welham *et al.* 2000, Hettick *et al.* 2008, Del Chierico *et al.* 2012, Chalupová *et al.* 2014) and the closely related *Aspergillus* (Bille

et al. 2011, De Carolis *et al.* 2011, Iriart *et al.* 2012, Verwer *et al.* 2013). Although these studies promisingly report that MALDI-TOF MS distinguish between species, a large degree of variation is observed within a species, even between duplicates of the same strain (Hettick *et al.* 2008). This means that data from a high number of strains will have to be included in the database to make robust identifications feasible. Difficulties with identifications are also reported in *Aspergillus*, where not all strains could be identified with 100 % accuracy (Iriart *et al.* 2012, Verwer *et al.* 2013). Thus we can conclude that although this technique shows promise, a lot of work remains to make routine identifications feasible.

TAXONOMY

Phylogenetic analyses revealed that *Hemicarpenteles paradoxus* should not be classified in *Aspergillus*, and together with its closest relatives *A. crystallinus* and *A. malodoratus*, should be transferred to *Penicillium* (Peterson 2000a, 2008, Houbraken & Samson 2011). Houbraken & Samson (2011) included them in *Penicillium* sect. *Paradoxa* together with *P. atramentosum*. The other *Hemicarpenteles* species were shown to belong in *Aspergillus* sect. *Clavati* (*H. acanthosporus* ≡ *A. acanthosporus* ≡ *Neocarpenteles acanthosporus* (Udagawa & Takada) Udagawa & Uchiy.) (Peterson 2000b, Tamura *et al.* 2000, Varga *et al.* 2007, Peterson 2008), while *H. ornatus* and *H. thaxteri* are classified in *Sclerocleista* (Pitt *et al.* 2000, Houbraken & Samson 2011).

Aspergillus paradoxus was described by Fennell & Raper (1955) for strains producing small clavate vesicles suggestive of those observed in aberrant strains of *A. clavatus*. Later, Raper & Fennell (1965) placed *A. paradoxus* in the *A. ornatus* species group, based on the resemblance of its sclerotia to young cleistothecia of *A. ornatus* and *A. citrinsporus*. Rai *et al.* (1964, 1967) also isolated *A. paradoxus* from soil in India, and found that conditions favouring sclerotia production also inhibited the formation of conidial heads and the yellow pigment in the medium. Sarbhoy & Elphick (1968) observed that a strain of *A. paradoxus* (CBS 793.68 = IMI 117502), isolated from dog dung in the United Kingdom, produced mature unilocular cleistothecia with tough peridial walls, embedding smooth-walled lenticular ascospores. They introduced the name *Hemicarpenteles* for the sexual morph, with *H. paradoxus* as type. Houbraken & Samson (2011) noted that the sexual morph most closely resembles *Eupenicillium*, both producing sclerotoid cleistothecia that ripen from the centre outwards (Sarbhoy & Elphick 1968, Pitt 1979, Stolk & Samson 1983). In our fresh isolates, some strains produce sclerotia abundantly while others have only a few sclerotia. Only two fresh isolates produced ascospores after incubation at 25 °C for three weeks on OA. *Aspergillus paradoxus*, *A. crystallinus* and *A. malodoratus* (Figs 3–6) have a striking similarity with members of section *Olsonii* and *Coronata* (Frisvad & Samson 2004, Samson *et al.* 2004), in particular the long, wide, smooth-walled conidiophores terminating in densely branched penicilli with metulae and phialides. Particularly *P. olsonii* bears a morphological resemblance with *A. paradoxus* and *A. crystallinus*.

Extrolite data also suggest that *A. paradoxus*, *A. crystallinus* and *A. malodoratus* belong in *Penicillium* (summarised in Table 4). Pseurotins are found in both *Penicillium* and *Aspergillus* species (Frisvad *et al.* 2004). Sorbicillins were previously only confirmed in *P. rubens* and *P. chrysogenum*, and not so far in




Fig. 3. *Penicillium paradoxum*. A. Colonies: top row left to right, obverse CYA, YES, DG18 and MEA; bottom row left to right, reverse CYA, reverse YES, reverse DG18 and CREA. B. Young sclerotia. C. Phototropic conidiophores after two weeks growth. D–H. Conidiophores. I. Conidia. Scale bars: D–I = 10 µm.




Fig. 4. *Penicillium paradoxum* sexual reproduction. A–C. Cleistothecia. D, E. Ascii. F. Ascospores. Scale bars: A, B = 500 µm; C–F = 10 µm.

Aspergillus, although Basaran & Demirbas (2010) reported sorbicillin production in *A. parasiticus*; we have never observed sorbicillins in any *A. parasiticus* strain. *Penicillium atramentosum*, also classified in section *Paradoxa*, produces andrastin A, atpenins, meleagrin and oxaline, roquefortine C & D and rugulovasine A & B (Frisvad et al. 2004, and reported here). Among *Penicillium* and *Aspergillus* species, the atpenins have only been found in *P. atramentosum* (Omura et al. 1988, Kawada et al. 2009), but the other listed extrolites are common in species of *Penicillium* sections *Paradoxa* and *Chrysogena*. Rugulovasins are an exception and have only been found in *P. commune* (Frisvad et al. 2004). *Aspergillus paradoxus* produces brefeldin, an extrolite also reported in *P. brefeldianum* (Hutchinson et al. 1983), but never in *Aspergillus* apart from a report for *A. clavatus* (Wang et al. 2002) that could not be confirmed (Varga et al. 2007). The latter strain was probably actually *A. ingratius*, first regarded as belonging to *Aspergillus* section *Clavati* (Yaguchi et al. 1993) but now considered a synonym of *A. paradoxus*. Xanthocillins are produced by *A. paradoxus* strains, and also by *P. chrysogenum* and *P. egyptiacum*. Meleagrin is commonly produced in *Penicillium* sections *Paradoxa* and *Chrysogena*, and species of several other sections (Frisvad et al. 2004). Only the closely related extrolite, neoxaline, has been reported in *A. japonicus* (Hirano et al. 1979). *Aspergillus crystallinus* produces chrysogine, typical of *P. rubens* and *P. chrysogenum* (Frisvad et al. 2004, Houbraken et al. 2012a), and a compound also reported in *A. nomius* (Varga et al. 2011). Chrysophanic acid, produced in *A. crystallinus*, was found in *Talaromyces islandicus* (Howard & Raistrick 1950), and pachybasin was found in *P. vulpinum* (Frisvad et al. 2004). *Aspergillus malodoratus* produces andrastin A, meleagrin and oxaline.

Andrastin A is produced by several *Penicillium* species including *P. rubens* and *P. chrysogenum* (Houbraken et al. 2011a, 2012a), but has not been found in *Aspergillus*. On balance, chemotaxonomic evidence points to the placement of *A. paradoxus*, *A. crystallinus* and *A. malodoratus* in *Penicillium*.

From the phenotypic, molecular and extrolite data we can conclude that the three species formerly placed in *Aspergillus* belong to *Penicillium*. Consequently we propose the following new combinations:

Penicillium paradoxum (Fennell & Raper) Samson, Houbraken, Visagie & Frisvad, **comb. nov.** MycoBank MB547045. **Figs 3, 4.**

Basionym: *Aspergillus paradoxus* Fennell & Raper, Mycologia 47: 69. 1955. MycoBank MB292853

= *Hemicarpenteles paradoxus* Sarbhoy & Elphick, Trans. Brit. Mycol. Soc. 51: 156. 1968. MycoBank MB265229

= *Aspergillus ingratius* Yaguchi, Someya & Udagawa, Trans. Mycol. Soc. Japan 34: 305. 1993. MycoBank MB361187

[Non *Trichocoma paradoxus* Jungh., Praemissa in floram cryptogamicam Javae insulae: 9. 1838. MycoBank MB161024]

Typus: New Zealand, Wellington, dung of opossum, 1948, isolated by J.H. Warcup, Neotype IMI 061446, culture ex-type CBS 527.65 = NRRL 2162 = ATCC 16918 = IMI 061446

Penicillium paradoxum (and *P. malodoratum*) deviate from other *Penicillium* species by being phototropic (Raper & Fennell 1965) and occasionally having a well-developed vesicle that makes the conidophore appear like an aspergillum (illustrated by Yaguchi et al. 1993 in the protologue of *Aspergillus ingratius*).




Fig. 5. *Penicillium crystallinum*. A. Colonies: top row left to right, obverse CYA, YES, DG18 and MEA; bottom row left to right, reverse CYA, reverse YES, reverse DG18 and CREA. B. Colony texture on MEA. C–G. Conidiophores. H. Conidia. Scale bars: C–H = 10 μm .




Fig. 6. *Penicillium malodoratum*. A. Colonies: top row left to right, obverse CYA, YES, DG18 and MEA; bottom row left to right, reverse CYA, reverse YES, reverse DG18 and CREA. B. Colony texture on MEA. C–G. Conidiophores. H. Conidia. Scale bars: C–H = 10 μm .

Table 4. Origin and extrolites of the strains examined in this study.

Species	Strain	Origin	Extrolites
<i>A. crystallinus</i>	CBS 479.65T = IBT 21947	Forest soil, Tilaran, Costa Rica	chrysophanic acid, chrysogine, meleagrin, pachybasin
<i>A. malodoratus</i>	CBS 490.65T = IBT 21948	Forest soil, Barranca, Costa Rica	andrastin A, oxaline, meleagrin
<i>A. paradoxus</i>	CBS 123898 = IBT 29765 = DTO 76F5	Dung of dog, the Netherlands	"astyl", brefeldin A, "SLOF"
	CBS 123898 = IBT 29768 = DTO 76F6	Dung of dog, the Netherlands	"astyl", brefeldin A, pseurotin, "SLOF", sorbicillins
	CBS 527.65T = IBT 19547	Dung of opossum, New Zealand	"astyl", brefeldin A, pseurotin, "SLOF", sorbicillins
	CBS 643.95 = IBT 17513 = PF 1116	Soil, USA, California (ex-type of <i>A. ingratus</i>)	"astyl", brefeldin A, pseurotin, "SLOF", sorbicillins
	CBS 793.68 = IBT 19572	Dung of dog, UK	"astyl", brefeldin A, "PAS", pseurotin, "SLOF", sorbicillins
	CBS 793.91 = IBT 12297	Dung of dog, Denmark	brefeldin A, "PAS", pseurotin, "SLOF", sorbicillins
	IBT 29766 = DTO 76F7	Dung of dog, the Netherlands	brefeldin A, pseurotin, "SLOF", sorbicillins
	IMI 173765 = IBT 19364	Jet fuel, New Zealand	"astyl", brefeldin A, pseurotin, sorbicillins
	IMI 320026 = IBT 19365	Air, UK	brefeldin A, pseurotin, sorbicillins

However, stipes were described as small, thin-walled and even septate, and the illustration by [Raper & Fennell \(1965\)](#) indicates a conidial head not unlike *P. crystallinum* and *P. malodoratum*. The latter authors also noted that *A. crystallinus* and *A. malodoratus* occasionally produced triseriate "sterigmata" and "conidial structures, particularly when small, tend to resemble those of the genus *Penicillium*". Most isolates of *P. paradoxum* are apparently from dung. *Penicillium paradoxum*, *P. malodoratus* and *P. crystallinum* produce strong penetrating odours, similar to those known from coprophilous species in *Penicillium* series *Claviformia*, apparently an adaptation to the dung habitat.

Penicillium crystallinum (Kwon-Chung & Fennell) Samson, Houbraken, Visagie & Frisvad, **comb. nov.** MycoBank MB809315. [Fig. 5](#).

Basionym: *Aspergillus crystallinus* Kwon-Chung & Fennell, The Genus *Aspergillus*: 471. 1965. MycoBank MB326624

Typus: **Costa Rica**, Province of Guanacaste, Tilaran, forest soil, 1962, isolated by K.J. Kwon & D.I. Fennell, Neotype IMI 139270, culture ex-type CBS 479.65 = NRRL 5082 = ATCC 16833 = IMI 139270

Penicillium crystallinum is only known from its ex-type strain (CBS 479.65 = NRRL 5082 = ATCC 16833 = IMI 139270), isolated from forest soil in Costa Rica. The strain shows typical sporulation, but the soft compacted masses of thin-walled cells suggesting sclerotia described by [Raper & Fennell \(1965\)](#) were not observed. The yellow needle crystals of pachybasin observed in the type strain are no longer conspicuously present, but the extrolite profile shows that the strain still produces this compound. The strain also produces the pungent odour noted in *P. paradoxum* and *P. malodoratum*.

Penicillium malodoratum (Kwon-Chung & Fennell) Samson, Houbraken, Visagie & Frisvad, **comb. nov.** MycoBank MB809316. [Fig. 6](#).

Basionym: *Aspergillus malodoratus* Kwon-Chung & Fennell, The genus *Aspergillus*: 468. 1965. MycoBank MB326644

Typus: **Costa Rica**, Province of Puntarenas, Barranca, forest soil, 1962, isolated by K.J. Kwon & D.I. Fennell, Neotype IMI

172289, culture ex-type CBS 490.65 = NRRL 5083 = IMI 172289 = ATCC 16834

Penicillium malodoratum is only known from its ex-type strain (CBS 490.65 = NRRL 5083 = IMI 172289 = ATCC 16834), isolated from forest soil in Costa Rica. Colonies contain numerous long, phototropic conidiophores. [Raper & Fennell \(1965\)](#) observed abundant soft sclerotium-like bodies with thick-walled cells, but we could not detect them. As the name suggests, this fungus produces an unpleasant odour on all media tested. Although *A. malodoratus* and *A. crystallinus* were isolated from forest soil in Costa Rica and have similar morphological characters, the taxa can be distinguished by the colony diameter, the length of the conidiophores and the size and shape of the conidia. Conidia of *A. crystallinus* are globose, 4–7 µm, echinulate, while those of *A. malodoratus* are subglobose to ellipsoidal, 3–3.5 × 3.5–4 µm and mainly smooth.

PROPOSED LIST OF ACCEPTED SPECIES IN THE GENUS *PENICILLIUM*

The following list includes species names that are accepted in the genus *Penicillium* on 8 August 2014. This list is updated from previous lists published by [Pitt & Samson \(1993\)](#) and [Pitt et al. \(2000\)](#). This revision was considered necessary following changes made to the ICN and the move to single name nomenclature ([McNeill et al. 2012](#)), the large number of species described since the 2000 list and new taxonomic information provided by molecular data.

The most dramatic changes in *Penicillium* are the incorporation of *Eupenicillium* and several other genera as synonyms ([Houbraken & Samson 2011](#)), and the transfer of species of the former *Penicillium* subgenus *Biverticillium* species to *Talaromyces* ([Samson et al. 2011](#)). *Aspergillus paradoxus*, *A. crystallinus* and *A. malodoratus* were shown to belong to *Penicillium* and were transferred above. Several species described as *Penicillium* belongs to other genera and not to *Penicillium* ([Houbraken & Samson 2011](#)). They are listed below the accepted species list.

Even though the updating of the accepted species list began for nomenclatural purposes, we aim to make this list more

functional by including additional information linked to the species names. The list thus includes MycoBank numbers where complete nomenclatural data can be obtained, collection numbers of ex-type strains for future taxonomists requiring authenticated reference material, the species current sectional classification, and GenBank accession numbers for ITS barcodes and, where available, alternative molecular identification markers for *BenA*, *CaM* and *RPB2*.

Despite the considerable amount of time and effort spent on completing this list, there is the possibility of errors or oversights. As such we solicit and gratefully accept any comments on missing names, errors or new data that has become available, especially when publishing a new species. We would also appreciate suggestions for making the list more useful. The active version of the list is currently hosted on the ICPA website (<http://www.aspergilluspenicillium.org>) where it will be updated as new information comes to light. The website contains a portal for comments, for the convenience of our users. The list published here contains only accepted species; the online version will in future also include data for synonyms. Similar lists are available for *Aspergillus* (Samson et al. 2014) and *Talaromyces* (Yilmaz et al. 2014).

Penicillium Link, Mag. Ges. Naturf. Freunde Berlin 3: 16. 1809. MycoBank MB9257.

= *Coremium* Link, Mag. Ges. Naturf. Freunde Berlin 3: 19. 1809, *fide Raper & Thom 1949*, Seifert & Samson 1985. [MB7782]. anamorphic synonym.
 = *Floccaria* Grev., Scott. crypt. fl.: pl. 301. 1827, *fide Seifert & Samson 1985*. [MB8260]. anamorphic synonym.
 ?= *Hormodendrum* Bonord., Handbuch allg. Mykol.: 76. 1851 *fide de Hoog & Hermanides-Nijhof 1977*, but see Hughes 1958. [MB8558]. anamorphic synonym.
 ?= *Walzia* Sorokin, Trudy Obshch. Ispyt. Prir. Imp. Kharkov: 47. 1871, *fide Constantin 1888*. [MB10429]. anamorphic synonym.
 = *Pritzeliella* Henn., Hedwigia Beibl.42: 88. 1903, *fide Clements & Shear 1931*. [MB9529]. anamorphic synonym.
 = *Eupenicillium* F. Ludw., Lehrbuch der Niederen Kryptogamen: 263. 1892, *fide Houbraken & Samson 2011*. [MB1933]. teleomorphic synonym.
 = *Citromyces* Wehmer, Ber. Deutsch. Bot. Ges. 11: 333. 1893, *fide Thom 1930*. [MB7672]. anamorphic synonym.
 = *Aspergillopsis* Sopp, Skr. Vidensk.-Selsk. Christiana Math.-Nat. Kl. 11: 204. 1912, non *Aspergillopsis* Speg. 1910, *fide Pitt 1979*. [MB22043]. anamorphic synonym.
 = *Carpenteles* Langeron, Crypt. Fr. Exs.: 344. 1922, *fide Stolk & Scott 1967*. [MB826]. teleomorphic synonym (= *Eupenicillium*).
 = *Torulomyces* Delitsch, Systematik der Schimmelpilze: 91. 1943, *fide Stolk & Samson 1983* and *Houbraken & Samson 2011*. [MB10252]. anamorphic synonym.
 = *Eladia* Smith, Trans. Brit. Mycol. Soc. 44: 47. 1961, *fide Samson 1981*, *Houbraken & Samson 2011*. [MB8134]. anamorphic synonym.
 = *Thysanophora* Kendrick, Can. J. Bot. 39: 820. 1961, *fide Houbraken & Samson 2011*. [MB10230]. anamorphic synonym.
 = *Hemicarpenteles* Sarbhoy & Elphick, Trans. Brit. Mycol. Soc. 51: 156. 1968, *fide Houbraken & Samson 2011*. [MB2279]. teleomorphic synonym.
 = *Penicillium* Link ex Gray sensu Pitt, The Genus *Penicillium*: 154. 1979 (nom. inval., art 13e). anamorphic synonym.
 = *Chromocleista* Yaguchi & Udagawa, Trans. Mycol. Soc. Japan 34: 101. 1993, *fide Houbraken & Samson 2011*. [MB25855]. teleomorphic synonym.

Penicillium abidjanum Stolk, Antonie van Leeuwenhoek 34: 49. 1968 ≡ *Eupenicillium abidjanum* Stolk, Antonie van Leeuwenhoek 34: 49. 1968. [MB335705]. — Herb.: CBS 246.67. Ex-type: CBS 246.67 = ATCC 18385 = FRR 1156 = IMI 136244. Section *Lanata-Divaricata*. ITS barcode: GU981582. (Alternative markers: *BenA* = GU981650; *RPB2* = JN121469; *CaM* = KF296383).

Penicillium admetzii K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat., 1927: 507. 1927. [MB119777]. — Herb.: IMI 39751. Ex-type: CBS 209.28 = ATCC 10407 = IMI 039751 = MUCL 29106 = NRRL 737. Section *Sclerotiora*. ITS barcode: JN714929. (Alternative markers: *BenA* = JN625957; *RPB2* = JN121455; *CaM* = KC773796).

Penicillium admetziooides S. Abe ex G. Sm., Trans. Brit. Mycol. Soc. 46: 335. 1963 ≡ *Penicillium admetziooides* S. Abe, J. Gen. Appl. Microbiol. 2: 68. 1956 (nom. inval., Art. 36). [MB302372]. — Herb.: CBS 313.59. Ex-type: CBS 313.59 = ATCC 18306 = FAT1302 = IFO 6055 = IMI 068227 = NRRL 3405 = QM 7312. Section *Sclerotiora*. ITS barcode: JN686433. (Alternative markers: *BenA* = JN799642; *RPB2* = JN406578; *CaM* = JN686387).

Penicillium albocoreum (Frissvad) Frissvad, Int. Mod. Tax. Meth. Pen. Asp. Clas.: 275. 2000 ≡ *Penicillium hirsutum* var. *albocoreum* Frissvad, Mycologia 81: 856. 1990. [MB459817]. — Herb.: IMI 285511. Ex-type: CBS 472.84 = FRR 2931 = IBT 10682 = IBT 21502 = IMI 285511. Section *Fasciculata*. ITS barcode: AJ004819. (Alternative markers: *BenA* = AY674326; *RPB2* = n.a.; *CaM* = n.a.).

Penicillium alexiae Visagie Houbraken & Samson, Persoonia 31: 59. 2013. [MB803785]. — Herb.: CBS H-21142. Ex-type: CBS 134558. Section *Sclerotiora*. ITS barcode: KC790400. (Alternative markers: *BenA* = KC773778; *RPB2* = n.a.; *CaM* = KC773803).

Penicillium alfredii Visagie, Seifert & Samson, Stud. Mycol. 78: 116. 2014. [MB801980]. — Herb.: CBS H-21800. Ex-type: CBS 138224 = DTO 269A4. Section unclassified. ITS barcode: KJ775684. (Alternative markers: *BenA* = KJ775177; *RPB2* = KJ834520; *CaM* = KJ775411).

Penicillium allii Vincent & Pitt, Mycologia 81: 300. 1989 ≡ *Penicillium hirsutum* var. *allii* (Vincent & Pitt) Frissvad, Mycologia 81: 855. 1989. [MB125498]. — Herb.: MU Vincent 114. Ex-type: CBS 131.89 = IMI 321505 = NRRL 13630 = ATCC 64636 = IMI 321506. Section *Fasciculata*. ITS barcode: AJ005484. (Alternative markers: *BenA* = AY674331; *RPB2* = n.a.; *CaM* = n.a.).

Penicillium allii-sativi Frissvad, Houbraken & Samson, Persoonia 29: 89. 2012. [MB801873]. — Herb.: CBS H-21058. Ex-type: CBS 132074 = DTO 149A8 = IBT 26507 = LJC 206. Section *Chrysogena*. ITS barcode: JX997021. (Alternative markers: *BenA* = JX996891; *RPB2* = JX996627; *CaM* = JX996232).

Penicillium alutaceum D.B. Scott, Mycopathol. Mycol. Appl. 36: 17. 1968 ≡ *Eupenicillium alutaceum* D.B. Scott, Mycopathol. Mycol. Appl. 36: 17. 1968. [MB335708]. — Herb.: CBS 317.67. Ex-type: CBS 317.67 = ATCC 18542 = FRR 1158 = IFO 31728 = IMI 136243. Section *Exilicaulis*. ITS barcode: AF033454. (Alternative markers: *BenA* = KJ834430; *RPB2* = JN121489; *CaM* = n.a.).

Penicillium amaliae Visagie, Houbraken & K. Jacobs, Persoonia 31: 52. 2013. [MB803784]. — Herb.: CBS H-21141. Ex-type: CBS 134209 = CV 1875 = DTO 183F3 = DAOM 241034. Section *Sclerotiora*. ITS barcode: JX091443. (Alternative markers: *BenA* = JX091563; *RPB2* = n.a.; *CaM* = JX141557).

Penicillium anatolicum Stolk, Antonie van Leeuwenhoek 34: 46. 1968 ≡ *Eupenicillium anatolicum* Stolk, Antonie van Leeuwenhoek 34: 46. 1968. [MB335710]. — Herb.: CBS 479.66. Ex-type: CBS 479.66 = IBT 30764. Section *Citrina*. ITS barcode: AF033425. (Alternative markers: *BenA* = JN606849; *RPB2* = JN606593; *CaM* = JN606571).

Penicillium angulare S.W. Peterson, E.M. Bayer & Wicklow, Mycologia 96: 1289. 2004. [MB487891]. — Herb.: BPI 842268. Ex-type: CBS 130293 = IBT 27051 = NRRL 28157. Section *Sclerotiora*. ITS barcode: AF125937. (Alternative markers: *BenA* = KC773779; *RPB2* = JN406554; *CaM* = KC773804).

Penicillium angustiporcatum Takada & Udagawa, Trans. Mycol. Soc. Japan 24: 143. 1983 ≡ *Eupenicillium angustiporcatum* Takada & Udagawa, Trans. Mycol. Soc. Japan 24: 143. 1983. [MB108322]. — Herb.: NHL 6481. Ex-type: CBS 202.84. Section *Gracilenta*. ITS barcode: KC411690. (Alternative markers: *BenA* = KJ834431; *RPB2* = JN406617; *CaM* = n.a.).

Penicillium antarcticum A.D. Hocking & C.F. McRae, Polar Biol. 21: 103. 1999. [MB482749]. — Herb.: DAR 72813. Ex-type: CBS 100492 = FRR 4989. Section *Canescens*. ITS barcode: KJ834503. (Alternative markers: *BenA* = KJ834432; *RPB2* = JN406653; *CaM* = n.a.).

Penicillium araracuaraense Houbraken, et al., Int. J. Syst. Evol. Microbiol. 61: 1469. 2011. [MB518025]. — Herb.: HUA 170334. Ex-type: CBS 113149 = IBT 23247. Section *Lanata-Divaricata*. ITS barcode: GU981597. (Alternative markers: *BenA* = GU981642; *RPB2* = KF296414; *CaM* = KF296373).

Penicillium ardesiacum Novobr., Novosti Sist. Nizsh. Rast. 11: 228. 1974. [MB319257]. — Herb.: IMI 174719. Ex-type: CBS 497.73 = ATCC 24719 = FRR 1479 = IFO 30540 = IMI 174719 = VKMF-1749. Section

- Aspergilloides*. ITS barcode: KM189565. (Alternative markers: *BenA* = KM088805; *RPB2* = KM089577; *CaM* = KM089190).
- Penicillium argentinense*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 78. 2011. [MB563185]. — Herb.: CBS H-20461. Ex-type: CBS 130371 = IBT 30761. Section *Citrina*. ITS barcode: JN831361. (Alternative markers: *BenA* = JN606815; *RPB2* = n.a.; *CaM* = JN60549).
- Penicillium arianeae*** Visagie, Houbraken & Samson, Persoonia 31: 59. 2013. [MB803786]. — Herb.: CBS H-21143. Ex-type: CBS 134559. Section *Sclerotiora*. ITS barcode: KC773833. (Alternative markers: *BenA* = KC773784; *RPB2* = n.a.; *CaM* = KC773811).
- Penicillium armarii*** Houbraken, et al. Stud. Mycol. 78: 410. 2014. [MB809955]. — Herb.: CBS H-21870. Ex-type: CBS 138171 = DTO 235-F1. Section *Aspergilloides*. ITS barcode: KM189758. (Alternative markers: *BenA* = KM089007; *RPB2* = KM089781; *CaM* = KM089394).
- Penicillium astrolabium*** R. Serra & S.W. Peterson, Mycologia 99: 80. 2007. [MB504766]. — Herb.: BPI 872160. Ex-type: CBS 122427 = NRRL 35611 = MUM 06.161. Section *Brevicompacta*. ITS barcode: DQ645804. (Alternative markers: *BenA* = DQ645793; *RPB2* = JN406634; *CaM* = DQ645808).
- Penicillium athertonense*** Houbraken, Stud. Mycol. 78: 412. 2014. [MB809956]. — Herb.: CBS H-21874. Ex-type: CBS 138161 = DTO 030-C2. Section *Aspergilloides*. ITS barcode: KM189462. (Alternative markers: *BenA* = KM088690; *RPB2* = KM089462; *CaM* = KM089075).
- Penicillium atramentosum*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 118: 65. 1910. [MB237291]. — Herb.: IMI 39752. Ex-type: CBS 291.48 = ATCC 10104 = FRR 795 = IBT 6616 = IFO 8137 = IMI 039752 = IMI 039752ii = LSHBP 1 = MUCL 29071 = MUCL 29126 = NRRL 795 = QM 7483. Section *Paradoxa*. ITS barcode: AF033483. (Alternative markers: *BenA* = AY674402; *RPB2* = JN406584; *CaM* = FJ530964).
- Penicillium atrofulvum*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 80. 2011. [MB563183]. — Herb.: CBS H-20650. Ex-type: CBS 109.66 = DTO 31B2 = FRR 799 = IBT 30032 = IBT 29667. Section *Citrina*. ITS barcode: JN617663. (Alternative markers: *BenA* = JN606677; *RPB2* = JN606620; *CaM* = JN606387).
- Penicillium atrosanguineum*** B.X. Dong, Ceská Mykol. 27: 174. 1973. [MB319260]. — Herb.: CBS H-15524. Ex-type: CBS 380.75 = FRR 1726 = IMI 197488. Section *Exilicaulis*. ITS barcode: JN617706. (Alternative markers: *BenA* = KJ834435; *RPB2* = JN406557; *CaM* = n.a.).
- Penicillium atrovenetum*** G. Sm., Trans. Brit. Mycol. Soc. 39: 112. 1956. [MB302377]. — Herb.: IMI 061837. Ex-type: CBS 241.56 = ATCC 13352 = FRR 2571 = IFO 8138 = IMI 061837 = LSHBSm683 = QM 6963. Section *Canescensia*. ITS barcode: AF033492. (Alternative markers: *BenA* = JX140944; *RPB2* = JN121467; *CaM* = KJ867004).
- Penicillium aurantiacobrunneum*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 80. 2011. [MB563206]. — Herb.: CBS H-20662. Ex-type: CBS 126228 = IBT 18753. Section *Citrina*. ITS barcode: JN617670. (Alternative markers: *BenA* = JN606702; *RPB2* = n.a.; *CaM* = JN606522).
- Penicillium aurantiogriseum*** Dierckx, Ann. Soc. Sci. Bruxelles 25: 88. 1901. [MB247956]. — Herb.: IMI 195050. Ex-type: CBS 249.89 = ATCC 48920 = FRR 971 = IBT 14016 = IMI 195050 = MUCL 29090 = NRRL 971. Section *Fasciculata*. ITS barcode: AF033476. (Alternative markers: *BenA* = AY674296; *RPB2* = JN406573; *CaM* = n.a.).
- Penicillium aurantioviolaceum*** Biourge, Cellule 33: 282. 1923. [MB257885]. — Herb.: CBS H-21954. Ex-type: CBS 137777 = NRRL 762 = ATCC 14974. Section *Aspergilloides*. ITS barcode: KM189756. (Alternative markers: *BenA* = KM089005; *RPB2* = KM089779; *CaM* = KM089392).
- Penicillium austroafricanum*** Houbraken & Visagie, Stud. Mycol. 78: 412. 2014. [MB809957]. — Herb.: CBS H-21864. Ex-type: CBS 137773 = DTO 133-G5. Section *Aspergilloides*. ITS barcode: KM189610. (Alternative markers: *BenA* = KM088854; *RPB2* = KM089628; *CaM* = KM089241).
- Penicillium bialowiezense*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 450. 1927. [MB258429]. — Herb.: unknown. Ex-type: CBS 227.28 = IBT 23044 = IMI 092237 = LSHBP 71 = NRRL 865. Section *Brevicompacta*. ITS barcode: EU587315. (Alternative markers: *BenA* = AY674439; *RPB2* = JN406604; *CaM* = AY484828).
- Penicillium bifforme*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 118: 54. 1910. [MB240764]. — Herb.: unknown. Ex-type: CBS 297.48 = ATCC 10416 = FRR 885 = IFO 7722 = IMI 039820 = LSHB P72 = MUCL 29165 = NRRL 885 = QM 7492. Section *Fasciculata*. ITS barcode: KC411731. (Alternative markers: *BenA* = FJ930944; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium biliae*** Chalab., Bot. Mater. Otd. Sporov. Rast. 6: 165. 1950. [MB302379]. — Herb.: IMI 113677. Ex-type: CBS 221.66 = ATCC 22348 = ATCC 48731 = CCRC 31675 = FRR 3391 = IJFM 5025 = IMI 113677 = MUCL 31187 = VKMF-854. Section *Sclerotiora*. ITS barcode:
- JN714937. (Alternative markers: *BenA* = JN625966; *RPB2* = JN406610; *CaM* = JN626009).
- Penicillium boreae*** S.W. Peterson & Sigler, Mycol. Res. 106: 1112. 2002. [MB483980]. — Herb.: BPI 841395. Ex-type: CBS 111717 = NRRL 31002 = UAMH 3896. Section *Stolkia*. ITS barcode: AF481122. (Alternative markers: *BenA* = JN617715; *RPB2* = n.a.; *CaM* = AF481138).
- Penicillium bovifimosum*** (Tuthill & Frisvad) Houbraken & Samson, Stud. Mycol. 70: 47. 2011. \equiv *Eupenicillium bovifimosum* Tuthill & Frisvad, Mycologia 94: 241. 2002. [MB561957]. — Herb.: WY RMF 82071. Ex-type: CBS 102825 = RMF 9598. Section *Turbata*. ITS barcode: AF263347. (Alternative markers: *BenA* = KJ834436; *RPB2* = JN406649; *CaM* = FJ530989).
- Penicillium brasiliianum*** Bat., Anais Soc. Biol. Pernambuco 15: 162. 1957. [MB302381]. — Herb.: URM IMUR 56. Ex-type: CBS 253.55 = ATCC 12072 = FRR 3466 = QM 6947. Section *Lanata-Divaricata*. ITS barcode: GU981577. (Alternative markers: *BenA* = GU981629; *RPB2* = KF296420; *CaM* = AB667857).
- Penicillium brefeldianum*** B.O. Dodge, Mycologia 25: 92. 1933 \equiv *Carpenteles brefeldianum* (B.O. Dodge) Shear, Mycologia 26: 107. 1934 \equiv *Eupenicillium brefeldianum* (B.O. Dodge) Stolk & D.B. Scott, Persoonia 4: 400. 1967 \equiv *Penicillium dodgei* Pitt, Genus *Penicillium*: 117. 1980. [MB258851]. — Herb.: IMI 216896. Ex-type: CBS 235.81 = NRRL 710 = FRR 710 = IFO 31731 = IMI 216896 = LCP 89.2573 = LCP 89.2578 = MUCL 38762 = QM 1872 = Thom 5296. Section *Lanata-Divaricata*. ITS barcode: AF033435. (Alternative markers: *BenA* = GU981623; *RPB2* = KF296421; *CaM* = EU021683).
- Penicillium brevicompactum*** Dierckx, Ann. Soc. Sci. Bruxelles 25: 88. 1901. [MB149773]. — Herb.: IMI 40225. Ex-type: CBS 257.29 = ATCC 10418 = ATCC 9056 = DSM3825 = FRR 862 = IBT 23045 = IMI 040225 = LSHBP 75 = MUCL 28647 = MUCL 28813 = MUCL 28935 = MUCL 30240 = MUCL 30241 = MUCL 30256 = MUCL 30257 = NRRL 2011 = NRRL 862 = NRRL 864 = QM 7496. Section *Brevicompacta*. ITS barcode: AY484912. (Alternative markers: *BenA* = AY674437; *RPB2* = JN406594; *CaM* = AY484813).
- Penicillium brevistipitatum*** L. Wang & W.Y. Zhuang, Mycotaxon 93: 234. 2005. [MB356064]. — Herb.: HMAS 130354-1-4. Ex-type: AS 3.6887. Section *Penicillium*. ITS barcode: DQ221696. (Alternative markers: *BenA* = DQ221695; *RPB2* = JN406528; *CaM* = n.a.).
- Penicillium brocae*** S.W. Peterson et al., Mycologia 95: 143. 2003. [MB373658]. — Herb.: BPI 841763. Ex-type: CBS 116113 = IBT 26293 = NRRL 31472. Section *Sclerotiora*. ITS barcode: AF484398. (Alternative markers: *BenA* = KC773787; *RPB2* = JN406639; *CaM* = KC773814).
- Penicillium brunneoconidiatum*** Visagie, Houbraken & K. Jacobs, Stud. Mycol. 78: 415. 2014. [MB809958]. — Herb.: CBS H-21873. Ex-type: CBS 137732 = DTO 182-E4 = CV 949 = DAOM 241359. Section *Aspergilloides*. ITS barcode: KM189666. (Alternative markers: *BenA* = KM088911; *RPB2* = KM089685; *CaM* = KM089298).
- Penicillium buchwaldii*** Frisvad & Samson, FEMS Microbiol. Lett. 339: 86. 2013. [MB800966]. — Herb.: IMI 304286. Ex-type: CBS 117181 = IBT 6005 = IMI 304286. Section *Brevicompacta*. ITS barcode: JX313164. (Alternative markers: *BenA* = JX313182; *RPB2* = JN406637; *CaM* = JX313148).
- Penicillium burgense*** Quintan., Av. Aliment. Majora Anim. 30: 176. 1990. [MB130241]. — Herb.: Sáez 1538. Ex-type: CBS 325.89. Section *Exilicaulis*. ITS barcode: KC411736. (Alternative markers: *BenA* = KJ834437; *RPB2* = JN406572; *CaM* = n.a.).
- Penicillium bussumense*** Houbraken, Stud. Mycol. 78: 415. 2014. [MB809959]. — Herb.: CBS H-21869. Ex-type: CBS 138160 = DTO 018-B2. Section *Aspergilloides*. ITS barcode: KM189458. (Alternative markers: *BenA* = KM088685; *RPB2* = KM089457; *CaM* = KM089070).
- Penicillium cainii*** K.G. Rivera, Malloch & Seifert, Stud. Mycol. 70: 147. 2011. [MB563159]. — Herb.: DAOM 239914. Ex-type: CCFC 239914. Section *Sclerotiora*. ITS barcode: JN686435. (Alternative markers: *BenA* = JN686366; *RPB2* = n.a.; *CaM* = JN686389).
- Penicillium cairnsense*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 83. 2011. [MB563184]. — Herb.: CBS H-20686. Ex-type: CBS 124325 = IBT 29042. Section *Citrina*. ITS barcode: JN617669. (Alternative markers: *BenA* = JN606692; *RPB2* = n.a.; *CaM* = JN606512).
- Penicillium camemberti*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 82: 33. 1906. [MB175171]. — Herb.: IMI 27831. Ex-type: CBS 299.48 = ATCC 1105 = ATCC 4845 = FRR 878 = IBT 21508 = IMI 027831 = IMI 092200 = LCP 66.584 = LSHBP 11 = MUCL 29790 = NCTC 582 = NRRL 877 = NRRL 878. Section *Fasciculata*. ITS barcode: AB479314. (Alternative markers: *BenA* = FJ930956; *RPB2* = JN121484; *CaM* = n.a.).
- Penicillium canariense*** S.W. Peterson & Sigler, Mycol. Res. 106: 1113. 2002. [MB483981]. — Herb.: BPI 841396. Ex-type: CBS 111720 = NRRL

- 31003 = IJFM 536 = UAMH 6403. Section *Stolkia*. ITS barcode: AF481121. (Alternative markers: *BenA* = JN617714; *RPB2* = n.a.; *CaM* = AF481137).
- Penicillium canescens*** Sopp, Skr. Vidensk.-Selsk. Christiana Math.-Nat. Kl. 11: 181. 1912. [MB153765]. — Herb.: IMI 28260. Ex-type: CBS 300.48 = ATCC 10419 = DSM1215 = FRR 910 = IMI 028260 = MUCL 29169 = NCTC 6607 = NRRL 910 = QM 7550 = VKMF-1148. Section *Canescencia*. ITS barcode: AF033493. (Alternative markers: *BenA* = JX140946; *RPB2* = JN121485; *CaM* = KJ867009).
- Penicillium canis*** S.W. Peterson, J. Clin. Microbiol. (in press). [MB807056]. — Herb.: BPI 892763. Ex-type: NRRL 62798. Section *Canescencia*. ITS barcode: KJ511291. (Alternative markers: *BenA* = KF900167; *RPB2* = KF900196; *CaM* = KF900177).
- Penicillium caperatum*** Udagawa & Y. Horie, Trans. Mycol. Soc. Japan 14: 371. 1973 ≡ *Eupenicillium caperatum* Udagawa and Y. Horie, Trans. Mycol. Soc. Japan 14: 371. 1973. [MB319262]. — Herb.: unknown. Ex-type: CBS 443.75 = ATCC 28046 = DSM2209 = NHL 6465. Section *Lanata-Divaricata*. ITS barcode: KC411761. (Alternative markers: *BenA* = GU981660; *RPB2* = KF296422; *CaM* = KF296392).
- Penicillium capsulatum*** Raper & Fennell, Mycologia 40: 528. 1948. [MB289079]. — Herb.: IMI 40576. Ex-type: CBS 301.48 = ATCC 10420 = DSM2210 = FRR 2056 = IJFM 5120 = IMI 040576 = NRRL 2056 = QM 4869 = VKMF-445. Section *Ramigena*. ITS barcode: AF033429. (Alternative markers: *BenA* = FJ004390; *RPB2* = JN406582; *CaM* = n.a.).
- Penicillium carneum*** (Frissvad) Frissvad, Microbiology, UK, 142: 546. 1996 ≡ *Penicillium roqueforti* var. *carneum* Frissvad, Mycologia 81: 858. 1990. [MB415652]. — Herb.: IMI 293204. Ex-type: CBS 112297 = IBT 6884 = IBT 18419 = IMI 293204. Section *Roquefortorum*. ITS barcode: HQ442338. (Alternative markers: *BenA* = AY674386; *RPB2* = JN406642; *CaM* = HQ442322).
- Penicillium cartierense*** Houbraken, Stud. Mycol. 78: 415. 2014. [MB809960]. — Herb.: CBS H-21861. Ex-type: CBS 137956 = DTO 092-H9. Section *Aspergilloides*. ITS barcode: KM189564. (Alternative markers: *BenA* = KM088804; *RPB2* = KM089576; *CaM* = KM089189).
- Penicillium caseifolivum*** Lund, Filt. & Frissvad, J. Food Mycol. 1: 97. 1998. [MB446013]. — Herb.: C 24999. Ex-type: CBS 101134 = IBT 18282 = IBT 21510. Section *Fasciculata*. ITS barcode: KJ834504. (Alternative markers: *BenA* = AY674372; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium catenatum*** D.B. Scott, Mycopathol. Mycol. Appl. 36: 24. 1968 ≡ *Eupenicillium catenatum* D.B. Scott, Mycopathol. Mycol. Appl. 36: 24. 1968. [MB335719]. — Herb.: CBS 352.67. Ex-type: CBS 352.67 = ATCC 18543 = CSIR 1097 = IFO 31774 = IMI 136241. Section *Exilicaulis*. ITS barcode: KC411754. (Alternative markers: *BenA* = KJ834438; *RPB2* = JN121504; *CaM* = n.a.).
- Penicillium cavernicola*** Frissvad & Samson, Stud. Mycol. 49: 31. 2004. [MB370976]. — Herb.: CBS H-13441. Ex-type: CBS 100540 = IBT 14499. Section *Fasciculata*. ITS barcode: KJ834505. (Alternative markers: *BenA* = KJ834439; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium chalybeum*** Pitt & A.D. Hocking, Mycotaxon 22: 204. 1985. [MB105608]. — Herb.: FRR 2660. Ex-type: CBS 254.87 = FRR 2660. Section *Exilicaulis*. ITS barcode: KC411713. (Alternative markers: *BenA* = KJ834440; *RPB2* = JN406596; *CaM* = n.a.).
- Penicillium charlesii*** G. Sm., Trans. Brit. Mycol. Soc. 18: 90. 1933. [MB260433]. — Herb.: unknown. Ex-type: CBS 304.48 = ATCC 8730 = CBS 342.51 = CECT 2277 = FRR 778 = IMI 040232 = LSHBBB127 = LSHBP 146 = NRRL 1887 = NRRL 778 = QM 6338 = QM 6838. Section *Charlesia*. ITS barcode: AF033400. (Alternative markers: *BenA* = JX091508; *RPB2* = JN121486; *CaM* = AY741727).
- Penicillium chermesinum*** Biourge, Cellule 33: 284. 1923. [MB260472]. — Herb.: IMI 191730. Ex-type: CBS 231.81 = NRRL 2048 = FRR 2048 = IFO 31745 = IMI 191730. Section *Cinnamopurpurea*. ITS barcode:AY742693. (Alternative markers: *BenA* = KJ834441; *RPB2* = JN406581; *CaM* = AY741728).
- Penicillium christenseniae*** Houbraken, Frissvad & Samson, Stud. Mycol. 70: 85. 2011. [MB563187]. — Herb.: CBS H-20656. Ex-type: CBS 126236 = IBT 23355. Section *Citrina*. ITS barcode: JN617674. (Alternative markers: *BenA* = JN606680; *RPB2* = JN606624; *CaM* = JN606373).
- Penicillium chrysogenum*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 118: 58. 1910. [MB165757]. — Herb.: IMI 24314. Ex-type: CBS 306.48 = ATCC 10106 = ATHUM2889 = CCRC 30564 = FRR 807 = IBT 5233 = IMI 024314 = IMI 092208 = LSHBAd 3 = LSHBP 19 = MUCL 29079 = MUCL 29145 = NCTC 589 = NRRL 807 = NRRL 810 = QM 7500. Section *Chrysogena*. ITS barcode: AF033465. (Alternative markers: *BenA* = AY495981; *RPB2* = JN121487; *CaM* = JX996273).
- Penicillium chrazszczii*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 464. 1927. [MB260609]. — Herb.: unknown. Ex-type: CBS 217.28 = FRR 903 = MUCL 29167 = NRRL 1741 = NRRL 903. Section *Citrina*. ITS barcode: GU944603. (Alternative markers: *BenA* = JN606758; *RPB2* = JN606628; *CaM* = JN606423).
- Penicillium cinerascens*** Biourge, Cellule 33: 308. 1923. [MB260785]. — Herb.: IMI 92234. Ex-type: NRRL 748 = ATCC 48693 = BIOURGE 90 = FRR 748 = IMI 92234 = QM 7555 = Thom 4733.34. Section *Exilicaulis*. ITS barcode: AF033455. (Alternative markers: *BenA* = JX141041; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium cinereotrum*** Chalab., Bot. Mater. Otd. Sporov. Rast. 6: 167. 1950. [MB302385]. — Herb.: CBS H-7469. Ex-type: CBS 222.66 = ATCC 22350 = FRR 3390 = IJFM 5024 = IMI 113676 = VKMF-856. Section *Exilicaulis*. ITS barcode: KC411700. (Alternative markers: *BenA* = KJ834442; *RPB2* = JN406608; *CaM* = KJ867002).
- Penicillium cinnamopurpureum*** Udagawa, J. Agric. Food Sci., Tokyo 5: 1. 1959. [MB302386]. — Herb.: unknown. Ex-type: CBS 429.65 = CBS 847.68 = NRRL 162 = ATCC 18489 = CSIR 936 = FAT 362 = IAM 7016 = IFO 6032 = NHL 6359 = QM 7888. Section *Cinnamopurpurea*. ITS barcode: EF626950. (Alternative markers: *BenA* = EF626948; *RPB2* = JN406533; *CaM* = EF626949).
- Penicillium citreonigrum*** Dierckx, Ann. Soc. Sci. Bruxelles 25: 86. 1901. [MB165197]. — Herb.: IMI 92209i. Ex-type: CBS 258.29 = ATCC 48736 = FRR 761 = IMI 092209 = LSHBP 20 = LSHBP 98 = MUCL 28648 = MUCL 29062 = MUCL 29116 = NRRL 761. Section *Exilicaulis*. ITS barcode: AF033456. (Alternative markers: *BenA* = EF198621; *RPB2* = JN121474; *CaM* = EF198628).
- Penicillium citrinum*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 118: 61. 1910. [MB165293]. — Herb.: IMI 92196ii. Ex-type: CBS 139.45 = ATCC 1109 = ATCC 36382 = CECT 2269 = FRR 1841 = IMI 091961 = IMI 092196 = LSHBAd 95 = LSHBP 25 = LSHBP 6 = MUCL 29781 = NRRL 1841 = NRRL 1842. Section *Citrina*. ITS barcode: AF033422. (Alternative markers: *BenA* = GU944545; *RPB2* = JF417416; *CaM* = GU944638).
- Penicillium clavigerum*** Demelius, Verh. Zool.-Bot. Ges. Wien 72: 74. 1923. [MB261069]. — Herb.: IMI 39807. Ex-type: CBS 310.48 = ATCC 10427 = CBS 255.94 = FRR 1003 = IMI 039807 = IMI 039807ii = MUCL 15623 = NRRL 1003 = QM 1918. Section *Penicillium*. ITS barcode: DQ339555. (Alternative markers: *BenA* = AY674427; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium clavistipitatum*** Visagie, Houbraken & K. Jacobs, Stud. Mycol. 78: 419. 2014. [MB809961]. — Herb.: CBS H-21882. Ex-type: CBS 138650 = DTO 182-E5 = CV 336 = KAS 4112 = DAOM 241092. Section *Aspergilloides*. ITS barcode: KM189667. (Alternative markers: *BenA* = KM088912; *RPB2* = KM089686; *CaM* = KM089299).
- Penicillium cluniae*** Quintan., Av. Aliment. Mejora Anim. 30: 174. 1990. [MB130240]. — Herb.: unknown. Ex-type: CBS 326.89. Section *Lanata-Divaricata*. ITS barcode: KF296406. (Alternative markers: *BenA* = KF296471; *RPB2* = KF296424; *CaM* = KF296402).
- Penicillium coeruleum*** Sopp *apud* Biourge, Cellule 33: 102. 1923. [MB446014]. — Herb.: unknown. Ex-type: CBS 141.45 = NCTC 6595. Section *Lanata-Divaricata*. ITS barcode: GU981606. (Alternative markers: *BenA* = GU981655; *RPB2* = KF296425; *CaM* = KF296393).
- Penicillium coffeeae*** S.W. Peterson et al., Mycologia 97: 662. 2005. [MB340281]. — Herb.: BPI 863480. Ex-type: CBS 119387 = IBT 27866 = NRRL 35363. Section *Charlesia*. ITS barcode: AY742702. (Alternative markers: *BenA* = KJ834443; *RPB2* = JN121436; *CaM* = AY741747).
- Penicillium commune*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 118: 56. 1910. [MB164241]. — Herb.: IMI 39812. Ex-type: CBS 311.48 = ATCC 10428 = ATCC 1111 = CCRC 31554 = DSM2211 = IBT 6200 = IFO 5763 = IMI 039812ii = IMI 039812iii = NRRL 890 = QM 1269 = VKMF-3233. Section *Fasciculata*. ITS barcode: AY213672. (Alternative markers: *BenA* = AY674366; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium concentricum*** Samson, Stolk & Hadlock, Stud. Mycol. 11: 17. 1976. [MB319263]. — Herb.: CBS 477.75. Ex-type: CBS 477.75 = IBT 14571 = IBT 6577. Section *Penicillium*. ITS barcode: KC411763. (Alternative markers: *BenA* = AY674413; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium confertum*** (Frissvad, Filt. & Wicklow) Frissvad, Mycologia 81: 852. 1990 ≡ *Penicillium glandicola* var. *confertum* Frissvad, Filt. & Wicklow, Can. J. Bot. 65: 769. 1987. [MB126404]. — Herb.: IMI 296930. Ex-type: CBS 171.87 = IBT 21515 = IBT 3098 = IBT 5672 = IMI 296930 = NRRL 13488 = NRRL A-26904. Section *Chrysogena*. ITS barcode: JX997081. (Alternative markers: *BenA* = AY674373; *RPB2* = JX996708; *CaM* = JX996963).
- Penicillium contaminatum*** Houbraken, Stud. Mycol. 78: 419. 2014. [MB809962]. — Herb.: CBS H-21866. Ex-type: CBS 345.52 = DTO 091-

- A3 = IMI 049057. Section *Aspergilloides*. ITS barcode: KM189554. (Alternative markers: *BenA* = KM088793; *RPB2* = KM089565; *CaM* = KM089178).
- Penicillium coprobum*** Frisvad, Mycologia 81: 853. 1990. [MB126405]. — Herb.: IMI 293209. Ex-type: CBS 561.90 = ATCC 58615 = IBT 21516 = IBT 4583 = IBT 6932. Section *Penicillium*. ITS barcode: DQ339559. (Alternative markers: *BenA* = AY674425; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium coprophilum*** (Berk. & M.A. Curtis) Seifert & Samson, Adv. *Penicillium Aspergillus* Syst.: 145. 1985 ≡ *Coremium coprophilum* Berk. & M.A. Curtis, J. Linn. Soc., Bot., 10: 363. 1868. [MB114760]. — Herb.: K Cuba Wright 666. Ex-type: CBS 110760 = IBT 5551 = IBT 3064 = NRRL 13627. Section *Penicillium*. ITS barcode: AF033469. (Alternative markers: *BenA* = AY674421; *RPB2* = JN406645; *CaM* = n.a.).
- Penicillium copicola*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 88. 2011. [MB563205]. — Herb.: CBS H-20643. Ex-type: CBS 127355 = IBT 30771. Section *Citrina*. ITS barcode: JN617685. (Alternative markers: *BenA* = JN606805; *RPB2* = JN606599; *CaM* = JN606553).
- Penicillium coralligerum*** Nicot & Pionnat, Bull. Soc. Mycol. France 78: 245. 1963 [1962]. [MB335721]. — Herb.: IMI 99159. Ex-type: CBS 123.65 = ATCC 16968 = FRR 3465 = IFO 9578 = IHEM 4511 = IMI 099159 = LCP 58.1674 = NRRL 3465. Section *Canescensia*. ITS barcode: JN617667. (Alternative markers: *BenA* = KJ834444; *RPB2* = JN406632; *CaM* = KJ866994).
- Penicillium corylophilum*** Dierckx, Ann. Soc. Sci. Bruxelles 25: 86. 1901. [MB178294]. — Herb.: IMI 39754. Ex-type: CBS 312.48 = TCC9784 = ATHUM2890 = CECT 2270 = FRR 802 = IMI 039754 = MUCL 28671 = MUCL 29073 = MUCL 29131 = NRRL 802 = QM 7510. Section *Exilicaulis*. ITS barcode: AF033450. (Alternative markers: *BenA* = JX141042; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium cosmopolitanum*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 91. 2011. [MB563188]. — Herb.: CBS H-20665. Ex-type: CBS 126995 = IBT 30681. Section *Citrina*. ITS barcode: JN617691. (Alternative markers: *BenA* = JN60733; *RPB2* = n.a.; *CaM* = JN606472).
- Penicillium cremeogriseum*** Chalab., Bot. Mater. Otd. Sporov. Rast. 6: 168. 1950. [MB302390]. — Herb.: CBS 223.60. Ex-type: CBS 223.66 = ATCC 18320 = ATCC 18323 = FRR 1734 = IJFM 5011 = IMI 197492 = NRRL 3389 = VKMF-1034. Section *Lanata-Divaricata*. ITS barcode: GU981586. (Alternative markers: *BenA* = GU981624; *RPB2* = KF296426; *CaM* = KF296403).
- Penicillium crocicola*** W. Yamam., Sci. Rep. Hyogo Univ. Agric. 2: 28. 1956. [MB302391]. — Herb.: CBS H-7528. Ex-type: CBS 745.70 = NRRL 6175 = ATCC 18313 = QM 7778. Section *Aspergilloides*. ITS barcode: KM189581. (Alternative markers: *BenA* = KJ834445; *RPB2* = JN406535; *CaM* = KM089210).
- Penicillium crustosum*** Thom, The Penicillia: 399. 1930. [MB262401]. — Herb.: IMI 91917. Ex-type: CBS 115503 = ATCC 52044 = FRR 1669 = IBT 5528 = IBT 6175 = IMI 091917 = NCTC 4002. Section *Fasciculata*. ITS barcode: AF033472. (Alternative markers: *BenA* = AY674353; *RPB2* = n.a.; *CaM* = DQ911132).
- Penicillium cryptum*** Goch., Mycotaxon 26: 349. 1986 ≡ *Eupenicillium cryptum* Goch., Mycotaxon 26: 349. 1986. [MB103648]. — Herb.: NY 769. Ex-type: CBS 271.89 = ATCC 60138 = IMI 296794 = NRRL 13460. Section *Torulomyces*. ITS barcode: KF303647. (Alternative markers: *BenA* = KF303608; *RPB2* = JN121478; *CaM* = KF303628).
- Penicillium crystallinum*** (Kwon-Chung & Fennell) Samson et al., (published here) ≡ *Aspergillus crystallinus* Kwon-Chung & Fennell, The Genus *Aspergillus*: 471. 1965. [MB809315]. — Herb.: IMI 139270. Ex-type: CBS 479.65 = NRRL 5082 = ATCC 16833 = IMI 139270. Section *Paradoxa*. ITS barcode: AF033486. (Alternative markers: *BenA* = EF669682; *RPB2* = EF669669; *CaM* = FJ530973).
- Penicillium cyaneum*** (Bainier & Sartory) Biourge, Cellule 33: 102. 1923 ≡ *Citromyces cyaneus* Bainier & Sartory, Bull. Soc. Mycol. France 29: 157. 1913. [MB251712]. — Herb.: IMI 39744. Ex-type: CBS 315.48 = ATCC 10432 = FRR 775 = IFO 5337 = IMI 039744 = NRRL 775 = QM 7516. Section *Ramigena*. ITS barcode: AF033427. (Alternative markers: *BenA* = JX091552; *RPB2* = JN406575; *CaM* = n.a.).
- Penicillium cyclopium*** Westling, Ark. Bot. 11: 90. 1911. [MB156739]. — Herb.: unknown. Ex-type: CBS 144.45 = ATCC 8731 = ATHUM2888 = CECT 2264 = DSM1250 = IBT 5130 = IMI 089372 = LSHBP 123 = MUCL 15613 = NRRL 1888 = QM 6839 = VKMF-265. Section *Fasciculata*. ITS barcode: JN097811. (Alternative markers: *BenA* = AY674310; *RPB2* = JN985388; *CaM* = n.a.).
- Penicillium daejeonium*** S.H. Yu & H.K. Sang, J. Microbiol. 51: 537. 2013. [MB561572]. — Herb.: KACC 46609. Ex-type: KACC 46609. Section *Sclerotiora*. ITS barcode: JX436489. (Alternative markers: *BenA* = JX436493; *RPB2* = n.a.; *CaM* = JX436491).
- Penicillium daleae*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 495. 1927. [MB262773]. — Herb.: IMI 89338. Ex-type: CBS 211.28 = ATCC 10435 = DSM 2449 = FRR 2025 = IFO 6087 = IFO 9072 = IMI 034910 = MUCL 29234 = NRRL 2025. Section *Lanata-Divaricata*. ITS barcode: GU981583. (Alternative markers: *BenA* = GU981649; *RPB2* = KF296427; *CaM* = KF296385).
- Penicillium decaturense*** S.W. Peterson, E.M. Bayer & Wicklow, Mycologia 96: 1290. 2004. [MB487890]. — Herb.: BPI 842267. Ex-type: CBS 117509 = NRRL 28152 = IBT 27117. Section *Citrina*. ITS barcode: GU944604. (Alternative markers: *BenA* = JN606685; *RPB2* = JN606621; *CaM* = JN606413).
- Penicillium decumbens*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 118: 71. 1910. [MB156582]. — Herb.: IMI 190875. Ex-type: CBS 230.81 = FRR 741 = IMI 190875 = MUCL 29107 = NRRL 741. Section *Exilicaulis*. ITS barcode: AY157490. (Alternative markers: *BenA* = KJ834446; *RPB2* = JN406601; *CaM* = n.a.).
- Penicillium desertorum*** Frisvad, Houbraken & Samson, Persoonia 29: 90. 2012. [MB801874]. — Herb.: CBS H-21056. Ex-type: CBS 131543 = IBT 16321 = DTO 14816. Section *Chrysogena*. ITS barcode: JX997010. (Alternative markers: *BenA* = JX996818; *RPB2* = JX996682; *CaM* = JX996937).
- Penicillium dierckxii*** Biourge, Cellule 33: 313. 1923. [MB263175]. — Herb.: IMI 92216. Ex-type: CBS 185.81 = IMI 092216 = LSHBP 32 = MUCL 28665 = NRRL 755. Section *Ramigena*. ITS barcode: EF634444. (Alternative markers: *BenA* = EF634442; *RPB2* = EF634445; *CaM* = EF634443).
- Penicillium digitatum*** (Pers.: Fr.) Sacc., Fung. Ital.: tab. 894. 1881 ≡ *Aspergillus digitatus* Pers., Disp. meth. Fung.: 41. 1794 ≡ *Monilia digitata* Pers., Syn. Meth. Fung.: 693. 1801: Fr., Syst. Mycol. 3: 411. 1832 ≡ *Mucor digitata* (Pers.) Mérat, Nouvelle flore des environs de Paris 1: 14. 1821. [MB169502]. — Herb.: Lectotype = icon in Saccardo, Fung. Ital.: tab. 894. Jul 1881. Ex-type: CBS 112082 = IBT 13068. Section *Digitata*. ITS barcode: KJ834506. (Alternative markers: *BenA* = KJ834447; *RPB2* = JN121426; *CaM* = n.a.).
- Penicillium dimorphosporum*** H.J. Swart, Trans. Brit. Mycol. Soc. 55: 310. 1970. [MB120334]. — Herb.: CBS 456.70. Ex-type: CBS 456.70 = ATCC 22783 = ATCC 52501 = FRR 1120 = IMI 149680. Section *Exilicaulis*. ITS barcode: AF081804. (Alternative markers: *BenA* = KJ834448; *RPB2* = JN121517; *CaM* = n.a.).
- Penicillium dipodomyicola*** (Frisvad, Filt. & Wicklow) Frisvad, Int. Mod. Meth. Pen. Asp. Clas.: 275. 2000. [MB459818]. — Herb.: IMI 296935. Ex-type: CBS 173.87 = IBT 21521 = IMI 296935 = ATCC 64187. Section *Penicillium*. ITS barcode: AY371616. (Alternative markers: *BenA* = AY674409; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium dipodomysis*** (Frisvad, Filt. & Wicklow) Banke, Firsvad & S. Rosend, Int. Mod. Meth. Pen. Asp. Clas. 270. 2000. [MB274307]. — Herb.: IMI 296926. Ex-type: IBT 5333 = CBS 110412 = NRRL 13485 = NRRL A-26836 = IMI 296926. Section *Chrysogena*. ITS barcode: AY371615. (Alternative markers: *BenA* = AY495991; *RPB2* = JF909932; *CaM* = JX996950).
- Penicillium discolor*** Frisvad & Samson, Antonie van Leeuwenhoek, 72: 120. 1997. [MB442902]. — Herb.: IMI 285513. Ex-type: CBS 474.84 = IBT 21523 = IBT 5738 = IBT 14440 = IMI 285513 = FRR 2933. Section *Fasciculata*. ITS barcode: AJ004816. (Alternative markers: *BenA* = AY674348; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium donkii*** Stolk, Persoonia 7: 333. 1973. [MB319267]. — Herb.: CBS 188.72. Ex-type: CBS 188.72 = NRRL 5562 = ATCC 48439 = CCRC 31694 = FRR 1738 = IFO 31746 = IMI 197489 = MUCL 31188. Section *Stolkia*. ITS barcode: AF033445. (Alternative markers: *BenA* = JN617718; *RPB2* = n.a.; *CaM* = AF481136).
- Penicillium dravuni*** Janso, Mycologia 97: 445. 2005. [MB501442]. — Herb.: BPI 844248. Ex-type: F01V25. Section *Exilicaulis*. ITS barcode: AY494856. (Alternative markers: *BenA* = n.a.; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium dunedinense*** Visagie, Seifert & Samson, Stud. Mycol. 78: 121. 2014. [MB809183]. — Herb.: CBS H-21803. Ex-type: CBS 138218 = DTO 244G1. Section *Canescensia*. ITS barcode: KJ775678. (Alternative markers: *BenA* = KJ775171; *RPB2* = n.a.; *CaM* = KJ775405).
- Penicillium echinulatum*** Raper & Thom ex Fassat, Acta Univ. Carol., Biol. 1974: 326. 1977. [MB319269]. — Herb.: PRM 778523. Ex-type: CBS 317.48 = IBT 6294 = IMI 040028 = ATCC 10434 = NRRL 1151 = FRR 1151 = IFO 7760 = MUCL 15615 = QM 7519. Section *Fasciculata*. ITS barcode: AF033473. (Alternative markers: *BenA* = AY674341; *RPB2* = n.a.; *CaM* = DQ911133).
- Penicillium egyptiacum*** J.F.H. Beyma, Zentralbl. Bakteriol. Parasitenk., Abt. 2 88: 137. 1933 ≡ *Eupenicillium egyptiacum* (J.F.H. Beyma) Stolk & D.B. Scott, Persoonia 4: 401. 1967. [MB263790]. — Herb.: IMI 040580. Ex-type: CBS

- 244.32 = ATCC 10441 = CSIR 707 = FRR 2090 = IBT 14684 = IFO 6094 = IFO 8141 = IFO 8847 = IMI 040580 = NRRL 2090 = QM 1875. Section *Chrysogena*. ITS barcode: AF033467. (Alternative markers: *BenA* = AY674374; *RPB2* = JN406598; *CaM* = JX996969).
- Penicillium ehrlichii** Kleb., Ber. Deutsch. Bot. Ges. 48: 374. 1930 ≡ *Eupenicillium ehrlichii* (Kleb.) Stolk & D.B. Scott, Persoonia 4: 400. 1967 ≡ *Penicillium klebahnii* Pitt, Genus *Penicillium*: 122. 1980. [MB319270]. — Herb.: IMI 039737. Ex-type: CBS 324.48 = ATCC 10442 = IMI 039737 = IMI 039737ii = NRRL 708 = QM 1874 = VKMF-273. Section *Lanata-Divaricata*. ITS barcode: AF033432. (Alternative markers: *BenA* = KF296464; *RPB2* = KF296428; *CaM* = KF296395).
- Penicillium elleniae** Houbenken et al., Int. J. Syst. Evol. Microbiol. 61: 1470. 2011. [MB518028]. — Herb.: HUA 170339. Ex-type: CBS 118135 = IBT 23229. Section *Lanata-Divaricata*. ITS barcode: GU981612. (Alternative markers: *BenA* = GU981663; *RPB2* = KF296429; *CaM* = KF296389).
- Penicillium ellipoideosporum** L. Wang & W.Y. Kong, Mycosistema 19: 463. 2000. [MB467721]. — Herb.: HMAS 71768. Ex-type: CBS 112493 = AS 3.5688. Section *Cinnamopurpurea*. ITS barcode: JX012224. (Alternative markers: *BenA* = JQ965104; *RPB2* = JN121427; *CaM* = AY678559).
- Penicillium erubescens** D.B. Scott, Mycopathol. Mycol. Appl. 36: 14. 1968 ≡ *Eupenicillium erubescens* D.B. Scott, Mycopathol. Mycol. Appl. 36: 14. 1968. [MB335726]. — Herb.: CBS 318.67. Ex-type: CBS 318.67 = ATCC 18544 = CSIR 1040 = FRR 814 = IFO 31734 = IMI 136204 = NRRL 6223. Section *Exilicaulis*. ITS barcode: AF033464. (Alternative markers: *BenA* = HQ646566; *RPB2* = JN121490; *CaM* = EU427281).
- Penicillium estinogenum** A. Komatsu & S. Abe ex G. Sm., Trans. Brit. Mycol. Soc. 46: 335. 1963 ≡ *Penicillium estinogenum* A. Komatsu & S. Abe, J. Gen. Appl. Microbiol., Tokyo 2: 132. 1956 (nom. inval., Art. 36). [MB302397]. — Herb.: IMI 68241. Ex-type: CBS 329.59 = ATCC 18310 = CCRC 31557 = FAT1196 = FRR 3428 = IFO 6230 = IMI 068241 = QM 8149 = VKMF-274. Section *Gracilenta*. ITS barcode: n.a. (Alternative markers: *BenA* = n.a.; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium euclaicum** J.F.H. Beyma, Antonie van Leeuwenhoek 6: 269. 1940 ≡ *Eupenicillium euclaicum* (J.F.H. Beyma) Stolk & Samson, Stud. Mycol. 23: 90. 1983. [MB289081]. — Herb.: CBS 323.71. Ex-type: CBS 323.71 = IBT 30767. Section *Citrina*. ITS barcode: JN617699. (Alternative markers: *BenA* = JN606856; *RPB2* = JN606594; *CaM* = JN606564).
- Penicillium expansum** Link, Mag. Ges. Naturf. Freunde Berlin 3: 16. 1809. [MB159382]. — Herb.: CBS H-7082. Ex-type: CBS 325.48 = ATCC 7861 = ATTHUM2891 = CCRC 30566 = FRR 976 = IBT 3486 = IBT 5101 = IMI 039761 = IMI 039761ii = MUCL 29192 = NRRL 976 = VKMF-275. Section *Penicillium*. ITS barcode: AY373912. (Alternative markers: *BenA* = AY674400; *RPB2* = JF417427; *CaM* = n.a.).
- Penicillium fagi** C. Ramírez & A.T. Martínez, Mycopathologia 63: 57. 1978. [MB283595]. — Herb.: IJFM 3049. Ex-type: CBS 689.77 = CCMF-696 = IJFM 3049 = IMI 253806 = VKMF-2178. Section *Exilicaulis*. ITS barcode: AF481124. (Alternative markers: *BenA* = KJ834449; *RPB2* = JN406540; *CaM* = n.a.).
- Penicillium fellutatum** Biourge, Cellule 33: 262. 1923. [MB264748]. — Herb.: IMI 39734. Ex-type: CBS 229.81 = CBS 326.48 = ATCC 10443 = FRR 746 = IFO 5761 = IMI 039734 = IMI 039734iii = NRRL 746 = QM 7554. Section *Charlesia*. ITS barcode: AF033399. (Alternative markers: *BenA* = KJ834450; *RPB2* = JN121460; *CaM* = AY741753).
- Penicillium fennelliae** Stolk, Antonie van Leeuwenhoek 35: 261. 1969. [MB335728]. — Herb.: CBS 711.68. Ex-type: CBS 711.68 = ATCC 22050 = ATCC 52492 = FRR 521 = IHEM 4389 = IMI 151747 = MUCL 31322. Section *Brevicompacta*. ITS barcode: JX313169. (Alternative markers: *BenA* = JX313185; *RPB2* = JN406536; *CaM* = JX313151).
- Penicillium flavidostipitatum** C. Ramírez & C.C. González, Mycopathologia 88: 3. 1984. [MB106338]. — Herb.: CBS 202.87. Ex-type: CBS 202.87 = IJFM 7824. Section *Exilicaulis*. ITS barcode: KC411691. (Alternative markers: *BenA* = KJ834451; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium flavigenum** Frisvad & Samson, Mycol. Res. 101: 620. 1997. [MB437441]. — Herb.: CBS 419.89. Ex-type: CBS 419.89 = BT21526 = IBT 3091 = IBT V1035 = IMI 293207. Section *Chrysogena*. ITS barcode: JX997105. (Alternative markers: *BenA* = AY495993; *RPB2* = JN406551; *CaM* = JX996281).
- Penicillium flavisclerotiatum** Visagie, Houbenken & K. Jacobs, Stud. Mycol. 78: 419. 2014. [MB809963]. — Herb.: CBS H-21879. Ex-type: CBS 137750 = DTO 180-I8 = CV 100 = DAOM 241157. Section *Aspergilloides*. ITS barcode: KM189644. (Alternative markers: *BenA* = KMO88888; *RPB2* = KM089662; *CaM* = KM089275).
- Penicillium formosanum** H.M. Hsieh, H.J. Su & Tzean, Trans. Mycol. Soc. Republ. China 2: 159. 1987. [MB126488]. — Herb.: PPEH 10001. Ex-type:
- CBS 211.92 = IBT 19748 = IBT 21527. Section *Penicillium*. ITS barcode: KC411696. (Alternative markers: *BenA* = AY674426; *RPB2* = JN406615; *CaM* = n.a.).
- Penicillium fractum** Udagawa, Trans. Mycol. Soc. Japan 9: 51. 1968 ≡ *Eupenicillium fractum* Udagawa, Trans. Mycol. Soc. Japan 9: 51. 1968. [MB335729]. — Herb.: CBS H-7086. Ex-type: CBS 124.68 = ATCC 18567 = FRR 3448 = IMI 136701 = NHL 6104 = NRRL 3448. Section *Fracta*. ITS barcode: KC411674. (Alternative markers: *BenA* = KJ834452; *RPB2* = JN121441; *CaM* = n.a.).
- Penicillium freii** Frisvad & Samson, Stud. Mycol. 49: 28. 2004. [MB369274]. — Herb.: IMI 285513. Ex-type: CBS 476.84 = IBT 5137 = IMI 285513. Section *Fasciculata*. ITS barcode: JN942696. (Alternative markers: *BenA* = AY674290; *RPB2* = JN985430; *CaM* = n.a.).
- Penicillium frequentans** Westling, Ark. Bot. 11: 133. 1911. [MB152118]. — Herb.: CBS 105.11. Ex-type: CBS 105.11. Section *Aspergilloides*. ITS barcode: KM189525. (Alternative markers: *BenA* = KM088762; *RPB2* = KM089534; *CaM* = KM089147).
- Penicillium fuscum** (Sopp) Biourge, Cellule 33: 103. 1923 ≡ *Citromyces fuscus* Sopp, Skr. Vidensk.-Selsk. Christiana Math.-Nat. Kl. 11: 120. 1912 ≡ *Eupenicillium pinetorum* Stolk, Antonie van Leeuwenhoek 34: 37. 1968. [MB289082]. — Herb.: WIS WSF 15-C. Ex-type: CBS 295.62 = ATCC 14770 = CCRC 31517 = DSM2438 = IFO 7743 = IMI 094209 = MUCL 31196 = NRRL 3008 = WSF15c. Section *Aspergilloides*. ITS barcode: AF033411. (Alternative markers: *BenA* = GQ367513; *RPB2* = JN121483; *CaM* = GQ367539).
- Penicillium fusisporum** L. Wang, PLoS ONE 9: e101454-P2. 2014. [MB806119]. — Herb.: HMAS 244961. Ex-type: CBS 137463 = NRRL 62805 = AS 3.15338. Section *Aspergilloides*. ITS barcode: KF769424. (Alternative markers: *BenA* = KF769400; *RPB2* = n.a.; *CaM* = KF769413).
- Penicillium galliacum** C. Ramírez, A.T. Martínez & Berer., Mycopathologia 72: 30. 1980. [MB113021]. — Herb.: CBS H-7464. Ex-type: CBS 167.81 = ATCC 42232 = IJFM 5597. Section *Citrina*. ITS barcode: JN617690. (Alternative markers: *BenA* = JN606837; *RPB2* = JN606609; *CaM* = JN606548).
- Penicillium georgiense** S.W. Peterson & B.W. Horn, Mycologia 101: 79. 2009. [MB509290]. — Herb.: BPI 877332. Ex-type: CBS 132826 = NRRL 35509. Section *Charlesii*. ITS barcode: EF422852. (Alternative markers: *BenA* = EF506223; *RPB2* = n.a.; *CaM* = EF506239).
- Penicillium glabrum** (Wehmer) Westling, Ark. Bot. 11: 131. 1911 ≡ *Citromyces glaber* Wehmer, Beitr. Einh. Pilze 1: 24. 1893. [MB120545]. — Herb.: IMI 91944. Ex-type: CBS 125543 = IBT 22658 = IMI 91944. Section *Aspergilloides*. ITS barcode: GU981567. (Alternative markers: *BenA* = GU981619; *RPB2* = JF417447; *CaM* = GQ367545).
- Penicillium gladioli** L. McCulloch & Thom, Science 67: 217. 1928 ≡ *Eupenicillium crustaceum* F. Ludw., Lehrb. Nied. Krypt.: 263. 1892. [MB266048]. — Herb.: IMI 34911. Ex-type: CBS 332.48 = ATCC 10448 = FRR 939 = IBT 14772 = IMI 034911 = IMI 034911ii = LCP 89.202 = MUCL 29174 = NRRL 939 = = QM 1955 = VKMF-2088. Section *Penicillium*. ITS barcode: AF033480. (Alternative markers: *BenA* = AY674287; *RPB2* = JN406567; *CaM* = n.a.).
- Penicillium glandicola** (Oudem.) Seifert & Samson, Adv. *Penicillium Aspergillus* Syst. 147. 1985 ≡ *Coremiun glandicola* Oudem., Ned. Kruidk. Arch. 2: 918. 1903. [MB114761]. — Herb.: Netherlands, Valkenburg, Jul 1901, Rick in herb. Oudemans (L). Ex-type: CBS 498.75 = IBT 21529 = IMI 154241. Section *Penicillium*. ITS barcode: AB479308. (Alternative markers: *BenA* = AY674415; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium glaucosorum** Demelius, Verh. Zool.-Bot. Ges. Wien 72: 72. 1923 (1922). [MB158423]. — Herb.: unknown. Ex-type: NRRL 908 (authentic acc. Raper & Thom 1949). Section *Lanata-Divaricata*. ITS barcode: KF296407. (Alternative markers: *BenA* = KF296469; *RPB2* = KF296430; *CaM* = KF296400).
- Penicillium glycyrrhizacola** A.J. Chen, B.D. Sun & W.W. Gao, PLoS ONE 8: e78285-P3. 2013. [MB804682]. — Herb.: HMAS 244707. Ex-type: CGMCC3.15271 (G4432). Section *Chrysogena*. ITS barcode: n.a. (Alternative markers: *BenA* = KF021538; *CaM* = n.a.; *RPB2* = KF021554).
- Penicillium godlewskii** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 466. 1927. [MB266206]. — Herb.: unknown. Ex-type: CBS 215.28 = ATCC 10449 = ATCC 48714 = FRR 2111 = IFO 7724 = IMI 040591 = MUCL 29243 = NRRL 2111 = QM 7566 = VKMF-1826. Section *Citrina*. ITS barcode: JN617692. (Alternative markers: *BenA* = JN606768; *RPB2* = JN606626; *CaM* = JN606443).
- Penicillium goetzii** J. Rogers et al., Persoonia 29: 92. 2012. [MB801876]. — Herb.: CBS H-21061. Ex-type: CBS 285.73 = DTO 88G6 = IBT 30199. Section *Chrysogena*. ITS barcode: JX997091. (Alternative markers: *BenA* = JX996847; *RPB2* = JX996716; *CaM* = JX996971).

- Penicillium gorlenkoanum*** Baghd., Novosti Sist. Nizsh. Rast. 5: 97. 1968. [MB335731]. — Herb.: CBS H-7490. Ex-type: CBS 408.69 = FRR 511 = IMI 140339 = VKMF-1079. Section *Citrina*. ITS barcode: GU944581. (Alternative markers: *BenA* = GU944520; *RPB2* = JN606601; *CaM* = GU944608).
- Penicillium gracilemum*** Udagawa & Y. Horie, Trans. Mycol. Soc. Japan 14: 373. 1973 ≡ *Eupenicillium gracilemum* Udagawa & Y. Horie, Trans. Mycol. Soc. Japan 14: 373. 1973. [MB319272]. — Herb.: NHL 6452. Ex-type: CBS 599.73 = ATCC 28047 = ATCC 48258 = FRR 1557 = IMI 216900 = NHL 6452. Section *Gracilenta*. ITS barcode: KC411768. (Alternative markers: *BenA* = KJ834453; *RPB2* = JN121537; *CaM* = n.a.).
- Penicillium grancanariae*** C. Ramírez, A.T. Martínez & Ferrer, Mycopathologia 66: 79. 1978. [MB319273]. — Herb.: IJFM 3745. Ex-type: CBS 687.77 = IJFM 3745 = IMI 253783. Section *Aspergilloides*. ITS barcode: KM189529. (Alternative markers: *BenA* = KM088766; *RPB2* = KM089538; *CaM* = KM089151).
- Penicillium grevilleicola*** Houbraken & Quaedvlieg, Stud. Mycol. 78: 423. 2014. [MB809964]. — Herb.: CBS H-21871. Ex-type: CBS 137775 = DTO 174-E6. Section *Aspergilloides*. ITS barcode: KM189630. (Alternative markers: *BenA* = KM088874; *RPB2* = KM089648; *CaM* = KM089261).
- Penicillium griseofulvum*** Dierckx, Ann. Soc. Sci. Bruxelles 25: 88. 1901. [MB120566]. — Herb.: IMI 75832. Ex-type: CBS 185.27 = ATCC 11885 = ATTHUM2893 = CECT 2605 = DSM896 = IBT 6740 = IFO 7640 = IFO 7641 = IMI 075832ii = LCP 79.3245 = LSHBP 68 = MUCL 28643 = NRRL 2152 = NRRL 2300 = QM 6902 = VKMF-286. Section *Penicillium*. ITS barcode: AF033468. (Alternative markers: *BenA* = AY674432; *RPB2* = JN121449; *CaM* = n.a.).
- Penicillium griseolum*** G. Sm., Trans. Br. Mycol. Soc. 40: 485. 1957. [MB302401]. — Herb.: unknown. Ex-type: CBS 277.58 = ATCC 18239 = FRR 2671 = IFO 8175 = IMI 071626 = LSH BB323 = NRRL 2671 = QM 7523. Section unclassified. ITS barcode: EF422848. (Alternative markers: *BenA* = EF506213; *CaM* = EF506232; *RPB2* = JN121480).
- Penicillium griseopurpureum*** G. Sm., Trans. Brit. Mycol. Soc. 48: 275. 1965. [MB335732]. — Herb.: IMI 96157. Ex-type: CBS 406.65 = ATCC 22353 = FRR 3429 = IFO 9147 = IMI 096157. Section *Lanata-Divaricata*. ITS barcode: KF296408. (Alternative markers: *BenA* = KF296467; *RPB2* = KF296431; *CaM* = KF296384).
- Penicillium guanacastense*** K.G. Rivera, Urb & Seifert, Mycotaxon 119: 324. 2011. [MB563044]. — Herb.: DAOM 239912. Ex-type: CCFC 239912. Section *Sclerotiora*. ITS barcode: JN626098. (Alternative markers: *BenA* = JN625967; *RPB2* = n.a.; *CaM* = JN626010).
- Penicillium guttulosum*** J.C. Gilman & E.V. Abbott, Iowa St. Coll. J. Sci. 1: 298. 1927. [MB266689]. — Herb.: unknown. Ex-type: ATCC 48734 = NRRL 907 = FRR 907 = Thom 4894.16. Section *Exilicaulis*. ITS barcode: HQ646592. (Alternative markers: *BenA* = HQ646576; *RPB2* = n.a.; *CaM* = HQ646587).
- Penicillium halotolerans*** Frisvad, Houbraken & Samson, Persoonia 29: 92. 2012. [MB801875]. — Herb.: CBS H-21060. Ex-type: CBS 131537 = DTO 148H9 = IBT 4315. Section *Chrysogena*. ITS barcode: JX997005. (Alternative markers: *BenA* = JX996816; *RPB2* = JX996680; *CaM* = JX996935).
- Penicillium hennebertii*** Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Thysanophora canadensis* Stolk & Hennebert, Persoonia 5: 189. 1968. [MB561964]. — Herb.: CBS H-7854. Ex-type: CBS 334.68 = ATCC 18741 = IMI 137644 = MUCL 21216 = VKMF-2999. Section *Thysanophora*. ITS barcode: KJ834507. (Alternative markers: *BenA* = KJ834454; *RPB2* = JN121493; *CaM* = n.a.).
- Penicillium herquei*** Bainier & Sartory, Bull. Soc. Mycol. France 28: 121. 1912. [MB536431]. — Herb.: IMI 28809. Ex-type: CBS 336.48 = NRRL 1040 = ATCC 10118 = BIOURGE 452 = FRR 1040 = IFO 31747 = IMI 28809 = MUCL 29213 = NCTC 1721 = QM 1926 = Thom 4640.447. Section *Sclerotiora*. ITS barcode: JN626101. (Alternative markers: *BenA* = JN625970; *RPB2* = JN121494; *CaM* = JN626013).
- Penicillium heteromorphum*** H.Z. Kong & Z.T. Qi, Mycosistema 1: 107. 1988. [MB135444]. — Herb.: CBS 226.89. Ex-type: CBS 226.89. Section *Exilicaulis*. ITS barcode: KC411702. (Alternative markers: *BenA* = KJ834455; *RPB2* = JN406605; *CaM* = n.a.).
- Penicillium hetheringtonii*** Houbraken, Frisvad & Samson, Fungal Divers. 44: 125. 2010. [MB518292]. — Herb.: CBS 122392. Ex-type: CBS 122392 = IBT 29057. Section *Citrina*. ITS barcode: GU944558. (Alternative markers: *BenA* = GU944538; *RPB2* = JN606606; *CaM* = GU944642).
- Penicillium hirayamae*** Udagawa, J. Agric. Soc. Tokyo 5: 6. 1959 ≡ *Eupenicillium hirayamae* D.B. Scott & Stolk, Antonie van Leeuwenhoek 33: 305. 1967. [MB302402]. — Herb.: IMI 78255. Ex-type: CBS 229.60 = ATCC 18312 = IFO 6435 = IMI 078255 = IMI 078255ii = NHL 6046 = NRRL 143 = QM 7885. Section *Sclerotiora*. ITS barcode: JN626095. (Alternative markers: *BenA* = JN625955; *RPB2* = JN121459; *CaM* = JN626003).
- Penicillium hirsutum*** Dierckx, Ann. Soc. Sci. Bruxelles 25: 89. 1901. [MB152720]. — Herb.: IMI 40213. Ex-type: CBS 135.41 = ATCC 10429 = FRR 2032 = IBT 21531 = IFO 6092 = IMI 040213 = MUCL 15622 = NRRL 2032. Section *Fasciculata*. ITS barcode: AY373918. (Alternative markers: *BenA* = AF003243; *RPB2* = JN406629; *CaM* = n.a.).
- Penicillium hispanicum*** C. Ramírez, A.T. Martínez & Ferrer, Mycopathologia 66: 77. 1978. [MB319274]. — Herb.: IJFM 3223. Ex-type: CBS 691.77 = ATCC 38667 = DSM2416 = IJFM 3223 = IMI 253785 = VKMF-2179. Section *Ramigena*. ITS barcode: JX841247. (Alternative markers: *BenA* = KJ834456; *RPB2* = JN406539; *CaM* = n.a.).
- Penicillium hoeksii*** Houbraken, Stud. Mycol. 78: 423. 2014. [MB809965]. — Herb.: CBS H-21860. Ex-type: CBS 137776 = DTO 192-H4. Section *Aspergilloides*. ITS barcode: KM189707. (Alternative markers: *BenA* = KM088954; *RPB2* = KM089728; *CaM* = KM089341).
- Penicillium hordei*** Stolk, Antonie van Leeuwenhoek 35: 270. 1969. [MB335734]. — Herb.: CBS 701.68. Ex-type: CBS 701.68 = ATCC 22053 = CECT 2290 = FRR 815 = IBT 17804 = IBT 6980 = IMI 151748 = MUCL 39559. Section *Fasciculata*. ITS barcode: n.a. (Alternative markers: *BenA* = AY674347; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium idahoense*** Paden, Mycopathol. Mycol. Appl. 43: 259. 1971 ≡ *Eupenicillium idahoense* Paden, Mycopathol. Mycol. appl. 43: 259. 1971. [MB319275]. — Herb.: UVIC JWP 66-32. Ex-type: CBS 341.68 = NRRL 5274 = ATCC 22055 = FRR 881 = IMI 148393. Section *Cinnamopurpurea*. ITS barcode: KC411747. (Alternative markers: *BenA* = EF626953; *RPB2* = JN121499; *CaM* = EF626954).
- Penicillium incoloratum*** L.Q. Huang & Z.T. Qi, Acta Mycol. Sin. 13: 264. 1994. [MB363421]. — Herb.: HMAS 65949. Ex-type: CBS 101753 = AS 3.4672. Section *Cinnamopurpurea*. ITS barcode: KJ834508. (Alternative markers: *BenA* = KJ834457; *RPB2* = JN406651; *CaM* = KJ866984).
- Penicillium indicum*** D.K. Sandhu & R.S. Sandhu, Can. J. Bot. 41: 1273. 1963. [MB335735]. — Herb.: CBS 7476. Ex-type: CBS 115.63 = NRRL 3387 = ATCC 18324 = FRR 3387 = IFO 31744 = IMI 166620. Section *Charlesia*. ITS barcode: AY742699. (Alternative markers: *BenA* = EU427263; *RPB2* = JN406640; *CaM* = AY741744).
- Penicillium infra-aurantiacum*** Visagie, Houbraken & K. Jacobs, Stud. Mycol. 78: 426. 2014. [MB809966]. — Herb.: CBS H-21880. Ex-type: CBS 137747 = DTO 183-C3 = CV 1518 = DAOM 241145. Section *Aspergilloides*. ITS barcode: KM189684. (Alternative markers: *BenA* = KM088930; *RPB2* = KM089704; *CaM* = KM089317).
- Penicillium infrapurpureum*** Visagie, Seifert & Samson, Stud. Mycol. 78: 116. 2014. [MB809181]. — Herb.: CBS H-21801. Ex-type: CBS 138219 = DTO 235F6. Section *Cinnamopurpurea*. ITS barcode: KJ775679. (Alternative markers: *BenA* = KJ775172; *RPB2* = n.a.; *CaM* = KJ775406).
- Penicillium inositatum*** D.B. Scott, Mycopathol. Mycol. Appl. 36: 20. 1968 ≡ *Eupenicillium inositatum* D.B. Scott, Mycopathol. Mycol. Appl. 36: 20. 1968. [MB335736]. — Herb.: CBS 351.67. Ex-type: CBS 351.67 = ATCC 18622 = CSIR 1096 = FRR 1163 = IMI 136214 = NRRL 5810. Section *Fracta*. ITS barcode: AF033431. (Alternative markers: *BenA* = KJ834458; *RPB2* = JN121503; *CaM* = n.a.).
- Penicillium isariiforme*** Stolk & J.A. Mey., Trans. Brit. Mycol. Soc. 40: 187. 1957. [MB302403]. — Herb.: IMI 60371. Ex-type: CBS 247.56 = ATCC 18425 = CCRC 31699 = IFO 6393 = IHEM 4376 = IMI 060371 = LSHBBB308 = MUCL 31191 = MUCL 31323 = NRRL 2638 = QM 1897. Section *Ochrosalmonea*. ITS barcode: AF454077. (Alternative markers: *BenA* = KJ834459; *RPB2* = JN121470; *CaM* = n.a.).
- Penicillium italicum*** Wehmer, Hedwigia 33: 211. 1894. [MB162660]. — Herb.: CBS 339.48. Ex-type: CBS 339.48 = ATCC 10454 = DSM2754 = FRR 983 = IBT 23029 = IMI 039760 = MUCL 15608 = NRRL 983 = QM 7572. Section *Penicillium*. ITS barcode: KJ834509. (Alternative markers: *BenA* = AY674398; *RPB2* = JN121496; *CaM* = n.a.).
- Penicillium jacksonii*** K.G. Rivera, Houbraken & Seifert, Stud. Mycol. 70: 151. 2011. [MB563160]. — Herb.: DAOM 239937. Ex-type: CCFC 239937. Section *Sclerotiora*. ITS barcode: JN686437. (Alternative markers: *BenA* = JN686368; *RPB2* = n.a.; *CaM* = JN686391).
- Penicillium jamesonlandense*** Frisvad & Overy, Int. J. Syst. Evol. Microbiol. 56: 1435. 2006. [MB521421]. — Herb.: DAOM 234087. Ex-type: CBS 102888 = DAOM 234087 = IBT 21984 = IBT 24411. Section *Ramosa*. ITS barcode: DQ267912. (Alternative markers: *BenA* = DQ309448; *RPB2* = n.a.; *CaM* = KJ866985).
- Penicillium janczewskii*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 488. 1927. [MB120703]. — Herb.: IMI 191499. Ex-type: CBS 221.28 = FRR 919 = IMI 191499 = NRRL 919. Section *Canescentia*. ITS barcode: AY157487. (Alternative markers: *BenA* = KJ834460; *RPB2* = JN406612; *CaM* = KJ867001).

- Penicillium janthinellum** Biourge, Cellule 33: 258. 1923. [MB119134]. — Herb.: IMI 40238. Ex-type: CBS 340.48 = ATCC 10455 = IMI 040238 = NRRL 2016 = QM 6865. Section *Lanata-Divaricata*. ITS barcode: GU981585. (Alternative markers: *BenA* = GU981625; *RPB2* = JN121497; *CaM* = KF296401).
- Penicillium javanicum** J.F.H. Beyma, Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk. 26: 17. 1929 ≡ *Carpenteles javanicum* (J.F.H. Beyma) Shear, Mycologia 26: 107. 1934 ≡ *Eupenicillium javanicum* (J.F.H. Beyma) Stolk & D.B. Scott, Persoonia 4: 398. 1967 ≡ *Penicillium indonesiae* Pitt, Genus *Penicillium*: 114. 1980. [MB268394]. — Herb.: CBS H-7088. Ex-type: CBS 341.48 = ATCC 9099 = CSIR 831 = FRR 707 = IFO 31735 = IMI 039733 = MUCL 29099 = NRRL 707 = QM 1876. Section *Lanata-Divaricata*. ITS barcode: GU981613. (Alternative markers: *BenA* = GU981657; *RPB2* = JN121498; *CaM* = KF296387).
- Penicillium jensenii** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 494. 1927. [MB120708]. — Herb.: IMI 39768. Ex-type: CBS 327.59 = ATCC 18317 = FRR 909 = IFO 5764 = IMI 039768 = LCP 89.1389 = NRRL 909 = QM 7587. Section *Canescensia*. ITS barcode: AY443470. (Alternative markers: *BenA* = JX140954; *RPB2* = JN406614; *CaM* = AY443490).
- Penicillium jiangxiense** H.Z. Kong & Z.Q. Liang, Mycosistema 22: 4. 2003. [MB489161]. — Herb.: HMAS 82540. Ex-type: AS 3.6521. Section *Cinnamopurpurea*. ITS barcode: KJ890411. (Alternative markers: *BenA* = KJ890409; *RPB2* = n.a.; *CaM* = KJ890407).
- Penicillium johnkrugii** K.G. Rivera, Houbraken & Seifert, Stud. Mycol. 70: 151. 2011. [MB563161]. — Herb.: DAOM 239943. Ex-type: CCFC 239943. Section *Sclerotiora*. ITS barcode: JN686447. (Alternative markers: *BenA* = JN686378; *RPB2* = n.a.; *CaM* = JN686401).
- Penicillium jugoslavicum** C. Ramírez & Munt.-Cvetk., Mycopathologia 88: 65. 1984. [MB124173]. — Herb.: CBS 192.87. Ex-type: CBS 192.87 = IJFM 7785 = IMI 314508. Section *Sclerotiora*. ITS barcode: KC773836. (Alternative markers: *BenA* = KC773789; *RPB2* = JN406618; *CaM* = KC773815).
- Penicillium kananaskense** Seifert, Frisvad & McLean, Can. J. Bot. 72: 20. 1994. [MB362160]. — Herb.: MU-F-39531. Ex-type: CBS 530.93 = ATCC 90282 = DAOM 216105 = IBT 11775 = IMI 356791. Section *Aspergilloides*. ITS barcode: KM189780. (Alternative markers: *BenA* = KM089030; *RPB2* = KM089804; *CaM* = KM089417).
- Penicillium katangense** Stolk, Antonie van Leeuwenhoek 34: 42. 1968 ≡ *Eupenicillium katangense* Stolk, Antonie van Leeuwenhoek 34: 42. 1968. [MB120725]. — Herb.: CBS 247.67. Ex-type: CBS 247.67 = ATCC 18388 = IMI 136206 = NRRL 5182. Section *Exilicaulis*. ITS barcode: AF033458. (Alternative markers: *BenA* = n.a.; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium kewense** G. Sm., Trans. Brit. Mycol. Soc. 44: 42. 1961. [MB335740]. — Herb.: CBS H-7077. Ex-type: CBS 344.61 = ATCC 18240 = FRR 3441 = IFO 8113 = IMI 086561 = LSHBBB400 = MUCL 2685 = NRRL 3332 = QM 7958. Section *Chrysogena*. ITS barcode: AF033466. (Alternative markers: *BenA* = JX996849; *RPB2* = JF417428; *CaM* = JX996973).
- Penicillium kamaense** Houbraken & Pitt, Stud. Mycol. 78: 426. 2014. [MB809967]. — Herb.: CBS H-21857. Ex-type: CBS 137947 = FRR 6087 = DTO 056-I6. Section *Aspergilloides*. ITS barcode: KM189506. (Alternative markers: *BenA* = KM088743; *RPB2* = KM089515; *CaM* = KM089128).
- Penicillium kojigenum** G. Sm., Trans. Brit. Mycol. Soc. 44: 43. 1961. [MB335741]. — Herb.: LSHTM BB394. Ex-type: CBS 345.61 = ATCC 18227 = CCRC 31515 = FRR 3442 = IFO 9581 = IMI 086562 = LSHBBB394 = MUCL 2457 = NRRL 3442 = QM 7957. Section *Ramosa*. ITS barcode: AF033489. (Alternative markers: *BenA* = KJ834463; *RPB2* = JN406564; *CaM* = KJ867011).
- Penicillium kongii** L. Wang, Mycologia 105: 1549. 2013. [MB803185]. — Herb.: HMAS 244382. Ex-type: AS 3.15329. Section *Brevicomacta*. ITS barcode: KC427191. (Alternative markers: *BenA* = KC427171; *RPB2* = n.a.; *CaM* = KC427151).
- Penicillium laeve** (K. Ando & Manoch) Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Torulomyces laevis* K. Ando & Manoch, Mycoscience 39: 317. 1998. [MB561960]. — Herb.: TNS-F-238517. Ex-type: CBS 136665 = KY 12727 = NBRC 109724. Section *Exilicaulis*. ITS barcode: KF667369. (Alternative markers: *BenA* = KF667365; *RPB2* = KF667371; *CaM* = KF667367).
- Penicillium lagena** (Delitsch) Stolk & Samson, Stud. Mycol. 23: 100. 1983 ≡ *Torulomyces lagena* Delitsch, Systematik der Schimmelpilze, Neudamm: 9. 1943. [MB109162]. — Herb.: CBS 185.65. Ex-type: CBS 185.65 = MUCL 8221 = JCM10149 = OAC10034. Section *Torulomyces*. ITS barcode: KF303665. (Alternative markers: *BenA* = KF303619; *RPB2* = JN121450; *CaM* = KF303634).
- Penicillium lanoscoeruleum** Thom, Penicillia: 322. 1930. [MB268949]. — Herb.: unknown. Ex-type: CBS 215.30 = CBS 334.48 = ATCC 10459 = IFO 7761 = IMI 039818 = NRRL 888 = QM 6755 = VKMF-3089. Section *Chrysogena*. ITS barcode: KC411740. (Alternative markers: *BenA* = JX996843; *RPB2* = JX996723; *CaM* = JX996967).
- Penicillium lanosum** Westling, Ark. Bot. 11: 97. 1911. [MB178497]. — Herb.: IMI 40224. Ex-type: CBS 106.11 = ATCC 10458 = FRR 2009 = IFO 5851 = IFO 6099 = IMI 040224 = LSHBP 86 = MUCL 29232 = NRRL 2009 = QM 7591. Section *Ramosa*. ITS barcode: DQ304540. (Alternative markers: *BenA* = DQ285627; *RPB2* = n.a.; *CaM* = FJ530974).
- Penicillium lapidosum** Raper & Fennell, Mycologia 40: 524. 1948 ≡ *Eupenicillium lapidosum* D.B. Scott & Stolk, Antonie van Leeuwenhoek 33: 298. 1967. [MB289094]. — Herb.: IMI 39743. Ex-type: CBS 343.48 = ATCC 10462 = CCT4477 = IFO 6100 = IMI 039743 = NRRL 718 = QM 1928. Section *Exilicaulis*. ITS barcode: AF033409. (Alternative markers: *BenA* = KJ834465; *RPB2* = JN121500; *CaM* = FJ530984).
- Penicillium lassenii** Paden, Mycopathol. Mycol. Appl. 43: 266. 1971 ≡ *Eupenicillium lassenii* Paden, Mycopathol. Mycol. Appl. 43: 266. 1971. [MB319281]. — Herb.: UVIC JWP 69-26. Ex-type: CBS 277.70 = NRRL 5272 = ATCC 22054 = FRR 858 = IMI 148395. Section *Torulomyces*. ITS barcode: KF303648. (Alternative markers: *BenA* = KF303607; *RPB2* = JN121481; *CaM* = KF303629).
- Penicillium lenticrescens** Visagie, Seifert & Samson, Stud. Mycol. 78: 123. 2014. [MB809184]. — Herb.: CBS H-21804. Ex-type: CBS 138215 = DTO 129A8. Section *Ramosa*. ITS barcode: KJ775675. (Alternative markers: *BenA* = KJ775168; *RPB2* = n.a.; *CaM* = KJ775404).
- Penicillium levitum** Raper & Fennell, Mycologia 40: 511. 1948 ≡ *Carpenteles levitum* (Raper & Fennell) C.R. Benj., Mycologia 47: 685. 1955 ≡ *Eupenicillium levitum* (Raper & Fennell) Stolk & D.B. Scott, Persoonia 4: 402. 1967 ≡ *Eupenicillium javanicum* var. *levitum* (Raper & Fennell) Stolk & Samson, Stud. Mycol. 23: 134. 1983. [MB289096]. — Herb.: IMI 039735. Ex-type: CBS 345.48 = ATCC 10464 = IFO 6101 = IFO 8849 = IMI 039735 = NRRL 705 = QM 1877. Section *Lanata-Divaricata*. ITS barcode: GU981607. (Alternative markers: *BenA* = GU981654; *RPB2* = KF296432; *CaM* = KF296394).
- Penicillium lilacinochinulatum** S. Abe ex G. Sm., Trans. Brit. Mycol. Soc. 46: 335. 1963 ≡ *Penicillium lilacinochinulatum* S. Abe, J. Gen. Appl. Microbiol., Tokyo 2: 54. 1956 (nom. inval., Art. 36). [MB120793]. — Herb.: unknown. Ex-type: CBS 454.93 = ATCC 18309 = FAT 84 = FRR 3451 = IFO 6231 = IMI 068211 = QM 7289. Section *Sclerotiora*. ITS barcode: AY157489. (Alternative markers: *BenA* = KC773790; *RPB2* = n.a.; *CaM* = KC773816).
- Penicillium limosum** S. Ueda, Mycoscience 36: 451. 1995 ≡ *Eupenicillium limosum* S. Ueda, Mycoscience 36: 451. 1995. [MB415136]. — Herb.: CBM NEI-5220. Ex-type: CBS 339.97 = NEI5220. Section *Lanata-Divaricata*. ITS barcode: GU981568. (Alternative markers: *BenA* = GU981621; *RPB2* = KF296433; *CaM* = KF296398).
- Penicillium lineolatum** Udagawa & Y. Horie, Mycotaxon 5: 493. 1977 ≡ *Eupenicillium lineolatum* Udagawa & Y. Horie, Mycotaxon 5: 493. 1977 ≡ *Eupenicillium javanicum* var. *lineolatum* (Udagawa & Y. Horie) Stolk & Samson, Stud. Mycol. 23: 134. 1983. [MB319283]. — Herb.: NHL 2776. Ex-type: CBS 188.77 = NHL 2776. Section *Lanata-Divaricata*. ITS barcode: GU981579. (Alternative markers: *BenA* = GU981620; *RPB2* = KF296434; *CaM* = KF296397).
- Penicillium lividum** Westling, Ark. Bot. 11: 134. 1911. [MB178817]. — Herb.: IMI 39736. Ex-type: CBS 347.48 = ATCC 10102 = CCRC 31286 = DSM1180 = IFO 6102 = IMI 039736 = NRRL 754 = QM 1930 = VKMF-303. Section *Aspergilloides*. ITS barcode: KM189582. (Alternative markers: *BenA* = KF088825; *RPB2* = KM089598; *CaM* = KF089211).
- Penicillium longicatenatum** Visagie, et al., Stud. Mycol. 78: 429. 2014. [MB809968]. — Herb.: CBS H-21875. Ex-type: CBS 137735 = DTO 180-D9 = CV 2847 = DAOM 241119. Section *Aspergilloides*. ITS barcode: KM189636. (Alternative markers: *BenA* = KM088880; *RPB2* = KM089654; *CaM* = KF089267).
- Penicillium longisporum** (W.B. Kend.) Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Thysanophora longispora* W.B. Kend., Can. J. Bot. 39: 826. 1961. [MB561966]. — Herb.: unknown. Ex-type: CBS 354.62 = DAOM 63073 = MUCL 4168. Section *Thysanophora*. ITS barcode: n.a. (Alternative markers: *BenA* = KJ834467; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium ludwigii** Udagawa, Trans. Mycol. Soc. Japan 10: 2. 1969 ≡ *Eupenicillium ludwigii* Udagawa, Trans. Mycol. Soc. Japan 10: 2. 1969. [MB335744]. — Herb.: NHL 6118. Ex-type: CBS 417.68 = FRR 559. Section *Lanata-Divaricata*. ITS barcode: KF296409. (Alternative markers: *BenA* = KF296468; *RPB2* = KF296435; *CaM* = KF296404).

- Penicillium macleanniae*** H.Y. Yip, Trans. Brit. Mycol. Soc. 77: 202. 1981. [MB112523]. — Herb.: DAR 35238. Ex-type: CBS 198.81 = DAR35238. Section *Exilicaulis*. ITS barcode: KC411689. (Alternative markers: *BenA* = KJ834468; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium macrosclerotiorum*** L. Wang, X.M. Zhang & W.Y. Zhuang, Mycol. Res. 111: 1244. 2007. [MB492622]. — Herb.: HMAS 133177-1-4. Ex-type: CBS 116871 = AS 3.6581. Section *Gracilenta*. ITS barcode: KJ834511. (Alternative markers: *BenA* = KJ834469; *RPB2* = JN121432; *CaM* = DQ911123).
- Penicillium madriti*** G. Sm., Trans. Brit. Mycol. Soc. 44: 44. 1961. [MB335747]. — Herb.: IMI 86563. Ex-type: CBS 347.61 = ATCC 18233 = CCRC 31672 = FRR 3452 = IFO 9148 = IMI 086563 = LSHBBB389 = MUCL 2456 = MUCL 31193 = NRRL 3452 = QM 7959. Section *Turbata*. ITS barcode: AF033482. (Alternative markers: *BenA* = KJ834470; *RPB2* = JN406561; *CaM* = EU644076).
- Penicillium magnielliptisporum*** Visagie, Seifert & Samson, Stud. Mycol. 78: 127. 2014. [MB809186]. — Herb.: CBS H-21806. Ex-type: CBS 138225 = DTO 128H8. Section *Paradoxa*. ITS barcode: KJ775686. (Alternative markers: *BenA* = KJ775179; *RPB2* = n.a.; *CaM* = KJ775413).
- Penicillium malacaense*** C. Ramírez & A.T. Martínez, Mycopathologia 72: 186. 1980. [MB113025]. — Herb.: IJFM 7093. Ex-type: CBS 160.81 = NRRL 35754 = ATCC 42241 = IJFM 7093 = IMI 253801 = VKMF-2197. Section *Cinnamopurpurea*. ITS barcode: EU427300. (Alternative markers: *BenA* = EU427268; *RPB2* = JN406626; *CaM* = KJ866997).
- Penicillium malachiteum*** (Yaguchi & Udagawa) Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Chromocleista malachitea* Yaguchi & Udagawa, Trans. Mycol. Soc. Japan 34: 102. 1993. [MB561971]. — Herb.: CBS 647.95. Ex-type: CBS 647.95 = IBT 17515. Section *Sclerotiora*. ITS barcode: KC773838. (Alternative markers: *BenA* = KC773794; *RPB2* = n.a.; *CaM* = KC773820).
- Penicillium mallochii*** K.G. Rivera, Urb & Seifert, Mycetaxon 119: 322. 2012. [MB563043]. — Herb.: DAOM 239917. Ex-type: CCFC 239917. Section *Sclerotiora*. ITS barcode: JN626104. (Alternative markers: *BenA* = JN625973; *RPB2* = n.a.; *CaM* = JN626016).
- Penicillium malmesburiense*** Visagie, Houbraken & K. Jacobs, Stud. Mycol. 78: 429. 2014. [MB809969]. — Herb.: CBS H-21872. Ex-type: CBS 137744 = DTO 182-H5 = CV 1180 = DAOM 241144. Section *Aspergilloides*. ITS barcode: KM189676. (Alternative markers: *BenA* = KM089821; *RPB2* = KM089695; *CaM* = KM089308).
- Penicillium malodoratum*** (Kwon-Chung & Fennell) Samson et al., (published here) ≡ *Aspergillus malodoratus* Kwon-Chung & Fennell, Gen. *Aspergillus*: 468. 1965. [MB809316]. — Herb.: IMI 172289. Ex-type: CBS 490.65 = NRRL 5083 = IMI 172289 = ATCC 16834. Section *Paradoxa*. ITS barcode: AF033485. (Alternative markers: *BenA* = EF669681; *RPB2* = EF669672; *CaM* = FJ530972).
- Penicillium manginii*** Duché & R. Heim, Trav. Cryptog.: 450. 1931. [MB270490]. — Herb.: CBS 253.31. Ex-type: CBS 253.31 = NRRL 2134. Section *Citrina*. ITS barcode: GU944599. (Alternative markers: *BenA* = JN606651; *RPB2* = JN606618; *CaM* = JN606381).
- Penicillium mariae-crucis*** Quintan., Av. Aliment. Majora Anim. 23: 334. 1982. [MB114171]. — Herb.: CBS 270.83. Ex-type: CBS 271.83 = IMI 256075. Section *Lanata-Divaricata*. ITS barcode: GU981593. (Alternative markers: *BenA* = GU981630; *RPB2* = KF296439; *CaM* = KF296374).
- Penicillium marinum*** Frisvad & Samson, Stud. Mycol. 49: 20. 2004. [MB370974]. — Herb.: CBS 109550. Ex-type: CBS 109550 = IBT 14360. Section *Penicillium*. ITS barcode: KJ834512. (Alternative markers: *BenA* = AY674392; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium maximae*** Visagie, Houbraken & Samson, Persoonia 31: 52. 2013. [MB803783]. — Herb.: CBS H-21144. Ex-type: CBS 134565 = NRRL 2060. Section *Sclerotiora*. ITS barcode: EU427298. (Alternative markers: *BenA* = KC773795; *RPB2* = n.a.; *CaM* = KC773821).
- Penicillium melanoconidium*** (Frisvad) Frisvad & Samson, Stud. Mycol. 49: 28. 2004 ≡ *Penicillium aurantiogriseum* var. *melanoconidium* Frisvad, Mycologia 81: 849. 1989. [MB368219]. — Herb.: IMI 321503. Ex-type: CBS 115506 = IBT 3444 = IMI 321503. Section *Fasciculata*. ITS barcode: AJ005483. (Alternative markers: *BenA* = AY674304; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium melinii*** Thom, Penicillia: 273. 1930. [MB270876]. — Herb.: IMI 40216. Ex-type: CBS 218.30 = ATCC 10469 = FRR 2041 = IFO 7675 = IMI 040216 = MUCL 29235 = NRRL 2041 = QM 7599. Section *Exilicaulis*. ITS barcode: AF033449. (Alternative markers: *BenA* = KJ834471; *RPB2* = JN406613; *CaM* = n.a.).
- Penicillium meloforme*** Udagawa & Y. Horie, Trans. Mycol. Soc. Japan 14: 376. 1973 ≡ *Eupenicillium meloforme* Udagawa & Y. Horie, Trans. Mycol. Soc. Japan 14: 376. 1973 ≡ *Eupenicillium javanicum* var. *meloforme* (Udagawa & Horie) Stolk & Samson, Stud. Mycol. 23: 136. 1983. [MB120882]. — Herb.: NHL 6468. Ex-type: CBS 445.74 = ATCC 28049 = IMI 216903 = NHL 6468. Section *Lanata-Divaricata*. ITS barcode: KC411762. (Alternative markers: *BenA* = GU981656; *RPB2* = KF296440; *CaM* = KF296396).
- Penicillium menonorum*** S.W. Peterson, IMA Fungus 2: 122. 2011. [MB519297]. — Herb.: BPI 881018. Ex-type: NRRL 50410. Section *Exilicaulis*. ITS barcode: HQ646591. (Alternative markers: *BenA* = HQ646573; *RPB2* = n.a.; *CaM* = HQ646584).
- Penicillium meridianum*** D.B. Scott, Mycopathol. Mycol. Appl. 36: 12. 1968 ≡ *Eupenicillium meridianum* D.B. Scott, Mycopathol. Mycol. Appl. 36: 12. 1968. [MB335750]. — Herb.: CBS 314.67. Ex-type: CBS 314.67 = ATCC 18545 = CSIR 1052 = IMI 136209. Section *Exilicaulis*. ITS barcode: AF033451. (Alternative markers: *BenA* = KJ834472; *RPB2* = JN406576; *CaM* = n.a.).
- Penicillium mexicanum*** Visagie, Seifert & Samson, Stud. Mycol. 78: 125. 2014. [MB809185]. — Herb.: CBS H-21805. Ex-type: CBS 138227 = DTO 270F1. Section *Paradoxa*. ITS barcode: KJ775685. (Alternative markers: *BenA* = KJ775178; *RPB2* = n.a.; *CaM* = KJ775412).
- Penicillium miczynskii*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 482. 1927. [MB271171]. — Herb.: IMI 40030. Ex-type: CBS 220.28 = ATCC 10470 = DSM2437 = FRR 1077 = IFO 7730 = IMI 040030 = MUCL 29228 = NRRL 1077 = QM 1957. Section *Citrina*. ITS barcode: GU944600. (Alternative markers: *BenA* = JN606706; *RPB2* = JN606623; *CaM* = JN606526).
- Penicillium mononematosum*** (Frisvad, Filt. & Wicklow) Frisvad, Mycologia 81: 857. 1990 ≡ *Penicillium glandicola* var. *mononematosum* Frisvad, Filt. & Wicklow, Can. J. Bot. 65: 767. 1987. [MB126406]. — Herb.: IMI 296925. Ex-type: CBS 172.87 = IBT 3072 = IBT 5518 = IBT 21535 = IMI 296925 = NRRL 13482 = NRRL A-26709. Section *Chrysogena*. ITS barcode: JX997082. (Alternative markers: *BenA* = AY495997; *RPB2* = JX996709; *CaM* = JX996964).
- Penicillium montanense*** M. Chr. & Backus, Mycologia 54: 574. 1962. [MB335752]. — Herb.: WIS Cryptogamic Herb. No. GW1-6. Ex-type: CBS 310.63 = ATCC 14941 = FRR 3407 = IFO 7740 = IHEM 4375 = IMI 099468 = MUCL 31326 = NRRL 3407. Section *Aspergilloides*. ITS barcode: KM189551. (Alternative markers: *BenA* = KM088789; *RPB2* = KM089561; *CaM* = KM089174).
- Penicillium multicolor*** Grig.-Man. & Porad., Arch. des Sciences Biol. Leningrad 19: 120. 1915. [MB119288]. — Herb.: unknown. Ex-type: CBS 501.73 = ATCC 24723 = IMI 174716 = VKM F-1745. Section *Charlesia*. ITS barcode: KC790402. (Alternative markers: *BenA* = JN799645; *RPB2* = EU427262; *CaM* = JN799646).
- Penicillium nalgiovense*** Laxa, Zentralbl. Bakteriol. Parasitenk., Abt. 2 86: 160. 1932. [MB114239]. — Herb.: CBS 352.48. Ex-type: CBS 352.48 = ATCC 10472 = IBT 21536 = IMI 039804 = MUCL 31194 = NRRL 911. Section *Chrysogena*. ITS barcode: AY371617. (Alternative markers: *BenA* = AY495999; *RPB2* = JX996719; *CaM* = JX996974).
- Penicillium namyslowskii*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Cl. Sci. Math., Sér. B, Sci. Nat. 1927: 479. 1927 ≡ *Geosmithia namyslowskii* (K.M. Zalessky) Pitt, Can. J. Bot. 57: 2024. 1979. [MB272006]. — Herb.: CBS 353.48. Ex-type: CBS 353.48 = ATCC 11127 = IMI 040033 = MUCL 29226 = NRRL 1070. Section *Exilicaulis*. ITS barcode: AF033463. (Alternative markers: *BenA* = JX141067; *RPB2* = JF417430; *CaM* = n.a.).
- Penicillium neocrassum*** R. Serra & S.W. Peterson, Mycologia 99: 81. 2007. [MB504767]. — Herb.: BPI 872161. Ex-type: CBS 122428 = NRRL 35639 = MUM 06.160. Section *Brevicompacta*. ITS barcode: DQ645805. (Alternative markers: *BenA* = DQ645794; *RPB2* = JN406633; *CaM* = DQ645809).
- Penicillium neoechinulatum*** (Frisvad, Filt. & Wicklow) Frisvad & Samson, Stud. Mycol. 49: 28. 2004 ≡ *Penicillium aurantiogriseum* var. *neoechinulatum* Frisvad, Filt. & Wicklow, Can. J. Bot. 65: 767. 1987. [MB368218]. — Herb.: IMI 296937. Ex-type: CBS 169.87 = CBS 101135 = IBT 3493 = IBT 21537 = IMI 296937 = NRRL 13486. Section *Fasciculata*. ITS barcode: JN942722. (Alternative markers: *BenA* = AF003237; *RPB2* = JN985406; *CaM* = n.a.).
- Penicillium neomiczynskii*** A.L.J. Cole et al., Stud. Mycol. 70: 105. 2011. [MB563192]. — Herb.: CBS H-20661. Ex-type: CBS 126231 = IBT 23560. Section *Citrina*. ITS barcode: JN617671. (Alternative markers: *BenA* = JN606705; *RPB2* = n.a.; *CaM* = JN606523).
- Penicillium nepalense*** Takada & Udagawa, Trans. Mycol. Soc. Japan 24: 146. 1983 ≡ *Eupenicillium nepalense* Takada & Udagawa, Trans. Mycol. Soc. Japan 24: 146. 1983. [MB108327]. — Herb.: NHL 6482. Ex-type: CBS 203.84 = NHL 6482. Section *Exilicaulis*. ITS barcode: KC411692. (Alternative markers: *BenA* = KJ834474; *RPB2* = JN121453; *CaM* = n.a.).

- Penicillium nodulum** H.Z. Kong & Z.T. Qi, Mycosistema 1: 108. 1988. [MB135445]. — Herb.: CBS 227.89. Ex-type: CBS 227.89. Section *Cinnamopurpurea*. ITS barcode: KC411703. (Alternative markers: *BenA* = KJ834475; *RPB2* = JN406603; *CaM* = KJ867003).
- Penicillium nordicum** Dragoni & Cantoni ex C. Ramirez, Adv. *Penicillium Aspergillus* Syst.: 139. 1986 ≡ *Penicillium nordicum* Dragoni & Cantoni, Ind. Aliment. 155: 283. 1979 (nom. inval., Art. 36). [MB114762]. — Herb.: ATCC 44219. Ex-type: ATCC 44219 = IBT 13307. Section *Fasciculata*. ITS barcode: KJ834513. (Alternative markers: *BenA* = KJ834476; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium nothofagi** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 105. 2011. [MB563189]. — Herb.: CBS H-20655. Ex-type: CBS 130383 = IBT 23018 = DTO 76C2. Section *Citrina*. ITS barcode: JN617712. (Alternative markers: *BenA* = JN606732; *RPB2* = n.a.; *CaM* = JN606507).
- Penicillium novae-zealandiae** J.F.H. Beyma, Antonie van Leeuwenhoek 6: 275. 1940. [MB522253]. — Herb.: IMI 40584ii. Ex-type: CBS 137.41 = ATCC 10473 = IFO 31748 = IMI 040584ii = NRRL 2128 = QM 1934 = VKMF-2886. Section *Canescens*. ITS barcode: JN617688. (Alternative markers: *BenA* = KJ834477; *RPB2* = JN406628; *CaM* = KJ866996).
- Penicillium ochrochloron** Biourge, Cellule 33: 269. 1923. [MB272701]. — Herb.: IMI 39806. Ex-type: CBS 357.48 = ATCC 10540 = IMI 039806 = NRRL 926 = QM 7604. Section *Lanata-Divaricata*. ITS barcode: GU981604. (Alternative markers: *BenA* = GU981672; *RPB2* = KF296445; *CaM* = KF296378).
- Penicillium ochrosalmoneum** Udagawa, J. Agric. Sci. Tokyo Nogyo Daig. 5: 10. 1959 ≡ *Eupenicillium ochrosalmoneum* D.B. Scott & Stolk, Antonie van Leeuwenhoek 33: 302. 1967. [MB302409]. — Herb.: NHL 6048. Ex-type: CBS 489.66 = ATCC 18338 = CSIR 145 = IMI 116248ii = NRRL 35499. Section *Ochrosalmonea*. ITS barcode: EF626961. (Alternative markers: *BenA* = EF506212; *RPB2* = JN121524; *CaM* = EF506237).
- Penicillium odoratum** M. Chr. & Backus, Mycologia 53: 459. 1961. [MB335755]. — Herb.: WSF 2000. Ex-type: CBS 294.62 = CBS 296.62 = ATCC 14769 = DSM2419 = IFO 7741 = IMI 094208ii = NRRL 3007 = WSF2000. Section *Aspergilloides*. ITS barcode: KC411730. (Alternative markers: *BenA* = KJ834478; *RPB2* = JN406583; *CaM* = KM089363).
- Penicillium olsonii** Bainier & Sartory, Ann. Mycol. 10: 398. 1912. [MB121021]. — Herb.: IMI 192502. Ex-type: CBS 232.60 = IBT 23473 = IMI 192502 = NRRL 13058 = NRRL 13716. Section *Brevicompacta*. ITS barcode: EU587341. (Alternative markers: *BenA* = AY674445; *RPB2* = JN121464; *CaM* = DQ658165).
- Penicillium onobense** C. Ramírez & A.T. Martínez, Mycopathologia 74: 44. 1981. [MB112525]. — Herb.: CBS 174.81. Ex-type: CBS 174.81 = ATCC 42225 = IJFM 3026 = VKMF-2183. Section *Lanata-Divaricata*. ITS barcode: GU981575. (Alternative markers: *BenA* = GU981627; *RPB2* = KF296447; *CaM* = KF296371).
- Penicillium ornatum** Udagawa, Trans. Mycol. Soc. Japan 9: 49. 1968 ≡ *Eupenicillium ornatum* Udagawa, Trans. Mycol. Soc. Japan 9: 49. 1968. [MB335756]. — Herb.: NHL 6101. Ex-type: CBS 190.68 = ATCC 18608 = IFO 31739 = IMI 137977 = NHL 6101 = NRRL 3471. Section *Ramigena*. ITS barcode: KC411687. (Alternative markers: *BenA* = KJ834479; *RPB2* = JN121451; *CaM* = n.a.).
- Penicillium osmophilum** Stolk & Veenb.-Rijks, Antonie van Leeuwenhoek 40: 1. 1974 ≡ *Eupenicillium osmophilum* Stolk & Veenb.-Rijks, Antonie van Leeuwenhoek 40: 1. 1974. [MB319288]. — Herb.: CBS 462.72. Ex-type: CBS 462.72 = CBS 439.73 = IBT 14678 = NRRL 5922. Section *Fasciculata*. ITS barcode: EU427295. (Alternative markers: *BenA* = AY674376; *RPB2* = JN121518; *CaM* = n.a.).
- Penicillium ovatum** (K. Ando & Nawawi) Houbraken & Samson, Stud. Mycol. 70: 48. 2011 ≡ *Torulomyces ovatus* K. Ando & Nawawi, Mycoscience 39: 317. 1998. [MB561961]. — Herb.: TNS-F-238518. Ex-type: CBS 136664 = KY 12726. Section *Exilicaulis*. ITS barcode: KF667370. (Alternative markers: *BenA* = KF667366; *RPB2* = KF667372; *CaM* = KF667368).
- Penicillium oxalicum** Currie & Thom, J. Biol. Chem. 22: 289. 1915. [MB121033]. — Herb.: IMI 192332. Ex-type: CBS 219.30 = ATCC 1126 = FRR 787 = IMI 192332 = MUCL 29047 = NRRL 787 = QM 7606. Section *Lanata-Divaricata*. ITS barcode: AF033438. (Alternative markers: *BenA* = KF296462; *RPB2* = JN121456; *CaM* = KF296367).
- Penicillium palitans** Westling, Ark. Bot. 11: 83. 1911. [MB203604]. — Herb.: CBS H-7531. Ex-type: CBS 107.11 = ATCC 10477 = IBT 23034 = IMI 040215 = NRRL 2033 = VKMF-3088. Section *Fasciculata*. ITS barcode: KJ834514. (Alternative markers: *BenA* = KJ834480; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium palmense** C. Ramírez & A.T. Martínez, Mycopathologia 66: 80. 1978. [MB319289]. — Herb.: IJFM 3840. Ex-type: CBS 336.79 = ATCC 38669 = IJFM 3840. Section *Aspergilloides*. ITS barcode: KJ834515. (Alternative markers: *BenA* = GQ367508; *RPB2* = JN406566; *CaM* = GQ367534).
- Penicillium pancosmium** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 108. 2011. [MB563191]. — Herb.: CBS H-20651. Ex-type: CBS 276.75 = DAOM 147467 = IBT 29991. Section *Citrina*. ITS barcode: JN617660. (Alternative markers: *BenA* = JN606790; *RPB2* = n.a.; *CaM* = JN606446).
- Penicillium paneum** Frisvad, Microbiology 142: 546. 1996. [MB415570]. — Herb.: C 25000. Ex-type: CBS 101032 = IBT 21541 = IBT 12407. Section *Roquefortorum*. ITS barcode: HQ442346. (Alternative markers: *BenA* = AY674387; *RPB2* = n.a.; *CaM* = HQ442331).
- Penicillium paradoxum** (Fennell & Raper) Samson et al., (published here) ≡ *Aspergillus paradoxus* Fennell & Raper, Mycologia 47: 69. 1955 ≡ *Hemicarpenteles paradoxus* Sarbhoy & Elphick, Trans. Brit. Mycol. Soc. 51: 156. 1968. [MB547045]. — Herb.: IMI 061446. Ex-type: CBS 527.65 = NRRL 2162 = ATCC 16918 = IMI 061446. Section *Paradoxa*. ITS barcode: EF669707. (Alternative markers: *BenA* = EF669683; *RPB2* = EF669670; *CaM* = EF669692).
- Penicillium paraherquei** S. Abe ex G. Sm., Trans. Brit. Mycol. Soc. 46: 335. 1963 ≡ *Penicillium paraherquei* S. Abe, J. Gen. Appl. Microbiol., Tokyo 2: 131. 1956 (nom. inval., Art. 36). [MB302412]. — Herb.: IMI 68220. Ex-type: CBS 338.59 = ATCC 22354 = ATCC 46903 = FAT964 = FRR 3454 = IFO 6234 = IMI 068220 = NRRL 3454. Section *Lanata-Divaricata*. ITS barcode: AF178511. (Alternative markers: *BenA* = KF296465; *RPB2* = KF296449; *CaM* = KF296372).
- Penicillium parvulum** S.W. Peterson & B.W. Horn, Mycologia 101: 75. 2009. [MB509289]. — Herb.: BPI 877331. Ex-type: CBS 132825 = NRRL 35504. Section *Cinnamopurpurea*. ITS barcode: EF422845. (Alternative markers: *BenA* = EF506218; *RPB2* = n.a.; *CaM* = EF506225).
- Penicillium parvum** Raper & Fennell, Mycologia 40: 508. 1948 ≡ *Carpenteles parvum* (Raper & Fennell) Udagawa, Trans. Mycol. Soc. Japan 6: 79. 1965 ≡ *Eupenicillium parvum* (Raper & Fennell) Stolk & D.B. Scott, Persoonia 4: 402. 1967. [MB289101]. — Herb.: CBS H-7105. Ex-type: CBS 359.48 = ATCC 10479 = IFO 7732 = IMI 040587 = NRRL 2095 = QM 1878. Section *Exilicaulis*. ITS barcode: AF033460. (Alternative markers: *BenA* = HQ646568; *RPB2* = JN406559; *CaM* = KF900173).
- Penicillium pasqualense** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 108. 2011. [MB563190]. — Herb.: CBS H-20663. Ex-type: CBS 126330 = IBT 14235. Section *Citrina*. ITS barcode: JN617676. (Alternative markers: *BenA* = JN606673; *RPB2* = JN606617; *CaM* = JN606394).
- Penicillium patens** Pitt & A.D. Hocking, Mycotaxon 22: 205. 1985. [MB105611]. — Herb.: FRR 2661. Ex-type: CBS 260.87 = FRR 2662. Section *Aspergilloides*. ITS barcode: KC411718. (Alternative markers: *BenA* = KJ834481; *RPB2* = JN406593; *CaM* = n.a.).
- Penicillium paxilli** Bainier, Bull. Soc. Mycol. France 23: 95. 1907. [MB203838]. — Herb.: IMI 40226. Ex-type: CBS 360.48 = ATCC 10480 = FRR 2008 = IMI 040226 = NRRL 2008 = QM 725. Section *Citrina*. ITS barcode: GU944577. (Alternative markers: *BenA* = JN606844; *RPB2* = JN606610; *CaM* = JN606566).
- Penicillium penarolense** Houbraken et al., Int. J. Syst. Evol. Microbiol. 61: 1471. 2011. [MB518024]. — Herb.: HUA 170335. Ex-type: CBS 113178 = IBT 23262. Section *Lanata-Divaricata*. ITS barcode: GU981570. (Alternative markers: *BenA* = GU981646; *RPB2* = KF296450; *CaM* = KF296381).
- Penicillium persicinum** L. Wang et al., Antonie van Leeuwenhoek 86: 177. 2004. [MB500259]. — Herb.: HMAS 80638-1-4. Ex-type: CBS 111235 = AS 3.5891 = IBT 24565. Section *Chrysogena*. ITS barcode: JX997072. (Alternative markers: *BenA* = AY495982; *RPB2* = JN406644; *CaM* = JX996954).
- Penicillium philippinense** Udagawa & Y. Horie, J. Jap. Bot. 47: 341. 1972. [MB319291]. — Herb.: CBS H-7099. Ex-type: CBS 623.72 = FRR 1532 = NHL 6130. Section *Exilicaulis*. ITS barcode: KC411770. (Alternative markers: *BenA* = KJ834482; *RPB2* = JN406543; *CaM* = n.a.).
- Penicillium phoeniceum** J.F.H. Beyma, Zentralbl. Bakteriol. Parasitenk., Abt. 2 88: 136. 1933 ≡ *Eupenicillium cinnamopurpureum* D.B. Scott & Stolk, Antonie van Leeuwenhoek 33: 308. 1967. [MB274284]. — Herb.: IMI 40585. Ex-type: CBS 249.32 = ATCC 10481 = IJFM 5122 = IMI 040585 = NRRL 2070 = QM 7608 = VKMF-321. Section *Charlesia*. ITS barcode: KC411711. (Alternative markers: *BenA* = KJ834483; *RPB2* = JN406597; *CaM* = AY741729).
- Penicillium pimeteouicense** S.W. Peterson, Mycologia 91: 271. 1999. [MB460126]. — Herb.: BPI 806262. Ex-type: CBS 102479 = NRRL 25542. Section *Exilicaulis*. ITS barcode: AF037431. (Alternative markers: *BenA* = HQ646569; *RPB2* = JN406650; *CaM* = HQ646580).
- Penicillium piscarium** Westling, Ark. Bot. 11: 86. 1911. [MB211321]. — Herb.: IMI 40032. Ex-type: CBS 362.48 = ATCC 10482 = FRR 1075 = IFO 8111 = IMI 040032 = NRRL 1075 = VKMF-1823. Section *Lanata-Divaricata*.

- ITS barcode: GU981600. (Alternative markers: *BenA* = GU981668; *RPB2* = KF296451; *CaM* = KF296379).
- Penicillium polonicum*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B, Sci. Nat. 1927: 445. 1927 ≡ *Penicillium aurantiogriseum* var. *polonicum* (K.M. Zalessky) Frisvad & Filz., Mycologia 81: 850. 1990. [MB274889]. — Herb.: CBS 222.28. Ex-type: CBS 222.28 = IBT 12821 = IMI 291194 = MUCL 29204 = NRRL 995. Section *Fasciculata*. ITS barcode: AF033475. (Alternative markers: *BenA* = AY674305; *RPB2* = JN406609; *CaM* = n.a.).
- Penicillium porphyreum*** Houben & Samson, Stud. Mycol. 70: 48. 2011 ≡ *Monocillium humicola* G.L. Barron, Can. J. Bot. 39: 1575. 1961 ≡ *Monocillium humicola* var. *brunneum* M. Chr. & Backus, Mycologia 56: 498. 1964 ≡ *Torulomyces brunneus* (M. Chr. & Backus) K. Ando, Mycoscience 39: 314. 1998. [MB561959]. — Herb.: CBS 382.64 = WSF 9-c. Ex-type: CBS 382.64 = KY 12723. Section *Torulomyces*. ITS barcode: KF303666. (Alternative markers: *BenA* = KF303621; *RPB2* = KF303677; *CaM* = KF303636).
- Penicillium psychosexualis*** Houben & Samson, IMA Fungus 1: 174. 2010. [MB519086]. — Herb.: CBS H-20501. Ex-type: CBS 128137 = IBT 29551. Section *Roquefortorum*. ITS barcode: HQ442345. (Alternative markers: *BenA* = HQ442356; *RPB2* = n.a.; *CaM* = HQ442330).
- Penicillium pullum*** S.W. Peterson & Sigler, Mycol. Res. 106: 1115. 2002. [MB483982]. — Herb.: BPI 841398. Ex-type: CBS 331.48 = ATCC 10447 = NRRL 721 = FRR 721 = IFO 6097 = IMI 39747 = QM 1925 = Thom 5179.4. Section *Stolka*. ITS barcode: AF033443. (Alternative markers: *BenA* = JN617719; *RPB2* = n.a.; *CaM* = AF481134).
- Penicillium pulvillorum*** Turfitt, Trans. Brit. Mycol. Soc. 23: 186. 1939. [MB275682]. — Herb.: CBS 280.39. Ex-type: CBS 280.39 = IFO 7763 = NRRL 2026. Section *Lanata-Divaricata*. ITS barcode: AF178517. (Alternative markers: *BenA* = GU981670; *RPB2* = KF296452; *CaM* = KF296377).
- Penicillium pulvis*** Houben, Visagie, Samson & Seifert, Stud. Mycol. 78: 429. 2014. [MB809970]. — Herb.: CBS H-21878. Ex-type: CBS 138432 = DTO 180-B7. Section *Aspergilloides*. ITS barcode: KM189632. (Alternative markers: *BenA* = KM088876; *RPB2* = KM089650; *CaM* = KM089263).
- Penicillium purpurescens*** (Sopp) Raper & Thom, A manual of the Penicillia: 177. 1949 ≡ *Citromyces purpurascens* Sopp, Skr. Vidensk.-Selsk. Christiana, Math.-Naturvidensk. Kl. 11: 117. 1912. [MB335761]. — Herb.: IMI 39745. Ex-type: CBS 366.48 = NRRL 720 = FRR 720 = ATCC 10485 = IMI 39745. Section *Aspergilloides*. ITS barcode: KM189561. (Alternative markers: *BenA* = KM088801; *RPB2* = KM089573; *CaM* = KM089186).
- Penicillium quebecense*** Houben, Frisvad & Samson, Stud. Mycol. 70: 111. 2011. [MB563202]. — Herb.: CBS H-20666. Ex-type: CBS 101623 = IBT 29050. Section *Citrina*. ITS barcode: JN617661. (Alternative markers: *BenA* = JN606700; *RPB2* = JN606622; *CaM* = JN606509).
- Penicillium queretorum*** Baghd., Novosti Sist. Nizsh. Rast. 5: 110. 1968. [MB335762]. — Herb.: CBS H-7527. Ex-type: CBS 417.69 = NRRL 3758 = ATCC 48727 = CCRC 31668 = FRR 516 = IFO 31749 = IMI 140342 = MUCL 31203 = VKMF-1074. Section *Aspergilloides*. ITS barcode: KM189556. (Alternative markers: *BenA* = KM088795; *RPB2* = KM089567; *CaM* = KM089180).
- Penicillium raciborskii*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B, Sci. Nat. 1927: 454. 1927. [MB276002]. — Herb.: IMI 40568. Ex-type: CBS 224.28 = ATCC 10488 = DSM2422 = FRR 2150 = IFO 7676 = IMI 040568 = LSHBP 92 = MUCL 29246 = NRRL 2150. Section *Exilicaulis*. ITS barcode: AF033447. (Alternative markers: *BenA* = JX141069; *RPB2* = JN406607; *CaM* = n.a.).
- Penicillium radicicola*** Overy & Frisvad, Syst. Appl. Microbiol. 26: 633. 2003. [MB488233]. — Herb.: C 00161. Ex-type: CBS 112430 = IBT 10696. Section *Fasciculata*. ITS barcode: KJ834516. (Alternative markers: *BenA* = AY674357; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium raistrickii*** G. Sm., Trans. Brit. Mycol. Soc. 18: 90. 1933. [MB276069]. — Herb.: IMI 40221. Ex-type: CBS 261.33 = ATCC 10490 = FRR 1044 = IFO 6104 = IMI 040221 = LSHBB100 = NRRL 1044 = NRRL 2039 = QM 1936 = VKMF-337. Section *Ramosa*. ITS barcode: AY373927. (Alternative markers: *BenA* = KJ834485; *RPB2* = JN406592; *CaM* = KJ867006).
- Penicillium ramuscum*** Bat. & H. Maia, Anais Soc. Biol. Pernambuco 13: 27. 1955. [MB302419]. — Herb.: unknown. Ex-type: CBS 251.56 = ATCC 12292 = FRR 3459 = IMI 063546 = IMUR478 = LSHBB324 = NRRL 3459 = QM 7057. Section *Ramigena*. ITS barcode: EF433765. (Alternative markers: *BenA* = EU427269; *RPB2* = JN121472; *CaM* = EU427278).
- Penicillium ranomafanaense*** Houben & Hagen, Stud. Mycol. 78: 433. 2014. [MB809971]. — Herb.: CBS H-21862. Ex-type: CBS 137953 = DTO 085-A5. Section *Aspergilloides*. ITS barcode: KM189541. (Alternative markers: *BenA* = KM088779; *RPB2* = KM089551; *CaM* = KM089164).
- Penicillium raperi*** G. Sm., Trans. Brit. Mycol. Soc. 40: 486. 1957. [MB302421]. — Herb.: IMI 71625. Ex-type: CBS 281.58 = ATCC 22355 = IFO 8179 = IMI 071625 = LSHBB338 = NRRL 2674 = QM 7527. Section *Lanata-Divaricata*. ITS barcode: AF033433. (Alternative markers: *BenA* = GU981622; *RPB2* = KF296453; *CaM* = KF296399).
- Penicillium raphiae*** Houben, Frisvad & Samson, Stud. Mycol. 70: 114. 2011. [MB563203]. — Herb.: CBS H-20660. Ex-type: CBS 126234 = IBT 22407. Section *Citrina*. ITS barcode: JN617673. (Alternative markers: *BenA* = JN606657; *RPB2* = JN606619; *CaM* = JN606409).
- Penicillium restrictum*** J.C. Gilman & E.V. Abbott, Iowa St. Coll. J. Sci. 1: 297. 1927. [MB276289]. — Herb.: IMI 40228. Ex-type: CBS 367.48 = ATCC 11257 = FRR 1748 = IMI 040228 = NRRL 1748 = QM 1962. Section *Exilicaulis*. ITS barcode: AF033457. (Alternative markers: *BenA* = KJ834486; *RPB2* = JN121506; *CaM* = n.a.).
- Penicillium restingae*** J.P. Andrade et al., Persoonia 32: 293. 2014. [MB807051]. — Herb.: CMR H-12. Ex-type: URM 7075. Section *Sclerotiora*. ITS barcode: KF803355. (Alternative markers: *BenA* = KF803349; *RPB2* = n.a.; *CaM* = KF803352).
- Penicillium reticulisorum*** Udagawa, Trans. Mycol. Soc. Japan 9: 52. 1968 ≡ *Eupenicillium reticulisorum* Udagawa, Trans. Mycol. Soc. Japan 9: 52. 1968. [MB335763]. — Herb.: NHL 6105. Ex-type: CBS 122.68 = ATCC 18566 = IFO 9024 = IMI 136700 = NHL 6105 = NRRL 3447. Section *Lanata-Divaricata*. ITS barcode: AF033437. (Alternative markers: *BenA* = GU981665; *RPB2* = KF296454; *CaM* = KF296391).
- Penicillium ribium*** Frisvad & Overy, Int. J. Syst. Evol. Microbiol. 56: 1436. 2006. [MB501061]. — Herb.: DAOM 234091. Ex-type: CBS 127809 = DAOM 234091 = IBT 16537 = IBT 24431. Section *Ramosa*. ITS barcode: DQ267916. (Alternative markers: *BenA* = DQ285625; *RPB2* = JN406631; *CaM* = KJ866995).
- Penicillium rolfsii*** Thom, Penicillia: 489. 1930. [MB276674]. — Herb.: IMI 40029. Ex-type: CBS 368.48 = ATCC 10491 = FRR 1078 = IFO 7735 = IMI 040029 = MUCL 29229 = NRRL 1078 = QM 1961. Section *Lanata-Divaricata*. ITS barcode: JN617705. (Alternative markers: *BenA* = GU981667; *RPB2* = KF296455; *CaM* = KF296375).
- Penicillium roqueforti*** Thom, U.S.D.A. Bur. Animal Industr. Bull. 82: 35. 1906. [MB213525]. — Herb.: IMI 24313. Ex-type: CBS 221.30 = ATCC 10110 = ATCC 1129 = CECT 2905 = IBT 6754 = IFO 5459 = IMI 024313 = LSHBP 93 = NCTC 588 = NRRL 849 = QM 1937. Section *Roquefortorum*. ITS barcode: EU427296. (Alternative markers: *BenA* = AF000303; *RPB2* = JN406611; *CaM* = HQ442332).
- Penicillium roseomaculatum*** Biourge, Cellule 33: 301. 1923. [MB276785]. — Herb.: unknown. Ex-type: CBS 137962 = IMI 189696 = NRRL 728 = FRR 728. Section *Aspergilloides*. ITS barcode: KM189755. (Alternative markers: *BenA* = KM089004; *RPB2* = KM089778; *CaM* = KM089391).
- Penicillium roseopurpureum*** Dierckx, Ann. Soc. Sci. Bruxelles 25: 86. 1901. [MB213447]. — Herb.: IMI 40573. Ex-type: CBS 226.29 = ATCC 10492 = ATHUM2895 = FRR 2064 = IMI 040573 = MUCL 28654 = MUCL 29237 = NRRL 2064 = NRRL 2064A. Section *Citrina*. ITS barcode: GU944605. (Alternative markers: *BenA* = JN606838; *RPB2* = JN606613; *CaM* = JN606556).
- Penicillium roseoviride*** Stapp & Bortels, Zentralbl. Bakteriol. Parasitenk., Abt. 2 93: 51. 1935. [MB492646]. — Herb.: unknown. Ex-type: CBS 267.35 = ATCC 10412 = IFO 6089 = IMI 039740i = NRRL 760 = QM 7485. Section *Aspergilloides*. ITS barcode: KM189549. (Alternative markers: *BenA* = KM088787; *RPB2* = KM089559; *CaM* = KM089172).
- Penicillium rubefaciens*** Quintan., Mycopathologia 80: 73. 1982. [MB109998]. — Herb.: CBS 145.83. Ex-type: CBS 145.83 = CECT 2752. Section *Exilicaulis*. ITS barcode: KC411677. (Alternative markers: *BenA* = KJ834487; *RPB2* = JN406627; *CaM* = n.a.).
- Penicillium rubens*** Biourge, Cellule 33: 265. 1923. [MB276884]. — Herb.: CBS H-20595. Ex-type: CBS 129667 = NRRL 792 = IBT 30129 = ATCC 9783. Section *Chrysogena*. ITS barcode: JX997057. (Alternative markers: *BenA* = JF909949; *RPB2* = JX996658; *CaM* = JX996263).
- Penicillium rubidurum*** Udagawa & Y. Horie, Trans. Mycol. Soc. Japan 14: 381. 1973 ≡ *Eupenicillium rubidurum* Udagawa & Y. Horie, Trans. Mycol. Soc. Japan 14: 381. 1973. [MB319295]. — Herb.: NHL 6460. Ex-type: CBS 609.73 = NRRL 6033 = ATCC 28051 = ATCC 48238 = FRR 1558 = IMI 228551 = NHL 6460. Section *Exilicaulis*. ITS barcode: AF033462. (Alternative markers: *BenA* = HQ646574; *RPB2* = JN406545; *CaM* = HQ646585).
- Penicillium rudallense*** Houben, Visagie & Pitt, Stud. Mycol. 78: 433. 2014. [MB809972]. — Herb.: CBS H-21867. Ex-type: CBS 138162 = FRR 6085 = DTO 056-I4. Section *Aspergilloides*. ITS barcode: KM088741. (Alternative markers: *BenA* = KM089126; *RPB2* = KM089513; *CaM* = KM189504).

- Penicillium sacculum** E. Dale, Ann. Mycol. 24: 137. 1926 ≡ *Eladia saccula* (E. Dale) G. Sm., Trans. Brit. Mycol. Soc. 44: 47. 1961. [MB277209]. — Herb.: CBS 231.61. Ex-type: CBS 231.61 = ATCC 18350 = IFO 8114 = IFO 9454 = IMI 051498 = LSHBBB298 = UC4505. Section *Eladia*. ITS barcode: KC411707. (Alternative markers: *BenA* = KJ834488; *RPB2* = JN121462; *CaM* = n.a.).
- Penicillium sajorovi** Quintan., Av. Aliment. Majora Anim. 22: 539. 1981. [MB114172]. — Herb.: CBS 277.83. Ex-type: CBS 277.83 = CECT 2751 = IMI 259992. Section *Ramosa*. ITS barcode: KC411724. (Alternative markers: *BenA* = KJ834489; *RPB2* = JN406588; *CaM* = KJ867007).
- Penicillium sanguifluum** (Sopp) Biourge, Cellule 33: 105. 1923 ≡ *Citromyces sanguifluus* Sopp, Skr. Vidensk.-Selsk. Christiana Math.-Nat. Kl. 11: 115. 1912. [MB356682]. — Herb.: CBS H-20645. Ex-type: CBS 127032 = IBT 29041. Section *Citrina*. ITS barcode: JN617681. (Alternative markers: *BenA* = JN606819; *RPB2* = n.a.; *CaM* = JN606555).
- Penicillium saturniforme** (L. Wang & W.Y. Zhuang) Houbraken & Samson, Stud. Mycol. 70: 48. 2011 ≡ *Eupenicillium saturniforme* L. Wang & W.Y. Zhuang Mycopathologia 167: 300. 2009. [MB561958]. — Herb.: AS 3.6886. Ex-type: CBS 122276 = AS 3.6886. Section *Aspergilloides*. ITS barcode: EU644081. (Alternative markers: *BenA* = EU644080; *RPB2* = JN121439; *CaM* = EU644062).
- Penicillium scabrosum** Frisvad, Samson & Stolk, Persoonia 14: 177. 1990. [MB136735]. — Herb.: IMI 285533. Ex-type: CBS 683.89 = FRR 2950 = IBT 3736 = IMI 285533 = DAOM 214786. Section *Ramosa*. ITS barcode: DQ267906. (Alternative markers: *BenA* = DQ285610; *RPB2* = JN406541; *CaM* = FJ530987).
- Penicillium sclerotigenum** W. Yamam., Sci. Rep. Hyogo Univ Agric. 1: 69. 1955. [MB302424]. — Herb.: IMI 68616. Ex-type: CBS 101033 = CBS 343.59 = ATCC 18488 = IBT 14346 = IFO 6167 = IMI 068616 = NRRL 3461 = QM 7779. Section *Penicillium*. ITS barcode: AF033470. (Alternative markers: *BenA* = AY674393; *RPB2* = JN406652; *CaM* = n.a.).
- Penicillium sclerotiorum** J.F.H. Beyma, Zentralbl. Bakteriol. Parasitenk., Abt. 2 96: 418. 1937. [MB277708]. — Herb.: IMI 40569. Ex-type: CBS 287.36 = ATCC 10494 = IFO 6105 = IMI 040569 = NRRL 2074 = QM 1938 = VKMF-353. Section *Sclerotiora*. ITS barcode: JN626132. (Alternative markers: *BenA* = JN626001; *RPB2* = JN406585; *CaM* = JN626044).
- Penicillium senticosum** D.B. Scott, Mycopathol. Mycol. Appl. 36: 5. 1968 ≡ *Eupenicillium senticosum* D.B. Scott, Mycopathol. Mycol. Appl. 36: 5. 1968. [MB335764]. — Herb.: CBS 316.67. Ex-type: CBS 316.67 = ATCC 18623 = CSIR 1042 = IMI 136211 = IMI 216905. Section *Eladia*. ITS barcode: KC411733. (Alternative markers: *BenA* = KJ834490; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium shearri** Stolk & D.B. Scott, Persoonia 4: 396. 1967 ≡ *Eupenicillium shearri* Stolk & D.B. Scott, Persoonia 4: 396. 1967. [MB335765]. — Herb.: CBS 290.48. Ex-type: CBS 290.48 = ATCC 10410 = IFO 6088 = IMI 039739 = IMI 039739iv = NRRL 715 = QM 1870. Section *Citrina*. ITS barcode: GU944606. (Alternative markers: *BenA* = JN606840; *RPB2* = JN121482; *CaM* = EU644068).
- Penicillium shenniangjianum** H.Z. Kong & Z.T. Qi, Mycosistema 1: 110. 1988. [MB119509]. — Herb.: CBS 228.89. Ex-type: CBS 228.89. Section *Cinnamopurpurea*. ITS barcode: KC411705. (Alternative markers: *BenA* = KJ834491; *RPB2* = JN121458; *CaM* = AY678561).
- Penicillium simile** Davolos et al., Int. J. Syst. Evol. Microbiol. 62: 457. 2012. [MB509645]. — Herb.: ATCC MYA-4591. Ex-type: CBS 129191 = ATCC MYA-4591. Section *Ramosa*. ITS barcode: FJ376592. (Alternative markers: *BenA* = FJ376595; *RPB2* = n.a.; *CaM* = GQ979710).
- Penicillium simplicissimum** (Oudem.) Thom, Penicillia: 335. 1930 ≡ *Spicaria simplicissima* Oudem., Ned. Kruidk. Arch. 2: 763. 1902. [MB278201]. — Herb.: CUP Jensen. 1912: No. 5921 (CUP). Ex-type: CBS 372.48 = ATCC 10495 = FRR 902 = IFO 5762 = IMI 039816 = QM 1939. Section *Lanata-Divaricata*. ITS barcode: GU981588. (Alternative markers: *BenA* = GU981632; *RPB2* = JN121507; *CaM* = KF296368).
- Penicillium sinicum** Udagawa & S. Ueda, Mycotaxon 14: 266. 1982 ≡ *Eupenicillium sinicum* Udagawa & S. Ueda, Mycotaxon 14: 266. 1982. [MB110862]. — Herb.: NHL 2894. Ex-type: CBS 279.82 = NHL 2894. Section *Chrysogena*. ITS barcode: JX997090. (Alternative markers: *BenA* = JX996846; *RPB2* = JN406587; *CaM* = JX996970).
- Penicillium singorense** Visagie, Seifert & Samson, Stud. Mycol. 78: 119. 2014. [MB809182]. — Herb.: CBS H-21802. Ex-type: CBS 138214 = DTO 133C6. Section *Lanata-Divaricata*. ITS barcode: KJ775674. (Alternative markers: *BenA* = KJ775167; *RPB2* = n.a.; *CaM* = KJ775403).
- Penicillium sizovae** Baghd., Novosti Sist. Nizsh. Rast. 1968: 103. 1968. [MB335767]. — Herb.: CBS 413.69. Ex-type: CBS 413.69 = FRR 518 = IMI 140344 = VKMF-1073. Section *Citrina*. ITS barcode: GU944588. (Alternative markers: *BenA* = GU944535; *RPB2* = JN606603; *CaM* = GU944618).
- Penicillium skrjabinii** Schmotina & Golovleva, Mikol. Fitopatol. 8: 530. 1974. [MB319296]. — Herb.: IMI 196528. Ex-type: CBS 439.75 = NRRL 13055 = FRR 1945 = IMI 196528 = VKMF-1940. Section *Lanata-Divaricata*. ITS barcode: GU981576. (Alternative markers: *BenA* = GU981626; *RPB2* = EU427252; *CaM* = KF296370).
- Penicillium smithii** Quintan., Av. Aliment. Majora Anim. 23: 340. 1982. [MB114173]. — Herb.: CBS 276.83. Ex-type: CBS 276.83 = CECT 2744 = IMI 259693. Section *Exilicaulis*. ITS barcode: KC411723. (Alternative markers: *BenA* = KJ834492; *RPB2* = JN406589; *CaM* = n.a.).
- Penicillium solitum** Westling, Ark. Bot. 11: 65. 1911. [MB206172]. — Herb.: CBS 424.89. Ex-type: CBS 424.89 = ATCC 9923 = CBS 288.36 = FRR 937 = IBT 3948 = IFO 7765 = IMI 039810 = IMI 092225 = LSHBP 52 = MUCL 28668 = MUCL 29173 = NRRL 937. Section *Fasciculata*. ITS barcode: AY373932. (Alternative markers: *BenA* = AY674354; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium soppii** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 476. 1927. [MB121424]. — Herb.: IMI 40217. Ex-type: CBS 226.28 = ATCC 10496 = FRR 2023 = IFO 7766 = IMI 040217 = MUCL 29233 = NRRL 2023 = QM 1964 = IBT 18220. Section *Ramosa*. ITS barcode: AF033488. (Alternative markers: *BenA* = DQ285616; *RPB2* = JN406606; *CaM* = n.a.).
- Penicillium spathulatum** Frisvad & Samson, FEMS Microbiol. Lett. 339: 88. 2013. [MB492650]. — Herb.: CBS 117192. Ex-type: CBS 117192 = IBT 22220. Section *Brevicompacta*. ITS barcode: JX313165. (Alternative markers: *BenA* = JX313183; *RPB2* = JN406636; *CaM* = JX313149).
- Penicillium spinulosum** Thom, U.S.D.A. Bur. Animal Industr. Bull. 118: 76. 1910. [MB215401]. — Herb.: IMI 243161. Ex-type: CBS 374.48 = ATCC 10498 = FRR 1750 = IMI 024316 = LSHBAd 29 = MUCL 13910 = MUCL 13911 = NCTC 591 = NRRL 1750 = QM 7654. Section *Aspergilloides*. ITS barcode: AF033410. (Alternative markers: *BenA* = KJ834493; *RPB2* = JN406558; *CaM* = GQ367524).
- Penicillium steckii** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 469. 1927. [MB278769]. — Herb.: IMI 40583. Ex-type: CBS 260.55 = ATCC 10499 = CECT 2268 = DSM1252 = IMI 040583 = NRRL 2140 = QM 6413. Section *Citrina*. ITS barcode: GU944597. (Alternative markers: *BenA* = GU944522; *RPB2* = JN606602; *CaM* = GU944611).
- Penicillium sterculinicola** Houbraken, Stud. Mycol. 78: 436. 2014. [MB809973]. — Herb.: CBS H-21877. Ex-type: CBS 122426 = DTO 031-A4. Section *Aspergilloides*. ITS barcode: KM189464. (Alternative markers: *BenA* = KM088693; *RPB2* = KM089465; *CaM* = KM089078).
- Penicillium stolkiae** D.B. Scott, Mycopathol. Mycol. Appl. 36: 8. 1968 ≡ *Eupenicillium stolkiae* D.B. Scott, Mycopathol. Mycol. Appl. 36: 8. 1968. [MB335768]. — Herb.: CBS 315.67. Ex-type: CBS 315.67 = ATCC 18546 = CSIR 1041 = FRR 534 = IMI 136210 = NRRL 5816. Section *Stolkia*. ITS barcode: AF033444. (Alternative markers: *BenA* = JN617717; *RPB2* = JN121488; *CaM* = AF481135).
- Penicillium striatisporum** Stolk, Antonie van Leeuwenhoek 35: 268. 1969. [MB335769]. — Herb.: CBS 705.68. Ex-type: CBS 705.68 = ATCC 22052 = CCRC 31679 = FRR 827 = IMI 151749 = MUCL 31202. Section *Exilicaulis*. ITS barcode: AF038938. (Alternative markers: *BenA* = JX141156; *RPB2* = JN406538; *CaM* = n.a.).
- Penicillium subarcticum** S.W. Peterson & Sigler, Mycol. Res. 106: 1116. 2002. [MB483983]. — Herb.: BPI 841397. Ex-type: CBS 111719 = NRRL 31108 = UAMH 3897. Section *Stolkia*. ITS barcode: AF481120. (Alternative markers: *BenA* = JN617716; *RPB2* = n.a.; *CaM* = AF481141).
- Penicillium sublateritium** Biourge, Cellule 33: 315. 1923. [MB279265]. — Herb.: IMI 40594. Ex-type: CBS 267.29 = ATCC 10502 = FRR 2071 = IFO 6107 = IMI 040594 = LSHB 55 = MUCL 28655 = NRRL 2071 = QM 7652 = NBRC 6107. Section *Ramigena*. ITS barcode: EU427288. (Alternative markers: *BenA* = JX091557; *RPB2* = JN406590; *CaM* = AB566102).
- Penicillium sublectaticum** Houbraken, et al., Stud. Mycol. 78: 436. 2014. [MB809974]. — Herb.: CBS H-21955. Ex-type: CBS 138217 = DTO 244-G2. Section *Aspergilloides*. ITS barcode: KM189761. (Alternative markers: *BenA* = KM089010; *RPB2* = KM089784; *CaM* = KM089397).
- Penicillium subrubescens** Houbraken, et al., Antonie van Leeuwenhoek 103: 1354. 2013. [MB801306]. — Herb.: CBS H-21029. Ex-type: CBS 132785 = DTO 188D6 = FBCC 1632 = IBT 31985. Section *Lanata-Divaricata*. ITS barcode: KC346350. (Alternative markers: *BenA* = KC346327; *RPB2* = KC346306; *CaM* = KC346330).
- Penicillium subspinulosum** Houbraken, Stud. Mycol. 78: 436. 2014. [MB809975]. — Herb.: CBS H-21856. Ex-type: CBS 137946 = DTO 041-F2.

- Section *Aspergilloides*. ITS barcode: KM189483. (Alternative markers: *BenA* = KM088719; *RPB2* = KM089491; *CaM* = KM089104).
- Penicillium sucrivorum*** Visagie & K. Jacobs, Mycologia 106: 546. 2014. [MB804723]. — Herb.: CBS H-21331. Ex-type: CBS 135116 = DAOM 241042 = DTO 183E5. Section *Citrina*. ITS barcode: JX140872. (Alternative markers: *BenA* = JX141015; *RPB2* = n.a.; *CaM* = JX141506).
- Penicillium sumatraense*** Szilvinyi [as 'sumatrense'], Archiv. Hydrobiol. 14, Suppl. 6: 535. 1936. [MB319297]. — Herb.: unknown. Ex-type: CBS 281.36 = NRRL 779 = FRR 779. Section *Citrina*. ITS barcode: GU944578. (Alternative markers: *BenA* = JN606639; *RPB2* = n.a.; *CaM* = JN606368).
- Penicillium svalbardense*** Frisvad, Sonjak & Gundae-Cim, Antonie van Leeuwenhoek 92: 48. 2007. [MB529943]. — Herb.: EX-F 1307. Ex-type: CBS 122416 = IBT 23856. Section *Lanata-Divaricata*. ITS barcode: GU981603. (Alternative markers: *BenA* = KC346325; *RPB2* = KF296457; *CaM* = KC346338).
- Penicillium swiecickii*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B, Sci. Nat. 1927: 474. 1927. [MB279618]. — Herb.: unknown. Ex-type: CBS 119391 = FRR 918 = IBT 27865 = IMI 191500 = NRRL 918. Section *Ramosa*. ITS barcode: AF033490. (Alternative markers: *BenA* = KJ834494; *RPB2* = n.a.; *CaM* = KJ866993).
- Penicillium tardochrysogenum*** Frisvad, Houbraken & Samson, Persoonia 29: 93. 2012. [MB801877]. — Herb.: CBS H-21057. Ex-type: CBS 132200 = IBT 30075 = DTO 149B9. Section *Chrysogena*. ITS barcode: JX997027. (Alternative markers: *BenA* = JX996898; *RPB2* = JX996634; *CaM* = JX996239).
- Penicillium taxi*** R. Schneid., Zentralbl. Bakteriol. Parasitenk., Abt. 2 110: 43. 1956 ≡ *Thysanophora taxi* (R. Schneid.) Stolk & Hennebert, Persoonia 5: 193. 1968. [MB282799]. — Herb.: unknown. Ex-type: CBS 206.57 = ATCC 18484 = BBA 7480 = MUCL 11402 = QM 8153. Section *Thysanophora*. ITS barcode: KJ834517. (Alternative markers: *BenA* = KJ834495; *RPB2* = JN121454; *CaM* = n.a.).
- Penicillium terrenum*** D.B. Scott, Mycopathol. Mycol. Appl. 36: 1. 1968 ≡ *Eupenicillium terrenum* D.B. Scott, Mycopathol. Mycol. Appl. 36: 1. 1968. [MB335771]. — Herb.: CBS 313.67. Ex-type: CBS 313.67 = ATCC 18547 = CSIR 1022 = IMI 136208. Section *Exilicaulis*. ITS barcode: AM992111. (Alternative markers: *BenA* = KJ834496; *RPB2* = JN406577; *CaM* = n.a.).
- Penicillium terrigenum*** Seifert et al., Stud. Mycol. 70: 125. 2011. [MB563204]. — Herb.: CBS H-20667. Ex-type: CBS 127354 = IBT 30769. Section *Citrina*. ITS barcode: JN617684. (Alternative markers: *BenA* = JN606810; *RPB2* = JN606600; *CaM* = JN606583).
- Penicillium thiersii*** S.W. Peterson, E.M. Bayer & Wicklow, Mycologia 96: 1283. 2004. [MB487738]. — Herb.: BPI 842269. Ex-type: CBS 117503 = IBT 27050 = NRRL 28162. Section *Aspergilloides*. ITS barcode: AF125936. (Alternative markers: *BenA* = KJ834497; *RPB2* = JN121434; *CaM* = AY741726).
- Penicillium thomii*** Maire, Bull. Soc. Hist. Nat. Afrique N. 8: 189. 1917 ≡ *Citromyces thomii* (Maire) Sacc., Syll. Fung. 25: 683. 1931 ≡ *Penicillium lividum* var. *thomii* (Maire) Stolk & Samson, Adv. *Penicillium Aspergillus* Syst.: 170. 1986. [MB202819]. — Herb.: IMI 189694. Ex-type: CBS 225.81 = IMI 189694 = NRRL 2077. Section *Aspergilloides*. ITS barcode: KM189560. (Alternative markers: *BenA* = KM088799; *RPB2* = KM089571; *CaM* = KM089184).
- Penicillium thymicola*** Frisvad & Samson, Stud. Mycol. 49: 29. 2004. [MB370969]. — Herb.: CBS 111225. Ex-type: CBS 111225 = IBT 5891. Section *Fasciculata*. ITS barcode: KJ834518. (Alternative markers: *BenA* = AY674321; *RPB2* = n.a.; *CaM* = FJ530990).
- Penicillium tricolor*** Frisvad et al., Can. J. Bot. 72: 937. 1994. [MB541710]. — Herb.: DAOM 216240. Ex-type: CBS 635.93 = IBT 12493 = DAOM 216240. Section *Fasciculata*. ITS barcode: JN942704. (Alternative markers: *BenA* = AY674313; *RPB2* = JN985422; *CaM* = n.a.).
- Penicillium tropicoides*** Houbraken, Frisvad & Samson, Fungal Divers. 44: 127. 2010. [MB518293]. — Herb.: CBS 122410. Ex-type: CBS 122410 = IBT 29043. Section *Citrina*. ITS barcode: GU944584. (Alternative markers: *BenA* = GU944531; *RPB2* = JN606608; *CaM* = GU944624).
- Penicillium tropicum*** Houbraken, Frisvad & Samson, Fungal Divers. 44: 129. 2010 ≡ *Eupenicillium tropicum* Tuthill & Frisvad, Mycol. Progress 3: 14. 2004. [MB518294]. — Herb.: SC42-1. Ex-type: CBS 112584 = IBT 24580. Section *Citrina*. ITS barcode: GU944582. (Alternative markers: *BenA* = GU944532; *RPB2* = JN606607; *CaM* = GU944625).
- Penicillium trzebinskii*** K.M. Zalessky, Bull. Int. Acad. Polon. Sci., Sér. B, Sci. Nat. 1927: 498. 1927. [MB280795]. — Herb.: unknown. Ex-type: CBS 382.48 = ATCC 10507 = FRR 731 = IFO 6110 = IMI 039749 = MUCL 29102 = NRRL 731 = QM 7678. Section *Aspergilloides*. ITS barcode:
- KM189784. (Alternative markers: *BenA* = KM089034; *RPB2* = KM089808; *CaM* = KM089421).
- Penicillium tsitsikammaense*** Houbraken, Stud. Mycol. 78: 440. 2014. [MB809976]. — Herb.: CBS H-21881. Ex-type: CBS 328.71 = DTO 006-13 = CSIR 1092. Section *Aspergilloides*. ITS barcode: KM189451. (Alternative markers: *BenA* = KM088675; *RPB2* = KM089447; *CaM* = KM089060).
- Penicillium tularensense*** Paden, Mycopathol. Mycol. Appl. 43: 264. 1971 ≡ *Eupenicillium tularensense* Paden, Mycopathol. Mycol. Appl. 43: 264. 1971. [MB319298]. — Herb.: UVIC JWP 68-31. Ex-type: CBS 430.69 = ATCC 22056 = FRR 899 = IFO 31740 = IMI 148394 = NRRL 5273 = AS 3.14006. Section *Brevicompacta*. ITS barcode: AF033487. (Alternative markers: *BenA* = KC427175; *RPB2* = JN121516; *CaM* = JX313135).
- Penicillium tulipae*** Overy & Frisvad, Syst. Appl. Microbiol. 26: 634. 2003. [MB488954]. — Herb.: C 60162. Ex-type: CBS 109555 = CBS 187.88 = IBT 3458. Section *Fasciculata*. ITS barcode: KJ834519. (Alternative markers: *BenA* = AY674344; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium turbatum*** Westling, Ark. Bot. 11: 128. 1911. [MB202895]. — Herb.: IMI 39738. Ex-type: CBS 383.48 = CBS 237.60 = ATCC 9782 = DSM2426 = FRR 757 = IFO 7767 = IMI 039738 = MUCL 29115 = NRRL 757 = NRRL 758 = QM 1941. Section *Turbata*. ITS barcode: AF034454. (Alternative markers: *BenA* = KJ834499; *RPB2* = JN406556; *CaM* = n.a.).
- Penicillium turcosoconidiatum*** Visagie, Houbraken & K. Jacobs, Stud. Mycol. 78: 440. 2014. [MB809977]. — Herb.: CBS H-21876. Ex-type: CBS 138557 = DTO 181-A3 = CV 110 = DAOM 241130. Section *Aspergilloides*. ITS barcode: KM189645. (Alternative markers: *BenA* = KM088889; *RPB2* = KM089663; *CaM* = KM089276).
- Penicillium ubiquetum*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 127. 2011. [MB563201]. — Herb.: CBS H-20659. Ex-type: CBS 126437 = IBT 22226. Section *Citrina*. ITS barcode: JN617680. (Alternative markers: *BenA* = JN606800; *RPB2* = n.a.; *CaM* = JN606460).
- Penicillium ulaiense*** H.M. Hsieh, H.J. Su & Tzean, Trans. Mycol. Soc. Rep. China 2: 161. 1987. [MB126489]. — Herb.: PPEH 29001.87. Ex-type: CBS 210.92 = CBS 261.94 = CCRC 32655 = IBT 18387 = IBT 23037. Section *Penicillium*. ITS barcode: KC411695. (Alternative markers: *BenA* = AY674408; *RPB2* = n.a.; *CaM* = n.a.).
- Penicillium vagum*** Houbraken, et al., Stud. Mycol. 78: 443. 2014. [MB809978]. — Herb.: CBS H-21926. Ex-type: CBS 137728 = DTO 180-G3 = CV 25 = DAOM 241357. Section *Aspergilloides*. ITS barcode: KM189642. (Alternative markers: *BenA* = KM088886; *RPB2* = KM089660; *CaM* = KM089273).
- Penicillium valentinum*** C. Ramírez and A.T. Martínez, Mycopathologia 72: 183. 1980. [MB113027]. — Herb.: unknown. Ex-type: CBS 172.81 = ATCC 42227 = IJFM 5071. Section *Aspergilloides*. ITS barcode: KM189550. (Alternative markers: *BenA* = KM088788; *RPB2* = KM089560; *CaM* = KM089173).
- Penicillium vancouverense*** Houbraken, Frisvad & Samson, Stud. Mycol. 70: 131. 2011. [MB563207]. — Herb.: CBS H-20646. Ex-type: CBS 126323 = IBT 20700. Section *Citrina*. ITS barcode: JN617675. (Alternative markers: *BenA* = JN606663; *RPB2* = n.a.; *CaM* = JN606399).
- Penicillium vanderhammenii*** Houbraken et al., Int. J. Syst. Evol. Microbiol. 61: 1473. 2011. [MB518027]. — Herb.: HUA 170337. Ex-type: CBS 126216 = IBT 23203. Section *Lanata-Divaricata*. ITS barcode: GU981574. (Alternative markers: *BenA* = GU981647; *RPB2* = KF296458; *CaM* = KF296382).
- Penicillium vanluykii*** Frisvad, Houbraken & Samson, Persoonia 29: 97. 2012. [MB801878]. — Herb.: CBS H-21059. Ex-type: CBS 131539 = DTO 14812 = IBT 14505. Section *Chrysogena*. ITS barcode: JX997007. (Alternative markers: *BenA* = JX996879; *RPB2* = JX996615; *CaM* = JX996220).
- Penicillium vanoranjei*** Visagie, Houbraken & Samson, Persoonia 31: 46. 2013. [MB803782]. — Herb.: CBS H-21145. Ex-type: CBS134406. Section *Sclerotioria*. ITS barcode: KC695696. (Alternative markers: *BenA* = KC695686; *RPB2* = n.a.; *CaM* = KC695691).
- Penicillium vasconiae*** C. Ramírez & A.T. Martínez, Mycopathologia 72: 189. 1980. [MB113028]. — Herb.: CBS 339.79. Ex-type: CBS 339.79 = ATCC 42224 = IJFM 3008. Section *Lanata-Divaricata*. ITS barcode: GU981599. (Alternative markers: *BenA* = GU981653; *RPB2* = KF296459; *CaM* = KF296386).
- Penicillium velutinum*** J.F.H. Beyma, Zentralbl. Bakteriol. Parasitenk., Abt. 2 91: 353. 1935. [MB283175]. — Herb.: IMI 40571. Ex-type: CBS 250.32 = ATCC 10510 = CECT 2318 = IJFM 5108 = IMI 040571 = NRRL 2069 = QM 7686 = VKMF-379. Section *Exilicaulis*. ITS barcode: AF033448. (Alternative markers: *BenA* = JX141170; *RPB2* = n.a.; *CaM* = n.a.).

- Penicillium venetum** (Frissvad) Frissvad, Adv. *Penicillium Aspergillus* Syst.: 275. 2000 ≡ *Penicillium hirsutum* var. *venetum* Frissvad, Mycologia 81: 856. 1990. [MB459816]. — Herb.: IMI 321520. Ex-type: IBT 10661 = IMI 321520. Section *Fasciculata*. ITS barcode: AJ005485. (Alternative markers: BenA = AY674335; RPB2 = n.a.; CaM = n.a.).
- Penicillium verhagenii** Houbraken, Stud. Mycol. 78: 443. 2014. [MB809979]. — Herb.: CBS H-21865. Ex-type: CBS 137959 = DTO 193-A1. Section *Aspergilloides*. ITS barcode: KM189708. (Alternative markers: BenA = KM088955; RPB2 = KM089729; CaM = KM089342).
- Penicillium verrucosum** Dierckx, Ann. Soc. Sci. Bruxelles 25: 88. 1901. [MB212252]. — Herb.: IMI 200310. Ex-type: CBS 603.74 = ATCC 48957 = ATHUM2897 = CECT 2906 = FRR 965 = IBT 12809 = IBT 4733 = IMI 200310 = IMI 200310ii = MUCL 28674 = MUCL 29089 = MUCL 29186 = NRRL 965. Section *Fasciculata*. ITS barcode: AY373938. (Alternative markers: BenA = AY674323; RPB2 = JN121539; CaM = DQ911138).
- Penicillium vinaceum** J.C. Gilman & E.V. Abbott, Iowa St. Coll. J. Sci. 1: 299. 1927. [MB281754]. — Herb.: IMI 29189. Ex-type: CBS 389.48 = ATCC 10514 = FRR 739 = IMI 029189 = NRRL 739 = QM 6746. Section *Exilicaulis*. ITS barcode: AF033461. (Alternative markers: BenA = HQ646575; RPB2 = JN406555; CaM = HQ646586).
- Penicillium virgatum** Nierenberg & Kwasna, Mycol. Res. 109: 977. 2005. [MB341488]. — Herb.: BBA 65745. Ex-type: CBS 114838 = BBA 65745. Section *Ramosa*. ITS barcode: AJ748692. (Alternative markers: BenA = KJ834500; RPB2 = JN406641; CaM = KJ866992).
- Penicillium viridicatum** Westling, Ark. Bot. 11: 88. 1911 ≡ *Penicillium aurantiogriseum* var. *viridicatum* (Westling) Frissvad & Filt., Mycologia 81: 850. 1990. [MB163349]. — Herb.: IMI 39758ii. Ex-type: CBS 390.48 = ATCC 10515 = IBT 23041 = IFO 7736 = IMI 039758 = IMI 039758ii = NRRL 963 = QM 7683. Section *Fasciculata*. ITS barcode: AY373939. (Alternative markers: BenA = AY674295; RPB2 = JN121511; CaM = n.a.).
- Penicillium viticola** Nonaka & Masuma, Mycoscience 52: 339. 2011. [MB516048]. — Herb.: TNS-F38702. Ex-type: JCM 17636 = FKI-4410. Section *Sclerotiora*. ITS barcode: AB606414. (Alternative markers: BenA = AB540174; RPB2 = n.a.; CaM = n.a.).
- Penicillium vulpinum** (Cooke & Massee) Seifert & Samson, Adv. *Penicillium Aspergillus* Syst.: 144. 1985 ≡ *Coremium vulpinum* Cooke & Massee, Grevillea 16: 81. 1888. [MB114763]. — Herb.: "on dung", s. coll., in herb. Cooke (K). Ex-type: CBS 126.23 = ATCC 10426 = IMI 040237 = NRRL 2031 = VKMF-257. Section *Penicillium*. ITS barcode: AF506012. (Alternative markers: BenA = KJ834501; RPB2 = n.a.; CaM = n.a.).
- Penicillium waksmanii** K.M. Zalewsky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat.: 468. 1927. [MB121677]. — Herb.: IMI 39746i. Ex-type: CBS 230.28 = ATCC 10516 = FRR 777 = IFO 7737 = IMI 039746 = IMI 039746i = MUCL 29120 = NRRL 777 = QM 7681. Section *Citrina*. ITS barcode: GU944602. (Alternative markers: BenA = JN606779; RPB2 = JN606627; CaM = JN606431).
- Penicillium wellingtonense** A.L.J. Cole et al., Stud. Mycol. 70: 133. 2011. [MB563208]. — Herb.: CBS H-20657. Ex-type: CBS 130375 = IBT 23557 = DTO 76C6. Section *Citrina*. ITS barcode: JN617713. (Alternative markers: BenA = JN606670; RPB2 = JN606616; CaM = JN606395).
- Penicillium westlingii** K.M. Zalewsky, Bull. Int. Acad. Polon. Sci., Sér. B., Sci. Nat. 1927: 473. 1927. [MB282076]. — Herb.: IMI 92272. Ex-type: CBS 231.28 = IMI 092272. Section *Citrina*. ITS barcode: GU944601. (Alternative markers: BenA = JN606718; RPB2 = JN606625; CaM = JN606500).
- Penicillium wotroi** Houbraken et al., Int. J. Syst. Evol. Microbiol. 61: 1474. 2011. [MB518026]. — Herb.: HUA 170336. Ex-type: CBS 118171 = IBT 23253. Section *Lanata-Divaricata*. ITS barcode: GU981591. (Alternative markers: BenA = GU981637; RPB2 = KF296460; CaM = KF296369).
- Penicillium yarmokense** Baghd., Novosti Sist. Nizsh. Rast. 5: 99. 1968. [MB335774]. — Herb.: CBS H-7536. Ex-type: CBS 410.69 = FRR 520 = IMI 140346 = VKMF-1076. Section *Canescensia*. ITS barcode: KC411757. (Alternative markers: BenA = KJ834502; RPB2 = JN406553; CaM = KJ867013).
- Penicillium yezoense** Hanzawa ex Houbraken, Stud. Mycol. 78: 443. 2014. [MB809980]. — Herb.: CBS H-21863. Ex-type: CBS 350.59 = ATCC 18333 = FRR 3395 = IFO 5362 = IMI 068615. Section *Aspergilloides*. ITS barcode: KM189553. (Alternative markers: BenA = KM088792; RPB2 = KM089564; CaM = KM089177).
- Penicillium zonatum** Hodges & J.J. Perry, Mycologia 65: 697. 1973 ≡ *Eupenicillium zonatum* Hodges & J.J. Perry, Mycologia 65: 697. 1973. [MB319303]. — Herb.: BPI FSL 525. Ex-type: CBS 992.72 = ATCC 24353. Section *Lanata-Divaricata*. ITS barcode: GU981581. (Alternative markers: BenA = GU981651; RPB2 = KF296461; CaM = KF296380).
- Penicillium zhuangii** L. Wang, PloS ONE 9: e101454-P4. 2014. [MB805945]. — Herb.: HMAS 244922. Ex-type: CBS 137464 = NRRL 62806 = AS 3.15341. Section *Aspergilloides*. ITS barcode: KF769435. (Alternative markers: BenA = KF769411; RPB2 = n.a.; CaM = KF769422).
- ## Doubtful species
- Penicillium asymmetricum** (Subram. & Sudha) Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Thysanophora asymmetrica* Subram. & Sudha, Kavaka 12: 88. 1985. [MB561963]. — Herb.: unknown. Ex-type: No culture available. Note: No material is available for this species. As such, we cannot confirm the taxonomic position of the species and consider it as doubtful.
- Penicillium coniferophilum** Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Thysanophora striatispora* Barron & Cooke, Mycopathol. Mycol. Appl. 40: 353. 1970. [MB561968]. — Herb.: unknown. Ex-type: No culture available. Note: No material is available for this species. As such, we cannot confirm the taxonomic position of the species and consider it as doubtful.
- Penicillium glaucoalbidum** (Desm.) Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Sclerotium glaucoalbidum* Desm., Ann. Sci. Nat., Bot. 16: 329. 1851 ≡ *Thysanophora glaucoalbida* (Desm.) M. Morelet, Ann. Soc. Sci. Nat. Archéol. Toulon Var 20: 104. 1968. [MB561965]. — Herb.: unknown. Ex-type: No culture available. Note: No material is available for this species. As such, we cannot confirm the taxonomic position of the species and consider it as doubtful.
- Penicillium melanostipe** Houbraken & Samson, Stud. Mycol. 70: 47. 2011 ≡ *Thysanophora verrucosa* Mercado, Gené & Guarro, Mycotaxon 67: 419. 1998. [MB561970]. — Herb.: HAC (M) 9165. Ex-type: No culture available. Note: No material is available for this species. As such, we cannot confirm the taxonomic position of the species and consider it as doubtful.
- Penicillium pariverrucosum** (K. Ando & Pitt) Houbraken & Samson, Stud. Mycol. 70: 48. 2011 ≡ *Torulomyces pariverrucosus* K. Ando & Pitt, Mycoscience 39: 317. 1998. — Herb.: TNS-F-238516. Ex-type: KY 12720. Note: Material for this species is not available for study. As such, we cannot confirm the taxonomic position of the species and consider it as doubtful.
- Penicillium taiwanense** (Matsush.) Houbraken & Samson, Stud. Mycol. 70: 48. 2011 ≡ *Phialomyces taiwanensis* Matsush., Matsushima Mycol. Mem. 4: 12. 1985 ≡ *Thysanophora taiwanensis* (Matsush.) Mercado, Gené & Guarro, Mycotaxon 67: 421. 1998. [MB561969]. — Herb.: unknown. Ex-type: No culture available. Section *Thysanophora*. Note: No material is available for this species. As such, we cannot confirm the taxonomic position of the species and consider it as doubtful.
- ## Excluded species
- Penicillium arenicola** Chalabuda, Bot. Mater. Otd. Sporov. Rast. 6: 162. 1950. [MB302375]. — Herb.: IMI 117658. Ex-type: CBS 220.66 = ATCC 18321 = ATCC 18330 = DSM 2435 = FRR 3392 = IMI 117658 = NRRL 3392 = VKM F-1035. ITS barcode: GU092964. (Alternative markers: BenA = n.a.; CaM = n.a.; RPB2 = JN121457). Note: Molecular data suggest that this species is related to *Phialomyces* (Houbraken & Samson 2011).
- Penicillium canadense** G. Sm., Trans. Br. Mycol. Soc. 39: 113. 1956 = *Penicillium arenicola* Chalab., Bot. Mater. Otd. Sporov. Rast. 6: 162. 1950. [MB302384]. — Herb.: unknown. Ex-type: CBS 245.56 = ATCC 18424 = FRR 2553 = IMI 061834 = LSHB BB300 = NRRL 2553 = QM 6970 = WB 4155. ITS barcode: GU092963. (Alternative markers: BenA = GU092758; CaM = n.a.; RPB2 = n.a.). Note: Molecular data suggest that this species is a synonym of *P. arenicola* and is related to *Phialomyces* (Houbraken & Samson 2011).
- Penicillium kabunicum** Baghdadi, Novosti Sist. Nizsh. Rast.: 98. 1968. [MB335738]. — Herb.: CBS 575.90. Ex-type: CBS 575.90 = CBS 409.69 = FRR 513 = IMI 140341 = VKM F-1072. ITS barcode: n.a.. (Alternative markers: BenA = n.a.; CaM = n.a.; RPB2 = n.a.). Note: Molecular data suggest that this species does not belong to *Penicillium* (Houbraken, unpubl. data).
- Penicillium moldavicum** Milko & Beliakova, Novosti Sist. Nizsh. Rast. 1967: 255. 1967. [MB335751]. — Herb.: unknown. Ex-type: CBS 574.90 = ATCC 18355 = CBS 627.67 = FRR 665 = IMI 129966 = VKM F-922. ITS barcode:

n.a.. (Alternative markers: *BenA* = n.a.; *CaM* = n.a.; *RPB2* = n.a.). Note: Molecular data suggest that this species does not belong to *Penicillium* (Houbraken, unpubl. data).

Penicillium nodosatum Valla, Pl. Soil 114: 146. 1989. [MB126535]. — Herb.: CBS 330.90. Ex-type: CBS 333.90. ITS barcode: KC790403. (Alternative markers: *BenA* = KC790399; *CaM* = n.a.; *RPB2* = n.a.). Note: Molecular data suggest that this species is closely related to *P. kabunicum* and does not belong in *Penicillium* (Visagie et al. 2013).

Penicillium resedanum McLennan & Ducker, Aust. J. Bot. 2: 360. 1954. [MB302422]. — Herb.: IMI 062877. Ex-type: CBS 181.71 = ATCC 22356 = FRR 578 = IMI 062877 = NRRL 578. ITS barcode: AF033398. (Alternative markers: *BenA* = n.a.; *CaM* = n.a.; *RPB2* = n.a.). Note: Pitt (1979) noted that this species form acerose phialides with weak growth on G25N, suggesting a relationship with *Talaromyces*, which is confirmed by ITS data (Houbraken & Samson 2011).

ACKNOWLEDGEMENTS

This research was supported by grants from the Alfred P. Sloan Foundation Program on the Microbiology of the Built Environment.

REFERENCES

- Abraham E, Chain E, Fletcher C, et al. (1941). Further observations on penicillin. *The Lancet* 238: 177–188.
- Adsul MG, Bastawde KB, Varma AJ, et al. (2007). Strain improvement of *Penicillium janthinellum* NCIM 1171 for increased cellulase production. *Bioresource Technology* 98: 1467–1473.
- Amiri-Eliasi B, Fenselau C (2001). Characterization of protein biomarkers desorbed by MALDI from whole fungal cells. *Analytical Chemistry* 73: 5228–5231.
- Barreto MC, Houbraken J, Samson RA, et al. (2011). Taxonomic studies of the *Penicillium glabrum* complex and the description of a new species *P. subericola*. *Fungal Diversity* 49: 23–33.
- Basaran P, Demirbas RM (2010). Spectroscopic detection of pharmaceutical compounds from an aflatoxigenic strain of *Aspergillus parasiticus*. *Microbiological Research* 165: 516–522.
- Berbee M, Yoshimura A, Sugiyama J, et al. (1995). Is *Penicillium* Monophyletic? An evaluation of phylogeny in the family *Trichocomaceae* from 18S, 5.8S and ITS ribosomal DNA sequence data. *Mycologia* 87: 210–222.
- Bille E, Dauphin B, Leto J, et al. (2011). MALDI-TOF MS Andromas strategy for the routine identification of bacteria, mycobacteria, yeasts, *Aspergillus* spp. and positive blood cultures. *Clinical Microbiology and Infection* 18: 1117–1125.
- Blakeslee A (1915). Lindner's roll tube method of separation cultures. *Phytopathology* 5: 68–69.
- Blaxter M (2003). Counting angels with DNA. *Nature* 421: 1–3.
- Blaxter M, Mann J, Chapman T, et al. (2005). Defining operational taxonomic units with DNA barcodes data. *Philosophical Transactions of the Royal Society B* 360: 1935–1943.
- Boysen M, Skouboe P, Frisvad JC, Rossen L (1996). Reclassification of the *Penicillium roqueforti* group into three species on the basis of molecular genetic and biochemical profiles. *Microbiology* 142: 541–549.
- Chain E, Florey H, Gardner A, et al. (1940). Penicillin as a chemotherapeutic agent. *The Lancet* 236: 226–228.
- Chalupová J, Raus M, Sedlářová M, et al. (2014). Identification of fungal microorganisms by MALDI-TOF mass spectrometry. *Biotechnology Advances* 32: 230–241.
- Chen AJ, Tang D, Zhou YQ, et al. (2013). Identification of Ochratoxin A producing fungi associated with fresh and dry liquorice. *PLoS One* 8: e78285.
- Christensen M, Frisvad JC, Tuthill DE (2000). *Penicillium* species diversity in soil and some taxonomic and ecological notes. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification* (Samson RA, Pitt JI, eds). Harwood Academic Publishers, Amsterdam: 309–321.
- Ciegler A, Pitt JI (1970). Survey of the genus *Penicillium* for tremorgenic toxin production. *Mycopathologia et Mycologia applicata* 42: 119–124.
- Clutterbuck PW, Raistrick H (1933). Studies in the biochemistry of microorganisms XXXI. The molecular constitution of the metabolic products of *Penicillium brevicompactum* Dierckx and related species. II. Mycophenolic acid. *Biochemical Journal* 27: 654–667.
- Cruickshank RH, Pitt JI (1987a). Identification of species in *Penicillium* subgenus *Penicillium* by enzyme electrophoresis. *Mycologia*, 614–620.
- Cruickshank RH, Pitt JI (1987b). The zymogram technique – isoenzyme patterns as an aid in *Penicillium* classification. *Microbiological Sciences* 4: 14–17.
- De Carolis E, Posteraro B, Lass-Flörl C, et al. (2011). Species identification of *Aspergillus*, *Fusarium* and *Mucorales* with direct surface analysis by matrix-assisted laser desorption ionization time-of-flight mass spectrometry. *Clinical Microbiology and Infection* 18: 475–484.
- Del Chierico F, Masotti A, Onori M, et al. (2012). MALDI-TOF MS proteomic phenotyping of filamentous and other fungi from clinical origin. *Journal of Proteomics* 75: 3314–3330.
- DeSalle R, Egan MG, Siddal M (2005). The unholy trinity: taxonomy, species delimitation and DNA barcoding. *Philosophical Transactions of the Royal Society B* 360: 1905–1916.
- El-Banna AA, Pitt JI, Leistner L (1987). Production of mycotoxins by *Penicillium* species. *Systematic and Applied Microbiology* 10: 42–46.
- Fennell DL, Raper KB (1955). New species and varieties of *Aspergillus*. *Mycologia* 47: 68–89.
- Fleming A (1929). On the antibacterial action of cultures of a *Penicillium*, with special reference to their use in the isolation of *B. influenzae*. *British Journal of Experimental Pathology* 10: 226–236.
- Frisvad JC (1981). Physiological criteria and mycotoxin production as aids in identification of common asymmetric *Penicillia*. *Applied and Environmental Microbiology* 41: 568–579.
- Frisvad JC, Andersen B, Thrane U (2008). The use of secondary metabolite profiling in fungal taxonomy. *Mycological Research* 112: 231–240.
- Frisvad JC, Filtenborg O (1983). Classification of terverticillate *Penicillia* based on profiles of mycotoxins and other secondary metabolites. *Applied and Environmental Microbiology* 46: 1301–1310.
- Frisvad JC, Filtenborg O (1989). Terverticillate *Penicillia*: chemotaxonomy and mycotoxin production. *Mycologia* 81: 837–861.
- Frisvad JC, Samson RA (2004). Polyphasic taxonomy of *Penicillium* subgenus *Penicillium*. A guide to identification of food and air-borne terverticillate *Penicillia* and their mycotoxins. *Studies in Mycology* 49: 1–174.
- Frisvad JC, Smedsgaard J, Larsen TO, et al. (2004). Mycotoxins, drugs and other extrolites produced by species in *Penicillium* subgenus *Penicillium*. *Studies in Mycology* 49: 201–241.
- Frisvad JC, Thrane U (1987). Standardized High Performance Liquid Chromatography of 182 mycotoxins and other fungal metabolites based on alkylphenone indices and UV VIS spectra (diode array detection). *Journal of Chromatography* 404: 195–214.
- Frisvad JC, Thrane U (1993). Liquid column chromatography of mycotoxins. In: *Chromatography of mycotoxins: techniques and applications* (Betina V, ed), *Journal of Chromatography Library*, 54. Elsevier, Amsterdam: 253–372.
- Geiser DM, Pitt JI, Taylor JW (1998). Cryptic speciation and recombination in the aflatoxin-producing fungus *Aspergillus flavus*. *Proceedings of the National Academy of Sciences of the United States of America* 95: 388–393.
- Giraud F, Giraud T, Aguilera G, et al. (2010). Microsatellite loci to recognize species for the cheese starter and contaminating strains associated with cheese manufacturing. *International Journal of Food Microbiology* 137: 204–213.
- Glass NL, Donaldson GC (1995). Development of premier sets designed for use with the PCR to amplify conserved genes from filamentous Ascomycetes. *Applied and Environmental Microbiology* 61: 1323–1330.
- Hebert PDN, Ratnasingham S, deWaard JR (2003). Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society London B (Suppl.)* 270: S96–S99.
- Hettick JM, Kashon ML, Slaven JE, et al. (2006). Discrimination of intact mycobacteria at the strain level: A combined MALDI-TOF MS and biostatistical analysis. *Proteomics* 6: 6416–6425.
- Hettick JM, Green BJ, Buskirk AD, et al. (2008). Discrimination of *Penicillium* isolates by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry fingerprinting. *Rapid Communications in Mass Spectrometry* 22: 2555–2560.
- Hirano A, Iwai Y, Masuma R, et al. (1979). Neoxaline, a new alkaloid produced by *Aspergillus japonicus*. Production, isolation and properties. *The Journal of Antibiotics* 32: 781–785.
- Hocking AD, Pitt JI (1980). Dichloran-glycerol medium for enumeration of xerophilic fungi from low-moisture foods. *Applied and Environmental Microbiology* 39: 488–492.
- Hoog GS de, Gerrits van den Ende AH (1998). Molecular diagnostics of clinical strains of filamentous *Basidiomycetes*. *Mycoses* 41: 183–189.

- Hong S-B, Cho H-S, Shin H-D, et al. (2006). Novel *Neosartorya* species isolated from soil in Korea. *International Journal of Systematic and Evolutionary Microbiology* **56**: 477–486.
- Houbraken J, Frisvad JC, Samson RA (2011a). Fleming's penicillin producing strain is not *Penicillium chrysogenum* but *P. rubens*. *IMA Fungus* **2**: 87–95.
- Houbraken J, Frisvad JC, Samson RA (2011b). Taxonomy of *Penicillium* section *Citrina*. *Studies in Mycology* **70**: 53–138.
- Houbraken J, Frisvad JC, Seifert KA, et al. (2012a). New penicillin-producing *Penicillium* species and an overview of section *Chrysogena*. *Persoonia* **29**: 78–100.
- Houbraken J, López-Quintero CA, Frisvad JC, et al. (2011c). *Penicillium aracuarensense* sp. nov., *Penicillium elleniae* sp. nov., *Penicillium penaroljense* sp. nov., *Penicillium vanderhammenii* sp. nov. and *Penicillium wotroi* sp. nov., isolated from leaf litter. *International Journal of Systematic and Evolutionary Microbiology* **61**: 1462–1475.
- Houbraken J, Samson RA (2011). Phylogeny of *Penicillium* and the segregation of *Trichocomaceae* into three families. *Studies in Mycology* **70**: 1–51.
- Houbraken J, Spierenburg H, Frisvad JC (2012b). *Rasamsonia*, a new genus comprising thermotolerant and thermophilic *Talaromyces* and *Geosmithia* species. *Antonie van Leeuwenhoek* **101**: 403–421.
- Howard BH, Raistrick H (1950). Studies in the biochemistry of micro-organisms. 81. The colouring matters of *Penicillium islandicum* Sopp. Part 2. Chrysophanic acid, 4:5-dihydroxy-2-methylanthrquinone. *Biochemical Journal* **46**: 49–53.
- Hubka V, Kolarik M (2012). β-tubulin parologue tubC is frequently misidentified as the benA gene in *Aspergillus* section *Nigri* taxonomy: primer specificity testing and taxonomic consequences. *Persoonia* **29**: 1–10.
- Hutchinson CR, Li S-W, McInnes AG, et al. (1983). Comparative biochemistry of fatty acid and macrolide antibiotic (brefeldin-A) formation in *Penicillium brefeldianum*. *Tetrahedron* **39**: 3507–3513.
- Iriart X, Lavergne RA, Filliaux J, et al. (2012). Routine identification of medical fungi by the new Vitek MS Matrix-Assisted Laser Desorption Ionization-Time of Flight System with a new time-effective strategy. *Journal of Clinical Microbiology* **50**: 2107–2110.
- Karahadian C, Josephson DB, Lindsay RC (1985). Volatile compounds from *Penicillium* sp. contributing musty-earthy notes to Brie and Camembert cheese flavors. *Journal of Agricultural and Food Chemistry* **33**: 339–343.
- Kawada M, Momose I, Someno T, et al. (2009). New atpenins, NBRI23477 A and B, inhibit the growth of human prostate cancer cells. *Journal of Antibiotics* **62**: 243–246.
- Kildgaard S, Mansson M, Dosen I, et al. (2014). Accurate dereplication of bioactive secondary metabolites from marine-derived fungi by UHPLC-DAD-QTOFMS and MS/HRMS library. *Marine Drugs* **12**: 3681–3705.
- Klitgaard A, Iversen A, Andersen MR, et al. (2014). Aggressive dereplication using UHPLC-DAD-QTOF – screening extracts for up to 3000 fungal secondary metabolites. *Analytical and Bioanalytical Chemistry* **406**: 1933–1943.
- Kolecka A, Khayhan K, Groenewald M, et al. (2013). Identification of medically relevant species of arthroconidial yeasts by use of matrix-assisted laser desorption ionization-time of flight mass spectrometry. *Journal of Clinical Microbiology* **51**: 2491–2500.
- Kornerup A, Wanscher JH (1967). *Methuen handbook of colour*. Methuen & Co Ltd, Copenhagen.
- Köljalg U, Larsson K-H, Abarenkov K, et al. (2005). UNITE: a database providing web-based methods for the molecular identification of ectomycorrhizal fungi. *New Phytologist* **166**: 1063–1068.
- Köljalg U, Nilsson RH, Abarenkov K, et al. (2013). Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* **22**: 5271–5277.
- Li Y, Cui F, Liu Z, et al. (2007). Improvement of xylanase production by *Penicillium oxalicum* ZH-30 using response surface methodology. *Enzyme and Microbial Technology* **40**: 1381–1388.
- Link HF (1809). Observations in Ordines plantarum naturales. *Dissertatio 1ma. Magazin der Gesellschaft Naturforschenden Freunde Berlin* **3**: 3–42.
- Liu YJ, Whelen S, Hall BD (1999). Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* **16**: 1799–1808.
- Liu W, Gu Q, Zhu W, et al. (2005). Penicillones A and B, two novel polyketides with tricyclo [5.3.1.03,8] undecane skeleton, from amarine-derived fungus *Penicillium terrestre*. *Tetrahedron Letters* **46**: 4993–4996.
- López-Díaz T-M, Santos J-A, García-López M-L, et al. (2001). Surface mycoflora of a Spanish fermented meat sausage and toxicogenicity of *Penicillium* isolates. *International Journal of Food Microbiology* **68**: 69–74.
- LoBuglio KF, Pitt JI, Taylor JW (1993). Phylogenetic analysis of two ribosomal DNA regions indicates multiple independent losses of a sexual *Talaromyces* state among asexual *Penicillium* species in subgenus *Biverticillium*. *Mycologia* **85**: 592–604.
- Ludemann V, Greco M, Rodriguez MP, et al. (2010). Conidial production by *Penicillium nalgioense* for use as starter cultures in dry fermented sausages by solid state fermentation. *LWT – Food Science and Technology* **43**: 315–318.
- Lund F (1995). Differentiating *Penicillium* species by detection of indole metabolites using a filter paper method. *Letters in Applied Microbiology* **20**: 228–231.
- Masclaux F, Guého E, Hoog GS de, et al. (1995). Phylogenetic relationships of human-pathogenic *Cladosporium* (*Xylohypha*) species inferred from partial LS rRNA sequences. *Medical Mycology* **33**: 327–338.
- McNeill J, Barrie FF, Buck WR, et al. (eds) (2012). *International code of nomenclature for algae, fungi, and plants (Melbourne Code)*. Koeltz Scientific Books, Königstein. [Regnum vegetable no. 154.]
- Min XJ, Hickey DA (2007a). Assessing the effect of varying sequence length on DNA barcoding of fungi. *Molecular Notes* **7**: 365–373.
- Min XJ, Hickey DA (2007b). DNA barcodes provide a quick preview of mitochondrial genome composition. *PLoS One* **3**: e325.
- Nelson JH (1970). Production of blue cheese flavor via submerged fermentation by *Penicillium roqueforti*. *Journal of Agricultural and Food Chemistry* **18**: 567–569.
- O'Donnell K, Cigelnik E, Nirenberg HI (1998). Molecular systematics and phylogeny of the *Gibberella fujikuroi* species complex. *Mycologia* **90**: 465.
- Okuda T (1994). Variation in colony characteristics of *Penicillium* strains resulting from minor variations in culture conditions. *Mycologia* **86**: 259–262.
- Okuda T, Klich MA, Seifert KA, et al. (2000). Media and incubation effect on morphological characteristics of *Penicillium* and *Aspergillus*. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification* (Samson RA, Pitt JI, eds). Harwood Academic Publishers, Amsterdam: 83–99.
- Omura S, Tomoda H, Kimura K, et al. (1988). Atpenins, new antifungal antibiotics produced by *Penicillium* sp. — production, isolation, physicochemical and biological properties. *Journal of Antibiotics* **41**: 1769–1774.
- Paterson RRM, Bridge PD, Crosswaite MJ, et al. (1989). A reappraisal of the terverticillate *Penicillia* using biochemical, physiological and morphological features iii. An evaluation of pectinase and amylase isoenzymes for species characterization. *Journal of General Microbiology Journal of General Microbiology* **135**: 2979–2991.
- Peterson SW (2000a). Phylogenetic analysis of *Penicillium* species based on ITS and LSU-rDNA nucleotide sequences. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification* (Samson RA, Pitt JI, eds). Harwood Academic Publishers, Amsterdam: 163–178.
- Peterson SW (2000b). Phylogenetic relationships in *Aspergillus* based on rDNA sequence analysis. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification* (Samson RA, Pitt JI, eds). Harwood Academic Publishers, Amsterdam: 323–355.
- Peterson SW (2008). Phylogenetic analysis of *Aspergillus* species using DNA sequences from four loci. *Mycologia* **100**: 205–226.
- Peterson SW, Jurjević Z (2013). *Talaromyces columbinus* sp. nov., and genealogical concordance analysis in *Talaromyces* Clade 2a. *PLoS One* **8**: e78084.
- Peterson SW, Orchard SS, Menon S (2011). *Penicillium menonorum*, a new species related to *P. pimiteiouiense*. *IMA Fungus* **2**: 121–125.
- Peterson SW, Vega F, Posada F, et al. (2005). *Penicillium coffeae*, a new endophytic species isolated from a coffee plant and its phylogenetic relationship to *P. fellutanum*, *P. thiersii* and *P. brocae* based on parsimony analysis of multilocus DNA sequences. *Mycologia* **97**: 659–666.
- Pitt JI (1973). An appraisal of identification methods for *Penicillium* species. Novel taxonomic criteria based on temperature and water relations. *Mycologia* **65**: 1135–1157.
- Pitt JI (1979). *The genus Penicillium and its teleomorphic states Eupenicillium and Talaromyces*. Academic Press Inc, London.
- Pitt JI, Hocking AD (2009). *Fungi and Food Spoilage*. Springer, New York.
- Pitt JI, Samson RA (1993). Names in current use in the family *Trichocomaceae*. In: *Names in current use in the family Trichocomaceae, Cladoniaceae, Pinaceae, and Lumnaceae* (Greuter W, ed). Koeltz Scientific Books, Königstein: 13–57.
- Pitt JI, Samson RA, Frisvad JC (2000). List of accepted species and their synonyms in the family *Trichocomaceae*. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification* (Samson RA, Pitt JI, eds). Harwood Academic Publishers, Amsterdam: 9–79.

- Rai JN, Tewari JP, Mukerji KG (1964). A new *Aspergillus* from Indian soils: *A. striatus* spec. nov. *Canadian Journal of Botany* **42**: 1521–1524.
- Rai JN, Tewari JP, Sinha AK (1967). Effect of environmental conditions on sclerotia and cleistothecia production in *Aspergillus*. *Mycopathologia et mycologia applicata* **31**: 209–224.
- Ramírez C (1982). *Manual and atlas of the Penicillia*. Elsevier Biomedical Press, Amsterdam.
- Raper KB, Fennell DI (1965). *The genus Aspergillus*. The Williams & Wilkins Company, Baltimore.
- Raper KB, Thom C (1949). *A manual of the penicillia*. The Williams & Wilkins Company, Baltimore.
- Ratnasingham S, Hebert PDN (2007). Bold: The Barcode of Life Data System. *Molecular Ecology Notes* **7**: 355–364.
- Ridgeway R (1912). *Color standards and color nomenclature*. Washington DC.
- Rivera KG, Diaz J, Chavarría-Díaz F, et al. (2012). *Penicillium mallochii* and *P. guanacastense*, two new species isolated from Costa Rican caterpillars. *Mycotaxon* **119**: 315–328.
- Rivera KG, Seifert KA (2011). A taxonomic and phylogenetic revision of the *Penicillium sclerotiorum* complex. *Studies in Mycology* **70**: 139–158.
- Samson RA, Houbraken J, Thrane U, et al. (2010). *Food and indoor fungi*. CBS KNAW Biodiversity Center, Utrecht.
- Samson RA, Pitt JI (1985). General recommendations. In: *Advances in Penicillium and Aspergillus systematics* (Samson RA, Pitt JI, eds). Plenum Press, London: 455–460.
- Samson RA, Seifert KA, Kuijpers A, et al. (2004). Phylogenetic analysis of *Penicillium* subgenus *Penicillium* using partial β -tubulin sequences. *Studies in Mycology* **49**: 175–200.
- Samson RA, Visagie CM, Houbraken J, et al. (2014). Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Studies in Mycology* **78**: 141–173.
- Samson RA, Yilmaz N, Houbraken J, et al. (2011). Phylogeny and nomenclature of the genus *Talaromyces* and taxa accommodated in *Penicillium* subgenus *Biverticillium*. *Studies in Mycology* **70**: 159–183.
- Santamaría M, Fosso B, Consiglio A, et al. (2012). Reference databases for taxonomic assignment in metagenomics. *Briefings in Bioinformatics* **13**: 682–695.
- Sarbhoy AK, Elphick JJ (1968). *Hemicarpenteles paradoxus* gen. & sp.nov.: the perfect state of *Aspergillus paradoxus*. *Transactions of the British Mycological Society* **51**: 155–157.
- Schoch CL, Robbertse B, Robert V, et al. (2014). Finding needles in haystacks: linking scientific names, reference specimens and molecular data for fungi. *Database* **2014**. <http://dx.doi.org/10.1093/database/bau061>.
- Schoch CL, Seifert KA, Huhndorf S, et al. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for fungi. *Proceedings of the National Academy of Sciences of the United States of America* **109**: 6241–6246.
- Seifert KA, Louis-Seize G (2000). Phylogeny and species concepts in the *Penicillium aurantiogriseum* complex as inferred from partial β -tubulin gene DNA sequences. In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification* (Samson RA, Pitt JI, eds). Harwood Academic Publishers, Amsterdam: 189–198.
- Seifert KA, Samson RA, deWaard JR, et al. (2007). Prospects for fungus identification using CO1 DNA barcodes, with *Penicillium* as a test case. *Proceedings of the National Academy of Sciences of the United States of America* **104**: 3901–3906.
- Serra R, Peterson SW, CTCOR, et al. (2008). Multilocus sequence identification of *Penicillium* species in cork bark during plank preparation for the manufacture of stoppers. *Research in Microbiology* **159**: 178–186.
- Shang Z, Li X, Meng L, et al. (2012). Chemical profile of the secondary metabolites produced by a deep-sea sediment derived fungus *Penicillium commune*. *Chinese Journal of Oceanology and Limnology* **30**: 305–314.
- Siegrist TJ, Anderson PD, Huen WH, et al. (2007). Discrimination and characterization of environmental strains of *Escherichia coli* by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF-MS). *Journal of Microbiological Methods* **68**: 554–562.
- Skouboe P, Frisvad JC, Taylor JW, et al. (1999). Phylogenetic analysis of nucleotide sequences from the ITS region of terverticillate *Penicillium* species. *Mycological Research* **103**: 873–881.
- Smedsgaard J (1997). Micro-scale extraction procedure for standardized screening of fungal metabolite production in cultures. *Journal of Chromatography A* **760**: 264–270.
- Stolk AC, Samson RA (1983). The Ascomycete genus *Eupenicillium* and related *Penicillium* anamorphs. *Studies in Mycology* **23**: 1–149.
- Tamura M, Kawahara K, Sugiyama J (2000). Molecular phylogeny of *Aspergillus* and associated teleomorphs in the *Trichocomaceae* (Eurotiales). In: *Integration of modern taxonomic methods for Penicillium and Aspergillus classification* (Samson RA, Pitt JI, eds). Harwood Academic Publishers, Amsterdam: 357–372.
- Tautz D, Arctander P, Minelli A, et al. (2003). A plea for DNA taxonomy. *Trends in Ecology and Evolution* **18**: 70–74.
- Taylor JW, Jacobson DJ, Kroken S, et al. (2000). Phylogenetic species recognition and species concepts in fungi. *Fungal Genetics and Biology* **31**: 21–32.
- Terrasan CRF, Temer B, Duarte MCT, et al. (2010). Production of xylanolytic enzymes by *Penicillium janczewskii*. *Bioresource Technology* **101**: 4139–4143.
- Thom C (1906). Fungi in cheese ripening: Camembert and Roquefort. *U.S. Department of Agriculture, Bureau of Animal Industry – Bulletin* **82**: 1–39.
- Thom C (1930). *The Penicillia*. The Williams & Wilkins Company, Baltimore.
- Thom C (1945). Mycology present penicillin. *Mycologia* **37**: 460–475.
- Thom C (1954). The evolution of species concepts in *Aspergillus* and *Penicillium*. *Annals of the New York Academy of Sciences* **60**: 24–34.
- Varga J, Due M, Frisvad JC, et al. (2007). Taxonomic revision of *Aspergillus* section *Clavati* based on molecular, morphological and physiological data. *Studies in Mycology* **59**: 89–106.
- Varga J, Frisvad JC, Samson RA (2011). Two new aflatoxin producing species, and an overview of *Aspergillus* section *Flavi*. *Studies in Mycology* **69**: 57–80.
- Verwer PEB, Leeuwen WB, Girard V, et al. (2013). Discrimination of *Aspergillus lentulus* from *Aspergillus fumigatus* by Raman spectroscopy and MALDI-TOF MS. *European Journal of Clinical Microbiology & Infectious Diseases* **33**: 245–251.
- Visagie CM, Hirooka Y, Tanney JB, et al. (2014a). *Aspergillus*, *Penicillium* and *Talaromyces* isolated from in house dust samples collected around the world. *Studies in Mycology* **78**: 63–139.
- Visagie CM, Houbraken J, Rodrigues C, et al. (2013). Five new *Penicillium* species in section *Sclerotiora*: a tribute to the Dutch Royal family. *Persoonia* **31**: 42–62.
- Visagie CM, Seifert KA, Houbraken J, et al. (2014b). Diversity of *Penicillium* section *Citrina* within the fynbos biome of South Africa, including a new species from a *Protea repens* inflorescence. *Mycologia* **106**: 537–552.
- Wang B, Wang L (2013). *Penicillium kongii*, a new terverticillate species isolated from plant leaves in China. *Mycologia* **105**: 1547–1554.
- Wang J, Huang Y, Fang M, et al. (2002). Brefeldin A, a cytotoxin produced by *Paecilomyces* sp. and *Aspergillus clavatus* isolated from *Taxus mairei* and *Torreya grandis*. *FEMS Immunology & Medical Microbiology* **34**: 51–57.
- White TJ, Bruns T, Lee S, et al. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: *PCR protocols: a guide to methods and applications* (Innis MA, Gelfand DH, Shinsky TJ, White TJ, eds). Academic Press Inc, New York: 315–322.
- Welham KJ, Domin MA, Johnson K, et al. (2000). Characterization of fungal spores by laser desorption/ionization time-of-flight mass spectrometry. *Rapid Communications in Mass Spectrometry* **14**: 307–310.
- Yaguchi T, Someya A, Miyadoh S, et al. (1993). *Aspergillus ingratus*, a new species in *Aspergillus* section *Clavati*. *Transactions of the Mycological Society of Japan* **34**: 305–310.
- Yilmaz N, Visagie CM, Houbraken J, et al. (2014). Polyphasic taxonomy of the genus *Talaromyces*. *Studies in Mycology* **78**: 175–341.
- Zhao Y, Chen H, Shang Z, et al. (2012). SD-118 – Xanthocillin X (1). A novel marine agent extracted from *Penicillium commune*, induces autophagy through the inhibition of the MEK/ERK pathway. *Marine Drugs* **10**: 1345–1359.