
Plant Propagation Protocol for Lotus aboriginus Jeps.

ESRM 412 – Native Plant Production

Protocol URL: https://courses.washington.edu/esrm412/protocols/LOAB.pdf



© 1995 Saint Mary's College of California⁷



TAXONOMY	
Plant Family	
Scientific Name	Fabaceae ¹
Common Name	Pea family ¹
Species Scientific Name	
Scientific Name	Lotus aboriginus Jeps. ¹
Varieties	
Sub-species	
Cultivar	
Common Synonym(s)	Hosackia rosea Eastw. ¹
	Lotus crassifolius (Benth.) Greene var. subglaber
	(Ottley) C.L. Hitchc. ¹
	Lotus stipularis (Benth.) Greene var. subglaber Ottley ¹
Common Name(s)	rosy bird's-foot trefoil ¹ , thicket trefoil ² , rose flowered
	lotus ³ , rosy lotus ⁷
Species Code (as per USDA Plants database)	LOAB

GENERAL INFORMATION		
Geographical range	In North America, <i>L. aboriginus</i> is only found along the west coast of the United States, in Washington, Oregon, and California ^{1,2} .	
	In Washington, <i>L. aboriginus</i> is found only in 7 counties on the western side of the state: Jefferson, Grays Harbor, Pacific, Mason, Kitsap, Thurston, and Pierce. It is similarly found in western counties of Oregon and northern California ¹ .	
	See maps above.	
Ecological distribution	L. aboriginus is found in grasslands and forests ⁴ .	
Climate and elevation range	60-1200 m ⁴	
Local habitat and abundance	<i>L. aboriginus</i> grows in woods, coastal mountains, and disturbed habitat such as along roadsides ⁴ , as well as on banks, streamsides, burns, and logged areas ⁵ .	
Plant strategy type / successional stage		
Plant characteristics	<i>L. aboriginus</i> is a forb, dicot, and perennial herb ^{1,3,4} .	
	The stem can be either spreading or erect, and 1-7 dm in height ⁵ with alternating leaves ⁸ . The stipules of its leaves are scarious, inconspicuous, and fragile ⁵ . Its leaflets are elliptical and 1.5-3 cm in length, about 2-3 times longer than they are wide. L. aboriginus flowers from May to July ^{5,6} , and its flowers grow in groups of 6-10 and are white or pink in color and striated ⁵ . The flowers darken with age, and have a claw coming out from the calyx tube. The calyx is 4-6 mm long, and the flower's lobes are shorter than the tube. Its fruit is 3-5 cm long, 2-4 mm wide, oblong, and glabrous, with several seeds ⁵ . The plant can be observed from the beginning of April	
DD OD	through the 3 rd week of September ⁸ .	
	AGATION DETAILS	
	No propagation protocols found for <i>Lotus aboriginus</i> or its close relatives	
Ecotype Propagation Goal		
Propagation Goal Propagation Method		
Product Type		
Stock Type		
Time to Grow		
Target Specifications		
Tangot operationations	I .	

Propagule Collection Instructions	
Propagule Processing/Propagule	
Characteristics	
Pre-Planting Propagule Treatments	
Growing Area Preparation / Annual	
Practices for Perennial Crops	
Establishment Phase Details	
Length of Establishment Phase	
Active Growth Phase	
Length of Active Growth Phase	
Hardening Phase	
Length of Hardening Phase	
Harvesting, Storage and Shipping	
Length of Storage	
Guidelines for Outplanting /	
Performance on Typical Sites	
Other Comments	
INFORMATION SOURCES	
References	See below
Other Sources Consulted	See below
Protocol Author	Kyra Woytek
Date Protocol Created or Updated	5/29/19

References:

- "Plants Profile for Lotus Aboriginus (Rosy Bird's-Foot Trefoil)." *USDA NRCS Plants Database*, plants.usda.gov/core/profile?symbol=LOAB.
- "Comprehensive Report Species Lotus Aboriginus." *NatureServe Explorer*, explorer.natureserve.org/servlet/NatureServe?searchName=Lotus%2Baboriginus.
- "Lotus Aboriginus." *Calflora*, www.calflora.org/cgi-bin/species_query.cgi?where-calrecnum=5018.
- Contu, S. "Lotus Aboriginus." *IUCN Red List of Threatened Species*, 2012, www.iucnredlist.org/species/19892492/20079094.
- Brouillet, Luc. "Hosackia Rosea." *The Jepson Herbarium University of California, Berkeley*, 2012, ucjeps.berkeley.edu/eflora/eflora_display.php?tid=28430.
- "Hosackia Rosea." *Calflora*, www.calflora.org/cgi-bin/species_query.cgi?where-taxon=Hosackia%2Brosea.

- Brousseau, Alfred. "Hosackia Rosea; Rosy Lotus." *CalPhotos Photo Database*, 1995, calphotos.berkeley.edu/cgi/img_query?seq_num=8289&one=T.
- "Hosackia Rosea | Rosy Bird's-Foot Trefoil." *WildflowerSearch.org*, wildflowersearch.org/search?oldstate=name%3AHosackia%3B&PlantName=&S__514876.x=87&S__514876.y=130.

Other Sources Consulted:

- European Bioinformatics InstituteProtein Information ResourceSIB Swiss Institute of Bioinformatics. "Taxonomy Hosackia Rosea." *UniProt*, European Bioinformatics Institute, Protein Information Resource, SIB Swiss Institute of Bioinformatics, www.uniprot.org/taxonomy/390714.
- "Hosackia Rosea Eastw." *Plants of the World Online | Kew Science*, www.plantsoftheworldonline.org/taxon/urn:lsid:ipni.org:names:124208-2.
- "Hosackia Rosea Eastw." *The Plant List A Working List for All Plant Species*, www.theplantlist.org/tpl/record/ild-14295.
- "Lotus Aboriginus Rosy Birds-Foot Trefoil." *Discover Life*, www.discoverlife.org/mp/20q?search=Lotus%2Baboriginus.
- "Lotus Aboriginus Jeps." *Tropicos*, www.tropicos.org/Name/13009993.
- "Lotus Aboriginus." EOL, eol.org/pages/48148611.
- "Lotus Aboriginus." *ITIS*, www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=503554#null .
- "Search Results Hosackia Rosea." *Consortium of Pacific Northwest Herbaria*, www.pnwherbaria.org/data/results.php?DisplayAs=WebPage&ExcludeCultivated=Y&Gro upBy=ungrouped&SortBy=Year&SortOrder=DESC&SearchAllHerbaria=Y&QueryCount =1&IncludeSynonyms1=Y&Genus1=Lotus&Species1=crassifolius&Zoom=4&Lat=55&L ng=-135&PolygonCount=0.
- "Taxon: Lotus Aboriginus Jeps." *Germplasm Resources Information Network*, USDA, Agricultural Research Service, National Plant Germplasm System, npgsweb.ars-grin.gov/gringlobal/taxonomydetail.aspx?id=408929.
- "Taxonomy Browser Lotus Aboriginus." *National Center for Biotechnology Information*, U.S. National Library of Medicine, www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=118905.