



Molecular systematics of Poaceae based on eight chloroplast markers, emphasizing the phylogenetic positions of Korean taxa

Jung-Hoon LEE¹, Ki-Joong KIM², Bo-Yun KIM³ and Young-Dong KIM^{1,4*}

¹Multidisciplinary Genome Institute, Hallym University, Chuncheon 24252, Korea

²Division of Life Sciences, Korea University, Seoul 02841, Korea

³National Institute of Biological Resources, Incheon 22689, Korea

⁴Department of Life Science, Hallym University, Chuncheon 24252, Korea

Corresponding author

Young-Dong KIM
E-mail: ydkim@hallym.ac.kr

ABSTRACT: This study was conducted to clarify the phylogenetic position and relationships of Korean Poaceae taxa. A total of 438 taxa including 155 accessions of Korean Poaceae (representing 92% and 72% of Korean Poaceae genera and species, respectively) were employed for phylogeny reconstruction. Sequence data of eight chloroplast DNA markers were used for molecular phylogenetic analyses. The resulted phylogeny was mostly concordant with previous phylogenetic hypotheses, especially in terms of subfamilial and tribal relationships. Several taxa-specific indels were detected in the molecular phylogeny, including a 45 bp deletion in *rps3* (PACMAD [Panicoidae, Arundinoideae, Chloridoideae, Micrairoideae, Aristidoideae, Danthonioideae] clade), a 15 bp deletion in *ndhF* (Oryzoae + Phyllorachideae), a 6 bp deletion in *trnL-F* (Poeae s.l.), and two (17 bp and 378 bp) deletions in *atpF-H* (Pooideae). The Korean Poaceae members were classified into 23 tribes, representing eight subfamilies. The subfamilial and tribal classifications of the Korean taxa were generally congruent with a recently published system, whereas some subtribes and genera were found to be non-monophyletic. The taxa included in the PACMAD clade (especially Andropogoneae) showed very weak and uncertain phylogenetic relationships, presumably to be due to evolutionary radiation and polyploidization. The reconstructed phylogeny can be utilized to update the taxonomic positions of the newly examined grass accessions.

Keywords: Andropogoneae, economic plants, indel, monocots, paraphyletic, polyphyletic, *rps3*

RECEIVED 27 July 2022; **REVISED** 17 September 2022; **ACCEPTED** 22 September 2022

OPEN ACCESS

© 2022 the Korean Society of Plant Taxonomists.
This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Poaceae (nom. alt. Gramineae) are the fifth largest family of angiosperms and the second largest family of monocots, containing about 700 genera and 11,000 to 12,000 species (Clayton and Renvoize, 1986; Watson and Dallwitz, 1992; Kellogg, 2015; Soreng et al., 2017). The members of Poaceae are distributed on all continents and are estimated to account for approximately 20% of the earth's vegetation (Shantz, 1954). They are also important food resources for mankind. For example, rice (*Oryza sativa* L.), wheat (*Triticum aestivum* L.), and maize (*Zea mays* L.) are vital food resources that account for more than half of the calories consumed by humans (Raven and Johnson, 1995).

Due to the tremendous ecological and economic importance of Poaceae, numerous taxonomic and phylogenetic studies focusing on this species-rich family have been conducted (Brown, 1810; Palisot de Beauvois, 1812; Kunth, 1833; Tateoka, 1957; Prat, 1960; Stebbins and Crampton, 1961; Jacques-Félix, 1962; Hilu and Wright, 1982; Caro, 1982; Clayton and Renvoize, 1986; Kellogg and Campbell, 1987; Hamby and Zimmer, 1988; Doebley et al., 1990; Watson and Dallwitz, 1992; Cummings et al., 1994; Nadot et al., 1994; Barker et al., 1995; Clark et al., 1995; Soreng and Davis, 1998; Clark et al., 2000; Grass Phylogeny Working Group, 2001; Duvall et al., 2007; Simon, 2007; Bouchenak-Khelladi et al., 2008; Saarela and Graham, 2010; Grass Phylogeny Working Group II, 2011; Kellogg, 2015; Soreng et al., 2017;

Saarela et al., 2018; Soreng et al., 2022). Although these studies have led to major changes thus far, the classification system of Poaceae has gradually been stabilizing. Currently, lineages of Poaceae are classified into 12 subfamilies, including three basal subfamilies (Anomochlooideae, Pharoideae, Puelioideae) and two major clades, namely the BOP (Bambusoideae, Oryzoideae, Pooideae) and PACMAD (Panicoideae, Arundinoideae, Chloridoideae, Micrairoideae, Aristidoideae, Danthonioideae) clades (Soreng et al., 2017; Saarela et al., 2018). Attempts have also been made to provide stronger evidence of the phylogeny of certain grass lineages that were not fully resolved in previous studies using plastomes or nuclear genes (Wu and Ge, 2012; Fisher et al., 2016; Dunning et al., 2019; Huang et al., 2022).

Among Korean flora, Poaceae are the third largest family, comprising 212 species representing six subfamilies, 21 tribes, and 94 genera (Lee, 2007). Relatively few taxonomic studies, however, have focused on Korean grasses, although there have been some morphological examinations of well-known genera, such as *Miscanthus* Andersson (Lee, 1964a, 1964b, 1964c, 1964d) and *Poa* L. (Jung and Chung, 2008). There have also been molecular phylogenetic studies of common genera, including *Echinochloa* P. Beauv. (Lee et al., 2014) and *Zoysia* Willd. (Cheon et al., 2021; Oh et al., 2021), but with limited taxon sampling. Thus far, researchers have disregarded the overall phylogenetic relationships and classifications of Korean Poaceae taxa. Therefore, it is necessary to review the current classification system based on detailed phylogenetic information of each plant species for the efficient management and conservation of Korean Poaceae taxa.

In this paper, we present the molecular phylogeny of

Poaceae based on the sequences of eight chloroplast markers (*atpF*-H, *psbK*-I, *trnH-psbA*, *trnL*-F, *matK*, *rbcL*, *ndhF*, *rps3*), emphasizing the phylogenetic relationships or positions of Korean taxa within the family. We address taxonomic issues pertaining to phylogenetically problematic poaceae groups for further comprehensive research.

MATERIALS AND METHODS

Taxon sampling

A total of 438 taxa (species or infraspecies) representing 12 subfamilies, 52 tribes, and 297 genera of Poaceae were included in the phylogenetic study. Of these, 155 Korean grass taxa representing 87 genera were newly sampled for the present study (Appendix 1). These correspond to 92% (at the genus level) and 72% (at the species level or below) of all known Poaceae taxa in Korea (Lee, 2007). Species lacking collecting records, rare species, or those only distributed in North Korea were not included in the present study. The names of subfamilies, tribes, and subtribes accepted in the phylogenetic classification system of the Poaceae (Soreng et al., 2017, 2022) were employed throughout the study. The acceptance of scientific names of several controversial taxa was based on The Plant List v1.1 (2013) (<http://www.theplantlist.org>), Tropicos v3.3.2 (2022) (<http://www.tropicos.org>), and Plants of the World Online (2022) (<http://www.plantsoftheworldonline.org>).

Most of the samples for DNA extraction were collected from the field. Fresh young leaves or leaves dried with silica gel were used for DNA extraction. The DNA of some species was obtained from dried specimens deposited at the herbarium of the National Arboretum (KH) and the herbarium

Table 1. Sequence and phylogenetic information of the eight plastid markers employed for the phylogenetic analyses of Poaceae (excluding outgroups).

Information	Number of examined taxa	Sequence type	Size range (bp)	Aligned length (bp)	Applied model for BI (AIC)
<i>atpF</i> -H	239	non-coding	439-738	1,087	TVM+I+G
<i>psbK</i> -I	215	non-coding	168-438	741	TPM1uf+G
<i>trnH-psbA</i>	249	non-coding	436-783	1,037	TPM1uf+I+G
<i>trnL</i> -F	316	non-coding	242-468	997	TIM1+G
<i>matK</i>	403	coding	632-1,096	1,117	GTR+I+G
<i>ndhF</i>	385	coding	206-805	1,010	GTR+I+G
<i>rbcL</i>	394	coding	438-743	743	TVM+I+G
<i>rps3</i>	247	coding	486-717	735	GTR+I+G
non-coding	349	non-coding	281-2,303	3,862	TPM1uf+I+G
coding	430	coding	292-3,343	3,605	GTR+I+G
All combined	438	mixed	292-5,556	7,467	GTR+I+G




Fig. 1. Structure of eight plastid regions examined for the present study. Arrows indicate the forward and reverse primers employed for polymerase chain reaction and sequencing. **A.** *atpF-H* (Lahaye et al., 2008). **B.** *trnH-psbA*. **C.** *psbK-I* (Lahaye et al., 2008). **D.** *trnL-F* (Taberlet et al., 1991). **E.** *matK* (Cuénoud et al., 2002; Johnson and Soltis., 1995). **F.** *ndhF* (Romaschenko et al., 2010). **G.** *rbcL* (Fay et al., 1997; Olmstead et al., 1992). **H.** *rps3* (Peterson et al., 2010).

of Hallym University (HHU). Voucher specimens of the plants used for DNA extraction were deposited at the HHU and the herbarium of the National Institute of Biological Resources of Korea (KB) (Appendix 1).

Eight plastid genome markers including four non-coding regions (*atpF-H*, *psbK-I*, *trnH-psbA*, and *trnL-F*) and four coding regions (*matK*, *ndhF*, *rbcL*, and *rps3*) were employed for the molecular phylogeny reconstruction (Fig. 1). A total of 154 *atpF-H* sequences, 130 *psbK-I* sequences, 152 *trnH-psbA* sequences, 130 *trnL-F* sequences, 129 *matK* sequences, 115 *ndhF* sequences, 131 *rbcL* sequences, 143 *rps3* sequences were newly obtained for the Korean Poaceae taxa (Appendix 1). Various number of sequence data (85 *atpF-H*, 85 *psbK-I*, 98 *trnH-psbA*, 192 *trnL-F*, 281 *matK*, 274 *ndhF*, 270 *rbcL*, and 108 *rps3*; see Appendix 1) were downloaded from the GenBank and combined with the newly obtained sequences. Altogether, variable number of nucleotide sequences (239 *atpF-H*, 215 *psbK-I*, 249 *trnH-psbA*, 316 *trnL-F*, 403 *matK*, 385 *ndhF*, 394 *rbcL*, and 247 *rps3*) were included for the

phylogenetic analyses (Table 1). Seven taxa in neighboring families, Joinvilleaceae and Ecteiocoleaceae were selected as outgroups in the phylogenetic analyses (Stevens, 2017).

DNA extraction and polymerase chain reaction

DNA was extracted from fresh leaves using CTAB (Doyle and Doyle, 1987) or DNeasy plant mini-kits (Qiagen Inc., Hilden, Germany) following the manufacturer's instructions in each case. The extracted DNA was electrophoresed on 1% agarose gel, stained with ethidium bromide, and compared with the markers on the UV trans-illuminator for DNA quality and for a concentration check. Polymerase chain reaction (PCR) was performed with a final volume of 50 μ L, and the concentration of the PCR reaction mixture was 5 units/ μ L TaKaRa Ex Taq, 20 mM Mg^{2+} plus 10 \times Ex Taq Buffer, 2.5 mM dNTP Mixture (TaKaRa Biotechnology Co., Dalian, China), template DNA 20–50 ng, 5 pmol-specific primer pair (Fig. 1), and distilled water. The PCR cycling conditions were pre-denaturation at 95 $^{\circ}$ C for 5 min, followed by 94–95 $^{\circ}$ C

denaturation for 40–60 s, annealing for 52–57°C for 40–60 s and 72°C for 60–90 s up to 35 cycles, with the final stabilization step carried out at 72°C for 10 min. The amplified PCR product was purified with a QIAquick PCR purification kit (Qiagen Inc., Valencia, CA, USA), and all procedures followed the supplier's instructions.

DNA sequencing and alignment

The PCR products were directly sequenced using an automatic sequencer, in this case an ABI Prism 3730xl DNA analyzer (Applied Biosystems, Foster City, CA, USA). Sequence alignments for each marker were performed using MUSCLE (multiple sequence comparison by log-expectation) (Edgar, 2004), which is implemented in Geneious Prime v. 2022.1.1 (Biomatters Ltd., Auckland, New Zealand) (Kearse et al., 2012). Subsequently, further alignment was conducted using MAFFT (Katoh et al., 2002), as needed, and the alignment was finally completed by manual adjustment.

Phylogenetic analyses

A total of 2,448 concatenated sequences of the eight plastid markers were used for the phylogenetic analyses (Table 1). Information about the aligned sequence lengths and the sequence divergence values of each marker are shown in Table 1. The validity of the sequence concatenation process was tested via an incongruence length difference test (Farris et al., 1994), in which the p-value ($p = 0.001$) for the partition homogeneity test was calculated using PAUP ver. 4.0a147 (Swofford, 2002) (performed with 1,000 replicates, simple addition sequence replicates, MulTrees, and TBR branch swapping, gaps treated as missing).

The phylogenetic tree was reconstructed using the Bayesian inference (BI) method. The BI analysis was conducted using MrBayes v3.2.2 (Ronquist and Huelsenbeck, 2003). jModelTest 2.1 (Guindon and Gascuel, 2003; Darriba et al., 2012) was used to evaluate the appropriate evolution model of the data combined with each dataset. The optimal model search for the BI analysis was selected using the Akaike information criterion (Akaike, 1974) method. Two independent analyses were performed using four Markov chain Monte Carlo = simulations, applying the general time reversible model, the gamma distribution of substitution rates (G), and estimates of invariant characters (I). Analyzing 10,000,000 generations, we sampled the trees every 200 generations and removed 25% of the sampled trees by burn-in processing. After burn-in removal, we calculated the 50% majority-rule sum tree and posterior probability (PP) using the remaining tree counts. The analyzed phylogenetic trees were confirmed by FigTree ver. 1.4 (Rambaut, 2012). Bootstrap percentage

(BP) values for each clade were calculated to provide additional information on the reliability of the clades observed in the BI tree. To do this, a maximum parsimony heuristic search was repeated 1,000 times using PAUP ver. 4.0a147 (Swofford, 2002) with the simple addition sequence replicate, tree-bisection-reconnection branch swapping, MulTrees, and ACCTRAN options.

RESULTS AND DISCUSSION

Subfamilial phylogenetic relationships

The subfamilial phylogenetic relationships in the BI tree were basically identical to those in previous results (Bouchenak-Khelladi et al., 2010; Edwards and Smith, 2010; Grass Phylogeny Working Group II, 2011; Wu and Ge, 2012; Cotton et al., 2015; Soreng et al., 2015), recognizing three small basal subfamilies (Anomochlooideae, Pharioideae, Puelioideae) and two large and strongly supported groups: the BOP and PACMAC clades (Fig. 2). All members of Korean Poaceae were included in the large clades; therefore, all subsequent discussions on the phylogenetic relationships of Poaceae will focus on the two major clades.

Phylogenetic relationships in the BOP clade

The phylogenetic relationships within the BOP clade differ depending on the author. A study by Gaut (2002), who estimated evolutionary patterns using the chromosome number and genome size of Poaceae, suggested a close relationship between Oryzoideae and Bambusoideae. In contrast, Mathews et al. (2000) found that Oryzoideae was more closely related to Pooideae based on the *phytochrome B* gene phylogeny employing 51 Poaceae taxa. In the present study, Bambusoideae and Pooideae formed a strong sister group with 1.00 PP and 89 BP support values (Fig. 2). This result is consistent with recent phylogenetic studies of BOP clades based on chloroplast genomes (Wu and Ge, 2012). Currently, the relationships among the subfamilies of the BOP clade identified in this study seem to be relatively firmly established (Hilu et al., 1999; Zhang, 2000; Salamin et al., 2002; Bouchenak-Khelladi et al., 2008, 2010; Saarela and Graham, 2010; Grass Phylogeny Working Group II, 2011).

Korean members of Poaceae belonging to the BOP clade were classified into 12 tribes (Fig. 2). The phylogenetic relationships and classification of each subfamily are as follows.

Oryzoideae: Oryzoideae was grouped into the three monophyletic tribes of Oryzeae, Phyllorachideae, and Ehrharteae. It is noteworthy that the “Oryzeae + Phyllorachideae” clade shared a 15 bp deletion in *ndhF*, although the clade





Fig. 2. Bayesian consensus tree based on the combined sequences of eight markers, emphasizing the phylogenetic relationships of the BOP (Bamboisoidae, Oryzoideae, Pooideae) clade. Posterior probabilities (● = 1.00) and bootstrap value (◆ = 100) are shown above or below. Korean taxa are in bold. Taxa with asterisks are either polyphyletic or paraphyletic. The tribal (blue) and subtribal (black or gray) classifications are from Soreng et al. (2015).



received rather weak support (PP 0.77, BP 64). Two Korean genera, *Leersia* and *Zinania*, within tribe Oryzeae formed distinct clades in the two subtribes of Oryzinae and Zizaniinae, respectively (Fig. 2).

Bambusoideae: Bambusoideae consists of three tribes: Arundinarieae, Bambuseae, and Olyreae. All Korean taxa (representing four genera) included in this subfamily are nested in Arundinarieae. Among the four genera, *Sasa* forms a monophyletic group, while *Phyllostachys*, *Pleioblastus*, and *Pseudosasa* were found to be polyphyletic groups, exhibiting unclear interspecies relationships (Fig. 2). Chloroplast DNA phylogeny of temperate bamboo identified six major clades (the *Bergbambos* clade, *Oldeania* clade, *Chimonocalamus* clade, *Shibataea* clade, *Phyllostachys* clade, and *Arundinaria* clade) (Triple and Clark, 2010; Stapleton, 2013) within the subfamily, but the intergeneric and interspecies relationships within each clade remain not well clarified. In the present phylogeny, Korean taxa are divided into the *Phyllostachys* clade and *Arundinaria* clade, but monophyletic or phylogenetic relationships among the genera or species within each clade are not evident. This is thought to be due to the active interspecific gene flow by hybridization that is known to occur in a significant number of temperate bamboos (Triple and Clark, 2010). Low sequence variations of the markers used may also be another reason for the weak relationship, causing some incongruence between the previous morphological classification and the molecular phylogeny. It will be necessary to obtain a large amount of gene sequence data to clarify the taxonomic positions of the Korean taxa. A phylogeny based on genome-wide SNP analyses would be helpful to resolve this problem.

Pooideae: This marks the largest and most strongly supported subfamily in the BOP clade. A 17 bp deletion in *atpF-H* was newly observed in this study, providing additional evidence of the monophyly of the subfamily (Fig. 2). The Korean members (39 genera) of Pooideae included in this study were nested in 18 subtribes representing ten tribes. It should be emphasized that there were two additional deletions, which could indicate the robustness of certain tribal or subtribal relationships (Fig. 2). These were a *trnL-F* 6 bp deletion and *atpF-H* 378 bp deletion, supporting a monophyletic Poaceae s.l. and a clade that consisted of four subtribes (Phleinae, Alopecurinae, Miliinae, and Poinae), respectively. Most of the Korean taxa were monophyletic except seven genera (*Achnatherum*, *Leymus*, *Festuca*, *Trisetum*, *Koeleria*, and *Calamagrostis*), which were found to be either paraphyletic or polyphyletic.

Achnatherum (exclusively represented by three Korean species in this study) was polyphyletic within tribe Stipeae

as *A. coreanum* (\equiv *Patis coreana* (Honda) Ohwi) clustered with *A. mongholicum* (\equiv *Ptilagrostis mongholica* (Turcz. ex Trin.) Griseb.), while *A. pekinense* (\equiv *Stipa pekinensis* Hance) formed a sister group with *Piptatherum miliaceum* (Fig. 2). It should be noted that the taxonomic boundary of *Achnatherum* has been controversial, as each of the three species has been treated as a member of the three different genera of *Stipa*, *Patis*, and *Ptilagrostis*, respectively (Clayton and Renvoiz, 1986; Tzvelev, 1989; Barkworth, 2007; Romaschenko et al., 2010, 2011). Expanded taxon sampling and data will clarify the taxonomic identity of *Achnatherum*.

Leymus in tribe Triticeae was found to be polyphyletic. *Leymus mollis*, a Korean accession, formed a clade with the group containing *Hordeum*, among others, whereas *L. angustus* appears as a sister species of the *Elymus* clade (Fig. 2). *Leymus* is known as one of the genera that underwent severe polyploidization in tribe Triticeae (Sha et al., 2010). Phylogenetic analyses and morphological investigations with expanded taxon sampling including *L. chinensis* and *L. secalinus*, which are known to be distributed on the Korean Peninsula, are needed to confirm their phylogenetic relationship.

In the subtribe Loliinae, *Festuca* formed a paraphyletic group. This is consistent with the results of molecular phylogenetic studies of this festucoid group (Torrecilla and Catalán, 2002; Torrecilla et al., 2004). Some species of *Festuca* have shown a close relationship with taxonomically controversial genera such as *Psilurus*, and *Vulpia* (Darbyshire and Warwick, 1992; Charmet et al., 1997; Gaut et al., 2000). Some *Festuca* species form a clade with *Psilurus* and *Vulpia* (Fig. 2). These *Festuca* species have been suggested as polyploids, and the relationships among them have not been clearly resolved. Additional data from chromosomal and phylogenetic analyses using expanded samples would be necessary to clarify the taxonomic issue.

Koeleria macrantha, a Korean accession of *Koeleria*, exhibited a closer relationship with *Gaudinia coarctata* than with *K. castellana*, meaning that *Koeleria* is a paraphyletic group. Similarly, the two Korean species of *Trisetum* (*T. bifidum*, *T. sibiricum*) sampled for this study did not form a clade with *T. cernuum*, rather the latter species showed a sister relationship with *Rostraria azorica*. This finding is concordant with a previous hypothesis based on nuclear internal transcribed spacer (ITS) and chloroplast DNA (*trnT-H*) phylogeny (Quintanar et al., 2007). *Rostraria* and *Trisetum* have been found to be very similar in terms of the epidermal characters of their lemma (Finot et al., 2006). The discrepancy between the results of molecular phylogenetic studies and the morphological traits necessitates a taxonomic reexamination

of these taxa.

Lastly, monophyly of *Calamagrostis* within subtribe Agrostidinae was uncertain as the phylogenetic relationships among two Korean accessions (*C. epigejos* and *C. arundinacea*) and *C. canadensis* remains unresolved (Fig. 2). Saarela et al. (2010) found that some species of *Calamagrostis* form geographically segregated lineages according to nuclear and chloroplast DNA sequence data. The present study, however, did not confirm any Eurasian lineage in which the Korean accessions were included. More comprehensive analyses with additional data, especially pertaining to the nuclear genome, would be necessary to resolve the phylogenetic relationships among *Calamagrostis* species.

Phylogenetic relationships in the PACMAD clade

The PACMAD clade has been considered as a robust phylogenetic group since it was initially proposed by Duvall et al. (2007). The clade was also evident, being supported by the additional character of a 45 bp deletion in *rps3*, which was newly observed in the present study (Fig. 3). We consider the deletion to be an important diagnostic molecular trait to distinguish the PACMAD clade from all other clades of Poaceae. The subfamilial relationship was identical to the *ndhF* + *rbcL* phylogeny, which employed 448 grass species (Soreng et al., 2015). The basal position of Aristidoideae within the PACMAD clade, however, was poorly supported in the present study (0.72 PP, 70 BP). In some studies, Aristidoideae has been found to have close relationships with Danthoioideae and Chloridoideae, but such relationships have not been well supported (Hsiao et al., 1999; Grass Phylogeny Working Group, 2001; Bouchenak-Khelladi et al., 2008, 2010). Cotton et al. (2015) performed a chloroplast genome analysis of the PACMAD clade using a limited number of representative taxa and argued that Panicoideae was the first to diverge in the clade. Lee (2016) also reconstructed the plastid genome (76 coding genes) phylogeny of Poaceae with expanded taxon sampling, but the basal position Aristidoideae varies depending on the phylogenetic method used.

Our study supported the subfamilial delimitation and relationships observed in previous studies (Clark et al., 1995; Mathews and Sharrock, 1996; Hilu et al., 1999; Sánchez-Ken et al., 2007; Vicentini et al., 2008; Grass Phylogeny Working Group II, 2011; Peterson et al., 2011; Soreng et al., 2017; Saarela et al., 2018). Among the six subfamilies in the PACMAD clade, Korean Poaceous taxa were classified into five subfamilies (excluding Aristidoideae) and 11 tribes (Fig. 3). The phylogenetic relationships and taxonomic considerations within each subfamily are described below.

Arundinoideae: All Korean accessions representing *Phragmites* (*P. australis*, *P. japonicus*) and *Molinia* (*M. japonica*) were nested in a highly robust subtribe, Molinieae (Fig. 3). The members of Molinieae shared a noteworthy 98 bp indel in *atpF-H*, possibly serving as a useful molecular marker defining the subtribe. The close relationship between the two genera is confirmed in the nuclear ribosomal internal transcribed spacer (nrITS) phylogeny based on a single accession for each genus (Hsiao et al., 1998). Two Korean accessions of *Phragmites* formed a clade with *P. mauritanus*. However, *M. japonica* formed a monophyletic group with *Phragmites* spp., and not with *M. caerulea*, making *Molinia* possibly paraphyletic. Additional taxon sampling and data will be necessary to clarify the taxonomic circumscription and delimitation of the two genera further.

Micraioideae: This subfamily includes three Korean species representing *Coelachne* and *Isachne* included in the subtribe Isachneae. Of the genera, monophyly of *Isachne* was evident as the two Korean accessions were closely allied with *I. distichophylla* (Fig. 3). It should be emphasized that these two genera have previously been included in Panicoideae (Clayton and Renvoize, 1986; Watson et al., 1992) and Chloridoideae as well (for example *C. japonica*) (Lee, 2007). The present study suggests that the Korean species should be included in Micraioideae.

Danthoioideae: There is no Korean native species included in this subfamily. The newly examined accession representing the subfamily is *Cortaderia selloana*, which has been introduced from South America for landscaping. There is another introduced species *Danthonia spicata*, which distributes in the middle part of Korea (Lee, 2016), but is not included in this study. Reimer (2006) studied the interspecific relationship of *Danthonia* using the *trnL-F*, confirming that *D. spicata* showed the closest relationship to *D. parryi* and *D. unispicata*.

Chloridoideae: The Korean accessions representing seven genera were classified into three tribes (Cynodonteae, Eragrostideae, Zoysieae) and five subtribes (Eleusininae, Eragrostidinae, Muhlenbergiinae, Sporobolinae, Zoysiinae) in Chloridoideae (Fig. 3). Among the seven genera, *Eragrostis* did not form a monophyletic group. Some of the species of *Eragrostis* showed a close phylogenetic relationship with *Pogonarthria* and *Ectrosia*, making *Eragrostis* polyphyletic. Similar results have been obtained from phylogenetic studies using the sequences of nrITS and *trnL-F* (Columbus et al., 2007) and *waxy* and *rps16* (Ingram and Doyle, 2004). Additional studies including samples of *E. multicaulis*, *E. ferruginea*, and *E. japonica* will clarify the taxonomic circumscription of the genera.





Fig. 3. Continued.

Panicoideae: The taxa previously classified as ‘Centothecoideae’ were included in Panicoideae (Fig. 3). These taxa had been classified as Arundinoideae based on morphological and anatomical investigations (Renvoize, 1981; Hilu and Wright, 1982; Davis and Soreng, 1993; Hsiao et al., 1998). Grass Phylogeny Working Group (2001) and Bouchenak-Khelladi et al. (2008) argued that ‘Centothecoideae’ was a monophyletic group which could be classified as a distinct subfamily, but several molecular phylogenetic studies have shown that the members of ‘Centothecoideae’ have closer relationships with Panicoideae (Hsiao et al., 1999; Zhang, 2000; Grass Phylogeny Working Group, 2001; Salamin et al., 2002; Duvall et al., 2007; Sánchez-Ken and Clark, 2007; Bouchenak-Khelladi et al., 2008, 2010). Sánchez-Ken and Clark (2007, 2010) and Grass Phylogeny Working Group II (2011) have also argued that the separation of ‘Centothecoideae’ as an independent subfamily was not appropriate, a contention supported by the present study. It is thought that the monophyly of ‘Centothecoideae’ observed in previous studies (Grass Phylogeny Working Group, 2001; Bouchenak-Khelladi et al., 2008) is due to the limited taxon sampling in those studies.

The Korean accessions (representatives of a total of 28 genera) included in this subfamily were nested in five tribes (Zeugiteae, Paniceae, Paspaleae, Arundinelleae, and Andropogoneae) (Fig. 3). Except for the accessions of three monophyletic genera (*Lophatherum*, *Paspalum*, and *Arundinella*) belonging to the relatively small tribes Zeugiteae, Paspaleae, and Arundinelleae, the remaining Korean accessions were divided into two large tribes, Paniceae (9 genera) and Andropogoneae (16 genera). Monophyly of many genera within these two large tribes is not evident, as discussed below.

Within the subtribe Cenchrinae (tribe Paniceae), *Pennisetum* and *Setaria* were found to be polyphyletic, as some members of each genus were grouped with species of other genera. Chemisquy et al. (2010) suggested merging *Pennisetum* into *Cenchrus* based on the results of morphological and molecular phylogenetic analyses. Meanwhile, *P. glaucum*, which showed a sister relationship with *S. pumilia*, was sometimes recognized as *S. glauca* (Hus, 1978; Lee, 2007). It is apparent from the molecular phylogeny that *P. glaucum* is distinct from other species in the genus, but the taxonomic treatment must be verified as the generic circumscriptions of related taxa have not yet been fully resolved. For example, *Setaria* also appears to be polyphyletic (Fig. 3). Polyphyly of the genus has also been suggested in molecular phylogenetic analyses and inflorescence differentiation pattern studies (Doust and Kellogg, 2002; Doust et al., 2007;

Kellogg et al., 2009). Korean *Setaria* species are divided into two groups; the first is an annual, with highly condensed panicle and spikelets with bristles (*S. chondrachne*, *S. pycnocomma*, *S. verticillata*, *S. italica*, *S. viridis*); the second group is an annual or perennial, with loose panicles, narrow spikelets, and wrinkled leaves (*S. pumila*). This is consistent with the results reported by Kellogg et al. (2009).

Eriochloa in the subtribe Melinidinae is paraphyletic, and the two representative species of the genus form a clade with *Urochloa panicoides* (Fig. 3). These two genera can be distinguished by the shape of the callus, i.e., the base of the spikelets; *Eriochloa* has a swollen cup shape, whereas *Urochloa* does not (Salaria et al., 2009). However, the other morphological features are very similar. Molecular phylogenetic studies using nuclear ITS sequences also did not support monophyly in *Eriochloa* (Torres Gonzalez and Morton, 2005). Because only one species (*E. villosa*) is currently distributed in Korea, a phylogenetic analysis including expanded taxon sampling from the genus as well as closely related genera would be necessary to address the taxonomic issue.

Two Korean accessions of *Digitaria* in the subtribe Anthephorinae are divided into two different clades: *D. sanguinalis* shows a sister group relationship with the “*Anthephora elongata* + *Chaetopoa pilosa*” clade and *D. violascens* is allied with *Megaloprotachne albescens* (Fig. 3). It is interesting to note that a close relationship between these genera and *Digitaria* has not yet been proposed. The two Korean species can be distinguished by the surfaces of the spikelets (smooth vs. the presence of protrusions) (Vega et al., 2009). Given that the phylogenetic positions of the two species are very different from each other, a more comprehensive study should be conducted to clarify the generic identity of *Digitaria*.

The Korean accessions classified as the subtribe Panicinae were *Panicum bisulcatum*, *P. acuminatum*, and *P. dichotomiflorum*, which were found to be polyphyletic (Fig. 3). To date, many taxa within *Panicum* have been transferred to other genera or reclassified into new genera based on morphological traits and molecular phylogenetic studies (Aliscioni et al., 2003; Bess et al., 2006; Morrone et al., 2007, 2008, 2012; Sede et al., 2008, 2009; Scataglini and Zuloaga, 2013). The three Korean taxa included in *Panicum* were placed at different positions in the phylogenetic tree. However, evident morphological characters to distinguish them have not yet been found. Washburn et al. (2015) reported that *P. bisulcatum* and *P. acuminatum* are C₃ plants, while *P. dichotomiflorum* will perform C₄ photosynthesis according to a comparative analysis of photosynthesis types.

It appears that more detailed morphological, physiological, and molecular phylogenetic investigations including *Panicum* and all relevant genera are required.

The Korean taxa classified as tribe Andropogoneae were divided into 16 genera, and their subtribal classification and interspecies relationship were found to be very complicated. Only two subtribes (Arthraxoninae and Coicinae) were monophyletic, while all other subtribes (Ischaeminae, Dimeriinae, Rottboelliinae, Saccharinae, Anthistiriinae, and Sorghinae) were found to be either paraphyletic or polyphyletic (Fig. 3). Most of the Korean taxa belonging to this tribe are also polyphyletic. This result is consistent with the findings of a phylogenetic study using ITS and *trnL-F* (Skendzic et al., 2007). Andropogoneae is known to diversify in a very short period, causing difficulty in establishing a stable classification system. Several studies have attempted to elucidate the phylogeny of this tribe using various data, but no meaningful progress has been made thus far. This is presumably because these taxa have undergone severe evolutionary radiation and frequent hybridization (Kellogg, 2000; Mathews et al., 2002; Skendzic et al., 2007; Doyle et al., 2008; Flagel and Wendel, 2010; Teerawatananon et al., 2011). To elucidate their relationship in more detail, it is thought that a large amount of genome-wide SNP data as well as intensive research on the major morphological traits that can be used to classify each subtribe are needed.

ORCID: Jung-Hoon LEE <https://orcid.org/0000-0001-6760-7913>; Ki-Joong KIM <https://orcid.org/0000-0002-0160-2180>; Bo-Yun Kim <https://orcid.org/0000-0002-1269-4068>; Young-Dong KIM <https://orcid.org/0000-0001-5904-4727>

Appendix 1 is available at <https://www.e-kjpt.org> as on-line supplemental data.

ACKNOWLEDGMENTS

This study was supported by a grant from the National Institute of Biological Resources of the Republic of Korea (NIBR202212101).

CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest.

LITERATURE CITED

Akaike, H. 1974. A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19: 716–723.

- Aliscioni, S. S., L. M. Giussani, F. O. Zuloaga and E. A. Kellogg. 2003. A molecular phylogeny of *Panicum* (Poaceae: Paniceae): Tests of monophyly and phylogenetic placement within the Panicoideae. *American Journal of Botany* 90: 796–821.
- Barker, N. P., H. P. Linder and E. H. Harley. 1995. Polyphyly of Arundinoideae (Poaceae): Evidence from *rbcL* sequence data. *Systematic Botany* 20: 423–435.
- Barkworth, M. E. 2007. Stipeae Dumort. *In* *Flora of North America, North of Mexico, Vol. 24, Magnoliophyta: Commelinidae (in part): Poaceae, Part 1.* Barkworth, M. E., K. M. Capels, S. Long, L. K. Anderton and M. B. Piep (eds.), Oxford University Press, New York. Pp. 109–186.
- Bess, E. C., A. N. Doust, G. Davidse and E. A. Kellogg. 2006. Zuloagaea, a new genus of neotropical grass within the “Bristle Clade” (Poaceae: Paniceae). *Systematic Botany* 31: 656–670.
- Bouchenak-Khelladi, Y., N. Salamin, V. Savolainen, F. Forest, M. van der Bank, M. W. Chase and T. R. Hodkinson. 2008. Large multi-gene phylogenetic trees of the grasses (Poaceae): Progress towards complete tribal and generic level sampling. *Molecular Phylogenetics and Evolution* 47: 488–505.
- Bouchenak-Khelladi, Y., G. A. Verboom, V. Savolainen and T. R. Hodkinson. 2010. Biogeography of the grasses (Poaceae): A phylogenetic approach to reveal evolutionary history in geographical space and geological time. *Botanical Journal of the Linnean Society* 162: 543–557.
- Brown, R. 1810. *Prodromus Florae Novae Hollandiae et Insulae Van-Diemen.* Vol. 1. J. Johnson, London, 590 pp. (in Latin)
- Caro, J. A. 1982. *Sinopsis taxonomica de las gramineas argentinas.* *Dominguezia* 4: 1–51. (in Spanish)
- Charmet, G., C. Ravel and F. Balfourier. 1997. Phylogenetic analysis in the *Festuca-Lolium* complex using molecular markers and ITS rDNA. *Theoretical and Applied Genetics* 94: 1038–1046.
- Chemisquy, M. A., L. M. Giussani, M. A. Scataglini, E. A. Kellogg and O. Morrone. 2010. Phylogenetic studies favour the unification of *Pennisetum*, *Cenchrus* and *Odontelytrum* (Poaceae): A combined nuclear, plastid and morphological analysis, and nomenclatural combinations in *Cenchrus*. *Annals of Botany* 106: 107–130.
- Cheon, S.-H., M.-A. Woo, S. Jo, Y.-K. Kim and K.-J. Kim. 2021. The chloroplast phylogenomics and systematics of *Zoysia* (Poaceae). *Plants* 10: 1517.
- Clark, L. G., M. Kobayashi, S. Mathews, R. E. Spangler and E. A. Kellogg. 2000. The Puelioideae, a new subfamily of Poaceae. *Systematic Botany* 25: 181–187.
- Clark, L. G., W. Zhang and J. F. Wendel. 1995. A phylogeny of the grass family (Poaceae) based on *ndhF* sequence data. *Systematic Botany* 20: 436–460.

- Clayton, W. D. and S. A. Renvoize. 1986. Genera Graminum: Grasses of the World. Her Majesty's Stationery Office, London, 389 pp.
- Columbus, J. T., R. Cerros-Tlatilpa, M. S. Kinney, M. E. Siqueiros-Delgado, H. L. Bell, M. P. Griffith and N. F. Refulio-Rodriguez. 2007. Phylogenetics of Chloridoideae (Gramineae): A preliminary study based on nuclear ribosomal internal transcribed spacer and chloroplast *trnL-F* sequences. *Aliso: A Journal of Systematic and Evolutionary Botany* 23: 565–579.
- Cotton, J. L., W. P. Wysocki, L. G. Clark, S. A. Kelchner, J. C. Pires, P. P. Edger, D. Mayfield-Jones and M. R. Duvall. 2015. Resolving deep relationships of PACMAD grasses: A phylogenomic approach. *BMC Plant Biology* 15: 178.
- Cuénoud, P., V. Savolainen, L. W. Chatrou, M. Powell, R. J. Grayer and M. W. Chase. 2002. Molecular phylogenetics of Caryophyllales based on nuclear 18S rDNA and plastid *rbcL*, *atpB*, and *matK* DNA sequences. *American Journal of Botany* 89: 132–144.
- Cummings, M. P., L. M. King and E. A. Kellogg. 1994. Slipped-strand mispairing in a plastid gene: *rpoC2* in grasses (Poaceae). *Molecular Biology and Evolution* 11: 1–8.
- Darbyshire, S. J. and S. I. Warwick. 1992. Phylogeny of North American *Festuca* (Poaceae) and related genera using chloroplast DNA restriction site variation. *Canadian Journal of Botany* 70: 2415–2429.
- Darriba, D., G. L. Taboada, R. Doallo and D. Posada. 2012. jModelTest 2: More models, new heuristics and parallel computing. *Nature Methods* 9: 772.
- Davis, J. I. and R. J. Soreng. 1993. Phylogenetic structure in the grass family (Poaceae) as inferred from chloroplast DNA restriction site variation. *American Journal of Botany* 80: 1444–1454.
- Doebley, J., M. Durbin, E. M. Golenberg, M. T. Clegg and D. P. Ma. 1990. Evolutionary analysis of the large subunit of carboxylase (*rbcL*) nucleotide sequence data among the grasses (Poaceae). *Evolution* 44: 1097–1108.
- Doust, A. N. and E. A. Kellogg. 2002. Inflorescence diversification in the panicoid “bristle grass” clade (Paniceae, Poaceae): Evidence from molecular phylogenies and developmental morphology. *American Journal of Botany* 89: 1203–1222.
- Doust, A. N., A. M. Penly, S. W. L. Jacobs and E. A. Kellogg. 2007. Congruence, conflict, and polyploidization shown by nuclear and chloroplast markers in the monophyletic “bristle clade” (Paniceae, Panicoideae, Poaceae). *Systematic Botany* 32: 531–544.
- Doyle, J. J. and J. L. Doyle. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemical Bulletin* 19: 11–15.
- Doyle, J. J., L. E. Flagel, A. H. Paterson, R. A., Rapp, D. E. Soltis, P. S. Soltis and J. F. Wendel. 2008. Evolutionary genetics of genome merger and doubling in plants. *Annual Review of Genetics* 42: 443–461.
- Dunning, L. T., J. K. Olofsson, C. Parisod, R. R. Choudhury, J. J. Moreno-Villena, Y. Yang, J. Dionora, W. P. Quick, M. Park, J. L. Bennetzen, G. Besnard, P. Nosil, C. P. Osborne and P.-A. Christin. 2019. Lateral transfers of large DNA fragments spread functional genes among grasses. *Proceedings of the National Academy of Sciences of the United States of America* 116: 4416–4425.
- Duvall, M. R., J. I. Davis, L. G. Clark, J. D. Noll, D. H. Goldman and J. G. Sánchez-Ken. 2007. Phylogeny of the grasses (Poaceae) revisited. *Aliso: A Journal of Systematic and Evolutionary Botany* 23: 237–247.
- Edgar, R. C. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* 32: 1792–1797.
- Edwards, E. J. and S. A. Smith. 2010. Phylogenetic analyses reveal the shady history of C_4 grasses. *Proceedings of the National Academy of Sciences of the United States of America* 107: 2532–2537.
- Farris, J. S., M. Källersjö, A. G. Kluge and C. Bult. 1994. Testing significance of incongruence. *Cladistics* 10: 315–319.
- Fay, M. F., K. M. Cameron, G. T. Prance, M. D. Lledó and M. W. Chase. 1997. Familial relationships of *Rhabdodendron* (Rhabdodendraceae): Plastid *rbcL* sequences indicate a caryophyllid placement. *Kew Bulletin* 52: 923–932.
- Finot, V. L., C. M. Baeza and O. Matthei. 2006. Epidermis micro-morphology of the lemma in *Trisetum* and related genera (Poaceae, Pooide). *Darwiniana* 44: 32–57. (in Spanish)
- Fisher, A. E., K. M. Hasenstab, H. L. Bell, E. Blaine, A. L. Ingram and J. T. Columbus. 2016. Evolutionary history of chloroid grasses estimated from 122 nuclear loci. *Molecular Phylogenetics and Evolution* 105: 1–14.
- Flagel, L. E. and J. F. Wendel. 2010. Evolutionary rate variation, genomic dominance and duplicate gene expression evolution during allotetraploid cotton speciation. *New Phytologist* 186: 184–193.
- Gaut, B. S. 2002. Evolutionary dynamics of grass genomes. *New Phytologist* 154: 15–28.
- Gaut, B. S., L. P. Tredway, C. Kubik, R. L. Gaut and W. Meyer. 2000. Phylogenetic relationships and genetic diversity among members of the *Festuca-Lolium* complex (Poaceae) based on ITS sequence data. *Plant Systematics and Evolution* 224: 33–53.
- Grass Phylogeny Working Group. 2001. Phylogeny and subfamilial classification of the grasses (Poaceae). *Annals of the Missouri Botanical Garden* 88: 373–457.

- Grass Phylogeny Working Group II. 2011. New grass phylogeny resolves deep evolutionary relationships and discovers C_4 origins. *New Phytologist* 193: 304–312.
- Guindon, S. and O. Gascuel. 2003. A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Systematic Biology* 52: 696–704.
- Hamby, R. K. and E. A. Zimmer. 1988. Ribosomal RNA sequences for inferring phylogeny within the grass family (Poaceae). *Plant Systematics and Evolution* 160: 29–37.
- Hilu, K. W., L. A. Alice and H. Liang. 1999. Phylogeny of Poaceae inferred from *matK* sequences. *Annals of the Missouri Botanical Garden* 86: 835–851.
- Hilu, K. W. and K. Wright. 1982. Systematics of Gramineae: A cluster analysis study. *Taxon* 31: 9–36.
- Hsiao, C., S. W. L. Jacobs, N. P. Barker and N. J. Chatterton. 1998. A molecular phylogeny of the subfamily Arundinoideae (Poaceae) based on sequences of rDNA. *Australian Systematic Botany* 11: 41–52.
- Hsiao, C., S. Jacobs, N. J. Chatterton and K. H. Asay. 1999. A molecular phylogeny of the grass family (Poaceae) based on the sequences of nuclear ribosomal DNA (ITS). *Australian Systematic Botany* 11: 667–688.
- Huang, W., L. Zhang, J. T. Columbus, Y. Hu, Y. Zhao, L. Tang, Z. Guo, W. Chen, M. McKain, M. Bartlett, C.-H. Huang, D.-Z. Li, S. Ge and H. Ma. 2022. A well-supported nuclear phylogeny of Poaceae and implications for the evolution of C_4 photosynthesis. *Molecular Plant* 15: 755–777.
- Hus, C.-C. 1978. Gramineae (Poaceae). *In* Flora of Taiwan. Vol. V, Angiospermae. Li, H.-L., T.-S. Liu, T.-C. Huang, T. Koyama and C. E. De Vol (eds.), Epoch Publishing Co., Ltd., Taipei. Pp. 373–783.
- Ingram, A. L. and J. J. Doyle. 2004. Is *Eragrostis* (Poaceae) monophyletic? Insights from nuclear and plastid sequence data. *Systematic Botany* 29: 545–552.
- Jacques-Félix, H. 1962. Les graminées (Poaceae) d'Afrique tropicale, 1: Généralités, classification, description des genres. Institut de Recherches Agronomiques Tropicales et des Cultures Vivrières: Bulletin Scientifique, Paris, 345 pp. (in French)
- Johnson, L. A. and D. E. Soltis. 1995. Phylogenetic inference in Saxifragaceae *sensu stricto* and *Gilia* (Polemoniaceae) using *matK* sequences. *Annals of the Missouri Botanical Garden* 82: 149–175.
- Jung, S. Y. and G. Y. Chung. 2008. A taxonomic study on the spikelet morphology of Korean *Poa* L. (Poaceae). *Korean Journal of Plant Taxonomy* 38: 477–502.
- Katoh, K., K. Misawa, K.-I. Kuma and T. Miyata. 2002. MAFFT: A novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* 30: 3059–3066.
- Kearse, M., R. Moir, A. Wilson, S. Stones-Havas, M. Cheung, S. Sturrock, S. Buxton, A. Cooper, S. Markowitz, C. Duran, T. Thierer, B. Ashton, P. Meintjes and A. Drummond. 2012. Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28: 1647–1649.
- Kellogg, E. A. 2000. Molecular and morphological evolution in the Andropogoneae. *In* Grasses: Systematics and Evolution. Jacobs, S. W. L. and J. Everett (eds.), CSIRO Publishing, Collingwood. Pp. 149–158.
- Kellogg, E. A. 2015. Flowering plants, Monocots, Poaceae. *In* The Families and Genera of Vascular Plants, Vol. 13. Kubitski, K. (ed.), Springer International Publishing, Cham. Pp. 1–416.
- Kellogg, E. A., S. S. Aliscioni, O. Morrone, J. Pensiero and F. Zuloaga. 2009. A phylogeny of *Setaria* (Poaceae, Panicoideae, Paniceae) and related genera based on the chloroplast gene *ndhF*. *International Journal of Plant Sciences* 170: 117–131.
- Kellogg, E. A. and C. S. Campbell. 1987. Phylogenetic analyses of the Gramineae. *In* Grass Systematics and Evolution. Soderstrom, T. R., K. W. Hilu, C. S. Campbell and M. E. Barkworth (eds.), Smithsonian Institution Press, Washington, D.C. Pp. 310–322.
- Kunth, C. S. 1833. Enumeratio plantarum, Omnium Hucusque Cognitarum: Secundum Familias Naturales Disposita, adjectis Characteribus, Differentiis et Synonymis, Vol. 1. Sumptibus J. G. Collae, Stutgardiae et Tubingae. Pp. 1–606. (in Latin)
- Lahaye, R., M. van der Bank, D. Bogarin, J. Warner, F. Pupulin, G. Gigot, O. Maurin, S. Duthoit, T. G. Barraclough and V. Savolainen. 2008. DNA barcoding the floras of biodiversity hotspots. *Proceedings of the National Academy of Sciences of the United States of America* 105: 2923–2928.
- Lee, J. H. 2016. Diversity and molecular phylogenetic relationship of Korean Poaceae. PhD dissertation, Hallym University, Chuncheon, Korea, 220 pp. (in Korean)
- Lee, J., C.-S. Kim and I.-Y. Lee. 2014. Taxonomic review of the genus *Echinochloa* in Korea (I): Inferred from sequences of cpDNA and nrDNA. *Weed and Turfgrass Science* 3: 183–189.
- Lee, Y. N. 1964a. Taxonomic studies on the genus *Miscanthus* II. Anatomical patterns of leaves. *Botanical Magazine, Tokyo* 77: 122–130.
- Lee, Y. N. 1964b. Taxonomic studies on the genus *Miscanthus*: Relationships among the section, subsections, and species, part 1. *Journal of Japanese Botany* 39: 196–205.
- Lee, Y. N. 1964c. Taxonomic studies on the genus *Miscanthus*: Relationships among the section, subsections, and species, part 2: Enumeration of the species and varieties. *Journal of Japanese Botany* 39: 257–265.

- Lee, Y. N. 1964d. Taxonomic studies on the genus *Miscanthus*. Relationships among the sections, subsections, and species, part 3: Enumeration of the species and varieties. *Journal of Japanese Botany* 39: 289–298.
- Lee, Y. N. 2007. Poaceae Barnhart. In *The Genera of Vascular Plants of Korea*. Park, C. W. (ed.), Academy Publishing Co., Seoul. Pp. 1182–1264.
- Mathews, S. and R. A. Sharrock. 1996. The phytochrome gene family in grasses (Poaceae): A phylogeny and evidence that grasses have a subset of the loci found in dicot angiosperms. *Molecular Biology and Evolution* 13: 1141–1150.
- Mathews, S., R. E. Spangler, R. J. Mason-Gamer and E. A. Kellogg. 2002. Phylogeny of Andropogoneae inferred from phytochrome B, GBSSI, and *ndhF*. *International Journal of Plant Science* 163: 441–450.
- Mathews, S., R. C. Tsai and E. A. Kellogg. 2000. Phylogenetic structure in the grass family (Poaceae): Evidence from the nuclear gene phytochrome B. *American Journal Botany* 87: 96–107.
- Morrone, O., L. Aagesen, M. A. Scataglini, D. L. Salariato, S. S. Denham, M. A. Chemisquy, S. M. Sede, L. M. Giussani, E. A. Kellogg and F. O. Zuloaga. 2012. Phylogeny of the Paniceae (Poaceae: Panicoideae): Integrating plastid DNA sequences and morphology into a new classification. *Cladistics* 28: 333–356.
- Morrone, O., M. A. Scataglini and F. O. Zuloaga. 2007. *Cyphoanthus*, a new genus segregated from *Panicum* (Poaceae: Panicoideae: Paniceae) based on morphological, anatomical and molecular data. *Taxon* 56: 521–532.
- Nadot, S., R. Bajon and B. Lejeune. 1994. The chloroplast gene *rps4* as a tool for the study of *Poaceae* phylogeny. *Plant Systematics and Evolution* 191: 27–38.
- Oh, S.-D., S.-K. Lee, D.-W. Yun, H.-J. Sun, H.-G. Kang, H.-Y. Lee, H. Xi, J. Park and B. Lee. 2021. The complete chloroplast genome of *Zoysia macrostachya* (Poaceae): Insights into intraspecific variations and species delimitation of the *Zoysia* species. *Korean Journal of Plant Taxonomy* 51: 326–331.
- Olmstead, R. G., H. J. Michaels, K. M. Scott and J. D. Palmer. 1992. Monophyly of the Asteridae and identification of their major lineages inferred from DNA sequences of *rbcL*. *Annals of the Missouri Botanical Garden* 79: 249–265.
- Palisot de Beauvois, A.-M.-F.-J. 1812. *Essai d'une Nouvelle Agrostographie, ou, Nouveaux Genres des Graminées: Avec Figures Représentant les Caractères de Tous les Genres*. Chez l'auteur, Paris. (in French)
- Peterson, P. M., K. Romaschenko and G. Johnson. 2010. A classification of the Chloridoideae (Poaceae) based on multi-gene phylogenetic trees. *Molecular Phylogenetics and Evolution* 55: 580–598.
- Peterson, P. M., K. Romaschenko, N. P. Barker and H. P. Linder. 2011. Centropodieae and *Ellisochloa*, a new tribe and genus in Chloridoideae (Poaceae). *Taxon* 60: 1113–1122.
- Plants of the World Online. 2022. Facilitated by the Royal Botanic Gardens, Kew. Retrieved Jun. 9, 2022, available from <http://www.plantsoftheworldonline.org/>.
- Prat, H. 1960. Vers une classification naturelle des Graminées. *Bulletin de la Société Botanique de France* 107: 32–79. (in French)
- Quintanar, A., S. Castroviejo and P. Catalán. 2007. Phylogeny of the tribe Aveneae (Pooideae, Poaceae) inferred from plastid *trnT-F* and nuclear ITS sequences. *American Journal Botany* 94: 1554–1569.
- Rambaut, A. 2012. FigTree v 1.4. Retrieved Dec. 5, 2012, available from <http://tree.bio.ed.ac.uk/software/figtree/>.
- Raven, P. H. and G. B. Johnson. 1995. *Understanding Biology*. 3rd ed. William C Brown Communications, New York, 536 pp.
- Reimer, E. 2006. A phylogenetic study of *Danthonia* DC. (Poaceae) in North America. MS thesis, University of Saskatchewan, Saskatoon, SK, Canada.
- Renvoize, S. A. 1981. The sub-family Arundinoideae and its position in relation to a general classification of the Gramineae. *Kew Bulletin* 36: 85–102.
- Romaschenko, K., P. M. Peterson, R. J. Soreng, O. Futorna and A. Susanna. 2011. Phylogenetics of *Piptatherum* s.l. (Poaceae: Stipeae): Evidence for a new genus, *Piptatheropsis*, and resurrection of *Patis*. *Taxon* 60: 1703–1716.
- Romaschenko, K., P. M. Peterson, R. J. Soreng, N. Garcia-Jacas and A. Susanna. 2010. Phylogenetics of Stipeae (Poaceae: Pooideae) based on plastid and nuclear DNA sequences. In *Diversity, Phylogeny, and Evolution in Monocotyledons*. Seberg, O., G. Petersen, A. S. Barfod and J. I. Davis (eds.), Aarhus University Press, Aarhus, Denmark. Pp. 511–537.
- Ronquist, F. and J. P. Huelsenbeck. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Saarela, J. M., S. V. Burke, W. P. Wysocki, M. D. Barrett, L. G. Clark, J. M. Craine, P. M. Peterson, R. J. Soreng, M. S. Vorontsova and M. R. Duvall. 2018. A 250 plastome phylogeny of the grass family (Poaceae): Topological support under different data partitions. *PeerJ* 6: e4299.
- Saarela, J. M. and S. W. Graham. 2010. Inference of phylogenetic relationships among the subfamilies of grasses (Poaceae: Poales) using meso-scale exemplar-based sampling of the plastid genome. *Botany* 88: 65–84.
- Saarela, J. M., Q. Liu, P. M. Peterson, R. J. Soreng and B. Paszko. 2010. Phylogenetics of the grass 'Aveneae-type plastid DNA clade' (Poaceae: Pooideae, Poaceae) based on plastid and nuclear ribosomal DNA sequence data. In *Diversity, Phylogeny, and*

- Evolution in the Monocotyledons. Seberg O., G. Petersen, A. S. Barfod and J. Davis (eds.), Aarhus University Press, Aarhus. Pp. 557–586.
- Salamin, N., T. R. Hodkinson and V. Savolainen. 2002. Building supertrees: An empirical assessment using the grass family (Poaceae). *Systematic Botany* 51: 136–150.
- Salariato, D. L., L. M. Giussani, O. Morrone and F. O. Zuloaga. 2009. *Rupichloa*, a new genus segregated from *Urochloa* (Poaceae) based on morphological and molecular data. *Taxon* 58: 381–391.
- Sánchez-Ken, J. G. and L. G. Clark. 2007. Phylogenetic relationships within the Centothecoideae + Panicoideae clade (Poaceae) based on *ndhF* and *rpl16* intron sequences and structural data. *Aliso: A Journal of Systematic and Evolutionary Botany* 23: 487–502.
- Sánchez-Ken, J. G. and L. G. Clark. 2010. Phylogeny and a new tribal classification of the Panicoideae s.l. (Poaceae) based on plastid and nuclear sequence data and structural data. *American Journal Botany* 97: 1732–1748.
- Sánchez-Ken, J. G., L. G. Clark, E. A. Kellogg and E. E. Kay. 2007. Reinstatement and emendation of subfamily Micrairoideae (Poaceae). *Systematic Botany* 32: 71–80.
- Scataglioni, M. A., F. O. Zuloaga and C. A. Wilson. 2013. *Morroña*, a new genus segregated from *Panicum* (Paniceae, Poaceae) based on morphological and molecular data. *Systematic Botany* 38: 1076–1086.
- Sede, S. M., O. Morrone, L. M. Giussani and F. O. Zuloaga. 2008. Phylogenetic studies in the Paniceae (Poaceae): A realignment of section *Lorea* of *Panicum*. *Systematic Botany* 33: 284–300.
- Sede, S. M., F. O. Zuloaga and O. Morrone. 2009. Phylogenetic studies in the Paniceae (Poaceae-Panicoideae): *Ocellochloa*, a new genus from the New World. *Systematic Botany* 34: 684–692.
- Sha, L., X. Fan, R. Yang, H. Kang, C. Ding, L. Zhang, Y. Zheng and Y. Zhou. 2010. Phylogenetic relationships between *Hystrix* and its closely related genera (Triticeae; Poaceae) based on nuclear *Acc1*, *DMC1* and chloroplast *trnL-F* sequences. *Molecular Phylogenetics and Evolution* 54: 327–335.
- Shantz, H. L. 1954. The place of grasslands in the Earth's cover. *Ecology* 35: 143–145.
- Simon, B. K. 2007. Grass phylogeny and classification: Conflict of morphology and molecules. *Aliso: A Journal of Systematic and Evolutionary Botany* 23: 259–266.
- Skendzic, E. M., J. T. Columbus and R. Cerros-Tlatilpa. 2007. Phylogenetics of Andropogoneae (Poaceae: Panicoideae) based on nuclear ribosomal internal transcribed spacer and chloroplast *trnL-F* sequences. *Aliso: A Journal of Systematic and Evolutionary Botany* 23: 530–544.
- Soreng, R. J. and J. I. Davis. 1998. Phylogenetics and character evolution in the grass family (Poaceae): Simultaneous analysis of morphological and chloroplast DNA restriction site character sets. *The Botanical Review* 64: 1.
- Soreng, R. J., P. M. Peterson, K. Romaschenko, G. Davidse, J. K. Teisher, L. G. Clark, P. Barberá, L. J. Gillespie and F. O. Zuloaga. 2017. A worldwide phylogenetic classification of the Poaceae (Gramineae) II. An update and a comparison of two 2015 classifications. *Journal of Systematics and Evolution* 55: 259–290.
- Soreng, R. J., P. M. Peterson, K. Romaschenko, G. Davidse, F. O. Zuloaga, E. J. Judziewicz, T. S. Filgueiras, J. I. Davis and O. Morrone. 2015. A worldwide phylogenetic classification of the Poaceae (Gramineae). *Journal of Systematics and Evolution* 53: 117–137.
- Soreng, R. J., P. M. Peterson, F. O. Zuloaga, K. Romaschenko, L. G. Clark, J. K. Teisher, L. J. Gillespie, P. Barberá, C. A. D. Welker, E. A. Kellogg, D.-Z. Li and G. Davidse. 2022. A worldwide phylogenetic classification of the Poaceae (Gramineae) III: An update. *Journal of Systematics and Evolution* 60: 476–521.
- Stapleton, C. M. A. 2013. *Bergbambos* and *Oldeania*, new genera of African bamboos (Poaceae, Bambusoideae). *PhytoKeys* 25: 87–103.
- Stebbins, G. L. and B. Crampton. 1961. A Suggested Revision of the Grass Genera of Temperate North America. *Recent Advances in Botany*, Vol. 1. University of Toronto Press, Toronto. Pp. 133–145.
- Stevens, P. F. 2017. Angiosperm phylogeny website, version 14. Retrieved Jul. 4, 2017, available from <http://www.mobot.org/mobot/research/apweb/>.
- Swofford, D. L. 2002. PAUP: Phylogenetic analysis using parsimony (and other methods), version 4.0 beta. Sinauer Associates, Sunderland, MA.
- Taberlet, P., L. Gielly, G. Pautou and J. Bouvet. 1991. Universal primers for amplification of three non-coding regions of chloroplast DNA. *Plant Molecular Biology* 17: 1105–1109.
- Tateoka, T. 1957. Miscellaneous papers on the phylogeny of the Poaceae (10). Proposition of a new phylogenetic system of Poaceae. *Journal of Japanese Botany* 32: 275–287.
- Teerawatananon, A., S. W. L. Jacobs and T. R. Hodkinson. 2011. Phylogenetics of *Panicoideae* (Poaceae) based on chloroplast and nuclear DNA sequences. *Telopea* 13: 115–142.
- The Plant List. 2013. Version 1.1. Published on the Internet. Retrieved Sep. 1, 2013, available from <http://www.theplantlist.org/>.
- Torrecilla, P. and P. Catalán. 2002. Phylogeny of broad-leaved and fine-leaved *Festuca* lineages (Poaceae) based on nuclear ITS sequences. *Systematic Botany* 27: 241–251.

- Torrecilla, P., J.-A. López-Rodríguez and P. Catalán. 2004. Phylogenetic relationships of *Vulpia* and related genera (Poeae, Poaceae) based on analysis of ITS and *trnL-F* sequences. *Annals of the Missouri Botanical Garden* 91: 124–158.
- Torres Gonzalez, A. M. and C. M. Morton. 2005. Molecular and morphological phylogenetic analysis of *Brachiaria* and *Urochloa* (Poaceae). *Molecular Phylogenetics and Evolution* 37: 36–44.
- Triple, J. K. and L. G. Clark. 2010. Phylogeny of the temperate bamboos (Poaceae: Bambusoideae: Bambuseae) with an emphasis on *Arundinaria* and allies. *Systematic Botany* 35: 102–120.
- Tropicos. 2022. Botanical information system at the Missouri Botanical Garden. Retrieved Jun. 9, 2022, available from <https://www.tropicos.org/>.
- Tzvelev, N. N. 1989. The system of grasses (Poaceae) and their evolution. *The Botanical Review* 55: 141–204.
- Véga, A. S., G. H. Rua, L. T. Fabbri and Z. E. R. de Agrasar. 2009. A morphology-based cladistic analysis of *Digitaria* (Poaceae, Panicoideae, Paniceae). *Systematic Botany* 34: 312–323.
- Vicentini, A., J. C. Barberb, S. S. Aliscioni, L. M. Giussani and E. A. Kellogg. 2008. The age of the grasses and clusters of origins of C₄ photosynthesis. *Global Change Biology* 14: 2963–2977.
- Washburn, J. D., J. C. Schnable, G. Davidse and J. C. Pires. 2015. Phylogeny and photosynthesis of the grass tribe Paniceae. *American Journal Botany* 102: 1493–1505.
- Watson, L. and M. J. Dallwitz. 1992. *The Grass Genera of the World*. CAB International, Wallingford, 1038 pp.
- Watson, L., T. D. Macfarlane and M. J. Dallwitz. 1992. *The grass genera of the world: Descriptions, illustrations, identification, and information retrieval; including synonyms, morphology, anatomy, physiology, phytochemistry, cytology, classification, pathogens, world and local distribution, and references*. Retrieved May 20, 2022, available from <http://www.delta-int-key.com>.
- Wu, Z.-Q. and S. Ge. 2012. The phylogeny of the BEP clade in grasses revisited: Evidence from the whole-genome sequences of chloroplasts. *Molecular Phylogenetics and Evolution* 62: 573–578.
- Zhang, W. 2000. Phylogeny of the grass family (Poaceae) from *rpl16* intron sequence data. *Molecular Phylogenetics and Evolution* 15: 135–146.