

The biology of Canadian weeds. 102. *Gaultheria shallon* Pursh.

Lauchlan Fraser¹, Roy Turkington¹, and C. P. Chanway²

¹Botany Department, University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z4; and ²Department of Forest Sciences, University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z4. Received 21 July 1992, accepted 24 March 1993.

Fraser, L., Turkington, R. and Chanway, C. P. 1993. **The biology of Canadian weeds. 102. *Gaultheria shallon* Pursh.** Can. J. Plant Sci. 73: 1233–1247. *Gaultheria shallon* Pursh., salal (Ericaceae), is a densely growing perennial evergreen shrub occurring only from the panhandle of Alaska along the entire coast of British Columbia to southern California; it is of native origin. Salal grows on a wide range of soil types and textures, and is abundant in open habitats near the coast particularly on rocky knolls and along bluffs. It is a persistent, pervasive woody perennial and is a serious competitor with coniferous species. The plant produces numerous seeds but the most significant and effective form of colonization is through vegetative spread. Several herbicides are recommended for control of the weed but it is both resistant and resilient to many herbicides. This contribution summarizes the known biological data for this species.

Key words: *Gaultheria shallon*, salal, evergreen shrub, weed biology, competition

Fraser, L., Turkington, R. et Chanway, C. P. 1993. **Biologie des mauvaises herbes du Canada. 102. *Gaultheria shallon* Pursh.** Can. J. Plant Sci. 73: 1233–1247. *Gaultheria shallon* Pursh., le salal (Ericacées), est un arbrisseau pérenne, touffu, à feuilles persistantes, que l'on ne retrouve, dans son habitat d'origine, que sur la côte du Pacifique, de la bande côtière sud (le "Manche-de-poêle") de l'Alaska à la Colombie-Britannique et jusqu'au sud de la Californie. Il pousse sur un vaste assortiment de types et de textures de sols. Il est abondant dans les habitats ouverts près de la côte, en particulier sur les buttes rocheuses et le long des escarpements. C'est une vivace ligneuse, persistante et envahissante, qui crée une grosse concurrence aux conifères. La production grainière est abondante, mais la plante se propage surtout par voie végétative. Plusieurs herbicides sont recommandés, mais la plante résiste à de nombreuses préparations. Le mémoire récapitule les données biologiques connues concernant cette espèce.

Mots clés: *Gaultheria shallon*, salal, arbrisseau à feuilles persistantes, biologie des mauvaises herbes, concurrence

1. Name

Gaultheria shallon Pursh. — **salal** (Scoggan 1978), **shallon** (Connors 1967), **lemon leaves** (Anonymous 1970); **wintergreen** genus, Ericaceae, Heath or Heather Family, (Hitchcock and Cronquist 1973), Éricacées.

2. Description and Account of Variation

(a) A sparse to densely growing perennial, evergreen shrub (Fig. 1) ranging from nearly prostrate to erect. *Gaultheria shallon* forms an extensive shallow root system with very fine roots (0.1–2.0 mm in diameter) and

rhizomatous-like structures spreading from a central point. Stem 0.5–2.5 m high, creeping to upright, smooth, but may become scored with age. Twigs green when young, pilose to hirsute, and often glandular. Mature twigs red or brown. Buds 1.0–1.5 cm long, usually green but sometimes tinged red, ovoid and acute-tipped. Leaves alternate and petiolate; persistent, leathery and shiny; ovate to ovate-elliptic, 3–10 cm × 3–5 cm, sharply serrulate; margins often finely toothed. Petioles 2–4 mm long. Flowers 5–15 in terminal and sub-terminal bracteate racemes that appear one-sided because pedicels are deflexed; racemes usually 5–12 cm long. Bracts reddish,

Can. J. Plant Sci. 73: 1233–1247 (Oct. 1993)




Fig. 1. *Gaultheria shallon* Pursh. (a) single flower (b) stamen (c) fruit (d) mature plant.

6–10 cm long and prominent. Calyx lobes triangular to lanceolate, glandular-pilose and red in color, usually 1/3 to 1/2 length of corolla, but often increasing in size with age.

Corolla white or pinkish, urn-shaped, with five short recurved lobes, glandular, and sticky, 6–11 mm. Stamens 10; filaments basally expanded; anthers with four slender

1966; Singleton 1976), mountain beaver (Banfield 1974), and small mammals (e.g. red squirrel (Dimock et al. 1974). Sheep will also eat young, tender salal leaves which has recently led to their introduction into salal-dominated areas. Salal is a very attractive plant used by landscapers and florists. Florists use the leaves as a foliage supplement (trade name "lemon leaves") for cut flower arrangements (Anonymous 1970). Salal is also grown commercially in greenhouses. In 1980 salal had an annual retail value of approximately \$2 million in BC (Hunt 1980). Salal berries are nutritious and plentiful and were known to be a staple food source for the native peoples of British Columbia (Turner 1975). The berry was prepared many ways but the main form was as a dried fruit for winter food. As well, native Indians smoked the leaves with kinnikinnick (*Arctostaphylos uva-ursi*) and used it for medicinal purposes (Gunter 1945).

4. Geographical Distribution

Gaultheria shallon occurs from the panhandle of Alaska (56°N) along the entire British Columbia coast continuing to southern California (34.5°N) (Fig. 2). All recorded populations in BC occur west of the Coast Mountains except for a single isolated population along the east shore of southern Kootenay Lake (Szczawinski 1962; Calder and Taylor 1968). *Gaultheria shallon* has




Fig. 2. The distribution of *Gaultheria shallon* in British Columbia (adapted from Szczawinski (1962), Haussler et al. (1990), and Hulien (1968). The shaded area indicates the general distribution, while the individual points represent the locations where herbarium samples were collected.

been introduced to England, but is found nowhere else in the world.

Gaultheria shallon is found mainly at lower elevations. In its southern range it can be found up to approximately 800 m (Lyons 1952), in the north salal rarely exceeds 100-200 m elevation.

5. Habitat

(a) *Climatic requirements* — Salal is restricted to a humid to perhumid coastal climate with mild temperatures, little snow, and unfrozen soils in winter. It is abundant on open habitats near the coast particularly on rocky knolls and along bluffs (Calder and Taylor 1968), where it can withstand gale force winds. Salal is sensitive to frost (Paul Bavis, personal communication). Sabhasri (1961) observed that a short period of freezing temperature in the middle of May killed nearly all germinants. Although salal can grow under dense forest canopies in 0.3% full sunlight, the vigour, growth rate and flowering of salal increases as the light intensity increases (Sabhasri 1961; Stanek et al. 1979; Koch 1983; Messier et al. 1989; Bunnell 1990; Messier 1993). However, best growth is in partial shade rather than in full sunlight (Anonymous 1970; Dimock et al. 1974; Koch 1983).

Characteristic of many trees, shrubs, and herbaceous plants salal leaf morphology differs dramatically under different light intensities. Sun leaves develop in bright light, whereas shade leaves develop under low light levels. Sun leaves characteristically have prostrate stems and small, thick blades. Shade leaves are found on taller, more erect plants with larger but thinner blades. Smith (1991) found that sun leaves had a mean specific leaf area less than $90 \text{ cm}^2 \text{ g}^{-1}$; and shade leaves were greater than $90 \text{ cm}^2 \text{ g}^{-1}$. At 1.8% full sunlight (80% canopy cover), leaves were almost three times larger in area than they were in a clearing (Messier et al. 1989). Under deep shade (0.3% full sunlight), Messier et al. (1989) reported that leaves were similar in size to those in the clearing, but only half the dry weight. Smith (1991) reported that sun leaves of salal require at

6. History

Gaultheria shallon is of native origin throughout its present range. Salal is the Coastal Indian name for the plant. When David Douglas landed in Oregon in 1825, Haskin (1934; cited in Szczawinski 1962) reported

usually grows in the Coastal Western Hemlock zone are shown (Table 1).

usually grows are *Pseudotsuga menziesii* var. Douglas-fir zone, the species with which salal occurs predominantly in the Coastal Western British Columbia (Krajina 1965), *G. shallon* — Using the biogeoclimatic classification for (c) *Communities in which the species occurs*

response.

tion of nutrients produces a definite growth grow on soils of low fertility, but the addition of nutrients produces a definite growth has demonstrated that salal can survive and climates (Anonymous 1970). Sabhasri (1961) ally on living trees in extremely humid in decaying wood, and will also grow epiphytic- often be found growing in old stumps and produces a low mat-forming habit. Salal can (Anonymous 1970). However, if the soil is poor salal often good soil and plenty of moisture (Anonymous vigorous upright growth in shaded conditions, (Anonymous 1970). Salal exhibits conditions (Anonymous 1970). Salal grows well in alkaline soil con- or peaty soil is the best medium for cultivation. Moist sandy organic soils (Klinka et al. 1979). Moist sandy but is also found on Fohsolic, Brumisolic and Podzols, often with deep mor humus forms, rocky soils. It is most commonly found on a wide range of soil types and textures including peat, glacial till, sand dunes, and shallow

(b) *Substratum* - *Gaultheria shallon* grows on in the formation of shade leaves.

Table 1. Zonal vegetation of subzones of the Coastal Western Hemlock biogeoclimatic zone (adapted from Pojar et al. (1991)). vh, very wet hypermaritime; xm, very dry maritime; dm, dry maritime; mm, moist maritime; vm, very wet maritime; ds (dry submaritime), ms (moist submaritime), and ws (wet submaritime) subzones. Percent cover classes, a — 0.1–1%, b — 2–5%, c — 6–10%, d — 11–25%, e — 26–99%

Common name	Botanical name	vh	xm	dm	mm	vm
salal	<i>Gaultheria shallon</i>	e				
amabilis fir	<i>Abies amabilis</i>	e				
vine maple	<i>Acer circinnatum</i>					
vanilla-leaf	<i>Achlys triphylla</i>					
deer fern	<i>Blechnum spicatum</i>					
yellow-cedar	<i>Chamaecyparis nootkatensis</i>					
prince's pine	<i>Chamaephila umbellata</i>					
fern-leaved goldthread	<i>Coptis asplenifolia</i>					
bunchberry	<i>Cornus canadensis</i>					
step moss	<i>Hylacomium splendens</i>					
heart-leaved twayblade	<i>Listera cordata</i>					
dull Oregon grape	<i>Mahonia nervosa</i>					
false lily-of-the-valley	<i>Maianthemum dilatatum</i>					
false azalea	<i>Menziesia ferruginea</i>					
Sitka spruce	<i>Picea sitchensis</i>					
flat moss	<i>Plagiocheilum undulatum</i>					
sword fern	<i>Polystichum munitum</i>					
Douglas fir	<i>Pseudotsuga menziesii</i>					
bracken fern	<i>Pteridium aquilinum</i>					
lanky moss	<i>Rhynidiadelphus loreus</i>					
five-leaved bramble	<i>Rubus pedatus</i>					
trailing blackberry	<i>Rubus ursinus</i>					
western redcedar	<i>Thuja plicata</i>					
three-leaved foamflower	<i>Tarella trifoliata</i>					
western hemlock	<i>Tsuga heterophylla</i>					
Alaskan blueberry	<i>Vaccinium alaskense</i>					
oval-leaved blueberry	<i>Vaccinium ovalifolium</i>					

mother shoot under sparse canopies, whereas daughter shoots were farther away from the oldest shoots under closed canopies. Salal accumulates considerable biomass in a growing season, but mainly through the expansion of rhizomes leading to the production of new shoots, not through significant increase in the height of pre-existing shoots (Sabhastri 1961).

Salal's ability to respond to a continuum of understory light conditions by producing either sun leaves or shade leaves (Messier et al. 1989; Smith 1991) is important to its survival. Under dense stands salal can persist by forming shade leaves; in clearings or in openings of otherwise dense stands, salal forms sun leaves. Another important adaptive characteristic to capture light is stem elongation which responds to high red light levels

that 'he was so impressed by the sight of salal he could scarcely see anything else. Throughout all his journey in the west, the salal was one of Douglas' favorites and he held great hopes of introducing it into England, and making of it a cultivated fruit.'

(a) *Morphology* — *Gaultheria shallon* has slow initial growth for approximately the first 2 yr, but once it is established it spreads vegetatively very rapidly by means of rhizomes (Messier 1991). The length and complexity of the rhizomes of an individual salal plant in an area dominated by salal has not been determined but Koch (1983) found new shoots up to 2 m from the parent plant. Bunnell (1990) observed a strong tendency for daughter shoots to be clumped around the

level tested ($24.24 \mu\text{mol m}^{-2} \text{s}^{-1}$) (Sabhasri 1961). Maximum growth occurs under red light (Sabhasri 1961). Bunnell (1990) reported that the cost of flowering was shared across connected shoots of a plant, showing there is physiological integration between ramets.

(d) *Phenology* — The maximal growth of salal (roots, rhizomes, shoots and leaves) is between late April and August, peaking in early June. Vegetative buds burst in early April (Sabhasri 1961). Flowering can occur anytime between March and July depending on the area (Dimock et al. 1974). In Alaska, flowering occurs between March and June (Viereck and Little 1972). Near Vancouver, BC, Pojar (1974) reported flowering between 12 June and 4 July. In Washington state, flowering begins in the 3rd week of June. Fruits ripen between August and October and remain on the stem until December (Dimock et al. 1974).

(e) *Mycorrhizae* — Largent et al. (1980) reported that *G. shallon* can be associated with three different kinds of mycorrhizae: arbutoid, ericoid, and ectomycorrhizae. Three different fungal species have been identified, one is *Oidiodendron griseum* Robak and the taxonomic position of the other two species are being studied (Xiao and Berch 1992). This is the first report of *O. griseum* as an ericoid mycorrhizal fungus of *Gaultheria*.

8. Reproduction

(a) *Floral biology* — *Gaultheria shallon* flowers are pollinated by insects, primarily bumblebees and flies (Pojar 1974). The fruit is a many-seeded capsule, and each inflorescence has 5–15 capsules. However, Bunnell (1990) found that salal will flower only when two conditions are met: on vigorous stems greater than 4 yr old, and at a mean crown completeness (a measure of forest canopy closure) less than 30%. Hebda (1982) found that *Gaultheria shallon* contributed less than 2% of the pollen and spore rain in areas dominated by salal, indicating how little salal flowers.

(Sabhasri 1961; Vales 1986) occurring in dense stands (Messier et al. 1989).

Leaves of *G. shallon* have a thin epicuticular layer that consists mostly of triterpenes, and always contains ursolic acid; this material also had traces of external flavonoid aglycones, namely galangin-3-methyl ether (Wollenweber and Kohorst 1984). Salasoo (1981, 1988) determined the patterns of alkane distribution in the epicuticular wax of *G. shallon* and detected rimuene. The wax was 28% hydrocarbon.

The pollen tetrad diameter of *Gaultheria shallon* has been calculated by Hebda (1978) from a sample size of 100 tetrads collected in Burns Bog, Delta, BC. The mean tetrad diameter is $46.97 \mu\text{m}$, with a range of $41\text{--}53 \mu\text{m}$ and a standard deviation of $2.50 \mu\text{m}$. Hebda (1978) postulated that small tetrads ($41\text{--}44.5 \mu\text{m}$) indicate growth of salal in dry conditions, intermediate tetrads ($44.6\text{--}49.5 \mu\text{m}$) imply dry to intermediate habitats or fire, and large tetrads ($49.6\text{--}53 \mu\text{m}$) reflect wet sites.

(b) *Perennation* — Individual shoots may survive for 10–15 yr, but leaves are rarely older than 4 yr before senescence (Haussler and Coates 1986). Shoots will bear leaves only during the first few years (Koch 1983). Rhizomes have the potential to continue reproducing vegetatively if there is sufficient light through the forest overstory (gaps and sunflecks) for shoot growth. Bunnell (1990) found extensive mats of salal rhizomes in conifer stands between 80 and 110 yr old, even when aboveground salal densities were low. During winter, salal is virtually dormant and is very sensitive to frozen soils and frost (Sabhasri 1961).

(c) *Physiological data* — Salal is at least a moderately shade-tolerant species (Sabhasri 1961; Koch 1983) but its photosynthetic and respiration characteristics are consistent with a shade-intolerant plant species (Sabhasri 1961). For an actively growing plant it was found that at low light intensities ($6.06 \mu\text{mol m}^{-2} \text{s}^{-1}$) respiration was greater than photosynthesis. Photosynthetic activity and seedling growth significantly increased with increased light intensities up to the maximum

(b) *Seed production and dispersal* — The

fruits of salal remain on the stem until December and those seeds remaining are viable for up to 1 yr following ripening (Dimock et al. 1974). Fruits have an average of 126 seeds each. When conditions are suitable for flowering, heavy crops of fruit are produced on a regular basis (Haussler et al. 1990). Seeds are dispersed mainly by the animals which feed upon the fruit: black-tailed deer, Roosevelt elk, mountain beaver, ruffed grouse, blue grouse, and other small mammals and birds (Halverson 1986).

(c) *Viability of seeds and germination* — Seeds

can remain viable for several years in cold, dry storage, but the viability of seeds under natural conditions is generally much reduced (Dimock et al. 1974). Sabhasri (1961) stored seeds at 4.4°C for 1 yr and found a decline in germination from 31 to 21%. Seeds do not require chilling (Haussler et al. 1990) or stratification (McKeever 1938) to induce germination. Successful germination requires moist, acidic sites under partial shade (Dimock et al. 1974) and light for 8 h or more per day is essential (McKeever 1938). In Washington, germination rates of 27–35% from fresh seed under lighted conditions were reported (Sabhasri 1961; Dimock et al. 1974). In British Columbia, Messier (personal communication) obtained germination rates of approximately 60% under the same conditions.

Despite the large quantity of seeds produced

and the many seeds which germinate, seedling survival is very low (Haussler et al. 1990). Sabhasri (1961) found that in Western Washington, seedling establishment is most successful in the understory of young Douglas-fir stands.

(d) *Vegetative reproduction* — The most significant and effective form of colonization is through vegetative spread, both in open habitats and in deep shade. Once salal is present on a site, further expansion is almost exclusively by vegetative means (Sabhasri 1961; Koch 1983; McGee 1988; Messier 1991) including layering and suckering from

10. Population Dynamics

The rate of increase of salal populations depends largely on the stage of succession of the area. Messier (1991) proposed a general growth model based on the work of Messier (1991), Messier and Kimmins (1991), Messier et al. (1989), Vogt et al. (1987), and Vales (1986) (Fig. 3) of the development of salal over a 60-yr period following a major disturbance (e.g. clear-cutting and slashburning) of old-growth western red cedar and western hemlock forests on northern Vancouver Island. The development of the live fine-root, leaf, stem and rhizome biomass of salal has three different stages. Stage one is between

9. Hybrids

Gaitheria shallon does not naturally hybridize with any other species (Pojar 1974). However, an artificial hybrid between *G. shallon* and *Pernettya mucronata* has been produced in England (Dimock et al. 1974).

54 naturally growing plants that Bunnell (1990) examined, no evidence of stem layering was found; rhizomes were the only means of producing new shoots. Bunnell (1990) forced stems into the organic mat and examined them 1 yr later to find that they had grown adventitious roots, and concluded that layering must also occur naturally. He found, while monitoring the colonization of salal, that 85% of the space occupied by salal after 9 yr of growth was occupied during the first 3 yr. Bunnell also observed that the vegetative reproduction of ramets was negatively associated with age ($r^2 = 0.95$) and that no new shoots were produced after an individual ramet reached 9 yr of age. Seed production may be significant in the initial colonization of a newly disturbed area (e.g. windthrow, clearcut), but considering that a plant must be at least 4 yr old before it will flower, and that most colonization of an area occurs within the first 3 yr, the vast majority of colonization occurs through vegetative spread. Bunnell (1990) found no seed production, only vegetative spread, at a mean crown completeness greater than 30%.


Bunnell et al. 1990; Messier et al. 1990; Messier 1991). The competitive influence of salal on the seedlings of Douglas-fir, western hemlock, and Sitka spruce is believed to be much stronger than on western red cedar and lodgepole pine (Bunnell et al. 1990). Competition is most severe during the early stages of stand development (Long 1977) but may continue through the rotation, in particular, if the canopy is open enough to allow a well-developed understory of salal to persist (Stanek et al. 1979).

Young stands of Douglas-fir on sub-xeric sites are limited by soil water potential (Tan et al. 1977). At low values of soil water potential, the reduction in stomatal conductance is greater for Douglas-fir than for salal, indicating that salal transpiration would account for a higher proportion of total stand transpiration (Tan et al. 1978). Price et al. (1986), found that the removal of the salal understory of thinned 32-yr-old Douglas-fir significantly increased rates of photosynthesis

1 and 15 yr and is characterized by a rapid growth rate of rhizomes and fine-roots, in particular, between 8 and 15 yr. Stage two, approximately between 16 and 45 yr, begins with a rapid decline in salal live fine-root biomass followed by a more gradual decrease in leaf and stem biomass. The decline of the population is caused by shading due to an increase in the overstory tree canopy. There are no data for rhizome biomass in the 10- to 60-yr period, but it is believed to be high (Messier 1991). The final stage, occurring from 46 yr until the next major disturbance, is the virtual exclusion of salal by the dense overstory canopy. However, it is not completely excluded and can quickly re-establish following a disturbance.

Much research effort has been focused on the competitive interactions of salal and coniferous tree species in low-elevation coastal British Columbia forests (Tan et al. 1978; Black et al. 1980; Price et al. 1986; Weetman et al. 1989a; Klinka et al. 1989;

Fig. 3. Hypothetical development of fine-root, leaf, stem and rhizome abundance of *Gaultheria shallon* over a 60-yr period following the clear-cutting and burning of old-growth forests of western redcedar and western hemlock on northern Vancouver Island (taken from Messier (1991)).



11. Response to Herbicides and Other

Chemicals

In general salal is both resistant and resilient to many herbicides. The most successful herbicide in controlling salal is triclopyr ester

(Garlon). Applying triclopyr ester at 4 kg a.i. ha⁻¹ reduced salal cover by 78% (Barker 1988). Combining diesel with triclopyr ester effectively controlled salal for three seasons in a Douglas-fir salal ecosystem on the south-east coast of Vancouver Island (Dunsworth 1986). In an experiment conducted in a dry cedar-hemlock ecosystem, 4 kg a.i. ha⁻¹ in triclopyr ester in diesel oil at 100 L ha⁻¹ in early spring or late summer reduced salal cover by 60-90%. When using mineral oil as the carrier instead of diesel oil, salal cover was reduced by only 40% (Haussler et al. 1990).

Stewart (1974b) tested 10 herbicides mixed in diesel-oil carriers. Salal plants were sprayed on 1 April, when flower buds were swelling and vegetative buds were dormant. Silvex, dichlorprop, 2,4,5-T, and dicamba produced at least 80% topkill and a 50% reduction in salal cover with silvex showing the best control. Salal is resistant to these same herbicides when applied in water, or applied later in the growing season (Gratkowski 1970; Stewart 1974a).

Other herbicides which are less effective but have been reported to cause slight damage to salal are glyphosate and 2,4-D (Conard and Emmingham 1984). Results from hexazinone applications are variable; Newton and Knight (1981) and Boeteng and Herring (1990) found it to be ineffective at controlling salal. However, Wellman and Harrison (1987) reported that an April ground application of 2 kg a.i. ha⁻¹ granular hexazinone resulted in control of salal. Two years after herbicide application, salal biomass on the treated areas was about half that on the untreated areas.

In all of the herbicidal tests to control salal, little is known about how the below-ground plant portion responds. D'Anjou (cited by Haussler et al. (1990)) showed that although aboveground parts of salal were well controlled by triclopyr ester, living roots (dry

and tree growth due to an increase in the soil water potential. Other researchers have proposed that salal may be competing for nutrients in the Douglas-fir forests (Stanek et al. 1979; Haussler and Coates 1986).

There is evidence to suggest that salal may also be competitive in moist ecosystems (Weetman et al. 1989b; Messier 1991). Two hypotheses have been proposed to account for this. First, is the resource exploitation hypothesis. Weetman et al. (1989a,b) have shown that on northern Vancouver Island western red cedar and western hemlock cutovers that are dominated with salal are nutrient deficient. Germain (1985) and Messier (1991) found that salal can have an impact on the nutrient budget for conifers. *Gaultheria shallon* forms three symbiotic fungal associations: ericoid-, arbutoid- and ecto-mycorrhizae (Largent et al. 1980), whereas western red cedar and western hemlock only form one fungal association each: ectomycorrhizae and ectomycorrhizae, respectively. Therefore, *G. shallon* might be more efficient at extracting nutrients such as nitrogen and phosphorus at low pH, and accessing nutrients in complex organic forms (Xiao, Cade-Menun and Berch, personal communication). Parke et al. (1983) suggest that dense salal can lower soil temperatures causing reduced conifer growth by inhibiting root growth and mycorrhizal infection. The second hypothesis involves the possible allelopathic properties of salal. Del Moral and Cates (1971) did not find convincing evidence for allelopathy in salal, but Rose et al. (1983) have shown that allelochemicals in salal litter may inhibit seedling growth. It has been suggested that salal may have an allelopathic effect through the production of tannins and phenolic acids (de Montigny, personal communication).

Gaultheria shallon can be classified as a stress-tolerator with a low R (the level of resource below which the population is unable to maintain itself) (Messier 1991). Evidence that salal has a low R is its ability to maintain a higher stomatal conductance than Douglas-fir under a low soil water potential (Tan et al. 1978).

Table 2. The percentage of total number of leaves infected by commonly observed fungal taxa on 1240 leaf discs of *Gaultheria shallon* Pursh. (adapted from Petrini et al. (1980)).

<i>Acronium</i> sp.	2.4
<i>Genticulosporium</i> sp.	8.4
<i>Leptothyrium berberidis</i> Cooke et Massee	3.6
<i>Nodulisporium</i> sp.	2.4
<i>Pezizula</i> sp.	6.4
<i>Phialophora</i> spp.	7.2
<i>Phomopsis</i> sp.	4.8
<i>Phyllosticta pyrolae</i> Ellis et Everh.	34.7
<i>Phyllosticta vaccinii</i> Earle	5.6
<i>Ramularia</i> sp.	4.8
<i>Septogloeum</i> sp.	4.4
<i>Sigmoida</i> sp.	1.2
<i>Xylaria hypoxylon</i> (L. ex Fr.) Grev. anamorph	2.0

southern Vancouver Island, B. Green (personal communication) has speculated that heavy soil scarification (i.e. removing the organic layer) on dry sites causes mortality of salal rhizomes due to desiccation.

13. Responses to Parasites

Salal is infected by numerous fungal parasites. The most common and serious is the leaf spot fungus (*Mycosphaerella gaultheriae*) (Conners 1967; Haussler et al. 1990). In a study conducted in western Oregon, Petrini et al. (1982) isolated 13 different species of endophytic fungi (Table 2). The most frequent was *Phyllosticta pyrolae* Ellis et Everh. which occurred on 34.7% of the observed leaves. They found that the frequency of endophyte infections diminishes with decreasing habitat moisture. Furthermore, rates of overall infection were higher in samples taken from densely wooded sites than in samples taken from more open sites. Other fungi reported on salal are *Asterella gaultheriae*, *Bulgaria melastoma*, *Lachnella gaultheriae*, *Lepidosphaeria gaultheriae*, *Phacidium gaultheriae*, *Sphaeria gaultheriae*, *Phacidium gaultheriae*, *Phyllosticta gaultheriae*, and *Poria ferruginea* (Conners 1967). Linderman and Zeitoun (1977) found *Phytophthora cinnamomi* Rands, a very widespread fungus, occurring on nursery-grown salal. Apothecia of *Valdensia heterodoxa* Peyr. (Sclerotiniaceae), have been found on fallen *G. shallon* leaves

weight basis) still comprised 89% of that in untreated controls, indicating that roots continue to survive despite substantial foliar control.

12. Responses to other Human Manipulations

Several studies have reported that salal will rapidly increase in cover and vigour following the removal or reduction of the forest canopy (Sabhasri 1961; Long and Turner 1975; Long 1977; Stanek et al. 1979; Black et al. 1980; Koch 1983; Gholz et al. 1985; Price et al. 1986; Vales and Bunnell 1988; Messier et al. 1989). Messier (1991) found that salal re-establishes relatively quickly below ground following clear-cut logging and slashburning, but the aboveground portion does not grow as rapidly and may take many years to become dominant.

Prescribed burning and logging can increase the growth of salal if the burn is light. Fire stimulates resprouting from roots and stem bases (Sabhasri 1961). Only severe burns that penetrate sufficiently deep to kill the roots can reduce salal cover. Vihanek (1985) reported that on dry sites on eastern Vancouver Island high severity burns decreased salal cover by 80% compared to adjacent unburned areas; whereas low to moderate burns decreased cover by only 40%. Fertilizer application, particularly nitrogen-rich fertilizer, increases both above and below-ground growth of salal (Sabhasri 1961; Anonymous 1970). However, in forest stands, applications of fertilizer that result in an increased tree canopy density may cause a decline in the vigour and cover of the salal understory due to shading (Long and Turner 1975; Stanek et al. 1979).

Salal readily forms new plants from cuttings of the stem and roots (Sabhasri 1961), so it can be expected that any form of soil disturbance that causes mechanical damage to the plant, but that does not physically remove it from the site, will stimulate resprouting. However, it has been reported by Muller (1989, cited in Haussler et al. (1990)) that heavy scarification on areas on southern Vancouver Island has resulted in very slow reinvasion of salal. Based on trials on

- (Redhead 1974). Salal is known to have at least one higher plant parasite, *Boschniakia hookeri* Walp. (Olson and Olsen 1981; Kujal and Toth 1985), a perennial root parasite. It is not known to what degree the parasite actually damages salal.
- This work was funded by a Natural Sciences and Engineering Research Council/Industry grant to C.P.C. and R.T. as part of the SCHIRP study.
- Anonymous, 1970.** *Gaultheria shallon* Pursh. Davidsonia I: 29-31.
- Bailey, A. W. 1966.** Forest associations and secondary plant succession in the southern Oregon Coast Range. Ph.D. thesis. Oregon State University, Corvallis, OR.
- Banfield, A. W. F. 1974.** The mammals of Canada. University of Toronto Press, Toronto, ON. (1988 update). Expert Committee on Weeds.
- Barker, J. E. 1988.** Control of salal with Garlon Research report, p. 194. Western Canada Section Meeting, Winnipeg, MB.
- Barker, J. E., Weetman, G. F. and Fournier, R. M. 1987.** Growth and nutrition of sitka spruce, western hemlock and western red cedar following fertilization of coastal cedar/hemlock sites. Western Forest Products. Vancouver, BC. 14 pp.
- Black, T. A., Tan, C. S. and Niyamah, J. U. 1980.** Transpiration rate of Douglas-fir trees in thinned and unthinned stands. Can. J. Soil Sci. 60: 625-631.
- Boateng, J. O. and Herring, L. 1990.** Site preparation: Chemical. Pages 164-178 in D. P. Lavender, R. Parish, C. A. Johnson, G. Montgomery, A. Vyse, and R. Willis, eds. Regenerating British Columbia's forests. University of British Columbia Press, Vancouver, B.C.
- Brown, R. L. and Hafenrichter, A. L. 1962.** Stabilizing sand dunes on the Pacific coast with woody plants. USDA, Washington, DC. Misc. Publ. 892, 18 pp.
- Bunnell, R. L. 1990.** Reproduction of salal (*Gaultheria shallon*) under forest canopy. Can. J. For. Res. 20: 91-100.
- Bunnell, F. L., McCann, R. K., Lindop, P. and Messier, C. 1990.** Salal — a competitor? A field guide. Ministry of Forest and Lands, Victoria, BC.
- Calder, J. A. and Taylor, R. L. 1968.** Flora of the Queen Charlotte Islands. Part I. Systematics of the vascular plants. Queen's Printer, Canadian Department of Agriculture, Ottawa, ON.
- BC. Deer in British Columbia
- Chambers, M. 1988.** Deer in British Columbia BC. Min. Environ., Fish Wildl. Br., Inf. and Education Section, Victoria, BC.
- Connors, I. L. 1967.** An annotated index of plant diseases in Canada and fungi recorded on plants in Alaska, Canada and Greenland. Canadian Department of Agriculture, Ottawa, ON. Publ. 1251. 381p.
- Conard, S. G. and Emmingham, W. H. 1984.** Herbicides for brush and fern control on forest sites in western Oregon and Washington. Special Publ. 8, For. Res. Lab., Oregon State University, Corvallis, OR.
- Crouch, G. L. 1979.** Food habits of black-tailed deer on forest habitats in the Pacific Northwest. Pages 53-59 in Wallmo, O. C. and Schoen, J. W. eds. Proceedings of a Conference on Sitka Black-tailed Deer. USDA Forest Service, Alaska Region, Juneau, AK.
- Del Moral, R. and Cates, R. G. 1971.** Allopathic potential of the dominant vegetation of western Washington. Ecology 52: 1030-1037.
- Dimock II, E. J., Johnston, W. F. and Stein, W. I. 1974.** *Gaultheria* L., wintergreen. In S. C. Schopmeyer, ed. Seeds of woody plants of the United States. Agric. Hand. 450. USDA For. Ser., Washington, DC.
- Dunsworth, B. G. 1986.** Salal removal and its impact on crop tree performance. Expert Committee on Weeds. Research report, pp. 197. Western Canada Section Meeting, Saskatoon, SK.
- Germain, A. 1985.** Fertilization of stagnated Sitka spruce plantations on northern Vancouver Island. M.Sc. thesis. University of British Columbia, Vancouver, BC.
- Gholz, H. L., Hawk, G. M., Campbell, A., Cromack, J.R., K. and Brown, A. T. 1985.** Early vegetative recovery and element cycles on a clear-cut watershed in western Oregon. Can. J. For. Res. 15: 400-409.
- Gratkowski, H. 1970.** Foliage sprays fail on salal. West. Soc. Weed Sci. Res. Prog. Rep. 1970: 18.
- Gunther, E. 1945.** Ethnobotany of Western Washington. University of Washington Press, Seattle, WA.
- Haussler, S. and Coates, D. 1986.** Autecological characteristics of selected species that compete with conifers in B.C.: a literature review. Land Manage. Rep. No. 33 BC. Ministry of Forests, Victoria, BC.
- Haussler, S., Coates, D. and Mather, J. 1990.** Autecology of common plants in British Columbia: a literature review. FRDA report #158, Victoria, BC.

- Halverson, N. M. (ed). 1986. Major indicator shrubs and herbs on national forests of western Oregon and southwestern Washington. USDA For. Serv., Pacific NW Region.
- Hebda, R. J. 1978. Size, productivity, and paleoecological implications of ericaceous pollen from Burns Bog, southern Fraser River Delta, British Columbia. *Can. J. Bot.* 57: 1712-1717.
- Hebda, R. J. 1982. Late-glacial and postglacial vegetation history at Bear Cove Bog, northeast Vancouver Island, British Columbia. *Can. J. Bot.* 61: 3172-3192.
- Hitchcock, C. L. and Cronquist, A. 1973. Flora of the Pacific Northwest. University of Washington Press, Seattle, WA.
- Hult ne E. 1968. Flora of Alaska and neighboring Territories: A manual of vascular plants. Stanford University Press, Stanford, CT.
- Hunt, R. S. 1980. Rhytismataceae on salal leaves. *Mycotaxonomy* 11: 233-240.
- Jones, G. W. and Bunnell, F. L. 1984. Response of black-tailed deer to winters of different severity on northern Vancouver Island. Pages 385-396 in W. R. Meehan, T. R. Merrell, Jr., and T. A. Hanley, eds. Proceedings of a Symposium on Fish and Wildlife Relationships in Old-Growth Forests. Am. Inst. Fish. Res. Biol. Morehead City, NC.
- King, R. D. 1969. Spring and summer foods of ruffed grouse on Vancouver Island. *J. Wildl. Manage.* 33: 440-442.
- King, R. D. and Bendall, J. F. 1982. Foods selected by blue grouse (*Dendragapus obscurus fuliginosus*). *Can. J. Zool.* 60: 3268-3281.
- Klinka, K., Nuszdorfer, F. C. and Skoda, L. 1979. Biogeoclimatic units of central and southern Vancouver Island. BC Min. For., Victoria, BC.
- Klinka, K., Carter, R. E., Feller, M. C. and Wang, Q. 1989. Relations between site index, salal, plant communities, and sites in coastal Douglas-fir ecosystems. Northwest Sci. 63(1): 19-28.
- Koch, J. M. 1983. Salal plant structure, growth and allometric relationships under several forest stand densities in south coastal British Columbia. B.S.F. thesis, University of British Columbia, Vancouver, BC.
- Krajina, V. J. 1965. Biogeoclimatic zones and classification of British Columbia. *Ecol. West. N. Am.* 1: 1-17.
- Kuift, J. and Toth, R. 1985. Structure of the host-parasite interphase of *Boschniakia hookeri* Walpers (Orobanchaceae). *Acta Bot. Neerl.* 34: 257-270.
- Largent, D. L., Sugihara, N. and Wisner, C. 1980. Occurrence of mycorrhizae on ericaceous and pyrolaceous plants in northern California. *Can. J. Bot.* 58: 2274-2279.
- Linderman, R. G. and Zetoum, F. 1977. *Pytophthora cinnamomi* causing root rot and wilt of nursery-grown native western azalea and salal. Plant Dis. Rep. 61: 1045-1048.
- Long, J. N. 1977. Trends in plant species diversity associated with development in a series of *Pseudotsuga menziesii*/*Gaultheria shallon* stands. Northwest Sci. 51: 119-130.
- Long, J. N. and Turner, J. 1975. Aboveground biomass of understory and overstory in an age sequence of four Douglas-fir stands. *J. Appl. Ecol.* 12: 179-188.
- Lyons, C. P. 1952. Trees, shrubs and flowers to know in British Columbia. J. M. Dent and Sons Ltd. Vancouver, BC.
- Martin, A. C., ZIM, H. S. and Nelson, A. L. 1951. American wildlife and plants. McGraw-Hill Inc., New York, NY.
- McGe, A. B. 1988. Effects of prescribed burning on vegetation and natural tree regeneration in mature cedar-hemlock forests in the Pacific Rim National Park. B.S.F. thesis. University of British Columbia, Vancouver, BC.
- McKeever, D. G. 1938. The effect of various methods of treatment on the germination of seeds of some plants valuable for game and erosion purposes. M.Sc. thesis, University of Idaho, Moscow, ID.
- McTaggart-Cowan, I. 1945. The ecological relationships of the food of the Columbia black-tailed deer, *Odocoileus hemionus columbianus* (Richardson), in the Coast Forest Region of southern Vancouver Island. British Columbia. *Ecol. Monogr.* 15: 110-139.
- Messier, C. 1991. Factors limiting early conifer growth in salal-dominated cutovers on northern Vancouver Island, British Columbia. Ph.D. thesis, University of British Columbia, Vancouver, BC.
- Messier, C. 1993. Effects of neutral shade and growing media on growth, biomass allocation, and competitive ability of *Gaultheria shallon*. *Can. J. Bot.* 70: 2271-2276.
- Messier, C., Honer, T. W. and Kimmings, J. P. 1989. Photosynthetic photon flux density, red:far-red ratio, and minimum light requirements for survival of *Gaultheria shallon* in western red cedar/western hemlock stands in coastal British Columbia. *Can. J. For. Res.* 19: 1470-1477.
- Messier, C., Kimmings, J. P., Bunnell, F. L. and McCann, R. K. 1990. Understanding salal as a competitor. In *Vegetation management: An integrated approach*. Proc. of the fourth annual BC Veg. Manag. Workshop. Victoria, BC.

- Vihank, R. E. 1985. The effects of slashburning on the growth and nutrition of young Douglas-fir plantations in some dry, salal dominated ecosystems. M.Sc. thesis, University of British Columbia, Vancouver, BC.
- Vogl, K. A., Vogt, D. J., Moore, E. E., Fatuga, B. A., Redlin, M. R., and Edmons, R. L. 1987. Conifer and angiosperm fine-root biomass in relation to stand age and site productivity in Douglas-fir forests. *J. Ecol.* 75: 857-870.
- Weetman, G. F., Fournier, R., Barker, J., Schnorbus-Panozzo, E., and Germain, A. 1989a. Foliar analysis and response of fertilized chlorotic Sitka spruce plantations on salal-dominated cedar-hemlock cutovers on Vancouver Island. *Can. For. Res.* 12: 1501-1511.
- Weetman, G. F., Fournier, R., Barker, J., and Schnorbus-Panozzo, E. 1989b. Foliar analysis and response of fertilized chlorotic western hemlock and western red cedar reproduction on salal-dominated cedar-hemlock cutovers on Vancouver Island. *Can. For. Res.* 12: 1512-1520.
- Wetman, G. F., Fournier, R. M., Schnorbus-Panozzo, E., and Barker, J. 1990. Post-burn nitrogen and phosphorous availability in coastal British Columbia cedar/hemlock forests and the use of fertilization and salal eradication to restore productivity. *In* S. P. Gessel, ed. Sustained productivity of forest lands. Proc. 7th North Amer. For. Soils Conf. Friesen Press, Vancouver, BC.
- Wellman, R. and Harrison, C. 1987. Granular hexazinone site prep. Expert Committee on Weeds. Research report, p. 207. Western Canada Section Meeting, Victoria, BC.
- Wollenweber, E. and Kohorst, G. 1984. Novel epicuticular leaf flavonoids from *Kalmia* and *Gaultheria* (Ericaceae). *Z. Naturforsch* 39c: 710-713.
- Xiao, G. and Berch, S. M. 1992. Erioid mycorrhizal fungi of *Gaultheria shallon*. *Mycologia* 84(3): 470-471.
- Zwickel, F. C., and Bendell, J. F. 1972. Observations on food habits of incubating female blue grouse. *Condor* 74: 493-494.