


SABAH FORESTRY DEPARTMENT [Prepared by Kalabakan Forestry Office]



ANALYSIS REPORT ON THE EFFECTIVENESS OF HCV MONITORING IN NGR FMU

2019

TABLE OF CONTENTS

	TAB	LE OF (CONTENTS	PAGE							
	EXE	CUTIVE	SUMMARY	3							
1.0	OBJE	ECTIVE		4							
2.0	HCV	MANA	GEMENT AND MONITORING PLAN	5							
3.0	EFFE	ECTIVE	NESS OF MONITORING PROGRAM AND	15							
	ENH	ANCEM	IENT BY EACH HCV ATTRIBUTES								
	3.1	HCV 1	BIODIVERSITY VALUES	15							
		3.1.1	HCV 1.1 Protected Area	15							
		3.1.2	HCV 1.2 Threatened and Endangered Species	19							
		3.1.3	HCV 1.3 Endemism	29							
		3.1.4	HCV 1.4 Critical Temporal Use	33							
	3.2	HCV 2	LANDSCAPE LEVEL FOREST	34							
	3.3	HCV 3	ECOSYSTEM	35							
	3.4	HCV 4	SERVICE OF NATURE	36							
		3.4.1	HCV 4.2 Erosion Control	36							
		3.4.2 HCV 4.3 Barriers to Destructive Fire									
	3.5	3.5 HCV 6 CULTURAL IDENTITY OF LOCAL COMMUNITIES									
4.0	RESU	JLTS O	R OUTCOME FOR HCV MANAGEMENT PRESCRIPTION	41							
	AND	EFFEC	TIVENESS OF MONITORING AND ENHANCEMENT								

Executive Summary

High Conservation Value (HCV) assessment for Northern Gunung Rara Sustainable Forest Management project area was carried out through field assessment in March 2015 and also attaining secondary data from management team and previous HCV report by Wong et al 2010. The main objective of this assessment is to enhance relevant information on the HCV elements within the NGR project area. A multidisciplinary team carried out the assessment with experienced assessors from various fields. Generally, all HCV elements are found to be present were elaborated for NGR project area. Appropriate management and monitoring actions have been recommended and discussed with the management team for further actions to be undertaken. Based on the High Conservation Values in Northern Gn Rara Forest Management Unit (July 2016), the HCV's identified present in NGR Project area is as follows:

- 1. HCV 1: Biodiversity Values
 - a. HCV 1.1: Protected Area
 - b. HCV 1.2: Threatened and Endangered Species
 - c. HCV 1.3: Endemic Species
 - d. HCV 1.4: Critical Temporal Use
- 2. HCV 2: Landscape Level Forest
- 3. HCV 3: Ecosystems
- 4. HCV 4: Services to Nature
 - a. HCV 4.2: Forest Critical to Erosion Control
 - b. HCV 4.3: Forest Providing Barriers to Destructive Fire
- 5. HCV 6: Cultural Identity of Local Community

All HCV identified were monitored according to the recommendation of the High Conservation Values in Northern Gn Rara Forest Management Unit (July 2016) report done by Forest Research Center. Periodic monitoring activities were established for all identified HCV's i.e. patrolling and surveillance activities, wildlife monitoring (5 methods established), and establishments of PSP Plots for HCV 1. On a landscape level (HCV 2), NGR project area forms a critical link that connects the three larger undisturbed natural forest of protected areas, namely Danum, Imbak Canyon and Maliau Basin forest reserves to support greater landscape connectivity of lowland areas Retaining the whole FMU under conservation and natural forest management functional zones is the best effort in maintaining the forest ecosystem function as forest corridor for plant dispersal and also acts as a transient wildlife migratory path between the different forest reserves it borders. Most of southern border of NGR project area is bordering oil palm estate. Furthermore, secondary vegetation dominates most of the peripheral area of the reserves that mostly susceptible to fire in comparison to pristine forest. A 50 m band of moderate to high forest structure inside the project area that border local communities land and oil palm estate are categorised as HCV 4.3. Nine teriti or heirs from Kg Kuamut carry out the edible nest collection at VJR Batu Timbang and they have obtained the collection permits from Sabah Wildlife Department. The continuous survival of the swiftlet that produce edible nests is highly dependent of the quality of food that are found outside of the limestone caves. The protection and preservation of mixed dipterocarp forest and limestone vegetation that is indicated as HCV 6 is crucial in order to support the cultural value of some Kg Kuamut villager. Continouous consultation with the stakeholder were established and strengthened through frequent meetings.

Analysis of the effectiveness of monitoring program was done and elaboration of the analysis was explained through this report. Improvements were seen in the effectiveness of conducting more frequent patrolling and surveillance in HCV 1 monitoring program i.e increase in patrolling frequency, extensive wildlife monitoring in all methods, and also maintaining all established PSP Plots. Frequent monitoring program covering the entire Project site for ensuring the link between the three larger undisturbed natural forests of protected areas, namely Danum, Imbak Canyon and Maliau Basin were done (HCV2). Reenumeration of all PSP Plots and establishing new PSP plot covers the activities done for HCV 3. Water sampling were done in 4 locations to monitor HCV 4.2 Erosion Control, while frequent patrolling along the Forest Barrier of the 50 m band of moderate to high forest structure inside the project area that border local communities land and oil palm estate covers HCV 4.3. The Project Team also strengthened the relationship with the Birds' Nest Collectors of Bt Timbang Cave through continuous consultation and conflict resolution, which covers HCV6.

1.0 Objectives:

- 1. To evaluate the effectiveness by which HCV management and protection measures to maintain and/ or enhance the pertinent conservation attributes
- 2. As a guidance for Forest Manager to modify/ adjust/ enhance HCV management prescription to cater for any weaknesses
- 3. To comply with FSC indicator 9.4.3.
- 4. To achieve the strategic management objective as stated in the Forest Management Plan.

References:

- Annual Report 2013
- Annual Report 2014
- Annual Report 2015
- Annual Report 2016
- Annual Report 2017
- Annual Report 2018
- Annual Report 2019
- Wildlife monitoring report 2014
- Wildlife monitoring report 2015
- Wildlife monitoring report 2016
- Wildlife monitoring report 2017
- Wildlife monitoring report 2018
- Wildlife monitoring report 2019
- Quarterly wildlife monitoring 2014-2019
- List of Enforcement Activities/ Arrests/ Prosecution Cases 2013 2019
- High Conservation Values in Northern Gunung Rara Forest Management Unit: Assessment Report and Management Recommendations (July 2016)
- Annual Environmental Monitoring Program Report 2017-2019
- Quarterly Report 2017-2019

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	ANAGEMENT MANAGEMENT STRATEGIES		OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
1.1	69,432 ha Consisting of Mt Magdelena FR, Gn. Rara Wildlife Corridor FR, Batu Timbang FR, Imbok FR (Class VI) and one as production forest (Gn. Rara FR).	The whole project site remained protected from serious environmental or ecological threat.	Encroachment into the Project site is controlled and reduced. *Indicators: 1. Number of encroaches arrested per year 2. Number of Aerial survey frequency (hours) conducted per year 3. Distance of boundary re- brushing per year	Illegal entry into NGR in a reducing trend.	Whole area	 No conversion of forests is permitted. No entry without permission. No poaching. Conduct periodic patrolling and surveillance in accessible HCV areas. Demarcation of HCV boundaries on the ground for all designated HCVs within the TPAs is not required since 100 % overlaps occured among other HCV elements. No open burning within the NGR SFM as well as right nearby/ boundary of NGR SFM Signboards to prohibit any illegal activities to be erected and visibly maintained at the strategic location as well as all identified HCV area. Awareness sessions to be conducted with neighbouring logging and plantation owners regarding the protection status of 	 Periodic monitoring and control should be carried out to prevent encroachment in the FMU. Monitoring to be conducted as documented target in AWP as follow: Aerial surveillance: 2 times per year. Gate control to be conducted 16 hours (6.00 am - 10.00 pm) Inspection of boundaries to be conducted 2 times a month. Enforcement and monitoring through Forest Checking Station Batu Timbang and Imbok. Signboards to prohibit any illegal activities erected and maintained at the strategic locations as well as all identified HCV area. 	 Aerial survey report Gate control report Stakeholder Consultation meeting minutes. List of stakeholders AWP, Compliance Report and Quarterly Report Communities and stakeholders grievance report Satellite imagery monitoring report.

2.0 HCV Management and Monitoring Plan:

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	MANAGEMENT STRATEGIES		DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
					Boundary NGR SFM (particularly neighbouring	 NGR SFM, applicable laws, forest fire, biodiversity and habitats within. Cooperation with World Wildlife Fund (WWF), Sabah Wildlife Department (SWD), Monitoring, Control, Environment and Enforcement (MCEE) Tawau, DAMAI Team, Yayasan Sabah (YS), Sabah Forestry Department (SFD) in preventing encroachment and poaching No entry without permission. No illegal felling 		 Aerial survey report Gate control report
					logging and estate) I. Empayar Kejora Sdn. Bhd II. Serijaya Industries Sdn Bhd III. Maliau Basin Conservation Area IV. Imbak Canyon Conservation Area V. Danum Valley	 No megal felling No poaching No illegal cultivation's No open burning within the NGR SFM as well as right nearby/ boundary of NGR SFM Signboards to prohibit any illegal activities to be erected and visibly maintained at the strategic location as well as all identified HCV area. Awareness sessions to 		 Gate control report Stakeholder Consultation meeting minutes. List of stakeholders AWP, Compliance Report and Quarterly Report Communities and stakeholders grievance report Satellite imagery monitoring report.

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	FRATEGIES	OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
					Conservation Area	be conducted with neighbouring logging and plantation owners regarding the protection status of NGR SFM, applicable laws, forest fire, biodiversity and habitats within.		
1.2	[Listed] Flora 28 species from 4 families of plants are rare, endangered or threatened plant species	Biodiversity in NGR is maintained or enhanced	Flora diversity in NGR is maintained *Indicators: 1. Number of RTE flora species (Refer: i. Plot similarities in species assemblages ii. Mortality and recruitment rates iii. Growth iv.Species change v. Above ground biomass	No deterioration detected in diversity and/or abundance of identified flora in NGR.	 Whole area 12 permanent sample plots: 12 x 0.13 ha 	 Conduct periodic patrolling and surveillance in designated HCV areas. No encroachment. No illegal felling and collecting trees species (except for authorized survey) Establish a long term biodiversity monitoring system for critical forest ecosystem, flora. Marked on the ground and on the maps of high conservation value plant species discover in PSP Plots and nature trails in NGR Project area. Field staff is required to attend training courses on plants to further enhance their botanical knowledge on species that are 	 Periodic monitoring and control. Monitoring to be conducted as documented target in AWP as follow: PSP Plot maintenance: once a year. Ensuring the tie point is still in position. Ensuring the plant indicating endemic tree species is still in position Re-paint all marked tree within the radius plot area of 20 m Re-paint each number on the tree. Periodic monitoring by conducting reenumeration of all trees in the permanent sample plots and to be conducted once every three year to get 	 AWP, Compliance Report and Quarterly Report PSP Plot Report. Wildlife monitoring report. Satellite imagery monitoring report. Annual monitoring report on wild animals utilise the integrated mosaic planted forest of Empayar Kejora Sdn Bhd

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	FRATEGIES	OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
						 currently listed in the threatened, endemic and forestry prohibited list to ensure they do not harvest or damage and also for monitoring purposes. Signboards of PSPs (No., Coordinate No, etc.), HCV 1.2 and list of RTEs species within the PSPs to be erected at the strategic locations. 	 indication of changes in tree structure and species assemblages. Analysis report of PSP re-enumeration to be conducted once every three year. 	
1.2	[Listed] Fauna 17 species of fauna	Biodiversity in NGR is maintained or enhanced	Fauna diversity in NGR is maintained *Indicators: 1. Number of RTE Species (Refer: IUCN Redlist), Wildlife Conservation Enactment 1997), CITES	No deterioration detected in diversity and/or abundance of identified fauna in NGR.	Whole area Adjacent forest reserve	 Conduct periodic patrolling and surveillance in designated HCV areas. No encroachment. No poaching. Establish a long term biodiversity monitoring system for critical forest ecosystem, fauna. Field staff is required to attend training courses on wildlife to further enhance their wildlife knowledge on species that are currently listed in the threatened, endemic and forestry prohibited list to ensure they do not harvest or damage and also for monitoring 	 Quarterly progress reports in reporting of the progress of activities as prescribed in the approved Annual Work Plan (AWP), encompassing reporting of monitoring results of know HCV attributes. Annual wildlife monitoring report from Empayar Kejora Sdn Bhd shall be submitted to Sabah Forestry Department once in a year. 	

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	RATEGIES	OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
						 purposes. Identifying migratory pathway of wildlife within the project area and adjacent to the project area. 		
1.3	[Listed] Flora 162 species endemics to Borneo- 16 of these endemic to Sabah.	Biodiversity in NGR is maintained or enhanced	Flora diversity in NGR is maintained *Indicators: 1. Number of Endemic flora species (Refer: i. Plot similarities in species assemblages ii. Mortality and recruitment rates iii. Growth iv.Species change v. Above ground biomass	No deterioration detected in diversity and/or abundance of identified flora in NGR.	Similar to HCV 1.2 (Flora)	Similar to HCV 1.2 (Flora)	Similar to HCV 1.2 (Flora)	 AWP, Compliance Report and Quarterly Report PSP Plot Report. Wildlife monitoring report. Annual monitoring report on wild animals utilise the integrated mosaic planted forest of Empayar Kejora Sdn Bhd
1.3	[Listed] Fauna 5 species of mammals, 4 species of birds, 4 species of frogs, 1 species of reptile and 12 species of insects	Biodiversity in NGR is maintained or enhanced	Fauna diversity in NGR is maintained *Indicators: 1. Number of RTE Species (Refer: IUCN Redlist), Wildlife	No deterioration detected in diversity and/or abundance of identified fauna in NGR.	Similar to HCV 1.2 (Fauna)	Similar to HCV 1.2 (Fauna)	Similar to HCV 1.2 (Fauna)	

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	RATEGIES	OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
	are endemic		Conservation Enactment 1997), Cites			N	M	
1.4	The limestone Karst in Batu Timbang VJR - nesting site for swiftlet, bats and other troglofauna.	Biodiversity of Batu Timbang VJR cave is maintained or enhanced	The limestone Karst in Batu Timbang VJR is maintained *Indicators: 1. Total weight of bird's nest harvested 2. Number of SFD revenue (RM) from COI endorsement 3. Number of Bird's Nest Stakeholder Consultation	Illegal harvesting of bird's nest in a reducing trend.	VJR Batu Timbang	 No entry without permission except: Approved by Persatuan Warisan Gua Batu Timbang Kuamut. Approved by Mahkamah Anak Negeri Approved by Sabah Wildlife Department No poaching. No illegal tree felling No illegal cultivation No littering and vandalism act No fishing No open burning Conduct periodic aerial survey to curb illegal activities such as encroachment and poaching. 	 Monitoring to be conducted as documented target in AWP as follow: Bird's nest Stakeholder Consultation: Once a year 	 AWP, Compliance Report and Quarterly Report Bird's Nest Stakeholder Consultation meeting minutes. Certificate of Identity (COI) records. Aerial survey report.
2	NGR connects Danum, Imbak Canyon and Maliau Basin	The whole project site remained protected from serious environmental or ecological threat.	Encroachment into the Project site is controlled and reduced. *Indicators: 1. Number of encroaches arrested per year / forest fire	Illegal entry into NGR in a reducing trend.	Whole area Adjacent forest reserve	Similar to HCV 1.1, 1.2 & 1.3	Similar to HCV 1.1, 1.2 & 1.3	 Aerial survey report Gate control report Stakeholder Consultation meeting minutes. List of stakeholders AWP, Compliance Report and

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT STRATEGIES		OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
			incidents. 2. Number of RTE & Endemic flora species (Refer: i. Plot similarities in species assemblages ii. Mortality and recruitment rates iii. Growth iv.Species change v. Above ground biomass 3. Number of RTE & Endemic Fauna Species (Refer: IUCN Redlist), CITES & Wildlife Conservation Enactment 1997					 Quarterly Report Communities and stakeholders grievance report Satellite imagery monitoring report. PSP Plot Report analysis from FRC. Wildlife monitoring report. Annual monitoring report on wild animals utilise the integrated mosaic planted forest of Empayar Kejora Sdn Bhd
3	Extreme lowland classified as mixed dipterocarp with mixture of kerangas forest and limestone vegetation.	Ecosystem and rich biodiversity in NGR SFM which is rich in significant diversity of ecosystem are maintained and enhanced.	Reduce the number of illegal activities and fire occur within the NGR SFM *Indicators: 1. Number of encroaches arrested per year / forest fire	Illegal entry into NGR in a reducing trend. No deterioration detected in diversity and/or abundance of identified flora and fauna in NGR.	Whole area	Similar to HCV 1.1, 1.2, 1.3	Similar to HCV 1.1, 1.2, 1.3	 Aerial survey report Gate control report Stakeholder Consultation meeting minutes. List of stakeholders AWP, Compliance Report and Quarterly Report Communities and


		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	RATEGIES	OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED
			incidents 4. Number of RTE & Endemic flora species (Refer: i. Plot similarities in species assemblages ii. Mortality and recruitment rates iii. Growth iv.Species change v. Above ground biomass 2. Number of RTE & Endemic Fauna Species (Refer: IUCN Redlist), CITES & Wildlife Conservation Enactment 1997					stakeholders grievance report • Satellite imagery monitoring report. • PSP Plot Report. • Wildlife monitoring report.
4.2	All areas with slopes >25° and 30 m riparian buffer strips	Areas with slopes >25° and river or streams banks are maintain to prevent soil erosion	Maintain the Areas with slopes >25° and riparian reserves to prevent soil erosion. *Indicators: 1. Total number of Water quality parameter	Zero encroachment, fire, illegal felling, illegal cultivation, and illegal occupation activities which are might contribute water pollution's.	• 30m buffer zone	 Conduct periodic patrolling and surveillance in designated HCV areas. No entry without permission. No poaching. No illegal felling No illegal fishing No open burning No chemical 	 Periodic monitoring and control to prevent encroachment in HCV areas. Quarterly progress report in reporting of the progress of activities as prescribed in the approved AWP encompassing reporting of monitoring results of 	 Water quality report. AWP, Compliance Report and Quarterly Report Environmental quarterly report. Annual Environmental Monitoring report

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	TRATEGIES	OPERATIONAL /	DOCUMENT TO BE
HCV	ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS	PRESCRIPTIONS	- STRATEGIC MONITORING	REVIEWED
	Buffer strips of 50 m inside the southern	Barriers to destructive fire	(Temperature, pH value, DO, Conductivity, TDS) Maintain the buffer zone along	Zero encroachment,	50m buffer zone	 treatment/application Warning signage to be erected and visibly maintained at the strategic location. Maintain the natural vegetation along the riparian buffer zone. Conduct water sampling along Sg. Lanap, Sg Kasuyan, Sg Kuamut and Sg. Imbok No encroachment. No entry without 	 known HCV attributes. Water sampling to be conducted twice yearly. Aerial surveillance: 2 times per year. 	• Aerial surveillance report
4.3	southern boundaries of NGR project area that bordering oil palm estate are categorised as HCV 4.3.	are maintained and enhance.	neighbouring boundaries of oil palm plantations *Indicators: Number of forest fire incidents	fire, poaching, illegal felling, illegal cultivation, and illegal occupation activities which are might contribute to fire.		 permission No poaching. No open burning. Maintain the natural vegetation along the buffer zone. The Forest Fire Management Plan has to be updated periodically. 	 Ground patrolling. Collecting FDRS data: Every day. Ensure that all fire prevention procedures (monitoring, fire drills, public awareness campaign etc.) to be practised on a regular basis (at least once a year), especially during the drought season. Main stakeholder consultation with neighbouring stakeholders and contractors to be conducted once a year for forest fire awareness. 	 Ground patrolling report FDRS report Fire drill training records AWP, Compliance Report and Quarterly Report Communities and stakeholders grievance report Satellite imagery monitoring report. Stakeholder Consultation meeting minutes.

		GENERAL HCV	SPECIFIC HCV	MANAGEMENT	MANAGEMENT ST	RATEGIES	OPERATIONAL /	DOCUMENT TO BE
HC	/ ID	MANAGEMENT OBJECTIVE	MANAGEMENT OBJECTIVE	TARGETS	AREAS PRESCRIPTIONS	STRATEGIC MONITORING	REVIEWED	
6	The mixed dipterocarp forest and limestone vegetation of CJR Batu Timbang are categorized as HCV 6 to depict the cultural value importance for the nine teriti of Kg Kuamut.	Cultural value in the NGR SFM are maintained and enhanced	Maintain the area of cultural value *Indicators: Similar to HCV 1.4	Zero encroachment within the significant HCV 6 area.	VJR Batu Timbang	Similar to HCV 1.4	Similar to HCV 1.4	 AWP, Compliance Report and Quarterly Report Bird's Nest Stakeholder Consultation meeting minutes. Certificate of Identity (COI) records. Aerial survey report.

3.0 Effectiveness of monitoring program and enhancements by each HCV Attributes:

Based on the summary and monitoring activities table above, the effectiveness of the monitoring activities can be elaborated as follows:



3.1 HCV 1 BIODIVERSITY VALUES

Map 1 of HCV 1: Biodiversity Value

3.1.1 HCV 1.1 Protected Areas

The protected areas activities in the NGR SFM project are divided into several activities, such as the following:

i. <u>Patrolling and enforcement, arrest, prosecution</u>

Based on the Annual Report 2013 – 2019, there were no arrests made from 2013 – 2016, but there were (5) arrest made in 2017, increased again to (10) arrest in 2018 and decreased to (1) arrest in 2019. There were no arrest made in the year 2020.




Figure 1: Number of Encroaches Arrested

Based on data from 2013 – 2019 (See Figure 1), the number of encroaches arrested was decreased. Patrolling and enforcement activities is intensified.

ii. <u>Aerial Surveillance</u>

Based on Figure 2, the data from year 2014 – 2019 shows data of the frequency of the aerial surveys done. As shown in the graph, the aerial surveillance had not been consistent in yearly hours as it was sometimes difficult to arrange the schedule with the Sabah Air. The frequency as per graph below:




Figure 2: Number of Aerial Surveillance Frequency (Hours)



Figure 3: Aerial Surveillance

iii. Inspection of boundaries and re-brushing of NGR SFM Project

As stated in the AWP 2014 – 2019 and also Compliance Report 2014 – 2019, the inspection and re-brushing of main boundaries was conducted 10,000 meter per year. This corresponds to the responsibility of the team Management to protect the project area and installing of proper signages to ensure that all the stakeholders are aware of entering the area. There are few signages had been placed as follows :

- Project signboards on all entries
- Warning signboards on all entries.
- Safety signage.
- Riparian signage
- HCV signage.
- PSP signage.
- Enforcement signage.
- Prohibited activities signage




Figure 4: Distance of Main Boundary Re-Brushing


Based on Figure 4, the boundary re-brushing activities were well maintained every year. Since 2014 until 2019, the distance of boundary re-brushing were achieved with a total distance of 45,000 metre at Mount Magdalena FR Class 1 and 15,000 metre at Northern Gn. Rara FR Class 1.



Figure 5: Inspection of Boundary

iv. Establishment of Forest Checking Station

The Project team has established enforcement gates and Forest Checking Stations, in regards to the team's responsibility to safeguard the HCVs as a whole. The two established Forest Checking Stations are FCS Batu Timbang and FCS Imbok. FCS Bt Timbang was established due to the frequent number of complaints received on poachers entering from the northwestern route of the project area, while FCS Imbok covers the eastern side of the project area.



Map 2: Map showing the aerial survey route covering the whole project area, ground patrolling route and road accessibility, and locations of the Forest Checking Stations



Figure 6: FCS Batu Timbang

Figure 7: FCS Imbok

3.1.2 HCV 1.2 Threatened and Endangered Species FLORA DIVERSITY

PSP Plot Analysis

The PSP Plots analysis will be prepared by FRC Sepilok. Eleven (11) 0.125 ha permanent sample plots (PSPs) were established end of 2014 and early 2015 (the report refer as 1st census 2015), and further re-census in 19th till 24th March 2018. An addition of one PSP Plot that is plot no. 12 were established in the year 2017. However, the re-enumeration for plot no.12 has not been conducted as it will be re-enumerate in the year 2020.

The overall objectives of the monitoring activities are to investigate changes of trees \geq 10 cm dbh according to the prescribe indicators:

- i. plot similarities in species assemblages
- ii. mortality and recruitment rates
- iii. growth
- iv. species change
- v. above ground biomass


Re-measurement of recorded trees

The location of eleven PSPs is shown in Table 1 and Map 3. All previously labelled-trees ≥ 10 cm diameter at breast height (dbh) within 20 m radius circular plot and newly recruited trees were labelled, measured and identify to species level.

Table 1: Summary of location, date of census and altitude of the eleven PSPs at Northern GunungRara Sustainable Forest Management Project Area in Sabah, Malaysia.

Plot No.	Location	Ist Census	2nd Census	Latitude	Longitude	Altitude (m)
1	Mt Magdelena	24-0ct-14	19-Mar-18	N 04 ⁰ 58' 26.0"	E 117 ⁰ 08' 56.6"	197
2	Mt Magdelena	24-0ct-14	19-Mar-18	N 04 ⁰ 58' 22.6"	E 117 ⁰ 09' 0.7"	190
3	Mt Magdelena	13-Nov-15	20-Mar-18	N 04 ⁰ 58' 47.3"	E 117 ⁰ 08' 57.2"	190
4	Mt Magdelena	13-Nov-15	20-Mar-18	N 04 ⁰ 59' 33.4"	E 117 ⁰ 08' 07.8"	223
5	Mt Magdelena	19-Nov-15	22-Mar-18	N 04 ⁰ 53' 41.7"	E 117 ⁰ 23' 45.7"	307
6	Mt Magdelena	19-Nov-15	22-Mar-18	N 04 ⁰ 53' 55.7"	E 117 ⁰ 23' 11.7"	283
7	Mt Magdelena	20-Nov-15	22-Mar-18	N 04 ⁰ 50' 30.5"	E 117 ⁰ 13' 54.8"	188
8	Mt Magdelena	22-Jan-15	23-Mar-18	N 04 ⁰ 56' 09.3"	E 117 ⁰ 10' 34.6"	350
9	Batu Timbang	12-Mar-15	23-Mar-18	N 04 ⁰ 58' 21.9"	E 117 ⁰ 05' 57.4"	455

10	Batu Timbang	12-Mar-15	24-Mar-18	N 04 ⁰ 58' 19.6"	E 117 ⁰ 05' 51.8"	532
11	Mt Magdelena	13-Mar-15	24-Mar-18	N 04 ⁰ 55' 43.9"	E 117 ⁰ 10' 51.4"	267




Map 3: The distribution of the eleven PSPs at Northern Gunung Rara Sustainable Forest Management Project Area in Sabah, Malaysia.


RESULTS

i. Plot similarities in relation to species assemblage

The cluster analysis based on species abundance indicated that all PSPs enumerated in 2015 and 2018 can be grouped into five forest types or species assemblages (Figure 8). The grouping can be differentiated in relation to level of past disturbances and edaphic factors. Group 1 and 3 are forest that were dominated by secondary tree species where the former natural climax forest could have been totally cleared in the past. Group 2 is mainly trees that are found associated with limestone substrate. The other groups are mainly advance regenerating lowland mixed dipterocarp forest. In the three-year interval (2015-2018), the species assemblages grouping has shown no deviation from the initial observation in 2015 (Figure 8).



Census 2015



Census 2018

Figure 8. The dendrogram derived from Bray-Curtis ordination analysis (1957) for trees recorded in eleven PSPs at Northern Gunung Rara Sustainable Forest Management Project Area in Sabah, Malaysia: A) year 2015 and B) year 2018. (Note: 0 = completely similar & 1 = completely dissimilar).

ii. Mortality, recruitment and turnover

A comparison of tree mortality and recruitment for all sample plots is presented in Table 2. Seven out of eleven plots had greater numbers of recruits than deaths over the interval 2015–2018. Two plots had slightly more mortality than recruitments, and one has similar count of mortality and recruitments. One plot has no observation of mortality and recruitment of trees. In total, tree recruitment is three times more than mortality. The highest number of tree deaths were recorded in Plot 3 and 7, whereas the greatest number of recruits were observed in Plot 1, 4 and 5.

Plot No	Tree	Tree	No of	No of	Mortality	Recruitment	Turnover
	2014	2017	Dead	Recruit	Rate (%	Rate (% per	Rate (%
					per year)	year)	per year)
1	75	84	3	12	1.19	4.41	2.80
2	69	70	1	2	0.43	0.84	0.64
3	52	56	5	9	2.96	5.08	4.02
4	65	83	0	18		7.01	3.51
5	60	72	2	14	1.01	6.26	3.63
6	34	40	0	6	0.00	4.74	2.37
7	50	48	4	2	2.46	1.26	1.86
8	62	62	7	7	3.71	3.71	3.71
9	57	55	2	0	1.17	0.00	0.58
10	47	47	0	0	0.00	0.00	0.00
11	53	59	1	7	0.63	4.08	2.35
Grand Total	624	676	25	77	1.23	3.40	2.32

Table 2 Dynamic of trees \geq 10 cm dbh in all 11 permanent sample plots (PSPs) between 2015–18 in Northern Gunung Rara Sustainable Forest Management project area, Sabah, Malaysia

iii. Growth

A comparison of tree growth on all sample plots is presented in Table 3. On average, all trees \geq 10 cm dbh for all plots have demonstrated positive growth within the interval 2015-2018. Trees in Plot 4 demonstrated the highest growth rate and Plot 9 as the lowest.

Plot			Min of	Max of		Min of	Max of
No.	Ν	Mean of AGR	AGR	AGR	Mean of RGR	RGR	RGR
1	71	0.31 ± 0.06	-0.38	2.55	1.31 ± 0.25	-1.90	9.79
2	67	0.38 ± 0.09	-0.59	4.10	2.11 ± 0.54	-2.70	23.30
3	45	0.44 ± 0.17	-5.06	2.74	2.27 ± 0.47	-4.81	13.68
4	64	0.93 ± 0.09	-0.27	3.93	4.90 ± 0.41	-1.03	17.30
5	57	0.60 ± 0.13	-0.75	4.90	3.99 ± 0.88	-4.98	37.70
6	34	0.74 ± 0.13	-0.36	2.78	2.93 ± 0.49	-2.11	9.67
7	46	0.55 ± 0.10	-1.44	3.20	2.71 ± 0.46	-3.68	16.84
8	55	0.54 ± 0.09	-0.85	3.19	3.28 ± 0.62	-4.48	24.51
9	54	0.23 ± 0.11	-2.24	3.95	0.91 ± 0.22	-3.35	5.90
10	47	0.49 ± 0.10	-0.03	3.59	2.28 ± 0.43	-0.21	17.10
11	50	0.51 ± 0.11	-0.33	4.29	2.63 ± 0.56	-1.82	18.64

Grand	500	0.51	+ 0.02	-5.06	4 00	2 66	+ 0.16	1 0 9	27 70
Total	390	0.51	± 0.03	-5.00	4.90	2.00	± 0.16	-4.90	37.70

Table 3 Growth of trees \geq 10 cm dbh in all 11 permanent sample plots (PSPs) between 2015–18 in Northern Gunung Rara Sustainable Forest Management project area, Sabah, Malaysia. (Annual Growth Rate, AGR; Relative Growth Rate, RGR)

iv. Species compositional changes

The number of species that were recruited is almost three times more than those recorded as dead in 2018, i.e. 52 species recorded as newly recruits and 19 species as dead (Table 4). This demonstrated that diverse tree species were recruited in all forest ecosystem throughout the three-year period. The composition of recruited trees also varies between successional group of trees from mixed climax and pioneer species, and also mixed structural canopy, such as main canopy, middle storey and understorey species. Based on plant traits, of the 52 species recruited, about 42% of the list are large stature climax species and 21% as pioneer. As for dead individuals, of the 19 species, both pioneer and climax contributed about 37% of total mortality, respectively.

Plot No	Recruits	Dead
1	Alangium javanicum	Nephelium maingayi
	Ardisia macrophylla	Shorea angustifolia
	Artocarpus kemando	Xanthophyllum vitellinum
	, Artocarpus lanceifolius	. ,
	Gluta wallichii	
	Lithocarpus beccarianus	
	Lithocarpus canfleyanus	
	Macaranga gigantea	
	Shorea ferruginea	
	Shorea parvifolia	
	Vatica albiramis	
2	Hopea beccariana	Lophopetalum subovatum
	Vatica chartacea	
3	Baccaurea tetrandra	Macaranga hypoleuca
	Cephalomappa malloticarpa	Macaranga indistincta
	Dendrocnide eliptica	Shorea ferruginea
	Knema latifolia	, ,
	Macaranga hypoleuca	
	Mallotus molissimus	
	Palaquium calophyllum	
	Shorea macrophylla	
4	Litsea garcia	
	Ludekia borneensis	
	Macaranga cf. grandibracteota	
	Macaranga pearsonii	
	Nauclea subdita	
	Neonauclea artocarpoides	
	Norrisia major	
	Terminalia citrina	
5	Aglaia rufinervis	Macaranga pearsonii
	Baccaurea macrocarpa	
	Brownlowia peltata	
	Dendrocnide elliptica	
	Garcinia gaudichaudi	
	Ludekia borneensis	
	Mallotus korthalsii	
	Paranephelium joannes	
	Parashorea tomentella	
	Polyalthia rumphii	
	Shorea johorensis	
	Syzygium tawahense	
6	Callicarpa pentandra	

	Dendrocnide elliptica Parashorea tomentella	
	Symplocos fasciculata	
7	Hopea ferruginea	Baccaurea bracteata
	Sindora irpicina	Bridelia sp.
		Callicarpa plumosa
		Litsea acceden
8	Aporosa acuminatissima	
	Artocarpus lanceifolius	
	Chisocheton medusae	
	Dacryodes rostrata	
	Pentace laxiflora	
9	-	Diospyros sp.
		Dryobalanops lanceolata
10	-	-
11	Dacryodes laxa	Macaranga hypoleuca
	Dacryodes rugosa	
	Shorea macroptera	
	Shorea parvifolia	
	Shorea pilosa	
	Swintonia acuta	
	Vatica albiramis	

Table 4 List of species recruited and dead of trees ≥ 10 cm dbh within the elven PSPs between 2015–18 in Northern Gunung Rara Sustainable Forest Management project area, Sabah, Malaysia

v. Above ground biomass changes


Throughout the period 2015–2018, the overall total AGB indicates an increase of 8 % from the initial observation value (Table 5). About nine PSPs demonstrated an increase of AGB with range from 6-33% from their initial values, and two PSPs indicated a decrease of 2-13%.

Plot No	AGB 2015		AGB	2018	Rate of change %
	N	Total (Mg/ha)	N	Total (Mg/ha)	
1	75	542	84	578	7
2	69	782	70	892	14
3	52	585	56	509	-13
4	65	272	83	362	33
5	60	488	71	478	-2
6	34	171	40	223	31
7	50	283	48	317	12
8	62	185	62	196	6
9	57	792	55	865	9
10	47	315	47	367	16
11	53	356	59	387	9
Grand Total	624	4769	675	5174	8

Table 5 Comparison of two census years (2014 and 201)7 on above ground biomass (AGB) per hectare of lived standing trees with ≥ 10 cm dbh within the 11 PSP in Timimbang Botition Sustainable Forest Management project area, Sabah, Malaysia.

FAUNA DIVERSITY

The NGR SFM Team established (5) methods of wildlife monitoring, that is; (a)Recce Walk Survey; (b)Camera Trapping; (c)Night Drive Survey; (d)Opportunistic Sightings; & (e)Gibbon Call. Of these (5) wildlife monitoring methodology, we analysed (5) of the methods. Below is a table of NGRSFM Wildlife RTE Indicator:



Map 4: Map of RTE Species (Flora&Fauna)

			Update NGR	SFM Wildlife	e List N	lac 2014-Dec2019				
No	Common name	Family	Scientific Name	WCE (SWD). 1997	CITES	Status IUCN Redlist	Endemic	HCV PRIORITY SPECIES	IUCN Red List	Comment
1	Asian Black Hombill	Bucerotidae	Anthracoceros malayanus			Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
2	Asian Giant Tortoise	Testudinidae	Manouria emys	Ш		Status: Endangered ver 2.3 Pop. trend: decreasing			EN	2019
3	Asian Glossy Starling	Sturnidae	Aplonis panayensis			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	2019
4	Asian Koel	Cuculidae	Eudynamis scolopacea			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
5	Asian Paradise Flycether	Monarchidae	Terpsiphone paradisi			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
6	Banded Linsang	Prionodon tidae	Prionodon linsang	I	I	Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
7	Banded Palm Civet	Viverridae	Hemigalus dermianus	I	I	Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
8	Bearded Pig	Suidae	Sus Barbatus	ш		Status: Vulnerable ver 3.1 Pop. trend: decreasing		1	VU	
9	Binturong	Viverridae	Arctictis binturong	I	III	Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
10	Black and Red Broadbill	Eurylaimidae	Cymbirhynchus macrorhynchos			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
11	Black and Yellow Broadbill	Eurylaimidae	Eurylaimus ochromalus			Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
12	Black Giant Squirrels	Sciuridae	Ratufa affinis			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
13	Blue-eared Barbet	Megalaimidae	Megalaima australis			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
14	Blue-Headed Pitta	pittidae	Pitta baudii			Status: Vulnerable ver 3.1 Pop. trend: decreasing	Endemic in Borneo(Bird's)		VU	
15	Bornean Banded Pitta	pittidae	Pitta guajana			Status: Least Concern ver 3.1 Pop. trend: decreasing	Endemic in Borneo(Bird's)		LC	
16	Bornean Banteng (Tambadau)	Bovidae	Bos Javanicus	1	I	Status: Endangered ver 3.1	Endemic in Borneo	4	EN	
17	Bornean Bay Cat	Felidae	Catopuma badia	I	1	Pop. trend: decreasing Status: Endangered ver 3.1	Endemic in Bomeo	1	EN	
18	Bornean Clouded Leopard	Felidae	Neofelis diardi	1	1	Pop. trend: decreasing Status: Vulnerable ver 3.1	Bonieo	1	VU	
19	Bornean Crested Fireback	Phasianidae	Lophura Ignita			Pop. trend: decreasing Status: Near Threatened ver 3.1 Pop. trend: decreasing	Endemic in Borneo(Bird's)		NT	
20	Bornean Gibbon	Hylobatidae	Hylobates funereus		1	Status: Endangered ver 3.1	Endemic in Borneo	1	EN	
21	Bornean Ground Cuckoo	Cuculidae	Carpococcyx radiatus			Pop. trend: decreasing Status: Near Threatened ver 3.1	Endemic in Borneo(Bird's)		NT	
22	Bornean Mountain Ground	Sciuridae	Dremomys everetti			Pop. trend: decreasing Status: Least Concern ver 3.1	Endemic in Bomeo		LC	
23	Squirrel Bornean Necklaced Partridge	Phasianidae	Arborophila graydoni			Pop. trend: decreasing Status: Least Concern ver 3.1	Endemic in Borneo(Bird's)		LC	
24	Bornean Orangutan	Hominidae	Pongo pygmaeus	1	I	Pop. trend: decreasing Status: Critically Endangered ver 3.1	Endemic in	1	CR	
25	Bornean Pygmy Elephant	Elephantidae	Elephas maximus bomeensis	II (I since	1	Pop. trend: decreasing Status: Endangered ver 3.1	Bomeo Endemic in	1	EN	
26	Bornean Striped Palm Civet	Viverridae	Arctogalidia stigmatica	June 2013)		Pop. trend: decreasing Status: Least Concern ver 3.1	Bomeo Endemic in		LC	
27	Bornean Sun Bear	Ursidae	Helarctos malayanus	1	1	Pop. trend: decreasing Status: Vulnerable ver 3.1	Bomeo Endemic in	1	VU	
28	Bornean-yellow Muntjac	Cervidae	Muntiacus atherodes			Pop. trend: decreasing Status: Near Threatened ver 3.1	Bomeo Endemic in		NT	
29	Brahminy Kite		Haliastur Indus			Pop. trend: decreasing	Bomeo		DD	2019
30	Brown Wood-old	Strigidae	Strix leptogrammica			Status: Least Concern ver 3.1			LC	
31	Common Kingfisher	Alcedinide	Alcedoatthis			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
32	Common Palm Civet (Island	Viverridae	Paradoxurus philippinensis	11		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
33	Palm Civet) Common Porcupine or	Hystricidae	Hystrix brachyura			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
34	Malayan Porcupine Crab-eating Mongoose	Herpestidae	Herpestes urva			Pop. trend: decreasing Status: Near Threatened ver 3.1			NT	
35	Crested Serpent Eagle	Accipitridae	Spilornis cheela			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
36	Emerald Dove	Columbidae	Chalcophaps indica			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
37	Four-striped Ground Squirrel	Sciuridae	Lariscus hosei			Pop. trend: decreasing Status: Near Threatened ver 3.1			NT	
38	Giant Squirrel	Sciuridae	Ratufa affinis			Pop. trend: decreasing			aa	
39	Great Argus	Phasianidae	Anhinga melanogaster			Status: Near Threatened ver 3.1			NT	
40	Great Egret	Ardeidae	Ardea alba			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
40	Greater Coucal	Cuculidae	Centropus sinensis			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
42	Greater Mouse-deer	Tragulidae	Tragulus napu			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
72		rragunuae				Pop. trend: decreasing				

43	Grey Night Jar	Caprimulgidae	Caprimulgus jotaka			Status: Least Concern ver 3.1			LC	
44	Helmeted Hombill	Bucerotidae	Buceros vigil			Pop. trend: decreasing Status: Critically Endangered ver 3.1			CR	
45	Hill Myna	Sturnidae	Gracula religiosa			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
40	Horse-tailed Squirrel	Sciuridae	Sundasciurus hippurus			Pop.trend:decreasing			20	
						Status: Least Concern ver 3.1				
47	Leopard Cat	Felidae	Prionailurus bengalensis			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
48	Lesser Mouse-deer	Tragulidae	Tragulus kanchil			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
49	Little Spider Hunter	Nectariniidae	Arachnothera longirostra			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
50	Long-tailed Macaque	Cercopithecidae	Macaca fascicularis		II	Pop. trend: decreasing			LC	
51	Long-tailed Pocupine	Hystricidae	Trichys fasciculata	1		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
52	MalayCivet	Viverridae	Viverra tangalunga			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
53	Malayan/Common Porcupine	Hystricidae	Hystrix brachyura	Ш		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
54	Marbled Cat	Felidae	Pardofelis marmorata	Ш	T	Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
55	Mask Palm Civet	Viverridae	Paguma larvata	Ш		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
56	Oriental Bay Owl	Tytonidae	Phodilus badius			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	2019
57	Oriental Darter	Anhingidae	Anthracoceros malayanus			Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
58	Oriental Pied Hombill	Bucerotidae	Anthracoceros albirostris			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
59	Otter Civet	Viverridae	Cynogale bennettii	1		Status: Endangered ver 3.1 Pop. trend: decreasing		1	EN	
60	Pig-tailed macaque	Cercopithecidae	Macaca nemestrina	11	1	Status: Vulnerable ver 3.1			VU	
61	Plantain Squirrel	Sciuridae	Callosciurus notatus			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
62	Prevost's Squirrel	Sciuridae	Callosciurus prevostii			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
	Purple Heron	Ardeidae	Ardea purpurea			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
64	Red Eyed Bulbul	Pycnonotidae	Pycnonotus brunneus			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
	Red Giant Flying Squirrel	Petauristinae		1		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
			Petaurista petaurista	"		Pop. trend: decreasing Status: Least Concern ver 3.1				
66	Red Junglefowl	Phasianidae	Galus-galus			Pop.trend:decreasing	Endemic in		LC	
	Red Langur		Presbytis rubicanda			Status: Least Concern ver 3.1	Borneo		040	2019
68	Red Muntjac or Barking Deer	Cervidae	Muntiacus muntjak	III		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
69	Reticulated Python	Pythonidae	Python reticulatus	Ш		Pop. trend: decreasing Status: Near Threatened ver 3.1			LC	
70	Rhinoceros Hornbill	Bucerotidae	Buceros rhinoceros			Pop. trend: decreasing Status: Near Threatened ver 3.1			NT	
71	Rouroul	Phasianidae	Rollulus rouroul			Pop.trend:decreasing	F ()))		NT	2019
72	Sabah Grey Langur	Cercopithecidae	Presbytis sabana	I		Status: Endangered ver 3.1 Pop. trend: decreasing	Endemicin Borneo		EN	2019
73	SambarDeer	Cervidae	Cervus unicolor	Ш	II	Status: Vulnerable ver 3.1 Pop.trend: decreasing		\checkmark	VU	
74	Short-tailed Mongoose	Herpestidae	Urva brachyurus	II		Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
75	Small-clawed Otter	Mustelidae	Aonyx cinerea	II	Ш	Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
76	Smooth Otter	Mustelidae	Lutra perspicillata	11		Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
77	Smooth-coated Otter	Mustelidae	Lutrogale perspicillata	11		Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
78	Sunda Giant Squirrel	Sciuridae	Ratufa affinis sandakanensis	11		Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
79	Sunda Pangolin	Manidae	Manis Javanica	13th Feb	11	Status: Critically Endangered ver 3.1 Pop. trend: decreasing		1	CR	
80	Sunda Skunk/Teludu	Mephitidae	Mydaus javanensis	2019)		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
81	Sunda Slow Loris	Lorisinae	Nycticebus menagensis		1	Status: Vulnerable ver 3.1			VU	
82	Thick-Spined Porcupine	Hystricidae	Hystrix crassispinis			Pop. trend: decreasing Status: Least Concern ver 3.1	Endemic in		LC	
83	Thomas's Flying Squirrel	Petauristinae	Aeromys thomasi			Pop. trend: decreasing Status: Least Concern ver 3.1	Borneo Endemic in		LC	
84	Tufted Ground squirrel	Sciuridae	Rheithrosciurus macrotis			Pop.trend: decreasing Status: Vulnerable ver 3.1	Borneo		VU	
85	Water Monitor	Varanidae	Varanus salvator			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
						Pop. trend: decreasing			LC	0010
86	Wreathed Hombill	Bucerotidae	Rhyticeros undulatus			Status: Data Deficient ver 3.1	Endemic in		00	2019
87	White-crowned Shama	Muscicapidae	Copsychus stricklandii			Pop. trend: unknown Status: Data Deficient ver 3.1	Borneo(Bird's)		DD	
88	Woodpeckersp		Unknown			Pop. trend: unknown Status: Least Concern ver 3.1			DD	
89	Yellow-throated Marten	Mustelidae	Martes flavigula	Ш		Pop. trend: decreasing			LC	

Table 6 #List Of Species RTE recorded from year 2014-2019 (Ref IUCN)

Based on the data collected from all the wildlife monitoring methods, the status of NGR RTE by percentage are as follows:

Status IUCN Redlist	No. of Widlife Species (Mammals)	Percentage (%)	Remarks	Percentage (%)
Critically Endangered	2	4%	Threatened	40%
Endangered	6	13%		
Vulnerable	11	23%		
Near Threatened	7	15%	Near Threatened	15%
Least Concern	22	46%	Least Concern	46%
Endemic	19	40%	Endemic	40%
HCV Priority	11	23%	HCV	23%
Total	48	100%		100%




Figure 9: Chart of RTE Species Recorded in NGR SFM (2014-2019)

Based on monitoring records form the year 2014-2019, the RTE indicator status shows that Least Concern (LC) dominated the chart with 50.6 %, Endangered (EN) + Critically Endangered (CR) + Vulnerable (VU) 25.8 %, Near Threatened (NT) 15.7 % and Data Deficient (DD) 7.9 %.


Eight new species have been evaluated and recorded through four monitoring methods namely:-

- 1. Mammals *Red Langur* and *Sabah Grey Langur*
- 2. Reptiles Asian Giant Tortoise
- 3. Bird's Asian Glossy Starling, Brahminy Kite, Oriental Bay Owl, Rouroul and Wreathed Hornbill

3.1.3 HCV 1.3 Endemism

FLORA DIVERSITY

Similar to HCV 1.2.



Map 5: Map of Endemic Species (Flora&Fauna)

There are nineteen known endemic species of wildlife found in the Project Area. Based on the analysis result of the wildlife monitoring, there were 13 endemic mammals species and 6 endemic bird species identified within the project area. Most of the identified species were recorded through camera traps.

Table 8: #List Of Endemic Species Recorded From the Year 2014-2019 (Ref. IUCN)

			· · · ·			lac 2014-Dec2019	, , , , , , , , , , , , , , , , , , ,			
No	Common name	Family	Scientific Name	WCE (SWD). 1997	CITES	Status IUCN Redlist	Endemic	HCV PRIORITY SPECIES	IUCN Red List	Comment
1	Asian Black Hombill	Bucerotidae	Anthracoceros malayanus			Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
2	Asian Giant Tortoise	Testudinidae	Manouria emys			Status: Endangered ver 2.3 Pop. trend: decreasing			EN	2019
3	Asian Glossy Starling	Sturnidae	Aplonis panayensis			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	2019
4	Asian Koel	Cuculidae	Eudynamis scolopacea			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
5	Asian Paradise Flycether	Monarchidae	Terpsiphone paradisi			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
6	Banded Linsang	Prionodon tidae	Prionodon linsang	I	I	Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
7	Banded Palm Civet	Viverridae	Hemigalus dermianus	I	I	Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
8	Bearded Pig	Suidae	Sus Barbatus			Status: Vulnerable ver 3.1 Pop. trend: decreasing		1	VU	
9	Binturong	Viverridae	Arctictis binturong	I	ш	Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
10	Black and Red Broadbill	Eurylaimidae	Cymbirhynchus macrorhynchos	3		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
11	Black and Yellow Broadbill	Eurylaimidae	Eurylaimus ochromalus			Status: Near Threatened ver 3.1			NT	
12	Black Giant Squirrels	Sciuridae	Ratufa affinis	I		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
13	Blue-eared Barbet	Megalaimidae	Megalaima australis			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
14	Blue-Headed Pitta	pittidae	Pitta baudii			Pop. trend: decreasing Status: Vulnerable ver 3.1	Endemic in		VU	
15	Bornean Banded Pitta	pittidae	Pitta guajana			Pop. trend: decreasing Status: Least Concern ver 3.1	Borneo(Bird's) Endemic in		LC	
16	Bornean Banteng (Tambadau)	Bovidae	Bos Javanicus	1	I	Pop. trend: decreasing Status: Endangered ver 3.1	Bomeo(Bird's) Endemic in	1	EN	
	Bornean Bay Cat	Felidae	Catopuma badia		-	Pop. trend: decreasing Status: Endangered ver 3.1	Borneo Endemic in		EN	
	Bornean Clouded Leopard	Felidae	Neofelis diardi	-	-	Pop. trend: decreasing Status: Vulnerable ver 3.1	Bomeo		VU	
10	Bornean Crested Fireback	Phasianidae		'		Pop. trend: decreasing Status: Near Threatened ver 3.1	Endemic in	*	NT	
			Lophura Ignita			Pop. trend: decreasing Status: Endangered ver 3.1	Bomeo(Bird's) Endemic in	1		
20	Bornean Gibbon	Hylobatidae	Hylobates funereus		1	Pop. trend: decreasing Status: Near Threatened ver 3.1	Borneo Endemic in	V	EN	
21	Bornean Ground Cuckoo Bornean Mountain Ground	Cuculidae	Carpococcyx radiatus			Pop. trend: decreasing Status: Least Concern ver 3.1	Bomeo(Bird's) Endemic in		NT	
22	Squirrel	Sciuridae	Dremomys everetti	I	8	Pop. trend: decreasing Status: Least Concern ver 3.1	Bomeo Endemic in		LC	
23	Bornean Necklaced Partridge	Phasianidae	Arborophila graydoni		9 - 18	Pop. trend: decreasing Status: Critically Endangered ver 3.1	Bomeo(Bird's) Endemic in		LC	
24	Bornean Orangutan	Hominidae	Pongo pygmaeus	l II (1 since	1	Pop. trend: decreasing Status: Endangered ver 3.1	Bomeo Endemic in	1	CR	
25	Bornean Pygmy Elephant	Elephantidae	Elephas maximus bomeensis	June 2013)	1	Pop. trend: decreasing	Borneo	1	EN	
26	Bornean Striped Palm Civet	Viverridae	Arctogalidia stigmatica	I		Status: Least Concern ver 3.1 Pop. trend: decreasing	Endemic in Borneo		LC	
27	Bornean Sun Bear	Ursidae	Helarctos malayanus	I	1	Status: Vulnerable ver 3.1 Pop. trend: decreasing	Endemic in Borneo	1	VU	
28	Bornean-yellow Muntjac	Cervidae	Muntiacus atherodes	111		Status: Near Threatened ver 3.1 Pop. trend: decreasing	Endemic in Borneo		NT	
29	Brahminy Kite		Haliastur Indus		я к				DD	2019
30	Brown Wood-old	Strigidae	Strix leptogrammica			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
31	Common Kingfisher	Alcedinide	Alcedoatthis			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
32	Common Palm Civet (Island Palm Civet)	Viverridae	Paradoxurus philippinensis	I		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
33	Common Porcupine or Malayan Porcupine	Hystricidae	Hystrix brachyura	111		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
34	Crab-eating Mongoose	Herpestidae	Herpestes urva	I		Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
35	Crested Serpent Eagle	Accipitridae	Spilornis cheela			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
36	Emerald Dove	Columbidae	Chalcophaps indica			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
37	Four-striped Ground Squirrel	Sciuridae	Lariscus hosei	I		Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
38	Giant Squirrel	Sciuridae	Ratufa affinis			. ep. sona. accredancy			00	
39	Great Argus	Phasianidae	Anhinga melanogaster			Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
40	Great Egret	Ardeidae	Ardea alba			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
41	Greater Coucal	Cuculidae	Centropus sinensis			Status: Least Concern ver 3.1			LC	
42	Greater Mouse-deer	Tragulidae	Tragulus napu			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
			•			Pop. trend: decreasing				

43	Grey Night Jar	Caprimulgidae	Caprimulgus jotaka			Status: Least Concern ver 3.1			LC	0
43	Helmeted Hombill	Bucerotidae	Buceros vigil			Pop. trend: decreasing Status: Critically Endangered ver 3.1			CR	
						Pop. trend: decreasing Status: Least Concern ver 3.1				
45	Hill Myna	Sturnidae	Gracula religiosa			Pop. trend: decreasing			LC	
46	Horse-tailed Squirrel	Sciuridae	Sundasciurus hippurus			Status: Least Concern ver 3.1			DO	
47	Leopard Cat	Felidae	Prionailurus bengalensis	I		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
48	Lesser Mouse-deer	Tragulidae	Tragulus kanchil			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
49	Little Spider Hunter	Nectariniidae	Arachnothera longirostra			Pop. trend: decreasing			LC	
50	Long-tailed Macaque	Cercopithecidae	Macaca fascicularis	II		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
51	Long-tailed Pocupine	Hystricidae	Trichys fasciculata	I		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
52	MalayCivet	Viverridae	Viverra tangalunga	II		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
53	Malayar/Common Porcupine	Hystricidae	Hystrix brachyura	Ш		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
54	Marbled Cat	Felidae	Pardofelis marmorata	I	I	Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
55	Mask Palm Civet	Viverridae	Paguma larvata	П		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
56	Oriental Bay Owl	Tytonidae	Phodilusbadius			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	2019
57	Oriental Darter	Anhingidae	Anthracoceros malayanus			Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
58	Oriental Pied Hombill	Bucerotidae	Anthracoceros albirostris			Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
59	Otter Civet	Viverridae	Cynogale bennettii	1		Status: Endangered ver 3.1		1	EN	
60	Pig-tailed macaque	Cercopithecidae	Macaca nemestrina		1	Pop. trend: decreasing Status: Vulnerable ver 3.1			VU	
61	Plantain Squirrel	Sciuridae	Callosciurus notatus	1		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
62	Prevost's Squirrel	Sciuridae	Callosciurus prevostii	1		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
63	Purple Heron	Ardeidae	Ardea purpurea			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
64	Red Eyed Bulbul	Pycnonotidae	Pycnonotus brunneus			Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
				11		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
	Red Giant Flying Squirrel	Petauristinae	Petaurista petaurista	1		Pop. trend: decreasing Status: Least Concern ver 3.1				
66	Red Junglefowl	Phasianidae	Galus-galus			Pop.trend:decreasing	Endemic in		LC	
	Red Langur		Presbytis rubicanda			Status: Least Concern ver 3.1	Borneo		040	2019
68	Red Muntjac or Barking Deer	Cervidae	Muntiacus muntjak	III		Pop. trend: decreasing Status: Least Concern ver 3.1			LC	
69	Reticulated Python	Pythonidae	Python reticulatus	Ш		Pop. trend: decreasing Status: Near Threatened ver 3.1			LC	
70	Rhinoceros Hornbill	Bucerotidae	Buceros rhinoceros			Pop. trend: decreasing Status: Near Threatened ver 3.1			NT	
71	Rouroul	Phasianidae	Rollulus rouroul			Pop.trend:decreasing			NT	2019
72	Sabah Grey Langur	Cercopithecidae	Presbytis sabana	II		Status: Endangered ver 3.1 Pop. trend: decreasing	Endemicin Borneo		EN	2019
73	SambarDeer	Cervidae	Cervus unicolor		Ш	Status: Vulnerable ver 3.1 Pop. trend: decreasing		\checkmark	VU	
74	Short-tailed Mongoose	Herpestidae	Urva brachyurus	Ш		Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
75	Small-clawed Otter	Mustelidae	Aonyx cinerea	I	Ш	Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
76	Smooth Otter	Mustelidae	Lutra perspicillata	I		Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
77	Smooth-coated Otter	Mustelidae	Lutrogale perspicillata	II		Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
78	Sunda Giant Squirrel	Sciuridae	Ratufa affinis sandakanensis	П		Status: Near Threatened ver 3.1 Pop. trend: decreasing			NT	
79	Sunda Pangolin	Manidae	Manis Javanica	13th Feb	Ш	Status: Critically Endangered ver 3.1 Pop. trend: decreasing		1	CR	
80	Sunda Skunk/Teludu	Mephitidae	Mydaus javanensis	2010)		Status: Least Concern ver 3.1 Pop. trend: decreasing			LC	
81	Sunda Slow Loris	Lorisinae	Nycticebus menagensis	II	I	Status: Vulnerable ver 3.1 Pop. trend: decreasing			VU	
82	Thick-Spined Porcupine	Hystricidae	Hystrix crassispinis	11		Status: Least Concern ver 3.1 Pop. trend: decreasing	Endemic in Borneo		LC	
83	Thomas's Flying Squirrel	Petauristinae	Aeromys thomasi	11		Status: Least Concern ver 3.1	Endemic in Borneo		LC	
84	Tufted Ground squirrel	Sciuridae	Rheithrosciurus macrotis	1		Pop. trend: decreasing Status: Vulnerable ver 3.1	USITION		VU	
85	Water Monitor	Varanidae	Varanus salvator	Ш		Pop.trend: decreasing Status: Least Concern ver 3.1			LC	
86	Wreathed Hombill	Bucerotidae	Rhyticeros undulatus			Pop.trend: decreasing			DD	2019
87	White-crowned Shama	Muscicapidae	Copsychus stricklandii			Status: Data Deficient ver 3.1	Endemic in		DD	
88	Woodpeckersp		Unknown			Pop. trend: unknown Status: Data Deficient ver 3.1	Borneo(Bird's)		DD	
89	Yellow-throated Marten	Mustelidae		11		Pop. trend: unknown Status: Least Concern ver 3.1			LC	
09	renow-throated Marten	Musiellae	Martes flavigula			Pop. trend: decreasing				



Figure 10: Sambar Deer – Camera Trap



Figure 11: Borneon Pygmy Elephants - Camera Trap




Figure 12: Sun Bear – Camera Trap



Figure 13: Bornean-yellow muntjac – Camera Trap

3.1.4 HCV 1.4 Critical Temporal Use



Map 6: HCV 1.4 Critical Temporal Use – VJR Batu Timbang

The limestone karst in Batu Timbang FR is an important nesting site for swiftlet, bats and other troglofauna. Team management is required to monitor this critical nesting site especially on illegal harvesting of bird's nest during the management period of the project area.


Therefore, it is significant for establishement of the enforcement gate, i.e FCS Bt Timbang to monitor the entering of the bird's nest collector to Bt. Timbang cave. The FCS is manned 16 hours seven days a week by rotation of staffs. The bird's nest scalling data such as the table 1 below:

Year	Cave in Bt. Timbang (kg)			
	Tendalaw	Intan	Mabpar	
2015	85	40	0	
2016	0	0	0	
2017	20	7	0	
2018	0	0	0	
2019	0	0	0	

 Table 9 The Harvesting of Bird's Nest Scalling in FCS Bt. Timbang

Based on the bird's nest scalling data, about 152 kg were taken out from the Bt. Timbang cave to date. From the data above, in year 2015 – 2019 harvesting of bird's nest was conducted at three caves in BT. Timbang (Tendalaw cave, Intan cave, and Mabpar cave). In 2016, 2018 and 2019, the teriti had not reported any bird's nest harvesting to FCS Bt. Timbang. Apart from that, the team has conducted meetings and consultations with the Bird's Nest Collector to give cooperation. We have raised this issue during the consultation process, in attendance of the Teriti's leader.

3.2 HCV 2 LANDSCAPE-LEVEL FOREST



Map 7: Map of HCV 2 - Landscape Level Forest


Periodic patrolling and surveillance was done as elaborated in HCV 1 activities done (page 14) in this report. Biodiversity monitoring were also done with the establishment of 12 PSP plots and five wildlife monitoring methods (1. Camera Trapping; 2. Recce walk; 3. Night drive survey; 4. Gibbon call; and 5. Opportunistic sightings). The results of the monitoring were elaborated in page 18 – 23 of this report under HCV 1.2 and HCV 1.3.

Map 7 shows the current status of all Forest Reserve surrounding the Project area. All activities pertaining to the fulfillment of monitoring for HCV were all done in tandem with all the other HCV's. In conjunction with the project area as the main connector of wildlife migratory pathway to other forest reserve, the FMU management cooperate with the adjacent stakeholder that is Empayar Kejora Sdn Bhd in wildlife monitoring data sharing. Based on the data derived from camera trap deployed in 2018-2019 form Empayar Kejora Sdn Bhd are shown as Figure 14.

 Image: series of the series

APPENDIX 4: MONITORING ON WILD ANIMALS UTILISE THE INTEGRATED MOSAIC PLANTED FOREST OF EMPAYAR KEJORA SDN. BHD. (DATA DERIVED FROM CAMERA TRAP DEVICES DEPLOYED IN 2018-2019)

Figure 14: Monitoring on Wild Animals Utilise the Intergrated Mosaic Planting Forest of Empayar Kejora Sdn Bhd



Map 8: HCV 3 - Forest Areas that are in or Contain Rare Threatening or Endangered Species

3.4 HCV 4 SERVICES OF NATURE

3.4.1 HCV 4.2 Erosion Control

- To date, there is no major erosion issues regarding of areas with slopes >25° and 30 m riparian buffer within the Project Area. The only potential erosion area of slope >25° is located at Bt Timbang area near the main roadside which is heading to FCS Bt Timbang. Continuous monitoring and installation of warning signage's were conducted not only at the potential erosion area but also the 30 m riparian buffer within the project area.
- 2. Three PSP Plots were establish within the slopes >25° area. To date, no erosion issues occurs at the PSP Plots area.
- 3. The report on the Assessment of water quality in Northern Gunung Rara SFM Area is shown as below:

REPORT ON THE ASSESSMENT OF WATER QUALITY IN NORTHERN GUNUNG RARA (NGR) FOREST RESERVE 2019

by

Noor Azmizah Binti Andaman, Reuben Nilus & Jabanus Miun

INTRODUCTION


An environmental baseline sampling was carried out by Hydrology Unit of Forest Research Centre to characterize the water quality of 4 rivers, which drained thru the Northern Gunung Rara (NGR) Sustainable Forest Management project area. These rivers are Sg.Lanap, Sg. Kasuyan, Sg. Kuamut and Sg. Imbok. This assessment is part of the study component required for the Forest Management Plan for NGR project area.

LOCATION OF STUDY AREA

A total of 4 sampling points represent the project watershed and its sub-catchment areas which predominantly drain through the project site (Map 9). These sampling points are labelled W1 to W4 (Table 10). The chemical analyses and water quality classes for all parameters tested for the sampling points in the project area are listed in Table 7.

Sampling Point	Location	GPS location		Date of	Surrounding
Tome	Location	Latitude	Longitude	Sampling	Condition
W1	Sg. Lanap	04°59'28. 7"	117°08'07.1"		Secondary forest
W2	Sg. Kasuyan	04°55'19.3"	117°11'18.0"	December	Secondary forest
W3	Sg. Kuamut	04°53'45.8"	117°14'24.3"	2019	Secondary forest
W4	Sg. Imbok	04°51'04.9"	117°22'18.7"		Secondary forest

Table 10. The location of water quality sampling points in NGR FR (see Map, Figure 1).



Map 9: Locations of Water Sampling Plots in NGR SFM Project Area

RESULTS

Water Quality

The chemical analyses and water quality classes for all parameters tested for four sampling points in the project area are listed in Table 11.

Table 11. The results of chemical analyses and water quality classes for all parameter tested for sampling location W1-W4 in NGR project area . Note: Dissolved Oxygen (D0 in mg/l), Conductivity (μ S/cm), Total Dissolved Solid (TDS in mg/l), and Temperature (°C)

Sampling Location	Location	Temperature (°C)	pH Value	Dissolved Oxygen, DO (mg/l)	Conductivity (µS/cm)	Total Dissolved Solid (mg/l)
W ₁	Sg. Lanap	24.51	7.23	7.16	31	16
W ₂	Sg. Kasuyan	24.59	7.12	7.08	28	14
W ₃	Sg. Kuamut	25.39	7.09	7.08	37	19
W4	Sg. Imbok	27.00	8.01	6.79	201	100
Mini	mum	24.51	7.09	6.79	28.00	14.00
Maxi	mum	27.00	8.01	7.16	201.00	100.00
Me	ean	25.37	7.36	7.03	74.25	37.25
NWQSM*		Normal	Class I	Class I	Class I	Class I
				to		
				Class II		

* National Water Quality Standards for Malaysia


Synthesis of assessment

According to the National Water Quality Standard (NWQS) the water quality of the rivers based on the physicochemical parameters (except for dissolved oxygen in certain sampling point) fall into Class 1. The pH for all rivers generally complied with the standards set for water under Class I of the NWQSM. The acceptable limit for river water pH is 6 to 9, thus the pH for all sampling points are in an acceptable limit.

For DO it is essential for the aquatic life within the river water. A low DO level would threaten the aquatic community whereas only DO level below 2 mg/l is considered harmful for aquatic life. The DO for all the sampling points are within the range of Class I and Class II of NWQSM. The temperatures for all sampling points vary as the sampling was done from early morning until in the afternoon and it is still in normal value of NWQSM.

The concentration of TDS (mg/L) and conductivity were low indicating that there is no leaching of organic or inorganic substances from the ground. Both parameter shows Class I of NWQSM.

It is recommended that the management team to always carry out periodic inspection and monitoring at all the sampling points to prevent deterioration of the water quality. The management team should install signage at all the sampling point to prevent visitors or passer by traversing the road from dumping waste into the watercourse. Nevertheless, the river water would require conventional treatment such as boiling before it can be used for domestic consumption.



3.4.2 HCV 4.3 Barriers to Destructive Fire

Map 10: HCV 4.3 - Forest Providing Barriers to Destructive Fire

Most of southern border of NGR FMU is bordering oil palm estate. Furthermore, secondary vegetation dominates most of the peripheral area of the reserves. A 50 m band of moderate to high forest structure inside the project area that border local communities land and oil palm estate are categorised as HCV 4.3.

This corresponds to the responsibility of the team Management to protect the project area is to ensure that all the stakeholders are aware of entering the area. The team management have conducted several measures such as:

- Forest Fire Management Plan (FFMP) was made available September 2016 and was revised on June 2018. The FFMP has been submitted to SFD HQ for review and feedback has been given and submitted back to the FMU for additional information improvement.
- Main Stakeholder Consultation was made all year round to discuss about forest fire awareness to stakeholders and contractors.
- Fire drill training was conducted on the 29th of August 2019.
- Patrolling was conducted all year round to curb illegal activities such as encroachment to steep areas and riparian.
- Collecting the data from Forest Fire Danger Rating System (FFDRS) to analysis the fire hotspot. It is compulsory for every Forestry Offices to submit FFDRS report every day to HQ. The data collected from our weather station everyday from our office in Brantian, directly to the SFM Office, SFD HQ.

Year	Number of Incidents	Forest Reserve
2014	1	Abandoned Sawmills of Kebamega Sdn Bhd (Mt Magdalena FR Class 1)
2015	0	-
2016	1	Abandoned Sawmills of Smart Oracle Sdn Bhd (Gunung Rara Wildlife Corridor FR)
2017	0	-
2018	0	-
2019	0	-

Table 12: Fire Incidents

3.5 HCV 6 CULTURAL IDENTITY OF LOCAL COMMUNITIES

Bird's nest collection in the limestone caves of VJR Batu Timbang is one of the main sources of income of the nine teriti. These caves are located at 4° 58' N, 117° 05' E on a 300 m height hill of the reserve and important nesting area of the edible bird'd nest of Aerodramus fuciphaga (White Nest Swiftlet) and Aerodramus maximus (Black Nest Swiftlet). Due the presence of these species the reserve is gazette as Virgin Jungle Reserve to protect their habitat.

The only access to the caves is by river and by foot. From the mouth of Kuamut on the Kinabatangan river (Kg. Kuamut) to the mouth of the Kasuyan River takes about 1 hour 30 minutes depending on river water level.

To monitor the bird's nest collectors, the team of management have conducted several measures such as:

1. Forming Bird's Nest stakeholder consultation commitee

Consultation with bird's nest collector and forming a committee. To date, seven times of meeting and consultation with the Bird's Nest Collector were conducted. The Bird's Nest Collectors Committee was formed and produced a resolution, which was agreed between SFD (NGR SFM Project and the Teritis' of the Bird's Nest Collectors). Certain issues (i.e Ruler & Regulation, State Forest Policy, etc), without disregarding the welfare of the Teritis' of the Bird's Nest Collectors had been included in the cooperation agreement.

2. Endorsement of Certificate Of Identity (COI)

Based on directivity from Chief Conservator of Forest vide RUJ:JPHTN/PP700-1/3/21/KLT6(87) dated 02.02.2016 the nine teriti must apply the COI permit when entering the forest serve to collect the bird's nest at The VJR Bt. Timbang.

To date the Sabah Forestry Department was collected RM 130 from the COI endorse from the teriti.

3. <u>Quarterly Monitoring at VJR Bt. Timbang</u>

To ensure the teriti complied with the Rules & Regulation, State Forest Policy, the team of management has conducted Quaterly monitoring to VJR Bt Timbang Cave to make inspections. Due to the bad road condition going into the Birds Nest Cave, VJR Bt. Timbang, it is far and difficult to arrive at the top of VJR Bt. Timbang, and thus resulting in difficulties to organize and do a proper inspection more often.

4.0 Result or Outcomes for HCV Management Prescription and Effectiveness of Monitoring and Enhancement:-

High Conservation	Effectiveness of Monitoring and Enhancement
Value (HCV)	
Value (HCV) HCV 1.1 Protected Areas	Program of Patrolling such as ground patrolling, aerial surveillance, Inspection of boundaries re-brushing of NGR SFM Project and Establishment of Forest Checking Station shows that the management have manage to implement the enforcement program adequately. Nevertheless the management still must enhance the enforcement practice i.e. increase the rate of patrolling within the area and outside the project area, identify possible route of poachers coming in the area via SMART Patrolling. Furthermore the frequency of aerial surveillance must be increased and installing of proper signage along the main boundaries of NGR-SFM must be maintained. Awareness sessions (Main Stakeholder Consultations) with neighbouring logging and plantation owners should be maintained as well
	as the cooperation of patrolling and operation with outside agencies such as WWF, SWD, MCEE Tawau, DAMAI Team and YS.
HCV 1.2	(Flora): PSP plots re-enumeration was done in 2017 by Forest Research Center. Project Team did maintenance every year. Additional PSP plots are to be established in Northern Gn Rara Forest Reserve. The existing method used for PSP Plots establishment shall be maintained. 2nd round of Scientific Expedition recommended for the whole project area covering new Forest Reserve. Produce a stand-based mapping of vegetation types and the extent of their qualities as baseline reference by using remote sensing technique. The GIS application can be effective management and monitoring tools to examine and evaluate spatial-temporal processes of changes in forest quality and conditions. Permanent plots require ongoing maintenance and when left unattended for long periods of time, they become increasingly difficult to relocate, re-establish, and to undertake accurate re-measurements. The maintenance of permanent plots consists of determining the presence of center post and tree labels, including inspection of damage to the trees in the plots, and investigate its cause. (Fauna): Four initial methods were established, with another additional
	method were established (Opportunistic Sightings) on 2017. Camera trap to be added for continuous data of wildlife, addition of manpower for monitoring works should be progressively maintained and continued
HCV 1.3	(Flora) Endemic plots were maintained and marked. Three plots of endemic flora species were identified. The existing method used for PSP Plots establishment shall be maintained as well as maintaining the existing PSP Plots within the project area. 2nd round of Scientific Expedition recommended for the whole project area covering new Forest Reserve. Produce a stand-based mapping of vegetation types and the extent of their qualities as baseline reference by using remote sensing technique. The GIS application can be effective management and monitoring tools to examine

	
	and evaluate spatial-temporal processes of changes in forest quality and conditions. Permanent plots require ongoing maintenance and when left unattended for long periods of time, they become increasingly difficult to relocate, re-establish, and to undertake accurate re-measurements. The maintenance of permanent plots consists of determining the presence of center post and tree labels, including inspection of damage to the trees in the plots, and investigate its cause.
	(Fauna) Monitoring should be progressively maintained and continued.
HCV 1.4 Critical	Based on the Table 9 (page 29) in the year 2016, 2018 and 2019 no data
Temporal Use	were recorded. It is because, the teriti not reported of bird's nest harvesting
	to FCS Bt. Timbang. A meeting and consultation with the Bird's Nest Collector must be maintained and should be conducted twice a year to aware them. Based on the Bird's Nest Stakeholder Consultation that were conducted, to control the entrance of illegal individual or groups that is not from the Teriti's, a mechanism has been made to tackle the illegal entrance. The mechanism consists of (i) approval letter from Persatuan Warisan Gua Batu Timbang Kuamut, (ii) approval letter form Mahkamah Anak Negeri Kinabatangan and (iii) approval letter from Sabah Wildlife Department Kinabatangan.
HCV 2	Continuous monitoring of HCV, PSP's and Wildlife, patrolling and
	surveillance of protected areas were done and should be maintained. Consider using drone monitoring/ drone mapping for HCV, and
	establishment of SMART Monitoring also should be considered. In conjunction with the project area as the main connector of wildlife
	migratory pathway to other forest reserve, the FMU management should
	cooperate with the adjacent stakeholder that is Empayar Kejora Sdn Bhd in wildlife monitoring data sharing.
HCV 3	PSP Plot re-enumeration were done in 2017, and PSP Plots maintenance were done every year. The existing method used for PSP Plots establishment shall be maintained as well as maintaining the existing PSP Plots within the project area. Analysis of PSP data should be done to provide insight of the forest growth data in site project.
HCV 4.2	Continuous monitoring should be done progressively. Water sampling plots are to be maintained and marked. Frequency of patrolling should be increased and reported and documented.
HCV 4.3 Forest	Main Stakeholder Consultation, Fire Drill Training, aerial surveillance and
Providing	ground patrolling should be maintained and continued. To enhance Forest
Barriers to	Fire awareness program, the management must conduct the Main
Destructive Fire	Stakeholder Consultation. Apart from that, the management staff and the all stakeholder staff outside the NGR SFM project area must be involved with
	the Fire Drill training.
	A meeting and consultation with the Bird's Nest Collector must be
HCV 6. Cultural	maintained and should be conducted twice a year to aware them. Based on
Identity of Local Communities	the Bird's Nest Stakeholder Consultation that were conducted, to control the entrance of illegal individual or groups that is not from the Teriti's, a
Communities	mechanism has been made to tackle the illegal entrance. The mechanism
	consists of (i) approval letter from Persatuan Warisan Gua Batu Timbang
	Kuamut, (ii) approval letter form Mahkamah Anak Negeri Kinabatangan and
	(iii) approval letter from Sabah Wildlife Department Kinabatangan