Report under the Article 17 of the Habitats Directive Period 2007-2012

European Environment Agency *European Topic Centre on Biological Diversity*



Scilla morrisii

Annex II, IV Priority Yes

Species group Vascular plants **Regions** Mediterranean


Morris Squill (*Scilla morrisii*) is a perennial plant with white-violet flower endemic to Cyprus (Mediterranean region). The taxonomy of the species is controversial. Genetic examination indicated that the similar species, *Scilla cilicica*, which occurs in crevices of limestone rocks along the Pentadactylos range and on the Akamas peninsula, may in fact belong to the *Scilla morrisii*. This assessment is however based on occurrence of *Scilla morrisii* sensu Meikle, the rare micro-endemic confined only to the vicinity of Chrysorrogiatissa monastery which was originally included in the Annexes of the Habitats Directive as a priority species. The species grows on moist, shaded places in the Cyprus oak (*Quercus infectoria* subsp. *veneris*) forest, in open maquis or cypress forest margins, at altitude 400-980 m. The species is assessed as Critically Endangered (CR) in the EU27 Red list, however.

All four parameters were evaluated as "Favourable"; therefore the overall conservation status resulted in "Favourable" as well. On the contrary, population and habitat were evaluated "Unfavourable Inadequate" in the previous reporting period, although the actual quantities have not changed much.

Expansion of cultivation is the main pressure affecting the Morris Squill population, followed by road construction and further grazing and succession as a low importance pressures. The population is threatened by genetic depression.

Changes in overall conservation status between 2001-06 and 2007-12 report are mostly caused by different methodical approach and better data rather than real change in conservation status.

Assessment of conservation status at the European biogeographical level



See the endnote for more informationⁱ

Assessment of conservation status at the Member State level



The map shows both Conservation Status and distribution using a $10 \text{ km} \times 10 \text{ km}$ grid. Conservation status is assessed at biogeographical level. Therefore the representation in each grid cell is only illustrative.

Biogeographical region

Unfavourable - bad

Unknown

Species: Scilla morrisii

Report under the Article 17 of the Habitats Directive

MS Region	Conservation status of parameters				Cumant	Tuondin	0/ im	Draviana	Reason
	Range	Population	Habitat	Future prospects	Current CS	Trend in CS	% in region	Previous CS	for change
CY MED	FV	FV	FV	FV	FV		100.0	U1	Better data

Knowing that not all changes in conservation status between the reporting periods were genuine, Member States were asked to give the reasons for changes in conservation status. Bulgaria and Romania only joined the EU in 2007 and Greece did not report for 2007-12 so no reason is given for change for these countries. Greek data shown above is from 2001-06.

Main pressures and threats reported by Member States

Member States were asked to report the 20 most important threats and pressures using an agreed hierarchical list which can be found on the Article 17 Reference Portal. Pressures are activities which are currently having an impact on the species and threats are activities expected to have an impact in the near future. Pressures and threats were ranked in three classes 'high, medium and low importance'; the tables below only show threats and pressures classed as 'high', for some species there were less than ten threats or pressures reported as highly important.

Ten most frequently reported 'highly important' pressures

Code	Activity	Frequency			
A01	Agricultural cultivation	100			
Ten most frequently reported 'highly important' threats					
Code	Activity	Frequency			

No 'highly important' threats were reported.

Proportion of population covered by the Natura 2000 network

For species listed in the Annex II of the Directive Member States were asked to report the population size within the Natura 2000 network. The percentage of species population covered by the network was estimated by comparing the population size within the network and the total population size in the biogeographical/marine region.

Percentage of coverage by Natura 2000 sites in biogeographical/marine region

	MED
CY	100

See the endnotes for more information ii

Species: Scilla morrisii

Report under the Article 17 of the Habitats Directive

Most frequently reported conservation measures

For species listed in the Annex II of the Directive Member States were asked to report up to 20 conservation measures being implemented for this species using an agreed list which can be found on the Article 17 Reference Portal. Member States were further requested to highlight up to five most important ('highly important') measures; the table below only shows measures classed as 'high', for many species there were less than ten measures reported as highly important.

Ten most frequently reported 'highly important' conservation measures

Code	Measure	Frequency
6.1	Establish protected areas/sites	50
6.3	Legal protection of habitats and species	50

This information is derived from the Member State national reports submitted to the European Commission under Article 17 of the Habitats Directive in 2013 and covering the period 2007-2012. More detailed information, including the MS reports, is available at: http://bd.eionet.europa.eu/article17/reports2012/species/summary/? group=Vascular+plants&period=3&subject=Scilla+morrisii

Species: Scilla morrisii

Report under the Article 17 of the Habitats Directive

Assessment of conservation status at the European biogeographical level: Current Conservation Status (Current CS) shows the status for the reporting period 2007-2012, Previous Conservation Status (Previous CS) for the reporting period 2000-2006. Reason for change in conservation status between the reporting periods indicates whether the changes in the status were genuine or not genuine. Previous Conservation Status was not assessed for Steppic, Black Sea and Marine Black Sea regions. For these regions the Previous status is therefore considered as 'unknown'. The percentage of the species population occurring within the biogeographical/marine region (% in region) is calculated based on the area of GIS distribution.

iiPercentage of coverage by Natura 2000 sites in biogeographical/marine region: In some cases the population size within the Natura 2000 network has been estimated using a different methodology to the estimate of overall population size and this can lead to percentage covers greater than 100%. In such case the value has been given as 100% and highlighted with an asterisk (*). The value 'x' indicates that the Member State has not reported the species population and/or the coverage by Natura 2000. No information is available for Greece. The values are only provided for regions, in which the occurrence of the species has been reported by the Member States.