

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/248612900>

Keys to fungi on Dung

Article · March 1968

DOI: 10.1016/S0007-1528(68)80026-4

CITATIONS

58

READS

316

2 authors, including:



Michael Richardson

71 PUBLICATIONS 1,546 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:




PREDICTS project [View project](#)

Keys to Fungi on Dung

by
M. J. Richardson
and
Roy Watling

British
Mycological
Society



KEYS TO FUNGI ON DUNG

KEYS TO FUNGI
ON DUNG

by

M. J. RICHARDSON
165 Braid Road,
EDINBURGH EH10 6JE

and

ROY WATLING
Royal Botanic Garden,
EDINBURGH EH3 5LR

Published by the British Mycological Society
PO Box 30, Stourbridge
West Midlands DY9 9PZ

© British Mycological Society 1997

Printed in Scotland by BPC-AUP Aberdeen Ltd

ISBN 0 9527704 2 3

The first edition of these keys was published in the *Bulletin of the British Mycological Society* 2, 18-43 (1968) and 3, 86-88, 121-124 (1969) in an attempt to bring together in one place information for the identification of coprophilous fungi which would be useful to teachers and others interested in these fungi. They were issued as a separate publication in 1972, and with corrections in 1974. They were reprinted in 1982 with additions. This latest edition is an update of all the earlier ones, with current nomenclature and recent references, and the inclusion of some additional species.

M.J.R.

R.W.

December 1996

INTRODUCTION

Coprophilous fungi are highly satisfactory for demonstrating the diversity and morphology of a group of related organisms within an ecological system. Representative genera of most major groups of fungi can usually be guaranteed to appear on dung after a period of incubation. There is no shortage of dung in our fields and woods, and this material will always produce characteristic fungi at whatever time of year it is collected.

Dung is best incubated in a light place, for example on a table in a warm room, on layers of moist filter paper or other absorbent material. For rabbit pellets, and samples of similar size, Petri dishes are ideal; for horse 'apples', and larger types of dung, large covered dishes such as glass casseroles, plastic sandwich boxes or yoghurt pots are needed. The top third cut from a plastic lemonade or mineral water bottle fits neatly in a Petri dish, and replacing the screw cap with a cotton wool plug allows aeration and gives adequate height for developing basidiomycetes. Samples should not be kept in airtight containers for any length of time after collection, as in such conditions insects and nematodes tend to break down the dung, and anaerobic conditions which do not favour the fungi rapidly develop. If they cannot be set to incubate soon after collection they can be gently air dried, as most dung fungi will remain alive after such treatment and grow out when the sample is eventually moistened. The absorbent material should be kept moist. Although free water will not allow the best development of ascomycetes, the succession of basidiomycetes appears to vary with the wetness of the dung. Earthworms and insect larvae should be excluded from the samples as far as possible, for they break up the dung too much; activity of the latter can be reduced by spraying lightly with a household insecticide. If space is limited and cultures are kept nearby, it is very important to prevent mite infestation. Containers can be isolated by placing on glass plates lightly smeared with Vaseline, to which an acaricide (e.g. methyl benzoate) can be added.

Fungi are best sought with a stereoscopic binocular microscope, when their full beauty will be seen, but a hand lens or simple magnifier, although less convenient, is sufficient for all but the smallest forms. The larger ascomycetes and most of the basidiomycetes are readily seen with the unaided eye, but the binocular microscope is still very useful for observing the gross features of the veil of the basidiomycetes. Perithecia, apothecia and similar structures can be removed with fine needles or forceps quite cleanly for mounting, initially in water, on slides. Subsequent irrigation with iodine solution will allow any reaction of ascus wall, tip or pore to be observed, and mounting in diluted Indian ink can enhance the visibility of appendages, caudae and sheaths which occur on some spores. Spore discharge in the ascomycetes often occurs from mature asci when material is

mounted in water, so mature spores can immediately be seen. Many of the coprophilous toadstools (agarics), because of their small size and/or rapidly deliquescent nature, often do not give spore prints in the normal way, but mature spores can usually be found on the stipe or in natural spore prints formed on the absorbent material on which the dung is supported. For accurate identification the ability to measure the size of spores and other structures will be necessary. Basic microscopical technique and mycological knowledge is assumed. Common species are well described and illustrated in popular books, and references are given to specialist works to allow descriptions of less common species to be found. It will be necessary to refer to these for critical taxa. Although this edition contains about one half more species than the 1982 edition, there are still many species to be described and new records and observations to be made, especially in the Ascomycotina.

Four keys are presented. Keys 1 and 2 (MJR) are to the coprophilous ascomycetes, a very diverse group which, although not covering all the possible types of reproductive structure found in the class, contains many of the important types. The information for the identification of these fungi is dispersed throughout the literature, and many new species are still being discovered and described. Some appear to be world-wide in their distribution, others more restricted, with a prevalence of reports from either arctic, temperate or tropical regions. These keys are not exhaustive, since there are far too many species to make it practical to include them all. They do, however, include most genera, and the commoner or well known species of temperate regions. Specific (and even generic) limits in some cases (e.g. *Coprotus/Ascophanus/Ryparobius/Thelebolus*) are still the subject of debate and the choice of names to use in the key for a few taxa has been a compromise. Key 2 includes the original 'plectomycete' key (RW), which contains fungi which may not be strictly coprophilous in the normal sense, but fungi which occur on hair, horn, bone and cadavers, and may thus be found on carnivore dung or pellets of owls and other birds of prey.

Key 3 (RW, p. 52) is to the basidiomycetes of dung and associated debris. The part of the key dealing with the agarics attempts to be as complete as possible. Since the toadstools have always been thought of as the best known of the coprophilous fungi, attention to their taxonomy has often been careless. In this key the opportunity has been taken to adopt a rather narrow species concept, and to provide in certain places indications of where distinct taxa, even autonomous species, may be found after further laboratory work. Many of these types have been cultured and appear to differ vegetatively in ways which support observations of gross morphology. Coprophilous agarics are popular material for genetic studies and additional information on veil structure, spore number etc. of individual species is given, even when these are not 'key characters'.

Key 4 (MJR, p. 63) is to the Zygomycota (phycomycetes) which are characteristic of dung and amongst the first to appear when freshly dropped dung is incubated. They soon disappear, however, but their fruiting can be prolonged by plating small portions of dung on a nutrient medium (e.g. potato carrot or potato dextrose agar) to which has been added a small amount of antibiotic to reduce bacterial growth. This method is especially suitable for the parasitic and predacious fungi. A cultural approach is essential for the identification of many of these fungi and the above media, and oatmeal agar, are suitable for culture as well as isolation. For this reason the study of this group of fungi is less easy than that of the ascomycetes and basidiomycetes but, because the asexual stages are characteristic, we have attempted to key out the commoner genera which might be found, with notes on common species. The asexual spores are sporangiospores formed in sporangia; some sporangia produce a single spore within a closely fitting sporangium, and have in the past been erroneously described as conidia. A great range of sporangial structure occurs within the orders concerned. The classical structure is the massive (up to 250µm diam.) multispored sporangium with an internal columella which remains after the spores have been dispersed (e.g. *Mucor*); those of *Mortierella* are similar, but smaller and without a columella. Other sporangia are much reduced and may be only 10-20µm diam., and contain only a small number of spores (*Thamnidium*) or one spore (*Chaetocladium*); these small globose structures are termed sporangioles. Spores may also form in chains; the chains are in terminal groups and are formed by the differentiation of the contents of cylindrical sporangia which are considered to be part-sporangia (merosporangia). When the sporangial wall has disappeared the 'spore chains' may remain discrete and intact, or they may collapse into a wet droplet of spores (*Syncephalastrum*, some *Piptocephalis*). Members of the Kickxellaceae (e.g. *Coemansia*, *Kickxella*) have single spored merosporangia produced in serried ranks on boat-shaped or swollen structures (sporoclastes). The sexual spores (zygospores) are rarely seen without culturing; oatmeal agar is one which favours their production. The key includes one member of the Entomophthorales, which also produces single-spored sporangia. Other members of this order may be found parasitising the various animals which live in dung; many other predacious fungi may also be seen, e.g. parasites of amoebae (*Acaulopage*). The key is of necessity far from complete, and omits members of the Dimargaritales, which have been found frequently on dung of small mammals in America.

Mitosporic fungi ('Fungi Imperfecti') and myxomycetes have been excluded, since they would expand the range of these keys beyond what was initially intended, although numerous species of both groups occur on dung when incubated in a damp chamber. For mitosporic fungi see Seifert, Kendrick & Murase (1983) and Ellis & Ellis (1988); for myxomycetes see Eliasson & Lundqvist (1979). As practical keys, rather than a taxonomic treatment, taxonomic authorities have not been cited. For ascomycetes, Cannon, Hawksworth & Sherwood-Pike (1985) have

been followed, unless there is a more recent treatment of a group. For the basidiomycetes the 'New Checklist of British Agarics and Boleti' (Dennis, Orton & Hora, 1960, *Supplement to the Transactions of the British Mycological Society* 43) has been followed, and *The British Fungus Flora* (Orton & Watling, 1979 and Watling, 1982).

ASCOMYCETE REFERENCES

- Ahmed, S.I. & Cain, R.F. (1972). Revision of the genera *Sporormia* and *Sporormiella*. *Canadian Journal of Botany* 50, 419-477. (Keys and descriptions of 66 spp.).
- Apinis, A.E. (1964). Revision of the British Gymnoascaceae. *Mycological Paper* 96.
- Arx, J.A. von (1971). On *Arachniotus* and related genera of the Gymnoascaceae. *Persoonia* 6, 371-380.
- Arx, J.A. von (1975). Revision of *Microascus* with the description of a new species. *Persoonia* 8, 191-197.
- Arx, J.A. von (1975). On *Thielavia* and some similar genera of Ascomycetes. *Studies in Mycology* 8.
- Arx, J.A. von (1982). A key to the species of *Gelasinospora*. *Persoonia* 11, 443-449.
- Arx, J.A. von (1986). The ascomycete genus *Gymnoascus*. *Persoonia* 13, 173-183.
- Arx, J.A. von (1987). A re-evaluation of the Eurotiales. *Persoonia* 13, 273-300. (Keys to families and genera).
- Arx, J.A. von, Dreyfuss, M. & Müller, E. (1984). A re-evaluation of *Chaetomium* and the Chaetomiaceae. *Persoonia* 12, 169-179. (Key to species).
- Arx, J.A. von, Figueras, M. J. & Guarro, J. (1988). Sordariaceous Ascomycetes without Ascospore Ejaculation. *Beihefte zur Nova Hedwigia* 94, 1-104.
- Arx, J.A. von, & Gams, W. (1967). Über *Pleurage verruculosa* und die zugehörige *Cladorrhinum*-Konidienform. *Nova Hedwigia* 13, 198-208.
- Arx, J. A. von, Guarro, J. & van der Aa, H. A. (1987). *Asordaria*, a new genus of the Sordariaceae, and a new species of *Melanocarpus*. *Persoonia* 13, 263-272.
- Barrasa, J. M. & Checa, J. (1990). Dothideales del Parque Natural de Monfrague Cáceres. I. *Boletín Sociedad Micológica de Madrid* 15, 91-102.
- Barrasa, J. M., Lundqvist, N. & Moreno, G. (1986). Notes on the genus *Sordaria* in Spain. *Persoonia* 13, 83-88.
- Bell, A. & Mahoney, D.P. (1995). Coprophilous fungi in New Zealand. I. *Podospora* species with swollen agglutinated perithecial hairs. *Mycologia* 87, 375-396. (Key and descriptions of 8 spp.).
- Bezerra, J.L. & Kimbrough, J.W. (1975). The genus *Lasiobolus* (Pezizales: Ascomycetes). *Canadian Journal of Botany* 53, 1206-1229. (Key and descriptions of 11 spp.).
- Booth, C. (1961). Studies of pyrenomycetes: VI. *Thielavia* with notes on some allied genera. *Mycological Paper* 83.
- Breton, A. & Faurel, L. (1968). Etudes des affinités du genre *Mycorhynchus* Sacc. et description de plusieurs especes nouvelles. *Revue de Mycologie* 32, 229-258.
- Brummelen, J. van (1962). Studies on Discomycetes - II. On four species of *Fimaria*. *Persoonia* 2, 321-330.
- Brummelen, J. van (1962). A World Monograph of the Genera *Ascobolus* and *Saccobolus*. *Persoonia*, Supplement Volume 1. (Key and descriptions of 66 spp., and a critical taxonomic treatment).

- Brummelen, J. van (1980). Two species of *Ascobolus* new to Britain. *Persoonia* **11**, 87-92.
- Brummelen, J. van (1981). The genus *Ascodesmis* (Pezizales, Ascomycetes). *Persoonia* **11**, 333-358.
- Brummelen, J. van (1984). Notes on cup-fungi -2. *Lasiobolus*. *Persoonia* **12**, 328-334.
- Brummelen, J. van (1986). Notes on cup-fungi -3. On three species of *Cheilymenia*. *Persoonia* **13**, 89-96.
- Brummelen, J. van (1990). Notes on cup-fungi -4. On two rare species of *Ascobolus*. *Persoonia* **14**, 203-207.
- Cailleux, R. (1971). Recherches sur la mycoflore coprophile centrafricaine. Les genres *Sordaria*, *Gelasinospora*, *Bombardia* (Biologie, Morphologie, Systématique). *Bulletin trimestriel de la Société Mycologique de France* **87**, 461-626 + 27 plates.
- Cain, R.F. (1934). Studies of Coprophilous Sphaeriales in Ontario. *University of Toronto Studies, Biological Series*, No.38. (Reprinted 1968 in *Bibliotheca Mycologica*, Band 9, by Cramer, Lehre).
- Cain, R.F. (1961). Studies of coprophilous Ascomycetes. VII. *Preussia*. *Canadian Journal of Botany* **39**, 1633-1666.
- Cain, R.F. (1962). Studies of coprophilous Ascomycetes. VIII. New species of *Podospora*. *Canadian Journal of Botany* **40**, 447-490.
- Cain, R.F. & Kimbrough, J.W. (1969). *Coprobolus*, a new genus of the tribe Thelebolae (Pezizaceae). *Canadian Journal of Botany* **47**, 1911-1914.
- Cain, R.F. & Mirza, J.H. (1972). Three new species of *Arnium*. *Canadian Journal of Botany* **50**, 333-336.
- Cannon, P.F. & Hawksworth, D.L. (1982). A re-evaluation of *Melanospora* Corda and similar Pyrenomycetes, with a revision of the British species. *Botanical Journal of the Linnean Society* **84**, 115-160.
- Cannon, P.F., Hawksworth, D.L. & Sherwood-Pike, M.A. (1985). *The British Ascomycotina. An Annotated Checklist*. Commonwealth Agricultural Bureaux, Slough, U.K.
- Cano, J. & Guarro, J. (1990). The genus *Aphanoascus*. *Mycological Research* **94**, 355-377. (Key to species).
- Currah, R.S. (1988). An annotated key to the genera of the Onygenales. *Systema Ascomycetum* **7**, 1-12.
- Dennis, R.W.G. (1978). *British Ascomycetes*. J. Cramer, Lehre. (or earlier edition, 1968 and 1960 (as *British Cup Fungi and their allies*, The Ray Society, London). (All groups).
- Dissing, H. (1987). Three 4-spored *Saccobolus* species from north east Greenland. In *Arctic and Alpine Mycology II* (ed. G.A. Laursen, J.F. Ammirati & S.A. Redhead), pp. 79-86.
- Dissing, H. (1989). Four new coprophilous species of *Ascobolus* and *Saccobolus* from Greenland (Pezizales). *Opera Botanica* **100**, 43-50.
- Dissing, H. (1992). Notes on the coprophilous pyrenomycete *Sporormia fimetaria*. *Persoonia* **14**, 389-394.
- Dissing, H. & Paulsen, M.D. (1976). *Trichophaeopsis tetraspora*, a New Coprophilous Discomycete from Denmark. *Botanisk Tidsskrift* **70**, 147 - 151.
- Elliott, M.E. (1967). *Rutstroemia cuniculi*, a coprophilous species of the Sclerotiniaceae. *Canadian Journal of Botany* **45**, 521-524.
- Guarro, J. & Arx, J. A. von (1987). The Ascomycete genus *Sordaria*. *Persoonia* **13**, 301-313. (Key to 14 species and checklist).
- Hawksworth, D.L. & Webster, J. (1977). Studies on *Mycorhynchus* in Britain. *Transactions of the British Mycological Society* **68**, 329-340. (Key to 12 spp. and descriptions of some).
- Jain, K. & Cain, R.F. (1973). *Mycoarctium*, a new genus in the Thelebolaceae. *Canadian Journal of Botany* **51**, 305-307.

- Jeng, R.S., Luck-Allen, E.R. & Cain, R.F. (1977). New species and new records of *Delitschia* from Venezuela. *Canadian Journal of Botany* **55**, 383-392.
- Khan, R.S. & Cain, R.F. (1972). Five new species of *Podospora* from East Africa. *Canadian Journal of Botany* **50**, 1649-1661.
- Kimbrough, J.W. (1969). North American species of *Thecotheus* (Pezizeae, Pezizaceae). *Mycologia* **61**, 99-114. (Key and description of 5 spp.).
- Kimbrough, J.W. & Korf, R.P. (1967). A synopsis of the genera and species of the tribe Thelebolae (Pseudoascobolaceae). *American Journal of Botany* **54**, 9-23.
- Kimbrough, J.W. & Luck-Allen, E.R. (1974). *Lasiothelebolus*, a new genus of the Thelebolaceae (Pezizales). *Mycologia* **66**, 588-592.
- Kimbrough, J.W., Luck-Allen, E.R. & Cain, R.F. (1969). *Iodophanus*, the Pezizeae segregate of *Ascophanus* (Pezizales). *American Journal of Botany* **56**, 1187-1202. (Key and description of 10 spp.).
- Kimbrough, J.W., Luck-Allen, E.R. & Cain, R.F. (1972). North American species of *Coprotus* (Thelebolaceae: Pezizales). *Canadian Journal of Botany* **50**, 957-972. (Key and description of 18 spp.).
- Krug, J.C. (1973). An enlarged concept of *Trichobolus* (Thelebolaceae, Pezizales) based on a new eight-spored species. *Canadian Journal of Botany* **51**, 1497-1501. (With key to 4 spp.).
- Krug, J.C. (1995). The genus *Fimetariella*. *Canadian Journal of Botany* **73**, 1905-1916. (With key to 8 spp.).
- Krug, J.C. & Cain, R.F. (1972). Additions to the genus *Arnim*. *Canadian Journal of Botany* **50**, 367-373. (Key to 25 spp.).
- Krug, J.C. & Cain, R.F. (1974). A preliminary treatment of the genus *Podosordaria*. *Canadian Journal of Botany* **52**, 589-605. (Key and descriptions of 10 spp.).
- Krug, J.C. & Cain, R.F. (1974). New species of *Hypocopra* (Xylariaceae). *Canadian Journal of Botany* **52**, 809-843. (Descriptions and synoptic key to 30 spp.).
- Krug, J.C. & Scott, J.A. (1994). The genus *Bombardioidea*. *Canadian Journal of Botany* **72**, 1302-1310. (Description and key to 4 spp.).
- Larsen, K. (1970). The Genus *Saccobolus* in Denmark. *Botanisk Tidsskrift* **65**, 371-389.
- Larsen, K. (1971). Danish Endocoprophilous Fungi and Their Sequence of Occurrence. *Botanisk Tidsskrift* **66**, 1-32.
- Lohmeyer, T. R. & Benkert, D. (1988). *Poronia erici* - eine neue Art der Xylariales (Ascomycetes). *Zeitschrift für Mykologie* **54**, 93-102.
- Luck-Allen, E.R. & Cain, R.F. (1975). Additions to the genus *Delitschia*. *Canadian Journal of Botany* **53**, 1827-1887. (Key to 46 spp. and descriptions/illustrations of most).
- Lundqvist, N. (1967). On spore ornamentation in the Sordariaceae, exemplified by the new cleistocarpous genus *Copromyces*. *Arkiv för Botanik*, Series 2, **6**(7), 327-337.
- Lundqvist, N. (1969). *Zygopleurage* and *ZygospERMella* (Sordariaceae s. lat., Pyrenomycetes). *Botaniska Notiser* **122**, 353-374.
- Lundqvist, N. (1970). New Podosporae (Sordariaceae s. lat., Pyrenomycetes). *Svensk Botanisk Tidsskrift* **64**, 409-420.
- Lundqvist, N. (1972). Nordic Sordariaceae s. lat. *Symbolae Botanicae Upsalienses* **XX.1**, 1-314. (Keys and descriptions of ca 100 spp., and critical taxonomic discussion).
- Lundqvist, N. (1980). On the genus *Pyxidiophora* sensu lato (Pyrenomycetes). *Botaniska Notiser* **133**, 121-144.
- Lundqvist, N. (1980). *Wawelia effusa* Lundqvist, spec. nov. (Xylariaceae). *Persoonia* **14**, 417-423.
- Malloch, D. & Cain, R.F. (1970). The genus *Arachnomyces*. *Canadian Journal of Botany* **48**, 839-845.

- Malloch, D. & Cain, R.F. (1970). Five new genera in the new family of Pseudeurotiaceae. *Canadian Journal of Botany* **48**, 1815-1825.
- Malloch, D. & Cain, R.F. (1971). New genera of the Onygenaceae. *Canadian Journal of Botany* **49**, 839-846.
- Malloch, D. & Cain, R.F. (1971). Four new genera of cleistothecial Ascomycetes with hyaline ascospores. *Canadian Journal of Botany* **49**, 847-854.
- Malloch, D. & Cain, R.F. (1971). New cleistothecial Sordariaceae and a new family, Coniochaetaceae. *Canadian Journal of Botany* **49**, 869-880.
- Malloch, D. & Cain, R.F. (1972). New species and combinations of cleistothecial Ascomycetes. *Canadian Journal of Botany* **50**, 61-72.
- Minter, D.W. & Webster, J. (1983). *Wavelia octospora* sp. nov., a xerophilous and coprophilous member of the Xylariaceae. *Transactions of the British Mycological Society* **80**, 370-373.
- Mirza, J.H. & Cain, R.F. (1969). Revision of the genus *Podospora*. *Canadian Journal of Botany* **47**, 1999-2048.
- Moravec, J. (1990). A taxonomic revision of the genus *Cheilymenia* - 3. A new generic and infrageneric classification of *Cheilymenia* in a new emendation. *Mycotaxon* **38**, 459-484. (Synopsis of genus, including *Coprobia*).
- Moravec, J. (1993). A taxonomic revision of the genus *Cheilymenia* - 5. The section *Cheilymenia*. *Czech Mycology* **47**, 7-37.
- Moreau, C. (1953). Les Genres *Sordaria* et *Pleurage*. *Encyclopédie mycologique* **25**, 1-330. (*Sordaria* and *Pleurage* (= *Podospora*/*Schizothecium*), and *Coniochaeta*, *Hypocopra*, *Sporormiella*, *Trichodelitschia*, and other pyrenomycetes for comparison).
- Munk, A. (1957). Danish Pyrenomycetes. *Dansk Botanisk Arkiv* **17**(1), 1-491.
- Orr, G.F. & Kuehn, H.H. (1971). Notes on Gymnoascaceae. I. A review of eight species. *Mycologia* **63**, 191-203.
- Orr, G.F., Kuehn, H.H. & Plunkett, O.A. (1963). A new genus of the Gymnoascaceae with swollen peridial septa. *Canadian Journal of Botany* **41**, 1439-1456. (Key to *Auxarthron* (*Gymnoascus*) species).
- Orr, G.F., Kuehn, H.H. & Plunkett, O.A. (1971). The genus *Myxotrichum* Kunze. *Canadian Journal of Botany* **41**, 1457-1480. (Key to species).
- Paulsen, M. D. & Dissing, H. (1979). The genus *Ascobolus* in Denmark. *Botanisk Tidsskrift* **74**, 67-78.
- Rehm, H. (1887-1895). Ascomyceten: Hysteriaceen und Discomyceten. Vol. 1, Abt. 3 of *Rabenhorst's Kryptogamen-Flora*. (Discomycetes).
- Renny, J. (1874). New species of the genus *Ascobolus*. *Journal of Botany* **12**, 353-357 and 4 plates. (Description and illustration of 6 *Ascozonus* spp.).
- Richardson, M.J. (1972). Coprophilous ascomycetes on different dung types. *Transactions of the British Mycological Society* **58**, 37-48.
- Samson, R.A. (1972). Notes on *Pseudogymnoascus*, *Gymnoascus* and related genera. *Acta botanica neerlandica* **21**, 517-527.
- Seth, H.K. (1970). The genus *Lophotrichus* Benjamin. *Nova Hedwigia* **19**, 591-599.
- Valldosera, M. & Guarro, J. (1987). Estudios sobre hongos coprófilos aislados en España. VI. Ascomycetes. *Boletín Sociedad Micológica de Madrid* **12**, 51-56.
- Valldosera, M. & Guarro, J. (1988). Some coprophilous ascomycetes from Chile. *Transactions of the British Mycological Society* **90**, 601-605.
- Valldosera, M. & Guarro, J. (1989). Estudios sobre hongos coprófilos aislados en España. XI. Ascomycetes. *Boletín Sociedad Micológica de Madrid* **14**, 75-80.
- Valldosera, M. & Guarro, J. (1989). Estudios sobre hongos coprófilos aislados en España. XV. El género *Preussia* (*Sporormiella*). *Boletín Sociedad Micológica de Madrid* **14**, 81-94.

- Valldosera, M. & Guarro, J. (1992). Estudios sobre hongos coprófilos en España. XVII. Ascomycotina. *Boletín Sociedad Micológica de Madrid* **17**, 19-37.
- Valldosera, M. & Guarro, J. (1992). Estudios sobre hongos coprófilos aislados en España. XVIII. Bibliographic catalogue of Ascomycotina. *Boletín Sociedad Micológica de Madrid* **17**, 39-55.
- Valldosera, M., Guarro, J. & Figueras, M.J. (1991). Two interesting coprophilous fungi from Spain. *Mycological Research* **95**, 243-246.
- Winter, G. (1884-1887). Ascomyceten: Gymnoasceen und Pyrenomyceten. Vol. 1, Abt. 2 of *Rabenhorst's Kryptogamen-Flora*. (Pyrenomycetes).
- Yao, Y-J. (1996). Notes on British species of *Lasiobolus*. *Mycological Research* **100**, 737-739.
- Yao, Y-J. & Spooner, B.M. (1996). Notes on British species of *Cheilymenia*. *Mycological Research* **100**, 361-367.

BASIDIOMYCETE REFERENCES

- Moser, M. (1978). in Gams, H. (ed.). *Kleine Kryptogamenflora von Mitteleuropa*. Fischer Verlag.
- Moser, M. (1983). *Keys to Agarics and Boleti* (English translation by S. Plant). Roger Phillips, London.
- Orton, P.D. & Watling, R. (1979). *British Fungus Flora: Coprinus*. Her Majesty's Stationery Office, Edinburgh.
- Phillips, R. (1981). *Mushrooms and other fungi of Great Britain and Europe*. Pan Books, London.
- Watling, R. (1982). *British Fungus Flora: Bolbitiaceae*. Her Majesty's Stationery Office, Edinburgh.

PHYCOMYCETE REFERENCES

- Benjamin, R.K. (1959). The merosporangiferous Mucorales. *Aliso* **4**, 321-433.
- Benjamin, R.K. (1961). Addenda to the merosporangiferous Mucorales. *Aliso* **5**, 11-19.
- Benjamin, R.K. (1963). Addenda to the merosporangiferous Mucorales. *Aliso* **5**, 273-288.
- Benjamin, R.K. (1965). Addenda to the merosporangiferous Mucorales. *Aliso* **6**, 1-10.
(The 4 papers above are an excellent account of *Syncephalis*, *Piptocephalis*, *Coemansia* and other unusual allied phycomycetes, republished (1967) as *Bibliotheca Mycologica* **5** by J. Cramer, Lehre).
- Gams, W. & Moreau, R. (1959). Le genre *Mortierella*. *Annales scientifiques de l'Université de Besançon*, Series 2 **3**, 95-105.
- Hesseltine, C.W. (1955). Genera of Mucorales with a note on their synonymy. *Mycologia* **47**, 344-363. (With good key; many other papers by Hesseltine, with others, in *Mycologia*, *American Journal of Botany*, *American Midland Naturalist* and *Lloydia*).
- Ingold, C.T. & Zoberi, M.H. (1963). The asexual apparatus of Mucorales in relation to spore liberation. *Transactions of the British Mycological Society* **46**, 115-134.
- Naumov, N.A. (1939). Clés des Mucorinées. *Encyclopédie mycologique* **9**, 1-137.
- Zycha, H., Siepmann, R. & Linneman, G. (1969). *Mucorales*. J. Cramer, Lehre. (A revision of Zycha, 1935).

GENERAL REFERENCES

- Bell, A. (1983). *Dung Fungi: an illustrated guide to coprophilous fungi in New Zealand*. Victoria University Press, Wellington.
- Bon, M. (1987). *The Mushrooms and Toadstools of Britain and North-western Europe*. Hodder & Stoughton, London.
- Cacialli, G., Caroti, V. & Doveri, F. (1995). *Funghi fimicoli e rari o interssanti del litorale Toscano*. Schede di Micologia vol. 1. Fondazione Centro Studi Micologici Dell' A.M.B., Vicenza, Italy.
- Domsch, K.H., Gams, W. & Anderson, T.H. (1980). *Compendium of soil fungi*. Academic Press, New York.
- Ellis, M.B. & Ellis, J.P. (1988). *Microfungi on Miscellaneous Substrates*. Croom Helm, London & Sydney.
- Gilman, J.C. (1957). *A Manual of Soil Fungi*. Iowa State College Press.
- Eliasson, U. & Lundqvist, N. (1979). Fimicolous Myxomycetes. *Botaniska Notiser* 132, 551-568. (A list of 34 spp., with some descriptions and illustrations).
- Hawksworth, D.L., Kirk, P.M., Sutton, B.C. & Pegler, D.N. (1995). *Ainsworth & Bisby's Dictionary of the Fungi*. 8th. edn. CAB International, Wallingford.
- Holden, M. (ed.) (1982). Guide to the literature for the identification of British fungi, 4th Edition. *Bulletin of the British Mycological Society* 16, 36-55; 92-112.
- Massee, G., & Salmon, E.S. (1901). Researches on coprophilous fungi. *Annals of Botany, London* 15, 313-357.
- Seifert, K.A., Kendrick, W.B. & Murase, G. (1983). *A key to hyphomycetes on dung*. University of Waterloo Biology Series No. 27.
- Webster, J. (1970). Coprophilous Fungi. *Transactions of the British Mycological Society* 54, 161-180.

Key 1. Ascomycota

- 1 Ascoma either globose to flask shaped, usually with an easily observable pore or neck (**perithecium** or **pseudothecium**, figs 16, 18, 19, 22, 27, 30, 32, 34-37), or discoid (**apothecium**, figs 1, 3, 4, 7, 11-14). Spores usually 8 in each ascus (less frequently 4, 16, 32, 64, 128 etc.). Asci ellipsoid to cylindrical, borne in a distinct hymenium, thus appearing in fascicles or distinct groups when the fruit body is squashed. 2
- Ascoma globose to subglobose, lacking a definite pore or neck (**cleistothecium** or **gymnothecium**, figs 38, 39, 46). Asci globose to subglobose, 8-spored, not in a distinct hymenium, appearing quite free when the fruit body is squashed. **Key 2, 148** (p. 45)
- 2(1) Ascoma a **perithecium** or **pseudothecium**, usually dark in some part, not opening to a disc but remaining globose or flask shaped. Asci unitunicate, not operculate but often with an apical pore, which may stain blue in iodine, or bitunicate. **Key 2, 1** (p. 24)
- Ascoma an **apothecium**, white or lightly coloured, soft fleshed, opening out to a disc or cushion shape when mature. Asci unitunicate. 3
- 3(2) Asci opening by an operculum (fig. 8), a bilabiate vertical split down to a subapical ring of thickening (fig. 15), or apparently just bursting. 4
- Asci inoperculate, with an apical pore. 96
- 4(3) Spores 8 (occasionally 4) in an ascus, colourless, purple or brown. 5
- Spores more than 8 in an ascus, colourless. 77
- 5(4) Spores remaining colourless. 6
- Spores purple or brown at maturity. 39
- 6(5) Apothecia with obvious hairs. 7
- Apothecia without obvious hairs (microscopic hairs up to 50µm long may be present). 14
- 7(6) Hairs brown. Apothecia orange, red orange or yellow orange. (*Cheilymenia*, fig. 1) 8
- Hairs colourless. Apothecia colourless or pinkish. (*Lasiobolus*, fig. 3) 12

- 8(7) Apothecia with stellate hairs. Spores $14-20 \times 8-11 \mu\text{m}$.
Cheilymenia stercorea (figs 1, 2)
- Apothecia without stellate hairs. 9
- 9(8) Spores $14.5-18 \times 8-9.5 \mu\text{m}$. Asci $10-13 \mu\text{m}$ diam. Apothecia 2mm
 diam. or more. *Cheilymenia coprinaria*
- Spores larger, $17 \times 10 \mu\text{m}$ or more. 10
- 10(9) Apothecia reddish orange, up to 1mm diam., marginal hairs rooting,
 wall $2-4 \mu\text{m}$ thick. Spores $21-26 \times 10-13.8 \mu\text{m}$. *Cheilymenia fimicola*
- Apothecia pale orange yellow, marginal hairs superficial, wall up
 to $2 \mu\text{m}$ thick. 11
- 11(10) Asci up to $22 \mu\text{m}$ diam. Spores $17-27 \times 10-14.5 \mu\text{m}$.
Cheilymenia pulcherrima
- Asci wider, $25 \mu\text{m}$ diam. or more. Spores $23-26.5 \times 13-16.5 \mu\text{m}$.
Cheilymenia raripila




Fig. 1. *Cheilymenia stercorea*, apothecium. Fig. 2. *C. stercorea*, stellate and rooted hairs.
 Fig. 3. *Lasiobolus ciliatus*, apothecium. Fig. 4. *Iodophanus carneus*, apothecium and spore.

- 12(7) Hairs 600 μ m or longer. Spores 19-23 \times 7-10 μ m. *Lasiobolus macrotrichus*
 - Hairs shorter, up to 600 μ m. 13
- 13(12) Asci clavate, 20 μ m diam. or wider. Spores 19-22 \times 10.5-13.5 μ m. *Lasiobolus cuniculi*
 - Asci cylindrical, up to 20 μ m diam. Spores 18-22.5 \times 9.5-11.5 μ m. *Lasiobolus ciliatus* (fig. 3)
- 14(6) Asci blue in iodine solution. 15
 - Asci not blue in iodine. 24
- 15(14) Spores large, 30-42 \times 15-18 μ m, warted, ellipsoid with acute apices. *Thecotheus cinereus*
 - Spores smaller, smooth or only finely ornamented. 16
- 16(15) Apothecia brownish, large, 1cm diam. or more. (*Peziza*) 21
 - Apothecia pale, up to 4mm diam. Asci protruding from hymenium when ripe. 17
- 17(16) Apothecia white to pink, up to 2mm diam. Spores finely verruculose, 18-25 \times 8-14 μ m. *Iodophanus carneus* (fig. 4)
 - Apothecia pale, variously coloured when fresh, but drying darker. Spores smooth. (*Thecotheus*) 18
- 18(17) Spores apiculate at each end, smooth. 19
 - Spores not apiculate, 20-22 \times 8-10 μ m. *Thecotheus agranulosus*
- 19(18) Spores with a collar at the base of the apiculus. 20
 - Spores without a collar at the base of the apiculus, 16-21 \times 8-12 μ m. *Thecotheus apiculatus*
- 20(19) Apothecia white. Spores 20-22 \times 10-12 μ m, apiculus 4-6 μ m diam. *Thecotheus perplexans*
 - Apothecia yellowish. Spores 12-15 \times 7.5-9 μ m, apiculus 2.5-3.5 μ m diam. *Thecotheus africanus*
- 21(16) Spores smooth, without guttules. 22
 - Spores verruculose or spinulose, 15-18 \times 8-9 μ m, with 1 guttule. Paraphyses with clavate apices, with brown contents. Apothecia asymmetrical, extended on one side. *Peziza pleurota*

- 22(21) Spores 19-24 × 10.5-14µm. Apothecia yellowish brown,
up to 10cm diam. *Peziza vesiculosa*
- Spores up to 10µm wide. 23
- 23(22) Apothecia ca 1cm diam., umber with a paler margin. Spores
15-22 × 9-10µm. *Peziza bovina*
- Apothecia up to 2 cm diam., pale brown. Spores 13-16 × 7-9µm.
Peziza fimeti
- 24(14) Apothecia robust, up to 4mm diam., orange or with brownish
or purple tints. 25
- Apothecia smaller, rarely more than 1mm, pale, yellowish
green, orange, grey or chestnut. 32
- 25(24) Apothecia orange or red. 26
- Apothecia discrete, brownish or purple. (*Fimaria*) 27
- 26(25) Apothecia crowded, 1-3mm diam., orange, with a granular surface.
Asci up to 190 × 15µm. Spores 15-18.5 × 7-9.5µm. Paraphyses
strongly clavate to apex up to 14µm diam., filled with orange
granules. *Coprobria granulata*
- Apothecia discrete, 1-2mm diam., orange or red. Asci 240 × 10-12µm.
Spores 12-15 × 7-8µm. Paraphyses yellow, only slightly swollen from
2µm to 3-4µm at apex. *Ascophanus bresadolae*
- 27(25) Spores 8-9.5 × 4-4.5µm. *Fimaria equina*
- Spores larger. 28
- 28(27) Spores 20-38 × 10-13µm. *Fimaria hepatica*
- Spores shorter. 29
- 29(28) Spores 10-13 × 7-9µm. *Fimaria porcina*
- Spores 13-17 × 7-11µm. 30
- 30(29) Disc punctate with asci. Paraphysis tips swollen up to 3-5µm.
Spores 14.5-16 × 9.5-11µm. *Fimaria leporum*
- Disc not punctate with asci. Paraphysis tips not or only slightly
swollen. 31
- 31(30) Apothecia pale yellowish. Spores 13-15.5 × 7.5-8.5µm.
Fimaria theioleuca
- Apothecia chestnut/purplish brown. Spores 14-17 × 7-8.5µm.
Fimaria cervaria

- 32(24) Spores less than 10 μ m long. 33
 - Spores mostly longer than 10 μ m. 36
- 33(32) Paraphyses markedly capitate to 5-6 μ m, with yellowish green contents. Apothecia dull at first, yellowish at maturity. Spores 7-10 \times 2-4.5 μ m. *Thelebolus microsporus* (fig. 5)
 - Paraphyses only slightly inflated above, without coloured contents. Apothecia whitish or grey. 34
- 34(33) Spores 5-7 \times 3-4 μ m. Asci 38-42 \times 6-7 μ m. Apothecia smoky grey, 0.3-0.4mm diam. *Ascophanus cinerellus*
 - Spores larger. Apothecia pale, white or yellowish. 35
- 35(34) Apothecia up to 1.2mm diam. Asci short stalked, 40-55 \times 8-12 μ m. Spores 7.5-9 \times 4.5-5.5 μ m. *Coprotus glaucellus*
 - Apothecia 0.2-0.5mm diam. Asci attenuate below, 65-85 \times 10-15 μ m. Spores 8-10 \times 5-6.5 μ m. *Coprotus lacteus*
- 36(32) Apothecia chestnut brown up to 1mm diam. Asci 160 \times 13 μ m. Spores 13-16 \times 8-11 μ m. Paraphyses forked, with swollen tips. *Ascophanus misturæ*
 - Apothecia lighter coloured. Asci less than 150 μ m long. 37
- 37(36) Spores 14-18 \times 9-11 μ m. Apothecia pale yellow/orange, up to 1.5mm diam. Asci cylindrical, 110-150 \times 12-15 μ m. Paraphyses yellowish, slightly inflated to 4-5 μ m at apices. *Coprotus ochraceus*
 - Spores less than 15 μ m long. Apothecia up to 0.6mm diam. Asci less than 100 μ m long. 38




Fig. 5. *Thelebolus microsporus*, ascus and paraphysis. **Fig. 6.** *Ascodesmis microscopica*, ascospores.

- 38(37) Apothecia bright yellow. Asci cylindrical clavate, attenuate below, 65-90 × 10-15µm. Spores 12-14 × 6-8.5µm. Paraphyses branched, apices inflated to 4-5µm, with yellow contents. *Coprotus aurorus*
- Apothecia white/pale yellow, with darker margin. Asci broadly clavate, stalked below 40-55 × 15-30µm. Spores 9-15 × 6.5-9.5µm. Paraphyses inflated above to 5-8µm, hyaline. *Coprotus granuliformis*
- 39(5) Spores spherical or broadly ellipsoid, brown, ornamented with warts, anastomosing ridges or a reticulum. Asci clavate. Apothecium without excipulum. *(Ascodesmis, fig. 6)* 40
- Spores ellipsoid or spherical, hyaline at first, then purple, becoming brown at maturity; episporium smooth, finely verruculose, warted or cracked. Asci cylindrical. Excipulum present. 45
- 40(39) Spores 18-21.5 × 13.5-17.5µm. *Ascodesmis macrospora*
- Spores up to 16µm. 41
- 41(40) Spores ± spherical, L/B ratio mostly up to 1.2. 42
- Spores ± broadly ellipsoidal, L/B ratio mostly 1.2 or more. 43
- 42(41) Spores ornamented with round warts, 8.5-11 × 8.3-10µm. *Ascodesmis nana*
- Spores ornamented with a network of ridges, 10.5-14 × 9-12µm. *Ascodesmis sphaerospora*
- 43(41) Spores with a prominent reticulum of ridges (fig. 6), 11-15.5 × 8-13.5µm. Apothecia 150-300µm diam. *Ascodesmis microscopica* (fig. 6)
- Spore ornament not a reticulum. 44
- 44(43) Spores with 1 simple or branched ridge and isolated or occasionally connected warts, 11-14.5 × 7-11.5µm. Apothecia up to 500µm diam. *Ascodesmis porcina*
- Spores with isolated warts, some joined to form short ridges, but not a reticulum, often capitate, 9.5-12.5 × 7.5-10µm. Apothecia 50-150µm diam. *Ascodesmis nigricans*
- 45(39) Spores separate in the ascus. *(Ascobolus)* 46
- Spores firmly joined together, both in the ascus and after ejection (fig. 10). *(Saccobolus)* 66
- 46(45) Spores spherical. 47
- Spores ellipsoid. 48

- 47(46) Spores 10.5-13.5 μ m, epispore with numerous but isolated warts.
Ascobolus brassicae (figs 8, 9)
- Spores 11.5-13.5(15) μ m, epispore with subparallel occasionally anastomosing lines. *Ascobolus crosslandii*
- 48(46) Spores very large, mostly 50-70 \times 25-35 μ m, almost oblong with rounded ends, typically with few cracks in the epispore.
Ascobolus immersus (figs 7, 9)
- Spores smaller, with epispore smooth, warted or with cracks. 49
- 49(48) Epispore strongly and irregularly wrinkled with a vesiculose layer of pigment, 11.6-16 \times 6.5-9.3 μ m. Paraphyses capitate up to 18 μ m. Apothecia up to 0.6mm diam. *Ascobolus rhytidiosporus*
- Epispore not strongly wrinkled/vesiculose. 50
- 50(49) Epispore basically smooth or warted, perhaps with a few irregular cracks. 51
- Epispore with a clear pattern of cracks or lines. 56




Fig. 7. Apothecia of, from left, *Ascobolus furfuraceus*, *A. immersus* and *A. albidus*.
 Fig. 8. *A. brassicae*, ascus with spores and detail of operculum. Fig. 9. Ascospores of, clockwise from left, *A. immersus*, *A. stictoideus*, *A. albidus*, *A. brassicae* and *A. crenulatus*.

- 51(50) Spores up to 25 μ m long. 52
 - Spores longer, 25 μ m or more. 54
- 52(51) Epispore smooth, finely granular or punctate. Gelatinous material
 unilateral, not surrounding spore. 53
 - Epispore warted, spores 18.5-21(22.5) \times (9)10-11.5 μ m,
 surrounded by gelatinous sheath. *Ascobolus hawaiiensis*
- 53(52) Spores 18-24 \times 10-13 μ m. Hymenial mucus greenish yellow.
 Excipulum not brown. *Ascobolus mancus*
 - Spores 20-25 \times 11-13 μ m. Hymenial mucus sulphur yellow.
 Excipulum with rich brown intercellular pigment. *Ascobolus boudieri*
- 54(51) Epispore smooth or finely granular, spores 23-29(32) \times 12-17 μ m.
Ascobolus elegans
 - Epispore warted. 55
- 55(54) Spores with a regular pattern of warts and intact epispore,
 26-32 \times 15-17.5 μ m. *Ascobolus stictoideus* (fig. 9)
 - Spores with irregular patches of thicker pigment, especially
 at the poles, 28-35 \times 16-18 μ m. *Ascobolus degluptus*
- 56(50) Spores mostly 18 \times 10 μ m or larger. 57
 - Spores mostly smaller than 20 \times 10 μ m. 61
- 57(56) Apothecia small, mostly up to 1mm diam., colourless. Spores 20-35
 \times 11-14 μ m, epispore cracks distant, irregular, often anastomosing.
Ascobolus albidus (figs 7, 9)
 - Apothecia larger, usually 1mm diam. or more, disc yellowish,
 greenish, purplish or brownish. 58
- 58(57) Apothecia crowded, purplish or purplish brown with intercellular
 pigment. Spores 18-28 \times 10-12 μ m, with longitudinal anastomosing
 cracks. *Ascobolus roseopurpurascens*
 - Apothecia yellowish or greenish. 59
- 59(58) Spores 17-22 \times 9.5-12 μ m with a few widely spaced and irregularly
 oriented cracks. *Ascobolus michaudii*
 - Spores with closely spaced, \pm longitudinal, cracks, with varying
 degrees of anastomosis. 60

- 60(59) Apothecia furfuraceous, sessile. Ascus wall blue in iodine.
Spores 19-28 × 10-14µm. *Ascobolus furfuraceus* (fig. 7)
- Apothecia smooth, substipitate. Ascus wall only faintly
blue in iodine. Spores 19-22 × 9.5-13µm. *Ascobolus perplexans*
- 61(56) Apothecia large, stipitate, 5-10mm diam. Spores 16-19.5 × 8.5-10µm,
with subparallel, longitudinal, only rarely anastomosing lines.
Ascobolus lignatilis
- Apothecia up to 2mm diam. 62
- 62(61) Apothecia white. 63
- Apothecia yellow, green or brownish. 64
- 63(62) Spores 13-17 × 7.5-8.5µm, with a coarse reticulum of fine cracks
when mature. Only recorded on grouse, capercaillie etc. (Tetraonidae)
dung. *Ascobolus carletonii*
- Spores 16-20 × 8-10µm, with a pattern of longitudinal anastomosing
cracks. Only recorded on deer dung. *Ascobolus sacchariferus*
- 64(62) Spores 14.5-16 × 8-9µm, episporic lines not densely crowded.
Ascobolus cervinus
- Spores smaller, episporic with densely crowded, rarely anastomosing
cracks. 65
- 65(64) Apothecia greenish yellow, furfuraceous, with crenulate margin.
Spores 9.5-15 × 6-8µm. *Ascobolus crenulatus* (fig. 9)
- Apothecia brownish yellow to brown, smooth, with undifferentiated
margin. Spores 12.5-14.5 × 7-8.5µm. *Ascobolus minutus*
- 66(45) Asci 4-spored. Spore clusters 42-58 × 14-20µm. Spores 16.5-23 ×
9.5-12µm, smooth to finely punctate, but with a thick cap or girdle of
reticulated or warted pigment. *Saccobolus quadrisporus*
- Asci 8-spored. 67
- 67(66) Spore clusters ± globular, 17-26(39) × 15-20µm. 68
- Spore clusters elongated, 2-3 times as long as wide. 69
- 68(67) Spore clusters compact, subglobose, with only the exposed surface of
spores pigmented, ornamented with small and coarse warts.
Saccobolus dilutellus
- Spores loosely united in cluster, ornamented with small isolated
warts
covering most of their surface. *Saccobolus globuliferellus*

- 69(67) Apothecia yellow. Spores in 4 rows of 2 longitudinally arranged spores (fig. 10). 70
 - Apothecia hyaline or violaceous (some mature darker). Spores in 2 rows of 3 and 1 row of 2 (fig. 10). 73
- 70(69) Spore clusters 40 μ m or longer. 71
 - Spore clusters up to 40 μ m long. 72
- 71(70) Spore clusters 50-71 \times 16-25 μ m. Spores 22-29 \times 8.5-14.5 μ m, smooth or rarely finely punctate, with distant irregular cracks. *Saccobolus glaber* (fig. 10)
 - Spore clusters 43-51 \times 14-17 μ m. Spores 16-22 \times 7.5-9 μ m, with fine isolated warts. *Saccobolus citrinus*
- 72(70) Spores 14-17.5(19.5) \times 7.5-8.5(10) μ m, easily separated at maturity. Spore clusters becoming shorter and more rounded with maturity. Apothecia up to 300 μ m diam., inconspicuous due to their solitary nature and the predominantly brownish colour due to the mature spores. *Saccobolus truncatus* (fig. 10)
 - Spores 11.5-13.5 \times 5.5-6.5 μ m. *Saccobolus minimus*
- 73(69) Apothecia white, covered with tapering squamules composed of septate hyphae. Spore clusters 38-43 \times 15-17 μ m. Spores 16-17.5 \times 7-8.5 μ m, smooth or finely punctate. *Saccobolus caesariatus*
 - Apothecia not white, without tapering scales. 74
- 74(73) Spore clusters mostly over 40 μ m long. 75
 - Spore clusters mostly under 40 μ m long. 76
- 75(74) Spore clusters 38-62 \times 14-19 μ m. Spores 13-21.5 \times 6.5-9.5 μ m, smooth, finely warted or with reticulate cracks. Apothecia 0.2-2mm diam. *Saccobolus versicolor* (fig. 10)
 - Spore clusters 42-60 \times 18-24 μ m. Spores very coarsely warted, 17.5-23 \times 8.5-10 μ m (inc. warts). *Saccobolus beckii*




Fig. 10. Spore clusters of, from left, *Saccobolus versicolor*, *S. glaber* and *S. truncatus*.

- 76(74) Spore clusters compact, 26-43 × 13-19µm. Spores 13.5-18 × 7.5-9.5µm, epispore with fine or coarse warts. Apothecia 0.3-0.8mm diam. *Saccobolus obscurus*
- Spore clusters elongated, 28-37 × 10-13µm. Spores 10-14.5 × 5-7.5µm, epispore smooth or very finely granular. Apothecia 0.1-0.3mm diam. *Saccobolus depauperatus*
- 77(4) Asci operculate or bursting, without a subapical ring. Spores ellipsoid. 78
- Apothecia white, often minutely hairy at the margin. Ascus dehiscing by a vertical slit; the slit is prevented from running right down the ascus by a subapical ring of thickening. Spores ellipsoid-fusiform. (*Ascozonus*, figs 14, 15) 90
- 78(77) Asci 16-spored. Spores ellipsoid, 11-16 × 7-10µm. *Coprotus sexdecemsporus*
- Asci more than 16-spored. 79
- 79(78) Asci 32-spored. 80
- Asci more than 32-spored. 84
- 80(79) Asci very large, nearly 0.5mm long, spores 30-35 × 13-17µm (32-40 × 20-24µm in Kimbrough, 1969). Apothecia pale coloured. *Thecotheus pelletieri*
- Asci and spores smaller. 81
- 81(80) Spores 10µm or longer. 83
- Spores up to 10µm long. 82
- 82(81) Spores ellipsoid, with minute scattered warts visible under oil-immersion, 7-9 × 4-4.5µm. Apothecia densely crowded, 90-120µm diam., with 8-13 asci. Asci 32-55 × 16-18µm with (24-)32 spores. Paraphyses 1.5-2µm, clavate to 4-4.5µm. *Thelebolus caninus*
- Spores subacute at apices, ca 6 × 4µm (described as 'minute'; this value is suggested by Boudier's comparison with *R. dubius*, for which measurements are given). Apothecia densely crowded, tawny yellowish-brown. *Ryparobius brunneus*
- 83(81) Spores 10-12.5 × 5-7.5µm. Asci clavate, 75-100 × 20-30µm. Paraphyses enlarged to 6µm at apex. *Coprotus albidus*
- Spores 13.5-17.5 × 7-8µm. Asci 10-15 per apothecium, 120-175 × 50-75µm. Paraphyses filiform. *Coprotus ryparobioides*

- 84(79) Asci with up to 64 spores. 85
 - Asci with many more than 64 spores - impractical to count. 86
- 85(84) Asci 64-spored, broad clavate with short stalk, 80-130 ×
 30-60µm. Spores 8-12 × 4-7µm. *Coprotus niveus*
 - Asci broadly clavate with up to 64 spores, 60-100 × 20-30µm.
 Spores 7-10 × 4.5-5.5µm. Apothecia superficial, on the surface of
 the substrate, yellowish brown, gregarious, united into a crust.
Thelebolus crustaceus
- 86(84) Apothecia superficial, 400-600µm diam., with prominent, acuminate,
 superficial, 1-2-septate hairs, 80-190µm long, often roughened towards
 their apex, with one 1000+-spored ascus, 110-240 × 15-27µm. Spores
 very variable, 6.5-16 × 3.7-8.8µm (mostly 7.5-13 × 4.5-7µm).
Lasiobolus monascus
 - Apothecia minute, rarely above 350µm diam., globose and
 immersed in substrate when young. Asci broad globose, with 100-
 200 spores. Usually only 1-3 asci in each apothecium, which dehisce
 by bursting at the apex. 87
 (Other *Ryparobius* spp. will key out here [e.g. *R. dubius*, *R. myriosporus*,
R. pachyascus and *R. polysporus*]. They all have scattered to gregarious,
 immersed to semi-immersed apothecia 100-200µm diam., with relatively
 few asci, each with 100-250 ellipsoid to subacuminate *ca* 5-7 × 3-4µm
 spores. There are insufficient modern observations to allow their
 identification and separation with confidence).
- 87(86) Apothecia with a few, but obvious, setae. Spores 9 × 7µm or larger. 88
 - Apothecia without setae. Spores ellipsoid, 6-9 × 3.5-4µm. 89
- 88(87) Spores ellipsoid, 9-11 × 7-9µm. Setae up to 600µm long.
Trichobolus zukalii
 - Spores subglobose, 11-12 × 10-11µm. Setae up to 300µm long.
Trichobolus sphaerosporus (fig. 11)
- 89(87) Apothecia and asci large, 170-250µm diam. *Thelebolus stercoreus* (fig. 12)
 - Apothecia and asci small, rarely above 80-90µm diam.
Thelebolus nanus (fig. 13)
- 90(77) Asci 16(-24)-spored. Spores not closely aggregated into an
 imbricated mass, 13-14 × 6µm (8-9 × 4µm)*. Apothecial hairs
 rough, subulate. *Ascozonus parvisporus*
 - Asci with 32 or more spores. 91

- 91(90) Asci 32-spored. Spores $16.5-18 \times 4.5-5\mu\text{m}$ ($11-12 \times 3-3.5\mu\text{m}$)*.
 Apothecia with a single row of sharp, pointed, roughened hairs.
Ascozonus crouanii 92
 - Asci more than 32-spored.
- 92(91) Asci 48-spored. Spores spindle-shaped, $12-14.5 \times 2.5-4\mu\text{m}$.
Ascozonus leveillei 93
 - Asci more than 48-spored.
- 93(92) Asci 64-spored. 94
 - Asci more than 64-spored. 95
- 94(93) Apothecia with a short base of globose cells, with minutely
 roughened marginal hairs up to $30 \times 8\mu\text{m}$. Spores elliptic-fusoid,
 $12-14 \times 3-5\mu\text{m}$. *Ascozonus woolhopensis* (figs 14, 15)
 - Apothecia sessile, with aseptate smooth hairs. Spores $21 \times 7.5\mu\text{m}$
 ($13-14 \times 4.5-5\mu\text{m}$)*. *Ascozonus cunicularis*




Fig. 11. *Trichobolus sphaerosporus*, apothecium. Fig. 12. *Thelebolus stercoreus*, apothecium.
 Fig. 13. *T. nanus*, mature and immature apothecia, and detail of ascus dehiscence.
 Fig. 14. *Ascozonus woolhopensis*, apothecium and apothecial hair. Fig. 15. *A. woolhopensis*,
 ascus with spores and detail of dehiscence.

*There are few reports of *Ascozonus*, apart from *A. woolhopensis*. Observed spore sizes of *A. woolhopensis* suggest that measurement of Renny's (1874) illustrations of spores leads to values which are too large ($19-20 \times 6-6.5\mu\text{m}$). Those in parentheses are what they might be, based on the discrepancy between observed values for *A. woolhopensis* and Renny's illustration.

- 95(93) Apothecia with a short base of globose cells, with short, irregular hairs. Asci 64-96-spored Spores elliptic-fusoid, $14-14.5 \times 5-5.5 \mu\text{m}$ ($10-15 \times 3.5-4 \mu\text{m}$)*. *Ascozonus leveillanus*
- Apothecia sessile, dotted with hairs in connate groups of 2-3. Asci with 128 or more spores. Spores $10 \times 5 \mu\text{m}$ ($7 \times 3.5 \mu\text{m}$)*. *Ascozonus subhirtus*
- 96(3) Apothecia stalked. 97
- Apothecia not stalked. 98
- 97(96) Apothecia up to 2mm diam., with a short cylindrical stalk, light brown. Asci $150 \times 10 \mu\text{m}$. Spores hyaline, with 2 oil drops, occasionally 1-septate, $13-15 \times 4.5 \mu\text{m}$. *Lanzia cuniculi*
- Apothecia up to 3mm diam., pale olivaceous to grey, with a long, slender, reddish-brown stalk arising from a sclerotium in the dung. Asci $30-40 \times 4-5 \mu\text{m}$. Spores ellipsoid, grey-brown, $4-4.5 \times 2 \mu\text{m}$. *Martininia panamaensis*
- 98(96) Spores $7-11(14) \times 1.75-2.75 \mu\text{m}$. ellipsoid, ellipsoid-fusiform or slightly clavate. Apothecia yellowish brown when fresh, drying darker, up to 1mm diam. Asci $42-60 \times 7.5-9 \mu\text{m}$, pore weakly blue in iodine. *Pezizella albula*
- Spores and asci smaller. 99
- 99(98) Spores linear, $3-5 \times 1 \mu\text{m}$. Asci $30 \times 5 \mu\text{m}$, cylindrical with a short stipe. Paraphyses not clavate but fused to form an epithecium. Apothecia pale pellucid, 0.5-1mm diam. *Orbilina leporina*
- Spores longer, subulate, curved. 100
- 100(99) Spores $7-8.5 \times 1.2-1.8 \mu\text{m}$. Asci $36-40 \times 3-5 \mu\text{m}$, gradually tapering to a short base. Paraphyses enlarged to $3 \mu\text{m}$ at apex, covered with brown granules. Apothecia light brown, 0.4-1.2mm diam. *Orbilina fimicola*
- Spores $8-10.5 \times 0.9-1 \mu\text{m}$. Asci $30-45 \times 3 \mu\text{m}$, cylindrical-clavate with narrow tapering base and truncate apex. Paraphyses $2 \mu\text{m}$ diam., the tips with a crust-like secretion fusing together to form a shiny epithecium. Apothecia white to yellowish, 180-700 μm diam. *Orbilina fimicoloides*

Key 2. Perithecial, pseudothecial, cleistothecial and gymnothecial fungi

- | | | |
|------|--|-----|
| 1 | Perithecia occurring singly or in groups, but directly on the dung (key 1,2) or buried in it (figs 16, 18, 19, 22, 27, 30, 32, 34-36). | 2 |
| - | Perithecia occurring in or on a mass of fungal tissue (stroma) growing in or on the dung (figs 32, 37). | 135 |
| 2(1) | Spores black, brown or dark olive-greenish. | 3 |
| - | Spores hyaline or pale coloured, at least under the microscope (may be coppery red <i>en masse</i>). | 117 |
| 3(2) | Spores smooth, without an ornamentation of hyaline pits. | 4 |
| - | Spores 1-celled, ornamented with hyaline pits. (<i>Gelasinospora</i>) | 114 |
| 4(3) | Perithecia dark, olive, brown or black. | 5 |
| - | Perithecia reddish brown, orange or golden, globose, with a neck. Spores black, limoniform. | 116 |
| 5(4) | Perithecia globose, surmounted by a dense tuft of greyish green hairs, which may be branched or simple, straight or curly. Spores olivaceous, limoniform. Asci clavate, soon disappearing. (A large genus not characteristic of dung, but occurring occasionally). | |
| | <i>Chaetomium</i> (fig. 16) | |
| - | Perithecia more pyriform, or if globose then with a distinct neck, may be setose but not densely hairy, with clavate or cylindrical asci. | 6 |
| 6(5) | Each spore composed of 4 or more cells in a row (figs 17, 21). Asci bitunicate (figs 20, 23). | 7 |
| - | Spores 1- or 2-celled. Asci bitunicate or unitunicate. | 29 |
| 7(6) | Spores 16-32-celled, united firmly together in a bundle both in the ascus and after discharge. Germ slits usually absent. (<i>Sporormia</i>) | 8 |
| - | Spores each with 4 or more cells, each spore free and surrounded by its own gelatinous sheath. Germ slits usually present. (<i>Sporormiella</i>) | 11 |
| 8(7) | Spores 16-20-celled. | 9 |
| - | Spores 29-32-celled, 130-160 × 4-6µm. (<i>Sporormia mirabilis</i>) | |

- 9(8) Spores 16-celled, $85-116 \times 5-6.5\mu\text{m}$. *Sporormia fimicola* 10
 - Spores smaller.
- 10(9) Spores 16-celled, $37-45 \times 3\mu\text{m}$. Asci $50-60 \times 10-12\mu\text{m}$. *Sporormia* sp. (fig. 17)
 [recorded as *S. fimetaria* by Richardson (1972); see also Bell (1983) and
 Dissing (1992)]
 - Spores 16-20-celled, $50-57 \times 3.5-4.5\mu\text{m}$. Asci $70-80 \times 12-16\mu\text{m}$. *Sporormia fimetaria*
 (These two taxa may represent the extremes of *S. fimetaria*).
- 11(7) Spores 4-celled. 12
 - Spores more than 4-celled. 22
- 12(11) Spores more than $65-70\mu\text{m}$ long. 13
 - Spores less than $65-70\mu\text{m}$ long. 15




Fig. 16. *Chaetomium* sp., perithecium and spore. Fig. 17. *Sporormia* sp., ascus and spores.
 Fig. 18. *Sporormiella ovina*, pseudothecium. Fig. 19. *S. intermedia*, pseudothecium.
 Fig. 20. *S. intermedia*, immature bitunicate ascus and mature ascus with outer layer ruptured.
 Fig. 21. Ascospores of, from left, *S. ovina*, *S. intermedia* (with gelatinous sheath characteristic of
 the genus), *S. lageniformis*, *S. vexans*, *S. bipartis* and *S. minima*.

- 13(12) Spores 65-95 × 15-18µm. *Sporormiella megalospora*
 - Spores longer than 90µm. 14
- 14(13) Spores 90-118 × 15-20µm. Asci tapering gradually from the
 broadest part near the apex to a 'stipe'. *Sporormiella ovina* (figs 18, 21)
 - Spores 91-114 × (14)18-21µm. Asci cylindrical, abruptly
 contracted below to a short 'stipe'. *Sporormiella borealis*
- 15(12) Spores mostly less than 35µm long. 16
 - Spores mostly between 35-60µm long. 19
- 16(15) Spores less than 25µm long. 17
 - Spores 25-35(38)µm long. 18
- 17(16) Spores (15)17-24(26) × 5-7µm, end cells broadly conical.
 Ascospores uniseriate. Asci 120-135µm long. Pseudothecia
 250-300µm diam. *Sporormiella pulchella*
 - Spores 16-22 × 4.5-5.5µm, end cells subovate. Ascospores
 biseriate. Asci 95-125µm long. Pseudothecia 300-350µm diam.
Sporormiella nigropurpurea
- 18(16) Spores 30-38.5 × 5.5-6.5µm. Asci clavate, tapering gradually
 below to a 'stipe'. *Sporormiella leporina*
 - Spores 27-36(38) × 4-6(8)µm, tending to break in two at the middle
 septum. Asci cylindrical, abruptly contracted below.
Sporormiella minima (fig. 21)
- 19(15) Spores with end cells rounded. Asci cylindrical, abruptly contracted
 below. 20
 - Spores with end cells tapered and slightly conical. Asci clavate,
 tapering gradually to a long stalk. 21
- 20(19) Spores 45-65 × 8-11.5µm. *Sporormiella intermedia* (figs 19-21)
 - Spores 38-46 × 6.5-8µm. *Sporormiella australis*
- 21(19) Spores 45-60 × 11.5-14µm, germ slits parallel with long axis.
Sporormiella grandispora
 - Spores 35-45(48) × 7-9(10)µm. *Sporormiella lageniformis* (fig. 21)
- 22(11) Spores 5-celled, 70-80 × 17-19µm. *Sporormiella pentamera*
 - Spores more than 5-celled. 23

23(22)	Spores 7- or 8-celled.	24
-	Spores 13-celled, 46-60 × 9-10µm.	<i>Sporormiella antarctica</i>
24(23)	Spores 7-celled.	25
-	Spores 8-celled.	26
25(24)	Spores 40-55 × 7-9µm, readily disarticulating, the end cells longer than wide, the rest shorter than wide.	<i>Sporormiella vexans</i> (fig. 21)
-	Spores 70-80 × 16-18µm, end cells rounded.	<i>Sporormiella heptamera</i>
26(24)	Spores mostly longer than 45µm.	27
-	Spores less than 50µm long, not disarticulating at the central septum.	28
27(26)	Spores 45-60 × 5-7.5µm, disarticulating at the central septum, all cells the same width.	<i>Sporormiella bipartis</i> (fig. 21)
-	Spores 50-59 × 10-12µm, not disarticulating, 3rd cell down wider than the others.	<i>Sporormiella corynespora</i>
28(26)	Spores (33)37-40(49) × 7-9µm, cylindrical. Asci abruptly contracted below.	<i>Sporormiella pascua</i>
-	Spores 40-48 × 7-8µm, fusiform cylindrical. Asci gradually tapered below.	<i>Sporormiella octomera</i>
29(6)	Spores obviously 2-celled at maturity.	30
-	Spores 1-celled, or appearing 1-celled at maturity. (Those of <i>Podospora</i> , <i>Schizothecium</i> etc. are 2-celled in early stages of their development, but only one cell matures to become pigmented; the other remains hyaline, often collapses, and may be difficult to see).	47
30(29)	Spores 23-28 × 13-17µm, upper cell dark, 15-19µm, with close, blunt spines giving the impression of a pitted spore surface, with apical germ pore, the lower cell hyaline, 6-8.5µm, smoky-brown. Asci unitunicate, 4-spored. Perithecia 400µm diam.	<i>Apiosordaria verruculosa</i> (fig. 24)
-	Both cells of spore similar in shape, size and colour.	31
31(30)	Asci unitunicate. Spores with a 'gelatinous' appendage at each end. Perithecial neck with setae.	32
-	Asci bitunicate. Spores without gelatinous appendages, although a sheath may be present.	33

- 32(31) Spores 38-48 × 11-14µm, appendages longitudinally
fibrillate. *Zygospermella striata*
- Spores 46-68 × 11-17µm, appendages hollow, not fibrillate.
Zygospermella insignis (fig. 25)
- 33(31) Spores with each end truncated by a germ pore. Pseudothecia
with dark bristles at neck. (*Trichodelitschia*) 34
- Spores with rounded ends and germ slits along the sides.
Pseudothecial neck smooth or hairy, but without setae.
(*Delitschia*, fig. 26) 36
- 34(33) Spores 28-34 × 9-12µm. *Trichodelitschia aedelphica*
- Spores smaller. 35
- 35(34) Spores 20-27.5 × 8-11µm. *Trichodelitschia bisporula* (figs 22, 23)
- Spores 18-21 × 6-7µm. *Trichodelitschia munkii*
- 36(33) Asci ca 256-spored. Spores 14-15 × 6-8µm. *Delitschia myriasporea*
- Asci 8-spored. 37
- 37(36) Spores less than 20µm long. 38
- Spores more than 20µm long. 41
- 38(37) Spores 8-11 × 3-5µm. *Delitschia perpusilla*
- Spores 10-20µm long. 39
- 39(38) Spores 10-14 × 5-6µm. *Delitschia marchalii*
- Spores longer. 40
- 40(39) Spores 14-18 × 6-10µm, uniseriate. Asci 70-90 × 7-16µm.
Delitschia niesslii
- Spores (16)18-20(22.5) × 6-7.5µm, biseriate. Asci 80-145 ×
20-25µm. *Delitschia consociata* (fig. 26)
- 41(37) Spores mostly wider than 20µm. 42
- Spores mostly less than 20µm wide. 43
- 42(41) Spores 50-64 × 19-23µm. *Delitschia furfuracea*
- Spores 50-70 × 25-33µm. *Delitschia winteri* (fig. 26)

- 43(41) Spores $20-25 \times 4.5-6\mu\text{m}$, the cells slightly tapered and almost completely separated. Pseudothecia hairless, globose, *ca* $200\mu\text{m}$ diam. *Delitschia leptospora* (fig. 26)
 - Spores longer and wider. 44
- 44(43) Spores transversely septate. 45
 - Spores obliquely septate, deeply constricted at the septum, $35-50 \times 15-18\mu\text{m}$. *Delitschia didyma*
- 45(44) Pseudothecia hairy. Spores $37-50 \times 17-20\mu\text{m}$, not deeply constricted at the septum. *Delitschia chaetomioides*
 - Pseudothecia smooth. 46
- 46(45) Spores biseriata, $45-55 \times 13-16\mu\text{m}$, one cell usually larger than the other, deeply constricted at the septum and readily separating. *Delitschia canina*
 - Spores uniseriate, $40-55 \times 16-21\mu\text{m}$, both cells equal. *Delitschia patagonica*




Fig. 22. *Trichodelitschia bisporula*, pseudothecium. **Fig. 23.** *T. bisporula*, expanded ascus broken through the outer wall, with spores. **Fig. 24.** *Apiosordaria verruculosa*, ascospores. **Fig. 25.** *Zygospermella insignis*, ascus and ascospore. **Fig. 26.** Ascospores of, from left, *Delitschia winteri*, *D. consociata* and *D. leptospora*.

- 47(29) Spores with colourless 'gelatinous' secondary appendages (caudae, fig. 28) at one or both ends (not always easy to see; mounting in Indian ink is useful, and essential for some). A hyaline (empty) cell, the primary appendage (fig. 28), may also be present. 48
 - Spores without caudae, although a colourless gelatinous sheath may be present. Primary appendages present or absent. 88
- 48(47) Perithecia often hairy or tomentose when young. Immature spores long, wavy cylindrical, with a row of globules, and more likely to be seen than mature spores (fig. 29). Secondary appendages thin, simple, up to $60 \times 3\mu\text{m}$. Mature spores with a dark cell $14-25 \times 7-13\mu\text{m}$ and pedicel (primary appendage) $25-50 \times 3-6\mu\text{m}$.
 (Cercophora) 49
 - Perithecia often with scales or setae at the neck or tomentose. Caudae, simple or compound. Immature spores clavate or ellipsoid, not long, wavy cylindrical. Mature spores readily observed. 51
- 49(48) Immature spores $45-70 \times 4-6\mu\text{m}$. 50
 - Immature spores smaller, $38-52 \times 3-3.5\mu\text{m}$. Mature spores with upper (dark) cell $14-18 \times 7-9\mu\text{m}$; hyaline pedicel $27-36 \times 3-3.5\mu\text{m}$.
Cercophora silvatica
- 50(49) Perithecia with white or grey tomentum. Young spores $45-65 \times 4.5-6\mu\text{m}$. Mature spores with upper cell $17-25 \times 8.5-13\mu\text{m}$ and pedicel $30-50\mu\text{m}$ long. *Cercophora coprophila* (fig. 29)
 - Perithecia with flexuose brown hairs and, at the neck, tufts of agglutinated, swollen, obtuse hairs. Young spores $52-68 \times 4-5\mu\text{m}$. Mature spores with upper cell $15-25 \times 9-11\mu\text{m}$ and pedicel $35-45\mu\text{m}$ long. *Cercophora mirabilis*
- 51(48) Primary appendage absent. (*Arnium*, fig. 28) 52
 - Primary appendage present. 60
- 52(51) Asci (64-)128-spored. Spores $18-26 \times (10)12-15\mu\text{m}$. Perithecial neck sometimes with rigid, brown, septate hairs up to $330\mu\text{m}$.
Arnium leporinum
 - Asci 4- or 8-spored. 53
- 53(52) Asci 4-spored. 54
 - Asci 8-spored. 55

- 54(53) Spores ellipsoid, sometimes inequilaterally flattened, $44-54 \times 22-30\mu\text{m}$, with 1 apical germ pore, caudae not swelling in water. Perithecium usually with lateral tufts of agglutinated hairs up to $550\mu\text{m}$ long. *Arnium arizonense*
- Spores evenly ellipsoid-fusiform, $31-55 \times 18-25\mu\text{m}$, with germ pore at each end, caudae covering germ pores, $35-60 \times 7-11\mu\text{m}$, but rupturing and swelling to up to $130 \times 50\mu\text{m}$, and becoming diffuse and irregular. Perithecial neck covered with rigid hairs up to $190 \times 2.5\mu\text{m}$. *Arnium hirtum*
- 55(53) Perithecial neck distinctly setose with rigid hairs. 56
- Perithecial neck without setae. 57
- 56(55) Spores evenly ellipsoid-fusiform, $31-55 \times 18-25\mu\text{m}$, with germ pore at each end, caudae covering germ pores, $35-60 \times 7-11\mu\text{m}$, but rupturing and swelling up to $130 \times 50\mu\text{m}$, and becoming diffuse and irregular. Perithecial neck covered with rigid hairs up to $190 \times 2.5\mu\text{m}$. *Arnium hirtum*
- Spores slightly inequilateral, $35-43 \times 17-23\mu\text{m}$, caudae $50-75 \times 5-8\mu\text{m}$, not covering germ pores. Perithecial neck with brown hairs up to $250\mu\text{m}$ long. *Arnium cervinum*
- 57(55) Perithecia covered with a dense tomentum of septate flexuous hairs. Spores mostly longer than $45\mu\text{m}$. Only occasionally fimicolous. 58
- Perithecia without a tomentum. Spores up to $45\mu\text{m}$. 59
- 58(57) Spores $(40)45-54 \times 25-35\mu\text{m}$, uniseriate. Tomentum pale or greyish. *Arnium olerum*
- Spores $47-70 \times 20-30\mu\text{m}$, biseriate above. Tomentum olivaceous brown. *Arnium tomentosum*
- 59(57) Spores somewhat inequilateral, rounded below, pointed above, $31-40 \times 18-24\mu\text{m}$, caudae $50-120 \times 6-10\mu\text{m}$, with 1 apical germ pore not covered by cauda. *Arnium caballinum*
- Spores equilateral, $36-44 \times 20-23\mu\text{m}$, caudae $50-80 \times 6-8\mu\text{m}$, covering germ pores. *Arnium mendax*
- 60(51) Perithecia with scales at the neck, composed of inflated and agglutinated cells (fig. 27, *S. conicum*). (*Schizothecium*) 61
- Perithecia setose or hairy at the neck, but not with inflated cells, or neck black but almost hairless. (*Podospora*) 70

61(60)	Asci 4-spored.		62
-	Asci 8-spored.		63
62(61)	Spores 11-14.5 × 6.5-9µm.	<i>Schizothecium nanum</i> (fig. 28)	
-	Spores 19-24 × 12-14.5µm.	<i>Schizothecium tetrasporum</i>	
63(61)	Spores more than 30µm long.		64
-	Spores less than 30µm long.		65
64(63)	Perithecia crowned with a fascicle of long agglutinated hairs at the neck, up to 335µm long. Spores 31-40 × 15-25µm, biseriate.	<i>Schizothecium aloides</i>	
-	Perithecia with shorter, less remarkable tufts. Spores 30-45 × 19-24µm, ± uniseriate.	<i>Schizothecium glutinans</i>	
65(63)	Perithecial neck with rigid setae, as well as agglutinated hairs (which may be greatly reduced). Asci 140-210 × 19-25µm, broadest at the markedly rounded apex. Spores 18-23 × 11-14µm.	<i>Schizothecium pilosum</i>	
-	Perithecial neck without rigid setae. Asci broadest in the middle.		66
66(65)	Spores mostly over 23µm long.		67
-	Spores up to 23µm long.		69
67(66)	Spores 22-25(27) × 11-13µm. Scales at neck distinct.	<i>Schizothecium hispidulum</i>	
-	Spores wider, 12-19µm.		68
68(67)	Perithecia 0.5-1mm high, scales at neck usually well developed. Spores (23)26-30 × 12-17µm.	<i>Schizothecium conicum</i> (fig. 27)	
-	Perithecia 1-2mm diam., subpyriform, neck velvety with indistinct scales. Spores 24-28 × 15-19µm.	<i>Schizothecium squamulosum</i>	
69(66)	Spores 17-23 × 8.5-13.5µm, primary appendage slender cylindrical, 6-8 × 2µm. Perithecia 0.25-0.7mm high, sometimes with poorly developed scales.	<i>Schizothecium vesticola</i> (fig. 28)	
-	Spores 11-14 × 6-8µm, primary appendage short, 2µm long, almost triangular. Perithecia 0.3-0.45mm high, with short agglutinated hairs.	<i>Schizothecium cervinum</i>	
70(60)	Asci 4-spored. Spores 35-40 × 18-19µm.	<i>Podospora pauciseta</i>	
-	Asci with more than 4 spores.		71




Fig. 27. Perithecia, from left, of *Podospora appendiculata*, *Schizothecium conicum*, *P. excentrica* and *P. decipiens*, with detail of hairs. **Fig. 28.** Ascospores of, from left, *Podospora excentrica*, *P. appendiculata*, *S. vesticola*, *S. nanum*, *P. decipiens*, '*P. dagobertii*' and *Arnium* sp. **Fig. 29.** *Cercophora coprophila*, immature (l) and mature (r) ascospores.

71(70)	Asci 8-spored.	72
-	Asci with more than 8 spores.	82
72(71)	Spores more than 45 μ m long.	73
-	Spores less than 45 μ m long.	74
73(72)	Spores 48-60 \times 27-31 μ m, caudae apparently striate. Perithecia superficial, covered with rigid, nonagglutinated hairs up to 120 μ m.	<i>Podospora fimiseda</i>
-	Spores 50-68 \times 22-32 μ m, caudae apparently segmented, with an intestine-like appearance. Perithecia immersed to superficial, with a long neck, tomentose with long flexuous hairs when young, more or less glabrous when mature.	<i>Podospora intestinacea</i>
74 (72)	Perithecia superficial, ovoid to globose, covered with short (up to 100 μ m), sparse, radiating, hyaline tipped, hairs. Spores 24-31 \times 11-15 μ m, with simple caudae.	<i>Podospora appendiculata</i> (figs 27, 28)
-	Perithecia with base immersed in substrate, pyriform, without such hairs.	75
75 (74)	Perithecial neck with short tubercular hairs, up to 20 μ m long. Spores 32-42 \times 17-22 μ m, with a long but withering primary appendage. Caudae in two rings, one inserted near the base of the primary appendage, the other at the spore apex. The individual filaments may be free, but often clump together to form an apparently broad appendage.	<i>Podospora decipiens</i> (figs 27, 28)
-	Perithecial hairs longer. Caudae single or 4 at each end.	76
76(75)	Spores with 4 caudae at each end.	77
-	Spores with a single cauda at each end.	78
77(76)	Spores 40-45 \times 22-25 μ m.	<i>Podospora gwynne-vaughaniae</i>
-	Spores 29-40 \times 16-25 μ m.	<i>Podospora communis</i>
78(76)	Spores less than 30 \times 15 μ m.	79
-	Spores larger than 30 \times 15 μ m.	80

- 79(78) Spores 21-28 × 11-14µm, primary appendage 12-14 × 4µm. Perithecia 0.3-0.5mm diam., neck setose with rigid cylindrical hairs. Asci 200-250 × 22-26µm, broadest in the middle. *Podospora ellisiana*
- Spores 18-23 × 11-14µm, primary appendage 4-8 × 3µm. Perithecia 0.2-0.3mm diam, neck setose with rigid hairs. Asci 140-210 × 19-25µm, broadest at the markedly rounded apex. *Schizothecium pilosum*
- 80(78) Perithecia ca 0.9-1.4mm high × 0.6-0.7(0.85)mm diam., neck not hairy. Spores (29)36-45 × (17.5)22-27µm, caudae ephemeral and difficult to see, even in Indian ink. *Podospora pyriformis*
- Perithecial neck with tufts of rigid hairs. 81
- 81(80) Perithecia 0.38-0.53mm high × 0.21-0.38mm diam., ± immersed, with hairs at the neck up to 335µm long, grouped in rigid fascicles. Spores slightly flattened on one side, 30-37 × 18-24µm, caudae invisible in water. *Podospora excentrica* (figs 27, 28)
- Perithecia ca 0.8-1.4mm high × 0.4-0.7mm diam., semi-immersed, hairy all over, flexuous below, rigid and pointed at the neck up to 170µm. Spores 33-45 × 22-27µm. *Podospora perplexens*
- 82(71) Asci 16-32-spored. Perithecial neck with short tubercular hairs. Spores 25-36 × 15-24µm. Caudae in two rings, one inserted at the base of the primary appendage, the other at the spore apex; individual filaments may be separate or clumped to appear as a broad single appendage (cf. *P. decipiens*). *Podospora pleiospora*
- Asci with more than 32 spores. 83
- 83(82) Perithecia with tufts of rigid hairs at neck. Asci with more than 64 spores. 84
- Perithecia without tufts of rigid hairs. Asci 64-spored. 87
- 84(83) Spores 14-17 × 9-11µm. Asci 256-spored. Perithecia ca 500µm diam., immersed, except for the neck, which has tapered tufts of hairs up to 300µm. *Podospora curvicolla*
- Spores larger. Perithecia semi-immersed. 85

- 85(84) Spores (18)20-26 × 12-16 μ m, caudae of 2-several filaments covered with granules. Asci 512-spored. Perithecia up to 1mm high × 0.95mm diam., neck with rigid but non-agglutinated hairs up to 130 μ m long. *Podospora granulostriata*
- Caudae simple, without granular appearance. Asci 128-spored. Perithecia not larger than 750 μ m high × 500 μ m diam., with rigid, non-agglutinated hairs up to 190 μ m long at neck. 86
- 86(85) Spores 17-19 × 10-12 μ m. *Podospora setosa*
- Spores 19-24 × 11-16 μ m. *Podospora tarvisina*
(See discussion in Lundqvist (1972) on these last three names)
- 87(83) Spores 24-34 × 14-19 μ m, caudae in two rings, one inserted at the base of the primary appendage, the other at the spore apex; individual filaments may be separate or clumped to appear as a broad single appendage (cf. *P. decipiens*/*P. pleiospora*). Perithecia ca 0.6-1.1mm high × 0.4-0.5mm diam., covered with flexuous hairs or rarely smooth. *Podospora myriasporea*
- Spores 15-20 × 10-15 μ m, caudae small, simple and evanescent. Perithecia 0.4-0.5mm high, covered with long flexuous hairs. *Podospora collapsa*
- 88(47) Spores with primary appendage. 89
- Spores without primary appendage. 93
- 89(88) Spores with primary appendage directed towards base of ascus. 90
- Spores with primary appendage directed towards apex of ascus. (*Anopodium*) 91




Fig. 30. Perithecia of, from left, *Coniochaeta ligniaria*, *C. scatigena* and *C. hansenii*.
Fig. 31. Ascospores of *C. scatigena* (l) and *C. ligniaria* (r).

- 90(89) Spores $34-45 \times 19-25\mu\text{m}$, without caudae but surrounded by a thin (*ca* $5\mu\text{m}$) gelatinous sheath. Perithecia *ca* $0.5-0.7\text{mm}$ diam., \pm smooth. *Podospora globosa*
- Spores $17-20 \times 8-9.5\mu\text{m}$, flattened on one side, convex on the other. Perithecia $0.3-0.45\text{mm}$ diam., with distal cells of agglutinated hairs fimbriate. *Podospora fimbriata*
- 91(89) Perithecia hairy. Spores $27-32 \times 16-19\mu\text{m}$, appendage $15-18 \times 2.5-3\mu\text{m}$. *Anopodium ampullaceum*
- Perithecia glabrous. 92
- 92(91) Spores $28-32 \times 16-21\mu\text{m}$, appendage $12-15 \times 3-3.8\mu\text{m}$. *Anopodium epile*
- Spores $30-37 \times 16-20\mu\text{m}$, appendage $24-27 \times 5\mu\text{m}$.
'*Podospora*' *dagobertii* (fig. 28)
(The combination in *Anopodium* has not been made; see Lundqvist, 1964,1972)
- 93(88) Spores flattened, disc shaped, with a germ slit around the edge. Perithecial neck with short (up to $120\mu\text{m}$) setae. (*Coniochaeta*, figs 30, 31) 94
- Spores ellipsoid. Perithecial neck without setae **or** with very prominent (up to $950\mu\text{m}$) tufts of agglutinated hairs. 99
- 94(93) Asci with numerous (64-128) spores. 95
- Asci 8-spored. 96
- 95(94) Spores $6-10 \times 5-9 \times 4-7\mu\text{m}$. Perithecial setae up to $120\mu\text{m}$ long. *Coniochaeta hansenii* (fig. 30)
- Spores $13-16 \times 9.5-13.5 \times 5.5-8\mu\text{m}$. Perithecial setae up to $35\mu\text{m}$ long. *Coniochaeta* sp.
- 96(94) Spores $7-9 \times 6-8 \times 5-6\mu\text{m}$, slightly flattened. *Coniochaeta leucoplaca*
- Spores larger. 97
- 97(96) Spores narrowly elliptical in face view (length more than $2 \times$ width), *ca* $13-18 \times 6-9 \times 4-6\mu\text{m}$. *Coniochaeta saccardoii*
- Spores broadly elliptical to nearly circular in face view (length less than $2 \times$ width). 98

- 98(97) Spores (9)10-16(20) × 7.5-10(15) × (4)5-8 μ m. Neck setae
20-50 μ m long. *Coniochaeta ligniaria* (figs 30, 31)
- Spores (16)17-23 × (10)13-19 × 7.5-10(15) μ m. Neck setae
40-80 μ m long. *Coniochaeta scatigena* (figs 30, 31)
- 99(93) Perithecial neck with prominent agglutinated tufts of rigid setae up
to 950 μ m long. Spores 43-54 × 20-29 μ m, with apical germ pore.
A gelatinous sheath which surrounds the whole spore swells in
water, and appears fringed at the margin and radially striate.
Arnium macrothecium
- Perithecial neck without setae. Gelatinous sheaths may be clearly
visible around spores, but are not complex in structure. 100
- 100(99) Spores with germ slit along the side. Ascus with a large and
complex plug at the tip staining blue or red in KI (other genera
have asci with blue staining ascus tips, but the feature is very
pronounced in this genus and is unlikely to be mistaken).
Perithecia form singly or severally in a stroma which is usually of
limited extent, often without a definite margin. [N.B. if orange and
with a stroma see *Selinia*, 119]. (*Hypocopra*, fig. 32) 101
- Spores without germ slits, but often asymmetrical, and with a
small papilla at the basal end. Asci without complex apical
plug. (*Sordaria*, fig. 33) 107
- 101(100) Spores mostly less than 25 μ m long. 102
- Spores more than 25 μ m long. 104
- 102(101) Spores 9-14 × 6-7 μ m. *Hypocopra parvula*
- Spores larger. 103
- 103(102) Stroma with a brown hyphal mat between perithecial necks.
Spores 19-27 × 10-14 μ m. *Hypocopra equorum* (fig. 32)
- Stroma with white hyphae between black perithecial necks,
becoming smooth. Spores 23-25 × 12-14 μ m. *Hypocopra brefeldii*
- 104(101) Ascospores up to 15 μ m wide. 105
- Ascospores 15 μ m or wider. 106
- 105(104) Ascospores 25-31 × 10-15 μ m, distinctly flattened on one side.
Ascus plug blue in KI, but becoming reddish. *Hypocopra planispora*
- Ascospores 26-32 × 13-14 μ m, ellipsoid and narrowed towards
their ends. *Hypocopra stephanophora*

- 106(104) Ascospores $27-43 \times 16-20\mu\text{m}$. *Hypocopra merdaria*
 - Ascospores $38-50 \times 19-24\mu\text{m}$. *Hypocopra stercoraria*
- 107(100) Spores up to $10\mu\text{m}$ long. 108
 - Spores $10\mu\text{m}$ or longer. 109
- 108(107) Asci 8-spored. Spores $8 \times 4\mu\text{m}$. *Sordaria minima*
 - Asci ca 128-spored. Spores $5-8 \times 4-5\mu\text{m}$. *Sordaria polyspora*
- 109(107) Spores relatively narrow, at least twice as long as wide,
 $22-26 \times 9-12\mu\text{m}$. Gelatinous sheath broad, distinct. *Sordaria alcina*
 - Spores relatively broad, less than twice as long as wide. 110
- 110(109) Spores mostly $25\mu\text{m}$ or longer. 111
 - Spores up to $25\mu\text{m}$ long. 112
- 111(110) Spores $(21)23-29(30) \times 14.5-17(18)\mu\text{m}$, with apiculate base.
 Gelatinous sheath broad, distinct. Asci $240-300 \times 20-24\mu\text{m}$. *Sordaria superba*
 - Spores $(26)28-35 \times (17)18-22\mu\text{m}$, with slightly apiculate base.
 Gelatinous sheath broad, distinct. Asci $280-350 \times 30-35\mu\text{m}$. *Sordaria macrospora*
- 112(110) Spores with gelatinous sheath absent or very thin, $19.5-25 \times$
 $15.5-19\mu\text{m}$. *Sordaria humana* (fig. 33)
 - Spores with gelatinous sheath, up to $15\mu\text{m}$ diam. 113
- 113(112) Spores obovoid to broadly ellipsoid, $18-23 \times 12-15\mu\text{m}$. *Sordaria lappae*
 - Spores ellipsoid, $17-25 \times 10-14\mu\text{m}$. *Sordaria fimicola* (fig. 33)




Fig. 32. *Hypocopra equorum*, perithecium with limited stroma, and detail of ascus tip with blue staining plug and spore. **Fig. 33.** Ascospores, from left, of *Sordaria fimicola*, *S. humana* and *Sphaerodes fimicola*.

- 114(3) Spores 20-28 × 12-16µm, with subacute ends, each with a germ pore. *Gelasinospora adjuncta*
 - Spores larger. 115
- 115(114) Asci 4-spored. Spores 24-29 × 15-18µm, with rounded ends and one germ pore. *Gelasinospora tetrasperma*
 - Asci 8-spored. Spores 26-35 × 22-27µm. *Gelasinospora cerealis*
- 116(4) Perithecia orange to golden, often gregarious, almost spherical, necks ca 50µm diam., 15µm high, setae at ostiole hyaline, up to 35 × 3µm. Spores limoniform, with a germ pore at each end, 15-25 × 9-16µm. *Sphaerodes fimicola* (fig. 33)
 - Perithecia yellow or reddish brown (darker when filled with mature spores), neck 50µm long, with setae at the ostiole 40-70µm long. Spores dark brown to black, limoniform, 20-34 × 11-17µm, with apical germ pore. *Melanospora brevirostris*
- 117(2) Asci more than 8-spored. see Key 1 at 86
 - Asci with 8 or fewer spores, or asci evanescent, not readily observed. 118
- 118(117) Perithecia orange/yellow, 500-1000µm diam. Spores long (over 45µm) or 2-celled if shorter. 119
 - Perithecia smaller, or black or with a neck. Spores shorter (less than 20µm) or septate if longer. 120
- 119(118) Perithecia orange, 500-1000µm diam., in small groups on a limited stroma. Spores thick walled, 48-60 × 22-26µm, with a gelatinous sheath. *Selinia pulchra*
 - Perithecia orange yellow, superficial, ca 500µm diam., with ostiole in a disc surrounded by silvery triangular tufts of hyphae ca 100µm long. Spores ellipsoid, 1-septate, 12-14 × 4-5µm. *Nectria suffulta*
- 120(118) Perithecia reddish brown or pale, hyaline, with a distinct neck. 121
 - Perithecia black. 131

- 121(120) Perithecia globose, up to 250 μ m diam., immersed, reddish brown, with a neck 1-3 mm long. Asci broad ellipsoid, 5-8.5 μ m, rapidly breaking down and difficult to see. Spores ellipsoid-allantoid, 5.5-7 \times 1.5-2 μ m, collecting in a pearly droplet at the fringed tip of the perithecial beak. *Viennotidia fimicola* (fig. 34)
- Perithecia pyriform, very pale in colour, 60-200 μ m diam., with a neck 60-700 μ m long. Asci rarely visible. Spores pointed-fusiform, 1-3 septate, often with a sheath and clumped together in fascicles. (*Pyxidiophora*, fig. 36)122
- 122(121) Neck 95-145 μ m long, brown, rugose, with cells arranged in 5-6 longitudinal rows visible in one view. Spores 38-52 μ m long. *Pyxidiophora badiorostris*
- Neck not brown or rugose, composed of hyaline, irregularly arranged cylindrical cells. 123
- 123(122) Spores less than 45 μ m long. 124
- Spores more than 45 μ m long. 125
- 124(123) Spores 35-45 μ m long, with brown apical or subapical patches of pigment. *Pyxidiophora brunneocapitatus*
- Spores 35-43 μ m long, without brown apical or subapical patches of pigment. *Pyxidiophora microsporus*
- 125(123) Spores mostly 45-60 μ m long. 126
- Spores mostly longer than 60 μ m. 129




Fig. 34. *Viennotidia fimicola*, perithecium and spores. Fig. 35. *Phomatospora coprophila*, perithecium, and ascus with spores. Fig. 36. *Pyxidiophora patchii*, perithecium and spores.

- 126(125) Perithecia 70-100 μ m diam., neck 100-190 μ m long. Spores
(43)48-58(65) μ m long. *Pyxidiophora grovei*
- Perithecia usually less than 80 μ m diam. 127
- 127(126) Perithecial necks mostly less than 100 μ m long. Spores
(45)48-57(60) μ m long. *Pyxidiophora arvernensis*
- Perithecial necks up to 200 μ m long. 128
- 128(127) Spores 45-53 μ m long. *Pyxidiophora petchii* (fig. 36)
- Spores 53-65 μ m long. *Pyxidiophora schotterianus*
- 129(125) Spores 60-70 μ m long. 130
- Spores (75)80-90(100) μ m. Perithecia 120-160 μ m diam.,
neck 220-370 μ m long. *Pyxidiophora bainemensis*
- 130(129) Perithecial necks 300-700 μ m long. Spores 60-70 μ m.
Perithecia 100-120 μ m diam. *Pyxidiophora spinuliformis*
- Perithecial necks 225-265 μ m long. Spores 65-70 μ m.
Perithecia 110-125 μ m diam. *Pyxidiophora marchalii*
- 131(120) Perithecia small, up to 400 μ m diam., with hairy necks. Spores
hyaline or pale, coppery-red *en masse*, extruded in tendrils. 132
- Perithecia larger, without hairy necks. If smaller than 200 μ m, with
spores smaller than 5 \times 3 μ m. 134
- 132(131) Spores reniform, with gelatinous sheath, 3-5 \times 2-3 μ m, yellow,
reddish brown *en masse* in extruded tendrils. Asci spherical,
evanescent. Perithecia black, spherical, 200-400 μ m diam., with
cylindrical neck up to 300 μ m long, with sparse pointed hairs.
Microascus longirostris
- Spores larger, not reniform. Perithecia up to 300 μ m diam. 133
- 133(132) Perithecial necks long, up to 750 μ m, with terminal hairs up to
1500 μ m, curved or circinate at tips. Spores limoniform,
7-10.5 \times 5.5-7 μ m. *Lophotrichus ampullus*
- Perithecial necks short, *ca* 50 μ m, with long straight tapering hairs.
Spore shape limoniform/variable, 6-7.5 \times 5-5.5 μ m, with prominent
germ pores. *Lophotrichus bartletti*

- 134(131) Perithecia up to 150 μm diam., immersed but for a conical neck
50-75 μm high. Asci 50 \times 2-2.5 μm . Spores minute, cylindrical,
3.5-4.5 \times 1.75-2.5 μm . *Phomatospora coprophila* (fig. 35)
- Perithecia more obvious, often hairy, or tomentose when young.
Immature spores up to 70 μm long, wavy cylindrical, with a row
of globules inside and a short thin appendage at each end.
(see *Cercophora*, 49)
- 135(1) Perithecia immersed, surrounded at the neck by a very limited -
flange-like stroma which is easily overlooked. see *Hypocopra*, 101
or if orange see *Selinia*, 119
- Stroma very conspicuous. 136
- 136(135) Perithecia in a subglobose group at the tip of the stromatic stalk.
Spores with germ slit and gelatinous sheath. (*Podosordaria*) 137
- Perithecia not in a terminal head. 139
- 137(136) Stalk short, 3-5mm. Spores (12)14-19 \times 6-9 μm , slightly flattened
on one side. *Podosordaria leporina*
- Stalk long, 1-6cm. Spores larger. 138
- 138(137) Spores 21-24 \times 11-12 μm . Stromatic stalk hairy. *Podosordaria tulasnei*
- Spores 40-60 \times 20-30 μm . Stromatic stalk not hairy.
Podosordaria pedunculata
- 139(136) Stroma externally black, rooted or partially immersed in the dung,
expanding at the surface to form a white disc up to 15mm diam.,
punctate with black perithecial ostioles. (*Poronia*) 140
- Stroma not as above. 141
- 140(139) Spores 18-26 \times 7-12 μm , bean shaped, with gelatinous sheath.
Stroma deeply rooted. Especially on horse dung. *Poronia punctata*
- Spores (22)25-32(35) \times (12)14-18 μm , oblong ellipsoid to slightly
fusiform. Stroma not deeply rooted. Especially on rabbit dung
near the sea. *Poronia erici*

- 141(139) Stroma spreading over surface of dung or filamentous. Spores ellipsoid to slightly flattened on one side, with germ slit. (Xerophilic fungi developing after long periods of relatively dry incubation). (*Wawelia*) 142
- Stroma clavate, black, partly immersed to superficial, usually aggregated in small groups, *ca* 1-1.5mm high \times 0.6-0.7mm diam., each containing a single perithecium. Spores ellipsoid with germ pore and gelatinous sheath. (*Bombardioidea*) 146
- 142(141) Stroma spreading on substrate, black brown, firm but not brittle. Ascumata globose, 0.5-1mm, with white hyphae at neck. Spores broad limoniform, 15-19 \times 9-10 μ m. *Wawelia effusa*
- Perithecia globose to pyriform, black, brown or dark grey, produced laterally along the length of fine stromatal strands growing from the dung. 143
- 143(142) Asci 4-spored. 144
- Asci 8-spored. 145
- 144(143) Spores 15-18 \times 9-12 μ m. Perithecia up to 400 μ m diam., dark grey at maturity, single or clustered, the ostiole with a crown of silvery white hyphae. Stromata up to 30 \times 0.1-0.5mm. *Wawelia* sp.
- Spores 6-8 \times 4-6 μ m. Stromata conical, white, 5-12 \times 1-2mm. *Wawelia regia*
- 145(143) Perithecia hairy, globose, 350-500 μ m diam., stromatal strands up to 25mm long. Spores ellipsoid, flattened on one side, 9-12 \times 6-8 μ m. *Wawelia octospora*
- Perithecia villose with conidiophores, globose, 230-420 μ m diam., produced laterally on stromatic filaments 20-30 \times 0.1-0.3mm. Filaments pink at first, with a white pointed tip, becoming brown, velvety with conidiophores. Spores ellipsoid to flattened on one side, 7.5-9.5 \times 3-4.5 μ m. *Wawelia* sp. (fig. 37)




Fig. 37. *Wawelia* sp., stromatic filaments with perithecia growing from a rabbit pellet, ascospores, and conidiophore and conidia.

- 146(141) Asci 8-spored. Spores $20-31 \times 9.5-15\mu\text{m}$.
Bombardioidea bombardioides
 - Asci 4-spored. 147
- 147(146) Spores $24-34 \times 15-19(20)\mu\text{m}$. Basal germ pore less distinct than
 the apical one. *Bombardioidea serignanensis*
 - Spores $34-43 \times 16-22\mu\text{m}$. Distinct germ pore at each end of spore.
Bombardioidea stercoreis
- 148 Fruit bodies solitary or in small groups, each a subglobose, fertile,
 (key 1,1) light brown head on a slender sterile stalk. Head soon bursting to
 expose the yellow ochraceous spore mass. On mixtures of bird
 droppings, cast pellets and decaying animal material. 149
 - Fruit bodies superficial, lacking a distinct stalk. 150
- 149(148) Spores $5-8 \times 2-3\mu\text{m}$. Head 1-2mm diam. *Onygena corvina* (fig. 38)
 - Spores $7-9 \times 4-6\mu\text{m}$. Head 2-4mm diam. *Onygena equina*
- 150(148) Fruit bodies with an external wall of loosely anastomosing and
 interwoven hyphae, and with \pm specialised terminal cells
 (*gymnothecia*, fig. 39). 151
 - Fruit bodies with a well defined parenchymatic wall
 (*cleistothecia*, fig. 46). 161
- 151(150) Gymnothecia with simple thin-walled, \pm uniform and poorly
 developed hyphae constituting the outer hyphal sheath. 152
 - Gymnothecia with thick-walled hyphae modified at their ends into
 appendages, or if thin-walled then always accompanied by
 appendages (i.e. curled, toothed or pointed hyphae). 155




Fig. 38. *Onygena corvina*, habit sketch, ascus and ascospore.

- 152(151) Gymnothecia red-orange to brick-red. Ascospores orange, subglobose to ellipsoid, with an equatorial furrow, smooth, $4.5-5.5 \times 3.5-4.5\mu\text{m}$. *Arachniotus ruber* (fig. 40)
- Gymnothecia white or yellow, never orange or brick-red. Ascospores without an equatorial furrow. 153
- 153(152) Gymnothecia white. Ascospores hyaline, ellipsoid, smooth, $3-4 \times 2-2.5\mu\text{m}$. *Arachniotus candidus*
- Gymnothecia distinctly pigmented, yellow or brown. Ascospores larger than $4\mu\text{m}$. 154
- 154(153) Gymnothecia yellow brown. Ascospores orange to brownish, slightly lenticular, smooth or slightly roughened, $5-6.5 \times 3.3-4.6\mu\text{m}$. *Arachniotus confluens*
- Gymnothecia lemon yellow. Ascospores lemon yellow, lenticular, smooth, $5-6 \times 3-4.5\mu\text{m}$. *Arachniotus citrinus*




Fig. 39. Habit sketch of a gymnothecium and ascus. Figs 40-45. Spores and peridial hyphae. Fig. 40. *Arachniotus ruber*. Fig. 41. *Myxotrichum chartarum*. Fig. 42. *Gymnoascus californiensis*. Fig. 43. *Gymnoascus reesii*. Fig. 44. *Ctenomyces serratus*. Fig. 45. *Arthroderma curreyi*.

- 155(151) Gymnothecia possessing only thick pigmented hyphae. 156
 - Gymnothecia possessing \pm thin, hyaline hyphae with only a few, although often distinctive, appendages (i.e. comb-shaped end cells or dumb-bell shaped asperulate cells accompanying twisted and bent hyphae). 160
- 156(155) Gymnothecia brown-black or dark greenish-grey, with external hyphae with spine-like branches and septate, hooked appendages. Ascospores orange brownish, ovate, delicately striate, $4-5.2 \times 2.4-3.3 \mu\text{m}$. *Myxotrichum chartarum* (fig. 41)
 - Gymnothecia never black, and, if possessing thick-walled hyphae, then appendages never septate. Ascospores smooth, or if ornamented then asperulate or echinulate. 157
- 157(156) Gymnothecia rose to orange-brown or yellowish. Appendages curved or irregularly branched and pointed, never verticillately branched. Ascospores smooth, or at most asperulate. 158
 - Gymnothecia red-brown with appendages verticillately branched. Ascospores $3-4.5 \times 2-2.8 \mu\text{m}$, yellowish brown, lenticular. *Actinodendron verticillatum*
- 158(157) Gymnothecia rosy pink when young, becoming browner, with spines and curved, non-septate hairs. Ascospores hyaline, globose to subglobose, asperulate, $3-5 \times 2.5-4 \mu\text{m}$. *Gymnoascus californiensis* (fig. 42)
 - Gymnothecia yellow. Ascospores smooth. 159
- 159(158) Gymnothecia yellow to yellow-brown, without elongated appendages but with thick-walled branches, few of which are pointed. Ascospores globose-ellipsoid, yellow to brownish, $3-4.5 \times 3.5 \mu\text{m}$. *Gymnoascus reesii* (fig. 43)
 - Gymnothecia golden yellow to reddish-brown, with acute-ended appendages. Ascospores lenticular, smooth, hyaline, $2.5-3.5 \times 2-2.5 \mu\text{m}$. *Pseudogymnoascus roseus*
- 160(155) Gymnothecia orange brown, with comb-like appendages. Ascospores slightly lenticular, pale orange, $3.3-3.6 \times 2-2.6 \mu\text{m}$. *Ctenomyces serratus* (fig. 44)
 - Gymnothecia whitish to pale ochraceous, particularly when dry, with few appendages but those present twisted and bent, and their branches constricted with regular or irregular dumb-bell shaped cells. Hyphal walls asperulate or with protuberances. Ascospores smooth, lenticular, hyaline, $2.4-3.3 \times 2 \mu\text{m}$. *Arthroderma curreyi* (fig. 45)

- 161(150) Asci relatively large, 100-200-spored, 1-3/fruit body.
 'Cleistothechia' minute, <100 (rarely <250) μ m diam., immersed.
see *Thelebolus* etc. (Key 1, 86)
 - Asci with 8 or fewer spores. 162
- 162(161) Ascospores purple at maturity, large, 50-70 \times 25-35 μ m, epispore
 with a few longitudinal cracks. see *Ascobolus immersus* (Key 1, 48)
 - Ascospores smaller, hyaline, yellow, olivaceous, brown or black. 163
- 163(162) Ascospores olivaceous, brown or black, at least in part. 164
 - Ascospores aseptate, hyaline, yellow or other pale colours. 174
- 164(163) Ascospores 4-celled (cf. *Sporormiella*), with germ slits, readily
 fragmenting. Asci clavate, bitunicate. Cleistothechia black, shiny,
 up to 500 μ m diam. 165
 - Ascospores 1- or 2-celled. 166
- 165(164) Ascus stalk up to 20 μ m long. Ascospores 25-32 \times 5 μ m.
Preussia vulgaris
 - Ascus stalk 30-60 μ m long. Ascospores 26-38 \times 5-7 μ m.
Preussia funiculata (fig. 47)
- 166(164) Ascospores 2-celled. 167
 - Ascospores 1-celled. 170
- 167(166) Spores unequally 2-celled, one brown ellipsoid, with an apical
 germ pore, 10-12 \times 6.5-7.5 μ m, the other a basal hyaline,
 cylindrical pedicel, 6-8 \times 3 μ m. Cleistothechia black, globose, up to
 250 μ m diam., covered with flexuous brown hairs up to 1mm long.
 Asci evanescent. *Zopfiella erostrata*
 - Spores equally 2-celled. 168
- 168(167) Spores not constricted at the septum, ellipsoid, golden-brown,
 25-30 \times 10-15 μ m with 1-3 guttules in each cell. Cleistothechia
 gregarious on a mycelial mat, whitish to pale orange, up to
 500 μ m diam. *Heleococcum aurantiacum* (fig. 48)
 - Spores hyaline, divided into two almost globose cells by the
 constricting septum. Ascomata superficial, globose, dark
 coloured. *(Mycoarachis)* 169

169(168) Asci 8-spored, 5.5-11 μ m diam. Spores 5-5.5 \times 3-3.5 μ m.

Mycoarachis inversa

- Asci 4-spored, 6-6.5 μ m diam. Spores 4.5-5 \times 2-2.5 μ m.

Mycoarachis tetraspora

170(166) Asci broad-clavate, (1)-2-(3)-spored, 30-50 \times 13-18 μ m. Spores brown-black with short ridges and warts, subglobose, 12-15.5 \times 11-12.5 μ m, with a single germ pore. *Copromyces bisporus* (fig. 49)

- Asci 8-spored.

171

171(170) Spores globose, sooty brown, 3 μ m diam. Cleistothecia gregarious, with basal spirally coiled appendages, black, 100-200 μ m diam., partially immersed in a white to red felty hyphal mat.

Pleuroascus nicholsonii

- Spores larger, ellipsoid or limoniform.

172




Fig. 46. Habit sketch of cleistothecia. Figs 47-54. Asci and spores. Fig. 47. *Preussia funiculata*. Fig. 48. *Heleococcum aurantiacum*. Fig. 49. *Copromyces bisporus*. Fig. 50. *Arachnomyces nitidus*. Fig. 51. *Orbicula parietina*. Fig. 52. *Roumegueriella rufula*. Fig. 53. *Aphanoascus stercoraria*. Fig. 54. *Pseudeurotium ovale*.

- 172(171) Spores olivaceous, limoniform, usually with an apical germ pore. Perithecia greyish or greenish, abundantly hairy, branched or simple, straight or curly. Asci pedicellate, soon disappearing. see *Chaetomium* at 5
- Spores darker, with 1 or more minute germ pores. Cleistothecia distinctly but not abundantly hairy. 173
- 173(172) Spores smoky brown, broadly ovoid, $9-14 \times 6-9\mu\text{m}$. Cleistothecial hairs short, up to $30\mu\text{m}$. *Thielavia wareingii*
- Spores dark brown, flattened limoniform, $13-16 \times 10-13 \times 8-9\mu\text{m}$. Cleistothecial hairs of two types, some smooth, dark brown, arising from the base up to 3mm long, others greyish green, rough, up to ca $120\mu\text{m}$. *Thielavia fimeti*
- 174(163) Cleistothecia produced within a common arachnoid mycelial mass. Spores smooth or minutely asperulate, yellow to yellow-brown, broadly ellipsoid, $4-5 \times 3-5\mu\text{m}$. *Aphanoascus fulvescens*
- Cleistothecia single or gregarious, but not on or in a mycelial mass. 175
- 175(174) Cleistothecia $170-750\mu\text{m}$ diam., covered with long (several mm when extended), thick-walled, aseptate, helical appendages. Asci clavate cylindrical, evanescent, $35-62 \times 12-21\mu\text{m}$. Spores ellipsoid, hyaline, $12-17 \times 9-12\mu\text{m}$. *Lasiobolium spirale*
- Cleistothecia without coiled appendages. 176
- 176(175) Cleistothecia with hairs or appendages. 177
- Cleistothecia smooth. 178
- 177(176) Cleistothecia black, shining, $100-200\mu\text{m}$ diam., with dark brown-black thick-walled hairs with hooked tips. Asci $8-15\mu\text{m}$ diam. Spores straw or copper coloured, ellipsoid, $4-7 \times 3.5-4.5\mu\text{m}$ with de Bary bubble and a germ pore at each end. *Kernia nitida*
- Cleistothecia reddish brown, less than 1mm diam., with long simple appendages curled at the tips. Spores hyaline, oblate, $3.5-5 \times 2-3\mu\text{m}$. *Arachnomyces nitidus* (fig. 50)
- 178(176) Ascospores globose, larger than $9\mu\text{m}$. 179
- Ascospores ellipsoid, up to $9\mu\text{m}$. Asci always subglobose. 180

- 179(178) Ascospores, smooth, 9-13 μ m. *Orbicula parietina* (fig. 51)
- Ascospores ornamented, 13-24 μ m. Asci subglobose. Cleistothecia
ochraceous, becoming yellowish brown or flushed cinnamon.
Roumegueriella rufula (fig. 52)
- 180(178) Ascospores hyaline, then faintly yellowish, minutely spiny, 2.5-3 \times
2-2.5 μ m. Cleistothecia pale, then dark brown.
Aphanoascus stercoraria (fig. 53)
- Ascospores hyaline, then brown, smooth, 5.5-6 \times 3.5-4 μ m.
Cleistothecia dark brown from the beginning.
Pseudeurotium ovale (fig. 54)

Key 3. Basidiomycota

- | | | |
|------|---|----|
| 1 | Basidia single-celled (fig. 55). | 2 |
| - | Basidia transversely or longitudinally septate (fig. 55), or difficult to observe. | 71 |
| 2(1) | Fruit body agaricoid, i.e. mushroom-shaped with gills underneath cap (figs 56, 67). | 3 |
| - | Fruit body not agaricoid, without gills (figs 65, 66). | 69 |
| 3(2) | Spore print white or pale coloured, hyaline s.m. (Usually on straw/dung mixtures, never on raw dung except when very old). | 5 |
| - | Spore print coloured. | 4 |
| 4(3) | Spore print pinkish or pale cinnamon, honey-coloured s.m. (Usually on straw/dung mixtures, never on raw dung). | 6 |
| - | Spore print darker, in shades of brown or black. | 8 |
| 5(3) | Stem eccentric. Fruit body pure white. Spores ellipsoid, smooth. <i>Pleurotellus s. lato</i>
(If gills pink and spores longitudinally ridged see <i>Clitopilus passackerianus</i> , fig. 67) | |
| - | Stem central | 7 |




Fig. 55. From left, sketches of holobasidium, with mature basidiospore showing germ pore; auriculariaceous basidium; tremellaceous basidium, lateral view and as often seen in sections.

- 6(4) Fruit body white, ivory or very pale tan, with a smell of cucumber. Gills decurrent. *Clitocybe augeana*
 - Fruit body yellow, with scaly cap. Gills free or just adnate. Fruit body with distinct ring and granular veil. (Commonly in plant pots. Probably associated with peaty material more than dung).
Leucocoprinus birnbaumii
 (*L. cepaestipes* and *L. lilacinogranulosus* occur in similar situations).
- 7(5) Fruit body with amethyst/purple shades, with eccentric stem. Spores subglobose, slightly ornamented to nearly smooth. (On compost heaps in gardens). *Lepista nuda*
 - Fruit body with pink gills and distinct volva at stem base. Cap white to pale hazel. Stem white. Spores broadly ellipsoid, smooth. *Volvariella speciosa*
- 8(4) Spore print distinctly brown (fulvous, tawny, rust coloured etc.). 9
 - Spore print some darker shade, fuscous, fuliginous or violaceous black. 20
- 9(8) Stem distinctly annulate, apex striate. *Conocybe percincta*
 (Has been found on straw/dung mixtures, never on raw dung).
 - Stem lacking a veil. 10
- 10(9) Cap rich chrome yellow, viscid, soon reduced to a sticky mass, easily collapsing. *Bolbitius vitellinus*
 - Cap in shades of brown, never brightly coloured and if collapsing then cap elongate-cylindric and white to pale cream. 11
- 11(10) Spore print dull, sepia or snuff-brown. On rabbit pellets in sand dunes. *Agrocybe subpediades*
 - Spore print brighter coloured, orange/ rust brown. (*Conocybe*) 12
- 12(11) Gill edge with irregularly fusoid cystidia with obtuse apices (lageniform). Cap viscid. *Conocybe coprophila*
 - Gill edge with distinctly capitate cells resembling a glass stoppered bottle (lecythiform). Cap never viscid, often pubescent under a lens. 13

- 13(12) Stem covered in long hairs. 14
 - Stem covered in lecythiform cells similar to those on gill edge, giving a farinaceous appearance under a lens. NEVER with long hairs. (Dung/straw mixtures). Large as in a *Cortinarius*. Spores smooth. *Conocybe intrusa*
 (*C. leucopus* has been found on manured soil in gardens; *C. antipus* has hexagonal spores and grows on dung piles).
- 14(13) Stem with both long hairs and lecythiform cystidia.  15
 - Stem with hairs and lageniform cystidia.  16
- 15(14) Spores 11-14 × 7-9µm. Taste and smell strong, of fresh meal. *Conocybe farinacea*
 - Spores large, over 15 × up to 10µm. Taste and smell none or slightly acidic. *Conocybe pubescens*
 (*C. subpubescens* might be found on straw/dung mixtures, and differs in spores 11-13 × 6-8µm).
- 16(14) Basidia 2-spored. *Conocybe rickenii*
 - Basidia 4-spored. 17
- 17(16) Spores ellipsoid. 18
 - Spores lentiform, angular in face view. *Conocybe lenticulospora*
- 18(17) Cap grey, contrasting with yellowish cream gills and pale stem. *Conocybe murinacea*
 - Spores 10.5-12.5 × 6-7µm. 19
 - Cap pinkish brown or tawny.
- 19(18) Spores 11-12 × 7.2-7.8µm. Cap sienna. On raw dung. *Conocybe fimetaria*
 - Spores 10-12 × 6-7µm. Cap pinkish to cinnamon brown. On manured soil or sewage sludge. *Conocybe fuscomarginata*
 (*Conocybe siennophylla* might be found on straw/dung mixtures or in soil in greenhouses. It differs in having smaller spores).
- 20(8) Cap deliquescing to some degree at maturity. Basidia of 2 or 3 different sizes. (*Coprinus*) 21
 - Cap not deliquescing. Basidia of one size only. 49
- 21(20) Veil on cap absent, cap either covered with small hairs (setules) or naked. 22
 - Cap covered with a granular, micaceous, powdery or fibrillar veil. 28




Fig. 56. Habit sketch of a stipitate agaric, *Psathyrella stercoraria*, with section. Fig. 57. Sketch of gill section of *Psathyrella* sp., showing position of marginal (m) and facial (f) cystidia. Fig. 58. *Coprinus pellucidus*, habit and vertical section of cap cuticle. Fig. 59. *C. pseudoradiatus*, habit and veil constituents. Fig. 60. *C. vermiculifer*, habit and veil constituents. Fig. 61. *C. filamentifer*, veil constituents. Fig. 62. *C. stercoreus*, habit. Fig. 63. *C. cordisporus*, vertical section of cap showing nature of veil cells on the cap cuticle. Fig. 64. Veil cells with structural (l) and superficial crystalline (r) ornamentation.

- 22(21) Cap without setules. 23
 - Cap with setules. 24
- 23(22) Cap minute, 1-5mm high before expanding, reddish orange at first, soon fading. Basidiospores almost globose to triangular in one view, elliptic in another, $7-10 \times 7-9 \times 5.5-6.5\mu\text{m}$. (2- and 4- spored forms have been found). *Coprinus miser*
 - Cap larger, up to 15mm when expanded. Basidiospores pip-shaped, $7.5-8.5 \times 9.5-11 \times 9.5-11.5\mu\text{m}$. (4-spored). *Coprinus nudiceps*
- 24(22) Spores hexagonal, $10-13 \times 6.5-7.5\mu\text{m}$. Cap purplish. *Coprinus hexagonosporus*
 - Spores ellipsoid. Cap brown or reddish, without purplish tints. 25
- 25(24) Basidia 4-spored. 26
 - Basidia 2-spored. Spores $11-13 \times 5.5-7\mu\text{m}$. Facial cystidia absent. *Coprinus bisporus*
 (*Coprinus sassii*, not yet recorded in British Isles, has 2-spored basidia with very large ellipsoid spores up to $20\mu\text{m}$ long).
- 26(25) Cap with a mixture of hyaline and brown thick-walled setules. Spores $9-10 \times 5.5-6\mu\text{m}$, with eccentric germ pore. Facial cystidia absent. *Coprinus heterosetulosus*
 - Cap with only one type of setule. Facial cystidia present or absent. 27
- 27(26) Facial cystidia present. Spores $7.9-13.3 \times 4.4-6.4\mu\text{m}$, with apical germ pore. *Coprinus stellatus*
 - Facial cystidia absent. Spores elongate and narrow, rarely greater than $5\mu\text{m}$ wide, with apical germ pore. Fruit body usually quite small, up to 6mm before expanding. *Coprinus pellucidus* (fig. 58)
 (Several species in the group, e.g. *C. congregatus* and *C. ephemerus* have been found on straw/dung mixtures).
- 28(21) Veil strongly adhering to cap. Spores elliptic ovate, $15-20 \times 8-12\mu\text{m}$. Stem with distinct ring. Usually on buried dung. *Coprinus sterquilinus*
 - Veil more floccose or powdery. Stem lacking ring or, if present (*C. ephemeroides*), fruit body small with 5-angled spores less than $10\mu\text{m}$ long. 29

- 29(28) Veil composed of filamentous units. 30
 - Filamentous units, if present, masked by a preponderance of rounded cells. 35
- 30(29) Veil composed of strings of sausage-shaped, thin-walled, hyaline cells. 31
 - Veil composed of rather narrow, slightly thickened hyphae. 32
- 31(30) Spores large, 11-14 × 6-7µm. Cap up to 1cm before expanding. Fruit body with or without a rooting base. *Coprinus radiatus*
 - Spores smaller, up to 9µm long. Cap up to 6mm before expanding. Fruit body without a rooting base. *Coprinus pseudoradiatus* (fig. 59) (*C. cinereus* is found on straw/dung mixture and *C. macrocephalus*, with large spores, has been recorded on raw dung).
- 32(30) Veil citrus- or lime-yellow, or a mixture of hyaline and brown strongly coloured hyphae. 33
 - Veil grey or whitish. 34
- 33(32) Veil of yellow hyphae. Spores 10.5-12.5 × 6-7.5µm. *Coprinus luteocephalus*
 - Veil with brown hyphae. Spores 7-9 × 3.5-5µm. *Coprinus poliommallus*
- 34(32) Veil hyphae thin-walled. Spores 6.5-7.5 × 5µm, 'shouldered' about the apiculus. *Coprinus filamentifer* (fig. 61)
 - Veil hyphae thin- and thick-walled, often with clamps. Spores elliptic-oblong, 9-10 × 5-6µm. *Coprinus vermiculifer* (fig. 60) (*Coprinus flocculosus*, with spores 11.5-16.5 × 6-9.5µm, can be found on straw/dung mixtures).
- 35(29) Stem with small, distinct ring. Spores subglobose to lentiform and 5-angled, 6-9 × 6.5-8 × 5-6µm. *Coprinus ephemeroides*
 - Stem at most with fibrils, even then rarely forming a faint ring zone. 36
- 36(35) Cap with setules in addition to veil. 37
 - Cap without setules. 38
- 37(36) Cap cystidia tapered. Spores 11-14 × 5-6.5µm. *Coprinus heptemerus*
 - Cap cystidia capitate. Spores 10-11 × 6-7µm. *Coprinus curtus*

- 38(36) Veil of inflated bladder-like cells attached to filamentous units.
Spores $7.5-8 \times 4.5-5.5\mu\text{m}$. *Coprinus utrifer*
- Veil of globose and subglobose cells and filamentous units often encrusted or with minute projections found sometimes at cap margin. 39
- 39(38) Globose cells, if ornamented then possessing crystalline or amorphous material (dissolved by 1N HCl, fig. 64). 40
- Globose cells covered in small fine blunt projections on the walls (not removed by 1N HCl, fig. 64). 45
- 40(39) Basidia 2-spored. 41
- Basidia 4-spored. 42
- 41(40) Spores $14-17 \times 8.5-10 \times 12.5-14\mu\text{m}$. *Coprinus pachyspermus*
- Spores smaller, $9-11 \times 6-6.5 \times 8-9\mu\text{m}$. *Coprinus cordisporus*
(2-spored form)
- 42(40) Spores less than $10\mu\text{m}$ long. *Coprinus cordisporus* (fig. 63)
(*C. patouillardii* is known on garden refuse, and an undescribed species with lemon-shaped spores has recently been found).
- Spores $10\mu\text{m}$ or more long. 43
- 43(42) Veil soon discolouring greyish, drab or buff. Spores $11.5-14.5 \times 6-8 \times 7.5-9\mu\text{m}$. *Coprinus cothurnatus*
- Veil remaining snowy white, only slowly discolouring greyish. 44
- 44(43) Fruit bodies several cm tall. Spores $15-19 \times 8.5-11.5 \times 11-13\mu\text{m}$. *Coprinus niveus*
- Cap small, 5-6mm at first. Spores $14-16 \times 8-9 \times 10-12.5\mu\text{m}$. *Coprinus latisporus*
- 45(39) Basidia 3-spored. 46
- Basidia 4-spored. 47
- 46(45) Spores narrow, $8.5-11 \times 5-6.2\mu\text{m}$. *Coprinus triplex*
- Spores broad, $9-10 \times 6-6.5 \times 6-7\mu\text{m}$, slightly flattened in face view. *Coprinus trisporus*
(These are possibly a single taxon).
- 47(45) Spores $7-8 \times 4-4.5\mu\text{m}$, perispore not visible in water or alkali mounts. *Coprinus stercoreus* (fig. 62)
- Spores $9\mu\text{m}$ or more long. 48

- 48(47) Spores 9-11 × 5.5-6µm. Perisporeal sac none or incomplete
or indistinct. *Coprinus foetidellus*
- Spores longer, 10.8-13.5 × 5.5-7µm, with distinct perispore with
dark lines and inclusions. Distinctive smell of gas. *Coprinus narcoticus*
(*C. sclerotiger* is found on straw/dung mixtures, and the smaller
C. tuberosus on garden refuse etc.).
- 49(20) Spores not discoloured in conc. H₂SO₄. 50
- Spores discolouring in conc. H₂SO₄. Gills not spotted at maturity. 56
- 50(49) Cap cuticle cellular. Gills spotted at maturity. (More often on rich,
'dungy', soils. *P. subbalteatus*, with copper coloured cap, drying
paler but retaining a dark marginal zone, occurs in gardens on
mulch etc.). (*Panaeolus*) 51
- Cap cuticle filamentous. 66
- 51(50) Velar remnants very obvious, either as an appendiculate veil or as
a distinct ring. 52
- Lacking all velar remnants. 54
- 52(51) Cap distinctly pigmented, with appendiculate veil. 53
- Cap pale coloured, smooth, semi-globate, soon cracking. Gills with
marginal cystidia only. *Panaeolus papilionaceus*
- 53(52) Cap brown, smooth, sometimes viscid, not exceedingly
wrinkled. *Panaeolus campanulatus*
- Cap grey, olivaceous, even black, with contrasting white
appendiculate veil. *Panaeolus sphinctrinus*
- 54(51) Cap with or without appendiculate veil, but always with distinct
ring. *Panaeolus semiovatus*
- Cap lacking veil. 55
- 55(54) Cap pinkish ochraceous to tawny-buff. Lacking facial
cystidia. *Panaeolus speciosus*
- Cap whitish or slightly yellowish. With facial cystidia.
Panaeolus antillarum

- 56(49) Gills with facial cystidia often containing yellow amorphous material when seen in ammonia solution or deep blue with cotton blue. (Stropharia) 57
 (Blue-green *S. cyanea* & *S. aeruginosa* often occur in rich garden soils).
 - Gills lacking facial cystidia. Never with yellowing cystidia in ammonia. (Psilocybe) 58
 (Red-capped *P. aurantia* can be found on straw/mulch mixtures in gardens).
- 57(56) Cap sticky, semi-globate ± expanding at maturity. On raw dung. *Stropharia semiglobata*
 - Cap plano-convex, often broad with a central umbo, margin flaring with age. On dungy mixtures in gardens. *Stropharia stercorearia*
- 58(56) Stipe bluing, with ring. Spores ellipsoid, 11-14 × 6.5-7.5µm. Fruit body with mealy smell and taste. *Psilocybe fimetaria*
 - Stipe lacking distinct ring, or if with ring or ring zone 2-spored and/or stem not bluing. Fruit body without mealy smell and taste. 59
- 59(58) Stem always with distinct ring. Basidia 2-spored. Spores 15-20µm long. *Psilocybe luteonitens*
 - Stem with or without ring. Basidia 4-spored. If with ring, spores smaller. 60
- 60(59) With ring zone. 61
 - Lacking velar remnants on stem, or only appendiculate teeth at cap margin. 62
- 61(60) Spores slightly angular/limoniform, 11-13(14) × 7-8µm. Often on sewage sludge. *Psilocybe merdaria*
 - Spores 13-14 × 7.5-8.5µm. *Psilocybe moelleri*
- 62(60) Spores 14-20 × 8-10µm. *Psilocybe subcoprophila*
 - Spores smaller. 63
- 63(62) Spores lentiform, angled, 6-8(8.5) × 4.5-5.5 × 3.75-4.5µm. *Psilocybe bullacea*
 (*P. crobula*, occasional on dung, differs in lacking purple colour in gills, and slightly smaller, ovoid, not angular, spores).
 - Spores larger. 64

- 64(63) Spores ellipsoid to slightly amygdaliform. *Psilocybe merdicola*
 - Spores lentiform, angular. 65
- 65(64) Spores 11-13(14) × 7-8(9)µm. see *Psilocybe merdaria*, 61
 - Spores 12-15 × 8-9.5µm. *Psilocybe coprophila*
- 66(50) Round cells on cap as a micaceous veil. (Re-examine gill face; if
 different sized basidia and facial cystidia separating the gills are
 present, go to *Coprinus* at 21). *Psathyrella sphaerocystis*
 - Cap lacking veil, or if present then fibrillar. 67
- 67(66) White copious veil at margin or also covering cap centre.
 Spores 10-12 × 5.5-6µm. *Psathyrella coprobia*
 - Lacking copious veil. 68
- 68(67) With red edge to gill. Spores 12-13 × 6-6.5µm, with central
 germ pore. *Psathyrella stercoraria*
 - Lacking red gill edge. Spores with eccentric germ pore.
Psathyrella coprophila
 (*P. fimetaria* differs in spore size: there are several members of the
P. prona group which grow on soil/straw mixtures).
- 69(2) Fruit body club-shaped. *Typhula setipes* (fig. 65)
 (*Clavaria acuta* often grows on peaty soil in pots in greenhouses).
 - Fruit bodies effuse, resupinate. 70
- 70(69) Fruit-body cobweb-like and greyish white. Basal hyphae 3-4.5µm
 wide. Spores sub-globose, 4.5µm diam. (Generally on old dung
 or straw/soil mixtures). *Athelia coprophila*
 (If with spiny spores 5-6µm diam., see the recently recorded
Tomentellopsis echinospora).
 - Fruit-body with pores, white or flushed slightly ochraceous,
 brownish or greyish. (On clods of soil in dunged land).
Cristella candidissima
- 71(1) Fruit body either a cup containing several 'eggs' or a single orange
 or yellowish gelatinous sphere. 72
 - Fruit-body effuse, without distinct shape. 73

- 72(71) Fruit-body whitish or pale yellow, up to 2.5mm diam., splitting at maturity to shoot away the orange/yellow spore mass.
Sphaerobolus stellatus (fig. 66)
- Fruit-body cup shaped, with silvery-grey 'eggs'. (Usually on dung and straw or attached to rabbit pellets).
Cyathus stercoreus
(Cyathus vernicosus often grows in plant pots on rich soil).
- 73(71) Basidia with transverse septa. Spores $11 \times 7\mu\text{m}$. Fruit body pinkish.
Platyglea fimicola
 (Not British; included for completeness. *Pilacrella solani*, with a glistening stipitate head, has been isolated from dungy soil).
- Basidia with longitudinal septa. Spores $14-18 \times 9-10\mu\text{m}$.
 Fruit body cream-white or ivory.
Sebacina incrustans





Fig. 65. Habit sketch of *Typhula* sp. Note attachment to sclerotium. **Fig. 66.** *Sphaerobolus stellatus*, habit. **Fig. 67.** *Clitopilus passackerianus*, a sessile agaric - habit sketch and section.

Key 4. Zygomycota

- 1 Spores formed in multispored sporangia (figs 68, 70, 72, 75, 76) or in few-spored sporangioles (figs 70, 73). 2
- Multispored sporangia and globose sporangioles absent. Spores formed singly on terminal, lateral or intermediate vesicles (figs 74, 79, 80, 82-86), or in short chains (figs 77, 78, 81). 11
- 2(1) Sporangiphore stout, simple, with a subsporangial swelling and a basal swelling buried in the substrate. Sporangia tough walled, black, projected some distance towards the light when mature, and sticking to whatever they hit. *Pilobolus* (fig. 76)
 e.g. spores pale yellow, 8-10 × 5-6µm - *P. crystallinus*
 spores orange, 12-20 × 6-10µm - *P. kleinii*
- Sporangiphores not stout; sporangia not violently discharged. 3
- 3(2) Sporangial wall black, tough, not readily broken when touched. Sporangia with a sticky base, becoming attached to whatever they contact after the marked elongation of the white sporangiophores at maturity. *Pilaira* (fig. 75)
 e.g. spores yellowish, 8-10 × 6µm - *P. anomala*
 spores colourless, 11-13 × 6-8µm - *P. moreaui*
- Sporangial wall diffluent, spores readily removed in a droplet, or fragile and then spores easily dispersed by external violence. 4
- 4(3) Sporangiphores stiff and metallic in appearance, growing towards the light and often to great length (5-30cm). *Phycomyces*
 e.g. spores 10.5-30 × 6.5-17µm; columella pyriform; sporangiophores up to 30cm - *P. nitens*
 spores 8-13 × 5-7.5µm; columella spherical or ovoid; sporangiophores up to 30cm - *P. blakesleeanus*
- Sporangiphores white, not reaching extreme lengths. 5
- 5(4) Small lateral sporangia (sporangioles) present. 10
- Sporangioles absent. 6
- 6(5) Sporangiphores usually grouped, less often single, connected by stolon-like hyphae. 7
- Sporangiphores arising singly, or if grouped then lacking stolon-like hyphae. 9

- 7(6) Stolons joining groups of sporangiophores often with rhizoids at the base of the group. 8
- Sporangiophores arising singly or in groups from stolons, which may be 'rooted' at intervals along their length, but rarely beneath the groups of sporangiophores. *Absidia* (fig. 71)
 - e.g. sporangiophores grouped, rhizoids poorly developed; spores 2.5-4.5 μ m diam. - *A. corymbifera*
 - sporangiophores grouped, rhizoids strongly developed; spores 2.5-3.5 μ m diam. - *A. orchidis*
- 8(7) Sporangiophores mostly unbranched. *Rhizopus* (fig. 69)
- e.g. spores irregularly angular-ovoid, 8-14 \times 11 μ m - *R. nigricans*
 - Sporangiophores with a whorl of branches beneath the main sporangium, each with a small columellate sporangium. Spores 6-8.5 μ m. *Actinomucor elegans*

Fig. 68. *Mucor*, habit and detail of sporangium before and after dehiscence. **Fig. 69.** *Rhizopus*, habit. **Fig. 70.** *Thamnidium elegans*, habit and detail of sporangioles. **Fig. 71.** *Absidia*, habit. **Fig. 72.** *Mortierella*, habit and sporangiophore tip after sporangial dehiscence. **Fig. 73.** *Helicostylum*, sporangioles. **Fig. 74.** *Chaetocladium*, sporangioles. **Fig. 75.** *Pilaira*, sporangiophores before and after elongation, and sporangium. **Fig. 76.** *Pilobolus*, sporangiophore. **Fig. 77.** *Syncephalis*, habit, sporangiophore and merosporangia. **Fig. 78.** *Piptocephalis*, habit and detail of final branch with head cell and merosporangia. **Fig. 79.** *Oedocephalum*, habit and sporing head. **Fig. 80.** *Rhopalomyces*, sporing head. **Fig. 81.** *Syncephalastrum*, habit and detail of merosporangium. **Fig. 82.** *Coemansia*, habit, sporoclade with sporangia and sporangium with spore inside. **Fig. 83.** *Kickxella*, habit and sporoclade. **Fig. 84.** *Cunninghamella*, habit and fertile head. **Fig. 85.** *Mycotypha* (l) and *Ostracoderma* (r) conidiophores. **Fig. 86.** *Ballocephala*, habit of sporangiophores growing from parasitised tardigrade, sporangiophore and sporangia.



- 9(6) Sporangia often with pigmented walls, yellowish when young, finally grey or black, with well marked columella left after spore dispersal. Individual sporangiophores observable with unaided eye, up to 20mm long. *Mucor* (fig. 68)
- e.g. spores smooth, $7-8 \times 2.5-4.5\mu\text{m}$ - *M. hiemalis*
 - spores smooth, $6-12 \times 3-6\mu\text{m}$ - *M.ucedo*
 - spores asperulate, $5-8.5\mu\text{m}$ diam. - *M. plumbeus*
- (N.B. *Zygorhynchus* would key out with *Mucor*. It is more often isolated from soil, and is distinguished from *Mucor* by the presence of zygosporangia with unequal suspensors)
- Sporangia white, without a columella, readily becoming a spore droplet. Sporangiophores delicate, often only $200-400\mu\text{m}$ long. Fine, white, garlic-smelling mycelium often present. *Mortierella* (fig. 72)
 - e.g. spores $16-27\mu\text{m}$ diam., few in each sporangium; sporangiophores *ca* $150\mu\text{m}$, with short lateral branches at right angles - *M. reticulata*
 - spores $6-10 \times 4-6\mu\text{m}$; sporangiophores 2-3mm high, with ascending branches - *M. bainieri*
 - spores $4-10\mu\text{m}$; sporangiophores richly branched - *M. candelabrum*
- 10(5) Sporangioles formed at the final tips of a densely dichotomous system of branchlets, originating some distance below a terminal sporangium (which may be absent in young specimens). Sporangioles up to $25\mu\text{m}$ diam., with up to 6 spores. Spores $8-12 \times 6-8\mu\text{m}$. *Thamnidium elegans* (fig.70)
- Sporangioles either at the curved tips of slender branches, or clustered in groups about halfway along tapering branches which radiate from the sporangiophore below the sporangium; the branch tips of the latter give the fertile portion of the sporangiophore a bristly appearance. *Helicostylum* (fig. 73)
 - e.g. spores $8-17 \times 3-7\mu\text{m}$; sporangioles on short secondary or tertiary branches; fertile region bristly with sterile branches - *H. fresenii*
 - spores $6-8 \times 4\mu\text{m}$; sporangioles reflexed, on slender primary or secondary branches; fertile region without sterile branches - *H. pyriforme*
- 11(1) Spores formed in chains. 12
- Spores formed singly. 14

- 12(11) Sporangiohores regularly and repeatedly dichotomously branched. Chains of 2-10 spores produced in small groups, which may be wet or dry, on deciduous heads, 4-15µm diam. Parasitic on other fungi, mostly other Mucorales. *Piptocephalis* (fig. 78)
- e.g. spores 4-5 × 2-3µm, in pairs; heads dry - *P. lepidula*
 spores 5-6 × 2-2.5µm, in chains of 4-9; heads dry - *P. cylindrospora*
 spores 4-8 × 2-4µm, in chains of 3-5; heads dry;
 sporangiophore without rhizoids - *P. freseniana*
 spores 4-6 × 4-4.5µm, in chains of 3-6; heads wet;
 sporangiophore with rhizoids - *P. repens*
 spores 3-5 × 2-2.5µm, in chains of 3-5, heads wet; head cell lyses, to leave only a fringe at the tip of the very fine sporangiophore - *P. fimbriata*
- Sporangiohores simple or irregularly branched. 13
- 13(12) A large conspicuous fungus, macroscopically Mucor-like, mycelium coarse. Sporangiohores with a distinct terminal swelling with crowded spore chains. Spores usually 5-10 in a chain, globose to ovoid, 2-8 × 4-6µm. *Syncephalastrum racemosum* (fig. 81)
- Sporangiohores less conspicuous, 100-1000µm high, with a 'holdfast' at the base attaching the sporangiophore to the substrate. Mycelium very fine. Parasitic on other Mucorales. *Syncephalis* (fig. 77)
- e.g. sporangiophores 100-200µm high, with three 'nodes' along their length; merosporangia often forked at the basal cell; spores 8-10 × 6µm - *S. nodosa*
 sporangiophores up to 750µm high; merosporangia usually subdivided at their base into several branches, each with 5-10 spores; spores 5-10 × 3-4µm - *S. depressa*
 (N.B. *Oedocephalum* spp. (fig. 79), the anamorphic states of many dung fungi (esp. Ascobolaceae and Pezizaceae), *Rhopalomyces* (fig. 80), and some *Aspergillus* spp. are superficially similar to *Syncephalis* at first sight).
- 14(11) Sporangia containing a single closely fitting elongated spore, produced in serried ranks on one side of a boat-shaped branch (sporoclade). 15
- Single-spored sporangia ('spores') globose, produced singly or in groups not on sporoclades. 16



Published by the British Mycological Society
© British Mycological Society 1997
ISBN 0 9527704 2 3