

doi.org/10.3114/fuse.2021.07.02

## A higher-rank classification for rust fungi, with notes on genera

M.C. Aime<sup>1\*</sup>, A.R. McTaggart<sup>2</sup>

<sup>1</sup>Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN 47907, USA

<sup>2</sup>Queensland Alliance for Agriculture and Food Innovation, The University of Queensland, Brisbane 4001, Australia

\*Corresponding author: maime@purdue.edu

### Key words:

host alternation

life cycles

*Uredinales*

*Urediniomycetes*

37 new taxa

**Abstract:** The rust fungi (*Pucciniales*) with 7000+ species comprise one of the largest orders of *Fungi*, and one for which taxonomy at all ranks remains problematic. Here we provide a taxonomic framework, based on 16 years of sampling that includes *ca.* 80 % of accepted genera including type species wherever possible, and three DNA loci used to resolve the deeper nodes of the rust fungus tree of life. *Pucciniales* are comprised of seven suborders – *Araucariomycetinae* subord. nov., *Melampsorinae*, *Mikronegeriinae*, *Raveneliinae* subord. nov., *Rogerpetersoniinae* subord. nov., *Skierkineae* subord. nov., and *Urediniinae* – and 18 families – *Araucariomycetaceae* fam. nov., *Coleosporiaceae*, *Crossosporaceae* fam. nov., *Gymnosporangiaceae*, *Melampsoraceae*, *Milesinaceae* fam. nov., *Ochrosporaceae* fam. & stat. nov., *Phakosporaceae*, *Phragmidaceae*, *Pileolariaceae*, *Pucciniaceae*, *Pucciniastriaceae*, *Raveneliaceae*, *Rogerpetersoniaceae* fam. nov., *Skierkaceae* fam. & stat. nov., *Sphaerophragmiaceae*, *Tranzscheliaceae* fam. & stat. nov., and *Zaghouaniaceae*. The new genera *Araucariomyces* (for *Aecidium fragiforme* and *Ae. balansae*), *Neolivea* (for *Olivea tectonae*), *Rogerpetersonia* (for *Caeoma torreyae*), and *Rossmatomyces* (for *Chrysomyxa monesis*, *Ch. pryrolae*, and *Ch. ramischiae*) are proposed. Twenty-one new combinations and one new name are introduced for: *Angiopsora apoda*, *Angiopsora chusqueae*, *Angiopsora paspalicola*, *Araucariomyces balansae*, *Araucariomyces fragiformis*, *Cephalotelium evansii*, *Cephalotelium neocaledoniense*, *Cephalotelium xanthophloeae*, *Ceropsora weirii*, *Gymnotelium speciosum*, *Lipocystis acaciae-pennatulae*, *Neolivea tectonae*, *Neophysopella kraunhia*, *Phakospora pipturi*, *Rogerpetersonia torreyae*, *Rossmatomyces monesis*, *Rossmatomyces pryrolae*, *Rossmatomyces ramischiae*, *Thekopsora americana*, *Thekopsora potentillae*, *Thekopsora pseudoagrimoniae*, and *Zaghouania notelaeae*. Higher ranks are newly defined with consideration of morphology, host range and life cycle. Finally, we discuss the evolutionary and diversification trends within *Pucciniales*.

**Citation:** Aime MC, McTaggart AR (2020). A higher-rank classification for rust fungi, with notes on genera. *Fungal Systematics and Evolution* 7: 21–47. doi: 10.3114/fuse.2021.07.02

**Received:** 10 August 2020; **Accepted:** 30 October 2020; **Effectively published online:** 13 November 2020

**Corresponding editor:** P.W. Crous

## INTRODUCTION

Rust fungi (*Pucciniomycotina*, *Pucciniales*) comprise one of the largest orders in *Fungi*, containing *ca.* 25 % of described *Basidiomycota*. All are obligate pathogens of plants and at *ca.* 7 000+ accepted species (Kirk *et al.* 2008) form the most species-rich group of plant pathogens. Diseases caused by rust fungi have impacted human agriculture and history through time. Rusts likely caused the earliest recognized diseases of agricultural plants (Carefoot & Sprott 1967), and have continued to impact anthropogenic ecosystems through epidemics and localized host extinctions (Carnegie & Pegg 2018). The Green Revolution in the mid to late 20<sup>th</sup> century that heralded the era of host resistance breeding targeted rust fungi (Philips 2013).

*Pucciniales* has a suite of characteristics that are rare or unique within *Fungi*, including alternation of generations with separate gametothalli (spermatogonia and aecia) and sporothalli (uredinia and telia) that may infect unrelated hosts (heteroecious); and the production of up to five different morphs within the life cycle. These characteristics, together with many

instances of convergent evolution within morphs, repeated evolution of derived life cycle variants, and varying taxonomic emphases on different morph characteristics, have contributed to the development of numerous classification schemes for rust fungi (Fig. S1). Further taxonomic confusion within *Pucciniales* at the species rank has been shaped by separate naming systems under prior nomenclatural codes for sexual and asexual morphs. For instance, prior to the use of molecular data to link morphs, only through painstaking inoculation studies could complete life cycles be elucidated (*e.g.*, Cummins 1978). Consequently, many asexual morphs were unplaceable within a sexual morph-based classification system. Recent changes to the nomenclatural code now allow the placement of taxa within natural genera, regardless of morph (McNeill *et al.* 2012, Turland *et al.* 2018). Although most asexual genera have been reduced to synonymy (Aime *et al.* 2018b), some, such as *Uredo* and *Aecidium* contain species that occur in over 50 sexual genera, and it will be non-trivial to assign these to natural genera.

Generic-rank classification, even for sexual morph species, is similarly difficult. At least 334 generic names have been described

in *Pucciniales*; most researchers accept *ca.* 130 of these (*e.g.*, Cummins & Hiratsuka 2003). Studies have shown that many diagnostic characters are homoplasious, such as the number of cells per teliospore (Aime 2006, Maier *et al.* 2007, van der Merwe *et al.* 2007, Yun *et al.* 2008, Beenken & Wood 2015). As a result, most taxon-rich genera – the largest being *Puccinia* (*ca.* 4 000 species), *Uromyces* (*ca.* 800 species), and *Ravenelia* (*ca.* 200 species) – are polyphyletic and will need thoughtful re-evaluation for how to reassign these species into monophyletic genera.

At the higher ranks, classification of rust fungi has varied through time as well (Fig. S1). Rust fungi were initially classified into families by characteristics of basidia and teliospores (*e.g.*, Cunningham 1931). This approach divided rusts into three (or four) families, *Melampsoraceae*, (*Coleosporiaceae*), *Pucciniaceae* and *Zaghouaniaceae* (Sydow & Sydow 1915, Cunningham 1931). Arthur (1907–1931), Sydow & Sydow (1915) and Dietel (1928) further classified rusts in subfamilies or tribes based on morphology of telia. Other workers, such as Hiratsuka & Cummins (1963) placed greater emphasis on the gametothallus, especially spermogonial morphology, resulting in conflicting taxonomic hypotheses. This approach was later combined with teliospore morphology (Cummins & Hiratsuka 1983, 2003) to achieve a 13-family classification that became the most broadly applied in the pre-molecular era.

The first molecular systematic study to test the familial classification of Cummins & Hiratsuka (2003) subdivided the rust fungi into three major radiations, *Mikronegeriineae*, *Melampsorineae*, and *Uredinineae*, that mostly correspond to the earlier three-family approach of Cunningham (1931) (Aime 2006). Within these radiations were (i) several lineages more or less corresponding to families circumscribed by Cummins & Hiratsuka (2003), such as *Coleosporiaceae*, *Melampsoraceae*, *Zaghouaniaceae* (as *Mikronegeriaceae*), *Phragmidiaceae*, *Pileolariaceae*, *Pucciniaceae*, *Pucciniastraceae* and *Raveneliaceae*; (ii) families, such as *Chaoniaceae* and *Phakopsoraceae* that were comprised of polyphyletic assemblages that could not be effectively resolved without data from type species; and (iii) several so-called “orphan” genera that could not confidently be assigned to families (Aime 2006).

Numerous subsequent studies have focused on resolution of single families, *e.g.*, *Sphaerophragmiaceae* (Beenken 2017); polyphyletic genera, *e.g.*, segregation of *Neophysopella* from *Phakopsora* (Ji *et al.* 2019); as well as conservation efforts to stabilize use of generic names (*e.g.*, Aime *et al.* 2018b, 2019a, b). Despite these efforts, a stable and resolved higher-rank classification for the rust fungi has not been achieved. A major bottleneck has been limited sampling of taxa that represent the type species of genera, especially for genera with convergent morphologies, that are polyphyletic, and/or contain species with multiple competing names for different morphs.

The purpose of the present study is to provide a stable higher-rank classification for *Pucciniales* that will serve as a framework for future systematic studies. We have assembled a dataset over the last 16 years that includes exemplars from 113 (*ca.* 80 %) rust genera, including 108 that are represented by sequences from type species (86) or type species proxies (22). Our phylogenetic hypotheses are based on DNA data from three loci (nuclear large subunit and small subunit rDNA, and Cytochrome-c-oxidase subunit 3) with varying evolutionary rates across *Pucciniales* (*e.g.*, Aime 2006, Vialle *et al.* 2009, Feau *et al.* 2011, Aime *et al.* 2018a, McTaggart & Aime 2018). We propose a natural classification for *Pucciniales* based on

combined evidence from morphology, life cycles, hosts, and phylogenetic data. Several new suborders, families, genera, and combinations are proposed, and suborders and families are redefined. Finally, we discuss the evolutionary trends that led to diversification within *Pucciniales* and highlight unresolved areas of the rust family tree for future research.

## MATERIALS AND METHODS

### Taxon selection

Priority was given to species that represent generic types of rust fungi. If type species were unavailable, wherever possible two congeneric species similar to the type in respect to host genus, morphology, and geography were chosen as proxies (*e.g.*, *Skierka*, *Tranzschelia*, *Uredopeltis* and *Uropyxis*). At least one exemplar was included for every major lineage of *Pucciniales* that had been previously identified (*e.g.*, Aime 2006, 2018a, Beenken 2017). Additional genera were targeted (i) from families that appeared polyphyletic in prior studies (*e.g.*, *Chaoniaceae*, *Phakopsoraceae*); (ii) from previously undersampled families, *e.g.*, *Uropyxidaceae*; and (iii) to broaden sampling of endocyclic species (*e.g.*, *Baeodromus*, *Chardoniella*, *Cionothrix*, *Dietelia*, *Puccinosira*). If possible, more than one species was included for genera (i) previously determined as orphaned taxa *sensu* Aime (2006) (*e.g.*, *Gymnosporangium*, *Prospodium*, *Ochropsora*, and *Tranzschelia*) or *incertae sedis sensu* Cummins & Hiratsuka (2003) (*Elateraecium*, *Masseëlla*); and (ii) previously demonstrated as polyphyletic (*e.g.*, *Maravalia*, *Phakopsora*, *Pucciniastrum*, *Ravenelia*). Additional taxa were also included for genera if complete data at the three sequenced loci were available (*e.g.*, *Gymnosporangium*, *Hamaspora*, *Melampsora*, *Neophysopella*, and *Phragmidium*). An initial dataset of 130 rust taxa and three loci (Table 1) was used to determine the familial placement of genera and the relationships between families in an overview tree. The overview tree was rooted with *Eocronartium muscicola*, from the sister order to *Pucciniales* (Aime *et al.* 2006).

With the overview tree as a guide, we divided the data into three subsets, *Melampsorineae* (73 species), *Raveneliineae* (77 species) and *Uredinineae* (164 species), for additional sampling and analyses (Table S1). In expanded sampling, we included taxa only sequenced for one of the three loci in order to broaden both generic representation and species representation for polyphyletic genera. Trees were rooted from the sister lineage as shown by the overview tree, or, in the case of *Raveneliineae*, midpoint rooted.

### DNA extraction, PCR and sequencing

DNA was extracted from fresh or herbarium material with the UltraClean Plant DNA Isolation Kit (MoBio Laboratories Inc., Solana Beach, CA, U.S.A.). The nuclear large subunit (28S) region of the ribosomal DNA repeat was amplified with Rust2INV (Aime 2006)/LR6 or LR7 (Vilgalys & Hester 1990) and, for weak products, nested with Rust28SF (Aime *et al.* 2018a)/LR5 or LR6 (Vilgalys & Hester 1990) following the protocols of Aime *et al.* (2018a). The small subunit (18S) region of the ribosomal DNA repeat was amplified with NS1 (White *et al.* 1990)/Rust 18S-R (Aime 2006) and nested with RustNS2-F (Aime *et al.* 2018a)/NS6 (White *et al.* 1990) following the protocols of Aime *et al.* (2018a). Cytochrome-

Table 1. Collection and accession data for sequences used in *Pucciniales* overview tree (Fig. 1).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	28S	18S	CO3	Host	Source
<i>Achroetium ichnocarpi</i>	T	BRIP 55685	KT199393	KT199381	KT199404	<i>Ichnocarpus frutescens</i>	McTaggart et al. (2016)
<i>Aecidium kalanchoe</i>		BPI 843633 (U18, HOLOTYPE)	AY463163	DQ354524	NA	<i>Kalanchoe blossfeldiana</i>	Hernandez et al. (2004)
<i>Allodus podophylli</i>	T	BPI 842277 (U2, NEOTYPE); 28S, 18S; PUR N16753:CO3	DQ354543	DQ354544	MG907270	<i>Podophyllum peltatum</i>	Aime (2006); Aime et al. (2018a)
<i>Angiopsora paspalicola</i>	*	BRIP 55625	<b>MW049243</b>	NA	<b>MW036496</b>	<i>Paspalum</i> sp.	this paper
<i>Aplopsora nyssae</i>	T	BPI 877823 (U1191)	<b>MW049244</b>	NA	NA	<i>Nyssa sylvatica</i>	this paper
<i>Araucariomyces fragiformis</i>	T	BRIP 68996	<b>MW049245</b>	<b>MW049292</b>	<b>MW036497</b>	<i>Agathis robusta</i>	this paper
<i>Austropuccinia psidii</i>	T	BRIP 58164	KF318449	KF318457	KT199419	<i>Rhodammia angustifolia</i>	Pegg et al. (2014); McTaggart et al. (2016)
<i>Baeodromus eupatorii</i>	*	PUR N16312 (U1386)	<b>MW049246</b>	NA	NA	<i>Ageratina</i> sp.	this paper
<i>Bibulocystis pulcherrima</i>	T	BRIP 58450	<b>MW049247</b>	NA	<b>MW036498</b>	<i>Daviesia latifolia</i>	this paper
<i>Blastospora smilacis</i>	T	PUR N270	DQ354568	DQ354567	NA	<i>Smilax sieboldii</i>	Aime (2006)
<i>Bubakia argentinensis</i> (as <i>Phakopsora argentinensis</i> )	*	ZT:RB 8248	KF528009	NA	NA	<i>Croton</i> cf. <i>anisodontus</i>	Beenken (2014)
<i>Calyptospora goeppertiana</i>	T	BPI 882188 (U866)	<b>MW147023</b>	NA	NA	<i>Abies balsamea</i>	this paper
<i>Catenulopsora flacourtae</i>	T	PUR N13865 (U669)	<b>MW049248</b>	<b>MW049293</b>	NA	<i>Flacourtia indica</i>	this paper
<i>Cephalotelium macowaniana</i> (as <i>Ravenelia macawaniana</i> )	T	PREM 61222	MG946007	NA	NA	<i>Vachellia karroo</i>	Ebinghaus et al. (2018a)
<i>Cephalotelium neocaledoniense</i> (as <i>Ravenelia neocaledoniensis</i> )		BRIP 56908	KJ862348	NA	KJ862460	<i>Vachellia farnesiana</i>	McTaggart et al. (2015)
<i>Ceratocoma jacksoniae</i>	T	BRIP 57717	KT199394	KT199382	KT199405	<i>Daviesia</i> sp.	McTaggart et al. (2016)
<i>Ceropsora weirii</i> (as <i>Chrysomyxa weirii</i> )	*	916CHWPCGS68	FJ666465	NA	NA	n.d.	Vialle et al. (2009)
<i>Chaonia ingae</i>	*	BPI 863575 (GUY74)	<b>MW049249</b>	NA	NA	<i>Inga</i> sp.	this paper
<i>Chardonella gynoxidis</i>	T	R15	<b>MW049250</b>	NA	NA	<i>Gynoxys</i> sp. (cf.)	this paper
<i>Chrysocelis lupini</i>	T	PUR N11562 (U1570)	<b>MW049251</b>	NA	NA	<i>Lupinus</i> sp.	this paper
<i>Chrysomyxa arctostaphylli</i>		CUW CFB 22246	AF522163	AY657009	NA	n.d.	Matheny et al. unpublished
<i>Cionothrix praelonga</i>	T	PUR 90104	<b>MW049252</b>	NA	NA	<i>Eupatorium</i> sp.	this paper
<i>Coleopuccinia sinensis</i>	T	BJFC R02506	MF802285	NA	NA	<i>Cotoneaster microphyllus</i>	Cao et al. (2018)
<i>Coleosporium senecionis</i>	T	PDD 98309	KJ716348	KJ746818	NA	<i>Senecio</i> sp.	Padamsee & McKenzie (2014)
<i>Cronartium flaccidum</i>	T	PUR N16561 (MCA4165)	<b>MW049253</b>	<b>MW049294</b>	NA	<i>Vincetoxicum hirsutinaria</i>	this paper
<i>Cronartium harknessii</i> (≡ <i>Endocronartium harknessii</i> )	(T)	CFB22250	AF522175	AY665785	NA	<i>Pinus</i> sp.	Szaro & Bruns unpublished; Matheny et al. unpublished
<i>Crossopora fici</i>		BRIP 58118	<b>MH047207</b>	<b>MH047212</b>	<b>MH047204</b>	<i>Ficus virens</i> var. <i>sublancoolata</i>	this paper

Table 1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	28S	18S	CO3	Host	Source
<i>Crossopora ziziphi</i>	T	BPI 877877 (U904)	MG744558	NA	NA	<i>Ziziphus mucronata</i>	Souza et al. (2018)
<i>Cumminsiiella mirabilissima</i>	T	BPI 871101 (U480)	DQ354531	DQ354530	NA	<i>Mahonia aquifolium</i>	Aime (2006)
<i>Dasyospora gregaria</i>	T	ZT Myc 3397	JF263477	JF263502	JF263518	<i>Xylophia cayennensis</i>	Beenken et al. (2012)
<i>Desmella aneimiae</i>	T	BRIP 60995	KM249867	NA	NA	<i>Nephrolepis hirsutula</i>	McTaggart et al. (2014)
<i>Diaphanopellis purpurea</i>	*	BJFC R02448	MK874622	NA	NA	<i>Picea brachytyla</i>	Yang & Wang unpublished
<i>Didymopora solani-argentei</i>	T	PUR N3728	<b>MW049254</b>	NA	NA	<i>Solanum argenteum</i>	this paper
<i>Dietelia codiae</i>	*	PUR N16488	<b>MW049255</b>	NA	NA	<i>Codiaeum variegatum</i>	this paper
<i>Dipyxis mexicana</i>	T	BPI 871906	<b>MW049256</b>	NA	NA	<i>Adenocalymna</i> sp.	this paper
<i>Edythea quitensis</i>	T	QCAM6453	MG596499	NA	NA	<i>Berberis hallii</i>	Barnes & Ordonez unpublished
<i>Elateraecium salaciicola</i>	T	PUR F17677	<b>MW049257</b>	<b>MW049295</b>	NA	<i>Salacia</i> sp.	this paper
<i>Endophylloides portoricensis</i> (as <i>Dietelia portoricensis</i> )	T	BPI 844288 (U322):28S; n.d.:18S	DQ354516	AY125389	NA	<i>Mikania micrantha</i>	Aime (2006); Wingfield et al. (2004)
<i>Endophyllum cassiae</i>	T	BPI 871369 (U525)	<b>MW049258</b>	NA	NA	<i>Cassia obtusifolia</i>	this paper
<i>Endophyllum circumscriptum</i>	T	BPI 872271	<b>MW049259</b>	NA	NA	<i>Cissus</i> sp.	this paper
<i>Endoraecium acaciae</i>	T	BPI 871098 (MCA2957)	DQ323916	DQ323917	NA	<i>Acacia koa</i>	Scholler & Aime (2006)
<i>Econartium muscicola</i>	*	MIN796447:28S; DUKE:DAH(e1):18S	AF014825	DQ241438	NA	NA	Bruns & Szaro unpublished; Henk & Vilgalys (2007)
<i>Gerwasia rubi</i>	T	BRIP 58440	KT199397	NA	KT199408	<i>Rubus</i> sp.	McTaggart et al. (2016)
<i>Gymnoconia interstitialis</i>	T	BPI 747600	JF907677	DQ521422	NA	<i>Rubus allegheniensis</i>	Yun et al. (2011); Matheny et al. unpublished
<i>Gymnosporangium clavariiforme</i> ( <i>=Podisoma clavariiforme</i> )	(T)	BRIP 59471	<b>MW049261</b>	<b>MW049296</b>	<b>MW036499</b>	<i>Crataegus</i> sp.	this paper
<i>Gymnosporangium sabiniae</i>	T	TNM F0030477	KY964764	KY964764	NA	<i>Pyrus communis</i>	Shen et al. (2018)
<i>Gymnotelium blasdaleanum</i>	*	PUR N10018 (U1469)	MG907218	MG907206	MG907269	<i>Amelanchier alnifolia</i>	Aime et al. (2018a)
<i>Hamaspora acutissima</i>	T	BRIP 56949	KT199398	KT199385	KT199409	<i>Rubus moluccanus</i>	McTaggart et al. (2016)
<i>Hamaspora longissima</i>	T	BPI 871506 (U305)	<b>MW049262</b>	<b>MW049297</b>	NA	<i>Rubus ludwigii</i>	this paper
<i>Haplophragmium derridis</i>	T	PUR N16494	<b>MW049263</b>	NA	NA	unidentified <i>Fabaceae</i>	this paper
<i>Hemileia vastatrix</i>	T	BRIP 61233	KT199399	DQ354565	KT199410	<i>Coffea robusta</i>	McTaggart et al. (2016); Aime (2006)
<i>Hyalopora aspidiotus</i>	T	PUR N4641	<b>MW049264</b>	NA	NA	<i>Gymnocarpium dryopteris</i>	this paper
<i>Kernkampella breyniae</i>	*	BRIP 56909	KJ862346	KJ862428	KJ862459	<i>Breynia cernua</i>	McTaggart et al. (2015)
<i>Kuehneola uredinis</i>	T	BPI 871104 (MCA2830)	DQ354551	DQ092919	NA	<i>Rubus argutus</i>	Aime (2006); Matheny & Hibbett unpublished

Table 1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	28S	18S	CO3	Host	Source
<i>Kwellingia bambusae</i>	T	PUR F18200	<b>MW147026</b>	NA	NA	<i>Bambusa</i> sp.	this paper
<i>Lipocystis acaciae-pennatulae</i> (as <i>Ravenelia acaciae-pennatulae</i> )		BPI 864189 (U115)	MG907213	MG907204	MG907264	<i>Vachellia pennatula</i>	Aime <i>et al.</i> (2018a)
<i>Lipocystis caesalpiniae</i>	T	BPI 863966	<b>MW049265</b>	NA	NA	<i>Mimosa ceratonia</i>	this paper
<i>Macruropyxis fraxini</i>	T	ZT Myc 56551	KP858145	KP858144	NA	<i>Fraxinus platypoda</i>	Beenken & Wood (2015)
<i>Maravalia limoniformis</i>	*	BRIP 59649	<b>MW049266</b>	NA	<b>MW036500</b>	<i>Austrosteenisia blackii</i>	this paper
<i>Masseëlla capparisi</i>	T	BRIP 56844	JX136798	NA	KT199413	<i>Flueggea virosa</i>	McTaggart <i>et al.</i> (2016)
<i>Melampsora euphorbiae</i>	T	BPI 863501 (U138)	DQ437504	DQ789986	<b>MW036501</b>	<i>Euphorbia macroclada</i>	Aime (2006); this paper
<i>Melampsora laricis-populina</i>		strain 98AG31	NW6768836	NW6768836	NW6768836	<i>Populus</i> sp.	Duplessis <i>et al.</i> unpublished
<i>Melampsorella caryophyllacearum</i>	T	PUR ex-MPPD-40507	MG907233	NA	NA	<i>Cerastium fontanum</i>	Aime <i>et al.</i> (2018a); this paper
<i>Melampsorium betulinum</i>	T	BPI 871107 (MCA2884):28S; n.d.: 18S	DQ354561	AY125391	NA	<i>Alnus</i> sp.	Aime (2006); Wingfield <i>et al.</i> (2004)
<i>Mikronegeria fagi</i>	T	PUR N16373	<b>MW049267</b>	<b>MW049298</b>	NA	<i>Nothofagus obliqua</i>	this paper
<i>Mikronegeria fuchsiae</i>	T	PDD 101517	KJ716350	KJ746826	NA	<i>Phyllocladus trichomanoides</i>	Padamsee & McKenzie (2014)
<i>Milesia polypodii</i> (as <i>Milesina polypodii</i> )	T	KRM0043190	MK302190	NA	NA	<i>Polypodium vulgare</i>	Bubner <i>et al.</i> (2019)
<i>Milesina kriegeriana</i>	T	KRM0048480	MK302207	NA	NA	<i>Dryopteris dilatata</i>	Bubner <i>et al.</i> (2019)
<i>Miyagia pseudosphaeria</i>	*	BPI 842230 (U63):28S; n.d.: 18S	DQ354517	AY125411	NA	<i>Sonchus oleraceus</i>	Aime (2006); Wingfield <i>et al.</i> (2004)
<i>Naohidemycetes vaccinii</i>	T	BPI 871754 (MCA2780)	DQ354563	DQ354562	NA	<i>Vaccinium ovatum</i>	Aime (2006)
<i>Neolivea tectonae</i>	T	PUR N15331 (MCA6480)	<b>MW049282</b>	<b>MW049307</b>	<b>MW036507</b>	<i>Tectona grandis</i>	this paper
<i>Neophysopella ampelopsidis</i> (as <i>Phakopsora ampelopsidis</i> )	T	IBA 8597	AB354738	NA	NA	<i>Ampelopsis brevipedunculata</i>	Chatasiri & Ono (2008)
<i>Neophysopella kraunhiae</i>		PUR N15073	<b>MW049242</b>	NA	NA	<i>Wisteria floribunda</i>	this paper
<i>Neophysopella meliosmae-myrianthae</i>		BRIP 58404	<b>MW049270</b>	NA	NA	<i>Vitrus</i> sp.	this paper
<i>Newinia heterophragmatis</i>	T	PUR N16505	<b>MW049271</b>	NA	NA	<i>Kigelia cf. africana</i>	this paper
<i>Nothoravenelia japonica</i>	T	HMJAU8598	MK296509	NA	NA	n.d.	Ji unpublished
<i>Nyssopora echinata</i>	T	KR-0012164 (U1022):28S; ESS244:18S	<b>MW049272</b>	U77061	NA	<i>Meum athamanticum</i>	this paper; Swann & Taylor (1995)
<i>Ochropsora ariae</i>	T	KR-0015027 (U1036)	<b>MW049273</b>	NA	NA	<i>Anemone nemorosa</i>	this paper
<i>Olivea capituliformis</i>	T	BPI 863670	<b>MW049274</b>	NA	NA	<i>Alchornea latifolia</i>	this paper
<i>Peridiopsis mori</i>		PUR N11676 (MCA4685)	<b>MW147025</b>	NA	<b>MW166323</b>	<i>Morus alba</i>	this paper
<i>Phakopsora crucis-filii</i>	T	ZT Myc 48990	KF528016	KF528041	KF528049	<i>Annona paludosa</i>	Beenken (2014)

Table 1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	28S	18S	CO3	Host	Source
<i>Phakopsora fici</i>		BRIP 59463	<b>MH047210</b>	<b>MW049299</b>	<b>MW036502</b>	<i>Ficus carica</i>	this paper
<i>Phakopsora pachyrhizi</i>	T	BRIP 56941	KP729475	<b>MW049300</b>	<b>MW036503</b>	<i>Neonotonia wightii</i>	Maier et al. (2016); this paper
<i>Phragmidium mucronatum</i>	T	BRIP 60097	<b>MW049275</b>	NA	NA	<i>Rosa rubiginosa</i>	this paper
<i>Phragmidium tormentillae</i> (≡ <i>Frommeëlla tormentillae</i> )	(T)	BPI 843392 (U3)	DQ354553	DQ354552	MG907265	<i>Potentilla canadensis</i>	Aime (2006); Aime et al. (2018a)
<i>Pileolaria brevipes</i>		PUR N16525 (MCA3477); 28S; CO3; BPI 877989 (MCA3223); 18S	MG907216	<b>MW049301</b>	MG907267	<i>Toxicodendron</i> sp.	Aime et al. (2018a); this paper
<i>Pileolaria shiriana</i>		BRIP 58344	KJ651957	NA	NA	<i>Rhus japonica</i>	Doungsa-ard et al. (2018)
<i>Pileolaria terebinthi</i>	T	PUR N11686 (U1282)	KY796222	NA	NA	<i>Pistacia terebinthus</i>	Ishaq et al. (2019)
<i>Porotenus biporus</i>	*	ZT Myc 3414	JF263494	JF263510	NA	<i>Memora flavida</i>	Beenken et al. (2012)
<i>Prospodium appendiculatum</i>	T	BPI 879956 (U753)	<b>MW049276</b>	NA	NA	<i>Tecoma stans</i>	this paper
<i>Prospodium lippiae</i>		BPI 843901 (U152)	DQ354555	DQ831024	NA	<i>Aloysia polystachya</i>	Aime (2006)
<i>Prospodium tuberculatum</i>		BRIP 57630	KJ396195	KJ396196	<b>MW036504</b>	<i>Lantana camara</i>	Pegg et al. (2014); this paper
<i>Puccinia graminis</i>	T	BRIP 60137	KM249852	<b>MW049302</b>	<b>MW036505</b>	<i>Glyceria maxima</i>	McTaggart et al. (2016); this paper
<i>Pucciniastrum epilobii</i>	T	PUR N11088 (MCA5308)	<b>MW049277</b>	NA	NA	<i>Epilobium angustifolium</i>	this paper
<i>Pucciniastrum minimum</i>		BRIP 57654	KC7633401	KT199391	KT199422	<i>Vaccinium corymbosum</i>	McTaggart et al. (2016)
<i>Puccinosira pallidula</i>	*	BPI 863541 (U282)	DQ354534	<b>MW049303</b>	NA	<i>Triumfetta semitriloba</i>	Aime (2006); this paper
<i>Puccinosira solani</i>		n.d.	EU851137	NA	NA	<i>Solanum aphyodendron</i>	Zuluaga et al. unpublished
<i>Puccorchidium polyalthiae</i>	T	ZT HeRB 251	JF263493	JF263509	JF263525	<i>Polyalthia longifolia</i>	Beenken & Wood (2015)
<i>Ravenelia</i> sp.	*	PUR F19717	<b>MW147024</b>	<b>MW166323</b>	<b>MW166322</b>	<i>Tephrosia</i> sp.	this paper
<i>Rogerpetersonia torreyae</i> (as <i>Caeoma torreyae</i> )	T	BPI 877825 (U1168); 28S; CO3; BPI 877824 (U808); 18S	MG907207	MG907197	MG907254	<i>Torreya californica</i>	Aime et al. (2018a)
<i>Rossmannomyces pyrolae</i> (as <i>Chrysomyxa pyrolae</i> )	T	390CHPPCGVF1	FJ666456	NA	NA	n.d.	Vialle et al. (2009)
<i>Skierka diploglottidis</i>	*	BRIP 59646	<b>MW049278</b>	<b>MW049304</b>	<b>MW036506</b>	<i>Dictyonera obtusa</i>	this paper
<i>Skierka robusta</i>	*	BPI 879954 (U747)	<b>MW049279</b>	<b>MW049305</b>	NA	<i>Rhoicissus rhomboidea</i>	this paper
<i>Sorataea arayatensis</i>		U416	<b>MW049280</b>	NA	NA	<i>Derris elliptica</i>	this paper
<i>Sphaerophragmium acaciae</i>	T	BRIP 56910	KJ862350	KJ862429	KJ862462	<i>Albizia</i> sp.	McTaggart et al. (2015)
<i>Sphenorchidium xylopiiae</i>	T	n.d.	KM217355	KM217372	NA	<i>Xylopiia aethiopica</i>	Beenken & Wood (2015)
<i>Sphenospora kevorkianii</i>		BPI 863558 (U10)	DQ354521	DQ354520	NA	<i>Stanhopea candida</i>	Aime (2006)

Table 1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	28S	18S	CO3	Host	Source
<i>Stereostromatum corticioides</i>	T	BPI 842314 (U27)	<b>MW049281</b>	<b>MW049306</b>	NA	<i>Bambusa</i> sp.	this paper
<i>Stomatiosora psychotriicola</i>	*	PREM 60886	NG059953	NA	NA	<i>Psychotria capensis</i>	Wood <i>et al.</i> (2014)
<i>Tegillum scitulum</i> (as <i>Olivea scitula</i> )	*	BPI 871108 (U668)	DQ354541	DQ354540	NA	<i>Vitex doniana</i>	Aime (2006)
<i>Thekopsora areolata</i>	T	n.d.	KJ546894	NA	NA	<i>Picea engelmannii</i>	Kaitera <i>et al.</i> unpublished
<i>Trachyspora intrusa</i>	T	BPI 84328 (MCA2384)	DQ354550	DQ354549	<b>MW036508</b>	<i>Alchemilla vulgaris</i>	Aime (2006); this paper
<i>Tranzschelia discolor</i>	*	BRIP 57662	KR995082	KR994969	KR995082	<i>Prunus persica</i>	Doungsa-ard <i>et al.</i> (2018)
<i>Tranzschelia mexicana</i>	*	KR-M-0040855	KP308391	NA	NA	<i>Prunus salicifolia</i>	Blomquist <i>et al.</i> (2015)
<i>Triphragmium ulmariae</i>	T	BPI 881364 (MCA2378); 28S; n.d.:18S	JF907676	AY125401	NA	<i>Filipendula ulmaria</i>	Yun <i>et al.</i> (2011); Wingfield <i>et al.</i> (2004)
<i>Uredinopsis filicina</i>	T	WM112	AF426237	NA	NA	<i>Phegopteris connectilis</i>	Maier <i>et al.</i> (2003)
<i>Uredo cryptostegiae</i> (as <i>Maravalia cryptostegiae</i> )	T	BRIP 56898	KT199401	KT199387	KT199412	<i>Cryptostegia grandiflora</i>	McTaggart <i>et al.</i> (2016)
<i>Uredo elephantopodis</i>		BRIP 58415	<b>MW049283</b>	NA	<b>MW036509</b>	<i>Elephantopus scaber</i>	this paper
<i>Uredo hiulca</i>		BRIP 53244	<b>MW049284</b>	NA	<b>MW036510</b>	<i>Dioscorea transversa</i>	this paper
<i>Uredo trichosanthis</i>		PUR N3445	<b>MW049285</b>	<b>MW049309</b>	NA	<i>Trichosanthes bracteata</i>	this paper
<i>Uredopeltis atrides</i>	*	PUR N13866 (U454)	<b>MW049286</b>	NA	NA	<i>Grewia flavescens</i>	this paper
<i>Uredopeltis chevalieri</i>	*	BRIP 56924	<b>MW049287</b>	NA	NA	<i>Grewia retusifolia</i>	this paper
<i>Uromyces appendiculatus</i>	T	BRIP 60020	KM249870	DQ354510	KX999933	<i>Phaseolus vulgaris</i>	Aime (2006); McTaggart <i>et al.</i> (2014)
<i>Uromykladium simplex</i>	T	BRIP 59214	KJ632990	KJ633029	KJ639078	<i>Acacia pycnantha</i>	Doungsa-ard <i>et al.</i> (2014)
<i>Uropyxis daleae</i>	*	BPI 910337	KY798364	NA	NA	<i>Dalea pringlei</i>	Demers & Castlebury unpublished
<i>Uropyxis diphyssae</i>	*	BPI 864148	<b>MW049288</b>	NA	NA	<i>Diphysa americana</i>	this paper
<i>Xenodochus carbonarius</i>	T	PUR N15566 (U1534)	<b>MW049289</b>	NA	NA	<i>Sanguisorba officinalis</i>	this paper
<i>Xenosteles litseae</i>	*	BRIP 53335	<b>MW049290</b>	<b>MW049310</b>	NA	<i>Neolitsea dealbata</i>	this paper
<i>Ypsilospora tucumanensis</i>	*	BPI 863688	<b>MW049291</b>	NA	NA	<i>Inga</i> sp.	this paper
<i>Zaghouania notelaeae</i> (as <i>Cystopsora notelaeae</i> )	*	BRIP 58325	KT199396	KT199384	KT199407	<i>Notelaea microcarpa</i>	McTaggart <i>et al.</i> (2016)

<sup>a</sup>Type Status: T = type species for the genus; \* = proxy for generic type (see methods for explanation).

<sup>b</sup>Numbers in parentheses are collection numbers, preceded by herbarium accession numbers. When sequences from more than one collection are used, data are separated by a /.

n.d. = no data.

NA = not applicable.

bold = new sequences generated for this paper.

c-oxidase subunit 3 (CO3) of the mitochondrial DNA was amplified with CO3\_F1/CO3\_R1 (Vialle *et al.* 2009) following the protocols of Vialle *et al.* (2009). PCR products were cleaned and sequenced with the amplification primers by Macrogen (Seoul, Korea) or Beckman Coulter Sequencing (Danvers, Massachusetts, USA). Sequences were edited in Sequencher v. 4.5–5.4 (Gene Codes Corp., Ann Arbor, Michigan, USA) and verified by BLASTn against the NCBI database (Altschul *et al.* 1990). Sequence accession numbers are provided in Tables 1 and S1.

## Phylogenetic analyses

The 28S, 18S and CO3 sequences were aligned in four datasets, (i) *Pucciniales* overview, (ii) *Melampsorineae*, (iii) *Raveneliineae*, and (iv) *Urediniineae* with the GUIDANCE2 webserver (Sela *et al.*, 2015; available at <http://guidance.tau.ac.il/ver2/credits.php>) (alignments are available from TreeBASE, study TB2:S27114). The aligned loci were concatenated and run as partitioned datasets with maximum likelihood (ML). We searched for the most likely tree in IQTree v. 1.7 beta (Nguyen *et al.* 2015) with a GTR gamma FreeRate heterogeneity model of evolution and a different rate for each partition (command -spp -m GTR+R), 10 000 ultrafast bootstraps (Hoang *et al.* 2018), an approximate likelihood ratio test with 10 000 replicates (Guindon *et al.* 2010) and genealogical concordance factors calculated from each locus (Minh *et al.* 2018).

We used the concatenated three-locus alignment of the familial-overview dataset to estimate the divergence dates of genera with BEAST v. 2.5 (Bouckaert *et al.* 2019). We calibrated the most recent common ancestor of the *Pucciniales* at 175 M yr and the *Melampsorineae* at 91 M yr based on Aime *et al.* (2018a). The dating analyses were constrained to the topology of the ML tree, and run for 150 M generations, with a BEAST model test for each partition and a relaxed log normal clock. Convergence of all priors was visualised in Tracer v. 1.7 (Rambaut *et al.* 2018) and 135 001 trees were summarised with TreeAnnotator, part of the BEAST v. 2.5 package.

We attempted to provide better resolution of genera and families within *Raveneliineae* by multiple means including removal of incongruent (rogue) taxa, constructing alignments with stricter and weaker gap opening penalties, pruning taxa with missing sequence data, removal of 18S and CO3 loci, and rooting with different outgroups from the *Melampsorineae* and *Urediniineae*. The 28S data of *Raveneliineae* were analysed with SplitsTree4 (Huson & Bryant 2005) to visualize the evolution as a network in order to determine if groups were supported when not constrained by dichotomous evolution as imposed by ML analyses.

## RESULTS

### Phylogenetic analyses

The ML tree based on three concatenated loci (Fig. 1) was mostly congruent with prior studies of more limited taxon and locus sampling (Aime 2006, Beenken & Wood 2015, McTaggart *et al.*

2016, Aime *et al.* 2017, 2018a, Beenken 2017, Souza *et al.* 2018). Sampled trees constrained to the ML topology in the dating analyses converged after 150 M generations, supported by all effective sample size values over 200. We recovered support for placement of previously unsupported or unplaceable taxa such as *Tranzschelia*. Newly sequenced taxa resolved include the rust fungi on *Agathis*, genera such as *Elateraecium*, *Masseëlla*, and *Skierka*, and most of the endocyclic *Puccinosiraceae*. Despite numerous attempts with different alignments and taxon selection, some families/genera could not be confidently resolved, namely: *Pucciniastrum* and *Pucciniastraceae*; *Raveneliaceae*; and *Allodus*, *Neopuccinia*, and *Nyssopsora* within *Urediniineae*. SplitsTree analysis of *Raveneliineae* recovered a star-shaped pattern of reticulate edges indicative of multiple competing hypotheses of evolution for this lineage (Fig. S2).

### Taxonomy

Families and sub-orders treated here show strong support at their most recent common ancestor, with the exception of *Pucciniastraceae* and *Raveneliaceae* (Figs 1–3). We propose four new suborders (*Araucariomycetinae*, *Raveneliineae*, *Rogerpetersoniineae*, and *Skierkineae*), seven new families (*Araucariomycetaceae*, *Crossosporaceae*, *Milesinaceae*, *Ochropsoraceae*, *Rogerpetersoniaceae*, *Skierkaceae*, and *Tranzscheliaceae*) and four new genera (*Araucariomyces*, *Neoolivea*, *Rogerpetersonia*, and *Rossmatomyces*); 21 new combinations and one new name are made for species. Suborders and families are arranged from earliest diverging to more recently derived (Fig. 1). We use the terms gametothallus and sporothallus as applied by Berndt (2018) and use the notation 0-I [for spermogonial (0) and aecial (I) stages] to denote the gametothallus, and II-III [for uredinial (II) and telial (III) stages] to denote the sporothallus. We follow the ontogenic system for sorus terminology, which emphasizes function in the life cycle and the nuclear cycle over morphology, as refined by Hiratsuka (1973). Morphological terms for spermogonia follow Hiratsuka & Cummins (1963); terms for aecial and uredinial sori follow the descriptions for asexual genera in Cummins & Hiratsuka (2003) but are indicated in lowercase, non-italics, to delineate use as descriptive terms from generic names.

***Rogerpetersoniineae*** Aime & McTaggart, *subord. nov.* MycoBank MB836604.

*Type family:* *Rogerpetersoniaceae* Aime & McTaggart, this paper.


*Diagnosis:* Differs from all other *Pucciniales* in that gametothalli are formed on *Taxaceae*.

*Description:* With the characteristics of *Rogerpetersoniaceae*.

*Included family:* *Rogerpetersoniaceae*.

**Fig. 1.** *Pucciniales*. Phylogram obtained from BEAST constrained to a ML topology from three concatenated loci (28S, 18S, and CO3). The tree is rooted with *Eocronartium muscicola*. Families are indicated by coloured blocks; dashed lines indicate uncertainty at the referenced nodes. Genera represented by types are indicated in bold; genera represented by type proxies (as explained in methods) are indicated by \*. Support for nodes is provided from an approximate likelihood ratio test ( $\geq 0.90$ ), ultrafast bootstraps ( $\geq 95\%$ ) and genealogical concordance factors for the three loci at each node as aLRT/UFBoot/gCF.





**Rogerpeteroniaceae** Aime & McTaggart, *fam. nov.* MycoBank MB836605.

*Type genus:* *Rogerpeteronia* Aime & McTaggart, this paper.

*Diagnosis:* Differs from all other *Pucciniales* in that gametothalli are formed on *Taxaceae*.

*Description:* With the characteristics of *Rogerpeteronia*.

*Included genus:* *Rogerpeteronia*.

*Host family:* *Taxaceae* (0-I); II-III unknown.

**Rogerpeteronia** Aime & McTaggart, *gen. nov.* MycoBank MB836606.

*Type species:* *Rogerpeteronia torreyae* (Bonar) Aime & McTaggart, this paper.

*Etymology:* In honour of Roger Peterson, botanist, ecologist, mycologist and plant pathologist, who pioneered studies on Southern Hemisphere conifer rusts.

*Diagnosis:* Differs from all other rust fungi in forming gametothalli on *Taxaceae* (*Torrea*).

*Description:* With the characteristics of *Rogerpeteronia torreyae*.

**Rogerpeteronia torreyae** (Bonar) Aime & McTaggart, *comb. nov.* MycoBank MB836608.

*Basionym:* *Caeoma torreyae* Bonar, *Mycologia* **43**: 62. 1951.

*Description:* *Rogerpeteronia torreyae* is described and illustrated as *C. torreyae* in Peterson (1974). Spermogonia are deep-seated, periphysate, otherwise similar to Group III (type 12). Aecia petersonia-like, *i.e.*, without peridium or intercalary cells. Sporothallus unknown.

*Notes:* *Caeoma*, as typified by *C. berberidis*, is a synonym of *Puccinia* (Aime *et al.* 2018b), necessitating a new name for the only known rust fungus that infects *Torrea*. Peterson (1974) hypothesized that *R. torreyae* belonged to an undescribed early diverging lineage of *Pucciniales*. Subsequent analyses have shown that *R. torreyae* is the earliest diverging extant rust sequenced to date and holds an isolated position within *Pucciniales* (Aime 2006, Aime *et al.* 2018a) (Fig. 1). No alternate host is known for this rust and it is likely that *R. torreyae* has adapted to cause systemic infections in the gametothallus host in order to compensate for loss of a sporothallus.

**Mikronegeriineae** Aime, *Mycoscience* **47**: 120. 2006.

*Description:* With the characteristics of the family.

*Included family:* *Zaghouaniaceae*.

**Zaghouaniaceae** P. Syd. & Syd., *Monogr. Uredin.* (Lipsiae) **3**(3): 586. 1915. *emend.* Aime & McTaggart

*Synonyms:* *Hemileieae* Dietel, *Uredinales* in Engler and Prantl., *Naturl.*: 51. 1928.

*Mikronegeriaceae* Cummins & Y. Hirats. (as '*Mikronegeriaceae*'), *Illustr. Gen. Rust Fungi*, rev. Edn (St. Paul): 13. 1983.

*Type genus:* *Zaghouania* Pat., *Bull. Soc. mycol. Fr.* **17**: 187. 1901.

*Description:* Spermogonia most often Group III (type 12) (deep seated and non-periphysate), but periphyses noted for some; aecia most commonly of the petersonia-type, *i.e.*, without peridium or intercalary cells, however in *Elateraecium* accompanied with specialized elaters; uredinia most often uredo-type, in *Elateraecium* with a weakly developed peridium in young sori; teliospores without dormancy, germinating externally by apical growth, or internally (*Achrotelium*). *Blastospora* and *Mikronegeria* are heteroecious and macrocyclic, *Elateraecium* and *Zaghouania* are autoecious and macro- or demi-cyclic; complete life cycles unknown for *Achrotelium*, *Botryorhiza* and *Hemileia*.

*Included genera:* *Achrotelium*, *Blastospora*, *Botryorhiza*, *Elateraecium* (= *Hiratsukamyces*), *Hemileia*, *Mikronegeria*, *Zaghouania* (= *Cystopsora*); likely includes *Desmosorus*.

*Host families:* *Araucariaceae*, *Betulaceae* (0-I heteroecious species); *Apocynaceae*, *Araliaceae*, *Capparidaceae*, *Celastraceae*, *Cupressaceae*, *Dioscoreaceae*, *Euphorbiaceae*, *Fagaceae*, *Oleaceae*, *Orchidaceae*, *Rubiaceae*, *Smilacaceae*, *Verbenaceae* (II-III and autoecious species).

*Notes:* The family *Mikronegeriaceae* accommodated the heteroecious rust genera *Mikronegeria*, *Blastospora*, and *Chrysocelis*, which have thin-walled basidia that germinate externally without dormancy (Cummins & Hiratsuka 2003). *Hemileia* and some *Maravalia* species formerly placed in *Chaconiaceae* that share the feature of substomatal sori without paraphyses or peridium, also belong here (Aime 2006). Two additional genera, *Achrotelium* and *Zaghouania* (as *Cystopsora*), were included by McTaggart *et al.* (2016). *Zaghouaniaceae*, long considered a synonym for *Pucciniaceae* (*e.g.*, Kirk *et al.* 2008), has priority over *Mikronegeriaceae* and the family is now referred to by the earlier name. The current study adds *Elateraecium* (syn. *Hiratsukamyces*; Aime *et al.* 2018b), whereas *Chrysocelis* is resolved within the *Pucciniaceae* (Fig. 4). The formation of basidia is primarily external by apical growth and spermogonia are primarily deep-seated Group III (type 12), or if periphysate, similar to Group V (type 4).

*Uredo cryptostegiae* (syn. *Maravalia cryptostegiae*; *Scopella cryptostegiae*), which has been used as a biocontrol agent for rubber-vine (*Cryptostegia grandiflora*) is placed in *Zaghouaniaceae* (Fig. 1). Cummins (1950) transferred *M. cryptostegiae* to *Scopella*, while hypothesizing that the rust might belong to *Hemileia*. Most later workers considered *Scopella* and *Maravalia* congeneric and *Scopella* fell out of use. The type of *Maravalia*, *M. pallida* occurs on *Fabaceae* and is now placed in *Raveneliineae*. The type of *Scopella*, *S. echinulata*, is a subepidermal rust on *Sapotaceae* (Mains 1939a). *Uredo cryptostegiae*, which is not congeneric with *Maravalia* (as represented by *M. limoniformis*, Fig. 1), is most appropriately retained in *Uredo* until type data from *S. echinulata* is obtained.

**Zaghouania notelaeae** (Syd.) Aime & McTaggart, *comb. nov.* MycoBank MB836655.

*Basionym:* *Cystopsora notelaeae* Syd., *Annls mycol.* **35**: 351. 1937.

*Notes:* *Zaghouania* contains two other species of rust fungi on *Oleaceae* with pale-walled teliospores that germinate without dormancy (Cummins & Hiratsuka 2003). There is little to differentiate *Cystopsora* and *Zaghouania* (Thirumalachar 1945, Cummins & Hiratsuka 2003) and we treat *Cystopsora* as a synonym of *Zaghouania*.

***Araucariomycetinae*** Aime & McTaggart, **subord. nov.** MycoBank MB836623.

*Type family:* *Araucariomycetaceae* Aime & McTaggart, this paper.

*Diagnosis:* Differs from all other *Pucciniales* in forming gametothalli on *Agathis*.

*Description:* With the characteristics of *Araucariomycetaceae*.

*Included family:* *Araucariomycetaceae*.

***Araucariomycetaceae*** Aime & McTaggart, **fam. nov.** MycoBank MB836624.

*Type genus:* *Araucariomyces* Aime & McTaggart, this paper.

*Diagnosis:* Differs from all other *Pucciniales* in forming gametothalli on *Agathis*.

*Description:* With the characteristics of *Araucariomyces*.

*Included genus:* *Araucariomyces*.

*Host family:* *Araucariaceae* (0-I); II-III unknown.

***Araucariomyces*** Aime & McTaggart, **gen. nov.** MycoBank MB836625.

*Type species:* *Araucariomyces fragiformis* (Ces.) McTaggart, R.G. Shivas & Aime, this paper.

*Entomology:* From the host family, *Araucariaceae*.

*Diagnosis:* Differs from all other rust genera in forming the gametothallus on species of *Agathis* (*Araucariaceae*).

*Description:* These species are described and illustrated in Peterson (1966). Spermogonia amphigenous, intra-epidermal becoming sub-epidermal as they break through host walls, convex hymenium; similar to Group 1 (type 1) but with scant periphyses not visible in all mounts, similar to *Rogerpetersonia*. Aecia peridiate, aecidium-type, deep-set within swollen host tissues. Sporothallus unknown. On leaves of *Agathis* (*Araucariaceae*). Two known species.

*Notes:* Two rust fungi with cupulate aecia on *Agathis* spp., formerly placed in the form-genus *Aecidium*, belong here. Our analyses consistently place these in a lineage separate from all other sequenced *Pucciniales* (Fig. 1). Despite over a decade of sampling rust fungi from Australia and Southeast Asia on hosts co-distributed with *Agathis* species, we have been unable to locate a telial state for these rusts. Peterson (1968) ruled out the possibility that *Araucariomyces* represents an endocyclic form, because aeciospores of *Ar. balansae* germinate to produce germ tubes rather than basidia. As is conjectured with

*Rogerpetersonia*, the life cycle may not produce a sporothallus, and instead has adapted to systemically infect their hosts possibly including a cryptic sexual or parasexual cycle.

***Araucariomyces balansae*** (Cornu) McTaggart, R.G. Shivas & Aime, **comb. nov.** MycoBank MB836626.

*Basionym:* *Aecidium balansae* Cornu., *Bull. Soc. mycol. Fr.* **3**: 173. 1887.

*Synonym:* *Peridermium balansae* (Cornu) Sacc., *Syll. Fung.* **9**: 326. 1891.

***Araucariomyces fragiformis*** (Ces.) McTaggart, R.G. Shivas & Aime, **comb. nov.** MycoBank MB836627.

*Basionym:* *Aecidium fragiforme* Ces., *Atti Accad. Sci. fis. mat. Napoli* **8**: 26. 1879.

***Skierkineae*** Aime & McTaggart, **subord. nov.** MycoBank MB836628.

*Type family:* *Skierkaceae* Aime & McTaggart, this paper.

*Diagnosis:* Differs from all other rust fungi in that sporothalli sori are deep-seated and subepidermal with mature uredinio- and teliospores single-celled and non-catenulate, these forced through a narrow sorus opening by the production of new spores from sporogenous cells from which they are detached before extrusion.

*Description:* With the characters of *Skierkaceae*.

*Included family:* *Skierkaceae*.

***Skierkaceae*** (Arthur) Aime & McTaggart, **fam. & stat. nov.** MycoBank MB836629.

*Basionym:* *Skierkatae* Arthur, *North American Flora* **7**(10): 704. 1926.

*Type genus:* *Skierka* Racib., *Parasit. Alg. Pilze Javas* (Jakarta) **2**: 30. 1900.

*Diagnosis:* Differs from all other rust fungi in that sporothalli sori are deep-seated and subepidermal with mature uredinio- and teliospores single-celled and non-catenulate, these forced through a narrow sorus opening by the production of new spores from sporogenous cells from which they are detached before extrusion.

*Description:* With the characteristics of *Skierka* as described and illustrated in Mains (1939b). Spermogonia deep-seated with convex hymenium, subepidermal, periphysate; aecia and uredinia uredo-type; teliospores strongly adherent, extruded in hair-like columns, germination external, without dormancy. Autoecious and macrocyclic.

*Included genus:* *Skierka*.

*Host families:* *Burseraceae*, *Euphorbiaceae*, *Sapindaceae*.

*Notes:* *Skierka* species are tropical and autoecious (Mains 1939c, Cummins & Hiratsuka 2003). All sori are subepidermal and deep-seated; non-catenulate teliospores are extruded in hair-like columns. *Skierka* has long held an isolated placement

within *Pucciniales*. Arthur (1907–1931) and Dietel (1928) placed *Skierka* in a separate subfamily or tribe, respectively, in the *Pucciniaceae*; Cummins & Hiratsuka (2003) treat it as *incertae sedis* within the rusts. Mains (1939c) hypothesised that *Skierka* represented an intermediate taxon between the *Melampsoraceae* and *Pucciniaceae* (equivalent to the subordinal ranks *Melampsorineae* and *Raveneliineae/Urediniineae*, under the present classification), a position largely congruent with our placement (Fig. 1).

**Melampsorineae** Aime, *Mycoscience* **47**: 120. 2006.

*Type family*: *Melampsoraceae* Dietel, in Engler & Prantl, *Nat. Pflanzenfam.* **1**(1): 38. 1897.

*Description*: Mostly macrocyclic and heteroecious, forming the gametothallus on species of *Pinaceae*. Teliospores germinate after a period of dormancy.

*Included families*: *Coleosporiaceae*, *Melampsoraceae*, *Milesinaceae*, *Pucciniastraceae*.

**Milesinaceae** Aime & McTaggart, *fam. nov.* MycoBank MB836630.

*Type genus*: *Milesina* Magnus, *Ber. Deutsch. Bot. Ges.* **27**: 325. 1909.

*Diagnosis*: Similar to other *Melampsorineae*, differing in either production of colourless urediniospores in species that infect ferns, or in production of milesia-type aecia in species that infect *Ericaceae*.

*Description*: With typically colourless sori, although urediniospores of *Naohidemycetes* are orange, otherwise similar to *Pucciniastraceae*. Spermogonia Group I (mostly type 1, also type 2 and 3); aecia peridermium-type, milesia-type in *Naohidemycetes*; uredinia milesia-type. Teliospores with dormant germination, 1- to many-celled, barely differentiated, sometimes laterally adherent, typically formed within host epidermal cells. Most species macrocyclic and heteroecious with sporothalli on ferns (excepting *Naohidemycetes* on *Ericaceae*), and gametothalli on *Pinaceae*.

*Included genera*: *Milesia*, *Milesina*, *Naohidemycetes*, *Uredinopsis*.

*Host families*: *Pinaceae* (*Abies*, *Tsuga*) (0-I); *Ericaceae* and some ferns in *Polypodiales* and *Lygodium* (II-III).

*Notes*: Early workers considered rust fungi on early diverging plant hosts (*i.e.*, ferns) to be the “ancestral” *Pucciniales*. Several molecular phylogenetic studies have shown this not to be the case (*e.g.*, Sjamsuridzal *et al.* 1999). However, the fern rusts are among the earliest diverging members of *Melampsorineae* (Fig. 2), the second major radiation of the rust fungi, and belong to the two earliest families in this suborder (*Milesinaceae* and *Pucciniastraceae*). Most of the species in *Milesinaceae* form sporothalli on fern species, except for *Naohidemycetes*, which alternates between *Tsuga* and *Vaccinium*.

Aime *et al.* (2018b) recommended protecting the name *Milesina* Magnus over *Milesia* F.B. White. However, our data show that the type of *Milesina*, *M. kriegiana* (Magnus)

Magnus, is not congeneric with the type of *Milesia*, *M. polypodii* F.B. White (Fig. 2), thus we recommend retaining both genera at this time. Should future work demonstrate that *Uredinopsis* is polyphyletic, then disposition of these taxa will need revision.

**Coleosporiaceae** Dietel, In: Engler & Prantl, *Nat. Pflanzenfam.*, Teil. I (Leipzig) **1**: 548. 1900. **emend.** Aime & McTaggart  
*Synonym*: *Cronartiaceae* Dietel, in Engler and Prantl, *Nat. Pflanzenfam.* **1**(1) (Suppl.): 548. 1900.

*Type genus*: *Coleosporium* Lév., *Ann. Sci. Nat. Bot.* III, Ser. **8**: 373. 1847.

*Description*: Spermogonia Group I (type 2 or 3) (but Group II, type 9 in *Cronartium*); aecia of peridermium-type; uredinia either of caeoma-type or milesia-type. Teliospores packed to loosely adherent, often extruded in columns and/or gelatinous; not dormant, with external germination. Most are heteroecious and macrocyclic, with some derived microcyclic or endocyclic species.

*Included genera*: *Chrysomyxa*, *Coleosporium*, *Cronartium*, *Diaphanopellis*, *Rossmatomyces*, *Thekopsora* (= *Quasi-pucciniastrum*).

*Host families*: *Pinaceae* (primarily *Pinus*) (0-I); various, including *Apocynaceae*, *Asteraceae*, *Campanulaceae*, *Convolvulaceae*, *Ericaceae*, *Lamiaceae*, *Ranunculaceae*, *Rosaceae*, *Rutaceae*, *Violaceae* (II-III).

*Notes*: *Coleosporiaceae* was shown to include *Cronartiaceae* (Aime 2006) as well as *Thekopsora s.s.* (Aime *et al.* 2018a). Aecia are peridermium-type in contrast to most *Milesinaceae*. Telial states show variable morphology but tend to form the sporothallus on herbaceous rather than woody plants (*cf.* *Pucciniastraceae*) or ferns. Dietel (1900) established both *Coleosporiaceae* and *Cronartiaceae* in the same publication. We follow Sydow & Sydow (1915) in applying *Coleosporiaceae* over *Cronartiaceae*, which is discussed in Aime (2006). *Endocronartium* is a later synonym of *Cronartium* (Aime *et al.* 2018b).

**Rossmatomyces** Aime & McTaggart, *gen. nov.* MycoBank MB836632.

*Type species*: *Rossmatomyces pyrolae* (Rostr.) Aime & McTaggart, this paper.

*Etymology*: In honour of Amy Rossman, biologist, mycologist, plant pathologist, and mentor.

*Diagnosis*: Similar to *Chrysomyxa* but differs in forming a systemic sporothallus; differs from all other rust fungi in forming sporothalli on *Moneses* and *Orthilia* (*Ericaceae*).

*Description*: See Saville (1950) and Feau *et al.* (2011). Gametothalli systemic in cones of *Picea* species; sporothalli systemic in *Moneses*, *Orthilia*, and *Pyrola* species.

**Rossmatomyces monesis** (Ziller) Aime & McTaggart, *comb. nov.* MycoBank MB836633.

*Basionym*: *Chrysomyxa monesis* Ziller, *Canad. J. Bot.* **32**: 435. 1954.




Fig. 2. *Melamporsorineae*. ML topology generated from three concatenated loci (28S, 18S, and CO3) with expanded taxon sampling. The tree is rooted with *Araucariomyces fragiformis* and *Zaghouania notelaeae*. Families are indicated by coloured blocks; *Pucciniastraceae* is recovered as a grade in these analyses. Genera represented by bold; genera represented by type proxies (as explained in methods) are indicated by \*. Support for nodes is provided from an approximate likelihood ratio test ( $\geq 0.90$ ) and ultrafast bootstrap ( $\geq 95\%$ ) as aLRT/UFBoot.

**Rossmatomyces pyrolae** (Rostr.) Aime & McTaggart, **comb. nov.** MycoBank MB836634.

*Basionym:* *Chrysomyxa pyrolae* Rostr., *Botan. Zbl.* **5**: 127. 1881.

**Rossmatomyces ramischiae** (Lagerh.) Aime & McTaggart, **comb. nov.** MycoBank MB836635.

*Basionym:* *Chrysomyxa ramischiae* Lagerh., *Svensk bot. Tidskr.* **3**: 26. 1909.

*Notes:* *Chrysomyxa* is typified by *C. abietis*, a microcyclic species for which there are no sequence data. In our analyses (Fig. 2) most species of *Chrysomyxa* were monophyletic, excluding *C. weirii* now placed in *Ceropsora*, and species that infect wintergreens, now placed in *Rossmatomyces*. The species of *Rossmatomyces* are the only known rust species that form sporothalli on species of *Moneses* and *Orthilia*, and the only *Coleosporiaceae* that form sporothalli on species of *Pyrola*. The gametothalli are produced on *Picea* and are systemic within the cones, in contrast to gametothalli of *Chrysomyxa* species, which infect needles.

**Thekopsora americana** (Farl.) Aime & McTaggart, **comb. et stat. nov.** MycoBank MB836637.

*Basionym:* *Pucciniastrum arcticum* var. *americanum* Farl., *Rhodora* **10**: 16. 1908.

*Synonym:* *Pucciniastrum americanum* (Farl.) Arthur, *Bull. Torrey bot. Club* **47**: 468. 1920.

**Thekopsora potentillae** (Korn.) Aime & McTaggart, **comb. nov.** MycoBank MB836636.

*Basionym:* *Pucciniastrum potentillae* Korn., in Jaczewski *et al.*, *Fungi Rossiae Exsicc. fasc.* **7**: 327. 1900 [1899].

*Notes:* Delimitation between *Thekopsora* and *Pucciniastrum* has never been satisfactory (e.g., Hiratsuka 1958, Sato *et al.* 1993). While prior works mostly consider these confamilial or even congeneric, *Thekopsora s.s.*, as typified by *T. areolata*, belongs to *Coleosporiaceae* (Aime *et al.* 2018a; Fig. 2). New combinations are proposed for ex-*Pucciniastrum* species. Other former *Thekopsora* species, such as *P. minima* and *P. rubiae* are placed in *Pucciniastraceae* (Fig. 2).

**Thekopsora pseudoagrimoniae** Aime & McTaggart, **nom. nov.** MycoBank MB836638.

*Basionym:* *Quasipucciniastrum agrimoniae* X.H. Qi *et al.*, *Mycology* **10**(3): 145. 2019.

*Description:* See Qi *et al.* (2019).

*Notes:* The recently described monotypic *Quasipucciniastrum* based on *Q. agrimoniae* is congeneric with *Thekopsora* (Fig. 2). In addition to the phylogenetic data, *Quasipucciniastrum* shares key morphological features, ecology, and hosts with *Thekopsora*. This paper highlights the importance of including type species and adequate sampling in phylogenetic studies of known polyphyletic genera. The name *Thekopsora agrimoniae* Dietel is already in use, thus a new name is proposed for this taxon. However, there is little to differentiate *T. pseudoagrimoniae* from *T. agrimoniae* and the two may be conspecific.

**Pucciniastraceae** Gäum. ex Leppik, *Ann. bot. fenn.* **9**: 139. 1972. **emend.** Aime & McTaggart

*Type genus:* *Pucciniastrum* G.H. Otth, *Mitt. Naturforsch. Ges. Bern* **1861**: 71. 1861.

*Description:* Similar to *Milesiaceae*, but most species with cytoplasmic pigmentation, at least within urediniospores. Spermogonia Group I (type 2 or 3). Aecia peridermium-type; uredinia milesia-type. Telia undergo dormancy with external germination; either formed within epidermal cells, or as a subepidermal crust, which is gelatinous in *Coleopuccinia*. Most species heteroecious, macrocyclic; *Calyptospora* is demicyclic, *Coleopuccinia* is microcyclic, producing only teliospores.

*Included genera:* *Calyptospora*, *Coleopuccinia*, *Hyalopsora*, *Melampsorella*, *Melampsidium*, *Pucciniastrum*.

*Host families:* *Pinaceae* (*Abies*, *Larix*, *Picea*, *Tsuga*) (0-I); *Aceraceae*, *Betulaceae*, *Caryophyllaceae*, *Ericaceae*, *Fagaceae*, *Onagraceae*, *Rosaceae*, *Rubiaceae* and some ferns in the *Polypodiales* (II-III).

*Notes:* Most species of *Pucciniastraceae* produce spores with pigmented cytoplasm and telia that may be subepidermal, in contrast to *Milesiaceae*. *Hyalopsora* is the only genus in *Pucciniastraceae* that infects ferns. *Coleopuccinia* is known only from teliospores (Cao *et al.* 2018). *Pucciniastraceae s.l.* has been difficult to resolve and appears polyphyletic with varying degrees of support in earlier studies (e.g., Maier *et al.* 2003, Aime 2006, Aime *et al.* 2016a, Ji *et al.* 2019). In this work, we find weak support for *Pucciniastraceae* in some analyses (data not shown) but not all (e.g., Fig. 1). In nearly all analyses *Pucciniastraceae* is resolved into two groups: (i) *Calyptospora*, *Melampsorella*, and *Pucciniastrum*; and (ii) *Coleopuccinia*, *Hyalopsora*, and *Melampsidium*. These often form a grade (Fig. 2) and may or may not represent separate family-rank lineages. Pending additional analyses, we broadly define *Pucciniastraceae* to include both groups. *Pucciniastrum* is also difficult to resolve with confidence, and is most likely paraphyletic, even after removing the ex-*Pucciniastrum* elements that were reassigned to *Thekopsora* (Fig. 2). We retain *Coleopuccinia*, *Calyptospora*, and *Melampsorella* at this time, although future work may show that the latter two are synonyms for *Pucciniastrum*.

**Melampsoraceae** Dietel, in Engler & Prantl, *Nat. Pflanzenfam.*, Teil. I (Leipzig) **1**: 38. 1897.

*Type genus:* *Melampsora* Castagne, *Obs. Plantes Acotylédonées Fam. Urédinié* **2**: 18. 1843.

*Description:* Spermogonia Group I (type 2 or 3). Aecia mostly caeoma-type; uredinia uredo-type. Teliospores subepidermal, laterally adherent in crusts, 1-celled, often with a sterile basal cell; germination external or semi-external (*Ceropsora*). Most species heteroecious, macrocyclic; *Ceropsora* species are microcyclic.

*Included genera:* *Melampsora*; likely includes *Ceropsora*.

*Host families:* Primarily *Pinaceae* (0-I); primarily *Salicaceae*, also *Apocynaceae*, *Asteraceae*, *Euphorbiaceae*, *Flacourtiaceae*, *Hypericaceae*, *Linaceae*, *Passifloraceae*, *Saxifragaceae*, *Scrophulariaceae*, *Thymelaeaceae* (II-III)

***Ceropsora weirii*** (H.S. Jacks.) Aime & McTaggart, **comb. nov.** MycoBank MB836631.

**Basionym:** *Chrysomyxa weirii* H.S. Jacks., *Phytopathology* **7**: 353. 1917.

**Notes:** Most of the ca. 30 species of *Chrysomyxa* are heteroecious with gametothalli on *Pinaceae* and are allied within *Coleosporiaceae* (Fig. 2). *Chrysomyxa weirii*, an autoecious microcyclic pathogen of *Picea* species, is unique among described *Chrysomyxa* in forming laterally adherent teliospores that act as diaspores, are adapted for dispersal in water, and germinate to produce 2-celled basidia (Crane 2000, Crane *et al.* 2000). Crane *et al.* (2000) conjectured that *Ch. weirii* is not a true *Chrysomyxa*, which is supported with molecular data (Feau *et al.* 2011, Aime *et al.* 2018a, Fig. 2). The type and only other species of *Ceropsora*, *C. picea*, is a teliospore-only species infecting *Picea* in India (Bakshi & Singh 1960). While we have been unable to sequence a representative of the type species, *C. weirii* and *C. picea* are both microcyclic producing telia on *Picea* species. In both species, the telia contain some thin-walled sterile cells on the sides that have been interpreted as remnants of a peridermium. And in both, teliospores are subtended by sterile basal cells forming initially adherent crusts that separate at dispersal; germination is semi-external (Bakshi & Singh 1960, Crane *et al.* 2000).

**Raveneliineae** Aime & McTaggart **subord. nov.** MycoBank MB836639.

**Type family:** *Raveneliaceae* Leppik, *Ann. Bot. Fenn.* **9**(3): 139. 1972.

**Diagnosis:** Similar to *Uredinineae* differing in that the majority of species form Group VI spermogonia whereas the majority of *Uredinineae* form Group V spermogonia.

**Description:** With the characteristics of the included families. Most species form Group VI spermogonia; many species form elaborate, multi-celled teliospores.

**Included families:** *Ochropsoraceae*, *Phakopsoraceae*, *Raveneliaceae*, *Tranzscheliaceae*.

**Notes:** The *Raveneliineae* is the most challenging suborder in which to resolve families due to: (i) a pattern of multiple, parallel radiations in this lineage (Fig. S2); (ii) multiple instances of convergent morphologies; (iii) polyphyly; and (iv) incomplete sampling and missing data in our analyses. *Raveneliineae* is the second richest suborder in terms of taxonomic diversity, with ca. 45 accepted genera, of which we were only able to sample representatives from about half and most of these with incomplete locus data.

Host range may be an informative character to place taxa of *Raveneliineae* in families. For example, Savile (1989) predicted *Maravalia sensu* Ono (1984) was polyphyletic, and hypothesised that species on *Fabaceae* belonged to *Raveneliaceae*, supported here with the placement of the *Fabaceae*-infecting *M. limoniformis* within *Raveneliineae* (Figs 1, 3) and the *Apocynaceae*-infecting *U. cryptostegiae* (syn. *M. cryptostegiae*) within *Zaghouaniaceae* (Fig. 1). Likewise, *Triphragmium* Link has evolved elaborate teliospores similar to those in some *Raveneliaceae* where it has been allied in the past

(e.g., Cummins & Hiratsuka 2003); *Triphragmium* species are now known to belong to *Phragmidiaceae* with other *Rosaceae*-infecting rusts (Aime 2006).

We treat four families within *Raveneliineae*, taking into account life cycle and host data, and have taken a conservative approach to assigning genera within families and species to genera until data from type species and/or exemplars from key missing taxa as well as additional loci can be obtained.

**Ochropsoraceae** (Arthur) Aime & McTaggart, **fam. & stat. nov.** MycoBank MB836640.

**Basionym:** *Ochropsoratae* Arthur, *Rés. Sci. Congr. Int. Bot. Vienne*: 336. 1906.

**Type genus:** *Ochropsora* Dietel, *Ber. Dtsch. Bot. Ges.* **13**: 401. 1895.

**Description:** Spermogonia Group VI (type 7). Aecia aecidium-type; uredinia malupa-type; aecial states systemic overwintering as mycelium; telia forming crusts, 1-cell deep, at first subepidermal, then erumpent; teliospores germinate without dormancy, either internally (*Ochropsora*) or externally (*Aplopsora*). Species likely macrocyclic and heteroecious, although gametothallus not known for *Aplopsora*.

**Included genera:** *Aplopsora*, *Ochropsora*; likely includes *Ceraceopsora*.

**Host families:** *Ranunculaceae* (0-I); *Rosaceae*, *Cornaceae* (II-III)

**Notes:** A monophyletic *Ochropsoraceae* as the earliest diverging lineage of *Raveneliineae* was recovered in all of our analyses. *Aplopsora* and *Ochropsora* were previously treated within the artificial *Chaconiaceae* (Cummins & Hiratsuka 2003) where they shared the convergent character of teliospore germination without dormancy.

**Tranzscheliaceae** (Arthur) Aime & McTaggart, **fam. & stat. nov.** MycoBank MB836641.

**Basionym:** *Tranzschelieae* Arthur, *Rés. Sci. Congr. Int. Bot. Vienne*: 340. 1906.


**Type genus:** *Tranzschelia* Arthur, *Rés. Sci. Congr. Int. Vienne*: 340. 1906.

**Description:** Spermogonia Group VI (type 7). Aecia aecidium-type; uredinia uredo-type. Teliospores 2-celled, pedicellate, produced from sterile basal cells. Species are macrocyclic and heteroecious, with some derived microcyclic species.

**Included genera:** *Leucotelium*, *Tranzschelia*.

**Host families:** *Ranunculaceae* (0-I and autoecious species); *Rosaceae* (II-III in heteroecious species).

**Notes:** *Tranzschelia* has held an isolated position within *Pucciniales* in prior molecular studies (Aime 2006) and appears as an independent lineage of *Raveneliineae* in this work (Fig. 1). *Leucotelium* has been treated as a synonym of *Sorataea* (Cummins & Hiratsuka 2003) but retained by Thirumalachar & Cummins (1940) due to the presence of a sterile basal cell layer from which the teliospores develop that is lacking in *Sorataea*.



**Fig. 3.** *Raveneliaceae*. ML topography generated from 28S with expanded taxon sampling. The tree is mid-point rooted. Families are indicated by colour blocks; *Raveneliaceae* is not resolved. Only 26 of the estimated 45+ genera in this suborder are represented by types (indicated in bold) and type proxies (indicated by \*), and poor resolution may be attributable to missing data (both locus and taxon sampling), combined with long branch lengths (Fig. S2) in this lineage. Support for nodes is provided from an approximate likelihood ratio test ( $\geq 0.90$ ) and ultrafast bootstraps ( $\geq 95\%$ ) as aLRT/UFBoot.



Two non-type species of *Sorataea* were included in our analyses and are referable to *Raveneliaceae* (Fig. 3). *Leucotelium* is the sister genus to *Tranzschelia* (Scholler *et al.* 2019), with which it shares a similar host range and teliospore production from sterile sporogenous cells (Thirumalachar & Cummins 1940, López-Franco & Hennen 1990). Many species of *Tranzschelia* are microcyclic on *Ranunculaceae* in accordance with Tranzschel's Law (Scholler *et al.* 2019).

**Phakopsoraceae** Cummins & Y. Hirats., *Illustr. Gen. Rust Fungi, rev. Edn* (St. Paul): 13. 1983. **emend.** Aime & McTaggart

*Type genus:* *Phakopsora* Dietel, *Ber. Deutsch. Bot. Ges.* **13**: 333. 1895.

*Description:* Spermogonia Group VI (type 7). Aecia caeoma-type, some *Masseeëlla* with aecidium-type aecia; uredinia lecythea- or uredo-type. Teliospores 1-celled. *Bubakia*, *Masseeëlla* and *Nothoravenelia* species are autoecious and macrocyclic. The majority of *Phakopsora* and *Uredopeltis* species are only known from the sporothallus.

*Included genera:* *Bubakia*, *Masseeëlla*, *Nothoravenelia*, *Phakopsora*, *Uredopeltis*; likely includes *Arthuria*, *Cerotelium*, *Dicheirinia*, *Monosporidium*, *Phragmidiella*, *Pucciniostele*, *Scalarispora*.

*Host families:* *Annonaceae*, *Bignoniaceae*, *Burseraceae*, *Commelinaceae*, *Euphorbiaceae*, *Fabaceae*, *Myrtaceae*, *Rubiaceae*, *Urticaceae* (0-III).

*Notes:* Both *Phakopsora* and the *Phakopsoraceae* are known to be polyphyletic (e.g., Aime 2006), with more than 100 species currently classified in *Phakopsora s.l.* However, lack of data and differing interpretations of the type have hampered taxonomic progress. The recent designation of a new type species for *Phakopsora*, *P. pachyrhizi* (Aime *et al.* 2019a, b), has stabilized use of the name as applied here, for those genera and species that share a common ancestor with *P. pachyrhizi*. *Phakopsora* remains poorly resolved with our data and consists of two supported clades, one containing *P. pachyrhizi* and its allies and the other containing most of the *Annonaceae*-infecting species, which may represent a separate genus, but were recovered as monophyletic in some analyses (not shown).

The name *Bubakia* is often treated as a synonym of *Phakopsora* (e.g., Cummins & Hiratsuka 2003). Our study shows that *Bubakia argentinensis* belongs to a distinct lineage within *Phakopsoraceae* (Figs 1, 3). Further, *B. argentinensis* shares similar hosts (*Croton* spp.) and characteristics with the type, *B. crotonis*, and we accept *Bubakia* for these species (Mundkur 1943). *Masseeëlla* has previously been treated as *incertae sedis* within *Pucciniales* (Cummins & Hiratsuka 2003), but our data place it within *Phakopsoraceae* (Figs 1, 3).

The majority of *Phakopsora* and *Uredopeltis* species are known only from sporothalli. It is unknown whether gametothalli occur on an alternate host, or whether these species are autoecious. Sporothalli have been described for a few *Phakopsora* species, i.e., *P. breyniae*, *P. innata*, *P. phyllanthidiscoidei*, and *P. stratosa*, which are all autoecious (Berndt & Wood 2012, Ono 2015b), although it is unclear whether these should be retained in *Phakopsora s.s.* or are allied with one of the segregate ex-*Phakopsora* genera.

**Phakopsora pipturi** (Syd.) Aime & McTaggart, **comb. nov.** MycoBank MB836642.

*Basionym:* *Pucciniastrum pipturi* Syd., *Annls mycol.* **29**(3/4):171. 1931.

*Synonym:* *Uredo pipturi* (Syd.) Hirats. f., *Trans. Mycol. Soc. Japan* **5**: 4. 1957.

**Raveneliaceae** Leppik, *Ann. bot. fenn.* **9**: 139. 1972. **emend.** Aime & McTaggart

*Synonyms:* *Chaconiaceae* Cummins & Y. Hirats., *Illustr. Gen. Rust Fungi, rev. Edn* (St. Paul): 14. 1983.

*Uropyxidaceae* Cummins & Y. Hirats., *Illustr. Gen. Rust Fungi, rev. Edn* (St. Paul): 14. 1983.

*Type genus:* *Ravenelia* Berk., *Gard. Chron.* **13**:132. 1853.

*Description:* Spermogonia Group VI (type 5 or 7); aecia uredo- (rarely aecidium-, caeoma-, or lecythea-) type; uredinia uredo-type. Teliospores 1- to many-celled, some species with elaborate compound or multi-celled teliospores. Majority of species autoecious and macrocyclic, with a few derived microcyclic species; many species on mimosoid (*Caesalpinioideae*) hosts.

*Included genera:* *Bibulocystis*, *Cephalotelium*, *Crossospora*, *Diorchidium*, *Endoraecium*, *Kernkampella*, *Lipocystis*, *Newinia*, *Olivea*, *Porotenus*, *Prospodium*, *Ravenelia*, *Sorataea*, *Uromycladium*, *Uropyxis*, *Ypsilospora*; likely includes *Allotelium*, *Anthomyces*, *Anthomycetella*, *Apra*, *Atelocauda*, *Chaconia*, *Cystomyces*, *Diabole*, *Diochordiella*, *Esalque*, *Hennenia*, *Maravalia*, *Mimema*, *Phragmopyxis*, *Spumula*, *Tegillum*.

*Host families:* *Bignoniaceae*, *Euphorbiaceae*, *Fabaceae*, *Ranunculaceae*, *Rosaceae*, *Sapotaceae*, *Verbenaceae* (0-III).

*Notes:* Leppik (1972) limited *Melampsoraceae* to rust species that are heteroecious and temperate, reassigning the autoecious and tropical species to a new family, *Raveneliaceae*. Savile (1989) provided an in-depth study of *Raveneliaceae* and hypothesised that the most recent common ancestor of *Raveneliaceae* was heteroecious, but that the family diversified as autoecious species on mimosoid (*Caesalpinioideae*) hosts after an environmental event severed their association with the initial sporothallus host. This hypothesis finds support in our work, which shows that the two early diverging families of *Raveneliaceae*, *Ochropsoraceae* and *Tranzscheliaceae* (Fig. 1), are heteroecious with sporothalli hosts in *Ranunculaceae*.

*Chaconia*, which we place within *Raveneliaceae*, has been placed variously in the *Melampsoraceae* or with other rust genera in the artificial *Chaconiaceae*. This and prior works have shown *Chaconiaceae*, and most of the genera therein, as polyphyletic. The morphological character on which they were based, specifically thin-walled, pale teliospores that germinate without dormancy, was derived multiple times within *Pucciniales* (Aime, 2006, Aime *et al.* 2018a), as a result of convergent morphologies in species adapted to tropical climates that do not need to overwinter (Savile 1989).

*Uropyxidaceae* consists of an artificial assemblage of rust fungi combining (mostly) 2-celled, transversely septate teliospores and Group VI (type 5) spermogonia (where present). In this study, we sampled nearly all genera of *Uropyxidaceae* as circumscribed by Cummins & Hiratsuka (1983, 2003), most of which had not been previously sequenced; from our results the family is clearly

polyphyletic. Many of the genera placed in *Uropyxidaceae* by Cummins & Hiratsuka (1983, 2003) were once considered allied within *Pucciniaceae* due to similarities in teliospore morphology. Our analysis shows that several of these, *i.e.*, *Desmella*, *Dipyxis*, *Edythea*, *Macruropyxis*, belong to *Pucciniaceae* (Fig. 4). *Dasyospora* is allied in *Sphaerophragmiaceae* and *Tranzschelia* in *Tranzscheliaceae* (Fig. 1). The remaining genera – *Newinia*, *Porotenus*, *Prospodium*, *Sorataea*, and *Uropyxis* – are included within a broadly defined *Raveneliaceae* (Fig. 3).

*Raveneliaceae* is not resolved in our analyses, with strong support for some genera with multiple sampling, but almost no support for infra-familial nodes (Figs 1, 3, S2). Branch lengths for species of *Raveneliaceae* are comparatively long (Figs 3, S2) and may indicate an accelerated evolutionary rate in this family. 28S data alone can be informative for other *Puccinales* lineages (*e.g.*, Ji *et al.* 2019), but are inadequate for resolving relationships of genera, and in many cases even species, within *Raveneliaceae* (Fig. S2).

No sequence data are available for the generic type, *R. glandulosa*, a Western Hemisphere rust of *Tephrosia*. *Ravenelia* sp. (PUR F19717, Fig. 3) shares a host with *R. glandulosa* and may be congeneric with the type. *Maravalia s.s.* as represented by *M. limoniformis* (Figs 1, 3) is likely to belong here.

The genus *Olivea*, as circumscribed in the past, contains a polyphyletic assemblage of species that form a hymenial layer of probasidia that germinate via apical extension. Three species formerly placed in *Olivea* were included in our analyses: (i) *O. capituliformis*, the type for the genus; (ii) *O. scitula*; and (iii) *O. tectonae*, none of which are related to each other (Figs 3 & S2). *Neolivea tectonae* (syn. *O. tectonae*) is placed in the *Crossosporaceae* and discussed there. *Olivea scitula* was considered by Mains (1940) as most similar to *Tegillum fimbriatum*, and we apply the name *T. scitulum* to this species, although further work is necessitated to determine if it is, indeed, congeneric with the type species, *T. fimbriatum*. *Olivea capituliformis* is the only described species in this complex that infects hosts in *Euphorbiaceae*; the ex-*Olivea* species that we treat infect hosts in *Verbenaceae* (Ono & Hennen 1983).

***Cephalotelium evansii*** (Syd. & P. Syd.) Aime & McTaggart, **comb. nov.** MycoBank MB836643.

*Basionym:* *Ravenelia evansii* Syd. & P. Syd. *Annls mycol.* **10**: 440. 1912.

*Synonym:* *Dendroecia evansii* (Syd. & P. Syd.) Syd., *Annls mycol.* **19**: 165. 1921.

***Cephalotelium neocaledoniense*** (B. Huguenin) Aime & McTaggart, **comb. nov.** MycoBank MB837616.

*Basionym:* *Ravenelia neocaledoniensis* B. Huguenin, *Bull. trimest. Soc. mycol. Fr.* **82**: 263 (1966).

***Cephalotelium xanthophloeae*** (M. Ebinghaus *et al.*) Aime & McTaggart, **comb. nov.** MycoBank MB836644.

*Basionym:* *Ravenelia xanthophloeae* M. Ebinghaus *et al.*, *MycoKeys* **43**: 11. 2018.

*Notes:* Of the *ca.* 200 species currently placed in *Ravenelia*, our data consistently resolved as congeneric those we now refer to *Cephalotelium* (Figs 3, S2). These species were also strongly supported as one of two monophyletic groups in *Ravenelia s.l.* by Ebinghaus *et al.* (2018b). *Cephalotelium macowanianum* (syn. *Ravenelia macowanianum*) is the type of *Cephalotelium*.

The formation of telial galls is sometimes induced by infection of *Ravenelia* species, but not by *Cephalotelium* species. In contrast, *C. evansii*, *C. macowanianum* and *C. xanthophloeae* induce aecial gall formation in host tissues, which is a trait that appears to be confined to the *Cephalotelium* lineage (Ebinghaus *et al.* 2018a, b). *Cephalotelium* species infect members of *Vachellia* (*Caesalpinioideae*) in the Eastern Hemisphere (Sydow 1921). *Cephalotelium* is possibly a later synonym for *Dendroecium*, however, the type, *D. farlowiana*, occurs on *Senegalia* (*Caesalpinioideae*) species in the Western Hemisphere (Dietel 1894).

***Lipocystis acaciae-pennatulae*** (Dietel) Aime & McTaggart, **comb. nov.** MycoBank MB836645.

*Basionym:* *Ravenelia acaciae-pennatulae* Dietel, *Beih. bot. Zbl., Abt. 2* **20**: 373. 1906.

*Notes:* *Lipocystis* with the type species *L. caesalpiniae* was described as a monotypic genus for a rust on *Mimosa* from the West Indies. A second species, *Lipocystis acaciae-pennatulae*, infects *Acacia* species in Central America and is congeneric with *L. caesalpiniae* (Figs 1, 3, S2).

***Urediniineae*** Engl., *Syllabus der Vorlesungen über spezielle und medizinisch-pharmazeutische Botanik*: 36. 1892. **emend.** Aime & McTaggart

*Synonym:* *Pucciniineae* Doweld, *Index Fungorum* **77**: 1. 2014.

*Type family:* *Pucciniaceae* Chevall.

*Description:* With the characteristics of the included families. Most species form Group V but also Group VI spermogonia and 1- or 2- celled teliospores but multi-celled telia formed in some or most *Nyssospora*, *Phragmidiaceae*, and *Sphaerophragmiaceae*.

*Included families:* *Crossosporaceae*, *Gymnosporangiaceae*, *Phragmidiaceae*, *Pileolariaceae*, *Pucciniaceae*, *Sphaerophragmiaceae*.

*Notes:* *Urediniineae* is the largest suborder in both species numbers and generic diversity. *Pucciniineae* is a superfluous name for the older *Urediniineae*. We were able to sample types or type representatives for 50 of the *ca.* 70 genera placed here as well as several species currently assigned to form-genera.

We were unable to resolve the placement for three genera: *Allodus*, *Neopuccinia*, and *Nyssospora*. *Allodus* was long considered a synonym of *Puccinia* due to its pedicellate, 2-celled teliospores. Minnis *et al.* (2012) resurrected *Allodus* as an orphan genus of uncertain placement. Our analyses occasionally resolved *Allodus* as sister to *Peridiopsora mori* with weak support (not shown). Only a single 28S sequence is available for the newly described *Neopuccinia*, which shares many similarities with *Kimuromyces* (Dianese *et al.* 1995). Connections between *Nyssospora* and *Sphaerophragmium* have been noted by Lohsomboon *et al.* (1994). *Nyssospora* was recovered as sister to *Sphaerophragmiaceae* in some but not all of our analyses (Figs 1, 4) and may represent a separate family lineage.

***Phragmidiaceae*** Corda *Icon. fung.* (Prague) **1**: 6. 1837.

*Type genus:* *Phragmidium* Link, *Mag. Ges. Naturfr. Freunde Berlin* **7**: 30. 1816.



**Fig. 4. Uredineae.** ML topography generated from three concatenated loci (28S, 18S, and CO3) with expanded taxon sampling. The tree is rooted with *Melampsorineae*. Six families are resolved and indicated by coloured blocks; three genera are unresolved to family and indicated as *incertae sedis*. Genera represented by types are indicated in bold; genera represented by type proxies (as explained in methods) are indicated by \*. Support for nodes is provided from an approximate likelihood ratio test ( $\geq 0.90$ ), ultrafast bootstraps ( $\geq 95\%$ ) and genealogical concordance factors for the three loci at each node as aLRT/UFBoot/gCF.

**Description:** Spermogonia of Group IV (various types); aecia variable, caeoma-, petersonia- or uredo-type; uredinia lecythea- or uredo-type. Teliospores mostly multi-celled, usually by transverse septa. Species autoecious on *Rosoideae* subfamily of *Rosaceae*.

**Included genera:** *Gerwasia*, *Gymnoconia*, *Hamaspora*, *Kuehneola*, *Phragmidium*, *Trachyspora*, *Triphragmium*, *Xenodochus*; likely includes *Joerstadia*.

**Host family:** *Rosaceae* (0-III).

**Notes:** Convergence in teliospore morphology between some genera of *Phragmidiaceae* and *Raveneliaceae* has been previously noted (e.g., Cummins & Hiratsuka 2003); Aime (2006) showed that *Phragmidiaceae* species are confined almost

exclusively to the *Rosoideae* in contrast to *Raveneliaceae*.

**Pileolariaceae** (Arthur) Cummins & Y. Hirats., *Illustr. Gen. Rust Fungi, rev. Edn* (St. Paul): 14. 1983. *emend.* Aime & McTaggart

**Type genus:** *Pileolaria* Castagne, *Obs. Plantes Acotylédonées Fam. Urédinées* 1: 22. 1842.

**Description:** Spermogonia Group VI (type 7). Aecia and uredinia uredo-type. Teliospores 1-celled with characteristic sculpted appearance; germination external after dormancy. Species mostly macrocyclic and autoecious.

**Included genus:** *Pileolaria*.

**Host family:** *Anacardiaceae* (0-III).

*Notes:* *Pileolariaceae* was established for autoecious rusts in *Pileolaria*, *Uromykladium* and *Endoraecium* (Arthur 1906, Cummins & Hiratsuka 2003). The latter two have been resolved within *Raveneliaceae*, while *Pileolaria* holds an isolated position within *Pucciniales* (Aime 2006, Scholler & Aime 2006, Figs 1, 4). *Pileolaria* species are autoecious on *Anacardiaceae*, with very characteristic sculpted discoid teliospores.

**Crossopsoraceae** Aime & McTaggart, **fam. nov.** MycoBank MB836646.

*Type genus:* *Crossopsora* Syd. & P. Syd., *Annls mycol.* **16**(3/6): 243. 1919.

*Diagnosis:* Similar to *Phakopsoraceae*, differing in that the majority of sporothalli infect *Poaceae*, *Vitaceae*, *Lamiaceae*, and *Rhamnaceae* with none known on *Annonaceae* and *Euphorbiaceae* and that some species are known to be heteroecious.

*Description:* Spermogonia Group VI (type 7) where known; aecia aecidium-type where known; uredinia typically paraphysate, malupa-type; teliospores germinate externally, with or without dormancy, 1-celled, compact, often produced in catenulate chains of a few to many cells. Most species only known from the sporothallus; *Neophysopella* is macrocyclic and heteroecious, as may be other species in this family.

*Included genera:* *Angiopsora*, *Catenulopsora*, *Crossopsora*, *Kweilingia* (= *Dasturella*), *Neolivea*, *Neophysopella*, *Stomatissora*.

*Host families:* *Papaveraceae*, *Sabiaceae*, *Rubiaceae* (0-I); *Lamiaceae*, *Fabaceae*, *Poaceae*, *Rhamnaceae*, *Rubiaceae*, *Salicaceae*, *Vitaceae* (II-III).

*Notes:* *Phakopsoraceae sensu* Cummins & Hiratsuka (2003) is a polyphyletic family, with multiple polyphyletic genera (Aime 2006, Aime *et al.* 2018a, 2019a, b, Ji *et al.* 2019). The phakopsoroid fungi share a convergent suite of characters including pale, subepidermal sori, and 1-celled, sessile teliospores with external germination, which have long been the subject of taxonomic debate, especially concerning application of the names *Angiopsora*, *Bubakia*, *Phakopsora*, and *Physopella*. *Phakopsoraceae s.s.* is now confined to species within *Raveneliineae* on various hosts including *Annonaceae*, *Euphorbiaceae* and *Fabaceae*. The remaining ex-*Phakopsoraceae* are now placed in *Crossopsoraceae* (Figs 1, 4), including the species that have radiated on *Poaceae* and *Vitaceae*. Some species in both families form teliospores in extruded columns, but these are produced in catenulate chains in *Crossopsoraceae*, versus tightly packed individual cells in *Phakopsoraceae*. Life cycles are unknown for many species. However, *Neophysopella* is heteroecious, alternating either between *Sabiaceae* and *Vitaceae* species (Ji *et al.* 2019) or *Papaveraceae* and *Fabaceae* (*N. kraunhiae*, Hiratsuka & Kaneko 1978); *K. divina* alternates between *Rubiaceae* (gametothallus) and *Poaceae* (sporothallus) (Thirumalachar *et al.* 1947).

*Physopella* Arthur (1906), although often applied to the species now assigned to *Neophysopella* and others, is a later homonym of *Physopella* G. Poirault (1905) and is therefore an illegitimate name (Xi *et al.* 2019). Our work (Fig. 4) shows that *Kweilingia* (type *K. bambusae*, syn. *Chrysomyxa bambusae*)

and *Dasturella* (type *D. divina*, syn. *Angiopsora divina*) are congeneric; *Kweilingia* (1940) has priority over *Dasturella* (1943) and we retain these species in the former genus.

The phakopsoroid species that form teliospore chains are difficult to diagnose and classify, as exemplified by the complex taxonomic histories of *Crossopsora*, *Cerotelium*, *Catenulopsora*, and *Kuehneola* (*e.g.*, Ono 2015a). In this work, *Crossopsora* and *Catenulopsora* are assigned to *Crossopsoraceae*. *Cerotelium* most likely belongs to *Phakopsoraceae s.s.*; the type, *C. canavaliae* parasitizes *Fabaceae* and the uredinia are peridiate in contrast to *C. fici* with paraphysate uredinia (Cummins 1941). Nonetheless, these genera still appear to be polyphyletic with little support for generic lineages and resolution will require additional taxon and locus sampling (Fig. 4).

Numerous *Uredo* species assigned to *Crossopsoraceae*, especially within the *Crossopsora/Catenulopsora* complex, could not be placed in other genera and we have retained use of names in anamorphic form-genera for these. *Crossopsora fici* and *U. moricola* form a distinct lineage within *Crossopsoraceae* and may require a new genus, pending examination of other critical types including those of *Mehtamyces*, *Phragmidiella*, *Pucciniostele*, and *Scalarispora*.

**Neophysopella kraunhiae** (Dietel) Aime & McTaggart, **comb. nov.** MycoBank MB837747.

*Basionym:* *Phakopsora kraunhiae* Dietel, *Hedwigia* **41**: 178. 1902.

*Synonyms:* *Ochropsora kraunhiae* (Dietel) Dietel., *Bot. Jahrb.* **37**: 106. 1905.

*Aecidium corydalinum* Syd. & P. Syd., *Monogr. Ured.* **4**: 235.1923.

*Notes:* *Neophysopella kraunhiae* is heteroecious and produces gametothalli on *Corydalis incisa* (*Papaveraceae*) and sporothalli on *Wisteria floribunda* (*Fabaceae*) (Hiratsuka & Kaneko 1978). Our data support its classification in the newly circumscribed *Neophysopella*, a genus that contains most other known heteroecious species in *Crossopsoraceae*.

**Neolivea** Aime & McTaggart, **gen. nov.** MycoBank MB837748.

*Type species:* *Neolivea tectonae* (T.S. Ramakr. & K. Ramakr.) Aime & McTaggart, this paper.

*Etymology:* New genus segregated from *Olivea*.

*Diagnosis:* Similar to *Olivea* and *Tegillum* but differs in having subglobose to ellipsoid, non-angular urediniospores with inconspicuous germ pores, and waxy telia.

*Description:* With characteristics of the type species *Neolivea tectonae*.

**Neolivea tectonae** (Racib.) Aime & McTaggart, **comb. nov.** MycoBank MB837749.

*Basionym:* *Uredo tectonae* Racib., *Parasit. Alg. Pilze Java's* (Jakarta) **1**: 28. 1900.

*Synonyms:* *Olivea tectonae* (Racib.) Thirum., *Curr. Sci.* **18**: 176. 1949.

*Tegillum tectonae* (Racib.) Doweld, *Index Fungorum* **36**: 1. 2013. *Chaonia tectonae* T.S. Ramakr. & K. Ramakr., *Indian Phytopath.* **2**: 19. 1949.

*Olivea tectonae* (T.S. Ramakr. & K. Ramakr.) R.L. Mulder, *CMI*

*Descriptions of Pathogenic Fungi and Bacteria* **37**: no. 365. 1973.  
*Olivea neotectonae* Buriticá & Salazar-Yepes, *Revista Fac. Agron. Medellín* **60**(1): 3652. 2007.  
*Tegillum neotectonae* (Buriticá & Salazar-Yepes) Doweld, *Index Fungorum* **36**: 1. 2013.

**Notes:** *Neolivea tectonae* causes leaf rust on teak (*Tectona grandis*, *Tectoneae*, *Verbenaceae*). It is described in Ono & Hennen (1983) and illustrated in Ramakrishnan & Ramakrishnan (1949). Our data show that *N. tectonae* is unrelated to the type of *Olivea* (*O. capituliformis*), which belongs to the *Raveneliaceae* (Figs 1, 3, 4). *Tegillum* was established for *Olivea*-like species, with the type *T. fimbriatum* (Mains 1940). While no type data exist for *T. fimbriata*, the species is most similar to *T. scitulum* (Mains 1940), sharing characteristics such as lobed or angular urediniospores with germ pores residing in the lobes and hosts in the *Vitex* group of *Verbenaceae* (Ono & Hennen 1983), which is placed in *Raveneliineae* (Fig. 1). *Neolivea tectonae* differs from other described *Olivea* and *Tegillum* species in producing rounded urediniospores with inconspicuous germ pores, as well as waxy, orange telia (Ono & Hennen 1983, Osorio *et al.* 2019).

***Angiopsora apoda*** (Har. & Pat.) Aime & McTaggart, **comb. nov.** MycoBank MB836647.

**Basionym:** *Puccinia apoda* Har. & Pat., *Bull. Mus. natn. Hist. nat.*, Paris **15**:199. 1909.

**Synonym:** *Phakopsora apoda* (Har. & Pat.) Mains, *Mycologia* **30**: 45. 1938.

***Angiopsora chusqueae*** (Pardo-Card.) Aime & McTaggart, **comb. nov.** MycoBank MB836648.

**Basionym:** *Uredo chusqueae* Pardo-Card., *Revta Acad. colomb. cienc. exact. fis. nat.* **20**: 205. 1996.

***Angiopsora paspalicola*** (Henn.) Aime & McTaggart, **comb. nov.** MycoBank MB836649.

**Basionym:** *Uredo paspalicola* Henn., *Hedwigia* **44**: 57. 1905.

**Synonyms:** *Puccinia paspalicola* (Henn.) Arthur, *Manual Rusts U.S. & Canada. Purdue Res. Found.*: 127. 1934.

*Physopella paspalicola* (Henn.) Buriticá & Hennen, *Buriticá Rev. I. C. N. E. (Medellín)* **5**: 179. 1994.

*Puccinia compressa* Arthur & Holway, *Proc. American Phil. Soc.* **64**:257. 1925.

*Angiopsora compressa* (Arthur & Holway) Mains, *Mycologia* **26**: 29. 1934.

*Physopella compressa* (Arthur & Holway) Cummins & Ramachar, *Mycologia* **50**: 742. 1958.

*Phakopsora compressa* (Arthur & Holway) Buriticá & Hennen, *Buriticá, Rev. I. C. N. E. (Medellín)* **5**: 179. 1994.

**Notes:** Mains (1934) established *Angiopsora* for *Poaceae*-infecting ex-*Pucciniaceae* species with similarities to *Phakopsora*. Although the name has been considered a synonym for *Phakopsora* (e.g., Cummins & Hiratsuka 2003), we find that it is applicable to numerous former *Phakopsora* species on grasses (Fig. 4).

***Gymnosporangiaceae*** P. Zhou & L. Cai, *Persoonia* **45**: 79. 2020. **emend.** Aime & McTaggart

**Synonym:** *Gymnosporangieae* Dietel, In: Engler & Prantl, *Naturlichen Pflanzenfamilien Band* **6**: 73. 1938.

**Type genus:** *Gymnosporangium* R. Hedw. ex DC., In: Lamarck & de Candolle, *Fl. franç.*, *Edn 3* (Paris) **2**: 216. 1805.

**Description:** Spermogonia Group V (type 4). Aecia roestelia-type (*Gymnosporangium*) or less frequently aecidium-type (*Gymnotelium*). Teliospores mostly 2-celled, germinating without dormancy via external basidia. Life cycles mostly demicyclic and heteroecious (*Gymnosporangium*).

**Included genera:** *Gymnosporangium*, *Gymnotelium*; likely includes *Peridiopsora*.

**Host families:** *Rosaceae*, *Hydrangeaceae*, *Myricaceae* (0-I); *Cupressaceae* (II-III); autoecious species on *Berberidaceae*, *Cupressaceae*, *Liliaceae*, *Moraceae*.

**Notes:** The genus *Gymnosporangium* is unusual in that temperate species form teliospores in the spring and aecia in the summer, in contrast to most other temperate rust species (Kern 1960). Most species of *Gymnosporangium* form sporothalli on *Juniperus* spp. and gametothalli on *Maloideae* (*Rosaceae*). Although traditionally placed in *Pucciniaceae*, *Gymnosporangium* has been treated as an “orphan” genus in molecular phylogenetic studies (Maier 2003, Aime 2006, Aime *et al.* 2018a) and recently established as a monotypic family of unresolved placement (Zhao *et al.* 2020). Although the older name *Gymnosporangieae* Dietel exists for this lineage, priority only applies within rank, thus we retain *Gymnosporangiaceae* P. Zhou & L. Cai for the family. In addition to the puccinioid character of 2-celled, pedicellate teliospores, most species are demicyclic. *Gymnosporangium* has been conserved against the older name *Roestelia* Rebent. (Aime *et al.* 2018b).

*Peridiopsora mori* (syn. *Caeoma mori*, *Aecidium mori*) causes a well-known disease in mulberries. It is unusual in having a true hemicyclic life cycle, wherein the mitospores appear to function as urediniospores although likely are derived from the aecial part of the life cycle (Mordue 1991). Spermogonia and teliospores are not known. *Peridiopsora* was erected to accommodate an unusual asexual rust that persisted as cyclical urediniospores, producing uredinia in peridiate sori that lack paraphyses, the latter being typical of aecia but rare in uredinia (Kamat & Sathe 1969) and which we interpret as uredinioid aecia. Whether *P. mori* will prove to be congeneric with the type, *P. adelocaryi*, or will be placed within *Gymnotelium*, remains uncertain.

***Gymnotelium speciosum*** (Peck) Aime & McTaggart, **comb. nov.** MycoBank MB836652.

**Basionym:** *Gymnosporangium speciosum* Peck, *Bot. Gaz.* (Crawfordsville) **4** (10): 217. 1879.

**Synonym:** *Tremella speciosa* (Peck) Arthur, *Proc. Indiana Acad. Sci.* **1900**: 135. 1901.

**Notes:** *Gymnotelium* was erected for *Gyt. nootkatense*, a macrocyclic (rather than demicyclic) species with a gametothallus host (*Chamaecyparis*) not within the usual *Maloideae/Juniperus* range of *Gymnosporangium*. Arthur (1929) considered the diagnostic character of aecidium-like rather than rostrate aecia as more important than presence of uredinia, transferring an additional two species, *Gyt. blasdaleanum* (syn. *G. libocedri*) and *Gyt. myricatum* (syn. *G. ellisii*) to *Gymnotelium*. In later works Arthur (1934) treated *Gymnotelium* as a subgenus of *Gymnosporangium*, including a fourth species, *G. speciosum*.

Novick (2008) showed that *G. nootkatense*, *G. blasdaleanum* (as *G. libocedri*), *G. myricatum* (as *G. elisii*), and *G. speciosum* are the earliest diverging members of *Gymnosporangium*. The cupulate aecia of *Gymnotelium* bear a strong resemblance to those of *P. mori*, and our work resolves these together as the sister lineage to *Gymnosporangium* (Figs 1, 4). In addition to producing cupulate (rather than rostrate) aecia, *Gymnotelium* species produce their sporothallus on hosts other than *Maloideae* spp., and/or the sporothallus on hosts other than *Juniperus* spp. (Kern 1960).

**Sphaerophragmiaceae** Cummins & Y. Hirats., *Illustr. Gen. Rust Fungi*, rev. Edn (St. Paul): 15. 1983. **emend.** Aime & McTaggart  
*Synonym:* *Dasysporatae* Arthur, *North American Flora* 7: 807. 1926.

*Type genus:* *Sphaerophragmium* Magnus Ber. dt. bot. Ges 9: 121. 1891.

*Description:* Spermogonia mostly lacking, Group V (type 4) in *Sphenorhynchium*. Aecia aecidium-type; uredinia resembling aecia, lecythea-type in *Sphenorhynchium*. Teliospores 2- to multi-celled. Species autoecious with variable life cycles.

*Included genera:* *Austropuccinia*, *Dasyspora*, *Puccorchidium*, *Sphaerophragmium*, *Sphenorhynchium*.

*Host families:* *Annonaceae*, *Fabaceae*, *Myrtaceae*.

*Notes:* *Sphaerophragmium* has been hypothesized as belonging to *Raveneliaceae* based on similarities in teliospores and host (Cummins & Hiratsuka 2003). More recently *Sphaerophragmium* was shown to belong to a monophyletic lineage (Beenken & Wood 2015) for which the oldest available name is *Dasysporatae*, but for which *Sphaerophragmiaceae* has been recently applied (e.g., McTaggart *et al.* 2016, Beenken 2017). Because priority only applies within rank, we retain the more widely used name for this family. *Austropuccinia psidii*, causes an economically important epidemic disease of *Myrtaceae* and was recently demonstrated to be autoecious and macrocyclic but lacking spermogonia (I-IV) and with uredinioid aecia (McTaggart *et al.* 2018, 2020).

**Pucciniaceae** Chevall., *Fl. gén. env. Paris* (Paris) 1: 413. 1826. **emend.** Aime & McTaggart

*Synonyms:* *Puccinosiraceae* (Dietel) Cummins & Y. Hirats., *Illustrated Genera of Rust Fungi*: 15. 1983.

*Endophyllaceae* Dietel, In: Engler & Prantl, *Nat. Pflanzenfam., Teil. I* (Leipzig) 1: 35. 1897.

*Type genus:* *Puccinia* Pers., *Synopsis methodica fungorum*: 225. 1801.

*Description:* Spermogonia Group V (type 4). Aecia aecidium-type; uredinia mostly uredo-type. Teliospores borne singly, mostly pedicellate, typically with 1 or 2 cells. Basidia external. Most species macrocyclic but many endocyclic and microcyclic species as well; heteroecious or autoecious.

*Included genera:* *Baeodromus*, *Ceratocoma*, *Chardonella*, *Chrycozelis*, *Cionothrix*, *Cumminsia*, *Desmella*, *Didymopsis*, *Dietelia*, *Dipyxis*, *Edythea*, *Endophylloides*, *Endophyllum*,

*Hapalophragmium*, *Leptopuccinia*, *Macruropyxis*, *Miyagia*, *Puccinia*, *Puccinosira*, *Sphenospora*, *Stereostratum*, *Uromyces*, *Xenosteles*; likely includes *Cerradoa*, *Chrysellia*, *Chrysocyclus*, *Chrysopsora*, *Cleptomycetes*, *Corbulopsora*, *Kernella*, *Polioma*, *Trichopsora*.

*Host families:* various including *Berberidaceae*, *Ranunculaceae* (0-I); various including *Asteraceae*, *Euphorbiaceae*, *Fabaceae*, *Malvaceae*, *Orchidaceae*, *Poaceae*, *Solanaceae* (II-III and autoecious species).

*Notes:* As with *Raveneliaceae*, infra-familial relationships in *Pucciniaceae* are difficult to resolve at least within the context of currently circumscribed genera (Fig. 4). However, whereas *Raveneliaceae* represents a lineage with multiple parallel radiations and differing rates of evolution, *Pucciniaceae* represents a recent and rapidly radiating lineage that has undergone multiple losses of teliospore septa and morphs. The majority of rust species and the two most speciose genera, *Puccinia* and *Uromyces*, belong here. Taxonomic changes that reflect natural genera will depend on sequencing of additional loci as well as representative types from genera such as *Cerradoa*, *Cleptomycetes*, *Corbulopsora*, *Didymopsis*, *Polioma*, *Kernella*, *Ramakrishnania*, and *Trichopsora*. *Pucciniaceae* includes many endocyclic derivatives, such as those once included in the family *Puccinosiraceae*, as hypothesized by Berndt (2018).

Species of *Pucciniaceae* can be roughly circumscribed into three radiations (Figs 1, 4). One, represented by *P. graminis*, the type of *Puccinia*, includes primarily species with 2-celled teliospores that are macrocyclic and heteroecious, and includes economically important species on *Poaceae* and *Asteraceae*. The second, represented by *U. appendiculatus*, the type of *Uromyces*, includes many species with 1-celled teliospores that have radiated on *Euphorbiaceae* and *Fabaceae*, and are primarily autoecious. A third, weakly supported in Fig. 1, but not in Fig. 4, includes, among others, the fern- and orchid-infecting species of *Pucciniaceae*.

## DISCUSSION

The proposed classification of rust fungi includes seven suborders and 18 families. Although we treat approximately the same number of families as Cummins & Hiratsuka (2003), the disposition of many genera varies considerably between our and earlier classifications (Fig. S1). We have excluded demonstrably artificial families such as *Chaconiaceae*, *Puccinosiraceae*, and *Uropyxidaceae*, while establishing new families for cohorts that have been repeatedly shown in our and other works to be strongly supported entities (e.g., *Crossosporaceae*, *Milesiaceae*, *Rogerpetersoniaceae*). While our three-locus approach has enabled resolution of most major lineages of *Puccinales*, providing a stable framework for future studies, resolution remains poor in some areas, especially in the *Pucciniastraceae* and *Raveneliaceae*, which will require additional strategic sampling of both taxa and loci. Additionally, three of our sampled genera – *Allodus*, *Neopuccinia*, and *Nyssopsora* – still cannot be placed to family and remain *incertae sedis* within *Urediniales*; placement of these is sensitive to taxon selection within analyses. For example, in our initial analyses (not shown) *Allodus* was weakly supported within *Gymnosporangiaceae*, however, inclusion of *Nyssopsora* (Fig. 1) and *Neopuccinia*

(Fig. 4) results in uncertain placement for all three genera and underscores the need for appropriate taxon selection in reconstructing phylogenetic hypotheses in *Pucciniales*. The necessity of including type species and adequate sampling when dealing with rust taxonomy, especially in polyphyletic and/or convergent genera and families, cannot be overemphasized.

The classification of rust fungi has undergone several epochs, each with emphasis on a different suite of characters including teliospores (e.g., Cunningham 1931) and telia (e.g., Dietel 1928), aecia (e.g., Leppik 1953), and spermogonia (Hiratsuka & Cummins 1963, Hiratsuka & Hiratsuka 1980) in attempts to circumscribe natural groups (Fig. S1). Our work shows that sporothallus characteristics can be useful at the generic and sometimes family ranks. Ontogenic characters, such as the presence of hymenial-like sporogenous cells from which teliospores develop, which distinguishes *Tranzscheliaceae* from *Ochropsoraceae*, may be useful for diagnosing some families. However, in general telial and uredinial characters are among the most homoplasious in rust taxonomy.

*The asexual genera* – At least 34 generic names for asexual rust morphs have been introduced. Of these, ca. 13 were in wide use (Cummins & Hiratsuka 2003) prior to changes in the nomenclatural code that now eliminate the use of dual nomenclature (McNeill *et al.* 2012, Turland *et al.* 2018). Most of these genera are recognized as later synonyms for sexual-morph genera (e.g., *Canasta* = *Prospodium*, *Endocronartium* = *Cronartium*, *Pelastoma* = *Blastospora*) or in cases where the asexual name has priority, the sexual name has been conserved (e.g., *Gymnosporangium* over *Roestelia*; *Melampsorella* over *Peridermium*) (Aime *et al.* 2018b). In many cases, species only known from an asexual morph can be recombined into sexual genera by a combination of host, morphology, and/or DNA sequence data. But for the largest asexual genera, *Aecidium* and *Uredo*, there remain hundreds of species that cannot reliably be recombined at this time, and the process to place these within natural genera will take painstaking work, even with DNA data. In this paper we were unable to assign several species (e.g., *A. kalanchoes*, *U. cryptostegiae*, *U. elephantopodis*, *U. hiulca* Cummins, *U. trichosanthes*) to sexual genera and recommend use of form-genera names for these species until they can be confidently reassigned.

*The endocyclic rusts*—Species with endocyclic life cycles, *i.e.*, reduced autoecious life cycles in which the aeciospores function as teliospores, were once treated collectively in heterogeneous families such as *Puccinosiraceae* and *Endophyllaceae* (Cummins & Hiratsuka 2003, Buriticá 1991). Perhaps not surprisingly we find that these are derived multiple times within *Pucciniales* as predicted by Berndt (2018) (e.g., *Baeodromus*, *Ceratocoma*, *Chardonella*, *Cionothrix*, *Didymopsisora*, *Dietelia*, *Endophyllum* and *Puccinosira*, Fig. 4). Interestingly, we see little evidence for expanded radiations of endocyclic rusts. Coupled with the fact that the majority of these species are found within the most recent radiation (*Pucciniaceae*), this life history strategy may represent an evolutionary dead-end for *Pucciniales*.

*The chaconiaceous rusts*—Classification of the primarily tropical rust species that produce thin-walled teliospores and germinate without dormancy has not received consensus in the past. Genera such as *Chaconia*, *Goplana*, *Hemileia*, *Maravalia*, *Ochropsora*, and *Olivea* were often treated as a single family, *Chaconiaceae*, hypothesized to represent the earliest diverging lineage of rusts by some urediniologists (e.g., Cummins & Hiratsuka 1983, Hiratsuka 1983). Our and prior works (e.g., Aime

2006, Aime *et al.* 2018a) show that these characters represent a syndrome, most likely as an adaptive response to tropical climates where teliospore dormancy or overwintering is not necessary, and that even the species within genera (with the probable exception of *Hemileia*) do not share a common recent ancestor (Figs 1–4).

*Rust evolution*—The study of rust fungi has been fertile ground for evolutionary theory. Researchers have posited co-evolution to explain rust success (Savile 1971, McTaggart *et al.* 2015), or alternatively host jumps (Hart 1988, McTaggart *et al.* 2016). However, most studies fail to take the heteroecious nature of many rust species into account. The most likely explanation for the success, in terms of species, of *Pucciniales* proposes a combination of both factors in a more complex interplay between the forces of biological specialization and biogenic radiation (Leppik 1953, van der Merwe *et al.* 2008, Aime *et al.* 2018a). A pattern of host jumps followed by rapid radiation to related or ecologically co-distributed host species, or biogenic radiation, is the best explanation for the relationship between rusts and their sporothallus hosts (van der Merwe *et al.* 2008, Aime *et al.* 2018). In contrast, a pattern of co-evolution or biological specialization, best explains the relationship between rust fungi and their gametothallus hosts (Aime *et al.* 2018a). This pattern becomes more evident in light of spermogonial evolution. The earliest rusts (*Rogerpetersoniaceae* and *Zaghouaniaceae*) produce spermogonia of Group III, deep-seated with an exit canal, convex hymenium, and indeterminate growth. Loss of the exit canal occurred approximately 145 mya coinciding with a gametothallus jump to *Pinaceae* hosts. These initially produced subepidermal convex hymenia [Group I (type 1); *Araucariomyces* and *Skierkaceae*], becoming flat and eventually subcuticular [Group I (types 2 & 3)] within the *Melampsorineae*. Well-developed periphyses (Group VI) are apparent approximately 115 mya, coinciding with a gametothallus jump to angiosperms. Group VI spermogonia have a flat hymenium and are retained in *Raveneliineae* and the earlier diverging *Urediniineae*. A well-defined bounding structure surrounding a convex hymenium (Group V) evolved last, ca. 85–90 mya, coinciding with the major angiosperm radiation, in the crown rust radiation that includes *Gymnosporangiaceae*, *Sphaerophragmiaceae*, and *Pucciniaceae*.

The complexities of the macrocyclic rust life cycle and its derivations is difficult to explain in an evolutionary context. One explanation, Tranzschel's law (Arthur 1929, Jackson 1931), posits that autoecious-microcyclic rusts are derived from heteroecious-macrocyclic life cycles that have been restricted to the gametothallus host. Evidence of this has been found at the scale of recently diverging, correlated species (Scholler *et al.* 2019). The authors posit that this plasticity of life cycle provides the template for multiple avenues of speciation, not just on the gametothallus host as demonstrated, but hypothetically from the sporothallus host as well (Scholler *et al.* 2019). At a larger scale, *Raveneliineae* is hypothesized as an initially heteroecious and macrocyclic lineage (Savile 1989) and the earliest diverging families in the suborder, *Ochropsoraceae* and *Tranzscheliaceae*, share this strategy, with gametothallus hosts in *Ranunculaceae*. Loss of the original gametothallus host may have occurred ca. 80 mya, driving the evolution of autoecious and macrocyclic lineages on the sporothallus hosts, with several contemporaneous radiations into what are now recognizable as the *Phakopsoraceae* and *Raveneliaceae* (Fig. 1).

Lineages restricted due to extinction or other forces to

gametothallus hosts compensate by becoming microcyclic (e.g., microcyclic *Tranzschelia* species), endocyclic (e.g., endocronartium-type species of *Cronartium*) or systemic infections (as hypothesized for *Rogerpetersonia* and *Araucariomyces*), but these show constraints, at least in terms of species diversification on that host and seem incapable of regaining all five spore stages. In contrast, lineages that have been restricted on the sporothallus hosts, e.g., *Raveneliaceae*, appear capable of more expansive radiation and in recovering lost spore stages.

**In conclusion** – We provide a rust tree of life resolved at the deeper nodes and use this framework to redefine the higher rank (suborder and family) classification for *Pucciniales*. The complexities of rust fungal biology that includes alternation of generations, heteroecism, and five developmental stages is mirrored in the taxonomic complexities encountered in this group. Taxonomy is often confounded by multiple parallel radiations, convergent morphologies, and the previous application of dual nomenclature making the necessity of consulting type species in taxonomic revision an imperative.

Our data support a model of rust evolution in which: 1) heteroecism favours diversity by allowing different speciation processes to act on different parts of the life cycle (biogenic radiation on the sporothallus, biologic specialization on the gametothallus); 2) the five developmental stages of a macrocyclic rust provide the templates for multiple avenues of speciation; and, 3) both unique features were present in the earliest extant rust radiation (*Mikronegeriineae*). There is evidence that the highly reduced endocyclic rusts on gametothallus hosts may represent an evolutionary dead-end but that rusts confined to the sporothallus host can regain lost spore stages. The heteroecious macrocyclic strategy is unique to *Pucciniales* and may help to explain the tremendous diversity in form and in species found in the rusts.

## ACKNOWLEDGEMENTS

MCA acknowledges the U.S. National Science Foundation, the U.S. Department of Agriculture, The Mellon Foundation, the Louisiana Board of Regents, and the Indiana Academy of Sciences for funding in support of various parts of this project. ARM acknowledges the University of Queensland Development Fellowships (UQFEL1718905) and support from the Department of the Environment and Energy under the Australian Biological Resources Study (grant numbers RG18-43 and RFL212-33). We are extremely grateful to Reinhard Berndt and Amy Rossman for numerous discussions on rust fungus classification and taxonomy, to Roger Shivas for his contributions of specimens and mentorship to ARM along the journey, to Amy Rossman and Roger Shivas for critical review of earlier versions of this manuscript, and to two anonymous reviewers for pointing out errors and providing helpful comments. Herbaria PUR, BPI, and BRIP provided loans of material without which this study would not have been possible.

**Conflict of interest:** The authors declare that there is no conflict of interest.

## REFERENCES

Aime MC (2006). Toward resolving family-level relationships in rust fungi (*Uredinales*). *Mycoscience* **47**: 112–122.

- Aime MC, Bell CD, Wilson AW (2018a). Deconstructing the evolutionary complexity between rust fungi (*Pucciniales*) and their plant hosts. *Studies in Mycology* **89**: 143–152.
- Aime MC, Castlebury LA, Abbasi M, et al. (2018b). Competing sexual and asexual generic names in *Pucciniomycotina* and *Ustilaginomycotina* (*Basidiomycota*) and recommendations for use. *IMA Fungus* **9**: 75–90.
- Aime MC, Matheny PB, Henk DA, et al. (2006). An overview of the higher-level classification of *Pucciniomycotina* based on combined analyses of nuclear large and small subunit rDNA sequences. *Mycologia* **98**: 896–905.
- Aime MC, McTaggart AR, Mondo SJ, et al. (2017). Phylogenetics and phylogenomics of rust fungi. *Advances in Genetics* **100**: 267–307.
- Aime MC, Rossman AY, Ono Y, et al. (2019a). (2688) Proposal to conserve the name *Phakopsora* (*Basidiomycota*, *Pucciniales*) with a conserved type. *Taxon* **68**: 592.
- Aime MC, Rossman AY, Ono Y, et al. (2019b). (2689–2690) Proposals to conserve the names *Phakopsora pachyrhizi* against *Uredo erythrinae* and *U. sojae* (*Malupa sojae*) and *Physopella meibomiaae* (*Phakopsora meibomiaae*) against *Aecidium crotalariicola*, *U. teramni*, and *U. vignae* (*M. vignae*) (*Basidiomycota*, *Pucciniales*). *Taxon* **68**: 593–594.
- Altschul SF, Gish W, Miller W, et al. (1990). Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403–410.
- Arthur JC (1906). *Résultats scientifiques du Congrès international de Botanique Wien 1905*: 331–348.
- Arthur JC (1907–1931). *Uredinales. North American Flora* **7**: 83–969.
- Arthur JC (1929). *The Plant Rusts (Uredinales)*. John Wiley & Sons, New York, USA.
- Arthur JC (1934). *Manual of the rusts in the US and Canada*. Purdue Res. Foundation, USA.
- Bakshi BK, Singh S (1960). A new genus in plant rusts. *Canadian Journal of Botany* **38**: 259–262.
- Beenken L (2014). *Pucciniales* on *Annona* (*Annonaceae*) with special focus on the genus *Phakopsora*. *Mycological Progress* **13**: 791–809.
- Beenken L (2017). *Austropuccinia*: a new genus name for the myrtle rust *Puccinia psidii* placed within the redefined family *Sphaerophragmiaceae* (*Pucciniales*). *Phytotaxa* **297**: 53–61.
- Beenken L, Wood AR (2015). *Puccorchidium* and *Sphenorchidium*, two new genera of *Pucciniales* on *Annonaceae* related to *Puccinia psidii* and the genus *Dasyospora*. *Mycological Progress* **14**: 1–13.
- Beenken L, Zoller S, Berndt R (2012). Rust fungi on *Annonaceae* II: the genus *Dasyospora* Berk. & M.A. Curtis. *Mycologia* **104**: 659–681.
- Berndt R (2018). The *Pucciniosiraceae*: taxonomy of a polyphyletic family of rust fungi (*Uredinales*). In: *Biodiversity and Ecology Fungi, Lichens and Mosses*. Österreichische Akademie der Wissenschaften, Germany: 245–269.
- Berndt R, Wood AR (2012). Additions to the rust fungi of South Africa. *Mycological Progress* **11**: 483–497.
- Blomquist CL, Scholler M, Scheck HJ (2015). Detection of rust caused by *Tranzschelia mexicana* on *Prunus salicifolia* in the United States. *Plant Disease* **99**: 1856.
- Bouckaert R, Vaughan TG, Barido-Sottani J, et al. (2019). BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLOS Computational Biology* **15**: e1006650.
- Bruns TD, Vilgalys R, Barns SM, et al. (1992). Evolutionary relationships within the fungi: Analyses of nuclear small subunit rRNA sequences. *Molecular Phylogenetics and Evolution* **1**: 231–241.
- Bubner B, Buchheit R, Friedrich F, et al. (2019). Species identification of European forest pathogens of the genus *Milesina* (*Pucciniales*) using urediniospore morphology and molecular barcoding including *M. woodwardiana* sp. nov. *MycoKeys* **48**: 1–40.



- Buriticá P (1991). Familias del orden *Uredinales* con ciclo de vida completamente reducido. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* **18**: 131–148.
- Carefoot GL, Sprott ER (1967). *Famine on the wind: plant diseases and human history*. Rand McNally & Company, USA.
- Cao B, Tao S, Tian C, et al. (2018). *Coleopuccinia* in China and its relationship to *Gymnosporangium*. *Phytotaxa* **347**: 235–242.
- Carnegie AJ, Pegg GS (2018). Lessons from the incursion of myrtle rust in Australia. *Annual Review of Phytopathology* **56**: 457–478.
- Chatasiri S, Ono Y (2008). Phylogeny and taxonomy of the Asian grapevine leaf rust fungus, *Phakopsora euvitis*, and its allies (*Uredinales*). *Mycoscience* **49**: 66–74.
- Crane PE (2000). *Systematics and Biology of the Genus Chrysomyxa*. Ph.D. thesis. University of Alberta, Canada.
- Crane PE, Hiratsuka Y, Currah RS (2000). Reproductive biology and evidence for water dispersal of teliospores in *Chrysomyxa weirii*, a microcyclic spruce needle rust. *Mycologia* **92**: 745–763.
- Cummins G (1950). The genus *Scopella* of the *Uredinales*. *Bulletin of the Torrey Botanical Club* **77**: 204–213.
- Cummins G (1978). J.C. Arthur: the man and his work. *Annual Review of Phytopathology* **16**: 19–30.
- Cummins GB, Hiratsuka Y (1983). *Illustrated genera of rust fungi*. Rev. ed. APS Press, USA.
- Cummins GB, Hiratsuka Y (2003). *Illustrated genera of rust fungi*. 3<sup>rd</sup> ed. APS Press, USA.
- Cunningham GH (1931). The rust fungi of New Zealand, together with the biology, cytology and therapeutics of the *Uredinales*. John McIndoe, NZ.
- Dietel P (1928). *Hemibasidii (Ustilaginales und Uredinales)*. A Engler, K Prantl: *Die natürlichen Pflanzenfamilien*, Germany.
- Doungsa-ard C, McTaggart AR, Geering ADW, et al. (2014). *Uromycladium falcatarium* sp. nov., the cause of gall rust of *Paraserianthes falcataria* in south-east Asia. *Australasian Plant Pathology* **44**: 25–30.
- Doungsa-ard C, McTaggart AR, Geering ADW, et al. (2018). Diversity of gall-forming rusts (*Uromycladium*, *Pucciniales*) on *Acacia* in Australia. *Persoonia* **40**: 221–238.
- Ebinghaus M, Maier W, Wingfield MJ, et al. (2018a). New host associations and a novel species for the gall-inducing acacia rust genus *Ravenelia* in South Africa. *MycKeys* **43**: 1–21.
- Ebinghaus M, Wingfield MJ, Begerow D, et al. (2018b). The genus *Ravenelia* (*Pucciniales*) in South Africa. *Mycological Progress* **19**: 259–290.
- Feau N, Vialle A, Allaire M, et al. (2011). DNA barcoding in the rust genus *Chrysomyxa* and its implications for the phylogeny of the genus. *Mycologia* **103**: 1250–1266.
- Gäumann E (1964). *Die Pilze. Grundzüge ihrer Entwicklungsgeschichte und Morphologie*. Birkhäuser, Basel.
- Guindon S, Dufayard JF, Lefort V, et al. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* **59**: 307–321.
- Hart JA (1988). Rust fungi and host plant coevolution: do primitive hosts harbor primitive parasites? *Cladistics* **4**: 339–366.
- Henk DA, Vilgalys R (2007). Molecular phylogeny suggests a single origin of insect symbiosis in the *Pucciniomycetes* with support for some relationships within genus *Septobasidium*. *American Journal of Botany* **94**: 1515–1526.
- Hernández J, Aime MC, Newbry B (2004). *Aecidium kalanchoe* sp. nov., a new rust on *Kalanchoe blossfeldiana* (*Crassulaceae*). *Mycological Research* **108**: 846–848.
- Hibbett DS, Blackwell M, James TY, et al. (2018). Phylogenetic taxon definitions for *Fungi*, *Dikarya*, *Ascomycota* and *Basidiomycota*. *IMA Fungus* **9**: 291–298.
- Hiratsuka N (1958). Revision of taxonomy of the *Pucciniastreae*. Kasai Publ. Print Co., Tokyo.
- Hiratsuka N, Kaneko S (1978). Heteroecism of the wisteria rust, *Ochropsoara kraunhiae* (Diet.) Dietel. *Proceedings of the Japan Academy, Series B* **54**: 300–303.
- Hiratsuka Y, Cummins GB (1963). Morphology of the spermogonia of the rust fungi. *Mycologia* **55**: 487–507.
- Hiratsuka Y (1983). The nuclear cycle and terminology of spore states in *Uredinales*. *Mycologia* **65**: 432–443.
- Hiratsuka Y, Hiratsuka N (1980). Morphology of spermogonia and taxonomy of rust fungi. *Reports of the Tottori Mycological Institute* **18**: 257–268.
- Hoang DT, Chernomor O, von Haeseler A, et al. (2018). UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**: 518–522.
- Huson DH, Bryant D (2005). Application of phylogenetic networks in evolutionary studies. *Molecular Biology and Evolution* **23**: 254–267.
- Ishaq A, Aime MC, Karleson-Ayaloa E, et al. (2019). First report of Asian pistachio rust (*Pileolaria pistachiae*) in Pakistan. *Canadian Journal of Plant Pathology* **42**: 210–217.
- Jackson HS (1931). Present evolution tendencies and the origin of life cycles in the *Uredinales*. *Memoirs of the Torrey Botanical Club* **18**: 1–108.
- Ji J-X, Li Z, Li Y, et al. (2019). Life cycle of *Nothoravenelia japonica* and its phylogenetic position in *Pucciniales*, with special reference to the genus *Phakopsora*. *Mycological Progress* **18**: 855–864.
- Kern FD (1960). Changing concepts of *Gymnosporangium*. *Mycologia* **52**: 837–844.
- Kirk PM, Cannon PF, Minter DW, et al. (2008). *Dictionary of the Fungi*. 10th Ed. CABI, UK.
- Leppik EE (1953). Some viewpoints on the phylogeny of rust fungi. I. Coniferous rusts. *Mycologia* **45**: 46–74.
- Leppik EE (1965). Some viewpoints on the phylogeny of rust fungi. V. Evolution of biological specialization. *Mycologia* **57**: 6–22.
- Leppik EE (1972). Evolutionary specialization of rust fungi (*Uredinales*) on the *Leguminosae*. *Annales Botanici Fennici* **9**: 135–148.
- López-Franco RM, Hennen JF (1990). The genus *Tranzschelia* (*Uredinales*) in the Americas. *Systematic Botany* **15**: 560–591.
- Maier W, Begerow D, Weiss M, et al. (2003). Phylogeny of the rust fungi: an approach using nuclear large subunit ribosomal DNA sequences. *Canadian Journal of Botany* **81**: 12–23.
- Mains EB (1934). *Angiopsora*, a new genus of rusts on grasses. *Mycologia* **26**: 122–132.
- Mains EB (1939a). Studies in the *Uredinales*, the genus *Maravalia*. *Bulletin of the Torrey Botanical Club* **66**: 173–179.
- Mains EB (1939b). *Scopella* gen. nov. of the *Pucciniaceae*. *Annales Mycologici* **37**: 57–60.
- Mains EB (1939c). The genera *Skierka* and *Ctenoderma*. *Mycologia* **31**: 175–190.
- Mains EB (1940). *Tegillum* a new genus of *Uredinales*. *Bulletin of the Torrey Botanical Club* **67**: 705–709.
- McNeill J, Barrie FR, Buck WR, et al. (2012). *International Code of Nomenclature for algae, fungi and plants (Melbourne Code) adopted by the Eighteenth International Botanical Congress Melbourne, Australia, July 2011*. Koeltz Scientific Books, Germany.
- McTaggart AR, Aime MC (2018). The species of *Coleosporium* (*Pucciniales*) on *Solidago* in North America. *Fungal Biology* **122**: 800–809.
- McTaggart AR, Doungsa-ard C, Geering ADW, et al. (2015). A co-evolutionary relationship exists between *Endoraecium* (*Pucciniales*) and its *Acacia* hosts in Australia. *Persoonia* **35**: 50–62.

- McTaggart AR, du Plessis E, Roux J, *et al.* (2020). Sexual reproduction in populations of *Austropuccinia psidii*. *European Journal of Plant Pathology* **156**: 537–545.
- McTaggart AR, Geering ADW, Shivas RG (2012). *Uredinopsis pteridis* and *Desmella aneimiae*, the first rust fungi (*Pucciniales*) reported on ferns (*Pteridophyta*) in Australia. *Australasian Plant Disease Notes* **9**: 149.
- McTaggart AR, Shivas RG, van der Nest MA, *et al.* (2016). Host jumps shaped the diversity of extant rust fungi (*Pucciniales*). *New Phytologist* **209**: 1149–1158.
- McTaggart AT, Shuey LS, Granados GM, *et al.* (2018). Evidence that *Austropuccinia psidii* may complete its sexual life cycle on *Myrtaceae*. *Plant Pathology* **67**: 729–734.
- Minh BQ, Hahn M, Lanfear R (2018). New methods to calculate concordance factors for phylogenomic datasets. *bioRxiv*: 487801.
- Minnis AM, McTaggart A, Rossman A, *et al.* (2012). Taxonomy of mayapple rust: the genus *Allodus* resurrected. *Mycologia* **104**: 942–950.
- Minnis AM, Yun HY, Rossman AY (2008). (1852) Proposal to conserve the name *Olivea tectonae* (T.S. Ramakr. & K. Ramakr.) R.L. Mulder against *Olivea tectonae* (Racib.) Thirum. (*Basidiomycota*). *Taxon* **57**: 1355–1356.
- Moore RT (1990). Order *Platygliales ord. nov.* *Mycotaxon* **39**: 245–248.
- Mordue JEM (1991). *Aecidium mori*. *C.M.I. Descriptions of Pathogenic Fungi and Bacteria* **1031**: 1–2.
- Mundkur BB (1943). Indian species of *Phakopsora* and *Bubakia*. *Mycologia* **35**: 538–545.
- Novick RS (2008). *Phylogeny, taxonomy, and life cycle evolution in the cedar rust fungi (Gymnosporangium)*. PhD Thesis, Yale University, New Haven, Connecticut, USA.
- Nguyen L-T, Schmidt HA, von Haeseler A, *et al.* (2015). IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**: 268–274.
- Ono Y (1984). A monograph of *Maravalia* (*Uredinales*). *Mycologia* **76**: 892–911.
- Ono Y (2015a). *Kuehneola* species (*Phragmidiaceae*, *Pucciniales*) on *Vitaceae* plants. *Mycological Progress* **14**: 50.
- Ono Y (2015b). *Phakopsora breyniae* (*Phakopsoraceae*, *Pucciniales*), an autoecious macrocyclic species on *Breynia cernua* from Australia. *Mycological Progress* **14**: 117.
- Ono Y, Hennen J (1983). Taxonomy of the chaconiaceous genera (*Uredinales*). *Transactions of the Mycological Society of Japan* **24**: 369–402.
- Padamsee M, McKenzie EHC (2014). A new species of rust fungus on the New Zealand endemic plant *Myosotidium* from the isolated Chatham Islands. *Phytotaxa* **174**: 223–230.
- Paradis E, Schliep K (2018). Ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* **35**: 526–528.
- Pegg GS, Giblin FR, McTaggart AR, *et al.* (2014). *Puccinia psidii* in Queensland, Australia: disease symptoms, distribution and impact. *Plant Pathology* **63**: 1005–1021.
- Persoon CH (1801). *Synopsis Methodica Fungorum*. Göttingen, Germany.
- Peterson RS (1968). Rust fungi on *Araucariaceae*. *Mycopathologia et mycologia applicata* **34**: 17–26.
- Peterson RS (1974). Rust fungi with *Caeoma*-like sori on conifers. *Mycologia* **66**: 242–255.
- Peterson RS, Oehrens E (1978) *Mikronegeria alba* (*Uredinales*). *Mycologia* **70**: 321–331.
- Qi X-H, Cai L, Zhao P (2019). *Quasipucciniastrum agrimoniae*, gen. et sp. nov. on *Agrimonia* (*Rosaceae*) from China. *Mycologia* **10**: 141–150.
- R\_Core\_Team (2014). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Austria.
- Ramakrishnan TS, Ramakrishnan K (1949). *Chaconia tectonae* Ramakrishnan T.S. & K. sp. nov. on teak. *Indian Phytopathology* **2**: 17–19.
- Rambaut A, Drummond AJ, Xie D, *et al.* (2018). Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67**: 901–904.
- Sato S, Katsuya K, Hiratsuka Y (1993). Morphology, taxonomy and nomenclature of *Tsuga-Ericaceae* rusts. *Transactions of the Mycological Society of Japan* **34**: 47–62.
- Savile DBO (1950). North American species of *Chrysomyxa*. *Canadian Journal of Botany* **28**: 318–330.
- Savile DBO (1971). Co-evolution of the rust fungi and their hosts. *The Quarterly Review of Biology* **46**: 211–218.
- Savile DBO (1989). *Raveneliaceae* revisited. *Canadian Journal of Botany* **67**: 2983–2994.
- Scholler M, Aime MC (2006). On some rust fungi (*Uredinales*) collected in an *Acacia koa*-*Metrosideros polymorpha* woodland, Mauna Loa Road, Big Island, Hawaii. *Mycoscience* **47**: 159–165.
- Scholler M, Lutz M, Aime MC (2019). Repeated formation of correlated species in *Tranzschelia* (*Pucciniales*). *Mycological Progress* **18**: 295–303.
- Sela I, Ashkenazy H, Katoh K, Pupko T (2015). GUIDANCE2: accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters. *Nucleic Acids Research* **43**: W7–W14.
- Shen YM, Chung WH, Huang TC, *et al.* (2018). Unveiling *Gymnosporangium corniforme*, *G. unicorne*, and *G. niitakayamense* sp. nov. in Taiwan. *Mycoscience* **59**: 218–228.
- Sjamsuridzal W, Nishida H, Ogawa H, *et al.* (1999). Phylogenetic positions of rust fungi parasitic on ferns: evidence from 18S rDNA sequence analysis. *Mycoscience* **40**: 21–27.
- Souza ESC, Aime MC, Elias SG, *et al.* (2018). *Crossopsorella*, a new tropical genus of rust fungi. *Phytotaxa* **375**: 189–202.
- Swann EC, Taylor JW (1995). Phylogenetic perspectives on basidiomycete systematics: evidence from the 19S rRNA gene. *Canadian Journal of Botany* **73**: 862–868.
- Sydow P, Sydow H (1915). *Monographia Uredinearum seu Specierum Omnium ad hunc usque Diem Descriptio et Adumbratio Systematica*. Fratres Borntraeger, Germany.
- Sydow H (1921). Die Verwertung der Verwandtschaftsverhältnisse und des gegenwärtigen Entwicklungsganges zur Umgrenzung der Gattungen bei den Uredineen. *Annales Mycologici* **19**: 161–175.
- Sydow H (1937). Novae fungorum species. XXV. *Annales Mycologici* **35**: 244–286.
- Thirumalachar MJ (1945). Development of spore-forms and nuclear cycle in the autoecious oopsis rust *Cystopsora oleae*. *Botanical Gazette* **107**: 74–86.
- Thirumalachar MJ, Cummins GB (1948). Status of *Allopuccinia*, *Leucotelium*, *Edythea*, and *Ypsilospora*. *Mycologia* **40**: 417–422.
- Thirumalachar MJ, Narasimhan MJ, Gopalkrishnan KS (1947). Morphology of spore forms and heteroecism of the giant bamboo rust, *Dasturella divina*. *Botanical Gazette* **108**: 371–379.
- Turland NJ, Wiersema JH, Barrie FR, *et al.* (2018). *International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code)* Regnum Vegetabile 159. Koeltz Botanical Books, Germany.
- Van der Merwe M, Ericson L, Walker J, *et al.* (2007). Evolutionary relationships among species of *Puccinia* and *Uromyces* (*Pucciniaceae*, *Uredinales*) inferred from partial protein coding gene phylogenies. *Mycological Research* **111**: 163–175.

- Vialle A, Feau N, Allaire M, *et al.* (2009). Evaluation of mitochondrial genes as DNA barcode for *Basidiomycota*. *Molecular Ecology Resources* **9**: 99–113.
- Vilgalys R, Hester M (1990). Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* **172**: 4238–4246.
- White TJ, Bruns TD, Lee S, *et al.* (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*. (Innis MA, Gelfand DH, Sninsky JJ, *et al.* eds). Academic Press Inc., USA: 315–322.
- Wingfield BD, Ericson L, Szaro T, *et al.* (2004). Phylogenetic patterns in the *Uredinales*. *Australasian Plant Pathology* **33**: 327–335.
- Wood AR, Lutz M, Bauer R, *et al.* (2014). Morphology and phylogenetics of *Stomatisora*, including *Stomatisora psychotricola* sp. nov. *Mycological Progress* **13**: 1097–1104.
- Yun HY, Minnis AM, Kim YH, *et al.* (2011). The rust genus *Frommeëlla* revisited: a later synonym of *Phragmidium* after all. *Mycologia* **103**: 1451–1463.

Supplementary Material: <http://fuse-journal.org/>

**Table S1.** Collection and accession data for additional sequences used in *Melampsorineae*, *Raveneliineae*, and *Uredinineae* analyses (Figs 2-4 & S2).

**Fig. S1.** Alluvial plot tracking generic placement at familial and subfamilial rank. Each colour represents the taxonomic hypotheses of an author. Tracks for genera with conflicting familial and subfamilial placement from multiple authors are dashed. The plot was made in R with the *ggalluvial* package.

**Fig. S2.** *Raveneliineae*. Network analysis generated with SplitsTree from 28S data. Generic types are indicated in bold, type proxies by \*.

**Table S1.** Collection and accession data for additional sequences used in *Melampsorineae*, *Raveneliineae*, and *Urediniineae* analyses (Figs 2-4 & S2).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Calyptospora goeppertiana</i>	T	TDB1497	n.d.	AF522180	NA	NA	<i>Melampsorineae</i>	Szaro & Bruns unpublished
<i>Ceropsora weirii</i> (as <i>Chrysomyxa weirii</i> )	*	916CHWP CGSG8	n.d.	FJ666465	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Ceropsora weirii</i> (as <i>Chrysomyxa weirii</i> )	*	545CHW-PC-ON31	n.d.	FJ666458	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa arctostaphyli</i>		n.d.	n.d.	AF522163	NA	NA	<i>Melampsorineae</i>	Szaro & Bruns unpublished
<i>Chrysomyxa cassandrae</i>		202CHCCHCLE3	n.d.	FJ666451	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa chiogenis</i>		224CHSGAHZM3	n.d.	FJ666452	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa empetri</i>		755CHE-EMN-RA1	n.d.	FJ666463	NA	FJ666440	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa ledi</i> (≡ <i>Melampsoropsis ledi</i> )	(T)	1307CHILLEPFIDAOM 149959	n.d.	FJ666468	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa ledicola</i>		272CH-PC-ZM9	n.d.	FJ666454	NA	FJ666431	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa nagodhii</i>		201CH-LE-LE2	n.d.	FJ666450	NA	FJ666427	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa nagodhii</i>		718CH-PC-ZM62	n.d.	FJ666461	NA	FJ666438	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa neoglandulosi</i>		510CHN LEL BC32	n.d.	GU049550	NA	NA	<i>Melampsorineae</i>	Feau et al. (2011)
<i>Chrysomyxa piperiana</i>		1214CHPP_RHM_BC 186CHRPCADJ1HT 1000	n.d.	GU049565	NA	NA	<i>Melampsorineae</i>	Feau et al. (2011)
<i>Chrysomyxa rhododendri</i>		n.d.	n.d.	FJ666447	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Chrysomyxa tsukubaense</i> (as <i>Caecoma tsukubaense</i> )		TSH-R22711(CFB 22259)	<i>Rhododendron kaempferi</i>	AB191241	NA	NA	<i>Melampsorineae</i>	Crane et al. (2005)
<i>Chrysomyxa vaccinii</i>		1356CHV_VAP_BC	n.d.	GU049562	NA	NA	<i>Melampsorineae</i>	Feau et al. (2011)
<i>Chrysomyxa woroninii</i>		464CHIPCGYU5	n.d.	GU049538	NA	NA	<i>Melampsorineae</i>	Feau et al. (2011)
<i>Coleopuccinia sinensis</i>	T	BJFC-R02364	n.d.	MF802284	NA	NA	<i>Melampsorineae</i>	Cao et al. (2018)
<i>Coleosporium cacaliae</i>		KUS F24178	<i>Syneilesis palmata</i>	JF273971	NA	NA	<i>Melampsorineae</i>	Park & Shin unpublished
<i>Coleosporium campanulae</i>		WM 1114	<i>Campanula scheuchzeri</i>	AF426244	NA	NA	<i>Melampsorineae</i>	Maier et al. (2003)
<i>Coleosporium delicatulum</i>		BPI 877848 (U347)	<i>Solidago</i> sp.	MG907221	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Coleosporium delicatulum</i>		BPI 877837 (MCA2846)	<i>Euthamia graminifolia</i>	MG907232	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Coleosporium eupatorii</i>		PUR N6727 (MCA4471)	<i>Eupatorium formosanum</i>	MG907208	MG907199	MG907256	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Coleosporium hedyotidis</i>		BPI 872227 (U896)	<i>Kadua</i> sp.	MG907222	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Coleosporium inulae</i>		BPI 871127 (U717)	<i>Inula fragilis</i>	MG907223	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Coleosporium ipomoeae</i>		PUR N6724 (MCA 4537) PUR N16579 (MCA4161)	<i>Ipomoea</i> sp. <i>Rhinanthus</i> <i>aristatus</i>	MF769677	<b>MW147070</b>	MG907272	<i>Melampsorineae</i>	Aime <i>et al.</i> (2018); this paper
<i>Coleosporium melampyri</i>				MG907224	NA	NA	<i>Melampsorineae</i>	Aime <i>et al.</i> (2018)
<i>Coleosporium montanum</i>		BPI 871769 (U627)	<i>Solidago</i> <i>canadensis</i>	MG907219	NA	MG907271	<i>Melampsorineae</i>	McTaggart & Aime (2018); Aime <i>et al.</i> (2018)
<i>Coleosporium montanum</i>		BPI 877845 (MCA3077)	<i>Solidago</i> sp.	MG907226	NA	NA	<i>Melampsorineae</i>	McTaggart & Aime (2018)
<i>Coleosporium montanum</i>		BPI 877833 (U848)	<i>Aster cordifolius</i> <i>Zanthoxylum</i> <i>ailanthoides</i>	MG907220	NA	NA	<i>Melampsorineae</i>	McTaggart & Aime (2018)
<i>Coleosporium phellodendri</i>		n.d.		AB639023	NA	NA	<i>Melampsorineae</i>	Jung <i>et al.</i> unpublished
<i>Coleosporium plectranthi</i>		CO14	<i>Perilla frutescens</i>	EF095711	NA	NA	<i>Melampsorineae</i>	Yun <i>et al.</i> unpublished
<i>Coleosporium plumeriae</i>		BPI 844177 (U50):LSU/ PUR N6729 (MCA4229):18S, CO3	<i>Plumeria rubri</i>	MG907225	MG907198	MG907255	<i>Melampsorineae</i>	Aime <i>et al.</i> (2018); McTaggart & Aime (2018)
<i>Coleosporium senecionis</i>	T	TUB RB2099	n.d.	AY512840	NA	NA	<i>Melampsorineae</i>	Begerow <i>et al.</i> unpublished
<i>Coleosporium tussilaginis</i> (≡ <i>Erannium</i> <i>tussilaginis</i> )	(T)	PUR N16713 (MCA2389)	<i>Sonchus</i> sp.	MG907228	NA	NA	<i>Melampsorineae</i>	McTaggart & Aime (2018)
<i>Coleosporium tussilaginis</i> (≡ <i>Erannium</i> <i>tussilaginis</i> )	(T)	WM 1113	<i>Tussilago farfara</i> <i>Verbesina</i> <i>gigantea</i>	AF426242	NA	NA	<i>Melampsorineae</i>	Maier <i>et al.</i> (2003)
<i>Coleosporium verbesinae</i>		BPI 864105	<i>Elephantopus</i> <i>mollis</i>	MG907229	NA	NA	<i>Melampsorineae</i>	Aime <i>et al.</i> (2018)
<i>Coleosporium vernoniae</i>		PUR N6723 (MCA4538) PUR N16565		MG907230	NA	MG907275	<i>Melampsorineae</i>	Aime <i>et al.</i> (2018)
<i>Coleosporium vernoniae</i>		(MCA3894) PUR N16561 (MCA4165)	<i>Pinus palustris</i> <i>Vincetoxicum</i> <i>hirundinaria</i>	MG907231	NA	MG907276	<i>Melampsorineae</i>	Aime <i>et al.</i> (2018)
<i>Cronartium flaccidum</i>	T			<b>MW147081</b>	<b>MW147076</b>	NA	<i>Melampsorineae</i>	this paper
<i>Cronartium harknessii</i> (≡ <i>Endocronartium harknessii</i> )	(T)	CFB22250	n.d.	AY700193	NA	NA	<i>Melampsorineae</i>	Matheny <i>et al.</i> unpublished
<i>Cronartium quercuum</i>		BPI 863626 BPI 871660 (U396):28S/ n.d.:18S	<i>Quercus alba</i> <i>Ribes</i> sp.	<b>DQ185028</b> DQ354560	NA	NA	<i>Melampsorineae</i>	this paper Aime (2006), Bruns Aime <i>et al.</i> (1992)
<i>Cronartium ribicola</i>		PUR N6734 (MCA5066)	<i>Ribes odoratum</i>	KC876675	NA	NA	<i>Melampsorineae</i>	Wilson <i>et al.</i> (2014)
<i>Diaphanopellis purpurea</i>	*	BJFC R02302	<i>Picea purpurea</i>	MK874623	NA	NA	<i>Melampsorineae</i>	Yang & Wang unpublished
<i>Goplana</i> sp.		PURN 16467 (U1399)	unidentified host	<b>MW049260</b>	NA	NA	<i>Melampsorineae</i>	this paper

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Hyalopsora nodispora</i>		BPI 893262	<i>Adiantum capillus-veneris</i>	KY798373	NA	NA	<i>Melampsorineae</i>	Demers & Castlebury unpublished
<i>Hyalopsora polypodii</i>		FO 47825:28S/HP1:18S	<i>Cystopteris fragilis/Athyrium yokoscense</i>	AF426229	AB011016	NA	<i>Melampsorineae</i>	Maier et al. (2003); Sjamsuridzal et al. 1999
<i>Hyalopsora polypodii</i>		PDD 71999	<i>Deparia petersenii</i>	KJ698627	KJ746817	NA	<i>Melampsorineae</i>	Padamsee & McKenzie (2014)
<i>Melampsora abietis-canadensis</i>		667X-TSC-SH12	n.d.	FJ666513	NA	FJ666543	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Melampsora abietis-canadensis</i>		666XTSCSH11	n.d.	FJ666512	NA	FJ666542	<i>Melampsorineae</i>	Vialle et al. (2009)
<i>Melampsora aecidioides</i>		PFH0910	<i>Populus alba</i> x <i>Populus tremula</i>	JN934983	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2013)
<i>Melampsora albertensis</i>		1344MEABPOTUSA	<i>Populus tremuloides</i>	JX416843	NA	NA	<i>Melampsorineae</i>	Vialle unpublished
<i>Melampsora amygdalinae</i>		BLACKMAULLG96	<i>Salix triandra</i>	AY444782	NA	NA	<i>Melampsorineae</i>	Pei et al. (2005)
<i>Melampsora arctica</i>		HMAS52899	<i>Salix alata</i>	KX386123	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora arctica</i>		HMAS67420	<i>Salix siuzevii</i>	KX386119	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora caprearum</i>		RFCR92-2	<i>Salix x hirtel</i>	AY444781	NA	NA	<i>Melampsorineae</i>	Pei et al. (2015)
<i>Melampsora caprearum</i>		NYSF003819	<i>Salix caprea</i>	KU550033	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora coleosporioides</i>		n.d.	<i>Salix</i> sp.	AY652950	NA	NA	<i>Melampsorineae</i>	Pei et al. (2005)
<i>Melampsora coleosporioides</i>		n.d.	<i>Salix babylonica</i>	AY652951	NA	NA	<i>Melampsorineae</i>	Pei et al. (2005)
<i>Melampsora epiphylla</i>		CJ01/2/01	<i>Salix viminalis</i>	AY444792	NA	NA	<i>Melampsorineae</i>	Pei et al. (2005)
<i>Melampsora epiphylla</i>		HMNWFCHM13	<i>Salix</i> sp.	KX386108	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora euphorbiae</i>	T	BPI 871135 (U681):28S/ BPI 863501 (U138):18S	<i>Euphorbia heterophylla</i>	DQ351722	NA	DQ789986	<i>Melampsorineae</i>	Deadman et al. (2006); Matheny et al. (2006)
<i>Melampsora euphorbiae-gerardiana</i>		BRIP 39560	<i>Euphorbia pepus</i>	<b>EF192199</b>	NA	NA	<i>Melampsorineae</i>	this paper
<i>Melampsora euphorbiae-gerardiana</i>		ECS48	n.d.	AF522172	NA	NA	<i>Melampsorineae</i>	Szaro and Bruns unpublished
<i>Melampsora ferrinii</i>		PUR N6740 (MCA3522) (HOLOTYPE)	<i>Salix babylonica</i>	NG060305	NA	NA	<i>Melampsorineae</i>	Toome & Aime (2005)
<i>Melampsora ferrinii</i>		SAG 21943	<i>Salix</i> sp.	KY053853	NA	NA	<i>Melampsorineae</i>	Zapata (2016)
<i>Melampsora helioscopiae</i>		WM 1029	<i>Euphorbia helioscopia</i>	AF426197	NA	NA	<i>Melampsorineae</i>	Maier et al. (2003)
<i>Melampsora hypericorum</i>		WM 1110	<i>Hypericum calycinum</i>	AF426196	NA	NA	<i>Melampsorineae</i>	Maier et al. (2003)
<i>Melampsora hypericorum</i>		PDD 97325	<i>Hypericum androsaemum</i>	KJ716353	NA	NA	<i>Melampsorineae</i>	Padamsee & McKenzie (2014)

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Melampsora idesiae</i>		KUSF25042	<i>Idesia polycarpa</i>	KX944284	NA	NA	<i>Melampsorineae</i>	Cho <i>et al.</i> unpublished
<i>Melampsora kupreviczii</i>		TSHR087	<i>Salix integra</i>	KX386110	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora kupreviczii</i>		TSHR6016	<i>Salix integra</i> <i>Populus adenopoda</i>	KX386109	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora laricis</i>		TSH R20045	<i>Salix viminalis</i>	AB116819	NA	NA	<i>Melampsorineae</i>	Tian <i>et al.</i> (2004)
<i>Melampsora laricis-epitea</i>		VMLG96-11	<i>Salix viminalis</i>	AY444790	NA	NA	<i>Melampsorineae</i>	Pei <i>et al.</i> (2005)
<i>Melampsora laricis-epitea</i> f.sp. <i>laricis-daphnoidis</i>		ACNI92	<i>Salix acutifolia</i>	AY444789	NA	NA	<i>Melampsorineae</i>	Pei <i>et al.</i> (2005)
<i>Melampsora laricis-miyabeana</i>		TSHR9620	<i>Salix futura</i>	KX386107	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora laricis-miyabeana</i>		TSHR960828	<i>Salix japonica</i>	KX386102	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora laricis-pentandrae</i>		PENTNWC97/093/03	<i>Salix pentandra</i>	AY444783	NA	NA	<i>Melampsorineae</i>	Pei <i>et al.</i> (2005)
<i>Melampsora lini</i>		n.d.	n.d.	L20283	NA	NA	<i>Melampsorineae</i>	Berres <i>et al.</i> unpublished
<i>Melampsora magnusiana</i>		1429MEGCRVDSD	<i>Corydalis cava</i> <i>Populus yunnanensis</i>	JN934929	NA	NA	<i>Melampsorineae</i>	Vialle <i>et al.</i> (2013)
<i>Melampsora nujiangensis</i>		1423MENPOYCHI	<i>Populus yunnanensis</i>	JN934933	NA	NA	<i>Melampsorineae</i>	Vialle <i>et al.</i> (2013)
<i>Melampsora nujiangensis</i>		1418MENPOYCHI	<i>Populus yunnanensis</i>	JN934932	NA	NA	<i>Melampsorineae</i>	Vialle <i>et al.</i> (2013)
<i>Melampsora occidentalis</i>		BPI 871537 (U395)	<i>Populus balsamifera</i> <i>Populus trichocarpa</i>	JQ042237	NA	NA	<i>Melampsorineae</i>	Busby <i>et al.</i> (2012)
<i>Melampsora occidentalis</i>		U1218	<i>Populus trichocarpa</i>	JQ042236	NA	NA	<i>Melampsorineae</i>	Busby <i>et al.</i> (2012)
<i>Melampsora pakistanica</i>		QAU BA13	<i>Euphorbia helioscopia</i> <i>Euphorbia helioscopia</i>	KU847978	NA	NA	<i>Melampsorineae</i>	Ali <i>et al.</i> (2016)
<i>Melampsora pakistanica</i>		BA13c	<i>Euphorbia helioscopia</i>	KX237556	NA	NA	<i>Melampsorineae</i>	Ali <i>et al.</i> (2016)
<i>Melampsora populnea</i>		AAH00-1	<i>Populus alba</i>	AY444786	NA	NA	<i>Melampsorineae</i>	Pei <i>et al.</i> (2005)
<i>Melampsora populnea</i> f.sp. <i>pinitorqua</i>		97MP08	n.d.	FJ666521	NA	NA	<i>Melampsorineae</i>	Vialle <i>et al.</i> (2009)
<i>Melampsora pruinosa</i>		1366MEPRPORUR	<i>Populus pruinosa</i> <i>Populus diversifolia</i>	JN934939	NA	NA	<i>Melampsorineae</i>	Vialle <i>et al.</i> (2013)
<i>Melampsora pruinosa</i>		1343MEPRPORCHI	<i>Populus diversifolia</i>	JN934938	NA	NA	<i>Melampsorineae</i>	Vialle <i>et al.</i> (2013)
<i>Melampsora ribesii-purpureae</i>		PURP897-1	<i>Salix purpurea</i>	AY444791	NA	NA	<i>Melampsorineae</i>	Pei <i>et al.</i> (2015)
<i>Melampsora ribesii-purpureae</i>		TSHR7549	<i>Salix purpurea</i>	KX386095	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora ribesii-viminalis</i>		HNMAP1698	<i>Salix viminalis</i>	KX386096	NA	NA	<i>Melampsorineae</i>	Zhao unpublished
<i>Melampsora ricini</i>		PDD 98363	<i>Ricinus communis</i>	KJ716352	NA	NA	<i>Melampsorineae</i>	Padamsee & McKenzie (2014)

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Melampsora rostrupii</i>		PFH083	<i>Populus alba</i>	JN934981	NA	NA	<i>Melampsorineae</i>	Vialle et al. (2013)
<i>Melampsora salicis-albae</i>		ALBANWC96/10	<i>Salix alba</i>	AY444788	NA	NA	<i>Melampsorineae</i>	Pei et al. (2005)
<i>Melampsorella caryophyllacearum</i>	T	WM 1092	<i>Abies alba</i>	AF426232	NA	NA	<i>Melampsorineae</i>	Maier et al. (2003)
<i>Melamporidium alni</i>		Pentti Alanko 31971	<i>Alnus mandshurica</i>	KF031534	NA	NA	<i>Melampsorineae</i>	McKenzie et al. (2013)
<i>Melamporidium betulinum</i>	T	BPI 871107 (MCA2884)	<i>Alnus</i> sp.	DQ354561	<b>MW147071</b>	NA	<i>Melampsorineae</i>	Aime (2006); this paper
<i>Melamporidium hiratsukanum</i>		PDR #1480181	<i>Alnus rhombifolia</i>	KC313888	NA	NA	<i>Melampsorineae</i>	Blomquist et al. (2014)
<i>Melamporidium hiratsukanum</i>		H 6035362	<i>Alnus incana</i>	KF031536	NA	NA	<i>Melampsorineae</i>	McKenzie et al. (2013)
<i>Milesia exigua</i>		KRM0050247	<i>Polystichum braunii</i>	MK302211	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Milesia murariae</i>		KRM0048132	<i>Asplenium ruta-muraria</i>	MK302196	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Milesia murariae</i>		KRM0048134	<i>Asplenium ruta-muraria</i>	MK302195	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Milesia scolopendrii</i>		HerB 4574	<i>Asplenium scolopendrium</i>	AF426236	NA	NA	<i>Melampsorineae</i>	McKenzie et al. (2013)
<i>Milesia scolopendrii</i>		KRM0049051	<i>Asplenium scolopendrium</i>	MK302209	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Milesia scolopendrii</i>		KRM0043186	<i>Asplenium scolopendrium</i>	MK302198	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Milesia</i> sp.		MCA3088	<i>Abies</i> sp.	MG907234	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Milesia vogesiaca</i>		KRM0043187	<i>Polystichum aculeatum</i>	MK302202	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Milesia vogesiaca</i>		PURN659a (PUR044)	<i>Polystichum aculeatum</i>	MG907235	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Milesina whitei</i>		KRM0049177	<i>Polystichum setiferum</i>	MK302212	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Naohidemycetes vaccinii</i>	T	BPI 871754 (MCA2780)	<i>Vaccinium ovatum</i>	DQ354563	DQ354562	NA	<i>Melampsorineae</i>	Aime (2006)
<i>Pucciniastrum actinidiae</i>		TSHR4268 IBA8002	<i>Actinidia rufa</i>	AB221407	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum boehmeriae</i>		TSH R21289	<i>Boehmeria tricuspis</i>	AB221393	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum circaeae</i>		RB 2098	<i>Circaea lutetiana</i>	AF426227	NA	NA	<i>Melampsorineae</i>	Maier et al. (2003)
<i>Pucciniastrum circaeae</i>		PUR N22374 (MCA2948)	<i>Circaea alpina</i>	MG907238	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Pucciniastrum corni</i>		TSHR13510	<i>Cornus kuosa</i>	AB221409	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)



Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Pucciniastrum coryli</i>		TSHR4233 IBA2582 PUR N11088 (MCA5308)	<i>Corylus sieboldiana</i>	AB221379	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum epilobii</i>	T		<i>Epilobium angustifolium</i>	<b>MW147083</b>	NA	NA	<i>Melampsorineae</i>	this paper
<i>Pucciniastrum fagi</i>		TSHR10724	<i>Fagus crenata</i>	AB221378	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006) Ishaq et al.
<i>Pucciniastrum guttatum</i>		PUR N11943	<i>Galium</i> sp.	KX254357	NA	NA	<i>Melampsorineae</i>	unpublished
<i>Pucciniastrum guttatum</i>		WM 1203	<i>Galium odoratum</i>	AF426231	NA	NA	<i>Melampsorineae</i>	Maier et al. (2003)
<i>Pucciniastrum hikosanense</i>		TSHR4288 IBA2569	<i>Acer rufinerve</i>	AB221390	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum hydrangeae</i>		MCA2837	<i>Hydrangea arborescens</i>	MG907240	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Pucciniastrum hydrangeae-petiolearis</i>		TSHR4264 IBA7881	<i>Hydrangea petiolearis</i>	AB221385	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum kusanoi</i>		HiroU21509	<i>Clethra barbinervis</i>	AB221402	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum minimum</i>		MCA2988	<i>Vaccinium angustifolium</i>	MG907243	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Pucciniastrum miyabeianum</i>		TSHR10202	<i>Viburnum furcatum</i>	AB221397	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006) Yang et al.
<i>Pucciniastrum nipponicum</i>		HMBFGS53	n.d.	KC416001	NA	NA	<i>Melampsorineae</i>	unpublished
<i>Pucciniastrum rubiae</i>		HMBFQH1	n.d.	KC416007	NA	NA	<i>Melampsorineae</i>	Yang et al.
<i>Pucciniastrum rubiae</i>		HMBFXZ1	n.d.	KC416010	NA	NA	<i>Melampsorineae</i>	Yang et al.
<i>Pucciniastrum styracinum</i>		TSHR1583	<i>Styrax japonica</i>	AB221418	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum symphyti</i>		HerB 4732	<i>Symphytum officinale</i>	AF426230	NA	NA	<i>Melampsorineae</i>	Maier et al. (2003)
<i>Pucciniastrum tiliae</i>		TSHR4295 IBA7878	<i>Tilia japonica</i>	AB221415	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Pucciniastrum yoshinagai</i>		TSHR4272 IBA8430	<i>Stewartia monadelphae</i>	AB221411	NA	NA	<i>Melampsorineae</i>	Liang et al. (2006)
<i>Rossmannomyces monesis</i> (as <i>Chrysomyxa monesis</i> )		1360CHM PCS BC	n.d.	GU049567	NA	NA	<i>Melampsorineae</i>	Feau et al. (2011)
<i>Rossmannomyces ramischiae</i> (as <i>Chrysomyxa ramischiae</i> )		1318CHH RMS FI	n.d.	GU049551	NA	NA	<i>Melampsorineae</i>	Feau et al. (2011)
<i>Thekopsora agrimoniae</i> (as <i>Pucciniastrum agrimoniae</i> )		BPI 878084 (MCA2826)	<i>Agrimonia parviflora</i>	MG907236	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Thekopsora americana</i> (as <i>Pucciniastrum americanum</i> )		BPI 871792 (U626)	<i>Rubus strigosus</i>	MG907237	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Thekopsora americana</i> (as <i>Pucciniastrum americanum</i> )		PUR N15345 (U1633)	<i>Rubus</i> sp.	<b>MW147084</b>	NA	NA	<i>Melampsorineae</i>	this paper

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Thekopsora potentillae</i>		BPI 878097 (U865)	<i>Potentilla</i> sp.	MG907241	<b>MW049308</b>	NA	<i>Melampsorineae</i>	Aime et al. (2018); this paper
<i>Thekopsora pseudoagrimoniae</i> (= <i>Quasipucciniastrum agrimoniae</i> )	(T)	HMAS248095	<i>Agrimonia pilosa</i> <i>Phegopteris</i> <i>connectilis</i>	MK208281	NA	NA	<i>Melampsorineae</i>	Qi et al. (2019)
<i>Uredinopsis filicina</i>	T	KRM0050313	<i>Osmunda</i>	MK302215	NA	NA	<i>Melampsorineae</i>	Bubner et al. (2019)
<i>Uredinopsis osmundae</i>		BPI 872258 (U856)	<i>claytoniana</i>	MG907245	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Uredinopsis osmundae</i>		PUR N16024 (U1188)	<i>Athyrium filix-</i> <i>femina</i>	MG907244	NA	NA	<i>Melampsorineae</i>	Aime et al. (2018)
<i>Uredinopsis pteridis</i>		BRIP 60091	<i>Pteridium</i> <i>esculentum</i>	KM249869	NA	NA	<i>Melampsorineae</i>	McTaggart et al. (2014)
<i>Cephalotelium evansii</i> (as <i>Ravenelia</i> <i>evansii</i> )		PREM 61846	<i>Vachellia</i> <i>luederitzii</i> var. <i>retinens</i>	MG945993	NA	NA	<i>Ravenellieae</i>	Ebinghaus et al. (2018)
<i>Cephalotelium evansii</i> (as <i>Ravenelia</i> <i>evansii</i> )		PREM 61028	<i>Vachellia swazica</i>	MG945996	NA	NA	<i>Ravenellieae</i>	Ebinghaus et al. (2018)
<i>Cephalotelium macowaniana</i> (as <i>Ravenelia macowaniana</i> )	T	PREM 61210	<i>Vachellia karroo</i>	MG946004	NA	NA	<i>Ravenellieae</i>	Ebinghaus et al. (2018)
<i>Cephalotelium neocaledoniense</i> (as <i>Ravenelia neocaledoniensis</i> )		BRIP 56907	<i>Vachellia</i> <i>farnesiana</i>	KJ862347	NA	NA	<i>Ravenellieae</i>	McTaggart et al. (2015)
<i>Cephalotelium xanthophloeae</i> (as <i>Ravenelia xanthophloeae</i> )		PREM 61215	<i>Vachellia</i> <i>xanthophloea</i>	MG946017	NA	NA	<i>Ravenellieae</i>	Ebinghaus et al. (2018)
<i>Crossoporella byrsonimae</i>	T	UB22384	<i>Byrsonima crassa</i>	MG250386	NA	NA	<i>Ravenellieae</i>	Souza et al. (2018)
<i>Diorchidium woodii</i>	T	U1475	<i>Milletia grandis</i>	<b>MW111533</b>	<b>MW111538</b>	NA	<i>Ravenellieae</i>	this paper
<i>Endoraecium falciforme</i>		BRIP 57583	<i>Acacia falciformis</i>	KJ862306	KJ862405	KJ862439	<i>Ravenellieae</i>	McTaggart et al. (2015)
<i>Endoraecium parvum</i>		BRIP 53616	<i>Acacia leiocalyx</i>	KJ862316	KJ862409	KJ862445	<i>Ravenellieae</i>	McTaggart et al. (2015)
<i>Endoraecium phyllodiorum</i>		BRIP 57516	<i>Acacia</i> <i>aulacocarpa</i>	KJ862324	KJ862411	KJ862447	<i>Ravenellieae</i>	McTaggart et al. (2015)
<i>Endoraecium podalyriifoliae</i>		BRIP 57576	<i>Acacia</i> <i>podalyriifolia</i>	KJ862334	KJ862414	KJ862449	<i>Ravenellieae</i>	McTaggart et al. (2015)
<i>Endoraecium tierneyi</i>		BRIP 27071	<i>Acacia</i> <i>harpophylla</i>	KJ862335	KJ862415	KJ862450	<i>Ravenellieae</i>	McTaggart et al. (2015)
<i>Endoraecium violae-faustiae</i>		BRIP 56547	<i>Acacia</i> <i>aulacocarpa</i>	KJ862338	KJ862418	KJ862453	<i>Ravenellieae</i>	McTaggart et al. (2015)
<i>Masseëlla capparidis</i>	T	BRIP 26745	<i>Flueggea virosa</i>	<b>MW147027</b>	NA	NA	<i>Ravenellieae</i>	this paper
<i>Nothoravenelia japonica</i>	T	HMJAU8600	n.d.	MK296511	NA	NA	<i>Ravenellieae</i>	Ji unpublished
<i>Phakopsora annonae-sylvaticae</i>		PUR 87311	<i>Annona sylvatica</i>	KF528008	KF528038	KF528046	<i>Ravenellieae</i>	Beenken (2014)

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Phakopsora arthuriana</i>		BRIP 59345	<i>Jatropha gossypifolia</i>	<b>MW147028</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Phakopsora cherimoliae</i>		PUR 89695	<i>Annona cherimola</i> <i>x squamosa</i>	KF528012	NA	NA	<i>Ravenelliaceae</i>	Beenken (2014)
<i>Phakopsora meibomiaae</i> (≡ <i>Malupa meibomiaae</i> )	(T)	R188	<i>Aeschynomene</i> sp.	EU851164	NA	NA	<i>Ravenelliaceae</i>	Zulaga <i>et al.</i> unpublished
<i>Phakopsora myrtacearum</i>		PREM 61156	<i>Eucalyptus grandis</i>	KP729473	NA	KT199414	<i>Ravenelliaceae</i>	Maier <i>et al.</i> (2003)
<i>Phakopsora phyllanthi</i>		ZT RB 8581	<i>Phyllanthus acidus</i>	KF528025	NA	NA	<i>Ravenelliaceae</i>	Beenken (2014)
<i>Phakopsora pipturi</i>		BRIP 56846	<i>Pipturus argenteus</i>	<b>MW147029</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Phakopsora pistila</i>		ZT Myc 48992	<i>Annona sericea</i>	KF528026	KF528043	KF528051	<i>Ravenelliaceae</i>	Beenken (2014)
<i>Phakopsora rollinae</i>		ZT Myc 49000	<i>Annona exsucca</i>	KF528034	KF528045	KF528054	<i>Ravenelliaceae</i>	Beenken (2014)
<i>Phakopsora tecta</i>		BRIP 56943	<i>Commelina</i> sp.	KP729476	NA	NA	<i>Ravenelliaceae</i>	Maier <i>et al.</i> (2015)
<i>Prosopidium cydistae</i>		BPI 871910	<i>Cyclista heterophylla</i>	<b>MW147030</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Prosopidium transformans</i>		PUR N11105 (U1506)	<i>Tecoma stans</i>	<b>MW147031</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Ravenelia arizonica</i>		BPI 878098 (U754)	<i>Prosopis</i> sp.	<b>MW147032</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Ravenelia berkeleyi</i>		BRIP 56904	<i>Cassia</i> sp.	KJ862349	NA	KJ862461	<i>Ravenelliaceae</i>	McTaggart <i>et al.</i> (2015)
<i>Ravenelia cohniana</i>		BPI 841187 (U85)	<i>Acacia praecox</i>	<b>MW147033</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Ravenelia cohniana</i>		BPI 841185 (U86)	<i>Senegalia praecox</i>	MG954487	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus <i>et al.</i> (2018)
<i>Ravenelia echinata</i> var. <i>ectypa</i>		BPI 841034	<i>Calliandra formosa</i>	DQ323925	NA	NA	<i>Ravenelliaceae</i>	Scholler & Aime (2006)
<i>Ravenelia escharoides</i>		WM3557	<i>Senegalia burkei</i>	MG954479	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus <i>et al.</i> (2018)
<i>Ravenelia escharoides</i>		WM3405	<i>Senegalia burkei</i>	MG954480	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus <i>et al.</i> (2018)
<i>Ravenelia halsei</i>		PREM 61855	<i>Senegalia ataxacantha</i>	MG954484	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus <i>et al.</i> (2018)
<i>Ravenelia havanensis</i>		ZT RB5788 (U19)	<i>Enterolobium contortisiliquum</i>	DQ354557	DQ354556	NA	<i>Ravenelliaceae</i>	Aime (2006)
<i>Ravenelia lonchocarpi</i>		BPI 863495 (U346)	<i>Lonchocarpus lilloi</i>	<b>MW147034</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Ravenelia macrocarpa</i>		BPI 841195 (U88)	<i>Senna subulata</i>	DQ323926	NA	NA	<i>Ravenelliaceae</i>	Scholler & Aime (2006)
<i>Ravenelia ornata</i>		BPI 910280	<i>Abrus</i> sp.	KY764175	NA	NA	<i>Ravenelliaceae</i>	Demers <i>et al.</i> unpublished

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Ravenelia pienaarii</i>		PREM 61892	<i>Senegalia caffra</i>	MG954482	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus et al. (2018)
<i>Ravenelia pienaarii</i>		WM3592	<i>Senegalia caffra</i>	MG954483	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus et al. (2018)
<i>Ravenelia piepenbringia</i>		MP5157	<i>Senegalia hayesii</i>	MG954489	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus et al. (2018)
<i>Ravenelia sessilis</i>		BRIP 59688	<i>Albizia procera</i>	<b>MW147035</b>	NA	NA	<i>Ravenelliaceae</i>	this paper Demers & Castlebury unpublished
<i>Ravenelia</i> sp.		BPI 893295	<i>Senna surattensis</i>	KY798359	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus et al. (2018)
<i>Ravenelia transvaalensis</i>		PREM 61858	<i>Senegalia mellifera</i>	MG954485	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus et al. (2018)
<i>Ravenelia transvaalensis</i>		PREM 61893	<i>Senegalia mellifera</i>	MG954486	NA	NA	<i>Ravenelliaceae</i>	Ebinghaus et al. (2018)
<i>Sorataea arayatensis</i>		BRIP 57452	<i>Derris elliptica</i>	<b>MW147036</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Sorataea</i> sp.		PUR N1652	<i>Baphia</i> sp.	<b>MW147037</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Tranzschelia thalictri</i>	*	PUR N13845 (U654)	<i>Thalictrum aquilegifolium</i>	<b>MW147038</b>	NA	NA	<i>Ravenelliaceae</i>	this paper
<i>Uromycladium fusisporum</i>		BRIP 57526	<i>Acacia salicina</i>	KJ632991	KJ633031	KJ639075	<i>Ravenelliaceae</i>	Doungsa-ard et al. (2014)
<i>Uromycladium morrisii</i>		BRIP 56962	<i>Acacia saligna</i>	KJ632985	NA	NA	<i>Ravenelliaceae</i>	Doungsa-ard et al. (2014)
<i>Uromycladium notabile</i>		BRIP 59234	<i>Acacia dealbata</i>	KJ632992	KJ633030	KJ639076	<i>Ravenelliaceae</i>	Doungsa-ard et al. (2014)
<i>Uromycladium robinsonii</i>		BRIP 57538	<i>Acacia implexa</i>	KJ632989	KJ633032	KJ639077	<i>Ravenelliaceae</i>	Doungsa-ard et al. (2014)
<i>Allodus proatii</i>		PUR N16267	<i>Tulipa</i> sp.	MT622756	<b>MW147072</b>	NA	<i>Urediniineae</i>	Ullah et al. (2020); this paper
<i>Angiopsora apoda</i>	*	PDD 72076	<i>Cenchrus clandestinus</i>	MG461668	NA	NA	<i>Urediniineae</i>	Demers et al. unpublished
<i>Angiopsora chusqueae</i>		R161	<i>Chusquea</i> sp.	EU851156	NA	NA	<i>Urediniineae</i>	Zulaga et al. unpublished
<i>Crossopora antidesmae-dioicae</i>		BRIP 56870	<i>Antidesma ghaesembilla</i>	<b>MW147039</b>	NA	<b>MW139652</b>	<i>Urediniineae</i>	this paper
<i>Dasyscypha amazonica</i>		BPI 0116382	<i>Xylopiamazonica</i>	JF263460	JF263496	JF263512	<i>Urediniineae</i>	Beenken et al. (2012)
<i>Dasyscypha echinata</i>		BPI 746651	<i>Xylopiemarginata</i>	JF263462	JF263497	JF263513	<i>Urediniineae</i>	Beenken et al. (2012)
<i>Dasyscypha guianensis</i>		ZTMyc3413	<i>Xylopiabenthamii</i>	JF263479	JF263503	JF263519	<i>Urediniineae</i>	Beenken et al. (2012)
<i>Dasyscypha mesoamericana</i>		PUR 42390	<i>Xylopiافرutescens</i> var. <i>frutescens</i>	JF263480	JF263504	JF263520	<i>Urediniineae</i>	Beenken et al. (2012)

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Dasyscypha nitida</i>		ZTMy3409	<i>Xylopi nitida</i>	JF263484	JF263505	JF263521	<i>Urediniineae</i>	Beenken et al. (2012)
<i>Dasyscypha segregaria</i>		PMA MP4941	<i>Xylopi aromatica</i>	JF263488	JF263507	JF263523	<i>Urediniineae</i>	Beenken et al. (2012)
<i>Dasyscypha winteri</i>		SF30078	<i>Xylopi sericea</i>	JF263492	JF263508	JF263524	<i>Urediniineae</i>	Beenken et al. (2012)
<i>Dietelia mesoamericana</i>		BPI 871533 (U321)	<i>Mikania micrantha</i>	<b>MW147040</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Endophyllum pampeanum</i>		BPI 843679 (U108)	<i>Salpichroa origanifolia</i>	<b>MW147041</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Endophyllum stachytarphetae</i>		BPI 844169 (GUY55)	<i>Stachytarpheta cayennensis</i>	<b>MW147042</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Gymnosporangium asiaticum</i>		TNMF0027942	<i>Pyrus pyrifolia</i>	KP308393	KP308393	NA	<i>Urediniineae</i>	Shen et al. (2018)
<i>Gymnosporangium betheli</i>		RSP01237	n.d.	KJ720163	KJ720163	NA	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnosporangium confusum</i>		BPI 880005:28S/ DAOM220748:18S	<i>Crataegus monogyna</i>	HM114219	KJ720165	NA	<i>Urediniineae</i>	Dervis et al. unpublished; Novick et al.
<i>Gymnosporangium cupressi</i>		RSP9998	n.d.	KJ720169	KJ720169	NA	<i>Urediniineae</i>	Novick et al. unpublished
<i>Gymnosporangium exiguum</i>		RSP0486 PUR N16632 (MCA3585)	n.d.	KJ720170	KJ720170	NA	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnosporangium juniperi-virginianae</i>		RSP0537	<i>Juniperus</i> sp.	MG907217	MG917687	MG907268	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnosporangium kernianum</i>		RSP0541	n.d.	KJ720177	KJ720177	NA	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnosporangium multiporum</i>		RSP9993	n.d.	KJ720179	KJ720179	NA	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnosporangium nelsonii</i>		RN91	n.d.	KJ720180	KJ720180	NA	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnosporangium trachysorum</i>		RSP0587	n.d.	KJ720184	KJ720184	NA	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnotelium vaqueliniae</i>		PUR 63656	n.d.	NA	KJ720159	NA	<i>Urediniineae</i>	Novik et al. unpublished
<i>Gymnotelium nootkatense</i> (as <i>Gymnosporangium nootkatense</i> )	T	BPI 871105 (MCA 2887)	<i>Poaceae</i>	DQ354554	MG907205	MG907266	<i>Urediniineae</i>	Novik et al. unpublished
<i>Kweilingia divina</i> (= <i>Dasturella divina</i> )	(T)	BRIP 57522	<i>Malva parviflora</i>	KU296888	NA	KX999924	<i>Urediniineae</i>	Novik et al. unpublished
<i>Leptopuccinia malvicearum</i> (as <i>Puccinia malvicearum</i> )	T	LM2016	<i>Saccharum</i> hybrid cultivar N41	JX036026	NA	NA	<i>Urediniineae</i>	Aime (2006); Aime et al. (2018) McTaggart et al. (2016)
<i>Macruronyxis fulva</i>							<i>Urediniineae</i>	Martin et al. (2017)

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Neophysopella meliosmae</i>		IBAR 9906	<i>Meliosma myriantha</i>	KC815644	NA	NA	<i>Urediniineae</i>	Pota <i>et al.</i> unpublished
<i>Neophysopella orientalis</i>		IBAR 9942	<i>Meliosma tenuis</i>	KC815656	NA	NA	<i>Urediniineae</i>	Pota <i>et al.</i> unpublished
<i>Neophysopella vitis</i>		IBAR 10117	<i>Meliosma myriantha</i>	KC815634	NA	NA	<i>Urediniineae</i>	Pota <i>et al.</i> unpublished
<i>Neopuccinia bursa</i>		RB:757071 (HOLOTYPE)	<i>Protium heptaphyllum</i>	MH047186	NA	NA	<i>Urediniineae</i>	Martins <i>et al.</i> unpublished
<i>Nyssopsora thwaitesii</i>		AMH 9528	<i>Schefflera wallichiana</i>	KF550283	NA	NA	<i>Urediniineae</i>	Baiswar <i>et al.</i> (2014)
<i>Phragmidium barnardii</i>		BRIP 56945	<i>Rubus</i> sp.	KT199402	NA	KT199415	<i>Urediniineae</i>	McTaggart <i>et al.</i> (2016)
<i>Phragmidium biloculare</i>		BPI 881121	<i>Potentilla flabellifolia</i>	JF907670	NA	NA	<i>Urediniineae</i>	Yun <i>et al.</i> (2011) Xu <i>et al.</i>
<i>Phragmidium fructigenum</i>		HMUT100472	<i>Rosa glomerata</i>	KU059168	NA	NA	<i>Urediniineae</i>	unpublished
<i>Phragmidium mexicanum</i>		BRIP 55383	<i>Potentilla indica</i>	<b>MW147043</b>	NA	<b>MW139653</b>	<i>Urediniineae</i>	this paper
<i>Phragmidium nambuanum</i>		HMUT100461	<i>Rubus innominatus</i>	KU059164	NA	NA	<i>Urediniineae</i>	Xu <i>et al.</i> unpublished
<i>Phragmidium potentillae</i>		BRIP 60089	<i>Acaena novae-zelandiae</i>	KT199403	NA	KT199416	<i>Urediniineae</i>	McTaggart <i>et al.</i> (2016)
<i>Phragmidium rosae-multiflorae</i>		HMAS14338	<i>Rosa multiflora</i>	KU059176	NA	NA	<i>Urediniineae</i>	Xu <i>et al.</i> unpublished
<i>Phragmidium rubi-idaei</i>		BRIP 59372	<i>Rubus idaeus</i>	<b>MW147044</b>	NA	<b>MW139654</b>	<i>Urediniineae</i>	this paper
<i>Phragmidium sanguisorbae</i>		BPI 872232	<i>Sanguisorba minor</i>	JF907674	NA	NA	<i>Urediniineae</i>	Yun <i>et al.</i> (2011)
<i>Phragmidium tuberculatum</i>		BPI 877981 (U486)	<i>Rosa</i> sp.	JF907675	NA	MG907280	<i>Urediniineae</i>	Yun <i>et al.</i> (2011); Aime <i>et al.</i> (2018)
<i>Pileolaria toxicodendri</i>		BPI 871761 (U607)	<i>Toxicodendron</i> sp.	DQ323924	NA	NA	<i>Urediniineae</i>	Scholler & Aime (2006)
<i>Puccinia arthrocnemi</i>		BRIP 57772	<i>Tecticornia</i> sp.	KX999865	<b>MW147073</b>	KX999908	<i>Urediniineae</i>	Marin-Felix <i>et al.</i> (2017)
<i>Puccinia arundinariae</i>		BPI 843964 (MCA2494)	<i>Arundinaria</i> sp.	DQ415277	NA	NA	<i>Urediniineae</i>	Aime (2006); this paper
<i>Puccinia bassiae</i>		BRIP 57788	<i>Sclerolaena eurotioides</i>	KX999867	NA	KX999910	<i>Urediniineae</i>	Marin-Felix <i>et al.</i> (2017)
<i>Puccinia boroniae</i>		BRIP 57810	<i>Boronia ramosa</i>	<b>MW147045</b>	<b>MW147074</b>	<b>MW139655</b>	<i>Urediniineae</i>	this paper
<i>Puccinia canaliculata</i>		BRIP 57789	<i>Cyperus</i> sp.	<b>MW147046</b>	NA	<b>MW139656</b>	<i>Urediniineae</i>	this paper
<i>Puccinia carissae</i>		BRIP 53242	<i>Carissa ovata</i>	KX999871	NA	NA	<i>Urediniineae</i>	Marin-Felix <i>et al.</i> (2017)
<i>Puccinia coronata</i> var. <i>avenae</i>		BRIP 57635	<i>Avena sativa</i>	<b>MW147047</b>	NA	<b>MW139657</b>	<i>Urediniineae</i>	this paper

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Puccinia cynodontis</i>		BRIP 57556	<i>Cynodon dactylon</i>	KX999873	NA	KX999913	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia dichondrae</i>		BRIP 60027	<i>Dichondra repens</i>	KX999874	NA	KX999914	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia engleriana</i>		BRIP 59644	<i>Tabernaemontana orientalis</i>	<b>MW147048</b>	<b>MW147075</b>	<b>MW139658</b>	<i>Uredinineae</i>	this paper
<i>Puccinia geitonoplesii</i>		BRIP 55679	<i>Geitonoplesium cymosum</i>	KM249860	NA	KX999916	<i>Uredinineae</i>	McTaggart et al. (2016); Marin-Felix et al. (2017)
<i>Puccinia gnaphallicola</i>		BRIP 58451	<i>Gamochaeta</i> sp.	KF690703	NA	KX999917	<i>Uredinineae</i>	McTaggart et al. (2014)
<i>Puccinia grevilleae</i>		BRIP 55600	<i>Grevillea mimosoides</i>	KX999878	NA	KX999918	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia helianthi</i>		BRIP 44673	<i>Helianthus annuus</i>	<b>MW147049</b>	NA	<b>MW139659</b>	<i>Uredinineae</i>	this paper
<i>Puccinia hemerocallidis</i>		BRIP 53476	<i>Hemerocallis</i> sp.	KM249855	NA	KX999920	<i>Uredinineae</i>	McTaggart et al. (2016)
<i>Puccinia heterospora</i>		BRIP 60937	<i>Abutilon</i> sp.	KU296886	NA	NA	<i>Uredinineae</i>	McTaggart et al. (2016)
<i>Puccinia iridis</i>		BRIP 56852	<i>Iris</i> sp.	KM249853	NA	NA	<i>Uredinineae</i>	McTaggart et al. (2016)
<i>Puccinia kuehnii</i>		BRIP 59200	<i>Saccharum officinarum</i>	KU296887	NA	KX999921	<i>Uredinineae</i>	McTaggart et al. (2016)
<i>Puccinia lagenophorae</i>		BRIP 57563	<i>Emilia sonchifolia</i>	KF690696	KT199388	KT199417	<i>Uredinineae</i>	McTaggart et al. (2014)
<i>Puccinia levis</i> var. <i>tricholaenae</i>		BRIP 56867	<i>Melinis repens</i>	KX999880	NA	NA	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia loranthicola</i>		BRIP 59685	<i>Amylotheca dictyophleba</i>	KX999882	NA	KX999923	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia melanocephala</i>		BRIP 59199	<i>Saccharum officinarum</i>	KU296889	NA	<b>MW139660</b>	<i>Uredinineae</i>	McTaggart et al. (2016)
<i>Puccinia menthae</i>		BRIP 59667	<i>Mentha spicata</i>	KU296890	NA	<b>MW139661</b>	<i>Uredinineae</i>	McTaggart et al. (2016)
<i>Puccinia merrilliana</i>		BRIP 56913	<i>Operculina aequisejala</i>	KX999885	NA	KX999926	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia myrsiphyllii</i>		BRIP 57782	<i>Asparagus asparagoides</i>	KM249854	NA	KT199418	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia nakanishikii</i>		BPI 879283	<i>Cymbopogon citratus</i>	GU058002	NA	NA	<i>Uredinineae</i>	Dixon et al. (2010)
<i>Puccinia paullula</i>		BRIP 60018	<i>Monstera deliciosa</i>	KX999886	NA	KX999927	<i>Uredinineae</i>	Marin-Felix et al. (2017)
<i>Puccinia pelargonii-zonalis</i>		BRIP 57414	<i>Pelargonium zonale</i>	KX999887	NA	NA	<i>Uredinineae</i>	Marin-Felix et al. (2017)

Table S1. (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Puccinia pritzeliana</i>		BRIP 57798 LSU-M121657 (MCA3793)	<i>Tremandra stelligera</i>	KX999888	NA	KX999928	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Puccinia sparganioidis</i>			<i>Spartina alterniflora</i>	GU327649	NA	NA	<i>Urediniineae</i>	Kauer et al. (2010) Marin-Felix et al. (2017)
<i>Puccinia styliidii</i>		BRIP 60107	<i>Styloidium armeria</i>	KJ622215	KT199389	KT199420	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Puccinia tetragoniae</i>		BRIP 59703	<i>Tetragonia implexicornia</i>	KX999894	NA	KX999931	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Puccinia unica</i>		BRIP 56930	<i>Aristida</i> sp.	KX999895	NA	KX999932	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Puccinia ursinae</i>		BRIP 57993	<i>Ursinia anthemoides</i>	KF690705	KT199390	KT199421	<i>Urediniineae</i>	McTaggart et al. (2016)
<i>Puccinia xanthii</i>		BRIP 56946	<i>Xanthium</i> sp.	KX999896	NA	NA	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Pucciniosira cornuta</i>		PUR N3902	<i>Mahonia leschenaultii</i>	<b>MW147050</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Pucciniosira tuberculata</i>	*	PUR N16031 (U748)	<i>Malvaceae</i>	<b>MW147051</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Schroeteriaaster elettariae</i>		BRIP 56855	<i>Alpinia caerulea</i>	<b>MW147052</b>	NA	<b>MW139662</b>	<i>Urediniineae</i>	this paper
<i>Sphaerophragmium longicorne</i>		PUR N16513	<i>Dalbergia hostilis</i>	<b>MW147053</b>	<b>MW147077</b>	NA	<i>Urediniineae</i>	this paper
<i>Sphenorhynchium</i> sp.		MCA7071	<i>Xylopia</i> sp.	<b>MW147054</b>	<b>MW147078</b>	NA	<i>Urediniineae</i>	this paper
<i>Sphenospora kevorikianii</i>		BPI 863558	<i>Stanhopea candida</i>	DQ354521	DQ354520	NA	<i>Urediniineae</i>	Aime (2006)
<i>Sphenospora saphena</i>		PUR 89514	<i>Orchidaceae</i>	<b>MW172290</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Stomatisora psychotricicola</i>	*	PREM 60886	<i>Psychotria capensis</i>	NG_059953	NA	NA	<i>Urediniineae</i>	Wood et al. (2014)
<i>Uredo ficicola</i>		BPI 863622	<i>Ficus goldmanii</i>	<b>MW147055</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Uredo moricola</i>		BRIP 57999	<i>Morus</i> sp.	<b>MH047206</b>	<b>MH047211</b>	<b>MH047203</b>	<i>Urediniineae</i>	this paper
<i>Uredo musae</i>		BRIP 59462	<i>Musa</i> sp.	<b>MW147056</b>	NA	<b>MW139663</b>	<i>Urediniineae</i>	this paper
<i>Uromyces ari-triphylli</i>		BPI 872261 (U617)	<i>Arisaema triphyllum</i>	<b>MW147085</b>	<b>MW147079</b>	NA	<i>Urediniineae</i>	this paper
<i>Uromyces bidenticola</i>		BRIP 57564	<i>Bidens pilosa</i>	KX999897	NA	NA	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Uromyces ciceris-arietini</i>		BRIP 53857	<i>Cicer arietinum</i>	<b>MW147057</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Uromyces lomandracearum</i>		BRIP 59022	<i>Lomandra</i> sp.	KM249862	KT199392	KT199423	<i>Urediniineae</i>	McTaggart et al. (2016)
<i>Uromyces orientalis</i>		BRIP 60934	<i>Indigofera linifolia</i>	KX999899	NA	KX999934	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Uromyces pisi-sativi</i>		BRIP 60151	<i>Genista monspessulana</i>	KX999900	NA	NA	<i>Urediniineae</i>	Marin-Felix et al. (2017)



**Table S1.** (Continued).

Taxon	Type status <sup>a</sup>	Voucher number (Collection number) <sup>b</sup>	Host	28S	18S	CO3	Tree	Source
<i>Uromyces saisolae</i>		BRIP 57696	<i>Salsola australis</i>	KX999901	<b>MW147080</b>	KX999935	<i>Urediniineae</i>	Marin-Felix et al. (2017); this paper
<i>Uromyces setariae-italicae</i>		BRIP 56931	<i>Brachiaria pubigera</i>	KX999902	NA	KX999936	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Uromyces trifolii-repentis</i>		BRIP 57653	<i>Trifolium repens</i>	KX999905	NA	KX999938	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Uromyces tripogonicola</i>		BRIP 58112	<i>Tripogon loliiiformis</i>	<b>MW147058</b>	<b>MW147081</b>	<b>MW139664</b>	<i>Urediniineae</i>	this paper
<i>Uromyces viciae-fabae</i>		BRIP 60149	<i>Vicia sativa</i>	<b>MW147059</b>	NA	NA	<i>Urediniineae</i>	this paper
<i>Uromyces vignae</i>		BRIP 60213	<i>Vigna marina</i>	KX999906	NA	NA	<i>Urediniineae</i>	Marin-Felix et al. (2017)
<i>Uromyces wedeliae</i>		BRIP 60000	<i>Wollastonia biflora</i>	KX999907	NA	KX999939	<i>Urediniineae</i>	Marin-Felix et al. (2017)

<sup>a</sup>Type Status: T = type species for the genus; \* = proxy for generic type (see methods for explanation).

<sup>b</sup>numbers in parentheses are collection numbers, preceded by herbarium accession numbers. When sequences from more than one collection are used, data are separated by a /.

n.d. = no data.

NA = not applicable.


bold = new sequences generated for this paper.

Additional sequence references (not cited in paper):

- Ali et al. (2016). Mycol. Prog. In press
- Baiswa et al. (2014). Australas. Plant Dis. Notes 9 (1) In press
- Blomquist et al. (2014). Plant Dis. 98 (1), 155
- Bruns et al. 1992. Mol. Phylog. Evol. 1, 231-241
- Busby et al. (2012). Fungal Biol 116 (7), 792-801
- Crane et al. (2005). Mycoscience 46, 143-147
- Deadman et al. (2006). J. Plant Pathol. 88, 121
- Dixon et al. (2010). Mycol. Prog. 9 (4), 459-468
- Kaur et al. (2010). Plant Dis. 94, 636
- Liang et al. (2006). Mycoscience 47, 137-144
- Marin-Felix et al. (2017). Stud. Mycol. 86, 99-216
- Matheny et al. (2006) Can. J. Bot. 84, 1794-1805
- McKenzie et al. (2013). Plant Pathol Quar J Fungal Biolog 3 (2), 59-65
- Pei et al. (2005). Mycol. Res. 109 (PT 4), 401-409
- Tian et al. (2004). Mycoscience 45, 56-66
- Toome & Aime (2015). Plant Pathol. 64, 216-224



**Figure S1**



**Supplementary Fig 1.** Alluvial plot tracking generic placement at familial and subfamilial rank. Each colour represents the taxonomic hypotheses of an author. Tracks for genera with conflicting familial and subfamilial placement from multiple authors are dashed. The plot was made in R with the *ggalluvial* package..

