

References

- Abu-Shanab, B., Adwan, G., Abu-Safiya, D., Jarrar, N. and Adwan, K. (2004). Antibacterial activities of some plant extracts utilized in popular medicine in Palestine. *Turk. J. Biol.* **28**:99-102
- Abdelraouf, A.E., Amany, A.A. and Nedaa, A.A. (2011). Antibacterial, antifungal and synergistic effect of *Lawsonia inermis*, *Punica granatum* and *Hibiscus sabdariffa*. *Annals of Alquds Medicine.* **7**:33-41
- Abu-Shanab, B., Adwan, G., Abu-Safiya, D., Jarrar, N. and Adwan, K. (2004). Antibacterial activities of some plant extracts utilized in popular medicine in Palestine. *Turk. J. Biol.* **28**:99-102
- Adedapo, A.A., Jimoh, F., Afolayan, A.J. and Masika, P.J. (2009). Antioxidant Properties of the Methanol Extracts of the Leaves and Stems of *Celtis africana*. *Rec. Nat. Prod.* **3(1)**:23-31
- Adesegun, S.A., Anyika, N.E., Adekoya, T.O. and Essien, S.G. (2012). Antibacterial and antioxidant investigations of *Hallea ledermannii* leaf extract. *Indian Soc. E. Indian J. Sc. Tech.* **5(1)**:1885-1887
- Ahmed, Z., Khan, S.S., Khan, M., Tanveer, A. and Lone, Z.A. (2010). Synergistic Effect of *Salvadora persica* Extracts, Tetracycline and Penicillin Against *Staphylococcus aureus*. *African J. Basic & Appl. Sci.* **2(1-2)**:25-29
- Ahmadjian, V. (1958). Guide for the identification of algae occurring as lichen symbionts. *Bot. Notiser.* **111(4)**:632-644
- Agboke, A.A. and Esimone, C.O. (2011). Antimicrobial evaluation of the interaction between methanol extract of the lichen, *Ramalina farinacea* (*Ramalinaceae*) and Ampicilin against clinical isolates of *Staphylococcus aureus*. *J. Med. Plants Res.* **5(4)**:644-648
- Akhand, A.A., Kato, M., Suzuki, H., Liu, W., Du, J., Hamaguchi, M., Miyata, T., Kurokawa, K. and Nakashima, I. (1999). Carbonyl compounds crosslink cellular proteins and activate protein-tyrosine kinase. *J. Cell. Biochem.* **60**:1-7
- Akhand, A.A., Kato, M., Suzuki, H., Miyata, T. and Nakashima, I. (1998). Level of HgCl₂-mediated phosphorylation of intracellular proteins determines death of thymic T-lymphocytes with or without DNA fragmentation. *J. Cell. Biochem.* **71**:243-253

- Akhand A.A., Chowdhury, M.K. and Ahsan, N. (2008). *Azadirachta indica* and *Terminalia arjuna*, leaf extracts induce death of bacterial cells involving aggregation of proteins. *Bangladesh J. Microbiol.* **25(2)**:115-12
- Almas, K. and Al-Zeid, Z. (2004). The immediate antimicrobial effect of a toothbrush and miswak on cariogenic bacteria: A clinical study. *J. Contemp. Dent. Pract.* **5**:105–14
- Alexopoulos, C.J. and Mims, C.W. (1979). *Introductory Mycology*, 3rd ed. John Wiley, NewYork
- Ali, S.S., Kasoju, N., Luthra, A., Singh, A., Sharanabasava, H., Sahu, A. and Bora, U. (2008). Indian medicinal herbs as sources of Antioxidants. *Food Res. Internat.* **41**: 1–15
- Alonso, P.E., Cerdeiras, M.P., Fernandez, J., Ferreira, F., Moyna, P., Soubes, M., Vazquez, A., Veros, S. and Zunno, L. (1995). Screening of Uruguayan medicinal plants for antimicrobial activity. *J. Ethnopharm.* **45**:67-70
- Ames, B.N., Shigenaga M.K. and Hagen T.M. (1993). Oxidants, antioxidants and the degenerative diseases of aging. *Proc. Nat. Acad. Sci. U.S.A.* **90**: 7915-22
- Anderson, K.J., Teuber, S.S., Gobeille, A., Cremin, P., Waterhouse, A.L. and Steinberg, F.M. (2001). Walnut polyphenolics inhibit invitro human plasma and LDL oxidation. *J. Nutr.* **131**: 2837-2842
- Aquino, R., Moreli, S., Lauro, M.R., Abdo, S., Saija, A. and Tomaino, A. (2001). Phenolic constituents and antioxidant activity of an extract of *Anthurium versicolor* leaves. *J. Natur. Prod.* **64(8)**:1019-1023
- Arnous, A., Makris. D.P. and Kefalas, P. (2001). Effect of principal polyphenolic components in relation to antioxidant characteristics of aged red wines. *J. Agric. Food Chem.* **49**:5736-5742
- Ark, P.A., Bottimi, A.T. and Thompson, J.P. (1960). Sodium usnate as an antibiotic for Plant diseases. *Plant Disease Report.* **44**:200-203
- Arora, D. and Kaur, J. (1999). Antimicrobial activity of spices. *International J. Antimicrob. Agents*, **12**: 257-262
- Arunachalam, K. (2011). Antioxidant and antimicrobial potential of methanolic extract of Indian sacred grove *Gymnostachyum febrifugum* Benth. root. *Int. J. Pharma. Biomed. Res.* **2(3)**:67-71
- Aslan, A., Gulluce, M., Sokmen, M., Adiguzel, A., Sahin, F., and Ozkan, H. (2006). Antioxidant and Antimicrobial Properties of the Lichens

- Cladonia foliacea.*, *Dermatocarpon miniatum.*, *Everinia divaricata.*,
Evernia prunastri and *Neofuscella pulla*. *Pharm. Biol.* **44(4)**:247-252
- Asada, K. (1999). The water-water cycle in chloroplasts: scavenging of active oxygen and dissipation of excess photons. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **50**:601-639
- Awasthi, D. and Agrawal, M.R. (1970). On enumeration of lichens from the tropical and subtropical regions of Darjeeling Districts. *J. Indian Bot. Soc.* **49**:122-136
- Awasthi, D.D. (1988). A key to macro lichens of India and Nepal. *J. Hattori Bot. Lab.* **65**:207-302
- Awasthi, D.D. (1991). A key to the microlichens of India, Nepal and Sri Lanka. *Biblio. Lichenol.* **40**:1-340
- Awani, A. H., Naser, J. and Kamel, A. (2009). Antibacterial activity of common *Varthemia iphionoides*; ethanol extract alone and in combination with Cefotaxime. *Adv. Biol. Res.* **3(56)**:144-147
- Aydin, S. and Kinalioglu, K. (2013). The investigation of antibacterial activities of ethanol and methanol extracts of *Flavoparmelia caperata* (L.) Hale (Parmeliaceae) and *Rocella phycopsis* Ach. (Roccellaceae). *J. Appl. Pharma. Sc.* **3**:2
- Balaji, P., Bharath, P., Satyan, R.S. and Hariharan, G.N. (2006). *In vitro* antimicrobial activity of *Rocella montagnei* thallus extracts. *J. Trop. Med. Plants.* **7(2)**:169-173
- Balaji, P. and Hariharan, G.N. (2007). *In vitro* antimicrobial activity of *Parmotrema praesorediosum* thallus extracts. *Res. J. Bot.* **2(1)**:54-59
- Bangajavalli, S. and Ramasubhramanian, V. (2015). GC-MS analysis of bioactive components of *Aglaia elaeagnoidea*. *Euro. J. Biomed. Pharma Sc.* **2(4)**:1248-1260
- Banerjee, R.D and Sen, S.P. (1979). Antibiotic activity of bryophytes. *Bryologist.* **82**:141- 53
- Baral, B. and Maharjan, B.L. (2011). Assessment of antimicrobial and phytochemical potentials of High altitudinal Nepalese lichens *J. Microbiol. Biotech. Food Sc.* **1(2)**:98-112
- Barlow, S.M. (1990). Toxicological aspects of antioxidants used as food additives. In: B.J.F, Hudson.(Ed.). *Food Antioxidants*. Elsevier, London. pp. 253-307
- Bartak, M., Hajek, J., Vrablikova, H. and Dubova, J. (2004). High- light

- stress and photoprotection in *Umbilicaria antarctica* monitored by chlorophyll fluorescence imaging and changes in zeaxanthin and glutathione. *Plant Biol.* **3**:331-341
- Bassam, A.S., Ghaleb, A., Naser, J., Awni, A., and Adwan, K. (2006). Antibacterial activity of four plant extracts used in Palestine in folkloric medicine against Methicillin-Resistant *Staphylococcus aureus*. *Turk. J. Biol.* **30**:195-198
- Baytop, T. v(1999). Therapy with medicinal plants in Turkey (past and present). Istanbul University Publications, Istanbul. Publication no 3255/40
- Beecken, H., Gottschalk, E.M., Gizycki, U., Kramer, H., Maassen Matthes, H.G., Musso, H., Rathjen, C. and Zahorsky, U.I. (1961). Orcein and lackmus. *Angew. Chem.* **73**:665-688
- Begg, W.R., Chester, D.O. and Elix, J.A. (1979). The structure of conorlobaridone and conloxodin: New depsidones from the lichen *Xanthoparmelia xanthosorediata*. *Austr. J. Chem.* **32**:927-929
- Behera, B.C., Verma, N., Sonone, A. and Makhija, U. (2005). Antioxidant and antibacterial activities of lichen *Usnea ghattensis in vitro*. *Biotechnol. Lett.* **27**:991-995
- Behera, B.C., Verma N., Sonone, A. and Makhija, U. (2008). Antioxidant and antibacterial properties of some cultured lichens. *Bioresour. Technol.* **99(4)**:776-84
- Behera, B.C., Verma, N., Sonone, A. and Makhija, U. (2006). Determination of antioxidative potential of lichen *Usnea ghattensis in vitro*. *Food Sc. Technol.* **39**:80-85
- Benedict, R.G. and Brady, L.R. (1972). Antimicrobial activity of mushroom metabolites. *J. Pharma. Sc.* **61**:1820-1822
- Benzie, I.F.F. and Strain, J.J. (1996). Ferric reducing ability of plasma (FRAP) as measure of antioxidant power of FRAP assay. *Anal. Biochem.* **239**:70-76
- Betoni, J.E.C., Mantovani, R.P., Barbosa, L.N., Distasi, L.C. and Fernandes A. (2006). Synergism between plant extract and antimicrobial drugs used on *Staphylococcus aureus* disease. *Mem. Inst. Oswaldo Cruz.* **101**:387-390
- Bhakuni, D.S. and Dhar, M.M. (1969). Crotosparinine, a dihydroproaporphine alkaloid from *Croton sparsiflorus*. *Experimenta.* **25**:354

- Bhattacharai, H.D., Paudel, B., Hong, G.S., Lee, K.H. and Yim, H.J. (2008). Thin layer chromatography analysis of antioxidant constituents of lichens from Antarctica. *J. Nat. Med.* **62**:481-484
- Block, K.G. and Patterson, B. (1992). Fruits, vegetables and cancer prevention: a review of the epidemiological evidence. *Nutr. Cancer.* **18**: 1-29
- Bobek, P. and Galbavy, S. (2001). Effect of pleuran (beta-glucan from *Pleurotus ostreatus*) on the antioxidant status of the organism and on dimethylhydrazine-induced precancerous lesions in rat colon. *British J. Biomed. Sc.* **58**:164-168
- Bown, D. (2001). Encyclopedia of herbs and their users. Darling-Kindersley, London
- Branen, A.L. (1975). Toxicology and biochemistry of butylated hydroxyanisole and butylated hydroxytoluene. *J. American Oil Chemists Soc.* **5**:59-63
- Brodo, I.M. and Sharnoff, S. (2001). Lichens of North America. Yale University Press, London
- Burkholder, P.R., Evans, A.W., McVeigh, I. and Thornton, H.K. (1944). Antibiotic activity of lichens. *Proc. Nat. Acad. Sc. USA.* **30(9)**:250-255
- Butler, M.S. (2004). The role of natural product chemistry in drug discovery. *J. Nat. Prod.* **67**:2141-2153
- Caccamese, S., Toscano, R.M., Bellesia, F. and Pinetti, A. (1985). Methyl β -Orcinolcarboxylate and Depsides from *Parmelia furfuracea*. *J. Nat. Prod.* **48**:157-158
- Caccamese, S., Compagnini, A., Toscano, R.M. and Cascio, O. (1986): Methyl β -Orcinolcarboxylate and Atranol from the Lichen *Stereocaulon vesuvianum*. *J. Nat. Prod.* **49**:1159-1160
- Candan, M. (2007). Antimicrobial activity of extracts of lichen *Parmelia sulcata* and its salazinic acid constituent. *Z. Naturforsch.* **62(7-8)**:619-621
- Candan, M. (2006). Antimicrobial activity of extracts of lichen *Xanthoparmelia pokornyi* and its gyrophoric and stenosporic acid constituents. *Z. Naturforsch.* **61(5-6)**:619-623
- Cansarana, D., Kahya, D., Yurdakulol, E. and Atakol, O. (2006). Identification and quantitation of usnic acid from the lichen *Usnea*

- species of Anatolia and antimicrobial activity. *Z. Naturforsch.* **61**:773-776
- Cao, G., Sofic, E.R. and Prior, R.L. (1996). Antioxidant capacity of tea and common vegetables. *J. Agric. Food Chem.* **44**:3426-3431
- Carbonero, E.R., Gracher, A.H.P., Smiderle, F.R., Rosado, F.R., Sasaki G.L., Gorin, P.A.J and Lacomini, M. (2006). A β -glucan from the fruit bodies of edible mushrooms *Pleurotus eryngii* and *Pleurotus ostreatoroseus*. *Carbohydrate Polymer.* **66**:252-257
- Cardarelli, M.A., Serino, G., Campanella, L., Ercole, P., De Cicco-Nardone, F., Alesiani, O. and Rossiello, F. (1997). Antimitotic effects of usnic acid on different biological systems. *Cell. Mol. Life. Sci.* **53**:667-672
- Chand, P., Singh, M. and Rai, M. (2009). Antibacterial activity of some Indian lichens. *Ecophysiol. Occup. Hlth. J.* **9**:23-29
- Chahardehi, A.M., Ibrahim, D., Sulaiman, S.F and Leila, M.(2012). Rev Screening antimicrobial activity of various extracts of *Urtica dioica*. *Biol. Trop.* **60**(4): 1567-1576
- Chandra, S. and Singh, A. (1971). A lichen crude drug (charila) from India. *J. Res. Indian Med.* **6**:209-206
- Chatterjee, S.K., Bhattacharjee, I. and Chandra G. (2009). *In vitro* synergistic effect of doxycycline & ofloxacin in combination with ethanolic leaf extract of *Vangueria spinosa* against four pathogenic bacteria. *Indian J. Med. Res.* **130**(4): 475-478
- Chao, G.R. (2001). Antioxidant properties and polysaccharide composition analysis of ear mushrooms. Master's Thesis, National Chung-Hsing University, Taichung, Taiwan
- Chester, D.O. and Elix, J.A. (1978). The identification of four new *meta*-depsides in the lichen *Ramalina asahinae*. *Aus. J. Chem.* **31**:2745-2749
- Chevallier, A. (1996). The Encyclopedia of Medicinal plants. Dorling Kindersley, London.
- Chihara, G. (1993). Medical aspects of lentinan isolated from *Lentinus edodes* (Berk.) Sing. In: S.T. Chang., J.A. Buswell., A.W. Chiu (Eds.). *Mushroom Biology and Mushroom Products*, The Chinese University Press, Hong Kong. pp. 261-266

- Chopra, G.L. (1934). Lichens of Himalayas. Part 1. Lichens of Darjeeling and the Sikkim Himalayas. *The University of Punjab, Lahore*. Reprinted in 1981
- Clix, J.A., Whitton, A.A. and Sargent, M.V. (1984). Recent progress in the chemistry of lichen substances. *Prog. Chem. Org. Nat. Prod.* **45**:207-211
- Cocchietto, M., Skert, N., Nimis P. L., and Sava, G. (2002). A review on usnic acid, an interesting natural compound. *Naturwissenschaften.* **89**:137-146
- Colonna, S., Gaggero, N. and Richelmic, P.P. (1999). Recent biotechnological developments in the use of peroxidases. *Trends. Biotechnol.* **17**:163-168
- Cotelle, N. (2001). Role of flavonoids in oxidative stress. *Curr. Topics Med. Chem.* **1**: 569-590
- Conchran, K.W. (1978). Medicinal effects of mushrooms. In: S.T. Chang and W.A. Hayes. (Eds.). *The Biology and Cultivation of Edible Mushroom*. Academic Press, New York. pp.160-187
- Connolly, J.D., Freer, A.A., Kalb, K. and Huneck, S. (1984). Eriodermin, a dichlorodepsidone from the lichen *Erioderma physcioides* – crystal structure analysis. *Phytochem.* **23**:857-858
- Cook, N and Samman, S. (1996). Flavonoids - chemistry, metabolism, cardioprotective effects and dietary sources. *J. Nutr. Biochem.* **7**(2):66-76
- Correche, E.R., Carrasco, M. and Escudero, M.E. (1998). Study of the cytotoxic and antimicrobial activities of usnic acid and derivatives. *Fitoterapia.* **69**: 493-501
- Cowan, M.M. (1999). Plant products as antimicrobial agents. *Clin. Microbiol. Rev.* **21**:564
- Crockett, M. (2003). Antibacterial properties of four Pacific Northern Lichens. Botany (Lichenology), Oregon State University, Oregon. pp. 465
- Culberson, C.F. and Amman, K. (1979). Standard methode zur Dunnschichtchromatographie von Flech setensubstanzen. *Herzogia.* **5**:1-24
- Culberson, C.F. (1969). Chemical and botanical guide to lichen products. The University of North Carolina Press, NC. pp.101, 114, 115, 116, 129, 137, 146, 156,159,160,161,163,170,180, 183,184,191,205, 210

- Culberson, C.F. and Esslinger, T.L. (1976). 4-O-Methylolivetic and loxodellic acids: new depsides from new species of brown Parmeliae. *Bryologist*. **79**:42-46
- Culberson C.F. (1972). Improved conditions and new data for the identification of lichen products by a standardized thin-layer chromatographic method. *J. Chromatogr*, **72**: 113-125
- Culberson, C.F. and Amman, K. (1979). Standardmethode zur Dunnschichtchromatographie von Flechtensubstanzen. *Herzogia*. **5**:1-24
- Culberson, C.F. and Esslinger, T.L. (1976). 4-O-Methylolivetic and loxodellic acids: new depsides from new species of brown Parmeliae. *Bryologist*. **79**:42-46
- Culberson, C.K., Culberson, W.L. and Esslinger, T.L. (1977). Chemosyndromic variation in the *Parmelia pulla* group. *Bryologist*. **80**:125-135
- Dahake, P.A., Chakma, C.R., Chakma, C. and Joshi D. (2010). Antimicrobial and anti-inflammatory activity of *Roccella belangeriana*. *Res. J. Pharma. Phytochem*. **2(1)**:18-21
- Dahanukar, S.A., Kulkarni, R.A. and Rege, N.N. (2000). Pharmacology of medicinal plants and natural products. *Indian J. Pharmacol*. **32**:S81-S118
- Das, A.K., Dutta, B.K. and Sharma, G.D. (2008). Medicinal plants used by different tribes of Cachar district, Assam. *Indian. J. Trad. Knowledge*. **7(3)**:446-454
- Das, K., Tiwari, R.K.S and Shrivastava. D.K. (2010). Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *J. Med. Plants. Res*. **4(2)**:104-111
- De Mule, M., Caire, G.D., Cano M.D. and Haperin, D. (1991). Bioactive compound from *Nostoc muscorum* (Cyanobacterium). *Cytobios*. **66**: 169-172
- Diplock, A. T. (1997). Will the “good fairies” please prove to us that vitamin E lessens human degenerative disease? *Free. Rad. Res*. **27(5)**:11-532
- Ding, T. J., Zhou, L.J.Xu. and Gao. Z.M. (2010). Evaluation of antimicrobial activity of endophytic fungi from *Camptotheca acuminata* (Nyssaceae)T. *Genet. Mol. Res*. **9** (4): 2104-2112

- Didem, S., Melike E., Meral O. and Mahmut K.S. (2007). Free Radical Scavenging and Antimicrobial Activities of Some *Geranium* Species **28(2)** pp. 115-124
- Djeridane, A., Yousfi, M., Nadjemi, B., Boutassouna, D., Stocker, P. and Vidal, N. (2006). Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. *Food Chem.* **97**:654–660
- Djura, P., Sargent, M.V., Elix, J.A., Engkaninan, U., Huneck, S. and Culberson, C.F. (1977). Depsidone synthesis. VIII. Isolation and structure determination of hydroxy- and methoxycoumaric acids: Synthesis of Methyl methoxy-O-methylcoumarate. *Aust. J. Chem.* **30**:599-607
- Dobescu, D. (1993). Contribution to the complex study of some lichens – *Usnea* genus: Pharmacological studies on *Usnea barbata* and *Usnea hirta* species. *Rom. J. Physiology.* **30(1-2)**:101-107
- Dorman, H.J., Bachmayer, O., Kosar, M. and Hiltunen, R. (2004). Antioxidant properties of aqueous extracts from selected Lamiaceae species grown in Turkey. *J. Agr. Food. Chem.* **52**:762-770
- Dubey, R.C. and Maheshwari, D.K. (2002). Practical Microbiology. S. Chand and Company Ltd., New Delhi
- Duh, P.D., Tu, Y.Y. and Yen, G.C. (1999). Antioxidant activity of water extract of harn jyr (*Chrysanthemum morifolium* Ramat). *Food Sci. Technol.* **32**:269-277
- Duh, P.D. (1998). Antioxidant activity of Budrock (*Arctium lappa* Linn.) its scavenging effect on free radical and active oxygen. *J. Am. Oil. Chem. Soc.* **75**:455-461
- Dulger, B., Hacıoğlu, N. and Uyar, G. (2009). Evaluation of antimicrobial activity of some mosses from Turkey. *Asian J. Chem.* **21**:4093-4096
- Duman, D.C. (2009). Evaluation of usnic acid in some lichens of Turkey by HPLC analysis and screening of their antimicrobial activity. *Turk. Hij. Den. Biyol. Derg.* **66(4)**:153-160
- Dzomba, P., Togarepi, E. and Musekiwa, C. (2012). Phytochemicals, antioxidant and antibacterial properties of a lichen species *Cladonia digitata*. *Afr. J. Biotech.* **11(31)**:7995-7999

- Eastwood, M.A. (1999). Interaction of dietary antioxidants *in vivo*: how fruit and vegetables prevent diseases? *Q. J. Med.* **92**: 527-530
- El mastal, Z.Y., Aera, A. and Aam, A. (2005). Antimicrobial activity of some medicinal plant extracts in Palestine. *Pak. J. Med. Sci.* **21(2)**:187-193
- El-Banna, N.M. (2007). Antifungal activity of *Comamonas acidovorans* isolated from water pond in South Jordan. *Afr. J. Biotech.* **6(19)**:2216-2219
- Elix, J.A. (1996). Biochemistry and secondary metabolites. In: T. H. Nash (Ed.). *Lichen Biology*, Cambridge University Press, Cambridge. pp. 154-181
- Elix, J.A and Wardlaw, J.H. (2000). 4-O-Methylconhypprotocetraric acid, a new β -orcinol depsidone from the lichen *Xanthoparmelia competitiva*. *Aus. J. Chem.* **53**:1009-1010
- Elix, J.A. (2014). A Catalogue of Standardized Chromatographic Data and Biosynthetic Relationships for Lichen Substances. 3rded. Pub.Canberra
- Elix, J.A., Chester, D.O., Wardlaw, J.H and Wilkins, A.L. (1990). Synthesis of two new β -orcinol *para*-depsides in the lichen *Pseudocyphellaria norvegica*. *Aus. J. Chem.* **43**:191-196
- Elix, J.A., Gaul, K.L., Sterns, M. and Samsudun, M.W. (1987). The structure of the novel lichen xanthone, thiomelin and its congenors. *Aus. J. Chem.* **40**:1169-1178
- Elix, J.A., Jenie, U.A., Arvidsson, L. and Jorgensen, P.M. (1986). New depsidones from the lichen genus *Erioderma*. *Aus. J. Chem.* **39**:719-722
- Elix, J.A., Musidlak, H.W., Sala, T. and Sargent, M.V. (1978). Structure and synthesis of some lichen xanthonones. *Aus. J. Chem.* **31**:145-155
- Elix, J.A., Wardlaw, J.H., Archer, A.W., Lumbsch, H.T. and Plumper, M. (1997). Four new lichen depsidones from *Pertusaria* and *Lecanora* lichens. *Aus. Lichenol.* **41**: 22-27
- Elix, J.A., Barclay, C.E., Wardlaw, J.H., Archer, A.W., Sen-hua, Y. and Kantvilas. G.(1999). Four new β -orcinol *meta*-depsides from *Pertusaria* and *Siphula* lichens. *Aus. J. Chem.* **52**:837-840
- Elix, J.A., Ferguson, B.A. and Sargent, M.V. (1974). The structure of alectronic acid and related lichen metabolites. *Aus. J. Chem.* **27**:2403-2411

- Elizabeth, K.M. (2005). Anti microbial Activity of *Terminalia belerica*. *Indian. J. Clinic. Biochem.* **20(2)**:150-153
- El-Massy, K.F., El-Ghorab, A.H., Shaaban, H.A. and Shibamoto, T. (2009). Chemical compositions and antioxidant/ antimicrobial activities of various samples prepared from *Schinus terebinthifolius* cultivated in Egypt. *J. Agric. Food Chem.* **57**:5265-5270
- Elmastas, M., Gucin, I., Ozturk, L., Gokce, I. (2005). Investigation of antioxidant properties of spearmint (*Mentha spicata* L.) *Asian J. Chem.* **17**:137-148
- Ernst-Russell, M.A., Elix, J.A., Chai, C.L.L., Rive, M.J. and Wardlaw, J.H. (2000). The structure and stereochemistry of coronatoquinone, a new pyranonaphthazarin from the lichen *Pseudocyphellaria coronata*. *Aus. J. Chem.* **53**:303-306
- Esimone C.O. and Adikwu M.U. (1999). Antimicrobial activity and cytotoxicity of *Ramalina farinacea*. *Fitoterapia.* **70**:428-431
- Esimone, C.O., Kenneth, C.O., Adikwu, M.U., Ibezim, Emmanuel C., Abonyi, D.O., Odaibo, G.N. and Olaleye, D.O. (2007). *In vitro* evaluation of the antiviral activity of extracts from the lichen *Parmelia perlata* (L.) Ach. against three RNA viruses. *J. Infect. Dev. Count.* **1(3)**:315-320
- Esimone, C.O., Iroha, I.R., Ibezim, E.C., Okeh, C.O. and Okpana, E.M. (2006). *In vitro* evaluation of the interaction between tea extracts and penicillin G against *Staphylococcus aureus*. *Afr. J. Biotechnol.* **5**:1082
- Eugene, W., Nester, D.G., Anderson, C., Evans, R. Jr., Nancy, P. and David, H. (2004). *Microbiology, A Human Perspective*. McGraw Hill (Higher Education), U.S.A.
- Evans, J.S., Pattison, E. and Morris, F. (1986). Antimicrobial agents from plant cell culture. *In: P. Morris., A. Scraggs., A. Stafford. and M. Fowler. (Eds.). Secondary metabolites in plant cell culture*. Cambridge University Press, Cambridge, UK
- Fathi-Azad, F., Garjani, A., Maleki, N. and Ranjdost S. (2005). Study of the hypoglycemic activity of the hydroalcoholic extract of *Urtica dioica* in normal and diabetic rats. *Pharma. Sci.* **94(2)**:65-69
- Faujan, H., Noriham, N., Norrakiah, A., and Babji, A.S. (2009). Antioxidant activity of plants methanolic extracts containing phenolic compounds. *Afr. J. Biotechnol.* **8(3)**:484-489

- Fazio, A.T., Adler, M.T., Bertoni, M.D., Sepulveda, C.S., Damonte, E.B. and Maier, M.S. (2007). Lichen secondary metabolites from the cultured lichen mycobionts of *Teloschistes chrysophthalmus* and *Ramalina celastri* and their antiviral activities. *Z. Natur.* **62**:543-549
- Fernandez. P. H., Izquierdo, M., Robaina, L., Valencia, A., Salhi, M. and Montero, D. (1996). The effect of dietary protein and lipid from squid and fish meals on egg quality of broodstock for gilthead seabream (*Sparus aurata*). *Aquaculture.* **148(2-3)**:233-246
- Freitas, M., Costa, N., Rodrigues, M., Marques, J., Vieira, De. and Silva. M. (2011). Lichens as bioindicators of atmospheric pollution in Porto, Portugal. *J. Biodiversity Ecol. Sci.* **1(1)**:30-39
- Fuji. M., Miyaichi, Y. and Tomimori, T. (1996b). Studies on Nepalese crude drugs. XXII on the phenolic constituents of the rhizome of *Bergenia ciliata* (Haw.) Stern *Nat. Med.* **50(6)**:404-407
- Fujii, Y., Furukawa, M., Hayakawa, Y., Sugawara, K. and Shibuya, T. (1991). Survey of Japanese medical plants for detection of allelopathic properties. *Weed Res. Jpn.* **36**:36-42
- Furmanowa, M. and Rapczewska, L. (1993). *Biotechnol. Agr. Fores.* **2**:18-33
- Gadow, A., Joubert, E. and Hansmann, C.F. (1997). Comparison of the antioxidant activity of aspalathin with that of other plant phenols of rooibos tea (*Aspalathus linearis*), β tocopherol, BHT, and BHA. *J. Agr. Food.* **45**:632-638
- Galun, M. (1988). CRC Handbook of Lichenology. CRC Press, Boca Raton, Florida, **3** pp. 95-107
- Gamez, E.J., Luyengi, L., Lee, S.K., Zhu, L.f., Zhou, B.N., Fong, H.H., Pezzuto, J.M. and Kinghorn, A.D. (1998). Antioxidant flavonoid glycosides from *Daphniphyllum calycinum*, *J. Natur. Prod.* **61(5)**: 706-708
- Garty, J. (2000). Environment and elemental content of lichens. In: B. Markert, and K. Friese. (Eds.). *Trace Elements-Their Distribution and Effects in the Environment.* Elsevier Sci. BV. pp. 245-276
- Gauslaa, Y. and Solhaug, K.A. (2001). Fungal melanins as a sun screen for symbiotic green algae in the lichen *Lobaria pulmonaria*. *Oncologia.* **126**:462-471

- Ghafar, M.F.A., Nagendra, P.K., Weng, K.K and Ismail, A. (2010). Flavonoid, hesperidine, total phenolic contents and antioxidant activities from Citrus species. *Afr. J. Biotechnol.* **9**:326-330
- Ghasemi, Y., Moradian, A., Mohagheghzadeh, A., Shokravi, S. and Morowvat, M.H. (2007). Antifungal and antibacterial activity of the microalgae collected from paddy fields of Iran: characterization of antimicrobial activity of *Chlorococcus disperses*. *J. Biol. Sc.* **7**:904-910
- Ghate,N.B., Chaudhuri, D.,Sarkar,R., Sajem,A.L., Panja,S, Rout, J and Mandal, N.(2013). An Antioxidant Extract of Tropical Lichen, *Parmotrema reticulatum*, Induces Cell Cycle Arrest and in Breast Carcinoma Cell Line. MCF-7. *PloS one.* **8 (12)**, e82293
- Gilani, A.H. and Rahman, A.U. (2005). Trends in Ethnopharmacology. *J. Ethnopharmacol.* **100**:43-49
- Goodwin, J. (1997). The limits of modern medicine. *J. Alt. Med.* **278**:1399–1400
- Gomathi. S., Ambikapathy, V. and Panneerselvam, A. (2011). Antimicrobial activity of some medical plants against *Pythium debaryanum* (Hesse). *J. Microbiol. Biotechnol. Res.* **1**:8-13
- Gonzales, A.G., Barrera, J.B., Rodrigues, E.M. and Hernandez , C.E. (1991). Chemical constituents of the lichen *Cladonia macronesica*. *Z. Naturforschung.* **46c**:12-18
- Gonzales, A.G., Martin, L.D, and Perez, C. (1974): Three new triterpenes from the lichen *Xanthoria resendei*. *Phytochem.* **13**:1547-1549
- Grice, H.C. (1986). Safety evaluation of butylated hydroxytoluene (BHT) in the liver, lung and gastrointestinal tract. *Food Chem.Toxicol.* **24**:1127-1130
- Grabley, S. and Thiericke, R. (1999). Drug discovery from nature. Springer, London. pp. 5-7
- Griffin, D.H. (1994). Molecular architecture. Fungal Physiology, 2nd ed., Wiley Liss, New York. pp. 65-74
- Gulcin, I., Elias, R., Gepdiremen, A., Boyerm, L. and Koksals, E. (2007). A comparative study on the antioxidant activity of fringe tree (*Chionanthus virginicus*, L.) extracts. *Afr. J. Biotech.* **64(4)**:410-418

- Gulcin, I., Kufrevioglu, O.I., Oktay, M. and Buyukokuroglu, M.E. (2004). Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.). *J. Ethnopharmacol.* **9**:205-215
- Gulcin, I., Oktay, M., Kufrevioglu, O.I. and Aslan, A. (2002). Determination Of antioxidant activity of lichen *Cetraria islandica* (L) Ach. *J. Ethnopharmacol.* **79**:325-329
- Gulcin, I., Oktay, M., Kyreççi, E., and Kufrevioglu, O.I. (2003). Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chem.* **83**:371-382
- Gulluce, M., Aslan. A., Sokmen, M., Sahin, F., Adiguzel, A., Agar, G., and Sokmen, A. (2006). Screening the antioxidant and antimicrobial properties of the lichens *Parmelia saxatilis*, *Platismatia glauca*, *Ramalina pollinaria*, *Ramalina polymorpha* and *Umbilicaria nylanderiana*. *Phytomed.* **13**:515-521
- Gulumser, A., Dogan, N.M., Duru, M.E. and Kivrak, I. (2010). Phenolic profiles, antimicrobial and antioxidant activity of the various extracts of *Crocus* species in Anatolia. *Afr. J. Microbiol. Res.* **4(11)**:1154-1161
- Gunde-Cimerman, N. (1999). Medicinal value of the genus *Pleurotus* (Fr.) P. Karst. (Agaricales, Basidiomycetes). *Int. J. Med. Mushroom.***1**: 69-80
- Gupta, V.K., Darokar, M.P., Saikia, D., Pal, A., Fatima, A. and Hanja S.P.S. (2007). Antimycobacterial activity of lichens. *Pharma. Biol.* **45**:200-204
- Gupta, D.R. and Ahmed B. (1990). Emarginatosides B and C: Two new saponins from *Sapindus emarginatus* fruits. *Indian J. Chem.* **29(B)**:268-270
- Gupta, S.K. and Paul, A.K. (1995). Studies on antimicrobial substances from lichens of Darjeeling hills. *J. Hill Res.* **8(1)**:61-66
- Gupta, V.K., Darokar, M.P., Saikia, D., Pal, A., Fatima, A. and Khanuja, S.P.S. (2007).Antimycobacterial activity of lichens. *Pharma. Biol.* **45**:200-204
- Gursoy, N. and Tepe, B. (2009). Determination of the antimicrobial and antioxidative properties and total phenolics of two "endemic" Lamiaceae species from Turkey: *Ballota rotundifolia* L. and *Teucrium chamaedrys* C. Koch., *Plant Foods Hum. Nutr.* **64(2)**:135-140
- Halama, P., Van, and Haluwin, C. (2004). Antifungal activity of lichen extracts and lichenic acids, *Bio. Control.* **49(1)**:95-107

- Halliwell, B. and Gutteridge, J.M. 1989. Free radicals in Biology and Medicine. Oxford: Clarendon Press. pp. 96-98
- Halliwell, B. and Gutteridge, J.M. (1990). Role of free radicals and catalytic metal ions in human disease: An overview 186:**1-85**
- Halliwell, B. (1995). How to characterize an antioxidant? An update. *Biochem. Soc. Symp.* **61**:73-101
- Halliwell, B. and Gutteridge, J.M.C. (1999). Free Radicals in Biology and Medicine, 3rd edn. Oxford: Clarendon Press
- Haq, M.U., Reshi, Z.A., Upreti, D.K. and Sheikh, M.A. (2012). Lichen wealth of Jammu and Kashmir- a promising plant source for bioprospection. *Life Sci.* **9(4)**:926-929
- Hickey, B.J., Lumsden, A.J., Cole, A.L.J. and Walker, J.R.L. (1990). Antibiotic compounds from New Zealand plants: methyl haematommate, an anti-fungal agent from *Stereocaulon ramulosum*. *New Zealand Nat. Sc.***17**:49-53
- Hidalgo, M.E., Quilhot, F.W. and Lissi, E. (1994). Antioxidant activity of depsides and depsidones. *Phytochem.* **37**:1585-1587
- Horiuchi, K., Shiota, S., Kuroda, T., Hatano, T., Yoshida, T. and Tsuchiya, T. (2007). Antimicrobial activity of oleanolic acid from *Salvia officinalis* and related compounds on vancomycin-resistant enterococci (VRE). *Biol. Pharm. Bull.* **30**:287
- Huneck, S. (1999). The significance of lichens and their metabolites. *Naturwissenschaften.* **86**:559-570
- Huneck, S. and Yoshimura, I. (1996). Identification of Lichen Substances. Springer-Verlag, Berlin, Heidelberg, New York. pp. 143, 166, 175, 183, 241, 254, 304, 335, 347, 349, 386, 399, 402, 493
- Hoskeri, H. Joy, Krishna. V. and Amruthavalli, C. (2010). Effects of extracts from lichen *Ramalina pacifica* against clinically infectious bacteria. *Researcher.* **2(3)**:81-85
- Hu, Z.Q., Zhao, W.H., Asano, N., Yoda, Y., Hara, Y. and Shimamura, T. (2002). Epigallocatechin gallate synergistically enhances the activity of carbapenems against methicillin-resistant *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **46(2)**:558-560
- Huda, Faujan. N., Noriham, A., Norrakiah, A.S. and Babji, A.S. (2007). Antioxidative activities of water extracts of some Malaysian herbs. *ASEAN Food. J.* **14**:61-68

- Hugenholtz, P. (2002). Exploring prokaryotic diversity in the genomic era. *Genome Biol.* **3** (2):3.1–3.8
- Hugo, W.B. and Russell, A.D. (1983). *Pharmaceutical Microbiology*, 3rd ed. Blackwell Scientific Publications. **51**:33-35
- Huneck, S. (1999). The significance of lichens and their metabolites. *Naturwissenschaften.* **86**:559-570.
- Huneck, S. and Yoshimura, I. (1996), Identification of Lichen Substances. Springer, Berlin, Heidelberg, p. 143, 166, 175, 183, 241, 254, 335, 304, 347, 349, 386,399, 402, 493
- Huneck, S. (1972). Flechteninhaltsstoffe – XCIII. Struktur der (-)-Placodiolsäure. *Tetrahedron.* **28**:4011-4017
- Huneck, S., Porzel, A., Schmidt, J., Feige, G.B. and Posner, B. (1993). Crustinic acid, a new tridepside from *Umbilicaria custulosa*. *Phytochem.* **32**:475-477
- Huneck, S., Tonsberg, T. and Bohlmann, F. I. (1986). (-)-allo-Pertusaric acid and (-)-dihydropertusaric acid from the lichen *Pertusaria albescens*. *Phytochem.* **25**:453-459
- Ibanez, E., Kubatova, A., Senorans, F.J., Cavero, S., Reglero, G., and Hawthorne, S.B. (2003). Sub-critical water extraction of antioxidant compounds from rosemary plants. *J. Agr. Food. Chem.* **51**:375-382
- Ignatowicz, E. and Rybczynska, M. (1994). Some biochemical and pharmacological aspects of free radical-mediated tissue damage. *Pol. J. Pharmacol.* **46**: 103-114
- Ingolfssdottir, K. (2002). Molecules of interest: usnic acid. *Phytochem.* **61**: 729-736
- Ingolfssdottir, K., Chung, G.A.C., Skulason, V.G., Gissurarson, S.R. and Vilhelmsdottir, M. (1998). Antimycobacterial activity of lichens metabolites *in vitro*. *Eur. J. Pharm. Sci.* **6**:141–144
- Ingolfssdottir, K., Hjalmarsdottir, M.A., Guojonsdottir, G.A., Brynjolfsdottir, A., Sigurdsson, A. and Stein-grimsson.O. (1997). *In vitro* susceptibility of *Helicobacter pylori* to protolichesterinic acid from *Cetraria islandica*. *Antimicrob. Agents Chemotherap.* **4**:215-217
- Ingondolfsdottir, K. (2002). Usnic acid. *Phytochem.* **61**:729-736
- Ishizaki, T., Kishi, T. and Sasaki, F. (1996). Effect of probucol, an oral hypocholesterolemic agent, on acute tobacco smoke inhalation in rats. *Clin. Sci.* **90**:517-523

- Islam, M.J., Barua, S., Das, S., Khan, M.S. and Ahmed, A. (2008). Antibacterial activity of some indigenous medicinal plants. *J .Soil .Nature.* **2(3)**:26-28
- Islam, U.M., Khan, I.A., Khan, U., Gill, M.A., Shahab. U.D. and Ahmad, A. (2002). Antifungal activity evaluation of *Berginia ciliata*. *Pak. J. Pharmacol.* **19(2)**:1-6
- Javeria, S., Shahi, S.K., Shahi, M.P. and Upreti, D.K. (2013). *Parmotrema nilgherrense*: potential antimicrobial activity against drug resistant pathogens. *Int. J. Microbial Resource Technol.* **2(1)**:36-40
- Jayamalar, P. and Suhaila, M. (1998). Antioxidative activities of Malaysian plants. *Malays. Appl. Biol.* **27**:56-58
- Jeffreys, D. (2005). Aspirin: the remarkable story of a wonder drug. Bloomsbury New York. pp. 38-40
- Jenner, P. (1994). Oxidative damage in neurodegenerative disease. *Lancet.* **344**:796-798
- Jerez, M., Selga, A., Sineiro, J., Torres, J.L. and Nune, M.J. (2007). A comparison between bark extracts from *Pinus pinaster* and *Pinus radiata*: Antioxidant activity and procyanidin composition. *Food. Chem.* **100**:439-444
- Johnson, O.O. and Ayoola G.A. (2015). Antioxidant activity among selected medicinal combinations (Multi-component Herbal Preparation). *Int. J. Pharma. Res. Health Sc.* **3(1)**:526-532
- Jose, N., Ajith, T.A. and Jananrdhanan K.K. (2002). Antioxidant, anti-inflammatory and antitumor activities of culinary-medicinal mushroom *Pleurotus pulmonarius* (Fr.) Quel. (Agaricomycetideae). *Int. J. Med. Mushrooms.* **4**:59-66
- Joseph, J.A., Shukitt-Hale, B., Denisova, N.A., Bielinski, D., Martin, A. McEwen, J. and Bickford, P.C. (1999). *J. Neurosci.* **19(18)**:8114-8121
- Kahkonen, M., Hopia, A.I., Vuorela, H.J., Rauha, J.P., Pihlaja, K. and Kujala, T.S. (1999). Antioxidant activity of plant extracts containing phenolic compounds. *J. Agric. Food. Chem.* **47(10)**:3954-62
- Kakwaro, J.O. (1976). Medicinal Plants of East Africa. East Africa Literature Bureau, Nairobi

- Kanatt, S.R., Chander, R. and Sharma, A. (2007). Antioxidant potential of mint (*Mentha spicata* L.) in radiation- processed lamb meat. *Food Chem.* **100**:451-458
- Karagoz, A., Dogruoz, N., Zeybek, Z. and Aslan, A. (2009). Antibacterial activity of some lichen extracts. *J. Med. Pl. Res.* **3**:1034-1039
- Karaman, I., Sahin, F., Gulluce, M., Ogutcu, H., Sengul, M. and Adiguzel A.(2003). Antimicrobial activity of aqueous and methanol extracts of *Juniperus oxycedrus* L. **85(2-3)**:231-5
- Karmegam, N., Karuppusamy, S., Prakash., M., Jayakumar, M and Rajasekar, K. (2003). Antibacterial potency and synergistic effect of certain plant extracts against certain food-borne diarrheagenic bacteria. *Int. J. Biomed. Pharma. Sc.* **2(2)**:88-93
- Karunaratne, V., Bombuwela, K., Kathirgamanathar, S. and Thadhani, V.M. (2005). Lichens: a chemically important biota. *J. Nat. Sc. Foundation of Sri Lanka.* **33 (3)**:169-186
- Katalinic, V., Milos, M., Kulisic, T. and Jukic, M. (2006). Screening of 70 medicinal plant extracts for antioxidant capacity and total phenols. *Food Chem.* **94(4)**:550-557
- Kaushik, R., Narayanan, P., Vasudevan, V., Muthukumaran, G. and Antony, U. (2010). Nutrient composition of cultivated *Stevia* leaves and the influence of polyphenols and plant pigments on sensory and antioxidant properties of leaf extracts. *J. Food Sci. Technol.* **47**:27-33
- Kehrer, J.P. (1993). Free radicals as mediators of tissue injury and disease. *Crit. Rev. Toxicol.* **23**:21-48
- Kekuda, K.S., Prashith, T.R., Kumar, P., Vinayaka, S.V., Sudharshan, S.J., Mallikarjun, N. and Swathi, D. (2010). Studies on antibacterial, anthelmintic and antioxidant activities of a Macrolichen *Parmotrema pseudotinctorum* (des. Abb.) Hale (Parmeliaceae) from Bhadra Wildlife Sanctuary, Karnataka. *Int. J. Pharm. Tech. Res.* **2(2)**:1207-1214
- Kekuda, P.T.R., Vinayaka, K.S., Praveen Kumar, S.V., Sudharshan, S.J. (2009). Antioxidant and antibacterial activity of lichen extracts, honey and their combination. *J. Pharm. Res.* **2**:1875-1878
- Kekuda, P.T.R., Vinayaka, K.S., Swathi. D., Suchitha, Y., Venugopal, and Mallikarjun, T.M. (2011). Mineral composition, total phenol content and antioxidant activity of a macrolichen *Everniastrum cirrhatum* (Fr.) Hale (Parmeliaceae), *E-J. Chem.* **8(4)**:1886-1894

- Keniya, M.V., Lukash, A.I. and Guskov, E.P. (1993). The role of low molecular weight antioxidants in oxidative stress, *Usp. Sov. Biol.* **113(4)**:456-470
- Krishnaraju, A.V., Rao, T.V.N., Sundararaju, D. Vanisree, M., Tsay, H and Subbaraju, G.V. (2006). Biological screening of medicinal plants collected from Eastern Ghats of India using *Artemia salina* (brine shrimp test). *Internat. J. Appl. Sci. Eng.* **4(2)**:115-125
- Krishna, D.R. and Venkataramana, D. (1992). Pharmacokinetics of d-(+)-usnic acid after intravenous and oral administration. *Drug Metabol. Dispos.* **20**, 909-911
- Keogh, M.F. (1978). New β -orcinol depsidones from the *Xanthoparmelia quintaria* and a *Thelotrema* species. *Phytochemistry.* **17**:1192-1193
- Khanam, R., Chaudhary, B.L., Khanam, S. and Kumar, P. (2011). Antibacterial activity of *Marchantia Palmata*. *Asian J. Biochem. Pharma. Res.* **2(1)**:27-36
- Kharel, M.K., Rai, N.P., Manandhar, M.D., Elix, J.A. and Wardlaw, J.H. (2000). Dehydrocollatolic acid, a new depsidone from the lichen *Parmotrema nilgherrense*. *Aust. J. Chem.* **53(10)**:891-92
- Kim, K.S., Lee, S., Lee, Y.S., Jung, S.H., Park, Y., Shin, K. and Kim, B.K. (2003). Antioxidant activities of the extracts from the herbs of *Artemisia apiacea*. *J. Ethnopharmacol.* **85**:67-72
- Kinoshita, K., Togawa, T., Hiraishi, A., Yuko Nakajima, Y., Koyama, K., Narui, T., Wang Li, S and Takahashi, K. (2010). Antioxidant activity of red pigments from the lichens *Lethariella seranderi*, *L. cashmeriana*, and *L. sinensis*. *J. Nat. Med.* **64**:85-88
- Kinoshita, K., Matsubara, H., Koyama, K., Takahashi, K., Yoshimura, I., Yamamoto, Y., Miura, Y., Kinoshita, Y. and Kawai, K.I. (1994). Topics in the chemistry of lichen compounds. *J. Hattori. Bot. Lab.* **76**:227-233
- Kinoshita, K., Narui, T., Koyama, K., Takahashi, K., Culberson, C.F. and Nishino, Y. (2005). Secondary metabolites from *Lethariella seranderi*. *Lichenol.* **4**:7-11
- Kinoshita, K., Takatori, K., Narui, T., Culberson, C.F., Hasumi, M., Nishino, Y., Koyama, K. and Takahashi, K. (2004). A novel secondary metabolite from *Lethariella seranderi*. *Heterocycles.* **63**:1023-1026

- Kirmizigul, S., Koz, O., Anil, H., and Icli, S. (2003). Isolation and structure elucidation of novel natural products from Turkish Lichens. *Turk. J. Chem.* **27**:493-500
- Kirk, P.M., Cannon P.F., David J.C. and Stalpers, J.A. (2001). Ainsworth and Bisby's dictionary of the fungi. 9th edn. CAB International, Oxford
- Kiselova, Y., Ivanova, D., Chervenkov, T., Gerova, D., Galunska, B. and Yankova, T.(2006). Correlation between the *in vitro* antioxidant activity and polyphenol content of aqueous extracts from Bulgarian herbs. *Phytother. Res.* **20**:961-965
- Komiya, T. and Kurokawa, S. (1970). Loxodin, a depsidone of lichens of *Parmelia* species. *Phytochem.* **9**:1139-1140
- Koptina, A., Shcherbakova, A., Soldati, F. and Ulrich-Merzenich, G. (2013). Total phenolic content and antioxidant capacity of lichen extracts. *Z. Phytother.* **34**:35
- Kosanic, M. and Rankovic B. (2011). Lichens as possible sources of antioxidants. *J. Pharm. Sci.* **24(2)**:165-170
- Kosanic, M., Manojlovic, N., Jankovic, S., Stanojkovic, T., Rankovic, B. (2013). *Evernia prunastri* and *Pseudoevernia furfuraceae* lichens and their major metabolites as antioxidant, antimicrobial and anticancer agents. *Food. Chem. Toxicol.* **53**:112-118
- Kosanic, M., Seklic, D. and Markovic, S. (2014b). Evaluation of antioxidant, antimicrobial and anticancer properties of selected lichens from Serbia. *Dig. J. Nanomater. Bios.* **9**:273-287
- Kosanic, M., Rankovic, B. and Vukojevic, J. (2011). Antioxidant properties of some lichen species. *J. Food Sci. Technol.* **48(5)**:584-590
- Kosanic, M., Rankovic, B. and Stanojkovic, T. (2013). Investigation of selected Serbian lichens for antioxidant, antimicrobial and anticancer properties. *J. Anim. Plant Sci.* **23**:1628-1633
- Kosanic, M. and Rankovic, B. (2015). Secondary Metabolites. B. Rankovic (ed.). Lichen Secondary Metabolites, DOI 10.1007/978-3-319-13374-4_4
- Krishnaraju, V.A., Rao, V.C., Rao N.V.T., Reddy, K.N. and Golakoti, T. (2009). *In vitro* and *In vivo* antioxidant activity of *Aphanamixis polystachya* bark. *Am. J. Infec. Dis.* **5(2)**:60-67

- Kristmundsdottir, T., Aradottir, H.A.E., Ingolfsdottir, K. and Ogmundsdottir, H.M. (2002). Solubilization of the lichen metabolite (+) -usnic acid for testing in tissue culture. *J. Pharm. Pharmacol.* **54**:1447-1452
- Kumar, B. (2009). Assessment of Lichen Species in a Temperate Region of Garhwal Himalaya, India. *J. Am. Sci.* **5(4)**:107-112
- Kumar, S.V.P., Kekuda, T.R.P and Vinayaka, K.S. (2010a). Studies on antibacterial, anthelmintic and antioxidant activities of a macrolichen *Parmotrema pseudotinctorum* (des. Abb.) Hale (Parmeliaceae) from Bhadra wildlife sanctuary, Karnataka. *Int. J. Pharm. Tech. Res.* **2**:1207-1214
- Kumar, K and Upreti, D.K. (2001). *Parmelia* sp. (lichens) in ancient medicinal plant.
- Awasthi, D. and Agrawal, M.R. (1970). On enumeration of lichens from the tropical and sub- tropical regions of Darjeeling District. *J. Indian Bot. Soc.* **49**:122-136
- Kumpulainen, J.T. and Salonen, J.T. (1999). Natural antioxidants and anticarcinogens in nutrition, health and disease, the Royal Society of Chemistry, UK, pp. 178-187.
- Kupchan, S.M. and Koppennan, H.L. (1975). *In vitro* activity of lichen secondary metabolite (usnic acid). *Experimenta.* **31**: 625
- Lange, O.L and Green, T.G.A. (2003). Photosynthetic performance of a foliose lichen of biological soil crust communities: long term monitoring of CO₂ exchange of *Cladonia convoluta* under temperate habitat conditions. *Bibliothica Lichenologica.* **86**:257-280
- Laloue, H., Weber-Lofti, F., Lucau-Danila, A. and Gullemat, P. (1997). Identification of ascorbate and guaiacol peroxidase in needle chloroplasts of spruce trees. *Plant Physiol. Biochem.* **35**:341-346
- Lauterwein, M., Oethinger, M., Belsner, K., Peters, T. and Marre, R. (1995). *In vitro* activities of the lichen secondary metabolites vulpinic acid, (+)-usnic acid and (Ð)- Usnic acid against aerobic and anaerobic microorganisms. *Antimicrob. Agents Chemother.* **39**:2541-2543
- Laskowska, E., Kuczynska-Wisnik, D., Bak, M. and Lipinska, B. (2003). Trimethoprim induces heat shock proteins and protein aggregation in *E. coli* cells. *Curr. Microbiol.* **47**:286-289

- Lawrey, J.D. (1983). Vulpinic and pinastric acids as lichen antiherbivore compounds: contrary evidence. *Bryologist*. **86**:365-369
- Lawrey, J.D. (1986). Biological role of lichen substances. *Bryologist*. **89**:11-122
- Leboritz, R.L., Zhang, H., Vogel, H., Catwright, J., Dionne, L., Lu, N., Husang, S. and Matzk, M.M. (1996). Neurodegeneration, myocardial injury and perinatal death in mitochondrial superoxide dismutase-deficient mice. *Proc. Nat. Acad. Sci.* **93**:9782-9787
- Liyana, Pathiranan., Chandrika, M. and Shahidi, F. (2005). Antioxidant activity of commercial soft and hard wheat (*Triticum aestivum* L.) as affected by gastric pH conditions. *J. Agri. Food Chem.* **53**:2433
- Longton, R.E. (1988). Biology of polar bryophytes and lichens. Cambridge Univ. Press
- Louwhoff, S.H.J.J. and Crisp, M.D. (2000). Phylogenetic analysis of *Parmotrema* (Parmeliaceae: Lichenized ascomycotina). *The Bryologist*. **103(3)**:541-54
- Luck, H. (1974). In: Methods in enzymatic analysis. 2nd ed. New York: Bergmeyer Academic Press.
- Luo, H., Ren, M., Lim, M.K., Koh, J.Y., Wang, S. and Hur, J.S. (2006). Antioxidative Activity of lichen *Thamnolia vermicularis* *in vitro*. *Mycobiol.* **34(3)**:124-127
- Madamombe, I.T. and Afolayan, A.J. (2003). Evaluation of antimicrobial activity of extracts from South African *Usnea barbata*. *Pharmaceut. Biol.* **41**:199-202
- Maestri, D.M., Nepote, V., Lamarque, A.L. and Zygadlo, J.A. (2006). Natural products as antioxidants. *Res. Signpost.* **1**:105-135
- Mahandru, M.M. and Tajbakhsh, A. (1983). Fulgoicin, a new depsidone from the lichen *Fulgensia fulgida*. *J. Chem. Soc. Perkin. Transactions.* **1**: 2249-2251
- Mahesh, B. and Satish, S. (2008). Antimicrobial activity of some important medicinal plant against plant and human pathogens. *World J. Agric. Sc.* **4(S)**:839-843
- Makabe, H., Maru, N., Kuwabara, A., Kamo, T. and Hirota, M. (2006). Antiinflammatory sesquiterpenes from *Curcuma zedoaria*. *Nat. Prod. Res.* **20**: 680-685

- Manjulatha, K., Jaishree, B. and Purohit, M.G. (2012). Antimicrobial activity of fruits of *Sapindus emarginatus*. *J. Pharmacog.* **3(2)**:55-58
- Manojlovic, N., Vasiljevic, P., Juskovic, M., Najman, S., Jankovic S, and Milenkovic-Andjelkovic, A. (2010). HPLC analysis and cytotoxic potential of extracts from the lichen, *Thamnolia vermicularis* var. *subuliformis*. *J. Med. Plants Res.* **4**:817-823
- Manojlovic, N.T. (2005). Antifungal activity of *Rubia tinctorum*, *Rhamnus frangula* and *Coloplaca cerina*. *Fitoterapia.* **76(2)**:244-46
- Manojlovic, N.T., Vasiljevic. P.J., Gritsanapan W., Supabphol.R and Ivana Manojlovic. (2010). Phytochemical and antioxidant studies of *Laurera benguelensis* growing in Thailand. *Biol. Res.* **43**:169-176
- Marijana, K. and Branislav, R. (2010). Screening of antimicrobial activity of some lichen species *in vitro*. *Kragujevac. J. Sci.* **32**:65-72
- Marshak , A. and Kushner, M. (1950). The action of Streptomycin and Usnic acid on development of tuberculosis in guinea pigs. *Pub. Health. Rep,* **65(5)**:131-162
- Matias, E.F., Santos K.K., Almeida T.S., Costa J.G. and Continuo, H.D. (2011). *Biomed. Prevent. Nutr.* **1**:57
- Mau, J.L., La, E.Y.C., Wang, N.P., Chen, C.C., Chang, C.H. and Chyau, C.C. (2003). Composition and antioxidant activity of the essential oil from *Curcuma zedoaria*. *Food Chem.* **82**:583-591
- Meda, A., Lamien, C. E., Romito, M., Millogo, J. and Nacoulma, O.G. (2005). Determination of the total phenolic, flavonoid and proline contents in burkina fasan honey, as well as their radical scavenging activity. *Food. Chem.* **91**:571-577
- Mehmet, A. and Sevda, K. (2009). Antimicrobial activity of *Pleurotus eryngii* var. *ferulae* grown on various agro-wastes. *Eur. Asia J. Bio.Sci.***3**:58-63
- Merzlyak, M.N. (1999). *Sorosovskiy Obrazovatel'nyi Zh.* **9**:20-26
- Miliauskas, G., Venskutonis, P.R. and Beek, T.A. (2004). Screening of radical scavenging activity of some medicinal and aromatic plant extracts. *Food Chem.* **85**:231-237
- Mohammed, F.A.G., Nagendra, P.K., Kong, K.W. and Amin, I. (2010). Flavonoid, hesperidine, total phenolic contents and antioxidant activities from Citrus species. *Afr. J. Biotechnol.* **9**:326-330
- Muggia, L., Schmitt, I. and Grube, M. (2009). Lichens as treasure chests of natural products. *SIM News.* **59**:85-97

- Muhammad, H.S and Muhammad S. (2005). The use of *Lawsonia inermis* Linn. (Henna) in the management of burn wound infection. *Afr. J. Biotechnol.* **4(9)**:934-937
- Mukerjee, T., Bhalla, N., Singh, G. and Aulakh, H.C. (1984). *Jain, Indian Drugs.* **21(6)**:224
- Mukherjee, P.K., Bhattacharya, S., Saha, K. Giri, S.N., Pal, M. and Saha, B.P. (1997). Antibacterial evaluation of *Drymaria cordata* Willd (Fam.Caryophyllaceae) extract. *Phytotherapy Res.***11(3)**:249-250
- Mukherjee, S., Pawar, N., Kulkarni, O., Nagarkar, B., Thopte, S., Bhujbal, A. and Pawar, P. (2011). Evaluation of free-radical quenching properties of standard Ayurvedic formulation Vayasthapana Rasayana. *BMC Complem. Altern. Med.***11**:38
- Muller, K. (2001). Pharmaceutically relevant metabolites from lichens. *Appl. Microbiol. Biotechnol.* **56(1-2)**:9-16
- Muller, K. (2002). Pharmaceutically relevant metabolites from lichens. *Appl. Microbiol. Biotechnol.* **56(1/2)**:9-16
- Munoz, M., Barrera, E., and Meza, I. (1981). El uso medicinal y alimenticio de plantas nativas y naturalizadas en Chile. *Museo National de Historia Natural Occasional Publication NO 33, :Santiago de Chile.*
- Muroi, H. and Kubo, I.(1996). Antibacterial activity of anacardic acid and total, alone and in combination with methicillin, against methicillin-resistant *Staphylococcus aureus*. *J.Appl. Bacteriol.* **80(4)**:387-94
- Nagarajan, K., Singh, D.K., Sharma, I. and Bodla, R.B. (2011). Activity guided fractionation and identification of fatty acid derivative from alcoholic extract of *Wrightia tomentosa*. *Der. Pharma. Chem.* **3(6)**:370-374
- Nagarajan, K., Mazumder, A and Ghosh, L.K. (2008). *In vitro* antioxidant activity of alcoholic extracts of *Wrightia tomentosa*. *Pharmacologyonline.* **1**:196-203
- Nagoshi, C., Shiota, S., Kuroda, T., Hatano, T., Yoshida, T., Kariyama, R. and Tsuchiya, T. (2006). Synergistic effect of [10]-gingerol and aminoglycosides against vancomycin-resistant enterococci (VRE). *Biol. Pharm. Bull.* **29**:44
- Naik, A.A., Hemavani, C. and Thippeswamy, B. (2012). Evaluation of antimicrobial property of *Spirogyra* species. *Int. Multidis. Res. J.* **2(2)**:13-15

- Najdenski, H.M., Gigova, L.G., Iliev I.I., Plamen., Pilarski, S., Lukavský , J., Iva, V.T. and Ninova, M.S. (2013). Antibacterial and antifungal activities of selected microalgae and cyanobacteria. *Int. J. Food Sc Tech.* **48** (7): 1533–1540
- Nakashima, T., Sekiguchi, T., Kuraoka, A., Fukushima, K., Shibata, Y., Komiyama, S. and Nishimoto, T. (1993). Molecular cloning of a human cDNA encoding a novel protein, DAD1, whose defect causes apoptotic cell death in hamster BHK21 cells. *Molec. Cell. Biol.* **3**:6367-6374
- Nascimento, S.C. Chiappeta, A and Lima, R.M. (1990). Antimicrobial and cytotoxic activities in plants from pernambuco. *Braz. Fitoter.* **61**:353-355
- Nash, T.H.III. and Wirth, V. (1988). Lichens, bryophytes and air quality. *Biblio. Lichenol.* **30**:231-267
- Nash, T.H.III. (1996). Lichen biology. Cambridge University Press, New York, USA
- Nash, T.H.III. and Gries, C. (2002). Lichens as bioindicators of sulfur dioxide. *Symbiosis.* **33(1)**:1-22
- Naveena, B.M., Sen, A.R., Kingsly, R.P., Singh, D.B. and Kondaiah, N. (2008). Antioxidant activity of pomegranate rind powder extract in cooked chicken patties. *Int. J. Food. Sci. Tech.* **43**:1807-1812
- Nedeljko, T., Perica, M., Vasiljevic, J., Gritsanapan, J., Supabphol, R and Manojlovic, I. (2010). Phytochemical and antioxidant studies of *Laurera benguelensis* growing in Thailand. *Bio. Res.* **43**: 169-176
- Negi, H.R. (2000). On the patterns of abundance and diversity of macrolichens of Chopta-Tunganath in the Garhwal Himalayas. *J. Biosci.* **25**:367-378
- Negi, H.R. and Kareem, K. (1996). Lichens: The unsung heroes. *Amrut.* **1**:3-6
- Nelsen, M.P., Lucking, R., Umana, L., Trest, M.T., Will, W.S., Chaves, J.L. and Gargas, A. (2007). *Multiclavula ichthyiformis* (Fungi: Basidiomycota: Cantharellales: Clavulinaceae), a remarkable new basidiolichen from Costa Rica. *Am. J. Bot.* **94**:1289–1296
- Ness, A.R. and Powles, J.W. (1997). Fruit and vegetables and cardiovascular disease: a review. *Int. J. Epidemiol.* **26**:1-13
- Newman, D.J., Cragg, G.M. and Snader, K.M. (2000). The influence of natural products upon drug discovery. *Nat. Prod. Reports.* **17**:175-285
- Nicollier, G., Tabacchi, R., Gavin, J., Breton, J.L. and Gonzales, A.G. (1979).

- Triterpenes de la 'mousse de chêne' (*Evernia prunastri* ((L.) Ach.). *Helvetica. Chimica. Acta.* **62**:807-810
- Nishitoba, Y., Nishimura, H., Nisshimaya, T. and Mijutami, J. (1987). Lichen acids, plant growth inhibitors from *Usnea longissima*. *Phytochem.* **26**:3181-3185
- Noaman, N.H., Khaleafa, A.F. and Zwky, S.H. (2004). Factors affecting antimicrobial activity of *Synechococcus leopoliensis*. *Microbiol. Res.* **56**:359-402
- Noriham, A., Babji, A.S. and Aminah, A. (2004). Determination of antioxidative activities of selected Malaysian plant extracts. *ASEAN Food. J.* **13**:193-199
- Nostro, A., Cellini, L., DiBartolomeo, S., Cannatelli, M.A., DiCampli, E., Procopio, F. and Grande, R. (2006). *Phytother. Res.* **20**:187
- Odabasoglu, F., Aslan, A., Cakir, A., Suleyman, H., Karagoz, Y., Halici, M. and Bayir, Y. (2004). Comparison of antioxidant activity and phenolic content of three lichen species. *Phytother. Res.* **18**:938-941
- Ogmundsdottir, H.M., Zoega, G.M., Gissurarson, S.R. and Ingolfsson, K. (1998). Anti-proliferative effects of lichen-derived inhibitors of 5-lipoxygenase on malignant cell lines and mytogen stimulated lymphocytes. *J. Pharm. Pharmacol.* **50**:107-115
- Oktay, M., Gulcin, I. and Kufrevioglu, O.I. (2003). Determination of *in vitro* antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *LWT. Food Sc. Technol.* **36**:263-271
- Okuyama, E., Umeyama, K., Yamazaki, M., Kinoshita, Y. and Yamamoto, Y. (1995). Usnic acid and diffractaic acid as analgesic and antipyretic components of *Usnea diffracta*. *Planta Med.* **61**:113-115
- Oliveira, I., Sousa, A., Ferreira, I.C., Bento, A., Estevinho, L and Pereira, J.A. (2008). Total phenols, antioxidant potential and antimicrobial activity of walnut (*Juglans regia* L.) green husks. *Food Chem. Toxicol.* **46(7)**:2326-2331
- Ordog, V., Stirk, W.A., Lenobel, R., Bancirova, M., Strnad, M., van Staden, J., Szigeti, J. and Nemeth, L. (2004). *J. Appl. Phyco.* **16(4)**: 309-314
- Ordon, Ez. A.A.L., Gomez, J.D., Vattuone, M.A. and Isla, M.I. (2006). Antioxidant activities of *Sechium edule* (Jacq). Swart extracts. *Food Chem.* **97**:452-458

- Orr, W.C. and Sohal, R.S. (1994). Extention of life, span by over expression of superoxide dismutase and catalase in *Drosophila melanogester*. *Science*. **263**:1128-1130
- Otzurk, S., Guvenc, S., Arikan, N, and Yylmaz, O. (1999). Effect of usnic acid on mitotic index in root tips of *Allium cepa* L. *Lagascalia*. **21(1)**: 47-52
- Owolabi, J., Omogbai, E.K.I. and Obasuyi, O. (2007). Antifungal and antibacterial activities of the ethanolic and aqueous extract of *Kigelia africana* (Bignoniaceae) stem bark. *Afr. J. Biotechnol.***6(14)**:882-8
- Oyaizu, M. (1986). Studies on products of browning reaction prepared from glucoseamine. *Jpn. J. Nutr.* **44**:307-314
- Parag. A.P. and Raman, B. (2013). Assessment of *Semecarpus anacardium* (Linn.f.) leaf methanolic extract for their antibacterial, antifungal and antioxidant activity. *Int. J. Pharma. and Pharmacol. Sc.* **5(1)**:201
- Patwardhan, B., Ashok, D., Vaidya, B. and Mukund, C. (2004). Ayurveda and natural products drug discovery. *Curr. Sci.* **86**:6
- Paudel, B., Bhattarai, H.D., Lee, J.S., Hong, S.G., Shin, S.W and Yim, J.H. (2008). Antibacterial potential of Antarctic lichens against human pathogenic Gram-positive Bacteria. *Phytother. Res.* **22**:1269-1271
- Paz, D.A., Raggio, J., Gomez-Serranillos, M.P., Palomino O.M., González, B., Carretero, M.E. and Crespo, A. (2010). HPLC isolation of antioxidant constituents from *Xanthoparmelia* sp. *J. Pharma. Biomed. Anal.* **53**:165–17
- Pejin, B., Lodice, C., Bogdanovic, G., Kojic, V. and Tesevic, V. (2013). Stictic acid inhibits cell growth on human colon adenocarcinoma HT-29 cells. *J. Arabic.* **10**:106
- Pereira, J.A., Oliveira, I., Sousa, A., Valentao, P., Andrade, P.B., Ferreira, I.C., Ferreres, F., Bento, A., Seabra, R. and Estevinho, L. (2007). Walnut(*Juglans regia* L.) leaves: Phenolic compounds, antibacterial activity and antioxidant potential of different cultivars, *Food Chem. Toxicol.* **45(11)**:2287-95
- Perez, C., Pauli, M. and Bazerque, P. (1990). An antibiotic assay by the agar-well diffusion method. *Acta. Biol. Med. Exp.* **15**:113-15
- Periasamy, K. (2005). Novel antibacterial compounds obtained from some edible mushrooms. *Int. J. Med. Mushrooms.* **7**:443-444

- Perry, N.B., Benn, M.H., Brennan, N.J., Burgess, E.J., Ellis, G., Galloway, D.J., Lorimer, S.D. and Tangney, R.S. (1999). Antimicrobial, antiviral and cytotoxic activity of New Zealand lichens. *Lichenologist*. **31**:627-636
- Pietta, P., Simonetti, P. and Mauri, P. (1998). Antioxidant activity of selected medicinal plants. *J. Agric. Food Chem.* **46**:4487-4490
- Piria.(1938). On new products extracted from salicine).*Comptes rendus* **6**: 620–624.
- Prashanthkumar, P., Angadi, S.B. and Vidyasagar, G.M.(2006) Antimicrobial activity of blue green algae and green algae. *Indian. J. Pharm. Sci.* **88(6)**: 647-648
- Pramoda, K.J., Mastan, A. and Sreedevi, B. (2014). Evaluation of the *in vitro* antioxidant and antibacterial activities of secondary metabolites produced from lichens. *Asian J. Pharm. Clin. Res.* **7(1)**:193-198
- Preeti, S. Babiah., Upreti, D.K and John, S.A. (2014). An *in vitro* analysis of antifungal potential of lichen species *Parmotrema reticulatum* against phytopathogenic fungi. *Int. J. Curr. Microbiol. App.Sci*, **3(12)**:511-518
- Prieto, P., Pineda, M. and Aguilar, M. (1999). Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *Anal. Biochem.* **269**:337-341
- Proska, B., Sturdikova, M., Pronayova, N. and Liptaj, T. (1996). (D)-Usnic acid and its derivatives. Their inhibition of fungal growth and enzyme activity. *Pharmazie.* **51**:195-196
- Purohit, S.S. and Mathur, S.K. (1999). Drugs in Biotechnology fundamentals and applications. S.S Purohit, Maximillan Publishers, India. p. 576
- Purushotham, K.G., Arun, P., Jayarani, J. J., Vasanthakumari, R., Sankar, P. and Reddy, R.B. (2010). Synergistic *in vitro* antibacterial activity of *Tectona grandis* leaves with tetracycline. *Int. J. Pharm. Tech. Res.* **1**:519-523
- Qi, H., Zhang, Q., Zhao, T., Hu, R., Zhang, K. and Li, Z. (2006). *Bioorg. Med. Chem. Lett.* **16 (9)**: 2441-2445
- Rahman, M.H., Alam, M.B., Hossain, S., Jha, M.K. and Islam, A. (2011b). Antioxidant, analgesic and toxic potentiality of methanolic extract of *Stephania japonica* (thunb.) miers. leaf, *Asian J. Pharm. Clin. Res.* **4(3)**:38-41

- Rahman, S.M.A., Abd-Ellatif, S.A., Deraz S .F and Khalil A.A. (2011a). Antibacterial activity of some wild medicinal plants collected from western Mediterranean coast, Egypt. Natural alternatives for infectious disease treatment. *African J .Biotech.* **10**(52):10733-10743
- Rai, P.C., Sarkar, A. Bhujel, R.B and Das, A.P. (1998). Ethnomedicinal studies in some fringe areas of Sikkim and Darjeeling Himalaya. *J. Hill Res.* **11**:12-21
- Rajkumar, V., Guha, G., Kumar, R.A. and Mathew, L. (2010). Evaluation of antioxidant activities of *Bergenia ciliata* Rhizome. *Rec. Nat. Prod.* **4**(1):38-48
- Ramtin, M., Massiha, A., Khoshkholgh-Pahlaviani M.R.M., Issazadeh, K., Assmar, M. and Zarrabi, S. (2014). Evaluation of the antibacterial activities of essential oils of *Iris pseudacorus* and *Urtica dioica*. *Zahedan .J. Res. Med .Sci.* **16**(3):35-39
- Rankovic, B.R., Kosanic, M.M. and Stanojkovic, T.P. (2011). Antioxidant, antimicrobial and anticancer activity of the lichens *Cladonia furcata*, *Lecanora atra* and *Lecanora muralis*. *BMC Complement. Alternat. Med.* **11**(97):1-8
- Rankovic, B., D. Rankovic, M. Kosanic and D, Maric. (2010). Antioxidant and antimicrobial properties of the lichen *Anaptychya ciliaris*, *Nephroma parile*, *Ochrolechia tartarea* and *Parmelia centrifuga*. *Cent. Eur. J. Biol*, **5**:649-655
- Rankovic, B., M. Misjic and S. Sukdolak. (2007). Antimicrobial activity of some lichens and their components. *J. Microbiol.* 279-286
- Rankovic, B., Misic, M. and Sukdolak, S. (2007). Antimicrobial activity of the lichens *Lasallia pustulata*, *Parmelia sulcata*, *Umbilicaria crustulosa* and *Umbilicaria cylindrica*. *Mikrobiologya.* **76**(6):817-821
- Rankovic, B., Misjic M. and Sukdolak, S. (2007). Evaluation of antimicrobial activity of the Lichens *Lasallia pustulata*, *Parmelia sulcata*, *Umbilicaria crustulosa* and *Umbilicaria cylindrica* *J. Microbiol.* **76**:723-727
- Rankovic, B., Rankovic, D., Kosanic, M and Maric, D. (2010). Antioxidant and antimicrobial properties of the lichens *Anaptychya ciliaris*, *Nephroma parile*, *Ochrolechia tartarea* and *Parmelia centrifuga*. *Cent. Eur. J. Biol.***5**:649-655

- Ray, S, Sinhababu, A. and Mandal, N.C. (2003). Antibacterial activity of three lichen specimen viz., *Usnea articulate*, *Ramalina jamesii*, and *Parmelia tinctorum* from the eastern Himalaya. *J. Hill Res.* **16(2)**:66-69
- Revathy, M., Sathya, S.S., Manimekala, N., Annadurai, G. and Ahila, M. (2015). Preliminary phytochemical investigation and antibacterial effects of lichen *Parmotrema perlatum* against human pathogens *European J. Biomed. Pharma. Sci.* **2 (4)**:336-347
- Rice-Evans, C.A, Miller, N.J., Bolwell, P.G, Bramley, P.M and Pridham, J.B. (1995). The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free. Radical Res.* **23**:375-383
- Rice-Evans, C., Miller, N. and Paganga, G. (1997). Antioxidant properties of phenolic compounds, **2(4)**:152-159
- Richardson, D.H.S. (1991). Lichen and man. *In: D. L. Hawksworth (Ed.). Frontiers in Mycology.*, CAB International, Wallingford pp. 187-210
- Richardson, D.H.S. (1988). Medicinal and other economic aspects of lichens. *In: M. Galun (Ed.). CRC Handbook of Lichenology*, CRC Press, Boca Raton, Florida. **1**:93-108
- Richardson, D.H.S. (1992). Pollution monitoring with lichens. *Naturalists' Handbooks* 19. Richmond Publishing Co., Ltd. Slough, England. 76
- Rios, J.L. and Recio, M.C. (2005). Medicinal plants and antimicrobial activity. *J. Ethnopharmacol.* **100**:1-2, 80-84
- Romagni, J.G., Meazza, G., Nanayakkara, N.P.D. and Dayan, F.E. (2000). The phytotoxic lichen metabolite, usnic acid, is a potent inhibitor of plant phydroxyphenylpyruvate dioxygenase. *FEBS Lett.* **480**:301-305
- Romagni, J.G. and Dayan, F.E. (2002). Structural diversity of lichen metabolites and their potential use. *In: R.K Updhyay (Ed.). Advances in Microbial Toxin Research and its Biotechnological Exploitation.* Kluwer Academic, Plenum Publishers, New York. pp.151-169
- Roman, M., Hradkova, I., Filip, V. and Smidrkal, J. (2010). Antimicrobial and antioxidant properties of phenolic acids alkyl esters, *Czech. J. Food Sci.* **28(4)**:275-279
- Rowe, J.G. (1999). Some lichen products have antimicrobial activity and identification of lichenic substances in some lichens from Southern Spain. *Ann. Pharm. Fr.* **49(5)**:278-285

- Rowe, J.G., Saez, M.T. and Garcia, M.D. (1989). Contribution à l'étude de l'activité antibactérienne de quelques lichens du sud de l'Espagne. *Annales pharmaceutique française*. **47**: 89-94
- Russo, A., Piovano, M., Lombardo, L., Garbarino, J. and Cardile, V. (2008). Lichen metabolites prevent UV light and nitric oxide-mediated plasmid DNA damage and induce apoptosis in human melanoma cells. *Life. Sci*. **83**:468-474
- Russo, A., Piovano, M., Lombardo, L., Vanella, L., Cardile, V. and Garbarino, J. (2006). Pannarin inhibits cell growth and induces cell death in human prostate carcinoma DU-145 cells. *Anticancer Drugs*. **17**:1163-1169
- Saklani, A. and Upreti, D.K. (1992). Folk uses of some lichens in Sikkim Himalaya. *J. Hill Res*. **11**:12-21
- Samuelsson, G. (2004). Drugs of Natural Origin, a Textbook of Pharmacognosy .5th ed. *Swedish Pharmaceutical Press, Stockholm*
- Santesson, J. (1970). Chemical studies on lichens. 30. Anthraquinoid pigments of *Trypetheliopsis boninensis* and *Ocellularia domingensis*. *Acta Chemica Scandanavica*. **24**:3331-3334
- Santos, P, Mondragon A. (1969). Studies on the Philippine lichens, II thin-layer chromatographic study of the constituents of some lichen species. *Philipp. J. Sci*. **98**:297-303
- Santiago, K.A.A., Jayne., Nicholei, C., Borricano, J, N. C., Canal, J.N., Marcelo, D.M.A., Perez, M.C.P and Cruz, T.E.E.D. (2010). Antibacterial activities of fruticose lichens collected from selected sites Luzon Island. *Philippine Sci. Lett*. **3 (2)**:18-29
- Sasikumar, J.M., Mathew, G.M. and Teepica, P.D.D. (2010). Comparative studies on antioxidant activity of methanol extract and flavonoid fraction of *Nyctanthes arbortristis* leaves. *EJEAF Che*. **9**:227-233
- Sathisha, A.D., Lingaraju, H.B. and Prasad, K.S. (2011). Evaluation of antioxidant activity of medicinal plant extracts produced for commercial purpose. *E-Journal Chem*. **8(2)**:882-886
- Sawa, T., Nakao, M., Akaike, T., Ono, K. and Maeda, H. (1999). Alkylperoxyl radical scavenging activity of various flavonoids and other phenolic compounds: Implications for the antitumor promoter effect of vegetables. *J. Agric. Food Chem*. **47**:397-492

- Saxena, G., McCutcheon, A.R., Farmer, S., Towers, G.H.N. and Hancock, R.E.W. (1994). Antimicrobial constituents of *Rhus glabra*. *J. Ethnopharm.* **42**:95–99
- Schinazi, R.F., Chu, C.K., Baby, J.R., Oswald, B.J., Saalman, D.L. and Nasai, M. (1990). Anthraquinones as a new class of antiviral agents against Human Immunodeficiency Virus. *Antiviral. Res.* **13**:265-272
- Schmeda, H., Tapia, G., Lima, A., Pertino, B., Sortino, M., Zacchino, S., Arias, D.R. and Feresin, A.G.E. (2008). A new antifungal and antiprotozoal depside from the Andean lichen *Protousnea poeppigii*. *Phytother. Res.* **22(3)**:349-355
- Schumm, F.(2002). Dunnschichtchromatogramme auch furr den Amateur moglich. *Aktuelle Lichenologische. Mitteilungen.* **9**: 8-22
- Sco, H.G., Takata, I., Nakamura, M., Tatsumi, H., Suzuki, K., Juji, J. and Taniguchi, N. (1995). Induction of nitric oxide synthase and concomitant suppression of superoxide dismutase in experimental colitis in rats. *Archs. Biochem. Biophys.* **324**:41-47
- Syvacy, A. and Sokmen, M. (2006). Seasonal changes in antioxidant activity, total phenolic andanthocyanin constituent of the stems of two *Morus* species (*Morus alba* L. and *Morus nigra* L.). *Plant Growth Regulation.* **44(5)**:251-256
- Seaman, T., Campbell, W., Lategan, C. and Smith, P. (2007). The antimicrobial activity of South African lichens and lichen-derived usnic acid. *Planta Med.* **73**:137
- Seaward, M.R.D. (1997). Major impacts made by lichens in biodeterioration processes. *Int. J. Biodeterior. Biodegrad.* **40**:269-273
- Sezik, E., Yesilada, F., Tabata, M., Honda, G., Takaishi, Y., Fujita, T., Tanaka, T. and Takeda, Y. (1997). Traditional medicine in Turkey VIII. Folk medicine in East Anatolia Erzurum Erzincan Aıgry, Kars, Ijgdyr provinces. *Economic Bot.* **51**:195–211
- Shahidi, F. and Wanasundara, P.K.J.P.D. (1992). Phenolic antioxidants. *Crit. Rev. Food Sci. Nutr.* **32**:67-103
- Shanmugam, H., Anil, K.K. and Mukesh, D. (2008). Synergism between naturalproducts and antibiotics against infectious diseases, *Phytomedicine.* **15**: 639-652

- Sharma, A.K., Sharma, M.C. and Dobhal, M.P. (2012). Phytochemical investigation of therapeutic important lichen *Parmelia perlata*. *J. Nat. Prod. Plant Resour.* **2(1)**:101-106
- Sharnoff, S.D. (1997). Lichens and People. www.lichen.com/people.html.
- Shimada, K., Fujikawa, K., Yahara, K. and Nakamura, T. (1992). Antioxidative properties of xanthan on the autoxidation of soybean oil in cyclodextrin emulsion. *J. Agric. Food Chem.* **40 (6)**, pp 945-948
- Shrestha, G., Raphael, J., Leavitt, D.S., and Clair, L.L. St. (2014). *In vitro* evaluation of the antibacterial activity of extracts from 34 species of North American lichens. *Pharma. Biol.* **52 (10)**:1262-1266
- Sibanda, T. and Okoh, A.I. (2007). The challenges of overcoming antibiotic resistance: Plant extracts as potential sources of antimicrobial and modifying agents. *Afr. J. Biotechnol.* **6**: 2886-2896
- Sibanda, T. and Okoh, A.I. (2008). In vitro evaluation of interaction between acetone extracts of *Garcinia kola* seeds and some antibiotics. *Afr. J. Biotechnol.* **7**:1672-1678
- Siddhuraju, P. and Becker, K. (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *J. Agric. Food. Chem.* **51**:2144-2155
- Sies, H. (1993). Strategies of antioxidant defense. *Eur. J. Biochem.* **215**:213-219
- Siew, S., Yeo, M. and Yes, F. (2012). Antiquorum sensing and antimicrobial activities of some traditional chinese medicinal plants commonly used in Southeast Asia. *Malaysian J. Microbiol.* **8(1)**:11-20
- Sisodia, R., Geol, M., Verma, S., Rani, A. and Dureja, P. (2013). Antibacterial and antioxidant activity of lichen species *Ramalina roesleri*. *Nat. Prod. Res.* **27(23)**:2235-9
- Slack, N.G. (1998). *Bib. Lichen.* **30**:6(9):23-53
- Slinkard, K and Slingleton, V.L. (1997). Total phenolic analyses: automation and comparison with manual method, *Am. J. Enol. Viticult.* **28**:49-55
- Soctanand, K.O. and Aiyelaagbe, O.O. (2009). The need for bioactivity-safety evaluation and conservation of medicinal plants-A review. *J. Med. Pl. Res.* **3(5)**:324-328

- Souri, E., Amin, G., Farsam, H., Jalalizadeh, H. and Barezi, S. (2008). Screening of thirteen medicinal plant extracts for antioxidant activity. *Iran. J. Pharm. Res.* **7**:149-154
- Squadriato, G. and Pelor, W.A. (1998). Free radical oxidative chemistry of nitric oxide: The roles of superoxide, peroxyxynitrite and carbon dioxide. *Biol. Med.* **25**:392-403
- Srivastava, P., Upreti, D.K., Dhole, T.N.K., Srivastava, K.A and Nayak. T.M. (2013). Antimicrobial Property of Extracts of Indian Lichen against Human Pathogenic Bacteria. *Mycosphere.* **4(4)**734-743
- Stefanovi, D., Olgica, D., Stanojevi, D. and Jiljana, R.O. (2012). Synergistic antibacterial activity of *Salvia officinalis* and *Cichorium intybus* extracts and antibiotics. *Aacta Poloniae Pharmaceutica Drug Research.* **69(3)**:457-463
- Steglich, W and Jedtke, K.F. (1976). Neue Anthachinonfarbstoffe aus *Solorina crocea*. *Z. Naturforsch.* **31C**:197-198
- Stolte, K., D. Mangis., R. Doty and K. Tonnessen. (1993). *Lichens as Bioindicators of Air Quality*. USDA-Forest Service, Rocky Mountain Forest and Range Experiment Station General Technical Report RM-224. Fort Collins, Colorado.131
- Stubler, D. and Buchenauer, H. (1996). Antiviral activity of the glucan lichenan (poly-β{1 → 3, 1 → 4} Danhydroglucose): Biological activity in tobacco plants. *J. Phytopathol.* **144**: 37-43
- Subramaniyan, V., Vijayakumar, M., Pitchai. D., Jeyacha, S. and Chockaiya Manoharan. (2011). Antimicrobial Activity of Cyanobacteria Isolated from Freshwater Lake. *Int. J. Microbiol. Res.* **2** (3): 213-216.
- Sydiskis, R.J., Owen, D.G., Lohr, J.L., Rosle, K.H.A. and Blomster, R.N. (1991). Inactivation of enveloped viruses by anthraquinones extracted from plants. *Antimicrob. Agents Chemother.* **35**:2463-2466
- Syvacy, A. and Sokmen, M.(2006). Seasonal changes in antioxidant activity, total phenolic and anthocyanin constituent of the stems of two *Morus* species (*Morus alba* L. and *Morus nigra* L.). *Plant Growth Regulation.* **44(5)**: 251-256
- Takahashi, R., Tanaka, O. and Shibata, S. (1970). The structure of phlebic acid B, a constituent of the lichen *Peltigera aphthosa*, and the

- occurrence of of 15 α -acetoxy- and β -acetoxy-22-hydroxyhopane in *P. dolichorrhiza*. *Phytochem.* **9**:2037-2040
- Takeda, T., Funatsu, M., Shibata, S. and Fukuoka, F. (1972). Lichen polysaccharides having antitumor activity. *Chem. Pharmacol. Bull.* **20**:2445
- Tepe, B., Donmez, E., Unlu, M., Candan, F., Daferera, D., Vardar-Unlu G., Polissiou, M. and Sokmen, A. (2004). Antimicrobial and antioxidative activities of the essential oils and methanol extracts of *Salvia cryptantha* (Montbret *et. Aucher ex Benth.*) and *Salvia multicaulis* (Vahl). *Food Chem.* **84(4)**:519-525
- Thabrew, M.I., Hughes, R.D. and Mc Farlane, I.G. (1998). Antioxidant activity of *Osbeckia aspera*. *Phytother. Res.* **12**:288-290
- Thatte, U., Bagadey, S. and Dhanukar, S. (2000). Modulation of programmed cell death by medicinal plants. *Mol. Cell. Biochem.* **46**:199-214
- Thippeswamy, B., Sushma, N.R. and Naveen Kumar, K.J. (2012). Antimicrobial property of bioactive factor isolated from *Parmelia perlata*. *Int. Multidiscipl. Res. J.* **2(2)**:01-05
- Thota, P., Bhogavalli, P.K., Bathula N.R.V., Gangula. S.R and Chenreddy, S.K. (2012). *In vitro* studies on antimicrobial screening of leaf extracts of *Sapindus saponaria* against common dental pathogens. *Plant. Sci. Feed.* **2 (2)**: 15- 18
- Tiwari, P., Himanshu, R., Upreti. D.K., Trivedi, S. and Shukla. P. (2011). Antifungal activity of a common Himalayan foliose lichen *Parmotrema tinctorum* (Despr. ex Nyl.) Hale. *Nat. Sc.* **9(9)**:167-171
- Tiwari, R.P., Bharti, S.K., Kaur, H.D., Dikshit R.P. and Hoondal, G.S. (2005). Synergistic antimicrobial activity of tea and antibiotics. *Indian J. Med. Res.* **122**:80-84
- Tolulope, O. (2007). Cytotoxicity and antibacterial activity of methanolic extract of *Hibiscus sabdariffa*. *J. Med. Plant. Res.* **1(1)**:009-013
- Trease, G.E. and Evans, W.C. (1978). *Pharmacology* .11th ed. Bailliere Tindall Ltd.London. 60-75
- Triggiani, D., Ceccarelli, D., Tiezzi, A., Pisani, T., Munzi, S., Gaggi, C. and Loppi, S. (2009). Antiproliferative activity of lichen extracts on murine myeloma cells. *Biologia.* **64**:59-62

- Turk, O.A., Yilmaz, M., Kivanc, M. and Turk, H. (2003). The antimicrobial activity of extracts of the lichen *Cetraria aculeata* and its protolichesterinic acid constituent. *Z. Naturforsch.* **58c**:850-854
- Turk, H., Kivnac, M., Turk, O.A. and Tay, T. (2004). The antimicrobial activity of extracts of the lichen *Cladonia foliacea* and its (-)-Usnic Acid, Atranorin, and Fumarprotocetraric Acid constituents. *Z. Naturforsch.* **59c**:249-254
- Turkoglu, S., Turkoglu, I., Kahyaoglu, M. and Celyk, S. (2010). Determination of antimicrobial and antioxidant activities of Turkish endemic *Ajuga chamaepitys* L.Schreber subsp. *Euphratica* P.H. Davis (Lamiaceae), *J. Med. Pl. Res.* **4(3)**:1260-1268
- Upreti, D.K. (1996). *Environ. Conserv.* **22**:362-363
- Ullah, M.O., Sultana, S., Haque, A. and Tasmin, S. (2009). Antimicrobial, cytotoxic and antioxidant activity of *Centella asiatica*. *Eur. J. Sci. Res.* **30(2)**:260-264
- Vartia, K.O. (1973). Antibiotics in lichens. *In*: V. Ahmadjian and M. E. Hale (eds.). *The Lichens* Academic Press, New York. 547-561
- Velioglu, Y.S., Mazza, G., Gao, L. and Oomah, B.D. (1998). Antioxidant activity and total phenolics in selected fruits, vegetables and grain products. *J. Agric. Food Chem.* **46**:4113-4117
- Vijaya, B.S.G. and Prabu, V.A. (2010). Antibacterial activity of cyanobacterial species from Adirampattinam Coast, South East Coast of Palk Bay. *Curr. Res. J. Biosci.* **2(1)**: 24-26
- Vinson, J. A., Yong, H., Xuchui, S. and Zubik, L. (1998). Phenol antioxidant quantity and quality in foods: vegetables. *J. Agric. Food. Chem.* **46**: 3630-3634
- Vries, F.A., Bitar, H.E., Green I.R., Klaasen, J.A., Mabusela, W.T., Bodo B. and Johnson, Q.(2005). An antifungal active extract from the aerial parts of *Galenia africana*. 11th Natural Products Research Network for Eastern and Central Africa.
- Wang, H., Nair, M.G., Straburg, G.M., Booren, A.M. and Gray, J.I. (1999). Antioxidant polyphenols from tart cherries (*Prunus cerasus*). *J. Agric. Food. Chem.* **47**:840-844
- Wang, L.F. and Zhang, H.Y. (2003). A theoretical investigation on DPPH radical scavenging mechanism of edaravone. *Bioorg. Med. Chem. Lett.* **13**:3789-3792

- Wang, H. and James, A.J. (1999). Quantifying cellular oxidative stress by dichlorofluorescein assay using microplate reader. *Free Radic. Biol. Med.* **27(5/6)**:612–616
- Wang, H. and NgT, B. (2004). Eryngin, a novel antifungal peptide from fruiting bodies of the edible mushroom *Pleurotus eryngii*. *Peptides*. **25**:1-5
- Wangensteen, H., Samuelsen, A.B. and Malterud, K.E. (2004). Antioxidant activity in extracts from coriander. *Food .Chem.* **88**:293-297
- Wasser, S.P. (2002) .Medicinal mushrooms as a source of antitumor and immunomodulating polysaccharides. *Appl. Microbiol. Biotechnol.* **60**: 258-274
- Weckesser, S., Engel, K., Simon-Haarhaus, B., Wittmer, A., Pelz, K. and Schempp, C.M. (2007). Screening of plant extracts for antimicrobial activity against bacteria and yeasts with dermatological relevance. *Phytomed.* **14**:508-516
- Weissman, L., Fraiberg, M., Shine, L., Garry, J. and Hochman, A. (2006). Responses of antioxidants in the lichen *Ramalina lacera* may serve as an early-warning bioindicator system for the detection of air pollution stress. *FEMS Microbiol. Ecol.* **58**:41–53
- Weissman, L., Garty, J. and Hochman, A. (2005). Characterization of enzymatic antioxidants in the lichen *Ramalina lacera* and their response to rehydration. *Appl .Environ. Microbiol.* **71**:6508–6514
- Williams, D.E.K., Bombuwala, E., Lobkovsky, E., Dilip, deSilva., Karunaratne, V., Allen, T.M., Clardy, J. and Anderson, R.J. (1998). Antineoplastic epidithiapiperazinediones isolated from the lichen *Usnea* sp. *Tetrahedron Lett.* **39**:9579-9582
- Wiseman, H. and Halliwell B. (1996). Damage to DNA by reactive oxygen and nitrogen species: role in inflammatory disease and progression to cancer. *Biochem. J.* **313**:17-29
- Witztum, J. L. (1994). Oxidative hypothesis of atherosclerosis. *Lancet.* **344**: 793-795
- Yagi, S., Chretien, F., Duval, R.E., Fontanay, S., Maldini, M., Piacente, S., Henry, M., Chapleur, Y. and Laurain-Mattar, D. (2012). Antibacterial activity, cytotoxicity and chemical constituents of *Hydnora johannis* roots. *South African. J. Bot.* **78**: 228-34

- Yamamoto, Y., Miura, Y., Kinoshita, Y., Higuchi, M., Yamada, Y., Murakami A., Ohigashi, H. and Koshimizu, K. (1995). Screening of tissue cultures and thalli of lichens and some of their active constituents for inhibition of tumor promoter-induced Epstein-Barr virus activation, *Chem. Pharm. Bull.* **43**:1388-1390
- Yamamoto, Y. (1995). Usnic acid and diffractaic acid as analgesic and antipyretic components of *Usnea diffracta*. *Planta Med.* **61**:113-115
- Yang, D.M., Takeda, N., Iitaka, Y., Sankawa, U. and Shibata, S. (1973). The structure of eumitrins A1, A2 and B. The yellow pigments of the lichen *Usnea baileyi* (Stirt.) Zahlbr. *Tetrahedron.* **29**: 518-529
- Yang, Y. and Anderson, E.J. (1999). Antimicrobial activity of a porcine myeloperoxidase against plant pathogenic bacteria and fungi. *J. Appl. Microbiol.* **86(2)**:211-220
- Yasar, S., Sagdic, O. and Kisioglu, A. (2004). *In vitro* antibacterial effects of single or combined plant extracts. **3(1)**: 39-43
- Yesilada, E., Honda, G., Sezik, E., Tabata, M., Goto, K. and Ikeshiro, Y. (1993). Traditional medicine in Turkey IV. Folk medicine in the Mediterranean subdivision. *J. Ethnopharmacol.* **39**:31-38
- Yesilada, E., Sezik, E., Honda, G., Takaishi, Y., Takeda, Y. and Tanaka, T. (2001). Traditional medicine in Turkey X. Folk medicine in Central Anatolia. *J. Ethnopharmacol.* **75**:95-115
- Yen, G.C. and Hsieh, C.L. (1998). Antioxidant activity of extracts from *Du-Zhong* (*Eucommia ulmoides*) toward various lipid peroxidation *in vitro*. *J. Agric. Food Chem.* **46**:3952-3957
- Yilam, M. (2005). The antimicrobial activity of extracts of lichen *Hypogymnia tubulosa* and its 3-hydroxy physodic acid constituent. *Z. Naturforsch.* **60c**:35-38
- Yilmaz, M., Ozdemir, T.A., Tay. T. and Kivanc, M. (2004). The antimicrobial activity of extracts of the lichen *Cladonia foliacea* and its (+)-Usnic Acid, Atranorin and Fumarprotocetraric Acid Constituents. *Z. Naturforsch.* **59c**:249-254
- Yoshioka, T., Fujii, E., Endo, M., Wada, K., Tokunaga, Y., Shiba, N., Hohsho, H., Shibuya, H. and Muraki, T. (1998). Antiinflammatory potency of dehydrocurdione, a zedoary derived sesquiterpene *Inflammation Res.* **47**:476-481

- Yosioka, I., Morita, Y. and Ebihara, K. (1970). The structure of constictic acid. *Chem. Pharm. Bull.* (Tokyo). **18**:2364-2366
- Yosioka, I., Yamauchi, H. and Kitagawa, I. (1972). Lichen triterpenoids. V: On the neutral triterpenes of *Pyxine endochrysin* Nyl *Chem. Pharm. Bulletin* (Tokyo). **20**:502-513
- Yu, B.P. (1994). Cellular defenses against damage from reactive oxygen species. *Physiological Rev.* **76**:139-162
- Zainol, M.K., Abdul-Hamid, A., Yusof, S. and Muse, R. (2003). Antioxidative activity and total phenolic compounds of leaf, root and petiole of four accessions of *Centella asiatica* (L.) Urban. *Food. Chem.* **49**:5165-5170
- Zenkov, N.K. and Menshikova. E.B. (1993). *Uspekhi sovremennoy biologii.* **113(3)**: 289-296
- Zhang, W.M., Li, B., Han, L. and Zhang, H.D. (2009). Antioxidant activities of extracts from areca (*Areca catechu* L.) flower, husk and seed. *Afr. J. Biotechnol.* **8**:3887-3892