

## Investigation of Fungal Spores of Tomato field at District Beed, M.S., India

Kshirsagar J. J.

Department of Botany, Vasantdada Patil College, Patoda, Dist.Beed. (M.S)  
Jyotikshirsagar1974@gmail.com

### Article Info

Received: 10-05-2021,

Revised: 22-06-2021,

Accepted: 30-07-2021

**Keywords:** Tomato (Solanum lycopersicum) field, Tilak air sampler, fungal spores.

### Abstract

Fungal survey was carried out in the Tomato field (*Solanum lycopersicum*) for a period of Rabbi Seasons from 10<sup>th</sup> July 2019 to 12<sup>th</sup> October 2019. For trapping the fungal spores, Tilak air sampler was used. During the investigation the aeromicroflora population includes large number of fungal spores, pollen grains, insect parts etc. The result showed incidence of varieties of fungal spores in the environment. In this investigation 62 spores were identified during the period of survey. In this seasons the most dominant spores were *Cladosporium* (26.44%), *Alternaria* (10.64%), *Cercospora* (1.96%), *Curvularia* (8.24%), *Helminthosporium* (7.98%), *Periconia* (2.11%), *Pithomyces* (2.76%), *Rust* spores (6.15%), *Nigrospora* (8.32%), *Smut* spore (5.94%), *Rhizopus* (3.96%).

### INTRODUCTION

The aerobiological studies are recent origin in India. In Maharashtra and Marathwada credit for developing the aerobiological research work goes to prof. Tilak S.T. Very few crops have been investigated so far. In Marathwada region, the climate is relatively moderate, average rainfall is 650 mm in monsoon. Temperature ranges from 20<sup>o</sup>c to 38<sup>o</sup>c, relative humidity varies from 30 to 70 % .For effective management of crop diseases, it is desirable to study the prevalence of air spora in this region. This is achieved by aerobiological study. Hence this observation could be helpful for the treatment of diseases (allergic as well as agriculture).

Crop diseases caused by airborne mycosporophytes constitute another important aspect of agriculture. Our agriculture crops, however continuously influence from various diseases, out of which fungal diseases are dominant in this region. In a study of airspora of Tomato fields, observed different types. Among them the *Alternaria*, *Cladosporium*, *Cercospora*, *Curvularia*,

*Rust* spores, *Helminthosporium*, *Periconia*, *Rhizopus*, *Nigrospora* , hyphal fragments, Pollen grains and insect parts were dominant ones. In view of the above facts qualitative and quantitative airborne spores was worked out.

### MATERIAL AND METHODS

In the present study, Tilak Air sampler was implemented to find out the availability of casual microbes of blight and leaf spot diseases in the Tomato field of 5 acres of land area. Tilak air sampler is an electrically operated machine which runs on electric power supply of (AC 230 V) & provides a continuous air sampling data for eight days. Sampler was kept with its orifice at constant height of 1 meter above the ground in the Tomato field. The air was sampled at the rate of 5 liters for minute & the transparent cellophane tape was fixed on the drum, coated uniformly with white petroleum jelly as adhesive. These cellophane brought to the laboratory, slides were made and scanned. Fungal spores isolation was made from these slides over Tomato Field.

### Scanning

Loaded tape on each slide was divided into six equal divisions by marking it over cover slip with a pointed ball pen. Each division representing two hours air sampling. Scanning of slides was carried out under the binocular research microscope using 10X X 45 x magnification, as per the procedure mentioned by (Tilak and Kulkarni, 1970). The identification of fungal spore type was made on the basis of size, shape septation of spores using standard keys and available authentic literature.


### Statistical Analysis

The total spores counted per day. The counted spores were multiplied by conversion factor 14 of Tilak Air Sampler.

### RESULTS AND DISCUSSION

Total 61 spores of different fungal spores were noted in August 2019 month. The abundant spores observed in the month of August 2019 were *Alternaria* (6314), *Nigrospora* (1428), *Cercospora* (490) *Curvularia* (1372), *Smut spores*


(1414), *Cladosporium* (12460), *Helminthosporium* (1316), *Rust spore* (1232), *Pithomyces* (630), *Periconia* (574), *Rhizopus* (770). The Figure below revealed dominant spores found in August 2019 month. In the month of September 2019, total 62 different spores were observed. The abundant spores found in September 2019 were *Rust spore* (6.15%), *Alternaria* (10.64%), *Nigrospora* (8.32%), *Curvularia* (8.24%), *Smut spores* (5.94%), *Cladosporium* (26.44%), *Cercospora* (1.96%), *Helminthosporium* (7.98%), *Rhizopus* (3.96%), *Periconia* (2.11%). The dominant spores types found in the month of September 2019 are shown in figure below. And most were *Alternaria*, *Cladosporium*, *Nigrospora*, *Rhizopus* and smut spores. *Cladosporium* as an allergen was at the top most in concentration and percentage contribution. Agarwal and Shivpuri (1974) reported role of *Cladosporium* bioaerosols in etiology of respiratory allergic disorders. *Alternaria*, *Curvularia*, *Periconia*, *Helminthosporium* and *Nigrospora* are known to be potentially allergenic. Allergenic diseases due to *Aspergilla* and *Penicillin* are recorded by Singh & Singh (1994).



In most of aeromycological survey, *Cladosporium* was as one of the most abundant aerospora reported all over world (oliveira *et al.*, 2007). The abundance of *Cladosporium* throughout the year may be attributed to the structural features of the spores such as small size and smooth wall which favors and facilitate the transport of airborne spores.

During the present investigation the 3 fungal spores belonged to other types viz. hyphal

fragments, Pollen grains and insect parts. These three types are well known aeroallergens responsible for nasobronchial allergy, respiratory allergy and cause allergenic reactions (Nair 1978, Kulkarni 1981, Shivpuri 1980). This study points out the prevalence of large percentage of aeroallergens which may be responsible for inducing allergenic reactions to sensitive individuals.



This investigations carried out indicates the significant allergenic nature of *Rhizopus*, *Cheotomium*, *Pleaspora*, *Alternaria*, *Aspergilles*, *Cladosporium*, *Curvularia*, *Epicoccum*, *Cercospora*, *Nigrospora*, *Helminthosporium*, *Heterosporium* and hyphal fragments (Tilak 1989). In the India significant allergenic fungi are *Curvularia*, *Alternaria*, *Helminthosporium*, *Cladosporium*, *Aspergillus* and *Rhizopus* (Shivpuri 1982).

A variation in the temperature, humidity, rainfall and wind was noted during the investigation period. *Cladosporium* species lives as sporophyte or parasite on many kinds of plants. Dry spores produced in excessive quantities can be transported over wide areas and during rainy season its concentration was low (Ebner *et al.*, 1989).

In European countries, *Alternaria* varies between 20,000-30000 spores/year (Oliveira *et al.*, 2007) to more than 200,000, only exceeding the levels of 300000 spores quoted for the north-western Iberian Peninsula in some areas (Mediavilla *et al.*, 1997). In several Italian cities, high quantities of *Cladosporium* and *Alternaria* are found from May to October, reaching their maximum levels in September (Zanca, 2003). However, in areas at lower latitudes where precipitation and humidity are limiting factors, but not temperature, the spores increase in the months before and after summer (Manoharachary *et al.*, 2005).

The month wise percentage contribution of each spore group to the total airspora revealed Deuteromycotina as highest, followed by Basidiomycotina, Ascomycotina and lowest was Zygomycotina.

The diurnal periodicity studies shows that *Chaetomium* and *Basidiospores* belongs to night spora group. The peak observed between 22 to 24 hrs in case of *Chaetomium* and 18 to 20, 22 to 24 hrs peak in case of Basidiospore. Patil (1985), while studying its circadian periodicity has showed that the *Chaetomium* was maximum at night. Hence, he was placed them “night spora” group. He was also reported 6.14 % basidiospores to be maximum in wet season. Thus, it belongs to “wet spora” group. Mishra and Kamal (1971) reported *Chaetomium globosum* during winter only.

#### Conclusion

Aerobiological studies are very important in relation to disease forecasting, so it must be carried out continuously year round in order to study transport of plant pathogenic spores type from place to place and their ultimate role in inciting plant diseases. Pathogenic spores like *Alternaria*, *Cladosporium*, *Cuvularia*, *Cercospora*, *Rust spore*, *Rhizopus* and *Helminthosporium* were observed in sufficiently high concentrations which were responsible for deterioration in Tomato field etc.

#### REFERENCES

- Ararwal MK, Mukharjee KG and Shivpuri DN, 1969. Studies on the allergenic fungal spores of Delhi, Indian Metropolitan area. *Botanical Aspects Journal of Allergy*, 44:193-203.
- Ararwal MK and Shivpuri DN, 1974. Fungal spores, their role in respiratory allergy, *Adv. Pollen Res.*, 1: 78-128.
- Ebner MR, Haselwandter K and Frank A, 1989. Seasonal fluctuations of airborne fungal allergens. *Journal of Mycology*, 92 : 170-176.

- Jacob WC, 1951.** Aerobiology in compendium of meteorology Society, *Boston*, 1103-1111.
- Lacey J, 1981.** Aerobiology of conidial fungi. In *Biology of conidial fungi*. Cole GT and Kendrick Academic Press Inc. B (eds.) Vol. 1, Pp. 273-416.
- Kularni RL, 1981,** Atmospheric concentration of hyphal fragments, *Proc.Nat.Conf.Env.Bio.*, pp. 209-214.
- Manoharachary C, Sridha K, Singh R, Adholeya A, Suryanarayana TS, Rawat S and Johri BN, 2005.** Fungal biodiversity : distribution, conservation and prospecting of fungi from India. *Current Science*, 89 : 58-71.
- Mediavilla A, Angulo J, Infante F and Dominguez E, 1997.** Influence of meteorological factors on the incidence of *Cladosporium* Link ex Fr. Conidia in the atmosphere of Cordoba (Spain). *Phenology in Seasonal Climates*, 1:117-126.
- Mishra RR and Kamal, 1971.** Aeromycology of Gorakhpur-III. Seasonal variation in air fungal spora. *Applied Journal of Mycopathology and Mycology*, 45 : 301-310.
- Nair PKK, 1978,** Pollen-problems and promises, *Science Reporter*, 15: 378-388.
- Oliveira M, Abreu I, Ribeiro H and Delgado L, 2007.** Fungal spores in the atmosphere in the city of Oporto and its allergological implications. *Review Port of Imunoalergol*, 15: 61-85.
- Shivpuri DN, 1980.** Clinically important pollen, funal and insect allergens for nasobronchial allergy patients in India. *Asp. Allergy Appl. Immunol*, 13 :19-23.
- Shivpuri DN, 1982.** Studies in allergy to fungi in India. *Asp. Allergy Appl. Immunol.*, 15 : 19-30.
- Singh A and Singh AB, 1994.** Airborne in bakery and prevalence of respiratory dysfunction among workers. *Grana*, 33 : 349-358.
- Patil BY, 1985.** Aeromucological studies at Aurangabad. Ph.D. Thesis, Marathwada University, Aurangabad.
- Tilak ST and Kulkarni RL, 1970.** A new air sampler, *Experientia*, 26 : 443.
- Tilak ST, 1989.** Airborne Pollen and Fungal Spores. *Vaijayanti Prakashan*, Aurangabad, pp. 316.
- Zanca M, 2003.** Aerobiological survey on Deuteromycetes from 1999 to 2002 of Mantua's area (Po Valley - North Italy). In : Emberlin et al. (Eds): *Proceeding Third European Symposium in Aerobiology*. Worcester, 30 August 2 September 2003. University College, Worcester 2003.

#### How to cite this article

---

**Kshirsagar J. J., 2021.** Investigation of Fungal Spores of Tomato field at District Beed, M.S., India. *Bioscience Discovery*, 12(3):138-141.

**Google Scholar citation:** <https://scholar.google.co.in/citations?user=vPzEyC8AAAAJ&hl=en>