

CHECKLIST OF INDIANA FUNGI I: MACROFUNGI

Scott T. Bates¹, Justin Golday, Rachel L. Kunnen² and Nathanael J. Pilla³: Department of Biological Sciences, Purdue University Northwest, Westville, IN 46391 USA

ABSTRACT. A checklist of macrofungi was compiled for Indiana as part of a larger effort to document fungi within the state. Our study compiled records of Indiana fungi from digitized specimen data available online through the Mycology Collections data Portal (<http://mycoportal.org>). These data were supplemented with records from the scientific literature. While several small checklists of Indiana fungi exist, the majority of these being published previously in the Proceeding of the Indiana Academy of Science, our study represents the first to comprehensively compile all the available data on Indiana fungi. Overall, more than 19,000 records of Indiana fungi were examined, with 1410 species of macrofungi being documented in this publication. These species represent 24 fungal orders from two major phyla, with 757 species in this checklist being reported in the literature for the first time here. Our study also recovered records documenting other groups of Indiana fungi, such as microfungi, which will be covered in subsequent publications.

Keywords: Ascomycota, Basidiomycota, checklist, Eumycota, fungi, taxonomy

INTRODUCTION

A wide range of natural communities exist in Indiana, with six general ecoregions recognized by Woods and colleagues (1998): the Eastern Corn Belt Plains, Huron/Erie Lake Plains, Interior Plateau, Interior River Valleys and Hills, Central Corn Belt Plains, and Southern Michigan/Northern Indiana Drift Plains. While over half the land area of the state is now devoted to agriculture (Vilsack & Clark 2012), it was once largely forested, and the previously dominant oak-hickory and beech-maple plant communities persist primarily as isolated stands (Friesner 1936; Clark 1994; Whitaker et al. 2012). Although the majority of agricultural land in Indiana is devoted to row crops (Vilsack & Clark 2012), some areas in the state support a high level of vascular plant diversity. For example, the Indiana Dunes National Lakeshore, encompassing some 6,000 ha, has nearly 1,200 native plant species (Wilhelm 1990) in a mosaic of habitats that include xeric sand dunes, sedge meadows, oak savanna, pannes, bogs, fens, kettle lakes, and mesic prairies. Considering the potential for tight linkages between vascular plants and their fungal counterparts (Van der Heijden et al. 1998), a high level of fungal diversity might also be expected across the state.

Members of the kingdom Eumycota (Fungi) are filamentous, or sometimes unicellular, heterotrophic organisms with absorptive nutrition. The diverse forms include mushrooms, rusts, smuts, and yeasts. Many fungal species (e.g., mycorrhizal forms) are essential symbiotic partners of vascular plants, providing them with nutrients such as phosphorus (Bolan 1991), while others are key in cycling carbon through the decay of organic matter (Lindahl et al. 2002). Recent research also suggests fungal pathogens are an increasing threat to the health of animals, plants, and even ecosystems (Fisher et al. 2012). Given this diversity of forms and of ecological roles, it is reasonable to expect fungal diversity to be vast, with estimates ranging from 1.5 to 5.1 million species worldwide (Blackwell 2011). In comparison to this extensive diversity, the number of described species (ca. 100,000) is proportionally low, and understanding of fungal diversity even in temperate regions, such as Indiana, is deficient. At the same time, recent projects aimed at digitizing our natural history collections (e.g., Baird 2010; Conway 2010; Beaman & Cellinese 2012; Blagoderov et al. 2012) have facilitated access to data on organismal diversity. Taking inspiration from these digitization efforts, this study relied largely on digital resources to document our current knowledge of fungal diversity in Indiana.

Over 19,000 digitized records of fungal specimens were examined online in public databases. To supplement these data, we carried out online searches of records of fungi from the state in the

¹ Corresponding author: Scott T. Bates, 219-785-5577 (phone), stbates@purdue.edu.

² Undergraduate student.

³ Graduate student.

scientific literature. While records of Indiana fungi have been published in the broader literature (e.g., Schmit et al. 1999; Harmon & Latin 2005; Wilson et al. 2014), a wealth of digitized publications (e.g., Van Hook 1911, 1912, 1913, 1916, 1921a, 1921b, 1922, 1924, 1925, 1926, 1928, 1929a, 1929b, 1930, 1935; Jackson 1918, 1921a, 1921b; Anderson & Anderson 1920; Fink & Fuson 1920; Van Hook & Busteed 1935) were discovered from the Proceedings of the Indiana Academy of Science, with documentation of fungal diversity in the state beginning with the first published volume (Underwood 1892) and continuing into the twenty-first century (Ruch et al. 2001). This study summarizes the sizable fungal diversity known to exist across Indiana. Due to length, the Indiana fungal checklist will be published in three parts: macrofungi (e.g., bracket fungi, mushrooms, and puffballs), microfungi (e.g., molds and rusts), and lichenized fungi (i.e., lichens) of Indiana.

METHODS

The preliminary fungal species list was compiled through a search of Indiana specimens housed in 72 public collections, universities, and natural history museums across the United States and Canada. The specimen records from these institutions are available online through the Mycology Collections data Portal (MyCoPortal, <http://mycoportal.org>). MyCoPortal is the official repository of data for the Macro- and Microfungi Collections Consortia (MaCC and MiCC, respectively) Thematic Collections Networks, digitization efforts funded by the NSF's Advancing Digitization of Biological Collections (ADBC) program. Data from MyCoPortal are served on the World Wide Web using Symbiota software (Gries et al. 2014). Its database currently contains over three million fungal specimen records from around the world, with more records being added daily and digitization efforts funded until 2018. Records obtained from MyCoPortal specimen data have been supplemented with records of Indiana fungi from the scientific literature. A number of historical mycological publications from the Proceedings of the Indiana Academy of Science (Arthur 1897; Lee 1929; Busteed 1936; Herre 1944; Cottingham 1947, 1948, 1949, 1950, 1951, 1952, 1954, 1956; Wolf 1972; Evans-Ruhl et al. 1982; Schoknecht 1982; Hyerczyk 1998) were accessed online through the Academy's digital archive (<https://journals.iupui.edu/index.php/ias/index>). Several additional pub-

lications from other journals (Reddick 1907; Cooke 1975; Avis et al. 2006, 2008, 2017; Gurung et al. 2011; Avis 2012; Leacock et al. 2016) were also used in compiling the complete list of Indiana fungi.

In order to verify current taxonomic status and check synonymy for each of the scientific names included in the checklist, the online fungal nomenclature databases MycoBank (<http://www.mycobank.org>) and Index Fungorum (<http://www.indexfungorum.org>) were consulted. The system of classification used here follows that outlined in MycoBank. The taxa included within this checklist of macrofungi generally have fruiting bodies that reach well over a centimeter in height or diameter (though see the discussion below). Each species entry in the checklist is annotated to indicate if the scientific name was sourced from MyCoPortal specimen records and/or a reference from the literature (see the 'Key' below). If the species had not been previously reported in the literature as occurring in Indiana, then the annotation 'NR' indicates a new published record for the state. All statistics (viz., numbers of new records and species) are considered only at the level of species; however, citations for new records are indicated (but not counted) for infraspecific taxa, such as varieties. Infraspecific taxa (e.g., *Helvella macropus* var. *brevis*) were included whenever encountered in databases or the literature and also were listed as the binomial (*Helvella macropus*) when it was not known if other species-level records represented a distinct infraspecific taxon (e.g., *Helvella macropus* var. *macropus*).

RESULTS AND DISCUSSION

Over 19,000 records were recovered in MyCoPortal and the scientific literature documenting a wide range of fungi from the state of Indiana (Figs. 1–3). After updating the taxonomic names and checking for synonymy, as well as excluding microfungi, lichens, and slime molds (protozoans traditionally included in mycological collections), 1410 species of macrofungi were recovered. Of these 757 (54%) are reported from Indiana for the first time, adding greatly to our knowledge of fungal diversity in the state. Overall, the species included in this macrofungi checklist represent two major phyla (Ascomycota, Basidiomycota) in the Eumycota. Within these phyla, 24 fungal orders are represented, most abundantly in the Basidiomycota (19 orders containing 1326 species). The complete list of all non-lichenized fungi



Figure 1–3.—Indiana macrofungi. A wide range of fungal diversity can be found within the state of Indiana, including macroscopic fruiting bodies recognizable as (1) mushrooms (*Marasmius* sp.), (2) turkey-tail fungi (*Trametes* sp.), and (3) stink horns (*Mutinus* sp.).

is being made available online through the MyCoPortal (<http://mycoportal.org>, see the Macrofungi of North America: Regions research checklist page), and will be updated regularly as more data become available.

Numerous basidiomycetous corticioid fungi (e.g., *Botryobasidium*, *Corticium*, *Ramaricium*, *Septobasidium*, *Subulicystidium*, *Tulasnella*), having little topology but growing to cover a large area, are included among the macrofungi following mycological tradition. Again following tradition, ascomycetous fungi recorded from Indiana (e.g., *Hypoxylon*, *Kretzschmaria*, *Rosellinia*) with perithecial ascomata (viz., pyrenomycetes) will be included in the checklist of microfungi, even though their growth habit is superficially similar to that of corticioid fungi. The same is true for a number of other minute ascomycetes (e.g., *Cordyceps*, *Dasyphyllum*, *Isaria*, *Mitrula*) that skirt the boundaries between macro- and microfungi, as well as those that cover fruiting bodies of

macrofungi as parasites (e.g., *Hypomyces*). However, included among the macrofungi are a few of the smaller ascomycetous genera (e.g., *Bulgaria*, *Camarops*, *Daldinia*, *Galiella*, *Leotia*, *Microglossum*, *Microstoma*, *Peziza*, *Xylaria*, *Xylocoremium*) known to generally produce somewhat conspicuous ascocarps.

There has been considerable mycological activity in the state, with the earliest specimens collected in 1804 and the first documentation in the literature dating to the late 1800's (Rose 1886). After a more intense period of activity, roughly 1885–1925, workers have consistently collected specimens in the field throughout the decades (Fig. 4). The fungal species in the Indiana checklist demonstrate the wide range of fungal diversity that exist in the state, and a number of species in the macrofungi checklist have importance to humans, such as deadly poisonous mushrooms (e.g., *Amanita phalloides*, see Pringle & Vellinga 2006) and choice edibles (e.g., *Cantharellus* and




Figure 4.—Number of fungal specimens collected per year from Indiana and deposited within various collections in the United States and Canada (an early specimen collected in 1804 is not included here in order to simplify the plot).

Morchella, see Molina et al. 1993). Estimates of the global diversity for fungi posit that there may be six fungal species for every vascular plant species (Hawksworth 1991; Blackwell 2011). If we apply this ratio to Indiana, then a comparison of the total number of fungal species (including macro-, microfungi, and lichens) recovered in our overall study (ca. 3000) against the number of vascular plant species known for state (ca. 2800; pers. comm. Kay Yatskievych) suggests that many more fungal species are in need of documentation. Many species of fungi absent from our Indiana checklists likely represent taxa already known to science. Indeed, the author's record for *Amanita muscaria* var. *guessowii*, never before reported from the state, was discovered without much effort on the campus of Purdue University Northwest (PNW) while preparing this manuscript (a voucher specimen was sent to Purdue's Arthur Fungarium). Noting the magnitude of this gap (perhaps thousands of ‘missing’ species of Indiana fungi), it is equally likely that a number of fungal species new to science await discovery in the state. These facts highlight the

great need for continued mycological research within the state, as suggested previously by Ruch and colleagues (2001).

We acknowledge that this checklist may contain inaccuracy given the scope of taxa included (and the restricted focus of the lead author) as well as the need to revise some of the species recorded here in light of new data (e.g., see O'Donnell et al. 2011 for the morels). Overall, however, we hope this publication represents solid progress toward a more complete understanding of fungal diversity in the state of Indiana.

ACKNOWLEDGMENTS

The authors thank Drs. Peter Avis and Patrick Leacock, as well as Stephen Russell, for making data from their mycological projects easily accessible for use in this publication, and we very much appreciate Mr. Russell's taxonomic review of the species included here. The lead author would also like to personally thank Drs. Barbara Thiers (MaCC) and Andrew N. Miller (MiCC) for spearheading the preparation of ADBC proposals (now

funded) and for serving as PIs overseeing the digitization efforts of fungal specimens, which have made literally millions of fungal specimen records available online.

CHECKLIST OF INDIANA MACROFUNGI

Annotation key.—Annotations [in brackets] follow each taxon, indicating records from Indiana that are published in the literature for the first time here (NR) and/or citing the source for each record, either from the literature (†) or from specimens (‡) deposited in collections (single or multiple records representing each taxon reported here can be searched individually at <http://mycoportal.org>).

ASCOMYCOTA

Boliniiales

Camarops petersii (Berk. & M.A. Curtis) Nannf. [†, ‡]

Geoglossales

Geoglossum difforme Fr. [‡, NR]

Geoglossum nigratum Cooke [‡, NR]

Trichoglossum farlowii (Cooke) E.J. Durand [‡, NR]

Trichoglossum hirsutum (Pers.) Boud. [†, ‡]

Trichoglossum variabile (E.J. Durand) Nannf. [‡, NR]

Trichoglossum velutipes (Peck) E.J. Durand [‡, NR]

Trichoglossum walteri (Berk.) E.J. Durand [‡, NR]

Helotiales

Bulgaria inquinans (Pers.) Fr. [†]

Leotia lubrica (Scop.) Pers. [†, ‡]

Microglossum olivaceum (Pers.) Gillet [‡, NR]

Microglossum rufum (Schwein.) Underw. [†, ‡]

Pezizales

Aleuria aurantia (Pers.) Fuckel [†, ‡]

Disciotis venosa (Pers.) Boud. [†]

Galiella rufa (Schwein.) Nannf. & Korf [†, ‡]

Gyromitra brunnea Underw. [†, ‡]

Gyromitra caroliniana (Bosc.) Fr. [‡, NR]

Gyromitra esculenta (Pers.) Fr. [†, ‡]

Gyromitra gigas (Krombh.) Cooke [†]

Gyromitra infula (Schaeff.) Quél. [‡, NR]

Helvella acetabulum (L.) Quél. [†, ‡]

Helvella albella Quél. [‡, NR]

- Helvella atra* Oeder [‡, NR]
- Helvella elastica* Bull. [†, ‡]
- Helvella lactea* Boud. [‡, NR]
- Helvella lacunosa* Afzel. [†, ‡]
- Helvella macropus* (Pers.) P. Karst. [‡, NR]
- Helvella macropus* var. *brevis* Peck [‡, NR]
- Helvella queletii* Bres. [‡, NR]
- Helvella subglabra* N.S. Weber [‡, NR]
- Helvella sulcata* Afzel. [‡, NR]
- Helvella villosa* Schaeff. [‡, NR]
- Jafnea semitosta* (Berk. & M.A. Curtis) Korf [‡, NR]
- Microstoma floccosum* (Schwein.) Raity. [†, ‡]
- Mitrophora semilibera* (DC.) Lév. [†, ‡]
- Morchella angusticeps* Peck [†, ‡]
- Morchella conica* Pers. [†, ‡]
- Morchella crassipes* (Vent.) Pers. [†]
- Morchella deliciosa* Fr. [‡, NR]
- Morchella esculenta* (L.) Pers. [†, ‡]
- Morchella hybrida* (Sowerby) Pers. [‡, NR]
- Otidea onotica* (Pers.) Fuckel [‡, NR]
- Pachyella clypeata* (Schwein.) Le Gal [‡, NR]
- Peziza aurantiaca* Bull. [‡, NR]
- Peziza badia* Pers. [†, ‡]
- Peziza badioconfusa* Korf [‡, NR]
- Peziza bronca* Peck [‡, NR]
- Peziza cerea* Sowerby [‡, NR]
- Peziza citrina* (Hedw.) Pers. [†, ‡]
- Peziza dehnii* Rabenh. [‡, NR]
- Peziza domiciliana* Cooke [†, ‡]
- Peziza odorata* Peck [‡, NR]
- Peziza repanda* Pers. [†, ‡]
- Peziza succosa* Berk. [†, ‡]
- Peziza sulcata* Pers. [†]
- Peziza sylvestris* (Boud.) Sacc. & Traverso [‡, NR]
- Peziza vesiculosa* Bull. [†, ‡]
- Plectania occidentalis* (Schwein.) Seaver [†, ‡]
- Sarcoscypha austriaca* (Beck ex Sacc.) Boud. [‡, NR]
- Sarcoscypha coccinea* (Scop.) Sacc. [†, ‡]
- Sarcoscypha occidentalis* (Schwein.) Sacc. [†, ‡]
- Tuber dryophilum* Tul. & C. Tul. [‡, NR]
- Tuber rufum* Pollini [†]
- Urnula craterium* (Schwein.) Fr. [†, ‡]
- Verpa bohemica* var. *bispora* (Sorokin) Syd. [†]
- Verpa conica* (O.F. Müll.) Sw. [†]

Xylariales

Daldinia childiae J.D. Rogers & Y.M. Ju [‡, NR]

Daldinia concentrica (Bolton) Ces. & De Not. [†, ‡]

Daldinia simulans Child [‡, NR]

Daldinia vernicosa Ces. & De Not. [†, ‡]

Xylaria castorea Berk. [†, ‡]

Xylaria comosa (Mont.) Mont. [†]
Xylaria corniformis (Fr.) Fr. [‡, NR]
Xylaria cornu-damae (Schwein.) Berk. [†, ‡]
Xylaria cubensis (Mont.) Fr. [‡, NR]
Xylaria digitata (L.) Grev. [‡, NR]
Xylaria hypoxylon (L.) Grev. [†, ‡]
Xylaria longiana Rehm [‡, NR]
Xylaria longipes Nitschke [‡, NR]
Xylaria mali Fromme [‡, NR]
Xylaria polymorpha (Pers.) Grev. [†, ‡]
Xylaria subterranea (Schwein.) Sacc. [†]
Xylaria tentaculata Ravenel ex Berk. [‡, NR]
Xylocoremium flabelliforme (Schwein.) J.D. Rogers [‡, NR]

BASIDIOMYCOTA

Agaricales

Agaricus abruptibulbus Peck [†, ‡]
Agaricus arvensis Schaeff. [†, ‡]
Agaricus bitorquis (Quél.) Sacc. [†, ‡]
Agaricus campestris L. [†, ‡]
Agaricus comatus Fr. [‡, NR]
Agaricus diminutivus Peck [‡, NR]
Agaricus dorsalis Peck [†]
Agaricus haerens Peck [†, ‡]
Agaricus magniceps Peck [‡, NR]
Agaricus moelleri Wasser [‡, NR]
Agaricus placomyces Peck [†, ‡]
Agaricus scitulus Massee [†]
Agaricus silvaticus Schaeffer [†, ‡]
Agaricus sylvicola (Vittad.) Peck [†, ‡]
Agaricus xanthoderma Genev. [‡, NR]
Agrocybe acericola (Peck) Singer [‡, NR]
Agrocybe arvalis (Fr.) Heim & Romagn. [‡, NR]
Agrocybe dura (Bolton) Singer [†, ‡]
Agrocybe erebia (Fr.) Singer [‡, NR]
Agrocybe firma (Peck) Singer [‡, NR]
Agrocybe gibberosa (Fr.) Fayod [‡, NR]
Agrocybe howeana (Peck) Singer [‡, NR]
Agrocybe pediades (Fr.) Fayod [†, ‡]
Agrocybe praecox (Pers.) Fayod [†, ‡]
Amanita abrupta Peck [†, ‡]
Amanita aestivalis Singer ex Singer [‡, NR]
Amanita bisporigera G.F. Atk. [†, ‡]
Amanita brunnescens var. *brunnescens* G.F. Atk. [†, ‡]
Amanita brunnescens var. *pallida* L. Krieg. [‡, NR]
Amanita caesarea (Scop.) Pers. [†]
Amanita ceciliae (Berk. & Broome) Bas [‡, NR]
Amanita chlorinosma (Peck) Lloyd [†, ‡]
Amanita citrina (Schaeff.) Pers. [‡, NR]
Amanita citrina f. *lavendula* (Coker) Veselý [‡, NR]

Amanita cokeri (E.-J. Gilbert & Kühner) E.-J. Gilbert [‡, NR]
Amanita cothurnata G.F. Atk. [†]
Amanita daucipes (Berk. & Mont.) Lloyd [‡, NR]
Amanita excelsa (Fr.) Bertill. [†, ‡]
Amanita farinosa Schwein. [†, ‡]
Amanita flavoconia G.F. Atk. [†, ‡]
Amanita flavorubescens G.F. Atk. [†, ‡]
Amanita floccocapitata G.F. Atk. [†]
Amanita fulva Pers. [‡, NR]
Amanita gemmata (Fr.) Bertill. [‡, NR]
Amanita junquillea Quél. [†]
Amanita magnivelaris Peck (Fr.) P. Kumm. [‡, NR]
Amanita multisquamosa Peck [‡, NR]
Amanita muscaria (L.) Lam. [†, ‡]
Amanita muscaria var. *guessowii* Veselý [‡, NR]
Amanita pantherina (DC.) Krombh. [‡, NR]
Amanita phalloides (Fr.) Link [†, ‡]
Amanita rubescens Pers. [†, ‡]
Amanita rubescens f. *alba* (Coker) Sartory & L. Maire [‡, NR]
Amanita solitaria (Bull.) Mérat [†, ‡]
Amanita spreta (Peck) Sacc. [†, ‡]
Amanita strobiliformis (Paulet ex Vittad.) Bertill. [†, ‡]
Amanita subphalloides (Murrill) Murrill [‡, NR]
Amanita thiersii Bas [‡, NR]
Amanita vaginata (Bull.) Lam [†, ‡]
Amanita verna (Bull.) Lam. [†]
Amanita virosa Bertill. [†, ‡]
Arachnion album Schwein. [‡, NR]
Armillaria caligata (Viv.) Gillet [‡, NR]
Armillaria calvescens Bérubé & Dessur. [‡, NR]
Armillaria gallica Marxm. & Romagn. [‡, NR]
Armillaria mellea (Vahl) P. Kumm. [†, ‡]
Armillaria nardosmia Ellis [†]
Armillaria tabescens (Scop.) Emel [†, ‡]
Arrhenia epichysium (Pers.) Redhead, Lutzoni, Moncalvo & Vilgalys [‡, NR]
Arrhenia peltigerina (Peck) Redhead [†]
Asterophora lycoperdoides (Bull.) Ditmar [‡, NR]
Baeospora myriadophylla (Peck) Singer [†]
Bolbitius coprophilus (Peck) Hongo [‡, NR]
Bolbitius tener Berk. [†]
Bolbitius titubans (Bull.) Fr. [†, ‡]
Bovista aestivalis (Bonord.) Demoulin [‡, NR]
Bovista dermoxantha (Vittad.) De Toni [‡, NR]
Bovista longispora Kreisel [‡, NR]
Bovista minor Morgan [‡, NR]
Bovista pilosa Berk. & M.A. Curtis [‡, NR]
Bovista plumbea Pers. [‡, NR]
Bovista pusilla (Batsch) Pers. [‡, NR]
Bovistella ohiensis (Ellis & Morgan) Morgan [†, ‡]

- Bovistella radicata* (Durieu & Mont.) Pat. [‡, NR]
Calvatia craniiformis (Schwein.) Fr. ex De Toni [†, ‡]
Calvatia cyathiformis (Bosc) Morgan [†, ‡]
Calvatia excipuliformis (Scop.) Perdeck [‡, NR]
Calvatia gigantea (Batsch) Lloyd [†, ‡]
Calvatia lepidophora (Ellis & Everh.) Coker & Couch [‡, NR]
Calvatia lilacina (Mont. & Berk.) Henn. [‡, NR]
Calvatia rugosa (Berk. & M.A. Curtis) D.A. Reid [†, ‡]
Calvatia utriformis (Bull.) Jaap [†]
Calyptella campanula (Nees) W.B. Cooke [‡, NR]
Cheimonophyllum candidissimum (Berk. & M.A. Curtis) Singer [‡, NR]
Chlorophyllum agaricoides (Czern.) Vellinga [‡, NR]
Chlorophyllum molybdites (G. Mey.) Massee [†, ‡]
Chondrostereum purpureum (Pers.) Pouzar [‡, NR]
Clavaria appalachiensis Coker [‡, NR]
Clavaria fragilis Holmsk. [†, ‡]
Clavaria inaequalis O.F. Müll. [†]
Clavaria rosea Dalman [‡, NR]
Clavaria tetragona Schwein. [†]
Clavulinopsis corniculata (Schaeff.) Corner [‡, NR]
Clavulinopsis fusiformis (Sowerby) Corner [†, ‡]
Clavulinopsis laeticolor (Berk. & M.A. Curtis) R.H. Petersen [‡, NR]
Clitocybe americana H.E. Bigelow [‡, NR]
Clitocybe aeruginosa H.E. Bigelow [‡, NR]
Clitocybe albidioides Peck [†]
Clitocybe candida Bres. [†, ‡]
Clitocybe cartilaginea (Bull. ex Pers.) Bres [†]
Clitocybe catinus (Fr.) Quél. [†]
Clitocybe clavipes (Pers.) P. Kumm. [‡, NR]
Clitocybe connata (Schumach.) Gillet [‡, NR]
Clitocybe dealbata (Sowerby) P. Kumm. [†, ‡]
Clitocybe diatreta (Fr.) P. Kumm. [‡, NR]
Clitocybe dilatata P. Karst. [‡, NR]
Clitocybe ditopa (Fr.) Gillet [†]
Clitocybe eccentrica X.H. Wang & P.G. Liu [†]
Clitocybe elephantina Murrill [‡, NR]
Clitocybe fragrans (With.) P. Kumm. [‡, NR]
Clitocybe gibba (Pers.) P. Kumm. [†, ‡]
Clitocybe intermedia Kauffman [‡, NR]
Clitocybe laccata var. *pallidifolia* Peck [†, ‡]
Clitocybe martiorum J. Favre [‡, NR]
Clitocybe multiceps Peck [†]
Clitocybe nebularis (Batsch) P. Kumm [†, ‡]
Clitocybe ochropurpurea (Berk.) Sacc. [†]
Clitocybe odora (Bull.) P. Kumm. [†, ‡]
Clitocybe patuloides (Peck) Peck [†]
- Clitocybe phaeophthalma* (Pers.) Kuyper [†, ‡]
Clitocybe phyllophila (Pers.) P. Kumm. [†]
Clitocybe piceina Peck [†]
Clitocybe regularis Peck [‡, NR]
Clitocybe sinopica (Fr.) P. Kumm. [†]
Clitocybe subconnexa Murrill [‡, NR]
Clitocybe trullisata Ellis [‡, NR]
Clitocybula abundans (Peck) Singer [†, ‡]
Clitocybula lacerata (Scop.) Singer ex Métrod [‡, NR]
Clitopilus caespitosus Peck [†]
Clitopilus hrbanovii (Velen.) Singer [‡, NR]
Clitopilus noveboracensis (Peck) Sacc. [†]
Clitopilus prunulus (Scop.) P. Kumm. [†, ‡]
Collybia amabilipes (Peck) Peck [†]
Collybia cookei (Bres.) J.D. Arnold [‡, NR]
Conocybe crispa (Longyear) Singer [†]
Conocybe cryptocystis (G.F. Atk.) Singer [‡, NR]
Conocybe tenera (Schaeff.) Fayod [†, ‡]
Coprinellus deliquescens (Bull.) P. Karst. [†]
Coprinellus disseminatus (Pers.) J.E. Lange [†, ‡]
Coprinellus domesticus (Bolton) Vilgalys [†]
Coprinellus ephemerus (Bull.) Redhead [†]
Coprinellus micaceus (Bull.) Vilgalys, Hopple & Jacq. Johnson [†, ‡]
Coprinopsis atramentaria (Bull.) Redhead, Vilgalys & Moncalvo [†, ‡]
Coprinopsis lagopus (Fr.) Redhead, Vilgalys & Moncalvo [†, ‡]
Coprinopsis nivea (Pers.) Redhead, Vilgalys & Moncalvo [‡, NR]
Coprinopsis variegata (Peck) Redhead, Vilgalys & Moncalvo [‡, NR]
Coprinus comatus (O.F. Müll.) Pers. [†, ‡]
Coprinus comatus var. *ovatus* (Scop.) Quél. [†]
Coprinus ebulbosus Peck [†]
Coprinus quadrifidus Peck [†]
Coprinus rotundisporus Peck [†]
Cortinarius alboviolaceus (Pers.) Fr. [†, ‡]
Cortinarius armillatus (Fr.) Fr. [†, ‡]
Cortinarius autumnalis Peck [†, ‡]
Cortinarius caperatus (Pers.) Fr. [†]
Cortinarius cinnabarinus Fr. [†, ‡]
Cortinarius cinnamomeus (L.) Gray [†]
Cortinarius claricolor (Fr.) Fr. [‡, NR]
Cortinarius collaris E. Horak & M.M. Moser [†]
Cortinarius collinitus (Sowerby) Gray [†]
Cortinarius corrugatus Peck [‡, NR]
Cortinarius cylindripes Kauffman [†]
Cortinarius distans Peck [†, ‡]

- Cortinarius duracinus* Fr. [†, ‡]
Cortinarius gentilis (Fr.) Fr. [‡, NR]
Cortinarius hinnuleus Fr. [†]
Cortinarius iodeoides Kauffman [‡, NR]
Cortinarius iodes Berk. & M.A. Curtis [‡, NR]
Cortinarius michiganensis Kauffman [†]
Cortinarius obliquus Peck [†]
Cortinarius ochraceus Peck [†]
Cortinarius olearioides Rob. [‡, NR]
Cortinarius semisanguineus (Fr.) Gillet [‡, NR]
Cortinarius sericeps Peck [‡, NR]
Cortinarius torvus (Fr.) Fr. [‡, NR]
Cortinarius violaceus (L.) Gray [‡, NR]
Crepidotus applanatus (Pers.) P. Kumm. [†, ‡]
Crepidotus applanatus var. *globiger* (Berk.) Pilát [‡, NR]
Crepidotus calolepis (Fr.) P. Karst. [†]
Crepidotus crocophyllus (Berk.) Sacc. [‡, NR]
Crepidotus ellipoideus Hesler & A.H. Sm. [‡, NR]
Crepidotus fulvotomentosus (Peck) Peck [†, ‡]
Crepidotus fusisporus Hesler & A.H. Sm. [†]
Crepidotus herbarum (Peck) Peck [‡, NR]
Crepidotus hygrophanus Murrill [‡, NR]
Crepidotus lundellii Pilát [†]
Crepidotus mollis (Schaeff.) Staude [†, ‡]
Crepidotus nephrodes (Berk. & M.A. Curtis) Sacc. [‡, NR]
Crepidotus versutus (Peck) Peck [†, ‡]
Crepidotus vulgaris Hesler & A.H. Sm. [‡, NR]
Crinipellis setipes (Peck) Singer [‡, NR]
Crinipellis zonata (Peck) Sacc. [†, ‡]
Crucibulum crucibuliforme (Scop.) V.S. White [†, ‡]
Cyathus olla (Batsch) Pers. [†, ‡]
Cyathus poeppigii Tul. & C. Tul. [‡, NR]
Cyathus stercoreus (Schwein.) De Toni [†, ‡]
Cyathus striatus (Huds.) Willd. [†, ‡]
Cylindrobasidium evolvens (Fr.) Jülich [‡, NR]
Cylindrobasidium torrendii (Bres.) Hjortstam [‡, NR]
Cyphella fasciculata Berk. & M.A. Curtis [‡, NR]
Cyphella mellea Burt [‡, NR]
Deconica subviscida Peck [‡, NR]
Disciseda candida (Schwein.) Lloyd [‡, NR]
Entoloma abortivum (Berk. & M.A. Curtis) Donk [†, ‡]
Entoloma alboumbonatum Hesler [‡, NR]
Entoloma asprellum (Fr.) Fayod [†]
Entoloma byssisedum (Pers.) Donk [‡, NR]
Entoloma carneogriseum (Berk. & Broome) Noordel. [†]
Entoloma clypeatum (L.) P. Kumm. [†]
Entoloma gracilipes (Peck) Malloch [‡, NR]
Entoloma grande Peck [†]
Entoloma jubatum (Fr.) P. Karst. [†]
Entoloma lividum (Bull.) Quél. [†]
Entoloma murrayi (Berk. & M.A. Curtis) Sacc. [‡, NR]
Entoloma rhodopolium (Fr.) P. Kumm. [†, ‡]
Entoloma salmonicum (Peck) Sacc. [‡, NR]
Entoloma sericellum (Fr.) P. Kumm. [‡, NR]
Entoloma serrulatum (Fr.) Hesler [‡, NR]
Entoloma sinuatum (Bull. ex Pers.) P. Kumm. [†]
Entoloma strictipes (Peck) Hesler [‡, NR]
Entoloma strictius (Peck) Sacc. [†, ‡]
Entoloma subcostatum G.F. Atk. [†]
Entoloma undatum (Gillet) M.M. Moser [†]
Entoloma vernum S. Lundell [‡, NR]
Fistulina hepatica (Schaeff.) With. [†]
Flammula betulina Peck [†]
Flammula praecox Peck [‡, NR]
Flammulaster erinaceellus (Peck) Watling [‡, NR]
Flammulaster gracilis (Quél.) Watling [‡, NR]
Flammulina velutipes (Curtis) Singer [†, ‡]
Galera ovalis (Fr.) Gillet [†]
Galerina autumnalis (Peck) A.H. Sm. & Singer [‡, NR]
Galerina marginata (Batsch) Kühner [‡, NR]
Galerina unicolor (Vahl) Singer [†, ‡]
Gerronema strombodes (Berk. & Mont.) Singer [‡, NR]
Gymnopilus aeruginosus (Peck) Singer [†]
Gymnopilus bellulus (Peck) Murrill [‡, NR]
Gymnopilus junonioides (Fr.) P.D. Orton [†, ‡]
Gymnopilus liquiritiae (Pers.) P. Karst. [‡, NR]
Gymnopilus sapineus (Fr.) Maire [‡, NR]
Gymnoporus acervatus (Fr.) Murrill [†, ‡]
Gymnoporus alkalivirens (Singer) Halling [‡, NR]
Gymnoporus androsaceus (L.) J.L. Mata & R.H. Petersen [‡, NR]
Gymnoporus aquosus (Bull.) Antonín & Noordel. [‡, NR]
Gymnoporus atratoides (Peck) Murrill [†]
Gymnoporus biformis (Peck) Halling [‡, NR]
Gymnoporus confluens (Pers.) Antonín, Halling & Noordel. [†, ‡]
Gymnoporus dichrous (Berk. & M.A. Curtis) Halling [‡, NR]
Gymnoporus dryophilus (Bull.) Murrill [†, ‡]
Gymnoporus erythropus (Pers.) Antonín [†, ‡]
Gymnoporus foetidus (Sowerby) J.L. Mata & R.H. Petersen [†, ‡]
Gymnoporus fusipes (Bull.) Gray [†]
Gymnoporus hariolorum (Bull.) Antonín [†]
Gymnoporus hygrophoroides (Peck) Murrill [†, ‡]
Gymnoporus iocephalus (Berk. & M.A. Curtis) Halling [‡, NR]
Gymnoporus luxurians (Peck) Murrill [‡, NR]

- Gymnopus peronatus* (Bolton) Gray [†, ‡]
Gymnopus polyphyllus (Peck) Halling [‡, NR]
Gymnopus spongiosus (Berk. & M.A. Curtis)
 Halling [‡, NR]
Gymnopus subnudus (Ellis ex Peck) Halling [‡, NR]
Gymnopus subsulphureus (Peck) Murrill [‡, NR]
Hebeloma albidulum Peck [†]
Hebeloma album Peck [†]
Hebeloma crustuliniforme (Bull.) Quél. [‡, NR]
Hebeloma mesophaeum (Pers.) Quél. [†]
Hebeloma sinapizans (Poulet) Gillet [‡, NR]
Henningssomyces candidus (Pers.) Kuntze [‡, NR]
Hohenbuehelia angustata (Berk.) Singer [‡, NR]
Hohenbuehelia approximans (Peck) Singer [‡, NR]
Hohenbuehelia horakii Courtec. [‡, NR]
Hohenbuehelia petalooides (Bull.) Schulzer [†, ‡]
Hydropus floccipes (Fr.) Singer [‡, NR]
Hygrocybe acutoconica (Clem.) Singer [‡, NR]
Hygrocybe acutoconica var. *cuspidata* (Peck)
 Arnolds [‡, NR]
Hygrocybe borealis (Peck) Bon [†]
Hygrocybe calytriformis (Berk. & Broome)
 Fayod [‡, NR]
Hygrocybe cantharellus (Schwein.) Murrill [‡, NR]
Hygrocybe ceracea (Wulfen) P. Kumm. [†, ‡]
Hygrocybe chlorophana (Fr.) Wünsche [‡, NR]
Hygrocybe coccinea (Schaeff.) P. Kumm. [†, ‡]
Hygrocybe colemanniana (A. Bloxam) P.D. Orton
 & Watling [‡, NR]
Hygrocybe conica (Schaeff.) P. Kumm. [†, ‡]
Hygrocybe irrigata (Pers.) Bon [‡, NR]
Hygrocybe laeta (Pers.) P. Kumm. [‡, NR]
Hygrocybe lepida Arnolds [‡, NR]
Hygrocybe marginata var. *concolor* (A.H. Smith)
 A.E. Bessette, A.R. Bessette, W.C. Roody &
 W.E. Sturgeon [‡, NR]
Hygrocybe marginata var. *marginata* (Peck)
 Murrill [‡, NR]
Hygrocybe marginata var. *olivacea* (A.H. Smith &
 Hesler) A.E. Bessette, A.R. Bessette, W.C.
 Roody & W.E. Sturgeon [‡, NR]
Hygrocybe miniata (Fr.) P. Kumm. [†, ‡]
Hygrocybe nitida (Berk. & M.A. Curtis) Murrill
 [‡, NR]
Hygrocybe obrussea (Fr.) Wünsche [‡, NR]
Hygrocybe pratensis (Schaeff.) Murrill [†, ‡]
Hygrocybe psittacina (Schaeff.) P. Kumm. [†, ‡]
Hygrocybe punicea (Fr.) P. Kumm. [†, ‡]
Hygrocybe virginea (Wulfen) P.D. Orton &
 Watling [‡, NR]
Hygrophorus eburneus (Bull.) Fr. [†]
Hygrophorus flavodiscus Frost [†]
- Hygrophorus laurae* Morg. [†]
Hygrophorus occidentalis A.H. Sm. & Hesler [‡, NR]
Hygrophorus pudorinus (Fr.) Fr. [†, ‡]
Hygrophorus roseobrunneus Murrill [†]
Hygrophorus russula (Schaeff.) Kauffman [†]
Hygrophorus sordidus Peck [†, ‡]
Hygrophorus tennesseensis A.H. Sm. & Hesler [‡, NR]
Hypholoma candolleanum (Fr.) Maire [‡, NR]
Hypholoma capnoides (Fr.) P. Kumm. [‡, NR]
Hypholoma cutifractum Peck [†]
Hypholoma echiniceps G.F. Atk. [‡, NR]
Hypholoma hydrophilum (Bull.) Quél. [‡, NR]
Hypholoma incertum (Peck) A.H. Sm. [‡, NR]
Hypholoma lateritium (Schaeff.) P. Kumm. [†, ‡]
Hypsizygus ulmarius (Bull.) Redhead [†, ‡]
Inocybe albodisca Peck [†, ‡]
Inocybe calospora Quél. [†, ‡]
Inocybe cookei Bres. [‡, NR]
Inocybe decipiensoides Peck [‡, NR]
Inocybe fibrillosa Peck [†]
Inocybe geophylla (Sowerby) P. Kumm. [†, ‡]
Inocybe griseovelata Kühner [‡, NR]
Inocybe lacera (Fr.) P. Kumm. [‡, NR]
Inocybe lilacina (Peck) Kauffman [†]
Inocybe maculata Boud. [‡, NR]
Inocybe pallidipes Ellis & Everh. [‡, NR]
Inocybe praetervisa Quél. [‡, NR]
Inocybe pyriodora (Pers.) P. Kumm. [‡, NR]
Inocybe radiata Peck [‡, NR]
Inocybe rimosa (Bull.) P. Kumm. [†, ‡]
Inocybe sororia Kauffman [‡, NR]
Inocybe unicolor Peck [‡, NR]
Kuehneromyces mutabilis (Schaeff.) Singer &
 A.H. Sm. [†]
Laccaria amethystina Cooke [‡, NR]
Laccaria bicolor (Maire) P.D. Orton [‡, NR]
Laccaria laccata (Scop.) Cooke [†, ‡]
Laccaria ochropurpurea (Berk.) Peck [†, ‡]
Laccaria proxima (Boud.) Pat. [‡, NR]
Laccaria tortilis (Bolton) Cooke [‡, NR]
Laccaria trullisata (Ellis) Peck [‡, NR]
Lachnella villosa (Pers.) Gillet [‡, NR]
Lacrymaria lacrymabunda (Bull.) Pat. [†, ‡]
Lepiota alluvina (Peck) Morgan [†]
Lepiota aspera (Pers.) Quél. [†, ‡]
Lepiota augustana (Britzelm.) Sacc. [†]
Lepiota castanea Quél. [‡, NR]
Lepiota conspurcata (Willd.) Morgan [†]
Lepiota clypeolaria (Bull.) P. Kumm. [‡, NR]
Lepiota cristata (Bolton) P. Kumm. [†, ‡]
Lepiota farinosa Peck [†]
Lepiota fischeri Kauffman [‡, NR]

- Lepiota granosa* Morgan [†]
Lepiota miamensis (Morgan) Sacc. [†]
Lepiota naucinoides (Peck) Peck [†]
Lepiota subincarnata J.E. Lange [‡, NR]
Lepiota virescens (Speg.) Morgan [†]
Lepista flaccida (Sowerby) Pat. [†, ‡]
Lepista irina (Fr.) H.E. Bigelow [‡, NR]
Lepista nuda (Bull.) Cooke [‡, NR]
Lepista personata (Fr.) Cooke [†, ‡]
Leptonia rosea var. *marginata* Largent [‡, NR]
Leucoagaricus americanus (Peck) Vellinga [†, ‡]
Leucoagaricus caeruleascens (Peck) J.F. Liang,
 Zhu L. Yang & J. Xu [†]
Leucoagaricus leucothites (Vittad.) Wasser [†, ‡]
Leucoagaricus rubrotinctus (Peck) Singer [†, ‡]
Leucocoprinus cepistipes (Sowerby) Pat. [†, ‡]
Leucocoprinus flavescens (Morgan) H.V. Sm. [‡, NR]
Leucopaxillus albissimus (Peck) Singer [†, ‡]
Leucopaxillus cerealis (Lasch) Singer [‡, NR]
Leucopaxillus giganteus (Sowerby) Singer [†]
Leucopaxillus laterarius (Peck) Singer & A.H. Sm.
 [‡, NR]
Leucopaxillus paradoxus (Costantin & L.M.
 Dufour) Boursier [‡, NR]
Lindtneria trachyspora (Bourdot & Galzin) Pilát
 [‡, NR]
Lycoperdon americanum Demoulin [‡, NR]
Lycoperdon echinatum Pers. [‡, NR]
Lycoperdon ericaeum Bonord. [‡, NR]
Lycoperdon fuscum Huds. [‡, NR]
Lycoperdon marginatum Vittad. [‡, NR]
Lycoperdon molle Pers. [‡, NR]
Lycoperdon nigrescens Wahlenb. [‡, NR]
Lycoperdon peckii J.B. Morgan [‡, NR]
Lycoperdon pedicellatum Gardezi [‡, NR]
Lycoperdon perlatum Pers. [†, ‡]
Lycoperdon pulcherrimum Berk. & M.A. Curtis [‡, NR]
Lycoperdon pyriforme Schaeff. [†, ‡]
Lycoperdon radicatum Reichard [‡, NR]
Lycoperdon rimulatum Peck [‡, NR]
Lycoperdon stellare (Peck) Lloyd [‡, NR]
Lycoperdon umbrinum Pers. [†, ‡]
Lycoperdon wrightii Berk. & M.A. Curtis [‡, NR]
Lyophyllum decastes (Fr.) Singer [‡, NR]
Lyophyllum plexipes (Fr.) Kühner & Romagn. [†]
Macrocytidia cucumis (Pers.) Joss. [‡, NR]
Macrolepiota procera (Scop.) Singer [†, ‡]
Marasmiellus candidus (Fr.) Singer [†]
Marasmiellus papillatus (Peck) Redhead & Hal-
 ling [‡, NR]
Marasmiellus vaillantii (Pers.) Singer [‡, NR]
Marasmius armeniacus Gilliam [‡, NR]
Marasmius capillaris Morgan [†, ‡]
Marasmius cohaerens (Pers.) Cooke & Quél. [†, ‡]
Marasmius copelandii var. *olidus* (Gilliam) Des-
 jardin [‡, NR]
Marasmius delectans Morgan [†, ‡]
Marasmius falcatipes Desjardin [‡, NR]
Marasmius felix Morgan [‡, NR]
Marasmius fulvoferrugineus Gilliam [‡, NR]
Marasmius glabellus Peck [†]
Marasmius graminum (Lib.) Berk. [†, ‡]
Marasmius haematocephalus (Mont.) Fr. [†]
Marasmius oreades (Bolton) Fr. [†, ‡]
Marasmius pulcherripes Peck [‡, NR]
Marasmius rotula (Scop.) Fr. [†, ‡]
Marasmius setosus (Sowerby) Noordel. [‡, NR]
Marasmius siccus Schwein. [†, ‡]
Marasmius spissus Gilliam [‡, NR]
Marasmius strictipes (Peck) Singer [‡, NR]
Marasmius sullivantii Mont. [‡, NR]
Marasmius velutipes Berk. & M.A. Curtis [†]
Marasmius vialis Peck [‡, NR]
Megacollybia platyphylla (Pers.) Kotl. & Pouzar
 [†, ‡]
Megacollybia rodmanii R.H. Petersen, K.W.
 Hughes & Lickey [‡, NR]
Megacollybia rodmanii f. *murina* R.H. Petersen &
 K.W. Hughes [‡, NR]
Melanoleuca alboflavida (Peck) Murrill [‡, NR]
Melanoleuca melaleuca (Pers.) Murrill [†, ‡]
Merismodes anomala (Pers.) Singer [‡, NR]
Morganella subincarnata (Peck) Kreisel & Dring
 [‡, NR]
Mycena acicula (Schaeff.) P. Kumm. [‡, NR]
Mycena albiceps (Peck) Gilliam [‡, NR]
Mycena alcalina (Fr.) P. Kumm. [†]
Mycena clavularis (Batsch) Sacc. [‡, NR]
Mycena corticola (Pers.) Gray [‡, NR]
Mycena epityrgia (Scop.) Gray [†, ‡]
Mycena epityrgia var. *griseoviridis* (A.H. Sm.)
 Maas Geest. [‡, NR]
Mycena filopes (Bull.) P. Kumm. [‡, NR]
Mycena galericulata (Scop.) Gray [†, ‡]
Mycena haematopus (Pers.) P. Kumm. [†, ‡]
Mycena hemisphaerica Peck [‡, NR]
Mycena inclinata (Fr.) Quél. [‡, NR]
Mycena leaiana (Berk.) Sacc. [†, ‡]
Mycena leptophylla (Peck) Sacc. [†]
Mycena luteopallens Peck [‡, NR]
Mycena murina (Murrill) Murrill [‡, NR]
Mycena niveipes (Murrill) Murrill [†]
Mycena parabolica (Fr.) Quél. [†]
Mycena pectinata (Murrill) Murrill [‡, NR]
Mycena polygramma (Bull.) Gray [†]
Mycena pulcherrima (Peck) Sacc. [‡, NR]

- Mycena pura* (Pers.) P. Kumm. [†, ‡]
Mycena rutilantiformis (Murrill) Murrill [‡, NR]
Mycena subcaerulea (Peck) Sacc. [‡, NR]
Mycetinis scorodonius (Fr.) A.W. Wilson & Desjardin [‡, NR]
Nolanea howellii Peck [‡, NR]
Omphalia alboflava Moorgan [†]
Omphalina pyxidata (Bull.) Quél. [‡, NR]
Omphalotus olearius (DC.) Singer [†, ‡]
Ossicaulis lignatilis (Pers.) Redhead & Ginns [†]
Ozonium omnivorum Shear [‡, NR]
Panaeolina foeniseccii (Pers.) Maire [†]
Panaeolus campanulatus (Bull.) Quél. [†]
Panaeolus papilionaceus (Bull.) Quél. [†, ‡]
Panaeolus semiovatus (Sowerby) S. Lundell & Nannf. [†]
Panaeolus solidipes (Peck) Sacc. [†]
Panellus pusillus (Pers. ex Lév.) Burds. & O.K. Mill. [†]
Panellus serotinus (Pers.) Kühner [‡, NR]
Panellus stipticus (Bull.) P. Karst. [†, ‡]
Parasola plicatilis (Curtis) Redhead, Vilgalys & Hopple [†, ‡]
Phaeosolenia inconspicua (Sacc.) Donk [‡, NR]
Pholiota adiposa (Batsch) P. Kumm. [†, ‡]
Pholiota aggericola (Peck) Sacc. [†]
Pholiota angustipes (Peck) Sacc. [†]
Pholiota aurivella (Batsch) P. Kumm. [‡, NR]
Pholiota cerasina Peck [†]
Pholiota curvipes (P. Kumm.) Quél. [†]
Pholiota flammans (Batsch) P. Kumm. [†]
Pholiota luteofolius (Peck) Sacc. [†]
Pholiota polychroa (Berk.) A.H. Sm. & H.J. Brodie [‡, NR]
Pholiota populnea (Pers.) Kuyper & Tjall.-Beuk. [†]
Pholiota squarrosoides (Peck) Sacc. [†]
Pholiota subsquarrosa (Fr.) Sacc. [†, ‡]
Pholiota veris A.H. Sm. & Hesler [‡, NR]
Pholiotina rugosa (Peck) Singer [†]
Phyllocladus nidulans (Pers.) Singer [†, ‡]
Pleurotus atropellitus Peck [†]
Pleurotus cystidiosus O.K. Mill. [‡, NR]
Pleurotus dryinus (Pers.) P. Kumm. [†]
Pleurotus ostreatus (Jacq.) P. Kumm. [†, ‡]
Pleurotus populinus O. Hilber & O.K. Mill. [‡, NR]
Pleurotus pulmonarius (Fr.) Quél. [†, ‡]
Pleurotus sapidus Quél. [†, ‡]
Pleurotus serotinoides (Peck) Sacc. [†]
Pleurotus serotinus (Pers.) P. Kumm. [†]
Pluteus americanus (P. Banerjee & Sundberg Justo, E.F. Malysheva & Minnis [‡, NR]
Pluteus cervinus (Schaeff.) P. Kumm. [†, ‡]
Pluteus cervinus var. *albus* Peck [†]
Pluteus chrysophaeus (Schaeff.) Quél. [‡, NR]
Pluteus chrysophlebius (Berk. & M.A. Curtis) Sacc. [‡, NR]
Pluteus cincus Cke. [†]
Pluteus fuliginosus Murrill [‡, NR]
Pluteus fulvobadius Murrill [‡, NR]
Pluteus leoninus (Schaeff.) P. Kumm. [†]
Pluteus longistriatus (Peck) Peck [†, ‡]
Pluteus nanus (Pers.) P. Kumm. [†, ‡]
Pluteus pellitus (Pers.) P. Kumm. [‡, NR]
Pluteus petasatus (Fr.) Gillet [‡, NR]
Pluteus podospileus Sacc. & Cub. [‡, NR]
Pluteus romellii (Britzelm.) Sacc. [‡, NR]
Pluteus salicinus (Pers.) P. Kumm. [†]
Pluteus umbrosus (Pers.) P. Kumm. [†]
Porodisculus pendulus (Schwein. ex Fr.) Murrill [‡, NR]
Porotheleum fimbriatum (Pers.) Fr. [‡, NR]
Psathyrella ammophila (Durieu & Lév.) P.D. Orton [‡, NR]
Psathyrella candolleana (Fr.) Maire [‡, NR]
Psathyrella cernua (Vahl) Hirsch [†]
Psathyrella conopilus (Fr.) A. Pearson & Dennis [†]
Psathyrella echiniceps (G.F. Atk.) A.H. Sm. [‡, NR]
Psathyrella epimyces (Peck) A.H. Sm. [‡, NR]
Psathyrella gracilis (Fr.) Quél. [‡, NR]
Psathyrella hydrophila (Bull.) Maire [‡, NR]
Psathyrella hymenocephala (Peck) A.H. Sm. [‡, NR]
Psathyrella incerta (Peck) A.H. Sm. [‡, NR]
Psathyrella longipes (Peck) A.H. Sm. [†]
Psathyrella piluliformis (Bull.) P.D. Orton [†]
Psathyrella sarcoccephala (Fr.) Singer [†, ‡]
Pseudoclitocybe cyathiformis (Bull.) Singer [†]
Pseudofistulina radicata (Schwein.) Burds. [‡, NR]
Pterula plumosa (Schwein.) Fr. [†, ‡]
Radulomyces confluens (Fr.) M.P. Christ. [‡, NR]
Resinomycena rhododendri (Peck) Redhead & Singer [‡, NR]
Resinipinatus alboniger (Pat.) Singer [‡, NR]
Resinipinatus applicatus (Batsch) Gray [†, ‡]
Rhizomarasmius pyrrhocephalus (Berk.) R.H. Petersen [‡, NR]
Rhodocybe popinalis (Fr.) Singer [†]
Rhodocollybia butyracea (Bull.) Lennox [‡, NR]
Rhodocollybia maculata (Alb. & Schwein.) Singer [†, ‡]
Rhodotus palmatus (Bull.) Maire [†, ‡]
Schizophyllum commune Fr. [†, ‡]
Simocybe centunculus (Fr.) P. Karst. [†, ‡]
Stropharia coronilla (Bull.) Quél. [‡, NR]

Stropharia hardii G.F. Atk. [‡, NR]
Stropharia rugosoannulata Farl. ex Murrill [‡, NR]
Stropharia semiglobata (Batsch) Quél. [†, ‡]
Stropharia viridula (Schaeff.) Morgan [†]
Tetrapyrgos nigripes (Fr.) E. Horak [‡, NR]
Tricholoma album (Schaeff.) P. Kumm. [†]
Tricholoma caligatum (Viv.) Ricken [‡, NR]
Tricholoma columbetta (Fr.) P. Kumm. [‡, NR]
Tricholoma equestre (L.) P. Kumm. [†]
Tricholoma fulvum (Fr.) Bigeard & H. Guill. [†]
Tricholoma fumescens (Peck) Sacc. [†, ‡]
Tricholoma leucophyllum Ovrebo & Tylutki [‡, NR]
Tricholoma resplendens (Fr.) P. Karst. [‡, NR]
Tricholoma saponaceum (Fr.) P. Kumm. [‡, NR]
Tricholoma sciodes (Pers.) C. Martín [‡, NR]
Tricholoma sejunctum (Sowerby) Quél. [†, ‡]
Tricholoma subresplendens (Murrill) Sacc. & Trotter [‡, NR]
Tricholoma subsaponaceum Peck [‡, NR]
Tricholoma sulphureum (Bull.) P. Kumm. [†]
Tricholoma terreum (Schaeff.) P. Kumm. [‡, NR]
Tricholoma virgatum (Fr.) P. Kumm. [‡, NR]
Tricholoma zelleri (D.E. Stuntz & A.H. Sm.) Ovrebo & Tylutki [‡, NR]
Tricholomopsis formosa (Murrill) Singer [‡, NR]
Tricholomopsis rutilans (Schaeff.) Singer [‡, NR]
Tubaria furfuracea (Pers.) Gillet [‡, NR]
Tubaria pellucida (Bull. & Vent.) Sacc. [‡, NR]
Tulostoma berkeleyi Lloyd [‡, NR]
Tulostoma brumale Pers. [‡, NR]
Tulostoma fimbriatum Fr. [‡, NR]
Tulostoma fimbriatum var. *campestre* (Morgan) G. Moreno [‡, NR]
Tulostoma longii Lloyd [‡, NR]
Tulostoma melanocyclum Bres. [‡, NR]
Tulostoma pygmaeum Lloyd [‡, NR]
Tulostoma simulans Lloyd [‡, NR]
Tulostoma striatum G. Cunn. [‡, NR]
Typhula subfasciculata Ellis & Everh [‡, NR]
Vascellum cruciatum (Rostk.) P. Ponce de León [†, ‡]
Vascellum curtisii (Berk.) Kreisel [‡, NR]
Vascellum pratense (Pers.) Kreisel [‡, NR]
Volvariella bombycinia (Schaeff.) Singer [†, ‡]
Volvariella gloiocephala (DC.) Boekhout & Enderle [†, ‡]
Volvariella hypopithys (Fr.) Shaffer [‡, NR]
Volvariella pusilla (Pers.) Singer [†, ‡]
Xeromphalina campanella (Batsch) Kühner & Maire [†, ‡]
Xeromphalina kauffmanii A.H. Sm. [‡, NR]

Xeromphalina tenuipes (Schwein.) A.H. Sm. [‡, NR]
Xerula furfuracea (Peck) Redhead, Ginns & Shoemaker [‡, NR]
Xerula megalospora (Clem.) Redhead, Ginns & Shoemaker [‡, NR]
Xerula limonispora R.H. Petersen [‡, NR]
Xerula longipes (Quél.) Maire [†]
Xerula radicata (Relhan) Dörfelt [†, ‡]
Xerula sinopudens R.H. Petersen & Nagas. [‡, NR]

Atheliales

Athelia arachnoidea (Berk.) Jülich [‡, NR]
Byssocorticium atrovirens (Fr.) Bondartsev & Singer ex Singer [‡, NR]

Auriculariales

Auricularia americana Parmasto & I. Parmasto ex Audet [‡, NR]
Auricularia auricula-judae (Bull.) J. Schröt. [†, ‡]
Seismosarca alba Lloyd [‡, NR]
Tremelodendron schweinitzii G.F. Atk. [†]

Boletales

Astraeus hygrometricus (Pers.) Morgan [†, ‡]
Aureoboletus auriporus (Peck) Pouzar [†, ‡]
Austroboletus gracilis (Peck) Wolfe [†]
Boletellus chrysenteroides (Snell) Snell [‡, NR]
Boletellus russellii (Frost) E.-J. Gilbert [‡, NR]
Boletinus cavipes (Opat.) Kalchbr. [‡, NR]
Boletus auripes Peck [†]
Boletus badius (Fr.) Fr. [†]
Boletus bicolor var. *bicolor* Raddi [†, ‡]
Boletus bicolor var. *subreticulatus* A.H. Sm. & Thiers [‡, NR]
Boletus calopus Pers. [†]
Boletus campestris A.H. Sm. & Thiers [†]
Boletus conicus Rav. [†]
Boletus curtisii Berk. [‡, NR]
Boletus edulis Bull. [†, ‡]
Boletus frostii J.L. Russell [†, ‡]
Boletus fumosiceps (Murrill) Murrill [†]
Boletus glabellus Peck [‡, NR]
Boletus illudens Peck [‡, NR]
Boletus indecisus Peck [†]
Boletus inedulis (Murrill) Murrill [†]
Boletus innixus Frost [†]
Boletus luridus Schaeff. [†]
Boletus luridiformis Rostk. [†]
Boletus magnisporus Frost [†]
Boletus mariae A.H. Sm. & Thiers [‡, NR]
Boletus miniatoolivaceus Frost [‡, NR]

- Boletus miniatopallescens* A.H. Sm. & Thiers [‡, NR]
Boletus modestus Peck [†, ‡]
Boletus nigrellus Peck [†]
Boletus ornatipes Peck [†, ‡]
Boletus pallidus Frost [†, ‡]
Boletus peckii Frost [‡, NR]
Boletus pseudosensibilis A.H. Sm. & Thiers [‡, NR]
Boletus reticulatus Schaeff. [†]
Boletus retipes Berk & M.A. Curtis [†]
Boletus rubropunctus Peck [‡, NR]
Boletus sensibilis Peck [‡, NR]
Boletus speciosus Frost [†]
Boletus subfraternus Coker & Beers [†, ‡]
Boletus subglabripes Peck [†]
Boletus subsanguineus Peck [†, ‡]
Boletus subvelutipes Peck [†, ‡]
Boletus tenax A.H. Sm. & Thiers [‡, NR]
Boletus variipes Peck [‡, NR]
Boletus vermiculosus Peck [†]
Bothia castanella (Peck) Halling [‡, NR]
Chalciporus piperatoides (A.H. Sm. & Thiers) T.J. Baroni & Both [‡, NR]
Chalciporus piperatus (Bull.) Bataille [‡, NR]
Chroogomphus ochraceus (Kauffman) O.K. Mill. [‡, NR]
Coniophora arida (Fr.) P. Karst. [‡, NR]
Coniophora olivacea (Fr.) P. Karst. [‡, NR]
Coniophora puteana (Schumach.) P. Karst. [†]
Cyanoboletus pulverulentus (Opat.) Gelardi [†]
Gomphidius flavipes Peck [‡, NR]
Gyroporus castaneus (Bull.) Quél [†, ‡]
Gyroporus cyanescens (Bull.) Quél [†, ‡]
Gyroporus cyanescens var. *violaceotinctus* Watling [‡, NR]
Gyrodon meruloides (Schwein.) Singer [†, ‡]
Hygrophoropsis aurantiaca (Wulfen) Maire [†, ‡]
Leccinum albellum (Peck) Singer [†, ‡]
Leccinum crocipodium (Letell.) Watling [‡, NR]
Leccinum holopus var. *holopus* (Rostk.) Watling [‡, NR]
Leccinum holopus var. *americanum* A.H. Sm. & Thiers [‡, NR]
Leccinum insigne A.H. Sm., Thiers & Watling [‡, NR]
Leccinum oxydabile (Singer) Singer [‡, NR]
Leccinum pseudoscabrum (Kallenb.) Šutara [‡, NR]
Leccinum rugosiceps (Peck) Singer [‡, NR]
Leccinum scabrum (Bull.) Gray [†, ‡]
Leccinum snellii A.H. Sm., Thiers & Watling [‡, NR]
Leccinum subglabripes (Peck) Singer [‡, NR]
Leccinum subleucophaeum E.A. Dick & Snell [‡, NR]
Leucogyrophana pinastri (Fr.) Ginns & Weresub [‡, NR]
Melanogaster broomeanus Berk. [‡, NR]
Paxillus involutus (Batsch) Fr. [†, ‡]
Phylloporus leucomycelinus Singer [‡, NR]
Phylloporus rhodoxanthus (Schwein.) Bres. [†, ‡]
Pisolithus arrhizus (Scop.) Rauschert [‡, NR]
Plicaturopsis crispa (Pers.) D.A. Reid [†, ‡]
Pseudoboletus parasiticus (Bull.) Šutara [‡, NR]
Pseudomerulius aureus (Fr.) Jülich [‡, NR]
Retiboletus griseus (Frost) Manfr. Binder & Bresinsky [‡, NR]
Rhizophogon roseolus (Corda) Th. Fr. [‡, NR]
Rubinoboletus ballouii (Peck) Heinem. & Ramme-
loo [‡, NR]
Scleroderma areolatum Ehrenb. [†, ‡]
Scleroderma aurantium (L.) Pers. [‡, NR]
Scleroderma bovista Fr. [‡, NR]
Scleroderma cepa Pers. [‡, NR]
Scleroderma citrinum Pers. [†, ‡]
Scleroderma flavidum Ellis & Everh. [‡, NR]
Scleroderma polyrhizum (J.F. Gmel.) Pers. [†, ‡]
Scleroderma septentrionale Jeppson [‡, NR]
Scleroderma verrucosum (Bull.) Pers. [†]
Serpula lacrymans (Wulff) P. Karst. [†, ‡]
Strobilomyces confusus Singer [‡, NR]
Strobilomyces strobilaceus (Scop.) Berk. [†, ‡]
Suillus acidus (Peck) Singer [‡, NR]
Suillus americanus (Peck) Snell [‡, NR]
Suillus collinitus (Fr.) Kuntze [†]
Suillus granulatus (L.) Roussel [‡, NR]
Suillus grevillei (Klotzsch) Singer [‡, NR]
Suillus hirtellus (Peck) Kuntze [‡, NR]
Suillus luteus (L.) Roussel [†, ‡]
Suillus pictus A.H. Sm. & Thiers [†, ‡]
Suillus purpureus (Fr. & Hök) Kuntze [†]
Suillus salmonicolor (Frost) Halling [‡, NR]
Suillus subaureus (Peck) Snell [‡, NR]
Tapinella atrotomentosa (Batsch) Šutara [‡, NR]
Tapinella panuoides (Fr.) E.-J. Gilbert [†, ‡]
Tylopilus alboater (Schwein.) Murrill [‡, NR]
Tylopilus eximius (Peck) Singer [†]
Tylopilus felleus (Bull.) P. Karst. [†, ‡]
Tylopilus indecisus (Peck) Murrill [‡, NR]
Tylopilus rubrobrunneus Mazzer & A.H. Sm. [‡, NR]
Tylopilus sordidus (Frost) A.H. Sm. & Thiers [‡, NR]
Veligaster nitidus (Berk.) Guzmán & Tapia [†]
Xanthoconium affine (Peck) Singer [†, ‡]
Xanthoconium affine var. *maculosus* (Peck) Singer [‡, NR]

- Xanthoconium purpureum* Snell & E.A. Dick [‡, NR]
Xanthoconium separans (Peck) Halling & Both [†, ‡]
Xerocomellus chrysenteron (Bull.) Šutara [†, ‡]
Xerocomus rubellus (Krombh.) Quél. [†, ‡]
Xerocomus subtomentosus (L.) Quél. [†, ‡]
Xerocomus truncatus Singer, Snell & E.A. Dick [‡, NR]
- Cantharellales**
- Botryobasidium medium* J. Erikss. [‡, NR]
Botryobasidium subcoronatum (Höhn. & Litsch.) Donk [‡, NR]
Botryobasidium vagum (Berk. & M.A. Curtis) D.P. Rogers [‡, NR]
Botryohypothecus isabellinus (Fr.) J. Erikss. [‡, NR]
Cantharellus chicagoensis Leacock, J. Riddell, Rui Zhang & G.M. Muell. [†, ‡]
Cantharellus cibarius Fr. [†, ‡]
Cantharellus cinnabarinus (Schwein.) Schwein. [†, ‡]
Cantharellus flavus M.J. Foltz & T.J. Volk [‡, NR]
Cantharellus ignicolor R.H. Petersen [‡, NR]
Cantharellus infundibuliformis (Scop.) Fr. [†, ‡]
Cantharellus lateritius (Berk.) Singer [†, ‡]
Cantharellus lutescens (Pers.) Fr. [†]
Cantharellus minor Peck [†, ‡]
Cantharellus tubaeformis Fr. [†]
Cantharellus wrightii Berk. & M.A. Curtis [†]
Ceratobasidium stevensii (Burt) P.H.B. Talbot [‡, NR]
Clavulina amethystina (Bull.) Donk [†]
Clavulina cinerea (Bull.) J. Schröt. [†, ‡]
Clavulina cristata (Holmsk.) J. Schröt. [†]
Craterellus convolvulatus (A.H. Sm.) Eyssart. & Buyck [‡, NR]
Craterellus cornucopioides (L.) Pers. [†, ‡]
Craterellus dubius Peck [‡, NR]
Craterellus fallax A.H. Sm. [‡, NR]
Haplotrichum croceum (Mont.) Partr. & Morgan-Jones [‡, NR]
Haplotrichum curtisii (Berk.) Hol.-Jech. [†, ‡]
Haplotrichum simile (Berk.) Hol.-Jech. [‡, NR]
Hydnellum aurantiacum (Batsch) P. Karst. [‡, NR]
Hydnellum caeruleum (Hornem.) P. Karst. [‡, NR]
Hydnellum concrescens (Pers.) Banker [‡, NR]
Hydnellum peckii Banker [‡, NR]
Hydnellum scrobiculatum (Fr.) P. Karst. [‡, NR]
Hydnellum spongiosipes (Peck) Pouzar [‡, NR]
Hydnellum albiceps Berk. & Ravenel ex Cooke [‡, NR]
- Hydnellum albidum* Peck [†, ‡]
Hydnellum combinans Peck [†]
Hydnellum flabelliforme Berk. [‡, NR]
Hydnellum membranaceum Bull. [‡, NR]
Hydnellum mucidum Pers. ex J.F. Gmel. [†]
Hydnellum repandum L. [†, ‡]
Hydnellum umbilicatum Peck [‡, NR]
Hydnellum velutinum Fr. [‡, NR]
Multiclavula mucida (Pers.) R.H. Petersen [†, ‡]
Pseudocraterellus undulatus (Pers.) Rauschert [‡, NR]
Rhizoctonia solani J.G. Kühn [†, ‡]
Tulasnella bifrons Bourdot & Galzin [‡, NR]
Tulasnella pinicola Bres. [‡, NR]
Tulasnella pruinosa Bourdot & Galzin [‡, NR]
Tulasnella violacea (Johan-Olsen) Juel [‡, NR]
Tulasnella violaea (Quél.) Bourdot & Galzin [‡, NR]
- Corticiales**
- Corticium roseocarneum* (Schwein.) Hjortstam [†, ‡]
Corticium roseum Pers. [‡, NR]
Cytidia salicina (Fr.) Burt [‡, NR]
Dendrothele acerina (Pers.) P.A. Lemke [‡, NR]
Dendrothele candida (Schwein.) P.A. Lemke [†, ‡]
Dendrothele macrodens (Coker) P.A. Lemke [‡, NR]
Dendrothele nivosa (Berk. & M.A. Curtis ex Höhn. & Litsch.) P.A. Lemke [†, ‡]
Mycolindneria trachyspora (Bourdot & Galzin) Rauschert [‡, NR]
Punctularia strigosozonata (Schwein.) P.H.B. Talbot [‡, NR]
- Dacrymycetales**
- Calocera cornea* (Batsch) Fr. [†, ‡]
Dacrymyces australis Lloyd [‡, NR]
Dacrymyces chrysospermus Berk. & M.A. Curtis [‡, NR]
Dacrymyces ellistii Coker [‡, NR]
Dacrymyces stillatus Nees [‡, NR]
Dacryopinax spathularia (Schwein.) G.W. Martin [†, ‡]
- Exobasidiales**
- Exobasidium symploci* Ellis & G. Martin [‡, NR]
Exobasidium vaccinii (Fuckel) Woronin [†, ‡]
- Gastrales**
- Gastrum arenarium* Lloyd [‡, NR]
Gastrum coronatum Pers. [‡, NR]
Gastrum elegans Vittad. [‡, NR]
Gastrum fimbriatum Fr. [‡, NR]

- Geastrum lageniforme* Vittad. [‡, NR]
Geastrum minimum Schwein. [†, ‡]
Geastrum morganii Lloyd [‡, NR]
Geastrum rufescens Pers. [‡, NR]
Geastrum saccatum Fr. [†, ‡]
Geastrum triplex Jungh. [†, ‡]
Geastrum velutinum Morgan [‡, NR]
Myriostoma coliforme (Dicks.) Corda [‡, NR]
Sphaerobolus stellatus Tode [‡, NR]

Gloeophyllales

- Gloeophyllum sepiarium* (Wulff) P. Karst. [†, ‡]
Gloeophyllum trabeum (Pers.) Murrill [†, ‡]
Neolentinus suffrutescens (Brot.) T.W. May &
A.E. Wood [†, ‡]

Gomphales

- Clavariadelphus pistillaris* (L.) Donk [†, ‡]
Corallium formosum (Pers.) G. Hahn [†]
Gomphus floccosus (Schwein.) Singer [‡, NR]
Kavinia himantia (Schwein.) J. Erikss. [‡, NR]
Lentaria byssiseda Corner [‡, NR]
Lentaria micheneri (Berk. & M.A. Curtis) Corner
[‡, NR]
Lentaria patouillardii (Bres.) Corner [‡, NR]
Ramaria abietina (Pers.) Quél. [‡, NR]
Ramaria apiculata (Fr.) Donk [‡, NR]
Ramaria aurea (Schaeff.) Quél. [†, ‡]
Ramaria botrytis (Pers.) Ricken [‡, NR]
Ramaria decurrens (Pers.) R.H. Petersen [‡, NR]
Ramaria gracilis (Pers.) Quél. [†]
Ramaria murrillii (Coker) Corner [†]
Ramaria obtusissima (Peck) Corner [‡, NR]
Ramaria stricta (Pers.) Quél. [‡, NR]
Ramaria stricta var. *concolor* Corner [‡, NR]
Ramaria suecica (Fr.) Donk [†]
Ramaricium polyporoideum (Berk. & M.A. Curtis)
Ginns [‡, NR]
Ramariopsis kunzei (Fr.) Corner [‡, NR]

Hymenochaetales

- Basidioradulum radula* (Fr.) Nobles [†]
Coltricia cinnamomea (Jacq.) Murrill [†, ‡]
Coltricia focicola (Berk. & M.A. Curtis) Murrill
[†]
Coltricia montagnei (Fr.) Murrill [‡, NR]
Coltricia perennis (L.) Murrill [†, ‡]
Cotylidia pallida (Pers.) Biodin [†]
Fibricium rude (P. Karst.) Jülich [‡, NR]
Fulvifomes johnsonianus (Murrill) Y.C. Dai [‡,
NR]
Fuscoporia gilva (Schwein.) T. Wagner & M.
Fisch. [†, ‡]
Hydnochaete olivacea (Schwein.) Banker [†, ‡]

- Hymenochaete cinnamomea* subsp. *spreta* (Peck)
Parmasto [†, ‡]
Hymenochaete corrugata (Fr.) Lév. [‡, NR]
Hymenochaete curtisii (Berk.) Morgan [‡, NR]
Hymenochaete fuliginosa (Pers.) Lév. [†, ‡]
Hymenochaete rubiginosa (Dicks.) Lév. [†, ‡]
Hyphodontia arguta (Fr.) J. Erikss. [‡, NR]
Hyphodontia aspera (Fr.) J. Erikss. [‡, NR]
Hyphodontia bugellensis (Ces.) J. Erikss. [‡, NR]
Hyphodontia latitans (Bourdot & Galzin) Ginns &
M.N.L. Lefebvre [‡, NR]
Hyphodontia quericina (Pers.) J. Erikss. [‡, NR]
Hyphodontia spathulata (Schrad.) Parmasto [‡,
NR]
Inonotus andersonii (Ellis & Everh.) Cerný [‡, NR]
Inonotus cuticularis (Bull.) P. Karst. [†, ‡]
Inonotus dryadeus Fr. [†]
Inonotus glomeratus (Peck) Murrill [‡, NR]
Inonotus hispidus (Bull.) P. Karst. [†, ‡]
Inonotus obliquus (Ach. ex Pers.) Pilát [†]
Inonotus radiatus (Sowerby) P. Karst. [†, ‡]
Onnia tomentosa (Fr.) P. Karst. [‡, NR]
Oxyporus corticola (Fr.) Ryvarden [†, ‡]
Oxyporus latemarginatus (Durieu & Mont.) Donk
[†, ‡]
Oxyporus pellicula (Jungh.) Ryvarden [‡, NR]
Oxyporus populinus (Schumach.) Donk [†, ‡]
Phellinus conchatus (Pers.) Quél. [†, ‡]
Phellinus contiguus (Pers.) Pat. [‡, NR]
Phellinus everhartii (Ellis & Galloway) A. Ames
[†, ‡]
Phellinus ferrugineofuscus (P. Karst.) Bourdot [‡,
NR]
Phellinus ferruginosus (Schrad.) Pat. [‡, NR]
Phellinus gilvus (Schwein.) Pat. [†, ‡]
Phellinus igniarius (L.) Quél. [†, ‡]
Phellinus laevigatus (Fr.) Bourdot & Galzin [‡,
NR]
Phellinus nigricans (Fr.) P. Karst. [†, ‡]
Phellinus prunicola (Murrill) Gilb. [‡, NR]
Phellinus punctatus (P. Karst.) Pilát [‡, NR]
Phellinus ribis (Schumach.) Quél. [†, ‡]
Phellinus rimosus (Berk.) Pilát [‡, NR]
Phellinus robiniae (Murrill) A. Ames [‡, NR]
Phellinus robustus (P. Karst.) Bourdot & Galzin
[‡, NR]
Phellinus viticola (Schwein.) Donk [‡, NR]
Polystictus hirsutus var. *marginatus* (Bres. ex
Schulzer) Sacc. [†, ‡]
Porodaedalea pini (Brot.) Murrill [†]
Pseudochaete tabacina (Sowerby) T. Wagner &
M. Fisch. [‡, NR]
Pycnoporus cinnabarinus (Jacq.) P. Karst. [†, ‡]
Rickenella fibula (Bull.) Raithelh. [‡, NR]

Schizopora paradoxa (Schrad.) Donk [†, ‡]
Schizopora radula (Pers.) Hallenb. [‡, NR]
Skvortzovia furfurella (Bres.) Bononi & Hjortstam
[‡, NR]

Phallales

Lysurus cruciatus (Lepr. & Mont.) Henn. [†, ‡]
Lysurus periphragmoides (Klotzsch) Dring [†, ‡]
Mutinus caninus (Huds.) Fr. [†, ‡]
Mutinus elegans (Mont.) E. Fisch [†]
Mutinus ravenelii (Berk. & M.A. Curtis) E. Fisch.
[†]
Phallus hadriani Vent. [‡, NR]
Phallus impudicus L. [†, ‡]
Phallus impudicus var. *togatus* (Kalchbr.) Cos-
tantin & L.M. Dufour [†, ‡]
Phallus ravenelii Berk. & M.A. Curtis [‡, NR]

Polyporales

Abortiporus biennis (Bull.) Singer [†, ‡]
Antrodia albida (Fr.) Donk [†, ‡]
Antrodia albobrunnea (Romell) Ryvarden [‡, NR]
Antrodia juniperina (Murrill) Niemelä & Ryvar-
den [‡, NR]
Antrodia lenis (P. Karst.) Ryvarden [‡, NR]
Antrodia malicola (Berk. & M.A. Curtis) Donk [‡,
NR]
Antrodia oleracea (R.W. Davidson & Lombard)
Ryvarden [‡, NR]
Antrodia sinuosa (Fr.) P. Karst. [‡, NR]
Antrodia xantha (Fr.) Ryvarden [‡, NR]
Antrodiella semisupina (Berk. & M.A. Curtis)
Ryvarden [‡, NR]
Aurantiporus fissilis (Berk. & M.A. Curtis) H.
Jahn ex Ryvarden [‡, NR]
Bjerkandera adusta (Willd.) P. Karst. [†, ‡]
Bjerkandera fumosa (Pers.) P. Karst. [†, ‡]
Byssomerulius incarnatus (Schwein.) Gilb. [†, ‡]
Ceriporia purpurea (Fr.) Donk [‡, NR]
Ceriporia spissa (Schwein. ex Fr.) Rajchenb. [‡,
NR]
Ceriporia tarda (Berk.) Ginns [†, ‡]
Ceriporia viridans (Berk. & Broome) Donk [‡,
NR]
Ceriporiopsis mucida (Pers.) Gilb. & Ryvarden [‡,
NR]
Ceriporiopsis subrufa (Ellis & Dearn.) Ginns [‡,
NR]
Cerrena unicolor (Bull.) Murrill [†, ‡]
Climacocystis borealis (Fr.) Kotl. & Pouzar [‡,
NR]
Climacodon pulcherrimus (Berk. & M.A. Curtis)
Nikol. [†, ‡]
Climacodon septentrionalis (Fr.) P. Karst. [†, ‡]

Coriolopsis floccosa (Jungh.) Ryvarden [†, ‡]
Coriolopsis gallica (Fr.) Ryvarden [†, ‡]
Coriolopsis trogii (Berk.) Domanski [‡, NR]
Dacryobolus sudans (Alb. & Schwein.) Fr. [‡, NR]
Daedalea quercina (L.) Pers. [†, ‡]
Daedaleopsis confragosa var. *confragosa* (Bolton)
J. Schröt. [†, ‡]
Daedaleopsis septentrionalis (P. Karst.) Niemelä
[‡, NR]
Datronia mollis (Sommerf.) Donk [‡, NR]
Datronia scutellata (Schwein.) Gilb. & Ryvarden
[‡, NR]
Diplomitoporus overholtsii (Pilát) Gilb. & Ryvar-
den [‡, NR]
Erastia salmonicolor (Berk. & M.A. Curtis)
Niemelä & Kinnunen [‡, NR]
Favolus albidus Massee [‡, NR]
Fibroporia radiculosa (Peck) Parmasto [‡, NR]
Fomes fomentarius (L.) Fr. [†, ‡]
Fomes fulvus (Scop.) Gillet [†, ‡]
Fomes meliae (Underw.) Murrill [‡, NR]
Fomitopsis cajanderi (P. Karst.) Kotl. & Pouzar [†]
Fomitopsis durescens (Overh. ex J. Lowe) Gilb. &
Ryvarden [‡, NR]
Fomitopsis palustris (Berk. & M.A. Curtis) Gilb.
& Ryvarden [‡, NR]
Fomitopsis spraguei (Berk. & M.A. Curtis) Gilb. &
Ryvarden [†, ‡]
Fuscocerrena portoricensis (Spreng. ex Fr.) Ry-
varden [‡, NR]
Ganoderma australe (Fr.) Pat. [‡, NR]
Ganoderma applanatum (Pers.) Pat. [†, ‡]
Ganoderma cupreolaccatum (Kalchbr.) Z.
Igmándy [‡, NR]
Ganoderma curtisii (Berk.) Murrill [‡, NR]
Ganoderma lipsiense (Batsch) G.F. Atk. [†, ‡]
Ganoderma lucidum (Curtis) P. Karst. [†, ‡]
Ganoderma resinaceum Boud. [‡, NR]
Ganoderma sessile Murrill [†, ‡]
Ganoderma tsugae Murrill [‡, NR]
Globiformes graveolens (Schwein.) Murrill [†, ‡]
Gloeoporus dichrous (Fr.) Bres. [†, ‡]
Gloeoporus thelephoroides (Hook.) G. Cunn. [†, ‡]
Grifola frondosa (Dicks.) Gray [†, ‡]
Hapalopilus croceus (Pers.) Bondartsev & Singer
[†, ‡]
Hapalopilus nidulans (Fr.) P. Karst. [‡, NR]
Hapalopilus rutilans (Pers.) Murrill [‡, NR]
Haploporus odorus (Sommerf.) Bondartsev &
Singer ex Singer [‡, NR]
Helicybe sulcata (Berk.) Redhead & Ginns [‡,
NR]
Hjortstamia crassa (Lév.) Boidin & Gilles [‡, NR]

- Hyphoderma litschaueri* (Burt) J. Erikss. & Å. Strid [‡, NR]
- Hyphoderma mutatum* (Peck) Donk [†, ‡]
- Hyphoderma populneum* (Peck) Donk [‡, NR]
- Hyphoderma puberum* (Fr.) Wallr. [‡, NR]
- Hyphoderma rubropallens* (Schwein.) Ginns [‡, NR]
- Hyphoderma setigerum* (Fr.) Donk [‡, NR]
- Hyphoderma tenue* (Pat.) Donk [‡, NR]
- Hypochnicium lundellii* (Bourdot) J. Erikss. [‡, NR]
- Inonotus andersonii* (Ellis & Everh.) Cerný [‡, NR]
- Inonotus cuticularis* (Bull.) P. Karst. [†]
- Inonotus hispidus* (Bull.) P. Karst. [†]
- Inocutis rheades* (Pers.) Fiasson & Niemelä [†]
- Irpea fuscescens* Schwein. [†, ‡]
- Irpea lacteus* (Fr.) Fr. [†, ‡]
- Ischnoderma benzoinum* (Wahlenb.) P. Karst. [†, ‡]
- Junguhnia nitida* (Pers.) Ryvarden [‡, NR]
- Laeitporus cincinnatus* (Morgan) Burds. [‡, NR]
- Laeitporus sulphureus* (Bull.) Murrill [†, ‡]
- Lentinus levis* (Berk. & M.A. Curtis) Murrill [‡, NR]
- Lentinus strigosus* Fr. [†, ‡]
- Lentinus tigrinus* (Bull.) Fr. [†, ‡]
- Lenzites betulina* (L.) Fr. [†, ‡]
- Lenzites elegans* (Spreng.) Pat. [†, ‡]
- Lenzites flaccida* Fr. [†]
- Lenzites stereoides* (Fr.) Ryvarden [‡, NR]
- Lopharia cinerascens* (Schwein.) G. Cunn. [†, ‡]
- Melanoporia nigra* (Berk.) Murrill [†, ‡]
- Meripilus giganteus* (Pers.) P. Karst. [†, ‡]
- Microporellus obovatus* (Jungh.) Ryvarden [‡, NR]
- Mutatoderma heterocystidiatum* (Burt) C.E. Gómez [‡, NR]
- Mycocacia aurea* (Fr.) J. Erikss. & Ryvarden [‡, NR]
- Mycocacia uda* (Fr.) Donk [‡, NR]
- Mycorrhaphium adustulum* (Banker) Ryvarden [‡, NR]
- Mycorrhaphium adustum* (Schwein.) Maas Geest. [†, ‡]
- Osteina obducta* (Berk.) Donk [‡, NR]
- Panus conchatus* (Bull.) Fr. [†, ‡]
- Perenniporia fraxinea* (Bull.) Ryvarden [‡, NR]
- Perenniporia fraxinophila* (Peck) Ryvarden [‡, NR]
- Perenniporia medulla-panis* (Jacq.) Donk [†, ‡]
- Perenniporia ohiensis* (Berk.) Ryvarden [‡, NR]
- Perenniporia robiniophila* (Murrill) Ryvarden [†, ‡]
- Perenniporia subacida* (Peck) Donk [‡, NR]
- Perenniporia tenuis* (Schwein.) Ryvarden [‡, NR]
- Perenniporia tenuis* var. *pulchella* (Schwein.) Gilb. & Ryvarden [‡, NR]
- Perenniporia variegata* Ryvarden & Gilb. [‡, NR]
- Phaeolus schweinitzii* (Fr.) Pat. [‡, NR]
- Phanerochaete affinis* (Burt) Parmasto [‡, NR]
- Phanerochaete avellanea* (Bres.) J. Erikss. & Hjortstam [‡, NR]
- Phanerochaete burtii* (Romell) Parmasto [‡, NR]
- Phanerochaete chrysorhiza* (Torr.) Budington & Gilb. [†, ‡]
- Phanerochaete filamentosa* (Berk. & M.A. Curtis) Burds. [‡, NR]
- Phanerochaete laevis* (Fr.) J. Erikss. & Ryvarden [‡, NR]
- Phanerochaete sanguinea* (Fr.) Pouzar [‡, NR]
- Phanerochaete sordida* (P. Karst.) J. Erikss. & Ryvarden [†, ‡]
- Phanerochaete velutina* (DC.) P. Karst. [‡, NR]
- Phlebia albida* H. Post [‡, NR]
- Phlebia chrysocreas* (Berk. & M.A. Curtis) Burds. [‡, NR]
- Phlebia cinnabarina* Fr. [‡, NR]
- Phlebia hydnoides* Schwein. [†]
- Phlebia livida* (Pers.) Bres. [‡, NR]
- Phlebia ludoviciana* (Burt) Nakasone & Burds. [‡, NR]
- Phlebia queletii* (Bourdot & Galzin) M.P. Christ. [‡, NR]
- Phlebia radiata* Fr. [†, ‡]
- Phlebia tremellosa* (Schrad.) Nakasone & Burds. [†, ‡]
- Phlebiella tulasnelloidea* (Höhn. & Litsch.) Ginns & M.N.L. Lefebvre [‡, NR]
- Phlebiella vaga* (Fr.) P. Karst. [‡, NR]
- Phlebiopsis flavidoolba* (Cooke) Hjortstam [‡, NR]
- Phlebiopsis gigantea* (Fr.) Jülich [‡, NR]
- Piptoporus betulinus* (Bull.) P. Karst. [‡, NR]
- Podoscypha ravenelii* (Berk. & M.A. Curtis) Pat. [†]
- Polyporus admirabilis* Peck [†, ‡]
- Polyporus albus* (Huds.) Fr. [‡, NR]
- Polyporus alveolaris* (DC.) Bondartsev & Singer [†, ‡]
- Polyporus arcularius* (Batsch) Fr. [†, ‡]
- Polyporus badius* (Pers.) Schwein. [†, ‡]
- Polyporus brumalis* (Pers.) Fr. [†, ‡]
- Polyporus epileucus* Fr. [†, ‡]
- Polyporus incertus* Pers. [†]
- Polyporus melanopus* (Pers.) Fr. [†, ‡]
- Polyporus osseus* Kalchbr. [†]
- Polyporus paragamenus* Fr. [†]
- Polyporus radicatus* Schwein. [†, ‡]

- Polyporus rigidus* Lév. [‡, NR]
Polyporus salignus (Fr.) Fr. [‡, NR]
Polyporus squamosus (Huds.) Fr. [†, ‡]
Polyporus varius (Pers.) Fr. [†, ‡]
Poria attenuata (Peck) Sacc. [†]
Poria bresadolae Bourdot & Galzin [‡, NR]
Poria discolor Overh. [‡, NR]
Poria homaema Sacc. [‡, NR]
Poria niphodes (Berk. & Broome) Sacc. [‡, NR]
Poria vitellina (Schwein.) Sacc. [‡, NR]
Poria xantholoma (Schwein.) Cooke [‡, NR]
Poronidulus conchifer (Schwein.) Murrill [†, ‡]
Postia caesia (Schrad.) P. Karst. [‡, NR]
Postia floriformis (Quél.) Jülich [‡, NR]
Postia fragilis (Fr.) Jülich [‡, NR]
Postia guttulata (Peck) Jülich [‡, NR]
Postia tephroleuca (Fr.) Jülich [†, ‡]
Pycnoporellus fulgens (Fr.) Donk [‡, NR]
Pycnoporus cinnabarinus (Jacq.) P. Karst. [†, ‡]
Pycnoporus sanguineus (L.) Murrill [†, ‡]
Rigidoporus crocatus (Pat.) Ryvarden [‡, NR]
Rigidoporus lineatus (Pers.) Ryvarden [‡, NR]
Rigidoporus undatus (Pers.) Donk [†, ‡]
Rigidoporus vitreus (Pers.) Donk [‡, NR]
Sarcodontia setosa (Pers.) Donk [‡, NR]
Scopuloides hydnoides (Cooke & Massee) Hjortstam & Ryvarden [‡, NR]
Skeletocutis alutacea (J. Lowe) Jean Keller [‡, NR]
Skeletocutis nivea (Jungh.) Jean Keller [†, ‡]
Skeletocutis odora (Peck ex Sacc.) Ginns [‡, NR]
Solenia conferta Burt [†]
Sparassis crispa (Wulfen) Fr. [†]
Sparassis laminosa Fr. [‡, NR]
Spongipellis delectans (Peck) Murrill [†, ‡]
Spongipellis pachyodon (Pers.) Kotl. & Pouzar [†, ‡]
Spongipellis spumeus (Sowerby) Pat. [‡, NR]
Spongipellis unicolor (Schwein.) Murrill [†, ‡]
Sporotrichum byssinum Link [‡, NR]
Sporotrichum chryseum Peck [†, ‡]
Steccherinum ciliolatum (Berk. & M.A. Curtis) Gilb. & Budington [‡, NR]
Steccherinum fimbriatum (Pers.) J. Erikss. [‡, NR]
Steccherinum ochraceum (Pers. ex J.F. Gmel.) Gray [†, ‡]
Terana caerulea (Schrad. ex Lam.) Kuntze [†]
Trametes aesculi (Fr.) Justo [‡, NR]
Trametes cubensis (Mont.) Sacc. [‡, NR]
Trametes gibbosa (Pers.) Fr. [‡, NR]
Trametes hirsuta (Wulfen) Pilát [†, ‡]
Trametes ochracea (Pers.) Gilb. & Ryvarden [†, ‡]
Trametes pubescens (Schumach.) Pilát [†, ‡]
Trametes suaveolens (L.) Fr. [‡, NR]
- Trametes versicolor* (L.) Lloyd [†, ‡]
Trametes villosa (Sw.) Kreisel [‡, NR]
Trametopsis cervina (Schwein.) Tomsovský [‡, NR]
Trichaptum abietinum (Pers. ex J.F. Gmel.) Ryvarden [†, ‡]
Trichaptum biforme (Fr.) Ryvarden [†, ‡]
Trichaptum subchartaceum (Murrill) Ryvarden [‡, NR]
Tyromyces chioneus (Fr.) P. Karst. [†, ‡]
Tyromyces fumidiceps G.F. Atk. [‡, NR]
Tyromyces galactinus (Berk.) J. Lowe [‡, NR]
Wolfiporia dilatohypa Ryvarden & Gilb. [‡, NR]
Wolfiporia extensa (Peck) Ginns [†, ‡]

Russulales

- Albatrellus cristatus* (Schaeff.) Kotl. & Pouzar [†, ‡]
Aleurodiscus andinus Núñez & Ryvarden [‡, NR]
Aleurodiscus oakesii (Berk. & M.A. Curtis) Pat. [†, ‡]
Artomyces pyxidatus (Pers.) Jülich [†, ‡]
Asterostroma andinum Pat. [‡, NR]
Auriscalpium vulgare Gray [†, ‡]
Bondarzewia berkeleyi (Fr.) Bondartsev & Singer [†, ‡]
Chondrostereum purpureum (Pers.) Pouzar [‡, NR]
Conferticium ochraceum (Fr.) Hallenb. [‡, NR]
Dichostereum granulosum (Fr.) Boidin & Lanq. [†]
Gloeopeniophorella convolvens (P. Karst.) Boidin, Lanq. & Gilles [‡, NR]
Gloiodon strigosus (Sw.) P. Karst. [†]
Hericium coralloides (Scop.) Pers. [†, ‡]
Hericium erinaceus (Bull.) Pers. [†, ‡]
Heterobasidion annosum (Fr.) Bref. [‡, NR]
Lactarius affinis Peck [†]
Lactarius aquizonatus Kytöv. [‡, NR]
Lactarius atroviridis Peck [‡, NR]
Lactarius camphoratus (Bull.) Fr. [†, ‡]
Lactarius carbonicola A.H. Sm. [‡, NR]
Lactarius chrysorrheus Fr. [†, ‡]
Lactarius corrugis Peck [†]
Lactarius deceptivus Peck [†, ‡]
Lactarius deliciosus (L.) Gray [†]
Lactarius distans Peck [†]
Lactarius fragilis (Burl.) Hesler & A.H. Sm. [‡, NR]
Lactarius fuliginellus A.H. Sm. & Hesler [‡, NR]
Lactarius fumosus Peck [‡, NR]
Lactarius gerardii Peck [†, ‡]
Lactarius glaucescens Crossl. [‡, NR]
Lactarius griseus Peck [‡, NR]

- Lactarius hygrophoroides* var. *hygrophoroides*
Berk. & M.A. Curtis [†, ♀]
Lactarius hygrophoroides var. *lavandulaceus* Hesler & A.H. Sm. [‡, NR]
Lactarius hysginus (Fr.) Fr. [†]
Lactarius imperceptus Beardslee & Burl. [‡, NR]
Lactarius indigo (Schwein.) Fr. [†, ♀]
Lactarius insulsus (Fr.) Fr. [†, ♀]
Lactarius lignyotus Fr. [†, ♀]
Lactarius maculatipes Burl. [‡, NR]
Lactarius maculatus Peck [‡, NR]
Lactarius mutabilis Peck [‡, NR]
Lactarius piperatus (L.) Pers. [†, ♀]
Lactarius plumbeus (Bull.) Fr. [†]
Lactarius psammicola A.H. Sm. [‡, NR]
Lactarius pubescens Fr. [‡, NR]
Lactarius pyrogalus (Bull.) Fr. [†, ♀]
Lactarius quietus (Fr.) Fr. [‡, NR]
Lactarius quietus var. *incanus* Hesler & A.H. Sm. [‡, NR]
Lactarius rufus (Scop.) Fr. [†]
Lactarius scrobiculatus (Scop.) Fr. [†, ♀]
Lactarius serifluus (DC.) Fr. [†, ♀]
Lactarius sordidus Peck [†]
Lactarius subplinthogalus Coker [‡, NR]
Lactarius subpurpureus Peck [‡, NR]
Lactarius subserifluus Longyear [‡, NR]
Lactarius subvernalis var. *cokeri* (A.H. Sm. & Hesler) Hesler & A.H. Sm. [‡, NR]
Lactarius tabidus Fr. [†]
Lactarius trivialis (Fr.) Fr. [†, ♀]
Lactarius uvidus (Fr.) Fr. [†]
Lactarius vellereus var. *vellereus* [†]
Lactarius vollemus var. *flavus* Hesler & A.H. Sm. [‡, NR]
Lactarius vollemus var. *volemus* (Fr.) Fr. [†, ♀]
Lactarius zonarius (Bull.) Fr. [‡, NR]
Lactifluus subvellereus (Peck) Nuytinck [‡, NR]
Lactifluus vellereus (Fr.) Kuntze [‡, NR]
Laxitextum bicolor (Pers.) Lentz [†, ♀]
Leia stratos (Berk.) Banker [‡, NR]
Lentinellus cochleatus (Pers.) P. Karst. [‡, NR]
Lentinellus flabelliformis (Bolton) S. Ito [‡, NR]
Lentinellus micheneri (Berk. & M.A. Curtis) Pegler [‡, NR]
Lentinellus ursinus (Fr.) Kühner [†, ♀]
Lentinellus vulpinus (Sowerby) Kühner & Maire [†]
Peniophora albobadia (Schwein.) Boidin [‡, NR]
Peniophora cinerea (Pers.) Cooke [†, ♀]
Peniophora erikssonii Boidin [‡, NR]
Peniophora erumpens (Burt) Boidin [‡, NR]
Peniophora filamentosa (Berk. & M.A. Curtis) Moffatt [‡, NR]
Peniophora limitata (Chaillet ex Fr.) Cooke [‡, NR]
Peniophora nuda (Fr.) Bres. [‡, NR]
Peniophora rufa (Fr.) Boidin [‡, NR]
Peniophora versiformis (Berk. & M.A. Curtis) Bourdot & Galzin [‡, NR]
Peniophora violaceolivida (Sommerf.) Massee [‡, NR]
Russula adulterina Fr. [‡, NR]
Russula adusta (Pers.) Fr. [‡, NR]
Russula aeruginea Lindblad [†, ♀]
Russula albonigra (Krombh.) Fr. [‡, NR]
Russula alutacea (Pers.) Fr. [†]
Russula amoena Romagn. [†, ♀]
Russula anomala Peck [‡, NR]
Russula atropurpurea (Krombh.) Britzelm. [†, ♀]
Russula ballouii Peck [‡, NR]
Russula barlae Quél. [‡, NR]
Russula basifurcata Peck [†]
Russula brevipes var. *brevipes* Peck [†, ♀]
Russula brunnealba De Marb. [‡, NR]
Russula brunneola Burl. [‡, NR]
Russula brunneoviolacea Crawshay [‡, NR]
Russula chloroides (Krombh.) Bres. [‡, NR]
Russula cinerascens Beardslee [‡, NR]
Russula claroflava Grove [‡, NR]
Russula compacta Frost [†, ♀]
Russula corinthii-rubra Burl. [‡, NR]
Russula cremeirosea Murrill [‡, NR]
Russula crustosa Peck [†, ♀]
Russula cyanoxantha (Schaeff.) Fr. [†, ♀]
Russula cyanoxantha var. *variata* (Banning) Singer [†, ♀]
Russula cystidiosa Murrill [‡, NR]
Russula decolorans (Fr.) Fr. [†]
Russula delica Fr. [†]
Russula densifolia Secr. ex Gillet [†]
Russula disparilis Burl. [‡, NR]
Russula emetica (Schaeff.) Pers. [†, ♀]
Russula flavidula Frost [‡, NR]
Russula flavisiccans Bills [‡, NR]
Russula foetens Pers. [†]
Russula font-queri Singer ex Singer [‡, NR]
Russula fragilis Fr. [†]
Russula fragiloides Murrill [‡, NR]
Russula fragrantissima Romagn. [‡, NR]
Russula fucosa Burl. [‡, NR]
Russula furcata (Lam.) Pers. [†]
Russula galochroa (Fr.) Fr. [‡, NR]
Russula granulata Peck [†]
Russula heterophylla (Fr.) Fr. [‡, NR]
Russula humidicola Burl. [‡, NR]
Russula incarnaticeps Murrill [‡, NR]

- Russula ionochlora* Romagn. [‡, NR]
Russula languida Cern. & H. Raab [†]
Russula laurocerasi Melzer [‡, NR]
Russula luteobasis Peck [‡, NR]
Russula luteofolia Fatto [‡, NR]
Russula macropoda Singer [‡, NR]
Russula mariae Peck [†, ‡]
Russula michiganensis Shaffer [‡, NR]
Russula modesta Peck [‡, NR]
Russula nigricans Fr. [†]
Russula ochricompacta Bills & O.K. Mill. [‡, NR]
Russula ochrophylla Peck [†, ‡]
Russula operata Burl. [‡, NR]
Russula ornaticeps Burl. [‡, NR]
Russula paludosa Britzelm. [‡, NR]
Russula pectinata Fr. [†]
Russula pectinatoides Peck [†, ‡]
Russula pseudointegra Arnould & Goris ex R. Maire [‡, NR]
Russula pseudolepida Singer [‡, NR]
Russula puellaris Fr. [‡, NR]
Russula pulchra Burl. [‡, NR]
Russula pulverulenta Peck [‡, NR]
Russula purpurina Quél. & Schulzer [†]
Russula pusilla Peck [‡, NR]
Russula redolens Burl. [‡, NR]
Russula rosea Pers. [†]
Russula rubescens Beardslee [‡, NR]
Russula rubriceps (Kauffman) Singer [‡, NR]
Russula seperina Dupain [‡, NR]
Russula sericeonitens Kauffman [‡, NR]
Russula silvicola Schaffer [‡, NR]
Russula sordida Peck [†]
Russula squamula Peck [†, ‡]
Russula subfoetens W.G. Sm. [†, ‡]
Russula subgraminicolor Murrill [‡, NR]
Russula subpunctata Kauffman [‡, NR]
Russula subsericeonitens Murrill [‡, NR]
Russula subtilis Burl. [‡, NR]
Russula uncialis Peck [‡, NR]
Russula vesca Fr. [†]
Russula veternosa Fr. [†]
Russula virescens (Schaeff.) Fr. [†, ‡]
Russula xerampelina (Schaeff.) Fr. [†, ‡]
Scytonostroma galactinum (Fr.) Donk [‡, NR]
Scytonostroma portentosum (Berk. & M.A. Curtis) Donk [‡, NR]
Scytonostromella protrusum (Burt) Nakasone [‡, NR]
Scytonostromella arachnoidea (Peck) Hjortstam [†]
Stereum complicatum (Fr.) Fr. [†, ‡]
Stereum gausapatum (Fr.) Fr. [†, ‡]
Stereum hirsutum (Willd.) Pers. [†, ‡]
- Stereum ochraceoflavum* (Schwein.) Sacc. [‡, NR]
Stereum ostrea (Blume & T. Nees) Fr. [†, ‡]
Stereum striatum (Fr.) Fr. [†, ‡]
Stereum versicolor (Sw.) Fr. [†, ‡]
Vararia investiens (Schwein.) P. Karst. [‡, NR]
Xylobolus frustulatus (Pers.) Boidin [†, ‡]
Xylobolus subpileatus (Berk. & M.A. Curtis) Boidin [‡, NR]
- Septobasidiales**
Septobasidium pteruloides (Mont.) Pat. [‡, NR]
- Thelephorales**
Hydnellum aurantiacum (Batsch) P. Karst. [‡, NR]
Hydnellum caeruleum (Hornem.) P. Karst. [‡, NR]
Hydnellum complicatum Banker [‡, NR]
Hydnellum concrescens (Pers.) Banker [†, ‡]
Hydnellum ferrugineum (Fr.) P. Karst. [‡, NR]
Hydnellum scrobiculatum (Fr.) P. Karst. [‡, NR]
Hydnellum spongiosipes (Peck) Pouzar [†, ‡]
Phellodon atratus K.A. Harrison [‡, NR]
Phellodon confluens (Pers.) Pouzar [‡, NR]
Phellodon melaleucus (Sw. ex Fr.) P. Karst. [‡, NR]
Phellodon niger (Fr.) P. Karst. [†, ‡]
Phellodon tomentosus (L.) Banker [‡, NR]
Sarcodon fennicus (P. Karst.) P. Karst. [‡, NR]
Sarcodon imbricatus (L.) P. Karst. [†, ‡]
Sarcodon scabrosus (Fr.) P. Karst. [‡, NR]
Sarcodon thwaitesii (Berk. & Broome) Maas Geest. [†]
Sarcodon versipellis (Fr.) Nikol. [‡, NR]
Thelephora albido Brunnea Schwein. [‡, NR]
Thelephora anthocephala (Bull.) Fr. [‡, NR]
Thelephora caryophyllea (Schaeff.) Pers. [†, ‡]
Thelephora cuticularis Berk. [‡, NR]
Thelephora intybacea Pers. [‡, NR]
Thelephora multipartita Schwein. [†, ‡]
Thelephora palmata (Scop.) Fr. [†, ‡]
Thelephora penicillata (Pers.) Fr. [‡, NR]
Thelephora regularis Schwein. [†, ‡]
Thelephora schweinitzii Peck [†]
Thelephora spongiosa Schwein. [‡, NR]
Thelephora terrestris Ehrh. [‡, NR]
Thelephora vialis Schwein. [‡, NR]
Tomentella botryoides (Schwein.) Bourdot & Galzin [‡, NR]
Tomentella bryophila (Pers.) M.J. Larsen [†, ‡]
Tomentella crinalis (Fr.) M.J. Larsen [†]
Tomentella fusca (Pers.) J. Schröt. [‡, NR]
Tomentella lapida (Pers.) Stalpers [‡, NR]

Tomentella rubiginosa (Bres.) Maire [‡, NR]

Trechisporales

- Subulicystidium brachysporum* (P.H.B. Talbot & V.C. Green) Jülich [‡, NR]
Subulicystidium longisporum (Pat.) Parmasto [‡, NR]
Trechispora farinacea (Pers.) Libert [‡, NR]
Trechispora fastidiosa (Pers.) Libert [‡, NR]
Trechispora mollusca (Pers.) Libert [‡, NR]
Trechispora stevensonii (Berk. & Broome) K.H. Larss. [‡, NR]

Tremellales

- Basidiocladon caesiocinereum* (Höhn. & Litsch.) Luck-Allen [‡, NR]
Basidiocladon eyrei (Wakef.) Luck-Allen [‡, NR]
Ductifera pululahuana (Pat.) Wells [‡, NR]
Eichlerella deglubens (Berk. & Broome) D.A. Reid [‡, NR]
Exidia alba (Lloyd) Burt [‡, NR]
Exidia glandulosa (Bull.) Fr. [†, ‡]
Exidia nucleata (Schwein.) Burt [‡, NR]
Exidia recisa (Ditmar) Fr. [†, ‡]
Gloeotromera alba (Lloyd) Ervin [‡, NR]
Gloeotromera pululahuana (Pat.) Ervin [‡, NR]
Myxarium hyalinum (Pers.) Donk [‡, NR]
Pseudohydnum gelatinosum (Scop.) P. Karst. [†]
Sebacina concrescens (Schwein.) P. Roberts [‡, NR]
Sebacina epigaea (Berk. & Broome) Neuhoff [†]
Sebacina incrustans (Pers.) Tul. & C. Tul. [†, ‡]
Syzygospora mycetophila (Peck) Ginns [†]
Tremella albida Huds. [†]
Tremella foliacea Pers. [†, ‡]
Tremella fuciformis Berk. [†, ‡]
Tremella mesenterica (Schaeff.) Retz. [†, ‡]
Tremella reticulata (Berk.) Farl. [‡, NR]
Tremella sparassoidea Lloyd [‡, NR]
Tremella vesicaria Bull. [‡, NR]

LITERATURE CITED

- Anderson, H.W. & P.J. Anderson. 1920. The parasitic fungi of Montgomery County I. Proceedings of the Indiana Academy of Science 29:175–222.
- Arthur, J.C. 1897. Additions to the cryptogamic flora of Indiana. Proceedings of the Indiana Academy of Science 6:214–216.
- Avis, P.G. 2012. Ectomycorrhizal iconoclasts: the ITS rDNA diversity and nitrophilic tendencies of fetid *Russula*. Mycologia 104:998–1007.
- Avis, P.G., I.A. Dickie & G.M. Mueller. 2006. A ‘dirty’ business: testing the limitations of terminal restriction fragment length polymorphism (TRFLP) analysis of soil fungi. Molecular Ecology 15:873–882.
- Avis, P.G., G.M. Mueller & J. Lussenhop. 2008. Ectomycorrhizal fungal communities in two North American oak forests respond to nitrogen addition. New Phytologist 179:472–483.
- Avis P.G., W.C. Gaswick, G.S. Tonkovich & P.R. Leacock. 2017. Monitoring fungi in ecological restorations of coastal Indiana, U.S.A. Restoration Ecology 25:92–100.
- Baird, R. 2010. Leveraging the fullest potential of scientific collections through digitization. Biodiversity Informatics 7:130–136.
- Beaman, R.S. & N. Cellinese. 2012. Mass digitization of scientific collections: new opportunities to transform the use of biological specimens and underwrite biodiversity science. Zookeys 209:7–17.
- Blackwell, M. 2011. The Fungi: 1, 2, 3 . . . 5.1 million species? American Journal of Botany 98:426–438.
- Blagoderov, V., I.J. Kitching, L. Livermore, T.J. Simonsen & V.S. Smith. 2012. No specimen left behind: industrial scale digitization of natural history collections. Zookeys 209:133–146.
- Bolan, N.S. 1991. A critical review on the role of mycorrhizal fungi in the uptake of phosphorus by plants. Plant and Soil 134:189–207.
- Busteed, R.C. 1936. James M. Van Hook. Proceedings of the Indiana Academy of Science 45:22–24.
- Clark, F. 1994. Indiana GAP analysis: implications for biodiversity conservation and restoration. Proceedings of the Indiana Academy of Science 103:215–224.
- Conway, P. 2010. Preservation in the age of Google: digitization, digital preservation, and dilemmas. Library Quarterly 80:61–79.
- Cooke, W.B. 1975. The 1970 Indiana foray. Mycologia 67:1065–1071.
- Cottingham, J.O. 1947. Higher fungi of Marion County, Indiana. Proceedings of the Indiana Academy of Science 57:74–76.
- Cottingham, J.O. 1948. Higher fungi of Marion County, Indiana. Proceedings of the Indiana Academy of Science 58:81.
- Cottingham, J.O. 1949. Higher fungi of Marion County, Indiana. Proceedings of the Indiana Academy of Science 59:58.
- Cottingham, J.O. 1950. Higher fungi of Marion County, Indiana. Proceedings of the Indiana Academy of Science 60:91–92.
- Cottingham, J.O. 1951. Higher fungi of Marion County, Indiana. Proceedings of the Indiana Academy of Science 61:77.
- Cottingham, J.O. 1952. Higher fungi of Marion County, Indiana. Proceedings of the Indiana Academy of Science 62:143.
- Cottingham, J.O. 1954. Higher fungi of Marion County, Indiana, VII. Proceedings of the Indiana Academy of Science 64:247.

- Cottingham, J.O. 1956. Higher fungi of Marion County, Indiana, VIII. Proceedings of the Indiana Academy of Science 66:305.
- Evans-Ruhl, G., R.X. Latin, P.C. Pecknold, D.H. Scott, B. Mitchell & L. Harmer. 1982. Compilation of plant diseases and disorders in Indiana. Proceedings of the Indiana Academy of Science 91:120–139.
- Fink, B. & S.C. Fuson. 1920. An arrangement of the Ascomycetes of Indiana. Proceedings of the Indiana Academy of Science 29:113–134.
- Fisher, M.C., D.A. Henk, C.J. Briggs, J.S. Brownstein, L.C. Madoff, S.L. McCraw & S.J. Gurr. 2012. Emerging fungal threats to animal, plant and ecosystem health. *Nature* 484:186–194.
- Friesner, R.C. 1936. Presidential Address: Indiana as a critical botanical area. Proceedings of the Indiana Academy of Science 46:28–45.
- Gries, C., E.E. Gilbert & N.M. Franz. 2014. Symbiota – a virtual platform for creating voucher-based biodiversity information communities. *Biodiversity Data Journal* 2:e1114.
- Gurung, S., S.B. Goodwin, M. Kabbage, W.W. Bockus & T.B. Adhikari. 2011. Genetic differentiation at microsatellite loci among populations of *Mycosphaerella graminicola* from California, Indiana, Kansas, and North Dakota. *Phytopathology* 101:1251–125.
- Harmon, P.F. & R. Latin. 2005. Winter survival of the perennial ryegrass pathogen *Magnaporthe oryzae* in north central Indiana. *Plant Disease* 89:412–418.
- Hawksworth, D.L. 1991. The fungal dimension of biodiversity: magnitude, significance, and conservation. *Mycological Research* 95:641–655.
- Herre, A.W.C.T. 1944. Lichens known from Indiana. *Proceedings of the Indiana Academy of Science* 53:81–95.
- Hyerczyk R.D. 1998. The lichen flora of Hoosier Prairie State Nature Preserve. *Proceedings of the Indiana Academy of Science* 106:25–32.
- Jackson, H.S. 1918. The Ustilaginales of Indiana. *Proceedings of the Indiana Academy of Science* 27:119–132.
- Jackson, H.S. 1921a. The Ustilaginales of Indiana II. *Proceedings of the Indiana Academy of Science* 30:157–164.
- Jackson, H.S. 1921b. The Ustilaginales of Indiana III. *Proceedings of the Indiana Academy of Science* 30:165–182.
- Leacock, P.R., J. Riddell, A.W. Wilson, R. Zhang, C. Chen Ning & G.M. Mueller. 2016. *Cantharellus chicagoensis* sp. nov. is supported by molecular and morphological analysis as a new yellow chanterelle in midwestern United States. *Mycologia* 108:765–772.
- Lee, H.J. 1929. New parasitic fungi of Montgomery County. *Proceedings of the Indiana Academy of Science* 39:135–136.
- Lindahl, B.O., A.F.S. Taylor & R.D. Finlay. 2002. Defining nutritional constraints on carbon cycling in boreal forests – towards a less ‘phytocentric’ perspective. *Plant and Soil* 242:123–135.
- Molina, R., T. O'Dell, D. Luoma, M. Amaranthus, M. Castellano & K. Russell. 1993. Biology, ecology, and social aspects of wild edible mushrooms in the forests of the Pacific Northwest: A preface to managing commercial harvest. General Technical Report PNW-GTR-309. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.
- O'Donnell, K., A.P. Rooney, G.L. Mills, M. Kuo, N.S. Weber & S.A. Rehner. 2011. Phylogeny and historical biogeography of true morels (*Morchella*) reveals an early Cretaceous origin and high continental endemism and provincialism in the Holarctic. *Fungal Genetics and Biology* 48:252–265.
- Pringle A. & E.C. Vellinga. 2006. Last chance to know? Using literature to explore the biogeography and invasion biology of the death cap mushroom *Amanita phalloides* (Vaill. ex Fr. :Fr.) Link. *Biological Invasions* 8:1131–1144.
- Reddick, D. 1907. A Preliminary List of the Hymenomycetes or Mushrooms of Indiana. Pp. 1191–1253. In Indiana Department of Geology and Natural Resources, Thirty-Second Annual Report. Indianapolis, Indiana.
- Rose, J.N. 1886. Mildews of Indiana. *Botanical Gazette* 11:60–63.
- Ruch, D.G., M.C. Robbins & K. Nurtjahja. 2001. The lignicolous fungi of Ginn Woods, Delaware County, Indiana. *Proceedings of the Indiana Academy of Science* 110:79–93.
- Schmit, J.P., J.F. Murphy & G.M. Mueller. 1999. Macrofungal diversity of a temperate oak forest: a test of species richness estimators. *Canadian Journal of Botany* 77:1014–1027.
- Schoknecht, J.D. 1982. Contribution to the flora of Indiana: fungi of west-central Indiana on the occasion of the sixth annual A. H. Smith Great Lakes Foray. *Proceedings of the Indiana Academy of Science* 91:140–143.
- Underwood, L.M. 1892. Some additions to the state flora from Putnam County. *Proceedings of the Indiana Academy of Science* 1:89–91.
- Van der Heijden, M.G., J.N. Klironomos, M. Ursic, P. Moutoglis, R. Streitwolf-Engel, T. Boller, A. Wiemken & I.R. Sanders. 1998. Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity. *Nature* 396:69–72.
- Van Hook, J.M. 1911. Indiana fungi. I. *Proceedings of the Indiana Academy of Science* 20:205–212.
- Van Hook, J.M. 1912. Indiana fungi. II. *Proceedings of the Indiana Academy of Science* 21:347–354.
- Van Hook, J.M. 1913. Indiana fungi. III. *Proceedings of the Indiana Academy of Science* 22:99–102.

- Van Hook, J.M. 1916. Indiana fungi. IV. Proceedings of the Indiana Academy of Science 25:141–148.
- Van Hook, J.M. 1921a. Indiana fungi. V. Proceedings of the Indiana Academy of Science 30:209–214.
- Van Hook, J.M. 1921b. The perithecium of *Cicinnobolis*. Proceedings of the Indiana Academy of Science 30:215–216.
- Van Hook, J.M. 1922. Indiana fungi. VI. Proceedings of the Indiana Academy of Science 31:143–148.
- Van Hook, J.M. 1924. Indiana fungi. VII. Proceedings of the Indiana Academy of Science 33:233–238.
- Van Hook, J.M. 1925. Indiana fungi. VIII. Proceedings of the Indiana Academy of Science 34:317–320.
- Van Hook, J.M. 1926. Indiana fungi. IX. Proceedings of the Indiana Academy of Science 35:333–336.
- Van Hook, J.M. 1928. Indiana fungi. X. Proceedings of the Indiana Academy of Science 37:365–371.
- Van Hook, J.M. 1929a. Indiana fungi. XI. Proceedings of the Indiana Academy of Science 38:127–131.
- Van Hook, J.M. 1929b. Some new species of fungi. Proceedings of the Indiana Academy of Science 38:131–132.
- Van Hook, J.M. 1930. Indiana fungi. XII. Proceedings of the Indiana Academy of Science 39:75–83.
- Van Hook, J.M. 1935. Indiana fungi. XIII. Proceedings of the Indiana Academy of Science 44:55–64.
- Van Hook, J.M. & R.C. Busteed 1935. Anthracnose of *Betula nigra*. Proceedings of the Indiana Academy of Science 44:81.
- Vilsack, T. & C.Z. Clark. 2012. Census of Agriculture: United States summary and state data. National Agriculture Statistics Service, United States Department of Agriculture, AC-12-A-51(1). Washington, DC.
- Wilhelm, G.S. 1990. Special vegetation of the Indiana Dunes National Lakeshore. Research program report 90–02. National Park Service–Midwest Region, Omaha, Nebraska. 373 pp.
- Wilson, A.W., J.L. Beckerman & M.C. Aime. 2014. First report of the white pine blister rust fungus, *Cronartium ribicola*, on *Ribes odoratum* in Indiana. Plant Disease 98:277–278.
- Whitaker Jr, J.O., C.J. Amlaner Jr, M.T. Jackson, G.R. Parker & P.E. Scott. 2012. Habitats and ecological communities of Indiana: presettlement to present. Indiana University Press. 512 pp.
- Wolf, S.C. 1972. Plant diseases in Indiana in 1972. Proceedings of the Indiana Academy of Science 82:101–108.
- Woods, A.J., J.M. Omernik, C.S. Brockman, T.D. Gerber, W.D. Hosteter & S.H. Azevedo. 1998. Ecoregions of Indiana and Ohio. At: ftp://newftp.epa.gov/EPADataCommons/ORD/Ecoregions/in/ohin_eco_lg.pdf (Accessed 16 January 2017).

Manuscript received 18 January 2017, revised 18 July 2017.