

Investigating the fungi responsible for the recent large-scale dieback of Blue Gum Eucalyptus (*Eucalyptus globulus*) in the San Francisco Bay Area

A report prepared by:

Matteo Garbelotto, Principal Investigator, U.C. Berkeley

For:

Mia Ingolia, San Francisco Public Utility Commission
Susan Frankel, Pacific Southwest Research Station, US Forest Service
Pamela Beitz, East Bay Regional Park District

Introduction and Materials and Methods

Since 2020, widespread decline and dieback of Eucalyptus trees has been reported in multiple parts of the San Francisco Bay Area in California. We visited a total of six sites around the Bay Area to examine dying and dead eucalyptus trees to assess the role of fungi in this tree health concern. For laboratory analysis we collected wood, foliage and soil samples from and around symptomatic trees.

Study sites

Two sites were located in the Crystal Springs reservoirs watershed (San Mateo County) on San Francisco Public Utility Commission (SFPUC) property. Site 1 (E1-E4) was accessed via the Trousdale gate. Site 2 (E5-8) and was located adjacent to Cañada Road. Site 3 (E9-12) was in Tilden Regional Park, off South Park Drive in Contra Costa County. Site 4 (E13-16) was on Albany Hill in Albany (Alameda County) and Site 5 (E17-20) was in Lake Chabot Regional Park near Marciel Road (Alameda County), while Site 6 (E21-24) was in Carquinez Strait Regional Park (Contra Costa County).

Sampling

At each site we sampled from four trees with dieback symptoms. We rated trees based on foliage browning/spotting levels, using a scale of 0-5; 0 being unaffected and 5 being dead. All the collections were from trees rated 2-4 (Fig. 1). For each tree we took samples - if lesions or staining were present - in the roots, root collar, stem, branches, twigs, and leaves (Fig. 2). In addition, we collected soil at the root collar to test for the presence of soilborne pathogens, including *Phytophthoras*. Trees were felled to enable access to the branches and leaves and to assess the condition of the entire bole (Fig. 3).



Figure 1. Examples of dieback in eucalyptus.



Figure 2. Examples of collected material.



Figure 3. Tree fallers at Tilden regional park, SFPUC and Albany Hill.

Isolations were made from all collected material on a variety of media: MEA, PARPH, Fusarium select, Leptographium select. We also placed a piece of plant tissue between two slices of fresh carrot (a method for retrieving *Ceratocystis* spp.) and incubated in the dark. For the stem and branch samples we plated from both bleached (30 s in 5% sodium hypochlorite followed by rinsing in sterile distilled water) and unbleached (sterile distilled water only) pieces.

Soil samples were collected to be tested for presence of *Phytophthora* using a soil baiting method. Soil was first dried for ten days, re-wetted and placed at 5 °C in the dark for three days. Samples were brought to room temperature and more water was added to about twice the volume of the soil. A green organic pear, organic oregano and a rhododendron leaf were floated in the container and checked for lesions after three, seven and ten days. Lesion segments, if any, were plated onto PARPH media.

All isolation plates were incubated at room temperature and checked three times for growth over a period of 2-3 weeks. Subcultures from mycelium were plated on MEA agar and grown at room temperature. Following 2-3 weeks of growth, the subcultures were grouped into morphotypes based on morphological characteristics. Two or three isolates from the same morphotypes were sequenced using primers ITS1f-ITS4 or DC6-ITS4.

Results

Samples collected are listed in Table 1, including tree location and DBH, sample type and symptoms observed. Fungal isolates obtained in culture and identification to the genus or species level are listed in Table 2. The map with details about location of the trees sampled and isolation outcomes in graphic format (a different color is assigned to each major fungus isolated) can be found by clicking the link below (clicking on it will download a kmz file you can view on Google Earth). Make sure you enlarge the image and click on the icon to view the “spiderfies” displaying all results

[Eucalyptus Dieback Project Map](#)

Table 1. List of 124 samples collected from the field. Note that twigs should read “twigs and leaves”.

| Date Sampled | Site | tree | Sample Type | spp | DBH/Size cm | Lat | Long | Canopy |
|--------------|-------------|------|---------------|---------------------|-------------|-----------|-------------|--------|
| 5/11/21 | SFPUC 1 | E1 | branch lesion | Eucalyptus globulus | 93 | 37.582646 | -122.406398 | 2 |
| 5/11/21 | SFPUC 1 | E1 | twig | Eucalyptus globulus | 93 | 37.582646 | -122.406398 | 2 |
| 5/11/21 | SFPUC 1 | E1 | leaf | Eucalyptus globulus | 93 | 37.582646 | -122.406398 | 2 |
| 5/11/21 | SFPUC 1 | E1 | root collar | Eucalyptus globulus | 93 | 37.582646 | -122.406398 | 2 |
| 5/11/21 | SFPUC 1 | E1 | soil | Eucalyptus globulus | 93 | 37.582646 | -122.406398 | 2 |
| 5/11/21 | SFPUC 1 | E2 | twig | Eucalyptus globulus | 93 | 37.582409 | -122.40632 | 2 |
| 5/11/21 | SFPUC 1 | E2 | leaf | Eucalyptus globulus | 93 | 37.582409 | -122.40632 | 2 |
| 5/11/21 | SFPUC 1 | E2 | stem | Eucalyptus globulus | 93 | 37.582409 | -122.40632 | 2 |
| 5/11/21 | SFPUC 1 | E2 | root collar | Eucalyptus globulus | 93 | 37.582409 | -122.40632 | 2 |
| 5/11/21 | SFPUC 1 | E2 | soil | Eucalyptus globulus | 93 | 37.582409 | -122.40632 | 2 |
| 5/11/21 | SFPUC 1 | E3 | twig | Eucalyptus globulus | 34 | 37.582288 | -122.406507 | 2 |
| 5/11/21 | SFPUC 1 | E3 | leaf | Eucalyptus globulus | 34 | 37.582288 | -122.406507 | 2 |
| 5/11/21 | SFPUC 1 | E3 | stem | Eucalyptus globulus | 34 | 37.582288 | -122.406507 | 2 |
| 5/11/21 | SFPUC 1 | E3 | root collar | Eucalyptus globulus | 34 | 37.582288 | -122.406507 | 2 |
| 5/11/21 | SFPUC 1 | E3 | soil | Eucalyptus globulus | 34 | 37.582288 | -122.406507 | 2 |
| 5/11/21 | SFPUC 1 | E4 | twig | Eucalyptus globulus | 5 | 37.582698 | -122.406605 | 2 |
| 5/11/21 | SFPUC 1 | E4 | leaf | Eucalyptus globulus | 5 | 37.582698 | -122.406605 | 2 |
| 5/11/21 | SFPUC 1 | E4 | stem | Eucalyptus globulus | 5 | 37.582698 | -122.406605 | 2 |
| 5/11/21 | SFPUC 1 | E4 | soil | Eucalyptus globulus | 5 | 37.582698 | -122.406605 | 2 |
| 5/11/21 | SFPUC 2 | E5 | twig | Eucalyptus globulus | 32 | 37.496194 | -122.326711 | 3 |
| 5/11/21 | SFPUC 2 | E5 | leaf | Eucalyptus globulus | 32 | 37.496194 | -122.326711 | 3 |
| 5/11/21 | SFPUC 2 | E5 | feeder root | Eucalyptus globulus | 32 | 37.496194 | -122.326711 | 3 |
| 5/11/21 | SFPUC 2 | E5 | stem | Eucalyptus globulus | 32 | 37.496194 | -122.326711 | 3 |
| 5/11/21 | SFPUC 2 | E5 | soil | Eucalyptus globulus | 32 | 37.496194 | -122.326711 | 3 |
| 5/11/21 | SFPUC 2 | E6 | twig | Eucalyptus globulus | 38 | 37.496315 | -122.326835 | 4 |
| 5/11/21 | SFPUC 2 | E6 | leaf | Eucalyptus globulus | 38 | 37.496315 | -122.326835 | 4 |
| 5/11/21 | SFPUC 2 | E6 | stem | Eucalyptus globulus | 38 | 37.496315 | -122.326835 | 4 |
| 5/11/21 | SFPUC 2 | E6 | soil | Eucalyptus globulus | 38 | 37.496315 | -122.326835 | 4 |
| 5/11/21 | SFPUC 2 | E7 | leaf | Eucalyptus globulus | 42 | 37.495979 | -122.326847 | 4 |
| 5/11/21 | SFPUC 2 | E7 | stem | Eucalyptus globulus | 42 | 37.495979 | -122.326847 | 4 |
| 5/11/21 | SFPUC 2 | E7 | soil | Eucalyptus globulus | 42 | 37.495979 | -122.326847 | 4 |
| 5/11/21 | SFPUC 2 | E7 | twig | Eucalyptus globulus | 42 | 37.495979 | -122.326847 | 4 |
| 5/11/21 | SFPUC 2 | E8 | twig | Eucalyptus globulus | 26 | 37.496183 | -122.326671 | 2 |
| 5/11/21 | SFPUC 2 | E8 | leaf | Eucalyptus globulus | 26 | 37.496183 | -122.326671 | 2 |
| 5/11/21 | SFPUC 2 | E8 | stem bark | Eucalyptus globulus | 26 | 37.496183 | -122.326671 | 2 |
| 5/11/21 | SFPUC 2 | E8 | soil | Eucalyptus globulus | 26 | 37.496183 | -122.326671 | 2 |
| 5/13/21 | Tilden | E9 | twig | Eucalyptus globulus | 97 | 37.891703 | -122.238327 | 2 |
| 5/13/21 | Tilden | E9 | leaf | Eucalyptus globulus | 97 | 37.891703 | -122.238327 | 2 |
| 5/13/21 | Tilden | E9 | soil | Eucalyptus globulus | 97 | 37.891703 | -122.238327 | 2 |
| 5/13/21 | Tilden | E10 | twig | Eucalyptus globulus | 37 | 37.891931 | -122.238192 | 2.5 |
| 5/13/21 | Tilden | E10 | leaf | Eucalyptus globulus | 37 | 37.891931 | -122.238192 | 2.5 |
| 5/13/21 | Tilden | E10 | stem | Eucalyptus globulus | 37 | 37.891931 | -122.238192 | 2.5 |
| 5/13/21 | Tilden | E10 | root collar | Eucalyptus globulus | 37 | 37.891931 | -122.238192 | 2.5 |
| 5/13/21 | Tilden | E10 | soil | Eucalyptus globulus | 37 | 37.891931 | -122.238192 | 2.5 |
| 5/13/21 | Tilden | E11 | twig | Eucalyptus globulus | 62 | 37.892104 | -122.238184 | 4 |
| 5/13/21 | Tilden | E11 | leaf | Eucalyptus globulus | 62 | 37.892104 | -122.238184 | 4 |
| 5/13/21 | Tilden | E11 | branch | Eucalyptus globulus | 62 | 37.892104 | -122.238184 | 4 |
| 5/13/21 | Tilden | E11 | stem | Eucalyptus globulus | 62 | 37.892104 | -122.238184 | 4 |
| 5/13/21 | Tilden | E11 | soil | Eucalyptus globulus | 62 | 37.892104 | -122.238184 | 4 |
| 5/13/21 | Tilden | E12 | twig | Eucalyptus globulus | 23 | 37.892243 | -122.238296 | 3 |
| 5/13/21 | Tilden | E12 | leaf | Eucalyptus globulus | 23 | 37.892243 | -122.238296 | 3 |
| 5/13/21 | Tilden | E12 | stem 1 | Eucalyptus globulus | 23 | 37.892243 | -122.238296 | 3 |
| 5/13/21 | Tilden | E12 | stem 2 | Eucalyptus globulus | 23 | 37.892243 | -122.238296 | 3 |
| 5/13/21 | Tilden | E12 | soil | Eucalyptus globulus | 23 | 37.892243 | -122.238296 | 3 |
| 5/19/21 | Albany Hill | E13 | leaf | Eucalyptus globulus | 29 | 37.895994 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E13 | twig | Eucalyptus globulus | 29 | 37.895994 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E13 | stem1 | Eucalyptus globulus | 29 | 37.895994 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E13 | stem2 | Eucalyptus globulus | 29 | 37.895994 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E13 | branch | Eucalyptus globulus | 29 | 37.895994 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E13 | root collar | Eucalyptus globulus | 29 | 37.895994 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E13 | soil | Eucalyptus globulus | 29 | 37.895994 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E14 | leaf | Eucalyptus globulus | 32 | 37.895961 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E14 | twig | Eucalyptus globulus | 32 | 37.895961 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E14 | root collar | Eucalyptus globulus | 32 | 37.895961 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E14 | branch 1 | Eucalyptus globulus | 32 | 37.895961 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E14 | branch 2 | Eucalyptus globulus | 32 | 37.895961 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E14 | stem | Eucalyptus globulus | 32 | 37.895961 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E14 | soil | Eucalyptus globulus | 32 | 37.895961 | -122.375166 | 2 |
| 5/19/21 | Albany Hill | E15 | leaf | Eucalyptus globulus | 52 | 37.896484 | -122.305234 | 3 |
| 5/19/21 | Albany Hill | E15 | twig | Eucalyptus globulus | 52 | 37.896484 | -122.305234 | 3 |
| 5/19/21 | Albany Hill | E15 | branch | Eucalyptus globulus | 52 | 37.896484 | -122.305234 | 3 |
| 5/19/21 | Albany Hill | E15 | stem | Eucalyptus globulus | 52 | 37.896484 | -122.305234 | 3 |
| 5/19/21 | Albany Hill | E15 | soil | Eucalyptus globulus | 52 | 37.896484 | -122.305234 | 3 |
| 5/19/21 | Albany Hill | E16 | leaf | Eucalyptus globulus | 57 | 37.89644 | -122.305141 | 3 |
| 5/19/21 | Albany Hill | E16 | twig | Eucalyptus globulus | 57 | 37.89644 | -122.305141 | 3 |
| 5/19/21 | Albany Hill | E16 | branch | Eucalyptus globulus | 57 | 37.89644 | -122.305141 | 3 |
| 5/19/21 | Albany Hill | E16 | stem | Eucalyptus globulus | 57 | 37.89644 | -122.305141 | 3 |
| 5/19/21 | Albany Hill | E16 | soil | Eucalyptus globulus | 57 | 37.89644 | -122.305141 | 3 |
| 5/19/21 | Chabot RP | E17 | stem 1 | Eucalyptus globulus | 12 | 37.742714 | -122.104369 | 4 |
| 5/19/21 | Chabot RP | E17 | stem2 | Eucalyptus globulus | 12 | 37.742714 | -122.104369 | 4 |
| 5/19/21 | Chabot RP | E17 | root collar | Eucalyptus globulus | 12 | 37.742714 | -122.104369 | 4 |
| 5/19/21 | Chabot RP | E17 | leaves | Eucalyptus globulus | 12 | 37.742714 | -122.104369 | 4 |
| 5/19/21 | Chabot RP | E17 | twigs | Eucalyptus globulus | 12 | 37.742714 | -122.104369 | 4 |
| 5/19/21 | Chabot RP | E17 | soil | Eucalyptus globulus | 12 | 37.742714 | -122.104369 | 4 |
| 5/19/21 | Chabot RP | E18 | branch 1 | Eucalyptus globulus | 28 | 37.741813 | -122.104596 | 4 |

| | | | | | | | | |
|---------|-----------|-----|--------------------|--------------------------|---------|-----------|-------------|---|
| 5/19/21 | Chabot RP | E18 | branch 2 | Eucalyptus globulus | 28 | 37.741813 | -122.104596 | 4 |
| 5/19/21 | Chabot RP | E18 | root collar | Eucalyptus globulus | 28 | 37.741813 | -122.104596 | 4 |
| 5/19/21 | Chabot RP | E18 | leaves | Eucalyptus globulus | 28 | 37.741813 | -122.104596 | 4 |
| 5/19/21 | Chabot RP | E18 | twigs | Eucalyptus globulus | 28 | 37.741813 | -122.104596 | 4 |
| 5/19/21 | Chabot RP | E18 | soil | Eucalyptus globulus | 28 | 37.741813 | -122.104596 | 4 |
| 5/19/21 | Chabot RP | E18 | stem | Eucalyptus globulus | 28 | 37.741813 | -122.104596 | 4 |
| 5/19/21 | Chabot RP | E19 | branch | Eucalyptus globulus | 31 | 37.741934 | -122.104645 | 4 |
| 5/19/21 | Chabot RP | E19 | leaves | Eucalyptus globulus | 31 | 37.741934 | -122.104645 | 4 |
| 5/19/21 | Chabot RP | E19 | twigs | Eucalyptus globulus | 31 | 37.741934 | -122.104645 | 4 |
| 5/19/21 | Chabot RP | E19 | soil | Eucalyptus globulus | 31 | 37.741934 | -122.104645 | 4 |
| 5/19/21 | Chabot RP | E20 | stem | Eucalyptus globulus | 30 | 37.742095 | -122.10404 | 3 |
| 5/19/21 | Chabot RP | E20 | branch | Eucalyptus globulus | 30 | 37.742095 | -122.10404 | 3 |
| 5/19/21 | Chabot RP | E20 | leaves | Eucalyptus globulus | 30 | 37.742095 | -122.10404 | 3 |
| 5/19/21 | Chabot RP | E20 | twigs | Eucalyptus globulus | 30 | 37.742095 | -122.10404 | 3 |
| 5/19/21 | Chabot RP | E20 | soil | Eucalyptus globulus | 30 | 37.742095 | -122.10404 | 3 |
| s | | | | | | | | |
| 5/27/21 | Carquinez | E21 | stem1 | Eucalyptus globulus | 24 | 38.047863 | 122.190632 | 2 |
| 5/27/21 | Carquinez | E21 | stem2 | Eucalyptus globulus | 24 | 38.047863 | 122.190632 | 2 |
| 5/27/21 | Carquinez | E21 | twigs | Eucalyptus globulus | 24 | 38.047863 | 122.190632 | 2 |
| 5/27/21 | Carquinez | E21 | leaves | Eucalyptus globulus | 24 | 38.047863 | 122.190632 | 2 |
| 5/27/21 | Carquinez | E21 | soil | Eucalyptus globulus | 24 | 38.047863 | 122.190632 | 2 |
| 5/27/21 | Carquinez | E21 | stem 3 (x-section) | Eucalyptus globulus | 24 | 38.047863 | 122.190632 | 2 |
| 5/27/21 | Carquinez | E22 | branch 1 | Eucalyptus globulus | 16 & 11 | 38.048104 | -122.190646 | 2 |
| 5/27/21 | Carquinez | E22 | stem 1 | Eucalyptus globulus | 16 & 11 | 38.048104 | -122.190646 | 2 |
| 5/27/21 | Carquinez | E22 | stem 2 | Eucalyptus globulus | 16 & 11 | 38.048104 | -122.190646 | 2 |
| 5/27/21 | Carquinez | E22 | twigs | Eucalyptus globulus | 16 & 11 | 38.048104 | -122.190646 | 2 |
| 5/27/21 | Carquinez | E22 | leaves | Eucalyptus globulus | 16 & 11 | 38.048104 | -122.190646 | 2 |
| 5/27/21 | Carquinez | E22 | soil | Eucalyptus globulus | 16 & 11 | 38.048104 | -122.190646 | 2 |
| 5/27/21 | Carquinez | E23 | stem 1 | Eucalyptus globulus | 24 | 38.046804 | -122.18696 | 4 |
| 5/27/21 | Carquinez | E23 | stem 2 | Eucalyptus globulus | 24 | 38.046804 | -122.18696 | 4 |
| 5/27/21 | Carquinez | E23 | root collar | Eucalyptus globulus | 24 | 38.046804 | -122.18696 | 4 |
| 5/27/21 | Carquinez | E23 | branch | Eucalyptus globulus | 24 | 38.046804 | -122.18696 | 4 |
| 5/27/21 | Carquinez | E23 | twigs | Eucalyptus globulus | 24 | 38.046804 | -122.18696 | 4 |
| 5/27/21 | Carquinez | E23 | leaves | Eucalyptus globulus | 24 | 38.046804 | -122.18696 | 4 |
| 5/27/21 | Carquinez | E23 | soil | Eucalyptus globulus | 24 | 38.046804 | -122.18696 | 4 |
| 5/27/21 | Carquinez | E24 | branch 1 | Eucalyptus camaldulensis | 14 | 38.046587 | -122.187324 | 3 |
| 5/27/21 | Carquinez | E24 | leaves | Eucalyptus camaldulensis | 14 | 38.046587 | -122.187324 | 3 |
| 5/27/21 | Carquinez | E24 | twigs | Eucalyptus camaldulensis | 14 | 38.046587 | -122.187324 | 3 |
| 5/27/21 | Carquinez | E24 | soil | Eucalyptus camaldulensis | 14 | 38.046587 | -122.187324 | 3 |

NOTE: Soil baiting for Phytophthoras has not been completed yet. Report will be amended when results are available.

Table 2. List of isolates obtained from Eucalyptus and their identification based on colony morphology and ITS DNA sequence.

| Site | Tree | Tree Species | Sample Type | Id |
|-----------|------|----------------------------|-----------------|---|
| SFPUC1 | E1 | <i>Eucalyptus globulus</i> | branch | <i>Cytospora eucalypticola</i> |
| SFPUC1 | E1 | <i>Eucalyptus globulus</i> | leaf | <i>Pseudosydowia eucalypti</i> |
| SFPUC1 | E1 | <i>Eucalyptus globulus</i> | root collar | <i>Pestalotiopsis</i> sp. |
| SFPUC1 | E1 | <i>Eucalyptus globulus</i> | twig | <i>Pseudosydowia eucalypti</i> |
| SFPUC1 | E2 | <i>Eucalyptus globulus</i> | leaf | <i>Cylindrium aeroginosum</i> like |
| SFPUC1 | E2 | <i>Eucalyptus globulus</i> | stem | <i>Chaetomium elatum</i> |
| SFPUC1 | E2 | <i>Eucalyptus globulus</i> | twig | <i>Graphostroma/Biscongiauxia</i> |
| SFPUC1 | E2 | <i>Eucalyptus globulus</i> | twig | <i>Coniochaeta</i> sp. |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | leaf | <i>Verrucoconiothyrium/Coniothyrium</i> |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | leaf | <i>Verrucoconiothyrium/Coniothyrium</i> |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | root collar | <i>Epicoccum nigrum</i> |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | stem | <i>Diplodia sapinea</i> |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | stem | <i>Cladosporium</i> sp. |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | stem | <i>Chaetomium cochliodes</i> |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | twig | <i>Stemphylium vesicarium</i> |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | twig | <i>Verrucoconiothyrium/Coniothyrium</i> |
| SFPUC1 | E3 | <i>Eucalyptus globulus</i> | twig | <i>Cytospora berkeleyi</i> |
| SFPUC1 | E4 | <i>Eucalyptus globulus</i> | leaf | <i>Pseudosydowia eucalypti</i> |
| SFPUC1 | E4 | <i>Eucalyptus globulus</i> | stem | <i>Cladosporium</i> sp. |
| SFPUC1 | E4 | <i>Eucalyptus globulus</i> | stem | <i>Diaporthe columnaris</i> |
| SFPUC1 | E4 | <i>Eucalyptus globulus</i> | twig | <i>Alternaria alternata</i> |
| SFPUC2 | E5 | <i>Eucalyptus globulus</i> | leaves | <i>Pseudosydowia eucalypti</i> |
| SFPUC2 | E5 | <i>Eucalyptus globulus</i> | stem | <i>Pseudosydowia eucalypti</i> |
| SFPUC2 | E5 | <i>Eucalyptus globulus</i> | stem | <i>Chaetomium cochliodes</i> |
| SFPUC2 | E5 | <i>Eucalyptus globulus</i> | stem | <i>Chaetomium elatum</i> |
| SFPUC2 | E5 | <i>Eucalyptus globulus</i> | twig | <i>Pseudosydowia eucalypti</i> |
| SFPUC2 | E6 | <i>Eucalyptus globulus</i> | leaves | <i>Pseudosydowia eucalypti</i> |
| SFPUC2 | E6 | <i>Eucalyptus globulus</i> | stem | <i>Beauveria</i> |
| SFPUC2 | E6 | <i>Eucalyptus globulus</i> | stem | <i>Beauveria</i> |
| SFPUC2 | E6 | <i>Eucalyptus globulus</i> | stem | <i>Cytospora berkeleyi</i> |
| SFPUC2 | E6 | <i>Eucalyptus globulus</i> | stem | <i>Chaetomium</i> |
| SFPUC2 | E6 | <i>Eucalyptus globulus</i> | stem | <i>Chaetomium cochliodes</i> |
| SFPUC2 | E6 | <i>Eucalyptus globulus</i> | twigs | <i>Pseudosydowia eucalypti</i> |
| SFPUC2 | E7 | <i>Eucalyptus globulus</i> | twig | <i>Cladosporium</i> sp. |
| SFPUC2 | E8 | <i>Eucalyptus globulus</i> | leaves | <i>Cladosporium</i> sp. |
| SFPUC2 | E8 | <i>Eucalyptus globulus</i> | leaves | <i>Pseudosydowia eucalypti</i> |
| SFPUC2 | E8 | <i>Eucalyptus globulus</i> | leaves | <i>Pseudosydowia eucalypti</i> |
| Tilden | E9 | <i>Eucalyptus globulus</i> | leaf | <i>Pseudosydowia eucalypti</i> |
| Tilden | E10 | <i>Eucalyptus globulus</i> | stem/fungal mat | <i>Phanerochaete martelliana</i> |
| Tilden | E11 | <i>Eucalyptus globulus</i> | branch | <i>Cladosporium</i> sp. |
| Tilden | E11 | <i>Eucalyptus globulus</i> | branch | <i>Biscaugniauxia</i> |
| Tilden | E11 | <i>Eucalyptus globulus</i> | stem | <i>Pseudosydowia eucalypti</i> |
| Tilden | E11 | <i>Eucalyptus globulus</i> | stem | <i>Phanerochaete martelliana</i> |
| Tilden | E11 | <i>Eucalyptus globulus</i> | twig | <i>Neofusicoccum eucalyptorum</i> |
| Tilden | E12 | <i>Eucalyptus globulus</i> | stem 2 | <i>Mortierella</i> sp. |
| Tilden | E12 | <i>Eucalyptus globulus</i> | twig | <i>Harknessia gibbosa</i> |
| Albany | E13 | <i>Eucalyptus globulus</i> | branch | <i>Chaetomium elatum</i> |
| Albany | E13 | <i>Eucalyptus globulus</i> | stem 1 | <i>Graphostroma/Biscongiauxia</i> |
| Albany | E13 | <i>Eucalyptus globulus</i> | stem 2 | <i>Graphostroma/Biscongiauxia</i> |
| Albany | E13 | <i>Eucalyptus globulus</i> | stem 2 | <i>Graphostroma/Biscongiauxia</i> |
| Albany | E13 | <i>Eucalyptus globulus</i> | twig | <i>Neofusicoccum eucalyptorum</i> |
| Albany | E14 | <i>Eucalyptus globulus</i> | branch 2 | <i>Stereum hirsutum</i> |
| Albany | E14 | <i>Eucalyptus globulus</i> | stem | <i>Scytalidium</i> sp. |
| Albany | E14 | <i>Eucalyptus globulus</i> | stem | <i>Scytalidium</i> sp. |
| Albany | E15 | <i>Eucalyptus globulus</i> | stem | <i>Mortierella</i> sp. |
| Albany | E16 | <i>Eucalyptus globulus</i> | branch | <i>Pseudosydowia eucalypti</i> |
| Albany | E16 | <i>Eucalyptus globulus</i> | stem | <i>Pseudosydowia eucalypti</i> |
| Chabot | E17 | <i>Eucalyptus globulus</i> | root collar | <i>Cytospora berkeleyi</i> |
| Chabot | E18 | <i>Eucalyptus globulus</i> | root collar | <i>Beauveria</i> |
| Chabot | E18 | <i>Eucalyptus globulus</i> | stem | <i>Beauveria</i> |
| Chabot | E20 | <i>Eucalyptus globulus</i> | stem | <i>Penicillium</i> |
| Chabot | E20 | <i>Eucalyptus globulus</i> | twig | <i>Cladosporium</i> |
| Carquinez | E23 | <i>Eucalyptus globulus</i> | leaf | <i>Pseudosydowia eucalypti</i> |
| Carquinez | E24 | <i>Eucalyptus globulus</i> | branch | <i>Neofusicoccum eucalyptorum</i> |

Study Rationale and Discussion

The Eucalyptus dieback being observed in the greater San Francisco Bay is comparable in many ways to the Acacia dieback recorded in the same region. The aims, the investigative approach and the methods of our Eucalyptus study were similar to those we have previously described for the Acacia study (see “An investigation into the causes of recent widespread *Acacia* spp. mortality in the San Francisco Bay Area” by Matteo Garbelotto).

In summary, our objectives were to determine:

- 1- What are the general symptoms and or signs (e.g. cankers, wood staining, fungal structures, etc.) associated with the observed dieback?
- 2- Are there fungi that are isolated from all study locations? Where are they isolated from in sampled trees (twigs, leaves, branches, etc.) and what are their putative roles in the dieback?
- 3- Are there other fungi that may be playing a role, and are they site-specific or are they shared among sites?
- 4- Are there primary players and secondary opportunistic players? Are secondary players only present in trees infected by primary players?
- 5- Are the fungi involved native or exotic?

Results of our sampling and isolation efforts on Eucalyptus suggest a different scenario from that observed in acacias, although with some important similarities. First, symptoms observed in Eucalyptus were more markedly limited to the foliage and twigs. Leaf blight and twig necrosis were the only symptoms common across all the six areas surveyed and sampled. Branch and stem cankers, wood discoloration and fungal mats were present, but generally were site-specific or shared by trees only in 2 or 3. Extensive heartrot (i.e. decay of the stem core) was not observed in any tree, although, some wood decay was observed both at the base of stems and on branches.

By far, the most common pathogen isolated and the only one present in all sites was *Pseudosydowia eucalypti* (synonyms *Sydowia/Sphaerulina eucalypti*). This is a little-studied pathogen only reported from Eucalyptus foliage (1), but unofficially thought to be present wherever Eucalyptus have been planted around the globe. Although there is little published on its biology and pathogenicity, the belief that it may be ubiquitous, including areas where Eucalyptus have been planted such as California, suggests the fungus must have an endophytic phase. Indeed, *Sphaerulina* spp are known to be endophytic (2). Our isolation effort, in fact, although mostly successful from foliage and twigs, was also successful from branches and stems. Given the isolation success was from bleached wood chips collected from stems and branches, we conclude this organism has an endophytic life stage not only in leaves, but also in woody parts of the tree, including the stem.

There are other foliar pathogens that, although not as widespread, may be contributing to the browning and early drop of the foliage currently observed in Eucalyptus across the Bay Area. The second most frequently isolated foliar pathogen was *Cladosporium* sp. (*Mycosphaerella* sensu lato) (present in four sites). In (3), Hunter et al. state these fungi are common and important foliar pathogens with the ability to infect and cause cankers in branches and stems, but also known to be endophytes and saprobes. Finally, other foliar pathogens, such as *Pestalotiopsis* sp. and *Alternaria alternata* were isolated, although each one was found only in a single tree and site.

Figure 5. *Pseudosydowia* symptoms in twigs, leaves and branches.



Other fungi

Lesions were found in twigs, branches and stems, with some fungi being isolated from 1-3 sites.

Cytospora spp. were isolated from three sites: *C. berkeleyi* was found in three sites and was isolated from a twig, a stem and a root collar. In one site (SFPUC1), both *C. berkeleyi* and *C. eucalypticola* were found on a twig and a branch, respectively. *Cytospora* cankers were visible from the outside as cankers with rounded margins and exudates (Figure 5). *C. berkeleyi* has been described in California where it has been reported multiple times from *Eucalyptus globulus* and once from *E. paniculata* (4). *Cytospora eucalypticola* has been isolated multiple times from the Southern Hemisphere, but is also listed in one California study on coast live oak (*Quercus*

agrifolia) (5). The development of *Cytospora* cankers has long been known to be associated with environmental stresses and defoliation (6).

The Botryosphaeriaceous (Bot) fungus *Neofusicoccum eucalyptorum* was isolated from three of four sites sampled in the East San Francisco Bay, twice from a twig and once from a branch. Branch and twigs displayed typical underbark lesions/cankers. This fungus has not been officially reported in California (4), and as most Bot fungi, it has a latent phase and canker development is associated with environmental stressors (7). Although mostly isolated from *Eucalyptus*, this fungus is known to have a rather broad host range, including multiple myrtaceous and ericaceous hosts, especially where *Eucalyptus* is grown outside of its native range (8, 9). The only other Bot fungus isolated was *Diplodia sapinea* from the stem of a single tree: incidence and severity of stem *Diplodia* cankers are known to be correlated with drought and other environmental stresses (10).

Figure 5. *Cytospora* canker.



Another fungus, from a genus known to include latent pathogens with both an endophytic and a saprobic phase, was *Diaporthe (Phomopsis) columnaris*, isolated from the stem of a single tree. This fungus is reported both as an endophyte (11) and as a pathogen (12) from various plant hosts. This fungus has already been reported from the US West Coast.

A xylareaceous fungus in the genus *Graphostroma/Biscogniauxia* was isolated in three sites: multiple times from the stem of the same tree, once from a branch and once from a twig. This genus contains latent pathogens with both an endophytic and a pathogenic phase, and drought is known to be correlated with a switch from the endophytic to the pathogenic phase.

Finally, three fungi known to be associated with wood decay activity were isolated. The first was *Phanerochaete martelliana* isolated from two trees in the same site. *P. martelliana* is a wood decay fungus mostly recognized as a saprobe or weak pathogen. In both cases, the fungus was isolated from the stem, and its presence was associated with xylem discoloration, incipient decay, and the presence of a fungal “mycelial” mat. Together, these symptoms and signs indicate that infection and host colonization were well under way and that trees were rather compromised healthwise. *Stereum hirsutum* was also isolated from a branch with obvious signs of decay, although localized to the branch itself. *Chaetomium* species are also wood inhabiting fungi and three species were isolated from the stems of 5 trees in 3 sites. Their isolation was associated with staining of the wood in the stem. *Chaetomium* fungi are associated with soft rot, a wood decay process that occurs when conditions are not conducive to white and brown rot decay processes. We have recently identified a high incidence of soft rot fungi in British Columbia aspen trees in early stages of decay (Matteo Garbelotto and Michael Johnson, personal communication): the role these fungi may be playing in Eucalyptus decline is unclear. They may be active on trees with compromised health, but whether their presence has any role in causation of the dieback or may simply be a consequence of their decline warrants further research. For instance, in the recently investigated Acacia dieback, we were able to uncover an unsuspected and unexpected pathogenic role for the zygomycete *Umbelopsis ramanniana*.

Figure 6. *Phanerochaete martelliana* mycelial mat (white).



Conclusions

Below are the answers to the questions that prompted this study, based on the preliminary and limited data available.

- 1- **What are the general symptoms (e.g. cankers, wood staining, fungal structures, etc.) associated with the observed dieback?** The most widespread symptoms in declining Eucalyptus were foliar blight characterized by coalescing brown spots on the foliage, twig cankers, defoliation and branch die-back. All other symptoms, including stem and branch cankers, xylem discoloration, and the presence of fungal structures were much less frequently observed.
- 2- **Are there fungi that are isolated from all locations? Where are they isolated from in sampled trees (twigs, leaves, branches, etc.) and what are their putative roles in the dieback?** Only *P. eucalypti* was common to all sites, and putatively it is involved in the foliar blight, twig cankers and branch dieback observed.
- 3- **Are there other fungi that may be playing a role, and are they site-specific or are they shared among sites?** Other secondary fungi are present and likely to contribute to the observed decline: Bot fungi and other endophytic latent pathogens, including *Cytospora* spp. and *Biscogniauxia* spp., are present, but they are mostly site-specific or present in a couple of sites and their activity is known to be associated with plant stress.
- 4- **Are there primary players and secondary opportunistic players? Are secondary players only present in trees infected by primary players?** Based on the literature and on the patterns of fungal distribution and symptoms uncovered by this study, we do not believe the fungi isolated are primary pathogens. We did not find any pattern of co-infection, suggesting infection by one pathogen facilitated infection by another pathogen.
- 5- **Are the fungi involved native or exotic?** Both native and exotic fungi were found, and at least two fungi (*P. eulcalypti* and *N. eucalyptorum*) have not officially been reported in California: however, these fungi appear to be common wherever Eucalyptus trees have been planted both within and outside their native range. Given their specificity to Eucalyptus and their endophytic nature, it appears that these fungi, although exotic, may not be invasive, given they are likely to be present wherever Eucalyptus are present.

The dieback in Eucalyptus appears to be strongly driven by environmental stress factors such as those caused by drought, increasing temperature and less fog days, combined with the expression of disease by endophytic latent pathogens or by opportunistic fungi. The only widespread

putative pathogen was *Pseudosydowia eucalypti*, causing mostly a leaf and twig blight, but also isolated from branches and stems. Other secondary opportunistic leaf pathogens such as *Cladosporium* and *Alternaria* were also isolated, without any clear association or co-infection with *P. eucalypti*, indicating that even the widespread *P. eucalypti* may not be predisposing trees to infection by other secondary pathogens.

Less widespread latent pathogens were *Cytospora*, *Neofusicoccum*, *Diplodia*, *Diaporthe* and *Biscogniauxia*, all known to cause disease in association with environmental stressors, especially drought. When analyzing the data cumulatively, one or more of these latent pathogens were found at all sites. Wood decay fungi were also isolated from symptomatic xylem in the stem and root collar of trees: their presence is unusual on trees that are still alive such as those sampled and suggests a condition of great stress for the plants in question.

Although technically some of the fungi have not been officially reported in California, most of them are known to be present wherever Eucalyptus trees are grown. The most widespread fungus was *P. eucalypti*, a putative pathogen that probably is universally present endophytically in Eucalyptus and whose role as a primary pathogen is still debated. This scenario differs from that recently suggested for the Acacia dieback observed in the San Francisco Bay Area. Two pathogens already reported in California on other hosts (*Dothiorella* and *Diaporthe* species) were found to be widespread in acacias, suggesting successful host jumps. Conversely, most, if not all, of the fungi identified on declining Eucalyptus in this study have been reported before on this tree genus, and many have been previously reported from California or the West Coast.

Stem cankers were caused by various site-specific fungi, and stem symptoms were less common than in acacias, suggesting Eucalyptus may resprout, if environmental stressors end within a reasonable time frame. However, stem cankers and decay were observed suggesting some trees are experiencing extreme stress conditions and may not be able to resprout unless their roots are grafted with those of trees experiencing less severe stress conditions.

Not much is known about the biology of *P. eucalypti*, so it is difficult to formulate recommendations, nonetheless this pathogen can be classified as an endophytic latent pathogen, capable of causing disease when plant stress becomes significant. Density may be in part an issue, but not as clearly as for the acacia dieback, given it is likely most Eucalyptus are already endophytically infected by *P. eucalypti* and other latent pathogens. Nonetheless, most infectious tree diseases show a positive density dependence, so its importance cannot be excluded. Tree density may be an issue particularly if the size of tree populations is over the carrying capacity of a site, keeping in mind that such carrying capacity is likely to have been lowered due to reduced water availability. Many of the other latent pathogens are known to sporulate on dead plant tissue, and even if none are widespread, each site has its site-specific array of such pathogens.

Preliminary recommendations (based on preliminary knowledge and not applicable to all sites).

Based on our results, the recent (2020-2021) dieback observed in Eucalyptus trees around the San Francisco Bay Area appears to be driven mostly by environmental stressors, including but not limited to drought, which predispose trees to disease caused by opportunistic pathogens. The best management option is to attempt to reduce such environmental stresses. Below, are some suggestions on how we can work towards maintaining healthier Eucalyptus stands.

- 1- Stanturf et al. (13) write:” *Eucalyptus* has potentially higher water use and water use efficiency compared to pasture, pine plantations, and native forests but water use is much lower in *Eucalyptus* plantings than in irrigated crops”. Eucalyptus can grow in drier sites not because they are xeric (drought-adapted) plants, but because they are extremely efficient in absorbing water. They actually require more water than truly xeric plants water, hence the dieback observed in 2021, during a severe drought, is not surprising. In Northern California, all Eucalyptus species are exotic and invasive, and the presence of decline could be used as an indicator that a stand should be removed, because ill adapted to the recent warmer and drier climate. On drier slopes or sites, the best option may be to remove a stand, possibly leaving a few better-looking individuals in wetter portions of the site, to serve as wildlife and recreation trees. As shown in our study, canopy decline/dieback can lead to other diseases including those responsible for wood decay, potentially making standing trees hazardous.
- 2- If a stand is to be preserved, reduce tree density where possible, selectively thinning out the weakest individuals characterized by: a) extensive foliar browning and canopy thinning (make sure the symptom is not caused by the Eucalyptus tortoise beetle <http://ipm.ucanr.edu/PMG/PESTNOTES/pn74104.html>), b) the presence of dead branch tips, c) the presence of stem and branch cankers, and, d) stem wood decay may be ongoing, at least based on the presence of symptoms and signs including but not limited to detached bark, underbark fungal mats, fruitbodies, and seepage through bark openings. This approach may be particularly valuable in recreational areas (campsites, picnic sites), near parking areas, visitors’ centers, etc.
- 3- To reduce fire hazard, adjacent to residential areas or in the proximity of buildings and recreational areas, parking areas, campsites, picnic sites eliminate dead woody debris from a site and dispose of it by burial in a landfill (or locally, see below), composting or burning. This will decrease the risk of increased fire spread rate and intensity, in the case of wildfire.
- 4- Any procedure aimed at reducing drought stress may be beneficial, but see below for caveats with regards to watering. (Also check for local water-use restrictions due to reduced water supply).
- 5- On high value trees: pruning of “dead wood” and watering may be beneficial, provided that no water directly reaches the canopy through overhead irrigation. Additional moisture on the canopy of drought stressed trees would further favor leaf blight.
- 6- Prune severely trees all the way to the first scaffold: this will reduce water needs and will eliminate much of the foliar inoculum.
- 7- In areas where Eucalyptus stands are key elements of the landscape and provide a desirable backdrop to recreational areas, concert venues, and even residential neighborhoods, it may be desirable to reduce the severity of leaf blight caused by *P. eucalypti* and other foliar pathogens. In order to reduce fungal inoculum (i.e. the number

of infectious fungal spores causing new leaf infections), bury leaf litter locally or broadcast soil on top of leaf litter, making sure no leaves are exposed. If pruning trees, chip branches and leaves and bury chips on site or remove from site and dispose as described in point 3 above.

- 8- NOTE: No major generalist (i.e. with a broad host range) infectious pathogen has been identified in ailing Eucalyptus we have examined in this study. The only widespread putative pathogen is *P. eucalypti* a fungus that may be endophytic in most Eucalyptus trees, and appears to be genus-specific (i.e. able to infect only Eucalyptus trees): we do not have sufficient information to determine whether special precautions and prescriptions normally enforced to curb the spread of exotic invasive plant pathogens may be needed to dispose of plant material infected by *P. eucalypti*. Nonetheless, in the preliminary recommendations above, we suggest how inoculum reduction may be achieved.

Possible useful future studies in the short term

In this study, a small number of sites and trees were sampled. The study is also a 'snapshot' of conditions in spring/summer 2021. Greenhouse and field inoculation studies are needed to corroborate some of the conclusions of this study. The pathogenicity of *P. eucalypti*, *N. eucalyptorum*, *C. berkeleyi* and of a *Chaetomium* sp. should be tested in the presence and absence of stress.

Acknowledgements

Funding for this study was provided by the San Francisco Public Utilities Commission (SFPUC) and by the East Bay Regional Parks Foundation. We appreciate access to sites and assistance from Margot Cunningham, City of Albany. We also appreciate the support and assistance provided by Mia Ingolia, SFPUC; Pamela Beitz, East Bay Regional Park District, and Susan Frankel, US Forest Service, Pacific Southwest Research Station. Tree cutting was possible thanks to funding, personnel and resources provided by the SFPUC, East Bay Regional Park District and the City of Albany.

References

- 1) Crous, P.W., Wingfield, M.J., Cheewangkoon, R., Carnegie, A.J., Burgess, T.I., Summerell, B.A., Edwards, J., Taylor, P.W.J. and Groenewald, J.Z., 2019. Foliar pathogens of eucalypts. *Studies in Mycology*, 94, pp.125-298.
- 2) Schlegel, M., Queloz, V. and Sieber, T.N., 2018. The endophytic mycobiome of European Ash and Sycamore Maple leaves—geographic patterns, host specificity and influence of Ash Dieback. *Frontiers in microbiology*, 9, p.2345.

- 3) Hunter, G.C., Crous, P.W., Carnegie, A.J., Burgess, T.I. and Wingfield, M.J., 2011. Mycosphaerella and Teratosphaeria diseases of Eucalyptus; easily confused and with serious consequences. *Fungal Diversity*, 50(1), pp.145-166.
- 4) Farr, D.F., & Rossman, A.Y. Fungal Databases, U.S. National Fungus Collections, ARS, USDA. Retrieved August 2, 2021, from <https://nt.ars-grin.gov/fungaldatabases/> .
- 5) Lynch, S.C., Zambino, P.J., Scott, T.A. and Eskalen, A., 2014. Occurrence, incidence and associations among fungal pathogens and *Agrilus auroguttatus*, and their roles in *Quercus agrifolia* decline in California. *Forest Pathology*, 44(1), pp.62-74.
- 6) Guyon, J.C., Jacobi, W.R. and McIntyre, G.A., 1996. Effects of environmental stress on the development of Cytospora canker of aspen. *Plant disease (USA)*.
- 7) Barradas, C., Pinto, G., Correia, B., Castro, B.B., Phillips, A.J.L. and Alves, A., 2018. Drought× disease interaction in Eucalyptus globulus under *Neofusicoccum eucalyptorum* infection. *Plant Pathology*, 67(1), pp.87-96.
- 8) Pérez, C.A., Wingfield, M.J., Slippers, B., Altier, N.A. and Blanchette, R.A., 2009. *Neofusicoccum eucalyptorum*, a Eucalyptus pathogen, on native Myrtaceae in Uruguay. *Plant Pathology*, 58(5), pp.964-970.
- 9) Hilário, S., Lopes, A., Santos, L. and Alves, A., 2020. Botryosphaeriaceae species associated with blueberry stem blight and dieback in the Centre Region of Portugal. *European Journal of Plant Pathology*, 156(1), pp.31-44.
- 10) Ferreira, S.L., Stauder, C.M., Martin, D.K. and Kasson, M.T., 2021. Morphological and Phylogenetic Resolution of *Diplodia corticola* and *D. quercivora*, Emerging Canker Pathogens of Oak (*Quercus* spp.), in the United States. *Plant Disease*, pp.PDIS-05. doi: <https://doi.org/10.1101/2020.05.08.084111>.
- 11) Martins, F., Pereira, J.A., Bota, P., Bento, A. and Baptista, P., 2016. Fungal endophyte communities in above-and belowground olive tree organs and the effect of season and geographic location on their structures. *Fungal Ecology*, 20, pp.193-201.
- 12) Castlebury, L.A., Rossman, A.Y. and Putnam, M.L., 2002. A new species of *Phomopsis* causing twig dieback of *Vaccinium vitis-idaea* (lingonberry). *Mycological Research*, 106(6), pp.745-752.
- 13) Stanturf, J.A., Vance, E.D., Fox, T.R. and Kirst, M., 2013. Eucalyptus beyond its native range: Environmental issues in exotic bioenergy plantations.