

Ponderosa pine (Pinus ponderosa) Tree up to 120' tall, 40' wide; much smaller in thickets. Bark black, turning yellow with age, smells like vanilla. 2-3 needles per cluster, cones 3-6" long.

Gambel oak (Quercus gambelii) • Small tree to 8-50' tall, crown 5-30' wide. Leaves lobed. Acorns long. Bark gray, checkered to flakey.

Juniper (Juniperus spp.) • Tree 20-40' high, 5-20' wide. Needles short, scaly. Cones ("berries") round, red or purple, ½" diameter, smell of gin. Utah juniper: bark shaggy. Rocky Mt., one-seed juniper: bark scaly. Alligator juniper: bark checkered.

Shrub up to 1' tall and 1' wide. Leaves 2+3" long and 1-2" wide, often mottled red, serrated evergreen. Flowers edaes. yellow, maturing to blue berries.

Buckbrush (Ceanothus fendlerii) Shrub to 3' tall, 3' wide. Branches have small spines. Leaves dark green, ½" long. Flowers white, in tipy clumps. Grows vigorously after fire.

Woods' rose (Rosa woodsii) • Shrub 1-6' tall and 3-6' wide. Leaves 2"/long, compound with 5-9 leaflets, serrated edges. Flowers pink, 2" wide, five petals, maturing to red rosehips, ½" long. Thorns to ½" long.

sumaç Skunkbush trilobata). • Shrub up to 7' tall. Leaves 1-2", lobed. Flowers yellow-white in small clusters, maturing to sticky red berries.

Mexico locust (Robina neomexicana) • Shrub 5-30' tall 5-20' wide. Stems light brown, covered with thorns. Leaves 12" long, compund with 9-15 leaflets. Flowers white-pink, 1/2" long in clusters. Bean-like seedpods.

Cliffrose (Purshia mexicana) • Shrub to 12, 8 wide, much smaller in harsh conditions. Bark gray, shreddy. Leaves 1/2-3/4" long, deeply lobed, white underneath. Flowers cream-colored, 1" wide. Fruits feather-tailed, clustered.

Mountain mahogany (Cercocarpus montanus) • Shrub to 9'. Leaves dark green, leathery, to ½" long. Flowers yellow and red, trumpet-shaped, 2" long.

Wax current (Ribes cereum) • Shrub to 4' tall. Red or tan twigs. Leaves lobed, long. Berries red, waxy.

arizonica) • Bunchgrass, 1–3' tall. Leaves (blades) tightly rolled, appear cylindrical. Seed heads golden.

Pine dropseed (Blepharoneuron tricholepis) • Bunchgrass, 6-8", leaf blades abundant, slender, erect, Seed heads blue-gray.

Mountain muhly (Muhlenbergia montana) • Bunchgrass 10-30" tall. Dead blades curl like wood shavings. Seed heads blue-grey.

Squirreltail (Elymus elymoides) Perennial grass up to 20" tall and 15" wide. Leaf blades come from base of plant, generally smooth. Seed heads resemble a "bottlebrush," seeds 1–3" long.

Cheatgrass (Broums tectorum) • Grass to 2'. Blades emerge in fall. Seedheads red/purple, drooping. Rapid spring growth, matures before other grasses; invasive species


## Trees, shrubs, and grasses






























Locoweed (Oxytropis lambertii) 3/4" long purple pea-like flowers in clusters to 9" tall. Leaves to 5" long with 11-17 leaflets, sometimes appearing gray. Seedstalk turns pale yellow and stays standing after releasing seed.

Rocky Mountain iris, blue flag (Iris missouriensis) • Common in moist sites, and into the Showy mixed conifer zone. purplish petals and white sepals with blue stripes on round stem to 24" tall. Long, narrow leaves from base of plant. Grows in groups. Growth depends on adequate snowfall.

gilia, Scarlet skyrocket (Ipomopsis aggregato) • Bright red, trumpet-shaped flowers along stalk to 3' tall. Dark green leaves with narrow lobes. First year it consists of leaves close to the ground, second year it grows higher and then flowers.

Yarrow (Achillea millefolium) • Feathery leaves 2-6" long, 1/2" wide in rosettes of many leaves. White flat-topped clusters of flowers on stalk 6-20" tall.

fern (Pteridium aquilinum) • A large fern, to 3' tall. Fronds triangular, usually with three leaflets with many tiny leaves. Grows in large colonies in fields, brushy areas, and woods. Thick root-like underground stems, to 6' long; many fronds may grow from each stem. Vigorous after disturbance. Cover for small animals and emerging plants.

Whitemargin pussytoes (Antennaria marginata) • Small plant lying close to ground forming mats. Necktie shaped leaves  $\frac{1}{2}$ -1" long, slightly hairy underneath, with a white underneath, with a white appearance. White clusters of flowers on stalks 2-4" tall.

Mullein (Verbascum thapsus) • Readily invades disturbed sites but only persists for a few years. First year forms basal rosette of large\_soft, fuzzy leaves, second year sends up seed stalk to 7' high. Hundreds of bright yellow flowers maturing to round seed pods of tiny black seeds. Seed stalk dries to a stout brown shaft that persists through winter.

lupine (Lupinus argenteus) • Branching flowers to 3' tall. Leaves fuzzy when young; 5-9 narrow lobes; pealike flowers in tall clusters; purple. Aggressive growth after fire.

rubberweed (Hymenoxys richardsonii) • To 12" tall, several to dozens of woody stems in a bunch, wooly at base. Yellow daisy-like flowers 1/2" wide maturing to wooly tufts of seeds. Very narrow leaves, poisonous to livestock.

Fleabane (Erigeron spp.) • Many species. Small white, pink, or light purple flowers with numerous narrow petals and yellow center.

Purple geranium (Geranium caespitosum) • Pink or purple flowers; up to 10" tall; few flowers per plant. Thin creeping stems, often red. Leaves fivelobed, smelly when crushed, red in autumn.

Mountain parslev (Pseudocymopterus montanus) Up to 30" tall. Yellow flowers in flat-topped clusters. Leaves green, broad, feathery, with many pointed lobes. Strong scent.

\* Cronquist, A. and A. H. Holmgren, N. H. Holmgren, J. L. Reveal and P. K. Holmgren. 1994. Intermountain Flora Vascular Plants of the Intermountain West, USA, The New York Botanical Garden. Bronx, NY. Volume 5.

## **Wildflowers**

