

## Land Unit 8c – Community 1

*Senna artemisioides* subsp. *sturtii*, *Eremophila gibsonii*, *Enchylaena tomentosa* mid open shrubland on limestone pediment plains.

No upper stratum was present.

The mid stratum occurs as *Senna artemisioides* subsp. *sturtii*, *Eremophila gibsonii* and *Enchylaena tomentosa* mid open shrubland.

The ground stratum occurs as low isolated forbs. Perennial species include *Dissocarpus paradoxus*, *Eriochiton sclerolaenoides*, *Goodenia fascicularis*, *Lepidium phlebopetalum*, *Maireana georgei*, *Ptilotus sessifolius*, *Sclerolaena cuneata*, *Tripogon loliiformis* and *Zygophyllum prismatothecum*. Annual species include *Atriplex spongiosa* and *Zygophyllum prismatothecum*.



| Structural and floristic summary |  |  |  |                               |
|----------------------------------|--|--|--|-------------------------------|
| Number of sites: 1               |  | Approximate coverage across total land unit area: 100% |  | Area: 58.4 ha                 |
| Strata                           | Growth Form  | Cover % (Range)  | Mean Height m (Range)                            | Basal Area m <sup>2</sup> /ha |
| Upper                            | Tree   | 0  | -  |                               |
| Mid                              | Shrub  | 20.0   | 1.0 (0.3-2)                                      |                               |
| Ground                           | Forb   | 4.0  | 0.2 (0-0.3)                                      |                               |
| Frequency of Species             |  |  |  |                               |
| Upper                            | None   |  |  |                               |
| Mid                              | 100%: <i>Senna artemisioides</i> subsp. <i>sturtii</i> , <i>Eremophila tomentosa</i> , <i>Enchylaena tomentosa</i> , <i>Acacia victoriae</i>   |  |  |                               |
| Ground                           | 100%: <i>Dissocarpus paradoxus</i> , <i>Eriochiton sclerolaenoides</i> , <i>Goodenia fascicularis</i> , <i>Lepidium phlebopetalum</i> , <i>Maireana georgei</i> , <i>Ptilotus sessifolius</i> , <i>Sclerolaena cuneata</i> , <i>Tripogon loliiformis</i> |  |  |                               |
|                                  | Annuals: <i>Atriplex spongiosa</i> , <i>Zygophyllum prismatothecum</i>   |  |  |                               |
| Other communities present        |  |  | Approximate coverage across total land unit area |                               |
| None                             |  |  | -  |                               |