

For: **Review of Palynology and Palaeobotany**

Special issue: **African palynology**

Article type: **Extended article**

Title: **Atlas of the tropical West African pollen flora**

Authors: **William D. Gosling¹*, Charlotte S. Miller¹, Daniel A. Livingstone²**

Address:

1 = Department of Environment, Earth & Ecosystems, Centre for Earth, Planetary, Space and Astronomical Research (CEPSAR), The Open University, Walton Hall, Milton Keynes, MK7 6AA, UK.

2 = Department of Biology, Duke University, Durham, North Carolina 27708, USA.

* = Corresponding author:

e-mail William.Gosling@open.ac.uk

tel. +44 (0)1908 655147

fax +44 (0)1908 655151

Abstract

The accurate and consistent identification of fossil pollen is essential to allow robust inferences to be drawn with regard to past climate and vegetation change. Identifications are best achieved through the direct inspection of reference material. Most substantial reference collections are held at prestigious university's in Europe or the United States of America which can restrict access for researchers trying to advance palynology in less developed countries. Digital imaging and fast spreading access to the internet means that it is now possible to produce and disseminate high quality images from pollen reference collections. In this paper we contribute to this growing body of work by presenting images of 364 pollen/spore taxa from West Tropical Africa both as printed plates with a key, and within an associated online searchable database.

Highlights

- 129 plates of 364 pollen/spore taxa
- Coverage of most pollen taxa identified in tropical West African fossil pollen records
- Comprehensive key to pollen identification

Keywords (x6): Africa, identification, key, images, pollen, tropical

1. Introduction

The accurate identification of fossil pollen and spores underpins our ability to reconstruct past vegetation and subsequently infer variation in the Earth's system. Palaeo-palynology can provide, among other things, evidence for past changes in climate (e.g. Davis and Shaw, 2001; Overpeck et al., 1990), biome (e.g. Jolly et al., 1998; Tarasov et al., 1998; Williams et al., 2004) and biodiversity (e.g. Brown, 1999; Odgaard, 1999; Rull, 1987) across the geological record. At one extreme the incredible resilience of pollen/spore chemistry (sporopollenin) allows palynology to provide evidence for the evolution of plants on timescales of millions of years (e.g. Crane et al., 1995; Wikström et al., 2001). Whilst at the other, the influence of human activity upon plants allows the actions of people within the landscape to be traced in the recent past (10-100's years ago) (e.g. Brugam, 1978; Piperno and Flannery, 2001). The integrity of the information supplied by palynologists to all these areas is reliant on consistent typing or, where possible, identification of the specimens found in the fossil record.

Since its inception morphological identification of pollen grains has been standard practice through comparison with modern material (von Post, 1916, reprinted 1967). Subsequently numerous atlases for pollen and spore identification have been produced from across the globe, including: Beug, 2004; Colinvaux et al., 1999; Faegri and Iversen, 1989; Moore et al., 1991; Reille, 1999; Roubik and Moreno, 1991. The proliferation of pollen/spore atlases has facilitated the expansion of the field of palynology and helped with international standardization of identifications. In addition, as computing power has developed, there has been an increase in the number, and sophistication, of online searchable pollen databases (e.g. Bush and Weng, 2007; Lezine, 2005; QPG, 2008). However, geographical coverage and access to images for pollen identification remains patchy.

The most substantial atlases for African pollen and spore identification are: (i) in print (Reille, 1998, 1999), and (ii) online the African pollen database (Lezine, 2005). Both contain thousands of images of species within hundreds of families, and therefore provide good general coverage of taxa likely to be found in the fossil record. In addition, pollen/spore atlases for specific regions have also been produced, e.g. East Africa (Riollet and Bonnefille, 1976), as well as for individual countries: (i) Chad (Maley, 1970), (ii) Ethiopia (Bonnefille, 1971a, b), (iii) Ivory Coast (Ybert, 1979), (iv) Nigeria (Sowunmi, 1973), (v) South Africa (Scott, 1982; van Zinderen Bakker, 1953, 1956; van Zinderen Bakker and Coetze, 1959), and (vi) Sudan (El Ghazali, 1993). In this paper we contribute to this body of knowledge by presenting c. 3000 images and identification keys for >450 pollen and spore taxa commonly found in tropical West Africa.

2. Materials and methods

2.1 Selection of pollen and spore taxa

Taxa were selected for inclusion within this tropical West African pollen atlas based upon two criteria: (i) they had been previously identified within fossil records obtained from terrestrial and marine settings within the region (Dupont *et al.*, 2000; Elenga *et al.*, 1994; Frédoux, 1994; Leroy and Dupont, 1994; Lezine and Vergnaud-Grazzini, 1993; Maley and Livingstone, 1983; Talbot *et al.*, 1984), and/or (ii) they had been identified as significant in the regional flora (Hall and Swaine, 1981).

2.2 Organisation and presentation of images

Images are presented on 129 plates showing both polar and equatorial views of the grain where it was possible to obtain images. Plates are organised by pollen morphology to allow

ease of reference when examining fossil pollen material; following (Beug, 2004). In addition, to facilitate interrogation of the images based on botanical classification the family, genera and species of the images are listed in Table 1 with reference to plate number. The paper is accompanied by a searchable online database which integrates the images presented here with the c. 6000 images of >1000 taxa from the Neotropical Pollen database (Bush and Weng, 2007). To access a Freeware version of the complete revised Tropical Pollen Database download and unpack the TopicalPollenDB2013.zip file from the supplementary online section of this paper. Please note due to the high number of pictures the file is large (2.77GB) and an Apple Macintosh computer running Mac OS X 10.6 or higher is required to run the software. We hope that the combination of our images with those of Bush and Weng (2007) within a freeware format online will promote further expansion by other research groups which will lead to the eventual development of a comprehensive pan-tropical searchable online pollen and spore identification tool.

2.3 Provenance of specimens and image capture

Images and descriptions of all pollen and spores were obtained from the reference collection of c. 30,000 specimens housed within the Department of Biology at Duke University which has been collected and curated by DAL. Reference material was sourced from herbaria and laboratories around the globe; full details for individual specimens can be found on the accompanying online database.

Images were obtained using QCapture software (v. 3.1.1) with a QImaging Micropublisher 3.3 RTV camera mounted on a Nikon Eclipse 50i microscope. All images were taken through a Nikon Plan Fluor 40x/0.75 DIC M/N2 objective.

3. Terminology

Terminology follows Punt *et al.* (2007).

4. Nomenclature

Taxonomic nomenclature follows conventions of Beug (2004).

5. Polyads

Plates I-III

Key

- | | |
|---|-----------------------------------|
| 1. Comprising eight grains | 2. |
| - Comprising twelve grains | 3. |
| - Comprising more than twelve grains | 5. |
| 2. Surface psilate (Plate I: 1-3) | <i>Calpocalyx brevibracteatus</i> |
| - Surface scabrate (Plate I: 4-6) | <i>Xylia evansii</i> |
| 3. Polyad diameter less than 40 μ m | 4. |
| - Polyad diameter greater than 40 μ m (Plate II: 1-3) | <i>Acacia clavigera</i> |
| 4. Surface scabrate (Plate I: 7-9) | <i>Acacia seyal</i> |
| - Surface psilate (Plate I: 10-12) | <i>Acacia eggelingii</i> |
| 5. Polyad arrangement non-uniform | 6. |
| - Polyad arrangement uniform (Plate III: 4-6) | <i>Parkia velutina</i> |
| 6. Polyad diameter less than 110 μ m (Plate II: 4-6) | <i>Parkia inundabilis</i> |
| - Polyad diameter greater than 110 μ m (Plate III: 1-3) | <i>Parkia bussei</i> |

6. Tetrads

Plate IV

Key

- | | |
|--------------------------------------|-----------------------------|
| 1. Visible columnae | 2. |
| - Invisible columnae (Plate IV: 1-4) | <i>Erica arborea</i> |
| 2. Surface verrucate (Plate IV: 5-7) | <i>Mimosa strigillosa</i> |
| - Surface scabrate (Plate IV: 8-9) | <i>Uvariopsis congensis</i> |

7. Vesiculate

Plate V

Podocarpus milanjianus (Plate V: 1-6)

8. Inaperturate, including Polypodiaceae

Plates VI-XVI

Key

- | | |
|--|---------------------------------|
| 1. Surface reticulate | 2. |
| - Surface echinate | 6. |
| - Surface psilate | 9. |
| - Surface reticulate | 14. |
| - Surface foveolate (Plate VII : 4-7) | <i>Trichomanes mandiocanum</i> |
| - Surface verrucate | 15. |
| - Surface gemmate (Plate VII: 1-3) | <i>Borassus aethiopum</i> |
| - Surface perforate (Plate VI: 5-7) | <i>Acidanthera brevicollis</i> |
| - Surface scabrate | 16. |
| 2. Equatorial grain shape circular | 3. |
| - Equatorial grain shape suboblate (Plate IX: 9-11) | <i>Xyris montana</i> |
| - Equatorial grain shape subprolate (Plate VIII: 10-13) | <i>Scaphopetalum thonneri</i> |
| - Equatorial grain shape rectangular (tall) (Plate IX: 1-8) | <i>Xyris</i> sp. (x2) |
| 3. Equatorial grain diameter ~9-11 μ m (Plate X: 1-4) | <i>Peperomia</i> sp. |
| - Equatorial grain diameter ~12-18 μ m (Plate XI: 8-10) | <i>Tiliacora funifera</i> |
| - Equatorial grain diameter ~27-33 μ m (Plate XI: 11-14) | <i>Scaphopetalum letestui</i> |
| - Equatorial grain diameter ~40-55 μ m | 4. |
| - Equatorial grain diameter ~63-77 μ m | 5. |
| - Equatorial grain diameter ~70-90 μ m (Plate XII: 7-10) | <i>Croton macrostachyus</i> |
| - Equatorial grain diameter ~90-110 (Plate XIII 1-2) | <i>Dicranolepis usambarica</i> |
| 4. Wall thickness ~2 μ m (Plate XII: 1-3) | <i>Psychotria goetzei</i> |
| - Wall thickness ~4 μ m (Plate XII: 4-6) | <i>Croton gratissimus</i> |
| 5. Wall thickness ~2 μ m (Plate XIII 5-7) | <i>Morinda citrifolia</i> |
| - Wall thickness ~6 μ m (Plate XIII: 3-4) | <i>Tribulus terrestris</i> |
| 6. Equatorial grain shape circular | 7. |
| - Equatorial grain shape suboblate (Plate VII: 8-11) | <i>Pycnanthus dinklagei</i> |
| - Equatorial grain shape oblate (Plate VI: 1-4) | <i>Illigera rhodantha</i> |
| - Equatorial grain shape subprolate (Plate VIII: 14-17) | <i>Pandanus livingstonianus</i> |

7. Equatorial grain diameter ~17-23µm - Equatorial grain diameter ~36-44µm (Plate XIV: 1-3)	8. <i>Iodes kamerunensis</i>
- Equatorial grain diameter ~63-77µm (Plate XIII: 8-10)	<i>Barteria acuminata</i>
- Equatorial grain diameter ~81-99µm (Plate XIV: 4-7)	<i>Illigera appendiculata</i>
8. Wall thickness ~0.5µm (Plate XIV: 8-11) - Wall thickness ~1.5µm (Plate XV: 1-4)	<i>Lemna gibba</i> <i>Pandanus kirkii</i>
9. Equatorial grain shape circular - Equatorial grain shape suboblate - Equatorial grain shape subprolate	10. 12. 13.
10. Polar grain shape circular (Plate XI: 1-4) - Polar grain shape quinquangular - Polar grain shape triangular (convex) (Plate X: 13-16)	<i>Nymphaea caerulea</i> 11. <i>Nymphaea lotus</i>
11. Wall thickness ~1µm (Plate VII: 16-19) - Wall thickness ~2µm (Plate VII: 12-15)	<i>Alternanthera repens</i> <i>Alternanthera nodiflora</i>
12. Equatorial grain diameter ~35-45µm (Plate XI: 5-7) - Equatorial grain diameter ~50-70µm (Plate VIII: 1-6)	<i>Uvaria kirkii</i> <i>Piptostigma mayumbense</i>
13. Equatorial grain diameter ~50-60µm (Plate X: 11-12)	<i>Lonchitis currori</i>
14. Wall thickness ~1µm (Plate X: 8-10) - Wall thickness ~2µm (Plate X: 5-7)	<i>Nephrolepis biserrata</i> <i>Nephrolepis exaltata</i>
15. Equatorial grain shape circular (Plate XVI: 4-10) - Equatorial grain shape suboblate (Plate VIII: 7-9)	<i>Dichrostachys</i> sp. (x2) <i>Dichrostachys unijuga</i>
16. Equatorial grain diameter ~150-200µm (Plate XV: 5-6) - Equatorial grain diameter unknown (Plate XVI: 1-3)	<i>Tylophora sylvatica</i> <i>Artabotrys likimensis</i>

9. Monoporate

Plate XVII

Key

1. Surface scabrate (Plate XVII: 1-2)	<i>Guaduella oblonga</i>
- Surface reticulate	2.
2. Pore size ~1.5µm (Plate XVII: 6-8)	<i>Typha angustifolia</i>
- Pore size ~2µm	3.
3. Wall thickness ~1µm (Plate XVII: 3-5)	<i>Typha australis</i>
- Wall thickness ~1.5µm (Plate XVII: 9-11)	<i>Typha capensis</i>

10. Monocolpate

Plates XVIII-XXI

Key

1. Surface psilate	2.
- Surface gemmate (Plate XIX: 6-8)	<i>Borassus aethiopum</i>
- Surface perforate (Plate XXI: 4-6)	<i>Acidanthera brevicollis</i>
- Surface foveolate	4.
- Surface scabrate	5.
- Surface echinate	8.
- Surface reticulate	10.
2. Equatorial grain shape rectangular (tall) (Plate XVIII: 1-2)	<i>Raphia ruffia</i>
- Equatorial grain shape rhombic (tall) (Plate XVIII: 6-8)	<i>Phoenix reclinata</i>
- Equatorial grain shape circular	3.
3. Polar grain shape triangular (convex) (Plate XX: 4-6)	<i>Nymphaea lotus</i>
- Polar grain shape circular (Plate XX: 1-3)	<i>Nymphaea caerulea</i>
4. Equatorial grain shape rectangular (tall) (Plate XIX: 9-10)	<i>Hyphaene natalensis</i>
- Equatorial grain shape rhombic (tall) (Plate XIX: 11-13)	<i>Hyphaene ventricosa</i>
5. Equatorial grain shape subprolate	6.
- Equatorial grain shape rectangular (tall) (Plate XVIII: 3-6, Plate XVIII: 17-18, Plate XVIII: 19-21)	<i>Raphia farinifera</i> <i>Calamus erectus</i> <i>Calamus gracilis</i>
6. Polar grain shape triangular (convex) (Plate XVIII: 9-11)	<i>Elaeis guineensis</i>
- Polar grain shape circular	7.

7. Columnae invisible (Plate XX: 7-9)	<i>Chlorophytum floribundum</i>
- Columnae visible (Plate XX: 10-11)	<i>Asparagus falcatus</i>
8. Wall thinner on pole (Plate XXI: 7-9)	<i>Aneilema johnstonii</i>
- Wall of even thickness	9.
9. Equatorial grain size ~50-60µm (Plate XXI: 10-12)	<i>Commelina africana</i>
- Equatorial grain size ~75-85µm (Plate XXI: 13- 15)	<i>Crinum powellii</i>
- Equatorial grain size ~85-95µm (Plate XXI: 16- 18)	<i>Crinum pauciflorum</i>
10. Colpus length full	11.
- Colpus length 2/3 (Plate XXI: 1-3)	<i>Dracaena reflexa</i>
- Colpus length 1/2 (Plate XX: 12-13)	<i>Dracaena camerooniana</i>
- Colpus length 1/3	12.
11. Columnae invisible (Plate XVIII: 12-14)	<i>Ancistrophyllum secundiflorum</i>
- Columnae visible (Plate XIX: 4-5)	<i>Borassus machadonis</i>
12. Equatorial grain size ~35-45µm (Plate XIX: 1-3)	<i>Eremospatha</i> sp.
- Equatorial grain size ~45-55µm (Plate XVIII: 15-16)	<i>Ancistrophyllum laurentii</i>

11. Syncorporate

Plate XXII

Key

1. Equatorial grain size ~15-20µm (Plate XXII: 7- 10)	<i>Syzygium guineense</i>
- Equatorial grain size ~20-25µm (Plate XXII: 1- 3)	<i>Eugenia michoacanensis</i>
- Equatorial grain size ~30-37µm (Plate XXII: 4- 6)	<i>Myrcia</i> sp.

12. Diporate

Plates XXIII-XXV

1. Equatorial grain shape rectangular (tall) (Plate XXIII: 1-4)	<i>Musanga smithii</i>
- Equatorial grain shape sub-prolate (Plate XXIII: 5-9)	<i>Musanga leo-errerae</i>

- Equatorial grain shape sub-oblanceolate	2.
- Equatorial grain shape circular	4.
2. Pore shape elliptic (tall) (Plate XXIII: 10-15)	<i>Chlorophora excelsa</i>
- Pore shape circular	3.
3. Pore morphology thickened pore (Plate XXIII: 16-17)	<i>Antiaris toxicaria</i>
- Pore morphology plain (Plate XXIII: 18-19)	<i>Ficus ingens</i>
4. Pore shape irregular (Plate XXIV: 1-5)	<i>Iodes ovalis</i>
- Pore shape circular with annulus	5.
- Pore shape circular	6.
5. Surface psilate (Plate XXIV: 6-9)	<i>Baissea multiflora</i>
- Surface scabrate (Plate XXIV: 10-12)	<i>Motandra guineensis</i>
6. Pore morphology thickened pore (Plate XXIV: 13-19)	<i>Trema orientalis</i>
- Pore morphology plain	7.
7. Equatorial grain diameter ~18-22µm (Plate XXIV: 20-22)	<i>Trema guinensis</i>
- Equatorial grain diameter ~63-77 µm (Plate XXV: 1-2)	<i>Morinda citrifolia</i>

13. Dicolporate

Plate XXVI

Key

1. Pore size ~10µm (Plate XXVI: 1-6)	<i>Justicia cordata</i>
- Pore size ~4µm (Plate XXVI: 7-11)	<i>Justicia flava</i>

14. Triporate

Plates XXVII-XXXIV

Key

1. Surface granulate (Plate XXVII: 1-7)	<i>Allophylus africanus</i>
- Surface echinate	2.
- Surface psilate	3.
- Surface reticulate	8.
- Surface scabrate	17.
2. Polar grain diameter ~18-21µm (Plate XXVII:	<i>Iodes ovalis</i>

8-11)		
- Polar grain diameter ~35-45µm (Plate XXVII: 12-14)		<i>Piliostigma reticulatum</i>
- Polar grain diameter ~50-70µm (Plate XXVIII: 1-3)		<i>Bombax brevicuspe</i>
3. Polar grain shape circular	4.	
- Polar grain shape triangular (convex)	5.	
4. Polar grain diameter ~12-16µm (Plate XXVIII: 4-6)		<i>Sesuvium sessile</i>
- Polar grain diameter ~20-30µm (Plate XXVIII: 10-12)		<i>Baissea multiflora</i>
5. Pore size ~1-3µm	6.	
- Pore size ~4-5µm (Plate XXVIII: 19-21)		<i>Protea susannae</i>
6. Pore morphology thickened pore (Plate XXVIII: 13-15)		<i>Sabicea floribunda</i>
- Pore morphology thinning sexine	7.	
7. Polar grain diameter ~8-12µm (Plate XXVIII: 7-9)		<i>Coula edulis</i>
- Polar grain diameter ~18-22µm (Plate XXVIII: 16-18)		<i>Heisteria parvifolia</i>
8. Pore morphology thickened pore	9.	
- Pore morphology slightly extruded	10.	
- Pore morphology extruded	11.	
- Pore morphology plain	12.	
- Pore morphology thinning sexine	15.	
9. Polar grain diameter ~25-35µm (Plate XXVIII: 22-24)		<i>Bombax buonopozense</i>
- Polar grain diameter ~40-50µm (Plate XXIX: 1- 6)		<i>Plectronia vulgaris</i>
10. Polar grain diameter ~14-16µm (Plate XXIX: 10-15)		<i>Lasianthus africanus</i>
- Polar grain diameter ~45-55µm (Plate XXIX: 16-18)		<i>Kirkia acuminata</i>
- Polar grain diameter ~60-80µm (Plate XXIX: 7- 9)		<i>Ceiba pentandra</i>
11. Polar grain diameter ~36-44µm (Plate XXX: 1-7)		<i>Vigna fischeri</i>
- Polar grain diameter ~54-66µm (Plate XXX: 8- 10)		<i>Cardiospermum grandiflorum</i>
12. Polar grain shape circular	13.	

- Polar grain shape triangular (convex)	14.
13. Pore size ~3µm (Plate XXX: 11-14)	<i>Triplochiton scleroxylon</i>
- Pore size ~5µm (Plate XXX: 15-18)	<i>Nesogordonia fertilis</i>
- Pore size ~6µm (Plate XXX: 19-21)	<i>Entada umbonata</i>
14. Pore size ~6µm (Plate XXXI: 1-6)	<i>Nesogordonia parvifolia</i>
- Pore size ~8µm (Plate XXXI: 13-15)	<i>Vigna luteola</i>
- Pore size ~25µm (Plate XXXI: 16-17)	<i>Psychotria fractinervata</i>
15. Polar grain diameter ~10-15µm (Plate XXXI: 7-12)	<i>Dichapetalum mossambicense</i>
- Polar grain diameter ~36-45µm (Plate XXXII: 1-3)	<i>Paullinia pinnata</i>
- Polar grain diameter ~45-55µm	16.
16. Pore shape circular (Plate XXXII: 4-6)	<i>Entada pursaetha</i>
- Pore shape elliptic (tall) (Plate XXXII: 7-13)	<i>Protea trichanthera</i>
17. Polar grain shape circular	18.
- Polar grain shape triangular (convex)	21.
18. Pore shape rectangular (broad) (Plate XXXII: 14-19)	<i>Leptaulus daphnoides</i>
- Pore shape circular	19.
- Pore shape circular (with annulus)	20.
19. Polar grain diameter ~18-22µm (Plate XXXIII: 1-4)	<i>Celtis zenkeri</i>
- Polar grain diameter ~27-33µm (Plate XXXIII: 5-7)	<i>Celtis mildbraedii</i>
20. Pore morphology sunken pore (Plate XXXIII: 8-13)	<i>Anthocleista grandiflora</i>
- Pore morphology slightly extruded (Plate XXXIII: 14-17)	<i>Hymenocardia acida</i>
21. Polar grain diameter ~10-15µm (Plate XXXIV: 1-3)	<i>Dichapetalum stuhlmannii</i>
- Polar grain diameter ~18-24µm	22.
- Polar grain diameter ~40-60µm (Plate XXXIV: 4-7)	<i>Turraeanthus africana</i>
22. Wall thickness ~0.25µm (Plate XXXIV: 8-9)	<i>Striga forbesii</i>
- Wall thickness ~1.5µm (Plate XXXIV: 10-14)	<i>Celtis integrifolia</i>

15. Tricolpate

Plates XXXV-XXXIX

Key

1. Surface psilate (Plate XXXV: 1-5)	<i>Gunnera chilensis</i>
- Surface clavate/psilate (Plate XXXVI: 6-11)	<i>Vitex amboniensis</i>
- Surface scabrate (Plate XXXV: 6-10)	<i>Napoleona imperialis</i>
- Surface striate (Plate XXXIX: 6-11)	<i>Berlinia grandiflora</i>
- Surface reticulate	2.
2. Equatorial grain shape rectangular (tall) (Plate XXXVII: 1-6)	<i>Scytopetalum tieghemii</i>
- Equatorial grain shape rhombic (tall) (Plate XXXVII: 7-12)	<i>Dichostemma</i> sp.
- Equatorial grain shape circular (Plate XXXVIII: 19-20)	<i>Paramacrolobium coeruleum</i>
- Equatorial grain shape rhombic (broad) (Plate XXXVIII: 7-12)	<i>Brachystegia leonensis</i>
- Equatorial grain shape suboblate (Plate XXXVIII: 1-6)	<i>Brachystegia spiciformis</i>
- Equatorial grain shape prolate	3.
- Equatorial grain shape subprolate	4.
3. Colp length 1 (Plate XXXVI: 1-5)	<i>Leucas calostachys</i>
- Colp length 2 (Plate XXXVI: 24-26)	<i>Aneulophus africanus</i>
4. Polar grain shape triangular (convex)	5.
- Polar grain shape circular	6.
- Polar grain shape tri-lobate	8.
5. Equatorial grain diameter ~15-25µm (Plate XXXVII: 13-18)	<i>Cissampelos mucronata</i>
- Equatorial grain diameter ~40-50µm (Plate XXXIX: 1-5)	<i>Petersia africana</i>
6. Wall thickness <2 µm (Plate XXXV: 17-21)	<i>Premna maxima</i>
- Wall thickness >2 µm	7.
7. Equatorial grain diameter ~27-37µm (Plate XXXVI: 18-23)	<i>Tetracera alnifolia</i>
- Equatorial grain diameter ~40-50µm (Plate XXXV: 22-27)	<i>Premna resinosa</i>
8. Wall thickness ~2µm (Plate XXXVI: 12-17)	<i>Vitex doniana</i>
- Wall thickness~ 3µm (Plate XXXV: 11-16)	<i>Farsetia stenoptera</i>
- Wall thickness ~10µm (Plate XXXVIII: 13-18)	<i>Afzelia bracteata</i>

16. Tricolporate

Plates XL-CXIV

Key

1. Equatorial grain shape circular	2.
- Equatorial grain shape oblate (Plate XL: 1-6)	<i>Gerrardina foliosa</i>
- Equatorial grain shape perprolate (Plate XL: 7-11)	<i>Thecacoris gymnoogyne</i>
- Equatorial grain shape prolate	3.
- Equatorial grain shape rectangular (tall)	4.
- Equatorial grain shape rhombic (broad)	5.
- Equatorial grain shape rhombic (tall)	6.
- Equatorial grain shape sub-oblate	7.
- Equatorial grain shape sub-prolate	8.
- Equatorial grain shape undetermined	87.
2. Surface psilate	11.
- Surface reticulate	12.
- Surface scabrate	13.
- Surface striate (Plate XL: 12-17)	<i>Vepris uguenensis</i>
3. Surface psilate	27.
- Surface reticulate	28.
- Surface striate	37.
4. Surface psilate	9.
- Surface reticulate	10.
- Surface scabrate	40.
5. Surface psilate (Plate XLI: 1-10)	<i>Alchornea floribunda</i>
- Surface clavate/pilate (Plate XLI: 11-16)	<i>Ilex mitis</i>
- Surface reticulate	42.
- Surface scabrate	43.
6. Surface psilate	44.
- Surface striate	46.
- Surface reticulate	49.
- Surface scabrate (Plate LXXVIII: 4-7)	<i>Daniella oliveri</i>
7. Surface psilate	53.
- Surface striate (Plate XLII: 1-7)	<i>Crudia bracteata</i>
- Surface reticulate	54.
- Surface scabrate (Plate XLII: 8-14)	<i>Parinari curatellifolia</i>
8. Surface echinate	60.
- Surface psilate	62.
- Surface reticulate	65.
- Surface scabrate	80.
- Surface striate	85.
9. Pore shape circular (Plate XLIII: 1-6)	<i>Diospyros mespiliformis</i>
- Pore shape elliptic (broad) (Plate XLIII: 7-9)	<i>Lotus chazaliei</i>

- Pore shape lalongate (Plate XLIII: 10-16)	<i>Canarium schweinfurthii</i>
- Pore shape lolongate (Plate XLIV: 1-14)	<i>Heliotropium</i> sp. x2
10. Pore shape circular	39.
- Pore shape elliptic (tall) (Plate XLV: 1-6)	<i>Balanites glaber</i>
11. Pore shape circular	14.
- Pore shape elliptic (broad) (Plate XLV: 7-12)	<i>Centroplacus glaucinus</i>
12. Pore shape circular (Plate XLV: 13-18)	
- Pore shape circular with annulus	<i>Indigofera leptoclada</i>
- Pore shape concave	15.
- Pore shape lalongate (Plate XLVI: 1-8)	16.
- Pore shape rectangular (broad)	<i>Millettia psilopetala</i>
- Pore shape lolongate	18.
- Pore shape rhombic (tall) (Plate XLVI: 9-16)	21.
- Pore shape squared	<i>Argomuellera macrophylla</i>
- Pore shape elliptic (tall) (Plate XLVII: 1-6)	23.
- Pore shape irregular (Plate XLVII: 7-12)	<i>Balanites aegyptiacus</i>
- Pore shape undetermined (Plate XLVIII: 1-3)	<i>Martretia quadricornis</i>
	<i>Cliffortia nitidula</i>
13. Pore shape circular	26.
- Pore shape rectangular (broad) (Plate XLVIII: 4-9)	<i>Parinari holstii</i>
- Pore shape lalongate (Plate XLVIII: 10-15)	<i>Macaranga schweinfurt</i>
- Pore shape lolongate (Plate XLVIII: 16-22)	<i>Prunus africana</i>
14. Pore morphology with an operculum (Plate XLIX: 1-8)	<i>Alchornea cordifolia</i>
- Pore morphology sunken pore (Plate XLIX: 9-12)	<i>Medusandra richardsiana</i>
15. Pore morphology thickened pore (Plate XLIX: 14-18)	
- Pore morphology sunken pore (Plate L: 1-7)	<i>Commiphora campestris</i>
16. Pore morphology thickened pore (Plate L: 8-16)	
- Pore morphology sunken pore	<i>Commiphora scheffleri</i>
- Pore morphology slightly extruded (Plate LI: 1-5)	<i>Erythrococca bongensis</i>
- Pore morphology extruded (Plate LI: 6-12)	17. <i>Amanoa strobilacea</i>
17. Pore position mid-wall (Plate LII: 1-8)	<i>Discoglypremna caloneura</i>
- Pore position corner (Plate LII: 9-15)	
18. Pore morphology thickened pore	19.
- Pore morphology sunken pore	20.
19. Colpus length full (Plate LIII: 1-6)	<i>Mallotus wrayi</i>

- Colpus length >2/3 (Plate LIII: 7-12)	<i>Fagara macrophylla</i>
20. Polar grain shape circular (Plate LIII: 13-18)	<i>Olea hochstetteri</i>
- Polar grain shape triangular (convex) (Plate LIV: 1-8)	<i>Ixora aneimenodesma</i>
21. Wall thickness even	22.
- Wall thinner on pole (Plate LIV: 9-16)	<i>Anthostema aubryananum</i>
22. Polar grain shape tri-lobate (Plate LV: 1-7)	<i>Agelaea heterophylla</i>
- Polar grain shape circular (Plate LV: 8-15)	<i>Adenia nicobarica</i>
23. Visible columnae	24.
- Invisible columnae (Plate LVI: 1-3)	<i>Salacia kraussii</i>
24. Polar grain shape tri-lobate (Plate LVI: 4-8)	<i>Maytenus senegalensis</i>
- Polar grain shape triangular (convex)	25.
25. Equatorial grain diameter ~17-25µm (Plate LVI: 9-14)	<i>Hippocratea affinis</i>
- Equatorial grain diameter ~25-35µm (Plate LVII: 1-8)	<i>Hippocratea africana</i>
26. Pore morphology extruded (Plate LVII: 9-14)	<i>Grandidiera boivinii</i>
- Pore morphology slightly extruded (Plate LVIII: 1-8)	<i>Cynometra alexandri</i>
27. Pore shape lalongate (Plate LVIII: 9-13)	<i>Calantica jalbertii</i>
- Pore shape rhombic (broad) (Plate LVIII: 14-20)	<i>Blighia wildemaniana</i>
28. Colpus length full	29.
- Colpus length >2/3	35.
29. Pore shape circular	30.
- Pore shape concave	31.
- Pore shape elliptic (tall)	32.
- Pore shape lalongate	33.
- Pore shape rhombic (broad) (Plate LIX: 1-6)	<i>Corchorus fascicularis</i>
- Pore shape squared (Plate LIX: 7-13)	<i>Strephonema pseudocola</i>
- Pore shape undetermined	34.
30. Equatorial grain diameter ~18-22µm (Plate LX: 1-6)	<i>Crossopteryx febrifuga</i>
- Equatorial grain diameter ~45-55µm (Plate LX: 7-9)	<i>Cissus petiolata</i>
31. Pore morphology sunken pore (Plate LXI: 1-6)	<i>Ritchiea capparoides</i>
- Pore morphology thinning sexine (Plate LX:	<i>Spondianthus preussii</i>

10-17)

32. Pore morphology plain (Plate LXI: 7-13)
 - Pore morphology slightly extruded (Plate LXII: 1-7) *Blighia unijugata*
Cissus quadrangularis
33. Pore morphology plain (Plate LXII: 8-13)
 - Pore morphology sunken pore (Plate LXIII: 1-5) *Grewia bicolor*
Vepris humbertii
34. Pore morphology plain (Plate LXIII: 6-11)
 - Pore morphology slightly extruded (Plate LXIV: 1-6) *Rubus scheffleri*
Drypetes gerrardii
35. Pore shape circular (Plate LXIV: 7-11)
 - Pore shape concave
 - Pore shape elliptic (broad) (Plate LXIV: 12-17)
 - Pore shape irregular (Plate LXV: 1-6)
 - Pore shape rectangular (broad) (Plate LXV: 7-15)
 - Pore shape rhombic (tall) (Plate LXV: 16-21) *Flacourtie indica*
 36. *Avicennia nitida*
Dasylepis assinensis
Hannoia sp. (x2) *Ormocarpum kirkii*
36. Pore morphology thickened (Plate LXVI: 1-7)
 - Pore morphology extruded (Plate LXVI: 8-14) *Maesobotrya hirtella*
Cassipourea flanaganii
37. Pore shape elliptic (tall) (Plate LXVI: 15-20,
 LXVII: 1-4)
 - Pore shape lolongate *Lannea* sp. (x2)
 38. *Isoberlinia angolensis*
Isoberlinia doka
38. Pore morphology thinning sexine (Plate
 LXVII: 5-9)
 - Pore morphology plain (Plate LXVIII: 1-5) *Scytopetalum tieghemii*
39. Pore morphology thinning sexine (Plate
 LXVIII: 6-12)
 - Pore morphology sunken pore (Plate LXIX: 1-7) *Euphorbia hypericifolia*
40. Pore shape elliptic (tall) (Plate LXIX: 8-13)
 - Pore shape lolongate (Plate LXIX: 14-19)
 - Pore shape rectangular (broad) *Pterocarpus abyssinicus*
Pterocarpus lucens
 41. *Dodonaea viscosa*
Tabernaemontana ventricosa
42. Pore shape circular (Plate LXXI: 1-6)
 - Pore shape concave (Plate LXXI: 7-12)
 - Pore shape lolongate (Plate LXXII: 1-8)
 - Pore shape rectangular (broad) (Plate LXXII: 9-16) *Sterculia tragacantha*
Prosopis africana
Hagenia abyssinica
Monotes kerstingii

43. Visible columnae (Plate LXXIII: 1-6)	<i>Copaifera gorskiana</i>
- Invisible columnae (Plate LXXIII: 7-12)	<i>Pygeum africanum</i>
44. Pore shape concave (Plate LXXIII: 13-18)	<i>Acridocarpus macrocalyx</i>
- Pore shape lolongate	45.
45. Wall thickness even (Plate LXXIV: 1-7)	<i>Guibourtia arnoldiana</i>
- Wall thicker on pole (Plate LXXIV: 8-14)	<i>Griffonia simplicifolia</i>
46. Pore morphology thinning sexine (Plate LXXV: 1-6)	<i>Lannea humilis</i>
- Pore morphology sunken pore	47.
- Pore morphology plain	48.
47. Colpus length full (Plate LXXV: 7-12)	<i>Sclerocarya birrea</i>
- Colpus length >2/3 (Plate LXXVI: 1-6)	<i>Spondias mombin</i>
48. Pore shape elliptic (tall) (Plate LXXVII: 1-6)	<i>Berlinia bifoliolata</i>
- Pore shape rhombic (broad) (Plate LXXVII: 7-12)	<i>Teclea villosa</i>
49. Pore shape circular	50.
- Pore shape elliptic (broad)	51.
- Pore shape rhombic (broad)	52.
- Pore shape zonorate (Plate LXXVIII: 1-3)	<i>Corchorus trilocularis</i>
50. Pore morphology thinning sexine (Plate LXXIX: 1-6)	<i>Aubrevillea platycarpa</i>
- Pore morphology sunken pore (Plate LXXX: 1-6)	<i>Caloncoba angolensis</i>
51. Pore morphology thinning sexine (Plate LXXX: 7-12)	<i>Avicennia officinalis</i>
- Pore morphology sunken pore (Plate LXXXI: 1-7)	<i>Culcasia dinklagei</i>
52. Pore morphology plain (Plate LXXXI: 8-12)	<i>Hildegardia barteri</i>
- Pore morphology sunken pore (Plate LXXXII: 1-7)	<i>Zizyphus mauritiana</i>
53. Pore morphology plain (Plate XXII: 7-10)	<i>Syzygium guineense</i>
- Pore morphology thinning sexine (Plate LXXXII: 8-13)	<i>Dialium guianense</i>
- Pore morphology extruded (Plate LXXXII: 14-17)	<i>Irvingia smithii</i>
54. Pore shape circular	55.
- Pore shape elliptic (broad)	58.
- Pore shape elliptic (tall) (Plate LXXXII: 18-23)	<i>Mitragyna inermis</i>

- Pore shape irregular	59.
- Pore shape lalongate (Plate LXXXIII: 1-7)	
- Pore shape squared (Plate LXXXIII: 8-13)	
- Pore shape rectangular (broad) (Plate LXXXIV: 1-3)	
55. Pore morphology extruded (Plate LXXIX: 7-10)	<i>Cardiospermum corindum</i>
- Pore morphology thinning sexine	56.
- Pore morphology sunken pore	57.
56. Polar grain shape triangular (convex) (Plate LXXXIV: 4-10)	<i>Pericopsis angolensis</i>
- Polar grain shape triangular (straight) (Plate LXXXIV: 11-18)	<i>Rhynchosia memnonia</i>
57. Wall thickness even (Plate LXXXV: 1-8)	<i>Piptadenia africana</i>
- Wall thinner on pole (Plate LXXXV: 9-14)	<i>Brachystegia spiciformis</i>
58. Pore morphology sunken pore (Plate LXXXVI: 1-20)	Sapindaceae sp. (x3)
- Pore morphology thinning sexine (Plate LXXXVII: 1-6)	<i>Sesbania goetzei</i>
- Pore morphology extruded (Plate LXXXVII: 7-14)	<i>Ixora brachypoda</i>
59. Visible columnae (Plate LXXXVIII: 1-6)	<i>Rhynchosia</i> sp.
- Invisible columnae (Plate LXXXVIII: 7-13)	<i>Prosopis alpataco</i>
60. Pore shape concave (Plate LXXXIX: 1-7)	<i>Centaurea perrottetii</i>
- Pore shape rectangular (broad) (Plate XC: 1-5)	<i>Centaurea dimorpha</i>
- Pore shape elliptic (tall)	61.
61. Wall thickness even (Plate XCI: 1-7)	<i>Artemisia judaica</i>
- Wall thinner on pole (Plate XCI: 8-13)	<i>Artemisia</i> sp.
62. Pore shape circular	90.
- Pore shape circular with annulus (Plate XCI: 14-19)	<i>Tabernaemontana retusa</i>
- Pore shape concave	63.
- Pore shape elliptic (broad)	64.
- Pore shape rectangular (broad) (Plate XCII: 1-5)	<i>Lophira alata</i>
- Pore shape rectangular (tall) (Plate XCII: 6-13)	<i>Trichodesma africanum</i>
- Pore shape rhombic (broad) (Plate XCII: 14-19)	<i>Diospyros austroafricana</i>
63. Pore morphology sunken pore (Plate XCIII: 1-7)	<i>Diospyros abyssinica</i>
- Pore morphology thinning sexine (Plate XCIII: 8-13)	<i>Afrolicania elaeosperma</i>

64. Colpus length 2/3 (Plate XCIII: 14-16)	<i>Casearia engleri</i>
- Colpus length 1/2 (Plate XCIV: 1-3)	<i>Neolemonniera clitandrifolia</i>
65. Pore shape circular	66.
- Pore shape concave	69.
- Pore shape elliptic (broad)	70.
- Pore shape elliptic (tall)	72.
- Pore shape irregular	75.
- Pore shape lalongate	77.
- Pore shape lolongate	79.
- Pore shape rectangular (broad) (Plate XCIV: 4-10)	<i>Elaeodendron buchananii</i>
- Pore shape rectangular (tall) (Plate XCV: 1-6)	<i>Euphorbia engleri</i>
- Pore shape rhombic (broad) (Plate XCVI: 1-7)	<i>Rhizophora mangle</i>
- Pore shape undetermined (Plate XCVI: 8-15)	<i>Euphorbia cussonioides</i>
66. Pore morphology sunken pore	67.
- Pore morphology thinning sexine (Plate XCVII: 1-6)	<i>Rhektophyllum congense</i>
67. Pore position mid-wall (Plate XCVII: 7-13)	<i>Securinega virosa</i>
- Pore position corner	68.
68. Equatorial grain diameter ~27-33µm (Plate XCVII: 14-17)	<i>Detarium senegalese</i>
- Equatorial grain diameter ~50-60µm (Plate XCVIII: 1-7)	<i>Cola nitida</i>
69. Pore morphology sunken pore (Plate XCVIII: 8-14)	<i>Rhizophora mucronata</i>
- Pore morphology thinning sexine (Plate XCIX: 1-7)	<i>Ritchiea fragariodora</i>
- Pore morphology slightly extruded (Plate XCIX: 8-13)	<i>Millettia oblata</i>
70. Pore morphology sunken pore	71.
- Pore morphology extruded (Plate C: 1-8)	<i>Cassine parvifolia</i>
- Pore morphology slightly extruded (Plate C: 9-15)	<i>Nauuclea esculenta</i>
71. Equatorial grain diameter ~25-30µm (Plate C: 16-23)	<i>Maesobotrya barteri</i>
- Equatorial grain diameter ~30-40µm (Plate CI: 1-6)	<i>Cola millenii</i>
72. Pore morphology thickened (Plate CI: 7-12)	<i>Cola gigantea</i>
- Pore morphology extruded (Plate CII: 1-8)	<i>Bridelia micrantha</i>
- Pore morphology slightly extruded (Plate CII: 9-14)	<i>Euphorbia grandicornis</i>
- Pore morphology plain	73.

73. Pore position mid-wall (Plate CIII: 1-7)	<i>Anthostema senegalense</i>
- Pore position corner	74.
74. Extine type tectate (Plate CIII: 8-14)	<i>Ormocarpum sennooides</i>
- Extine type semitectate (Plate CIV: 1-7)	<i>Salacia pyriformis</i>
75. Pore morphology sunken pore	76.
- Pore morphology thickened (Plate CIV: 8-13)	<i>Grewia glandulosa</i>
76. Polar grain shape circular (Plate CV: 1-6)	<i>Lotus arabicus</i>
- Polar grain shape triangular (convex) (Plate CV: 7-14)	<i>Uapaca heudelotii</i>
77. Pore position corner (Plate CV: 15-20)	78.
- Pore position mid-wall	<i>Vepris gossweileri</i>
78. Pore size ~8µm (Plate CVI: 1-6)	<i>Zanthoxylum procerum</i>
- Pore size ~2µm (Plate CVI: 7-12)	<i>Odyendea gabunensis</i>
79. Pore morphology sunken pore (Plate CVI: 13-20)	<i>Convolvulus trabutianus</i>
- Pore morphology thinning sexine (Plate CVII: 1-8)	<i>Millettia tanaensis</i>
- Pore morphology plain (Plate CVII: 9-15)	<i>Oncoba dentata</i>
80. Pore shape concave (Plate CVII: 16-23)	<i>Detarium le-testui</i>
- Pore shape elliptic (broad)	81.
- Pore shape irregular (Plate CVIII: 1-7)	<i>Kiggelaria africana</i>
- Pore shape lalongate	83.
- Pore shape lolongate	84.
- Pore shape rectangular (broad) (Plate CVIII: 8-12)	<i>Sarcophrynum brachystachyum</i>
81. Pore morphology extruded	82.
- Pore morphology thinning sexine (Plate CVIII: 13-20)	<i>Cassia longiracemosa</i>
82. Wall thickness ~3µm (Plate CIX: 1-7)	<i>Tephrosia nana</i>
- Wall thickness ~1µm (Plate CIX: 8-14)	<i>Tephrosia elata</i>
83. Pore morphology plain (Plate CIX: 15-19)	<i>Dissomeria crenata</i>
- Pore morphology sunken pore (Plate CX: 1-8)	<i>Homalium buchholzii</i>
84. Colpi width ~15µm (Plate CX: 9-14)	<i>Cassia burttii</i>
- Colpi width ~ 35µm (Plate CX: 15-20)	<i>Copaifera carrisoana</i>
85. Pore shape circular	86.
- Pore shape lalongate (Plate CXI: 1-7)	<i>Vepris eugenifolia</i>
- Pore shape lolongate (Plate CXII: 1-6)	<i>Hymenostegia afzelii</i>

86. Equatorial grain diameter ~27-33µm (Plate CXI: 8-15)	<i>Cynometra pedicellata</i>
- Equatorial grain diameter ~35-50µm (Plate CXI: 16-20)	<i>Aphloia theiformis</i>
87. Pore morphology plain	88.
- Pore morphology sunken pore (Plate CXII: 7-9)	<i>Placodiscus amaniensis</i>
- Pore morphology thickened	89.
88. Polar grain shape triangular (convex) (Plate CXII: 10-12)	<i>Nauclea diderrichii</i>
- Polar grain shape tri-lobate (Plate CXIII: 1-3)	<i>Microdesmis</i> sp.
89. Visible columnae (Plate CXIII: 6-7)	<i>Rinorea oblongifolia</i>
- Invisible columnae (Plate CXIII: 4-5)	<i>Rinorea welwitschii</i>
90. Pore morphology sunken pore (Plate CXIII: 8-15)	<i>Pericopsis laxiflora</i>
- Pore morphology thinning sexine (Plate CXIV: 1-6)	<i>Baphia obovata</i>

17. Stephanoporate

Plate CXV

Key

Funtumia latifolia (Plate CXV: 1-3)

18. Stephanocolpate

Plates CXVI-CXX

Key

1. Full length of grain	2.
- 2/3 the full length of grain	3.
- 1/2 the full length of grain	4.
- < 1/3 the full length of grain (Plate CXVI: 1-7)	<i>Borreria densiflora</i>
- Colpi length undetermined (Plate CXVI: 8-9, Plate CXVI: 10-11)	<i>Borreria ruelliae</i> <i>Sesamum indicum</i>
2. Equatorial grain shape subprolate (Plate CXVII: 1-4)	<i>Afzelia quanzensis</i>
- Equatorial grain shape rectangular (broad)	<i>Diodia scandens</i>

(Plate CXVIII: 1-6)

- | | |
|--|----------------------------------|
| 3. Equatorial grain shape circular (Plate CXIX: 1-5) | <i>Mitracarpus hirtus</i> |
| - Equatorial grain shape suboblate (Plate CXIX: 6-9) | <i>Sesamum angustifolium</i> |
| 4. Surface baculate (Plate CXX: 1-6) | <i>Diodia aulacosperma</i> |
| - Surface reticulate (Plate CXX: 7-14) | <i>Mitracarpus verticillatus</i> |

19. Stephanocolporate

Plates CXXI-CXXII

Key

- | | |
|---|------------------------------------|
| 1. Polar grain shape elliptic (Plate CXXI: 1-7) | <i>Atroxima afzeliana</i> |
| - Polar grain shape circular (Plate CXXII: 1-7) | <i>Securidaca longepedunculata</i> |

20. Heterocolporate

Plate CXXIII

Key

- | | |
|--|----------------------------|
| 1. Pore shape elliptic (tall) (Plate CXXIII: 1-6) | <i>Combretum aculeatum</i> |
| - Pore shape elliptic (broad) (Plate CXXIII: 7-12) | <i>Terminalia brownii</i> |
| - Pore shape rectangular (broad) (Plate CXXIII: 13-17) | <i>Combretum gueinzii</i> |
| - Pore shape concave | 2. |
| - Pore shape squared (Plate CXXIII: 18-23) | <i>Terminalia aemula</i> |
| 2. Pore morphology extruded (Plate CXXIII: 24-29) | <i>Guiera senegalensis</i> |
| - Pore morphology sunken pore (Plate CXXIII: 30-35) | <i>Pteleopsis diptera</i> |

21. Periporate

Plates CXXIV-CXXIX

Key

- | | |
|--|----------------------------|
| 1. Pore shape irregular | 2. |
| - Pore shape circular | 4. |
| - Pore shape circular with annulus | 8. |
| - Pore shape elliptic broad (Plate CXXIV: 1-4) | <i>Plantago lanceolata</i> |

2. Surface pattern reticulate	3.
- Surface pattern scabrate (Plate CXXIV: 5-8)	<i>Celosia stuhlmanniana</i>
- Surface pattern verrucate (Plate CXXV: 1-3)	<i>Calystegia sepium</i>
- Surface pattern psilate (Plate CXXV: 4-6)	<i>Costus spectabilis</i>
3. Pore section plain (Plate CXXIV: 9-14)	<i>Celosia</i> sp. (x2)
- Pore section thinning sexine (Plate CXXIV: 15-18)	<i>Drymaria cordata</i>
4. Surface pattern reticulate	5.
- Surface pattern echinate	6.
5. Equatorial grain diameter ~27-33µm (Plate CXXIV: 19-22)	<i>Plantago major</i>
- Equatorial grain diameter ~36-44µm (Plate CXXVI: 1-3)	<i>Plantago palmata</i>
- Equatorial grain diameter ~90-110µm (Plate CXXVI: 4-7)	<i>Dicranolepis oligantha</i>
6. Equatorial grain diameter ~20-30µm (Plate CXXVII: 1-4)	<i>Bosqueia manongarivensis</i>
- Equatorial grain diameter ~60-80µm (Plate CXXVII: 5-8)	<i>Ipomoea donaldsonii</i>
- Equatorial grain diameter ~80-120µm	7.
7. Wall thickness ~5µm (Plate CXXVIII: 5-6)	<i>Hewittia sublobata</i>
- Wall thickness ~12µm (Plate CXXIX: 1-2)	<i>Ipomoea ochracea</i>
8. Equatorial grain diameter ~35-45µm (Plate CXXVIII: 1-4)	<i>Lynchnis</i> sp.
- Equatorial grain diameter ~45-55µm (Plate CXXIX: 3-6)	<i>Cerastium indicum</i>

Acknowledgements

The authors would like to thank the two reviewers for constructive comments which improved the quality of this manuscript. Collection and collation of the pollen images, development of the keys and preparation of this manuscript was funded by awards from the Royal Society (International Travel grant, #TG090002, WDG and DAL) and the Natural Environments Research Council (New Investigator Award, #NE/G000824/1, WDG; and studentship #M82525J, CSM).

References

- Beug, H.-J., 2004. Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete. Verlag Dr. Friedrich Pfeil.
- Bonnefille, R., 1971a. Atlas des pollens d'Ethiopie. *Adansonia* 11, 463-518.
- Bonnefille, R., 1971b. Atlas des pollens d'Ethiopie principales especes des forests de montagne. *Pollen et Spore* 13, 15-72.
- Brown, A.G., 1999. Biodiversity and Pollen Analysis: Modern Pollen Studies and the Recent History of a Floodplain Woodland in S. W. Ireland. *Journal of Biogeography* 26, 19-32.
- Brugam, R.B., 1978. Pollen indicators of land-use change in southern Connecticut. *Quaternary Research* 9, 349-362.
- Bush, M.B., Weng, C., 2007. Introducing a new (freeware) tool for palynology. *Journal of Biogeography* 34, 377-380.
- Colinvaux, P.A., De Oliveira, P.E., Moreno, J.E., 1999. Amazon pollen manual and atlas. Harwood Academic Press., New York.
- Crane, P.R., Friis, E.M., Pedersen, K.R., 1995. The origin and early diversification of angiosperms. *Nature* 374, 27-33.
- Davis, M.B., Shaw, R.G., 2001. Range shifts and adaptive responses to quaternary climate change. *Science* 292, 673-679.
- Dupont, L.M., Jahns, S., Marret, F., Ning, S., 2000. Vegetation change in equatorial West Africa: Time-slices for the last 150 ka. *Palaeogeography, Palaeoclimatology, Palaeoecology* 155, 95-122.
- El Ghazali, G.E.B., 1993. A study on the pollen flora of Sudan. *Review of Palaeobotany and Palynology* 76, 99-345.

- Elenga, H., Schwartz, D., Vincens, A., 1994. Pollen evidence of late Quaternary vegetation and inferred climate changes in Congo. *Palaeogeography, Palaeoclimatology, Palaeoecology* 109, 345-356.
- Faegri, K., Iversen, J., 1989. Textbook of pollen analysis, 4th ed. Wiley, Chichester.
- Frédoux, A., 1994. Pollen analysis of a deep-sea core in the Gulf of Guinea: vegetation and climatic changes during the last 225,000 years B.P. *Palaeogeography, Palaeoclimatology, Palaeoecology* 109, 317-330.
- Hall, J.B., Swaine, M.D., 1981. Distribution and Ecology of Vascular Plants in a Tropical Rain Forest. Forest Vegetation in Ghana. The Hague, Junk.
- Jolly, D., Prentice, I.C., Bonnefille, R., Ballouche, A., Bengo, M., Brenac, P., Buchet, G., Burney, D., Cazet, J.P., Cheddadi, R., Edorh, T., Elenga, H., Elmoutaki, S., Guiot, J., Laarif, F., Lamb, H., Lezine, A.M., Maley, J., Mbenza, M., Peyron, O., Reille, M., Reynaud-Farrera, I., Riollet, G., Ritchie, J.C., Roche, E., Scott, L., Ssemmanda, I., Straka, H., Umer, M., Van Campo, E., Vilimumbalo, S., Vincens, A., Waller, M., 1998. Biome reconstruction from pollen and plant macrofossil data for Africa and the Arabian peninsula at 0 and 6000 years. *Journal of Biogeography* 25, 1007-1027.
- Leroy, S., Dupont, L., 1994. Development of vegetation and continental aridity in northwestern Africa during the Late Pliocene: the pollen record of ODP site 658. *Palaeogeography, Palaeoclimatology, Palaeoecology* 109, 295-316.
- Lezine, A.-M., 2005. African Pollen Database, <http://medias.obs-mip.fr/apd/accueil.htm>.
- Lezine, A.M., Vergnaud-Grazzini, C., 1993. Evidence of forest extension in West Africa since 22,000 BP: A pollen record from the eastern tropical Atlantic. *Quaternary Science Reviews* 12, 203-210.
- Maley, J., 1970. Contributions a l'étude du Bassin tchadien Atlas de pollens du Tchad. *Bulletin du Jardin botanique national de Belgique* 40, 29-48.

- Maley, J., Livingstone, D.A., 1983. Extension d'un element montagnard dans le sud du Ghana (Afrique de l'Ouest) au Pleistocene superieur et a l'Holocene inferieur: premieres donnees polliniques. Comptes Rendus des Seances - Academie, Serie II 296, 1287-1292.
- Moore, P.D., Webb, J.A., Collinson, M.E., 1991. Pollen analysis, 2nd ed. Blackwell Scientific, Oxford.
- Odgaard, B.V., 1999. Fossil pollen as a record of past biodiversity. Journal of Biogeography 26, 7-17.
- Overpeck, J.T., Rind, D., Goldberg, R., 1990. Climate-induced changes in forest disturbance and vegetation. Nature 343, 51-53.
- Piperno, D.R., Flannery, K.V., 2001. The earliest archaeological maize (*Zea mays L.*) from highland Mexico: new accelerator mass spectrometry dates and their implications. Proceedings of the National Academy of Sciences 98, 2101-2103.
- Punt, W., Hoen, P.P., Blackmore, S., Nilsson, S., Le Thomas, A., 2007. Glossary of pollen and spore terminology. Review of Palaeobotany and Palynology 143, 1-81.
- QPG, 2008. Cambridge University Palynological Online Database. University of Cambridge, Cambridge.
- Reille, M., 1998. Pollen et spores d'Europe et d'Afrique du Nord (supplement 2). Laboratoire de Botanique Historique et Palynologie, Marseille.
- Reille, M., 1999. Pollen et Spores d'Europe et d'Afrique du nord (2nd edition). Laboratoire de Botanique Historique et Palynologie, Marseille.
- Riollet, G., Bonnefille, R., 1976. Pollen des Amaranthacées du Lac Rodolphe (Afrique orientale). Déterminations générique et spécifique. Pollen et Spore 18, 67-92.
- Roubik, D.W., Moreno, P.J.E., 1991. Pollen and Spores of Barro Colorado Island. Monographs in Systematic Botany 36, Missouri Botanical Garden.

- Rull, V., 1987. A note on pollen counting in palaeoecology. *Pollen et Spores* XXIX, 471-480.
- Scott, L., 1982. Late quaternary fossil pollen grains from the Transvaal, South Africa. *Review of Palaeobotany and Palynology* 36, 241-278.
- Sowunmi, M.A., 1973. Pollen grains of Nigerian plants. *Grana Palynologica* 13, 145-186.
- Talbot, M.R., Livingstone, D.A., Palmer, P.G., Maley, J., Melack, J.M., Delibrias, G., Gulliksen, S., 1984. Preliminary results from sediment cores from Lake Bosumtwi, Ghana. *Palaeoecology of Africa* 16, 173-192.
- Tarasov, P.E., Webb III, T., Andreev, A.A., Afanas'eva, N.B., Berezina, N.A., Bezusko, L.G., Blyakharchuk, T.A., Bolikhovskaya, N.S., Cheddadi, R., Chernavskaya, M.M., Chernova, G.M., Dorofeyuk, N.I., Dirksen, V.G., Elina, G.A., Filimonova, L.V., Glebov, F.Z., Guiot, J., Gunova, V.S., Harrison, S.P., Jolly, D., Khomutova, V.I., Kvavadze, E.V., Osipova, I.M., Panova, N.K., Prentice, I.C., Saarse, L., Sevastyanov, D.V., Volkova, V.S., Zernitskaya, V.P., 1998. Present-day and mid-Holocene biomes reconstructed from pollen and plant macrofossil data from the former Soviet Union and Mongolia. *Journal of Biogeography* 25, 1029-1053.
- van Zinderen Bakker, E.M., 1953. South African pollen grains and spores I. A.A. Balkema, Amsterdam/Cape Town.
- van Zinderen Bakker, E.M., 1956. South African Pollen Grains and Spores II. A.A. Balkema, Amsterdam/Cape Town.
- van Zinderen Bakker, E.M., Coetzee, J.A., 1959. South African Pollen Grains and Spores III. A.A. Balkema, Amsterdam/Cape Town.
- von Post, L., 1916, reprinted 1967. Forest tree pollen in south Swedish peat bog deposits. *Pollen et Spore* 9, 375-401 (English translation by M.B. Davis and K. Faegri from original 1916 lecture; Introduction by K. Faegri and J. Iversen).

- Wikström, N., Savolainen, V., Chase, M.W., 2001. Evolution of the angiosperms: Calibrating the family tree. *Proceedings of the Royal Society B: Biological Sciences* 268, 2211-2220.
- Williams, J.W., Shuman, B.N., Webb Iii, T., Bartlein, P.J., Leduc, P.L., 2004. Late-Quaternary vegetation dynamics in north America: Scaling from taxa to biomes. *Ecological Monographs* 74, 309-334.
- Ybert, J.P., 1979. Altas des pollens de Côte d'Ivoire. ORSTOM, Paris.

Table caption

Table 1: List of pollen species shown on plates ordered alphabetically by family, genus and species.

Plate captions

5. Polyads

Plate I: 1-3 *Calpocalyx brevibracteatus*, 4-6 *Xylia evansii*, 7-9 *Acacia seyal*, 10-12 *Acacia eggelingii*.

Plate II: 1-3 *Acacia clavigera*, 4-6 *Parkia inundabilis*.

Plate III: 1-3 *Parkia bussei*, 4-6 *Parkia velutina*.

6. Tetrad

Plate IV: 1-4 *Erica arborea*, 5-7 *Mimosa strigillosa*, 8-9 *Uvariopsis congensis*.

7. Vesiculate

Plate V: 1-6 *Podocarpus milanjanus*.

8. Inaperturate, including Polypodiaceae

Plate VI: 1-4 *Illigera rhodantha*, 5-7 *Acidanthera brevicollis*.

Plate VII: 1-3 *Borassus aethiopum*, 4-7 *Trichomanes mandiocanum*, 8-11 *Pycnanthus dinklagei*, 12-15 *Alternanthera nodiflora*, 16-19 *Alternanthera repens*.

Plate VIII: 1-6 *Piptostigma mayumbense*, 7-9 *Dichrostachys unijuga*, 10-13 *Scaphopetalum thonneri*, 14-17 *Pandanus livingstonianus*.

Plate IX: 1-4 *Xyris welwitschii*, 5-8 *Xyris aristata*, 9-11 *Xyris montana*.

Plate X: 1-4 *Peperomia* sp., 5-7 *Nephrolepis exaltata*, 8-10 *Nephrolepis biserrata*, 11-12 *Lonchitis currori*, 13-16 *Nymphaea lotus*.

Plate XI: 1-4 *Nymphaea caerulea*, 5-7 *Uvaria kirkii*, 8-10 *Tiliacora funifera*, 11-14 *Scaphopetalum letestui*.

Plate XII: 1-3 *Psychotria goetzei*, 4-6 *Croton gratissimus*, 7-10 *Croton macrostachyus*.

Plate XIII: 1-2 *Dicranolepis usambarica*, 3-4 *Tribulus terrestris*, 5-7 *Morinda citrifolia*, 8-10 *Barteria acuminata*.

Plate XIV: 1-3 *Iodes kamerunensis*, 4-7 *Illigera appendiculata*, 8-11 *Lemna gibba*.

Plate XV: 1-4 *Pandanus kirkii*, 5-6 *Tylophora sylvatica*.

Plate XVI: 1-3 *Artabotrys likimensis*, 4-6 *Dichrostachys cinerea*, 7-10 *Dichrostachys glomerata*.

9. Monoporate

Plate XVII: 1-2 *Guaduella oblonga*, 3-5 *Typha australis*, 6-8 *Typha angustifolia*, 9-11 *Typha capensis*.

10. Monocolpate

Plate XVIII: 1-2 *Raphia ruffia*, 3-6 *Raphia farinifera*, 6-8 *Phoenix reclinata*, 9-11 *Elaeis guineensis*, 12-14 *Ancistrophyllum secundiflorum*, 15-16 *Ancistrophyllum laurentii*, 17-18 *Calamus erectus*, 19-21 *Calamus gracilis*.

Plate XIX: 1-3 *Eremospatha* sp., 4-6 *Borassus machadonis*, 7-8 *Borassus aethiopum*, 10-11 *Hyphaene natalensis*, 12-13 *Hyphaene ventricosa*.

Plate XX: 1-3 *Nymphaea caerulea*, 4-6 *Nymphaea lotus*, 7-9 *Chlorophytum floribundum*, 10-11 *Asparagus falcatus*, 12-13 *Dracaena camerooniana*.

Plate XXI: 1-3 *Dracaena reflexa*, 4-6 *Acidanthera brevicollis*, 7-9 *Aneilema johnstonii*, 10-12 *Commelina africana*, 13-15 *Crinum powellii*, 16-18 *Crinum pauciflorum*.

11. Syncolporate

Plate XXII: 1-3 *Eugenia michoacanensis*, 4-6 *Myrica* sp., 7-10 *Syzygium guineense*.

12. Diporate

Plate XXIII: 1-4 *Musanga smithii*, 5-9 *Musanga leo-errerae*, 10-15 *Chlorophora excelsa*, 16-17 *Antiaris toxicaria*, 18-19 *Ficus ingens*.

Plate XXIV: 1-5 *Iodes ovalis*, 6-9 *Baissea multiflora*, 10-12 *Motandra guineensis*, 13-19 *Trema orientalis*, 20-22 *Trema guinensis*.

Plate XXV: 1-2 *Morinda citrifolia*.

13. Dicolporate

Plate XXVI: 1-6 *Justicia cordata*, 7-11 *Justicia flava*.

14. Triporate

Plate XXVII: 1-7 *Allophylus africanus*, 8-11 *Iodes ovalis*, 12-14 *Piliostigma reticulatum*.

Plate XXVIII: 1-3 *Bombax brevicuspe*, 4-6 *Sesuvium sessile*, 7-9 *Coula edulis*, 10-12 *Baissea multiflora*, 13-15 *Sabicea floribunda*, 16-18 *Heisteria parvifolia*, 19-21 *Protea susannae*, 22-24 *Bombax buonopozense*.

Plate XXIX: 1-6 *Plectronia vulgaris*, 7-9 *Ceiba pentandra*, 10-15 *Lasianthus africanus*, 16-18 *Kirkia acuminata*.

Plate XXX: 1-7 *Vigna fischeri*, 8-10 *Cardiospermum grandiflorum*, 11-14 *Triplochiton scleroxylon*, 15-18 *Nesogordonia fertilis*, 19-21 *Entada umbonata*.

Plate XXXI: 1-6 *Nesogordonia parvifolia*, 7-12 *Dichapetalum mossambicense*, 13-15 *Vigna luteola*, 16-17 *Psychotria fractinervata*.

Plate XXXII: 1-3 *Paullinia pinnata*, 4-6 *Entada pursaetha*, 7-13 *Protea trichanthera*, 14-19 *Leptaulus daphnoides*.

Plate XXXIII: 1-4 *Celtis zenkeri*, 5-7 *Celtis mildbraedii*, 8-13 *Anthocleista grandiflora*, 14-17 *Hymenocardia acida*.

Plate XXXIV: 1-3 *Dichapetalum stuhlmannii*, 4-7 *Turraeanthus africana*, 8-9 *Striga forbesii*, 10-14 *Celtis integrifolia*.

15. Tricolpate

Plate XXXV: 1-5 *Gunnera chilensis*, 6-10 *Napoleona imperialis*, 11-16 *Farsetia stenoptera*, 17-21 *Premna maxima*, 22-27 *Premna resinosa*.

Plate XXXVI: 1-5 *Leucas calostachys*, 6-11 *Vitex amboniensis*, 12-17 *Vitex doniana*, 18-23 *Tetracera alnifolia*, 24-26 *Aneulophus africanus*.

Plate XXXVII: 1-6 *Scytopetalum tieghemii*, 7-12 *Dichostemma* sp., 13-18 *Cissampelos mucronata*.

Plate XXXVIII: 1-6 *Brachystegia spiciformis*, 7-12 *Brachystegia leonensis*, 13-18 *Afzelia bracteata*, 19-20 *Paramacrolobium coeruleum*.

Plate XXXIX: 1-5 *Petersia africana*, 6-11 *Berlinia grandiflora*.

16. Tricolporate

Plate XL: 1-6 *Gerrardina foliosa*, 7-11 *Thecacoris gymnogyne*, 12-17 *Vepris uguenensis*.

Plate XLI: 1-10 *Alchornea floribunda*, 11-16 *Ilex mitis*.

Plate XLII: 1-7 *Crudia bracteata*, 8-14 *Parinari curatellifolia*.

Plate XLIII: 1-6 *Diospyros mespiliformis*, 7-9 *Lotus chazaliei*, 10-16 *Canarium schweinfurthii*.

Plate XLIV: 1-6 *Heliotropium bacciferum*, 7-14 *Heliotropium subulatum*.

Plate XLV: 1-6 *Balanites glaber*, 7-12 *Centroplacus glaucinus*, 13-18 *Indigofera leptoclada*.

Plate XLVI: 1-8 *Millettia psilopetala*, 9-16 *Argomuellera macrophylla*.

Plate XLVII: 1-6 *Balanites aegyptiacus*, 7-12 *Martretia quadricornis*.

Plate XLVIII: 1-3 *Cliffortia nitidula*, 4-9 *Parinari holstii*, 10-15 *Macaranga schweinfurt*, 16-22 *Prunus africana*.

Plate XLIX: 1-8 *Alchornea cordifolia*, 9-12 *Medusandra richardsiana*, 14-18 *Commiphora campestris*.

Plate L: 1-7 *Commiphora scheffleri*, 8-16 *Erythrococca bongensis*.

Plate LI: 1-5 *Amania strobilacea*, 6-12 *Discoglypremna caloneura*.

Plate LII: 1-8 *Entada abyssinica*, 9-15 *Baphia massaiensis*.

Plate LIII: 1-6 *Mallotus wrayi*, 7-12 *Fagara macrophylla*, 13-18 *Olea hochstetteri*.

Plate LIV: 1-8 *Ixora aneimenodesma*, 9-16 *Anthostema aubryanum*.

Plate LV: 1-6 *Agelaea heterophylla*, 7-14 *Adenia nicobarica*.

Plate LVI: 1-3 *Salacia kraussii*, 4-8 *Maytenus senegalensis*, 9-14 *Hippocratea affinis*.

Plate LVII: 1-8 *Hippocratea africana*, 9-14 *Grandidiera boivinii*.

Plate LVIII: 1-8 *Cynometra alexandri*, 9-13 *Calantica jalbertii*, 14-20 *Blighia wildemaniana*.

Plate LIX: 1-6 *Corchorus fascicularis*, 7-13 *Strephonema pseudocola*.

Plate LX: 1-6 *Crossopteryx febrifuga*, 7-9 *Cissus petiolata*, 10-17 *Spondianthus preussii*.

Plate LXI: 1-6 *Ritchiea capparoides*, 7-13 *Blighia unijugata*.

Plate LXII: 1-7 *Cissus quadrangularis*, 8-13 *Grewia bicolor*.

Plate LXIII: 1-5 *Vepris humbertii*, 6-11 *Rubus scheffleri*.

Plate LXIV: 1-6 *Drypetes gerrardii*, 7-11 *Flacourtia indica*, 12-17 *Avicennia nitida*.

Plate LXV: 1-6 *Dasylepis assinensis*, 7-9 *Hannoa klaineana*, 10-15 *Hannoa undulata*, 16-21 *Ormocarpum kirkii*.

Plate LXVI: 1-7 *Maesobotrya hirtella*, 8-14 *Cassipourea flanaganii*, 15-20 *Lannea stuhlmannii*.

Plate LXVII: 1-4 *Lannea triphylla*, 5-9 *Isoberlinia angolensis*.

Plate LXVIII: 1-5 *Isoberlinia doka*, 6-12 *Scytopetalum tieghemii*.

Plate LXIX: 1-7 *Euphorbia hypericifolia*, 8-13 *Pterocarpus abyssinicus*, 14-19 *Pterocarpus lucens*.

Plate LXX: 1-7 *Dodonaea viscosa*, 8-13 *Tabernaemontana ventricosa*.

Plate LXXI: 1-6 *Sterculia tragacantha*, 7-12 *Prosopis africana*.

Plate LXXII: 1-8 *Hagenia abyssinica*, 9-16 *Monotes kerstingii*.

Plate LXXIII: 1-6 *Copaifera gorskiana*, 7-12 *Pygeum africanum*, 13-18 *Acridocarpus macrocalyx*.

Plate LXXIV: 1-7 *Guibourtia arnoldiana*, 8-14 *Griffonia simplicifolia*.

Plate LXXV: 1-6 *Lannea humilis*, 7-12 *Sclerocarya birrea*.

Plate LXXVI: 1-6 *Spondias mombin*.

Plate LXXVII: 1-6 *Berlinia bifoliolata*, 7-12 *Teclea villosa*.

Plate LXXVIII: 1-3 *Corchorus trilocularis*, 4-7 *Daniella oliveri*.

Plate LXXIX: 1-6 *Aubrevillea platycarpa*, 7-10 *Cardiospermum corindum*.

Plate LXXX: 1-6 *Caloncoba angolensis*, 7-12 *Avicennia officinalis*.

Plate LXXXI: 1-7 *Culcasia dinklagei*, 8-12 *Hildegardia barteri*.

Plate LXXXII: 1-7 *Zizyphus mauritiana*, 8-13 *Dialium guianense*, 14-17 *Irvingia smithii*, 18-23 *Mitragyna inermis*.

Plate LXXXIII: 1-7 *Tetrorchidium didymostemon*, 8-13 *Gaertnera paniculata*.

Plate LXXXIV: 1-3 *Uapaca bojeri*, 4-10 *Pericopsis angolensis*, 11-18 *Rhynchosia memnonia*.

Plate LXXXV: 1-8 *Piptadenia africana*, 9-14 *Brachystegia spiciformis*.

Plate LXXXVI: 1-8 *Lecaniodiscus cupanioides*, 9-15 *Chytranthus obliquinervis*, 16-20 *Chytranthus sacleuxii*.

Plate LXXXVII: 1-6 *Sesbania goetzei*, 7-14 *Ixora brachypoda*.

Plate LXXXVIII: 1-6 *Rhynchosia* sp., 7-13 *Prosopis alpataco*.

Plate LXXXIX: 1-7 *Centaurea perrotetii*.

Plate XC: 1-5 *Centaurea dimorpha*.

Plate XCI: 1-7 *Artemisia judaica*, 8-13 *Artemisia* sp., 14-19 *Tabernaemontana retusa*.

Plate XCII: 1-5 *Lophira alata*, 6-13 *Trichodesma africanum*, 14-19 *Diospyros austroafricana*.

Plate XCIII: 1-7 *Diospyros abyssinica*, 8-13 *Afrolicania elaeosperma*, 14-16 *Casearia engleri*.

Plate XCIV: 1-3 *Neolemonniera clitandrifolia*, 4-10 *Elaeodendron buchananii*.

Plate XCV: 1-6 *Euphorbia engleri*.

Plate XCVI: 1-7 *Rhizophora mangle*, 8-15 *Euphorbia cussonioides*.

Plate XCVII: 1-6 *Rhektophyllum congense*, 7-13 *Securinega virosa*, 14-17 *Detarium senegalense*.

Plate XCVIII: 1-7 *Cola nitida*, 8-14 *Rhizophora mucronata*.

Plate XCIX: 1-7 *Ritchiea fragariodora*, 8-13 *Millettia oblate*.

Plate C: 1-8 *Cassine parvifolia*, 9-15 *Nauclea esculenta*, 16-23 *Maesobotrya barteri*.

Plate CI: 1-6 *Cola millenii*, 7-12 *Cola gigantea*.

Plate CII: 1-8 *Bridelia micrantha*, 9-14 *Euphorbia grandicornis*.

Plate CIII: 1-7 *Anthonstema senegalense*, 8-14 *Ormocarpum sennoides*.

Plate CIV: 1-7 *Salacia pyriformis*, 8-13 *Grewia glandulosa*.

Plate CV: 1-6 *Lotus arabicus*, 7-14 *Uapaca heudelotii*, 15-20 *Vepris gossweileri*.

Plate CVI: 1-6 *Zanthoxylum procerum*, 7-12 *Odyendea gabunensis*, 13-20 *Convolvulus trabutianus*.

Plate CVII: 1-8 *Millettia tanaensis*, 9-15 *Oncoba dentata*, 16-23 *Detarium le-testui*.

Plate CVIII: 1-7 *Kiggelaria africana*, 8-12 *Sarcophrynum brachystachyum*, 13-20 *Cassia longiracemosa*.

Plate CIX: 1-7 *Tephrosia nana*, 8-14 *Tephrosia elata*, 15-19 *Dissomeria crenata*.

Plate CX: 1-8 *Homalium buchholzii*, 9-14 *Cassia burttii*, 15-20 *Copaifera carrissoana*.

Plate CXI: 1-7 *Vepris eugeniifolia*, 8-15 *Cynometra pedicellata*, 16-20 *Aphloia theiformis*.

Plate CXII: 1-6 *Hymenostegia afzelii*, 7-9 *Placodiscus amaniensis*, 10-12 *Nauclea diderrichii*.

Plate CXIII: 1-3 *Microdesmis* sp., 4-5 *Rinorea welwitschii*, 6-7 *Rinorea oblongifolia*, 8-15 *Pericopsis laxiflora*.

Plate CXIV: 1-6 *Baphia obovata*.

17. Stephanoporate

Plate CXV: 1-3 *Funtumia latifolia*.

18. Stephanocolpate

Plate CXVI: 1-7 *Borreria densiflora*, 8-9 *Borreria ruelliae*, 10-11 *Sesamum indicum*.

Plate CXVII: 1-4 *Afzelia quanzensis*.

Plate CXVIII: 1-6 *Diodia scandens*.

Plate CXIX: 1-5 *Mitracarpus hirtus*, 6-9 *Sesamum angustifolium*.

Plate CXX: 1-6 *Diodia aulacosperma*, 7-14 *Mitracarpus verticillatus*.

19. Stephanocolporate

Plate CXXI: 1-7 *Atroxima afzeliana*.

Plate CXXII: 1-7 *Securidaca longepedunculata*.

20. Heterocolporate

Plate CXXIII: 1-6 *Combretum aculeatum*, 7-12 *Terminalia brownii*, 13-17 *Combretum gueinzii*, 18-23 *Terminalia aemula*, 24-29 *Guiera senegalensis*, 30-35 *Pteleopsis diptera*.

21. Periporate

Plate CXXIV: 1-4 *Plantago lanceolata*, 5-8 *Celosia stuhlmanniana*, 9-11 *Celosia patentiloba*, 12-14 *Celosia trigyna*, 15-18 *Drymaria cordata*, 19-22 *Plantago major*.

Plate CXXV: 1-3 *Calystegia sepium*, 4-5 *Costus spectabilis*.

Plate CXXVI: 1-3 *Plantago palmata*, 4-7 *Dicranolepis oligantha*.

Plate CXXVII: 1-4 *Bosqueia manongarivensis*, 5-8 *Ipomoea donaldsonii*.

Plate CXXVIII: 1-4 *Lychnis* sp., 5-6 *Hewittia sublobata*.

Plate CXXIX: 1-2 *Ipomoea ochracea*, 3-6 *Cerastium indicum*.

Table 1: List of pollen species shown on plates ordered alphabetically by family, genus and species.

Family	Species	Plate
Acanthaceae	<i>Avicennia nitida</i>	LXIV
Acanthaceae	<i>Avicennia officinalis</i>	LXXX
Acanthaceae	<i>Justicia cordata</i>	XXVI
Acanthaceae	<i>Justicia flava</i>	XXVI
Achariaceae	<i>Caloncoba angolensis</i>	LXXX
Achariaceae	<i>Dasylepis assinensis</i>	LV
Achariaceae	<i>Grandidiera boivinii</i>	LVII
Achariaceae	<i>Kiggelaria africana</i>	CVIII
Aizoaceae	<i>Sesuvium sessile</i>	XXVIII
Amaranthaceae	<i>Alternanthera nodiflora</i>	VII
Amaranthaceae	<i>Alternanthera repens</i>	VII
Amaranthaceae	<i>Celosia patentiloba</i>	CXXIV
Amaranthaceae	<i>Celosia stuhlmanniana</i>	CXXIV
Amaranthaceae	<i>Celosia trigyna</i>	CXXIV
Amaryllidaceae	<i>Crinum pauciflorum</i>	XXI
Amaryllidaceae	<i>Crinum powellii</i>	XXI
Anacardiaceae	<i>Lannea humilis</i>	LXXV
Anacardiaceae	<i>Lannea stuhlmannii</i>	LXVI
Anacardiaceae	<i>Lannea triphylla</i>	LXVII
Anacardiaceae	<i>Sclerocarya birrea</i>	LXXV
Anacardiaceae	<i>Spondias mombin</i>	LXXVI
Annonaceae	<i>Artobotrys likimensis</i>	XVI
Annonaceae	<i>Piptostigma mayumbense</i>	VIII
Annonaceae	<i>Uvaria kirkii</i>	XI
Annonaceae	<i>Uvariopsis congensis</i>	IV
Aphloiacae	<i>Aphloia theiformis</i>	CXI
Apocynaceae	<i>Baissea multiflora</i>	XXIV
Apocynaceae	<i>Baissea multiflora</i>	XXVIII
Apocynaceae	<i>Funtumia latifolia</i>	CXV
Apocynaceae	<i>Motandra guineensis</i>	XXIV
Apocynaceae	<i>Tabernaemontana retusa</i>	XCI
Apocynaceae	<i>Tabernaemontana ventricosa</i>	LXX
Apocynaceae	<i>Tylophora sylvatica</i>	XV
Aquifoliaceae	<i>Ilex mitis</i>	XLI
Araceae	<i>Culcasia dinklagei</i>	LXXXI
Araceae	<i>Lemna gibba</i>	XIV
Araceae	<i>Rhektophyllum congense</i>	XCVII
Arecaceae	<i>Ancistrophyllum laurentii</i>	XVIII
Arecaceae	<i>Ancistrophyllum secundiflorum</i>	XVIII
Arecaceae	<i>Borassus aethiopum</i>	VII
Arecaceae	<i>Borassus aethiopum</i>	XIX
Arecaceae	<i>Borassus machadonis</i>	XIX
Arecaceae	<i>Calamus erectus</i>	XVIII
Arecaceae	<i>Calamus gracilis</i>	XVIII
Arecaceae	<i>Elaeis guineensis</i>	XVIII
Arecaceae	<i>Eremospatha sp.</i>	XIX
Arecaceae	<i>Hyphaene natalensis</i>	XIX
Arecaceae	<i>Hyphaene ventricosa</i>	XIX

Arecaceae	<i>Phoenix reclinata</i>	XVIII
Arecaceae	<i>Raphia farinifera</i>	XVIII
Arecaceae	<i>Raphia ruffia</i>	XVIII
Asparagaceae	<i>Asparagus falcatus</i>	XX
Asparagaceae	<i>Chlorophytum floribundum</i>	XX
Asparagaceae	<i>Dracaena camerooniana</i>	XX
Asparagaceae	<i>Dracaena reflexa</i>	XXI
Asteraceae	<i>Artemisia judaica</i>	XCI
Asteraceae	<i>Artemisia sp.</i>	XCI
Asteraceae	<i>Centaurea dimorpha</i>	XC
Asteraceae	<i>Centaurea perrottetii</i>	LXXXIX
Boraginaceae	<i>Heliotropium bacciferum</i>	XLIV
Boraginaceae	<i>Heliotropium subulatum</i>	XLIV
Boraginaceae	<i>Trichodesma africanum</i>	XCII
Boraginaceae	<i>Trichomanes mandiocanum</i>	VII
Brassicaceae	<i>Farsetia stenoptera</i>	XXXV
Burseraceae	<i>Canarium schweinfurthii</i>	XLIII
Burseraceae	<i>Commiphora campestris</i>	XLIX
Burseraceae	<i>Commiphora scheffleri</i>	L
Cannabaceae	<i>Celtis integrifolia</i>	XXXIV
Cannabaceae	<i>Celtis mildbraedii</i>	XXXIII
Cannabaceae	<i>Celtis zenkeri</i>	XXXIII
Cannabaceae	<i>Trema guinensis</i>	XXIV
Cannabaceae	<i>Trema orientalis</i>	XXIV
Capparaceae	<i>Ritchiea capparoides</i>	LXI
Capparaceae	<i>Ritchiea fragariodora</i>	XCIX
Cardioteridaceae	<i>Leptaulus daphnoides</i>	XXXII
Caryophyllaceae	<i>Cerastium indicum</i>	CXXIX
Caryophyllaceae	<i>Drymaria cordata</i>	CXXIV
Caryophyllaceae	<i>Lychnis sp.</i>	CXXVIII
Celastraceae	<i>Cassine parvifolia</i>	C
Celastraceae	<i>Elaeodendron buchananii</i>	XCIV
Celastraceae	<i>Hippocratea affinis</i>	LVI
Celastraceae	<i>Hippocratea africana</i>	LVII
Celastraceae	<i>Maytenus senegalensis</i>	LVI
Celastraceae	<i>Salacia kraussii</i>	LVI
Celastraceae	<i>Salacia pyriformis</i>	CIV
Centroplacaceae	<i>Centroplacus glaucinus</i>	XLV
Chrysobalanaceae	<i>Afrolicania elaeosperma</i>	XCIII
Chrysobalanaceae	<i>Parinari curatellifolia</i>	XLII
Chrysobalanaceae	<i>Parinari holstii</i>	XLVIII
Combretaceae	<i>Combretum aculeatum</i>	CXXIII
Combretaceae	<i>Combretum gueinzii</i>	CXXIII
Combretaceae	<i>Guiera senegalensis</i>	CXXIII
Combretaceae	<i>Pteleopsis diptera</i>	CXXIII
Combretaceae	<i>Strephonema pseudocola</i>	LIX
Combretaceae	<i>Terminalia aemula</i>	CXXIII
Combretaceae	<i>Terminalia brownii</i>	CXXIII
Commelinaceae	<i>Aneilema johnstonii</i>	XXI
Commelinaceae	<i>Commelina africana</i>	XXI
Connaraceae	<i>Agelaea heterophylla</i>	LV
Convolvulaceae	<i>Calystegia sepium</i>	CXXV
Convolvulaceae	<i>Convolvulus trabutianus</i>	CVI
Convolvulaceae	<i>Hewittia sublobata</i>	CXXVIII

Convolvulaceae	<i>Ipomoea donaldsonii</i>	CXXVII
Convolvulaceae	<i>Ipomoea ochracea</i>	CXXIX
Costaceae	<i>Costus spectabilis</i>	CXXV
Davalliaceae	<i>Nephrolepis biserrata</i>	X
Davalliaceae	<i>Nephrolepis exaltata</i>	X
Dichapetalaceae	<i>Dichapetalum mossambicense</i>	XXXI
Dichapetalaceae	<i>Dichapetalum stuhlmannii</i>	XXXIV
Dilleniaceae	<i>Tetracera alnifolia</i>	XXXVI
Dipterocarpaceae	<i>Monotes kerstingii</i>	LXXII
Ebenaceae	<i>Diospyros abyssinica</i>	XCIII
Ebenaceae	<i>Diospyros austroafricana</i>	XCII
Ebenaceae	<i>Diospyros mespiliformis</i>	XLIII
Ericaceae	<i>Erica arborea</i>	IV
Erythroxylaceae	<i>Aneulophus africanus</i>	XXXVI
Euphorbiaceae	<i>Alchornea cordifolia</i>	XLIX
Euphorbiaceae	<i>Alchornea floribunda</i>	XLI
Euphorbiaceae	<i>Anthostema aubryanum</i>	LIV
Euphorbiaceae	<i>Anthostema senegalense</i>	CIII
Euphorbiaceae	<i>Argomuellera macrophylla</i>	XLVI
Euphorbiaceae	<i>Croton gratissimus</i>	XII
Euphorbiaceae	<i>Croton macrostachyus</i>	XII
Euphorbiaceae	<i>Dichostemma</i> sp.	XXXVII
Euphorbiaceae	<i>Discoglypremma caloneura</i>	LI
Euphorbiaceae	<i>Erythrococca bongensis</i>	L
Euphorbiaceae	<i>Euphorbia cussonioides</i>	XCVI
Euphorbiaceae	<i>Euphorbia engleri</i>	XCV
Euphorbiaceae	<i>Euphorbia grandicornis</i>	CII
Euphorbiaceae	<i>Euphorbia hypericifolia</i>	LXIX
Euphorbiaceae	<i>Macaranga schweinfurt</i>	XLVIII
Euphorbiaceae	<i>Mallotus wrayi</i>	LIII
Euphorbiaceae	<i>Martretia quadricornis</i>	XLVII
Euphorbiaceae	<i>Tetrorchidium didymostemon</i>	LXXXIII
Fabaceae (Ceasalpinioideae)	<i>Afzelia bracteata</i>	XXXVIII
Fabaceae (Ceasalpinioideae)	<i>Afzelia quanzensis</i>	CXVII
Fabaceae (Ceasalpinioideae)	<i>Berlinia bifoliolata</i>	LXXVII
Fabaceae (Ceasalpinioideae)	<i>Berlinia grandiflora</i>	XXXIX
Fabaceae (Ceasalpinioideae)	<i>Brachystegia leonensis</i>	XXXVIII
Fabaceae (Ceasalpinioideae)	<i>Brachystegia spiciformis</i>	XXXVIII
Fabaceae (Ceasalpinioideae)	<i>Brachystegia spiciformis</i>	LXXXV
Fabaceae (Ceasalpinioideae)	<i>Cassia burttii</i>	CX
Fabaceae (Ceasalpinioideae)	<i>Cassia longiracemosa</i>	CVIII
Fabaceae (Ceasalpinioideae)	<i>Copaifera carrisoana</i>	CX
Fabaceae (Ceasalpinioideae)	<i>Copaifera gorskiana</i>	LXXIII
Fabaceae (Ceasalpinioideae)	<i>Crudia bracteata</i>	XLII
Fabaceae (Ceasalpinioideae)	<i>Cynometra alexandri</i>	LVIII
Fabaceae (Ceasalpinioideae)	<i>Cynometra pedicellata</i>	CXI
Fabaceae (Ceasalpinioideae)	<i>Daniella oliveri</i>	LXXVIII
Fabaceae (Ceasalpinioideae)	<i>Detarium le-testui</i>	CVII
Fabaceae (Ceasalpinioideae)	<i>Detarium senegalense</i>	XCVII
Fabaceae (Ceasalpinioideae)	<i>Dialium guianense</i>	LXXXII
Fabaceae (Ceasalpinioideae)	<i>Griffonia simplicifolia</i>	LXXIV
Fabaceae (Ceasalpinioideae)	<i>Guibourtia arnoldiana</i>	LXXIV
Fabaceae (Ceasalpinioideae)	<i>Hymenostegia afzelii</i>	CXII
Fabaceae (Ceasalpinioideae)	<i>Isoberlinia angolensis</i>	LXVII

Fabaceae (Ceasalpiniodeae)	<i>Isoberlinia doka</i>	LXVIII
Fabaceae (Ceasalpiniodeae)	<i>Paramacrolobium coeruleum</i>	XXXVIII
Fabaceae (Ceasalpiniodeae)	<i>Piliostigma reticulatum</i>	XXVII
Fabaceae (Faboideae)	<i>Baphia massaiensis</i>	LII
Fabaceae (Faboideae)	<i>Baphia obovata</i>	CXIV
Fabaceae (Faboideae)	<i>Indigofera leptoclada</i>	XLV
Fabaceae (Faboideae)	<i>Lotus arabicus</i>	CV
Fabaceae (Faboideae)	<i>Lotus chazaliei</i>	XLIII
Fabaceae (Faboideae)	<i>Millettia oblata</i>	XCIX
Fabaceae (Faboideae)	<i>Millettia psilopetala</i>	XLVI
Fabaceae (Faboideae)	<i>Millettia tanaensis</i>	CVII
Fabaceae (Faboideae)	<i>Ormocarpum kirkii</i>	LXV
Fabaceae (Faboideae)	<i>Ormocarpum sennoides</i>	CIII
Fabaceae (Faboideae)	<i>Pericopsis angolensis</i>	LXXXIV
Fabaceae (Faboideae)	<i>Pericopsis laxiflora</i>	CXIII
Fabaceae (Faboideae)	<i>Pterocarpus abyssinicus</i>	LXIX
Fabaceae (Faboideae)	<i>Pterocarpus lucens</i>	LXIX
Fabaceae (Faboideae)	<i>Rhynchosia memnonia</i>	LXXXIV
Fabaceae (Faboideae)	<i>Rhynchosia</i> sp.	LXXXVIII
Fabaceae (Faboideae)	<i>Sesbania goetzei</i>	LXXXVII
Fabaceae (Faboideae)	<i>Tephrosia elata</i>	CIX
Fabaceae (Faboideae)	<i>Tephrosia nana</i>	CIX
Fabaceae (Faboideae)	<i>Vigna fischeri</i>	XXX
Fabaceae (Faboideae)	<i>Vigna luteola</i>	XXXI
Fabaceae (Mimosoideae)	<i>Acacia clavigera</i>	II
Fabaceae (Mimosoideae)	<i>Acacia eggelingii</i>	I
Fabaceae (Mimosoideae)	<i>Acacia seyal</i>	I
Fabaceae (Mimosoideae)	<i>Aubrevillea platycarpa</i>	LXXIX
Fabaceae (Mimosoideae)	<i>Calpocalyx brevibracteatus</i>	I
Fabaceae (Mimosoideae)	<i>Dichrostachys cinerea</i>	XVI
Fabaceae (Mimosoideae)	<i>Dichrostachys glomerata</i>	XVI
Fabaceae (Mimosoideae)	<i>Dichrostachys unijuga</i>	VIII
Fabaceae (Mimosoideae)	<i>Entada abyssinica</i>	LII
Fabaceae (Mimosoideae)	<i>Entada pursaetha</i>	XXXII
Fabaceae (Mimosoideae)	<i>Entada umbonata</i>	XXX
Fabaceae (Mimosoideae)	<i>Mimosa strigillosa</i>	IV
Fabaceae (Mimosoideae)	<i>Parkia bussei</i>	III
Fabaceae (Mimosoideae)	<i>Parkia inundabilis</i>	II
Fabaceae (Mimosoideae)	<i>Parkia velutina</i>	III
Fabaceae (Mimosoideae)	<i>Piptadenia africana</i>	LXXXV
Fabaceae (Mimosoideae)	<i>Prosopis africana</i>	LXXI
Fabaceae (Mimosoideae)	<i>Prosopis alpataco</i>	LXXXVIII
Fabaceae (Mimosoideae)	<i>Xylia evansii</i>	I
Gentianaceae	<i>Anthocleista grandiflora</i>	XXXIII
Gerrardinaceae	<i>Gerrardina foliosa</i>	XL
Gunneraceae	<i>Gunnera chilensis</i>	XXXV
Hernandiaceae	<i>Illicera appendiculata</i>	XIV
Hernandiaceae	<i>Illicera rhodantha</i>	VI
Icacinaceae	<i>Iodes kamerunensis</i>	XIV
Icacinaceae	<i>Iodes ovalis</i>	XXIV
Icacinaceae	<i>Iodes ovalis</i>	XXVII
Iridaceae	<i>Acidanthera brevicollis</i>	VI
Iridaceae	<i>Acidanthera brevicollis</i>	XXI
Irvingiaceae	<i>Irvingia smithii</i>	LXXXII

Kirkiaceae	<i>Kirkia acuminata</i>	XXIX
Lamiaceae	<i>Leucas calostachys</i>	XXXVI
Lamiaceae	<i>Premna maxima</i>	XXXV
Lamiaceae	<i>Premna resinosa</i>	XXXV
Lamiaceae	<i>Vitex amboniensis</i>	XXXVI
Lamiaceae	<i>Vitex doniana</i>	XXXVI
Lecythidaceae	<i>Napoleona imperialis</i>	XXXV
Lecythidaceae	<i>Petersia africana</i>	XXXIX
Lecythidaceae	<i>Scytopetalum tieghemii</i>	XXXVII
Lecythidaceae	<i>Scytopetalum tieghemii</i>	LXVIII
Lindsaeaceae	<i>Lonchitis currori</i>	X
Malpighiaceae	<i>Acridocarpus macrocalyx</i>	LXXIII
Malvaceae	<i>Bombax brevicuspe</i>	XXVIII
Malvaceae	<i>Bombax buonopozense</i>	XXVIII
Malvaceae	<i>Ceiba pentandra</i>	XXIX
Malvaceae	<i>Cola gigantea</i>	CI
Malvaceae	<i>Cola millenii</i>	CI
Malvaceae	<i>Cola nitida</i>	XCVIII
Malvaceae	<i>Corchorus fascicularis</i>	LIX
Malvaceae	<i>Corchorus trilocularis</i>	LXXVIII
Malvaceae	<i>Grewia bicolor</i>	LXII
Malvaceae	<i>Grewia glandulosa</i>	CIV
Malvaceae	<i>Hildegardia barteri</i>	LXXXI
Malvaceae	<i>Nesogordonia fertilis</i>	XXX
Malvaceae	<i>Nesogordonia parvifolia</i>	XXXI
Malvaceae	<i>Scaphopetalum letestui</i>	XI
Malvaceae	<i>Scaphopetalum thonneri</i>	VIII
Malvaceae	<i>Sterculia tragacantha</i>	LXXI
Malvaceae	<i>Triplochiton scleroxylon</i>	XXX
Marantaceae	<i>Sarcophrynum brachystachyum</i>	CVIII
Meliaceae	<i>Turraeanthus africana</i>	XXXIV
Menispermaceae	<i>Cissampelos mucronata</i>	XXXVII
Menispermaceae	<i>Tiliacora funifera</i>	XI
Moraceae	<i>Antiaris toxicaria</i>	XXIII
Moraceae	<i>Bosqueia manongarivensis</i>	CXXVII
Moraceae	<i>Chlorophora excelsa</i>	XXIII
Moraceae	<i>Ficus ingens</i>	XXIII
Myricaceae	<i>Myrica sp.</i>	XXII
Myristicaceae	<i>Pycnanthus dinklagei</i>	VII
Myrtaceae	<i>Eugenia michoacanensis</i>	XXII
Myrtaceae	<i>Syzygium guineense</i>	XXII
Nymphaeaceae	<i>Nymphaea caerulea</i>	XI
Nymphaeaceae	<i>Nymphaea caerulea</i>	XX
Nymphaeaceae	<i>Nymphaea lotus</i>	X
Nymphaeaceae	<i>Nymphaea lotus</i>	XX
Ochnaceae	<i>Lophira alata</i>	XCII
Olacaceae	<i>Coula edulis</i>	XXVIII
Olacaceae	<i>Heisteria parvifolia</i>	XXVIII
Oleaceae	<i>Olea hochstetteri</i>	LIII
Orobanchaceae	<i>Striga forbesii</i>	XXXIV
Pandaceae	<i>Microdesmis sp.</i>	CXIII
Pandanaceae	<i>Pandanus kirkii</i>	XV
Pandanaceae	<i>Pandanus livingstonianus</i>	VIII
Passifloraceae	<i>Adenia nicobarica</i>	LV

Passifloraceae	<i>Barteria acuminata</i>	XIII
Pedaliaceae	<i>Sesamum angustifolium</i>	CXIX
Pedaliaceae	<i>Sesamum indicum</i>	CXVI
Periscaceae	<i>Medusandra richardsiana</i>	XLIX
Phyllanthaceae	<i>Amanoa strobilacea</i>	LI
Phyllanthaceae	<i>Bridelia micrantha</i>	CII
Phyllanthaceae	<i>Hymenocardia acida</i>	XXXIII
Phyllanthaceae	<i>Maesobotrya barteri</i>	C
Phyllanthaceae	<i>Maesobotrya hirtella</i>	LXVI
Phyllanthaceae	<i>Securinega virosa</i>	XCVII
Phyllanthaceae	<i>Spondianthus preussii</i>	LX
Phyllanthaceae	<i>Thecacoris gymnogyne</i>	XL
Phyllanthaceae	<i>Uapaca bojeri</i>	LXXXIV
Phyllanthaceae	<i>Uapaca heudelotii</i>	CV
Piperaceae	<i>Peperomia</i> sp.	X
Plantaginaceae	<i>Plantago lanceolata</i>	CXXIV
Plantaginaceae	<i>Plantago major</i>	CXXIV
Plantaginaceae	<i>Plantago palmata</i>	CXXVI
Poaceae	<i>Guaduella oblonga</i>	XVII
Podocarpaceae	<i>Podocarpus milanjanus</i>	V
Polygalaceae	<i>Atroxima afzeliana</i>	CXXI
Polygalaceae	<i>Securidaca longepedunculata</i>	CXXI
Proteaceae	<i>Protea susannae</i>	XXVIII
Proteaceae	<i>Protea trichanthera</i>	XXXII
Putranjivaceae	<i>Drypetes gerrardii</i>	LXIV
Rhamnaceae	<i>Zizyphus mauritiana</i>	LXXXII
Rhizophoraceae	<i>Cassipourea flanaganii</i>	LXVI
Rhizophoraceae	<i>Rhizophora mangle</i>	XCVI
Rhizophoraceae	<i>Rhizophora mucronata</i>	XCVIII
Rosaceae	<i>Cliffortia nitidula</i>	XLVIII
Rosaceae	<i>Hagenia abyssinica</i>	LXXII
Rosaceae	<i>Prunus africana</i>	XLVIII
Rosaceae	<i>Pygeum africanum</i>	LXXIII
Rosaceae	<i>Rubus scheffleri</i>	XLII
Rubiaceae	<i>Borreria densiflora</i>	CXVI
Rubiaceae	<i>Borreria ruelliae</i>	CXVI
Rubiaceae	<i>Crossopteryx febrifuga</i>	LX
Rubiaceae	<i>Diodia aulacosperma</i>	CXX
Rubiaceae	<i>Diodia scandens</i>	CXVIII
Rubiaceae	<i>Gaertnera paniculata</i>	LXXXIII
Rubiaceae	<i>Ixora aneimenodesma</i>	LIV
Rubiaceae	<i>Ixora brachypoda</i>	LXXXVII
Rubiaceae	<i>Lasianthus africanus</i>	XXIX
Rubiaceae	<i>Mitracarpus hirtus</i>	CXIX
Rubiaceae	<i>Mitracarpus verticillatus</i>	CXX
Rubiaceae	<i>Mitragyna inermis</i>	LXXXII
Rubiaceae	<i>Morinda citrifolia</i>	XIII
Rubiaceae	<i>Morinda citrifolia</i>	XXV
Rubiaceae	<i>Nauclea diderrichii</i>	CXII
Rubiaceae	<i>Nauclea esculenta</i>	C
Rubiaceae	<i>Plectronia vulgaris</i>	XXIX
Rubiaceae	<i>Psychotria fractinervata</i>	XXXI
Rubiaceae	<i>Psychotria goetzei</i>	XII
Rubiaceae	<i>Sabicea floribunda</i>	XXVIII

Rutaceae	<i>Fagara macrophylla</i>	LIII
Rutaceae	<i>Teclea villosa</i>	LXXVII
Rutaceae	<i>Vepris eugeniiifolia</i>	CXI
Rutaceae	<i>Vepris gossweileri</i>	CV
Rutaceae	<i>Vepris humbertii</i>	LXII
Rutaceae	<i>Vepris uguenensis</i>	XL
Rutaceae	<i>Zanthoxylum procerum</i>	CVI
Salicaceae	<i>Calantica jalbertii</i>	LVIII
Salicaceae	<i>Casearia engleri</i>	XCIII
Salicaceae	<i>Dissomeria crenata</i>	CIX
Salicaceae	<i>Flacourtie indica</i>	LXIV
Salicaceae	<i>Homalium buchholzii</i>	CX
Salicaceae	<i>Oncoba dentata</i>	CVII
Sapindaceae	<i>Allophylus africanus</i>	XXVII
Sapindaceae	<i>Blighia unijugata</i>	LXI
Sapindaceae	<i>Blighia wildemaniana</i>	LVIII
Sapindaceae	<i>Cardiospermum corindum</i>	LXXIX
Sapindaceae	<i>Cardiospermum grandiflorum</i>	XXX
Sapindaceae	<i>Chytranthus obliquinervis</i>	LXXXVI
Sapindaceae	<i>Chytranthus sacleuxii</i>	LXXXVI
Sapindaceae	<i>Dodonaea viscosa</i>	LXX
Sapindaceae	<i>Lecaniodiscus cupanioides</i>	LXXXVI
Sapindaceae	<i>Paullinia pinnata</i>	XXXII
Sapindaceae	<i>Placodiscus amaniensis</i>	CXII
Sapotaceae	<i>Neolemonniera clitandrifolia</i>	XCIV
Simaroubaceae	<i>Hannoa klaineana</i>	LXV
Simaroubaceae	<i>Hannoa undulata</i>	LXV
Simaroubaceae	<i>Odyendea gabunensis</i>	CVI
Thymelaeaceae	<i>Dicranolepis oligantha</i>	CXXVI
Thymelaeaceae	<i>Dicranolepis usambarica</i>	XIII
Typhaceae	<i>Typha angustifolia</i>	XVII
Typhaceae	<i>Typha australis</i>	XVII
Typhaceae	<i>Typha capensis</i>	XVII
Urticaceae	<i>Musanga leo-errerae</i>	XXIII
Urticaceae	<i>Musanga smithii</i>	XXIII
Violaceae	<i>Rinorea oblongifolia</i>	CXIII
Violaceae	<i>Rinorea welwitschii</i>	CXIII
Vitaceae	<i>Cissus petiolata</i>	LX
Vitaceae	<i>Cissus quadrangularis</i>	LXII
Xyridaceae	<i>Xyris aristata</i>	IX
Xyridaceae	<i>Xyris montana</i>	IX
Xyridaceae	<i>Xyris welwitschii</i>	IX
Zygophyllaceae	<i>Balanites aegyptiacus</i>	XLVII
Zygophyllaceae	<i>Balanites glaber</i>	XLV
Zygophyllaceae	<i>Tribulus terrestris</i>	XIII