



(51) International Patent Classification:

AOIC 1/06 (2006.01) *A01N 25/00* (2006.01)
C12N 1/04 (2006.01) *A01N 25/14* (2006.01)
C12N 11/10 (2006.01) *A01P 21/00* (2006.01)
C12R 1/80 (2006.01) *C09D 105/00* (2006.01)
A01N 63/04 (2006.01)

(21) International Application Number:

PCT/US20 16/050647

(22) International Filing Date:

8 September 2016 (08.09.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/217,250 11 September 2015 (11.09.2015) US
62/273,054 30 December 2015 (30.12.2015) US
62/296,766 18 February 2016 (18.02.2016) US
62/343,250 31 May 2016 (31.05.2016) US

(71) Applicant: **NOVOZYMES BIOAG A/S** [DK/DK];
Krogshoejvej 36, DK-2880 Bagsvaerd (DK).

(72) Inventor; and

(71) Applicant (for BW only): **KELLAR, Kenneth Edmund** [US/US]; 108 TW Alexander Dr., Bldg. 1A, Durham, North Carolina 27709 (US).

(72) Inventors: **KANG, Yaowei**; 108 TW Alexander Dr., Bldg. 1A, Durham, North Carolina 27709 (US). **PELLIGRA, Claire**; 108 TW Alexander Dr., Bldg. 1A, Durham, North Carolina 27709 (US). **LOOZE, Emily**; 5400 Corporate

Circle, Salem, Virginia 24153 (US). **BURKLEW, Caitlin**; 5400 Corporate Circle, Salem, Virginia 24153 (US). **WYSINSKI, Anna**; 5400 Corporate Circle, Salem, Virginia 24153 (US). **LELAND, Jarrod**; 5400 Corporate Circle, Salem, Virginia 24153 (US). **DOUGHAN, Ben**; 5400 Corporate Circle, Salem, Virginia 24153 (US). **FETHE, Michael Harrison**; 108 TW Alexander Dr., Bldg. 1A, Durham, North Carolina 27709 (US). **TRAHAN, Ashley Delanie**; 108 TW Alexander Dr., Bldg. 1A, Durham, North Carolina 27709 (US). **GREENSHIELDS, Dave**; 702 McPherson Ave, Saskatoon, Saskatchewan S7N0X8 (CA). **WOODS, Kristi**; 5400 Corporate Circle, Salem, Virginia 24153 (US).

(74) Agents: **RUCKER, Adam L.** et al; Novozymes North America, Inc., 60 E. 42nd St., Suite 700, New York, New York 10165 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on nextpage]

(54) Title: STABLE INOCULANT COMPOSITIONS AND METHODS FOR PRODUCING SAME




FIG. 1

(57) Abstract: The present disclosure provides non-aqueous inoculant compositions and methods for enhancing the survival and/or stability of microbial spores in an inoculant composition. In some embodiments, inoculant compositions of the present disclosure comprise microbial spores, one or more dispersants, one or more dust suppressants and a solid non-aqueous carrier.

WO 2017/044545 A1

GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT,
LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE,

SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

STABLE INOCULANT COMPOSITIONS AND METHODS FOR PRODUCING SAME**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application claims priority under 35 U.S.C. § 119 to U.S. Provisional Patent Application Nos. 62/217,250, filed September 11, 2015; 62/273,054, filed December 30, 2015; 62/296,766, filed February 18, 2016; and 62/343,250, filed May 31, 2016, the disclosure of each of which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present disclosure relates to compositions and methods for enhancing the stability and survival of microbial spores in inoculant compositions.

BACKGROUND OF THE INVENTION

Inoculant compositions comprising agriculturally beneficial microorganisms are well known in the art. *See, e.g.*, U.S. Patent Nos. 5,484,464; 5,586,411; 5,695,541; 5,804,208; 5,916,029; 6,569,425; 6,808,917; 6,824,772; 7,429,477; 8,148,138; 8,278,247; 8,445,256; 8,883,679; 8,921,089; 8,999,698; 9,017,442; 9,101,088; 9,234,251; 9,340,464.

Because the effectiveness of such inoculant compositions generally depends on the ability of the microorganisms therein to survive and propagate following application, much effort has been made to increase the stability of agriculturally beneficial microorganisms in inoculant compositions. *See, e.g.*, U.S. Patent Nos. 8,011,132 (describing a method of adding trehalose, sucrose or glycerol to the substantially stationary phase of fermentation) and 9,090,884 (describing the microencapsulation of microorganisms in a water-soluble encapsulating material).

Nevertheless, there remains a need for improved compositions and methods for enhancing the stability and survival of microorganisms in inoculant compositions.

SUMMARY OF THE CLAIMED INVENTION

The present disclosure provides stable inoculant compositions and methods for enhancing the survival and/or stability of microbial spores in inoculant compositions.

A first aspect of the present disclosure is a non-aqueous inoculant composition comprising *Penicillium* spores, one or more dispersants and a solid non-aqueous carrier. In some embodiments, the inoculant composition comprises one or more dust suppressants, one or more pesticides, one or more lipo-chitooligosaccharides, one or more chitooligosaccharides, one or more chitinous compounds, one or more flavonoids and/or one or more drying agents.

A second aspect of the present disclosure is a coated plant propagation material comprising a plant propagation material and a coating that covers at least a portion of an outer surface of the plant propagation material, said coating comprising a non-aqueous inoculant composition of the present disclosure.

A third aspect of the present disclosure is a kit comprising a coated plant propagation material of the present disclosure and a container housing the coated plant propagation material.

A fourth aspect of the present disclosure is a plant germinated from a coated plant propagation material of the present disclosure.

A fifth aspect of the present disclosure is a plant part harvested from a plant that was germinated from a coated plant propagation material of the present disclosure.

A sixth aspect of the present disclosure is a processed product derived from a plant that was germinated from a coated plant propagation material of the present disclosure.

A seventh aspect of the present disclosure is a crop comprising a plurality of plants germinated from coated plant propagation materials of the present disclosure.

An eighth aspect of the present disclosure is a method that comprises applying a non-aqueous inoculant composition of the present disclosure to a plant propagation material.

A ninth aspect of the present disclosure is a method that comprises, consists essentially of or consisting of planting a coated plant propagation material of the present disclosure.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1-3 are graphs showing the survivability of spray-dried *Penicillium bilaiae* on corn seeds and 50% relative humidity and 10°C, 20°C or 30°C, respectively. Triangles = corn seeds coated with a spray-dried composition comprising *Penicillium bilaiae* spores (10% w/w), MALTRIN QD® M580 (78.469% w/w), maltose monohydrate (8.719% w/w) and BIOSOFT® N23-3 (2.813% w/w). Squares = corn seeds coated with a spray-dried composition comprising *Penicillium bilaiae* spores (10% w/w), MALTRIN QD® M580 (74.879% w/w), maltose monohydrate (8.320% w/w), MULTIWET® MO-85P-PW-(AP) (2.750% w/w) and SUNSPRAY® 6N (4.051% w/w). Diamonds = corn seeds coated with a commercially available wettable powder comprising *Penicillium bilaiae* spores.

DETAILED DESCRIPTION

The present disclosure is explained in greater detail below. This description is not intended to be a detailed catalog of all the different ways in which the invention may be implemented or of all the features that may be added to the instant invention. For example, features illustrated with respect to one embodiment may be incorporated into other embodiments and features illustrated with respect to a particular embodiment may be deleted from that embodiment. In addition, numerous variations and additions to the various embodiments suggested herein, which do not depart from the instant invention, will be apparent to those skilled in the art in light of the instant disclosure. Hence, the following specification is intended to illustrate some particular embodiments of the invention and not to exhaustively specify all permutations, combinations and variations thereof.

The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention.

Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the specification and relevant art and should not be interpreted in an idealized or overly formal sense unless expressly so defined herein. For the sake of brevity and/or clarity, well-known functions or constructions may not be described in detail.

As used herein, the singular forms "a," "an," and "the" are intended to include the plural forms as well, unless the context clearly indicates otherwise. Thus, unless the context clearly indicates otherwise, "a maltodextrin" is to be interpreted as "one or more maltodextrins," "a microorganism" is to be interpreted as "one or more microorganisms," "a lipo-chitooligosaccharide" is to be interpreted as "one or more lipo-chitooligosaccharides," etc.

As used herein, the term "about," when used in reference to a measurable value such as an amount of mass, dose, time, temperature and the like, is meant to encompass variations of 0.1%, 0.25%, 0.5%, 0.75%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the specified amount.

Unless otherwise indicated, all numerical values in the specification are to be understood as being modified by the term "about."

As used herein, the term "agriculturally beneficial agent" refers to any agent (e.g., chemical or biological agent) or combination of agents the application of which causes or provides a beneficial and/or useful effect in agriculture including, but not limited to, agriculturally beneficial microorganisms, biostimulants, nutrients, pesticides (e.g., fungicides, herbicides, insecticides, and nematocides) and plant signal molecules.

As used herein, the term "agriculturally beneficial microorganism" refers to a microorganism having at least one agriculturally beneficial property (e.g., the ability to fix nitrogen, the ability to solubilize phosphate and/or the ability to produce an agriculturally beneficial agent, such as a plant signal molecule).

As used herein, the term "agriculturally acceptable carrier" refers to a material that can be used to deliver an agriculturally beneficial agent to a plant, plant part or plant growth medium (e.g., soil). As used herein, the term "soil-compatible carrier" refers to a material that can be added to a soil without causing/having an unduly adverse effect on plant growth, soil structure, soil drainage, or the like. As used herein, the term "seed-compatible carrier" refers to a material that can be added to a seed without causing/having an unduly adverse effect on the seed, the plant that grows from the seed, seed germination, or the like. As used herein, the term "foliar-compatible carrier" refers to a material that can be added to a plant or plant part without causing/having an unduly adverse effect on the plant, plant part, plant growth, plant health, or the like.

As used herein, the term "and/or" is intended to include any and all combinations of one or more of the associated listed items, as well as the lack of combinations when interpreted in the alternative ("or").

As used herein, the term "aqueous" refers to a composition that contains more than a trace amount of water (i.e., more than 0.5% water by weight, based upon the total weight of the composition).

As used herein, the term "biostimulant" refers to an agent or combination of agents the application of which enhances one or more metabolic and/or physiological processes of a plant or plant part (e.g., carbohydrate biosynthesis, ion uptake, nucleic acid uptake, nutrient delivery, photosynthesis and/or respiration).

As used herein, the term "colony forming unit" refers to a microbial cell/spore capable of propagating on or in a substrate (e.g., a soil) when conditions (e.g., temperature, moisture, nutrient availability, pH, etc.) are favorable for microbial growth.

As used herein, the terms "comprise," "comprises," "comprising," "include," "includes" and "including" specify the presence of stated features, steps, operations, elements and/or components, but do not preclude the presence or addition of one or more other features, steps, operations, elements, components and/or groups thereof.

As used herein, the term "consists essentially of" (and grammatical variants thereof), as applied to the compositions and methods of the present disclosure, means that the compositions/methods may contain additional components so long as the additional components do not materially alter the composition/method. The term "materially alter," as applied to a composition/method, refers to an increase or decrease in the effectiveness of the composition/method of at least about 20% or more. For example, a component added to an inoculant composition of the present disclosure "materially alters" the composition if it increases or decreases the composition's ability to enhance microbial survival by at least about 50%.

As used herein, the terms "effective amount," "effective concentration," and "effective dosage" (and grammatical variants thereof) refer to an amount, concentration or dosage that is sufficient to cause a desired effect (e.g., enhanced microbial survival). The absolute value of the amount/concentration/dosage that is sufficient to cause the desired effect may be affected by factors such as the type and magnitude of effect desired, the type, size and volume of seeds to which the inoculant composition will be applied, the stability of the microorganisms in the

inoculant composition and the storage conditions (e.g., temperature, relative humidity, duration). Those skilled in the art will understand how to select an effective amount/concentration/dosage using routine dose-response experiments.

As used herein, the term "enhanced stability" refers to an improvement in one or more characteristics of microbial stability as compared to one or more controls (e.g., a control composition that is identical to an inoculant composition of the present disclosure except that it lacks one or more of the components found in the inoculant composition of the present disclosure). Exemplary microbial stability characteristics include, but are not limited to, the ability to cause a desired effect (e.g., enhanced plant yield and/or increased pesticidal activity) after being coated on a seed and/or stored for a defined period of time and survival rate after being coated on a seed and/or stored for a defined period of time. A microorganism that exhibits improvement in one or more microbial stability characteristics as compared to a control microorganism when each is subjected to the same conditions (e.g., seed coating and storage conditions) displays enhanced stability and can be referred to as a "stable microorganism." An inoculant composition that improves one or more microbial stability characteristics of the microorganism(s) contained therein as compared to a control composition (e.g., a control composition that is identical to the inoculant composition except that it lacks one or more of the components found in the inoculant composition) provides enhanced stability and can be referred to as a "stable inoculant composition."

As used herein, the term "enhanced survival" refers to an improvement in the survival rate of one or more microorganisms in an inoculant composition as compared to one or more microorganisms in a control composition (e.g., a control composition that is identical to an inoculant composition of the present disclosure except that it lacks one or more of the components found in the inoculant composition of the present disclosure). An inoculant composition that improves the survival rate of one or more of the microorganisms contained therein as compared to a control composition (e.g., a control composition that is identical to the inoculant composition except that it lacks one or more of the components found in the inoculant composition) provides enhanced survival and can be referred to as a stable inoculant composition.

As used herein, the terms "enhanced yield" and "enhanced plant yield" refer to an improvement in one or more characteristics of plant yield as compared to one or more control plants (e.g., a control plant germinated from an untreated seed). Exemplary plant yield characteristics include, but are not limited to, biomass; bushels per acre; grain weight per plot (GWTPP); nutritional content; percentage of plants in a given area (e.g., plot) that fail to produce grain; yield at standard moisture percentage (YSMP), such as grain yield at standard moisture percentage (GYSMP); yield per plot (YPP), such as grain weight per plot (GWTPP); and yield reduction (YRED). Unless otherwise indicated, references to enhanced plant yield are to be interpreted as meaning that compositions and methods of the present disclosure may be capable of enhancing plant yield by enhancing nutrient availability, improving soil characteristics, etc. and are not to be interpreted as suggesting that compositions and methods of the present disclosure act as plant growth regulators. As used herein, the term "foliage" refers to those portions of a plant that normally grow above the ground, including, but not limited to, leaves, stalks, stems, flowers, fruiting bodies and fruits.

As used herein, the terms "foliar application," "foliarly applied" and grammatical variations thereof, refer to the application of one or more active ingredients to the foliage of a plant (e.g., to the leaves of the plant). Application may be effected by any suitable means, including, but not limited to, spraying the plant with a composition comprising the active ingredient(s). In some embodiments, the active ingredient(s) is/are applied to the leaves, stems and/or stalk of the plant and not to the flowers, fruiting bodies or fruits of the plant.

As used herein, the term "glass transition temperature" and its abbreviation "T_g" refer to the midpoint of the temperature range over which a composition transitions from a glassy state to a rubbery state.

As used herein, the term "glassy state" refers to an amorphous solid.

As used herein, the terms "inoculant composition" and "inoculum" refer to compositions comprising microbial cells and/or spores, said cells/spores being capable of propagating on or in a substrate (e.g., a soil) when conditions (e.g., temperature, moisture, nutrient availability, pH, etc.) are favorable for microbial growth.

As used herein, the term "isomer" includes all stereoisomers of the compounds and/or molecules to which it refers, including enantiomers and diastereomers, as well as all conformers, rotamers and tautomers, unless otherwise indicated. Compounds and/or molecules disclosed herein include all enantiomers in either substantially pure levorotatory or dextrorotatory form, or in a racemic mixture, or in any ratio of enantiomers. Where embodiments disclose a (D)-enantiomer, that embodiment also includes the (L)-enantiomer; where embodiments disclose a (L)-enantiomer, that embodiment also includes the (D)-enantiomer. Where embodiments disclose a (+)-enantiomer, that embodiment also includes the (-)-enantiomer; where embodiments disclose a (-)-enantiomer, that embodiment also includes the (+)-enantiomer. Where embodiments disclose a (S)-enantiomer, that embodiment also includes the (R)-enantiomer; where embodiments disclose a (R)-enantiomer, that embodiment also includes the (S)-enantiomer. Embodiments are intended to include any diastereomers of the compounds and/or molecules referred to herein in diastereomerically pure form and in the form of mixtures in all ratios. Unless stereochemistry is explicitly indicated in a chemical structure or chemical name, the chemical structure or chemical name is intended to embrace all possible stereoisomers, conformers, rotamers and tautomers of compounds and/or molecules depicted.

As used herein, the term "nitrogen fixing organism" refers to an organism capable of converting atmospheric nitrogen (N₂) into a form that may be utilized by a plant or plant part (e.g., ammonia (NH₃), ammonium (NH₄⁺), etc.).

As used herein, the term "non-aqueous" refers to a composition that comprises no more than a trace amount of water (i.e., no more than 0.5% water by weight, based upon the total weight of the composition).

As used herein, the term "nutrient" refers to a compound or element useful for nourishing a plant (e.g., vitamins, macrominerals, micronutrients, trace minerals, organic acids, etc. that are necessary for plant growth and/or development).

As used herein, the term "onset temperature" refers to the temperature at which a composition begins the transition from a glassy state to a rubbery state.

As used herein, the term "*Penicillium bilaiae*" is intended to include all iterations of the species name, such as "*Penicillium bilaji*" and "*Penicillium bilaii*."

As used herein, the term "pest" includes any organism or virus that negatively affects a plant, including, but not limited to, organisms and viruses that spread disease, damage host plants and/or compete for soil nutrients. The term "pest" encompasses organisms and viruses that are known to associate with plants and to cause a detrimental effect on the plant's health and/or vigor. Plant pests include, but are not limited to, arachnids (e.g., mites, ticks, spiders, etc.), bacteria, fungi, gastropods (e.g., slugs, snails, etc.), invasive plants (e.g., weeds), insects (e.g., white flies, thrips, weevils, etc.), nematodes (e.g., root-knot nematode, soybean cyst nematode, etc.), rodents and viruses (e.g., tobacco mosaic virus (TMV), tomato spotted wilt virus (TSWV), cauliflower mosaic virus (CaMV), etc.).

As used herein, the terms "pesticide" and "pesticidal" refer to agents or combinations of agents the application of which is toxic to a pest (i.e., kills a pest, inhibits the growth of a pest and/or inhibits the reproduction of a pest). Non-limiting examples of pesticides include fungicides, herbicides, insecticides, and nematicides, etc.

As used herein, the term "phosphate-solubilizing microorganism" refers to a microorganism capable of converting insoluble phosphate into a soluble form of phosphate.

As used herein, the term "plant" includes all plant populations, including, but not limited to, agricultural, horticultural and silvicultural plants. The term "plant" encompasses plants obtained by conventional plant breeding and optimization methods (e.g., marker-assisted selection) and plants obtained by genetic engineering, including cultivars protectable and not protectable by plant breeders' rights.

As used herein, the term "plant cell" refers to a cell of an intact plant, a cell taken from a plant, or a cell derived from a cell taken from a plant. Thus, the term "plant cell" includes cells within seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves, shoots, gametophytes, sporophytes, pollen and microspores.

As used herein, the term "plant part" refers to any part of a plant, including cells and tissues derived from plants. Thus, the term "plant part" may refer to any of plant components or organs (e.g., leaves, stems, roots, etc.), plant tissues, plant cells and seeds. Examples of plant parts, include, but are not limited to, anthers, embryos, flowers, fruits, fruiting bodies, leaves, ovules, pollen, rhizomes, roots, seeds, shoots, stems and tubers, as well as scions, rootstocks, protoplasts, calli and the like.

As used herein, the term "plant propagation material" refers to a plant part from which a whole plant can be generated. Examples of plant propagation materials include, but are not limited to, cuttings (e.g., leaves, stems), rhizomes, seeds, tubers and cells/tissues that can be cultured into a whole plant.

As used herein, the term "protectant" refers to an agent or combination of agents the application of which enhances the survival and/or stability of a microorganism in an inoculant composition.

As used herein, the term "rubbery state" refers to an amorphous, viscoelastic liquid.

As used herein, the terms "signal molecule" and "plant signal molecule" refer to an agent that, when applied to a plant or plant part, results in enhanced growth and/or development as compared to untreated plants or plant parts (e.g., seeds and plants harvested from untreated seeds). Non-limiting examples of signal molecules include lipooligosaccharides, chitooligosaccharides, chitinous compounds, flavonoids, jasmonic acid or derivatives thereof, linoleic acid or derivatives thereof, linolenic acid or derivatives thereof, karrikins, etc.

As used herein, the term "solid" refers to a composition that is neither a gas nor a liquid.

As used herein, the terms "spore" and "microbial spore" refer to a microorganism in its dormant, protected state.

As used herein with respect to inoculant compositions, the term "stable" refers to an inoculant composition in which microorganisms exhibit enhanced stability and/or enhanced survival. In general, an inoculant composition may be labeled "stable" if it improves the survival rate and/or at least one microbial stability characteristic of at least one microorganism contained therein.

As used herein with respect to microbial spores, the term "survival rate" refers to the percentage of microbial spores that are viable (i.e., capable of propagating on or in a substrate (e.g., on a seed and/or in a soil) when conditions (e.g., temperature, moisture, nutrient availability, pH, etc.) are favorable for microbial growth) at a given period of time.

While certain aspects of the present disclosure will hereinafter be described with reference to embodiments thereof, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope of the present disclosure as defined by the claims.

All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety, except insofar as they contradict any disclosure expressly set forth herein.

The present disclosure provides inoculant compositions and methods for enhancing the stability and/or survival of microbial spores.

The present disclosure provides non-aqueous inoculant compositions comprising, consisting essentially of, or consisting of one or more microbial spores and a non-aqueous carrier.

In some embodiments, inoculant compositions of the present disclosure improve the stability of one or more microbial spores contained therein by at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition (e.g., a control composition that is identical to the inoculant composition of the present disclosure except that it lacks one or more of the components found in the inoculant composition and/or contains one or more components not found in the inoculant composition of the present disclosure). For example, inoculant compositions of the present disclosure may improve one or more microbial stability characteristics of one or more of the microbial spores contained therein by at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition that is identical to the inoculant composition except that lacks one or more of the protectants/dispersants found in the inoculant composition and/or comprises a reduced amount of one or more of the protectants/dispersants found in the inoculant composition.

In some embodiments, microbial spores remain viable in inoculant compositions of the present disclosure for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more (e.g., at least 12, 16, 20, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more when stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity).

In some embodiments, inoculant compositions of the present disclosure improve the survival rate of one or more microbial spores contained therein by at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition (e.g., a control composition that is identical to the inoculant composition of the present disclosure except that it lacks one or more of the components found in the inoculant composition and/or contains one or more components not found in the inoculant composition of the present disclosure). For example, inoculant compositions of the present disclosure may improve the survival rate of one or more of the microbial spores contained therein by at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition that is identical to the inoculant composition except that lacks one or more of the protectants/dispersants found in the inoculant composition and/or comprises a reduced amount of one or more of the protectants/dispersants found in the inoculant composition.

In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of the microbial spores survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70% or more of the microbial spores survive when the inoculant composition is stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25,

30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of the microbial spores survive when the inoculant composition is desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70% or more of the microbial spores survive when the inoculant composition is coated on a seed, dried and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/gram or milliliter or more of the microbial spores survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 cfu/seed or more of the microbial spores survive when the inoculant composition is stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30° for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/gram or milliliter or more of the microbial spores survive when the inoculant composition is desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, inoculant compositions of the present disclosure improve the survival of one or more of the microbial spores contained therein to the extent that at least 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 cfu/seed or more of the microbial spores survive when the inoculant composition is coated on a seed, dried and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure improve both the survival rate and one, two, three, four, five, six, seven, eight, nine, ten or more microbial stability characteristics of the microbial spore(s) contained therein.

In some embodiments, inoculant compositions of the present disclosure improve the dispersion of one or more microbial spores contained therein by at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65,

70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition (e.g., a control composition that is identical to the inoculant composition of the present disclosure except that it lacks one or more of the components found in the inoculant composition and/or contains one or more components not found in the inoculant composition of the present disclosure). For example, inoculant compositions of the present disclosure may improve the dispersion of one or more of the microbial spores contained therein by at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition that is identical to the inoculant composition except that lacks one or more of the protectants/dispersants found in the inoculant composition and/or comprises a reduced amount of one or more of the protectants/dispersants found in the inoculant composition.

In some embodiments, inoculant compositions of the present disclosure improve the dispersion of one or more of the microbial spores contained therein to the extent that at least about 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of the microbial spores are present as single spores (rather than as members of a clump comprising two or more microbial spores).

Inoculant compositions of the present disclosure may be used to improve any suitable microbial stability characteristic(s), including, but not limited to, the ability of microbial spores therein to enhance plant yield after being coated on a seed and stored for a defined period of time prior to planting the seed. For example, in some embodiments, inoculant compositions of the present disclosure improve the ability of the microbial spores therein to propagate and increase yield after being coated on a plant propagation material (e.g., seed) and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

Inoculant compositions of the present disclosure may comprise any suitable spores(s), including, but not limited to, the spores of agriculturally beneficial microorganisms such as diazotrophs, phosphate-solubilizing microorganisms and biopesticides.

In some embodiments, inoculant compositions of the present disclosure comprise spores from one or more Gram-negative bacteria.

In some embodiments, inoculant compositions of the present disclosure comprise spores from one or more Gram-positive bacteria.

Non-limiting examples of bacterial spores that may be useful in compositions of the present disclosure include spores of *Bacillus amyloliquefaciens* D747, *Bacillus amyloliquefaciens* NRRL B-50349, *Bacillus amyloliquefaciens* TJ1000, *Bacillus amyloliquefaciens* FZB24, *Bacillus amyloliquefaciens* FZB42, *Bacillus amyloliquefaciens* IN937a, *Bacillus amyloliquefaciens* IT-45, *Bacillus amyloliquefaciens* TJIOOO, *Bacillus amyloliquefaciens* MBI600, *Bacillus amyloliquefaciens* BS27 (deposited as NRRL B-5015), *Bacillus amyloliquefaciens* BS2084 (deposited as NRRL B-50013), *Bacillus amyloliquefaciens* 15AP4 (deposited as ATCC PTA-6507), *Bacillus amyloliquefaciens* 3AP4 (deposited as ATCC PTA-6506), *Bacillus amyloliquefaciens* LSSA01 (deposited as NRRL B-50104), *Bacillus amyloliquefaciens* ABP278 (deposited as NRRL B-50634), *Bacillus amyloliquefaciens* 1013 (deposited as NRRL B-50509), *Bacillus amyloliquefaciens* 918 (deposited as NRRL B-50508), *Bacillus amyloliquefaciens* 22CP1 (deposited as ATCC PTA-6508) and *Bacillus amyloliquefaciens* BS18 (deposited as NRRL B-50633), *Bacillus cereus* 1-1562, *Bacillus firmus* 1-1582, *Bacillus lichenformis* BA842 (deposited as NRRL B-50516), *Bacillus lichenformis* BL21 (deposited as NRRL B-50134), *Bacillus mycoides* NRRL B-21664, *Bacillus pumilus* NRRL B-21662, *Bacillus pumilus* NRRL B-30087, *Bacillus pumilus* ATCC 55608, *Bacillus pumilus* ATCC 55609, *Bacillus*

pumilus GB34, *Bacillus pumilus* KFP9F, *Bacillus pumilus* QST 2808, *Bacillus subtilis* ATCC 55078, *Bacillus subtilis* ATCC 55079, *Bacillus subtilis* MBI 600, *Bacillus subtilis* NRRL B-21661, *Bacillus subtilis* NRRL B-21665, *Bacillus subtilis* CX-9060, *Bacillus subtilis* GB03, *Bacillus subtilis* GB07, *Bacillus subtilis* QST-713, *Bacillus subtilis* FZB24, *Bacillus subtilis* D747, *Bacillus subtilis* 3BP5 (deposited as NRRL B-505 10), *Bacillus thuringiensis* ATCC 13367, *Bacillus thuringiensis* GC-91, *Bacillus thuringiensis* NRRL B-21619, *Bacillus thuringiensis* ABTS-1857, *Bacillus thuringiensis* SAN 401 I, *Bacillus thuringiensis* ABG-6305, *Bacillus thuringiensis* ABG-6346, *Bacillus thuringiensis* AM65-52, *Bacillus thuringiensis* SA-12, *Bacillus thuringiensis* SB4, *Bacillus thuringiensis* ABTS-35 1, *Bacillus thuringiensis* HD-1, *Bacillus thuringiensis* EG 2348, *Bacillus thuringiensis* EG 7826, *Bacillus thuringiensis* EG 7841, *Bacillus thuringiensis* DSM 2803, *Bacillus thuringiensis* NB-125, *Bacillus thuringiensis* NB-176 and combinations thereof, as well as spores of microorganisms having at least at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or more identical to any of the aforementioned strains on the basis of 16S rDNA sequence identity.

In some embodiments, inoculant compositions of the present disclosure comprise one or more fungal spores.

Non-limiting examples of fungal spores that may be useful in compositions of the present disclosure include spores of *Gliocladium virens* ATCC 52045, *Gliocladium virens* GL-21, *Glomus intraradices* RTI-801, *Metarhizium anisopliae* F52, *Penicillium bilaiae* (formerly known as *P. bilaii* and *P. bilaji*) ATCC 18309, *Penicillium bilaiae* ATCC 2085 1, *Penicillium bilaiae* ATCC 22348, *Penicillium bilaiae* NRRL 50162, *Penicillium bilaiae* NRRL 50169, *Penicillium bilaiae* NRRL 50776, *Penicillium bilaiae* NRRL 50777, *Penicillium bilaiae* NRRL 50778, *Penicillium bilaiae* NRRL 50779, *Penicillium bilaiae* NRRL 50780, *Penicillium bilaiae* NRRL 50781, *Penicillium bilaiae* NRRL 50782, *Penicillium bilaiae* NRRL 50783, *Penicillium bilaiae* NRRL 50784, *Penicillium bilaiae* NRRL 50785, *Penicillium bilaiae* NRRL 50786, *Penicillium bilaiae* NRRL 50787, *Penicillium bilaiae* NRRL 50788, *Penicillium bilaiae* RS7B-SD1, *Penicillium brevicompactum* AgRF18, *Penicillium canescens* ATCC 10419, *Penicillium expansum* ATCC 24692, *Penicillium expansum* YT02, *Penicillium fellatanum* ATCC 48694, *Penicillium gastrivorus* NRRL 50170 , *Penicillium glabrum* DAOM 239074, *Penicillium glabrum* CBS 229.28, *Penicillium janthinellum* ATCC 10455, *Penicillium lanosocoeruleum* ATCC 48919, *Penicillium radicum* ATCC 201836, *Penicillium radicum* FRR 4717, *Penicillium radicum* FRR 4719, *Penicillium radicum* N93/47267, *Penicillium raistrickii* ATCC 10490, *Trichoderma asperellum* SKT-1, *Trichoderma asperellum* ICC 012, *Trichoderma atroviride* LC52, *Trichoderma atroviride* CNCM 1-1237, *Trichoderma fertile* JM41R *Trichoderma gamsii* ICC 080, *Trichoderma hamatum* ATCC 52198, *Trichoderma harzianum* ATCC 52445, *Trichoderma harzianum* KRL-AG2, *Trichoderma harzianum* T-22, *Trichoderma harzianum* TH-35, *Trichoderma harzianum* T-39, *Trichoderma harzianum* ICC012, *Trichoderma reesi* ATCC 28217, *Trichoderma virens* ATCC 57678, *Trichoderma virens* GI-3, *Trichoderma virens* GL-21, *Trichoderma virens* G-41, *Trichoderma viridae* ATCC 52440, *Trichoderma viridae* ICC080, *Trichoderma viridae* TVland combinations thereof, as well as spores of microorganisms having at least at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or more identical to any of the aforementioned strains on the basis of internal transcribed spacer (ITS) and/or cytochrome c oxidase (COI) sequence identity.

In some embodiments, inoculant compositions of the present disclosure comprise spores from one or more mycorrhizal fungi (e.g., one or more endomycorrhizal fungi and/or one or more ectomycorrhizal fungi).

Non-limiting examples of mycorrhizal spores that may be useful in compositions of the present disclosure includespores of mycorrhizal strains such as *Gigaspora margarita*, *Glomus aggregatum*, *Glomus brasilianum*, *Glomus clarum*, *Glomus deserticola*, *Glomus etunicatum*, *Glomus intraradices*, *Glomus monosporum*, *Glomus mosseae*, *Laccaria bicolor*, *Laccaria laccata*, *Paraglomus brazilianum*, *Pisolithus tinctorius*, *Rhizopogon*

amylopogon, *Rhizopogon fiilvigleba*, *Rhizopogon luteolus*, *Rhizopogon villosuli*, *Scleroderma cepa* and *Scleroderma citrinum* and combinations thereof.

In some embodiments, inoculant compositions of the present disclosure comprise spores from one or more diazotrophs.

In some embodiments, inoculant compositions of the present disclosure comprise spores from one or more phosphate-solubilizing microorganisms.

In some embodiments, inoculant compositions of the present disclosure comprise spores from one or more biofungicides, bioherbicides, bioinsecticides and/or bionematicides. *See generally* BURGESS, FORMULATION OF MICROBIAL BIOPESTICIDES: BENEFICIAL MICROORGANISMS, NEMATODES AND SEED TREATMENTS (Springer Science & Business Media) (2012); HALL & MENN, BIOPESTICIDES: USE AND DELIVERY (Humana Press) (1998); McCoy, et al., *Entomogenous fungi*, in CRC HANDBOOK OF NATURAL PESTICIDES. MICROBIAL PESTICIDES, PART A. ENTOMOGENOUS PROTOZOA AND FUNGI (C. M. Inoffo, ed.), Vol. 5:151-236 (1988); SAMSON, et al., ATLAS OF ENTOMOPATHOGENIC FUNGI (Springer-Verlag, Berlin) (1988); and deFaria and Wraight, *Mycoinsecticides and Mycoacaricides: A comprehensive list with worldwide coverage and international classification of formulation types*, BIOL. CONTROL (2007), doi: 10.1016/j.biocontrol.2007.08.001.

Microbial spores may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, microbial spores comprise about 0.1 to about 30% (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30% or more (by weight) of one or more microbial spores. In some embodiments, the microbial spore amount/concentration is about 1, 2, 3, 4 or 5 to about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25% (by weight) of the inoculant composition.

In some embodiments, microbial spores are present in an amount ranging from about 1×10^1 to about 1×10^{20} colony-forming units (cfu) per gram and/or milliliter of inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} , 1×10^{13} , 1×10^{14} , 1×10^{15} or more microbial spores per gram and/or milliliter of inoculant composition (e.g., about 1×10^4 to about 1×10^9 *Bacillus amyloliquefaciens* TJIOOO (also known as 1BE, isolate ATCC BAA-390), *Metarhizium anisopliae* F52, *Penicillium bilaiae* ATCC 2085 1, *Penicillium bilaiae* RS7B-SD1 and/or *Trichoderma virens* GI-3 spores per gram/milliliter).

In some embodiments, the amount/concentration of spores is that amount/concentration which is effective to enhance the yield of the plant or plant part to which the inoculant composition is applied.

Microbial spores may be produced by any suitable method(s), including, but not limited to, liquid fermentation and solid state fermentation. *See, e.g.*, Cunningham et al., CAN. J. BOT. 68:2270 (1990); Friesen et al., APPL. MICROBIOL. BIOTECH. 68:397 (2005).

Although the primary foci of the present disclosure are compositions and methods for enhancing the stability and/or survival of microbial spores, it is to be understood that inoculant compositions of the present disclosure may comprise one or more microorganisms (e.g., diazotrophs, phosphate-solubilizing microorganisms and/or biopesticides) in a vegetative state. *See* International Patent Application No. PCT/US20 16/050529, the disclosure of each of which is incorporated herein by reference in its entirety.

Vegetative cells may be incorporated into inoeulant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, inoeulant compositions of the present disclosure are devoid of vegetative cells.

In some embodiments, vegetative cells are present in an amount ranging from about 1×10^1 to about 1×10^{20} colony-forming units (cfu) per gram and/or milliliter of inoeulant composition. For example, inoeulant compositions of the present disclosure may comprise about 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} , 1×10^{13} , 1×10^{14} , 1×10^{15} or more cfu of agriculturally beneficial microorganisms per gram and/or milliliter of inoeulant composition.

In some embodiments, the amount/concentration of vegetative cells is an amount effective to enhance the yield of the plant or plant part to which the inoeulant composition is applied.

In some embodiments, the amount/concentration of vegetative cells is 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more cfu per gram/milliliter of inoeulant composition.

Inoeulant compositions of the present disclosure may comprise any suitable non-aqueous carrier(s), including, but not limited to, foliar-compatible carriers, seed-compatible carriers and soil-compatible carriers. Selection of appropriate carrier materials will depend on the intended application[^]) and the microorganism(s) present in the inoeulant composition.

In some embodiments, the carrier material(s) are selected to provide an inoeulant composition in the form of a powder or granule. For example, carrier materials may be selected to provide inoeulant compositions that are formulated as freeze-dried powders, spray-dried powders, spray-freeze-dried powders and combinations thereof.

Non-limiting examples of solid carriers that may be useful in compositions of the present disclosure include disaccharides, maltodextrins, monosaccharides, oligosaccharides, peat-based powders and granules, and agriculturally acceptable polymers.

Additional examples of solid carriers that may be included in inoeulant compositions of the present disclosure may be found in **BURGES, FORMULATION OF MICROBIAL BIOPESTICIDES: BENEFICIAL MICROORGANISMS, NEMATODES AND SEED TREATMENTS** (Springer Science & Business Media) (2012).

Carriers incorporated into inoeulant compositions of the present disclosure may comprise a growth medium suitable for culturing one or more of the microorganisms in the inoeulant composition. For example, in some embodiments, inoeulant compositions of the present disclosure comprise Czapek-Dox medium, glycerol yeast extract, mannitol yeast extract, potato dextrose broth and/or YEM media.

The non-aqueous carrier may constitute any suitable portion of the inoeulant composition. In some embodiments, the non-aqueous carrier(s) comprise(s) about 1 to about 99 % (by weight) of the inoeulant composition. For example, inoeulant compositions of the present disclosure, the non-aqueous carrier(s) constitute(s) about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, 99.5% or more (by weight) of the inoeulant composition. In some embodiments, the carrier amount/concentration is about 50 to about 99%, about 55% to about 95%, about 60% to about 95%, about 65% to about 90%, about 70 to about 90% , about 75 to about 90%, about 80 to about 90% or about 80 to about 85% (by weight) of the inoeulant composition. In some embodiments, inoeulant compositions of the present disclosure comprise one or more commercial carriers used in accordance with the manufacturer's recommended amounts/concentrations.

In some embodiments, the non-aqueous carrier comprises, consists essentially of, or consists of one or more maltodextrins.

Inoeulant compositions of the present disclosure may comprise any suitable maltodextrins, including, but not limited to, maltodextrins having a dextrose equivalent value (DEV) of about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15,

16, 17, 18, 19, 20, 21, 22, 23, 24, or 25. In some embodiments, inoculant compositions of the present disclosure comprise one or more maltodextrins having a DEV of about 5 to about 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, about 10 to about 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or about 15 to about 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25. In some embodiments, the non-aqueous carrier comprises a combination of maltodextrins having a DEV of about 5 to about 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, about 10 to about 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or about 15 to about 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25. In some embodiments, the non-aqueous carrier comprises one or more maltodextrins having a DEV of about 10 to about 25 (e.g., one or more maltodextrins having a DEV of about 15 to about 20). In some embodiments, the non-aqueous carrier comprises a combination of maltodextrins having a DEV of about 10 to about 25 (e.g., a combination of maltodextrins having a DEV of about 15 to about 20).

Non-limiting examples of maltodextrins that may be useful in compositions of the present disclosure include MALTRIN® M150 (DEV = 15; molecular weight = 1200; Grain Processing Corporation, Muscatine, IA), MALTRIN® M180 (DEV = 18; molecular weight = 1050; Grain Processing Corporation, Muscatine, IA), MALTRIN® M200 (DEV = 20; molecular weight = 900; Grain Processing Corporation, Muscatine, IA), MALTRIN® M250 (DEV = 25; molecular weight = 720; Grain Processing Corporation, Muscatine, IA); MALTRIN QD® M580 (DEV = 16.5-19.9; Grain Processing Corporation, Muscatine, IA); MALTRIN QD® M585 (DEV = 15.0-19.9; Grain Processing Corporation, Muscatine, IA); MALTRIN QD® M600 (DEV = 20.0-23.0; Grain Processing Corporation, Muscatine, IA); GLOBE® Plus 15 DE (Ingredion Inc., Westchester, IL); and combinations thereof. In some embodiments, the maltodextrin (or combination of maltodextrins) has a DEV of 15 to 20.

Maltodextrins may be incorporated into inoculant compositions of the present disclosure in any suitable form. In some embodiments, the maltodextrin(s) included in inoculant compositions of the present disclosure is/are at least 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 99.5% or more pure.

Maltodextrins may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the maltodextrin(s) comprise about 5 to about 99 % or more (by weight) of the inoculant composition. For example, in some embodiments, the non-aqueous carrier comprises about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99% or more of one or more maltodextrins (each and/or collectively) having a DEV value of about 15 to about 20. In some embodiments, the maltodextrin amount/concentration is about 50 to about 95%, about 55% to about 90%, about 60% to about 85%, about 65% to about 80%, or about 70 to about 80% (by weight) of the non-aqueous carrier.

In some embodiments, the non-aqueous carrier comprises, consists essentially of, or consists of one or more monosaccharides, disaccharides and/or oligosaccharides.

Inoculant compositions of the present disclosure may comprise any suitable monosaccharide, including, but limited to, allose, altrose, arabinose, fructose, galactose, glucose, gulose, iodose, lyxose, mannose, ribose, talose, threose and xylose. In some embodiments, the non-aqueous carrier comprises glucose. In some embodiments, the non-aqueous carrier does not comprise glucose.

Inoculant compositions of the present disclosure may comprise any suitable disaccharide, including, but limited to, cellobiose, chitobiose, gentiobiose, gentiobiulose, isomaltose, kojibiose, lactose, lactulose, laminaribiose, maltose (e.g., maltose monohydrate, anhydrous maltose), maltulose, mannobiose, melibiose, melibiulose, nigerose, palatinose, rutinose, rutinulose, sophorose, sucrose, trehalose, turanose and xylobiose. In some embodiments, the non-aqueous carrier comprises maltose. In some embodiments, the non-aqueous carrier comprises sucrose. In some embodiments, the non-aqueous carrier comprises trehalose. In some embodiments, the non-aqueous carrier does not comprise trehalose.

Inoculant compositions of the present disclosure may comprise any suitable oligosaccharide, including, but limited to, fructo-oligosaccharides, galacto-oligosaccharides, mannon-oligosaccharides and raffinose.

Mono-, di- and/or oligosaccharides may be incorporated into inoculant compositions of the present disclosure in any suitable form. In some embodiments, the mono-, di- and/or oligosaccharide(s) included in inoculant compositions of the present disclosure is/are at least 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 99.5% or more pure.

Mono-, di- and oligosaccharides may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the mono-, di- and/or oligosaccharide(s) comprise(s) about 5 to about 95% (by weight) of the inoculant composition. For example, in some embodiments, the non-aqueous carrier comprises about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more (by weight) of one or more mono-, di- and/or oligosaccharides (e.g., maltose and/or trehalose). In some embodiments, the mono-, di- and/or oligosaccharide (e.g., maltose) amount/concentration is about 1 to about 65%, about 5% to about 20%, about 10% to about 25%, about 20% to about 50%, or about 30 to about 60% (by weight) of the non-aqueous carrier.

In some embodiments, the non-aqueous carrier comprises, consists essentially of, or consists of one or more malt extracts.

Inoculant compositions of the present disclosure may comprise any suitable malt extract(s).

Malt extracts may be incorporated into inoculant compositions of the present disclosure in any suitable form. In some embodiments, the malt extract(s) included in inoculant compositions of the present disclosure is/are at least 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 99.5% or more pure.

Malt extracts may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the malt extract(s) comprise(s) about 5 to about 95 % (by weight) of the inoculant composition. For example, in some embodiments, the non-aqueous carrier comprises about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90% or more (by weight) of one or more malt extracts. In some embodiments, the malt extract amount/concentration is about 50 to about 99%, about 55% to about 95%, about 60% to about 90%, about 65% to about 90%, or about 70 to about 90% (by weight) of the non-aqueous carrier.

In some embodiments, the non-aqueous carrier comprises, consists essentially of, or consists of one or more peat-based powders and/or granuales.

Inoculant compositions of the present disclosure may comprise any suitable peat-based powder(s) and/or granuale(s).

Peat-based powders and/or granuales extracts may be incorporated into inoculant compositions of the present disclosure in any suitable form. In some embodiments, the peat-based powder(s) and/or granuale(s) included in inoculant compositions of the present disclosure is/are at least 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 99.5% or more pure.

Peat-based powders and/or granuales may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the peat-based powder(s) and/or granuale(s) comprise(s) about 5 to about 95 % (by weight) of the inoculant composition. For example, in some embodiments, the non-aqueous carrier comprises about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more (by weight) of one or more peat-based powders and/or granuales. In some embodiments, the peat extract amount/concentration is about 50 to about 99%, about 55% to about 95%, about 60% to about 90%, about 65% to about 90%, or about 70 to about 90% (by weight) of the non-aqueous carrier.

In some embodiments, the non-aqueous carrier comprises, consists essentially of, or consists of one or more agriculturally acceptable polymers.

Inoculant compositions of the present disclosure may comprise any suitable agriculturally acceptable polymer(s), including, but not limited to, biodegradable polymers and synthetic polymers. For example, in some embodiments, inoculant compositions of the present disclosure comprise agar, alginate, carrageenan, cellulose, guar gum, locust bean gum, methylcellulose, pectin, polycaprolactone, polylactide, polyvinyl alcohol, polyvinyl pyrrolidone, sodium carboxymethyl cellulose, starch and xanthan gum.

Non-limiting examples of polymers that may be useful in compositions of the present disclosure include TICAXAN® xanthan powders, such as PRE-HYDRATED® TICAXAN® Rapid-3 Powder (TIC Gums, White Marsh, MD) and combinations thereof.

Additional examples of polymers that may be included in inoculant compositions of the present disclosure may be found in Pouci, et al. AM. J. AGRIC. BIOL. SCI. 3(1):299 (2008).

Agriculturally acceptable polymers may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the agriculturally acceptable polymer(s) comprise(s) about 5 to about 95 % (by weight) of the inoculant composition. For example, in some embodiments, the non-aqueous carrier comprises about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more (by weight) of one or more agriculturally acceptable polymers. In some embodiments, the agriculturally acceptable polymers amount/concentration is about 50 to about 99%, about 55% to about 95%, about 60% to about 90%, about 65% to about 90%, or about 70 to about 90% (by weight) of the non-aqueous carrier.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial polymers used in accordance with the manufacturer's recommended amounts/concentrations.

It is to be understood that monosaccharides, disaccharides, oligosaccharides, maltodextrins, malt extracts, agriculturally acceptable polymers and peat-based powders/granules may be combined to form dry non-aqueous carriers having beneficial properties, including, but not limited to, enhanced stabilization of microbial spores. Thus, in some embodiments, inoculant compositions of the present disclosure comprise a dry non-aqueous carrier that comprises one or more maltodextrins in combination with one or more mono-, di- and/or oligosaccharides, one or more sugar alcohols, one or more malt extracts, one or more agriculturally acceptable polymers and/or one or more peat-based powders and/or granules.

Maltodextrins and mono-, di- and/or oligosaccharides may be incorporated into inoculant compositions of the present disclosure in any suitable ratio(s). In some embodiments, the non-aqueous carrier has a maltodextrin:(mono-, di- and/or oligosaccharide) (e.g., maltodextrin:maltose) ratio of about 1:99 to about 99:1 (by weight, based upon the respective weight percentages of the maltodextrin(s) and mono-, di- and/or oligosaccharide(s) in the non-aqueous carrier). For example, in some embodiments, the non-aqueous carrier has a maltodextrin:(mono-, di- and/or oligosaccharide) (e.g., maltodextrin:maltose) ratio of about 5:95, 10:90, 15:85, 20:80, 25:75, 30:70, 35:65, 40:60, 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5 or more. In some embodiments, the maltodextrin:(mono-, di- and/or oligosaccharide) (e.g., maltodextrin:maltose) ratio is about 45:55 to about 95:5. Thus, in some embodiments, inoculant compositions of the present disclosure comprise one or more maltodextrins having a DEV of about 15 to about 20 and one or more mono-, di- and/or oligosaccharides (e.g., maltose) in a ratio of about 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5.

Maltodextrins and malt extracts may be incorporated into inoculant compositions of the present disclosure in any suitable ratio(s). In some embodiments, the non-aqueous carrier has a maltodextrin:malt extract ratio of about 1:99 to about 99:1 (by weight, based upon the respective weight percentages of the maltodextrin(s) and malt extract(s) in the

non-aqueous carrier). For example, in some embodiments, the non-aqueous carrier has a maltodextrin:malt extract ratio of about 5:95, 10:90, 15:85, 20:80, 25:75, 30:70, 35:65, 40:60, 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5 or more. In some embodiments, the maltodextrin:malt extract ratio is about 45:55 to about 95:5. Thus, in some embodiments, inoculant compositions of the present disclosure comprise one or more maltodextrins having a DEV of about 15 to about 20 and one or more malt extracts in a ratio of about 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5.

Maltodextrins and peat-based powders and/or granules may be incorporated into inoculant compositions of the present disclosure in any suitable ratio(s). In some embodiments, the non-aqueous carrier has a maltodextrin:peat powder/granule ratio of about 1:99 to about 99:1 (by weight, based upon the respective weight percentages of the maltodextrin(s) and peat powder(s)/granule(s) in the non-aqueous carrier). For example, in some embodiments, the non-aqueous carrier has a maltodextrin:peat powder/granule ratio of about 5:95, 10:90, 15:85, 20:80, 25:75, 30:70, 35:65, 40:60, 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5 or more. In some embodiments, the maltodextrin:peat powder/granule ratio is about 45:55 to about 95:5. Thus, in some embodiments, inoculant compositions of the present disclosure comprise one or more maltodextrins having a DEV of about 15 to about 20 and one or more peat-based powders and/or granules in a ratio of about 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5.

Non-aqueous carriers may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, the non-aqueous carrier(s) comprise(s) about 5 to about 99.9% (by weight) of the inoculant composition. For example, in some embodiments, the non-aqueous carrier(s) constitute(s) about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, 99.5% or more (by weight) of the inoculant composition. In some embodiments, the non-aqueous carrier amount/concentration is about 50 to about 99%, about 55% to about 95%, about 60% to about 95%, about 65% to about 90%, about 70 to about 90%, about 75 to about 90%, about 80 to about 90% or about 80 to about 85% (by weight) of the inoculant composition.

In some embodiments, the amount/concentration of the non-aqueous carrier(s) in the inoculant composition is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores/vegetative cells in the inoculant composition survive following desiccation (of about 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more) and storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the amount/concentration of the non-aqueous carrier(s) in the inoculant composition is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores/vegetative cells in the inoculant composition survive following storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial carriers used in accordance with the manufacturer's recommended amounts/concentrations.

In the interest of maximizing the teaching of the present application and without intending to be limited by any particular theory, applicants submit that the aforementioned carriers may enhance the stability and survival of microbial spores in an inoculant composition by encapsulating the microbial spores, thereby reducing their exposure

to elements that would otherwise cause the spores to germinate or destabilize. In such instances, the non-aqueous carrier may be referred to as a protectant.

In some embodiments, the non-aqueous carrier at least partially encapsulate 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99% or more of the microbial spores in the inoculant composition. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more monosaccharides, disaccharides, oligosaccharides, maltodextrins and/or malt extracts that fully or partially encapsulate at least 50, 55, 60, 65, 70, 75, 80, 85, 90 or 95% of the microbial spores in the inoculant composition.

Inoculant compositions of the present disclosure may comprise any suitable agriculturally acceptable dispersant(s), including, but not limited to, surfactants and wetting agents.

In some embodiments, inoculant compositions of the present disclosure comprise one or more anionic surfactants. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more water-soluble anionic surfactants and/or one or more water-insoluble anionic surfactants, optionally one or more anionic surfactants selected from the group consisting of alkyl carboxylates (e.g., sodium stearate), alkyl sulfates (e.g., alkyl lauryl sulfate, sodium lauryl sulfate), alkyl ether sulfates, alkyl amido ether sulfates, alkyl aryl polyether sulfates, alkyl aryl sulfates, alkyl aryl sulfonates, alkyl sulfonates, alkyl amide sulfonates, alkyl aryl sulfonates, alkyl benzene sulfonates, alkyl diphenyloxide sulfonate, alpha-olefin sulfonates, alkyl naphthalene sulfonates, paraffin sulfonates, alkyl sulfosuccinates, alkyl ether sulfosuccinates, alkylamide sulfosuccinates, alkyl sulfosuccinamates, alkyl sulfoacetates, alkyl phosphates, alkyl ether phosphates, acyl sarconsinates, acyl isethionates, N-acyl taurates, N-acyl-N-alkyltaurates, benzene sulfonates, cumene sulfonates, dioctyl sodium sulfosuccinate, ethoxylated sulfosuccinates, lignin sulfonates, linear alkylbenzene sulfonates, monoglyceride sulfates, perfluorobutanesulfonate, perfluorooctanesulfonate, phosphate ester, styrene acrylic polymers, toluene sulfonates and xylene sulfonates.

In some embodiments, inoculant compositions of the present disclosure comprise one or more cationic surfactants. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more pH-dependent amines and/or one or more quaternary ammonium cations, optionally one or more cationic surfactants selected from the group consisting of alkyltrimethylammonium salts (e.g., cetyl trimethylammonium bromide, cetyl trimethylammonium chloride), cetylpyridinium chloride, benzalkonium chloride, benzethonium chloride, 5-Bromo-5-nitro-1,3-dioxane, dimethyldioctadecylammonium chloride, cetrimonium bromide, dioctadecyldimethylammonium bromide and/or octenidine dihydrochloride.

In some embodiments, inoculant compositions of the present disclosure comprise one or more nonionic surfactants. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more water-soluble nonionic surfactants and/or one or more water-insoluble nonionic surfactants, optionally one or more nonionic surfactants selected from the group consisting of alcohol ethoxylates (e.g., TERGITOL™ 15-S surfactants, such as TERGITOL™15-S-9 (The Dow Chemical Company, Midland, MI)), alkanolamides, alkanolamine condensates, carboxylic acid esters, cetostearyl alcohol, cetyl alcohol, cocamide DEA, dodecyldimethylamine oxides, ethanolamides, ethoxylates of glycerol ester and glycol esters, ethylene oxide polymers, ethylene oxide-propylene oxide copolymers, glucoside alkyl ethers, glycerol alkyl ethers (e.g.,), glycerol esters, glycol alkyl ethers (e.g., polyoxyethylene glycol alkyl ethers, polyoxypropylene glycol alkyl ethers, glycol alkylphenol ethers (e.g., polyoxyethylene glycol alkylphenol ethers, glycol esters, monolaurin, pentaethylene glycol monododecyl ethers, poloxamer, polyamines, polyglycerol polyricinoleate, polysorbate, polyoxyethylenated fatty acids, polyoxyethylenated mercaptans, polyoxyethylenated polyoxypropylene glycols, polyoxyethylene glycol sorbitan alkyl esters, polyethylene glycol-polypropylene glycol copolymers, polyoxyethylene glycol octylphenol

ethers, polyvinyl pyrrolidones, sugar-based alkyl polyglycosides, sulfoanilamides, sorbitan fatty acid alcohol ethoxylates, sorbitan fatty acid ester ethoxylates, sorbitan fatty acid ester and/or tertiary acetylenic glycols.

In some embodiments, inoculant compositions of the present disclosure comprise at least one nonionic surfactant. In some embodiments, inoculant compositions of the present disclosure comprise at least one water insoluble nonionic surfactant and at least one water soluble nonionic surfactant. In some embodiments, inoculant compositions of the present disclosure comprise a combination of nonionic surfactants having hydrocarbon chains of substantially the same length.

In some embodiments, inoculant compositions of the present disclosure comprise one or more zwitterionic surfactants. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more betaines and/or one or more sultaines, optionally one or more zwitterionic surfactants selected from the group consisting of 3-[(3-cholamidopropyl)dimethylammonio]-l-propanesulfonate, cocamidopropyl betaine, cocamidopropyl hydroxysultaine, phosphatidylserine, phosphatidylethanolamine, phosphatidylcholine and/or one or more sphingomyelins.

In some embodiments, inoculant compositions of the present disclosure comprise one or more soaps and/or organosilicone surfactants. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more alkali metal salts of fatty acids.

In some embodiments, inoculant compositions of the present disclosure comprise one or more wetting agents. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more naphthalene sulfonates, optionally one or more alkyl naphthalene sulfonates (e.g., sodium alkyl naphthalene sulfonate), one or more isopropyl naphthalene sulfonates (e.g., sodium isopropyl naphthalene sulfonate) and/or one or more butyl naphthalene sulfonates (e.g., sodium n-butyl naphthalene sulfonate).

Selection of appropriate surfactants will depend on the intended application(s) and the microorganism(s) present in the inoculant composition. In general, the surfactant(s) will have low toxicity for the microorganism(s) in the inoculant composition and for the plant part(s) to which the inoculant composition is to be applied. In some embodiments, the surfactant(s) will be selected to wet and/or emulsify one or more soils.

Non-limiting examples of dispersants that may be useful in compositions of the present disclosure include Atlox™ (e.g., 4916, 4991; Croda International PLC, Edison, NJ), BIO-SOFT® (e.g., N series, such as NI-3, NI-7, NI-5, NI-9, N23-3, N2.3-6.5, N25-3, N25-7, N25-9, N91-2.5, N91-6, N91-8; Stepan Company, Northfield, IL), MAKON® nonionic surfactants (e.g., DA-4, DA-6 and DA-9; Stepan Company, Northfield, IL), MORWET® powders (Akzo Nobel Surface Chemistry LLC, Chicago, IL), MULTIWET™ surfactants (e.g., MO-70R, MO-85P, MO-85P-PW-(AP); Croda International PLC, Edison, NJ), SILWET® L-77 (Helena Chemical Company, Collierville, TN), SPAN™ surfactants (e.g., 20, 40, 60, 65, 80 and 85; Croda Inc., Edison NJ), TAMOL™ dispersants (The Dow Chemical Company, Midland, MI), TERGITOL™ surfactants (e.g., 15-S-9, TMN-6, TMN-100X and XD; The Dow Chemical Company, Midland, MI), TERSPERSE surfactants (e.g., 2001, 2020, 2100, 2105, 2158, 2700, 4894 and 4896; Hunstman Corp., The Woodlands, TX), TRITON™ surfactants (e.g., X-100; The Dow Chemical Company, Midland, MI), TWEEN® surfactants (e.g., TWEEN® 20, 21, 22, 23, 28, 40, 60, 61, 65, 80, 81 and 85; Croda International PLC, Edison, NJ) and combinations thereof.

Dispersants may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the dispersant(s) comprise(s) about 0.1 to about 25 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 15, 20, 25% or more (by weight) of one or more dispersants. In some embodiments, the dispersant (s) comprise(s) about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6,

0.7, 0.8, 0.9 or 1 to about 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 % (by weight) of the inoculant composition.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial wetting agents and/or one or more surfactants used in accordance with the manufacturer's recommended amounts/concentrations.

Additional examples of dispersants that may be included in inoculant compositions of the present disclosure may be found in BAIRD & ZUBLENA. 1993. SOIL FACTS: USING WETTING AGENTS (NONIONIC SURFACTANTS) ON SOIL (North Carolina Cooperative Extension Service Publication AG-439-25) (1993); BURGESS, FORMULATION OF MICROBIAL BIOPESTICIDES: BENEFICIAL MICROORGANISMS, NEMATODES AND SEED TREATMENTS (Springer Science & Business Media) (2012); MCCARTY, WETTING AGENTS (Clemson University Cooperative Extension Service Publication) (2001).

Inoculant compositions of the present disclosure may comprise any suitable excipient(s), including, but not limited to, anti-freezing agents, drying agents, safeners and pH buffers.

Inoculant compositions of the present disclosure may comprise any suitable agriculturally acceptable anti-freezing agent(s), including, but not limited to, ethylene glycol, glycerin, propylene glycol and urea.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial anti-freezing agents used in accordance with the manufacturer's recommended amounts/concentrations.

Inoculant compositions of the present disclosure may comprise any suitable agriculturally acceptable drying agent(s), including, but not limited to, drying powders. For example, in some embodiments, inoculant compositions of the present disclosure comprise calcium stearate, clay (e.g., attapulgite clay, montmorillonite clay), graphite, magnesium stearate, magnesium sulfate, powdered milk, silica (e.g., fumed silica, hydrophobically-coated silica, precipitated silica), soy lecithin and/or talc.

Non-limiting examples of drying agents that may be useful in compositions of the present disclosure include AEROSIL® hydrophobic fumed silica powders (Evonik Corporation, Parsippany, NJ), BENTOLITE® powders (BYK-Chemie GmbH, Wesel, Germany), SIPERNAT® silica powders (Evonik Corporation, Parsippany, NJ) and combinations thereof.

Additional examples of drying agents that may be included in inoculant compositions of the present disclosure may be found in BURGESS, FORMULATION OF MICROBIAL BIOPESTICIDES: BENEFICIAL MICROORGANISMS, NEMATODES AND SEED TREATMENTS (Springer Science & Business Media) (2012).

Drying agents may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the inoculant compositions of the present disclosure comprise about 0.5 to about 10 grams of drying powder per liter of inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.5, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10 grams or more of drying powder per liter of inoculant composition. In some embodiments, the amount/concentration of drying agent(s) comprise(s) calcium stearate, attapulgite clay, montmorillonite clay, graphite, magnesium stearate, silica (e.g., fumed silica, hydrophobically-coated silica and/or precipitated silica) and/or talc.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial drying agents used in accordance with the manufacturer's recommended amounts/concentrations.

Inoculant compositions of the present disclosure may comprise any suitable agriculturally acceptable dust suppressant(s), including, but not limited to, adhesives, glycerin, mineral oils, paraffinic oils, vegetable oils and synthetic polymers. It is to be understood that some compounds may act as both an adhesive and a dust suppressant.

Indeed, the dust suppressant activity of many compounds arises from their ability to adhere dust particles to other heavier particles. For example, in some embodiments of the present disclosure, one or more oils is used to adhere microbial spores in the inoculant composition to larger particles comprising one or more maltodextrins.

Non-limiting examples of dust suppressants that may be useful in compositions of the present disclosure include ARENAPRO, BIORAIN and ROADKILL (Dustkill LLC, Columbus, IN), BIO-SOFT® surfactants (e.g., N series, such as NI-3, NI-7, NI-5, NI-9, N23-3, N2.3-6.5, N25-3, N25-7, N25-9, N91-2.5, N91-6, N91-8; Stepan Company, Northfield, IL), DURASOIL (Soilworks, LLC, Scottsdale, AZ), DUSGON (Dupont (Australia) Ltd, Macquarie Park, Australia), DUSTRX (Seaco Technologies, Inc., Bakersfield, CA), SUN AG® oils (HollyFrontier Refining & Marketing, LLC, Plymouth Meeting, PA), SUNSPRAY® oils (e.g., 6N, 6E, 8N, UN; HollyFrontier Refining & Marketing, LLC, Plymouth Meeting, PA), SUNSPRAY® ULTRA-FINE® oils (HollyFrontier Refining & Marketing, LLC, Plymouth Meeting, PA), TOMADOL® surfactants (e.g., 23-1, 23-3, 23-5, 23-6.5; Air Products and Chemicals, Inc., Allentown, PA) and combinations thereof.

Dust suppressants may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, dust suppressants comprise about 0.1 to about 30% (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 6, 7, 8, 9, 10% or more (by weight) of one or more dust suppressants. In some embodiments, the dust suppressant amount/concentration is about 1 to about 2, 3, 4, 5, 6, 7, 8, 9 or 10%, about 1.5 to about 2, 3, 4, 5, 6, 7, 8, 9 or 10%, about 2 to about 3, 4, 5, 6, 7, 8, 9 or 10%, about 2.5 to about 3, 4, 5, 6, 7, 8, 9 or 10% or about 3 to about 4, 5, 6, 7, 8, 9 or 10% (by weight) of the inoculant composition.

It is to be understood that some compounds may act as both a dispersant and a dust suppressant. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more surfactants, such as BIO-SOFT® N23-3, that disperse microorganisms within the inoculant composition and suppress dusting.

As noted above, the non-aqueous carrier in inoculant compositions of the present invention may act as a protectant. It is to be understood that inoculant compositions of the present disclosure may comprise additional protectants.

Protectants may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, the protectant(s) comprise(s) about 0.1 to about 25% (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 15, 20, 25% or more (by weight) of one or more protectants. In some embodiments, the protectant amount/concentration is about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1 to about 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10% (by weight) of the inoculant composition.

In some embodiments, the protectant amount/concentration is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores/vegetative cells in the inoculant composition survive following storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the protectant amount/concentration is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores/vegetative cells

in the inoculant composition survive following desiccation (of about 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more) and storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the protectant amount/concentration is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores/vegetative cells in the inoculant composition survive following cryopreservation at or below -80°C for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure comprise one or more sugar alcohols (e.g., sorbitol).

Sugar alcohols may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the sugar alcohol(s) comprise(s) about 0.005 to about 10 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.75, 1, 1.25, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5% or more (by weight) of one or more sugar alcohols (e.g., sorbitol).

In some embodiments, inoculant compositions of the present disclosure comprise one or more peat extracts.

Peat extracts may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the peat extract(s) comprise(s) about 0.1 to about 25 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 15, 20, 25% or more (by weight) of one or more peat extracts. In some embodiments, the peat extract amount/concentration is about 0.5 to about 10% (by weight) of the inoculant composition.

In some embodiments, inoculant compositions of the present disclosure comprise one or more peptones.

Peptones may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the peptone(s) comprise(s) about 0.1 to about 25 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 15, 20, 25% or more (by weight) of one or more peptones. In some embodiments, the peptone amount/concentration is about 0.5 to about 10% (by weight) of the inoculant composition.

In some embodiments, inoculant compositions of the present disclosure comprise one or more skim milk extracts.

Skim milk extracts may be incorporated into inoculant compositions of the present disclosure in any suitable amount/concentration. In some embodiments, skim milk extracts comprise about 0.1 to about 25 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 15, 20, 25% or more (by weight) of skim milk extracts. In some embodiments, the skim milk extract amount/concentration is about 0.5 to about 10% (by weight) of the inoculant composition.

In some embodiments, inoculant compositions of the present disclosure comprise one or more hygroscopic polymers. For example, inoculant compositions of the present disclosure may comprise one or more albumins, alginates, celluloses, gums (e.g., cellulose gum, guar gum, gum arabic, gum combretum, xanthan gum), methyl celluloses, nylons, pectins, polyacrylic acids, polycarbonates, polyethylene glycols (PEG), polyethylenimines (PEI),

polylactides, polymethylacrylates (PMA), polyurethanes, polyvinyl alcohols (PVA), polyvinylpyrrolidones (PVP), propylene glycols, sodium carboxymethyl celluloses and/or starches.

Non-limiting examples of hygroscopic polymers that may be useful in compositions of the present disclosure include AGRIMER™ polymers (e.g., 30, AL-10 LC, AL-22, AT/ATF, VA 3E, VA 3I, VA 5E, VA 5I, VA 6, VA 6E, VA 7E, VA 7I, VEMA AN-216, VEMA AN-990, VEMA AN-1200, VEMA AN-1980, VEMA H-815MS; Ashland Specialty Ingredients, Wilmington, DE), EASYSperse™ polymers (Ashland Specialty Ingredients, Wilmington, DE); DISCO™ AG polymers (e.g., L-250, L-280, L-285, L-286, L-320, L-323, L-5 17, L-5 19, L-520, L800; Incotec Inc., Salinas, CA), KELZAN® polymers (Bri-Chem Supply Ltd., Calgary, Alberta, CA), SEEDWORX™ polymers (e.g., Bio 200; Aginnovation, LLC, Walnut Grove, CA), TABULOSE® gels (e.g., SC-580, SC-612, SC-613, SC-681; Blanver Farmquimica, Boca Raton, FL), TICAXAN® xanthan powders (TIC Gums, White Marsh, MD) and combinations thereof.

Additional examples of hygroscopic polymers that may be included in inoculant compositions of the present disclosure may be found in Pouci, et al. *AM. J. AGRIC. BIOL. SCI.* 3(1):299 (2008).

Hygroscopic polymers may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the hygroscopic polymer(s) comprise(s) about 0.1 to about 25 % (by weight) of the inoculant composition. In some embodiments, the hygroscopic polymer(s) comprise(s) about 0.5 to about 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 15, 20, 25% or more (by weight) of one or more hygroscopic polymers. In some embodiments, the hygroscopic polymer amount/concentration is about 0.5 to about 10% (by weight) of the inoculant composition. In some embodiments, the hygroscopic polymer amount/concentration is about 0.5 to about 5% (by weight) of the inoculant composition.

In some embodiments, the hygroscopic polymer amount/concentration is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the hygroscopic polymer amount/concentration is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following desiccation (of about 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more) and storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the hygroscopic polymer amount/concentration is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following cryopreservation at or below -80°C for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial hygroscopic polymers used in accordance with the manufacturer's recommended amounts/concentrations.

In some embodiments, inoculant compositions of the present disclosure comprise one or more oxidation control components.

Inoculant compositions of the present disclosure may comprise any suitable oxidation control component(s), including, but not limited to, antioxidants and/or oxygen scavengers. In some embodiments, the oxidation control component is/comprises ascorbic acid and/or glutathione.

In some embodiments, inoculant compositions comprise one or more antioxidants. For example, in some embodiments, inoculant compositions of the present disclosure comprise ascorbic acid, ascorbyl palmitate, ascorbyl stearate, calcium ascorbate, carotenoids, lipoic acid, phenolic compounds (e.g., flavonoids, flavones, flavonols), potassium ascorbate, sodium ascorbate, thiols (e.g., glutathione, lipoic acid, N-acetyl cysteine), tocopherols, tocotrienols, ubiquinone and/or uric acid.

Non-limiting examples of antioxidants that may be useful in compositions of the present disclosure include those that are soluble in the cell membrane (e.g., alpha tocopherol (vitamin E), ascorbyl palmitate), those that are soluble in alcohols (e.g., IRGANOX® antioxidants (BASF Schweiz AG, Basel, Switzerland)) and those that are soluble in water (e.g., ascorbic acid and isomers or ascorbic acid, sodium or potassium salts of ascorbic acid or isomers or ascorbic acid, glutathione, sodium or potassium salts of glutathione). In some embodiments, use of a membrane-soluble antioxidant necessitates the addition of one or more surfactants to adequately disperse the antioxidant within the inoculant composition.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial antioxidants used in accordance with the manufacturer's recommended amounts/concentrations.

In some embodiments, inoculant compositions comprise one or more naturally occurring or synthetic oxygen scavengers. For example, in some embodiments, inoculant compositions of the present disclosure comprise ascorbic acid, ascorbate salts, catechol and/or sodium hydrogen carbonate.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial oxygen scavengers used in accordance with the manufacturer's recommended amounts/concentrations.

Oxidation control components may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, the oxidation control component(s) comprise(s) about 0.0001 to about 5 % or more (by weight) of the composition. For example, inoculant compositions of the present disclosure may comprise about 0.0005, 0.001, 0.002, 0.003, 0.004, 0.005, 0.0075, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.75, 1, 1.25, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5% or more of one or more oxidation control components. In some embodiments, the amount/concentration of oxidation control components is about 0.005 to about 2% (by weight) of the composition. In some embodiments, the oxidation control component(s) is/are present in a concentration ranging from about 1×10^{-20} M to about 1×10^{-1} M. For example, one or more oxidation control components may be added at a concentration of 1×10^{-20} M, 1×10^{-19} M, 1×10^{-18} M, 1×10^{-17} M, 1×10^{-16} M, 1×10^{-15} M, 1×10^{-14} M, 1×10^{-13} M, 1×10^{-12} M, 1×10^{-11} M, 1×10^{-10} M, 1×10^{-9} M, 1×10^{-8} M, 1×10^{-7} M, 1×10^{-6} M, 1×10^{-5} M, 1×10^{-4} M, 1×10^{-3} M, 1×10^{-2} M, 1×10^{-1} M or more.

In some embodiments, the amount/concentration of oxidation control components is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the amount/concentration of oxidation control components is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following desiccation (of about 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50,

55, 60, 65, 70, 75, 80, 85, 90, 95% or more) and storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the amount/concentration of oxidation control components is effective to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following cryopreservation at or below -80°C for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

Inoculant compositions of the present disclosure may comprise any suitable safener(s), including, but not limited to, naphthalic anhydride.

Safeners may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial safeners used in accordance with the manufacturer's recommended amounts/concentrations.

Inoculant compositions of the present disclosure may comprise any suitable pH buffer(s), including, but not limited to, potassium phosphate monobasic and potassium phosphate dibasic.

Inoculant compositions of the present disclosure may comprise any suitable adhesive(s), including, but not limited to, adhesive compositions comprising one or more maltodextrins and/or one or more mono-, di- or oligosaccharides. Inoculant compositions of the present disclosure may be formulated into any suitable type of composition, including, but not limited to, seed coatings, soil inoculants and foliar inoculants.

Inoculant compositions of the present disclosure may comprise any suitable anti-settling agent(s), including, but not limited to, polyvinyl acetate, polyvinyl alcohols with different degrees of hydrolysis, polyvinylpyrrolidones, polyacrylates, acrylate-, polyol- or polyester-based paint system binders which are soluble or dispersible in water, moreover copolymers of two or more monomers such as acrylic acid, methacrylic acid, itaconic acid, maleic acid, fumaric acid, maleic anhydride, vinylpyrrolidone, ethylenically unsaturated monomers such as ethylene, butadiene, isoprene, chloroprene, styrene, divinylbenzene, *o*-methylstyrene or *p*-methylstyrene, further vinyl halides such as vinyl chloride and vinylidene chloride, additionally vinyl esters such as vinyl acetate, vinyl propionate or vinyl stearate, moreover vinyl methyl ketone or esters of acrylic acid or methacrylic acid with monohydric alcohols or polyols such as methyl acrylate, methyl methacrylate, ethyl acrylate, ethylene methacrylate, lauryl acrylate, lauryl methacrylate, decyl acrylate, *N,N*-dimethylamino-ethyl methacrylate, 2-hydroxyethyl methacrylate, 2-hydroxypropyl methacrylate or glycidyl methacrylate, furthermore diethyl esters or monoesters of unsaturated dicarboxylic acids, furthermore (meth)acrylamido-*N*-methylol methyl ether, amides or nitriles such as acrylamide, methacrylamide, *N*-methylol(meth)acrylamide, acrylonitrile, methacrylonitrile, and also *N*-substituted maleimides and ethers such as vinyl butyl ether, vinyl isobutyl ether or vinyl phenyl ether, and combinations thereof.

Anti-settling agents may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, inoculant compositions of the present disclosure comprise about 0.0001 to about 10% or more (by weight) of the composition. For example, inoculant compositions of the present disclosure may comprise about 0.0005, 0.001, 0.002, 0.003, 0.004, 0.005, 0.0075, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.75, 1, 1.25, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10% or more of one or more anti-settling agents. In some embodiments, the amount/concentration of anti-settling agents is about 0.01 to about 5% (by weight) of the composition.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial anti-settling agents used in accordance with the manufacturer's recommended amounts/concentrations.

As noted above, inoculant compositions of the present invention may comprise myriad agriculturally beneficial microbial spores. It is to be understood that inoculant compositions of the present disclosure may comprise other agriculturally beneficial constituents such as biostimulants, microbial extracts, nutrients, pesticides and plant signal molecules.

Inoculant compositions of the present disclosure may comprise any suitable biostimulant(s), including, but not limited to, seaweed extracts (e.g., *Ascophyllum nodosum* extracts, such as alginate, *Ecklonia maxima* extracts, etc.), humic acids (e.g., potassium humate), fulvic acids, myo-inositol, glycine and combinations thereof.

Biostimulants may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the biostimulant(s) comprise(s) about 0.01 to about 5 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4, 4.1, 4.2, 4.3, 4.4., 4.5, 4.6, 4.7, 4.8, 4.9, 5% or more (by weight) of one or more biostimulants. In some embodiments, the biostimulant(s) comprise(s) about 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1 to about 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4, 4.1, 4.2, 4.3, 4.4., 4.5, 4.6, 4.7, 4.8, 4.9, 5% (by weight) of the inoculant composition.

Inoculant compositions of the present disclosure may comprise any suitable microbial extract(s), including, but not limited to, bacterial extracts (e.g., extracts of media comprising one or more diazotrophs, phosphate-solubilizing microorganisms and/or biopesticides), fungal extracts and combinations thereof. In some embodiments, inoculant compositions of the present disclosure comprise one or more *Azospirillum* extracts (e.g., an extract of media comprising *A. brasilense* INTA Az-39), one or more *Bradyrhizobium* extracts (e.g., an extract of media comprising *B. elkanii* SEMIA 501, *B. elkanii* SEMIA 587, *B. elkanii* SEMIA 5019, *B. japonicum* NRRL B-50586 (also deposited as NRRL B-59565), *B. japonicum* NRRL B-50587 (also deposited as NRRL B-59566), *B. japonicum* NRRL B-50588 (also deposited as NRRL B-59567), *B. japonicum* NRRL B-50589 (also deposited as NRRL B-59568), *B. japonicum* NRRL B-50590 (also deposited as NRRL B-59569), *B. japonicum* NRRL B-50591 (also deposited as NRRL B-59570), *B. japonicum* NRRL B-50592 (also deposited as NRRL B-59571), *B. japonicum* NRRL B-50593 (also deposited as NRRL B-59572), *B. japonicum* NRRL B-50594 (also deposited as NRRL B-50493), *B. japonicum* NRRL B-50608, *B. japonicum* NRRL B-50609, *B. japonicum* NRRL B-50610, *B. japonicum* NRRL B-50611, *B. japonicum* NRRL B-50612, *B. japonicum* NRRL B-50726, *B. japonicum* NRRL B-50727, *B. japonicum* NRRL B-50728, *B. japonicum* NRRL B-50729, *B. japonicum* NRRL B-50730, *B. japonicum* SEMIA 566, *B. japonicum* SEMIA 5079, *B. japonicum* SEMIA 5080, *B. japonicum* USDA 6, *B. japonicum* USDA 110, *B. japonicum* USDA 122, *B. japonicum* USDA 123, *B. japonicum* USDA 127, *B. japonicum* USDA 129 and/or *B. japonicum* USDA 532C), one or more *Rhizobium* extracts (e.g., an extract of media comprising *R. leguminosarum* S012A-2), one or more *Sinorhizobium* extracts (e.g., an extract of media comprising *S. fredii* CCBAU1 14 and/or *S. fredii* USDA 205), one or more *Penicillium* extracts (e.g., an extract of media comprising *P. bilaiae* ATCC 18309, *P. bilaiae* ATCC 20851, *P. bilaiae* ATCC 22348, *P. bilaiae* NRRL 50162, *P. bilaiae* NRRL 50169, *P. bilaiae* NRRL 50776, *P. bilaiae* NRRL 50777, *P. bilaiae* NRRL 50778, *P. bilaiae* NRRL 50777, *P. bilaiae* NRRL 50778, *P. bilaiae* NRRL 50779, *P. bilaiae* NRRL 50780, *P. bilaiae* NRRL 50781, *P. bilaiae* NRRL 50782, *P. bilaiae* NRRL 50783, *P. bilaiae* NRRL 50784, *P. bilaiae* NRRL 50785, *P. bilaiae* NRRL 50786, *P. bilaiae* NRRL 50787, *P.*

bilaiiae NRRL 50788, *P. bilaiiae* RS7B-SD1, *P. brevicompactum* AgRF18, *P. canescens* ATCC 10419, *P. expansum* ATCC 24692, *P. expansum* YT02, *P. fellatanum* ATCC 48694, *P. gaestrivorus* NRRL 50170, *P. glabrum* DAOM 239074, *P. glabrum* CBS 229.28, *P. janthinellum* ATCC 10455, *P. lanosocoeruleum* ATCC 48919, *P. radicum* ATCC 201836, *P. radicum* FRR 4717, *P. radicum* FRR 4719, *P. radicum* N93/47267 and/or *P. raistrickii* ATCC 10490), one or more *Pseudomonas* extracts (e.g., an extract of media comprising *P. jessenii* PS06), one or more acaricidal, insecticidal and/or nematocidal extracts (e.g., an extract of media comprising *Bacillus firmus* 1-1582, *Bacillus mycoides* AQ726, NRRL B-21664; *Beauveria bassiana* ATCC-74040, *Beauveria bassiana* ATCC-74250, *Burkholderia* sp. A396 sp. nov. rinojensis, NRRL B-503 19, *Chromobacterium subtsugae* NRRL B-30655, *Chromobacterium vaccinii* NRRL B-50880, *Flavobacterium* H492, NRRL B-50584, *Metarhizium anisopliae* F52 (also known as *Metarhizium anisopliae* strain 52, *Metarhizium anisopliae* strain 7, *Metarhizium anisopliae* strain 43 and *Metarhizium anisopliae* BIO-1020, TAE-001; deposited as DSM 3884, DSM 3885, ATCC 90448, SD 170 and ARSEF 771 1) and/or *Paecilomyces fumosoroseus* FE991), and/or one or more fungicidal extracts (e.g., an extract of media comprising *Ampelomyces quisqualis* AQ 10® (Intrachem Bio GmbH & Co. KG, Germany), *Aspergillus flavus* AFLA-GUARD® (Syngenta Crop Protection, Inc., CH), *Aureobasidium pullulans* BOTECTOR® (bio-ferm GmbH, Germany), *Bacillus pumilus* AQlll (NRRL B-21662), *Bacillus pumilus* NRRL B-30087, *Bacillus* AQ175 (ATCC 55608), *Bacillus* AQ177 (ATCC 55609), *Bacillus subtilis* AQ713 (NRRL B-21661), *Bacillus subtilis* AQ743 (NRRL B-21665), *Bacillus amyloliquefaciens* FZB24, *Bacillus amyloliquefaciens* NRRL B-50349, *Bacillus amyloliquefaciens* TJ1000 (also known as 1BE, isolate ATCC BAA-390), *Bacillus thuringiensis* AQ52 (NRRL B-21619), *Candida oleophila* 1-82 (e.g., ASPIRE® from Ecogen Inc., USA), *Candida saitoana* BIOCURE® (in mixture with lysozyme; BASF, USA) and BIOCOAT® (ArystaLife Science, Ltd., Cary, NC), *Clonostachys rosea* f. *catenulata* (also referred to as *Gliocladium catenulatum*) J1446 (PRESTOP®, Verdera, Finland), *Coniothyrium minitans* CONTANS® (Prophyta, Germany), *Cryphonectria parasitica* (CNICM, France), *Cryptococcus albidus* YIELD PLUS® (Anchor Bio-Technologies, South Africa), *Fusarium oxysporum* BIOFOX® (from S.I.A.P.A., Italy) and FUSACLEAN® (Natural Plant Protection, France), *Metschnikowia fructicola* SHEMER® (Agrogreen, Israel), *Microdochium dimerum* ANTIBOT® (Agrauxine, France), *Muscodor albus* NRRL 30547, *Muscodor roseus* NRRL 30548, *Phlebiopsis gigantea* ROTSOP® (Verdera, Finland), *Pseudozyma flocculosa* SPORODEX® (Plant Products Co. Ltd., Canada), *Pythium oligandrum* DV74 (POLYVERSUM®, Remeslo SSRO, Biopreparaty, Czech Rep.), *Reynoutria sachlinensis* (e.g., REGALIA® from Marrone BioInnovations, USA), *Streptomyces* NRRL B-30145, *Streptomyces* M1064, *Streptomyces galbus* NRRL 30232, *Streptomyces lydicus* WYEC 108 (ATCC 55445), *Streptomyces violaceusniger* YCED 9 (ATCC 55660; DE-THATCH -9®, DECOMP-9® and THATCH CONTROL®, Idaho Research Foundation, USA), *Streptomyces* WYE 53 (ATCC 55750; DE-THATCH-9®, DECOMP-9® and THATCH CONTROL®, Idaho Research Foundation, USA), *Talaromyces flavus* VI 17b (PROTUS®, Prophyta, Germany), *Trichoderma asperellum* SKT-1 (ECO-HOPE®, Kumiai Chemical Industry Co., Ltd., Japan), *Trichoderma atroviride* LC52 (SENTINEL®, Agrimm Technologies Ltd, NZ), *Trichoderma harzianum* T-22 (PLANTSHIELD®, der Firma BioWorks Inc., USA), *Trichoderma harzianum* TH-35 (ROOT PRO®, from Mycontrol Ltd., Israel), *Trichoderma harzianum* T-39 (TRICHODEX®, Mycontrol Ltd., Israel; TRICHODERMA 2000®, Makhteshim Ltd., Israel), *Trichoderma harzianum* ICC012 and *Trichoderma viride* TRICHOPEL (Agrimm Technologies Ltd, NZ), *Trichoderma harzianum* ICC012 and *Trichoderma viride* ICC080 (REMEDIER® WP, Isagro Ricerca, Italy), *Trichoderma polysporum* and *Trichoderma harzianum* (BINAB®, BINAB Bio-Innovation AB, Sweden), *Trichoderma stromaticum* TRICOVAB® (C.E.P.L.A.C., Brazil), *Trichoderma virens* GL-21 (SOILGARD®, Certis LLC, USA), *Trichoderma virens* GI-3, ATCC 57678, *Trichoderma virens* GI-21 (Thermo Trilogy Corporation, Wasco, CA), *Trichoderma virens* GI-3 and *Bacillus*

amyloliquefaciens FZB2, *Trichoderma virens* GI-3 and *Bacillus amyloliquefaciens* NRRL B-50349, *Trichoderma virens* GI-3 and *Bacillus amyloliquefaciens* TJIOOO, *Trichoderma virens* GI-21 and *Bacillus amyloliquefaciens* FZB24, *Trichoderma virens* GI-21 and *Bacillus amyloliquefaciens* NRRL B-50349, *Trichoderma virens* GI-21 and *Bacillus amyloliquefaciens* TJIOOO, *Trichoderma viride* TRIECO® (Ecosense Labs. (India) Pvt. Ltd., Indien, BIO-CURE® F from T. Stanes & Co. Ltd., Indien), *Trichoderma viride* TV1 (Agribiotec srl, Italy), *Trichoderma viride* ICC080, and/or *Ulocladium oudemansii* HRU3 (BOTRY-ZEN®, Botry-Zen Ltd, NZ)).

Microbial extracts may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the microbial extract(s) comprise(s) about 0.01 to about 5 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4, 4.1, 4.2, 4.3, 4.4., 4.5, 4.6, 4.7, 4.8, 4.9, 5% or more (by weight) of one or more microbial extracts. In some embodiments, the microbial extract(s) comprise(s) about 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1 to about 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4, 4.1, 4.2, 4.3, 4.4., 4.5, 4.6, 4.7, 4.8, 4.9, 5% (by weight) of the inoculant composition.

Inoculant compositions of the present disclosure may comprise any suitable nutrient(s), including, but not limited to, organic acids (e.g., acetic acid, citric acid, lactic acid, malic acid, taurine, etc.), macrominerals (e.g., phosphorous, calcium, magnesium, potassium, sodium, iron, etc.), trace minerals (e.g., boron, cobalt, chloride, chromium, copper, fluoride, iodine, iron, manganese, molybdenum, selenium, zinc, etc.), vitamins, (e.g., vitamin A, vitamin B complex (i.e., vitamin B₁, vitamin B₂, vitamin B₃, vitamin B₅, vitamin B₆, vitamin B₇, vitamin B₈, vitamin B₉, vitamin B₁₂, choline) vitamin C, vitamin D, vitamin E, vitamin K, carotenoids (α-carotene, β-carotene, cryptoxanthin, lutein, lycopene, zeaxanthin, etc.) and combinations thereof. In some embodiments, inoculant compositions of the present disclosure comprise phosphorous, boron, chlorine, copper, iron, manganese, molybdenum and/or zinc.

Inoculant compositions of the present disclosure may comprise any suitable pesticide(s), including, but not limited to, fungicides, herbicides, insecticides, and nematocides. In some embodiments, inoculant compositions of the present disclosure comprise one or more biopesticides (e.g., one or more biofungicides, bioinsecticides and/or bionematicides).

Inoculant compositions of the present disclosure may comprise any suitable insecticide(s), including, but not limited to, biological insecticides and chemical insecticides. Insecticides may be selected so as to provide effective control against a broad spectrum of insects, including, but not limited to, insects from the orders Coleoptera, Dermaptera, Diptera, Hemiptera, Homoptera, Hymenoptera, Lepidoptera, Orthoptera and Thysanoptera. For example, inoculant compositions of the present disclosure may comprise one or more insecticides toxic to insects from the families Acrididae, Aleyrodidae, Anobiidae, Anthomyiidae, Aphididae, Bostrichidae, Bruchidae, Cecidomyiidae, Cerambycidae, Cercopidae, Chrysomelidae, Cicadellidae, Coccinellidae, Crylotalpidae, Cucujidae, Curculionidae, Dermestidae, Elateridae, Gelechiidae, Lygaeidae, Meloidae, Membracidae, Miridae, Noctuidae, Pentatomidae, Pyralidae, Scarabaeidae, Silvanidae, Spingidae, Tenebrionidae and/or Thripidae.

In some embodiments, inoculant compositions of the present disclosure comprise an insecticide (or combination of insecticides) that is toxic to one or more species of *Acalymma*, *Acanthaoscelides* (e.g., *A. obtectus*,), *Anasa* (e.g., *A. tristis*), *Anastrepha* (e.g., *A. ludens*), *Anoplophora* (e.g., *A. glabripennis*), *Anthonomus* (e.g., *A. eugeni*), *Acyrtosiphon* (e.g., *A. pisum*), *Bactrocera* (e.g., *B. dosalis*), *Bemisia* (e.g., *B. argentifolii*, *B. tabaci*), *Brevicoryne*

(e.g., *B. brassicae*), *Bruchidius* (e.g., *B. atrolineatus*), *Bruchus* (e.g., *B. atomarius*, *B. dentipes*, *B. lentis*, *B. pisorum* and/or *B. rufipes*), *Callosobruchus* (e.g., *C. chinensis*, *C. maculatus*, *C. rhodesianus*, *C. subinnotatus*, *C. theobromae*), *Caryedon* (e.g., *C. serratus*), *Cassidinae*, *Ceratitid* (e.g., *C. capitata*), *Chrysomelinae*, *Circulifer* (e.g., *C. tenellus*), *Criocerinae*, *Cryptocephalinae*, *Cryptolestes* (e.g., *C. ferrugineus*, *C. pusillus*, *C. pussilloides*), *Cylas* (e.g., *C. formicarius*), *Delia* (e.g., *D. antiqua*), *Diabrotica*, *Diaphania* (e.g., *D. nitidalis*), *Diaphorina* (e.g., *D. citri*), *Donaciinae*, *Ephestia* (e.g., *E. cautella*, *E. elutella*, *E. keuhniellid*), *Epilachna* (e.g., *E. varivestris*), *Epiphyas* (e.g., *E. postvittana*), *Eumolpinae*, *Galerucinae*, *Helicoverpa* (e.g., *H. zed*), *Heteroligus* (e.g., *H. meles*), *Iobesia* (e.g., *I. botrand*), *Lamprosomatinae*, *Lasioderma* (e.g., *L. serricorne*), *Leptinotarsa* (e.g., *L. decemlineata*), *Leptoglossus*, *Liriomyza* (e.g., *L. trifolii*), *Manducca*, *Melittia* (e.g., *M. cucurbitae*), *Myzus* (e.g., *M. persicae*), *Nezara* (e.g., *N. viridula*), *Orzaephilus* (e.g., *O. merator*, *O. surinamensis*), *Ostrinia* (e.g., *O. nubilalis*), *Phthorimaea* (e.g., *P. operculella*), *Pieris* (e.g., *P. rapae*), *Plodia* (e.g., *P. interpunctella*), *Plutella* (e.g., *P. xylostella*), *Popillia* (e.g., *P. japonica*), *Prostephanus* (e.g., *P. truncatus*), *Psila*, *Rhizopertha* (e.g., *R. dominica*), *Rhopalosiphum* (e.g., *R. maidis*), *Sagrinae*, *Solenopsis* (e.g., *S. invicta*), *Spilopyrinae*, *Sitophilus* (e.g., *S. granaries*, *S. oryzae* and/or *S. zeamais*), *Sitotroga* (e.g., *S. cerealella*), *Spodoptera* (e.g., *S. frugiperda*), *Stegobium* (e.g., *S. paniceum*), *Synetinae*, *Tenebrio* (e.g., *T. malens* and/or *T. molitor*), *Thrips* (e.g., *T. tabaci*), *Trialeurodes* (e.g., *T. vaporariorum*), *Tribolium* (e.g., *T. castaneum* and/or *T. confusum*), *Trichoplusia* (e.g., *T. ni*), *Trogoderma* (e.g., *T. granarium*) and *Trogossitidae* (e.g., *T. mauritanicus*).

Additional species of insects that may be targeted by inoculant compositions of the present disclosure may be found in CAPINERA, HANDBOOK OF VEGETABLE PESTS (2001) and Steffey and Gray, *Managing Insect Pests*, in ILLINOIS AGRONOMY HANDBOOK (2008).

Inoculant compositions of the present disclosure may comprise any suitable nematicide(s) including, but not limited to, biological nematicides and chemical nematicides. Nematicides may be selected so as to provide effective control against a broad spectrum of nematodes, including, but not limited to, phytoparasitic nematodes from the classes Chromadorea and Enoplea.

In some embodiments, inoculant compositions of the present disclosure comprise a nematicide (or combination of nematicides) that is toxic to one or more strains of *Anguina*, *Aphelenchoides*, *Belonolaimus*, *Bursaphelenchus*, *Ditylenchus*, *Globodera*, *Helicotylenchus*, *Heterodera*, *Hirschmanniella*, *Meloidogyne*, *Nacobus*, *Pratylenchus*, *Radopholus*, *Rotylenchulus*, *Trichodorus*, *Tylenchulus* and/or *Xiphinema*.

Additional examples of nematodes that may be targeted by inoculant compositions of the present disclosure may be found in CAPINERA, HANDBOOK OF VEGETABLE PESTS (2001) and Niblack, *Nematodes*, in ILLINOIS AGRONOMY HANDBOOK (2008).

As discussed above, inoculant compositions of the present disclosure may comprise one or more biological insecticides and/or nematicides (i.e., one or more microorganisms the presence and/or output of which is toxic to an acarid, insect and/or nematode).

In some embodiments, inoculant compositions of the present disclosure comprise one or more chemical insecticides and/or nematicides. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more carbamates, diamides, macrocyclic lactones, neonicotinoids, organophosphates, phenylpyrazoles, pyrethrins, spinosyns, synthetic pyrethroids, tetrionic acids and/or tetramic acids.

Non-limiting examples of chemical insecticides and nematicides that may be useful in inoculant compositions of the present disclosure include acrinathrin, alpha-cypermethrin, beta-cyfluthrin, cyhalothrin, cypermethrin, deltamethrin, esfenvalerate, etofenprox, fenprothrin, fenvalerate, flucythrinate, fosthiazate, lambda-cyhalothrin, gamma-cyhalothrin, permethrin, tau-fluvalinate, transfluthrin, zeta-cypermethrin, cyfluthrin, bifenthrin, tefluthrin,

e-flusilanat, fubfenprox, pyrethrin, resmethrin, imidacloprid, acetamiprid, thiamethoxam, nitenpyram, thiacloprid, dinotefuran, clothianidin, imidaclothiz, chlorfluazuron, diflubenzuron, lufenuron, teflubenzuron, triflumuron, novaluron, flufenoxuron, hexaflumuron, bistrifluoron, noviflumuron, buprofezin, cyromazine, methoxyfenozide, tebufenozide, halofenozide, chromafenozide, endosulfan, fipronil, ethiprole, pyrafluprole, pyriprole, flubendiamide, chlorantraniliprole (e.g., *Rynaxypyr*), cyazypyr, emamectin, emamectin benzoate, abamectin, ivermectin, milbemectin, lepimectin, tebufenpyrad, fenpyroximate, pyridaben, fenazaquin, pyrimidifen, tolfenpyrad, dicofol, cyenopyrafen, cyflumetofen, acequinocyl, fluacrypyrin, bifenazate, diafenthiuron, etoxazole, clofentezine, spinosad, triarathen, tetradifon, propargite, hexythiazox, bromopropylate, chinomethionat, amitraz, pyrifluquinazon, pymetrozine, flonicamid, pyriproxyfen, diofenolan, chlorfenapyr, metaflumizone, indoxacarb, chlo ϕ yrifos, spiroadiclofen, spiromesifen, spirotetramat, pyridalyl, spinetoram, acephate, triazophos, profenofos, oxamyl, spinetoram, fenamiphos, fenamipclothiahos, 4-[(6-cUoropyrid-3-yl)methyl](2,2 -difluoroemyl)arnino}furan -2(5H)-one, cadusaphos, carbaryl, carbofuran, ethoprophos, thiodicarb, aldicarb, aldoxycarb, metamidophos, methiocarb, sulfoxaflor, cyantraniliprole and tioxazafen and combinations thereof. In some embodiments, inoculant compositions of the present disclosure comprise abamectin, aldicarb, aldoxycarb, bifenthrin, carbofuran, chlorantraniliprole, chlothianidin, cyfluthrin, cyhalothrin, cypermethrin, cyantraniliprole, deltamethrin, dinotefuran, emamectin, ethiprole, fenamiphos, fipronil, flubendiamide, fosthiazate, imidacloprid, ivermectin, lambda-cyhalothrin, milbemectin, nitenpyram, oxamyl, permethrin, spinetoram, spinosad, spiroadiclofen, spirotetramat, tefluthrin, thiacloprid, thiamethoxam and/or thiodicarb. In some embodiments, inoculant compositions of the present disclosure comprise an insecticide selected from the group consisting of clothianidin, thiamethoxam, imidacloprid, cyantraniliprole, chlorantraniliprole, fluopyram and tioxazafen.

Additional examples of insecticides and nematicides that may be included in inoculant compositions of the present disclosure may be found in Steffey and Gray, *Managing Insect Pests, in ILLINOIS AGRONOMY HANDBOOK (2008)* and Niblack, *Nematodes, in ILLINOIS AGRONOMY HANDBOOK (2008)*.

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial insecticides and nematicides used in accordance with the manufacturer's recommended amounts/concentrations.

Inoculant compositions of the present disclosure may comprise any suitable fungicide(s), including, but not limited to, biological fungicides and chemical fungicides. Fungicides may be selected so as to provide effective control against a broad spectrum of phytopathogenic fungi (and fungus-like organisms), including, but not limited to, soil-borne fungi from the classes Ascomycetes, Basidiomycetes, Chytridiomycetes, Deuteromycetes (syn. Fungi imperfecti), Peronosporomycetes (syn. Oomycetes), Plasmodiophoromycetes and Zygomycetes.

In some embodiments, inoculant compositions of the present disclosure comprise a fungicide (or combination of fungicides) that is toxic to one or more strains of *Albugo* (e.g., *A. Candida*), *Alternaria* (e.g., *A. alternata*), *Aspergillus* (e.g., *A. candidus*, *A. clavatus*, *A. flavus*, *A. fumigatus*, *A. parasiticus*, *A. restrictus*, *A. sojae*, *A. solani*), *Blumeria* (e.g., *B. graminis*), *Botrytis* (e.g., *B. cinerea*), *Cladosporium* (e.g., *C. cladosporioides*), *Colletotrichum* (e.g., *C. acutatum*, *C. boninense*, *C. capsici*, *C. caudatum*, *C. coccodes*, *C. crassipes*, *C. dematium*, *C. destructivum*, *C. fragariae*, *C. gloeosporioides*, *C. graminicola*, *C. kehawee*, *C. lindemuthianum*, *C. musae*, *C. orbiculare*, *C. spinaceae*, *C. sublineolum*, *C. trifolii*, *C. truncatum*), *Fusarium* (e.g., *F. graminearum*, *F. moniliforme*, *F. oxysporum*, *F. roseum*, *F. tricinctum*), *Helminthosporium*, *Magnaporthe* (e.g., *M. grisea*, *M. oryzae*), *Melampsora* (e.g., *M. lini*), *Mycosphaerella* (e.g., *M. graminicola*), *Nematospora*, *Penicillium* (e.g., *P. rugulosum*, *P. verrucosum*), *Phakopsora* (e.g., *P. pachyrhizi*), *Phomopsis*, *Phytophthora* (e.g., *P. infestans*), *Puccinia* (e.g., *P. graminis*, *P. striiformis*, *P. tritici*, *P. triticina*), *Puccinia* (e.g., *P. graministice*), *Pythium*, *Pytophthora*, *Rhizoctonia* (e.g., *R. solani*), *Scopulariopsis*, *Sclerotinia*, *Thielaviopsis* and/or *Ustilago* (e.g., *U. maydis*).

Additional examples of fungi that may be targeted by inoculant compositions of the present disclosure may be found in Bradley, *Managing Diseases, in ILLINOIS AGRONOMY HANDBOOK* (2008).

As discussed above, inoculant compositions of the present disclosure may comprise one or more biological fungicides (i.e., one or more microorganisms the presence and/or output of which is toxic to a fungus).

In some embodiments, inoculant compositions of the present disclosure comprise one or more chemical fungicides. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more aromatic hydrocarbons, benzimidazoles, benzthiadiazole, carboxamides, carboxylic acid amides, morpholines, phenylamides, phosphonates, quinone outside inhibitors (e.g. strobilurins), thiazolidines, thiophanates, thiophene carboxamides and/or triazoles.

Non-limiting examples of chemical fungicides that may be useful in inoculant compositions of the present disclosure include strobilurins, such as azoxystrobin, coumethoxystrobin, coumoxystrobin, dimoxystrobin, enestroburin, fluoxastrobin, kresoxim-methyl, metominostrobin, oryastrobin, picoxystrobin, pyraclostrobin, pyrametostrobin, pyraoxystrobin, pyribencarb, trifloxystrobin, 2-[2-(2,5-dimethyl-phenoxy-methyl)-phenyl]-3-methoxy-acrylic acid methyl ester and 2-(2-(3-(2,6-dichlorophenyl)-1-methyl-allylideneamino-oxymethyl)-phenyl)-2-methoxyimino-N-methyl-acetamide; carboxamides, such as carboxanilides (e.g., benalaxyl, benalaxyl-M, benodanil, bixafen, boscalid, carboxin, fenfuram, fenhexamid, flutolanil, fluxapyroxad, furametpyr, isopyrazam, isotianil, kiralaxyl, mepronil, metalaxyl, metalaxyl-M (mefenoxam), ofurace, oxadixyl, oxycarboxin, penflufen, penthiopyrad, sedaxane, tecloftalam, thifluzamide, tiadinil, 2-amino-4-methyl-thiazole-5-carboxanilide, N-(4'-trifluoromethylthiobiphenyl-2-yl)-3-difluoromethyl-1-methyl-1H-pyrazole-4-carboxamide, N-(2-(1,3,3-trimethylbutyl)-phenyl)-1,3-dimethyl-5-fluoro-1H-pyrazole-4-carboxamide), carboxylic acid amides (e.g., flumorph, pyrimorph, benzoic acid amides (e.g., flumetover, fluopicolide, fluopyram, zoxamide), carpropamid, dicyclomet, mandiproamid, oxytetracyclin, siltiofam and N-(6-methoxy-pyridin-3-yl) cyclopropanecarboxylic acid amide; azoles, such as triazoles (e.g., azaconazole, bitertanol, bromuconazole, cyproconazole, difenoconazole, diniconazole, diniconazole-M, epoxiconazole, fenbuconazole, fluquinconazole, flusilazole, flutriafol, hexaconazole, imibenconazole, ipconazole, metconazole, myclobutanil, oxpoconazole, paclobutrazole, penconazole, propiconazole, prothioconazole, simeconazole, tebuconazole, tetraconazole, triadimefon, triadimenol, triticonazole, uniconazole) and imidazoles (e.g., cyazofamid, imazalil, pefurazoate, prochloraz, triflumizol); heterocyclic compounds, such as pyridines (e.g., fluazinam, pyrifenoxy (cf.D1b), 3-[5-(4-chloro-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine, 3-[5-(4-methyl-phenyl)-2,3-dimethyl-isoxazolidin-3-yl]-pyridine), pyrimidines (e.g., bupirimate, cyprodinil, diflometorim, fenarimol, ferimzone, mepanipyrim, nitrapyrin, nuarimol, pyrimethanil), piperazines (e.g., triforine), pyrroles (e.g., fenpiclonil, fludioxonil), naphthalenes (e.g., aldimoxy, dodemoxy, dodemoxy-acetate, fenpropimox, fenpropidin), piperidines (e.g., fenpropidin); dicarboximides (e.g., fluoroimid, iprodione, procymidone, vinclozolin), non-aromatic 5-membered heterocycles (e.g., famoxadone, fenamidone, flutianil, othilone, probenazole, 5-amino-2-isopropyl-3-oxo-4-ortho-tolyl-2,3-dihydro-pyrazole-1-carbothioic acid S-allyl ester), acibenzolar-S-methyl, ametoctradin, amisulbrom, anilazin, blasticidin-S, captan, captan, chinomethionat, dazomet, debacarb, diclomezine, difenzoquat, difenzoquat-methylsulfate, fenoxanil, Folpet, oxolinic acid, piperalin, proquinazid, pyroquilon, quinoxifen, triazoxide, tricyclazole, 2-butoxy-6-iodo-3-propylchromen-4-one, 5-chloro-1-(4,6-dimethoxy-pyrimidin-2-yl)-2-methyl-1H-benzimidazole and 5-cyano-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)-[1,2,4]triazolo-[1,5-a]pyrimidine; benzimidazoles, such as carbendazim; and other active substances, such as guanidines (e.g., guanidine, dodine, dodine free base, guazatine, guazatine-acetate, iminoctadine), iminoctadine-triacetate and iminoctadine-tris(albesilate); antibiotics (e.g., kasugamycin, kasugamycin hydrochloride-hydrate, streptomycin, polyoxine and

validamycin A), nitrophenyl derivatives (e.g., binapacryl, dicloran, dinobuton, dinocap, nitrothal-isopropyl, tecnazen), organometal compounds (e.g., fentin salts, such as fentin-acetate, fentin chloride, fentin hydroxide); sulfur-containing heterocyclyl compounds (e.g., dithianon, isoprothiolane), organophosphorus compounds (e.g., edifenphos, fosetyl, fosetyl-aluminum, iprobenfos, phosphorus acid and its salts, pyrazophos, tolclofos-methyl), organochlorine compounds (e.g., chlorothalonil, dichlofluanid, dichlorophen, flusulfamide, hexachlorobenzene, pencycuron, pentachloro phenole and its salts, phthalide, quintozene, thiophanate-methyl, thiophanate, tolylfluanid, N-(4-chloro-2-nitro-phenyl)-N-ethyl-4-methyl-benzenesulfonamide) and inorganic active substances (e.g., Bordeaux mixture, copper acetate, copper hydroxide, copper oxychloride, basic copper sulfate, sulfur) and combinations thereof. In some embodiments, inoculant compositions of the present disclosure comprise acibenzolar-S-methyl, azoxystrobin, benalaxyl, bixafen, boscalid, carbendazim, cyproconazole, dimethomorph, epoxiconazole, fludioxonil, fluopyram, fluoxastrobin, flutianil, flutolanil, fluxapyroxad, fosetyl-Al, ipconazole, isopyrazam, kresoxim-methyl, mefenoxam, metalaxyl, metconazole, myclobutanil, orysastrobin, penflufen, penthiopyrad, picoxystrobin, propiconazole, prothioconazole, pyraclostrobin, sedaxane, silthiofam, tebuconazole, thiabendazole, thifluzamide, thiophanate, tolclofos-methyl, trifloxystrobin and triticonazole. In some embodiments, inoculant compositions of the present disclosure comprise azoxystrobin, pyraclostrobin, fluoxastrobin, trifloxystrobin, ipconazole, prothioconazole, sedaxane, fludioxonil, metalaxyl, mefenoxam, thiabendazole, fluxapyroxad and/or fluopyram.

Additional examples of fungicides that may be included in inoculant compositions of the present disclosure may be found in Bradley, *Managing Diseases, in ILLINOIS AGRONOMY HANDBOOK* (2008).

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial fungicides used in accordance with the manufacturer's recommended amounts/concentrations.

Inoculant compositions of the present disclosure may comprise any suitable herbicide(s), including, but not limited to, biological herbicides and chemical herbicides. Herbicides may be selected so as to provide effective control against a broad spectrum of plants, including, but not limited to, plants from the families Asteraceae, Caryophyllaceae, Poaceae and Polygonaceae.

In some embodiments, inoculant compositions of the present disclosure comprise an herbicide (or combination of herbicides) that is toxic to one or more strains of *Echinochloa* (e.g., *E. brevipedicellata*, *E. callopus*, *E. chacoensis*, *E. colona*, *E. crus-galli*, *E. crus-pavonis*, *E. elliptica*, *E. esculenta*, *E. frumentacea*, *E. glabrescens*, *E. haploclada*, *E. helodes*, *E. holciformis*, *E. inundata*, *E. jaliscana*, *E. Jubata*, *E. kimberleyensis*, *E. lacunaria*, *E. macrandra*, *E. muricata*, *E. obtusiflora*, *E. oplismenoides*, *E. orzyoides*, *E. paludigena*, *E. picta*, *E. pithopus*, *E. polystachya*, *E. praestans*, *E. pyramidalis*, *E. rotundiflora*, *E. stagnina*, *E. telmatophila*, *E. turneriana*, *E. ugandensis*, *E. walteri*), *Fallopia* (e.g., *F. baldschuanica*, *F. japonica*, *F. sachalinensis*), *Stellaria* (e.g., *S. media*) and/or *Taraxacum* (e.g., *T. albidum*, *T. aphrogenes*, *T. brevicorniculatum*, *T. californicum*, *T. centrasiatum*, *T. ceratophorum*, *T. erythrospermum*, *T. farinosum*, *T. holmboei*, *T. japonicum*, *T. kok-saghyz*, *T. laevigatum*, *T. officinale*, *T. platycarpum*).

Additional species of plants that may be targeted by inoculant compositions of the present disclosure may be found in Hager, *Weed Management, in ILLINOIS AGRONOMY HANDBOOK* (2008) and Loux ET AL., *WEED CONTROL GUIDE FOR OHIO, INDIANA AND ILLINOIS* (2015).

As discussed above, inoculant compositions of the present disclosure may comprise one or more biological herbicides (i.e., one or more microorganisms the presence and/or output of which is toxic to a plant).

In some embodiments, inoculant compositions of the present disclosure comprise one or more chemical herbicides. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more acetyl CoA carboxylase (ACCCase) inhibitors, acetolactate synthase (ALS) inhibitors, acetohydroxy acid

synthase (AHAS) inhibitors, photosystem II inhibitors, photosystem I inhibitors, protoporphyrinogen oxidase (PPO or Protox) inhibitors, carotenoid biosynthesis inhibitors, enolpyruvyl shikimate-3-phosphate (EPSP) synthase inhibitor, glutamine synthetase inhibitor, dihydropteroate synthetase inhibitor, mitosis inhibitors, 4-hydroxyphenylpyruvate-dioxygenase (4-HPPD) inhibitors, synthetic auxins, auxin herbicide salts, auxin transport inhibitors, nucleic acid inhibitors and/or one or more salts, esters, racemic mixtures and/or resolved isomers thereof. Non-limiting examples of chemical herbicides that may be useful in inoculant compositions of the present disclosure include 2,4-dichlorophenoxyacetic acid (2,4-D), 2,4,5-trichlorophenoxyacetic acid (2,4,5-T), ametryn, amicarbazone, aminocyclopyrachlor, acetochlor, acifluorfen, alachlor, atrazine, azafenidin, bentazon, benzofenap, bifenox, bromacil, bromoxynil, butachlor, butafenacil, butoxydim, carfentrazone-ethyl, chlorimuron, chlorotoluro, clethodim, clodinafop, clomazone, cyanazine, cycloxydim, cyhalofop, desmedipham, desmetryn, dicamba, diclofop, dimefuron, diuron, dithiopyr, fenoxaprop, fluzifop, fluzifop-P, fluometuron, flufenpyr-ethyl, flumiclorac-pentyl, flumioxazin, fluoroglycofen, fluthiacet-methyl, fomesafen, fomesafen, glyphosate, glufosinate, haloxyfop, hexazinone, imazamox, imazaquin, imazethapyr, ioxynil, isoproturon, isoxaflutole, lactofen, linuron, mecoprop, mecoprop-P, mesotrione, metamilon, metazachlor, methibenzuron, metolachlor (and S-metolachlor), metoxuron, metribuzin, monolinuron, oxadiargyl, oxadiazon, oxyfluorfen, phenmedipham, pretilachlor, profoxydim, prometon, prometry, propachlor, propanil, propaquizafop, propisochlor, pyraflufen-ethyl, pyrazon, pyrazolynate, pyrazoxyfen, pyridate, quizalofop, quizalofop-P (e.g., quizalofop-ethyl, quizalofop-P-ethyl, clodinafop-propargyl, cyhalofop-butyl, diclofop-methyl, fenoxaprop-P-ethyl, fluzifop-P-butyl, haloxyfop-methyl, haloxyfop-R-methyl), saflufenacil, sethoxydim, siduron, simazine, simetryn, sulcotrione, sulfentrazone, tebuthiuron, tembotrione, tepraloxym, terbacil, terbutolone, terbuthylazine, thaxtomin (e.g., the thaxtomins described in US Patent No.: 7,989,393), thenylchlor, tralkoxydim, triclopyr, trietazine, tropamezone, salts and esters thereof; racemic mixtures and resolved isomers thereof and combinations thereof. In some embodiments, inoculant compositions of the present disclosure comprise acetochlor, clethodim, dicamba, flumioxazin, fomesafen, glyphosate, glufosinate, mesotrione, quizalofop, saflufenacil, sulcotrione, ethyl 2-((3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-2,3-dihydropyrimidin-1(6H)-yl)phenoxy)pyridin-2-yl)oxy)acetate and/or 2,4-D. In some embodiments, inoculant compositions of the present disclosure comprise glyphosate, glufosinate, dicamba, 2,4-D, acetochlor, metolachlor, pyroxasulfone, flumioxazin, fomesafen, lactofen, metribuzin, mesotrione, and/or ethyl 2-((3-(2-chloro-4-fluoro-5-(3-methyl-2,6-dioxo-4-(trifluoromethyl)-2,3-dihydropyrimidin-1(6H)-yl)phenoxy)pyridin-2-yl)oxy)acetate.

Additional examples of herbicides that may be included in inoculant compositions of the present disclosure may be found in Hager, *Weed Management, in ILLINOIS AGRONOMY HANDBOOK* (2008) and Loux ET AL., *WEED CONTROL GUIDE FOR OHIO, INDIANA AND ILLINOIS* (2015).

In some embodiments, inoculant compositions of the present disclosure comprise one or more commercial herbicides used in accordance with the manufacturer's recommended amounts/concentrations.


Inoculant compositions of the present disclosure may comprise any suitable plant signal molecule(s), including, but not limited to, lipo-chitooligosaccharides (LCOs), chitooligosaccharides (COs), chitinous compounds, flavonoids, jasmonic acid or derivatives thereof, linoleic acid or derivatives thereof, linolenic acid or derivatives thereof and karrikins.

Inoculant compositions of the present disclosure may comprise any suitable LCO(s).

LCOs, sometimes referred to as symbiotic nodulation (Nod) signals or Nod factors, consist of an oligosaccharide backbone of β -1,4-linked *N*-acetyl-D-glucosamine ("GlcNAc") residues with an N-linked fatty acyl chain condensed at the non-reducing end. LCOs differ in the number of GlcNAc residues in the backbone, in the length and degree of saturation of the fatty acyl chain and in the substitutions of reducing and non-reducing sugar

residues. See, e.g., Denarie, *et al*, ANN. REV. BIOCHEM. 65:503 (1996); Hamel, *et al*, PLANTA 232:787 (2010); Prome, *et al*, PURE & APPL. CHEM. 70(1):55 (1998).


In some embodiments, inoculant compositions of the present disclosure comprise one or more LCOs represented by formula I:



(I)

in which G is a hexosamine which can be substituted, for example, by an acetyl group on the nitrogen, a sulfate group, an acetyl group and/or an ether group on an oxygen; R₁, R₂, R₃, R₅, R₆ and R₇, which may be identical or different, represent H, CH₃ CO-, C_xH_y CO-- where x is an integer between 0 and 17 and y is an integer between 1 and 35, or any other acyl group such as, for example, a carbamoyl; R₄ represents a saturated or mono-, di- or tri-unsaturated aliphatic chain containing at least 12 carbon atoms; and n is an integer between 1 and 4.


In some embodiments, inoculant compositions of the present disclosure comprise one or more LCOs represented by formula II:



(II)

in which R represents H or CH₃CO-- and n is equal to 2 or 3. See, e.g., U.S. Patent No. 5,549,718. A number of *Bradyrhizobium japonicum*-derived LCOs have also been described, including BjNod-V (Ci_{8,i}), BjNod-V (Ac, Ci_{8,i}), BjNod-V (Ci_{6,i}) and BjNod-V (Ac, Ci_{6,i}) (with "V" indicating the presence of five N-acetylglucosamines, "Ac" an acetylation, the number following the "C" indicating the number of carbons in the fatty acid side chain and the number following the ":" indicating the number of double bonds). See, e.g., U.S. Patent Nos. 5,175,149 and 5,321,011. Additional LCOs obtained from bacterial strains include NodRM, NodRM-1, NodRM-3. When acetylated (the R=CH₃CO--), they become AcNodRM-1 and AcNodRM-3, respectively (U.S. Patent No. 5,545,718).

In some embodiments, inoculant compositions of the present disclosure comprise one or more LCOs represented by formula III:



in which n = 1 or 2; Ri represents C16, C16:0, C16:1, C16:2, C18:0, C18:1A9Z or C18:1A11Z; and R₂ represents hydrogen or SO₃H.


LCOs included in compositions and methods of the present disclosure may be obtained from any suitable source.

In some embodiments, the LCO is obtained (i.e., isolated and/or purified) from a bacterial strain. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more LCOs obtained from a of *Azorhizobium*, *Bradyrhizobium* (e.g., *B. japonicum*), *Mesorhizobium*, *Rhizobium* (e.g., *R. leguminosarum*), or *Sinorhizobium* (e.g., *S. meliloti*).

In some embodiments, the LCO is obtained (i.e., isolated and/or purified) from a mycorrhizal fungus. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more LCOs obtained from a strain of Glomerocycota (e.g., *Glomus intraradieux*). See, e.g., WO 2010/049751 (in which the LCOs are referred to as "Myc factors").


In some embodiments, the LCO is synthetic. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more of the synthetic LCOs described in WO 2005/063784, WO 2007/117500 and/or WO 2008/071674. In some embodiments, the synthetic LCO contains one or more modifications or substitutions, such as those described in Spaink, *CRIT. REV. PLANT SCI.* 54:257 (2000) and D'Haeze, supra. LCOs and precursors for the construction of LCOs (e.g., COs, which are themselves useful as plant signal molecules) may be synthesized by genetically engineered organisms. See, e.g., Samain et al., *CARBOHYDRATE RES.* 302:35 (1997); Cottaz, et al., *METH. ENG.* 7(4):311 (2005); and Samain, et al., *J. BIOTECHNOL.* 72:33 (1999) (e.g., Fig. 1 therein, which shows structures of COs that can be made recombinantly in *E. coli* harboring different combinations of genes *nodBCHL*).


Further examples of LCOs (and derivatives thereof) that may be useful in compositions and methods of the present disclosure are provided below as formula IV:




in which R_i represents C14:0, 3OH-C14:0, iso-C15:0, C16:0, 3-OH-C16:0, iso-C15:0, C16:1, C16:2, C16:3, iso-C17:0, iso-C17:1, C18:0, 3OH-C18:0, C18:0/3-OH, C18:1, OH-C18:1, C18:2, C18:3, C18:4, C19:1 carbamoyl, C20:0, C20:1, 3-OH-C20:1, C20:1/3-OH, C20:2, C20:3, C22:1 and C18-26(co-l)-OH (which according to D'Haese, *et al.*, *Glycobiology* 72:79R-105R (2002), includes C18, C20, C22, C24 and C26 hydroxylated species and C16:1A9, C16:2 (Δ 2,9) and C16:3 (Δ 2,4,9)); R_2 represents hydrogen or methyl; R_3 represents hydrogen, acetyl or carbamoyl; R_4 represents hydrogen, acetyl or carbamoyl; R_5 represents hydrogen, acetyl or carbamoyl; R_6 represents hydrogen, arabinosyl, fucosyl, acetyl, SO_3H , sulfate ester, 3-O-S-2-O-MeFuc, 2-O-MeFuc and 4-O-AcFuc; R_7 represents hydrogen, mannosyl or glycerol; R_8 represents hydrogen, methyl, or $-CH_2OH$; R_9 represents hydrogen, arabinosyl, or fucosyl; R_{i0} represents hydrogen, acetyl or fucosyl; and n represents 0, 1, 2 or 3. Naturally occurring LCOs embraced by this structure are described in D'Haese, *et al.*, *supra*.

Further examples of LCOs (and derivatives thereof) that may be useful in compositions and methods of the present disclosure are provided below as structures V-XXXIII:







(VI)




(VII)




(VIII)




(IX)




(X)




(XI)




(XII)




(XIII)




(XIV)




(XV)




(XVI)




(XVII)




(XVIII)




(XIX)




(XX)




(XXI)




(XXII)




(XXIII)




(XXIV)




(XXV)




(XXVI)




(XXVII)




(XXVIII)




(XXIX)




(XXX)



(XXXI)



(XXXII)



(XXXIII)

It is to be understood that compositions and methods of the present disclosure may comprise analogues, derivatives, hydrates, isomers, salts and/or solvates of LCOs.

Thus, in some embodiments, inoculant compositions of the present disclosure comprise one, two, three, four, five, six, seven, eight, nine, ten, or more LCOs represented by one or more of formulas I-IV and/or structures V-XXXIII and/or one, two, three, four, five, six, seven, eight, nine, ten, or more analogues, derivatives, hydrates, isomers, salts and/or solvates of LCOs represented by one or more of formulas I-IV and/or structures V-XXXIII.


LCOs may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, the inoculant compositions of the present disclosure comprise about 1×10^{-20} M to about 1×10^{-1} M LCO. For example, inoculant compositions of the present disclosure may comprise about 1×10^{-20} M, 1×10^{-19} M, 1×10^{-18} M, 1×10^{-17} M, 1×10^{-16} M, 1×10^{-15} M, 1×10^{-14} M, 1×10^{-13} M, 1×10^{-12} M, 1×10^{-11} M, 1×10^{-10} M, 1×10^{-9} M, 1×10^{-8} M, 1×10^{-7} M, 1×10^{-6} M, 1×10^{-5} M, 1×10^{-4} M, 1×10^{-3} M, 1×10^{-2} M, 1×10^{-1} M of one or more LCOs. In some embodiments, the LCO concentration is 1×10^{-14} M to 1×10^{-5} M, 1×10^{-12} M to 1×10^{-6} M, or 1×10^{-10} M to 1×10^{-7} M. In some embodiments, the LCO concentration is 1×10^{-14} M to 1×10^{-5} M, 1×10^{-12} M to 1×10^{-6} M, or 1×10^{-10} M to 1×10^{-7} M.

Inoculant compositions of the present disclosure may comprise any suitable CO(s).

COs, sometimes referred to as N-acetylchitooligosaccharides, are also composed of GlcNAc residues but have side chain decorations that make them different from chitin molecules $[(C_8H_{13}NO_5)_n]$, CAS No. 1398-61-4 and chitosan molecules $[(C_5H_9NO_4)_n]$, CAS No. 9012-76-4. See, e.g., D'Haese et al., *GLYCOBIOL.* 12(6):79R (2002); Demont-Caulet et al., *PLANT PHYSIOL.* 120(1):83 (1999); Hanel et al., *PLANTA* 232:787 (2010); MuUer et al., *PLANT PHYSIOL.* 124:733 (2000); Robina et al., *TETRAHEDRON* 58:521-530 (2002); Rouge et al., *Docking of Chitin Oligomers and Nod Factors on Lectin Domains of the LysM-RLK Receptors in the Medicago-Rhizobium Symbiosis, in THE MOLECULAR IMMUNOLOGY OF COMPLEX CARBOHYDRATES-3* (Springer Science, 2011); Van der Hoist et al., *CURR. OPIN. STRUC. BIOL.* 11:608 (2001); and Wan et al., *PLANT CELL* 21:1053 (2009); PCT/F 100/00803 (2000). COs differ from LCOs in that they lack the pendant fatty acid chain that is characteristic of LCOs.

In some embodiments, inoculant compositions of the present disclosure comprise one or more COs represented by formula XXXIV:




in which R_1 represents hydrogen or methyl; R_2 represents hydrogen or methyl; R_3 represents hydrogen, acetyl or carbamoyl; R_4 represents hydrogen, acetyl or carbamoyl; R_5 represents hydrogen, acetyl or carbamoyl; R_6 represents hydrogen, arabinosyl, fucosyl, acetyl, sulfate ester, 3-0-S-2-0-MeFuc, 2-0-MeFuc and 4-0-AcFuc; R_7 represents hydrogen, mannosyl or glycerol; R_8 represents hydrogen, methyl, or $-CH_2OH$; R_9 represents hydrogen, arabinosyl, or fucosyl; R_{10} represents hydrogen, acetyl or fucosyl; and n represents 0, 1, 2 or 3.

COs included in compositions and methods of the present disclosure may be obtained from any suitable source.

In some embodiments, the CO is derived from an LCO. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more COs derived from an LCO obtained (i.e., isolated and/or purified) from a strain of *Azorhizobium*, *Bradyrhizobium* (e.g., *B. japonicum*), *Mesorhizobium*, *Rhizobium* (e.g., *R. leguminosarum*), *Sinorhizobium* (e.g., *S. meliloti*), or mycorrhizal fungus (e.g., *Glomus intraradices*). In some embodiments, the CO is derived from an LCO represented by one or more of formulas I-IV and/or structures V-XXXIII. Thus, in some embodiments, inoculant compositions of the present disclosure may comprise one or more COs represented by one or more of formulas I-IV and/or structures V-XXXIII except that the pendant fatty acid is replaced with a hydrogen or methyl group.


In some embodiments, the CO is synthetic. Methods for the preparation of recombinant COs are known in the art. See, e.g., Cottaz et al., METH. ENG. 7(4):3 11 (2005); Samain et al., CARBOHYDRATE RES. 302:35 (1997.); and Samain et al., J. BIOTECHNOL. 72:33 (1999).


Examples of COs (and derivatives thereof) that may be useful in compositions and methods of the present disclosure are provided below as formula XXXV:




in which $n = 1$ or 2 ; R_1 represents hydrogen or methyl; and R_2 represents hydrogen or SO_3H .

Further examples of COs (and derivatives thereof) that may be useful in compositions and methods of the present disclosure are provided below as structures XXXVI-XXXIX:






(XXXVII)



(XXXVIII)



(XXXIX)

COs (and derivatives thereof) may be utilized in various forms of purity and may be used alone or in the form of a culture of CO-producing bacteria or fungi. In some embodiments, the CO(s) included in inoculant compositions of the present disclosure is/are at least 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more pure.

It is to be understood that compositions and methods of the present disclosure may comprise hydrates, isomers, salts and/or solvates of COs.

Thus, in some embodiments, inoculant compositions of the present disclosure comprise one, two, three, four, five, six, seven, eight, nine, ten, or more COs represented by one or more of formulas XXXIV-XXXV and/or structures XXXVI-XXXIX and/or one, two, three, four, five, six, seven, eight, nine, ten, or more analogues, derivatives, hydrates, isomers, salts and/or solvates of COs represented by one or more of formulas XXXIV-XXXV and/or structures XXXVI-XXXIX.

COs may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). In some embodiments, the inoculant compositions of the present disclosure comprise about 1×10^{-20} M to about 1×10^{-1} M CO. For example, inoculant compositions of the present disclosure may comprise about 1×10^{-20} M, 1×10^{-19} M, 1×10^{-18} M, 1×10^{-17} M, 1×10^{-16} M, 1×10^{-15} M, 1×10^{-14} M, 1×10^{-13} M, 1

1×10^{-12} M, 1×10^{-11} M, 1×10^{-10} M, 1×10^{-9} M, 1×10^{-8} M, 1×10^{-7} M, 1×10^{-6} M, 1×10^{-5} M, 1×10^{-4} M, 1×10^{-3} M, 1×10^{-2} M, 1×10^{-1} M of one or more COs. In some embodiments, the CO concentration is 1×10^{-14} M to 1×10^{-5} M, 1×10^{-12} M to 1×10^{-6} M, or 1×10^{-10} M to 1×10^{-7} M. In some embodiments, the CO concentration is 1×10^{-14} M to 1×10^{-5} M, 1×10^{-12} M to 1×10^{-6} M, or 1×10^{-10} M to 1×10^{-7} M.

Inoculant compositions of the present disclosure may comprise any suitable chitinous compound(s), including, but not limited to, chitin (IUPAC: N-[5-[[3-acetylamino-4,5-dihydroxy-6-(hydroxymethyl)oxan-2yl]methoxymethyl]-2-[[5-acetylamino-4,6-dihydroxy-2-(hydroxymethyl)oxan-3-yl]methoxymethyl]-4-hydroxy-6-(hydroxymethyl)oxan-3-yl]ethanamide), chitosan (IUPAC: 5-amino-6-[5-amino-6-[5-amino-4,6-dihydroxy-2-(hydroxymethyl)oxan-3-yl]oxy-4-hydroxy-2-(hydroxymethyl)oxan-3-yl]oxy-2-(hydroxymethyl)oxane-3,4-diol) and isomers, salts and solvates thereof.

Chitins and chitosans, which are major components of the cell walls of fungi and the exoskeletons of insects and crustaceans, are composed of GlcNAc residues.

Chitins and chitosans may be obtained commercially or prepared from insects, crustacean shells, or fungal cell walls. Methods for the preparation of chitin and chitosan are known in the art. *See, e.g.*, U.S. Patent Nos. 4,536,207 (preparation from crustacean shells) and 5,965,545 (preparation from crab shells and hydrolysis of commercial chitosan); Pochanavanich, et al., LETT. APPL. MICROBIOL. 35: 17 (2002) (preparation from fungal cell walls).

Deacetylated chitins and chitosans may be obtained that range from less than 35% to greater than 90% deacetylation and cover a broad spectrum of molecular weights, e.g., low molecular weight chitosan oligomers of less than 15kD and chitin oligomers of 0.5 to 2kD; "practical grade" chitosan with a molecular weight of about 15kD; and high molecular weight chitosan of up to 70kD. Chitin and chitosan compositions formulated for seed treatment are commercially available. Commercial products include, for example, ELEXA® (Plant Defense Boosters, Inc.) and BEYOND™ (Agrihouse, Inc.).

Inoculant compositions of the present disclosure may comprise any suitable flavonoid(s), including, but not limited to, anthocyanidins, anthoxanthins, chalcones, coumarins, flavanones, flavanonols, flavans and isoflavonoids, as well as analogues, derivatives, hydrates, isomers, polymers, salts and solvates thereof.

Flavonoids are phenolic compounds having the general structure of two aromatic rings connected by a three-carbon bridge. Classes of flavonoids include are known in the art. *See, e.g.*, Jain et al., J. PLANT BIOCHEM. & BIOTECHNOL. 11:1 (2002); Shaw et al., ENVIRON. MICROBIOL. 11:1867 (2006). Flavonoid compounds are commercially available, e.g., from Novozymes BioAg, Saskatoon, Canada; Natland International Corp., Research Triangle Park, NC; MP Biomedicals, Irvine, CA; LC Laboratories, Woburn MA. Flavonoid compounds may be isolated from plants or seeds, e.g., as described in U.S. Patents 5,702,752; 5,990,291; and 6,146,668. Flavonoid compounds may also be produced by genetically engineered organisms, such as yeast, as described in Ralston et al., PLANT PHYSIOL. 137:1375 (2005).

In some embodiments, inoculant compositions of the present disclosure comprise one or more anthocyanidins. For example, in some embodiments, inoculant compositions of the present disclosure comprise cyanidin, delphinidin, malvidin, pelargonidin, peonidin and/or petunidin.

In some embodiments, inoculant compositions of the present disclosure comprise one or more anthoxanthins. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more flavones (e.g., apigenin, baicalein, chrysin, 7,8-dihydroxyflavone, diosmin, flavoxate, 6—hydroxyflavone, luteolin, scutellarein, tangeritin and/or wogonin) and/or flavonols (e.g., amurensin, astragalins, azaleatin, azalein, fisetin, furanoflavonols galangin, gossypetin, 3-hydroxyflavone, hyperoside, icariin, isoquercetin, kaempferide,

kaempferitrin, kaempferol, isorhamnetin, morin, myricetin, myricitrin, natsudaidain, pachypodol, pyranoflavonols quercetin, quercitin, rhamnazin, rhamnetin, robinin, rutin, spiraeoside, troxerutin and/or zanthorhamnin).

In some embodiments, inoculant compositions of the present disclosure comprise one or more flavanones. For example, in some embodiments, inoculant compositions of the present disclosure comprise butin, eriodictyol, hesperetin, hesperidin, homoeriodictyol, isosakuranetin, naringenin, naringin, pinocembrin, poncirin, sakuranetin, sakuranin and/or sterubin.

In some embodiments, inoculant compositions of the present disclosure comprise one or more flavanonols. For example, in some embodiments, inoculant compositions of the present disclosure comprise dihydrokaempferol and/or taxifolin.

In some embodiments, inoculant compositions of the present disclosure comprise one or more flavans. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more flavan-3-ols (e.g., catechin (C), catechin 3-gallate (Cg), epicatechins (EC), epigallocatechin (EGC) epicatechin 3-gallate (ECg), epigallocatechin 3-gallate (EGCg), epiafzelechin, fisetinidol, gallocatechin (GC), gallcatechin 3-gallate (GCg), guibourtinidol, mesquitol, robinetinidol, theaflavin-3-gallate, theaflavin-3'-gallate, theflavin-3,3'-digallate, thearubigin), flavan-4-ols (e.g., apiforol and/or luteoforol) and/or flavan-3,4-diols (e.g., leucocyanidin, leucodelphinidin, leucofisetinidin, leucomalvidin, luecopelargonidin, leucopaeonidin, leucorobinetinidin, melacacidin and/or teracacidin) and/or dimers, trimers, oligomers and/or polymers thereof (e.g., one or more proanthocyanidins).

In some embodiments, inoculant compositions of the present disclosure comprise one or more isoflavonoids. For example, in some embodiments, inoculant compositions of the present disclosure comprise one or more isoflavones (e.g. biochanin A, daidzein, formononetin, genistein and/or glycitein), isoflavanes (e.g., equol, ionchocarpane and/or laxifloorane), isoflavandiols, isoflavenes (e.g., glabrene, haginin D and/or 2-methoxyjudaicin), coumestans (e.g., coumestrol, plicadin and/or wedelolactone), pterocarpan and/or roetionoids.

Inoculant compositions of the present disclosure may comprise any suitable flavonoid derivative, including, but not limited to, neoflavonoids (e.g. calophyllolide, coutareagenin, dalbergichromene, dalbergin, nivetin) and pterocarpan (e.g., bitucarpin A, bituca ϕ in **B**, erybraedin A, erybraedin **B**, erythrabysin II, erthrabissin-1, erycristagallin, glycinol, glyceollidins, glyceollins, glycyrrhizol, maackiain, medicarpin, morisianine, orientanol, phaseolin, pisatin, striatine, trifolirhizin).

Flavonoids and derivatives thereof may be incorporated into inoculant compositions of the present disclosure in any suitable form, including, but not limited to, polymorphic and crystalline forms.

Flavonoids may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s). Inoculant compositions of the present disclosure may comprise any suitable non-flavonoid node-gene inducer(s), including, but not limited to, jasmonic acid ([1Pv-[1 α ,2 β (Z)]]-3-oxo-2-(pentenyl)cyclopentaneacetic acid; JA), linoleic acid ((Z,Z)-9,12-Octadecadienoic acid) and linolenic acid ((Z,Z,Z)-9,12,15-octadecatrienoic acid), as well as analogues, derivatives, hydrates, isomers, polymers, salts and solvates thereof.


Jasmonic acid and its methyl ester, methyl jasmonate (MeJA), collectively known as jasmonates, are octadecanoid-based compounds that occur naturally in some plants (e.g., wheat), fungi (e.g., *Botryodiplodia theobromae*, *Gibberella fujikuroi*), yeast (e.g., *Saccharomyces cerevisiae*) and bacteria (e.g., *Escherichia coli*). Linoleic acid and linolenic acid may be produced in the course of the biosynthesis of jasmonic acid. Jasmonates, linoleic acid and linolenic acid (and their derivatives) are reported to be inducers of nod gene expression or LCO production by rhizobacteria. See, e.g., Mabood, et al. **PLANT PHYSIOL. BIOCHEM.** 44(1 1):759 (2006); Mabood et al.,

AGR. J. 98(2):289 (2006); Mabood, et al., FIELD CROPS RES.95(2-3):412 (2006); Mabood & Smith, *Linoleic and linolenic acid induce the expression of nod genes in Bradyrhizobium japonicum* USDA 3, PLANT BIOL. (2001).

Useful derivatives of jasmonic acid, linoleic acid, linolenic acid that may be useful in compositions of the present disclosure include esters, amides, glycosides and salts. Representative esters are compounds in which the carboxyl group of linoleic acid, linolenic acid, or jasmonic acid has been replaced with a --COR group, where R is an --OR¹ group, in which R¹ is: an alkyl group, such as a C₁-C₈ unbranched or branched alkyl group, e.g., a methyl, ethyl or propyl group; an alkenyl group, such as a C₂-C₈ unbranched or branched alkenyl group; an alkynyl group, such as a C₂-C₈ unbranched or branched alkynyl group; an aryl group having, for example, 6 to 10 carbon atoms; or a heteroaryl group having, for example, 4 to 9 carbon atoms, wherein the heteroatoms in the heteroaryl group can be, for example, N, O, P, or S. Representative amides are compounds in which the carboxyl group of linoleic acid, linolenic acid, or jasmonic acid has been replaced with a --COR group, where R is an NR²R³ group, in which R² and R³ are independently: hydrogen; an alkyl group, such as a C₁-C₈ unbranched or branched alkyl group, e.g., a methyl, ethyl or propyl group; an alkenyl group, such as a C₂-C₈ unbranched or branched alkenyl group; an alkynyl group, such as a C₂-C₈ unbranched or branched alkynyl group; an aryl group having, for example, 6 to 10 carbon atoms; or a heteroaryl group having, for example, 4 to 9 carbon atoms, wherein the heteroatoms in the heteroaryl group can be, for example, N, O, P, or S. Esters may be prepared by known methods, such as acid-catalyzed nucleophilic addition, wherein the carboxylic acid is reacted with an alcohol in the presence of a catalytic amount of a mineral acid. Amides may also be prepared by known methods, such as by reacting the carboxylic acid with the appropriate amine in the presence of a coupling agent such as dicyclohexyl carbodiimide (DCC), under neutral conditions. Suitable salts of linoleic acid, linolenic acid and jasmonic acid include e.g., base addition salts. The bases that may be used as reagents to prepare metabolically acceptable base salts of these compounds include those derived from cations such as alkali metal cations (e.g., potassium and sodium) and alkaline earth metal cations (e.g., calcium and magnesium). These salts may be readily prepared by mixing together a solution of linoleic acid, linolenic acid, or jasmonic acid with a solution of the base. The salts may be precipitated from solution and be collected by filtration or may be recovered by other means such as by evaporation of the solvent.

Non-flavonoid node-gene inducers may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

Inoculant compositions of the present disclosure may comprise karrakin(s), including, but not limited to, 2H-furo[2,3-c]pyran-2-ones, as well as analogues, derivatives, hydrates, isomers, polymers, salts and solvates thereof. In some embodiments, the inoculant composition comprises one or more karrakins represented by formula XXXX:



in which Z is O, S or NR₅; R₁, R₂, R₃ and R₄ are each independently H, alkyl, alkenyl, alkynyl, phenyl, benzyl, hydroxy, hydroxyalkyl, alkoxy, phenoxy, benzyloxy, CN, COR₆, COOR₆, halogen, NR₆R₇, or NO₂; and R₅, R₆ and R₇ are each independently H, alkyl or alkenyl, or a biologically acceptable salt thereof.

Examples of biologically acceptable salts of karrakins may include acid addition salts formed with biologically acceptable acids, examples of which include hydrochloride, hydrobromide, sulphate or bisulphate, phosphate or hydrogen phosphate, acetate, benzoate, succinate, fumarate, maleate, lactate, citrate, tartrate, gluconate; methanesulphonate, benzenesulphonate and p-toluenesulphonic acid. Additional biologically acceptable metal salts may include alkali metal salts, with bases, examples of which include the sodium and potassium salts. Examples of compounds embraced by formula XXXX and which may be suitable for use in the present disclosure include 3-methyl-2H-furo[2,3-c]pyran-2-one (where $R_i=CH_3$, $R_2, R_3, R_4=H$), 2H-furo[2,3-c]pyran-2-one (where $R_1, R_2, R_3, R_4=H$), 7-methyl-2H-furo[2,3-c]pyran-2-one (where $R_1, R_2, R_4=H, R_3=CH_3$), 5-methyl-2H-furo[2,3-c]pyran-2-one (where $R_1, R_2, R_3=H, R_4=CH_3$), 3,7-dimethyl-2H-furo[2,3-c]pyran-2-one (where $R_1, R_3=CH_3, R_2, R_4=H$), 3,5-dimethyl-2H-furo[2,3-c]pyran-2-one (where $R_1, R_4=CH_3, R_2, R_3=H$), 3,5,7-trimethyl-2H-furo[2,3-c]pyran-2-one (where $R_1, R_3, R_4=CH_3, R_2=H$), 5-methoxymethyl-3-methyl-2H-furo[2,3-c]pyran-2-one (where $R_i=CH_3, R_2, R_3=H, R_4=CH_2OCH_3$), 4-bromo-3,7-dimethyl-2H-furo[2,3-c]pyran-2-one (where $R_1, R_3=CH_3, R_2=Br, R_4=H$), 3-methylfuro[2,3-c]pyridin-2(3H)-one (where $Z=NH, R_i=CH_3, R_2, R_3, R_4=H$) and 3,6-dimethylfuro[2,3-c]pyridin-2(6H)-one (where $Z=N-CH_3, R_i=CH_3, R_2, R_3, R_4=H$). See, e.g., U.S. Patent No. 7,576,213; Halford, *Smoke Signals*, in CHEM. ENG. NEWS (April 12, 2010) (reporting that karrakins or butenolides contained in smoke act as growth stimulants and spur seed germination after a forest fire and can invigorate seeds such as corn, tomatoes, lettuce and onions that had been stored).

Karrakins may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

Inoculant compositions of the present disclosure may comprise gluconolactone and/or one or more analogues, derivatives, hydrates, isomers, polymers, salts and/or solvates thereof.

Gluconolactone may be incorporated into inoculant compositions of the present disclosure in any suitable amount(s)/concentration(s).

In some embodiments, inoculant compositions of the present disclosure are formulated as non-aqueous formulations in which at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of the microbial spores therein survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, inoculant compositions of the present disclosure are formulated such that at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70% or more of the microorganisms therein survive when the inoculant composition is coated on a seed and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 50, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure are formulated as non-aqueous formulations in which at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/gram or more of the microbial spores therein survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative

humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure are formulated as non-aqueous formulations in which at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/ml or more of the microbial spores therein survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure are formulated such that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cru/seed or more of the microorganisms therein survive when the inoculant composition is coated on a seed and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, inoculant compositions of the present disclosure are formulated as amorphous solids.

In some embodiments, inoculant compositions of the present disclosure are formulated as wettable powders.

In some embodiments, inoculant compositions of the present disclosure are formulated as granules.

In some embodiments, inoculant compositions of the present disclosure comprise no water.

In some embodiments, inoculant compositions of the present disclosure comprise a trace amount of water.

In some embodiments, inoculant compositions of the present disclosure comprise less than 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5% water by weight, based upon the total weight of the composition.

In some embodiments, inoculant compositions of the present disclosure have a pH of about 4.5 to about 9.5. In some embodiments, inoculant compositions of the present disclosure have a pH of about 6 to about 7.5. In some embodiments, inoculant compositions of the present disclosure have a pH of about 5, 5.5, 6, 6.5, 7, 7.5, 8 or 8.5.

As noted above, inoculant compositions of the present disclosure may comprise agriculturally beneficial constituents, such as biostimulants, microbial extracts, nutrients, pesticides and plant signal molecules. It is to be understood that agriculturally beneficial constituents may also be used in conjunction with inoculant compositions of the present disclosure. Thus, the present disclosure extends to systems and methods of using inoculant compositions of the present disclosure in conjunction with compositions comprising one or more agriculturally beneficial constituents (e.g., a second composition comprising one or more LCOs and/or COs, a third composition comprising one or more fungicides, herbicides, insecticides and/or nematocides, etc.).

Inoculant compositions of the present disclosure may be formulated for the treatment of any suitable plant type, including, but not limited to, row crops and vegetables. In some embodiments, inoculant compositions of the present disclosure are formulated for the treatment of one or more plants selected from Amaranthaceae (e.g., chard, spinach, sugar beet, quinoa), Asteraceae (e.g., artichoke, asters, chamomile, chicory, chrysanthemums, dahlias, daisies, echinacea, goldenrod, guarayule, lettuce, marigolds, safflower, sunflowers, zinnias), Brassicaceae (e.g., arugula, broccoli, bok choy, Brussels sprouts, cabbage, cauliflower, canola, collard greens, daikon, garden cress, horseradish, kale, mustard, radish, rapeseed, rutabaga, turnip, wasabi, watercress, *Arabidopsis thaliana*), Cucurbitaceae (e.g., cantaloupe, cucumber, honeydew, melon, pumpkin, squash (e.g., acorn squash, butternut squash, summer squash), watermelon, zucchini), Fabaceae (e.g., alfalfa, beans, carob, clover, guar, lentils, mesquite, peas, peanuts, soybeans,

tamarind, tragacanth, vetch), Malvaceae (e.g., cacao, cotton, durian, hibiscus, kenaf, kola, okra), Poaceae (e.g., bamboo, barley, corn, fonio, lawn grass (e.g., Bahia grass, Bermudagrass, bluegrass, Buffalograss, Centipede grass, Fescue, or Zoysia), millet, oats, ornamental grasses, rice, rye, sorghum, sugar cane, triticale, wheat), Polygonaceae (e.g., buckwheat), Rosaceae (e.g., almonds, apples, apricots, blackberry, blueberry, cherries, peaches, plums, quinces, raspberries, roses, strawberries), Solanaceae (e.g., bell peppers, chili peppers, eggplant, petunia, potato, tobacco, tomato) and Vitaceae (e.g., grape). In some embodiments, inoculant compositions of the present disclosure are formulated for the treatment of one or more fungicide-, herbicide-, insecticide- and/or nematocide-resistant plants (e.g., one or more plants resistant to acetolactate synthase inhibitors).

Non-limiting examples of plants that may be treated with inoculant compositions of the present disclosure include plants sold by Monsanto Company (St. Louis, MO) under the BOLLGARD II®, DROUGHTGARD®, GENUITY®, RIB COMPLETE®, ROUNDUP READY®, ROUNDUP READY 2 YIELD®, ROUNDUP READY 2 EXTEND™, SMARTSTAX®, VT DOUBLE PRO®, VT TRIPLE PRO®, YIELDGARD®, YIELDGARD VT ROOTWORM/RR2®, YIELDGARD VT TRIPLE® and/or XTENDFLEX™ tradenames.

The present disclosure extends to plants that have been treated with inoculant compositions of the present disclosure, plant parts harvested from plants that have been treated with inoculant compositions of the present disclosure, processed products derived from plants that have been treated with inoculant compositions of the present disclosure and crops comprising a plurality of plants that have been treated with inoculant compositions of the present disclosure.

The present disclosure also provides coated plant propagation materials comprising, consisting essentially of, or consisting of a plant propagation material and a coating that covers at least a portion of the outer surface of the plant propagation material, said coating comprising, consisting essentially of, or consisting of an inoculant composition of the present disclosure.

The coating may comprise one, two, three, four, five or more layers. In some embodiments, the coating comprises an inner layer that contains one or more microorganisms and one or more outer layers free or substantially free of microorganisms.

In some embodiments, the inner layer of the coating is an inoculant composition of the present disclosure and the outer layer is equivalent to an inoculant composition of the present disclosure but it lacks one or more microorganisms. For example, coated plant propagation materials of the present disclosure may comprise a seed that is coated with an inner layer that comprises microbial spores (e.g., spores of one or more strains of *Bacillus*, one or more strains of *Gliocladium*, one or more strains of *Glomus*, one or more strains of *Metarhizium*, one or more strains of *Penicillium* and/or one or more strains of *Trichoderma*), one or more protectants and one or more dispersants in a non-aqueous carrier and an outer layer that comprises the same carrier but is free of microbial spores.

In some embodiments, the coating comprises, consists essentially of, or consists of an inoculant composition of the present disclosure and a drying powder. For example, coated plant propagation materials of the present disclosure may comprise a seed that is coated with an inoculant composition comprising microbial spores (e.g., spores of one or more strains of *Bacillus*, one or more strains of *Gliocladium*, one or more strains of *Glomus*, one or more strains of *Metarhizium*, one or more strains of *Penicillium* and/or one or more strains of *Trichoderma*), one or more protectants and one or more dispersants in a non-aqueous carrier and is then covered with a drying powder (e.g., a drying powder that comprises calcium stearate, one or more clays, graphite, magnesium stearate, magnesium sulfate, powdered milk, silica, soy lecithin and/or talc).

Inoculant compositions of the present disclosure may be coated on plant propagation material in any suitable amount(s)/concentration(s).

In some embodiments, the inoculant composition is applied in an amount ranging from about 0.5 to about 10 milliliters of inoculant composition per kilogram of plant propagation material. For example, in some embodiments, about 0.5, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10 milliliters or more of inoculant composition is applied per kilogram of seed. In some embodiments, an inoculant composition comprising one or more maltodextrins (e.g., one or more maltodextrins having a DEV of about 15 to about 20) and one or more microbial spores (e.g., spores of one or more strains of *Bacillus*, one or more strains of *Gliocladium*, one or more strains of *Glomus*, one or more strains of *Metarhizium*, one or more strains of *Penicillium* and/or one or more strains of *Trichoderma*) is applied at a rate of about 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, or 4 milliliters per kilogram of seed.

Drying powders may be applied in any suitable amount(s)/concentration(s).

In some embodiments, the drying powder is applied in an amount ranging from about 0.5 to about 10 grams of drying powder per kilogram of plant propagation material. For example, in some embodiments, about 0.5, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10 grams or more of drying powder (e.g., drying powder comprising magnesium stearate, magnesium sulfate, powdered milk, silica, soy lecithin and/or talc) is applied per kilogram of seed. In some embodiments, a drying powder comprising calcium stearate, attapulgite clay, montmorillonite clay, graphite, magnesium stearate, silica (e.g., fumed silica, hydrophobically-coated silica and/or precipitated silica) and/or talc is applied to seeds coated with an inoculant composition comprising one or more maltodextrins (e.g., one or more maltodextrins having a DEV of about 15 to about 20) and one or more microbial spores (e.g., spores of one or more strains of *Bacillus*, one or more strains of *Gliocladium*, one or more strains of *Glomus*, one or more strains of *Metarhizium*, one or more strains of *Penicillium* and/or one or more strains of *Trichoderma*) at a rate of about 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, or 3 grams per kilogram of seed.

In some embodiments, the coating covers about 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, 99, 99.5% or more of the outer surface of the plant propagation material.

In some embodiments, the average thickness of the coating is at least 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5 μm or more. In some embodiments, the average thickness of the coating is about 1.5 to about 3.0 μm .

In some embodiments, the coating comprises one or more effect pigments. Effect pigments, which are sometimes also referred to in the art as "pearl pigments," are a class of materials that provide reflectivity, shine, and/or a pearlescent effect when applied as a coating. In some instances, the effect pigment is in the form of a powder comprising a substrate material and a metal oxide coating. For example, the effect pigment may comprise a substrate material including but not limited to talc, silicate materials (e.g., mica), clay minerals, calcium carbonate, kaolin, phlogopite, alumina, and similar substances. In some instances, the substrate material comprises a hydrophilic material. The substrate material may be coated with a semi-transparent layer of a metal oxide, including but not limited to titanium dioxide, iron oxide, chromium oxide, or zirconium oxide. Alternatively, in some instances, the effect pigment comprises metal powder or metal flakes. The metal powder or metal flakes may comprise a metal including, but not limited to aluminum, copper, silver, or bronze. In some instances, the effect pigment comprises a silicate based substrate. Non-limiting examples of particulate silicates that can be incorporated into the dry powder coating include mica coated with titanium dioxide (e.g., SUNMICA FINE WHITE 2800102, which is commercially available from Sun Chemical Corp.). Other non-limiting examples of commercially available effect pigments that can be incorporated into the dry powder include MAGNA PEARL, LUMINA and MEARLIN

pigments from BASF Corporation; PHIBRO PEARL from PhibroChem; and IRIDESIUM 120 from Aakash Chemicals. In some instances, the dry powder has a mean particle size of from about 1 to about 25 microns.

Inoculant compositions of the present disclosure may be used to coat any suitable plant propagation materials, including, but not limited to, cuttings (e.g., leaves, stems), rhizomes, seeds and tubers. In some embodiments, the plant propagation material is a seed.

Inoculant compositions of the present disclosure may be used to coat plant propagation materials of any suitable plant type, including, but not limited to, row crops and vegetables. In some embodiments, inoculant compositions of the present disclosure are coated on propagation material derived from one or more plants selected from Amaranthaceae (e.g., chard, spinach, sugar beet, quinoa), Asteraceae (e.g., artichoke, asters, chamomile, chicory, chrysanthemums, dahlias, daisies, echinacea, goldenrod, guayule, lettuce, marigolds, safflower, sunflowers, zinnias), Brassicaceae (e.g., arugula, broccoli, bok choy, Brussels sprouts, cabbage, cauliflower, canola, collard greens, daikon, garden cress, horseradish, kale, mustard, radish, rapeseed, rutabaga, turnip, wasabi, watercress, *Arabidopsis thaliana*), Cucurbitaceae (e.g., cantaloupe, cucumber, honeydew, melon, pumpkin, squash (e.g., acorn squash, butternut squash, summer squash), watermelon, zucchini), Fabaceae (e.g., alfalfa, beans, carob, clover, guar, lentils, mesquite, peas, peanuts, soybeans, tamarind, tragacanth, vetch), Malvaceae (e.g., cacao, cotton, durian, hibiscus, kenaf, kola, okra), Poaceae (e.g., bamboo, barley, corn, fonio, lawn grass (e.g., Bahia grass, Bermudagrass, bluegrass, Buffalograss, Centipede grass, Fescue, or Zoysia), millet, oats, ornamental grasses, rice, rye, sorghum, sugar cane, triticale, wheat), Polygonaceae (e.g., buckwheat), Rosaceae (e.g., almonds, apples, apricots, blackberry, blueberry, cherries, peaches, plums, quinces, raspberries, roses, strawberries), Solanaceae (e.g., bell peppers, chili peppers, eggplant, petunia, potato, tobacco, tomato) and Vitaceae (e.g., grape).

Non-limiting examples of plant propagation materials that may be coated with inoculant compositions of the present disclosure include seeds sold by Monsanto Company (St. Louis, MO) under the BOLLGARD II®, DROUGHTGARD®, GENUITY®, RIB COMPLETE®, ROUNDUP READY®, ROUNDUP READY 2 YIELD®, ROUNDUP READY 2 EXTEND™, SMARTSTAX®, VT DOUBLE PRO®, VT TRIPLE PRO®, YIELDGARD®, YIELDGARD VT ROOTWORM/RR2®, YIELDGARD VT TRIPLE® and/or XTENDFLEX™ tradenames.

The present disclosure extends to plants that grow from coated plant propagation materials of the present disclosure, plant parts harvested from plants that grow from coated plant propagation materials of the present disclosure, processed products derived from plants that grow from coated plant propagation materials of the present disclosure and crops comprising a plurality of plants that grow from coated plant propagation materials of the present disclosure.

The present disclosure also provides kits comprising, consisting essentially of, or consisting of a coated plant propagation material of the present disclosure and a container housing the coated plant propagation material. In some embodiments, the kit further comprises one or more oxygen scavengers, such as activated carbon, ascorbic acid, iron powder, mixtures of ferrous carbonate and metal halide catalysts, sodium chloride and/or sodium hydrogen carbonate.

The container may comprise any suitable material(s), including, but not limited to, materials that reduce the amount of light, moisture and/or oxygen that contact the coated plant propagation material when the container is sealed.

In some embodiments, the container comprises, consists essentially of, or consists of a material having light permeability of less than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70 or 75%.

In some embodiments, the container reduces the amount of ambient light that reaches said coated plant propagation material by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% when sealed.

In some embodiments, the container reduces the amount of ambient moisture that reaches said plant propagation material by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% when sealed.

In some embodiments, the container comprises, consists essentially of, or consists of a material having an oxygen transmission rate of less than about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, or 500 cm³/m²·day (as measured in accordance with ASTM D3985).

In some embodiments, the container reduces the amount of ambient oxygen that reaches said plant propagation material by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% when sealed.

In some embodiments, kits of the present disclosure comprise 1, 2, 3, 4, 5 or more additional containers. The additional containers may comprise any suitable component(s) or composition(s), including, but not limited to, agriculturally beneficial microorganisms, biostimulants, drying agents, nutrients, oxidation control components and pesticides. Examples of agriculturally beneficial microorganisms, biostimulants, drying agents, nutrients, oxidation control components and pesticides that may be included in the additional containers are described above.

The present disclosure also provides animal feed compositions comprising, consisting essentially of, or consisting of a food component and a microbial component, said microbial component comprising, consisting essentially of, or consisting of an inoculant composition of the present disclosure.

Animal feed compositions of the present disclosure may comprise any suitable food component, including, but not limited to, fodder (e.g., grains, hay, legumes, silage and/or straw) and forage (e.g., grass).

Animal feed compositions of the present disclosure may be fed to any suitable animal, including, but not limited to, farm animals, zoo animals, laboratory animals and/or companion animals. In some embodiments, the animal feed composition is formulated to meet the dietary needs of birds (e.g., chickens, ducks, quails and/or turkeys), bovids (e.g., antelopes, bison, cattle, gazelles, goats, impala, oxen, sheep and/or wildebeests), canines, cervids (e.g., caribou, deer, elk and/or moose), equines (e.g., donkeys, horses and/or zebras), felines, fish, pigs, rabbits, rodents (e.g., guinea pigs, hamsters, mice and/or rats) and the like.

The present disclosure also provides methods comprising, consisting essentially of, or consisting of applying an inoculant composition of the present disclosure to a plant or plant part.

Inoculant compositions of the present disclosure may be applied in any suitable manner, including, but not limited to, on-seed application, in-furrow application, soil application and foliar application.

In some embodiments, inoculant compositions of the present disclosure are applied to plant propagation materials (e.g., seeds) at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more prior to planting.

In some embodiments, inoculant compositions of the present disclosure are applied to plant propagation materials (e.g., seeds) at the time of planting.

Plant propagation materials may be treated using any suitable method(s), including, but not limited to, coating, dripping, spraying and soaking. Batch systems, in which predetermined batch sizes of material and inoculant composition are delivered into a mixer, may be employed. Continuous treatment systems, which are calibrated to apply inoculant composition at a predefined rate in proportion to a continuous flow of material, may also be employed.

In some embodiments, plant propagation materials are soaked for at least 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.25, 1.5, 1.75, 2, 3, 4, 5, 6, 9, 12, 15, 18, 21, 24, 36, 48 hours or more in an inoculant composition of the present disclosure.

In some embodiments, plant propagation materials (e.g., seeds) are coated by applying an inoculant composition of the present disclosure to the inside wall of a round container, adding the seeds, then rotating the container such that the seeds come into contact with the composition, a process known in the art as "container coating".

In some embodiments, an inoculant composition of the present disclosure is freeze- spray- or spray-freeze-dried and then applied to plant propagation material. For examples, in some embodiments, an inoculant composition comprising one or more maltodextrins (e.g., one or more maltodextrins having a DEV of about 15 to about 20), one or more microbial spores (e.g., spores of one or more strains of *Bacillus*, one or more strains of *Gliocladium*, one or more strains of *Glomus*, one or more strains of *Metarhizium*, one or more strains of *Penicillium* and/or one or more strains of *Trichoderma*) and one or more disaccharides (e.g., maltose) is freeze- spray- or spray-freeze-dried, mixed with a drying powder (e.g., a drying powder comprising calcium stearate, attapulgite clay, montmorillonite clay, graphite, magnesium stearate, silica (e.g., fumed silica, hydrophobically-coated silica and/or precipitated silica) and/or talc), then coated on seed that was been pre-treated with one or more adhesives (e.g., an adhesive composition comprising one or more maltodextrins, one or more mono-, di- or oligosaccharides, one or more peptones, etc.), one or more pesticides and/or one or more plant signal molecules (e.g., one or more LCOs).

The present disclosure also provides methods comprising, consisting essentially of, or consisting of planting a coated seed of the present disclosure.

The present disclosure also provides methods of enhancing the survival and/or stability of microbial spores in a composition, said methods comprising, consisting essentially of, or consisting of adding an effective amount of one or more protectants to said composition.

Protectants may be used to improve any suitable microbial stability characteristic(s) of the microbial spores in a composition, including, but not limited to, the ability of the microbial spores in a composition to enhance plant yield after being coated on a seed and stored for a defined period of time prior to planting the seed. For example, the addition of one or more protectants to a composition enhances the ability of the microbial spores therein to propagate and increase yield after being coated on a plant propagation material (e.g., seed) and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the addition of one or more protectants to a composition improves the stability of one or more microbial spores therein by at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to microbial spores in a control composition (e.g., a control composition that is identical to the inoculant composition of the present disclosure except that it lacks one or more of the components found in the inoculant composition and/or contains one or more components not found in the inoculant composition of the present disclosure). For example, the addition of one or more protectants to a composition may improve one or more microbial stability characteristics of one or more microbial spores therein by at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition that is identical to the inoculant composition except that lacks one or more of the protectants found in the inoculant composition and/or comprises a reduced amount of one or more of the protectants found in the inoculant composition.

In some embodiments, the addition of one or more protectants to a composition improves the stability of one or more microbial spores therein by at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition (e.g., a control composition that is identical to the inoculant composition of the present disclosure except that it lacks one or more of the components found in the inoculant composition and/or contains one or more components not found in the inoculant composition of the present disclosure). For example, the addition of one or more protectants to a composition may improve the survival rate of one or more of the microbial spores contained therein by at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 150, 175, 200% or more as compared to a control composition that is identical to the inoculant composition except that lacks one or more of the protectants found in the inoculant composition and/or comprises a reduced amount of one or more of the protectants found in the inoculant composition.

In some embodiments, the addition of one or more protectants to a composition improves the survival of one or more of the microbial spores in an inoculant composition to the extent that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of the microbial spores survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, the addition of one or more protectants to a composition improves the survival of one or more of the microbial spores in an inoculant composition to the extent that at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70% or more of the microbial spores survive when the inoculant composition is coated on a seed, dried and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the addition of one or more protectants to a composition improves the survival of one or more of the microbial spores in an inoculant composition to the extent that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/seed or more of the microbial spores survive when the inoculant composition is coated on a seed and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, the addition of one or more protectants to a composition improves the survival of one or more of the microbial spores in an inoculant composition to the extent that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 cfu/seed or more of the microbial spores survive when the inoculant composition is coated on a seed and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the addition of one or more maltodextrins to a composition improves both the survival rate and one, two, three, four, five, six, seven, eight, nine, ten or more microbial stability characteristics of the microbial spore(s) contained therein.

Any suitable protectant(s) may be added to the composition, including, but not limited to maltodextrins and oxidation control components.

In some embodiments, one or more maltodextrins is added to the inoculant composition.

Any suitable maltodextrin(s) may be added to the composition, including, but not limited to, maltodextrins having a dextrose equivalent value (DEV) of about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25. In some embodiments, one or more maltodextrins having a DEV of about 5 to about 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, about 10 to about 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or about 15 to about 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 is added to the composition. In some embodiments, a combination of maltodextrins having a DEV of about 5 to about 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, about 10 to about 11, 12, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25, or about 15 to about 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 is added to the composition. In some embodiments, one or more maltodextrins having a DEV of about 10 to about 25 (e.g., one or more maltodextrins having a DEV of about 15 to about 20) is added to the composition. In some embodiments, a combination of maltodextrins having a DEV of about 10 to about 25 (e.g., a combination of maltodextrins having a DEV of about 15 to about 20) is added to the composition.

Non-limiting examples of maltodextrins that may be useful in methods of the present disclosure include MALTRIN® M150 (DEV = 15; molecular weight = 1200; Grain Processing Corporation, Muscatine, IA), MALTRIN® M200 (DEV = 20; molecular weight = 900; Grain Processing Corporation, Muscatine, IA), MALTRIN® M250 (DEV = 25; molecular weight = 720; Grain Processing Corporation, Muscatine, IA); MALTRIN QD® M580 (DEV = 16.5-19.9; Grain Processing Corporation, Muscatine, IA); MALTRIN QD® M585 (DEV = 15.0-19.9; Grain Processing Corporation, Muscatine, IA); MALTRIN QD® M600 (DEV = 20.0-23.0; Grain Processing Corporation, Muscatine, IA); GLOBE® Plus 15 DE (Ingredion Inc., Westchester, IL) and combinations thereof. In some embodiments, the maltodextrin (or combination of maltodextrins) has a DEV of 15 to 20.

Maltodextrins may be added to the composition in any suitable form. In some embodiments, the maltodextrin(s) added to the composition is/are at least 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 99.5% or more pure.

The absolute value of the amount/concentration/dosage of maltodextrins that is sufficient to cause the desired effect may be affected by factors such as the type, size and volume of material to which the composition will be applied, the stability of the microbial spores in the composition, the identity and amounts/concentrations of other components in the inoculant composition (e.g., monosaccharides, disaccharides, sugar alcohols) and the storage conditions (e.g., temperature, relative humidity, duration). Those skilled in the art will understand how to select an effective amount/concentration/dosage using routine dose-response experiments.

In some embodiments, the maltodextrin(s) is/are added to the composition until they comprise about 5 to about 95 % or more (by weight) of the composition. For example, one or more maltodextrins may be added until the composition comprises about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of one or more maltodextrins (each and/or collectively) having a DEV value of about 15 to about 20. In some embodiments, the maltodextrin amount/concentration is about 1 to about 65%, about 10% to about 30%, about 20% to about 40%, about 20% to about 50%, or about 30 to about 60% (by weight) of the composition.

In some embodiments, the maltodextrin(s) is/are added to the composition in an amount/concentration sufficient to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of the microbial spores therein survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a

period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, the maltodextrin(s) is/are added to the composition in an amount/concentration sufficient to ensure that at least about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70% or more of the microbial spores therein survive when the inoculant composition is coated on a seed, dried and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the maltodextrin(s) is/are added to the composition in an amount/concentration sufficient to ensure that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/seed or more of the microbial spores therein survive when the inoculant composition is coated on a seed and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, the maltodextrin(s) is/are added to the composition in an amount/concentration sufficient to ensure that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 cfu/seed or more of the microbial spores therein survive when the inoculant composition is coated on a seed and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, one or more oxidation control components is added to the inoculant composition.

Any suitable oxidation control component(s) may be added to the composition, including, but not limited to, antioxidants and/or oxygen scavengers. In some embodiments, ascorbic acid, ascorbyl palmitate, ascorbyl stearate, calcium ascorbate, carotenoids, glutathione, lipoic acid, phenolic compounds (e.g., flavonoids, flavones, flavonols), potassium ascorbate, sodium ascorbate, sodium hydrogen carbonate, thiols (e.g., glutathione, lipoic acid, N-acetyl cysteine), tocopherols, tocotrienols, ubiquinone and/or uric acid is added to the composition. In some embodiments, ascorbic acid is added to the composition.

The absolute value of the amount/concentration/dosage of oxidation control components that is sufficient to cause the desired effect may be affected by factors such as the type, size and volume of material to which the composition will be applied, the stability of the microbial spores in the composition and the storage conditions (e.g., temperature, relative humidity, duration). Those skilled in the art will understand how to select an effective amount/concentration/dosage using routine dose-response experiments.

In some embodiments, the oxidation control component(s) is/are added to the composition in an amount/concentration ranging from about 0.0001 to about 5 % or more (by weight) of the composition. For example, one or more oxidation control components may be added until it/they comprise(s) about 0.0005, 0.001, 0.002, 0.003, 0.004, 0.005, 0.0075, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.75, 1, 1.25, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5% or more of the composition.

In some embodiments, the oxidation control component(s) is/are added to the composition at a concentration ranging from about 1×10^{-20} M to about 1×10^{-1} M. For example, one or more oxidation control components may be added until the composition comprises about 1×10^{-20} M, 1×10^{-19} M, 1×10^{-18} M, 1×10^{-17} M, 1×10^{-16} M, 1×10^{-15} M, 1×10^{-14} M, 1×10^{-13} M, 1×10^{-12} M, 1×10^{-11} M, 1×10^{-10} M, 1×10^{-9} M, 1×10^{-8} M, 1×10^{-7} M, 1×10^{-6} M, 1×10^{-5} M, 1×10^{-4} M, 1×10^{-3} M, 1×10^{-2} M, 1×10^{-1} M of one or more oxidation control components.

In some embodiments, the oxidation control component (s) is/are added to the composition in an amount/concentration sufficient to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of the microbial spores therein survive when the inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, the oxidation control component (s) is/are added to the composition in an amount/concentration sufficient to ensure that at least about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70% or more of the microbial spores therein survive when the inoculant composition is coated on a seed, dried and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the oxidation control component (s) is/are added to the composition in an amount/concentration sufficient to ensure that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/seed or more of the microbial spores therein survive when the inoculant composition is coated on a seed and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, the oxidation control component (s) is/are added to the composition in an amount/concentration sufficient to ensure that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 cfu/seed or more of the microbial spores therein survive when the inoculant composition is coated on a seed and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, one or more hygroscopic polymers is added to the inoculant composition. For example, in some embodiments, albumin, alginate, cellulose, gums (e.g., cellulose gum, guar gum, gum arabic, gum combretum, xanthan gum), methyl cellulose, nylons, pectin, polyacrylic acid, polycarbonates, polyethylene glycol (PEG), polyethylenimine (PEI), polylactides, polymethylacrylate (PMA), polyurethanes, polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), propylene glycol, sodium carboxymethyl cellulose and/or starch is added to the inoculant composition.

Non-limiting examples of hygroscopic polymers that may be useful in methods of the present disclosure include AGRIMER™ polymers (e.g., 30, AL-10 LC, AL-22, AT/ATF, VA 3E, VA 31, VA 5E, VA 51, VA 6, VA 6E, VA 7E, VA 71, VEMA AN-216, VEMA AN-990, VEMA AN-1200, VEMA AN-1980, VEMA H-815MS; Ashland Specialty Ingredients, Wilmington, DE), EASYSERSE™ polymers (Ashland Specialty Ingredients, Wilmington, DE); DISCO™ AG polymers (e.g., L-250, L-280, L-285, L-286, L-320, L-323, L-517, L-519, L-520, L800; Incotec Inc., Salinas, CA), KELZAN® polymers (Bri-Chem Supply Ltd., Calgary, Alberta, CA), SEEDWORX™ polymers (e.g., Bio 200; Aginnovation, LLC, Walnut Grove, CA), TABULOSE® gels (e.g., SC-580, SC-612, SC-613, SC-681; Blanver Fermoquimica, Boca Raton, FL), TICAXAN® xanthan powders (TIC Gums, White Marsh, MD) and combinations thereof.

Additional examples of hygroscopic polymers that may be added to inoculant compositions of the present disclosure may be found in Pouci, et al. AM. J. AGRIC. BIOL. SCI. 3(1):299 (2008).

The absolute value of the amount/concentration/dosage of maltodextrins that is sufficient to cause the desired effect may be affected by factors such as the type, size and volume of material to which the composition will be applied, the stability of the microbial spores in the composition, the identity and amounts/concentrations of other components in the inoculant composition (e.g., monosaccharides, disaccharides, sugar alcohols) and the storage conditions (e.g., temperature, relative humidity, duration). Those skilled in the art will understand how to select an effective amount/concentration/dosage using routine dose-response experiments.

Hygroscopic polymers may be added to the inoculant composition in any suitable amount(s)/concentration(s). In some embodiments, the hygroscopic polymer(s) is/are added to the composition until it/they comprise(s) about 0.1 to about 25 % (by weight) of the inoculant composition. For example, inoculant compositions of the present disclosure may comprise about 0.1, 0.2, 0.3, 0.4, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 15, 20, 25% or more (by weight) of one or more hygroscopic polymers. In some embodiments, the hygroscopic polymer amount/concentration is about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1 to about 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5 or 10 % (by weight) of the inoculant composition.

In some embodiments, the hygroscopic polymer(s) is/are added to the composition in an amount/concentration sufficient to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following desiccation (of about 0, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more) and storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the hygroscopic polymer(s) is/are added to the composition in an amount/concentration sufficient to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65% or more of the microbial spores in the inoculant composition survive following storage at 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

In some embodiments, the hygroscopic polymer (s) is/are added to the composition in an amount/concentration sufficient to ensure that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} cfu/seed or more of the microbial spores therein survive when the inoculant composition is coated on a seed and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more. In some embodiments, the hygroscopic polymer (s) is/are added to the composition in an amount/concentration sufficient to ensure that at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 cfu/seed or more of the microbial spores therein survive when the inoculant composition is coated on a seed and stored at 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 and/or 30°C and 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

The present disclosure also provides systems and methods of using inoculant compositions of the present disclosure in conjunction with additional compositions comprising one or more agriculturally beneficial constituents. The additional composition(s) may comprise any suitable agriculturally beneficial constituent(s), including, but not limited to, the agriculturally beneficial constituents described above.

In some embodiments, inoculant compositions of the present disclosure are used in conjunction with one or more on-seed compositions, one or more in-furrow compositions, one or more soil-applied compositions and/or one or more foliar-applied compositions.

In some embodiments, inoculant compositions of the present disclosure are used as part of an integrated disease and/or pest management system.

Particular embodiments of the present disclosure are described in the following numbered paragraphs:

1. An inoculant composition, comprising, consisting essentially of, or consisting of:
microbial spores; and
a non-aqueous carrier.
2. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise about 0.1% to about 30% (by weight) of said inoculant composition, optionally about 5 to about 15% (by weight) of said composition, optionally about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25% (by weight) of said inoculant composition.
3. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores are present in said inoculant composition in a concentration ranging from about 1×10^1 to about 1×10^{20} colony-forming units per gram and/or milliliter of said inoculant composition, optionally 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} , 1×10^{13} , 1×10^{14} , 1×10^{15} or more colony-forming units per gram and/or milliliter of said inoculant composition.
4. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more diazotrophic microorganisms.
5. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains of *Bacillus*, optionally one or more strains of *Bacillus circulans*, *Bacillus licheniformis*, *Bacillus macerans*, *Bacillus megatarium*, *Bacillus polymyxa* and/or *Bacillus pumilus*.
6. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more phosphate-solubilizing microorganisms.
7. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains of *Penicillium*, optionally one or more strains of *P. bilaiae* and/or *P. gaestrivorus*.

8. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains of *Trichoderma*, optionally one or more strains of *T. asperellum*, *T. atroviride*, *T. fertile*, *T. gamsii*, *T. hamatum*, *T. harzianum*, *T. reesi*, *T. virens* and/or *T. viridae*.

9. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more mycorrhizal fungi.

10. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains of *Gliocladium*, optionally one or more strains of *Gliocladium virens*, one or more strains of *Glomus*, optionally one or more strains of *Glomus intraradices*, and/or one or more strains of *Metarhizium*, optionally, one or more strains of *Metarhizium anisopliae*.

11. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains of *Bacillus*, optionally *Bacillus amyloliquefaciens* D747, *Bacillus amyloliquefaciens* NRRL B-50349, *Bacillus amyloliquefaciens* TJ1000, *Bacillus amyloliquefaciens* FZB24, *Bacillus amyloliquefaciens* FZB42, *Bacillus amyloliquefaciens* IN937a, *Bacillus amyloliquefaciens* IT-45, *Bacillus amyloliquefaciens* TJ1000, *Bacillus amyloliquefaciens* MBI600, *Bacillus amyloliquefaciens* BS27 (deposited as NRRL B-5015), *Bacillus amyloliquefaciens* BS2084 (deposited as NRRL B-50013), *Bacillus amyloliquefaciens* 15AP4 (deposited as ATCC PTA-6507), *Bacillus amyloliquefaciens* 3AP4 (deposited as ATCC PTA-6506), *Bacillus amyloliquefaciens* LSSA01 (deposited as NRRL B-50104), *Bacillus amyloliquefaciens* ABP278 (deposited as NRRL B-50634), *Bacillus amyloliquefaciens* 1013 (deposited as NRRL B-50509), *Bacillus amyloliquefaciens* 918 (deposited as NRRL B-50508), *Bacillus amyloliquefaciens* 22CP1 (deposited as ATCC PTA-6508) and *Bacillus amyloliquefaciens* BS18 (deposited as NRRL B-50633), *Bacillus cereus* 1-1562, *Bacillus firmus* 1-1582, *Bacillus lichenformis* BA842 (deposited as NRRL B-50516), *Bacillus lichenformis* BL21 (deposited as NRRL B-50134), *Bacillus mycoides* NRRL B-21664, *Bacillus pumilus* NRRL B-21662, *Bacillus pumilus* NRRL B-30087, *Bacillus pumilus* ATCC 55608, *Bacillus pumilus* ATCC 55609, *Bacillus pumilus* GB34, *Bacillus pumilus* KFP9F, *Bacillus pumilus* QST 2808, *Bacillus subtilis* ATCC 55078, *Bacillus subtilis* ATCC 55079, *Bacillus subtilis* MBI 600, *Bacillus subtilis* NRRL B-21661, *Bacillus subtilis* NRRL B-21665, *Bacillus subtilis* CX-9060, *Bacillus subtilis* GB03, *Bacillus subtilis* GB07, *Bacillus subtilis* QST-713, *Bacillus subtilis* FZB24, *Bacillus subtilis* D747, *Bacillus subtilis* 3BP5 (deposited as NRRL B-50510), *Bacillus thuringiensis* ATCC 13367, *Bacillus thuringiensis* GC-91, *Bacillus thuringiensis* NRRL B-21619, *Bacillus thuringiensis* ABTS-1857, *Bacillus thuringiensis* SAN 401 I, *Bacillus thuringiensis* ABG-6305, *Bacillus thuringiensis* ABG-6346, *Bacillus thuringiensis* AM65-52, *Bacillus thuringiensis* SA-12, *Bacillus thuringiensis* SB4, *Bacillus thuringiensis* ABTS-35 1, *Bacillus thuringiensis* HD-1, *Bacillus thuringiensis* EG 2348, *Bacillus thuringiensis* EG 7826, *Bacillus thuringiensis* EG 7841, *Bacillus thuringiensis* DSM 2803, *Bacillus thuringiensis* NB-125 and/or *Bacillus thuringiensis* NB-176.

12. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains of *Gliocladium*, optionally *Gliocladium virens* ATCC 52045 and/or *Gliocladium virens* GL-21, one or more strains of *Glomus*, optionally *Glomus intraradices* RTI-801, one or more strains of *Metarhizium*, optionally *Metarhizium anisopliae* F52, *Penicillium*,

optionally *Penicillium bilaiae* ATCC 18309, *Penicillium bilaiae* ATCC 20851, *Penicillium bilaiae* ATCC 22348, *Penicillium bilaiae* NRRL 50162, *Penicillium bilaiae* NRRL 50169, *Penicillium bilaiae* NRRL 50776, *Penicillium bilaiae* NRRL 50777, *Penicillium bilaiae* NRRL 50778, *Penicillium bilaiae* NRRL 50777, *Penicillium bilaiae* NRRL 50778, *Penicillium bilaiae* NRRL 50779, *Penicillium bilaiae* NRRL 50780, *Penicillium bilaiae* NRRL 50781, *Penicillium bilaiae* NRRL 50782, *Penicillium bilaiae* NRRL 50783, *Penicillium bilaiae* NRRL 50784, *Penicillium bilaiae* NRRL 50785, *Penicillium bilaiae* NRRL 50786, *Penicillium bilaiae* NRRL 50787, *Penicillium bilaiae* NRRL 50788, *Penicillium bilaiae* RS7B-SD1, *Penicillium brevicompactum* AgRF18, *Penicillium canescens* ATCC 10419, *Penicillium expansum* ATCC 24692, *Penicillium expansum* YT02, *Penicillium fellatanum* ATCC 48694, *Penicillium gaestrivorus* NRRL 50170, *Penicillium glabrum* DAOM 239074, *Penicillium glabrum* CBS 229.28, *Penicillium janthinellum* ATCC 10455, *Penicillium lanosocoeruleum* ATCC 48919, *Penicillium radicum* ATCC 201836, *Penicillium radicum* FRR 4717, *Penicillium radicum* FRR 4719, *Penicillium radicum* N93/47267 and/or *Penicillium raistrickii* ATCC 10490, and/or one or more strains of *Trichoderma*, optionally *Trichoderma asperellum* SKT-1, *Trichoderma asperellum* ICC 012, *Trichoderma atroviride* LC52, *Trichoderma atroviride* CNCM 1-1237, *Trichodermafertile* JM41R, *Trichoderma gamsii* ICC 080, *Trichoderma hamatum* ATCC 52198, *Trichoderma harzianum* ATCC 52445, *Trichoderma harzianum* KRL-AG2, *Trichoderma harzianum* T-22, *Trichoderma harzianum* TH-35, *Trichoderma harzianum* T-39, *Trichoderma harzianum* ICC012, *Trichoderma reesi* ATCC 28217, *Trichoderma virens* ATCC 57678, *Trichoderma virens* GI-3, *Trichoderma virens* GL-21, *Trichoderma virens* G-41, *Trichoderma viridae* ATCC 52440, *Trichoderma viridae* ICC080 and/or *Trichoderma viridae* TV1.

13. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains having a genomic sequence that is at least 75, 80, 85, 90, 95, 96, 97, 98, 99% or more identical to any of the strains recited in paragraph 12 on the basis of 16S rDNA sequence identity.

14. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprise, consist essentially of, or consist of spores of one or more strains having a genomic sequence that is at least 75, 80, 85, 90, 95, 96, 97, 98, 99% or more identical to any of the strains recited in paragraph 13 on the basis of 16S rDNA, internal transcribed spacer (ITS) and/or cytochrome c oxidase (COI) sequence identity.

15. The inoculant composition of any one of the preceding paragraphs, wherein said microbial spores comprises, consists essentially of, or consists of spores of one or more biopesticides, optionally one or more biofungicides, one or more bioinsecticides and/or one or more bionematicides.

16. The inoculant composition of any one of the preceding paragraphs, wherein said non-aqueous carrier comprises about 70 to about 99% (by weight) of said inoculant composition, optionally about 75 to about 95% (by weight) of said composition, optionally about 80, 80.5, 81, 81.5, 82, 82.5, 83, 83.5, 84, 84.5, 85, 85.5, 86, 86.5, 87, 87.5, 88, 88.5, 89, 89.5 or 90% (by weight) of said inoculant composition.

17. The inoculant composition of any one of the preceding paragraphs, wherein said non-aqueous carrier comprises, consists essentially of or consists of a solid non-aqueous carrier.

18. The inoculant composition of any one of the preceding paragraphs, wherein said non-aqueous carrier comprises less than 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5% water by weight, based upon the total weight of the composition.

19. The inoculant composition of any one of the preceding paragraphs, wherein said non-aqueous carrier comprises no water.

20. The inoculant composition of any one of the preceding paragraphs, wherein said non-aqueous carrier comprise, consist essentially of or consists of a seed- and/or soil-compatible carrier.

21. The inoculant composition of any one of the preceding paragraphs, wherein said non-aqueous carrier comprises, consists essentially of, or consists of one or more maltodextrins having a dextrose equivalent value of about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25.

22. The inoculant composition of paragraph 1, wherein said non-aqueous carrier comprises, consists essentially of, or consists of one or more maltodextrins having a dextrose equivalent value of about 10 to about 25, optionally about 15 to about 20.

23. The inoculant composition of any one of paragraphs 21-22, wherein said one or more maltodextrins comprise about 5 to about 95 % (by weight), optionally about 50 to about 95%, about 55% to about 90%, about 60% to about 85%, about 65% to about 80%, or about 70 to about 80% (by weight), of said inoculant composition.

24. The inoculant composition of any one of paragraphs 21-23, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more.

25. The inoculant composition of any one of paragraphs 21-24, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more and stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

26. The inoculant composition of any one of paragraphs 21-25, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is coated on a plant propagation material and desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more.

27. The inoculant composition of any one of paragraphs 21-26, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is coated on a plant propagation material, desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more and stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

28. The inoculant composition of any one of paragraphs 21-27, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

29. The inoculant composition of any one of paragraphs 21-28, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is coated on a plant propagation material.

30. The inoculant composition of any one of paragraphs 21-29, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is coated on a plant propagation material and stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

31. The inoculant composition of any one of paragraphs 21-30, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units, of said one or more microbial spores per gram and/or milliliter of said inoculant composition remain viable when said inoculant composition is desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more.

32. The inoculant composition of any one of paragraphs 21-31, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units, of said one or more microbial spores per gram and/or milliliter of said inoculant composition remain viable when said inoculant composition is desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60,

65, 70, 75, 80, 85, 90, 95% or more and stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

33. The inoculant composition of any one of paragraphs 21-32, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units, of said one or more microbial spores per seed remain viable when said inoculant composition is coated on a seed and desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more.

34. The inoculant composition of any one of paragraphs 21-33, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units, of said one or more microbial spores per seed remain viable when said inoculant composition is coated on a seed, desiccated by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more and stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30°C and 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

35. The inoculant composition of any one of paragraphs 21-34, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units, of said one or more microbial spores per gram and/or milliliter of said inoculant composition remain viable when said inoculant composition is stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

36. The inoculant composition of any one of paragraphs 21-35, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units, of said one or more microbial spores per seed remain viable when said inoculant composition is coated on a seed.

37. The inoculant composition of any one of paragraphs 21-36, wherein said one or more maltodextrins is present in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units, of said one or more microbial spores per seed remain viable when said inoculant composition is coated on a seed and stored at 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27,

28, 29 or 30°C and 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

38. The inoculant composition of any one of the preceding paragraphs, wherein said non-aqueous carrier comprises one or more disaccharides, optionally maltose, trehalose, lactose, sucrose and/or cellobiose.

39. The inoculant composition of paragraph 38, wherein said one or more disaccharides comprise about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90% or more (by weight) of said inoculant composition.

40. The inoculant composition of paragraph 38, wherein said one or more disaccharides comprises about 5 to about 90% (by weight), optionally about 1 to about 20%, about 1% to about 15%, about 1% to about 15%, about 5% to about 15%, or about 5 to about 10% (by weight), of said inoculant composition.

41. The inoculant composition of paragraph 38, wherein said one or more disaccharides constitutes about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25% of the said inoculant composition (by weight).

42. The inoculant composition of any one of paragraphs 38 to 41, wherein said one or more maltodextrins and said one or more disaccharides collectively comprise about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more (by weight) of said inoculant composition.

43. The inoculant composition of any one of paragraphs 38 to 41, wherein said one or more maltodextrins and said one or more disaccharides collectively comprise about 5 to about 95% (by weight), optionally about 50 to about 99%, about 55% to about 95%, about 60% to about 90%, about 65% to about 90%, about 70% to about 90%, about 75% to about 90%, or about 80 to about 90% (by weight), of said inoculant composition.

44. The inoculant composition of any one of paragraphs 38 to 41, wherein said one or more maltodextrins and said one or more disaccharides are present in said inoculant composition in a maltodextrin:disaccharide ratio of about 5:95, 10:90, 15:85, 20:80, 25:75, 30:70, 35:65, 40:60, 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5 or more, optionally about 70:30, 75:25, 80:20, 85:15, 90:10, 95:5, 96:4, 97:3, 98:2, 99:1 or more.

45. The inoculant composition of any one of paragraphs 1-37, wherein said inoculant composition comprises one or more malt extracts.

46. The inoculant composition of paragraph 45, wherein said one or more malt extracts comprise about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90% or more (by weight) of said inoculant composition.

47. The inoculant composition of paragraph 45, wherein said one or more malt extracts comprises about 5 to about 90% (by weight), optionally about 1 to about 20%, about 1% to about 15%, about 1% to about 15%, about 5% to about 15%, or about 5 to about 10% (by weight), of said inoculant composition.

48. The inoculant composition of paragraph 45, wherein said one or more malt extracts constitutes about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25% of the said inoculant composition (by weight).

49. The inoculant composition of any one of paragraphs 45 to 48, wherein said one or more maltodextrins and said one or more malt extracts collectively comprise about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more (by weight) of said inoculant composition.

50. The inoculant composition of any one of paragraphs 45 to 48, wherein said one or more maltodextrins and said one or more malt extracts collectively comprise about 5 to about 95% (by weight), optionally about 50 to about 99%, about 55% to about 95%, about 60% to about 90%, about 65% to about 90%, about 70% to about 90%, about 75% to about 90%, or about 80 to about 90% (by weight), of said inoculant composition.

51. The inoculant composition of any one of paragraphs 45 to 48, wherein said one or more maltodextrins and said one or more malt extracts are present in said inoculant composition in a maltodextrin:disaccharide ratio of about 5:95, 10:90, 15:85, 20:80, 25:75, 30:70, 35:65, 40:60, 45:55, 50:50, 55:45, 60:40, 65:35, 70:30, 75:25, 80:20, 85:15, 90:10, 95:5 or more, optionally about 70:30, 75:25, 80:20, 85:15, 90:10, 95:5, 96:4, 97:3, 98:2, 99:1 or more.

52. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more dispersants.

53. The inoculant composition of paragraph 52, wherein said one or more dispersants comprise about 0.1 to about 5% (by weight) of said composition, optionally about 0.1 to about 2% (by weight) of said composition, optionally about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5 or 5% (by weight) of said composition.

54. The inoculant composition of any one of paragraphs 52-53, wherein said one or more dispersants comprises:

one or more anionic surfactants, optionally one or more alkyl carboxylates, alkyl sulfates, alkyl ether sulfates, alkyl amido ether sulfates, alkyl aryl polyether sulfates, alkyl aryl sulfates, alkyl aryl sulfonates, alkyl sulfonates, alkyl amide sulfonates, alkyl aryl sulfonates, alkyl benzene sulfonates, alkyl diphenyloxide sulfonate, alpha-olefin sulfonates, alkyl naphthalene sulfonates, paraffin sulfonates, alkyl sulfosuccinates, alkyl ether sulfosuccinates, alkylamide sulfosuccinates, alkyl sulfosuccinamates, alkyl sulfoacetates, alkyl phosphates, alkyl ether phosphates, acyl sarconsinates, acyl isethionates, N-acyl tartrates, N-acyl-N-alkyltaurates, benzene sulfonates, cumene sulfonates, dioctyl sodium sulfosuccinate, ethoxylated sulfosuccinates, lignin sulfonates, linear alkylbenzene sulfonates, monoglyceride sulfates, perfluorobutanesulfonate, perfluorooctanesulfonate, phosphate ester, styrene acrylic polymers, toluene sulfonates and/or xylene sulfonates

one or more cationic surfactants, optionally one or more alkyltrimethylammonium salts, cetylpyridinium chloride, benzalkonium chloride, benzethonium chloride, 5-Bromo-5-nitro-1,3-dioxane, dimethyldioctadecylammomum chloride, cetrimonium bromide, dioctadecyldimethylammonium bromide and/or octenidine dihydrochloride; and/or

one or more non-ionic surfactants, optionally one or more alcohol ethoxylates, alkanolamides, alkanolamine condensates, carboxylic acid esters, cetostearyl alcohol, cetyl alcohol, cocamide DEA, dodecyltrimethylamine oxides, ethanolamides, ethoxylates of glycerol ester and glycol esters, ethylene oxide polymers, ethylene oxide-propylene oxide copolymers, glucoside alkyl ethers, glycerol alkyl ethers, glycerol esters, glycol alkyl ethers, glycol alkylphenol ethers, glycol esters, monolaurin, pentaethylene glycol monododecyl ethers, poloxamer, polyamines, polyglycerol polyricinoleate, polysorbate, polyoxyethylenated fatty acids, polyoxyethylenated mercaptans, polyoxyethylenated polyoxypropylene glycols, polyoxyethylene glycol sorbitan alkyl esters, polyethylene glycol-polypropylene glycol copolymers, polyoxyethylene glycol octylphenol ethers, polyvinyl pyrrolidones, sugar-based alkyl polyglycosides, sulfoanilamides, sorbitan fatty acid alcohol ethoxylates, sorbitan fatty acid ester ethoxylates, sorbitan fatty acid ester and/or tertiary acetylenic glycols; and/or

one or more wetting agents, optionally one or more naphthalene sulfonates, optionally one or more alkyl naphthalene sulfonates, one or more isopropyl naphthalene sulfonates and/or one or more butyl naphthalene sulfonates.

55. The inoculant composition of any one of paragraphs 52-54, wherein said one or more dispersants comprises one or more polyoxyethylene alkyl ethers, one or more acrylic copolymers, one or more polyoxyethylene sorbitan trioleates and/or one or more secondary alcohol ethoxylates.

56. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more dust suppressants, optionally one or more adhesives, glycerin, mineral oils, paraffinic oils, vegetable oils and/or synthetic polymers.

57. The inoculant composition of paragraph 56, wherein said one or more dust suppressants comprises about 0.5 to about 15%, optionally about 1 to about 10%, about 1 to about 5%, about 2 to about 5%, about 2 to about 4% or about 2 to about 3 (by weight) of the inoculant composition.

58. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more protectants.

59. The inoculant composition of paragraph 58, wherein said one or more protectants comprise about 0.0001 to about 10% (by weight) of said composition, optionally about 2 to about 6% (by weight) of said composition, optionally about 0.0005, 0.001, 0.002, 0.003, 0.004, 0.005, 0.0075, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 7, 7.5, 8, 8.5, 9, 9.5 or 10% (by weight) of said composition.

60. The inoculant composition of any one of paragraphs 58-59, wherein said one or more protectants comprises one or more hygroscopic polymers, optionally one or more albumins, alginates, celluloses, gums (e.g., cellulose gum, guar gum, gum arabic, gum combretum, xanthan gum), methyl celluloses, nylons, pectins, polyacrylic acids, polycarbonates, polyethylene glycols (PEG), polyethylenimines (PEI), polylactides, polymethylacrylates (PMA), polyurethanes, polyvinyl alcohols (PVA), polyvinylpyrrolidones (PVP), propylene glycols, sodium carboxymethyl celluloses and/or starches.

61. The inoculant composition of any one of paragraphs 58-60, wherein said one or more protectants comprises one or more oxidation control components, optionally one or more antioxidants (e.g., ascorbic acid, ascorbyl palmitate, ascorbyl stearate, calcium ascorbate, one or more carotenoids, lipoic acid, one or more phenolic compounds (e.g., one or more flavonoids, flavones and/or flavonols), potassium ascorbate, sodium ascorbate, one or more thiols (e.g., glutathione, lipoic acid and/or N-acetyl cysteine), one or more tocopherols, one or more tocotrienols, ubiquinone and/or uric acid) and/or one or more oxygen scavengers, optionally ascorbic acid and/or sodium hydrogen carbonate.

62. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more pesticides, optionally:

one or more insecticides and/or nematicides, optionally one or more carbamates, diamides, macrocyclic lactones, neonicotinoids, organophosphates, phenylpyrazoles, pyrethrins, spinosyns, synthetic pyrethroids, tetrionic acids and/or tetramic acids;

one or more fungicides, optionally one or more aromatic hydrocarbons, benzimidazoles, benzthiadiazole, carboxamides, carboxylic acid amides, morpholines, phenylamides, phosphonates, quinone outside inhibitors, thiazolidines, thiophanates, thiophene carboxamides and/or triazoles; and/or

one or more herbicides, optionally one or more acetyl CoA carboxylase (ACCase) inhibitors, acetolactate synthase (ALS) inhibitors, acetohydroxy acid synthase (AHAS) inhibitors, photosystem II inhibitors, photosystem I inhibitors, protoporphyrinogen oxidase (PPO or Protox) inhibitors, carotenoid biosynthesis inhibitors, enolpyruvyl shikimate-3-phosphate (EPSP) synthase inhibitor, glutamine synthetase inhibitor, dihydropteroate synthetase inhibitor, mitosis inhibitors, 4-hydroxyphenyl-pyruvate-dioxygenase (4-HPPD) inhibitors, synthetic auxins, auxin herbicide salts, auxin transport inhibitors, and/or nucleic acid inhibitors.

63. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one more lipo-chitooligosaccharides, optionally one or more lipo-chitooligosaccharides represented by formulas I-IV.

64. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one more lipo-chitooligosaccharides, optionally one or more of the lipo-chitooligosaccharides represented by structures V-XXXIII.

65. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more chitooligosaccharides, optionally one or more chitooligosaccharides represented by formulas XXXIV-XXXV.

66. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more chitooligosaccharides, optionally one or more chitooligosaccharides represented by structures XXXVI-XXXIX.

67. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more chitinous compounds, optionally one or more chitins and/or one or more chitosans.

68. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more flavonoids, optionally:

one or more anthocyanidins, optionally cyanidin, delphinidin, malvidin, pelargonidin, peonidin and/or petunidin;

one or more anthoxanthins, optionally one or more flavones, such as apigenin, baicalein, chrysin, 7,8-dihydroxyflavone, diosmin, flavoxate, 6—hydroxyflavone, luteolin, scutellarein, tangeritin and/or wogonin; and/or flavonols, such as amurensin, astragalin, azaleatin, azalein, fisetin, furanoflavonols galangin, gossypetin, 3-hydroxyflavone, hyperoside, icariin, isoquercetin, kaempferide, kaempferitrin, kaempferol, isorhamnetin, morin, myricetin, myricitrin, natsudaicain, pachypodol, pyranoflavonols quercetin, quercitin, rhamnazin, rhamnetin, robinin, rutin, spiraeoside, troxerutin and/or zanthorhamnin;

one or more flavanones, optionally butin, eriodictyol, hesperetin, hesperidin, homoeriodictyol, isosakuranetin, naringenin, naringin, pinocembrin, poncirin, sakuranetin, sakuranin and/or sterubin;

one or more flavanonols, optionally dihydrokaempferol and/or taxifolin; flavans, such as flavan-3-ols (e.g., catechin (C), catechin 3-gallate (Cg), epicatechins (EC), epigallocatechin (EGC) epicatechin 3-gallate (ECg), epigallocatechin 3-gallate (EGCg), epiafzelechin, fisetinidol, gallocatechin (GC), gallcatechin 3-gallate (GCg), guibourtinidol, mesquitol, robinetinidol, theaflavin-3-gallate, theaflavin-3'-gallate, theflavin-3,3'-digallate, thearubigin), flavan-4-ols (e.g., apiforol and/or luteoforol) and/or flavan-3,4-diols (e.g., leucocyanidin, leucodelphinidin, leucofisetinidin, leucomalvidin, luecopelargonidin, leucoppeonidin, leucorobinetinidin, melacacidin and/or teracacidin); and/or

one or more isoflavonoids, optionally one or more isoflavones, such as biochanin A, daidzein, formononetin, genistein and/or glycitein; isoflavanes, such as equol, ionchocarpane and/or laxifloorane; isoflavandiols; isoflavenes, such as glabrene, haginin **D** and/or 2-methoxyjudaicin; coumestans, such as coumestrol, plicadin and/or wedelolactone; pterocarpans; and/or rotonoids; and/or

one or more neoflavonoids, optionally calophyllolide, coutareagenin, dalbergichromene, dalbergin and/or nivetin; and/or

one or more pterocarpans, optionally bitucarpin A, bitucarpin B, erybraedin A, erybraedin B, erythrabysin II, erythrabysin-1, erycristagallin, glycinol, glyceollidins, glyceollins, glycyrrhizol, maackiain, medica ϕ in, morisianine, orientanol, phaseolin, pisatin, striatine and/or trifolirhizin.

69. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises jasmonic acid and/or one or more derivatives thereof.

70. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises linoleic acid and/or one or more derivatives thereof.

71. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises linolenic acid and/or one or more derivatives thereof.

72. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more karrakins, optionally one or more karrakins represented by formula XXXX.

73. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises gluconolactone.
74. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more drying agents, optionally calcium stearate, one or more clays, graphite, magnesium stearate, magnesium sulfate, powdered milk, one or more silica powders, soy lecithin and/or talc.
75. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more biostimulants, optionally one or more seaweed extracts, one or more humic acids, one or more fulvic acids, myo-inositol and/or glycine.
76. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more microbial extracts, optionally one or more extracts from media comprising one or more diazotrophic, phosphosphate-solubilizing and/or biopesticidal microorganisms.
77. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more nutrients, optionally one or more vitamins (e.g., vitamin A, vitamin **B** complex (i.e., vitamin **Bi**, vitamin **B**₂, vitamin **B**₃, vitamin **B**₅, vitamin **B**₆, vitamin **B**₇, vitamin **B**₈, vitamin **B**₉, vitamin **B**₁₂, choline) vitamin C, vitamin D, vitamin E, vitamin K, carotenoids (α-carotene, β-carotene, cryptoxanthin, lutein, lycopene and/or zeaxanthin), macrominerals (e.g., calcium, iron, magnesium, nitrogen, phosphorous, potassium and/or sodium), trace minerals (e.g., boron, cobalt, chloride, chromium, copper, fluoride, iodine, iron, manganese, molybdenum, selenium and/or zinc) and/or organic acids (e.g., acetic acid, citric acid, lactic acid, malic acid and/or taurine).
78. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more growth media, optionally YEM media, mannitol yeast extract, glycerol yeast extract, Czapek-Dox media and/or potato dextrose broth.
79. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises one or more agriculturally acceptable anti-freezing agents, optionally ethylene glycol, glycerin, propylene glycol and/or urea.
80. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises a trace amount of water.
81. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises, consists essentially of or is a solid.
82. The inoculant composition of any one of the preceding paragraphs, wherein said inoculant composition comprises, consists essentially of or is a wettable powder.
83. The inoculant composition of any one of the preceding paragraphs, wherein at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more

microbial spores remain viable when said inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

84. The inoculant composition of any one of the preceding paragraphs, wherein at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is coated on a plant propagation material.

85. The inoculant composition of any one of the preceding paragraphs, wherein at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said one or more microbial spores remain viable when said inoculant composition is coated on a plant propagation material and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

86. The inoculant composition of any one of the preceding paragraphs, wherein at least about 1×10^1 to about 1×10^{10} colony-forming units of said one or more microbial spores per gram and/or milliliter of said inoculant composition remain viable when said inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more, optionally at least 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 or more colony-forming units per gram and/or milliliter of said inoculant composition.

87. The inoculant composition of any one of the preceding paragraphs, wherein at least about 1×10^1 to about 1×10^{10} colony-forming units of said one or more microbial spores per seed remain viable when said inoculant composition is coated on a seed and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more, optionally 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 or more colony-forming units per seed.

88. A coated plant propagation material, comprising, consisting essentially of, or consisting of:
a plant propagation material; and
a coating that covers at least a portion of an outer surface of said seed, said coating comprising, consisting essentially of, or consisting of the inoculant composition of any one of claims 1-87.

89. The coated plant propagation material paragraph 88, wherein said coating comprises, consists essentially of, or consists of an inner coating layer that comprises said microbial spores and an outer coating layer that is devoid of said microbial spores.

90. The coated plant propagation material of any one of paragraphs 88-89, wherein said coating comprises, consists essentially of or is an amorphous solid.

91. The coated plant propagation material of any one of paragraphs 88-89, wherein said coating comprises, consists essentially of or is a wettable powder.

92. The coated plant propagation material of any one of paragraphs 88-91, wherein said coating comprises about 1×10^1 to about 1×10^{15} colony-forming units of said microbial spores, optionally 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units.

93. The coated plant propagation material any one of paragraphs 88-92, wherein said plant propagation material is a seed.

94. The coated plant propagation material of paragraph 93, wherein said seed is a monocot seed.

95. The coated plant propagation material of paragraph 93, wherein said seed is a dicot seed.

96. The coated plant propagation material of paragraph 93, wherein said seed is a leguminous seed.

97. The coated plant propagation material of paragraph 93, wherein said seed is a non-leguminous seed.

98. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Amaranthaceae, optionally chard, spinach, sugar beet, or quinoa.

99. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Asteraceae, optionally artichoke, asters, chamomile, chicory, chrysanthemums, dahlias, daisies, echinacea, goldenrod, guayule, lettuce, marigolds, safflower, sunflowers, or zinnias.

100. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Brassicaceae, optionally arugula, broccoli, bok choy, Brussels sprouts, cabbage, cauliflower, canola, collard greens, daikon, garden cress, horseradish, kale, mustard, radish, rapeseed, rutabaga, turnip, wasabi, watercress, or *Arabidopsis thaliana*.

101. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Cucurbitaceae, optionally cantaloupe, cucumber, honeydew, melon, pumpkin, squash (e.g., acorn squash, butternut squash, summer squash), watermelon, or zucchini.

102. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Fabaceae, optionally alfalfa, beans, carob, clover, guar, lentils, mesquite, peas, peanuts, soybeans, tamarind, tragacanth, or vetch.

103. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Malvaceae, optionally cacao, cotton, durian, hibiscus, kenaf, kola, or okra.

104. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Poaceae, optionally bamboo, barley, corn, fonio, lawn grass (e.g., Bahia grass, Bermudagrass, bluegrass, Buffalograss, Centipede grass, Fescue, or Zoysia), millet, oats, ornamental grasses, rice, rye, sorghum, sugar cane, triticale, or wheat.

105. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Polygonaceae, optionally buckwheat.

106. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Rosaceae, optionally almonds, apples, apricots, blackberry, blueberry, cherries, peaches, plums, quinces, raspberries, roses, or strawberries.

107. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Solanaceae, optionally bell peppers, chili peppers, eggplant, petunia, potato, tobacco, or tomato.

108. The coated plant propagation material of any one of paragraphs 88-93, wherein said plant propagation material is of the family Vitaceae, optionally grape.

109. A kit, comprising:

the coated plant propagation material of any one of paragraphs 88-108; and
a container housing said coated plant propagation material.

110. The kit of claim 109, wherein said container reduces the amount of ambient light that reaches said coated plant propagation material by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% when sealed.

111. The kit of any one of paragraphs 109-110, wherein said container reduces the amount of ambient oxygen that reaches said plant propagation material by about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100% when sealed.

112. The kit of any one of paragraphs 109-111, wherein said container comprises, consists essentially of, or consists of a material having light permeability of less than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70 or 75%.

113. The kit of any one of paragraphs 109-112, wherein said container comprises, consists essentially of, or consists of a material having an oxygen transmission rate of less than about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, or 500 cm³/m²·day (as measured in accordance with ASTM D3985).

114. The kit of any one of paragraphs 109-113, wherein said kit further comprises one or more oxygen-absorbing compound, optionally activated carbon, iron powder, sodium chloride, ferrous carbonate, one or more metal halide catalysts and/or sodium hydrogen carbonate.

115. A plant treated with the inoculant composition of any one of paragraphs 1-87.

116. A plant germinated from the coated plant propagation material of any one of paragraphs 88-108.

117. A plant part harvested from the plant of any one of paragraphs 115-116.

118. A processed product produced from the plant part of paragraph 117.

119. A crop comprising, consisting essentially of, or consisting of a plurality of the plant or plant part of any one of paragraphs 115-117.

120. A method, comprising, consisting essentially of, or consisting of:
applying the inoculant composition of any one of paragraphs 1-87 to a plant propagation material.

121. The method of paragraph 120, further comprising planting said plant propagation material in a growth medium, optionally soil.

122. The method of paragraph 121, wherein said plant propagation material is planted in soil in which plants of the same genus were cultivated in at least one of the three years prior to said planting, optionally in each of the one, two or three years immediately preceding said planting.

123. The method of any one of paragraphs 121-122, wherein said inoculant composition is applied to the plant propagation material at the time of planting.

124. The method of any one of paragraphs 121-122, wherein said inoculant composition is applied to the plant propagation material at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15, 18, 21, 24, 27, 30, 33, 36, 39, 42, 45, 48 hours or more prior to planting.

125. The method of any one of paragraphs 121-122, wherein said inoculant composition is applied to the plant propagation material at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more prior to planting.

126. The method of any one of paragraphs 121-122, wherein said inoculant composition is applied to the plant propagation material about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36 months or more prior to planting
127. The method of any one of paragraphs 121-126, wherein said plant propagation material is a seed.
128. The method of any one of paragraphs 121-126, wherein said plant propagation material is a monocot seed.
129. The method of any one of paragraphs 121-126, wherein said plant propagation material is a dicot seed.
130. The method of any one of paragraphs 121-126, wherein said plant propagation material is a leguminous seed.
131. The method of any one of paragraphs 121-126, wherein said plant propagation material is a non-leguminous seed.
132. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Amaranthaceae, optionally chard, spinach, sugar beet, or quinoa.
133. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Asteraceae, optionally artichoke, asters, chamomile, chicory, chrysanthemums, dahlias, daisies, echinacea, goldenrod, guayule, lettuce, marigolds, safflower, sunflowers, or zinnias.
134. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Brassicaceae, optionally arugula, broccoli, bok choy, Brussels sprouts, cabbage, cauliflower, canola, collard greens, daikon, garden cress, horseradish, kale, mustard, radish, rapeseed, rutabaga, turnip, wasabi, watercress, or *Arabidopsis thaliana*.
135. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Cucurbitaceae, optionally cantaloupe, cucumber, honeydew, melon, pumpkin, squash (e.g., acorn squash, butternut squash, summer squash), watermelon, or zucchini.
136. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Fabaceae, optionally alfalfa, beans, carob, clover, guar, lentils, mesquite, peas, peanuts, soybeans, tamarind, tragacanth, or vetch.
137. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Malvaceae, optionally cacao, cotton, durian, hibiscus, kenaf, kola, or okra.
138. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Poaceae, optionally bamboo, barley, corn, fonio, lawn grass (e.g., Bahia grass, Bermudagrass, bluegrass,

Buffalograss, Centipede grass, Fescue, or Zoysia), millet, oats, ornamental grasses, rice, rye, sorghum, sugar cane, triticale, or wheat.

139. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Polygonaceae, optionally buckwheat.

140. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Rosaceae, optionally almonds, apples, apricots, blackberry, blueberry, cherries, peaches, plums, quinces, raspberries, roses, or strawberries.

141. The method of any one of paragraphs 121-131, wherein said plant propagation material is of the family Solanaceae, optionally bell peppers, chili peppers, eggplant, petunia, potato, tobacco, or tomato.

142. The method of any one of paragraphs 121-131, wherein plant propagation material seed is of the family Vitaceae, optionally grape.

143. A method comprising, consisting essentially of, or consisting of:
planting the coated plant propagation material of any one of paragraphs 88-108 in a growth medium, optionally soil.

144. The method of any one of paragraphs 121-143, further comprising applying the inoculant composition of any one of paragraphs 1-87 to the plant that grows from the plant propagation material.

145. A method of enhancing the stability and/or survivability of one or more microorganisms in a composition, comprising, consisting essentially of, or consisting of:

adding one or more protectants to said composition.

146. The method of paragraph 145, wherein said one or more protectants comprises, consists essentially of, or consists of:

one or more maltodextrins having a dextrose equivalent value of about 10 to about 25, optionally about 15 to about 20;

one or more hygroscopic polymers, optionally one or more albumins, alginates, celluloses, gums (e.g., cellulose gum, guar gum, gum arabic, gum combretum, xanthan gum), methyl celluloses, nylons, pectins, polyacrylic acids, polycarbonates, polyethylene glycols (PEG), polyethylenimines (PEI), polylactides, polymethylacrylates (PMA), polyurethanes, polyvinyl alcohols (PVA), polyvinylpyrrolidones (PVP), propylene glycols, sodium carboxymethyl celluloses and/or starches; and/or

oxidation control components, optionally one or more antioxidants (e.g., ascorbic acid, ascorbyl palmitate, ascorbyl stearate, calcium ascorbate, one or more carotenoids, lipoic acid, one or more phenolic compounds (e.g., one or more flavonoids, flavones and/or flavonols), potassium ascorbate, sodium ascorbate, one or more thiols (e.g., glutathione, lipoic acid and/or N-acetyl cysteine), one or more tocopherols, one or more tocotrienols, ubiquinone and/or uric acid) and/or one or more oxygen scavengers, optionally ascorbic acid and/or sodium hydrogen carbonate.

147. The method of any one of paragraphs 145-146, wherein said one or more protectants is added until it comprises about 0.0001 to about 10% (by weight) of said composition, optionally about 2 to about 6% (by weight) of said composition, optionally about 0.0005, 0.001, 0.002, 0.003, 0.004, 0.005, 0.0075, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 7, 7.5, 8, 8.5, 9, 9.5 or 10% (by weight) of said composition.

148. The method of any one of paragraphs 145-147, wherein said one or more protectants is added in an amount sufficient to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said microbial spores remain viable when said inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

149. The method of any one of paragraphs 145-148, wherein said one or more protectants is added in an amount sufficient to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said microbial spores remain viable when said inoculant composition is coated on a plant propagation material.

150. The method of any one of paragraphs 145-149, wherein said one or more protectants is added in an amount sufficient to ensure that at least about 0.01, 0.05, 0.1, 0.5, 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95% or more of said microbial spores remain viable when said inoculant composition is coated on a plant propagation material and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more.

151. The method of any one of paragraphs 145-150, wherein said one or more protectants is added in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units of said microbial spores per gram and/or milliliter of said inoculant composition remain viable when said inoculant composition is stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units per gram and/or milliliter of said inoculant composition.

152. The method of any one of paragraphs 145-151, wherein said one or more protectants is added in an amount sufficient to ensure that at least about 1×10^1 to about 1×10^{15} colony-forming units of said microbial spores per seed remain viable when said inoculant composition is coated on a seed and stored at 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 and/or 40°C and 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85% or more relative humidity for a period of 1, 2, 3,

4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 28, 32, 36, 40, 44, 48, 52, 56, 60, 64, 68, 72, 76, 80, 84, 88, 92, 96, 100, 104 weeks or more, optionally 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} or more colony-forming units per seed.

153. The method of any of paragraphs 145-152, further comprising:

adding one or more dispersants to said composition.

154. The method of paragraph 153, wherein said one or more dispersants comprises:

one or more anionic surfactants, optionally one or more alkyl carboxylates, alkyl sulfates, alkyl ether sulfates, alkyl amido ether sulfates, alkyl aryl polyether sulfates, alkyl aryl sulfates, alkyl aryl sulfonates, alkyl sulfonates, alkyl amide sulfonates, alkyl aryl sulfonates, alkyl benzene sulfonates, alkyl diphenyloxide sulfonate, alpha-olefin sulfonates, alkyl naphthalene sulfonates, paraffin sulfonates, alkyl sulfosuccinates, alkyl ether sulfosuccinates, alkylamide sulfosuccinates, alkyl sulfosuccinamates, alkyl sulfoacetates, alkyl phosphates, alkyl ether phosphates, acyl sarconsinates, acyl isethionates, N-acyl tartrates, N-acyl-N-alkyltaurates, benzene sulfonates, cumene sulfonates, dioctyl sodium sulfosuccinate, ethoxylated sulfosuccinates, lignin sulfonates, linear alkylbenzene sulfonates, monoglyceride sulfates, perfluorobutanesulfonate, perfluorooctanesulfonate, phosphate ester, styrene acrylic polymers, toluene sulfonates and/or xylene sulfonates

one or more cationic surfactants, optionally one or more alkyltrimethylammonium salts, cetylpyridinium chloride, benzalkonium chloride, benzethonium chloride, 5-bromo-5-nitro-1,3-dioxane, dimethyldioctadecylammomum chloride, cetrimonium bromide, dioctadecyldimethylammonium bromide and/or octenidine dihydrochloride; and/or

one or more non-ionic surfactants, optionally one or more alcohol ethoxylates, alkanolamides, alkanolamine condensates, carboxylic acid esters, cetostearyl alcohol, cetyl alcohol, cocamide DEA, dodecyldimethylamine oxides, ethanolamides, ethoxylates of glycerol ester and glycol esters, ethylene oxide polymers, ethylene oxide-propylene oxide copolymers, glucoside alkyl ethers, glycerol alkyl ethers, glycerol esters, glycol alkyl ethers, glycol alkylphenol ethers, glycol esters, monolaurin, pentaethylene glycol monododecyl ethers, poloxamer, polyamines, polyglycerol polyricinoleate, polysorbate, polyoxyethylenated fatty acids, polyoxyethylenated mercaptans, polyoxyethylenated polyoxypropylene glycols, polyoxyethylene glycol sorbitan alkyl esters, polyethylene glycol-polypropylene glycol copolymers, polyoxyethylene glycol octylphenol ethers, polyvinyl pynolidones, sugar-based alkyl polyglycosides, sulfoanlamides, sorbitan fatty acid alcohol ethoxylates, sorbitan fatty acid ester ethoxylates, sorbitan fatty acid ester and/or tertiary acetylenic glycols; and/or

one or more wetting agents, optionally one or more naphthalene sulfonates, optionally one or more alkyl naphthalene sulfonates, one or more isopropyl naphthalene sulfonates and/or one or more butyl naphthalene sulfonates.

155. The inoculant composition of any one of paragraphs 153-154, wherein said one or more dispersants comprises one or more polyoxyethylene alkyl ethers, one or more acrylic copolymers, one or more polyoxyethylene sorbitan trioleates and/or one or more secondary alcohol ethoxylates.

156. The method of any one of claims 145-155, further:

adding one or more non-aqueous microbial extracts to said composition.

157. The method of paragraph 156, wherein said one or more non-aqueous microbial extracts comprises:

one or more *Bacillus* extracts, optionally an extract of media comprising *B. amyloliquefaciens* D747, *B. amyloliquefaciens* NRRL B-50349, *B. amyloliquefaciens* TJ1000, *B. amyloliquefaciens* FZB24, *B. amyloliquefaciens* FZB42, *B. amyloliquefaciens* IN937a, *B. amyloliquefaciens* IT-45, *B. amyloliquefaciens* TJ1000, *B. amyloliquefaciens* MBI600, *B. amyloliquefaciens* BS27 (deposited as NRRL B-5015), *B. amyloliquefaciens* BS2084 (deposited as NRRL B-50013), *B. amyloliquefaciens* 15AP4 (deposited as ATCC PTA-6507), *B. amyloliquefaciens* 3AP4 (deposited as ATCC PTA-6506), *B. amyloliquefaciens* LSSA01 (deposited as NRRL B-50104), *B. amyloliquefaciens* ABP278 (deposited as NRRL B-50634), *B. amyloliquefaciens* 1013 (deposited as NRRL B-50509), *B. amyloliquefaciens* 918 (deposited as NRRL B-50508), *B. amyloliquefaciens* 22CP1 (deposited as ATCC PTA-6508) and *B. amyloliquefaciens* BS18 (deposited as NRRL B-50633), *B. cereus* 1-1562, *B. firmus* 1-1582, *B. lichenformis* BA842 (deposited as NRRL B-50516), *B. lichenformis* BL21 (deposited as NRRL B-50134), *B. mycoides* NRRL B-21664, *B. pumilus* NRRL B-21662, *B. pumilus* NRRL B-30087, *B. pumilus* ATCC 55608, *B. pumilus* ATCC 55609, *B. pumilus* GB34, *B. pumilus* KFP9F, *B. pumilus* QST 2808, *B. subtilis* ATCC 55078, *B. subtilis* ATCC 55079, *B. subtilis* MBI 600, *B. subtilis* NRRL B-21661, *B. subtilis* NRRL B-21665, *B. subtilis* CX-9060, *B. subtilis* GB03, *B. subtilis* GB07, *B. subtilis* QST-713, *B. subtilis* FZB24, *B. subtilis* D747, *B. subtilis* 3BP5 (deposited as NRRL B-50510), *B. thuringiensis* ATCC 13367, *B. thuringiensis* GC-91, *B. thuringiensis* NRRL B-21619, *B. thuringiensis* ABTS-1857, *B. thuringiensis* SAN 401 I, *B. thuringiensis* ABG-6305, *B. thuringiensis* ABG-6346, *B. thuringiensis* AM65-52, *B. thuringiensis* SA-12, *B. thuringiensis* SB4, *B. thuringiensis* ABTS-35 I, *B. thuringiensis* HD-1, *B. thuringiensis* EG 2348, *B. thuringiensis* EG 7826, *B. thuringiensis* EG 7841, *B. thuringiensis* DSM 2803, *B. thuringiensis* NB-125 and/or *B. thuringiensis* NB-176;

one or more *Bradyrhizobium* extracts, optionally an extract of media comprising *B. elkanii* SEMIA 501, *B. elkanii* SEMIA 587, *B. elkanii* SEMIA 5019, *B. japonicum* NRRL B-50586 (also deposited as NRRL B-59565), *B. japonicum* NRRL B-50587 (also deposited as NRRL B-59566), *B. japonicum* NRRL B-50588 (also deposited as NRRL B-59567), *B. japonicum* NRRL B-50589 (also deposited as NRRL B-59568), *B. japonicum* NRRL B-50590 (also deposited as NRRL B-59569), *B. japonicum* NRRL B-50591 (also deposited as NRRL B-59570), *B. japonicum* NRRL B-50592 (also deposited as NRRL B-59571), *B. japonicum* NRRL B-50593 (also deposited as NRRL B-59572), *B. japonicum* NRRL B-50594 (also deposited as NRRL B-50493), *B. japonicum* NRRL B-50608, *B. japonicum* NRRL B-50609, *B. japonicum* NRRL B-50610, *B. japonicum* NRRL B-50611, *B. japonicum* NRRL B-50612, *B. japonicum* NRRL B-50726, *B. japonicum* NRRL B-50727, *B. japonicum* NRRL B-50728, *B. japonicum* NRRL B-50729, *B. japonicum* NRRL B-50730, *B. japonicum* SEMIA 566, *B. japonicum* SEMIA 5079, *B. japonicum* SEMIA 5080, *B. japonicum* USDA 6, *B. japonicum* USDA 110, *B. japonicum* USDA 122, *B. japonicum* USDA 123, *B. japonicum* USDA 127, *B. japonicum* USDA 129 and/or *B. japonicum* USDA 532C;

one or more *Rhizobium* extracts, optionally an extract of media comprising *R. leguminosarum* S012A-2;

one or more *Sinorhizobium* extracts, optionally an extract of media comprising *S. fredii* CCBAU114 and/or *S. fredii* USDA 205;

one or more *Penicillium* extracts, optionally an extract of media comprising *P. bilaiae* ATCC 18309, *P. bilaiae* ATCC 2085 I, *P. bilaiae* ATCC 22348, *P. bilaiae* NRRL 50162, *P. bilaiae* NRRL 50169, *P. bilaiae* NRRL 50776, *P. bilaiae* NRRL 50777, *P. bilaiae* NRRL 50778, *P. bilaiae* NRRL 50777, *P. bilaiae* NRRL 50778, *P. bilaiae* NRRL 50779, *P. bilaiae* NRRL 50780, *P. bilaiae* NRRL 50781, *P. bilaiae* NRRL 50782, *P. bilaiae* NRRL 50783, *P. bilaiae* NRRL 50784, *P. bilaiae* NRRL 50785, *P. bilaiae* NRRL 50786, *P. bilaiae* NRRL 50787, *P. bilaiae* NRRL 50788, *P. bilaiae* RS7B-SD1, *P. brevicompactum* AgRF18, *P. canescens* ATCC 10419, *P. expansum* ATCC 24692, *P. expansum* YT02, *P. fellatanum* ATCC 48694, *P. gaestrivorus* NRRL 50170, *P. glabrum* DAOM 239074,

P. glabrum CBS 229.28, *P. janthinellum* ATCC 10455, *P. lanosocoeruleum* ATCC 48919, *P. radicum* ATCC 201836, *P. radicum* FRR 4717, *P. radicum* FRR 4719, *P. radicum* N93/47267 and/or *P. raistrickii* ATCC 10490;

one or more *Streptomyces* extracts, optionally an extract of media comprising *Streptomyces* NRRL B-30145, *Streptomyces* M1064, *S. galbus* NRRL 30232, *S. lydicus* WYEC 108 (ATCC 55445), *S. violaceusniger* YCED 9 (ATCC 55660) and/or *Streptomyces* WYE 53 (ATCC 55750); and/or

one or more *Trichoderma* extracts, optionally an extract of media comprising *T. asperellum* SKT-1 (ECO-HOPE®, Kumiai Chemical Industry Co., Ltd., Japan), *T. atroviride* LC52 (SENTINEL®, Agrimm Technologies Ltd, NZ), *T. harzianum* T-22 (PLANTSHIELD®, der Firma BioWorks Inc., USA), *T. harzianum* TH-35 (ROOT PRO®, from Mycontrol Ltd., Israel), *T. harzianum* T-39 (TRICHODEX®, Mycontrol Ltd., Israel; T. 2000®, Makhteshim Ltd., Israel), *T. harzianum* ICC012 and *T. viride* TRICHOPEL (Agrimm Technologies Ltd, NZ), *T. harzianum* ICC012 and *T. viride* ICC080 (REMEDIER® WP, Isagro Ricerca, Italy), *T. polysporum* and *T. harzianum* (BINAB®, BINAB Bio-Innovation AB, Sweden), *T. stromaticum* TRICOVAB® (C.E.P.L.A.C., Brazil), *T. virens* GL-21 (SOILGARD®, Certis LLC, USA), *T. virens* GI-3, ATCC 57678, *T. virens* GI-21 (Thermo Trilog Corporation, Wasco, CA), *T. virens* GI-3 and *Bacillus amyloliquefaciens* FZB2, *T. virens* GI-3 and *Bacillus amyloliquefaciens* NRRL B-50349, *T. virens* GI-3 and *Bacillus amyloliquefaciens* TJ1000, *T. virens* GI-21 and *Bacillus amyloliquefaciens* FZB24, *T. virens* GI-21 and *Bacillus amyloliquefaciens* NRRL B-50349, *T. virens* GI-21 and *Bacillus amyloliquefaciens* TJ1000, *T. viride* TRIECO® (Ecosense Labs. (India) Pvt. Ltd., Indien, BIO-CURE® F from T. Stanes & Co. Ltd., Indien), *T. viride* TV1 (Agribiotec srl, Italy), *T. viride* ICC080.

158. The inoculant composition of any one of paragraphs 156-157, wherein said one or more non-aqueous microbial extracts comprises less than 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5% water by weight, based upon the total weight of the composition.

159. The method of any one of paragraphs 156-158, wherein said one or more non-aqueous microbial extracts is added until it comprises about 0.1 to about 5% (by weight) of said composition, optionally about 0.1 to about 2% (by weight) of said composition, optionally about 0.1, 0.15, 0.2, 0.25, 0.3, 0.35, 0.4, 0.45, 0.5, 0.55, 0.6, 0.65, 0.7, 0.75, 0.8, 0.85, 0.9, 0.95, 1, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4, 4.1, 4.2, 4.3, 4.4., 4.5, 4.6, 4.7, 4.8, 4.9, 5% or more (by weight) of said composition.

160. A method, comprising, consisting essentially of or consisting of:

applying the inoculant composition of any one of paragraphs 1-87 to a seed and/or to the plant that grows from said seed;

applying a second composition to said seed and/or to the plant that grows from said seed, said second composition comprising:

one or more agriculturally beneficial microorganisms, optionally one or more diazotrophs, one or more phosphate-solubilizing microorganisms, one or more mycorrhizal fungi and/or one or more biopesticides, optionally one or more biofungicides, one or more bioinsecticides and/or one or more bionematicides;

one or more biostimulants, optionally one or more seaweed extracts, one or more humic acids, one or more fulvic acids, myo-inositol and/or glycine;

one or more nutrients, optionally one or more vitamins (e.g., vitamin A, vitamin B complex (i.e., vitamin B1, vitamin B2, vitamin B3, vitamin B5, vitamin B6, vitamin B7, vitamin B8, vitamin B9, vitamin B12,

choline) vitamin C, vitamin D, vitamin E, vitamin K, carotenoids (α -carotene, β -carotene, cryptoxanthin, lutein, lycopene and/or zeaxanthin), macrominerals (e.g., calcium, iron, magnesium, nitrogen, phosphorous, potassium and/or sodium), trace minerals (e.g., boron, cobalt, chloride, chromium, copper, fluoride, iodine, iron, manganese, molybdenum, selenium and/or zinc) and/or organic acids (e.g., acetic acid, citric acid, lactic acid, malic acid and/or taurine);

one or more fungicides, optionally one or more of the fungicides disclosed on pages 31-32 above;

one or more herbicides, optionally one or more of the herbicides disclosed on page 33-34 above;

one or more insecticides, optionally one or more of the insecticides disclosed on page 30 above;

one or more nematicides, optionally one or more of the nematicides disclosed on page 30 above;

one or more lipo-chitooligosaccharides, optionally one or more of the lipo-chitooligosaccharides represented by formulas I-IV and/or one or more of the lipo-chitooligosaccharides represented by structures V-XXXIII;

one or more chitooligosaccharides, optionally one or more of the chitooligosaccharides represented by formulas XXXIV-XXXV and/or one or more of the chitooligosaccharides represented by structures XXXVI-XXXIX,

one or more chitinous compounds, optionally one or more chitins and/or one or more chitosans;

one or more flavonoids, optionally one or more anthocyanidins, such as cyanidin, delphinidin, malvidin, pelargonidin, peonidin and/or petunidin; anthoxanthins, such as flavones (e.g., apigenin, baicalein, chrysin, 7,8-dihydroxyflavone, diosmin, flavoxate, 6—hydroxyflavone, luteolin, scutellarein, tangeritin and/or wogonin) and/or flavonols (e.g., amurensin, astragalin, azaleatin, azalein, fisetin, furanoflavonols galangin, gossypetin, 3-hydroxyflavone, hyperoside, icariin, isoquercetin, kaempferide, kaempferitrin, kaempferol, isorhamnetin, morin, myricetin, myricitrin, natsudaicain, pachypodol, pyranoflavonols quercetin, quericitin, rhamnazin, rhamnetin, robinin, rutin, spiraeoside, troxerutin and/or zanthorhamnin); flavanones, such as butin, eriodictyol, hesperetin, hesperidin, homoeriodictyol, isosakuranetin, naringenin, naringin, pinocembrin, poncirin, sakuranetin, sakuranin and/or sterubin; flavanonols, such as dihydrokaempferol and/or taxifolin; flavans, such as flavan-3-ols (e.g., catechin (C), catechin 3-gallate (Cg), epicatechins (EC), epigallocatechin (EGC) epicatechin 3-gallate (ECg), epigallocatechin 3-gallate (EGCg), epiafzelechin, fisetinidol, galocatechin (GC), gallcatechin 3-gallate (GCg), guibourtinidol, mesquitol, robinetinidol, theaflavin-3-gallate, theaflavin-3'-gallate, theflavin-3,3'-digallate, thearubigin), flavan-4-ols (e.g., apiforol and/or luteoforol) and/or flavan-3,4-diols (e.g., leucocyanidin, leucodelphinidin, leucofisetinidin, leucomalvidin, leucopelargonidin, leucoppeonidin, leucorobinetinidin, melacacidin and/or teracacidin); and/or isoflavonoids, such as isoflavones (e.g., biochanin A, daidzein, formononetin, genistein and/or glycitein), isoflavanes (e.g., equol, ionchocarpane and/or laxifloorane), isoflavandiols, isoflavenes (e.g., glabrene, hagin D and/or 2-methoxyjudaicin), coumestans (e.g., coumestrol, plicadin and/or wedelolactone), pterocarpanes and/or rotonoids; and/or one or more analogues, derivatives, hydrates, isomers, polymers, salts and solvates thereof, such as neoflavonoids (e.g., calophyllolide, coutareagenin, dalbergichromene, dalbergin and/or nivetin) and/or pterocarpanes (e.g., bitucarpin A, bitucarpin B, erybraedin A, erybraedin B, erythrabyscin II, erythrabyscin-1, erycristagallin, glycinol, glyceollidins, glyceollins, glycyrrhizol, maackiain, medicarpin, morisianine, orientanol, phaseolin, pisatin, striatine and/or trifolirhizin);

jasmonic acid and/or one or more derivatives thereof;

linoleic acid and/or one or more derivatives thereof;

linolenic acid and/or one or more derivatives thereof;

one or more karrakins, optionally one or more karrakins represented by formula XXXX;

gluconolactone; and/or

one or more oxidation control components, optionally one or more antioxidants (e.g., ascorbic acid, ascorbyl palmitate, ascorbyl stearate, calcium ascorbate, one or more carotenoids, lipoic acid, one or more phenolic compounds (e.g., one or more flavonoids, flavones and/or flavonols), potassium ascorbate, sodium ascorbate, one or more thiols (e.g., glutathione, lipoic acid and/or N-acetyl cysteine), one or more tocopherols, one or more tocotrienols, ubiquinone and/or uric acid) and/or one or more oxygen scavengers, optionally ascorbic acid and/or sodium hydrogen carbonate.

EXAMPLES

The following examples are not intended to be a detailed catalogue of all the different ways in which the present disclosure may be implemented or of all the features that may be added to the present disclosure. Subjects skilled in the art will appreciate that numerous variations and additions to the various embodiments may be made without departing from the present disclosure. Hence, the following descriptions are intended to illustrate some particular embodiments of the invention and not to exhaustively specify all permutations, combinations and variations thereof.

Example 1

Stable Wettable Powders Comprising *Penicillium* Spores

Solid non-aqueous inoculant compositions of the present disclosure comprising *Penicillium bilaiae* spores (ATCC 2085 1 and/or RS7B-SD1) were stored for four weeks at 40° C or for twenty weeks at 20°C alongside a commercially available wettable powder comprising *Penicillium bilaiae* spores (ATCC 2085 1 and RS7B-SD1). The survival rate of *Penicillium* spores was greater in each of the solid non-aqueous inoculant compositions of the present disclosure than in the commercially available wettable powder. **Table 1.**

Table 1.

Inoculant Composition		Viable spores after 20 weeks at 20°C ¹	Viable spores after 4 weeks at 40°C ¹
A	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (74.879% w/w) + maltose monohydrate (8.320% w/w) + MULTIWET MO-85P-PW-(AP) (2.750% w/w) + SUNSPRAY® 6N (4.051% w/w)	51%	11%
B	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (62.40% w/w) + maltose monohydrate (20.80% w/w) + MULTIWET MO-85P-PW-(AP) (2.75% w/w) + SUNSPRAY® 6N (4.05% w/w)	54%	13%
C	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (49.92% w/w) + maltose monohydrate (33.28% w/w) + MULTIWET MO-85P-PW-(AP) (2.5% w/w) + SUNSPRAY® 6N (4.05% w/w)	40%	12%
D	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (52.31% w/w) + maltose monohydrate (34.88% w/w) + BIOSOFT® N23-3 (2.81% w/w)	79%	28%
E	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (65.39% w/w) + maltose monohydrate (21.80% w/w) + BIOSOFT® N23-3 (2.81% w/w)	93%	32%
F	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (78.469% w/w) + maltose monohydrate (8.719% w/w) + BIOSOFT® N23-3 (2.813% w/w)	68%	34%
Control	Commercially available wettable powder comprising <i>P. bilaiae</i> spores	52%	4%

¹ Expressed as a percentage of the spore content (cfu per gram of inoculant composition) measured at time zero.

The abilities of the aforementioned inoculant compositions to disperse *Penicillium bilaiae* spores are tested by measuring the ease with which each inoculant composition can be mixed into 100 ml of water in a 250 ml Erlenmeyer flask at 130 rpm on an orbital shaker. As a follow up to this qualitative testing, the percentage of single spores (compared to clumps of >2 spores) in each inoculant composition is calculated by observing the spores under a microscope at 200X magnification. Each of the solid non-aqueous inoculant compositions of the present disclosure exhibits greater spore dispersion than the commercially available wettable powder.

Example 2

Stable Wettable Powders Enhance On-Seed Stability

Solid non-aqueous inoculant compositions of the present disclosure comprising spores of *Penicillium bilaiae* and a commercially available wettable powder comprising corresponding spores were coated on corn seeds treated

with a commercially available pesticide. The coated seeds were stored at 10, 20 or 30°C and 50% relative humidity for 1, 2, 3, 4, 6, 8, 10, 12, 16, 18, 20, 22, 26, 30, 32, 34 or 42 weeks and then assayed for on-seed survivability. As indicated in **Table 2** and **Figures 1-3**, the survival rate of *Penicillium* spores was greater on seeds coated with the solid non-aqueous inoculant compositions of the present disclosure than on seeds coated with the commercially available wettable powder.

Table 2.

Inoculant Composition		Viable spores after 42 weeks at 10°C ¹	Viable spores after 20 weeks at 20°C ¹	Viable spores after 12 weeks at 30°C ¹
F	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (78.469% w/w) + maltose monohydrate (8.719% w/w) + BIOSOFT® N23-3 (2.813% w/w)	8%	28%	11%
A	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (74.879% w/w) + maltose monohydrate (8.320% w/w) + MULTIWET MO-85P-PW-(AP) (2.750% w/w) + SUNSPRAY® 6N (4.051% w/w)	7%	23%	17%
Control	Commercially available wettable powder comprising <i>P. bilaiae</i> spores	2%	14%	2%

¹ Expressed as a percentage of the spore content (cfu per seed) measured at time zero.

Example 3

Stable Wettable Powders Enhance On-Seed Stability

Solid non-aqueous inoculant compositions of the present disclosure comprising spores of *Penicillium bilaiae* and a commercially available wettable powder comprising corresponding spores were coated on corn seeds treated with a commercially available pesticide. The coated seeds were stored at 20 or 30°C and 50% relative humidity for 12 weeks and then assayed for on-seed survivability. As indicated in **Table 3**, the survival rate of *Penicillium* spores was greater on seeds coated with the solid non-aqueous inoculant compositions of the present disclosure than on seeds coated with the commercially available wettable powder.

Table 3.

Inoculant Composition		Viable spores after 12 weeks at 20°C ¹	Viable spores after 12 weeks at 30°C ¹
E	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (65.39% w/w) + maltose monohydrate (21.80% w/w) + BIOSOFT® N23-3 (2.81% w/w)	58%	8%
D	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (52.31% w/w) + maltose monohydrate (34.88% w/w) + BIOSOFT® N23-3 (2.81% w/w)	55%	5%
B	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (62.40% w/w) + maltose monohydrate (20.80% w/w) + MULTIWET MO-85P-PW-(AP) (2.75% w/w) + SUNSPRAY® 6N (4.05% w/w)	38%	8%
C	<i>P. bilaiae</i> spores (10% w/w) + MALTRIN QD® M580 (49.92% w/w) + maltose monohydrate (33.28% w/w) + MULTIWET MO-85P-PW-(AP) (2.5% w/w) + SUNSPRAY® 6N (4.05% w/w)	41%	6%
Control	Commercially available wettable powder comprising <i>P. bilaiae</i> spores	33%	2%

¹ Expressed as a percentage of the spore content (cfu per seed) measured at time zero.

THAT WHICH IS CLAIMED:

1. A non-aqueous inoculant composition, comprising:

Penicillium spores comprising about 0.1 to about 30% (by weight) of said inoculant composition, optionally about 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24 or 25% (by weight) of said inoculant composition;

one or more dispersants comprising about 0.1 to about 5% (by weight) of said composition, optionally about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5 or 5% (by weight) of said composition; and

a solid non-aqueous carrier comprising about 70 to about 95% (by weight) of said inoculant composition, optionally about 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94 or 95% (by weight) of said inoculant composition, said solid non-aqueous carrier comprising one or more maltodextrins having a dextrose equivalent value of about 15 to about 20 and one or more disaccharides, optionally maltose and/or trehalose.

2. The inoculant composition of claim 1, wherein said *Penicillium* spores comprise about 8 to about 12% (by weight) of said inoculant composition, optionally about 8, 8.5, 9, 9.5, 10, 10.5, 11, 11.5 or 12% (by weight) of said inoculant composition.

3. The inoculant composition of any one of claims 1-2, wherein said *Penicillium* spores comprise about 1×10^1 to about 1×10^{15} colony-forming units, optionally at least 1×10^1 , 1×10^2 , 1×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} colony-forming units.

4. The inoculant composition of any one of claims 1-3, wherein said *Penicillium* spores comprise spores of *Penicillium bilaiae* ATCC 18309, *Penicillium bilaiae* ATCC 20851, *Penicillium bilaiae* ATCC 22348, *Penicillium bilaiae* NRRL 50162, *Penicillium bilaiae* NRRL 50169, *Penicillium bilaiae* NRRL 50776, *Penicillium bilaiae* NRRL 50777, *Penicillium bilaiae* NRRL 50778, *Penicillium bilaiae* NRRL 50779, *Penicillium bilaiae* NRRL 50780, *Penicillium bilaiae* NRRL 50781, *Penicillium bilaiae* NRRL 50782, *Penicillium bilaiae* NRRL 50783, *Penicillium bilaiae* NRRL 50784, *Penicillium bilaiae* NRRL 50785, *Penicillium bilaiae* NRRL 50786, *Penicillium bilaiae* NRRL 50787, *Penicillium bilaiae* NRRL 50788, *Penicillium bilaiae* RS7B-SD1, *Penicillium brevicompactum* AgRF18, *Penicillium canescens* ATCC 10419, *Penicillium expansum* ATCC 24692, *Penicillium expansum* YT02, *Penicillium fellatanum* ATCC 48694, *Penicillium gastrivorus* NRRL 50170, *Penicillium glabrum* DAOM 239074, *Penicillium glabrum* CBS 229.28, *Penicillium janthinellum* ATCC 10455, *Penicillium lanosocoeruleum* ATCC 48919, *Penicillium radicum* ATCC 201836, *Penicillium radicum* FRR 4717, *Penicillium radicum* FRR 4719, *Penicillium radicum* N93/47267 and/or *Penicillium raistrickii* ATCC 10490.

5. The inoculant composition of any one of claims 1-4, wherein said solid non-aqueous carrier comprises about 80 to about 90% (by weight) of said inoculant composition, optionally about 80, 80.5, 81, 81.5, 82, 82.5, 83, 83.5, 84, 84.5, 85, 85.5, 86, 86.5, 87, 87.5, 88, 88.5, 89, 89.5 or 90% (by weight) of said inoculant composition.

6. The inoculant composition of any one of claims 1-5, wherein said one or more dispersants comprise about 0.1 to about 2% (by weight) of said composition, optionally about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5 or 2% (by weight) of said composition.

7. The inoculant composition of any one of claims 1-6, wherein said one or more dispersants comprises one or more polyoxyethylene alkyl ethers, one or more acrylic copolymers, one or more polyoxyethylene sorbitan trioleates and/or one or more secondary alcohol ethoxylates.

8. The inoculant composition of any one of claims 1-7, wherein said inoculant composition comprises one or more dust suppressants comprising about 0.1 to about 30% (by weight) of said composition, optionally about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75, 5, 6, 7, 8, 9 or 10% (by weight) of said composition.

9. The inoculant composition of any one of claim 8, wherein said one or more dust suppressants comprises one or more adhesives, one or more mineral oils, one or more paraffinic oils, one or more vegetable oils, one or more synthetic polymers and/or glycerin.

10. The inoculant composition of any one of claims 1-9, wherein said inoculant composition comprises one or more protectants comprising about 0.5 to about 10% (by weight) of said composition, optionally about 0.5, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 7, 7.5, 8, 8.5, 9, 9.5 or 10% (by weight) of said composition.

11. The inoculant composition of claim 10, wherein said one or more protectants comprises one or more hygroscopic polymers, optionally one or more albumins, alginates, celluloses, gums (e.g., cellulose gum, guar gum, gum arabic, gum combretum, xantham gum), methyl celluloses, nylons, pectins, polyacrylic acids, polycarbonates, polyethylene glycols (PEG), polyethylenimines (PEI), polylactides, polymethylacrylates (PMA), polyurethanes, polyvinyl alcohols (PVA), polyvinylpyrrolidones (PVP), propylene glycols, sodium carboxymethyl celluloses and/or starches.

12. The inoculant composition of any one of claims 1-11, wherein said inoculant composition comprises one or more pesticides, optionally:

one or more insecticides and/or nematicides, optionally one or more carbamates, diamides, macrocyclic lactones, neonicotinoids, organophosphates, phenylpyrazoles, pyrethrins, spinosyns, synthetic pyrethroids, tetric acids and/or tetramic acids;

one or more fungicides, optionally one or more aromatic hydrocarbons, benzimidazoles, benzthiadiazole, carboxamides, carboxylic acid amides, morpholines, phenylamides, phosphonates, quinone outside inhibitors, thiazolidines, thiophanates, thiophene carboxamides and/or triazoles;

and/or

one or more herbicides, optionally one or more acetyl CoA carboxylase (ACCase) inhibitors, acetolactate synthase (ALS) inhibitors, acetohydroxy acid synthase (AHAS) inhibitors, photosystem II inhibitors, photosystem I inhibitors, protoporphyrinogen oxidase (PPO or Protox) inhibitors, carotenoid biosynthesis inhibitors, enolpyruvyl shikimate-3-phosphate (EPSP) synthase inhibitor, glutamine synthetase inhibitor, dihydropteroate synthetase inhibitor, mitosis inhibitors, 4-hydroxyphenyl-pyruvate-dioxygenase (4-HPPD) inhibitors, synthetic auxins, auxin herbicide salts, auxin transport inhibitors, and/or nucleic acid inhibitors.

13. The inoculant composition of any one of claims 1-12, wherein said inoculant composition comprises one more lipo-chitooligosaccharides, optionally one or more lipo-chitooligosaccharides represented by formulas I-IV.
14. The inoculant composition of any one of claims 1-13, wherein said inoculant composition comprises one more lipo-chitooligosaccharides, optionally one or more of the lipo-chitooligosaccharides represented by structures V-XXXIII.
15. The inoculant composition of any one of claims 1-14, wherein said inoculant composition comprises one or more chitooligosaccharides, optionally one or more chitooligosaccharides represented by formulas XXXIV-XXXV.
16. The inoculant composition of any one of claims 1-15, wherein said inoculant composition comprises one or more chitooligosaccharides, optionally one or more chitooligosaccharides represented by structures XXXVI-XXXIX.
17. The inoculant composition of any one of claims 1-16, wherein said inoculant composition comprises one or more chitinous compounds, optionally one or more chitins and/or one or more chitosans.
18. The inoculant composition of any one of claims 1-17, wherein said inoculant composition comprises one or more flavonoids, optionally:
- one or more anthocyanidins, optionally cyanidin, delphinidin, malvidin, pelargonidin, peonidin and/or petunidin;
 - one or more anthoxanthins, optionally one or more flavones, such as apigenin, baicalein, chrysin, 7,8-dihydroxyflavone, diosmin, flavoxate, 6—hydroxyflavone, luteolin, scutellarein, tangeritin and/or wogonin; and/or flavonols, such as amurensin, astragalin, azaleatin, azalein, fisetin, furanoflavonols galangin, gossypetin, 3-hydroxyflavone, hyperoside, icariin, isoquercetin, kaempferide, kaempferitrin, kaempferol, isorhamnetin, morin, myricetin, myricitrin, natsudaicain, pachypodol, pyranoflavonols quercetin, quericitin, rhamnazin, rhamnetin, robinin, rutin, spiraeoside, troxerutin and/or zanthorhamnin;
 - one or more flavanones, optionally butin, eriodictyol, hesperetin, hesperidin, homoeriodictyol, isosakuranetin, naringenin, naringin, pinocembrin, poncirin, sakuranetin, sakuranin and/or sterubin;
 - one or more flavanonols, optionally dihydrokaempferol and/or taxifolin; flavans, such as flavan-3-ols (e.g., catechin (C), catechin 3-gallate (Cg), epicatechins (EC), epigallocatechin (EGC) epicatechin 3-gallate (ECg), epigallocatechin 3-gallate (EGCg), epiafzelechin, fisetinidol, galocatechin (GC), gallcatechin 3-gallate (GCg), guibourtinidol, mesquitol, robinetinidol, theaflavin-3-gallate, theaflavin-3'-gallate, theflavin-3,3'-digallate, thearubigin), flavan-4-ols (e.g., apiforol and/or luteoforol) and/or flavan-3,4-diols (e.g., leucocyanidin, leucodelphinidin, leucofisetinidin, leucomalvidin, luecopelargonidin, leucoppeonidin, leucorobinetinidin, melacacidin and/or teracacidin); and/or
 - one or more isoflavonoids, optionally one or more isoflavones, such as biochanin A, daidzein, formononetin, genistein and/or glycitein; isoflavanes, such as equol, ionchocarpene and/or laxifloorene; isoflavandiols; isoflavenes, such as glabrene, haginin **D** and/or 2-methoxyjudaicin; coumestans, such as coumestrol, plicadin and/or wedelolactone; pterocarpans; and/or rotonoids; and/or

one or more neoflavonoids, optionally calophyllolide, coutareagenin, dalbergichromene, dalbergin and/or nivetin; and/or

one or more pterocarpanes, optionally bitucarpin A, bitucarpin B, erybraedin A, erybraedin B, erythrabysin II, erythrabysin-1, erycristagallin, glycinol, glyceollidins, glyceollins, glycyrrhizol, maackiain, medicarpin, morisianine, orientanol, phaseolin, pisatin, striatine and/or trifolirhizin.

19. The inoculant composition of any one of claims 1-18, wherein said inoculant composition comprises one or more drying agents, optionally calcium stearate, one or more clays, graphite, magnesium stearate, magnesium sulfate, powdered milk, one or more silica powders, soy lecithin and/or talc.

20. The inoculant composition of any one of claims 1-19, wherein said solid non-aqueous carrier encapsulates at least 50% of said *Penicillium* spores, optionally at least 55, 60, 65, 70, 75, 80, 85, 90 or 95% of said *Penicillium* spores.

21. A coated plant propagation material, comprising:
a plant propagation material, optionally a seed; and
a coating that covers at least a portion of an outer surface of said seed, said coating comprising the inoculant composition of any one of claims 1-20.

22. A kit, comprising:
the coated plant propagation material of claim 21; and
a container housing said coated plant propagation material.

23. A plant germinated from the coated plant propagation material of claim 21.

24. A plant part harvested from the plant of claim 23.

25. A processed product produced from the plant part of claim 24.

26. A crop comprising a plurality of the plant of claim 23.

27. A method, comprising applying the inoculation composition of any one of claims 1-20 to a plant propagation material, optionally a seed.

28. A method, comprising planting the coated plant propagation material of claim 21 in a growth medium, optionally a soil.

1/2




FIG. 1




FIG. 2

2/2




FIG.3

INTERNATIONAL SEARCH REPORT

International application No PCT/US2016/05Q647

A. CLASSIFICATION OF SUBJECT MATTER
 INV. AOICI/06 C12N1/04 C12N11/10 C12R1/80 A01N63/04
 AO1N25/00 A01N25/14 A01P21/00 C09D105/Q0
 ADD.
 According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED
 Minimum documentation searched (classification system followed by classification symbols)
 C12N A01C A01N C12R C09D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 EPO-Internal , BIOSIS, CHEM ABS Data, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	wo 2010/037228 AI (NOVOZYMES AS [DK] ; NOVOZYMES BIOLOG LTD [CA] ; HNATOWICH GARRY LAWRENCE) 8 April 2010 (2010-04-08) cited in the application claims examples	1-28
Y	US 2014/143909 AI (GREENSHIELDS DAVID [CA] ET AL) 22 May 2014 (2014-05-22) claims examples	1-28

Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 30 November 2016	Date of mailing of the international search report 08/12/2016
-------------------------------------------------------------------------------	------------------------------------------------------------------

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Galley, Carl
----------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/05Q647

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>Anonymous : "JumpStart LCO - phosphate-solubilizing inoculant and LCO Promotor Technology for corn and wheat" , Extended Label , 23 July 2013 (2013-07-23) , XP055324682, Retrieved from the Internet: URL: http://www.monsanto.com/site/col/lection/document/s/bi oag/uni ted- states/1 abel s/j sico _corn_2ozextl ab_usa_2_13133.pdf [retrieved on 2016-11-30] the whole document</p>	1-28
Y	<p>TIMOTHY FRIESEN ET AL: "Experimental determination of viability loss of Penicillium bilaiae conidia during convective air-drying" , APPLIED MICROBIOLOGY AND BIOTECHNOLOGY, SPRINGER, BERLIN, DE, vol . 68, no. 3, 1 August 2005 (2005-08-01) , pages 397-404, XP019331913 , ISSN: 1432-0614, DOI : 10.1007/S00253-004-1866-1 cited in the application page 397, right-hand column "Formulation product ion" ; page 398, right-hand column</p>	1-28
Y	<p>wo 2009/010561 AI (NOVOZYMES AS [DK] ; MAZEAUD ISABELLE [FR] ; JOERGENSEN KAARE [DK] ; PEDER) 22 January 2009 (2009-01-22) claims examples</p>	1-28
Y	<p>NAN FU ET AL: "Towards a maximal cell survival in convective thermal drying processes " , FOOD RESEARCH INTERNATIONAL, ELSEVIER, AMSTERDAM, NL, vol . 44, no. 5, 27 March 2011 (2011-03-27) , pages 1127-1149 , XP028375002, ISSN: 0963-9969, DOI : 10.1016/J.FOODRES.2011.03.053 [retrieved on 2011-04-03] page 1128; table 1 pages 1130-1 ; table 2 Section 4.1.2.1; pages 1136-7</p>	1-28

-/--

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/05Q647

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>SOLMAZ BEHBOUDI -JOBBEHDAR ET AL: "Optimi zation of Spray-Drying Process Condi tions for the Production of Maximal ly Vi abl e Mi croencapsulated L. acidophi lus NCIMB 701748" , DRYING TECHNOLOGY. , vol . 31, no. 11, 18 August 2013 (2013-08-18) , pages 1274-1283 , XP055324405, US ISSN : 0737-3937, DOI : 10.1080/07373937.2013.788509 Materi als and Methods ; page 1275 Concl usions; page 1281</p>	1-28
Y	<p>----- OLDENHOF H ET AL: "Effect of sucrose and mal to dextrin on the physi cal properti es and survi val of air-dried Lactobaci llus bulgari cus: An in situ Fourier Transform Infrared Spectroscopy Study" , BIOTECHNOLOGY PROGRESS, AMERICAN INSTITUTE OF CHEMICAL ENGINEERS, US, vol . 21, no. 3, 1 January 2005 (2005-01-01) , pages 885-892, XP002597860, ISSN : 8756-7938, DOI : 10.1021/BP049559J [retri eved on 2005-05-18] Di scussion; page 891 Concl usions; page 891</p>	1-28
Y	<p>----- SHAHIDI S ET AL: "ENCAPSULATION OF FOOD INGREDI ENTS" , CRITICAL REVI EWS IN FOOD SCI ENCE AND NUTRITION , TAYLOR & FRANCIS, USA, vol . 33, no. 6, 1 January 1993 (1993-01-01) , pages 501-547, XP009038399 , ISSN : 1040-8398 Mal to dextrins and Corn Syrup Sol i ds; page 508</p> <p>----- -/--</p>	1-28

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2016/05Q647

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>COLACO C ET AL: "EXTRAORDINARY STABILITY OF ENZYMES DRIED IN TREHALOSE: SIMPLIFIED MOLECULAR BIOLOGY", BIOTECHNOLOGY. THE INTERNATIONAL MONTHLY FOR INDUSTRIAL BIOLOGY, NATURE PUBLISHING GROUP, US, vol . 10, no . 9, 1 September 1992 (1992 -09 -01), pages 1007 - 1011, XP00G568746 , ISSN: 0733 -222X , DOI: 10.1038/NBT0992 - 1007 the whole document</p>	1-28
Y	<p>----- W0 2009/049747 A2 (MERCK PATENT GMBH [DE] ; SMITH RAYMOND STEWART [US] ; OSBURN ROBERT M [U] 23 April 2009 (2009 -04 -23) claims</p>	13 - 18
Y	<p>----- W0 2013/0442 14 A1 (NOVOZYMES BIOLOG HOLDINGS AS [DK] ; NOVOZYMES BIOLOGICALS INC [US]) 28 March 2013 (2013 -03 -28) claims</p> <p>-----</p>	13 - 18

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2016/05Q647

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
wo 2010037228	AI	08-04-2010	AR 073718 AI 24-11-2010
			AU 2009299020 AI 08-04-2010
			AU 2014203771 AI 31-07-2014
			BR PI0919373 A2 12-01-2016
			CA 2747841 AI 08-04-2010
			EP 2344610 AI 20-07-2011
			US 2010099560 AI 22-04-2010
			US 2012322655 AI 20-12-2012
			US 2015307409 AI 29-10-2015
			US 2016229762 AI 11-08-2016
			Wo 2010037228 AI 08-04-2010
US 2014143909	AI	22-05-2014	AR 093781 AI 24-06-2015
			AU 2013344562 AI 07-05-2015
			CA 2890280 AI 22-05-2014
			CN 104797701 A 22-07-2015
			EP 2920293 AI 23-09-2015
			US 2014143909 AI 22-05-2014
			US 2016251275 AI 01-09-2016
			wo 2014078647 AI 22-05-2014
wo 2009010561	AI	22-01-2009	NONE
wo 2009049747	A2	23-04-2009	AR 068871 AI 09-12-2009
			BR PI0817934 A2 30-09-2014
			wo 2009049747 A2 23-04-2009
wo 2013044214	AI	28-03-2013	AR 088163 AI 14-05-2014
			AU 2012312009 AI 03-04-2014
			AU 2016201191 AI 17-03-2016
			CA 2849889 AI 28-03-2013
			CN 104066327 A 24-09-2014
			EP 2747568 AI 02-07-2014
			NZ 622560 A 28-10-2016
			RU 2014116238 A 27-10-2015
			US 2013096002 AI 18-04-2013
			wo 2013044214 AI 28-03-2013