



Pyrenula chlorospila (Nyl.) Arnold (Pyrenulaceae), a new addition to lichen biota of India

Suparna Biswas¹, Rebecca Daimari^{1*}, Pungbili Islary¹, Sanjeeva Nayaka², Dalip Kumar Upreti², Pranjit Kumar Sarma³

¹ Department of Botany, Bodoland University, Kokrajhar, Assam, India • SB: suparnabiswas886@gmail.com • RD: publicationBU@gmail.com • PI: pungbilii@gmail.com

² Lichenology Laboratory, CSIR – National Botanical Research Institute, Rana Pratap Marg, Lucknow, Uttar Pradesh, India • SN: nayaka.sanjeeva@gmail.com • DKU: upretidknbri@gmail.com

³ Department of Geography, Mangaldai College, Upahupara, Assam, India • PKS: prangis@gmail.com

* Corresponding author

Abstract

Climatic conditions and physical features of the Eastern Himalayan region encompasses a rich diversity of lichen biota. However, the region's lichens are still far from extensively explored. Here, we study the genus *Pyrenula* Ach. from the Dhubri district of Assam in North-East India. Twelve species of *Pyrenula* have been recorded, of which one of the species, *P. chlorospila* (Nyl.) Arnold, is a newly reported from India. Morphotaxonomic comments on this species, emphasizing its distribution, are provided.

Keywords

New record, North-East India, taxonomy

Academic editor: Marcos Kitaura | Received 8 November 2021 | Accepted 30 June 2022 | Published 12 July 2022

Citation: Biswas S, Daimari R, Islary P, Nayaka S, Upreti DK, Sarma PK (2022) *Pyrenula chlorospila* (Nyl.) Arnold (Pyrenulaceae), a new addition to lichen biota of India. Check List 18 (4): 773–777. <https://doi.org/10.15560/18.4.773>

Introduction

The number of lichens known from India continues to increase and currently 2963 species in 468 genera and 84 families are known (Sinha 2021). India has rich diversity of pyrenocarpous lichens, with about 382 species in 49 genera and 12 families. The warm, moist conditions in the Eastern Himalaya, including Assam, favours a rich diversity of pyrenocarpous lichens of about 295 species (Mishra et al. 2020). Generally, these lichens grow on tree bark, rocks, soil, and leaves. Trees having smooth, shaded bark are the most preferred. Among the pyrenocarpous lichens, the family Pyrenulaceae is dominant

with *Pyrenula* Ach. the dominant genus. Aptroot (2012) listed 745 species of *Pyrenula* worldwide. The genus is characterized by an olivaceous green or brownish to yellowish thallus. Pseudocyphellae and maculae may be or may not be present. The ascocarp is of the perithecia type, which may be covered by the thallus or naked in upper part solitary or rarely 2 or 3 confluent. The peridium is carbonized and either spreading laterally or not. The ostioles are apical or lateral, and the paraphyses are simple and rarely branched. Each asci bears eight spores. Ascospores are brown and transversely septate or

muriform, with or without lichexanthone and anthraquinones (Awasthi 1991, Ingle 2018). Upreti (1990, 1991a, 1991b, 1992, 1993) made extensive revisionary studies on the *Pyrenula* of India and described several new species, as well as reports of species from the country. Jagadesh Ram et al. (2005) newly described *Pyrenula subcylindrica* Jagadesh & Upreti from India. Thus, India is represented by 83 species of *Pyrenula* and greatest diversity of species is reported from the Western Ghats and the Eastern Himalayan Region (Mishra et al. 2020; Rajaprabu et al. 2021). Among the northeastern states, Assam is one of the states with significant number of lichen studies and novel species reports from India. Altogether, 525 species of lichens have been reported from Assam, of there are 59 *Pyrenula* species (Daimari 2016; Gupta and Sinha 2018; Gogoi et al. 2019, 2020; Behera et al. 2021). Gupta and Sinha (2018) compiled lichens from the state and mentioned *P. acutalis* R.C. Harris found in the Dhubri district as part of an ongoing study.

Methods

Lichen specimens were collected from 15 locations in the Dhubri district in Assam state. The specimens were collected from the bark of trees, then air dried and stored in acid free paper packets. They were identified morphologically, anatomically and chemically. The morphological characters were studied under a Leica EZ4W stereozoom microscope. For examining anatomical characters, thin sections of the perithecia were mounted in water and observed under a Leica DM 750 compound microscope. The presence of chemical substances was analyzed by performing thin-layer chromatography (Orange et al. 2001) and colour tests

using KOH (K), para-phenylenediamine (P) and Calcium hypochlorite (C) solutions. Lichen thallus were also observed under ultraviolet light. The specimens were identified to species following Aptroot (2012) and Upreti (1998) and deposited in the Bodoland University Botanical Herbarium, (BUBH), Department of Botany, Bodoland University (Kokrajhar, Assam, India). A set of voucher specimen is deposited in the herbarium of CSIR–National Botanical Research Institute, LWG; Lucknow Uttar-Pradesh, India

Results

Pyrenula chlorospila (Nyl.) Arnold

Figure 2A–C

Specimens examined. INDIA – Assam • Dhubri district, Bidyadabri part v; 26°26'11"N, 089°47'11"E; 34m alt; 21. XII. 2020; S. Biswas & P. Biswas leg.; on the bark of *Lannea* sp.; 2020-0280 (BUBH) • Dhubri district, Debotar hasdaha part 4; 26°05'79"N, 089°83'083"E; 30.23 m alt; 22. XI. 2020; S. Biswas & P. Biswas leg.; on the bark of *Lannea* sp.; 2020-0400 (BUBH), 61701 (LWG).

Description. Thallus crustose, corticolous, yellowish; pseudocyphellae present; maculae absent; perithecia numerous, black; ascomata 0.15–0.20 mm in diameter; ostiole white, apical to lateral; peridium black, carbonized, spreading laterally; hamathecium not interspersed, ; hymenium clear; asci 8-spored; ascospores brown, transversely 3-septate; middle locules diamond-shaped and terminal locules triangular with their base towards the end, 26–30 × 10–12 µm.

Chemistry: Thallus K–, C–, P–, TLC–no chemical present.




Figure 1. Geographic distribution of *Pyrenula chlorospila* around the world. Yellow circles: occurrence data from GBIF. Red circle: new record of *P. chlorospila* in Assam, India.




Figure 2. *Pyrenula chlorospila* from Dhubri district, Assam, India. **A.** Habitus with perithecia. **B.** Vertical section of ascomata. **C.** Ascospores (26–30 μm).

Ecology. The species was found growing on smooth bark of *Lannea* sp.

Remarks. *Pyrenula chlorospila* is similar with *P. macrospora* (Degel.) Coppins & P. James. (Weerakoon et al. 2012), in its thallus colour, presence of pseudocyphellae on the thallus and absence of chemicals in TLC. Both of the species luxuriously grow on smooth bark of the trees. On the other hand, both of the species differs with their ascomata size. *Pyrenula chlorospila* has 0.1–0.4 mm diameter ascomata but *P. macrospora* has ascomata size approximately 0.5–0.9 mm in diameter (Orange 2013). Globally, *P. chlorospila* is reported from Denmark, France, Germany, Greece, Iran, Ireland, Isle of Man, Malta, the Netherlands, Russian Federation, Spain, Ukraine (GBIF 2022) (Fig. 1).

Discussion

Pyrenula species mostly prefer to grow on the smooth bark of *Alnus*, *Quercus*, and *Ficus* species (Ingle et al. 2018), and in India, most *Pyrenula* species have been found on the smooth bark of *Lannea* sp. and some on the bark of species of *Artocarpus*, *Mangifera*, *Michelia*, *Polyalthia*, and *Ricinus*. Based on the literature (Awasthi

1961; Rout et al. 2005, 2010, 2012; Singh and Sinha 2010; Das et al. 2012, 2013; Gupta et al. 2013; Daimari et al. 2014; Dey et al. 2015; Choudhury et al. 2016; Gupta and Sinha 2018; Gogoi et al. 2019; Joseph et al. 2020; Behera et al. 2021), our record of *P. chlorospila* represents a new addition to the lichen biota of India. With this addition, 12 species of *Pyrenula* have been recorded from Dhubri district (Table 1). As the district is mainly plain with deciduous vegetation, crustose lichens are dominant, although a few foliose genera, *Dirinaria*, *Pyxine*, and *Parmotrema*, also occur.

Acknowledgements

We are thankful to the Botany Department, Bodoland University for providing the facilities to carry out our research and the Director of CSIR – National Botanical Research Institute, Lucknow for permitting access to the LWG herbarium and library. We are also thankful to Mr. Sanswring Basumatary for his help and suggestions. Rebecca Daimari is thankful to Department of Science and Technology – Science and Engineering Research Board, New Delhi, India for financial assistance under the EMEQ scheme (EEQ/2019/000547).

Table 1. Distribution of *Pyrenula* species in different localities of Dhubri district, Assam, India. Locations: 1 = Barobaluchar, 2 = Bidyadabri part 5, 3=Bhalukmari, 4 = Bhelupara part 2, 5 = Brahmin para, 6 = Debotar hasdaha part 4, 7 = Dhubri town, 8 = Falimari, 9 = Gaurangtari part 2, 10 = Gauripur, 11 = Golakganj, 12 = Hatipota part 2, 13 = Khajurbari part 1, 14 = Khalilpur, 15 = Kherbari part 2. (+) presence and (–) absence.

<i>Pyrenula</i> species	Locations														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
<i>P. acutalis</i> R.C. Harris	–	–	–	–	+	–	–	–	+	–	–	–	–	–	–
<i>P. anomala</i> (Ach.) Vain.	–	–	–	–	–	–	–	–	–	–	–	–	+	–	–
<i>P. approximans</i> (Kremp.) Müll. Arg.	–	–	–	–	–	–	–	–	–	–	–	–	+	–	–
<i>P. breutelii</i> (Müll. Arg.) Aptroot	–	–	–	–	–	–	–	–	–	–	–	–	–	+	–
<i>P. chlorospila</i> (Nyl.) Arnold	–	+	–	–	–	+	–	–	–	–	–	–	–	–	–
<i>P. leucostoma</i> Ach.	–	–	–	–	–	–	+	–	–	+	–	–	–	–	–
<i>P. leucotrypa</i> (Nyl.) Upreti	–	–	–	+	–	–	–	–	–	–	–	–	–	–	–
<i>P. oculata</i> Ajay Singh & Upreti	+	–	–	–	–	–	–	–	–	–	–	–	–	–	–
<i>P. quassicola</i> (Fée)	–	–	–	–	–	–	–	–	–	–	+	–	–	–	+
<i>P. sublaevigata</i> (Patw. & Makhija) Upreti	–	–	–	–	–	–	–	+	–	–	–	–	–	–	–
<i>P. submastophora</i> Ajay Singh & Upreti	–	–	–	–	–	–	–	–	–	–	–	+	–	–	–
<i>P. wrightii</i> (Müll. Arg.) R.C. Harris	–	–	+	–	–	–	–	–	–	–	–	–	–	–	–

Authors' Contributions

Conceptualization: SB. Data curation: SB, PI. Formal analysis: SN, DKU, PKS. Funding acquisition: RD.

References

- Aptroot A (2012) A world key to the species of *Anthracotheceum* and *Pyrenula*. The Lichenologist 44 (1): 5–53. <https://doi.org/10.1017/s0024282911000624>
- Awasthi DD (1961) Some foliose and fruticose lichens from Assam and North-East frontier agency of India. Proceedings of the Indian Academy of Sciences, Section B 54 (1): 24–44. <https://doi.org/10.1007/bf03053852>
- Awasthi DD (1991) A key to the microlichens of India, Nepal and Sri Lanka. Bibliotheca Lichenologica 40: 1–336. <https://doi.org/10.1006/lich.1993.1017>
- Behera PK, Nayaka S, Upreti DK, Chauhan RJ (2021) New distributional records to lichen biota of Assam, India. Indian Forester 147 (4): 400–404.
- Choudhury MP, Sarma M, Nayaka S, Upreti DK (2016) Distribution of Lichens on few ancient monuments of Sonitpur district, Assam, North East India. International Journal of Biodiversity and Conservation 8(11): 291–296. <https://doi.org/10.5897/ijbc2016.0971>
- Daimari R, Hazarika N, Hoque RR, Nayaka S, Upreti DK (2014) New records of epiphytic lichens from three districts of Assam, India. Indian Forester 140 (8): 807–811.
- Daimari R (2016) Epiphytic lichens, their diversity, distribution and air pollution sensitivity in Sonitpur district, Assam. PhD thesis, Department of Environmental Science, Tezpur University, Assam, India, 259 pp.
- Das P, Joshi S, Rout J, Upreti DK (2012) Exploration of homegardens as important lichen conservation areas in Dargakona Village of Southern Assam, Northeast India. Journal of Functional and Environmental Botany 2 (2): 87–95. <https://doi.org/10.5958/j.2231-1742.2.2.013>
- Das P, Joshi S, Rout J, Upreti DK (2013) Impact of anthropogenic factors on abundance variability among Lichen species in southern Assam, North East India. Tropical Ecology 54 (1): 65–70.
- Dey AK, Mishra GK, Rout J, Upreti DK (2015) An enumeration of epiphytic lichens from Hojai sub-division of Nagaon district, Assam, India. International Journal of Advanced Research in Biological Sciences 2 (10): 111–115.
- Gogoi R, Joseph S, Nayaka S, Yasmin F (2019) Additions to the lichen biota of Assam state, India. Journal of Threatened Taxa 11(6): 13765–13781. <https://doi.org/10.11609/jott.4642.11.6.13765-13781>
- Gogoi R, Joseph S, Choudhury MP, Nayaka S (2020) Crustose lichens new to India. Mycotaxon 135 (3): 657–663.
- Gupta P, Sinha GP, Solanki CM (2013) Epiphytic lichens in tea gardens of Assam, India. Indian Journal of Forestry 36 (2): 279–284.
- Gupta P, Sinha GP (2018) Lichen flora of Assam. Bishen Singh Mahendra Pal Singh, Dehradun, India, 274 pp.
- Ingle KK, Uppadhyay V, Nayaka S, Trivedi S, Sahoo D (2018) New records and an updated key of *Pyrenula* from India. Cryptogam Biodiversity and Assessment (Special Volume 2018): 37–46.
- Jagadesh Ram TAM, Upreti DK, Sinha GP (2005) *Pyrenula subcylindrica*, a new pyrenocarpous lichen from India. The Lichenologist 37(2): 109–110. <https://doi.org/10.1017/s0024282904014604>
- Joseph S, Nayaka S, Sinha GP (2020) Additions to the bibliography of Indian lichens in the years 2018 and 2019. Cryptogam Biodiversity and Assessment 4 (2): 7–13.
- Mishra GK, Nayaka S, Upreti DK (2020) Distribution, biomonitoring and conservation studies of pyrenocarpous lichens in India. G-Journal of Environmental Science and Technology 7 (5): 54–59.
- Orange AP, James W, White FJ (2001) Microchemical methods for the identification of lichens. British Lichen Society, London, UK, 101 pp.
- Orange A (2013) British and other pyrenocarpous lichens. Version 2. Department of Biodiversity and Systematic Biology, National Museum of Wales, Cardiff, UK, 250 pp. <https://museum.wales/media/13849/Orange-A-2013-British-and-other-pyrenocarpous-lichens.pdf>. Accessed on 2022-7-4.
- Rajaprabu N, Ponmurugan P, Mishra GK (2021) A new record of pyrenocarpous lichen to the Indian biota. Journal of Threatened Taxa 13 (1): 17607–17610.
- Rout J, Rongmei R, Das P (2005) Epiphytic lichen flora of a pristine habitat (Nit Campus) in Southern Assam, India. Phytotaxonomy 5: 117–119.
- Rout J, Das P, Upreti DK (2010) Epiphytic lichen diversity in a Reserve Forest in southern Assam, northeast India. Tropical Ecology 51(2): 281–288.
- Rout J, Singha AB, Upreti DK (2012) Lichen flora on Betel Nut (*Areca catechu*) palm tree from a pristine habitat in Southern Assam, India. International Journal of Plant Research 25 (1): 198–201.
- Sinha KP, Sinha GP (2010) Indian lichens: an annotated checklist. Botanical Survey of India, Ministry of Environmental and Forests, Kolkata, India, 571pp.
- Sinha GP (2021) An overview of the current status of lichen diversity in India and identification of gap areas. In: Nayaka S, Joseph S, Mishra GK (Eds.) Accelerate Vigyan Karyashala Workshop Manual, integrative systematics of lichens – identification to bio-prospecting. CSIR –National Botanical Research Institute, Lucknow, India, 5 pp.

- Upreti DK (1990) Lichen genus *Pyrenula* in India: I *Pyrenula subducta* spore type. Journal of the Hattori Botanical Laboratory 68: 269–278. https://doi.org/10.18968/jhbl.68.0_269
- Upreti DK (1991a) Lichen genus *Pyrenula* from India: The species with spores of *Pyrenula brunnea* type. Bulletin de la Société Botanique de France. Letters Botaniques 138 (3): 241–247. <https://doi.org/10.1080/01811797.1991.10824926>
- Upreti DK (1991b) Lichen genus *Pyrenula* from India: IV. *Pyrenula cayennensis* spore type. Cryptogamie Bryologie Lichenologie 12 (1): 41–46.
- Upreti DK (1992) Lichen genus *Pyrenula* from India: VII. *Pyrenula mastophora* spore type. Feddes Repertorium 103 (3–4): 279–296.
- Upreti DK (1993) Lichen genus *Pyrenula* from India: II. *Pyrenula camptospora* spore type, III. *Pyrenula pinguis* spore type. Acta Botanica Gallica 140 (5): 519–523. <https://doi.org/10.1080/12538078.1993.10515628>
- Upreti DK (1998) A key to the lichen genus *Pyrenula* from India, with nomenclatural notes. Nova Hedwigia 66: 557–576.
- Weerakoon G, Aptroot A, Lumbsch HT, Wolseley PA, Wijeyaratne CS, Gueidan C (2012) New molecular data on Pyrenulaceae from Sri Lanka reveal two well-supported groups within this family. The Lichenologist 44 (5): 639–647. <https://doi.org/10.1017/S0024282912000333>