

# Index

Indexed items are referenced by section number, which consists of the chapter number followed by the number of a major topic, such as a disease or pest description. **The number, in bold italic, of each illustration is the same as that of the corresponding section in the text;** for sections having more than one illustration, letters follow the section number; for example, the color illustrations for Colorado potato beetle (text section 16.44) are figures *16.44a* to *16.44d*. For text figures (line drawings and halftones), the figure number includes the letter T (e.g. *16.44T1*). To aid in finding items in the text, the running heads for each two-page spread identify the inclusive section numbers beginning on those pages; the number in the running head for a left-hand page is that of the first section beginning on that page, while the running head for the facing (right-hand) page carries the number of the last section beginning on that page.

## A

- Abutilon theophrasti*, 2.3
- Acalymma vittatum*, 9.21; 22.35
- acid scab
  - on potato, 16.5
- Acremonium apii*, 7.2
- Acrobelloides* spp., 26.28
- Actebia fennica*, 18.35
- Actinomyces scabies*
  - (see *Streptomyces scabies*)
- actinomycetes
  - as vegetable pathogens, 2.3; 26.15
  - (see bacteria)
- Aculops lycopersici*, 25.31
- Acyrtosiphon pisum*, 15A.14; 23.17
- Agriotes mancus*, 16.50; **12.21T1**
- Agriotes obscurus*, 16.50
- Agrobacterium radiobacter*, 3.5; 6.3
- Agrobacterium rhizogenes*, 6.3
- Agrobacterium rubi*, 6.3
- Agrobacterium tumefaciens*, 6.3; 17.1
- Agropyron repens*, 2.3
- Agrotis ipsilon*, 18.35
- Agrotis orthogonia*, 18.35
- Agrotis subterranea*, 18.35
- Albugo candida*, 8.15; 10.10
- Albugo crueiferarum*
  - (see *Albugo candida*)
- Albugo occidentalis*, 5.9
- Aleochara bilineata*, 13.26
- alfalfa looper
  - larva, **8.38**
  - on crucifers, 8.38
  - on pea, 15A.15
- alfalfa mosaic
  - on pea, 15A.9
  - on pepper, 18.20
  - on potato, 16.24
- alfalfa mosaic virus
  - (see alfalfa mosaic)
- alfalfa sprout rot, 27.1
- allelopathy, 3.6
- Alternaria alternata*, 9.8; 18.8; 25.9; **18.8f,g**
- alternaria blight
  - on ginseng, 20.1 ; **20.1a-c**
- Alternaria brassicae*, 8.5
- Alternaria brassicicola*, 8.5
- Alternaria cucumerina*, 9.8
- Alternaria cucurbitae*

(see *Ulocladium cucurbitae*)

*Alternaria dauci*, 6.5; 10.4  
  conidium, **6.5b**

alternaria diseases  
  on crucifers, 8.5

alternaria fruit rot  
  on pepper, 18.8; **18.8g**  
  on tomato, 18.8; **18.8f**

alternaria leaf blight  
  on carrot, 6.5; **6.5a**  
  on greenhouse cucumber, 22.12; **22.12a,b**

alternaria leaf spot  
  on Brussels sprouts, **8.5b**  
  on cauliflower, **8.5a**  
  on crucifers, 8.5; **8.5a,b**  
  on cucumber, **9.8a**  
  on cucurbits, 9.8; **9.8a**  
  on parsley, 10.4

*Alternaria panax*, 20.1; 20.3; **20.3a**  
  conidia, **20.1c**

*Alternaria porri*, 13.9

*Alternaria radicina*, 6.7; 10.3; **6.7**

*Alternaria raphani*, 8.5

*Alternaria solani*, 16.8; 18.8; 18.42; 25.9

*Alternaria* spp., 10.10; 22.12; **16.8c**

*Alternaria tenuissima*, 9.8

aluminum toxicity  
  on bean, 15B.12

amitrole  
  on potato, **16.34c**  
  (see herbicide injury)

*Amaranthus graecizans*, 2.3

*Amaranthus retroflexus*, 2.3

*Amblyseius barkeri*  
  a beneficial mite, 3.7; 22.34

*Amblyseius cucumeris*  
  adult, **22.34h**  
  a beneficial mite, 3.7; 25.31  
  on western flower thrips, 22.34; 24.14

*Ambrosia artemisiifolia*, 2.3

*Amplimerlinius* spp.  
  as vegetable pests, 2.3

*Anagrapha falcifera*, 7.22

*Anaphes sordidatus*, 6.24

*Anasa tristis*, 9.22

Andean potato latent virus  
  a foreign pathogen, 3.10

Andean potato mottle virus  
  a foreign pathogen, 3.10

angular leaf spot  
  on cucumber, **9.1a,b**  
  on cucurbits, 9.1; **9.1a,b**  
  on greenhouse cucumber, 22.1; **22.1**

*Anthonomus eugenii*, 3.10; 24.13

anthracnose  
  on bean, 15B.2; **15B.2a-d**  
  on coriander, 10.10  
  on cucumber, **9.3b**  
  on cucurbits, 9.3; **9.3a-c**  
  on eggplant, 18.6  
  on greenhouse cucumber, 22.3  
  on greenhouse lettuce, 23.4

- on lettuce, 11.4; **11.4**
- on muskmelon, **9.3a,c**
- on pepper, 18.6; **18.6b**
- on tomato, 18.6; **18.6a,c**

ants

- as beneficial organisms, 3.7

*Apanteles glomeratus*  
(see *Cotesia glomerata*)

*Apanteles rubeculus*  
(see *Cotesia rubecula*)

*Aphaereta pallipes*, 13.26

*Aphanocladium album*, 26.10

aphanocladium cap spot

- on mushroom, 26.10

*Aphanomyces cochlioides*, 5.2

*Aphanomyces euteiches*, 3.6; 15A.3

*Aphanomyces raphani*, 8.7

aphanomyces root rot

- on beet, 5.2; **5.2**
- on chard, 5.2
- on spinach, 5.2

*Aphelenchoides* spp., 26.27

*Aphidius matricariae*

- a beneficial insect, 3.7; 24.14
- on aphids, 24.12

*Aphidius* sp., **3.7s**

*Aphidoletes aphidimyza*, 3.7; 22.33; 24.12

*Aphidoletes* sp., 22.31 ; **24.12d**

aphids

- as vegetable pests, 2.2
- monitoring for, 3.2; **3.2TI**
- on asparagus, 4.9; 4.11 ; **4.9**
- on beet, 5.17
- on Brussels sprouts, **8.39a**
- on cabbage, **8.39b**
- on celery, 7.17
- on crucifers, 8.39
- on cucurbits, 9.22
- on eggplant, 18.33
- on greenhouse cucumber, 22.33
- on greenhouse lettuce, 23.17
- on greenhouse pepper, 24.12
- on greenhouse tomato, 25.30
- on herbs and spices, 10.13
- on hop, 10.13
- on lettuce, 11.24; 11.25
- on maize, 12.14; **12.16b**
- on onion, 13.28
- on parsley, 10.13
- on pea, 15 A. 14
- on pepper, 18.33
- on potato, 2.2; 16.40-16.43; **16.43T1**
- on potato, key to wingless females, 16.43; **16.43T2**
- on rhubarb, 17.13
- on tomato, 18.33
- parasite of, 3.7; **3.7s; 24.12c**
- predator of, 3.7; **3.7b-i; 24.12d**
- (see asparagus aphid)
- (see black bean aphid)
- (see buckthorn aphid)
- (see bulb and potato aphid)
- (see cabbage aphid)

(see carrot-willow aphid)  
 (see corn leaf aphid)  
 (see crescent-marked lily aphid)  
 (see foxglove aphid)  
 (see green peach aphid)  
 (see hop aphid)  
 (see lettuce aphid)  
 (see melon aphid)  
 (see pea aphid)  
 (see potato aphid)  
 (see shallot aphid)  
 (see sugarbeet root aphid)  
 (see turnip aphid)  
 (see turnip root aphid)

*Aphis abbreviata*  
 (see *Aphis nasturtii*)

*Aphis fabae*, 5.17; 16.43; 17.13; 23.17  
*Aphis gossypii*, 9.22; 16.43; 22.33; 24.15  
*Aphis nasturtii*, 16.40

apical chlorosis  
 on Jerusalem artichoke, 21.1; **21.1a,b**

*Apis mellifera*, 1.4

arabis mosaic  
 on rhubarb, 17.9  
 (see hop nettle head)

arabis mosaic virus  
 (see arabis mosaic)

*Arion ater*, 11.27

army cutworm, 18.35

army worm  
 adult, **12.12c**  
 larva, **12.12b**  
 on maize, 12.12; **12.12a,b**

army worms  
 pathogen of, 3.5  
 on maize, 12.12; 12.17  
 (see armyworm)  
 (see fall armyworm)

arracacha virus B  
 a foreign pathogen, 3.10

*Arthrobotrys* spp., 26.21

artichoke Italian latent virus  
 on endive, 11.18

*Artogeia rapae*  
 (see *Pieris rapae*)

*Aschersonia aleyrodis*  
 a beneficial pathogen, 3.5; 3.7  
 on greenhouse whitefly, 22.32; 25.27; **3.7w**

*Asclepias syriaca*, 2.3

*Ascochyta cucumis*, 22.11  
 pycnidia, **22.11a,e**  
 (see *Didymella bryoniae*)

ascochyta foot rot  
 on pea, 15A.2; **15A.2c,d**

ascochyta leaf spot  
 on rhubarb, 17.6

ascochyta leaf and pod spot  
 on pea, 15A.2; **15A.2a,b**

*Ascochyta pinodella*  
 (see *Phoma medicaginis* var. *pinodella*)

*Ascochyta pinodes*  
 (see *Mycosphaerella pinodes*)

*Ascochyta pisi*, 15A.2  
     conidia, **15A.2T1**  
*Ascochyta rhei*, 17.6  
 asparagus aphid  
     on asparagus, 4.9; **4.9**  
 asparagus beetle  
     adult, **4.10a,e**  
     egg, **4.10e**  
     larva, **4.10b**  
     on asparagus, 4.10; **4.10a,b,e**  
 asparagus beetles  
     on asparagus, 4.10; **4.10a-e**  
     (see asparagus beetle)  
     (see spotted asparagus beetle)  
 asparagus virus I  
     on asparagus, 4.7  
 asparagus virus II  
     on asparagus, 4.7  
*Aspergillus* spp., 26.16  
 aster leafhopper  
     adult, **11.23a, b**  
     on carrot, 6.22  
     on celery, 7.18  
     on lettuce, 11.23  
     on potato, 16.51  
 aster yellows  
     on carrot, 6.17; **6.17a-c**  
     on celery, 7.8; **7.8a,b**  
     on cucurbits, 9.17  
     on dill, 10.11; **10.11a,b**  
     on eggplant, 18.16  
     on greenhouse lettuce, 23.11  
     on lettuce, 11.15; **11.15a,b**  
     on onion, 13.13; **13.13**  
     on parsley, 10.11  
     on pea, 15A.9; **15A.9a**  
     on pepper, 18.16  
     on potato, 16.23; **16.23**  
     on sage, 10.11  
     on tomato, 18.16  
 aster yellows mycoplasma-like organism  
     (see aster yellows)  
*Athelia arachnoidea*  
     (see *Rhizoctonia carotae*)  
*Aulacorthum circumflexum*, 16.43  
*Aulacorthum solani*, 16.43  
*Aureobasidium zeae*  
     (see *Kabatiella zeae*)  
*Autographa californica*, 8.38  
 autotoxicity  
     on asparagus, 4.8

## **B**

*Bacillus subtilis*, 13.12  
*Bacillus thuringiensis*, 3.1; 3.5; 3.7; 3.14; 6.24; 16.44; 16.51; 18.34; 22.35; 23.18; 25.30  
     on cabbage looper, 3.5; 8.40; **3.7y,z**  
     on Colorado potato beetle, 3.5  
     on diamondback moth, 3.5; 8.42  
     on imported cabbageworm, 3.5; 8.45  
*Bacillus thuringiensis* var. *israelensis*, 26.29  
*Bacillus* spp., 16.2

bacteria, 1.3; 11.7; 22.7  
  as insect pathogens, 3.7  
  as vegetable pathogens, 2.3

bacterial blight  
  on pea, 15A.1; **15A.1a,b**

bacterial blotch  
  on mushroom, 26.1

bacterial brown spot  
  on bean, 15B.1; **15B.1a**

bacterial canker  
  on greenhouse tomato, 25.1; **25.1**  
  on tomato, 18.1; **18.1a-c**

bacterial leaf blight  
  on carrot, 6.1; **6.1a,b**

bacterial leaf spot  
  on celery, 7.1; **7.1a,b**  
  on crucifers, 8.1; **8.1a,b**

bacterial ring rot  
  an introduced disease, 3.11  
  on potato, 16.1; **16.1a-e**

bacterial soft rot  
  disease cycle, **16.2T1**  
  on alfalfa sprouts, 27.1; **27.1a,b**  
  on bean sprouts, 27.2; 27.2; **27.2T1**  
  on carrot, 6.2; **6.2**  
  on chicory, 11.1  
  on eggplant, 18.2  
  on endive, 11.1  
  on lettuce, 11.1  
  on onion, 13.2; **13.2a,b**  
  on pepper, 18.2  
  on potato, 16.2; **16.2a,b**  
  on rhubarb, 17.2  
  on tomato, 18.2; **18.2**

bacterial speck  
  on greenhouse tomato, 25.2  
  on tomato, 18.3; **18.3a,b**

bacterial spot  
  on pepper, 18.4; **18.4d-f**  
  on tomato, 2.2; 18.4; **18.4a-c**

bacterial stem rot  
  on greenhouse tomato, 25.3; **25.3**

bacterial wilt  
  on chicory, 11.1  
  on cucumber, **9.2a,b**  
  on cucurbits, 9.2; **9.2a,b**  
  on eggplant, 18.5  
  on endive, 11.1  
  on greenhouse cucumber, 22.2  
  on lettuce, 11.1  
  on pepper, 18.5  
  on tomato, 18.5

*Bacterium rhapontici*  
  (see *Erwinia rhapontici*)

baldhead  
  on bean, 15B.13; **15B.13a**

barley  
  a weed pest, 2.3

barnyard grass  
  a weed pest, 2.3; **2.3a**

basal rot  
  on onion, 13.4; **13.4**

bean aphid  
(see black bean aphid)

bean common mosaic  
on bean, 15B.10; **15B.10**

bean common mosaic virus  
(see bean common mosaic)

bean yellow mosaic  
on bean, 15B. 11 ; **15B.11**  
on pea, 15A.9; **15A.9b**

bean yellow mosaic virus  
(see bean yellow mosaic)

*Beauveria bassiana*  
a beneficial pathogen, 3.7; 6.24  
on Colorado potato beetle, 3.7

*Beauveria* spp.  
as beneficial pathogens, 3.7

bees  
as pollinators of vegetable crops, 1.4  
(see bumblebees)  
(see honeybees)  
(see leafcutting bees)  
(see orchard bees)  
(see squash bees)

beet leafhopper  
adult, 5.15; 5.15T1  
on beet, 5.15

beet leafminer  
adult, **5.17d**  
on beet, 5.17; **5.17e**  
on chard, **5.17f**

beetles  
as beneficial organisms, 3.7  
(see asparagus beetle)  
(see blister beetles)  
(see common June beetle)  
(see Colorado potato beetle)  
(see common June beetle)  
(see flea beetles)  
(see ground beetles)  
(see Japanese beetle)  
(see Mexican bean beetle)  
(see red turnip beetle)  
(see rove beetles)  
(see sap beetles)  
(see spotted asparagus beetle)  
(see spotted cucumber beetle)  
(see white grubs)

beet mild yellowing  
on endive, 11.18

beet mild yellowing virus  
(see beet mild yellowing)

beet pseudo-yellows  
on greenhouse cucumber, 22.19; **22.19**

beet pseudo-yellows virus  
(see beet pseudo-yellows)

beet webworm  
adult, **5.17c**  
larvae and pupa, **5.17a,b**  
on beet, 5.17

beet western yellows  
on endive, 11.18  
on greenhouse lettuce, 23.15

- on lettuce, 11.18
- beet western yellows virus
  - (see beet western yellows)
- Bembidion* spp., 13.26
- Bemisia tabaci*, 3.10
- beneficial insects
  - for management of insect pests, 3.7; **3.7a-v**
- beneficial mites
  - for management of insect pests, 3.7
  - for management of mite pests, 3.7
- beneficial pathogens
  - for management of insect pests, 3.7; **3.7w-z**
  - for management of plant pathogens, 3.5
- beneficial plants
  - for pest management, 3.6
- Benincasa cerifera*
  - a resistant rootstock, 1.2
- Benincasa hispida*
  - a resistant rootstock, 1.2
  - a virus indicator, 22.22
- bentazon
  - on pea, **15A. 11a**
  - (see herbicide injury)
- Beta vulgaris*
  - a virus indicator, 4.7
- Bigonicheto spinipennis*
  - (see *Triarthria setipennis*)
- big vein
  - on greenhouse lettuce, 23.12
  - on lettuce, 11.16; **11.16**
- big-vein virus
  - (see big vein)
- bindweed, field
  - a weed pest, 2.3
- biological control of pests and diseases
  - in commercial vegetable crops, 3.5
  - in home gardens, 3.14
- BioMal
  - a mycoherbicide, 3.13
- birdcherry-oat aphid, 12.14
- black army cutworm, 18.35
- black bean aphid
  - on beet, 5.17
  - on greenhouse lettuce, 23.17
  - on potato, 16.43; **16.43TI**
  - on rhubarb, 17.13
- black blotching
  - on crucifers, 8.32; **8.32a**
- black cutworm, 18.35
  - larva, **6.25a**
- black dot
  - on potato, 16.6; **16.6a,b**
- blackheart
  - on celeriac, 7.10
  - on celery, 7.10; **7.10a,b**
  - on potato, 16.30; **16.30**
- black joint
  - on celery, 7.21
- black leaf spot
  - on crucifers, 8.5
- blackleg
  - on cabbage, **8.6a,b**



- on crucifers, 8.6; **8.6a-d**
- on potato, 16.3; **16.3a-c**
- on rutabaga, **8.6c,d**
- black midrib
  - on cabbage, **8.27**
  - on crucifers, 8.27; **8.27**
- black mold
  - on carrot, 6.6
- black root
  - on crucifers, 8.7; **8.7**
  - on radish, **8.7**
- black root rot
  - on bean, 15B.4; **15B.4**
  - on beet, 5.2
  - on carrot, 6.6; **6.6a-c**
  - on chard, 5.2
  - on chicory, 11.5; **11.5**
  - on greenhouse cucumber, 22.4; **22.4a,b**
  - on spinach, 5.2
- black rot
  - on broccoli, **8.2d**
  - on cabbage, **8.2a,b**
  - on carrot, 6.7; **6.7**
  - on crucifers, 8.2; **8.2a-f**
  - on greenhouse cucumber, 22.5
  - on rutabaga, **8.2e**
- black scurf
  - on potato, 16.15
- black slug
  - on lettuce, 11.27; **11.27a**
- black sooty mold
  - on greenhouse cucumber, **22.32b**
  - on greenhouse pepper, **24.12e**
  - on greenhouse tomato, **25.27g**
  - (see aphids)
  - (see greenhouse whitefly)
- black speck
  - on cabbage, 8.28; **8.28a,b**
  - on cauliflower, 8.17; **8.17**
- black spot
  - on cabbage, **8.32b**
  - on crucifers, 8.32; **8.32b**
- black swallowtails
  - larva, **10.15a; 14.7**
  - on dill, 10.15; **10.15a**
  - on parsley, 10.15
  - on parsnip, 14.7
- black whisker mold
  - on mushroom, 26.16; **26.16**
- blight
  - on dill, 10.10
- blister beetles
  - on potato, 16.51
- BLIGHT-ALERT
  - a predictive program for onion, 13.5
  - (see botrytis leaf blight)
- BLITECAST
  - a predictive program for potato, 3.1
  - (see late blight)
- blossom-end rot
  - on eggplant, 18.21
  - on greenhouse pepper, 24.9; **24.9**

- on greenhouse tomato, 25.23; **25.23**
  - on pepper, 18.21; **18.21c,d**
  - on tomato, 18.21; **18.21a,b**
- blotchy ripening
  - on greenhouse tomato, 25.25; **25.25a**
  - on tomato, 18.22; **18.22a,b**
- blow flies
  - as pollinators of vegetable crops, 1.4
- Bombus terrestris*, 1.4
- borers
  - (see European corn borer)
  - (see potato stem borer)
  - (see sesiid borers)
  - (see squash vine borer)
  - (see stalk borer)
  - (see stem borers)
- boron deficiency
  - on beet, 5.11
  - on cauliflower, **8.23a,b**
  - on celeriac, 7.12
  - on celery, 7.12
  - on crucifers, 8.23; **8.23a-d**
  - on greenhouse cucumber, 22.26; **22.26a**
  - on pea, 15A.10
  - on rutabaga, **8.23c,d**
- boron toxicity
  - on bean, 15B.12
  - on pea, 15A.10; **15A.10**
- BOTCAST
  - a predictive program for onion, 3.1; 13.5
  - (see botrytis leaf blight)
- Botryotinia allii*
  - (see *Botrytis byssoidea*)
- Botryotinia fuckeliana*, 2.3
  - (see *Botrytis cinerea*)
- Botryotinia squamosa*
  - (see *Botrytis squamosa*)
- Botryotrichum piluliferum*, 26.19
- Botrytis aclada*, 13.5; 13.7
- Botrytis allii*
  - (see *Botrytis aclada*)
- botrytis blight
  - on asparagus, 4.1 ; **4.1a,b**
  - on ginseng, 20.2; **20.2**
- Botrytis byssoidea*, 13.7
- Botrytis cinerea*, 2.2; 2.3; 4.1; 9.7; 10.10; 11.10; 13.5; 15B.3; 16.10; 17.5; 18.11; 20.2; 22.10; 23.9; 24.3; 25.12; **11.10c**
  - conidia, **11.10f**
  - sclerotia, **11.10d**; **22.10d**
  - spores, **9.7**; **22.10b**
  - sporulation, **15B.3a**; **22.10.a**; **25.12a**
- Botrytis fulva*, 26.20
- botrytis leaf blight
  - on onion, 13.5; **13.5a,b**
- Botrytis* spp.
  - parasite of, 3.5
  - sclerotia, **13.7c**
- Botrytis squamosa*, 13.5; 13.7
- bottom rot
  - on chicory, 11.6
  - on crucifers, 8.13
  - on eggplant, 18.21
  - on endive, 11.6

- on greenhouse lettuce, 23.5
- on lettuce, 11.6; **11.6a,b**
- on pepper, 18.21
- on tomato, 18.21

*Brachycorynella asparagi*, 4.9

*Bracon montrealensis*, 8.46

*Bradysia* spp., 22.31

*Brassica kaber*, 2.3

*Brassica* spp.

- as beneficial plants, 3.6
- as virus indicators, 25.21

*Bremia lactucae*, 3.4; 11.8; 23.7

- sporulation, **11.8a**

*Brevicoryne brassicae*, 8.39; 23.17

broad bean wilt

- on parsley, 10.12

broad bean wilt virus

- (see broad bean wilt)

brown bead

- on broccoli, **8.18**
- on crucifers, 8.18; **8.18**

brown blotch

- on mushroom, 26.1 ; **26.1**

brown eye

- on potato, 16.4

brown garden snail

- an introduced pest, 3.11
- adult shell, **3.11e**
- on lettuce, 11.27

brown heart

- on crucifers, 8.23

brown mold

- on mushroom, 26.17

brown plaster mold

- on mushroom, 26.19; **26.19**
- (see plaster molds)

brown root rot

- on greenhouse tomato, 25.6

brown rot

- on chicory, 11.3
- on eggplant, 18.5
- on endive, 11.3
- on lettuce, 11.3
- on pepper, 18.5
- on tomato, 18.5

brown spot

- on celery, 7.2; **7.2**

*Bruchus pisorum*, 15A.15

buckthorn aphid

- adult, 16.40a,b
- on potato, 16.40; 16.40a

buckwheat, wild

- a weed pest, 2.3; **2.3b**

bulb and potato aphid

- on potato, 16.43; **16.43TI**

bumblebees

- as pollinators of vegetable crops, 1.4

button-hole rot

- on potato, 16.13

butt rot

- on chicory, 11.3
- on endive, 11.3

on greenhouse lettuce, 23.1  
on lettuce, 11.3

## C

- cabbage aphid
  - on Brussels sprouts, **8.39a**
  - on crucifers, 8.39; **8.39a**
  - on greenhouse lettuce, 23.17
- cabbage butterfly
  - (*see* imported cabbageworm)
- cabbage flea beetle
  - on crucifers, 8.44
- cabbage looper
  - adult, **8.40f**
  - egg, **8.40b**
  - larva, **8.40c,d**
  - pupa, **8.40e**
  - on broccoli, 2.2
  - on Brussels sprouts, 2.2
  - on cabbage, 2.2; **8.40a**
  - on cauliflower, 2.2
  - on celery, 7.22
  - on crucifers, 8.40; **8.40a-f**
  - on greenhouse lettuce, 23.18
  - on greenhouse tomato, 25.30
  - on lettuce, 11.26
  - on pea, 15A.15
  - on tomato, 18.37
  - pathogens of, **3.7x-z**
- cabbage maggot
  - adult, **8.41g**
  - egg, **8.41d**
  - larva, **8.41e**
  - pupa, **8.41f**
  - on cabbage, 2.2; **8.41b**
  - on crucifers, 8.41; **8.41a-g**
  - on radish, **8.41a**
  - on rutabaga, **8.41c**
- cabbage root maggot
  - (*see* cabbage maggot)
- cabbage white
  - (*see* imported cabbageworm)
- Caenorhabditis* spp., 26.28
- calcium deficiency
  - on greenhouse cucumber, 22.26
  - on greenhouse lettuce (*see* tipburn, 23.16; **23.16**)
  - on pea, 15A.10
  - (*see also* tipburn)
- calico
  - on potato, 16.24; **16.24**
- Calosoma frigidum*, 8.49
- canker
  - on hop, 10.1
  - on potato, 16.21
- Capsella bursa-pastoris*, 2.3; 8.10; 8.46; 16.26
- Carpophilus* sp., 10.15
- carrot motley dwarf
  - on parsley, 10.12; **10.12**
- carrot mottle virus
  - (*see* carrot motley dwarf)
- carrot red leaf virus

(*see* carrot motley dwarf)

carrot rust fly  
adult, **6.23e**  
larvae and pupa, **6.23a,c**  
monitoring for, **3.2T1**  
on carrot, 2.2; 6.23; **6.23a,b,d**  
on celeriac, 7.19  
on celery, 7.19  
on parsley, 10.15; **10.15b**  
on parsnip, 14.5

carrot weevil  
adult, **6.24c**  
larva and pupa, **6.24d**  
monitoring for, **3.2T2**  
on carrot, 6.24; **6.24a,b**  
on celeriac, 7.20  
on celery, 7.20; **7.20**  
on parsnip, 14.6

carrot-willow aphid  
on parsley, 10.13

caterpillars  
on bean, 15B.17; 15B.19  
on beet, 5.17  
on carrot, 6.25  
on celery, 7.22  
on crucifers, 8.38; 8.40; 8.42; 8.45; 8.46  
on cucurbits, 9.22  
on dill, 10.15; **10.15a**  
on ginseng, 20.11  
on greenhouse lettuce, 23.18  
on greenhouse cucumber, 22.35  
on greenhouse pepper, 24.15  
on greenhouse tomato, 25.30  
on herbs and spices, 10.15  
on Jerusalem artichoke, 21.6  
on lettuce, 11.26  
on maize, 12.12; 12.13; 12.16; 12.17; 12.22  
on parsnip, 14.7  
on pea, 15A.15  
on pepper, 18.35; 18.36  
on potato, 16.47; 16.51  
on rhubarb, 17.13  
on tomato, 18.35; 18.37  
(*see* alfalfa looper)  
(*see* armyworms)  
(*see* beet webworm)  
(*see* black swallowtails)  
(*see* cabbage looper)  
(*see* celery looper)  
(*see* celery stalkworm)  
(*see* corn earworm)  
(*see* cutworms)  
(*see* diamondback moth)  
(*see* European corn borer)  
(*see* fall armyworm)  
(*see* hornworms)  
(*see* loopers)  
(*see* tomato pinworm)  
(*see* imported cabbageworm)  
(*see* pea moth)  
(*see* potato stem borer)  
(*see* potato tuberworm)

(see purple-backed cabbageworm)  
 (see stalk borer)

catface  
   on greenhouse tomato, 25.25  
   on tomato, 18.23; **18.23**

*Cavariella aegopodii*, 10.13

cavity spot  
   on carrot, 6.8; **6.8**

celery looper  
   larva, **7.22a**  
   on celery, 7.22

celery mosaic  
   on parsley, 10.12

celery mosaic virus  
   (see celery mosaic)

celery stalkworm  
   larva, **7.22b**  
   on celery, 7.22

celery webworm  
   (see celery stalkworm)

celeryworms  
   (see black swallowtails)  
   (see celery looper)  
   (see celery stalkworm)

centipedes  
   as beneficial organisms, 2.3; 3.7

*Cephalosporium apii*  
   (see *Acremonium apii*)

*Cephalosporium* spp., 26.10

*Ceratobasidium cereale*  
   (see *Rhizoctonia cerealis*)

*Cercospora apii*, 7.3

*Cercospora armoraciae*, 10.10

*Cercospora beticola*, 5.3

cercospora blight  
   on celeriac, 7.3  
   on celery, 7.3; **7.3a,b**

*Cercospora carotae*, 6.9  
   conidia, **6.9d**

cercospora leaf blight  
   on carrot, 6.9; **6.9a-d**

cercospora leaf spot  
   on beet, 5.3; **5.3**

*Cercospora malkoffii*, 10.10

*Cercospora traversiana*, 10.10

*Cercosporidium punctum*  
   (see *Phorna anethi*)

*Chaetocnema denticulata*, 12.18

*Chaetocnema pulicaria*, 12.18

*Chaetomium globosum*, 26.13

*Chaetomium olivaceum*, 26.13

*Chaetomium piluliferum*  
   (see *Botryotrichum piluliferum*)

*Chaetomium* spp., 26.13; 26.16  
   fruiting bodies, **26.13**

*Chalara elegans*, 6.6; 10.10; 11.5; 15B.4; **6.6c**

chemical control  
   in commercial vegetable production, 3.8  
   in home gardens, 3.14

chemical injury  
   on vegetable crops, 2.4

*Chenopodium album*, 2.3; 5.4; 6.13; 7.21

- a virus indicator, 4.7
- Chenopodium amaranticolor*
  - a virus indicator, 4.7; 9.16; 22.20; 22.21
- Chenopodium capitatum*, 5.9
  - a virus indicator, 4.7
- Chenopodium quinoa*
  - a virus indicator, 4.7; 9.16; 22.20
- Chenopodium* spp.
  - as virus indicators, 2.3; 15B.11; 25.20; 25.21
- cherry leaf roll
  - on rhubarb, 17.9
- cherry leaf roll virus
  - (see cherry leaf roll)
- chickweed
  - a weed pest, 2.3
- chicory yellow mottle
  - on chicory, 11.18
  - on endive, 11.18
- chicory yellow mottle virus
  - (see chicory yellow mottle)
- chilling injury
  - on greenhouse cucumber, 22.25; **22.25b**
  - (see cold injury)
- chlorsulfuron, 3.13
- chlorosis
  - on celery, 7.11; **7.11**
- Choanephora cucurbitarum*, 9.4; 22.6
- choanephora rot
  - on cucurbits, 9.4
  - on greenhouse cucumber, 22.6
- Choriorhabditis* spp., 26.28
- Chromosporium fulvum*, 26.20
- chrysanthemum leafminer
  - adult, **22.35b**; **25.28a,d**
  - egg, **25.28a**
  - on greenhouse cucumber, 22.35; **22.35a,b**
  - on greenhouse pepper, 24.15
  - on greenhouse tomato, 25.28; **25.28a-e**
- Chrysosporium luteum*, 26.5
- Chrysosporium merdarium*, 26.5
- Chrysosporium* sp., **26.5**
- cinnamon brown mold
  - on mushroom, 26.20; **26.20**
- Circulifer tenellus*
  - (see *Neoaliturus tenellus*)
- Cirsium arvense*, 2.3
- Cladobotryum dendroides*, 26.3; **26.3**
- Cladosporium cucumerinum*, 9.13; 22.16
- Cladosporium fulvum*
  - (see *Fulvia fulva*)
- Clavibacter michiganensis*
  - subsp. *michiganensis*, 18.1 ; 25.1
  - subsp. *sepedonicus*, 3.11; 16.1
- Claytonia perfoliata*
  - (see *Montia perfoliata*)
- click beetles
  - adult, **12.21b**
  - (see wireworms)
- climatic effects
  - on pathogens and pests, 2.4
- climbing cutworm, 18.35
- Clostridium* spp., 16.2; 22.28

cloudy spot  
on tomato (stink bug injury), 18.40; **18.40a,b**

clover root curculio  
adult, **15A.15**  
on pea, 15A.15

clubroot  
on broccoli, **8.8b**  
on cabbage, **8.8a**  
on crucifers, 8.8; **8.8a-c**  
on rutabaga, **8.8c**

cobweb  
on mushroom, 26.3; **26.3**

*Coenosia tigrina*, 13.26

cold injury  
on asparagus, 4.8; **4.8**  
on cucumber, **9.18a,b**  
on cucurbits, 9.18; **9.18a-c**  
on greenhouse cucumber, 22.25; **22.25a**  
on melon, **9.18c**  
on pea, 15A.11  
on potato, 16.34; **16.34n**  
on tomato, 18.24  
(see chilling injury)

*Coleomegilla maculata*, 16.44

collar rot  
on tomato, **18.15b**  
(see white mold)

*Colletotrichum atramentarium*  
(see *Colletotrichum coccodes*)

*Colletotrichum circinans*, 13.10

*Colletotrichum coccodes*, 16.6; 18.6

*Colletotrichum dertnatum* f. sp. *circinans*  
(see *Colletotrichum circinans*)

*Colletotrichum gloeosporioides* f. sp.  
malvae, 3.13

*Colletotrichum lagenarium*  
(see *Colletotrichum orbiculare*)

*Colletotrichum lindemuthianum*, 15B.2

*Colletotrichum orbiculare*, 9.3; 22.3

*Colletotrichum* sp.  
sclerotia, **16.6a,b**

Colorado potato beetle  
an introduced pest, 3.11  
adult, **16.44a**  
egg, **16.44b**  
larva, **16.44c,d**  
life cycle, **16.44T1**  
on potato, 2.2; 3.1; 16.44; **16.44a-c**  
on tomato, 3.1; 18.34  
predator of, **3.7k,m**

Columbia root-knot nematode  
eggs, **3.10b**  
galls, **3.10a**  
a foreign pest, 3.10

common blight  
on bean, 15B.1 ; **15B.1b-f**

common June beetle  
life cycle, **16.49T2**  
on potato, 16.49

common rust  
on maize, 12.5; **12.5**

common scab



on potato, 16.5; **16.5a-c; 16.5T1**

common smut  
on maize, 12.6; **12.6a,b**

confetti  
on mushroom, 26.5

*Coniotherium minitans*, 3.5

*Convolvulus arvensis*, 2.3

copper deficiency  
on greenhouse cucumber, 22.26

*Coprinus comatus*, 26.12

*Coprinus niveus*, 26.12

*Coprinus* spp., **26.12**

corky ring spot  
on potato, 16.25; **16.25a,b**

corky root (infectious)  
on greenhouse tomato, 25.6; **25.6a,b**  
on lettuce, 11.2

corky root (non-infectious)  
on lettuce, 11.20

corn borer  
(see European corn borer)

corn earworm  
adult, **12.13d**  
larva, **12.13b,c**  
on greenhouse tomato, 25.30  
on maize, 12.13; **12.13a-c**  
on tomato, 18.37

corn flea beetle  
on maize, 12.18

corn leaf aphid  
on maize, 12.14; **12.14**

corn rootworms  
on maize, 12.15

corn wireworm  
on maize, 12.21

*Corynebacterium michiganense*  
(see *Clavibacter michiganensis*)

*Corynebacterium sepedonicum*  
(see *Clavibacter michiganensis* subsp. *sepedonicus*)

*Corynoptera* spp., 22.31

*Cotesia glomerata*, 8.45

*Cotesia rubecula*, 8.45

cotton aphid  
(see melon aphid)

cottony soft rot  
on crucifers, 8.14

crabgrass  
a weed pest, 2.3

cracked stem  
on celeriac, 7.12; **7.12b**  
on celery, 7.12; **7.12a**

crane flies, 8.48

crater rot  
on carrot, 6.10; **6.10**  
on crucifers, 8.13; **8.13c**

crescent-marked lily aphid  
on potato, 16.43; **16.43T1**

cress, creeping yellow  
a weed pest, 2.3

crickets  
on tomato, 18.42

*Crioceris asparagi*, 4.10

*Crioceris duodecimpunctata*, 4.10  
crop losses  
    in commercial vegetable crops, 2.1  
    in home gardens, 3.14  
crown and root rot  
    on asparagus, 4.2; **4.2a-e**  
    on greenhouse cucumber, 22.7; **22.7a,c,d**  
    on greenhouse tomato, 25.10; **25.10a-d**  
    on tomato, 18.9  
    (see fusarium crown and root rot)  
crown gall  
    on carrot, 6.3; **6.3**  
    on rhubarb, 17.1; **77.7**  
crown rot  
    on carrot, 6.11; **6.11a-c**  
    on rhubarb, 17.3  
crucifer-feeding flea beetles  
    on crucifers, 8.44; **8.44a-e**  
    on herbs and spices, 10.14  
crucifer flea beetle  
    adult, **8.44a,c; 10.14a**  
    on broccoli, **8.44a,e**  
    on cabbage, **8.44d**  
    on crucifers, 8.44; **8.44a,c-e; 10.14b**  
    on herbs and spices, 10.14  
    on rutabaga, **8.44c**  
*Crymodes devastator*, 18.35  
cucumber beetles  
    on cucurbits, 9.21; **9.21**  
    on greenhouse cucumber, 22.35  
    (see spotted cucumber beetle)  
    (see striped cucumber beetle)  
cucumber mosaic  
    on cucurbits, 9.15; **9.15**  
    on greenhouse cucumber, 22.20; **22.20a,b**  
    on greenhouse lettuce, 23.13  
    on greenhouse pepper, 24.4  
    on greenhouse tomato, 25.18; **25.18a,b**  
    on lettuce, 11.18  
    on parsley, 10.12  
    on pepper, 18.17; **18.17**  
    on rhubarb, 17.9  
    on tomato, 18.17  
    on zucchini, **9.15**  
cucumber mosaic virus, 5.10; 7.9  
    (see cucumber mosaic)  
    (see heart mosaic)  
    (see spinach blight)  
cucumber necrosis  
    on cucurbits, 9.17  
    on greenhouse cucumber, 22.21  
cucumber necrosis virus  
    (see cucumber necrosis)  
cucumber pale fruit  
    on greenhouse cucumber, 22.22; **22.22**  
cucumber pale fruit viroid  
    (see cucumber pale fruit)  
cucumber, wild, 2.2  
*Cucumis melo*  
    a virus indicator, 9.16  
*Cucumis sativus*  
    a virus indicator, 4.7

*Cucumis* spp.  
 a virus indicator, 25.21; 25.22

*Cucurbita ficifolia*  
 a resistant rootstock, 1.2; 22.9

*Cucurbita pepo*  
 a virus indicator, 4.7; 9.16; 11.18

cudweed, low  
 a weed pest, 2.3

cultural practices  
 in commercial vegetable crops, 3.3; 3.12; 3.13  
 in home gardens, 3.14

*Cuscuta* sp., 2.3; **2.3TI**

cutworms  
 pathogen of, 3.5  
 on asparagus, 4.11  
 on bean, 15B.19  
 on carrot, 6.25  
 on cucurbits, 9.22  
 on eggplant, 18.35  
 on ginseng, 20.11  
 on lettuce, 11.26  
 on maize, 12.22  
 on parsnip, 14.7  
 on pea, 15A.15  
 on pepper, 18.35  
 on potato, 16.51  
 on tomato, 2.2; 18.35  
 species commonly found in Canada, 18.35  
 (*see* black cutworm)  
 (*see* pale western cutworm)  
 (*see* redbacked cutworm)  
 (*see* variegated cutworm)

*Cydia nigricana*, 15A.15

*Cydia rusticella*  
 (*see* *Cydia nigricana*)

*Cylindrocarpon destructans*, 20.4

*Cylindrocarpon* spp., 10.10; 20.6

*Cyphomandra betacea*  
 a virus indicator, 16.29

## D

*Dacnusa gracilis*, 6.23

*Dacnusa sibirica*, 25.28

*Dactylium dendroides*  
 (*see* *Cladobotryum dendroides*)

daddy longlegs, 2.3; 3.7

dagger nematodes  
 as ectoparasitic nematodes, 2.3

damping-off  
 on bean, 15B.4  
 on cabbage, **8.9**  
 on celeriac, 7.4  
 on celery, 7.4; **7.4**  
 on crucifers, 8.9; 8.13; **8.9**  
 on eggplant, 18.7  
 on ginseng, 20.3; **20.3a,b**  
 on greenhouse cucumber, 22.7; **22.7b**  
 on greenhouse lettuce, 23.6; **23.6a,b**  
 on greenhouse pepper, 24.1; **24.1**  
 on greenhouse tomato, 25.7; **25.7**  
 on lettuce, 11.7; **11.7a**

on maize, 12.2; **12.2a,b**  
on pea, 15A.3  
on pepper, 18.7  
on tomato, 18.7  
dark-sided cutworm, 18.35  
  adult, **18.35f**  
  egg, **18.35g**  
  larva, **6.25b; 18.35d**  
  pupa, **18.35e**  
dark-winged fungus gnat  
  adult, **26.29TI**  
  larva, **26.29TI**  
  on mushroom, 26.29  
*Datura* spp.  
  a virus indicator, 25.21  
*Delia antiqua*, 13.26  
*Delia platura*, 9.22; 12.20; 15B.18; 16.51  
*Delia radicum*, 8.41  
*Deroceras reticulatum*, 8.49; 11.27  
*Diabrotica barberi*, 12.15  
*Diabrotica longicornis*, 12.15  
*Diabrotica undecimpunctata howardi*, 9.21; 12.15; 22.35  
*Diabrotica virgifera virgifera*, 12.15  
*Diadegma insulare*, 8.42  
*Diadromus subtilicornis*, 8.42  
diamondback moth  
  adult, **8.42fg**  
  egg, **8.42b**  
  larva, **8.42c,d**  
  pupa in cocoon, **8.42e**  
  on cabbage, **8.42a**  
  on cmcifers, 8.42; **8.42a-g**  
dicamba, 3.13  
  on bean, **15B.13b**  
  (see herbicide injury)  
*Dictyna* sp., 10.16  
*Didymella bryoniae*, 2.2; 22.11  
  pseudothecia, **22.11c**  
*Didymella lycopersici*, 25.8  
didymella stem canker  
  on greenhouse tomato, 25.8; **25.8**  
*Diehliomyces microsporus*, 26.7; **26.7**  
*Digitaria sanguinalis*, 2.3  
*Digitaria* spp., 2.3  
dingy cutworm, 18.35  
*Diglyphus isaea*, 25.28  
*Diplodia maydis*  
  (see *Stenocarpella maydis*)  
diplodia stalk rot  
  on maize, 12.8; **12.8a**  
*Diplodia zeae*  
  (see *Stenocarpella maydis*)  
*Diplodina lycopersici*  
  (see *Didymella lycopersici*)  
disappearing root rot  
  on ginseng, 20.4; **20.4a,b**  
*Ditylenchus destructor*, 3.10; 16.37  
*Ditylenchus dipsaci*, 2.3; 13.24; 15A.13  
*Ditylenchus* spp., 3.12; 26.27  
dodder  
  a parasitic higher plant, 2.3; **2.3TI**  
*Doratomyces microsporus*, 26.16

dormancy  
 in crucifers, 8.33  
*Doryphorophaga doryphorae*  
 (see *Myiopharus doryphorae*)

double streak  
 on greenhouse tomato, 25.19; **25.19a,b**  
 on tomato, 18.18

DOWNCAST  
 a predictive program for onion, 13.6  
 (see downy mildew)

downy mildew, **5.4b**  
 on beet, 5.4  
 on broccoli, **8.10a,b**  
 on cauliflower, **8.10d**  
 on chives, 13.6  
 on crucifers, 8.10; **8.10a-e**  
 on garlic, 13.6  
 on greenhouse cucumber, 22.8; **22.8a,b**  
 on greenhouse lettuce, 23.7  
 on hop, 10.2; **10.2**  
 on Jerusalem artichoke, 21.2; **21.2**  
 on leek, 13.6  
 on lettuce, 3.4; 11.8; **11.8a,b**  
 on onion, 13.6; **13.5b**; **13.6a-c**  
 on pea, 15A.4; **15A.4a-c**  
 on radish, **8.10e**  
 on rhubarb, 17.4  
 on rutabaga, **8.10c**  
 on shallot, 13.6  
 on spinach, 5.4; **5.4a**

drop  
 on greenhouse lettuce, 23.8  
 on lettuce, 11.9; **11.9a-e**

*Drosophila* spp., 18.42  
 adult, **18.42g**  
 larva, **18.42f**

dry bubble  
 on mushroom, 26.8; **26.8**

dry leaf spot  
 on chicory, 11.1  
 on endive, 11.1  
 on lettuce, 11.1

dry rot  
 on potato, 16.7; **16.7a,b**

dusky wireworm  
 on potato, 16.50

Dygal  
 a microbial bactericide, 3.5

## E

ear and kernel rots  
 on maize, 12.3; **12.3a-c**

early blight  
 on celeriac, 7.3  
 on celery, 7.3  
 on eggplant, 18.8, **18.8a,d**  
 on greenhouse tomato, 25.9; **25.9a,b**  
 on pepper, 18.8; **18.8e**  
 on potato, 16.8; **16.8a-c**  
 on tomato, 3.1; 18.8; **18.8b,c**

early dying

(see verticillium wilt)

earwigs  
 in home gardens, 3.14  
 (see European earwig)

earworms  
 (see corn earworm)

eastern field wireworm  
 on potato, 16.50

*Echinochloa crusgalli*, 2.3

*Echinocystis lobata*, 2.2

ectoparasitic nematodes, 2.3

edema  
 on crucifers, 8.21  
 on greenhouse tomato, 25.25; **25.25b-d**

*Empoasca fabae*, 9.22; 16.46

enation  
 on crucifers, 8.21

*Encarsia formosa*  
 adult, **3.7q**; **25.27f**  
 a beneficial insect, 3.5; 3.7  
 on greenhouse whitefly, 22.32; 25.27; **22.32d**; **25.27e**  
 on sweetpotato whitefly, **3.10g**

endoparasitic nematodes, 2.3

enlarged lenticels  
 on potato, 16.34; **16.34m**

entomopathogenic nematodes, 3.7

*Entomophthora forficulae*, 8.43

*Entomophthora muscae*, 13.26

*Entomophthora* spp.  
 as beneficial pathogens, 3.7

*Entomoscelis americana*, 8.47

*Entylomella armoraciae*, 10.10

environmental disorders  
 on vegetable crops, 2.4

epidermal detachment  
 on cabbage, **8.32c**  
 on crucifers, 8.32; **8.32c**

*Epilachna varivestis*, 15B.19

*Epitrix cucumeris*, 16.45

*Epitrix tuberis*, 16.48

*Erwinia carotovora*  
 subsp. *atroseptica*, 8.3; 16.2; 16.3; 16.17  
 subsp. *carotovora*, 2.3; 6.2; 8.3; 11.1; 13.2; 16.2; 16.3; 16.17; 18.2; 25.3; 27.2; **11.1b**

*Erwinia chrysanthemi*, 27.1; 27.2

*Erwinia rhapontici*, 17.2

*Erwinia* spp.  
 disease cycle, **16.2TI**

*Erwinia stewartii*, 12.1

*Erwinia tracheiphila*, 9.2; 22.2

*Erysimum cheiranthoides*, 2.3

*Erysiphe cichoracearum*, 10.5; 10.10; 11.12; 21.3; 22.15; 23.10  
 cleistothecia, **22.15d**

*Erysiphe cruciferarum*  
 (see *Erysiphe polygoni*)

*Erysiphe heraclei*, 10.5

*Erysiphe pisi*  
 (see *Erysiphe polygoni*)

*Erysiphe polygoni*, 8.12; 10.10; 15A.5; 17.7  
 cleistothecia, **15A.5b**

ethylene  
 in crucifers, 8.33

*Eucosma* sp., 21.6

*Eumegachile pugnata*  
as a pollinator, 1.4

*Eumerus strigatus*, 13.25

European corn borer  
adult, **12.16g**  
egg, **12.16h**  
larva, **12.16ef; 18.36a**  
an introduced pest, 3.11  
crop residue practices, 3.3  
monitoring for, 3.2  
on bean, 15B.17  
on greenhouse tomato, 25.30  
on maize, 2.2; 3.4; 12.16; **12.16a-h**  
on pepper, 2.2; 18.36; **18.36a,b**  
on potato, 16.51  
parasite of, **3.7v**

European earwig  
adult, **8.43a,b**  
egg, **8.43c**  
nymph, **8.43d**  
on basil, 10.15  
on bean, 15B.19; **15B.19a**  
on cabbage, **8.43a**  
on crucifers, 8.43; **8.43a**  
on cucurbits, 9.22; **3.14TI**  
on maize, 12.22  
on parsley, 10.15; **10.15c**  
on rhubarb, 17.13

European wireworm  
(see dusky wireworm, 16.50)

*Euxoa auxiliaris*, 18.35  
*Euxoa detersa*, 18.35  
*Euxoa messoria*, 18.35  
*Euxoa ochrogaster*, 18.35  
*Euxoa scandens*, 18.35  
*Euxoa tessellata*, 18.35  
*Evergestis pallidata*, 8.46

exclusion  
of foreign diseases and pests, 3.9; 3.10

eyespot  
on maize, 12.4; **12.4**

## F

fall army worm  
adult, **12.17e**  
larva, **12.17a-c**  
pupa, **12.17d**  
on maize, 12.17; **12.17a**

*Feltia jaculifera*, 18.35

femleaf  
on greenhouse tomato, 25.18; 25.21; **25.18a; 25.21c**  
on tomato, 18.18  
(see also cucumber mosaic)

fire  
of endive, 11.4; 23.4

*Flavobacterium* spp., 16.2

flea beetles  
on beet, 5.16  
on crucifers, 8.44; 10.14  
on eggplant, 18.42  
on hop, 10.14

- on horseradish, 10.14
- on maize, 12.18
- on mustard, 10.14
- on pepper, 18.42
- on potato, 16.45; 16.48
- on tomato, 18.42
- (see cabbage flea beetle)
- (see corn flea beetle)
- (see crucifer-feeding flea beetles)
- (see crucifer flea beetle)
- (see garden flea beetle)
- (see hop flea beetle)
- (see horseradish flea beetle)
- (see potato flea beetle)
- (see redheaded flea beetle)
- (see toothed flea beetle)
- (see tuber flea beetle)

flies

- as pollinators of vegetable crops, 1.4
- (see blow flies)
- (see crane flies)
- (see fungus gnats)
- (see house flies)
- (see hover flies)
- (see moth flies)
- (see phorid flies)
- (see shore flies)
- (see tachinid flies)
- (see vinegar flies)

flower flies

- (see hover flies)

flower-of-an-hour

- a weed pest, 2.3

fluazifop-p-butyl, 3.13

foot rot

- on pea, 15A.2; **15A.2c,d**

*Forficula auricularia*, 8.43; 9.22; 10.15; 12.22; 15B.19; 17.13

four-spotted sap beetle

- adult, 12.3c; 12.19; 18.39
- on maize, 12.19; 12.3c; 12.19

foxglove aphid

- on potato, 16.43; **16.43**; **16.43TI**

foxtail, green

- a weed pest, 2.3; **2.3c**

foxtail, yellow

- a weed pest, 2.3

*Frankliniella occidentalis*, 18.42; 22.34; 24.14; 25.29

frost blemishing

- on cabbage, **8.32d**
- on crucifers, 8.32; **8.32d**

frost-induced injury

- on crucifers, 8.32

fruit flies

- (see vinegar flies)

*Fulvia fulva*, 25.14

*Fumago vagans*, 10.8

fungi

- as insect pathogens, 3.5; 3.7; **3.7w**
- as vegetable pathogens, 2.3
- monitoring for, 3.2

fungus gnats

- adult, **22.31b**; **22.31TI**



larva, **22.31b**  
 pupa, **22.31a**  
 on greenhouse cucumber, 22.31  
 on greenhouse lettuce, 23.19  
 on mushroom, 26.29; 26.32  
 predator of, **22.31c**  
 (see dark-winged fungus gnat)

*Fusarium acuminatum*, 6.12; 9.5  
*Fusarium avenaceum*, 6.12; 16.7; 16.9  
*Fusarium conglutinans*  
 (see *Fusarium oxysporum* f. sp. *conglutinans*)

*Fusarium culmorum*, 12.3  
 fusarium crown and root rot  
 on asparagus, 4.2; **4.2a-e**  
 on greenhouse tomato, 25.10; **25.10a-d**  
 on tomato, 18.9

fusarium dry rot  
 on carrot, 6.12; **6.12**

*Fusarium equiseti*, 6.12; 9.5  
 fusarium foot rot  
 on cucurbits, 9.5; **9.5**  
 on muskmelon, **9.5**

*Fusarium graminearum*, 12.3; 12.8; **12.3a**  
 fusarium kernel rot  
 on maize, 12.3; **12.3b**

*Fusarium moniliforme*, 4.2; 12.3; 12.8  
*Fusarium moniliforme* var. *subglutinans*  
 (see *Fusarium subglutinans*)

*Fusarium oxysporum*, 6.12; 16.9; 25.10; 25.11  
 f. sp. *apii*, 7.5  
 f. sp. *asparagi*, 4.2  
 f. sp. *cepae*, 13.4; **13.4**  
 f. sp. *conglutinans*, 3.6; 8.11  
 f. sp. *cucumerinum*  
 (see *Fusarium oxysporum* f. sp. *cucurbitacearum*)  
 f. sp. *cucurbitacearum*, 9.6; 22.9  
 f. sp. *lycopersici*, 3.4; 18.10; 25.11  
 f. sp. *melonis*  
 (see *Fusarium oxysporum* f. sp. *cucurbitacearum*)  
 f. sp. *pisi*, 15A.3  
 f. sp. *radicis-lycopersici*, 3.6; 18.9; 25.10; **25.10c**  
 f. sp. *spinaciae*, 5.5; **5.5d**

*Fusarium oxysporum* var. *redolens*  
 (see *Fusarium redolens*)

*Fusarium poae*, 9.5; 12.3  
*Fusarium redolens*, 6.12; 9.5  
 fusarium root rot  
 of bean, 15B.5; **15B.5a-c**

*Fusarium sambucinum*, 10.1; 16.7  
*Fusarium solani*, 6.12; 9.5; 16.9; 24.2  
 f. sp. *cucurbitae*, 9.5  
 f. sp. *eumartii*, 16.9  
 f. sp. *phaseoli*, 15B.5  
 f. sp. *pisi*, 15A.3  
 var. *coeruleum*, 16.7

*Fusarium sporotrichioides*, 12.3  
*Fusarium* spp., 1.3; 10.10; 12.2; 12.3; 15A.3; 16.17; 20.3; 22.7; **12.3c**  
 mycelium, **16.7b**

fusarium stalk rot  
 on maize, 12.8

fusarium stem and fruit rot  
 on greenhouse pepper, 24.2; **24.2a-c**

*Fusarium subglutinans*, 12.3; 12.8  
*Fusarium sulphureum*  
(see *Fusarium sambucinum*)  
fusarium wilt  
on beet, 5.5  
on cabbage, **8.11a,b**  
on chard, 5.5  
on crucifers, 8.11; **8.11a,b**  
on cucurbits, 9.6; **9.6a,b**  
on greenhouse cucumber, 22.9; **22.9a,b**  
on greenhouse tomato, 25.11 ; **25.11a-c**  
on muskmelon, **9.6a,b**  
on potato, 16.9; **16.9**  
on spinach, 5.5; **5.5a-d**  
on tomato, 18.10  
fusarium yellows  
of celery, 7.5; **7.5a-c**  
of crucifers 8.11; **8.11a,b**  
fuscous blight  
on bean, 15B.1

## G

*Galeopsis tetrahit*, 2.3  
galinsoga, hairy  
a weed pest, 2.3  
*Galinsoga ciliata*, 2.3  
gall midges  
on mushroom, 26.30  
gangrene  
on fiddlehead, 19.1; **19.1**  
on potato, 3.10  
(see potato gangrene)  
garden flea beetle  
on crucifers, 8.44  
garlic mosaic  
on garlic, 13.14; **13.14**  
garlic mosaic virus  
(see garlic mosaic)  
genetic abnormalities  
on potato, 16.34; **16.34a**  
*Geolaelaps* sp.  
(see *Hypoaspis* sp.)  
*Geotrichum candidum*, 26.18  
ghost spot  
on greenhouse pepper (thrips injury), 24.14; **24.14c**  
on greenhouse tomato, 25.12; **25.12d**  
on pepper (thrips injury), 18.42; **24.14c**  
on tomato, 2.2; 18.11; **18.11b**  
(see gray mold)  
(see western flower thrips)  
gibberella ear rot  
on maize, 12.3; **12.3a**  
*Gibberella fujikuroi*  
(see *Fusarium moniliforme*)  
*Gibberella pulicaris*  
(see *Fusarium sambucinum*)  
*Gibberella roseum* f. sp. *cerealis*  
(see *Fusarium graminearum*)  
gibberella stalk rot  
on maize, 12.8; **12.8b,c**  
*Gibberella subglutinans*

(see *Fusarium subglutinans*)  
*Gibberella zeae*  
(see *Fusarium graminearum*)  
gill mildew  
    on mushroom, 26.10  
glassy cutworm, 18.35  
*Gliocladium* spp.  
    in biological control, 3.5  
*Glischrochilus quadrisignatus*, 12.19  
*Globodera pallida*, 2.3; 3.11; 16.36  
*Globodera rostochiensis*, 2.3; 3.11; 16.36  
*Gloeosporium* spp., 10.10  
*Glomerella lagenaria*  
(see *Colletotrichum orbiculare*)  
*Glomerella lindemuthiana*  
(see *Colletotrichum lindemuthianum*)  
*Gnaphalium uliginosum*, 2.3  
gnats  
    (see dark-winged fungus gnat)  
    (see potato scab gnat)  
*Gnorimoschema operculella*  
(see *Phthorimaea operculella*)  
golden nematode  
    an endoparasitic nematode, 2.3  
    an introduced pest, 3.11  
    on potato, 16.36; **16.36**  
*Gomphrena globosa*  
    a virus indicator, 9.16; 11.18; 22.21  
*Gomphrena* spp.  
    a virus indicator, 25.21  
*Gracilacus* spp.  
    as vegetable pests, 2.3  
granulate cutworm, 18.35  
granulosis virus  
    a beneficial pathogen, 3.7  
    on imported cabbageworm, 3.5; 8.45  
grasshoppers  
    on maize, 12.22; **12.22a**  
    on potato, 16.51  
    on tomato, 18.42  
    (see two-striped grasshopper)  
gray garden slug on crucifers, 8.49  
    on dill, **11.27c**  
    on lettuce, 11.27  
    on tomato, **18.43**  
gray leaf spot  
    on crucifers, 8.5  
gray mold  
    on asparagus, 4.1  
    on bean, 15B.3; **15B.3a,b**  
    on chicory, 11.10  
    on cucumber, **9.7**  
    on cucurbits, 9.7; **9.7**  
    on eggplant, 18.11  
    on endive, 11.10  
    on greenhouse cucumber, 22.10; **22.10a-d**  
    on greenhouse lettuce, 23.9; **23.9**  
    on greenhouse pepper, 24.3  
    on greenhouse tomato, 25.12; **25.12a-d**  
    on lemon balm, 10.10  
    on lettuce, 11.10; **11.10a-f**  
    on pepper, 18.11

- on potato, 16.10; **16.10**
- on rhubarb, 17.5
- on thyme, 10.10
- on tomato, 18.11 ; **18.11a-c**
- gray mold neck rot
  - on chives, 13.7
  - on garlic, 13.7
  - on leek, *13.7*; **13.7d**
  - on onion, 13.7; **13.7a-c**
  - on shallot, 13.7
- gray speck
  - on cabbage, **8.29**
  - on crucifers, 8.29; **8.29**
- gray wall
  - on tomato, 18.22
- greenhouse whitefly
  - adult, **3.10e**; **22.32a,c**; **25.27b**
  - egg, **25.27c**
  - nymph, **25.27d**
  - pupa, **22.32d**; **25.27e**
  - on greenhouse cucumber, 2.2; 22.32; **22.32a-d**
  - on greenhouse tomato, 2.2; 25.27; **25.27a-e,g**
  - on tomato, 18.42
  - parasite of, **3.7g**; **22.32d**; **25.27ef**
  - pathogens of, **3.7w**
- green mold
  - on mushroom, 26.4; **26.4a,b**
- green muscardine fungus, 3.7
- green peach aphid
  - adult, **16.41a,b**; **24.12b,c**
  - nymph, **24.12b**
  - on asparagus, 4.11
  - on beet, 5.17
  - on celery, 7.17
  - on crucifers, 8.39
  - on greenhouse lettuce, 23.17
  - on greenhouse pepper, 24.12; **24.12a-c**
  - on greenhouse tomato, 25.30
  - on parsley, 10.13
  - on pepper, 18.33
  - on potato, 16.41; **16.41a**
  - on spinach, 5.17
  - on tomato, 18.33
- ground beetles
  - adult, **3.7a**
  - as beneficial insects, 3.7
- ground cherry, 18.38
  - a reservoir host of PVY<sup>N</sup>, 16.27; **3.11c**
- groundsel, common
  - a weed pest, 2.3; **2.3d**
  - herbicide resistance in, 3.13
- growth cracks
  - on carrot, 6.18; **6.18**
  - on crucifers, 8.19; **8.19**
  - on greenhouse tomato, 25.25; **25.25e**
  - on potato, 16.31; **16.31**
  - on rutabaga, **8.19**
  - on tomato, 18.25; **18.25**
- growth regulator
  - on pea, **15A.11c**
  - (see herbicide injury)
- gummosis

on cucurbits, 9.13  
on greenhouse cucumber, 22.16  
gummy stem blight  
on cucumber, 2.2  
on greenhouse cucumber, 22.11; **22.11a-e**  
on melon, 2.2  
*Gymnoascus uncinatus*  
(see *Chrysosporium merdarium*)

## H

hail injury  
on onion, 13.21  
halo blight  
on bean, 15B.1; **15B.1g-i**  
hardcap  
on mushroom, 26.22  
hardgill  
on mushroom, 26.22  
harvestmen, 2.3  
(see daddy longlegs)  
haywire  
on potato, 16.23  
head rot  
on broccoli, **8.3a,b**  
on cabbage, **8.3c**  
on chicory, 11.1  
on crucifers, 8.13; **8.3a-c; 8.13e**  
on endive, 11.1  
on greenhouse lettuce, 23.1  
on lettuce, 11.1; 11.3; **11.1a**  
head smut  
on maize, 12.7; **12.7a,b**  
heart mosaic  
on celery, 7.9; **7.9a-c**  
heart rot  
on beet, 5.11  
heat canker  
on carrot, 6.19; **6.19a-c**  
*Helicobasidium brebissonii*  
(see *Rhizoctonia crocorum*)  
*Helicotylenchus* spp.  
as vegetable pests, 2.3  
*Helicoverpa zea*, 12.13; 18.37  
*Heliothis zea*  
(see *Helicoverpa zea*)  
*Helix aspersa*, 3.11; 11.27  
*Helminthosporium solani*, 16.18  
hemp nettle  
a weed pest, 2.3  
herbicide injury, 3.13  
on bean, 15B.13; **15B.13b**  
on onion, 13.15; **13.15a-c**  
on pea, 15A.11; **3.13; 15A.11a-c**  
on pepper, 18.26; **18.26c**  
on potato, 16.34; **16.34b,c**  
on tomato, 18.26; **18.26b,d**  
*Heterodera schachtii*, 2.3; 5.14; 8.37; 17.12  
*Heterodera* spp., 3.12  
*Heterorhabditis* spp., 22.31  
*Hibiscus trionum*, 2.3  
hollow core

- on mushroom, 26.26
- hollow heart
  - on potato, 16.32; **16.32**
- hollow stem
  - on broccoli, **8.20**
  - on crucifers, 8.20; **8.20**
- honeybees
  - as pollinators of vegetable crops, 1.4
- hop aphid
  - on hop, 10.13
- hop flea beetle
  - on crucifers, 8.44
  - on hop, 10.14
  - on mustard, 10.14
- hop latent virus
  - (see hop mosaic virus)
- hop mosaic
  - on hop, 10.12
- hop mosaic virus
  - (see hop mosaic)
- hop nettle head
  - on hop, 10.12
- hop stunt viroid, 22.22
- Hordeum vulgare*, 2.3
- horizontal lesions
  - (see cavity spot)
- Hormiactis alba*, 26.10
- hormiactis cap spot
  - on mushroom, 26.10
- hornets
  - as beneficial insects, 3.7
- hornworms
  - adult moth, **18.37a**
  - larva, **18.37b**
  - on tomato, 2.2; 18.37; **18.37c**
- horse nettle
  - an alternative host, 18.38; **18.38a**
  - (see pepper maggot)
- horseradish flea beetle
  - on crucifers, 8.44
  - on horseradish, 10.14
- house flies
  - as pollinators of vegetable crops, 1.4
- hover flies
  - adult, **3.7f**
  - larva, **3.7g,h**
  - as beneficial insects, 3.7
  - as pollinators of vegetable crops, 1.4
- Howardula husseyi*, 26.31
- Humicola* spp., 26.15; 26.32
- Hydraecia micacea*, 12.22; 16.47; 17.13
- Hypoaspis* sp.
  - adult, **22.31c**
  - a beneficial mite, 3.7
  - on fungus gnats, 22.31
  - on western flower thrips, 22.34
- Hypocrea ceramica*
  - (see *Trichoderma koningii*)
- Hypocrea rufa*
  - (see *Trichoderma viride*)
- Hypocrea vinosa*
  - (see *Trichoderma harzianum*)

*Hypomyces rosellus*  
(see *Cladobotryum dendroides*)  
*Hypomyces* sp.  
(see *Mycogone pernicioso*)  
(see *Sepedonicum niveum*)  
*Hypomyces trichothecioides*  
(see *Trichothecium roseum*)

## I

imazamethabenz, 3.13  
imazethapyr, 3.13  
imidazolinone, 3.13  
imported cabbageworm  
  adult, **8.45f**  
  egg, **8.45b,c**  
  larva, **8.45c,d**  
  pupa, **8.45e**  
  on broccoli, 2.2  
  on Brussels sprouts, 2.2  
  on cabbage, 2.2; 3.4; **8.45a**  
  on cauliflower, 2.2  
  on crucifers, 8.45; **8.45a-f**  
  parasite of, **3.7p**  
infectious corky root  
  on lettuce, 11.2  
ink caps  
  on mushroom, 26.12; **26.12**  
insects  
  as beneficial organisms, 3.1; 3.7  
  as pollinators of vegetable crops, 1.4  
  as vegetable pests, 2.2; 2.3  
  on vegetable crops, key to orders, 2.3  
integrated pest management  
  in commercial vegetable crops, 3.1  
  in home gardens, 3.14  
internal black spot  
  on potato, 16.34  
internal browning  
  on crucifers, 8.22  
  on tomato, 18.18; 25.21; **18.18d,e**  
internal sprouting  
  on potato, 16.34; **16.34d**  
intumescence  
  on cabbage, **8.21a-c**  
  on crucifers, 8.21; **8.21a-c**  
*Iodophanus testaceus*  
(see *Oedocephalum glomerulosum*)  
iron deficiency  
  on bean, 15B.12; **15B.12a**  
  on greenhouse cucumber, 22.26; **22.26d**  
  on pea, 15A.10  
itersonilia canker  
  on parsnip, 14.2; **14.2a-c**  
*Itersonilia pastinaceae*, 14.2  
*Itersonilia perplexans*, 14.2; **14.2b**

## J

Japanese beetle  
  an introduced pest, 3.11  
  adult, **3.11d**

jelly end rot  
on potato, 16.33  
June beetles  
on potato, 16.49; **16.49b,c**  
(see common June beetle)  
(see white grubs)

## K

*Kabatiella zaeae*, 12.4  
*Keiferia lycopersicella*, 3.10  
kochia, 2.3; **2.3e**  
*Kochia scoparia*, 2.3

## L

lacewings  
as beneficial insects, 3.7  
lacewings, green  
adult, **3.7e**  
larva, **3.7d,e**  
as beneficial insects, 3.7  
on aphids, **3.7d,e**  
lady beetles  
adult, **3.7b**  
larva, **12.16b**  
pupa, **3.7c**  
as beneficial insects, 3.7  
ladybirds, 3.7  
(see lady beetles)  
La France  
on mushroom, 26.11  
lamb's-quarters  
a weed pest, 2.3; **2.3f,q**  
*Laspeyresia nigricana*  
(see *Cydia nigricana*)  
late blight  
disease cycle, **16.11T1**  
on celeriac, 7.7  
on celery, 7.7  
on greenhouse tomato, 25.13; **25.13a,b**  
on potato, 3.1; 16.11; **16.1 la-d; 16.11T1**  
on tomato, 18.12; **18.12a-d**  
leaf and pod spot  
on pea, 15A.2  
leaf blight  
on cucurbits, 9.8  
on greenhouse cucumber, 22.12  
on onion, 13.5  
(see botrytis leaf blight)  
leaf blotch  
on fiddlehead, 19.2  
leafcutting bees  
as pollinators of vegetable crops, 1.4  
leaf flecking  
on potato, **16.34e**  
leafhoppers  
as vegetable pests, 2.2  
monitoring for, **3.2T1**  
on beet, 5.15  
on carrot, 6.22  
on celery, 7.18



- on cucurbits, 9.22
- on lettuce, 11.23
- on potato, 16.46; 16.51
  - (see aster leafhopper)
  - (see beet leafhopper)
  - (see potato leafhopper)
  - (see *Scleroracrus* spp.)
- leafminers
  - on beet, 5.17; **5.17e**
  - on chard, **5.17f**
  - on greenhouse cucumber, 22.35
  - on greenhouse pepper, 24.15
  - on greenhouse tomato, 25.28
  - on spinach, 5.17
    - (see beet leafminer)
    - (see chrysanthemum leafminer)
    - (see vegetable leafminer)
- leaf mold
  - on greenhouse tomato, 25.14; **25.14**
- leafroll
  - on eggplant, 18.26
  - on pepper, 18.26
  - on potato, 16.26; **16.26a,b**
  - on tomato, 18.26; **18.26a**
- leafrollers
  - on horsemint, 10.15
- leaf rot
  - on cucurbits, 9.9
  - on greenhouse cucumber, 22.13; **22.13**
- leaf scorch
  - on parsley, 10.3; **10.3a,b**
- leaf spot
  - on anise, 10.10
  - on borage, 10.10
  - on corn-salad, 10.10
  - on dill, 10.10
  - on fennel, 10.10
  - on fenugreek, 10.10
  - on horseradish, 10.10
  - on lavender, 10.10
  - on lemon balm, 10.10
  - on parsley, 10.4
  - on rhubarb, 17.6
- leak
  - on cucurbits, 9.11
  - on potato, 16.12; **16.12a,b**
- leatherjackets
  - on crucifers, 8.48
- Lebia* spp., 16.44
- leek yellow stripe
  - on onion, 13.14
- leek yellow stripe virus
  - (see leek yellow stripe)
- Leptinotarsa decemlineata*, 3.11; 16.44; 18.34
- Leptinotarsa juncta*, 16.44
- Leptosphaeria maculans*
  - (see *Phoma lingam*)
- lesser bulb fly, 13.25
  - (see onion bulb fly)
- lettuce aphid
  - adult, **11.24T1**, T2
  - on greenhouse lettuce, 23.17

- on lettuce, 11.24
- lettuce big vein virus
  - (see big vein, 11.16; 23.12)
- lettuce infectious yellows
  - on chicory, 11.18
  - on endive, 11.18
  - on greenhouse lettuce, 23.15
  - on lettuce, 11.18
- lettuce infectious yellows virus
  - (see lettuce infectious yellows)
- lettuce mosaic
  - on endive, 11.17
  - on greenhouse lettuce, 23.14; **23.14**
  - on lettuce, 11.17; **11.17a,b**
- lettuce mosaic virus
  - (see lettuce mosaic)
- Limax maximus*, 11.27
- Limonius agonus*, 16.50
- Lipaphis erysimi*, 8.39
- lipstick mold
  - on mushroom, 26.18; **26.18**
- Liriomyza sativae*, 22.35; 24.15; 25.28
- Liriomyza trifolii*, 22.35; 24.15; 25.28
- Listronotus oregonensis*, 6.24; 7.20; 14.6
- Lixus concavus*, 17.13
- Longidorus apulus*, 11.18
- Longidorus* spp.
  - as vegetable pests, 2.3
  - as virus vectors, 2.3; 11.18
- loopers
  - on celery, 7.22
  - on crucifers, 8.38; 8.40
  - on greenhouse lettuce, 23.18
  - on greenhouse pepper, 24.15
  - on lettuce, 11.26
  - on pea, 15A.15
  - on pepper, 18.37
  - on tomato, 18.37
  - (see alfalfa looper)
  - (see cabbage looper)
  - (see celery looper)
- Loxostege sticticalis*, 5.17
- Loxotropa tritoma*, 6.23
- Lycopersicon* spp.
  - a virus indicator, 18.18
- Lycoriella mali*, 26.29
- Lydella radialis*, 16.47
- Lygus lineolaris*, 7.21; 9.22; 11.26; 16.51; 18.42
- Lygus* spp., 22.35; 24.15
  - adult, **22.35d**

## M

- Macrocentrus* sp.
  - a beneficial insect, 3.7
  - on potato stem borer, 3.7u
- Macrosiphum euphorbiae*, 10.13; 16.42; 18.33
- Macrosteles fascifrons*
  - (see *Macrosteles quadrilineatus*)
- Macrosteles quadrilineatus*, 6.22; 7.18; 11.23; 16.51
- maggots
  - (see cabbage maggot)

(see onion bulb fly)  
(see onion maggot)  
(see pepper maggot)  
(see seedcorn maggot)

magnesium deficiency  
  on broccoli, **8.24**  
  on celery, 7.11; **7.11**  
  on crucifers, 8.24; **8.24**  
  on greenhouse cucumber, 22.26; **22.26b**  
  on greenhouse tomato, 25.24; **25.24a,b**  
  on pea, 15A.10

maize dwarf mosaic  
  on maize, 12.10; **12.10a,b**

maize dwarf mosaic virus  
  (see maize dwarf mosaic)

*Malbranchea* spp., 26.15

maleic hydrazide  
  on onion, 13.17; **13.17**  
  (see sprout inhibitor injury)

mallow, round-leaved  
  a weed pest, 2.3  
  control of, 3.13

*Malva rotundifolia*, 2.3

management  
  by exclusion and regulation, 3.9

mancozeb  
  on ginseng, 20.9b  
  (see phytotoxicity)

*Manduca* spp., 18.37

manganese deficiency  
  on bean, 15B.12; **15B.12b**  
  on celery, 7.11  
  on greenhouse cucumber, 22.26  
  on lettuce, 11.19; **11.19a**  
  on pea, 15A.10  
  on potato, 16.34; **16.34f**

manganese toxicity  
  on lettuce, 11.19; **11.19b**  
  on pea, 15A.10

mantids  
  as beneficial insects, 3.7  
  (see praying mantis)

*Mantis religiosa*, 3.7

marigolds  
  African, 3.6; 3.12  
  French, 3.6; 3.12  
  as beneficial plants, 3.6  
  in home gardens, 3.14  
  in nematode management, 3.12

*Marssonina panattoniana*  
  (see *Microdochium panattonianum*)

mat  
  on mushroom, 26.5; **26.5**

maturity  
  in crucifers, 8.33

*Megachile* spp.  
  as pollinators of vegetable crops, 1.4

*Megaselia halterata*, 26.31

*Melanotus communis*, 12.21

*Melittia cucurbitae*, 9.22

*Meloidogyne arenaria*, 2.3; 6.20; 22.30; 25.26

*Meloidogyne chitwoodi*, 3.10

*Meloidogyne hapla*, 2.3; 5.12; 6.20; 7.15; 8.34; 9.19; 11.21; 13.22; 14.4; 15A.12; 15B.14; 16.35; 17.10; 18.30; 20.10; 22.30; 24.11; 25.26; **6.20**  
*Meloidogyne incognita*, 2.3; 6.20; 22.30; 25.26  
*Meloidogyne javanica*, 2.3; 6.20; 22.30; 25.26  
*Meloidogyne* sp., 20.3  
melon aphid  
    adult, **22.33a,b**  
    nymph, **22.33b**  
    on cucurbits, 9.22  
    on greenhouse cucumber, 22.33; **22.33b**  
    on greenhouse pepper, 24.15  
    on potato, 16.43  
*Mentha arvensis*, 2.3  
*Merlinius* spp.  
    as vegetable pests, 2.3  
*Mermis nigrescens*, 8.43  
mermithid nematodes, 8.43  
*Metarhizium anisopliae*  
    a beneficial pathogen, 3.7; 6.24  
    on white grubs, 3.7; 16.49  
*Metarhizium* spp.  
    as beneficial pathogens, 3.7  
*Metaseiulus occidentalis*  
    a beneficial mite, 3.7; 25.31  
*Metaseiulus* spp., 3.7  
*Meteorus autographae*, 8.46  
metribuzin  
    on pea, **15A.11b**  
    (see herbicide injury)  
metsulfuron methyl, 3.13  
Mexican bean beetle  
    adult, **15B.19b**  
    on bean, 15B.19  
*Microctonus bicolor*  
    (see *Townesilitus bicolor*)  
*Microdochium panattonianum*, 11.4; 23.4  
*Microplitis plutellae*, 8.42  
microsporidia  
    as beneficial pathogens, 3.7  
midges  
    larva, **3.7i; 24.12d**  
    as beneficial insects, 3.7  
    on aphids, **3.7i; 24.12d**  
milkweed, common  
    a weed pest, 2.3  
millipedes  
    as vegetable pests, 2.3; **12.21T1**  
    on potato, 16.52  
mint, field  
    a weed pest, 2.3  
minute pirate bug  
    adult, **22.34i**  
    a beneficial insect, 3.7  
    on western flower thrips, 22.34; 24.14  
mites  
    as beneficial organisms, 3.1; 3.7  
    as vegetable pests, 2.2; 2.3  
    in pest management, 3.4  
    on greenhouse cucumber, 22.36  
    on greenhouse pepper, 24.16  
    on greenhouse tomato, 25.31; 25.32  
    on herbs and spices, 10.16

- on horsemint, 10.16
- on mushroom, 26.32
- on parsnip, 14.8
  - (see red pepper mites)
  - (see tomato russet mite)
  - (see two-spotted spider mite)
- molybdenum deficiency
  - on cabbage, **8.25b**
  - on cauliflower, **8.25a**
  - on crucifers, 8.25; **8.25a,b**
  - on greenhouse cucumber, 22.26; **22.26c**
- Monilia* spp., 26.32
- monitoring for diseases and pests
  - in commercial vegetable crops, 3.2; **3.2T1**; **3.2T2**
  - in home gardens, 3.14
- Montia perfoliata*
  - a virus indicator, 11.18
- Mortierella bainieri*, 26.10
- mosaic diseases
  - on potato, 16.27; **16.27a,b**
- moth flies, 22.31
  - adult, **22.31T1**
- moths
  - as pollinators of vegetable crops, 1.4
- monitoring for, 3.2
- mottled arum aphid
  - (see crescent-marked lily aphid)
- mottled heart
  - on crucifers, 8.23
- Mucor* spp., 26.15
- mulching, 3.13
  - in home gardens, 3.14
  - (see solarization)
- mummy
  - on mushroom, 26.2; **26.2**
- muscid flies
  - as pollinators of vegetable crops, 1.4
  - (see house flies)
- muskmelon yellow stunt virus
  - (see zucchini yellow mosaic virus)
- mustard, wild
  - a weed pest, 2.3
- mustard, wormseed
  - a weed pest, 2.3
- mycelial neck rot
  - on chives, 13.7
  - on garlic, 13.7
  - on leek, 13.7
  - on onion, 13.7
  - on shallot, 13.7
- Mycogone pernicioso*, 26.9
- Mycophila* spp., 26.30
- mycosphaerella blight
  - on pea, 15A.2; **15A.2e-g**
- mycoplasma-like organisms
  - as vegetable pathogens, 2.3
- Mycosphaerella citrullina*
  - (see *Didymella bryoniae*)
- Mycosphaerella melonis*
  - (see *Didymella bryoniae*)
- Mycosphaerella pinodes*, 15A.2
  - ascospores, **15A.2T1**

conidia, **15A.2T1**  
*Myiopharus doryphorae*, 16.44  
*Myriococcum* spp., 26.15  
*Mythimna unipuncta*, 12.12  
*Myzus ascalonicus*, 13.28  
*Myzus nicotianae*, 24.12  
*Myzus persicae*, 4.11; 5.17; 7.17; 8.39; 10.13; 16.41; 18.33; 23.17; 24.12

## N

nail-head, 9.3  
(see anthracnose)  
*Nasonovia ribisnigri*, 11.24; 23.17; **11.24T1T2**  
neck rot  
    on onion, 13.7  
necrotic spot  
    on cabbage, **8.30**  
    on crucifers, 8.30; **8.30**  
*Nectria haematococca*, 24.2  
    perithecia, **24.2b,c**  
    (see *Fusarium solani*)  
*Nectria radiculicola*  
    (see *Cylindrocarpon destructans*)  
needle nematodes  
    as ectoparasitic nematodes, 2.3  
nematodes  
    as beneficial organisms, 3.7  
    as cause of vegetable crop losses, 2.2  
    as ectoparasites on vegetable crops, 2.3  
    as endoparasites on vegetable crops, 2.3  
    management of, 3.12; **3.12**  
    monitoring for, 3.2; 3.12  
    (see Columbia root-knot nematode)  
    (see dagger nematodes)  
    (see ectoparasitic nematodes)  
    (see endoparasitic nematodes)  
    (see golden nematode)  
    (see needle nematodes)  
    (see northern root-knot nematode)  
    (see pale cyst nematode)  
    (see pin nematodes)  
    (see potato cyst nematodes)  
    (see potato-rot nematode)  
    (see root-knot nematodes)  
    (see root-lesion nematode)  
    (see southern root-knot nematodes)  
    (see spiral nematodes)  
    (see stubby-root nematodes)  
    (see stunt nematodes)  
    (see sugarbeet cyst nematode)  
nematode-trapping fungi  
    on mushroom, 26.21  
*Neolittoridius tenellus*, 5.15  
*Neoalectana carpocapsae*  
    (see *Steinernema carpocapsae*)  
neoplasm  
    on crucifers, 8.21  
*Neoseiulus* (see *Amblyseius*)  
nesting  
    of carrot, 6.15  
    of chicory, lettuce, 11.10  
*Nicotiana clevelandii*

- a virus indicator, 22.20
- Nicotiana glutinosa*
  - a virus indicator, 22.20; 25.20; 25.21
- Nicotiana* spp.
  - as virus indicators, 2.3; 25.20; 25.21; 25.22
- Nicotiana sylvestris*
  - a virus indicator, 25.20
- Nicotiana tabacum*
  - a virus indicator, 22.20; 24.5; 25.20
- nightshade, black
  - a weed pest, 2.3
- nightshade, eastern black
  - a weed pest, 2.3
- nightshade, hairy
  - a weed pest, 2.3
- Nitidulidae (*see* sap beetles)
- nitrogen deficiency
  - on bean, 15B.12
  - on greenhouse cucumber, 22.26; **22.26ef**
  - on pea, 15A.10
- Nomophila nearctica*, 7.22
- non-infectious corky root
  - on lettuce, 11.20
- northern bacterial blight
  - on celery, 7.1
- northern corn rootworm
  - adult, **12.15a,b**
  - on maize, 12.15; **12.15a**
- northern root-knot nematode
  - an endoparasitic nematode, 2.3
  - on bean, 15B.14
  - on carrot, 6.20; **6.20**
  - on celeriac, 7.15
  - on celery, 7.15; **7.15a,b**
  - on crucifers, 8.34
  - on cucurbits, 9.19
  - on eggplant, 18.30
  - on ginseng, 20.10
  - on greenhouse cucumber, 22.30; **22.30d**
  - on greenhouse pepper, 24.11
  - on greenhouse tomato, 25.26; **25.26**
  - on lettuce, 11.21
  - on onion, 13.22
  - on parsnip, 14.4
  - on pea, 15A.12
  - on pepper, 18.30
  - on potato, 16.35; **16.35**
  - on rhubarb, 17.10
  - on spinach, 5.12
  - on tomato, 18.30; **18.30**
- Nosema locustae*
  - a beneficial pathogen, 3.7
  - on grasshoppers, 3.7
- nuclear polyhedrosis virus
  - (*see* polyhedrosis viruses)
- nutritional disorders, 2.4
  - on bean, 15B.12
  - on crucifers, 8.23-8.26
  - on eggplant, 18.27
  - on ginseng, 20.9
  - on greenhouse cucumber, 22.26
  - on greenhouse tomato, 25.24

on pea, 15 A. 10  
on pepper, 18.27  
on potato, 16.34  
on tomato, 18.27

## O

oedema

(see edema)

*Oedocephalum glomerulosum*, 26.17

olive-green mold

on mushroom, 26.13; **26.13**

*Olpidium brassicae*, 11.16; 23.12

*Olpidium* spp.

as virus vectors, 1.3; 22.21; 23.12

*Olpidium radicale*, 22.21

onion bulb fly

adult, **13.25**

on onion, 13.25

onion maggot

adult, **13.26b**

larva, **13.26a,c**

pupa, **13.26d**

on onion, 3.14; 13.26; **13.26a,e**

onion thrips

adult, **22.35c**

on asparagus, 4.11

on greenhouse cucumber, 22.35; **22.35c**

on greenhouse pepper, 24.15

on greenhouse tomato, 25.29

on onion, 13.27

onion yellow dwarf

on garlic, 13.14

on leek, 13.14

on onion, 13.14

on shallot, 13.14

onion yellow dwarf virus

(see onion yellow dwarf)

*Oospora pustulans*

(see *Polyscytalum pustulans*)

open veil

on mushroom, 26.23

*Opius sanguineus*, 18.38

orchard bees

as pollinators of vegetable crops, 1.4

*Orius insidiosus*, 22.34; 24.14

*Orius tristicolor*, 3.7; 22.34; 24.14; 25.31

*Osmia* spp.

as pollinators of vegetable crops, 1.4

*Ostrinia nubilalis*, 311; 12.16; 15B.17; 16.51; 18.36

oxyfluorfen

on onion, **13.15a**

(see herbicide injury)

ozone injury

on bean, 15B.13; **15B.13c**

on onion, 13.16; **13.16**

on potato, 16.34

## P

pale cyst nematode



- an endoparasitic nematode, 2.3
  - an introduced pest, 3.11
  - on potato, 16.36
- pale western cutworm, 18.35
- Papaipema nebris*, 16.51
- Papilio brevicauda*, 14.7
- Papilio* spp., 10.15; 14.7
- Papulaspora byssina*, 26.19
- parasitic nematodes
  - on mushroom, 26.27
- Paratrichodorus allii*, 2.3; 8.36; 12.11; 15B.16; 16.39; 18.32
- Paratrichodorus pachydermus*, 2.3; 8.36; 12.11; 15B.16; 16.39; 18.32
- Paratrichodorus* spp., 2.3; 8.36; 12.11; 15B.16; 16.39; 18.32
- Paratrioza cockerelli*, 16.51
- Paratylenchus* spp., 17.11
  - as vegetable pests, 2.3
- parsleyworms
  - (see black swallowtails)
- pathogens
  - as beneficial organisms, 3.7
- pea aphid
  - on greenhouse lettuce, 23.17
  - on pea, 15A.14; **15A.14**
- pea enation mosaic
  - on pea, 15A.9; **15A.9c**
- pea enation mosaic virus
  - (see pea enation mosaic)
- pea leaf weevil
  - adult, **15A.15**
  - on pea, 15A.15
- pea moth
  - on pea, 15A.15
- pea seed-borne mosaic
  - on pea, 15A.9; **15A.9f**
- pea seed-borne mosaic virus
  - (see pea seed-borne mosaic)
- pea streak
  - on pea, 15A.9; **15A.9d**
- pea streak virus
  - (see pea streak)
- pea stunt
  - on pea, 15A.9; **15A.9e,g**
- peat mold
  - on mushroom, 26.20
- pea weevil
  - on pea, 15A.15
- Pegomya betae*, 5.17
- Pegomya hyoscyami*, 5.17
- pelting rain injury
  - on onion, 13.21; **13.21**
- Pemphigus populitransversus*, 8.39
- Pemphigus populivenae*, 5.17
- penicillium mold
  - on mushroom, 26.14
- Penicillium crustosum*, 22.14
- Penicillium janczewskii*, 26.14
- Penicillium nigricans*
  - (see *Penicillium janczewskii*)
- Penicillium oxalicum*, 12.9; 22.14
- Penicillium* spp., 12.2; 12.9; 26.14; 26.16; **12.9c**
- penicillium stem rot
  - on greenhouse cucumber, 22.14; **22.14a,b**

*Peponapis pruinosa*, 1.4  
 pepper maggot  
     adult flies, **18.38b,d**  
     egg, **18.38ef**  
     larva, **18.38f**  
     pupa, **18.38g**  
     on pepper, 18.38; **18.38c,e**  
 pepper mild mottle  
     on greenhouse pepper, 24.5; **24.5a,b**  
 pepper mild mottle virus  
     (see pepper mild mottle)  
 pepper spot  
     on crucifers, 8.28  
 pepper weevil  
     adult, **24.13e**  
     larva, **24.13c**  
     pupa, **24.13d**  
     a foreign pest, 3.10  
     on greenhouse pepper, 24.13; **24.13a-e**  
 peppery leaf spot  
     on crucifers, 8.1 ; **8.1b**  
*Peridroma saucia*, 4.11 ; 18.35  
*Perillus bioculatus*, 16.44  
*Peristenus pallipes*, 7.21  
*Peristenus pseudopallipes*, 7.21  
*Peronospora destructor*, 10.10; 13.6  
     sporangiophores, **13.6d**  
*Peronospora eijfusa*  
     (see *Peronospora farinosa* f. sp.  
     *spinaciae*)  
*Peronospora farinosa*  
     (see *Peronospora farinosa* f. sp.  
     *spinaciae*)  
*Peronospora farinosa* f. sp. *spinaciae*, 5.4  
*Peronospora parasitica*, 8.10; 8.15  
     sporulation of, **8.10b**  
*Peronospora pisi*  
     (see *Peronospora viciae*)  
*Peronospora rumicis*, 17.4  
*Peronospora schachtii*  
     (see *Peronospora farinosa* f. sp. *betae*)  
*Peronospora viciae*, 15A.4  
     mycelium, **15A.4c**  
     sporulation, **15A.4a**  
 PESTCASTER  
     (see PREDICTOR)  
*Petunia* spp.  
     a virus indicator, 25.22  
*Peziza ostracoderma*  
     fruiting bodies, **26.20**  
     (see *Chromelosporium fulva*)  
*Phaseolus* spp.  
     a virus indicator, 15B. 10; 25.20; 25.21  
*Phaseolus vulgaris*  
     a virus indicator, 4.7; 11.18; 22.20; 25.20  
*Phoma anethi*, 10.4; 10.10  
*Phoma asparagi*  
     (see *Phomopsis asparagi*)  
*Phoma betae*, 5.6 disease cycle, **5.6TI**  
 phoma blight  
     on dill, 10.10; **10.10a**  
 phoma canker

on parsnip, 14.3; **14.3a-d**  
*Phoma complanata*, 14.3  
*Phoma exigua*, 10.10  
*Phoma exigua* var. *exigua*, 11.11; 16.13  
*Phoma exigua vai.foveata*, 3.10; 11.11; 19.1  
phoma leaf spot  
    on parsley, 10.4  
phoma leaf spot and root rot  
    disease cycle, **5.6TI**  
    on beet, 5.6  
    on chard, 5.6  
*Phoma lingam*, 8.6  
*Phoma mattheuccicola*, 19.1  
*Phoma medicaginis* var. *pinodella*, 15A.2  
    conidia, **15A.2TI**  
phoma rot  
    on chicory, 11.11  
    on potato, 16.13; **16.13**  
*Phoma terrestris*, 13.8  
*Phomopsis asparagi*, 4.3  
phomopsis blight  
    on asparagus, 4.3  
*Phomopsis cucurbitae*, 22.5  
*Phomopsis sclerotoides*, 22.4  
    sclerotia, **22.4b**  
phorid flies  
    adult, **26.3ITI**  
    larva, **26.3ITI**  
    on mushroom, 26.31  
*Phorodon humuli*, 10.13  
phosphorus deficiency  
    on bean, 15B.12  
    on greenhouse cucumber, 22.26  
    on pea, 15A.10  
*Phryxe vulgaris*, 8.45  
*Phthorimaea operculella*, 3.10  
*Phyllophaga anxia*, 16.49  
    life cycle, **16.49T2**  
*Phyllophaga* spp., 5.17; 16.49  
*Phyllosticta rhei*  
    (*see Ascochyta rhei*)  
*Phyllotreta albionica*, 8.44  
*Phyllotreta armoraciae*, 8.44; 10.14  
*Phyllotreta cruciferae*, 8.44  
*Phyllotreta robusta*, 8.44  
*Phyllotreta* spp., 10.14  
*Phyllotreta striolata*, 8.44  
*Physalis* sp., **3.11c**  
physiological collapse  
    on bean sprouts, 27.2  
*Phytophthora cactorum*, 20.5  
*Phytophthora cryptogea*, 4.4  
*Phytophthora erythroseptica*, 16.14  
*Phytophthora infestans*, 16.11; 18.12; 25.13; **18.12d**  
    disease cycle, **16.11TI**  
*Phytophthora megasperma* f. sp. *glycinea*, 4.4  
*Phytophthora megasperma* var. *sojae*  
    (*see Phytophthora megasperma* f. sp. *glycinea*)  
phytophthora mildew and root rot  
    on ginseng, 20.5; **20.5**  
*Phytophthora porri*, 6.14  
phytophthora spear rot

- on asparagus, 4.4
- Phytophthora spp.*, 1.3; 17.3; 18.7; 25.7
- Phytoseiulus persimilis*
  - adult, **22.36g**; **25.32**
  - a beneficial mite, 3.7
  - on two-spotted spider mite, 22.36; 24.16; 25.32
- phytotoxicity
  - on ginseng, 20.9; **20.9b,c**
- picloram, 3.13
  - on potato, 16.34; **16.34b**
  - (see herbicide injury)
- picnic beetles
  - (see four-spotted sap beetle)
- Pieris rapae*, 8.45
- pigmy mites, 26.32
  - (see red pepper mites)
- pigweed, redroot
  - a weed pest, 2.3; **2.3g,q**
- pigweed, prostrate
  - a weed pest, 2.3
- pillbugs
  - as vegetable pests, 2.3
- pin nematodes
  - as ectoparasitic nematodes, 2.3
  - on rhubarb, 2.3; 17.11
- pink eye
  - on potato, 16.4; **16.4a,b**
- pink mold rot
  - on cucurbits, 9.9
  - on greenhouse cucumber, 22.13; **22.13**
- pink rib
  - on lettuce, 11.20; **11.20**
- pink root
  - on chives, 13.8
  - on garlic, 13.8
  - on leek, 13.8
  - on onion, 13.8; **13.8a,b**
  - on shallot, 13.8
- pink rot
  - on celeriac, 7.6
  - on celery, 7.6; **7.6a,b**
  - on potato, 16.14; **16.14**
- pinworms
  - (see tomato pinworm)
- pith necrosis
  - on greenhouse tomato, 25.4; **25.4a-c**
- pithiness
  - on celery, 7.13
- plant bugs
  - as beneficial insects, 3.7
  - as vegetable pests, 3.7
  - on celery, 7.21
  - on cucurbits, 9.22
  - on greenhouse cucumber, 22.35; **22.35d**
  - on greenhouse pepper, 24.15
  - on lettuce, 11.26
  - on pepper, 18.40; 18.42
  - on potato, 16.51
  - on tomato, 18.40; 18.42
  - (see minute pirate bug)
  - (see squash bug)
  - (see stink bugs)

(see tarnished plant bug)  
*Plasmodiophora brassicae*, 8.8  
*Plasmopara halstedii*, 21.2  
plaster molds  
    on mushroom, 26.19  
*Pleospora allii*  
    (see *Stemphylium vesicarium*)  
*Pleospora betae*  
    (see *Phoma betae*)  
*Pleospora bjoerlingii*  
    (see *Phoma betae*)  
*Pleospora herbarum*  
    (see *Stemphylium botryosum*)  
*Plutella xylostella*, 8.42  
*Pnyxia scabiei*, 16.51  
pocket rot  
    on potato, 16.13  
pollination  
    by insects, 1.4  
*Polygonum convolvulus*, 2.3  
*Polygonum* spp., 2.3; 6.13  
polyhedrosis viruses  
    as beneficial pathogens, 3.7  
    on cabbage looper, 8.40; **3.7.x**  
*Poly scytalum pustulans*, 16.19  
*Popillia japonica*, 3.11  
*Portulaca oleracea*, 2.3; 6.13  
potassium deficiency  
    on greenhouse cucumber, 22.26; **22.26g**  
    on pea, 15A.10  
potato aphid  
    adult, **16.42b,c**  
    on greenhouse tomato, 25.30  
    on hop, 10.13  
    on pepper, 18.33  
    on potato, 16.42; **16.42a**  
    on tomato, 18.33  
potato beetles  
    (see Colorado potato beetle)  
potato cyst nematodes  
    as introduced pests, 3.11  
    on potato, 16.36  
    (see golden nematode)  
    (see pale cyst nematode)  
potato deforming mosaic virus  
    a foreign pathogen, 3.10  
potato flea beetle  
    adult, **16.45c**  
    on potato, 16.45; **16.45a,b**  
potato gangrene,  
    a foreign disease, 3.10  
potato leafhopper  
    adult, **16.46b**  
    on cucurbits, 9.22  
    on potato, 16.46; **16.46a**  
potato leafroll virus  
    on potato, 16.26  
potato mop top virus  
    a foreign pathogen, 3.10  
potato necrotic ring necrosis  
    (see corky ring spot, 16.25)  
potato psyllid

- on potato, 16.51
- potato-rot nematode
  - a foreign pest, 3.10
  - on potato, 16.37; **16.37**
- potato scab gnat
  - on potato, 16.51
- potato spindle tuber
  - an introduced disease, 3.11
  - (see spindle tuber, 16.28)
- potato spindle tuber viroid, 3.11; 16.28
- potato stem borer
  - adult moths, **16.47a**
  - egg, **16.47a,b**
  - larva, **12.22b; 16.47b**
  - on maize, 12.22; **12.22b**
  - on potato, 16.47
  - on rhubarb, 17.13
  - parasite of, **3.7u**
- potato tuberworm
  - larva, **3.10c**
  - a foreign pest, 3.10
- potato virus A
  - on potato, 16.27
- potato virus M
  - on potato, 16.27
- potato virus S
  - on potato, 16.27
- potato virus T
  - a foreign pathogen, 3.10
- potato virus V
  - a foreign pathogen, 3.10
- potato virus X
  - on greenhouse tomato, 25.19
  - on pepper, 18.20
  - on potato, 16.27
  - on tomato (see double streak, 18.18)
- potato virus X, resistance breaking strain
  - a foreign pathogen, 3.10
- potato virus Y
  - on pepper, 18.20
  - on potato, 16.27
  - on tomato, 18.20
- potato virus Y, necrotic strain
  - (see potato virus Y<sup>N</sup>)
- potato virus Y<sup>N</sup>
  - an introduced pathogen, 3.11
  - on ground cherry, **3.11c**
  - on tobacco, **3.1 la,b**
- potato wart
  - an introduced disease, 3.11
  - on potato, 16.21; **16.21a-d**
- potato witches'-broom mycoplasma-like organism
  - (see witches'-broom, 16.29)
- powdery mildew
  - on basil, 10.10
  - on bumet, 10.10
  - on crucifers, 8.12; **8.12**
  - on cucurbits, 9.10; **9.10**
  - on fenugreek, 10.10
  - on greenhouse cucumber, 3.1; 3.5; 22.15; **22.15a-d; 22.15T1**
  - on greenhouse lettuce, 23.10; **23.10**
  - on hop, 10.5

- on Jerusalem artichoke, 21.3; **21.3**
- on lettuce, 11.12
- on mint, 10.5
- on parsley, 10.5; **10.5**
- on pea, 15A.5; **15A.5a,b**
- on rhubarb, 17.7
- on rutabaga, **8.12**
- on sage, 10.5
- on squash, **9.10**
- powdery scab
  - on potato, 16.16; **16.16**
- Pratylenchus penetrans*, 2.3; 5.13; 6.21; 7.16; 8.35; 9.20; 11.22; 13.23; 15B.15; 16.38; 18.31; 20.3
- Pratylenchus* spp.
  - inhibition of, 3.6
- praying mantis, European
  - adult, **3.7j**
  - egg mass, **3.7j**
  - a beneficial insect, 3.7
- PREDICTOR
  - a predictive program for onion, 13.5
  - (see botrytis leaf blight)
- premature fruit yellowing
  - on greenhouse cucumber, 22.27
- protozoa
  - as beneficial organisms, 3.7
- prunus necrotic ringspot virus
  - (see hop nettle head, 10.12)
- Pseudaletia unipuncta*
  - (see *Wlythimna unipuncta*)
- Pseudomonas apii*
  - (see *Pseudomonas syringae* pv. *apii*)
- Pseudomonas cepacia*, 13.3
- Pseudomonas cichorii*, 11.3; 23.2; 25.5
- Pseudomonas corrugata*, 25.4
- pseudomonas diseases
  - on chicory, 11.3
  - on endive, 11.3
  - on lettuce, 11.3; **11.3a-d**
- Pseudomonas fluorescens*, 8.3; 11.3; 16.4; 23.1; **11.3b**
- Pseudomonas gladioli* pv. *alliiicola*, 13.1
- Pseudomonas lachrymans*
  - (see *Pseudomonas syringae* pv. *lachrymans*)
- Pseudomonas marginalis*
  - (see *Pseudomonas fluorescens*)
- Pseudomonas pisi*
  - (see *Pseudomonas syringae* pv. *pisii*)
- Pseudomonas solanacearum*, 18.5
- Pseudomonas* spp., 11.3; 16.2; 25.5; 26.2
- Pseudomonas syringae*, 11.3
  - pv. *apii*, 7.1
  - pv. *lachrymans*, 9.1; 22.1
  - pv. *maculicola*, 8.1
  - pv. *phaseolicola*, 15B.1
  - pv. *pisii*, 15A.1
  - pv. *syringae*, 15B.1
  - pv. *tagetis*, 21.1
  - pv. *tomato*, 18.3; 25.2
- Pseudomonas tagetis*
  - (see *Pseudomonas syringae* pv. *tagetis*)
- Pseudomonas tolaasii*, 26.1 ; 26.2
- Pseudomonas viridiflava*, 8.3; 11.3
- Pseudomonas viridilivida*

(see *Pseudomonas viridiflava*)  
*Pseudoperonospora cubensis*, 22.8  
 sporangiophores, **22.8b**  
*Pseudoperonospora humuli*, 10.2  
*Psila rosae*, 6.23; 7.19; 10.15; 14.5  
 psyllids  
 (see potato psyllid)  
*Psylliodes punctulata*, 8.44; 10.14  
*Pterostichus* spp., 16.44  
*Pteromalus puparum*, 8.45  
*Puccinia allii*, 10.10  
*Puccinia angustata*, 10.7  
*Puccinia asparagi*, 4.6  
 disease cycle, **4.6T1**  
*Puccinia dioicae*, 11.13  
 aecia, **11.13b**  
*Puccinia extensicola*  
 (see *Puccinia dioicae*)  
*Puccinia helianthi*, 21.4  
*Puccinia hieracii* f. sp. *cichoriae*, 11.13  
*Puccinia menthae*, 10.7  
*Puccinia patruelis*  
 (see *Puccinia hieracii* f. sp. *cichoriae*)  
*Puccinia phragmitis*, 17.8  
*Puccinia pimpinellae*, 10.10  
*Puccinia sorghi*, 12.5  
*Puccinia tanaceri* var. *dracunculina*, 10.10  
 puffiness  
 on greenhouse tomato, 25.25  
 on tomato, 18.28; **18.28**  
 purple-backed cabbageworm  
 adult, **8.46g**  
 egg mass, **8.46c**  
 larva, **8.46df**  
 pupa in cocoon, **8.46e**  
 on cabbage, **8.46b**  
 on crucifers, 8.46; **8.46a,b**  
 on rutabaga, **8.46a**  
 purple blotch  
 on leek, 13.9  
 on onion, 13.9; **13.5b**; **13.9a,b**  
 on shallot, 13.9  
 purple dwarf on potato, 16.23  
 purple spot  
 on asparagus, 4.5; **4.5a-c**  
 purple stem  
 on mushroom, 26.26  
 purple-top wilt  
 on potato, 16.23; **16.23**  
 purslane, common  
 a weed pest, 2.3; **2.3q**  
 pyemotid mites, 26.32  
 (see red pepper mites)  
*Pygmephorus* spp., 26.32  
*Pyrausta* sp., 10.15  
*Pyrenochaeta lycopersici*, 25.6  
*Pyrenochaeta terrestris*  
 (see *Phoma terrestris*)  
*Pythium acanthicum*, 9.11  
*Pythium anandrum*, 9.11  
*Pythium aphanidermatum*, 5.7; 9.11; 9.12; 12.8; 15B.6; 23.6  
*Pythium butleri*



(see *Pythium aphanidermatum*)

*Pythium coloratum*, 6.13

*Pythium debaryanum*, 7.4; 8.9; 9.11

pythium diseases

on bean, 15B.6; **15B.6**

pythium fruit rot

on cucurbits, 9.11

*Pythium dissotocum*, 23.6

*Pythium helicoides*, 9.11

*Pythium intermedium*, 6.8

*Pythium irregulare*, 6.8; 6.13; 9.11; 9.12; 15B.6

*Pythium mammilatum*, 9.11

*Pythium myriotylum*, 15B.6

*Pythium oligandrum*, 10.10

*Pythium paroecandrum*, 10.6; 15B.6

*Pythium periplocum*, 9.11

pythium root dieback

on carrot, 6.13; **6.13a,b**

pythium root rot

disease cycle, **5.7T1**

on beet, 5.7

on cucurbits, 9.12; **22.7a-d**

on parsley, 10.6; **10.6a,b**

*Pythium spp.*, 1.3; 5.7; 9.11; 9.12; 10.6; 11.7; 12.2; 12.8; 15A.3; 16.12; 16.17; 17.3; 18.7; 20.3; 22.7; 23.6; 24.1; 25.7; **11.7b,c**;  
**22.7a,c**

disease cycle, **5.7T1**

pythium stalk rot

on maize, 12.8; **12.8d**

*Pythium sulcatum*, 6.8; 6.13

*Pythium sylvaticum*, 6.8; 6.13

*Pythium ultimum*, 5.7; 6.8; 6.13; 7.4; 8.9; 9.11; 9.12; 15B.6; 16.12

*Pythium violae*, 6.8

## Q

quack grass

a weed pest, 2.3; 3.13; **2.3i,j,m**

## R

raan

on crucifers, 8.23

radish, wild

a weed pest, 2.3

ragweed, common a weed pest, 2.3; **2.3n**

ragworms

(see black swallowtails)

*Ramosia rileyana*, 10.15

ramularia leaf spot

on rhubarb, 17.6; **17.6a,b**

*Ramularia rhei*, 17.6

*Ramularia spp.*, 10.10

*Raphanus raphanistrum*, 2.3

redbacked cutworm, 18.35

larva, 6.25c; **11.26**

on lettuce, **11.26**

red clover vein mosaic virus

on pea (see pea stunt, 15A.9)

redheaded flea beetle

adult, **5.16**

on beet, 5.16

redheart

- on cabbage, **8.32e**
  - on crucifers, 8.32; **8.32e**
- red leaf
  - on rhubarb, 17.2; **17.2a,b**
- red leg
  - (see bottom rot)
- red pepper mites
  - on mushroom, 26.32
- red turnip beetle
  - adult, **8.47a**
  - egg, **8.47c**
  - larva, **8.47b**
  - pupa, **8.47c**
  - on crucifers, 8.47; **8.47b**
- regulation
  - of disease and pests, 3.9
- resistant cultivars
  - in commercial vegetable production, 3.4
  - in home gardens, 3.14
  - in tomato, VFN seed, 3.14
- resistant rootstocks, 1.2
- Rhabditis* spp., 26.21; 26.28
- Rhizobium* spp., 3.12
- rhizoctonia canker
  - on carrot, 6.11
  - on potato, 16.15; **16.15a-g**
- rhizoctonia damping-off and canker
  - on vegetable crops, **15B.7TI**
- Rhizoctonia carotae*, 6.10
- Rhizoctonia cerealis*, 5.8
- Rhizoctonia crocorum*, 6.16
- rhizoctonia diseases
  - on crucifers, 8.13
- rhizoctonia root rot
  - on bean, 15B.7; **15B.7**
  - on beet, 5.8; **5.8a,b**
  - on crucifers, **SA3\ 8.13d**
  - on rutabaga, **8.13c,d**
  - on spinach, 5.8
  - (see crater rot)
- Rhizoctonia solani*, 3.6; 5.8; 6.11; 7.4; 8.9; 8.13; 11.6; 15A.3; 15B.7; 16.15; 17.3; 18.7; 20.3; 22.7; 23.5; 24.1 ; 25.7; **20.3b**
- disease cycle, **15B.7TI**
- Rhizoctonia violaceae*
  - (see *Rhizoctonia crocorum*)
- Rhizomonas suberifaciens*, 11.2; 11.20
- Rhopalosiphoninus latysiphon*, 16.43
- Rhopalosiphum maidis*, 12.14
- Rhopalosiphum padi*, 12.14
- rhubarb curculio
  - on rhubarb, 17.13
- Rhynchosia minima*, 15B.10
- ring rot
  - an introduced pest, 3.11
  - on potato, 16.1
- ring spot
  - on chicory, 11.4
  - on endive, 11.4
  - on greenhouse lettuce, 23.4
  - on lettuce, 11.4
- root aphids
  - on lettuce, 11.25
- root death

on greenhouse cucumber, 22.28; **22.28**

root-knot nematodes  
monitoring for, 3.2  
on greenhouse vegetables, 22.30; 24.11; 25.26  
(see northern root-knot nematode)  
(see southern root-knot nematodes)

root-lesion nematode  
adult female, **16.38T1**  
monitoring for, 3.2  
on bean, 15B.15  
on beet, 5.13  
on carrot, 6.21  
on celeriac, 7.16  
on crucifers, 8.35  
on cucurbits, 9.20  
on eggplant, 18.31  
on lettuce, 11.22  
on onion, 13.23; **13.23**  
on pepper, 18.31  
on potato, 16.38; **16.38**  
on spinach, 5.13

root rot  
on ginseng, 20.3  
on maize, **12.9a**  
on pea, 3.1; 15A.3; **15A.3d-f**  
on tarragon, 10.10; **10.10c**  
(see aphanomyces root rotO)  
(see black root rot)  
(see brown root rot)  
(see disappearing root rot)  
(see fusarium crown and root rot)  
(see fusarium root rot)  
(see phoma leaf spot and root rot)  
(see phytophthora mildew and root rot)  
(see pythium root rot)  
(see rhizoctonia root rot)  
(see thielaviopsis root rot)  
(see violet root rot)

rootworms  
on maize, 12.15  
(see corn rootworms)  
(see northern corn rootworm)  
(see southern corn rootworm)  
(see southwestern corn rootworm)  
(see western corn rootworm)

*Rorippa sylvestris*, 2.3

rose comb  
on mushroom, 26.24

*Rotylenchus* spp.  
as vegetable pests, 2.3  
inhibition of, 3.6

rove beetles  
as beneficial insects, 3.7

rubbery brown rot  
on carrot, 6.14; **6.14**

rugose mosaic  
on potato, 16.27; **16.27b**

russeting  
on greenhouse tomato, 25.25

russet scab  
on potato, 16.5; **16.5c**

russet spot

on lettuce, 11.20; **11.20**  
 rust  
   on anise, 10.10  
   on asparagus, 4.6; **4.6a,b; 4.6T1**  
   on bean, 15B.8; **15B.8a,b**  
   on chicory, 11.13  
   on endive, 11.13  
   on fiddlehead, 19.2  
   on Jerusalem artichoke, 21.4; **21.4**  
   on lettuce, 11.13; **11.13a,b**  
   on maize, 12.5; **12.5**  
   on mint, 10.7; **10.7a,b**  
   on pea, 15A.6; **15A.6**  
   on rhubarb, 17.8  
   on savory, 10.10  
   on tarragon, 10.10  
 rusted root  
   on ginseng, 20.6; **20.6**  
 rust fly  
   (*see* carrot rust fly)  
 rusty root  
   on dill, 10.10  
   on ginseng, 20.6

## S

saggy socks  
   on mushroom, 26.26  
 sandhill cutworm, 18.35  
 sap beetles  
   on horsemint, 10.15  
   on maize, 12.19; **12.3c; 12.19**  
   on tomato, 18.39  
   (*see* four-spotted sap beetle)  
 saprophytic nematodes  
   on mushroom, 26.28  
 scab  
   on beet, 2.3; 5.1 ; **5.1**  
   on carrot, 2.3; 6.4; **6.4**  
   on crucifers, 8.4; **8.4a-c**  
   on cucumber, **9.13**  
   on cucurbits, 9.13; **9.13**  
   on greenhouse cucumber, 22.16; **22.16**  
   on parsnip, 2.3; 14.1  
   on potato, 2.3; 16.5; 16.16; **16.5a-c; 16.16**  
   on radish, 2.3; **8.4c**  
   on rutabaga, 2.3; **8.4a,b**  
 scab gnats  
   (*see* potato scab gnat)  
*Scleroracrus* spp., 16.29  
*Sclerotinia fuckeliana*  
   (*see* *Botrytis cinerea*)  
*Sclerotinia minor*, 2.2; 9.14; 11.9; 18.15; 22.18; 23.8; 25.17; **11.9a,d,e; 18.15a,b**  
   apothecia, **11.9e**  
   sclerotia, **11.9d,e**  
   sclerotia, parasite of, 3.5  
 Sclerotinia rot, **15B.9T1**  
   on carrot, 6.15; **6.15a,b**  
   on crucifers, 8.14; **8.14**  
   on cucurbits, 9.14  
   on greenhouse lettuce, 23.8  
   on lettuce, 11.9; **11.9a-f**

(see white mold)

*Sclerotinia sclerotiorum*, 2.2; 6.15; 7.6; 8.14; 9.14; 10.10; 11.9; 15A.7; 15B.9; 16.22; 18.15; 20.7; 21.5; 22.18; 23.8; 25.17; **16.22**  
disease cycle, **15B.9TI**  
mycelium and sclerotia, **6.15b; 9.14b; 11.9b; 18.15d**

*Sclerotinia* spp.  
parasite of, 3.5

sclerotinia stem rot  
on greenhouse cucumber, 22.18  
on pea, 15A.7; **15A.7**  
(see white mold)

sclerotinia white rot  
on ginseng, 20.7

sclerotinia wilt  
on Jerusalem artichoke, 21.5; **21.5a,b**

sclerotiniöse  
on chicory, 11.9  
on endive, 11.9

*Sclerotium cepivorum*, 3.6; 13.12  
sclerotia, **13.12e**  
sclerotia, parasite of, 3.5  
on garlic, **13.12c,d**  
on onion, **13.12b**

*Scopulariopsis brevicaulis*, 26.19  
*Scopulariopsis fimicola*, 26.19

secondary tubers  
on potato, 16.34; **16.34gj**

seedcorn maggot  
adult, **12.20c**  
larva, **9.22a; 12.20a,b**  
on bean, 15B.18; **15B.18**  
on cucumber, **9.22a,b**  
on cucurbits, 9.22; **9.22a,b**  
on maize, 12.20; **12.20a**  
on potato, 16.51

seed decay  
on bean, 15B.4  
on ginseng, 20.3  
on pea, 15A.3; **15A.3a**

seedling blight  
on pea, 15A.3; **15A.3b,c**

seed-piece decay  
on potato, 16.17; **16.17a-c**

*Senecio vulgaris*, 2.3

senescent black speck  
of cabbage, 18.28; **18.28b**

*Sepedonium niveum*, 26.6

sepedonium yellow mold  
on mushroom, 26.6; **26.6**

*Septoria apii*  
(see *Septoria apiicola*)

*Septoria apiicola*, 7.7 conidia, 7.7g

*Septoria apii-graveolentis*  
(see *Septoria apiicola*)

septoria blight  
on celeriac, 7.7  
on celery, 7.7; **7.7a-f**  
on greenhouse tomato, 25.15; **25.15**  
on tomato (see septoria leaf spot, 18.13)

*Septoria lactucae*, 11.14

*Septoria lavendulae*, 10.10

septoria leaf blotch  
on pea, 15A.8; **15A.8**

septoria leaf spot  
 on celery (*see* septoria blight, 7.7)  
 on greenhouse tomato, 25.15; **25.9b**  
 on lettuce, 11.14  
 on parsley, 10.4; **10.4**  
 on tomato, 18.13; **18.13a-c**

*Septoria lycopersici*, 18.13; 25.15  
*Septoria petroselini*, 10.4  
*Septoria pisi*, 15A.8  
*Septoria* spp., 10.10

sesiid borers  
 on horsemint, 10.15

*Setaria glauca*, 2.3  
*Setaria viridis*, 2.3  
*Setosphaeria turcica*, 12.1

sethoxydim, 3.13

shaggy stipe  
 on mushroom, 26.10; 26.26

shallot aphid  
 on onion, 13.28

shallot latent virus  
 on onion, 13.14

shepherd' s-purse  
 a weed pest, 2.3; **2.3h,q**

shoestring  
 of cucurbits (zucchini), 9.16  
 of greenhouse tomato, 25.18; **25.18a**  
 of tomato, 18.17

shore flies, 22.31  
 adult, **22.3IT 1**

short-tailed swallowtail, 14.7  
 (*see* black swallowtails)

silver scurf  
 on potato, 16.18; **16.18a-c**

simazine, 3.13

single streak  
 on greenhouse tomato, 25.21  
 on tomato, 18.18

*Sitona hispidulus*, 15A.15  
*Sitona lineatus*, 15A.15

six-spotted leafhopper  
 (*see* aster leafhopper)

skin spot  
 on potato, 16.19; **16.19**

slime  
 on chicory, 11.3  
 on endive, 11.3  
 on lettuce, 11.3

slime rot  
 on chicory, 11.1  
 on endive, 11.1  
 on lettuce, 11.1; **11.1b**

slippery skin  
 on onion, 13.1; **13.1**

slugs  
 as vegetable pests, 2.3  
 in home gardens, 3.14  
 on celery, 7.23  
 on crucifers, 8.49  
 on dill, 10.16; **11.27c**  
 on ginseng, 20.12  
 on greenhouse lettuce, 23.20

- on lettuce, 11.27; **11.27a,b**
- on pepper, 18.43
- on potato, 16.53
- on rhubarb, 17.14
- on tomato, 18.43; **18.43**
- (see black slug)
- (see gray garden slug)
- (see spotted garden slug)
- small narcissus fly, 13.25
- (see onion bulb fly)
- small sclerotial neck rot
- on onion, 13.7
- smartweed, annual
- a weed pest, 2.3; **2.3p**
- smudge
- on onion, 13.10; **13.10**
- smut
- on chives, 13.11
- on garlic, 13.11; **13.11c**
- on leek, 13.11
- on maize, 12.6; 12.7; **12.6a,b; 12.7a,b**
- on onion, 13.11; **13.1 la,b**
- on shallot, 13.11
- snails
- as vegetable pests, 2.3
- on lettuce, 11.27
- (see brown garden snail)
- social wasps
- as beneficial insects, 3.7
- soft mildew
- on mushroom, 26.3
- soft rot
- on onion, 13.2; **13.2a,b**
- on potato, 16.2; **16.2a,b; 16.2T1**
- Solanum nigrum*, 2.3; 16.26
- Solanum ptycanthum*, 2.3
- Solanum sarrachoides*, 2.3
- solarization
- to control nematodes, 3.12
- to control weeds, 3.13
- (see mulching)
- Sonchus arvensis*, 2.3
- sooty mold
- on greenhouse cucumber, **22.32b**
- on greenhouse pepper, 24.12; **24.12e**
- on greenhouse tomato, **25.27g**
- on hop, 10.8
- (see black sooty mold)
- Sorghum halepense*
- a virus indicator, 12.10
- sour skin
- on onion, 13.3; **13.3**
- southern bacterial wilt
- on pepper, 18.5
- on tomato, 18.5
- southern corn rootworm
- on maize, 12.15
- (see spotted cucumber beetle)
- southern root-knot nematodes
- as endoparasitic nematodes, 2.3
- on greenhouse cucumber, 22.30; **3.12; 22.30a-c**
- on greenhouse tomato, 25.26; **25.26**

southwestern corn rootworm, 12.15  
sowbugs  
    as vegetable pests, 2.3  
spear rot  
    on asparagus, 4.4  
*Spergula arvensis*, 2.3  
*Sphacelotheca reiliana*  
    (see *Sporisorium holci-sorghii*)  
*Sphaerotheca fuliginea*, 22.15  
    disease cycle, **22.15TI**  
*Sphaerotheca humuli*  
    (see *Sphaerotheca macularis*)  
*Sphaerotheca macularis*, 10.5; 10.10  
sphecid wasps  
    as beneficial insects, 3.7  
spider mites  
    as vegetable pests, 2.3  
    on parsnip, 14.8  
    on greenhouse cucumber, 22.36  
    on greenhouse pepper, 24.16  
    on greenhouse tomato, 25.32  
    (see two-spotted spider mite)  
spiders  
    as beneficial organisms, 3.7  
    in vegetable production, 2.3  
spinach blight  
    on beet, 5.10; **5.10**  
    on spinach, 5.10  
spinach leafminer  
    on beet, 5.17  
    on spinach, 5.17  
spindle tuber  
    an introduced disease, 3.11  
    on potato, 16.28; **16.28a,b**  
spiral nematodes  
    as ectoparasitic nematodes, 2.3  
split stipe  
    on mushroom, 26.8  
*Spodoptera frugiperda*, 12.17  
*Spondylocladium atrovirens*  
    (see *Helminthosporium solani*)  
*Spongospora subterranea*, 16.16  
spongy petiole  
    on celery, 7.13; **7.13**  
*Sporendonema purpurascens*, 26.18; **26.18**  
*Sporidesmium sclerotivorum*, 3.5  
*Sporisorium holci-sorghii*, 12.7  
spotted asparagus beetle  
    adult, **4.10c**  
    on asparagus, 4.10; **4.10c,d**  
spotted cucumber beetle  
    adult, **9.21**  
    on cucurbits, 9.21  
    on greenhouse cucumber, 22.35  
    on maize, 12.15  
spotted cutworm  
    (see climbing cutworm)  
spotted garden slug  
    on lettuce, 11.27; **11.27b**  
spotted lady beetles, 3.7  
    (see lady beetles)  
spotted necrosis



- on crucifers, 8.28
- spraing
  - on potato, 16.25
- sprouting
  - on potato, **16.34i**
- sprout inhibitor injury
  - on onion, 13.17; **13.17**
  - on potato, **16.34d**
- spurry, corn
  - a weed pest, 2.3
- squash bees
  - as pollinators of vegetable crops, 1.4
- squash bug
  - on cucurbits, 9.22
- squash vine borer
  - adult, **9.22c**
  - on cucurbits, 9.22
- stalk and tuber rot
  - on Jerusalem artichoke, 21.5; **21.5a,b**
- stalk borer
  - on potato, 16.51
- stalk rots
  - on maize, 12.8; **12.8a-d**
- stalkworms
  - (see celery stalkworm)
- Steinernema carpocapsae*, 6.24; 16.49
- Steinernema feltiae*
  - (see *Steinernema carpocapsae*)
- Stellaria media*, 2.3
- stem and bulb nematode
  - an endoparasitic nematode, 2.3
  - on chives, 13.24
  - on garlic, 13.24
  - on leek, 13.24
  - on onion, 13.24; **13.24**
  - on pea, 15A.13
  - on shallot, 13.24
- stem blight
  - on asparagus, 4.3
- stem borers
  - on Jerusalem artichoke, 21.6; **21.6**
- stem-end browning
  - on potato, 16.34; **16.34k**
- stem mottle
  - on potato, 16.25
- stem necrosis
  - on greenhouse tomato, 25.5; **25.5**
- stem rot
  - on anise, 10.10
  - on caraway, 10.10
  - on dill, 10.10
  - on greenhouse lettuce, 23.2
  - on sage, 10.10
- Stemphylium atrum*
  - (see *Ulocladium atrum*)
- Stemphylium botryosum*, 9.8; 13.19
- stemphylium leaf spot
  - on asparagus, 4.5
- Stemphylium radicinum*
  - (see *Alternaria radicina*)
- Stemphylium vesicarium*, 4.5
- spores, **4.5c**

*Stenocarpella maydis*, 12.2; 12.8  
Stewart's wilt  
    on maize, 12.1; **12.1a-c**  
*Stilbella thermophila*, 26.15  
stink bugs  
    adult, **3.7m**; **18.40a**  
    as beneficial insects, 3.7; **3.7k,m**  
    on tomato, 18.40; **18.40a,b**  
storage disorders  
    on crucifers, 8.27-8.33; **8.27-8.32**  
*Strauzia longipennis*, 21.6  
strawberry latent ringspot  
    on rhubarb, 17.9  
strawberry latent ringspot virus  
    (see strawberry latent ringspot)  
*Streptomyces acidiscabies*, 16.5  
*Streptomyces aureofaciens*, 16.5  
*Streptomyces scabies*, 2.3; 5.1; 6.4; 8.4; 14.1; Î 6.5  
    disease cycle, **16.5T1**  
*Streptomyces* spp., 16.5; 26.15  
stringiness  
    on celery, 7.14  
striped cucumber beetle  
    adult, **9.21**  
    on cucurbits, 9.21  
    on greenhouse cucumber, 22.35  
striped cutworm, 18.35  
striped flea beetle  
    adult, **8.44b**  
    on crucifers, 8.44  
stroma  
    on mushroom, 26.25  
stubby-root nematodes  
    as ectoparasitic nematodes, 2.3  
    on bean, 15B.16  
    on crucifers, 8.36  
    on eggplant, 18.32  
    on maize, 12.11  
    on pepper, 18.32  
    on potato, 16.39  
    on tomato, 18.32  
stunt  
    on greenhouse lettuce, 23.6  
    on lettuce, 11.7; **11.7b,c**  
    on pea, 15A.9  
stunting  
    on maize, **12.9a**  
stunt nematodes  
    as ectoparasitic nematodes, 2.3  
stylar cork  
    on tomato 18.23; **18.23**  
sudden wilting  
    on greenhouse cucumber, 22.7; 22.29  
sugarbeet cyst nematode  
    an endoparasitic nematode, 2.3  
    on beet, 5.14; **5.14a,b**  
    on crucifers, 8.37  
    on spinach, 5.14  
    on rhubarb, 17.12  
sugarbeet root aphid  
    on beet, 5.17  
sugarcane mosaic virus

on maize, 12.10; **12.10b**  
(see maize dwarf mosaic)  
sulfur deficiency  
on cauliflower, **8.26**  
on crucifers, 8.26; **8.26**  
sulphonylurea, 3.13  
sunflower maggot  
on Jerusalem artichoke, 21.6; **21.6**  
sunscald  
on bean, 15B. 13; **15B.13d**  
on eggplant, 18.29  
on ginseng, 20.9; **20.9b,d**  
on greenhouse pepper, 24.10  
on onion, 13.18  
on parsley, 10.10  
on pepper, 18.29; **18.29b**  
on tomato, 18.29; **18.29a**  
sweetpotato whitefly  
a foreign pest, 3.10; **3.10d-g**  
parasite of, **3.10g**  
symphylans  
as vegetable pests, 2.3  
*Synchytrium endobioticum*, 3.11; 16.21  
syrphid flies (see hover flies, 3.7; **3.7f-h**)  
*Systema frontalis*, 5.16

## T

2,4-D, 3.13  
on pepper, **18.26c**  
on tomato, **18.26b**  
(see herbicide injury)  
tachinid flies  
as beneficial insects, 3.7  
on European corn borer, **3.7v**  
on European earwig, 8.43  
*Tagetes erecta*, 3.6; 3.12  
*Tagetes patula*, 3.6; 3.12  
*Talaromyces* sp., 26.15  
*Taphrina struthiopteris*, 19.2  
target spot  
on greenhouse tomato, 25.9; **25.9a,b**  
on tomato, 18.8; **18.8b**  
tarnished plant bug  
adult, **7.21d,e; 18.42d**  
egg, **18.42e**  
nymph, **7.21b,d; 18.42b,c**  
on celery, 7.21; **7.21a-c**  
on cucurbits, 9.22  
on eggplant, 18.42  
on lettuce, 11.26  
on pepper, 18.42  
on potato, 16.51  
on tomato, 18.42; **18.42a,e**  
*Tetragnatha* sp., 10.16  
*Tetragonia* spp.  
a virus indicator, 25.21  
*Tetranychus urticae*, 10.16; 14.8; 22.36; 24.16; 25.32  
*Thanatephorus cucumeris*  
on potato, **16.15g**  
(see *Rhizoctonia solani*)  
*Thermoactinomyces* spp., 26.15

*Thermoascus* spp., 26.15  
*Thermomonospora* spp., 26.15  
*Thermomyces* spp., 26.15  
*Thielaviopsis basicola*  
(see *Chalara elegans*)  
thielaviopsis root rot  
    on fenugreek, 10.10; **10.10b**  
    (for other crops, see black root rot)  
thistle, Canada  
    a weed pest, 2.3; **2.3k**  
thistle, sow  
    a weed pest, 2.3  
three- to five-leaf dieback  
    on maize, 12.9; **12.9a-d**  
thrips  
    monitoring for, 3.2; **3.2TI**; **3.7t**  
    on asparagus, 4.11  
    on cabbage, 2.2; **8.21c**  
    on crucifers (thrips pustule), 8.21; **8.21c**  
    on ginseng, 20.11  
    on greenhouse cucumber, 22.34; 22.35  
    on greenhouse pepper, 24.14; 24.15  
    on greenhouse tomato, 25.29  
    on onion, 13.27  
    on pepper, 18.42  
    on tomato, 18.42  
    (see onion thrips)  
    (see western flower thrips)  
thrips pustule  
    on crucifers, 8.21 ; **8.21c**  
*Thrips tabaci*, 4.11; 13.27; 22.35; 24.15; 25.29  
tipburn  
    on Brussels sprouts, **8.22b**  
    on cabbage, **8.22a**  
    on crucifers, 8.22; **8.22a,b**  
    on greenhouse lettuce, 23.16; **23.16**  
    on lettuce, 11.19; **11.19c**  
    on onion, 13.19; **13.19**  
tip dieback  
    on onion, 13.19; **13.19**  
tobacco  
    a virus indicator, 2.3  
tobacco etch  
    on pepper, 18.20; **18.17**  
    on tomato, 18.20  
tobacco etch virus  
    (see tobacco etch)  
tobacco mosaic  
    on greenhouse pepper, 24.6  
    on greenhouse tomato, 25.20  
    on onion, 13.14  
    on tomato, 18.18  
tobacco mosaic virus  
    (see tobacco mosaic)  
tobacco rattle virus  
    on potato, 16.25  
tobacco ringspot virus, potato calico strain  
    a foreign pathogen, 3.10  
tobacco streak virus  
    on asparagus, 4.7  
tobacco veinal necrosis strain of potato virus Y  
    (see potato virus Y<sup>N</sup>)

tomato black ring  
 on onion, 13.14

tomato black ring virus  
 (*see* tomato black ring)

tomato fruitworm  
 (*see* corn earworm)

tomato mosaic  
 on eggplant, 18.18  
 on greenhouse pepper, 24.7  
 on greenhouse tomato, 25.21; **25.21a-e**  
 on pepper, 18.18  
 on tomato, 18.18; **18.18a-e**

tomato mosaic virus, 3.3  
 on greenhouse tomato, 25.19; 25.21  
 (*see* double streak)  
 (*see* single streak)  
 (*see* tomato mosaic)

tomato pinworm  
 a foreign pest, 3.10, **3.10h**

tomato russet mite  
 on greenhouse tomato, 25.31; **25.31**

tomato spotted wilt  
 on greenhouse lettuce, 23.15; **23.15**  
 on greenhouse pepper, 24.8; **24.8a-c**  
 on greenhouse tomato, 25.22; **25.22a-d**  
 on lemon balm, 10.12  
 on lettuce, 11.18  
 on parsley, 10.12  
 on pepper, 18.19  
 on peppermint, 10.12  
 on sage, 10.12  
 on tomato, 18.19

tomato spotted wilt virus  
 (*see* tomato spotted wilt)

TOM-CAST  
 a predictive program for tomato, 3.1; 18.6; 18.8; 18.13; 18.42

toothed flea beetle  
 on maize, 12.18

*Torula* sp., 26.15

*Townesilitus bicolor*, 8.44

translucent scale (not an insect)  
 on onion, 13.20; **13.20**

*Trialeurodes vaporariorum*, 18.42; 22.32; 25.27

*Triarthria setipennis*, 8.43

*Trichocladium basicola*  
 (*see* *Chalara elegans*)

*Trichoderma harzianum*, 26.4

*Trichoderma koningii*, 26.4

*Trichoderma* spp., 12.2; 26.4; 26.14; 26.32; **26.4a,b**  
 in biological control, 3.5

*Trichoderma viride*, 26.4

*Trichodoros* spp., 2.3; 8.36; 12.11; 15B.16; 16.39; 18.32  
 as vegetable pests, 2.3  
 as virus vectors, 2.3

*Trichogramma* spp., 12.16; 25.30

*Trichoplusia*, 7.22; 8.40; 11.26; 18.37; 23.18

*Trichothecium roseum*, 9.9; 22.13; 26.19

*Trifolium pratense*  
 a virus indicator, 15A.9

*Triticum aestivum*, 2.3

*Tropaeolum* spp.  
 a virus indicator, 25.22

true bugs  
  nymph, **3.7k**  
  adult, **3.7m**  
  as beneficial insects, 3.7  
  on Colorado potato beetle, **3.7k,m**

truffle  
  on mushroom, 26.7; **26.7**

tuber flea beetle  
  adult, **16.48c**  
  on potato, 16.48; **16.48a,b**

tuber greening  
  on potato, 16.34; **16.34h**

tuber rot  
  on Jerusalem artichoke, 21.5

tuberworms, 3.10  
  (*see* potato tuberworm)

turnip aphid  
  on crucifers, 8.39

turnip mosaic  
  on crucifers, 8.16; **8.16a-c**  
  on rhubarb, 17.9  
  on rutabaga, **8.16a-c**

turnip mosaic virus  
  (*see* turnip mosaic)

turnip root aphid  
  on crucifers, 8.39

two-spotted spider mite  
  adult, summer, **22.36c,e**; **25.32**  
  adult, winter, **22.36d-f**  
  damage indices, **22.36TI**  
  on greenhouse cucumber, 2.2; 22.36; **22.36a-d**; **22.36TI**  
  on greenhouse pepper, 24.16  
  on greenhouse tomato, 2.2; 25.32  
  on hop, 10.16  
  on parsnip, 14.8  
  predator of, **22.36g**; **25.32**

two-striped grasshopper  
  adult, **12.22a**

*Tylenchorhynchus* spp.  
  as vegetable pests, 2.3  
  inhibition of, 3.6

*Typhlodromus*  
  (*see* *Metaseiulus*)

*Tyrophagus* spp., 26.8

## U

*Ulocladium atrum*, 9.8  
*Ulocladium consortiale*, 9.8  
*Ulocladium cucurbitae*, 9.8  
ulocladium leaf spot  
  on cucumber, **9.8b**  
  on cucurbits, 9.8; **9.8b**  
  on greenhouse cucumber, 22.12  
*Ulocladium* spp., 22.12  
*Uredinopsis struthiopteridis*, 19.2  
*Urocystis cepulae*  
  (*see* *Urocystis magica*)  
*Urocystis colchici* var. *cepulae*  
  (*see* *Urocystis magica*)  
*Urocystis magica*, 13.11  
*Uromyces appendiculatus*, 15B.8

*Uromyces fabae*, 15A.6  
*Uromyces phaseoli*  
(see *Uromyces appendiculatus*)  
*Uromyces viciae-fabae*  
(see *Uromyces fabae*)  
*Ustilago reiliana*  
(see *Sporosorium holci-sorghii*)  
*Ustilago maydis*  
(see *Ustilago zaeae*)  
*Ustilago zaeae*, 12.6

## V

variegated cutworm  
larva, 18.35b,c  
on asparagus, 4.11  
on parsnip, 14.7.  
on tomato, 18.35; 18.35a,b

varnish spot  
on chicory, 11.3  
on endive, 11.3  
on lettuce, 11.3

Vedalia lady beetle, 3.7

vegetable crops in Canada  
description of, 1.2  
importance of, 1.1  
in field production, 1.2  
in home garden production, 1.2  
in protected production, 1.3  
loss of, due to diseases and pests, 1.1  
pollination of, 1.4

vegetable leafminer  
on greenhouse cucumber, 22.35  
on greenhouse pepper, 24.15  
on greenhouse tomato, 25.28

vegetable sprout rot  
of alfalfa, 27.1; **27.1a,b**  
of bean, 27.2; **27.2**; **27.2T1**

vein streaking  
on cabbage, **8.31**  
on crucifers, 8.31; **8.31**

vert de gris  
on mushroom, 26.5; **26.5**  
(see mat)

*Verticillium albo-atrum*, 10.9; 16.20; 18.14; 22.17; 25.16  
*Verticillium dahliae*, 10.9; 10.10; 16.20; 18.14; 20.8; 22.17; 25.16

verticillium disease  
on mushroom, 26.8; **26.8**

*Verticillium fungicola*, 26.8; 26.31

*Verticillium lecanii*  
a beneficial pathogen, 3.5; 3.7  
on greenhouse whitefly, 22.32; 25.27

*Verticillium malthousei*  
(see *Verticillium fungicola*)

verticillium spot  
on mushroom, 26.8

verticillium wilt  
on eggplant, 18.14; **18.14a-c**  
on ginseng, 20.8  
on greenhouse cucumber, 22.17; **22.17a,b**  
on greenhouse tomato, 25.16; **25.16a,b**  
on hop, 10.9; **10.9b**

- on mint, 10.9; **10.9a,c,d**
  - on pepper, 18.14
  - on potato, 16.20; **16.20a-c**
  - on savory, 10.10
  - on tomato, 18.14
- Vicia faba*
  - a virus indicator, 15A.9
- Vigna* spp.
  - a virus indicator, 25.21
- Vigna unguiculata*
  - a virus indicator, 22.20
- Vinca* spp.
  - a virus indicator, 25.22
- vinegar flies
  - adult, **18.42g**
  - larva, **18.42f**
  - on tomato, 18.42
- violet root rot
  - on carrot, 6.16; **6.16**
- viral diseases
  - as bio-insecticides, 3.5; 3.7
  - as vegetable pests, 2.3; 3.1
  - on mushroom, 26.11 ; **26.11**
  - on pepper, 18.17; **18.17**
  - on rhubarb, 17.9; **17.9**
  - vectors of, 1.3; 2.2
- viroids
  - as vegetable pathogens, 2.3
- viruses
  - as vegetable pathogens, 2.3
  
- W**
- wart
  - an introduced disease, 3.11
  - on potato, 16.21; **16.21a-d**
- wasps
  - as beneficial insects, 3.7; **3.7n-u**
  - as parasites, 3.7; **3.7p-u; 24.12c; 25.28e**
  - as predators, 3.7; **3.7n**
  - on aphids, 3.7s; **24.12c**
  - on chrysanthemum leafminer, **25.28e**
  - on greenhouse whitefly, **3.7q; 25.27ef**
  - on horn worm, **3.7r**
  - on imported cabbageworm, **3.7p**
  - on potato stem borer, **3.7u**
- water congestion
  - on pea, 15A.11; **15A.11d**
- water core
  - of crucifers, 8.23; **8.23c**
- watermelon mosaic
  - on cucumber, **9.17a,b**
  - on cucurbits, 9.17; **9.17a,b**
  - on greenhouse cucumber, 22.23; **22.23**
- watermelon mosaic virus
  - (see watermelon mosaic)
- watery soft rot
  - of lettuce, 11.9; 23.8; **11.9a-e**
- webworms
  - on beet, 5.17; **5.17a-c**
  - on celery, 7.22; **7.22b**
  - (see beet webworm)



(see celery stalkworm)

weepers  
on mushroom, 26.26

weeds  
as alternative hosts, 2.2  
as vegetable pests, 2.2; 2.3; **2.3a-q**  
management of, 3.13; **3.13**  
monitoring for, 3.13

weevils  
on carrot, 6.24; **6.24a-d**  
on celery, 7.20; **7.20**  
on ginseng, 20.11  
on greenhouse pepper, 24.13; **24.13a-e**  
on parsnip, 14.6  
on pea, 15A.15; **15A.15**  
on rhubarb, 17.13  
(see carrot weevil)  
(see clover root curculio)  
(see pea leaf weevil)  
(see pea weevil)  
(see pepper weevil)  
(see rhubarb curculio)

western corn rootworm  
adult, **12.15c**  
on maize, 12.15

western flower thrips  
adult, **18.42k; 22.34g; 25.29**  
egg, **18.42m; 24.14c,d**  
propupa, **18.42j; 22.34e**  
pupa, **22.34f**  
monitoring for, **3.2TI; 3.7i; 22.34d**  
on greenhouse cucumber, 22.34; **22.34a-d**  
on greenhouse pepper, 24.14; **24.14a-d**  
on greenhouse tomato, 25.29  
on pepper, 18.42; **18.42i**  
on tomato, 18.42; **18.42h**  
predator of, **22.31c; 22.34h,i**

wet bubble  
on mushroom, 26.9; **26.9**

wheat  
a weed pest, 2.3

wheat wireworm  
on potato, 16.50

*Whetzelinia sclerotiorum*  
(see *Sclerotinia sclerotiorum*)

whiptail  
on crucifers, 8.25; **8.25a**

white cutworm, 18.35

whiteflies  
as foreign pests, 3.10  
monitoring in greenhouses for, 3.2  
on tomato, 18.42  
on greenhouse cucumber, 22.32; **22.32a-d**  
on greenhouse tomato, 25.27; **25.27a-e**  
(see greenhouse whitefly)  
(see sweetpotato whitefly)

white grubs  
adult beetles, **16.49b,c**  
egg, **16.49d**  
larva, **6.26; 16.49b,d,e; 16.49T1**  
on beet, 5.17  
on carrot, 6.26

- on crucifers, 8.48
- on maize, 12.22
- on potato, 16.49; **16.49a,b**
- white mold
  - on bean, 3.5; 15B.9; **15B.9a,b; 15B.9T1**
  - on carrot, 6.15
  - on celery, 7.6
  - on chicory, 11.9; **11.9f**
  - on cucumber, **9.14b**
  - on cucurbits, 9.14; **9.14a-c**
  - on eggplant, 18.15
  - on endive, 11.9
  - on greenhouse cucumber, 22.18; **22.18a-d**
  - on greenhouse tomato, 25.17
  - on lettuce (*see drop*, 11.9, 23.8)
  - on pepper, 18.15
  - on potato, 16.22; **16.22**
  - on pumpkin, **9.14a,c**
  - on tomato, 18.15; **18.15a-e**  
(*see sclerotinia rot*)  
(*see sclerotinia stem rot*)
- white pickle, of cucumber, 22.20
- white plaster mold
  - on mushroom, 26.19  
(*see plaster molds*)
- white rot
  - on chives, 13.12
  - on garlic, 13.12; **13.12a,c,d**
  - on leek, 13.12
  - on onion, 13.12; **13.12b**
  - on shallot, 13.12
- white rust
  - on crucifers, 8.15; **8.15**
  - on horseradish, 10.10; **8.15**
  - on spinach, 5.9; **5.9**
- wild cucumber
  - an alternative host of *Didymella bryoniae*, 2.2
- wilt
  - on pea, 15A.3
  - on savory, 10.10  
(*see verticillium wilt*)
- wind injury
  - on bean, 15B.13; **15B.13e**
  - on onion, 13.21
- wirestem
  - on broccoli, **8.13b**
  - on cauliflower, **8.13a**
  - on crucifers, 8.13; **8.13a,b**
  - on pepper, 24.1
- wireworms
  - adult, **12.21b**
  - larva, **12.21a; 12.21T1; 16.50**
  - in home gardens, 3.14
  - on carrot, 6.26
  - on cucurbits, 9.22
  - on eggplant, 18.41
  - on ginseng, 20.11
  - on maize, 12.21
  - on parsnip, 14.7
  - on pepper, 18.41
  - on potato, 2.2; 16.50; **16.50**
  - on tomato, 18.41

(see corn wireworm)  
(see dusky wireworm)  
(see eastern field wireworm)  
(see wheat wireworm)

witches'-broom  
on potato, 16.29; **16.29a,b**

## **X**

*Xanthomonas campestris*

pv. *campestris*, 8.2; **8.2f**  
pv. *carotae*, 6.1  
pv. *phaseoli*, 15B.1  
pv. *vesicatoria*, 18.4  
pv. *vitians*, 11.1

*Xanthomonas phaseoli*

(see *Xanthomonas campestris* pv. *phaseoli*)

*Xanthomonas phaseoli* var. *fuscans*

(see *Xanthomonas campestris* pv. *phaseoli*)

*Xanthomonas stewartii*

(see *Erwinia stewartii*)

*Xestia adela*, 18.35

*Xiphinema diversicaudatum*, 10.12

*Xiphinema* spp.

as vegetable pests, 2.3  
as virus vectors, 2.3; 17.9

## **Y**

yellow jacket wasp

adult, **3.7n**  
a beneficial insect, 3.7

yellows

on crucifers, 8.11; **8.11a,b**

yellow streak

on garlic, 13.14; **13.14**

yellow streak virus

(see yellow streak)

## **Z**

zinc deficiency

on bean, 15B.12; **15B.12c**  
on ginseng, 20.9; **20.9a**

*Zonosemata electa*, 18.38

zucchini yellow mosaic

on cucumber, **9.16a,b**  
on cucurbits, 9.16; **9.16a-c**  
on greenhouse cucumber, 22.24; **22.24a,b**  
on squash, **9.16c**

zucchini yellow mosaic virus

(see zucchini yellow mosaic)



## About the book

An illustrated guide to identifying destructive diseases and pests affecting vegetable crops in fields, gardens, greenhouses and other environments.

An indispensable manual for commercial growers, crop advisors, market gardeners, diagnosticians, teachers, master gardeners and students.

A unique collection of more than 1000 full-color photographs of infectious diseases, environmental disorders, nematode injury, and damage from insects, mites, slugs and snails.

A valuable source of plant health management strategies for all major vegetable crops, from asparagus to zucchini; also including herbs and spices, mushrooms, vegetable sprouts, and such native crops as ginseng, Jerusalem artichoke and fiddlehead.

## About the contributors

Each chapter has been written and reviewed by researchers and extension specialists on vegetable diseases and pests at universities, colleges, and federal and provincial departments of agriculture across Canada.

## About the editors

*Ronald J. Howard, Ph.D., PAg.*, a graduate in plant pathology of the University of Saskatchewan and the University of Wisconsin, is a vegetable disease specialist at the Alberta Special Crops and Horticultural Research Center, Alberta Agriculture, Food and Rural Development, Brooks, Alberta.

*J. Allan Garland, Ph.D., PAg.*, a graduate in entomology of the University of Manitoba, University of Wisconsin, and McGill University, is a biologist with the Food Production and Inspection Branch, Agriculture and Agri-Food Canada, Ottawa, Ontario.

*W. Lloyd Seaman, Ph.D.*, a graduate in plant pathology of McGill University and the University of Wisconsin, is a research scientist at the Plant Research Centre, Agriculture and Agri-Food Canada, Ottawa, Ontario.

## About the publishers

The *Canadian Phytopathological Society* and the *Entomological Society of Canada* are the national associations of Canada's professional plant pathologists and entomologists.



The Canadian Phytopathological Society



Entomological Society of Canada

ISBN 0-9691627-2-3 (hard cover)

ISBN 0-9691627-3-1 (soft cover)