



Jurnal Ilmiah Kimia

MOLEKUL

Volume 18, Issue 1, March 2023




Indexed by Scopus

Department of Chemistry
Faculty of Mathematics and Natural Sciences
Universitas Jenderal Soedirman
Purwokerto

Molekul	Volume 18	Issue 1	pp. 1 - 169	March 2023	P-ISSN 1907-9761	E-ISSN 2503-0310
---------	--------------	------------	-------------	---------------	---------------------	---------------------

HOME / ARCHIVES / Vol 18 No 1 (2023)



PUBLISHED: 2023-03-21

ARTICLES

Adsorption of Rhodamine B on Spherical Activated Carbon synthesized from Waste Bagasse Liquid using Hydrothermal Process

Uyi Sulaeman, Baynuri Ikhyia Ulumuddin, Roy Andreas, Irmanto Irmanto, Ponco Iswanto

1-10

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5510>

 PDF

Ointment Formulation of Arumanis Mango (*Mangifera indica* L.) Leaf Extract with Chitosan Tripoliphosphate Matrix as Antibacterial

Dian Riana Ningsih, Anung Riapanitra, Zufahair Zufahair, Uyi Sulaeman, Istinganatun Khoeriyah
11-20

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5725>

 PDF

Detoxification of Cadmium on Albino Rats (*Rattus norvegicus*) with Natural Chelator of Fruiting Body Extract of *Ganoderma lucidum*

Hamedha Dhaka Kusuma Taufiq, Hernayanti Hernayanti, Nuniek Ina Ratnaningtyas

21-28

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5844>

 PDF

Molecularly Imprinted Polymer Solid-Phase Extraction (MISPE) for the Determination of Dinitro-ortho-cresol (DNOC) and Its Risk Assessment in Tomatoes

Raden Tina Rosmalina, Muhammad Bachri Amran, Bimastyaji Surya Ramadan, Sri Priatni, Novi Fitria

29-40

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5852>

 PDF

Use of Calixarenes Bearing Benzoyl Groups for the Adsorption of Heavy Metals Cations: Equilibrium and Kinetics Studies

Busroni Busroni, Dwi Siswanta, Jumina Jumina, Sri Juari Santosa, Chairil Anwar

41-49

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5911>

 PDF

Synthesis of Ionophore from p-t-Butyl-(carboxymethoxy)calix[4]arene Substituted Amide

Nasriadi Dali, Seniwati Dali, Armadi Chairunnas, Hilda Ayu Melvi Amalia, Sri Ayu Andini Puspitasari

50-58

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5927>

 PDF

Bioactivity of Endophytic Fungi Isolated from Branch of Jambu Mawar (*Syzygium jambos* (L.) Alston)

Kurratul Aini, Elfita Elfita, Hary Widjajanti, Arum Setiawan

59-69

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5931>



PDF

The Isolation, Immobilization, and Characterization of Urease from The Seeds of Winged Bean (*Psophocarpus tetragonolobus* (L.) DC.

Zusfahair Zusfahair, Dian Riana Ningsih, Amin Fatoni, Bilalodin Bilalodin, Aprilia Nafi Nuraini

70-79

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.5932>



PDF

Essential Oils of *Alpinia monopleura* and Their Antibacterial and Antioxidant Activity

Agung Wibawa Mahatva Yodha, Esti Badia, Musdalipah Musdalipah, Muhammad Azdar Setiawan, Nur Saadah Daud, Angriani Fusvita, Adryan Fristiohady, Sahidin Sahidin

80-88

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.6265>



PDF

An Insight of *Cryptocarya* Secondary Metabolites as Anticancer P388: Study of Molecular Docking and ADMET Properties

Herlina Rasyid, Riska Mardiyanti, Ihsanul Arief, Wahyu Dita Saputri

89-97

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.6364>



PDF

Effect of LiTFSI Electrolyte Salt Composition on Characteristics of PVDF-PEO-LiTFSI-Based Solid Polymer Electrolyte (SPE) for Lithium-Ion Battery

Hasim Muzadi, Nayla Zahra Kamalia, Titik Lestariningsih, Yayuk Astuti

98-106

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.6446>



PDF

Determination of The Adsorption Kinetics for Adsorption Methylene Blue Dye with C-4-Hydroxy-3-Methoxyphenylcalix[4]resorcinarene

Santi Nur Handayani, Irmanto Irmanto, Nastiti Nur Indriyani

107-116

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.6768>



PDF

Chemical Profiling and Histamine Inhibitory Activity Assessment of Merremia Vitifolia and Bidens Pilosa Extracts

Candra Yulius Tahya, Karnelasatri Karnelasatri, Nelson Gaspersz

117-130

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.6833>

 PDF

A Novel Enzyme Biosensor Based on Ag/Reduced Graphene Oxide/Chitosan Membrane with Potentiometer for Pesticide Detection

Mashuni Mashuni, Nur Arfa Yanti, Irnawati Irnawati, Fitri Handayani Hamid, Muhammad Jahiding

131-139

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.6864>

 PDF

Does Divacancy Defect Combine with N,S-Codoping Enhance the Electronic Properties of Graphene to Its Interaction with K⁺ Ion?

Yuniawan Hidayat, Fitria Rahmawati, Khoirina Dwi Nugrahaningtyas, Abduro'uf Althof Abiyyi

140-146

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.7077>

 PDF

Virtual Screening of Natural Compounds Against Six Protein Receptors Coded by The SARS-CoV-2 Genome

Fikry Awaluddin, Irmanida Batubara, Setyanto Tri Wahyudi

147-158

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.7884>

 PDF

Photodegradation and Photoelectrodegradation of Methyl Orange and Methyl Violet Dyes using Graphite/PbTiO₃ Composite under Visible Light Irradiation

Candra Purnawan, Sayekti Wahyuningsih, Ozi Adi Saputra, Erick Ryan Yulianto

159-169

DOI: <https://doi.org/10.20884/1.jm.2023.18.1.8123>

 PDF

[MAKE A SUBMISSION](#)

FOCUS AND SCOPE
GUIDE FOR AUTHORS
VIEW ARTICLES
EDITORIAL BOARD
PUBLICATION ETHICS
PEER REVIEW PROCESS
PLAGIARISM POLICY
CONTACT



Indexing and Abstracting:

Scopus[®]

INDEX COPERNICUS
INTERNATIONAL



DOAJ DIRECTORY OF
OPEN ACCESS
JOURNALS



Google
scholar



GOOGLE ANALYTICS

Google Analytics

Matomo



Molekul

Jurnal Ilmiah Kimia

Department of Chemistry, Faculty of Mathematics
and Natural Sciences,

Universitas Jenderal Soedirman, Purwokerto,
Indonesia



This work is licensed under
a [Creative Commons
Attribution 4.0
International License](https://creativecommons.org/licenses/by/4.0/).

Platform & workflow by

OJS

creating a better future

PKP | PUBLIC KNOWLEDGE PROJECT

Bioactivity of Endophytic Fungi Isolated from Branch of Jambu Mawar (*Syzygium jambos* (L.) Alston)Kurratul 'Aini^{1,2*}, Elfita^{3,*}, Hary Widjajanti⁴, Arum Setiawan⁴¹Graduate School of Sciences, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Palembang 30139, Indonesia.²Universitas Islam Negeri Raden Fatah, Palembang, Palembang 30126, South Sumatra, Indonesia³Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia⁴Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Sriwijaya, Indralaya, Ogan Ilir 30662, South Sumatra, Indonesia*Corresponding author email: elfita.elfita.69@gmail.com; **: kurratulaini_uin@radenfatah.ac.id

Received June 04, 2022; Accepted December 20, 2022; Available online March 20, 2023

ABSTRACT. Endophytic fungi isolated from medicinal plants have high diversity and the secondary metabolites produced have biological activity. Endophytic fungi isolated from medicinal plants have been the subject of many studies on their diversity and biological activity, one of which is jambu mawar (*Syzygium jambos* (L.) Alston). *S. jambos* has long been used as a traditional medicine to treat infections caused by pathogenic bacteria in many parts of the world, including South Sumatra. This study purposed to explore the diversity of endophytic fungi isolated from the branches of *S. jambos*, the antibacterial activity of endophytic fungi extracts, the determination of the structure of the compounds, and the activity of the active compounds of the selected endophytic fungi. Observations of the morphological characteristics of endophytic fungi were macroscopically and microscopically. The endophytic fungal extracts were then tested for antimicrobial activity against test bacteria *Staphylococcus aureus*, *Bacillus subtilis*, *Salmonella typhi*, and *Escherichia coli* using the Kirby-Bauer paper disk diffusion method. Isolation of pure compounds using a gravity chromatography column, the determination of the structure of the compound is determined based on ¹H-NMR, ¹³C-NMR, HMQC, and HMBC spectroscopy. Four endophytic fungi, code SJC1–4, were isolated from a branch of *S. jambos*. The results of macroscopic and microscopic morphological characterization showed three genera of *Botryosphaeria*, *Trichothecium*, and *Aspergillus*. The endophytic fungal isolate SJC1 exhibited strong activity against Gram-positive bacteria and moderate activity against Gram-negative bacteria, while SJC2–4 showed moderate activity against the four bacteria. Molecular identification of SJC1 revealed that it was *Botryosphaeria mamane*. Isolation of SJC1's pure compound yielded compound 1, which was identified as 5-acetyl-6-hydroxy-3-methyl-2H-pyran-2-one. This compound is thought to have antibacterial properties.

Keywords: Antibacterial activity, *Botryosphaeria mamane*, Endophytic fungi, *Syzygium jambos*

INTRODUCTION

Endophytic fungi are a type of fungus in plants that do not cause significant plant disease (Jia et al., 2016). These fungi and their host plants have a very complex relationship. Some endophytic fungi can produce hormones that promote plant growth, such as antiphagocytes; help the host resist biological feeding; develop medicinal ingredients; and produce many plant by products in their biological activity (Khan et al., 2013). Endophytic fungi also possess natural antimicrobial activity that can compensate for plant resource shortages, limit the length of the regeneration cycle, and feature in industrial fermentation to mass produce natural active compounds at low cost and without pollution (Yuan et al., 2017). Endophytes associated with medicinal plants have been extensively studied. These fungi

grow through natural openings in plants, such as stomata and natural wounds. Endophytes also belong to the group of dark septate endophytes and play an important role in the production of secondary metabolites and enzymes. Endophytes and the secondary metabolites they produce have great potential for medicine and pharmacy (Devi et al., 2020; Tiwari & Bae, 2022). In therapeutic applications, endophytic fungi act as antibacterial agents (Deshmukh et al., 2015; Gagana & Shivanna, 2020; Pansanit & Pripdeevech, 2018). For instance, the antibacterial activity of the endophyte *Tritirachium oryzae* isolated from *Syzygium malaccense* contains 2-(4-hydroxyphenyl)-4-methoxytetrahydrofuran-3-ol (Hapida et al., 2022); *Fusarium verticillioides*, isolated from the stem bark of *S. jambos*, contains 3-hydroxy-4-(hydroxy(4-hydroxyphenyl)methyl)dihydrofuran-2-on

(Aini et al., 2022); and *Penicillium brefeldianum* from the root bark of *Syzygium zeylanicum* contains p-hydroxybenzaldehyde (Syarifah et al., 2021).

Antibacterial resistance in turn occurs when microorganisms do not react to antibiotics that previously actively treated infections caused by bacteria. Resistance can occur through DNA mutations during replication, when bacterial species are naturally resistant to certain antibiotics, or the overuse of antibiotics (Mancuso et al., 2021). Natural sources of antibiotics are thus needed to overcome resistance to synthetic antibiotics. Numerous studies have been conducted to derive raw natural antibiotic materials from medicinal plants, including jambu mawar (*S. jambos* (L.) Alston). *S. jambos* commonly known as rose apple or jambu mawar, belongs to the family Myrtaceae and is widespread across sub-Saharan Africa, Central America, and Asia (Sharma, et al., 2013).

This plant is used for various ailments given its many active substances against pathogenic bacteria (Lim, 2012). All parts of the plant are reported to have medicinal value. In Indo-China, the whole plant is used to treat for digestive and dental ailments. Decoction of the leaves is used as a diuretic, medicine for sore eyes, and rheumatism. The seeds are used to treat diarrhea, dysentery, diabetes, and similar to nasal mucus. A decoction of the bark can also relieve asthma and bronchitis (Sharma et al., 2013). Methanol extracts of *S. jambos* leaves and stem bark have shown activity against *S. aureus* (Fathima et al., 2017; Wamba et al., 2018). However, *S. jambos* is difficult to find and requires a long period of cultivation before it can be used as a raw material. Therefore, another approach is to isolate the endophytic fungi associated with it. Studies on the isolation of endophytic fungi from specifically the stem bark (Aini et al., 2022; Roux et al., 2020), from leave and root bark (Aini et al., 2022) of *S. jambos* have been previously performed, which have antibacterial activity. Other endophytic fungi can also be found in *S. aqueum*, such as *Aspergillus*, *Trichoderma*, *Penicillium*, and others, which have antibacterial properties (Habisukan et al., 2021). Accordingly, this study explored diverse endophytic fungi isolated from *S. jambos* branches, their antibacterial and isolated pure compounds.

EXPERIMENTAL SECTION

Sampling

Samples of *S. jambos* branches were collected at the Kencana Damai Housing Complex multipurpose yard, Sako, Palembang in March 2021. The samples' morphologies were identified at the Biosystematics Laboratory, Department of Biology, Faculty of Mathematics and Natural Science, Universitas Sriwijaya, Palembang with the letter number 233/UN9.1.7/4/EP/2021.

Isolation and Identification of Endophytic Fungi

To prepare the samples, the *S. jambos* branches were rinsed 5–10 times with sterile water, with 70% ethanol for 3 minutes. They were rinsed again with sterile water before being set in a sodium hypochlorite solution for 5 minutes. Sterile specimens were cut into approximately 1–2 cm pieces and cultured in a potato dextrose agar (PDA) medium for 6 days (Manias et al., 2020; Shi et al., 2013).

The endophytic fungal isolates were then identified via macroscopic and microscopic morphology. In macroscopic morphology, each sample's surface and reverse colony color, shape, texture, and morphology were observed. Microscopic observation was carried out under a microscope (Hirox MXB-2500REZ) with glass slide cultures to confirm the type and shape of the spores, hyphae, and specific characteristics. These observations were then compared to fungus identification books from Walsh et al. (2018) and Watanabe (2010); and related articles.

Cultivation and Extraction of Endophytic Fungi

After the endophytic fungus samples' maturation, up to 5 x 300 ml of pure isolate in potato dextrose broth (PDB) was placed in 6 pure culture flasks (1x1 cm). The PDB's composition was 20 g monohydrate dextrose, 200 g potato, and 1000 mL aquadest (Gustianingtyas et al., 2020). The cultures were then incubated under static conditions at room temperature for 4 weeks (\pm 28 days). After the incubation period, the endophyte mycelium was separated from the liquid culture and dissolved in ethyl acetate at a 1:1 ratio. The ethyl acetate extract was further isolated from the liquid culture and then evaporated using a rotary evaporator until a thick extract was obtained (Nagarajan, 2019), which weigh SJC1 3.9 g, SJC2 3.7 g, SJC3 3.9 g, and SJC4 3.2 g, respectively.

Antibacterial Activity of Endophytic Fungi Extract

The endophyte isolates were generated by screening for antibacterial activity using the Kirby-Bauer paper disk diffusion method. The endophyte extract was at a concentration of 400 μ g/mL, and the antibiotic tetracycline was used at a concentration of 30 μ g/mL for comparisons. The test bacteria are represented by Gram-negative – *E. coli* (InaCCB5) and *S. typhi* (ATCC 1408), while Gram-positive – *S. aureus* (InaCCB4) and *B. subtilis* (InaCCB4). A clear area around the tray indicated antibacterial activity. The isolates' antibacterial activity characterization was determined by comparing the clear zones of the endophytic fungus extract (A) to the clear zone of the antibiotic (B) and reported as weak < 50% < moderate < 70% < strong (Aini et al., 2022).

$$\text{Antibacterial Activity (\%)} = \frac{A}{B} \times 100\%$$

Molecular Identification

Endophytic fungus isolates that exhibited strong antibacterial activity were identified at the molecular

level at the Genetika Science Indonesia Laboratory using genomic DNA extraction with a Quick-DNA Fungal Miniprep Kit (Zymo Research, D6005) and twice via polymerase chain reaction (PCR) amplification with MyTaq HS Red Mix (Bioline, BIO-25048). ITS1 and ITS4 were used as standard PCR primers (Singha et al., 2016). The samples' DNA structures were analyzed using Molecular Evolution Genetics Analysis Version 11 (Tamura et al., 2021).

Isolation of Chemical Compounds of Endophytic Fungi

The mycelium was removed from the endophytic fungal cultures after 8 weeks of culture, with the medium then filtered. The medium was extracted 3 times with ethyl acetate (1:1), then evaporated under a vacuum to obtain a concentrated extract. The concentrated extracts (1.0 g) were separated via column chromatography on silica gel 60 (70-230 mesh) in the stationary phase (1:30). The eluent had been previously determined via thin layer chromatography using silica gel 60 F254. The selected eluent, according to increasing polarity, was *n*-hexane:EtOAc at a ratio of 10:0 to 0:10 (v/v). The eluates were collected and then combined into column fractions using thin layer chromatography. Each fraction was evaporated and purified using chromatographic techniques to obtain purified compound. The chemical structures of the compounds were determined via ¹H-NMR, ¹³C-NMR, HMQC, and HMBC spectroscopy (Fadhillah et al., 2019).

RESULTS AND DISCUSSION

The isolation of the endophytic fungi from the *S. jambos* branches yielded 4 isolates encoded as SJC1–4. Based on macroscopic (Figure 1, Table 1) and microscopic (Figure 1, Table 2) characterization, they fell under three different genera, namely *Botryosphaeria*, *Trichothecium*, and *Aspergillus*. Isolate SJC3 had erect hyphae, septate, and black conidia on the tip. The inverted colonies were white, the phialides were scattered around the vesicles and twins, and the conidia were rough and dark.

SJC3 was then identified as *A. niger* (Walsh et al., 2018). SJC4's surface color was yellow-green with yellow spots, and the reverse color was brown. The hyphae had septa, and the conidia were long. Additionally, phialides uniserat and bisserat were formed in the vesicles, a trait identified in *A. flavus*. SJC2 was characterized by septate hyphae and long conidiophore, and was unbranched. The conidia were two-celled and pear-shaped, with attachment points cut into "legs." This feature indicated it was *Trichothecium roseum*. Lastly, SJC1 was identified as purposive *Botryosphaeria* with *Diplodia* (Watanabe, 2010); *B. mamane* is biosynonymous with *Cophinforma mamane* (Phillips et al., 2013) and morphologically anamorphic with *Fusicoccum mamane* (Crous et al., 2006; Mohali et al., 2007).

The antibacterial activity of the ethyl acetate extracts of SJC1–4 were tested by forming a clear area around the paper disk. The results showed that SJC1–4 all exhibited antibacterial activity (Table 3). SJC1 showed strong activity against Gram-positive bacteria and moderate activity against Gram-negative bacteria. SJC2 had moderate activity against the four test bacteria (*E. coli* 53.6%, *S. typhi* 52%, *S. aureus* 51.9%, *B. subtilis* 52%). Based on its characterization, SJC2 was identified as *Trichothecium* sp. Several *Trichothecium* sp species produce trichothecinol-A, which has antifungal, anticancer, and antimetastatic effects (Taware et al., 2014). The compounds 3 β -hydroxyartemisinic acid, 3 β ,15-dihydroxyartemisinic acid, and 3-Oxoartemisinic acid produced by the biotransformation of *T. roseum*'s artemisinic acid have antifungal properties as well (Singh et al., 2019). Additionally, it produces cuspidatol compounds belonging to the sesquiterpene group, trichomide cyclodepsopeptides which exhibit immunosuppressive activity (Zhou et al., 2018). However, *T. roseum* can cause crop damage to apples, melons, oranges, and other crops by secreting harmful mycotoxins (Zhang et al., 2020).




Figure 1. Microscopic and macroscopic characterization

Table 1. Macroscopic characteristics of endophytic fungi isolated from jambu mawar branches

Isolate	Colony color	Reverse colony color	Texture	Topography	Pattern	Radial line	Concentric circle
SJC1	Black	Black	cottony	Umbonate	Zonate	-	-
SJC2	Grey	Dark brown	cottony	Rugose	Zonate	√	√
SJC3	Grey and	White	cottony	Raised	Radiated	-	√
SJC4	Golden yellow	White	Powdery	Umbonate	Spread	-	-

Note: (-) = characteristic doesn't appear; (√) = characteristic appear

Table 2. Microscopic characteristics of endophytic fungi isolated from jambu mawar branches

Isolate	Type of spore	Shape of spore	Hyphae	Specific characteristic	Genus
SJC1	Conidia	Globose	Aerial	Pycnidia are globose, black, no observed of conidiospores	<i>Botryosphaeria</i> sp.
SJC2	Conidia	Globose	Septate	Conidiospores are long, unbranched. Conidia with two celled	<i>Trichothecium</i> sp.
SJC3	Conidia	Globose	Septate	Codiospores long. Phialides radiate around entire vesicle and are bisertae. Conidia are dark	<i>Aspergillus</i> sp.
SJC4	Conidia	Globose	Septate	Phialides are uniserate with club-shaped vesicle on surface. Conidia are oval	<i>Aspergillus</i> sp.

Table 3. Antibacterial activity of ethyl acetate extract of endophytic fungi SJC1 – SJC4

Sample	Genus/Species	Antibacterial Activity (%)			
		<i>E. coli</i>	<i>S. typhi</i>	<i>S. aureus</i>	<i>B. subtilis</i>
Tetracycline	-	21.3 ± 0.17 100.0	21.67 ± 0.56 100.0	21.77 ± 0.25 100.0	22.1 ± 0.16 100.0
SJC1	<i>Botryosphaeria</i> sp.	14.57 ± 0.21 68.9 **	15.13 ± 0.17 69.8**	15.57 ± 0.42 71.5***	15.73 ± 0.09 71.2***
SJC2	<i>Trichothecium</i> sp.	11.33 ± 0.74 53.6**	11.27 ± 0.65 52.0**	11.30 ± 0.16 51.9**	11.5 ± 0.24 52.0**
SJC3	<i>Aspergillus</i> sp.	11.97 ± 0.25 56.6**	11.17 ± 0.21 51.5**	12.10 ± 0.16 55.6**	11.23 ± 0.17 50.8**
SJC4	<i>Aspergillus</i> sp.	11.0 ± 0.22 52.1**	11.77 ± 0.12 54.3**	11.17 ± 0.21 51.3**	11.47 ± 0.31 51.9**

SJC3 and SJC4 both demonstrated moderate activity against all four bacteria. Based on macroscopic and microscopic characterization, these isolates were identified as *Aspergillus* varieties. Some of *Aspergillus* species, such as *A. flocculus* contains diorcinol and 3-Hydroxymellein compounds inhibit the parasite *Trypanosoma brucei brucei* (Tawfike et al., 2019). In turn, secondary metabolites of *A. aculeatus* are ergosterol peroxide, secalonic acid D and F, variecolin, variecolactone, and ergosterol, all of which have antimalarial (Yodsing et al., 2018). The bioactive

compounds eicosane, eicosane 2-methyl, phenol, 2,6-bis(1,1-dimethyl ethyl)-4-methyl, hexadecane 2, and 11-octadecenoic acid, the methyl ester produced by *A. fumigatus*, show antibacterial activity against *S. aureus* and *E. coli* (Octarya et al., 2021). Alkaloid compounds from the genus *Aspergillus* are generally antimicrobial, cytotoxic, and antioxidant as well (Youssef et al., 2021). Bioactive compounds 1,2,3,4,6-Penta-trimethylsilyl glucopyranose, Fmoc-L-3-(2-Naphthyl)-alanine, D-(-)-fructopyranose, pentakis (trimethylsilyl) ether, bis (2-ethylhexyl) phthalate,

trimethylsilyl ether-glucitol, octadecanamide, and N-(2-methylpropyl)-N-nitroso, as produced by *A. niger*, are antioxidants that prevent genotoxicity (Abdel-Wahhab et al., 2020).

SJC1 had strong antibacterial activity against *S. aureus* (71.5%) and *B. subtilis* (71.2%) and moderate activity against *E. coli* (68.9%) and *S. typhi* (69.8%). SJC1 was also selected for molecular identification. Its evolutionary history was derived using the Neighbor-Joining method (Saitou & Nei, 1987). The percentage of replicate trees (1000 replicates), in which related taxa are clustered together in a bootstrap test, is given next to the branches (Felsenstein, 1985). Each tree is drawn to scale, and the units of branch length are the same as the evolutionary distances used to derive the phylogenetic tree. Evolutionary distances were calculated using the number-of-difference method (Nei & Kumar, 2000) and measured in the number of base differences per sequence. The analysis included 20 nucleotide sequences. All positions with blank and missing information were removed. The final dataset contained a total of 455 locations and underwent evolutionary analysis in MEGA11 (Tamura et al., 2021). Based on PCR amplification, the result was 539bp, percentage identity > 97.5%, similar with KF531822, then SJC1 is identified as *B. mamane* (Figure 2).

It is known that *Botryosphaeria* sp. contains phenyl derivatives (Ju et al., 2016) and botryoisocoumarin-A inhibitors COX-2 (Ju et al., 2015). *B. dothidea* contains pycnophorin which has antibacterial activity against *B. subtilis* and *S. aureus*; compound stemphyperlenol, which exhibits antifungal activity against *Alternaria solani*; and the antioxidant

compounds altenusin and djalonensone (Xiao et al., 2014). Druzian et al. (2020) and Valente et al. (2018) reported that *B. dothidea* contains antioxidants. *B. mamane* also contains sesquiterpen (Oliveira et al., 2015), thiodiketopiperazines (botryosulfuranols A-C) (Barakat et al., 2019), botryomaman, 2,4-dimethoxy-6-pentylphenol, (R)-(-)-mellein, primin, *cis*-4-hydroxymellein, *trans*-4-hydroxymellein and 4,5-dihydroxy-2-hexenoic acid which show antibacterial properties against *S. aureus* (Pongcharoen et al., 2007). *B. fabicerciana* contains mellein and b-orcinaldehyde compounds, which are useful against *B. cereus*, *S. aureus*, *B. subtilis* bacteria and have antioxidants properties (Silva et al., 2021). Isocoumarin and tryptamin are antifungals produced by *B. ramose* (Hu et al., 2020; Wu et al., 2019).

Pure compound isolation (1.0 g) was performed to determine the active ingredient of the endophytic fungus isolate with the strongest antibacterial activity, which in this case was SJC1. The ¹H-NMR spectrum of compound 1 (Figure 3) showed three singlet signals, namely one vinyl proton signal at δ_H 5.92 ppm (1H, s) and two methyl proton signals at δ_H 2.26 (3H, s) and 2.66 ppm (3H, s). Also according to this spectrum, compound 1 was not an aromatic compound. The ¹³C-NMR spectrum showed another eight signals: two methyl carbon signals at δ_C 20.8 and 30.1 ppm; one methane sp² carbon signal at δ_C 101.5; and five sp² quaternary carbons. There were also three sp² quaternary carbon compounds, δ_C 169.1, 181.2, and 205.3 ppm, in a low-field chemical shift, indicating the presence of cyclic ester carbonic carbon and ketone carbonyl.




Figure 2. Phylogenetic tree of SJC1




Figure 3. The $^1\text{H-NMR}$ (A) and $^{13}\text{C-NMR}$ (B) spectra of compound 1 (^1H -500 MHz; ^{13}C -125 MHz in CDCl_3)




Figure 4. The HMQC spectra of compound 1





Figure 5. The HMBC spectra of compound 1

Table 4. The NMR data of compound 1, recorded at ^1H -500 MHz; ^{13}C -125 MHz in CDCl_3

No. C	δ_{C} ppm	Type of C	δ_{H} ppm (ΣH , multiplicity, J (Hz))	HMBC
2	169.1	C		
3	161.3	C		
4	101.5	CH	5.92 (1H, s)	169.1; 181.2; 20.8; 100.0
5	100.0	C		
6	181.2	C		
7	205.3	C		
8	30.1	CH_3	2.66 (3H, s)	205.3; 100.0
9	20.8	CH_3	2.26 (3H, s)	101.5; 169.1

**Figure 6.** The structure of compound 1 as 5-acetyl-6-hydroxy-3-methyl-2H-pyran-2-one**Table 5.** Antibacterial activity of pure compound

Sample	Concentration (ppm)	Diameter of inhibitor zone (mm)			
		<i>S. typhi</i>	<i>E. coli</i>	<i>B. subtilis</i>	<i>S. aureus</i>
Compound 1	500	7.25	7.30	7.05	7.05
	250	7.00	6.65	6.35	6.28
	125	6.68	-	-	-
	62.5	6.43	-	-	-
	31.25	6.20	-	-	-
	15.625	-	-	-	-

The HMBC spectrum of compound 1 (Figure 5, Table 4) showed a ^1H - ^{13}C three-bond correlation between two or three methyl proton bonds at δ_{H} 2.26 ppm, carbonyl ester carbon, and carbon methine sp^2 . The proton methine vinyl sp^2 5.92 ppm was correlated with three bonds to the carbonyl ester (169.1 ppm) and carbonyloxy (181.2 ppm) in the alkyl ester; methyl carbon (20.8 ppm); and quaternary carbon sp^2 through two bonds (100.0 ppm). Furthermore, there was a triple bond correlation between methyl proton (δ_{H} 2.66 ppm) and both ketone carbonyl carbon (205.3 ppm) and quaternary carbon sp^2 (100.0 ppm). Based on the spectroscopic ^1H -NMR, ^{13}C -NMR, HMQC, and HMBC data, compound 1 had a cyclic lactone or pyran ring attached to methyl, hydroxyl, and acetyl groups. The structure of the compound was thus identified as 5-acetyl-6-hydroxy-3-methyl-2H-pyran-2-one (Figure 6).

The antibacterial activity of pure compound was determined using Kirby-Bauer's disc paper diffusion method at concentrations of 500, 250, 125, 62.25,

31.25 and 15.625 ppm. The test bacteria were again *E. coli*, *S. typhi*, *B. subtilis*, and *S. aureus*. The antibacterial activity test also determined the minimum inhibitory concentration (MIC) of compound (Table 5). MIC is the lowest concentration of an antimicrobial that will inhibit the growth of microorganisms after the incubation period (Andrews, 2001). The MIC parameter value uses a breakpoint based on a modification from EUCAST (1998) with the categories of susceptible (≤ 62.5), resistant (> 62.5) ppm, and area of technical uncertainty (ATU). The MIC value of compound 1 was at a concentration of 31.25 ppm for *S. typhi* and 250 ppm for all other bacteria. This means that *S. typhi* is more susceptible and other three bacteria is more resistant to compound 1.

Based on spectroscopic ^1H -NMR, ^{13}C -NMR, HMQC, and HMBC data, compound 1 had a cyclic lactone or pyran ring attached to methyl, hydroxyl, and acetyl groups. The structure of the compound was then identified as 5-acetyl-6-hydroxy-3-methyl-2H-pyran-2-one (Figure 6), confirming it as a lactone.

Lactone compounds are widely distributed in nature as various biologically active substances and metabolic intermediates. Lactones are cyclic compounds that have intramolecular ester bonds (Kataoka et al., 2007). Compounds in the lactone group are known to exhibit antibacterial activity (Sartori et al., 2021). Accordingly, the lactone penicilactones A compound produced by *Penicillium* sp. showed antibacterial activity against *S. aureus* (Bai et al., 2019). However, lactone compounds' antibacterial activity depends on if the structure contains substituent fluorine and the length of the bacterial lipopolysaccharide (Kowalczyk et al., 2021). As such, unsaturated and hydroxy lactone compounds have antibacterial activity against *S. aureus* (Wińska et al., 2018). ϵ -lactone compounds also show antibacterial activity (Mazur & Maslowiec, 2022). From the results of this study, compound 1 can be used as a drug candidate in the future.

CONCLUSIONS

This study provides important knowledge about endophytic fungi derived from the plant *S. jambos* and their secondary metabolites as promising sources of alternative medicine. Four endophytic fungi were isolated from stem branches of *S. jambos*, namely *Botryosphaeria*, *Trichothecium*, and two *Aspergillus* species. Antibacterial tests showed that the four species exhibited antibacterial activity, with the strongest shown by *B. mamane*. *B. mamane* was found to contain 5-acetyl-6-hydroxy-3-methyl-2H-pyran-2-one, which is show antibacterial activity. This study therefore plays an important role in future examinations of herbal antibacterial medicines, with further clinical trials already underway.

ACKNOWLEDGEMENTS

The authors would like to thanks to the DRPM Kemenristek Republik Indonesia, which provided research funding through Hibah Disertasi Doktor 2022, contract no. 057/E5/PG.02.00.PT/2022 and derivative contract number: 0064.02/UN9.3.1/PL/2022.

REFERENCES

- Abdel-Wahhab, M. A., El-Nekeety, A. A., Hathout, A. S., Salman, A. S., Abdel-Aziem, S. H., Sabry, B. A., Hassan, N. S., Abdel-Aziz, M. S., Aly, S. E., & Jaswir, I. (2020). Bioactive compounds from *Aspergillus niger* extract enhance the antioxidant activity and prevent the genotoxicity in aflatoxin B1-treated rats. *Toxicon*, 181(March), 57–68. <https://doi.org/10.1016/j.toxicon.2020.04.103>
- Aini, K., Elfita, Widjajanti, H., & Setiawan, A. (2022). Diversity and antibacterial activity of endophytic fungi isolated from the medicinal plant of *Syzygium jambos*. *Biodiversitas Journal of Biological Diversity*, 23(6), 2981–2989. <https://doi.org/10.13057/biodiv/d230625>
- Aini, K., Elfita, Widjajanti, H., Setiawan, A., & Kurniawati, A. R. (2022). Antibacterial activity of endophytic fungi isolated from the stem bark of jambu mawar (*Syzygium jambos*). *Biodiversitas*, 23(1), 521–532. <https://doi.org/10.13057/biodiv/d230156>
- Andrews, J. M. (2001). Determination of minimum inhibitory concentrations. *Journal of Antimicrobial Chemotherapy*, 48(1), 5–16.
- Bai, M., Huang, G. L., Mei, R. Q., Wang, B., Luo, Y. P., Nong, X. H., Chen, G. Y., & Zheng, C. J. (2019). Bioactive lactones from the mangrove-derived fungus *Penicillium* sp. TGM112. *Marine Drugs*, 17(8), 1–7. <https://doi.org/10.3390/md17080433>
- Barakat, F., Vansteelandt, M., Triastuti, A., Jargeat, P., Jacquemin, D., Graton, J., Mejia, K., Cabanillas, B., Vendier, L., Stigliani, J. L., Haddad, M., & Fabre, N. (2019). Thiodiketopiperazines with two spirocyclic centers extracted from *Botryosphaeria mamane*, an endophytic fungus isolated from *Bixa orellana* L. *Phytochemistry*, 158, 142–148. <https://doi.org/10.1016/j.phytochem.2018.11.007>
- Crous, P. W., Slippers, B., Wingfield, M. J., Rheeder, J., Marasas, W. F. O., Philips, A. J. L., Alves, A., Burgess, T., Barber, P., & Groenewald, J. Z. (2006). Phylogenetic lineages in the Botryosphaeriaceae. *Studies in Mycology*, 55, 235–253. <https://doi.org/10.3114/sim.55.1.235>
- Deshmukh, S. K., Verekar, S. A., & Bhave, S. V. (2015). Endophytic fungi: A reservoir of antibacterials. *Frontiers in Microbiology*, 5, 1–44. <https://doi.org/10.3389/fmicb.2014.00715>
- Devi, R., Kaur, T., Guleria, G., Rana, K. L., Kour, D., Yadav, N., Yadav, A. N., & Saxena, A. K. (2020). Fungal secondary metabolites and their biotechnological applications for human health. In *New and Future Developments in Microbial Biotechnology and Bioengineering* (bll 147–161). Elsevier Inc. <https://doi.org/10.1016/b978-0-12-820528-0.00010-7>
- Druzian, S. P., Pinheiro, L. N., Susin, N. M. B., Dal Prá, V., Mazutti, M. A., Kuhn, R. C., & Terra, L. de M. (2020). Production of metabolites with antioxidant activity by *Botryosphaeria dothidea* in submerged fermentation. *Bioprocess and Biosystems Engineering*, 43(1), 13–20. <https://doi.org/10.1007/s00449-019-02200-y>
- EUCAST Devinitive Document. (1998). Methods for the determination of susceptibility of bacteria to antimicrobial agents. Terminology. *Clinical Microbiol. Infect*, 4(5), 291–296. <https://doi.org/10.1111/j.1469-0691.1998.tb00061.x>
- Fadhillah, Elfita, Muharni, Yohandini, H., & Widjajanti, H. (2019). Chemical compound

- isolated from antioxidant active extract of endophytic fungus *Cladosporium tenuissimum* in *Swietenia mahagoni* leaf stalks. *Biodiversitas*, 20(9), 2645–2650. <https://doi.org/10.13057/biodiv/d200929>
- Fathima, S. B. N., Pandian, R., & Perumal, P. V. (2017). Antimicrobial activity of methanol and ethyl acetate extracts of three medicinal plants. *International Journal of Advances in Science Engineering and Technology*, 5(1), 16–22.
- Felsenstein, J. (1985). Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, 39(4), 783–791.
- Gagana, S. L., & Shivanna, M. B. (2020). Diversity and antibacterial activity of endophytic fungi in *Memecylon umbellatum* Burm. F. a medicinal plant in the Western Ghats of Karnataka, India. *Indian Journal of Ecology*, 47(1), 171–180.
- Gustianingtyas, M., Herlinda, S., Suwandi, Suparman, Hamidson, H., Hasbi, Setiawan, A., Verawaty, M., Elfita, & Arsi. (2020). Toxicity of entomopathogenic fungal culture filtrate of lowland and highland soil of South Sumatra (Indonesia) against *Spodoptera litura* larvae. *Biodiversitas*, 21(5), 1839–1849. <https://doi.org/10.13057/biodiv/d210510>
- Habisukan, U. H., Elfita, Widjajanti, H., Setiawan, A., & Kurniawati, A. R. (2021). Diversity of endophytic fungi in *Syzygium aqueum*. *Biodiversitas*, 22(3), 1129–1137. <https://doi.org/10.13057/biodiv/d220307>
- Hapida, Y., Elfita, Widjajanti, H., & Salni. (2022). *Tritirachium oryzae* and other endophytic mediated jambu bol (*Syzygium malaccense*) are potential as an antioxidant. *Science and Technology Indonesia*, 7(2), 220–227. <https://doi.org/10.26554/sti.2022.7.2.220-227>
- Hu, Z., Wu, Z., Su, Q., Li, M., Wu, S., Meng, R., Ding, W., & Li, C. (2020). Metabolites with phytopathogenic fungi inhibitory activities from the mangrove endophytic fungus *Botryosphaeria ramose*. *Bioorganic Chemistry*, 104, 1–8. <https://doi.org/10.1016/j.bioorg.2020.104300>
- Ju, Z., Lin, X., Lu, X., Tu, Z., Wang, J., Kaliyaperumal, K., Liu, J., Tian, Y., Xu, S., & Liu, Y. (2015). Botryoisocoumarin A, a new COX-2 inhibitor from the mangrove *Kandelia candel* endophytic fungus *Botryosphaeria* sp. KcF6. *The Journal of Antibiotics*, 68(10), 653–656. <https://doi.org/10.1038/ja.2015.46>
- Ju, Z., Qin, X., Lin, X., Wang, J., Kaliyaperumal, K., Tian, Y., Liu, J., Liu, F., Tu, Z., Xu, S., & Liu, Y. (2016). New phenyl derivatives from endophytic fungus *Botryosphaeria* sp. SCSIO KcF6 derived of mangrove plant *Kandelia candel*. *Natural Product Research*, 30(2), 192–198. <https://doi.org/10.1080/14786419.2015.1050670>
- Kataoka, M., Honda, K., Sakamoto, K., & Shimizu, S. (2007). Microbial enzymes involved in lactone compound metabolism and their biotechnological applications. *Applied Microbiology and Biotechnology*, 75(2), 257–266. <https://doi.org/10.1007/S00253-007-0896-X>
- Kowalczyk, P., Gawdzik, B., Trzepizur, D., Szymczak, M., Skiba, G., Raj, S., Kramkowski, K., Lizut, R., & Ostaszewski, R. (2021). δ -Lactones—A new class of compounds that are toxic to *E. coli* K12 and R2–R4 strains. *Materials*, 14(11). <https://doi.org/10.3390/MA14112956>
- Lim, T. K. (2012). *Syzygium jambos*. In *Edible medicinal and non medicinal plants* (bll 760–766). Springer Netherlands. https://doi.org/10.1007/978-94-007-2534-8_101
- Mancuso, G., Midiri, A., Gerace, E., & Biondo, C. (2021). Bacterial antibiotic resistance: The most critical pathogens. *Pathogens*, 10(10), 1–14. <https://doi.org/10.3390/pathogens10101310>
- Manias, D., Verma, A., & Soni, D. K. (2020). Isolation and characterization of endophytes: Biochemical and molecular approach. In *Microbial endophytes*. Elsevier Inc. <https://doi.org/10.1016/b978-0-12-818734-0.00001-2>
- Mazur, M., & Maslowiec, D. (2022). Antimicrobial activity of lactones. *Antibiotics*, 11(10), 1–27.
- Mohali, S. R., Slippers, B., & Wingfield, M. J. (2007). Identification of Botryosphaeriaceae from eucalyptus, acacia and pinus in Venezuela. *Fungal Diversity*, 25, 103–125.
- Nagarajan, D. (2019). In vitro antioxidant potential of endophytic fungi isolated from *Enicostemma axillare* (Lam.) Raynal. and *Ormocarpum cochinchinense* (Lour.) Merr. *Journal of Pharmacognosy and Phytochemistry*, 8(1), 1356–1363.
- Nei, M., & Kumar, S. (2000). *Molecular evolution and phylogenetics*. Oxford University Press, Inc. <https://doi.org/10.1002/wilm.10463>
- Octarya, Z., Novianty, R., Suraya, N., & Saryono. (2021). Antimicrobial activity and GC-MS analysis of bioactive constituents of *Aspergillus fumigatus* 269 isolated from Sungai Pinang hot spring, Riau, Indonesia. *Biodiversitas*, 22(4), 1839–1845. <https://doi.org/10.13057/biodiv/d220429>
- Oliveira, F. C., Barbosa, F. G., Mafezoli, J., Oliveira, M. C. F., Camelo, A. L. M., Longhinotti, E., Lima, A. C. A., Câmara, M. P. S., Gonçalves, F. J. T., & Freire, F. C. O. (2015). Volatile organic compounds from filamentous fungi: A chemotaxonomic tool of the Botryosphaeriaceae family. *Journal of the Brazilian Chemical Society*, 26(11), 2189–2194. <https://doi.org/10.5935/0103-5053>

- 20150204
- Pansanit, A., & Pripdeevech, P. (2018). Antibacterial secondary metabolites from an endophytic fungus, *Arthrinium* sp. MFLUCC16-1053 isolated from *Zingiber cassumunar*. *Mycology*, 9(4), 264–272. <https://doi.org/10.1080/21501203.2018.1481154>
- Phillips, A. J. L., Alves, A., Abdollahzadeh, J., Slippers, B., Wingfield, M. J., Groenewald, J. Z., & Crous, P. W. (2013). The Botryosphaeriaceae: Genera and species known from culture. *Studies in Mycology*, 76, 51–167. <https://doi.org/10.3114/sim0021>
- Pongcharoen, W., Rukachaisirikul, V., Phongpaichit, S., & Sakayaroj, J. (2007). A new dihydrobenzofuran derivative from the endophytic fungus *Botryosphaeria mamane* PSU-M76. *Chemical and Pharmaceutical Bulletin*, 55(9), 1404–1405. <https://doi.org/10.1248/cpb.55.1404>
- Roux, J., Nkuekam, G. K., Marincowitz, S., van der Merwe, N. A., Uchida, J., Wingfield, M. J., & Chen, S. F. (2020). Cryphonectriaceae associated with rust-infected *Syzygium jambos* in Hawaii. *MycKeys*, 76, 49–79. <https://doi.org/10.3897/MYCOKEYS.76.58406>
- Saitou, N., & Nei, M. (1987). The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Molecular biology and evolution*, 4(4), 406–425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>
- Sartori, S. K., Diaz, M. A. N., & Diaz-Muñoz, G. (2021). Lactones: Classification, synthesis, biological activities, and industrial applications. *Tetrahedron*, 84, 1–39. <https://doi.org/10.1016/j.tet.2021.132001>
- Sharma, R., Kishore, N., Hussein, A., & Lall, N. (2013). Antibacterial and anti-inflammatory effects of *Syzygium jambos* L. (Alston) and isolated compounds on *Acne vulgaris*. *BMC Complementary and Alternative Medicine*, 13(292), 1–10. <https://doi.org/10.1186/1472-6882-13-292>
- Shi, Y., Zhang, X., & Lou, K. (2013). Isolation, characterization, and insecticidal activity of an endophyte of drunken horse grass, *Achnatherum inebrians*. *Journal of Insect Science*, 13(151), 1–12. <https://doi.org/10.1673/031.013.15101>
- Silva, A. A. da, Polonio, J. C., Bulla, A. M., Polli, A. D., Castro, J. C., Soares, L. C., Oliveira-Junior, V. A. de, Vicentini, V. E. P., Oliveira, A. J. B. de, Gonçalves, J. E., Gonçalves, R. A. C., Azevedo, J. L., Abreu-Filho, B. A. de, & Pamphile, J. A. (2021). Antimicrobial and antioxidant activities of secondary metabolites from endophytic fungus *Botryosphaeria fabiceriana* (MGN23-3) associated to *Morus nigra* L. *Natural Product Research*, 0(0), 1–5. <https://doi.org/10.1080/14786419.2021.1947272>
- Singh, P., Kapkoti, D. S., Singh, N., Tewari, R., Saikia, D., Rout, P. K., Pandey, R., & Bhakuni, R. S. (2019). Biotransformation of artemisinic acid by the fungus *Trichothecium roseum* and anti-candidal activity of its metabolites. *Biocatalysis and Biotransformation*, 37(4), 304–309. <https://doi.org/10.1080/10242422.2018.1552265>
- Singha, I. M., Kakoty, Y., Unni, B. G., Das, J., & Kalita, M. C. (2016). Identification and characterization of *Fusarium* sp. using ITS and RAPD causing Fusarium wilt of tomato isolated from Assam, North East India. *Journal of Genetic Engineering and Biotechnology*, 14(1), 99–105. <https://doi.org/10.1016/j.jgeb.2016.07.001>
- Slama, H. Ben, Bouket, A. C., Alenezi, F. N., Pourhassan, Z., Golińska, P., Oszako, T., & Belbahri, L. (2021). Potentials of endophytic fungi in the biosynthesis of versatile secondary metabolites and enzymes. *Forests*, 12(12), 1–13. <https://doi.org/10.3390/f12121784>
- Syarifah, Elfita, Widjajanti, H., Setiawan, A., & Kurniawati, A. R. (2021). Diversity of endophytic fungi from the root bark of *Syzygium zeylanicum*, and the antibacterial activity of fungal extracts, and secondary metabolite. *Biodiversitas*, 22(10), 4572–4582. <https://doi.org/10.13057/biodiv/d221051>
- Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, 38(7), 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Taware, R., Abnave, P., Patil, D., Rajamohananan, P. R., Raja, R., Soundararajan, G., Kundu, G. C., & Ahmad, A. (2014). Isolation, purification and characterization of trichothecinol-A produced by endophytic fungus *Trichothecium* sp. and its antifungal, anticancer and antimetastatic activities. *Sustainable Chemical Processes*, 2(8), 1–9. <https://doi.org/10.1186/2043-7129-2-8>
- Tawfike, A. F., Romli, M., Clements, C., Abbott, G., Young, L., Schumacher, M., Diederich, M., Farag, M., & Edrada-Ebel, R. A. (2019). Isolation of anticancer and anti-trypanosome secondary metabolites from the endophytic fungus *Aspergillus flocculus* via bioactivity guided isolation and MS based metabolomics. *Journal of Chromatography B*, 1106–1107(October 2018), 71–83. <https://doi.org/10.1016/j.jchromb.2018.12.032>
- Tiwari, P., & Bae, H. (2022). Endophytic fungi: key insights, emerging prospects, and challenges in natural product drug discovery. *Microorganisms*, 10(2), 1–34. <https://doi.org/10.3390/microorganisms10020360>
- Valente, I. de L., Confortin, T. C., Luft, L., Ugalde, G.

- A., Zabot, G. L., Mazutti, M. A., & Terra, L. de M. (2018). Extraction of bioactive compounds from *Botryosphaeria dothidea* using supercritical carbon dioxide and compressed liquefied petroleum gas. *Journal of Supercritical Fluids*, 136, 52–59. <https://doi.org/10.1016/j.supflu.2018.02.013>
- Walsh, T. H., Hayden, R. T., & Larone, D. H. (2018). *Larone's medically important fungi: A guide to identification* (6th Edition). ASM Press, Washington, DC. <https://doi.org/10.1128/9781555819880> Printed
- Wamba, B. E. N., Nayim, P., Mbaveng, A. T., Voukeng, I. K., Dzotam, J. K., Ngalani, O. J. T., & Kuete, V. (2018). Syzygium jambos displayed antibacterial and antibiotic-modulating activities against resistant phenotypes. *Evidence-based Complementary and Alternative Medicine*, 2018. <https://doi.org/10.1155/2018/5124735>
- Watanabe, T. (2010). *Pictorial atlas of soil and seed fungi: Morphologies of cultural fungi and key to species* (3th Edition). CRC Press: Taylor & Francis Group. <https://doi.org/10.1201/ebk1439804193>
- Wawrosch, C., & Zotchev, S. B. (2021). Production of bioactive plant secondary metabolites through in vitro technologies—status and outlook. *Applied Microbiology and Biotechnology*, 105(18), 6649–6668. <https://doi.org/10.1007/s00253-021-11539-w>
- Wińska, K., Grabarczyk, M., Mączka, W., Żarowska, B., Maciejewska, G., & Anioł, M. (2018). Antimicrobial activity of new bicyclic lactones with three or four methyl groups obtained both synthetically and biosynthetically. *Journal of Saudi Chemical Society*, 22(3), 363–371. <https://doi.org/10.1016/J.JSCS.2016.04.004>
- Wu, Z., Chen, J., Zhang, X., Chen, Z., Li, T., She, Z., Ding, W., & Li, C. (2019). Four new isocoumarins and a new natural tryptamine with antifungal activities from a mangrove endophytic fungus *Botryosphaeria ramosa* L29. *Marine Drugs*, 17(88), 1–9. <https://doi.org/10.3390/md17020088>
- Xiao, J., Zhang, Q., Gao, Y. Q., Tang, J. J., Zhang, A. L., & Gao, J. M. (2014). Secondary metabolites from the endophytic *Botryosphaeria dothidea* of *Melia azedarach* and their antifungal, antibacterial, antioxidant, and cytotoxic activities. *Journal of Agricultural and Food Chemistry*, 62(16), 3584–3590. <https://doi.org/10.1021/jf500054f>
- Yodsing, N., Lekphrom, R., Sangsopha, W., Aimi, T., & Boonlue, S. (2018). Secondary metabolites and their biological activity from *Aspergillus aculeatus* KKU-CT2. *Current Microbiology*, 75(5), 513–518. <https://doi.org/10.1007/s00284-017-1411-y>
- Youssef, F. S., Alshammari, E., & Ashour, M. L. (2021). Bioactive alkaloids from genus *Aspergillus*: Mechanistic interpretation of their antimicrobial and potential SARS-CoV-2 inhibitory activity using molecular modelling. *International Journal of Molecular Sciences*, 22(4), 1–23. <https://doi.org/10.3390/ijms22041866>
- Zhang, Z., Zhang, W., Bi, Y., Han, Y., Zong, Y., & Prusky, D. (2020). Cuminal inhibits *Trichothecium roseum* growth by triggering cell starvation: Transcriptome and proteome analysis. *Microorganisms*, 8(256), 1–33. <https://doi.org/10.3390/microorganisms8020256>
- Zhou, Y. M., Ju, G. L., Xiao, L., Zhang, X. F., & Du, F. Y. (2018). Cyclodepsipeptides and sesquiterpenes from marine-derived fungus *Trichothecium roseum* and their biological functions. *Marine Drugs*, 16(12), 1–12. <https://doi.org/10.3390/md16120519>



Molekul

MOLEKUL is migrating to a new website. For articles published in 2022 and forward, please submit on our new website at <http://jos.unsoed.ac.id/index.php/jm/index>

Metrics

	2019	2020	2021
Scopus CiteScore	0.1	0.3	0.7
Copernicus ICV	99.19	100	
Google Scholar h-index	10	12	13
Acceptance rate	72%	46%	40%

ISSN: 1907-9761 (print) , 2503-0310 (electronic)

MOLEKUL is a scientific journal in chemistry, **focus on** the regional resources, environment or natural products of Indonesia and nearby tropical country.

The scope of this journal including :

- Theoretical chemistry and environmental chemistry study
- Material synthesis for energy and environmental
- Isolation, purification and modification of biomolecules, including their applications

[Guide for Authors](#)

[View Articles](#)

[Submit your paper](#)

[Editorial Board](#)

[Focus and Scope](#)

[Publication Ethics](#)

[Peer Review Process](#)

[Plagiarism Policy](#)

Molekul

Q3

Chemistry
(miscellaneous)

best quartile

SJR 2021

0.26

powered by scimagojr.com

Indexing and Abstracting:

Scopus®

- Fabrication, development and validation of analytical methods

MOLEKUL is published by the Department of Chemistry, Faculty of Mathematics and Natural Sciences, Jenderal Soedirman University, Indonesia. Previously, the Molekul published 2 issues per year, on May and November. Since 2020, Molekul publishes 3 issues per year, on March, July, and November.

History

Molekul has been established since 2006 with volume 1 no 1 (November 2006). The name of the birth of certificate from the Universitas Jenderal Soedirman was "Jurnal Ilmiah Kimia "Molekul". Jurnal Ilmiah Kimia mean Scientific Journal in Chemistry. The journal name in the quotes was Molekul. Furthermore, the journal cover used the name MOLEKUL and a small Jurnal Ilmiah Kimia. The running journal name in all articles is Molekul. Print ISSN name was Molekul, electronic ISSN was Molekul: Jurnal Ilmiah Kimia. The name of the journal in the National Accreditation certificate was Jurnal Ilmiah Kimia Molekul. The final name of the journal should be only **Molekul**.



Molekul

Jurnal Ilmiah Kimia
Department of
Chemistry, Faculty of
Mathematics and
Natural Sciences,
Universitas Jenderal
Soedirman,
Purwokerto, Indonesia



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

INDEX COPERNICUS INTERNATIONAL

DOAJ DIRECTORY OF OPEN ACCESS JOURNALS

CAS[®]
 A DIVISION OF THE AMERICAN CHEMICAL SOCIETY

Google scholar

Crossref

SINTA
 Science and Technology Index



[Detail Statistic](#)

Visitors		
ID 80,787	RU 326	
US 4,528	TR 236	
IN 1,649	KR 226	
PH 1,025	IR 220	
CN 983	TW 205	
MY 750	DE 205	
TH 593	GB 196	
JP 502	NG 186	
BR 450	SG 180	
VN 335	PK 180	
Pageviews: 274,269		
FLAG counter		

Dokumen Akreditasi

- [Sertifikat Akreditasi 2015](#)
- [SK Akreditasi 2018](#)
- [SK Akreditasi 2020](#)
- [Sertifikat Akreditasi 2020](#)

USER

Username

Password

Remember me

HOME / Editorial Team

Editor-in-Chief

- [Dr. Amin Fatoni](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia, Indonesia

Managing Editor

- Ely Setiawan, Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

Editorial Board

- [Prof. Shin-ichiro Suye](#), Department of Applied Chemistry and Biotechnology, University of Fukui, Fukui, Japan
- [Prof. Dr. Wan Aini Wan Ibrahim](#), Universiti Teknologi Malaysia, Malaysia
- [Prof. Pierre Doumeng](#), Aix Marseille University, France.
- [Prof. Mohammed M. Rahman](#), King Abdulaziz University, Saudi Arabia
- [Prof. Dr. Kristian H Sugijarto](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Yogyakarta State University, Indonesia
- [Prof. Dr. drh. Maria Bintang](#), Department of Biochemistry, Faculty of Mathematics and Natural Sciences, Institut Pertanian Bogor,, Indonesia
- [Prof. Dr. Hassan Y Aboul-Enein](#), National Research Centre, Cairo, Egypt
- [Prof. Dr. Hadi Nur](#), Centre for Sustainable Nanomaterials, Ibnu Sina Institute for Scientific and Industrial, Universiti Teknologi Malaysia, Malaysia
- [Dr. Arief Cahyo Wibowo](#), King Fahd University of Petroleum & Minerals, Saudi Arabia
- [Dr. Adi Darmawan](#), Chemistry Department, Faculty of Sciences and Mathematics, Diponegoro University Jl. Prof. Soedarto, SH., Tembalang, Semarang, Indonesia
- [Assist. Prof. Dr. Apon Numnuam](#), Department of Chemistry, Faculty of Science, Prince of Songkla University, Hat Yai, Thailand.

- [Dr. Uyi Sulaeman](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia, Indonesia
- [Dr. Jas Raj Subba](#), Royal University of Bhutan, Bhutan
- [Dr. Hartiwi Diastuti](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia
- [Dr. Saluma Samanman](#), Faculty of Science and Technology, Princess of Naradhiwas University, Narathiwas, Thailand
- [Mardiyah Kurniasih](#), Department of Chemistry, Faculty of Mathematic and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia
- [Zusfahair](#), Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

Layout Editor

-
- Irmanto Irmanto, Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia

MAKE A SUBMISSION

FOCUS AND SCOPE
GUIDE FOR AUTHORS
VIEW ARTICLES
EDITORIAL BOARD
PUBLICATION ETHICS
PEER REVIEW PROCESS
PLAGIARISM POLICY



Ads by Google

[Stop seeing this ad](#)[Why this ad? ⓘ](#)

Molekul

COUNTRY

Indonesia

Universities and
research institutions in
IndonesiaMedia Ranking in
Indonesia**SUBJECT AREA AND CATEGORY**Biochemistry, Genetics and
Molecular Biology
BiochemistryChemistry
Chemistry
(miscellaneous)Environmental Science
Environmental
ChemistryMaterials Science
Biomaterials**PUBLISHER**Universitas Jenderal
Soedirman**H-INDEX**

4

PUBLICATION TYPE

Journals

ISSN

19079761, 25030310

COVERAGE

2019-2021

INFORMATION[Homepage](#)[How to publish in this
journal](#)j.molekul@gmail.com**SCOPE**

MOLEKUL is a scientific journal in chemistry, focus on the regional resources, environment or natural products of Indonesia and nearby tropical country. The scope of this journal including : • Theoretical chemistry and environmental chemistry study • Material synthesis for energy and environmental • Isolation, purification and modification of biomolecules, including their





applications • Fabrication, development and validation of analytical methods MOLEKUL is published by the Department of Chemistry, Faculty of Mathematics and Natural Sciences, Jenderal Soedirman University, Indonesia. Previously, the Molekul published 2 issues per year, on May and November. Since 2020, Molekul publishes 3 issues per year, on March, July, and November.





Join the conversation about this journal

Quartiles

FIND SIMILAR JOURNALS ?

Rank	Journal Name	Country	Similarity
1	Pharmacognosy Journal	IND	38% similarity
2	Natural Products Journal	NLD	37% similarity
3	Journal of Biologically Active Products from Nature	GBR	35% similarity
4	Revista Brasileira de Farmacognosia	BRA	34% similarity





Molekul

Q3
Chemistry (miscellaneous)
best quartile

SJR 2021
0.26

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

`<a href="https://www.scimaç`

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2022

Leave a comment