

## COMPARATIVE STUDY ON THE PHYTOCHEMICAL AND *IN VITRO* ANTIOXIDANT PROPERTIES OF METHANOLIC LEAF EXTRACT OF *JUSTICIA SECUNDA* VAHL

Osioma, E.<sup>1</sup> and Hamilton- Amachree, A.<sup>2\*</sup>

<sup>1</sup>Department of Biochemistry, Faculty of Science, Federal University Otuoke, Bayelsa State, Nigeria.

<sup>2</sup>Department of Chemical Sciences, Faculty of Science, Federal University Otuoke, Bayelsa State, Nigeria.

\*Corresponding author. E-mail: hamiltonaa@fuotuo.ke.edu.ng.

Accepted 22nd July, 2017

Medicinal plants are a primary source of healing and widely used for the treatment of various ailments. This present study evaluates phytochemicals and *in vitro* antioxidant activities of the methanolic leaf extract of *Justicia secunda*, *Acanthaceae* family. Fresh leaves of the plant were collected, and shade dried. The dry leaves were pulverized, and the resultant powder extracted with 0.1% hydrogen chloride (HCl) in methanol, and 1% HCl in methanol respectively. Extracts obtained were subjected to qualitative phytochemical screening and *in vitro* antioxidant assays. Results obtained confirm the presence of saponins, tannins, steroids, flavonoids, and alkaloids in equal degree in both 0.1% and 1% HCl methanolic leaf extract. Concentrations of anthocyanins, phenols and phlobatannins were quantitatively higher in 0.1% HCl extract as compared to 1% HCl extract. However, the presence of carbohydrates was not confirmed in either extracts. The results also indicated that phenolic and anthocyanins content of the leaf extracts were significantly higher ( $p < 0.05$ ) in 0.1% HCl extract as compared to the 1% HCl leaf extract. Also, flavonoids contents were comparable ( $p > 0.05$ ) in both extracts. The antioxidant scavenging efficacy showed that higher percentage inhibition was observed from the 0.1% HCl methanolic extract of *Justicia secunda* in its ability to scavenge hydrogen peroxide, 1,1 – diphenylpicrylhydrazyl, and superoxide anion. Overall, from this study, it could be concluded that *Justicia secunda* may be a good source of phytochemicals and natural antioxidants that will be useful in the food, medicine, and pharmaceutical industries.

**Key words:** *Justicia secunda*, anthocyanins, phenols, flavonoids, phlobataninins, phytochemicals.

### INTRODUCTION

*Justicia* is a creeping perennial herb which grows on sand in shaded forest floor, and it is about 1 to 1.5 m height. Ethno pharmacological information according to Correa et al. (2012), suggests that lignans are the major components of the active extracts of the species of *Justicia* exhibiting important pharmacological properties such as antiviral, antitumoral, anti-inflammatory and anti-platelet – aggregation.

Mpiana et al. (2010a) opine that *Justicia tenella*, *Justicia gendarussa*, and *Justicia insularis* possess high anti-sickling potency as a result of the high levels of anthocyanins they contain; while Mpiana et al. (2010b) reported the *in vitro* effects of anthocyanin extracted from *Justicia secunda* vahl on the solubility of sickle erythrocyte.

Three compounds: lupeol, 16-hydroxyl lupeol and stigmasterol have also been isolated from *J. gendarussa* with promising antioxidant properties (Uddin et al., 2011).

The plant *J. secunda* belongs to the *Acanthaceae* family. It is locally called ‘*Asindiri*’ or ‘*Ohowaazara*’ (meaning medicine that gives blood) by the Ogbia people of Otuoke-Otuaba community in Ogbia Local Government Area of Bayelsa State, and other communities in the Niger Delta region of Nigeria. It is used for the treatment of anaemia. Other common names are ‘*Hospital too far*’, ‘*Blood leaf*’ or *Blood tonic* (Roberts, 2016). Its aqueous extract is normally served as a tea drink to the anaemic patients in these communities.

Phytochemicals are naturally present in the medicinal plants, leaves, vegetables and roots that have protective mechanism and defend plants

from various infections; also called secondary metabolites. They are non – nutritive compounds that confer flavor and colour to the plants (Agbafor and Nwachukwu, 2011). These include alkaloids, terpenoids, and phenolic compounds.

Over the centuries, secondary metabolites in plants have played considerable role in medicine, being used in their natural state or their derivatives (Kinghorn et al., 2011; Newman and Cragg, 2012). One principal characteristic of phytochemicals that support its importance in medicine, food, pharmaceuticals, and nutraceuticals is their ability to function as antioxidants (Potterat, 1997; Khan et al., 2003; Prakash et al., 2012)

Antioxidants, according to Husseni et al. (2011) are substances that are capable of quenching the oxidation reaction of free radicals. Its mechanism may be through playing a direct role in radical scavenging or indirectly through destroying consumed substances that easily generate free radicals, thus preventing the occurrence of further reactions. However, there is a global shift from the use of synthetic antioxidants to naturally occurring antioxidants due to the former's harmful effect to human health due to their potential toxicity and carcinogenicity (Thaipong et al., 2006). Owing to the abundance of *J. secunda* in the lowland rainforests of the Niger Delta region of Nigeria, and its use by the indigenous people for curing certain ailments, this research reports its preliminary phytochemical and *in vitro* antioxidants properties.

## MATERIALS AND METHODS

### Plant materials

Fresh *J. secunda* leaves were collected from a garden in Otuoke village, Bayelsa State, Nigeria on May 2016. It was authenticated in the University of Port Harcourt herbarium by Dr. Ekeke Chimezie with voucher number UPH/V/1279.

### Preparation of extract

The leaves of the plant were carefully cleaned and washed with tap water to remove impurities followed by shade drying. The air-dried leaves were pulverized and the resultant

powder was stored in air-tight bottles at 20°C. One part (600mg) of the powdered sample was soaked in 1500ml of 0.1% HCl methanol; while another part was soaked in 1% HCl methanol for 72 h at room temperature (25°C), and then filtered through a cotton plug followed by Whatman filter paper No.1. The parent extract was concentrated with a Rotary Evaporator and stored in the refrigerator until required for use.

### Qualitative phytochemical screening

Phytochemical screening was performed on parent plant extracts using standard procedures to qualitatively identify its constituents. The re-constituted aqueous crude extract (10mg / 20ml of distilled H<sub>2</sub>O) obtained was used for the experiment. The methods of Evans (1997) and Sofowara (2006) were used for the detection of alkaloids and flavonoids respectively. The presence of tannins, phenolics, and saponins were detected by employing standard methods by Segelman et al. (1969), Mukherjee (2002) and Kokate (1999) respectively. The method of Edeogal et al. (2005) was employed to confirm the presence of phlobatannins and steroids in the plant extract. Screening for anthocyanins and carbohydrates were done using standard methods by Paris and Moyses (1969), and Rajagopal and Ramakrishnan (1993) respectively.

### Estimation of *In vitro* Antioxidant Properties in Methanolic Leaf Extract of *J. secunda*

Employing previously described standard procedures, flavonoids and total phenolic content in the methanolic leaf extract of *J. secunda* were estimated according to Chang et al. (2002) and Singleton and Rossi (1965). The concentration of anthocyanins was determined according to the method by Lapornik et al. (2005). The hydroxyl radical scavenging ability was assayed by the method of Kaur et al. (2006), while the methods of Abdel – Hameed (2009) and Fontana et al. (2001) were respectively employed for the assay of DPPH radical and superoxide anion scavenging activities.

### Statistical analysis

The values of all the parameters determined are expressed as Mean  $\pm$  SD for triplicate determinations, and all data were analyzed using one-way ANOVA. The level of significance was placed at  $p < 0.05$ . All statistical

analyses were performed using SPSS version 16.0.

## RESULTS

### Qualitative phytochemical screening

The results of the qualitative phytochemical screening of the methanolic leaf extracts of *J. secunda* are presented in Table 1. The presence of flavonoids, tannins, phenols,

saponins, and steroids were qualitatively confirmed to the same degree for both 0.1% and 1% HCl methanolic leaf extracts of *Justicia secunda*. Alkaloids, phlobatannins, and anthocyanins were observed more in 0.1% HCl methanolic leaf extract as compared with 1% HCl methanolic leaf extract of *Justicia tenuipes*. The table also revealed that carbohydrate was not observed in either 0.1% or 1.0% HCl extract of the plant by the screening procedure used.

**Table 1.** Phytochemical constituent of the methanolic leaf extracts of *J. secunda*.

Phytochemicals	0.1% HCl in methanol extract	1.0% HCl in methanol extract
Saponins	+ + +	+ + +
Tannins	+ + +	+ + +
Carbohydrate	- - -	- - -
Steroids	+ + +	+ + +
Flavonoids	+ + +	+ + +
Anthocyanins	+ + +	+ +
Phenols	+ + +	+ +
Alkaloids	+ + +	+ + +
Phlobatannins	+ + +	+ +

+ = Presence of constituent in the leaf extract; - = Absence of constituent in the leaf extract.

### Concentrations of flavonoids, phenols and anthocyanins

The concentrations of flavonoids, total phenolic and anthocyanin contents in both 0.1 and 1.0% HCl methanolic leaf extracts of *J. secunda* are depicted in Table 2. The results

indicated that 0.1% HCl methanolic leaf extract of the plant showed a significantly ( $p < 0.05$ ) higher total phenolic and anthocyanin content as compared with 1.0% HCl plant extract. However, the concentrations of flavonoids were comparable between 0.1 and 1.0% HCl methanolic leaf extracts.

**Table 2.** Flavonoids, total phenolic and anthocyanin contents in 0.1 and 1% HCl methanolic leaf extracts of *J. secunda*.


Phytochemicals	0.1% extract	1.0% extract
Flavonoid content (mg/g of rutin)	2.67±0.31 <sup>a</sup>	2.56±0.1 <sup>a</sup>
Phenolic Content (mg/100ml of extract)	6.88±0.33 <sup>a</sup>	4.47±0.41 <sup>b</sup>
Anthocyanin content (µg/g cyanidin)	9.72 ± 0.41 <sup>a</sup>	8.59±0.50 <sup>b</sup>

Values are expressed as Mean ±SD for triplicate determinations; Means sharing different superscript alphabet on same row differ significantly at  $p < 0.05$ .

### In vitro free radical scavenging efficacy

*In vitro* free radical scavenging efficacy of 0.1 and 1.0% HCl methanolic leaf extract of *J. secunda* were assessed by its ability to scavenge hydrogen peroxide, diphenylpicrylhydrazyl (DPPH) and superoxide anion radical. Figure 1 illustrates

the % inhibition of H<sub>2</sub>O<sub>2</sub>, DPPH, and superoxide anion radicals. It was observed from the results that 0.1% HCl methanolic extracts showed a significantly ( $p < 0.05$ ) higher % inhibition in both H<sub>2</sub>O<sub>2</sub>, DPPH, and superoxide anion scavenging abilities as compared with the 0.1% HCl methanolic leaf extracts of *Justicia secunda*.



**Figure 1.** Hydrogen peroxide scavenging ability, DPPH radical scavenging ability and superoxide anion scavenging ability of methanolic leaf extracts of *J. secunda* (Values are expressed as Mean ± SD % inhibition of triplicate determinations; Group bars having the same alphabet does not differ significantly at  $P < 0.05$ ).

## DISCUSSION

In this present study, the qualitative phytochemical screening of the acidified methanolic leaf extract of *J. secunda* confirmed the presence of flavonoids, alkaloids, phenols, saponins, tannins, steroids, anthocyanins, and phlobatannins; but the absence of carbohydrates. The protective properties ascribed to phytochemicals are as a result of their ability to either react with free radicals or inhibit the process of oxidation in cells. In other words, phytochemicals act as either antiradical or antioxidant compounds (Chanda and Dave, 2009; Samatha et al., 2012; Suprava et al., 2013).

Phenols according to Shahidi et al. (1992) are a class of antioxidant agents which act as free radical terminators and their bioactivities may be related to their ability to chelate radicals, inhibit lipoxygenase and scavenge free radicals. The results from this study indicate high phenolic content in 0.1% HCl methanolic leaf extract. Flavonoid concentration was also high in the experimental plant. Flavonoids represent one of the most important sources of natural phenols. They possess a broad spectrum of chemical and biological activities including radical scavenging properties. Flavonoids has also been associated with possible role in the

prevention of several chronic diseases involving oxidative stress as well as their protective effect against low – density lipoprotein oxidation (Das and Pereira, 1990; Shanaz et al., 2011)

Generally, acidified alcoholic solvents are the choice solvents for the extraction of anthocyanins (Devi et al., 2011; Kang et al., 2013; Puertolas et al., 2013). The acid contributes to stabilize anthocyanins but it may also transform the original form of the pigment in the tissue by breaking associations with metals, co-pigments or other factors (Strack and Wray, 1994).

Higher ( $p < 0.05$ ) anthocyanin content observed in the 0.1% HCl methanol leaf extract as compared with the 1% HCl methanolic leaf extract in this study could be due to minimal anthocyanin degradation during concentration procedures in solvent recovering (Jackman et al., 1987). The high anthocyanin content observed could enhance the plant's potential as a good source for food colourant production and natural antioxidant because anthocyanins are responsible for colouration in plants; and their consumption has been linked as protective agent against many chronic diseases (Welch et al., 2008)

Assay of the *in vitro* antioxidant scavenging abilities in this study revealed that *J. secunda* may possess free radical scavenging activities. Hydrogen peroxide, DPPH and

Superoxide anion scavenging radical abilities has been previously employed to assess the *in vitro* antioxidant scavenging properties of plants (Malik et al., 2011; Spreena et al., 2011; Singh et al., 2012; Nwaoguikpe et al., 2014 and Seo et al., 2014).

Results of this study showed significantly higher scavenging properties (that is, H<sub>2</sub>O<sub>2</sub>, DPPH, Superoxide anion radical scavenging properties). The observed high antioxidant scavenging properties in *J. secunda* leaf extract could be attributed to the higher concentrations of flavonoids, phenols, and anthocyanins in the leaf extract of the plant (Table 2). Together, the phytochemicals contained in the leaf of *J. secunda* coupled with the antioxidant activities as exhibited by the plant's ability to scavenge H<sub>2</sub>O<sub>2</sub>, DPPH, and superoxide anion radical could justify its use locally for medicinal purposes.

Conclusively, the leaves of *J. secunda* may serve as a significant source of natural antioxidants which may be helpful in preventing the progress of various oxidative stresses. However, the components responsible for the anti-oxidative activity are currently unclear. Therefore, further investigation is needed to isolate and identify the antioxidant components present in the plant leaf extract. Also, *in vivo* anti-oxidant of the leaf extracts need to be assessed prior to clinical use.

### CONFLICT OF INTERESTS

The authors declare that there is no conflict of interests.

### ACKNOWLEDGEMENTS

This research was supported by the Year 2014 TETFund Institution Based Research (IBR) 4<sup>th</sup> Batch Intervention granted to Dr (Mrs) Hamilton Amachree Akens of the Department of Chemistry, Faculty of Science, Federal University Otuoke, Bayelsa State, Nigeria.

### REFERENCES

- Abdel – Hameed, E.S.S. (2009). Total Phenolic contents and free radical Scavenging activity of certain Egyptian Ficus Species leaf Samples. *Food Chem.* Pp: 1271 – 1277.
- Agbafor, K.N. and Nwachukwu, N. (2011). Phytochemical Analysis and Antioxidant property of Extracts of *Vitex doniana* and *Mucuna pruriens*. *Biochem. Res. Intern*10: 1155 – 1159.
- Chanda, S., and Dave, R. (2009). *In vivo* models for antioxidant activity evaluations and some medicinal plants possessing antioxidant properties: An overview. *Afr. J. Microbiol. Res.* 3(13): 981 – 996.
- Chang, C., Yang, M., Wen, H and Chern, J. (2002). Estimation of total flavonoid content in Propolis by two complementary colorimetric methods. *J. Food & Drug Anal.* 10(3): 178-182.
- Correa, G.M. and Alcantara, A.F. de C. (2012). Chemical constituents and biological activities of species of *Justicia* – a review. *Braz. J. Pharmacognosy.* 22(1): 220 – 238.
- Das, N. P. and Pereira, T. A. (1990). Effect of flavonoids on thermal auto oxidation of Palm oil: structure–activity relationship. *J. Amer. Oil Chem. Soc.* 44: 255-258.
- Devi, P. S., Saravanakumar, M. and Mohandas, S. (2011). Identification of 3 – deoxyanthocyanins from red sorghum (*Sorghum bicolor*) bran and its biological properties. *Afr. J. Pure Appl. Chem.* 5(7): 181 – 193.
- Edeoga, H.A., Okwu, D.E. and Mbaebie, B.O. (2005). Phytochemical constituent of some Nigerian Medicinal Plants. *Afr. J. Biotechnol.* 4: 685 – 688.
- Evans, W.C. (1997). *Pharmacology*. Harcourt Brace and Company. Asia, Singapore.pp: 226
- Fontana, M., Mosca, L. and Rosel, M.A. (2001). Interaction of enkephalines with oxyradicals. *Biochem Pharmacol.* 61: 1253 – 1257.
- Hussein, S. Z., Yusoff, K. M., Makpol, S. and Yusuf, Y. A. M. (2011). Antioxidant capacities and total phenolic contents increase with gamma irradiation in two types of Malaysian honey. *Molecules.* 16(8), 6378-6395.
- Jackman, R.L., Yada, R.Y. and Tung, M.A. (1987). A Review: Separation and chemical properties of anthocyanins used

- for their qualitative and quantitative analysis. *J. Food Biochem.* 11: 279 – 308.
- Kang, Y. J., Jung, S. W. and Lee, S. J. (2013).** An optimal extraction solvent and purification adsorbent to produce anthocyanins from black rice (*Oryza sativa* cv. Heugjinjubyeo). *Food Sci. Biotechnol.* 23: 97 – 106.
- Kaur, G., Alam, M.S., Jabbar, Z., Javed, K. and Athar, M. (2006).** Evaluation of antioxidant activity of *Cassia siamea* flowers. *J. Ethnopharmacol.* 108: 340 – 348.
- Khan, M.R., Kihara, M. and Omoloso, A.D. (2003).** Antimicrobial activity of alkaloidal constituents of the root bark of *Eupomatia laurina*. *Pharm Biol.* 41(4): 277 – 280.
- Kinghorn, A. D., Pan, L., Fletcher, J.N. and Chai, H. (2011).** The relevance of higher plants in lead compound discovery programs. *J. Nat. Prod.* 74: 1539 – 1555.
- Kokate, C.K.(1999).** Practical Pharmacognosy. 4<sup>th</sup> edn. Vallabh Prakashan Publication. New Delhi India. Pp: 111 – 116.
- Lapornik, B., Prosek, M., and Wondra, A. (2005).** Comparison of extracts prepared from plant by-products using different solvents and extraction time. *J. Food Engr.* 71:214-222.
- Malik, A., Kushnoor, A., Sanini, V., Singhai, S., Kumar, S. and Yadav, Y.C. (2011).** *In vitro* Antioxidant properties of *Scopoletin*. *J. Chem. Pharm. Res.* 3(3): 659 - 665
- Mpiana, P.T., Bokota, M.T., Ndjele, M.B.L., Mudogo, V., Tshibangu, D.S.T., Ngbolua, K.N., Atibu, E.K., Kwembe, J.T.K. and Makelele, L.K. (2010a).** Antisickling activity of three species of *Justicia* from Kisangani (D.R. Congo): *J. tenella*, *J. gendarussa* and *J. insularis*. *Int. J. Biol. Chem. Sci.* 4(6): 1953-1961.
- Mpiana, P.T., Ngbolua, K.N., Bokota, M.T., Kasonga, T.K., Atibu, E.K. and Mudogo, V. (2010b).** *In vitro* Effects of anthocyanins extracts from *Justicia secunda* Vahl on the solubility of hemoglobin S and membrane stability of sickle erythrocytes. *Blood Transfusion.* 4: 1 - 8.
- Mukherjee, P.K. (2002).** Quality control of Herbal Drugs, Business Horizons Pharmaceutical Publishers, New Delhi, India. pp: 356 – 358.
- Newman, D.J. and Cragg, G.M. (2012).** Natural products as sources of new drugs over the 30 years from 1981 to 2010. *J. Nat. Prod.* 75: 311 - 335
- Nwaoguikpe, R. N., Ujowundu, C. O. and Emejulu, A. A.(2014).** The Antioxidant and free radical scavenging effects of extracts of seeds of some neglected legumes of South - East Nigeria. *Sch. Acad. J. Biosci.* 2(1): 51 – 59.
- Paris, R. and Moyse, H. (1969).** *Precis de matiere medicinale.* Paris: Masson.
- Potterat, O. (1997).** Antioxidants and free radical scavengers of natural origin. *Curr. Organic Chem.* 14: 415 – 440.
- Prakash, D., Gupta, C. and Sharma, G. (2012).** Importance of phytochemicals in nutraceuticals. *J. Chinese Med. Res & Dev.* 1(3): 70 – 78.
- Puertolas, E., Cregenzan, O., Luengo, E., Alvarez, I. and Raso, J. (2013).** Pulsed electric-field-assisted extraction of anthocyanins from purple-fleshed potato. *Food Chem.* 136: 1330 - 1336.
- Rajagopal, G. and Ramakrishnan, S. (1993).** Qualitative analysis of Carbohydrates. Practical biochemistry for medical students. Orient Longman. Pp: 1 -3.
- Samatha, T., Acharya, R.S., Srinivas, P. and Ramaswamy, N. (2012).** Quantification of total phenolics and total flavonoid contents in extract of *Oroxylum indicum* L. *Kurz. Asian J. Pharm Clin. Res.* 5(4): 177 – 179.
- Segelman, A.B., Farnsworth, N.R. and Quimby, M.D. (1969).** False-negative saponins test results induced by the presence of tannins. *Lloydia:* 32:52-58.
- Seo, J., Lee, S., Elam, M.L., Johnson, S.A., Kang, J. and Arjmandi, B.H. (2014).** Study to find the yeast extraction solvent for use with guava leaves (*Psidium guajava* L.) for high antioxidant efficacy. *Food Sci. Nutr.* 2(2): 174 – 180.

- Shahidi, F., Janitha, P.K. and Wanasundara, P.D. (1992).** Phenolic antioxidants. *Critical Rev. Food Sci. Nutr.* 32: 67-103
- Shanaz, B., Arunachalam, G.1., Jayaveera, K. N., Ashoka – Babu, V. L. and K. B. Premakumari, K. B. (2011).** Estimation of total phenolic content and in vitro antioxidant activity of *Barleria Montana*. *Der Pharmacia Lettre.* 3(4): 178 – 182.
- Singh, P., Vishwakarma, S.P., Singh, U., Shukla, M., Singh, R., Singh, R. K., Singh, R.B., Wilson, D. W. and Singh, R.L. (2012).** Quantification and evaluation of antioxidant activity of some bioactive phytochemicals in different medicinal plants. *The Open Nutraceut. J.* 5: 179 - 186
- Singleton, V.L. and Rossi, J. A. (1965).** Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am. J. Enolo. Viticul.* 16: 144-158
- Sofowara, N. A. (2006).** Medicinal Plants and Traditional Medicine in Africa. (Rep. edn). Spectrum Books Ltd: Ibadan;pp: 150 – 160.
- Spreena, K.P., Poongothai, A., Soundariya, S.V., Srirekha, G., Santhi, R. and Annapoorani, S. (2011).** Evaluation of *in vitro* free radical scavenging efficacy of different organic extracts of *Morinda tinctoria* leaves. *Int. J. Pharm. Pharm. Sci.* 3(Suppl.3): 207 – 209.
- Strack, D. and Wray, V. (1994).** The anthocyanins. In *The Flavonoids: Advances in Research Since 1986.* (J. B. Harborne. Ed). Chapman and Hall, London. pp: 1 – 9.
- Suprava, S., Goutam, G., Debagyoti, D. and Sanghamitra, N. (2013).** Phytochemical investigation and *in vitro* antioxidant activity of an indigenous medicinal plant *Alpinia nigra* B. L. Burtt. *Asian Pac. J. Trop. Biomed.* 3(11): 871 – 876.
- Thaipong, K., Boonprakob, U., Crosby, K., Cisneros-Zevallos, L., and Hawkins Byrne, D. (2006).** Comparison of ABTS, DPPH, FRAP, and ORAC assays for estimating antioxidant activity from guava fruit extracts. *J. Food Comp. Anal.* 19(6): 669-675.
- Uddin, R., Sinha, S., Hossain, A., Kaiser, A., Hossain, K. and Rashid, M.A. (2011).** Chemical and Biological Investigation of *Justicia gendarussa* (Burm. f). *Dhaka Univ. J. Pharm. Sci.* 10(1): 53 – 57.
- Welch, C.R., Wu, Q. and Simon, J.E. (2008).** Recent advances in anthocyanin analysis and characterization. *Curr. Anal. Chem.* 4(2): 75 – 101.