

Ref.  
QH1  
.A47573

Annals  
of the  
Missouri  
Botanical  
Garden  
1997



Volume 84  
Number 3

---

**The Annals**, published quarterly, contains papers, primarily in systematic botany, contributed from the Missouri Botanical Garden, St. Louis. Papers originating outside the Garden will also be accepted. All manuscripts are reviewed by qualified, independent reviewers. Authors should write the Managing Editor for information concerning arrangements for publishing in the ANNALS. Instructions to Authors are printed in the back of the last issue of each volume.

---

### Editorial Committee

Michael H. Grayum

*Editor,*

*Missouri Botanical Garden*

Amy Scheuler McPherson

*Managing Editor,*

*Missouri Botanical Garden*

Diana Gunter

*Editorial Assistant,*

*Missouri Botanical Garden*

Vicki Couture

*Secretary*

Teresa Johnson

*Publications Order Processor*

Ihsan A. Al-Shehbaz

*Missouri Botanical Garden*

Gerrit Davidse

*Missouri Botanical Garden*

Roy E. Gereau

*Missouri Botanical Garden*

Peter Goldblatt

*Missouri Botanical Garden*

Gordon McPherson

*Missouri Botanical Garden*

P. Mick Richardson

*Missouri Botanical Garden*

Henk van der Werff

*Missouri Botanical Garden*

---

For subscription information contact Department Eleven, P.O. Box 299, St. Louis, MO 63166-0299. Subscription price is \$110 per volume U.S., \$115 Canada and Mexico, \$135 all other countries. Four issues per volume. The journal *Novon* is included in the subscription price of the ANNALS.

amcpher@admin.mobot.org (editorial queries)

dept11@mobot.org (orders)

http://www.mobot.org

© Missouri Botanical Garden 1997

The mission of the Missouri Botanical Garden is to discover and share knowledge about plants and their environment, in order to preserve and enrich life.

Ⓜ This paper meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

---

THE ANNALS OF THE MISSOURI BOTANICAL GARDEN (ISSN 0026-6493) is published quarterly by the Missouri Botanical Garden, 2345 Tower Grove Avenue, St. Louis, MO 63110. Periodicals postage paid at St. Louis, MO and additional mailing offices. POSTMASTER: Send address changes to ANNALS OF THE MISSOURI BOTANICAL GARDEN, Department Eleven, P.O. Box 299, St. Louis, MO 63166-0299.

---

Volume 84  
Number 3  
1997

Annals  
of the  
Missouri  
Botanical  
Garden

---

A REVISION OF  
*PHILODENDRON* SUBGENUS  
*PHILODENDRON* (ARACEAE)  
FOR MEXICO AND CENTRAL  
AMERICA<sup>1</sup>

Thomas B. Croat<sup>2</sup>

MISSOURI BOTANICAL

OCT 3 1 1997

GARDEN LIBRARY

---

ABSTRACT

This is the first revision of *Philodendron* subg. *Philodendron* since that of K. Krause in *Das Pflanzenreich* in 1913. *Philodendron* subg. *Philodendron*, the largest of the three subgenera, includes 103 taxa (95 species and 8 subspecies or varieties) for Central America. Sixty-eight taxa are new to science. These include 62 species: *P. albisuccus* Croat, *P. alticola* Croat & Grayum, *P. angustilobum* Croat & Grayum, *P. annulatum* Croat & Grayum, *P. antioianum* Croat, *P. aromaticum* Croat & Grayum, *P. bakeri* Croat & Grayum, *P. breedlovei* Croat, *P. breusterense* Croat, *P. brunneicaule* Croat & Grayum, *P. chiriquense* Croat, *P. chiripoense* Croat & Grayum, *P. clewellii* Croat, *P. coloradense* Croat, *P. copense* Croat, *P. correae* Croat, *P. cotobrusense* Croat & Grayum, *P. colonense* Croat & Grayum, *P. crassispatum* Croat & Grayum, *P. cretotum* Croat & Grayum, *P. dodsonii* Croat & Grayum, *P. dolichophyllum* Croat, *P. dominicalense* Croat & Grayum, *P. duyeri* Croat, *P. odenudatum* Croat, *P. ferrugineum* Croat, *P. findens* Croat & Grayum, *P. foliomii* Croat, *P. fortunei* Croat, *P. gigas* Croat, *P. granulare* Croat, *P. grayumii* Croat, *P. hammelii* Croat, *P. hebetatum* Croat, *P. heleniae* Croat, *P. immixtum* Croat, *P. jefense* Croat, *P. knappiae* Croat, *P. lazarii* Croat, *P. lentii* Croat & Grayum, *P. llanense* Croat, *P. madronense* Croat, *P. malesevichiae* Croat, *P. morii* Croat, *P. niqueanum* Croat, *P. pirrenae* Croat, *P. pseudauriculatum* Croat, *P. parvifolium* Croat, *P. roseospathum* Croat, *P. scalarinense* Croat & Grayum, *P. sousae* Croat, *P. squamicaule* Croat & Grayum, *P. squamipetiolatum* Croat, *P. stramineicaule* Croat, *P. sulcicaule* Croat & Grayum, *P. thalassicum* Croat & Grayum, *P. tysonii* Croat, *P. ubigantapense* Croat, *P. utleyanum* Croat, *P. verapazense* Croat, *P. wilburii* Croat & Grayum, *P. zhuangum*; 6 infraspecific taxa: *P. davidsonii* Croat subsp. *bocatoranum* Croat, *P. hederaceum* (Jacq.) Schott var. *kirkbridei* Croat, *P. ligulatum* Schott var. *heracleionum* Croat, *P. ligulatum* Schott var. *ovatum* Croat, *P. roseospathum* Croat var. *angustilaminatum* Croat, *P. wilburii* var. *longipedunculatum* Croat & Grayum; and 2 new combinations: *P. hederaceum* (Jacq.) Schott var. *oxycardium* (Schott) Croat, *P. radiatum* Schott var. *pseudoradiatum* (Matuda) Croat. Species diversity of *P.* subg. *Philodendron* in Central America shows a general diminution from Mexico to Middle America, followed by a marked increase closer to South America. Mexico has 21 taxa, Guatemala 15, Belize 9, El Salvador 5, Honduras 13, Nicaragua 18, Costa Rica 49, and Panama 80, respectively. Endemism is high, especially for Panama where 39 taxa are currently considered endemic. Mexico and Costa Rica each have 7 endemic species. With the exception of Belize, which has 1 endemic, no other country in Middle America has any endemic species. Only 26 species (28 taxa), a total of 26% of all Central American species, range into South America, 8 species (7% of the total) only to Colombia.

---

<sup>1</sup> This study was completed with support from National Science Foundation grant BSR-8614777.

<sup>2</sup> P. A. Schulze Curator of Botany, Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri 63166-0299, U.S.A.

## INTRODUCTION

*Philodendron* is, after *Anthurium*, the largest genus in the Araceae, with 700 or more species (Croat, 1979, 1983a, 1988, 1990). This is the first major revision of *P. subg. Philodendron* for Central America since Kurt Krause's (1913) generic treatment in *Das Pflanzenreich*. *Philodendron* is one of the most important genera in the Neotropics, inhabiting a wide range of mesic habitats from sea level to over 2000 m and in life zones (Holdridge, 1967) ranging from *Tropical moist forest* to *Premontane rain forest*. While most species occur in virgin humid forests, the genus is known from freshwater swamps, stream banks, regrowth forest, rock outcrops, and roadbanks. It is not only one of the largest genera in the Neotropics but also often constitutes the most conspicuous element of the vegetation because of its abundance, primarily climbing habit, and frequently large, showy leaves. The genus provides a wide variety of choice ornamental plants for horticulture, including most of the species treated here. Unfortunately, it is also still poorly known taxonomically, especially in the South American Andes.

*Philodendron* has 119 Central American species comprising 128 taxa, distributed in two subgenera of *Philodendron*. This Central America revision encompasses only members of *P. subg. Philodendron*, with 103 taxa, including 95 species and 8 varieties or subspecies (Appendix 1, Geographic Distribution of Central American *Philodendron subg. Philodendron*). A total of 68 taxa are new to science: 62 species, 6 subspecies or varieties, and 2 combinations. Alternatively, *P. subg. Pteromischum*, revised separately by Grayum (1996), contains 21 species (26 taxa) for Central America. That revision encompasses all species from Pacific and Caribbean tropical America, ignoring only species from the region of the Guianas and from the Amazon drainage of South America.

*Philodendron* is a distinct genus, not easily confused with any other, though closest to *Homalomena*, which differs in having a consistently terrestrial habit, frequently spiny petioles, a sap smelling of anise, and staminodia among the pistillate flowers.

Species diversity of *P. subg. Philodendron* in Central America shows a general diminution from Mexico to Middle America with the lowest totals just north of the San Juan depression, followed by a marked increase approaching the South American continent. Mexico has 21 taxa, Guatemala 15, Belize 9, El Salvador 5, Honduras 13, Nicaragua 18, Costa Rica 48, and Panama 82. Endemism is high, especially for Panama where 39 taxa are currently

considered endemic. Mexico and Costa Rica each have 7 endemic species. With the exception of Belize, which has 1 endemic, no other country in Middle America has any endemic species.

Most of the Central American species of *P. subg. Philodendron* (Appendix 3, Sectional Composition of *Philodendron subg. Philodendron* in Central America) are in *P. sect. Calostigma* (Schott) Engl. with 52 taxa (48 species) and *P. sect. Philodendron* with 62 taxa (58 species). Other sections represented in Central America are: *P. sect. Tritomophyllum* (Schott) Engl. with 6 species; *P. sect. Polytomium* (Schott) Engl. with 3 species (4 taxa); and *P. sect. Macrogynium* Engl. with 1 species. Sections not represented in Central America are: *Schizophyllum* (Schott) Engl.; *Camptogynium* K. Krause; and *Philopsammis* G. S. Bunting. A key for the sections of *P. subg. Philodendron* is included under "Taxonomy."

## MATERIALS AND METHODS

This revision is based on more than 25 years of field studies in Central and South America, between 1967 and 1993. All but 15 of the 95 species were studied in the field or under cultivation at the Missouri Botanical Garden. Those only known from herbarium material are: *P. breedlovei*, *P. brewsterense*, *P. chirripoense* Croat & Grayum, *P. cotobruense*, *P. dwyeri*, *P. folsomii*, *P. hammelii*, *P. jefense*, *P. madronense*, *P. roseospathum* var. *angustilaminatum*, *P. sousae*, *P. ubigantupense*, *P. utleyanum*, and *P. verapazense* Croat. Except for these, all descriptions have been prepared from both living and dried specimens. The use of ("dried") preceding all or any part of the description is an indication that all that follows is based on herbarium material only. Morphological characters were coded directly into a computerized database to ensure parallel and sortable descriptions. The aroid descriptions database, completely rewritten since the publication of my revision of *Anthurium* sect. *Pachyneurium* (Croat, 1991), contains 892 character states used to describe the morphological diversity expressed in *Philodendron*. A total of 108 of these are used exclusively for description of the bisexual inflorescence (and thus were not used in the descriptions of *Philodendron*), while 220 describe unisexual inflorescences. The database also allows for sorting of characters for use in writing keys or in providing useful lists of characters for preparing a cladistic survey. In addition, the database can be put to future use for the preparation of floristic surveys or for adding additional newly discovered species. The description database is tied directly to the nomen-

clatural database in TROPICOS (Crosby, 1986; Crosby & Magill, 1986). Finally, discussions and references to illustrations as well as exsiccatae are stored separately but tied to a particular species description and to the nomenclatural information by a unique taxon number. Specimens can be added to the exsiccatae almost until the time of publication because they are automatically presorted to localities before being printed. Species descriptions are decoded into narrative text automatically before final editing for style.

Terminology and usage in the descriptions in this revision are largely defined by Croat and Bunting (1979). Further definitions of petiole cross-sectional shapes are defined and illustrated in Croat (1983a). The colors reported in the description frequently are taken from the color chart by Berlin and Kay (1969) and are referenced in the text as B & K. This color chart, available from the University of California Press, is a reproduction of the Munsell Color Array of 40 hues, at maximum saturation, with nine degrees of brightness. This represents 40 hues in the vertical columns and 9 degrees of brightness in the horizontal columns. Colors are arranged in 10 basic clusters with 4 different hues per cluster, ranging from red through yellow, green, blue, purple, and finally red-purple. The four columns for each color cluster are numbered 2.5, 5, 7.5, and 10. These numbers are repeated for each basic color type. The colors from the B & K color chart are read by first reporting the color, then the row followed by the column. For example, the third color in the fifth row is Red 5/7.5. The second color in the eighth row is Red 8/5.

Ecological zones, though sometimes estimated from my own experience with Central American vegetation, are largely taken from Holdridge life-zone maps (Holdridge, 1967; Holdridge et al., 1971), where they exist for Central American countries. Vegetation types for Mexico are taken from the "Mapa de tipos de vegetación de la República de México" (Flores et al., 1971).

Each life zone is represented by a full textual statement and abbreviation which appear on life zone maps. The Holdridge Life Zones of Central America and areas where *Philodendron* occur are listed here, arranged in a generally drier to wetter order: *Tropical thorn woodland* (T-tw); *Tropical dry forest* (T-df); *Tropical moist forest* (T-mf); *Tropical wet forest* (T-wf); *Tropical rain forest* (T-rf); *Premon-tane thorn woodland* (P-tw); *Premon-tane moist forest* (P-mf); *Premon-tane wet forest* (P-wf); *Premon-tane rain forest* (P-rf); *Tropical Lower Montane wet forest*; (TLM-wf); *Tropical Lower Montane rain forest* (TLM-rf); *Tropical wet forest transition to Premon-*

*tane wet forest* (T-wf/P-wf); *Premon-tane wet forest transition to moist forest* (P-wf/ml); and *Premon-tane wet forest transition to rain forest* (P-wf/rf).

Herbarium material has been widely distributed, and original field vouchers are cited for all herbaria whose material was seen. Herbarium material may consist of one of three kinds: (1) complete original sets (wild collected); (2) sterile original material with an inflorescence added from a cultivated plant of the same number; and (3) material collected entirely from cultivated plants. Specimens based entirely or in part on cultivated material are clearly indicated on the herbarium label.

Herbarium specimens were borrowed from most major herbaria including: AAU, B, BBS, BISH, BM, BR, CAS, CAY, CM, COL, CR, DAV, DUKE, DS, EAP, ECON, ENCB, F, FLAS, FSU, FTG, G, GH, HBG, ISC, K, L, LA, LL, LE, M, MEXU, MICH, NY, PMA, RSA, S, SCZ, SEL, TEX, U, UC, UMO, US, VEN, and WIS.

Descriptions are mostly parallel and as complete as possible. Descriptions of the pistils, vitally important in the infrageneric classification of *Philodendron*, are particularly detailed. In order to avoid repetition, description references are made to Style Types discussed by Mayo (1986). These style types are discussed and illustrated (Fig. 469) in the introduction under the section on "Morphology of Reproductive Structures—Gynoecium."

Intraspecific categories in this work adhere to the following definitions. Subspecies are those infraspecific taxa that are mostly or entirely allopatric either geographically, elevationally, or ecologically. Varieties, though morphologically distinct in one or several characters, are apparently geographically, elevationally, or ecologically sympatric. All infraspecific taxa within Central American *Philodendron* subg. *Philodendron* are ecologically sympatric.

#### ACKNOWLEDGMENTS

This revision could not have been completed without the help of many people, most important among them my dedicated and able co-worker, Petra Malesevic, who participated in nearly every phase of the work. I could count on her to continue the research unabated during my long field trips. Along with Bob Magill, she revised, updated, and improved the more powerful description program that we used for the *Philodendron* revision. Jo Ann Beiser played an important role for nearly two years, providing assistance in the preparation of descriptions. Kathy Pickett Upton, former Research Greenhouse manager, played a major role in maintaining the live collections, making pollinations,

and in dealing with label problems. Very special thanks also go to my colleague and fellow aroid specialist Mike Grayum, whose knowledge of the Costa Rican Araceae is without parallel. He has not only made many of the best collections of *Philodendron*, but his breadth of experience in that part of Central America and his vivid insights into the interrelationships of species were very rewarding. In addition, his skills as an editor and his knowledge of Latin and rules of nomenclature contributed much to this work. Simon Mayo, like Mike Grayum, conducted extensive work on *Philodendron*. Their work was concurrent or preceded my own, and their ideas and interpretations of phenomena in the genus were ever insightful. My own work has benefited greatly from their work. Dan Nicolson was always generous with his time in helping to solve nomenclatural problems as well as problems with translations of difficult Latin phraseology. Gordon McPherson provided pickled material of many species of *Philodendron* during his tenure as our resident botanist in Panama. Eleanor Sauer has proof-read or written Latin diagnoses for nearly all of the new species. Joseph Tosi, of the Tropical Science Center in San José, Costa Rica, assisted us in standardizing our use of the Holdridge Lifezone System, which we have added to our computerized database. Leland Russell, summer volunteer and later summer employee while vacationing from his undergraduate studies at Carleton College in Minnesota, became very skillful at dissections of *Philodendron* inflorescences. His critical comparative study of the Central American species of *Philodendron* went a long way in increasing our understanding of the differences within the genus. Finally, I would like to thank my wife, Patricia, who has without complaint allowed me to spend several months each year wandering the tropics of Central and South America for almost three decades. She was always available to solve every computer glitch and has also helped with the computerized sorts of information which led to the construction of the keys.

#### HISTORY OF THE GENUS *PHILODENDRON*

Although *Philodendron* apparently figured in pre-Columbian folklore, art, and medicine during the 16th century, and herbarium material was collected by Georg Marcgraf as early as 1644 (Mayo, 1990), it was Charles Plumier who made the first effective introduction of the genus to European scientists (Mayo, 1990). He collected five or six species from Martinique, St. Thomas, and Hispaniola, giving phrase names beginning with "Arum" or "Dracunculus." See Mayo (1990) for a detailed ac-

count of *Philodendron* collections made by other late 17th-century and 18th-century explorers, including Hans Sloane in Jamaica and N. J. Jacquin in the West Indies, Colombia, and Venezuela.

Plumier's expeditions resulted in the first species of *Philodendron* to be published as new to science, albeit as an *Arum*, *A. lingulatum* L., a member of *Philodendron* subg. *Pteromischum*. His trips also resulted in *Philodendron hederaceum*, the first member of *P.* subg. *Philodendron* to be published new to science, as *Arum hederaceum* Jacq.

#### HEINRICH WILHELM SCHOTT

One hundred thirty-six years passed from the time Plumier first introduced plants of what came to be known as *Philodendron* to European scientists in 1693 and the first circumscription of the genus by Schott in 1829. During this interim a number of workers, including Carl Linnaeus, worked with generic concepts and decided that not all aroids belonged in the same genus. *Arum* came to be used only for the European plants it now represents and *Arisaema*, *Dracunculus*, and *Colocasia* were also separated by the early 18th century (Hermann, 1698; Tournefort, 1700). By the fifth edition of *Genera Plantarum* Linnaeus (1754) had recognized also *Calla*, *Dracontium*, and *Pothos* as well as *Orontium*, *Pistia*, and *Acorus*, although not recognizing the last three as related to *Arum*.

By the middle of the 18th century the exploration of the Neotropics was well under way. The introduction of so many new plants from the New World tropics, including many *Philodendron* species, made it clear that further separation was needed. Because of the uncritical acceptance of many aberrant elements into *Arum* during the course of the 18th century a number of taxa now recognized as *Caladium*, *Philodendron*, and *Syngonium* were incorporated into *Arum*. Ventenat (1800) solved part of the problem by recognizing *Caladium*, but Willdenow (1805) erred in transferring into *Caladium* four of Jacquin's West Indian *Philodendron* species (all considered *Arum* at the time).

Even by the time of the 16th edition of *Systema Vegetabilium* Sprengel (1826) did not distinguish *Philodendron* but had increased the number of genera to 12, adding *Ambrosina*, *Arisarum*, *Caladium*, *Gymnostochys*, and *Zantedeschia*.

Resolution of many of the remaining generic problems with the Araceae awaited Austrian botanist Heinrich Wilhelm Schott, who was the first to focus on the taxonomy of the Araceae (Nicolson, 1960). Schott was uniquely qualified in this regard, having spent four years in Brazil collecting plants

and with access to the large collection of living plants at the Imperial Gardens [of the Hapsburg's] at Schönbrunn Palace in Vienna where he worked as the director. These included his own Brazilian collections as well as those of N. J. Jacquin from the Caribbean. In 1829 Schott described the genus *Philodendron* (published as *Philodendrum*) Schott in one of his first publications after returning from Brazil in 1821. The first species placed in the genus were: *P. grandifolium* (Jacq.) Schott, *P. lacinosum* Schott, *P. tripartitum* (Jacq.) Schott, *P. lacrerum* (Jacq.) Schott, *P. pinnatifidum* (Jacq.) Schott, and *P. hederaceum* (Jacq.) Schott. In 1832 Schott published a preliminary classification of the Araceae in a work entitled *Meletemata Botanica*, which included many plant families and was done in conjunction with his colleague S. Endlicher. In this work he recognized 39 genera of Araceae including *Philodendron* and some other by now well known species-rich genera such as *Anthurium*, *Syngonium*, *Dieffenbachia*, *Aglaonema*, and *Spathiphyllum*. In this work he also published the first infrageneric system of classification for *Philodendron*, although it contained only three groups still recognized today. Though Schott subsequently continued to work on his circumscription of the genera and species of the Araceae for the remainder of his career, he did not publish any major revisions of Araceae for 24 years.

Endlicher (1837), publishing alone but with the obvious assistance of Schott (Mayo, 1990), produced a revision of *Philodendron* that differed from the *Meletemata Botanica* account by Schott only in having a more complete generic description, including vegetative details which were presented for the first time.

Shortly after Endlicher's revision, K. S. Kunth (1841) published in his "Enumeratio Plantarum. . ." the first species-level revision of *Philodendron*. This revision included new species and new combinations and for the first time included species of the genus that would later be recognized as members of *P. subg. Pteromischum*.

It was not until Schott's (1856) publication of a work entitled *Synopsis Aroidearum* that a fully developed infrageneric system of classification was realized. This was Schott's first species-level revision of the genus, and the *Synopsis* divided the genus into 22 "greges" grouped in 7 unnamed categories arranged in increasing order of complexity of leaf blade shape (Mayo, 1990). A total of 99 species (which was recognized here for the first time as grex *Pteromischum*) and 3 species of *P. subg. Meconostigma* were included (two of them reported as members of grex *Sphincterostigma*).

Schott's final classification of *Philodendron* was published four years later in the *Prodromus Systematis Aroidearum* (1860), a more rigorous work that came to be his last comprehensive self-published work, since he died at the age of 71 in 1865. This revision included 110 genera; almost all are still recognized as genera or subgenera. The fact that his work has stood the test of time is a testimony to the serious nature of the research he had done in Vienna during his long period of seeming inactivity.

Schott's treatment of *Philodendron* in the *Prodromus* differed from his revision in the *Synopsis* in having more complete descriptions, using more inflorescence characters; in having six rather than seven unnamed categories of leaf shape to group the "greges"; and in beginning to make use of the cataphyll (prophyll in the strictest modern sense) as a character. Schott used the term "subopposite stipule" for the feature that came to be known as the cataphyll in Engler's usage. This has proven to be one of the best and most reliable characters for the genus. For a detailed listing of the key characters for *Philodendron* used in Schott's *Prodromus* revision, refer to Mayo (1990: 50).

The *Prodromus* treatment included 135 species of *Philodendron* included in the same "greges." Twenty species of the total were members of *P. subg. Pteromischum* Schott and 6 species are now placed in *P. subg. Meconostigma* (Schott) Engl. (3 in grex *Meconostigma* and 3 in grex *Sphincterostigma*). Thus Schott included 110 names of *P. sect. Philodendron*. Taking synonymy into account, only 76 species of *P. sect. Philodendron* were included in this 1860 revision. Only 18 names pertained to Central America. Of these, only 8 are accepted members of *P. sect. Pteromischum*.

The species included by Schott (1860) in each grex of the *Prodromus* for Central America are listed below:

- Grex *Boursia* Rchb. ex Schott: *P. wendlandii* Schott
- Grex *Pteromischum* Schott: *P. aurantiifolium* Schott, *P. ligulatum* (L.) K. Koch, *P. seguine* Schott, *P. inaequilaterum* Liebm.
- Grex *Canniphyllum* Schott: (no species represented)
- Grex *Glossophyllum* Schott: (no species represented)
- Grex *Solenostergma* Klotzsch: *P. oxycardium* Schott, *P. micans* K. Koch, *P. scandens* K. Koch & Sello = (*P. hederaceum*)
- Grex *Psoropodium* Schott: *P. tenue* K. Koch & Augustin, *P. gracile* Schott

- Grex Achyropodium* Schott: *P. verrucosum* L. Mathieu ex Schott
- Grex Platypodium* Schott: *P. pterotum* K. Koch & Augustin, *P. fragrantissimum* (Hook.) Kunth
- Grex Cardiophylacium* Schott: *P. brevispatum* Schott, *P. hederaceum* Schott (= *P. jacquinii* Schott)
- Grex Belocardium* Schott: *P. hoffmannii* Schott, *P. advena* Schott, *P. acrocardium* Schott
- Grex Cardiobellium* Schott: (no species represented)
- Grex Meconostigma* Schott: (no species represented)
- Grex Eubelium* Schott: (no species represented)
- Grex Macrolonchium* Schott: (no species represented)
- Grex Macrobilium* Schott: *P. daemonum* Liebm. = *P. sagittifolium* Liebm., *P. sagittifolium* Schott, *P. tanyphyllum* Schott = *P. sagittifolium* Liebm.
- Grex Imbéa*: (no species represented)
- Grex Oligophlebium* Poepp.: (no species represented)
- Grex Doratophyllum* Schott: (no species represented)
- Grex Schizophyllum* Schott: (no species represented)
- Grex Tritomophyllum* Schott: *P. tripartitum* (Jacq.) Schott, *P. dagilla* Schott = *P. tripartitum* (Jacq.) Schott, *P. anisotomum* Schott
- Grex Polytomium* Schott: *P. subincisum* Schott, *P. impositum* Schott = *P. radiatum* Schott, *P. polytomum* Schott = *P. radiatum* Schott, *P. warszewiczii* K. Koch & Bouché
- Grex Sphincterostigma* Schott: (no species represented)

In all, only 18 of the 135 species of *Philodendron* included in this 1860 revision were from Central America, and 4 of the total were members of *P. grex Pteromischum*. One of the four species, *P. ligulatum* Schott, is a member of *P. sect. Philodendron*, which Schott had inadvertently placed in his *grex Pteromischum*. Five of Schott's names became *P. hederaceum* (*P. acrocardium*, *P. hoffmannii*, *P. oxycardium*, *P. scandens*, and *P. micans*), but the count was reduced by only four since the name *P. hederaceum* was involved in two currently accepted species (namely, *P. hederaceum* and *P. jacquinii*). Other reductions are *P. gracile*, which is a synonym of *P. tenue*; *P. daemonum* and *P. tanyphyllum*, which are synonyms of *P. sagittifolium*; *P. dagilla*, a synonym of *P. tripartitum*; and *P. impositum* and *P. polytomum*, synonyms of *P. radiatum*. Thus

Schott was dealing with only 16 (16.5%) of the 96 currently known Central American species.

## ADOLF ENGLER

Little was done with *Philodendron* following Schott's death in 1865 until Adolf Engler, working at the Universities of Kiel and Breslau (and finally at the Berlin Botanical Garden), began his revisionary work on the Araceae. Schott had laid the groundwork, describing most of the genera that still exist today, but he was dealing with only a small portion of the species. Taking up his first positions at Kiel and Breslau in 1871, at age 27, Engler worked with various tropical families on the *Flora Brasiliensis* project, publishing and working on a general review of the vegetative and floral morphology for the entire family (Engler, 1877). Later, in his powerful position as Director of the Berlin Botanical Garden, Engler commanded attention and a wealth of herbarium specimens and living material from all over the world during Germany's preeminent period of domination in the botanical world preceding World War II. Since Engler was only 21 at the time of Schott's death, it is not likely that the two ever met, but Engler would have had access to some of the same material, including living material from the Schönbrunn greenhouses, as well as Schott's notes and illustrations made over a 40-year period (Engler, 1876). Unlike modern workers, he had access to Schott's herbarium in Vienna before it was destroyed during World War II. This is important for a continuum of species concepts in groups often described from inadequate material of unknown origin and worse yet sometimes destroyed by war.

Engler's (1878) treatment of the Araceae in Martius's *Flora Brasiliensis* included 116 species of *Philodendron*, 95 of which were members of *P. sect. Philodendron* (13 were in *P. sect. Pteromischum* and 8 in *P. sect. Meconostigma* (Schott) Engl.). Only 47 species of *Philodendron* were then known from Brazil. The work also included sectional and species descriptions as well as a key to all existing species of *Philodendron*. In this work Engler modified Schott's system of classification for *Philodendron* by recasting Schott's "greges" as sections and reducing the number from 22 to 10. He also synonymized a number of species, reducing the total from 132 to 116 species. Only a single Central American species, *P. oxycardium* (= *P. hederaceum* var. *oxycardium*) was reported for Brazil, a fairly accurate picture as we know today. Only a couple of other species, namely *P. fragrantissimum* and *P. verrucosum*, have been found to range into the Am-



azon drainage of South America. Two additional species, *P. glanduliferum* Matuda and *P. brevispathum*, have a subspecies or variety that occurs in the Amazon basin of South America, but the same do not occur in Central America.

In the following year Engler's treatment of *Philodendron* for A. and C. DeCandolle's *Monographie Phanerogamarum* (Engler, 1879) was essentially unchanged, adding only 3 species to bring the total to 120. Of these, 93 species were members of *P. subg. Philodendron* and 20 of the epithets represented species currently known from Central America (now reduced to 15 through synonymy).

The final revisionary effort by Engler on *Philodendron* published 20 years later (Engler, 1899) was changed only slightly at the subgeneric level from the 1878 work. One section was raised to subgeneric status, and the remaining nine sections were included in *P. subg. Euphilodendron* Eng. (now *P. subg. Philodendron*). Despite minor changes made by Krause (1913), it is essentially Engler's classification that persists almost a century later. Engler's (1899) revision was substantially larger than Schott's last revision. Engler's revision contained 167 species, 134 of them in *P. subg. Philodendron*, with 23 reported for Central America (reduced to 15 species through synonymy in this revision). One species, *P. purpureoviride* Eng., reported for Ecuador, is now known for Central America.

The species of *P. subg. "Euphilodendron"* included in Engler's (1899) revision broken down by section for Central America are listed below:

*P. sect. Pteromischum* Schott: *P. aurantiifolium* Schott (as synonym of *P. guttiferum* Kunth), *P. guatemalense* Eng., *P. inaequilaterum*, *P. seguine*, *P. talamancae* Eng.

*P. sect. Boursia* (Rchb. ex Schott) Eng.: *P. wendlandii* Schott

*P. sect. Polyspermium* Eng.

"Gruppe" *Platypodium* Schott: *P. pterotum*

"Gruppe" *Solenostegima* Klotzsch: *P. oxycardium* (= *P. hederaceum* var. *oxycardium*), *P. purpureoviride* (as *P. purpureoviridis* from South America), *P. micans* K. Koch (= *P. hederaceum* var. *hederaceum* forma *micans*), *P. scandens* (= *P. hederaceum* var. *hederaceum*)

"Gruppe" *Cardiobelium* Schott: *P. brevispathum*, *P. gracile* (= *P. tenue*), *P. schottianum*, *P. tenue*

"Gruppe" *Achyropodium* Schott: *P. verrucosum*

*P. sect. Oligospermium* Eng.

"Gruppe" *Macrobilium* Schott: *P. sagittifolium*, *P. daemonum*, *P. sanguineum* Regel (= *P. sagittifolium*), *P. mexicanum* Eng.

"Gruppe" *Belocardium* Schott: *P. ligulatum* Schott, *P. advena*, *P. subovatum* Schott (= *P. advena*), *P. smithii* Eng.

"Gruppe" *Oligocarpidium* Eng.: *P. pittieri* Eng. (= *P. hederaceum*)

*P. sect. Tritomophyllum* (Schott) Eng.: *P. anisotomum*, *P. tripartitum*, *P. fenzlii* Eng. (= *P. tripartitum*)

*P. sect. Schizophyllum* (Schott) Eng.: no species represented

*P. sect. Polytomium* (Schott) Eng.: *P. augustinum* K. Koch (= *P. radiatum*), *P. radiatum* Schott, *P. warszewiczii* K. Koch & Bouché

*P. sect. Macrolonchium* (Schott) Eng.: *P. fragrantissimum*

*P. sect. Macrogynium* Eng.: *P. hoffmannii* Schott sensu Eng. (= *P. jacquinii* Schott)

The turn of the century saw major activity with *Philodendron*, no doubt due to Engler's just published revision. Engler made no changes in his revision but went on to publish 26 additional species (Engler, 1905b). In addition, seven species were described by Alfred Barton Rendle, Ignaz Urban, Ambroise Gentil, and N. E. Brown between 1901 and 1908.

#### KURT KRAUSE

Kurt Krause, who began working with Engler at the Berlin Botanical Garden on 1 January 1905, described two additional species before preparing his revision of *Philodendron* for *Das Pflanzenreich* (Krause, 1913). The latter remains the most recent revision of the whole genus. Krause's revision is a slightly reworked version of Engler's (1899) revision but did include the description of a new section, *P. sect. Campogynium*, with a single species in *P. subg. Philodendron* ("*Euphilodendron*") and included 55 more species. There were 32 other new species published in *P. subg. Philodendron*. Six of these were in *P. sect. Pteromischum* (*P. subg. Pteromischum*), while one was in *P. subg. Meconostigma*. The remaining 25 were in *P. subg. Philodendron*. Most were members of *P. sect. Boursia* and *P. sect. Polyspermium* (*Philodendron*) with a single species each in the following sections: *Oligospermium* (*Calostigma*), *Schizoplacium*, *Macrolonchium*; and three species in *P. sect. Polytomium*. Only two species, *P. grandipes* K. Krause and *P. panamense* K. Krause (both in current *P. sect. Philodendron*), were from Central America. Krause's treatment of *P. subg. Philodendron* included the following sections and species for Central America (or at least now known from Central America):

Names used by Krause (1913) and their current status. Species numbers refer to those in Krause's revision.

2. *P.* sect. *Baurisia* (Rehb. ex Schott) Engl.
39. *P. wendlandii* Schott
3. *P.* sect. *Polyspermium* Engl.
67. *P. coerulescens* Engl. (member of *P.* subg. *Pteromischum* mistakenly placed here)
70. *P. pterotum* K. Koch & Augustin
77. *P. grandipes* K. Krause
90. *P. purpureoviride* Engl. (reported for Ecuador)
95. *P. scandens* K. Koch & Sello = *P. hederaceum*
96. *P. oxycardium* Schott = *P. hederaceum* var. *oxycardium*
99. *P. micans* K. Koch = *P. hederaceum* var. *hederaceum* forma *micans*
108. *P. schottianum* H. Wendl. ex Schott
110. *P. panamense* K. Krause
117. *P. gracile* Schott = *P. tenue* K. Koch & Augustin
116. *P. tenue* K. Koch & Augustin
125. *P. verrucosum* L. Mathieu ex Schott
4. *P.* sect. *Oligospermium* Engl.
134. *P. sagittifolium* Liebm.
140. *P. sanguineum* Regel = *P. sagittifolium* Liebm.
147. *P. mexicanum* Engl.
150. *P. ligulatum* Schott
164. *P. subrotatum* Schott = *P. sagittifolium* Liebm.
168. *P. smithii* Engl.
173. *P. advena* Schott
176. *P. pittieri* Engl. = *P. hederaceum* (Jacq.) Schott
5. *P.* sect. *Tritomophyllum* (Schott) Engl.:
181. *P. fendii* Engl. = *P. tripartitum* (Jacq.) Schott
182. *P. anisotomum* Schott
183. *P. tripartitum* (Jacq.) Schott
6. *P.* sect. *Schizophyllum* (Schott) Engl.: (no species represented in Central America).<sup>3</sup>
7. *P.* sect. *Polytomium* (Schott) Engl.:
195. *P. radiatum* Schott
198. *P. augustinum* K. Koch = *P. radiatum* Schott
199. *P. polytomum* Schott = *P. radiatum* Schott
200. *P. warszewiczii* K. Koch & Bouché
8. *P.* sect. *Macrolonchium* (Schott) Engl.

202. *P. fragrantissimum* (Hook.) Kunth

9. *P.* sect. *Macrogynium* Engl.

207. *P. hoffmannii* Schott sensu Engl. = *P. jacquinii* Schott

10. *P.* sect. *Campitogynium* K. Krause: (no species represented in Mesoamerica)

The 28 Central American species names for *Philodendron* subg. *Philodendron* represented only 11% of the 222 species in Krause's revision. Of the 28 taxa Krause reported for Central America, *P. coerulescens* was a member of *P.* subg. *Pteromischum*, 10 are now considered synonyms of other Central American species which he also reported or which have been subsumed into other species as varieties or subspecies. Of the ten, two, *P. tenue* and *P. jacquinii*, have been given older names that he did not report but still represent Central American species. Thus, Krause's revision of *Philodendron* of Central America treated only 19 species, roughly 1/3 of the total currently known for Central America. One additional Central American species, *P. purpureoviride*, was treated by Krause but not reported for Central America.

#### MODERN WORK

Since no revisionary work took place on *Philodendron* between the time of Krause and the advent of *Philodendron* specialist George Bunting's (1963a) revisionary work, most species were described as the result of floristic work and most of these were from South America. In all, 142 new species epithets for *Philodendron* have been introduced to date since the time of Krause's revision (including 5 species published by Grayum, 1992b). In addition, 11 new varieties, 7 new subspecies, and 1 new form were published during this period. Six new combinations were made. Of this total only 33 species of *Philodendron* were reported for Central America, and 11 of these were in *P.* subg. *Pteromischum*. Thus 22 species of *P.* subg. *Philodendron* were published between the last full revision of the genus by Krause and the completion of the current work (not including those species of *P.* subg. *Pteromischum* in Grayum (1996)). The Mesoamerican taxa of *P.* subg. *Philodendron* described between the time of Krause's revision and the completion of this work are as follows: *Philodendron apocarpum* Matuda (= *P. jacquinii* Schott), *P. auriculatum* Standl. & L. O. Williams, *P. basii* Matuda, *P. brenesii* Standl., *P. davidsonii* Croat, *P. dressleri* G. S. Bunting, *P. erlanstonii* I. M. Johnston (= *P. jacquinii*), *P. glanduliferum* Matuda, *P. harlowii* I. M. Johnston, = *P. hederaceum*, *P. jamapanum* G. S. Bunting (= *P. sagittifolium* Liebm.), *P. jo-*

<sup>3</sup> *Philodendron pedatum* (Hook.) Kunth was reported by John Hall of Costa Rica but no live or dried material is available for confirmation.

*davisianum* G. S. Bunting, *P. lancigerum* Standl. & L. O. Williams (= *P. sagittifolium*), *P. latisagittatum* Matuda (= *P. mexicanum*), *P. lundellii* Bartlett ex Lundell (= *P. jacquinii*), *P. microstictum* Standl. & L. O. Williams, *P. miduhoi* Matuda = *P. hederaceum*, *P. mirificum* Standl. & L. O. Williams (= *P. pterotum* K. Koch & Augustin), *P. monticola* Matuda = *P. advena*, *P. platypetiolatum* Madison, *P. pleistoneurum* Standl. & L. O. Williams (= *P. grandipes* K. Krause), *P. pseudoradiatum* Matuda (= *P. radiatum* var. *pseudoradiatum* (Matuda) Croat), and *P. trisectum* Standl. (= *P. anisotomum*).

Three additional species were described in *Philodendron* but actually pertain to other genera (see under "Excluded Names").

Many of the other taxa in Krause's revision were synonymized or reduced in rank, and only 11 taxa in *P. subg. Philodendron* remained. These were: *P. brenesii*, *P. radiatum* var. *pseudoradiatum*, *P. auriculatum*, *P. microstictum*, *P. basii*, *P. glanduliferum*, *P. dressleri*, *P. jodavisianum*, *P. platypetiolatum*, *P. davidsonii*, and *P. strictum*.

Most of the Central American floristic work outside of Mexico was carried out by Paul C. Standley, often working with his associate Louis O. Williams. Standley worked initially at the Smithsonian and later at the Field Museum in Chicago, then at the herbarium of the Escuela Agrícola Panamericana, where he died at Zamorano in Honduras. Standley described *P. brenesii* and *P. trisectum* (= *P. anisotomum* Schott) alone, and with L. O. Williams he also provided the following epithets: *P. armigerum* (= *Syngonium armigerum* (Standl. & L. O. Williams) Croat), *P. auriculatum*, *P. brevinodum* (= *Monstera tuberculata* Lundell var. *brevinoda* (Standl. & L. O. Williams) Madison), *P. hastiferum* (= *Syngonium hastiferum*), *P. lancigerum* (= *P. sagittifolium*), *P. microstictum*, *P. mirificum* (= *P. pterotum*), and *P. pleistoneurum* (= *P. sagittifolium*). It is unusual that despite the fact that there were many undescribed species of *Philodendron* subg. *Philodendron* in Costa Rica and Panama, these workers did not succeed in describing many of them since of the nine described, three proved to belong to other genera, and three others proved to be synonyms of existing *Philodendron* names. This is particularly surprising since both Standley and L. O. Williams were astute observers who were very familiar with the Central American flora in general. Their mistakes point out the complexity of the taxonomy of Araceae and the bewildering array of material available to them at that time. Even when I began my own work with the Araceae in the late 1960s there were few specimens (aside from types) that proved to have the correct names. Specimens

were often poorly preserved and inadequate, and virtually all lacked good field notes. Either field observations or well-prepared field notes describing aspects of the plants not available on dried plants are essential to the proper understanding of *Philodendron* and other Araceae. I believe that it is possible that this confusion, coupled with the dearth of well-prepared specimens and the paucity of types of Araceae, discouraged workers from making decisions about which species were new.

Perhaps owing to the difficulty with the genus, local floras in Central America were often inadequately done, even if one considers that fewer specimens were available to the authors than today. The treatment of the *Flora of Panama* (Standley, 1944), for example, is woefully inadequate considering the small percentage of the total aroid flora that is covered compared to what is now known to exist.

In his treatment of some genera Standley (1944) seemed too willing to accept epithets of species described in Colombia, regardless of how well they "fit" Panamanian species. As a result many species names in genera such as *Anthurium*, for example, were wrong. His treatment of *Philodendron* subg. *Philodendron* was much better from the standpoint of correct names, but he treated only 8 of the 104 taxa (7%) of *Philodendron* subg. *Philodendron* now known for Panama. Only one species, *P. hoffmannii* (now *P. jacquinii*), had the wrong name. *Philodendron jacquinii* was also misapplied, being intended by Standley for *P. hederaceum*. The other species he included in the *Flora of Panama* were: *P. brenesii*, *P. brevispatum*, *P. grandipes*, *P. panamense*, *P. radiatum*, *P. tripartitum*, and *P. wendlandii*. He did not do so well with members of *P. subg. Pteromischum*, where *P. karstenianum* Schott was a mixture of two species and *P. guttiferum* was a mixture of three species. Perhaps the most curious thing about Standley's *Flora of Panama* treatment is that by 1944, after Robert Woodson and his collaborators had already made several expeditions to Panama, so few of the new species included in the present revision had been collected. Standley had collected some of the new species but failed to recognize them as new. These included: *P. crassispatum* (Standley & Valerio 51910), *P. findens* (Standley & Torres 52355), *P. purulhense* (Standley 89902), *P. strictum* (Standley 51371), *P. verapazense* (Standley 91978), and *P. wilburii* var. *wilburii* (Standley 38300). See discussion of those species for additional details. See also section on "Collecting History of *P. subg. Philodendron*."

The *Flora of Guatemala* (Standley & Steyermark, 1958b) was much more accurate and complete in the percentage of the total taxa of *P. subg. Philo-*

*dendron* included. However, this probably has less to do with the fact that it was published 14 years later than it does with the fact that there are fewer, generally more widespread species occurring there than in Panama. The *Flora of Guatemala* treated 11 species of *Philodendron*, 8 of them members of *P. subg. Philodendron*. These were: *P. anisotomum*, *P. hederaceum*, *P. hoffmannii* (= *P. jacquinii*), *P. radiatum*, *P. sagittifolium*, *P. smithii*, *P. tripartitum*, and *P. warszewiczii*. With 50% of the present total taxa included and with all but one of the taxa still properly bearing the name, the Guatemala treatment remains generally more useful than that of the *Flora of Panama*. Added to the flora since the 1958 revision are: *P. advena*, *P. fragrantissimum*, *P. glanduliferum*, *P. jodavisanum*, *P. mexicanum*, *P. purulhense*, and *P. verapazense*.

The *Flora of Guatemala* (Standley & Steyermark, 1958b) treated *P. hederaceum* correctly but treated *P. jacquinii* under its synonym, *P. hoffmannii*. Curiously, however, the illustration used represents both species. The leaf and stem seem clearly to be *P. hederaceum*, but the inflorescence clearly shows the long-protruded styles of *P. jacquinii*.

Standley's (1937) treatment of the Araceae of Costa Rica was reasonably good, partly because many species had been described by Schott from collections made by H. Wendland in Costa Rica. Other widespread species whose taxonomy had been well established were also a part of the flora. Properly named Costa Rican species recognized by Standley were: *P. brenesii*, *P. ligulatum*, *P. pterotum*, *P. radiatum*, *P. schottianum*, *P. tripartitum*, *P. verrucosum*, and *P. wendlandii*. Species now synonymized are: *P. gracile* (= *P. tenue*), *P. hoffmannii* (= *P. jacquinii*), *P. pittieri* (= *P. hederaceum*), and *P. trisectum* Standl. (= *P. anisotomum*). It has never been determined which species he included under the name *P. panamense*, but that species is not known for Costa Rica. Thus, with 8 out of the 13 names correct and 3 additional species that at least represent synonyms of currently recognized species, Standley did pretty well. However, with only 13 of the current 49 species treated (26%), the treatment was no more complete than that of the *Flora of Panama*, which was written a few years later.

Though no other floristic taxonomist had such a prominent role with Central American *Philodendron* as Standley, there were others who described *Philodendron* during the course of their floristic work.

Ivan M. Johnston of the Arnold Arboretum described *Philodendron erlansonii* (= *P. jacquinii*) and *P. harlowii* (= *P. hederaceum*) while working

on the flora of San José Island (Johnston, 1949) of Panama.

In Mexico, Eizi Matuda, the local aroid specialist, described seven species (Matuda, 1954): *P. apocarpum* (= *P. jacquinii*), *P. basii*, *P. glanduliferum*, *P. latisagittatum* (= *P. mexicanum*), *P. miduhoi* (= *P. hederaceum*), *P. monticola* (= *P. advena*), and *P. pseudoradiatum* (= *P. radiatum* var. *pseudoradiatum*); and George Bunting described four species during his investigations of Mexican Araceae (Bunting, 1965): *P. dressleri*, *P. jamapanum* (= *P. advena*), *P. jodavisanum*, and *P. tuxtlanum* G. S. Bunting (= *P. sagittifolium*).

Matuda's treatment of the *Philodendron* in Mexico (Matuda, 1954) dealt with 16 species, 13 of them in *P. subg. Philodendron*. While a number of the species had the correct names, e.g., *P. advena*, *P. mexicanum*, *P. pseudoradiatum*, *P. radiatum*, and *P. tripartitum*, others had synonymous names, e.g., *P. sanguineum* and *P. daemonium* (both currently recognized as *P. sagittifolium*). Matuda also recognized *P. mexicanum* under three different names, namely *P. mexicanum*, *P. latisagittatum*, and *P. sagittifolium* (a distinct species that he treated as both *P. daemonium* and *P. sanguineum* (a currently recognized name improperly used; see above). Matuda redescribed two species using the names *P. apocarpum* and *P. miduhoi* (currently *P. jacquinii* and *P. hederaceum*, respectively). Finally, he described *P. monticola* (now considered synonymous with *P. advena*).

Other Central American species of *P. subg. Philodendron* published since the time of Krause's revision and prior to the beginning of this project in 1986 are *P. davidsonii* and *P. platypetalatum*, the latter described from Ecuador during Mike Madison's extensive work with the flora of Ecuador during his tenure at the Selby Botanical Garden. Thus, up to the initiation of the current study, only 26 species in *P. subg. Philodendron* were described for Central America. With 96 species of *P. subg. Philodendron* now known for Central America, this registers an increase of 70 species or a 270% increase.

Of the species of *P. subg. Philodendron* described since Krause's revision (not including the present effort) only seven were described by non-specialists (I. M. Johnston, P. C. Standley, and Standley and L. O. Williams). Standley, together with L. O. Williams, described most of the new Central American species and wrote most of the floristic treatments of the Araceae for Central America. Despite his extensive experience, I do not believe that Standley fully understood the diversity of the Araceae. This is because he failed to recognize a rel-

atively large number of species that were new to science. In Standley's defense it must be stated that the Araceae are a particularly complex family with so much interspecific variation that proper decisions often cannot be easily made without direct comparisons of living material. In addition, collections in the past often had few or no field notes to use for study.

#### COLLECTING HISTORY

In addition to the history of revisionary efforts it is instructive to consider the collecting efforts in Central America that have laid the groundwork for the scientific work done with *Philodendron* subg. *Philodendron*. Perhaps owing to their sometimes intimidating size, the difficulty of retrieval, and certainly due to the difficulty of preparing and drying specimens, members of *P.* subg. *Philodendron* have never been popular with botanical collectors. The early neotropical collectors, i.e., Ruiz and Pavon, Sessé and Moçiffo, Triana and Planchon, and others collected few Araceae (or at least few survived to modern times). Eduard Poeppig, working in Peru, and Richard Spruce, working in the Amazon region of Brazil, did somewhat better, collecting a number of new species. In Central America few collectors made many collections until modern times. Even Standley and Julian A. Steyermark, two of the most prodigious collectors in the region, did not make many collections of Araceae. Both gathered well over 100,000 collections in their careers. Yet in all, Standley made only 146 collections of *P.* subg. *Philodendron*, comprising 21 species in all of Central America. Of this total only 6 of the collections (these previously mentioned) proved to be undescribed species, none of which Standley recognized as new. Standley was not avoiding collecting *Philodendron* because they constituted a lot of work to press. This is demonstrated by the fact that he collected *P. radiatum* 13 times and *P. warszewiczii* 12 times. These are among the most difficult plants to prepare owing to their huge size and fleshy parts.

That so few new species of *Philodendron* were collected was apparently due to the fact that in earlier times relatively few roads led into areas of wet forest.

Matuda, working exclusively in Mexico and concentrating on Araceae, accrued 27 collections of *P.* subg. *Philodendron* and only 1 of these, *P. glanduliferum*, proved ultimately to be new to science. George Bunting, also working only in Mexico (under the numbers of Harold Moore of Cornell University), made 36 collections of *P.* subg. *Philoden-*

*dron*, including 2 new species (*P. dressleri* and *P. jodavisianum*).

Steyermark, though collecting many more *Philodendron* in Venezuela later in his career, collected only 26 Central American *P.* subg. *Philodendron*, comprising 9 species, none of which were new. L. O. Williams, who also worked on the *Flora of Guatemala* and collected for many years in Guatemala, Honduras, and Costa Rica, made only 14 collections of *P.* subg. *Philodendron*. Two of these were species that remained undescribed until this work, namely Williams 628 (*P. sulcicaule*) and Williams 28387 (*P. wilburii* var. *longipedunculatum* Croat & Grayum). Percy Gentle, collecting in Belize, made 20 collections. The *Philodendron* subg. *Philodendron* collections of Pittier in Costa Rica totaled nine, none representing new species. Adolf Tonduz, also working in Costa Rica, made only five collections of this group. Alberto Brenes made four collections in Costa Rica between 1926 and 1932. Two of them, *P. brenesii* and *P. bakeri*, proved to be new. Paul Allen made five collections of *Philodendron* in Costa Rica, four of which were described as new (though two were subsequently synonymized). Aside from these few collections of new species mentioned above, no other new species were collected until the early 1960s. Roy Lent, living in Costa Rica and collecting between 1964 and 1971, made 24 collections of *P.* subg. *Philodendron*, including 5 new taxa. These are: *P. lentii* Croat & Grayum, *P. hederaceum* var. *kirkbridei*, *P. strictum*, *P. thalassicum* Croat & Grayum, and *P. wilburii* var. *wilburii*. W. C. Burger, collecting in Costa Rica between 1968 and 1986, in part with one-time aroid specialist Richard Baker, made 56 collections of *P.* subg. *Philodendron*, including 5 new species, *P. bakeri*, *P. chirripoense*, *P. crassispatum*, *P. thalassicum*, and *P. wilburii*.

Though Panama is even richer in Araceae than Costa Rica, the collecting activity there was not particularly rewarding. The *Philodendron* collections of H. von Wedel, who worked in Bocas del Toro Province in Panama, totaled only seven. Robert Woodson and his collaborators, Paul Allen, and Carrol Dodge collectively made only 11 collections before the *Flora of Panama* project was begun. This occurred in a country that proved to have 96 species, 65 new to science.

Even James Duke, who regularly got into areas of wet forest in many parts of Panama, made only four collections of *P.* subg. *Philodendron*, none of them new.

Collecting activities begun by Walter H. Lewis and staff from the Missouri Botanical Garden in the early 1960s were more aggressive by using heli-

copters, and made it into areas not previously accessible. Despite this greater penetration, even these expeditions resulted in few specimens of *Philodendron*. John Dwyer, one of the principal participants in these early Panama expeditions (and later Belize), made 14 collections of *Philodendron* (mostly from Panama), 2 of which [*P. dwyeri* (from Belize) and *P. stramineicaule* Croat]] were new to science. During the same period, Edwin Tyson, while teaching for Florida State University in the former Canal Zone, made 16 collections, 5 of them new to science: *P. annulatum* Croat & Grayum, *P. dolichophyllum* Croat, *P. lazarii* Croat, *P. llanense* Croat, and *P. tysonii* Croat. Even the late Alwyn Gentry, who made more than 80,000 collections of tropical American plants during his career and who went to many remote and interesting areas, made only 24 collections of *P. subg. Philodendron* in Central America. Three of them, *P. llanense*, *P. pseudauriculatum*, and *P. wilburii*, were undescribed at the time.

Collecting in Honduras, Costa Rica, and Panama between 1973 and 1993, Ron Liessner collected 43 species of *P. subg. Philodendron*, 4 of them, *P. alticola* Croat & Grayum, *P. crassipathum*, *P. findens*, and *P. llanense*, new to science.

Michael Grayum, an aroid specialist working primarily with *P. subg. Pteromischum*, was involved personally or in conjunction with other collectors, in making (since 1984) 426 *Philodendron* collections (in both Central and South America). Of this total, 153 were of *P. subg. Philodendron* for Central America. His collections are particularly useful, since they are accompanied by excellent field notes. He was responsible for collecting 13 undescribed species of *P. subg. Philodendron*: *P. angustilobum* Croat & Grayum, *P. aromaticum* Croat & Grayum, *P. bakeri*, *P. brunneicaule* Croat & Grayum, *P. cotobrusense*, *P. crassipathum*, *P. cretosum* Croat & Grayum, *P. dodonii* Croat & Grayum, *P. grayumii* Croat, *P. scalarinerve* Croat & Grayum, *P. stramineicaule*, *P. thalassicum*, and *P. wilburii*.

My own collecting activities with *Philodendron* began in 1967 with work on the *Flora of Barro Colorado Island* (Croat, 1978). Collecting on BCI and taking other trips in Panama with Walter Lewis, John Dwyer, and others gave me my first real experience gathering Araceae. Many of the areas visited were not previously explored by botanists, mostly in wet or very wet areas where numerous new Araceae occurred. Many species were collected for the first time even in areas where considerable collecting had been carried out in the past. In some cases nearly all the species collected on a given day proved to be new to science. Later, when

serving as the Missouri Botanical Garden's resident botanist in Panama, during 1970-1971, and while continuing my work on Barro Colorado Island, I took many field trips to other areas of Panama. By this time I had become seriously interested in collecting Araceae, and many plants were brought back to Summit Garden for cultivation and study. Upon my return to St. Louis, most of these collections were shipped to the Missouri Botanical Garden where many have persisted. In all, 3582 collections of *Philodendron* have been made by myself (or sometimes in conjunction with another collector) in Central and South America; 1715 of these were made in Central America, and 1594 of these were members of *P. subg. Philodendron*: these Central American collections represented 52 undescribed species.

#### FOSSIL HISTORY

The fossil history of the Araceae was reviewed recently by Grayum (1990). The records are scant and inconclusive, but macrofossils of Araceae showed the family to be relatively diverse by the early Tertiary (Daghljan, 1981; Gregor & Bogner, 1984), and at least four modern subfamilies were present by the Oligocene. No fossil pollen definitely ascribed to *Philodendron* has yet been encountered (Grayum, 1990). A plant fossil from Tennessee believed to be a *Philodendron*, and described as *P. limnestis* Dilcher & Daghljan (Dilcher & Daghljan, 1977), has proven to be a member of Peltandreae, either *Peltandra* or *Typhonodorum* (J. Bogner, pers. comm.). This fossil was believed by the authors to be a member of *P. subg. Meconostigma*; Mayo (1991) discussed in detail why he believed that it could not be a *Philodendron*, instead comparing it most closely with *Typhonodorum*.

#### INTRAGENERIC RELATIONSHIPS

The last thorough taxonomic revision of the Araceae was that by Engler (1905-1920), conducted in part by K. Krause (Engler & Krause, 1908, 1920; Krause, 1913). That revision included eight subfamilies of Araceae with *Philodendron* in its own subfamily and tribe, *Philodendroideae* and *Philodendreae*, respectively. *Philodendreae* (Engler, 1911) shared the subfamily with seven other tribes, namely, *Schismatoglottidinae*, *Anubiadeae*, *Aglaonemateae*, *Dieffenbachieae*, *Zantedeschieae*, *Typhonodorea*, and *Peltandreae*. Engler (1912) grouped *Philodendron* with *Homalomena* and the genera of the largely Asian subtribe *Schismatoglottidinae* (*Bucephalandra*, *Gamogyne*, *Microcasia*, and *Piptospatha*) into subtribe *Philodendrinae*.

Bogner and Nicolson (1991) left Engler's subfamily Philodendroideae intact, but Grayum (1984) made substantive changes including an incorporation of the subfamily Calloioideae and the Calocasioideae (thus forcing a change in the subfamilial name to Calloioideae, because of nomenclatural priority). The subfamily is divided into *Nepthytis*, *Aglaonema*, *Peltandra*, and *Philodendron* Alliances. In Grayum's system, *Philodendron*, in its own tribe Philodendreae, originally shared the alliance with the tribes Spathicarpeae, Dieffenbachieae, and Bogneriae. Both *Dieffenbachia* and *Bognera* are in their own tribes, but the Spathicarpeae have eight small genera mostly occurring in southern South America: *Asterostigma*, *Gearum*, *Gorgonidium*, *Mangonia*, *Spathantheum*, *Spathicarpa*, *Synandropadix*, and *Taccarum*. Grayum (1990) later placed *Philodendron* close to Homalomeninae and the African genera *Culcasia* and *Cercestis* (which had been placed in the Pothoideae and Lasiodeae, respectively, by Engler). They all share similar stem and stamen vasculature as well as extrafloral nectaries and resin canals in their roots and a sclerotic root hypodermis. Most (excluding Homalomeninae) also lack an anther endothecium that is present in all other Araceae (Grayum, 1990). Grayum believed that the Philodendroideae are a sister group to the Pothoideae (including Engler's Monsteroideae), which have in common the exclusive characteristics of geniculate petioles, cork formation in aerial roots, compound vascular bundles, collateral bundles, and other features (Grayum, 1984).

While there is reasonable agreement on the classification of tribes and subtribes, subfamilial concepts are still evolving. A comparison of the major systems of classification at the suprageneric level was made by Croat (1990[1992]). It included the systems of Hotta (1970), Grayum (1990), and Bogner and Nicolson (1991).

Phenetic analysis on the Philodendroideae by Mayo (1986) shows *Philodendron* to be distinct but without any definitive diagnostic features with which to distinguish it completely from other genera. He reported the genus to be only distantly related to other genera in the subfamily but that its closest relatives were the African genera *Culcasia* and *Cercestis*. In his survey of sclerotic hypodermis in the roots of Araceae, French (1987a) provided evidence to link *Philodendron* to the West African genera *Anubias*, *Culcasia*, and *Cercestis* and the neotropical *Montrichardia*.

More recently, in an attempt to bridge differences in the systems of Bogner and Nicolson, Grayum, and Hay and Mabberley (1991), Mayo et al. (1995) conducted another sweeping survey and produced

a cladistic analysis. While maintaining essentially the same alliances suggested by Grayum (1990), Mayo et al. (1995) placed all araceous genera with unisexual flowers in subfamily Aroideae. *Philodendron* is placed in tribe Philodendreae in the *Philodendron* Alliance, along with tribe Homalomenaeae with *Furtadoa* and *Homalomena* and tribe Anubiadeae (*Anubias* only). Their cladistic analysis also shows *Bognera* to be a close ally of *Dieffenbachia*.

Another cladistic analysis resulting from a study of chloroplast DNA restriction site variation in the Ariflorae by French et al. (1995) places *Philodendron* as a sister group to *Homalomena*, suggesting, according to Grayum (1996), that Homalomeninae is paraphyletic. According to French's findings, *Anubias* is a sister taxon of *Homalomena*, *Furtadoa*, and *Philodendron* taken together, while *Montrichardia* is a sister taxon to all four of these genera.

While questions still remain about the closest generic relatives of *Philodendron*, the classification by Mayo et al. (1995) has taken into account all the evidence to date including the extensive molecular studies by French.

The genera of Araceae most easily confused with *P. subg. Philodendron* are *Homalomena* and *Dieffenbachia*, with *Schismatoglottis* coming a distant third. Despite the somewhat more distant placement of *Dieffenbachia* from *Homalomena*, herbarium material of *Dieffenbachia* is most easily confused with *Philodendron*, sometimes requiring the opening of the spathe to determine the genus. Since the pistillate flowers of *Dieffenbachia* are distant from one another, surrounded by clavate staminodia and borne on a spadix that is fused throughout its length to the spathe, while those of *Philodendron* are closely compacted, devoid of staminodia, and borne on a spadix that is largely free from the spathe, separation from *Dieffenbachia* is immediate as long as the flowers are visible. Dried sterile material without field notes denoting terrestrial habit (consistently true of *Dieffenbachia*, but rarely so of *Philodendron*) or scent (usually foul and of oxalic acid in *Dieffenbachia*) are much more problematic. *Dieffenbachia* leaf blades are rarely ovate and never truly cordate, whereas this blade shape is common in *Philodendron*. *Philodendron* may, however, have blade shapes that closely match those of some *Dieffenbachia*. If the petiole is well preserved the presence of the petiole sheath is the best means to separate *Dieffenbachia* and *P. subg. Philodendron*; the latter generally has a very short sheath, while it is rare that the sheath of *Dieffenbachia* does not extend to the middle or above the middle of the petiole.

Live material of neotropical *Homalomena* is not easily confused with *Philodendron* because the for-

mer usually has anise-scented sap, while *Philodendron* usually has a distinct turpentine-like aroma, sometimes also like fresh carrots, but never anise-scented. *Philodendron* and *Homalomena* may have similar leaves, but the latter often has pubescence on the blades and pubescence and/or scales and spines on the petioles. These two genera have similar inflorescences with unisexual flowers, sterile and fertile sections of the staminate portion of the spadix, and a close arrangement of pistillate flowers, not to mention the similar constricted spathe that persists after anthesis. However, *Homalomena* can usually be determined by the presence of minute, club-shaped staminodia sparsely scattered among the pistils.

Sterile specimens of neotropical *Schimatoglottis* may be confused with *Philodendron* by the novice because the two genera share similar venation. However, *Schimatoglottis* always occurs terrestrially, often in somewhat marshy situations where *Philodendron* rarely occurs. In fertile condition they are easily separated by the spathe promptly dehiscent above the tube in *Schimatoglottis*, with the staminate portion of the spadix falling free. By contrast, in *Philodendron* the spathe is thick and persistent, usually reclosing over the staminate portion of the spadix, which in turn rots away inside, never really falling free until the spathe opens in fruit.

Sterile material of *Spathiphyllum* has been confused with *Philodendron*, but that genus differs by its consistently terrestrial habit, long-sheathed petioles (exhibited in *Philodendron* only in *P. subg. Pteromischum*), the presence of trichoscherids, and by its distinctive, closely spaced primary lateral veins.

#### INFRAGENERIC RELATIONSHIPS

*Philodendron* is currently divided into three subgenera. A subgeneric system of classification for *Philodendron* was proposed as early as 1832 by Schott, who recognized four unranked groups: *Euphilodendron*, *Calostigma*, *Meconostigma*, and *Sphinctrostigma*. The latter two were combined by Engler (1899) as *P. subg. Meconostigma*. Schott's *Calostigma* was later called *P. sect. Oligospermium* Engler (Engler, 1878) and is once more called *P. sect. Calostigma* [(Schott) Pfeiffer] (Mayo, 1990). *Euphilodendron* became *P. sect. Polyspermium* in Engler's *Flora Brasiliense* treatment in 1878 and must now be treated according to the Code (Greuter et al., 1994) as *P. sect. Philodendron* (Mayo, 1990).

It was not until Kunth's (1841) treatment that members of what are now called *P. subg. Pterom-*

*ischum* were removed from *Monstera* and placed in *Philodendron*. Schott recognized *Pteromischum* as a *grecx* in his 1860 *Prodromus*, and Engler first recognized the species occurring in this group as *P. sect. Pteromischum* in his *Flora Brasiliensis* treatment (Engler, 1878).

Phylogenetic and phenetic analyses by Mayo (1986, 1988) have shown *Philodendron* to have three subgenera distinct in vegetative and floral morphology, floral anatomy, and to some extent by distribution. *Philodendron subg. Meconostigma*, with a predominantly southeastern South American distribution, is highly apomorphic but cladistically primitive in the genus (Mayo, 1990). Based on a study of gynoecial morphology Mayo considers *P. subg. Meconostigma* to have evolved in eastern Brazil as a group adapted to open habitats and later spread into the more humid Amazon basin. By the same standard he assumed that *P. subg. Philodendron* and *P. subg. Pteromischum* also arose later and became predominant as hemiepiphytes in humid forests. He considered *P. subg. Pteromischum* to be a sister group to *P. subg. Philodendron* and that *P. subg. Philodendron* is the most advanced of the three subgenera. The geological history of the continent would probably support this since eroded mountain plateaus of eastern Brazil are much older than the current land surfaces of the Amazonian basin. Most of the species of the genus, now so rich on the Andean slopes of northern and western South America, surely must have evolved since the Andes arose during the late Cenozoic.

Mayo elevated *P. sect. Pteromischum* to the status of subgenus (Mayo, 1989) and Grayum (1996) subdivided the subgenus into two sections, *P. sect. Pteromischum* (Schott) Engler with sylleptic<sup>4</sup> sym-

<sup>4</sup> *Sylleptic shoots* are shoots that develop from lateral meristems without any cessation of activity after initiation, i.e., growth is continuous with the lateral shoot usually taking over from the main axis, which terminates in flowering in most Araceae genera. *Sylleptic sympodial growth*, characteristic of *P. subg. Philodendron*, results when branching occurs from a non-resting lateral bud such that the existing stem with its terminal inflorescence is immediately overtopped by continued growth of the axillary branch in a manner that makes it appear that the growth of the stem is indeterminate and that the inflorescences produced appear to be axillary to the leaf produced by the new stem segment. This type of growth, though in reality a series of branches, each with a single cataphyll (prophyll of Ray), a single stem segment, and a single leaf (sympodial foliage leaf of Ray; metaphyll of Grayum) and terminated by an inflorescence, appears to be an unbranched stem producing a continuous series of cataphylls, and a continuous series of what appears to be laterally attached leaves each with an axillary inflorescence (see fig. 3 in Ray, 1987b).



podial growth and *P.* sect. *Fruticosa* Grayum with proleptic<sup>1</sup> sympodial growth (Ray, 1987b), a growth form that is rare in the family, known only in *Alocasia* and a few species of *Monstera* (Grayum, 1996).

*Philodendron* subg. *Philodendron* is difficult to define and is primarily distinguished by its negatives, i.e., it lacks the specific characteristics of *P.* subgenera *Pteromisichum* and *Meconostigma* (see key to subgenera below). There are relatively few members of *P.* subg. *Philodendron* with a pachycaulous habit common to so many members of *P.* subg. *Meconostigma*, i.e., with very stout, generally erect stems and possessing conspicuous leaf scars. *Philodendron* subg. *Philodendron* also lacks the conspicuous, more or less triangular scales borne in the leaf axils of *P.* subg. *Meconostigma*. Though sometimes obvious (Fig. 11) in *P.* subg. *Philodendron*, they are usually small and inconspicuous and fall early. The species most similar to *P.* subg. *Meconostigma* is *P. warszewiczii*, but another species, *P. basii*, is similar in being large with a thick, erect stem.

Species of *P.* subg. *Pteromisichum* have subtle characteristics that to the expert permit immediate recognition. These characters include the slender, somewhat woody, brittle stem, a conspicuous petiole sheath, thinner blades with rather pronounced primary lateral veins, the presence of interprimary veins, and the frequent presence of raphide cells or stitch-like markings. Another feature that is often useful in separating *P.* subg. *Pteromisichum* from *P.* subg. *Philodendron* is the much higher incidence of asymmetrical leaf blades in the former. Species with asymmetrical (especially at the base) blades are not common in *P.* subg. *Philodendron*. Some species of *P.* subg. *Philodendron* may have oblong to elliptic, non-cordate blades like those of *P.* subg. *Pteromisichum*, but they never have the fully sheathed petioles of the latter (except in juvenile condition, which may confuse the non-expert).

The three subgenera of *Philodendron* in general can be most easily separated by the characters presented in the following key (modified after Mayo, 1991).

KEY TO THE SUBGENERA OF *PHILODENDRON*

- 1a. Stem of mature flowering plants with a succession of many leaves terminated by a solitary or commonly several inflorescences; petioles with long sheaths narrowly encircling the stem at their base ..... *P.* subg. *Pteromisichum*
- 1b. Stem of mature flowering plants with a succession of short sympodial segments each bearing a cataphyll and a single leaf with the inflorescence(s) 1 to many and appearing to be borne in the leaf axils; petioles of adult plants with short, usually inconspicuous petiole sheath and borne on the side of the stem, not encircling it at the base.
  - 2a. Stems often arborescent; staminodial zone between staminate and pistillate zones of the spadix subequal or longer than fertile zone; stamens usually at least 3 times longer than broad ..... *P.* subg. *Meconostigma*
  - 2b. Stems rarely arborescent, often scandent; staminodial zone between staminate and pistillate zones of spadix much shorter than the fertile staminate zone; stamens less than 3 times longer than broad ..... *P.* subg. *Philodendron*

There are also a number of anatomical characteristics separating the subgenera. Vegetative buds of *Philodendron* subg. *Philodendron* are always located below the point of overlap in the sheath margins of the cataphyll, whereas they are lacking in *P.* subg. *Pteromisichum* (Ray, 1987b). *Philodendron* subg. *Pteromisichum* is distinct in having a style with a shallow compitum<sup>4</sup> with a subepidermal concentration of raphide crystals (Mayo, 1986, 1989) and a total lack of tannin cells in the stamens (Mayo, 1986). In addition, while hypophyllous<sup>5</sup> stem segments are typical for *P.* subg. *Philodendron* they are ambiphylous,<sup>6</sup> hyperphylous,<sup>7</sup> or peraphyllous<sup>10</sup> in *P.* subg. *Pteromisichum*. *Philodendron* subg. *Philodendron* is characterized by having continuous parenchyma from the cortex to the center of the stem. In contrast, *P.* subg. *Pteromisichum* has a central cylinder with a solid ring of fibers around it.

While not definitive, there are a number of other features that normally are useful to separate *P.*

<sup>4</sup> The common depression that leads to individual stylar canals.

<sup>5</sup> Hypophyllous stem segments are those those in which the petiole scar borders the lower edge of the segment (see figs. 3, 7 in Ray, 1987b).

<sup>6</sup> Ambiphylous stem segments are so short that the petiole scars border on both ends of the stem segment (see figs. 9-11 in Ray, 1987b).

<sup>7</sup> Hyperphylous stem segments have the petiole scar at the upper edge of the stem segment (see figs. 2, 8 in Ray, 1987b).

<sup>10</sup> Peraphyllous stem scars are those in which the internode subtending the petiole "is elongated and supercedes the point of attachment" (Ray, 1987b).

<sup>1</sup> Proleptic sympodial growth, characteristic of members of *Philodendron* sect. *Fruticosa* (Grayum, 1996), results when branching occurs from a bud that has rested, relative to the parent shoot, so that the lateral branch does not appear until well after the inflorescence has set fruit and fallen off. Thus the inflorescence appears to be terminal, not axillary as in sylleptic sympodial growth. The first few internodes in proleptic growth are very short, lack buds, and produce a consecutive series of cataphylls or reduced leaves (see fig. 2 in Ray, 1987b).

subg. *Pteromischum*. Its blades are typically more or less oblong, moderately thin, and typically more inequilateral than those of *P. subg. Philodendron*. The stems of *P. subg. Pteromischum* are commonly less than 1 cm in diameter and frequently with rather long internodes. While there are many members of *P. subg. Philodendron* with more or less oblong leaf blades, blades of most species are cordate or subcordate at the base. Many members of *P. subg. Pteromischum* have stems that branch and spread away from their support before flowering; this behavior is rare in *P. subg. Philodendron*.

Because of the usually conspicuously sheathed petioles of *P. subg. Pteromischum*, the subgenus is more likely to be confused with sterile specimens of *Rhodospatha* than with the oblong-bladed species of *P. subg. Philodendron* (and in such cases the presence of trichosclereids beneath the epidermis of *Rhodospatha* easily distinguishes it from *Philodendron*).

#### RELATIONSHIPS WITHIN *P. SUBG. PHILODENDRON*

*Discussion of subgeneric classification.* Any attempt to revise the subgeneric classification of *Philodendron subg. Philodendron* is frustrated by the lack of morphological characteristics that correlate with one another throughout the subgenus. Engler (1899) separated the species of *P. subg. Philodendron* primarily on the basis of leaf shape, leaf blade venation, and the nature of the pistil (i.e., number of locules per ovary, type of placentation, and number of ovules per locule). Most sections, all moderately small ones, were separated on the basis of leaf morphology. These are: *P. sect. Tritomophyllum*, *P. sect. Schizophyllum*, and *P. sect. Polytomium*. Two sections, *P. sect. Macrogygium* and *P. sect. Campogynium*, are based on the nature of the pistil. Following a cladistic analysis of a relatively small number of species, including some in *P. subg. Philodendron*, Mayo (1986) concluded that *Philodendron* should be divided into two to three sections instead of the existing nine sections in the genus. While I agree that *P. sect. Macrolonchium* should be reduced, I think that a cladistic analysis making use of the leaves as well would justify the existence of the remaining sections recognized by Krause with the possible exception of *P. sect. Campogynium*, which was not studied by Mayo.

The most time-honored way to separate species in *P. subg. Philodendron* is based on number of ovules per locule, a system first devised by Engler stemming from his first revision of *Philodendron* (Engler, 1878). Engler used the number of ovules

per locule to separate two large groups, which he called *P. sect. Polyspermium* and *P. sect. Oligospermium* (now *P. sect. Philodendron* and *P. sect. Calostigma*, respectively). These two sections together comprise the largest percentage of species in the subgenus. As they are constituted they are very diverse morphologically, and it is possible that the number of ovules per locule will not prove to be reliable for separation at the sectional level. It is possible that species with relatively few ovules or solitary ovules may have evolved independently several times from ancestors with numerous ovules having axile placentation. Since it has not yet been determined if this is the case, the classification system used here will in general be conservative.

Krause (1913) closely followed Engler's sectional revision. His *P. sect. Philodendron* (as *P. sect. Polyspermium*) consisted of species with axile placentation and "many" ovules per locule, while *P. sect. Calostigma* (as *P. sect. Oligospermium*) consisted of those species with sub-basal (or less frequently basal) placentation with "1 or few" ovules per locule. Since the time of the last revision of *Philodendron* many species have been added, and the distribution of ovules per locule for all species now forms a more complete continuum. There is still a significant correlation between axile placentation and moderately large numbers of ovules per locule and the converse, basal and sub-basal placentation and relatively low numbers of ovules. Both *P. sect. Philodendron* and *P. sect. Calostigma* will be discussed below.

While these two groups, *P. sect. Philodendron* and *P. sect. Calostigma*, constitute the largest percentage of Central American species, several other sections are separated on the basis of leaf shape, venation, and style type (see above key). Each of these will be discussed in turn, despite the fact that some do not occur in Central America. Krause (1913) treated ten sections in his revision of *Philodendron*. As previously discussed, *P. sect. Pteromischum* has been elevated to the status of subgenus, and *P. subg. Macrolonchium* Engl. has been reduced to a subsection of *P. subg. Philodendron*.

1. *Philodendron sect. Baurisia* (Rchb. ex Schott) Engl., in Mart., Fl. bras. 3(2): 134. 1878. *Philodendron grex Baurisia* Rchb. ex Schott, Syn. Aroid. 73. 1856. TYPE: *Philodendron crassinervium* Lindl. (lectotype, designated by Mayo, 1990).

As defined by Engler and Krause, *P. sect. Baurisia* consists of species with generally inconspicuous primary lateral veins, but the group as constituted

by Krause remains highly variable in terms of its ovules, habit, and leaf shape. The group comprises species with moderately many ovules and axile placentation, moderately many ovules with basal placentation, a few ovules with basal placentation, and solitary ovules with basal placentation. All of these species have leaves purportedly devoid of primary lateral veins. In reality this is not true of all species included in the group. Most species have elongate, simple blades, but three have 3-lobed or tripartite blades.

With 33 species included by Krause the section was third in size of the three major sections (*P. sect. Philodendron* with 64 and *P. sect. Calostigma* with 53 species). One species, *P. acraeanum* K. Krause, is actually a member of *P. sect. Pteromischum*. Of the remaining species in the section, those that best fit the description of the group occur principally in eastern South America and in the upper Amazon basin and have more or less oblong leaf blades. *Philodendron crassinervium* is the type of the section. Except for *P. crassinervium*, *P. longilaminatum* Schott (with axile placentation), *P. bahiense* Engl., and *P. paxianum* K. Krause (each with solitary ovules per locule), species of *P. sect. Boursia* have a few basal ovules per locule and oblong to oblong-elliptic blades. Many but not all have indistinct primary lateral veins.

Some members of Krause's *P. sect. Boursia*, especially the species that are vines with a solitary ovule per locule and occur in the Andes west of the Continental Divide, such as *P. lehmannii* Engl., *P. ellipticum* Engl., *P. pachycaule* K. Krause, *P. chimboanum* Engl., *P. longipes* Engl., and *P. graveolens* Engl., do not seem to belong with the remainder and should perhaps be put into another section. The same is true of the three-lobed and tripartite species, *P. deltoideum* Poepp. & Endl., *P. panduriforme* (Kunth) Kunth (Krause also included here *P. reichenbachianum* Schott, now a variety of *P. panduriforme*), and *P. micranthum* Poepp. ex Schott. With the exception of *P. micranthum*, which has primary lateral veins lacking or weak, these species have primary lateral veins at least some of the time (though they are indistinct in *P. deltoideum*). None of the three species appear to have any other features in common with the more typical members of *P. sect. Boursia*, e.g., *P. crassinervium* Lindl., *P. linnaei* Kunth, and *P. callosum* K. Krause, among others.

Mayo (1986) believed that *P. sect. Boursia* contained two groups of species, and he would also separate *P. deltoideum* and related species from the remainder, suggesting that Schott's grex

*Oligophlebium* be recognized to accommodate these species.

Some species that were placed in *P. sect. Boursia* will have to be reinvestigated to determine if they belong instead in *P. sect. Philopsammos* G. S. Bunting (1986). That group is often similar in having elongated leaf blades, but it differs in having bilocular ovaries whereas those of *P. sect. Boursia* are plurilocular.

By no means all of the species with more or less oblong blades in *P. sect. Boursia* have primary lateral veins weak or lacking. At least one species, *P. wendlandii*, the only Central American species placed in *P. sect. Boursia* by Engler, should be placed in *P. sect. Calostigma*. It has distinct primary lateral veins and a spongy petiole with a distinct dark green annular ring like the other members of *P. subsect. Glossophyllum* in Central America. This leaves Central America without members of *P. sect. Boursia*.

2. *Philodendron* sect. *Philopsammos* G. S. Bunting, *Phytologia* 60: 306. 1986. TYPE: *Philodendron ptarianum* Steyer., Fieldiana, Bot. 28: 99. 1956. [*Philodendron callosum* K. Krause subsp. *ptarianum* (Steyer.) G. S. Bunting, *Phytologia* 64: 467. 1988.]

*Philodendron* sect. *Philopsammos* is restricted to South America, known largely from the region of the Guiana highlands with extensions into the Amazon basin, occurring principally on white sand savannas, sandstone outcrops, and on tepuis, rarely in alluvium in lowland forests. It is characterized by having usually terrestrial or epipetric, thick, creeping stems with mostly short internodes, mostly long persistent, mostly intact cataphylls, moderately long petioles, mostly erect, geniculate petioles, more or less oblong, elliptic to narrowly ovate, coriaceous blades, usually lacking any prominent posterior lobes and with usually distinct, sometimes moderately obscure primary lateral veins. Inflorescences are moderately large with pistils bilocular, rarely 3-locular, and ovaries moderately numerous with axile placentation.

Included in the section are the following species: *P. canaimae* G. S. Bunting, *P. craspedodromum* R. E. Schult., *P. dunstervilleorum* G. S. Bunting, *P. dyscarpium* R. E. Schult., *P. peraiense* G. S. Bunting, *P. phlebodes* G. S. Bunting, *P. pimichinense* G. S. Bunting, *P. callosum* K. Krause, *P. pulchrum* G. M. Barroso, *P. remifolium* R. E. Schult., *P. sabulosum* G. S. Bunting, *P. steyermarkii* G. S. Bunting, and *P. tatei* K. Krause. Perhaps also to be included

in this group is *P. englerianum* Steyerl. No species in the group occur in Central America.

In describing this section Bunting (1986) made no mention of how the section is distinguished from *P. sect. Boursia* (or any other section). This is a critical point since both sections have species with elongated blades and at least sometimes have primary lateral veins not markedly more prominent than the interprimary veins (secondary lateral veins). The section is presumably distinguished from *P. sect. Boursia* on the basis of having mostly two locules per ovary. As an indication of its affiliation with *P. sect. Boursia*, Bunting, at the time he described *P. sect. Philopsammos*, specifically mentioned *P. callosum* (a species included by Krause in *P. sect. Boursia* and initially considered by Bunting to be distinct from *P. ptarianum*) as being a possible member. *Philodendron ptarianum* has proven to be closely related to *P. callosum*, but that species was described by Krause as being "plurilocula," i.e., with many locules per ovary and "pauciovulata," i.e., with few ovules per locule. If this is true, the single character separating *P. sect. Philopsammos* from *P. sect. Boursia*, namely the small number of locules per ovary, would break down even in two subspecies (as now recognized by Bunting, 1995), one of which is the type of *P. sect. Philopsammos*. One collection of *P. callosum*, *Davidse & Miller 27269*, had 2-locular ovaries with 8–10 ovules per locule with unusual black, shiny seeds. It would clearly appear to be a member of *P. sect. Philopsammos*.

**3. *Philodendron* sect. *Philodendron*.** TYPE: *P. grandifolium* (Jacq.) Schott

*Philodendron* sect. *Philodendron* in Central America is both large and diverse, and like *P. sect. Calostigma* (a discussion of which follows) it is further subdivided here into subsections.

With 38 species (40 taxa) *P. sect. Philodendron* is the second largest section in *P. subg. Philodendron* in Central America. *Philodendron* sect. *Philodendron* is characterized by having axile placentation and typically many ovules per locule but, owing to its size and diversity, there are no other characters that completely characterize the group. *Philodendron* sect. *Polyspermium* (according to the Code (Art. 21) it must now be *P. sect. Philodendron*) was subdivided by Krause into six groups, "Gruppen" (termed subsections by Mayo, 1990). These subsections will be discussed here, especially in relation to the Central American species.

SUBSECTIONS OF *P. SECT. PHILODENDRON*

**1. *Philodendron* subsect. *Macrolonchium*** (Schott) Engl., in Mart., Fl. bras. 3: 139. 1878. *Philodendron* grex *Macrolonchium* Schott, Prodr. syst. Aroid. 269. 1860. TYPE: *Philodendron simsii* (Hook.) G. Don (lectotype, designated by Mayo, 1990: 64).

This is a small group of species characterized by D-shaped or broadly and sharply sulcate petioles and the presence of short stems with short internodes. The leaf blades typically are ovate-triangular. The cataphylls typically persist as fibers. The ovaries are 5–10-locular with numerous ovules per locule. This group was treated by Engler (1899) and Krause (1913) as a section based on stems with internodes shorter than broad versus scandent stems, but the group is in no way warranted at the sectional level based on this or any other character.

Though neither short stems nor D-shaped petioles are unique to this group, it appears to be natural at the subsectional level. All of the species have similar ovate-triangular blades and coarse reddish brown persistent cataphyll fibers. The two pinnately lobed species in the group, *P. pinnatifidum* (Jacq.) Schott and *P. robustum* Schott, seem unlikely members of this subsection. Those species are very similar to *P. fendleri* K. Krause, which Engler placed in his *P. sect. Polytomium*. *Philodendron pinnatifidum* and *P. robustum* should probably be placed there as well. They seem to have little in common with the other simple-leaved species. *Philodendron melinonii* Brongn., *P. fragrantissimum*, *P. simsii*, and *P. roraimae* K. Krause all appear to be related. The only Central American species in this subsection is *P. fragrantissimum*.

**2. *Philodendron* subsect. *Canniphyllum*** (Schott) Mayo, J. Linn. Soc., Bot. 100. 168. 1989. *Philodendron* grex *Canniphyllum* Schott, Syn. Aroid. 76. 1856. TYPE: *Philodendron fibrillosum* Poepp.

This is a small group that included only five species in Krause's revision and one of these, *P. coerulescens*, proved to be a synonym of *P. inaequilaterum*, a member of *P. subg. Pteromischum*. *Philodendron juninense* Engl., *P. fibrillosum*, and *P. cuneatum* Engl. are all Andean species, while the no doubt distantly related *P. blanchetianum* Schott is a species from eastern Brazil in Bahia. *Philodendron* subsect. *Canniphyllum* has only a few species in Central America. Both *Philodendron cretsum* and *P. roseospathum* have persistent fibrous cataphylls and resemble the type species, *P. fibrillosum*.

*losum* (selected by Mayo, 1990). The only other Central American species that has characteristics to fit in the subsection is *P. chirripoense*, which is tentatively placed here, though it differs in having longer, more slender internodes and lacks persistent cataphylls. The presence or absence of persistent cataphylls is highly correlated with the length of internodes (persistent when internodes are short, deciduous when internodes are long) and thus it is not surprising that *P. chirripoense* would lack cataphylls. Still, the latter is a most peculiar species so its placement remains in doubt.

**3. *Philodendron* subsect. *Platypodium*** (Schott) Engl., in Mart., Fl. bras. 3: 137. 1878. *Philodendron* grex *Platypodium* Schott, Syn. Aroid. 85. 1856. TYPE: *Philodendron pterotum* K. Koch & Augustin (lectotype, designated by Mayo, 1990: 61).

This group was characterized by Schott (1856) as having D-shaped petioles. While he included both *P. pterotum* and *P. fragrantissimum*, Engler later transferred the latter to his *P. sect. Macrolonchium* (Schott) Engl. Mayo (1990) designated the only remaining species in Schott's section, namely *P. pterotum*, as the type species.

In addition to *P. pterotum*, Krause (1913) placed in *P. sect. Platypodium* four more species, *P. clementis* Wright, now considered a synonym of *P. fragrantissimum* (a member of *P. subsect. Macrolonchium*); *P. splügerberi* Schott, a possible synonym of *P. fragrantissimum*; *P. lechlerianum* Schott, a close relative of *P. purpureoviride* (a member of *P. subsect. Solenosterrigina*); and *P. ernestii* Engl. Thus, of the species listed by Krause, probably only *P. pterotum* and *P. ernestii* belong here. It is unlikely that *P. ernestii* is very closely related to *P. pterotum*, but it does appear to be closely related to *P. brunneicaule*, another species placed in this section.

In addition to *P. pterotum*, other species of *P. subsect. Platypodium* that occur in Central America are: *P. brunneicaule*, *P. copense*, *P. findens*, and *P. fortunense* Croat.

**4. *Philodendron* subsect. *Psoropodium*** (Schott) Engl., in Mart., Fl. bras. 3: 138. 1878. *Philodendron* grex *Psoropodium* Schott, Syn. Aroid. 84. 1856. TYPE: *Philodendron ornatum* Schott (lectotype, designated by Mayo, 1990: 61).

As defined by Schott (1856), this is a group that has petioles verrucose at the apex. Mayo (1991) appropriately selected *P. ornatum* as the lectotype.

Krause's description, "petioles semirounded above, smooth or asperous, blade base cordate or sagittate," would accommodate more than half of the species in the genus *Philodendron*, but Schott's intention for the subsection was clear. The only other species included by Schott was *P. rubens* Schott, now considered by some authors to be a synonym of *P. ornatum*. Krause also expanded the group substantially, adding a number of plants that do not have glandular petioles and others that have proven to be synonyms of *P. ornatum*. The latter are: *P. muschlerianum* K. Krause, *P. dolosum* Schott, *P. asperatum* K. Koch, and *P. tobagoense* Engl. The aberrant elements which, I believe, are unrelated to *P. ornatum* include *P. brevilaminatum* Schott and *P. traunii* Engl. (now both synonyms of *P. fragrantissimum* in *P. subsect. Macrolonchium*), *P. grandipes*, and *P. maximum* K. Krause, a gigantic species from the southwest Amazon basin that is unlikely to be related in any way to *P. ornatum*. I have moved *P. grandipes* from *P. subsect. Psoropodium* to *P. subsect. Philodendron*, where it more appropriately belongs.

Krause also included in his *Psoropodium* group three other poorly known species, *P. thaliifolium* Schott, *P. brandtianum* K. Krause, and *P. bertae* K. Krause. These do not appear to be in any way related to *P. ornatum*. It is clear from these as well as the other species included by Krause that the section as defined by Engler and by Krause no longer resembles Schott's original description. Probably a number of the species will need to be moved into other groups, or the section will need to be more well defined. As currently defined no species in *P. subsect. Psoropodium* occur in Central America.

**5. *Philodendron* subsect. *Solenosterrigina*** (Klotzsch ex Schott) Engl., in Mart., Fl. bras. 3: 139. 1878. *Philodendron* grex *Solenosterrigina* Klotzsch, Syn. Aroid. 81. 1856. TYPE: *Philodendron scandens* K. Koch & Sello [(=*Philodendron hederaceum* (Jacq.) Schott (lectotype, designated by Mayo, 1990: 61)].

As treated by Krause (1913), this appears to be a natural group, consisting of *P. hederaceum*, the type, as well as *P. consanguineum* Schott and a number of relatives. *Philodendron fuertesii* K. Krause, *P. krebsii* Schott, and *P. urbanianum* K. Krause all closely resemble *P. consanguineum*, and *P. marginatum* Urb., *P. prievianum* Schott, *P. oxycardium*, *P. micans*, and *P. melanochrysum* Linden & André are all synonyms or subspecies of *P. hederaceum*. These species share long, slender inter-

nodes, deciduous, mostly unribbed cataphylls, more or less terete petioles, ovate-cordate leaf blades, and solitary inflorescences.

Other species included by Krause in *P.* subsect. *Solenostergma* are *P. purpureoviride* (aside from *P. hederaceum* and *P. brevispathum*, the only other species of the group that occurs in Central America), *P. jenmanii* K. Krause, and *P. scabrum* K. Krause. The latter two species are now considered synonyms of *P. muricatum* Willd. ex Schott. Though surely belonging in *P.* subsect. *Solenostergma* (along with the two synonyms already placed here) Krause placed *P. muricatum* in *P. sect. Calostigma*.

*Philodendron brevispathum* is transferred here from *P.* subsect. *Cardiobelium* (Schott) Engl., since it appears to be closely related to *P. muricatum*.

## 6. *Philodendron* subsect. *Philodendron*

*Philodendron grex Cardiobelium* Schott, Syn. Aroid. 88. 1856. *Philodendron* subsect. *Cardiobelium* (Schott) Engl., in Mart., Fl. bras. 3: 139. 1878. *Philodendron* "Gruppe" *Cardiobelium* (Schott) Engl., Bot. Jahrb. Syst. 26: 522, 529. 1899. TYPE: *Philodendron giganteum* Schott (lectotype, designated by Mayo, 1990: 60).

*Philodendron grex Eubelium* Schott, Syn. Aroid. 92. 1856. *Philodendron* subsect. *Eubelium* (Schott) Engl., in Mart., Fl. bras. 3: 140. 1878. TYPE: *Philodendron grandifolium* (Jacq.) Schott.

As defined by Schott (1860), *grex Cardiobelium* consisted of a single species, *P. giganteum*. The group was greatly expanded by Engler (1899) and by Krause (1913) as a subgroup within section *Polyspermium*. Now with 20 species, it constitutes the largest subsection in *P. sect. Philodendron*. The characterization, as expanded by Krause, "petioles smooth or lightly striate, asperate; blade cordate to sagittate, with the primary lateral veins much more conspicuous than the secondary veins," is so broad that many unrelated species might easily be contained within it. Certainly to be excluded is *P. rubens* Schott (now a synonym of *P. ornatum*) and the core species in *P.* subsect. *Psoropodium*. Among the well known and seemingly distinct taxa included by Krause are *P. grandifolium*, *P. acutatum* Schott, and *P. fraternum* Schott, all with deciduous cataphylls, and *P. tenue*, *P. schottianum*, and *P. panamense*, with persistent cataphylls. The type, *P. giganteum*, also has conspicuous persistent cataphylls. A particularly unusual species included by Krause is *P. quitense* Engl., with deeply three-lobed leaves. It is probably a synonym of *P. acuminatissimum* Engl. in *P.* subsect. *Doratophyllum*.

The only Central American species included in the section by Krause were: *P. brevispathum*, *P. panamense*, *P. schottianum*, and *P. tenue*. *Philoden-*

*dron brevispathum*, with its scaly stems, is best accommodated with *P. muricatum* in *P.* subsect. *Solenostergma*.

*Philodendron* subsect. *Philodendron* has 20 species in Central America. The size and diversity of *P.* subsect. *Philodendron* warrant the recognition of the five series presented below.

### SERIES OF *P.* SUBSECTION *PHILODENDRON*

#### 1. *Philodendron* ser. *Philodendron*. TYPE: *P. grandifolium* (Jacq.) Schott

The series is not known from Central America, but has several species in South America. This group is characterized by thick stems, deciduous cataphylls, subterete petioles, large ovate-triangular blades conspicuously lobed at the base, several inflorescences per axil, and 5-7-locular ovaries with many ovules per locule. In addition to the type, *P. acutatum* Schott, *P. billietiae* Croat, and *P. megalophyllum* Schott appear to be typical members of this group, having many ovules per locule, deciduous cataphylls, more or less terete petioles, and ovate-triangular leaf blades.

#### 2. *Philodendron* ser. *Impolita* Croat, ser. nov. TYPE: *P. strictum* G. S. Bunting

*Internodia brevia*; cataphylla persistentia; petiolus teres vel D-formatus; lamina palida adaxialiter, glaucescens.

*Etymology.* From *impolitus* meaning unpolished, i.e., matte, in reference to the matte and pale lower blade surfaces.

Three Central American taxa, *P. hebetatum*, *P. strictum*, and *P. thalassicum*, and at least one additional undescribed species from South America belong in this group. It is characterized by having leaf blades dark to medium green above but very pale, almost white beneath and covered with a minute waxy covering making the surface matte. The species have relatively short internodes, persistent cataphylls that often dry yellowish or have patches of yellowish epidermis persistent (not always true for *P. thalassicum*). Petioles may be obtusely somewhat flattened to D-shaped and dry with a yellowish epidermis (not always true for *P. thalassicum*). Blades are ovate-cordate to ovate-triangular.

#### 3. *Philodendron* ser. *Velvetina* Croat, ser. nov. TYPE: *P. gigas* Croat

*Internodia brevia*, maxime crassa; cataphylla D-formata, persistentia in fibris tenuibus; petiolus subteres; lamina ovata-cordata, bicolorata, velutina adaxialiter, *impolita* abaxialiter, 81-125 cm longa, 37-90 cm lata.

*Etymology.* From *velutinus* meaning velvet, referring to the velvety upper surfaces of the blades.

The subsection consists of a single species, *P. gigas*, in Central America but would also include *P. andreanum* Devansaye from northern Colombia. The group is characterized by its stout stems, short internodes, cataphylls that persist as thin fibers, subterete petioles, and especially by the very large, narrowly ovate-cordate blades, which are velvety and bicolored on the upper surface with the midrib and primary lateral veins paler, and paler and matte on the lower surface. Inflorescences range up to 7 per axil.

**4. *Philodendron* ser. *Fibrosa* Croat, ser. nov.**  
TYPE: *P. jodavisanum* G. S. Bunting

Internodia brevia, cataphylla persistentia ut fibrae; petiolum plerumque teres aut subterete; lamina ovato-cordata; pistilla cum placentatione axiali; loculi pluriovulati.

*Etymology.* *Fibrosa* = composed of separable fibers, in reference to the cataphylls persisting as fibers.

This represents the largest series in *P.* subsect. *Philodendron*. It is characterized by thick stems, short internodes, cataphylls that usually persist as a mass of fibers on the stem, usually subterete petioles, and more or less ovate-cordate blades. Pistils have axile placentation and many ovules per locule.

The following species of *P.* ser. *Fibrosa* occur in Central America: *P. alticola*, *P. antonioanum*, *P. breedlovei*, *P. chiriquense*, *P. dodsonii*, *P. grandipes*, *P. jodavisanum*, *P. lazorii*, *P. llanense*, *P. panamense*, *P. pirrense*, *P. purulhense*, *P. scalarinerve*, *P. schottianum*, and *P. tenue*. Of these a few are still doubtful. *Philodendron grandipes*, with a D-shaped petiole, and *P. jodavisanum*, with a U-shaped petiole, perhaps belong in *P.* subsect. *Platypodium* but also do not seem to be related to the species in that group. *Philodendron breedlovei*, which appears to lack persistent cataphylls (specimen very incomplete), is at odds with the other species but fits nowhere else. A few species are in related clusters within the series; for example, *P. lazorii* and *P. panamense* seem to be closely related as do *P. grandipes* and *P. jodavisanum*.

Grayum (pers. comm.) believes that *P. dodsonii* perhaps belongs with *P. pterotum* in *P.* subsect. *Platypodium*.

**5. *Philodendron* ser. *Albisuccosa* Croat, ser. nov.** TYPE: *P. albisuccus* Croat

Internodia brevia; succus albus, calcareus; cataphylla persistentia in fibris tenuibus pallidis; petiolum subterete;

lamina ovato-cordata; inflorescentia solitaria; pistilla 5-6-locularia; loculi cum 18-20 seminibus.

*Etymology.* *Albus* = white; *succus* = juice, sap, in reference to the white sap of cut parts.

The series consists of a single species, *P. albisuccus*, characterized by having copious white sap, which turns chalky on drying. Only one other species in Central America, *P. cretosum*, shares this unusual feature. It may belong in the same series despite its linear to oblanceolate leaf blades. It currently is placed in *P.* subsect. *Canniphyllum*.

**7. *Philodendron* subsect. *Achyropodium***  
(Schott) Engl., in Mart., Fl. bras. 3: 139. 1878.  
*P. grex* *Achyropodium* Schott, Syn. Aroid. 85. 1856. TYPE: *P. verrucosum* L. Mathieu ex Schott

This is a seemingly quite natural group characterized by scaly petioles. As defined by Schott (1856) it was represented only by the type, *P. verrucosum*. Krause (1913) included six species (one of which, *P. arcuatum* K. Krause, is a synonym of *P. brevispathum* in *P.* subsect. *Solenostigma*). The others in the subsection are: *P. nanegalense* Engl., *P. pilatonense* Engl., and *P. gualeanum* Engl. (all of which, I believe, represent a single species), as well as *P. serpens* Hook. f. All but *P. verrucosum* were known only from South America. There are several other undescribed species in *P.* subsect. *Achyropodium* now known from South America.

*Philodendron* subsect. *Achyropodium* is largely restricted to the northern Andes and lower Central America (Costa Rica and Panama) but has one species, *P. verrucosum*, that ranges as far south as Peru. The subsection is represented in Central America by *P. glanduliferum* subsp. *glanduliferum*, *P. hammelii*, *P. malesovichiae*, *P. squamicaulis*, *P. squamipetiolatum*, and *P. verrucosum*.

**4. *Philodendron* sect. *Calostigma* (Schott) Pfeiffer, Nomencl. Bot. 2: 674. 1874. *P.* [rankless]**  
b. *Calostigma* Schott, in Schott & Endl., Melet. Bot. 19. 1832. TYPE: *P. imbe* Schott

*Philodendron* sect. *Calostigma* is the largest section in Central America with 48 species comprising 52 taxa. It is characterized by having basal or sub-basal placentation and typically solitary or few ovules per locule. Owing to its size and diversity, there are no other characters that completely characterize the group.

There are some differences between *P.* sect. *Philodendron* and *P.* sect. *Calostigma* in Central America that are statistically significant, even if not without exception. For example, 81% of those spe-

Table 1. Leaf blade breakdown by section. # = number of species.

Section	Blades entire										Section total
	With basal lobes				Lacking basal lobes				Blades not entire		
	Cordate		Subcordate		Ovate		Oblong		Incised-lobate		
#	% of sect.	#	% of sect.	#	% of sect.	#	% of sect.	#	% of sect.		
<i>Calostigma</i>	28	56%	7	14%	1	2%	14	28%	0	0%	50
<i>Macrogynium</i>	1	100%	0	0%	0	0%	0	0%	0	0%	1
<i>Philodendron</i>	34	81%	3	7%	1	2%	3	7%	1	2%	42
<i>Polytomium</i>	0	0%	0	0%	0	0%	0	0%	3	100%	3
<i>Tritomophyllum</i>	3	38%	0	0%	0	0%	0	0%	5	63%	8

cies in *P.* sect. *Philodendron* have cordate blades, while only 7% have subcordate blades, and 12% lack posterior lobes. In contrast, *P.* sect. *Calostigma* has only 56% of its species with cordate blades, 14% have subcordate blades, and 30% lack posterior lobes. See Table 1 for a complete breakdown by section.

See other comparisons between *P.* sect. *Calostigma* and *P.* sect. *Philodendron* under sections entitled "Ovary Locule Number" and "Ovules Per Locule" as well as in Appendix 2, Technical Data on Pistils.

*Philodendron* sect. *Calostigma* was subdivided by Krause into five groups (termed subsections by Mayo, 1990). These subsections will be discussed here, especially in relation to the Central American species.

**1. *Philodendron* subsect. *Macrobelum***  
(Schott) Engl., in Mart., Fl. bras. 3: 143. 1878.  
*P.* grex *Macrobelum* Schott, Syn. Aroid. 96.  
1856. TYPE: *P. sagittifolium* Liebm.

*Philodendron* subsect. *Belocardium* (Schott) Engl., in Mart., Fl. bras. 3: 141. 1878. *P.* grex *Belocardium* Schott, Prodr. Syst. Aroid. 255. 1860. TYPE: *P. advena* Schott (lectotype, designated by Mayo, 1990).

As reported by Krause, *Philodendron* subsect. *Macrobelum* was the largest subsection in *P.* sect. *Calostigma*, with 22 species. It was poorly defined by Krause as consisting of species with somewhat spongy petioles, 6–12-locular ovaries with relatively few ovules per locule, and basal to sub-basal placentation. In addition (although not stated by Krause), the blades are all cordate, sagittate, or hastate. Although Krause described the subsection as having 2–5 ovules per locule, many more species have been added to the group since the last revision and several species that logically belong in *P.* subsect. *Macrobelum* have more than 10 ovules per locule, despite having sub-basal placentation.

*Philodendron* subsect. *Belocardium* must be synonymized with *P.* subsect. *Macrobelum* since *P. advena*, clearly a close relative of *P. sagittifolium*, was designated the lectotype of the former subsection, and Schott's definition of grex *Macrobelum* differs little from grex *Belocardium*, primarily since the blades are described as elongate-sagittate with 6–7 primary lateral veins instead of ovate-cordate blades with 3–4 primary lateral veins for grex *Belocardium*. As defined originally by Schott (1860), *P.* grex *Belocardium* comprised vining plants with ovate-cordate blades with the posterior rib mostly lacking and not at all naked along the sinus and bearing solitary inflorescences with 1–2 ovules per locule. The group consisted of *P. subovatum* Schott (= *P. advena*), *P. hoffmannii* (= *P. hederaceum*), *P. deviatum* Schott (= *P. jacquini*), *P. advena*, *P. populneum* K. Koch, *P. erubescens* Linden, *P. jacquini*, *P. lindenii* Schott, *P. acrocardium* Schott, and *P. consanguineum* Schott. Several of these species are now known to be synonyms of *P.* sect. *Oligocarpidium* or *P.* sect. *Macrogynium*. *Philodendron lindenii* would have better served as a lectotype for the group and, along with *P. erubescens*, *P. deviatum*, and *P. populneum*, might have to be formed into a group distinct from *P.* sect. *Belocardium*.

Engler (1913) synonymized Schott's grex *Glossophyllum* with *P.* "Gruppe" *Belocardium*, but the former group, as defined by Schott, appears to be distinct and will be resurrected in this work. It comprises more or less scandent plants with tumid petioles and more or less oblong blades that are frequently subcordate or cordulate at the base (see that section for more details).

Because of its size (28 species) and the diverse composition of the species comprising *P.* subsect. *Macrobelum*, it was deemed necessary to divide the subsection into four series. A discussion of the new series of *P.* subsect. *Macrobelum* follows.



**1. *Philodendron* ser. *Macrobelyum* (Schott)  
Croat, ser. nov. TYPE: *P. sagittifolium* Liebm.**

*Philodendron* ser. *Macrobelyum* is the largest group of species in *P.* subsect. *Macrobelyum* and includes the type. *Philodendron* ser. *Macrobelyum* is distinguished by having moderately coriaceous, cordate to sagittate blades with the basal posterior rib usually not naked or only weakly naked near its base. In addition, cataphylls are usually deciduous or only briefly persistent. Most of the species in the series have a strong resemblance to the widespread *P. sagittifolium*. All members have type D styles (see section on style type), relatively few ovules per locule (usually 1–5, rarely to 6, 7, or 8). In most cases the ovules are contained within a translucent or transparent ovule sac.

Central American representatives of *P.* ser. *Macrobelyum* are: *P. advena*, *P. annulatum*, *P. aromaticum*, *P. coloradense*, *P. dwyeri*, *P. edematatum*, *P. ferrugineum*, *P. grayumii*, *P. knappiae*, *P. mexicanum*, *P. platypetiolatum*, *P. sagittifolium*, *P. subincisum*, *P. sousae*, *P. verapazense*, and *P. zhuangum*. *Philodendron platypetiolatum* is unusual in having a much-flattened petiole. Perhaps it warrants recognition as a separate series.

*Philodendron mexicanum*, long considered a member of the group, is unusual in not greatly resembling *P. sagittifolium* and in having blades sometimes more or less hastate.

It is possible that *P. annulatum* belongs in *P.* subsect. *Glossophyllum* because it sometimes has a purple annular ring at the apex of the petiole (one of the features characterizing this subsection) and usually has only 1–2 ovules per locule (though sometimes 4–5 ovules per locule).

**2. *Philodendron* ser. *Ecordata* Croat, ser. nov. TYPE: *P. brenesii* Standl.**

Internodia elongata vel brevica; cataphylla decidua; petioli subteres; lamina acuta vel subcordata ad basin; nervia minores saepe "etched" in superficie supra; pistilla 5–14 locularia; loculi plerumque 4–14 ovulati.

*Philodendron* ser. *Ecordata* represents a group of species with elongate stems, internodes often longer than broad, deciduous cataphylls, elongate petioles, and blades that are ovate to ovate-elliptic, acute, or at most subcordate at the base (hence the name "ecordata," meaning without a cordate blade), often with the minor veins on the upper surface weakly etched. The basal veins are either free to the base or if they are united into a posterior rib, the latter is not naked or is naked for only a short distance. Pistils are 5–14-locular and locules are 4–14-ovulate.

The Central American species of *P.* ser. *Ecordata* are: *P. brenesii*, *P. crassispatum*, *P. davidsonii*, *P. lentii*, and *P. niqueanum*. A relationship between *P. brenesii* and *P.* ser. *Impolita* is possible, based on the glaucous lower blade surface in *P. brenesii*, but no other member of the group has glaucous leaves. Moreover, all the members of the above group usually have internodes longer than wide and have deciduous, rather than persistent cataphylls.

**3. *Philodendron* ser. *Reticulata* Croat, ser. nov. TYPE: *P. tysonii* Croat**

Internodia brevica; cataphylla persistentia in fibris tenuibus; petiolus lamina subaequalis aut longior, teres aut subteres; lamina ovato-cordata; pistilla cum loculis 5–9-ovulatis; ovulae plerumque 5–7 per ovaria.

*Etymology.* Reticulus, meaning netted, referring to the dried network of cataphyll fibers characterizing this series.

*Philodendron* ser. *Reticulata* is the only group of species in *P.* subsect. *Macrobelyum* with a few ovules per locule, stems with short internodes, and persistent cataphylls. Most species with short internodes and persistent cataphyll fibers are members of *P.* sect. *Philodendron*. Species in this group have terete or subterete petioles about as long as the ovate-cordate blades and pistils with 5–9 locules, each with 5–7 ovules.

*Philodendron* ser. *Reticulata* is represented in Central America by only two species, *P. jefense* and *P. tysonii*.

**4. *Philodendron* ser. *Pachycaulia* Croat, ser. nov. TYPE: *P. basii* Matuda**

Caudex succulentus; internodia brevica, crassa; cataphylla 2-costata, persistentia intacta; petiolus teres; lamina ovato-cordata, costae postice desinuae secus sinum ad 4 cm; pistilla 4–6-locularia; loculi 4–6 ovulati.

*Etymology.* Pachy = thick, caule = stem, referring to the thick, succulent stems.

*Philodendron* ser. *Pachycaulia* is represented by a single species and is characterized by its very stout succulent stems with intact persistent cataphylls, terete petioles, ovate-cordate blades with the posterior rib naked along the sinus to 4 cm. The pistil has a type D style and is 4–6-locular with 4–6 ovules per locule. It is believed that the succulent stems evolved to store water, allowing the plant to survive the long dry season in western Mexico. The series is represented only by *P. basii* from western Mexico.

**2. *Philodendron* subsect. *Glossophyllum*** (Schott) Croat, comb. nov. Basionym: *Philodendron* grex *Glossophyllum* Schott, Syn. Aroid. 80. 1856. TYPE: *P. elaphoglossoides* Schott (lectotype, designated by Mayo, 1990).

**1. *Philodendron* ser. *Glossophyllum*** Croat, ser. nov. TYPE: *P. elaphoglossoides* Schott

*Philodendron* subsect. *Glossophyllum* consists of two new series, *Glossophyllum* and *Ovata*.

*Philodendron* ser. *Glossophyllum*, as defined here, has the appearance of being a natural group of species with more or less oblong leaves acute or frequently cordulate or subcordate at the base. The primary lateral veins are usually distinct. Stems are typically somewhat scandent, though some members of the group, such as *P. auriculatum*, *P. bakeri*, *P. dolichophyllum*, *P. ligulatum*, *P. morii*, *P. pseudauriculatum*, *P. utleyanum*, and *P. wendlandii*, sometimes have the internodes scarcely longer than broad. The petioles are usually spongy or subspungy, usually subterete, and often bear a purplish or greenish annulus around the circumference where the petiole joins the blade; the cataphylls may be unribbed or sharply 1-2-ribbed and are typically deciduous (though persisting for a time in some species with short internodes, e.g., *P. auriculatum* and *P. wendlandii*). The style type is variable in the group, with most having B or D type styles but with one species, *P. granulare*, having an unusual type E style. Many species in the series have orange berries. *Philodendron* ser. *Glossophyllum* ranges from Nicaragua to Colombia and Ecuador on the Pacific slope and to the Guianas and the Amazon basin.

Krause included this group of plants with his section *Belocardium*, comprising both Schott's grex *Belocardium* and grex *Glossophyllum*. The former group consisted of plants with elongate internodes and ovate-cordate to sagittate blades, and Krause included *P. subovatum* Schott (= *P. advena*), *P. lindeni* Schott, *P. weberbaueri* Engl., *P. smithii*, *P. subhastatum* Engl. & K. Krause, *P. myrmecophyllum* Engl., *P. pachyphyllum* K. Krause, *P. advena*, and *P. viride* Engl. The latter group included species with mostly oblong blades. He characterized his section *Belocardium* as having tumid petioles and unilocular ovaries. Most of the species in the cordate-bladed group are not believed to be closely related to *P.* ser. *Glossophyllum* and have been referred here to *P.* subsect. *Macrobelyum* (see discussion of *P.* subsect. *Macrobelyum*). What remains is a group that usually has more or less oblong blades

with tumid petioles that are purple-ringed at the apex and ovaries usually unilocular.

*Philodendron smithii*, the only other Central American species among those mentioned above, does indeed have tumid petioles and only a single ovule per locule like most members of *P.* subsect. *Glossophyllum*, but the great difference in leaf shape in this species and others placed here warrant their separation into another series within *P.* subsect. *Belocardium* (see below).

Though Krause's revision characterized *P.* subsect. *Belocardium* as having a single ovule per locule, that is not in itself the defining feature of the subsection. For example, several species typical of the group have oblong leaves, purple-ringed petioles, and the same general appearance but have more than one ovule per locule. These are: *P. auriculatum*, with (3)4 ovules per locule; *P. ligulatum* var. *heracleianum*, and *P. ligulatum* var. *ovatum*, both with 2; *P. pseudauriculatum*, with 1-2(4); and *P. wendlandii*, with 2. *Philodendron bakeri* sometimes has 2 ovules per locule. Other Central American species, each with 1 ovule per locule are *P. brewsterense*, *P. correae*, *P. dolichophyllum*, *P. flosomii*, *P. granulare*, *P. immixtum*, *P. ligulatum*, *P. morii*, *P. ubigantupense*, and *P. utleyanum*.

Typical South American species in *P.* subsect. *Glossophyllum* presented by Krause (1913) are: *P. cyrtocleum* Diels (= *P. ruizii* Schott), *P. longipetiolatum* Engl., *P. heterophyllum* Poepp., *P. uleanum* Engl., *P. adhatodifolium* Schott, *P. elaphoglossoides* Schott, *P. ruizii* Schott (erroneously placed in sect. *Boursia* by Schott), *P. wittianum* Engl., and *P. angustialatum* Engler. *Philodendron tenuipes* Engl., placed in *P.* subsect. *Belocardium* by Engler, appears to be closely related to *P. fibrillosum* Poepp. and probably belongs in *P.* subsect. *Canniphyllum*.

Other species of *P.* subsect. *Glossophyllum* described since the last revision by Krause are *P. acutifolium* K. Krause, *P. buntingianum* Croat, *P. liessneri* Croat, and *P. wardackii* G. S. Bunting.

Though, as defined here, *P.* subsect. *Glossophyllum* ser. *Glossophyllum* consists only of species with more or less oblong blades, some of the species are somewhat anomalous. *Philodendron granulare* is much like other species in *P.* subsect. *Glossophyllum*, but it has a style that is unique in the group (Style Type E; see discussion of this under that species). *Philodendron brewsterense* and *P. ubigantupense* are still poorly known but appear to belong here. Perhaps the most doubtful is *P. dolichophyllum*, with 3-7 ovules, which is certainly high for this group; however, it is otherwise similar in most aspects to other members of the subsection. *Philodendron wendlandii*, placed by Krause in *P.* sect.

*Boursia*, seems to fit best in *P.* subsect. *Glossophyllum*. It differs from other members of subsection *Glossophyllum* in having a petiole that is usually broader than thick and sharply flattened adaxially and lacks an annular ring.

**2. *Philodendron* ser. *Ovata* Croat, ser. nov.**

TYPE: *P. smithii* Engl.

Caudex longus vel brevis; internodia plerumque longiora quam lata; cataphylla plerumque decidua; petioli subaequantur laminam, teres vel subteres, cum annulo deficienti apice; lamina ovata vel ovato-triangularata, cordata vel sagittata ad basi; pistilla cum stylo plerumque "type B," rare C; ovaria 4-8-locularia; loculi uniovulati aut raro 2-5.

*Philodendron* ser. *Ovata* is a somewhat heterogeneous group characterized by blades ovate to ovate-triangular and cordate to subcordate at the base, with terete or subterete petioles lacking an annular ring at the apex. Locule number varies from 4 to 8, and each usually has a single ovule. Berry color is greenish white to white, yellowish to lavender for most species known, but *P. cotonense* and *P. wilburii* have orange fruits. Though only seven Central American species are known, the series undoubtedly has South American representatives.

The relationship with other members of *P.* subsect. *Glossophyllum* is uncertain. Aside from the usually solitary ovule per locule and usually somewhat spongy petiole, there is little resemblance between this group and typical members of *P.* subsect. *Glossophyllum*, which have elongated, rather than more or less ovate, blades. *Philodendron dominicalense* is the most doubtful member of the group; with the strongest possible similarity to *P. dodsonii* except for having a solitary ovule per locule (rather than about 20 ovules per locule as in *P. dodsonii*), it would appear that it might belong elsewhere. *Philodendron cotonense*, despite having 4-5 ovules per locule, appears also to belong to this section, considering its marked resemblance to *P. wilburii* and even *P. smithii*.

Central American species of *P.* ser. *Ovata* are: *P. cotonense*, *P. dominicalense*, *P. microstictum*, *P. smithii*, *P. stramineale*, *P. sulcicale*, and *P. wilburii*. With the exception of *P. smithii*, which ranges from Mexico to Nicaragua, *P.* ser. *Ovata* in Central America ranges from Nicaragua to Panama.

**3. *Philodendron* subsect. *Oligocarpidium***

(Engl.) Mayo, J. Linn. Soc., Bot. 100: 168. 1989. TYPE: *P. multispadicum* Engl. (lectotype, designated by Mayo, 1990).

As defined by Krause, this was a group of four

species that differed greatly from one another. *Philodendron deviatum* Schott has proven to be a synonym of *P. jacquinii* (which was placed by Krause in its own *P.* sect. *Macrogynium*). Another species that he included, *P. pittieri*, is just poorly preserved material of *P. hederaceum*, which Krause placed in his *P.* "Gruppe" *Solenosterigma* in *P.* sect. *Philodendron*. This leaves only *P. multispadicum* Engl. and *P. muricatum* which, in my opinion, are quite unrelated. *Philodendron muricatum* is the oldest name for two other synonyms, *P. jenmanii* K. Krause and *P. scabrum*, which were placed by Krause in *P.* subsect. *Solenosterigma*. Though I am doubtful of the affinity between *P. hederaceum* and *P. muricatum*, it seems best to include *P. muricatum* in subsect. *Solenosterigma* because *P. muricatum* is clearly a member of *P.* sect. *Philodendron* and not *P.* sect. *Calostigma*.

This leaves only *P. multispadicum*, which Mayo (1990) wisely chose as the type of *P.* subsect. *Oligocarpidium*. There are two Central American representatives of the subsection and a number of other undescribed South American species as well. The Central American species are *P. clewellii* and *P. heleniae*.

The subsection is distinguished by having a relatively large number of small (usually 4-10 cm long) inflorescences per axil. The plants are vines with long internodes and long-petiolate, narrowly ovate to ovate-cordate blades that have the basal veins either lacking or all free to the base and without the development of a posterior rib. Both Central American species represented have a type B style and an ovule sac surrounding the ovules.

**4. *Philodendron* subsect. *Bulaoana* Mayo** [as

"*Bulaoanum*"], J. Linn. Soc., Bot. 100: 168. 1989. TYPE: *P. bulaoanum* Engl. (lectotype, designated by Mayo, 1989).

This subsection includes two species treated by Krause, *P. bulaoanum* and *P. acuminatissimum*. They probably represent the same species, a plant with deeply 3-lobed leaves and persistent, reddish brown cataphyll fibers. Currently the subsection is known only from South America. Krause treated this subsection as *P.* "Gruppe" *Doratophyllum*, but that name has no priority at the subsectional level (Mayo, 1990).

This subsection can be easily confused with and is perhaps inseparable from *P.* sect. *Tritomophyllum*. A moderately large number of South American species, including *P. barrosoanum* G. S. Bunting, *P. cataniapoense* G. S. Bunting (Bunting placed it in *P.* "Gruppe" *Doratophyllum*), *P. effusilobum*

Croat, *P. holtonianum* Schott, *P. hylaeae* G. S. Bunting (*P. sect. Calostigma*), *P. levelii* G. S. Bunting (similar to *P. barrosoanum*), *P. panduriforme* Schott, and *P. victoricae* G. S. Bunting might all belong here.

*Philodendron* subsect. *Bulaoana* and *P. sect. Tritomophyllum* are separated on weak features. Krause described his "Gruppe" *Doratomyllum* as having somewhat succulent petioles longer than the blade, a hastate blade, and ovaries that are several-ovulate near the base of the locule, whereas he described *P. sect. Tritomophyllum* as having tripartite blades with the lateral lobes erect or spreading, and the primary lateral veins more prominent than the minor veins. The ovary was described as 5–11-locular with the locules 1- or more-ovulate. The Central American species that appear to be closely related have one or two ovules per locule, whereas those of *P. subsect. Bulaoana* are described as having several ovules per locule. These distinctions are pretty weak, and further study of the members of both groups is necessary to decide whether the two groups should be merged.

5. *Philodendron* subsect. *Eucardium* (Engl.) Mayo, J. Linn. Soc., Bot. 100: 168. 1989. *Philodendron* "Gruppe" *Eucardium* Engl., Bot. Jahrb. Syst. 26: 535, 542. 1899. TYPE: *P. wallisii* Regel ex Engl.

This is a subsection of dubious status, based only on *P. wallisii*. The original description characterizes the subsection as having scarcely succulent petioles that are flattened to sulcate abaxially, cordiform blades, and 5–6-locular ovaries with a few sub-basal ovules per locule. Unfortunately, *P. wallisii* is a poorly known taxon for which, so far as is known, no material is extant.

Following his treatment of *Eucardium*, Krause discussed several poorly known species, all of which lacked inflorescences and for which no proper sectional placement was possible. Among these were *P. andreanum*, which is almost certainly related to *P. gigas*, a member of *P. ser. Velutina*. Listed among these dubious species are *P. latilobum* Schott and *P. obtusilobum* Miq. The former is a synonym of *P. panduriforme*, possibly a member of *P. subsect. Bulaoana*, which it most resembles. *Philodendron obtusilobum* is a poorly known species of unknown origin, known only from a single leaf. Its affinities remain unclear, but it appears similar to *P. lindenii* Schott or *P. rubens* Schott.

Also described among this group of unassigned species were *P. gloriosum* André and *P. mamei* André. These two species, along with *P. sodiroi* Hort., appear to be closely related and probably constitute

a new section. *Philodendron pastazanum* K. Krause has similar features and probably belongs here as well. Another species known from the Amazonian lowlands of Peru and believed to be new is also in this group. All the species in this putative new section are terrestrial plants with a unique growth form for *Philodendron*. All have a short, repent, creeping stem with erect leaves clustered near the apex. The internodes are usually much broader than long and have cataphylls persistent, sometimes persisting somewhat intact. The petioles are frequently winged or undulate-winged along adaxial margins (but not *P. gloriosum*) and blades are typically quite attractive, sometimes mottled with paler green, sometimes (as in *P. gloriosum*) somewhat velvety. The group is restricted to South America chiefly in the upper Amazon region. Studies of the ovules are necessary to confirm these speculations, and investigations will be carried out as material becomes available.

5. *Philodendron* sect. *Tritomophyllum* (Schott) Engl., in Mart., Fl. bras. 3: 144. 1878. *Philodendron* *group. Tritomophyllum* Schott, Syn. Aroid. 107. 1856. *Baursia* sect. *Tritomophyllum* (Schott) T. Post & Kuntze, Lex. gen. phan. 63. 1903. TYPE: *P. tripartitum* (Jacq.) Schott (lectotype, designated by Mayo, 1990: 63).

The section is distinguished by having a scandent habit, three-lobed leaf blades and 1–2 ovules per locule. The section ranges from Mexico to South America, ranging as far as Ecuador on the Pacific slope and to Venezuela, the Guianas, and the Amazon basin. It is represented in Central America by six species: *P. angustilobum*, *P. anisotomum*, *P. cotobrusense*, *P. madronense*, *P. rothschuhianum* (Engl. & K. Krause) Croat & Grayum, and *P. tripartitum*. In South America there are a number of species with 3-lobed leaves, including: *P. barrosoanum*, *P. cataniapoense*, *P. effusilobum*, *P. holtonianum*, *P. hylaeae*, *P. levelii*, *P. panduriforme*, and *P. victoricae*, all now tentatively placed in *P. subsect. Bulaoana*, which might belong in *P. sect. Tritomophyllum*. Certainly *P. hylaeae*, with 1–2 ovules per locule and a strong similarity with *P. tripartitum*, would appear to be closely related to *P. sect. Tritomophyllum*.

Bunting (1986) placed *P. cataniapoense* in *P. sect. Oligospermium* "Gruppe" *Doratomyllum* (now *P. subsect. Bulaoana*) but this species has a single ovule per locule, elongate internodes, and deciduous cataphylls. It may properly belong with *P. sect. Tritomophyllum*.

Some of the above-mentioned South American species are doubtfully included here, since they

have several (more than 1 or 2) ovules per locule. *Philodendron levelii* has 2-4 sub-basal ovules, and *P. barrosoanum* has 4-6 ovules attached above the base. *Philodendron victoriae* is apparently closely related to *P. barrosoanum* so it probably has similar ovules. It is possible that none of these species belong to *P. sect. Tritomophyllum*, but on the other hand they do not closely match *P. bulaoanum*, the type of *P. subsect. Bulaoana*, either.

6. **Philodendron sect. Schizophyllum** (Schott) Engl., in Mart., Fl. bras. 3: 144. 1878. *Philodendron grex Schizophyllum* Schott, Syn. Aroid. 104. 1856. TYPE: *P. pedatum* (Hook.) Kunth (lectotype, designated by Mayo, 1990: 63).

The section is a small but natural group of 6-7 scandent species with irregularly pedatisect leaves, 3-4 ovules per locule, and sometimes scaly petioles, primarily restricted to eastern South America and the Amazon basin, but with one species, *P. pedatum*, more widespread and ranging to northwestern Colombia. The species may occur in Central America, according to horticulturist John Hall of Costa Rica, who illustrated such a plant supposedly from the Osa Peninsula. Characteristically, no material was collected and this claim cannot be refuted.

7. **Philodendron sect. Polytomium** (Schott) Engl., in Mart., Fl. bras. 3: 145. 1878. *Philodendron grex Polytomium* Schott, Syn. Aroid. 108. 1856. TYPE: *P. radiatum* Schott (lectotype, designated by Mayo, 1990: 63).

The section is a small group of 7-8 species with pinnately or bipinnately lobed leaves from the West Indies, Central America, and northern South America. Plants are vines or appressed climbers with more or less terete petioles, generally deeply lobed, large, moderately coriaceous blades, and flowers with several axillary or sub-basal ovules per locule. Only one species, *P. distantilobum* K. Krause, was reported for the Amazon basin, but another species, *P. pinnatifidum*, placed erroneously, I believe, in *P. sect. Macrogygium*, really belongs here as well. It is also a species occurring in the upper Amazon basin. Two additional species, *P. angustisectum* Engl. and *P. elegans* K. Krause, occur in Colombia. *Philodendron fendleri* occurs in Trinidad and northern Venezuela, and *P. lacerum* (Jacq.) Schott occurs in the Greater Antilles (Cuba, Jamaica, and Hispaniola). One poorly known species, *P. houlettianum* Engl., has been reported from French Guiana but no material exists to confirm what it is. The most widespread species in the section, *P. radiatum*, occurs in Central America, ranging from Mex-

ico (San Luis Potosí) to Colombia (Antioquia). *Philodendron radiatum* var. *pseudoradiatum* is endemic to the State of Chiapas in Mexico. Central American species in *P. sect. Polytomium* are *P. warszewiczii*, ranging from Mexico to Nicaragua, and *P. dressleri*, which is endemic to Mexico.

Mayo (1986), following a cladistic survey of inflorescence types, concluded that *P. fendleri* belonged in a group with *P. melinonii* and *P. pedatum*. Based on overall morphology I would conclude that the three species are not closely related. The latter is, in my opinion, a member of a distinctive section, *P. sect. Schizophyllum*. *Philodendron melinonii* Brongn., a distinctive species with cordate blades and placed by Krause in *P. sect. Macrolonchium*, is in my opinion, not related to either of the sections with lobed leaves.

8. **Philodendron sect. Macrogygium** Engl., Bot. Jahrb. Syst. 26: 553. 1899. TYPE: *P. hoffmannii* Schott.

*Philodendron sect. Macrogygium* is one of two small sections treated by Krause (1913) (the other being *P. sect. Camptogygium*). *Philodendron sect. Macrogygium* consists of a single species, *P. jacquinii* (treated by Krause as *P. hoffmannii*). Both sections are represented by scandent species, distinguished by having the style prolonged and much narrower than the ovary and a single ovule per locule. Despite these similarities the two sections are very different from one another. *Philodendron jacquinii* (*P. sect. Macrogygium*) has setose stems and thin, veiny, ovate-cordate blades which may be deciduous during the dry season. Its spathe is bulbous and roomy inside, quite unlike most species. The style, though narrowed to the apex, has a typical, hemispheroid stigma. For differences with *P. sect. Camptogygium*, see below.

*Philodendron brevispathum*, with its similarity to *P. jacquinii*, might be considered a relative, but the former has branched scales, not simple trichomes on stems and petioles; a normal, sessile style; and 6-14 ovules per locule with axile placentation, rather than 2 ovules per locule and sub-basal placentation for *P. jacquinii*.

This section appears, at least on the surface, to be natural. No other species known is alike either morphologically or ecologically. Its thin, veiny leaves are deciduous in the dry season and the large colorful infructescences are prominently displayed in a mostly deciduous environment.

9. **Philodendron sect. Camptogygium** K. Krause in Engl., Das Pflanzenreich IV. 23Db (Heft 60): 3, 127. 1913. TYPE: *P. longistilum* K. Krause

In contrast to *P. jacquinii*, which has setose stems and thin cordate blades with prominent veins, *P. longistilum* (*P. sect. Campotogynium*) has glabrous stems and oblanceolate, subcoriaceous blades with obscure primary lateral veins. The chief reason for its status as the only member of a section is the peculiar prolonged style, which is deflected to one side and has a cupular apex. It is perhaps not as unique as Krause assumed. Other species are now

known to have prolonged styles deflected to one side, among them two undescribed species with cordate blades from western Ecuador, one based on *Camp 3701* and the other on *Jaramillo et al. 25449*. However, neither has a cupular style apex. These perhaps represent another section, or more likely Krause's *P. sect. Campotogynium* may have to be incorporated in *P. sect. Calostigma*, where it might be easily accommodated.

KEY TO THE SECTIONS AND SUBSECTIONS OF *P.* SUBG. *PHILODENDRON*

- 1a. Pistils usually 2-locular, rarely 3-locular; eastern South America and Amazon basin ..... *P. sect. Philopsamma*  
 1b. Pistils 4-10-locular; throughout the range of the genus.  
 2a. Plants with primary lateral leaf veins moderately obscure; all South American (excludes *P. wendlandii*, which is transferred to *P. sect. Calostigma*) ..... *P. sect. Baarsia*  
 2b. Plants with primary lateral leaf veins usually much more prominent than the minor veins.  
 3a. Pistil with style much narrowed and much prolonged beyond the body of the ovary and much narrower than ovary.  
 4a. Style turned toward apex, perpendicular to the body of the ovary; South American species only ..... *P. sect. Campotogynium*  
 4b. Style straight, directed in the same axis as the ovary; Central and South America ..... *P. sect. Macrogynium (P. jacquinii)*  
 3b. Pistil with style about as broad as the ovary, scarcely prolonged beyond body of ovary.  
 5a. Blades 3-lobed or deeply incised-lobate.  
 6a. Blades 3-lobed; ovules 1-3 per locule, basal or sub-basal. [Note: *P. subsect. Doratophyllum* with 3-lobed blades would also key here. See the discussion after that subsection.] .....  
     ..... *P. sect. Tritomophyllum (P. angustilobum (perhaps this belongs in P. subsect. Macrobolium (Schott) Engl.), P. anisotomum, P. cotobriense, P. madronense, P. rothschuhianum, P. tripartitum)*  
 6b. Blades incised-lobate; ovules 2-8 per locule, sub-basal or axile.  
 7a. Blades divided along the anterior lobe into two or more divisions, each division in turn terminating with much smaller lobes; petioles sometimes with trichome-like scales; South America only ..... *P. sect. Schizophyllum*  
 7b. Blades pinnately or bipinnately lobed, the divisions  $\pm$  uniform along the anterior lobe; petiole never scaly; Central and South American species .....  
     ..... *P. sect. Polytonium (P. dressleri, P. radiatum, P. warzewiczii)*  
 5b. Blades entire, sometimes with the anterior lobes markedly concave, but the blade not markedly 3-lobed or incised-lobate.  
 8a. Pistils with axile placentation; ovules usually 15 or more per locule, seldom with as few as 10 per locule, rarely fewer ..... *P. sect. Philodendron*  
 9a. Blades cordate to subcordate or sagittate at base; petioles various.  
 10a. Petioles  $\pm$  D-shaped in cross section, sharply flattened adaxially, sometimes also with the lateral margins raised.  
 11a. Stems with internodes much broader than long .....  
     ..... *P. subsect. Macrolonchium (Schott) Engl. (P. fragrantissimum)*  
 11b. Stems with internodes about as broad as long or longer than broad .....  
     ..... *P. subsect. Platypodium (Schott) Engl. (P. brunneicaule, P. copense, P. fendens, P. fortuneense, P. pterotum)*  
 10b. Petioles terete or subterete.  
 12a. Petioles warty at apex; South American species (excluding *P. grandipes* erroneously placed here by Krause); *P. ornatum* Schott and relatives .....  
     ..... *P. subsect. Psoropodium (Schott) Engl.*  
 12b. Petioles smooth or conspicuously scaly but not merely warty at apex.  
 13a. Plants scandent; internodes much longer than broad; cataphylls usually deciduous; blades cordate ..... *P. subsect. Solenasterigma (Klotzsch ex Schott) Engl. (P. bresipathum, P. purpleovoides, P. hederaceum)*  
 13b. Plants usually not scandent, either terrestrial or appressed-climbing with internodes broader than long or not much longer than broad; cataphylls usually persistent, rarely deciduous; blades ovate to sagittate.  
 14a. Plants with petioles scaly ..... *P. subsect. Achyropodium (Schott) Engl. (P. glanduliferum, P. hammelii, P. malesevichiae, P. squamipetiolatum, P. squamicaule, P. verrucosum)*  
 14b. Plants with petioles smooth, not scaly ..... *P. subsect. Philodendron*  
 15a. Cataphylls deciduous intact or sometimes becoming fibrous but soon deciduous ..... *P. sec. Philodendron Schott (P. giganteum (not in Central America))*  
 15b. Cataphylls persisting as a mass of fibers.

- 16a. Blades matte on lower surface.  
17a. Blades matte, not velvety on upper surface, whitish and glaucous on lower surface .....  
*P. ser. Impolita* Croat [*P. hebetatum*, *P. strictum*,  
*P. thalassicum*]  
17b. Blades velvety on upper surface, not whitish and glaucous on lower surface .....  
*P. ser. Velutina* Croat [*P. gigas*]
- 16b. Blades usually semiglossy, never whitish or glaucous, on lower surface.  
18a. Sap brown to clear, never chalky .....  
*P. ser. Fibrosa* Croat [*P. alticola*,  
*P. antonioanum*, *P. breudlowei*, *P. chiriquense*, *P. dodonii*, *P. grandipes*, *P. jordanianum*, *P. lazarii*,  
*P. llanense*, *P. panamense*, *P. pirrense*, *P. purulhense*, *P. scalarinerve*, *P. schottianum*, *P. tenue*]  
18b. Sap white and chalky .....  
*P. ser. Albisuccosa* Croat [*P. albisuccus*]
- 9b. Blades acute to obtuse at base; petioles subterete and smooth .....  
*P. subsect. Canniphyllum* (Schott) Mayo [*P. chiripoense*, *P. cretatum*, *P. roseospathum*]
- 8b. Pistils with basal or sub-basal placentation; ovules few per locule, usually fewer than 5, seldom up to 8, rarely as many as 12 (but with some locules in the same inflorescence with as few as 6 ovules per locule) .....  
*P. sect. Calastigma*
- 19a. Plants usually appressed hemiepiphytic climbers; internodes frequently as long as wide or longer than wide; widespread in both Central and South America.  
20a. Blades deeply 3-lobed; cataphylls persisting in a dense, reddish brown layer of fibers; South American species only; *P. acuminatissimum* Engl., *P. bulaoanum* Engl .....  
*P. subsect. Bulaoana* Mayo
- 20b. Blades not at all 3-lobed, cordate, sagittate, or  $\pm$  oblong; cataphylls deciduous or persistent, fibrous or intact; widespread in both Central and South America.  
21a. Inflorescences both small and numerous, usually more than 6 per axil, less than 5 cm long .....  
*P. subsect. Oligocarpidium* (Engl.) Mayo [*P. clewellii*, *P. helenaiae*]
- 21b. Inflorescences of normal size, typically 1-few and more than 5 cm long.  
22a. Pistils with several ovules per locule; petioles rarely with purple ring at apex; blades sagittate or cordate at base .....  
*P. subsect. Macrobelyum*
- 23a. Cataphylls deciduous.  
24a. Leaf blades with minor veins not at all etched into upper surface .....  
*P. ser. Macrobelyum* (Schott) Croat [*P. advena*, *P. annulatum*, *P. aromaticum*, *P. coloradense*, *P. duyeri*, *P. edendatum*, *P. ferrugineum*, *P. graymii*, *P. knappiae*, *P. mexicanum*, *P. platypetiolatum*, *P. sagitti-fo-  
lium*, *P. souae*, *P. subincisum*, *P. verapazense*, *P. zhuangum*]  
24b. Leaf blades with minor veins etched into upper surface .....  
*P. ser. Ecordata* Croat [*P. brenesii*, *P. crassipathum*, *P. davidsonii*, *P. lentii*, *P. niqueanum*]
- 23b. Cataphylls persistent.  
25a. Cataphylls weathering and persisting as fibers .....  
*P. ser. Reticulata* Croat [*P. jefense*, *P. tysonii*]  
25b. Cataphylls persisting intact; stems markedly succulent .....  
*P. ser. Pachycaulia* Croat [*P. bosii*]
- 22b. Pistils with ovules solitary (or sometimes 2, as in *P. wilburii*) in each locule; petioles frequently with purple ring at apex (*P. subsect. Glossophyllum*); blades elongate (*P. ser. Glossophyllum*) or  $\pm$  ovate (*P. subsect. Glossophyllum* *ser. Ovata*).  
26a. Blades mostly  $\pm$  oblong and acute to narrowly subcordate at base; petioles frequently with a purple ring at apex .....  
*P. subsect. Glossophyllum* (Schott) Engl.; *P. ser. Glossophyllum* (Schott) Croat [*P. auriculatum*, *P. bakeri*, *P. brewsterense*, *P. corvae*, *P. dolichophyllum*, *P. folsomii*, *P. granulare*, *P. inmixtum*, *P. ligulatum*, *P. morii*, *P. pseudauriculatum*, *P. ubiganupense*, *P. utleyanum*, *P. wendlandii*]
- 26b. Blades ovate to ovate-triangular, cordate to subcordate at base; petioles usually lacking a purple ring at apex .....  
*P. subsect. Glossophyllum*; *P. ser. Ovata* Croat [*P. cotonense*, *P. dominicalense*, *P. microstictum*, *P. smithii*, *P. straminicaule*, *P. sulcicaule*, *P. wilburii*]
- 19b. Plant habit unknown; internodes much broader than long; Colombia (exact locality unknown) .....  
*P. subsect. Eucardium* (Engl.) Mayo

## MORPHOLOGY OF VEGETATIVE STRUCTURES

## ANATOMY

*Vegetative anatomy.* (Section on Vegetative Anatomy contributed by R. C. Keating; vouchers will be cited by Keating (in press)).

*General comments.* *Philodendron* has secretory ducts present in all morphological parts. These occur mostly as 2-4 "files" in vascular bundles. Laticifers are of the non-anastomosing type, simple and articulated (French, 1988). The genus also has secretory ducts present in roots, stems, leaves, and inflorescences. The roots have a sclerotic hypodermis.

*Leaf surfaces.* Cuticle smooth or occasionally rough on both surfaces. *Epidermis:* adaxial cells polygonal (1-2:1), straight-sided. Abaxial cells similar to or larger than adaxial cells or more elongate (2-4:1 l/w). *Stomata:* abaxial, randomly oriented, brachyparacytic to brachyparaxacytic. *Venation:* secondary veins parallel with reticulate higher order veins.

*Leaf Cross Section.* Cuticle of medium thickness, may intrude deeply along anticlinal walls of epidermal cells. Abaxial cuticle may be thinner and smooth or striate. *Epidermal cells:* large or small, square to columnar or tile-like adaxial cells, larger abaxially, or same size; outer and inner walls flat. *Stomata* level with surface with conspicuous double cuticular flanges often present. *Hypodermis:* absent or as 1-3 layers present on adaxial side of midrib, or as isodiametric cells beneath both epidermal layers (in *P. crassispatum*, hypoderm of 1-4 layers adaxially and 2-3 layers abaxially), cells may contain dark material. *Mesophyll:* palisade 1-4 layers of elongated or rounded cells, usually packed with plastids; occupying 10-30% of lamina. Spongy layers up to 15 cells deep. *Air spaces:* large, substomatal cylinders perpendicular to surface; 30-90% space in spongy layer; or air cavities irregular in some species. Paradermally, spongy tissue with one layer of unarmed cells over abaxial epidermis, then small cells or short-armed cells surrounding large air cavities extending from stomata to palisade cells; 3-5 spongy cells per uniseriate partition. Large cavities in midrib separated by uniseriate partitions of ground tissue, partitions 3-4 cells long between junctions. *Collenchyma:* 4-5 discrete subepidermal layers of small cells on abaxial side of midrib, often extending to vascular bundles.

*Vascular bundles.* One to several large bundles in the midrib, often at corners of a net-like aeren-

chyma; well delimited, with 1-5 large metaxylem cells, next to a small, round or elliptic phloem strand. *Sclerenchyma:* fibers, usually as a phloem cap 1-3 layers deep, and less commonly as a xylem cap. Some laminar bundles ensheathed with fibers. Fibers having up to 5 discrete wall layers. Xylem cells often ringed with a single layer of small parenchyma cells. *Secretory ducts:* occasional in midlamina or midrib with 1-2 layers of epithelium. *Laticifers:* lateral to vascular bundles, 2-3 per bundle, outside fibers, often with short processes protruding between adjacent mesophyll cells; containing dark amorphous contents. *Starch:* abundant in midrib ground tissue, not seen in lamina. *Crystals:* raphides common or uncommon in single packets in short or elongate cells, ends protruding across partitions into both adjacent air spaces, or in large rounded cells of upper or lower mesophyll. Raphide packets occasionally surrounded by dark sheaths within the cell (in *P. crassispatum* crystal cells elongated parallel with vascular bundles). Druses rare or common, often coarse and sharp-pointed, in rounded cells in palisade and spongy tissue partitions. Crystal sand absent or common in most mesophyll cells.

*Petiole Cross Section.* Cuticle smooth and thin. Epidermis of small cells, level or rounded on surface and inner walls rounded to angular. Hypodermis of 2-4 layers. Collenchyma beneath hypodermis as a continuous band of 4-8 layers of angularly thickened, longitudinally elongated cells (8-10:1). *Chlorenchyma:* up to 5 rows of cells inside collenchyma, and in addition, in *P. jodavianum*, as perpendicular partitions dividing the collenchyma band, and reaching subepidermal layers. Ground tissue of various sizes of parenchyma cells among numerous small to medium-sized air cavities separated by uniseriate partitions of up to 3 cells long. Cells often longitudinally elongated (3-6:1 l/w). Vascular bundles numerous and scattered, mostly oriented with phloem facing perimeter. Bundles are parallel in longitudinal section with some higher order angular connections. Xylem of 1-3 protoxylem cells per bundle, with spiral to annular thickenings, and occasionally one metaxylem element. Phloem of sieve cells and companion cells forming organized grid. In longitudinal section, sieve elements with horizontal to 45° oblique end walls. *Sclerenchyma:* fibers forming phloem cap, 2-4 cells deep, usually not fully surrounding phloem laterally. In some species, fiber caps surrounding xylem and phloem or ensheathing entire bundle.

Secretory ducts surrounded by epithelium and 1-2 rows of additional small cells, very common in



collenchyma and common in ground tissue. *Laticifers*: 1-3, found laterally on one or both flanks of vascular bundles; non-articulated. Starch common throughout ground tissue, rare in collenchyma. *Crystals*: narrow raphide cells occasional, oriented across aerenchyma partitions with ends pointing into air cavities, or cells randomly oriented. Druses present subepidermally and in cells bordering air cavities. Small groups of small prisms common in ground tissue.

**Stem Cross Section.** 1-3 cm diam. Cuticle of medium thickness, smooth to rough-surfaced. Epidermis of very small cells, outer walls rounded. *Cork*: cells may be present subepidermally. *Hypodermis*: a subepidermal layer of large cells, over 1-2 layers of thick-walled fibers, followed by thin-walled periderm cells. *Collenchyma*: angularly-thickened cells, 3-10 layers deep, beneath periderm. Cortical ground tissue of loosely packed parenchyma. *Air cavities*: cell-sized or smaller in all ground tissue. *Central cylinder*: weakly delimited by a circle of vascular bundles, some of which tangentially fused in pairs or threes by phloem fibers and confluent phloem strands. Cortical bundles randomly scattered. Vascular bundles collateral and highly variable in form and orientation. Compound bundles numerous in central cylinder with irregular fiber strands in center of groups of 2-6 bundles. Small phloem strands outside fibers facing groups of 1-2 metaxylem elements on outer side. Simple collateral bundles may have a large wide strand of fibers surrounding phloem, a very small irregularly shaped strand of fibers, or no fibers. Fibers having up to 7 discrete wall layers visible.

Secretory ducts small and numerous in collenchyma and outer cortex, less common elsewhere. Duct cavities surrounded by two modified layers of small cells; contents of cavities testing positively for lipids. *Laticifers*: absent or present as 1-3 cells lateral to many vascular bundles, occasional in phloem. *Crystals*: raphides as single packets in elongate cells, most common in central cylinder. Druses of all sizes common in small cells in cortex.

**Vascular anatomy.** The configuration of shoot vasculature of *Philodendron* has been extensively studied by French and Tomlinson (1980, 1981, 1984). They reported the genus to be one of the most diverse in the family from an anatomical point of view. The axis of the vascular system of *Philodendron* is continuous throughout the stem because the renewal shoots develop precociously and because the morphologically terminal parts of the stem soon become branches (French & Tomlinson, 1984). All species of *Philodendron* examined by

French and Tomlinson have an "independent cortical vascular system of traces that pass from the leaf into the cortex, but without entering the central cylinder." Generally the cortex is wide with 3-5 or more indistinct series of bundles. Cortical bundles are collateral, typically with a fibrous sheath next to the phloem. Major leaf traces enter the cortex at an acute angle and promptly enter the central cylinder, while smaller traces enter the cortex at a less acute angle and may enter the central cylinder well below where they entered the cortex. No endodermis was observed by French and Tomlinson for any of the species.

*Philodendron* subg. *Philodendron* has secretory resin canals occurring in the cortex. These consist of a schizogenous space lined with 2-3 layers of epithelial cells and contain a Sudan IV-staining resin (French & Tomlinson, 1984). The central cylinder is separated from the cortex by an arrangement of vascular bundles into which the leaf traces merge. Bud traces are equal on both sides and form an arc near the periphery of the central cylinder before joining with the axial bundles.

In their survey of 3-dimensional arrangement of vascular bundles, leaf traces and axial bundles were distinct, with the leaf traces consistently collateral with protoxylem and usually with a prominent sheath of sclerenchyma next to the phloem. The axial bundles were divided roughly into five groups, four of which pertain to *P.* subg. *Philodendron*.

In the *P. hederaceum* pattern (French & Tomlinson pattern 1) there are simple collateral bundles with some bipolar bundles in the central cylinder of the internodes with thin-walled and non-lignified ground tissue.

Another relatively rare pattern involving *P.* sect. *Baurisia* (but not a Central American representative of the section) and *P. jacquinii* (pattern 3 of French & Tomlinson) has compound bundles throughout the central cylinder with individual compound bundles consisting of strands of xylem and phloem separated from each other by sclerenchyma "in the form of a partial or complete sheath" (French & Tomlinson, 1984). In *P. jacquinii* leaf traces enter the central cylinder but make a variety of configurations, sometimes including pairing before joining the compound bundles. Compound bundles do not make a particularly straight course, sometimes bundling and pairing within and between compound bundles.

A pattern of vascular bundle traces exhibited by *P. fragrantissimum* and *P. roseospathum* (French & Tomlinson pattern 4) is similar to pattern 3 except that "the pattern of the vascular components with

compound bundles is less clear because components are not separated by sclerenchyma" (French & Tomlinson, 1984). The sclerenchyma occupies the central core but does not isolate individual bundle components.

The pattern for *P. mexicanum* and *P. sagittifolium* (French & Tomlinson pattern 5), described as the most common in the genus, has axial bundles strictly amphivasal with a central core of phloem and a peripheral region of xylem. The tracheary elements may form a more or less continuous cylinder or may be arranged in clusters around the phloem. Leaf traces penetrate the central bundle and fuse with the central cylinder, while the sclerenchyma sheath of the trace migrates to the center of the axial bundle.

**Habit and growth patterns.** In terms of growth habit *Philodendron* is clearly one of the most variable genera in the Araceae (Blanc, 1977a, 1977b, 1978, 1980; French & Tomlinson, 1981). The habit ranges from terrestrial to epiphytic or hemiepiphytic (primary or secondary). Secondary hemiepiphytes may be vines or appressed-climbers.

Relatively few *Philodendron* species are terrestrial, although a few are consistently terrestrial. These include *P. glanduliferum*, *P. grandipes*, *P. malesevichiae*, and perhaps *P. hammelii* (known only from a single collection). The stem of *P. glanduliferum* (Fig. 198) is usually repent. Other species are mostly terrestrial, but sometimes hemiepiphytic: e.g., *P. basii*, *P. roseospathum* var. *roseospathum*, and *P. warszewiczii*. *Philodendron knappiae* is about equally terrestrial or hemiepiphytic. Label data regarding habit on herbarium collections are often suspect, with many herbarium labels using the term "epiphyte" when in fact the collections were probably rooted in the soil and were therefore technically hemiepiphytic. In addition, although many species have collections reporting them to be terrestrial, the majority were probably found on the ground as a result of accidentally falling from trees. Many collections are naturally made in virgin forest areas where man-made disturbances such as road building give access to the forest. In such areas, and especially in the regrowth along road cuts, aroids are very common. Many *Philodendron* species, e.g., *P. lentii* (Fig. 261) and *P. squamicaule* (Fig. 382), persist in excellent condition on steep roadbanks because the steep clay slopes allow adequate drainage for the plants to survive. Still, these are somewhat unusual situations. On the other hand, stream banks often provide similar habitats.

Many members of *P. subg. Philodendron* are

hemiepiphytic, meaning that they are growing on trees as appressed climbers or as vines, while being rooted in the soil. There are two types of hemiepiphytes (Putz & Holbrook, 1986). Primary hemiepiphytes begin their lives when seeds germinate on the host tree, in the same way as true epiphytes, then go on to develop slowly until they reach sufficient age and size to begin developing long aerial roots that may eventually reach the ground. This growth strategy enables the species to quickly attain a height where light is adequate but nutrients are more difficult to obtain. Not surprisingly, one of the most successful species with this life form is *P. megalophyllum* Schott, which often lives on ant nests from which it obtains extra nutrients. No Central American species are known to live exclusively with associated ants, although occasional ant nests occur among the cluster of roots of many species. The roots hold the otherwise fragile ant nest to prevent it from being washed away in the rain. Examples of this type of hemiepiphyte are *P. goeldii* G. Barroso, *P. solimoense* A. C. Sm., *P. megalophyllum* (all from South America), and *P. radiatum* (Fig. 1), the only species known with this growth form in Central America.

Secondary hemiepiphytes (Putz & Holbrook, 1986) start their lives on the ground or on tree trunks near the ground (where they may send roots to the ground) and climb trees where they become adults and may lose their connection with the ground. Most *Philodendron* sect. *Philodendron* are members of this group.

As is the case with most Araceae, adult plants of *P. subg. Philodendron* have homeophyllous monopodial growth<sup>11</sup> (monophyllous sympodial of Ritterbusch, 1971) with each article bearing a cataphyll and a leaf. When the plant is mature, each article is terminated by an inflorescence (Ray, 1986, 1987b) or its aborted remains (Engler, 1877; Ritterbusch, 1971). The branching pattern for *P. subg. Philodendron* has the prophyll following each internode suppressed, with a single foliage leaf followed by the inflorescences and with the internode subtending the prophyll of the elongation shoot being suppressed. Growth patterns for *P. subg. Philodendron* were diagrammed schematically by Engler

<sup>11</sup> "In monopodial growth, flowering occurs on axillary short shoots and does not interrupt the formation of the main shoot by a single apex. In monophyllous sympodial growth, after the shoot terminates in an inflorescence (which often aborts at the primordial stage), each continuation shoot will produce only a single foliage leaf before terminating in another inflorescence, and being replaced by another continuation shoot with one foliage leaf" (Ray, 1987b).

(1877) and Ray (1988) and discussed in detail by French and Tomlinson (1981). In contrast to adult plants of *P. subg. Philodendron*, the juvenile growth phase (with a few exceptions) has monopodial (actually anisophyllous sympodial) growth with an article bearing an indeterminate number of foliage leaves before terminating in the first (likely aborted) inflorescence, marking the transition into homeophyllous sympodial growth where each article has a fixed number of leaves and is terminated by an inflorescence or its aborted remains. This transition does not mean that stem has reached maturity. It has only made a transition from monopodial (actually anisophyllous sympodial) growth to homeophyllous sympodial growth. At this point, the plant is probably still not mature, and while every article will end in an inflorescence, these will all be aborted. Later, after further thickening of the stem, the shoot will become mature, and this will be indicated by the fact that the inflorescences do not always abort (T. Ray, pers. comm.).

In contrast to *P. subg. Philodendron*, *P. subg. Pteromischum* has monopodial growth only up until the time of flowering when it is followed by anisophyllous sympodial growth, wherein the stem articles bear a variable number of leaves before producing another inflorescence. Each new branch forms from a bud in the "axil of the penultimate leaf of the previous unit" (French & Tomlinson, 1981). Both *P. subg. Pteromischum* and *P. subg. Philodendron* have development of the stem beyond each terminal inflorescence. Thus although *P. subg. Philodendron* appears to have an unbranched stem with an inflorescence in each leaf axil it is really producing a new branch after producing each leaf. Close examination will show that a bud for the renewal shoot occurs on each article just below the point of overlap of the sheath edges of the cataphyll (prophyll of Ray, 1987a). It is from this point that the new branch will form. There is also a second, supernumerary bud which lies below (proximal to) the bud for the renewal shoot. This acts as a reserve meristem, lying dormant unless the stem is severed just distally to it (French & Tomlinson, 1984).

Unlike *P. subg. Philodendron*, *P. subg. Pteromischum* typically does not branch after the production of each new leaf and therefore lacks comparable branch buds. The inflorescences are never produced terminally on determinate lateral branches (Grayum, 1996).

Internode length and width are altered markedly as the plant climbs, with the internodes getting ever shorter and thicker. Ray (1986) has shown a direct correlation between the length and width of an article (one segment of the stem) demonstrating that

the length-width ratio is fixed and varies according to a set pattern. Ray (1987b) categorized four different types of stem segments based on where the leaf was attached to the stem. In *P. subg. Philodendron* all species had the petiole attached to the lower end of the stem segments, and these are referred to as "hypophyllous segments." Most species of *P. subg. Pteromischum* have "hyperphyllous segments" where the petiole is attached to the upper end of the stem segment. "Ambiphylous segments," where the stem segment is so short that the petiole is attached across the entire segment, is also known in *P. subg. Pteromischum* and is also the most common type elsewhere in the family. The fourth type of sympodial stem segment, referred to as "peraphyllous segments" and presently known in the family only in *P. subg. Pteromischum*, has the stem segments elongated and extended below the point of attachment of the cataphyll.

Most members of *P. subg. Philodendron* have leaves that turn from juvenile to adult gradually as the plant climbs so that there is no abrupt transition to adult foliage.

Although not as pronounced as in *Monstera* or *Syngonium*, leaf dimorphism is present in some species of *Philodendron*. For example, in *P. hederaceum* var. *hederaceum* juvenile leaves have short petioles tightly appressed to the tree, more or less like the "shingle" leaves of *Monstera*. They are also dark blackish green and velvety above and often somewhat purplish violet beneath. The adult plants have spreading leaves with longer petioles and smooth, semiglossy blades. The velvety blades seem to be associated only with the earliest growth. Once the plant grows up high enough on the tree the leaves become smooth and semiglossy. Further growth, even when it represents a reversion to smaller-leaved forms, results in the same smooth, semiglossy texture as that of the adult.

*Philodendron hederaceum* and other scandent species tend to climb high up the trunk of the tree then spread into the canopy and finally often hang down from branches before they flower. *Philodendron jacquini* often has a similar habit. Some scandent species, such as *P. brevispatum* and *P. sulcicaule*, typically sprawl over lower vegetation rather than high in the canopy.

The amount of internode elongation varies immensely in some vining species, such as *P. immixtum*, *P. hederaceum*, and *P. sulcicaule*, with internodes 10–20 cm long (even longer on plants that are juvenile or have reverted to a "searching mode" as the result of dislodgement from the tree). Usually internode length varies considerably depending on the light and nutrients available for growth. Even

fully adult plants with very short internodes can be induced to produce longer internodes in cases where the plant is accidentally displaced from its growing situation or if it becomes heavily shaded.

Some species have evolved the ability to produce flagellate branches in an attempt to reposition themselves. These have long slender internodes and reduced leaves. At first the flagelliform growth spreads laterally, but if no other growth support is in the immediate vicinity the branch inclines toward the ground where it may creep across the surface and ascend another tree. Blanc (1980) referred to these as "flagelles" or "stolons." Ray (1987b) indicated that these flagellate branches are usually developed when a plant overgrows its support or is accidentally dislodged, but in some species, e.g., *P. fragrantissimum*, the flagellate branches may develop from a normally growing plant (Croat, 1978). *Philodendron linnæi* Kunth, a South American species, has developed this method of locomotion to an extreme. That species produces a series of rosulate clusters of leaves all interconnected by more slender flagellate stems, which ascend the tree trunk. After a period of rapid growth resulting in long, narrow internodes with the leaves reduced to small scales, the plant produces a series of short thick internodes, each of which is associated with an increasingly larger leaf, and leaves arranged in a tight rosette (Ray, 1987b). Ray (pers. comm.) reports that "with only a few exceptions, flagellar shoots do not occur in species with homeophyllous growth. Flagellar shoots are almost universal among climbing species with anisophyllous or intermittent homeophyllous growth, but almost totally lacking among climbing species with homeophyllous growth."

**Stems.** Stems vary considerably in length depending on the nature of growth. Vines, such as *P. hederaeum*, may produce stems well over 30 m long, whereas the more slow-growing appressed climbing hemiepiphytes rarely have stems more than about 2 m long. The youngest part of the stem bears most of the leaves and has the newest root growth including the most active feeder roots. Older portions of the stem have thick, strong roots usually tightly fastened to the tree, anchoring it in place. Some roots all along the stem extend downward along the stem and may lead all the way to the ground. The older portion of the stem is often devoid of any leaves and usually lacks even cataphyll fibers. The bare stem clearly shows the petiolar and cataphyll scars (Figs. 9–12). See discussion below. Inspection usually shows that the older portions of the stem have been at some time attacked by root borers so

there may be surface damage or more likely active root borers in the center of the stem. Eventually the lower part of the stem rots away as the stem climbs higher on the side of the tree.

Stems of *Philodendron* are typically unbranched, but may be branched naturally, as in many species of *P.* subg. *Pteromischum*, or through injury. Commonly internodes become increasingly shorter and thicker toward the apex of the stem on appressed-climbing plants, but some species, e.g., *P. fragrantissimum*, may from time to time begin to produce long slender internodes to enable the plant to enter a colonizing mode (see above section on "Habit and Growth Patterns").

Mayo (1991) described differences in the branching pattern of *P.* subg. *Philodendron* and *P.* subg. *Meconostigma*. While most *P.* subg. *Meconostigma* have very short internodes, some species, e.g., *P. leaf-costae* Mayo & Barroso and *P. corcovadense* Kunth, have elongated internodes. In these species the elongated internode of each article is the one between the cataphyll (prophyll of Ray, 1987a) and the succeeding foliage leaf, while the internode preceding the cataphyll is abbreviated. In *P.* subg. *Philodendron* the elongated internode is the hypopodial internode (which precedes the cataphyll), while the internode between the cataphyll and the foliage leaf is the abbreviated one.

Stems of *Philodendron* are typically rich in taxonomic characters, and together with their associated cataphylls, yield some of the best key characters for identification. Normally these stem characteristics are useful only at the specific level, but Grayum (1996) has found the dried stem color to be useful in separating the two sections of *P.* subg. *Pteromischum*. Unlike most *Anthurium* species, which have internodes so short that the epidermal features are for the most part obscure or very uniform, *Philodendron* has features important both before and after drying. The internodes of most appressed-climbing hemiepiphytes are actually broader than long at maturity (Figs. 10, 394, 443), but some species, such as *P. advena* (Fig. 38) and *P. stramineicaule* (Fig. 392), have internodes typically longer than broad (though short enough that the plants are not considered scandent). Only nine species in Central America have internodes longer than 25 cm on adult or preadult plants. Alternatively, many species have short internodes, with 33 Central American species having internodes to only 5 cm long or less.

Internode width is somewhat less variable, ranging from less than 3 mm diam. in *P. chirripense* to 10 cm diam. in *P. gigas*. Only nine additional species (*P. chiriquense*, *P. copense*, *P. dressleri*, *P. fer-*

*rugineum*, *P. fortuneense*, *P. grayumii*, *P. pterotum*, *P. schottianum*, and *P. warszewiczii*) have stems that attain diameters greater than 7 cm. Most species have relatively narrow internodes, with 44 species having internodes of no more than 3 cm diam., and 16 species with internodes of less than 1.5 cm diam.

Fresh stems often have characteristic surface features. Surfaces are frequently speckled with a lighter or darker pattern of green. The surface may also be short-lineate (Fig. 2) or may present a combination of short lines and speckles (Fig. 3). At other times the surface may be weakly to prominently striate as in *P. anisotomum*, *P. hederaceum*, and *P. wilburii*. The stems may be glaucous as sometimes in *P. dodsonii*, *P. fortuneense*, *P. immixtum*, *P. ligulatum*, *P. mexicanum*, and *P. wendlandii*. Frequently the internodes are coarsely short-costate (Fig. 2) near the apex of the internode, as in *P. chiriquense*, *P. grandipes*, and *P. schottianum*. These apparently represent areas at which roots will emerge later.

Stems are commonly smooth but may be minutely wrinkled, as in *P. verrucosum*, or sparsely warty, as in *P. brunneicaule*, or setose, as in *P. jacquinii*, or with often branched, trichome-like scales (Fig. 5), as in *P. brevispathum*. Other species with scaly stems are: *P. hammelii*, *P. malesevichiae*, *P. squamicaule*, *P. squamipetiolatum*, and *P. verrucosum*. Even stems of species with smooth epidermis are often weakly fissured in fresh condition (Fig. 2).

Coloration of fresh stems may also be characteristic; typically, they are medium to dark green when fresh, turning gray-green to brownish, yellow-brown or reddish brown in age. Often several transitional stages are involved, e.g., medium green to gray-green, to yellowish green and finally yellowish or reddish brown. This final stem color is often the same as the color of the stem of artificially dried herbarium collections, but at other times the dried stem color of the herbarium collection is not the same. Thus it may be important to note the color of the fresh stem before drying takes place. Natural aging of the stems often causes a scurfy condition, which results from numerous close cracks (Figs. 4, 5). Sometimes these minute fissures are restricted to a specific point of stress yet not visible elsewhere (Fig. 5).

Another important feature is the extent to which the stem wrinkles or cracks or is otherwise distorted by the drying process. The results of the drying process are usually quite consistent from collection to collection, such that the dried stem and the features it exhibits provide useful recognition characteristics.

Stems commonly dry with irregular (or sometimes regular) ribs and intervening sulcae, reflecting shrinkage of the relatively indurate outer surface of the stem compared to the rather soft, somewhat aerenchymatous stem interior. In some cases the regularity and severity of this ribbing provide especially useful characteristics, e.g., in *P. sulcicaule* the stems of which become prominently ribbed in the course of normal development, and in *P. verapazense* where the stems become regularly and conspicuously ribbed on herbarium collections.

In addition to frequently present longitudinal ribs, in e.g., *P. findens*, *P. fortuneense*, and *P. heleniae*, stems may be transversely fissured or checked with small to large fissures. These may be quite regular or irregular in severity or spacing. While sometimes characteristic, these transverse stem fissures tend to be somewhat less consistent than the longitudinal fissures. However, for some species such as *P. edendatum*, *P. ferrugineum*, *P. findens*, *P. fortuneense*, *P. ligulatum*, *P. malesevichiae*, *P. mexicanum*, and *P. wendlandii* (among others), the stems are frequently transversely fissured. Such stems are particularly noteworthy in an undescribed South American species, i.e., Croat 62785 from the Pacific coast of Colombia.

Another feature exhibited by some dried stems is an exfoliating epidermis, present in *P. angustilobum*, *P. cotonense*, *P. dodsonii*, *P. hederaceum*, *P. schottianum*, *P. smithii*, *P. stramineicaule*, *P. subincisum*, and *P. tripartitum*. In some cases the epidermis not only cracks but may begin to loosen and fall off or protrude away from the stem (Fig. 8), such as in *P. brevispathum*, *P. cotonense*, *P. dodsonii*, *P. ligulatum*, and *P. purpureoviride*. Occasionally this feature is exhibited on fresh stems as well. This is especially true if the stem is forced to bend by falling from its support, such as in *P. immixtum* and *P. sulcicaule*. The epidermis appears to be hard and brittle, while the underlying stem appears to be green and supple. In some cases the epidermis seems to be naturally shed and replaced by another epidermal layer on the fresh stem.

*Petiole scars.* Although less conspicuous in general than in *Anthurium* or in *P.* subg. *Meconostigma*, the petiole scars on the stems of *P.* subg. *Philodendron* are nonetheless clearly visible unless covered with persisting cataphylls. In contrast to *Anthurium* the petiole scars of *P.* subg. *Philodendron* are generally much less indented but rather are more or less flush with the general contour of the stem surface. They may be moderately inconspicuous as in *P. rothschuhianum* (Fig. 9) or moderately conspicuous as in *P. davidsonii* (Fig. 10).

*Philodendron* petioles are typically much swollen at the base, so the scar is usually broader than the rest of the petiole itself. Petiole scars vary from (0.5-)1 to 4(-7.5) cm high and (0.7-)1 to 5(-7) cm diam., but they are rarely more than 3 cm high and 3.5 cm diam. The average height and width for the 32 species studied is 2 cm high and 2.4 cm diam. The peduncular scar is often quite conspicuous (Fig. 5) and deep. Of ecological significance is the fact that these deep holes left when the inflorescences fall off are the points of entry for phytophagous insects, especially stem borers, which infect the older and sometimes the younger portions of the stems.

Intravaginal squamulae (Dahlgren & Clifford, 1982), so prominent in *P. subg. Meconostigma*, are usually present but often inconspicuous in *P. subg. Philodendron* (Fig. 11). Mayo (1991) stressed this as evidence of differences in two contrasting patterns of stem elongation (see discussion under stem above). In *P. subg. Philodendron* the intravaginal squamulae (Fig. 11) are always immediately above the cataphyll (cataphyll of Ray, 1987a), whereas in *P. subg. Meconostigma* the intravaginal squamulae are immediately below the cataphyll scars and often around the scar for the foliage leaf as well.

**Roots.** *Philodendron* roots have an anatomically distinct layer of exodermis beneath the epidermis, distinguished, among other things, by a long-cell/short-cell pattern (French, 1987a). "Thick-walled, pitted sclereids form a cylinder adjacent to the endodermis and similar sclereids also occur singly or in bands with suberized cork cells in the periderm of older roots" (French, 1987a). Like those of other members of tribe Philodendreae, *Philodendron* roots have a sclerotic hypodermis. French reported the sclerotic hypodermis to be distinctive because of its position next to the exodermis and its occurrence in the primary axis. Another distinctive anatomical feature of the roots of *Philodendron* is resin canals with sclerotic sheaths (French, 1987a).

All species of *Philodendron* produce adventitious roots at some or all nodes. The number of roots developed seems to have more to do with the environment than with the species involved. Plants that are appressed-climbing and in close contact with the substrate generally produce the largest number of adventitious roots. Roots may be of two types, which differ both morphologically and anatomically (Lierau, 1888; Porsch, 1911), either for anchoring the plant to the substrate or for feeding (van Tieghem, 1907; Went, 1893). The anchor roots (Fig. 54) tend to be more numerous and shorter, often with a dense layer of root hairs (they are

sometimes restricted to the side of the root that contacts the substrate). They also have a proportionately much smaller central cylinder and more mechanical tissue to give them strength (Engler & Krause, 1908) than those of feeder roots. They arise principally at the nodes but may arise all along the internodes. Anchoring roots may spread from the nodes as in *P. auriculatum* (Fig. 12) or closely appressed to the surface of the tree on which they are growing as in *P. gigas* (Fig. 13). In contrast to the anchoring roots, the feeder roots (Fig. 14) tend to be much thicker and longer and usually extend toward the ground. This behavior is to be expected since Goebel and Sandt (1930) reported that feeder roots of aroids are negatively heliotropic and positively hydrotropic. Feeder roots have a much broader central cylinder and broader vessels and sieve tubes. Feeder roots arise only at the nodes (Graham, 1990). Normally, the feeder roots are 2-4 times thicker than the anchor roots, and in *P. gigas* the feeder roots may be somewhat woody and up to 3.5 cm diam.

Some species, such as *P. auriculatum* (Fig. 12), have spine-like branch buds sparsely scattered along the length of the root, especially near the stem. Some hemiepiphytes, such as *P. solimoesense* A. C. Sm. in South America, have roots that may become markedly roughened with warty tubercles. In such cases the only portion of the root that has root hairs is the apex where the roots branch as they near the ground. It is unknown whether these roots are capable of absorbing atmospheric humidity, as is true for some *Anthurium* species, but certainly they must be able to take in the free water that runs down the root.

French (1987a) reported that in *P. subg. Philodendron* a sclerotic hypodermis is entirely absent in subterranean roots but present in the aerial roots. *Philodendron subg. Philodendron* has elongated, infrequently anastomosing resin canals that extend lengthwise through the root cortex (French, 1987c). They are lined with a layer of epithelial cells that consist of thin-walled, un lignified cells (parenchyma). In *P. subg. Philodendron* and *P. subg. Pteromischum* the epithelium is surrounded by a sheath with lignified cell walls. In contrast, *P. subg. Meconostigma* has resin canal sheaths that lack a sclerenchyma and instead have 2-5 layers of (un lignified) collenchymatous cells, which are easily distinguished from the ground tissue (French, 1987c).

While seldom used taxonomically, roots are variable to some extent from species to species. Fresh root coloration (ranging from whitish to green to brownish), length, diameter, and surface texture

Table 2. Persistent condition of cataphylls by section.

Section	Deciduous			Persisting				
	Total	% of sect.	Intact	Semi-intact	As fibers	Total	% of sect.	Section total
Calostigma	34	69%	6	6	3	15	31%	49
Macrogynium	1	100%	0	0	0	0	0%	1
Philodendron	10	24%	0	18	14	32	76%	42
Polytomium	3	100%	0	0	0	0	0%	3
Tritomophyllum	7	88%	0	1	0	1	13%	8
Total	55	53%	6	25	17	48	47%	103

(smooth, coarse, or even warty), as well as the dried color and degree to which they are fissured or folded, are all features that may be recorded. These features have not been used extensively since the roots are generally removed from the stems before the herbarium specimens are prepared.

**Cataphylls.** In the taxonomy of *Philodendron* probably no morphological feature is more diagnostic than the cataphyll (sympetiole prophyll of Ray, 1987a). *Philodendron* exhibits considerable variation and remarkable consistency in cataphyll characters. Among the most valuable characters is whether the cataphylls are deciduous or persistent. This is the single feature that makes the preparation of keys to species of *Philodendron* easier than those for *Anthurium*. In *P.* subg. *Philodendron* 56 taxa have deciduous cataphylls, while 48 have persistent cataphylls. Of the latter, 6 have cataphylls that persist intact, 25 have cataphylls that persist semi-intact, and 17 have cataphylls that persist as fibers. While a few species have cataphylls tardily deciduous and others have persistent cataphylls that eventually fall off, relatively few species are difficult to place in one or the other category. See Table 2 for a breakdown of persistent condition of cataphylls by section.

Generally speaking cataphylls are deciduous in vines and persistent on appressed-climbing plants, but there are exceptions in both groups. Cataphylls are bract-like modified leaves which function in the protection of newly emerging leaves. For vines, the cataphyll becomes functionless once the leaf has emerged and generally is promptly deciduous. Typically it recurls away from the stem on vines and eventually becomes loosened from the base and falls free, such as on *P. purpureoviride* (Fig. 320). Alternatively, on epiphytes or appressed-climbing hemiepiphytes with short internodes the cataphylls cannot easily fall free (Figs. 15, 115, 312). Although they may fall free from the stem and invariably promptly dry and usually become withered

or weather into fibers, they are often prevented from falling by the tight cluster of petioles generally found on plants with short internodes. They do regularly fall free on some species with short internodes, such as *P. warszewiczii* (Fig. 443) or *P. dresleri* (Fig. 160), but typically cataphylls persist on species with short internodes (Fig. 302, *P. pirrense*; Fig. 358, *P. scalarinerve*). Cataphylls often persist in an organized mass around the apex of the stem (Fig. 382, *P. squamicaule*). They may persist intact in *P. grandipes* (Fig. 204), *P. jodavisanum* (Fig. 248), and *P. roseospathum* (Figs. 341, 343), or have a very thin, flaky epidermis that remains intact in large pieces as in *P. dodsonii*, *P. hebetatum*, or *P. strictum* (Fig. 398); more frequently they decompose at least partially to expose a network of fibers beneath the epidermis (Fig. 258, *P. lazarii*), and even more frequently the epidermis disappears altogether (Fig. 202, *P. glanduliferum*; Fig. 253, *P. jodavisanum*; Fig. 370, *P. schottianum*). Ecologically the mass of cataphyll fibers serves a useful purpose, namely to prevent desiccation of the stem apex and especially the young roots which emerge through the moist cataphyll mass. In some species the layered cluster of cataphylls forms a sodden mass (Fig. 370, *P. schottianum*; Fig. 63, *P. antioanum*; Fig. 99, *P. chiriquense*), and in some cases the youngest cataphylls are protected by a gelatinous, mucilaginous fluid that fills the interstices of the cataphyll mass. Cataphylls are generally rigid and firm when fresh, which protects the young leaf from physical damage. Some are thick and fleshy with considerable amounts of liquid in their tissues. As these begin to decompose they may yield large amounts of watery or even gelatinous sap. This fluid may be important as a lubricant for the emerging leaf to prevent damage to the tender tissues or to prevent desiccation. Since epiphytes are often subject to harsh conditions, this is probably important to protect the growing point of the plant and the newly emergent leaves from

extremes in temperature and humidity. Although the cataphylls soon lose their moisture after they have opened, many persist semi-intact or as an organized network of fibers so they continue to add protection to the stem apex. Even after the cataphylls are reduced to a mass of fibers they retain moisture much as a pile of straw does after a rain.

A typical cataphyll is narrowly triangular, 2.3–4 times longer than wide, with the base as broad as the circumference of the stem. It may be more than 1 cm thick toward the base on larger species. Cataphyll size is somewhat proportional to leaf size on any species, ranging from (5–)10 to 50(–70) cm long and (1.5–)2.5 to 18(–31) cm wide. Since the cataphyll is affixed around the complete circumference of the stem and eventually becomes deciduous, it leaves a distinct scar on the stem. These cataphyll scars alternate with the petiolar scars and may be conspicuous (Fig. 10, *P. davidsonii* subsp. *davidsonii*) or inconspicuous (Fig. 9, *P. rothschildianum*). Fresh cataphylls are usually green, though in some cases almost white, as in *P. wilburii*; bright red, or heavily tinged red on exposed parts, such as in *P. antioquiense*, *P. niqueanum*, and *P. tysonii*; or reddish as in *P. glanduliferum*, *P. grandipes*, and *P. pirrense*. Cataphylls are often variously ribbed. In cross section they are frequently sharply D-shaped in outline but may be subterete. When subterete they may be unribbed or bluntly 1- or 2-ribbed (Fig. 60, *P. annulatum*; Fig. 16, *P. warszewiczii*; Fig. 17, *P. schottianum*) or sharply 1-ribbed. Those cataphylls which are D-shaped are often bluntly to sharply raised along the edges of the flat side so that they appear bluntly to sharply 2-ribbed, as in *P. pseudauriculatum* (Fig. 313). The ribs may be moderately low and closely spaced as in *P. panamense* (Fig. 298). In some cases the ribs are slender and knife-edged and may be as much as 1 cm high, as in *P. auriculatum* (Fig. 72) and *P. annulatum* (Fig. 60).

Useful taxonomic characters are found in the manner in which the cataphylls weather and persist. Cataphylls may be quite distinctive in the degree to which they persist intact or weather to fibers. The coloration of the cataphyll fibers is also variable, with colors ranging from almost white to tan, various shades of brown or yellow to reddish or reddish brown. Cataphylls are often short-lineate in the same manner as the petioles (Fig. 16). Some species have the network of fibers highly organized, while in other species the fibers are very disheveled, such as in *P. tenue* or *P. panamense*. In most species the epidermis of the cataphyll promptly falls off, while in some species all or parts of the epidermis persist. In *P. schottianum* the distinctive

yellowish epidermis of the cataphylls is one of the most characteristic features.

In time the lowermost cataphylls simply rot away, even for species with a large cataphyll mass, so that ultimately the oldest part of the stem is visible. By this time the stem is quite old and few visible features are worth noting.

For many species the entire cataphyll does not persist, but rather only the basal part remains. Despite the fact that the stem is not covered with a large cataphyll mass, the stem is not easily visible. In such species the cataphyll bases seem to persist longer than in those cases where the entire cataphyll eventually falls free after persisting for a time.

**Leaves.** Leaves of *P.* subg. *Philodendron*, like those of *P.* sect. *Meconostigma* and *P.* sect. *Pteromischum* (indeed, most Araceae), have supervolute vernation (Cullen, 1978). Like most other aroids, *Philodendron* leaves undergo heteroblastic development, a gradual change in morphology from juvenile to adult forms (Grayum, 1990). However, heteroblasty in *P.* subg. *Philodendron* is not so severe as in other Araceae, especially in subfamilies Lasiodeae and Monsteroideae. Typically, the juvenile blades are not dramatically different from those of the adults; commonly they are of similar shape as the adult blades, typically ovate to oblong, and virtually always lack posterior lobes. In rare cases, such as with *P. heteraceum* var. *kirkbridei* (Fig. 223), the juvenile blades are velvety, a feature that is caused by the markedly convex or even somewhat cone-shaped epidermal cells. Ecologically, this lack of glossiness has the effect of allowing nearly all light that falls on the leaf to be absorbed. In all but a few species, notably *P. gigas*, the velvety juvenile leaves are transformed to glossy or semiglossy adult leaves.

In terms of adult blade shape *P.* subg. *Philodendron* is exceedingly diverse, encompassing more morphological variation than is exhibited in *P.* subg. *Meconostigma* or *P.* subg. *Pteromischum* and indeed all the variation exhibited in the much larger genus *Anthurium*. Leaves are typically clustered at the end of the stem, and rarely more than ten are present at a time. Older leaves are eventually deciduous, falling free along with the petiole. In some rare cases the blades fall free first, followed by the somewhat more persistent petioles. The resulting petiolar scar (discussed above under stem) may not be apparent initially on species with persistent cataphylls, but even in these cases they are typically apparent on the older stems. Still, they are never so conspicuous as the petiolar scars in *P.* subg. *Meconostigma*.



In his review of leaf morphology and function Ray (1987a) divided leaves into three main types: foliage leaves, reduced leaves (10–70% the size of the foliage leaves), and cataphylls (reduced to less than 10% of a foliage leaf). The first leaf on a vegetative axis he refers to as a prophyll (in *Philodendron* always a cataphyll). This structure is usually 2-ribbed and is in this regard similar to the bracts that subtend inflorescences. The latter are referred to as bracteoles by Ray (1987a).

The leaf immediately following the prophyll is a fully developed leaf, referred to as a mesophyll (metaphyll of Grayum, 1996). All three types of leaves mentioned above may be modified by the terms proleptic and sylleptic, depending on the type of growth involved. Sympodial growth has petioles borne on the side of the stem, whereas monopodial growth has conspicuously sheathed petioles that encircle the stem. Any leaf subtending an inflorescence or an aborted inflorescence primordium is termed a sympodial leaf (Ray, 1987a). In *P.* subg. *Philodendron* all adult leaves are of this type. Juvenile leaves of *P.* subg. *Philodendron*, on the other hand, are all monopodial leaves (Grayum, 1990).

**Petioles.** Petioles of *P.* subg. *Philodendron* are usually elongated, sheathed only at the base in adult foliage and rarely with a conspicuous geniculum. Most species have blades as long as or longer than the petioles, but 63 species have blades at least sometimes shorter than the petioles. Twenty-one species have petioles reported as being as long as the blades, while 93 taxa have blades at least sometimes longer than the petioles. Petioles range in length from 4 cm in *P. bakeri* to 137 cm in *P. gigas*. Twenty-two species have petioles always less than 30 cm long, 11 less than 20 cm long, and only 4 species, namely *P. bakeri*, *P. brewsterense*, *P. chirripoense*, and *P. ubigantupense*, have petioles less than 15 cm long. Four taxa, *P. copense*, *P. radiatum* var. *radiatum*, *P. chiriquense*, and *P. tenae*, have petioles more than 1 m long.

Petioles range from less than 5 mm diam. on fresh petioles of *P. brewsterense* to ca. 7.5 cm diam. on *P. wendlandii* and *P. radiatum*.

Usually the petioles are erect-spreading from the stem with the blades either extending initially in the same plane or, more frequently, somewhat pendent from the end of the petiole.

Petiole cross section is not so variable in shape as in *Anthurium*, but the variation is important taxonomically. All too often herbarium collections make no mention of this frequently critical diagnostic feature. Typically petioles are obtusely somewhat flattened or sometimes broadly and obtusely

sulcate at the base (Fig. 212). In either event they typically become more terete toward the middle and then obtusely somewhat flattened toward the apex (Fig. 18). Frequently also the petiole has an obtuse medial rib toward the apex where it becomes somewhat flattened.

The majority of *Philodendron* species have terete or subterete petioles. Sometimes the petioles are C-shaped, that is, subterete, yet also sharply adaxially sulcate. Forty-six taxa are described as having petioles subterete in cross section. Some species have subterete petioles with the adaxial surface also narrowly and obtusely sulcate as in *P. cretosum* (Fig. 126), *P. jacquinii*, *P. roseospathum*, and sometimes in *P. sagittifolium*.

Twenty-one Central American species of *P.* subg. *Philodendron* are described as having petioles variously D-shaped in cross section. Examples of species with D-shaped petioles are: *P. copense*, *P. findens*, *P. fortunense*, *P. fragrantissimum* (Fig. 187), *P. grandipes*, *P. jodavisanum*, *P. knappiae*, *P. lentii* (Fig. 264), *P. ligulatum* var. *heracleoanum* (Figs. 275, 276), *P. thalassicum*, and *P. verapazense*. Petioles of *Philodendron advena*, *P. pterotum*, *P. immixtum*, and *P. ligulatum* var. *onatum* are also sometimes sharply flattened adaxially. Considerable variation exists in petioles described as D-shaped in cross section. Several species have petioles broadly and sharply sulcate, e.g., *P. bakeri*, *P. davidsonii*, *P. heleniae*, *P. lentii* (Fig. 264), *P. ligulatum*, *P. jodavisanum*, and *P. scalarinerve*. Rarely the adaxial surface is sharply and broadly concave and with the petiole often much broader than thick, as in *P. wendlandii* (Fig. 453). In the latter, the lateral margins may be very acute and directed outward. D-shaped petioles generally have the lateral margins weakly to prominently raised with the marginal rib either acute or obtuse. Taxa with the petiole margins acute comprise: *P. davidsonii*, *P. ligulatum* var. *heracleoanum*, and *P. wendlandii*. They are sometimes also acute on *P. chiriquense*, *P. llanense*, and *P. warszewiczii*. Sometimes, as in *P. findens*, *P. fortunense*, and *P. pterotum*, the margins are prominently winged. The wing is erect-spreading and may be markedly undulate in the area of the geniculum as in *P. fortunense* (Fig. 183) and *P. findens* (Fig. 171). Sometimes the petioles of *P. ligulatum* var. *heracleoanum* also may be weakly winged (Figs. 275, 276). Even petioles not D-shaped often have a slightly thickened, slightly raised lateral margin on the adaxial surface. Examples include *P. annulatum*, *P. cretosum*, *P. microstictum*, *P. thalassicum*, and *P. wallburii*.

Sometimes in addition to the raised margins of the petioles, there may be a prominent medial rib,

such as in *P. jodavisianum* and occasionally *P. grandipes*, giving the petiole a three-ribbed appearance adaxially. More commonly the medial rib is broad and obtuse, such as in *P. angustilobum*, *P. annulatum*, *P. copense*, *P. crassispatum*, *P. ferrugineum*, *P. ligulatum*, *P. llanense*, *P. schottianum*, *P. smithii*, *P. tenue*, *P. thalassicum*, and *P. tripartitum*.

Rarely the petioles may be markedly flattened dorso-ventrally, as in *P. platypetiolatum*. In this case the petioles are much broader than thick with the lateral margins nearly acute.

A few species have petioles markedly thicker than broad and U-shaped in cross section. Examples include *P. roseospathum*, *P. jodavisianum*, and sometimes *P. davidsonii*. Petioles of the latter species are more commonly obtusely V-shaped in cross section.

Petioles of *P.* subg. *Philodendron* typically have very short sheaths when the inflorescence is aborted and much longer sheaths (to accommodate the emerging inflorescences) when subtending an inflorescence (Figs. 38, 67, 93, 150). Generally the sheath is markedly closed with one margin overlapping the other, making it inconspicuous (Figs. 83, 160). Occasionally the petiole sheath may be conspicuous, such as in *P. correae* (Fig. 119) and some members of *P.* subsect. *Glossophyllum*.

A geniculum is typically not apparent, but the genicular area serves the same purpose as it does in *Anthurium*. It is involved with inclining or twisting the plane of the blade, presumably optimizing exposure to sunlight. Usually the genicular area is somewhat firmer than the remainder of the petiole, and sometimes it is differently colored, but it is usually neither swollen nor as conspicuously distinct as in *Anthurium*. Typically the genicular area is the same shape as the remainder of the petiole, but sometimes the cross-sectional shape is different, being more frequently bluntly ribbed and often rather deeply cracked or scurfy around the circumference (perhaps owing to the bending in response to light). The geniculum, when apparent, is sometimes thicker than the remainder of the petiole as in *P. brunneicaule*, *P. ferrugineum*, *P. heleniae*, and *P. tenue*. It may be darker than the petiole as sometimes in *P. bakeri* or *P. glanduliferum*, or slightly paler as in *P. ferrugineum* or *P. scalarinerve*.

Most species of *Philodendron*, like Araceae in general, have glabrous petioles, but there are conspicuous exceptions. Several species have conspicuously scaly petioles: *P. hammelii*, *P. glanduliferum*, *P. malesevichiae* (Fig. 283), *P. squamipetiolatum* (Fig. 387), *P. squamicaule* (Fig. 382), and *P. verrucosum*. The type of scales is also variable. In some species,

such as *P. malesevichiae*, *P. glanduliferum*, and *P. squamipetiolatum*, the scales are acicular and more or less terete. The scales of *P. glanduliferum* and *P. malesevichiae* are spreading, while those of *P. squamipetiolatum* are sometimes retrorse near the apex of the petiole. In addition, they are densely granular-scurfy on the surface. In *P. hammelii* the scales are short and broadened laterally, similar to fish scales, and less than three times longer than broad. In *P. verrucosum* and *P. squamicaule* the scales are of two distinct types. *Philodendron verrucosum* has short, broad, and often lacerate scales interspersed with long-acicular scales. *Philodendron squamicaule* has short, purplish, deltoid, broader than high (about 2 mm high) scales interspersed among acicular, greenish scales 3–5 mm long.

Petiole scales are not uniformly distributed along the petioles. In *P. squamipetiolatum*, *P. squamicaule*, and *P. verrucosum* the scales are usually scattered throughout the petioles. In *P. malesevichiae* they are present in the upper ½ to ⅓ of the petioles, becoming increasingly denser toward the base. In *P. glanduliferum* they are similar but more closely aggregated toward the apex. In *P. hammelii* the scales are restricted to a small area near the apex. *Philodendron ornatum* in South America is the most extreme example of this reduction. In that species the scales are reduced to stubby protuberances restricted to the apex of the petiole.

Surface features of both fresh and dried petioles are often conspicuous and frequently diagnostic. Though usually solid green, petioles may be irregularly purplish-spotted as in *P. edenudatum* (Fig. 163) and *P. sagittifolium*. This purplish spotting may continue onto the lower midrib, a feature common to other types of markings that continue onto at least the proximal portions of the midrib. The surfaces of petioles may be minutely speckled but more commonly bear a light pattern of short dashes or streaks usually uniformly distributed throughout the petiole. They may be densely to sparsely marked with short dashes, which in turn may be either darker or lighter than the surface. I have referred to this feature in the descriptions as "short-lineate" or "lineate." Though the lineations are usually short, usually less than 3 mm long, they may sometimes be longer and sometimes are variable in length with both short and long lines interspersed (Fig. 19). Examples of species with short-lineate petioles include: *P. annulatum*, *P. auriculatum*, *P. copense*, *P. cotonense*, *P. edenudatum*, *P. fortunense*, *P. immixtum*, *P. ligulatum*, *P. pirrense*, *P. purulhense*, *P. smithii*, *P. subincisum*, *P. thalassicum*, *P. warszewiczii*, and *P. zhuianum*. The peti-

oles of *P. brunneicaule* are red short-lineate. The short lineations may be associated as well with much longer, continuous, and frequently weakly raised striations or ribs. Species with petioles described as short-lineate to striate include: *P. antioceanum*, *P. clewellii*, *P. gigas*, *P. glanduliferum*, and *P. panamense*. Still other species have more or less continuous striations throughout: e.g., *P. anisotomum*, *P. davidsonii*, *P. dressleri*, *P. findens*, *P. madronense*, *P. rothschuhianum*, *P. stramineicaule*, *P. tenue*, and *P. warszewiczii*.

Fresh petioles of *P. subsect. Glossophyllum* usually have a distinct purplish (or rarely dark green) ring (Fig. 20) around the entire petiole just below its junction with the blade. The species that share this feature, mostly members of *P. subsect. Glossophyllum*, include *P. annulatum* (Fig. 59), *P. bakeri*, *P. correae*, and *P. ligulatum* (all three varieties; see Figs. 271, 275) with a purplish or purple-black ring, and *P. auriculatum* (Fig. 20), *P. immixtum*, *P. dolichophyllum*, *P. pseudauriculatum*, and *P. wendlandii* with a dark green ring at the apex of the petiole.

Other apparently unrelated species may likewise have purple rings around the apexes of the petioles. These include *P. brenesii*, *P. davidsonii*, *P. dressleri*, *P. ferrugineum*, *P. microstictum*, *P. smithii*, *P. warszewiczii*, and sometimes *P. grayumii*.

Petioles may produce a cluster of viscid droplets of a sweet, sugary solution on the abaxial surface at the apex of the youngest leaves. This has been observed on *P. davidsonii* subsp. *bocatoranum* (Fig. 19) and *P. megalophyllum* Schott, a common South American species. These droplets appear to have no function in pollination or in blade orientation, and I speculate that they act as a food source for ant guards, which serve to prevent the young blades from predation by phytophagous insects. This seems all the more likely because the droplets are associated with new leaves, which are typically very tender and thus most easily damaged.

Freshly cut petioles usually form resin droplets in the same manner as the stems, and may in time become completely covered with resin. Of rare occurrence are slender strands of latex on broken and partially severed petioles as on *P. malesevichiae* (Fig. 21).

Dried petioles yield another suite of characteristics generally unrelated to those exhibited on fresh plants.

The dried petiole is sometimes diagnostic and in some cases, such as *P. hebetatum* and *P. schottianum*, petioles have a dried epidermis that is so conspicuously yellow-brown as to be nearly unique. Though less conspicuous, the petioles of *P. schot-*

*tianum* and *P. thalassicum* also dry quite yellowish brown.

Petiole firmness is variable in *P. subg. Philodendron*, but this character is difficult to quantify. Unfortunately Engler used this character as one of his major key characters to separate subsections of *P. sect. Cardiobelium*, *P. subsect. Macrobelyum*, and *P. subsect. Glossophyllum* with very fleshy petioles from *P. Gruppe Oligocarpidium*, and *P. Gruppe Doratophyllum* (now *P. subsect. Bulaoana*) with firm petioles. This character is difficult to describe and petioles appear to range from quite firm to quite spongy without any major discontinuities. Most species have petioles firm to weakly spongy when squeezed, and generally they are quite flexible, capable of being bent to a great extent without breaking. However, spongy petioles, such as those in *P. ligulatum* which can be easily crushed by squeezing, are also more likely to be brittle.

**Blades. Blade shape.** Virtually the entire range of morphological variation in leaf shape in the genus as a whole is exhibited among the Central American species of *P. subg. Philodendron*. The only blade shapes of *Philodendron* not shown by Central American species are those seen on *P. goeldii* G. Barroso in *P. subg. Meconostigma*, which has blades reniform in outline and pedately compound, and members of *P. sect. Schizoplacium*, which have blades pinnately lobed with a few, prominently falcate divisions. While *P. subg. Pteromischum* has distinctive leaves (even discounting the winged petiole on adult plants, which is definitive), their overall shape and size is matched by some other species of *P. subg. Philodendron*. Blade shape in *P. subg. Philodendron* relates in some cases to sectional differences. For example, species with three-lobed blades are either members of *P. sect. Tritomophyllum* or *P. subsect. Bulaoana*. Pinnately lobed species are members of *P. sect. Polytomium*. Some sections, especially *P. sect. Philodendron* and *P. sect. Calostigma*, are highly variable in blade shape, with that of *P. sect. Philodendron* ranging from oblong to variously ovate to ovate-sagittate and that of *P. sect. Calostigma* ranging from oblong to subcordate, prominently cordate, cordate-sagittate, or even 3-lobed. *Philodendron subsect. Glossophyllum*, relatively common in Central America, has more or less oblong, frequently subcordate or cordulate blades. See also section on "Discussion of Subgeneric Classification" for additional details.

While lobed leaves in some Araceae, e.g., *Monstera*, are due to necrotic processes in which growth stops and tissue rots away (Madison, 1977), the

lobed leaves of *Philodendron* are due to differential growth of leaf tissue.

In Central America more species have ovate-cordate blades than any other shape. In all variations, 38 species have some sort of ovate-cordate leaf blade. Twenty-two species are described as having ovate-cordate blades, and 11 as broadly cordate. Species with ovate to ovate-triangular leaves are most abundant among *P.* subg. *Philodendron* in Central America. At least 55 species have leaf blades with well-developed posterior lobes, including cordate, sagittate, and hastate. Only 15 species have non-cordate blades prominently longer than wide. Eight species have lobed blades, six trilobed, and two pinnately lobed. Species with trilobed leaves are: *P. anisotomum*, *P. cotobrusense*, *P. madronense*, *P. rothschuhianum*, *P. tripartitum*, and *P. angustilobum*. Species with pinnately lobed leaves are: *P. radiatum* and *P. warszewiczii*.

**Posterior lobes.** *Philodendron* subg. *Philodendron* species vary greatly in the shape of posterior lobes: usually rounded for those species with cordate leaf bases, e.g., *P. glanduliferum* (Fig. 197) or *P. platypetiolatum* (Fig. 306), or cordulate<sup>12</sup> leaf bases, e.g., *P. auriculatum* (Fig. 69), *P. immixtum* (Fig. 236), and *P. pseudauriculatum* (Fig. 314); to more or less oblong to narrowly triangular and much longer than wide as in *P. verapazense* (Fig. 434), or sometimes in *P. sagittifolium* (Fig. 346) and *P. tenue* (Fig. 404). Even in species with well-developed posterior lobes the lobes may be about as long as broad, as in *P. purulhense* (Fig. 322), *P. tysonii* (Fig. 426), or *P. schottianum* (Fig. 365). Posterior lobes may be sagittate as in *P. brunneicaule* (Fig. 95), *P. hebetatum* (Fig. 218), or *P. gigas* (Fig. 194); to hastate as in *P. angustilobum* (Fig. 49), *P. anisotomum* (Fig. 55), and *P. mexicanum* (Fig. 287); or bluntly triangular as in *P. wilburii* var. *longipedunculatum* (Fig. 461).

**Sinus shape.** *Philodendron* species with posterior lobes have remarkable variation in the shape of the sinus: merely arcuate, in *P. microstictum* (Fig. 289); arcuate with decurrent petioles in *P. morii* (Fig. 292); V-shaped in *P. glanduliferum* (Fig. 200), *P. lentii* (Fig. 263), *P. squamipetiolatum* (Fig. 386), and *P. tenue* (Fig. 406); oblong in *P. duyeryi* (Fig. 155); spatulate in *P. basii* (Fig. 76), *P. hebetatum* (Fig. 218), *P. smithii* (Fig. 372), and *P. rothschu-*

*hianum* (Fig. 339); parabolic in *P. sulcicaule*; hippocrepiform in *P. hebetatum* (Fig. 225), *P. lazorii*, *P. panamense* (Fig. 297), and *P. squamicaule* (Fig. 374); sometimes closed on live plants as in *P. fortuneense* (Fig. 182), *P. gigas* (Fig. 194), *P. pterotum* (Fig. 311), and *P. schottianum* (Fig. 366). In *P. fortuneense* and *P. subincisum* (Fig. 401) the sinus may be closed even on pressed plants. The shape of the sinus varies greatly between live and flattened dried plants, since the shape of the sinus varies depending on the angle at which the posterior lobes are turned up in relation to the midrib. Many species have the posterior lobes turned upward at an angle to the midrib on live plants (see, e.g., Figs. 104, 130, 159, 186, 366, 374), causing the inner margins of the posterior lobes to become closer to each other and thus decreasing the apparent size of the sinus. For example, when the angle of the posterior lobes is extreme the posterior lobes might be closed, but when the same blade is flattened and dried the sinus might become spatulate or hippocrepiform. The description of the sinus as presented in this work for Central American *P.* subg. *Philodendron* is exclusively that of the flattened sinus unless otherwise stated.

The sinus shape may be intraspecifically variable as in *P. sousae*, for example, with the sinus ranging from spatulate (Fig. 379) to parabolic (Fig. 380). This is in part related to the age of the plant, with older plants bearing larger blades that have larger, more well developed posterior lobes.

**Blade size.** Blades of *P.* subg. *Philodendron* in Central America as elsewhere are highly variable in size, but relatively few have blades that are considered huge. Only 19 Central American species of *P.* subg. *Philodendron* have blades that exceed 75 cm long (although many others approach that size). Of these, only *P. gigas* has blades that regularly exceed 1 m in length, with the maximum recorded at 137 cm. The blades of *Philodendron ferrugineum* may rarely exceed 1 m in length. By the same token few species in Central America have small leaves. Only 37 species have blades less than 25 cm long on adult plants, and of these only 13 have blades less than 15 cm long. *Philodendron brewsterense*, with blades a maximum of 11 cm long, is the species with the smallest leaves, and *P. chirripoense*, with blades up to 11.6 cm long, has leaves about as small. The median leaf length for *Philodendron* in Central America is 57 cm.

The widest blades belong to *P. gigas* and *P. radiatum* (to 90 cm long), but *P. pterotum* (to 84 cm wide), *P. finders* (to 70 cm wide), *P. dodsonii* (to 66

<sup>12</sup> With miniature lobes held very near the petiole (see photos of *P. auriculatum*) as compared to subcordate, wherein the lobes, though short, are usually broadly rounded with each lobe encompassing about half the width of the entire blade.

cm), *P. schottianum* (to 64 cm), and *P. warszewiczii* (to 62 cm) also have quite broad blades.

While blade size of *Philodendron* is often not reliable for taxonomic separation and even blade shape is sometimes unreliable, the length-width ratio is, in general, more reliable as a taxonomic character. Blades range from being much broader than long, to more than eight times longer than broad (in *P. cretosum*), to broader than long. Blades are broader than long in a number of species, e.g., averaging only 0.62 times as long as broad in *P. anisotomum* and *P. brunneicaule* and 0.7 times as long as broad in *P. cotabruense* and *P. graymii*. The average blade length-width ratio for species with simple, unlobed leaves ranges from 1.6 to 2.5 times longer than wide. Exceptions include *P. cretosum* and *P. folsomii* (8.3 and 5.2 times, respectively).

Blade margins are typically entire on simple leaves, never toothed and only rarely sinuate, such as in *P. subincisum*. Species with weakly sinuate blades are: *P. basii*, *P. dreyeri*, and *P. jefense*. On the other hand, blade margins are frequently undulate to a certain extent, especially on larger blades. Species that have coarsely undulate leaves are: *P. annulatum*, *P. antonioanum*, *P. copense*, *P. ferrugineum*, *P. fortunense*, *P. grandipes*, *P. hebetatum*, *P. panamense*, and *P. sulcicaule*.

Blade margins are frequently hyaline and also often weakly revolute. Examples of species with hyaline blade margins are: *P. advena*, *P. crassispatum*, *P. davidsonii*, *P. ferrugineum*, *P. gigas*, *P. immixtum*, *P. hederaceum*, *P. smithii*, *P. squamicale*, *P. straminicaule*, *P. subincisum*, and *P. wilburii* var. *longipedunculatum*.

The coloration of leaf blades is highly variable and largely dependent on the habitat, but some species, such as *P. roseospathum*, *P. scalarinerve*, and *P. tysonii*, have very dark green blades while others (e.g., *P. lazoria*) have lighter green blades. A few species, such as *P. chiriquense*, have purple coloration on young leaf blades but most lose this coloration in age. An exception is *P. verrucosum*, which has adult blades generally purplish on the lower surface. A few taxa such as *P. ligulatum* var. *heraclioanum* (Fig. 274) and *P. sagittifolium* have leaf blades purplish-spotted on the lower surface, especially when young. Coloration of lower blade surfaces is more often restricted to the major veins, which are frequently reddish (along with the young petiole) as, for example, in *P. antonioanum*.

**Venation.** *Midribs.* Leaf midribs of *P.* subg. *Philodendron* are more variable in cross-sectional shape than those of other aroid genera, such as *Senospermaton*, *Rhodospatha*, *Spathiphyllum*, and

*Monstera*, yet not as taxonomically significant as in the related genus *Dieffenbachia*.

Midribs on the upper (adaxial) blade surface of *P.* subg. *Philodendron* are highly variable, being flattened or variously sunken or raised (sometimes even in the same species). Fifty-six species have adaxial midribs at least sometimes raised, 26 have upper midribs at least sometimes sunken (only *P. cretosum* has a deeply sunken midrib), and 54 have adaxial midribs sometimes flat. Only 12 taxa, *P. antonioanum*, *P. aromaticum*, *P. albisuccum*, *P. coloradense*, *P. dressleri*, *P. granulare*, *P. pirrense*, *P. purulhense*, *P. strictum*, *P. warszewiczii*, *P. wilburii* var. *longipedunculatum*, and *P. zhanum*, are described as having strictly flattened adaxial midribs. Fifteen species are described as having only convex midribs, and 15 are described as having only sunken midribs. Only 2 species, *P. morii* and *P. niqueanum*, are described as having the adaxial midribs prominently raised.

The adaxial midrib is nearly always to some extent paler than the surface. In only 12 species are the midrib and the blade concolorous. An additional 8 species have midribs concolorous to slightly paler.

The lower (abaxial) midrib of *Philodendron* blades exhibits less variability. All species have adaxial midribs to some extent raised. A few species, e.g., *P. crassispatum*, have the abaxial midrib so broadly convex as to appear nearly flat, but most species have the midrib noticeably raised. The abaxial midrib is generally convex but often narrowly rounded, thicker than broad, and sometimes bluntly acute. Species with the abaxial midrib weakly raised are rare in *P.* subg. *Philodendron*. Only seven species have abaxial midribs at least sometimes described as broadly convex or weakly raised: *P. anisotomum*, *P. crassispatum*, *P. fortunense*, *P. microstictum*, *P. platypetiolatum*, *P. purpureoviride*, and *P. straminicaule*.

Many more species have abaxial midribs more prominently raised. In the largest category, 47 species have abaxial midribs more or less convex, i.e., more or less hemispherical in cross section. Thirty-six species have abaxial midribs described as thicker than broad, narrowly convex, narrowly rounded or bluntly acute at least part of the time. A number of these species have abaxial midribs so prominently raised as to be noticeably thicker than broad: *P. copense*, *P. cretosum*, *P. dolichophyllum*, *P. hebetatum*, *P. heleniae*, *P. jodavisanum*, *P. panamense*, and *P. roseospathum*. Taxa that have abaxial midribs sometimes thicker than broad are *P. scalarinerve* and *P. wilburii* var. *longipedunculatum*.

The most extreme type of raised adaxial midribs

is so raised as to appear to be a cylinder attached tangentially to the surface of the leaf. This type of midrib has been referred to as "round-raised." Examples of species with such midribs are *P. brunneicaule*, *P. dressleri*, *P. madronense*, *P. squamipetiolatum*, and *P. radiatum*. In cases where the midrib is round-raised, the primary lateral veins are also sometimes round-raised.

While the abaxial midrib is more likely than the adaxial midrib to be concolorous with the rest of the blade or darker than the remaining blade surface, 45 species still are reported as having the midrib paler than the surface. Twenty-four species have the abaxial midrib described as darker than the surface.

A number of species, including *P. edenudatum*, *P. grayumii*, *P. ligulatum* var. *ligulatum*, and *P. sagittifolium*, have one or both midribs maroon- or purplish-spotted.

**Primary lateral veins.** The primary lateral veins (referred to by some authors as "secondary veins") are those that branch off the midrib and extend to the margins usually without additional branching. This use of primary lateral vein is consistent with that of early aroid workers Engler, Krause, and Sodiro. Engler and later Krause referred to the primary lateral veins as "nervis primariis." Sodiro also referred to the primary lateral veins as "nervis laterales I."

On larger cordate blades there is frequent branching of the primary lateral veins in the lower part of the blade where primary lateral veins are widely spaced (Fig. 26) (see also Croat & Bunting, 1979). The presence of these veins, referred to here as "secondary veins," is not commonly indicated in this work, but in shape and aspect they are virtually the same as the primary lateral veins (Fig. 24).

Both surfaces of the blade typically have primary lateral veins similar to the associated midrib. Usually they are somewhat less prominently raised than the midrib. The primary lateral veins on the upper (adaxial) surface, while usually sunken, may be essentially flat with the surface. Sometimes they are raised but at the same time contained within a broad or narrow valley so that the veins appear to be sunken. Primary lateral veins may sometimes be "quilted" (Figs. 23, 200, 361), i.e., with the veins deeply sunken and with the intervening leaf tissue of the blade that lies between the primary lateral veins being broadly raised, making the surface appear like that of a plush quilt.

Primary lateral veins of *P.* subg. *Philodendron* are rarely either absent or so inconspicuous as to appear to be absent, such as in *P. brewsterense*,

though they are sometimes not at all conspicuous, as in *P. microstictum* (Fig. 289) or *P. sulcicaule* (Fig. 403). The number of primary lateral veins varies from 2 pairs in *P. chirripoense* to 25 pairs in *P. madronense*. Most species have 3 to 6 pairs of primary lateral veins. These generally arise at an acute angle with the midrib and, after extending along or near the midrib, spread at an angle of generally 40–80° toward the margins, generally forming a broad arc in the process. The angle of primary lateral veins for all species may be as little as 5° to as much as 100°. When species of Central American *P.* subg. *Philodendron* are divided into a series of range categories based on the angle of their primary lateral veins, more species (eight) have primary lateral veins spreading at 50°–60° angle or at 60°–70° (also eight species) than any other group.

**Basal veins.** Primary lateral veins that contact the plexus at the base of the blade and the apex of the petiole are here referred to as "basal veins" (Fig. 22) (see also Croat & Bunting, 1979). While the uppermost 1 or 2 pairs usually extend upward and into the anterior lobe where they join the antemarginal vein, most extend into the posterior lobes. Generally, basal veins are best developed on plants that have posterior lobes and are not present, or at least are few in number, when the blade lacks posterior lobes. Basal veins, like the primary lateral veins, are typically paired, with usually an equal number on either side of the midrib.

Seventy-three species (75 taxa) of *P.* subg. *Philodendron* in Central America have basal veins. The number of pairs of basal veins ranges from 1 to 15, but only eight species (*P. angustilobum*, *P. copense* (to 15), *P. findens*, *P. grandipes*, *P. thalassicum*, *P. pirrense*, *P. squamicaule*, and *P. strictum*) have 10 or more pairs. Most species have 3–6 pairs of basal veins, though up to 16 species may have only 2 pairs of basal veins. Only three species, namely *P. lentii*, *P. mexicanum*, and *P. scalarinerve*, regularly may have a solitary pair of basal veins.

**Posterior ribs.** The basal veins are generally to some extent coalesced near their union with the petiole. The union or coalescence of basal veins is here referred to as the "posterior rib" (Figs. 22, 25) (see also Croat & Bunting, 1979). Sixty-three species (65 taxa) have the basal veins at least in part united and thus possess posterior ribs, while 34 species (38 taxa) have the basal veins free to the base (or lack basal veins all together) and thus lack posterior ribs. Both the number and nature of basal veins and the length of the posterior rib are good characters for distinguishing taxa. The basal veins

may be completely coalesced for a distance (Fig. 22) or loosely coalesced. The posterior rib may be prominently naked for a distance (Figs. 22, 169) or barely (Figs. 26, 211) or not at all naked (Figs. 161, 163, 287). The basal veins are free or nearly so in most species with rounded or cordulate leaf bases (Fig. 20). Such species are: *P. auriculatum*, *P. bakeri*, *P. corraeae*, *P. chirripoense*, *P. cretosum*, *P. dolichophyllum*, *P. granulare*, *P. immixtum*, *P. lentii*, *P. ligulatum*, *P. pseudauriculatum*, *P. roseospathum*, *P. scalarinerve*, *P. ubigantupense*, *P. utleyanum*, and *P. wendlandii*. Even species with short or poorly developed posterior lobes, such as *P. annulatum*, *P. clewellii*, *P. davidsonii*, *P. microstictum*, *P. niqueanum*, and *P. platypetiolatum* usually have the basal veins free to the base. On the other hand, species with well-developed posterior lobes usually have the basal veins coalesced to some degree. Posterior ribs range from less than 0.5 cm long (in *P. straminicaule*) and 1 cm long (in *P. angustilobum*, *P. basii*, and *P. alticola*) to more than 14 cm long in *P. gigas*.

Another important taxonomic character is the extent to which the posterior rib is naked along the sinus. In general, the longer the posterior rib the more likely it is to be naked at least to some extent along the sinus. However, *P. tenue* is exceptional in having a well-developed posterior rib and being not naked along the sinus. The degree to which the posterior rib is naked along the sinus varies from as little as 0.5 cm in *P. straminicaule* to 6 cm in *P. squamicaule*. Examples of species with prominently naked posterior ribs are: *P. albisuccus*, *P. alticola*, *P. aromaticum*, *P. basii*, *P. brunneicaule*, *P. copense*, *P. dodsonii*, *P. dressleri*, *P. findens*, *P. hebetatum*, *P. knappiae*, *P. panamense*, *P. purulhense*, *P. strictum*, *P. thalassicum*, and *P. verapazense*.

**Lesser order veins.** Between successive primary lateral veins there are possibilities for two additional orders of veins. Frequently present are intermediate primary lateral veins here called "interprimary veins" (Figs. 24, 26) (see also Croat & Bunting, 1979). These veins, while decidedly less conspicuous than the primary lateral veins, are nevertheless too prominent to be classified as the smallest-order veins. To qualify as an interprimary, the vein must extend continuously from the midrib to very near the margin without major branching. Generally there is no more than one pair of interprimary veins between alternate primary lateral veins. They are akin to the primary lateral veins in all aspects except for their reduced size. Like the primary lateral veins they may bear minor veins, which may form all along their margins.

The smallest-order veins are here referred to as "minor veins" (Fig. 24) and may be close, fine, and conspicuous as in *P. sulcicaule*, *P. tripartitum*, and *P. radiatum*, to thick, well-spaced, and inconspicuous in *P. gigas*, *P. granulare*, *P. grayumii*, and *P. ligulatum*. The minor veins are not all equally distinct and sometimes, as in *P. dominicalense*, the minor veins are alternately weakly visible and strongly visible.

The minor veins may arise from the midrib or from the primary lateral and interprimary veins, but in either case they form a generally close, uniform, and parallel array, which extends without interruption to near the margin of the blade. In most species the minor veins arise from both the midrib and the primary lateral veins, but some species have the minor veins arising from only the midrib. A total of 77 species (81 taxa) have the minor veins arising from both the midrib and the primary lateral veins (Fig. 26). In such cases the minor veins are not always equally arising from one of the two entities but may, as in the case of *P. brenesii*, *P. davidsonii* subsp. *davidsonii*, *P. ferrugineum*, and *P. tripartitum*, be more heavily arising from the midrib rather than the primary lateral veins. In *P. auriculatum*, *P. glanduliferum*, *P. lentii*, and *P. ligulatum* the minor veins arising from the primary lateral veins are many fewer than those arising from the midrib. In *P. heleniae* the primary lateral veins are only rarely arising from the midrib.

In another variation of this venation type, some species, while having minor veins arising from the midrib as well as both adjacent primary lateral veins, have considerably more veins arising from the distal primary lateral vein than from the proximal vein.

A total of 24 species have the minor veins arising from only the midrib, and in this case they course along the primary lateral veins but do not join with it. Many of the species that have the minor veins arising only from the midrib are species with oblong blades, such as: *P. bakeri*, *P. cretosum*, *P. dolichophyllum*, *P. granulare*, *P. heleniae*, *P. roseospathum*, *P. ubigantupense*, *P. utleyanum*, and *P. wendlandii*. However, this group also has blades ovate or nearly so as in *P. brewsterense*, *P. chirripoense*, *P. cotobrusense*, *P. crassispatum*, *P. folsomii*, *P. knappiae*, *P. microstictum*, *P. niqueanum*, *P. sulcicaule*, and *P. verapazense*. Interestingly, *P. anisotomum*, with deeply 3-lobed leaf blades, also has the minor veins arising from only the midrib, whereas *P. tripartitum*, a similar species with 3-lobed blades, has the minor veins arising from both the midrib and the primary lateral veins.

*Philodendron dressleri*, a species with deeply di-

vided leaves, has a more complex venation pattern. Although the minor veins arise from both the midrib and the primary lateral veins, they also arise from short secondary veins that regularly branch off the primary lateral veins. In addition, the minor veins that arise from the midrib are considerably fewer in number and weaker than in most species with this venation pattern. Generally, the confluent minor veins that arise from the primary lateral veins and make a broad sweep before continuing to the margin leave little area for the minor veins, which arise from the midrib. The latter tend to merge imperceptibly with those from the primary lateral veins. A similar pattern with weak midrib-borne minor veins is present with *P. basii*.

At or very near the margin both the primary lateral veins and the minor veins generally turn sharply toward the apex of the blade. The minor veins join with other minor veins and with the primary veins before finally merging into an inconspicuous and somewhat opaque marginal plexus. This narrow band is usually chlorophyllous and apparently veinless. Frequently the outer margin of this chlorophyllous band is a hyaline edge, which is colorless and typically revolute.

The minor veins are sometimes noticeably interconnected by inconspicuous to conspicuous veins, referred to here as "cross-veins" (Fig. 26). Generally the cross-veins are markedly perpendicular to consecutive minor veins where they are sufficiently prominent to be noticeable, but in some cases the cross-veins cross transversely from one minor vein to the next. While *P. scalarinerve* has cross-veins so prominent, even on fresh material, as to be easily visible, other species such as *P. chiriquense* and *P. copense* have cross-veins easily visible only when the blades are dry.

Secretory ducts and other secretory tissues are frequently present on Central American members of *P. subg. Philodendron*. The contents of the secretory ducts are either latex or tanniferous compounds (Solereder & Meyer, 1928). No thorough survey has been made of the nature of the secretory canals in Central American species, so it is not always apparent whether these structures are resin canals or secretory ducts. Secretory canals (also referred to as secretory files) in *Philodendron* are always non-anastomosing and consist of a linear sequence of secretion cells, each separated from the next by cell walls (Solereder & Meyer, 1928). On fresh and dried leaves these can usually be recognized by being darker, usually blackened, and in being intermittent rather than continuous as is generally true of veins. While the distribution of secretory ducts in *Philodendron* may be more com-

mon than is apparent from surface examination, not all species exhibit the secretory ducts clearly. Thus the presence or absence of distinct secretory ducts can be useful taxonomically. They are distinctly visible on *P. alticola*, *P. cotonense*, *P. grayunii*, *P. heleniae*, and *P. zhuangum*, but obscurely visible on *P. antonioanum*, and *P. bakeri*. They are clearly visible on *P. schottianum*, somewhat visible on *P. llanense*, and obscure on *P. findens*, even though these three species have very similar leaf blades.

While perhaps no more reliable than blade shape, blade size, or other features, blade surfaces at a magnification of 10× or higher often reveal another suite of characters, which often yields another degree of confidence (or forewarns of misidentification) when making determinations. The surface between the minor veins is frequently marked by pale sub-surface granulations (perhaps indicating the presence of druse crystals), short, pale lineations, gland-like punctiform markings, reddish or brownish speckling, and also what might be referred to as "stitching," pale intermittent short lines appearing on the surface of the blade as though the blade was loosely sewn with a needle and thread. Though this phenomenon is much more common in *P. subg. Pteromischum*, it is also exhibited in *P. subg. Philodendron*.

## MORPHOLOGY OF REPRODUCTIVE STRUCTURES

### INFLORESCENCES

When a plant of *P. subg. Philodendron* flowers, the stem apex is terminated by an inflorescence (or frequently an aborted inflorescence primordium) (Ritterbusch, 1971; Blanc, 1977c; Madison, 1978) and new growth is from a bud in the axil of the penultimate leaf. This new growth pushes aside the newly developing leaf and the inflorescence that is developing in its axil (Ray, 1987a). When the inflorescences abort, the petiole sheath remains quite small and unexpanded, but when the inflorescences develop, the sheath may be much larger. *Philodendron* inflorescences are sympodial, with each sympodium consisting of 1-10 (to 11 in South America) inflorescences arising typically in what appears to be the leaf axil of usually one of the upper leaves (but almost never in the uppermost leaf axil). Often inflorescences emerge from the mass of cataphyll fibers, enabling the inflorescence to remain moist (Figs. 63, 115, 188, 371, 430). The individual inflorescences are subtended by and enclosed in a series of moderately coriaceous, whitish to pinkish, 2-ribbed structures (Fig. 195, *P. gigas*, Fig. 428, *P. tysonii*) called bracteoles (Ray, 1987a). These have elsewhere been referred to as prophylls (Usher,



1966; Jackson, 1971). At anthesis the bracteoles are typically deciduous, although sometimes they persist for a time simply because they are held so tightly by the developing infructescences.

Typically the inflorescences are borne in the erect or semi-erect position regardless of the position of the stem. Even when the inflorescence is borne on the end of a pendent stem, the stem usually curves upright so that the opening inflorescence is erect at anthesis (Fig. 273, *P. ligulatum* var. *ovatum*; Fig. 269, *P. ligulatum* var. *ligulatum*).

The number of inflorescences per axil is taxonomically significant. More species (38) have a solitary inflorescence per axil (or are believed to have a solitary inflorescence) than any other combination. Many others may be found with only a single inflorescence if they are seen in the early stages of flowering or if inflorescences have aborted owing to lack of pollination, but plants frequently have two or more inflorescences per axil later in the flowering season. Careful dissection of the leaf axil usually can provide evidence of the peduncular scar indicating a lost inflorescence. Taxa known to usually have only a single inflorescence per axil are: *P. albisuccum*, *P. anisotomum*, *P. aromaticum*, *P. bakeri*, *P. basii*, *P. breedlovei*, *P. brevispathum*, *P. brewsterense*, *P. brunneicaule*, *P. chirripoense*, *P. coreae*, *P. cotonense* (rarely 2), *P. crassispalum*, *P. dressleri*, *P. dwyeri*, *P. edendatum*, *P. folsomii*, *P. glanduliferum*, *P. granulare*, *P. hammelii*, *P. hederaceum*, *P. immixtum*, *P. jacquinii*, *P. jefense*, *P. knappiae*, *P. ligulatum* var. *heracleoanum*, *P. ligulatum* var. *ligulatum* (sometimes 2, rarely to 5), *P. ligulatum* var. *ovatum*, *P. madronense*, *P. mexicanum*, *P. microstictum*, *P. platypetalatum* (rarely 2), *P. purpureoviride*, *P. purulhense*, *P. radiatum* var. *pseudoradiatum*, *P. squamipetalatum*, *P. utleyanum*, *P. verapazense*, and *P. warszewiczii* (rarely to 3).

Most species with a single inflorescence per axil are either vines or appressed-climbing hemiepiphytes with internodes longer than broad.

The number of upper axils bearing inflorescences is also variable. Some species, e.g., *P. lentii*, rarely have more than one leaf axil bearing an inflorescence, while other species, e.g., *P. annulatum* and *P. fragrantissimum*, bear inflorescences in two or more of the closely spaced upper internodes, making it appear that the plants have a large number of inflorescences.

Species that regularly have up to four or more inflorescences per axil are: *P. copense* (to 6), *P. cotobrusense* (to 5), *P. clewellii* (to 6), *P. dodsonii* (2-5), *P. ferrugineum* (to 6), *P. fortunense*, *P. gigas* (to 7), *P. graymii* (to 5), *P. heleniae* (to 10), *P. jodaviansium* (2-6), *P. panamense* (to 6), *P. pirrense*, *P.*

*squamicaule* (to 5), *P. sousae*, *P. tenue*, *P. tysonii* (to 5), and *P. verrucosum*.

Species with five or more inflorescences per axil are rare, totaling only eleven. Only two species, *P. gigas* and *P. heleniae*, have up to seven inflorescences per axil.

#### PEDUNCLES

Typically the peduncle is terete but it may be somewhat flattened laterally, especially when there are clusters of inflorescences per axil. The peduncle is usually whitish or pale green at the base where it is often hidden by the leaf sheath. The remaining portion of the peduncle is usually dark to medium green with the surface often shortly pale-lineate like the petioles. The peduncle is often coarsely pale-ribbed near the apex, as in *P. copense*, *P. dodsonii*, and *P. findens*. Rarely is the peduncle pinkish red or reddish as in *P. malesevichiae*, *P. schottianum*, and *P. roscospathum* (or sometimes in *P. sulcicaule*) or tinged with red or purple as in *P. chiriquense*, *P. davidsonii* subsp. *bocatoranum*, *P. heleniae*, *P. grandipes*, *P. purpureoviride*, and *P. verrucosum*.

Most peduncles, like other parts of the average *Philodendron*, are glabrous, but some species have peduncles conspicuously covered with trichome-like glands, e.g., *P. squamicaule* (Fig. 384), *P. squamipetalatum* (Fig. 388), and *P. verrucosum* (Fig. 440).

The peduncle is usually much narrower than the spathe, commonly no more than  $\frac{1}{2}$  to  $\frac{3}{4}$  the width of the spathe tube, and is almost always broadened distally, merging almost imperceptibly with the spathe tube. Usually the color distinction is also gradual, but in some cases, e.g., *P. annulatum*, *P. dodsonii*, and *P. dolichophyllum*, there is an abrupt transition between the green peduncle and the colored spathe. In *P. ligulatum* there is a purple ring at the apex of the petiole, much like that at the apex of the petiole (Fig. 276).

The length of the peduncle, especially relative to the length of the spathe, may be important taxonomically. Some species, e.g., *P. antonioanum*, *P. crassispalum*, *P. davidsonii*, *P. ferrugineum*, *P. findens*, *P. malesevichiae*, *P. purulhense*, *P. radiatum*, *P. schottianum*, and *P. zhuangum*, have very short peduncles relative to the length of the spathe. In contrast, other taxa, e.g., *P. advena*, *P. angustilobum*, *P. dodsonii*, *P. hebetatum*, *P. heleniae*, *P. immixtum*, *P. mexicanum*, *P. pterotum*, *P. rothschildianum*, and *P. wilburii* var. *wilburii*, have peduncles as long as or longer than the spathes.

Peduncles range in length from 1 to 25 cm, with

four taxa, *P. microstictum*, *P. pseudauriculatum*, *P. wilburii* var. *longipedunculatum*, and *P. verrucosum*, all having peduncles that attain the maximum length of 25 cm. Several additional species have peduncles that may be more than 20 cm long. These include *P. anisotomum*, *P. brunneicaule*, *P. panamense*, *P. pseudauriculatum*, *P. rothschuhianum*, and *P. scalarinerve*.

Many species of *P.* subg. *Philodendron* have peduncles more than 10 cm long. In addition to those mentioned above, 37 Central American species have peduncles more than 10 cm long, while 38 taxa (35 species) have peduncles less than 10 cm long. Sixteen species have petioles that may be less than 3 cm long, but only *P. knappiae* has a peduncle which does not exceed 3 cm in length.

#### SPATHES

The spathe of *P.* subg. *Philodendron* is highly variable in many regards. The spathe of *Philodendron* is typically coriaceous and constructed in such a way that the spathe may open and reclose without major alteration in its shape. The margins are invariably much thinner, with the apical portion of the spathe usually tightly rolled and frequently acuminate (Fig. 178, *P. findens*) and the basal portion more conspicuously convolute. Because the margins of the base are more conspicuously overlapped, when the spathe opens the lower portion remains convolute (Fig. 70, *P. auriculatum*) whereas the apical portion separates to very near the tip. Some species have spathes that open more fully so that the opening is almost rounded (Fig. 298, *P. panamense*) or broadly elliptic (Fig. 288, *P. mexicanum*). In such cases more of the inner tube surface and the pistillate portion of the spadix are exposed. On the other hand, some species have spathes that open only slightly so the opening is merely elliptic with the pistillate portion of spadix not visible without peering down through the top of the opening (Fig. 63, *P. antioianum*).

Species vary greatly in the extent to which the spathe remains convolute at the base during anthesis. Spathes of some species, e.g., *P. angustilobum* (Fig. 51) and *P. mexicanum* (Fig. 288), unfold to very near the base. Those of other species, e.g., *P. antioianum* (Fig. 63), do not open very broadly. The spathe tubes of *P. auriculatum* (Fig. 70), *P. panamense* (Fig. 298), and *P. wendlandii* (Fig. 451) remain convolute only in the lower half. The spathe of *P. davidsonii* (Figs. 134, 136) may be convolute only in the lower  $\frac{1}{2}$  to  $\frac{3}{4}$  in *P. squamicaule*. That of *P. hebetatum* (Fig. 226) is convolute to the upper  $\frac{1}{4}$  and the spathe tubes of *P. aromaticum* (Fig. 68),

*P. brenesii* (Fig. 89), *P. copense* (Fig. 110), *P. scalarinerve* (Fig. 140), *P. findens* (Fig. 177), *P. rothschuhianum* (Fig. 340), *P. stramineicaule* (Fig. 360), and *P. schottianum* (Fig. 371) remain essentially closed to the apex of the tube during anthesis.

Mayo (1986) presented a classification of inflorescence types based principally on South American *Philodendron* species. Eleven inflorescence types were characterized, and a key was prepared to separate them. The characters chosen for use in the key were: (1) presence or absence of resin ducts in the inner surface of the spathe; (2) solitary vs. two or more inflorescences per axil; (3) relative length of the sterile staminate portion of the spadix (equal to or longer than the staminate portion vs. shorter than staminate zone); (4) presence or absence of colorful contrasts on inner surface of spathe tube; (5) prominently constricted vs. weakly constricted spathes; (6) presence or absence of resin ducts in the spathe; (7) presence of resin ducts in the sterile staminate portion of spadix vs. with resin ducts in either the fertile staminate portion or in both sterile and fertile staminate portions; (8) presence of non-functional resin canals in the spathe vs. resin canals lacking in the spathe (usually correlated with contrasting spathe colors); and (9) presence or absence of a terminal sterile zone on the spadix.

While I believe that this classification may be somewhat artificial, it demonstrates many evolutionarily important morphological features and shows the complexity of the *Philodendron* spathe.

Species of *P.* subg. *Philodendron* have the spathe tube relatively well demarcated from the blade either by being constricted at the apex or by its contrasting color. Some species, e.g., *P. crassispatum* (Fig. 124), lack any constriction above the tube, thus the spathe is more or less elliptic. Species that have a weakly constricted spathe include *P. advena* (Fig. 40), *P. alticola* (Fig. 45), *P. breedlovei* (Fig. 85), *P. purulhense* (Fig. 323), *P. schottianum* (Fig. 368), and *P. antioianum* (Fig. 63). Despite these exceptions, most members of *P.* subg. *Philodendron* have spathes to some extent constricted. The subgenera *Meconostigma* and *Pteromischum*, on the other hand, have spathes not likely to be constricted or are only weakly constricted (Mayo, 1986).

Spathe length is relatively variable depending on the age of the plant and the stage of development. Elongation of the spathe is typically more or less arrested after anthesis, but sometimes a considerable amount of elongation takes place in both the peduncle and spathe after anthesis, presumably to accommodate the considerable expansion of the developing fruits. Spathe length varies from as little

as 4.5 cm in *P. clewellii* and 4.8 cm in *P. heleniae* to 29 cm in *P. pterotum* and 30 cm in *P. warszewiczii*. Relatively few species have spathes that ever exceed 25 cm in length even when in fruit.

Relatively few species have spathes less than 10 cm long. These are: *P. brewsterense*, *P. chirripoense*, *P. clewellii*, *P. dwyeri*, *P. knappiae*, *P. roseospathum* var. *angustilaminatum*, and *P. ubigantupense*.

The shape of the spathe and the relative disposition of the spadix at anthesis are taxonomically significant, but seldom described. This is because many species are still poorly known at anthesis, no doubt due to the fact that anthesis in *Philodendron* lasts only one or two days for each inflorescence.

One of the features that determines the shape of the spathe at anthesis is the degree to which the spathe opens. Spathes of some species, such as *P. aromaticum*, *P. copense*, *P. dodsonii*, *P. findens*, *P. fragrantissimum*, *P. grandipes*, *P. hebetatum*, *P. jordanianum*, *P. stramineicaule*, and *P. strictum*, open along the blade portion (see discussion above), leaving an oblong-elliptic opening (Fig. 110, *P. copense*; Fig. 140, *P. dodsonii*; Fig. 378, *P. smithii*). In these examples the spadix is either included within the spathe or is barely exerted, stiffly erect yet held slightly in front of the top edge of the spathe. On the other hand, spathes of some species, such as *P. angustilobum*, *P. mexicanum*, and *P. tripartitum* (Fig. 423), may open so broadly as to expose much of the pistillate portion of spadix. The pistillate portion of the spadix is usually for the most part obscured by the prominently convolute tubular portion of the spathe, as in *P. antonioanum*, *P. aromaticum*, *P. copense*, and *P. dodsonii*. Those species with spathes opening more broadly may also have the spadix protruding forward somewhat out of the spathe (see section on Spadix).

The definition of the spathe tube is somewhat imprecise, in terms of both morphology and color, but the spathe tube may be defined as the lower portion of the spathe (that portion which covers the pistillate portion of the spadix). In reality it usually extends somewhat above the pistillate portion to include the lowermost or sterile zone of the staminate portion of the spadix. In cases where the spathe has an obvious constriction it is that portion below the constriction (the balance being the spathe blade) that is referred to as the spathe tube. Though the ending of the spathe tube is imprecise, it generally terminates where the greatest constriction of the staminate portion of the spadix occurs, usually immediately above the sterile staminate portion of the spadix. Functionally, the constriction of the spathe, coinciding with the narrowest portion of the staminate portion of the spadix, prevents much of

the pollen from the fertile upper portion of the spadix from falling directly into the chamber of the pistillate portion of the spathe. This probably serves more to prevent wastage of the pollen (by containing the pollen in a place where it can most easily be carried away by beetles) than it does to prevent self-pollination, since most species are so markedly protogynous that no self-pollination is possible, i.e., the pistils are probably no longer receptive when the pollen is shed. Testing for pistil receptivity with the use of peroxidase paper, Grayum (1996) found stigmas receptive throughout the period of staminal dehiscence in *P.* subg. *Pteromischum*, but numerous attempts at self-pollinations of members of *P.* subg. *Philodendron* by me and others failed to produce berries and instead resulted in aborted inflorescences. Not surprisingly those species, e.g., *P. advena*, *P. crassispatum* (Fig. 124), and *P. purulhense*, that have spathes scarcely constricted above the tube also have spadices not markedly narrowed above the sterile staminate flowers. It is not known whether the pollination behaviors of these species differ.

For *Philodendron* species with a prominent spathe constriction, the partially closed spathe during staminal anthesis, coupled with the swollen area of the staminate portion of the spadix immediately below it, provides a "pollen well" through which the beetles must squeeze on their departure. This probably ensures a greater coverage of pollen for the departing beetle pollinators.

Spathe tube shape and length are not particularly useful taxonomically. Tubes range in shape from virtually oblong (e.g., in *P. heleniae*, *P. ligulatum*, and *P. tripartitum*), to oblong-elliptic (e.g., in *P. aromaticum*, *P. copense*, *P. dodsonii*, *P. findens*, *P. gigas*, *P. hebetatum*, and *P. pseudauriculatum*) to elliptic (e.g., *P. rothschuhianum* and *P. smithii*), and range from only 2 cm long in *P. clewellii* to 14 cm long in *P. davidsonii* and *P. warszewiczii*. Species with spathe tubes longer than 10 cm are: *P. auriculatum*, *P. ferrugineum*, *P. schottianum*, *P. sousae*, and *P. tysonii*. Seventeen species have spathe tubes less than 5 cm long.

Even though the spathe tube often remains totally closed at anthesis it is, nonetheless, still somewhat expanded owing to the separation caused by the flaring of the spathe blade. Generally this provides considerable room around the spadix to accommodate the pollinating beetles (Figs. 100, 128, 242, 268). At anthesis the spathe tube is considerably more voluminous than before anthesis.

*Spathe color.* Spathe coloration is generally taxonomically significant though variable depending on

the age of the inflorescence. Before anthesis spathe are frequently green on the outside, but reddish coloration often develops well in advance of anthesis. Coloration of the spathe tube and blade often differs, with the spathe tube more commonly various shades of green and the spathe blade more commonly white, whitish, or whitish green. Even when the spathe is green throughout, the spathe blade is typically paler green than the tube. The tube portion of the spathe is commonly colored or tinged with red, maroon, or purplish violet (sometimes with other shades of red, orange, or purple) on one or both surfaces. Strong color contrasts (other than merely green vs. white), so common in *Philodendron*, are absent in other ostensibly related genera, namely *Furtadoa*, *Homalomena*, and *Anubias* (based on the *Philodendron* Alliance of Mayo, Bogner & Boyce, 1995) and also *Cercestis* and *Culcasia* (included in the *Philodendron* Alliance of Grayum, 1990). This would also be true of other presumed relatives including *Dieffenbachia*, *Montrichardia*, *Nephtytis*, *Peltandra*, *Typhonodorum*, and even *Zantedeschia*.

Generally if the spathe is colored on the outer surface it is also colored on the inner surface (but the reverse is not true). The inner surface is typically much more intensely colored than the outer surface. The coloration of the inner surface of the spathe, though commonly more or less restricted to the tube, may extend well onto the blade, sometimes even to very near the apex, such as in *P. breedlovei*, *P. davidsonii*, *P. mexicanum*, and sometimes *P. sagittifolium*. In some cases, such as *P. findens* and *P. schottianum*, the color is merely weakly diffused onto the inside of the blade. At other times only the very base of the tube is colored inside, such as in *P. ligulatum* and *P. hederaceum*. The spathe blade is commonly more extensively colored on the outer than on the inner surface, although blade coloration is highly variable, both inter- and intraspecifically. Nevertheless, most species have spathe blades some shade of green or white on the outer surface, often tinged with red, pink, purple, or yellow. A few species, such as *P. sagittifolium* and *P. ferrugineum* (Fig. 167), have purplish spots or blotches throughout much of the exterior of the spathe surface. Although the external coloration of the spathe is usually restricted to the tube portion, it often extends onto the blade, a situation that happens more frequently on the outer surface of the spathe than on the inner surface. Species that have coloration extending well above the spathe tube are: *P. antioianum*, *P. breedlovei*, *P. dolichophyllum*, *P. subincisum*, and *P. roseospathum*. Sometimes the coloration of the spathe may

be restricted to the tube, but only near the margins of the open edge such as in *P. grayumii* and *P. malesewicziae*.

While some species, e.g., *P. alticola*, *P. heleniae*, and *P. findens*, show little or no distinction in the coloration of the spathe tube and spathe blade, other species have distinctly different colors. In some cases, e.g., *P. antioianum*, *P. cretosum*, *P. edensdatum*, *P. findens*, *P. grandipes*, *P. lazorii*, *P. ligulatum*, *P. llanense*, and *P. roseospathum*, the transition from one color to the next is very gradual. In other cases, such as *P. hebetatum*, *P. immixtum*, *P. jodavisanum*, *P. panamense*, *P. pterotum*, *P. strictum*, *P. radiatum*, *P. rothschuhianum*, *P. scalarinerve*, *P. smithii*, *P. tenue*, *P. wendlandii*, and *P. zhuangum*, the transition is less gradual but by no means abrupt. Still other species, e.g., *P. ligulatum* var. *ovatum*, *P. dodsonii*, *P. gigas*, *P. strictum*, and *P. fragrantissimum*, show a distinct and abrupt transition in the spathe tube and blade colors.

The biological significance of the frequently darker colors on the inside of the spathe tube in contrast to the paler colors of the spathe blade is uncertain. It seems unlikely that either the contrasting colors or the dark color of the spathe tube act in attracting pollinators. The presumed pollinators, dynastine scarab beetles (see section on "Pollination Biology"), are believed to orient more by smell than sight (Faegri & van der Pijl, 1979; Gottsberger & Silberbauer-Gottsberger, 1991). Moreover, their normal arrival time, near dark, would probably preclude their seeing contrasting colors in any event. In addition, the dark coloration is usually restricted to the spathe tube and often not even visible in good light from near the mouth of an open spathe. It is more likely that the dark-colored spathe tube acts to encourage these crepuscular beetles to stay for an extended period of time. Mayo (1986) pointed out that, since neither *P.* subg. *Pteromischum* nor *P.* sect. *Meconostigma* have color contrasts in the spathe blade and tube, *P.* subg. *Philodendron* may have different pollinators. Although not enough pollinators have yet been identified to confirm this possibility, there seems to be little evidence in favor of Mayo's hypothesis (see section on "Pollination Biology"). Some beetle species, such as *Erioscelis proba* Sharp, are known to visit species of both *P.* subg. *Philodendron* and *P.* subg. *Pteromischum* (Grayum, 1996) (see also Table 3, "Pollinators of *P.* subg. *Philodendron*").

Lenticel-like structures on both the exterior of the spathe tube and the peduncle in some *Philodendron* species secrete large droplets of a viscous, somewhat sweet substance. Mayo (1986) has shown that these may consist of clusters of stomata, and

Table 3. Pollinators of *Philodendron* subg. *Philodendron*.

Plant species	Voucher	Beetle species
<i>P. anisotomum</i>		* <i>Cyclocephala amblyopsis</i> Bates * <i>Erioscelis columbica</i> Endrödi
<i>P. brenesii</i>	Croat 35519	<i>Cyclocephala nigerrima</i> Bates
<i>P. brevspathum</i>		<i>Erioscelis proba</i> Sharp
<i>P. callosum</i>	No voucher	<i>Cyclocephala rustica</i> (Olivier)
<i>P. correae</i>	Croat 66653	<i>Cyclocephala conspicua</i> Sharp
<i>P. grandipes</i>	Croat 76594 Jiménez 6 Jiménez 6 Croat 43289	* <i>Cyclocephala gravis</i> Bates <i>Cyclocephala sexpunctata</i> Castelnau <i>Erioscelis columbica</i> Endrödi
<i>P. grayumii</i>	Croat 74840 Croat 74840	<i>Cyclocephala rubescens</i> Bates <i>Cyclocephala sexpunctata</i> Castelnau
<i>P. jodavisiianum</i>	Croat 35950 Croat 35950	<i>Cyclocephala ligyrina</i> Bates <i>Cyclocephala mafaffo</i> Burmeister * <i>Erioscelis columbica</i> Endrödi
<i>P. pearianum</i>	Ramírez 1163	<i>Cyclocephala rustica</i> (Olivier)
<i>P. pterotum</i>	Croat 10903	* <i>Cyclocephala ampliata</i> Bates * <i>Cyclocephala amblyopsis</i> Bates <i>Cyclocephala ligyrina</i> Bates * <i>Cyclocephala sexpunctata</i> Castelnau
<i>P. radiatum</i>		* <i>Cyclocephala ampliata</i> Bates * <i>Erioscelis columbica</i> Endrödi * <i>Cyclocephala amblyopsis</i> Bates <i>Cyclocephala ligyrina</i> Bates <i>Cyclocephala kaszabi</i> Endrödi
<i>P. rothschildianum</i>	(Young, 1987)	* <i>Cyclocephala amblyopsis</i> Bates <i>Cyclocephala kaszabi</i> Endrödi * <i>Erioscelis columbica</i> Endrödi
<i>P. sagittifolium</i>	Thompson 4636	<i>Cyclocephala sexpunctata</i> Castelnau
<i>P. schottianum</i>	Croat 36110	<i>Cyclocephala melane</i> Bates
<i>P. tripartitum</i>		* <i>Cyclocephala amblyopsis</i> Bates * <i>Cyclocephala kaszabi</i> Endrödi * <i>Erioscelis columbica</i> Endrödi
<i>P. tysonii</i>	Croat 66711	<i>Cyclocephala nigerrima</i> Bates

\*Entries were provided by Helen Young, Barnard College, New York, and were based on unpublished observations made during 1984–1985 at La Selva (O.T.S. Field Station) in Costa Rica (with the assistance of George Schatz). Entries without an asterisk were based on determinations made by John Rawlins at Carnegie Museum in Pittsburgh. Collections by S. Thompson were contributed independently to Rawlins.

pointed out their possible ecological significance as extrafloral nectaries.

The secretion of resin from the inner surface of the spathe is apparently unique, in Araceae, to *Philodendron* (Mayo, 1991). Mayo (1986) enumerated four different types of inflorescence resin canals, at least two of which occur among Central American *Philodendron*. One type, found in *P. tripartitum*, has large-diameter resin canals imbedded just beneath the epidermis on the inner surface of the spathe. Resin is secreted directly onto the surface. Another type, represented by *P. smithii*, lacks resin canals on the spathe but instead has resin canals in the staminate zone of the spadix.

The two other types of resin canals pertain to *P.* subg. *Meconostigma* and *P.* subg. *Pteromischum*. In the former, exemplified by *P. bipinnatifidum* Schott ex Endl., the resin canals are "J"-shaped and tangential, arising in the aerenchyma of the spathe and extending to the surface. In the latter, exemplified by *P. sonderianum* Schott, the resin canals are evenly distributed between the aerenchyma and epidermis and are parallel to the surface of the epidermis rather than arising in a J-shaped fashion.

Resin canals usually are present in the proximal 1/2 to 3/5 of the spathe blade and sometimes also the distal portion of the tube. In *P.* subg. *Philodendron*, these resin canals are generally reddish, red-pur-

ple, or orange to brownish and generally somewhat intermittent, as in *P. sulcicaule*, though sometimes continuous, as in *P. immixtum*, *P. mexicanum*, *P. sagittifolium*, and *P. wilburii*. They are always oriented vertically, paralleling the veins of the spathe. The resin canals actively secrete a resinous liquid at anthesis that wells up onto the surface of the spathe and sometimes runs down the spathe below the resin ducts. The resin is generally very sticky and probably functions in causing pollen, sometimes itself not particularly sticky, to adhere to the bodies of the beetle pollinators as they leave the spathe.

## SPADIX

*Philodendron* species are monoecious, with the spadix bearing naked unisexual flowers. The flowers lack a perigon and are closely arranged on a more or less cylindrical spadix, with the staminate flowers comprising the upper  $\frac{3}{5}$ - $\frac{4}{5}$  of the spadix and the pistillate flowers comprising the remaining basal portion. The average spadix for *P.* subg. *Philodendron* in Central America has the staminate portion 2.2 times longer than the pistillate.

The spadix is usually contained for the most part inside the spathe at anthesis (see section above on the Spathe), although in the majority of species the spadix is weakly exerted in front of the spathe. The majority of plants at anthesis have their spathes erect or tipped slightly forward so they can provide shelter to spadices from water dripping straight overhead.

Spadices of some species, especially those with broadly opened spathes (e.g., *P. angustilobum* (Fig. 51), *P. ligulatum* (Fig. 266), *P. mexicanum*, and *P. tripartitum*) protrude prominently forward at anthesis. Other species whose spathes remain convolute to about the middle may also have spadices that protrude prominently forward; these are: *P. auriculatum*, *P. davidsonii*, *P. grayumii*, *P. hebetatum*, *P. heleniae* (Fig. 229), *P. jodavianum*, *P. ligulatum*, *P. llanense* (Fig. 268), and *P. panamense* (Fig. 299).

The spadix of *Philodendron* is usually only slightly shorter than the spathe. The average length of the spathe is 16.5 cm, while the average spadix is 15.8 cm long. Thus on average, the spathe is nearly 1 cm longer than the spadix. This difference may be as little as a few millimeters or as much as 4 cm.

The general morphology of the spadix of *Philodendron*, though highly variable in detail, is more or less the same for all species in Central America. The pistillate portion is usually pale green to green-

ish white and obliquely attached to the spathe, and sometimes markedly stipitate at the base. The floriferous part of the spadix is thus longer on the adaxial (front) than on the abaxial (back) surface (Fig. 40). The differences in the length of the adaxial and abaxial surfaces of the pistillate portion of the spadix depend on the angle at which it is attached to the spathe. These are referred to as "front side" and "back side" of the spadix in the descriptions presented in this revision.

The staminate portion of the spadix characteristically has a swollen sterile section at the base. Typically the staminate portion is broadest at the sterile section and gradually constricted just above this. Above the constriction the spadix is fertile and is gradually broadened usually to a point about  $\frac{3}{5}$  of its length, then narrowed gradually or abruptly to the tip.

The staminate portion is usually uniformly white to creamy white on the outside, although sometimes with the axis pinkish. Immature spadices are usually green, though they become white well before anthesis. The sterile section of the staminate portion is commonly a different color than the fertile section. Often it is more nearly white at an earlier stage of development when the fertile flowers are typically green. At anthesis the sterile staminate portion is more likely to be tinged with yellow or tan, reflecting the higher concentration of oil in these flowers than in the fertile flowers (see below). However, sometimes at anthesis the sterile and the fertile staminate flowers are identical in color. The sterile staminate portion frequently can be distinguished easily after anthesis with the sterile portion maintaining a white color (Fig. 28) while the fertile portion turns grayish or brownish. The sterile staminate portion can usually be distinguished even in these cases by the difference in the size of the spadix (usually larger than the adjacent fertile staminate portion), by the usually larger and more irregularly shaped flowers, and because the sterile staminate section typically ends just before the narrowly constricted portion of the fertile staminate portion (which itself coincides with the constricted portion of the spathe). The sterile staminate flowers are often more easily discernible on dried specimens than they are on live ones since they tend to dry a different color, usually somewhat more brownish.

The much higher oil content of the sterile staminate portion was reported by Pohl (1932), and it is probably owing to this high oil content that the sterile staminate flowers are often eaten by the beetle pollinators (Fig. 29). A high concentration of

lipids in the mitochondria of the staminodial cells has been documented (Walker et al., 1983).

The fertile staminate portion of the spadix is typically more or less clavate and bluntly tapered toward the apex. In a few species, such as *P. advena*, *P. crassispatum*, and *P. davidsonii*, it is more nearly oblong-ellipsoid, often broadest at the base and bluntly tapered toward the apex. The spadix of *P. ferrugineum* is similarly shaped but is somewhat more constricted above the sterile staminate portion. Some species, e.g., *P. heleniae*, have the spadix broadest in the upper  $\frac{1}{3}$  rather than in the sterile staminate portion.

**Male flowers.** The androecium of *P.* subg. *Philodendron* is truncate at the apex, prismatic to obpyramidal and usually irregularly 4–5-sided (Fig. 30). It consists of 2–6 sessile stamens (mostly 3–4). These are always distinct for most of their length although often weakly fused at the base. Stamen number varies within a single spadix and is never constant (Mayo, 1986). Despite the clear grouping of stamens it is sometimes difficult to discern clear floral groupings, especially when the stamens are distinct. Mayo (1986) reported that some species have stamens dehiscent laterally rather than extrorsely.

The androecia in *P.* subg. *Philodendron* range from 2 to 6 mm diam. and show little interspecific variation. Stamens of all species studied by Mayo (1986) had druses in their apices. Anthers are sessile to subsessile with a thick connective that is truncate at the apex and overtops the thecae. The connective of all species studied by Mayo (1986) had raphides present. The thecae are ellipsoid, oblong or linear, emarginate at the base, and each opens by a short lateral slit or subapical pore. The two thecae of a stamen are generally adjacent in *P.* subg. *Philodendron* but positioned far apart in *P.* subg. *Pteromischum* and *P.* subg. *Meconostigma*. Anther thecae lack cell wall thickenings in the endothecium in *P.* subg. *Philodendron* (French, 1985).

Pollen typically emerges in long, slender, somewhat viscid filamentous strands (Fig. 31) (see section on "Pollination" for a description of its emergence). These slender strands of pollen do not persist for many hours, and ultimately the pollen becomes matted in irregular clusters (Fig. 391, *P. straminicaule*).

In *P.* subg. *Philodendron* resin canals in the staminate portion of the spadix are situated beneath the stamens and secrete resin onto the surface of the stamens. The same is not true of *P.* subg. *Pteromischum* and *P.* sect. *Meconostigma*, where the

resin canals are borne at a deeper level in the axis and do not secrete resin onto the surface of the stamens (Mayo, 1986) (Fig. 128, *P. crassispatum*).

French (1986a) reported that the stamen vasculature of most species of *P.* subg. *Philodendron* consists of a single forked bundle with widely divergent branches. Carvell (1989) reported that a single unbranched trace supplies each stamen. According to Mayo (1986) the staminodia have a more distinctive floral receptacle than do the fertile staminate flowers and often have a multiple-branched vascular trace, whereas the latter have an indistinct floral envelope and only a once-branched vascular trace. In contrast, members of *P.* subg. *Meconostigma* have traces that lack branches or have only short branches that spread at an acute angle.

Stamens in *P.* subg. *Philodendron* have both druses and raphide idioblasts with secondarily thickened walls and tanniferous idioblasts occurring throughout the ground tissue (Carvell, 1989).

#### POLLEN

Anthers of *Philodendron* (Grayum, 1991) have a two- or more-layer tapetum of a periplasmoidal type. Pollen mother cell cytokinesis is probably successive.

*Philodendron* pollen (Fig. 30) is binucleate (Zavada, 1983; Grayum, 1985, 1986, 1992a), inaperturate, starchy and of moderately large size (averaging 40  $\mu\text{m}$ , ranging 28  $\mu\text{m}$ –40  $\mu\text{m}$ ), with subsipolar polarity (Grayum, 1985, 1991). As in most aroid genera, it is shed in monads. Pollen is typically boat-shaped-elliptic to oblong, or occasionally elongate as in *P. radiatum*. It is usually round in cross section, but may be very obscurely keeled in *P. hederaceum* to moderately keeled in *P. jacquinii*, *P. jodavisanum*, and *P. wendlandii*, or prominently keeled in *P. mexicanum*. The exine sculpturing is usually psilate, but sometimes minutely verruculate, scabrate, or foveolate (*P. fragrantissimum*, *P. grandipes*, *P. jacquinii*, *P. jodavisanum*, and *P. pterotum*) to punctate, subfoveolate, subfoveate, or subverrucate (*P. mexicanum* and *P. wendlandii*).

#### FEMALE FLOWERS

The pistils of *P.* subg. *Philodendron* are closely aggregated on the spadix in a series of irregular spirals. Gynoecial characters have long been considered important in the subgeneric classification of *Philodendron*, and lobed stigmas were used as early as 1832 by Schott in the recognition of his greges *Meconostigma* and *Sphinctrostigma* (the latter now a synonym of the former). The number of

ovules per locule was used by Engler (1878) in part to characterize the two largest sections of the genus, *Oligospermum* Engl. and *Polyspermium* (now *Calostigma* and *Philodendron*, respectively) (Mayo, 1990).

The pistillate flowers consist of a single naked pistil lacking staminodia. Typically the ovaries are ovoid to obovoid or elliptic, and terete in cross section, or with the sides often somewhat irregularly angular by compression owing to their close proximity with adjacent pistils. Embryo sac development is of the *Polygonum*-type (Grayum, 1991). Each ovary for the genus is syncarpous, superior, and contains 2–47 locules (4–10 in *P.* subg. *Philodendron* of Central America). Ovaries range in size from 0.5 mm (as in *P. sousae*) to 9.2 mm long (as in *P. advena*). The locules are typically oblong, with thin translucent walls that extend  $\frac{2}{3}$ – $\frac{3}{4}$  the length of the ovary. The style is barely distinguishable from the remainder of the ovary, mostly by being slightly thicker and opaque, rather than translucent. While Dahlgren and Clifford (Dahlgren et al., 1985) reported no style in Araceae, both Eyde et al. (1967) and Mayo (1986, 1989) indicated that the region has a distinct anatomy. Mayo (1989) defined the style of *Philodendron* as "that portion of the gynoecium between the base of the stigmatic epidermis and the ovary locules." Slender, conspicuous styles are rare in *Philodendron*, but do exist, as in *P. jacquinii* (Fig. 242).

Each carpel is connected to the stigma apex or compitum (common styler canal) by a styler canal. The compitum (Endress, 1982) is a cavity or complexly shaped channel into which the pollen may be inadvertently packed by the beetle pollinators. This no doubt allows for enough grains to be left to insure pollination, and to make sure not all are removed from the stigma by movements of the beetles. The stigmatic epidermis extends into the compitum from the stigma apex (Mayo, 1989). At the base of the channel or cavity there is a ring of holes that leads into the styler canals. Where no compitum is present, such as with "Type B" and "Type D" styles (see section on "Style Types" below), the styler canals lead directly onto the surface of the style. These canals are readily visible on the dried stigmas of many species if the preservation is adequate. They are particularly easy to see in fruiting collections.

The gynoecium has a separate styler canal for each carpel (see fig. 1 in Mayo, 1989). Each canal may open at its upper end into a compitum. A compitum is rare among Central American species (see section on Style Types below), but is present in *P. correae*, *P. ligulatum* var. *heracleoanum*, *P. smithii*,

*P. straminicaule*, and *P. warszewiczii*. A compitum has also been seen on an unusual collection of *P. radiatum* and some populations of *P. tripartitum*.

The vascular anatomy of *Philodendron* ovules has been studied by French (1986b). Of the five species he studied, three were South American while two, *P. jacquinii* ("*P. hederaceum*") and *P. inconcinnum* ("*P. immixtum*"), occur in Central America. French reported a single vascular trace for all of these except *P. jacquinii*, which has multiple traces. Carvell (1989) reported that the vasculature of the gynoecium in *Philodendron* arises from a single trace, which divides centrifugally to yield a single branch trace for each locule. The carpel traces branch once to form a connection with the placenta as well as a single dorsal trace. The placental trace forms individual connections with each ovule (Carvell, 1989).

Finally, a few miscellaneous anatomical features of the gynoecium should be mentioned. Both tanniferous idioblasts as well as raphide idioblasts are lacking in the gynoecium of *Philodendron*, but the gynoecium does contain druse idioblasts (Carvell, 1989). Like most Araceae, *Philodendron* has unicellular ovular and placental trichomes (French, 1987b). These function in secreting mucilage presumably for the protection of ovules.

Ovules of *Philodendron* are bitegmic, with the inner integument forming the micropyle. The integuments are usually completely free from one another (Grayum, 1991).

Seemingly the most important morphological features of the pistil from the standpoint of systematic importance are the quantity and distribution of the ovules and the type of style, discussed in subsequent sections.

**Placentation.** Placentation type for *P.* subg. *Philodendron* in Central America is usually axile or sub-basal, with 50 species having sub-basal placentation and five species with basal placentation. Those with basal placentation are: *P. dwyeri*, *P. granulare*, *P. smithii* (also sometimes sub-basal), *P. sousae* (also sometimes sub-basal), and *P. zhuangum*. Forty-seven species have axile placentation and two additional species, *P. ferrugineum* and *P. sagittifolium*, with mostly sub-basal placentation, sometimes have axile placentation when they have especially numerous ovules per locule.

Some general rules regarding placentation type and number of ovules per locule are the following: species with basal placentation have only a solitary ovule per locule and tend to have a few large seeds per berry. Species with sub-basal placentation usually have only 1 to a few ovules per locule (aver-



aging 2.1 to 3.5 respectively for the minimum and maximum number in the range) and typically fewer than 6 but rarely to 10 or to 12 (as in *P. brenesii* and *P. davidsonii* subsp. *bocatoranum*). Species with basal and sub-basal ovules tend to have rather large seeds as well, although sometimes their seeds are quite small. On the other hand, species with axile placentation usually have 10 or more ovules per locule (averaging 14–18 respectively for the minimum and maximum in the range) but sometimes as few as 3 (*P. dressleri* and *P. warszewiczii*), or 4.

For those species with only a few ovules per locule, the disposition of the ovules seems to be unorganized or digitate, but for those with more than a few ovules, the placentas may be uniseriate or more generally biseriate or, less frequently, in 1–2 or 2–3 series (see Appendix 2, "Technical Data on Pistils"). Mayo (1989) reported that, based on his studies of the gynoecial morphology and anatomy of 15 mostly South American *Philodendron* species, the placentas of even those species whose ovules appear to be in a single row might actually be biseriate, with the funicles inserted alternately along the placental ridge. This is also true in Central American species. Forty Central American taxa of *Philodendron* proved to have biseriate placentas, while much smaller numbers appeared uniseriate (13), 1–2-seriate (9), 2–3-seriate (11), or digitate (4). Twenty-six species lacked any appreciable organization owing to the small number of ovules, and one species, *P. niqueanum*, was not studied due to inadequate material.

*Ovary locule number.* Mayo (1991) reported that ovaries in *P. subg. Meconostigma* range from 3- to 47-locular, and Bunting (1986) reported 2 (rarely 3) locules per ovary as characteristic of *P. sect. Philpsammos*.

Though the number of ovules per ovary is highly significant in separating *P. sect. Philodendron* and *P. sect. Calostigma*, the number of locules is not a good indicator of relationship. While the average number of locules per ovary has a slightly lower range for *P. sect. Philodendron* (4.6–6.1 vs. 5.5–7.4 for *P. sect. Calostigma*), the difference is not great. In *P. sect. Calostigma* the number of locules per ovary ranges from 1 to 10. In *P. sect. Philodendron* it ranges from 3 to 10. While *P. sect. Philodendron* rarely has more than 8 locules per ovary, and *P. sect. Calostigma* rarely has fewer than 5, more species (14 in all) have 8 locules per ovary than any other number.

No species of *P. subg. Philodendron* in Central America were found to have bilocular ovaries,

though two species, *P. sagittifolium* and *P. zhuangum*, may have only a single locule. In the latter species, the number of locules ranged only from 1 to 3, but in *P. sagittifolium* the number of locules ranged up to 8. Often such low locule numbers occur only near the base of the spadix, where the ovaries are often somewhat irregular in shape. Frequently, the lowermost pistils on the spadix are larger or smaller than those in the middle of the spadix and are more widely spaced and of irregular cross-sectional shape. Three-locular ovaries are rare, known only in two species: *P. copense* and *P. sagittifolium*. Five other species, *P. chirripoense*, *P. hederaceum*, *P. immixtum*, *P. morii*, and *P. squamicaule*, may rarely have 3-locular ovaries.

Four-celled ovaries are fairly common in *P. subg. Philodendron*. Four species, *P. cretosum*, *P. jacquinii*, *P. knappiae*, and *P. subincisum*, so far have demonstrated only 4-locular ovaries. Most species with 4-locular ovaries also have 5-locular ovaries on the same spadix. Four species showed exclusively 5-locular ovaries. Only eight species have locule numbers greater than 10; all but one of these are in *P. sect. Calostigma*.

*Ovules per locule.* The number of ovules in each locule varies from 1 to numerous. Mayo reported numbers as high as 51 for *P. subg. Meconostigma*, but my studies of *P. subg. Philodendron* show the highest number found for any Central American species is 36, in *P. fragrantissimum*. Only 4 other species, *P. antioquianum*, *P. panamense*, *P. squamipetiolatum*, and *P. verrucosum*, have 30 or more ovules per locule, and 17 species have 20–29 (see Appendix 2, "Technical Data on Pistils"). Fifty-six species (62 taxa) have fewer than 10 ovules per locule.

Along with leaf morphology, the number of ovules per locule has long been used as a principal means of assigning sectional affinity in *Philodendron* (Engler, 1878, 1879, 1899; Krause, 1913; Mayo, 1989). Central American members of *P. subg. Philodendron* generally fall into three categories in terms of number of ovules per locule. Many species, including most members of *P. sect. Calostigma*, *P. sect. Tritomophyllum* and *P. sect. Boursia*, have one to few (rarely to 4 or 5, but always with basal or sub-basal placentation). Forty-four species and four subspecies belong to this category. A second group, with axile or sub-basal placentation, generally has 4–10 ovules per locule, but sometimes up to 14. Nineteen species fall into this category. The third group has exclusively axile placentation, mostly with many (15+) ovules per locule. These groups may not be natural, since two

species (*P. davidsonii* and *P. roseospathum*) have one variety in each of the aforementioned groups.

It might be expected that, since the pistils of different species of *Philodendron* are roughly the same size (mostly ranging from 1 to 4 mm long and about  $\frac{1}{2}$  to  $\frac{1}{3}$  as wide), the size of the ovules might be larger in those species with fewer ovules per locule than in those with more. Though there are differences in ovule size (see Appendix 2, "Technical Data on Pistils"), they are not great. Of 44 species of *Philodendron* studied that had axile placentation, the ovules ranged from 0.1 to 2.1 mm long. Alternatively, 48 species with basal or sub-basal placentation had only slightly larger ovules, ranging from 0.31 to 0.5 mm long.

Ovules in *Philodendron* are hemiorthisotropous or orthotropous (Grayum, 1991), rarely hemianatropous (Mayo et al., 1997) and bilaterally symmetrical with an eccentric attachment of the funiculus (French, 1986b). The funicles are usually as long as or longer than the ovules. The ovules that are axile usually have funicles of different lengths, since they are positioned increasingly higher on the axis. After meeting with the wall of the locule the funicle is lightly fused to the wall all the way to the base and can be easily removed intact by pulling it free. This might argue against the assumption by Mayo (1989) that basal and sub-basal placentation were derived from axile placentation, since axile placentation could have developed by a lengthening of the funicular plexus and a fusion to the wall. While no anatomical study was made of this phenomenon, the frequent presence of basal funicles in axile placentas makes it appear that at least in the case of some Central American species of *P. subg. Philodendron*, species with axile placentation may have evolved from species with basal or sub-basal placentas.

The funicles in *P. subg. Philodendron* are frequently densely covered with short, usually inconspicuous, gland-like trichomes from near the base, sometimes extending to about half their total length. The secretory trichomes are continuous with those of the placentas and lining of the styler canals (Mayo, 1989).

Funicles often bear a band of glandular trichomes at or near the base. These were well illustrated by Krause (1913) who reported them commonly in *P. sect. Boursia* and *P. sect. Calostigma* but with a few in *P. sect. Philodendron* as well. These small glands can only be seen under high magnification and probably secrete mucous into the ovary, preventing the ovules from drying out.

Funicles are often fused into a thickened, sometimes ramified, more or less translucent placenta.

The entire placenta and its associated funicles and ovules may be removed, making counting them less difficult. Sometimes, however, adjacent locules share a common trunk so that care must be taken to insure that one is not removing the contents of two locules.

The free portion of the funicle on species with axil placentation seems to be proportionately shorter than the free portion of those with sub-basal placentation. This is perhaps because those species with axile placentation have somewhat longer locules allowing greater spacing of the ovules. Those species with sub-basal placentation and three or more ovules per locule have the funicles arising from one small area near the base of the axile wall, and often have much smaller locules than those with axile placentation. Ovules need to be separated for proper development, and funicles of different lengths allow for this. Thus, in many cases, funicles are longer in those species with sub-basal placentation. The generally smaller locule size for species with sub-basal or basal placentation is confirmed by a survey of locule size. Species with basal or sub-basal placentation have locules ranging in length from 0.32 mm in *P. jefense* to 9.5 mm in *P. warszewiczii*, with the average minimum length 1.9 mm and the average maximum length 3.5 mm. On the other hand, locules range in length from 0.40 (*P. bakeri*, *P. sousae*, *P. sulcicaule*, and *P. wilburii* var. *longipedunculatum*) to 0.7 mm (*P. advena*); the average minimum length is 1.12 mm while the average maximum is 1.75 mm.

The funicles of a single ovary are usually not of equal length if the ovules are basal or sub-basal, since the ovules are usually positioned at slightly different heights off the floor of the locule. Even when the ovules are scattered along much of the length of the axis of the locule wall, the funicles are of slightly different sizes.

**Style and stigma morphology.** This treatment follows the classification of styles and stigmas of Mayo (1989). Mayo (1986) defined the style in *Philodendron* as "that portion of the gynoecium between the ovary locules and the base of the stigmatic epidermis." Although there is considerable diversity at the microscopic level, much of the detail of the style is easily visible only by dissection (see section entitled "Style Types"). The fresh styles of *P. subg. Philodendron* are relatively uniform microscopically. Usually they are hemispherical or sometimes globose or depressed-globose. The microscopically visible portion is the stigma, a "single continuous area in which the epidermal cells are greatly elongated into secretory papillae" (Mayo, 1986). This

dense layer of stigmatic papillae is so closely packed and so engulfed in a gelatinous fluid as to appear almost solid. Still, a needle can be passed easily across and through much of its volume in any direction without disturbing its shape. The stigmatic papillae are slender, many times longer than broad, and attached to the surface of the style. Mayo (1986) reported that the stigmatic papillae often contain tannin cells.

The stigmatic papillae typically dry to form a thin, flat, often translucent, wafer-like mantle around the now somewhat visible style apex. In time the stigmatic mantle may fall off altogether allowing easy inspection of the style apex. Sometimes the stigma dries erect and disheveled. In these cases the appearance of the stigmas is more difficult to determine but, regardless of how the stigma dries, the style apex often can be made easily visible on a dried specimen by lightly scraping away the stigmatic papillae. The complexities of the style types of *Philodendron* are discussed below.

**Style types.** The present work adopts the gynoeical classification of Mayo (1989) in which six distinct style types were described and illustrated (Fig. 469, "Style Types in Central America"; see Appendix 2, "Technical Data on Pistils," which summarizes the style types for *P.* subg. *Philodendron* of Central America). Only two species, *P. niqueanum* and *P. utleyanum*, have style types not yet known. The table also includes the number of locules per ovary, placentation type, number of ovules per locule, ovary size, disposition of ovary, and the nature of the ovular sac when present.

Style type is one of the most important gynoeical features in *Philodendron*. While the stigma is moderately uniform in its superficial appearance, the style is highly diverse in morphological features. Unfortunately, these features are largely hidden by the stigmatic papillae when the pistil is fresh. Despite the shape of the style, the stigmatic papillae, which cover all or part of the style, may form a stigma of more or less uniform shape.

**Style Type A** has a compitum (common funnel into which the pollen may be packed, defined as the space between the upper stigmatic papillae and the level at which the stylar canals emerge onto the style surface) with ridged inner walls and a lobed apex, with each lobe corresponding to the apex of one carpel. Style Type A is restricted to *P.* subg. *Meconostigma* and thus will not be considered further here.

**Style Type B** (Fig. 469A) lacks a compitum. Instead, the stylar canals open into relatively

broad concavities on the style apex. These concavities are arranged in a ring with one hole per locule. Style Type B also has the stylar canal entering directly into the apex of the locular cavity. Although the style apex may be completely flattened or broadly concave, it is sometimes weakly ridged between the apertures of the stylar canals. These ridges meet in the middle of the style and may even form a weak central beak. Some styles also have well-developed stigmatic papillae associated with the stylar canals causing the surface to be at least weakly lobed, with a single lobe for each locule. In dried condition Style Type B sometimes appears as a button-like structure, somewhat resembling Style Type D. It is therefore important in determining style type to make comparisons of material at or near anthesis. Style Type B seems to be most closely related to Style Type D and shares with that type the relatively large stylar canal pores on the surface of the style apex relatively near its periphery. Style Type B is the most common type in Central America, known in at least 70 species. One species, *P. tripartitum*, though usually having Type D styles, also has Type B and E styles in some populations of the species (see that species for a discussion of its style types).

**Style Type C** (Fig. 469B) is characterized by being decidedly concave or funnel-shaped at the apex with no lobes on the margins of the rim and with the stylar canals arising in a narrow cluster at the base of the compitum (funnel). Since the stylar canals are closely clustered there is no central dome (defined as any stylar tissue that lies above the level at which the stylar canals emerge onto the style surface). In contrast to style Types B and D, both of which have rather prominent stylar pores, the stylar pores of Type C are small, sometimes barely visible, and in a generally smaller circle nearer the middle of the style apex. This style type is rare in Central American *P.* subg. *Philodendron*, and is known primarily in *P.* subg. *Pteromischum*. Only five taxa of *P.* subg. *Philodendron*, (*P. correae*, *P. cotonense*, *P. ligulatum* var. *heracleioanum*, *P. stramineale*, and *P. warszewiczii*) have exclusively Type C styles. Though their pistils are funnel-shaped at the apex, these species have funnels generally not as deep as those of *P.* subg. *Pteromischum* as illustrated by Mayo (1989). One other species, *P. radiatum*, has at least one collection (Croat & Hannon 63414) with the

style type that also has a funnellform apex and looks like a Type C style. While the dried style of this collection is distinct and button-like, the pores are central in a shallow concavity.

**Style Type D** (Fig. 469C) is similar to Type B in that it lacks a compitum and has thick styler canals emerging in a circle on a flat styler apex relatively close to the margin of the stigma apex. It differs from Type B in that the style is constricted around the circumference to form a protruded flat "style boss" (defined as a  $\pm$  domed, circular, stigma-bearing projection that extends beyond the main part of the style and is separated from it by a short neck) that rises above the general level of the style apex. Thus the style appears to have a short flat neck at the apex. It is from the "style boss" that the styler canals emerge. The styler pores are relatively large and borne relatively near the margin of the style apex. At least 23 species of *P.* subg. *Philodendron* in Central America have Style Type D, the second most common style type.

It is easy to confuse or misinterpret Style Types B and D if the specimens are not well preserved, especially if the material studied is not fresh but rather rehydrated. Species with Type B styles sometimes have styles that dry with a button-like apex resembling that of Type D. At least one collection of *P. advena* has both Type D and Type B styles. Some populations of *P. tripartitum* have not only Type D styles but Type B and Type E as well (see the discussion of that species for details).

**Style Type E** (Fig. 469D) has a slender funnel-shaped to cylindrical compitum with a distinct raised annulus around the upper rim. The styler canals arise in a small cluster at the base of the compitum just as in Type C styles. The latter differ, however, in lacking the rim on the style apex. Type E styles are rare in Central American *P.* subg. *Philodendron*, found in only *P. granulare* and perhaps *P. smithii*. *Philodendron granulare* has such an unusual form of the Type E style that it should perhaps warrant its own status. In *P. granulare* the annulus of the styler funnel actually protrudes well above the surface of the style (at least in its dried state). *Philodendron smithii* was reported by Mayo (1989) as having a Type E style, but no rim is obvious in fresh material of the species. It is more appropriately a Type C style. *Philodendron tripartitum*, though usually with Type D styles, has Type E styles in

some parts of its range (see the discussion following that species for details).

**Style Type F** is narrowly funnel-shaped with a small dome at the base of the compitum around which the styler canals arise. This type is not known among the Central American *Philodendron*. It is known only from *P. burlemarxii* G. M. Barroso, a member of *P.* sect. *Boursia* (Mayo, 1989) from Amazonian Brazil.

Mayo (1989) has shown that the course of individual styler canals is correlated with the type of placentation. Those species with basal or sub-basal ovules, such as *P. tripartitum* and *P. smithii*, have styler canals that course down the center of the axis of the pistil and enter the locule near its base. Those species with axile placentation have styler canals that also course down the pistil axis and enter the locule somewhere above the base of the locule, but only rarely at the very apex. The styler canals are lined with papillose epithelial cells, which are contiguous with the stigmatic epidermis (Mayo, 1989). These secrete the gelatinous mucous that keeps the stigmatic surface moist.

#### FRUITS AND SEEDS

In *Philodendron* the developing pistils remain within the reclosed spathe after anthesis and pollination until they are fully ripe [except in the rare case of *P. surinamense* (Miq.) Engl., a South American member of *P.* subg. *Pteromischum* that promptly loses its spathe after anthesis]. The spathe enlarges somewhat to accommodate the enlarging berries. When the berries are mature the spathe once more begins the process of reopening, but it breaks completely free at the base where it is obliquely attached to the peduncle. Sometimes the spathe falls completely free after loosening even before it opens, but in general it breaks up beginning at the base and falls off (Fig. 33) eventually falling completely free and leaving a scar just above the peduncle (Fig. 34). Generally the old, withered staminate portion of the spadix falls free at this time as well and the berries are exposed on the remaining pistillate portion of the spadix (Fig. 33, *Philodendron hebetatum*). In the process of unfolding, the spathe often develops deep longitudinal fissures, which apparently enable it to unfold. Sometimes the old persistent spathe persists on the peduncle with fragments of the inner surface exposed (Figs. 35, 36).

The berries of *Philodendron* are cylindrical to obovoid, generally with a thickened cap-like apex and not markedly colored, though berry colors from pale yellow to bright orange or even red to purple

do exist (see section on berry color). Berries in *P.* subg. *Philodendron* are universally soft and fleshy except for the frequently thickened apex. The seeds can be seen easily through the sides of the berries. When fully mature the apical portion of the berry is easily torn free, and the thin, fragile sides of the berries are easily ruptured (see section on seeds).

Though little is known about fruit dispersal, the mesocarp surrounding the seeds contained within each locule is juicy or gelatinous and is usually sweet and sticky, making it logically animal dispersed. Inflorescences that appear to have been pecked apart by birds (Fig. 36) are frequently seen. Certainly the sticky seeds, often many per berry, would logically be easily dispersed on birds' beaks. Alternatively, the infructescence is large, and even faintly scented when fully mature, making it an appealing meal even for mammals such as monkeys. Grayum (1996) theorized that those species of *P.* subg. *Pteromischum* with whitish fruits, which may produce a garlic-like or pepper-like odor at night, are dispersed by bats. Those with orange fruits may be dispersed by diurnal animals. Certainly the manner in which many species of *P.* subg. *Pteromischum* flower, e.g., on the ends of short, spreading branches some meters above the ground, would make them superbly positioned as bat fruits. There are also species of *P.* subg. *Philodendron*, such as *P. lentii*, which have their branches held in a similar manner.

Ants are also probable dispersers of *Philodendron* seeds. I have seen two different species of ants carrying away individual seeds of *Philodendron*. Ant dispersal is certainly important for those species, such as *P. megalophyllum* in South America, that live almost exclusively on ant nests. One cultivated individual of that species even set fruit repeatedly without being pollinated. Doing so in nature would provide a steady, abundant supply of berries for its ant dispersers and assure the species widespread dispersal. Indeed, the species is particularly successful even in areas of white sand soil where soil nutrients are very low.

Although berry color is known for only a relatively small number of *Philodendron* species (53 out of 96), some general comments can be made. Berry color in *Philodendron* is not so important taxonomically as it appears to be in *Anthurium* (Croat, 1983a, 1986a, 1991). Grayum (1996) reported that for *P.* subg. *Pteromischum* the fruits sometimes provided taxonomically significant characters, with species related to *P. inequilaterum* having orange fruits while other species have whitish fruits.

Mature berries of most *Philodendron* species are generally described as "white," but are usually

more nearly ochraceous or somewhat greenish white. A total of 33 species have either white, creamy white, or greenish white fruits at least some of the time. In some species, such as *P. advena* and *P. smithii*, the whitish berries turn somewhat yellowish when fully mature, and the mature fruits of *P. brevispatum* are pale yellow. The berries of *P. sagittifolium* are usually pale yellowish but rarely have been reported as orange. While the majority of *Philodendron* berries are whitish, 17 species of Central American *P.* subg. *Philodendron* are known to have berries at least sometimes pale orange to orange. An additional three species have yellow or yellowish berries. *Philodendron fragrantissimum* has bright red to purple-red berries.

Known species with orange fruits are members of *P.* subsect. *Glossophyllum*. These are: *P. auriculatum*, *P. bakeri*, *P. cotonense*, *P. pseudauriculatum*, *P. wendlandii*, and *P. wilburii*. Other species with orange fruits are *P. anisotomum*, *P. breneisii*, *P. crassispatum*, *P. ferrugineum*, *P. lentii*, as well as, at least sometimes, *P. jacquinii* and *P. sagittifolium*. Three of the above, *P. breneisii*, *P. crassispatum*, and *P. lentii*, are members of *P. ser. Ecordata*. Thus, it may be significant that most species with decidedly orange fruits are members of *P. sect. Calostigma*. Even *P. ferrugineum*, with orange berries, and *P. sagittifolium*, sometimes with orange berries, are members of *P. sect. Calostigma*. Only *P. anisotomum* (*P. sect. Tritomophyllum*) and *P. jacquinii* (*P. sect. Macrogynium*) are not members of *P. sect. Calostigma*.

The seeds of *P.* subg. *Philodendron* are few to many per berry and are mostly oblong to oblong-ellipsoid, ellipsoid, ovoid-oblong, or less frequently ovoid. Typically they are minute (see below). The seeds are typically more or less terete, though they may be somewhat flattened, e.g., in *P. findens*. They have a rather thick, smooth (e.g., *P. granulare*) or striate-costate testa, as well as copious endosperm. The longitudinally oriented striations may be weak (*P. advena*, *P. findens*, *P. cotonense*, *P. purpureoviride*, and *P. sagittifolium*) to strong (*P. ferrugineum* and *P. grandipes*). Seeds of *Philodendron microstictum* have both longitudinal striations and much finer cross-etching. A number of species are reported to have seeds with pale raphide cells on the surface. These include *P. hederaceum* and *P. warzewiczii*. Seeds of some species, e.g., *P. anisotomum* and *P. hederaceum*, sometimes have a constriction on the end opposite the funicle.

*Philodendron* seeds are usually dramatically smaller than those of *Anthurium*, which typically have only two seeds per berry. Seeds of species of *P.* subg. *Philodendron* studied (a total of 48 species)

ranged from 0.5 mm to 5 mm and averaged 1.67 mm long; seed diameter ranged from 0.1 to 3 mm, averaging 0.7 mm. The longest seeds were those of *P. jacquinii* and *P. hederaceum*, at 5 mm. Relatively few species have seeds longer than 2 mm: *P. advena*, *P. anisotomum*, *P. findens*, *P. dolichophyllum*, *P. grayunii*, *P. purulhense*, *P. rothschuhianum*, *P. sagittifolium*, and *P. warszewiczii*.

The number of seeds in each locule is often many fewer than the average number of ovules per locule, presumably owing to the fact that not all are properly pollinated. A careful examination sometimes shows the aborted undeveloped ovules that were present at the time of flowering. The percentage of ovules in each locule that develops into seeds varies from species to species. Species of *P. sect. Calostigma* that have only one or a few ovules per locule are more likely to have an equal number of seeds. On the other hand, species in *P. sect. Philodendron*, especially those with large numbers of ovules per locule, rarely develop all their ovules.

Because species of *P. sect. Calostigma* have fewer ovules per locule and thus fewer seeds, one might expect them to have larger seeds. Though admittedly the sample size was small (only 21 species studied for *P. sect. Calostigma* and 17 for *P. sect. Philodendron*), the unexpected results were that the seeds of species of *P. sect. Philodendron* in Central America averaged slightly larger than those of *P. sect. Calostigma* (to 1.69 mm long for *P. sect. Philodendron* and to 1.66 for *P. sect. Calostigma*).

*Philodendron* seeds are largely pale in color, mostly shades of brown, tan, or white, though for most species mature seeds have not been observed. Seeds vary in color from medium green (*P. rothschuhianum*) to whitish (*P. roseospathum*), tan (*P. alticola*, *P. annulatum*, *P. granulare*, *P. morii*, *P. smithii*), yellowish or pale yellow (*P. brenesii*, *P. llanense*, and *P. strictum*), yellow-orange (*P. crassispathum*, *P. purpureoviride*, *P. wilburii*, and *P. mexicanum*), brown (*P. ferrugineum*), or reddish brown (*P. heleniae*). The dried seeds of two species, *P. findens* and *P. jefense*, are described as translucent.

#### POLLINATION BIOLOGY

Although there are frequent insect visitors to *Philodendron*, especially small Hemiptera in the genus *Neelia*, which appear not to feed or mate on the inflorescences (H. Young, pers. comm.), only the larger beetles are known to be pollinators. The system of pollination is nearly identical to that of *Dieffenbachia* (Croat, 1983b; Young, 1986, 1990). Pollinators are members of subfamily Dynastinae in

the family Scarabaeidae (Fig. 32). All determined beetles collected from either Central American or South American *Philodendron* are members of the genera *Cyclocephala* and *Erioscelis*. Some species of beetles are not particularly host-specific, visiting members of both *P. subg. Philodendron* and *P. subg. Pteromischum*; other genera such as *Dieffenbachia*, *Homalomena*, *Syngonium*, and *Xanthosoma*; and sometimes even other families including some palms (Arecaceae), *Cyclanthus bipartitus* Poit. ex A. Rich (Cyclanthaceae), as well as *Annona* and *Cymbopetalum* in the Annonaceae (M. Grayum, pers. comm.; Schatz, 1990). *Cyclocephala negerima* Bates, for example, has been found visiting *P. brenesii* and *P. tysonii*, but also *P. standleyi* Grayum, a member of *P. subg. Pteromischum* (see Table 3). Though beetles are not very species-specific pollinators, individual beetles of some species tend to be somewhat stratum-specific, visiting only those species growing at particular height ranges above the ground (Schatz, 1990; Helen Young, pers. comm.). These beetles are attracted to the *Philodendron* inflorescence, usually late in the day or at dusk. Attractants and/or rewards are apparently a combination of scent (at least in many cases), a source of food (oil-bearing sterile staminate flowers), warmth (thermogenesis), and shelter.

Scents produced by *Philodendron* species are not always obvious, at least in *P. subg. Philodendron*. George Schatz and Helen Young (pers. comm.) have documented floral odors for species of *Philodendron* and have identified the principal constituents of these aromas. Some species have noticeably sweet aromas in the early evening hours, while other species have no noticeable scent, at least during the early evening hours on the first day of anthesis. *Philodendron megalophyllum* (in cultivation at the Missouri Botanical Garden), a South American species, had a faint spicy aroma detectable directly at the spathe during the evening, but even this faint aroma was absent the following morning on day two of the flowering event. At the same time the stigmas were soft, juicy, and sticky but without an obvious flavor or taste both in the evening and the following morning. Schatz (1990) believed that the pattern of visitation to *Philodendron* exhibited by beetles at La Selva in Costa Rica was to a great degree explained by odor. He pointed, for example, to the high degree of specificity exhibited by *Philodendron radiatum* and an undescribed species of *Cyclocephala*. The aroma given off by *P. radiatum* was made up of compounds unique to that species.

At anthesis the open spathe of *Philodendron* provides ample space at the base in the area surrounding the pistillate portion of the spadix. The spathe

blade may provide a certain amount of protection against rain, as mentioned previously. Beetles typically spend the first night and most of the following day inside the spathe where they remove the sticky exudate from the pistils, eat pollen (Gottsberger & Silberbauer-Gottsberger, 1991), feed on the sterile staminate flowers, and mate. Studying *P. bipinnatifidum*, Gottsberger and Silberbauer-Gottsberger (1991) found that the beetles were active mostly during the first 10 to 20 minutes after arrival and during the strongest production of scent. Copulation was most active immediately after arrival, when the spadix was warmest; grazing on the sterile male flowers also subsided by the time the spadix had cooled off. Old inflorescences that have been pollinated often have the sterile staminate portion of the spadix entirely eaten away.

Typically the number of beetles found in a single inflorescence is modest, frequently no more than five and sometimes up to a dozen beetles. However, sometimes the numbers are simply astounding, with Gottsberger and Amaral (1984) reporting as many as 200 beetles in a single inflorescence of *P. bipinnatifidum*. While visiting beetles are often of the same species, sometimes more than one species of beetle may be found in the inflorescence. Though beetles typically do not leave the spathe until about dusk, they will leave if the inflorescence is sufficiently disturbed. When beetles are disturbed, they may crawl slowly up the spathe or spadix and appear at the rim of the open spathe, at which point they generally fall promptly to the ground where they disappear in the leaf litter or soil with remarkable speed.

Thermogenesis, the production of heat in the spadix by the rapid oxidation of stored starch or lipids (Walker et al., 1983; Gottsberger, 1990), plays an important role in the pollination of *Philodendron* (van Herk, 1937a-c; Van der Pijl, 1937; Knutsen, 1974; Seymour et al., 1984). Gottsberger (1984, 1986), studying *P. bipinnatifidum*, has shown that, although oxidation of carbohydrates takes place during preheating of the spadix, lipids are oxidized thereafter during maximum heating and are consumed directly, not after conversion to carbohydrates. This makes the biochemistry of this species similar to that accompanying heat production in some animals. The thermogenic reaction occurs principally in the staminodial region of the staminate portion of the spadix (James & Beevers, 1950; Henry & Nyns, 1975) involving the inner surface of mitochondrial membranes (Urdentlich et al., 1991) and is triggered by an accumulation of salicylic acid (Meeuse & Buggeln, 1969; Raskin et al., 1987, 1989; Meeuse, 1975, 1978; Raskin,

1992). The end result of this high rate of respiration in these plants is the production of heat rather than ATP as in animals (Meeuse, 1966). To accomplish this high increase in metabolism the plant's mitochondria in the inflorescence switch to an electron transport pathway commonly referred to as the "cyanide-resistant pathway."

The thermogenic heat rise in *Philodendron* is sometimes dramatic, with temperatures rising well above ambient temperature (Nagy et al., 1972; Knutsen, 1974). Gottsberger and Silberbauer-Gottsberger (1991), working with *P. bipinnatifidum*, reported temperatures of spadices occasionally to 46°C with the highest and most efficient temperatures for the emission of scents being maintained for 20-40 minutes. Thermogenesis does not create even or constant temperatures, but rather produces fluctuations depending on the time of day with definite peaks (Leick, 1910, 1916; Engler, 1920a, 1920b; Foster, 1949; Nagy et al., 1972; Sheridan, 1960; Gottsberger & Amaral, 1984), the greatest occurring when beetle visitation and odor is most intense (Gottsberger & Amaral, 1984). Temperature peaks may occur on two or more successive days. While the increased temperature is presumably responsible for the production of scent compounds (Nagy et al., 1972), and whereas the production of heat and scent appears closely correlated with peaks in temperature, there is still controversy over the exact function of the heat production at least as it pertains to genera that produce foul odors. Moodie (1976) suggested that heat production and the higher levels of carbon dioxide production are components of a carrion, dung, and mammal mimicry syndrome and that heat production aids in providing sufficient warmth in colder climates for the activity of pollinating organisms. The subject of thermogenesis and its role in pollination has been reviewed in great detail by Mayo (1986), Grayum (1990), and Bay (1995/1996).

The exact role that thermogenesis plays in the pollination of *Philodendron* is still poorly known, and rather few plants have been studied on an experimental basis. Despite its probable occurrence in all *Philodendron* species, thermogenesis has thus been documented only for *P. selloum* K. Koch and *P. bipinnatifidum*, now considered by Mayo (1991) to be synonymous. My own measurements with a recording thermometer on *P. glanduliferum* and *P. advena* in the field in Chiapas, Mexico, indicated a definite heating during the early evening hours, usually peaking between 18:00 and 19:00 hours.

The site of the heat production on the spadix is another possible difference between *P. subg. Meconostigma* and *P. subg. Philodendron*. Mayo's

(1986) investigations on *P.* subg. *Meconostigma* indicated that heat production was centered in the sterile staminate section of the spadix. In *P.* subg. *Meconostigma*, that section is as large as or larger than the fertile staminate part of the spadix, a situation unknown in *P.* subg. *Philodendron* where the staminodial segment is always a small percentage of the total spadix.

Contrary to Mayo, Leick (1916) reported that heating took place in the "middle and upper part" of the spadix in *P. sellowii*, presumably implying at least a part of the fertile staminate portion. Concurring with this view, Ron Weeks (pers. comm.) reported that a perceived temperature rise occurred in both the sterile and fertile staminate spadix portions of those species of *P.* subg. *Meconostigma* that he studied. Mayo (1986) theorized that, because of the major morphological differences in the relative lengths of the sterile and fertile staminate portions of the spadices, the two subgenera would likely have different thermogenetic patterns.

In the Central American *Philodendron* species observed by the author, the spathe opens broadly late in the afternoon of the first day of flowering; it closes slightly the morning after the first night of anthesis. The evolutionary significance of this slight closure is uncertain but, even in the most extreme cases, it occludes only the pistillate portion. The spathe continues to close on the evening of the second day after having been open for about 24 hours (see discussion below). The beetles, which generally enter the spathe on the first night of opening, spend about 24 hours in the spathe tube. This second partial closing process usually corresponds with staminal dehiscence. The now crowded condition of the spathe tube, and the probable desire on the part of the beetles to seek a new food source with the onset of dusk, encourages the beetles to leave the inflorescence. The beetles emerge from the spathe tube by either climbing up the side of the spadix or up the inside wall of the spathe. By the time they emerge from the spathe, it is quite constricted and they must squeeze through the constriction which fits rather tightly around the spadix just above the sterile staminate portion. In order to depart the beetles must literally crawl through the copious strands of pollen that emerge from the apical pores of the stamens. The constriction of the spathe and its corresponding constricted area on the spadix help to insure that most of the pollen that falls into the spathe tube accumulates and is carried forward and out of the spathe by the departing beetles. Araceae pollen is not very tacky and probably does not adhere well to the smooth, hard surfaces of the beetles. However, the beetles

themselves are usually quite sticky from the sugary secretions of the styles and especially from the resin, which arises usually from the inner surface of the spathe or sometimes from the spadix itself (Fig. 128). Once the beetles have emerged they fly off in search of another place to spend the night, usually another open inflorescence. The beetles apparently have a keen perception of infra-red radiation or of scent because they are often seen in a "homing-in" pattern, which is quite direct to the next available inflorescence (John Rawlins, pers. comm.). Gottsberger and Silberbauer-Gottsberger (1991) reported that beetles flew in a zig-zag pattern toward the center of fragrance concentration, indicating that they were very sensitive to the aroma being produced. They also reported that once the beetles were within sight of the inflorescence they shifted to a straight line of flight until they hit the inner surface of the spathe blade, whence they moved into the lower portion of the spathe. Gottsberger and Silberbauer-Gottsberger (1991) have proven experimentally in the case of *P. bipinnatifidum* (a member of *P.* subg. *Meconostigma*) that the beetles use only visual references for location as they near the inflorescence. Shelter, warmth, food, and copulation are the driving forces behind this pollination strategy, and although selectivity is not perfect in such beetle-pollinated systems (Young, 1986, 1988a) fruit-set in undisturbed populations is high. The precision and high degree of synchrony of thermogenesis gives evidence of a highly evolved system of pollination.

Even though all detailed observations thus far have been made with *P. bipinnatifidum*, there is considerable confusion regarding the results. Horticulturist Ron Weeks (Homestead, Florida) reported (pers. comm.) that three members of *P.* subg. *Philodendron*, *P. bipinnatifidum*, *P. speciosum* Schott, and *P. williamsii* Hook. f., showed no variation in the schedule of spathe opening, the capability of the plants being hand-pollinated on the first evening of opening, or in the shedding of pollen on the evening of the second day. On the other hand, he reported that *P. eichleri* Engl. showed great variation in opening periods, temperature changes, fragrance, and pollen shed, perhaps owing to weather conditions. Scientific studies carried out on other plants in *P.* subg. *Meconostigma* showed considerable variability. Four separate and conflicting reports were made on material determined as belonging to *P. bipinnatifidum* in Brazil. Warming (1867, 1883) reported a two-day pollination event with two heat peaks (early evening and late morning, respectively) with the spathe closing then re-opening during the first night. Gottsberger and



Amaral (1984) reported on two plants, one as *P. selloum* (now considered to be a synonym of *P. bipinnatifidum*) with a three-day pollination event with two unequal early evening heat peaks, and one as *P. bipinnatifidum* with a four-day pollination event with three unequal early-evening heat peaks. The spathe was not reported to close during the event. Confirming the complexity of the thermogenesis riddle is the fact that Seymour et al. (1983), studying a cultivated but similar plant believed to be the same species (Mayo, 1986), found both types of pollination events that had previously been described by Gottsberger and Amaral but this time in a single plant. Clearly more investigation must take place, at least in *P. subg. Meconostigma*, to determine the pollination behavior.

Leick (1916), reviewing work done by Kraus (1894, 1896) with *P. bipinnatifidum*, a member of *P. subg. Philodendron* from Venezuela, reported a two-day pollination event with temperature peaks in the evening of two consecutive days. While it is not certain that most Central American *Philodendron* have an elevated temperature on two consecutive days, the general pattern of opening and closing of the spathe and the beetle visitation in *P. bipinnatifidum* would appear to match the events of Central American species observed in the field and under greenhouse conditions at the Missouri Botanical Garden. Further detailed studies of this phenomenon, including a much broader survey of Central American species, will be carried out by my student, Jane Whitehill, during graduate studies at the Missouri Botanical Garden.

Grayum's (1996) observations with *Philodendron* subg. *Pteromischum* showed a similar pattern, with most species having the spathe beginning to loosen by early afternoon and being fully open by mid-afternoon. He reported that for the species of *P. subg. Pteromischum* he observed, the pollinators appeared at the opened inflorescence during a relatively brief time, usually between 19:00 and 19:15 hours. An important feature in the pollination story reported by Grayum (1996) for the first time is that resin secretion from the inner spathe surface does not begin until 21:00 to 22:00 hours on the first day of anthesis, and that it then continues until the end of anthesis.

Once opened, the spathes of *P. subg. Philodendron* apparently remained open during the night and were always open the next morning at the beginning of day two, remaining open during the course of most of the day. During the latter part of the same day, usually in late afternoon of day two, the spathe began to close and pollen began to shed in long filaments. The spathe did not fully close at

this time but remained open near the apex. It remained in this condition into the beginning of the evening of day two. By the beginning of day three the spathe was generally fully closed, and the only evidence that it had ever opened was often some loose pollen remaining on the closed edges of the spathe. In addition, the closed spathe is somewhat less turgid than before anthesis, sometimes allowing it to be forced open without breaking the margins of the spathe. Doing the same with an unopened spathe is impossible without breaking the stiff and brittle spathe margins.

The entire pollination episode usually requires little more than 24 hours, counting just the time that the beetles are present. The time that the spathe is to any extent open could be as much as 8 hours longer, since it may open late in the afternoon and remain open for some hours after the pollen has been shed. Grayum (1996) reported that for the species of *P. subg. Pteromischum* studied in Costa Rica the average pollination event required about 30 hours (i.e., from spathe opening to closing).

That the intensity of light must play an important role in flowering behavior is indicated by the fact that on cloudy days spathes in cultivated collections open earlier than usual, sometimes as early as noon on day one of the flowering sequence. This may support the argument of Buggeln et al. (1971) that darkness induces opening of the spathe and an elevated respiration rate in *Sauromatum tenosum* Kunth.

Armbruster (1984), studying the role of resin in angiosperm pollination, has questioned the efficacy of floral resin in the transport of pollen, citing its possible toxicity and the difficulty of transporting pollen embedded in resin. While he stressed the role of resin for other purposes, mainly in nest-building by bees, it must be pointed out that bees which use resin for nest building play no role whatever in *Philodendron* pollination. In contrast, the general availability of resin, its close association with pollen delivery, the non-tacky nature of *Philodendron* pollen, and the availability of resin only at anthesis all point to a strong role for resin in *Philodendron* pollination. In species with resiniferous spadices (Fig. 128), the pollen is shed with and incorporated in the resin from the moment of theca dehiscence. Alternatively, species that lack staminal resin and instead have resin only on the inner spathe surface have pollen presented as slender filaments.

Breeding studies (see section on Breeding Behavior) have shown that *Philodendron* species have few if any genetic barriers to cross-pollination, ow-

ing perhaps to the fact that there are other physical and temporal barriers to self-pollination. Even when two species of *Philodendron* are in flower simultaneously, there are other parameters that effect separation. Beetles tend to fly at certain elevations above the ground (Schatz, 1990), helping to prevent cross-pollination of species that flower at different strata. In addition, specific attractants, i.e., species-specific pheromones, may exist in some species that attract principally a single beetle species (H. Young, pers. comm.). The sloppiness in the system, when it occurs, is owing to opportunistic beetle visitors (G. Schatz & H. Young, pers. comm.), and this might produce some hybridization.

Perhaps because of the substantial barriers already present, *Philodendron* appears to have developed the ability to cross between sections. In *Anthurium*, relatively little cross-breeding was possible between different sections in the genus (Croat, 1991). In contrast, quite unrelated species of *Philodendron*, even species in different sections, readily cross-pollinate and produce intermediate offspring (Keith Henderson, Cairns, Queensland, Australia, pers. comm.). For this reason pre-zygotic separation may be critical to maintaining distinct species lines. The pollinators of *Philodendron*, dynastine scarab beetles, are for the most part not very species-specific and frequently switch from one species to any other in flower at the same time. Not only will some species of beetles switch from one *Philodendron* species to the next, as is known for certain in *Dieffenbachia* (Young, 1986), but some individuals will also switch to another genus. For example, beetles that regularly visit *D. longispatha* Engl. & K. Krause at La Selva may visit *P. grandipes*, another species that is terrestrial and about the same height above the ground (G. Schatz & H. Young, pers. comm.). Beetles are also reported to move from *Dieffenbachia longispatha* to *Xanthosoma undipes* (K. Koch & Bouché) K. Koch. Some individuals of beetle species that regularly visit *D. longispatha* will even switch to *Cyclanthus bipartitus* Poit. (Cyclanthaceae). Schatz believes that this is owing to the fact a small component of the pheromone emitted by *Cyclanthus* Poit. ex A. Rich is the principal component of the scent given off by *Dieffenbachia longispatha*. He believes that during the end of the flowering season of *Dieffenbachia* and the beginning of the flowering season of *Cyclanthus* some confusion occurs in the pollinators' behavior.

The unpublished observations of G. Schatz and H. Young (pers. comm.), and the published results reported by Helen Young (1986, 1988a, 1988b) principally for *Dieffenbachia*, probably are comparable to what is happening in *Philodendron*. Beetles

that visit any particular species of *Dieffenbachia* are often predominantly of one species, but they are often accompanied by other opportunistic beetles. Schatz believes that these opportunistic species are not likely to be effective pollinators, since they are so catholic in their tastes that they are not likely to make their next visit to a receptive *Dieffenbachia*. Likewise, Helen Young (Young, 1988a) indicated that the most common species of beetles are not the most effective pollinators. However, they may be responsible for the occasional hybrids seen in *Dieffenbachia*, at La Selva, where the studies of both Schatz and Young were carried out. The pollination system described for *Dieffenbachia* by Young and Schatz is apparently similar to that of *Philodendron*. Schatz (pers. comm.) reported that while one undescribed beetle species (determined as *Cyclocephala ampliata* by H. Young), was found to visit only *Philodendron radiatum*, it was accompanied occasionally by another more opportunistic species. Despite the presence of opportunistic beetle species, some beetle pollinators of *Philodendron* species are probably much more species-specific. For example, Grayum (1996), citing unpublished data collected by George Schatz, reported that two unrelated species of *Philodendron* subg. *Pteromisium* were pollinated by the same species of beetles, and that both *Philodendron* species have floral odors featuring the same two principal components. This leads to the conclusion that there is a degree of specificity among pollinators for certain species based on their floral odors. In addition, in the list of pollinators known for *P.* subg. *Philodendron* (see Table 3), only three *Philodendron* species were observed to have more than a single species of beetle present at any one time. In each case, two species of beetles were present. As can be seen from *P. grandipes* (Table 3), the beetle species need not always be the same. Though more studies must be made on pollination biology of *Philodendron*, and even though the beetle pollination system is somewhat sloppy and imprecise, a combination of a moderately strong beetle-plant specificity, coupled with severe phenological constraints and narrow windows of pollination opportunities (perhaps as little as a few hours per year), works to reduce interspecific hybridization. Although hybrids can be readily produced under greenhouse conditions, evidence for hybridization is not usually apparent among wild populations.

#### BREEDING BEHAVIOR

In comparison to *Anthurium*, where breeding studies were easy to conduct (Croat, 1980, 1983a,

1986a, 1991), *Philodendron* pollination was difficult. Cross-pollination attempts were easy in *Anthurium* owing to their hermaphroditic flowers and because plants often had several inflorescences per plant in different stages of development. In addition, the plants reached anthesis during the day when greenhouse personnel and volunteers were available to make cross-pollinations. In *Philodendron*, the number of inflorescences available was always fewer than in *Anthurium*; *Philodendron* sometimes produced only a single inflorescence per season. In addition, *Philodendron* was very seasonal in its flowering behavior (unlike *Anthurium*, which sometimes flowered all year), making pollination all the more difficult. When flowering did occur it was often unexpected since it is difficult to tell when the spathe is ready to open. Moreover, the spathe generally opens for one day only. Opening usually took place late in the afternoon after greenhouse personnel left. Even if the opening inflorescence was found in time, it was generally impossible to find another plant with fresh pollen to use for purposes of experimental crosses. *Philodendron* pollen does not remain viable very long, though it can be kept viable for a time in glassine envelopes. Ron Weeks, a grower from Homestead, Florida, reports (pers. comm.), that he stores pollen in film canisters at refrigerator temperatures and that it remains viable for several weeks. He also reports that inflorescences cannot be pollinated after the first evening they open. After the spathe opens (generally late in the day) it is only during the evening and night of the first night that the pistillate flowers are believed to be receptive. Attempts to pollinate plants with their own pollen have always failed if one waits until the pollen emerges. It should be noted, however, that Grayum (1996) reported that, based on the use of peridoxase paper, which purportedly indicates the receptivity of stigmas (Young, 1986), the stigmas were receptive for up to 24 hours after anthesis. Though it seems unlikely that pistillate flowers are receptive after the first evening of anthesis, the pistillate flowers are receptive for an unknown period of time before the spathe opens, so that most successful pollinations usually involved cutting a hole in the spathe after obtaining very fresh pollen from a plant in the staminate phase of flowering. Though one can completely remove the spathe then protect the developing pistils with a plastic bag, it is better to simply cut a window in the spathe large enough to see most of the spathe. Then with a small brush one can spread pollen over as many of the pistils as possible, again covering the spathe for a time with a plastic bag to insure that the pollen does not

dry out and fail to germinate. An effective means of spreading the pollen to insure adequate and uniform coverage is to mix the pollen with water.

Failure to remove or at least loosen the bag used to cover the pollinated spadix later may result in mold developing in the spathe. Unpollinated inflorescences usually fall off within a week or two. Ron Weeks (pers. comm.) reported that for *P.* subg. *Meconostigma* in Florida the unpollinated inflorescences may persist for up to a month. Development time for fruits ranges from only a few weeks or more generally a few months and sometimes nearly a whole year. Ron Weeks (pers. comm.) reports that in *P.* subg. *Meconostigma* fruits ripen in South Florida in 2.5 to 3 months.

#### PHENOLOGY

Based on a field-oriented study of phenology and pollination behavior carried out at La Selva in Heredia Province, Costa Rica, Grayum (1996) reported species of *P.* subg. *Pteromischum* flowered for periods of 4–8 weeks. Obviously, since most members of *P.* subg. *Pteromischum* have only one or two inflorescences per axil, vs. sometimes 4 or more for *P.* subg. *Philodendron*, the flowering episodes of the latter might be longer than two months. The flowering events, however short or long, are not necessarily the same year after year. I suspect that, like understory vegetation (Croat, 1975), their flowering phenology may be affected by the onset of the rainy season.

Based primarily on a phenological survey of herbarium collections, flowering (and to a lesser extent fruiting) behavior has been studied here for Central American species of *P.* subg. *Philodendron*. These studies resulted in a phenological statement for each species. This statement follows the description of each species as a part of the discussion. Some general comments regarding phenology are important.

In studying herbarium material for phenological variation it is often difficult to determine the exact state of the inflorescence. However, with experience it is relatively easy to distinguish inflorescences that have never opened, i.e., pre-anthesis, from those which have already opened. Spathes that have never opened are very tightly closed, whereas those that have already undergone anthesis are not so tightly closed. By dissecting the spathe one can quickly determine if the pollen has emerged. If so, the spathe has already opened and reclosed over the spadix.

Spadices at anthesis when they are collected are usually easy to discern as well because they are

typically opened when pressed. It is difficult, however, to easily predict the age of an inflorescence beyond anthesis and before swelling due to the enlargement of the pistils. Since an inflorescence at anthesis is rare (open less than 24 hours), while every successfully pollinated inflorescence persists for one to many months, there will always be many more collections with "post-anthesis" inflorescences than those described as "in flower."

Central American members of *P.* subg. *Philodendron* fall into several phenological groups. All of these categories can have variations, and flowering is rarely consistent throughout any period. Some categorizations are tentative, as listed in Appendix 4, "Phenological Patterns of Central American *Philodendron* subg. *Philodendron*."

The flowering patterns of Central American members of *P.* subg. *Philodendron* are as follows:

#### A. FLOWERING IN DRY SEASON AND WET SEASON

This is the largest flowering category among Central American *P.* subg. *Philodendron*. A total of 47 species fall into this category, including *P. fragrantissimum*, *P. panamense*, *P. sagittifolium*, *P. grandipes*, *P. pterotum*, and *P. radiatum*. Although these species begin flowering during the dry season, the dry season rarely constitutes the period of greatest flowering activity (except perhaps in the case of *P. hederaceum*, a species which though apparently flowering all year and thus aseasonal appears to have more flowering collections made during the dry season than at any other time of the year). Low flowering activity is particularly characteristic of species inhabiting regions of *Tropical moist forest* or other areas where marked seasonal changes are apparent and affect the availability of beetle pollinators.

Those species occurring in cloud forest habitats (usually *Premontane rain forest* or *Lower montane wet forest*) also seem to have more flowering in the rainy season even though they are more apt to be in flower in the dry season than their *Tropical moist forest* counterparts.

Though perhaps it is merely a matter of poor sampling, species more common, widespread, or locally abundant tend to have flowering seasons that extend from the dry season to the wet season, whereas rare species seem much more likely to have flowering restricted to either the dry or the wet season.

#### B. FLOWERING ONLY IN WET SEASON

This is the second largest flowering category, comprising 30 species (32 taxa) believed to flower

exclusively in the wet season (roughly between May and December in Central America). They represent largely rare or narrowly distributed species for the most part and are often species that inhabit the wettest and frequently the coolest forest types such as *Tropical wet forest*, *Premontane wet forest*, *Premontane rain forest*, and *Lower montane wet forest*. Examples of species in this flowering type are: *P. albisuccus*, *P. antonioanum*, *P. chiriquense*, *P. coloradense*, *P. correae*, *P. cotobrusense*, *P. dominicense*, *P. dodsonii*, *P. ferrugineum*, *P. gigas*, *P. hammelii*, *P. jefense*, *P. madroneum*, *P. niqueanum*, *P. pirrense*, *P. purulhense*, *P. squamicaula*, and *P. ubigantupense*.

Not all species that flower exclusively in the wet season occur in very wet or cool forest. A few species flower only in the wet season because the dry season in the region where they occur is often too severe, perhaps so severe as to limit the beetle pollinators. These include several Mexican species, e.g., *P. basii*, *P. breedlovei*, *P. dressleri*, and *P. sousoe*.

#### C. FLOWERING ONLY IN DRY SEASON

One of the most unusual flowering categories, and a relatively small one with only 7 species, is a group believed to flower only during the dry season (January to April in Central America). This group is diverse and difficult to characterize. Some of the species, such as *P. bakeri*, *P. brewsterense*, *P. chiripoense*, *P. edenudatum*, *P. folsomii*, and *P. knappiae*, occur in wet to very wet areas, some in areas where weather conditions are so bad in general during the rainy season that it may be more efficient to compete for pollinators during the dry season (which would not be very dry in any event). *Philodendron dwyeri* is unusual in that it flowered at the beginning of the dry season in an area that in general is quite arid in this season. It is known from only a single individual and may have represented an unusual, out-of-phase flowering.

#### D. FLOWERING ALL YEAR

Species that flower aseasonally are usually common species such as *P. hederaceum* (though it apparently flowers more frequently in the dry season) and *P. jacquinii* (which is also often edaphically versatile or less subject to the pressures of the environment). Though not as widespread as the two aforementioned species, *P. advena*, another member of this group, is a tough, highly variable and ecologically adaptive species. *Philodendron purpureoviride* has much the same edaphic preferences as *P. hederaceum*, although it is much less widespread.

Both are highly adaptable, scandent species. *Philodendron hederaceum*, though flowering all year, is even more commonly in flower in the dry season than in the wet season. *Philodendron radiatum*, also a member of this group, is as widespread and even more common than *P. hederaceum*.

#### E. FLOWERING BIMODALLY

Based on the historical record of herbarium collections, only a few species are expected to flower twice per year. This pattern may be much more common and simply masked by the year-to-year variation in flowering behavior. Grayum (1996) reported bimodal flowering to be common with *P. subg. Pteromischum*. He indicated that the two modes were quite unequal, that one of them involved far fewer individuals and lasted for a shorter period of time. Among species of *P. subg. Philodendron*, *P. aromaticum* appears to flower in the mid-dry season and mid-wet season. It is possible this species is just too poorly known to determine its phenology. *Philodendron morii* may also flower bimodally, with flowering collections seen in March and November, and with immature fruiting collections in December, February, and June. *Philodendron wilburii* appears to flower at the beginning of the dry season and primarily later at the beginning of the rainy season.

Several species are poorly known phenologically because of sparsity of flowering collections: e.g., *P. dominicalense*, *P. niqueanum*, and *P. ubigantupense* (all seen in flower only once during the early rainy season). All of these are assumed to be species that flower entirely during the rainy season.

Fruiting phenology in *Philodendron* subg. *Philodendron* is too poorly known to report on here. Most species appear to develop fruits from between one and four months after the time of pollination, depending on the size of the infructescence, but too few mature fruiting collections were observed to determine the phenological period. Fruit development time is affected by the size of the infructescence. Species with small spadices, such as *P. heleniae*, produce mature berries faster than species with larger spadices.

#### CYTOLOGY

No karyological studies were made for this revision, but Petersen (1989) reviewed all chromosomal literature and made new studies. The chromosomes of *P. subg. Philodendron* are small, with counts of  $2n = 30, 32, 34,$  and  $36$  (rarely  $26$  and  $48$ ). Petersen (1989) speculated that the base number for the ge-

nus is 18. Very few of the 29 species (other names were synonyms or hybrids) for which chromosome counts have been reported are members of the genus from Central America. Those Central American species for which chromosome counts have been reported are: *P. radiatum*  $2n = 32$ , *P. microstictum* (as *P. pittieri*)  $2n = 34$ , *P. verrucosum*  $2n = 34$ , and *P. hederaceum* (as *P. scandens*)  $2n = 36$ . *Philodendron uendlandii* was counted at  $2n = 54$ , but Petersen believed this to be either an error in counting or not a *Philodendron*.

#### GEOGRAPHICAL DISTRIBUTION AND ENDEMISM

The *Philodendron* subg. *Philodendron* flora in Central America is diverse but heavily concentrated in the southeastern part near South America. Despite this, relatively few Central American species actually enter South America. Most species that do enter the South American continent range along the Pacific slope into northwestern Colombia and Ecuador, with relatively few occurring east of the Andes and even fewer entering the Amazon basin.

As was shown for *Anthurium* (Croat, 1983a, 1986a, 1986b), species diversity of *P. subg. Philodendron* shows a general diminution from Mexico to Middle America, followed by a marked increase approaching South America (Appendix 3, "Sectional Classification of *Philodendron*"). Distribution of *Philodendron* in Central America is as follows: Mexico has 21 taxa, Guatemala 15, Belize 9, El Salvador 5, Honduras 13, Nicaragua 18, Costa Rica 48, and Panama 82. Thus species diversity is greatest in Costa Rica and Panama, with 46% and 79% of the total Central American species, respectively. Collectively the number of species in these two countries comprises 90% of the species total.

Fifty-three species (58 taxa) of *P. subg. Philodendron* (nearly 59% of the total) are endemic to Costa Rica and Panama. Endemism is particularly high in Panama, where 38 taxa (34 species) of 81 (47%) are endemic. In Costa Rica 7 of 47 taxa (46 species) (15%) are endemic. Costa Rican endemics are: *P. aromaticum*, *P. auriculatum*, *P. chirripense*, *P. cotobrusense*, *P. dominicalense*, *P. microstictum*, and *P. wilburii* var. *wilburii*. Mexico has a higher rate of endemism, with 7 of 21 (20 species) (33%) endemic. Mexican endemics are: *P. basii*, *P. breedlovei*, *P. dressleri*, *P. radiatum* var. *pseudoradiatum*, *P. hederaceum* var. *oxycardium*, *P. sousae*, and *P. subincisum*. In Middle America little endemism occurs. With the exception of Belize, which has one endemic (*P. dwyeri*), no other country in Middle America has any endemic species.

The distribution of Central American *Philodendron* reflects the trend for endemism in the genus. Only 27 species (28 taxa) (a total of 26% of all Central American species of *P. subg. Philodendron*) range into South America, eight (7% of the total) only to Colombia (Appendix 1). These are: *P. heleniae*, *P. immixtum*, *P. ligulatum* var. *ligulatum*, *P. malesevichiae*, *P. mexicanum*, *P. pseudauriculatum*, *P. radiatum* var. *radiatum*, and *P. squamicaule*.

Sixteen Central American taxa occur in Ecuador (Appendix 1), all but two (*P. brevispathum* and *P. jacquinii*), ranging along the Pacific slope of the Andes. These are: *P. brunneicaule*, *P. dodsonii*, *P. grandipes*, *P. hebetatum*, *P. platypetiolatum*, *P. purpureoviride*, *P. hederaceum* var. *kirkbridei*, *P. hederaceum* var. *hederaceum*, *P. squamipetiolatum*, *P. scalarinerve*, *P. squamicaule*, *P. tenue*, *P. tripartitum*, and *P. verrucosum*. Of these 14 taxa, 1, *P. hederaceum* var. *kirkbridei*, skips Colombia or has not yet been collected there. *Philodendron dodsonii* is particularly unusual in being absent from Panama.

Eight Central American species, *P. brevispathum*, *P. fragrantissimum*, *P. hederaceum*, *P. jacquinii*, *P. jodavisanum*, *P. sagittifolium*, *P. strictum*, and *P. tenue*, range to Venezuela. The ranges of *P. brevispathum*, *P. fragrantissimum*, and *P. hederaceum* also extend into the Amazon drainage, while the others occur either on the northern slope of the coastal cordillera or otherwise in the drainage of the Orinoco River Basin. Curiously only four species, *P. brevispathum*, *P. fragrantissimum*, *P. hederaceum*, and *P. verrucosum*, occur in the Amazon drainage. Three additional species, *P. jacquinii*, *P. strictum*, and *P. tenue*, occur east of the Andes, but only along the Cordillera de Mérida, the Cordillera de la Costa or the northern part of the Guiana Highlands and within the drainage of the Río Orinoco.

The only truly widespread Central American species of *P. subg. Philodendron* is *P. hederaceum*, which occurs virtually throughout the Neotropics, and is one of only two species (the other being *P. verrucosum*) that occurs on both slopes of the Andes. *Philodendron fragrantissimum* is probably the next most widespread species, ranging from Belize to the West Indies and into South America to the Guianas, northern Brazil, and to southern Peru.

Further collecting in Colombia, especially along the western slope of the Andes, will probably change these statistics but the figures most likely reflect the realities of life zone ecology and geologic history of the area rather than under-collecting. Since relatively few species of Araceae are known to occur at lower elevations on both the eastern and western side of the Andes, it can probably be pre-

sumed that the evolution of the respective Amazonian and Pacific coastal floras occurred independently after the Andes began to arise toward the end of the Cretaceous (Raven & Axelrod, 1974). The relatively few truly wide-ranging species, i.e., those ranging from Mexico to Brazil, appear to attest to this isolation. In Central America only one species, *P. hederaceum*, really falls into such a category, and it is also common in the West Indies indicating that it may have an ancient origin (or be easily dispersed). The high rates of endemism in Costa Rica, Panama, and Mexico perhaps reflect the isolation of these areas during periods when the oceans were at much higher levels than they are today and when the area that is now central Panama and Costa Rica was disconnected from South America. Much of the present area of Central America was submerged during early times. At the close of the Tertiary, 800,000 years ago, sea level was about 100 m higher than today (Holmes, 1969). The land mass of what is now Central America began to emerge as a series of islands during the Oligocene with further uplifting during the Middle Miocene. It was not until the Upper Miocene and Pliocene that the final portions of the isthmus of Panama emerged above sea level (Torre, 1965), and the final connection of Central and South America was made about 5.7 million years ago. In order to place these geological events in relation to the modern aroid flora, it should be noted that even during this era precursors to the existing flora probably already existed, since the angiosperm floras of the Oligocene were believed to have consisted almost entirely of existing genera, and the floras of the Oligocene and Pliocene probably already had existing species (Takhtajan, 1969).

Just as important as geology, from the standpoint of the isolation of the Central American aroid flora, are ecological factors that would cause Central American species to be isolated from those of South America. Much of eastern Panama consists of broad expanses of *Tropical moist forest* with other, generally smaller areas of *Premontane wet* and *Tropical wet forest*. In contrast to Panama, much of the area of northwestern Colombia in the Department of Chocó consists of much wetter pluvial forest with annual precipitation often exceeding 11,700 mm in some parts of the region (Gentry, 1982). This broad band of pluvial forest with its own suite of unique endemic species no doubt acts as a barrier for species from regions with lesser rainfall amounts. It probably also accounts for the Panamanian and Costa Rican species that skip the wettest areas of northwest South America but recur in the relatively drier areas of mesic western Ecuador.

Just as the Central American *P.* subg. *Philodendron* flora is rather isolated from that of South America, there is a certain amount of isolation within different parts of Central America. In comparison to Mexico and Costa Rica/Panama, Middle America (Appendix 1) has low species diversity, with Guatemala having only 15 species, Honduras 13 species, and Nicaragua 18. Most of the species in Honduras, excepting *P. anisotomum*, *P. mexicanum*, and *P. warszewiczii*, are shared with Nicaragua (see Appendix 1). Nicaragua has eight additional taxa not shared with Honduras: *P. brevispathum*, *P. grandipes*, *P. immixtum*, *P. ligulatum* var. *ligulatum*, *P. platypetiolatum*, *P. pterotum*, *P. tenue*, and *P. wendlandii*. All of the latter are shared with Costa Rica and Panama. Guatemala shares only about half of its species with Honduras and Nicaragua, namely *P. fragrantissimum*, *P. jacquinii*, *P. jodavisanum*, *P. radiatum*, *P. sagittifolium*, *P. smithii*, and *P. tripartitum*. Its other species are shared only with Mexico (or rarely with Costa Rica and Panama, e.g., *P. mexicanum*). These are: *P. advena*, *P. anisotomum*, *P. glanduliferum*, *P. mexicanum*, *P. purulhense*, *P. verapazense*, and *P. warszewiczii*.

The low species diversity and the very low endemism in Middle America are perhaps explained by the fact that Central America is rather more remote from existing large land masses to the north and the south, leaving it isolated from the independent evolution that must have been taking place in both of these larger areas (see below for a discussion of the possible origins of the respective species in Central American *Philodendron*). There is strong evidence, at least based on the distribution of modern aroid species, that the northwestern part of Middle America may have been isolated from Costa Rica by the San Juan Depression. Many of the species that occur in Costa Rica or Panama enter into Nicaragua in only a small area in the southeastern part of the country. Although the contemporary flora of Guatemala does not reflect isolation from Mexico to the same degree, it is possible that the more elevated portions of Guatemala, Nicaragua, and Honduras were isolated from major portions of Mexico at the Isthmus of Tehuantepec.

Certainly the Mexican aroid flora appears to be quite isolated, even when compared to the western parts of Middle America (here defined as Guatemala to Nicaragua). Mexico, in addition to having one-third of its species endemic, has relatively few species of *Philodendron* that range throughout Central America. Aside from the aforementioned *P. hederaceum*, only *P. jacquinii*, *P. jodavisanum*, *P. radiatum*, *P. sagittifolium*, and *P. tripartitum* range to northern South America. Two additional species,

*P. anisotomum* and *P. mexicanum*, range as far as Panama. Several taxa, *P. advena*, *P. glanduliferum* var. *glanduliferum*, and *P. verapazense*, range only to Guatemala. *Philodendron purulhense* and *P. warszewiczii* range to Honduras, and *P. smithii* ranges to Nicaragua.

Taken together, Honduras and Nicaragua have 21 species of *Philodendron*. Of these, 8 species, *P. advena*, *P. anisotomum*, *P. mexicanum*, *P. radiatum*, *P. sagittifolium*, *P. smithii*, *P. tripartitum*, and *P. warszewiczii*, appear to be of Mexican origin, or in the case of the more widespread and variable species, namely *P. radiatum*, *P. sagittifolium*, and *P. tripartitum*, they may have originated in Panama or Costa Rica and ranged to both Mexico and South America. Certainly, in terms of morphological variation, all of these species are much more variable in Panama and Costa Rica than they are further north. *Philodendron jacquinii* has a circum-Caribbean distribution, indicating that it may be of West Indian origin. It is difficult to determine the origin of *P. hederaceum* given its extensive distribution. Three species, *P. brevispathum*, *P. fragrantissimum*, and *P. tenue*, probably originated in South America considering their widespread distribution there. *Philodendron platypetiolatum*, ranging from Ecuador and barely entering Nicaragua, may be another South American derivative. The remainder, *P. angustilobum*, *P. grandipes*, *P. immixtum*, *P. jodavisanum* (ranging barely to Chiapas and rare there), *P. ligulatum* var. *ligulatum*, *P. pterotum*, *P. rothschuhianum*, and *P. wendlandii*, are probably of Panamanian or perhaps Costa Rican origin.

The Costa Rican and Panamanian species not already discussed above appear not to have strong affinities with South American species, and clearly did not originate in areas of Middle America. Much of the flora of adjacent Nicaragua is closely related to that of Costa Rica. Except for those rather widespread species mentioned above, i.e., *P. brevispathum*, *P. fragrantissimum*, and *P. tenue*, as well as *P. platypetiolatum* (already discussed), there are relatively few species likely to be of South American origin. *Philodendron verrucosum* is almost certainly a South American species, since it is relatively widespread there, occurring on both sides of the Andes. In addition, it seems to have more related species in parts of South America especially in the Andes of western Colombia.

*Philodendron dodsonii*, which occurs in Ecuador but not Colombia, is just as likely to have originated in South America as in Central America. The same is true of *P. strictum*, which is known from eastern Venezuela and has relatives in the Andes of central Colombia, as well as *P. hebetatum*, which

is rather widespread in western South America as far south as Ecuador. Moreover, the latter apparently does not even reach Costa Rica, making the case for a South American origin even more likely.

*Philodendron grandipes*, known from western South America as far south as Ecuador, might conceivably be of South American origin despite being very widespread and common as far north as Nicaragua. More probably of South American origin is *P. heleniae*, which is common in Colombia and ranges only to western Panama. Moreover, it seems to have a close relative on the eastern slopes of the Andes.

Three species with scaly parts, *P. malesevichiae*, *P. squamicaulis*, and *P. squamipetiolatum*, are moderately rare in Panama, and though all are still poorly known in Colombia, they are more likely to have originated in northwestern Colombia where there are several other relatives with scaly parts.

Some taxa, such as *P. ligulatum* var. *ligulatum*, *P. pseudauriculatum*, and *P. scalarinerve*, either barely enter Colombia or are rare there and are more likely therefore to have originated in Central America. All the remaining species of *P.* subg. *Philodendron* occur in Panama or Costa Rica, with 11 species shared between the two countries.

#### TAXONOMIC TREATMENT

**Philodendron** ("Philodendrum") Schott, Wiener Z. Kunst. 1829: 780. 1829, nom. et orth. cons.

TYPE: *P. grandifolium* (Jacq.) Schott (*Arum grandifolium* Jacq.)

*Telipodus* Raf., Fl. tellur. 3: 66. 1836[1837]. TYPE: *T. grandifolium* (Jacq.) Raf. (*Arum grandifolium* Jacq.)

*Thaumatophyllum* Schott, Bonplandia 7: 31. 1859. TYPE: *T. spruceanum* Schott

*Elopium* H. W. Schott, Oesterr. Bot. Z. 15: 34. 1865. TYPE: *E. surinamense* (Miq.) Schott (*Anthurium surinamense* Miq.)

*Baurisia* T. Post & Kuntze, Lex. gen. phan. 62. 1903. TYPE: *Caladium bauerii* Rchb.

Appressed-climbing or scandent hemiepiphytes, sometimes epiphytes or terrestrial herbs (sometimes stout and arborescent); growth sympodial, anisophyllous, or (more commonly) diphyllous; trichosclereids absent, biforines and secretion files present. Leaves with spiral phyllotaxy; petioles rarely geniculate, sheath obsolete or extensive; blades with parallel-pinnate venation, hypostomatic with stomates paracytic, simple to trifoliate, pinnatifid, bipinnatifid, or (rarely) pedately compound. Inflorescence terminal (appearing axillary), solitary to multiple in leaf axils; spathe enveloping the spadix and constricted or not, usually persistent; spadix monoecious, with sterile male flowers below or both below and (rarely) above the fertile male region;

flowers naked, unisexual. Male flowers with 2-6 free stamens, connective enlarged; anther dehiscence by apical slits or pores; pollen inaperturate, boat-shaped, subsopolar, bilaterally symmetrical, shed in monads, medium-sized (mean 40  $\mu\text{m}$ ; range 28-54  $\mu\text{m}$ ), the exine coarsely verrucate to subfoveolate or (most frequently) psilate; pollen at anthesis binucleate and starch-bearing. Female flowers virtually always lacking staminodes, ovary (2-3)-6(47)-locular, each locule with axile or (rarely) basal placentation and (1-)-4-51 or more orthotropous or hemiantropous, endospermous ovules. Fruit a berry, usually white or orange. Chromosome numbers  $2n = 30, 32, 34, 36$ . [Description adapted from Grayum (1996).]

#### KEY TO PHILODENDRON SUBGENERA

- 1a. Stem of mature flowering plants with a succession of many leaves terminated by solitary or paired (rarely 3) inflorescences; petioles with long sheaths narrowly encircling the stem at base, winged to at least midway on adult plants; up to 5 leaves produced on the stem between each successive inflorescence ..... subg. *Pteromischum*
- 1b. Stem of mature flowering plants with a succession of short sympodial segments each bearing a cataphyll and a single leaf with the inflorescence(s) 1 to 10 and appearing to be borne in the leaf axils; petioles of leaves of adult plants with a short, usually inconspicuous petiole sheath borne on the side of the stem, not encircling it at the base (sometimes fully winged on juvenile plants); inflorescences produced with each new leaf (though frequently aborted).
  - 2a. Stems often arborescent, with conspicuous leaf scars and frequently interpetiolar scales persisting around at least the upper margins of the petiolar scars; male flowers conspicuously elongate, up to 10 times longer than wide; staminodial zone between staminate and pistillate zones of the spadix subequal or longer than fertile zone ..... subg. *Mecostigma*
  - 2b. Stems rarely arborescent, often scandent, stout or slender and with interpetiolar scales lacking; male flowers only 2-3 times longer than wide; staminodial zone between staminate and pistillate zones of spadix much shorter than the fertile staminate zone ..... subg. *Philodendron*

**Philodendron** subg. **Philodendron** Schott.  
*Philodendron* subg. *Euphilodendron* Engl., Bot. Jahrb. Syst. 26: 509. 1899. *Baurisia* subg. *Eubaurisia* Kuntze in T. Post & Kuntze, Lex. gen. phan. 62.

Appressed hemiepiphytic climbers or vines with aerial roots, less frequently terrestrial with creeping rhizomatous or deeply rooted stems, rarely short-



stemmed true epiphytes, rarely somewhat arborescent in Central America; sometimes with flagelliform shoots; sap usually tanniferous and drying dark, rarely with latex and drying white; stems of monophyllous sympodia with elongated hypopodial internodes, densely rooted at nodes; internodes often much longer than broad or about as long as broad, sometimes broader than long at anthesis, sometimes flattened on one side, often coarsely pale-streaked just below the node, usually green and semiglossy, but often turning gray-green to brownish or reddish brown in age; juvenile plants terrestrial or epiphytic and scandent, the petioles conspicuously sheathed and subtended by inconspicuous intravaginal squamulae; cataphylls of mature stems unribbed or variously ribbed, caducous, marcescent and deciduous or persistent and membranaceous to moderately coriaceous, remaining intact or more commonly decomposing to net-like, persistent fibrous reticulum. Leaves usually long-petiolate; petioles usually with ligulate sheath in juvenile plants, on adult plants usually sheathed only at base, variously shaped in cross section, firm or spongy, usually smooth, frequently densely pale-short-linear or pale-striate throughout, sometimes warty or covered with scale-like processes, rarely glicate apically; blades simple and entire, ovate, cordate, hastate, sagittate, oblong to elliptic or variously divided, trifid, trifoliolate, or pinnatifid; midrib raised or sunken above, raised below; primary lateral veins pinnate, usually conspicuous, spreading to the margins and running into an antemarginal collective vein; lowermost primary lateral veins (basal veins) often coalesced on cordate blades, the posterior rib (coalesced basal veins) naked along the sinus or not; interprimary veins sometimes present; secondary lateral and higher order veins transversely reticulate between the secondary veins, sometimes all veins slender with no distinct primary lateral veins; minor veins conspicuous or obscure, usually fine and closely parallel; cross-veins (minute veins extending transversely between the minor veins) sometimes visible; secretory ducts sometimes appearing like veins, linear, short to long, obscure to very distinct on lower surface. Inflorescences 1-several per axil, usually much shorter than the petioles; peduncles shorter or longer than the spathe; spathe erect, usually coriaceous, entirely persistent, often with large superficial resin canals on inner surface which exude resin, opening widely at anthesis (usually for about one day), then reclosing and persisting in fruit, deciduous only on ripening of fruit, frequently colorful, often bicolorous on outside, typically somewhat constricted between tube and blade, convolute at base; tube cylindrical to inflated, uniformly greenish to red or violet-purple within; blade usually opening widely, becoming  $\pm$  boat-

shaped at anthesis, usually white within, sometimes tinged reddish; spadix sessile to stipitate, divided into pistillate and staminate portions, each with unisexual flowers; pistillate zone usually greenish, obliquely fused at its base to the spathe, free above, usually much shorter than the staminate portion and separated from it by a sterile zone of staminodial flowers; intermediate sterile zone cylindrical to ellipsoid, much shorter than staminate zone in Central America, usually thicker than staminate zone; staminate zone terete to clavate, white, usually somewhat constricted above the sterile staminate zone; flowers unisexual, naked, closely aggregated in several spirals; staminate flowers 2-6 (usually 4-5)-androus; stamens free, adjacent, lacking stomial groove, prismatic to obpyramidal; anthers tetrasporangiate (with microsporangia embedded in the abaxial surface of the anther), columnar in shape, elliptic, ovate to rhombic in cross section, sessile to subsessile; connective thick, apically truncate, usually irregularly 4-5-sided, overtopping thecae; thecae oblong or elliptic, emarginate at the base, dehiscing apically by short, ragged lateral pores; endothelial thickenings lacking; pollen extruded in strands or mixed with resin secretion or exuded in amorphous masses, inaperturate, ellipsoid or oblong or occasionally elongate, medium sized (mean 40  $\mu\text{m}$ , range 28-54  $\mu\text{m}$ ), mostly perfectly psilate, sometimes minutely verruculate, scabrate or foveolate to clearly punctate, subfossulate, subfoveolate or subverrucate; sterile staminate flowers naked, usually prismatic, truncate and usually more irregular than fertile flowers and lacking thecae; pistillate flowers with ovary syncarpous, ovoid, subcylindrical, cylindrical or obovoid, 3-9(14)-locular in Central America (2-locular in *P. sect. Philopsammis* in South America); carpels presumably equal to number of locules; placentation axile, sub-basal or basal; ovules 1 to numerous (to ca. 30) per locule, usually hemicarpotropous, rarely hemianatropous, ascending on moderately long or sometimes short funicles; stylar region as broad as or sometimes slightly narrower than ovary; style short, unlobed, with or without boss (see definition under Style Type D), funnel, or annulus; central style dome usually lacking in Central America; stigma sessile, hemispherical to lobulate. Berries subcylindrical to obovoid, exposed by the re-opening of the spathe, white, whitish translucent to red or orange; seeds few to many per berry, oblong to ellipsoid or ovoid-oblong, testa rather thick, striate-costate, rarely sarcotestate; embryo axile, straight, elongate, endosperm copious; chromosomes:  $2n = 30, 32, 34, 36, (26, 48)$ . Species ca. 700, Central Mexico to Argentina; West Indies (occurring in all countries of Central and South America except Chile and Uruguay).

KEY 1: BLADES DEEPLY LOBED OR DIVIDED .....	382
KEY 2: BLADES SIMPLE, NON-CORDATE, EITHER OBLONG OR OVATE; SOMETIMES SUBCORDATE TO CORDULATE AT BASE, THE POSTERIOR LOBES UP TO $\frac{1}{4}$ AS LONG AS ANTERIOR LOBES .....	383
KEY 3: CORDATE BLADES WITH PERSISTENT CATAPHYLLS .....	385
KEY 4: VINES WITH CORDATE BLADES AND DECIDUOUS CATAPHYLLS .....	388

KEY TO *PHILODENDRON* SUBGENUS *PHILODENDRON* OF CENTRAL AMERICA

A. Blades deeply divided, either pinnately lobed, or 3-lobed to trifoliate .....	KEY 1
A'. Blades not deeply divided, at most with the margins merely sinuate, the lateral margins of the anterior lobe sometimes deeply concave, but not to such an extent that the blade looks decidedly 3-lobed (blades at first entire, but naturally splitting into irregular segments in <i>P. findens</i> ).	
B. Blades acute, obtuse or truncate at base or sometimes weakly cordulate or subcordate but with the posterior lobes only $\frac{1}{4}$ as long as the anterior lobe or less .....	KEY 2
B'. Blades cordate, sagittate, or hastate at base, the posterior lobes usually more than $\frac{1}{4}$ as long as the anterior lobes.	
C. Cataphylls persistent on stem, either intact or as fibers .....	KEY 3
C'. Cataphylls ultimately deciduous, though sometimes persisting .....	KEY 4

## KEY 1: BLADES DEEPLY LOBED OR DIVIDED

- 1a. Blades deeply pinnately lobed, the divisions extending at least  $\frac{1}{4}$  of the way to the midrib.
- 2a. Leaves bipinnately lobed, the blade divisions near the middle of the blade pinnately lobed, usually divided  
more than  $\frac{1}{2}$ -way to the midrib.
- 3a. Blades lobed only  $\frac{1}{16}$ - $\frac{1}{12}$  the way to the midrib; Mexico (Sinaloa & Nayarit); 0-370 m .....  
*P. dressleri* G. S. Bunting
- 3b. Blade deeply lobed, the divisions extending more than  $\frac{1}{16}$  of the way to the midrib.
- 4a. Blades deciduous during the dry season; leaf segments thin, drying papyraceous, pale yellow-  
green to yellow-brown, with the minor veins distinctly visible and darker than surface; lateral  
segments with usually 3 or more lobes per side, extending to below the middle of the pinnae;  
Mexico (Jalisco & Chiapas) to El Salvador, Guatemala, Honduras, and Nicaragua, 300-1500 m .....  
*P. searszevicii* K. Koch & Bouché
- 4b. Blades evergreen; leaf blade segments subcoriaceous, drying moderately thick, dark brown or  
sometimes reddish brown, the minor veins not distinctly visible on drying, not markedly darker  
than surface; lateral segments with usually 1-2 lobes per side, usually restricted to near the apex  
of the pinnae; Mexico to Colombia, 0-700, rarely to 1860 m ..... *P. radiatum* var. *radiatum*
- 2b. Leaves merely pinnately lobed, the blade divisions near the middle of the blade not pinnately lobed, or  
if so, divided less than  $\frac{1}{2}$ -way to the midrib ..... *P. radiatum* var. *pseudoradiatum* Croat
- 1b. Blades deeply 3-lobed or trifoliate.
- 5a. Blade with the medial lobe less than 1.7 times longer than the lateral lobes; lateral lobes directed  $\pm$   
toward the apex.
- 6a. Lateral lobes of the blade broadly confluent with the medial lobe by at least 2 cm from the base of  
the blade (as measured from the depths of the incised area and the apex of the petiole); inflorescences  
usually 3-9 per axil.
- 7a. Medial lobe of the blade with 5-12 pairs of primary lateral veins; spathe tube uniformly greenish  
to whitish within; Honduras to Panama, usually below 1000 m .....  
*P. ruthachianum* (Engl.) Croat & Grayum
- 7b. Medial lobe of the blade with 18-19 pairs of primary lateral veins; spathe tube red-violet or  
purplish within; Costa Rica (Puntarenas, San José), 1000-1250 m .....  
*P. catobrusense* Croat & Grayum
- 6b. Lateral lobes of the blade not confluent with or only weakly confluent with the medial lobe to no  
more than 1 cm from the base of the blade; inflorescences usually 1, rarely to 3 per axil.
- 8a. Medial lobe of blades with more than 18 pairs of primary lateral veins, elliptic; central Panama,  
350-450 m .....  
*P. madroense* Croat
- 8b. Medial lobe of blade with fewer than 12 pairs of primary lateral veins, usually oblanceolate to  
oblong-lanceolate; widespread species, Mexico to Ecuador.
- 9a. Medial lobe mostly 3-3.5 times longer than broad (rarely as little as 1.7 times longer than  
broad); primary lateral veins of medial lobe 4-12 per side, prominently sunken; lateral lobes  
directed  $\pm$  toward the apex; minor veins arising from both the midrib and the primary lateral  
veins; ripe fruits whitish; Mexico to Ecuador, 0-1300(1500) m ..... *P. tripartitum* (Jacq.) Schott
- 9b. Medial lobe mostly 1.5-2 times longer than broad; primary lateral veins of medial lobe  
mostly 2-4, rarely 5 per side, weakly sunken; lateral lobes directed  $\pm$  outward; minor veins  
arising only from the midrib; ripe fruits orange; Mexico to Costa Rica, 30-1800 m .....  
*P. anisotomum* Schott

- 5b. Blades with the medial lobe more than 1.7 times longer than the lateral lobes; lateral lobes directed laterally (spreading) outward or even downward toward the base of the petiole (reflexed).
- 10a. Blades sagittate, the lateral lobes directed downward somewhat toward the base of blade (in direction of petiole); Mexico to Panama and South America, 0-1900 m ..... *P. mexicanum* Engl.
- 10b. Blades hastate, the lateral lobes directed outward at ca. 90° angle.
  - 11a. Lateral lobes confluent with the medial lobe by usually for more than 4 cm (rarely to 3.3 cm); basal veins 8-10 per side; Honduras to Panama, 0-680 m ..... *P. angustilobum* Croat & Grayum
  - 11b. Lateral lobes confluent with the medial lobe by usually less than 3 cm (rarely to 8 cm); basal veins 4-7 per side; Mexico to Costa Rica, 30-1800 m ..... *P. anisotomum* Schott

KEY 2: BLADES ENTIRE OR SHALLOWLY DIVIDED, NON-CORDATE, EITHER OBLONG OR OVATE; SOMETIMES SUBCORDATE OR CORDULATE AT BASE, THE POSTERIOR LOBES UP TO 3/4 AS LONG AS ANTERIOR LOBES

1a. Plants vines or at least with internodes much longer than broad.

VINES OR SCANDENT: INTERNODES MUCH LONGER THAN BROAD; BLADES OVATE

- 2a. Blades ± ovate, less than 2 times longer than wide.
- 3a. Blades with primary lateral veins either 2 or fewer or with the primary lateral veins inconspicuous, scarcely more distinct than the minor veins.
  - 4a. Blades more than 11 cm wide; peduncles more than 10 cm long; Costa Rica, 50-450 m ..... *P. microstictum* Standl. & L. O. Williams
  - 4b. Blades less than 11 cm wide; peduncles less than 10 cm long.
    - 5a. Internodes less than 10 cm long; blades less than 1.4 times longer than wide, drying greenish, lacking primary lateral veins; Panama, Cerro Brewster, 850 m ..... *P. brewsterense* Croat
    - 5b. Internodes more than 10 cm long; blades ca. 2 times longer than wide, drying brown, with up to 2 obscure primary lateral veins; Costa Rica, Río Chirripó del Pacífico, San José Prov., 1000 m ..... *P. chirripoense* Croat & Grayum
- 3b. Blades with the primary lateral veins more than 2 and much more prominent than the minor veins.
  - 6a. Blades ovate to broadly ovate, mostly less than 1.7 times longer than wide.
    - 7a. Stems, petioles, and inflorescences densely scaly (petiolar scales slender and spreading); major veins on lower surface densely puberulent; Panama to Ecuador, 0-1300 m ..... *P. squamipetiolatum* Croat
    - 7b. Stems, petioles, and inflorescences glabrous; major veins on lower surface glabrous; Panama (Coclé, Veraguas), 770-1200 m ..... *P. ligulatum* var. *ovatum* Croat
  - 6b. Blades narrowly ovate to oblong-elliptic, usually more than 2 times longer than wide.
    - 8a. Blades less than 12 cm wide; posterior lobes about as long as broad; Panama to Colombia, 0-140 m ..... *P. immixtum* Croat
    - 8b. Blades usually more than 12 cm wide; posterior lobes much broader than long; Costa Rica to Panama, (210)670-1800 m ..... *P. lentii* Croat & Grayum
- 2b. Blades ± oblong to oblong-elliptic (rarely narrowly ovate) or oblanceolate, usually more than 2.5 times longer than wide.

VINES WITH ± OBLONG BLADES

- 9a. Blades lacking distinct primary veins at base of blade; primary lateral veins obscure or lacking.
- 10a. Blades drying gray-green; primary lateral veins not at all apparent; Panama (San Blas), near sea level ..... *P. ubigantupense* Croat
- 10b. Blades drying reddish brown or blackened; primary lateral veins 3-6, obscure but still visible.
  - 11a. Blades drying blackened, usually more than 25 cm long; spathe more than 12 cm long; Panama (Bocas del Toro and Chiriquí), 780-1400 m ..... *P. coreae* Croat
  - 11b. Blades drying reddish brown, usually less than 25 cm long; spathe less than 12 cm long; Costa Rica (100-900 m) to Panama, 900-1420 m ..... *P. bakeri* Croat & Grayum
- 9b. Blades with one or more distinct primary veins at base of blade; primary lateral veins distinct.
  - 12a. Petioles encircled distally with a dark purplish (or dark green) ring separating the petiole and the blade.
  - 13a. Blades with posterior lobes usually narrowly rounded and somewhat spreading or broader than long, rarely about as long as broad; inflorescences 1-3 per axil; Panama, 100-970 m ..... *P. annulatum* Croat
  - 13b. Blades with posterior lobes about as broad as long and held closely near the petiole; inflorescence usually solitary (sometimes 2 in *P. immixtum*).
    - 14a. Blades typically drying greenish to greenish brown, rarely more than 30 cm long; Panama to Colombia, 0-140 m ..... *P. immixtum* Croat
    - 14b. Blades typically drying blackened, rarely less than 30 cm long; Nicaragua to Colombia, 0-1200 m ..... *P. ligulatum* Schott var. *ligulatum*
  - 12b. Petioles lacking a dark purple distal ring.

- 15a. Blades less than 10 cm wide; stem minutely pale granular-puberulent at high magnification; pistil with a funnel-shaped stigma (type E); ovules 1 per locule; Panama (Darién), 75 m ..... *P. granulare* Croat
- 15b. Blades usually more than 15 cm wide (rarely to as little as 10 cm wide in *P. heleniae*); stems smooth to irregularly ridged or folded at high magnification (sometimes minutely warty but not pale granular-puberulent); pistil with a flat style apex (type B); ovules 3 or more per locule.
- 16a. Blades drying coriaceous, lacking secretory ducts; spathes greenish white to white outside, more than 10 cm long; Costa Rica to Panama, (210)670-1800 m ..... *P. lentii* Croat & Grayum
- 16b. Blades drying subcoriaceous, with conspicuous secretory ducts; spathes red outside, mostly less than 10 cm long; Panama and Ecuador, mostly 100-1040 m ..... *P. heleniae* Croat

1b. Plants appressed-climbing or sometimes terrestrial, with the internodes typically broader than long or only slightly longer than broad.

#### NON-VINING PLANTS; BLADES OBLONG

##### 17a. Cataphylls deciduous.

- 18a. Blades usually prominently lobed at base, the lobes typically longer than broad or at least usually spreading away from the petiole.
- 19a. Blades thin-drying; sinus arcuate with blade tissue somewhat decurrent on petiole; petiole drying 2-4 mm diam.; Panama, E. of Canal Area, 450-850 m ..... *P. morii* Croat
- 19b. Blades thick-drying; sinus oblong to parabolic or arcuate but with blade tissue not at all decurrent onto petiole; Panama, 100-970 m ..... *P. anisulatum* Croat
- 18b. Blades not lobed at base or merely cordulate with posterior lobes round, as broad as long, held closely against the petioles.
- 20a. Largest leaf blades more than 50 cm long.
- 21a. Blades usually more than 30 cm wide; petioles usually about 85% the length of the blades or even longer than the blades; Costa Rica and Panama, 0-200 m ..... *P. davidsonii* Croat
- 21b. Blades less than 30 cm wide; petioles typically much shorter than the blade.
- 22a. Blades drying reddish brown, 4.4-4.9 times longer than wide; petioles lacking a purple distal ring; Panama, 325-650 m ..... *P. dolichophyllum* Croat
- 22b. Blades drying dark gray-green to blackened, usually less than 4.3 times longer than wide; petiole with a purple or green distal ring.
- 23a. Petioles subterete adaxially, not at all winged marginally; Panama and Colombia, 20-1400 m ..... *P. pseudauriculatum* Croat
- 23b. Petioles sharply D-shaped with slender wings on the adaxial margins; Panama, Darién, 50-200 m ..... *P. ligulatum* var. *heracleioides* Croat
- 20b. Largest leaf blades usually less than 50 cm long.
- 24a. Blades usually drying green to yellowish green, thin.
- 25a. Blades oblong-lanceolate, less than 7 cm wide, obtuse to almost rounded at base and then decurrent along petiole; Panama (Coclé), 700-800 m ..... *P. jolsomii* Croat
- 25b. Blades ovate-triangular, 11-16 cm wide, subcordate at base; eastern Panama, 450-850 m ..... *P. morii* Croat
- 24b. Blades drying yellow-brown or brownish, gray-brown to reddish brown, moderately coriaceous.
- 26a. Inflorescences 2-10 per axil; peduncle less than 6 cm long; spathe tube reddish maroon outside and within; leaf blades with conspicuous secretory ducts visible on the lower dried surface; Panama to Colombia and Ecuador, 20-1040(1450) m ..... *P. heleniae* Croat
- 26b. Inflorescence solitary; peduncle ca. 9 cm long; spathe tube green outside, white within; leaf blades lacking any sign of secretory ducts; Panama (Colón), ca. 200 m ..... *P. utleyanum* Croat

##### 17b. Cataphylls persistent.

- 27a. Plants usually terrestrial and rosulate; petioles deeper than wide; blades usually elliptic to oblong-elliptic, acute to narrowly rounded at base; Panama to Colombia, 100-1000 m ..... *P. roseospathum* Croat
- 27b. Plants epiphytic to hemiepiphytic, not rosulate; petioles as deep as wide; blades linear to oblanceolate or oblong-elliptic, sometimes weakly cordate to cordulate at base.
- 28a. Blades linear to oblanceolate, attenuate at base; primary lateral veins arising at 25-45° angle from midrib; cross-veins not visible on drying; sap chalky white; Costa Rica to Panama, 0-900 m ..... *P. cretatum* Croat & Grayum
- 28b. Blades oblong-elliptic to narrowly ovate, obtuse to subcordate at base; primary lateral veins arising at 60-75° angle; cross-veins very conspicuous on drying; sap not colored, or if so not white; Costa Rica to Colombia, 0-1325 m ..... *P. scalarinerve* Croat & Grayum
- 29a. Blades usually broadest above the middle, acute, obtuse, narrowly rounded to merely minutely cordulate at base.
- 30a. Petioles sharply flattened adaxially (with acute lateral margins); blades acute to narrowly rounded (never cordulate) at base, mostly more than 2.5 times longer

- than petioles; internodes broader than long; Costa Rica, Nicaragua, and Panama, Atlantic slope; 10–1250 m ..... *P. wendlandii* Schott
- 30b. Petioles subterete; blades narrowly cordulate at base, less than 1.8 times longer than petioles; internodes often longer than broad; southwestern Costa Rica, Pacific lowlands, 0–1200 m ..... *P. auriculatum* Standl. & L. O. Williams
- 29b. Blades broadest below the middle, cordate to subcordate at base (acute to truncate in *P. davidsonii* subsp. *bocatoranum*).
- 31a. Blades ovate to broadly ovate, less than 60 cm long, drying dark brown; primary lateral veins 5–9 per side; petioles terete; Panama (Darién), ca. 1500 m ..... *P. niquense* Croat
- 31b. Blades ovate-oblong, usually more than 60 cm long, drying light brown; primary lateral veins 18–21 per side; petioles thicker than broad (i.e., with the thickest dimension perpendicular to the plane of the blade) and broadly sulcate adaxially; Costa Rica to Panama, 0–200 m ..... *P. davidsonii* Croat

KEY 3: CORDATE BLADES WITH PERSISTENT CATAPHYLLS

- 1a. Blades with posterior rib (union of basal veins) naked along the edge of the sinus.
- 2a. Petioles with conspicuous, elongate, hair-like scales.
- 3a. Petiolar scales scattered and mostly in the distal 1/2 of petiole; plants terrestrial; Panama and Colombia, 830–860 m in Panama, 50–150 m in Colombia ..... *P. malesinchiae* Croat
- 3b. Petiole scales dense, covering the entire petiole; plants usually hemiepiphytic.
- 4a. Blades ovate-triangular, typically more than 1.3 times longer than broad, semiglossy above, paler and solid light green to silvery-green beneath, Costa Rica to Ecuador, <100–1250 m ..... *P. squamicale* Croat & Grayum
- 4b. Blades ovate, typically less than 1.3 times longer than broad, matte and subvelvety above, paler and tinged with purplish violet between the major veins below; Costa Rica to Peru, (200)500–1500 m ..... *P. serrucosum* L. Mathieu ex Schott
- 2b. Petioles glabrous or least lacking scales.
- 5a. Petioles conspicuously and sharply flattened adaxially, with lateral margins sharply edged or with a slender wing.
- 6a. Petioles merely sharply edged, lacking an actual wing at the margin.
- 7a. Blades whitish and matte below; cataphylls persisting in semi-intact fragments of thin epidermis; spathe tube green outside, at most tinged pink within; Costa Rica to Panama (Chiriquí), (775)1000–2100 m ..... *P. thalassicum* Croat & Grayum
- 7b. Blades green and semiglossy below; cataphylls persisting in a dense reddish brown, semi-intact mass; spathe tube green or bright red outside, dark red to maroon within, blade white; Belize to South America, 0–900 m.
- 8a. Blades ovate-triangular, 1.8–2.7 times longer than wide; cross-veins conspicuous between minor veins on dried blades; spathe tube green on outside; W Panama (Bocas del Toro to Coclé), 590–1300 m ..... *P. copense* Croat
- 8b. Blades mostly ovate, 1.1–1.7 times longer than wide; cross-veins between minor veins not at all apparent; spathe tube bright red outside; Belize to South America, 0–1000 m ..... *P. fragrantissimum* (Hook.) G. Don
- 6b. Petioles with a narrow marginal wing.
- 9a. Leaf blades promptly splitting into slender segments laterally, drying blackened; petiole wing undulate distally; Costa Rica to Panama, Atlantic slope, 0–1400 m ..... *P. findens* Croat & Grayum
- 9b. Leaf blades remaining intact, drying yellow-green; petiole wing usually undulate throughout its entire length; Nicaragua to central Panama, mostly less than 700 m ..... *P. pterotum* K. Koch & Augustin
- 5b. Petioles terete to obtusely flattened or U-shaped but not sharply flattened adaxially, if sulcate adaxially the margins merely obtuse.
- 10a. Blades whitish and matte on lower surface even on fresh leaves; petioles usually drying light yellowish brown (except sometimes not in *P. strictum*).
- 11a. Blades narrowly ovate, usually more than 1.8 times longer than wide; plants commonly hemiepiphytes; Panama to Ecuador, usually <1400 m ..... *P. hebetatum* Croat
- 11b. Blades ovate to broadly ovate, averaging 1.3 times longer than wide; typically terrestrial; Costa Rica to western Panama, Colombia, and Venezuela, mostly at 850–1665 m in Central America ..... *P. strictum* G. S. Bunting
- 10b. Blades green to yellow-green beneath, usually semiglossy to glossy; petioles drying various colors, not light yellowish brown (except *P. copense*, *P. schottianum*, and *P. thalassicum*).

NARROWLY OVATE BLADES

- 12a. Blades usually 1.8 times or more longer than wide.
- 13a. Spathe tube greenish to yellow-green or whitish inside.
- 14a. Leaf blades with lower surface matte, frequently bluish green; style apex prolonged into a short but distinct neck (style type D); central Costa Rica to western Panama, (775)1000–2100 m ..... *P. thalassicum* Croat & Grayum

- 14b. Leaf blades with lower surface glossy to semiglossy, not at all bluish green; style apex flat, not prolonged into a distinct neck (style type B; rarely type C); Panama, except *P. jodavisanum* (Mexico to Venezuela).
- 15a. Blades drying yellow-green; basal veins fewer than 5; sap milky white, drying chalky; inflorescence solitary; cataphylls to 40 cm long; Panama (Darién), 800-1480 m ..... *P. albissimus* Croat
- 15b. Blades drying dark gray-brown to olive-green, typically somewhat blackened; basal veins usually more than 5; sap usually clear, drying dark brown to reddish; inflorescences 2-6 per axil; cataphylls less than 20 cm long.
- 16a. Petioles terete to obtusely flattened adaxially, lacking prominently raised lateral margins; peduncles frequently longer than the spathe; central and eastern Panama, 0-800 m ..... *P. panamense* K. Krause
- 16b. Petioles usually D-shaped or U-shaped, usually flattened adaxially with prominently raised margins, often thicker than broad, rarely C-shaped to subterete; peduncles usually much shorter than the spathe; S Mexico to Panama and Venezuela (Mérida), 0-1500 m ..... *P. jodavisanum* G. S. Bunting
- 13b. Spathe tube red to maroon or violet-purple on inside.
- 17a. Blades drying greenish to yellow-green (sometimes yellow-brownish in *P. alticola*), lacking conspicuous cross-veins.
- 18a. Pistils with 1 ovule per locule; leaf blades with secretory ducts moderately obscure, the abaxial surface densely and minutely granular on drying; staminate portion of the spadix constricted above the sterile portion, the fertile portion clavate, the sterile staminate portion only slightly thicker than the pistillate portion; dried style base doughnut-shaped with stylar canals at the bottom of a deep concavity; Panama (Chiriquí to Coclé, (750) 950-2200 m) and Costa Rica ..... *P. stramineocala* Croat
- 18b. Pistils with 12-18 ovules per locule; leaf blades with secretory ducts conspicuous, abaxial surface smooth on drying; staminate portion of the spadix scarcely constricted above the sterile portion, the fertile portion stubby and evenly tapered to the apex, the sterile staminate portion much thicker than the pistillate (approximately ½ thicker); dried style base raised but flattened at apex with a narrow pale ring around its outer margin, the stylar canals exerted like minute funnels and extending above the surface; eastern Costa Rica and western Panama, 800-2500 m ..... *P. alticola* Croat & Grayum
- 17b. Blades drying reddish brown with conspicuous cross-veins.
- 19a. Petioles drying with a conspicuous light reddish brown or yellow-brown epidermis, this smooth and often flaking; cataphylls less than 25 cm long; primary lateral veins 11-16 per side; basal veins frequently more than 8 per side; Panama (Bocas del Toro and Coclé Provinces), 590-930 m ..... *P. copense* Croat
- 19b. Petioles drying dark brown, the epidermis not peeling; cataphylls more than 55 cm long; primary lateral veins fewer than 8 per side; basal veins up to 8 per side; Panama (Chiriquí, Bocas del Toro, Coclé, and Veraguas Provinces), 500-1630 m ..... *P. chiriquense* Croat

## OVATE TO BROADLY OVATE BLADES

- 12b. Blades usually about 1.5 times longer than wide or less (sometimes wider than long).
- 20a. Leaf blades with lower surface matte (but never velvety), frequently bluish green, the dried waxy surface forming an areolate pattern; central Costa Rica to western Panama, (775) 1000-2100 m ..... *P. thalassicum* Croat & Grayum
- 20b. Leaf blades with lower surface semiglossy (or, if matte, then velvety), never bluish green.
- 21a. Leaf blades matte and velvety above; central Panama in the region of the isthmus, 300-375 m ..... *P. gigas* Croat
- 21b. Leaf blades semiglossy to glossy above.
- 22a. Peduncles usually less than 10 cm long (rarely longer in fruiting peduncles of *P. llanense* and *P. dodsonii*).
- 23a. Spathe tube solid bright red on outside; central Panama (Veraguas and Coclé), 800-1200 m ..... *P. antonioanum* Croat
- 23b. Spathe tube green outside (sometimes tinged weakly red on *P. schottianum*).
- 24a. Spathe weakly or not at all constricted above the tube.
- 25a. Cataphylls persisting usually with large fragments of glossy, yellowish epidermis; petioles drying yellowish to yellowish brown and glossy as if with a layer of shellac; Costa Rica to Panama, 600-2200 m ..... *P. schottianum* H. Wendl. ex Schott
- 25b. Cataphylls persisting semi-intact and brown, not glossy or yellowish; petioles usually drying brown to blackened, usually not yellowish and glossy; Guatemala, 1360-1870 m ..... *P. parulense* Croat
- 24b. Spathe moderately to prominently constricted above the tube.
- 26a. Leaf blades drying yellow-brown below, coriaceous; ovules mostly (6)12-(20) per locule; Isthmus of Panama, mostly below 500 m ..... *P. llanense* Croat

- 26b. Leaf blades drying reddish brown, moderately thin; ovules 1-6 per locule (except *P. dodsonii* with ca. 20 ovules per locule); Costa Rica or Mexico.
- 27a. Stems usually growing over rocks in dry habitats, rarely on trees; cataphylls eventually deciduous, intact on the older stem; ovules 4-6 per locule; western Mexico (Jalisco & Guerrero), 350-1250 m ..... *P. basii* Matuda
- 27b. Stems appressed-climbing on trees in humid habitats; cataphylls decomposing and breaking up before falling off; ovules either solitary or ca. 20 per locule; Costa Rica and/or Ecuador.
- 28a. Spathe tube green; ovules 1 per locule; posterior rib bare-ly or not at all naked along the sinus (up to at most 2.5 cm); southwestern Costa Rica, 1000 m .....  
..... *P. dominicense* Croat & Grayum
- 28b. Spathe tube dark reddish to red-purple; ovules ca. 20 per locule; posterior rib naked to ca. 6 cm along the sinus; Costa Rica, principally on the Atlantic slope at 300-850 m and Colombia and Ecuador at 200-1300 m .....  
..... *P. dodsonii* Croat & Grayum
- 22b. Peduncles usually more than 10 cm long (rarely less than 10 cm long) when not at anthesis.
- 29a. Stems clothed with a dense series of overlapping cataphylls, each persisting as a reticulum of coarse fibers often overlain with a thin, fragmented epidermis.
- 30a. Blades 1.2-1.5 times longer than wide, drying reddish brown above; Panama (Darién), 1000-1560 m ..... *P. pirense* Croat
- 30b. Blades 0.8-1.4 times longer than wide, drying grayish green above.
- 31a. Dried blades with lower surface semiglossy, sparsely granular, with moderately conspicuous cross-veins, not speckled, the upper surface lacking raphide cells; ovules 6 per locule; Panama, Panama Province (Cerro Jefe), 550-800 m ..... *P. jefense* Croat
- 31b. Dried blades with lower surface matte, smooth and minutely brownish to whitish speckled, lacking obvious cross-veins, the upper surface usually with short whitish raphide cells visible; ovules 14-20 per locule; Panama, Canal Area to Darién Province (Cerro Sapo), 140-800 m ..... *P. lazarii* Croat
- 29b. Stems with only a few cataphylls, these usually semi-intact or with a few, thin, disorganized fibers.
- 32a. Blades drying reddish brown to dark yellow-brown; peduncles usually less than half as long as the spathe, straight below the spathe; spathe tube reddish purple to red outside; Costa Rica, Colombia, and Ecuador, 200-1300 m ..... *P. dodsonii* Croat & Grayum
- 32b. Blades drying dark gray-brown to olive-green, typically somewhat blackened above; peduncles usually as long as or longer than the spathe, frequently bent just below the spathe; spathe tube greenish on outside; Panama, 0-800 m ..... *P. panamense* K. Krause
- 1b. Blades with posterior rib (union of basal veins) not naked along the sinus, flanked with laminar tissue.
- 33a. Petioles conspicuously scaly distally; terrestrial (except *P. serrucosum*).
- 34a. Blades less than 25 cm long; petiolar scales close and overlapping, broad, usually less than 3 times longer than wide; endemic to El Copé region of Panama, ca. 800 m ..... *P. hamnelii* Croat
- 34b. Blades usually more than 30 cm long; petiolar scales more widely scattered, many times longer than wide.
- 35a. Plants terrestrial; stems usually creeping laterally across the ground; spathes glabrous; Mexico and Guatemala, 580-1900 m ..... *P. glanduliferum* Matuda subsp. *glanduliferum*
- 35b. Plants appressed hemiepiphytic climbers; stems usually climbing trees; spathes conspicuously covered with slender scales, Costa Rica to Peru, 200-1500 m (mostly >500 m) .....  
..... *P. verrucosum* L. Mathieu ex Schott
- 33b. Petioles glabrous and smooth distally, at least not conspicuously scaly; usually hemiepiphytic.
- 36a. Plants consistently terrestrial; petioles D-shaped with raised margins and a medial rib; spathe tube greenish to whitish within (or usually reddish on Pacific slope of Costa Rica); Nicaragua (Zelaya) to Panama, Colombia, and Ecuador; mostly 0-750 m ..... *P. grandipes* K. Krause
- 36b. Plants usually hemiepiphytic (*P. tysonii* frequently terrestrial in some habitats); petioles terete to subterete; spathe tube maroon to red or violet-purple within.
- 37a. Spathe not at all constricted above the tube, the spathe wall usually to ca. 1 cm thick midway; leaf blades very coriaceous, minor veins etched in the upper surface of fresh leaves; central Costa Rica to western Panama, 1100-2600 m ..... *P. crassipathum* Croat & Grayum
- 37b. Spathe visibly constricted above the tube (except only weakly constricted for *P. antioianum*), the wall usually less than 3 mm thick except at very base; leaf blades not markedly coriaceous, minor veins never etched into surface of fresh leaves.

- 38a. Primary lateral veins frequently 10 or more per side; spathe tube white to pale green within (sometimes tinged red at very base); Nicaragua to S Ecuador and to Venezuela; 20-1400 m ..... *P. tenae* K. Koch & Augustin
- 38b. Primary lateral veins usually fewer than 10 per side (to 10 in *P. tysonii*); spathe tube red to red-violet, maroon, or violet-purple throughout within.
- 39a. Blades markedly sinuate along the margins; Mexico (Veracruz), less than 500 m ..... *P. subincisum* Schott
- 39b. Blades with the margins entire, not at all sinuate.
- 40a. Fresh cataphylls usually red; petioles and midribs lacking conspicuous purplish spots; basal veins 5-10 per side.
- 41a. Blades with primary lateral veins typically 6-10 per side, upper surface drying somewhat blackened and smooth; cataphylls unribbed to weakly 1-ribbed; spathe tube green or merely tinged reddish outside; ovaries with fewer than 7 sub-basal ovules contained in an ovule sac; western and central Panama, 600-1500 m ..... *P. tysonii* Croat
- 41b. Blades with primary lateral veins usually 5-6 per side, upper surface drying dark brown to gray-brown with prominent cross-veining; cataphylls sharply 2-ribbed; spathe tube dark maroon outside; ovaries with about 30 axile ovules not contained in an envelope; endemic to Veraguas and Coelé Provinces at 850-1150 m ..... *P. antonioanum* Croat
- 40b. Fresh cataphylls green (purplish-spotted in *P. edenudatum* and *P. grayumii*); petioles and midrib frequently purplish-spotted; basal veins usually 3-5 per side.
- 42a. Blades mostly more than 1.8 times longer than wide; petioles 0.85-0.90 times as long as the blade; dried blades without secretory ducts visible on lower surface; Panama, 110-1150 m ..... *P. edenudatum* Croat
- 42b. Blades mostly less than 1.8 times longer than wide; petioles 1-1.4 times longer than the blade; dried blades with secretory ducts clearly visible on the lower surface; Costa Rica to central Panama, 0-1630 m ..... *P. grayumii* Croat

## KEY 4: VINES WITH CORDATE BLADES AND DECIDUOUS CATAPHYLLS

[Note: None of the species in this group have the posterior rib of the leaf blades naked along the sinus except *P. brunneicaule*, *P. hederaceum*, *P. mexicanum*, and sometimes (weakly so) *P. coloradense*, *P. cottonense*, *P. sulcicaule*, *P. wilburii*, and *P. zhuangum*.]

- 1a. Internodes elongate, usually much longer than broad; plants usually vines or at least prominently scandent.
- 2a. Blades with the posterior rib (coalesced basal veins) not naked (i.e., not running on the margin of the sinus).
- 3a. Stems coarsely asperous, covered with branched scales or setose-pubescent.
- 4a. Blades ovate-triangular, subcoriaceous; posterior lobes slender, usually flaring; stems reddish brown, covered with trichome-like, frequently branched scales; berries pale yellow; styles very short; occurring usually in wet habitats; Nicaragua to Panama and South America, 60-280(1300) m ..... *P. brevspathum* Schott
- 4b. Blades ovate-cordate, thin; posterior lobes directed toward one another at maturity; stems greenish, covered with elongate setae; berries pale orange to red or reddish orange; usually occurring in dry habitats; Mexico to Panama, the West Indies, and South America, 0-1500(2500) m ..... *P. jacquinii* Schott
- 3b. Stems smooth, not covered with branched scales.
- 5a. Petioles prominently flattened dorsiventrally with the lateral margins sometimes acute; Nicaragua to Ecuador, 10-1500 m ..... *P. platypetiolatum* Madison
- 5b. Petioles terete to subterete, not prominently flattened dorsiventrally.
- 6a. Peduncles usually much shorter than the spathe, usually less than 7 cm long at anthesis, drying 6-10 mm diam.; ovules many (usually 10-20) per locule.
- 7a. Leaf blades narrowly ovate-elliptic or triangular-sagittate or triangular-hastate, usually 1.8 times or more longer than wide; posterior rib more than 3 cm long.
- 8a. Leaf blades narrowly ovate-elliptic; posterior lobes about as broad as long, directed toward the base; sinus  $\pm$  V-shaped; ovules ca. 20 per locule; Mexico (Chiapas), 0-1000 m ..... *P. breedlovei* Croat
- 8b. Leaf blades triangular-sagittate to triangular-hastate; posterior lobes typically much longer than broad, directed somewhat outward; sinus usually not V-shaped but parabolic to spatulate; ovules usually 1-2 per locule; Mexico to Colombia, 0-1900 m ..... *P. mexicanum* Engl.
- 7b. Leaf blades ovate, usually only slightly longer than wide (rarely to 1.9 times longer than wide in *P. hederaceum*); posterior rib virtually lacking or to at most 1 cm long.
- 9a. Stems drying dark reddish brown, prominently ribbed, usually densely warty; Costa Rica to Ecuador and Suriname; (250)300-900 m ..... *P. hederaceum* var. *kirkbridei* Croat
- 9b. Stems drying either green and only weakly striate, or light yellowish brown and deeply fissured on drying.



- 10a. Stems drying yellowish brown, prominently ridged, smooth and glossy; spathe tube violet-purple outside; Costa Rica to Ecuador, 0-1600 m ..... *P. purpureoviride* Engl.
- 10b. Stems drying greenish, weakly striate, matte; spathe tube green outside; Mexico to West Indies, widespread in South America ..... *P. heteraceum* (Jacq.) Schott
- 6b. Peduncles frequently as long as or longer than the spathe, sometimes more than 7 cm long, usually drying 3-4 mm diam.; ovules 1 or 2 per locule.
- 11a. Leaf blades lacking any obvious primary lateral veins; stems prominently sulcate-ribbed even when fresh; ovules 1 per locule; SE Costa Rica to central Panama; 100-700 m ..... *P. sulcicaule* Croat
- 11b. Leaf blades with obvious primary lateral veins; stems smooth (or at least not regularly and prominently sulcate-ribbed longitudinally when fresh); ovules 2 per locule; Panama (Darién); 850-1560 m ..... *P. clevelandii* Croat

CORDATE VINES WITH DECIDUOUS CATAPHYLLS AND BLADES WITH NAKED POSTERIOR RIBS

- 2b. Blades with the posterior rib naked and running on the margin of the sinus for usually 1.5 cm or more (sometimes to as little as 1 cm in *P. smithii*).
- 12a. Stems setose-pubescent; blades membranaceous on drying; spathe tube inflated, with considerable space between the inner surface of the spathe and the spadix; pistils elongated at apex into a style to ca. 5 mm long; Mexico to Panama, Cuba, and northern South America; 0-1500(2500) m ..... *P. jacquinii* Schott
- 12b. Stems glabrous; blades coriaceous to subcoriaceous on drying; spathe not inflated-bulbous at base, fitting tightly over spadix; pistils not at all elongated into a distinct styler region.
- 13a. Stems prominently and regularly sulcate-ribbed when fresh.
- 14a. Blades less than 24 cm long, lacking obvious primary lateral veins; petioles to 21 cm long; NE Costa Rica to Isthmus of Panama; 100-700 m ..... *P. sulcicaule* Croat
- 14b. Blades more than 26 cm long, with 3-4 obvious pairs of primary lateral veins; petioles more than 29 cm long; S Mexico to Guatemala; 700-1525 m ..... *P. serapazense* Croat
- 13b. Stems smooth or irregularly fissured but not regularly sulcate-ribbed when fresh.
- 15a. Blades mostly broadly ovate, usually more than 25 cm wide; posterior lobes usually broadly rounded, about as wide as long; stems drying light reddish brown with flaking epidermis; Costa Rica to Ecuador, 100-1300 m ..... *P. bracteicaule* Croat & Grayum
- 15b. Blades mostly ovate-triangular or triangular-sagittate to triangular-hastate, mostly less than 25 cm wide (sometimes broader in *P. smithii*); Mexico to Panama; 0-2000 m.
- 16a. Blades sagittate to hastate at base; posterior lobes 2-3 times longer than broad; posterior rib directed straight toward the apex of the posterior lobe and 1.5-3.5 cm distant from the posterior margin of the blade; Mexico to Panama, 0-1900 m ..... *P. mexicanum* Engl.
- 16b. Blades cordate or sagittate (rarely hastate in some forms of *P. wilburii*); posterior lobes usually less than 2 times longer than wide (except sometimes more than 2 times longer in *P. wilburii*, and then blades drying reddish brown, not greenish as in *P. mexicanum*); posterior rib mostly curved along and near the margin of the sinus, rarely more than 1 cm from the posterior margin of the blade.
- 17a. Cataphylls 2-ribbed; blades usually more than 35 cm long; petiole often with a purple distal ring; peduncle usually longer than the spathe; Mexico to Nicaragua, 40-1630 m ..... *P. smithii* Engl.
- 17b. Cataphylls 1-ribbed (rarely 2-ribbed); blades usually less than 36 cm long; petioles lacking a purple distal ring; peduncle usually shorter than the spathe (except often longer in *P. wilburii* var. *longipedunculatum*); central Costa Rica to central Panama, 0-2000 m ..... *P. wilburii* Croat & Grayum

CORDATE NON-VINES WITH DECIDUOUS CATAPHYLLS AND NON-NAKED POSTERIOR RIBS

- 1b. Internodes of mature stems broader than long or sometimes somewhat longer than broad but not elongate with scandent stems (possibly somewhat scandent in *P. broadlovei* and *P. souae*); plants mostly appressed-climbing hemiepiphytes.
- 18a. Petioles either densely covered with scales or D-shaped with undulate-winged margins.
- 19a. Petioles terete, less than 1 cm diam., densely covered with conspicuous, spreading scales; blades almost as broad as long, the major veins on the lower surface densely puberulent; Panama to Ecuador; 0-1300 m ..... *P. squamipetiolatum* Croat
- 19b. Petioles sharply D-shaped or sharply flattened with undulate-winged margins, more than 1.5 cm diam., glabrous; blades much longer than broad; veins of lower surface glabrous; Panama (Chiriquí); 1100-1300 m ..... *P. fortuneense* Croat
- 18b. Petioles glabrous, terete to merely obtusely flattened.
- 20a. Posterior rib naked along the sinus.
- 21a. Peduncle usually less than 10 cm long.
- 22a. Blades less than 25 cm long, the adaxial surface drying dark brown to yellow-brown.
- 23a. Blades ovate-triangular, 1.8-3 times longer than wide; posterior lobes 1.4-1.8

- times longer than wide; inflorescence solitary; Panama (Chiriquí), 1750–2100 m ..... *P. knappiae* Croat
- 23b. Blades ovate, 1.3–1.6 times longer than wide; posterior lobes about as wide as long; inflorescences 2–3 per axil; Panama (Chiriquí), 1600 m ..... *P. coloradense* Croat
- 22b. Blades more than 35 cm long, the adaxial surface usually drying green to brownish green to grayish green to dark olive-green (sometimes dark brown in *P. zhuangum* or blackish in *P. smithii*).
- 24a. Petioles on drying not markedly flattened, 8–15 mm diam., usually greenish to dark brown; less than 360 m along the Caribbean coast in Costa Rica ..... *P. aromaticum* Croat & Graym
- 24b. Petioles on drying markedly flattened, to 3–6 cm diam., often yellowish; 700–900 m, Coclé Province, Panama ..... *P. zhuangum* Croat
- 21b. Peduncle usually 10 or more cm long.
- 25a. Leaf blades with margins convex in lower ½ of blade.
- 26a. Stem drying light brown; epidermis often flaking; petioles spongy, but drying moderately smooth, flattening to ca. 1 cm wide; blades drying thin and lacking conspicuous secretory canals; 40–1630 m (mostly less than 600 m), Mexico to Nicaragua ..... *P. smithii* Engl.
- 26b. Stem drying dark brown; epidermis sulcate or cracked but not flaking; petioles firm, drying heavily wrinkled, flattening to 3–6 cm wide; blades drying coriaceous with conspicuous secretory canals; 700–900 m, Panama (Coclé) ..... *P. zhuangum* Croat
- 25b. Blades with margins straight to concave in lower ½ of blade; Costa Rica and Panama.
- 27a. Leaf blades drying pale yellow-green, lacking any sign of secretory ducts on lower surface; ovules 1 per locule; Costa Rica, 1350–1400 m ..... *P. stramineicula* Croat
- 27b. Leaf blades drying dark olive-green or dark brown, with secretory ducts visible alternating with the minor veins on lower surface; ovules 4–5 per locule; near Costa Rican–Panamanian border, 1100–1950 m ..... *P. cotoniense* Croat
- 20b. Posterior rib not naked along the sinus, or weakly and obscurely naked near its base.
- 28a. Blades with posterior lobes turned inward and overlapping; peduncles slender, drying ca. 2–4 mm diam.; 850–1560 m, Serranía de Pirre, Darién Province, Panama ..... *P. clewellii* Croat
- 28a. Blades with posterior lobes never overlapping (except sometimes in *P. ferrugineum*), usually directed backward or somewhat outward; peduncles stout, usually 5–10 mm diam. on drying.
- 29a. Spathe barely or not at all constricted near the middle or above the tube portion; sterile staminate portion of the spadix often conspicuously (50%) broader than the pistillate portion at anthesis; Mexico to Guatemala, 70–2500 m ..... *P. advena* Schott
- 29b. Spathe usually visibly constricted above the tube (usually somewhat below the middle); sterile staminate portion of the spadix narrower to only slightly broader than the pistillate portion at anthesis.
- 30a. Inflorescences 1–3 per axil; spathes less than 11 cm long (except 9.5–18 cm long in *P. brevesii*).
- 31a. Primary lateral veins mostly more than 8 pairs; minor veins etched-sunken on upper blade surface; lower blade surface ± bluish green; 800–2200 m, Costa Rica and Panama ..... *P. brevesii* Standl.
- 31b. Primary lateral veins 4–8 pairs (sometimes to 9 pairs in *P. sagittifolium*); minor veins flat or raised on upper blade surface; lower blade surface medium green to yellow-green, not bluish green; Mexico and Belize.
- 32a. Leaf blades ca. 1.6 times longer than wide; ovules 1–7 per locule.
- 33a. Blades drying gray-green below; sap white; ovules 1 per locule; Belize, less than 500 m ..... *P. dreyeri* Croat
- 33b. Blades drying dark yellow-brown below; sap clear, turning brown?; ovules 4–7 per locule; Panama, 1600 m ..... *P. coloradense* Croat
- 32b. Leaf blades ca. 1.8–2 times longer than wide; ovules 2–4(5–8) or ca. 20 per locule; Mexico (Chiapas, 1300 m) or widespread, Mexico to South America, 150–2700 m.
- 34a. Blades ca. 1.8 times longer than wide; inflorescence 1 per axil; ovules 20 per locule; Mexico (Chiapas), 1300 m ..... *P. breedlovei* Croat
- 34b. Blades 1.85–2 times longer than wide, inflorescences 1–3 per axil; Mexico (Veracruz) to Colombia, 0–1800 m ..... *P. sagittifolium* Liebm.
- 30b. Inflorescences 4–6 per axil; spathes 15–23 cm long.

- 35a. Cataphylls usually more than 25 cm long; blades usually more than 55 cm long, drying coriaceous, reddish brown, with minute interrupted ridges on lower surface; Panama, 100–770 m ..... *P. ferrugineum* Croat
- 35b. Cataphylls less than 25 cm long; blades usually less than 55 cm long, drying subcoriaceous, brown to olive-green, smooth or minutely ridged but not with interrupted ridges on lower surface; Mexico (Chiapas), 490–1400 m ..... *P. souae* Croat

***Philodendron advena*** Schott, Oesterr. Bot. Wochenbl. 5: 289. 1855. TYPE: Mexico. Exact location uncertain, described from living material, Schott s.n. (lectotype, here designated, L 900230 (89889152)). Figures 37–40, 73.

*Philodendron subovatum* Schott, Oesterr. Bot. Wochenbl. 5: 289. 1855. TYPE: Western Mexico: a cultivated collection prepared by Schott (holotype, W now destroyed). Field Museum Photo 29864 (neotype, here designated; duplicate photo at MO).

*Philodendron monticola* Matuda, Madroño 10: 170. 1950. TYPE: Mexico. Chiapas: Sierra Madre, Pacific slope, Colonia San Juan Panamá, 50 km E of Escuintla, 1600 m, Matuda 18169 (holotype, MEXU; isotypes, DS, NY, UCLA).

*Philodendron jamapatum* G. S. Bunting, Gentes Herb. 9: 336, fig. 242. 1965. TYPE: Mexico. Veracruz: Coscomatepec-Huatusco, Route 155 at Riá Jamapa, Moore & Bunting 8872 (holotype, BH).

Terrestrial, epilithic, or hemiepiphytic; stem appressed-climbing, to 2 m long, sap brownish orange; internodes weakly glossy, 2–4 cm long, 1.5–2.5(5.5) cm diam., longer than broad, sometimes obtusely flattened on one side, medium green to gray-green, drying brown to yellow-brown, epidermis smooth to closely fissured; roots few per node, drying dark brown to yellow-brown, epidermis sometimes flaking; cataphylls subcoriaceous, 7.7–45 cm long, sharply D-shaped to sharply 2-ribbed, rarely weakly 2-ribbed, light green to whitish, tinged reddish, generally deciduous intact, weakly emarginate at apex, margins clear. LEAVES erect-spreading to spreading with blades pendent; petioles 30–45(65) cm long, 7–18 mm diam., erect-spreading, subterete to D-shaped, somewhat spongy to moderately firm, obtusely flattened adaxially, sheath 3–11 cm long; blades ovate-cordate to sagittate, subcoriaceous, moderately bicolorous, gradually acuminate at apex, cordate at base, (32)41–47(64) cm long, 16.5–40 cm wide (1.2–2.5 times longer than wide; 1–1.5 times longer than petioles), margins hyaline, upper surface dark green, drying dark brown to blackened, sometimes yellow-green, lower surface slightly more glossy, paler, drying yellow-brown to reddish brown, sometimes greenish brown; anterior lobe 20.5–53 cm long, 12–40 cm wide (3.4–3.8 times longer than posterior lobes); posterior lobes rounded, 6–14(20) cm long, 5.4–15(20) cm wide, directed toward base; sinus  $\pm$  parabolic to V-shaped or spatulate, 8–15

cm deep; midrib broadly convex to flat, slightly paler than surface above, convex and slightly paler below; basal veins 3–6 per side, with 0–1(2) free to base, coalesced (0.4)1.3–4(8.5) cm; posterior rib not naked or obscurely naked to 1 cm at base; primary lateral veins (3)4–6 per side, departing midrib at a (45)50–60(80) $^\circ$  angle, straight or weakly arcuate to the margins, flat to sunken and paler than surface above, convex below; minor veins weakly raised, moderately visible, arising from both the midrib and primary lateral veins. INFLORESCENCES spreading-pendent, 1–2(4) per axil; peduncle 5–13.5(17) cm long, 7–11 mm diam.; spathe coriaceous, (6.5)12–19 cm long ((0.7)1.4–2.4 times longer than peduncle), weakly or not at all constricted above the tube, acuminate, usually greenish throughout, broadest at or below the middle; spathe blade greenish white outside, spathe red to maroon or greenish white, glossy inside; spathe tube sometimes reddish outside, 5–8 cm long, 2.5–3.5(5) cm diam., red to maroon or dark violet-purple, pale-lineate, glossy inside; spadix stipitate to 4 mm long; clavate to cylindrical, bluntly pointed at apex, 11.5–14 cm long, broadest at the middle or  $\pm$  uniform throughout; pistillate portion 3.3–6.9 cm long, 1.2–2.7 mm diam. at middle, narrowed somewhat at both ends; staminate portion 5–9.2 cm long; fertile staminate portion tapered to cylindrical or clavate, usually longer than pistillate portion, 11–14 mm diam. at base, 8–15 mm diam. at middle, 5–7 mm diam. ca. 1 cm from apex, broadest below middle, equal in length to pistillate portion, narrower than pistillate and sterile portions; sterile staminate portion, 11–30 mm diam., usually broader than pistillate portion at anthesis; pistils 1.7–8.5 mm long, 1.3–2.9 mm diam., ovary (5)6–8-locular, with basal to sub-basal placentation; ovules 1–4 per locule, arranged digitately in translucent ovule sac, 0.2–0.6(1.5) mm long, equal in length to funicle; funicle adnate to lower part of partition, style 0.9–3 mm long, similar to style type B; style apex flat; stigma discoid to subdiscoid; thecae cylindrical to elliptical, 0.5 mm wide, contiguous. INFRUCTESCENCE with spathe green outside, dark purple-violet inside, 11.5–14 cm long; berries white to yellowish; seeds 1(3–4) per locule, oblong-ellipsoid, sometimes obovoid, 1.7–2.2 mm

long, 0.7–1.8 mm diam., with clear longitudinal and faint latitudinal striations and speckled with shiny, silver raphide cells; funicular scar knob-like, clear, thicker than the seed body.

Flowering specimens of *Philodendron advena* have been collected in post-anthesis condition virtually throughout the entire year. Mature fruits have been collected less consistently with July and August the only consecutive months without fruits reported.

*Philodendron advena* ranges from Central Mexico (Veracruz) along the Atlantic slope to Chiapas and then along both the Pacific and Atlantic slopes to southern Guatemala (Santa Rosa and Chiquimula Departments). It occurs at 70 to 2500 (mostly above 1000) m elevation in a wide variety of vegetation types, including "Selva Alta Perennifolia," "Bosque Caducifolia," "Selva Mediana Subperennifolia," and "Bosque Caducifolia."

*Philodendron advena* is a member of *Philodendron* sect. *Calostigma* subsect. *Macrobelum* ser. *Macrobelum*. The species is distinguished by internodes longer than broad, its two-ribbed, generally deciduous cataphylls, terete to obtusely flattened, somewhat spongy petioles (about as long as the blades), and by its generally coriaceous, ovate-cordate, usually dark brown to blackened-dried to somewhat sagittate-lobed blades with rounded lobes with usually four to six primary lateral veins, and usually one to two green inflorescences with the inner surface reddish to purplish.

Material from the Pacific slope in Chiapas State, Mexico, and in adjacent Guatemala often dries somewhat more greenish. This was described by Matuda as *P. monticola* Matuda. Considering the variability in the species, that taxon is not warranted. Though not closely related to *P. purulhense* Croat, *P. advena* is perhaps most easily confused with that species, because both may have similar ovate, black-drying leaves. *Philodendron purulhense* differs from *P. advena* in its persistent cataphylls, a more or less elliptic spathe scarcely constricted above the tube, and ovaries with axile placentation and 13–20 ovules per locule.

*Philodendron advena* is probably most closely related to the equally ecologically versatile *P. sagittifolium*, which occurs throughout much of its range. *Philodendron sagittifolium* is distinguished by having usually longer, more prominently sagittate blades (averaging 1.5–3 times longer than broad), which dry typically reddish brown rather than blackened. In contrast, *P. advena* has blades averaging 1.5 times longer than broad (ranging from 1.3 to 1.8 times longer). Both *P. advena* and *P. sagittifolium* have five to eight locules per pistil

with up to three more or less basal ovules (or they have the funicle adnate to the lower part of the axillary wall but extending down to the base).

Two Oaxaca collections are noteworthy. Moore & Bunting 8891 from above Valle Nacional in the Serranía de Juárez of Oaxaca dried greenish brown on the lower surface and gray-brown above. Also unusual is Moore & Bunting 8919 (collected near 8891), which dried reddish brown. This collection was discussed as possibly distinct by Bunting (1965). Aside from the color of the blades upon drying, there is nothing else out of the ordinary with the collection (see fig. 235 in Bunting, 1965).

*Additional specimens examined.* GUATEMALA. Alta Verapaz: near Chirriacté, on the Petén Highway, ca. 900 m, Standley 92192 (F); ca. 6 km NE of Panzós, 800 m, Croat 41689 (MO); 7 mi. up the road to Oxec, 700 m, 41651 (MO); Finca Argentina, 15 mi. W of Telemán, 550–650 m, 41571 (MO); 2–4 mi. E of Tamahú, 9–11 mi. E of Hwy. CA-14 to Cobán, 700–800 m, 41498 (MO); near Tactic, ca. 1500 m, Standley 70495 (F); near Tactic, above Río Frío, 1400–1500 m, 90484 (F). Chiquimula: Cerro Tixit, 3–5 mi. N of Jocotán, 500–1500 m, Steyermark 31563 (F). Quezaltenango: in reserve INDE "Santa María," km 99, 14°45'N, 91°32'W, Croat & Hanson 63430 (K, MO, US, USCG); El Pocito, S of San Martín Chile Verde, on road to Colombia, ca. 2200 m, Standley 85039 (F), 85093 (F); between San Martín Chile Verde and Colombia, above Mujulú, ca. 1800 m, 85723 (F), 85561 (F); between Finca Pirineos and Finca Patulún, 1200–1400 m, 86875 (F); Río Samalá, near Santa María de Jesús, 1500–1650 m, 84571 (F), 84669 (F). San Marcos: Canjulá-La Unión Juárez, near SE portion of Volcán Tacaná, 2000–3000 m, Steyermark 36407 (F); Volcán Tajumulco, 1400–1700 m, 37391 (F); ca. 3.3 mi. above Finca Armenia, above San Rafael, 1600 m, Croat 40963 (MO, NY); San Rafael-Pic de la Cuesta-Palo Gordo, near Aldes Fraternidad, 1800–2400 m, Williams et al. 26014 (F, NY). Santa Rosa: Cenaguilla, 4000 ft., Heyde & Lux 4281 (G, NY, US). Zacapa: summit of Sierra de las Minas, vic. of Finca Alejandria, 2500 m, Steyermark 29855 (F); Río Lima, Sierra de las Minas, below Finca Alejandria, 2000 m, 30026 (F). MEXICO. Chiapas: Lagos de Montebello, 3 mi. W of Dos Lagunas, 1460 m, Croat 46636 (MEXU, MO), 46655 (MO); ca. 6 mi. NW of Pueblo Nuevo Solistahuacán, vic. km 99, 1900–1950 m, 17°07'N, 92°52'W, Croat & Hanson 65193 (MEXU, MO); 8 mi. NW of Pueblo Nuevo Solistahuacán, 1900–1950 m, Croat 46429 (MEXU, MO), 46430 (MEXU, MO); Ixtapa-Pichucalco, 4777 (B, K, MO); Bochil-Pichucalco, 3.4 km W of El Anepe, 1860 m, Croat 78610 (CHIP, CM, K, MO, NY); San Cristóbal-Palenque, 97 km NE of San Cristóbal, 1130 m, 17°27'N, 92°04'W, Hammel et al. 15624 (MO); Palenque-Bonampak, 89–90 mi. SW of Palenque, 350–370 m, Croat 40221 (MO); San Juan Panamá, Escuintla, 1800 m, Matuda 18485 (MEXU, NY); Cascada, Siltepec, 1800 m, 19643 (MEXU); Paso Pashital, Sierra Madre de Chiapas, 1800 m, 37585 (MEXU); Cerro Brujo, Ocozacoatlán de Espinosa, MacDougall 415 (MEXU); ca. 6 mi. SW of Motozintla de Mendoza, ca. 1850 m, Croat 47246 (MO); 26–30 mi. N of Motozintla de Mendoza, 9.5–4.5 mi. S of Siltepec, 1000–1300 m, 47462 (MO); Ojo de Agua, Soconusco, Nakamura 38 (DS); San Fernando-Merzillas,

4-66 mi. NW of San Fernando, 840-940 m, 16°53'N, 93°16'W, *Croat & Hannon 65006* (B, MEXU, MO, US); Mpio. Angel Albino Corzo, above Finca Cuxtepec, 1380 m, *Breedlove 52073* (DS); Mpio. Berriozábal, 13 km N of Berriozábal, 900 m, 35438 (DS); 16 km NW of Rizo de Oro, 1600 m, *Breedlove & Smith 31382* (DS); Mpio. Ocosingo, Laguna Ocotal Grande, 3300 ft., *Breedlove 15683* (F, MICH); Mpio. Rayón, Selva Negra, 10 km above Rayón Mezcalapa, along road to Jitotol, 1700 m, 23291 (DS); Mpio. Unión Juárez, Volcán Tacaná, above Talquiam, 2200 m, *Breedlove & Almeda 47712* (CAS); Mpio. Villa Corzo, E base of Cerro Tres Picos, near Cerro Bola, 1500 m, 16°10'N, 93°15'W, *Breedlove & Thorne 30193* (DS). Oaxaca: Sierra de Juárez, Highway 175, 6-14 mi. from bridge at Valle Nacional, ca. 580 m, *Moore & Bunting 8891* (BH); Vista Hermosa, 17.9 mi. from bridge at Valle Nacional on road to Oaxaca, 8919 (BH); 660 m, *Croat 78721* (CHIP, MO); 4.3-6 mi. above bridge at Valle Nacional, 705 m, 17°44'N, 96°19'W, *Croat & Hannon 65549* (CM, MEXU, MO, NY, US); 14 mi. W of Valle Nacional, 1210 m, *Croat 39783* (MEXU, MO), 39796 (MO); 21.5 mi. above Valle Nacional, 1660-1670 m, 48108 (MO); Teotitlán del Camino-Chilchocla, 2.2 mi. beyond turnoff to Huautla de Jiménez, 2270 m, 48335 (CM, MO); 3.8 mi. past turnoff to Huautla de Jiménez, 2265 m, 48369 (MO); Ixtlán, Río Soyalapan watershed, 765-805 m, 17°37'N, 96°17'W, *Boyle et al. 3943* (MO). Tabasco: Teapa-Tacotalpa, 3.1 mi. E of Teapa, ca. 0.25 mi. S of highway, 150 m, 17°33'N, 92°59'W, *Croat & Hannon 65376* (MO); Mpio. Teapa, 7 km SE of Teapa on road to Tacotalpa, Rancho San Enas, Sierra Madragal, 70 m, 17°35'N, 92°50'W, *Hammel & Movello 15519* (MO, NY); Río Puyacatengo, Sierra Madragal, 7 km E of Teapa, trail from Centro Regional del Sureste de Universidad Autónoma Chapingo, 300 m, 18°32'N, 92°55'W, *Schatz et al. 1178* (MO). Veracruz: Veracruz-Oaxaca, Uxpanapa, 150 m, 17°05'N, 84°35'W, *Hammel & Merello 15563* (MO); above San José de Gracia, 1 mi. S of hwy. between Córdoba and Veracruz, 750 m, *Croat 39618* (MO); Conejo-Huatusco, at km 45, barranca de Santa María, across hwy. from Hacienda El Mirador, *Moore & Bunting 8856* (BH); ca. 1200 m, *Croat 44013* (MO); Mendoza-Orizaba, Sierra San Cristóbal, Highway 150-D, ca. 3 km SW of Orizaba, 1260-1400 m, *Croat 39549* (CM, MEXU, MO); Fortín-Huatusco, Highway 125, 300 m, 19°06'N, 97°02'W, *Croat & Hannon 63107* (B, COL, CR, CM, F, G, INPA, K, LL, M, MO, NY, QCA, P, PMA, US, VEN); Acacajete, 1750 m, *Cházaro & Robles 3115* (WIS); Mpio. Coacoatzintla, 1300 m, *Cházaro 1053* (F); Mpio. Ixhuacán, Río Ixhuacán, El Aguacate-Patlalanan, Puebla, 1500 m, *Cházaro & Cházaro 4068* (MO, WIS); Mpio. Jalapa, 8 km NW of Consolapan, 1525 m, ca. 19°30'N, 96°59'W, *Nee 29683* (MO, NY); Coapexpan, 1470 m, *Young 35* (F, MO); Mpio. Tlanelhuyocan, Barranca del Piquisac, Rancho Viejo-Ya Vega, 1650 m, *Cházaro & Ramarillo 3895* (MO, WIS); Mpio. Yecuatla, Los Capulines, near Paz de Enriquez, ca. 8 km N of Chiconquico, 1400-1600 m, 19°47'N, 96°49'W, *Taylor et al. 147* (F, MO, NY).

***Philodendron albisuecus* Croat, sp. nov. TYPE:**

Panama. Darién: Parque Nacional del Darién, middle slopes on W side of Cerro Pirre, 800-1500 m, 7°56'N, 77°45'W, 29 June 1988, *Croat 68940* (holotype, MO-3589989; isotypes, B, COL, K, PMA, US). Figures 41-43.

Planta hemiepiphytica; internodia 2-3 cm longa, 2.8 cm diam.; succus albus, in sicco calcareus; cataphylla leviter 1-costata vel acute 2-costata, persistentia ut fibræ tenues, pallidae; petiolus usque 51-72 cm longus, subteretes, leviter complanatus basi usque ad medium; lamina anguste ovata, cordata basi, 50-66 cm longa, 24.5-42 cm lata; inflorescentia 1; pedunculus (3.7)7-12 cm longus; spathe 13.5-14.5 cm longa, extus viridis, intus pallide flaviviridis; pistilla 5-6-locularia; locules cum 18-20 seminibus; baccae albae.

Hemiepiphytic; stem ± appressed-climbing, to 75 cm long, to 2.8 cm diam., sap chalky white; internodes short, usually sparsely covered with cataphyll fibers, 2-3 cm long, 2.8 cm diam., dark green; roots brownish, few per node, epidermis smooth to flaking and peeling; cataphylls thin, to 40 cm long, weakly 1-ribbed to sharply 2-ribbed, sparsely short-lineate, persisting as thin, pale fibers at upper nodes, then deciduous; petioles 51-72 cm long, 3-7 mm diam., erect-spreading, subterete, dark green, weakly flattened from near base to middle, obtusely sulcate at the base adaxially, surface semiglossy; blades narrow ovate, subcoriaceous, splitting, long-acuminate at apex (the acumen in-rolled), cordate at base, 50-66 cm long, 24.5-42 cm wide (1.6-2 times longer than wide), (ca. about equal in length to petiole), broadest just below point of petiole attachment, upper surface dark green, semiglossy, lower surface semiglossy, moderately paler, drying yellow-green; anterior lobe 37-48(54.5) cm long, 23.1-27.4(42.4) cm wide (4.2-4.4 times longer than posterior lobes); posterior lobes 8.5-11.5(16.5) cm long, 9.7-12(18.5) cm wide; sinus hippocrepiform; midrib flat, concolorous above, bluntly acute, slightly paler than surface below; basal veins 3-5 per side, with 1 free to base, third and higher order veins coalesced 0.5-2 cm long, the fifth and sixth veins sometimes coalesced to 5 cm long; posterior rib naked, raised; primary lateral veins (4)6-7(8) per side, departing midrib at a 60-65° angle, sunken above, raised to convex below; interprimary veins darker than surface; minor veins arising from both the midrib and primary lateral veins; lesser veins obscure to visible. INFLORESCENCES 1 per axil; peduncle (3.7)7-12 cm long, 3-3.5 mm diam., pale green; spathe semiglossy, 13.5-14.5 cm long (1.2-1.8 times longer than peduncle), weakly constricted above the tube, green throughout, pale yellow-green (greenish brown post-anthesis) within; spathe tube densely short pale lineate outside, 5-7 cm long; spadix 12 cm long at anthesis; pistillate portion 4.3 cm long in front, 3 cm long in back, 8 mm diam. midway, 7 mm diam. at apex; staminate portion to 10 cm long; fertile staminate portion to 9 mm diam. toward apex, somewhat narrower just

above the sterile portion, to 5 mm diam. 1 cm from apex; sterile staminate portion 9 mm diam.; pistils 6–6.3 mm long, 1.8–2.5 mm diam., ovary 5–6-locular, with axile placentation; ovules 2-seriate, style similar to style type B; style apex flat; stigma subdiscoid. INFRACTESCENCE 6.4 cm long, 3 cm diam.; berries white; seeds 18–20 per locule.

Flowering in *Philodendron albisuccus* is poorly known with a flowering collection seen in October, immature fruits in June, and nearly mature fruits in July.

*Philodendron albisuccus* is endemic to Panama, known only from the Serranía de Pirre in Darién Province, at 800 to 1400 m elevation in *Premontane rain forest*.

*Philodendron albisuccus* is a member of *Philodendron* sect. *Philodendron* subsect. *Cardiobelium* ser. *Albisuccosa*. The species is characterized by its short internodes; thin, obscurely one-ribbed cataphylls, which persist at the upper nodes as thin, pale fibers; subterete petioles (about as long as the blades); the narrowly ovate, moderately cordate blades with a hippocrepiform sinus; and especially by the sap in all the cut plant parts, which promptly turns white and chalky upon exposure to air. Only one other species in Panama, *P. cretosum* Croat & Grayum, has chalky sap. Though both species have chalky sap (a rare feature for *Philodendron*) and 18–20 ovules per locule, I do not believe them to be closely related as they show no other features in common. *Philodendron cretosum* differs most particularly in having slender oblong-lanceolate blades that are 5.1–8.3 times longer than wide (vs. 1.6–2 times longer than wide for *P. albisuccus*). *Philodendron albisuccus* also has the much longer petioles (51–72 cm long vs. 10–23 cm long for *P. cretosum*), more broadly spreading primary lateral veins (60–65° vs. 25–45° angle for *P. cretosum*). In addition, *P. albisuccus* has minor veins more distinct than those of *P. cretosum*.

The species is superficially similar to *P. alticola* Croat and *P. straminea* Croat & Grayum, both of which have blades of similar size, shape, and color upon drying. Both differ by having the inner surface of the spathe red rather than green.

*Additional specimens examined.* PANAMA, Darién: Cerro Sapo, ca. 5 km S of Garachiné, 600–800 m, 7°59'N, 78°25'W, Hammel et al. 14820 (MO, US); Cerro Pirre region, Altos de Nique–Cerro Pirre, ca. 8 km N of Alturas de Nique, ca. 8 km W of Cana Gold Mine, 1430–1480 m, Croat 37851 (MO, PMA); Río Cana–Río Escucha Ruido, along ascent of Serranía de Pirre above Cana Gold Mine, 1310–1430 m, 37830 (MO, US); Río Cana, SW of Cerro Pirre, vic. of old gold mine at Cana, 1400 m, 27319 (MO); Parque Nacional Darién, slopes of Cerro Mali, headwaters of S branch of Río Pucuro, ca. 22 km E of

Pucuro, 1300–1400 m, 8°04'30"N, 77°14'W, Cuadros et al. 3961 (MO).

***Philodendron alticola* Croat & Grayum, sp. nov.**

TYPE: Panama. Chiriquí: Parque La Amistad, 3.5 mi. W of Cerro Punta, 2 km inside park along old abandoned roads and trails, 800 m, 8°53'N, 82°35'W, 28 Mar. 1993, Croat 74906 (holotype, MO–4343624–5; isotypes, B, CR, F, K, PMA, US). Figures 45–48.

Planta epiphytica aut hemiepiphytica; internodia (1.5–2)4–10 cm longa, 2–4 cm diam., in sicco pallide flavibrunnea, subtiliter costata; cataphylla usque 28 cm longa, incostata, in sicco pallide brunnea, decidua; petiolus 42–70 cm longus, 7–10 mm diam.; lamina ovato-sagittata, 44–72 cm longa, 21–44 cm lata, sagittata basi, in sicco brunnea supra, flavibrunnea infra; inflorescentia 1–2; pedunculus 5–12 cm longus; spatha 11.5–14(18) cm longa, viridis; pistilla (4)5–6(7)-locularia; loculi 12–18-ovulati.

Epiphytic or hemiepiphytic; stem appressed-climbing; internodes dark green, promptly gray-green, finally light brown, semiglossy, finely ribbed, (1.5–2)4–10 cm long, 2–4 cm diam., drying pale yellow-brown, epidermis sometimes loosening and flaking free in small patches; cataphylls sharply D-shaped with weakly raised margins, pale green, weakly glossy, drying thin, to 28 cm long, unribbed, drying pale brown, deciduous at lower nodes, persisting semi-intact at upper nodes with a few exposed pale fibers, in part persisting as pale fibers at lower nodes with a few pale fibers and small, thin fragments of pale brown epidermis. LEAVES erect-spreading to spreading; petioles 42–70 cm long, 7–10 mm diam., subterete, medium green, weakly glossy, faintly lineate, obtusely flattened adaxially, weakly spongy, drying greenish to blackened, slightly flattened toward the apex adaxially, surface drying ± matte, sometimes with portions of epidermis loose and bubbled, tan and translucent; sheathing subtending an inflorescence, to 6 cm long; blades narrowly ovate-sagittate, subcoriaceous to moderately coriaceous, semiglossy, somewhat bicolorous, weakly and shortly acuminate at apex, sagittate at base, 44–72 cm long, 21–44 cm wide (1.6–2 times longer than wide), (about as long as petioles), broadest somewhat above point of petiole attachment, upper surface dark green, matte, drying brown to greenish brown and almost matte, lower surface moderately paler and weakly glossy, drying yellow-brown; anterior lobe 34–35.5 cm long, margins convex; posterior lobes 13–26 cm long, 11–17 cm wide, directed downward; sinus obovate, 10–19 cm deep; midrib speckled, slightly paler, drying broadly convex, concolorous above, narrowly rounded, slightly paler, drying prominently convex, often flat and irregularly ridged, yellow-

ish brown below; basal veins 5-6 per side, with the first free to base, the second coalesced 1-5 cm, third and higher order veins coalesced 4-7.5 cm long; posterior rib naked for 1-3 cm; primary lateral veins 5-8 per side, those near end of blade departing midrib at a 50-65° angle, those in the lower one-half of blade departing at 80-95° angle, broadly curved to the margins, obtusely sunken and slightly paler, drying weakly raised above, convex and paler than surface, splayed out and downturned at the midrib, drying weakly raised below; interprimary veins inconspicuous; minor veins moderately distinct, fine, arising from both the midrib and primary lateral veins, drying prominulous, alternating with secretory canals appearing as intermittent lines, sometimes branched, blackened and minutely sunken. INFLORESCENCES erect, 1-2 per axil; peduncle  $\pm$  terete, coarsely striate (coarser on spathe and base of tube, pale short-lineate otherwise), weaker toward apex, 5-12 cm long, drying 7-10 mm diam., pale green, drying dark brown; spathe coriaceous, 11.5-14(18) cm long, scarcely or not at all constricted midway, green to yellowish green throughout, acuminate; spathe blade dark green throughout, drying dark brown and unmarked, smooth outside, pale greenish white to white on upper two-thirds of blade inside; spathe tube 2.8-4 cm long, 2-3 cm diam., red to red-violet to purplish on lower one-third of tube inside; spadix 9-14 cm long; pistillate portion pale green, 0.8-1.2 cm long, 1-1.5 cm diam. throughout; staminate portion 8-12.5 cm long; fertile staminate portion creamy white, stubby, evenly and bluntly tapered to apex, 1.5-2 cm diam. throughout, broadest at the base, much broader than the pistillate portion; sterile staminate portion 1.5 cm diam.; pistils 3-4 mm long, 1.7-1.8 mm diam., ovary (4)5-6(7)-locular, with axile placentation; ovules 12-18 per locule, 2-seriate, weakly translucent, 0.4-0.5 mm long, longer than funicle; funicle 0.1-0.2 mm long, adnate to lower part of partition, style similar to style type B; central style dome sometimes weakly developed; dried style base raised but flattened apically with a narrow, pale ring around its outer margin; stylar canals emerging as tiny funnels at base of small apical depressions and arranged separately in a ring, drying as minute funnels extending above the surface of the style boss; style apex weakly concave; central domes and small depressions surrounding stylar canal exits; stigma covering entire style apex; androecium truncate, oblong prismatic, margins irregularly 4-5-sided or 4-6-sided, 0.9-1 mm long; thecae  $\pm$  oblong, ca. 0.3 mm wide; sterile staminate flowers in part prismatic, in part weakly clavate or irregularly 4-6-sided,

1.2-1.8 mm wide. INFRUCTESCENCE in early fruit to 14 cm long; spathe becoming dark brown with paler lines in tube; berries white with dried style bases ca. 2 mm long, brown, with a narrow smooth ring around its margin; seeds tan, ovoid, weakly mammiliform, 1.4-1.8 mm long, 1-1.2 mm diam.

Flowering in *Philodendron alticola* is documented by just a few collections and might be bimodal. Specimens in Panama were collected both in bud and immature fruit in February (nearly mid-dry season in Panama), but post-anthesis collections were also made in August and September. No mature fruits were seen. The region where *P. alticola* occurs is decidedly less seasonal (at least in terms of totally rainless days) than is much of the rest of Panama. This might induce a less seasonal flowering in *Philodendron*.

*Philodendron alticola* is known only from eastern Costa Rica and western Panama in the region adjacent to the frontier at 800 to 2500 m elevation in *Tropical Lower Montane wet forest* or possibly *Tropical Lower Montane rain forest*. It is one of the most high-ranging species of *Philodendron* subg. *Philodendron* Schott in Central America, hence the name "alticola" (meaning high-dweller).

*Philodendron alticola* is a member of *P. sect. Philodendron* subsect. *Philodendron ser. Fibrosa*. In addition to the high elevations at which it occurs, the species is characterized by having the internodes longer than broad and drying yellowish brown and finely ribbed; by its subterete petioles about twice as long as the blades, drying darkened and sometimes with bubbly epidermis; the ovate-sagittate blades, which dry mostly dark brown above and yellowish green below with conspicuous blackened, interrupted and sometimes branched secretory ducts; and by the usually paired, short-pedunculate inflorescences with the spathe scarcely constricted above the tube, green outside, white inside on the blade and red to purple on the tube within. The species is most similar to *P. stramineicaule*, which is also subscandent and has similarly colored blades. That species differs in lacking prominent intermittent laticifers and the fertile staminate portion of the spadix constricted above the sterile portion; the sterile staminate portion of the spadix only slightly thicker than the pistillate portion; and a doughnut-shaped style with the style tubes in the bottom of a concavity upon drying. In addition, it has only 1 ovule per locule (vs. 12-18 per locule for *P. alticola*). In contrast, *P. alticola* has conspicuous laticifers, the staminate spadix scarcely or not at all constricted above the sterile

portion, and the sterile portion of the spadix much thicker than the sterile staminate portion. The dried style base of *P. alticola* is flat at the apex with erect, tubular-extended styler pores.

The species is also similar to *P. schottianum* H. Wendl. ex Schott, which also shares the conspicuous latex canals on the lower blade surface and a similarly stubby spadix, but that species differs in having shorter internodes, cataphylls that persist as a dense mass of fibers (vs. semi-intact or as only a few pale fibers interspersed with small fragments of epidermis in *P. alticola*), by its petioles, which dry smoother and more matte, and by its more broadly ovate blades, which often have conspicuous cross-veins.

*Additional specimens examined.* COSTA RICA. **Lámón:** Cordillera de Talamanca, Atlantic slope, Valle de Silencio, along Río Terbi, 0.5–1.5 airline km W of Costa Rican–Panamanian border, 2300–2400 m, 9°8'N, 82°57'W, *Davidse et al.* 28735 (CR, MO); S side of unnamed cordillera between Río Terbi and Río Sinf, 2–4 airline km W of Costa Rican–Panamanian border, 2300–2500 m, 9°9–11'N, 82°57–58'W, *Davidse et al.* 28931 (MO). PANAMA. **Chiriquí:** vic. of Las Nubes, 2.7 mi. NW of Río Chiriquí Viejo, W of Cerro Punta, 2200 m, *Liesner* 323 (MO, US), 325 (MO, US); Cerro Punta, Las Nubes, 2000 m, *Croat* 26492 (MO).

***Philodendron angustilobum*** Croat & Grayum, sp. nov. TYPE: Costa Rica. Heredia: Estación Biológica La Selva, confluence of Río Sarapiquí and Río Puerto Viejo, 10°26'N, 84°01'W, 50–80 m, *Grayum & Chavarría* 8302 (holotype, MO). Figures 49–52, 74.

Planta hemiepiphytica; internodia 3–6(15) cm longa, 1.5–3 cm diam.; cataphylla 12–33 cm longa, acute 2-costata; petiolus subteres, 17–54 cm longus, 5–10 mm diam.; lamina profunde 3-lobata, 30–48 cm longa, 20–48 cm lata; in sicco atricanua vel stribunnea supra, flavibrunnea vel atricanibrunnea infra; lobis lateralibus manifeste patentibus, in sicco denigratis infra; inflorescentia 1–3; pedunculus (4.5/9–11(15) cm longus; spathe (11)13–18 cm longa, lamina spatheae extus interdum suffusa alba; tubo spatheae pallide viridi, intus interdum suffusa marronino basi; pistilla 7–10-locularia; locali (1)2-ovulati.

Hemiepiphytic; stem appressed-climbing, growing to 2.5–6 m high in trees; internodes matte to weakly glossy, 3–6(15) cm long, 1.5–3 cm diam., dark green, drying gray (within 4–6 internodes), epidermis drying light brown, semiglossy, often peeling (underlying stem blackened), fissured transversely, longitudinally ridged; cataphylls soft (especially near base), 12–33 cm long, bluntly to sharply 2-ribbed or sharply 1-ribbed to near the apex then 2-ribbed below, green, sometimes tinged red, deciduous, persisting deciduous, intact; petioles 17–54 cm long, 5–10 mm diam., subterete,

somewhat spongy, dark green, weakly flattened toward apex with an obscure medial rib adaxially, with adaxial margins rounded to weakly angular, surface weakly glossy, drying blackened; sheathing 3–9 cm long; blades deeply 3-lobed, subcoriaceous, moderately to conspicuously bicolorous, 30–48 cm long, 20–48 cm wide (1–1.5 times longer than wide), (0.8–1.8 times longer than petioles), upper surface dark green, drying dark gray to dark brown, semiglossy, lower surface drying dark yellow-brown to dark gray-brown, weakly glossy, moderately paler; anterior lobe 28–39 cm long, 8.5–14(17) cm wide (1.8–1.9 times longer than lateral lobes), rarely broadest at the base, usually narrowed toward the base (4–12 cm wide just above the base); sinus arcuate to broadly V-shaped; lateral lobes (7.5)15–22(27) cm long, 3–8(12) cm wide, broadest usually toward the apex, directed outward, broadly spreading (90–127° angle), broadly confluent with medial lobe 3.3–6.5 (rarely to 14) cm, acute; midrib broadly convex to flat-sunken above, convex, slightly paler than surface, drying blackened below; basal veins 8–10 per side, pinately arranged along a stout medial rib; posterior rib not at all naked or rarely naked for 1–1.5 cm; primary lateral veins 5–6 per side (45–75° angle on median lobe), gradually curved to the margins, sunken above, convex, drying blackened below; interprimary veins weakly raised and darker than surface below; minor veins distinct, darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES 1–3 per axil; peduncle (4.5)9–11(15) cm long; spathe (11)13–18 cm long, 8–10 mm diam. (1.4–1.6(2.4) times longer than peduncles), greenish white outside, greenish white to white within; spathe opening broadly at anthesis, convolute in only lower one-half of tube, barely or not at all constricted above the tube; spathe blade sometimes tinged white outside; spathe tube light green, sometimes tinged maroon at base outside; spadix held erect, not protruded forward, oblong to oblong-tapered, rounded to bluntly acute at apex, (9.5)11–15.5 cm long, broadest near the middle, constricted weakly above sterile staminate portion; pistillate portion pale green, oblong to oblong-tapered, 4.5–5.7 cm long, 8–10 mm diam. at apex, 8–12 mm diam. at middle, (5)11–13 mm wide at base; staminate portion 8–11 cm long; fertile staminate portion oblong-tapered, 8–13 mm diam. at base, (6)12–13 mm diam. at middle, 8–9 mm diam. ca. 1 cm from apex, usually broader than the pistillate portion; sterile staminate portion ca. 1 cm diam.; pistils 1.5–3 mm long, 1.1–1.8 mm diam., ovary 7–10-locular, (0.8)1.1 mm diam., with sub-basal placentation; ovules (1)2 per



locule, contained within transparent ovule sac, 0.6 mm long, usually shorter than funicle, style similar to style type B; style apex rounded; stigma button-like, subdiscoid, 1 mm diam., 0.2–0.4 mm high, covering center of style apex; thecae oblong to oblong-ovate, contiguous, sometimes divaricate. JUVENILE plants with petioles 8–10 cm long; blades oblong, rounded to weakly subcordate.

Flowering in *Philodendron angustilobum* has been recorded from the early dry season through the mid-wet season, including February, March, May (the greatest number), and August, but too few collections exist to be certain of phenology.

*Philodendron angustilobum* ranges from Honduras (Olancho) to Panama, apparently being restricted to the Atlantic slope from Honduras to Costa Rica and ranging from near sea level to 680 m elevation. In Panama it has been collected on both slopes near the Continental Divide from 800 to 1430 m elevation, but a juvenile collection from near sea level in the Canal Area is probably also this species. It is known from *Tropical wet forest* and *Premontane wet forest* life zones.

*Philodendron angustilobum* is apparently a member of *P.* sect. *Tritomophyllum* [though M. Grayum (pers. comm.) believes it is closest to *P. ligulatum*, a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*]. This species is characterized by its moderately long internodes, spongy petioles, and 3-lobed blades, which dry blackened with broadly spreading lateral lobes that are much narrower and shorter than the medial lobes. It is superficially similar to *P. mexicanum* Engl., which has sagittate blades that dry green to brown and occurs typically in much drier habitats.

The sole Honduran collection might prove to represent another species or perhaps even a hybrid. It differs from material collected from Nicaragua, Costa Rica, and Panama in having the medial lobe broader, more triangular, and not at all constricted at the base.

Two collections (Croat & Grayum 59933 and Croat 67616) from the same plant from southwestern Costa Rica near Golfito differ in having the posterior lobes less narrowed; they may prove to represent another species.

*Additional specimens examined.* COSTA RICA. Cartago: CATIE, near Quebrada Molina, SE of Florencia de Turrialba, ca. 680 m, 9°52'30"N, 83°40'W, Grayum 3877 (MO). Heredia: Río Guacimal downstream from Monte Verde, 1300 m, Grayum 5418 (CR, MO). Puntarenas: Coto Brus, Las Cruces Botanical Garden, near San Vito de Java, ca. 4000 ft., Croat 32960 (MO); 6 km W of San Vito de Java, 1200 m, 8°49'N, 82°58'W, 57230 (MO); ca. 1 km NW of Golfito, 11 km SW of Interamerican Highway,

<100 m, 8°11'N, 83°12'W, Croat & Grayum 59933 (CR, F, K, MEXU, MO, US); Golfito-Villa Briceño, 3.1 mi. NW of center of Golfito, 30 m, 8°11'N, 83°12'W, Croat 67616 (CM, CR, MEXU, MO). HONDURAS. Olancho: San Esteban-Bonito Oriental, along Río Olancho, 3.3 mi. SW of border with Colón Dept., along Río Grande, 350–400 m, 15°31'N, 85°42'W, Croat & Hannon 64522 (CAS, CR, EAP, HNMN, K, MO, PMA, US). NICARAGUA. Zelaya: Río Sucio, 2 km E of Bonanza, 140 m, Neill 4024 (MO). PANAMA. Canal Area: ca. 1 mi. E of Fort Sherman, <25 m, 9°19'N, 79°57'30"W, Croat & Zhu 76283 (MO). Chiriquí: Cerro Colorado, 15.6 mi. above bridge over Río San Félix, 1330 m, Croat 48439 (MO), 24 mi. above bridge over Río San Félix, 1430–1500 m, 48489 (MO), 800–1000 m, 33180 (MO); Gualaca-Chiriquí Grande, near Lago Fortuna, trail to Río Hornito, 8°45'N, 82°18'W, 763724 (MO). Coeléc: vic. of La Mesa, N of El Valle de Antón, 800–900 m, 8°38'N, 80°09'W, 67209 (CM, MO). Panamá: Cerro Campana, above Su Lin Motel, 14759 (MO, NY, SCZ).

***Philodendron anisotomum* Schott, Oesterr. Bot.**

Z. 8: 179. 1858. *Philodendron fenlii* Schott var. *anisotomum* (Schott) Engl., in A. DC. & C. DC., Monogr. Phan. 2: 412. 1879. TYPE: Guatemala. Las Nubes, *Wendland* 321 (lectotype, here designated, GOET). Figures 44, 53–56, 75.

*Philodendron affine* Hemsl., Diagn. Pl. Nov. Mexic.: 37. 1879. TYPE: Guatemala. Barranca Honda, Volcán de Fuego, 3800 ft., Salvin s.n. (holotype, K).

*Philodendron dagilla* Schott, Oesterr. Bot. Z. 5: 179. 1858. TYPE: Costa Rica. Cartago and Aguacate, Oersted s.n. (holotype, destroyed). Schott ic. 2592 (neotype, here designated).

*Philodendron trisetum* Standl., Publ. Field. Mus. Nat. Hist., Bot. Ser. 18: 137. 1937. TYPE: Costa Rica. Alajuela: La Palma de San Ramón, 1050 m, Brenes 5762 (holotype, F).

Usually a hemiepiphytic vine or sometimes on rocks; stem appressed-climbing, green as juvenile, graying with maturity, minutely and densely striate, sap watery, unscented, leaf scars conspicuous, 8–10 mm long, 7–9 mm wide; internodes smooth, semiglossy, to 2.5 cm long, 4–10 mm diam., usually longer than broad, medium green to olive-green, epidermis thin, tan, peeling; roots olive green, smooth, few per node; cataphylls thin, semispongy, to 10 cm long, unribbed, bluntly or sharply 1-ribbed, green, drying pale yellow-green, deciduous. LEAVES erect to spreading; petioles 21–57 cm long, (2)3–9 mm diam., terete, moderately spongy, whitish toward apex, slightly flattened toward apex adaxially, surface dark green striate at base; blades triangular in outline, deeply 3-lobed, subcoriaceous, moderately bicolorous, long-acuminate at apex (the acumen apiculate, to 3 mm long), hastate at base, 20–35 cm long, 20–42 cm wide (0.8–1 times longer than wide), (0.6–1 times the

petiole length), upper surface dark green, semiglossy to glossy, drying dark brown to dark gray-green, lower surface semiglossy, paler, drying yellow-brown to yellow-green; anterior lobe oblong-lanceolate to oblanceolate, almost elliptic, 16–30 cm long, 7–15 cm wide (1–1.3(1.5) times longer than lateral lobes); lateral lobes broadly confluent 1–3(8) cm with medial lobe, 10–25 cm long, 3–8.7 cm wide, directed outward (90° angle from midrib), acute to bluntly acute; sinus arcuate; midrib  $\pm$  flat to sunken, paler than surface above, broadly convex below; basal veins 4–7 per side, sometimes with last vein free to base, most veins coalesced 1.5–13 cm, 2 veins coalesced to 17 cm, drying reddish brown to yellowish brown; posterior rib naked; primary lateral veins 4–5 per side, departing midrib at a 50–60° angle, narrowly sunken above, convex below; interprimary veins weakly sunken and concolorous above, weakly raised and darker than surface below; tertiary veins visible, darker than surface below; minor veins fine, close, weakly visible to distinct below, arising from both the midrib and primary lateral veins but mostly from the midrib. INFLORESCENCES erect-spreading, 1 per axil; peduncle 5.5–19.5 cm long, 3–8 mm diam.; **spathe** 7.4–16.6 cm long (0.8–1.4(1.7) times longer than peduncle); spathe blade green to greenish white to creamy yellow, tinged with violet-purple outside, tinged with violet-purple, at least sometimes, with yellowish resin canals visible inside; spathe tube green outside, 4–6 cm long, 2–3.5 cm diam., dark violet-purple inside; **spadix** sessile; white, drying golden-yellow throughout, tapered,  $\pm$  rounded at apex, 7.5–10.8 cm long, broadest near the base; pistillate portion pale green (post-anthesis), ellipsoid, 3.6 cm long, 1.3 cm diam. at apex, 1.2 cm diam. at middle, 6 mm wide at base; staminate portion to 8.4 cm long; fertile staminate portion white, ellipsoid, tapered at apex, 8 mm diam. at base, 5 mm diam. at middle, 5 mm diam. ca. 1 cm from apex, broadest at base, much narrower than the pistillate portion; sterile staminate portion tannish, 8 mm diam.; pistils (1.5)2.9–4.4 mm long, 1.7–2.3 mm diam., ovary (6)7–8-locular, (1)2.2–3.2 mm long, ovule sac 0.7 mm long, with sub-basal placentation; ovules 3 per locule, contained within translucent, gelatinous ovule sac, ca. 0.4 mm long, longer than funicle; funicle 0.2–0.3 mm long (can be pulled free to base), style similar to style type B; style apex flat; stigma discoid, 5–6 mm diam., 0.1 mm high, covering center of style apex; the androecium  $\pm$  prismatic, margins irregularly 4–6-sided, 0.7–1.1 mm long; thecae oblong, 0.3 mm wide,  $\pm$  divaricate; sterile staminate flowers irregularly 5–6-sided, 1 mm wide. IN-

FRUITESCENCE 9.5–16 cm long, peduncle to 14 cm long; spadix to 5.3 cm long, to 3 cm wide; berries orange, rhomboid; seeds 1–2(3) per locule, (16)21–22(29) per berry, light brown, 1.5–2 mm long, 0.5 mm diam., with weak constriction (nipple) opposite funicular end of seed.

Flowering in *Philodendron anisotomum* appears to be restricted to the rainy season in Central America. One collection was seen in flower in May but most are from July through December (the latter month is sometimes the beginning of the dry season in parts of Middle America). Immature fruits were collected from December through July, with mature fruits known only from March, April, and July.

*Philodendron anisotomum* ranges from Mexico to Costa Rica, at 30 to 1800 m elevation. In Mexico, the species ranges from Nayarit to Oaxaca and Chiapas mostly along the Pacific coast, but also occurs in Puebla (Ajencibre) and Morelos (Cuernavaca). One collection, *Moore & Bunting 8874*, from near Córdoba in Veracruz state, appears out of range for the species.

Collections from Guatemala are few, but all are from the Pacific slope except one collection from Baja Verapaz between El Chol and Rabinal (*Groat & Hannon 63670*). In Honduras and Costa Rica, the species is nearly restricted to the Pacific slope. In Costa Rica the species occurs in *Premontane moist forest*. *Philodendron anisotomum* is a member of *P. sect. Tritomophyllum*.

This species is distinguished by its deeply three-lobed blades with frequently much smaller, falcate lateral lobes broadly confluent with the medial lobe.

*Philodendron anisotomum* is easily confused with *P. tripartitum* (Jacq.) Schott, which differs in having proportionately narrower medial lobes (mostly 3–3.5, rarely to 1.7 times longer than broad) with 4–9 prominently sunken primary lateral veins and lateral lobes typically directly more or less toward the apex. It has only one ovule per locule and white fruits. In contrast, *P. anisotomum* has medial lobes less than 1.5–2.8 times longer than broad with 2–4(5) weakly sunken primary lateral veins and usually much smaller lateral lobes typically directed outward, as well as 3 ovules per locule, and orange fruits.

Material from Nayarit (*McVaugh 13363*, *Moore & Bunting 8703*) is not only geographically isolated from populations in Puebla, Cuernavaca, and Oaxaca, but also differs morphologically by having much larger blades (medial lobe >23 cm wide) with the lateral lobes narrowly rounded rather than pointed.

Costa Rican collections differ in sometimes having two inflorescences per axil, rather than solitary inflorescences, which is more typical.

*Philodendron dagilla* Schott, published on the same page as *P. anisotomum*, was considered a synonym of *P. tripartitum* by Krause (1913), but it is clearly synonymous with *P. anisotomum*. Schott distinguished it from *P. anisotomum* by its longer, less spreading lateral lobes, but they are well within the norm.

*Additional specimens examined.* COSTA RICA. 1911, without locality, *Pittier & Durand 3090* (BR); *Worthen s.n.* (MO). **Alajuela:** San Pedro de San Ramón, 1075 m, *Brenes 4874/33* (F); San Ramón, 1500–1600 m, *Tanduz 17719* (BM, K, P); San Ramón-Balsa, 2.3 km N of Río Balsa, 1050–1150 m, 10°11'N, 84°30'W, *Stevens 14193* (MO); ca. 5.7 km N of Quebrada Volio, 1100–1150 m, ca. 10°08'N, 84°29'W, *14165* (CR, MO); San Ramón-Balsa, 5.7 mi. N of San Ramón, 1200 m, *Croat 46837* (MO); San Ramón-Bajo Rodríguez, 940 m, *Croat 78892* (CR, INB, MO); San Ramón-San Lorenzo, 1 km S of Balsa, 1100 m, 10°10'N, 84°29'W, *Liesner & Judziewicz 14937* (CR, MO); 1–3 km E of San Ramón, ca. 1000 m, 10°05'N, 84°27'W, *Liesner 14198* (B, CR, MO); above Río San Luis, 15 km NW of San Ramón, 800 m, 10°14'N, 83°31'W, *Lent 3108* (F); Finca Los Ensayos, ca. 11 mi. NW of Zarcero, 900 m, *Croat 43522* (MO); Zarcero region, ca. 100 m, *Smith 4447* (MO); ca. 7.5 mi. N of Zarcero, ca. 1000 m, *Croat 43499* (MO); ca. 15 km N of Zarcero, 1350 m, *Williams et al. 29041* (F); along Rte. 9, ca. 2 km N of Cariblanco de Suraipiquí, 800 m, 10°17'N, 84°12'W, *Grayum et al. 8098* (MO); Atenas, near old Pan-American Highway, *Gentry 7828* (CR, MO); San Luis de Zarcero, Cantón Alfaro Ruiz, 1425 m, *A. Smith H1178* (F, MO); N of Carrizal, 1440 m, *Grayum 3085* (DUKE); Cantón San Carlos at Sucre, 975 m, *A. Smith H1669* (F, US); San Isidro de San Ramón, 1259 m, 10°04'46"N, 84°26'30"W, *Herrera 58* (AAU, M, MEXU, MO); Monteverde Biological Reserve, Río Peñas Blancas, 900 m, 10°18'N, 84°45'W, *Haber & Bello 7181* (MO); 1250–1350 m, 9°17'N, 84°84'W, *Burger et al. 10770* (CHAPA, CR); Río Peje, 4 km S of Ciudad Quesada, 960 m, *Lent 1274* (BM, CR, F, GH); Río Trojes, 2 km N of La Luisa, 1380 m, *1674* (CR, US). **Cartago:** Juan Viñas, 1300 m, *Carpenter 608* (US); S slope of Volcán Irazú, *Standley 36638* (US); *Stevens 48* (US). **Guanacaste:** La Cruz de Abangares, 1400 m, *Haber & Bello 2881* (MO); Monteverde, 1400 m, 10°22'N, 84°49'W, *Haber & Zuchowksi 8734* (CR, ENCB, L, MO, QCA); 8 km NW of Monteverde, 1200 m, 10°22'N, 84°51'W, *Haber & Zuchowksi 9518* (INB, MO). **Puntarenas:** Cordillera de Talamanca, foothills around Tres Colinas, 1800–1850 m, 9°07'N, 83°04'W, *Davidse et al. 25635* (MO); Cantón de Buenos Aires, Quebrada Dorora (tributary of Río Kuyé), ca. 9 km NE of Ujarrás, 1500 m, 9°17'30"N, 83°16'W, *Grayum 10275* (CAS, CR, INB, K, MO, US); Reserva Biológica Carara Estación Quebrada Bonita, 30 m, 9°46'N, 84°36'W, *Bello & Rojas 2286* (CR, INB, MO). **San José:** ca. 25 km N of San Isidro del General, along the Inter-American Highway, 1800 m, 9°29'N, 83°41'W, *Burger & Baker 10078* (CR, F); Finca Micos-Llano Limón, ca. 8.5 km by road W of Ciudad Colón, 550–650 m, 9°56'N, 84°18'W, *Grayum et al. 6096* (MO); San José, *Standley 47358* (US), ca. 1130 m, *41210* (US), *47334* (US), 1150 m, *33262* (US); 1.4 km NW of Brazil de Santa

Ana, 800 m, *Taylor 17378* (NY, US); San Sebastián, S of San José, 1160 m, *Standley 49293* (US); San Pedro Montes de Oca-Carrizabal, ca. 1200 m, *32830* (US), *1250 m, 41307* (US); Río María Aguilar, near San José, 1200 m, *38952* (US); Las Pavas, 1070 m, *36085* (US); camino de Hatillo, near San José, 1200 m, *32171* (US); Acosta, Z.P. Cerros de Escuzú, Río Tabarcia, 1600–1700 m, 9°50'52"N, 84°04'40"W, *Morales 2745* (CR, MO); Valle del Candelaria, 1000–1050 m, *Morales & González 4596* (CR, INB); Santiago de Puriscal, *Echeverría 838* (F). **EL SALVADOR. Abuchapán:** *Padilla 60297* (US), 1922, 297 (US), *60* (US); 2–3 mi. NE of Bridge Imposible, 1000–1250 m, *Croat 42162* (MO). **San Salvador:** Toncatepeque, *Caldén 200* (US); 650–850 m, *Standley 19199* (GH, NY, US). **GUATEMALA. Alta Verapaz:** Cobán, ca. 1300 m, *Standley 90275* (F); Semococh, 17 km from Seból, on Cobán Road, *Contreras 4714* (LL); 5 mi. N of Cobán, along Highway CA-14, 1300 m, *Croat 41406* (MO); 5 mi. S of Cobán, along Highway CA-14, 1300 m, *41364* (MO); San Juan Chamelco, *Wilson 41008* (F). **Baja Verapaz:** Mpio. Rabinal, El Chol-Rabinal, Highway 6, 8.7 mi. N of El Chol, 1330 m, 15°03'N, 90°29'W, *Croat & Hannon 63670* (GH, LL, MO, TEX). **Chiquimula:** Quezaltenango, 1200–1500 m, *Steyermark 31198* (F). **Escuintla:** Escuintla-Santa Lucía Colz, Río Jute-Río Pantealeón, *Standley 63390* (F); Río Burrión, NE of Escuintla, *89605* (F, MEXU). **Jalapa:** Los Chorrás along Río Pinule, 1 mi. W of San Pedro Pinula, 1400 m, *Steyermark 32931* (F). **Jutiapa:** San José Acatempo-Río de los Escalones, Cuesta de la Conora, 900–1200 m, *Standley 60616* (F). **Quezaltenango:** Reserve INDE, "Santa María," km 199, 1200–1300 m, 14°45'N, 91°32'W, *Croat & Hannon 63435* (B, BM, CM, MO, NY, US); Calahuaché, 1020 m, *Standley 67137* (F); below Santa María de Jesús, 1350–1380 m, *68386* (F); Finca Pirineos-Patzún, 1200–1400 m, *86844* (F), *86918* (F). **Retalhuleu:** Río Coyote, 4 km W of Retalhuleu, 300 m, *87496* (F). **Sacatepéquez:** near Las Lajas, 1200 m, *58284* (F); Pueblo Nuevo, 750 m, *66976* (F). **San Marcos:** Finca Armenia, near La Trinidad, above San Rafael, 1100–1250 m, *Croat 40826* (MO); La Trinidad, ca. 2 km from Finca Armenia, above San Rafael, 1100–1250 m, *40864* (MO); Finca El Porvenir "Número 6," Volcán Tajumulco, *Steyermark 37140* (F, MEXU); Volcán Tajumulco, above Finca Porvenir, 1300–1500 m, *37358* (F, US). **Santa Rosa:** El Molino, ca. 600 m, *Standley 78508* (F); Volcán Jujuytepeque, 6000 pp, *Heyde & Lux 4283* (CM, GH, NY, US). **HONDURAS. Olancha:** Río Olancha, Gualaco-San Bonito Oriental, 7.4 mi. NE of San Esteban, 540 m, 15°20'N, 85°42'W, *Croat & Hannon 64364* (MO, VEN). **MEXICO. Teas s.n.** (MO). **Chiapas:** 8.5 mi. NE of Escuintla, on gravel road to El Triunfo, 250 m, *Croat 43823* (MO); ca. 4 km N of Ovarado Turquia, 450–850 m, *47575* (MO); Tapachula-Unión Juárez, at km 13.5, 1.3 mi. N of Trinidad, ca. 1000 m, *47212* (MO); San Fernando-Moravillas, near Lago Malpaso, 4–6 mi. NW of San Fernando, 840–940 m, 16°53'N, 93°06'W, *Croat & Hannon 65029* (CM, MO); Mpio. Angel Albino Corzo, along Río Cuxtepeques near Finca Gadlow, 1270 m, *Breedlove 40162* (CAS, DS); Barr. Aguas Calientes, *Miranda 1740* (MEXU); finca between Finca Cuxtepeque and Finca Cabañas, 1100 m, *Breedlove & Daniel 71160* (CAS); Las Nubes, Guatimoc, *Miranda 1720* (MEXU); Cascada, Siltpeec, 1800 m, *Matuda 38677* (MEXU); Motozintla de Mendoza-Huixtla, 15 mi. S of Motozintla de Mendoza, 900 m, *Croat 40764* (MO); Esperanza, Escuintla, *Matuda 17791* (MEXU, MO), *Matuda s.n.* (MO), *16661* (F, MEXU, MICH, MO), *17768* (F, GH, MEXU); Zacatonal, Acacoy-

*agras*, 1200 m, 18364 (MEXU); Cerro Ovando, 900–1050 m, Croat 78479 (CHIP, CM, MO, NY); Escuintla, Turquía, Matuda 17066 (F, MEXU); Mpio. Angel Albino Corzo, 1380 m, *Breedlove & Almeda* 56917 (MO); 1380 m, *Breedlove & Strother* 46700 (MO); Mpio. Mapastepec, Sierra de Socomusco, new unfinished road to Tuxtla Gutiérrez from Hwy. 200, (5.5 mi. NW of turnoff to Mapastepec), 6.5–8.5 mi. up road, 15°32'N, 92°48'W, Croat & Hannan 63339 (B, BM, CM, MO, NY, PMA, US); Mpio. Villa Corzo, E. base of Cerro Tres Picos, near Cerro Baul, 1500 m, *Breedlove* 23968 (DS). **Jalisco**: 20–22 km S of Talpa de Allende, 1200–1450 m, *McVaugh* 23308 (MICH). **Morelos**: near Cuernavaca, *Pringle* 7015 (GH); Cuernavaca, *Rose & Hough* 4438 (US). **Nayarit**: Tepic–Jaliscoacán, near km 20, 700–800 m, *Moore & Bunting* 8703 (BH); at km 15, ca. 900 m, 8694 (BH); ca. 4 mi. E of Jaliscoacán on road to Tepic, 750 m, *McVaugh* 13363 (MICH). **Oaxaca**: Oaxaca–Pochutla, along Highway 175, 55.1 mi. S of Miahuatlán, 21.9 mi. S of Suchixtepec, 39.7 mi. N of turnoff to Pluma Hidalgo, 1540 m, Croat 46108 (MEXU, MO); Finotepa Nacional–Tlaxiaco, along Highway 125, 5.8 mi. N of Putla de Guerrero, ca. 1000 m, 45904 (MO); Oaxaca–Pochutla, ca. 56 mi. S of Miahuatlán, 6.9 mi. N of turnoff to Pluma Hidalgo, 1480 m, 46116 (DUKE, F, MO). **Puebla**: Ajenjibre, *Bravo* 1-4516 (MEXU). **Veracruz**: Córdoba–Veracruz, Ejido San José de Gracia, below Peñuelo, *Moore & Bunting* 8874 (BH).

***Philodendron annulatum* Croat, sp. nov. TYPE:**

Panama. Panamá: El Llano–Cartí road, 5 mi. from Pan American Hwy., virgin forest on steep slopes, along trail through forest (W side of road), 350 m, 9°17'N, 78°58'W, 16 July 1987, Croat 67346 (holotype, MO 3609133; isotypes, B, CM, COL, CR, K, PMA, RSA, NY, US). Figures 57–60.

Planta hemiepiphytica, interdum epiphytica aut terrestri; internodia 1–4(11) cm longa, 1.5–3.5 cm diam.; cataphylla obtuse usque acute 2-costata aut incostata; petiolus subteres, 14–42 cm longus, subspongiosus cum annulo purpureo apice; lamina oblonga vel oblongo-ovata, subcordata, 24–73 cm longa, 8–29 cm lata; inflorescentia 1–2(3); pedunculus 3.6–11.5(17) cm longus; spathe 12.3–22 cm longa, lamina spathe extus alba vel viridialba, intus pallide viridis, tubo spathe extus viridi, intus viridi, striato rubello vel marconino basi; pistilla (5)7–8-locularia; loculi 1–2(4–5)-ovulati; baccae pallide virides vel abae.

Usually hemiepiphytic, sometimes epiphytic or terrestrial; stem appressed-climbing, sometimes creeping, to 1 m long, sap clear, sticky, sometimes black pepper-scented, leaf scars conspicuous, to 2.5 cm long, to 1.5 cm wide; internodes semiglossy, sometimes becoming matte, sometimes scurfy, 1–4(11) cm long, 1.5–3.5 cm diam., usually longer than broad, dark green to grayish or grayish green to brown, epidermis thin, drying brownish, irregularly ridged, frequently transverse fissured, cracking off; roots olive-green to brownish, slender, 0.6–100 cm long, to 3 mm diam., semiglossy, somewhat twisting, drying pale; cataphylls to 34 cm

long, bluntly to sharply 2-ribbed (ribs to 5 mm high), green, densely short, dark green lineate or speckled, sometimes persisting as a rotting mass, eventually deciduous. LEAVES erect-spreading to spreading to weakly arching; petioles 14–42 cm long, ± terete, slightly spongy, medium green, obtusely flattened and sometimes with an obscure medial rib near apex adaxially, sometimes with adaxial margins erect, blunt, surface sparsely dark green short-lineate or unmarked, sometimes with a purple ring at apex, sheath 4–9.5 cm long; blades oblong to oblong-ovate, subcoriaceous, moderately bicolorous, semiglossy, subcordate at base, 24–73 cm long, 8–29 cm wide (2.5–3 times longer than wide), (ca. 1.7 times longer than petiole), margins moderately undulate, upper surface dark green, lower surface paler; anterior lobe 23.5–68 cm long, 11–27.5 cm wide (10.5–15.6 times longer than posterior lobes); posterior lobes usually narrowly rounded and somewhat spreading or broader than long, rarely cordulate and about as long as broad, 1.5–6.5 cm long, 4.1–11.6 cm wide; sinus open or closed with lobes overlapping; midrib flat to broadly convex, slightly paler than surface or concolorous above, convex to bluntly acute, mostly concolorous, sometimes minutely speckled, matte or slightly semiglossy below; basal veins (1)2–3(4) per side, usually all free to base, sometimes with 2 united for up to 2 cm; posterior rib usually absent, never naked when present; primary lateral veins 6–10 per side, departing midrib at a 60–70° angle, ± straight to the margins, obtusely sunken, paler than surface above, convex to weakly raised and darker than surface below; interprimary veins flat and darker than surface below; minor veins moderately distinct to ± obscure above and below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect to spreading, 1–2(3) per axil; peduncle 3.6–11.5(17) cm long, to 1.8 cm diam., somewhat flattened adaxially, green, sometimes tinged red, weakly lineate or unmarked; spathe 12.3–22 cm long (1.7–3.4 times longer than peduncle), to 3 cm diam. (when closed), oblong-linear, acute at apex, weakly constricted above the tube; spathe blade white to greenish white outside (opening 8.5 cm long, 5.5 cm wide), pale green, with orange resin canals inside; spathe tube green outside, 5–7 cm long, 2.3 cm diam., green, reddish to maroon or maroon-streaked at base inside; spadix stipitate to 4 mm long; white, 9–14.7 cm long, broadest above the middle, weakly constricted above sterile staminate portion; pistillate portion pale green, cylindrical, 2.9–5.4 cm long, 0.9–1.3 mm diam., weakly narrowed toward both ends; staminate portion 6.8–12.5 cm long; fertile staminate

portion white, cylindrical to clavate, weakly tapered at apex, (5-7)12-15 mm diam. at base, 8-16 mm diam. at middle, (4)6-9 mm diam. ca. 1 cm from apex, broadest at middle or at base; sterile staminate portion narrower or broader than pistillate portion, white, 4-15 mm diam.; pistils 1.5-2.1(3.5) mm long, ovary (5)7-8-locular, 0.8-1.2(2) mm diam., with sub-basal or basal placentation (or axile in *Croat 69250*); ovules 1-2(4-5) per locule, 1-seriate, contained within transparent ovule sac, 0.3-1.7 mm long, longer than or equal in length to funicle, style similar to style type B; style apex with the shallow depression completely covered by stigma; style boss sometimes shallow, broad; stigma discoid to button-like (brush-like), 0.5-1 mm diam., less than 0.3 mm high, drying with the crown button-shaped and flat to concave with moderately large styler pores around its periphery; the androecium margins 5(4-6)-sided, somewhat scalloped, 2 mm long; thecae oblong, 0.5 mm wide, nearly contiguous, sometimes divaricate; sterile staminate flowers  $\pm$  rounded. **INFRUCTESCENCE** (immature), pale yellow-green outside; spadix to 3 cm wide, pistillate spadix 8-11 cm long; berries pale green to white; seeds tan, 1 mm long, 0.6 mm diam. **JUVENILE** petioles sheathed nearly throughout; blades oblong, obtuse to rounded at base, lower surface sometimes maroon.

Flowering in *Philodendron annulatum* is vouchered by one collection in January, with the vast majority of post-anthesis collections made primarily between February through May, and single collections in August and October. Mature fruits have been seen for only June and July.

*Philodendron annulatum* is endemic to Panama, ranging from Veraguas (Santa Fe) to Panamá and San Blas (El Llano-Cartí Road), at 100 to 970 m elevation in *Tropical wet forest* and *Premontane rain forest*.

*Philodendron annulatum* is a member of *P.* sect. *Calostigma* subsect. *Macrobelyium* ser. *Macrobelyium*. The species is distinguished by having internodes somewhat longer than broad and drying light brown; usually sharply two-ribbed, deciduous cataphylls; somewhat spongy obtusely somewhat flattened petioles with a purple distal ring (hence the name "annulatum" from *annulus*, meaning "having a ring"); oblong to oblong-ovate, subcordate blades; and one to two inflorescences per axil with the spathe blade white and the tube green outside (usually colored or tinged red to maroon within).

*Philodendron annulatum* is most easily confused with *P. ligulatum* Schott. *Philodendron correae* Croat also has narrow, usually cordulate leaf bases

and petioles with a purplish annular distal ring. Both differ from *P. annulatum* in having narrower blades broadest at or above the middle and narrowest just above the narrowly cordulate base. Alternatively, blades of *P. annulatum* are broadest at or near the base. *Philodendron annulatum* can also be confused with narrow-leaved representatives of *P. sagittifolium*. That species differs in having typically purple-spotted petioles (and often midribs) that lack a purplish distal ring.

**Additional specimens examined.** PANAMA. Cocle: 46 km N from Penonomé on road to Coclesito, 100 ft., *Hammel 1698* (MO); El Copé region, Alto Calvario, on Continental Divide, 5.2 mi. above El Copé, 930 m, *Croat 49186* (COL, K, MO); Alto Calvario above El Copé, ca. 6 km N of El Copé, 710-800 m, 8°39'N, 80°36'W, 68728 (AAU, CAS, K, L, M, MEXU, MO, PMA, TEX, US); 860 m, *Folsom 1257* (MO); 4.1 mi. N of El Copé, 680-770 m, 8°39'N, 80°36'W, 74831 (CAS, CM, F, MO, PMA), 74855 (L, MO, PMA); E of sawmill along old logging road, 2600 ft., *Hammel 4076* (MO); 4.5 mi. N of El Copé, 2.5 mi. N of Escuela Barrigón, 580-740 m, 8°38'N, 80°36'W, *Croat 67524* (AAU, COL, MO, US); El Valle region, vic. of La Mesa, N of El Valle de Antón, along E edge of Cerro Gaital, 900-1000 m, 8°37'N, 80°08'W, *Croat 67225* (COL, MO, US); La Mesa, above El Valle, *Croat 13467* (MO, SCZ); 775 m, 8°36'N, 80°07'W, 74893 (AAU, CAS, MO, PMA); 800 m, *Croat & Zhu 76676* (MO). Colón: Río Guaniche, ca. 1 km upstream from bridge at the Colón-Portobelo Road, 100 m, 9°30'N, 79°39'W, *Croat 75191* (CM, MO, TEX); Río Iguanita, ca. 3 km above the bridge on Portobelo road, <100 m, 9°27'N, 79°40'W, 49760 (CM, MO). Panamá: El Llano-Cartí, 10 mi. from Pan-American Highway, 350 m, 9°14'N, 79°W, *Knapp et al. 4727* (MO); 450 m, 9°14'N, 79°W, *Knapp & Huff 4431* (MO); 7.5 km N of highway, *Folsom 2555* (MO); 5-6 mi. N of highway, 350-375 m, *Croat 34762* (MO); ca. 16-18 km from highway, 400 m, *Fyson & Nee 7344* (F, MO), 7357 (MO, US); 4 mi. from highway, *Croat 33731* (MO, NY); 3-3.5 mi. NE of Altos de Pacora, 700-750 m, 9°15'N, 79°25'W, 68696 (CM, KYO, MO, PMA, RSA); Cerro Jefe region, 750-800 m, 9°14'N, 79°22'W, 67088 (MO); 850 m, 67054 (MO); ca. 0.5 mi. below tower, 700-800 m, 9°15'N, 79°30'W, *Thompson 4789* (MO); 800 m, *Sullivan 211* (MO); 21 km above Pan-American Highway, ca. 600 m, *Croat 35889* (MO); 4.6 km beyond peak on road to Altos de Pacora, 26.3 km from the Inter-American Highway, 600 m, 35895 (MO); 0.8 mi. beyond turnoff to Altos de Pacora, 970 m, 8°43'N, 82°17'W, *Croat & Zhu 76620* (MO); Río Teralde, on road 8 km from Pan-American Hwy., 400 m, 9°08'N, 79°W, *Knapp 1825* (MO). San Blas: El Llano-Cartí Road, 14 mi. N of Pan-American Highway, 300 m, 9°15'N, 79°W, *Croat 69247* (CM, DUKE, MO, TEX), 69250 (CM, MO, NY); 14.5 mi. N of highway, 350 m, 9°15'N, 79°W, *McPherson 9508* (MO, PMA); 10.1 mi. N of highway, 300 m, *Croat & Zhu 76540* (MO); 1 mi. S of Nusagandi, 9 mi. N of Interamerican Highway, 350 m, 9°20'N, 79°W, *Croat & Zhu 77013* (MO); 76568 (MO, US); 76569A (MO); 10.1 mi. N of main highway, 300 m, 9°20'N, 79°W, *Croat & Zhu 76540* (MO); near Nusagandi, along Sendero Nusagandi, 300-350 m, 9°15'N, 79°W, *McPherson 10756* (MO); 1-5 mi. N of Nusagandi, 250-300 m, 9°16'N, 79°W, *Thompson 4661* (CM). Veraguas: Santa Fe region, Santa Fe-Río San Luis, past Escuela Agrícola

Alto de Piedra, 8 mi. N of school, 450 m, 8°33'N, 81°08'W, Croat 66954 (K, MO, NY, PMA, QCA); 0.2 mi. beyond fork in road at Escuela Agrícola Alto Piedra, 750 m, 33971 (MO).

***Philodendron antonioanum*** Croat, sp. nov.

TYPE: Panama. Veraguas: vic. of Santa Fe, along road between Alto Piedra and Calvébora, 0.5 mi. N of Alto Piedra, on slopes of Cerro Tute, Parque Nacional Cerro Tute, 1030 m, 15 July 1994, Croat & Zhu 76909 (holotype, MO-4619868-9, MO-4619871-2; isotypes, B, CAS, COL, CR, F, GH, K, MEXU, MO, NY, PMA, SCZ, US, VEN). Figures 61-64.

Planta hemiepiphytica; internodia brevia, 4-7 cm diam.; cataphylla rubra, acute 2-costata, persistentia ut fibrae tenuis; petiolus 28-91 cm longus, obtuse complanatus adaxialiter; lamina ovata, 26-86 cm longa, 18-57 cm lata, nervis primariis lateralibus 5-6 utroque; inflorescentia 2; pedunculus 6-7 cm longus; spathe 15-16 cm longa; lamina spatheae extus viridialba, intus alba suffusa rubella in dimidio inferiore; tubo spatheae in superficialibus ambabus marronino; pistilla 5-locularia; loculi 25-30-ovulati.

Hemiepiphytic; larger stems drying reddish brown, smooth, and coarsely striate at nodes, smaller stems drying light yellow-brown, glossy, and finely longitudinally ribbed, finally dark brown and transversely finely cracked; internodes short, 4-7 cm diam., broader than long; roots to 30 cm long, numerous per node, drying reddish brown, weakly glossy, and somewhat ridged; cataphylls 35-39 cm long, sharply 2-ribbed, red, broadly sulcate abaxially, persisting as pale brown fibers densely clothing the entire apex of the stem with tiny fragments of reddish brown epidermis persisting among the fibers; petioles 28-91 cm long, 1.5-2 cm diam., subterete, obtusely flattened adaxially, weakly 1-ribbed near apex, broadly and obtusely sulcate, to 3.5 cm diam. near base, dark green (reddish brown when young), drying reddish brown and matte, surface moderately glossy, prominently striate-lineate throughout; blades broadly ovate, moderately coriaceous, bicolorous, semiglossy, 26-86 cm long, 18-57 cm wide (1.2-1.5 times longer than wide), margins broadly and sparsely undulate, upper surface drying dark brown to gray-brown, lower surface drying reddish brown; sinus hippocrepiform to obovate, 4-18 cm deep; midrib flat and slightly paler than surface, drying convex and darker than surface above, convex, sometimes tinged reddish, drying obtusely angular, grayish brown, and sparsely short pale-lineate below; basal veins 7 per side, with the first and second free to base, the third through sixth veins coalesced to 7 cm, posterior rib barely naked near the base, weakly curved; primary

lateral veins 5-6 per side, departing midrib at a 40-50° angle, drying reddish brown below, obtusely and deeply sunken, concolorous, drying raised and somewhat darker above, convex, ± concolorous, drying concolorous, matte below; minor veins moderately distinct, drying prominulous above and below, arising from both the midrib and primary lateral veins, secretory canals appearing but not obvious below; "cross-veins" numerous, ± transverse. INFLORESCENCES 2 per axil; peduncle 6-7 cm long, 1.5 cm diam., magenta, conspicuously white-striate toward the apex; spathe 15-16 cm long, weakly constricted midway, weakly glossy, moderately coriaceous, drying reddish brown on both surfaces, acuminate at apex; spathe blade greenish white outside, white, tinged reddish in lower half inside; spathe tube inflated, dark maroon outside, to 7 cm long, 5.8 cm diam., dark maroon inside; spadix 15 cm long; pistillate portion 3.5-5 cm long in front, 3.3-3.7 cm long in back, 2-2.5 cm diam. at apex, 2.3-3 cm wide at base; staminate portion 9-11.7 cm long; fertile staminate portion gradually tapered toward apex; sterile staminate portion 2.5-2.6 cm diam.; pistils 3.2-3.5 mm long, creamy brown, ovary 5-locular, ca. 0.4 mm long, with parietal placentation; ovules 25-30 per locule, 2-3-seriate; funicle ca. 0.2 mm long, style 1 mm diam., similar to style type B; style apex flat, with small depressions; stigma 1.3-1.5 mm diam., 0.4 mm high, covering entire style apex; the androecium margins sharply 4-6-sided, 0.6-1.2 mm diam. at apex; sterile staminate flowers rounded to bluntly 4-6-sided, larger toward the base, 0.6-2.8 mm wide.

Flowering in *Philodendron antonioanum* probably takes place in the early rainy season. Two fertile collections have been seen, one with immature fruits in June and one post-anthesis in July.

*Philodendron antonioanum* is endemic to central Panama and known only from the type locality at Cerro Tute and at El Copé at 800 to 1200 m elevation in *Tropical Lower Montane rain forest*. *Philodendron antonioanum* is a member of *P. sect. Philodendron* subsect. *Philodendron ser. Fibrosa*. The species is distinguished by its red, sharply two-ribbed cataphylls, which persist as fibers; its subterete petioles longer than the blades; broadly ovate reddish brown drying blades with 5-6 primary lateral veins; and by the stout inflorescences with scarcely constricted spathes having the tube magenta on both surfaces.

*Philodendron antonioanum* is most easily confused with *Philodendron tysonii*, a species with which it may occur. Both species occur at higher

elevations, and have subterete petioles longer than blades, large, ovate, glossy blades, and red or reddish cataphylls persisting as fibers. *Philodendron tysonii* differs in having mostly unribbed cataphylls and blades with usually 6–10 primary lateral veins and with the upper surfaces drying usually more or less blackened and smooth, in contrast to blades with 4–6 primary lateral veins and with upper surfaces drying gray-green with prominulous cross-veins as in *P. antonioanum*. In addition, *P. tysonii* has more prominently pedunculate inflorescences with green spathes, which are merely tinged red on the tube outside and are proportionately more slender and more constricted above the tube rather than being dark magenta on the tube and barely constricted above the tube as in *P. antonioanum*. Finally, the species differ in the nature of ovules, with *P. tysonii* having ovaries (5)6–8(9)-locular with sub-basal placentation and (4)5–7 ovules per locule borne in an envelope, while *P. antonioanum* has ovaries 5-locular with parietal placentation and about 30 ovules per locule not borne within an envelope.

The species is named in honor of Thomas Antonio of the Chicago Botanic Garden, who was resident botanist in Panama for the Missouri Botanical Garden and who first collected the species in June 1980.

*Additional specimens examined.* PANAMA. Coelá: Alto Calvario above El Copé, ca. 6 km N of El Copé, 710–800 m, 8°39'N, 80°36'W, Croat 687664 (MO); 930 m, 49197 (CM, M, MO); along Continental Divide, 900–1000 m, 8°39'N, 80°36'W, 75051 (MO). Veraguas: vic. of Santa Fe, along road between Alto Piedra and Calovebona, 0.5 mi. N of Alto Piedra, on slopes of Cerro Tate, 1150 m, Antonio 4945 (MO, PMA).

***Philodendron aromaticum*** Croat & Grayum, sp. nov. TYPE: Costa Rica. Limón: 3.5 airline km S of Islas Buena Vista in the Río Colorado, 16 airline km SW of Barra del Colorado, 10–120 m, 10°39'N, 83°40'40"W, 15–16 Sep. 1986, *Davidse & Herrera 31212* (holotype, MO-3582209–10; isotype, US). Figures 66–68.

Planta plerumque epiphytica; internodia 2–3 cm longa, 1.5–3.5 cm diam.; cataphylla acute 2-costata, decidua; petioli subteretes, 33.5–66 cm longus, subspongiosus; lamina ovato-cordata, 33–55 cm longa, 17–26 cm lata, in sicco canoviridis saepe, viridis vel brunneiviridis infra; inflorescentia 1; pedunculus 4.5–9.4 cm longus; spathe 15–20 cm longa, lamina spathe extus atriviridis, intus pallide marronina vel alba; tubo spathe atrimarronino; pistilla 9–10(11)-locularia; loculi 2–3-ovulati.

Usually epiphytic, sometimes hemiepiphytic (often occurring high in canopy), rarely terrestrial (accidentally); stem usually appressed-climbing,

creeping, brownish, sap sweet, turpentine- to burseraceous-scented; internodes weakly glossy, minutely striate longitudinally, 2–3 cm long, 1.5–3.5 cm diam., usually broader than long, dark to medium green, drying brown, epidermis fissured longitudinally; roots moderately few, drying reddish brown; cataphylls sharply 2-ribbed, D-shaped, deciduous, margins sharply raised. LEAVES erect-spreading to spreading; petioles 33.5–66 cm long, ± terete, somewhat spongy, dark green, obtusely to weakly flattened adaxially, surface weakly glossy, not noticeably marked, drying 8–15 mm diam., usually greenish dark brown; blades ovate cordate, subcoriaceous, glossy, weakly bicolorous, acuminate to long-acuminate at apex (the acumen inrolled, 7–11 mm long), cordate at base, 35–55 cm long, 17–26 cm wide (2.1 times longer than wide), (0.8–1 times longer than petiole), margins weakly undulate; upper surface dark green, semiglossy, drying gray-green, lower surface glossy to semiglossy, paler, drying green to brownish green; anterior lobe 30.5–41 cm long, 16–26 cm wide (2.6–2.9 times longer than posterior lobes); posterior lobes 10.5–15.8 cm long, 7–13 cm wide, rounded at apex; sinus spatulate, 12 cm deep; midrib flattened, slightly paler than surface above, raised to convex, paler than surface below; basal veins 5–6 per side, with none or rarely 1 free to base, two to three veins coalesced 2.7–5.5(6.3) cm; posterior rib naked for 4 cm; primary lateral veins 5–7 per side, departing midrib at a 70–80° angle, straight to the margins, weakly and narrowly sunken above, convex and paler than surface below; minor veins moderately obscure to clearly visible below, arising from both the midrib and primary lateral veins. INFLORESCENCES 1 per axil; peduncle 4.5–9.4 cm long, 0.08–0.15 cm diam., green, weakly striate; spathe semiglossy, 15–20 cm long (2.1–3.3 times longer than peduncle), constricted above the tube, dark green throughout, pale maroon to white inside; spathe tube oblong-ellipsoid, weakly wrinkled longitudinally outside, 6–9 cm long, dark maroon, suffused onto blade inside; spadix sessile; greenish white to white throughout, oblong, 12–16.9 cm long; pistillate portion creamy white to pale green, cylindrical, 5–13 cm long, 6–12 mm diam. at apex, 12 mm diam. at middle, 5–10 mm wide at base; staminate portion 7–11 cm long; fertile staminate portion white, ellipsoid, 1 cm diam. at base, 1.4 cm diam. at middle, 6 mm diam. ca. 1 cm from apex, broadest at the middle, slightly broader than the pistillate portion, narrower than the sterile portion; sterile staminate portion broader than the pistillate portion, white, 1.5 cm diam.; pistils 3 mm long, 1.1–1.3 mm diam., ovary 9–10(11)-

locular, 1.8 mm long, 1.2 mm diam., with sub-basal placentation; ovules 2-3 per locule, contained within translucent or transparent ovule sac, 0.3 mm long,  $\pm$  equal in length to funicle, style similar to style type B; style crown about as broad as ovary; central style dome well developed; style apex with central domes; stigma discoid, weakly lobed, 1.1 mm diam., 0.2-0.3 mm high, covering entire style apex, depressed medially; the androecium margins mostly 5-sided, sometimes 4-6-sided; thecae oblong to ovate, 0.4 mm wide.

Flowering phenology in *Philodendron aromaticum* is unclear since too few collections exist, although present collections might suggest bimodality. Post-anthesis collections have been made in September, October, and in February. Immature fruiting collections have also been made in the same months, and mature fruiting collections have been made in September.

*Philodendron aromaticum* is endemic to the Atlantic slope of northwestern Costa Rica from sea level to 360 m elevation in a *Tropical wet forest* life zone.

*Philodendron aromaticum* is a member of *P.* sect. *Calostigma* subsect. *Macrobolium* ser. *Macrobolium*. The species is characterized by its thick, bare, short internodes; sharply two-ribbed, deciduous cataphylls; moderately spongy, obtusely flattened petioles longer than the blades; ovate-cordate blades drying gray-green on the upper surface and green to brownish green below; and by its solitary inflorescence with a short peduncle and green spathe, which is maroon on the tube within. Also characteristic is the macilaginous, aromatic sap of the petioles (hence the name).

*Philodendron aromaticum* is most easily confused with *P. sagittifolium*, but that species typically has more elongate blades, which dry usually reddish brown; basal veins free or at least not forming a posterior rib that is naked at the sinus; and often more than a single inflorescence per axil. In contrast, *P. aromaticum* has blades with a prominent posterior rib naked along much of the sinus.

*Additional specimens examined.* COSTA RICA. **Carriago:** Turrialba, Dress 6331 (BH; cultivated). **Heredia:** San José-Pto. Viejo, vic. of Chilamate, 11.6 mi. N of Carriago, 100 m, 10°27'N, 84°05'W, Croat 68382 (CAS, CR, F, G, L, M, MO); La Selva Field Station, Puerto Viejo de Sarapiquí, along El Sará Trail, 100-150 m, 44315 (MO); ca. 100 m, Folsom 8963 (DUKE); 9438 (DUKE); Grayum 2686 (DUKE); Hammel 10733 (DUKE); Croat 61214 (MO). **Limón:** 2-3.5 airline km SSE of Isla Buena Vista in the Río Colorado, 14 airline km SW of Barra del Colorado, 10-120 m, 10°40'N, 83°40'W, Davidse & Herrera 31136 (CR, MO, US); Cerro Corozal, 10-80 m, 83°39'30"W, 10°40'30"N, 31481 (CR, MO); 20-170 m,

10°41'N, 83°38'W, Stevens 23764 (MO); 2 km W of Río Toro Amarillo, on road W from Guápiles, 275 m, 10°13'N, 83°50'W, Thompson & Raudins 1197 (CM); Río Sierpe, 5 km NE of La Aurora, Guápiles, 30 m, 10°22'N, 83°31'W, Robles 2189 (CR); 2238 (CR, MO); Río Frio-Limón, vic. of Río Blanco, W of Guápiles, 360 m, 10°12'N, 83°49'W, Croat 68423 (B, CR, K, MO); Refugio Nacional de Vida Silvestre, Río Chirripocito-Río Sardana, 12 m, 10°38'N, 83°45'W, Grayum 9778 (CR, MO).

***Philodendron auriculatum*** Standl. & L. O. Williams, Ceiba 3: 108. 1952. TYPE: Costa Rica. Puntarenas: Esquinas forest preserve, Esquinas Experiment Station, region between Río Esquinas and Palmar Sur de Osa, 60 m, Allen 5697 (holotype, EAP; isotypes, F, US). Figures 12, 20, 23, 65, 69-72.

Usually epiphytic, sometimes epilithic, rarely terrestrial; stem appressed-climbing, creeping, leaf scars conspicuous, 2.5-3 cm long, 2.5-3.5 cm wide; internodes short, scurfy, 1-4 cm long, 1.5-5.5 cm diam., usually broader than long, dark green, drying becoming light tan with age; roots purplish maroon at base, brownish green or yellow at apex, pubescent, tapered, slender, to 27 cm long, 3-4 mm diam., few per node, sometimes emerging through the cataphylls at the nodes; cataphylls subcoriaceous, sometimes moderately spongy, 13-50 cm long, sharply 2-ribbed, pale green, drying brown, several persisting intact, eventually deciduous, emarginate with subapical apiculum at apex. LEAVES mostly erect or erect-spreading; petioles 24-53 cm long, 7-23 mm diam., erect to spreading, subterete,  $\pm$  spongy, green, somewhat flattened at base adaxially, rounded abaxially, surface short, dark, green-lineate and with dark green ring at apex; blades usually oblong-oblancoolate, rarely oblong-elliptic, broadest from middle to above middle, with narrow auriculate lobes, subcoriaceous, bicolorous, acuminate at apex (the acumen inrolled), usually narrowly cordulate or auriculate at base, 37.5-75 cm long, 9-23 cm wide (3.3-4.2 times longer than wide), (1.4-1.6 times longer than petiole), broadest at the middle, upper surface dark green, drying pale yellow-green, semiglossy, lower surface slightly paler green, semiglossy; sinus 7-14 mm deep; midrib broadly convex at base becoming flattened toward the middle, eventually weakly sunken at apex above, green, with dark green lineations, midrib convex at base, eventually narrowly convex at apex below, green-striate, paler than surface; primary lateral veins 7-10(11) per side, departing midrib at a 80-85° angle,  $\pm$  straight to the margins, sunken, concolorous above, raised to round-raised, paler than surface below; interprimary veins numerous, usually as conspicuous as



primary lateral veins; minor veins arising mostly from the midrib, fewer from the primary lateral veins; lesser veins obscure above, fine, moderately conspicuous and darker than surface below. INFLORESCENCES  $\pm$  erect, 1–2 per axil; peduncle 7–11.3 cm long, 9–11 mm diam., medium to pale green, unmarked, semiglossy; spathe erect-spreading, subcoriaceous, 13–14.5 cm long (1.2–1.8 times longer than peduncle), weakly constricted midway (6 cm above base); spathe blade broadly flattened, curved weakly forward, pale greenish outside, to 9 cm long (opening elliptic in face view, 5.5 cm wide), pale green, heavily suffused with red (B & K Red-Purple 3/10) throughout inside; spathe tube oblong-ellipsoid, medium to dark green outside, densely short lineate throughout outside, 6 cm long, 3.5 cm diam., red to maroon (B & K Red-Purple 3/10) at base, weakly so toward apex inside; spadix oblong (weakly tapered or weakly clavate), 11–24 cm long, broadest usually at the middle; pistillate portion medium to dark green, tapered toward the apex, 2.3–4.2 cm long, 2.7–4.3 cm long in front, 2.4–2.7 cm long in back, 8–12 mm diam. at apex, 9–15 mm diam. at middle, 5–7 mm wide at base; staminate portion 9.5–12 cm long; fertile staminate portion weakly tapered or weakly ellipsoid, 8–16 mm diam. at base, 10–16 mm diam. at middle, 6–10 mm diam. ca. 1 cm from apex, broadest at the middle or  $\pm$  uniform, broader than or as broad as the pistillate portion, as broad as the sterile portion; sterile staminate portion broader than or as broad as the pistillate portion; sterile staminate portion 0.7–1.5 cm diam.; pistils to 2.6 mm long, to 1.6 mm diam.; ovary 5–7(8–9)-locular, 0.7–1.2(2.1) mm long, (0.7)1.1–1.3 mm diam., with sub-basal placentation; ovules (3)4 per locule, 1-seriate (or in 2 series of 2), contained within translucent ovule sac, if present, 0.2–0.4 mm long, usually longer than funicle, style similar to style type B; style crown usually as broad as ovary; stigma discoid, 0.7–1 mm diam., 0.1–0.3 mm high, covering entire style apex; the androecium margins 4–6-sided and scalloped, 1–2.3 mm diam. at apex; thecae oblong (elliptical, slightly obovate); sterile staminate flowers irregularly rounded to bluntly 5-sided, 2.3–3 mm long, 1–1.5 mm wide, white. INFRUCTESCENCE with berries orange.

Flowering phenology in *Philodendron auriculatum* is unclear, but possibly flowering is initiated in the late rainy season. Only a single flowering collection exists (October), but there are a modest number of collections collected post-anthesis during the dry season and early rainy season (January through June). A single mature fruiting collection was made in January.

*Philodendron auriculatum* occurs only in southwestern Costa Rica on the Pacific slope from San José and Puntarenas Provinces (as far west as Carrara), ranging from near sea level to 1200 m elevation in *Tropical wet forest* life zones.

*Philodendron auriculatum* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is characterized by short internodes, sharply two-ribbed, deciduous cataphylls, moderately long, markedly spongy, somewhat flattened petioles (averaging slightly shorter than the blades), oblong-elliptic to oblong-oblancoate, pale yellow-green-drying blades, which are usually narrowly cordulate at base, and one to two greenish inflorescences, which are red to maroon within at base.

*Philodendron auriculatum* is probably most closely related to *P. pseudauriculatum* Croft, which ranges along the Atlantic slope of Central America from Nicaragua to Panama and also on the Pacific slope of Panama near the Continental Divide (as well as the Serranía de Cañasas and Serranía de Pirre). That species differs in its usually darker gray-green-drying leaf blades with a dark green ring at the apex of the petiole, and in having the leaf base acute, rounded, or broadly subcordate at base (lacking the narrow auriculate lobes so common for *P. auriculatum*). In addition, *P. pseudauriculatum* has the whitish spathe clearly demarcated from the contrasting green peduncle. In contrast, *P. auriculatum* has a yellowish green spathe that is not at all demarcated from the peduncle.

*Additional specimens examined.* COSTA RICA. Puntarenas: near Inter-American Highway in vic. of Piedras Blancas, Croft 32956 (CR, MO); hills N of Palmar Norte, along trail to Jalisco, 50–700 m, Croft 35205 (MO); cove at NE base of peninsula, 3 km S of Puerto Quepos, 0 m, 9°24'N, 84°10'W, Grayson & Sleeper 6612 (CR, K, MO, US); El General Valley, along Río Sonador, near Pan-American Highway, 600 m, Williams et al. 28908 (F); along the Río Cacao above Pan-American Highway, 900 m, Williams et al. 28687 (F); Las Cruces Tropical Botanical Garden, Cantón Coto Brus, 6 km W of San Vito de Java, 1200 m, 8°49'N, 82°58'W, Croft 57268 (MO); 8.8 mi. N of Villa Neily, 1010 m, 66171 (MO); Osa Peninsula, vic. Boscosa, Croft & Hannon 79297 (INB, MO); Esquinas Ridge, 150–250 m, Gómez 19677 (MO, NY, RSA, US); Rincón de Osa, Liesner 1819 (F, MO); ca. 5 km W of Rincón de Osa, 50–200 m, 8°42'N, 83°31'W, Burger & Liesner 7298 (F, MO, PMA); 7306 (CR, F); 50–200 m, 8°42'N, 83°31'W, Burger & Gentry 8978 (CR, F); 8°42'N, 83°31'W, Aguilar 1507 (INB, MO); Piedras Blancas–Rincón, 3.7 mi. W of Pan-American Highway, 90–105 m, 8°46'N, 83°18'W, Croft 67691 (MO); Quebrada Aparicio–Quebrada Aguabuena, Rincón de Osa, 200–400 m, 8°42'N, 83°31'W, Grayson et al. 4013 (MO); Chacarita–Rincón de Osa, ca. 6 km W of Inter-American Highway at Chacarita, 100 m, 8°45'N, 83°18'W, Croft & Grayson 39730 (CM, CR, K, MO, SAR); Parque Nacional Corco-

vado, Sirena, 0-150 m, 8°27'-30'N, 83°33'-38'W, Kernan 25 (CR, MO); Estación Sirena, S of Río Sirena along Río Camaronel, 0 m, 8°28'N, 83°35'W, Knapp 2189 (CR, L, MO, US); Talamancas Range, Pacific slope, forested foothills of mountains E of Quepos, 150-250 m, 9°29'N, 84°03'W, Burger et al. 10603 (F, MO). **San José:** Puriscal, Z.P. La Cangreja, 300 m, Morales et al. 3239 (CR, INB); Carara-El Sur de Turubares, 280-370 m, 9°45'30"N, 84°32'W, Grayum 10445 (CR, INB, MO).

**Philodendron bakeri** Croat & Grayum, sp. nov.

**TYPE:** Costa Rica. Guanacaste: W slope of Cerro Nubes, ca. 2 km E of Silencio de Tilarán, large patch of remnant primary forest, 900 m, 10°28'N, 84°53'W, Grayum, Herrera & Sleeper 4992 (holotype, MO-3392250; isotypes, CR, DUKE). Figures 77, 78.

Planta hemiepiphytica; internodia pleurumque longiora quam lata, (2)7-9 cm longa, (0.7)1-1.5 cm lata; cataphylla 5-10 cm longa, acuta 1-costata aut raro vel acuta 2-costata, decidua; petiolus subteres, 4-12 cm longus, 2-4(6) mm diam.; lamina plus minusve oblonga aut anguste oblonga vel oblanceolata, basi cuneata vel rotundata, 12-25.5 cm longus; inflorescentia 1; pedunculus 2.6-5.2 cm longus, 2-5 mm diam.; spathe 7-11.5 cm longa; lamina spathe extus pallide viridi vel crema suffusa rubra vel marronina, intus viridi vel albidula suffusa rubra; tubo spathe intus atrimarronino, rubro aut carmesino; pistilla 5-8(10)-locularia; loculi 1(2)-ovulati; baccae aurantiaceae.

Hemiepiphytic; stem scandent, slender, to 1.5 m long; internodes (2)7-9 cm long, (0.7)1-1.5 cm diam., usually longer than broad, medium green, ± matte, becoming light brown, epidermis fissured minutely longitudinally upon drying; roots dark brown, thin; cataphylls subcoriaceous, 5-10 cm long, unribbed, bluntly to sharply 1-ribbed or rarely sharply 2-ribbed, green to reddish or maroon, deciduous, narrowly rounded at apex, margins clear. **LEAVES** spreading; **petioles** 4-12 cm long, 2-4(6) mm diam., spreading, subterete to C-shaped, sometimes tinged reddish, sometimes maroon at base, obtusely somewhat flattened or bluntly sulcate adaxially, rounded abaxially, surface semiglossy, with narrow purple ring at apex; sheathing 2.7-3.2 cm long, sheathing to ca. 2 cm but for ½ to ⅓ of its petiole length when subtending inflorescences; geniculum subterete, maroon, 2.7-4 cm long, darker than petiole; **blades** ± oblong or narrowly oblong to oblanceolate, moderately coriaceous, acuminate to long acuminate, sometimes acute at apex (the acumen sometimes short apiculate, 1-3 mm long), cuneate to rounded at base, 12-25.5 cm long, (2.7)4-10 cm wide (2.5-3 times longer than wide), (2-3 times longer than petiole), margins thin, narrow, reddish, upper surface drying reddish brown, semiglossy, lower surface weakly glossy; midrib flat to sunken, slightly paler than surface above, nar-

rowly convex, reddish violet below; basal veins lacking; posterior rib lacking; primary lateral veins 3-4 per side, departing midrib at a 45-55° angle, ascending to the apex, inconspicuous or slightly sunken above, not distinct below; tertiary veins visible, darker than surface; minor veins obscurely visible to moderately distinct, fine, close, arising from midrib only; secretory ducts usually obscurely visible on lower surface of dried blade. **INFLORESCENCES** 1 per axil; peduncle 2.6-5.2 cm long, 2-5 mm diam., obscured by petiole sheath; **spathe** moderately coriaceous, 7-11.5 cm long (2.2-2.7 times longer than peduncle), scarcely or not at all constricted, pale green to cream, often tinged lightly or heavily with reddish lineations, sometimes solid red or maroon, or with red speckles or lines (rarely seen in Panamanian specimens), green to whitish, suffused with red inside; resin canals appearing medially, especially near the constriction; spathe tube cylindrical, semiglossy outside, 4-5 cm long, deep maroon, red, or crimson inside; **spadix** sessile; cylindrical, sometimes clavate or weakly tapered, 7.2-8.5(9.5) cm long, ± uniform throughout; pistillate portion pale greenish, cylindrical, 1.8-2.4 cm long, 5.8-8 mm diam. at apex, 6-9 mm diam. at middle, 5-6 mm wide at base; staminate portion 4.8-8.9 cm long; fertile staminate portion white, cylindrical, clavate or weakly tapered, 6-11 mm diam. at base, 8-9 mm diam. at middle, 7 mm diam. ca. 1 cm from apex, as broad as or sometimes broader than the pistillate portion, if detectable, narrower than the sterile portion; sterile staminate portion broader than the pistillate portion, white, 6 mm diam.; pistils 1-2.2 mm long, 0.6-1.4 mm diam.; ovary 5-8(10)-locular, 0.4-1 mm long, 0.8-1 mm diam., with sub-basal placentation; ovules 1(2) per locule, sometimes contained within transparent, gelatinous ovule sac, 0.2-0.5 mm long, equal in length to or longer than funicle; funicle 0.2 mm long, style 0.7 mm diam., similar to style type B; style apex with depressions surrounding stylar canal exits; stigma brush-like, discoid, 0.5(1.5) mm diam., 0.2-0.6 mm high, covering entire style apex; the androecium truncate, margins irregularly 4-5-sided, 1.3 mm long, 0.3-1.4 mm diam. at apex; thecae oblong, sometimes oblong-elliptical; pollen spheroidal to ellipsoidal or elongate, less than 0.2 mm long, 0.1 mm diam.; sterile staminate flowers 3-6-sided or irregularly 4-sided, 0.9-3.9 mm long, 0.4-1.3 mm wide. **INFRUDESCENCE** pendent; seeds 1(2) per locule. **JUVENILE** leaves and spathes tinged red.

Flowering in *Philodendron bakeri* occurs from the early dry season, December through April, with post-anthesis collections from January through May.

Immature fruiting collections were made in May and June.

*Philodendron bakeri* is known along the Atlantic slope of Costa Rica from 100 to 900 m elevation in *Premontane wet forest* and *Tropical wet forest* life zones and also occurs in Central Panama at 900 to 1420 m in *Tropical wet forest* and *Premontane rain forest* life zones. It is expected to be found in intervening areas.

*Philodendron bakeri* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. The species is characterized by its scandent habit, slender stems (frequently pendulous on flowering portions), reddish, deciduous cataphylls, subterete petioles (one-fourth to one-third as long as the blades), more or less oblong blades, and by the solitary inflorescence with the spathe green, suffused with red or maroon outside, and heavily suffused with red within, especially at the base. Also characteristic are the bright orange fruits.

*Philodendron bakeri* is most similar to *P. immixtum*, also a vine with more or less elongate blades. That species differs in having thicker internodes drying with broad fissures and an often loose, flaking epidermis, and thinner blades that dry greenish or greenish brown and are more typically subcordate at the base. In contrast, *P. bakeri* has minutely fissured stems that lack a loose, flaking epidermis and blades that dry typically reddish brown, moderately coriaceous, and are merely rounded to obtuse at the base.

Though both species have spathe tubes that are reddish on the inner surface, *P. immixtum* differs in that the tube is not reddish on the outside.

Both *P. bakeri* and *P. immixtum* are members of *P.* sect. *Calostigma* subsect. *Glossophyllum*, with one ovule per locule, but locules of *P. immixtum* differ in being longer, with the transparent envelope enclosing the ovules being about twice as long as the ovules, whereas those of *P. bakeri* have the ovule(s) embedded in a gelatinous matrix completely filling the envelope.

*Philodendron bakeri* may also be confused with precociously flowering shoots of *P. sagittifolium*. Both species have petioles that may be tinged purplish violet on both ends. *Philodendron sagittifolium* differs in having pistils with 7–8 locules with 2–4 ovules per locule in a transparent envelope (vs. 1 ovule per locule in a gelatinous matrix).

Croat 44312 from La Selva, Costa Rica, differs in drying dark gray-brown above, dark brown below, and in lacking secretory ducts. It is perhaps a juvenile of some other species.

The species is named in honor of Richard Baker,

an aroid specialist previously at the Field Museum, who made the first collection in 1974.

*Additional specimens examined.* COSTA RICA. **Alajuela:** Cariblanco-Laguna Hule, ca. 2 km W of Costa Rica Highway 9, 10°17'N, 84°13'W, Baker et al. 227 (F, MO); Monteverde Biological Reserve, Río Peñas Blancas, 900 m, 10°19'N, 84°44'W, Haber & Bello 6932 (MO), 2907 (CR); Volcán Miravalles, W of Bijagua, near Río Zapote, ca. 600 m, 10°44'N, 85°05'W, Burger et al. 11628 (F, MO); Cañas-Upala, 4 km NNE of Bijagua, on slopes leading into Río Zapote, ca. 400 m, Croat 36263 (MO); Upala, 2 km NE of Colonia Libertad along Río Caño Negro, 300 m, 10°50'N, 85°16'W, Herrera 1958 (MO); San Ramón, 800 m, Bello 1973 (CR); Bello et al. 4537 (CR, INB); Reserva Forestal de San Ramón, 1000 m, 10°12'40"N, 84°36'20"W, Herrera 6748 (CR, MO). **Guanacaste:** N side of Lake Arenal, 1 km NW of dam, 650 m, 10°30'N, 84°46'W, Haber et al. 4915 (CR, MO); 4916 (MO). **Heredia:** ca. 7 km SW of Las Horquetas, ca. 400 m, 10°18'N, 84°01'W, Grayum 5018 (MO); ca. 8 km SW of Las Horquetas, 450–550 m, 10°18'N, 84°02'W, Grayum et al. 6543 (MO); Río Peje–Río Sardinalito, Volcán Barva, 700–750 m, 10°17'30"N, 84°04'30"W, Grayum & Jeremy 6784 (MO); La Zona Protectora, Río Peje–Río Guacimo, northern slopes of Volcán Barva, along trail from main road across Quebrada Cantarana to Río Guacimo, 250 m, Grayum & Schatz 3219 (DUKE); La Selva Field Station, 100 m, Hammel 7805 (DUKE, MO); 100–150 m, Croat 44312 (MO); 6 km by road from Río Peje crossing, 5 km SSE of Magsasay, 10°21'N, 84°03'–04'W, Schatz & Grayum 667 (DUKE); Magsasay, 700 m, I. Chacón 182 (MO). **Limón:** Hacienda Tapozo–Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, Davidson et al. 6982 (LAM); Tortuguero National Park, 0 m, 10°34'N, 83°31'W, Croat 61210 (MO); 2 km W of Río Toro Amarillo on road W from Guápiles, 275 m, 10°13'N, 83°50'W, Thompson & Rawlins 1225 (CM). **PANAMA.** **Chiriquí:** Cerro Colorado, 19.7 mi. N of Río San Félix, 1420 m, 8°31'N, 81°46'W, Croat 74997 (MO). **Coclé:** Penonomé, Llano Grande–Coclequito, 4.3 mi. N of Llano Grande, 410 m, 8°42'N, 80°26'W, Croat 67456 (CM, L, MO, PMA); El Copé region, Alto Calvario, 5.2 mi. above El Copé, 930 m, 49159 (MO); above Alto Calvario, 1200–1300 m, Szymka & Anderson 4546 (MO); El Valle region, La Mesa, above El Valle de Antón, 860–900 m, Croat 37419 (MO), 37508 (MO). **Panamá:** El Llano–Cartí Road, Km 19.1, 350 m, 9°19'N, 78°55'W, de Nevers & Herrera 5875 (MO, PMA); El Llano–Cartí, Nusagandi, 325–350 m, Croat 67395 (MO); Km 10, 33710 (MO); 0.5 mi. E of El Llano, D'Arcy 5202 (MO).

**Philodendron basii** Matuda, *Anales Inst. Biol. Univ. Nac. Méxic.* 32: 153, Fig. 8. 1961. TYPE: Mexico. México: Barranca de Malinaltenango, NE (published as SE) of Zacualpan, 1200 m, 18°43'N, 99°36'W, Matuda 37244 (holotype, MEXU). Figures 76, 81–84.

Almost always terrestrial, or trailing over rocks; stem green, stout, succulent, leaf scars conspicuous, 2–4 cm long, 4–5 cm wide; internodes 1–4 cm long, 2.5–7 cm diam., broader than long, drying olive-green, epidermis peeling, light brownish,

roots long, 5 mm diam., light reddish tan; cataphylls 20–28 cm long, weakly to sharply 2-ribbed, medium green with darker ribs and dark green speckles in lower one-half of cataphyll, drying tannish, persisting intact, eventually deciduous; **petioles** 22–70 cm long, 8–15 mm diam., terete, medium green, surface dark striate; **blades** ovate-cordate, drying moderately thin, weakly bicolorous, semiglossy, acute at apex, cordate at base, 16–56 cm long, 9–38 cm wide (1.5–1.8 times longer than wide), (0.7–0.9 times the petiole length), broadest at or above middle; upper surface medium green, weakly glossy, drying medium yellow-brown; lower surface slightly paler, semiglossy, drying yellow-green to yellow-brown; margins sinuate or weakly undulate; anterior lobe 35–48 cm long, 37.5–46 cm wide (2.5–8.5 times longer than posterior lobes); posterior lobes 4–19 cm long, 18–23 cm wide, directed inward, rounded to obtuse at apex; sinus spatulate, 10–13 cm deep; midrib weakly sunken to flat, heavily dark green striate, paler than surface above, convex, sparsely dark-lineate, paler than surface below; basal veins 3–6 per side, with 1 pair free to base, the third through fifth coalesced 3–3.5(4) cm; posterior rib naked for 1–2.5 cm; primary lateral veins 4–7 per side, departing midrib at a 40–60° angle, ± straight to the margins, sunken and paler than surface above, convex and paler than surface below; interprimary veins sunken and concolorous above, flat and darker than surface below; branches of primary lateral veins ± ruffled-sunken and concolorous above, raised and paler than surface below; minor veins predominately arising from the primary lateral veins, those arising from the midrib seem to disappear soon after leaving the midrib. **INFLORESCENCES** probably 1 per axil; peduncle 7–8 cm long, 7–8 mm diam., green; **spathe** 12.6–13.7 cm long (1.7–1.8 times longer than peduncle), apiculate at apex; spathe blade pale green or white, tinged red near base outside, 8 cm long, white tinged with red near base, pale-speckled inside; spathe tube ellipsoid, dark green outside, 6 cm long, 3 cm diam., maroon-red (B & K Red-Purple 10/3) inside; **spadix** sessile; tapered, 11.8–12 cm long, broadest at the base; pistillate portion cylindrical, 2.5–3.8 cm long, 1.5–1.9 cm diam. midway, slightly narrower on both ends; staminate portion 7.6–13 cm long; fertile staminate portion clavate, 9–15 mm diam. at base, 11–15 mm diam. at middle, 5–10 mm diam. ca. 1 cm from apex, broadest above the middle, narrower than the pistillate portion, broader than the sterile portion; sterile staminate portion narrower than the pistillate portion, 1–1.8 cm diam.; pistils 2.9–3.9(5.7) mm long,

1.4–2 mm diam.; ovary 4–6-locular, 2 mm long, 1.4 mm diam., with sub-basal placentation; ovules 4–6 per locule, 2–3-seriate, contained within transparent ovule sac, 0.3 mm long, equal in length to funicle, style similar to style type D; style apex rounded, with style boss and with depressions surrounding stylar canal exits; style boss narrow but pronounced; stigma slightly discoid to hemispheroid, 1.6–2.2 mm diam., 0.6–1.4 mm high, covering almost entire style apex; the androecium 4–6-sided; thecae oblong, 0.3 mm wide; sterile staminate flowers 4–6-sided, 1.7 mm long, 0.9–1.6 mm wide. **JUVENILE** leaves ± sagittate at base.

Flowering in *Philodendron basii* is recorded during the early wet season with post-anthesis collections made during September and in January, although too few collections overall have been seen.

*Philodendron basii* is endemic to Mexico, ranging from western Jalisco, south and east to the states of Colima, México, and Guerrero, at 350 to 1200 m elevation in “Bosque Pino-Encino” and “Selva Baja Caducifolia.” Matuda (1962) reported it from as low as 200 m elevation.

*Philodendron basii* is a member of *P.* sect. *Colastigma* subsect. *Macrobolium* ser. *Pachycaulia*. The species is characterized by its very stout, green, succulent stems, which usually trail over rocks; weakly two-ribbed cataphylls, which remain intact and are usually soon deciduous; long-petiole, moderately thin, ovate cordate blades with a sinuate margin; and by the green spathes with the tube reddish purple within.

Although *Philodendron basii* is not easily confused with any other species, it is ecologically very similar to *P. warszewiczii*, which also has thick, succulent, bare stems and occurs in very dry habitats. The latter is distinguished by its deeply dissected leaf blades. Matuda compared *P. basii* with *P. smithii*, to which it bears only a superficial resemblance. That species differs in having more slender internodes, which are usually longer than broad; more narrowly ovate blades; and (usually) two inflorescences per axil with longer peduncles.

Though no specimens have been seen from Colima State in Mexico, Matuda (1962) reported having seen this species there.

*Additional specimens examined.* MEXICO. **Guerrero:** km 337–338 beyond Acapulco on highway to Acapulco, 3000 ft., *Moore 5100* (BH, CM); road to Xalpatlahuac between bridge over Río Omilán and confluence of Río Papagayo with Río Omilán, 850 m, *Croat 45766* (MO). **Jalisco:** Highway 200, 10–13 km SE of El Tuito, valley of Río las Justas, 250–330 m, *McVaugh 25396* (MICH); ca. 6 mi. S of Tuito, 350 m, *Croat 45442* (COL, MEXU, MO, US).

***Philodendron breedlovei*** Croat, sp. nov. TYPE: Mexico. Chiapas: Mun. La Trinitaria, Monte Bello National Park, E of Lago Tz'ikaw, 13 May 1973, *Breedlove 35181* (holotype, DS; isotype, MEXU). Figures 79, 80, 85.

Planta hemiepiphytica; internodia usque plus quam, 7 cm, ca. 1.5 cm diam.; cataphylla decidua; petiolus subteres, 34 cm longus, 7 mm diam.; lamina anguste ovato-cordata, usque 37 cm longa, 20 cm lata, in sicco flavo-brunnea; inflorescentia 1; pedunculus 6 cm longus, 6 mm diam.; spathe 10.5 cm longa; lamina spathe viridialba vel alba; tubo spathe extus virenti, intus rubro; pistilla 5-6-locularia; loculi 20-ovulati.

Hemiepiphytic; internodes to 7 cm or more long, ca. 1.5 cm diam., usually longer than broad, drying light brown, semiglossy, weakly and irregularly ribbed; cataphylls deciduous; **petioles** 34 cm long (slightly longer than blades), 7 mm diam., subterete; **blades** narrowly ovate-cordate, acuminate and slightly inequilateral at apex, sagittate at base, 37 cm long, 20 cm wide (1.8 times longer than wide; about equal in length to petiole), semiglossy, upper surface drying dark olive-green, lower surface drying yellowish brown; anterior lobe 31.5 cm long, 21 cm wide (4 times longer than posterior lobes); posterior lobes 7.5 cm long, 8.7 cm wide, narrowly rounded, directed toward base; sinus  $\pm$  V-shaped, 5-6 cm deep; midrib drying narrowly raised and darker below; basal veins 3-4 per side, with 0 free to base, 1 pair coalesced for 2.8 cm; posterior rib to 3 cm long, not naked; primary lateral veins 5-6 per side, arising initially at an acute angle then forming a gradual arch to margin, departing midrib at a 55° angle toward apex, 60-70° angle midway, to 80° angle near the base, weakly arcuate to the margins, drying weakly raised and paler below; minor veins arising from both the midrib and primary lateral veins. INFLORESCENCES 1 per axil; peduncle 6 cm long, 6 mm diam.; **spathe** 10.5 cm long (1.75 times longer than peduncle), slightly constricted midway, densely speckled-lineate, especially on the tube, red within, darker red on the tube within, weakly acuminate at apex, convolute to half its length at base; spathe blade greenish white to white, paler along the margins outside; spathe tube medium green outside, 5 cm long; **spadix** sessile; 6.5-8 cm long, broadest above the middle; pistillate portion 2.9 cm long, 1.7-1.9 cm diam., broadest at the middle; staminate portion 7.2 cm long; fertile staminate portion to 2.2 cm diam. midway, usually broader than the pistillate portion; sterile staminate portion 1.7 cm diam.; pistils 3.3-3.8 mm long, 1.7-1.8 mm diam.; ovary 5-6-locular, with axile placentation; ovules 20 per locule, 0.3-0.4 mm long, equal in length or slightly

longer than funicle; style similar to style type B, 1.3-1.5 mm diam.; style apex truncate, minutely warty, irregularly 3-5-sided; stigma 2.3-2.5 mm diam., depressed with 5-6 small stylar canals; sterile staminate flowers 2.7-3.1 mm long, 0.9-1.6 mm wide.

Flowering in *Philodendron breedlovei* is based on a single collection in post-anthesis condition made in May. Flowering is probably in the early wet season, which begins in May in Mexico.

*Philodendron breedlovei* is known only from the type locality in Mexico (Chiapas) in "Bosque Pino-Encino" or "Bosque Caducifolio," at 1300 m elevation. Since its type locality is very near Guatemala, it is expected to be found there as well.

*Philodendron breedlovei* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. The species is characterized by its long internodes; subterete petioles about equaling the blades; narrowly ovate-cordate, yellow-brown drying blades; solitary, short-pedunculate spathes with the outer surface white on the blade and reddish on the tube; and especially by its ovaries, which have parietal placentation and about 20 ovules per locule.

*Philodendron breedlovei* is most easily confused with *P. sousae* Croat, which has similarly shaped blades with weakly coalesced basal veins that often dry a similar yellow-brown color. That species differs in having only 1-3 basal ovules per locule and persistent stigmas bearing a conspicuous rim. The old stigmas of *P. breedlovei* instead are truncate and smooth with up to six more or less equally spaced stylar canals in a ring around the center.

***Philodendron brenesii*** Standl., Publ. Field Mus. Nat. Hist., Bot. Ser. 18: 140. 1937. TYPE: Costa Rica. Alajuela: La Palma de San Ramón, 1050-1100 m, *Brenes 5110* (holotype, F). Figures 36, 86-89.

Hemiepiphytic, sometimes terrestrial; stem gray-green, to 2 m long, glaucous; sap watery, spicyscented; internodes weakly glossy, becoming matte, 2.5 cm long, 2.5-5 cm diam., about as long as broad, medium to dark green, weakly glossy to semiglossy, drying gray, epidermis brown, crisp; cataphylls to 24 cm long, sharply 1-ribbed to sharply 2-ribbed, sharply and broadly sulcate, pale green to greenish brown to reddish or weakly tinged red near base, densely short dark lineate, drying tannish brown, promptly deciduous, obtuse at apex, margins clear to pale; **petioles** 20-53 cm long, 0.6-1.7 cm diam., subterete, somewhat spongy, yellowish green, very broadly convex to weakly flattened to obtusely flattened with obtuse medial rib

adaxially, rounded to convex abaxially, with adaxial margins rounded, surface sparsely short, dark green or reddish lineate, sometimes with green to reddish ring around apex; **blades** narrowly ovate, subcoriaceous, short- to long-acuminate at apex,  $\pm$  sagittate at base, 28–79 cm long, 8–38 cm wide, (1.6)1.8–2(2.4) times longer than wide, 1.1–1.4 times longer than petioles, upper surface medium to dark green, semiglossy to glossy, lower surface pale green to bluish green, matte to glaucous; anterior lobe 30–52 cm long, 20–30 cm wide (4.3–5.5 times longer than posterior lobes); posterior lobes 5.5–12 cm long, 8.8–13 cm wide, rounded, directed toward base; sinus narrowly V-shaped; midrib very broadly convex to flat, whitish to pale green to pale reddish green, sometimes sparsely short red-lineate, at least near base above, narrowly convex to convex, pale green to reddish, matte, sometimes dark red-lineate below; basal veins 5–6 per side, 2 free to base, the third and fourth sometimes coalesced to 3.5 cm; posterior rib not naked; primary lateral veins (5–6)8–12(18) per side, departing midrib at a 60–95° angle, sunken above, convex and paler than surface below; minor veins visible, etched-sunken and paler above, slightly raised and slightly darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, 1–2 per axil; peduncle 4.5–10.5 cm long, 0.5–1 cm diam., convex adaxially, rounded to angular abaxially, medium green, semiglossy; **spathe** thin, 9.5–18 cm long (1.7–2.1 times longer than peduncle), constricted  $\pm$  at the middle, pale yellowish green throughout, spathe blade cream inside; spathe tube 4–6 cm long, semiglossy outside, deep magenta with resin canals inside; **spadix** sessile,  $\pm$  tapered, 11.5–17.3 cm long, broadest at the base; pistillate portion pale yellow, cylindrical, 2.8–8 cm long, 0.8–1.4 cm diam., tapered toward base; staminate portion 8.6–10.8 cm long, only slightly broader than pistillate portion; fertile staminate portion cream, generally tapered, 9–14 mm diam. at base, 1.1–1.2 cm diam. at middle, 4–6 mm diam. ca. 1 cm from apex, broadest at base, usually narrower than the pistillate portion; sterile staminate portion 1–1.5 cm diam.; pistils 3.1–4.7 mm long, 1.7–2.8 mm diam.; ovary 5–8-locular, 1.8 mm long, 1.7 mm diam., with sub-basal placentation; ovules 6–12 per locule, arranged in 2 series of 6 ovules, contained within gelatinous matrix (no true envelope), 0.4 mm long,  $\pm$  equal in length to funicle, style 1 mm diam., similar to style type B; style apex flat to weakly rounded; stigma subdiscoid, 1.5–2.1 mm diam., 0.3–1.7 mm high, inserted on center of style apex, shallowly depressed medially; the androecium trun-

cate, 4–6-sided; thecae oblong to ovate, 1.3 mm wide, contiguous; pollen cream, elongate to subspheroidal, 0.1–0.2 mm long, to 0.1 mm diam. INFLORESCENCE with berries yellow maturing to orange; seeds 2 per locule, very pale yellow, 1.4–1.7 mm long, 0.7 mm diam. JUVENILE plants creeping, appressed-climbing; internodes 8–15 mm long, 2–2.5 cm wide; petioles 7–8 mm diam, D-shaped, flattened adaxially, weakly striate; blades narrowly ovate, 19–23.3 cm long, 9–12 mm wide; basal veins 1–2; posterior rib not naked; primary lateral veins 5–10 per side.

Flowering in *Philodendron brenesii* may be aseasonal. Flowering collections have been made in July and August, and many post-anthesis collections have been made between March and November. Fruiting collections have been made mostly during what is the dry season and early wet season in Costa Rica, December through May. Mature fruiting collections have been made in December, January, February, and May.

*Philodendron brenesii* ranges from Costa Rica to central Panama, at 800 to 2200 m elevation in *Premontane rain forest* and *Tropical Lower Montane rain forest* life zones.

*Philodendron brenesii* is a member of *P.* sect. *Calostigma* subsect. *Macrobelyum* ser. *Ecordata*. The species is distinguished by its mid-elevation habitat, bright green internodes about as long as broad, sharply two-ribbed, promptly deciduous cataphylls, and narrowly ovate blades with a narrow V-shaped sinus, more or less free basal veins, and a more or less bluish green lower surface.

*Philodendron brenesii* is most easily confused with *P. crassispatum* Croat & Grayum, which has similar venation but more broadly ovate blades and a nearly ellipsoid spathe with very thickened walls (usually more than 1 cm thick) and no sign of a constriction between the blade and tube portions. The peduncle on the latter is sometimes so short that it appears to be lacking. In contrast, the spathe of *P. brenesii* is of normal thickness and shape (e.g. the spathe is divided into a tube and blade portion). *Philodendron straminicaule* can sometimes be confused with the more narrow-leaved forms of *P. brenesii*, but that species differs in having a hippocrepiform, rather than a narrowly V-shaped, sinus.

The leaf blades of *Philodendron brenesii* closely resemble those of *P. validinervium* Engl. from coastal Ecuador between Nanegal and Gualaes. That species differs in having proportionately longer and narrower posterior lobes, more closely spaced and prominently raised primary lateral veins, a long-pedunculate spathe (1.2 times longer than the pe-

duncle vs. 1.6–3.1 times longer), and a 4-locular ovary with about 4 ovules per locule (vs. a 5–7-locular ovary with up to 12 ovules per locule).

*Philodendron brenesii* is one of the most commonly cultivated species throughout the Meseta Central in Costa Rica. A specimen collected in Coclé Province of Panama (Croat 67578) is somewhat disjunct from the nearest populations in Chiriquí Province, but perhaps belongs here as well. The material has juvenile blades more broadly ovate than in other populations of the species. It also lacks the bluish green coloration on the lower blade surface and has the dried midrib dark rather than pale. The adult blades of the Coclé collections are remarkably similar to those of the Chiriquí collections, and the other differences may be due to the fact that these plants occur near the lower part of the elevational range. A number of differences in the pistil argue that this collection may represent a different species. Pistil differences in the Coclé collection include, among other things, the presence of a style funnel and a style dome (lacking in the Chiriquí populations). The Coclé collection also has eight locules per ovary and two seeds per locule, perhaps another important difference.

A Costa Rican collection (Herrera et al. 8789) from Tarrazú in San José Province is unusual in having smaller leaves with indistinct primary lateral veins and petioles drying minutely wrinkled. The dried blade color, minor venation, and dried stem characters otherwise match *P. brenesii*. While this might represent a new species, more information is needed. Herrera et al. 8789 also is similar in stature and blade shape with *P. knappiae* from Chiriquí Province in Panama. The collection differs from *P. knappiae* in its pale gray-green drying (vs. dark brown) lower blade surface, lack of secretory ducts between the minor veins, and a deeply sunk-into style into the apex of the pistil on drying (vs. a style held above the apex of the pistil on drying).

*Additional specimens examined.* COSTA RICA. **Alajuela:** San Ramón–Balsa, ca. 5.7 km N of Quebrada Volio, 1100–1150 m, 10°08'N, 84°29'W, Stevens 14124 (CR, F, MO, NY, US); Cerro el Chayote–Zarcero, Poveda 1187 (CR, F, MO); 8.9 mi. NW of San Ramón, 1100 m, 10°10'30"N, 84°30'W, Croat 68084 (B, CM, DUKE, F, K, M, MEXU, MO, NY, US); 1.3 mi. N of Angeles Norte, 1200 m, Croat 46879 (MO); ca. 11 mi. NW of Zarcero, 900 m, Croat 43521 (CR, MO); 1.5 km past Zarcero, 1800 m, Hoover 1350 (CR, MO); Volcán Poás–Volcán Barba, 7 mi. N of Carrizal, 1850 m, Croat 35491 (F, MO). **Cartago:** ca. 7.3 km NE of Pacayas, 5200 ft., Wilbur et al. 16066 (MO); road to Moravia, ca. 20 km E of Río Pacuare, 1150 m, 9°50'N, 83°24'W, Thompson & Rawlins 1230 (CM); Cerro de la Muerte, between summit & Empalme, 1800 m, Croat 35415 (CR, L, MEXU, MO); La Cangreja, 10 km S of El Tejar, 1850 m, Williams et al. 24191 (F, NY);

Río Grande de Oro, 15 km S of Tapantí, E slope above río, 1500 m, Burger & Liesner 6713 (F); Río Naranjo, 3.5 km E of Cachí, 1360 m, Lent 1431 (F); Tapantí Hydroelectric Reserve, along Río Grande de Oro, 4.5 km beyond small bridge, 1500–1700 m, Croat 36111 (MO); ca. 1 km S of jct. of Quebrada Salto & Río Grande de Oro, 1500–1800 m, 9°43'N, 83°47'W, Croat & Grayum 682264 (MO); 1200 m, Lent 990 (F). **Guanacaste:** Guachipelín–El Volcán de la Vieja, Brenes 15565 (F, NY); SW slopes of Volcán Rincón de la Vieja and Volcán Santa María, trail from Hacienda Guachipelín, 1400 m, 10°48'N, 85°21'W, Burger & Pohl 1471 (CR, F, MO, PMA); Filá del Volcán Cacao, 1400–1520 m, Chacón & Chacón 2302 (MO); 1 km N of Las Nubes village, 8 km NW of Monteverde, 1200 m, 10°22'N, 84°51'W, Haber & Zuchowski 9519 (CR, INB, MO, MV); Río Colorado, 820 m, 10°46'30"N, 85°20'35"W, Rivera 660 (CR, MO). **Heredia:** 2 km S of Vara Blanca, 1900 m, Wilbur et al. 15711 (DUKE); NW slopes of Volcán Barba, Río San Rafael, Lent 1299 (CR, F, US); Volcán Poás–Vara Blanca, 1.5 km past divide in road, 1930 m, 10°12'N, 84°10'W, Hoover 13474 (MO, W); La Zona Protectora, Río Peje–Río Guacimo, N slopes Volcán Barba, 800 m, Grayum & Schatz 3232 (DUKE); Río San Rafael, Atlantic slope of Volcán Barba, 1500 m, 10°13'N, 84°05'W, Grayum et al. 7750 (MO). **Limón:** Moravia, 1300 m, Williams 16171 (EAP); Cantón de Talamanca, Bratsi, Amubri, Alto Lari, Kivut, between Río Dapari and Río Lari, 1350 m, 9°23'50"N, 83°05'10"W, Herrera 5384 (INB, MO); Talamanca, Río Lari, 1700 m, Fernandez 816 (CR, INB). **Puntarenas:** Monteverde Cloud Forest Reserve, 1450–1650 m, 10°18'N, 84°47'W, Burger & Baker 9767 (CR, F); 1700 m, 10°20'N, 84°50'W, Haber & Bello 4115 (MO); 1700 m, Haber 2413 (MO); 1550–1600 m, Haber & Zuchowski 9827 (INB, MO); Zarcero region, Palmira, 5700 ft., A. Smith 143 (F); Osa, vic. Boscosic, Croat 78806 (CR, INB, MO). **San José:** vic. of Vara Blanca, 1880 m, Croat 35519 (MO); Río Parí Blanca, Cerros de Zurquí, 1600–1800 m, 10°13'N, 84°01'W, Burger et al. 10242 (F, MO); SW fork of Río Parí Blanca, lower slopes of Cerro Zurquí, Utley & Utley 1270 (F); La Palma–San Jerónimo, Utley & Utley 532 (F); 2 km N of Highway 12, ca. 10 km W of Inter-American Highway, 2200 m, Croat 43387 (CR, MO); along CA-2, Cerro de la Muerte, N of turn off for road 222, 2000 m, Croat 32857 (MO); Patarrá, Cerro El Espino, 1600–1800 m, 9°53'N, 84°02'W, Chacón & Herrera 1586 (CR, MO); Aserri, Cerros de Escarú, 1950–2100 m, Morales 1305 (CR, INB); Rincón de la Vieja, Boucler 233 (CR); Tarrazú, Herrera et al. 8789 (MO). **PANAMA. Chiriquí:** Callejón Seco, Volcán de Chiriquí, 1700 m, Woodson & Schery 510 (F, GH, MO); 4 km past divide in road to Alto Quiel from Boquete, 1600 m, 8°49'N, 82°28'W, Hoover 1337 (MO); Gualaca–Chiriquí Grande, 5.9 mi. beyond Los Planes de Hornito, 1225 m, 8°45'N, 82°14'W, Croat 67793 (AAU, CM, COL, CR, F, G, K, L, MEXU, MO, OOM, QCA, SAR, TEX, US); 5.5 mi. NW of Los Planes de Hornito, 1320 m, 8°40'N, 82°14'W, Croat 74914 (MO); Cerro Colorado, ca. 13 mi. N of Río San Félix bridge, 800–1200 m, Croat 33507 (MO); 24 mi. N of Río San Félix, 1430–1500 m, Croat 48487 (MO); above San Félix, 33184 (MO); Los Planes de Hornito beyond Gualaca, 1400–1900 m, Croat 48879 (MO, SAR, US); Boquete region, W of Río Caldera, ca. 2 km NW of Bajo Mono, 1700 m, 8°49'N, 82°28'W, Grayum et al. 6448 (MO); 8.3 mi. W of Chame, 1300 m, 8°35'N, 81°50'W, Croat 69090 (AAU, CR, MO, NY, PMA, VDB); near Continental Divide, 1500 m, Antonio 1497 (MO, NY); Cerro

Hornito, S slope approached from Los Planes de Hornito, 1640 m, 8°41'N, 82°11'W, *Croat* 67937 (CAS, MO); Cerro Horqueta, 1650 m, *Croat* 27000 (CM, MO). **Costa Rica:** Alto Calvario along summit of Continental Divide 5.5 mi. N of El Copé, 3.5 mi. N of Escuela Barrigón, 850 m, 8°39'N, 80°36'W, *Croat* 67578 (CM, CR, K, MO, NY, PMA, US); 1.5 mi. N of El Copé, ca. 900 m, *Croat* 44577 (MO).

***Philodendron brevispathum*** Schott, *Bonplandia* 7: 29. 1859. TYPE: Panama. Canal Zone: Chagres River, *Fendler* 431 (holotype, MO; isotype, K). Figures 90, 91.

*Philodendron arcuatum* K. Krause, in Engl. & K. Krause, *Pflanzenr.* IV. 23Db (Heft 60): 72. 1913. TYPE: Bolivia. Pando: Río Acre, Cobija (on Brazilian border, SW of Río Branco), 11°02'S, 68°44'W, *Ule* 8819 (holotype, B).

*Philodendron holmquistii* G. S. Bunting, *Acta Bot. Venez.* 10: 297. 1975. *Philodendron brevispathum* subsp. *holmquistii* (G. S. Bunting) G. S. Bunting, *Phytologia* 64: 466. 1988. TYPE: Venezuela. Amazonas: Pueblo Viejo, open zone between the Río Pacimoni and the forest, 100 m, 1°50'N, 66°30'W, *Steyermark & Bunting* 102495 (holotype, VEN; isotype, MY).

Terrestrial or hemiepiphytic; stem scandent, coarsely scabrous, densely covered with trichome-like, often branched scales; internodes elongate, 7.5–14(23) cm long, 8–15 mm diam., longer than broad, epidermis reddish brown, loosening and flaking, without fissures; roots drying tan to dark brown, few per node; cataphylls 6–9 cm long, soft, unribbed, green, drying dark reddish brown, eventually deciduous; **petioles** 10–43 cm long, 3–7 mm diam., subterete to bluntly C-shaped, somewhat spongy; **blades** ovate to ovate-triangular, concolorous, semiglossy, thin, acuminate at apex, cordate at base, 16–36 cm long, 11–26 cm wide (1.4–2 times longer than wide), (0.8–1.6 times longer than petiole), broadest at or above middle, upper surface semiglossy, lower surface semiglossy; anterior lobe 12–30 cm long, 10.8–24 cm wide (1.3–2.4 times longer than posterior lobes); posterior lobes somewhat triangular to narrowly triangular to rounded, or rounded to with hastate or flaring lobes, 5–15 cm long, 4.9–11.5 cm wide, subacute; sinus narrow or sometimes V-shaped, 9–11 cm deep; midrib sunken above, convex to raised, drying slightly paler below; basal veins 3–4 per side, with 1 free to base, coalesced 4–10 mm, or with 2–3 veins coalesced to 3 or 4 cm; posterior rib not naked or obscurely and briefly naked to 0.5 cm long; primary lateral veins (3)4–5 per side, departing midrib at a 55–70° angle, gradually ascending to the margins, weakly sunken above, convex and drying slightly darker below; interprimary veins almost as conspicuous as primary lateral veins; minor veins arising from both the midrib and primary lateral veins;

tertiary veins visible and darker than surface below. **INFLORESCENCES** erect, 1 per axil; peduncle 1–7 cm long, 2.5–5 mm diam., subterete; **spathe** 6–10.5 cm long (1.4–4.6(9) times longer than peduncle); spathe blade green outside, red inside; spathe tube green to greenish white outside, 4–4.5 cm long, 2–2.5 cm diam., green to white inside; **spadix** sessile; 7–9.5 cm long, whitish to yellowish; pistillate portion 2–3 cm long, 3.5 cm diam. throughout; staminate portion 5–7 cm long; fertile staminate portion 5–6 mm diam.; ovary ca. 6-locular, with axile placentation; ovules 6–14 per locule, 0.3–0.4 mm long, longer than funicle; funicle 0.2 mm long, adnate to lower part of partition, style similar to style type B; stilar canals emerging at base of pronounced apical depressions; style apex drying dark brown with a pale undulate margin and a central solitary stigmatic pad; the androecium truncate, prismatic, margins distinctly scalloped, irregularly 4–5-sided, 0.9 mm long; thecae oblong, 0.4 mm wide, nearly contiguous. **INFURCESCENCE** with peduncle 4 cm long, spathe 5–6 cm long; spadix, 3.5–4 cm long, 2.5 cm wide; berries pale yellow, 6 mm long, 3 mm diam.; seeds 4–6 per berry, oblong to weakly ovoid, 1.4 mm long, 0.4–0.5 mm diam.

Flowering in *Philodendron brevispathum* is apparently aseasonal, perhaps owing to its habitat along and near streams. Flowering collections have been made in February through April, July, August, and November. Immature fruiting collections were also made throughout most of the year, January through April, July, September, and November.

Though long known only from the Caribbean coast of Panama, *Philodendron brevispathum* is now known to be a locally rare but more widespread species with special ecological requirements. It ranges from Nicaragua to Panama at 0–280 m with disjunct populations in South America, there ranging from Venezuela to Brazil (Amazonas), Colombia (Vichada, Meta), Ecuador (Napó), Peru (Loreto), and Bolivia, at elevations of 60 to 230 m.

*Philodendron brevispathum* is a member of *P.* sect. *Philodendron* subsect. *Solenostegma*. It grows as a low hemiepiphyte or in drier areas as a terrestrial herb, usually along streams, on river banks, or in swamps. *Maguire* 36157 is unusual in being reported as a vine growing to 13 m in trees. The species is recognized by its scandent habit, its thin ovate to ovate-triangular leaf blades with rounded to hastate posterior lobes and a generally narrow (sometimes V-shaped) sinus, and especially by the stems that are densely covered with trichome-like, often branched scales and dry with a



flaky brown periderm. While the posterior leaf lobes are commonly somewhat triangular (and may even be narrowly triangular and subacute at apex), they may be nearly round on the same collection (Davidse & Gonzalez 12973).

Bunting (1988) distinguished the South American material as *P. brevispathum* subsp. *holmquistii*, characterized by having 7–14 ovules per locule (vs. 14–18 for the typical material in Central America) and pistils with the apex convex with 3–4 stigmatic pads. Yet my observations indicate that Central American material also has 6–14 ovules per locule.

This species is closest to *P. muricatum* Willd. ex Schott (an older name) and may not be separable from it. The latter species is distinguished by having densely verrucose-warty petioles and usually smaller blades with more rounded lobes. However, some collections with verrucose petioles such as Steyermark et al. 115076 and Liesner 9056 from Delta Amacuro, Venezuela, and Krukoff 7250 from southern Amazonas State, Brazil (the type of *P. amplexans* A. C. Sm., a synonym of *P. muricatum*), have narrow, more or less triangular blades like those of *P. brevispathum*. On the other hand, not all specimens of *P. brevispathum* have similar lobes. Central American material commonly has more rounded or elongate lobes that are turned somewhat inward, but some sheets (e.g., Burger & Antonio 11236) have blades identical to those of South American plants. In addition, some South American collections (Croat 58586, Davidse 4294) have the posterior lobes noticeably rounded and scarcely longer than broad. Despite this variation, the verrucose petiole character is adequate to separate *P. brevispathum* from *P. muricatum*.

*Philodendron brevispathum* has also been confused with *P. jacquinii*, but that species has merely puberulent stems (with trichomes simple and unbranched), thinner, more broadly ovate leaves (often also puberulent on the petiole and lower midrib), a conspicuously bulging spathe tube, and more elongate styles.

**Additional specimens examined.** COSTA RICA. **Heredia:** Puerto Viejo-Guápiles, along Río Puerto Viejo, 7 km N of Buenos Aires, 10°23'30"N, 83°48'30"W, Croat 68401 (K, MO); La Selva Field Station, Grayum 2300 (DUKE, MO); Grayum 2642 (DUKE, F, MO), 100 m, Hammel & Drainer 10810 (DUKE); Río Frío de Sarapiquí, between Río Sucio and railroad tracks, SW of Finca Zona Ocho, 110 m, 10°18'N, 83°52'30"W, Grayum & Hammel 5568 (MO). **Limón:** Río Bananito-Cahuíta, near Punta Vargas ca. 4 km S of Cahuíta, 0–10 m, Burger et al. 10493 (F, MO); 16 airline km SW of Barra del Colorado, 10–120 m, 10°39'N, 83°40'40"W, Davidse & Herrera 31254 (MO); Refugio Barra del Colorado, area between Río Chirripocito and Río Sardina, 12 m, 10°38'N, 83°45'W, Grayum et al. 9742 (AAU, CR, MO); Finca Tapezco-La Suerte, 29 air

km W of Tortuguero, 40 m, 10°30'N, 83°47'W, Davidson & Donahue 8842 (MO); 40 m, 10°30'N, 83°47'W, 8970 (RSA, MO); Gandoca (slightly to N of trail from Mata de Limón), 0–5 m, 9°36'N, 82°36'30"W, Grayum et al. 8024 (CR, K, M, MO); Cerro Coconel, 10–40 m, 10°40'N, 83°40'W, Stevens et al. 24670 (CR, MO); Parque Tortuguero National Park, Tortuguero, 2 m, Robles 1410 (CR, MO); 4 m, 10°32'N, 83°30'W, Robles 1877 (CR, F, MO); near Boca de las Lagunas de Tortuguero, 0–30 m, 10°34'N, 83°32'W, Burger & Antonio 11236 (F, MO, U). **NICARAGUA. Matagalpa:** Rancherta, 11 km al NE de Muy Muy, 280 m, 12°46'N, 85°31'W, Moreno 24433 (MO). **Río San Juan:** Quebrada Santa Crucita, 50 m, 11°02'N, 84°25'W, Moreno 23445 (MO); Sábalo, 1 km N of Río San Juan, 100 m, 11°02'N, 84°27'W, 26111 (MO); Boca de Sábalo, cerca de "La Toboba," 70–90 m, 11°03'–04'N, 84°28'–29'W, Robledo 1833 (MO, US). **Rivas:** Isla de Ometepe, La Argentina, 300–800 m, 11°27'N, 85°32'W, Moreno 22112 (CM, MO). **Zelaya:** La Barra de Punta Gorda, 0–2 m, 11°30'N, 83°47'W, Moreno & Sandino 13298 (MO); Caño Monte Cristo, "La Grapera," 10 m, 11°33'N, 87°48'W, Moreno & Sandino 14739 (MO); Puerto Cabezas-Río Wawa, Ibo Tingni, drainage of Caño Sung Sung, 10 m, 14°9'–11'N, 83°29'–31'W, Stevens 10636 (ENCB, MO); Santa Marta, 5 m, 14°18'N, 83°37'W, Stevens & Moreno 19623 (MO); SW of Bluefields, 10–40 m, 11°59'N, 83°46'W, Stevens 19736 (CAS, L, MO, UWL). **PANAMA. Colón:** Miguel de la Borda, Croat 10012 (MO, SCZ). **San Blas:** Playón Chico, road to Iraisakun, 0–10 m, 9°20'N, 78°13'W, Herrera 596 (MO).

#### *Philodendron brewsterense* Croat, sp. nov.

**TYPE:** Panama. Comarca de San Blas: Cerro Brewster, 850 m, 9°18'N, 79°16'W, de Nevers, Herrera, Hammel & Charnley 5545 (holotype, MO-3246994). Figure 92.

Planta epiphytica; caulis scandens; internodia 5–6.5 cm longa, 4–5 mm diam.; cataphylla leniter vel acute 2-costata, decidua; petiolus 9–11 cm longus, usque 4 mm diam., anguste et obtuse sulcatus; lamina ovata, leniter subcordata, 9.8–11 cm longa, 7–8 cm lata, in sicco flavibrunnea; nervis lateralibus 1 absentibus; inflorescentia 1; peduncululus 4.7 cm longus, 3.5 mm latus; spatha omnino in superficiebus ambabus maronina, 9–9.3 cm longa; pistilla 5-locularia; loculi 2-ovulati.

Epiphytic; stem scandent; leaf scars inconspicuous; internodes terete, 5–6.5 cm long, 4–5 mm diam., much longer than broad, drying yellowish brown, epidermis fissured narrowly; roots dark brown, drying moderately fuzzy, slender, 1 mm diam., very few per node; cataphylls weakly to sharply 2-ribbed, glossy, deciduous; petioles 9–11 cm long, to 4 mm diam., narrowly and obtusely sulcate adaxially, surface minutely wrinkled; sheath conspicuous, the tip free-ending (2–3 mm long); blades ovate, acuminate at apex (acumen inrolled, 1–2 mm long), weakly subcordate at base, 9.8–11 cm long, 7–8 cm wide (1–1.4 times longer than wide; about equal in length to petiole), broadest in lower one-third, margins revolute, drying reddish brown, upper surface medium green, drying grayish

green, semiglossy, lower surface weakly glossy, moderately paler, drying yellowish green; anterior lobe 9–10.5 cm long, 7–8 cm wide (5.9–7.1 times longer than posterior lobes), broadest in lower one-third; posterior lobes 2–2.4 cm long, 2.8 cm wide, rounded; sinus obtusely angular, 5 mm deep; posterior rib lacking; midrib concave above, moderately raised, drying paler than surface below; basal veins numerous on each side but none of them outstanding, primary lateral veins lacking or indistinguishable from minor veins; minor veins close, arising from the midrib only, arising from both the midrib and primary lateral veins. INFLORESCENCES 1 per axil; peduncle 4.7 cm long, 3.5 mm diam., dark brown; spathe erect, subcoriaceous, 9–9.3 cm long (2.1 times longer than peduncle), maroon throughout on both surfaces; spathe blade 6.9 cm long; spathe tube 4.5 cm long; spadix green throughout (probably juvenile), narrowly rounded at apex, 7.3 cm long, constricted below sterile staminate portion; pistillate portion drying grayish, slightly tapered toward the apex, 2.1–2.5 cm long in front, 1.2 cm long in back, 4.5–5 mm diam. at apex, 5–6.8 mm diam. at middle; staminate portion 4.5–6.1 cm long; fertile staminate portion white, gradually tapered towards apex, 6.5–6.8 mm diam.; pistils 1.1 mm long; ovary 5-locular, 0.9 mm diam., ovule sac ca. 0.5 mm long, with sub-basal placentation; ovules 2 per locule, contained within transparent ovule sac, ca. 0.2–0.3 mm long, longer than funicle; funicle 0.1–0.2 mm long (can be pulled free to base), style 0.7–0.9 mm diam., similar to style type D; style apex rounded; style boss broadly and gently rounded; stigma button-like, distinctly lobed, 1 mm diam., 0.3 mm high, covering entire style apex, inserted on style boss; the androecium truncate, prismatic, margins irregularly 4–6-sided, 0.9–1.1 mm long, 3–5 mm diam. at apex; thecae oblong to weakly elliptical, 0.4 mm wide,  $\pm$  parallel to one another; sterile staminate flowers subrounded, 5–9 mm wide, grayish to pale orange.

Flowering in *Philodendron brewsterense* is based on a single flowering collection made in April in an area that is somewhat aseasonal.

*Philodendron brewsterense* is endemic to Panama, where it is known only from the type specimen collected on Cerro Brewster in Comarca de San Blas Province, at 850 m elevation in a *Premontane rain forest* life zone.

*Philodendron brewsterense* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. The species is recognized by its scandent habit with internodes much longer than broad, drying yellowish brown and narrowly fissured; small (<8

cm diam., <11 cm long), coriaceous, ovate, yellowish brown blades lacking primary lateral veins; and solitary inflorescences with the outer surface surface maroon.

*Philodendron brewsterense* is apparently unrelated to any other species but is most similar to *P. crassispatum* and resembles juvenile plants of that species. The latter differs, however, in having juvenile leaves with an arcuate sinus and adult leaf blades usually more than 20 cm long and with four to six pairs of primary lateral veins.

***Philodendron brunneicaule*** Croat & Grayum, sp. nov. TYPE: Panama. Coclé: Alto Calvario, 6 mi. N of El Copé, Atlantic slope along Continental Divide, along trail which heads off old lumber trail and leads down to Las Ricas, Limón, and San Juan, 710–800 m, 8°39'N, 80°36'W, 22 June 1988, Croat 68713 (holotype, MO-3591332). Figures 93–97.

Planta hemiepiphytica; internodia 6–20 cm longa, 1–1.2 cm diam., ruberbrunnea; cataphylla 20–40 cm longa, incostata, decidua; petiolus teres vel C-formatus, leviter subcomplanatus adaxialiter 21–63 cm longus, 4–11 mm diam.; lamina ovata, cordata aut interdum sagittata basi, 25–62 cm longa, 16–52 cm lata; inflorescentia 1; pedunculus 4–20 cm longus, 8–9 mm diam.; spatha 11–21 cm longa; lamina spathae extus cremaea, intus viridialba; tubo spathae extus viridi, intus cerasino; pistilla (5)(7)-locularia; loculi (13)15–18-ovulati; baccae albae.

Hemiepiphytic; stem light reddish brown to rusty red, leaf scars conspicuous, 8 mm long, 12 mm wide; internodes sparsely short dark-lineate, sometimes weakly warty, semiglossy, 6–20 cm long, 1–1.2 cm diam., longer than broad, dark green to olive-green to brown, epidermis loosening and flaking, reddish brown; cataphylls soft, 20–40 cm long, unribbed, sometimes bluntly 1-ribbed, greenish white to white with margins pinkish, drying reddish, deciduous; petioles 21–63 cm long, 4–11 mm diam., erect-spreading to spreading, terete, to C-shaped, somewhat flattened adaxially, medium green, surface matte, sometimes striate and dark green- to red-lineate; geniculum 3 cm long, 2.5–3 cm diam., thinner and paler than petiole; blades ovate, subcoriaceous, bicolorous, acuminate at apex, cordate or sometimes sagittate at base, 25–62 cm long, 16–52 cm wide (0.6–1.75 times longer than wide), (0.7–1.6 times longer than petiole), upper surface dark green, glossy to semiglossy, lower surface glossy to semiglossy, paler; anterior lobe 23–65 cm long, 19–52 cm wide (0.88–1.5 times longer than wide); posterior lobes rounded to obtuse, 8–20 cm long, 9–26 cm wide; sinus spatulate to hippocrepiform; midrib flat to sunken, paler than

surface above, narrowly convex to round-raised, paler than surface, sometimes tinged maroon below; basal veins 5–9 per side, with 0–2 free to base, (2)3–4 coalesced 1.9–4.5 cm; posterior rib naked for 1–4.5 cm; primary lateral veins 4–6 per side, departing midrib at a 45–55° angle, conspicuously sunken above, narrowly convex and tinged maroon below; interprimary veins weakly raised and darker than surface below; minor veins arising from both the midrib and primary lateral veins; tertiary veins visible and sometimes darker than surface below. INFLORESCENCES erect, 1 per axil; peduncle 4–20 cm long, 8–9 mm diam., medium green, whitish at base, short dark green lineate; **spathe** glossy, 11–21 cm long (1.05–2.85 times longer than peduncle), constricted midway above the tube; spathe blade cream outside, greenish white (at anthesis) inside; spathe tube 6.5–9 cm long, green outside, cherry-red inside; **spadix** 9–17 cm long, broadest above the middle or ± uniform throughout; pistillate portion cylindrical to obovoid, 3.8–7 cm long, 7–13 mm diam. at apex, 8–14 mm diam. at middle, 10–14 mm wide at base; staminate portion 4.6–17.5 cm long; fertile staminate portion clavate, 9–16 mm diam. at base, 12–20 mm diam. at middle, 7–10 mm diam. ca. 1 cm from apex, broadest at middle, broader than the pistillate portion, narrower than the sterile portion; sterile staminate portion as broad as or narrower than the pistillate portion, 0.9–1.6 cm diam.; pistils 1.3–3.2 mm long, 1–1.8 mm diam.; ovary 5(6)7-locular, 0.9–2.1 mm long, 0.9 mm diam., with axile placentation; ovules (13)15–18 per locule, 2-series, 0.1–1.7 mm long, longer than or equal in length to funicle, style 1.2 mm diam., similar to style type B; style apex rounded; stigma subdiscoid, lobed, 1.5 mm diam., 0.3 mm high, covering not quite entire style apex; the androecium truncate, 3–6-sided, 2.2 mm long; thecae oblong, 0.2–0.5 mm wide; sterile staminate flowers blunt with one side scalloped, 1–2.2 mm long, 0.7–1.9 mm wide. INFRUCTION with berries white. JUVENILE plants with internodes to 10 cm long, 4–10 mm diam.; cataphylls reddish, persisting at upper nodes, sharply C-shaped in cross section.

Flowering in *Philodendron brunneicaule* probably occurs in the early rainy season, but a single (post-anthesis) collection was made in January. Most post-flowering collections have been made between April and July, with immature fruits collected in August and October.

*Philodendron brunneicaule* ranges from Costa Rica to Panama, Colombia (Valle), and Ecuador

(Esmeraldas), at 50 to 1300 m elevation in *Tropical wet forest* and *Premontane rain forest*.

*Philodendron brunneicaule* is a member of *P.* sect. *Philodendron* subsect. *Platyopodium*. The species is characterized by its appressed-climbing habit, long internodes with thin, flaking reddish brown epidermis (hence the name "brunneicaule," from "brunneus" meaning brown, and "caulis" meaning stem), white, unribbed, deciduous cataphylls, somewhat adaxially flattened and red-lineate petioles, ovate blades with reddish-brown-drying veins, large, solitary inflorescences borne at several of the uppermost internodes, and green spathes colored cherry-red inside on the tube.

*Philodendron brunneicaule* is probably related to *P. ernstii* Engl. from Amazonian Ecuador and Peru. That species shares long internodes with flaking brown epidermis and similar, solitary inflorescences. It differs, however, in having stems that are often warty and petioles that are undulate-winged vs. more nearly terete for *P. brunneicaule*. It is noteworthy, however, that a single collection from Amazonas Department, Peru (Vásquez & Aparu 19051, MO), appears to lack a petiolar wing. This may prove to be *P. brunneicaule*, but if so it would be the first collection from east of the Andes.

In Central America *Philodendron brunneicaule* is most easily confused with *P. copense*. See that species for the differences.

*Additional specimens examined.* COSTA RICA. **Alajuela:** San Ramón-Balsa, ca. 16.7 km N of bridge over Quebrada Volio and ca. 7.5 km N of Río Balsa, 700–800 m, 10°10'–15'N, 84°30'–35'W, Stevens 13859 (CR, F, MO); 4.6 km N of bridge over Río Balsa, 900–1000 m, 10°12'N, 84°31'W, Stevens 13735 (MO); 4–7 km N of Balsa, 750 m, 10°13'N, 84°32'W, Liesner & Judziewicz 14667 (CR, MO); 18–19 km N of San Ramón, 950 m, 10°10'N, 84°28'W, Hammel et al. 15234 (MO); Reserva Biológica Monteverde Río Peñas Blancas, 900 m, Bello 369 (CR, INB, MO); Río Sarapiquí, crossing to Colonia Virgen del Socorro, 740 m, 10°5.5'N, 81°10.5'W, Grayum & Hammel 5517 (MO); 830 m, 10°16'N, 84°11'W, Croat 68302 (CM, G, M, MO, NY). **Heredia:** Puerto Viejo-Guápiles, 7 km N of Buenos Aires, <100 m, 10°23'30"N, 83°48'30"W, Croat 68402 (MEXU, MO); La Selva Field Station, Grayum 2790 (F, MO). ECUADOR. **Esmeraldas:** Quinindé, Balsa Biological Station, Montañas de Mache, 35 km W of Quinindé, 5 km W of Santa Isabela, 400–600 m, 10°21'N, 79°44'W, Pitman & Bass 1085 (MEXU, MO, NY, QCNE). PANAMA. **Bocas del Toro:** Fortuna Dam area, Fortuna Dam-Chiriquí Grande, 2.8 road mi. N of Divide, 850–950 m, 8°45'N, 82°15'W, McPherson 9661 (MO); 1.2 mi. N of Divide, 5.3 mi. N of bridge over Fortuna Dam, 910 m, 8°44'N, 82°17'W, Croat 60468 (AAU, CM, CR, MEXU, MO, PMA, US); gravel road near Continental Divide, 1170 m, 8°44'N, 81°17'W, Croat 66655 (L, MO). **Chiriquí:** Gualaca-Fortuna Dam, 10.1 mi. NW of Los Planes de Hornito, 1300 m, 82°17'W, 8°45'N, Croat 49836 (COL, K, MO, NY); 9.4 km above El Copé, 750–900 m, Croat 44733 (MO). **Coeló:** El Copé, Alto

Calvario above El Copé, 4.5 mi. N of El Copé, 580–740 m, 8°38'N, 80°36'W, *Croat* 67532 (AAU, MO, W); 930 m, 49183 (MO, US). **Panamá:** El Llano–Cartí road, 10 km from Inter-American Highway, near El Llano, 330 m, *Croat* 33823 (MO); Km 14, 350–500 m, *Folsom et al.* 1483 (MO); 10.1 mi. above highway, 325–350 m, *Croat* 67365 (MO). **San Blas:** El Llano–Cartí Road, km 19, 350 m, 9°19'N, 78°55'W, *de Nevers et al.* 5599 (B, K, MO, PMA); 1–2 mi. N of Nusagandi, 250–275 m, 9°20'N, 79°W, *Croat & Zhu* 76581 (MO). **Veraguas:** Santa Fe region, Santa Fe–Río San Luis, past Escuela Agrícola Alto de Piedra, at Río Segundo Brazo, 480 m, 8°33'N, 81°08'W, *Croat* 66909 (MO).

**COLOMBIA:** Valle: Bajo Calima region, Buenaventura–Málaga, Km 28, 50–150 m, 3°59'N, 77°03'W, *Bay* 240 (CUCV, MO).

**Philodendron chiriquense** *Croat, sp. nov.* TYPE:

Panama. Bocas del Toro: Cerro Colorado, 9.2 km W of Chame, along trail E of road leading down to stream, 1450–1480 m, 8°35'N, 81°50'W, 6 July 1988, *Croat* 69068 (holotype, MO–3599857; isotypes, B, CAS, CM, COL, CR, F, GH, K, MEXU, PMA, US). Figures 98–100, 107.

Planta hemiepiphytica; internodia 1–4 cm longa, (1.5–2)3–4(6–10) cm diam.; cataphylla (20)55–70 cm longa, acute 2-costata, suffusa rubra, in sicco rubrobrunnea, persistentia semi-intacta; petioli subteres vel D-formati, 27–108 cm longi, in sicco 9–15 mm diam.; lamina triangulari-sagittata, cordata basi, 32.5–91 cm longa, 11–44 mm lata; inflorescentia 3; pedunculus 9 cm longus, 8–9 mm diam., albus suffusus ruber; spathe 14.2 cm longa, lamina spathe extus alba, intus albida, suffusa marronina usque ad dimidium; tubo spathe extus pallide viridi, intus marronino; pistilla 5-locularia; loculi circa 14-ovulati.

Hemiepiphytic; stem appressed-climbing to 4 m high or creeping, to 30 cm long; internodes short, very thick, semiglossy, 1–4 cm long, (1.5–2)3–4(6–10) cm diam., much broader than long, medium to pale green, coarse white-ribbed at upper edge, completely hidden by cataphyll fibers, drying brown; roots several per node, to 4 mm diam., light reddish brown to dark brown with flaky lighter brown epidermis, weakly glossy; cataphylls (20)55–70 cm long, sharply 2-ribbed, sharply sulcate with acute marginal ribs, green, heavily tinged with red toward base, sometimes red throughout, drying reddish brown, with two low ribs, persisting semi-intact at upper nodes, as pale fibers at base, eventually as dark brown to reddish brown fibers; **petioles** 27–108 cm long, 9–15 mm diam., subterete to obtusely D-shaped, firm and flexible, obtusely to weakly flattened adaxially, with adaxial margins sharp to bluntly rounded, dark green, base reddish, weakly glossy, surface faintly and densely white striate to coarsely pale striate, drying dark brown; **blades** triangular-sagittate, moderately coriaceous to subcoriaceous, short- to long-acuminate at apex (the acumen some-

times inrolled), cordate at base, 32.5–91 cm long, 11–44 cm wide (1.34–3.28 times longer than wide), (0.58–1.05 times longer than petiole), about equal in length to petiole, broadest at or near point of petiole attachment, upper surface dark green, with velvety sheen, drying gray-green to yellow-brown above, lower surface yellow-green, semiglossy, paler, drying reddish brown below; anterior lobe 39.5–68 cm long, 22–53 cm wide (1.2–1.8 times longer than posterior lobes); posterior lobes 6.2–29.6 cm long, 10.3–26.5 cm wide, directed inward, obtuse to rounded; sinus parabolic, hippocrepiform or spatulate; midrib narrowly rounded to almost flat to weakly sunken, drying gray-brown, paler than surface above, narrow-rounded to convex, concolorous or paler than surface, drying reddish brown below; basal veins (3–4)5–8 per side, with 0–1 free to base, (1)2–3 coalesced 0.9–7.5(11.5) cm, posterior rib naked 2–4 cm, sometimes not naked but with the rib sometimes running very near the margin; primary lateral veins 7–8 per side, departing midrib at a 60–70° angle, nearly straight to the margins, obtusely sunken, concolorous or paler than surface above, prominently convex to narrowly rounded, concolorous or paler than surface below; interprimary veins many, distinct; minor veins fine, moderately distinct; “cross-veins” visible on lower surface, sometimes weakly visible above. **INFLORESCENCES** ± erect, 3 per axil; peduncle 9 cm long, 8–9 mm diam., white tinged red, pale white striate toward apex, turned at 100° angle to spathe at anthesis; **spathe** 14.2 cm long (1.57 times longer than peduncle), weakly constricted above the tube; spathe blade white outside, whitish, suffused maroon to about midway inside; spathe tube globose, 6 cm long, pale green, tinged red on front outside, densely short, pale streaked throughout outside, maroon inside, **spadix** stipitate to 5 mm long; ovate, 11.6–12.5 cm long, broadest below the middle; pistillate portion greenish white, clavate, 2.7 cm long, 1.6 cm diam. at apex, 1.5 cm diam. at middle, 1.1 cm wide at base; staminate portion 10.9 cm long; fertile staminate portion weakly tapered, 1.8 cm diam. at base, 1.6 cm diam. at middle, 1 cm diam. ca. 1 cm from apex, broadest at base, broader than the pistillate portion, narrower than the sterile portion; sterile staminate portion broader than the pistillate portion, 1.8 cm diam.; pistils 4.4 cm long, 1.8 mm diam.; ovary 5-locular, with axile placentation; ovules ca. 14 per locule, 2-seriate, contained within gelatinous matrix (no true envelope), 0.6 mm long, longer than funicle, style 2 mm diam., similar to style type B; style apex rounded; stigma subdiscooid, lobed, 1.5 mm diam., 0.3 mm high, drying lobed, covering center of style apex; the androecium truncate, margins 4–6-sided; thecae oblong, 0.2 mm

wide; sterile staminate flowers blunt, 4 mm long, 1.6 mm wide. JUVENILE blades with lower surface dark maroon.

Flowering in *Philodendron chiriquense* is known only during July. While this is, in general, the early wet season in Panama, the region around Fortuna where the species is common is somewhat less seasonal than other areas of Panama.

*Philodendron chiriquense* is endemic to Panama, known from Bocas del Toro, Chiriquí, and Coclé Provinces in *Tropical Lower Montane rain forest* and *Premontane rain forest* life zones at 500 to 1630 m elevation. It is frequent in the cloud forests near the Continental Divide in the Fortuna Dam area.

*Philodendron chiriquense* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. The species is distinguished by its short, very thick internodes; sharply two-ribbed, red-tinged cataphylls, which persist as semi-intact, dark brown fibers; obtusely flattened to D-shaped, densely pale-striate petioles; huge, triangular-sagittate blades; and up to three inflorescences per axil with the spathe tube pale green outside and maroon within.

*Philodendron chiriquense* is similar to *P. copense*, which also has persistent reddish brown cataphyll fibers and reddish-brown-drying, more or less triangular blades, but that species differs in having petioles drying reddish brown, smooth and matte, with peeling periderm (usually blackened, minutely fissured and semiglossy in *P. chiriquense*), primary lateral leaf veins drying reddish brown and darker than the surface (pale and lighter than the surface in this species), and minor veins and "cross-veins" both distinct (minor not distinct and "cross-veins" scarcely or not at all visible in *P. chiriquense*).

The only collection from Coclé, *Croat 44566*, may represent another species. This collection differs from Chiriquí material in having the primary lateral veins scarcely paler than the surface and more prominent "cross-veins."

*Additional specimens examined.* PANAMA. **Bocas del Toro:** Cerro Colorado, 8.6 mi. W of Chame, 1450–1480 m, 8°35'N, 81°50'W, *Croat 69133* (CM, MO); 7 mi. W of Chame, 1500 m, 8°35'N, 81°50'W, *69215* (MO, NY). **Chiriquí:** Cerro Colorado, 28 mi. above San Félix, 1200–1500 m, *Croat 33266* (MO); 20.5 mi. N of Río San Félix, 8.3 mi. W of Chame, 1630 m, 8°32'N, 81°46'W, *Croat 75047* (MO, PMA); 6.6 mi. W of Chame, 1475–1485 m, *Croat 75011* (CM, MO, PMA); Fortuna Dam area, Gualaca–Chiriquí Grande, 1 km S of Continental Divide, 1075 m, 8°45'N, 82°18'W, *Croat 66865* (MO); Los Planes de Hornito, 1400 m, *Croat 48761* (CAS, L, MEXU, MO, NY, PMA); 7.9 mi. NW of Los Planes de Hornito, 1300 m, 8°44'N, 82°16'W, *Croat 49905* (K, MO, US); 8.3 mi. NW of Los Planes de Hornito, 1260 m, 8°44'N, 82°16'W,

*Croat 49940* (MO); 10 mi. NW of Los Planes de Hornito, 1260 m, 82°17'W, 8°45'N, *Croat 50101* (MO); Fortuna–Chiriquí Grande, 1170 m, 8°44'N, 81°17'W, *Croat 66666* (AAU, CAS, CM, COL, F, KYO, L, MEXU, MO, NY, PMA, QCA, RSA, SAR, TEX, US); 8°44'N, 82°17'W, *Croat & Zhu 76345* (MO); Fortuna Lake area, 3.4 km N of Quebrada Chorro, 1.6 mi. N of center of bridge over lake, 1205 m, 8°43'N, 82°14'W, *Croat 74956* (MO). **Coclé:** El Copé region, near Continental Divide, ca. 1.5 mi. N of El Copé, 900 m, *Croat 44566* (MO); Alto Calvario along summit of Continental Divide 5.5 mi. N of El Copé, 850 m, 8°39'N, 80°36'W, *Croat 67573* (MO, NY); ca. 6 mi. N of El Copé, 770 m, 8°38'N, 80°35'W, *Croat & Zhu 76790* (MO, PMA). **Veraguas:** Santa Fe region, ca. 15 km past Escuela Agrícola Alto Piedra above Santa Fe, Río Caloveborita, Atlantic watershed, 500 m, *Sytova & Andersson 4772* (MO).

***Philodendron chirripoense*** *Croat & Grayum*, sp. nov. TYPE: Costa Rica. San José: Canaán–Chirimol, above Río Chirripó del Pacífico, remnant forest along river and on steep slopes above river, 9°27'N, 83°37'W, 30 Dec. 1969, *Burger & Liesner 7139* (holotype, F; isotype, CR). Figure 101.

Planta hemiepiphytica; caulis scandens; internodia 20 cm longa, minus quam 3 mm diam.; cataphylla decidua; petiolus teres, 7–7.4 cm longus, 1 mm diam., obtuse complanatus adaxialiter; lamina anguste ovata, subcordata basi, 11–11.6 cm longa, 5–5.5 cm lata; nervis basilibus liberis ad basin; inflorescentia 1; pedunculus 6 cm longus, 4 mm diam.; spathe 7.4 cm longa, intus viridi vel rubella; pistilla (3)5–6(8)-locularia; loculi 8–12-ovulati.

Hemiepiphytic; stem scandent; internodes very long and slender, 20 cm long, less than 3 mm diam.; roots drying dark brown; cataphylls unknown, probably <10 cm long, lanceolate, deciduous; petioles 7–7.4 cm long, 1 mm diam., obtusely flattened abaxially; blades narrowly ovate, very long-acuminate at apex (the acumen 2.4–2.6 cm long), subcordate at base, 11–11.6 cm long, 5–5.5 cm wide (ca. 2 times longer than wide), (ca. 1.5 times longer than petiole), about equal in length to petiole, drying brown; posterior lobes rounded to obtuse, 2–3 mm long, 1.4–1.7 cm wide; sinus arcuate with blade decurrent on petiole, 2–3 mm deep; midrib drying with up to 5 ribs above; basal veins 2–3 per side, with all free to base, obscure; primary lateral veins 2 per side, departing midrib at a 40–45° angle, straight to the margins; minor veins arising from the midrib only. INFLORESCENCES nearly as long as leaves, probably 1 per axil; peduncle 6 cm long, 4 mm diam., 0.83 times the petiole length; spathe 7.4 cm long (1.2 times longer than peduncle), green to reddish within; spadix pinkish throughout (fide field notes, in part lost); pistils 2.2 mm long, 1.1–1.2 mm diam.; ovary (3)5–6(8)-locular, with axile placentation; ovules

8–12 per locule, mostly 1-seriate, 0.1–0.2 mm long, slightly longer than funicle; funicle ca. 0.1 mm long, adnate to lower part of partition, style similar to style type B; style apex domed; the androecium truncate, oblong, prismatic, margins irregularly 4–5-sided, 0.7–1 mm long; thecae oblong, 0.3 mm wide,  $\pm$  parallel to one another; sterile staminate flowers irregularly 4–5-sided, 0.8–1.2 mm wide.

Flowering in *Philodendron chirripoense* is based on a single fertile collection made in December, which is the early dry season on the western slope of Costa Rica.

*Philodendron chirripoense* is known only from the type specimen from the Pacific slope of southern Costa Rica, in *Premontane wet forest* at 1000 m elevation.

*Philodendron chirripoense* is tentatively placed in *P. sect. Philodendron subsect. Canniophyllum*. The species is characterized by its scandent habit, very long and slender internodes (drying <3 mm diam.), slender petioles about as long as the small, narrowly ovate, subcordate blades, and especially by the inflorescence being longer than the petioles.

*Philodendron chirripoense* is not similar to any other species in either Central or South America, but it might be confused with *P. microstictum*, the only other species that has the inflorescence longer than the leaves. The latter has larger blades (more than 13 cm wide), which are often broader than long.

***Philodendron clewellii* Croat, sp. nov. TYPE:**

Panama, Darién: middle slopes of W side of Cerro Pirre, cloud forest, 850–1050 m, 29 June 1988, *Croat 68945* (holotype, MO–3589994; isotypes, F, K, PMA, US). Figure 102.

Planta epiphytica; internodia 12–18 cm longa, 3.5 cm diam., moderate nitida, fissurata; cataphylla mollia, acute 2-costata, decidua; petiolus subteres, obtuse complanatus adaxialiter, 24.5–53 cm longus, (2)7–14 mm diam.; lamina ovato-cordata, 30–59 cm longa, 20–45 cm lata, in sicco brunnea, nervis basalibus liberis (aut leniter coalatis); inflorescentia usque 6; pedunculus 3.5–16 cm longus, 3–6 mm diam.; spathe 4.5–7 cm longa; lamina spathe extus cremaea, intus purpurascens cum margine albido; tubo spathe in superficiebus ambabus purpurascens; pistilla 6(7)-locularia; loculi 2-ovulati.

Epiphytic; stem scandent, smooth, thick, moderately glossy; internodes moderately glossy, 12–18 cm long, 3.5 cm diam., longer than broad, medium green to dark gray-green, epidermis fissured weakly longitudinally; cataphylls 18–25 cm long, soft, sharply 2-ribbed, green, caducous; **petioles** 24.5–53 cm long, (2)7–14 mm diam., subterete, spongy,

dark green, obtusely flattened adaxially, surface green or white streaked-lineate; **blades** broadly ovate-cordate, subcoriaceous, moderately bicolorous, acuminate at apex (the acumen 1–2 cm long), cordate at base, 30–59 cm long, 20–45 cm wide (1.25–1.83 times longer than wide), (0.95–1.65 times longer than petiole), upper surface dark green, drying brown, semiglossy, lower surface slightly paler, drying brown, sometimes red-brown or green-brown, semiglossy, paler; anterior lobe 26.7–42.2 cm long, 23–45 cm wide (2.36–3.95(5.28) times longer than posterior lobes); posterior lobes broadly rounded, often overlapping, 7–18 cm long, 12–21 cm wide, obtuse to rounded; sinus narrowly closed; midrib weakly raised to almost flat, purple-spotted, paler than surface above, convex, darker than surface below; basal veins 3–5 per side, all free to base (or weakly fused); posterior rib absent or to 1.5 cm long, not at all naked; primary lateral veins 4–7 per side, departing midrib at a 45–60° angle, straight to the margins, weakly sunken, concolorous above, raised below, secondary veins weakly visible; minor veins visible, sparse, discontinuous below, arising from both the midrib and primary lateral veins. INFLORESCENCES to 6 per axil; peduncle 3.5–16 cm long, 3–6 mm diam., drying 2–4 mm diam., green, heavily red-stripped; **spathe** 4.5–7 cm long ((0.62)1.28–2.33 times longer than peduncle); spathe blade cream outside, purplish, with margins whitish inside; spathe tube purple outside, 2.5–3 cm long, 6–10 mm diam., purple inside; **spadix** sessile; clavate, 6 cm long; pistillate portion green, tapered toward the apex, 2.5 cm long, 6 mm diam. at apex, 7 mm diam. at middle, 8 mm wide at base; staminate portion 3.7 cm long; fertile staminate portion white, clavate, 7 mm diam. at base, 9 mm diam. at middle, 8 mm diam. ca. 1 cm from apex, broader than the pistillate or sterile portions; sterile staminate portion 7 mm diam.; pistils 1.6 mm long, 1.1 mm diam.; ovary 6(7)-locular, with sub-basal placentation; ovules 2 per locule, contained within translucent ovule sac, 0.2 mm long, equal in length to funicle, style similar to style type B; style apex concave; stigma discoid, unlobed, 0.6 mm diam., 0.1 mm high, covering interior faces of stylar funnel; the androecium truncate, 4–6-sided; thecae oblong to elliptical, 0.1–0.2 mm wide; sterile staminate flowers irregularly shaped, 2.1 mm long, 9 mm wide, the margins blunt. JUVENILE plants with internodes 5–6 cm long, 8 mm diam., dark green; petioles terete, sheathing to midway.

Flowering phenology in *Philodendron clewellii* is unclear because there are too few fertile collections. It has been collected in flower in December

and post-anthesis in June, giving a hint of a bimodal flowering. Though Cerro Pirre is not markedly seasonal, these two months in general would mark extremes, with December the general beginning of the dry season and June the beginning of the rainy season in Panama.

*Philodendron clewellii* is endemic to Panama, known only from the Serranía de Pirre, from 850 to 1560 m elevation in Premontane wet forest.

*Philodendron clewellii* is a member of *P.* sect. *Calostigma* subsect. *Oligocarpidium*. The species is distinguished by its thick, smooth stems; usually long internodes; sharply two-ribbed, deciduous cataphylls; subterete petioles; ovate-cordate, dark-brown-drying blades with free or weakly fused basal veins and narrow, closed sinus with often overlapping posterior lobes; and especially by its clusters of up to six small inflorescences per axil with long peduncles, and spathe purple on the outside of the tube and cream on the blade.

In fertile condition, *Philodendron clewellii* does not resemble any other described species, but the leaf blades are similar to those of *P. tysonii* Croat. That species differs in having no more than three, much larger inflorescences at anthesis (to 12 cm or more long). In addition, the leaves of *P. tysonii* usually dry darker. In the blade shape and lack of a naked posterior rib, it is similar to *P. grayumii* Croat. That species differs in having generally more reddish brown blades with a broader sinus and much stouter peduncles.

The species is named in honor of Andrew Clewell, who made several expeditions to Panama while working for Florida State University and who helped collect the type specimen.

*Additional specimens examined.* PANAMA. Darién: Cerro Pirre region, 1000–1400 m, Gentry & Clewell 7028 (K, MO); Río Cans-Río Escucha Ruido, Croat 37831 (MO); ca. 12 km N of Alto de Nique, 1520–1560 m, Croat 37915 (F, MO); ca. 16 km N of Alto de Nique, 1530–1550 m, Croat 37946 (MO); middle slopes on W side, 800–1050 m, 7°56'N, 77°45'W, Croat 68957 (MO).

***Philodendron coloradense* Croat, sp. nov.**

TYPE: Panama. Chiriquí: Cerro Colorado, along mining road 31.6 km beyond bridge over Río San Félix (10.6 km beyond the turnoff to Escopeta), 1690 m, ca. 8°30'N, 81°45'W, 15 July 1976, Croat 37168 (holotype, MO-2395435). Figures 103–106.

Planta hemiepiphytica; internodia usque 10 cm longa, 3 cm diam., sparsim rimosa; cataphylla usque 20 cm longa, incostata, viridia, decidua; petioli subteretes, 35–37 cm longus, 1.5 cm diam.; lamina ovato-cordata, 26–33 cm longa, 17.5–25 cm lata; inflorescentia 2–3; pedunculus 8

cm longus, ad angulum ca. 140° infra spatham flexus; spathe usque 10 cm longa, in tubo viridis, in lamina extus viridialba, intus viridis; pistilla 4–5-locularia; loculi 4–7-ovulati.

Hemiepiphytic; appressed-climbing; internodes smooth, sparsely cracked but only weakly or not at all ribbed, somewhat flattened on one side (at least at the upper nodes), to 10 cm long, 3 cm diam., medium green, soon turning gray, drying light brown; cataphylls to 20 cm long, unribbed, green, deciduous; petioles 35–37 cm long, 1.5 cm diam., subterete, firm, drying reddish brown, obtusely flattened adaxially, surface drying finely and irregularly striate; sheathing at base, 4–5 cm long; blades ovate-cordate, subcoriaceous, semiglossy, weakly bicolorous, acuminate at apex, prominently cordate at base, 26–33 cm long, 17.5–25 cm wide, 1.3–1.6 times longer than broad, upper surface drying dark brown, lower surface drying dark yellow-brown; anterior lobe 20–23.5 cm long, margins convex; posterior lobes 9–12 cm long, directed somewhat upward at an angle to the midrib, drying directed toward the base; sinus obovate, 6.5–9 cm deep; midrib flat, pale green above, convex, paler than surface below; basal veins 4–6 per side, and with the first free to base, third and fourth veins coalesced 1.5–3.5 cm, pale green; posterior rib not at all naked or naked up to 2 cm, only weakly curved; primary lateral veins 4–5 per side, departing midrib at a 45–55° angle, weakly curved to the margins, pale green, weakly sunken above, weakly raised below, drying paler than surface, flattened, with acute margins below; interprimary veins only occasionally present; minor veins easily visible on both surfaces, arising from both the midrib and primary lateral veins, drying weak and paler than surface and usually alternating with blackened secretory ducts, surface minutely granular upon magnification. INFLORESCENCES 2–3 per axil; peduncle to 8 cm long, drying dark reddish brown, minutely striate, bent at ca. 140° angle beneath the spathe; spathe to 10 cm long, to 5 cm wide when flattened, drying dark reddish brown throughout within, spathe blade pale green outside, drying reddish brown with prominent resin canals extending from the blade well into the tube within; spathe tube green, finely striate outside; spadix bluntly pointed at apex, 9 cm long; pistillate portion 2.5–2.7 cm long in front, 1–1.1 cm long in back, 10 mm diam. at apex, 9 mm diam. at middle; staminate portion 8.2 cm long; fertile staminate portion 1 cm diam. at base, 1.2 cm diam. at middle, 9 mm diam. ca. 1 cm from apex, sterile staminate portion 9 mm diam.; pistils 2.5 mm long, 1.4 mm diam.; ovary 4–5-locular, with axile placentation; ovules 4–7 per

locule, 2-seriate, 0.2 mm long, contained within transparent matrix; funicle 0.1–0.2 mm long, adnate to lower part of partition, style similar to style type B; style apex flat; stigma discoid, covering most of style apex except for center, drying irregularly 5-lobed, 0.9–1.3 mm diam.; the androecium truncate, weakly oblong, probably prismatic, margins irregularly 4–5-sided, mostly 4-sided, 1.3–1.6 mm diam. at apex; sterile staminate flowers irregularly 4–6-sided, prismatic, 1.4 mm wide. INFLORESCENCE with seeds 5–7 per locule, 0.4–0.5 mm long.

Flowering in *Philodendron coloradense* is documented by a single collection made in July, but the plant has three inflorescences, probably all of which opened after the onset of the rainy season in May. The region where the collection was made, though at a high elevation, is on the western slope of the Continental Divide, usually much affected by the dry season.

*Philodendron coloradense* is endemic to Panama, known only from near the Continental Divide at Cerro Colorado in Chiriquí Province at 1600 m elevation.

*Philodendron coloradense* is a member of *P.* sect. *Calostigma* subsect. *Macrobelyum* ser. *Macrobelyum*. It is characterized by having grayish internodes longer than broad; green, unribbed, deciduous cataphylls; obtusely flattened petioles; ovate-cordate reddish-brown-drying blades with a spatulate sinus and a weakly developed posterior rib that is barely or not at all naked; and paired, green inflorescences bent somewhat at the apex of the peduncle.

The species is perhaps closest to *P. grayumii*, which differs in having 7–8 (vs. 4–5) locules per ovary and 3–4 (vs. 5–7) ovules per locule. In addition, the lower leaf surface of *P. grayumii* dries glossy and is epunctate with dense secretory ducts at higher magnifications. In contrast, the blades of *P. coloradense* dry more or less matte on the lower surface and are minutely speckled with only sparse secretory ducts at higher magnification.

*Additional specimen examined.* PANAMA, Chiriquí: Cerro Colorado, along mining road 20.5 mi. N of bridge over Río San Félix, 8.3 mi. beyond Chame and turnoff to Escopeta, 1630 m, Croat 75039 (MEXU, MO, PMA, US).

***Philodendron copense*** Croat, sp. nov. TYPE: Panama, Coclé: Alto Calvario, 6 mi. N of El Copé, Atlantic slope, along Continental Divide, along trail which heads off old lumber road and leads down to Las Ricas, Limón, and San Juan, 710–800 m, 8°39'N, 80°36'W, 22 June 1988, Croat 68765 (holotype, MO–3584056–8; isotypes, AAU, B, CAS, CM, COL, CR, K, PMA, US). Figures 109–111, 113–115.

Planta hemiepiphytica; internodia 1.5–2.5 cm longa; 2–7 cm diam.; cataphylla 7–25 cm longa, plerumque 1-costata, in sicco flavibrunnea vel atribrunnea, persistentia semi-intacta; petiolus acute D-formatus, 53–109 cm longus, 8–20 mm diam., epidermide interdum dense brevi-lineata, in sicco atriflavibrunneus; lamina ovato-triangularis, cordata, 56–84.5 cm longa, 22–47 cm lata, in sicco rubrobrunnea; inflorescentia usque 6; pedunculus 7–7.5 cm longus, 7–15 mm diam.; spatha 13–18.5 cm longa, lamina spathae viridi; tubo spathae viridi, extus suffuso purpureoviolaceo, pallidius secus marginem, intus marronino; pistilla 3–5-locularia; loculi 20–22-ovulati; baccae albae.

Hemiepiphytic; stem appressed-climbing, scandent; leaf scars inconspicuous; internodes short, thick, 1.5–2.5 cm long, 2–7 cm diam., broader than long, drying yellow-brown, epidermis moderately glossy, finely and acutely ridged on younger stems, lighter and more matte on older stems; roots drying brown, 15–75 cm long, 1–5 mm diam., numerous per node, epidermis peeling; cataphylls drying chartaceous, 7–25 cm long, usually 1-ribbed, pale red to reddish brown, drying yellowish brown to dark brown, persisting semi-intact as fibers. LEAVES erect to erect-spreading; petioles 53–109 cm long, 8–20 mm diam., sharply D-shaped, with broad medial rib adaxially, surface sometimes densely short-lineate, drying smooth and matte to semiglossy, light reddish brown or dark yellow-brown, the epidermis smooth and often flaking; blades ovate-triangular, moderately bicolorous to concolorous, abruptly acuminate at apex (the acumens) 1.17–2.5 cm long, cordate at base, 56–84.5 cm long, 22–47 cm wide (1.8–2.7 times longer than wide), (0.77–1.1 times longer than petiole), broadest below point of petiole attachment, margins broadly undulate, occasionally downturned, upper surface dark green, glossy to semiglossy, drying reddish brown, gray-green or olive-green, matte to weakly semiglossy, lower surface drying semiglossy, yellow-brown to reddish yellow-green; anterior lobe 32–64 cm long, 20–46 cm wide (2.6–3.3 times longer than posterior lobes), broadest at or near base, margins concave; posterior lobes 14.5–21.5 cm long, 22–47 cm wide, broadest at or near base, directed downward and inward to downward and outward, obtuse to semirounded; sinus spatulate to widely hippocrepiform; midrib weakly raised to flat above, thicker than broad, paler than surface, drying concolorous or darker than surface below; basal veins 6–10(15) per side, with 1 free to base, numbers 4–7 coalesced 6–9 cm, prominently raised below; posterior rib naked for 3.5–4 cm; primary lateral veins 11–16 per side, departing midrib at a 55–90° angle, straight to weakly arcuate, drying reddish brown, sunken and paler than surface



above, raised, drying slightly darker than surface below; interprimary veins as conspicuous as primary lateral veins; minor veins arising from both the midrib and primary lateral veins; lesser veins drying prominulous; "cross-veins" moderately conspicuous on both surfaces. INFLORESCENCES erect, to 6 per axil; peduncle 7–7.5 cm long, 7–15 mm diam., green, coarsely lineate; spathe coriaceous, 13–18.5 cm long (1.8–2.6 times longer than peduncle), ± constricted midway on the tube, acute at apex; spathe blade green, short pale-lineate toward base, striate near apex outside, 10.5 cm long (opening broadly oblong-ovate in face view); spathe tube oblong-ellipsoid, green, tinged purple-violet, paler along margin outside, 8.5–9 cm long, 3.5 cm diam., maroon, conspicuously short-lineate inside; spadix sessile; oblong, 12.5 cm long, constricted at base of fertile staminate portion and tapering at both ends; pistillate portion white, cylindrical to ellipsoid, 6.5–7.5 cm long, 2.2–2.5 cm diam. throughout; staminate portion creamy white, 9 cm long, 9 mm diam. throughout, broadest in lower one-third; fertile portion 9–11 mm diam.; sterile portion narrower than the pistillate portion, 9–15 mm diam.; pistils 1.9–2.3 mm long, 1.9–2.3 mm diam.; ovary 3–5-locular, drying irregularly angled, walls embedded with granular, crystal-like particles, with axile placentation; ovules 20–22 per locule, style similar to style type B; style apex broadly and shallowly concave; stigma button-like, 0.5–0.7 mm long; sterile staminate flowers 2–2.4 mm long, 1–1.5 mm wide. INFRUCTESCENCE with berries white; seeds 0.8–0.9 mm long, 0.3–0.4 mm diam.

Specimens of *Philodendron copense* hint at bimodal flowering. A single flowering collection in post-anthesis condition was made in June, immature fruits were collected in January, and mature fruits were collected in December. The mature fruits in December could be the result of a flowering in the early rainy season, but immature fruits in January probably were the result of flowering in the late rainy season or early dry season. More flowering specimens are needed for more conclusive statements on phenology.

*Philodendron copense* is endemic to Panama (but should be expected in eastern Costa Rica) and is known from Bocas del Toro and Coclé (El Copé) at 590 to 930 m elevation in *Premontane rain forest*.

*Philodendron copense* is a member of *P.* sect. *Philodendron* subsect. *Platypodium*. This species is characterized by its appressed climbing habit; short, thick internodes (2–7 cm diam.); usually one-ribbed cataphylls which persist semi-intact; D-shaped petioles with flaky, yellow to yellow-brown epi-

dermis upon drying; large, ovate-triangular blades drying reddish brown with many conspicuous "cross-veins"; and up to six inflorescences per axil with the spathes green or reddish tinged outside and maroon within on the tube.

*Philodendron copense* is most easily confused with *P. brunneicaule*, with which it occurs. Though the latter species has blades that dry a similar color, it is distinguished by having long internodes; deciduous cataphylls; petioles that lack the yellow-brown, peeling epidermis; ovate blades that lack the minute "cross-veins" upon drying; and stouter inflorescences that are solitary in each axil.

*Philodendron copense* is also similar to *P. chiriquense*, another species with an ovate-triangular leaf that dries reddish brown. *Philodendron chiriquense* differs by having much shorter, thicker internodes, conspicuous reddish brown persistent fibers (rather than semi-intact), and dark brown to blackened striate petioles, which lack an exfoliating smooth epidermis.

*Additional specimens examined.* PANAMA. Bocas del Toro: Fortuna Dam area, Fortuna-Chiriquí Grande, 4.3 km N of the Continental Divide, 590 m, 8°46'N, 82°14'W, Croat 60159 (CM, DUKE, F, MO, OOM, QC). Coclé: near Continental Divide, along lumber road, ca. 1.5 mi. N of El Copé, 900 m, Croat 44563 (MO, RSA, U); 44593 (COL, MO); 5.2 mi. above El Copé, 930 m, Croat 49156 (L, MO, SAR, US); 5.6 mi. N of El Copé, 800 m, 8°39'N, 80°36'W, Croat 75064 (M, MEXU, MO, NY, SCZ); 75091 (MO); ca. 5.6 mi. N of El Copé, 800 m, 75091 (MO); 9.4 km above El Copé, 750–900 m, 44729 (B, CR, K, MO, PMA, TEX).

***Philodendron correae* Croat, sp. nov. TYPE:**

Panama. Chiriquí: Gualaca-Chiriquí Grande, vic. Lago Fortuna, along trail to meteorological station on Río Hornito departing from N side of highway, ca. 0.5 km S of Centro de Científicos, 8°45'N, 82°18'W, 24 July 1994, Croat & Zhu 76395 (holotype, MO-4619416; isotypes, B, CAS, COL, CR, F, K, NY, PMA, US, VEN). Figures 116–119.

Planta hemiepiphytica aut raro terrestris; internodia obtuse compressata usque latera, usque 7 cm longa, 1.5 m diam.; cataphylla (8)13–16 cm longa, incostata, decidua; petiolus D-formatus, 7–16.5 cm longus, plus minusve spongiosus; lamina plus minusve oblonga, rotunda vel leniter subcordata basi, 21–46 cm longa, 5–12 cm lata; nervis lateralibus 1 4 utroque, leniter visibilibus supra; inflorescentia 1; pedunculus 10.5–12 cm longus; spathe 15–18.4 cm longa, extus flavialba, intus leviter pallidior; interdum viridis basi, tubo intus suffuso obscure rubro; pistilla 5–6-locularia; loculi 1-ovulati.

Hemiepiphytic or rarely terrestrial; stem appressed-climbing or spreading; sap viscous, unscented; internodes obtusely flattened on one side,

moderately glossy, to 7 cm long, to 1.5 cm diam., usually much longer than broad, medium green to tannish, drying gray, epidermis becoming grayish brown and peeling; roots mostly 15–30 cm long, drying 2–3 mm diam.; cataphylls (8)13–16 cm long, unribbed, green, deciduous, turning mushy; **petioles** 7–16.5 cm long, (3)6–15 cm diam., D-shaped, obtusely flattened (drying usually sulcate) adaxially,  $\pm$  spongy, narrowly rounded abaxially, with adaxial margins raised, surface with a conspicuous purple-black ring around apex; sheathing usually to near the apex, with margins erect and incurved, the tip free-ending; **blades**  $\pm$  oblong, subcoriaceous, semiglossy, conspicuously to moderately bicolorous, abruptly long-acuminate or sometimes acute at apex (the acumen inrolled if present), rounded to weakly subcordate at base, 21–46 cm long, 5–12 cm wide (3–5 times wider than long), (ca. 2–4 times longer than petiole); upper surface semiglossy, dark green to medium green, drying dark brown to blackish brown, lower surface glossy and slightly to moderately paler; sinus 5–12 mm deep; midrib flat to broadly convex, paler than surface above, convex or thicker than broad and bluntly acute, darker than surface below; basal veins lacking; primary lateral veins weak, 4–6 per side when present but sometimes not apparent, departing midrib at a 60–70° angle, arcuate to the margins, slightly sunken, weakly visible above, convex, slightly darker than surface, weakly visible below; minor veins fine, moderately visible to distinct, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 1 per axil; peduncle 10.5–12 cm long, 5–6(14) mm diam., subterete; **spathe** coriaceous, 15–18.4 cm long (1.43–1.75 times longer than peduncle), weakly or not at all constricted, spathe blade yellow-white (B & K yellow-red 9/7.5–10 (at anthesis)) throughout, slightly paler inside; spathe tube sometimes green outside, 6.5–7.5 cm long, 1.6–2 cm diam., slightly paler and tinged dull red inside; **spadix** 8–19 cm long; stipitate 7–10 mm long; pistillate portion green to pale yellow, cylindrical, 7.9 cm long, 1.3–1.4 cm diam. throughout; staminate portion 4–8.4 cm long; fertile staminate portion tapered; pistils glossy, 3.3–3.8 mm long, 2.3–2.6 mm diam.; ovary 5–6-locular, with sub-basal placentation; ovules 1 per locule, style similar to style type C; style funnel shallow, sometimes deep subcylindric; style apex rounded, somewhat flat, with simple funnel; stigma  $\pm$  discoid, covering interior faces of stylar funnel. **INFRUDESCENCE** (post-anthesis) with seeds 1 per locule, 1.5–1.7 mm long, 7–8 mm diam.

Flowering in *Philodendron correae* has been recorded only during the early rainy season, June

through September, but relatively few fertile collections have been seen overall.

*Philodendron correae* is endemic to western Panama in Chiriquí and Bocas del Toro Provinces at 780 to 1400 m in *Premontane rain forest* and *Tropical Lower Montane rain forest*.

*Philodendron correae* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is distinguished by its internodes generally longer than broad; more or less oblong, dark brown- to black-drying blades with rounded to weakly cordulate bases; primary lateral veins only weakly visible; and solitary inflorescences with the spathe blade yellowish white on both inner and outer surfaces.

*Philodendron correae* is closest to *P. ligulatum*, which differs in having rather prominent primary lateral veins and by occurring at generally lower elevations (sea level to 900 m). *Folsom et al. 5471* is noteworthy in having blades that dry greener than is usual for the species. It also has a Type B style type; the only other specimen studied for ovules (*Croat 66748*) had a Type C style. The difference may have been due to the age of the flowers, but this situation warrants further investigation.

*Philodendron correae* is named in honor of Mirya D. Correa A., among the first scientists to collect in the Fortuna area, and the first to collect this species.

*Additional specimens examined.* PANAMA. **Bocas del Toro:** Fortuna Dam area, Chiriquí Grande–Fortuna, 1.2 mi. N of Continental Divide, 910 m, 8°44'N, 82°17'W, *Croat 60436* (CAS, L, MO); 6.6 mi. N of middle of bridge over Fortuna Lake, 780 m, 8°45'N, 82°18'W, *Croat 66779* (MO); along gravel road near Continental Divide, 1170 m, 8°44'N, 82°17'W, *Croat 66653* (MO, PMA, US). **Chiriquí:** Cerro Colorado, above San Félix, along mining road 18–27 mi. N of Pan-American Highway, 1200–1500 m, *Croat 33151* (MO); 13–14.6 km N of Chame, 1390–1410 m, *Croat 37217* (MO); Fortuna Dam area, N of Gualaca, 11.8 mi. N of Los Planes de Hornito, 1400 m, *Croat 48676* (MO, US); Gualaca–Chiriquí Grande, at junction of road to IRHE headquarters, 1200 m, 8°45'N, 82°18'W, *Croat 66748* (CM, CR, HMNM, HUA, MO); 1.4 mi. W of Centro de Operaciones, trail to Río Hornito, 1010–1130 m, 8°44'N, 82°14'30"W, *Croat 67917* (AAU, MEXU, MO, NY); Los Planes de Hornito, 1000–1200 m, *Correa 2256* (PMA); ca. 10.7 mi. from Planes de Hornito, 4000 ft., *Antonio 5148* (MEXU, MO); along aqueduct to IRHE water source near dam, 1200–1300 m, 8°45'N, 82°18'W, *Croat 66626* (MO); 1400–1500 m, *Folsom et al. 5471* (MO).

**Philodendron cotobrusense** *Croat & Grayum*, sp. nov. TYPE: Costa Rica. Puntarenas: along road about halfway between Flor del Roble and Las Alturas de Coto Brus, 1250 m, 8°54'N, 82°51'W, 13 July 1985, *Grayum & Hammel 5689* (holotype, MO–3446320; isotypes, CR, K, US). Figure 120.

Planta hemiepiphytica; internodia 1–3 cm longa, 1.5–2 cm diam., cum epidermide argenteocana; cataphylla usque 14 cm longa, acute 2-costata, decidua; petiolus 32.5–35 cm longus; lamina profunde 3-lobata, 37–38 cm longa, 46–52 cm lata, venis confertis; nervis lateralibus 1 18–19 utroque; inflorescentia usque 5; pedunculus 6–9 cm longus; spathe 6–14 cm longa, lamina spathe extus cremeo-flava, intus eburnea; tubo spathe extus viridi, intus rubro-violacio; pistilla 6–7-locularia; loculi 3-ovulati.

Hemiepiphyte; stem appressed-climbing; internodes short, thick, 1–3 cm long, 1.5–2 cm diam., mostly longer than broad, epidermis silvery gray; cataphylls to 14 cm long, sharply 2-ribbed, yellow-green, tinged pinkish, deciduous; **petioles** 32.5–35 cm long, 4–9 mm diam., subterete, firm, obtusely flattened adaxially, surface matte; **blades** deeply 3-lobed, margins incised to within 1.5–2 cm of base, subcoriaceous, weakly bicolorous, long-acuminate at apex (the acumen probably downturned),  $\pm$  cordate at base, 37–39 cm long, 46–52 cm wide (0.7–0.8 times longer than wide), (1–1.2 times longer than petiole), usually about equal to petiole, upper surface dark green, semiglossy, lower surface weakly glossy, somewhat paler; median lobes 30.5–32 cm long, 13–17 cm wide; lateral lobes 24–27 cm long, 9–12.6 cm wide (2.14–2.88 times longer than wide), directed  $\pm$  upward, broadly confluent to 2.5 cm with medial lobes; midrib broadly convex, closely veined above, convex below; basal veins lacking; primary lateral veins 18–19 per side, departing midrib at a 50–65° angle, broadly curved to the margins, sunken above, convex below; interprimary veins almost as conspicuous as primary lateral veins; tertiary veins visible, slightly darker than surface; minor veins arising from both the midrib and primary lateral veins but mostly from midrib in upper half of blade. **INFLORESCENCES** to 5 per axil; peduncle 6–9 cm long, 3–4 mm diam., terete; **spathe** 6–14 cm long (1.8–3.5 times longer than peduncle); spathe blade creamy yellow outside, creamy white inside; spathe tube green outside, red-violet to purplish inside; **spadix** 9.5–12 cm long; pistillate portion yellow-green, 7–10 mm diam.; staminate portion 5–8 cm long; fertile staminate portion white, 8–12 mm diam.; ovary 6–7-locular, with sub-basal placentation; ovule sac 1 mm long; ovules 3 per locule, digitately arranged in series of 2–3, contained within transparent ovule sac, 0.2–0.3 mm long, slightly shorter than funicle; funicle ca. 0.3 mm long (can be pulled free to base), style similar to style type B; style apex flat; stigma subdiscoid, 1 mm diam., 0.2–0.3 mm high, covering entire style apex; the androecium truncate,  $\pm$  oblong, prismatic, irregularly 4–6-sided, 0.7–1.1 mm long; thecae oblong, 0.3 mm wide,  $\pm$  parallel to one another.

Flowering in *Philodendron cotobrusense* has been recorded only during the early rainy season in July.

*Philodendron cotobrusense* is known only from eastern Costa Rica in the Cordillera de Talamanca, at 1000 to 1250 m elevation in a *Premontane wet forest* transition to rain forest life zone.

*Philodendron cotobrusense* is a member of *P.* sect. *Tritomophyllum*. This species is characterized by its appressed-climbing habit; sharply two-ribbed, deciduous cataphylls; short, thick internodes with a silvery-gray epidermis; petioles about equaling the deeply three-lobed, closely veined blades (with more than 16 pairs of primary lateral veins); and especially by the inflorescences in clusters of up to five per leaf axil with the spathe tube green outside, red-violet within, and the blade creamy yellow on the outside and creamy white within.

*Philodendron cotobrusense* is most similar to *P. madronense* Croat. That species differs in having the medial segment even more closely veined (to 18 or more) and not at all confluent with the lateral lobes. In addition, it has solitary inflorescences.

*Philodendron cotobrusense* is also similar to *P. tripartitum*, but that species differs in usually having narrower, oblanceolate to oblanceolate-elliptic blades with more widely spaced primary lateral veins (usually 4–10, rarely to 12), and fewer inflorescences per axil (up to three) with longer peduncles. In addition, *P. cotobrusense* has 3 ovules per locule whereas *P. tripartitum* has 1.

*Davidse* 24202, tentatively placed in *P. tripartitum*, seems to be somewhat intermediate with *P. cotobrusense*. See the discussion of that species.

*Additional specimen examined.* COSTA RICA. **San José:** General Valley, between Canadú and Chiriquí, Rto Chiriquí del Pacífico, 1000 m, 9°27'N, 83°37'W, *Burger & Liesner* 7119 (F).

***Philodendron cotonense*** Croat & Grayum, sp. nov. TYPE: Panama. Chiriquí: vic. of El Hato del Volcán, at Las Lagunas, 3 mi. from Nueva California, 1.7 mi. past new airstrip, 1380 m, 8°46'N, 82°40'W, 17 Sep. 1987, *Croat 67708* (holotype, MO-3584981; isotypes, AAU, B, CAS, CM, COL, CR, DUKE, F, K, L, G, NY, OAM, PMA, QCA, RSA, SAR, TEX, U, US, VEN). Figures 108, 121–123.

Planta plerumque hemiepiphytica; internodia 1–3 cm longa, 2.5–3 cm diam.; cataphylla 18–31 cm longa, incostata vel acute 1-costata, decidua intacta; petiolus subteretes, 17.5–53.5 cm longus, 5–8 mm diam.; lamina anguste ovato-triangularis, 23–45 cm longa, 11.5–20.5 cm lata; inflorescentia 1–2; pedunculus (6.5)10–17.5 cm longus, spathe 11–17 cm longa; lamina spathe extus di-

lute viridi vel cremeoflava vel alba, intus pallide viridi vel eburnea vel alba; tubo spathae extus striviridi vel viridi, intus marronino vel carmesino pallide rubroviolaceo aut atripurpureo; pistilla 5-6-locularia; loculi 4-5-ovulatii; baccae vivide aurantiacae.

Hemiepiphytic vine or sometimes terrestrial; stem appressed-climbing but with ends often divergent, fertile to at least 5 m; leaf scars conspicuous, 1-1.3 cm long, 1.2-1.3 cm wide; internodes terete, semiglossy, sometimes matte, drying irregularly ridged, 1-3 cm long, 2.5-3 cm diam., usually broader than long, green, soon becoming gray to silvery-gray, drying yellowish brown, epidermis sometimes flaking free; roots short, stubby, light-colored at tips; cataphylls 18-31 cm long, unribbed to sharply 1-ribbed, yellow-green to green, promptly deciduous intact; **petioles** 17.5-53.5 cm long (averaging 34.4 cm long), 5-8 mm diam., subterete, weakly spongy, dark green, obtusely flattened, especially toward the apex adaxially, with abaxial margins broadly rounded, surface pale greenish lineate, semiglossy; **blades** narrowly ovate-triangular, subcoriaceous, usually gradually acuminate at apex (the acumen 1-3.5 cm long), prominently lobed at base, 23-45 cm long (averaging 35 cm long), 11.5-20.5 cm wide (averaging 15 cm wide), (0.7-1.4 times as long as petiole), broadest near point of petiole attachment, moderately bicolorous to almost concolorous, upper surface drying dark brown to dark olive-green, semiglossy, lower surface weakly glossy, usually yellow-brown, sometimes yellow-green; anterior lobe 16-31 cm long, margins weakly convex to flat or concave; posterior lobes 6-16.5 cm long, 3-8 cm wide, broadest at or near the middle, sometimes narrower at point of petiole attachment; sinus parabolic to obovate or almost oblong, 5-7.5 cm deep; midrib flat to weakly sunken, concolorous to paler than surface above, convex, paler than surface, often tinged purple (with coloration continuing onto petiole apex) below; basal veins 3-4 per side, the first often free to base (sometimes all free to base), the second and third coalesced 1-2.5(4.5) cm; posterior rib usually not naked except sometimes for up to 5 mm at the base; primary lateral veins (3)4-5(6) per side, departing midrib at a 30-65° angle (70-90° angle near base), weakly curved to the margins, sometimes drying undulate, often prominently downturned at midrib, sometimes tinged purplish, scarcely to weakly sunken above, narrowly raised below; minor veins obscure to clearly visible, arising from both the midrib and primary lateral veins, close and almost indistinguishable from the secretory canals, secretory canals appearing as intermittent lines, usually more prominently downturned at

the midrib and appearing to cross over the minor veins, drying blackened. INFLORESCENCES 1-2 (usually solitary, sometimes 2) per axil; peduncle (6.5)10-17.5 cm long, subterete, pale-striate; **spathe** 11-17 cm long (0.94-1.6 times longer than peduncle), somewhat constricted above the tube; spathe blade light green to cream-yellow to white outside, pale green to cream-white to white inside; spathe tube dark to medium green, sometimes pale striate-lineate outside, 3-5.5 cm long, 1.3-2 cm diam., maroon to crimson pale red-violet or dark purple inside; **spadix** sessile, (6.7)10.5-16.5 cm long; pistillate portion green, cylindrical to ovoid, 2 cm long, 4-7 mm diam.; staminate portion 4.2-7.1 cm long; fertile staminate portion to 5 mm diam. throughout; sterile staminate portion whitish to cream-white, to 5 mm diam.; pistils 0.9 mm long; ovary 5-6-locular, 0.6 mm diam., with sub-basal placentation; ovules 4 or 5 per locule, arranged digitately, <0.1 mm long, longer than funicle; funicle <0.1 mm long, style 0.9 mm long, 0.5 mm diam., similar to style type C; style apex flat, drying with 5-6 depressions alternating with ribs from a raised center; stigma hemispherical, inserted on stylar funnel; the androecium irregularly 3-6-sided; thecae ovate to cylindrical; sterile staminate flowers 0.8 mm long. INFRUCTESCENCE with berries bright orange; seeds drying pale yellow-brown, narrowly ellipsoid, 1.3-1.5 mm long, 0.6-0.7 mm diam., with weak striations, appendages absent. JUVENILE plants with petioles 4.7-7.7 cm long; blades lanceolate (acumen 6.5-10.5 cm long), 1.7-2.5 cm wide.

Flowering in *Philodendron cotonense* has been recorded in both Costa Rica and Panama during June, and post-anthesis or early fruiting collections have been made from July through December (except October). Mature fruits have been collected only in March. However, several post-anthesis collections have been made during March, indicating perhaps a bimodal flowering period. Too few data are available to be certain.

*Philodendron cotonense* is known only from eastern Costa Rica and western Panama in *Premontane wet forest* and *Premontane rain forest*, *Tropical Lower Montane wet forest*, and *Tropical Lower Montane rain forest* life zones, as well as the transition zone between these zones, at 1100 to 1950 m elevation.

*Philodendron cotonense* is a member of *P. sect. Calostigma* subsect. *Glossophyllum* ser. *Ovata*. This species is characterized by having internodes usually longer than broad (except distally), subterete petioles, and more or less triangular blades averaging over twice as long as broad, with posterior

lobes typically much longer than broad and often directed somewhat outward. Also characteristic are the conspicuous secretory canals visible on the dried blade surface and the solitary long-pedunculate inflorescences in each leaf axil.

*Philodendron cotonense* is most easily confused with *P. wilburii* var. *longipedunculatum*, which differs in having longer and more slender internodes (3–8 cm long and 8–15(20) mm diam.); sharply 2-ribbed cataphylls; and more typically hastate leaves ranging from 1.03 to 1.4 (averaging 1.29) times longer than wide, with proportionately narrower lobes directed outward at a broader angle (averaging 91° vs. 121° for *P. cotonense*). *Philodendron wilburii* var. *longipedunculatum* also differs in having mostly two inflorescences per axil with proportionately much longer peduncles, averaging 1.35 times longer than the spathe (vs. rarely longer than the spathe in *P. cotonense*).

*Philodendron cotonense* might be confused with smaller plants of *P. alticola*, which also have secretory canals visible on the blades. The latter species differs in typically having much larger blades (minimum 44 cm × 21 cm) that dry greener and thicker; persistent cataphyll fibers; and a short-pedunculate inflorescence with the spathe usually not at all constricted.

Noteworthy is *Davidse et al.* 28367, which has a narrower sinus and the basal veins all free to the base with the uppermost primary lateral veins more or less obscure. Otherwise, it fits well into *P. cotonense*.

*Additional specimens examined.* COSTA RICA. **Puntarenas:** vic. of San Vito de Java, ca. 1 km S of San Vito, on road to Villa Neily, 1100 m, *Croat 66169* (CR, K, MO, NY, US); 4000 ft., *Croat 32905* (MO); Zona Protectora Las Tablas, San Vito Coto Brus–Sabalito, Finca Neblinas, 1300–1800 m, 8°53'20"N, 82°50'30"W, *Mora et al.* 54 (CR, MO); Flor del Roble–Las Alturas de Coto Brus, ca. 1250 m, 8°54'N, 82°51'W, *Grayum & Hammel 5690* (MO); Cerro Pando, ridges above Río Cotón and Río Negro, 1000–1800 m, 8°55'N, 82°45'W, *Barringer & Gómez 1606* (CR, F); Laguna Zoncho, San Vito, Coto Brus, 1200 m, *Gómez-Laurito 10774* (F); Las Alturas, along Río Cotón, 1340 m, 8°56'30"N, 82°50'W, *Davidse 24093* (CR, MEXU, MO); Cordillera de Talamanca, area around Río Canasta, 9.5 airline km NW of Agua Caliente, between Cerro Frantizias and Cerro Pittier, 1500–1600 m, 9°02'N, 82°59'W, *Davidse et al.* 28367 (MO); sitio Cotón–Mellizas, 1300–1450 m, ca. 8°54'N, 82°46'W, *Davidse et al.* 25562 (CR, K, MO); N of Santa Elena on Filá Cotón, S of Agua Caliente, 1100 m, 8°57'N, 82°56'W, *Davidse et al.* 28236 (COL, CR, MO); N of Alturas, 1400–1500 m, 8°57'N, 82°50'W, *Davidse 24166* (CR, MO, US); around Tres Colinas, 1800–1850 m, 9°07'N, 83°04'W, *Davidse et al.* 25666 (CR, MO); Las Cruces Botanical Garden, Coto Brus, ca. 4 km SE of San Vito, 1150–1200 m, 8°47'30"N, 82°58'W, *Grayum et al.* 8113 (CM, CR, L, MO, QCA); Río Coto Brus, near Cotón, 23 km N of La Unión, *Croat 26692*

(MO); Río Cotón, vic. of first large concrete culvert before Finca Las Alturas at Cotón, ca. 1300 m, *Croat 44349* (MO); Río Piedras Blancas, Cerro Anguciana, Filá Costeña, Filá Cruces, 950–1150 m, 8°49'18"N, 83°11'15"W, *Grayum 10646* (CR, INB, MO, VEN). PANAMA. **Chiriquí:** Volcán–Río Sereno, 7 mi. N of Volcán, 8°50'N, 82°38'W, *Croat 66226* (CM, MO, PMA, US); 13.7 mi. W of Volcán, 1200 m, 8°51'N, 82°43'W, *Croat 66343* (MO); vic. of Boquete, Cerro Pate Macho, upper NE slopes and summit, 1900–2000 m, 8°46'N, 82°25'W, *Croat 66504* (CAS, L, MEXU, MO, PMA).

***Philodendron crassipathum* Croat & Grayum, sp. nov.** TYPE: Panama. Chiriquí: Cerro Colorado, above San Félix along mining road, 18–27 mi. off Pan-American Hwy., above Chame (turnoff to Escopeta), 1200–1500 m, 8°35'N, 81°50'W, 12 Mar. 1976, *Croat 33150* (holotype, MO–2395067; isotypes, B, F, K, PMA, US). Figures 112, 124, 127, 128, 149.

Planta hemiepiphytica, interdum terrestris; internodia 1.5–8 cm longa, (1–1.8)2.5–3.5 cm diam.; cataphylla carnososa, 8.5–24 cm longa, persistentia semi-intacta, denique decidua; petiolos subtteres, obtuse complanatus, 20–31(48) cm longas, 5–19 mm diam.; lamina ovato-cordata, 14–29 cm longa, 11–24 cm lata; inflorescentia 1; pedunculus 3.5–7 cm longas, 1–2.5 cm longas; spatha carnososa, haud constricta, (6.8)10–14 cm lata; lamina spatheae extus rubra extus, rubra vel alba intus; tubo spatheae extus viridi, interdum extus flavido aut aurantiacorebulo, intus violaceopurpureo vel rubro; pistilla 6–7-locularia; loculi 7–10-ovulati; bacca virides, aurantiacentes vel albidae.

Hemiepiphytic, sometimes terrestrial; stem appressed-climbing or scandent, creeping, becoming mat and gray, smooth when fresh, sap watery, spicy-scented, leaf scars conspicuous, 9–20 mm long, 2–3.5 cm wide; internodes drying conspicuously wrinkled, epidermis weakly glossy, 1.5–8 cm long, (1–1.8)2.5–3.5 cm diam., usually broader than long, dark green, epidermis brownish and cracking; roots dark brown, smooth upon drying, elongate, 3–50 cm long, 1–3 mm diam., few per node; cataphylls fleshy, 8.5–24 cm long, sharply 2-ribbed, green to dark green, dark green short-linear, drying yellowish brown to pale green, persisting semi-intact at upper nodes then eventually fibrous then deciduous. LEAVES erect-spreading to spreading; **petioles** 20–31(48) cm long, 5–19 mm diam., subterete, somewhat spongy, sometimes tinged brown near apex, obtusely flattened with obtuse medial rib adaxially, convexly rounded abaxially; **blades** broadly ovate-cordate, very coriaceous, acuminate at apex (the acumen inrolled, 2–5 mm long), cordate at base, 14–29 cm long, 11–24 cm wide (1–1.56 times longer than wide), broadest at or near the middle, margins hyaline, whitish or reddish, tightly curled under when dried, upper surface dark green, glossy, drying semiglossy to

matte, lower surface paler, drying greenish gray to yellowish green to brownish, drying semiglossy to matte; anterior lobe 10–25 cm long, 11–33 cm wide (1.6–2.5 times longer than than posterior lobes); posterior lobes 4.5–10 cm long, 5–16.5 cm wide, obtuse to rounded; sinus spatulate to hippocrepiform; midrib flat to very broadly convex, drying yellowish green, paler than surface above, weakly convexly raised, paler than surface below; basal veins 3–4(6) per side, with 0–1 free to base, most of the remainder coalesced 1–2.5(4.4) cm, flat above, weakly raised below; posterior rib not naked if present; primary lateral veins 4–6 per side, departing midrib at a 40–70° angle,  $\pm$  straight to the margins, flat, paler than surface, drying obscure above, raised and paler below; interprimary veins obscure above, weakly visible below; minor veins etched above, visible and darker than surface below, arising from midrib only. INFLORESCENCES shorter than leaves, 1 per axil; peduncle 3.5–7 cm long, 1–2.5 cm diam. (dried), terete, drying brown; spathe fleshy, (6.8)10–14 cm long, 4.5–10 cm diam., not at all constricted,  $\pm$  ellipsoid, bluntly acute to rounded at apex, the walls to 1 cm or more thick midway; spathe blade red outside, red to white inside; spathe tube green, sometimes yellowish to orange-red outside, violet-purple to red inside; spadix (6.5)9–14 cm long; pistillate portion grayish to golden-yellow, cylindrical to slightly ellipsoid, 1.5–7 cm long, 10–25 mm diam. throughout, with 13–15 flowers visible per spiral; staminate portion 5–9 cm long; fertile staminate portion white with orange-brown droplets, cylindrical, weakly constricted above sterile portion, tapered toward apex, 15–23 mm diam. throughout, broader than pistillate and sterile portions, 22–27 flowers per spiral; sterile staminate portion 17–24 mm diam.; pistils 4–4.5 mm long, 4–4.5 mm diam.; ovary 6–7-locular, with axile placentation; ovules 7–10 per locule, 0.6–1 mm long, 0.3 mm diam.; funicles with free portion ca. 1 mm long, the remainder loosely fused to the wall of the locule; style similar to style type D, button-like and concave on drying, 1.3–2 mm diam., the margins pale; androecium margins irregular, 0.6–1.3 mm long; sterile staminate flowers  $\pm$  globose, 1.4–2.2 mm long. INFRUCTESCENCE erect; pistillate spadix 4.5–5 cm long, 4 cm diam.; berries green becoming orange to whitish; pericarp white; mesocarp yellowish white; seeds yellow-orange, 2.9–3.5 mm long, 0.7–1 mm diam.

Flowering in *Philodendron crassispathum* appears to occur during the dry season and early rainy season based on flowering collections made during February through April and in August. Post-anthesis collections have been made from January

through October. Mature fruiting collections have been made only from January through March, indicating that they may take up to almost one year to develop.

*Philodendron crassispathum* ranges from central Costa Rica to western Panama, at 1100 to 2600 m elevation in *Premontane rain forest* and *Tropical Lower Montane rain forest* life zones.

*Philodendron crassispathum* is a member of *P.* sect. *Calostigma* subsect. *Macrobelyum* ser. *Ecorata*. The species is characterized by its high elevational habitat, fleshy, deciduous, sharply two-ribbed cataphylls, subterete, obtusely flattened, somewhat spongy petioles, coriaceous ovate-cordate blades with the minor veins etched above, and especially by the very fleshy more or less ellipsoid, green spathe with walls typically 1 cm or more thick, colored white within on the blade and red in the tube.

*Philodendron crassispathum* is most closely related to *P. brenesii*, which differs in having narrowly ovate blades with a narrow V-shaped sinus and mostly free basal veins, and a comparatively thin spathe clearly constricted above the tube portion. In contrast, *P. crassispathum* has blades with a typically spatulate to hippocrepiform sinus and a usually obvious posterior rib.

*Additional specimens examined.* COSTA RICA. **Alajuela:** Río San Pedro, Cerro Azahar, 15 km NW of San Ramón by air, 1400–1500 m, 10°9'30"N, 84°34'–35"W, *Liesner et al.* 15502 (CR, K, MEXU, MO, WIS); Monteverde Reserve, Cerro Chomogo, 1600–1690 m, *Dryer 1470* (CR, F). **Cartago:** Río Dos Amigos–Río Villegas, narrow ridge W of Río Grande de Orosi, 1650 m, 9°42'N, 83°47'W, *Grayum et al.* 3762 (MO, PMA, US); 4.5 km past town of Río Macho, along road to Humo, 1633 m, 9°17'N, 83°45'W, *Hoover 1346* (CR, MO); Tapanti Reserve, Quebrada Salto–Río Grande de Orosi, ca. 1 km S of jct., 1500–1800 m, 9°43'N, 83°47'W, *Croat & Grayum 68226* (CR, MO), *Grayum et al.* 6301 (MO); 1380 m, *Croat 79077* (CR, INB, MO); Tapanti Watershed Preserve, 20 mi. SW of Paraiso, *Croat 47045* (CM, MO), 47043 (MO, SCZ). **Heredia:** Río Las Vueltas–Río Nuevo, E slopes of Volcán Barba, 2000 m, 10°6'N, 84°03'W, *Burger & Baker 9500* (CR, F, ISC, MO, NY, PMA); 9 km SE of Virgen del Socorro, 9 km E of Isla Bonita, 1530 m, 10°14'N, 84°05'W, *Loiselle 228* (MO); Cerro Chompipe, N of San Rafael, 2000 m, *Lems s.n.* (F, NY, US); Cerro de las Caricias, N of San Isidro, 2000–2400 m, *Standley & Valerio 51910* (US); San Rafael–Río Las Vueltas, N slope of Cerro Chompipe, 2100–2200 m, 10°05'N, 84°04'W, *Stevens 13989* (MO); Volcán Barva, Cerro Chompipe, near Ermita Santa Cruz, *Lems s.n.* (F, US); Río Las Vueltas, 12 km NE of San Rafael, 2000 m, *Wilbur et al.* 15986 (DUKE); Río Santo Domingo, ca. 3 km E of San Rafael de Vara Blanca, N slope of Volcán Barva, 2060–2100 m, 10°11'N, 84°07'W, *Grayum 7335* (MO). **Limon:** Cantón de Talamanca, Bratsi, Amubri, Alto Lari, Kivot, between Río Lari and Río Dapari, 1900 m, 9°22'45"N, 83°06'15"W, *Herrera 5504* (INB, MO); Cordillera de Tal-

amanca, Cerro Biricuacua, 2600 m, 9°23'55"N, 83°10'10"W, *Herrera 6219* (CR, INB, MO); 9°20'20"N, 83°13'33"W, *Bittner 1864* (INB, MO). **Puntarenas:** Monteverde Reserve, 1500 m, ca. 10°17'N, 84°48'W, *Croat 61195* (MO); Cerro Negro, 1500–1600 m, *Haber & Bello 2843* (MO); Brillante Trail to Veracruz, river valley along Continental Divide, 1600 m, 10°20'N, 84°50'W, *Haber et al. 4584* (MO). **San José:** S slopes of Cerro Zurquí, ca. 4–4.5 km N of San Isidro de San José, 1500–1800 m, *Uley & Uley 408* (DUKE); 4 km N of Cascajal, 7 km N of Las Nubes, on CR-216, 1500–1600 m, *Uley & Uley 5255* (MO). **PANAMA. Bocas del Toro:** Cerro Colorado, 12 km W of Chame, 3000–4000 ft., *Kress et al. 86-1949* (MO, SEL); 9.2 mi. W of Chame, 1450–1480 m, 8°35'N, 81°50'W, *Croat 69069* (L, MEXU, MO, P, US); near Continental Divide, 9.4 mi. from Chame, ca. 1700 m, ca. 8°35'N, 81°45'W, *McPherson 8918* (CAS, MO); Fortuna Dam area, along trail on Continental Divide, ca. 1200 m, *McPherson 9031* (MO). **Bocas del Toro-Chiriquí:** Chiriquicito-Calderas, Elfin forest, at Divide, on trail, *Kirkbride & Duke 975* (MO); Cerro Colorado, 1300–1400 m, 8°35'N, 81°50'W, *McPherson 13640* (MO, NY). **Chiriquí:** Cerro Colorado, 20 mi. N of Río San Félix, 1660 m, 8°30'N, 81°46'W, *Croat 74987* (MO, US); 1110–1750 m, 8°35'N, 81°54'W, *Hammel & Trainer 14932* (MO); Boquete region, SW slope of Cerro Pate Macho, 1630–1780 m, 8°46'N, 82°25'W, *Croat 66382* (MO, QCA, SCZ); Fortuna Dam area, Fortuna Lake-Chiriquí Grande, 4.5–5 km N of dam over Fortuna Lake, 1100–1134 m, 8°43'N, 82°17'W, *Croat & Grayum 59979* (AAU, B, CM, GH, K, MO, RSA); Gualaca-Chiriquí Grande, 4.8 mi. beyond IRHE facilities at Dam, 4 mi. N of bridge over Bayano Lake, 8°46'N, 82°16'W, *Croat 68027* (COL, G, MEXU, MO).

***Philodendron cretosum*** Croat & Grayum, sp. nov. TYPE: Panama. Cooclé: vicinity El Valle de Antón, at forested flat area near Finca Macarenita at La Mesa, 800 m, 8°36'N, 80°07'W, 6 July 1994, *Croat & Zhu 76661* (holotype, MO-04619350-51; isotypes, AAU, B, CAS, CM, COL, CR, F, K, L, M, MEXU, NY, OOM, P, PMA, QCA, SEL, TEX, US, VEN). Figures 125, 126, 144.

Planta epiphytica aut hemiepiphytica; intermedia brevis, 1–3 cm diam.; succus calcareus, in sicco mox calcareus; cataphylla 23–42 cm longa, obtuse 1-costata, in sicco pallide flavibrunnea, persistentia ut fibrae pallidae; petiolus obtuse et late aut profunde et anguste sulcatus adaxialiter, 10.5–23.5 cm longus, 3–6 mm diam.; lamina linearis vel oblanceolata, 29–84.5 cm longa, 5–12 cm lata; inflorescentia 1–3; pedunculus (2.5)4–8.9 cm longus, ca. 10 mm diam.; spatula 9–13.5 cm longa; lamina spatulae extus pallide viridi vel viridialba, intus viridialba; tubo spatulae extus viridi, intus albis; pistilla 4-locularia; loculi cum 18–20-seminibus.

Epiphytic or hemiepiphytic; stem appressed-climbing, elongate, sap chalk-white; internodes short, 4–9 cm long on lower stems, short on flowering plants, 1–3 cm diam., as broad as long or slightly longer than broad, gray-green; roots drying dark brown, few per node; cataphylls thin, mem-

branous, 23–42 cm long, bluntly 1-ribbed, green, drying pale yellowish brown, persisting as pale fibers; **petioles** 10.5–23.5 cm long, 3–6 mm diam., medium green, semiglossy, bluntly and broadly or deeply and narrowly sulcate adaxially, with adaxial margins obtuse or sometimes acute; **blades** linear to oblanceolate, chartaceous to weakly subcoriaceous, weakly bicolorous, drying greenish to yellowish brown on both surfaces, matte and paler below, acuminate to weakly acute at apex (the acumen apiculate), attenuate at base, 29–84.5 cm long, 5–12 cm wide (5.1–8.3 times longer than wide), (2.6–4.7 times longer than petiole), much longer than petioles, broadest in upper one-third; midrib prominently and narrowly sunken above, thicker than broad, bluntly acute, glossy and darker than surface below; basal veins lacking; primary lateral veins 6–9 per side, departing midrib at a 25–45° angle, straight to weakly arcuate to the margins, sunken above, convex below; interprimary veins sunken above, slightly raised below; minor veins arising from midrib only; tertiary veins visible, darker than surface below. **INFLORESCENCES** 1–3 per axil; peduncle (2.5)4–8.9 cm long, ca. 10 mm diam., drying 3–5 mm diam., shorter than petiole, subterete or irregularly angled, white-striate; **spathe** subcoriaceous, semiglossy, 9–13.5 cm long (1.8–3.3 times longer than peduncle), moderately constricted midway or just below the middle, abruptly acuminate at apex; spathe blade lanceolate, pale green to greenish white outside, 5.5–8 cm long, greenish white inside; spathe tube ellipsoid, green, short white lineate outside, 3.5–5.5 cm long, 1.8–2 cm diam., white (at anthesis) inside; **spadix** stipitate to 5 mm long; 11–15 cm long; pistillate portion white to pale greenish, weakly ellipsoid, 3–3.5 cm long; staminate portion 4.5–7.8 cm long; fertile staminate portion white; sterile staminate portion 1.2 cm diam.; pistils 0.6–0.9 mm long, 3 mm diam., drying blackish; ovary 4-locular, with axile placentation; ovules ca. 20 per locule, 2–3-seriate, 0.3–0.4 mm long; funicle ca. 0.4 mm long, adnate to lower part of partition, style similar to style type D; style apex flat, with a small style boss; stigma hemispherical; the androecium oblong, 3–5-sided, 0.8–1.2 mm long, 0.4–0.6 mm diam. at apex; sterile staminate flowers irregularly 4–6-sided, 1.5–2.1 mm long, 0.7–1 mm wide. **INFRUESCENCE** 6 cm long, 2.5 cm diam.; seeds 18–20 per locule, obovoid to weakly ellipsoid, 0.8–1 mm long, 0.4–0.5 mm diam.

Flowering in *Philodendron cretosum* occurs in both Costa Rica and Panama from March through July (except May), mid-dry season to early rainy season. It probably flowers over a broader period

because post-anthesis collections have been made in January, April, June, July, and November. The post-anthesis collection in January means that the species may have flowered later than July or that there is some bimodality in the flowering. Immature fruits have been collected in November.

*Philodendron cretosum* ranges from Costa Rica to Panama, from near sea level to 900 m elevation in *Tropical wet forest* and *Premontane rain forest* life zones. The species is perhaps restricted to the Atlantic drainage but has been collected principally from areas along the Continental Divide in Panama.

*Philodendron cretosum* is a member of *P.* sect. *Philodendron* subsect. *Canniphyllum*. This species is recognized by its short internodes, white chalky sap (hence the name "cretosum," meaning "full of chalk"), persistent, thin, pale cataphyll fibers, bluntly sulcate petioles much shorter than the blades, and especially its linear to oblanceolate blades. Few other species in Central America have white chalky sap, and no other species with white sap have narrow, non-cordate blades.

*Philodendron cretosum* is not easily confused with any other species in Central America. It resembles *P. tenuipes* Engl. from Ecuador, which also has elongated blades with acute bases and persistent cataphyll fibers, but that species differs in having blades drying darker brown with the primary lateral veins closer and more numerous (more than 15 pairs) and smaller spathes (usually less than 7.5 cm long).

One collection, *Gómez et al.* 19121, reportedly from the slopes of Volcán Miravalles along the Guanacaste-Alajuela border at about 1500 m, would be well above the confirmed elevational range and in a *Lower Montane rain forest* life zone. It is doubtful if it was collected in this life zone or at that elevation. Neither Mike Grayum nor I have encountered this species above 600 m in Costa Rica.

*Additional specimens examined.* COSTA RICA. Alajuela: Reserva Biológica Monteverde Río Peñas Blancas, 500–800 m, *Bello 1070* (CR, MO); 950 m, *1092* (INB, MO); Naranjo-Aguas Zarcas, along Highway 15, 8.5 km NE of Villa Quesada, 600 m, *Croat 46978* (MO); Cañas-Upala, 4 km NNE of Bijagua, Río Zapote, 400 m, *Croat 36295* (MO); San Ramón, *Bittner & G. Herrera 2129* (CR). Guanacaste: Parque Nacional Guanacaste, Estación Pituilla, 550 m, 11°02'N, 85°25'20"W, *Hammel et al.* 17497 (CR, MO); slopes of Miravalles, above Bijagua, *Gómez et al.* 19121 (F, MO, US). Heredia: Cerros Sardinal, ca. 2–2.5 km N of Chilamate de Sarapiquí, 80–160 m, 10°28'N, 84°04'W, *Grayum et al.* 6148 (MO); La Selva Field Station, 50–100 m, 10°26'N, 84°01'W, *Grayum 1894* (MO), 8546 (MO); *Croat 61212* (MO). Limón: Barra de Colorado Refugio, Río Chirripico-Río Sardina, 10–15 m, 10°38'N, 83°45'W, *Grayum 9022* (CR, MO); Río

Segundo, Asunción, Estribaciones Fila Matama, Cerro Matama, 300 m, *Gómez et al.* 23424 (MO). PANAMA. Coelí: 9.4 km above El Copé, 750–900 m, *Croat 44743* (MO); El Valle region, La Mesa, above El Valle de Antón, 860–900 m, 8°38'N, 80°09'W, *Croat 37344* (F, MO); 800–900 m, 67123 (F, MO, US); 13344 (MO); 775 m, *Croat 74782* (M, MO); 900–1000 m, 8°40'N, 80°07'W, *Knapp 5799* (MO, NY). Colón: Nuevo Tonosí-Río Indio, ca. 0 m, *Croat 33555* (MO). Panamá: Cerro Campana, 400 m, *Hutchison & Dressler 2974* (BH, F, M, US); *Croat 17164* (F, MO, PMA); 150 m, 35965 (MO); 780–875 m, 25254 (MO); 800 m, 8°41'N, 79°56'W, 74775 (MO); Cerro Jefe region, 3–3.5 mi. NE of Altos de Pacora, 700–750 m, 9°15'N, 79°25'W, *Croat 68697* (MO); along road to summit, 750–800 m, 9°14'N, 79°22'W, *Croat 67090* (MO); El Llano-Carri, 6.8 mi. from the highway, 350 m, *Croat 49123* (MO); Campamento de los guardabosques de INRENARE, 800–900 m, 8°40'N, 79°55'W, *Correa et al.* 9516 (STRI). San Blas: Nusagandi, km 19.1, *de Nevers & Herrera 7950* (MO). Veraguas: Santa Fe region, Escuela Agrícola Alto Piedra-Río Dos Bocas, 5–8 km from school, 730–770 m, *Croat 25910A* (MO).

***Philodendron davidsonii*** Croat, *Aroideana* 6: 39–41. 1983. TYPE: Costa Rica. Limón: N of Siquirres, originally collected by Jim Tally of Miami, Florida, *Croat 52232A* (holotype, MO-3000000–1; isotypes, B, COL, CR, K, MEXU, NY, US). Figures 10, 129, 130, 136.

Epiphytic climber, often occurring high in canopy; sap clear; stem appressed-climbing, thick, creeping; leaf scars conspicuous, 1.5–2 cm long, 1–3 cm wide; internodes short, thick, broader than long, 1.5–5 cm long, 3–9 cm diam., olive-green to gray-green, becoming tannish white to brownish, scurfy; epidermis peeling and cracking with age; roots few per node, to 7 mm diam., light reddish brown, drying dark reddish brown, smooth, epidermis peeling; cataphylls soft, 28–54 cm long, unribbed to bluntly ribbed to sharply 2-ribbed near apex, green to reddish, sometimes dark green-striate, promptly deciduous; petioles 50–85 cm long, 1–2 cm diam., thicker than broad, obtusely V-shaped, broadly sulcate adaxially, rounded abaxially, with adaxial margins sharp, surface medium to dark green-striate with a weak maroon to dark green ring around apex; blades ovate-oblong, coriaceous, acute to ± acuminate at apex (the acumen inrolled), subcordate to cordate at base, 50–76 cm long, 21–43 cm wide (1.7–2.6 times longer than wide), upper surface dark green, semi-glossy to glossy, lower surface matte, much paler (often tinged purplish violet when young); margins hyaline to yellowish green, sometimes revolute; anterior lobe 48–77.5 cm long, 20–50 cm wide (3.4–6.3 times longer than posterior lobes); posterior lobes 9–19 cm long, 9.6–17.8(24) cm wide, rounded; midrib flat to weakly raised, yellowish green to cream, paler than surface above, convex, weakly striate, pal-



er than surface below; basal veins (1)3-6, mostly 4 per side, with 1-2 (or sometimes all) free to base, 4th to 6th veins coalesced 3-4 cm; posterior rib 0.8-4 cm long, not at all naked or naked to 4 cm; primary lateral veins (10)18-21 per side, departing midrib at a 50-75° angle, straight or weakly arcuate to the margins, sunken to weakly quilted and paler than surface above, convex to raised and paler than surface below; interprimary veins weakly sunken and paler than surface above, weakly raised and darker than surface below; minor veins darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES ± erect, 1-3 per axil; peduncle (3.5)5-9 cm long, 7-27 mm diam., flattened adaxially, pale green to reddish green, or white, often dark green-striate; **spathe** coriaceous, 15-27 cm long ((2.1)2.9-4.6 times longer than peduncle), weakly constricted near the middle (opening 16.5 cm long); spathe blade at anthesis pale green, pale speckled, with margins reddish pink outside, 10-19 cm long, 2-6 cm diam. (the opening to 9 cm long), tinged reddish to maroon (B & K red-purple 2/7) inside; spathe tube pale green, minutely white-short-lineate outside, 5.9-14 cm long, 3-5.5 cm diam., reddish to maroon (B & K red-purple 10/3 to 2/10) inside; **spadix** sessile or weakly stipitate; tapered, 14.5-25 cm long, tapered, broadest near the base; pistillate portion green to yellow-green, slightly tapered toward the apex to cylindrical, 3.3-10 cm long, 1.1-2.5 cm diam. at apex, 1.4-1.7 cm diam. at middle, 1.5-3 cm wide at base, with 19-25 flowers per spiral; staminate portion 9.2-20 cm long; fertile staminate portion creamy white, becoming orange-red (post-anthesis), tapered, 1.6-2.8 cm diam. at base, 1.5-2 cm diam. at middle, 7-12 mm diam. ca. 1 cm from apex, broadest at the base, narrower than the pistillate or sterile portions; sterile staminate portion narrower than the pistillate portion, white, 17-28 mm diam.; pistils 3.3-4.7 mm long, 1.6-2.5 mm diam.; ovary 8-14-locular, 3.7-4 mm long, 2-2.6 mm diam., with sub-basal placentation; ovules 4-8(12) per locule, striate, 2-seriate, usually contained within translucent ovule sac, 0.2-0.6 mm long, about equal in length or shorter than funicle; funicle 0.2 mm long, style 1.2-2.3 mm long, 1.6-2.1 mm diam., similar to style type B or D; style apex usually flat; style apex flat or with a broad boss, pronounced (with divot in center); stigma subdiscoid to globular, unlobed to weakly lobed, 1.8-1.9 mm diam., 0.3-0.7 mm high, covering entire style apex, depressed medially; the androecium truncate, margins 4-6-sided; thecae elliptical to oblong-obovate, 0.1 mm wide; sterile staminate flowers blunt, 4-5-sided, 3.5-4.5 mm long, 1.5 mm wide. Berries greenish (nearly ripe).

*Philodendron davidsonii* ranges from Costa Rica (no doubt into adjacent southeastern Nicaragua as well) to Panama on the Atlantic slope, from near sea level to about 200 m elevation in *Premontane wet forest* and *Tropical wet forest* life zones.

*Philodendron davidsonii* is a member of *P.* sect. *Calostigma* subsect. *Macrobelum* ser. *Ecordata*. This species is characterized by its generally short-creeping habit often high in the canopy (reportedly to 30 m); short, thick internodes; sharply two-ribbed cataphylls, which are deciduous intact; long, stout, broadly sulcate petioles; large ovate-oblong blades with many sunken primary lateral veins; and one to three inflorescences per axil with the coriaceous spathes green outside and reddish to maroon within.

*Philodendron davidsonii* has two subspecies; the typical subspecies is found throughout the range of the species, and *P. davidsonii* subsp. *bocatoranum* is known only from the type locality in Bocas del Toro Province.

KEY TO THE SUBSPECIES OF *PHILODENDRON DAVIDSONII*

- Base of blade cordate to subcordate; primary lateral veins 16-21 per side, arising at 65-75° angle; Costa Rica to Panama, 0-100 m elevation ..... subsp. *davidsonii*
- Base of blade acute; primary lateral veins 10-16 per side, arising at 50-55° angle; Panama (Bocas del Toro), 190-220 m elevation ..... subsp. *bocatoranum*

*Philodendron davidsonii* Croat subsp. *davidsonii*

Internodes 1-5 cm long, 3-8 cm diam.; cataphylls 38-54 cm long, sharply 2-ribbed; **petioles** 56-85 cm long, obtusely V-shaped, broadly sulcate adaxially, with adaxial margins sharp, with a dark ring at apex; blades 49.5-67 cm long, 21-43 cm wide; basal veins 3-6, mostly 4 per side, with 1-2 free to base, 4th to 6th veins coalesced 3-4 cm; posterior rib 0.8-4 cm long, not at all naked to 4 cm; primary lateral veins (16)18-21 per side, departing midrib at 65-75° angle. INFLORESCENCES 1-3 per axil; peduncle (3.5)5-9 cm long, flowers with style similar to style type D; style apex usually flat with a broad style boss bearing a medial divot; stigma subdiscoid to globular, weakly lobed. Berries greenish, nearly ripe.

Flowering in *Philodendron davidsonii* subsp. *davidsonii* occurs in the late dry season and early wet season with post-anthesis collections from March through August (except April) and immature fruiting collections from June and November. Cultivated collections at the Missouri Botanical Garden flowered in April, May, June, and July.

*Philodendron davidsonii* subsp. *davidsonii* ranges from northeastern Costa Rica (no doubt into southeastern Nicaragua as well) to Panama on the Atlantic slope, from near sea level to about 100 m elevation in *Premontane wet forest* and *Tropical wet forest* life zones. No specimens have been seen in Costa Rica above 100 m elevation.

*Additional specimens examined for P. davidsonii subsp. davidsonii.* COSTA RICA. **Heredia:** La Selva Field Station, ca. 100 m, *Jacobs* 2708 (MO), *Grayum* 2931 (MO), 2829 (MO). **Límón:** Hacienda Tapezco-Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, *Davidson* 3956 (CM, F, MO); 7097 (L, MO, RSA), 8782 (MO); Manzanillo de Talamanca, ca. 5 m, 9°38'N, 82°39'W, *Grayum & Burton* 4326 (MO); Refugio Barra del Colorado, between Río Chirripocito and Río Sardina, 12 m, 10°38'N, 83°45'W, *Grayum* 9807 (CR, MO). PANAMA. **Bocas del Toro:** Fortuna Dam-Chiriquí Grande, 10 mi. below the divide off highway 1.2 mi., 300–400 ft., *Kress et al.* 86-1992 (MO, SEL).

***Philodendron davidsonii* subsp. *bocatoranum***

Croat, subsp. nov. TYPE: Panama. Bocas del Toro: Ojo de Agua, 7 km W of Almirante, 190–220 m, 9°16'N, 82°28'W, 4 Aug. 1976, *Croat* 56853 (holotype, MO–3636098–99). Type live at MO. Type plant is a re-collection of a sterile live collection vouchered as *Croat* 38177 on 8 July 1983. Figures 19, 131–135.

Internodia 4–9 cm diam.; cataphylla obtuse vel acute 2-costata, persistentia intacta nodis superioribus, tum decidua aut semi-intacta basi; petiolus complanatus vel acute sulcatus adaxialiter, 50.5–67 cm longus; lamina 58–76.5 cm longa, 25–35 cm lata; inflorescentia 2; pedunculus 5.5–6 cm longus; spathe 23–25 cm longa.

Internodes 1–2 cm long, 4–9 cm diam.; cataphylls 28–45 cm long, ca. 6 cm wide, unribbed to bluntly ribbed to sharply 2-ribbed near apex, persisting semi-intact at base; **petioles** 50.5–67 cm long, flattened to sharply sulcate adaxially with adaxial margins conspicuous and erect, maroon ring at apex weakly apparent; **blades** 58–76.5 cm long, 25–35 cm wide; basal veins 1–4, prominently ascending, all free to base or weakly coalesced at base to 1 cm; primary lateral veins 10–16 per side, departing midrib at 50–55° angle. INFLORESCENCES 2 per axil; peduncle 5.5–6 cm long; **spathe** 23–25 cm long; flowers with style similar to style type B, lacking a boss; style apex flat; stigma subdiscoid, unlobed. Berries not seen.

Flowering in *Philodendron davidsonii* subsp. *bocatoranum* occurs during July (based on only two collections).

*Philodendron davidsonii* subsp. *bocatoranum* is endemic to Panama, known only from the type lo-

cality in Bocas del Toro, at 190 to 220 m elevation in *Premontane wet forest*.

The taxon is distinguished by its thick, short internodes, two-ribbed, deciduous cataphylls, long, sharply sulcate petioles, and ovate-oblong blades acute at the base.

*Philodendron davidsonii* subsp. *bocatoranum* differs from the typical subspecies in having blades acute rather than decidedly lobed at the base and 11 or fewer primary lateral veins (vs. about 16 or more) arising at more acute angles with the midrib (50–55° vs. 65–75°).

*Additional specimen examined.* PANAMA. **Bocas del Toro:** vic. of Ojo de Agua, 7 km W of Almirante, 190–220 m, 9°06'N, 82°28'W, *Croat* 38177 (CAS, HUA, M, MO, SCZ, VEN).

***Philodendron dodsonii* Croat & Grayum, sp. nov.**

TYPE: Ecuador. Pichincha: along Río Blanco across from Villa Hermosa on road 1.9 km N of main Sto. Domingo de Los Colorados–Esmeraldas Highway, departing main highway 25 km NW of Santo Domingo, 410 m, 0°5'S, 79°15'W, 13 Mar. 1992, *Croat* 72982 (holotype, MO–04658574–5; isotypes, AAU, B, COL, R, K, M, MEXU, NY, PMA, QCA, QCNE, US). Figures 137–143, 150.

Planta hemiepiphytica; internodia brevia 2.5–3 cm diam.; cataphylla usque 20–33 cm longa, leniter vel acute 2-costata, interdum incostata, persistentia, demum fibrosa; petiolus subteres, 52–93 cm longus; lamina ovata, 36–87 cm longa, 28–66 cm lata, atrivridis supra, moderate pallidior et nitida infra, in sicco atriflaviviridum et impolita vel leniter nitida; costa postica manifeste nuda; inflorescentia 2–5; pedunculus 5–9.5(14) cm longus, 6–10 mm diam.; spathe 16–18 cm longa; lamina spatheae extus alba, intus rubella; tubo spatheae extus rubropurpureo vel atrirubella vel viridi, intus rubro-purpureo vel atrirubello; pistilla (4)5-locularia; loculi ca. 20-ovulati.

Hemiepiphytic; stem appressed-climbing, leaf scars conspicuous, 1–2.7 cm long, 1.5–3.5 cm wide; internodes short on adults, stout, matte to glaucous, 2.5–3 cm diam., frequently longer than broad on nonflowering plants, gray-green, becoming whitish to grayish with age, epidermis flaking; cataphylls thin, spongy, to 20–33 cm long, weakly to sharply 2-ribbed, sometimes unribbed, light green, dark green-striate, persisting in parchment-like mats, eventually fibrous, rarely deciduous intact, margins clear; **petioles** 52–93 cm long, 9–25 mm diam., subterete to obtusely flattened abaxially, soft, drying black, surface pale, dull whitish-streaked, thinly dark green-striate, drying black; **blades** ovate, subcoriaceous, semiglossy, slightly bicolorous, very short acuminate to ± acute at apex, weakly cordate to sagittate at base, 36–87 cm long.

28–66 cm wide (1.3–1.6 times longer than wide), (0.7–1.2 times longer than petiole), about equal in length to petiole, upper surface dark green, drying dark brown to dark yellow-brown, lower surface moderately paler and glossy, drying yellow-brown and matte to weakly glossy, margins weakly undulate and upturned, hyaline; anterior lobe 31–76 cm long, 39–70 cm wide (1.65–2.9(4) times longer than posterior lobes); posterior lobes 11–29 cm long, 9–31 cm wide, broadly rounded to broadly obtuse; midrib flat to sunken, slightly paler than surface above, convex, concolorous below; basal veins 7–8 per side, and with the first free to base, numbers 3–7 coalesced 4–12 cm; posterior rib prominently naked to 6 cm along the sinus; primary lateral veins (4)7–8 per side, departing midrib at a 50–60° angle, to the margins, sunken and paler than surface above, convex to round-raised, usually darker than surface, sometimes paler than surface near base below; interprimary veins sunken and concolorous above, raised and concolorous below; tertiary veins visible, slightly darker than surface below; minor veins darker than surface, drying smooth below, arising from both the midrib and primary lateral veins. INFLORESCENCES 2–5 per axil; peduncles 5–9.5(14) cm long, 6–10 mm diam., subterete, pale green, white-streaked; **spathe** 16–18 cm long ((1)1.8–3.5 times longer than peduncle), moderately constricted above the tube; spathe blade white outside, reddish inside; spathe tube red-purple to dark reddish (B & K red-purple 3/7.5) outside, red-purple to dark reddish inside, sap mango-scented; **spadix** 13–15(18) cm long; pistillate portion 6 cm long in front, 5 cm long in back, 2.5 cm diam. at middle, 2.2 cm wide at base; staminate portion 11 cm long; fertile staminate portion ca. 1 cm diam.; sterile staminate portion 2.5–3 cm diam.; pistils 4 mm long, 1.7–2.3 mm diam.; ovary (4)5-locular, with axile placentation; ovules ca. 20 per locule, 2-seriate, 0.3–0.4 mm long, slightly longer than funicle; funicle 0.2–0.4 mm long, adnate to lower part of partition, style similar to style type B; central style dome fairly well developed; style apex broadly domed; stigma inserted on entire style apex; the androecium truncate, ± prismatic, oblong, margins irregularly 4–5-sided, 1–2.5 mm long; thecae oblong, 0.3 mm wide, contiguous, ± parallel to one another; sterile staminate flowers clavate, irregularly 3–5-sided, 2.3–3.7 mm wide.

Flowering in *Philodendron dodsonii* probably occurs during the rainy season, June–September, in Central America. It is known to flower only during September there, with post-anthesis collections also made from July through September and mature fruits in March. In South America, flowering col-

lections have been made in June and July, post-anthesis from May through September, immature fruiting from February through November (except May, August, and October), and mature fruiting in September.

*Philodendron dodsonii* is known from Costa Rica and Ecuador and is expected to be found on the Caribbean slope of Panama and probably on the Pacific slope of Colombia. In Costa Rica, it occurs at 240 to 1300 m elevation, principally on the Atlantic slope but also on the Pacific slope on the Fila Costeña in *Tropical wet forest* and especially in *Premontane rain forest* life zones. In Ecuador, this species is known from *Tropical wet forest* life zones at 20 to 1750 m in Pichincha Province.

*Philodendron dodsonii* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is distinguished by its stout, short internodes, weakly two-ribbed cataphylls that persist in parchment-like mats and are eventually fibrous; more or less terete petioles; large ovate, yellowish-brown-drying blades with naked posterior ribs; and short-pedunculate spathes with the tube dark green to reddish outside and red within. Another feature useful for determination is the presence of tiny granulations or protuberances on the lower leaf surface. Juvenile blades often have minute pale dots on the surface.

*Philodendron dodsonii* is most easily confused with *P. dominicalense*, a species with very similar features, which differs in having a single ovule per locule (vs. ca. 20 for *P. dodsonii*) and an outer green spathe tube (vs. red for *P. dodsonii*).

*Philodendron dodsonii* also appears close to *P. schottianum*, but that species differs in having cataphylls drying with a much thicker, yellowish epidermis and coarser fibers; petioles drying yellowish brown (vs. almost black for *P. dodsonii*); blades drying with raised minor veins (vs. essentially smooth for *P. dodsonii*); and inflorescences usually scarcely constricted above the tube.

Another species that might be confused with *P. dodsonii* is *P. grayumii*. It has blades similar in shape, size, and color upon drying but which differ in having the upper surface glossy upon drying and in usually having laticifers visible on the lower surface, and especially in having mostly free basal veins with the posterior rib (when present) rarely naked along the sinus.

A collection near San Vito and the only wild-collected specimen from the Pacific slope in Costa Rica differs in having longer internodes. It perhaps represents another species, but in other respects it matches well with *P. dodsonii*.

This species was recognized as distinct by Sodiro (in herb.), under the name *P. robustum* Sodiro ined., previously published for a different species by Schott (1860).

*Philodendron dodsonii* is named in honor of Calaway Dodson, Director of the Centro Científico Río Palenque and staff member of the Missouri Botanical Garden, who first studied the species for the *Flora of Río Palenque* (Dodson & Gentry, 1978). The species was erroneously included in the above-mentioned work as *P. devansayanum* L. Linden, a poorly known entity purportedly collected in the eastern Andes of Peru. Though the last-mentioned species was poorly described and is represented only by a somewhat stylized painting (but no bona fide herbarium specimens), there is little likelihood that it could be the same as a species occurring on the western slopes of the Andes. The illustration appears to show leaf blades with basically free basal veins and little or no posterior ribs, whereas *P. dodsonii* has well-developed posterior ribs. A collection from Ecuador in Sucumbios (Croat 50317) may prove to be this species. Though the inflorescence is immature its blades match those of *P. dodsonii*. Two other collections, Gentry 71004 from Bolivia (La Paz Province, 1500–1550 m) and Croat 55455 from Colombia (Cundinamarca Dept., 2000 m), may prove to be *P. dodsonii*. If so, the range statement could be affected.

**Additional specimens examined.** COSTA RICA. **Alajuela:** Río Sarapiquí, at bridge on road to Colonia Virgen del Socorro, 830 m, 10°16'N, 84°11'W, Croat 68138 (CR, MO); 17 km NW of San Ramón, 785 m, 10°14'15"N, 84°33'W, Croat 68138 (MO); Río Cataratas, ca. 20 km NW of San Ramón, 850 m, 10°12'30"N, 84°32'W, Grayum et al. 6327 (MO). **Cartago:** Río Reventazón, CATIE, Turrialba, ca. 600 m, 9°54'N, 83°39'W, Grayum et al. 3820 (MO); along Camino Raíz de Hule, SE of Platanillo (Tapi-ri), 1200–1400 m, Croat 36796 (MO, US). **Heredia:** Zona Protectora La Selva, Río Peje–Río Guácimo, along Quebrada Cantarras, 300–400 m, 10°22'N, 84°30'W, Grayum & Jacobs 3592 (CR, MO). **Limón:** Cordillera de Talamanca, along Río Madre de Dios, 240 m, 10°03'N, 83°26'W, Grayum et al. 8662 (MO); Río Blanco, Río Frio-Limón, W of Guápiles, Quebrada Danta, 3 mi. S of main highway, 360 m, 10°12'N, 83°49'W, Croat 68422 (CM, MO). **Puntarenas:** Las Cruces Botanical Garden, 1300 m, Croat 44420 (MO); 44464 (MO); ca. 4 km SE of San Vito, 1150 m, 8°47'30"N, 82°58'W, Grayum 8112 (INB, MO); Las Cruces Botanical Garden–Río Jaba, ca. 3.5 km SE of San Vito de Coto Brus, ca. 1160–1200 m, 8°47'30"N, 82°58'W, Grayum 5980 (MO). **San José:** San Isidro del General–Dominical, SW of San Isidro, 4.8 mi. from Río Pacuare, 1000 m, Croat 35260 (MO); 990–1100 m, Croat & Hannon 79114 (CR, INB, MO); Tarrazú, vic. Hormiguero, 1100–1200 m, Croat 78974 (CR, INB, MO).

**COLOMBIA.** **Valle:** Cali–Buenaventura Highway, 1.2 km E of Cisneros, 220–260 m, Croat 62829 (COL, MO, NY). **ECUADOR.** **Cotopaxi:** Quevedo–Latacunga, 3 km E of El Palmar, 800 m, Dodson & Gentry 10253 (MO); La

Mana, 1 km N of Pucayacu, 750 m, Croat 73276 (MO); 2 km N of Pucayacu near Río San Francisco, 690 m, Croat 57080 (MO, QCA); Río Guapara, ca. 20 km NW El Corazón, 250 m, Sparre 17106 (S), 17164 (S), 17150 (S), 17108 (S), 17145 (S); Río Pilaló, Teneferste, km 52–53, 750–900 m, Dodson & Dodson 12914 (MO). **Cotopaxi–Cañar–Chimborazo–Bolívar:** Bucay, 1000–1250 ft., Camp E-3714 (MO, NY, S), E-3656 (MO, NY, S). **Cañar:** Azuques–El Triunfo Road, 1 km S of La Delicia, 227'S, 79°10'W, Croat 50867 (MO, QCA). **El Oro:** Pitias–Santa Rosa, above El Placer, 400–700 m, Harling et al. 15524 (GB); Machala–Loja Road, 890 m, Croat 50718 (MO, QCA). **Esmeraldas:** Quimindé, Bilsa Biological Station, Montañas de Macho, 35 km W of Quimindé, 5 km W of Sta. Isabela, 400–600 m, 0°21'N, 79°44'W, Pitman & Marsh 1146 (MO, QCA); Vieche, Asplund 16514 (S); Fila de Bilsa, 7 km E of San José de Bilsa, ca. 80 km SW of Esmeraldas, 12 km SE of El Salto on Atacames–Muiste Road, 280 m, 0°37'N, 79°51'W, Gentry et al. 72977 (MO); Río Esmeraldas, opposite Quimindé (Rosa Zárate), Asplund 16361 (S); near San Mateo, 80 m, Croat 55632 (CAS, GH, MO, QCA); Río Lita, near Lita, 600–650 m, Croat 38944 (MO). **Guayas:** Teresita, 2 km W of Bucay, 270 m, Hitchcock 20537 (NY, US); 3 km W of Bucay, 270 m, Hitchcock 20489 (NY, US), 20440 (US); Cordillera Chongon–Colonche, 600 m, 1°48'S, 80°47'N, Cornejo & Bonifaz 5237 (GUAY, MO); Huigra–El Triunfo, 160 m, Croat 61593 (F, MO, QCA). **Imbabura:** 13–15 km E of Lita, 800 m, Croat 38918 (MO); Cachaco, 9 km E of Lita, 630 m, Croat 39000 (MO). **Los Ríos:** Babahoyo–Montalve, Hacienda Clementina, 20 m, Sparre 17966 (S); Centro Científico Río Palenque, 230 m, Croat 38651 (MO); 250 m, 50658 (MO, QCA); 220 m, 0°35'S, 79°12'W, 73803 (MO, QCA, US). **Pichincha:** Reserva ENDESA, Río Cabayales, ca. 700 m, 0°05'N, 79°02'W, Croat & Rodríguez 61503 (CAS, CR, MO, NY, QCA); 61516 (CM, MO, PMA, QCA, US); 710 m, 0°03'N, 79°07'W, Croat 73185 (CM, MO, QCA); Alóag–Santo Domingo, Río Toachi, 850 m, Sparre 17816 (S); Tinalandia, above Río Toachi, 700 m, Croat 55745 (MO), 55738 (MO, QCA, SEL); Nono–Nanegal, 13 km SE of Nanegal, 1440 m, Croat 38895 (MO); Nanegal, Sodiro 13 (G); 35 km NW of Santa Domingo, Río Blanco, ca. 250 m, Gentry 9619 (MO); Quito, Parroquia Nanegal, along Río Umachaca near Hacienda El Carmen, 1250 m, 0°07'–7.5'N, 78°38'W, Webster et al. 28781 (DAV, MO); Reserva Maquipucuna, Hacienda Esparragos–Cerro de Sosa, ca. 6 km airline SE of Nanegal, 1500–1600 m, 0°7'N, 78°38'W, Webster & Boland 27500 (DAV, MO); Cerro Campana, 5–6 km (airline) E of Nanegal, 1750 m, 0°09'N, 78°37'W, Webster et al. 30069 (DAV); La Independencia–Río Caoní, 210 m, Croat 55650 (MEXU, MO, QCA, WIS); Santo Domingo de Los Colorados, Hacienda Zaracay, 500 m, Sparre 15190 (S), 15191 (S), 15192 (S); Rancho Brahman, ca. 10 km NW of Santo Domingo de los Colorados on road to Esmeraldas, 400 m, Sparre 14090 (S); El Paraiso–Saguangal Road, 3 km from El Paraiso, 1500 m, 78°46'W, 0°10'N, Øllgaard et al. 37799 (QCA).

#### *Philodendron dolichophyllum* Croat, sp. nov.

TYPE: Panama, San Blas: Nusagandi, El Llano–Cartí road, 9 mi. N of main highway, Nergan Igar (Nergan Trail), 350 m, 9°20'N, 79°W, 2 July 1994, Croat & Zhu 76569 (holotype; MO–4619523–25; isotypes, B, COL, CR, F, K, NY, PMA, US, VEN). Figures 145–148, 151.

Planta plerumque hemiepiphytica; internodia 4–6 cm longa, 2.5–4 cm diam.; cataphylla usque 20–60 cm longa, acute 2-costata, decidua; petiolas subteres, 28–41 cm longas, 7–20 cm diam.; lamina oblongo-lanceolata, anguste rotunda vel leniter subcordata basi, 53–75 cm longa, 12–15.2 cm lata, in sicco rubrobrunnea; nervis lateralibus 17–20 utroque; inflorescentia 2–3; pedunculus 9–12.5 cm longus, 5–17 mm diam.; spathe 14–20 cm longa; lamina spathe extus viridi vel purpureo, intus albida; tubo spathe extus purpureo aut marronino, intus purpureo vel rubello; pistilla 6–8(9)-locularia; loculi 3–7-ovulati; baccae aurantiacae.

Hemiepiphytic, rarely terrestrial on steep banks; stem appressed-climbing, to 3 m long, elongate, leaf scars conspicuous, 1–1.4 cm long, 1–1.3 cm wide; roots few, dark yellow-green, faintly ridged, becoming reddish brown, semiglossy, to ca. 30 cm long, 3–4 mm diam.; internodes short, semiglossy, somewhat scurfy, 4–6 cm long, 2.5–4 cm diam., as long as broad or slightly longer than broad, dark green, drying brown; cataphylls coriaceous, 20–60 cm long, sharply 2-ribbed, pale greenish white, sometimes with reddish base, drying red-brown, deciduous. LEAVES erect-spreading to spreading, the lowermost somewhat pendent; petioles 28–41 cm long, 7–20 mm diam., subterete, somewhat obtusely flattened adaxially, weakly glossy, moderately spongy, dark green to gray-green with a moderately conspicuous dark ring at apex, drying brownish; blades large oblong-lanceolate, subcoriaceous to weakly coriaceous, acuminate to gradually acuminate at apex, gradually tapered toward base or narrowly rounded to weakly subcordate at base, 53–75 cm long, 12–15.2 cm wide (4.4–4.9 times longer than wide), (1.6–2.3 times longer than petiole), broadest at or near middle, upper surface medium green, moderately bicolorous, drying reddish brown, semiglossy, lower surface semiglossy, paler; midrib raised to slightly convexly raised to broadly convex, paler than surface when fresh, drying darker than surface above, rounded, somewhat thicker than broad to acute, paler than surface below; basal veins lacking; primary lateral veins 17–20 per side, departing midrib at a 70–80° angle, slightly arcuate to margin, weakly sunken to moderately obscure above, weakly raised and paler than surface below; interprimary veins slightly less conspicuous than primary lateral veins; minor veins distinct, prominently visible below upon drying, arising from the midrib only. INFLORESCENCES 2–3 per axil; peduncle 9–12.5 cm long, 5–17 mm diam., medium green, minutely white striate, clearly demarcated from the colored spathe; spathe coriaceous, 14–20 cm long (1.6–1.85 times longer than peduncle), acute at apex; spathe blade green to purplish outside, whitish inside; spathe tube ob-

long-ellipsoid, purple or maroon outside, purple to reddish inside; spadix sessile, (6)9.5–19 cm long, broadest near the base or above the middle; pistillate portion cylindrical to ellipsoid, 3.2–5.8 cm long, 1.1–1.5 cm diam. at apex, 1.1–1.7 cm diam. at middle, 1.1–1.2 cm wide at base, with 15–17 flowers visible per spiral; staminate portion 6–9.8 cm long; fertile staminate portion white, tapered to clavate, 8–11 mm diam. at base, 7–10 mm diam. at middle, 5–18 mm diam. ca. 1 cm from apex, narrower than or as broad as the pistillate portion; sterile staminate portion as broad as the pistillate portion, 8–11 mm diam.; pistils 1.4–3 mm long, 8–25 mm diam.; ovary 6–8(9)-locular, 9–25 mm diam., with sub-basal, sometimes basal placentation; ovule sac 0.5 mm long; ovules 3–7 per locule, contained within transparent ovule sac, 0.1–0.9 mm long, almost equal in length to funicle, style similar to style type B; style apex flat; stigma discoid, unlobed, 0.6–1.3 mm diam., 0.1–0.4 mm high, covering center of style apex; the androecium truncate, irregularly 4–6-sided, 0.4–0.8 mm diam. at apex; thecae ovate or oblong, 0.2 mm wide; sterile staminate flowers irregularly blunt, 1–2 mm long, 1 mm wide. INFRUCTESCENCE with berries orange, narrowly ovoid-ellipsoid, tapered and obscurely beaked at apex, 1.2 mm long, 9 mm diam.; seeds 2–3 per locule, 0.8–2.8 mm long, 0.2–2.5 mm diam.

Flowering and phenology in *Philodendron dolichophyllum* are unclear. Flowering collections have been made in February, June, and August, with post-anthesis collections from March, July, August, and October, and immature fruiting collections from March, June, July, and September.

*Philodendron dolichophyllum* is endemic to Panama (though to be expected in eastern Costa Rica), from 325 to 650 m elevation in *Tropical wet forest* life zones.

*Philodendron dolichophyllum* is a member of *P. sect. Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is distinguished by its appressed-climbing habit with at least the apical internodes short; sharply two-ribbed, deciduous cataphylls; subterete petioles (about half as long as the blade); and especially by its long, slender spreading leaf blades (hence the name “dolichophyllum”), which dry typically reddish brown and have weakly sunken primary lateral veins.

*Philodendron dolichophyllum* is not easily confused with any other species in Central America, but bears a superficial resemblance to *P. pseudauriculatum* with which it may occur. Both species have petioles and blades of comparatively equal

lengths and spathes clearly demarcated from the peduncle. *Philodendron pseudauriculatum* differs in having proportionately much shorter leaves with blades that dry gray-green and have prominently sunken primary lateral veins. In addition, the outer surface of the spathe tube in *P. dolichophyllum* is typically maroon or purple, while the blade is green to purplish. *Philodendron pseudauriculatum* has the spathe tube white to pinkish white at anthesis.

*Philodendron dolichophyllum* might also be confused with *P. auriculatum*, which differs in having proportionately longer petioles and blades drying mostly yellow-green and with a lower blade length/petiole ratio (1.4–1.6 vs. 1.6–2.3 times longer than petiole) than *P. dolichophyllum*. In addition, the spathe of *P. auriculatum* is yellowish green outside and not clearly demarcated from the peduncle (clearly demarcated from peduncle in *P. dolichophyllum*).

*Additional specimens examined.* PANAMA. **Boeas del Toro:** Fortuna Dam area, road to Chiriquí Grande, N of Fortuna Dam, 650–700 m, 8°45'N, 82°15'W, *McPherson 11136* (MO). **Coeló:** El Copé region, Alto Calvario, 200–400 m, 8°45'N, 80°35'W, *Hamilton & Davidge 2638* (MO). **Panamá:** El Llano-Cartí, 10.1 mi. N highway, 325–350 m, 3°20'N, 78°58'W, *Croat 67350* (F, MO, RSA, US); Mile 12, 200–500 m, *Croat 22908* (F, DUKE, MO, NY); Mile 7, 460 m, 9°19'N, 79°59'W, *Croat 75112* (MO); Km 18, 900–1000 ft., 9°16'N, 78°58'W, *Sytama 1068* (MO); ca. 16–18 km N, 400 m, *Fyson & Nee 7359* (L, MO, QCA); El Llano-Cartí, near Nusagandí, 300–400 m, 9°20'N, 79°W, *Hamilton & Stockwell 1075* (MO). **San Blas:** Nusagandí-Cartí, 400 m, 9°18'N, 78°58'W, *McDonagh et al. 126* (BM, MO); El Llano-Cartí, 23–29 km N of Pan-American Hwy, 300–400 m, 9°22'N, 78°69'W, *Knapp 1878* (M, MO, NY); Km 22, 350 m, 9°19'N, 78°55'W, *de Nevers & Herrera 7849* (MO). **Veraguas:** Santa Fe region, Santa Fe-Río San Luis, past Escuela Agrícola Alto de Piedra, at Río Segundo Brazo, 480 m, 8°33'N, 81°08'W, *Croat 66895* (CM, MO, PMA, SAR).

***Philodendron dominicense*** Croat & Grayum, sp. nov. TYPE: Costa Rica. San José: along road between San Isidro del General and Dominical, ca. 4.8 mi. E of Río Pacuar, ca. 1000 m, ca. 9°19'N, 83°46'W, 22 May 1976, *Croat 35268*, (holotype, MO-2395109, MO-2395111). Figures 153, 154.

Planta hemiepiphytica; internodia 2.5–10 cm longa, 2–5 cm diam., cano-viridis, in sicco flavibrunnea; cataphylla mollia, 16–29 cm longa, leniter vel acute 2-cosata, persistente semi-intacta; petioli subteres, 37–74 cm longus, 1.5–2 cm diam.; lamina ovato-cordata, 41–46 cm longa, 19–37.5 cm lata, in sicco atribrunnea; costa postica haud nuda aut usque 2.5 cm; inflorescentia 2; pedunculus 8–9 cm longus; spathe 14–14.5 cm longa, omnino viridis; pistilla 5–6-locularia; loculi 1-ovulati.

Hemiepiphytic; internodes coarsely white-striate beneath each node, somewhat soft, drying semi-

glossy, 2.5–10 cm long, 2–5 cm diam., gray-green, drying yellow-brown, epidermis fissured closely; roots moderately few per node, to ca. 30 cm long, drying 2–3 mm diam., reddish brown, semiglossy, sharply ridged; cataphylls soft, 16–29 cm long, weakly to sharply 2-ribbed, green to pale red, drying thin, brown, persisting semi-intact at upper nodes; **petioles** 37–74 cm long (averaging 51 cm long), 1.5–2 cm diam., subterete, weakly spongy, medium green, weakly flattened near apex adaxially, surface light green streaked; **blades** ovate-cordate, acuminate at apex, prominently lobed at base, 41–46 cm long, 19–37.5 cm wide (1.1–1.7 times longer than wide), upper surface dark green, semiglossy to weakly glossy, drying dark brown, lower surface semiglossy, much paler, drying yellow-brown; posterior lobes rounded, 12.5–15 cm long, about as broad as long; sinus hipocrepiform, 8–11 cm deep; midrib flat to raised, paler than surface above; basal veins 5–6 per side, with the uppermost free to base, second basal vein coalesced no more than 1 cm, (2)3–4 coalesced 2.5–5 cm; posterior rib not at all naked or naked up to 2.5 cm; primary lateral veins 4–6 per side, raised or sunken, darker than surface, drying dark brown below; interprimary veins in part sunken, the remainder flat but visible below; minor veins alternately strongly or weakly visible, with the more prominent veins weakly stitched below, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 2 per axil; peduncle 8–9 cm long, 1–1.5 cm diam., fleshy, drying dark brown; **spathe** 14–14.5 cm long, weakly constricted above the tube, green throughout, narrowly acuminate at apex, drying dark brown outside; spathe tube 5.5–6.5 cm long, 2–2.3 cm diam.; **spadix** sessile; to 11.7 cm long; pistillate portion cylindrical, broadest midway, 1.2–1.3 cm diam., weakly tapered in both directions; staminate portion 8.5–9 cm long; fertile staminate spadix broadest at sterile portion, constricted to 9–10 mm ca. 1.5 cm above base, then clavate upward, bluntly tapered at apex, 10–11 mm diam. in upper one-third, 7–8 mm diam. 1 cm from apex; sterile staminate portion 12–13 mm diam.; pistils 1.6 mm long; ovary 5–6-locular, 0.9 mm diam., with sub-basal placentation; ovules 1 per locule, contained within transparent, gelatinous matrix (no true envelope), 0.3–0.4 mm long, longer than funicle; funicle 0.2–0.3 mm long, with a tuft of trichomes near base, style 0.7 mm diam., similar to style type B; style apex flat to concave; the androecium truncate, ± prismatic, margins mostly irregularly 5-sided, 0.7–1.1 mm long; thecae oblong, 0.4 mm wide, ± parallel to one another, contiguous.

PREADULT leaves with petioles 17–20 cm long, blades to 31 cm long and 15 cm wide.

Flowering phenology of *Philodendron dominicalense* is unclear owing to its rarity, but since it was found in flower in May, during the early wet season, it probably flowers and fruits in the wet season.

*Philodendron dominicalense* is endemic to southwest Costa Rica on the Pacific slope in the vicinity of the type locality along San Isidro–Dominical road, at ca. 1000 m, in *Premontane rain forest*.

*Philodendron dominicalense* is a member of *P. sect. Calostigma* subsect. *Glossophyllum* ser. *Ovata*. This species is distinguished by its appressed epiphytic habit; thick grayish green stems with internodes mostly longer than broad; soft, weakly 2-ribbed, semi-persistent, intact cataphylls; subtetere petioles about as long as the blades; ovate-cordate brown-drying blades with the posterior ribs naked for a short distance to the sinus; and paired short-pedunculate inflorescences with green outer spathe surfaces and one ovule per locule.

*Philodendron dominicalense* is vegetatively almost identical to *P. dodsonii*, but that species differs in having the spathe red-purple to dark reddish outside and ovaries with axile placentation and about 20 ovules per locule (vs. spathe tube green outside, ovaries with basal to sub-basal placentation, and 1 ovule per locule). *Philodendron dodsonii* also differs in usually having persistent fibrous cataphylls, a sunken upper midrib (vs. flat to raised), and posterior ribs usually prominently naked for 6 cm (vs. not at all naked or naked to only 2.5 cm).

This species is also similar to *P. schottianum*, especially in shape and color of the dried blades. The latter species differs in having shorter internodes with thicker cataphylls, which dry yellow and weather promptly into a coarse network of fibers (vs. persisting semi-intact at upper nodes then deciduous).

*Additional specimen examined.* COSTA RICA. San José: San Isidro del General–Dominical, SW of San Isidro, 6 mi. from Río Pacuare, 1000 m, Croat 35454 (MO).

***Philodendron dressleri*** G. S. Bunting, *Ann. Missouri Bot. Gard.* 50: 25. Fig. 2. 1963. TYPE: Mexico. Nayarit: Tepic–San Blas, along Hwy. 54, 14–16 mi. SW of junction with Hwy. 15 (Tepic–Mazatlán), 75 m, 5 Sep. 1961, Moore & Bunting 8688 (holotype, BH; isotypes, K, MO, US). Figures 152, 157–160.

Hemiepiphytic; stem scandent, 3–3.5 cm long, 4–4.6 cm diam., leaf scars conspicuous, 3–3.5 cm long, 4–4.6 cm wide; internodes short, stout, succulent, scurfy, 3–6.5 cm long, 4–7 cm diam., broad-

er than long, pale olive-green to gray-green, becoming grayish white with age, epidermis peeling; roots few per node, brownish, smooth, long, to 7 mm diam.; cataphylls 16–22 cm long, sharply 2-ribbed, drying weakly 2-ribbed, densely short dark striate, sometimes deciduous in dry season; **petioles** 34.5–70 cm long, 1–1.8 cm diam., terete, somewhat spongy, surface dark green-striate, dark green ring around apex; **blades** ovate in outline, deeply incised-lobate, bipinnatifid, weakly bicolorous, matte to semiglossy, acute to weakly obtuse at apex (the acumen apiculate), cordate at base, 30–46.5 cm long, 27–40.5 cm wide (0.99–1.2 times longer than wide), (0.9–1.2 times longer than petiole), broadest near the middle; anterior lobe 20.5–30 cm long with up to 6 segments, 12–19 cm long and each 3–5-lobed; posterior lobes rounded in outline, turned up at an angle to midrib with ca. 5 similarly lobed segments; sinus closed or nearly so; segments pinnatifid, 12.3–27 cm long, divided to within 2–15 cm from the midrib; the interlobed sinus divided 0.4–0.7 the length of the lobe; midrib flat, dark green-striate, paler than surface above, round-raised, pale striate, paler than surface below; basal veins 4 per side, 3–4 coalesced 5–10 mm; posterior rib naked for 2.5 cm; primary lateral veins 5–6 per side, departing midrib at a 45–60° angle, ± straight, eventually branching to the margins, flat to weakly raised above, round-raised and paler than surface below; reticulate veins visible, darker than surface below; minor veins arising from both the midrib and primary lateral veins; tertiary veins sunken and paler than surface above, raised and paler than surface below. **INFLORESCENCES** 1 per axil; peduncle 9–16 cm long, 7–9 mm diam. (dried), green; **spathe** 12–21 cm long (1.3 times longer than peduncle), weakly constricted above the middle, ± obtuse at apex; spathe blade dark green outside, pinkish, with darker punctations inside; spathe tube green outside, to 4.5 cm long, purplish violet to wine-red or crimson inside; **spadix** 10–15 cm long; pistillate portion to 3.5 cm long, 2.2 cm diam.; staminate portion to 14 cm long; fertile staminate portion clavate, to 1.9 cm diam. at base, 2.4 cm diam. ca. 1 cm from apex; sterile staminate portion 2.4 cm diam.; pistils 3.7(7) mm long; ovary 4–5-locular, with axile placentation; ovule sac 2.5 mm long; ovules 3–4 per locule, 2-seriate, contained within translucent envelope, 0.4 mm long, longer than funicle; funicle 0.2–0.3 mm long, adnate to lower part of axillary wall, style similar to style type B; central style dome sometimes present; style apex flat; stigma subdiscoid to hemispheroid, 1.8 mm diam., 0.7 mm high, covering entire style apex; the androecium truncate, ±

prismatic, irregularly 5–6-sided, ca. 1.6 mm long; thecae  $\pm$  oblong, 0.4 mm wide, nearly contiguous,  $\pm$  parallel to one another. INFRUCTESCENCE with spathe 10–13 cm long, pistillate spadix 4–6.5 cm long, 3.5–4 cm diam.; seeds 3–4 per locule, 2 mm long, 0.8 mm diam.

Flowering in *Philodendron dressleri* is probably during the rainy season. Post-anthesis collections are known from July and September with immature fruiting collections known from December and January.

*Philodendron dressleri* is endemic to west-central Mexico from coastal Nayarit including Tres Marias Islands (off the western coast of Mexico) to southern Sinaloa, from sea level to 370 m elevation in "Selva Baja Caducifolia."

*Philodendron dressleri* is a member of *P.* sect. *Polytomium*. This species is characterized by its stout, succulent stems with short internodes, weakly two-ribbed, deciduous cataphylls, terete, somewhat spongy petioles (about as long as the blades), bipinnatifid blades divided about midway to the midrib, and solitary green inflorescences with the spathe tube purplish violet to wine-red within.

*Philodendron dressleri* is the most northerly-ranging species of *Philodendron*, extending almost to the Tropic of Cancer. It is probably most closely related to *P. warszewiczii*, but is also similar to *P. radiatum*, both of which differ in having gradually much more deeply divided (almost to the midrib) blades (vs. pinnatifid 0.4–0.7 the way to the midrib in *P. dressleri*). *Philodendron warszewiczii* ranges from Honduras to western Mexico, but no further north than the state of Jalisco. *Philodendron radiatum* ranges no further north than Chiapas on the Pacific slope.

The species is superficially most similar to *P. radiatum* var. *pseudoradiatum*, which also has blades divided less than halfway to the midrib. However, that taxon differs in comprising more scandent plants with slender stems having internodes longer than broad (2–12  $\times$  1–2.5 cm for *P. radiatum* var. *pseudoradiatum* vs. 3–3.5  $\times$  4–4.6 cm for *P. dressleri*).

For an additional photo of this species see Bunting (1965: 332).

*Additional specimens examined.* MEXICO. Nayarit: near Sangaité, E of San Blas, Philbrick 414 (BH); Tepic-Puerto Vallarta, along Hwy. 200, 33 mi. S of Tepic, 4 mi. N of Las Varas, 370 m, Croat 45360 (CM, MO); 6 mi. S of Mazatán, Dressler & Wirth 2732 (UC, US); W of Jaliscoacán, Dressler 1051 (UC); 5–6 mi. E of San Blas along highway, Gentry et al. 19477 (US); Tres Marias Islands, María Madre, Ferris 6249 (DS, US). Sinaloa: Labradas vic., Ferris & Mexia 5290 (CAS, DS); Culiacán, Gónza-

lez-Ortega 6632 (US); Mpio. Concordia, Sindicatura Mesillas, Cañada La Calera, 200 m, Trejo 1112 (US); Sindicatura Pantuco, La Calera, González-Ortega 271 (MEXU); Mpio. Mazatlán, González-Ortega 7393 (US).

***Philodendron dwyeri*** Croat, sp. nov. TYPE: Belize. Cayo: Macal (Macaw) River, Guacamallo Bridge, 16°52'N, 89°05'W, Dwyer & Liesner 12334 (holotype, MO-2179389). Figure 155.

Planta epiphytica aut epilithica; succus albus; internodia longior quam lata, ca. 2 cm diam.; cataphylla probabiliter decidua; petiolus teres, 36.5 cm longus, 4 mm diam., leniter longior quam laminae; lamina ovata, 33 cm longa, 21 cm lata, in sicco cana-viridis supra, flaviviridis infra; sinus 7–8 cm profundus; inflorescentia 1; pedunculus 7 cm longus, 5 mm diam.; spathe 9.5 cm longa, omnino viridis, in sicco intus rubrobrunnea; pistilla 7–8-locularia; loculi 1-ovulati.

Epiphytic or epilithic; sap white; internodes slightly longer than broad, ca. 2 cm diam., semiglossy, epidermis light brown, drying conspicuously wrinkled and folded into sharp, irregular ridges; roots 3–4 mm diam., drying reddish brown with prominent ridges, semiglossy, with thin broad scales; cataphylls not seen, probably deciduous; **petioles** 36.5 cm long, 4 mm diam., terete; **blades** ovate, acute at apex (the acumen  $\pm$  inrolled, 5 mm long), cordate at base, 33 cm long, 21 cm wide (1.5 times longer than wide), (0.9 times the petiole length), slightly shorter petioles, margins sinuate, upper surface drying gray-green, lower surface paler, drying yellow-green; anterior lobe 24.6 cm long, 21 cm wide (1.2 times longer than wide); posterior lobes 9 cm long, 9.5 cm wide, rounded, directed toward base; sinus 7–8 cm deep; midrib convex and colorous above, convex and paler, drying yellowish, closely fissured and minutely warty below; basal veins 3 per side, with 1 free to base, 1 coalesced, second and third veins coalesced 1.9 cm; posterior rib not naked; primary lateral veins 5 per side, departing midrib at a 50–60(75)° angle; minor veins moderately distinct, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 1 per axil; peduncle 7 cm long, 5 mm diam.; **spathe** 9.5 cm long (1.3 times longer than peduncle), visibly constricted above the tube, elliptic and to 5.5 cm wide when flattened, green throughout, green, drying reddish brown within; spathe tube ca. 4 cm long; **spadix** sessile; 7.2 cm long, broadest  $\pm$  uniform throughout; pistillate portion cylindrical, 1.3 cm long, 1 cm diam.; staminate portion 6 cm long; fertile staminate portion cylindrical, 1 cm diam., broader than the pistillate portion, sterile staminate portion not detectable; pistils 1.1 mm long, 0.6 mm diam.; ovary 7–8-locular, 0.6 mm diam., with basal placentation; ovules 1 per



locule, 0.1 mm long,  $\pm$  equal in length to funicle, style 0.6–0.8 mm diam., similar to style type D; style apex  $\pm$  flat, with style boss; stigma subdiscoid, unlobed, 0.7–1.1 mm diam., 0.1 mm high, covering interior faces of stylar funnel; the androecium truncate, margins 4–6-sided; thecae oblong, very elongated, 0.2 mm wide.

Flowering in *Philodendron dwyeri* is documented by a single post-anthesis collection from January, early in the dry season.

*Philodendron dwyeri* is endemic to Belize, known only from the type collection in Cayo District of Belize at the Macal (Macaw) River near the Guacamallo Bridge at less than 500 m elevation.

*Philodendron dwyeri* is a member of *P.* sect. *Calostigma* subsect. *Macrobelum* ser. *Macrobelum*. This species is characterized by having internodes longer than broad; white sap; terete petioles (slightly longer than blades); ovate blades drying gray-green above and much paler gray-green below with a very narrow sinus; solitary inflorescences with a green spathe; and ovaries with seven to eight locules and one ovule per locule.

*Philodendron dwyeri* is probably most easily confused with *P. breddlovei* from adjacent Chiapas, which is also a vine with blades of similar size and shape. That species differs, however, in having blades that dry dark yellow-brown with a more or less V-shaped sinus and ovaries with about 20 ovules per locule.

*Additional specimen examined.* BELIZE, Cayo: S of Guacamallo Bridge, *Whiteford* 2837A (BM).

***Philodendron edenudatum* Croat, sp. nov.**

TYPE: Panama, Veraguas: along road from Santa Fe to Río Calovéhora, vic. of Alto Piedra, on Atlantic slope, 0.6 mi. N of Escuela Agrícola Alto Piedra (now Escuela Primer Agrícola Alto Piedra), 735 m, 4 Apr. 1976, *Croat & Folsom* 33988 (holotype, MO-2389069; isotypes, B, COL, CR, F, K, NY, US). Figures 156, 161–164.

Planta hemiepiphytica aut epiphytica; internodia 2.5–7.5 cm longa, 2–5 cm diam., viridia vel cana, in sicco pallide flavibrunnea; cataphylla 20–31 cm longa, acute 2-costata, decidua; petiolus subterete aut obtuse D-formatus, 34–41 cm longus, 1–1.5 cm diam., virens, guttatis purpureis; lamina ovata vel anguste ovata, cordata basi, 34–63 cm longa, 17–33 cm lata; costa postica haud nuda; inflorescentia 1; pedunculus 7–9 mm longus, 9–12 mm diam.; spathe 10–13.5 cm longa; lamina spatheae extus viridiflava, intus pallide viridi; tubo spatheae extus atriviolaceo-purpureo, intus atrisaronino; pistilla 7-locularia; loculi (3/4)-ovulati.

Hemiepiphytic to epiphytic; appressed-climbing, leaf scars inconspicuous, 2 cm long, 2–2.5 cm

wide; internodes moderately smooth and glossy, loosely and irregularly ribbed and grooved, 2.5–7.5 cm long, 2–5 cm diam., green to gray, soon brownish and  $\pm$  densely transversed-fissured to scurfy and glossy to semiglossy, drying pale yellowish brown; epidermis flaking; roots 1–few per node, reddish brown; cataphylls 20–31 cm long, D-shaped to sharply 2-ribbed, green, spotted with purple, persisting intact at upper nodes, eventually deciduous; **petioles** 34–41 cm long, 1–1.5 cm diam., subterete to obtusely flattened or obtusely D-shaped adaxially, especially toward apex, medium to dark green, surface densely pale lineate, purple-spotted; sheathing up to  $\frac{3}{4}$  its length; **blades** ovate to narrowly ovate, subcoriaceous, semiglossy, moderately bicolorous, abruptly acuminate at apex, cordate to subcordate at base, 34–63 cm long, 17–33 cm wide (1.8–2.2 times longer than wide), longer than petiole, upper surface drying gray to gray-brown, lower surface drying yellow-brown; anterior lobe 29–51 cm long, margins convex; posterior lobes 6–17 cm long, 6–13 cm wide, directed downward; sinus arcuate to V-shaped or rarely parabolic, (4)6–12 cm deep; midrib flat to concave, paler than surface above, convex to narrowly rounded, maroon-spotted and darker than surface, drying paler than surface below; basal veins 3–4 per side, second and higher (or sometimes only third and fourth) coalesced 0.5–4.5 cm; posterior rib not at all naked; primary lateral veins 5–6 per side, departing midrib at a 60–75° angle, weakly curved to gradually curved downward just before reaching the midrib, weakly sunken and paler above, convex and darker below; minor veins moderately distinct to  $\pm$  obscure, arising from both the midrib and primary lateral veins, drying minutely undulate. **INFLORESCENCES** 1 per axil; peduncle 7–9 mm long, 9–12 mm diam.,  $\pm$  terete, light green, sometimes faintly tinged reddish medially on one side, clearly demarcated from spathe; **spathe** semiglossy, 10–13.5 cm long, 2–2.5 cm diam. midway, weakly constricted above the tube, oblong-ellipsoid, abruptly acuminate at apex (the acumen ca. 8 mm long); spathe blade greenish yellow outside, pale green, faintly striate inside, resin droplets forming on blade surface within; spathe tube 5 cm long, dark violet-purple with thin greenish margin (ca. 5 mm wide) and along a narrow band adaxially, faintly pale striate-speckled to faintly pale lineate outside, dark maroon inside; **spadix** sessile; gradually tapered to a blunt apex, 7.3–11.5 cm long; pistillate portion greenish, 4 cm long in front, 3.2 cm long in back, 1.2 cm diam.; staminate portion 7.3 cm long; fertile staminate portion 1.4 cm diam. at base, 1.1 cm diam. at middle, 8 mm diam. ca. 1 cm from

apex, broadest at the base; sterile staminate portion 1.4 cm diam.; pistils 2.3 mm long; ovary 7-locular, 1.7 mm diam., with sub-basal placentation; ovules 3(4) per locule, contained within gelatinous matrix (no true envelope), 0.5 mm long; funicle 0.3 mm long (can be pulled free to base), style 1 mm diam., similar to style type B; style apex flat; stigma lobed, 1 mm diam., 0.2 mm high, covering apical depressions (forming a ring without papillae in the center of the apex), depressed medially; the androecium truncate, prismatic, irregularly 4-6-sided, mostly 4-5-sided, 0.9-1.2 mm long; sterile staminate flowers irregularly 4-5-sided, prismatic, 1.3-1.8 mm long.

Flowering in *Philodendron edenudatum* possibly occurs in the dry season, based on a single, post-anthesis collection made in April.

*Philodendron edenudatum* is endemic to Panama, known in *Premontane rain forest* life zones at 110 to 1150 m elevation. It perhaps occurs in the adjacent Chocó of Colombia.

*Philodendron edenudatum* is a member of *P.* sect. *Calostigma* subsect. *Macrobolium* ser. *Macrobolium*. This species is characterized by its slightly elongate internodes; sharply 2-ribbed cataphylls; somewhat D-shaped, usually purple-spotted petioles shorter than the blades; and ovate leaf blades with maroon-spotted midribs and posterior ribs which are never naked (hence the epithet "edenudatum," meaning not naked). Also characteristic are the solitary greenish inflorescences with the spathe tube dark purple-violet inside.

*Philodendron edenudatum* was first collected in 1979 at Alto de Piedra in Veraguas and more recently on Cerro Pirre in the Darién.

The species appears closest to *P. graymii* and is perhaps only subspecifically distinct from that species. The latter differs in having more regularly and conspicuously ridged dried stems with a glossy, brownish yellow epidermis; petioles always longer than the blades (1.09-1.44 times longer); usually larger, more broadly ovate leaf blades (1.2-1.5 times longer than wide) with secretory ducts easily visible on the lower surface; and much larger inflorescences with ovaries having axile placentation (vs. sub-basal in *P. edenudatum*).

*Additional specimens examined.* PANAMA, Darién: Parque Nacional Cerro Pirre, W side, 550-760 m, 7°57'N, 77°46'W, Croat 68891 (CM, MO, PMA, US); 68901 (CM, MO); 68903 (MO); vic. of field station along Río Perisencio, 17 km N of El Real, 110 m, 8°01'N, 77°44'W, Croat & Zhu 77087 (MO, NY, US); 100 m, 77185 (CAS, CM, MEXU, MO, VEN); vic. Cerro Pirre, trail to Rancho Frío on slopes of Cerro Pirre, 200-450 m, 7°58'N, 77°43'W, Croat & Zhu 77157 (AAU, CM, MO, US); Serranía de

Pirre above Cana Gold Mine, Río Cana-Río Escucha Ruido, 600-1000 m, Croat 37741 (MO). Veraguas: Santa Fe-Río Calovebora, 1.7 mi. past Escuela Agrícola Alto Piedra, 570 m, 8°38'N, 81°08'W, Croat & Zhu 76861 (CM, MO); trail to top of Cerro Tute, 1050-1150 m, Croat 48904 (MO); 48906 (MO, US).

***Philodendron ferrugineum* Croat, sp. nov.**

TYPE: Panama. Panamá: along El Llano-Cartí road, 8.3 mi. above Inter-American Hwy., 380 m, 17 July 1987, Croat 67400 (holotype, MO-3582221-22; isotypes, AAU, B, CM, COL, CR, F, K, L, MEXU, PMA, US). Figures 165-168, 173, 174, 185.

Planta hemiepiphytica; internodia brevia, 3-5 mm diam.; cataphylla (15)26-48 cm longa, incostata aut obtuse 2-costata prope apicem; petiolus teres aut semiteres, 41-67 cm longus, 1-1.3 cm diam., circa tam longus quam lamina; lamina anguste ovata, manifeste cordata basi, (38)56-85(102) cm longa, 17-56(62) cm lata, in sicco rubrilinea; venis minoribus obscuris, arcte dispositis, ut videtur intermittentibus; inflorescentia 4-6; pedunculus 2.5-11 cm longus, 1-2 cm diam.; spathe 16-23 cm longa; lamina spatheae extus pallide viridi, intus albidia; tubo spatheae viridi, extus dense maculato violaceo-purpureo, intus purpureo-violaceo vel pallide marronino; pistilla 8-10-locularia; locali 4-7-ovulata; baccae aurantiacae.

Hemiepiphytic; stem appressed-climbing, pale to dark green, soon brown, to 1 m long, leaf scars conspicuous, 2.5-3 cm long, 2-5 cm wide; internodes semiglossy, short on adult plants, 3-5 cm diam., sometimes longer than broad, pale to dark green, becoming gray, finally brown, somewhat scurfy, sometimes transversely fissured; roots dark brown, to ca. 30 cm long, 1-2 cm diam., drying reddish brown; cataphylls (15)26-48 cm long, unribbed, except bluntly or sharply 2-ribbed near apex, green, blotched with purple-violet, to paler green or whitish to densely dark green speckled, maroon spotted or tinged, sulcate between ribs, drying reddish brown, deciduous, persisting weakly at upper nodes, rounded at apex margins clear to hyaline. LEAVES spreading-pendent, scattered evenly along stem, clustered at or near stem apex; petioles 41-67 cm long, 1-1.3 cm diam. (about as long as the blade), erect-spreading, terete or subterete, dark green, firm, weakly and obscurely sulcate adaxially, often obtusely flattened and obtusely ribbed toward apex, to bluntly and broadly sulcate near base adaxially, surface weakly glossy to matte, weakly and densely light green-lineate or striate, purplish red ring around apex; sheathing to 6 cm long; geniculum thicker than petiole, 2.5-3.5 cm long, slightly paler than petiole; blades pendent, narrowly ovate, moderately coriaceous, bicolorous, gradually to strongly acuminate at apex (the acumen tightly inrolled), cordate at base, (38)56-

85(102) cm long, 17–56(62) cm wide (1.3–2.3 times longer than wide), (0.8–1.3 times longer than petiole), about equal in length to petiole, broadest somewhat above point of petiole attachment, or at lower one-third, margins hyaline, weakly to strongly undulate; upper surface dark to medium green, drying coriaceous, reddish brown, semiglossy to glossy, weakly arched along midrib; lower surface very weakly glossy to matte, much paler, drying with minute uninterrupted ridges; anterior lobe 42–70 cm long, 9–22.2 cm wide ((7.5)12–27 times longer than posterior lobes), margins broadly convex; posterior lobes 14–27 cm long, directed downward or inward, sometimes overlapping, obtuse to rounded; sinus strongly spatulate to hippocrepiform; midrib flat to weakly convex, paler than surface above, convex to narrowly rounded, lineate, sometimes maroon-spotted, dark green, paler than surface, drying dark yellowish brown below; basal veins 2–3–5 per side, with 1(2) free to base, 2–5 coalesced 2–4.5 cm, convex and paler than surface above, bluntly acute to convex below; posterior rib not naked, straight to weakly curved; primary lateral veins 5–8 per side, departing midrib at a 45–70° angle, ± straight, slightly curved toward apex, pale green to whitish, usually obtusely sunken, sometimes to weakly and bluntly raised above, convex and darker than surface below; interprimary veins weakly visible to inconspicuous above and below; secondary veins drying inconspicuous; minor veins moderately obscure, close and apparently intermittent, giving veins a bumpy look, arising from both the midrib and primary lateral veins. INFLORESCENCES semi-erect to erect, 4–6 per axil; peduncle 2.5–11 cm long, 1–2 cm diam., whitish at base; spathe coriaceous, 16–23 cm long (1.6–4.2 times longer than peduncle), slightly constricted above the tube, acuminate (the acumen inrolled), dark green outside; spathe blade with lateral margins rolled back, sometimes pale green outside, whitish inside; spathe tube green, heavily spotted with purple-violet outside, 6–8(12) cm long, purple-violet to light maroon, at least the lower ½, otherwise greenish white inside; spadix sessile; clavate, weakly protruding forward at anthesis, 12–20 cm long, broadest above the middle, constricted weakly above sterile staminate portion; pistillate portion pale green to yellow, cylindrical or tapered toward the apex to narrowly ellipsoid, 3.7–6.5(8) cm long, 1.3–1.5 cm diam. at apex, 1.6 cm diam. at middle, 1.1–1.8 cm wide at base; staminate portion 8.6–16 cm long; fertile staminate portion white to yellowish, ± ellipsoid to clavate, 1.1–1.5 cm diam. at base, 1.3–1.9 cm diam. at middle, 1.1–1.5 cm diam. ca. 1 cm from apex, broadest at the mid-

dle or at the base, broader than the pistillate portion, broader than the sterile portion; sterile staminate portion narrower than the pistillate portion, white, 1.5 cm diam.; pistils 2–3.4 mm long, (0.7–0.8)1.6–2 mm diam.; ovary 8–10-locular, 1.6 mm diam., with sub-basal (axile) placentation; ovules 4–7 per locule, 1-seriate, 0.4–0.5 mm long, longer than (or equal in length to) funicle; funicle adnate to lower part of axile wall; style 1.1–1.6 mm diam., similar to style type D; style apex flat, with raised annulus; style boss broad; stigma button-like with medial depression, 1.3 mm diam., 0.2–0.5 mm high, covering entire style apex except (including annulus) in center, medially and shallowly depressed; the androecium truncate, margins 4–6-sided, irregularly scalloped or lobed on at least one margin; thecae oblong, 0.4–0.5 mm wide, contiguous; sterile staminate flowers rounded or blunt, 2.2–2.5(4) mm long, (1.9)2.2–3.2 mm wide, 2.2–2.5(4) mm long. INFRACTESCENCE erect to semi-erect; berries turning orange, obovoid-ellipsoid, apex blunt, 0.8–1.2 mm long, 0.5 mm diam.; mesocarp pale yellow to orange; seeds 3–7 per locule, pale and striate or brown and smooth, strongly sulcate, 1–1.5 mm long, 0.5–0.6 mm diam., enclosed in a translucent envelope. JUVENILE blades narrowly ovate, rounded at base, gradually acuminate at apex, ± shorter than petiole.

Flowering in *Philodendron ferrugineum* is recorded by only one collection in August, although post-anthesis collections range from May through September, indicating that the species probably flowers in the rainy season.

*Philodendron ferrugineum* is currently known only from Panama, from 0 to 770 m elevation in *Tropical wet forest* and *Premontane wet forest* life zones.

*Philodendron ferrugineum* is a member of *P. sect. Calostigma* subsect. *Macrobetium* ser. *Macrobetium*. This species is distinguished by its short internodes; unribbed, deciduous cataphylls; terete to subterete petioles about equaling the blades; and large, thick, cordate blades drying reddish brown with the minor veins obscure, close, and apparently intermittent with a bumpy look. Also characteristic are the 4–6 inflorescences per axil with the spathe tube green outside and maroon inside.

*Philodendron ferrugineum* can be confused with *P. lanense* Croat, with which it occurs in both Bocas del Toro and on the El Llano Cartí Road and Cerro Jefe in Panamá Province and which has similar blades. The latter species differs in having cataphylls becoming fibrous and blades drying greenish brown to somewhat blackened, never

conspicuously reddish brown. *Philodendron ferrugineum* has also been confused with *P. grayumii*. See the discussion of that species for differences.

A similar, apparently undescribed species from Colombia resembles *P. ferrugineum* in its large ovate, thick blades (especially in living condition) and in having three to four inflorescences per axil. Examples include *Croat 56246, 56708*, and *Mon-salve 911* from Bajo Calima near Buenaventura in Valle Province. These differ, however, in drying grayish brown above and yellowish brown beneath and in having interrupted secretory ducts between the minor veins. In addition, they do not have the undulated and puckered minor veins on drying that are so characteristic of specimens of *P. ferrugineum*.

A noteworthy collection is *McPherson 11479*, which is somewhat intermediate between *P. ferrugineum* and *P. grayumii*. That collection has blades that dry more reddish brown and have a bumpy surface but also secretory ducts. It may prove to be a new species.

*Additional specimens examined.* PANAMA. **Bocas del Toro:** Chiriquí Lagoon, on Cayo Agua, 5 m, 9°10'N, 82°W, *McPherson 11479* (AAU, MO); Escudo de Veraguas Island, 5 m, 9°05'N, 81°35'W, *McPherson 11410* (MO, PMA). **Colón:** Santa Rita Ridge Road, along trail to Río Indio, 10.6 km from Transisthmian Highway, 380 m, *Croat 34349* (MO); along route between Sabanita and Portobello, 3.9 mi. from highway, 250 m, 9°22'30"N, 79°41'30"W, *Croat 75155* (AAU, BR, CM, I, MO, PMA); Río Agustín, Río Guanche, ca. 0 m, 9°30'N, 79°40'W, *Churchill et al. 6018* (MO); Río Iguanita, ca. 3 km above bridge on Portobelo road, <100 m, 9°27'N, 79°40'W, *Croat 49749* (MO). **Panamá:** El Llano-Cartí Road, *Croat 33145A* (MO); Km 7-12 km, 360-400 m, 25098 (MO); 5-6 mi. N of highway, 350-375 m, 34787 (MO); Mile 3.4, 1000 ft., 49098 (MO); Mile 4, 33732 (CAS, F, MO, NY); Km 7-12, 360-400 m, 25173 (MO); Km ca. 17, 400 m, 9°15'N, 78°50'W, *Knaap 1376* (MEXU, MO); Cerro Jefe region, 9°15'N, 79°30'W, *Croat & Zhu 76216* (CM, MO); 21 km above Pan-American Highway, 600 m, *Croat 35886* (MO); 750-800 m, 9°14'N, 79°22'W, *Croat 67092A* (CM, MO, PMA); 4.6 km beyond peak on road to Altos de Pacora, 26.3 km from Inter-American Highway, 600 m, *Croat 35923* (MO); at Altos de Pacora, 750 m, 9°15'N, 79°29'W, *Croat & Zhu 76607* (CM, MO); 0.8 mi. beyond turnoff to Altos de Pacora, 770 m, 9°15'N, 79°29'W, *Croat & Zhu 76612* (CR, MO, NY). **San Blas:** Nusgandi, along El Llano-Cartí Rd., 0.7 mi. beyond Cuna Headquarters, located 10.9 mi. N of Pan-American Highway, 450 m, 9°18'N, 79°59'W, *Croat 75116* (CM, MO, PMA, TEX); 300-400 m, 9°20'N, 79°W, *Hamilton & Stockwell 1073* (MO); 1-2 mi. N of Nusgandi on road to Cartí, 250-275 m, 9°20'N, 79°W, *Croat & Zhu 76577*, (CAS, I, MO, PMA, US); 76580 (CM, MO, SEL, WIS); Mile 10.1, 300 m, 9°20'N, 79°W, *Croat & Zhu 77029* (COL, CR, MEXU, MO); 77030 (MO, PMA).

### *Philodendron findens* Croat & Grayum, sp. nov.

TYPE: Panama. Chiriquí: Fortuna Dam area, Fortuna-Chiriquí Grande, 1.8 mi. NW of center of dam, 1080 m, 8°45'N, 82°18'W, 27 June 1994, *Croat & Zhu 76502* (holotype, MO-4619581-84; isotypes, B, COL, CR, F, K, NY, PMA, US, VEN). Figures 7, 34, 169-171, 175, 177, 178.

Planta plerumque hemiepiphytica, variis terrestribus; internodia 4-9 cm longa, 2-4 cm diam.; cataphylla usque 35 cm longa, acute 2-costata, persistentia semi-intacta nodis superioribus, fibrosentia, demum decidua; petioli D-formati, (25)43-119 cm longi, 0.6-35(45) cm diam., cum ala marginali tenui, erecta; lamina ovato-cordata, (25)40-76 cm longa, (20.5)28-70 cm diam., findens inter nervos laterales I sic videtur fere pinnata, in sicco denigrata; inflorescentia 2; pedunculus 4-12(15) cm longus, 0.3-1.4 cm diam.; spathe 13-22.8 cm longa; lamina spathe extus virens vel pallide viridi; intus viridialba; tubo spathe extus viridi, intus rubro; pistilla 5-6-localaria; loculi 15-20(22)-ovulati.

Usually hemiepiphytic, rarely terrestrial on steep banks; stem appressed-climbing, medium green, glossy, sap watery, spicy-scented, leaf scars conspicuous, 3.5-4 cm long, 5 cm wide; internodes short, thick, semiglossy, 4-9 cm long, 2-4 cm diam., longer than broad, dark olive-green, epidermis cracking, yellowish, fissured longitudinally and transversely; roots moderately few per node, drying dark brown, semiglossy, acutely ridged; cataphylls thin, fleshy, somewhat spongy, to 35 cm long, sharply 2-ribbed (ribs 2-3 mm high), D-shaped, thicker than broad, medium green, light or pale yellow-green toward margins, semiglossy, broadly sulcate abaxially, persisting semi-intact at upper nodes, becoming fibrous and eventually deciduous, blunt to acute at apex, margins weakly raised adaxially; petioles (25)43-119 cm long, 0.6-35(45) cm diam., sharply D-shaped, with a thin, erect marginal wing, the wing undulate toward apex, surface densely light striate near base; blades large ovate-cordate, subcoriaceous, strongly bicolorous, short acuminate at apex, cordate at base, (25)40-76 cm long, (20.5)28-70 cm wide (1.1-1.5 times longer than wide), (0.6-1.1 times longer than petiole), margins usually promptly splitting into segments by dividing between the primary lateral veins, making blades appear almost pinnate, upper surface dark green, semiglossy, drying blackened to dark reddish brown, nearly matte, lower surface semiglossy to matte, paler; anterior lobe 18-57.5 cm long, 15.5-55(79) cm wide (2.2-3.2 times longer than posterior lobes); posterior lobes (7.2)11.5-32 cm long, 8.6-28(37.5) cm wide, obtuse to rounded; sinus  $\pm$  parabolic; midrib weakly to deeply sunken, paler than surface above, con-

vex, weakly striate, slightly paler than surface below; basal veins 5–12 per side, with 0–1(2–3) free to base, in part coalesced 5–10 cm; posterior rib naked for 1–7 cm; primary lateral veins 5–7 per side, departing midrib at a 40–50° angle, deeply sunken and paler than surface above, convex, usually concolorous below; interprimary veins visible, few near apex above and below; tertiary veins visible, darker than surface below; minor veins weakly raised and darker than surface, arising from both the midrib and primary lateral veins; secretory canals obscurely visible. INFLORESCENCES erect, 2 per axil; peduncle 4–12(15) cm long, 0.3–1.4 cm diam., pale to medium green, prominently, densely white-streaked; spathe 13–22.8 cm long (1.4–2.4 times longer than peduncle), constricted midway slightly above the tube; spathe blade medium to pale green outside, greenish white inside; spathe tube oblong-ellipsoid, green, densely and minutely lineate-speckled outside, 7–9 cm long, red (B & K red-purple 4/10, 2/10) inside; spadix sessile, tapered, 13–23 cm long, broadest just below the middle; pistillate portion gray-white, tapered toward the apex, slightly curved, 4–8.5 cm long, 1.4–2.1 cm diam. at apex, 1.6 cm diam. at middle, 1.9 cm wide at base; staminate portion 14.3–17.5 cm long; fertile staminate portion creamy white, tapered, 1.8–2 cm diam. at base, 1.3 cm diam. at middle, 9 mm diam. ca. 1 cm from apex, broadest at base, narrower than the pistillate portion, narrower than the sterile portion; sterile staminate portion broader than the pistillate portion, 1.6 cm diam.; pistils (3)7.7–8.4 mm long, 2–2.5 mm diam.; ovary 5–6-locular, locules (1.6)5.8–6.1 mm long, with axile placentation; ovules 15–20(22) per locule, 2-seriate, 0.3–0.5 mm long, longer than funicle, style similar to style type B; style apex rounded; stigma subdiscoid to discoid, ± lobed, 1.2–2 mm diam., 0.1–0.5 mm high, covering entire style apex; the androecium truncate, 4–6-sided; thecae oblong, 0.5 mm wide, contiguous; sterile staminate flowers blunt, 3.1 mm long, 1.2 mm wide. INFRUCTESCENCE 12 cm long, 3.7 cm diam., 19.5 cm long; stipe 4.5 cm diam.; berries irregularly quadrangular to bluntly 4–5-sided, with a kiwi-fruit scent, 3.5–4.2 mm diam.; seeds 1–2 per locule, ca. 17 per berry, somewhat flattened, 1.5–2.2 mm long, 0.3–0.4 mm diam., translucent and with fine striations. JUVENILE stems dark green to yellow-brown, semi-glossy to glossy, 2.4 cm long, 5 mm diam.; blades thin, entire. PRE-ADULT petioles broadly convex adaxially, the margins spreading winged, incurved, erect; blades moderately bicolorous.

Flowering in *Philodendron findens* occurs in May and July, but post-anthesis and immature fruiting

collections from March through November (except October) suggest a broader flowering period, with plants flowering earlier in the dry season or much later in the rainy season.

*Philodendron findens* ranges from Costa Rica to Panama on the Atlantic slope, from near sea level to 1400 m elevation in mostly *Tropical wet forest* and *Premontane rain forest* but also in *Premontane wet forest*. It probably also occurs in Colombia.

*Philodendron findens* is a member of *P. sect. Philodendron* subsect. *Platypodium*. This species is characterized by its generally appressed-climbing habit and short, thick internodes with sharply two-ribbed cataphylls, which are finally fibrous and persistent; D-shaped petioles with marginal wings adaxially; and especially by its usually black-drying, large, ovate-cordate, leaf blades, which promptly shred into segments, making the blade appear almost pinnate (hence the name "findens," meaning "tearing or splitting"). Also characteristic are the paired, green inflorescences with the spathe tube red within.

*Philodendron findens* can be confused with *P. pterotum*, which has similarly shaped blades and D-shaped petioles with narrowly winged margins. That species differs, however, in having blades drying yellow-green, and major veins drying paler than the surface below (in contrast to blackened and darker than the surface in *P. findens*) and do not rip into segments, and also by occurring more commonly in *Tropical moist forest* and at elevations of usually less than 400 m. In addition, the spathe tube in *P. pterotum* is purplish on the outside, vs. solid green (sometimes reddish) for *P. findens*.

*Cuatrecasas 14948*, from Valle Department along Río Digua at Piedra de Moler at 900–1180 m, is apparently this species. It differs in no significant way, but there are no field notes and the petiole is improperly preserved so its cross-sectional shape cannot be confirmed.

*Additional specimens examined.* COSTA RICA. **Alajuela:** ca. 7.5 km N of bridge over Río Balsa, 700–800 m, 10°10'–15'N, 84°30'–35'W, *Stevens 13892* (MO); ca. 5.7 km N of bridge over Quebrada Volio, 1100–1150 m, 10°06'N, 84°29'W, *Stevens 14115* (MO); San Ramón Forest Reserve, ca. 10 km W of Laguitos, along Río San Laurencio, 850–1100 m, 10°18'N, 84°34'W, *Hammel et al. 15283* (MO); ca. 1.3 mi. N of Angeles Norte, 1200 m, *Croat 46876* (MEXU, MO); 800–1000 m, *Nilsen 458* (CR); Reserva Biológica Monteverde, 950 m, 10°18'N, 84°45'W, *Bello & Méndez 2667* (MO); 2668 (CR); 820 m, 10°19'N, 84°43'W, *Bello & Eyal 2682* (JNB, MO). **Cartago:** Moravia-Quebrada Platanillo, Moravia, 3–5 km from Finca Racine, 1200–1300 m, *Croat 36627* (MO); Río Navarro, El Muñeco, 1400–1500 m, *Standley & Torres 51355* (US). **Heredia:** 3 mi. S of Cariblanco, 760 m, *Croat 35796* (MO); 35813 (MO); 4 mi. N of Vars Blanca, 1350

m, *Croat* 35606 (MO); Río Frío de Sarapiquí, Río Sució-Finca Zona Ocho, 110 m, 10°18'N, 83°52.5'W, *Grayum & Hammel* 5565 (CR, L, MO); Río Frío, W of Finca Zona Nueva, ca. 110 m, 10°18'N, 83°53'W, *Grayum* 3561 (K, M, MO, RSA). **LIMÓN:** Fila Lleskila Talamanca, 1160 m, *Gómez et al.* 23061 (CM, K, MO). **San José:** Braulio Carrillo National Park, 600-700 m, *Croat* 78751 (CM, CR, INB, M, MO). **PANAMA:** **Bocas del Toro:** Fortuna Dam area, Chiriquí Grande-Fortuna, 470 m, 8°50'N, 82°15'W, *McPherson* 11646 (MO); ca. 0 m, 8°55'N, 82°09'W, *Croat* 66811 (MO); Almirante-Ojo de Agua, 3-6 km W of Almirante, 30-200 m, *Croat* 38218 (BR, CAS, F, K, MO, PMA, US); 2.8 mi. N of Divide, 850-950 m, 8°45'N, 82°15'W, *McPherson* 9674 (MO, NY). **Chiriquí:** Gualaca-Chiriquí Grande, 1.4 mi. W of Centro de Operaciones, trail to Río Hornito, 1010-1130 m, 8°44'N, 82°14'30"W, *Croat* 67919 (B, F, K, M, MO); *Croat & Zhu* 76293B (MO); Río Chiriquí, near La Sierpe, ca. 0.5 km N of river, IRHE Fortuna Hydroelectric Project, 1000-1100 m, 8°46'N, 82°12'W, *Knapp* 5052 (MO). **Corlé:** El Valle de Antón region, at La Mesa, 860-900 m, *Croat* 37410 (MO); 775 m, 8°36'N, 80°07'W, 74795 (MO); 800-900 m, 8°38'N, 80°09'W, 67153 (MO); *Luteyn & Kennedy* 1616 (MO); El Copé region, N of El Copé, 1200-1300 m, *Sytzma & Anderson* 4622 (MO). **Colón:** 10 mi. SW of Portobelo, 2-4 mi. from coast, 10-200 m, *Liesner* 1115 (F, MO, NY, US); Río Guanche, ca. 2 km E of bridge on main Puerto Pilón-Portobelo Road, 100 m, ca. 9°30'N, 79°39'W, *Croat* 75175 (B, CAS, COL, F, K, MO, NY, PMA, US). **Darién:** along headwater of Río Tuquesa, ca. 2 km air distance from Continental Divide, Tyler Kätredge gold mine, *Croat* 27148 (MO). **Veraguas:** Santa Fe region, Santa Fe-Río San Luis, beyond Escuela Agrícola Alto de Piedra, 5.9 mi. N of school, 480 m, 8°33'N, 81°08'W, *Croat* 66937 (MO, PMA, US); trail up Cerro Tute, 1050-1150 m, *Croat* 48899 (CM, MO); Escuela Agrícola Alto de Piedra-Río San Luis, vic. of Santa Fe, along Río Primero Brazo, on Atlantic Coast, 490 m, 8°33'N, 81°08'W, *Croat* 66879A (MO); Escuela Agrícola Alto Piedra-Calovobora, 15.6 km NW of Santa Fe, along trail to Santa Fe, E of Río Dos Bocas, 450-550 m, *Croat* 27653 (MO); beyond Tres Beavos River, 11 km beyond Santa Fe, 650 m, *Croat* 25625 (MO).

**COLOMBIA:** Valle: Cordillera Occidental, Río Digua, Piedra de Moler, 900-1180 m, *Cuatrecasas* 14948 (F).

***Philodendron folsomii* Croat, sp. nov. TYPE:**

Panama. Coclé: Alto Calvario, 9 km N of El Copé, N of Continental Divide along path to W, ca. 800 m, 8°40'N, 80°37'W, 24 Jan. 1989, *McPherson* 13619 (holotype, MO 3693168; isotypes, B, K, PMA, US). Figures 172, 176.

Planta epiphytica; internodia brevia, 1-1.5 cm longa, 8-20 mm diam.; cataphylla 11-16 cm longa, acute 2-costata, decidua; petiolius subteretes, obtuse sulcatus, 12-17 cm longus; lamina oblongo-lanceolata, 28-33 cm longa, 5-7 cm lata, rotundata aut truncata basi; inflorescentia 1-2; pedunculus 10-11.5 cm longus, 2-5 mm diam.; spathe 10-12 cm longa, extus alba; pistilla 6-7-localaria; loculi 1-ovulati.

Epiphytic; stem scandent; internodes short, 1-1.5 cm long, 8-20 mm diam., about as long as broad; cataphylls 11-16 cm long, sharply 2-ribbed,

green, deciduous; roots not seen; petioles 12-17 cm long, 3-5 mm diam., subterete, obtusely sulcate; blades oblong-lanceolate, narrowly acuminate at apex, obtuse to almost rounded and decurrent, sometimes nearly truncate at base, 28-33 cm long, 5-7 cm wide (4.6-5.2 times longer than wide), (1.8-2.3 times longer than petiole), upper surface drying gray-green to brownish green; lower surface paler, drying yellowish green; midrib sunken above, convex, drying paler than surface below; basal veins 1-2, free to base; primary lateral veins 8-10 per side, departing midrib at a 35° angle, arcuate to the margins, distinct above; minor veins arising from the midrib only. INFLORESCENCES 1-2 per axil; peduncle 10-11.5 cm long, 2-5 mm diam.; spathe 10-12 cm long (about as long as the peduncle), white throughout outside; spathe tube 5.5 cm long, 2.5 mm diam.; spadix short stipitate; 7-8 cm long; pistillate portion 3-4.1 cm long, 6-7 mm diam.; staminate portion 3.8-4 cm long; sterile staminate portion whitish; pistils 0.8 mm long, 0.6 mm diam.; ovary 6-7-locular, locules 0.6 mm long, 0.3 mm diam., ovule sac 0.6 mm long, with sub-basal placentation; ovules 1 per locule, contained within gelatinous transparent matrix (no true envelope), 0.2-0.35 mm long, as long as funicle; funicle 0.2-0.3 mm long (can be pulled free to base), with glands at base, style similar to style type B; style apex flat; stigma subdiscoid, unlobed, ± truncate, 0.9 mm diam., 0.3 mm high, covering almost entire style apex; the androecium truncate, prismatic, margins irregularly 5-6-sided, 0.6-0.8 mm long.

Flowering phenology in *Philodendron folsomii* is uncertain because few collections exist. Flowering collections are known from January and April, both within the dry season but in an area that is not markedly seasonal.

*Philodendron folsomii* is endemic to Panama, known only from the type locality in Coclé Department, at 700 to 800 m elevation in *Premontane rain forest*.

*Philodendron folsomii* is a member of *P.* sect. *Costigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is characterized by internodes about as long as broad; sharply two-ribbed, deciduous cataphylls; oblong, narrowly acuminate, oblong-lanceolate, green-drying blades with obtuse-attenuate bases; and solitary, long, slender-pedunculate inflorescences with white spathes.

*Philodendron folsomii* is most easily confused with *P. sphaerum* Schott from the Guianas and eastern Venezuela, which has similar leaves. The latter species differs in having smaller, generally

more numerous (up to four) inflorescences per axil with more slender peduncles (drying scarcely 2 mm diam. vs. 4 mm or more in *P. folsomii*) and spathes less than 7 cm long (vs. 10–13.5 cm long in *P. folsomii*).

*Philodendron folsomii* is also similar to *P. correae*, which has similarly shaped leaf blades that also sometimes dry green. The latter species differs in having generally longer internodes; petioles usually sheathed to near the apex; blades usually drying black and lacking conspicuous primary lateral veins; and spathes not so conspicuously constricted above the tube.

*Additional specimen examined.* PANAMA. Coclé: El Copé region, 700 m, *Folsom & Robinson 2447* (MO).

***Philodendron fortuneae* Croat, sp. nov. TYPE:**

Panama. Chiriquí: Gualaca-Chiriquí Grande Road, vic. of Fortuna Dam, at junction of road to IRHE headquarters, 1200 m, 8°45'N, 82°18'W, 23 June 1987, *Croat 66714* (holotype, MO-3635169; isotypes, AAU, B, CAS, CM, COL, CR, DUKE, EAP, F, GH, HUA, K, L, M, MEXU, NY, P, PMA, QCNE, RSA, S, SEL, TEX, US, VEN). Figures 179, 181–184.

Planta terrestris aut hemiepiphytica; internodia 3–6 cm longa, 4–7 cm diam.; cataphylla 18–32 cm longa, acute 2-costata, glauca, decidua; petiolus plus minusve D-formatus, 49–70 cm longus, 1.2–2.5 cm diam., glaucus, marginibus manifeste alatus, alis undulatus; lamina ovato-cordata vel oblongato-ovata, cordata basi, 36–60 cm longa, 22–44 cm lata; inflorescentia 4; pedunculus 7.7–8 cm longus, 8 mm diam.; spatha 20–21 cm longa, omnino vires, tubo spathae marronino, intus suffuso marronino in laminam; pistilla 7–8-locularia; loculi 9–10-ovulati.

Terrestrial or hemiepiphytic; stem appressed-climbing, 2–3 m long, thick, leaf scars conspicuous, 1.2–1.7 cm long, 2–2.5 cm wide; internodes smooth, to weakly pruinose, 3–6 cm long, 4–7 cm diam., about as long as broad, green to olive-green, epidermis drying smooth, semiglossy, light yellow-brown (B & K yellow-red 9/10), cracking with age, fissured transversely, sometimes with minute cracks perpendicular to axis; roots few per node, elongate, to 4 mm diam., light brown, smooth, weakly glossy, the epidermis cracking free upon drying; cataphylls thick, 18–32 cm long, sharply 2-ribbed (ribs 4 mm high), pale green, glaucous, dark short-lineate, deciduous. LEAVES erect-spreading, clustered at or near stem apex; petioles 49–70 cm long, 1.2–2.5 cm diam., ± D-shaped, becoming flattened and much broader than thick toward apex, medium green, rounded to flattened abaxially, with adaxial margins prominently winged (to 3 mm diam.), undulate to broadly flattened, surface dark short-li-

neate, glaucous; blades ovate-cordate to oblong-ovate cordate, moderately coriaceous to subcoriaceous, gradually acuminate to abruptly acuminate at apex, cordate at base, 36–60 cm long, 22–44 cm wide (1.3–1.8 times longer than wide), (0.85–0.88 times the petiole length), broadest slightly above point of petiole attachment, margins moderately undulate, upper surface medium green, semiglossy, drying greenish yellow, lower surface medium to pale green, glaucous, matte, paler, drying greenish yellow; anterior lobe 28–44 cm long, 10–20 cm wide (0.9–1.3 times longer than wide), (1.8–2.9 times longer than posterior lobes), broadest at base; posterior lobes usually overlapping, 10–20 cm long, 10–20 cm wide, directed downward and slightly inward, obtuse to rounded; sinus oblong-triangular or closed, 10–19 cm deep; midrib flat to weakly raised, pale green, concolorous above, weakly raised, paler than surface below; basal veins 5–10 per side, 1–2 pairs free to base, the remainder mostly free, sometimes weakly coalesced 1–4 cm, drying weakly raised; posterior rib usually not present, naked and straight if present; primary lateral veins 7–11(15) per side, departing midrib at a 30–45° angle, straight or slightly arcuate to the margins, flat to sunken and drying darker than the surface above, convex and darker than surface below; interprimary veins prominulous, weakly sunken to flat, concolorous above, flat and darker than surface below; lesser veins visible when dried; minor veins conspicuous, fine, darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES 4 per axil; peduncle 7.7–8 cm long, 8 mm diam., medium green; spathe ± erect, 20–21 cm long (2.5–2.7 times longer than peduncle), medium green throughout, glaucous, ± acute at apex; spathe tube maroon, suffused maroon onto blade inside; spadix sessile; tapered, 20–21 cm long; pistillate portion tapered toward the apex, 5.2 cm long, 1.3 cm diam. at apex, 1.5 cm diam. at middle, 1.7 cm wide at base; staminate portion 14.7 cm long; fertile staminate portion tapered, 1.4 cm diam. at base, 1.3 cm diam. at middle, 1.1 cm diam. ca. 1 cm from apex, broadest at the base, narrower than the pistillate portion, narrower than the sterile portion; sterile staminate portion narrower than the pistillate portion, 1.4 cm diam.; pistils 4.5 mm long, 2.1–2.4 mm diam.; ovary 7–8-locular, locules 1.9 mm long, with sub-basal placentation; ovules 9–10 per locule, 0.4–0.5 mm long, slightly longer than funicle, style 2 mm long, 1.8 mm diam., similar to style type B; style apex flat; stigma subdiscoid, unlobed, 2 mm diam., 0.4 mm high, covering entire style apex; the androecium truncate, margins 4–6-sided; thecae oblong, 0.5 mm wide,

contiguous; sterile staminate flowers blunt, 3.2 mm long, 1.7 mm wide.

Flowering in *Philodendron fortuneuse* is known only from April, and immature fruits from June. This corresponds to the late dry season and early rainy season in most of Panama, though this species occurs in an area with little seasonality.

*Philodendron fortuneuse* is endemic to Panama, known only from Chiriquí Province in the Fortuna Dam region (hence the name), from 1100 to 1300 m in elevation in regions of *Premontane rain forest*.

*Philodendron fortuneuse* is a member of *P.* sect. *Philodendron* subsect. *Platyodidium*. This species is recognized by its thick stems with short, smooth (when fresh), green internodes; sharply two-ribbed, glaucous, deciduous cataphylls; ovate-cordate blades with usually overlapping posterior lobes, and pale, glaucous lower surfaces; and especially by the glaucous, broadly flattened petioles with undulate-winged margins.

*Philodendron fortuneuse*, one of the showiest, most distinctive species in the Central American region, is not easily confused with any other species. Its greatest similarity is to *P. brenesii*, with which it may occur, but that species differs in having proportionately longer, more or less terete petioles, and one to two inflorescences per axil (vs. up to four inflorescences per axil in *P. fortuneuse*).

*Additional specimens examined.* PANAMA, Chiriquí: Fortuna Dam area, Gualaca-Chiriquí Grande, jct. of road to IRHE headquarters, 1200–1300 m, 8°45'N, 82°18'W, Croat 66529 (CM, ENCB, HNMN, JAUM, MO, MY, QCA, RB, W); 9.6 mi. beyond Los Planes de Hornito, 1300 m, Croat 48727 (AAU, COL, F, MO, OOM, QCNE); 48728A (MO); 10.1 mi. NW of Los Planes de Hornito, 1300 m, 82°17'W, 8°45'N, Croat 49814 (MO); 50000 (BR, CAS, CM, L, MO, QCA, SCZ, TEX); 4.5–5 km N of dam over Fortuna Lake, 1100–1135 m, 8°43'N, 82°17'W, Croat & Grayum 60000 (B, CM, K, MO, NY, RSA, US); 8 mi. beyond Los Planes de Hornito, 1.4 mi. W of Centro de Operaciones, trail to Río Hornito, 1010–1130 m, 8°44'N, 82°14'30"W, Croat 67921 (G, M, MO).

*Philodendron fragrantissimum* (Hook.) G. Don, in Sweet, Hort. Brit. ed. 3: 632. 1839. Figures 30, 186–191. *Caladium fragrantissimum* Hook., Bot. Mag. 61: t. 3314. 1834. TYPE: Guayana (specimen introduced by C. S. Parker in 1834 to Liverpool Botanical Garden) (holotype, K).

*Philodendron latipes* K. Koch & Augustin, in A. Braun et al., Append. Gen. Sp. Hort. Berol. 1854: 6. 1854–1855. TYPE: origin unknown (holotype, B destroyed); *Buchell* 9452 (neotype, here designated, K).

*Philodendron poeppigii* Schott, Syn. Aroid. 84. 1860. TYPE: Brazil. Amazonas: (Ega) Tefé, *Poeppig* s.n. (holotype, B).

*Philodendron brevilaminatum* Schott, J. Bot. 2: 4. 1864. TYPE: Brazil. Bahia: Ilheos, *Archduke F. Maximilianus* s.n. (holotype, lost); Schott ic. 3592 (neotype, here designated).

*Philodendron clementis* C. Wright, in Griseb., Cat. Pl. Cub., 220. 1866. TYPE: Cuba. Near Retimo, *Wright* 3212 (lectotype, here designated, K).

*Philodendron demerarae* Gleason, Bull. Torrey Bot. Club 56: 11. 1929. TYPE: Guyana. SE of Lamaha stop-off, 27 Nov. 1919, *Hitchcock* 16987 (holotype, NY).

*Philodendron accrescens* N. W. Simmonds, Kew Bull. 1951: 402. 1951. TYPE: Trinidad. Long Stretch, 15 Jan. 1949, *Simmonds* 14256 (holotype, TRIN).

Hemiepiphytic; acaulescent or caulescent, stem appressed-climbing or rarely scandent with slender, whip-like branches bearing small leaves, to 1–6 m long, sap orangish to brownish, sticky, spicy-scented; internodes short, 1–4 cm diam., usually thicker than long, dark green, semiglossy, obscured by cataphyll fibers, roots often many per node, drying dark brown, 2–3 mm diam.; cataphylls sharply 2-ribbed or sharply D-shaped, greenish white, sometimes drying reddish brown, persisting as fibers, margins weakly upturned below. LEAVES erect-spreading; **petioles** 22–70 cm long, 4–11 mm diam., ± D-shaped to sharply C-shaped, broadly sulcate adaxially; **blades** ovate to ovate-triangular, subcoriaceous, moderately bicolorous, acuminate at apex (the acumen sometimes inrolled or very short apiculate, 2–5 mm long), cordate at base, 21.6–59 cm long, 17–37.5 cm wide (1.1–1.7 times longer than wide), (0.7–1.3 times longer than petiole), about equal in length to petiole, upper surface semiglossy, lower surface semiglossy; anterior lobe 19.5–49 cm long, 17–39 cm wide, (2.1–3.9 times longer than posterior lobes); posterior lobes 5–16.5 cm long, 10–18.3 cm wide, obtuse to rounded; midrib broadly sunken, paler than surface above, convex, bluntly angular, paler than surface below; basal veins (3/4/5) per side, with (0/1/2) free to base, 1–2 coalesced 1–4 cm; posterior rib not naked or naked 1–4 cm along the sinus; primary lateral veins 3–6 per side, etched-sunken above, convex below; interprimary veins visible and discontinuous above; minor veins visible and darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect to semi-erect, 2 per axil; peduncle 3–13.5(17) cm long, 3–15 mm diam.; **spathe** 8.5–19 cm long, (1.2–3.6 times longer than peduncle); spathe blade white to greenish, rarely reddish outside, white to greenish inside; spathe tube reddish to dark maroon outside, red to maroon inside; **spadix** stipitate 3–4 mm long; cylindrical, 9–16 cm long, ± uniform throughout; pistillate portion cylindrical, 2.5–5 cm long, 1.4 cm diam. throughout, 1.6 diam. at apex, 1.8 cm diam.



at middle, 1.5 cm wide at base; staminate portion 6.3–7.8 cm long; fertile staminate portion creamy white, cylindrical, to 1.3 cm diam. at base, 1.4 cm diam. at middle, 1 cm diam. ca. 1 cm from apex, broadest in the middle and as broad as the pistillate and sterile portions; sterile staminate portion as broad as or broader than the pistillate portion, 1.2–1.5 cm diam.; pistils 3.2–3.5 mm long, 2.1–2.9 mm diam.; ovary 6–10-locular, with axile placentation; locules 1.6–2.3 mm long; ovules (24)32–36 per locule, 2-seriate, 0.2–0.3 mm long,  $\pm$  equal in length to funicle; funicle 0.2–0.3 mm long, style similar to style type D; stylar canals emerging into conspicuous depressions; style apex slightly concave to flat; stigma subdiscoïd (brush-like), unlobed, 2.1–2.7 mm diam., 0.5–0.7 mm high, covering entire style apex, inserted on entire style apex; the androecium 4–6-sided; thecae slightly obovate, 0.2–0.4 mm wide; sterile staminate flowers ellipsoidal to globose, 2.2–3 mm long. INFRUCTESCENCE with stipe of spadix maroon; berries usually bright red to purple-red, sometimes orange, rarely yellowish white (McPherson 14496) to white (McPherson 11380).

Flowering in *Philodendron fragrantissimum* in both Panama and Middle America occurs during the rainy season from May to October (except September). Some post-anthesis collections from Middle America are scattered in the dry season as well (January and March). Post-anthesis collections are also principally from May through December (except October) as are immature fruiting collections (July through November). Mature fruits have been collected during May through December (except July and August).

*Philodendron fragrantissimum* ranges from Belize to Panama along the Caribbean coast and to Pacific Colombia (Chocó), then disjunctly to the Amazon basin, where it ranges from Venezuela to the Guianas, northern Brazil (Roraima and Amazonas), southern Colombia (Meta, Caquetá, Guajira, and Vaupes), and Peru (Amazonas, Loreto, Ucayali, and Madre de Dios). It is also known from Cuba and Trinidad, and it is to be expected in Ecuador and western Brazil. Ranging from near sea level to 1000 m elevation, this species occurs in *Tropical moist forest*, *Premontane wet forest*, and *Tropical wet forest* life zones.

*Philodendron fragrantissimum* is a member of *P. sect. Philodendron* subsect. *Macrolonchium*. This species is distinguished by its short internodes; a tendency to produce slender, whip-like branches from near the apex; persistent, reddish brown cataphyll fibers; more or less D-shaped petioles with

somewhat elevated, lateral margins; ovate to ovate-triangular, cordate blades (about equal to the petioles in length); and colorful inflorescences with the spathes bright red on the tube and white on the blade.

*Philodendron fragrantissimum* is not easily confused with any other species in Central America, since the combination of D-shaped petioles, persistent cataphylls, and red and white inflorescences is unique, but it has been confused with the Venezuelan *P. chimantae* G. S. Bunting, which differs in having typically more triangular petioles that are actually winged (not merely ribbed) on the lateral adaxial margins, brown cataphylls persisting intact for a time before falling off, and blades with a much narrower sinus.

*Philodendron fragrantissimum* populations in South America, particularly in the Guiana region (the type locality), have nearly triangular rather than D-shaped petioles typical of those in Central America. In addition, the blade shape is more nearly triangular than ovate as in Central America. I believe, nevertheless, that differences between Central American and South American populations do not warrant taxonomic recognition.

*Additional specimens examined.* BELIZE. Cayo: along Hummingbird Highway at Mile 28, Dwyer 11411 (MO); Mile 34, Gentle 9180 (LL). STANN CREEK: Middlesex, Gentle 2951 (CM, MICH, RSA). Toledo: Punta Gorda-San Antonio, 1 mi. E of junction with road, 100 ft., Croat 24510 (MO). COSTA RICA. Alajuela: Cañas-Uapala, near Río Zapote, 1.8–2.7 km S of Río Canalete, 100 m, Croat 36361 (MO). Cartago: Tucurrique, Las Vueltas, 635–700 m, Tondúz 13313 (G, US). Heredia: Zona Protectora, N slopes of Volcán Barba, along Quebrada Cantarera, 300–400 m, Grayum & Schatz 3179 (DUKE); La Selva, 100 m, Grayum 2436 (CR, MO); Río Sarapiquí, S base of Cerros Sardinial, Chilamate de Sarapiquí, 80–110 m, 10°27'30"N, 84°04'W, Grayum & Hammel 5541 (MO). LIMÓN: Cantón de Talamanca, Amubri, 100 m, 9°31'N, 82°58'W, Hammel et al. 17512 (CR, MO); Barra del Colorado, 0–2 m, 10°47'N, 83°55'W, Stevens 24119 (CR, MO); Río Colorado, 1–5 m, 10°47'40"N, 8°35'30"W, Daviase & Herrera 30922 (CR, MO); Manzanillo de Talamanca, ca. 5 m, 9°38'N, 82°39'W, Grayum & Burton 4327 (MO); Fila Dimat-Río Uren, Finca de Hermógenes Pereira, Gómez et al. 23760 (MO, PMA); Hacienda Tapexco-Hacienda La Saerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, Davidson & Donahue 8872 (RSA), 8641 (RSA), 8370 (MO, RSA), Davidson et al. 6766 (MO, RSA). Puntarenas: Piedras Blancas-Rincón, 3.7 mi. W of Pan-American Highway, 90–105 m, 8°46'N, 83°18'W, Croat 67652 (MO); 10 km W of highway, 200 m, 79169 (INB). GUATEMALA. Izabal: ca. 7 mi. S of Puerto Barrios, 50 m, Croat 41810 (GUAT, MO). HONDURAS. Atlántida: Parque Nacional Pico Bonito, ca. 10 km SW of La Ceiba, 220 m, 15°42'N, 86°50'W, Evans 1647 (MO); 80–180 m, 15°42'N, 86°51'W, Liesner 26183 (MO). Yoro: Cordillera Nombre de Dios, S of San José de Tesiguat, 250–350 m, 15°30'N, 87°26'W, Daviase et al. 34398 (MO). NICARAGUA. Jinotega: Río Bocay, Caño Litut, somewhat elevated, lateral margins; ovate to ovate-triangular, cordate blades (about equal to the petioles in length); and colorful inflorescences with the spathes bright red on the tube and white on the blade.

175 m, 13°58'N, 85°21'W, *Stevens 16680* (MO). **Zelaya:** Comarca del Cabo, Miguel Bikou, *Robbins 5874* (F, GH, MO, NY); Bluefields, *Neill 2587* (MO); Monkey Point, Caño El Pato, 10 m, 11°35'N, 83°42'W, *Moreno 12464* (MO); ca. 13 km above Kururia, 200 m, *Pipoly 3833* (MO); ca. 13 km above Kururia (ca. 14°39'N, 84°04'W), on road to San Jerónimo (14°42'N, 84°11'W), 200 m, *Pipoly 3835* (MO); 3838 (MO); Colonia Kururia, 50 m, 14°41'N, 84°04'W, *Pipoly 3884* (MO); Cerro Baká, ca. 6.5 km E of Río Coperna, 200–300 m, 13°40'N, 84°30'W, *Pipoly 4967* (MO); near Tala Has and Puente Mango (over Río Kisalya), 40–60 m, 14°41'N, 84°03'W, *Stevens 7627* (MO); Rosita–Puerto Cabezas, ca. 15.7 km SW of Río Kukalaya, 100 m, 13°58'N, 84°12'W, *Stevens 8526* (MO); Mina Nueva América Road, ca. Km 10, *Stevens 12663* (MO); El Empalme–Limbaika, vic. of road to Alamikamba, ca. 25 m, 13°32'N, 84°30'W, *Stevens 12738* (MO); ca. 1.5 km SE of Palmera, 60 m, 13°35'N, 84°20'W, *Stevens 12847* (MO); Bahía de Bluefields, Round Cay, 0–15 m, 11°56'N, 83°45'W, *Stevens 20069* (MO); Bluefields, Río Escondido, *Molina 2019* (F); at junction of road to Alamikamba, ca. 25 m, 13°32'N, 84°30'W, *Stevens 21747* (MO); Río Tronceta, Waspam–Puerto Cabezas, 200 m, 14°43'N, 84°06'W, *Pipoly 4056* (MO); Comarca El Cabo, Pine Savannas, *Molina 15043* (F). PANAMA. **Bocas del Toro:** Chiriquí Grande, near Rambala, 250 m, 8°45'N, 82°15'W, *McPherson 11148* (M, MISSA, MO); Chiriquí Lagoon, von Wedel 1148 (GH, MO). **Canal Area:** Pipeline Road, Km 5.6, *Witherspoon 8606* (MO, SEL); 150 m, *Nee 6389* (MO, RSA, US); 6591 (MO); Summit Gardens, *Croat 10891* (F, MO, NY, SCZ); Frijoles, *Standley 27477* (US); Ft. Randolph, *Standley 28728* (US); Cerro Pelado, 1 km N of Gamboa, 200–220 m, *Nee 7769* (MO, US); Cerro Viejo vicinity, on K16C, *Blum 1273* (FSU, MO, PMA); Camp Piña, Ft. Sherman, *Duke 4427* (MO); Barro Colorado Island, *Kenoyer 184* (MO); 185 (US); *Fairchild 3080* (US); *Croat 6769* (MO); 4529 (MO); 11857 (MO); 11007 (MO); 10911 (MO, SCZ); 9042 (MO); 9410 (MO); *Netting 44* (MO); ca. 400 m, *Wetmore & Abbe 1104* (MO); *Shattuck 33* (F); *Croat 9003* (MO); 6658 (MO, PMA, SCZ). **Coclé:** Llano Grande–Coclesito Road, Mile 12, 200 m, 8°47'N, 80°28'W, *Churchill et al. 3995* (MO); Penonomé–Coclequito, above Río Cascajal, 5.7 mi. N of Llano Grande, 210 m, 8°40'N, 80°26'W, *Croat 67542* (MO); El Valle region, La Mesa, 880 m, *Croat 37574* (MO); 850 m, *Wilbur et al. 15669* (MO). **Colón:** Miguel de la Borda, *Croat 10014* (MO); Nuevo Tonosí–Río Indio, on road from Portobelo and Nombre de Dios, ca. 0 m, *Croat 33539* (MO); Río Indio–Miguel de la Borda, *Croat 36922* (MO); Tyson *et al.* 4546 (MO, SCZ); Santa Rita Ridge Road, Km 21–16, 500–550 m, 9°25'N, 79°37'W, *Knapp 5851* (B, MO); Km 21, 400–500 m, 9°26'N, 79°38'W, *Knapp & Schmalzer 1797* (MO); Km 18–20, 1000–1200 ft., 9°24'N, 79°39'W, *Sytma 2035* (MO, PMA); Km 16–18 km from highway, 300–400 m, 9°26'N, 79°37'W, *Knapp 1773* (CM, K, MO). **Darién:** Puerto St. Dorothea, *Dwyer 2256* (MO); Alturas de Nique Region, S of El Real, near Cana Mine, 550 m, 7°45'N, 77°40'W, *McPherson 11496* (CAS, MO, PMA, RSA); trail NW of Cana, 600 m, *Sullivan 669* (MO); Cerro Pirre region, along trail from base camp to Rancho Frío on slope of Cerro Pirre, 200–450 m, 7°58'N, 77°43'W, *Croat & Zhu 77120* (MEXU, MO, PMA). **Panamá:** Cerro Jefe region, 5 mi. above Inter-American Highway on road to Cerro Azul, *Croat 11526* (F, MO, NY, SCZ, U); 3.5 mi. W of Lago Cerro Azul, *Croat 11566* (MO); 5.8 mi. above Lago Cerro Azul, 840 m, 9°13'N, 79°22'W, *J. Miller & L. Miller 888* (MO, NY); Cerro Jefe region, 10 mi. from

Tocumen Circle, 800–1000 m, *Gentry 2894* (F, MO); 10 mi. from highway, *Croat 15188* (MO); Km 10.1, 300 m, 9°20'N, 79°W, *Croat & Zhu 76548* (MO); near summit, 750–800 m, 9°14'N, 79°22'W, *Croat 67095* (MO); *Witherspoon & Witherspoon 8551* (MO); 770 m, 9°15'N, 79°29'W, *Croat & Zhu 76613* (MO, PMA, US); El Llano–Cartí Road, Km 5–6, 350–375 m, *Croat 348004* (MO); 7 mi. from highway, 460 m, 9°19'N, 79°59'W, *Croat 75107* (MO, PMA); ca. 8 km N of turnoff, 300 m, 9°16'N, 78°57'W, *Churchill 3810* (MO, RSA); 5 km N of highway, 300 m, *Nee 7915* (GH, MO); along trail to Cerro Brewster from Río Pacora Valley, 670 m, 9°20'N, 79°15'W, *McPherson 7503* (MO). **San Blas:** El Llano–Cartí Road, 34–38 km from Pan-American Highway, 100–200 m, 9°25'N, 79°W, *Knapp & Schmalzer 5467* (MO); km 26.5, 200 m, 9°19'N, 78°55'W, *de Nevers et al. 7839* (GH, MO, US); NW of Nusagandi on Sendero Wedar, 150–200 m, 9°15'N, 79°W, *McPherson 11049* (MO). **Veraguas:** Boca de Concepción, in Golfo de los Mosquitos, 10 m, 8°50'N, 81°W, *McPherson 11380* (CM, M, MBM, MO).

**Philodendron gigas** Croat, sp. nov. TYPE: Panama. San Blas: El Llano–Cartí road, 10.1 mi. N of Inter-American Highway, then 1.9 mi. N of Nusagandi, 310 m, 9°20'N, 79°W, 20 July 1994, *Croat & Zhu 76988* (holotype, MO-4619563–69; isotypes, B, CAS, CM, COL, CR, F, K, MEXU, NY, PMA, QCNE, VEN, SCZ, US). Figures 192–196, 205.

Planta hemiepiphytica; internodia 1–2 cm longa, (3)6–10 cm diam.; cataphylla acute D-formata, 30–50 cm longa, in sicco rubrilinea persistentia ut grasso reticulo fibrarum et frustra epidermidia; petiolum semiteres, (5)99–137 cm longus; lamina ovata vel late ovata, cordata vel sagittata basi, (4)81–125 cm longa, (3)37–90 cm lata, in superficie superiore velutina; inflorescentia usque 7; pedunculus 13–16.5 cm longus; spatha 13–16.5 cm longa; lamina spatha alba extus, suffusa rubella atrirubriovaleacea; tubo spathae extus purpurascensivialeacea, intus atriparascensivialeacea; pistilla 4–5-locularia; loculi 20-ovulati; baccae viridilabae aut albae.

Hemiepiphytic, growing to 15–20 m; stem appressed-climbing (or scandent when pre-adult), sap reddish, viscous, leaf scars conspicuous, to 4 cm wide; internodes short, 1–2 cm long, (pre-adult length to 15 cm long), (3)6–10 cm diam., glaucous, olive-green to dark green, coarsely white striate near apex, the epidermis with reddish brown patches and reddish brown raised ridges, becoming brown and scurfy, ridged; roots scurfy, short and blunt, dark reddish brown, 5–11 mm diam., feeder root to 3 cm diam., grayish, sparsely and conspicuously warty; cataphylls 30–50 cm long, to 25–30 cm broad at base (when flattened out), ± D-shaped, unribbed to sharply 2-ribbed (ribs low), greenish white to reddish, thick, juicy, drying reddish brown, persisting semi-intact, eventually as a reticulum of fibers and patches of epidermis; **petioles** (5)99–137 cm long, (1.5)2–3 cm diam., subterete, medium

to dark green, semiglossy, weakly and obtusely flattened adaxially, especially near apex, surface densely and finely short-lineate to faintly striate; **blades** ovate to narrowly ovate, moderately coriaceous, bicolorous, acuminate at apex (the acumen weakly inrolled), cordate to sagittate at base, (44)81–125 cm long, (30)37–90 cm wide (1.4–1.7 times longer than wide), (0.9–1.2 times longer than petiole), subcoriaceous, margins hyaline, upper surface dark green, velvety, matte, lower surface slightly paler, semiglossy to matte, sometimes tinged pale reddish between veins; anterior lobe 33–102 cm long (3.5–6.1 times longer than posterior lobes); posterior lobes 13–26 cm long, 13–24 cm wide, incurved and overlapping with sinus closed on adult plants, obtuse; midrib flat to weakly convex, silvery-white to yellowish green, paler than surface above, narrowly rounded and slightly paler than surface below; basal veins (3)9–12 per side, first and second free to base, most of remainder coalesced to 4 cm, 2 coalesced to 14 cm; posterior rib naked 1.5–4 cm; primary lateral veins (4)7–10 per side, departing midrib at a 40–60° angle, arcuate to the margins, narrowly weakly sunken and slightly paler than surface and marginally discolored above, prominently round-raised, paler than surface below; minor veins weakly visible but not distinct below, arising from both the midrib and primary lateral veins. **INFLORESCENCES** (post-anthesis) to 7 per axil; peduncle 13–16.5 cm long, green, white-striate, especially at apex; prophylls to 22 cm long; **spathe** 13–16.5 cm long, 5 cm diam., moderately constricted above the tube, weakly glossy; spathe blade 5.8 cm long, white, tinged reddish outside, dark red-violet inside; spathe tube oblong-ellipsoid, 7.5 cm long, purplish violet, short white lineate outside, dark violet-purple, short white lineate inside; **spadix** sessile; to 12.5 cm long, ca. 1.5 cm diam., constricted above sterile portion; pistillate portion pale green,  $\pm$  ovoid, 4.2 cm long in front, 3.4 cm long in back, 2.2 cm diam.; staminate portion 11 cm long; fertile staminate portion to 2.5 cm diam. midway; sterile staminate portion 2.4 cm diam., slightly broader than pistillate portion; pistils 4–6 mm long, 1.4 mm diam.; ovary 4–5-locular, with  $\pm$  axile placentation; locules 2.8–5.2 mm long; ovules ca. 20 per locule, 0.1–0.25 mm long, funicle shorter than to equal in length to ovules, style similar to style type B; style crown domed, irregularly lobed; stigma hemispheroid, 1–1.2 mm diam., 0.4–0.7 mm high. **INFRUCTESCENCE** with berries greenish white; seeds 3–4 per locule, 1.6 mm long.

Flowering in *Philodendron gigas* occurs during the rainy season beginning in about July and prob-

ably lasting for more than one month, perhaps as much as two months. Immature fruits have been collected in March.

*Philodendron gigas* is endemic to the Canal Area, where it is known only along the El Llano-Cartí road in both Panamá Province and in the Comarca de San Blas at 300 to 375 m in *Premontane wet forest* and *Tropical wet forest*.

*Philodendron gigas* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Velutina*. This species is distinguished by its appressed-climbing habit; short, thick internodes (6–10 cm diam.); sharply D-shaped, semi-persistent cataphylls; subterete petioles; thick, scaly roots; and especially by its huge, ovate, velvety, dark green leaf blades. It is found growing high on trees, to about 20 m.

*Philodendron gigas* is apparently most closely related to *P. andreanum* Devansaye from Colombia. That species shares with *P. gigas* dark green, velvety leaf blades, but differs in having more elongate ovate-triangular blades with the lobes much longer than broad (vs. rounded and about as long as broad on *P. gigas*). *Philodendron gigas* has blades about 1.6–2.5 times longer than broad, vs. 2.6–3 times for *P. andreanum*. In addition, *P. andreanum* is described as having the spathe tube green outside and becoming whitish within. In contrast, *P. gigas* has the post-anthesis spathe tube purplish both inside and outside.

*Philodendron gigas* is not easily confused with any of the other velvety-bladed species of *Philodendron* from Central America. *Philodendron verrucosum* L. Mathieu ex Schott and *P. squamipetiolatum* Croat have velvety blades, but both have scaly petioles. In South America, two other species with velvety, ovate leaf blades, *P. gloriosum* André and *P. mamei* André, differ in being terrestrial with repent stems. Both are probably restricted to the eastern slopes of the Andes.

*Additional specimens examined.* PANAMA, Panamá: El Llano-Cartí Road, Mile 8.2, 300–350 m, Croat 33680 (B, F, L, MEXU, MO, PMA, QCA, US); Mile 5–6, 350–375 m, Croat 34778 (F, MO); Mile 10, Croat 33714 (F, MO). San Blas: El Llano-Cartí road, Mile 14, 300 m, 9°15'N, 79°W, Croat 69242 (CM, MO).

***Philodendron glanduliferum*** Matuda, Bol. Soc. Bot. México 27: 47. 1962. TYPE: Mexico, Oaxaca: Sierra de Juárez, along Hwy. 175, 1900 m, 15 Sep. 1961, Matuda 37247 (holotype, MEXU). Figures 180, 200–202.

Terrestrial; stem to 1 m long; internodes short, to 8 cm long, to 2 cm diam., about as long as broad

or sometimes broader than long, dense; roots drying 2–3 mm diam., dark reddish brown, folded into irregular longitudinal ribs; cataphylls fleshy, sharply 2-ribbed, reddish, drying pale yellowish brown to brown, persisting semi-intact as a reticulum of fibers which are persistent; **petioles** 44–51 cm long, 3–6 mm diam., subterete, obtusely flattened near base adaxially, often purple at apex, surface inconspicuously short-lineate to striate, prominently so toward apex, with moderately dense, hair-like scales, at least near apex, scales many times longer than wide; geniculum darker than petiole; **blades** ovate-cordate, subcoriaceous, bicolorous, abruptly acuminate at apex (the acumen sometimes apiculate, 2–4 mm long), cordate at base, 34–47 cm long, 1.2–1.4 cm wide (1.2–1.4 times longer than wide), (0.6–0.8 times the petiole length), upper surface semiglossy, lower surface glossy, paler; anterior lobe 24–36 cm long, 11–28(38) cm wide (0.9–1 times longer than wide), (1.9–2.7 times longer than posterior lobes); posterior lobes 11–14.5 cm long, broadly rounded to obtuse; sinus spatulate or V-shaped; midrib flat to convex, paler than surface above; basal veins 5–7 per side, with 1–2 free to base, 2–3 coalesced (0.7)1–2.2 cm; posterior rib not naked, to 1.5 cm; primary lateral veins 2–4 per side, departing midrib at a 50–65° angle, sunken, raised below; minor veins arising from both the midrib and primary lateral veins; tertiary veins moderately distinct, darker than surface below. **INFLORESCENCES** 1–2 per axil; peduncle 9 cm long, 6 mm diam.; **spathe** smooth, 12–13 cm long (1.5–2.6 times longer than peduncle); spathe blade pinkish white outside; spathe tube magenta outside; **spadix** to 12 cm long, remaining in the spathe at anthesis; pistillate portion to 2 cm long, 10 mm diam.; staminate portion to 10 cm long; fertile staminate portion gradually tapered to apex, 10 mm diam. at constriction, scarcely constricted above the sterile staminate portion, narrowly rounded at apex, 11 mm diam. near the middle, 10 mm diam. at constriction; pistils 2 mm long, 1.9 mm diam.; ovary (5)6(7)-locular, locules 1.3 mm long, 0.6 mm diam., with axile placentation; ovules ca. 10 per locule, 2-seriate, 0.5 mm long, longer than funicle; funicle 0.3 mm long, adnate to lower part of partition; style similar to style type D; style apex somewhat rounded to flat; the androecium truncate, margins irregularly 4–5-sided, ca. 0.9 mm long. **JUVENILE** plants with lower blade surface purplish.

Flowering in *Philodendron glanduliferum* occurs during the rainy season, in June and September (Moore & Bunting 8889). Fruiting collections have not been seen.

*Philodendron glanduliferum* occurs in Mexico, Guatemala, and Venezuela, with *P. glanduliferum* subsp. *glanduliferum* endemic to Mexico and Guatemala. In Mexico, it is known only from the Sierra de Juárez, at 580 to 1900 m elevation in "Bosque mesófilo." It has been collected recently in Guatemala in the Sierra de las Minas in the province of Zacapa. *Philodendron glanduliferum* subsp. *camiloanum* Croat is endemic to Venezuela, where it is known only from the states of Táchira and Apure at 250 to 1200 m elevation in *Tropical wet forest* and *Premontane wet forest* life zones.

*Philodendron glanduliferum* is a member of *P.* sect. *Philodendron* subsect. *Achyropodium*. This species is characterized by its terrestrial habit; short internodes; sharply two-ribbed cataphylls persisting semi-intact as a reticulum of pale yellowish brown fibers; subterete petioles with an area of moderately dense hair-like glands at least near the apex; and ovate-cordate blades (about three-fourths as long as petioles) with a spatulate or V-shaped sinus (often overlapping in *P. glanduliferum* subsp. *camiloanum*).

*Philodendron glanduliferum* subsp. *glanduliferum* is not easily confused with any other species in Central America. *Philodendron glanduliferum* subsp. *camiloanum*, endemic to Venezuela, is distinguished from the typical subspecies by having sharply 2-ribbed cataphylls; petioles glandular more or less throughout; and blades with the sinus closed or nearly so with the margins of the lobes overlapping (at least slightly). In contrast, *P. glanduliferum* subsp. *glanduliferum* has unribbed or only weakly 2-ribbed cataphylls, petioles glandular only near the apex, and blades with the sinus V-shaped and the lobes not at all overlapping.

*Philodendron glanduliferum* may be confused with *P. ornatum* Schott where they occur together in Venezuela. The latter species is distinguished by its appressed-climbing epiphytic habit and typically larger leaves with the petioles merely warty-verrucose (never glandular) near the apex.

**Additional specimens examined.** GUATEMALA. **Isabul:** Sierra de las Minas, 700 m, Forther s.n. (M). MEXICO. **Oaxaca:** Ixtlán, S side of Río Soyalapan, 1230–1260 m, 17°34'40"N, 96°20'W, Boyle et al. 2583 (CM, MO); Comaltepec, Highway 185, trail from Puerto Antonio to Cuaje, 1280–1725 m, 17°39'50"N, 96°18'05"W, Boyle & Acosta 3980 (MO); Valle Nacional–Oaxaca, 660 m, Croat 78709 (CHIP, MO); 13.9 mi. above bridge at Valle Nacional, 2.6 mi. S of Metates, 1165 m, Croat 47936 (F, MO); 660 m, 39753 (MO); 1220 m, 43909 (F, MO); 39780 (MO); 4.3–6 mi. above bridge, 705 m, 17°44'N, 96°19'W, Croat & Hannan 65544 (B, MEXU, MO, US); 6–14 mi. from bridge, 580 m, Moore & Bunting 8889 (BH, MO); Coatlán–Mazatlan, Cerro Carnero, MacDougal 335 (MEXU); 409 (MEXU); Tuxtpec–Oaxaca, 55 km S of Tuxtpec, 409 (MEXU); Tuxtpec–Oaxaca, 55 km S of Tuxtpec, 409 (MEXU); Tuxtpec–Oaxaca, 55 km S of Tuxtpec, 409 (MEXU).

tpec, 630 m, 17°37'N, 96°20'W, Hammel & Merello 15481 (MO); Tuxtpec, Tuxtpec-Oaxaca, 4 km SE of Metates, Torres et al. 7811 (MO).

***Philodendron grandipes*** K. Krause, in Engl. & K. Krause, *Das Pflanzenr.* IV. 23Db (Heft 60): 48. 1913. TYPE: Panama. Colón: Rfo Fató (Pató) at Dos Bocas (E of Nombre de Dios), 40–80 m, ca. 9°35'N, 79°28'W, Pittier 4228 (holotype, US). Figures 203, 204, 206, 216.

*Philodendron pleistoneurum* Standl. & L. O. Williams, *Ceiba* 3. 109. 1952. TYPE: Costa Rica. Puntarenas: Esquinas Forest, 60 m, 27 Mar. 1951, Allen 6036 (holotype, EAP; isotypes, GH, US).

Terrestrial; stem creeping over soil, 20–100 cm long, leaf scars obscured by cataphylls; internodes short, 1–4 cm long, 2.8 cm diam., usually broader than long, medium green, semiglossy, coarsely white-streaked at apex; roots  $\pm$  smooth, slender, elongate, 2–4 mm diam., few per node, descending; cataphylls moderately coriaceous, 16–22 cm long, sharply 2-ribbed, green to pale green or reddish or pinkish, drying brown to tan, persisting semi-intact or as fibers at lower nodes, acuminate at apex. LEAVES arching to pendent; **petioles** 25.5–73 cm long, 10–12 mm diam., D-shaped, spreading, pale to medium green to reddish at base, with a medial rib adaxially, rounded abaxially, with adaxial margins erect, surface finely and weakly striate; sheathing 3.5–5.5 cm long; geniculum slightly thicker than petiole when apparent, 1.4–1.7 cm long, paler than petiole; **blades** broadly ovate-cordate, subcoriaceous, moderately bicolorous, acute to abruptly acuminate at apex (the acumen apiculate and downturned), cordate at base, 20–50 cm long, 15.5–36 cm wide (0.9–1.7 times longer than wide), (0.5–0.9 times the petiole length), broadest at or near the middle, margins broadly undulate, upper surface dark green, semiglossy to glossy, sometimes matte or subvelvety, drying brown to greenish brown, lower surface semiglossy, moderately paler, drying brown to greenish brown; anterior lobe 16–37 cm long, 15–38 cm wide (1.8–3.7 times longer than posterior lobes), broadest slightly above or at point of petiole attachment; posterior lobes 5.5–15.5 cm long, 7.4–18.6 cm wide, directed downward, usually held up somewhat at an angle from the midrib, rounded to obtuse; sinus parabolic to hippocrepiform, sometimes closed with lobes overlapping before being pressed; midrib concolorous, flat at base, becoming weakly sunken toward apex above, thicker than broad at base, becoming convexly raised, concolorous below; basal veins 8–10 per side, 1(2) free to base, several remaining coalesced 0.5–2(4) cm, prominently sunken above,

convexly raised below; posterior rib usually not naked, sometimes briefly so at base; primary lateral veins (8–10)11–17 per side, departing midrib at a 20–30° angle, spreading to 50–70° angle,  $\pm$  straight to the margins, quilted-sunken and concolorous above, prominently convex, matte, and darker than surface below; interprimary veins about as conspicuous as primary lateral veins above, flat and darker than surface below; minor veins arising from both the midrib and primary lateral veins; tertiary veins distinct, weakly raised above, darker than surface below. INFLORESCENCES erect, 2–4 per axil; peduncle 2.5–14 cm long, 5–8 mm diam., terete, green to reddish, white-lineate; **spathe** 6.6–11.8 cm long (0.6–2.5 times longer than peduncle), constricted  $\pm$  midway; spathe blade lanceolate, pinkish red, tinged green, pale green to white or green, short white lineate outside, 4–6 cm long (opening broadly elliptic in face view), pinkish red to white, pale greenish white or pale green inside; spathe tube oblong-ellipsoid, dark reddish maroon to reddish purple to green, weakly short dark lineate outside, 2.5–4 cm long, 1.4–3.8 cm diam., pale green to greenish white to pinkish red inside; **spadix** sessile or very short stipitate; tapered, 6.6–11.8 cm long, broadest near the base; pistillate portion white, yellowish white (post-anthesis), slightly tapered toward the apex and base, (1.6)2–2.7 mm long, 9–11 mm diam. at apex, 1–1.5 cm diam. at middle, (6)10–11 mm wide at base; staminate portion 3.5–7.1 cm long; fertile staminate portion white, cylindrical to tapered, 7–10 mm diam. throughout, 1–1.5 cm diam. at base, 7–10 mm diam. at middle, 4–6 mm diam. ca. 1 cm from apex, narrower than the pistillate portion, as broad as the sterile portion; sterile staminate portion about as the pistillate portion, white, 1–1.5 cm diam.; pistils 1.8–2.3 mm long, 1.1–1.5 mm diam.; ovary (4)5–6-locular, 1.3–1.5 mm diam., locules 1–1.5 mm long, 0.4–0.6 mm diam., with axile placentation; ovules (7–10)16–22 per locule, 2-seriate, 0.2–0.4 mm long, longer than funicle, style similar to style type B; style apex  $\pm$  concave, sometimes weakly lobed; stigma subsdiscoid, brushlike, lobed or unlobed, 0.9–1.4 mm diam., 0.2–0.5 mm high, covering center of style apex, depressed medially; the androecium truncate, margins irregularly 3–6-sided; thecae oblong, 0.2–0.3 mm wide; sterile staminate flowers blunt, margins irregularly elongate, 13–18 mm long, 9–13 mm wide. Berries creamy white, obovoid, apex truncate; seeds 20 per locule, 1.9 mm long, 0.2 mm diam., with prominent striations running from funicle to apex and slightly spiraling, perpendicular to larger veins. JUVENILE

blades narrowly elliptic to ovate, acute to weakly cordate at base.

Flowering in *Philodendron grandipes* probably occurs throughout most of the year (February through November, except October) but primarily in July and August. Post-anthesis collections have also been made throughout the year, and immature fruiting collections are known from throughout the year (except February, March, and June).

*Philodendron grandipes* ranges from Nicaragua (Zelaya) to Panama, Colombia, and Ecuador (Esmeraldas), from near sea level to mostly less than 750 m (sometimes to 1200 m) elevation in *Tropical moist forest*, *Premontane wet forest*, and *Tropical wet forest* life zones.

*Philodendron grandipes* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is characterized by its terrestrial habit, short internodes, persistent cataphyll fibers, broadly ovate-cordate leaf blades, usually green spathes, and, especially, by its D-shaped petioles with erect margins and a medial adaxial rib.

*Philodendron grandipes* is closest to *P. jodaviansum*, which has a similarly shaped petiole and other general features in common. The latter species differs in being an appressed-climbing hemi-epiphyte with ovate-triangular (rather than broadly ovate) blades.

This species frequents stream banks in central Panama and is one of the few consistently terrestrial *Philodendron* species in Central America. Habitat in Costa Rica, however, is variable. *Philodendron grandipes* occurs along stream banks on the Osa Peninsula, but at La Selva (Heredia) this species is widely scattered in the forest understory (M. Grayum, pers. comm.).

Spathe tube color in this species is variable to some extent geographically. For example, on the Atlantic slope of Costa Rica the spathe tube is usually greenish, whereas on the Pacific slope it is generally reddish on both surfaces.

*Additional specimens examined.* COSTA RICA. **Alajuela:** 36–37 km NW of San Ramón, 500–515 m, 10°15'N, 84°34'W, *Croat* 68198 (MO); Río Chiquito, Upland, 800 m, *Gómez* 18639 (MO, PMA). **Heredia:** Tirimbina, 230 m, *Proctor* 32347 (LL); 1 km SE of Tirimbina, 220 m, *Lent* 2136 (CR, F, MEXU, MO); near Río Bijagual, 2 km E of Tirimbina, 150 m, *Maas* 1329 (U); Porto Viejo-Río Sucio, 20 m, *Croat* 35676 (CR, MO); La Selva Field Station, 100 m, *Grayum* 3028 (DUKE), *Folsom* et al. 8806 (DUKE), *Hammel* 9126 (DUKE), *Sperry* 855 (DUKE), *Croat* 44254A (MO), 44224 (MO), *Cannarella* 8806 (DUKE), *McDowell* 687 (MO), *Grayum* et al. 1758 (F, MO), *Jimenez IV* (MO), *VII* (MO). **Limón:** Braulio Carrillo-Gudipiles, 250–270 m, *Croat* 78745 (CR, INB); Toro Amarillo, 10 km W of Gudipiles, 300 m, *Lent* 73 (F); Ha-

cienda Tapezco-Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, *Davidson* & *Donahue* 8741 (MO, RSA); Turrialba-Limón, along Hwy. 32, ca. 11 mi. S of Siquirres, 650 m, *Croat* 43330 (MO); Río Chirripó-Río Corinto, N of Quebrada El Molinete, 400 m, 10°12'N, 83°54'W, *Grayum* & *Jacobs* 3525 (MO); Quebrada Cañabral-Río Barbilla, 200–400 m, 10°02'N, 83°26'W, *Grayum* et al. 8739 (CR, MBM, MO); Hitoy Cerere Reserve, 140 m, 9°42'N, 83°02'W, *Hammel* & *Grayum* 14349 (MO); Parque Nacional Tortuguero Lomas de Sierpe, near Río Sierpe, 100 m, 10°24'N, 83°33'W, *Robles* et al. 2006 (CR, G, MO); Río Blanco, Río Frío-Limón, W of Gudipiles, Quebrada Danta, 360 m, 10°12'N, 83°49'W, *Croat* 68419 (MO); Río Catarata, 50–100 m, 9°37'N, 82°49'W, *Burger* & *Antonio* 10888 (CR, F, MO, PMA, U); Río Colorado, 16 airline km SW of Barra del Colorado, 10–120 m, 10°39'N, 83°40'40"W, *Davide* & *Herrera* 31214 (CR, K, MO); Río Sixasola, ca. 3 mi. NE of Bratsi, 15 m, *Croat* 43253 (CR, MO); 0.3 mi. E of Bratsi, ca. 35 mi. SE of Limón, 30 m, *Croat* 43289 (MO); Limón-Shiroles, 6.5 mi. SW of Bratsi, 50 m, *Croat* 43299 (CR, MO); Pocot, Barra del Colorado, Llanura de Tortuguero Saránas, 15–20 m, 10°38'38"N, 83°44'10"W, *Araya* 596 (INB). **Puntarenas:** Osa Peninsula, Piedras Blancas-Rincón, 3.7 mi. W of Pan-American Highway, 90–105 m, 8°46'N, 83°18'W, *Croat* 67651 (K, MO); 2.5 mi. SW of Rincón, 8°42'N, 83°29'W, *Kennedy* 1622 (MO); *Croat* 76751 (MO); Rincón de Osa, 250–540 m, 8°42'N, 83°31'W, *Croat* & *Grayum* 59838 (MO); SW of Rincón de Osa, 40–200 m, 8°42'N, 83°30'W, *Grayum* et al. 7555 (CR, M, MO); Fila Costeña, 950–1150 m, *Grayum* 10651 (CR, INB); Corto-La Unión, near San Miguel, *Croat* 26519 (MO); Finca Loma Linda, 1 mi. SW of Cañas Gordas, 1150 m, *Croat* 22262 (MO); *Allen* 6036 (F, GH); Finca El Edén, km 183, ca. 400 m from Santa Marta, *Gómez* 22954 (MO); Corcovado National Park, Sirena, 0–200 m, 8°29'N, 83°36'W, *Liesner* 2903 (MO); 5–25 m, 8°29'N, 83°34'W, *Hammel* & *Kernan* 16662 (CR, MO); 150 m, *Kernan* & *Phillips* 512 (CR, MO); Rancho Quemado, 250–350 m, *Marín* 39 (CR, INB, MO); vic. Boscosa, 50 m, *Croat* & *Hanson* 79244 (MO); Las Cruces Tropical Botanical Garden, 1200 m, 8°49'N, 82°58'W, *Croat* 57234 (CR, MO); Río Jaba, Las Cruces, 1200 m, *Meerow* et al. 2017 (SEL). **San José:** San Isidro del General-Dominical, SW of San Isidro, 4.8 mi. from Río Pacuare, 1000 m, *Croat* 35215 (MO); 9 mi. SW of Río Pacuare, 680 m, *Croat* 35346 (MO); 12 km SW of San Isidro de El General, along CR-223, 900–1000 m, *Uiley* & *Uiley* 4925 (MO); 14.5 km S of San Isidro del General, 500 m, *Kress* & *Gómez* 88-2467 (SEL); Zona Protectora La Cangreja, ca. 2 km NNE of Mastatal de Puriscal, 400–540 m, 9°42'N, 84°22'W, *Grayum* 8644 (MO); Parque Nacional Braulio Carrillo, above Río Sucio, 5–600 m, *Pennington* 11533 (K); 1100 m, 10°24'04"N, 85°03'03"W, *Carballo* 87 (CR, MO). **NICARAGUA.** **Zelaya:** 150–180 m, 11°43'N, 84°18'W, *Stevens* 4965 (MO); Río Rama, at Salto La Oropendula, 15–25 m, 11°57'N, 84°17'W, *Stevens* 8960 (MO). **PANAMA.** **Bocas del Toro:** Chiriquí Grande-Fortuna, 13.2 mi. W of Chiriquí Grande, 310 m, 8°45'N, 82°10'W, *Croat* & *Grayum* 60141 (MO); ca. 10 km SW of Chiriquí Grande, ca. 300 m, 8°52'N, 82°10'W, *Thompson* 4938 (CM, MO); 3.2 mi. N of Divide, 700 m, 8°45'N, 82°15'W, *Croat* 60264 (CM, MO); 450 m, 8°45'N, 82°15'W, *McPherson* 7369 (MO); 3 mi. N of Continental Divide, 650 m, 8°47'N, 82°11'W, *Churchill* & *Churchill* 6209 (MO); 6210 (MO); Chiriquí Lagoon, von Wedel 1049 (F, GH, MO). **Canal Area:** Gamboa, *Standley* 28401 (US); Frijoles, *Standley* 27471 (US);

Barro Colorado Island, *Standley 31361* (US); *40888* (US); *Croat 16574* (MO); *12300* (MO, SCZ); *11886* (MO); *11194* (MO); *11077* (MO); *9526* (MO); *6512* (MO); *5117* (MO); *Schmaltz 549* (MO); Summit Gardens, *Croat 11491* (MO). **Chiriquí:** Chiriquí Grande-Fortuna, 7.7 mi. W of Chiriquí Grande, 80 m, 8°50'N, 82°10'W, *Croat & Grayum 60114* (MO); Burica Peninsula, San Bartolomé Limite, 12 mi. W of Puerto Armuelles, 400–500 m, *Croat 22188* (MO). **Coclé:** La Junta-Limón, 5 hours walk N of Alto Calvario, 800–1000 m, *Folsom 5861* (MO); Coclesito-Llano Grande, 200 m, 8°47'N, 80°28'W, *Churchill et al. 4170* (MO); El Valle region, 800–900 m, 8°36'N, 80°07'W, *Croat & Zhu 76664* (MO, PMA); *67212* (F, MO); 800 m, *25406* (F, MO); ca. 1000 m, *Gearty 5662* (F, MO); *Croat 14391* (MO); 860 m, 8°37'N, 80°08'W, *Croat & Zhu 76710A* (MO); 860–900 m, *Croat 37398* (MO, RSA); Cerro Pilón, *Duke & Dyer 13977* (MO). **Colón:** Portobelo-Nombre de Dios, 6–8 km from Peluca Hydrographic Station, *Kennedy & Dressler 3331* (US); Portobelo-Río Cascajal, vic. of Nuevo Tonosí, *Croat 33648* (K, MO, US); 6 mi. S of Portobelo, *Croat 11401* (MO); Portobelo-Nombre de Dios, 1.2 mi. beyond junction of road to Isla Grande, 9°40'N, 79°35'W, *Croat 49795* (MO); Altos de Pacora-Cerro Brewster trail, 700 m, 9°18'N, 79°16'W, *de Nevers et al. 6234* (MO); Río Guanche, 0.5–1 km upstream from Puerto Pilón-Portobelo road, 6 km S of Portobelo, 5–30 m, *Nee 7153* (MO, US); ca. 3–5 mi. inland, 10–100 m, *Croat 26205* (MO); 30–100 m, *79345* (PMA, MO); 1.5 mi. upstream, ca. 10 m, *Kennedy & Dressler 1513* (SEL); <100 m, 9°27'N, 79°40'W, *Croat & Zhu 76245* (MO); Río Miguel de la Borda, vic. of Guásimo, *Croat 9940* (MO). **Darién:** Cerro Pirre region, vic. Cama gold mine, 500–600 m, *Croat 37637* (MO); 480 m, *37957* (MO); 17 km N of El Real, trail from base camp along Río Perisencio, 100 m, 8°01'N, 77°44'W, *Croat & Zhu 77177* (MO); Río Cocalito, *Whitford & Eddy 132* (BM); Río Tuquesa, ca. 2 km by air from Continental Divide, vic. of Tyler Kittredge gold mine, *Croat 27191* (MO). **Panamá:** El Llano-Cartí Road, 9.6 km from Pan-American Highway, 410 m, *Mori & Kallunki 1835* (MO, PMA); Km 12.4, 300–400 m, *Folsom et al. 6178* (MO, PMA); Km 13.2, 370 m, 9°16'N, 78°57'W, *Hooser 1313* (MO); Mile 5–6, 350–375 m, *Croat 34902* (MO); vic. of Gorgas Mosquito Control Project Site at km 12, *26058* (MO); Serranía de Majé, S of Ipetí, 500–650 m, *Hufl et al. 1692* (MO); Serranía de Cañazas, Rancho Chorro, above Tortí Arriba, 400–700 m, *Folsom 6748* (MO); Cerro Campana, ca. 1 mi. from Inter-American Highway, ca. 150 m, *Croat 35969* (MO, RSA). **San Blas:** El Llano-Cartí Road, trail along Continental Divide, 400 m, 9°20'N, 78°56'W, *McDonagh et al. 323* (BM); SE of Puerto Obaldía, *Croat 16762* (MO); 14 mi. N of Pan-American Highway, 300 m, 9°15'N, 79°W, *Croat 69249* (MO); Miria Ubigandup Island, Digole, 0–20 m, 9°26'N, 78°54'W, *Herrera 292* (MO); Río Playón Chico, vic. NEBA DUMMAT, 100–450 m, 9°14.5'N, 78°15'W, *Herrera et al. 1399* (MO, PMA); 1 mi. S of Nusagandi, Mile 9, 350 m, 9°20'N, 79°W, *Croat & Zhu 76997* (MO, PMA); Nusagandi, 275–300 m, 9°20'N, 79°W, *Croat 76594* (MEXU, MO). **Veraguas:** Santa Fe region, Cerro Tute, 1050–1150 m, *Croat 48907* (MO); Escuela Agrícola Alto Piedra-Calovébora, Río Dos Bocas Valley, N of Santa Fe, 350–400 m, *Croat 27398* (MO); 450 m, *27550* (MO); Escuela Agrícola Alto Piedra-Río Dos Bocas, ca. 10 km from the school, 530–620 m, *Croat 25903* (MO).

***Philodendron granulare*** Croat, sp. nov. TYPE: Panama. Darién: Río Pirre, 14 July 1971, *Croat & Porter 15543* (holotype, MO-2059344; isotype, PMA). Figures 207, 208.

Planta hemiepiphytica; internodia 3.5–4 cm longa, ca. 1 cm diam.; cataphylla acuta 2-costata, 15 cm longa, decidua; petiolus 8–9 cm longus, 7–8 mm diam., convexus adaxialiter cum 2 costis marginalibus; lamina oblongo-oblancoolata, debiliter cordata basi, 19–28 cm longa, 4–6 cm lata; nervis primariis lateralibus 5–6 utroque; inflorescentia 1; pedunculus 10.5 cm longus; spathe usque 15.7 cm longa, viridis; lamina spathe intus viridi; tubo spathe extus suffuso rubro, intus rubro; pistilla 3–5-locularia; loculi 1-ovulati.

Hemiepiphytic vine; internodes semiglossy, 3.5–4 cm long, ca. 1 cm diam., longer than broad, drying light brown, epidermis longitudinally deeply ridged, finely striate on magnification and densely granular to almost scabrous (but not harsh to the touch), the vestiture raised to ca. 1.5–2 times longer than wide; roots drying dark brown, <15 cm long, 1 mm diam., few per node; cataphylls 15 cm long, sharply 2-ribbed (ribs low), green, deciduous. LEAVES with petioles 8–9 cm long, 7–8 mm diam., convex with acute margins adaxially, drying dark brown and somewhat sulcate adaxially with distinct acute marginal rib; sheath short, inconspicuous, 1–2 cm long, for <0.25 its petiole length; blades oblong-oblancoolate, acuminate at apex, weakly cordate at base, 19–28 cm long, 4–6 cm wide (3.8–4.7 times longer than wide), upper surface medium green, weakly glossy, drying dark brown, lower surface slightly paler, drying yellowish brown; midrib flat, concolorous above, convex, slightly paler than surface below; basal veins 1–2 per side, with 1–2 pairs free to base, indistinct; primary lateral veins 5–6 per side, weakly sunken above, convex, slightly paler than surface, drying paler below; minor veins moderately obscure, barely visible on drying, the larger weakly undulate, arising from the midrib only, secretory canals appearing on magnification as short, pale raphide cells along the minor veins. INFLORESCENCES (post-anthesis), 1 per axil; peduncle 10.5 cm long, 1 cm diam.; spathe to 15.7 cm long, to 7 cm wide when flat, weakly constricted above the tube, green throughout; spathe blade green inside with prominent striations from lower 1.5 cm to 9 cm up from base; spathe tube tinged red outside, red inside; fertile staminate portion to 5.5 cm long, 9 mm diam.; spadix with pistils 1.5–3 cm diam., drying light brown, sparsely granular on the sides; ovary 3–5-locular; ovules with 1 per locule, style similar to style type E; style funnel 0.5–0.7 mm, distinctly raised above the surface. INFRACTESCENCE 10

cm long, 2.7 cm diam. in lower  $\frac{1}{2}$ , prominently tapered toward apex to 1.3 cm diam. at apex; berries whitish; seeds oblong-elliptic, tan, smooth, 1.3–1.9 mm long.

Flowering in *Philodendron granulare* probably occurs in the early wet season. Fruiting is recorded by a single collection made in July.

*Philodendron granulare* is known only from the Panamanian type specimen, from Darién Province near El Real along the Río Pirre at less than 75 m elevation in *Premontane wet forest*.

*Philodendron granulare* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is distinguished by its scandent habit, more or less oblong, short-petioled blades with subcordate bases, and especially by its densely granular dried stem (hence the name) and distinct funnel-shaped styles.

*Philodendron granulare* is most easily confused with *P. bakeri*, which it resembles in a superficial way. The latter species differs in having totally smooth stems, petioles with conspicuous sheaths often extending to the middle or beyond, leaf blades with more conspicuous minor veins, and pistils with a truncate, type B style which lacks a funnel, having instead the styler canals emerging directly onto the flat truncate apex.

***Philodendron grayumii* Croat, sp. nov. TYPE:**

Panama. Bocas del Toro: Fortuna Lake area, Fortuna-Chiriquí Grande, 0.3 km N of Continental Divide, 970 m, 8°43'N, 82°17'W, 27 June 1994, *Croat & Zhu 76524* (holotype, MO-4619417-20; isotypes, AAU, B, CAS, CM, CR, COL, DUKE, F, GH, K, M, MEXU, NY, PMA, S, SCZ, SEL, TEX, US). Figures 209–215.

Planta hemiepiphytica; internodia 1–10 cm longa, (2.5)3.5–7 cm diam.; cataphyllum 11–26 cm longa, incostata vel leniter 1-costata, demum decidua aut acuta complanta vel 2-costata; petiolus 40–97 cm longus, subteres, obtuse planatus prope apicem; lamina ovata vel late ovata, (31)50–73 cm longa, 24–44(58) cm lata; cordata basi; nervis basalibus 4–5(6) utroque; nervis lateralibus 1–4–6 utroque; inflorescentia 1–5 per axillam; pedunculus (2)4–6 cm longus; spathe (11)13–16(18) cm longa, extus pallide viridis vel virens, albida vel pallide flava vs. apicem, minimum interdum maculata purpurea-violacea, tubo spathae marronino vel rubro intus; pistilla 7–8-locularia; loculi (2)3–4(5)-ovulati.

Hemiepiphytic, appressed-climbing; internodes semiglossy to matte, 1–10 cm long, (2.5)3.5–7 cm diam., dark green to gray-green or tannish gray, drying brown to yellow-brown or reddish brown, epidermis closely ridged or fissured, sometimes

peeling; roots usually 1–2 per node, ca. 3–4 mm diam., drying dark reddish brown; cataphylls thick, somewhat spongy, 11–26 cm long, unribbed to weakly 1-ribbed near apex or sharply flattened to sharply 2-ribbed (ribs close), green, sometimes weakly maroon-spotted, often persisting intact for a short time then deciduous, intact; **petioles** 40–97 cm long, 1–1.5 cm diam., subterete, firm, dark to medium green, obtusely and often weakly flattened near apex, sometimes weakly ribbed near apex adaxially, surface semiglossy, weakly and densely short pale green lineate, drying yellowish brown to dark brown, smooth to finely striate, sheath 6–15 cm long; **blades** ovate to broadly ovate, coriaceous to subcoriaceous, moderately bicolorous, semiglossy, acute to gradually or abruptly acuminate at apex, deeply cordate at base, (31)50–73 cm long, 24–44(58) cm wide ((0.7)1–1.9 times longer than wide), three-fourths as long as to somewhat longer than the petiole, upper surface dark green, drying dark brown to yellow-brown or grayish brown, semiglossy, lower surface conspicuously paler, drying yellow-brown, weakly to semiglossy; anterior lobe (25)31–49(60) cm long, (13)24–32(42) cm wide (2.3–4.6(3) times longer than posterior lobes), margins broadly rounded, sometimes sinuate; posterior lobes rounded to narrowly rounded, sometimes overlapping, (6)9–13(17) cm long, 12–18(22) cm wide; sinus hippocrepiform to oblong spatulate, obovate, or sometimes triangular to parabolic, rounded to narrowly rounded, (8)10.5–13 cm deep; midrib flat to broadly raised above, paler than surface above, convex to obtusely acute and sometimes purplish spotted below; basal veins 4–5(6) per side, with 1–2 free to base, third and higher order veins coalesced 3–6(8) cm long; posterior rib usually not naked, rarely naked for 1.5 cm; primary lateral veins 4–6 per side, departing midrib at a (45)55–70(75°) angle, downturned and splayed out when joining midrib, obtusely to narrowly sunken and paler than surface, sometimes weakly raised, drying weakly ridged near the midrib above, convex and paler than surface, sometimes reddish below; minor veins moderately indistinct, drying weakly prominent (surface often with minute purplish raised areas seen on high magnification), arising from both the midrib and primary lateral veins, secretory ducts appearing as intermittent lines (less conspicuous on highland forms); **INFLORESCENCES** 1–5 per axil; peduncle (2)4–6 cm long, 1–1.5 cm diam., pale green, finely short-lineate; **spathe** (11)13–16(18) cm long, 2–3 cm diam. (2.1–2.6(3.5) times longer than peduncle), weakly constricted  $\pm$  above the middle, pale to medium green, sometimes whitish to pale yellowish toward apex, often tinged or



spotted violet-purple throughout, especially near base, sometimes green dorsally with only the front edges of tube maroon, margins white, outer surface drying reddish brown throughout inside, violet-purple to maroon in lower one-half to two-thirds, especially in lower half; spathe tube 7–8.5 cm long, 2–3.6 cm diam., maroon to red, with pale lineations extending onto blade area inside; **spadix** sessile; cylindrical, 9.5–16 cm long, broadest above the middle or below the middle; pistillate portion pale green, cylindrical to weakly tapered toward the apex, 3.3–3.5 cm long, 4.6 cm long in front, 3.2–3.5 cm long in back, 9–20 mm diam. throughout; staminate portion 6.3–12.5 cm long; fertile staminate portion creamy white, cylindrical, 1–1.4 cm diam. at base, 1–1.4 cm diam. at middle, 8–10 mm diam. ca. 1 cm from apex, broadest at the base or at the base broader than or as broad as the pistillate portion, sterile staminate portion usually not detectable; pistils 1.4–1.9 mm long, 0.8–1.1 mm diam.; ovary 7–8-locular, 1–1.4 mm long, 0.8–1 mm diam., walls thin and membranous, locules 1–1.3 mm long, 0.3–0.4 mm diam., ovule sac sometimes present, to 0.8 mm long, with sub-basal placentation; ovules (2)3–4(5) per locule, 2-seriate, contained within transparent, gelatinous ovule sac, 0.3–0.5 mm long, usually as long as funicle; funicle 0.2–0.3 mm long (can be pulled free to base), style 0.5 mm long, 1.2 mm diam., similar to style type B; style apex sloping; stigma brushlike, sometimes cupulate, subdiscoid, unlobed, sometimes truncate or weakly domed, 0.9–1.3 mm diam., 0.2–0.6 mm high, covering entire style apex; the androecium truncate, prismatic, oblong, margins irregularly 4–6-sided, 0.9 mm long, 1.1 mm diam. at apex; thecae cylindrical to weakly elliptical, 0.3–0.4 mm wide,  $\pm$  parallel to one another; sterile staminate flowers trapezoidal to 4–6-sided, 1.1 mm long, 1.3 mm wide. **INFRACTESCENCE** with peduncle to 11.5 cm long; spathe to 26 cm long, pistillate spadix to 13.5 cm long, 2.5 cm diam.; staminate spadix to 13.5 cm long; berries pale ochraceous, 3–3.3 mm long; seeds 1.9–2 mm long, 0.7–1 mm diam. **PRE-ADULT** petioles sheathing throughout; blades oblong, subcordate, 26–31 cm long, 9–13 cm wide.

Flowering in *Philodendron grayumii* apparently occurs primarily in the rainy season (March as well as June and September), but also in the late dry season. Post-anthesis inflorescences have been collected in March and June through August and November. Immature fruits are known from June and November.

*Philodendron grayumii* ranges from Costa Rica to central Panama from sea level to 1630 m ele-

vation in *Premontane wet forest*, *Tropical wet forest*, and *Premontane rain forest* life zones, as well as *Premontane wet forest* transition to *Tropical moist forest*.

*Philodendron grayumii* is a member of *P.* sect. *Calostigma* subsect. *Macrobelum* ser. *Macrobelum*. This species is characterized by its yellow-brown-drying stems; cataphylls deciduous intact; more or less ovate blades drying yellow-brown with the basal veins scarcely or not at all united into posterior ribs and rarely naked on the sinus; and primary lateral leaf veins splayed out, somewhat ridged, and downturned near the midrib. The spathe is often tinged reddish to purplish outside and dark red to maroon in the tube within.

This species is polymorphic, especially in the nature of the resin canals in the leaves and the number of inflorescences per axil. Collections from near sea level in Limón Province of Costa Rica (*Grayum* 8467, 8469) are unusual in lacking any clearly visible secretory ducts (normally easily visible at least on the dried lower leaf surfaces).

*Philodendron grayumii* may be confused with *P. edenudatum*, and the two may prove to be conspecific. *Philodendron edenudatum* differs in having proportionately longer leaf blades (1.8–2.1 times longer than broad) and proportionately shorter petioles (0.77–0.89 times as long as the blades).

*Philodendron grayumii* may also be confused with *P. ferrugineum*. The latter, occurring principally in the Canal Area, shares with *P. grayumii* brownish-drying blades with the basal veins usually not at all or only weakly naked. *Philodendron ferrugineum* differs in having generally shorter internodes; larger cataphylls (26–48 vs. 11–26 cm long); typically much larger, more coriaceous blades drying reddish brown (rather than yellowish brown); and minor veins usually drying markedly longitudinally raised but interrupted, giving the leaf surface a markedly bumpy, irregular, and uneven appearance.

Also perhaps confused with *P. grayumii* is *P. dodsonii*, which has dried leaf blades of a similar yellowish brown color. The latter species differs in having conspicuously naked posterior ribs, and blades drying matte on the upper surface and lacking any obvious secretory ducts on the lower surface.

Perhaps also confused with *P. grayumii* is *P. aromaticum*, which has similarly cordate blades that may dry a similar color and also has deciduous cataphylls. The latter species is distinguished by being a shorter-stemmed canopy epiphyte, and by having spongy petioles and proportionately more elongated, usually more gray-drying leaf blades

(usually more than 1.7 times longer than wide) with the naked portion of the posterior ribs much longer (usually 2.5–5 cm long).

A noteworthy collection is *Croat* 67525, which is somewhat intermediate between *P. grayumii* and *P. edenudatum*. It has narrowly ovate leaf blades about 1.6 times longer than wide, mottled petioles and lower midribs, and relatively small (possibly immature) inflorescences (to 13 cm long). In these characters, the specimen is closest to *P. edenudatum*, but it has prominent secretory ducts visible on the lower blade surface, and two inflorescences per axil, features not known in *P. edenudatum*. In addition, the blades dry dark brown. In this latter regard, this collection differs from both *P. grayumii* and *P. edenudatum*. It perhaps represents a new species, although it is being included with *P. grayumii* for now.

Another noteworthy collection is *Croat & Zhu* 76613, the only collection from east of the Canal Area. It is unusual in having the cataphylls semi-persistent as fibers, rather than deciduous while still intact. This may prove to represent a different taxon.

A sterile collection (*Croat* 66712) is noteworthy in having stems that dry smoother, darker reddish brown, and semiglossy; much glossier petioles than those of *P. grayumii*; and blades that dry more coriaceous, glossier, and with the minor veins prominent on both surfaces. It might represent another new species, but is tentatively included with this species.

The new species is named in honor of Michael H. Grayum, a colleague and aroid specialist, whose excellent collections of Araceae from Costa Rica and Panama have played an important role in this revision. He has collected the only Costa Rican material of the species.

*Additional specimens examined.* COSTA RICA. **Limón:** along road between Puerto Viejo de Talamanca and Manzanillo, vic. of Punta Cocles, ca. 5 m, 9°38'N, 82°43'W, 5 m, *Grayum & Sleeper* 4302 (CR, MO); 10–20 m, 10°38–39'N, 82°40–44'W, *Grayum* 3637 (MO); Piuta, ca. 1.5 km NW of Puerto Limón along highway to Portete, 5 m, 10°01'N, 83°03'W, *Grayum & Hammel* 8467 (MO), 8469 (MO). PANAMA. **Boocas del Toro:** Fortuna–Chiriquí Grande, 1.6 mi. N of Continental Divide, 770–790 m, 8°45'N, 82°17'W, *Croat* 76456 (AAU, CM, MEXU, MO, NY, OOM); 1.2 mi. N of Divide, 5.3 mi. N of Fortuna Dam, 910 m, 8°44'N, 82°17'W, *Croat* 60476 (AAU, CR, MO, PMA), 1170 m, *Croat* 666534 (MO); 5.8 mi. N of Fortuna Lake, 600 m, 8°45'N, 82°18'W, *Croat* 66794 (B, CAS, CM, COL, CR, F, K, L, MEXU, MO, NY, PMA, US, WIS); along Continental Divide (4.5 mi. N of middle of bridge over Fortuna Lake), 1170 m, 8°44'N, 81°17'W, *Croat* 66712 (MO); Gualaca–Chiriquí Grande, 1.6 mi. N of Continental Divide, 850 m, 8°48'N, 82°13'W, *Croat* 74933 (MO); 4.2 mi. E of Chiriquí Grande, ca. 0 m,

8°55'N, 82°09'W, *Croat* 66814 (MO, TEX); Station Mills 7.5 on Changuinola–Almirante Railroad, <100 m, *Croat* 38105 (B, F, K, MO, NY, PMA, US); Cerro Colorado, 6.5 mi. W of Chame, 1630 m, 8°35'N, 81°50'W, *Croat* 69164 (MO); Chiriquí Grande, Palo Seco–El Guayabo, km 112, *Gordon* 344 (PMA). **Chiriquí:** Gualaca–Chiriquí Grande, 4 mi. N of bridge over Fortuna Lake, 1200 m, 8°46'N, 82°16'W, *Croat* 68026 (CAS, CR, DUKE, F, G, MO, QCA, TEX, W); 1170 m, 8°44'N, 81°17'W, *Croat* 66712 (CAS, K, L, MEXU, MO, PMA); Río Hornito, ca. 0.5 km S of Centro de Científicos, 8°45'N, 82°18'W, *Croat* 76420 (CM, MO); 8.4 mi. beyond Los Planes de Hornito, 1130 m, 8°44'N, 82°14'35'W, *Croat* 67869 (DUKE, MO, NY, PMA). **Coclé:** 4.5 mi. N of El Copé, 580–740 m, *Croat* 67525 (C, ENCB, MO, P, VDB); 4.1 mi. N of El Copé, 770–680 m, 8°39'N, 80°36'W, 74840 (CM, CR, MO, PMA, US); 5–6 mi. N of El Copé, 600–800 m, 8°38'N, 80°35'W, *Croat & Zhu* 77202 (IBE, M, MO, MY, W); Llano Grande–Cocleito, 4.3 mi. N of stream in Llano Grande, 330 m, *Croat* 49235 (CM, MO). **Veraguas:** vic. of Santa Fe, 5 mi. past Escuela Agrícola Alto de Piedra, 670 m, 8°33'N, 81°08'W, *Croat* 66922 (BR, COL, M, MO), 66969 (AAU, MO, NY, OOM, PMA).

***Philodendron hammelii* Croat, sp. nov. TYPE:**

Panama. Coclé: Alto Calvario, ca. 6 km N of El Copé, Pacific drainage, on ridge W of sawmill, 850–1100 m, 8°39'N, 80°36'W, *Hammel* 3551 (holotype, MO-2658909). Figure 217.

Planta terrestis; internodia 1–1.5 cm longa, 8–12 mm diam.; cataphylla 11.5 cm longa, 1-costata, persistentia ut reticulum tenue fibrarum; petioli subteretes, 31–32 cm longus, 2–3 mm diam., dense obiecta squamis latis, pallidis prope apicem; lamina ovato-cordata, 21–22 cm longa, 13–16 cm lata, in sicco canoviridis supra, flavibrunnea infra; inflorescentia 1; pedunculus 4.5 cm longus, 4 mm diam.; spatula 11 cm longa; pistilla 6-locularia; loculi 14-ovulati.

Terrestrial; internodes 1–1.5 cm long, 8–12 mm diam., drying yellow-brown, semiglossy, conspicuously ridged; roots drying reddish brown to brown, slender, numerous per node; cataphylls to 11.5 cm long, 1-ribbed, drying yellowish brown, thin, smooth (not scaly), persisting as a thin, pale reticulum of branched tan fibers; petioles 31–32 cm long, 2–3 mm diam., subterete, drying weakly glossy, smooth or at most weakly granular, surface densely covered with broad pale scales near apex, the scales closed and overlapping, mostly less than three times longer than broad; blades ovate-cordate, subcoriaceous, shortly acuminate at apex, cordate at base, 21–22 cm long, 13–16 cm wide (1.4–1.5 times longer than wide), (0.66–0.67 as long as petiole), upper surface dark green, subvelvety and glistening, drying grayish green, lower surface much paler, semiglossy, drying yellowish brown and moderately glossy; anterior lobe 15–16.7 cm long, 15.3 cm wide (2.2–2.6 times longer than

posterior lobes); posterior lobes 6.5–6.7 cm long, 7 cm wide, rounded; sinus obovate to narrow hippocrepiform and closed; midrib drying sunken and concolorous above, convex, matte below; basal veins 6 per side, with 2 free to base, third and higher order veins coalesced 1 cm, sunken above, convex below; posterior rib not naked; primary lateral veins 3 per side, departing midrib at a 40–45° angle, quilted-sunken above, round-raised or raised, darker below; minor veins distinct, arising from both the midrib and primary lateral veins; “cross-veins” prominently raised upon drying. INFLORESCENCES probably 1 per axil; peduncle 4.5 cm long, 4 mm diam., ± terete, glabrous; spathe 11 cm long (2.4 times longer than peduncle), green, drying dark reddish brown; spadix sessile, 9.5–10.5 cm long; pistillate portion 3 cm long, 1 cm diam.; staminate portion 6.5–7.5 cm long, 1–1.5 cm diam.; pistils 1.5 mm long, 1.9 mm diam.; ovary 6-locular, 1.2 mm diam., locules 1 mm long, 0.3 mm diam., ovule sac 0.1–0.2 mm long, with axile placentation; ovules ca. 14 per locule, 2-seriate, contained within small, cloudy, gelatinous ovule sac, 0.1–0.2 mm long, longer than funicle; funicle to 0.1 mm long, adnate to lower part of partition, style similar to style type B; style apex flat; stigma hemispheroid, lobed, 1 mm diam., 0.3–0.4 mm high, covering entire style apex, depressed shallowly at center; the androecium truncate, ± prismatic, margins irregularly 4–5-sided, ca. 1 mm long; sterile staminate flowers irregularly 4–5-sided, 1.6–2.1 mm long.

Flowering in *Philodendron hammelii* apparently occurs in the early wet season, based on a single collection made in June.

*Philodendron hammelii* is endemic to Panama, known only from the type in Coclé Province in *Pre-montane rain forest* at 850 to 1100 m elevation.

*Philodendron hammelii* is a member of *P.* sect. *Philodendron* subsect. *Achyropodium*. This species is characterized by its short, slender internodes; persistent, thin, pale network of branched cataphyll fibers; petioles longer than the blades and densely covered with broad, pale scales near the apex; and ovate-cordate, yellow-green-drying blades.

*Philodendron hammelii* is superficially most similar to *P. colombianum* R. E. Schult. from the Amazon basin. These species have similarly shaped blades drying yellow-green with prominent “cross-veins” and persistent, pale cataphyll fibers. *Philodendron colombianum* differs, however, in having the blade sinus acute at apex and petioles lacking scales.

In Panama, this species might be confused with

*P. squamipetiolatum*, which also has scaly petioles. It differs in having blades with a more arcuate sinus and petioles conspicuously long-scaly throughout most of their length.

The species is named in honor of Barry Hammel of the Missouri Botanical Garden who has collected many important Araceae in Panama and Costa Rica, including the type specimen.

***Philodendron hebetatum* Croat, sp. nov. TYPE:**

Panama. Coclé: vicinity el Valle de Antón, at forested flat area near Finca Macarenita at La Mesa, 800 m, 8°36'N, 80°07'W, 6 July 1994, Croat & Zhu 76693 (holotype, MO-4619514-15; isotypes, B, CAS, COL, CR, F, K, NY, PMA, US). Figures 33, 218–220, 225, 226.

Planta hemiepiphytica; internodia 1–4 cm longa, 1.5–2.5 cm diam.; cataphylla ad 12–35 cm longa, incostata, persistentia semi-intacta; petiolus subteres, 27–75 cm longus, 0.8–1.8 cm diam., obtuse complanatus adaxialiter, in sicco conspicue flavibrunneus; lamina anguste ovato-triangularis, cordato-sagittata basi, 24–63 cm longa, 19–44 cm lata, atriviridis supra, opace albida, hebetata infra; inflorescentia 1–5; pedunculus 4–11 cm longus, 4–11 mm diam.; spathe 11–17.5 cm longa; lamina spathae extus viridi vel albida; tubo spathae extus atriviridis aut interdum purpureo vel violaceipurpureo vel rubeo aut subroseo, intus albidus vel rubellus vel pallide violaceipurpureo; pistilla (4)5–8-locularia; loculi (4)20–24-ovulati.

Hemiepiphytic; stem appressed-climbing; internodes short, semiglossy, 1–4 cm long, 1.5–2.5 cm diam., longer than broad, green to gray-green, drying yellow-brown; roots moderately numerous per node, drying dark brown, ridged; cataphylls 12–35 cm long, unribbed, sometimes sharply 1-ribbed near apex (sharply 2-ribbed in South America), medium green, drying light to medium brown, persisting semi-intact at upper nodes, then deciduous with large patches of yellow-brown periderm remaining intact; petioles 27–75 cm long, 0.8–1.8 cm diam., ± terete, obtusely flattened adaxially, spongy (in South America), dark green to gray-green, surface semiglossy to matte, transversely fissured near apex, drying pale yellow-brown. LEAVES erect-spreading; blades ovate-triangular, subcoriaceous, abruptly acuminate at apex, cordate-sagittate at base, 24–63 cm long, 19–44 cm wide (1.1–2.3 times longer than wide), (0.7–1.2 times longer than petiole), equal to or longer than petiole, margins concave, broadly undulate, upper surface semiglossy, dark green, drying conspicuously yellow-brown, semiglossy, lower surface opaque whitish, matte, paler; anterior lobe 19.5–49.5 cm long, 14.5–42 cm wide (1.9–3.6 times longer than posterior lobes); posterior lobes 7.1–20.2

cm long, 7.3–20.5 cm wide, obtuse to rounded; midrib broadly concave, paler than surface above, thicker than broad, weakly glossy, darker than surface below; basal veins 6 per side, with 1–2 free to base, second and third veins coalesced ca. 3 cm; posterior rib naked for 2.5–3.5 cm long; primary lateral veins 6–11 per side, departing midrib at a 35–50° angle, weakly sunken above, convex and darker than surface below; minor veins moderately distinct, fine, intermittent below, arising from both the midrib and primary lateral veins. INFLORESCENCES 1–5 per axil; peduncle 4–11 cm long, 4–11 mm diam., white-lineate to coarsely white-streaked toward apex and on spathe base; **spathe** 11–17.5 cm long, 2–3.5 cm diam. (1.3–3.1 times longer than peduncle), white-speckled, semiglossy, constricted above the tube, 9–11 mm diam. at constriction; spathe blade green to whitish, sometimes rose-red outside, green to whitish or sometimes rose-red and white-speckled inside; spathe tube oblong-ellipsoid, 4–7 cm long, usually dark green or sometimes purple to violet-purple to red or pinkish outside, reddish to pale violet-purple (dark maroon in South America) inside; **spadix** white, cylindrical, 12.9–16.6 cm long, broadest near the base, constricted near the middle; pistillate portion whitish, cylindrical to ovoid, 3.8–8.9 cm long, 7–16 mm diam.; staminate portion 9.3–13.4 cm long; fertile staminate portion cylindrical, 8–13 mm diam., broadest at the base, narrower than pistillate and sterile portions; sterile staminate portion as broad as or narrower than the pistillate portion, 1–1.3 cm diam.; pistils glossy, 2.1–4(5.8) mm long, 1.2–2.6 mm diam.; ovary (4)5–8-locular, walls embedded with granular, crystal-like particles, locules 1.3–3.2(4.5) mm long, 0.4–0.6 mm diam., with axile placentation; ovules (4)20–24 per locule, 2-seriate, 0.2 mm long, about equal in length to funicle; funicle 0.1 mm long, adnate to lower part of axillar wall, style 0.4–0.7 mm long, 1.2–2.6 mm diam., similar to style type B; style apex flat or rounded; stigma subdiscoid, 1–1.5 mm diam., 0.2–0.5 mm high, covering center of style apex; the androecium truncate, margins 4–6-sided, sometimes scalloped; thecae oblong, 0.3–0.4 mm wide, contiguous,  $\pm$  parallel to one another; sterile staminate flowers 3–6-sided, 1.5–2.5 mm long, 1.4–1.8 mm wide. Berries white, with purple stigmas; seeds 2–3, 6, 10–12, 18–25 per locule, (1–5)8–11 mm long, 2–3 mm diam.

Flowering in *Philodendron hebetatum* apparently occurs throughout the dry season and first part of the rainy season in Panama, from December through July. Post-anthesis inflorescences have been collected from January through October (ex-

cept September). Immature fruits have been collected in June and July. South American collections follow more or less the same pattern as those from Panama.

*Philodendron hebetatum* ranges from Panama to Colombia (Chocó) and Ecuador (Carchi, Esmeraldas, and Pichincha) along the Pacific coast, at 10 to 1880 m elevation in *Premontane wet forest*, *Tropical wet forest*, and *Premontane rain forest* life zones. It is to be expected in adjacent Costa Rica. In Panama, most collections have been made at or less than 1400 m, rarely to about 1630 m.

*Philodendron hebetatum* is a member of *P. sect. Philodendron* subsect. *Philodendron* ser. *Impolita*. This species is characterized by its generally appressed-climbing hemiepiphytic habit; short internodes; usually unribbed, semi-intact persistent cataphylls with large patches of yellow-brown epidermis remaining intact; petioles equaling or exceeding the blade in length and drying conspicuously yellow-brown and smooth (thus clearly demarcated from the blade); and markedly bicolorous, narrowly ovate-triangular blades usually concave along the margin and matte and whitish on the lower surface.

The petioles of *P. hebetatum* are variable, being typically terete to obtusely D-shaped in Central America but sharply C-shaped to sharply D-shaped with acute to bluntly raised lateral margins in South America. South American populations also have the major leaf veins drying yellowish to pale red vs. dark red in Central America. It is possible that the South American elements may ultimately prove to be at least subspecifically distinct.

Some South American specimens of *P. hebetatum* are particularly noteworthy. *Holm-Nielsen et al.* 25665, from Ecuador, differs from other collections in having the petiole epidermis drying paler yellow and the immature spathes drying a paler reddish brown. A collection from Nariño Department, Colombia (*Croat 72425A*), is unusual in having the margins of the anterior lobe markedly concave, but perhaps belongs also to this species.

*Philodendron hebetatum* is easily confused, especially in live condition, with *P. thalassicum*, which may also have triangular-ovate blades much paler and matte on the lower surface. The latter differs, however, in having petioles that are sharply D-shaped and dry blackened rather than conspicuously yellow-brown. In addition, *P. hebetatum* has spathes reddish within, while in *P. thalassicum* they are greenish within.

*Philodendron hebetatum* is also similar to *P. strictum*, which differs by its usually terrestrial hab-

it, more or less reclining stems, and broadly ovate blades. Both species have pale, matte lower blade surfaces, yellow-brown drying petioles, and cataphylls with persistent patches of intact, yellowish brown epidermis. The lower blade surface in both species is covered with a wax-like substance, and dries with a fine reticulate pattern (areolate at 7× or higher magnification). *Philodendron strictum* ranges from Costa Rica to Panama and Venezuela at 650 to 1900 m elevation. In Panama, where both species occur together along the Fortuna Dam Road, *P. strictum* tends to occur at higher elevations than *P. hebetatum*.

Juvenile plants (Croat 56049, 56174, 56213, 69291; Valle and Chocó Departments, Colombia) of *P. hebetatum* have oblong-elliptic leaf blades acute at the base, but the petiolar epidermis already is colored as in adult plants. Leaves of older plants become increasingly rounder and broader at the base and finally subcordate to cordate.

An unusual collection is Croat 49298 from Chocó Department, Colombia, with leaves only shallowly cordate, even though they are in the upper end of the size range for adult plants of this species.

Sánchez et al. 553, a collection from Colombia (Antioquia), is noteworthy in being from 1880 m elevation and in being terrestrial. Holm-Nielsen et al. 25665 from Ecuador (Esmeraldas) differs from other South American collections in having a paler, flakier epidermis on the petioles and smaller, more or less sessile (perhaps immature) inflorescences. Perhaps it represents another species. South American collections of *P. hebetatum* differ from those of Panamanian material in that the petioles dry dark brown rather than yellowish.

Croat 61396 is unusual in apparently having only four ovules per locule.

**Additional specimens examined.** PANAMA. **Bocas del Toro:** Fortuna Dam area, Chiriquí Grande-Fortuna, 2.8 mi. N of Divide, 850-950 m, ca. 8°45'N, 82°15'W, McPherson 9663 (MO, NY); 3.2 mi. N of Divide, 700 m, 8°45'N, 82°15'W, Croat & Grayum 60263 (CM, MO); 13.2 mi. W of Chiriquí Grande, 310 m, 8°45'N, 82°10'W, 60140 (AAU, MO). **Chiriquí:** Cerro Colorado, 18-27 mi. off Pan-American Highway along mining road, 1200-1500 m, Croat 33141 (MO); 1390-1410 m, Croat 37223 (MO); 9.2 mi. W of Chame, 1450-1480 m, 8°35'N, 81°50'W, Croat 69064 (CM, MO, PMA); 8.3 mi. beyond Chame, 1630 m, Croat 75042 (MO); 75046 (CM, MO); Fortuna Dam area, 4.5-5 km N of dam over Fortuna Lake, 1100-1135 m, 8°43'N, 82°07'W, Croat & Grayum 60073 (B, CM, MO); 11.8 mi. N of Los Planes de Hornito, Croat CM 48668 (MO). **Coelís:** El Copé region, 9.4 km above El Copé, 750-900 m, Croat 44686 (CAS, MO); Alto Calvario, 800 m, 8°39'N, 80°36'W, Croat 75075 (CM, MO); 710-800 m, 8°39'N, 80°36'W, Croat 68712 (ENCB, MO); El Valle region, vic. of La Mesa, N of El Valle de Antón, 900-1000 m, 8°37'N, 80°08'W, Croat 67279 (MO);

8°37'N, 80°07'W, McPherson 11203 (CM, M, MO); 8°40'N, 80°07'W, Knapp 5337 (MO); 5753 (MEXU, MO); La Mesa, 775 m, 8°36'N, 80°07'W, Croat 74783 (MO, QCA); 14372 (MO); 2000-2700 ft., Dreyer et al. 4567 (MO, NY); 860-900 m, Croat 37364 (MO); ca. 800 m, 25407 (MO); 800-900 m, 8°38'N, 80°09'W, 67204 (DUKE, MO, US). **Darién:** Parque Nacional Darién, near gold mine at headwaters of N branch of Río Pucuro, slopes of Cerro Tacarcuna, ca. 6 km N of Cerro Mali, 1300-1500 m, 8°09'5"N, 77°15'W, Hammel et al. 16541 (COL, MO); Río Tuquesa, Tyler Kittredge gold mine, ca. 2 air km from Continental Divide, Croat 27240 (MO). **Panamá:** Cerro Jefe region, vic. of summit, 850 m, 9°14'N, 79°22'W, Croat 67060 (MO, NY); 0.8 mi. beyond turnoff to Altos de Pacora, 770 m, 9°15'N, 79°29'W, Croat & Zhu 76611 (CM, K, M, MEXU, MO, W); 4.6 km beyond peak on road to Alto de Pacora, 26.3 km from InterAmerican Highway, 600 m, Croat 35934 (MO); 3-3.5 mi. NE of Altos de Pacora, 7.8-8.2 mi. above highway, 700-750 m, 9°15'N, 79°25'W, Croat 68661 (IBE, MO); 5-10 km NE of Altos de Pacora, 700-800 m, Mori & Kallunki 6031 (MO); El Llano-Carri Road, Mile 10, near El Llano, 330 m, Croat 33773 (MO, US); Cerro Campana, ca. 850 m, 8°40'N, 79°50'W, Thompson 4598 (MO), Croat 17243 (MO); 780-875 m, 25224 (F, MO, PMA); 1000 m, Lutesyn 3193 (DUKE); ca. 1 mi. from Inter-American Highway, ca. 150 m, Croat 35961 (F, MO); above Su Lin Motel, Croat 14726 (MO, PMA); lower slopes above FSU cabin, 850 m, Croat 227954 (MO). **San Blas:** El Llano-Carri Road, Mile 14, 300 m, 9°15'N, 79°W, Croat 69231 (MO, PMA, SAR); sendero de Interpretación, 1 km al este del Campamento de los guardabosques de INRENA, 800-900 m, 8°40'N, 79°55'W, Correa et al. 9489 (MO). **Veraguas:** Santa Fe region, Santa Fe-Río Calovebora, 0.6 mi. beyond Escuela Agrícola Alto Piedra, 735 m, Croat & Folsom 33996 (MO); Cerro Tute, 800-1100 m, 8°35'N, 81°5'W, Hamilton & Krager 3934 (MO); Río Tercero Brusó Valley, beyond Escuela Agrícola Alto Piedra, above Santa Fe, Croat 27326 (MO).

**COLOMBIA. Antioquia:** Mpio. Frontino, Corregimiento Nutibara, valley of the upper Río Cuevas, 1880 m, Sánchez et al. 553 (MEDEL); Río Anorí, confluence of Quebrada La Tirana and Río Anorí, ca. 3 km upriver from Planta Providencia, 28 km SW of Zaragoza, 400-700 m, Abernethy et al. 299 (COL, WIS); Río Anorí valley near Planta Providencia, 350-600 m, 7°30'N, 74°50'W, Shepherd 924 (COL, WIS). **Chocó:** Quibdó-Tutunendo Road, ca. 3 km W of Tutunendo, 80 m, Gentry et al. 30108 (MO); 1 km E of Tutunendo, 100 m, 5°46'N, 76°35'W, Gentry et al. 30079 (COL); Medellín-Quibdó, Km 136.4, 63 km E of Tutunendo, 960 m, 5°47'N, 76°22'W, Croat 56348 (COL, JAUM, MO); Km 208.5, 9 km W of Tutunendo, <100 m, 5°39'N, 76°40'W, Croat 56213 (MO); Río Atrato, 39 km W of Bolívar, 1600 m, Croat 49261 (MO); 78 km W of Bolívar, 466 m, Croat 49298 (MO); 27 mi. W of Bolívar, 1190 m, 5°50'N, 76°16'W, Croat & Cogollo 52089 (MO); Quibdó-Bolívar, Km 137-138, 79-80 km E of Quibdó, 910-920 m, 5°45'N, 76°21'30"W, Croat 57345 (COL, JAUM, MO); Quibdó-Las Animas, ca. 1 km N of Las Animas, ca. 100 m, 5°14'N, 76°40'W, Croat 55958 (COL, MO); San José del Palmir-Cartago, Vereda La Bella between San José del Palmir and turnoff to El Cairo, 1430 m, 4°53'N, 76°13'W, Croat 56715 (COL, MO, QCA); Pueblo Rico (Risaralda)-Istmina (Chocó), Quebrada Antón, 15 km W of Santa Cecilia, 6 km W of Chocó-Risaralda border, 240-350 m, 5°20'30"N, 76°13'45"W, Croat 70961 (MO); Quibdó-Istmina, Km 4, <100 m, 6°28'N,

76°36'W, *Croat & Cogollo* 52234 (MO); Serranía de Baudo, Las Animas-Pato on Río Pato, ca. 150 m, 5°30'N, 76°46'W, *Croat* 56131 (COL, JAUM, MO, PMA); 16 km NW of junction with Quibdó-Istmina road near Las Animas, 100 m, 5°20'N, 76°42'W, *Croat* 56174 (COL, JAUM, MO); 5°16'N, 76°41'W, *Croat* 56049 (COL, JAUM, MO); Río San Juan, Docordó, 0 m, 4°15'N, 77°22'W, *Forero et al.* 4352 (MO). **Nariño:** along road between Junín and Barbacons, 1.9 km NE of Junín, 1130 m, 1°21'N, 87°6'W, *Croat* 724254 (CM, MO). **Risaralda:** Mpio. Mistrato, Chirrincha, Río Aguita, 950 m, *Franco et al.* 3522 (MO); corregimientos de Geguadas—Puerto de Oro, 1550 m, *Alonso et al.* 9819 (MO); 9742 (MO). **Valle:** Bajo Calima area, *Croat* 62761 (CM, MO); ca. 15 km N of Buenaventura, Cartón de Colombia concession, Juanchaco region, 500 m, 3°56'N, 77°08'W, *Gentry et al.* 53711 (MO); Lijal-Gasolina road junction, SSW of San Isidro, 50 m, *Gentry et al.* 62926 (MO); 11 km N of main Cali—Buenaventura Highway, ca. 50 m, 3°56'30"N, 77°01'W, *Croat & Monsalve* 61396 (MO, QCA); Buenaventura—Río Calima, 33.3 km N of Cali—Buenaventura Highway, ca. 50 m, 4°02'N, 77°07'W, *Croat* 61279 (COL, MO, NY, TEX); Km 11, 50–80 m, 3°56'30"N, 77°01'W, *Croat* 69291 (CAS, F, L, MO, US); ca. 4 km from Río Calima, near Km 14 marker, <50 m, 3°56'N, 76°59'W, *Croat* 57530 (AAU, DUKE, K, MEXU, MO); Buenaventura—Málaga Road, 185 m, 3°56'N, 77°01'W, *Croat* 70103 (B, CAS, COL, F, K, MO, PMA, US); Km 17.5, 130 m, 3°57'N, 77°01'W, *Croat & Bay* 75629 (MO); Km 18–20 on Cali—Buenaventura Highway, Finca Zinzara, 1500–2000 m, *Cabrera & van der Werff* 15779 (MO); Río Cajambre, 5–80 m, *Cuatrecasas* 17054 (US); Río Naya, upriver from Puerto Merizalde, ca. 10 m, 3°15'N, 77°25'W, *Gentry & Juncosa* 40673 (MO). **ECUADOR. Carchi:** Tulcán Cantón, Reserva Indígena Awá, sector Sabalera, parroquia Tobar Donoso, 650–100 m, 1°N, 78°24'W, *Tipaz et al.* 1298 (MO); 1462 (MO). **Esmeraldas:** Lita—San Lorenzo, 15.5 km W of Lita, 705 m, 0°55'N, 78°28'W, *Croat* 72372 (MO); San José de Cayapas, 80 m, 0°52'N, 78°57'W, *Holm-Nielsen et al.* 25665 (AAU, MO); Eloy Alfaro, Reserva Ecológica Cotacachi-Cayapas, Charco Vicente, Río San Miguel, afluente del Río Cayapas, 150 m, 0°43'N, 78°53'W, *Palacios & Trado* 11276 (MO, QCNE). **Pichincha:** Reserva ENDESA, Quito—Puerto Quito, Km 11.3, 0°05'N, 79°02'W, 750 m, *Croat & Rodríguez* 61454 (CM, K, M, MO, QCA); 61455 (K, MO, QCA); 710 m, 0°03'N, 79°07'W, *Croat* 73155 (AAU, B, COL, F, K, MO, NY, QCA, US).

**Philodendron hederaceum** (Jacq.) Schott, *Wiener Z. Kunst.* 1829: 780. 1829. *Arum hederaceum* Jacq., *Enum. Syst. Pl.* 31. 1760. TYPE: t. 51, fig. D in *Plum.*, *Pl. Amer.*, 1756 (holotype). Figures 221–223.

Hemiepiphytic vine; growing to often high in trees, stem appressed-climbing, eventually scandent, often pendent, sap clear, turning honey-colored, leaf scars 7–15 mm long; internodes weakly flattened on one side, sometimes with 2 sharply raised on the side above the petioles, usually weakly glossy, sometimes matte, pale to medium green, minutely speckled to striate, usually smooth when fresh, but drying minutely ridged, sometimes prominently ribbed throughout its circumference (the

ribs smooth to prominently warty), (2)10–28 cm long, 1–2.5(3.5) cm diam., dark green, usually drying green, sometimes reddish; roots brown, to 10 cm long, many at nodes; cataphylls 6–10 cm long, unribbed, weakly 1-ribbed, or bluntly to sharply 2-ribbed, pale green, deciduous; **petioles** (6)9.7–27(33) cm long, 6–10 mm diam., terete to subterete, pale green, firm, flattened adaxially, pale green, surface smooth, weakly glossy to matte; **blades** broadly ovate, subcoriaceous to coriaceous, semi-glossy to matte, acuminate to long acuminate, sometimes cuspidate at apex (the acumen inrolled, 0.5–0.9 mm long), 11–40(50) cm long, 8–24(34) cm wide (1.2–1.9 times longer than wide), (ca. 1.3(1.5) times longer than petiole), margins hyaline to pale yellowish, upper surface dark green, sometimes subvelvety, lower surface slightly paler, often purplish violet, drying gray-green to yellow-green; anterior lobe 9–30(41) cm long, 9–24(29) cm wide ((2)2.4–3.7(6.3) times longer than posterior lobes); posterior lobes 3–10(14) cm long, 4.3–15.7 cm wide, directed inward and sometimes overlapped, obtuse to rounded; sinus usually deeper than broad, mostly spatulate, rarely hippocrepiform, 3–7 cm deep; midrib convex to flat or sunken, concolorous or slightly paler than surface above, convex, concolorous below; basal veins (3)4–5(6) per side, with (0)1–2 free to base, part of the remainder coalesced to 1.5–2 cm; posterior rib not naked; primary lateral veins 2–6 per side, departing midrib at a 35–55° angle, ± straight to the margins, sunken to weakly raised, slightly paler than surface above, convex and paler than surface below; minor veins obscured to moderately distinct, arising from both the midrib and primary lateral veins. **INFLORESCENCES** erect or pendent, 1 per axil; peduncle (2–)3)4–15.7 cm long, 8–12 mm diam., pale green, sometimes tinged purple, matte; **spathe** subcoriaceous to coriaceous, 9–16.6(24) cm long, (0.9–)2.6(3.3) times longer than peduncle, weakly constricted above the tube, 1.3–3.6(5.7) cm diam. at constriction, usually green, sometimes yellowish white, yellowish green, or cream to creamy-white throughout; spathe blade sometimes purple tinged outside, 1.2–3.6 cm diam. when furled, pale green, greenish yellow, sometimes tinged red inside; spathe tube dark green, sometimes tinged reddish maroon outside, 5–6 cm long, 1.5–4.9(6.9) cm diam., maroon, dark red, crimson, or purple at base inside; **spadix** stipitate to 5–10 mm long, dark maroon; 12–20 cm long; pistillate portion pale greenish white to green, 3.5–6 cm long, 1.5 cm diam. at base, 1.7 cm diam. at middle and near apex; staminate portion 7–11 cm long; fertile staminate portion creamy white to pinkish, broadest at base,

weakly constricted ca. 1 cm above sterile portion then  $\pm$  uniform to near apex, 1.4–2.2 cm diam. at middle, 9 mm diam. ca. 1 cm from apex; sterile staminate portion 1.6 cm diam.; pistils 4–9.2 mm long, 1.8–3.1 mm diam.; ovary 4–6(7)-locular, 8 mm long, locules 8 mm long, 1.4 mm diam., with axile placentation; ovules 20–25 per locule, 0.1 mm long, 2-seriate; funicle 0.1–0.3 mm long, adnate to lower part of partition, style 1.1 mm long, 2.9 mm diam., similar to style type B; style apex flat or somewhat rounded, drying concave with a pale margin and 4–6 paler, flat to weakly sunken circular areas associated with the stylar pores; stigma 3–4-sided, light brown to reddish, drying light brown, 1.8–3.3 mm diam., 0.35 mm high, margins thin; sterile staminate flowers blunt, irregularly 4–5-sided, 2.2 mm long, 1.3 mm wide. INFRUDESCENCE pendent, often on leafless stems; spathe dark green, weakly glossy outside; pistillate spadix 5–8 cm long, 3.5–4 cm diam.; berries greenish white; seeds 1–2 per locule, somewhat orange, many per berry,  $\pm$  ovoid to oblong ellipsoid, (1.5)3–5 mm long, 2.5–4 mm diam., with weak constriction (nipple) and densely covered with raphide cells. JUVENILE plants with upper blade surface dark green, sometimes reddish green, with glistening minute close papillations, lower surface somewhat maroon; veins less conspicuous.

*Philodendron hederaceum* ranges throughout the West Indies and from Mexico throughout Central America and much of South America, at sea level to 1200(1500) m elevation. In South America it extends from as far south as Los Ríos Province in coastal Ecuador to Trinidad, Venezuela, the Guianas, Brazil, Ecuador, Peru, and Bolivia on the Atlantic drainage of the continent.

*Philodendron hederaceum* is a member of *P.* sect. *Philodendron* subsect. *Solenostegma*. This species is distinguished by its scandent habit, long internodes, deciduous cataphylls, ovate-cordate, long-petiole leaves, and solitary inflorescence with usually green spathes with the tube reddish to purplish within.

*Philodendron hederaceum* is most easily confused with *P. purpureosviride*, which is a vine with similar leaves. See that species for differences.

*Philodendron hederaceum* is also somewhat similar to *P. jacquinii*, but the latter differs in its generally pubescent stems, petioles and major veins of the lower blade surface, its thinner leaf blades, swollen spathe tube, and broad pistillate portion of the spadix with elongate styles.

The taxa here treated as *Philodendron hederaceum* and *P. jacquinii* have long been confused no-

menclaturally. The former name is based on *Arum hederaceum*, first validly published by N. J. Jacquin (1760) in his *Enumeratio Systematica Plantarum*. Jacquin cited only a Plumier (1756) plate (t. 51, fig. d), which thus must be accepted as the holotype of the name (Greuter et al., 1994; Art. 9.1, Note 1). Three years later, Jacquin (1763), in his *Selectarum Stirpium Americanarum Historia*, published his own plate (t. 152), identified as *A. hederaceum* but actually depicting a different species from Cartagena, Colombia, namely, that treated in this revision as *P. jacquinii* Schott. These two species are fortunately sufficiently different in appearance that even drawings of sterile plants, such as that of Plumier, are unmistakable. This mistake by Jacquin has caused considerable confusion, since several authors, including Kunth (1841), Engler (1899), Krause (1913), Dugand (1945), and Bunting (1963b, 1995), have misapplied the name *P. hederaceum* based on Jacquin's (1763) publication.

Schott, however, understood the problem. He had already transferred *Arum hederaceum* to *Philodendron* (Schott, 1829), and in his treatment for *Synopsis Aroidearum* (Schott, 1856) he described *P. jacquinii*, explicitly basing it on Jacquin's (1763) plate. Schott (1856) also placed *P. hederaceum* sensu Kunth into synonymy under his newly described *P. jacquinii*. However, he included *P. hederaceum* in his grex *Macrobolium*, while treating other synonyms of *P. hederaceum* in grex *Solenostegma*. Names so treated were: *P. scandens*, *P. prieurianum*, *P. oxycardium*, *P. cuspidatum*, and *P. micans*.

Engler (1899), followed by Krause (1913), treated the species herein called *P. hederaceum* as four distinct species: *P. prieurianum*, *P. scandens*, *P. oxycardium*, and *P. micans*. Both Engler and Krause erred in treating *Arum hederaceum* Jacq. as a questionable synonym of *P. hoffmannii* (= *P. jacquinii*), citing Jacquin's (1763) t. 152 as the type.

Despite the confusion by Engler and Krause, Standley and Steyermark (1958b) correctly dealt with the taxonomy of *P. hederaceum*, citing *P. scandens*, *P. oxycardium*, and *P. miduhoi* in synonymy. Their treatment of *P. jacquinii* was incorrect, since they cited that name under the later synonym *P. hoffmannii* Schott (1858). In this regard they followed Krause (1913). Thus, despite the confusion by Engler and by Krause, the nomenclature of these species was essentially rectified as early as 1958 to the species as *P. oxycardium* or *P. cordatum* hort. (non Vell.).

Bunting (1963b), apparently following the lead of Dugand (1945), was aware of the Plumier illustration cited by Jacquin but seemed to believe that it had no bearing on the application of the name.

He also referred to Jacquin (1763) as the "initial" publication of *Arum hederaceum*, presumably in the erroneous belief that the name had not been validly published in Jacquin (1760). Bunting (1963a) accordingly applied the name *P. hederaceum* to the species here called *P. jacquinii* and the obscure *P. scandens* K. Koch (1853) for the species herein called *P. hederaceum*. Certainly the epithet *P. scandens* was not in general use up until that time because most horticultural works (Birdsey, 1951) still referred to the species as *P. oxycardium* Schott.

Other, more practical matters substantiate that Plumier's (1756) rather crude drawing depicts the species here treated as *P. hederaceum*, rather than the one called *P. jacquinii* (accurately illustrated by Jacquin, 1763). Plumier's text associated with t. 51, fig. D states that the plant grew in Martinique, known to have been visited by Plumier (Urban, 1898: 123). *Philodendron hederaceum* is a widespread species in the West Indies and occurs on Martinique, while *P. jacquinii*, though also widespread, is not known from the Lesser Antilles. The epithet *hederaceum* connotes an ivy-like growth habit and aspect, apt for the species to which the epithet is here applied, but not for *P. jacquinii*. The application of the name *P. hederaceum*, as explained above, is unambiguous, whereas that of *P. scandens*, the name used during the last 30 years for this plant, is highly dubious, as it is based on a sterile Koch specimen of unknown origin, lacking an extant type specimen or even illustrations.

Article 57.1 of the Tokyo Code (Greuter et al., 1994) states that "A name that has been widely and persistently used for a taxon or taxa not including its type is not to be used in a sense that conflicts with current usage unless and until a proposal to deal with it under Art. 14.1 or 56.1 has been submitted and rejected." I consider that this Article does not apply in the present case, since the name *P. hederaceum* was used in the sense, including its type, as recently as 1958, in a higher regional flora (*Flora of Guatemala*), which is the most recently published treatment of Araceae for any Central American country and still reigns as the standard work throughout the region.

In Central America, *P. hederaceum* can be divided into three varieties. *Philodendron hederaceum* var. *hederaceum* and *P. hederaceum* var. *oxycardium* (Schott) Croat are only distinguishable on the basis of their juvenile leaf blades. In *P. hederaceum* var. *hederaceum*, the juvenile blades are velvety with a silky sheen on the upper surface, whereas the juvenile blades of *P. hederaceum* var. *oxycardium* are glossy on the upper surface. These juvenile forms

were formally treated by Bunting (1968) as *P. scandens* forma *micans*.

The third newly recognized variety, *P. hederaceum* var. *kirkbridei* Croat, is distinguishable by its adult stems that dry brown and are deeply sulcate with prominent ridges. This taxon also differs in occurring at higher elevations.

The following key separates the three varieties of *P. hederaceum* in Central America. For anatomical differences see Bunting (1968).

#### KEY TO THE VARIETIES OF *P. HEDERACEUM*

- 1a. Adult stems weakly sulcate on living plants, moderately to conspicuously sulcate and reddish brown upon drying, usually densely warty; spathe tube dark red to red-purple inside; Costa Rica, Panama, Ecuador, and Suriname, mostly 450–900 m elevation ..... var. *kirkbridei*
- 1b. Adult stems smooth, drying mostly green, weakly striate, never minutely warty; spathe tube green (sometimes weakly tinged reddish) inside.
  - 2a. Juvenile blades velvety with a lustrous sheen on the upper surface, often tinged purplish on the lower surface; apparently ranging throughout the range of the species in Mexico, Central America, and to the West Indies and South America (including Amazonian basin) ..... var. *hederaceum*
  - 2b. Juvenile blades glossy on the upper surface, green on the lower surface; known only from the Gulf slope of Mexico in Veracruz, northern Oaxaca, and Tabasco, but possibly also found in Jamaica ..... var. *oxycardium*

#### *Philodendron hederaceum* (Jacq.) Schott var. *hederaceum*

*Philodendron scandens* K. Koch & Sello, in A. Braun et al., *Append. sp. Hort. berol.* 1853: 14. 1853–1854. TYPE: A cultivated plant grown in Berlin, origin not stated (holotype, B lost).

*Philodendron harlowii* I. M. Johnston, *Sargentia* 8: 91, t. 14, fig. 1. 1949. TYPE: Panama, Panamá: San José Island, area 11B, Johnston 1030 (holotype, GH; isotype, MO).

*Philodendron midahoi* Matuda, *Revista Soc. Mex. Hist. Nat.* 11: 95, figs. 6–7. 1950. TYPE: Mexico, Chiapas: Finca Esperanza, 160 m, 23 Dec. 1949, Matuda 18721 (holotype, MEXU; isotype, UC).

*Philodendron cuspidatum* K. Koch & Bouché, in A. Braun et al., *Append. Gen. sp. Hort. berol.* 1854: 7. 1854–1850. *Philodendron scandens* var. *cuspidatum* (K. Koch & Bouché) Engl., *Bot. Jahrb. Syst.* 26: 528. 1899. TYPE: cultivated at Berlin Botanical Garden (holotype, B? destroyed). Schott ic. 2679 (neotype, here designated).

*Philodendron micans* K. Koch, in A. Braun et al., *Append. Gen. sp. Hort. berol.* 1854: 7. 1854–1855. *Philodendron scandens* f. *micans* (K. Koch) G. S. Bunting, *Gentes Herb.* 10: 63. 1968. TYPE: original locality unknown (holotype, B? destroyed). Schott ic. 2709 (neotype, here designated, W).



*Philodendron microphyllum* K. Koch, in A. Braun et al., *Append. Gen. sp. Hort. berol.* 1854: 7, 1854-1855. TYPE: cultivated collection at Berlin (holotype, B? now lost).

*Philodendron oxypurum* Schott, *Syn. Aroid.* 82. 1856. TYPE: Venezuela. [Synonymized by Engler (1899) but no specimen or illustration now exists.]

*Philodendron acrocardium* Schott, *Oesterr. Bot. Z.* 8: 179. 1858. TYPE: Guatemala, *Wendland* (holotype, W? destroyed). Schott *ic.* 2498 (neotype, here designated).

*Philodendron pittieri* Engl., *Bot. Jahrb. Syst.* 26: 541. 1899. TYPE: Costa Rica, Puntarenas: Ujarrás de Buenos Aires, 9°12'N, 83°17'W, Feb. 1897, *Pittier 11132* (holotype, B; isotype, BR).

Hemiepiphyte; internodes 10-25 cm long, 1-2.5(3.5) cm diam., weakly flattened on one side, medium green, minutely speckled to striate or smooth when fresh but drying minutely ridged, greenish; cataphylls 6-10 cm long, unribbed, weakly 1-ribbed or bluntly to sharply 2-ribbed, deciduous intact; **petioles** (6)10-27(33) cm long, 6-10 mm diam.; blades (11)16-40(50) cm long, 8-24(34) cm wide; upper surface medium to dark green, velvety when juvenile, drying brown to greenish brown, lower surface medium green, glossy, drying gray-green to yellow-green; primary lateral veins 2-6 per side, departing midrib at a 35-55° angle. INFLORESCENCES with peduncle (2)4-16 cm long; **spathe** 9-17(20) cm long, spathe usually green, sometimes yellowish white, spathe blade sometimes tinged purple outside; spathe tube dark, sometimes tinged reddish maroon outside; **spadix** stipitate to 5 mm, 12-18 cm long; ovary 4-6-locular, 20-25 ovules per locule. INFRUITES-CENCES with many seeds per locule.

Flowering in *Philodendron hederaceum* var. *hederaceum* is rare (known only in October and November), although the species has been found in pre-anthesis condition in every month of the year. Post-anthesis collections predominate in the dry season and earliest part of the rainy season from December through May, but there are a surprisingly large number of post-anthesis collections made in November, a month when very little flowering generally takes place (Croat, 1975, 1978). Mature fruiting collections have been made from April and May, also indicating that the species may flower predominantly in the dry season.

The range of *P. hederaceum* var. *hederaceum* is essentially that of the species. It is the most widespread taxon of *Philodendron* and, indeed, perhaps of all neotropical Araceae, ranging from San Luis Potosí State in Mexico to the Greater and Lesser Antilles, and down both slopes of the Andes, east to the Guianas and south to Brazil and Bolivia. In

Central America, the variety occurs on both slopes of the Continental Divide. In Mexico, it ranges from sea level to 1200(1500) m elevation, whereas in Middle America and Panama it ranges mostly to 450 (rarely to 900) m. It occurs principally in *Tropical moist forest* but ranges into *Premontane wet forest* and even *Tropical wet forest*. A single sterile pre-adult collection from *Tropical wet forest* transition to *Premontane wet forest* in Bajo Calima (Bay 237) is apparently also this species.

*Additional specimens examined for P. hederaceum* var. *hederaceum*. BELIZE. *Gentle 617A* (MICH). Cayo: S of Guacamallo Bridge, *Whiteford 2837* (BM); Millionario-Grano de Oro, 1700 ft., *Croat 23697* (MO); 0-10 mi. S on Hummingbird Highway, Belmopan, *Dwyer 12718* (MO); Río Frío, 1.5 mi. W of Augustine, 450 m, *Sutton et al.* 205 (BM); 0.9 km before Caracol, 16°46'N, 89°07'W, *Balick et al.* 3149 (MO). STANN CREEK: Cockscomb Basin, Jaguar Preserve, 10 km W of Maya Center, off Southern Highway, 400 m, 16°45'N, 88°35'W, *Balick et al.* 2603 (MO). TOLEDO: along highway to Punta Gorda, 1 mi. E of road to San Antonio, ca. 100 ft., *Croat 24511* (MO); Union Camp, 750 m, *Whiteford 1773* (BM, MO); Columbia River Forest Reserve, Gloria Camp, *Holst 4422* (MO); Southern Maya Mountains, Bladen Nature Reserve, 16°29'31"N, 88°54'37"W, *Davidse & Meadows 35763* (MO). COSTA RICA. *Monell s.n.* (MO); *Tondiz 13322* (G); 800 m, *Rojas 178* (MO). ALAJUELA: Cañas-Upala, 4 km NNE of Bijagua, ca. 400 m, *Croat 36320* (MO); Escuela Centroamericana de Ganadería, (near Atenas), 425 m, 9°58'N, 84°22'W, *Thompson & Raulins 1256* (CM). GUANACASTE: Tilarín, 500-650 m, *Standley & Valerio 44563* (US); 44162 (US); 44984 (US); 46563 (US); 7 km N of Cañas, *Janzen T-24* (MO); Quebrada Desprendimiento-Quebrada Sangajuela, along Río Las Flores, ca. 450 m, 10°40'N, 85°04'5"W, *Grayum et al.* 4912 (B, CAS, CR, K, MO, US); Nandayure, Península de Nicoya, 120 m, A. *Rodríguez & Estrada 141* (CR, INB). HEREDIA: La Selva Field Station, ca. 100 m, *McDowell 810* (MO); ca. 100 m; 852 (MO); *Grayum 2565* (MO). LIMÓN: 20 mi. SE of Limón, road to Punta Cahuita, near sea level, *Croat 43174* (MO); ca. 1 km N of Shiroles, ca. 50 m, *Croat 43278* (MO); Siquirres-Río Pacuare, S of Río Pacuare, 50-100 m, 10°5'N, 83°29'W, *Burger & Liesner 6953* (MO, NY); La Bomba-Cahuita, 20 m, *Gómez & Hampshire 20129* (MO); Limón-Bomba, 0-50 m, *Taylor & Shatak 4443* (DUKE); Río Jiménez, S of Guápiles, 10°17'N, 83°44'W, *Barringer 2393* (CR, MO). PUNTARENAS: Cantón de Buenos Aires, along Río Ceibo, Ujarrás, 500 m, 9°14'N, 83°18'W, *Grayum 10229* (CR, INB, K, MO, US); Gólfito, 20-200 m, 8°38'N, 83°11'15"W, *H. Schmidt 603* (CR, MO); San Vito, *Benzecry CR.CB.7C.109* (CR); Monteverde, Cordillera de Tilarín, 1500-1620 m, *Pounds 94* (MO); Cantón de Turubares, along Río del Sur, between Río Carara and Quebrada Cimarrada, 130-150 m, 9°46'30"N, 84°32'W, *Grayum 10420* (CR, MO). EL SALVADOR. ABUACHAPÁN: San Francisco Menéndez-Tacuba, 1-3 mi. above road to Río Clara Sucia, 1000-1250 m, *Croat 42135* (MO). SAN MIGUEL: ca. 50 mi. NW of San Miguel, along highway CA-1, *Croat 32799* (MO). SAN SALVADOR: vic. of Tonatepeque, *Standley 19533* (GH, NY, US). SONSONATE:

Pedral de San Isidro, ca. 3 mi. S of Lake Coatepeque, ca. 850 m, *Croat 42241* (MO). GUATEMALA. Los Amates, *Kellerman s.n.* (US); *Watson 427* (GH). **Alta Verapaz:** road to El Estor (Lago Izabal), 7 mi. E of Highway CA-14 to Cobán, 1000 m, *Croat 41480* (MO). **Escuintla:** Santa Lucia, 1045 ft., *Kellerman 4547* (US); 5285 (US). **Izabal:** Montaña del Mico, between Milla 49.5 and ridge 6 mi. from Izabal, 65–600 m, *Seyermark 38486* (F); ca. 7 mi. S of Puerto Barrios, 50 m, *Croat 41811* (MO). **Quezaltenango:** CA-2 between Coatepeque and turnoff to Colombia, 1.9 mi. W of turnoff, 580 m, 14°41'N, 91°48'W, *Croat 63394* (B, BM, MO, US). **San Marcos:** near San Rafael, 600 m, *Croat 40769* (MO). **Suchitepéquez:** 1 mi. E of Mazatenango, <500 m, *Croat 43757* (MO). HONDURAS. **Atlántida:** Lancetilla Valley, Tonacatepeque, *Pfeifer 2130* (BH, US); San José de Textiquat–El Chorzio, 100 m, *Nelson 10565* (TEFH); 4 km S of Tela, 0–100 m, *Irias 190* (TEFH, UNAH); ca. 10 mi. SE of Tela, along Río Lancetilla, 10–150 m, *Croat 42639* (MO); Quebrada Grande, ca. 10 km SW of La Ceiba, 80–140 m, 15°42'N, 86°51'W, *Liesner 26335* (MO). **Colón:** Río Selen, 7 km E of Trujillo, *Howler Site, Saunders 192* (MO). **Comayagua:** junction Río Yure–Río Humuya, 200 m, *Nelson et al. 6162* (MO). **Copán:** 13 mi. E of Copán, road to La Entrada, 750 m, *Croat 42529* (MO); Sta. Rita village, 650 m, *Molina 33668* (MO). **Cortés:** Puerto Cortés–Guatemalan border, 2–3 mi. SW of Omoa, sea level, *Croat 42555* (MO); N of Lago de Yojoa, 2–6 mi. from highway, 600 m, *Croat 42739* (MO). **Gracias a Dios:** Ahuas Bila, 200 km SW of Puerto Limpira, 100 m, *Nelson & Cruz 9316* (UNAH); 9292 (TEFH, UNAH). **Olancha:** Mpio. San Estebán, near Santa María del Carbón, 21 mi. NE of San Estebán, along road to Bonito Oriental, 440 m, 15°25'25"N, 85°34'45"W, *Davidse et al. 35571* (MO); Río Guyape, San Pedro de Catacamas–Poncaya, *Blackmore & Heath 1984* (BM); Río Wampá, 8 km S of Pisijire, 500–700 m, 15°15'N, 85°25'W, *Nelson & Clewett 594* (FSU, MO). MEXICO. **Chiapas:** El Triunfo, ca. 10 mi. NE of Escuintla, 300 m, *Croat 43859* (MO); 2 mi. SW of Guatemalan border, Highway 200 to Tapachula, 300 m, *Croat 43771* (MO); Bonampak, near ruins, 500 m, *Matuda 38715* (MO); Mpio. Ocosingo, 5 km SW of Santo Domingo, 600 m, *Davidse et al. 20425* (MO); Esperanza, Escuintla, 150 m, *Matuda 17789* (NY); 6 mi. N of Ocozacoautla, 1000 m, *Croat 40584* (MO); Palenque–Bonampak, 60 mi. SE of Palenque, ca. 400 m, *Croat 40167* (MO); Palenque–Ocosingo, Highway 199, 27 mi. SW of Palenque, 210 m, *Croat 40302* (MO); Cerro Vernal, NW side, 25–30 km SE of Tonalá, 400–600 m, *Breedlove 25617* (DS). **Guerrero:** Tierra Colorada–Xalpatlahuac, Tierra Colorada, Río Comitán, 900–1000 m, *Croat 45755* (MO); Pinotepa Nacional–Tlaxiaco, Highway 125, ca. 8.4 mi. S of Putla de Guerrero, ca. 1000 m, *Croat 45807* (MO); Tierra Colorada–Acapulco, kms 366–367, ca. 380 m, *Moore & Bunting 8840* (BH). **Jalisco:** Puerto Vallarta, 100 m, *Mexía 1314* (UC); 24.1 mi. from Autlán, ca. 300 m, *Moore & Bunting 8737* (BH); Quimixto, *Mexía 1201* (BM, CAS, DS, G, GH, MO, NY, UC, US). **Nayarit:** Miramar, ca. 10 km W of Jalcoatlán, *Dressler & Wirth 2703* (US); Singaita, *Lewis s.n.* (BH); *Phillbrick 785* (BH); San Blas, *Lewis s.n.* (BH). **Oaxaca:** Pinotepa–Tlaxiaco, Highway 125, 4.4 km S of Putla de Guerrero, 850 m, *Croat 45835* (MO); 12 km from Hwy. 200, road to Chayuco, 220 m, *Miller & Tenorio L. 524* (MO); Tuxtepec, Rincón del Tigre, Mpio. Acatlán, 2 km from Acatlán on road to Capilla, ca. 100 m, ca. 18°31'N, 96°36'W, *Gerazu et al. 2190* (CAS, MO, RSA); Tuxtepec–Oaxaca, 10 mi. S of Valle Nacional, 700 m, *Croat 39802* (MO); 0.5 mi. S of

Valle Nacional, 120 m, *Croat 39694* (MO); Esmeralda–Río Verde, Uxpanapa region, 1.1 mi. S of Esmeralda, 100 m, 17°10'N, 94°45'W, *Croat & Hannon 63233* (MO). **Puebla:** Teziutlán–Nautla, Rancho Las Margaritas, Hueytamalco, near border with Veracruz, 19°57'N, 97°16'W, *Conrad 218* (MEXU). **San Luis Potosí:** 6 mi. NW of Tamazunchale, 250 m, *Croat 39271* (MO); Tamazunchale, *Lundell & Lundell 7115* (CM); 6.5 mi. S of Matlapa, *Itiame*, 325 m, 21°18'N, 98°47'W, *Thompson et al. 1320* (CM); 1321 (CM). **Tabasco:** Tacotalpa, *Cowan 1999* (MO). **Veracruz:** 927 m, *Birdsey 226* (UC); Huatusco–Puente Nacional, El Mirador, 21 km E of Huatusco at km 45, ca. 1200 m, *Croat 44014* (MO); near Fortín, Cervecería Moxtezuma, 1000–1150 m, *Croat 39408* (MO); Mpio. Nautla, Cuatro Caminos, near Cerro Chico, 30 m, *Ventura 3690* (DS); Mpio. San Andrés Tuxtla, Estación de Biología Tropical Los Tuxtles, LOTE 71, 400 m, *Ibarra & Colín 3126* (MO), 400 m, 18°34'–36"N, 95°04'–09"W, *Masrué & Colín 3126* (MO). NICARAGUA. **Boaco:** along Hwy. 33 from Río Quilán, ca. 300–310 m, 12°35'N, 85°32'W, *Stevens 9335* (MO); Cerro Mombachito, 500–900 m, ca. 12°24'–25'N, 85°32'–33'W, *Stevens & Grigsha 14749* (MO); Quebrada Río Grande, NE del Cerro Mombachito, 600–700 m, 12°25'N, 85°32'W, *Moreno 354* (MO). **Chinandega:** Río Chiquito, El Viejo, 0–100 m, *Atwood 2635* (MO). **Chontales:** Río Bizocho–Río El Jordán, 350–550 m, ca. 12°12'–16'N, 85°15'–17'W, *Stevens & Maniel 22589* (MO); ca. 2.8 km N of Cuapa, 400–500 m, ca. 12°17'N, 85°23'W, *Stevens 3696* (MO); 3 km N of Santo Tomás, 280–300 m, 12°05'N, 85°07'W, *Moreno 16066* (MO); Juigalpa, La Libertad, Río El Bizocho, ca. 17.4 km NE of Río Mayales, 350–400 m, ca. 12°12'N, 85°17'W, *Stevens 4090* (BM, MO); 4 km NW of Villa Sandino, 100 m, *Davidse et al. 30786* (MO). **Managua:** ca. 4 km from Highway 12, near bridge of Río La Aduana, 80–100 m, ca. 12°02'N, 86°31'W, *Stevens 5394* (BM, LL, MO); Highway 8, ca. 2.4 km SW of intersection with Hwy. 2, km 2B, ca. 700 m, 11°57'N, 86°20'W, *Stevens 3990* (MO); 4 km from Highway 8 to Highway 2 intersection, 800–860 m, 11°58'N, 86°18'–19'W, *Stevens 4539* (MO); Escuela Nacional de Agricultura and Ganadería, Route 1, 12 km E of Managua, *Atwood 2930* (MO). **Matagalpa:** Matagalpa–Jinotega, km 140, ca. 900–1000 m, *Guzmán et al. 229* (MO); Cerro Musín, ca. 300 m, ca. 12°55'N, 85°16'W, *Stevens 12032* (MO); Quebrada Malacal, Hacienda La Bonanza, ca. 20 km from Matagalpa, 560 m, 13°01'N, 85°47'W, *Castro 2391* (MO). **Nueva Segovia:** 7 km SE of Santa Clara, 600–700 m, 13°40'N, 86°14'W, *Araquistain & Moreno 2191* (HNMN, LE, MO); El Terreno, 4 kms NE of El Jicaró, 500–600 m, 13°45'N, 86°07'W, *Stevens & Moreno 2215* (MO). **Río San Juan:** Río Indio, San Juan del Norte, 2 m, *Araquistain 3312* (K, M, MBM, MEXU, MO). **Rivas:** Volcán Concepción, La Esperanza, 200–400 m, 11°31'N, 85°37'W, *Robledo 1618* (ENCB, MO); Isla Ometepe, 140–350 m, 11°33'–34'N, 85°36'W, *Robledo 997* (MO); SE of “La Flor,” 300–800 m, 11°32'–34'N, 85°37'–38'W, *Robledo 1915* (MO); 400–460 m, 11°33'N, 85°37'W, *Sandino 503* (MO). **Zelaya:** road to Mina Nueva America, ca. 10 km from main road, *Stevens 12672* (MO); Ibo Tingni, N of road between Puerto Cabezas and Río Wawa, <10 m, ca. 14°9'–11'N, 83°29'–31'W, *Stevens 10663* (CM, MO); Puerto Cabezas, 0–20 m, 14°01'N, 83°22'–23'W, *Stevens 10684* (MO); Río Blanco–Río Copalar, ca. 29 km E of Río Blanco, 200–400 m, ca. 12°50'–55'N, 85°0'–05'W, *Stevens 12179* (MO); “Las Mercedes,” 160–180 m, *Vincelli 331A* (MO); Matagalpa–Waslala, near Río Las Carpas and Río Babasca, 540–580 m,

13°15'N, 85°32'W, *Sandino 2426* (MEXU, MO); La Luz, ca. 200 m, 13°44'N, 84°47'W, *Ortiz 1606* (F, MO); Cerro Waylawas, E side of Central Range, ca. 100–200 m, 13°38–39'N, 84°48–49'W, *Pipoly 4172* (MO); 6 km SE of Waslala, 520–560 m, 13°16'N, 85°24'W, *Moreno 17279* (MO); Estación Experimental "El Recreo," ca. 15 m, 12°09'N, 84°17'W, *Rios 176* (MO); ca. 80 m, 13°39'N, 84°48'W, *Pipoly 4450* (MO); Río Prinzapolka–Quebrada San Rafael, Waní, ca. 100 m, ca. 13°42'N, 84°50'W, *Pipoly 4593* (MO); ca. 0–100 m, ca. 13°42'N, 84°50'W, 4739 (MO); Río Waní, Boca de Waspuke, 90–100 m, ca. 13°44'N, 84°53'W, *Stevens 7195* (MO); Rama, Atwood & Moore 897 (MO); Caño Monte Cristo, "La Grapera," ca. 10 m, 11°33'N, 87°48'W, *Moreno & Sandino 14709* (MO); E of campo Germán Pomares, ca. 60–90 m, 11°36'N, 83°52'W, *Moreno 15137B* (MO); ca. 10 m, 11°36'N, 83°51'W, *15187* (MO); Caño Monte Cristo, 1 km before the camp Germán Pomares, ca. 10 m, 12°35'N, 83°51'W, *Moreno 14839* (MO); *14843B* (MO); "Kurinwacito," 80–100 m, 13°08'N, 84°55'W, *Moreno 23699* (F, MO); Cerro El Ocote, 700 m, 13°38'43"N, 85°07'06"W, *Ortiz 1093* (MO); Comarca El Hormiguero, W of Río Uli, *Neill 1921* (MO); Mpio. Rama, Loma Buena Vista, 100–150 m, 12°08'N, 84°12'W, *Robledo 620* (MO); Mpio. Siuna, Siunawás, ca. 200 m, 13°43'N, 84°45'W, *Ortiz 1481* (MO); Santa Rosa, *Ortiz 102* (MO); Negro Wás Sector, El Empalme–Rosita, <200 m, ca. 13°45'N, 84°25'W, *Ortiz 2163* (MO); Río Kuhlali, ca. 7 km W of Río Iyas, ca. 200 m, ca. 13°29'N, 85°16'W, *Stevens & Moreno 19278* (MO); Río Matis, Waní–Siuna, ca. 0–100 m, ca. 13°43'N, 84°49'W, *Pipoly 4685* (MO); ca. 0–100 m, *Stevens 4686* (MO). PANAMA. **Bocas del Toro:** S end, Cayo García, 0–5 m, *Peterson & Annable 7287* (MO); Chiriquí Lagoon, *von Wedel 1253* (GH, MO); 2668 (GH, MO); Gualaca–Chiriquí Grande, 4.2 mi. S of Chiriquí Grande, near sea level, 8°55'N, 82°09'W, *Croat 66813* (MO, TEX); Changainola–Almirante, Station Millá 7.5, <100 m, *Croat 38130* (MO); Río Oeste, S of Almirante, 5 m, *Peterson & Annable 7226* (MO). **Canal Area:** end of Pipeline Road, 19 km NW of Gamboa, 25–50 m, *Nee & J. D. Smith 11070* (MO, US); Fort San Lorenzo, *Maxon & Valentine 7001* (US); 40 m, 9°03'N, 79°37'W, *Croat 69834* (B, CAS, CR, MO); near Madden Dam and along Azote Caballo Road, near Alahueta, 90–100 m, *Dodge 16592* (MO); road to Fort Gulic, *Lazar & Blum 5412* (MO, NY); Barro Colorado Island, *Croat 6586* (MO, NY, US); *Wetmore & Woodworth 874* (GH); *Welch 19833* (MO, NY); *Croat 6066* (MO); *11758* (MO); *9223* (MO); *5878* (MO); *10357* (MO); *12802* (MO); *10744* (MO); *Starry 24* (MO); *Netting 60* (MO); *Hutchison & Wright 2920* (NY, US); *Croat 11744* (MO, SCZ); *10383* (MO, SCZ); *7129* (MO, SCZ); Madden Forest Preserve, vic. George W. Green Park, *Welch 19853* (MO, NY, RSA); Río Pequet, near Salamanca Hydrographic Station, 70–80 m, *Dodge & Ortiz 16592a* (MO); Curimdu, 8°58'N, 79°32'55"W, *Croat & Zhu 76204* (MO). **Chiriquí:** W of Gualaca, 100 m, 8°29'N, 82°17'W, *Churchill & de Nevers 4482* (MO, RSA). **Coelé:** El Valle de Antón, *Folsom & Kauke 2776* (MO); *Kennedy et al. 2209* (MO, US). **Colón:** near Palmas Bellas, *Thomas 47* (MO); vic. of Miguel de la Borda, *Croat 9895* (MO); *9885* (MO); 10 mi. SW of Portobelo, 2–4 m, from coast, 10–200 m, *Liener 1103* (MO, NY, US); Portobello, 5–100 m, *Plazier 2424* (US); France Field–Catal, *Standley 30439* (US); Río Guanache, *D'Arcy 9724* (MO); *Gentry 6313* (MO); 100 m, 9°30'N, 79°39'W, *Croat 75193* (MO). **Darién:** 3 mi. N of Santa Fe, *Tyson et al. 4616* (MO, SCZ); Canglón–Yaviza–Río Chucunaque, 7.7 mi. E of Canglón, 50 m, 8°20'N, 77°50'W, *Knapp &*

*Mallet 3944* (MO, US). **Los Santos:** Tonosí–Jobero, 50–80 m, *Croat 34454* (MO). **Panamá:** Corozal Road, near Panamá, *Standley 26861* (US); *26841* (US); Río Tapia, *28112* (US); 4 mi. E of Panama City, road to Tocumen, *Tyson 3483* (FSU, MO, SCZ); Cerro Jefe region, *D'Arcy et al. 15520* (MO); 9°15'N, 79°30'W, *Croat & Zhu 76215* (CR, MO, P, TEX); San José Island, *Johnston 718* (GH); *919* (GH); *921* (GH, US). **San Blas:** Nusagandi, El Llano–Certi road, Mile 10.3, 300 m, 9°20'N, 79°W, *Croat & Zhu 76576* (MO). **Veraguas:** S of Santa Fe, ca. 450 m, *Nee 8041* (MO); Bahía Honda, *Elmore 830* (US); ca. 1 km above Cahasas on road to Los Valles, 230 m, *Croat 37063* (MO).

***Philodendron hederaceum* var. *kirkbridei***

Croat, var. nov. TYPE: Costa Rica. Alajuela: eastern rim of Laguna Hule, ca. 1.5 km N of Angeles, along road through pasturelands, 840–860 m, 8 July 1972, *Luteyn 3350* (holotype, MO–2173874; isotype, DUKE). Figure 224.

Planta hemiepiphytica; internodia 2–28 cm longa, 1–2.5 cm diam.; cataphylla 6–19 cm longa, incoastata aut interdum leniter 2-costata, decidua intacta; petiolus (6.5)8–22.5 cm longus, 3–6 mm diam.; lamina (11)16–29.5(42) cm longa, (8)10.3–23(35) cm lata; inflorescentia 1; pedunculus 5.5–7.5 cm longus; spatha 14–24 cm longa, lamina spatulae flaviviridis, extus cremascenti vis. apicem, intus pallidius viridis; pistilla 5-locularia; loculi 20–25-ovulati; loculi cum ca. 24-semibus.

Hemiepiphyte; internodes 2–28 cm long, 1–2.5 cm diam., weakly flattened on one side with two sharply raised ribs on the side above petiole, very weakly sulcate on the opposite rounded side, prominently ribbed throughout its circumference, the ribs smooth or prominently warty, 2–28 cm long, 1–2.5 cm diam., medium to dark green, matte, drying reddish brown; cataphylls 6–19 cm long, unribbed or sometimes weakly 2-ribbed, deciduous intact; **petioles** (6.5)8–22.5 cm long, 3–6 mm diam.; **blades** (11)16–29.5(42) cm long, (8)10.3–23(35) cm wide; upper surface medium to dark green, drying brown to greenish brown, lower surface medium green, glossy, drying brown to greenish brown; primary lateral veins 3–4 per side, departing midrib at a 45–50° angle. **INFLORESCENCES** 1 per axil; peduncle 5.5–7.5 cm long; **spathe** 14–24 cm long, spathe blades yellow-green becoming cream-colored toward apex outside, lighter green inside; **spadix** stipitate to 1 cm; 13–20 cm long; ovary 5-locular, 20–25 ovules per locule. **INFRACTESCENCE** with about 24 seeds per locule.

Flowering in *Philodendron hederaceum* var. *kirkbridei* occurs during the late dry season and early rainy season (probably as early as March and as late as September) based on post-anthesis and early fruiting collections. No collections have been made

at anthesis. Immature fruits have been collected in April, July, and August.

*Philodendron hederaceum* var. *kirkbridei* ranges from Costa Rica to Ecuador at (250)300 to 900 m in *Premontane wet forest* and *Tropical wet forest*. The variety has been collected only once in both Suriname and Ecuador. It was collected in Ecuador at Reserva ENDESA (0°5'N, 79°02'W), an area of *Premontane rain forest*. It is to be expected in Colombia.

This variety is recognized by its scandent habit, slender stems, petioles shorter than the blades, ovate-cordate brownish-drying blades and especially by its dried stems, which are reddish brown and conspicuously ribbed. Specimens from Panama have stems also conspicuously and densely warty while specimens in Costa Rica have stems reddish brown and conspicuously ribbed but are smooth rather than warty.

The variety differs from the autonymic variety primarily in occurring at generally higher elevations in wetter forests and by its reddish brown, conspicuously ribbed stems. In parts of Central America and in South America where both species occur, *P. hederaceum* var. *kirkbridei* occurs usually above 500 m in *Premontane wet* and *Tropical wet* forests. In contrast, *P. hederaceum* var. *hederaceum* has stems that dry smooth and green and occur primarily in tropical moist forest. Though there is overlap in the elevation range and even life zone (both may sometimes be found in *Premontane wet forest*), the two varieties do not occur in the same sites and can be readily distinguished by their stems.

*Philodendron hederaceum* var. *kirkbridei* is named in honor of Joseph Kirkbride (BARC), who first collected the taxon in 1968 on Cerro Caracoral in Coclé Province in Panama while a graduate student at the Missouri Botanical Garden.

*Additional specimens examined.* COSTA RICA. **Alajuela:** Catus-Upala, 8 km N of Bijagua, 300 m, *Croat* 36498 (MO); Cariblanco, 900 m, 10°15'N, 84°11'W, *Lent* 3591 (F). **Heredia:** 3 mi. S of Cariblanco, 760 m, *Croat* 35837 (MO). PANAMA. **Coclé:** 710–800 m, 8°39'N, 80°36'W, *Croat* 68758 (MO, PMA, US); Cerro Caracoral, *Kirkbride* 1098 (MO); Alto Calvario, 9.4 km above El Copé, 750–900 m, *Croat* 44741 (MO); La Mesa, above El Valle de Antón, 900–930 m, *Croat* 37471 (MO); Cerro Gaital, 900–1000 m, 8°37'N, 80°08'W, *Croat* 67236 (MO). **Colón:** Río Piedras, Río Piedras Lumber Road, 6.7 mi. E of Sabanita, 250 m, 9°22'30"N, 79°41'30"W, *Croat* 75158 (MO, PMA). **Panamá:** El Llano–Cartí Road, 12 mi. above Pan-American Highway, 200–500 m, *Croat* 22909 (MO); Mile 7, near El Llano, 460 m, 9°19'N, 79°59'W, *Croat* 75108 (MO, PMA, US); Km 12–16, *Kennedy et al.* 3146 (MO, US). **San Blas:** 1 mi. S of Nusa-

gandi, 9 mi. N of Interamerican Highway, 350 m, 9°20'N, 79°W, *Croat & Zhu* 77001 (CM, MO).

ECUADOR. **Pichincha:** between Quito and Puerto Quito, Km 113, 800 m, 0°5'N, 79°2'W, *Rodríguez* 282 (MO).

#### *Philodendron hederaceum* var. *oxycardium*

(Schott) Croat, stat. et. comb. nov. Basionym: *Philodendron oxycardium* Schott, Syn. Arid. *Philodendron scandens* subsp. *oxycardium* (Schott) G. S. Bunting, Gentes Herb. 10: 163. 1968. TYPE: Mexico: exact origin unknown (holotype, W? lost). Schott ic. 2714 (neotype, here designated, W). [See fig. 17 in Bunting, 1968.]

Internodes smooth, sometimes with 2 ridges or weakly angular on one side, semiglossy, 5–26 cm long, to 1 cm diam., usually longer than broad, drying greenish to pale brownish green; cataphylls 19 cm long, unribbed, green; petioles 8–23 cm long; blades broadly ovate, semiglossy, long-acuminate to abruptly acuminate at apex, cordate at base, 26–32 cm long, 18–22 cm wide (ca. 1.5 times longer than wide), 2.5–3 times longer than petiole, broadest near the middle, upper surface bright green, lower surface yellow-green; sinus narrowly triangular to closed, to 5 cm deep; basal veins 2–3 per side, with 1–2 free to base, and with the first and or second free to base, 2–3 coalesced to 1 cm; primary lateral veins 2–3 per side, departing midrib at a 45° angle. INFLORESCENCES (based on dried specimen) with peduncle 4 cm long, 7 mm diam.; spathe 14 cm long; spathe blade 6.5 cm long; spathe tube 7.5 cm long; spadix sessile, cylindrical, 12 cm long; pistillate portion 4 cm long; staminate portion 8 cm long; fertile staminate portion drying reddish brown.

Flowering in *Philodendron hederaceum* var. *oxycardium* is poorly known. A single fertile collection with immature fruits in January was seen.

*Philodendron hederaceum* var. *oxycardium* is apparently endemic to Mexico, known only from the Gulf slope in the states of Jalisco, Nayarit, Oaxaca, San Luis Potosí, and Veracruz at 120 to 580 m, principally in areas of *Tropical moist forest*. Material of *P. hederaceum* [as *P. scandens*] collected in Jamaica (Bunting, 1968) is sterile.

*Philodendron hederaceum* var. *oxycardium* is similar in appearance to variety *hederaceum*, except that leaves of the former are firmer in texture and glossy in both the juvenile and adult forms. In outline, the leaf blades of *P. hederaceum* var. *oxycardium* are often somewhat broader and more abruptly acuminate than for variety *hederaceum* (Bunting, 1968). The epidermal cells have a more or less

flattened epidermis on the upper surface (Bunting, 1968), which causes it to be glossy. In contrast, the epidermal cells of *P. hederaceum* var. *hederaceum* are rounded and sometimes conical, giving leaves of that variety their characteristic velvety sheen.

Though Bunting (1968) treated this taxon as a subspecies, it appears to overlap geographically with the typical material so it will be treated here as a variety.

*Additional specimens examined.* MEXICO. **Nayarit:** San Blas, Ferris 5440 (DS). **Oaxaca:** Tuxtepec-Oaxaca, 6-14 mi. from bridge at Valle Nacional, ca. 580 m, Moore & Bunting 8905 (BH). **San Luis Potosí:** N of Tamazunchale, Clark 7408 (MO, NY); Tamazunchale, 120 m, Aguirre & Reko 334 (NY). **Veracruz:** 5.7-6 mi. from Catemaco on road to Sontecomapan, ca. 380 m, Moore & Bunting 8937 (BH); just before Papantla on road from Tecolutla, Moore & Bunting 8953 (BH); NW of Misantla, 2.4 mi. on road to Martínez de la Torre, ca. 320 m, Moore & Bunting 8944 (BH); Conejo-Huatusco, Km 45, near Puente Nacional, slopes of barranca de Santa María across highway from Hacienda El Mirador, Moore & Bunting 8869 (BH); Dist. Papantla, Kelly 16 (BH).

***Philodendron heleniae* Croat, sp. nov. TYPE:**

Panama. San Blas: El Llano-Carri road, 14 km N of Pan-American Hwy., 300 m, 9°15'N, 79°W, 12 July 1988, Croat 69244 (holotype, MO-3599872; isotypes, B, CAS, COL, CR, F, K, MEXU, NY, PMA, SCZ, US, VEN). Figures 227-230.

Planta plerumque hemiepiphytica, raro terrestris; internodia 4-12 cm longa, 2-3 cm diam.; cataphylla 15-36 cm longa, 1-2-costata, decidua; petiolus teres aut D-formatus, 14-41.5 cm longus, 6-10 mm diam.; lamina anguste ovata, 25-52 cm longa, 6-26 cm lata, in sicco flavibrunnea; inflorescentia 2-10; pedunculus 2-6 cm longus, 4-6 mm diam.; spathe 5-10 cm longa, rubrimaronea; lamina spatheae extus pallidiori vs. apicem, intus alba, dilute suffusa marronina; tubo spatheae intus strimarronino; pistilla (5)6-9-locularia; locali (1)3-4-ovulati.

Usually hemiepiphytic, rarely terrestrial; vining or appressed-climbing, stem appressed-climbing to somewhat scandent, sap tannish; internodes scurfy, 4-12 cm long, 2-3 cm diam., much longer than broad, medium green, drying reddish brown, epidermis smooth or irregularly folded and ridged, sometimes fissured with minute cracks perpendicular to axis, drying with folds longitudinally and usually with fissures transversely; roots dark brown, to 50 cm long, 1-2 mm diam.; cataphylls thin, 15-36 cm long, 1-ribbed to sharply 1-ribbed, or sharply 2-ribbed, green, tinged red, speckled violet-purple, deciduous, rounded, apiculate at apex; petioles 14-41.5 cm long, 6-10 mm diam., ± terete or to slightly thicker than broad, or bluntly D-shaped, firm, medium green, tinged violet-purple

at base and apex, bluntly flattened adaxially; geniculum 9 mm long, thicker and paler than petiole; blades narrowly ovate, subcoriaceous to coriaceous, strongly bicolorous, acuminate at apex (the acumens sometimes apiculate, inrolled, 5 mm long), usually weakly subcordate, sometimes obtuse, truncate or rounded at base, 25-52 cm long, 6-26 cm wide (2-3.37(4.2) times longer than wide), (1.1-1.8(2.30) times longer than petiole), upper surface dark green, semiglossy, drying yellow-brown, lower surface much paler and with dark secretory canals, weakly glossy to matte, drying yellowish to reddish brown; sinus arcuate, 1-1.5 cm deep when present; midrib broadly convex or weakly raised to flat, pale green above, bluntly acute and thicker than broad, reddish slightly paler than surface below; basal veins lacking; primary lateral veins (6-7)8-10(11-13) per side, departing midrib at a 30-40° angle, weakly and narrowly sunken above, raised and paler than surface below; minor veins weakly visible, arising from the midrib only, often interspersed with intermittent secretory ducts, the surface often minutely granular below. INFLORESCENCES 2-10 per axil; peduncle 2-6 cm long, 4-6 mm diam., green, tinged red to reddish; spathe recurved (curved downward), semiglossy, 5-10 cm long (1.2-2.5 times longer than peduncle), reddish maroon throughout; spathe blade pale toward apex outside, white, weakly tinged maroon inside; spathe tube 3-4 cm long, dark maroon inside; spadix sessile; cylindrical to clavate, protruding forward at anthesis, 5.8-7.5 cm long, ± uniform throughout; pistillate portion pale green to yellowish green, cylindrical to slightly ovoid, 2-3.5 cm long, 5-7 mm diam. throughout; staminate portion cream to white, cylindrical or clavate, 3.4-5 cm long, 6-9 mm diam. at base, 9-10 mm diam. at middle, 5-9 mm diam. ca. 1 cm from apex, broadest at or above the middle, or equally broad throughout, as broad as or slightly broader than the pistillate portion; sterile staminate portion not detectable; pistils 1.4-1.7 mm long, 0.5-1.4 mm diam.; ovary (5)6-9-locular, 0.7 mm long, 0.7 mm diam., locules 0.9-1.2 mm long, 0.3-0.4 mm diam., ovule sac 0.4-1 mm long, with sub-basal placentation; ovules (1)3-4 per locule, 1-seriate, contained within translucent ovule sac, 0.2-0.4 mm long, longer than funicle; funicle 0.1-0.2 mm long (can be pulled free to base), style 0.2-0.4 mm long, 0.4-0.6 mm diam., similar to style type B; styler canals tiny, difficult to see emerging; style apex flat; stigma discoid or subdiscoid, unlobed, 0.4-0.5 mm diam., 0.1-0.2 mm high, covering entire style apex except for the center; the androecium truncate, margins 4-6-sided, sometimes weakly scalloped; thecae oblong, 0.1-

0.3 mm wide,  $\pm$  parallel to one another. INFRU-  
TESCENCE with berries white, 3.3 mm long, 2.4  
mm diam.; seeds ca. 6 per berry, reddish brown,  
1–1.1 mm long, 0.3–0.5 mm diam.

Flowering in *Philodendron heleniae* occurs in  
Panama during the late dry season and much of the  
rainy season, March through October (except April,  
May, and September). South American collections  
broaden that range to include the whole year (Janu-  
ary through December, except May and Septem-  
ber). Post-anthesis collections have been made  
from June to October.

*Philodendron heleniae* ranges from Panama (to  
be expected in the Cordillera de Talamanca in east-  
ern Costa Rica) to Ecuador, from 20 to about  
1040(–1450) m elevation in *Tropical wet forest* and  
*Premontane rain forest* in Panama, Colombia, and  
Ecuador. In addition, it occurs in pluvial wet forest  
in Colombia.

*Philodendron heleniae* is a member of *P.* sect.  
*Calostigma* subsect. *Oligocarpidium*. This species  
is characterized by its scandent habit; internodes  
much longer than broad drying with longitudinal  
folds and usually transverse fissures; petioles terete  
to bluntly D-shaped; and narrowly ovate, subcor-  
date, yellow-brown-drying blades with dark secre-  
tory canals visible on the lower surface. Especially  
characteristic are the clusters of 2–10 small inflo-  
rescences with externally red spathes.

*Philodendron heleniae* can be confused with *P.*  
*lentii*, which also comprises more or less scandent  
plants with subcordate leaf blades. That species  
differs, however, in having one to two large, mostly  
white inflorescences per axil and more broadly  
ovate blades drying with many, pale, sunken veins  
on the upper surface and no dark secretory vessels  
on the lower surface.

Specimens from the Pacific slope of South Amer-  
ica (e.g., *Croat* 58424, *Tipaz et al.* 1318) are on  
average larger, with leaves ranging up to 60 cm  
long and 30 cm wide and with spathes occasionally  
to 12 cm long. Still, these probably do not differ  
sufficiently to represent even another subspecies.

*Lawrance* 817 from Boyacá Department at El  
Humbo (at 914 m in the Río Magdalena drainage)  
in Colombia closely matches the material from the  
Pacific slope. If it proves to be *P. heleniae* it would  
be the first collection from the Eastern Cordillera  
of the Andes.

Some South American collections from the Am-  
azon basin may belong to this species. These col-  
lections are from both Colombia (*Pipoly et al.*  
15892, 16027 in Amazonas) and Ecuador (*Gudiño*  
1160 and *Thomas & Ríos* 6708 in Pastaza Province

and *Cerón* 3360 and *Bennett et al.* 4526 in Napo  
Province). However, most of the collections from  
the Amazon basin do not mention spathe color, al-  
though *Gudiño* 1160 indicates the spathes to be  
greenish red.

Particularly interesting is *Thomas & Ríos* 6708,  
with spathes described as becoming creamy white,  
something that never happens with Central Amer-  
ican material of *Philodendron heleniae*.

The species was first collected in 1970 by Jim  
Luteyn and Helen Kennedy in the vicinity of El  
Valle in Coclé Province, Panama. It is named in  
honor of one of the collectors, Helen Kennedy, who  
at the time worked for the Missouri Botanical Gar-  
den as the Curator of Summit Herbarium.

*Additional specimens examined.* PANAMA. **Bocas  
del Toro:** Fortuna Dam area, Gualaca–Chiriquí Grande,  
6.6 mi. N of bridge over Fortuna Lake, 780 m, 8°45'N,  
82°18'W, *Croat* 66780 (MO). **Coclé:** El Valle region, La  
Mesa, 800–900 m, 8°38'N, 80°09'W, *Croat* 67154 (AAU,  
CM, KYO, L, MEXU, MO, PMA, TEX, US); El Valle, 5  
mi. N of town, 2200 ft., *Hammel & Kress* 13427 (DUKE);  
8 mi. N of El Valle ca. 880 m, *Luteyn & Kennedy* 1734  
(DUKE); 880 m, *Croat* 37576 (MO); 860 m, 8°37'N,  
80°08'W, *Croat & Zhu* 76738 (AAU, CM, DUKE, M, MO,  
S, SEL, TEX); slopes of Cerro Gaital, 800–900 m, 8°37'N,  
80°07'W, *McPherson* 11209 (L, MO, NY, P). **Darién:** Río  
Tuquesa, 2 km by air from Continental Divide, vic. of  
mining camp of Tyler Kittredge, *Croat* 27097 (CAS, F, K,  
MO, PMA, US). **Panamá:** El Llano–Cartí Road, 10.1 mi.  
above Inter-American Highway, 325–350 m, *Croat* 67351  
(MO, NY, PMA, US, WIS); Mile 6.8, 350 m, 49106 (MO);  
Mile 10, 330 m, 33778 (F, MO), 33817 (MO); Mile 7, 460  
m, 9°19'N, 79°59'W, 75114 (CAS, MO); 550 m, 9°43'N,  
78°60'W, 60502 (MO); Mile 10.6, ca. 400 m, 9°17'N,  
78°58'W, *Miller et al.* 869 (HUA, MO, PMA); El Valle de  
Madroño–La Saena, 2.5 mi. N of El Valle de Madroño,  
180 m, 9°14'25"N, 79°05'W, *Croat & Zhu* 77048A (MO);  
Cerro Jefe–Cerro Brewster ["Panama/San Blas"], 600–800  
m, 9°17'N, 79°17'W, *Hammel & de Nevers* 13553 (MO).  
**San Blas:** El Llano–Cartí Road, 1–5 mi. N of Nusagandi,  
250–300 m, 9°16'N, 79°W, *Thompson* 4660 (CM, MO);  
Nusagandi–Cartí Road, 400 m, 9°18'N, 78°58'W, *Mc-  
Donagh et al.* 93 (BM, MO); ca. 350 m, ca. 9°15'N, 79°W,  
*McPherson* 9762 (MO); 350 m, 78°15'W, 9°19'N, *de Ne-  
vers & Todzia* 3530 (COL, MO, PMA, QCA, RSA, TEX);  
10.1 mi. N of Interamerican Highway, then ca. 0.5 mi. N,  
Paseo Mariska near road, 300 m, 9°20'N, 79°W, *Croat &  
Zhu* 77020 (MO); El Llano–Cartí Road, 23–29 km from  
Pan-American Highway, 300–400 m, 9°22'N, 78°69'W,  
*Knapp* 1830 (MO); 1840 (MO, PMA); Km 15, 350 m,  
9°20'N, 78°58'W, *Galdames et al.* 1288 (PMA); km 17–  
19, 350 m, 9°19'N, 78°55'W, *de Nevers & Herrera* 7963  
(CM, MO, US); Cerro Obu, 400–500 m, *de Nevers et al.*  
8068 (MO, NY). **Veraguas:** Santa Fe region, Alto Piedra-  
Calvebora, Río Dos Bocas Valley, 350–400 m, *Croat*  
27370 (F, K, MO, PMA); 27397 (MO).

COLOMBIA. **Antioquia:** Medellín–Bogotá, 12.4 km  
from entrance to San Luis 800 m, 5°59'N, 74°59'74"W,  
*Betancur et al.* 622 (MO); 4.8 km E of San Rafael, 1040  
m, 6°18'N, 75°W, *Brant & Roldán* 1486 (HUA); Quebada  
Honda, Río Guatapé, *Orozco et al.* 898 (COL); Río Gua-  
tapé, 1250 m, *Orozco et al.* 955 (COL); San Carlos, Finca

"El Desespero," Alto de Samaná, near Miraflores, 4 hrs. SW of Alto de Samaná, 710–820 m, 6°5'N, 74°56'W, *Callegas et al.* 8628 (NY); San Luis, Medellín–Bogotá, La Tebaida, 1010–1060 m, 6°8'N, 75°10'W, *Callegas et al.* 4015 (MO); Sector Río Samaná–Río Claro, 750 m, *Hernández et al.* 551A (HUA); San Luis–Puerto Triunfo, SE of Granadá, S of San Carlos, ca. 3 km SE of San Luis, ca. 800 m, 5°57'N, 74°57'W, *Croat* 52038 (COL, MO, NY); Gómez Plata–Yolombo border, Sector La Cañana, along Río Porce, km 14 via Amalí, 1030 m, *Callegas et al.* 2338 (MO); Sonson, Rioverde region, Hacienda "La Soledad," on the banks of Quebrada "Curubital," *Gutiérrez* 35552 (UC); Amalí–Fraguas, NE of Salazar, 23–26.5 km from center of Amalí, 1220–1300 m, 6°58'N, 74°59'W, *MacDougal et al.* 4042 (MO); **Chocó:** Bahía Solano, S of airport, ca. 100 m, 5°13'30"N, 76°21'30"W, *Croat* 57458 (CHOCO, COL, JAUM, MO, US); ca. 2 km E of Playa de Oro, ca. 200 m, 5°20'N, 76°23'W, *Croat* 57424 (CHOCO, COL, MO); Mecana (N of Bahía Solano), Quebrada Resaquita, 30 m, *Juncosa* 2536 (MO); Quibdó–Istmina, 6.6 km S of Quibdó, <100 m, 5°33'N, 76°37'W, *Croat* & *Cogollo* 52156 (MO); Río Pató, Las Animas–Pató, Serranía de Baudó, 10 km SE of Pató, 5°17'N, 76°45'W, *Croat* 56071 (MO), *Croat* 56107 (MO); Río Sucio, Cerro del Cuchillo, zona de Urabá, 520 m, *Cárdenas* 468 (MO); Parque Nacional de Utría, Río San Pichí, 0–100 m, 6°20'N, 77°20'W, *García & Aguilimpia* 458 (MO); Río San Juan, Quebrada del Taparal, 5–20 m, *Castrocasas* 21462 (F). **Risaralda:** Mpio. Mistrato, corregimiento San Antonio de Chamí, vereda la Unión, carretera hacia La Mesenia, 1450–1700 m, 5°28'N, 75°54'W, *Betancur et al.* 3520 (MO). **Valle:** Cali–Buenaventura, Río Digua valley, 1 km E of La Cascada, 340 m, *Croat* 38579 (MO, PMA); Bajo Calima area, Quebrada La Sierpe, 5 m, 4°10'N, 77°10'W, *Forero et al.* 4006 (MO); Bahía Málaga, 0–20 m, 4°N, 77°30'W, *Gentry et al.* 53362 (MO); carretera Gasolina 6 km S of main road between Cali–Buenaventura highway and Málaga, 50–80 m, 3°56'N, 77°07'30"W, *Croat* 69411 (CM, MO); 22.3 km beyond Pulpapel Headquarters, 4°02'N, 77°07'W, *Croat* 61275 (CUV, MO), ECUADOR. **Carehí:** Tulcán, Reserva Indígena Awá, 650–100 m, 1°N, 78°24'W, *Tipaz et al.* 1318 (MO, QCNE). **Esmeraldas:** Lita, 550–650 m, *Madison et al.* 4990 (F, K, SEL); Lita–San Lorenzo, Km 18, 0°55'N, 78°28'W, *Croat* 72389 (MO); San Lorenzo, Mun. Ricaurte, Reserva Indígena Awá, 300 m, 1°10'N, 78°32'W, *Tipaz et al.* 2068 (MO); Río Jordán, NE of Las Golondrinas, near San Isidro, 300 m, 0°20'N, 79°12'W, *Palacios* 11485 (MO, QCNE). **Imbabura:** Lita, 600 m, *Cobb* 29 (MO). **Pichincha:** Río Verde, 2 km SE of Santo Domingo de Los Colorados in Cooperativa Santa Marta #2, 530 m, *Dodson* 7403 (MO, QCNE); Santo Domingo de los Colorados, Río Baba, 28 km S, 350 m, *Dodson & Thien* 1187 (MO).

**Philodendron immixtum** Croat, sp. nov. TYPE:

Panamá. Panamá: Comarca de San Blas, Río Playón Chico, 50–100 m, *H. Herrera et al.* 1167 (holotype, MO–4256423; isotypes, AAU, K, MEXU, PMA, US). Figures 233–236.

Planta hemiephytica; internodia (2)10–15 cm longa, (7 mm) 1–1.5 cm diam.; petiolus subtere, actue D-formatus ad debiliiter planatus; lamina anguste ovata vel ovato-elliptica, 16–36 cm longa, 5–12 cm lata; inflorescentia solitaria; pedunculus 5–13 cm longus; spathe 9–19 cm longa, lamina albo-viridis vel alba, tubo viridis extus, rubo vel

purpureo intus; spadix 10–11 cm longus; ovarium (3)4–(6)8-loculare; loculi (1)2) ovulati.

Hemiephytic; stem scandent, sap reddish, watery; internodes long, slender, ± glaucous, (2)10–15 cm long, (0.7)1–1.5 cm diam., much longer than broad, dark olive-green matte to weakly glossy, becoming gray-green, drying light brown, epidermis sometimes cracking with loose flakes; roots ca. 5 per node, to 15 cm long, 2–3 mm diam., tannish gray, drying reddish brown, ± smooth, long; cataphylls somewhat spongy, to 17 cm long, unribbed, weakly 2-ribbed or sharply (1)–2-ribbed (ribs 2 mm high), pale green, dark short-lineate, drying pale brown to yellowish brown, usually some deciduous, sometimes persisting somewhat intact, eventually fibrous, obtuse at apex, margins paler, yellowish; **petioles** 7.5–16(21) cm long, 2–10 mm diam., subterete, sharply D-shaped to slightly flattened adaxially, rounded abaxially, surface dark green-striate, matte to semiglossy, shortly dark lineate, with dark green ring around apex; **blades** narrowly ovate to ovate-elliptic, or rarely oblong-elliptic, subcoriaceous, semiglossy, strongly bicolorous, abruptly acuminate at apex (the acumen tightly inrolled to apiculate, 2–4 mm long), subcordate, rarely rounded at base, 16–36 cm long, 5–12 cm wide (1.9–3.7 times longer than wide), (1.3–3.1(3.8) times longer than petiole), about twice as long as petiole, broadest below middle, margins narrowly hyaline, upper surface drying green to grayish to brownish green; posterior lobes short, about as broad as long and held close to petiole; sinus shallow, 0–2.5(3) cm deep; midrib flat to weakly convex, concolorous or slightly paler than surface above, convex to bluntly acute, paler than surface below; basal veins lacking; primary lateral veins 4–8 per side, departing midrib at a 50–60° angle, ± straight to weakly arcuate to the margins, sunken and paler than surface above, weakly raised and darker than surface below; interprimary veins weakly raised and darker than surface; minor veins distinct to weakly visible, darker than surface below, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 1 per axil; peduncle 5–13 cm long, 3–7(11) mm diam., dark green; **spathe** 9–19 cm long, (1.1–2.1 times longer than peduncle), green, densely and obscurely short-lineate, narrowly acuminate at apex; spathe blade pale greenish white to white outside, pale greenish white, dark lineate inside; resin canals appearing as continuous lines, drying reddish brown, resin droplets forming on spathe surface within; spathe tube dark green in back, paler in front on open edges outside, red or violet-purple inside; **spadix** gradually tapered to apex,

protruding out of the end of spathe, bluntly rounded to obtuse at apex, 10–11 cm long, broadest above the middle, constricted slightly above sterile staminate portion; pistillate portion pale green, cylindrical, 3.4–4.2 cm long, 3.7 cm long in front, 2.7 cm long in back, 9–11 mm diam. at apex, 10–11 mm diam. at middle, 9–11 mm wide at base; staminate portion 6.8–7.5 cm long; fertile staminate portion white, cylindrical to clavate, 9–12 mm diam. at base, 1.1–1.3 cm diam. at middle, 7–12 mm diam. ca. 1 cm from apex, broadest at the middle, broader than the pistillate portion, broader than the sterile portion; sterile staminate portion as broad as, narrower than, or broader than the pistillate portion,  $\pm$  12 mm diam.; pistils 1.9–2.3 mm long, 1.3–1.9 mm diam.; ovary (3)4–6(8)-locular, walls sometimes embedded with granular, crystal-like particles, locules 0.7–1.3 mm long, 0.4–0.7 mm diam., ovule sac 0.6–1.3 mm long, with sub-basal placentation; ovules 1(2) per locule, contained within transparent ovule sac, 0.4–0.7 mm long, longer than funicle; funicle 0.2–0.6 mm long (can be pulled free to base), style 0.3 mm long, 0.9–1.4 mm diam., similar to style type B; stigma subdiscoid, unlobed, 0.7–0.9 mm diam., 0.2 mm high; the androecium truncate, margins 4–6-sided; thecae oblong to slightly elliptical, 0.4 mm wide,  $\pm$  parallel to one another; sterile staminate flowers 4–6-sided, 1–1.8 mm long, 1.1–1.7 mm wide. INFLORESCENCE with spadix yellowish orange, 8 cm long, 3 cm wide; berries concave at apex; seeds 1 per locule, 1.5–1.8 mm long, 0.6–0.7 mm diam., yellow-brown.

Flowering in *Philodendron immixtum* occurs during the last half of the dry season and the first part of the rainy season, March through May, with post-anthesis inflorescences seen from March through August, and fruits seen in July.

*Philodendron immixtum* ranges from Panama to Colombia (Golfo de Urabá), occurring at or near sea level (to ca. 140 m elevation), primarily along the Atlantic slope in Colón and Panamá Provinces but also in Panamá Province on the Pacific slope along the Río Majé. It occurs in *Tropical wet forest*, *Premontane wet forest*, and in wetter parts of *Tropical moist forest* life zones.

*Philodendron immixtum* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. The species is characterized by its scandent habit; long, slender, mostly light-brown-drying internodes; slightly flattened petioles (about one-half or less as long as the blade); narrowly ovate to ovate-elliptic, green-drying blades, which are usually cordulate at the base; as well as by the solitary

inflorescence with a green spathe, the base white on the blade at anthesis and red within on the tube.

The species has been confused with *P. inconcinnum* Schott (1856) (hence the name "immixtum," meaning "intmixed or mingled with") from Venezuela (Standley, 1944; Croat, 1978). The Schott paintings of *P. inconcinnum* show a single unattached leaf and unattached inflorescences and probably represent the same species as *P. sphaerum* Schott (1860). Schott paintings of *P. sphaerum* show some leaves that are virtually identical to the above-mentioned drawings. *Philodendron sphaerum* differs from *P. immixtum* in having several small inflorescences per axil (rather than one, somewhat larger inflorescence for *P. inconcinnum*). Schott's original description of *P. sphaerum* did not mention an inflorescence, so his drawing of that must have come later than his description. Schott's original description of *P. inconcinnum* cited Venezuela as the type locality, not Panama, and the solid greenish inner spathe surface depicted in his paintings corresponds well to *P. sphaerum* but not to the material in Panama, which has a reddish interior spathe surface.

*Philodendron immixtum* has also been confused with *P. ligulatum*. The latter species differs in having blades that are typically more nearly oblong-elliptic to narrowly obovate-elliptic, frequently broadest above the middle, and drying usually much blackened. In addition, it has petioles that bear a conspicuous purple ring around the petiole at the apex. In contrast, the blades of *P. immixtum* typically dry grayish to brownish green and are broadest frequently below the middle. The species also differs in having petioles that lack the purple ring at the apex.

A number of sterile or immature collections from Nicaragua may represent this species, but if so it is curious that the species has not been collected in Costa Rica (or for that matter, not further west in Panama than Miguel de La Borda in Colón Province).

*Additional specimens examined.* PANAMA. Canal Area: Balboa, Standley 28542 (US); Ft. Randolph, 28616 (US); road between Gatón Locks and Ft. Sherman, ca. 1 mi. E of Ft. Sherman, <25 m, 9°19'N, 79°57'30"W, Croat & Zhu 76282 (CR, GB, MO, PMA, TEX, US); Ft. Sherman, from Pina to 3 mi. NE, Liesner 1382 (MO, US); Barro Colorado Island, Croat 5064 (MO), 10865 (MO, SCZ); Shattuck 851 (F, MO); Wetmore & Woodworth 11 (F, GH); Aviles 89 (F); Fairchild 3079 (US); Parque Nacional Saborana, 140 m, 9°06'20"N, 79°37'20"W, Croat & Zhu 76987 (MO). Colón: 0.25 mi. N of María Chiquita on road to Portobelo, 11360 (F, MO); Río Indio-Miguel de la Borda, along ocean trail, 0 m, 36896 (MO); vic. of San Miguel de la Borda, 9896 (MICH, MO, NY, PMA, SCZ); 2 mi. W of Portobelo, 33574 (BR, DUKE, MO); Palma Bellas-Sal-



ud, near sea level, 36865 (MO); 4.5 km SW of Piña, 0-5 m, *Nee 11713* (MO); 3 km S of Piña, 50 m, *Sullivan 137* (MO); Portobelo, *Croat & Porter 15613* (MO); Nuevo Tonosí-Río Indio, near sea level, *Croat 33545* (MO). **Darién:** Cerro Pirre, 17 km N of El Real, along Río Perisénica, 100 m, 8°01'N, 77°44'W, *Croat & Zhu 77093* (CM, MO); 77162 (CM, MEXU, MO, PMA, US). **Panamá:** Río Maje, W of Bayano Lake, ca. 2 mi. upstream, 30-60 m, *Croat 34592* (MO).

**COLOMBIA.** **Chocó.** Mpio. Acandí, Golfo de Urabá, Bahía Zardí, 0-5 m, 8°23'N, 77°07'W, *Betancur et al. 1238* (HUA, MO).

***Philodendron jacquinii*** Schott, Syn. Aroid. 90. 1856. TYPE: Plate 152 in Jacq., *Select Stirp. Amer. Hist.* 1763 (holotype). Figures 231-232, 237, 241-242.

*Philodendron hoffmannii* Schott, Oesterr. Bot. Z. 8: 178. 1858 (as "*P. hoffmanni*"). TYPE: Costa Rica. *Hoffmann s.n.* (holotype, B destroyed). Schott ic. 2507 (neotype, here designated, W).

*Philodendron desiatum* Schott, Bonplandia 7: 29. 1859. TYPE: Venezuela, *Fendler 1329* (lectotype, here designated, K).

*Philodendron erlansonii* I. M. Johnston, Sargentia 8: 89, t. 13. 1949. TYPE: Panama. Panamá: Perlas Archipelago, San José Island, along road about one-half mi. E of Red Hill, 31 Dec. 1945, *Johnston 967* (holotype, GH).

*Philodendron apocarpum* Matuda, Madroño 10: 171. 1950. TYPE: Mexico. Chiapas: along Río Cintalapa, at Gilguero, 15 km E of Escuintla, Chiapas, ca. 200 m, 10 Aug. 1948, *Matuda 18313* (holotype, MEXU; isotypes, F, MEXU).

Hemiepiphytic vine; stem scandent, green to gray-brown, glossy, setose-pubescent; leaf scars conspicuous, 1 cm long, 1 cm wide; internodes elongate, 6-25 cm long, 0.8-1.5 cm diam., longer than broad, brownish gray, semiglossy, weakly asperous, epidermis paper-thin, sometimes peeling with age, drying gray-green to yellowish green, matte, sometimes almost blackened, larger stems drying yellow-brown and finely ridged; roots few per node, short, ca. 1 mm diam., light brown, sparsely scaly; cataphylls unribbed, semiglossy, 6-10 cm long, light green, translucent, drying cream-colored, deciduous intact; **petioles** 7.5-44.5 cm long, 5-10 mm diam., subterete, medium green, weakly glossy, narrowly and obtusely sulcate adaxially, surface setose-pubescent, often puberulous; **blades** broadly ovate-cordate, thin, chartaceous, acute to shortly acuminate, sometimes mucronate at apex (the acumen sometimes apiculate, 1-3 mm long), cordate at base, (10.5)11.5-39 cm long, 7.5-28 cm wide (0.99-1.8 times longer than wide), (0.6-1.85 times longer than petiole), usually about equal in length to petiole, broadest near point of petiole attachment, upper surface dark green, matte to subvelvety to weakly glossy, lower surface mod-

erately paler, weakly glossy to almost matte, moderately paler; anterior lobe 8.3-27 cm long, 9-28 cm wide (1.6-3.4 times longer than posterior lobes); posterior lobes  $\pm$  rounded, 3-13.5 cm long, 3.4-13.6 cm wide, directed inward at maturity, rounded or obtuse; sinus  $\pm$  narrowly parabolic; midrib broadly convex, concolorous above, broadly convex to round-raised, often setose-pubescent near base, slightly paler below; basal veins 3-6 per side, with 0-1 free to base, 1-2 coalesced 0.5-2.5 cm; posterior rib not naked or briefly naked for 0.5 cm (rarely to 1.5 cm); primary lateral veins 2-3(4-5) per side, departing midrib at a 45-55° angle,  $\pm$  straight to the margins, prominently impressed to weakly convex above, convex to round-raised below; minor veins arising from both the midrib and primary lateral veins; tertiary veins obscurely visible to raised and conspicuous, darker than surface below. **INFLORESCENCES** erect, 1 per axil; peduncle (2)4-15 cm long, (2)3-6(7-13) mm diam., green, dark-striate, glabrous; **spathe** 10-16.5 cm long, (0.96-2.3(2.7-3) times longer than peduncle); spathe blade green outside, red inside; spathe tube inflated bulbous, green outside, 4 cm long, white or green inside; **spadix** sessile; 8-13 cm long, narrowly rounded at apex; pistillate portion 3-3.5 cm long, 3.5 cm long in front, 3 cm long in back, 3.5 cm diam. throughout; staminate portion 9.2 cm long; fertile staminate portion 1.5-1.6 cm diam. at base, 1.1 cm diam. at middle, 1 cm diam. ca. 1 cm from apex; sterile staminate portion narrower than the pistillate portion, 1.2-1.5 cm diam.; pistils 6.5-7.2 mm long, 2 mm diam.; ovary 4-locular, locules ca. 1.3 mm long, ca. 0.6 mm diam., with sub-basal placentation; ovules 2 per locule, contained within transparent, gelatinous matrix (no true envelope), 0.6-1 mm long, as long as or longer than funicle; funicle 0.4 mm long (can be pulled free to base), style similar to style type B; style apex quite rounded; styles usually elongated to 5 mm long, sometimes essentially sessile; stigma hemispheroid to sometimes somewhat globular, lobed, 3.3 mm diam., 1.5 mm high, covering entire style apex; the androecium truncate, prismatic, some oblong, margins irregularly 4-6-sided, 1-1.7 mm long; thecae oblong, 0.5 mm wide,  $\pm$  contiguous; sterile staminate flowers irregularly 4-5-sided, 2.2 mm wide. **INFRACTESCENCE** usually pendent, often from leafless stems; peduncle 7-15 cm long, recurved; spathe falling free; spadix (2.5)5-9(14) cm long, (1.5)2.5-7 cm wide; berries greenish when immature, pale orange to red to reddish orange when mature, irregular, 1 cm long, 5 mm diam.; seeds 4-6 per berry, white, ovoid, 4 mm long, 3-4 mm diam., moderately sticky. **JUVENILE**

plants with petiole margins vaginate-winged almost to one-half their length; blades narrowly ovate-cordate.

Flowering in *Philodendron jacquinii* occurs in August but most post-anthesis collections are from the rainy season, mostly June through September (but also November, January, and March). Immature fruits are known from virtually throughout the year, and mature fruits from March, June, and July. There seems to be little phenological variation in this species.

*Philodendron jacquinii* ranges from Mexico to Panama on both slopes of the Continental Divide to northern Colombia, Venezuela, and the Guianas, as well as western Cuba and the Cayman Islands, from sea level to 1500(2500)m elevation (mostly below 300 m), mostly in *Tropical moist forest* but also in *Tropical dry forest*, *Premontane moist forest*, and *Premontane wet forest* life zones.

This species, the only member of *P.* sect. *Macrogynium*, is distinguished by occurring in moderately low, dry habitats, and by its scandent habit, elongate internodes, setose-pubescent stems, petioles and veins of lower blade surfaces, subterete petioles (about equal the blades in length), and thin, ovate-cordate veiny blades. Especially characteristic are the generally swollen spathe tubes and elongated styles, making the pistillate portion of the spadix much broader than the staminate portion.

*Philodendron jacquinii* may be confused with *P. brevispatum*, another species with scaly petioles and thin blades with prominent posterior lobes. The latter species differs in having ovate-triangular blades with more slender, frequently flaring posterior lobes and a reddish brown stem with simple or branched scales lacking the elongated setae typical of *P. jacquinii*, which has typically greenish stems. In addition, the berries of *P. jacquinii* are pale orange to reddish, and the spathe is reddish within (vs. berries whitish and the spathe green within for *P. brevispatum*).

For a discussion of the long-standing nomenclatural confusion involving *P. hederaceum*, see under the latter name. Although Standley and Steyermark (1958b), in their treatment of Araceae for the *Flora of Guatemala*, had the taxonomy correct in regards to this confusion, their illustration labeled *P. hederaceum* (fig. 58) appears to be a mixture of the two species, with the leaves of *P. hederaceum* and an inflorescence of *P. jacquinii*. The former species differs in lacking the setose-pubescent scales characteristic of *P. jacquinii*, and in having more coriaceous blades as well as normal, short, closely

compacted pistils. The spathe tube of the latter is also not markedly inflated.

A collection from El Salvador (Croat 42075) is unusual in lacking the usual trichomes on the stems.

*Philodendron lundellii* Bartlett ex Lundell (Bartlett, 1937) also corresponds to this species, but the name was invalidly published (without a description or Latin diagnosis).

*Additional specimens examined.* BELIZE. Belize: Western Highway, Mile 35, Croat 24784 (F, MO); Belmopan, Gentry 8423 (MO); 9.5 mi. S of Georgeville, road to Augustine, Croat 23479 (MO); Hummingbird Highway, 7 mi. S of junction with Western Highway, N boundary of Rosling River Est., ca. 110 m, Spellman & Newey 1967 (GH, MO). **Corozal:** Gentle 530 (CM, MICH); Cerro Maya Ruins, Lowry's Bight, Crane 510 (LL), 418 (LL, MO); Gentle 345 (MICH). **Orange Walk:** Honey Camp, Lundell 99 (F, US); 2 mi. N of Orange Walk, Sutton et al. 122 (BM). **Toledo:** Maya Mountains, lower slopes of Richardson Peak, N of junction of Richardson Creek and Bladen Branch, 300-620 m, 88°46'30"W, 16°33'-35'N, Davids & Brant 31955 (B, MO). COSTA RICA. **Alajuela:** Orotina, Valle del Tárcoles, 100 m, Hammel & Grayum 19932 (CR, INB). **Guancaste:** El Mirador, 450 m, Aguilar et al. 1343 (INB, MO). **Heredia:** La Selva Field Station, ca. 100 m, Hammel 10500 (MO); 1.4 km NW of Puerto Viejo, 75 m, Anderson & Mori 25 (CM). **Puntarenas:** Barranca Site, 15 mi. N of Puntarenas, Janzen 10742 (MO); Los Barrancas, along Pan-American Highway, near Miramir turnoff, Gentry 774 (MO); Río Gaseimal, San Luis, Monteverde, 700 m, 10°16'N, 84°49'W, Bello et al. 20 (CR, MO); Monteverde Reserve, 1500 m, ca. 10°17'N, 84°48'W, Croat 61192 (CM, MO); 1300 m, Haber & Bello 7983 (CR, MO); Cantón de Buenos Aires, along Río Ceibo, Ujarrás, 500 m, 9°14'N, 83°18'W, Grayum 10228 (CR, INB, MO, US). **San José:** Mora, Colón-Hacienda El Rodeo, 800 m, 9°55'N, 84°16'W, Hammel 18579 (INB, MO). EL SALVADOR. **Ahuachapán:** San Francisco Menéndez-Tacuba, 0-2 mi. NE of San Francisco Menéndez, 200-450 m, Croat 42075 (MEXU, MO). GUATEMALA. **Without locality:** Heyde 463 (US); Aguilar 309 (F). **Chiquimula:** Río Chiquimula, Santa Bárbara-Petapilla, 4-6 mi. N of Chiquimula, 350-420 m, Standley 30245 (F); Las Mamacas, 16 mi. S of Guatemala City on CA-9, 3800 ft., Dieckhaus et al. 3472 (UMO). **Jalapa:** Jalapa-San Pedro Pinula, 1400-1800 m, Standley 77051 (F). **Jutiapa:** Standley 75385 (F); 75695 (F); 75212 (F); El Barrial, E of Jutiapa, ca. 800 m, Standley 75764 (F); Jutiapa-Las Tunas, NW of Jutiapa, 850-900 m, Standley 76294 (F). **Petén:** 31 km S of Flores, 175 m, Harmon & Dwyer 2797 (MO); La Libertad, Lundell 2551 (MICH); 1 km S on old trail to Mahaquils, Contreras 2912 (LL); Tikal National Park, Lundell 15292 (LL); 16792 (LL); Contreras 343 (LL); Dolores, Contreras 3763 (LL). **Retalhuleu:** Nueva Linda-Chasperico, 120 m, Standley 87774 (F). **Santa Rosa:** SE of Barberena, 1100-1800 m, Standley 77740 (F); Caillapilla, 900 m, Standley 78087 (F); La Sepultura region, W of Chiquimulilla, 220 m, Standley 79331 (F); Chiapas, 3500 pp, Heyde & Lux 3867 (US); Yaxoquimela, floodplain of Río Santa Cruz (upper tributary of Río Usamacinta), 560 m, 16°58'N, 91°47'W, Rawlin 93 (CM). **Zacapa:** trail between Río Honda and waterfall, 250-400 m, Standley 29396 (F); Quebrada

Alejandria, summit of Sierra de Las Minas, Finca Alejandria, 2500 m, *Steyermark* 30856 (F); Mpio. Ocozacoatlán, E edge of Selva del Ocote, 900 m, *Breedlove* 70760 (CAS, MO). HONDURAS. **Colón:** Trujillo, Capuchin Monkey Site West, *Saunders* 523 (MO); Bonito Oriental-Limón road, ca. 50 m SW of Río Piedra Blanca, 40 m, *Rivas* 46'30"N, 85°41'W, *Evans* 1086 (MO). **Copán:** 10 mi. W of Copán, road to La Entrada, 700 m, *Croat* 42517 (MO). **Morazán:** Río Yeguaré drainage, ca. 14°N, 87°W, *Williams* 15908 (F); Río Yeguaré, E of El Zamarano, ca. 750 m, *Standley* 15448 (F); Río de la Orilla region, SE of El Zamarano, 900-950 m, *Standley* 22404 (F). **Olancho:** Juticalpa, 380-480 m, *Standley* 17524 (F); Río Olancho, Gualaco-San Bonito Oriental, 7.4 mi. NE of San Esteban, 540 m, 15°20'N, 85°42'W, *Croat* & *Hannon* 64363 (MO). **Yoro:** Río Aguán, Coyoles, *Yuncker* et al. 8663 (F, GH, MICH, MO, NY, US); Río Guaymón, La Cieba-San Pedro Sula, SW of Tela, near border of Yoro and Atlántida, 50 m, 15°30'N, 87°43'W, *Croat* & *Hannon* 64666 (MO). MEXICO. *Hahn* s.n. (P). **Campeche:** Yucatán Peninsula, Champoton, *Steere* 1742 (CM); Maskall, *Gentle* 1186 (CM, MICH, NY); Tuxtepa, *Lundell* 1074 (F). **Chiapas:** Arriaga-Las Cruces, 5.7 mi., ca. 160 m, *Moore* & *Bunting* 8923 (HB); 51 mi. NE of Chiapas-Oaxaca border on Hwy. 190 to Tuxtla Gutiérrez, 660 m, 16°47'N, 93°18'W, *Thompson* et al. 432 (CM, MO); 5 mi. N of Ocozacoatlán, along road to Apatzaco, 1000 m, *Croat* 40544 (MO); Río Cintalapa, Aguas Calientes, Escuintla, *Matuda* 18312 (NY); Esperanza, Escuintla, *Matuda* 18314 (MEXU, MO); Mpio. Berriozábal, 5 km E of Berriozábal, along Mexican Hwy. 190, 800 m, *Breedlove* & *Thorne* 30412 (MO); Mpio. Ocosingo, Río Usamacinta, 3 km. S of Frontera Corozal, 120 m, *Martínez* S. 7309 (MO); Mpio. Ocozacoatlán, km 103-104 Hwy. 190, 15.7 mi. E of Cintalapa, 3050 ft., *Baswell* & *Kinnoch* 527 (US); Río Grijalva, 10 km S of Hwy. 190 on road to Acala, at Nandaburri, 1600 ft., *Laughlin* 2811 (F). **Guerrero:** Acapulco, *Palmer* 462 (US). **Oaxaca:** Tuxtepec-Matías Romero, <100 m, *Croat* 78724 (CHIP, MO); 7.5 mi. SW of Tuxtepec on road to Valle Nacional, *Moore* & *Bunting* 8908 (HB); 14 mi. N of Puerto Escondido on Rte. 131 to Oaxaca, 300 m, 16°13'-47"N, 97°5'-8"W, *Thompson* et al. 424 (CM, MO); 5 mi. N of Palmar, 54 mi. N of junction of Highways 185 and 190, along Hwy. 185, 65 m, *Croat* 40002 (MO); Parque Nacional de Laguna Temazcal, Temazcal, 150 m, 18°25'N, 96°25'W, *Hammel* & *Metcalfe* 15454 (MO). **Quintana Roo:** Mpio. Chetumal, 6.5-7 km N of Tomás Garrido on the road joining Hwy. 186 W of Nicolás Bravo, 150 m, 18°6'N, 89°3'W, *Saunders* et al. 9950 (MO); 1 km E of Chanca Veracruz, *E. Cabrera* & *H. Cabrera* 4981 (MEXU). **San Luis Potosí:** Tamazunchale, *Edwards* 549 (F, MO). **Veracruz:** Córdoba-Veracruz, Ejido San José de Gracia below Penuela, *Moore* & *Bunting* 8877 (HB); San Lorenzo Tenochtitlán, 22-75 m, *Wing* 50 (GH). **Yucatán:** Schott 489 (BM); Mérida, *Gaumer* 489 (BM, F); Izamal, *Gaumer* 23174 (F) *Gaumer* 1422 (F); *Greenman* 376 (F); *Gaumer* 23823 (F, US). NICARAGUA. **Boaco:** Las Pitas, Camoapa, 400 m, 12°28'N, 85°35'W, *Moreno* 10643 (MO); Río Las Cañas, along Hwy. 33, ca. 3.1 km N of Hwy. 35 intersection, ca. 275 m, 12°38'N, 85°33'W, *Stevens* 5838 (MO). **Chontales:** Route 7, 5 km SE of Juticalpa, *Moore* 1622 (MO). **Granada:** Laguna de Apoyo, 110-180 m, 11°53'N, 86°01'W, *Moreno* 11178 (MO). **Masaya:** Parque Nacional Volcán Masaya, N slope of Volcán Santiago, ca. 375 m, 11°59'N, 86°10'W, *Stevens* 5267 (BM, MO); summit of Volcán Masaya, 500 m, *Stevens* 2950 (MO); Piedra Quemada, 2 km E of Volcán Masaya, 250 m, *Stevens* 4606

(BM, MO). **Matagalpa:** Ranchería, 11 km NE de Muy Muy, ca. 280 m, 12°46'N, 85°31'W, *Moreno* 24483 (AAU, MO); Río Yacica, Tuma, ca. 28 km NE Matagalpa, El Diamante, ca. 350-400 m, 13°04'N, 85°46'W, *Guzmán* et al. 915 (MO). **Río San Juan:** Boca de Sábalo, "La Toboba," 70-90 m, 11°03'-04'N, 84°28'-29'W, *Robledo* 1833 (US). **Rivas:** Tola-Las Salinas, El Coyol, ca. 2.3 km beyond entrance of Hda. Miramar, ca. 30-40 m, 11°23'N, 85°58'W, *Stevens* 9749 (BM, MO). **Rivas-Carazo-Granada:** Río La Pita-Río Escalante, ca. 20 m, 12°34'N, 86°08'W, *Stevens* 9718 (BM, MO). **Zelaya:** SW of Puerto Cabezas, 0-2 m, 14°01'N, 83°24'W, *Stevens* 7880 (MO); Mpio. Siuna, *Wany*, *Ortiz* 59 (MO). PANAMA. **Canal Area:** Balboa, *Standley* 29256 (US); road S-10 N of Escobal, *Croat* 12458 (MO, NY, U); Rodman Marine Base, Rodman Naval Ammunition Supply Depot, W of Balboa, 70-80 m, 8°58'N, 79°36'25"W, *Howard* 147 (MO); road to Cerro Pelado Radar Station, 0.5 km NW of Gamboa, 75-150 m, *See* 7760 (MO, RSA); Curundú, Parque Metropolitano, 8°58'N, 79°32'55"W, *Croat* & *Zhu* 76202 (MO); Gamboa, *Croat* 74755 (F, MO); Geologic Test Site N of Paraíso, *Croat* 12977 (MO); near Summit Hills Golf Course, *Croat* 10956 (MO); Gaillard Hwy., vic. Summit Golf Club, 40 m, 9°03'N, 79°37'W, *Croat* 69835 (AAU, MO, US); Radar Station Road, 1 mi. N of Summit Gardens, *Croat* 9080 (MO); Barro Colorado Island, *Shattuck* 115 (F, GH); *Zetek* 4675 (MO, US); Wheeler 5, *Shattuck* 215 (MO); laboratory clearing, *Croat* 10261 (MO); 9259 (MO); Colorado Point, *Croat* 6138 (MO); Vista Alegre, *Zetek* 5576 (MO); Chagres River, vic. of Juan Mina, Flat Rock, *Bartlett* & *Lasser* 16839 (MICH, MO); Fort Clayton, vic. of end of C-16 road, *Blum* 2243 (MO, SCZ); Ft. Sherman, *Standley* 31020 (US); Madden Dam Area, Boy Scout Road, *Porter* et al. 4062 (MO); Madden Forest, Las Cruces Trail, *Croat* 11878 (MO); 140 m, 9°06'20"N, 79°37'20"W, *Croat* & *Zhu* 77072 (MO); Pipeline Road, 2-4 mi. N of Gamboa, ca. 100 m, *Gentry* 6543 (MO); Río Cocolí, road K-9, *Stern* et al. 348 (GH, MO, US); Paraíso, *Tyson* & *Lazar* 6194 (F). **Coelá:** Penonomé and vic., 50-1000 ft., *Williams* 381 (NY). **Darién:** Cerro Pirre region, El Real, *Croat* & *Porter* 15460 (MO); Santa Fe region, Univ. of Georgia Cuipe Forest Site 02, 15 m, *Duke* 14258 (F, MO). **Herrera:** Orú, *Edinger* 1090 (MO, US). **Los Santos:** ca. 5 m S of Las Tablas, *Burch* et al. 1236 (MO, UC); Pocerí, *Dwyer* 11894 (MO); Las Tablas, *Dwyer* 1189 (MO). **Panamá:** along road to Bique, 5 km SW of Arriaján, 20-40 m, *See* 7699 (MO, US); Panamá Viejo, *Rose* 18505 (NY, US); road K-15 near Huile, vic. Gatún Lake, *Smith* et al. 3277 (F, US); El Llano-Chepo, *Gentry* & *Tyson* 1727 (MO, SCZ); Tocumen, *Dwyer* 4220 (MO); vic. Macambo, *Croat* 14911 (MO); Cerro Campana, *Croat* 12018 (MO, SCZ); Penonomé and vic., 50-1000 ft., *Williams* 381 (NY); San José Island, Pearl Archipelago, *Erlanson* 234 (US); 402 (GH, US); *Johnston* 974 (GH, MO), 918 (GH), 12 (GH), 960 (GH).

**Philodendron jefense** *Croat*, sp. nov. TYPE: Panama. Panamá: Cerro Jefe, along road short of summit, 550-800 m, 9°15'04"N, 79°30'04"W, *McPherson* 10038 (holotype, MO-3475849; isotypes, K, PMA, US). Figures 238, 239.

Planta hemiepiphytica; internodia brevia, in sicco usque 3 cm diam.; cataphylla 20-24 cm longa, acute 2-costata, mox decomposita et persistentia ut fibrae pallide brunneae; petiolus teres, 38-76 cm longus, 4 mm

diam.; lamina late ovata, 39–54 cm longa, (22.5)34–49 cm lata, cordata basi, in sicco canoviridis; inflorescentia 1; pedunculus 10–13 cm longus, 4–5 mm diam.; spathe 9.5–14 cm longa, omnino viridis, in sicco cum magnis maculis albis; pistilla (6)7–8-locularia; locules cum 6 seminibus; baccae albae.

Hemiepiphytic; stem appressed-climbing, internodes short, to 3 cm diam.; roots moderately few per node, drying to 3 mm diam., sharply ridged, brown; cataphylls 20–24 cm long, sharply 2-ribbed, persisting, promptly weathering to light brown, semi-organized fibers with small, thin fragments of epidermis persisting; **petioles** 38–76 cm long, 4 mm diam., terete, drying black; **blades** broadly ovate, short acuminate at base, cordate at base, 39–54 cm long, (22.5)34–49 cm wide (0.8–1.4 times longer than wide), (0.5–0.6 times the petiole length), about one-half as long as the petiole, margins sinuate, upper surface glossy, drying gray-green and semiglossy, lower surface glossy, drying minutely granular; anterior lobe 22–25 cm long, 24.5–29.7 cm wide (1.8–2 times longer than posterior lobes); posterior lobes 12–14 cm long, 12.2–13.2 cm wide, obtuse; midrib drying darker than surface above; basal veins 5 pairs per side, first and second free to base, the remainder coalesced 3–6 cm; posterior rib naked to 3–3.5 cm long; primary lateral veins 5–6 per side, departing midrib at a 35–45° angle; minor veins frequently branched, arising from both the midrib and primary lateral veins, minute pustules visible between veins on both surfaces; "cross-veins" many. INFLORESCENCES 1 per axil; peduncle 10–13 cm long, 4–5 mm diam., terete, black-drying; **spathe** 9.5–14 cm long (ca. 1 time longer than peduncle), green throughout, drying with large white flecks; **spathe blade** 6 cm long; **spathe tube** 6 cm long; **pistillate portion** 5 cm long; **spadix** to 11 cm long; staminate portion 5.5–6 cm long, 7–10 mm diam.; fertile staminate portion to 1 cm diam.; sterile staminate portion 7 mm diam.; **pistils** 3.9 mm long; **ovary** (6)7–8-locular, locules 3.2 mm long, 0.4–0.5 mm diam., with axile placentation; **ovules** 2-seriate; **funicle** 0.3–0.4 mm long, adnate to lower part of partition, style similar to style type D; style apex flat; style boss small. INFRUCTESCENCE with berries white (immature); seeds 6 per locule, 1–1.1 mm long, 0.2–0.3 mm diam., translucent.

Flowering in *Philodendron jefense* is documented by a single collection with immature fruits collected in December. Flowering apparently occurs during the wet season.

*Philodendron jefense* is endemic to Panama on Cerro Jefe in Premontane rain forest at about 800 m elevation.

*Philodendron jefense* is a member of *P.* sect. *Colostigma* subsect. *Macrobeltium* ser. *Reticulata*. This species is distinguished by its appressed-climbing habit, short, thick internodes, sharply 2-ribbed cataphylls promptly weathering to light brown, semi-organized fibers with small, thin fragments of epidermis persisting, terete petioles (about twice as long as the blades), broadly ovate, gray-green-drying blades, and solitary long-pedunculate inflorescences with the spathes green outside.

*Philodendron jefense* is probably closest to *P. lazorii* Croat, which shares broadly ovate, gray-green-drying leaf blades and semi-organized, brownish masses of persistent cataphyll fibers. The latter species differs in having matte-drying blades with the lower surface smooth and minutely brownish- or whitish-speckled, with the minor veins drying darker than the surface and weakly puckered without any sign of branching or cross-veins. In contrast, *P. jefense* has blades semiglossy on drying with the lower surface sparsely granular with frequently branched minor veins and with cross-veins not drying darker than the surface. The upper surface lacks any signs of raphide cells.

The species name refers to the type locality, Cerro Jefe, which has proven to be one of the most endemic-rich areas for its size of any site in Panama.

***Philodendron jodavisanum*** G. S. Bunting, Genes Herb. 9: 337. 1965. TYPE: Mexico. Chiapas: Cerro de Madrugal, along Teapa-Tacotalpa (Tabasco) Highway, at km 4, W of hwy., on rock face of steep slopes, *Bunting 1526* (holotype, US). Figures 243, 245–248, 253.

Usually hemiepiphytic, terrestrial; stem appressed-climbing, gray-green to brown or green, sap sweet-scented, drying blackened, leaf scars conspicuous, 2 cm long, 1.7 cm wide; internodes weakly flattened on one side, 2-ribbed, weakly glossy, to 8 cm long, but surely longer than wide and 2.5–3 cm diam. at apex, dark to medium green, soon gray, epidermis brown, flaking, fissured sometimes; roots whitish, few per node; cataphylls 10–20 cm long, bluntly or sharply 2-ribbed or unribbed, sharply D-shaped, somewhat spongy, light to medium green or reddish to brownish, sometimes streaked pinkish, semiglossy, drying brown, persisting semi-intact as pale fibers at upper nodes, disorganized below; **petioles** (18)29–91 cm long, (2)3–7(8–12) mm diam., sometimes terete, usually D- or U-shaped, spongy or firm, dark green, flattened or somewhat sulcate to convex, often with medial rib, margins raised adaxially, rounded abaxially, surface semiglossy, minutely and short-lineate; **blades**

triangular-ovate,  $\pm$  subcoriaceous, moderately bicolorous, long acuminate at apex, deeply cordate at base, 25–74 cm long, 18–34 cm wide (1.1–2.8 times longer than wide), (0.5–2 times longer than petiole), broadest  $\pm$  near the middle, upper surface dark green, drying blackish, semiglossy, lower surface pale green to yellow-greenish, drying blackish, glossy to semiglossy or matte; anterior lobe 18–58 cm long, 10.1–40.4 cm wide (1.7–4.3 times longer than posterior lobes); posterior lobes 8–17 cm long, (3.7)5–17 cm wide, obtuse to rounded; midrib broadly sunken to broadly convex or flat, dark green above, raised to narrowly convex, thicker than broad, concolorous or darker than surface below; basal veins 5–10 per side, in part coalesced to 4 cm, barely naked or naked to 2 cm; primary lateral veins 7–11 per side, departing midrib at a 50–60° angle,  $\pm$  straight to the margins, obtusely to weakly or narrowly sunken and sometimes weakly quilted above, convex and darker than surface below; interprimary veins almost as prominent as primary lateral veins; minor veins in part discontinuous and darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, 2–6 per axil; peduncle 3–13 cm long (to 20 cm long in South America), 2–6 mm diam., pale green to whitish, with raised, prominent white striations; spathe semiglossy, 6–14.5 cm long (0.9–3 times longer than peduncle), sometimes greenish with short white lineations throughout; spathe blade white to pale green (at anthesis), reddish (post-anthesis) (B & K red-purple 7.5/7), short-lineate outside (opening 4.6 cm long, 3.6 cm wide), very pale green to white, matte, few, sparse, dark-lineate inside; spathe tube semiglossy and greenish to greenish brown or reddish (maroon or reddish at anthesis) outside, green to white inside; spadix weakly exerted from the spathe, 7.6–9.2 cm long, constricted near base of fertile staminate portion; pistillate portion pale green or whitish, cylindrical, 1.4–2.4 cm long, 1 cm diam. at apex, 1.1 cm diam. at middle; staminate portion 5.3–7.6 cm long; fertile staminate portion creamy white, cylindrical to weakly tapered, 0.85–1.5 cm diam. at base, 0.7–1.3 cm diam. at middle, 6–10 mm diam. ca. 1 cm from apex, broadest at the base, narrower than the pistillate portion, narrower than the sterile portion; sterile staminate portion broader than the apical area of the pistillate portion, gray-white, 1.1–1.5 cm diam.; pistils 1.8–2.5 mm long, 1–1.4 mm diam.; ovary 4–6(5–7)-locular; locules 1.3–1.7 mm long, 0.5–0.7 mm diam., with axile placentation; ovules 18–26(23–29) per locule, 2-seriate, 0.2–0.4 mm long, longer than funicle; funicle 0.1–0.2 mm long, adnate to lower part of partition, style 0.2–0.3 mm long, 1–1.4 mm diam., similar to style type B; style apex flat to slightly rounded; stigma subdiscoid, brushlike, unlobed, 1–

1.3 mm diam., 0.2–0.6 mm high, covering entire style apex; the androecium truncate, margins 4–6-sided; thecae oblong to obovate, 0.3–0.5 mm wide,  $\pm$  parallel to one another, contiguous; sterile staminate flowers blunt, 1.6–1.8 mm long, 1.4–1.5 mm wide. INFLORESCENCE with berries white (mature), 6 mm long, 2.6 mm diam.; seeds 14–16(24–28) per locule, 1.3–1.7 mm long, 0.2–0.3 mm diam. JUVENILE plants with internodes 8 cm long, 5 mm diam.; petioles sharply U-shaped in cross section, usually lacking medial rib adaxially, margins erect.

Flowering in *Philodendron jodavisanium* occurs in the late dry season and throughout much of the rainy season (April through October) in Panama. In Costa Rica it has been collected in flower in February, and post-anthesis collections have been made in January. Immature fruits have been collected August through November.

*Philodendron jodavisanium* ranges from southern Mexico to Colombia, Ecuador, and Venezuela (Mérida) from near sea level to 1500 m elevation. It occurs in *Tropical wet forest* and *Premontane wet forest* in Panama and Costa Rica but also in *Tropical moist forest* in some parts of Central America.

*Philodendron jodavisanium* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is characterized by its short internodes, sharply 2-ribbed cataphylls persisting as pale fibers, petioles flattened or somewhat sulcate above and usually with a medial rib, usually triangular-ovate, blackish-drying blades with many impressed primary lateral veins, and spathes usually greenish on the tube and white on the blade.

*Philodendron jodavisanium* is quite variable throughout its range, and the type (being one of the most northerly collections) differs from most in having more elliptic blades.

*Philodendron jodavisanium* is most easily confused with *P. grandipes*, which differs in comprising terrestrial plants with broadly ovate leaf blades. It has also been confused with *P. panamense* K. Krause, which differs in having subterete petioles, promptly deciduous cataphylls, and longer peduncles often bent beneath the spathe (see *P. panamense* for additional details).

Similar collections from South America, especially Croat 57441 from Bahía Solano, Chocó Department, Colombia, differ in sometimes having distinct "cross-veins." Collections from Pichincha Province, Ecuador (Croat 55737 and Grayum & Zamora 9431), differ in having peduncles 16–20 cm long.

Two Darién collections (*Hammel et al.* 16252 and *McPherson* 11517) are unusual in having leaf blades with somewhat flaring posterior lobes and markedly

concave margins. However, the leaves on the second sheet of the McPherson collection are typically shaped with a convex blade margin.

*Additional specimens examined.* BELIZE: Toledo: Blue Creek, Whiteford 3234 (BM). COSTA RICA. **Alajuela:** Los Esfrazos, Buena Vista de San Carlos, *Barquero* 10 (UC); Finca Los Esfrazos, ca. 11 mi. NW of Zarco, ca. 850 m, *Croat* 43583 (MO); Reserva Biológica Monteverde, along Río Peñas Blancas, 820 m, 10°21'N, 84°40'W, *Bello & Haber* 2881 (INB, MO); Río Zapote, 5 km S of Canalete, along new road to Upala, 100–200 m, 10°48'N, 85°02'W, *Burger & Baker* 9961 (F, MO, SEL); Cañas–Upala, 4 km NNE of Bijagua, ca. 400 m, *Croat* 36279 (MO); San Ramón, *Bittner & G. Herrera* 2134 (CR). **Cartago:** Moravia–Quebrada Platamillo, Moravia, 3–5 km from Finca Racine, 1200–1300 m, *Croat* 36648 (MO); along Camino Raíz de Hule, SE of Platamillo, 1200–1400 m, *Croat* 36756 (MO); Jicotea Valley, Jicotea–Río Pacuare, *Croat* 36539 (MO). **Guanacaste:** slopes of Miravalles, above Bijagua, ca. 1500 m, *Gómez* 19048 (MO). **Heredia:** 11 km E of Cariblanco, 1060 m, *Loiselle* 255 (MO); Paracito–Río Claro, Bajo La Honduras, 1100–1400 m, *Croat* 44516 (MO); Parque Nacional Brusillo Carrillo, 1000 m, 10°16'38"N, 84°04'57"W, *Boyle* 1285 (MO); Zona Protectora "La Selva," along trail from main road across Quebrada Cantarrana to Río Guacimo, 300 m, *Grayson & Schatz* 3216 (DUKE); La Selva Field Station, ca. 100 m, *Croat* 44300 (MO); 100–150 m, 44320 (MO); *Grayson* 1915 (MO); *Jiménez* V (MO); *Murrell* 1710 (DUKE); Río Frío, W of Finca Zona Nueva, ca. 110 m, 10°18'N, 83°53'W, *Grayson* 3562 (MO); Río Sucio, near Puerto Viejo, 2 m, *Croat* 35703 (MO). **Limón:** Manzanillo de Talamanca, ca. 5 m, 9°38'N, 82°39'W, *Grayson & Burton* 4340 (MO); ca. 11 mi. S of Siquirres, 650 m, *Croat* 43329 (MO); Cerro Coronel, 10–40 m, 10°40'N, 83°40'W, *Stevens* 24346 (MO); Río Sixaola, ca. 0.5 mi. SW of Bambú, ca. 3 mi. NE of Bratsi, ca. 15 m, *Croat* 43254 (MO). **Puntarenas:** ca. 1 km S and 2 km W of Cañasas, 60 m, 8°34'N, 83°25'W, *Croat & Grayson* 59807 (CM, CR, K, MO); Osa Peninsula, 200 m, *Aguilar* 2201 (CR, INB); Golfo Dulce Forest Reserve, Rancho Quemado, 200 m, 8°42'N, 83°33'W, *Hammel & Robles* 16834 (CR, MO); Piedras Blancas, 3.7 mi. W of Pan-American Highway, 90–105 m, 8°46'N, 83°18'W, *Croat* 67696 (CR, MO); 67693 (CR, F, MO); *Golfito, G. Herrera & Rivera* 7042 (CR). **San José:** above Río Sucio, 5–6000 m, *Pennington et al.* 11529 (K); San Isidro del General–Dominical, SW of San Isidro, 4.8 mi. from Río Pacuare, 1000 m, *Croat* 35253 (CR, F, MO); 9 mi. SW of Río Pacuare, 680 m, *Croat* 35344 (MO); Canara Reserve, SW part of Montañas Jamaica, ca. 2.5 km NE of Bijagua de Turrualeses, 460–520 m, 9°45'N, 84°33'30"W, *Grayson et al.* 5488 (MO); vic. of San Isidro El General, 900 m, *Molina et al.* 18063 (EAP, F, GH, NY, UC, US); Pariscal, Z.P. La Cangreja, 800 m, *Morales* 2020 (CR, INB); Tarrazú, vic. Homigüero, 1100–1200 m, *Croat* 78962 (INB, MO). GUATEMALA. **Alta Verapaz:** Tacurú–El Estor, 4–9 mi. up road to Oxec, gravel road N off Hwy. 7E, ca. 6 km NE of Parán, 500–800 m, *Croat* 41607 (MO); 41698 (K, MO, NY). **Izabal:** 12 km NW of El Estor, 650 m, *Martínez & Stevens* 23305 (MO). HONDURAS. **Atlántida:** Lancetilla Botanical Gardens, 2 mi. WSW of Tela, 10–150 m, *Croat* 42640 (K, MO); 70–90 m, 15°44'N, 87°27'W, *Croat & Hannon* 64612 (CM, CR, EAP, G, MEXU, MO); Lancetilla, 100 ft., *Yuncker* 5008 (F, MICH, MO, NY). MEXICO. **Tlaxcala:** Mpio. Teapa, 7 km SE of Teapa on road to Tacotalpa, Rancho San Eneas, 70 m, 17°35'N, 92°50'W, *Hammel & Merello* 15516 (MO). NICARAGUA. **Chontales:** 4 km NNW of Caapa, Cerro Oluma, 700–775 m, 12°18'N,

85°23'30"W, *Neer* 28413 (MO, NY). **Río San Juan:** Coto Chontaleño, 20 km NE of El Castillo, 200 m, *Neill & Vincelli* 3623 (MO); Río Indio, 5 hours upriver from San Juan del Norte, 11°07'N, 83°50'–52'W, *Risener* 247 (MO). **Zelaya:** ca. 6 km upriver from Barra de Punta Gorda, 8–10 m, 11°30'N, 83°49'W, *Stevens* 20756-a (MO); Mpio. Siuna, Santa Rosa, *Ortiz* 67 (MO). PANAMA. **Bocas del Toro:** Escudo de Veraguas Island, N coast, 5 m, 9°05'N, 81°35'W, McPherson 11412 (MO, US); Fortuna Dam area, road along Continental Divide, 2.8 mi. S of Continental Divide, 850–950 m, 8°45'N, 82°15'W, *McPherson* 9676 (MO, US); Chiriquí Grande–Fortuna, 1.2 mi. N of Divide, 5.3 mi. N of bridge over Fortuna Dam, 910 m, 8°44'N, 82°17'W, *Croat* 60450 (MO, PMA, RSA); along Divide, 1.1 mi. W of highway, 1200 m, 8°44'N, 82°17'W, *Croat* 60368 (MO); E of camp Corriente Grande (IRHE), 200 m, *Correa et al.* 3697 (PMA). **Chiriquí:** Río Chiriquí Valley, vic. of Fortuna Dam, 1100–1200 m, 8°45'N, 82°18'W, *Croat* 66587 (F, IBE, M, MEXU, MO, US); 1000–1200 m, 8°45'N, 82°15'W, *Correa et al.* 2686 (MO, PMA); 4.5–5 km N of dam over Fortuna Lake, 1100–1135 m, 8°43'N, 82°17'W, *Croat & Grayson* 60080 (CM, MO); Gualaca–Fortuna, 10 mi. NW of Los Planes de Hornito, 1260 m, 8°45'N, 82°17'W, *Croat* 50074 (MO); Gualaca–Chiriquí Grande, Río Hornito, ca. 0.5 km S of Centro de Científicos, 1010–1130 m, 8°44'N, 82°13'30"W, *Croat* 67922 (MO); 8°45'N, 82°18'W, *Croat & Zhu* 76376 (MO, SEL); 1100–1200 m, 8°45'N, 82°15'W, *Thompson* 5026 (CM); Río Chiriquí, beyond Gualaca, 10.8 mi. beyond Los Planes de Hornito, ca. 1400 m, *Croat* 48721 (MO). **Coelá:** Santa Rita Ridge Road, 4–6 km from Transisthmian Highway, 150–200 m, *Croat* 34280 (MO); Mile 6.5, 370 m, 9°21'15"N, 79°44'W, *Croat & Zhu* 76960 (MO); El Copé region, near Continental Divide, ca. Mile 1.5, ca. 900 m, *Croat* 44572 (MO); ca. Mile 5.6, 800 m, *Croat* 75068 (MO); La Pintada–El Copé, 3000 ft., *Hammel* 2633 (MO); El Valle region, N of El Valle de Antón, La Mesa, 900–1000 m, 8°40'N, 80°10'W, *Hamilton et al.* 4111 (INB, MO); 800–900 m, 8°38'N, 80°09'W, *Croat* 67122 (CAS, CM, COL, HNML, L, MEXU, MO, PMA); 860–900 m, 37396 (MO); 67211 (AAU, DUKE, MO, NY, PMA, TEX); 67213 (AAU, CM, EAP, MO, PMA, TEX, USCG); 67121 (F, HNML, MO); 67129 (AAU, CAS, CM, COL, CR, F, HUA, K, L, MEXU, MO, NY, P, PMA, SEL, TEX); 775 m, 8°36'N, 80°07'W, 74793 (KUN, MO); 860 m, 8°37'N, 80°08'W, *Croat & Zhu* 76742 (MO, SEL); 76710 (MO, SEL). **Colón:** Río Guanche, ca. 3–5 mi. inland, 10–100 m, *Croat* 26175 (MO). **Darién:** Cerro Pirre region, Alturas de Nique region, S of El Real, near Cana mine, 650–800 m, 7°45'N, 77°40'W, *McPherson* 11526 (MO); 11517 (B, MO); Cano gold mine, 480 m, *Croat* 38003 (MO); 38039 (MO); 600–100 m, *Croat* 37743 (MO); Parque Nacional Darién, N of Taracuna, ca. 18 km E of Pacuro, 600–800 m, 8°05'N, 77°16'W, *Hammel et al.* 16481 (MO); ca. 17 km E of Pacuro, 850 m, 8°03.5'N, 77°17'W, 16252 (MO). **Panamá:** El Llano–Carti road, 10 m from highway, near El Llano, 330 m, *Croat* 33779 (F, MO, NY); Mile 5–6, 350–375 m, 34771 (F, MO); Mile 6.8, 350 m, 49128 (AAU, CAS, CM, MEXU, MO); Mile 10.1, 325–350 m, 67366 (MO); Río Maje–Quebrada Brava, ca. 60 m, *Croat* 34657 (F, MO); Cerro Campana, *Croat* 25252 (MO); 25204 (MO); along trail to summit, 780–875 m, *Croat* 17207 (F, MO, NY); at end of road beyond Su Lin Hotel, *Croat* 14227 (MO); 800 m, 8°41'N, 79°56'W, 74772 (CM, MO); summit and upper trail, *Witherspoon & Witherspoon* 8462 (MO); ca. 1 mi. from highway, 450 m, *Croat* 35950 (MO, PMA); Cerro Jefe region, Campo Tres, ca. 700 m, *Croat* 27064 (F, MO); 700–750 m, 9°15'N, 79°25'W, *Croat* 68687 (IBE, MO, TEX). **San Blas:** El Llano–Carti, 23–29 km from Pan-American Highway, 300–400 m, 9°22'N, 78°69'W,

*Knapp 1839* (MO, PMA); El Llano-Carri Road, vicinity Naguandí, 300–350 m, 9°15'N, 79°W, *Croat 69279* (DUKE, MO); 300 m, 9°20'N, 79°W, *Croat & Zhu 76542* (MO); 350 m, 9°20'N, 79°W, *Croat & Zhu 77011* (MO); 450 m, 9°18'N, 79°59'W, *Croat 75121* (MO). **Veraguas:** Río Concepción-Río Barrera, 300–600 ft., *Hammel 5252* (MO); Santa Fe region, Alto Piedra-Calovehora, 0.5 mi. N of Escuela Alto Piedra, Parque Nacional Cerro Tute, 800–1030 m, *Croat & Zhu 76914* (MO); 5–8 km from school, 730–770 m, *Croat 25967* (MO); Río Primero Braso, 5 mi. NW of Santa Fe, 700–1200 m, *Croat 23114* (MO); 15.6 km NW of Santa Fe, 450–550 m, *Croat 27639* (MO); 0.6 mi. beyond Escuela Agrícola Alto Piedra, 730 m, *Croat & Folsom 33997A* (MO); 34062 (MO); ca. 1200 m, 8°32'N, 81°07'W, *Hamilton et al. 1280* (MO); Escuela Agrícola Alto Piedra, *Mori & Kallunki 2529* (MO); Santa Fe-Río San Luis, past Escuela Agrícola Alto Piedra, Río Segundo Braso, 480 m, 8°33'N, 81°08'W, *Croat 66901* (CM, MO).

***Philodendron knappiae*** Croat, sp. nov. TYPE: Panama. Chiriquí: Cerro Hornito, in elfin forest on ridges and summit, approached from Los Planes de Hornito, 2100 m, 8°42'N, 82°06'W, 14 Mar. 1982, *Knapp, Kress & Hammel 4219* (holotype, MO-3043611). Figures 240, 244, 249, 254.

Planta terrestris aut hemiepiphytica; internodia 0.5–7 cm longa, 1–1.7 cm diam., in sicco brunnea; cataphylla 10–18 cm longa, acute 2-costata, sulcata adaxialiter, viridia, decidua; petiolus obtuse D-formatus, 14–19 cm longus, in sicco 2–7 mm diam.; lamina ovato-triangularis, cordata basi, 16–23.5 cm longa, 7.5–11.5 cm lata; inflorescentia 1; pedunculus 2–3 cm longus, 5–6 mm diam.; spathe 7–8 cm longa, estus viridis, intus subrosea; pistilla 4-locularia; loculi 4-ovulati.

Terrestrial or hemiepiphytic; stems usually erect, internodes short, somewhat flattened on one side with marginal ribs, moderately glossy, 0.5–7 cm long, 1–1.7 cm diam., longer than broad below, short near apex, medium green, epidermis smooth, light brown; cataphylls 10–18 cm long, sharply 2-ribbed, green, sulcate adaxially, margins weakly raised abaxially, deciduous; **petioles** 14–19 cm long, 2–7 mm diam., bluntly D-shaped, green, tinged reddish; **blades** ovate-triangular, subcoriaceous, bicolorous, long-apiculate at apex, cordate at base, 16–23.5 cm long, 7.5–11.5 cm wide (1.8–3 times longer than wide), (1.1–1.4 times longer than petiole), about equal in length to petiole, upper surface semiglossy, drying dark brown, lower surface slightly paler, drying dark yellow-brown or dark yellow-green; anterior lobe 12–18.5 cm long, 10.6–13 cm wide (2.2–3.3 times longer than posterior lobes); posterior lobes 5–8 cm long, 5–6.5 cm wide (1.4–1.8 times longer than wide), narrowly to obtusely rounded; sinus hippocrepiform, 3–5.5 cm deep; midrib broadly convex, paler than surface above, convex, reddish or paler than surface below; basal veins 3(5) per side, with 0–1 free to base, 2–3 coalesced 1–1.5 cm; posterior rib naked; primary lateral veins 5–7 per side, departing midrib at a

(40)55–70° (lowermost to 80°) angle, arcuate-ascending to the margins, weakly visible above, weakly raised, often reddish below; minor veins distinct below, arising from the midrib only; secretory ducts present but inconspicuous. **INFLORESCENCES** 1 per axil; peduncle 2–3 cm long, 5–6 mm diam.; **spathe** 7–8 cm long (2.3–3.6 times longer than peduncle), weakly constricted above the tube, 1.5 cm diam. at constriction, green throughout, pinkish within; spathe tube 3–4 cm long, 1.9 cm diam.; **spadix** 5.5–7 cm long, white throughout; pistillate portion 1.2 cm long, 1 cm diam.; staminate portion 4.3–5.8 cm long, 8–12 mm diam.; pistils 1–1.3 mm long, 0.7 mm diam.; ovary 4-locular; locules 0.8 mm long, 0.3–0.4 mm diam., with axile placentation; ovules 4 per locule, 2-seriate, 0.2 mm long, longer than funicle; funicle 0.1 mm long, adnate to lower part of partition, style similar to style type B; style apex flat to concave; stigma ± hemispheroid, unlobed, 0.8 mm diam., 0.2 mm high, covering entire style apex; the androecium truncate, ± prismatic, margins irregularly 4–6-sided, mostly 4–5-sided, 0.8–1.1 mm long; thecae oblong, 0.4 mm wide, ± parallel to one another, not contiguous; sterile staminate flowers irregularly 4–6-sided, 0.9–1.1 mm long.

Flowering in *Philodendron knappiae* apparently occurs in the dry season and is documented by only two fertile collections, one flowering and one post-anthesis, both made in March.

*Philodendron knappiae* is endemic to western Panama, known only from the type locality in Chiriquí Province on Cerro Hornito in *Tropical Lower Montane rain forest* at 2100 m elevation.

*Philodendron knappiae* is a member of *P.* sect. *Castigima* subsect. *Macrobolium* ser. *Macrobolium*. This species is recognized by its relatively small stature; short internodes (distally) with smooth, brown epidermis; adaxially sulcate, sharply two-ribbed, green but densely short-red-lineate, deciduous cataphylls; bluntly D-shaped petioles (about as long as the blades); small, ovate-triangular dark brown-drying blades with a hippocrepiform sinus; and solitary inflorescences with the spathe green outside and pink inside.

The species is perhaps most easily confused with *P. wilburii*, which has leaf blades of similar size and shape. The latter species differs in having longer internodes drying with a tan, glossy epidermis, thicker blades drying yellowish green to dark olive brown above, 2–3 inflorescences per axil, and 2 ovules per locule (vs. 4 per locule for *P. knappiae*).

*Additional specimens examined.* PANAMA, Chiriquí Cerro Hornito, above Los Planes de Hornito, 1750 m, 8°41'N, 82°10'W, *Croat 67982* (CM, MO); 2100 m, *Kress et al. 82-1363* (DUKE).

***Philodendron lazorii*** Croat, sp. nov. TYPE: Panama. Canal Area: vic. Madden Lake, along both sides of stream SSE of pumping station (SE of dam), 140 m, 9°13'N, 79°37'W, 18 Jan. 1990, Croat 69833 (holotype, MO-3789003-5; isotypes, AAU, B, CAS, CM, COL, CR, F, GH, K, MEXU, NY, PMA, QCNE, RSA, SCZ, TEX, US, VEN). Figures 13, 14, 250-252, 255-258.

Plants hemiepiphytic; internodia 1-4 cm longa, 5 cm diam.; cataphylla acute D-formata, persistentia semi-intacta nodis superioribus, demum decidua; petiolus teres, 37-63(76) cm longus, 5-6 mm diam.; lamina late ovata, 32-54 cm longa, (22)34-49 cm lata, cordata basi, in sicco canoviridis; sinus hippocrepiformis vel obovatus; inflorescentia 2; pedunculus 8.6-17 cm longus; spathe 12-13.5 cm longa; pistilla (4)5-6-locularia; loculi 10-14(18)-ovulati.

Hemiepiphytic; stem appressed-climbing (scandent as juvenile); internodes weakly striate, 1-4 cm long, 5 cm diam.,  $\pm$  broader than long, light olive-green to dark yellow-green, glossy, roots 1-3 per node, dark brown to reddish brown, matte, smooth to densely scaly, to 30 cm or longer, 3-5 mm diam., feeder roots to 8 mm diam., densely scaly; cataphylls 17-19 cm long, pale yellow-green, sharply D-shaped, margins acutely raised, persisting semi-intact at upper nodes with a dense reticulum of coarse fibers, often overlain with a thin, fragmented epidermis, becoming dilacerated, eventually deciduous; petioles 37-63(76) cm long, 5-6 mm diam., terete, dark green, firm, flexible, surface matte, faintly dark green striate; blades broadly ovate, subcoriaceous, long-acuminate at apex, cordate at base, 32-54 cm long, (22)34-49 cm wide (0.97-1.4 times longer than wide, averaging 1.13 times), (0.65-0.9 times the petiole length), slightly shorter in length than petiole, upper surface dark green, drying gray-green, weakly to semiglossy, lower surface drying gray-green, matte, much paler; anterior lobe 27-33 cm long, 36 cm wide (1.6-2.2 times longer than posterior lobes); posterior lobes 13-17 cm long, 16-19.5 cm wide, obtuse; sinus hippocrepiform to obovate, 9.5-11.5 cm deep; midrib flat to broadly raised, concolorous to slightly paler than surface above, narrow-rounded to bluntly acute, darker than surface below; basal veins (5)6-7 per side, with 0-1 free to base, 3rd and higher order veins coalesced 2-4.5 cm, obscure; posterior rib 2-3 cm long along the sinus; primary lateral veins 3-4 per side, departing midrib at a 40-50° then to 70° angle, straight to weakly arcuate to the margins, weakly and obtusely sunken and concolorous or paler above, convex and darker than surface below; mi-

nor veins moderately distinct, arising from both the midrib and primary lateral veins, drying weakly puckered and darker than surface below, upper surface usually drying with whitish cells visible, lower surface drying smooth and minutely brownish to whitish speckled. INFLORESCENCES 2 per axil; peduncle 8.6-17 cm long, to 1.1 cm diam., medium green, lightly white-striate, heavily white-striate nearest apex; spathe 12-13.5 cm long (spathe equal in length to or slightly shorter than peduncle), constricted above the tube, 2.9 cm diam. at constriction, margins pale to creamy; spathe blade yellowish green outside, 8 cm long (opening 4 cm wide), creamy on outer margins, glossy inside; spathe tube medium green, finely white-striate with pale margins outside, 5 cm long, 3.3 cm diam., pale yellowish green, glossy inside; spadix bluntly pointed at apex, 10.9-11.3 cm long, constricted 1.5 cm above base of fertile staminate portion; pistillate portion pale yellowish green, 3.3-4 cm long, 1 cm diam. at base, 1.5 cm diam. midway, 1.4 cm diam. at apex; staminate portion 7.8-9.3 cm long; fertile staminate portion creamy white, 1.7 cm diam. at middle, 11 mm diam. ca. 1 cm from apex; sterile staminate portion barely detectable, creamy white, drying darker than fertile staminate flowers, 1.7 cm diam.; pistils 1.9-2.5 mm long; ovary (4)5-6-locular, 1.1 mm diam., locules 1.1-1.6 mm long, 0.3-0.4 mm diam., with axile placentation; ovules 10-14(18) per locule, 2-seriate, 0.3 mm long, longer than funicle; funicle 0.1-0.2 mm long, adnate to lower part of partition, style 0.9 mm diam., similar to style type D; style apex  $\pm$  rounded to bluntly pointed; style boss broad and pronounced; stigma subdiscoid, unlobed,  $\pm$  truncate, 1.5 mm diam., 0.3 mm high, covering entire style apex, depressed shallowly and medially; the androecium truncate, prismatic, oblong, margins irregularly 4-6-sided, 0.7-1 mm long; thecae oblong, 0.4-0.5 mm wide, not contiguous,  $\pm$  parallel to one another; sterile staminate flowers irregularly 4-6-sided, slightly clavate to prismatic, 1.2-1.9 mm long. JUVENILE plants with internodes matte, gray-green, 6 cm long, 7 mm diam.; petioles terete; blades weakly velvety; upper surface  $\pm$  glistening-glossy, lower surface matte, much paler, with flecks of brilliance; minor veins very distinct, darker than surface.

Flowering in *Philodendron lazorii* occurs in the late dry season and early rainy season (March through June), with mature fruits in August. Immature fruits have been collected in March and June, which indicates that there must be flowers in the late wet season as well (or perhaps it indicates bimodal flowering).



*Philodendron lazorii* is endemic to Panama, known from the type locality, a region of limestone outcrops near Madden Lake, at about 100 m elevation in *Tropical moist forest*, and in Darién Province at 250 to 1050 m in *Tropical moist forest* and *Premontane wet forest*.

*Philodendron lazorii* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is distinguished by its thick, short internodes, cataphylls persisting semi-intact with an organized network of yellow-brown fibers, terete petioles averaging 1.25 times longer than the blade, and broadly ovate, gray-green-drying blades matte on the lower surface.

*Philodendron lazorii* is probably most closely related to *P. panamense*, which has similar inflorescences. The latter species differs in having semiglossy, ovate-triangular blades usually 1.3 or more times longer than wide (averaging 1.5 times longer than wide), and petioles commonly shorter than the blade. In addition, it has generally shorter peduncles (usually shorter than the spathe).

This species is also similar to *P. jefense*. See that species for a discussion of the differences.

*Philodendron lazorii* is named in honor of one of its original collectors, Robert Lazor (Army Corps of Engineers, Vicksburg), who collected in Panama while a student at Florida State University.

*Additional specimens examined.* PANAMA. Canal Area: Madden Lake Cave area, ca. 9°13'N, 79°37'W, *Tyros & Lazor 6266* (FSU, IBE); Salvador Hill, near Juan Mina, *Bartlett & Lasser 16785* (MICH, MO). DARIÉN: along trail from base camp to Rancho Frio on slopes of Cerro Pirre, 200–450 m, 7°58'N, 77°43'W, *Croat & Zhu 77126* (CAS, CM, CR, MO); W side of Cerro Pirre, 800–1050 m, 7°56'N, 77°45'W, *Croat 68953* (DUKE, HUA, M, MO).

*Philodendron lentii* Croat & Grayum, sp. nov.

TYPE: Panama. Coclé: El Valle region, between Finca Mandarinas and Finca Furlong, ca. 1 km off road, vic. La Mesa, N of El Valle de Antón, 800–900 m, 8°38'N, 80°09'W, 11 June 1987, *Croat 67163* (holotype, MO-3582921; isotypes, AAU, B, BR, CAS, CM, COL, CR, CUV, DUKE, EAP, F, G, GH, HMNM, HUA, IBE, K, MEXU, NY, P, PMA, QCA, RSA, S, SCZ, TEX, US, VEN). Figures 259–265.

Planta hemiepiphytica; caulis scandens; internodia 4–12 cm longa, 1.5–4.5 cm diam.; cataphylla 20–23(40) cm longa, plerumque incostata, interdum 1–2 costata, decidua; petiolus teres vel late D-formatus, interdum late sulcatus, 22–44 cm longus, 4–12 mm diam.; lamina ovato-cordata, 21.6–51 cm longa, 9–25 cm lata, truncata vel subcordata basi; nervis basalibus 1–3(4) utroque, liberis

basi; inflorescentia 1–2; pedunculus 3–10 cm longus; spathe 12–21 cm longa, lamina spatheae extus viridialba vel alba, intus atrimarronina vel creamea; tubo spatheae extus albo vel viridialbo, interdum suffuso subroseo, intus atrimarronino vel violaceipurpureo; pistilla (5)6–8-locularia; loculi (1–2)4–6-ovulati; baccae aurantiacae.

Hemiepiphytic; stem scandent, creeping, assurgent, sap soapy-scented; internodes elongate, semiglossy to matte, 4–12 cm long, 1.5–4.5 cm diam., usually slightly longer than broad, medium to dark green, conspicuously pale striate at upper edge of each node, drying brownish, often narrowly ribbed (ribs irregularly ridged, sometimes warty), epidermis weakly fissured transversely; roots moderately few, to ca. 30 cm long, drying to 2 mm diam., reddish brown; cataphylls 20–30(40) cm long, usually unribbed, sometimes 1-ribbed, sharply 1-ribbed, or weakly to sharply 2-ribbed, greenish, tinged reddish, deciduous, intact; petioles 22–44 cm long, 4–12 mm diam., terete to broadly D-shaped, firm, dark green, sometimes purplish below, broadly sulcate, flattened or broadly convex adaxially, convex abaxially, with adaxial margins obtuse to rounded, surface semiglossy, densely and minutely white or dark striate-lineate; blades ovate to narrowly ovate or ovate-elliptic, coriaceous to subcoriaceous, conspicuously bicolorous, acuminate to narrowly acuminate at apex, rounded to truncate or subcordate at base, 21.6–51 cm long, 9–25 cm wide (1.4–2.7(3.3) times longer than wide), (0.7–1.4(2.25) times longer than petiole), usually about equal in length to petiole, upper surface dark green, semiglossy, drying dark grayish brown to yellowish brown, lower surface semiglossy to matte, paler; anterior lobe 28–45.5 cm long, 9–25 cm wide (4.9–10.2 times longer than than posterior lobes); posterior lobes 3–7 cm long, broader than long, broadly rounded to obtuse; sinus ± V-shaped to arcuate, to 4 cm deep; midrib flat to broadly convex, paler than surface above, convex, tinged reddish or darker than surface, drying minutely granular below; basal veins 1–3(4) per side, with all free to base; primary lateral veins 8–14 per side, departing midrib at a 60–70° angle, weakly arcuate, ascending to the margins, weakly to narrowly sunken or weakly quilted, drying paler than surface, raised along the margins with the center collapsed (forming a channel) above, convex to weakly raised or weakly pleated, darker than surface, drying minutely granular, paler than surface below; interprimary veins weakly sunken or obscure above, visible below; minor veins visible, few, darker than surface, arising from both the midrib and primary lateral veins, minutely etched in upper surface of fresh leaves, drying raised. INFLORES-

CENCES 1-2 per axil; peduncle 3-10 cm long, 3-7(14) mm diam., pale green; **spathe** 12-21 cm long (1.5-3.98(4.5-5.1) times longer than peduncle), margins reddish; spathe blade greenish white to white outside, 4 cm diam., dark maroon to cream inside; spathe tube white to greenish white, sometimes tinged pinkish or red, densely white-lineate outside, 6-9 cm long, dark maroon to violet-purple, densely white-lineate inside; **spadix** sessile; pale greenish to white throughout, cylindrical, tapered, 11-14(18) cm long, broadest near the base or slightly above the middle, constricted below the middle; pistillate portion green, cylindrical to ovoid-tapered, 6.6 cm long in front, 4.5 cm long in back, 1.1 cm diam. at apex, 1.3-1.4 cm diam. at middle, 8-11 mm wide at base; fertile staminate portion greenish white, tapered to clavate or cylindrical, 6.8-9.4 cm long, 9-16 mm diam. at base, 9-15 mm diam. at middle, 8-10 mm diam. ca. 1 cm from apex, broadest at or near the base, broader than the pistillate portion, sterile staminate portion not detectable; pistils 1.6-2.4 mm long, 1-1.8 mm diam.; ovary (5)6-8-locular; locules 1.2-2 mm long, 0.3-0.6 mm diam., ovule sac 0.8-1.5 mm long, with basal or sub-basal placentation; ovules (1-2)4-6 per locule, 1- or 2-seriate, 0.3-0.5 mm long, longer than or equal in length to funicle; funicle 0.1-0.4 mm long (can be pulled free to base); style 0.1-0.5 mm long, 1-1.6 mm diam., similar to style type B; style apex flat or weakly rounded; stigma discoid, brushlike, unlobed, 1-1.3 mm diam., 0.1-0.2(0.6) mm high, covering almost entire style apex or just the center of style apex; the androecium truncate, margins 4-6-sided (4-5-sided); thecae oblong, 0.3-0.5 mm wide, contiguous or  $\pm$  parallel to one another; sterile staminate flowers 4-6-sided, 1.1-1.5 mm long, 0.8-1.3 mm wide. INFRUCESCENCE with berries orange; seeds 3-4 per locule, 1-1.1 mm long, 0.5-0.6 mm diam.

Flowering in *Philodendron lentii* appears to occur during the dry season and early rainy season (February, March, June, and July). Post-anthesis collections exist from January, March through July, and November, indicating a somewhat broader range of flowering. Most post-anthesis inflorescences were collected between March through July. Immature fruits are known from January, April, and July while mature fruits are known from April, June, and July.

*Philodendron lentii* ranges from Costa Rica (Cartago) to Panama (Chiriquí to Coclé), from (210)670 to 1800 m elevation in *Premontane rain forest* and *Tropical Lower Montane rain forest* life zones. In Costa Rica, this species is apparently rare, having

been collected only near Tapantí and Moravia in Cartago Province and near Cariblanco in Alajuela Province. In Panama, it has been collected mostly in the Fortuna Dam region, at Cerro Colorado (both Chiriquí), at Santa Fe in Veraguas and at El Valle and El Copé in Coclé Province.

*Philodendron lentii* is a member of *P.* sect. *Calostigma* subsect. *Macrobilium* ser. *Ecordata*. This species is distinguished by its somewhat scandent habit; elongate internodes; deciduous cataphylls; D-shaped to broadly sulcate petioles (about as long as the blades); ovate, subcordate blades with the few basal veins free; and one to two large inflorescences with the spathe white externally and purplish internally on the tube. Also characteristic are the orange berries. Especially useful for recognition is the upper dried blade surface, which usually dries dark with the lateral primary and minor veins weakly raised.

*Philodendron lentii* may be confused with *P. heleniae*, which has blades of similar size and shape with few pairs of free basal veins and petioles about as long as the blades. That species differs in having more slender stems, subterete petioles, and several to many small, red inflorescences. Although a few collections of *P. lentii* (Knapp 4970, Thompson 5022) report the spathe to be red, these cannot be confused with *P. heleniae*, owing to the much larger size of the spathes (mostly more than 12 cm long vs. mostly less than 10 cm long for *P. heleniae*).

Possibly also belonging to this species is a sterile collection from Ecuador (Esmeraldas), Croat 72298, which differs in having internodes to 30 cm long, more numerous primary lateral veins (to ca. 20), and interprimary veins also sunken on live plants giving the blade the appearance of having very close primary lateral veins. This collection also differs in having the cross-veins prominulous on the lower surface toward the margin. Despite these differences, this Ecuadorian collection surely represents either *P. lentii* or another closely related new species.

*Philodendron lentii* is named in honor of Roy Lent, who first collected the species in 1967. Lent, a resident of Costa Rica, has been an important collector over many years and has collected many new species of Araceae.

*Additional specimens examined.* COSTA RICA. Alajuela: 3 mi. S of Cariblanco, 760 m, Croat 35777 (MO). Cartago: Moravia-Quebrada Platanillo, Moravia, 3-5 km from Finca Racine, 1200-1300 m, Croat 36592 (F. MEXU, MO, US); Río Grande de Oroá, 0.5 km W of Tapantí, 1200 m, Lent 909 (CR, F). PANAMA. Bocas del Toro: Fortuna Dam area, Chiriquí Grande-Fortuna, near Continental Divide, 1170 m, 8°44'N, 82°17'W, Croat 66647 (B, CAS, CR, K, MO, PMA, US); 1200 m, 60360

(CAS, MO); 1.2 mi. N of Divide, 910 m, 8°44'N, 82°17'W, Croat 60441 (K, MO, US). **Chiriquí:** Cerro Colorado, vic. of Cerro Colorado Copper Mine Development, 28 mi. above San Félix, 1200–1500 m, Croat 33273 (MO); Cerro Pate Macho, ca. 5 mi. NE of Boquete, 1800–2200 m, Croat 48567A (MO); 1600–1700 m, 8°48'N, 82°23.5'W, Grayum et al. 6358 (MO); vic. of Boquete, 1630–1780 m, 8°46'N, 82°25'W, Croat 66395A (MO); 66391 (MO); 66355 (CM, MO); Fortuna Dam area, 1100–1200 m, 8°45'N, 82°05'W, Thompson 5022 (CM, MO); 8 km N of Los Planes de Hornito, 1250–1300 m, 8°45'N, 82°12'W, Knapp 4970 (MO, NY); 11.8 mi. N of Los Planes de Hornito, 1400 m, Croat 48682 (CM, MO); 10.8 mi. beyond Los Planes de Hornito, ca. 1400 m, Croat 48712 (MO); 10.1 mi. NW of Los Planes de Hornito, 1250 m, 8°45'N, 82°17'W, Croat 50041 (CM, MO); 10 mi. NW of Los Planes de Hornito, 1260 m, Croat 50102 (MO, NY, PMA); Chiriquí Grande–Fortuna, 4.5–5 km N of dam over Fortuna Lake, 1100–1135 m, 8°43'N, 82°17'W, Croat & Grayum 60007 (K, MO, US); 60069 (K, MO, US); Gualaca–Chiriquí Grande, 8 mi. N of Los Planes de Hornito, 1.4 mi. W of Centro de Operaciones, along trail to Río Hornito, 1010–1130 m, 8°44'N, 82°14'30"W, Croat 67923 (CM, MO); near Fortuna Lake, 8°45'N, 82°18'W, Croat & Zhu 76388 (MEXU, MO); N edge of lake, ca. 1100 m, ca. 8°45'N, 82°15'W, McPherson 9078 (MO, NY); Fortuna Dam, 1200 m, Hammel 2196 (MO); Quebrada Arena, 1050 m, 8°45'N, 82°16'W, Hammel et al. 14706 (CR, MO); 26 km past Gualaca, 670 m, 8°35'N, 82°19'W, Hooser 1329 (CAS, MO). **Coeló:** Penonomé–Coelécito, above Río Cascajal, 5.7 mi. N of Llano Grande, 210 m, 8°40'N, 80°26'W, Croat 67538 (MO, PMA); El Copé region, Alto Calvario, Continental Divide, 5.2 mi. above El Copé, 930 m, Croat 49204 (MO); Alto Calvario, 710–800 m, 8°39'N, 80°36'W, Croat 68724 (MO); El Valle region, Croat 14367 (MO); 2700 ft., *Sytima* et al. 4369 (CR, MO); ca. 800 m, Croat 25434 (F, MO); *Gentry & Dwyer* 3683 (MO); 900–930 m, Croat 37481 (F, MO); 860 m, 8°37'N, 80°08'W, Croat & Zhu 76740 (MO, US); Cerro Gaital, 800–900 m, 8°37'N, 80°07'W, McPherson 11211 (L, MO); 830–900 m, 8°36'N, 80°07'W, Croat 74812 (MO, PMA); *Hartman* 3962 (OS); 1000–1100 m, 8°37'N, 80°07'W, McPherson 12150 (MO); N slope of Cerro Gaital, 800 m, 8°38'N, 80°07'W, *Churchill* 3924 (F, MO). **Verrugas:** Santa Fe region, Alto de Piedra, 800–950 m, 8°33'N, 81°08'W, Croat 67004 (MO); Parque Nacional Cerro Tute, Alto Piedra–Calovebora, 0.5 mi. N of Alto Piedra, 800–1030 m, Croat & Zhu 76897 (MO); 1000–1250 m, Croat 48931 (MO).

***Philodendron ligulatum*** Schott, *Prod. Syst. Arid.* 224. 1860. TYPE: Cultivated from Central America, *Wendland* s.n. (holotype, W? destroyed). Costa Rica. Limón: Ref. Nac. Barra del Colorado, between Río Chirripico and Río Sardina (Sardinal), 10°38'N, 83°45'W, 12 m, 22 Apr. 1990, *Grayum* 9823 (neotype, MO; isoneotype, CR). Figures 269–271.

Hemiepiphytic, usually scandent or appressed-climbing or sometimes epiphytic, sometimes occurring high into canopy, rarely terrestrial; stem appressed-climbing or scandent (flowering stems often loose, semi-erect or spreading), green, sometimes

with white, waxy coating; leaf scars conspicuous, 1.5–2 cm long, 1–1.8(3) cm wide; internodes sometimes weakly flattened on one side, (1)3–9(20) cm long, 0.5–3 cm diam., usually longer than broad, medium to dark green or gray-green to brownish, weakly glossy, drying gray-green to pale yellow-brown, sometimes irregularly ridged or cracked, sometimes closely transverse-fissured; epidermis peeling, bubbling or with loose flakes; roots several per node, 15–45 cm long, drying 1–2 mm diam.; cataphylls usually sharply 2-ribbed, sometimes weakly and bluntly 1–2-ribbed, sometimes sharply D-shaped with adaxial margins winged to 6 mm high, 14–19 cm long, sharply flattened, with obtuse medial rib, green to whitish, sometimes tinged reddish, densely dark-lineate, sometimes densely dark-speckled, deciduous intact, margins minutely undulate; **petioles** (7)20–38 cm long, 5–15 mm diam., subterete to obtusely flattened, rarely D-shaped, sometimes with a thin, medial rib toward apex, sometimes with adaxial margins winged to 6 mm high, spongy but brittle (fresh), medium to dark green, obtusely to sharply flattened to sulcate adaxially, the margins at least sometimes acute, broadly rounded abaxially, surface matte to semiglossy, densely short dark green-lineate or speckled, with a deep maroon to purple ring around apex; sheath flattened, unopened, to 3–10 cm long, for up to half the length of the petiole; **blades** oblong-elliptic to narrowly ovate or narrowly oblanceolate-elliptic, subcoriaceous, weakly to moderately bicolorous, acuminate to long-acuminate at apex (the acumen inrolled or apiculate and sometimes downturned, 1–4 mm long), subcordate, obtuse to narrowly rounded or weakly cordate at base, (14)18–61 cm long, 8–19 cm wide (1.57–5.4 times longer than wide), (1.1–3.89(6.3) times longer than petiole), margins weakly undulate; upper surface dark green, semiglossy to moderately glossy, drying blackish brown to dark grayish or grayish brown, lower surface moderately paler, matte to weakly glossy, mottled violet-purple or maroon, drying dark olive-green to yellow-brown, dark olive-brown or dark brownish black; posterior lobes, when present, rounded to narrowly rounded, about as broad as long, 4–5 cm long, 5.5–6.6 cm wide, held close to petiole; sinus somewhat V-shaped, 0.5–3 cm (mostly 2.5 cm) deep; midrib flat to broadly convex, concolorous to paler than surface, sometimes sparsely purple-speckled near base above, narrowly rounded to convex to broadly convex, sometimes round-shouldered, paler or darker than surface, or sometimes concolorous below, sometimes sparsely purple-speckled near base below; basal veins not evident or 1–3 per side; primary lateral veins 4–8(12) per

side, departing midrib at a 40–70(75)° angle, ± straight or slightly curved to the margins, dark green, weakly sunken to etched, concolorous above, weakly raised to convex and paler than surface below (sometimes with purple spots on older leaves), drying brownish; interprimary veins obscure to impressed, paler than surface, weakly sunken above, weakly raised, darker than surface below; minor veins weakly visible above, very close, obscurely and weakly raised, darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, 1(2) per axil; peduncle (5)7.5–17.5(28) cm long, 7–10 mm diam., subterete, obtusely angular to 3-sided, pale to medium green; spathe 10–19.5 cm long, 1.5–2 cm diam. (to 6.5 cm wide when flattened), ((0.8–1.3)1.4–1.6(1.8) times shorter than peduncle), semiglossy, acuminate at apex, barely or not at all constricted, ribbed on back, margins revolute; spathe blade white or cream, sometimes sparsely red-spotted outside, sometimes tinged purple-violet on outer margin, 7–8.5 cm long, 1.7 cm diam. (opening 7.5–7.8 cm long, 3.2–4 cm wide), whitish inside; resin canals bright orange, appearing throughout lower ¼ of spathe and to near base of tube inside; spathe tube oblong, pale to medium green with pale violet nearest base, sparsely red-spotted outside, 6.5–7 cm long, 1.2–1.5 cm diam., glossy, whitish or pale green, sometimes red to violet-purple (B & K purple 5/10) near base or throughout tube inside; **spadix** ± tapered with slight bend, held ± erect, protruding forward from spathe, rounded at apex, (8.3)11–15.5 cm long; pistillate portion light green, (1.4)3.7–6.5 cm long, 9–13(16) mm diam. at apex, 10–14(17) mm diam. at middle, 9–12(15) mm diam. at base; staminate portion (6.8–7.2)10.5–11.2 cm long; fertile staminate portion (9)11–16 mm diam. at base, (7)11–16 mm diam. at middle, (5)9–11 mm diam. ca. 1 cm from apex, broadest slightly above the middle, as broad as or broader than the pistillate portion, as broad as or slightly narrower than the sterile portion; sterile staminate portion generally broader than pistillate portion, whitish or tinged faintly violet, 1–1.2 cm diam.; pistils 1.2–2.7(3.5) mm long, (0.8)1.1–1.9(2.6) mm diam.; ovary (5)6–8-locular, 0.8–1.8(2.4) mm long, (0.8)1.1–1.9(2.6) mm diam., walls embedded with granular, crystal-like particles; locules (0.8)1.1–1.8(2.4) mm long, (0.1–0.2)0.4–0.7 mm diam., ovule sac 0.5–0.7(1–1.3) mm long, with sub-basal placentation; ovules 1 per locule, contained within translucent, gelatinous envelope, 0.2–0.6 mm long, usually longer than funicle; funicle 0.1–0.2(0.4) mm long, style 1–6 mm long, (0.8–10)14–19(26) mm diam., similar to style type B or type D (rarely

to type C); central style dome sometimes present; style apex flat to rounded or sloping; style boss sometimes present; funnel shallow when present; stigma somewhat cupulate, truncate, unlobed to subdiscoid, (0.4)0.9–1.2 mm diam., 0.1–0.3 mm high, covering center of or entire style apex, inserted on center of style apex or style boss or funnel if present, sessile, papillate, semiglossy; the androecium truncate, margins irregularly 4–6-sided, 1–3 mm long; thecae oblong to ovate, sometimes elliptical, 0.3–0.6 mm wide, ± parallel to one another or contiguous; pollen subspheroidal to oblong or obovoid; sterile staminate flowers usually 4–6-sided, 1.2–2.2 mm long, 1.3–2 mm wide. INFRUCTION with berries oblong-elliptic, 3 mm long; seeds oblong, light yellow-brown, 1.4 mm long, 0.4 mm diam., narrowly ribbed longitudinally. JUVENILE blades with lower surface sometimes tinged maroon.

*Philodendron ligulatum* ranges from Nicaragua (Chontales and Zelaya) along the Atlantic slope of Costa Rica and along the Atlantic slope and Continental Divide of Panama to Colombia (Antioquia and Chocó). It occurs in *Tropical wet forest* and *Premontane rain forest* life zones, from sea level to 1200 m elevation (to 600 m in Nicaragua and 650 m in Costa Rica).

*Philodendron ligulatum* is a member of *P. sect. Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is characterized by its somewhat scandent to appressed-climbing habit; generally elongate internodes; sharply 2-ribbed to obtusely 1-ribbed cataphylls; more or less spongy, obtusely flattened to somewhat D-shaped petioles clearly demarcated from the blade by a purple ring; usually more or less oblong blades narrowly cordulate at the base (or with the blades ovate and cordate) with usually 1–2 free basal veins; and usually 1–2 inflorescences per axil.

This species comprises three varieties. *Philodendron ligulatum* var. *ligulatum* exists throughout the range of the species, whereas varieties *ovatum* and *heraclioanum* are endemic to Panama. See under those varieties for the differences separating them from variety *ligulatum*. *Philodendron ligulatum* var. *ovatum* is endemic to central Panama in Coclé and Veraguas Provinces, while *P. ligulatum* var. *heraclioanum* is known from far eastern Panama in San Blas and Darién.

*Philodendron ligulatum* is probably close to *P. correae* from upland Chiriquí in western Panama. The latter species differs in lacking distinct primary lateral veins. It also differs in occurring at generally higher elevations (780–1400 m).

*Philodendron ligulatum* may also be confused with *P. immixtum*, also a vine but with smaller leaves. See under the latter species for a discussion of the differences. *Philodendron ligulatum* may also be confused with *P. wendlandii*. The latter has similar blades and spongy petioles, but differs in being a true epiphyte with short internodes and petioles sharply flattened adaxially and usually much broader than thick.

The protologue of *P. ligulatum* describes a sterile plant, whereas Schott's color plates (Icones) depict fertile material. None can therefore be used as type material. In addition, Engler's *Araceae* No. 180 prepared in 1883 and said to have been collected from the living type plant was not studied by Schott. This species is represented by excellent illustrations of fertile material by Schott, one of which could have served as a neotype, but it was deemed best to select a new fertile specimen for the neotype.

The names *Philodendron ligulatum* and *P. ligulatum* (L.) K. Koch (*P.* subg. *Pteromischum*) have been confused in the past. Pursuant to an unpublished request for a recommendation by the Committee for Spermatophyta, the two names have officially been ruled not confusable, hence not homonymous (see Nicolson, 1994: 280).

KEY TO THE VARIETIES OF *P. LIGULATUM*

- 1a. Petioles sharply D-shaped with undulate lateral margins adaxially; lower leaf blade surface mottled purple-violet or maroon; ovaries with 2 ovules per locule (lacking an obvious ovule sac); eastern Panama, 0–200 m ..... var. *heracleiostium* Croat
- 1b. Petioles subterete, at most obtusely somewhat flattened adaxially; lower leaf blade surface not colored; ovaries with 1 ovule per locule, contained within an obvious ovule sac; Nicaragua to Panama.
  - 2a. Leaf blades 1.5–1.7 times longer than wide; peduncles longer than spathe; Panama, 770–1200 m ..... var. *ovatum* Croat
  - 2b. Leaf blades 2–4.5 times longer than wide; peduncles shorter than or about as long as spathe; usually occurring below 800 m (rarely to 1200 m) ..... var. *ligulatum*

*Philodendron ligulatum* Schott var. *ligulatum*

Internodes sometimes weakly flattened on one side, (1)3–9(2) cm long, 0.5–3 cm diam.; cataphylls usually sharply 2-ribbed, sometimes weakly 2-ribbed; petioles 7–38 cm long, 5–15 cm diam., subterete, usually at most obtusely somewhat flattened adaxially, sometimes sulcate, sometimes sharply flattened on one margin adaxially, rarely D-shaped; lower leaf blade surface not colored; blades oblong-elliptic, 24–60 cm long, 8–19 cm wide (2.1–4.4 times longer than wide), (1.1–

3.9(5.1) times longer than petiole), upper surface drying dark olive-green to brownish; sinus 0.5–3 cm (mostly 2.5 cm) deep; basal veins not evident or 1–3 per side; primary lateral veins 5–8(9) per side, departing midrib at 60–70° angle. INFLORESCENCES 1 per axil; peduncle 7.5–17(28) cm long; spathe 10–19.5 cm long, spathe blade white or cream, sometimes sparsely red-dotted outside, spathe blade whitish or sometimes violet-purple near base of tube or throughout inside; spathe tube pale green throughout, pale violet nearest base, sometimes sparsely red-violet outside; spadix (8.3)11–15.5 cm long; pistillate portion (1.4)3.7–6.5 cm long; ovules 1 per locule, contained within an obvious ovule sac; style similar to style type B or D, central dome sometimes present; stigma somewhat cupulate, truncate, unlobed to subdiscoid.

Flowering in *Philodendron ligulatum* var. *ligulatum* occurs in the early rainy season, between May and August, with a few flowering collections from October through March at the end of the rainy season and early dry season. Post-anthesis inflorescences have been collected from February through August (as well as November). It is possible that this species flowers bimodally, once near the beginning of the rainy season and again near the end of the rainy season or the early dry season. Immature fruits are known from July, August, and December, but mature fruits only from December.

*Philodendron ligulatum* is typically somewhat scandent in Costa Rica but is often more nearly an appressed hemiepiphyte in Panama. In Costa Rica, the number of inflorescences per axil is never more than two but in some areas of Panama up to three inflorescences per axil may be encountered, and in the Santa Rita Ridge area of Panama, a particularly unusual specimen (Croat 76954) has up to five inflorescences per axil. That collection also has proportionately somewhat longer petioles than other collections. This unusual specimen is otherwise identical to other plants of the species, even those from the same region. It is perhaps of hybrid origin.

A few Panamanian specimens of *P. ligulatum* var. *ligulatum* from the vicinity of Santa Fe in Veraguas Province (Croat 25692, 48906A, 66914) and in Coelá (Croat 49195) differ in drying yellow-green and having somewhat more coarse venation. These are somewhat intermediate with *P. ligulatum* var. *ovatum*.

Croat 33306, from 1200 m elevation on Cerro Colorado in eastern Chiriquí Province, has black-drying blades averaging slightly more than two

times longer than broad, and is also somewhat intermediate with *P. ligulatum* var. *ovatum*.

A collection from the southern slopes of Panamá Province in the basin of the Río Madroño (Croat & Zhu 77041) is intermediate between *P. ligulatum* var. *heracleoanum* and variety *ligulatum* in having sharply D-shaped petioles lacking undulate-winged margins.

A noteworthy collection from Antioquia Department in Colombia (Fonnegra et al. 1957) perhaps also represents *P. ligulatum* var. *ligulatum*, but differs in having the primary lateral veins drying paler than the surface below rather than darker as is the general case.

*Additional specimens examined for P. ligulatum* var. *ligulatum*. COSTA RICA. **Alajuela:** 3 mi. N of San Miguel, 380 m, Croat 35643 (MO); Cañas-Upala, 4 km NNE of Bijagua, ca. 400 m, Croat 36262 (MO); near Río Zapote, 1.8–2.7 km S of Río Canalete, ca. 100 m, Croat 36360 (MO); Naranjo-Aguas Zarcas, 8.5 km NE of Villa Quezada, ca. 600 m, Croat 46971 (MO). **Guanacaste:** Parque Nacional Rincon de la Vieja Colecta, 400 m, Ruera 1136 (CR, MO). **Heredia:** La Zona Protectora, SSE of Magsasay, Schatz & Grayum 707 (DUKE); Finca La Selva, at confluence of Río Sarapiquí and Río Puerto Viejo, Atlantic slope, 50–80 m, 10°26'N, 84°01'W, Grayum 7673 (CR, MO); Folsom 9973 (MO); 100 m, Grayum 2800 (MO); Hammel 8827 (MO); Croat 61220 (MO); between Río Sució and railroad tracks, SW of Finca Zona Ocho, Río Frio de Sarapiquí, ca. 110 m, 10°18'N, 83°52'30"W, Grayum & Hammel 5567 (CR, MO); near Porto Viejo along road near Río Sució, 20 m, Croat 35730 (MO). **Limon:** Hitoy Cerere reserve, SW of Valle La Estrella, 150–550 m, 9°39'N, 83°02'W, Grayum et al. 5806 (CAS, CR, MO, PMA, US); Turrillalba-Limón, ca. 11 mi. S of Siquirres, 650 m, Croat 43331 (MO); Refugio Nacional de Vida Silvestre Barra del Colorado, Río Chirripocito-Río Sardina, 12 m, 10°38'N, 83°45'W, Grayum 9823 (CR, MO). NICARAGUA. **Chontales:** Cerro Las Nubes, El Tamagón-Loma San Gregorio, ca. 2 km N of Santo Domingo, 600 m, *Grijalva & Ríos* 3455 (MO). **Matagalpa:** NW of Cerro Mustín, 500–800 m, Araquistain & Moreno 2570 (MO). **Zelaya:** Colonia Kururia, <50 m, ca. 14°41'N, 84°04'W, Pipoly 3974 (MO); plantel of Neptune Mining Company, NE Bonanza, 200–350 m, ca. 14°01'N, 84°35'W, Stevens 13029 (MO); Río Iyas, between Quebrada El Toro, 260–280 m, Vincelli 371 (MO); Río Sució, 2 km E of Bonanza, 140 m, Neill 4011 (K, MO). PANAMA. **Bocas del Toro:** Station Milla 7.5 on Changuinola-Almirante Railroad, 100 m, Croat 38098 (CM, COL, CR, F, MO, NY); Almirante-Ojo del Agua, 3–6 km W of Almirante, 30–200 m, Croat 38212 (F, K, MO); vic. of Chiriquí Lagoon, Wedel 1479 (F, GH, MO). **Canal Area:** along Río Indio de Gatún, near sea level, Pittier 2794 (US). **Chiriquí:** Cerro Colorado, along Continental Divide 20–28 mi. from San Félix, 1200–1500 m, Croat 33306 (MO). **Coelá:** La Mesa, N of El Valle de Antón, 785 m, 8°37'N, 80°08'W, Croat 67113 (MO, PMA); 800 m, 25370 (F, MO, US); 800 m, 8°36'N, 80°07'W, Croat & Zhu 76669 (MO); 76694 (CM, MO, NY, PMA); Alto Calvario, 5.2 mi. above El Copé, 930 m, Croat 49195 (BR, CM, DUKE, F, GH, HNMM, MO, PMA, TEX, WIS); 750–900 m, 44734 (MO); 2700 ft., Sytama et al. 4343 (MO); Quebrada Mollejón, ca. 5

mi. N of El Copé, 700 m, Croat 75050 (MO); road to Colesito, 12 mi. from Llano Grande, 200 m, 8°47'N, 80°28'W, Churchill et al. 4018 (F, MO). **Colón:** Santa Rita Ridge road, ca. 22 km from Transisthmian Highway, 500 m, 9°25'N, 79°40'W, Hammel et al. 14474 (L, MO); along trail to Río Indio, beginning 10.6 km from highway, 380 m, Croat 34302 (MO, NY, PMA); 26 km from highway, 500 m, 9°26'N, 79°57'W, Knapp et al. 1737 (MO); Mile 6.5, 370 m, 9°21'15"N, 79°44'W, Croat & Zhu 76954 (B, CAS, COL, CR, F, K, MEXU, MO, NY, PMA, SCZ, US, VEN); Sahánitas-Portobelo, Río Piedras Lumber Road, 250 m, 9°22'30"N, 79°41'30"W, 250 m, Croat 75164 (MO, PMA); 75171 (MO); Portobelo-Nombre de Dios, 4.1 mi. W of Nombre de Dios, 50 m, 9°35'N, 79°33'W, Croat 67318 (AAU, CAS, CM, COL, CR, DUKE, F, G, HNMM, K, KYO, L, LE, MBM, MO, NY, P, PMA, QCA, RSA, TEX, US, VDB); Portobelo-Nombre de Dios, vic. Río Indio, 50 m, Croat 33603 (MO, PMA); above road 18 km past Sardinilla on road to Nombre de Dios, 150–300 km, Croat 26100 (MO); Río Boquerón, past Salamanca, 300–400 m, 9°20'N, 79°35'W, Knapp & Sytama 2389 (MO); Río San Agustín flowing into Río Guanche, ca. 0 m, 9°30'N, 79°40'W, Churchill et al. 6017 (CAS, MO, NY, PMA, RSA); Río Iguanita-Portobello Road, <50 m, 9°27'N, 79°42'W, Croat 49778 (F, MO); Río Guanche above bridge on Portobello Road, ca. 3–5 km above bridge, 50–200 m, Croat 36970 (MO); 4 km up Río Guanche, 0–50 m, 9°30'N, 79°40'W, Knapp 997 (MO); ca. 1.5 mi. S of road, <100 m, 9°27'N, 79°40'W, Croat & Zhu 76237 (MO); vic. of Guáimino on Río Miguel de la Borda, Knapp 9983 (MO, SCZ); Río Fató, 10–100 m, Pittier 3867 (NY, US). **Panamá:** El Llano-Cartí Road, 6.8 mi. from highway, 350 m, Croat 49102 (MO); Cerro Jefe, 15.4 mi. from Panamerican Highway, ca. 1000 m, Croat 49092 (MO); Serranía de Majé, high point of ridges S of Ipeti, 650–800 m, 8°45'N, 77°30'W, Knapp et al. 4490 (MO); El Valle de Madroño-La Saena, 2.5 mi. N of El Valle de Madroño, 180 m, 9°14'25"N, 79°05'W, Croat & Zhu 77041 (MO). **San Blas:** El Llano-Cartí Road, 1–2 mi. N of Nusagán on road to Cartí, 250–275 m, 9°20'N, 79°W, Croat & Zhu 765804 (MO). **Verraguá:** vic. of Santa Fe, past Escuela Agrícola Alto de Piedra, at Río Segundo Brazo, 480 m, 8°33'N, 81°08'W, Croat 66914 (MO, US); ca. 5–8 km beyond agricultural school, 730–770 m, Croat 25962 (F, MO); Santa Fe-Calovebora, 1.7 mi. past Alto Piedra School, 1.5 mi. beyond Quebrada Casilla, 570 m, 8°33'N, 81°08'W, Croat & Zhu 76862 (MO); Río Concepción, Lewis 2793 (MO, NY).

#### *Philodendron ligulatum* var. *heracleoanum*

Croat, var. nov. TYPE: Panama. Darién: Parque Nacional Cerro Pirre, vicinity of station along Río Perinico, 110 m, 8°01'N, 77°44'W, 26 July 1994, Croat & Zhu 77098 (holotype, MO—04619347–9; isotypes, AAU, B, CAS, CM, COL, CR, DUKE, F, GB, GH, HUA, IBE, K, L, M, MEXU, NY, OOM, P, PMA, QCA, SEL, TEX, U, US, VEN, WIS). Figures 274–276.

Internodia 1 cm longa, 1.1–2.5 cm diam.; cataphylla 15–20 cm longa, obtuse 2-cosata, decidua; petiolus acute D-formati, (8.5)20–28 cm longus, marginibus alatis, lamina (29)46–61 cm longa, (8.5)12–15 cm lata, oblongo-elliptica vel anguste oblanceolata-elliptica, obtusa basi, in

siccio nigrescens; inflorescentia 1; pedunculus (5)13-15 cm longus, 1 cm diam.; spathe 10-12 cm longa.

Internodes short, to 1 cm long, 1.1-2.5 cm diam., usually longer than broad, medium green, semi-glossy, drying pale yellow-brown, conspicuously and irregularly ridged and cracked; roots moderately and irregularly ridged and cracked; roots moderately few, to 30 cm long, 4 mm diam., reddish brown, weakly glossy, finely scaly; cataphylls 15-20 cm long, bluntly 1-2-ribbed, sharply 2-ribbed near apex; **petioles** (8.5)20-28 cm long, sharply D-shaped, with adaxial margins winged (to 6 mm high); **blades** oblong-elliptic to narrowly oblanceolate-elliptic, obtuse to narrowly rounded at base, (29)46-61 cm long, (8.5)12-15 cm wide ((2.9)4.3-5.4 times longer than wide), (2-2.6(6.3) times longer than petiole), upper drying blackish, lower surface matte, heavily tinged or mottled violet-purple or maroon, drying dark brownish black; midrib narrowly rounded and paler than surface below with maroon spots on older veins; basal veins lacking; primary lateral veins 5-12 per side, departing midrib at a 40-50° angle to the margins, green with maroon spots on older plants. **INFLORESCENCES** 1 per axil; peduncle (5)13-15 cm long, 1 cm diam.; **spathe** pale green, 10-12 cm long, to 6.5 cm wide when flattened, tinged violet-purple on outer margin and at base outside and at base inside; spathe blade greenish inside; spathe tube pale green, glossy inside; **spadix** 9.5-12 cm long; pistillate portion 4.3 cm long in front, 3.6 cm long in back; ovules 2 per locule, contained within transparent matrix, style similar to style type C; style funnel shallow to moderately deep. **INFRUTESCENCE** with berries oblong-elliptic, 3 mm long; seeds oblong, light yellow-brown, 1.4 mm long, 0.4 mm diam., narrowly ribbed longitudinally. **JUVENILE** blades with lower surface heavily tinged maroon.

Flowering in *Philodendron ligulatum* var. *heraclioanum* is poorly known. Pre-anthesis flowering collections were seen from March and in July. The inflorescence on the July collection was nearly fully emerged so that it would certainly have opened in less than one month. Immature fruits were seen in October.

*Philodendron ligulatum* var. *heraclioanum* is endemic to Panama, known definitely only from the type locality at the base of Cerro Pirre in Darién Province from 50 to 200 m elevation in a *Premontane wet forest* life zone.

This variety is characterized by its bluntly two-ribbed cataphylls; sharply D-shaped, marginally winged petioles; and the oblong-elliptic to narrowly oblanceolate-elliptic, blackish drying blades heavily

ly tinged with maroon on the lower surface when young.

*Philodendron ligulatum* var. *heraclioanum* differs from both varieties *ligulatum* and *ovatum* in its sharply D-shaped petioles with undulate-winged margins; in having 2 ovules per locule without an obvious ovule sac (vs. 1 ovule per locule contained in an ovule sac for the other two varieties); and in the maroon coloration of the lower surface of the young blades. In contrast, the other two varieties have petioles that are typically terete or subterete, merely obtusely flattened adaxially.

*Philodendron ligulatum* var. *heraclioanum* may be confused with *P. pseudauriculatum*, but the latter differs in having the petioles unwinged and, at most, obtusely flattened adaxially. In addition, *P. pseudauriculatum* has never been reported to have the blades purplish on the lower surfaces.

A collection from San Blas in far eastern Panama is probably also this variety; it differs in having much longer internodes (to 10 cm long and drying ca. 5 mm diam.) and proportionately shorter petioles.

*Philodendron ligulatum* var. *heraclioanum* is named in honor of Panamanian collector Heraclio Herrera, who was one of the first to collect the taxon.

*Additional specimens examined.* PANAMA, Darién: Cerro Pirre National Park, W side of Cerro Pirre, base camp, 50 m, 8°N, 77°48'W, Croat 68963 (MO); Parque Nacional Darién, Estación Rancho Frío, N base of Cerro Pirre, ca. 9 km S of El Real, Quebrada Perisencio, 70-270 m, 8°01'N, 77°44'W, Hammel et al. 16145 (MO). **San Blas:** Quebrada Musargandi, Isla de Tubualá, 200-400 m, 8°54'N, 77°46'W, Herrera 1304 (CAS, CM, MO, PMA).

***Philodendron ligulatum* var. *ovatum*** Croat, var. nov. **TYPE:** Panama, Veraguas: vicinity of Santa Fe, along road between Alto Piedra and Calovebora, 0.5 mi. N of Alto Piedra, on slopes of Cerro Tute, Parque Nacional Cerro Tute, 800-1030 m, 15 July 1994, Croat & Zhu 76888 (holotype, MO- 4619517; isotypes, B, COL, CR, F, K, NY, PMA, US, VEN). Figure 273.

Internodia 1-6(8) cm longa, 1-1.7(3) cm diam.; cataphylla 13-16 cm longa, incostata vel acute 2-costata; petioli (10)15-19 cm longus, 7 mm diam.; lamina anguste ovata, subcordata basi, (14)18-26(39) cm longa, (8.5)12.5-16 cm lata, in sicco viridibrunnea supra, flavibrunnea infra; inflorescentia 1; pedunculus 12-16.5 cm longus, 7-10 mm diam.; spathe 11.5-18.5 cm longa, extus omnino viridialba; tubo spathae intus rubro basi.

Internodes 1-6(8) cm long, 1-1.7(3) cm diam., sometimes to 3 m or more long, sometimes covered with a thin layer of translucent white wax; cata-

phylls 13–16 cm long, unribbed to bluntly 1-ribbed or sharply 2-ribbed (ribs to ca. 4 mm high, incurled); **petioles** (10)15–19 cm long, 7 mm diam., subterete and obtusely flattened toward apex; **blades** narrowly ovate, weakly cordate at base, (14)18–26(39) cm long, (8.5)12.5–16 cm wide (1.57–1.65 times longer than wide), upper surface drying greenish brown to blackened, lower surface drying yellowish brown to dark olive-brown; posterior lobes rounded to narrowly rounded, 4–5 cm long, 5.5–6.6 cm wide, narrowly rounded to obtuse; sinus somewhat V-shaped, 2.5–3 cm deep; midrib convex to round-raised below; basal veins 2–3 per side, free to base; primary lateral veins 4–6 per side, departing midrib at a 55–65(75°) angle. **INFLORESCENCES** 1 per axil; peduncle 12–16.5 cm long, 7–10 mm diam.; **spathe** 11.5–18.5 cm long, 1.5–2 cm diam. (0.8–0.95 times as long as peduncle), greenish white throughout; spathe tube sometimes pale to medium green outside, pale to medium green, sometimes red to purplish violet at base inside; **spadix** 10–12.5 cm long; pistillate portion 4–4.5 cm long; pistils 1.7–1.8 mm long, 1.7–1.8 mm diam., whitish; ovules 1 per locule, contained within transparent, gelatinous ovule sac, style similar to style type B; stigma subdiscoid, unlobed, often truncate.

Flowering in *Philodendron ligulatum* var. *ovatum* occurs in the rainy season. Collections at or near anthesis have been made in July, October, and November. Observations on a single plant in July showed a series of inflorescences with one at anthesis and one other on the same stem in post-anthesis condition. No fruiting collections are known.

*Philodendron ligulatum* var. *ovatum* is endemic to Panama, known only from Santa Fe in Veraguas and at El Copé in Coclé Province in *Premontane rain forest* at 770 to 1200 m elevation.

This taxon is characterized by its smooth, brown-drying stems, elongate internodes, sharply 2-ribbed to bluntly 1-ribbed deciduous cataphylls, subterete to D-shaped petioles about as long as the blades, ovate, subcordate blades with two to three pairs of free basal veins, and solitary inflorescences with the peduncle longer than the spathe.

*Philodendron ligulatum* var. *ovatum* differs from variety *ligulatum* in having blades 1.5–1.7 times longer than wide (vs. 2–4.5 times longer than wide in var. *ligulatum*); peduncles longer than the spathe (vs. shorter than or about as long as the spathe for var. *ligulatum*); and two ovules (rather than one) per locule. In addition, *P. ligulatum* var. *ovatum* usually occurs at higher elevations (770 to 1200 m)

vs. usually below 800 m for *P. ligulatum* var. *ligulatum*. Curiously, leaves of a few specimens of *P. ligulatum* var. *ovatum* dry yellowish green or brown, as opposed to the somewhat blackened color typically associated with the species.

*Philodendron ligulatum* var. *ovatum* may also be confused with *P. lentii* Croat & Grayum with which it also occurs in the El Copé region. That species differs in having usually more than ten pairs of primary lateral veins with several pairs of interprimary veins visible between them on the dried upper blade surfaces (vs. only about five pairs of primary lateral veins and no interprimaries visible on the upper dried blade surface). In addition, the spathes of *P. lentii* are typically short-pedunculate.

*Additional specimens examined.* PANAMA. Coclé: Alto Calvario, ca. 6 mi. N of El Copé, 770 m, 8°38'N, 80°35'W, Croat & Zhu 76754 (MO); Croat 67572 (MO); 68767A (CM, MO); 1200–1300 m, 8°38'N, 80°36'W, *Sytna* 1903 (MO); 650–850 m, *Fobom* 6221 (MO, PMA); ca. 5 mi. N of El Copé, 900–1000 m, Croat & Zhu 75055 (K, MO, SCZ). Veraguas: vic. Escuela Agrícola Alto Piedra near Santa Fe, 1050–1150 m, Croat 48906A (MO).

***Philodendron lanense* Croat, sp. nov. TYPE:**

Panama. Panamá: vic. of Cerro Jefe, along road to summit which leads S off main road to La Eneida, 750–800 m, 9°14'N, 79°22'W, Croat 67092 (holotype, MO-3582669-71; isotypes, B, CAS, COL, CR, F, G, HUA, K, M, MEXU, NY, P, PMA, SCZ, SEL, TEX, US, VEN). Figures 268, 277, 278.

Planta plerumque hemiepiphytica, raro terrestris; internodia 5–10 cm longa, 4–6 cm diam.; cataphylla 30–41 cm longa, plerumque incostata, persistentia semi-intacta aut ut fibrae nodis supremis; petiolus subterete, 49–76 cm longus, 1–2 cm diam.; lamina 51–77 cm longa, 39–60.4 cm lata, late ovato-cordata; nervi lateralibus 1 (4)5–6(7) utroque; inflorescentia (1)2–3; pedunculus 3–10.5 cm longus; spathe 12.5–24 cm longa, lamina spathe extus atriviridi, intus alba; tubo spathe atriviridivirescente basi; pistilla (5)6–8(9)-locularia; loculi (6)12(20)-ovulati.

Usually hemiepiphytic, rarely terrestrial; stem appressed-climbing, to 1 m long, sap clear, viscous, sticky, spicy- to turpentine-like-scented; internodes usually short, dark green, soon gray-green, finally light brown, moderately scurfy, semiglossy, 5–10 cm long, 4–6 cm diam., usually broader than long, sometimes longer than broad; roots greenish, becoming brown, 3–4 mm diam.; cataphylls 30–41 cm long, usually unribbed, sometimes weakly 2-ribbed, light to medium green, sharply and broadly D-shaped, persisting semi-intact or as fibers at uppermost nodes, fibrous below, the fibers thin, pale, disorganized, light gray. LEAVES erect-spreading, clustered at or near stem apex; **petioles** 49–76 cm



long, 1–2 cm diam., subterete, somewhat spongy, pale to medium green, rarely with medial rib adaxially, with adaxial margins sometimes acute, surface semiglossy to almost matte, densely and in conspicuously short-lineate, at least sometimes with a dark green ring at apex; **blades** broadly ovate-cordate, moderately coriaceous, abruptly acuminate, sometimes acute at apex, conspicuously cordate at base, 51–77 cm long, 39–60.4 cm wide (1.2–1.6 times longer than wide), (0.9–1.2 times longer than petiole), about equal in length to petiole, margins  $\pm$  hyaline, upper surface dark green, semiglossy to moderately glossy, drying dark brown, lower surface semiglossy to almost matte, paler, drying yellow-brown; anterior lobe 41–60 cm long, 39–60.4 cm wide (0.9–1.4 times longer than wide), (2.2–3 times longer than posterior lobes); posterior lobes 18.5–24 cm long, 16–31 cm wide, obtuse; midrib almost flat to broadly convex to broadly flattened, paler than surface above, convex, paler than surface below; basal veins 4–6 per side, with 0–1 free to base, the remaining coalesced 6–9 cm; posterior rib usually naked for 2–3 cm; primary lateral veins (4)5–6(7) per side, flat to weakly sunken, paler than surface above, convex, paler than surface below; minor veins moderately obscure above, darker than surface, arising from both the midrib and primary lateral veins, drying prominulous above and below; secretory canals distinct. **INFLORESCENCES** (1)2–3 per axil; peduncle 3–10.5 cm long, 1.1–1.2 cm diam., coarsely white striate near apex; **spathe** 12.5–24 cm long (1.7–5 times longer than peduncle), oblong, dark to medium green throughout outside, markedly white-lineate near the base, faintly so above the base, moderately constricted above the tube; spathe blade weakly and densely lineate outside (opening broadly elliptic in face view), white inside; spathe tube cylindrical, markedly white-lineate near the base outside, 6–8.5 cm long, 3–5.5 cm diam., dark reddish violet, suffused onto base of blade inside; **spadix** sessile, greenish white throughout,  $\pm$  cylindrical, protruding prominently forward from and out of the end of spathe (at anthesis), 10–17 cm long; pistillate portion creamy white (anthesis), reddish (pre-anthesis), weakly tapered toward apex, 4.1–5 cm long, 2.5–5 cm long in front, 1 cm diam. throughout, 1.1–1.5 cm diam. at apex, 1.4–1.6 cm diam. at middle, 1–1.5 cm wide at base; staminate portion 6–14.4 cm long; fertile staminate portion white,  $\pm$  cylindrical 1.1–1.7 cm diam. at base, 1.1–1.9 cm diam. at middle, 7–10 mm diam. ca. 1 cm from apex, broadest  $\pm$  at the middle, slightly broader than the pistillate portion, narrower than the sterile portion; sterile staminate portion 1.1–1.9 mm diam.; pistils 1.2–

4.8(7.2) mm long, 1.1–2.4(4.3) mm diam.; ovary (5)6–8(9)-locular, 1.5–3.5(6.8) mm long, 1.1–2.4(4.3) mm diam., walls sometimes embedded with granular, crystal-like particles, locules 1.5–3.5(6.8) mm long, (0.2)0.6–1.5 mm diam., thin and membranous, ovule sac 1.9–2 mm long, with axile placentation; ovules (6)12(20) per locule, 1–2-seriate, contained within gelatinous ovule sac, 0.2–0.45 mm long, slightly longer than funicle; funicle 0.1–0.3 mm long, style 0.3–0.7 mm long, 1.3–1.6(2.5, 4.5) mm diam., similar to style type B; style crown truncate at apex with lobed depression; style apex rounded; stigma button-like, unlobed or weakly lobed, truncate, 1.1–2.2 mm diam., 0.5–1.0 mm high, covering entire style apex, sometimes depressed medially; the androecium truncate, margins irregularly 4–6-sided; thecae oblong to cylindrical, 0.4 mm wide,  $\pm$  parallel to one another,  $\pm$  contiguous; sterile staminate flowers acutely to bluntly 4–6-sided, 2.1–2.4 mm long, 1.1–2 mm wide. **INFLORESCENCE** with spathe green outside, orange-brown inside; berries white, 7.2–10 mm long, 3.6 mm diam.; seeds more than 10 per locule, pale yellow. **JUVENILE** plants terrestrial or epiphytic; internodes short, brown, weakly glossy; petioles 29.2 cm long,  $\pm$  terete, moderately spongy; blades 28.6–37.7 cm long, 17–28 cm wide; midrib broadly convex above; minor veins distinct below.

Flowering in *Philodendron llanense* occurs at the beginning of the rainy season, in July. Post-anthesis collections exist from January through August, with immature fruits and mature fruits found only in the dry season and early rainy season, mostly January and March but also in June.

*Philodendron llanense* is endemic to Panama (though to be expected in adjacent Colombia), ranging from 250 to 800 (mostly below 500) m elevation in *Tropical wet forest* and *Premontane rain forest* life zones. This species appears restricted to the region of the Cerro Jefe and the El Llano-Cartí Road (hence the name "llanense").

*Philodendron llanense* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is characterized by its short, thick internodes; mostly unribbed, semi-intact, or fibrous cataphylls; obtusely flattened to D-shaped petioles (about as long as the blades); thick, broadly ovate-cordate blades with a deeply spatulate to closed sinus and with prominently raised minor veins upon drying; and two to three green inflorescences with the spathe tube dark reddish violet inside.

*Philodendron llanense* is perhaps most easily confused with *P. schottianum*, which occurs at

mostly higher elevations (generally 1000 to 1600 m) in Costa Rica, rarely to as low as 490 m or as high as 2200 m elevation. It differs from *P. llanense* in having sharply two-ribbed cataphylls persisting more intact and with a prominent yellow epidermis, and blades usually drying with the lower surface more yellowish with the secretory ducts more outstanding and usually with partial "cross-veins" between the minor veins. In addition, *P. schottianum* also has blades proportionately somewhat longer (averaging 1.65 times longer than broad vs. 1.4 times longer than broad for *P. llanense*) and spathes shorter and nearly elliptic with almost no constriction (vs. oblong and noticeably constricted above the tube for *P. llanense*).

*Philodendron llanense* looks superficially much like *P. ferrugineum* in live condition, and since they occur together, they can be confused. However, *P. ferrugineum* differs in having promptly deciduous cataphylls and somewhat more elongate blades, which dry conspicuously reddish brown.

A single sterile collection (Croat & Grayum 60209) from the Atlantic slope near the Continental Divide at 590 m elevation in Bocas del Toro Province may also belong to this species. In addition to being out of range, this collection has the major veins much darker than the surface below.

*Additional specimens examined.* PANAMA. **Bocas del Toro:** Fortuna-Chiriqui Grande, 4.3 km N of the Continental Divide, 590 m, 8°46'N, 82°14'W, *Croat & Grayum 60209* (MO). **Panamá:** El Llano-Carti Road, 10-12 km from El Llano, 400 m, *Moas & Dressler 1709* (U); Km 9.5 of Pan-American Highway, 200-300 m, 9°15'N, 79°W, *McPherson 10822* (B, MO); Km 12, 350 m, *Mori et al. 4628* (MO); Mile 7, 460 m, 9°19'N, 79°59'W, *Croat 75104* (CAS, MEXU, MO, NY); Km 8-12, ca. 400-450 m, *Nee et al. 8775* (MO, NY, SCZ); Mile 5-9, 200-250 m, 9°15'-16'N, 78°59'W, *Thompson 4621* (CM); ca. Km 16-18, 400 m, *Tyson & Nee 7361* (MO, PMA, US); Mile 10.1, 325-350 m, *Croat 67367* (DUKE, K, M, MEXU, MO, NY, US); Mile 12, 200-500 m, *Croat 22910* (MO, PMA); Km 7-12, 360-400 m, *Croat 25118* (MO), *25172* (MO); Mile 10, 330 m, *Croat 33774* (MO), *33819* (MO); Mile 6.8, 350 m, *Croat 49127* (MO); Km 12, *Croat 26031* (MO); Cerro Jefe, near summit, <2900 ft., *Gentry et al. 3501* (MO, NY); 12 km E of Lago Azul, 800-1000 m, *Gentry & Mori 13450* (MO). **San Blas:** El Llano-Carti Road, 300-500 m, *Liesner 1317* (F, MO, NY, US); Mile 14, 300 m, 9°15'N, 79°W, *Croat 69251* (F, MO); Mile 7, 550 m, 9°43'N, 78°68'W, *60505* (CM, MO, PMA); Km 22, 350 m, 9°19'N, 78°55'W, *de Nevers & Herrera 7859* (MO, PMA); Nusagandi, Sendero Wedar, 300-400 m, 9°18'N, 78°58'W, *McDonagh et al. 216* (BM, MO); 1-2 mi. N of Nusagandi on road to Carti, 250-275 m, 9°20'N, 79°W, *Croat & Zhu 76577A* (CM, MO); Mile 10.1, 300 m, 9°20'N, 79°W, *76541* (MO, SEL); 1.9 mi. N of Nusagandi, 310 m, *76993* (AAU, CR, GB, MO, NY).

### *Philodendron madronense* Croat, sp. nov.

TYPE: Panama. Panamá: Valle de Madroño, ca. 10 road mi. N of La Margarita (near Chepo), just S of Continental Divide along main trail to Cangandí, 350-500 m, 9°19'N, 79°08'W, 21 Feb. 1986, *Hammel & McPherson 14526* (holotype, MO-3490432; isotypes, CM, M, PMA). Figures 279, 280.

Planta hemiepiphytica; internodia usque 2.5 cm diam., longiora quam lata; cataphylla decidua; petiolus subteres, usque 68 cm longus, 1.5 cm diam., aliquantum spongiosus; lamina trisectus, 39.5 cm longis, 15 cm latis; segmentis medianis ellipticis; segmentis lateralibus valde inequaliter, 32-32.8 cm longis, 12.7-13 cm latis; inflorescentia 1. Inflorescentia 1; pedunculus usque 22 cm longus, 1 cm diam., in sicco demigratus; spatha usque 20 cm longa, ca. 4 cm diam.; spadix pistillata 11.8 cm longa, ca. 3 cm diam.; spadix staminata ca. 7.5 cm longa, ca. 1 cm diam.; 1 semen in quoque loculo.

Hemiepiphytic vine; internodes to 2.5 cm diam., longer than broad, epidermis drying light brown and semiglossy but conspicuously and irregularly folded; roots several per node, drying dark brown; cataphylls deciduous; **petioles** to 68 cm long, 1.5 cm diam., subterete, somewhat spongy, drying blackened, surface closely and finely ridged; sheathing to 8.5 cm long; **blades** trisect, subcoriaceous, moderately bicolorous, drying weakly glossy, upper surface dark green, drying dark brown, semiglossy, lower surface drying yellow-green, matte; median segments elliptic, somewhat inequilateral, 39.5 cm long, 15 cm wide (ca. 2.5 times longer than wide), (1.5 times longer than lateral lobes), somewhat acuminate at apex, base attenuate and somewhat inequilateral with leafy tissue extending to base on one side, ending 2.5 cm above the base of the other side; lateral segments markedly inequilateral, 32-32.8 cm long, 12.7-13 cm wide, acuminate; the outermost lobes with the inner margins attenuate, tissue ending 1-2.5 cm from base of petioles; outer margins rounded, barely or not at all naked along the sinus; midrib weakly raised, concolorous above, ± round-raised, drying blackened and irregularly ridged below; basal veins lacking; primary lateral veins ca. 25 per side, ca. 4-5 mm apart, departing midrib at a 65-70° angle toward apex, 85-100° angle toward base, gradually curved to the margins, weakly sunken, scarcely more conspicuous than interprimaries above, prominently raised below; interprimary veins numerous, 1-3 between each pair of primary lateral veins; minor veins in part undulate upon drying, close, moderately visible, arising from both the midrib and primary lateral veins. **INFLORESCENCES** (post-anthesis) 1 per axil; peduncle to 22 cm long, 1 cm diam., drying blackened, somewhat spongy; **spathe**

to 20 cm long, ca. 4 cm diam., semiglossy, green outside, red at base inside, drying blackened; pistillate spadix 11.8 cm long, ca. 3 cm diam.; staminate spadix 7.5 cm long, ca. 1 cm diam., moderately tapered to a bluntly acute apex; pistils  $\pm$  cylindrical, 3–4 mm long, 2.5 mm diam.; ovary 6–7-locular, with sub-basal placentation; locules 2.3 mm long, 0.5 mm diam.; ovules 1 per locule, contained within a transparent envelope; funicle ca. 0.4 mm long (can be pulled free to base), style similar to style type D; style apex flat; style boss pronounced but narrow; seeds 1 per locule, 1.4 mm long.

Flowering in *Philodendron madronense* apparently occurs in the rainy season, since immature fruits have been collected in January.

*Philodendron madronense* is endemic to central Panama along the border of Panamá and San Blas at 350 to 450 m elevation in a *Tropical wet forest* life zone.

*Philodendron madronense* is a member of *P.* sect. *Tritomophyllum*. This species is recognized by its scandent habit; elongate internodes; subterete, more or less spongy petioles; and especially by its trisect leaf blades with the medial segments elliptic, inequilaterally attenuate at the base, and with many close prominent primary lateral veins only 4–5 mm apart and departing the midrib at about a 90° angle.

*Philodendron madronense* is superficially most similar to *P. cotobrusense*, which has deeply 3-lobed blades with more or less elliptic, closely veined divisions. The latter species differs in having the lobes united and confluent at the base and up to five inflorescences per axil. It is also similar to some broad-leaved forms of *P. tripartitum* (e.g., *Whiteford & Eddy* 223), which occur in eastern Panama, but those differ in having no more than 12 pairs of primary lateral veins, which depart the midrib at a 50–60° angle.

***Philodendron malesevichiae* Croat, sp. nov.**

TYPE: Panama, Coclé: vicinity El Valle de Antón, La Mesa, 4 mi. E of El Valle at base of Cerro Gaital, along trail which goes to the S edge and leading to the summit, 830–900 m, 8°36'N, 80°07'W, 25 Mar. 1993, *Croat 74818* (holotype, MO-4342656–57; isotypes, B, COL, CR, F, K, MEXU, NY, PMA, QCNE, US, VEN). Figures 3, 21, 25, 27, 28, 281–284.

Planta terrestris; caulis repens; internodia 2–4 cm longa, 1.5–5 cm diam.; cataphylla 21–29 cm longa, leniter 1–2-costata, in sicco rubrobrunnea, persistentia semi-intacta; petiolus teres, 50–56 cm longus, 8–19 mm diam.,

anguste sulcatus, sparsim squamatus in triente superiore; lamina ovato-cordata, 33–65 cm longa, 23–36 cm lata, in sicco atribrunnea supra, flavibrunnea infra, nervis basilibus 6–9 paribus, superioribus 2–3 liberis ad aliquot basin; reliquiarum coarctatis 2–4.5 cm; inflorescentia 1–3; peduncululus 4.5–8 cm longus; spatula 8–20 cm longa; pistilla 5–6-locularia; locali ca. 19–25-ovulata.

Terrestrial, usually less than 1 m tall, sometimes to 1.5 m, reclining against trees; stem creeping over the ground but well rooted in the soil; internodes 2–4 cm long, 1.5–5 cm diam., pale green to whitish, moderately glossy, usually totally hidden by cataphylls, coarsely but faintly white-short-lineate at apex, drying pale brown, epidermis closely and mostly bluntly ridged with ridges close and cracked transversely; roots moderately few per node, drying 2–3 mm diam., dark brown, closely ridged and weakly scaly; cataphylla 21–29 cm long, unribbed, weakly 1-ribbed or weakly 2-ribbed, whitish to maroon or russet, heavily tinged pink toward lower half, drying dark brown to reddish brown, persisting semi-intact at least toward apex with an underlying network of pale, anastomosing fibers, the outer surface becoming fibrous toward the base. LEAVES erect to erect-spreading; petioles 50–56 cm long, 8–19 mm diam., terete or subterete, dark green, tinged purple-violet, weakly glossy, drying dark brown, weakly, obtusely and narrowly sulcate to obtusely flattened adaxially, weakly flattened and often tinged purplish toward apex, faintly striate on surface, sparsely scaly in upper one-third to one-half, scales fine,  $\pm$  terete, 1–2 mm long, 0.1 mm diam., green; blades ovate-cordate, subcoriaceous, moderately bicolorous, acuminate to long-acuminate at apex, prominently cordate at base, 33–65 cm long, 23–36 cm wide, upper surface subvelvety-matte, dark green, drying dark brown, lower surface semiglossy, moderately paler, drying medium yellow-brown; anterior lobe 23–48 cm long, 23–36 cm wide; posterior lobes narrowly rounded, 8–25 cm long, 12–14 cm wide; sinus hippocrepiform, 8–14 cm deep; midrib concave, concolorous or paler above, convex or thicker than broad, much darker olive-green and matte, sometimes tinged maroon near base below, drying darker than surface; basal veins 6–9 per side, with upper 2–3 free to base, part of the remainder coalesced 2–4.5 cm; posterior rib naked for 0.5–3 cm, moderately straight; primary lateral veins (3)5–10 per side, 1.7–2.2 cm apart, departing midrib at a 45–60° angle, downturned acutely at the midrib, weakly sunken to weakly quilted above, convex and darker below, often branching toward the margins, drying blackened; interprimary veins persistent in lower half of blades; minor veins moderately visible but not dis-

tinct when fresh, moderately distinct on drying, weakly undulate, arising from both the midrib and primary lateral veins. INFLORESCENCES 1–3 per axil; peduncle 4.5–8 cm long, to 4 mm diam., pinkish red, coarsely white-lineate toward apex, drying blackened; **spathe** 8–20 cm long, 1–2 cm diam. when furled, to 5.5 cm wide when open (7–11 cm wide when fully flattened), ((2.5)3–3.7 times longer than peduncles), acuminate at apex (acumen 2.5 cm long), semiglossy, in Central America pale yellow-green, darker green toward middle, white along margins, sometimes tinged with pink, the open margin sometimes violet-purple outside, greenish white and matte throughout within, in South America spathe tube sometimes reddish outside, dark red to red-violet, suffused onto lower one-half of blade inside; spathe blade to 15 cm long, white within; spathe tube 7.5–8.5 cm long, to 4.3 cm diam. at anthesis; **spadix** weakly stipitate; 14–15.5 cm long; pistillate spadix (3.4)4–4.3 cm long in front, 2.8–4 cm long in back, 11–16 mm diam. at apex, 1.2–1.8 cm diam. at middle and base; staminate portion 10–13 cm long, 1.2–1.5 cm diam. at base, 1.5–1.8 mm diam. midway, 8–12 mm diam. 1 cm from apex, constricted to 1.5 cm diam. above the sterile portion; sterile staminate portion to 3.4 cm long, 20 cm diam. at base, 10–18 mm diam. at apex; pistils 1.1–2.4(3.5) mm long, 1.2–5 mm diam.; ovary 5–6-locular, with axile placentation; locules 1.3–1.6 mm long; ovules 19–25 per locule, 2–3-seriate, ca. (0.1)0.25–0.3 mm long; funicle as long as or shorter than ovules,  $\frac{1}{2}$ – $\frac{3}{5}$  as long as the ovule; stigma 1–1.5 mm diam., depressed medially; style similar to style type B, 1 mm diam.; style apex flat to weakly concave; styler canals arising at base of apical depressions; stigma discoid, 0.8 mm thick, 1.5–1.8 mm diam.; the androecium truncate to  $\pm$  prismatic, margins irregularly 4–6-sided, many 6-sided, ca. 0.6 mm long; thecae oblong,  $\pm$  parallel to one another; sterile staminate flowers clavate, irregularly 4–5-sided to ovoid, 0.9–1.3 mm long.

Flowering phenology in *Philodendron malesevichiae* in Panama is uncertain. All flowering collections are from South America. Flowering collections have been made in February and March and post-anthesis collections in March, July, and December. Cultivated collections from Panama grown at the Missouri Botanical Garden flowered in May, June, July, November, and December.

*Philodendron malesevichiae* ranges from Panama to Colombia. In Panama it is only known from the type locality in *Tropical wet forest* at 830 to 860 m in Coclé Province. In Colombia it has been col-

lected only along the Pacific Coast in Chocó and Valle at 50 to 150 m elevation.

*Philodendron malesevichiae* is a member of *P.* sect. *Philodendron* subsect. *Achyropodium*. This species is characterized by its terrestrial, creeping habit; short internodes; weakly 1–2-ribbed cataphylls drying reddish brown and persisting semi-intact; subterete, narrowly sulcate petioles, which are conspicuously scaly in the upper one-third; and ovate-cordate brown-drying blades with up to eight basal veins, largely coalesced and naked along the hippocrepiform sinus.

This species is most easily confused with *P. glanduliferum*. The latter species differs in having fewer primary lateral veins (2–4 vs. 6–10 for *P. malesevichiae*); posterior ribs that are not at all naked; and a narrow closed or spatulate sinus (vs. hippocrepiform in *P. malesevichiae*).

In Panama there are three other species that have petiolar glands of some form and thus might be confused with *P. malesevichiae*. These are: *P. hammelii*, *P. verrucosum*, and *P. squamipetiolatum*. *Philodendron hammelii* differs in its smaller, green-drying blades and petiolar scales mostly less than three times longer than broad. *Philodendron verrucosum* differs in its mostly appressed-climbing habit; scaly cataphylls, inflorescences, and even parts of the lower blade surface; and velvety and matte (rather than semiglossy) upper blade surface. *Philodendron squamipetiolatum* differs in its appressed-climbing habit, long internodes, deciduous cataphylls, green-drying blades, and scaly inflorescences.

*Zarucchi & Escheverry 4776*, from 2000 m elevation in Antioquia Department, Colombia, may also belong to this species. It is described as having a deep wine-red (rather than green) spathe.

This species is named in honor of Petra S. Malesevich, who has loyally worked with me on all aspects of the *Philodendron* revision. This species is in cultivation at the Missouri Botanical Garden and is deemed a beautiful addition to horticulture.

*Additional specimens examined.* PANAMA. Coclé: La Mesa, above El Valle de Antón, 860–900 m, *Croat 37319* (MO); base of Cerro Gatital, 860 m, 8°37'N, 80°08'W, *Croat & Zhu 76707* (CAS, MO, PMA).

COLOMBIA. Chocó: hills above junction of Río Capá and Río Mumbá, up river from Lloré, 80–120 m, 5°37'N, 76°25'W, *Juncosa 1448* (MO). Valle: Buenaventura, Bajo Calima region, Buenaventura–Málaga, Km 65–66, 40–65 m, 4°10'N, 77°12'W, *Croat 71057* (COL, MO); 100 m, 4°4'N, 77°09'W, *70162* (CAS, COL, L, MEXU, MO, NY, PMA); Km 49, 150 m, 4°02'N, 77°04'W, *Croat & Boy 75810* (B, CM, COL, F, GB, M, MEXU, MO, PMA, US); Buenaventura–Río Calima, 33.3 km beyond main Cali–Buenaventura Highway, 50 m, 4°02'N, 77°07'W, *Croat 61377* (AAU, COL, MO, PMA, TEX, US); near Km 14

marker, <50 m, 3°56'N, 76°59'W, Croat 57547 (CM, COL, G, JAUM, MO).

***Philodendron mexicanum*** Engl., in Mart., Fl. Bras. 3(2): 143. 1878. TYPE: Mexico. Veracruz: vic. of Córdoba, *Bourgeau 2176* (holotype, P; isotype, G). Figures 285–288.

*Philodendron latisagittum* Matuda, Anales Inst. Biol. Univ. Nac. México 22: 371, Fig. 2. 1951. TYPE: Mexico. Chiapas: Mpio. Siltepec, Cascada, mixed humid forest, over humid rocks, 1200 m, 4 Mar. 1951, *Nakamura 31* (holotype, MEXU).

Usually hemiepiphytic, sometimes terrestrial or epilithic; stem scandent, leaf scars conspicuous, 1–1.5 cm long, 1.3–1.6 cm wide; internodes scurfy, glaucous to semiglossy, 18–21 cm long, 1–2 cm diam., longer than broad, medium green to gray-green, epidermis blistering, fissured  $\pm$  transversely; roots branched at tips, sometimes with swollen nodes along length; cataphylls fleshy, 10–23 cm long, unribbed or bluntly 1-ribbed, pale green, glossy, drying yellowish tan to yellowish green, deciduous intact; **petioles** 22–66.5 cm long, 2–13 mm diam., terete, moderately spongy, medium green, somewhat flattened adaxially, surface semiglossy; **blades** narrowly triangular-sagittate to triangular-hastate, subcoriaceous, acuminate to long-acuminate at apex, sagittate to hastate at base, 23–46 cm long, 14–38 cm wide (1.3–2.7 times longer than wide), (0.6–1.5 times longer than petiole), about equal in length to petiole, margins weakly undulate, upper surface dark green, drying dark brownish green, semiglossy, lower surface drying yellowish green, weakly glossy, paler; medial lobe 19–38 cm long, 8–20 cm wide (1.7–2.7 times longer than posterior lobes), usually 3–3.5 times longer than wide (rarely to 1.7 times longer than wide); posterior lobes 7–19.5 cm long, 4–12.6 cm wide, directed somewhat toward the base, rounded to rarely rounded; sinus parabolic to hippocrepiform or spatulate; midrib broadly sunken, concave above, convex, sparsely orange-spotted, slightly paler than surface below; basal veins 0–1(2–5) per side, with 0–1(2–5) free to base, or 1 coalesced, the third and fourth coalesced 3.6–4(13) cm; posterior rib not naked or naked for 0.5–2 cm, and directed straight toward the tip of the blade and remaining 1.5–3.5 cm distant from blade margin; primary lateral veins (2)4–5(6) per side, departing midrib at a 55–65° angle,  $\pm$  straight to the margins, weakly sunken above, raised below; minor veins moderately distinct below, arising from both the midrib and primary lateral veins. INFLORESCENCES 1 per axil; peduncle 4–15 cm long, 4–12 mm diam.; **spathe** glossy, 8–16.5 cm long, 1.9–2.3

cm diam. (0.8–1.4(1.5–2.4) times longer than peduncle), acute at apex, the margins paler to clear within; spathe blade greenish to whitish outside; resin canals pale range and appearing as continuous lines inside; spathe tube greenish, sometimes pale reddish tinged outside; red-violet to maroon (B & K red-purple 5/7.5) inside; **spadix** sessile; bluntly rounded to somewhat acute at apex, 10.5–14.7 cm long, broadest toward the apex, constricted below the middle of fertile staminate portion; pistillate portion pale green to green to pale yellow, cylindrical to obovoid, 2–6 cm long, 1 cm diam. at apex, 1.1 cm diam. at middle, 1.3 cm wide at base; staminate portion (6.4)10.3–12.7 cm long; fertile staminate portion creamy white,  $\pm$  cylindrical, 6–17 mm diam. at base, 7–15 mm diam. at middle, 1.1 cm diam. ca. 1 cm from apex, about as broad as the pistillate and the sterile portions; sterile staminate portion usually broader than the pistillate portion, white to light gray, 1–1.2 cm diam.; pistils 1.6–2(3.5) mm long, 1.1–1.3(2.6) mm diam., transparent white; ovary 5–7-locular, 0.9–2.7 mm long, 1.2–1.4(2.3–2.6) mm diam., with sub-basal placentation; locules 0.9–1.1(2.5) mm long, 0.2–0.4(0.6–0.7) mm diam.; ovule sac 0.6–0.7 mm long; ovules 1–2(3) per locule, digitate, contained within transparent ovule sac, 0.3–0.6(1.1) mm long, longer than funicle; funicle 0.3 mm long, style 0.7(1) mm long, 1.2–1.5(2.4) mm diam., similar to style type B; style apex domed; stigma discoid, at least sometimes lobed both laterally and vertically, sometimes  $\pm$  cylindrical, 0.7–1.0 mm diam., 0.1–0.3 mm high, covering center of style apex, at least sometimes drying with radial arms sunken between the central peak and the peaks on the end of the arms (Croat & Hannon 64520); the androecium truncate, margins irregularly 4–6-sided; thecae  $\pm$  cylindrical, 0.3–0.4 mm wide,  $\pm$  parallel to one another; sterile staminate flowers bluntly or acutely 4–6-sided, 1.3–1.9 mm long, 1.3–1.8 mm wide. INFLORESCENCE with seeds 1 per locule, yellowish orange, 1.5–1.9 mm long, 0.6–0.8 mm diam.

Flowering in *Philodendron mexicanum* occurs in the mid-dry season and early rainy season (February through May), with post-anthesis collections known February through June (except May) and also in November. Mature fruits are not known. A cultivated collection at Missouri Botanical Garden (Croat 59933) flowered perhaps twice per year, in March and in October.

*Philodendron mexicanum* ranges from Mexico to Colombia, from near sea level to 1900 m elevation, ranging from dry habitats in west-central Mexico (both "Selva Baja Caducifolia" and "Bosque Pino-

Encino") to more humid sites on the Atlantic slope in Vera Cruz to as far south as Honduras in *Tropical moist forest* and to Costa Rica in *Premontane wet forest*. Though widespread, this species is apparently rare and has been collected from relatively few localities, often restricted to mesic enclaves in otherwise arid regions as in the state of Morelos, Mexico. It is one of the most ecologically versatile species in the genus.

*Philodendron mexicanum* is a member of *P.* sect. *Calostigma* subsect. *Macrobilium* ser. *Macrobilium*. This species is recognized by its scandent habit; internodes longer than broad; unribbed, deciduous cataphylls; moderately spongy, somewhat flattened petioles (about as long as the blades); and especially by its narrowly triangular-sagittate to triangular-hastate blades, which dry dark brownish green above and yellowish green below. Sterile collections from Los Ríos and Guayas Provinces of Ecuador (*Dodson & Valverde 6959*) at Jaunche and in Guayaquil Cantón (*Rubio et al. 2008*) may also represent this species. Dodson et al. (1985) erroneously reported the former collection as *P. barrosoanum* G. S. Bunting, a species restricted to the eastern side of the Andes.

*Philodendron mexicanum* is most similar to *P. angustilobum*, a species ranging from Honduras to Panama. The latter species differs in having leaves that dry more blackened (rather than green to yellow-green as in *P. mexicanum*), and which are more decidedly three-lobed with the medial lobe proportionately narrower and broadest at the middle or even above the middle.

Although the type specimen of *P. mexicanum* was collected in Mexico near Córdoba, most Mexican collections are from the Pacific slope. These have somewhat less prominently narrowed posterior lobes than the type. *Moore & Bunting 8873*, from near the type locality, has the posterior lobes even more conspicuously narrowed than those of the type specimen. In this regard, its blades approach those of *P. angustilobum* in overall shape, but differ in drying greenish to yellowish brown rather than blackened.

A collection from Olancho Department, Honduras (*Croat & Hannon 64520*), is unusual in having a style that dries with radiating arms from a central peak.

The sole Costa Rican collection (*Grayum 5418*) is unusual in having prominently hastate blades. This collection has only a juvenile inflorescence. Further collections may prove it represents a distinct species.

*Additional specimens examined.* COSTA RICA. **Puntarenas:** Río Guacimal downstream from Monteverde, Cordillera de Tilarán, 1300 m, 10°18'N, 84°49'W, *Grayum*

*5418* (MO). GUATEMALA. **Quezaltenango:** Finca Pi-reneos-Patzún, 1200–1400 m, *Standley 86917* (F); *87007* (F); *87022* (F). **San Marcos:** Volcán Tajumulco, 1300–1500 m, *Steyermark 37968* (F). HONDURAS. **Atlántida:** Quebrada Grande, ca. 10 km SW of La Ceiba, 80–180 m, 15°42'N, 86°51'W, *Liesner 26138* (MO). **Olancho:** San Esteban-Bonito Oriental, Río Grande, 350–400 m, 15°31'N, 85°42'W, *Croat & Hannon 64520* (B, CAS, CM, CR, HNMN, K, L, MEXU, MO, NY, USCC). **Yoro:** Puente Grande, Río Puente Grande (tributary of the Río Agua), *Blackmore & Chorley 4077* (BM, MO). MEXICO. Cultivated at Cornell University, Ithaca, New York, *Moore 7437* (BH); Ixtapan de La Sal, 1900 m, 18°50'N, 99°41'W, *Matuda et al. 32130* (MEXU). **Guerrero:** Atzac region, above Fila de Caballo, El Paraíso in Parque Nacional de Guerrero, *Croat 67442* (MO). **Morelos:** Cuernavaca, *Boegeus 1420* (K, P); 1350 m, *Quarles van Ufford 95* (U); 5000 ft., *Pringle 8093* (BH, BM, BR, CM, F, G, GH, H, HBG, ISC, K, LL, MASS, MEXU, MO, NY, P, POM, RSA, UC, US); *Rose & Hough 4439* (US); Río Pollo, below Salto San Antonio, Fraccionamiento San Antonio, W of Colonia Carolina, NW of Center of Cuernavaca, 1500 m, 18°57'N, 99°15'W, *Croat & Hannon 65778* (B, CM, F, GH, K, MEXU, MO, NY, US); *Matuda 26030* (MEXU); *25982* (F, MEXU); *Moore & Bunting 8820* (BH); Barranca Santa Clara, N de Acatilpa, 1450–1550 m, *Vázquez 3094* (MEXU). **Veracruz:** Córdoba–Veracruz, Ejido San José de Gracia, below Peñuelo, ca. 730? m, *Moore & Bunting 8873* (BH, MO).

***Philodendron microstictum* Standl. & L. O. Williams, Ceiba 3: 108–109, 1952. TYPE:** Costa Rica. Puntarenas: Esquinas Forest Reserve, sea level, 10 Jan. 1951, *Allen 5755* (holotype, EAP; isotypes, F, GH, US). Figures 289, 290.

Hemiepiphytic vine, stem scandent, green to gray-green, drying pale yellowish brown, unscented, leaf scars inconspicuous, 1.2 cm long, 8 mm wide, obscured by cataphylls; internodes smooth, semiglossy, 6–12 cm long, 1–2 cm diam., longer than broad, moderately green, drying khaki-colored, epidermis fissured transversely; roots drying light reddish brown, smooth, weakly glossy, 20–30 cm long, 2–3 mm diam., 4–6, at the nodes; cataphylls subcoriaceous, bluntly 2-ribbed or unribbed, pale green to cream-colored, promptly deciduous; **petioles** 12–24 cm long, 2–4 mm diam., subterete to broader than thick to broadly D-shaped, weakly spongy, bluntly flattened to broadly sulcate adaxially, rounded abaxially, with adaxial margins blunt, surface pale or dark green streaked and demarcated from blade by dark green ring around apex; **blades** broadly ovate, moderately coriaceous, abruptly acuminate at apex (the acumen tightly inrolled, 1–4 mm long), weakly subcordate at base, 12.5–30 cm long, 13–21.5 cm wide (0.7–1(1.7) times longer than wide), (0.65–2 times longer than petiole), about equal in length to petiole, broadest ± near the middle, margins straight, upper surface dark green, semiglossy, somewhat pruinose, lower sur-

face glossy to weakly glossy, paler; anterior lobe 12–20.5 cm long, 10.7–22 cm wide (1.85–3 times longer than posterior lobes); posterior lobes 4–8 cm long, 3.5–9.5 cm wide, broadly rounded to obtuse; sinus arcuate, sometimes parabolic, 3.4–6 cm deep; midrib broadly convex to weakly raised, concolorous to paler than surface above, weakly to broadly convex, paler than surface below; basal veins about 3 per side, obscurely and scarcely more conspicuous than primary laterals; posterior rib lacking; primary lateral veins (3)4–5 per side, obscure above; minor veins about as conspicuous as the primary laterals, arising from the midrib only. INFLORESCENCES spreading, as long as leaves, 1 per axil; peduncle 14–25 cm long, 2–6 mm diam., subterete, green; spathe 6.5–15 cm long (0.3–1 times longer than peduncle); spathe blade green outside, pale green inside; spathe tube ca. 5 cm long, green outside, red or violet-purple inside; spadix sessile, 8 cm long, broadest below the middle; pistillate portion medium green to pale greenish yellow,  $\pm$  cylindrical, 1.9–2 cm long, 6.5–7 mm diam. at apex, 7–8 mm diam. at middle, 6.5–7 mm wide at base, with 13 (per cm) flowers per spiral; staminate portion 5.9–8.3 cm long; fertile staminate portion white, tapered toward apex, 5–9 mm diam. at base, 5–6 mm diam. at middle, 0.9 mm diam. ca. 1 cm from apex, broadest at the base, mostly narrower than the pistillate and sterile portions; sterile staminate portion broader than the pistillate portion, 6–9 mm diam.; pistils 1.2–1.4 mm long, 0.8–1.1 mm diam.; ovary 6–7-locular, 0.8–1 mm long, 0.8–1.1 mm diam., with sub-basal placentation, walls drying weakly warty; locules 0.8 mm long, 0.3 mm diam., ovule sac 0.7–0.8 mm long; ovules 1 per locule, contained within transparent ovule sac, 0.4 mm long, as long as funicle; funicle 0.4 mm long; style 0.2 mm long, 1 mm diam., similar to style type B; style apex flat; stigma unlobed, subdiscoid to somewhat cylindrical, 0.8–0.9 mm diam., 0.2 mm high, covering almost entire style apex, centered on stylar canal pores; the androecium truncate, margins irregularly to bluntly 4–6-sided, 1.2 mm long, 0.8 mm diam. at apex; thecae  $\pm$  cylindrical, 0.3 mm wide,  $\pm$  parallel to one another, contiguous; sterile staminate flowers acutely or bluntly and irregularly 4–6-sided, 1.2–2 mm long, 0.7–1.1 mm wide. INFRUDESCENCE with fruits whitish. Seeds (dried) ca. 20 per locule, tan, 1–1.2 mm long, 0.5–0.6 mm diam., finely ridged with much finer cross-etching.

Flowering in *Philodendron microstictum* occurs during the dry season and early rainy season, with flowering collections known from January, April, and May, and post-anthesis collections from Feb-

ruary, March, May, and July. Immature fruits have been collected in May.

*Philodendron microstictum* is endemic to Costa Rica (though to be expected on the Burica Peninsula in adjacent Panama), where it is restricted to the Pacific slope, at 50 to 450 m elevation in wetter parts of *Tropical moist forest*, *Tropical wet forest*, and *Tropical wet forest basal belt transition* life zones.

*Philodendron microstictum* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Ovata*. This species is characterized by its scandent habit; long, moderately slender stems drying pale yellowish brown; subterete petioles, which are about as long as the blades; and broadly ovate, weakly subcordate blades (which may be as wide or wider than long) with an arcuate sinus and obscure primary lateral and basal veins (sometimes with only the basal veins visible). Also characteristic is the externally green spathe (red within the tube) with the peduncle as long as or longer than the petiole.

*Philodendron microstictum* is not easily confused with any other species. It has been confused by some with *P. scandens*, which differs in having more narrowly ovate blades with prominent major veins and a spatulate to parabolic sinus, and much more short-pedunculate inflorescences (with the peduncles much shorter than the petioles).

This species is perhaps related to *P. chirripoense*, which is also a vine with inflorescences longer than leaves and has blades of similar color and texture. That species differs in having longer, more slender internodes that dry dark brown and by its narrowly ovate, subcordate blades, which are more than twice as long as broad and have more prominent primary lateral veins.

*Additional specimens examined.* COSTA RICA. **Puntarenas:** Palmar Norte, 100–200 m, *Croat 35107* (CR, MO); Palmar Norte, along trail to Jalisco, 50–700 m, *Croat 35189* (F, MO); 110 m, *Croat & Hannon 79210* (CR, INB, MO); Villa Neily–San Vito de Coto Brus, Cuesta Fila de Cal, 300 m, 8°41'N, 82°57'W, *Grayson et al. 7580* (CR, MO); El General Valley, Río Angol–Río Volcán, 450–500 m, *Williams et al. 24214* (CR, F); Osa Peninsula, 20–300 m, *Liesner 1836* (B, MO); ca. 5 km W of Rincón de Osa, 50–200 m, 8°42'N, 83°31'W, *Burger & Liesner 7300* (CR, F, MO, PMA); Fila Huacas, ca. 4 km NE of Las Huacas ('Venecia') along road to Sinat, 450–500 m, 8°52'N, 83°17'W, *Grayson & Hammel 10132* (CR, MO); Parque Nacional Corcovado, La Palma, 100 m, 8°35'N, 83°30'40"W, *Herreza 4066* (CR, L, MO, NY); Sirena, 0–150 m, 8°27'–30'N, 83°33'–38'W, *Kernan 463* (CR); in Monkey Woods (just W of airstrip), 5 m, 8°29'N, 83°35'W, *Hammel et al. 16643* (CR, MO); Claro Ridge, 1–10 m, 8°28'N, 83°35'W, *Kernan & Phillips 1028* (CR, MO); Sirena Woods, 1–50 m, 8°28'N, 83°35'W, *Kernan & Phillips 1089* (CR, MO); Río Sorpresa, ca. 1 km NE of Gólfito, E of microwave towers, ca. 400 m, 8°39'N, 83°10'W, *Croat & Grayson 59941* (CR, MO); Gólfito, Cerro Las Torres, 500

m, G. Herrera 5045 (CR, INB, MO); Reserva Forestal Golfo Dulce, W of Rancho Quemado, 1-300 m, 8°44'N, 83°36'W, Saborio *et al.* 127 (INB, MO); Río Volcán, 48 km SE of San Isidro General, 300 m, Molina *et al.* 18162 (F. GH, NY, US). **San José:** Río Negro, ZP La Cangreja, ca. 1.5 km E of Santa Rosa de Puriscal, 320 m, 9°42'N, 84°23'30"W, Grayson *et al.* 8340 (MO); Parque Nacional, sector Esquinas, vic. Fila Gamba, 200-300 m, Croat & Hannon 79288 (MO).

**Philodendron morii** Croat, sp. nov. TYPE: Panama. Panamá: Valle de Madroño, ca. 10 mi. N of La Margarita (near Chepo), in forest S of and on Continental Divide, near border of Comarca de San Blas, along trail to Cangandí, 350-450 m, 9°19'N, 79°08'W, 21 Feb. 1986, Hammel & McPherson 14530 (holotype, MO-3398570). Figures 291, 292.

Planta epiphytica; internodia 1-1.5 cm longa, 1.5-2 cm diam.; cataphylla 13 cm longa, obtuse 2-costata, decidua; petiolum subterete, 23-37 cm longus, (2)4-6 mm diam., subspongiosus; lamina ovato-triangularis, leniter cordata basi, 25-27.5 cm longa, 11.5-16 cm lata, in sicco canaviridis; inflorescentia 2; pedunculus 9.5-14 cm longus, 3-6 mm diam.; spathe 8.7-12 cm longa, viridis vel flaviviridis omnino; pistilla (3)4-5(6)-locularia; loculi cum 3-6 seminibus; baccae albae.

Epiphytic; stem appressed-climbing; internodes semiglossy, 1-1.5 cm long, 1.5-2 cm diam., about as long as broad, dark green, drying light brown; roots drying reddish brown; cataphylls 13 cm long, bluntly 2-ribbed, green, deciduous, intact; petioles 23-37 cm long, (2)4-6 mm diam., subterete, somewhat spongy, dark green, obscurely flattened adaxially, surface unmarked, often dries with loose, puffy epidermis; blades ovate-triangular, narrowly acuminate at apex, weakly cordate at base, 25-27.5 cm long, 11.5-16 cm wide (1.7-2.4 times longer than wide), (0.7-1.2 times longer than petiole), about equal in length to petiole, upper surface semiglossy, lower surface drying green, weakly glossy, moderately paler; anterior lobe 24-28 cm long, 12.6-16 cm wide; posterior lobes broadly rounded, 5-7 mm long, 3-7.5 cm wide, broadly rounded to obtuse; sinus arcuate with blade decurrent on petiole; midrib prominently raised above, slightly paler than surface below; basal veins 2-3 per side, with 0-1 free to base, 0-1 coalesced less than 1 cm; posterior rib weak, to 1.3 cm long, naked throughout its length; primary lateral veins (2)5-8 per side, departing midrib at a 40-50° angle, straight to the margins, sunken and concolorous above, convex and darker than surface below; minor veins fine, numerous, and distinct below, arising from both the midrib and primary lateral veins. INFLORESCENCES (post-anthesis) 2 per axil; peduncle 9.5-14 cm long, 3-6 mm diam.; spathe 8.7-12 cm

long, (0.8-1.1 times longer than peduncle), green to yellowish green throughout; spathe tube 3.5-5 cm long; spadix 8-11 cm long; pistillate portion (post anthesis) 5.7 cm long in front, 4.8 cm long in back, 1.5 cm diam. midway, 1.2 cm diam. near apex, 8 mm diam. near base; fertile staminate portion 5 cm long, narrowly tapered to apex, the narrowest portion to 5 mm diam., ca. 1.3 cm above base; sterile staminate portion broader than constricted area, to 5.5 mm diam.; pistils 1.4-1.6 mm long; ovary (3)4-5(6)-locular, with sub-basal placentation; locules 1.3 mm long, 0.5 mm diam.; ovules 1 per locule, 0.3 mm long; funicle 0.2-0.3 mm long (can be pulled free to base), style similar to style type D; style apex flat to weakly rounded, 0.6-0.7 mm long, style boss small; stigma covering entire style apex and inserted on style boss; the androecium truncate, margins irregularly 4-6-sided, 0.7 mm long. INFRUCTESCENCE with pistillate spadix 3-5 cm long; berries white, ± oblong-ellipsoid, 1.1-2 mm long, 0.4-0.8 mm diam.; seeds 3-6 per locule, tan.

Flowering phenology in *Philodendron morii* is poorly known, but it is perhaps bimodal with flowering collections known in March and November and immature fruits in December, February, and June.

*Philodendron morii* is endemic to Panama, known only from highlands east of the Canal Area in Panamá Province and Comarca de San Blas and from Cerro Pirre in Darién Province, at 450 to 850 m elevation in *Tropical wet forest* and *Premontane rain forest* life zones.

*Philodendron morii* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is characterized by its slender internodes (about as long as broad); bluntly two-ribbed, deciduous cataphylls; terete petioles (about as long as the blades), which often dry with loose, puffy epidermis; ovate-triangular, scarcely cordate, green-drying blades; and peduncles equaling or exceeding the green spathes.

*Philodendron morii* is most similar to *P. sphalerum* Schott from the Guianas, which has similar green-drying, long-petiolate leaves. The latter species differs in having the leaf blades acute to rounded or truncate at the base and proportionately longer petioles (fully as long as or much longer than the blades). In addition, *P. sphalerum* has up to four much smaller inflorescences with spathes 5.5-7 cm long, whereas *P. morii* has one to two much larger inflorescences per axil (spathes 9-12.5 cm long). *Philodendron morii* is also similar to *P. wilburii*, especially in terms of size and color of its leaf



blades. The latter species differs, however, in comprising more or less scandent plants with usually long internodes.

*Philodendron morii* is not easily confused with any other Central American species. It is named in honor of one of its earliest collectors, Scott Mori (NY), who collected for the Missouri Botanical Garden during 1974–1975.

*Additional specimens examined.* PANAMA. **Darién:** Parque Nacional Darién, W side of Cerro Pirre, 800–1050 m, 7°56'N, 77°45'W, Croat 68700B (MO). **Panamá:** Cerro Jefe region, Altos de Azul-Río Chagres, 700–850 m, 9°15'N, 79°30'W, McPherson 11899 (MO, PMA, US); 3–3.5 mi. NE of Altos de Pacora, 11.1–11.6 mi. beyond Lago Cerro Azul, 700–750 m, 9°15'N, 79°25'W, Croat 68693 (CM, MO); Campos Tres, 3 mi. NE of Altos de Pacora, 500–800 m, Liesner 567 (MO); Gorgas Memorial Labs "Campamento Cuatro," 5–10 km NE of Altos de Pacora, ca. 600 m, Mori & Kallunki 3441 (MO). **San Blas:** Río Diablo, Cordillera de Ibedón, 350–480 m, 9°21'N, 78°34'W, Herrera et al. 1489 (MO, PMA, US).

***Philodendron niqueanum* Croat, sp. nov. TYPE:**

Panama, Darién: Serranía de Pirre, along headwaters of Río Escucha Ruido, ca. 16 km due N of Alto de Nique, ca. 7°47'N, 77°45'W, 27 July 1976, Croat 37942 (holotype, MO-2416709). Figures 293–296.

Planta hemiepiphytica; internodia brevia, usque 5 cm diam.; cataphylla incostata vel leniter l-costata, rubra, persistentia plus minusve intacta; petiolus teres, (32)46–59 cm longus, in sicco 6–8 mm diam.; lamina ovata vel late ovata, (28)42–55.5 cm longa, (14)25–28.5 cm lata, circa tam longa quam petioli; sinus plus minusve V-formati; inflorescentia immatura; pedunculus 4 cm longus; spathe virides.

Hemiepiphytic; stem scandent when young, appressed-climbing when adult; internodes short, to 5 cm diam. (younger stems with internodes to 8 cm long, 1.5 cm diam.), epidermis moderately smooth, dark reddish brown; cataphylls unribbed to weakly 1-ribbed, red, persisting ± intact; **petioles** (32)46–59 cm long, 6–8 mm diam., terete; **blades** ovate to broadly ovate, acuminate at apex, truncate to subcordate at base, (28)42–55.5 cm long, (14)25–28.5 cm wide (1.7–2 times longer than wide), (ca. 0.9 times the petiole length), about equal in length to petiole, upper surface weakly glossy, drying dark brown, lower surface paler, drying dark yellowish brown; anterior lobe (27)37–49.5 cm long, 25–28.5 cm wide (4.7–5.4(13.5) times longer than posterior lobes); posterior lobes (2)7–9 cm long, 9–12 cm wide, broadly rounded to obtuse; sinus ± V-shaped, to 4 cm deep; midrib prominently raised, paler than surface above, raised below; basal veins 3 per side, with 0 free to base, some of the lowermost coalesced to ca. 1 cm; pos-

terior rib never naked; primary lateral veins 5–9 per side, departing midrib at a 45–65°(70°) angle, weakly arcuate to the margins, sunken above, raised below; minor veins obscurely visible, slightly raised on drying below, arising from the midrib only. INFLORESCENCES immature; peduncle 4 cm long; **spathe** green, 7 cm long; **spadix** immature.

Flowering in *Philodendron niqueanum* is poorly known owing to too few collections overall. The species was collected with flower buds in late July and probably both flowers and fruits within the rainy season (although, since it flowers so late, the fruits may mature in the dry season).

*Philodendron niqueanum* is endemic to Panama, known only from the type locality on the Serranía de Pirre, at 1530 to 1550 m elevation in *Tropical Lower Montane wet forest*.

*Philodendron niqueanum* is a member of *P.* sect. *Calostigma* subsect. *Macrobolium* ser. *Ecordata*. This species is characterized by its thick, short internodes; red, weakly one-ribbed cataphylls persisting mostly intact; terete petioles; and narrowly ovate, dark brown-drying blades about as long as the petioles.

*Philodendron niqueanum* is apparently close to *P. lentii*, which ranges from Costa Rica to central Panama, but no further east than the province of Coclé. Both species are similar in having truncate-to-subcordate-based leaf blades with more or less V-shaped sinuses. *Philodendron lentii* differs in having the primary lateral veins sunken, paler, and much more conspicuous on the upper dried blade surface. In contrast, the primary lateral veins of *P. niqueanum* are scarcely or not at all paler than the surface on dried leaves and are raised rather than sunken. In addition, the epidermal pattern is alveolate and moderately smooth at 10× magnification on the upper blade surface of *P. niqueanum*, whereas *P. lentii* lacks an alveolate pattern and the adaxial surface is densely covered with round, pale inclusions.

*Philodendron niqueanum* is named for the type locality near the Alto de Nique, hence the name.

*Additional specimen examined.* PANAMA. **Darién:** Cerro Pirre region, ca. 9 km from Alto de Nique, 1480–1520 m, Croat 37886 (MO, PMA, US).

***Philodendron panamense* K. Krause, in Engl. & K. Krause, Pflanzenr. IV. 23Db (Heft 60): 65. 1913. TYPE:** Panama. Canal Area: at Frijoles, 25–30 m, 9°10'N, 79°48'W, Pittier 3753 (holotype, US). Figures 297–299.

Usually hemiepiphytic; stem appressed-climb-

ing, to 1.3 m long, sap reddish, sticky; internodes short, semiglossy, 2.5–4 cm diam., sometimes longer than broad, dark green; roots several per node, drying 2–4 mm diam., dark brown, semiglossy, sparsely scaly; cataphylls 18–20 cm long, sharply 1-ribbed to sharply 2-ribbed, green to whitish, drying light brown, persisting briefly  $\pm$  intact, eventually fibrous, sometimes persisting for a time, eventually deciduous; **petioles** erect-spreading, (23)34–70(79) cm long, 4–13 mm diam., terete to subterete, sometimes weakly flattened or with narrow flattened rib adaxially, sometimes weakly and narrowly sulcate at base, dark green, sometimes pink at base, surface sparsely to densely pale greenish striate or striate-lineate, minutely grooved upon drying, geniculum to 6.5 cm long, sheath 1–4 cm long, usually inconspicuous, to 10 cm long when subtending an inflorescence; **blades** broadly triangular-ovate or more infrequently ovate, subcoriaceous, semiglossy, moderately bicolorous, abruptly acuminate, sometimes acute at apex (the acumens sometimes inrolled, to 4 mm long), deeply cordate at base, 32–72 cm long, 24–38 cm wide (1.2–2.3 times longer than wide, averaging 1.5), ((0.3)0.7–1.4 times longer than petiole, averaging 1.07), broadest near point of petiole attachment; upper surface dark green and glossy, drying semiglossy, dark gray-brown to olive-green, often somewhat blackened, lower surface slightly paler, semiglossy; margins sometimes broadly undulate; anterior lobe 25–41(58) cm long, (15)20–36(41) cm wide (1.9–4 times longer than posterior lobes); posterior lobes 7.5–18 cm long, 4.5–17.7 cm wide, rounded to broadly rounded to broadly obtuse; sinus hippocrepiform to parabolic (arcuate on younger blades), 4–12 cm deep; midrib broadly sunken, concolorous or paler than surface above, weakly asperous, thicker than broad, matte, sometimes short-white-striate, darker than surface below; basal veins 5–8 per side, with 1 free to base or nearly so, third and higher order veins coalesced 4–7 cm long; posterior rib naked for 2–3 cm long; primary lateral veins 4–7 per side, departing midrib at a 55–65° angle, spreading to a 65–75° angle, usually curved down gradually before merging with the midrib, narrowly sunken, concolorous or paler than surface above, convex, matte, slightly darker than surface below; interprimary veins narrowly sunken above; minor veins distinct, darker than surface below, arising from both the midrib and primary lateral veins; secretory ducts moderately visible on lower dried surface, alternating with minor veins. **INFLORESCENCES**  $\pm$  erect, 4(6) per axil; peduncle (4.5)6.5–20 (most more than 15) cm long, 4–12 mm diam., pale green, strongly white-lineate,

slightly to moderately bent just below the spathe; **spathe** 10.5–18.5 cm long, (0.6–1.7(2.7) times longer than peduncle), acute at apex; spathe blade white outside, (opening 4–7.5 cm wide), pale green, moderately glossy to pale-punctate inside; spathe tube ellipsoid, medium green, densely pale-speckled outside, 6–9 cm long, to 4.5 cm diam., pale green, moderately glossy to pale-punctate inside, **spadix** sessile; protruding forward at anthesis, 12–16 cm long, broadest at upper two-thirds constricted to ca. 1.5 cm diam. between sterile staminate portion and fertile staminate portion; pistillate portion pale green, cylindrical, 3–6.5 cm long in front, 2.7–3.6 cm long in back, 1.3–1.7 mm diam. at apex, 1.3–1.9 mm diam. at middle, 1.1–1.5 mm wide at base; staminate portion 10.5–14.7 cm long; fertile staminate portion creamy white,  $\pm$  clavate, 1.5–1.8 cm diam. at base, 1.3–2 cm diam. at middle, 9–12 mm diam. ca. 1 cm from apex, about as broad as the pistillate portion; sterile staminate portion 1–1.9 cm diam.; pistils 2.1–3.4 mm long, 1–1.6 mm diam.; ovary 6(7)-locular, 1.5–3.1 mm long, 1–1.6 mm diam., with axile placentation, walls sometimes embedded with granular, crystal-like particles; locules 1.5–3 mm long, 0.5–0.7 mm diam.; ovules 20–31 per locule, 2-seriate, 0.3 mm long; funicle 0.2 mm long, adnate to lower part of partition, style 0.5–0.6 mm long, 1.3–1.6 mm diam., similar to style type B; style apex rounded or domed; stigma truncate, hemispheroid, 1–1.3 mm diam., 0.3–0.5 mm high, covering entire style apex; the androecium truncate, prismatic, oblong, margins acutely and regularly 4–6-sided, 0.9–1 mm long, 0.7–2.2 mm diam. at apex; thecae oblong to cylindrical, 0.3–0.5 mm wide,  $\pm$  parallel to one another, contiguous; sterile staminate flowers bluntly, irregularly 4–6-sided, 1.9–4.1 mm long, 1.1–2 mm wide. **INFRACTESCENCE** with seeds many per berry, white, narrowly cylindrical, 1.3 mm long, sticky. **JUVENILE** petioles terete, sheathing broadly, for  $\frac{1}{2}$  to  $\frac{3}{4}$  its petiole length, acute to rounded, eventually weakly to strongly cordate; blades broadest at the middle.

Flowering in *Philodendron panamense* occurs during the dry season and early rainy season (March through May), with post-anthesis inflorescences collected from May through August and immature fruits from July through November.

*Philodendron panamense* is endemic to Panama, but it is likely to occur also in adjacent Colombia. In Panama, it occurs in *Tropical moist forest* on both slopes of the Canal Zone, and *Premontane wet forest* and *Tropical wet forest* in Panamá, Colón, and Da-

rién Provinces, at sea level to 800 m elevation (but mostly below 300 m).

*Philodendron panamense* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. *Philodendron panamense* is characterized by its appressed-climbing hemiepiphytic habit; short internodes; weakly one-ribbed, mostly deciduous cataphylls (sometimes persisting intact or as fibers); terete, pale-striate petioles; usually broadly triangular-ovate, moderately thin blades with parabolic to hippocrepiform sinus; and by the long-pedunculate spathe with externally greenish tube and white blade (greenish white throughout within). Also characteristic is the peduncle, which is often bent abruptly just beneath the spathe.

This species is most easily confused with *P. lazarii*, which has similar greenish inflorescences. The latter species differs in having more broadly ovate blades (0.97–1.4 times longer than wide and averaging 1.1 times) matte on the lower surface. In contrast, the blades of *P. panamense* are usually ovate-triangular, 1.3 or more times longer than broad (averaging 1.5 times longer) with the lower surface semiglossy. In addition, the peduncles of *P. lazarii* are proportionately shorter and usually shorter than the spathe (vs. often longer than the spathe in *P. panamense*).

*Philodendron panamense* has been confused with *P. jodavisanum*, which has leaves drying a similar, somewhat blackened color. The latter species differs, however, in having D- to U-shaped petioles, typically with a medial rib adaxially, as well as more typically persistent cataphyll fibers and much shorter peduncles (typically less than 10 cm long in *P. jodavisanum*, vs. typically more than 10 cm long in *P. panamense*). In addition, the peduncles of *P. jodavisanum* are usually straight, not bent.

A collection from the Serranía del Pirre at Cana, Croat 37600, is unusual in having several persistent cataphylls and shorter-than-usual peduncles.

Collections from Cerro Sapo and Cerro Pirre (Croat 55184 and 68951 respectively) differ from typical material of *P. panamense* in having ovate (rather than triangular ovate) leaves and more short-pedunculate inflorescences. In addition, they have sharply two-ribbed cataphylls that persist intact (in the case of Croat 55184, on plants in the living collection at MO) or semi-intact (Croat 68951, collected in the wild on Cerro Pirre). Most dried collections of *P. panamense* have no cataphylls and longer peduncles.

Earlier (Croat, 1978), the species was reported to be much more widely distributed (to Honduras and Ecuador), but collections from outside of Pan-

ama have since proven to be misidentified (most now *P. jodavisanum*).

*Additional specimens examined.* PANAMA. Canal Area: Barro Colorado Island, Croat 5101 (MO); 5530 (MO); 5840 (F, MO, SCZ); 6188 (MO); 8819 (MO, PMA); 8993 (MO); 9292 (MO, PMA, SCZ); 10083 (MO, SCZ); 10264 (MO, SCZ, US); 10819 (MO); 10894 (MO); 11016 (MO); 14876 (MO); Bailey & Bailey 199 (BH); Parque Nacional Soberanía, Pipeline Road, N of Gamboa, Latcyn & Foster 1556 (MO); Río Macho bridge, 10 km NW of Gamboa, 115 m, Nee 7869 (MO, US); 6 mi. N of Gamboa, Río Mendoza, 9°11'N, 79°46'W, Croat & Zhu 77084 (MO); Summit Gardens, Croat 10867 (F, MO). COLOMBIA: Portobelo-Nombré de Dios, 0.5 mi. beyond junction of road to Isla Grande, 9°40'N, 79°35'W, Croat 49804 (MO); near Nuevo Tonosí, <100 m, Croat 33517 (MO, RSA). DARIÉN: Cerro Sapo, Croat 55184 (AAU, K, MEXU, MO, PMA, US); ca. 5 km S of Garachiné, Río San Antonio, 130 m, 7°59'N, 78°25'W, Hammel et al. 1481 (MO); Cerro Pirre region, NW of Cana, 600 m, Sullinan 672 (MO); vic. of gold mine at Cana, 500–600 m, Croat 37600 (MO); Serranía Sapo above Casa Vieja along boundary trail of Darién National park, 150–300 m, 7°58'N, 78°23'W, McPherson et al. 15359 (COL, F, MO, NY, PMA), 550–830 m, 15378 (MO, US); Parque Nacional Darién, vic. of Cerro Pirre base camp, trail E, side of Río Paracida, 0–80 m, 8°N, 77°48'W, Croat 68991 (CM, L, MEXU, MO, PMA, US); Estación Rancho Frío, at N base of Cerro Pirre, ca. 9 km S of El Real, 70–270 m, 8°01'N, 77°44'W, Hammel et al. 16131 (MO); Cerro Pirre, 800–1050 m, 7°56'N, 77°45'W, Croat 68951 (CAS, COL, MO, NY, PMA); Río Tupuesa, ca. 2 km air distance from Continental Divide, vic. of Tyler Kiltredge gold mine, Croat 27193 (MO); Parque Nacional Cerro Pirre, Río Perisemico, 110 m, 8°01'N, 77°44'W, Croat & Zhu 77100 (CAS, CM, F, MO). PANAMA: road to Cerro Azul, Mile 5, Croat 11515 (F, MO, PMA); 720 m, 9°04'N, 79°29'W, Croat 75152 (CM, MO); Cerro Campana, Dwyer et al. 4848 (MO), Croat 12074 (MO, SCZ).

**Philodendron pirrese** Croat, sp. nov. TYPE:

Panama. Darién: Cerro Pirre, middle slopes on western approach, 800–1050 m, 7°56'N, 77°45'W, 29 June 1988, Croat 68952 (holotype, MO–3610823–24; isotypes, B, COL, F, K, NY, PMA, US). Figures 301–305.

Planta hemiepiphytica aut terrestris; internodia 1–2 cm longa, 4–5 cm diam.; cataphylla 25–30 cm diam., acute 2-costata, persistentia ut fibrae grossae pallidae, cum frustis rubrobrunneae epidermidis; petiolus 56.5–100 cm longus, 5–8 mm diam., aliquantum complanatus adaxialiter cum costa inconspicua; lamina late ovato-cordata, 35–58 cm longa, 27–46 cm lata, in sicco brunnea; costa postica nuda 1–5 cm; inflorescentia 4; pedunculus manifeste albostriatus, 8.5–12 cm longus, 6–8 mm diam.; spathe 12–16.7 cm longa; lamina spathae extus viridis, suffusa maronina, intus alba suffusa maronina; tubo spathae extus rubicarronino, intus atrimarronino; pistilla 5–6-locularia; loculi 20-ovulati; baccae albae.

Hemiepiphytic or sometimes terrestrial; stem to 5 cm long; internodes short, semiglossy, closely ribbed, completely enclosed in cataphyll fibers,

1–2 cm long, 4–5 cm diam.; roots to ca. 30 cm long, drying reddish brown, smooth, semiglossy, ca. 2 mm diam., closely ridged; cataphylls 25–30 cm long, sharply 2-ribbed (ribs prominently raised), reddish, drying reddish brown, persisting semi-intact, ultimately as coarse pale fibers with fragments of reddish brown epidermis; **petioles** 56.5–100 cm long, 5–8 mm diam., subterete, somewhat flattened near base, weakly flattened toward apex, with faint medial rib adaxially, surface densely short-lineate; **blades** broadly ovate-cordate, acuminate to narrowly acuminate at apex, cordate at base, 35–58 cm long, 27–46 cm wide (1.2–1.5 times longer than wide), (0.6–0.9 times longer than petiole), upper surface drying reddish brown, semiglossy, lower surface much paler; anterior lobe 24.5–48.5 cm long, 27–50 cm wide (1.7–2.3 times longer than posterior lobes); posterior lobes  $\pm$  rounded, 12.5–19 cm long, 12.5–23.5 cm wide, broadly obtuse; sinus hippocrepiform, 14 cm deep; midrib flat, paler than surface above, drying reddish brown below; basal veins 7–11 per side, with 0–1 free to base, 1–2 coalesce (4)6–8(10) cm, posterior rib well developed, naked for 1–5 cm; primary lateral veins 6–10 per side, departing midrib at a 55–65° angle,  $\pm$  straight to the margins, prominently sunken above, prominently raised below; interprimary veins distinct, fine, mostly continuous, drying darker than surface below; minor veins arising from both the midrib and primary lateral veins. **INFLORESCENCES** 4 per axil; peduncle 8.5–12 cm long, 6–8 mm diam., prominently white-striate, clearly demarcated from spathe; **spathe** 12–16.7 cm long (1.3–1.5 times longer than peduncle), moderately constricted above the tube; spathe blade green, tinged maroon, conspicuously and densely pale lineate-striate outside, white, tinged maroon inside; spathe tube red-maroon, inconspicuously short-lineate outside, 4–6 cm long, dark maroon inside; **spadix** sessile;  $\pm$  ovate, 11–13.4 cm long, broadest below the middle; pistillate portion greenish white, 2.7 cm long, 1.4 cm diam. at apex, 1.3 cm wide at base; staminate portion 10.8 cm long; staminate portion 5.7–8.3 cm long; fertile staminate portion white, tapered toward apex, 1.4 cm diam. at base, 1.3 cm diam. at middle, 8 mm diam. ca. 1 cm from apex, broadest at base, about as broad as pistillate portion; sterile staminate portion broader than the pistillate portion, 1.4–1.5 cm diam., pistils 2.5 mm long, 1.2–1.4 mm diam.; ovary 5–6-locular, 1.7 mm long, 1.3 mm diam., with axile placentation, walls embedded with granular, crystal-like particles; locules 1.7 mm long, 0.4 mm diam.; ovules 20 per locule, 2-seriate, contained within gelatinous matrix (no true envelope), 0.2–0.3 mm long, longer

than funicle; funicle 0.1–0.2 mm long, style 0.3 mm long, 1.3 mm diam., similar to style type B; style apex flat; stigma subdiscoid to slightly hemispheroid, weakly lobed, 1.2 mm diam., 0.1–0.3 mm high, covering entire style apex; the androecium truncate, prismatic, oblong, margins irregularly 4–6-sided; thecae oblong, 0.5 mm wide,  $\pm$  parallel to one another; sterile staminate flowers blunt, irregular, 4–6-sided, 2.1–3 mm long, 1.7 mm wide. **Berries** white.

Flowering in *Philodendron pirrense* occurs during the early rainy season, judging by post-anthesis collections from June and July.

*Philodendron pirrense* is endemic to Panama, where it is known for certain only from the Serranía de Pirre, at 1000 to 1560 m elevation in *Premontane rain forest*.

*Philodendron pirrense* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrusa*. This species is distinguished by its short internodes; sharply two-high-ribbed cataphylls persisting as pale fibers with tiny fragments of thin, reddish brown epidermis; petioles somewhat flattened adaxially with a faint medial rib; broadly ovate-cordate blades with a well-developed posterior rib naked to the sinus up to half its length; inflorescences up to four per axil; and whitish peduncles clearly demarcated from the red spathe tube.

*Philodendron pirrense* is similar to *P. copense*, with which it shares dried leaves of similar color and persistent, reddish brown cataphylls. The latter species differs in having longer blades with the anterior lobes frequently concave along their margins and petioles with a yellowish or reddish brown flaking periderm.

*Additional specimens examined.* PANAMA. Darién: Serranía de Pirre, ca. 12 km N of Alto de Nique, 1520–1560 m, Croat 37916 (MO); ca. 9 km from Alto de Nique, 1480–1520 m, Croat 37887 (MO); Río Esucho Ruído, ca. 16 km N of Alto de Nique, 1530–1550 m, Croat 37944 (MO).

***Philodendron platypetiolatum*** Madison, Selbyana 2: 22. 1977. TYPE: Ecuador. Los Ríos: Río Palenque Science Center, km 56 on road to Santo Domingo and Quevedo, 150–220 m, ca. 0°35'S, 79°22'W, Dodson 6638 (holotype, SEL; isotypes, MO, QCA, US). Figures 306–308.

Hemiepiphytic; stem  $\pm$  scandent, loosely appressed-climbing, to 3 m long, semiglossy, sap clear, watery, sticky, leaf scars to 2 cm wide; internodes slender, 14–20 cm long, 1–2 cm diam., longer than broad, sometimes somewhat flattened on

side above, semiglossy, green to grayish green, epidermis drying tannish brown, cracking, loosening and flaking; roots pale to brownish, less than 20 cm long, thin, 2 mm diam., smooth; cataphylls (6)10–17 cm long, sharply 2-ribbed, sharply and deeply sulcate with margins flared, pale to medium green, drying yellowish to olive-green, deciduous; **petioles** 15.5–56 cm long, 3–9(12) mm diam., broadly flattened to markedly flattened and turned slightly upward adaxially, broadly convex abaxially, firm, medium to dark green, surface  $\pm$  unmarked; **blades** ovate-triangular to broadly ovate, subcoriaceous, semiglossy to glossy, weakly to moderately bicolorous, acuminate, sometimes long, narrowly acuminate at apex (the acumen tightly inrolled, 2–3 mm long), weakly cordate at base, 17–39 cm long, 12.7–29 cm wide (1–1.5 times longer than wide), (0.6–1.4 times longer than petiole), usually about equal in length to petiole; anterior lobe 15–34 cm long, 13–29 cm wide (2.4–4.5(5.2–5.4) times longer than posterior lobes); posterior lobes 4–10 cm long, 5–12 cm wide, broadly rounded to obtuse; sinus hippocrepiform, rarely arcuate with blade decurrent on petiole; midrib sunken or flat to broadly convex above, slightly paler than surface to concolorous above, bluntly low-triangular to convex below, paler than surface below; basal veins (2)3–4(5) per side, with 0–1 free to base, second and third coalesced 1–2 cm; posterior rib naked for 0–1 cm; primary lateral veins 3–5 per side, departing midrib at a 50–60° angle, straight to the margins, sunken to weakly sunken above, convex below; minor veins moderately indistinct, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 1(2) per axil; peduncle 5–20 cm long, 2–11 mm diam., subterete or bluntly 1–2-angled, pale green, semiglossy, unmarked; **spathe** semiglossy, (8)11–18 cm long (0.7–2(2.8) times longer than peduncle); spathe blade green or red outside, pale yellow-green (dark red in South America), inside; spathe tube green to olive-green, usually tinged red or maroon outside, 4.5–7.5 cm long, tinged red or maroon inside; **spadix** sessile; weakly tapered,  $\pm$  acute at apex, 8–9.8 cm long, broadest at the base or  $\pm$  uniform throughout, constricted weakly between sterile and fertile staminate portions; pistillate portion pale greenish, cylindrical or weakly tapered toward the apex, 2.5(3.5) cm long, 9–11 mm diam. at apex, 10–13 mm diam. at middle, 10–13 mm wide at base; staminate portion 5.7–7.3 cm long; fertile staminate portion white, weakly tapered or cylindrical, 5–7 mm diam. at base, 4–5 mm diam. ca. 1 cm from apex, broadest at the base or  $\pm$  uniform, narrower than the pistillate portion, broader than or as broad as the sterile portion; ster-

ile staminate portion as broad as or slightly narrower than the pistillate portion, 5–10 mm diam.; pistils (0.8)1.7–2.5 mm long, (0.6)0.9–1.4 mm diam., white; ovary (6)7–8-locular, (0.7)1.1–1.8 mm long, 0.8–1.2 mm diam., with sub-basal placentation; locules 0.7–1.5 mm long, 0.3 mm diam., ovule sac 0.8 mm long; ovules 3 per locule, 1-seriate, contained within translucent to transparent, gelatinous envelope, (0.2–0.3)0.4–0.5 mm long, longer than funicle; funicle 0.2–0.3 mm long (can be pulled free to base), style (0.1)0.4–0.7 mm long, 0.6–0.8 mm diam., similar to style type B; style apex steeply sloping, sometimes with small medial depression; stigma subdiscoid to hemispheroid, somewhat cupulate, 0.7–0.9(1.3) mm diam., 0.2–0.3 mm high, covering entire style apex; the androecium prismatic, truncate, oblong, margins irregularly 4–6-sided, sometimes weakly scalloped; thecae cylindrical to oblong, 0.3–0.4 mm wide,  $\pm$  parallel to one another, sometimes  $\pm$  contiguous; sterile staminate flowers irregularly 4–6-sided or rounded, 0.9–1.5 mm long, 0.7–1.2 mm wide. **INFRUITESCENCE** turning red outside; berries green (immature).

Flowering in *Philodendron platypetiolatum* occurs in the early rainy season, June and September, in Central America based on the few available flowering collections. Post-anthesis collections from Costa Rica and Panama are mainly from August through November, but also from March. Immature fruits have been collected only in January, February, and November. Ecuadorian populations have a similar phenology, but flowering collections have been made earlier, in March and April, with post-anthesis collections from March through August.

*Philodendron platypetiolatum* ranges from Nicaragua to Ecuador, from sea level to 1500 (though most collections are from below 400) m elevation, in *Premontane wet forest*, *Tropical wet forest* transition to *Premontane wet forest*, and *Tropical wet forest* life zones. This species has been collected at several localities in Chocó and Valle Departments; in Ecuador, it has only been collected at and near the type locality.

*Philodendron platypetiolatum* is a member of *P. sect. Calostigma* subsect. *Macrobelium* ser. *Macrobelium*. This species is characterized by its scandent habit; moderately long, slender internodes with tannish brown, frequently flaking epidermis; markedly flattened petioles about as long as the blade; and ovate-triangular to broadly ovate, weakly cordate blades. Though the spathes may be entirely green outside, they usually are heavily tinged with red or maroon outside and the tube within is likewise colored.

There are no other species in Central America with which *P. platypetiolatum* might be confused. Madison compared this species to *P. lechlerianum* Schott, a species from Peru and Bolivia that has terete petioles.

In Panama, this species ranges along the Atlantic slope but may also occur on the Pacific slope near the Continental Divide. In Ecuador, it is restricted to the Pacific slope.

Ecuadorian specimens have blades that are more broadly ovate than those from Panama, but no other differences have been detected.

*Additional specimens examined.* COSTA RICA. **Alajuela:** Upala Road, 3 km NNE of Bijagua, 450 m, 10°45'N, 85°3'W, *Burger & Baker* 9890 (CR, F, MO, NY, SEL); 5 km S of Canalete, near Río Zapote, 100–200 m, 10°48'N, 85°2'W, *Burger & Baker* 9972 (F, MO); Dos Ríos, 5 km S de Brasilia, Río Pizote, 500 m, 10°55'N, 85°20'W, *Herrera* 1001 (CR, MO, SAR); 17 km NW of San Ramón, 785 m, 10°14'14"N, 84°33'W, *Croat* 68136 (F, MO); Bajo Rodríguez-La Tigra, vic. of La Tigra, 330 m, 10°22'N, 84°38'W, *Croat* 68205 (B, CM, G, M, MO, SAR, US); Monteverde Cloud Forest Nature Reserve, Río Peñas Blancas, 1250–1350 m, 9°17'N, 84°86'W, *Burger et al.* 10745 (F, MO); Vara Blanca—Puerto Viejo, 3 mi. N of San Miguel, 380 m, *Croat* 35666 (MO); Cañas—Upala, 13.8 km N of Bijagua, 100–150 m, *Croat* 36433 (MO); 4 km NNE of Bijagua, ca. 400 m, *Croat* 36269 (MO); slopes of Miravalles, above Bijagua, *Gómez et al.* 19066 (MO); Naranjo—Aguas Zarcas, along Hwy. 15, 8.5 km NE of Villa Quesada, 600 m, *Croat* 46973 (CR, MO). **Heredia:** "Starkey Road," 4.5 km SE of bridge at Puerto Viejo, ca. 50 m, 10°26'N, 83°58'W, *Stevens* 13489 (MO); La Selva Field Station, *Grayum* 2228 (MO); 100 m, *McDouell* 329 (MO); 100 m, 647 (CAS); 50–80 m, 10°26'N, 84°01'W, *Grayum* 7665 (CR, MO); 6 km from Río Peje crossing, 5 km SSE of Magassay, 340 m, 10°21'N, 84°04'W, *Schetz & Grayum* 634 (CR, MO); Puerto Viejo de Sarapiquí, 100 m, *Croat* 44248 (MO); 10°26'N, 84°01'W, 61213 (MO); Parque Nacional Braulio Carrillo, Río Peje—Río Sardinalito, Volcán Barva, 700–800 m, 10°17'30"N, 84°05'W, *Grayum & Herrera* 7968 (CR, F, MO, VEN). **Limón:** Hacienda Tapasco—Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, *Davidson* 6744 (RSA), *Davidson & Donahue* 8383 (F, MO, RSA); *Davidson & Donahue* 8802 (MO); 8874 (MO, RSA); Barra del Colorado, 0–2 m, 10°47'N, 83°35'W, *Stevens* 24214 (CR, MO); Cerro Coronel, 20–170 m, 10°41'N, 83°38'W, *Stevens* 24623 (CR, MO); Quebrada Danta, W of Guápiles, 360 m, 10°12'N, 83°49'W, *Croat* 68426 (MO, NY); Braulio Carrillo—Guápiles, 250–270 m, *Croat* 78744 (CR, INB, MO); Río Colorado, 14 km by air SW of Barra del Colorado, 10–120 m, 10°40'N, 83°40'W, *Davidse & Herrera* 31060 (CR, MO); 16 km by air SW of Barra del Colorado, 10°39'N, 83°40'40"W, 31211 (MO). **Puntarenas:** Osa Peninsula, Piedras Blancas, 3.7 mi. W of Pan-American Highway, 90–105 m, 8°46'N, 83°18'W, *Croat* 67687 (CR, K, MO); Rincón de Osa-Esquinas, 100 m, *Pennington et al.* 11386 (K); Fila Gamba, ca. 6 km from Golfo airport, <100 m, 8°41'30"N, 83°12'W, *Croat* 59926 (CM, K, MO); 200–300 m, *Croat & Hannon* 79290 (MO). **San José:** El General Valley, Finca Volcán Angel, *Schubert & Rogerson* 769 (A, GH); Braulio Carrillo National park, 600–700 m, *Croat*

78778 (CR, INB, MO). **NICARAGUA.** Río San Juan: Río Santa Cruz—Caño Santa Cruzita, La Palma, 40–60 m, 11°2–4'N, 84°24–26'W, *Stevens* 23496 (MO). **Zelaya:** Río Sució, E of Bonanza, ca. 140 m, 14°01'N, 84°34'W, *Stevens* 12347 (MO); Mpio. Siuna, Comarca Danlí, 100–130 m, *Ortiz* 218 (MO). **PANAMA.** **Bocas del Toro:** Fortuna Dam area, Gualaca—Chiriquí Grande, 9.4 mi. N of Continental Divide, 175 m, 8°46'N, 82°16'W, *Croat* 66822 (AAU, MO). **Colón:** Río Guanche, ca. 3–4 km above bridge on road to Portobelo, *J. Wütherspoon & F. Wütherspoon* 8662 (CAS, MO); Sabanita—Portobello, Río Piedras Lumber Road, 6.7 mi. E of Sabanita, 250 m, 9°22'30"N, 79°41'30"W, *Croat* 75166 (MO, PMA). **Darién:** Parque Nacional Cerro Pirre region, Cana, 500–600 m, *Croat* 37661 (MO); near station along Río Perisénico, 110 m, 8°01'N, 77°44'W, *Croat & Zhu* 77116 (MO); Parque Nacional Darién, Río Topalisa—Río Pucuro, ca. 17 km E of Pucuro, Mi Casita—La Laguna, 600–850 m, 8°03.5'N, 77°17'W, *de Nevers et al.* 8337 (CAS, MO); ca. 5 km E of Pucuro, Quebrada Maskia, 200 m, 8°01'N, 77°25'W, *Hammel et al.* 16167 (MO). **Panamá:** El Llano—Cartí, 13.8 km N of Pan-American Highway, *Folsom et al.* 5788 (MO); Mile 6.8, 350 m, *Croat* 49125 (MO, PMA); Cerro Jefe region, 0.8 mi. beyond turnout to Altos de Pacora, 770 m, 9°15'N, 79°29'W, *Croat & Zhu* 76645 (MO); 4.6 km beyond peak on road to Altos de Pacora, ca. 600 m, *Croat* 35936 (MO); near summit, 750–800 m, 9°14'N, 79°22'W, *Croat* 67083 (MO); 3–3.5 mi. NE of Altos de Pacora, 700–750 m, 9°15'N, 79°25'W, *Croat* 68680 (CM, MO). **San Blas:** El Llano—Cartí Road, vic. Nusugrúti, 300–350 m, 9°15'N, 79°W, *Croat* 69278 (CM, MO); 450 m, 9°18'N, 79°59'W, *Croat* 75118 (CAS, MO, NY, PMA); 10.1 mi. N of main highway, 300 m, 9°20'N, 79°W, *Croat & Zhu* 76539 (CM, M, MO); Mile 9, 350 m, 9°20'N, 79°W, *Croat* 76999 (MO); Río Playón Chico, 80–200 m, 9°13.5'N, 78°15'W, *Herrera & Arasmena* 1778 (AAU, COL, CR, K, MEXU, MO, NY, P, PMA, SIRI, US).

#### *Philodendron pseudauriculatum* Croat, sp. nov.

TYPE: Panama. Panamá: El Llano—Cartí road, 4 mi. from Inter-American Hwy. near El Llano, ca. 300 m, 27 Mar. 1976, *Croat* 33730 (holotype, MO—2381528; isotypes, PMA, RSA, SEL). Figures 35, 300, 309, 310, 313, 314.

Planta hemiepiphytica; internodia 1.4(9) cm longa, 2–4 cm diam.; cataphylla 18–27 cm longa, acute 2-costata, decidua; petiolus subspongiosus, 11–42 cm longus, 0.7–2.6 cm diam., subteter, leviter complanatus adaxialiter, late et obtuse sulcatus in medio, cum annulo viridi apice; lamina oblongo-elliptica vel oblanceolata-elliptica, plurimum acuta vel anguste rotundata, internum subcordata truncata basi, 27–80 cm longa, 7.5–25 cm lata; in sicco atricaniviridis; inflorescentia 2–3; pedunculus 5.5–21(25) cm longus, 8–12 mm diam.; spatia (10.6)12–23 cm longa, omnino alba vel subrosea, marginibus cremeis; lamina spathae extus eburnea vel flaviviridi, striata flavaurantiaca, intus pallide viridi vel alba; tubo spathae extus vel atriviridi, internum extus leniter suffuso purpureo-roseo; cecus margines, vivide rubrivioleaceo vel atrimarmerino vel flavaurantiaco; pistilla 5–8(9)-locularia; loculi 1–2(4)-ovulati; baccae aurantiacae.

Hemiepiphytic; stem appressed-climbing, to 1 m long; internodes gray-green, glossy to semiglossy.

1-4(9) cm long, 2-4 cm diam., usually somewhat flattened on one side, frequently with a short series of transverse ridges just below the nodes on both sides of the rounded portion of internodes, usually about as broad as long, or broader than long on flowering plants, sometimes slightly longer than broad; roots dark brown, slender, few per node; cataphylls moderately spongy, 18-27 cm long, sharply 2-ribbed (ribs to ca. 1 cm high), medium green, densely dark green short-lineate, deciduous, apiculate (apiculum >1 cm long) at apex. LEAVES erect,  $\pm$  rosulate, somewhat clustered at or near stem apex; **petioles** 11-42 cm long, 0.7-2.6 cm diam., subterete, somewhat spongy, slightly flattened, bluntly and broadly sulcate midway, less so toward apex, more so toward base adaxially, surface medium green, semiglossy, with dark green ring around apex; **blades** oblong-elliptic to oblanceolate-elliptic, subcoriaceous to coriaceous, somewhat to markedly bicolorous, semiglossy, acuminate, sometimes long-acuminate at apex, mostly acute to rounded, sometimes broadly subcordate, or sometimes truncate at base, 27-80 cm long, 7.5-25 cm wide (2.2-4(5) times longer than wide), (0.97-3.6 times longer than petiole), upper surface dark green, drying dark gray-green, lower surface much paler, yellow-green, drying yellow-green to yellow-brown; sinus (when present) to 2.5 cm deep; midrib flattened at base, slightly sulcate midway, broadly convex at apex and concolorous above, convex, short-green-lineate, paler than surface below; basal veins (0)1-3(4), all free to base; posterior rib lacking; primary lateral veins 8-14 per side, departing midrib at a 65-75° angle (45-55° angle at apex),  $\pm$  straight to the margins, sunken above, convex and slightly paler than surface or darker than surface below; interprimary veins flat, darker than surface below; minor veins moderately distinct and fine below, arising from both the midrib and primary lateral veins. INFLORESCENCES 2-3 per axil; peduncle 5.5-21(25) cm long, 8-12 mm diam., pale to medium green, finely white- or dark-striate; **spathe** (10.6)12-23 cm long, 1.7 cm diam. (0.7-2.6 times longer than peduncle), constricted midway above the tube, white to pinkish throughout, margins cream; spathe blade creamy white to yellowish green with faint green tinge along center on backside (B & K yellow-red 7.5/9), yellow-orange striate outside, 8.5-9 cm long (opening elliptic in face view, 7.5-9 cm long, 2-5.5 cm wide), pale green to white inside; resin canals orange; spathe tube oblong-ellipsoid, medium to dark green, sometimes weakly tinged purplish along margins outside, densely short white-lineate, semi-glossy outside, 5.5-10.5 cm long, 1.8-4 cm

diam., bright red-violet to dark maroon (weakly so toward apex) to yellowish orange, sometimes broad, white-lineate or orange striate inside; **spadix** weakly stipitate to 3-5 mm long, cylindrical to weakly tapered, 7.5-15.3 cm long, broadest below the middle; pistillate portion cylindrical to clavate, pale lime green to pale yellow to medium or dark green, 3-5.5 cm long in front, 1.7-4 cm long in back, 1-1.2 cm diam. at apex, 1.5 cm diam. at middle, 8-11 mm wide at base; staminate portion 4.5-10.7(12) cm long; fertile staminate portion broadest in middle, slightly tapered toward both ends and broadened before the sterile portion, 9-16 mm diam. at base, 9-10 mm diam. at middle, 5-10 mm diam. ca. 1 cm from apex, broadest at the base, as broad as the pistillate portion, narrower than the sterile portion; sterile staminate portion broader than the pistillate portion, white, 8-15 mm diam.; pistils 1.7-3.2 mm long, 1.3-2 mm diam.; ovary 5-8(9)-locular, 1.4-2.3 mm long, 1.5-2 mm diam., with sub-basal placentation, walls embedded with granular, crystal-like particles; locules 1.4-2.1 mm long, 0.3-0.5 mm diam.; ovule sac 1-1.2 mm long; ovules 1-2(4) per locule, contained within transparent, gelatinous ovule sac, 0.4-0.5 mm long, longer than funicle; funicle 0.1-0.5 mm long (can be pulled free to base), style 0.4-1.4 mm long, 0.9-2.1 mm diam., similar to style type B; style apex sloping to rounded, with small medial depression; stigma brush-like, cupulate, subdiscoid, 0.9-1.2 mm diam., 0.2-0.6 mm high, covering entire style apex; the androecium truncate, prismatic, oblong, margins irregularly 4-6-sided, 1.3-1.4 mm long, 1.6-1.8 mm diam. at apex; thecae oblong to cylindrical, 0.4-0.5 mm wide, nearly contiguous and  $\pm$  parallel to one another; pollen ellipsoidal to spheroidal, <1 mm long, <1 mm diam.; sterile staminate flowers irregularly 4-5-sided, margins bluntly rounded, 1.2-1.9 mm long, 1.8 mm wide. Berries orange (mature) or white; seeds 5 per berry, 1.3-1.9 mm long, 0.5-0.7 mm diam.

Flowering in *Philodendron pseudauriculatum* occurs during the dry season and the first half of the rainy season (January through September). Immature fruits have been collected during March, June, and July, with mature fruits known only from September.

*Philodendron pseudauriculatum* is definitely known only from Panama and adjacent Colombia (N Antioquia), ranging from 20 to 1400 m elevation in *Premontane wet forest* and *Tropical wet forest* life zones. Most collections have been made at La Mesa (Coclé), Cerro Campana (Panamá), and along the El Llano-Cartí Road (Panamá), but the species is

also known from a single collection in Bocas del Toro and in the Serranía de Cafazas along the Pacific Ocean.

*Philodendron pseudauriculatum* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is recognized by its appressed-climbing habit; short internodes; more or less rosulate habit; somewhat spongy, subterete petioles; oblong-elliptic to oblanceolate-elliptic, dark gray-green-drying blades with mostly narrowly rounded bases, and two to three inflorescences per axil; and white to pinkish spathes clearly demarcated from the peduncles.

*Philodendron pseudauriculatum* is most easily confused with *P. ligulatum*, especially *P. ligulatum* var. *heracioanum* and *ligulatum*, which have similarly shaped blades. *Philodendron ligulatum* var. *heracioanum* differs in having sharply D-shaped petioles with undulate-margined wings. While the aforementioned varieties of *P. ligulatum* differ in their usually vining habit and typically elongate internodes, these features are particularly apparent in *P. ligulatum* var. *ligulatum*, which has internodes much longer than wide (vs. about as long as broad or scarcely longer than broad as in *P. pseudauriculatum*). In addition, the leaves of *P. ligulatum* frequently dry much darker, mostly somewhat blackened, rather than the typical yellowish or brownish green of *P. pseudauriculatum*. Another feature separating live material of the species is the line of demarcation at the apex of the petiole, which is purple in *P. ligulatum* rather than green as in *P. pseudauriculatum*.

*Philodendron pseudauriculatum* may also be confused with both *P. auriculatum* (hence the epithet "pseudauriculatum"), from the Pacific slope of southwestern Costa Rica and *P. wendlandii* Schott from the Atlantic slope of Costa Rica and Panama. *Philodendron auriculatum* is distinguished by having leaf blades drying a paler yellow-green color and more narrowed toward the base with minute narrow auriculate posterior lobes. It also has proportionately somewhat longer petioles. *Philodendron wendlandii* differs in having a much shorter stem and petioles typically broader than thick with sharp margins.

Two collections (*Croat 16908* and *Knapp & Mallet 4658*) from San Blas at Puerto Obaldía are probably also this species. They differ in having broader blades and drying somewhat blacker. If these prove to represent *P. pseudauriculatum*, the species is most assuredly present in adjacent Chocó. A collection from the Department of Santander in Colombia at 1460 to 1700 m (*García-Barriga & Jaramillo 19671*) may also be this species.

*Additional specimens examined.* PANAMA. Bocas del Toro: Quebrada Huron, Duwebdubup Peak, N of Río Terere, 300–900 ft., *Kirkbride & Duke 558* (MO). Canal Area: Summit Gardens, *Croat 10791* (F, MO, SCZ). Colé: El Valle region, La Mesa, 650–710 m, *Mori et al. 1916* (MO); *Luteyn & Kennedy 1660* (DUKE); *Hammel 3844* (MO); 860–900 m, *Croat 37420* (MO); 900 m, 22964 (MO); 14362 (MO); 800 m, *Croat & Zhu 76692* (MO); *Bartlett & Lasser 16694* (F, MICH, MO); *Kennedy et al. 3189* (MO, PMA); 785 m, 8°37'N, 80°08'W, *Croat 67112* (MO); summit of Cerro Catacoral, near La Mesa, N of El Valle de Antón, 1100 m, *Knapp 1118* (K, MO); Río Cascajal, Penonomé-Cocleito, 5.6 mi. N of Llano Grande, 150 m, 8°46'N, 80°27'W, *Croat 67484* (AAU, CAS, COL, F, L, MEXU, MO, NY, PMA, QCA, TEX). Colón: near Peluca, on road to Nombre de Dios, *Kennedy 2774* (F, MO); Portobelo-Nombre de Dios, Nuevo Tonosí, <100 m, *Croat 33526* (CM, COL, K, L, MO, NY, PMA, QCA, RSA, US); Sabanita-Portobelo, Río Piedras drainage, 250 m, 9°22'30"N, 79°41'30"W, *Croat 75160* (MO); Portobelo, *Croat & Porter 15606* (MO); Río Boquerón, near No. 1 (manganese mine), E of Salamanca, 50 m, 9°35'N, 79°32'W, *Knapp et al. 5832* (B, K, MO); Puerto Pilón-Portobelo, ca. 1.5 mi. above bridge, <100 m, 9°27'N, 79°4'W, *Croat & Zhu 76251* (CR, MO, PMA); Río Guanche, 30–100 m, *Croat 79322* (PMA); ca. 5 km upstream from road to Portobelo, 50 m, 9°30'N, 79°40'W, *Hammel & Trainer 14765* (MO); ca. 3–5 km above bridge, 50–200 m, *Croat 37002* (MO); 10–100 m, 26147 (BR, F, MO); ca. 50 m, ca. 9°30'N, 79°40'W, *McPherson 8507* (MO); Río Iguanita, near bridge along Portobelo Road, <50 m, 9°27'N, 79°42'W, *Croat 497784* (MO); Río Miguel de la Boeda, vic. of Guásimo, *Croat 9985* (MO, SCZ); Río Sierri, Trinidad Basin, 20–50 m, *Pitiner 4015* (US). Darién: Cerro Pirre National Park, W side of Cerro Pirre, base camp, 50 m, 8°N, 77°48'W, *Croat 68962* (M, MO, NY, US); near station along Río Perisenoa, 110 m, 8°01'N, 77°44'W, *Croat & Zhu 77094* (CM, MO); Cerro Pirre region, trail NW of Cana, 600 m, *Sullivan 712* (MO); Cana, near Río Setigandí, 540–580 m, *Gentry et al. 28542* (MO); Cana gold mine, vic. of airstrip, 480 m, *Croat 38038* (MO); 500–600 m, 37595 (MO), along Río Cana, SW of Cerro Pirre, 1400 m, *Croat 27292* (MO); Río Coasi, 0–2 mi. E of Tres Bocas, *Kirkbride & Duke 1203* (MO, NY); Río Cocalito, *Whiteford & Eddy 224* (BM); 162 (BM, MO); Río Jaqué Valley, Quebrada Luksa, 100–200 m, 7°27'N, 78°05'W, *Knapp & Mallet 3199* (MO, NY); Río Turgesca, ca. 2 km by air from Continental Divide, *Croat 27160* (F, MO); *Clezio 168* (MO). Panamá: El Llano-Cartí Road, 5 mi. from Pan-American Highway, 350 m, *Croat 67345* (CAS, MO); Km 8–12, ca. 400–450 m, *Nee et al. 8802* (MO); ca. Mile 8, 225–275 m, 9°15'04"N, 79°00'04"W, *McPherson 10492* (AAU, MEXU, MO); Km 19.1, 350 m, 9°19'N, 78°55'W, *de Nevers et al. 7342* (MO); Río Terabe Valley, El Llano-Cartí, 8 km from highway, 300–400 m, 9°16'N, 79°W, *Knapp & Schmalzer 5493* (MO); area around Pilota de Toro, *Folsom et al. 6819* (MO, PMA); Tortí-Pilota del Toro, above Tortí Arriba, *Folsom et al. 4993* (MO, PMA); Cerro Campana, 6.1 mi. above Pan-American Highway, 800 m, 8°41'N, 79°56'W, *Croat 74762* (F, MO); ca. 1 mi. from highway, ca. 150 m, 35991 (MO); along trail to summit, 780–875 m, 25253 (MO, NY, PMA, US); ca. 850 m, 8°42'N, 79°56'W, *Miller et al. 754* (MO); upper slopes, 207 m, *LeDoux 2595* (MO); above Su Lin Motel, *Porter et al. 4250* (MO); Río Tortí, base of Serranía de Cafazas, ca. 15 km SW of Casaza, 150 m, 8°52'N, 78°22'W, *Stein 1342* (MO); Sendero de Intercep-



tación. 1 km al este del Campamento de los guardabosques de INRENARE, 800–900 m, 8°40'N, 79°55'W, *Correa & Montenegro 10681b* (STRI). **San Blas:** El Llano-Carti Road, Km 19, 350 m, 9°19'N, 78°55'W, *de Nevers et al. 5598* (MO); Puerto Obaldía, 0–50 m, *Pittier 4398* (US); beach E of Puerto Obaldía, *Croat 16908* (MO); Puerto Obaldía-La Bonga, ca. 2 hours walk from Puerto Obaldía, 0–50 m, 8°40'N, 77°25'W, *Knapp & Mallet 4658* (MO); Nusagandí, El Llano-Carti Road, 10.1 mi. N of main highway, 300 m, 9°20'N, 79°W, *Croat & Zhu 76554* (CM, MO); Río Playón Chico, 80–200 m, 9°13'05"N, 78°15'W, *Herrera & Arosemena 1784* (MO, PMA, STRI).

**COLOMBIA. Antioquia:** Mpio. Turbo, carretera tapón del Darién, sector Río León-lomas aisladas, km 37, 20 m, *Braud 1070* (COL, MO). **Chocó:** Mecana, N of Bahía Solano, 1–100 m, 5°16'N, 77°21'W, *Juncosa 1609* (MO). **Risaralda:** Misstrató, Jiguasdas-Santa Cecilia, 800–850 m, 5°24'N, 76°01'W, *Betanour et al. 3459* (MO); corregimiento de Santa Cecilia, 500–550 m, 5°17'N, 76°13'W, *Betanour et al. 2930* (MO); Pueblo Rico, Santa Cecilia-Pueblo Rico, Km 13, Quebrada Pionda, 700–900 m, 5°17'N, 76°13'W, *Betanour et al. 3052* (MO).

***Philodendron pterotum*** K. Koch & Augustin, in A. Braun et al., *Append. gen. sp. Hort. berol.* 1854: 6. 1854–1855. TYPE: Cultivated at Berlin [received from Warszewicz in Venezuela] (holotype, B? lost). Panama. Canal Area: vic. Fort Sherman, along road between Gatún Locks and Fort Sherman, ca. 3 mi. W of Gatún Locks, <50 m, 9°19'N, 79°57'30"W, 17 July 1994, *Croat & Zhu 76982* (neotype, MO-4619421–26, here designated; isoneotypes, AAU, B, CAS, CM, COL, CR, DUKE, F, GB, GH, K, MEXU, P, PMA, QCA, RSA, SEL, VEN, W). Figures 311, 312, 315, 316.

*Philodendron mirificum* Standl. & L. O. Williams, *Ceiba* 3: 38. 1952. TYPE: Costa Rica. Puntarenas: near Palmar Sur de Osa, 75 m, 24 Mar. 1951, *Allen 6031* (holotype, EAP; isotype, F).

Usually hemiepiphytic or epiphytic; stem appressed-climbing, semiglossy, sap watery, weakly turpentine-scented, leaf scars conspicuous, 2 cm long, 1 cm wide; internodes sometimes obscured by cataphylls, sparsely short-striate, about as long as broad or sometimes longer than broad, 2–8 cm diam., dark green to gray-green, eventually brown, epidermis sometimes cracking, fissured longitudinally; roots short, few per node; cataphylls 20–30 cm long, sharply 2-ribbed, C-shaped, semiglossy, dark green, densely pale lineate, persisting as reddish brown, semi-intact fibers, eventually deciduous; **petioles** (37)47–111 cm long, (3–)5(6–)13(16–22) mm diam., erect-spreading, D-shaped, marginally winged, broadly convex adaxially, with adaxial margins slender, erect, undulate, medium green, weakly glossy, conspicuously pale striate; sheath with margins involute; **blades** ovate, subcoriaceous,

semiglossy, moderately bicolorous, abruptly acuminate at apex (the acumen inrolled, 2–4(6) mm long), broadly cordate at base, 36–93 cm long, 22–84 cm wide (0.7–1.5(2.9) times longer than wide), (0.6–0.9 times the petiole length), margins weakly undulate, upper surface dark green, lower surface much paler, glossy, drying yellow-green; anterior lobe 23.5–65(84) cm long, 22.1–75(86.4) cm wide (1.1–2.8 times longer than posterior lobes); posterior lobes 12–33 cm long, (11)15–34 cm wide, directed inward, broadly rounded to broadly obtuse; sinus spatulate to rhombic; basal veins 7–10 per side, first free to base, part of remainder coalesced 1–6 cm, loosely so distally; posterior rib naked to 6 cm; midrib flat to broadly sunken, paler than surface above, convex to narrowly rounded, paler than surface below; primary lateral veins 3–6 per side, departing midrib at a 45–55° angle, ± straight to weakly arcuate to the margins, deeply sunken, paler than surface above, raised to convex, paler than surface below; interprimary veins weakly raised, darker than surface below; the minor veins moderately distinct, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, 2–4 per axil; peduncle (2)4–12 cm long, 4–13 mm diam., medium green, coarsely white streaked toward apex; **spathe** 11–29 cm long (1.4–3.8(5) times longer than peduncle), ± cuspidate at apex; spathe blade light green outside, greenish white inside; spathe tube abruptly delineated from tube, reddish to purplish to dark purple-violet (B & K purple 2/10) or dark green and raised-white-striate at base outside, red to magenta inside; **spadix** weakly stipitate, exerted from the spathe, constricted above sterile staminate portion; pistillate portion pale green, 4.3 cm long in front, 3.8 cm long in back, 2 cm diam. at apex, 1.7 cm wide at base; staminate portion 16–18 cm long; fertile staminate portion white, 1.7 cm diam. at middle; sterile staminate portion 1.4–2.2 cm diam.; pistils 2.2–2.8 mm long, 1.1–1.4 mm diam.; ovary 6-locular, with axile placentation; locules 1.5–1.9 mm long, 0.4 mm diam.; ovules ca. 20 per locule, 2-seriate, somewhat translucent, 0.2–0.3 mm long, longer than funicle; funicle 0.1–0.2 mm long, adnate to lower part of partition, style similar to style type B; style apex flat to weakly rounded; stigma subdiscooid to weakly hemispheroid, sometimes weakly lobed, 1.5 mm diam., 0.5 mm high, covering entire style apex; the androecium truncate, prismatic, margins irregularly 4–6-sided, 0.7–1.5 mm long; thecae oblong, 0.4 mm wide, not contiguous, ± parallel to one another; sterile staminate flowers irregularly 4–5-sided, 1.5–2.5 mm long. INFRUCTIONESCENCE with spathe green at base and tip, pur-

ple around fruiting area when ripe, berries pale brown to white. JUVENILE and PRE-ADULT plants with petioles flattened adaxially; blades broadly ovate. PRE-ADULT blades broadly ovate, 28.5 cm long, 22.5 cm wide.

Flowering in *Philodendron pterotum* occurs in the early rainy season from May through August based on both specimens and field observations; it certainly must flower during part of the dry season as well, since post-anthesis collections have been made as early as March. Other post-anthesis collections have been made in April and May but especially in June. Immature fruits have been collected in February, March, May, July, October, and November, especially October. Mature fruits have been collected only in November.

*Philodendron pterotum* ranges from Nicaragua to Central Panama, from sea level to 1900 (mostly below 700) m elevation in *Tropical moist forest* and *Tropical wet forest* life zones. In Nicaragua it occurs only on the Atlantic slope, but in Costa Rica and Panama it occurs on both slopes.

*Philodendron pterotum* is a member of *P.* sect. *Philodendron* subsect. *Platyopodium*. This species is characterized by its broadly ovate juvenile leaves with flattened petioles; appressed-climbing adult habit with short internodes; persistent cataphyll fibers; D-shaped petioles with slender, erect, undulate, marginal wings; and large, ovate, yellow-green-drying blades with large inflorescences with the spathe tube reddish to purplish on the outside and much darker magenta within.

No type material has survived for *P. pterotum*. The species was described from cultivated material of a juvenile plant purportedly received from the "little garden of Cl. Augustin," which the tireless traveler (Mr. Augustin) collected from Warszewicz in Venezuela. It is important to note that it was not said to have been collected in Venezuela (where it does not occur), but only that it was obtained from Warszewicz in Venezuela. There is much uncertainty involved, since the extrapolation of juvenile forms to adult plants is at best risky, even at the type locality. Given the frequency of mishaps involving the dispersal of living material among horticulturists and botanical gardens, there is always the chance of a mislabeled plant.

Still, while it is not possible to confirm the true nature of Koch and Augustin's plant, it is certain that the plant which Schott illustrated in detail (Icones #2478, #2480, #725, and #726) really does represent the plant currently being called *P. pterotum*. Although this Central American species does not occur in Venezuela, Koch and Augustin seemed

not to state that the plant received from Venezuela had been collected there, only that it had been received from Warszewicz in Venezuela.

Krause (1913), in his revision of *Philodendron*, cited only a Wendland collection from Costa Rica. That collection is neither at Göttingen (GOET) nor Berlin (B) and must be lost. Since Engler reported the Wendland collection to be alive at the Berlin Botanical Garden and since Wendland made much of his material available to Schott, it might have been that it was the Wendland material from Costa Rica that Schott illustrated, rather than the Koch material received from Venezuela.

Whether these two elements corresponded to the same species is conjectural, but Schott would have conceivably had the opportunity to compare both, and it is reasonable to assume that he made the correct interpretation and that his use of *P. pterotum* for the Central American species is correct. In any event, there is a need for a neotype since no specimens actually seen by Koch, Schott, or Engler and Krause still exists. Therefore, a modern collection has been chosen here.

*Philodendron pterotum* may be confused with *P. findens*, but that species differs in having blades that dry usually blackened and promptly split pinately into segments.

*Additional specimens examined.* COSTA RICA. **Alajuela:** Cañas-Upala, 13.8 km N of Bijagua, 100–150 m, *Croat 36448* (MO); Llanura de San Carlos, 18–22 km N of Aguas Zarcas, 60 m, 10°31'N, 84°24'W, *Burger & Stolze 5186* (CR, F, US). **Heredia:** La Selva Field Station, *Grayum 2064* (MO); *Hammel 8150* (MO). **Puntarenas:** Gollito, 90 m, 8°39'N, 83°11'W, *Croat 67613* (MO, US); N of Palmar Norte, trail to Jalisco, 50–700 m, *Croat 35170* (MO); Palmar Norte–Panamanian border, 110 m, *Croat 79197* (CR, INB, MO); San Vito de Coto Brus–Neily, Fila de Cal and Cuesta Fila de Cal, 300–600 m, 8°41'N, 82°56.5'W, *Hammel 14161* (MO); Carara Reserve, Quebrada Bonita, ca. 35–80 m, 9°47'N, 84°36'W, *Grayum et al. 5721* (CR, MO); Isla del Caño, 40 km NW of Corcovado National Park, *Gómez 19963* (MO); Corcovado National Park, 0–39 m, 8°42'N, 83°52'W, *Janzen 11546* (MO); *11600* (MO); 0–200 m, 8°29'N, 83°36'W, *Liesner 2850* (CR, MO); 1–10 m, *Kernan & Phillips 1026* (CR, MO); Osa Peninsula, ca. 5 km W of Rincón de Osa, 50–200 m, 8°42'N, 83°31'W, *Burger & Gentry 8967* (CR, F, MO, NY); Quebrada Agasbuena–Quebrada Banegas, ca. 5 km W of Rincón de Osa, 300–400 m, 8°42'N, 83°33'W, *Grayum 4066* (CR, MO); vic. Boscosa, *Croat & Hannon 79245* (INB, MO); Río Claro, along Inter-American Highway, 30 m, *Croat 32945* (MO). **San José:** San Isidro del General–Dominical, 9 mi. SW of Río Pacuar, 680 m, *Croat 35372* (MO); Puriscal, Z.P. La Cangreja, 800 m, *Morales 2031* (CR). NICARAGUA. **Zelaya:** Siana–Matagalpa, ca. 12.9 km before Caño Piedra del Balsamo, <200 m, *Stevens 8812* (MO); near Bil Tingnia, 6 km NW of Bonanza, 150 m, *Neill 3995* (MO); Cerro Baká, ca. 6.5 km E of Río Coperna, 200–300 m, 13°40'N, 84°30'W, *Pipoly 4927* (MO); *4844* (MO); Cerro Waylawás, ca.

100–200 m, ca. 13°38'–39'N, 84°48'–49'W, *Pipoly 4201* (MO); Siuna–Empalme, Caño Calcamo, ca. 5 km al SE de Siuna, ca. 13°40'N, 84°45'W, *Grijalva & Burgos 1536* (MO); El Empalme–Limbaika, ca. 65 m, ca. 13°39'N, 84°24'W, *Stevens 12902* (MO); Cerro Livico, 7 km NE of Siuna, 500 m, *Neill 3633* (MO). PANAMA. Bahía Soldado, *Couell 224* (NY). **Canal Area:** Gatún–Piña, ca. 3 km S of Piña, ca. 50 m, *Croat 36929* (MO); Gatún Lake, *Hutchison & Wright 2885* (BH, UC, US); Barro Colorado Island, *Elmore X20* (F, RSA); *Fairchild 3081* (US); *Croat 10903* (MO); *10265* (MO, SCZ); *7143* (MO); *6640* (MO, PMA); *6581* (MO, SCZ); *5136* (MO); *Bailey & Bailey 328* (BH); along road between Gatún Locks and Fort Sherman, ca. 3 mi. W of Gatún Locks, 1.4 mi. E of Ft. Sherman, 9°18'N, 79°38'W, *Croat 69860* (CM, MO); Summit Gardens, *Croat 10792* (MO, SCZ). **Chiriquí:** Puerto Armuelles–San Bartolomé Límite, 7 mi. W of Puerto Armuelles, ca. 120 m, *Croat 35044* (MO); 1.6 m W of Puerto Armuelles, ca. 50 m, *Croat 21933* (MO). **Colón:** 4 km E of Buena Vista, Quebrada Ancha, 80 m, *Nee 7781* (MO, US); Portobelo–Nombre de Dios, 1.2 mi. beyond the junction of the road to Isla Grande, 79°35'W, 9°40'N, *Croat 49810* (MO); Santa Rita Ridge Road, 6.5 mi. E of Boyd–Roosevelt Highway, 370 m, 9°21'15"W, 79°44'W, *Croat & Zhu 76965* (MO). **Panamá:** 26.8 km E of Bayano bridge, *Folsom 3529* (MO).

***Philodendron purpureoviride* Engl., Bot. Jahrb.**

Syst. 26: 526. 1899. TYPE: Ecuador. Guayas:

Balao, *Eggers 14710* (holotype, B). Figures 317–320.

Hemiepiphytic; stem appressed-climbing, scandent, often pendent, green becoming brownish to gray-green; internodes glossy, to 25 cm long, (0.8)1–2 cm diam., longer than broad, gray-green, semiglossy, ± terete, epidermis drying light yellow-brown, conspicuously fissured or ridged but smooth, frequently flaking free; roots thin, ± twisting or sinuous, few per node; cataphylls 10–29 cm long, unribbed to obtusely 1-ribbed or bluntly to sharply 2-ribbed, cream to medium green, magenta speckled, quickly deciduous, fragile; **petioles** 9–24 cm long, 8–10 mm diam., ± terete, somewhat spongy, somewhat flattened adaxially, surface semiglossy to glossy, frequently fissured, medium green, sometimes maroon-spotted; **blades** narrowly ovate-cordate, subcoriaceous, concolorous or weakly bicolorous, acuminate to long-acuminate at apex (the acumen inrolled), cordate at base, 12–23(27) cm long, 7.4–19 cm wide (1.3–1.8 times longer than wide), (0.8–2.2 times longer than petiole), broadest just below point of petiole attachment, upper surface drying yellow-green, semiglossy, lower surface glossy; anterior lobe 10–21 cm long, 7.4–19 cm wide (2.5–4.2 times longer than posterior lobes); posterior lobes (2.8)3.6–7(8.2) cm long, (3.1)4.3–7.7 cm wide; sinus hippocrepiform; midrib convex to broadly convex and slightly paler above, convex to broadly convex and paler below; basal

veins ca. 3 per side, with 0–1 free to base, 2–3 coalesced to 5 mm long, flattened to raised; posterior rib 0.5–1 cm long, never naked; primary lateral veins about 3(4) per side, departing midrib at a 55–65° angle, ± straight to the margins, convex to weakly raised above, convex below; interprimary veins drying darker than surface below; tertiary veins ± obscure to visible and darker than surface below; minor veins fine below, arising from both the midrib and primary lateral veins; “cross-veins” conspicuous (in Central America). INFLORESCENCES erect, 1 per axil; peduncle (3.5)5–7(11.5) cm long, 5–7 mm diam., subterete, purplish tinged, whitish streaked; **spathe** 11–15 cm long (1.2–3(4.3–4.7) times longer than peduncle), constricted only slightly midway above the tube; spathe blade green to greenish white, with reddish speckling outside, 7–7.5 cm long, pale greenish cream inside; spathe tube violet-purple, short-lineate outside, 6.7–7 cm long, dark violet-purple inside; **spadix** sessile; ca. 13 cm long; pistillate portion white to pale greenish white, 4.5–5 cm long, 1.5 cm diam. throughout; staminate portion 8.5–9.2 cm long; fertile staminate portion white, drying reddish brown, 1.2 cm diam. throughout; sterile staminate portion 1.5 cm diam.; pistils 7.5 mm long, 1.6 mm diam.; ovary 4–5-locular, 6.2 mm long, 1.6 mm diam., with axile placentation; locules ca. 6.2 mm long; ovules 15–25 per locule, 0.2–0.25 mm long, 2–3-seriate, style similar to style type D; style apex with low style boss. INFRUCTESCENCE with pistillate spadix 6.5–7.5 cm long, 2.5–3.5 cm wide; berries 5.9 cm long, 2.5 cm diam.; seeds 24–25 per locule, yellow-orange, 1.2 mm long, 0.5 mm diam., thin and faintly striate.

Flowering in *Philodendron purpureoviride* is apparently aseasonal with post-anthesis material collected virtually year-round. Post-anthesis or early fruiting collections have been made in every month except September, but mature fruits have been collected only in January and August.

*Philodendron purpureoviride* ranges from Costa Rica and Panama to the Pacific slope of Colombia and Ecuador (to Los Ríos and Guayas Provinces) from sea level to 1600 m elevation in *Premontane rain forest*, *Tropical wet forest*, *Tropical wet forest transition*, *Premontane wet forest*, and *Premontane wet forest*. In Central America, it is known primarily from the Pacific slope of Costa Rica and adjacent Panama, but also from the Atlantic slope in both countries.

*Philodendron purpureoviride* is a member of *P.* sect. *Philodendron* subsect. *Solenostergma*. This species is reportedly the dominant climber in west-

ern Ecuador in forests around San Sebastián, south of Jipijapa in the province of Manabí (A. Gentry, pers. comm.).

*Philodendron purpureoviride* is recognized by its scandent habit; stems with the epidermis drying yellow-brown, conspicuously exfoliating and frequently fissured; terete to somewhat flattened petioles about four-fifths as long as the blades; narrowly ovate-cordate blades drying yellow-green; solitary inflorescences, with the spathe tube violet-purple on both surfaces.

*Philodendron purpureoviride* is closest to and perhaps inseparable from *P. lechlerianum* Schott from Peru. The latter species is known from the type collected at "San Gaván" (San Gabon) in the Department of Puno, Carabaya Province, as well as a more recent collection (*D. N. Smith* 6386) from Pasco Department. *Philodendron lechlerianum* differs in having more conspicuous and prominent cross-veins. If these names prove to be synonymous, *P. lechlerianum* is older and would extend the range of the species into the Amazon basin.

Among sympatric species, *Philodendron purpureoviride* is most easily confused with *P. hederaceum*, also a vine with ovate-cordate, greenish-drying blades. The latter species is distinguished by having stems usually drying green or dark brown and without a peeling epidermis. In addition, the blades are more coriaceous, typically more broadly ovate with more (four to six) pairs of basal veins (vs. two to three pairs for *P. purpureoviride*), and generally have more prominent primary lateral veins. *Philodendron purpureoviride* may also be confused with some sterile specimens of *P. wilburii*. The latter species has much longer peduncles and stems drying darker brown and more closely fissured with the epidermis seldom exfoliating.

Central American material of *P. purpureoviride* differs from the Ecuadorian type by having conspicuous "cross-veins" throughout most of the blade. The Ecuadorian material, by contrast, has less conspicuous minor veins which lack "cross-veins" except at or near the margins. In addition, Engler's description of the inflorescence was based solely on Eggers's field label. Since the holotype is sterile and no fertile material associated with Eggers's original collection is apparently extant, further studies should investigate whether these specimens all represent the same species.

*Additional specimens examined.* COSTA RICA. **Alajuela:** 3.5 km W of Fortuna, 2.5 km NW of New Volcán Arenal, 1500 m, 10°28'N, 84°41'W, *Taylor & Taylor* 11706 (MO, NY, US); Cahas-Upala, 4 km NNE of Bijagua, 400 m, *Croat* 36267 (MO); Upala, 600 m, *Rivera* 1559 (INB, MO). **Cartago:** 12 km S of Turrialba by air,

4 km SE of Pejibaye along Río Gato, 700 m, 9°48'N, 83°42'W, *Liesner* 14355 (CR, MO); Río Reventazón, Turrialba, 500–600 m, 9°53.5'N, 83°38.5'W, *Grayum & Schatz* 5242 (CR, MO); Tacurrique, Las Vueltas, 635–700 m, *Tanduz* 13312 (US). **Limon:** Turrialba–Limon, along Highway 32, ca. 11 mi. S of Siquirres, 650 m, *Croat* 43332 (MO); Río Telire, Bajo Telire, 400–600 m, *Gómez* 24119 (MO). **Puntarenas:** Zona Protectora Las Tablas, Parque Internacional La Amistad, Finca Cafrosa, 1600–1800 m, 8°53'20"N, 82°50'30"W, *Mora* 139 (CR, MO); 1680 m, *Alfaro & Navarro* 29 (INB); Cerro Anguciana, 950–1150 m, 8°49'18"N, 83°11'15"W, *Grayum* 10647 (CR, MO); Palmar Norte to Jalisco, 780–960 m, 8°59.5'N, 83°28'W, *Grayum* 9141 (CR, F, K, MO, US); 50–700 m, *Croat* 35203 (MO); Las Cruces–Neily, Finca de Cal, 1000–1400 m, *Gómez* 19635 (MO, US); Cantón Golfito, 100–500 m, *Morales et al.* 1903 (CR, INB); 9 km W of La Palma, along Río Rincón, *Grant & Rundell* 92-02203 (CR, MO); Golfo Dulce area, Cantón de Osa, vic. of Esquinas Experiment Station, 0 m, *Allen* 5370 (MO, UC, US); W of Rincón de Osa, 250–540 m, 8°42'N, 83°31'W, *Croat & Grayum* 59857 (CAS, CM, CR, MO, NY); Villa Briceno–Golfito, Fila Gamba, ca. 6 km from Golfito airport, <100 m, 8°41'30"N, 83°12'W, *Croat* 59902 (CR, MO); Parque Nacional Corcovado, Dos Bracos de Río Tigre, Jiménez, along Río Madrigal, 600 m, 8°29'50"N, 83°28'55"W, *G. Herrera* 4728 (CR, MO). **San José:** San Isidro del General–Dominical, 9 mi. SW of Río Pacuar, 680 m, *Croat* 35348 (MO); Carara Reserve, W Montañas Jamaica, ca. 3 km NE of Bijagua de Turruabares, 500–600 m, 9°45'30"N, 84°33'W, *Grayum et al.* 5851 (CR, F, MO). **PANAMA.** **Bocas del Toro:** Fortuna Dam area, near road to Chiriquí Grande, 650 m, 8°45'N, 82°15'W, *McPherson* 9925 (MO). **Chiriquí:** Burica Peninsula, 11 mi. W of Puerto Armuelles, vic. of San Bartolomé Limite, 100–500 m, *Liesner* 84 (F, MO, US); 450 m, *Basey* 595 (F, MO); "Ojo de Agua," Finca Hartmann, vicinity of Santa Clara (between Volcán and Río Sereno), 1520–1750 m, 8°50'N, 82°45'W, *Croat* 66290 (MO, PMA, US); ca. 13 km from Río Sereno, *McPherson & Richardson* 15968 (B, K, MEXU, MO, PMA, US). **Coelá:** El Valle region, La Mesa, above El Valle de Antón, 860–900 m, *Croat* 37421 (MO); Finca Macarenita, 800 m, 8°36'N, 80°07'W, *Croat & Zhu* 76677 (MO). **Darién:** Cerro Pirre region, Cuna, 500–600 m, *Croat* 37631 (MO); W slope along Río Periseno, 110 m, 8°01'N, 77°44'W, *Croat & Zhu* 77117 (MO). **Veraguas:** Santa Fe area, between Santa Fe and Calovébora, 1.7 mi. past Alto Piedra School, 570 m, 8°33'N, 81°08'W, *Croat & Zhu* 76865 (MO).

***Philodendron purulhense* Croat, sp. nov. TYPE:**

Guatemala. Alta Verapaz: El Progreso–Cobán, on Hwy. CA-14, 2–3 mi. S of Purulhá, 1500–1720 m, 15°13'S, 90°12'W, 21 July 1977, *Croat* 41752 (holotype, MO-2582045; isotype, GUAT). Figures 321–323, 325, 326.

Planta hemiepiphytica aut raro terrestria; internodia brevia, usque 6 cm diam.; cataphylla 20–30 cm longa, 2-costata, acute D-formata, persistentia semi-intacta; petiolus subteres, aliquantum spongiosus, obtuse complanatus adaxialiter, 32–52 cm longus, 1.5–2 cm diam.; lamina ovata vel ovato-cordata, 25–48 cm longa, 18.5–40 cm lata, in sicco atribrunnea vel nigrescens; nervis basalibus liberis ad basim; inflorescentia 1; pedunculus 2.5–7 cm longus; spathe 10–17 cm longa; lamina spathe extus vir-

idi, breve lineata; intus viridi; pistilla 6-7-locularia; loculi 13-20-ovulati.

Hemiepiphytic or rarely terrestrial, growing 3-6 m high in trees; stem appressed-climbing or creeping; internodes short, semiglossy, to 6 cm diam., broader than long, dark green, epidermis intact; roots moderately dense, drying dark brown, ca. 3 mm diam.; cataphylls 20-30 cm long, 2-ribbed, sharply D-shaped, persisting semi-intact with brownish fibers at upper nodes; **petioles** 32-52 cm long, 1.5-2 cm diam., subterete, somewhat spongy, obtusely flattened adaxially, surface semiglossy, short-lineate, usually drying brown to blackened; **blades** ovate to ovate-cordate, subcoriaceous, slightly bicolorous, acute to acuminate at apex, cordate at base, 25-48 cm long, 18.5-40 cm wide (1.2-1.97 times longer than wide), (0.6-0.98 times the petiole length), about equal in length to petiole, upper surface somewhat silvery, semiglossy, lower surface drying dark brown to blackish; anterior lobe 20-38 cm long, 18.7-38 cm wide (2.1-3.1(3.6) times longer than posterior lobes); posterior lobes rounded, 7-17 cm long, 8-17 cm wide; sinus usually hippocrepiform, 12 cm deep; midrib flat, slightly paler than surface above, convex, paler than surface below; basal veins 4-7(8) per side, with all free to base, third and higher order veins coalesced 1.5-4 cm long; posterior rib naked for 1.5 cm; primary lateral veins 3-6 per side, departing midrib at a 35-40° angle, ± straight to the margins, sunken above, convex and paler than surface below; minor veins moderately distinct below, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 1 per axil; peduncle 2.5-7 cm long, 8-10 mm diam., whitish streaked; **spathe** 10-17 cm long (2.4-3.4(4) times longer than peduncle), constricted slightly above the tube, ± ellipsoid; spathe green, short-lineate outside, reddish purple throughout inside; spathe tube 5 cm long; **spadix** sessile; 10-13 cm long; pistillate portion 4-4.5 cm long, 1.7-2.3 cm diam. at base, 2-2.7 cm diam. midway, 1.7-1.9 cm diam. near apex; staminate portion 8-8.5 cm long; fertile staminate portion 1.8-2 cm diam., sterile staminate portion usually much broader than pistillate portion at anthesis, weakly tapered toward apex, scarcely or not at all constricted above the sterile portion, 1.4-1.8 cm diam. midway, 11-12 mm diam. 1 cm from apex, bluntly acute at apex; sterile staminate portion not very obvious, ca. 1.3 cm long, abruptly contracted at base, as broad as or narrower than the pistillate portion; pistils 6.2 mm long, 3.4 mm diam.; ovary 6-7-locular, 5 mm long, 3 mm diam., with axile placentation; locules 5 mm long, 1 mm

diam.; ovules 13-20 per locule, somewhat transparent, 1 mm long, much longer than funicle; funicle 0.2 mm long, adnate to lower part of partition, style 1 mm long, 3 mm diam., similar to style type D; style apex ± rounded; style boss broad, pronounced; stigma 1-1.1 mm diam.; the androecium truncate, oblong, prismatic, margins irregularly 4-5-sided, 1 mm long, 2-2.5 mm diam. at apex. **INFLORESCENCE** with pistillate spadix 4.5-5 cm long, 3-4 cm diam.; berries 1 cm long, 4 mm diam.; seeds 6-8 per locule, oblong-ellipsoid, 2.2-2.5 mm long, 0.7-0.9 mm diam., sticky.

The flowering phenology in *Philodendron purulhense* is poorly known. No flowering collections have been made, and only one post-anthesis collection is known (July), but immature fruiting collections are known from March, July, September, October, and November. This might suggest that the species flowers throughout at least a substantial part of the rainy season and perhaps also in the dry season (December through April).

*Philodendron purulhense* ranges from southern Mexico (Chiapas) to Guatemala (Alta Verapaz) and Honduras (Cortés and Olancho), at 1360 to 1870 m elevation in *Tropical Lower Montane wet forest* and *Premontane wet forest* life zones.

*Philodendron purulhense* is a member of *P. sect. Philodendron* subsect. *Philodendron ser. Fibrosa*. This species is characterized by its short internodes; sharply D-shaped to two-ribbed cataphylls persisting semi-intact; subterete petioles (about as long as the blades), dark brown to blackish-drying; ovate-cordate blades with basal veins free or weakly coalesced and scarcely or not at all naked on the usually hippocrepiform sinus; solitary, short-pedunculate inflorescence, more or less elliptic green spathe; and ovaries with 13-20 ovules per locule.

*Philodendron purulhense* may be confused with *P. advena* because leaves of the latter species dry a similar color. *Philodendron advena* differs in having long internodes, deciduous cataphylls, more or less sagittate blades, and one to three (four) ovules per locule.

*Additional specimens examined.* GUATEMALA. **Alta Verapaz:** Tactic, Río Frío, 1400-1500 m, Standley 90510 (F); Tactic-Tamahá, 1500-1600 m, Standley 90820 (F); Río Carchá, Cobán-San Pedro Carchá, ca. 1360 m, Standley 89902 (F). **Baja Verapaz:** Highway CA-14 to Cobán, Croat 41189 (MO); Biotopo del Quetzal, WNW of Purulhá, 1500-1600 m, Stevens et al. 25446 (MO); Mpio. Purulhá, El Progreso-Cobán, 1620-1720 m, 15°13'N, 90°12'W, Croat & Hannon 63765 (CAS, CM, K, L, MEXU, MO, US, USCG). HONDURAS. **Cortés:** Parque Nacional Cusuco, Río de Cusuco, ca. 22 km W of San Pedro Sula, 1500 m, 15°30'N, 88°13'W, Evans 1490 (EAP, MO). **Ocotepaque:** Cordillera de Celaque, 3 mi. N of Belén Gualcho,

1870 m, 14°30'06"N, 88°48'02"W, *Davidse et al.* 35340 (MO), MEXICO, Chiapas: Mpio. Jitotol, 5 km SE of Jitotol, along road to Bochil, 1600 m, *Breedlove & Davidse* 55077 (MO); Mpio. La Trinitaria, 4 km E of Laguna, near Dos Lagos, 1300 m, *Breedlove* 38810 (CAS); Mpio. Las Margaritas, 12 km E of Tzucaco, 1200–1300 m, *Davidse et al.* 29865 (MO).

***Philodendron radiatum*** Schott, Oesterr. Bot. Wochenbl. 3: 378. 1853. TYPE: Mexico (holotype lost). Schott ic. 2623 (neotype, here designated). Figures 1, 324, 331–335.

Epiphytic or hemiepiphytic, rarely terrestrial; stem appressed-climbing, creeping if terrestrial, sap clear to orange, watery, sticky, leaf scars conspicuous, 2–4(7.5) cm long, 2.5–6(7) cm wide; internodes thick, 3–7(12) cm long, (1)3–8 cm diam., about as long as broad or longer than broad, dark green to gray-green, sometimes scurfy and light brownish tan, transversally lined (raised 2 mm); roots 4 mm diam., with swollen nodes and branched tips; cataphylls to 38 cm long, soft, bluntly to sharply 2-ribbed, rarely unribbed, green, sometimes conspicuously reddish-lineate, deciduous; **petioles** 28–108 cm long, (2–3)4–17 mm diam., terete or subterete to obtusely flattened adaxially, dark green, surface dark greenish or to greenish red-lineate; sheath persisting, sometimes moderately spongy; **blades** ± triangular-ovate in outline, (15)27–101 cm long, (11)25–90 cm wide (ca. 0.8–2.1 times longer than wide), (0.5–1.7 times longer than petiole), broadest at point of petiole attachment, thinly coriaceous, weakly to moderately bicolorous, semiglossy, acuminate to long-acuminate at apex, upper surface dark green, lower surface slightly paler, usually drying yellow-brown to reddish brown on both surfaces; margins weakly incised-lobate to usually deeply incised-lobate to within 1–8 cm of midrib, rarely entire and sagittate; apex often acute, sometimes acuminate, very short acuminate or ± rounded (the acumen tightly inrolled if present, 1–8 mm long), lobed-cordate at base; basal segments pinnatifid, lateral segments entire, sinuate or the lowermost of the anterior lobe pinnately lobed with 1–2 lobes on each side, final divisions linear-lanceolate in shape (0.9–2.1 times longer than wide), segment apex obtuse to broadly obtuse; interlobal sinuses 0.7–0.97 the length of the lobes; basal sinus hippocrepiform to obovate or closed, 3–15 cm deep; midrib slightly raised to convex, weakly reddish green-lineate, concolorous to paler than surface above, convex to prominently raised to round-raised below; basal veins 4–5(8) per side, directed into the segments of the posterior lobe, 0–1 free to base, (1)2–4 coalesced to 3 cm long, naked 1–6 cm; posterior rib absent; primary lateral veins 3–8 per side, departing midrib at a 50–60° angle, straight to mar-

gin, weakly raised above, raised to convex and paler than surface below; minor veins moderately visible, paler and slightly raised below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, 1–3(4) per axil; peduncle 2–12 cm long, (3)5–10(19) mm diam.; **spathe** erect, 11–25 cm long (1.3–6.3 times longer than peduncle), obscurely striate, margins paler; spathe blade cuspidate at apex, pale green to yellowish green, semiglossy, sparsely purplish-dotted outside, pink to dull red or pale brownish (post-anthesis) inside; spathe tube dark purple-violet or dark green, sometimes tinged purple-violet outside, 4 cm diam., dark purple-violet inside; **spadix** sessile to weakly stipitate, cylindrical to weakly tapered; staminate portion creamy white, protruding forward at anthesis, pointed at apex, 10–17 cm long; pistillate portion whitish, weakly obovoid, 3.5–5.2 cm long (1 cm shorter on back side), 1.1–1.8 mm diam. at apex, 1.5–1.9 mm diam. at middle, (7)17 mm wide at base; staminate portion 6.3–8.1 cm long; fertile staminate portion tapered, 1–1.9 cm diam. at base, 1.2–1.6 cm diam. at middle, 0.9–1.4 cm diam. ca. 1 cm from apex, broadest at the base, slightly constricted above the base; sterile staminate portion as or slightly broader than the pistillate portion, white with tan ring around apex, 0.7–2 cm diam.; pistils (1.6)4–5(8) mm long, (1)2.8–3.2(5.2) mm diam.; ovary (7)8-locular, (1.6)3.3–4(6.7) mm long, 1.5–3(5.2) mm diam., with axile or sub-basal placentation; locules (0.6)1.9–3.4(6.5) mm long, 0.5–0.8(1) mm diam.; ovule sac 1.8(3.5) mm long; ovules 8 per locule, 1–2-seriate, contained within translucent, gelatinous ovule sac, 0.1–0.3 mm long, longer than funicle; funicle 0.1–0.2 mm long, style 0.3–0.5 mm long, 1.2–3 mm diam., similar to style type B (rarely C); style apex flat to weakly rounded, sometimes domed; stigma usually hemispheroid, sometimes subdiscoid, pink, 1.2–1.5 mm diam., 0.6–0.7 mm high, covering entire style apex; the androecium truncate, oblong, prismatic, margins irregularly 3–5-sided, sometimes weakly scalloped, 1 mm long, (0.7)2.5–2.7 mm diam. at apex; thecae cylindrical, 0.3 mm wide, ± parallel to one another and contiguous; sterile staminate flowers usually truncate, sometimes clavate or irregularly 5–6-sided, 1.1–2.9 mm long, 0.6–1.8 mm wide. INFRACTESCENCE 11–17 cm long, 4.5 cm diam.; pistillate spadix 6–8 cm long, 2.5–3.5 cm diam.; berries white, somewhat translucent, 4 per locule, oblong, sticky.

*Philodendron radiatum* ranges from Mexico (San Luis Potosí to Chiapas and on both Atlantic and Pacific slopes in Chiapas) to Colombia (Antioquia), from sea level to usually no more than 700 m (rare-

ly 1250 to 1860 m; most collections are from below 100 m) elevation. It is perhaps more widespread in South America than collections indicate (only two are known). The plants are very large and difficult to collect and may have been overlooked in other areas.

This species is highly variable morphologically and ecologically versatile as well. In Mexico, it occurs in "Selva Alta Perennifolia," "Selva Mediana," and "Selva Baja Caducifolia" in mesic areas on the Atlantic slope as well as in "Selva Mediana Subperennifolia" and "Selva Baja Subperennifolia" in the drier Yucatán Peninsula. On the Pacific slope it also occurs in areas of "Bosque Pino-Encino." In Central America, this taxon occurs principally in *Tropical moist forest* but also in *Premontane wet forest* and drier parts of *Tropical wet forest*.

*Philodendron radiatum* is a member of *P. sect. Polytomium*. *Philodendron radiatum* has two varieties, the typical variety having more deeply divided leaf blades with the segments often incised almost to the midrib and variety *pseudoradiatum* with the blades only weakly incised-lobate (less than half the distance to the midrib). While the latter variety is restricted to southwestern Chiapas, even populations of the typical variety on the western slope of Central America have less deeply incised-lobate blades than those on the Atlantic slope. There is also considerable clinal variation in the width of the divisions throughout Central America, with plants in Mexico having, on average, broader pinnae (averaging 4.5 cm wide) than those in Panama, for example, where the pinnae average only about 3 cm wide. Leaves of Mexican populations of *P. radiatum* also consistently lack free basal veins, whereas elsewhere in Central America such veins are present. In western Mexico and Guatemala, plants are also more likely to have longer internodes and to occur more frequently on rocks or creeping over the ground than plants on the more mesic Atlantic slope of Central America.

KEY TO THE VARIETIES OF *P. radiatum*

- 1a. Blades deeply lobed, the divisions near the middle of the blade extending more than  $\frac{1}{2}$  the way to the midrib ..... var. *radiatum*
- 1b. Blades shallowly lobed, the divisions near the middle of the blade extending less than  $\frac{1}{2}$  the way to the midrib ..... var. *pseudoradiatum* (Mutuda)  
Croat

*Philodendron radiatum* Schott var. *radiatum*

*Philodendron angustatum* K. Koch, in A. Braun et al.,  
Append. sp. Hort. berol. 1853: 4. 1853-1854.  
TYPE: Central America. Without exact locality (ho-

lotype, B? lost). Schott ic. 2605 (neotype, here designated).

*Philodendron polytomum* Schott, Bonplandia 7: 164. 1859. TYPE: Mexico. Veracruz: Colipa, Hac. de Sta. Bárbara, Liebmann s.n. (holotype, C; isotype, K not seen).

*Philodendron impositum* Schott, Prodr. Syst. Aroid. 291. 1860. TYPE: Costa Rica, *Wendland* s.n. (not seen).

Internodes 3-7 cm long, 3-8 cm diam.; cataphylls bluntly to sharply 2-ribbed; petioles 33-108 cm long; blades deeply lobed, the divisions near the middle of the blade extending more than  $\frac{1}{2}$  the way to the midrib, 21-101 cm long, 25-90 cm wide; posterior lobes (8) 12.5-25 cm long, (9) 16-18 cm wide; primary lateral veins 3-8 cm long; basal veins 4(5) per side, (1) 2-4 coalesced to 3 cm long, naked 1-6 cm. INFLORESCENCES 1-3(4) per axil; spathe 11-25 cm long; spadix 10-13 cm long; pistillate portion 3.5-5.2 cm long; pistils (3) 4-5(8) mm long; ovaries with axile placentation; style similar to style type B (rarely C).

Flowering in *Philodendron radiatum* var. *radiatum* is documented by few flowering collections made in February, August, and September. Post-anthesis collections are common, from every month of the year, and there are no clearly marked differences in the phenology of this species in different parts of Central America. Material was seen at various stages of development from all parts of Central America in about equal numbers. Mature fruits are known only from February and May.

*Philodendron radiatum* var. *radiatum* is characterized by its moderately thick, mostly short internodes; sharply two-ribbed, deciduous cataphylls; terete petioles; and especially by its deeply incised-lobate leaf blades that mostly linear to linear-lanceolate segments, themselves often lobate toward the base of the blades.

*Philodendron radiatum* var. *radiatum* might be confused with *P. dressleri* and *P. warszewiczii*, Central American species with incised-lobate leaf blades. *Philodendron dressleri* differs in having thicker leaf blades, more succulent stems with shorter internodes, and especially by having blades that are incised lobate only about midway to the midrib. In addition, *P. dressleri* has one inflorescence per axil and 3-4 locules per ovary, while *P. radiatum* has 1-3(4) inflorescences per axil and 7-8 locules per ovary. *Philodendron warszewiczii* differs by having thinner blades, drying papryaceous with the minor veins distinctly visible. In addition, the lateral leaf blade segments usually have three or more lobes per side. *Philodendron radiatum*, in contrast, has blades drying subcoriaceous with the minor veins rather indistinct and with the

lateral leaf blade segments usually 1–2-lobed per side.

*Philodendron radiatum* var. *radiatum* has a unique attractant, whose odor profile is made up almost entirely of compounds unique to it, and is pollinated by a new species of *Cyclocephala* (Schatz, 1990).

*Additional specimens examined for P. var. radiatum.* BELIZE. Honey Camp, *Lundell s.n.* (US); Maskall, *Gentle 1141* (MICH, NY); Gracie Rock, 1.5–4 mi. S of Mile 22 on Western Highway, 100 m, *Liesner & Dwyer 1460* (MO); Crooked Tree Village, 60 m, 17°47'N, 88°32'W, *Davidse & Brant 33146* (MO). Cayo: Roaring Creek, *Dwyer 12677* (MO); 1.5 mi. W of Augustine, Río Frío, ca. 450 m, *Sutton et al. 211* (BM); Hummingbird Highway, Mile 20, *Dwyer & Liesner 12130* (MO); Chiquibul National Park, Caracol Archaeological Reserve, 550 m, 16°45'N, 89°07'W, *Ingram & Ferrell-Ingram 1935* (MO); Río Ma Cal, San Luis-Cuevas, 1050 ft., *Croat 23534* (MO); Yaca, *Gentle 2552* (MICH). COROZAL: edge of New River, Pueblo Nuevo, *Gentle 473* (CM, MICH); Cerro Maya Ruins, Lowry's Bright, *Crane 348* (LL); *Gentle 617* (MICH). **Orange Walk:** road to Trinidad, ca. 5 km N of August Pine Ridge, 100 m, 18°N, 88°42'W, 100 m, *Davidse & Brant 32791* (MO); Pulltrouser Swamp, *Lincoln 20* (MO). **Toledo:** Punta Gorda highway, 1 mi. E of junction to San Antonio, ca. 100 ft., *Croat 24512* (MO); Solomon Camp, vic. of Richardson Creek and Bladen Branch junction, Maya Mountains foothills, 80–420 m, 16°32'–33'N, 88°45'–46'W, *Davidse & Brant 32198* (MO); Columbia Forest Station Area, 1.5 mi. S of Mayan Village of San José, *Croat 24335* (MO); *Holst 4303* (MO). COSTA RICA. **Alajuela:** Cañas-Upala, 10 km N of Bijagua, 200 m, *Croat 36488* (MO). **Heredia:** La Selva Field Station, *Graham 25344* (DUKE); 100 m, 3022 (DUKE); *Jacobs 2290* (DUKE); 2707 (MO); 10°26'N, 84°W, *Burger & Stolze 5921* (BM, CR, US); 100 m, 10°26'N, 84°01'W, *Gearty & Ortiz 78615* (MO); Río Sucio, near Puerto Viejo, 20 m, *Croat 35680* (MO); Puerto Viejo-Guápiles, 7 km N of Buenos Aires, 10°23'30"N, 83°48'30"W, *Croat 68386* (MO); Río Sarapiquí, S of La Virgen, 200 m, *Lent 33* (MO); La Tirimbina, 220 m, *Hunter 821* (WIS). **Limón:** Barra del Colorado, N side, 0–2 m, 10°47'N, 83°35'W, *Stevens 24133* (CR, MO); Barra del Colorado-Río San Juan, Laguna de Atrás, 5 m, 10°48'–52'N, 83°38'W, *Davidse & Herrera 31512* (CR, MO); Cerro Coronel, E of Laguna Danto, 20–170 m, 10°41'N, 83°38'W, *Stevens 23845* (CR, MO); Parque Nacional Tortuguero, Estación Agua Fría, 2 m, *Robles 1398* (CR, MO); 40 m, 10°26'N, 83°35'W, *1730* (CR, MO); 600 m S, 4 m, 10°32'N, 83°30'W, *1846* (CR, MO); Cahuita, *Poveda 1166* (CR, MO); Cahuita-Limón, 0–10 m, 9°44'N, 83°20'W, *Baker & Burger 157* (CR, NY); Hacienda Tapasco-Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, *Davidson & Donahue 8729* (MO, RSA); 8437 (RSA); Río Colorado, 3.5 air km S of Islas Buena Vista, 10–120 m, 10°39'N, 83°40'40"W, *Davidse & Herrera 31268* (MO); Barra del Colorado, 1–5 m, 10°47'40"N, 83°25'30"W, *Davidse & Herrera 30882* (MO); Río Pacuara, 50–100 m, 10°15'N, 83°29'W, *Burger & Liesner 6894* (MO); 7 km SE of Bribrí, 100–250 m, *Gomez 20315* (MO); 1 mi. SW of Bribrí, <50 m, *Croat 43228* (CR, MO); vic. of Moín, 0 m, 10°N, 83°04'W, *Croat 61207* (MO); Bahía de Portete Parque Nacional, 0–10 m, 10°N, 83°05'W, *Thompson & Rawlins 1175* (CM); Río Reventazón, Finca Montecristo, below

Cairo, 25 m, *Standley & Valerio 48969* (US). EL SALVADOR. La Cebadilla, *Calderón 1245* (US). **San Salvador:** Tonacatepec vicinity, *Standley 19538* (GH, NY, US); *Calderón 201* (GH, NY, US). GUATEMALA. **Alta Verapaz:** *Tuerckheim 8330* (US); eastern portions of Verapaz and Chiquimula, *Watson 194* (GH); near Finca Sepacuite, *Cook & Griggs 724* (US); Cubilquitz, 350 m, *Tuerckheim 8330* (US); Guatemala-El Estor, 5 mi. W of Tzurú, 600 m, *Croat 41509* (MO); Tzurú-El Estor, Finca Argentina, above Papalpa, 15 mi. W of Telemán, 550–650 m, *Croat 41550* (MO); ca. 6 km NE of Panzós, 500 m, *Croat 41616* (MO). **Izabal:** Quirigua vicinity, 75–225 m, *Standley 23939* (US); ca. 7 mi. S of Puerto Barrios, 50 m, *Croat 41803* (MO); Río Frío, Cerro San Gil, 75–150 m, *Steyermark 41534* (MO, NY); W of El Estor and abandoned nickel mine, 1–10 m, *Stevens & Martínez 25283* (MO). **Petén:** La Libertad, *Lundell 2645* (MICH, NY); El Paso, *Lundell 1571* (MICH); Uxactun, *Bartlett 12702* (MICH); Tikal National Park, *Contreras 105* (LL). **Quezaltenango:** along CA-2, 4 mi. NW of turnoff to Coloba, *Croat 32767* (MO); Coatepeque-Retalbuleu, Hwy. CA-2, 3 mi. S of turnoff to Coloba, 600 m, *Croat & Hannon 63414* (MO, US). **Sacatepéquez:** Volcán Santa Clara, 1250–2650 m, *Steyermark 46621* (MO). HONDURAS. Puente Sierra, *Wilson 321* (NY). **Atlántida:** Lantecilla Valley, near Tela, *Standley 52945* (US); 53668 (US); 53985 (US); ca. 10 mi. SE of Tela, Río Lancetilla, 10–150 m, *Croat 42631* (MO); Río Cangrejal, 5 km inland from La Ceiba, *Blackmore & Charley 4155* (BM, MO, UNAH); Río Sambo, Tocoa-La Cieba, 11.3 mi. from Río Cangreja Bridge at La Ceiba, 30 m, 15°47'N, 86°30'W, *Croat & Hannon 64581* (BM, MO); Parque Nacional Pico Bonito, ca. 10 km SW of La Ceiba, 160 m, 15°42'N, 86°51'W, *Evans 1675* (MO). **Colón:** Río Negro, Trujillo, *Clewell et al. 4345* (MO). **Copán:** ca. 4 mi. E of Copán, 800 m, *Croat 42502* (MO); 10 mi. W of Copán, road to La Entrada, 700 m, *Croat 42516* (MO). **Cortés:** Lago Yojoa, Punta del Cacao, 650 m, 14°05'30"N, 87°58'W, *MacDougal et al. 3079* (MO); Ocote Arrancado, 50 km N Lago de Yojoa, *Nelson et al. 5868* (MO, VBD); Puerto Cortés-Guatemalan border, 2–3 mi. SW of Omasa, *Croat 42558* (MO); N of Lago de Yojoa, along old Highway 1, ca. 2–6 mi. from junction with new Highway 1, SW of Santa Cruz de Yojoa, 600 m, *Croat 42745* (MO). **Gracias a Dios:** Río Platano Biosphere Reserve, Las Marías (Baúl Tuk), 30 m, *Knees et al. 2825* (BM). **Morazán:** El Zamorano, 800 m, *Molina 34353* (MO). **Olancho:** El Jocondo-Cerro El Mulato, *Blackmore & Heath 1686* (BM, MO); Gualaco-San Esteban, Río Olancho, 7.4 mi. NE of San Esteban, 540 m, 15°20'N, 85°42'W, *Croat 64365* (K, MO, NY). **Santa Bárbara:** Lago Yojoa, Punta Gorda, 650 m, 14°52'N, 88°W, *MacDougal et al. 3126* (MEXU, MO, NY); 630 m, 14°53'N, 88°W, *Liesner 26769* (MO); 700 m, *Croat 42752* (MO); El Novillo, 640 m, 14°53'N, 88°00'30"W, *Evans 1044* (MO). **Yoro:** Aguán River Valley, Coyoles, *Yunker et al. 8624* (GH, MO, NY, UC, US). MEXICO. Location unspecified, *Reko 3690* (US); *Sandoz 14* (US). **Campeche:** Mpio. Hopelchén, Rancho El Carmen, 33–35 km S of Xmalben on road from Hopelchén to Xpujil, near Xpanzil, 200 m, 18°58'N, 89°20'W, *Sanders et al. 9735* (MO); Huastusco, *Sandalan 14* (US); Patria, *Engler 197* (BM, GH); Santa Leonor, E of Río San Pedro, *Barlow 16/8* (BH). **Chiapas:** San Manuel, ca. 15 km above Palenque, on road to Ocosingo, ca. 500 m, *Madison 7312* (SEL); Escuintla-El Triunfo, 1 mi. N of Escuintla, 100 m, *Croat 43811* (MEXU, MO); Esperanza-Escuintla, *Matado 16662* (F, MEXU); *17788* (NY); 150 m, *18045* (MEXU);



Tapachula-Nueva Alemán, 4 mi. N of Tapachula, 250 m, *Croat* 43794 (MO); Escuintla-El Triunfo, ca. 8.5 mi. NE of Escuintla, 250 m, *Croat* 43820 (MO); Escuintla-Monte Ovando, 2.8 km NW of Turquiz, ca. 100 m, *Croat* 47511 (MO); Acacoyagua, Cerro Ovando, 800-900 m, *Croat* 78548 (CHIP, MO); Huixtla-Motozintla de Mendoza, 4.8 km N of Huixtla, ca. 200 m, *Croat* 47523 (MO); 2 mi. S of Chiapas border, along Hwy. 195, 8 mi. N of Pichucalco, 80 m, *Croat* 40088 (MO); 5 mi. SE of Palenque, on road to Ocosingo, 200 m, *Croat* 40135 (MO); Palenque-Bonampak, 89-90 mi. SW of Palenque, 350-370 m, *Croat* 40217 (MO); 73 mi. SE of Palenque, 460 m, *Croat* 40267 (MO); Ocozacoatlán-Apitae, 20 mi. N of Ocozacoatlán, 700 m, *Croat* 40650 (MO); 5 mi. N of Ocozacoatlán, 1000 m, *Croat* 40543 (MO); Motozintla de Mendoza-Huixtla, 15 mi. S of Motozintla de Mendoza, 900 m, *Croat* 40765 (MO); Mpio. Ixtacomitán, 7 km SW of Ixtacomitán, 250 m, *Breedlove* 45915 (MO); Mpio. Mapastepec, Sierra de Socuassoco, new unfinished road to Tuxtla Gutiérrez, 200 m, 15°31'N, 92°50'W, *Croat* 63382 (B, K, MO, NY, US); Mpio. Ocosingo, Lacanja-Chanzayab, Palenque-Boca Lacantum, 340 m, *Martínez* 15071 (MO); Laguna Ocotallón, 12 km N of Monte Libano, trail to Chancala, 980 m, *Martínez* 17029 (MO); Bonampak, 520 m, *Breedlove* & *Almeida* 58051 (MO); Mpio. Palenque, 8-9 km S of Palenque, 300 m, *Breedlove* & *Strother* 46895 (CAS); 25 km S of Palenque, 300 m, *Breedlove* & *Almeida* 57318 (CAS); Mpio. Tzimol, 15 km S of Comitán, 1200 m, *Breedlove* 53732 (CAS). **Oaxaca:** *Williams* 9173 (US); 14 mi. S of Tuxtepec, *Moore* & *Bunting* 8904 (BH, MO); 8910 (BH); Tuxtepec-Oaxaca, 0.5 mi. S of Valle Nacional, Highway 175, 120 m, *Croat* 39700 (MO); 3 km S of Hidroeléctrica Temascal, 50 m, *Cortés et al.* 875 (MO); Uxpanapa region, Esmeralda-Río Verde, 1.1 mi. S of Esmeralda, 100 m, 17°10'N, 94°45'W, *Croat* & *Hannon* 63234 (MO); Jalpan, 2-3 km E of La Baquilla, Río Santa María, 270-320 m, *Carranza* & *Zamudio* 4548 (MO); ca. 1.5 km E of La Boquilla, 270-320 m, *Carranza* & *Díaz* 4721 (MO). **San Luis Potosí:** Highway 85, 6 mi. NW of Tamazunchale, 250 m, *Croat* 39262 (MO); Tamazunchale, *Edwards* 626 (DS, MO). **Tehuacan:** 30 mi. E of Minatitlán, *Barkley* & *Carr* 36221 (GH); Alemán, 10 mi. NW of Cardenas, *Barlow* 2625 (BH); Mpio. Cárdenas, vic. Cárdenas, *Cosson* 2021 (MO); Mpio. Huimanguillo, km 12.6 de la desviación de Huimanguillo, *Cosson* & *Magaña* 3255 (CAS, NY). **Veracruz:** Huatusco-Puerto Nacional, El Mirador, 21 km E of Huatusco, ca. 1200 m, *Croat* 43974 (MO); *Moore* & *Bunting* 8855 (BH); Playa Escondida, N of Sontecomapan, along Caribbean cliffs, 10-60 m, *Genry et al.* 32610 (MO); Córdoba-Veracruz, San José de Gracia, ca. 750 m, *Croat* 39612 (MO); Highway 180, 6 mi. E of Costacacalcos, *Croat* 40064 (MO); NW of Misantla, 2.4 mi. on road to Martínez de la Torre, ca. 320 m, *Moore* & *Bunting* 8946 (BH); Dto. Papantla, *Kelly* 126 (BH); Mpio. Coatepec, 5 km by air SE of Tuzamapan, 680 m, 19°21'N, 96°50'W, *Nee* & *Taylor* 26026 (NY); Mpio. Cosamalopan, 10 km by air NW of Tres Valles, 25 m, 18°17'N, 96°13'W, *Nee* & *Taylor* 29306 (NY); Mpio. Hidalgotitlán, 1 km SE of Agustín Melgar, 100 m, 17°15'N, 94°33'W, *Nee* 29765 (NY); Mpio. Nolinco, 2 km from Concha al Espinal, 900 m, *Ortega* 630 (MO). **Yucatán:** Izamal, 20°56'N, 89°01'W, *Gaumer* 23200 (GH, NY, US); 1413 (GH, MO); Uayma, Schott 766 (BM). **NICARAGUA.** **Boaco:** Boaco-Campusa, N slope of Cerro Mombachita, 500-900 m, ca. 12°24'-25'N, 85°32'-33'W, *Stevens* & *Grijalva* 14699 (MO); Hwy. 33, vic. Río Quilán bridge, ca. 300-310 m, 12°35'N, 85°32'W, *Stevens* 9330 (BM, MO). **Chontales:**

Juigalpa-La Libertad, ca. 17.4 km NE of Río Mayales, at ford of Río El Bizcocho, 350-400 m, ca. 12°12'N, 85°17'W, *Stevens* 4093 (BM, MO); 4095 (BM, MO); ca. 2.8 km above Caspa, 400-500 m, ca. 12°17'N, 85°23'W, *Stevens* 3671 (MO, PMA). **Estelí:** km 167 on Hwy. 1, ca. 15.8 km E of entrance to Estelí, 825-850 m, ca. 13°15'N, 86°22'W, *Stevens* 5788 (MO). **Jinotega:** Río Bocay, Salto Acatulú, ca. 130 m, 14°13'N, 85°10'W, *Stevens et al.* 16749 (MO). **Madriz:** Cerro Quisúva, 1100-1250 m, ca. 13°30'N, 86°31'W, *Stevens* & *Grijalva* 16063 (BM, MO). **Managua:** Ciudad Managua, Reparto Bolonia, *Guzmán et al.* 1071 (MO). **Nueva Segovia:** Río San Fernando Valley, ca. 5.2 km N of San Fernando, 13°2-3'N, 86°19'-20'W, *Stevens* 3245 (BM, MO). **Río San Juan:** Río Santa Cruz-Caño Santa Cruzita, La Palma, 40-60 m, 11°2-4'N, 84°24'-26'W, *Stevens* 23427 (MO); Caño Chontaleño, 20 km NE of El Castillo, 200 m, *Neill* & *Vincelli* 3620 (MO). **Zelaya:** La Barra de Punta Gorda, 0-2 m, 11°30'N, 83°46'W, *Moreno* 13225 (MO); Caño Montecristo, mouth of Caño El Consuelo, ca. 10 m, 11°35'N, 83°51'W, *Moreno* 15027 (MO); 6.5 km al SE de Waslala, 520-560 m, 13°16'N, 85°24'W, *Moreno* 17288 (MO); Puerto Cabezas-Rosita, ca. km 47, ca. 5.3 km W of Río Wawa Ferry, <10 m, 14°06'N, 83°35'W, *Stevens* 8568 (MO); along road to Panua, 2.3-3.1 km NW of Panua, 4.2-5 km from main road, 30-40 m, 14°18'-19'N, 83°41'-42'W, *Stevens* 7796 (MO); Puerto Cabezas-Río Wawa, Ibo Tingni, drainage of Caño Sung Sung, <10 m, 14°9'-11'N, 83°29'-31'W, *Stevens* 10658-a (MO); Awasstara vicinity, <10 m, ca. 14°19'N, 83°12'-13'W, *Stevens* 10441 (MO); SW of Bluefields, 10-40 m, 12°N, 83°46'W, *Stevens* 19781 (MO); 6.3 km S of bridge at Colonia Yolania, on road to Colonia Manantiales de Nueva Guinea, 200-300 m, *Vincelli* 249 (MO); 40-45 km SW of Waspam, 10-100 m, *Atwood* 3717 (MO); Cerro Livico, 7 km NE of Siuna, 500 m, *Neill* 3634 (BM, MO); Cerro Wayladas, ca. 80 m, 13°39'N, 84°49'W, *Pipoly* 4370 (MO); ca. 100-200 m, 13°38'-39'N, 84°48'-49'W, 4239 (MO); 4162 (MO); ca. 100-268 m, ca. 13°39'N, 84°48'-49'W, *Stevens* 7378 (MO); Caño Costa Rica, ca. 1.8 km SW of Colonia Naciones Unidas, 150-180 m, ca. 11°43'N, 84°18'W, *Stevens* 5081 (MO); Cerro Saslaya-San José del Hormiguero, between Caño Socio and Loma Molléjones, ca. 300-450 m, 13°45'N, 84°58'-59'W, *Stevens* 7011 (MO); Cayo Palmeta, 0-10 m, 11°34'N, 83°39'W, *Stevens* 20765 (MO); San Juan del Norte, *Smith* 22 (MO); *Seymour* 5906 (MO); Cerro Baká, ca. 6.5 km E of Río Coperna, 200-320 m, 13°40'N, 84°30'W, *Pipoly* 4846 (MO); Estación Experimental El Recreo, W de Cerro La Ceiba, 12°10'N, 84°18'W, *Sandino* 1656 (MO); Monkey Point, Caño El Pato, ca. 10 m, 11°35'N, 83°42'W, *Moreno* 12399 (MEXU, MO); 1-5 m, 11°36'N, 83°40'W, 12318 (MO); 1-5 m, 11°36'N, 83°38'W, *Moreno* & *Sandino* 12007 (MO); 0-20 m, 11°35'N, 83°39'W, *Stevens* 20031 (MO); 1 km S of Monkey Point, ca. 5 m, 83°39'W, 11°35'N, *Moreno* & *Sandino* 12541 (MO); Río Likas, near Silima Lila, 50 m, ca. 14°30'N, 83°50'W, *Pipoly* 4097 (MO); Río Prinzapolka, 0.3-1.9 km N of Limbaika, 8-10 m, ca. 13°29'N, 84°13'W, *Stevens* 8264 (MO); ca. 2 km S of Waní, ca. 0-100 m, ca. 13°42'N, 84°50'W, *Pipoly* 4729 (MO); Río Punta Gorda, Atlanta, 10 m, 11°33'N, 84°02'W, *Moreno* & *Sandino* 12808 (MO); Atlanta, 2 km S of Carolina del Sur, ca. 60 m, 11°32'N, 84°01'W, *Moreno* & *Sandino* 128904 (MO); Río Rama, Caño Zamora, ca. 10 m, 11°57'N, 84°16'W, *Stevens* 8549 (MO); Río Socio, E of Bonanza, ca. 140 m, 14°01'N, 84°34'W, *Stevens* 12345 (MO); Río Waspak, 75-100 m, 14°15'N, 84°36'W, *Stevens*

13091 (MO). PANAMA. Ahorca Lagarto, *Cowell 262* (NY). **Bocas del Toro:** Gualaca-Chiriquí Grande, 4.2 mi. S of Chiriquí Grande, 8°55'N, 82°09'W, *Croat 66816* (AAU, CM, F, MO); Changuinola-Almirante Railroad, Milla 7.5, 100 m, *Croat & Porter 16433* (MO); *Croat 38122* (MO); Chiriquí Lagoon, Water Valley, non *Wedel 989* (GH, MO); Isla Colón, Swan Key, 2 km N, *Tyson & Loflin 6307* (FSU, MO); Río Cricumola, Finca St. Louis-Konkintoe, *Woodson et al. 1901* (GH, MO, NY); Río Changuinola, ca. 1 km above mouth of Río Teribe, <100 m, 9°21'40"N, 82°31'40"W, *Croat & Zhu 76446* (CM, MO). **Canal Area:** *Hayes 805* (NY); Gorgona-Gatún, 10–15 m, *Pittier 2300* (US); Gatún, *Standley 27023* (US); Ft. Sherman, *Standley 30992* (US); Frijoles, *Standley 27467* (US); Ft. Randolph, *Standley 28626* (US); 66–70 m, *Dodge et al. 16870* (MO, U, UC); near Gatún, *Standley 27203* (US); vic. Summit Garden, *Tyson & Lazo 6121* (FSU, RSA); Barro Colorado Island, Gatún Lake, *Standley 31328* (US); 40818 (US); *Kenover 178a* (US); *Elmore X14* (MICH, RSA, US); *Croat 7178* (MO); 79°49'46"W, 9°09'56"N, 6124 (MO, NY, SCZ); 6060 (MO); 4655 (MO, SCZ); *Aviles 44* (MO); *Shattuck 266* (GH, MO); Navy Pipeline Road, *Smith & Smith 3313* (US); 0.25 km NW of Río Macho bridge, 10 km NW of Gamboa, 125 m, *Nees 7863* (MO, RSA, TEX). **Chiriquí:** Gualaca-Chiriquí Grande, 13.6 mi. N of Continental Divide, 120 m, 8°57'N, 80°56'W, *Croat 74934* (MO). **Coclé:** 27 km N of Penonomé, on road to Coeslito, at Continental Divide, 1500 ft., *Hammel 1648* (SEL, MO); Llano Grande-Coeslito, 12 mi. from Llano Grande, 200 m, 8°47'N, 80°28'W, *Churchill et al. 3987* (RSA). **Colón:** ca. 8 km E of Pita, 50–100 m, 9°17'N, 80°W, *Thompson 4815* (CM, MO); Río Indio-Miguel de la Borda, 0 m, *Croat 36927* (MO); near Nuevo Tomosf, 2 mi. from Portobelo, on road to Nombre de Dios, <100 m, *Croat 33527* (MO); Santa Rita Ridge, Transisthmian Highway, ca. 10 mi. from hwy., *Porter et al. 4738* (MO); ca. 7 mi. from Transisthmian Highway, ca. 650 ft., *Wilbur et al. 15083* (MO); Santa Rita Trail, *Cowell 106* (NY). **Panamá:** ca. 20 km from Inter-American Highway, N of Cerro Azul, *Mori et al. 3786* (MO); Río Terabe Valley, El Llano-Cartí, 8 km from Pan-American Highway, 300–400 m, 9°16'N, 79°W, *Knapp & Schmalzer 5476* (MO). **San Blas:** Isla Playón Chico, 0–50 m, 9°20'N, 78°13'W, *Herrera 369* (MO); Nusagardi, El Llano-Cartí Road, 9 mi. N of main highway, 350 m, 9°20'N, 79°W, *Croat & Zhu 76566* (MO, SEL). **Veraguas:** Santa Fe-Río Calovehora, 0.6 mi. beyond agricultural school near Santa Fe, 735 m, *Croat 32962* (MO).

### *Philodendron radiatum* var. *pseudoradiatum*

(Matuda) Croat, stat. et comb. nov. Basionym: *Philodendron pseudoradiatum* Matuda, *Revista Soc. Mex. Hist. Nat.* 1: 96, fig. 8. 1950. TYPE: Mexico, Chiapas: Mun. Escuintla: Turquía, Salto de Agua, 8 Sep. 1947, *Matuda 17787* (holotype, MEXU; isotypes, DS, NY, UC). Figures 8, 327–330.

Internodes 2–12 cm long, 1–2.5 cm diam.; cataphylls unribbed; petioles 28–78 cm long; blades shallowly lobed, the divisions near the middle of the blade extending less than ½ the way to midrib, 11–60 cm long, 11–42 cm wide; posterior lobes (8)12.5–25 cm long, (9)16.2–18 cm wide; basal

veins 5(7–8) per side, with 0–1 free to base, posterior rib absent; primary lateral veins 3–5 per side. INFLORESCENCES 1 per axil; spathe 12–18 cm long; spadix 12.8–17 cm long; pistillate portion 4–4.5 cm long; pistils 1.6–2.2 mm long; ovary with sub-basal placentation; style similar to style type B.

The flowering phenology of *Philodendron radiatum* var. *pseudoradiatum* is still poorly known. Flowering collections are known only from January and September, but the flowering pattern is probably just poorly documented rather than bimodal.

*Philodendron radiatum* var. *pseudoradiatum* is endemic to Mexico, where it is known only from southwestern Chiapas, at 200 to 350 m elevation in "Selva Alta Perennifolia."

*Philodendron radiatum* var. *pseudoradiatum* is distinguished by its incised-lobate blades with the lobes divided less than halfway to the midrib. The typical variety generally has the blades incised more than three-fourths of the way to the midrib.

This taxon might be confused with *P. dressleri*, which also has leaf blades divided only about halfway to the midrib. That species differs in having much thicker stems (typically to 6 cm diam.), sharply 2-ribbed cataphylls, and by occurring in much drier areas where it loses its leaves during the dry periods. In contrast, *P. radiatum* var. *pseudoradiatum* has rather slender stems, unribbed cataphylls and occurs in mesic areas.

Until recently, *Philodendron radiatum* var. *pseudoradiatum* was represented by only two collections, *Matuda 17787* and *18718*, from the same locality. Several collections are noteworthy. *Matuda 18045* consists of two sheets, one of which has a leaf lobed in much the same manner as the type, while the other has a leaf lobed to more than two-thirds of the way to the midrib. This may be an indication that the type of *P. radiatum* var. *pseudoradiatum* is merely an aberration in the degree of lobing for *P. radiatum*.

Another noteworthy collection is *Croat & Hannon 63381* from Sierra de Soconusco, northwest of Mapastepec. It is perhaps also *P. pseudoradiatum* but has smaller and narrower leaves. It looks much like a hybrid between *P. radiatum* var. *pseudoradiatum* and some other species, but if so, it is not clear which other species might be involved in the putative hybrid plant.

*Additional specimens examined.* MEXICO. Chiapas: Escuintla, Salto de Agua, 350 m, *Matuda 18718* (MEXU, UC); Mpio. Mapastepec, Sierra de Soconusco, new unfinished road to Tuxtla Gutiérrez, 200 m, 15°31'N, 92°50'W, *Croat & Hannon 63381* (CAS, CM, F, MEXU, MO, TEX).

***Philodendron roseospathum* Croat, sp. nov.**

TYPE: Panama. Panamá: Cerro Jefe, along road just below the summit, 9°15'N, 79°30'W, 17 June 1976, *Croat & Zhu 76211* (holotype, MO-4619415; isotypes, AAU, B, CAS, CM, COL, CR, DUKE, F, GH, HUA, K, M, MEXU, NY, P, PMA, RSA, S, SEL, TEX, US, VEN). Figures 341-345.

Planta terrestris, raro hemiepiphytica; internodia 1-4 cm longa, 1.5-2.7 cm diam.; cataphylla 12-27 cm longa, acute 2-costata, persistentis semi-intacta nodis superioribus; petioli U-formati, obtuse sulcati, 10-23 cm longi, 3-6 mm diam., crassior quam latus; lamina (22)30-49 cm longa, 9.8-21 cm lata, elliptica vel oblongo-elliptica, interdum oblanceolata, interdum attenuata apice, acute vel anguste rotundata, interdum attenuata basi; inflorescentia 1-2; pedunculus 2.5-6 cm longus; spathe 6.5-10 cm longa; lamina spathe extas saepe rubella plerumque atriviridi, intus pallide viridi vel marronina; tubo spathe extas pallido-lineato, intus pallide viridi vel marronino; pistilla (4)5-7(8)-locularia; loculi 14-18-ovulati; baccæ albae.

Terrestrial or rarely hemiepiphytic; stem appressed-climbing or creeping, leaf scars inconspicuous, obscured by cataphyll fibers; internodes short, semiglossy, 1-4 cm long, 1.5-2.7 cm diam., longer than broad, dark green to gray-green to reddish, obscured by cataphyll fibers; roots many, thin, wiry, descending, greenish brown; cataphylls 12-27 cm long, sharply 2-ribbed (unribbed to weakly to sharply 2-ribbed in Colombia), pale red to purple-violet, drying dark brown, sharply D-shaped, persisting semi-intact at upper nodes, as fibers below; petioles 10-27 cm long, 3-6 mm diam., U-shaped, thicker than broad, firm, dark green to reddish, tinged purple, somewhat flattened to obtusely or narrowly sulcate adaxially; geniculum pale green to reddish violet, 1 cm long, 1.5 cm diam.; blades elliptic to oblong-elliptic, sometimes oblanceolate, rarely elliptic, coriaceous to subcoriaceous, conspicuously bicolorous, sometimes acuminate at apex (the acumen inrolled, 1-3 mm long), acute to narrowly rounded, sometimes attenuate at base, (22)30-49 cm long, 9.5-21 cm wide (1.8-3.5(4.5) times longer than broad), (1.6-3.2 times longer than petiole), about twice as long as wide, upper surface dark green, matte to semiglossy, lower surface moderately paler, semiglossy to glossy; midrib narrowly sunken, paler than surface above, narrowly convex, thicker than broad, matte, paler than surface and sometimes reddish green below; basal veins lacking or sometimes 1-2 per side, free to base; primary lateral veins 8-14(16) per side, departing midrib at a 40-55° angle, sunken to narrowly sunken, concolorous to

paler than surface above, convex and darker than surface below; interprimary veins few, sunken to narrowly sunken, concolorous to paler than surface above, raised and darker than surface below; minor veins few, moderately obscure above, darker than surface below, arising from midrib only, sometimes prominent and weakly undulate on drying. INFLORESCENCES 1-2 per axil; peduncle 2.5-6 cm long, 2.5-5 mm diam., reddish, prominently green streaked; spathe 6.5-10 cm long (1.3-3.3(4.6) times longer than peduncle), red to maroon or violet-purple or sometimes green heavily tinged red, sometimes greenish white throughout; spathe blade frequently dark green (red in Colombia) outside, pale green to maroon inside; spathe tube short pale-lineate outside, 3-4 cm long, 2 cm diam., pale green to maroon inside; spadix ± tapered, 6.8-8 cm long, broadest at the base; pistillate portion white to pale green, cylindrical, 2-3.7 mm long, 1-1.5 mm diam. at apex, 11-7.2 mm diam. at middle, 9-11(17) mm wide at base; staminate portion 3.9-4.9(6) cm long; fertile staminate portion cream, cylindrical to somewhat tapered, 10 mm diam. at base, 8 mm diam. at middle, 8 mm diam. ca. 1 cm from apex, broadest usually at base, narrower than the pistillate or sterile portion; sterile staminate portion as broad as the pistillate portion, 10 mm diam.; pistils 2.5 mm long, 1.6-1.9 mm diam., tinged reddish; ovary (4)5-7(8)-locular, 1.4 mm long, 1.6 mm diam., with axile placentation; locules 1.4 mm long, 0.5 mm diam.; ovules 14-18 per locule, contained within a gelatinous matrix, 2-seriate, 0.3-0.5 mm long, longer than funicle; funicle 0.2-0.4 mm long, adnate to lower part of partition, style 0.4-0.6 mm long, 1.6-1.8 mm diam., similar to style type B; style apex flat to weakly rounded; stigma usually subdiscoid, 1.1-1.3 mm diam., 0.2-0.5 mm high, covering entire style apex; the androecium truncate, prismatic, oblong, margins irregularly 4-6-sided, 0.7 mm long, 1.2-1.5 mm diam. at apex; thecae oblong to cylindrical, 0.5 mm wide, ± parallel to one another and contiguous; sterile staminate flowers clavate, blunt, irregularly 4-6-sided, 2.3 mm long, 2 mm wide. INFRUCTESCENCE with pistillate spadix 4-4.5 cm long, 1.5-2.5 cm diam.; berries white (immature), 3.5 mm long, 2.6 mm diam., 12-14 per locule; seeds 7-9 per locule, whitish, oblong, 1.8 mm long, sharply tapered on one end.

*Philodendron roseospathum* ranges from Panama to Colombia. In Panama, this taxon is restricted to both sides of the Panama Canal, ranging from Ver-

aguas to San Blas, mostly in *Premontane rain forest* or less frequently in *Tropical wet forest* life zones, at 350 to 1000 m elevation. In Colombia, it has been collected only in tropical pluvial forest at 100 to 900 m.

*Philodendron roseospathum* is a member of *P.* sect. *Philodendron* subsect. *Canniphyllum*. The typical variety is recognized by its generally erect, frequently terrestrial habit, short internodes, and rosulate cluster of leaves with moderately short, bluntly sulcate, thicker-than-broad petioles. The dark green, elliptic-oblancoate leaf blades are narrowly rounded to acute at the base. It is especially recognizable by its masses of weathered persistent cataphylls and clusters of inflorescences with reddish spathes (hence the name "roseospathum"). This species is unusual for the genus in that it is generally terrestrial. It has prospered in cultivation and is deemed an important horticultural introduction.

*Philodendron roseospathum* appears to be most closely related to *P. cuneatum* Engl. from the Pacific slope of Colombia, but the latter species differs in having acutely sulcate, C-shaped petioles and smaller, more narrowly pedunculate inflorescences with white spathe blades. It is also related to several other undescribed species from the Amazon basin.

*Philodendron roseospathum* is here divided into two varieties, with variety *angustilaminatum* Croat differing from the typical variety in having somewhat longer petioles and proportionately much narrower blades (4.5 times longer than broad vs. an average of 2.5 times longer than broad for the typical variety). See *P. roseospathum* var. *angustilaminatum* for additional differences.

#### KEY TO THE VARIETIES OF *P. ROSEOSPATUM*

- 1a. Blades oblong, ca. 4.5 times longer than wide; Panama (Colón), 915 m elevation ..... var. *angustilaminatum*  
 1b. Blades  $\pm$  elliptic, ca. 2.5 times longer than wide; Panama to Colombia, 400–900 m elevation ..... var. *roseospathum*

#### *Philodendron roseospathum* Croat var. *roseospathum*

Flowering in *Philodendron roseospathum* var. *roseospathum* occurs in the late dry season and early rainy season (March through July) with post-anthesis collections known from March through September and immature fruits from July to December. The few South American collections concur with this pattern. Flowering collections are known from Colombia in July.

In Central America, *P. roseospathum* var. *roseospathum* is common locally in cloud forests along the Continental Divide as far west as Veraguas and as far east as the El Llano-Cartí Road. It is certainly to be expected in Darién Province.

For differences with *P.* var. *angustilaminatum*, see the preceding key.

*Additional specimens examined for P. roseospathum var. roseospathum.* PANAMA: Coelá: Continental Divide ridge, Cclesito Road, *Hammel 2540* (CR, MO); El Copé region, 7 km N of Copé, Alto Calvario, 800 m, *Folsom & Maushet 7948* (MO); 750–900 m, *Croat 44738* (F, MO); 710–800 m, 8°39'N, 80°36'W, 68763 (MO); 900–1000 m, 8°39'N, 80°36'W, *Croat 75057* (MO); El Valle region, La Mesa, N of El Valle de Antón, 800–900 m, 8°38'N, 80°09'W, *Croat 67130* (CAS, CM, K, MEXU, MO, NY, PMA, TEX); 860–900 m, 37346 (F, MO); 830–900 m, 8°36'N, 80°07'W, 74810 (CM, LE, MO, WIS); ca. 800 m, 25435 (F, MO); *Luteyn & Kennedy 1652* (MO); 850 m, 8°37'N, 80°06'W, *de Nevers et al. 3513* (MO); 2700 ft., *Sytama et al. 4350* (CM, MO); 840–880 m, 8°38'N, 80°7.5'W, *Grayum & Evans 9908* (MO); Cerro Gaital, 900–1000 m, 8°40'N, 80°07'W, *Knapp 5311* (MO); 860 m, 8°37'N, 80°08'W, *Croat & Zhu 76734* (AAU, CAS, CM, MEXU, MO); N slope and summit of Cerro Pilón, 900–1173 m, *Croat 22951* (MO). PANAMÁ: El Llano-Cartí Road, 4 mi. beyond the highway, 500 m, *Croat 49135* (MO); Cerro Jefe region, 750–850 m, 9°15'N, 79°30'W, *McPherson 11166* (MO), *11197* (CM, MO); 1000 m, *Croat 49088* (MO); ca. 1000 m, *Mori et al. 3795* (MO); ca. 950 m, ca. 9°15'N, 79°30'W, *McPherson 7107* (B, K, MO, US); 2500–3000 ft., *Hammel 3704* (MO); 750–800 m, 9°14'N, 79°22'W, *Croat 67089* (MO); vic. Altos de Pacora, 800–1000 m, *Croat 22672* (CM, L, MO, WIS); 3–3.5 mi. NE of Altos de Pacora, 700–750 m, 9°15'N, 79°25'W, *Croat 68635* (MO); La Envidia, ca. 800 m, *Kennedy et al. 3374* (US); *Plouman & Weil 3158* (ECON); 1000 m, *Folsom & Page 5938* (MO); 6 mi. above Lago Cerro Azul, *Croat 15225* (MO); Campo Tres, ca. 700 m, *Croat 27069* (F, MO); 4.6 km beyond peak, on road to Altos de Pacora, 26.3 km from the Inter-American Highway, ca. 600 m, *Croat 35931* (CAS, CM, F, GH, M, MEXU, MO, P). SAN BLAS: El Llano-Cartí Road, Nusagandi, 300–350 m, 9°15'N, 79°W, *Croat 69282* (MO); Mile 10.9, 450 m, 9°18'N, 79°59'W, *Croat 751234* (MO); Cerro Obu, 400–500 m, 400–500 m, *de Nevers et al. 8054* (MO). VERAGUAS: Santa Fe region, Santa Fe-Río San Luis, past Escuela Agrícola Alto de Piedra, ca. 5 mi. N of school, 670 m, 8°33'N, 81°08'W, *Croat 66972* (GB, MO, RSA); Santa Fe-Calovebora, 1.7 mi. past Alto Piedra School, 570 m, 8°33'N, 81°08'W, *Croat & Zhu 76857B* (MO).

COLOMBIA. Antioquia: Parque Nacional Natural "Las Orquídeas," Sector Venados, 900 m, 6°33'N, 76°19'W, *Cogollo et al. 3344* (MO). Chocó: Quibdó-Istmina, vicinity of Quibdó, <100 m, 6°28'N, 76°36'W, *Croat & Cogollo 52233* (MO); Quibdó-Bolivar, 117–118 km E of Quibdó, 465 m, 5°44'N, 76°28'W, *Croat 57515* (CHOCO, CM, COL, JAUM, MO, NY, PMA); Río Baudó, *Fuchs et al. 22048* (COL); Medellín-Quibdó at Km 208.5, 9 km W of Tutusendo, ca. 9 km E of Quibdó, <100 m, 5°39'N, 76°40'W, *Croat 56202*.

***Philodendron roseospathum*** Croat var. ***angustilaminatum*** Croat, var. nov. TYPE: Panama. Colón: Cerro Bruja as approached from Río Escandaloso, 915 m, 18 May 1978, Hammel 3133 (holotype, MO-2639732). Figure 349.

*Intermedia brevia*; cataphylla ignota; petiolus 24–27 cm long; lamina oblonga, 42–46 cm longa, 9.5–10 cm lata, circa duplo longior quam petiolus; pedunculus 3.8 cm long; spatha 7 cm longa, omnino viridialba; pistilla (4)5-locularia; loculi 6–8-ovulati.

Internodes short; cataphylls not seen; petioles 24–27 cm long; blades oblong, acuminate at apex, rounded at base, 42–46 cm long, 9.5–10 cm wide (4.2–4.8 times longer than wide), (1.75 times longer than petiole); basal veins 2 per side, free to base; primary lateral veins 10 per side. INFLORESCENCES with peduncle 3.8 cm long; spathe 7 cm long greenish white throughout; spathe tube 3 cm long; spadix 6.3 cm long; pistillate portion 2 cm long in front; staminate portion 4.3 cm long; pistils 1.7 mm long; ovary 4(5)-locular; ovules 6–8 per locule, 1-seriate, ca. 0.3 mm long; style similar to style type D. Berries unknown.

Flowering in *Philodendron roseospathum* var. *angustilaminatum* probably occurs in the wet season, with the only flowering collection from May.

*Philodendron roseospathum* var. *angustilaminatum* is known only from the type locality near the Canal Area in Colón Province, at 915 m elevation in Premontane rain forest.

This variety is distinguished by its long-petiole, oblong leaf blades with rounded bases and greenish white spathes. It is distinguished from the typical variety by its proportionately much narrower blades (4.5 times longer than wide in *P. roseospathum* var. *angustilaminatum* vs. about 2.5 times longer than wide in var. *roseospathum*) and proportionately longer petioles (only 1.7 times longer than petiole vs. 1.6–3.2 times longer in var. *roseospathum*). In addition, *P. roseospathum* var. *angustilaminatum* has only 6–8 ovules per locule (vs. 16–18 for *P. roseospathum* var. *roseospathum*).

It is unusual that *P. roseospathum* var. *angustilaminatum* has style type D while the autonymic variety has style type B. Perhaps it is an artifact of the poorly preserved nature of the material.

One collection, Croat 57592, from near Quere-mal in Valle Department, at 1300 m in an area of either Premontane moist forest or Tropical Lower Montane wet forest, may also represent this variety but has unribbed cataphylls. Croat 57036, a sterile collection from Cotopaxi Province, Ecuador, also has unribbed cataphylls and may represent the same taxon.

***Philodendron rothschuhianum*** (Engl.) Croat & Grayum, Ann. Missouri Bot. Gard. 74: 659. 1987. *Syngonium rothschuhianum* Engl., in Engl. & K. Krause, Pflanzenr. IV. 23E (Heft 71): 124. 1920. TYPE: Nicaragua. Matagalpa: Matagalpa, 1000 m, 12°55'N, 85°55'W, Roths-chuh 229 (holotype, B). Figures 4, 9, 31, 336–340, 347, 348.

Hemiepiphytic to epiphytic or rarely terrestrial; stem appressed-climbing, creeping, sap watery, leaf scars conspicuous, 2 cm long, 2.5 cm wide; internodes short, semiglossy, 1.8–3 cm long, 1.5–3 cm diam., about as long as broad, gray-green, epidermis fissured; roots scurfy, few per node; cataphylls fleshy, soft, 16–23 cm long, unribbed then sharply 2-ribbed at apex, or bluntly 2-ribbed throughout or weakly 1-ribbed throughout, green, short dark striate, semiglossy, deciduous, emarginate with subapical apiculum at union of ribs at apex. LEAVES erect-spreading to spreading; petioles 33–71 cm long, 3–13(20) mm diam., subterete, moderately spongy, medium green, obtusely 1-ribbed near apex adaxially, surface weakly glossy, densely green striate; sheath inconspicuous; blades deeply 3-lobed, subcoriaceous, weakly bicolorous, gradually acuminate, long to very long acuminate at apex (the acumen inrolled, 1–2.5 mm long), ± hastate at base, 22–42 cm long, 25–57 cm wide (0.6–1 times longer than wide), (0.5–1 times the petiole length), about two-thirds as long as petiole, upper surface semiglossy, lower surface weakly glossy, slightly paler; sinus hippocrepiform; median lobes 16–39.5 cm long, 7–20 cm wide; lateral lobes narrowly ovate, 14–28 cm long, 5–17 cm wide, directed toward apex, broadly confluent 2–5 cm with medial lobe, the margins undulate; midrib sunken to narrowly concave, paler than surface above, almost round-raised, paler than surface below; basal veins 10–15 per side, the remainder arranged in a ± regular manner in the posterior lobe, coalesced in part throughout much of their length into a central rib; posterior rib usually not naked, sometimes naked for 1 cm, rarely to 2.5 cm; primary lateral veins 5–12 per side, departing midrib at a 50–65° angle, narrowly sunken, concolorous above, convex below; interprimary veins sunken and concolorous above, raised, almost as conspicuous as primary lateral veins below; minor veins darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, 2–3(9) per axil; peduncle 5.2–21 cm long, 5–10 mm diam., medium green, densely lineate; spathe (6.5)8–14(16) cm long, (0.6–2.2 times longer than peduncle), markedly constricted above the tube; spathe

blade pale green to greenish white to yellowish green outside, to ca. 5 cm wide when open, ca. 2.4 cm diam. at constriction, greenish white to creamy white inside; spathe tube ellipsoid, 4–7 cm long, pale green to yellowish green outside, greenish to greenish white to creamy white inside; **spadix** very short stipitate; somewhat protruding forward at anthesis, 10.5 cm long, constricted at base of fertile staminate portion; pistillate portion pale lime-green, 3.7 cm long in front, 2.5 cm long in back, 1–1.3 cm diam. at apex, 1.2–1.4 cm diam. at middle, 1.2 cm wide at base; staminate portion 6.4–9.3 cm long; fertile staminate portion creamy white, clavate, 1 cm diam. at base, 1.3 cm diam. at middle, 1 cm diam. ca. 1 cm from apex, usually broader than the pistillate portion; sterile staminate portion broader than fertile or pistillate portion, creamy white, 1.6–1.9 cm diam. at base, 1.3–1.4 cm diam. at apex; pistils 1.6–2.6 cm long, 1.2 mm diam.; ovary 5–7-locular, 0.8 mm long, 1.1 mm diam., with sub-basal placentation; locules 0.8 mm long, 0.3–0.4 mm diam.; ovule sac 0.6 mm long; ovules 1 per locule, contained within translucent, gelatinous ovule sac, 0.3–0.5 mm long, longer than funicle; funicle 0.1–0.3 mm long (can be pulled free to base), style 0.6 mm long, 1.2 mm diam., similar to style type B; style apex sloping to rounded; stigma brush-like, hemispheroid, 0.7–1 mm diam., 0.3–0.6 mm high, covering entire style apex; apex drying button-like, with or without radial ridges from center; the androecium truncate, oblong,  $\pm$  prismatic, margins irregularly 4–6-sided, 1–1.6 mm long, 1.5–1.9 mm diam. at apex; thecae oblong, 0.4 mm wide,  $\pm$  parallel to one another, nearly contiguous; sterile staminate flowers blunt, sometimes clavate or prismatic or irregularly 5–6-sided or 3–6-sided, 1.7–2.3 mm long, 1.6 mm wide. Berries pale green or pale yellow-green to greenish white, 4 mm long, 2.5 mm diam.; seeds 1–3, medium green, (1.2/2.1–2.3 mm long, 0.6–0.8 mm diam., within thin envelope. JUVENILE plants with petioles sheathed to about midway; blades with posterior lobes hastate, acuminate at apex, promptly auriculate on posterior margin, broadly confluent with anterior margin on anterior margin.

Flowering in *Philodendron rothschuhianum* occurs during the dry season and early rainy season. Most collections have been made in March, but flowering occurs as late as August. Post-anthesis collections have been made from January through November (except October), but are particularly abundant from March through May. Immature fruiting collections have been made from April through September (also December), but mature fruits are known from only August.

*Philodendron rothschuhianum* ranges from Honduras (Gracias a Dios) and from Nicaragua (Río San Juan, Zelaya, Jinotega, and Matagalpa) to Panama (Bocas del Toro, Veraguas, and Coclé), principally on the Atlantic slope, from sea level to 1450 (mostly less than 1000) m elevation. It occurs in *Premontane wet forest*, *Tropical wet forest*, and less frequently in wetter parts of *Tropical wet forest* and *Premontane rain forest* life zones.

*Philodendron rothschuhianum* is a member of *P. sect. Tritomophyllum*. This species is distinguished by its short internodes; unribbed, deciduous cataphylls; subterete, moderately spongy petioles; and especially by its deeply three-lobed blades about two-thirds as long as the petioles with the lateral lobes broadly confluent with the medial lobe and the sinus hippocrepiform and naked along the posterior rib for a short distance from the petiole.

*Philodendron rothschuhianum* is most frequently confused with *P. tripartitum*, which differs in having less auriculate lateral lobes that are scarcely or not at all naked along the posterior rib. In addition, *P. tripartitum* has a much more slender, scarcely constricted, spathe tube reddish within (vs. a markedly inflated, uniformly greenish tube in *P. rothschuhianum*).

In terms of leaf shape, *P. rothschuhianum* is closest to *P. cotobrusense*. The latter differs in medial lobes with 18–19 pairs of primary lateral veins (vs. 5–12 pairs for *P. rothschuhianum*).

The style apex is apparently variable in this species, with *Croat 66772* and *Neill 1569* having type B styles while *Croat 35657* has style type D. The latter has the style apex prolonged into a distinct boss separated from the rest of the style apex by a distinct neck.

*Additional specimens examined.* COSTA RICA. **Alajuela:** Quebrada Guillermina, on N side of Volcán Arenal, 500 m, 10°29'N, 84°42'W, *Lent et al. 3415* (F, U); Naranjo-Aguas Zarcas, 8 km NE of Quesada, 600 m, *Croat 46942* (MO); Finca Los Ensayos, ca. 11 mi. NW of Zarcero, ca. 850 m, *Croat 43567* (MO); 43637 (MO); Monteverde Biological Reserve, Río Peñas Blancas, 800 m, *Bello 1980* (CR, INB); 850–900 m, 10°20'N, 84°43'W, *Haber & Bello 6836* (MO); 850 m, 10°19'N, 84°43'W, *Haber & Cruz 8409* (MO); Finca de Tomás Guindón, 900 m, 10°19'N, 84°43'W, *Bello 763* (CR, MO); Río Sarapiquí, road to Colonia Virgen del Socorro, ca. 740 m, 10°5.5'N, 84°10.5'W, *Grayum & Hammel 5516* (B, CR, MO); ca. 800 m, *Burger & Antonio 11101* (CR, F); 11097 (F); 830 m, *Croat 68301* (MO); Vara Blanca-Puerto Viejo, 3 mi. N of San Miguel, 380 m, *Croat 35657* (CR, MO); *Cattus-Upala*, 3 km NNE of Bijagua, 450 m, 10°45'N, 85°3'W, *Burger & Baker 9881* (CR, F); 8 km N of Bijagua, 300 m, *Croat 36502* (CR, MO); 13.8 km N of Bijagua, 100–150 m, *Croat 36438* (MO); 36402 (CR, MO); near Río Zapote, 1.8–2.7 km S of Río Canalete, 100 m, *Croat 36402* (MO); 13 km W of Fortuna, Río Tabacón, 500–550 m, 10°29'N, 84°43'W, *Lesner et al. 15250* (MO); 15 km

NW of Arenal by air, 700 m, 10°34.5'N, 84°54'W, *Liesner et al.* 15088 (MO); 15065 (MO); San Ramón, *Bittner & Venschott* 1941 (CR); *Nilson* 460 (CR). **Cartago:** 4 km SE of Pejibaye, 700 m, 9°48'N, 83°42'W, *Liesner et al.* 14330 (MO); Turrialba, León 393 (US). **Guacacaste:** Parque Nacional Guanacaste Estación Pitilla, 600 m, 11°02'N, 85°25'18"W, *Hammel et al.* 17372 (CR, MEXU, MO, NY, U, USJ). **Heredia:** near Tirimbina, E of Río Sarapiquí, 150–250 m, 10°24'N, 84°7'W, *Burger & Burger* 8026 (F); Tirimbina, 700 ft., *Proctor* 32148 (LL); La Selva Field Station, 100 m, *Jacobs* 2826 (DUKE); *Folsom* 9874 (DUKE); 50–80 m, 10°26'N, 84°01'W, *Grayum & Chaurria* 8300 (MO); Puerto Viejo de Sarapiquí, *Grayum* 2120 (DUKE); ca. 100 m, *Grayum* 2856 (MO); *Croat* 57199 (MO, voucher of live plant of *Grayum* 2856); Río Puerto Viejo, ca. 2 km upstream from confluence with Río Sarapiquí, 100 m, 10°26'N, 84°10'W, *Burger & Stolze* 5756 (CR, US); Volcán Barva, Río Peje–Río Sardinalito, 700–950 m, 10°17.5'N, 84°04.5'W, *Grayum* 6981 (CR, MO); 700–800 m, 10°17.5'N, 84°04.5'W, 6720 (CR, MO). **Limon:** ca. 1 mi. NE of Bribrí, ca. 40 mi. SW of Limón, above Río Catarata, 50–100 m, *Croat* 43245 (MO); 7 km SW of Bribrí, 100–250 m, *Gómez et al.* 20320 (MO); Río Pacuare, 50–100 m, 10°05'N, 83°29'W, *Burger & Liesner* 6895 (F, MO); vic. of Guápiles, 300–500 m, *Standley* 37469 (US); 2 km W of Río Toro Amarillo, W of Guápiles, 275 m, 10°13'N, 83°50'W, *Thompson & Rawlinson* 1216 (CM); Hacienda Tapezco–Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, *Davidson et al.* 7138 (RSA); Bahía de Portete Parque Nacional, 0–10 m, 10°N, 83°05'W, *Thompson & Rawlinson* 1176 (CM); Cerro Corónel, E of Río Zapote, along and above new road, within 1 km of Río Colorado, 10–40 m, 10°40'N, 83°40'W, *Stevens* 24277 (CR, MO); E of Río Zapote, 10–100 m, 10°40'N, 84°40'W, *Stevens* 23974 (CR, MO); E of Laguna Danto, 20–170 m, 10°41'N, 83°38'W, *Stevens* 24453 (CR, MO); Parque Nacional Tortuguero, Puesto Cuatro Esquinas, 4 m, 10°32'N, 83°30'W, *Robles* 1837 (CR, MO); Barra del Colorado, 0–2 m, 10°47'N, 83°35'W, *Stevens* 24073 (CR, MO); Río Colorado, Barra del Colorado, 1–5 m, 10°47'40"N, 83°35'30"W, *Davidse & Herrera* 30869 (MO); 2 airline km SSE of Islas Buena Vista, 10–120 m, 10°40'N, 83°40'W, *Davidse & Herrera* 31035 (MO); Río Reventazón, Finca Montecristo, below Cairo, ca. 25 m, *Standley & Valerio* 49027 (US); Parque La Amistad, Fila Tsurábeta, between Río Urén and Río Larí, 800 m, 9°27'30"N, 83°W, *A. Chacón* 294 (CR, MO). **Puntarenas:** hills above Santa Elena, 3 km N of Monteverde, 1450 m, 10°20'N, 84°50'W, *Haber & Bello* 5067 (MO). **San José:** S of San José, *Greenman & Greenman* 5353 (MO); Braulio Carrillo National Park, Fila Carrillo, 700 m, *Gómez et al.* 21149 (CR, MO); 600–700 m, *Croat* 78777 (CR, INB). **HONDURAS. Gracias a Dios:** Ahuas Bils, 200 km SW of Puerto Lempira, 100 m, *Nelson & Cruz* 9146 (CM, MO, NY, UNAH). **NICARAGUA. Chontales:** Cerro Buenavista, 5 km N of Cuapa, *Neill* 637 (MO); 4 km NW of Santo Domingo, 280 m, ca. 12°17'N, 85°06'W, *Grijalva et al.* 3771 (MO). **Jimotega:** Finca San Luis, 21 km from Valle del Cuá, NW de El Cedro, 700 m, 13°30'N, 85°38'W, *Moreno* 959 (MO); Río Bocay, Caño Litutus, ca. 175 m, ca. 13°58'N, 85°21'W, *Stevens et al.* 16617 (MO); Salto Kayaska, 190–340 m, ca. 13°51'N, 85°22'W, *Stevens et al.* 16494 (F, MO). **Matagalpa:** Macizos de Peñas Blancas, SE side, drainage of Quebrada El Quebradón, slopes NW of Hacienda San Sebastián, 800–1100 m, 13°14'–15'N, 85°38'W, *Stevens* 21258 (MO); summit of El Toms Road, *Neill* 1569

(HNMN, MO); 10.5 km NW of Matagalpa, ca. 1000 m, 12°57'N, 85°51'W, *Moreno* 10233 (MO); El Paraíso, 10 km de Matagalpa, 800–820 m, 12°59'N, 85°51'W, *Moreno* 6607 (MO); N of Cerro Musán, near Wanawás, *Araquistain & Moreno* 2741 (LE, MO); NW of Cerro Musán, near Paylo, 500–800 m, *Araquistain & Moreno* 2572 (MO); 800–1200 m, 2510 (MO); 2495 (CAS, HNMN, MO); 300–600 m, 2471A (MO); Río Bilampí, NW of Cerro Musán, 4 km SW of Wanawás, 200–500 m, 13°00'–01'N, 85°14'W, *Araquistain & Moreno* 2614 (HNMN, MO, US); Comerca Wanawás, 180–200 m, 12°02'–03'N, 85°13'W, *Araquistain & Moreno* 2398 (HNMN, K, MO); near Cerro Musán, 8 km from Población Wanawás, Palán Grande, 500–800 m, 12°59'–13°N, 85°14'W, *Araquistain & Moreno* 2355 (HNMN, MO, NY); Río Bilampí Valley, Cerro Musán, Salto Grande de Quebrada Negra, 500–800 m, *Neill* 1800 (MO). **Río San Juan:** Caño Chontaleño, 20 km NE of El Castillo (Río Indio watershed), 200 m, *Neill* 3367 (MO); *Neill & Vincelli* 3541 (BM, MO); Río Indio Valley, 6 km upstream from the junction with Caño La Pimienta, 300–600 m, *Neill* 1557 (MO); Río Sábalo, Buenos Aires, 70 m, 11°02'N, 84°28'W, *Moreno* 25595 (MO). **Zelaya:** ca. 1.5 km NE of Las Esperanza de Las Quebradas, 300–350 m, ca. 13°38'N, 85°02'W, *Stevens & Moreno* 19308 (MO); Limbaika–El Empalme, ca. 3.9 km SE of El Empalme, ca. 65 m, ca. 13°39'N, 84°24'W, *Stevens* 12930 (MO); road to Mina Nueva América, ca. 10 km from main road, *Stevens* 12687 (MO); Rosita–Puerto Cabezas, ca. 15.7 km SW of Río Kukulaya, <100 m, ca. 13°58'N, 84°12'W, *Stevens* 8500 (MO); Cerro El Inocente, near Caño Majagua, ca. 800–1000 m, ca. 13°45'N, 85°W, *Stevens* 6805 (MO); Siuna–Matagalpa, ca. 31.4 km beyond Río Uli, ca. 8.9 km beyond Rosa Grande La Balsama, <200 m, *Stevens* 7456 (MO); Cerro Suslaya–San José del Hormiguero, from Loma Mollejones eastward, ca. 200–400 m, ca. 13°44'–45'N, 84°57'–58'W, *Stevens* 7049 (BM, MO); Caño Majagua, ca. 750–850 m, ca. 13°45'N, 85°00'–01'W, *Stevens* 6936 (MO); Caño Majagua–Caño Sucio, ca. 600–800 m, ca. 13°45'N, 84°59'–85°W, *Stevens* 6821 (MO, PMA); vic. of La Pimienta, ca. 13°45'N, 84°59'W, *Pipoly* 6225 (MO); Cerro La Pimienta–El Hormiguero, ca. 800–1000 m, ca. 13°45'N, 84°59'W, *Pipoly* 6012 (MO); Cerro La Pimienta, number 1, ca. 900–980 m, ca. 13°45'N, 84°59'W, *Pipoly* 5129 (MO); ca. 13 km above Kururia, on road to San Jerónimo, <200 m, *Pipoly* 3794 (MO); Risco de Oro, ca. 40 m, *Pipoly* 5043 (MO); 10 km NE of Siuna, along Caño Madriguera, 250 m, *Neill* 3754 (BM, MO); 4 km NE of Siuna, road to El Dos, 300 m, 13°45'N, 84°45'W, *Moreno & Robledo* 20856 (MO); 1 km W of El Naranjo, 200–210 m, 13°34'N, 85°11'W, *Moreno & Robledo* 20616 (MO, QCA); SE of Cerro El Hormiguero, ca. 900–1000 m, 13°44'10"N, 84°59'50"W, *Grijalva* 473 (MO); Río Iyas, Quebrada El Toro, 260–280 m, *Vincelli* 370 (MO); near San Juan del Norte, C. Smith 5 (F). **PANAMA. Bocas del Toro:** Guslaca–Chiriquí Grande, ca. 10 km SW of Chiriquí Grande, ca. 300 m, 8°52'N, 82°10'W, *Thompson* 4936 (CM); 4.2 mi. S of Chiriquí Grande, ca. 0 m, 8°55'N, 82°09'W, *Croat* 66815 (MO); Fortuna Lake area, Continental Divide, 900 m, 8°48'04"N, 82°15'04"W, *McPherson & Aranda* 10185 (MO, PMA, W); 3.2 mi. N of Divide, 700 m, 8°45'N, 82°15'W, *Croat & Grayum* 60276 (CM, MO, RSA); 850–950 m, 8°40'04"N, 79°50'04"W, *McPherson* 10546 (AAU, M, MEXU, MO, US); ca. 300 m, ca. 8°50'N, 82°15'W, *McPherson* 8522 (K, MO, US); 6.6 mi. N of bridge over Fortuna Lake, 780 m, 8°45'N, 82°18'W,

Croat 66772 (MO); 1.2 mi. N of Divide, 910 m, 8°44'N, 82°17'W, Croat 66462 (MO, PMA); 1.6 mi. N of Divide, 850 m, 8°48'N, 82°13'W, Croat 74931 (MO); 1.6 mi. N of Divide, 770 m, 8°45'N, 82°17'W, Croat & Zhu 76534 (MO); Río Cricamola, Finca St. Louis-Konkintot, ca. 10-50 m, Woodson et al. 1919 (MO). Coelá: El Copé region, 4.5 mi. N of El Copé, 750 m, 8°38'N, 80°35'W, Thompson 4760 (CM, MO); 5-6 mi. N of El Copé, 600-800 m, 8°38'N, 8°35'W, Croat & Zhu 77224 (CM, MO); 680-770 m, 8°39'N, 80°36'W, Croat 74828 (CM, MO); Alto Calvario, Croat 68848 (MO). Veraguas: 15.6 km NW of Santa Fe, 450-550 m, Croat 27704 (MO); 350-400 m, 27385 (MO).

**Philodendron sagittifolium** Liebm., Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn 1849: 17. 1849. TYPE: Mexico. Veracruz: Río Nautla at Pital, Apr. 1841, Liebm. s.n. (holotype, C). Figures 346, 350-356.

*Philodendron daemonium* Liebm., Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn 1849: 17. 1849. TYPE: Mexico. Veracruz: vic. Colipa at Potrero de Consoquita, Liebm. s.n. (holotype, C; isotype, K).

*Philodendron tanyphyllum* Schott, Prod. Syst. Aroid.: 272. 1856. TYPE: Mexico. Liebm. s.n. (W destroyed). Schott's ic. 2557 (neotype, here designated, W).

*Philodendron sanguineum* Regel, Ind. Sem. Hort. Petrop. 82. 1868. TYPE: Mexico. Locality unknown: Karwinsky s.n. (holotype, LE? no longer extant); t. 621 in Regel, Gartenflora 18. 1869 (neotype, here designated).

*Philodendron lancigerum* Standl. & L. O. Williams, Ceiba 1. 232. 1951. TYPE: Costa Rica. Puntarenas: vic. Palmar Norte, Río Térraba, 30 m, Allen 5612 (holotype, US). Costa Rica. Puntarenas: along road between Chacarita and Rincón de Osa, ca. 6 km W of Interamerican Highway at Chacarita, 160 m, ca. 8°45'N, 83°18'W, Croat & Grayum 59728 (epitype, here designated, MO-3319112; isotypes, B, CR, F, K, NY, PMA, US).

*Philodendron taxilanum* G. S. Bunting, Gentes Herb. 9. 353. 1965. TYPE: Mexico. Veracruz: vic. Santiago Tuxtla, Bunting & Davies 162 (holotype, US).

Usually hemiepiphytic, rarely terrestrial or epilithic; stem appressed-climbing, parchment-white, sap turning blackish, slimy; internodes usually terete, weakly angular, sometimes obscurely flattened on one side or closely and acutely ribbed, semiglossy to matte, 1-20 cm long, 1.5-4 cm diam., dark green, becoming usually gray-green to brown, sometimes pale yellow-green, epidermis somewhat ridged ("wrinkled"), sometimes fissured transversely; roots reddish brown, 6-10 mm long, 3-5 mm diam., arising from and along the node on one side; cataphylls 18-39 cm long, usually weakly 1-ribbed, sometimes unribbed or weakly to sharply 2-ribbed or sharply 1-ribbed (ribs to 1 cm high), soft, green, sometimes reddish to pinkish, sometimes sparsely green-spotted, purple-maroon or darker striate near base, drying reddish brown, margins sometimes prominently

and thinly raised, usually soon deciduous, rarely persisting as a rotting mass, rounded at apex; **petioles** 20-65(91) cm long, 3-15 mm diam., usually subterete, moderately spongy to firm, medium green, drying greenish brown, obtusely flattened, usually obtusely and narrowly sulcate, rarely obtusely and broadly sulcate adaxially, surface minutely and densely short purple- or occasionally white-striate, sometimes dark green or violet-purple blotched, sometimes smooth to irregularly ribbed and folded; sheath subtending an inflorescence, 5-8 cm long; **blades** ovate to ovate-triangular, semiglossy, moderately coriaceous, weakly to moderately bicolorous, acuminate to narrowly acuminate, sometimes short-acuminate at apex, prominently cordate to  $\pm$  sagittate at base, 30-72 cm long, 15-39 cm wide (1.85-2 times longer than wide), (0.6-1.6 times longer than petiole), margins somewhat hyaline, weakly revolute, upper surface medium green, drying brownish green to greenish brown, semiglossy, lower surface pale yellow-green, sometimes reddish, drying usually reddish brown, sometimes yellowish brown, weakly glossy to matte; anterior lobe 24-61 cm long, 13-34.5 cm wide (2-5.1(5.5-6.4) times longer than posterior lobes); posterior lobes usually narrowly rounded, 6-20.5 cm long, 5-17.6 cm wide; sinus  $\pm$  V-shaped to almost closed, 6-15 cm deep; midrib flat to weakly raised, paler than surface, drying broadly raised and  $\pm$  concolorous above, convex to narrowly raised, sometimes maroon-spotted or with white flecks, concolorous to darker than surface, drying broadly raised and reddish below; basal veins 3-5(6) per side, with 0-1(2) free to base, third and higher order veins sometimes coalesced 2.5-4.5 cm, a few additional veins sometimes coalesced to 6.5 cm; posterior rib not at all naked or only briefly so; primary lateral veins 4-9 per side, departing midrib at a 60-70° angle, weakly curved to the margins, but usually turned prominently up just before the margin, rather prominently downturned just before the midrib, round-raised to flat to obtusely sunken and paler than surface above, convex, concolorous to darker than surface below; interprimary veins weakly sunken, concolorous above, flat, darker than surface below; minor veins weakly visible below, arising from both the midrib and primary lateral veins, drying moderately prominent, weakly undulate, alternating with dark, mostly contiguous secretory ducts. **INFLORESCENCES** erect, 1-3 per axil; peduncle 4-15 cm long, 1.6-1.8 cm diam., somewhat flattened to terete, green, sometimes tinged reddish, drying greenish, densely short and broad striate; **spathe** 8-22 cm long, 2-3 cm diam. (0.8-2.2 times longer than peduncle), weakly to obscurely constricted above the tube, semiglossy, usually green, sometimes



plum-red, often purple-spotted, densely short pale lineate throughout, weakly so near apex, blunt to narrowly cuspidate-acuminate to prominently acuminate, frequently tinged purplish violet at base; spathe blade green to pale yellow-green, 8–11 cm long (opening 3–4.3 cm wide, sometimes opening to near the base), greenish white, weakly tinged red-purple in throat to pale yellow-green or white inside, sometimes reddish throughout in age; resin canals appearing as continuous lines, red-purple to orange in color; spathe tube green, sometimes moderately to heavily tinged red-purple to red (B & K red-purple 3/7.5), with sparse, dark purplish spots (mostly medially) throughout outside, 4–7 cm long, 2.5–3.5 cm diam., red to reddish purple (B & K red-purple 3/7.5), white striate inside; **spadix** stipitate to 7 mm long; tapered toward apex, (8)9–16 cm long, broadest below the middle or near the base, usually protruding somewhat forward at anthesis but not curved; pistillate portion pale green (anthesis) to greenish white to yellow-green (post-anthesis), uniformly wide throughout or weakly tapered toward both ends, 2.5–6.2 cm long in front, 2–4.7 cm long in back, 1–1.2 cm diam. at apex, 1.1–1.7(2.1) cm diam. at middle, 1.1–1.7 cm wide at base; staminate portion 6.3–10.3(13) cm long; fertile staminate portion usually creamy white, sometimes pale green to pinkish, 9–13 mm diam. at base, 11–13 mm diam. at middle, 8–10 mm diam. ca. 1 cm from apex, broadest at base or middle, about as broad as the pistillate and sterile portions; sterile staminate portion often broader than the pistillate portion, white, (0.9–1)1.2–1.9 cm diam.; pistils (0.9)1.8–4.4 mm long, (0.9)1.3–2.3 mm diam.; ovary 6–9-locular, 1–1.7(3) mm long, 1.3–2.3 mm diam., with sub-basal placentation; locules 1–1.7(3) mm long, 0.2–0.4 mm diam.; ovule sac (0.6)0.8–1.2 mm long; ovules usually 2–4, rarely 5–8 per locule, 1-seriate (2-seriate, if 4 or more ovules), usually contained within translucent, gelatinous ovule sac, sometimes contained within gelatinous matrix (no true envelope), 0.1–0.5 mm long, usually longer than funicle; funicle 0.1–0.3 mm long (can be pulled free to base), sometimes adnate to lower part of partition, style (0.4)0.6–0.8 mm long, (0.8)1.2–1.6 mm diam., similar to style type B; style apex flat to sloping; stigma discoid or subdiscoid, truncate, (0.7)1–1.5 mm diam., (0.1)0.3–0.5 mm high, covering entire style apex, sometimes shallowly depressed at middle; the androecium truncate, prismatic, oblong, margins irregularly 4–6-sided to weakly ovate, ca. 1 mm long, 1.6–2 mm diam. at apex; thecae oblong, 0.4–0.6 mm wide,  $\pm$  parallel to one another, sometimes contiguous; sterile staminate flowers blunt, irregularly 4–6-sided, sometimes clavate or prismatic, 1.4–2.1 mm long, 1.4–2

mm wide. INFRUCTESCENCE pink, green at base with reddish spots outside; berries pale yellowish, rarely orange, with stigmas reddish brown, 7 mm long; seeds 2–3 per locule, drying pale brown to tannish brown, narrowly ellipsoid to oblong-ellipsoid, 1.4–2 mm long, 0.7–0.9 mm diam., with faint striations.

Flowering in *Philodendron sagittifolium* occurs almost throughout the year, principally after the onset of the dry season and continuing throughout much of the rainy season. There is a slight geographical shift, with flowering beginning about one month earlier in Mexico and Guatemala (January through August, less frequently in September and October) and continuing somewhat longer in Panama (February through September, but also rarely in December). Fruits apparently mature in about two months' time, but mature fruits have seldom been collected, only from January and July.

*Philodendron sagittifolium* ranges from Mexico (Veracruz) to Colombia (and probably also to Venezuela), from sea level to 1800 m elevation. It is probably the most morphologically variable, and one of the most ecologically versatile, species in Central America. In Mexico, this species occurs in "Selva Alta Perennifolia," "Selva Mediana Subperennifolia," "Selva Baja Caducifolia," and "Bosque Caducifolio." In the remainder of Central America, it occurs principally in *Tropical moist forest* and *Premontane wet forest* but also in *Premontane rain forest* and *Tropical wet forest* life zones.

*Philodendron sagittifolium* is a member of *P.* sect. *Calostigma* subsect. *Macrobelyum* ser. *Macrobelyum*. This species, though highly variable in most regards, can be characterized by its appressed-climbing habit; short, stout internodes; sharply two-ribbed, deciduous cataphylls; obtusely flattened, firm petioles usually spotted with violet-purple; and ovate-triangular, moderately coriaceous blades with the posterior rib not at all or only briefly naked along the sinus. Also characteristic is the externally green, frequently purple-spotted spathe, which is reddish purple on the tube within.

In Mexico and Guatemala, *Philodendron sagittifolium* is most easily confused with *P. advena* and *P. purulhense*, differing from both in having the blades somewhat triangular and drying reddish brown rather than generally ovate and drying blackened. See *P. advena* for additional discussion.

In Panama, *P. sagittifolium* may sometimes be confused with *P. annulatum*, which differs in having the petioles somewhat spongy with a purple distal ring, blades typically ovate-oblong, and spathes commonly white on the blade portion. It can be

confused at some stages of development with sterile specimens of *P. bakeri*.

Considering the highly variable nature of this species there are many noteworthy collections, only a few of which can be discussed here.

*Whitefoord & Eddy 222* from Panama has the lower blade surfaces drying yellowish brown rather than reddish brown and spathe solid plum-red outside. Also noteworthy is *Hammel et al. 14598*, which reports fruit color to be orange.

A few collections from Puntarenas Province, Costa Rica (*Croat 57243, 67697, Grayum & Hammel 10066*), differ in having the primary lateral veins paler, rather than darker on the lower surface. Sterile specimens from Cocos island (*W. Klawe s.n., Foster 4177*) differ in having the minor veins minutely raised on the upper surface. These specimens may prove to represent distinct species.

A large number of sterile and ostensibly juvenile collections from Nicaragua are of uncertain identity. One series, including *Pipoly 3826, 4055, 5144, Stevens 7628, 12673, 12739*, may ultimately prove to be *P. bakeri*. Two other collections (*Pipoly 5190, 5194*) might prove to be still another species.

*Croat 60804*, from the coastal cordillera of Venezuela, reported as *P. cf. sagittifolium* (*Croat & Lambert, 1986*) is either this species or a very close relative.

When Regel described *P. sanguineum* Regel, he cited no specimens or country of origin, but Krause (1913) cited a collection he had prepared from the Berlin Botanical Garden of a cultivated Karwinsky collection from Córdoba in Veracruz State, Mexico. While no such collection still exists, Karwinsky collected in Mexico only a few years (1840-1843) before Regel described *P. sanguineum*. Thus it is possible that both Regel and Engler could have been dealing with the same material originally collected by Karwinsky at Córdoba.

Another synonym that deserves mention is *P. lancigerum* Standl. & L. O. Williams, corresponding to a narrow-leaved form of *P. sagittifolium* restricted to the Pacific lowlands of Costa Rica. Epitypification was necessary because the type specimen (*Allen 5612*) consists of only an inflorescence.

*Additional specimens examined.* BELIZE. Cayo: Mountain Pine Ridge, *Bartlett 13052* (MICH); Blancaneaux Lodge, *Wiley 460* (MO); Water Hole, near Vaca, *Gentle 2447* (MICH); Río Ma Cal, San Luis-Cuevas, 1050 ft., *Croat 23497* (MO); 1050 ft., 23539 (MO). Toledo: Punta Gorda, *Cosminsky 227* (F); Richardson Creek, lower part, affluent of Bladen Branch, lower part of Maya Mountains, 100-250 m, 88°46'-48'W, 16°33'N, *Davidse & Brant 32334* (MO); Columbia road, *Gentle 6118* (LL); San Antonio, *Gentle 5510* (LL); Punta Gorda, *Cosminsky 226*

(F); Columbia River Forest Reserve, SW Maya Mountains, trail between Gloria Camp and Edwards Camp to the S, 16°22'N, 89°10'W, *Holst 4499* (MO). COSTA RICA. Alajuela: 3 mi. N of San Miguel, 380 m, *Croat 35642* (CR, MO); 8.9 mi. NW of San Ramón, 1100 m, 10°10'30"N, 84°30'W, *Croat 68070* (L, MO); Río Sarapiquí, road to Colonia Virgen del Socorro, 830 m, 10°1'6"N, 84°11'W, *Croat 68338* (MO). Cartago: Tapanti, ca. 1350 m, 9°47'N, 83°47.5'W, *Grayum 3890* (MO); 1400 m, 9°47'N, 83°48'W, 3322 (K, MO); 31 km S of Siquirres, 850 m, 9°57'N, 83°36'W, *Thompson & Rasdwin 1159* (CM); 1160 (CM); 7 km W of Turrialba, *Croat 36829* (MO); Turrialba, Moravia de Chirripó, 1200 m, *Campes & Chasarría 40* (CR, INB); Río Peñibaye, 2 km SW of Taus, 750 m, 9°29'N, 83°41'W, *Lent 2967* (CR, F). Heredia: La Selva Field Station, *Hammel 10244* (MO); *Grayum 2306* (MO); *Hammel 9641* (DUKE); La Zona Protectora, SSE of Magnessay, *Schatz 722* (CR, DUKE, MO); 4 mi. N of Vars Blanca, 1350 m, *Croat 35579* (MO). Limón: vic. of Moín, 0 m, 10N, 83.04W, *Croat 61206* (MO); Parque Nacional Tortuguero, 2 km W of Tortuguero, 3 m, 10°31'N, 83°30'W, *Robles 2166* (CR, MO); 2178 (CR, MO). Puntarenas: Palmar Sur-Piedras Blancas, 20 m, *Croat 32917* (MO); Quebrada Aparicio-Quebrada Aguabuenas, Rincón de Osa, 200-400 m, 8°42'N, 83°31'W, *Grayum et al. 4017* (CR, MO); Rincón de Osa, region to W of airstrip, 40-100 m, *Uley & Uley 1119* (F); San Vito de Coto Brus-Ciudad Neily, 300-600 m, 8°41'N, 82°56.5'W, *Hammel 14159* (CR, MO); El Campo, Aguabuenas, 350 m, 8°42'50"N, 83°31'42"W, *Herrera 3959* (INB, MO); Las Brisas-Las Juntas de Coto Brus, ca. 900 m, 8°52'N, 82°57'W, *Grayum & Hammel 5692* (INB, MO); Parque Nacional Corcovado, Sirens, 1-10 m, 8°28'N, 83°35'W, *Kernan & Phillips 1028* (MO); 1-50 m, *Kernan & Phillips 1089* (MO); *Kernan 461* (CR, MO), 574 (CR, MO); 10 m, *Aguilar 3396* (INB); Río Claro, 5 m, *Kernan 783* (CR, MO); Las Cruces Botanical Garden-Río Jaba, ca. 3 km SW of San Vito de Coto Brus, ca. 1050-1200 m, 8°47'N, 82°58'W, *Grayum 5624* (CM, CR, INB, MEXU, MO, U); San Vito de Java, ca. 4000 ft., *Croat 32904* (MO); 1 mi. S of San Vito de Java, ca. 3500 ft., *Raven 21882* (BM, DS, F); 6 km S of San Vito de Java, 4000 ft., *Raven 22009* (DS, F, WIS); Las Cruces Botanical Garden, ca. 1300 m, *Croat 44440* (MO); Gollito, near TV transmission tower, 2000 ft., 8°49'N, 82°58'W, *Croat 57243* (MO), 57231 (MO); Finca Loma Linda, 1 mi. SW of Cañas Gordas, 1150 m, *Croat 22287* (F, MO), 22257 (MO); hills above Palmar Norte, 100-200 m, *Croat 35141* (MO); Punta Cathedral, ca. 7 km SE of Quepos, 20-70 m, 9°22.5'N, 84°09'W, *Grayum & Sleeper 5903* (MO); Punta Quepos (3 km S of Puerto Quepos), 0 m, 9°24'N, 84°10'W, *Grayum 6613* (CR, MO, PMA, U); road to Rincón de Osa, 16.5 km W of Inter-American Highway, 280 m, 8°45'N, 83°22'W, *Grayum et al. 7550* (INB, MO); Reserva Forestal Golfo Dulce, S of Rincón de Osa, 150 m, 8°37'N, 83°28'W, *Hammel & Robles 16728* (CR, MO); 150-200 m, 16739 (CR, MO); 450 m, *G. Herrera 3953* (CR, INB); ca. 0.7 km N of Gollito-Villa Bricheño road, 160-260 m, 8°40'30"N, 83°12'W, *Grayum & Hammel 10066* (CR, MO); Cantón de Gollito, Jiménez, Alto de la Carbonera, road to Cerro de Osa, 200-350 m, 8°25'30"N, 83°19'W, A. *Chacón 1069* (CR, INB, MO); 3.1 mi. NW of center of Gollito, 30 m, 8°11'N, 83°12'W, *Croat 67632* (CM, CR, MEXU, MO, WIS); Piedras Blancas-Rincón Road, Mile 3.7, 90-105 m, *Croat 67697* (CR, MO); Cocos Island, valley behind Bahía Iglesias, *Foster 4177* (F, MO); *Klawe s.n.* (POM). San José: San Isidro de El General-Dominical, SW of San Isidro,

4.8 mi. from Río Pacuare, 1000 m, *Croato* 35254 (MO); 9 mi. SW of Río Pacuare, 680 m, 35343 (MO); ca. 0.5 mi. above turnout to Cañanán at Rivas, 900 m, *Croato* 43430 (CR, MO); 43415 (MO); Estación Carrillo, Fila Cañón del Río Sucio 450–700 m, *A. Chacón & Herrera* 1705 (CR, MO); Carara—El Sur de Turruabares, 200–310 m, 9°45'15"N, 84°21'30"W, *Grayum* 10451 (CR, INB, MO); Z. P. La Cangreja, Santa Rosa de puriscal, 400 m, 9°42'50"N, 84°23'30"W, *Morales* 1468 (CR, MO); Acosta, Valle del Candelaria, 1000–1050 m, *Morales & González* 4581 (CR, INB); Tarraza, vic. Hormiguero, 1100–1200 m, *Croato* 78938 (INB, MO); 78959 (CR, INB, MO); 990–1100 m, *Croato & Hannon* 79110 (MO); Parque Nacional, sector Esquinas, vic. Fila Gamba, 200–300 m, *Croato & Hannon* 79294 (INB, MO). GUATEMALA. Eastern portions of Verapaz and Chiquimula, *Watson* 106 (GH). **Alta Verapaz:** E of Tactic, 1300 m, *Steyermark* 43993 (F, US); near Chiriacé, on the Petén Highway, ca. 900 m, *Standley* 92189 (F); Chapultepec Farm, 62 km from Cobán on Sebol Road, *Contreras* 4823 (LL); Río Careh, Cobán—San Pedro Careh, ca. 1360 m, *Standley* 89891 (F). **Baja Verapaz:** Mpio. Purulhá, El Progreso—Cobán, Hwy. CA-14, 17 mi. N of junction with Hwy. 17, 1620–1720 m, 15°13'N, 90°12'W, *Croato & Hannon* 63766 (CAS, CR, EAP, MO, SAR, USCG). **Chimaltenango:** Quisacah, 1800 m, *Standley* 62029 (F). **Izabal:** Escobas—Santo Tomás, Santo Tomás Bay, 0–2 m, *Steyermark* 39331 (F); between Milla 49.5 and ridge 6 mi. from Izabal, Montaña del Mico, 65–600 m, *Steyermark* 38506 (F, US); vic. EXMIBAL Camp 2 (La Gloria), NW of Lake Izabal, 400–500 m, *Jones & Facey* 3283 (NY); Sierra de las Minas, 13 km E of Doña María, 240 m, ca. 15°14'N, 89°20'W, *Harmon & Fuentes* 5857 (UMO); Río Frío, 75–150 m, *Steyermark* 41533 (F, LL). **Petén:** Canchacán, near San Luis, *Landell* 16325 (LL); Dolores, 300 m, *Contreras* 2562 (LL). **Quezaltenango:** La Shuya, SW of San Martín Chile Verde, ca. 1620 m, *Standley* 67882 (F). **HONDURAS.** Mont. de la Flor, 3300 ft., *von Hagen & von Hagen* 1304 (F, NY). **Atlántida:** Quebrada Grande, ca. 10 km SW of La Ceiba, 80–200 m, 15°42'N, 86°51'W, *Liesner & Mejía* 26011 (MO); 26352 (MO); 80–180 m, 15°42'N, 86°51'W, *Liesner* 26179 (MO). **Comayagua:** El Achote, above Siguatepeque, 1350 m, *Yuncker et al.* 5994 (F, GH, K, MO, NY, U). **El Paraíso:** El Jurquillo—El Robledal, Sierra el Chile, 1300 m, *Molina* 14142 (EAP, F, NY). **Olancho:** Catacamas—La Presa, N of Catacamas, 500–600 m, *Standley* 18698 (EAP); Neñón Hagen—TEFH, US); Río Olancho, San Esteban—Bonito Oriental, 14.8 mi. NE of San Esteban, 635 m, 15°25'N, 85°47'W, *Croato & Hannon* 64405 (DUKE, LL, MO); San Francisco de la Paz—Gualaco, 7.3 mi. NE of San Francisco de la Paz, 1130 m, 14°58'N, 86°12'W, *Croato & Hannon* 64197 (B, BM, CM, MICH, MO, NY, QCA); 13.6 mi. SW of Gualaco, 1300 m, 15°N, 86°07'W, *Croato & Hannon* 64295 (F, K, MO, NY); Mpio. La Unión, 6 mi. E of La Unión along road to Olanchito, 950 m, 15°03'N, 86°35'W, *Davidse et al.* 35428 (MO). **Yoro:** Cordillera Nombre de Dios, El Guaymón—San José de Texiguat, Cerro Aguacatalas, 830 m, 15°29'N, 87°37'W, *Davidse et al.* 34524 (EAD, MO, TEFM); 34524 (EAD, MO, TEFM); Río Pijal Valley, 6–7 km S of Nueva Esperanza, 1570–1670 m, 15°12'N, 87°35'W, *Liesner* 26581 (MO). **MEXICO.** **Patria, Engler** 198 (GH, P). **Chiapas:** Mpio. Ocosingo, 16 km NW of Boca Lacantún, 900 m, *Martínez* 21345 (MEXU, MO); 12 km N of Monte Líbano, 950 m, *Martínez* 21252 (MEXU, MO); Cascada Corralita, SW of Ocosingo, 600 m, 16°59'N, 92°08'W, *Hammel et al.* 15677 (MO); Palenque—Bonampak, 69 mi. SW

of Palenque, 450 m, *Croato* 40197 (F, MO), 6 mi. N of Ocozacoautla, 1000 m, *Croato* 40581 (B, MEXU, MO); Palenque—Ocosingo, Hwy. 199, 43 mi. SW of Palenque, 810 m, *Croato* 40337 (MO); 13 mi. N of Ocozacoautla, 900 m, *Croato* 40632 (MO); San Cristóbal de Las Casas—Palenque, Villa Paraiso, 136 km NE of San Cristóbal, 450 m, 17°29'N, 92°05'W, *Hammel et al.* 15641 (MO); 18–20 km N of Ocozacoautla, 800 m, *Breedlove* 25202 (DS); 3 mi. N of Pueblo Nuevo Solistahuacán, 1700 m, *Breedlove & Smith* 32436 (DS); 6700 ft., *Thorne & Lathrop* 40267 (RSA). **Morales:** Cuernavaca, Barranca del Salto de San Antonio, *Moore & Bunting* 8823 (BH); Iurbide, *Bougeau* 1417 (K, P). **Oaxaca:** Latani, Dto. Choapan, 900 m, 17°24'N, 95°48'W, *Schulze & Reko* 900 (ECON, GH); Sierra de Juárez, Tuxtpec—Oaxaca, Highway 67, 6–14 mi. above bridge at Valle Nacional, ca. 580 m, *Moore & Bunting* 8902 (BH, CM); 8892 (BH); 660 m, *Croato* 39752 (MO); 39757 (MO); 43917 (CM, MO); 1400 m, *Croato* 48000 (L, MEXU, MO); 705 m, 17°44'N, 96°19'W, *Croato & Hannon* 65537 (B, MO); 55 km S of Tuxtpec, 630 m, 17°37'N, 96°20'W, *Hammel & Merello* 15482 (MO, NY); Uxpanapa region, Esmeralda—Río Verde, 100 m, 17°10'N, 94°45'W, *Croato & Hannon* 63237 (CR, MO, VEN); 63299 (MO); Palomares—Matías Romero, road to Uxpanapa, 44 km E of entrance at Sarabia, on Rte. 185, 125 m, 17°09'N, 94°37'W, *Hammel & Merello* 15556 (MO); 15555 (MO); Uxpanapa, 14 km E of Campamento La Laguna (Poblado D.S.), 150 m, 17°05'N, 94°35'W, *Hammel & Merello* 15578 (MO); Río Uxpanapa, *Maranda* 38666 (MEXU); Ixtlán, Mpio. Comaltepec, Puerto Antonio, ca. 1300 m, 17°45'N, 96°30'W, *Luna* 436 (MO); Highway 175, 1 km above Vista Hermosa, 1255 m, 17°35'N, 96°21'W, *Boyle & Boyle* 684 (MO). **Veracruz:** Conejo—Huatusco, Hacienda El Mirador vicinity, km 45 on highway, *Moore & Bunting* 8862 (BH, MO); 8863 (BH, MO); El Mirador, 8857 (BH, MO); Fortín de las Flores (Posada Loma), 850 m, *Nesling & Gómez-Pompa* 287 (cultivated, XAL); Catemaco—Acayucan, ca. 8 mi. S of Catemaco, near Zapopan, ca. 400 m, *Moore & Bunting* 8925 (BH); Fortín, Cervecería Motezuma, 1000–1150 m, *Croato* 39415 (CM, F, MO); 900 m, *Croato* 44038 (MO); 5.7–6 mi. from Catemaco, road to Sontecomapan, ca. 380 m, *Moore & Bunting* 8941 (BH); Playa Montepío, 0 m, 18°38'N, 95°05'W, *Hammel & Merello* 15498 (BR, CM, MO); Coatzacoalcos River, Fortuño, 30–50 m, *Williams* 8702 (F); Estación de Biología Triocacac “Los Tuxtles,” ca. 31 km by road N of Catemaco, along road to Montepío, 250–300 m, 18°60'N, 95°07'W, *Grayum & Steeper* 8355 (MO); 300 m, 18°44'N, 85°10'W, *Hammel & Merello* 15490 (CM, MO); 170–200 m, *Gentry et al.* 32488 (MO); 250 m, *Ibarra* 589 (MO); Mpio. Coatepec, Tuzamapan—Jalcacmulco, Cerro de Achicubca, 700 m, 19°23'N, 96°48'W, *Castillo & Tapia* 678 (F, MO); Mpio. Hidalgotitlán, toward La Laguna, 160 m, 17°16.6'N, 94°33.4'W, *Valdivia* 744 (XAL); Hidalgotitlán, 150 m, *Valdivia* 1351 (XAL); near La Escudera, 160 m, 17°19'N, 94°38.5'W, *Valdivia* 1330 (XAL); Mpio. J. de Ferrer, Cerro de Villa Rica, near Plan de la Flor, 1250 m, 19°48'N, 96°46'W, *Castillo et al.* 1715 (F); Mpio. Jesús Carranza, 2 km N del Poblado 2, Ejido F. J. Mina, 120 m, 17°16'N, 94°40'W, *Vásquez & Novarrese* 2383 (MO); Mpio. Pajapan, Volcán San Martín Pajapan, 6 km NW of Pajapan, 1200 m, 18°18'20'N, 94°43'30'W, *Nee et al.* 24974 (F, MO); Sontecomapan—Montepío, N of San Andrés Tuxtla, 150–200 m, 18°42'N, 95°10'W, *Croato & Hannon* 63158 (MO, NY, US); Mpio. Tlapacoyan, Río Tablazos, Tlapacoyan—Altoingá, 6 km SSW of Tlapacoyan, 19°56'N, 97°13'W, *Nee et al.* 26090

(F. MO, NY); Vallé de Cédova, *Bourgeau* 2296 (K, P). NICARAGUA. **Estelí:** El Zacatón (Plan Helado), ca. 1300 m, 13°13'N, 86°14'W, *Moreno* 17521 (MO); Laguna de Miraflores, ca. 1200 m, *Grijalva* 952 (MO).  **Jinotega:** ca. 5–8 mi. SW of Jinotega, 1500 m, *Croat* 43084 (CM, MO); San Ramón–Paricutin, Comarca Kilambé, SE of Cerro Kilambé, 800–900 m, 13°35'N, 85°40'W, *Moreno* 7687 (MO); along Hwy. 3 ca. 1 km NW of La Fundadora entrance, 1450–1520 m, 13°01'N, 85°56'W, *Stevens* & *Henrich* 20435 (MO), Las Camelias–La Salvadora, 1100–1150 m, 13°05–06'N, 85°53–54'W, *Stevens et al.* 15327 (MO); N slope of Volcán Yalí, 1200–1400 m, 13°15'N, 86°10'W, *Stevens et al.* 15091 (MO); Matagalpa–Jinotega Road, Km 146, 1200–1400 m, *Moreno* 571 (MO); Laguna Miraflores, ca. 26.1 km NE of Hwy. 1 at Estelí, 1250–1300 m, ca. 13°15'N, 86°15'W, *Henrich* & *Stevens* 243 (MO); Las Camelias–La Salvadora, road from Hwy. 3 through La Fundadora, 1100–1150 m, ca. 13°05–06'N, 85°53–54'W, *Stevens* & *Grijalva* 15322 (MO); Macizos de Peñas Blancas, vic. of Finca de Manuel Estrada (El Cielo), 1200–1400 m, ca. 13°15'N, 85°41'W, *Stevens* 11665 (MO); 1500–1650 m, ca. 13°15–16'N, 85°41'W, *Stevens* 11430 (MO); Río Bocay, Salto Kayaska, 190–340 m, ca. 13°51'N, 85°22'W, *Stevens et al.* 16479 (MO). **Matagalpa:** Valle de Arenal, on road to Sanatorio de Aranjuez, 1400 m, 13°02'N, 85°55'W, *Davidse et al.* 30444 (MO); 13°01'N, 85°54'W, *Moreno* 9583 (MO); Fuente Pura, km 142, 1400–1450 m, 13°N, 85°55'W, *Moreno* 17017 (MO); road to Cerro La Carlota, 2 km from Tuma, 1040–1100 m, 12°58'N, 85°52'W, *Moreno* 15659 (MO); Cerro Picacho, behind La Selva Negra Hotel, 1200–1540 m, 13°N, 85°55'W, *Davidse et al.* 30311 (MO). **Nueva Segovia:** Río San Fernando Valley, Cerro El Peñascal, 800–1125 m, 13°2–3'N, 86°19–20'W, *Stevens* 3297 (BM, MO). **Río San Juan:** Bocas de Sábalo, 70–100 m, 11°03'N, 84°27'W, *Moreno* 26692 (MO). **Zelaya:** El Empalme–Limbaika, junction of road to Alamikamba, ca. 25 m, ca. 13°32'N, 84°30'W, *Stevens* 12739 (MO); road to Mina Nueva América, ca. 10 km N of main road, *Stevens* 12673 (MO); ca. 11.3 km N of main road, *Pipoly* 5283 (MO); ca. 6.3 km S of bridge at Colonia Yolania, ca. 200–300 m, ca. 11°36–37'N, 84°22'W, *Stevens* 4824 (MO); Cerro El Horniguero, W range, ca. 1100–1183 m, ca. 13°44'N, 85°W, *Pipoly* 5198 (MO); 5194 (MO); 5190 (MO); Cerro La Pimienta number 1, ca. 900–980 m, ca. 13°45'N, 84°59'W, 5107 (MO); 5114 (MO); ca. 13 km above Kururia, on road to San Jerónimo, <200 m, *Pipoly* 3826 (MO); 1 km S of Monkey Point, ca. 5 m, 11°35'N, 83°39'W, *Moreno* & *Sandino* 12531 (MO); Monkey Point, 1.5 km NW, 1–5 m, 11°36'N, 83°38'W, *Moreno* & *Sandino* 11957 (MO); Río Kisalaya, near Tala Has and Puente Mango, 40–60 m, ca. 14°41'N, 84°03'W, *Stevens* 7628 (BM, MO); Río Mico, near El Recreo, ca. 30 m, *Standley* 19404 (F); 19525 (EAR, F), 19614 (F); Río Pis Pis, 0.5–1.5 km from Plantel El Salto, road to Bonanza, ca. 140 m, ca. 14°03'N, 84°37'W, *Stevens* 18871 (MO); Río Punta Gorda, Atlanta, ca. 10 m, 11°34'N, 84°01'W, *Moreno* & *Sandino* 12770 (MO); Waspan–Puerto Cabezas, Río Troncera, <200 m, ca. 14°43'N, 84°06'W, *Pipoly* 4055 (MO). PANAMA. **Bocas del Toro:** Gualaca–Chiriquí Grande, 4.2 mi. E of Chiriquí Grande, ca. 0 m, 8°55'N, 82°09'W, *Croat* 66814 (MO); 0.4 mi. SE of Punta Peña, 120 m, 8°57'N, 83°56'W, *Croat* 74939 (AAU, CR, MO, NY); Fortuna Dam area, N of Fortuna Dam, 650–700 m, 8°45'N, 82°15'W, *McPherson* 11131 (MO); Chiriquí Grande–Fortuna, 1.6 mi. N of Continental Divide, 770–790 m, 8°45'N, 82°17'W, *Croat* & *Zhu* 76453A (MO), 76456A (CM, MO); 10 mi. N Divide,

1 mi. N from highway, 130 m, 8°46'N, 82°11'W, *Hammel et al.* 14598 (MO); 4.3 km N of Divide, 590 m, 8°46'N, 82°14'W, *Croat* & *Grayum* 60210 (CM, MO); 3.2 mi. N of Divide, 700 m, 8°45'N, 82°15'W, *Croat* & *Grayum* 60259 (CAS, CR, F, KYO, M, MEXU, MO, PMA). **Canal Area:** Frijoles–Monte Lirio, 30 m, *Killip* 12133 (US); hills N of Frijoles, *Standley* 27598 (US); Pipeline Road, 14 mi. NW of Gamboa gate, *Croat* 12363 (MO); Barro Colorado Island, *Croat* 6472 (F, MO, RSA, SCZ); *Aviles* 25 (F); 46 (F); Gatún Lake, *Standley* 31398 (US); *Croat* 6531 (F, MO, SCZ), 6334 (MO), 5052 (MO), 10901 (MO), 8290 (MO), 10659 (MO), 9018 (MO), 8999 (MO), 10076 (MO); *Silvestre s.n.* (MO); *Croat* 6023 (MO); Summit Gardens, *Croat* 59139 (MO), 17155 (MO), 10768 (MO), 17060A (MO). **Chiriquí:** Finca Ojo de Agua, 1300 m, 8°51'N, 82°46'W, *Knapp* 1590 (MO); Cañas Gordas–Volcán, 1 m E of Cañas Gordas, ca. 1000 m, *Croat* 22304 (MO); 8 mi. W of Puerto Armuelles, in vic. of San Bartolo Límite, ca. 600 m, *Croat* 22022 (MO), 22026 (MO); Cerro Colorado, above San Félix, 15–18 mi. N of Pan-American Highway, 800–1000 m, *Croat* 33179 (CM, MO); Gualaca–Chiriquí Grande, Fortuna Dam area, 7.2 mi. beyond Los Planes de Hornito, 1165–1200 m, 8°44'N, 82°14'W, *Croat* 67843 (CM, MO); 5.9 mi. NW of Los Planes de Hornito, 1225 m, 8°45'N, 82°14'W, *Croat* 67795 (CR, MO, PMA); 1400 m, *Croat* 48719 (MO); 48697 (MO); 48681 (MO); Gualaca–Chiriquí Grande, 8 mi. beyond Los Planes de Hornito, trail to Río Hornito, 1010–1130 m, 8°44'N, 82°14'30"W, *Croat* 67918 (CAS, COL, L, MEXU, MO, PMA, US); 10.1 mi. NW of Los Planes de Hornito, 1250 m, 82°17'W, 8°45'N, *Croat* 50040 (CM, MO); Quebrada Los Chorros–Quebrada Honda, N of Fortuna Lake, 1100 m, 8°45'N, 82°14'W, *Churchill* & *Churchill* 6107 (MO); 4.5–5 km N of dam over Fortuna Lake, 1100–1135 m, 8°43'N, 82°17'W, *Croat* & *Grayum* 60056 (B, CAS, F, L, MBM, MO, P, PMA); Cerro Colorado, 2.3 km above Chacra, 1000 m, *Croat* 37067 (MO). **Coclé:** Penonomé–Coclequito, Río Cascajal, 5.7 mi. N of Llano Grande, 210 m, 8°40'N, 80°26'W, *Croat* 67541 (CR, MO); El Copé region, 9.4 km above El Copé, 750–900 m, *Croat* 44726 (MO); Alto Calvario, 710–800 m, *Croat* 68716 (EAP, G, HNMN, MO, TEX); 8°39'N, 80°36'W, 68849 (MO); 580–740 m, 8°38'N, 80°36'W, *Croat* 67522 (MO, US); El Valle region, La Mesa, above El Valle de Amón, 860–900 m, *Croat* 37407 (MO); 870 m, 8°50'N, 80°07'W, *Hoover* 1320 (MO); 800–900 m, 8°38'N, 80°09'W, *Croat* 67152 (B, CAS, CM, COL, DUKE, F, K, L, MEXU, MO, NY, PMA, QCA, TEX, VDB, WIS); 800 m, 8°36'N, 80°07'W, *Croat* & *Zhu* 76691 (MO); Cerro Gaital, *Churchill* 3907 (MO). **Colón:** Sabanas–Portobelo, Río Piedras Lumber Road, 250 m, 9°22'30"N, 79°41'30"W, *Croat* 75159 (MO); Río Indio–Miguel de la Borda, 0 m, *Croat* 36906 (MO); Miguel de la Borda, *Croat* 9856 (F, MO, RSA, US); Santa Rita Ridge Road, 10.6 km from highway, ca. 380 m, *Croat* 34345 (MO); Mile 6.5, 370 m, 9°21'15"N, 79°44'W, *Croat* & *Zhu* 76964 (MO); along Río Guaniche ca. 2 km E of bridge on the main Puerto Pilón–Portobelo Road, 100 m, 9°30'N, 79°39'W, *Croat* 75176 (MO). **Darién:** Cerro Pirre region, middle slopes on W side, 550–760 m, 7°57'N, 77°46'W, *Croat* 68958 (MO); 68893 (MO); Río Cana–Río Escucha Ruido, above Cana Gold Mine, 1310–1430 m, *Croat* 37755 (MO); 37818 (MO); 37827 (MO); S of El Real, Alturas de Nique, near Cana mine, along old Camino Real toward Colombia, 800–1000 m, 7°45'N, 77°40'W, *McPherson* 11536 (MO); Río Cocalito, ca. 5 mi. from P. Cocalito, *Whiteford* & *Eddy* 2222 (BM, MO); Parque Nacional del Darién, Panama–Colombia border, Río Pucuro, Cerro Tacarcuna, ca. 6

km N of Cerro Mali, 1300–1500 m, 8°09.5'N, 77°15'W, *Hammel et al.* 16537 (MO, PMA). **Herrera:** Las Minas, Primer Ciclo de Chepo, 900 m, 7°43'N, 80°50'W, *Galdames et al.* 1765 (MO); 18 km W of Las Minas, Alto Higu, N slope, 2400 ft., *Hammel* 4227 (MO). **Los Santos:** Azuero Peninsula, Jobero-Río Pedregal, 300–700 m, *Croat* 34479 (MO); Río Guanico valley, 600 m, 7°18'N, 80°30'W, *McPherson* 9245 (MO, NY, SCZ). **Panamá:** Cerro Azul, *Croat* 17281 (MO); El Llano-Cartí Road, ca. 16–18 km N of Pan-American Highway, 400 m, *Tyson & Nec* 7352 (MO); Mile 5–9, 200–250 m, 9°15–16'N, 78°59'N, *Thompson* 4624 (CM); Mile 10, 330 m, *Croat* 33823 (CM, F, MO, TEX, WIS); Mile 7, 460 m, 75109 (MO); Km 7–12, 360–400 m, 25096 (MO); Mile 8.2, 450 m, 9°14'N, 79°W, *Knapp & Hufi* 4412 (MO); Mile 12, 200–500 m, *Licner* 695 (MO); Mile 5–9, 200–250 m, 9°15–16'N, 78°59'W, *Thompson* 4636 (CM); Cerro Campana, 800 m, 8°41'N, 79°56'W, *Croat* 74770 (AAU, MO); 74771 (MO, PMA); 17177 (MO); ca. 850 m, 8°40'N, 79°50'W, *Thompson* 4577 (CM); 700 m, 79°55'W, 8°40'N, *Hamilton et al.* 1174 (MO); *Hammel* 5561 (MO); ca. 1 mi. from Inter-American Highway, ca. 150 m, *Croat* 35983 (MO); 12063 (MO); 800 m, 8°41'N, 79°56'W, 74770 (MO); Cerro Jefe region, near summit 750–800 m, 9°14'N, 79°22'W, *Croat* 67087 (F, MO); 800–1000 m, *Gentry* 2892 (GH, MO); 770 m, 9°15'N, 79°29'W, *Croat & Zhu* 76608 (CR, GB, MBM, MEXU, MO, NY); 4.6 km beyond peak, on road to Altos de Pacora, 26.3 km from highway, ca. 600 m, *Croat* 35933 (MO); 3–3.5 mi. NE of Altos de Pacora, 11.1–11.6 mi. beyond Lago Cerro Azul, 700–750 m, 9°15'N, 79°25'W, *Croat* 68695 (MO, PMA). **San Blas:** beach E of Puerto Obaldia, *Croat* 16910 (MO); 16911 (MO); 16914 (MO); 7 mi. N of Inter-American Highway, 550 m, 9°43'N, 78°68'W, *Croat* 60506 (MO); Cerro Brewster, 800–850 m, 9°18'N, 79°16'W, *de Nevers et al.* 6287 (MO); El Llano-Cartí Road, vic. Nusapandi, 450 m, 9°18'N, 79°59'W, *Croat* 75150 (CM, MO, PMA); 350 m, 9°20'N, 79°W, *Croat & Zhu* 77012 (CM, MO). **Veraguas:** Dist. Montijo, Isla de Coiba, 7°37'N, 81°44'W, *Galdames et al.* 2213 (PMA); 2073 (CM, MO, PMA); 400 ft., *Antonio* 2383 (MO); 2343 (MO, PMA); 400 m, 7°31'N, 81°51'W, *Galdames et al.* 2284 (MO, PMA); 50 m, 7°35'N, 81°47'W, *Galdames* 2256 (MO, PMA); Santa Fe region, 7 km W of Santa Fe, 2900 ft., *Nec* 11192 (MO, RSA); Cerro Tute, to 1200 m, *Witherspoon et al.* 8873 (MO); Santa Fe-Río San Luis, at Río Segundo Brazo, 480 m, 8°33'N, 81°08'W, *Croat* 66886 (MO, PMA); 66916 (MO, PMA), 66920 (MO).

***Philodendron scalarinerve*** Croat & Grayum, sp. nov. TYPE: Panama. Panamá: road past and 3–3.5 mi. NE of Altos de Pacora, 7.8–8.2 mi. above Pan-American Hwy., 11.1–11.6 mi. beyond Lago Cerro Azul, 700–750 m, 9°15'N, 79°25'W, *Croat* 68692 (holotype, MO-3585744; isotypes, B, CAS, COL, CR, F, K, NY, PMA, US). Figures 357–363.

Planta hemiepiphytica; internodia 1–1.5(3) cm longa, 1.5–3.5(4) cm diam.; cataphylla usque 22 cm longa, in-costata vel obtuse 1-costata, raro acute 2-costata, persistentia semi-intacta; petiolus subteres, 20–75 cm longus, 0.7–1.5 cm diam.; lamina oblong-elliptica vel leniter cordata, 33–67 cm longa, 11.4–40 cm lata, cum nervis conspicue scalariformibus inter nervos minores; inflores-

centia 1–4; pedunculus 6–20 cm longus, subteres; spatha 16–21.5 cm longa, pallide viridis; lamina spathae seminitida, exius striata alba apice, intus viridi diluta; tubo spathae hebetato, exius striato albo basi, intus viridibulo; pistilla 4–6(7–8)-locularia; loculi (10)12–14-ovulati.

Appressed-climbing hemiepiphyte; stems moderately short; leaf scars conspicuous, 2–4 cm wide; internodes 1–1.5(3) cm long, 1.5–3.5(4) cm diam., about as long as broad; roots moderately few per node, short, drying dark brown, matte, longitudinally ridged; cataphylls moderately thick, to 22 cm long, unribbed to bluntly 1-ribbed, rarely sharply 2-ribbed, drying dark brown, persisting semi-intact, eventually fibrous at upper nodes. LEAVES erect-spreading, clustered at or near stem apex; petioles 20–75 cm long, 0.7–1.5 cm diam., subterete, stiff, green, obtusely flattened, sulcate adaxially; sheathing 1–2.5 cm long; geniculum scarcely thicker than petiole, 6–7.8 cm long slightly paler than petiole; blades oblong-elliptic to narrowly ovate, moderately coriaceous, acuminate at apex (the acumen sometimes inrolled, to 2 mm long), obtuse to subcordate at base, 33–67 cm long, 11.4–40 cm wide (1.2–4 times longer than wide), (0.82–2.11 times longer than petiole), broadest at the middle, margins weakly undulate, upper surface dark green, semiglossy, lower surface much paler, matte to semiglossy, obscurely dark-punctate at 10× magnification; midrib convex to narrowly raised, sometimes paler than surface above, convex to thicker than broad and concolorous below; basal veins 1–4 per side, with all free to base; primary lateral veins 5–18 per side, departing midrib at a 60–75° angle, arcuate-ascending to the margins, sunken above; minor veins arising from the midrib only; with scalariform “cross-veins” weakly visible when fresh, conspicuously raised above on drying, barely visible below. INFLORESCENCES spreading, pendent, shorter than leaves, 1–4 per axil; peduncle 6–20 cm long, subterete, white-streaked near apex, spongy; spathae 16–21.5 cm long (0.6–1.4 times longer than peduncle), pale green, acuminate at apex, obtuse at base, convolute to about the middle at anthesis; spathae blade semiglossy, white-streaked at apex outside, light green inside; spathae tube matte, white-streaked at base outside, 5–7 cm long, 2–3.5 cm diam., greenish white inside; spadix sessile, erect, (7.6)9–12 cm long, broadest at the base or at the middle or below the middle; pistillate portion light green, slightly tapered above middle or toward both ends, (1.7)2.6–4.3(7.8) cm long, 0.8–1.3 cm diam. at apex, 0.8–1.4 cm diam. at middle, 0.9–1.4 cm wide at base; staminate portion 4.8–8.7 cm long; fertile staminate portion (9)12 mm diam. at base, (9)12 mm diam. at middle, (6)10 mm diam.

ca. 1 cm from apex, broadest at the base, slightly narrower than the pistillate portion, slightly narrower than the sterile portion; sterile staminate portion as broad as the pistillate portion, 1.2–1.4 cm diam.; pistils (1.4)2.3–2.8(3.2) mm long, (1)1.4–1.9 mm diam.; ovary 4–6(7–8)-locular, (0.7)1.5–1.8(3) mm long, (1.1)1.5–1.9 mm diam., with axile placentation; locules (0.7)1.5–1.8(3) mm long, 0.4–0.6 mm diam.; ovules (10)12–14 per locule, 2-seriate, (0.1)0.3–0.4 mm long, longer than funicle; funicle 0.1–0.2(0.3) mm long, adnate to lower part of partition, style 0.5(0.7) mm long, 1.1–1.4(2.2) mm diam., similar to style type B; style apex usually flat, sometimes rounded or domed, with stigma lobes over depressions (if apex is flat or domed); stigma usually lobed, sometimes subdiscoid or truncate, lobed, 0.9–1.2(1.4–1.7) mm diam., (0.1)0.2–0.3 mm high, covering entire style apex, inserted on center of style apex, sometimes medially depressed; the androecium truncate, prismatic, sometimes weakly oblong to oblong, margins 4–6-sided, sometimes scalloped, (0.6)0.8–1.1 mm long, 1–1.8 mm diam. at apex; thecae oblong, 0.3–0.5 mm wide,  $\pm$  parallel to one another, contiguous or nearly contiguous; pollen ellipsoidal,  $<0.1$  mm long,  $<0.1$  mm diam.; sterile staminate flowers irregularly 4–6-sided, clavate or weakly rounded, blunt, rarely irregularly 3–5-sided or depressed with a furrow, (0.8–1.2)1.7 mm long, 0.9–1.5(1.5–2 mm) wide.

Flowering in *Philodendron scalarinerve* occurs principally in the early rainy season (June through August) but also in March. Post-anthesis inflorescences have been collected primarily from June through September, but one such Costa Rican collection was made in January and one South American collection in December. Immature fruiting collections are known from July through December.

*Philodendron scalarinerve* ranges from Costa Rica to Ecuador, from sea level to 1325 m in *Tropical wet forest*, *Premontane rain forest*, *Tropical Lower Montane rain forest*, and *Tropical rain forest* (Colombia).

*Philodendron scalarinerve* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is distinguished by its short internodes, bluntly 1-ribbed to unribbed cataphylls persisting as pale fibers with small fragments of thin, pale epidermis, terete to subterete petioles, narrowly ovate to oblong-ovate blades drying dark with the "cross-veins" (tertiary veins that extend between the minor veins) numerous and conspicuous on both surfaces and uniformly greenish spathes.

*Additional specimens examined.* COSTA RICA. He-

**redia:** Finca El Bejuco, N of Río Sarapiquí, Chilamate de Sarapiquí, ca. 100 m, 10°27'N, 84°04'W, Grayum & Ray 5564 (CR, MO); Cerros Sardinal, ca. 2–2.5 km N of Chilamate de Sarapiquí, 80–160 m, 10°28'N, 84°04'W, Grayum et al. 6147 (CR, MO). **Limon:** Cerro Coronel, S of Río Colorado, 10–80 m, 10°40'30"N, 83°39'30"W, Durside & Herrera 31475 (CR, MO). PANAMA. **Bocas del Toro:** Fortuna Dam area, Chiriquí Grande-Fortuna, 4.3 km N of the Continental Divide, 590 m, 8°46'N, 82°14'W, Croat & Grayum 60173 (CM, MO); 3.2 mi. N of Divide, 700 m, 8°45'N, 82°15'W, Croat & Grayum 60251 (MO, NY, PMA, QCA). **Chiriquí:** Fortuna Lake area, Río Chiriquí Valley, vic. IRHE facilities, 1100–1200 m, 8°45'N, 82°18'W, Croat 66593 (AAU, CAS, CM, COL, DUKE, G, GH, HUA, K, L, MEXU, MO, NY, PMA, QCA, TEX, US, VDB, VEN); Lago Fortuna, along trail to Río Hornito, 8°45'N, 82°18'W, Croat & Zhu 76419 (MEXU, MO, SCZ, US). **Coelé:** El Copé region, 5–6 mi. N of El Copé, 600–800 m, 8°38'N, 8°35'W, Croat & Zhu 77222 (HUA, MO, MY, QCNE); ca. 0.5 mi. N of Continental Divide at Alto Calvario, ca. 5.5 mi. N of El Copé, 800 m, 8°39'N, 80°36'W, Croat 75081 (MO, US); Alto Calvario, ca. 6 km N of El Copé, 710–800 m, 8°39'N, 80°36'W, Croat 68723 (MO, PMA, RSA); 720–800 m, 8°38'N, 80°35'W, McPherson 12856 (BR, CAS, CR, MO, PMA); 930 m, Croat 49196 (CAS, L, MO); 49174 (MO), 49154 (MO); 68722 (DUKE, F, MO, PMA); 750–900 m, 44737 (CM, MO); 44720 (MO, PMA); El Valle region, La Mesa, N of El Valle de Antón, 800–900 m, 8°38'N, 80°09'W, Croat 67203 (CR, MO, PMA, QCA, US); 25436 (F, MO); 880 m, 37569 (F, MO); ca. 1000 m, Kennedy & Dressler 1759 (MO); Lutney & Kennedy 1717 (DUKE); Cerro Gaital, 800–900 m, 8°37'N, 80°07'W, McPherson 11196 (MO, PMA). **Colón:** Santa Rita Ridge Road, ca. 22 km from Highway, 500 m, 9°25'N, 79°40'W, Hammel et al. 14478 (MO). **Panamá:** Vertiente Pacífica, 150–200 m, 9°13.5'N, 78°15'W, Herrera 1604 (K, MO, PMA, STRI, US); Cerro Jefe region, 26.3 km from highway, ca. 600 m, Croat 35919 (MO); Campo Tres, 3 mi. NE of Altos de Pacora, 500–800 m, Croat 22701A (CM, MO, NY); 700 m, Croat 27072 (F, MO), 27093 (MO); ca. 1 mi. beyond Alto de Pacora, 2600 ft., Hammel & Kress 13420 (DUKE); El Llano-Carrt, 5–6 mi. N of Inter-American Highway at El Llano, 350–375 m, Croat 34805A (F, MO, NY, QCA, WIS); Mile 6.8, 350 m, Croat 49112 (MO); Mile 7, 9°19'N, 79°59'W, Croat 75103 (CM, MO). **San Blas:** Nusagandi, 300 m, 9°15'N, 79°W, McPherson 11066 (MO, PMA); Río Urgandi-Cerro Óbu, on trail, 100–300 m, 9°23'N, 78°48'W, de Nevers et al. 8019 (MO, PMA); Río Diabolo, 40 m, 9°23'N, 78°34'W, Herrera et al. 1714 (MEXU, MO, PMA, STRI, US). **Veraguas:** Santa Fe region, slopes of Cerro Tute, NW of Santa Fe, 1250–1350 m, Croat 48971 (F, MO, W); Santa Fe-Río San Luis, 8 mi. N of school, 450 m, 8°33'N, 81°08'W, Croat 66959 (CM, ENCB, MBM, MO, PMA, QCA); Alto Piedra-Calovebona, Río Dos Bocas Valley, 350–400 m, Croat 27367 (F, MO, PMA, US).

**COLOMBIA.** **Chocó:** Medellín-Quibdó, 78 km W of Bolívar, 466 m, Croat 49286 (MO); Km 175–176, 117–118 km E of Quibdó, 465 m, 5°44'N, 76°28'W, Croat 57485 (CHOCO, COL, JAUM, K, MO, US); Quibdó-Istmina, Km 14,  $<100$  m, 5°32'N, 67°37'W, Croat & Cogollo 52215 (MO); S of Río Rancherita, Km 31–32,  $<50$  m, 5°27'N, 76°39'W, Croat 57372 (CHOCO, COL, JAUM, MO); Quibdó-Lloro, vic. Río Atrato, ca. 150 m, 5°29'N, 76°35'W, Croat 55979 (COL, JAUM, MO, NY); Río Capá-Río Mumbá, upriver from Lloro, 80–120 m, 5°37'N, 76°25'W, Jancosa 1457 (MO); Quibdó-Yuto, 12 km S of

Quibdó, 60 m, 5°38'N, 76°40'W, *Croat 56261A* (CM, MO); Serranía de Baudó, Las Animas-Pato (Río Pato), 4 km S of Pato, 150 m, 5°30'N, 76°46'W, *Croat 56151* (MO); Jequédo, 41 km W of Las Animas, ca. 10 km E of Río Pato, 220 m, *Gentry & Rentería 24112* (MO). **Nariño:** Mpio. Barbacoas, Reserva Natural Río Nambí, 1325 m, 1°18'N, 78°08'W, *Betancur et al. 4778* (MO). **Valle:** Bajo Calima region, Buenaventura-Río Calima, 6.5 km beyond Portón Tomar, 50 m, 4°02'N, 77°07'W, *Croat 61278* (AAU, B, CAS, CM, COL, CR, F, K, L, MO, NY, QCA, US); 61380 (CM, MO); road to Málaga, 6 km S of main road, 50-80 m, 3°56'N, 77°07'30"W, *Croat 69417* (AAU, COL, CR, GB, MEXU, MO, NY, QCA); 100 m, 3°55'N, 77°W, *Monsalve 892* (MO); Buenaventura-Málaga, Km 65-66, 40-65 m, 4°10'N, 77°12'W, *Croat 71055* (MO); at Km 17.5, 3°57'N, 77°01'W, *Croat & Bay 75630* (MO); Km 28, W of Buenaventura-Málaga road, 3°59'N, 77°03'W, *Bay 269* (MO); Km 35.2, 100 m, 4°N, 77°03'W, *Croat & Bay 75759* (MO); Km 44, <100 m, 4°03'N, 77°08'W, *Croat & Watt 70199* (COL, MO, NY, US); Km 52.4, 140 m, 4°03'N, 77°05'W, *Croat & Bay 75724* (MO); 75728 (MO); 11 km NW of Cali-Buenaventura Highway, 3°56'30"N, 77°01'W, *Croat 69321* (CM, MO). **ECUADOR. Carchi:** Chical, 1200-1250 m, 0°56'N, 78°11'W, *Thompson & Rawlins 761* (CM, MO).

***Philodendron schottianum*** H. Wendl. ex Schott, *Oesterr. Bot. Z.* 15: 72. 1865. TYPE: Costa Rica. *Wendland s.n.* (holotype, W? lost). Schott ic. 2735-36 (neotype, here designated, W). Figures 2, 17, 365-371.

Usually epiphytic or hemiepiphytic; stem appressed-climbing, grayish green, sap strongly thyme-scented; internodes short, thick, semiglossy, 4-7.5 cm diam., broader than long, pale green to gray, epidermis thin, yellow, fragmented, without fissures; roots moderately few per node, drying dark brown to ca. 5 mm diam., epidermis semiglossy, flaking; cataphylls 16-46 cm long, sharply 2-ribbed (ribs to 1.5 cm high), usually tinged red, drying brownish yellow, often glossy (as if surface is shellacked), broadly concave to broadly D-shaped adaxially, persisting semi-intact, finally as a dense mass of whitish fibers; margins acute; **petioles** 35-83 cm long, 1-2.2 cm diam., subterete to D-shaped, firm to moderately spongy, medium green, drying yellowish brown, obtusely flattened with obtuse medial rib toward apex adaxially, surface semiglossy and obtusely striate; **blades** ovate, subcoriaceous, semiglossy, moderately bicolorous, acuminate to abruptly acuminate at apex (the acuminate strongly inrolled, 2-8 mm long), cordate to sagittate at base, 30-77 cm long, (17.5)23-64 cm wide (1-1.7(2.7) times longer than wide), (0.6-1.2 times longer than petiole); upper surface dark green, semiglossy to subvelvety-matte; lower surface semiglossy or rarely matte, moderately paler; anterior lobe 23-57 cm long, 24.5-64 cm wide (1.6-2.7 times longer than posterior lobes); poste-

rior lobes 8.5-28 cm long, 10-29.6 cm wide, obtuse to broadly obtuse; sinus usually spatulate, 8-20 cm deep; midrib flat to broadly convex, paler than surface above, convex to narrowly rounded, concolorous or slightly darker than surface below; basal veins (1)6-7(8-9) per side, with 0-1(2) free to base, most of the remainder coalesced 1-5.5 cm, 2 coalesced to 11 cm; posterior rib usually naked, 1-3 cm long; primary lateral veins 3-8 per side, departing midrib at a usually 40-70° angle, quilted-sunken to sunken, paler than surface above, convex and slightly paler than surface below; tertiary veins visible and darker than surface below; minor veins conspicuous, arising from both the midrib and primary lateral veins, moderately prominent on drying, alternating with secretory ducts perpendicular or more frequently oblique, sometimes branching; "cross-veins" weakly parted. **INFLORESCENCES** erect to erect-spreading, (1)2-3 per axil; peduncle 2-9 cm long, 8-14 cm diam., pinkish red, white striate, especially toward apex; **spathe** 10-17 cm long (1.9-7 times longer than peduncle), weakly constricted, oblong-ellipsoid; spathe blade light green outside, cream, pale lineate in upper one-half inside; spathe tube green, tinged red outside, 6-9 cm long, red with conspicuous resin canals inside; **spadix** tapered to somewhat ovate, weakly protruding, 8.8-14.4 cm long; pistillate portion weakly ovoid, whitish, (2)3-4.9 cm long, 2-2.4 cm diam. throughout, 1.4-1.5 cm diam. at apex, 1.4-1.9 cm diam. at middle, 1.4-1.8 cm wide at base; staminate portion 5.7-10.8 cm long; fertile staminate portion bluntly tapered at apex, 1.1-1.4 cm diam. at base, 1.2-2.3 cm diam. at middle, 1(1.7) cm diam. ca. 1 cm from apex, broadest at upper two-thirds, broader than the pistillate portion, slightly narrower than the sterile portion; sterile staminate portion broader than the pistillate portion, 1.4-2.5 cm diam.; pistils (1)3-3.9 mm long, 1.5-1.9(3.8) mm diam., margins broadly rounded and slightly raised above the apex; ovary 4-6-locular, 1.9-2.5 mm long, 1.3-1.9(3.8) mm diam., with axile placentation; locules 1.9-2.5 mm long, 0.4-0.7 mm diam.; ovule sac not present or to 1.9 mm long; ovules 10-14(18) per locule, 2-seriate, rarely contained within translucent or transparent envelope, 0.1-0.3 mm long, longer than funicle; funicle 0.1-0.3 mm long, adnate to lower part of partition, style 0.8-1.2 mm long, 1.5-1.9(3.6) mm diam., similar to style type B; style apex barely raised, button-like, broadly concave, medial apex with a whitened margin, raised and apparently like type D on drying; stigma subdiscoid, truncate, 1.4-1.6 mm diam., 0.3-0.4 mm high, covering ± entire style apex, sometimes de-

pressed shallowly and medially; the androecium truncate, oblong, prismatic, margins irregularly 4-5-sided, sometimes scalloped, 1 mm long, 1.8-2.1 mm diam. at apex; thecae  $\pm$  oblong, 0.4-0.5 mm wide,  $\pm$  parallel to one another; sterile staminate flowers blunt, irregularly 4-6-sided, prismatic, 2.5-3.5 mm long, (1.4)1.8-2.2 mm wide. INFRUCTION with pistillate spadix 5-6 cm long, 3.5 cm diam.; berries white, 1.1 cm long, 4-4.6 mm diam.; seeds 1.4 mm long, 0.5 mm diam., cream-colored.

Flowering specimens of *Philodendron schottianum* have been collected only from March and June, but post-anthesis collections have been made from March through August. Immature fruiting collections have been collected from January, May, September, and November. The immature January fruiting collection is a clear indication that the species must flower much earlier in the dry season than March (as indicated above). Perhaps it flowers throughout the dry season. In the cloud forest regions where this species occurs the dry season would not be very severe.

*Philodendron schottianum* ranges from Costa Rica to Panama at (490)730 to 2250 m in *Premontane rain forest* and *Tropical Lower Montane wet forest* life zones. In Panama, this species ranges no further east than Veraguas (Cerro Tute), except for a disjunct occurrence on Cerro Jefe in Panamá Province. All Costa Rican collections are from the northern slopes of the Cordillera Central in Alajuela, Heredia, and San José, and the northern end of the Cordillera de Talamanca in the Tapantí region of Cartago Province. It is to be expected throughout much of the Cordillera de Talamanca.

*Philodendron schottianum* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is distinguished by its short, thick internodes; sharply 2-ribbed cataphylls persisting as a dense mass of fibers (frequently with patches of glossy, yellowish to orange-brown epidermis); obtusely to sharply D-shaped petioles drying somewhat grayish or rarely yellowish and glossy; large, broadly ovate blades with a deep, usually spatulate sinus, usually pale-drying primary lateral veins; and rather conspicuous secretory canals between the veins; 2-3 short-pedunculate inflorescences per axil; and scarcely constricted, externally green spathes red on the tube within.

In Costa Rica and at higher elevations in Panama, such as on Cerro Colorado and on Cerro Pate de Macho (1000 to 2200 m), the petioles are subterete or obtusely flattened adaxially. At lower elevations in Bocas del Toro, Veraguas, and Coclé,

the petioles become D-shaped to sharply D-shaped with erect margins, and at the lowest elevations they are nearly always wing-margined.

In Bocas del Toro Province, Panama, at middle elevations and in mesic situations, this species is most easily confused with *P. findens*, which also has sharply D-shaped petioles. *Philodendron findens* also has spathes which, like those of *P. schottianum*, are barely constricted midway. In rare situations where the blades of *P. findens* do not promptly tear into narrow segments, *P. findens* can be distinguished from *P. schottianum* by having primary lateral veins of the lower surface drying darker than the surface.

In central Panama, *P. schottianum* can also be confused with *P. llanense*. Both *P. schottianum* and *P. llanense* occur in the Cerro Jefe region, though *P. schottianum* has been collected there only once, northeast of Altos de Pacora. This collection (Croat 6869I) exhibits most of the diagnostic features of *P. schottianum*, especially the persistent yellowish, semi-intact cataphylls, and the acutely D-shaped petioles with an obtuse medial rib (unknown in *P. llanense*), but has a blade shape midway between that of *P. schottianum* and *P. llanense* (blade length/width ratio 1.5 vs. an average of 1.4 for *P. llanense* and 1.65 for *P. schottianum*). *Philodendron llanense* differs in having at most obtusely flattened petioles and in lacking the conspicuous yellowish cataphylls of *P. schottianum*. *Philodendron llanense* also rarely occurs above 500 m (to 950 m), whereas *P. schottianum* only rarely occurs to as low as 500 m.

*Philodendron schottianum* can be confused with *P. thalassicum* and *P. alticola*, especially in Costa Rica. *Philodendron thalassicum* differs in having bluish green leaf blades that are glaucous beneath and sharply D-shaped petioles drying somewhat blackened rather than merely obtusely flattened and light yellow-brown, as in *P. schottianum*. *Philodendron alticola* differs in usually having narrower leaf blades (usually more than 1.8 times longer than wide) and stigma tubes exerted as minute funnels on the dried stigma.

*Philodendron dodsonii* may also be confused with *P. schottianum*. The former differs in occurring usually at lower elevations principally in tropical wet forest and premontane wet forest, and in having larger leaf blades, more long-pedunculate inflorescences, and longer spathes with a normal constriction above the spathe tube.

One collection, *Grayum* 7333, differs from more typical collections in having blades reportedly matte on both surfaces and lacking prominent minor veins and conspicuous secretory ducts (so evident in other material of this species where they



alternate with the minor veins). In addition, this specimen has darker-drying petioles and more fragmentary old cataphylls as well as intact cataphylls (perhaps juvenile?) drying brown (rather than the typical brownish yellow) and a spathe with a narrowly acuminate portion extending a full 5 cm beyond the end of the spadix. On other specimens, the spadix ends only about 1 cm or less short of the end of the spathe.

*Hammel et al. 14705* is unusual in lacking cataphylls, suggesting that they might have been deciduous (or forcibly removed during preparation).

**Additional specimens examined.** COSTA RICA. **Alajuela:** 3.5–4 mi. W of center of San Ramón, ca. 800 m, *Croat 46813* (MO); Atlantic side of Alto Palomo, 1900 m, *Leur 1845* (CR, F); Río Gorrón–Río Toro, 1700 m, 10°12'N, 84°19'W, 2996 (CR, F); Río Cariblanco Canyon, Río Cariblanco–Quebrada Quicual, SW of Cariblanco, 840–950 m, 10°16'N, 84°12'W, *Grayum et al. 6186* (CR, MO, NY); Río Sarapiquí, road to Colonia Virgen del Socorro, 830 m, 10°16'N, 84°11'W, *Croat 68341* (CR, MO). **Cartago:** 31 km from San José, SE of CR-2, 1750 m, *Harrison & Fuentes 6081* (UMO); Tapanti Watershed Preserve, ca. 20 mi. SW of Paraiso, 1500–1800 m, 9°43'N, 83°47'W, *Croat 47047* (MO); 1500–1800 m, *Croat & Grayum 68293* (B, K, MO); 1500–1700 m, *Croat 36110* (MO); 1250–1350 m, 79018 (CR, INB, MO); *Grayum & French 5820* (MO); *Nilsen et al. 383* (CR); along Camino Raíz de Hule, SE of Platanillo, 1200–1400 m, *Croat 36824* (MO). **Heredia:** 3 mi. S of Cariblanco, 760 m, *Croat 35778* (MO); 4 mi. N of V. Bara Blanca, 1350 m, *Croat 35620* (F, MO, PMA); Río La Paz Grande, 7.5 km N of Vara Blanca, 1270–1350 m, *Croat 36017* (MO); Río Santo Domingo, ca. 3 km E of San Rafael de Vara Blanca, Volcán Barva, 2060–2100 m, 10°11'N, 84°07'W, *Grayum 7333* (CR, MO, US); Volcán Barva, Finca Montreal, Río Volcán–Río San Fernando, 1740 m, 10°12'39"N, 84°06'45"W, *Boyle et al. 1155* (MO, NY); Volcán Barva, along Río Vueltas, 1900 m, 10°06'N, 84°04'W, *Burger & Gentry 9042* (F, US); Parque Nacional Braulio Carrillo, 1000 m, 10°16'38"N, 84°04'57"W, *Boyle 964* (CM, MO, TEX); 1223 (CR, MO); 1990 m, 10°11'03"N, 84°06'27"W, *J058* (CR, MO); 1750 m, 10°03'40"N, 84°01'W, 2432 (CR, MO, NY). **Puntarenas:** Monteverde Reserve, 1500 m, ca. 10°17'N, 84°48'W, *Croat 61194* (CM, MO, PMA); Coto Brus, Cordillera de Talamanca, 730 m, 9°01'30"N, 82°57'40"W, *Mora 14* (INB). **San José:** 4 km N of Cascajal, 7 km N of Las Nubes on CR 216, 1500–1600 m, *Uiley & Uiley 5259* (DUKE); Cerro de la Muerte, ca. 2000 m, *Croat 32859* (MO); Braulio Carrillo National Park, 1000–1500 m, 10°05'N, 83°57'W, *Croat 61226* (MO). PANAMA. **Bocas del Toro:** Cerro Colorado, 9.2 mi. W of Chame, 1450–1480 m, 8°35'N, 81°50'W, *Croat 69066* (CM, CR, MO, PMA); Fortuna Dam area, Chiriquí Grande–Fortuna, 1.2 mi N of Continental Divide, 910 m, 8°44'N, 82°17'W, *Croat 60447* (CR, MO, NY, PMA, TEX); 0.3 km N of Divide, 970 m, 8°43'N, 82°17'W, *Croat & Zhu 76518* (CR, DUKE, MO, WIS). **Chiriquí:** Cerro Colorado, above San Félix, 18–27 mi. off Pan-American Highway, 1200–1500 m, *Croat 33153* (MO); 20 mi. N of Río San Félix, 2000 m, *Croat 48460* (MO, TEX); 18.5 mi. N of Río San Félix, 1660 m, 8°30'N, 81°46'W, *Croat 74991* (AAU, CAS, CM, COL, CR, G, K, MEXU, MO,

QCA, US); Cerro Hornito, above Los Planes de Hornito, 1750–1900 m, 8°41'N, 82°10'W, *Croat 67979* (F, MEXU, MO, PMA); Cerro Pate Macho, NE of Boquete, 1630–1780 m, 8°46'N, 82°25'W, *Croat 66395* (F, MO); 1800–2200 m, 48560 (MO); 1900–2000 m, 66505 (B, CM, COL, CR, DUKE, F, L, MEXU, MO, QCA, TEX, US, VDB, VEN); ca. 2200 m, *Croat 48545* (MO); 1650–2000 m, 8°50'N, 82°25'W, *McPherson 11315* (MO, PMA); Cerro Punta, above town, 2250 m, *Croat 48601* (MO); Fortuna Dam area, Gualaca–Chiriquí Grande, Río Chiriquí, 9.1 mi. beyond Los Planes de Hornito, 1300 m, *Croat 48758* (MO); Quebrada Arena, 1050 m, 8°45'N, 82°16'W, *Hammel et al. 14705* (MO). **Coeló:** El Valle region, La Mesa, N of El Valle de Antón, E edge of Cerro Gaital, 900–1000 m, 8°37'N, 80°08'W, *Croat 67234* (MO, PMA); 900–930 m, 37433 (F, MO); El Copé area, on western slope, just S of the old saw-works area, 700 m, 8°38'N, 80°35'W, *Croat & Zhu 76795* (MO). **Panamá:** Cerro Jefe region, 3–3.5 mi. NE of Altos de Pacora, 7.8–8.2 mi. above Pan-American Highway, 700–750 m, 9°15'N, 79°25'W, *Croat 68691* (AAU, CAS, K, MO, NY). **Veraguas:** Santa Fe region, Santa Fe–Calovébora, beyond Escuela Agrícola Alto Piedra, beyond Río Tercero Brazo, 600 m, 8°31'N, 81°08'W, *Croat 66930* (CM, MEXU, MO, NY, PMA); Río Primero Brazo, 490 m, 8°33'N, 81°08'W, 66879 (CM, CR, HUA, IBE, JBGP, MO, PMA, QCA, US); ca. 3 mi. N of the school, ca. 700 m, *Croat 49002* (MO); 1.7 mi. past the school, 570 m, 8°33'N, 81°08'W, *Croat & Zhu 76832* (MO); slopes of Cerro Tute, near Escuela Agrícola Alto Piedra, 800–1030 m, *Croat & Zhu 76872* (CAS, MO); 1050–1150 m, *Croat 48896* (CM, MO); 0.6 mi. beyond school, 730 m, *Croat & Folsom 33954* (F, MO, PMA); 34041 (MO); Cerro Tute, 1250–1350 m, *Croat 48958* (CAS, MO); 800–950 m, 8°33'N, 81°08'W, *Croat 67002* (CAS, CR, L, MO, NY, PMA).

**Philodendron smithii** Engl., Bot. Jahrb. Syst. 26: 540. 1899. TYPE: Guatemala. Izabal: Río Dulce at Livingston, sea level, 5°49'N, 88°45'W, *J. D. Smith 1535* (lectotype, here designated, US; isolectotype, K). Figures 364, 372, 373, 377, 378.

Usually hemiepiphytic, sometimes terrestrial or epilithic; stem usually appressed-climbing, leaf scars conspicuous, 7–15 mm long, 7–16 mm wide; internodes glossy, (1)2.2–3 cm long, 2–3 cm diam., usually longer than broad or about as long as broad, tan or gray-green to dark green, epidermis drying yellow-brown with loose flakes, fissured conspicuously; cataphylls soft, spongy, 12–20 cm long, bluntly to sharply 2-ribbed, green, dark green short-lineate, rounded at apex, drying tannish to yellowish, deciduous, intact; petioles 21.5–63 cm long, 4–16 mm diam., subterete, spongy, medium green, obtusely flattened with obtuse angle toward apex adaxially, surface dark short-lineate, dark with a dark purple-black ring around apex; blades narrowly ovate, subcoriaceous, glossy, moderately bicolorous, acuminate to long-acuminate at apex (the acumen inrolled, 1–4 mm long), cordate at base, 26.5–53 cm long, 11–31 cm wide (1.2–2.2 times

longer than wide), (0.7–1.4 times longer than petiole), margins hyaline, convex in lower half of blade, upper surface dark green, drying dark gray-green to yellow-brown, lower surface light greenish, usually drying yellow-green to yellow-brown, sometimes olive-green; anterior lobe 18–41 cm long, 18–30 cm wide, (1.8–2.9 times longer than posterior lobes); posterior lobes (3.2)7.5–16 cm long, 8.7–14.2 cm wide, obtuse to broadly rounded; midrib flat to broadly convex, paler than surface above, prominently raised, concolorous below; basal veins (3)4–5(6) per side, with 0–1 free to base, 1–2 coalesced (1.8)2.5–5.5(7.5) cm; posterior rib not at all naked or briefly so, usually 1–1.5 cm (rarely to 2 cm); primary lateral veins 3–6 per side, darker than surface below, departing midrib at a 50–70° angle, ± straight, eventually weakly arcuate to the margins, sunken above, convex below; interprimary veins weakly raised, darker than surface below; minor veins numerous, sparsely visible, darker than surface below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, 2 per axil; peduncle 9–22 cm long, (1)9–12 mm diam., medium green, moderately spongy; **spathe** ± erect, 8.8–19.8 cm long, (0.5–1.4 times longer than peduncle), prominently constricted at anthesis; spathe blade yellowish green to green, glossy outside, 7 cm long, 1.7 cm diam., whitish green, minutely white-dotted, sometimes orange-striate inside; spathe tube ellipsoid, green outside, 8.5 cm long, 2.5 cm diam., light red to maroon (B & K red-purple 2/10) inside; **spadix** stipitate 6–15 mm long, gradually to bluntly tapered to apex, 9.3–16.9 cm long; pistillate portion pale green to greenish yellow, 3.3–5.2 cm long in front, 2.2–2.5 cm long in back, 1.4 cm diam. at apex, 1.3–1.6 cm diam. at middle, 1.6 cm wide at base; staminate portion 6.5–11.3 cm long; fertile staminate portion creamy white, 9–12 mm diam. at base, 1–1.2 cm diam. at middle, 5–7 mm diam. ca. 1 cm from apex, broadest at the middle, narrower than the pistillate and sterile portions; sterile staminate portion broader than the pistillate portion, creamy white, (9)1.1–1.5 cm diam.; pistils 1.8–2.3 mm long, 1.1 mm diam., creamy white; ovary 6–8-locular, 0.9–1.2 mm long, 1 mm diam., usually with basal, sometimes sub-basal placentation; locules 0.9–1.1 mm long, 0.3–0.4 mm diam.; ovule sac 0.9(1.2) mm long; ovules 1 per locule, contained within transparent ovule sac, 0.3–0.4 mm long, about as long as funicle; funicle 0.3–0.4 mm long (can be pulled free to base), style 0.6–0.8 mm long, 1.1 mm diam., similar to style type B; style apex ± sloping; stigma subdiscoid, greenish yellow, 1 mm diam., 0.3 mm high, covering entire style apex;

the androecium irregularly 5–6-sided, sometimes oblong or quadrangular, 1.6–1.9 mm long, 0.9–1.2 mm diam. at apex; thecae oblong, 0.4–0.5 mm wide, ± parallel to one another and nearly contiguous; sterile staminate flowers blunt, irregularly 4–6-sided, sometimes prismatic, (1)1.6–2 mm long, (1.2)1.7–1.8 mm wide. Berries white or sometimes yellowish; seeds 1 per locule, tan, oblong, 2 mm long, 0.8 mm diam.

Flowering in *Philodendron smithii* occurs principally in the dry season, and flowers have been collected from December to April (but also July). Post-anthesis collections are common from February through July but with a few in October, November, and December. The latter, especially those in October and November, are the most aberrant, since even if this species is cued to flower by the onset of the dry season (which appears to be the case), they would appear to have flowered too early. This may be explained by the fact that the dry season starts much earlier in Mexico and Guatemala than in Costa Rica and Panama.

*Philodendron smithii* ranges from Mexico (mostly in Chiapas, Tabasco, and Quintana Roo) to Guatemala, Honduras, and Nicaragua on the Atlantic slope at 40 to 1630 (mostly less than 600) m elevation. It is known from "Selva Alta Perennifolia" and "Selva Mediana Subperennifolia" in Mexico as well as from *Tropical moist forest* and *Subtropical wet forest* in Guatemala.

*Philodendron smithii* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Ovata*. This species is characterized by its conspicuously fissured tan internodes, which are about as long as to longer than broad; sharply two-ribbed, deciduous cataphylls; spongy subterete petioles; usually blackish, narrowly ovate-cordate leaf blades; long-pedunculate inflorescences; and externally green spathes that are maroon inside the tube.

*Philodendron smithii* is not easily confused with any other species in Central America. It is most similar to *P. panamense*, which is endemic to Panama, and shares with that species similarly dark-drying ovate-triangular, long-petiole blades and long-pedunculate inflorescences. *Philodendron panamense* differs in having much shorter internodes (shorter than broad on flowering plants), weakly one-ribbed cataphylls, moderately firm petioles, and spathes green on the tube outside with a white blade and greenish white throughout within (vs. green throughout outside and maroon on the tube within for *P. smithii*).

*Philodendron smithii* is also similar to *P. stramineicaule* and *P. wilburii*, both from Costa Rica and

Panama and both differing in having stems drying gray-green to grayish or stramineous rather than conspicuously yellow-brown as in *P. smithii*.

Mayo (1986) reported that *P. smithii* lacks superficial resin canals in the spathe and that instead resin production has entirely shifted to the spadix. However, we observed resin canals on the inner surface of both the spathe and the blade in this species.

A noteworthy collection is *Standley 52776* from the Lancetilla Valley in Honduras. It has leaf blades with broadly flaring lobes, but otherwise does not differ from other collections.

One outlying collection in Mexico, from the state of Querétaro at 450 m (*López 539*), is believed also to be this species.

Mayo (1989) reported style type E for *Croat 47913*, but my investigations of flowers of that collection showed no evidence of an annulus on the ovary, and while there was a slight depression in the style apex, the latter was not deemed sufficiently funnel-shaped to qualify for anything except a type B style.

The type specimens (*J. D. Smith 1535*), though unequivocal, were distributed by Engler with a printed label bearing the name *Philodendron donnell-smithii*, but no such name appears in Engler's publications.

*Additional specimens examined.* BELIZE. Camp 1, flood plain of the Caves Branch River, Whiteford 1263 (BM); Sibun River, Gracie Rock, *Gentry 1715* (MICH). Cayo: near Camp 6, *Gentry 2424* (CM, MICH). Stann Creek: Stann Creek Valley, 250–300 ft., *Schipp S-304* (F, GH, MO); Leoniel Valley, *Gentry 3513* (MICH, NY); Cockscomb Mountains, 2 km N of Victoria Peak, 300–500 ft., *Gentry 8033* (MO). Toledo: Richardson Creek, affluent of Bladen Branch, Maya Mountains, 100–250 m, 16°33'N, 88°47'W, *Davidse & Brant 31877* (BM, GH, MO); 80–420 m, 16°32'–33'N, 88°45'–46'W, *32052* (MO); Richardson Creek–Quebrada de Oro, canyon along Bladen Branch, 100–200 m, 16°31'–33'N, 88°46'–49'W, *Davidse & Brant 32230* (MO); Columbia Forest Station, 1.5 mi. from Maya village of San José, *Gentry 8132* (MO); *Croat 24360* (MO); *Dwyer 9891* (MO); Southern Mayo Mountains, Bladen Nature Reserve, 400–500 m, 16°30'22"N, 88°54'54"W, *Davidse & Meadows 35827* (BRH, CM, MO, SEL); Big Fall Estate, ca. 3 airline km NE of village of Big Fall, 40 m, 16°16'22"N, 88°52'27"W, *Davidse 35628* (BRH, MO). GUATEMALA. Alta Verapaz: Tzurucú–El Estor, 9 mi. up road to Oxec, 700 m, *Croat 41652* (MO); 800 m, *41694* (MO); 4 mi. up road to Oxec, 500 m, *41611* (MO); road to El Estor, 5 mi. W of Tzurucú, 12 mi. E of Hwy. CA-14 to Cobán, 600 m, *Croat 41512* (MO); Pantín, below Tamahú, ca. 600 m, *Standley 70512* (F); *70513* (F); along Quiché Highway, ca. 12 km W of San Cristóbal, ca. 1100 m, *Standley 89738* (F); near Tactic, ca. 1500 m, *Standley 92022* (F); vic. of Cobán, ca. 1300 m, *Standley 92676* (F); ca. 20 km W of Cobán at Cruz Max, 1500 m, *Castillo & Hodel 1100* (MO); vic. of Cubilguitz, 1.5–2 mi. S, 300–350 m, *Steyermark 44404* (F, MO); Río Carchá, Cobán–

San Pedro Carchá, ca. 1360 m, *Standley 90086* (F); Río Polochic, below Tamahú, ca. 975 m, *Standley 91990* (F). Baja Verapaz: Biotope del Quetzal, 1630 m, *Martínez et al. 23011* (MEXU). Izabal: Quirigua vicinity, 75–225 m, *Standley 23911* (US); ca. 4 mi. SW of Puerto Barrios, 50 m, *Croat 41823* (MO); ca. 7 mi. S of Puerto Barrios, 50 m, *Croat 41804* (MO); Lago Izabal vicinity, 0–600 m, 89°0'–25'W, 15°15'–35'N, *Jones et al. 3046* (F, NY); near Entre Ríos, ca. 18 m, *Standley 72746* (F); Bananea–La Presa, Montaña del Mico, 40–300 m, *Steyermark 38184* (F); Montaña del Mico, between Mills 49.5 and ridge 6 mi. from Izabal, 65–600 m, *Steyermark 38509* (F); *38613* (F). Petén: La Libertad, *Lundell 2994* (MICH, US); Uaxactun, *Bartlett 12426* (MICH, UC, US); El Paso, *Lundell 1572* (CM, MICH); 2929 (MICH); Santa Teresa, *Lundell 2888* (GH, MICH); Tikal National Park, 2 mi. S of entrance, 500 ft., *Croat 24700* (MO); Finca Yalpomech, Río San Diego, 50–150 m, *Steyermark 45315* (CM, F, NY, US). Zacapa: 41 mi. S of turnoff to Petén Morales, 150 m, *Croat 41874* (MO). HONDURAS. Potrero, along Highland Creek, *Wilson 484* (NY). Atlántida: San Alejo vicinity, near Río San Alejo, 150–270 m, *Standley 7699* (F); San José de Texigant, SE of Tela, *Nelson 10732* (TEFH); Lancetilla Reserve, ca. 2 mi. WSW of Tela and S of main hwy., 70–90 m, 15°44'N, 87°27'W, *Croat & Hanson 64588* (CM, F, K, MO, US); 100 m, *Molina & Molina 25650* (EAP, F, NY); 20–600 m, *Standley 52776* (F, US); 20–600 m, *Standley 52980* (F, US); 45 m, *Allen 6143* (EAP, F); 10–150 m, *Croat 42656* (MO), *42675* (MO); 19.2 km E of Tela on Tela–Ceiba Highway, near San Francisco de Soco, 50–100 m, 15°44'N, 87°21'W, *Brant & Zúñiga 2845* (EAP, MO, US); Campamento Quebrada Grande, ca. 10 km SW of La Ceiba, 80–180 m, 15°42'N, 86°51'W, *Liesner 261794* (MO). Cortés: Lago de Yojoa, 650 m, *Croat 42761* (MO); Agua Azul, *Allen 6448* (EAP, F). Lempira: Montaña de Celacán, between Gracias and top of Cerro Celacán, 14°33'N, 88°39'W, *Davidse & Zúñiga 34540* (MO). Olanchito: Río Wampusito, Dulce Nombre de Cuimí, 500–700 m, 15°15'N, 85°25'W, *Nelson & Clewell 439* (EAP, FSU, MO); San Esteban–Bonito Oriental, Río Grande, 3.3 mi. SW of border with Colón Dept., 350–400 m, 15°31'N, 85°42'W, *Croat 64524* (AAU, CAS, EAP, GB, K, MEXU, MO, NY, P). MEXICO. Chiapas: Mpio. La Trinitaria, Lagos de Montebello National Park, at Cinco Lagunas, 1600 m, *Breedlove 68620* (CAS); 10 km ENE of Dos Lagos, above Santa Elena, 1170 m, *Breedlove & Almada 57585* (MO); 1000 m, *Breedlove 56538* (CAS); Mpio. Las Margaritas, confluence of Río Ixcán and Río Lacantum, 300 m, *Breedlove & McClintock 34198* (MO); 12 km E of Taiscao, 1200–1300 m, *Davidse et al. 29886* (MO); Palenque–Boca Lacantum, Mpio. Ocosingo, Nuevo Guerrero, 340 m, *Martínez 18108* (MO); Ojo de Agua de San Javier, a 23 km al SE de Nuevo Guerrero, road to Boca Lacantum, 370 m, *Martínez 16929* (MO); 2 km N of Naja, trail to Chancala, 900 m, *Martínez 21320* (MEXU), *21328* (MEXU), *21331* (MEXU); San Cristóbal–Palenque, San Cristóbal de las Casas, 97 km NE of San Cristóbal, 1130 m, 17°27'N, 92°04'W, *Hammel et al. 15625* (K, MEXU); 70 km SW of Palenque, *Breedlove 47180* (CAS, MO); Laguna Ocotlito, 12 km N of Monte Libano, trail to Chancala, 950 m, *Martínez 17605* (MO); near lake at Naja, 300 m, *Breedlove 49951* (CAS); Mpio. Pueblo Nuevo Solistahuacán, 3 km NW of town, 6700 ft., 17°30'N, 92°40'W, *Will 412* (CAS, DH); San Fernando–Maravillas, 840–940 m, 16°53'N, 93°16'W, *Croat & Hanson 65003* (COL, F, K, MEXU, MO, NY, US); 20–30 mi. SW of El Jocote, road to Motozintla de Mendoza, 700–900 m, *Croat 40707* (B,

MEXU, MO); 1 mi. N of Escuintla, 100 m, *Croat 43802* (MO); 200 m, *Croat 40134* (MO); 160 m, *40295* (B, K, MEXU, MO); 210 m, *Croat 40317* (MO); 1000 m, *Croat 40588* (MO); 700 m, *Croat 40660* (B, MEXU, MO); 550 m, *Breedlove 56538* (CAS); ca. 400 m, *Croat 40190* (MO, NY); 460 m, *40271* (MO); 350–370 m, *40223* (MO); 0.6 mi. N of Tapulula, 35.6 mi. SSE of Pichucalco, 750 m, 17°16'N, 93°01'W, *Croat & Hannon 65285* (B, MO); 8 mi. N of Pichucalco, 80 m, *Croat 40079* (CM, L, MEXU, MO, PMA, US). **Oaxaca:** Uxpanapas region near Sarabia, 52 km E of main highway, 100 m, 17°10'N, 94°32'W, *Hammel & Merello 15579* (MEXU, MO); Chiltepec, 30 m, *Sousa 1732* (MEXU). **Queretaro:** Mpio. Jalpan, Río Las Pagnas, S of Tanchanaguito, 450 m, *López 539* (MO). **Quintana Roo:** 110 km SW of Chetumal, 100 m, *Davidse et al. 20172* (MEXU, MO). **Tabasco:** 3.1 mi. E of Teapa, 150 m, 17°33'N, 92°59'W, *Croat & Horsman 65364* (MO); ca. 9 km E of Teapa, cultivated, *Mayo 85* (K, MO); base of Cerro de Madrugal, ca. 4 km SE of Teapa, 40 m, *Croat 40133* (MO); ca. 300 m, *47913* (CM, G, MEXU, MO, US); Mpio. Macuspana, Centro Recreativo de Agua Blanca, 7 km from the Villahermosa-Escarecega Highway, *Cowan & Zamudio 3351* (CAS, MEXU, MO, NY); Mpio. Tacotalpa, 0.2 km NW of Tapajulapa, *Cowan et al. 3530* (CSAT). **NICARAGUA. Jinotega:** Jinotega-Matagalpa, ca. 5–8 mi. SW of Jinotega, 1500 m, *Croat 43059* (MO). **Río San Juan:** Río Santa Cruz—Caño Santa Cruzita, La Palma, 40–60 m, 11°2–4'N, 84°24–26'W, *Stevens 23432* (MO). **Zelaya:** Caño Monte Cristo, vic. Campamento Germán Pomares, ca. 10 m, 13°35'N, 83°51'W, *Morero 14841* (MO); Cerro La Calera, 4 km N of Siuna, 350 m, *Neill 4290* (BM, MO); Cerro Waylawás, ca. 100–268 m, 13°39'N, 84°48–49'W, *Stevens 7380* (MO); ca. 80 m, ca. 13°39'N, 84°49'W, *Pipoly 4406* (MO).

***Philodendron sousae*** Croat, sp. nov. TYPE: Mexico. Chiapas: Mpio. Ocosingo, Río Santo Domingo at Santo Domingo, 490 m, *Davidse et al. 20450* (holotype, MO-2946607; isotype, MEXU). Figures 379, 380.

Planta hemiepiphytica; internodia usque 3.5 cm longa, 8–15 mm diam.; cataphylla incostata, decidua; petiolus semiteres, 29–54 cm longus, 8–12 mm diam.; lamina ovato-cordata, 28–55 cm longa, 12.5–28 cm lata, in sicco brunnea vel olivacea supra, virella vel flavibrunnea infra; nervis lateralibus 1 3–4; inflorescentia usque 4; spathe 15–21 cm longa, lamina spatheae extus cremea, tubo spatheae albo, in superficiebus ambabus saturate suffuso rubro aut purpureo; pistilla 6–8-locularia; loculi 2-ovulati.

Hemiepiphytic; stem probably scandent; internodes to 3.5 cm long, 8–15 mm diam., yellow-brown to dark brown, epidermis drying smooth to wrinkled, sometimes fissured closely; roots drying reddish brown, thin, elongate, drying 3–4 mm diam., few per node; cataphylls 14 cm long, unribbed, deciduous; petioles 29–54 cm long, 8–12 mm diam., ± terete, drying yellow-brown; sheathing 3–6.5 cm; blades ovate-cordate, subcoriaceous, gradually acuminate at apex, conspicuously cordate at base, 28–55 cm long, 12.5–28 cm wide (2.2 times longer than wide), (about as long as petiole),

drying subcoriaceous, upper surface drying brown to olive-green, lower surface drying greenish to yellowish brown, smooth or minutely ridged; anterior lobe 23–36 cm long, 21–27 cm wide (2.8–3 times longer than posterior lobes); posterior lobes 8–12 cm long, 10–13 cm wide; sinus parabolic; midrib drying broadly convex and usually paler than surface above, drying narrowly convex and usually concolorous below; basal veins 3–5 per side, free to base, several coalesced to 1–2 cm, occasionally 2 fused to 3.5 cm; posterior rib usually not at all naked, sometimes obscurely so for about 1 cm; primary lateral veins 3–4 per side, departing midrib at a 60–70° angle, weakly arcuate to the margins, drying weakly raised and usually paler than surface above, drying raised below; minor veins arising from both the midrib and primary lateral veins. INFLORESCENCES to 4 per axil; peduncle 6.5–13 cm long, ca. 1 cm diam.; spathe 15–21 cm long (1.6–2.3 times longer than peduncle), with resin canals on inner surface, visibly constricted above the tube; spathe blade cream outside; spathe tube white, heavily tinged red or purple on both surfaces, spathe tube 5–11 cm long; spadix sessile; 6–14 cm long; pistillate portion cylindrical-tapered toward the apex, 2.4–4.1 cm long, 4–9 mm diam. at apex, 5–10 mm diam. at middle, 6–10 mm wide at base; staminate portion 3.6–10 cm long; fertile staminate portion (4/9 mm diam. at base, 6–12 mm diam. at middle, 3–7 mm diam. ca. 1 cm from apex, broadest at or below middle, broader than the pistillate portion; sterile staminate portion not obvious; pistils 0.5–2.7 mm long, (0.7)1.8–2.7 mm diam.; ovary 6–8-locular, 0.4–1.4 mm long, 0.7–2.7 mm diam., with basal (to sub-basal) placentation; locules 0.4–1.4 mm long, 0.9 mm diam.; ovule sac 0.3 mm long; ovules 2 per locule, contained within a translucent, gelatinous ovule sac, 0.1–0.2 mm long, much shorter than funicle; funicle 0.7–0.9 mm long, style 0.4 mm long, 0.4–1.5 mm diam., similar to style type D; stilar canals emerging at base of apical depressions and arranged separately in a ring on orifices around center; style apex flat; style boss moderately shallow to moderately prominent; stigma discoid, 0.4 mm diam., 0.1 mm high, covering entire style apex; thecae oblong to weakly ovate, 0.2–0.4 mm wide, divaricate.

Flowering in *Philodendron sousae* occurs in the early rainy season, and specimens have been collected in May and August. Fruits are not known.

*Philodendron sousae* is endemic to Mexico in Chiapas, at 490 to 1400 m elevation on the Atlantic slope in regions of "Selva Alta Perennifolia."

*Philodendron sousae* is a member of *P.* sect. *Co-*

*lostigma* subsect. *Macrobelyium* ser. *Macrobelyium*. This species is characterized by its ovate-cordate, light greenish-brown-drying blades with three to four primary lateral veins per side and three to five free to weakly coalesced basal veins, which are usually not at all naked. Also characteristic are the inflorescences, ranging up to four per axil and with the spathe more or less oblong and constricted somewhat above the tube. The spathe tube is white and heavily tinged with red or purple on both surfaces.

*Philodendron sousae* is most easily confused with some material of *P. advena* (from the Pacific slope), which dries a similar light brown to greenish brown (not dark brown typical of most material). The latter differs in having a more or less ellipsoid spathe, scarcely or not at all constricted above the tube.

*Philodendron sousae* may also be confused with *P. breedlovei*, which has somewhat similar blades that dry a similar color. The latter species has narrower blades, to 1.8 cm long, a single inflorescence per axil, and about 20 ovules per locule (vs. blades usually broadly ovate, averaging 1.4 times longer than wide, four inflorescences per axil, and 1-3 ovules per locule).

*Philodendron sousae* was first collected by D. E. Breedlove in 1972. It is named in honor of Mario Sousa (MEXU), co-editor of *Flora Mesoamericana* (Davidse et al., 1995), who (in the company of Gerit Davidse) collected the type specimen.

*Additional specimens examined.* MEXICO. Chiapas: Mpio. Cintalapa: 4 km W of La Ciénega, 38 km W of Las Cruces, Oaxaca-Chiapas border, 1400 m, *Breedlove 25142* (DS); Mpio. Ocosingo, vic. Centro de Población Velasco Suárez (Selva Lacandona), 570 m, 16°47'N, 91°17'W, *Calzada et al. 744* (MO); Laguna Ocotol Grande, ca. 25-30 km SE of Monte Líbano (ca. 45 km E of Ocosingo), 950 m, *Dressler 1589* (GH).

***Philodendron squamicaulis*** Croat & Grayum, sp. nov. TYPE: Panama. Coclé: vic. El Copé, on western slope, just S of the old saw-works area, 700 m, 8°38'N, 80°35'W, 12 July 1994, *Croat & Zhu 76798* (holotype, MO-4613255-8; isotypes, B, CAS, CM, COL, CR, F, G, K, MEXU, NY, PMA, US, VEN). Figures 374, 381-384.

Planta hemiepiphytica; internodia brevia, 1-8 cm longa, 2-3.5 cm diam., dense squamata; cataphylla 27-34 cm longa, incostata, rosea vel rubella aut rubribrunnea, persistentia semi-intacta; petioli teres vel subteres, 28-61 cm longus, 1 cm diam., dense squamatus, squamae purpurascens; lamina ovato-triangularis, 26-53 cm longa, 17.5-40 cm lata, sagittato-cordata basi; nervis lateralibus 1 dense puberulis infra; inflorescentia usque 5; pedunculus 3.5-8 cm longus, olivaceus, dense squamatus; spathe 15-17 cm longa; lamina spatheae extus viridialba, intus diluta viridi, suffusa rubra, dense squamata; tubo

spatheae omnino viridi, suffuso rubro; pistilla (3/4)(5)-locularia; loculi 20-28-ovulati.

Hemiepiphytic; appressed-climbing, stem 3-20 cm long; internodes short, densely scaly with  $\pm$  deltoid, sometimes bifurcated scales, mixed with fewer acicular scales, drying reddish brown, 1-8 cm long, 2-3.5 cm diam., longer than broad, semiglossy, dark green, soon gray-green, finally tannish to brownish, drying straw-colored to reddish brown to brown, completely hidden by old cataphylls; roots few per node, slender, drying brownish; cataphylls fleshy, 27-34 cm long, unribbed to bluntly 2-ribbed, yellow-green, pink to reddish, or reddish brown, densely long-scaly throughout with a sparse underlay of tuberculate structures, drying reddish brown, persisting as semi-intact pale fibers, sometimes with patches of epidermis, soon deciduous; margins folded in to form groove. LEAVES erect-spreading; petioles 28-61 cm long, ca. 1 cm diam., erect-spreading, terete to subterete, purplish to dark brown, surface densely scaly, 1.5-4 mm long, flattened, with a sparse underlay of tubercles; sheathing briefly only near or at the base; petiolar scales of two types, short purplish scales deltoid or broader than high, less than 0.2 mm high, these interspersed with much longer, spreading light green acicular scales, 3-5 mm long, both types of scales drying reddish brown; blades ovate-triangular, thinly coriaceous to subcoriaceous, conspicuously bicolorous, short acuminate to acute at apex, sagittate-cordate at base, 26-53 cm long, 17.5-40 cm wide (1-1.5 times longer than wide), (0.7-1.1 times longer than petiole), broadest at the petiole attachment, margins hyaline, upper surface medium to dark green, semiglossy to sometimes almost matte above, slightly paler, light to silvery-green, glossy to semiglossy below, drying reddish brown on both surfaces; sometimes with sparse whitish raphide cells; anterior lobe 21-28 cm long, 18-26 cm wide (2.2-2.8 times longer than posterior lobes), margins straight to weakly concave; posterior lobes 8-10.7 cm long, 7.1-11.5 cm wide, directed toward the base to somewhat outward, broadly rounded; sinus parabolic to hippocrepiform on younger leaves, mitered on older leaves, 3.5-17 cm deep; midrib flat to sunken or deeply sunken and concolorous above, convex to broadly round-raised, concolorous to paler to light reddish below; basal veins 5-12 per side, with 0(1) free to base, third and higher order veins coalesced 2-6 cm long; posterior rib naked for 3-4.5 cm, densely scaly on both upper and lower edges; primary lateral veins 5-9 per side, departing midrib at a 30-45° angle (55° angle near base), moderately straight to the margins

(the lowermost veins somewhat branched near the margins), deeply sunken and concolorous above, brown to reddish, round-raised and densely puberulent below; tertiary veins visible, slightly raised, darker than surface; minor veins moderately distinct below, arising from both the midrib and primary lateral veins; "cross-veins" sometimes moderately distinct on lower surfaces on drying. INFLORESCENCES to 5 per axil; peduncle 3.5–8 cm long, olive-green, densely scaly, especially near apex, peduncle and spathe with scales of 2 lengths, both long and acicular, 1–2 mm long, much shorter and broader deltoid to tuberculate; **spathe** 15–17 cm long, 2–4.8 times longer than peduncle, moderately constricted above the tube, green, densely to sparsely scaly outside, shortly acuminate at apex; spathe blade sometimes greenish white outside, to 5.5 cm diam., light green, tinged red near base inside; spathe tube  $\pm$  ellipsoid, green, tinged red, covered with greenish scales (except for 5 mm along margin to 3 cm along infolded edge near base) outside, ca. 6 cm long, to 3.5 cm diam., reddish violet inside except greenish near tip inside; **spadix** 13.2 cm long; pistillate portion pale green to dirty-white, to 4.2 cm long in front, 3.6 cm long in back, 12 mm diam. at apex, 14 mm diam. at middle; staminate portion 11.5 cm long; fertile staminate portion 1.2 cm diam. at base, 1 cm diam. at middle, 7 mm diam. ca. 1 cm from apex; sterile staminate portion 1.2 cm diam.; pistils 1.22 mm diam.; ovary (3/4)5-locular, with axile placentation; locules 1–1.2 mm long; ovules 20–28 per locule, 2-seriate, ca. 0.2 mm long, longer than funicle; style similar to style type D; style apex flat; style boss narrow, fairly shallow; stigma hemispheroid, 0.7–0.8 mm diam., 0.2 mm high; the androecium prismatic, margins irregularly 4–6-sided, 1.4–1.6 mm long, 1.4–1.6 mm diam.; sterile staminate flowers prismatic, margins irregularly 4–6-sided, 1.1–1.2 mm diam. INFRUCTESCENCE with peduncle to 11 cm long, 8–11 mm diam.; spathe to 23 cm long, 1.7 cm diam., dark punctate in part and drying light brown inside; spathe tube 9–9.5 cm long, reddish, spathe blade to 14 cm long; spadix to 21 cm long; pistillate spadix 6–8.5 cm long, 3.2 cm wide; pistils 4.2 mm long, 2.7 mm diam.; locules to 3.3 mm long; fertile staminate spadix 11.5 cm long, 1.1 cm diam. midway; sterile staminate spadix 2.5 cm long, 1.4 cm diam.; staminate flowers to 3.3 mm long; berries white; seeds to 18 per locule, 1 mm long. JUVENILE plants terrestrial, scandent; lower surface of blades purplish violet. PRE-ADULT plants with internodes 2–30 cm long, 0.5–1.2 cm diam.; petioles 8.5–18.5 cm long, 2–3 mm diam.; blades 14–28 cm long, 15–22 cm wide.

Flowering in *Philodendron squamicale* occurs in the early rainy season from May through August. Mature fruits have been collected in early December.

*Philodendron squamicale* ranges from Costa Rica to Esmeraldas Province, Ecuador, from <100 to 1250 m elevation in pluvial forest, *Premontane wet forest*, and *Premontane rain forest* life zones. In Central America it has not been collected below 550 m elevation.

*Philodendron squamicale* is a member of *P.* sect. *Philodendron* subsect. *Achyropodium*. This species is recognized by its appressed-climbing habit, moderately thin, triangular-ovate, semiglossy blades, and especially by its scaly stems, cataphylls, petioles, peduncles, and outer surfaces of spathes and densely puberulent major veins on the lower blade surface.

*Philodendron squamicale* is apparently closest to *P. serpens* Hook. f., which was described from a cultivated plant from an unknown Colombian locality. The latter species differs in having oblong-ovate blades, sometimes with a constriction above the posterior lobes, with 4–5 basal veins on the most well developed blades (in contrast to 6–12 basal veins). In addition, *P. squamicale* has a coarse row of scales extending along both edges of the posterior rib and densely puberulent primary lateral veins, neither of which was described for *P. serpens*. *Philodendron serpens* also differs in having glabrous, rather than scaly, peduncles and spathes.

*Philodendron squamicale* is probably related to *P. verrucosum*, which also has densely scaly petioles. That species differs, however, in having longer petioles (more than 30 cm long vs. less than 30 cm for *P. squamicale*), and broadly ovate blades that are velvety and matte above and matte with frequently purplish areas along the veins below (vs. usually solid green below in *P. squamicale*). Both species have a similar number of primary lateral veins, but the veins are much more closely spaced in *P. squamicale* than *P. verrucosum*.

*Additional specimens examined.* COSTA RICA. **Heredia:** N of Quebrada Tigre, from Finca El Plástico (ca. 8 km SW of Las Horquetas), 450–550 m, 10°18'N, 84°02'W, Grayum & Sleeper 6523 (CR, MO, US); Zona Protectora, between Río Peje and Río Güacimo, Volcán Barba, 750 m, Grayum & Schatz 3250 (DUKE); Sarapiquí, P. N. Braulio Carrillo, El Ceibo, 750 m, 10°17'50"N, 84°04'25"W, Boyle et al. 2897 (CR, MO). PANAMA. **Bocas del Toro:** Fortuna Dam area, Chiriquí Grande-Fortuna, 4.3 km N of the Continental Divide, 590 m, 8°46'N, 82°14'W, Croat & Grayum 60197 (K, MO, PMA). **Coelé:** 5–5.5 mi. N of El Copé, 850 m, 8°38'N, 80°35'W, Thompson 4731 (CM); 680–770 m, 8°39'N, 80°36'W, Croat 74836 (MO, PMA, US); 710–800 m, 68764 (MO, NY).

**Veraguas:** Alto Piedra-Calovebora, 0.5 mi. N of Alto Piedra, Parque Nacional Cerro Tute, 800–1030 m, *Croat & Zhu 76906* (CR, MO, NY); ca. 5 mi. N of Alto Piedra, 670 m, 8°33'N, 81°08'W, *Croat 66962* (CM, F, MO); 1250–1350 m, *Croat 48959* (MO).

**COLOMBIA.** **Chocó:** between Medellín and Quibdó, 60 km W of Bolívar, 800 m, *Croat 49271* (MO); 5.5 km E of Tutunendo, 23.5 km E of Quibdó, 150 m, 5°44'N, 76°29'W, *Croat 56227* (COL, JAUM, MO); km 208.5, 9 km W of Tutunendo, ca. 9 km E of Quibdó, <100 m, 5°39'N, 76°40'W, *Croat 56204* (COL, JAUM, MO); Quibdó-Bolívar, Km 175–176, 117–118 km E of Quibdó, 5°44'N, 76°20'W, *Croat 57481* (CHOCO, COL, MO); Quibdó-Lloro, vic. Río Atrato, ca. 150 m, 5°29'N, 76°35'W, *Croat 55996* (COL, JAUM, MO, PMA). **Valle:** Buenaventura-Cali, 14 km SE of Río Sabaletas, 53 km ESE of Querebral, 270, 3°42'N, 77°51'W, *Croat & Wuz 70458* (MO). **ECUADOR.** **Esmeraldas:** Lita-San Lorenzo road (based on *Dodson 18645*), cultivated at Río Palenque, *Croat 73857* (MO).

***Philodendron squamipetiolatum* Croat, sp. nov.**

**TYPE:** Panama. Coclé: El Calvario above El Copé, ca. 6 mi. N of El Copé, 710–800 m, 8°39'N, 80°36'W, 23 June 1994, *Croat 68767* (holotype, MO-3591312–4; isotypes, AAU, B, CAS, CM, COL, CR, DUKE, F, G, GH, HMNM, K, M, MEXU, NY, PMA, RSA, S, SCZ, TEX, US, VEN). Figures 375, 385–388.

Planta hemiepiphytica; internodia 3–17 cm longa, 6–15 mm diam., atriviridis, squamis approximatis, scalariformibus, caespitosis, transverse orientibus; cataphylla usque 29 cm longa, incostata, squamata, persistentia nodis superioribus, mox decidua; petiolus 6–16 cm longus, usque ca. 6 mm diam., teres; lamina ovata, rotundata vel subcordata basi, 15–37.5 cm longa, 9.5–29 cm lata; inflorescentia 1–2; pedaculus subteres, 6–8 cm longus, dense squamatus; spatha 11–15.5 cm longa, 2–2.5 cm diam.; lamina spathae extus viridi, dense squamata, intus atriviridis, tubo spathae extus viridi, intus atriviridis; pistilla 5–6-locularia; loculi 20–30-ovulati.

Appressed-climbing hemiepiphyte, occurring on understory trees in primary forest, fertile ca. 6 cm above the ground; stems trailing when juvenile, loosely appressed-climbing as adults; internodes 3–17 cm long, 6–15 mm diam., dark green, soon dark brown, drying yellow-brown, closely ridged-fissured with the surface moderately glossy, granular and with close, scalariform, tufted transversely oriented scales; cataphylls to 29 cm long, terete, unribbed, green and densely whitish scaly, promptly weathering to a loose semiorganized network of slender, pale fibers with fragments of epidermis, the fibers persisting at upper nodes, but soon completely deciduous; **petioles** erect to spreading, 6–16 cm long, to ca. 6 mm diam., terete, matte, dark green, surface densely covered with antrorse light green scales throughout (these ca. 6 mm long), densely granular-scurfy; **blades** somewhat pendent when

young, spreading with petioles when mature, ovate, subcoriaceous, acuminate at apex, rounded to subcordate at base, 15–37.5 cm long, 9.5–29 cm wide (ca. 1.2–2.1 times longer than wide), (1.6–2.3 times longer than petiole), upper surface dark green, weakly glossy (subvelvety), drying dark brown, lower surface matte and much paler, drying yellow-brown; posterior lobes broadly rounded; sinus arcuate; midrib narrowly sunken and paler above, round-raised to thicker than broad, darker and densely puberulent below; basal veins 2–6(9) per side, with all free to base, or sometimes 3–5 pairs coalesced up to 1.5 cm, posterior rib sometimes lacking; primary lateral veins 5–7 per side, departing midrib at a 50–55° angle, weakly quilt-sunken above, round-raised to thicker than broad, darker and densely puberulent below; minor veins flat but distinct and darker than surface below, some intermittent; “cross-veins” distinct, moderately raised on drying. **INFLORESCENCES** 1–2 per axil; peduncle subterete, 6–8 cm long, densely scaly; **spathe** 11–15.5 cm long, 2–2.5 cm diam. when furled, to 6.5 cm wide, and constricted somewhat when flattened, green outside, especially below, densely scaly (the scales whitish), deep reddish inside, especially in the tube; **spadix** yellowish white, 9.7–13.5 cm long, to 1.3 cm diam., only weakly constricted midway; pistillate portion 2.7–4.7 cm long in the front, 0.7–1.4 cm diam.; staminate portion 7–8.8 cm long; fertile staminate portion broadest at upper one-third and bluntly pointed at apex, 6 mm diam. 1 cm from apex; sterile staminate portion 1.2 cm diam.; pistils 2.2 mm long; ovary 5–6-locular, ca. 1.4 mm long, with axile placentation, the sides covered throughout, especially toward apex, with short pale raphide cells; locules 1–1.2 mm long; ovules 20–30 per locule, 0.1–0.2 mm long, oblong, arranged all along the length of the locules; style similar to style type B; short, 0.9–1 mm diam., drying with 2–5 depressions at apex; stigma 1–1.1 mm diam. when flattened and dried. Inflorescence not seen.

Flowering in *Philodendron squamipetiolatum* occurs in the rainy season, during June in Central America. Ecuadorian collections at post-anthesis have been seen from February and April, with an early fruiting collection from August.

*Philodendron squamipetiolatum* ranges from Panama to Ecuador from sea level to 1300 m elevation in *Premontane rain forest* and *Premontane wet forest* transition to *rain forest* and *Tropical rain forest*. In Panama this species is known only from along the Continental Divide in Coclé Province,

whereas in South America it occurs along the Pacific slope of the Andes.

*Philodendron squamipetiolatum* is a member of *P.* sect. *Philodendron* subsect. *Achyropodium*. This species is distinguished by its long, inconspicuously scaly internodes; long, scaly, terete petioles (hence the name); long, scaly, deciduous cataphylls; rounded to subcordate blades with an arcuate sinus and puberulent major veins below; 1–2 dark green, scaly inflorescences; and spathe tube reddish within.

In Central America, *Philodendron squamipetiolatum* is most easily confused with *P. hammelii*, a species with which it occurs in the vicinity of the type locality. Both species have small, ovate-cordate blades with scaly petioles and persistent, fibrous cataphylls. *Philodendron hammelii* is distinguished by having leaves drying grayish green above and yellowish brown below and closely arranged broad scales only near the apex of the petioles.

This species might also be confused with two others with scaly petioles. It is easily distinguished from both *P. verrucosum* and *P. squamicale* by its much smaller, subcordate vs. deeply cordate leaf bases, and from *P. squamicale* by its more broadly ovate blades (vs. triangular ovate).

*Madison & Besse 17185* from Esmeraldas Province, Ecuador, probably also belongs to this species, but field notes indicate that the stems are smooth. Unfortunately, there is no stem with the collection. It is not known whether the intent was to indicate that the stems lacked scales or whether it lacked other features. In addition, the stems were reported to be 10–14 cm long and to 1.5 cm diam.

*Additional specimens examined.* PANAMA. **Coclé:** Alto Calvario, ca. 4.6 km above El Copé, 800 m, 8°39'N, 80°36'W, *Croat 74857* (AAU, B, CAS, CM, COL, DUKE, F, G, GH, HMN, K, M, MEXU, MO, NY, PMA, RSA, SCZ, S, TEX, US, VEN); 75098 (MO). **Colón:** Río Guanche, 30–100 m, *Croat 79327* (PMA, MO), 79350 (PMA, SCZ).

COLOMBIA. **Chocó:** Serranía de Baudó, Las Animas-Pato on Río Pato, ca. 4 km SW of Pato, ca. 150 m, 5°30'N, 76°46'W, *Croat 56140* (MO); Serranía de Baudó, Río Pato, 10 km SW of Pato, 5°17'N, 76°45'W, *Croat 56074* (MO); San José del Palmar–Novita, ca. 3 km W of San José del Palmar, 930 m, 4°56'N, 76°29'W, *Croat 56650* (COL, JAUM, MO); Pueblo Rico–Istmina, along Quebrada Antón, 15 km W of Santa Cecilia, 240–350 m, 5°20'30"N, 76°13'45"W, *Croat 70972* (MO); 300 m downstream and across the river from Catrú, 50–100 m, *Warner & White 123* (COL). **Valle:** Buenaventura–Cali, Sabaletas, km 29, 25 m, *Killip & Cuatrecasas 38775* (US); 100 m, *Croat 38567* (F, MO); vic. Bajo Calima, <100 m, 4°03'N, 77°08'W, *Croat 70218* (CM, MO, NY); 100 m, *Montalvo 3193* (MO). ECUADOR. **Carchi:** El Paillón, ca. 45 km below Maldonado, 800 m, *Madison & Besse 7185* (SEL);

Tulcán Cantón, Awá Reservation, Gualpí Chico area, 1300 m, 0°58'N, 78°16'W, *Hoover et al. 3710* (MO, QCA). **Esmeraldas:** Lita, 550–650 m, *Madison et al. 5057* (SEL); Cerro de Río Bravo de Cayapas, 250 m, 0°41'N, 78°56'W, *Holm-Nielsen et al. 25535* (AAU); 40.1 km W of Lita, 350 m, 0°56'N, 78°40'W, *Croat 72309* (L, MO, QCA, US). **Imbabura:** Lita, 812 m, *D'Arcy 14850* (MO). **Pichincha:** Reserva ENDESA, 9 km N of Km 113 on Quito–Pto. Quito Highway, 750–800 m, 0°05'N, 79°02'W, *Rodríguez 247* (MO, QCA); *Croat & Rodríguez 61483* (MO, QCA); 650–800 m, 0°03'N, 79°07'W, *Cerdán & Ayala 10169* (MO); along tributary of Río Guayllabamba, 600 m, 0°10'N, 79°03'W, *Grayum et al. 9353* (MO). **Los Ríos:** La Centinela at Km 12, N of Patricia Pilar, 600 m, *Dodson & Dodson 6771* (MO, SEL); Río Verde, 2 km SE of Sto. Domingo de Los Colorados, 530 m, *Dodson 7435* (SEL).

**Philodendron straminicaule** Croat, sp. nov.

TYPE: Panama. Chiriquí: Fortuna Dam area, trail to meteorological station of Río Hornito, ca. 0.5 km S of Centro de Científicos, 8°45'N, 82°18'W, 23 June 1994, *Croat & Zhu 76302* (holotype, MO–4610957–9; isotypes, AAU, B, CAS, COL, CR, F, GH, K, M, MEXU, NY, PMA, QCNE, S, SEL, US, VEN). Figures 6, 376, 389–392.

Planta hemiepiphytica; internodia 1.5–5(12) cm longa, 2–3.5 cm diam., plus minusve fragilia, canoviridia vel atriviridia, in sicco straminea; cataphylla (17)20–35 cm longa, decidua; petiolus subteretes, (27)32–59 cm longus, 5–12(17) mm diam.; lamina triangulari-ovato-cordata, (27)33–55 cm longa, (11)19.7–31 cm lata, cordata basi; inflorescentia 2–3(5); pedunculus (4.5)5.6–18 cm longus; spathe 7–18.5 cm longa; lamina spatheae alba in superficiebus ambabus; tubo spatheae extus pallide viridi, intus rubroviolaceo vel marronino; pistula 4–6(10)-locularis; loculi 1-ovulati; baccae viridulae vel lavandulae.

Hemiepiphytic; stem appressed-climbing, leaf scars conspicuous, 1–1.6 cm long, 1.2–1.7 cm wide; internodes glossy to semiglossy, ± brittle, somewhat scurfy, 1–5(12) cm long, 2–3.5 mm diam., longer than broad, dark green, soon gray-green, finally light tan, drying straw-colored, sometimes with a narrow ring of yellow-brown periderm at the nodes; epidermis transversely fissured in part, sometimes deeply fissured and cracking free on drying; roots blunt, to 30 cm or longer, 4 mm diam., yellow-green, weakly glossy (fresh), drying reddish brown with minute scales in age; cataphylls somewhat spongy, to (17)20–35 cm long, variously ribbed, sometimes D-shaped or unribbed, more frequently bluntly to sharply 1- or 2-ribbed (sometimes unribbed in Coclé), emarginate at apex, pale green to pale yellowish to whitish, margins clear, sometimes closely dark green short-lineate, sometimes moderately red-purple-spotted (more so at base) throughout, ribs darker, drying greenish or brownish, deciduous, persisting intact; **petioles** (27)32–59 cm long, 5–12(17) mm diam., subterete,



± spongy, medium green, obtusely flattened adaxially, rounded abaxially, surface semiglossy, faintly dark lineate; **blades** triangular-ovate-cordate, subcoriaceous, semiglossy, bicolorous, long acuminate at apex, cordate at base, (27)33–55 cm long, (11)19.7–31 cm wide (1.3–2.3 times longer than wide), (0.7–1.5 times longer than petiole), margins hyaline, lower surface drying green to yellow-green, densely glaucous with secretory ducts visible; anterior lobe (18.7)27–38.6(47) cm long, 17.6–27.6 cm wide (2.4–4.5 times longer than posterior lobes), margins straight to moderately concave; posterior lobes directed toward base, rarely reflexed-spreading, sometimes overlapping when posterior lobes are held somewhat upward at an angle to the midrib, (7)8.5–15.5 cm long, 7.5–13 cm wide, obtuse to rounded; sinus hippocrepiform, rarely closed, 2.6–14 cm deep; midrib flat to broadly rounded, paler than surface above, convex to narrowly rounded, paler than surface below; basal veins 4–6(7) per side, with 0–1 free to base, the remainder coalesced 1.2–5 cm, 2 coalesced 3.5–7 cm; posterior rib naked 0.5–1.5 cm; primary lateral veins 3–6 per side, departing midrib at a 45–65° angle, acutely ascending then weakly arcuate to straight to the margins, ± obscure, sunken and concolorous above, convex, darker than surface or paler than surface below; interprimary veins raised and darker than surface below; minor veins moderately distinct, fine, darker than surface, arising from both the midrib and primary lateral veins, drying minutely granular, prominulous and alternating with secretory ducts. **INFLORESCENCES** erect, usually 2–3 (sometimes to 5) per axil; peduncle (4.5)5.6–18 cm long, terete, pale to medium green, faintly short-lineate to densely speckled; **spathe** 7–18.5 cm long (0.9–1.8 times longer than peduncle), moderately constricted above the tube; spathe blade pale green to white on both surfaces, acuminate at apex; spathe tube dark green, finely pale lineate, glossy outside, 4–8.5 cm long, 1.5–3 cm diam., reddish violet to maroon or magenta inside; **spadix** sessile; erect at anthesis, 7.6–13 cm long; pistillate portion 2.2–3.7(6.5) cm long, 1–1.4(2) cm diam. at middle, tapered slightly toward both ends; staminate portion clavate, 5.2–7.7 cm long; fertile staminate portion white, 8–10 mm diam. at base, 1–1.5 cm diam. at middle, 4–8 mm diam. ca. 1 cm from apex, broadest at the middle, abruptly tapered toward apex, weakly constricted just above the sterile portion; sterile staminate portion usually broader than the pistillate portion at anthesis, 8–15 mm diam., the lowermost flowers sometimes drying whitened; pistils 1.1–1.7(3.4) mm long, 7–9(17) mm diam.; ovary 4–6(10)-locular, 0.5–0.7(2.2) mm

long, 0.7–0.9(1.7) mm diam., with sub-basal placentation; locules 0.5–0.7(2.1) mm long, 0.2–0.3(0.6) mm diam.; ovule sac 0.5–0.7 mm long; ovules 1 per locule, contained within sticky, transparent, gelatinous ovule sac, 0.2–0.3 mm long, as long as funicle; funicle 0.2–0.3 mm long (can be pulled free to base), style 0.5–0.8(1) mm long, 0.6–0.7(1.7) mm diam., drying doughnut-shaped, similar to style type C; style funnel shallow with a small ring of stylar canals near the base; stigma subdiscoid, truncate, 0.4–0.5(0.8) mm diam., 0.1–0.3 mm high, covering almost entire style apex, with 1–5 depressions when dried, papillate, sessile, matte; the androecium truncate, prismatic, margins ± 4–5-sided, 0.6–1 mm long, 1.1–1.3 mm diam. at apex; thecae oblong, 0.4–0.5 mm wide, ± parallel to one another and nearly contiguous to contiguous; pollen ± spheroidal, <0.1 mm long, <0.1 mm diam., scarce; sterile staminate flowers irregularly 4–6-sided, blunt, prismatic, 1–1.3 mm long, 0.9–1.2 mm wide. **INFRACTESCENCE** with berries pale green (immature), greenish white to lavender.

Flowering in *Philodendron straminicaule* is believed to occur throughout much of the dry season to the mid-rainy season (January through August) in Panama, but flowering specimens have been seen from only April through August. Post-anthesis inflorescences have been seen from as early as March, and immature fruits as early as February, making it obvious that the plants have to be in flower as early as January. Post-anthesis inflorescences have also been collected from May through September and November, and immature fruits occur in March and July. The post-anthesis inflorescences from November suggest that flowering occurs as late as September or October.

*Philodendron straminicaule* is known only from the Cordillera de Guanacaste in northwest Costa Rica and in western Panama, from Chiriquí and Bocas del Toro to Coclé Province, at (710)950 to 2200 m elevation, generally in *Tropical Lower Montane rain forest* and *Premontane rain forest*, rarely in *Tropical wet forest*.

*Philodendron straminicaule* is a member of *P. sect. Calostigma* subsect. *Glossophyllum* ser. *Ovata*. This species is characterized by its grayish (drying straw-colored), more or less brittle, glossy internodes, which are longer than broad; deciduous cataphylls; obtusely flattened, more or less spongy petioles; and triangular-ovate-cordate, green-drying blades with the lower surface drying densely and minutely granular and secretory ducts usually prominently visible and alternating with the minor veins. Also characteristic is the white spathe blade

and tube green outside and reddish violet to maroon inside.

*Philodendron straminicaule* is similar to *P. wilburii* var. *longipedunculatum*, but the latter species differs in having longer, more slender internodes; longer, unribbed to bluntly 1-ribbed cataphylls; typically smaller blades that dry brownish with mostly 3-4 primary lateral veins; and especially by having the peduncle usually as long as or even longer than the spathe (vs. usually less than length of spathe). It also differs in having orangish (rather than greenish white to lavender) berries.

The species may be confused with *P. alticola* and *P. smithii*. See those species for separation.

A noteworthy collection is *Croat 66503*, which has the veins on the lower leaf surface drying darker than the surface rather than paler, which is normally the case.

Collections from Costa Rica differ from those in Panama in having slightly smaller leaves with both the petioles and the blades shorter than those in Panama (petioles to 28 cm long and blades to 31 cm long in Costa Rica vs. petioles to 59 cm long and blades to 55 cm long in Panama). Nevertheless, the smaller leaves in Panama range down to only slightly larger than those in Costa Rica (with petioles as short as 32 cm long and blades to as small as 33 cm long).

One collection from Guanacaste (*Herrera 1473*) is unusual in having more narrowly triangular blades with concave margins, narrower posterior lobes, a parabolic sinus, and a D-type style. However, it is otherwise similar, having the same dried stem characteristics and 1 ovule per locule. More collections are needed to determine if this entity is distinct from *P. straminicaule*.

This species perhaps also occurs in Colombia based on collections from Valle Department (*Junco 2054*) at 580 m and from Gorgonilla Island, Cauca Department (*Killip & García 33060*). These collections differ from those in Panama in having blades with no obvious secretory ducts visible on the abaxial surface. In addition, the latter collection is from just 130 to 200 m elevation, substantially lower than where it is found in Panama.

*Additional specimens examined.* COSTA RICA. **Alejuela:** above Bijagua, slopes of Miravalles, Gómez et al. 19171 (MO). **Guanacaste:** Parque Nacional Guanacaste, Estación Caño, 1100 m, 10°55'45"N, 85°28'15"W, Chávez 80 (INBIO, MO); Estación Mengo, 1100 m, 10°55'N, 85°28'W, INBio 219 (CR, MO); Parque Nacional Rincón de la Vieja, SE slopes of Volcán Santa María, above Estación Hacienda Santa María, 900-1200 m, 10°47'N, 85°18'W, Davides et al. 23397 (CR, MEXU, MO, TEX); 1350-1400 m, 10°46'N, 85°49'W, Herrera 1473 (CR, MO). PANAMA. **Bocas del Toro:** Fortuna Dam area,

Gualaca-Chiriquí Grande, 21.4 km past Gualaca, 8°32'N, 82°19'W, Hoover 1326 (MO); Continental Divide, 1200 m, 8°44'N, 82°17'W, Croat 60350 (CM, MO); 60364 (AAU, CM, COL, CR, F, M, MO, PMA, UC, WIS), 60365 (K, MO, PMA, US); 8°45'N, 82°15'W, McPherson 10867 (CR, MO, PMA); 2.8 mi. from Divide, 850-950 m, ca. 8°45'N, 82°15'W, 9675 (L, MO, PMA, US). **Bocas del Toro-Chiriquí:** Fortuna Dam area, Continental Divide, above Quebrada Arena, road to Oleoducto, 1150-1200 m, Knapp & Vodicka 5658 (MO, PMA). **Chiriquí:** Cerro Colorado, above San Félix, 18-27 mi. off of Pan-American Highway, 1200-1500 m, Croat 33146 (MO); 19.7 mi. N of Río San Félix, 1420 m, 8°31'N, 81°46'W, Croat 74996 (MO); 33.1 km N of Río San Félix, ca. 1400 m, Croat 37203 (MO, PMA); 20 mi. N of Río San Félix, 2000 m, Croat 48446 (COL, K, MEXU, MO, PMA, VEN, W); 1420 m, 75010 (MO, PMA); vic. of Chame, 800-1200 m, Croat 33440 (MO); 11.2 km from Chame, 1700 m, Folsom 4889 (CAS, MO); 9.2 mi. W of Chame, 1450-1480 m, 8°35'N, 81°50'W, Croat 69013 (B, CAS, CM, COL, CR, F, G, K, L, MEXU, MO, PMA, TEX, US); Cerro Horquetas, 7000 ft., Blum & Dwyer 2591 (SCZ); Boquete, 5000-6000 ft., Dwyer & Hayden 7736 (MO); 6500 ft., von Hagen & von Hagen 2073 (MO, NY); 2164 (MO, NY); Volcán Barú, along old road to Boquete, 1750-1900 m, 8°50'N, 82°30'W, McPherson 11340 (K, MO); Cerro Pate Macho, ca. 5 mi. NE of Boquete, 1800-2200 m, Croat 48562 (MO); 1900-2000 m, 8°46'N, 82°25'W, 66503 (MO); ca. 6 km NE of Boquete, 8°49'N, 82°23'W, Grayum 6408 (CM, MO, PMA); 1600-1700 m, Grayum et al. 6397 (MO, US); Río Palo Alto, 1300-1800 m, 8°47'N, 82°22'W, Knapp et al. 2042 (MO); Fortuna Lake area, Gualaca-Chiriquí Grande, along Continental Divide, 1200 m, 8°44'N, 81°17'W, Croat 74972 (MO, PMA); 4.5-5 km N of dam over Fortuna Lake, 1100-1135 m, 8°43'N, 82°17'W, Croat & Grayum 60057 (AAU, B, CAS, F, K, L, MEXU, MO, PMA, SEL, U); ca. 4.6 mi. N of Los Planes de Hornito, Stevens 18437 (MO); behind Vivero Forestal (now Centro de Científicos), 12 km N of Los Planes de Hornito, 1200-1300 m, 8°45'N, 82°12'W, Knapp 4950 (MO); 10.1 mi. NW of Los Planes de Hornito, 1250 m, 82°17'W, 8°45'N, Croat 49837 (BR, COL, MO, NY, TEX, UC), 50030 (MO, NY, TEX); vic. IRHE facilities, 1100-1200 m, 8°45'N, 81°18'W, Croat 66542 (MO, PMA), 66584 (AAU, BR, CAS, CDBI, CM, DUKE, F, IBE, K, KYO, LE, MBM, MEXU, MO, NY, PE, PMA, RB, SCZ, UC, US); Río Chiriquí, 9.6 mi. beyond Los Planes de Hornito, 1300 m, Croat 48733 (CR, MO); 11.8 mi. NW, 1400 m, 48686 (MO), 48703 (CM, MO), 48701 (CAS, MO); Quebrada Bonita, 1100 m, 8°45'N, 82°13'W, Churchill 5262 (MO); Quebrada Ortega, 1200 m, 8°41'N, 82°14'W, Churchill 5283 (MO); 8 mi. beyond Los Planes de Hornito, 1010-1130 m, 8°44'N, 82°14'30"W, Croat 67920 (AAU, CAS, MO, PMA, US). **Coelá:** El Copé region, Alto Calvario, ca. 6 km N of El Copé, 770 m, 8°38'N, 80°35'W, Croat & Zhu 76761 (MO, PMA); 710-800 m, 8°39'N, 80°36'W, Croat 68816 (MO); 900 m, 44576 (MO); 5.5 mi. N of El Copé, 850 m, 8°39'N, 80°36'W, 67574 (CM, MO, PMA, TEX). **Vereguanos:** Santa Fe region, Escuela Agrícola Alto Piedra, ca. 5-8 km NE of school, 730-770 m, Croat 25909 (MO, NY).

**Philodendron strictum** G. S. Bunting, *Phytologia* 60: 328. 1986. TYPE: Venezuela. Táchira: San Cristóbal-Chorro del Indio-Caño Seco-La Florida, km 20-22 (E of San Cristóbal), 1100-1125 m, 6 Mar. 1977, Bunting & Borges 5001 (holotype, NY; isotypes, PT, VEN). Figures 393-395, 397, 398.

Terrestrial or hemiepiphytic; stems stout; internodes 3–4 cm long, 2.5–5 cm diam., usually broader than long, sometimes longer than broad, dark green to gray-green, semiglossy; roots moderately few, drying dark brown, semiglossy, sparsely scaly; cataphylls 25–40 cm long, unribbed to bluntly 1-ribbed, rarely bluntly 2-ribbed or sharply 1-ribbed, pale green, soft, turning yellowish and persisting semi-intact at upper nodes; **petioles** (47–56/63–105 cm long, subterete, obtusely D-shaped with faint medial rib, obtusely flattened adaxially, light green to gray-green, weakly glossy, sparsely dark lineate, drying usually light yellow-brown, sometimes blackened; sheath inconspicuous; **blades** ovate-cordate, conspicuously bicolorous, acuminate at apex (the acumen sometimes inrolled), cordate, sometimes sagittate at base, (24)38–66(74) cm long, (16)27–52 cm wide (1–1.3 times longer than wide), (0.5–1 times the petiole length), broadest near the middle; upper surface dark green, semiglossy, lower surface much paler, whitish, matte; anterior lobe (19–28)31–51(57–62) cm long, (19)27–48(53–56) cm wide, (1.7–2.8)(3.3) times longer than posterior lobes; posterior lobes 12–23 cm long, 8.6–26 cm wide, obtuse to broadly rounded; sinus usually spatulate, sometimes hippocrepiform; midrib flat, paler than surface above, convex and darker below; basal veins 7–10 per side, with 0–2 free to base, part of the remainder coalesced 1–12.4 cm; posterior rib sometimes not naked, often obscurely naked for 1.5 cm, rarely 3.5 cm; primary lateral veins 5–11 per side, departing midrib at a 50–65° angle, deeply sunken above, convex and darker below, usually prominently downturned before meeting midrib; minor veins numerous, fine, indistinct below, arising from both the midrib and primary lateral veins. **INFLORESCENCES** erect-spreading, 1–4 or more per axil; peduncle (6)11–15 cm long, 1–1.5 cm diam., with thin, yellowish epidermis; **spathe** 13–17.7 cm long, 2.5–4.3 cm diam. (0.9–2.3 times longer than peduncle), moderately constricted above the tube (very slick inside), 2.7 cm diam. at constriction; spathe blade light green to whitish, tinged purple-violet (B & K red-purple 3/2.5), sparsely short-white-lineate medially outside, margins paler, (opening elliptic in face view, 10.7 cm long, 4.8 cm wide), light green to whitish and suffused red inside, drying dark to reddish brown; spathe tube green to purple-violet (B & K red-purple 3/2.5) outside, 4 cm diam., maroon or violet-purple inside; **spadix** sessile; bluntly pointed at apex, 9.2–16 cm long; pistillate portion weakly tapered toward apex, 2.1–3.9 cm long in front, (1.6)3–3.3 cm long in back, 0.6–1.4 cm diam. at apex, 1–1.5 cm diam.

at middle, 1.3–1.5 cm wide at base; staminate portion 8.9–11.9 cm long; fertile staminate portion somewhat ellipsoid, sometimes ovate to tapered, 1.1 cm diam. at base, 1.2–1.3 cm diam. at middle, 8–9 mm diam. ca. 1 cm from apex, broadest in the middle or sometimes just above the base, narrower than the pistillate portion, as broad as the sterile portion; sterile staminate portion narrower than the pistillate portion, 1–1.4 cm diam.; pistils (1.1)2.7–3.4(5.4) mm long, 1.3–1.6 mm diam.; ovary (4)5–6-locular, 1.8 mm long, 1.3–1.7 mm diam., with axile placentation; locules 1.8 mm long, 0.6 mm diam.; ovules 20–28 per locule, 2-seriate, 0.2–0.4 mm long, longer than funicle; funicle 0.1 mm long, adnate to lower part of partition, style 0.7–0.9 mm long, 1.3–1.6 mm diam., similar to style type B; style apex flat; stigma subdiscoid, truncate, 1.1–1.3 mm diam., 0.3–0.5 mm high, covering entire style apex; the androecium prismatic, truncate, oblong, irregularly 4–6-sided at apex 0.9 mm long, 1.6–2 mm diam. at apex; thecae oblong, ± parallel to one another, nearly contiguous; sterile staminate flowers blunt, prismatic, irregularly 4–6-sided, 1.8–2.9 mm long, 0.9–2.4 mm wide. **INFRACTESCENCE** 2 cm wide; seeds pale yellowish, 0.9 mm long, 0.3 mm diam.

Flowering in *Philodendron strictum* apparently occurs throughout the dry season and first half of the rainy season in Central America (January through September, though no flowers were seen in May). South American flowering collections have been seen from February and July, and post-anthesis inflorescences (or immature fruits) from January through November. Mature fruits have been seen only from September.

*Philodendron strictum* ranges from Costa Rica to western Panama, Venezuela (Táchira), Colombia (Antioquia, Chocó), and Ecuador (Carchi, Esmeraldas). In Costa Rica, this species occurs at 850 to 1525 m, and in Panama at 680 to 1665 m elevation in *Lower Montane rain*, *Premontane rain*, and *Tropical wet forest* life zones. In Colombia, this species has been collected at 100 to 150 m in Chocó and Valle Departments, and in Antioquia at 1560 m. In Venezuela, it is known only from the state of Táchira in the southwestern part of the country, at 1000 to 1250 m elevation in *Premontane wet forest*. It was reported erroneously (owing to a typographical error) by Croat and Lambert (1986) from 110 to 1330 m.

*Philodendron strictum* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Impolita*. This species is characterized by its usually terrestrial habit; thick stems; short internodes; thick, yellowish

lowish, unribbed to bluntly one-ribbed cataphylls persisting semi-intact at the upper nodes; obtusely flattened to D-shaped petioles usually drying pale yellow-brown; and ovate-cordate blades with the lower surface whitish and matte.

*Philodendron strictum* is most easily confused with *P. hebetatum*, which shares blades with whitish, matte lower surfaces, as well as yellow-drying cataphylls and petioles. Both species are easily identified by these features alone. The two species are sympatric in at least one area along the Fortuna Dam road in Panama but remain distinct by virtue of their respective habits and blade shapes. *Philodendron hebetatum* differs in being consistently an appressed epiphyte and in having a triangular-ovate blade vs. a generally terrestrial habit and consistently ovate blades for *P. strictum*. Juvenile plants of the two species, at this stage both terrestrial, are quite distinct with the blades of *P. hebetatum* more elongate (2.5–3 times longer than broad), while those of *P. strictum* are more broadly ovate (1.2–2.5 times longer than wide).

This species is also easily confused with the often syntypic *P. thalassicum*, which differs in having blackened rather than pale yellow-brown petioles, cataphylls promptly weathering to pale fibers vs. usually persisting yellowish and semi-intact in *P. strictum* and internally greenish to white spathes (vs. maroon or violet-purple in *P. strictum*). In addition (at least in Panama), the petioles of *P. thalassicum* are more sharply D-shaped vs. obtusely flattened in *P. strictum*.

A collection from 1875 m elevation in Antioquia Department, Colombia (McPherson 12939), possibly belongs to this species but differs in having reddish brown cataphylls that are more fragmented on drying. It also has the petioles drying dark brown, not yellowish as is typical for the species in Colombia and Ecuador.

*Additional specimens examined.* COSTA RICA. **Alajuela:** San Ramón, Bajo Rodríguez, 1025–1100 m, *Croat* 78988 (CR, INB, MO). **Cartago:** ca. 11 mi. NE of Turrialba, 850 m, *Croat* 43354 (MO); Turrialba–Limón, Hwy. 32, ca. 11 mi. S of Siquirres, 650 m, *Croat* 43335 (MO); 31 km S of Siquirres on road to Turrialba (CR-10), 850 m, 9°57'N, 83°36'W, *Thompson & Raulins* 1167 (CM); Shipití, Moravia de Chirripó, 900–1000 m, *I. Chacón* 9 (MO); Tapantí, Nilsson & Chacón 236 (CR); *Croat* 79039 (MO). **Puntarenas:** ca. 1 km S of San Vito, 1100 m, *Croat* 66168 (CR, G, MO, NY, PMA); Las Cruces, near San Vito de Java, ca. 4000 ft., *Croat* 32959 (MO); N of Palmar Norte, trail to Jalisco, 50–700 m, *Croat* 35208 (MO). **San José:** San Isidro de El General–Dominical, 4.8 mi. from Rio Pacuare, 1000 m, *Croat* 32550 (MO); 990–1100 m, *Croat* & *Hannon* 79105 (CR, INB, MO); Braulio Carrillo National Park, 1400 m, *Croat* 78811 (CR, INB, MO); Tarrazú, vic. Hormiguero, 1100–1200 m, *Croat* 78939 (CR,

INB, MO). **PANAMA. Bocas del Toro:** Fortuna Dam area, Continental Divide, 1170 m, 8°44'N, 82°17'W, *Croat* 66649 (AAU, B, CAS, CM, COL, CR, F, G, K, L, MEXU, MO, NY, PMA, US, VEN); 1000 m, 8°48'N, 82°12'W, *Churchill et al.* 4646 (MO); ca. 1200 m, 8°44'N, 82°17'W, *Croat* & *Grayum* 60326 (B, CM, K, MO, RSA, US); 2.2 mi. N of Continental Divide, 820 m, 8°45'N, 82°16'W, *Croat* 60401 (MO); 825 m, 8°45'N, 82°15'W, *McPherson* 7361 (B, K, MO, PMA, RSA); 1.2 mi. N of Continental Divide, 910 m, 8°44'N, 82°17'W, *Croat* 60475 (MO, PMA). **Chiriquí:** Cerro Colorado, 9–10 mi. road to Chame, 1200–1500 m, *Croat* 33272 (MO); 7.7 mi. beyond Chame, 1420 m, 8°31'N, 81°46'W, 74995 (MO); Chame, 1600 m, *Kress et al.* 86-1925 (SEL); 9.2 mi. W of Chame, 1450–1480 m, 8°35'N, 81°50'W, *Croat* 69064 (MO); 6.5 km beyond Chame, 1660 m, 8°30'N, 81°46'W, *Croat* 74991 (MO); Fortuna Dam area, Gualaca–Chiriquí, 1200 m, 8°46'N, 82°16'W, *Croat* 68930 (F, MO); E of main camp at dam site, 1400–1500 m, *Folsom et al.* 5448 (MO); trail to Rio Hornito weather station, 1100–1200 m, 8°45'N, 82°15'W, *Thompson* 5041 (CM); 8°45'N, 82°18'W, *Croat* & *Zhu* 76311 (AAU, CM, COL, CR, DUKE, EAP, GH, IBE, MEXU, MO, P, QCNE, RSA, US); 76375 (CAS, CM, COL, CR, DUKE, ENCB, F, GB, K, L, MEXU, MO, NY, P, SEL, WIS); Continental Divide, 1200 m, 8°44'N, 81°17'W, *Croat* 74971 (MO, PMA); Quebrada Arena, just S of Continental Divide, 1050 m, 8°45'N, 82°16'W, *Hammel et al.* 14707 (MO); Quebrada Los Chorros–Quebrada Honda, to N of reservoir, 8°45'N, 82°14'W, *H. Churchill* & *A. Churchill* 6166 (MO, NY); vic. IRHE facilities at dam, 1200–1300 m, 8°45'N, 82°18'W, *Croat* 66543 (AAU, DUKE, F, M, MEXU, MO, PMA); Gualaca–Chiriquí Grande, 7.2 mi. beyond Los Planes de Hornito, 1165–1200 m, 8°44'N, 82°14'W, *Croat* 67840 (CM, MO, PMA); 8.4 mi. beyond Los Planes de Hornito, 1130 m, 8°44'N, 82°14'35'W, 67870 (G, M, MO). **Coeló:** El Copé region, Alto Calvario, ca. 6 km N of El Copé, 8°39'N, 80°36'W, *Croat* 68818 (MO); 680–770 m, 8°39'N, 80°36'W, 74852 (MO, W); 700 m, 8°38'N, 80°35'W, *Croat* & *Zhu* 76796 (MO, NY); El Valle region, vic. of La Mesa, N of El Valle de Antón, Cerro Gaital, 900–1000 m, 8°37'N, 80°08'W, *Croat* 67227 (B, CAS, CM, K, MO). **Veraguas:** Santa Fe region, Alto de Piedra, 800–950 m, 8°33'N, 81°08'W, *Croat* 67003 (CAS, MO); Parque Nacional Cerro Tute, 0.5 mi. N of Alto Piedra, 800–1030 m, *Croat* & *Zhu* 76885 (MO); 1000–1250 m, *Croat* 48920 (MO).

**Philodendron subincium** Schott, Oesterr. Bot. Z. 9: 99. 1859. TYPE: Mexico. Veracruz: Papantla, 20°27'N, 97°19'W, *Karwinski* s.n. (holotype, LE? destroyed?); Schott ic. 2636 (neotype, here designated, W). Figures 396, 401, 402.

Hemiepiphytic; leaf scars conspicuous, 1–1.5 cm long, 1–2 cm wide; internodes 1.2–6 cm long, 2.7–3 cm diam., broader than long, tan to reddish brown, epidermis with loose flakes; roots few to numerous per node; cataphylls moderately spongy, 23–34 cm long, unribbed, drying sharply 1-ribbed in lower one-third, pale green, drying tan to reddish brown, persisting semi-intact, eventually fibrous; petioles 55.5–68 cm long, 1.4–1.5 cm diam., terete, medium green, drying yellowish tan to reddish

tan, surface dark green diffuse-lineate; **blades** narrowly ovate, coriaceous, semiglossy, moderately bicolorous, long acuminate at apex, sagittate at base, (40)57.5–72 cm long, (24)32.5–37 cm wide (1.7–1.8 times longer than wide), (ca. 1 time longer than petiole), margins hyaline, markedly sinuate, upper surface drying brown, lower surface drying yellow-red to brown; anterior lobe (32.5)40–49 cm long, (10)17.6–18 cm wide (2.3–3.3 times longer than posterior lobes); posterior lobes (10)17–18 cm long, 16–16.5 cm wide, obtuse; sinus hippocrepiform or closed and obovate with lobes overlapping, acute when young; midrib broad, flat, paler than surface above, convex, concolorous below; basal veins 4–9 per side, first (second) pair free to base, most coalesced 1–4 cm, 2–3 coalesced to 5.5 cm; posterior rib not naked or obscurely naked for ca. 1 cm; primary lateral veins 5–7 per side, departing midrib at a 50–60° angle, sunken, paler than surface above, convex, concolorous below; minor veins flat, darker than surface below, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 2 per axil, ± erect; peduncle 7.5–10 cm long, 1.3 cm diam., medium green; **spathe** 16.5–20 cm long, 3.5 cm diam. (closed), 5 cm diam. (opened), constricted above the tube, acuminate at apex, margins paler; spathe blade semiglossy, greenish outside, white inside; spathe tube more ellipsoid, maroon-purple outside, bright cherry-red inside; **spadix** not seen.

The flowering phenology of *Philodendron subincisum* is unknown. The only modern collection (Moore & Bunting 8952) is sterile.

*Philodendron subincisum* is endemic to Mexico, known only from northern Veracruz in the Poza Rica region, at less than 500 m elevation in "Selva Mediana Subperennifolia."

*Philodendron subincisum* is a member of *P.* sect. *Calostigma* subsect. *Macrobolium* ser. *Macrobolium*. This species is characterized by its large, thick stems, short internodes, deciduous cataphylls, subterete petioles, ovate-cordate blades with markedly sinuate margins, two short-pedunculate inflorescences per axil, spathe tube maroon-purple outside and bright cherry-red inside, and spathe blade green outside and white within.

*Philodendron subincisum* is apparently rare, having been collected only once (Moore & Bunting 8952) since the type gathering.

*Philodendron subincisum* appears to be most closely related to *P. sagittifolium* based on most features, but it may represent a hybrid. Bunting (1965) even suggested that *P. radiatum* was closely allied to *P. subincisum*. Judging from its rarity, the

latter taxon might be a hybrid between *P. radiatum* and *P. sagittifolium*. Because of its ovate blade with sinuate margins, it is not confused with either of the above species. Perhaps most easily confused with *P. subincisum* is *P. radiatum* var. *pseudoradiatum*. That taxon differs in occurring only on the Pacific slope and in having ovate-triangular blades with proportionately narrower and longer lobes.

The inflorescence of this species remains poorly known. Collected in sterile condition by Moore and Bunting (8956), the species later flowered in cultivation at Cornell University. The flowering specimen was apparently vouchered by Bunting, but the whereabouts of the collection are unknown. Much of the information concerning the species is based on photographs published by Bunting (1965).

*Additional specimen examined.* MEXICO, Veracruz: Rancho El Huasteco, 14.6 mi. from Tuxpan on road to Tihuatlán, Moore & Bunting 8956 (BH, MO).

***Philodendron sulcicaule*** Croat & Grayum, sp. nov. TYPE: Costa Rica. Limón: 7 km SW of Bribrí, 100–250 m, ca. 9°36'N, 82°54'W, 4 May 1983, Gómez, Liesner & Judziewicz 20473 (holotype, MO-3160203; isotypes, B, CR, K, PMA, US). Figures 399, 400, 403.

Planta hemiepiphytica; internodia 6–16 cm longa, 5–13 mm diam., profunde sulcata; cataphylla 12 cm longa, incostata, vel acute 1-costata, raro acute 2-costata, decidua intacta; petioli 11–20 cm longus, subterete, leniter complanatus adaxialiter; lamina ovato-cordata, abrupte vel longa acuminata, 14.5–24 cm longa, 9.6–15 cm lata; nervis basalibus 3 utroque, infirmis; nervis lateralibus 1 obscuris; inflorescentia 1–4(5); pedunculus tenuis, 3.5–10 cm longus, 1–3 mm diam.; spathe (4.5)5.8–11 cm longa, omnino alba, tubo spatheae intus rubro, roseo aut purpureo basi; pistilla 4–5-locularia; loculi 1-ovulati.

Hemiepiphytic vine; stem scandent, drying yellow, glossy; internodes deeply sulcate, prominently ribbed, semiglossy, 6–16 cm long, 5–13 mm diam., longer than broad, olive-green; epidermis becoming brown, breaking free on bending stem; roots reddish-brown, slender, curled, 8–20 cm long; cataphylls 12 cm long, unribbed to sharply 1-ribbed, rarely sharply 2-ribbed, green rarely pink, deciduous intact; **petioles** 11–20 cm long, 4–8 mm diam., subterete, dark green, weakly flattened adaxially, semiglossy; **blades** ovate-cordate, subcoriaceous, semiglossy, moderately bicolorous, abruptly- to long-acuminate at apex (the acumen apiculate, to 4 mm long), cordate at base, 14.5–24 cm long, 9.6–15 cm wide (1.3–1.9 times longer than wide), (0.9–1.6 times longer than petiole), about two-thirds as long as petiole, margins drying moderately undulate, upper surface dark green, drying usually

dark brown, semiglossy to matte, lower surface paler, drying light brown to greenish brown, sometimes yellowish brown, semiglossy; anterior lobe 10–19.6 cm long, 7–15 cm wide (2.6–3.8 times longer than posterior lobes); posterior lobes 3.5–6 cm long, 2.7–6.4 cm wide, rounded-obtuse to nearly acute; sinus arcuate with blade decurrent on petiole to short-hippocrepiform; midrib flat to weakly raised, concolorous above, convex, concolorous to paler than surface below; basal veins to 3 per side, weak, 1 sometimes free to base, the remainder coalesced 0.5–2 cm; posterior rib naked for all its length; primary lateral veins obscure above, obscure below; minor veins distinct, fine, close, arising from the midrib only; "cross-veins" sometimes visible. INFLORESCENCES 1–4(5) per axil; peduncle 3.5–10 cm long, 1–3 mm diam., reddish, whitish or purplish, slender; **spathe** (4.5)5.8–11 cm long, (0.9–1.1)(2) times longer than peduncle), weakly constricted midway, 5 mm diam. at constriction, white throughout, becoming greenish (post-anthesis), rarely reddish (Gómez *et al.* 20473), cuspidate-acuminate at apex (the acumen ca. 1.5 mm long), white inside; resin canals appearing as intermittent lines in lower two-thirds of spathe, extending into upper part of spathe tube, drying blackened; spathe tube 3–4 cm long, red, pink, or purple at base inside; **spadix** sessile; slender, acute at apex, 4.5–7.7 cm long, broadest at the middle of the staminate portion, constricted scarcely or not at all above the sterile staminate portion; pistillate portion white, 2.5 cm long, 4.5 mm diam. midway, 3.5–4.5 mm diam. at apex and at base; staminate portion 3–5.2 cm long; sterile staminate portion cream, 3–4 mm diam.; pistils white; ovary 4–5-locular, with  $\pm$  sub-basal placentation; locules 0.4 mm long, 0.2–0.3 mm diam.; ovule sac 0.4 mm long; ovules 1 per locule, contained within translucent or transparent ovule sac, 0.2–0.3 mm long; funicle 0.1–0.2 mm long (can be pulled free to base), style similar to style type B; style apex flat; stigma subdiscoid, unlobed,  $\pm$  truncate, 0.4–0.5 mm diam., 0.1 mm high, covering center of style apex; the androecium margins irregularly 4–6-sided, 0.6–1.2 mm long. INFLORESCENCE peduncle, 8 cm long, 1 cm diam.

Flowering in *Philodendron sulcicaule* occurs during the dry season and early wet season (December through May, though no flowering collections have been seen from April). Post-anthesis collections have been seen from March and April. No fruits have been seen.

*Philodendron sulcicaule* ranges from extreme southeastern Costa Rica to the Atlantic slope of

Panama, from 100 to about 700 m elevation (though generally less than 300 m) in *Premontane wet forest* transition to *Tropical wet forest* life zones.

*Philodendron sulcicaule* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Ovata*. The species is characterized by its scandent habit; deeply sulcate stems drying yellow and glossy; unribbed, deciduous cataphylls; subterete petioles weakly flattened adaxially; and ovate-cordate blades with obscure primary lateral veins and up to four pairs of weak basal veins. Also characteristic are the numerous (up to five per axil), small, slenderly pedunculate inflorescences with spathes white outside (sometimes tinged red) and red to purplish in the base of the tube.

*Philodendron sulcicaule* appears closest to an undescribed species from Bajo Calima, Valle Department, Colombia (e.g., *Croat* 70137). Though the latter species has the same habit, petiole shape, leaf size and shape, and ribbed yellowish stems, it differs in having white cataphylls, usually 3–4 primary lateral leaf veins, and 3–5 basal veins with a well-developed posterior rib. In addition, the spathe tube is reddish outside and the entire inner surface is creamy-white (vs. usually white outside and reddish to purplish inside the tube in *P. sulcicaule*).

In Central America, *Philodendron sulcicaule* resembles *P. purpureoviride* and *P. microstictum*, both of which are vines with similar leaves. *Philodendron purpureoviride* differs in having solitary, stout inflorescences (instead with up to three or four smaller inflorescences) and leaf blades with distinct primary lateral veins. *Philodendron microstictum* differs in having blades broader than long (vs. longer than broad in *P. sulcicaule*).

*Additional specimens examined.* PANAMA. **Bocas del Toro:** Fortuna Dam area, Chiriquí Grande–Fortuna, 3.2 mi. N of Continental Divide, 700 m, 8°45'N, 82°15'W, *Croat* & *Gayman* 60249 (MO). **Coelá:** Bismarck, 2000–3000 ft., *Williams* 628 (NY); 16.7 km N of turnoff to Colesito from Llano Grande, 700 ft., *Hammel* 1791 (MO). **Colón:** Santa Rita Ridge Road, ca. 22 km from Transistmian Highway, 500 m, 9°25'N, 79°40'W, *Hammel et al.* 14504 (B, CAS, K, MO, PMA, US); E Santa Rita Ridge, *Correa* 670 (MO, PMA); Río Iguañita, ca. 1 mi. upstream, *Kennedy & Dressler* 3481 (F); Sabanitas–Portobello, Río Piedras Lumber Road, 250 m, 9°22'30"N, 79°41'30"W, *Croat* 75167 (CM, MO, PMA, US); Río Guanche, 30–100 m, *Croat* 79344 (PMA, MO). **Panamá:** El Llano–Cartí, 12–16 km N of Pan-American Highway, 150–400 m, *Kennedy et al.* 3162 (MO); Mile 10, 330 m, *Croat* 33734 (CM, F, MO); Mile 12, 200–500 m, *Liesner* 1154 (MO, US); ca. 4.6 mi. N, ca. 350 m, 9°15'N, 79°W, *McPherson & Merello* 8163 (B, F, MO); Mile 4, 500 m, *Croat* 49140 (MO). **San Blas:** El Llano–Cartí road, Mile 12, ca. 1000 ft., *Antonio* 3781 (MO); Mile 14, 300 m, 9°15'N, 79°W, *Croat* 69246 (CAS, CM, CR, L, MO, NY); Mile 9, 350 m, 9°20'N,

79°W, *Croat & Zhu* 76565 (MO). **San Blas-Panamá:** Kuna Divide trail, W of El Llano-Carti Road, 250 m, 9°20'N, 79°W, *McPherson* 11868 (MO); Cordillera de San Blas, Pacific slope headwaters of Río Piria, 200–400 m, 9°12'N, 78°16'W, *H. Herrera et al.* 1143 (K, MO, PMA, US).

***Philodendron tenue*** K. Koch & Augustin, in A. Braun et al., *Append. gen. sp. Hort. berol.* 1854: 7. 1854–1855. TYPE: Cultivated at Berlin, origin unknown (holotype, B? lost); Schott ic. 2465 and 2466 (neotype, here designated, W). Figures 404–407, 413, 414.

*Philodendron gracile* Schott, *Prod. syst. Aroid.* 244. 1860. TYPE: Costa Rica. *Wendland s.n.*, not seen (holotype, W? now lost; impression of type is at K); Schott ic. 2687–2688 (neotype, here designated).

*Philodendron ecuadorensis* Engl., *Bot. Jahrb. Syst.* 26: 531. 1899. TYPE: Ecuador. *Manabi, Eggers* 15221 (holotype, B; isotype, F).

*Philodendron sodiroanum* Engl., *Bot. Jahrb. Syst.* 26: 531. 1899. TYPE: Ecuador. *Pichincha: Nanegal, Sodiro* 47 (holotype, B).

Usually hemiepiphytic, rarely terrestrial or epilithic; stem usually appressed-climbing, thick, short, leaf scars inconspicuous; internodes striate, 2–3.5 cm long, 1–2.5 cm diam., about as long as broad, dark green to olive-green; roots drying tan to brownish red to blackish, smooth, elongate, 7–28 cm long, 0.2–0.5 cm diam., usually few per node, rarely numerous per node, dense, spreading at the nodes, epidermis peeling; cataphylls membranous, 28–31 cm long, 8 cm broad at base, usually unribbed, sometimes bluntly or sharply 1-ribbed to sharply 2-ribbed, green, red or purplish green, pale white-striate, drying brownish to reddish brown, persisting intact, eventually fibrous. LEAVES ± erect; petioles 29–107 cm long, 2–11 mm diam., erect, ± terete, stiff, firm, dark green, sometimes somewhat flattened with a low medial rib adaxially, surface weakly semiglossy, weakly light green striate with many small, fine ridges; geniculum thicker than petiole, 3–4 cm long, scurfy, slightly paler than petiole; blades ovate to ovate-triangular with prominent posterior lobes, chartaceous to subcoriaceous, moderately bicolorous, narrowly acuminate, sometimes long-acuminate at apex (the acumen inrolled, 1–4 mm long), cordate at base, 31–78 cm long, 16–52 cm wide (1.2–3.1 times longer than wide), (0.6–1.4 times longer than petiole), margins weakly undulate, upper surface gray-green to dark green, glossy, lower surface semiglossy, paler; anterior lobe 30.5–60 cm long, 14–34 cm wide (1.5–3.8 times longer than posterior lobes), broadest at point of petiole attachment; posterior lobes 10–29.5 cm long, 6–16.5 cm wide, broadest at or near the middle, directed downward

and outward, sometimes inward, obtuse to subrounded to bluntly acute; sinus triangular, acute to almost obtuse, sometimes closed, 4.5–17 cm deep; midrib flat to sunken, concolorous to slightly paler than surface above, raised, concolorous, drying somewhat paler than surface below; basal veins 5–7 per side, with 0(1) free to base, part of the remainder coalesced 2–10.5 cm; posterior rib not naked; primary lateral veins (6)8–14(20) per side, departing midrib at a 65–90° angle, ± straight or weakly arcuate to the margins, prominently to narrowly sunken above, raised to convex below; interprimary veins as conspicuous as primary lateral veins, weakly sunken and concolorous above, weakly raised and concolorous below; minor veins moderately visible to distinct below, arising from both the midrib and primary lateral veins. INFLORESCENCES erect, to 4 per axil; peduncle 2–11 cm long, 3–18 mm diam., pale green, drying reddish brown, heavily white-lineate, drying longitudinally fissured; spathe erect to erect-spreading, coriaceous, acuminate at apex, 7–14 cm long, (1.3–4.4 times longer than peduncle), moderately constricted above the tube, 2.1 cm diam. at constriction, short-lineate throughout; spathe blade weakly lanceolate, greenish white, sometimes heavily tinged red outside, 4–8 cm long, 2.1 cm diam., (opening 3.6 cm long), white to pale green inside; spathe tube ovoid, medium to dark green, usually tinged red or sometimes dark violet outside, white-striate with clear margins in front, short, pale green lineate in back outside, 3–7 cm long, 2–3.5 cm diam., white to pale green, sometimes red at base inside, sometimes sparsely white-spotted; spadix sessile; 6–12 cm long; pistillate portion yellowish green to pale green, weakly tapered upward, 2.8–4 cm long in front, 1.9–2.2 cm long in back, 1.1–1.6 cm diam. at middle, 1.75 cm wide at base, 0.9–1.5 cm ca. 1 cm from apex, with 12–14 flowers per spiral; staminate portion 5.3–7.1 cm long; fertile staminate portion white, drying orange reddish to light reddish, 1–1.1 cm diam. at base, 1.2–1.5 cm diam. at middle, 7–10 mm diam. ca. 1 cm from apex, broadest in the middle, as broad as to slightly broader than the pistillate portion, as broad as to narrower than the sterile portion, 27–35 flowers visible per spiral; sterile staminate portion as broad as to broader than the pistillate portion, pale brown, 11–17 mm diam.; pistils 2.7–3.1(6.2) mm long, 1.2–1.7(3.9) mm diam.; ovary 4–5-locular, 1.6–2(5.9) mm long, 1.2–1.7(3.9) mm diam., with axile placentation; locules 1.6–2(4.2–5.9) mm long, 0.4–0.7(1.1) mm diam.; ovules 12–14 per locule, 2-seriate, 0.4 mm long, longer than funicle; funicle 0.2(0.5) mm long, adnate to lower part of partition,

style 0.5–0.6 mm long, 1.2–1.7 mm diam., similar to style type B; style apex flat; stigma cupulate,  $\pm$  hemispheroid, 1.1 mm diam., 0.4–0.6 mm high, covering center of or entire style apex, depressed medially with 6–8 holes; the androecium truncate, prismatic, margins irregularly 4–6-sided, (0.3)0.7–0.9 mm long, 1.1–1.4 mm diam. at apex; thecae oblong, 0.3–0.4 mm wide,  $\pm$  parallel to one another, nearly contiguous; sterile staminate flowers irregularly 4–6-sided, mostly prismatic, sometimes clavate, blunt, 1.3–2.5 mm long, 1.4–2.5 mm wide. **INFRUCTESCENCE** with fruits purplish.

Flowering in *Philodendron tenue* occurs during the dry season and early rainy season (January through August, perhaps especially April and May). Post-anthesis collections are from January to October, and immature fruits are known from April through November.

*Philodendron tenue* ranges from Nicaragua to southern Ecuador (El Oro) on the Pacific slope, and in Venezuela along the foothills of the Sierra de Perijá (Zulia), through the Cordillera de la Costa (Yaracuy to the Distrito Federal at Cerro Naiguatá) and in the foothills of the Cordillera de Mérida south to Apure. In Central America it ranges from 20 to 1400 m in elevation in *Premontane wet forest*, rarely in *Tropical moist forest*. In Colombia it has been collected to 2300 m, and in Ecuador to 1930 m. *Philodendron tenue* is ecologically quite versatile in South America, where it occurs in *Tropical thorn woodland*, *Premontane thorn woodland*, *Tropical dry forest*, *Tropical moist forest*, and *Premontane wet forest* life zones.

*Philodendron tenue* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Fibrosa*. This species is characterized by its epiphytic habit, short internodes, persistent cataphyll fibers, more or less terete petioles (equal to or longer than the blades), and especially by its ovate to ovate-triangular blades with prominent posterior lobes often directed outward, narrow to almost closed V-shaped sinus with the posterior ribs not at all naked, and anterior lobe with numerous close primary lateral veins.

*Philodendron tenue* is similar and perhaps related to *P. maguirei* G. S. Bunting, but that species is terrestrial, has fewer primary lateral veins (typically 3–6 vs. 8–20 for *P. tenue*), and petioles drying straw-yellow (vs. green to brown for *P. tenue*).

The species can be confused with narrow-leaved epiphytic forms of *P. thalassicum* in Costa Rica, but that species has a somewhat glaucous epidermis on the abaxial blade surface and also uniformly greenish spathes.

It is strange that *P. tenue* is found on both sides of the Andes (an unusual pattern except for the most weedy of species), but still does not extend southward from Venezuela. It should certainly be expected at least in Amazonian Colombia.

**Additional specimens examined.** COSTA RICA. **Alajuela:** Cañas–Upala, 4 km NNE of Bijagua, ca. 400 m, *Croat 36000* (MO); 36294 (MO); 13.8 km N of Bijagua, 100–150 m, 36450 (MO); 8.5 km NE of Villa Quesada, ca. 600 m, *Croat 46969* (MO); 7.5 km N of Río Balsas, 700–800 m, ca. 10°10'–15'N, 84°30'–35'W, *Stevens 13861* (MO); 17 km NW of San Ramón, 785 m, 10°14.15'N, 84°33'W, *Croat 68161* (CR, F, M, MO); 3.5–4 mi. W of center of San Ramón, 800 m, 46778 (MO); Río San Rafael, W of La Marina, 500 m, 10°23'N, 84°23'W, *Burger & Stolze 5003* (CR, F, GH, NY, US); 5013 (CR, GH, US); San Carlos–San Pedro, ca. 150 m, *Horich s.n.* (M). **Cartago:** Río Gato, 12 km S of Turrialba by air, 4 km SE of Pejibaye, 700 m, 9°48'N, 83°42'W, *Liesner 14360* (CR, MO); Río Reventazón valley, 3 km SE of Turrialba, 500–600 m, *Holm & Itis 190* (A, BM, G, GH, U); 525–600 m, 9°54'N, 83°39'W, *Liesner et al. 15340* (CR, MO, WIS); Cartago, 4250 ft., *J. D. Smith 5964* (US). **Guanacaste:** Lake Arenal, 550–850 m, 10°27'N, 84°50'W, *Hammel et al. 15140* (MO); Continental Divide, La Chirripa Ridge, 4 km NE El Dos de Tilarán, 1000 m, 10°25'N, 84°53'W, *Haber et al. 4859* (MO); Río Chiquito, Tilarán–Arenal, Zona Monteverde, 700 m, 10°23'N, 84°51'W, *Haber & Bello 8241* (CR, MO); Río Aguas Verdes, 600 m, *Rivera & Dennis 1050* (CR, INB, MO); Río Negro, 1100 m, *Rivera 718* (CR, INB, MO). **Heredia:** vic. Porto Viejo, near Río Sucio, 20 m, *Croat 35691* (F, MO); Río Peje–Río Sardinalito, Volcán Barva, 700–950 m, 10°17'30"N, 84°04'30"W, *Grayum 6924* (MO); 6886 (CR, MO); La Selva Field Station, ca. 100 m, *Grayum 2626* (DUKE); 2901 (DUKE); Parque Nacional Braulio Carrillo, above Río Sucio, 5–600 m, *Pennington et al. 11536* (K). **Limón:** Guápiles, 300–500 m, *Standley 37415* (US); 37451 (US); Hacienda Tapezco–Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, *Davidson & Donahue 83664* (RSA); 8742 (MO, RSA); W of Guápiles, Quebrada Danta, 360 m, 10°12'N, 83°49'W, *Croat 68427* (MO); Finca Castilla, 30 m, *Dodge & Goerger 9270* (MO); Cerro Coronel, 20–170 m, 10°41'N, 83°38'W, *Stevens 23637* (CR, MO); 24462 (CR, MO); Pococi, Sardinas, 15–20 m, *Araya 510* (CR, INB). **Puntarenas:** Osa Peninsula, 500 m, *G. Herrera 4013* (CR, INB, MO); Chacarita–Rincón de Osa, 10 km W of Chacarita, ca. 100 m, 8°45'N, 83°18'W, *Croat & Grayum 59893* (CR, MO); Piedras Blancas, 3.7 mi. W of Pan-American Highway, 90–105 m, 8°46'N, 83°18'W, *Croat 67698* (CAS, CR, MO); vic. Golfito, 100 m, 8°40'20"N, 83°12'10"W, *Hammel 18399* (CR, MO); 6 km S of San Vito de Java, 3500 ft., *Raven 21882* (BM, CR, DS, F); 4000 ft., 22009 (CR, DS, F, WIS). **San José:** Without locality, *Yielding 235* (CR, F). **HONDURAS. Gracias a Dios:** Ahuas Bila, 200 km SW of Puerto Lempira, 100 m, *Nelson & Cruz 9293* (MO, TEFH, UNAH). **Olancho:** Poncaya, *Blackmore & Heath 1994* (BM); Río Olancho, 19 mi. NE of San Esteban, 550 m, 15°29'N, 85°43'W, *Croat & Hannos 64475* (B, MO). **NICARAGUA. Boaco:** Cerro Mombachito, 700–1000 m, *Grijalva & Araquistain 80* (MO). **Chontales:** La Libertad, 500–700 m, *Standley 9056* (F); 4 km NW of Santo Domingo, 280 m, ca. 85°06'W.



12°17'N, *Grijalva* 3806 (MO). **Jimotea:** La Salvadora-La Palestina, Hwy. 3, 1100–1150 m, ca. 13°04'–05'N, 85°53'–54'W, *Stevens & Grijalva* 15369 (MO); Río Bocay, Salto Kayaska, 190–340 m, ca. 13°51'N, 85°22'W, *Stevens et al.* 16464 (CAS, MO); Caño Litutus, ca. 175 m, ca. 13°58'N, 85°21'W, *Stevens et al.* 16665 (MO). **Matagalpa:** NW of Cerro Musún, trail to Palwás, 800–1200 m, *Araquistain & Moreno* 2535 (MO); 300–600 m, 2470 (MO); trail to Paylo, 500–800 m, 2568 (K, MO); Macizos de Peñas Blancas, 1000–1400 m, ca. 13°14'–15'N, 85°38'–39'W, *Stevens et al.* 21074 (MO); Cerro Musún, Río Bilampí, 500–800 m, 13°N, 85°14'W, *Araquistain & Moreno* 2351 (MO); NW of Cerro Musún, 4 km SW of Wanawás, 200–500 m, 13°00'–01'N, 85°14'W, 2613 (HNMN, MO, NY); Cerro Musún, above Salto Grande de Quebrada Negra, 500 m, *Neill* 1817 (MO). **Río San Juan:** El Castillo-Caño de Oro, Gigante, 10 m, 10°43'15"N, 84°54'30"W, *Martínez* 2191 (MO); Caño Chontaleño, 20 km NE of El Castillo, 200 m, *Neill & Vincelli* 3608 (BM, CAS, MO). **Rivas:** Isla Ometepe, Volcán Maderas, 600–1000 m, 11°26'–27'N, 83°31'–32'W, *Robledo* 1674 (ENCB, MO); 800–1000 m, 11°26'N, 85°29'W, 2065 (MO). **Zelaya:** El Cerro La Pimienta, ca. 800–1000 m, 13°44'50"N, 84°59'45"W, *Grijalva* 396 (CM, HNMN, MO); "Kurinwacito," 80–100 m, 13°08'N, 84°55'W, *Moreno* 23753 (MO); 2 km N of Kuikainita, S of Siuna, 100 m, *Neill* 4150 (MO); Cerro Livico, 7 km NE of Siuna, 500 m, *Neill* 3630 (MO); ca. 13 km above Kururia, road to San Jerónimo, <200 m, *Pipoly* 3831 (MO); Cerro Baká, ca. 6.5 km E of Río Coperna, 200–300 m, 13°40'N, 84°30'W, *Pipoly* 4918 (MO); Cerro La Pimienta number 1, ca. 900–980 m, ca. 13°45'N, 84°59'W, *Pipoly* 5118 (MO); ca. 6.3 km S of Colonia Yolaina, ca. 200–300 m, 11°36'–37'N, 84°22'W, *Stevens* 6419 (MO); along Caño Majagua, ca. 750–850 m, ca. 13°45'N, 85°00'–01'W, *Stevens* 6881 (BM, LL, MO); Cerro Saslaya-San José del Hormiguero, ca. 200–400 m, 13°44'–45'N, 84°57'–58'W, *Stevens* 7076 (MO); road to Mina Nueva América, ca. 14.3 km N of El Empalme, *Stevens* 8435 (MO); Colonia San José-Tomás Mejía Caño, ca. 40–150 m, ca. 11°53'–54'N, 84°20'W, *Stevens* 9011 (BM, MO); Bonanza-Constancia, 160–360 m, ca. 13°58'–14°01'N, 84°37'–40'W, *Stevens* 12466 (MO); ca. 1.5 km NE of Las Esperanza de Las Quebradas, 300–350 m, ca. 13°38'N, 85°02'W, *Stevens & Moreno* 19353 (MO); Río Mico, El Recreo, ca. 30 m, *Standley* 19749 (F). **PANAMA. Bocas del Toro:** Campamento Changuinola 1, 20 m, *Correa et al.* 4067 (MO, PMA). **Coeló:** La Pintada-Cocleito, 5.3 mi. N of Llano Grande, 513 m, *Croat* 49240 (MO); 5.6 mi. N of Llano Grande, 1.4 mi. N of Continental Divide, 150 m, 8°46'N, 80°27'W, *Croat* 67478 (CM, L, MEXU, MO, PMA); El Valle de Anton, behind Club Campestre, *Croat* 14278 (MO). **Darién:** Parque Nacional Cerro Pirre, Alturas de Nique region, near Cana Mine, 500 m, 7°45'N, 77°40'W, *McPherson* 11584 (MO); Cerro Pirre region, Cana, 480 m, *Croat* 38009 (MO, RSA); 550–760 m, 7°57'N, 77°46'W, 68871 (MO); 68954 (CAS, MO, NY); *Folsom* 4417 (MO, PMA); near station along Río Perisénico, 110 m, 8°01'N, 77°44'W, *Croat & Zhu* 77107 (MO); Río Paracida, 0–40 m, 8°N, 77°48'W, *Croat* 68998 (MO); ca. 13 km E of Pucuro, 450–600 m, 8°03'N, 77°20'W, *Hammel et al.* 16195 (MO, US). **Panamá:** El Llano-Cartí, 6.8 mi. N, 350 m, *Croat* 49111 (MO).

***Philodendron thalassicum* Croat & Grayum, sp. nov.** TYPE: Panama. Chiriquí: Fortuna Dam area, trail to meteorological station of Río Hornito, ca. 0.5 km S of Centro de Científicos, 1010–1030 m, 8°45'N, 82°18'W, 23 June 1994, *Croat & Zhu* 76297 (holotype, MO-4619518-22; isotypes, AAU, B, CAS, CM, COL, CR, DUKE, EAP, F, GB, GH, HUA, IBE, K, LE, M, MEXU, NY, P, PMA, QCA, RSA, S, SCZ, SEL, TEX, U, US, VEN). Figures 408–410, 415.

Planta terrestris aut hemiepiphytica; internodia brevia longa, 2–4 cm diam.; cataphylla 14–46 cm longa, acuta 2-costata, persistentia ut fibra; petiolus D-formatus, 38–109 cm longus, 2–14 mm diam., plerumque longior quam laminae; lamina triangulari-sagittata vel ovata, 42–77 cm longa, 16–52 cm lata, atriviridis supra, saepe venata infra; inflorescentia 1–3; pedunculosa 3–14 cm longus, 2–9 mm diam.; spathe 11–20 cm longa; lamina spatheae extus alba vel creamea, intus viridialba; tubo spatheae 3–7 cm longo, extus pallide viridi, intus virello vel albo; pistilla (4)5–7(8)-locularia; loculi 16–22-ovulati.

Terrestrial or hemiepiphytic, growing to 4 m high in trees; stem appressed-climbing; internodes short, 2–4 cm diam., dark green, semiglossy, drying light yellowish brown, epidermis fragmented, reddish brown; roots several per node, <30 cm long, drying dark brown with epidermis ridged and splitting; cataphylls soft, 14–46 cm long, sharply 2-ribbed (ribs low), green or tinged pink when young to reddish brown or brown, drying brownish, persisting semi-intact, soon dilacerating; petioles 38–109 cm long, 2–14 mm diam., D-shaped, firm, flexible, sometimes with low, medial rib adaxially and with adaxial margins obtuse to rounded, medium green, weakly glossy, faintly to conspicuously dark- or striate-lineate, drying yellow-brown, dark brown to sometimes blackened; blades triangular-sagittate to ovate-triangular, subcoriaceous, conspicuously bicolorous, acuminate to long-acuminate at apex, sagittate at base, 42–77 cm long, 16–52 cm wide (0.8–1.8 times longer than wide), (0.6–1.4 times longer than petiole), usually shorter than petiole (averaging 0.9 times as long), upper surface dark green, matte or weakly glossy, sometimes glistening, drying dark brown, lower surface much paler (forming an areolate pattern on high magnification) and matte, often bluish green to whitish, drying bluish green if triangular-sagittate or yellowish brown if ovate-triangular, glaucous; anterior lobe 28–58 cm long, 18–51 cm wide (1.6–3.3 times longer than posterior lobes); posterior lobes broadly obtuse to narrowly rounded, 13.5–25 cm long, 10–23 cm wide, rounded; sinus spatulate to parabolic, 9–12 cm deep; midrib broadly sunken to flat above, slightly paler, obtusely triangular and darker below;

basal veins 5–11 per side, with 1–2(3) free to base, with a few veins coalesced for 5–9 cm; posterior rib prominently naked to 2.5–5 cm; primary lateral veins (4)12–18 per side, departing midrib at a 50–60° angle, weakly arcuate to ± straight to the margins, C-shaped to sunken and slightly paler above, convex and concolorous below; minor veins conspicuous, darker than surface to obscure below, arising from both the midrib and primary lateral veins. INFLORESCENCES ± erect, 1–3 per axil; peduncle 3–14 cm long, 2–9 mm diam., coarsely white-striate; **spathe** 11–20 cm long (1.15–2.8(4.7) times longer than peduncle); spathe blade white to cream and densely short-lineate outside, greenish white inside; spathe tube pale green and short-lineate outside, 3–7 cm long, greenish to white, sometimes faintly pinkish inside; **spadix** sessile to weakly stipitate; 12.8–16.6 cm long; pistillate portion 5.5 cm long in front, 3–3.5 cm long in back, 1.6–2 cm diam. midway, 1.5–1.6 cm diam. ca. 1 cm from apex; staminate portion 6.8–9.4 cm long; fertile staminate portion tapered toward apex, constricted (to 8–10 mm, 1.5–2 cm above base of sterile section) above sterile male flowers, 1 cm diam. at base, 1–1.6 cm diam. at middle, 5–7 mm diam. near apex; sterile staminate portion 1–1.7 cm diam. at base, 1 cm diam. at apex; pistils 2.8–3.8(6.2) mm long, 1.2 mm diam.; ovary (4)5–7(8)-locular, 1.7–2.3 mm diam., with axile placentation, wall sometimes embedded with angular crystal-like particles; locules 2.7(4.9) mm long, 0.5–0.6 mm diam.; ovules ca. 16–22 per locule, 2-seriate, 0.2–0.4 mm long, longer than funicle; funicle 0.1–0.2 mm long, adnate to lower part of partition; stigma lobed; style 1.3–2.1 mm diam., similar to style type B; style apex sloping to somewhat rounded, drying granular with an acute, turned up margin; stylar pores thickened, usually fused with the crown, sometimes free or nearly so, each with its own fringe of stigmatic papillae; style boss broad, shallow to quite shallow; the androecium truncate, prismatic, oblong, margins irregularly 4–6-sided, 0.9–1.2 mm long; thecae oblong, ± parallel to one another; sterile staminate flowers prismatic to weakly clavate, irregularly 5–6-sided, 1.2–1.6 mm long. INFRUCTION with berries whitish (maturing).

Flowering in *Philodendron thalassicum* occurs throughout the dry season to the mid-rainy season (January through August), though no flowering collections are known from February (although post-anthesis collections do exist). Post-anthesis collections are known from April and May, and especially June, July, and August, and also from November (indicating that flowering may occur later than August). Immature fruiting collections are known from

January, February, June, September, and November.

*Philodendron thalassicum* ranges from central Costa Rica to western Panama (Chiriquí) at (775)1000 to 2100 m elevation in *Premontane rain forest*, *Tropical Lower Montane wet forest*, and *Tropical Lower Montane rain forest* life zones.

*Philodendron thalassicum* is a member of *P.* sect. *Philodendron* subsect. *Philodendron* ser. *Impolita*. This species is characterized by its short thick internodes, two-ribbed cataphylls persisting as fibers, more or less D-shaped petioles (averaging longer than the blades), internally greenish to white spathe tube, and especially by the pale matte, often bluish green lower leaf surfaces. The epithet "thalassicum" (from *thalassicus*, meaning sea green or bluish green) is derived from this coloration of the lower blade surface.

*Philodendron thalassicum* may be confused with *P. hebetatum* and *P. strictum*, which also have matte, much paler lower blade surfaces. Both differ in having subterete petioles drying conspicuously light yellow-brown, rather than sharply D-shaped and drying somewhat blackened as in *P. thalassicum*. In addition, both *P. hebetatum* and *P. strictum* have cataphylls drying yellowish and semi-intact and spathe tubes colored reddish, maroon, or violet-purple within.

Plants of *P. thalassicum* with sagittate blades might be confused with narrow-leaved plants of *P. jodavisanum* or *P. tenue*, but both of those species have blades that are semiglossy to glossy (not at all glaucous) below. More ovate blade-forms of *P. thalassicum* might be confused with *P. schottianum*, but the latter also usually has a semiglossy lower blade surface, cataphylls that persist semi-intact, and spathe tubes reddish internally.

*Philodendron thalassicum* is variable in blade shape, with some triangular-sagittate (e.g., Grayum et al. 3760, Croat 15876, 15745) and others ovate (e.g., Burger & Burger 7541, Burger & Stolze 5661, Burger & Liesner 6745). Those with ovate leaves sometimes dry brownish on the lower surface while those with the triangular-sagittate blades dry bluish green. No other differences between these two forms have been detected.

*Additional specimens examined.* COSTA RICA. Alajuela: Cantón Alfaro Ruiz, Tapasco de Zarcero, 1650 m, A. Smith NY1232 (F, NY); Alto Palomo, 1900 m, Lent 1844 (CR, F); Naranjo-Quesada, along Hwy. 15, 3.2 mi. N of Zapote, 1560 m, Croat 46908 (MO); Cantón San Ramón, R. B. Monteverde, Cordillera de Tilarán, valle del Río Peñas Blancas, 1600 m, 10°19'15"N, 84°46'30"W, Bello 5151 (CR, INB, MO, NY); Monteverde reserve, 10°20'N, 85°50'W, Croat 61191 (cultivated) (MO). Car-

**tago:** Moravia, 3.5 km from Finca Racine, 1200–1300 m, Croat 36634 (CR, MO); Moravia de Chirripó-Turrialba, 1400–1600 m, *I. Chacón* 269 (MO); Río Dos Amigos–Río Villegas, W of Río Grande de Orosi, ca. 1650 m, 9°42'N, 83°47'W, *Grayum et al.* 3760 (MO, RSA); Quebrada Honda–Río Sombrero, ca. 1–2 km above El Muñeco, ca. 1400 m, *Luteyn* 3238 (DUKE); ca. 20 km E of Río Pacuare, on road toward Moravia, 1150 m, 9°50'N, 83°24'W, *Thompson & Rawlins* 1234 (CM); Quebrada Cangreja, 3 km S of Pan-American Highway, 1620–1650 m, 9°46'N, 83°57'W, *Liesner & Judziowiec* 14498 (CR, MO); Tapantí Hydroelectric Reserve, ca. 1 km S of junction of Quebrada Salto and Río Grande de Orosi, 1500–1800 m, 9°43'N, 83°47'W, Croat 36211 (MO); 1500–1700 m, 36078 (MO); 1525–1595 m, 79049 (CR, INB, MO); Croat & Grayum 68290 (MO, US); *Grayum & French* 5821 (INB, MO); 1600 m, *Lent* 964 (CR, F, US); 1500 m, 9°42'N, 83°47'W, *Burger & Liesner* 6745 (F, US); 1600 m, 9°42'N, 83°47'W, *Burger & Stolze* 5661 (CR, DS, F, WS); 5,200 ft., *Utey & Utey* 7284 (MO); 1400–1700 m, 5184 (DUKE, MO); ca. 1600 m, 9°42'N, 83°46'W, *Burger & Burger* 7541 (F, MO); 1400–1600 m, 9°42'N, 83°47'W, *Baker & Utey* 201 (F, MO); Río Navarro, El Muñeco, 1400–1500 m, *Standley & Torres* 51371 (US). **Heredia:** 4 mi. N of Vara Blanca, 1350 m, Croat 35595 (MO); 35621 (MO); 9 km SE of San Ramón [de Sarapiquí], 1000 m, 10°16'N, 84°05'W, *Loiselle* 138 (MO); Paracito–Río Claro, vic. of Bajo La Honda, 1100–1400 m, Croat 44500 (MO); Parque Nacional Braulio Carrillo, San Rafael de Vara Blanca, 1830 m, 10°11'50"N, 84°06'35"W, *Herrera* 246 (MO); Sarapiquí, 1750 m, 10°12'47"N, 84°06'05"W, *Boyle & Boyle* 2695 (MO); Finca Montreal, between headwaters of Río Volcán and Río San Fernando, 1800 m, 10°12'39"N, 84°06'45"W, *Boyle* 1084 (CR, MO); Volcán Barva, Río Peje–Río Sardiná, Atlantic slope, 1200–1400 m, 10°15'30"N, 84°05'W, *Grayum & Herrera* 7846 (MO); Río San Rafael, 1500 m, 10°13'N, 84°05'W, *Grayum* 7050 (MO). **Limón:** Cordillera de Talamanca, Kamuk massif, 1900–2300 m, 9°14–15'N, 82°59'W, *Davidse & Herrera* 29201 (MO). **Puntarenas:** Las Cruces Tropical Botanical Garden, 6 km W of San Vito de Java, 1200 m, 8°49'N, 82°58'W, Croat 57264 (CR, MO); Río Coto Brus, near Cotoón, 23 km N of La Unión (on Panama border), Croat 26685 (CR, MO); Cerro Frantzius–Cerro Pittier, around Río Canasta, 9.5 airline km NW of Agua Caliente, 1500–1600 m, 9°02'N, 82°59'W, *Davidse et al.* 28365 (CR, MO, NY); Quebrada Kuisa, near crossing of Ujarrás–San José Cabécar trail, 2100 m, 9°20'30"N, 83°14'W, *Grayum* 10282 (CR, INB, MO); Monteverde Reserve, 1500 m, 10°17'N, 84°48'W, *Hammel et al.* 14206 (MO); 1500–1600 m, 10°18'N, 84°47'W, *Haber & Zuchowksi* 10020 (CR, INB, MO, MV); 1500–1620 m, *Pounds* 334 (CM, MO); 357 (MO); 1450–1650 m, *Burger & Baker* 9772 (F, MO); Cerros Centinelas, 1550–1600 m, 10°18'N, 84°47'W, *Grayum & Sleeper* 3857 (CR, MO). **San José:** trail beyond Bajo La Honda, towards Río Claro, 1100–1200 m, *Luteyn* 3318 (DUKE, MO); NE slope of Altos Tablazo, between Quebrada Tablazo and summit of ridge, 1700–1875 m, 9°50'N, 84°02'W, *Grayum & Schatz* 5181 (CR, MO); Parque Nacional Braulio Carrillo, Estación La Montura, *Gómez et al.* 20827 (CR, MO); General Valley, 1500 m, 9°27'N, 83°43'W, *Burger & Barringer* 11601 (F, MO); Cerro Honda, between Río Patria and Río Zarquí, 1500–1600 m, 10°04'N, 84°01'W, *Grayum & Sleeper* 6131 (CM, MO). **PANAMA.** **Chiriquí:** NW side of Cerro Pando, Croat 15966 (MO); Fortuna Lake area, along Continental Divide, 1200 m, 8°44'N, 81°17'W, Croat 74975 (L, MO, NY,

PMA, QCA); Río Hornito–Río Chiriquí, 1050–1100 m, ca. 8°44'N, 82°13'W, *Stevens* 18371 (MO); Boquete region, Monte Rey, Croat 15876 (MO), 15745 (MO); SW slope of Cerro Pate Macho, 1630–1780 m, 8°46'N, 82°25'W, Croat 66401 (CAS, F, K, MBM, MO, TEX, W); Quiel road, 10 km above Boquete, 5500 ft., *Proctor* 31831 (LJ); Río Palo Alto–Cerro Pate Macho, ca. 6 km NE of Boquete, 1600–1700 m, 8°48'N, 82°23.5'W, *Grayum et al.* 6359 (MO, PMA, US); Cerro Horqueta, ca. 1650 m, Croat 270004 (MO).

***Philodendron tripartitum*** (Jacq.) Schott, Wiener Z. Kunst 1829: 780. 1829. *Arum tripartitum* Jacq., Pl. hort. schoenbr. 2: 33, t. 190. 1797. *Caladium tripartitum* (Jacq.) Willd., Sp. pl. 4: 491. 1805. TYPE: t. 190 in Jacq., Pl. hort. schoenbr. 1797 (holotype). Figures 411, 412, 416–424.

*Philodendron fenalii* Engl., in Mart., Fl. Bras. 3(2): 144. 1878. TYPE: Mexico (holotype, B? lost). Schott ic. 2599 (neotype, here designated, W).

*Philodendron affine* Hemsl., Diagn. Pl. Nov. Mexic. 37. 1879. TYPE: Guatemala. Barranca Honda, Volcán de Fuego, Salván s.n. (holotype, K).

*Philodendron tripartitum* var. *tricuspidatum* Engl., in Mart., Fl. Bras. 3(2): 144. 1878. TYPE: Costa Rica. San José: 1300 m, May 1857, C. Hoffmann s.n. (B? not seen).

Hemiepiphytic; appressed-climbing, stem often several meters long, sap watery, very aromatic-scented; internodes moderately elongate except near the apex, semiglossy, drying coarsely and irregularly ribbed (the ribs usually acute), 3–14 cm long, 1–2 cm diam. (to 3.5 cm diam. in Colombia), green to dark green to gray-green, soon drying light brown, epidermis sometimes loose and papery, roots usually 3–6 per node, usually less than 20 cm long, drying brown, less than 3 mm diam., semiglossy, sparsely scaly; cataphylls thin, (10)18–33 cm long usually unribbed, sometimes sharply 1-ribbed, greenish, sometimes tinged reddish, deciduous intact. LEAVES erect-spreading to spreading; **petioles** 20–61 cm long, 6–15 mm diam., terete to subterete, moderately spongy, sometimes obtusely somewhat flattened, sometimes with an obtuse medial rib, dark green, surface glossy to weakly glossy; **blades** deeply 3-lobed almost to the base or rarely trisect, ca. as broad as long in outline, thinly coriaceous, semiglossy, moderately bicolorous, acuminate at apex, lower surface sometimes heavily tinged purplish or reddish; typically rather dissimilar, median segment usually oblanceolate, sometimes nearly elliptic, rarely oblong to linear, 15–45 cm long, (4.5)7–14(19) cm wide, 1.9–7.8 times longer than wide (averaging 4.3 times longer than wide), ((0.9)1.5(1.7) times longer than lateral segments); the lateral segments conspicuously in-

quilateral, arcuate-spreading toward apex, (12)18–35 cm long, 1.3–15 cm wide, the inner margin always much narrower than the outer margin and weakly confluent with medial lobe, usually to 1 cm, rarely to 2 cm, the outer margin 1.5–5 times wider than the inner margin where the difference is most severe; midrib broadly convex, concolorous above, convex to round-raised, darker below; basal veins lacking; primary lateral veins (4)6–10(12) per side, departing midrib to a 25° angle on narrow blades or to 90° angle on broader blades, weakly curved on narrow blades or markedly curved on broader blades to the margins, sunken and usually concolorous above, convex and darker than surface below; minor veins fine and conspicuously visible, arising from both the midrib and primary lateral veins. INFLORESCENCES usually solitary, sometimes 2–3 per axil; peduncle 3–16 cm long, 6–13 cm diam., medium green, moderately recurved with reddish ring around the apex; spathe (8)13–21 cm long, 1.2–4 cm diam., weakly constricted above the tube, to 2.5 cm diam. at constriction, usually medium green to yellow-green, rarely creamy white, with dark purple dots (most to 3 mm circumference) heavily so on tube, less so on blade throughout, white to creamy white or greenish white within, cucullate at apex, convolute in lower 3/4 of tube at base, margins whitish or colorless, recurved at anthesis; spathe blade glossy, smooth outside, (the opening 7.5–8.5 cm long, 3.4–5.2 cm wide), creamy white, glossy; resin canals intermittent, orange, appearing medially and extending into spathe tube inside; spathe tube oblong, glossy outside, 5 cm long, 3 cm diam., maroon, deep red, or purplish-violet inside; spadix sessile or shortly stipitate (to 2 mm); clavate, ± erect, 12.3 cm long, broadest above the middle, weakly constricted between sterile and fertile portions; pistillate portion greenish white to pale yellowish green, 2.5–4.5(6.2) cm long, 1–1.8 cm diam. throughout; staminate portion 7.5–8.5 cm long; fertile staminate portion creamy white, 1.2 cm diam. at base, 1.3 cm diam. at middle, 1 cm diam. ca. 1 cm from apex, broadest in the middle, usually narrower than the pistillate and sterile portions; sterile staminate portion usually broader than the pistillate portion, creamy white, drying yellowish, 1.3 mm diam.; pistils 1.7–2.5(3.4) mm long, 0.6–1.1(2.3) mm diam.; ovary 6–8(10)-locular, 0.8–2.3 mm long, 1.1–2.3 mm diam., with sub-basal placentation; locules 0.8–1.1(2.3) mm long, 0.2–0.4(0.7) mm diam.; ovule sac 0.7–0.9(1.2) mm long; ovules 1–2 per locule, contained within a translucent and gelatinous matrix, 0.4–0.6 mm long, longer than funicle; funicle 0.2–0.4 mm long (can be pulled free to base), style 0.4–

0.8 mm long, 1–1.4(2.5) mm diam., usually similar to style type D (rarely B or E); style apex ± rounded; stigma subdiscoid, truncate, 0.7 mm diam., 0.2–0.3 mm high, covering center of style apex, inserted on style boss; the androecium truncate, prismatic, margins usually irregularly 5-sided, 1.4 mm long, 2.2 mm diam. at apex; thecae oblong, 0.5 mm wide, ± parallel to one another; sterile staminate flowers blunt, prismatic, irregularly 4–6-sided, 1.9 mm long, 1.5–2 mm wide. Berries whitish, irregularly angular, 4 mm long, 2–2.7 mm diam.; seeds usually 6–8 per berry, cylindrical, 1.5 mm long, sticky. JUVENILE plants with blades maroon below.

Flowering in *Philodendron tripartitum* occurs throughout the dry season and the first half of the rainy season (mostly May through July), with seemingly little phenological variation throughout its range. Most flowering collections have been made from January through July with a few from August and also November. Post-anthesis inflorescences have been collected throughout the whole year with the greatest concentration from May to July. Immature fruits are known from January, April, July, August, September, and November, while mature fruits are known only from October.

*Philodendron tripartitum* ranges from Mexico to Ecuador from near sea level to 1300(1500) m elevation. In Central America, this species ranges along the Caribbean slope in Mexico (Veracruz to Tabasco) and throughout Middle America, but occurs on both slopes in Costa Rica and Panama thence along the Pacific slope into lowland Antioquia and Chocó Departments, Colombia. In Mexico, *Philodendron tripartitum* occurs in Selva Alta Perennifolia and the transition forest to Bosque Caducifolia. In Costa Rica and Panama, it occurs in Tropical moist forest to Premontane wet forest, Tropical wet forest, and Premontane rain forest. In Colombia, it is known from Tropical wet forest and Tropical rain forest.

*Philodendron tripartitum* is a member of *P.* sect. *Tritomophyllum*. This species is characterized by its elongate internodes, deeply trilobate leaf blades only weakly confluent at the base or sometimes completely trisect, and spathes scarcely constricted above the tube.

*Philodendron tripartitum* is most likely confused with the common *P. anisotomum*. See under the latter species for the differences. It is probably most closely related to *P. costobrusense* from Costa Rica, which differs in having elliptic medial lobes with 18 or more pairs of primary lateral veins. *Philodendron tripartitum* may be confused also with *P.*

*madronoense* from Panama. That species also differs in having elliptic medial lobes with about 25 pair of primary lateral veins.

Along with *P. sagittifolium*, *P. tripartitum* is one of the most highly variable and widespread species in the genus, with complex patterns of variation in the shape of the lobes, the degree to which the lobes are confluent, the number of primary lateral veins, and the size and number of inflorescences per axil. Clinal variations range both north and south throughout Mexico and Central America and also elevationally. Most Mexican specimens have the lobes conspicuously confluent at the base with the lateral lobes moderately to prominently auriculate, and all have one inflorescence per axil. This form is well characterized by Engler's drawings of Araceae Drawing no. 48 of the synonym, *P. fendlii*. In Mexico, *P. tripartitum* ranges from sea level to usually about 900 m elevation.

To the south and east in Central America, especially at lower elevations, the blades of *P. tripartitum* dry paler green and are more weakly confluent at the base or completely trisect with the lateral lobes only weakly auriculate. This form is encountered in eastern Chiapas and Tabasco and ranges through Belize and the remainder of Caribbean coastal Central America, as well as on the Pacific coast of El Salvador. It reaches its most deeply trisect and pale green expression in *Tropical Moist forest* along the Caribbean coast of Panama, but also at lower elevations along the Pacific slope in Darién Province. In contrast to material from Mexico and Belize, this form of the species generally has but a single inflorescence per axil, though in Panama it may have from one to three inflorescences per axil. Noteworthy collections of this form are Dressler 1513 and Breedlove 10330 from eastern Chiapas, which have blades drying pale yellow-green and weakly confluent with only weakly auriculate lateral lobes, yet occur at between 950 and 1266 m (3800 ft.) elevation.

Most collections of *P. tripartitum* from above 500 m and from wet life zones (e.g., *Premontane wet forest* and *Tropical wet forest*) in Nicaragua, Costa Rica, and western Panama (Chiriquí Province) have typically dark-drying blades with moderately confluent and auriculate lateral lobes. Collections from the highlands of Nicaragua and Costa Rica may have one to three inflorescences per axil, and each inflorescence is typically smaller than when but a single inflorescence is present.

Collections from lowland southwestern Costa Rica on the Pacific Coast differ from those of lowland Pacific coastal Panama and El Salvador in having the lobes more broadly confluent; more

prominently auriculate lateral lobes; broader medial lobes, with more numerous primary lateral veins; and in often drying more yellowish brown. Particularly noteworthy is Davids 24202 from the Costa Rican highlands southeast of Las Alturas. It has comparably broadly convolute lateral lobes and approaches *P. cotobrusense* in having medial lobes with closer, more numerous primary lateral veins and five inflorescences per axil. It is conceivably a hybrid with that species.

Collections from central Panama, especially El Valle and the vicinity of Cerro Jefe, exhibit extreme variation in leaf morphology and even spathe coloration. This is true to a lesser extent elsewhere in Panama, at Santa Fe, along the El Llano-Cartí Road, and on the Santa Rita Ridge Road. In Coclé, typical plants have somewhat blackened leaf blades with slender, weakly auriculate and weakly confluent lobes. These collections have the medial lobe ranging from 4.4 to 7.2 times longer than wide. This same narrow-lobed variation occurs also on Cerro Jefe and the Santa Rita Ridge Road where the lobes may be proportionately even more slender, ranging from 7.3 to 13 times longer than wide. Two collections from the El Llano-Cartí Road, Thompson 4704 and Miller et al. 862, have somewhat larger blades with the medial lobes five to six times longer than wide. These narrow-leaved forms vary from having 1-2 relatively large inflorescences per axil (Luteyn 4066) or with up to three smaller inflorescences per axil (Gentry 7433). The spathes are variously reported as reddish green or red-brown, to pink, red, or purple.

Spathe color is not variable throughout most of Central America, where the spathe tube is generally green outside but maroon in the lower two-thirds within. Spathe color becomes more variable in Panama, where in the narrow-lobed forms (mentioned above) it may be reported as purplish or red. As one approaches South America, spathe color becomes more variable. In eastern Panama at the Serranía de Majé and in Darién Province, the spathe is reported as green mottled purple (Whiteford & Eddy 223) or yellow-green with purple flecks (Churchill & de Nevers 4480), or even cream with red spots (Knapp & Mallet 3160). There seems to be no correlation with either elevation or the degree to which the lobes are confluent or auriculate.

Style type in *P. tripartitum* is also variable. Throughout most of its range, from Mexico to Panama along the Caribbean coast, the style is raised and button-shaped (Mayo style type D). Generally, the stylar canals emerge onto the surface near the outer margin of the style through moderately large pores on a generally flattened apex. However, some

collections, notably *Grayum et al.* 5034 from 600 m in Heredia Province in Costa Rica, have pores somewhat pie-shaped and the general level of the apex somewhat sunken. More extreme manifestations of this phenomenon are present in *Grayum* 6979 (studied in reconstituted condition), from Heredia Province at 700–900 m and *Croat & Porter* 16241, from Bocas del Toro Province, Panama, near sea level. In these collections the style type might more appropriately be referred to as a Mayo style type E since it is definitely funnel-shaped with a distinct marginal rim. In *Grayum* 6979 the style is broadly bowl-shaped with moderately large pores situated about midway between the annulus and the center. In *Croat & Porter* 16241 the funnel is held above the surface of the style in dried condition, with the margins curled under, and there are radial ribs on the inside of the funnel just as there are in the *Grayum et al.* 5034 collection. It is also similar to the latter collection and the remainder of the type D styles in having large pores positioned along the outer edge of the nearly straight-walled funnel. Indicating that these style types are probably merely variations of the D-type style is the fact that *Gómez et al.* 21988 from the highlands of Puntarenas Province has both extremes of style types represented on both of the above-mentioned collections.

Three other style types are also present in *P. tripartitum* and may ultimately allow separation of distinct subspecies because they occur in more isolated populations. One of these is Mayo style type E, represented by specimens collected from the Pacific slope of Costa Rica, e.g., *Croat* 35248 from San Isidro–Dominical Road at 1000 m, and *Burger & Stolze* 5473 from Rincón de Osa. In this type of style the stylar pores are smaller than in type D and are arranged in a smaller circle at the base of the funnel. Similar to the material from southwestern Costa Rica in this respect is a collection from Darién Province, Panama, near the Colombian border (*Whiteford & Eddy* 223).

The second novel type of style is present on collections from higher elevations in Panama, e.g., *Croat* 66228 from Chiriquí Province and *Croat* 69721 from Darién Province. This style type is perhaps sufficiently different to warrant the recognition of these collections as a distinct subspecies. It is similar to Mayo style type B, except that the style is weakly sunken rather than flattened. The stigmatic papillae may cover the entire concave area or they may arise primarily from the center. The stylar pores are smaller than in style type D, and are arranged in a tight circle at the center of the funnel, much as in the case of style type E in south-

western Costa Rica. In this case, however, there is no sign of a protruded funnel, so apparent on collections from the latter area.

While the style types mentioned here seem to warrant recognition of two or more subspecies in *P. tripartitum*, the highly variable general morphology of the species and the general state of preservation of the ovaries of most collections precludes the thorough analysis needed.

Standley and Steyermark (1958a) reported the fruits of *P. tripartitum* as red, but this is unsubstantiated. Bunting (1965) stated that the locules are strictly 1-ovulate, whereas Standley and Steyermark (1958a) described them as 1–3-ovulate. Bunting's findings have here been confirmed.

Many of the variations discussed above, if considered in isolation, are distinctive and would appear to be worthy of recognition. However, considering the immense variation within populations, the overlapping characteristics, and the generally similar structure of the flowers, it seems best to consider this group a single but highly variable species.

This species was reported in Jacquin's (1797) original description for Venezuela, but it does not occur there. Since the type illustration (t. 190) clearly represents this species it must be presumed that Jacquin, who apparently described the plant from a cultivated collection, must have been confused about the origin of the material. Schott (1829), who transferred *Arum tripartitum* to *Philodendron*, claimed to have seen only living material, perhaps the same collection seen by Jacquin at Schönbrunn. Bunting (1979) excluded *P. tripartitum* from his treatment of the Araceae for Venezuela and used the name *P. holtonianum* Schott for a Venezuelan species with three-lobed species. The Jacquin plate (t. 190) clearly does not match *P. holtonianum*, which differs in having the lateral lobes more broadly confluent and not at all auriculate at the base with the posterior rib naked 3 to 5 cm at the base. In contrast, leaves of *P. tripartitum* are often noticeably auriculate with the posterior rib not at all naked. Krause (1913) treated *P. holtonianum* as a variety of *P. tripartitum* but these species, while superficially similar in blade shape, are probably not closely related.

*Caladium trifoliatum* Desf. was questionably included by Krause as a synonym of this species, but the description is of a plant with the lateral lobes petiolulate, whereas leaf blades in *P. tripartitum* are at most trisect, with the lateral lobes sessile.

*Additional specimens examined.* BELIZE. Cayo: Lundell 6336 (CM, MICH); McCaw River–Cuevas, vic. of Milonario, 1900 ft., *Croat* 23670 (MO). *Stann Creek*: Mid-

dlesex, *Gentle* 2936 (CM, MICH). **Toledo:** Edwards Road beyond Columbia, *Gentle* 6217 (LL); Jacinto Creek, *Gentle* 6002 (LL); Monkey River, Monkey River-Cockscorb, *Gentle* 4458 (LL, MICH); Punta Gorda, 1 mi. E of junction with road to San Antonio, 100 ft., *Croat* 24509 (MO); SW Maya Mountains, Columbia River Forest Reserve, trail between Union and Gloria Camps, 700–750 m, 16°23'22"N, 89°08'10"W, *Holst* 4388 (MO); Southern Maya Mountains, Bladen Nature Reserve, 260 m, 16°30'15"N, 88°55'07"W, *Davidse* 35794 (BRH, MO). **COSTA RICA.** **Alajuela:** 22 km NE of Quesada by air, 4 km W of Muelle San Carlos, 10°28'N, 84°30'W, *Liesner* 14121 (B, CR, MO); Río Peñas Blancas, below Monteverde Cloud Forest Nature Reserve, 1250–1350 m, 9°17'N, 84°86'W, *Burger et al.* 10770 (CHAPA, NY); 850 m, *Bello* 362 (CR, MO); 700–900 m, *Huber & Zuchowski* 9468 (CR, MO); San Ramon, 820 m, 10°19'N, 84°43'W, *Bello & Eyo* 2684 (INB, MO); 840 m, *Bello et al.* 10870 (CR, INB); 15 km N of Boca Arenal, ca. 100 m, 10°38'N, 84°31'W, *Liesner et al.* 15043 (CR, MO); Cañas-Upala, 13.8 km N of Bijagua, 100–150 m, *Croat* 36447 (MO); 4 km NNE of Bijagua, ca. 400 m, 36253 (MO); Upala, 550 m, *García* 187 (CR, INB); Finca Los Ensayos, ca. 11 mi. NW of Zarco, *Croat* 43593 (MO). **Cartago:** Turrialba, *Cook & Doyle* 378 (US); Moravia—Quebrada Platanillo (Tspirí), ca. 1250 m, *Croat* 36659 (MO); 4 km SE of Pejibaye along Río Gato, 700 m, 9°48'N, 83°42'W, *Liesner* 14342 (CR, MO). **Guanaacaste:** Volcán Miravalles, ca. 800 m, 10°42'N, 85°07'W, *Burger & Geary* 9127 (CR, F, U); Parque Nacional Rincón de la Vieja, Volcán Santa María, 900–1200 m, 10°47'N 85°18'W, *Davidse et al.* 23313 (CR, MO); 840 m, *Risera* 479 (CR, MO). **Heredia:** La Selva Field Station, 100 m, *Grayum* 3019 (DUKE); 3033 (DUKE); 2444 (DUKE); 100 m, 2863 (DUKE); *Croat* 44227 (MO); *Grayum & Sperry* 2212 (DUKE); *Sperry* 962 (DUKE); *Folsom* 10011 (DUKE); Quebrada Tigre—Río Sardinal, ca. 9 km SW of Las Horquetas, ca. 600 m, 10°17'N, 84°02'W, *Grayum et al.* 5034 (MO); Volcán Barva, 700–950 m, 10°17'30"N, 84°04'30"W, *Grayum* 6979 (MBM, MO, W); Route 9, 3 km S of La Virgen de Sarapiquí, 200 m, *Woodruff* 188 (FLAS); 9 km SE of San Ramón, 1000 m, 10°16'N, 84°05'W, *Loiselle* 161 (MO); San Ramón, *Tondus* 177194 (K, P). **Limón:** ca. 4–5 airline km S of Islas Buena Vista in the Río Colorado, 100–180 m, 10°38'40"N, 83°41'W, *Davidse & Herrera* 31293 (CR, MO); Hitoy Cerere reserve, SW of Valle La Estrella, 150–550 m, 9°39'N, 83°02'W, *Grayum et al.* 5809 (MO); 2 km W of Río Toro Amarillo, W of Guápiles, 275 m, 10°13'N, 83°50'W, *Thompson & Rawlins* 1217 (CM); Hone Creek—Cahuíta, *Gómez et al.* 20523 (MO); ca. 1 km N of Shiroules, ca. 50 m, *Croat* 43277 (MO); ca. 10 mi. S of Punta Cahuíta, ca. 3 mi. S of turnoff to Briber, ca. 70 m, *Croat* 43202 (MO); vic. Laguna Pereira, 2–4 m, 10°47'–49'N, 83°37'W, *Steens et al.* 25184 (MO); Guápiles, 850 ft., *J. D. Smith* 4981 (US); 30 m, *Robles* 2241 (CR, MO); Los Angeles, San Miguel, 1100 m, 10°06'40"N, 83°59'40"W, *Herrera & Schik* 3826 (CR, INB, MO); Parque Nacional Tortuguero, 40 m, 10°28'N, 83°34'W, *Robles* 1738 (CR, MO). **Puntarenas:** Cordillera de Talamanca, 2160 m, 9°20'20"N, 83°13'55"W, *Bütner* 1787 (INB, MO); Parque Nacional Corcovado Estación Sirena, sea level, 8°28'N, 83°35'W, *Knapp* 2189 (F, MO, NY, US); 1–50 m, *Kernan & Phillips* 1090 (CR, MO); 220 m, 8°34'N, 83°31'W, *Aguilar* 2848 (CR, INB); Rincón de Osa, 20–150 m, *Liesner* 1991 (MO, US); Fila El Tigre, SE of Las Alturas, 1350–1450 m, 8°56'N, 82°51'W, *Davidse* 24202 (B, MO); Río Canasta, 9.5 airline km NW of Aguas Calientes, between Cerro

Frantzius and Cerro Pittier, 1500–1600 m, 9°02'N, 82°59'W, *Davidse et al.* 28422 (CR, MO); Osa Peninsula, Rancho Quemado, ca. 15 km W of Rincón, 200 m, 8°40'N, 83°34'W, *Hammel et al.* 16981 (CR, MO); Chacarita—Rincón de Osa, 10 km W of Chacarita, ca. 100 m, 8°45'N, 83°18'W, *Croat & Grayum* 59881 (MO); Osa, Agua Buena, 60–70 m, 8°41'32"N, 83°30'21"W, *Ramírez* 223 (CR, MO); vic. Rincón de Osa, 50–200 m, 8°42'N, 83°31'W, *Burger & Geary* 8922 (F); 20–60 m, *Uley & Uley* 1234 (F, MO); 30 m, 8°42'N, 83°31'W, *Burger & Stolze* 5473 (CR, F, US); Quebrada Vaquedano, 500 m, 8°38'45"N, 83°35'25"W, *Herrera* 4013 (CR, MO); La Tigra—Las Mellizas, 1280 m, *Gómez et al.* 21988 (MO); Palmar Norte, trail to Jalisco, 50–700 m, *Croat* 35193 (MO); 6 km S of San Vito de Java, ca. 5000 ft., *Raven* 21852 (CR, DH, DS); Río Bella Vista near Las Alturas, 1300 m, 8°56'N, 82°51'W, *Lent* 2686 (CR, F); Finca Las Alturas, at Cotón, Río Cotón, ca. 1300 m, *Croat* 44362 (MO); La Unión—Cotón, Río Negro—Río Coto Brus, *Croat* 26582 (MO); Finca Loma Linda, 1 mi. SW of Cañas Gordas, 1150 m, *Croat* 22295 (MO); Burica Peninsula, Quebrada Palito, 20–270 m, *Croat* 22617 (MO). **San José:** El General Valley, vic. of San Isidro El General, 730 m, *Molina et al.* 18308 (EAP, F, NY, US); San Isidro del General—Dominical, above Alfombra, 1000 m, 9°18'N, 83°46'W, *Burger & Baker* 10113 (F); 990–1100 m, *Croat & Hannon* 79108 (INB, MO); 2 km N of Dominical along CR-223, 40–100 m, *Uley & Uley* 4944 (DUKE); ca. 4.5 mi. SW of Canán road from Rivas, ca. 900 m, *Croat* 43438 (MO), 43446 (MO); 9 mi. SW of Río Pacuar, 680 m, *Croat* 35342 (MO); SW of San Isidro, 4.8 mi. from Río Pacuar, 1000 m, *Croat* 35248 (MO). **EL SALVADOR.** Jardín Botánico, Zona 23SC, 800 m, 13°40'N, 89°15'W, *Villacorta* 317 (LAGU, MO). **GUATEMALA.** **Alta Verapaz:** near Finca Sepacuita, *Cook & Griggs* 743 (US); Chamá, *Johnson* 404 (CM, US); Cibilquitz, 350 m, *Tuerckheim* 8023 (US); *J. D. Smith* 8023 (US); Pantín, below Tamahú, 600 m, *Standley* 70543 (F); 4 mi. up road to Oxeo, N off Highway 7E, ca. 6 km NE of Panzacán, 500 m, *Croat* 41610 (MO); 9 mi. up road to Oxeo, 800 m, 41684 (MO). **Izabal:** near Entre Ríos, 18 m, *Standley* 72745 (F). **Petén:** Dolores, km 83, Machaquilla Road, Contreras 2115 (LL). **HONDURAS.** **Atlántida:** Lancetilla Reserve, near Tela, 70–150 m, 15°44'N, 87°27'W, *Croat* 42670 (MO); *Croat & Hannon* 64640 (MO, NY); *Standley* 53381 (F, US); S of San Alejo near Río San Alejo, 150–270 m, *Standley* 7962 (F); ca. 10 km SW of La Ceiba, 80–150 m, 15°42'N, 86°51'W, *Liesner* 26077 (MO). **Gracias a Dios:** Ahuas Bila, 200 km SW of Puerto Lempira, 100 m, *Nelson & Cruz* 9181 (MO, UNAH); Río Plátano, 84°40'–85'N, 15°30'–55'W, *Clewell & Cruz* 4062 (CHAPA, MO, UC). **Morazan:** vic. El Zamorano, 800 m, *Molina* 34354 (MO). **Yoro:** Tuxtiguit River, 1 km SW of San José de Tuxtiguit, 100–400 m, 15°30'N, 87°26'W, *Hazlett & Brant* 8047 (MO); Río Guán Guán, ca. 3 km S of San José de Tuxtiguit, 380 m, 15°30'N, 87°27'W, *Evans* 1323 (MO). **MEXICO.** **Chiapas:** 27–43 km SW of Palenque, 210–810 m, *Croat* 40335 (MO), 40306 (MO); 460 m, *Croat* 40269 (MO); 20 mi. N of Ocozacoatlán, 700 m, *Croat* 40634 (MO, NY); 13 mi. N of Ocozacoatlán, 900 m, *Croat* 40620 (MO, NY); Mpio. Cintalapa, Oaxaca—Chiapas border, 4 km W of La Ciénega, 38 km W of Las Cruces, 1400 m, *Breedlove* 25132 (MO); 5 km W of Bonampak, 350 m, *Martínez* 6972 (MO); 2 km NE of Bonampak above Cojolite, 350 m, 7939 (MO); Laguna Ocoatl Grande, 45 km E of Ocoasingo, *Dressler* 1513 (GH, US); 10 mi. N of Ocozacoatlán, 3800 ft., *Breedlove* 10330 (F); km 18 on road to Malpasco (N of Ocozo-

cautla), 2700 ft., *Kennedy 1426* (SEL); Bochil-Pichucalco, 17.1 km SW of Pichucalco, 430 m, *Croat 78662* (CHIP, MO). **Oaxaca:** Uxpanapa region, 17 km E of Sarabia, 100 m, 17°10'N, 94°45'W, *Croat & Hannon 63212* (B, BM, K, MO, NY, US); 150 m, 17°33'N, 92°59'W, 65367 (MO); Tuxtutepe-Oxaca, 6 mi. W of Valle Nacional on Highway 175, 660 m, *Croat 39741* (MO); 8 km SW of Valle Nacional, 400 m, *Hernández & Chacón 491* (INB, MO). **Tabasco:** Grutas de Ocona near Teapa, 20–100 m, *Davidse et al. 29509* (CM, MO); Teapa, cultivated, *Croat 59160* (MO); Mpio. Tacotalpa, 0.2 km NW of Tapijulapa, *Cowan et al. 3496* (CSAT, MO). **Veracruz:** Valle de Córdoba, *Bourgeois s.n.* (P); Route 185, 4 km NE of Minatitlán, 50 m, *King 1113* (MICH); 6 mi. E of Coatzacoalcas, 10 m, *Croat 40058* (MO); 7 km NE of Sontecomapan, 0–50 m, 18°33'30"N, 94°59'W, *Nee 22607* (F, MO); 5 km NW of Pajapan, Cerro San Martín, 750–900 m, 18°17'20"N, 94°43'W, *Nee & Calzada 22697* (F, MO, NY); Los Tuxtlas, 200 m, 18°42'N, 95°10'W, *Croat 63163* (MICH, MO, US). **NICARAGUA:** 40–45 km SW of Waspmam, 10–100 ft., *Seymour 3776* (MO). **Jimotepe:** Río Bocay, Salto Acatulú, ca. 130 m, 14°13'N, 85°10'W, *Stevens et al. 16750* (MO). **Matagalpa:** Comarca Wanawás, Río Bilampf, 180–200 m, 12°03'–02'N, 85°13'W, *Araguistain & Moreno 2401* (MO, SAR); Cerro Musún, path to Palán, 3000–600 m, *Araguistain & Moreno 2471* (AAU, LE, MO); 20 km E of Matagalpa, along Río Yasicá, 700 m, *Neill 1963* (MO); Cerro Musún, 500–800 m, *Araguistain & Moreno 2571* (MO); Cerro Musún, 8 km from village of Wanawás, 500–800 m, 12°59'–13'N, 85°14'W, *Araguistain & Moreno 2359* (MO); Quebrada Negra, Río Bilampf, 500 m, *Neill 1816* (MO). **Río San Juan:** Bocas de Sábalo, 70–100 m, 11°03'N, 84°27'W, *Moreno 26748* (MO); above Río Sábalo, 40 m, 11°03'N, 84°28'W, *Moreno & Robledo 26016* (MO). **Zelaya:** ca. 14.3 km N of El Empalme, *Stevens 8303* (BM, MO); El Empalme-Rosita, along new road to Mina Nueva América, *Stevens 12667* (MO); 12668 (MO); *Pipoly 5293* (MO); 15 km W of Rama, 100 ft., *Harmon & Fuentes 5079* (MO); Kuikuinta, S of Siama, 160 m, *Neill 4132* (MO); Río Prinzapolka-Quebrada San Rafael, Wanf, ca. 100 m, 13°42'N, 84°50'W, *Pipoly 4578* (MO); 4587 (MO); 4588 (MO); 4589 (MO); Wanf-Siama, near Río Matías, ca. 0–100 m, 13°43'N, 84°49'W, *Pipoly 4682* (MO); 4701 (MO); Colonia Yolania-Colonia Manantiales (Colonia Somoza), ca. 200–300 m, ca. 11°36'–37'N, 84°22'W, *Stevens 4825* (MO); Rosita-Puerto Cabezas, ca. 15.7 km SW of Río Kukulaya, <100 m, 13°58'N, 84°12'W, *Stevens 8502* (MO); Caño Zamora on Río Rama, ca. 10 m, ca. 11°57'N, 84°16'W, *Stevens 8821* (MO); Bonanza-Constancia, 160–360 m, ca. 13°58'–14°01'N, 84°37'–40'W, *Stevens 12482* (MO); 12455 (MO); El Empalme-Limbaika, ca. 3.9 km SE, ca. 65 m, 13°39'N, 84°24'W, *Stevens 12883* (MO); 12886 (MO); vic. of Bonanza, 200–350 m, 14°01'N, 84°35'W, *Stevens 13031* (MO); 13053 (MO); 250–350 m, ca. 14°02'N, 84°34'–35'W, 18786 (MO); 7.4 m SE, ca. 130 m, 13°38'N, 84°23'W, 21752 (MEXU, MO); 12346 (MO); ca. 140 m, ca. 14°01'N, 84°34'W, *Stevens 9056* (BM, MO); 1 km N of "Las Mercedes" settlement, 160–180 m, *Vincelli 3344* (MO); "Bodega" region, 30 km NE of Río Blanco, 80–100 m, 13°03'N, 84°58'W, *Moreno 24078* (MO); "San Agustín," SE of Rama, ca. 60 m, 12°09'N, 84°12'W, *Robledo 564* (MO); *Neill 3632* (MO); Bonanza-El Salto Grande, Laguna Santa Rosita, ca. 140–160 m, 14°03'N, 84°37'W, *Pipoly 3687* (MO); Mpio. Siama, La Pimentada, *Ortiz 1966* (MO); Río Mico, Experimental Station El Recreo, 30 m, 12°10'N, 84°18'W, *Davidse et al. 30768* (MO); Río Prinzapolka, ca. 2 km S of

Wanf, ca. 0–100 m, 13°42'N, 84°50'W, *Pipoly 4733* (MO), 4738 (MO), 4747 (MO); Río Punta Gorda, Atlanta, 10 m, 11°34'N, 84°01'W, *Moreno & Sandino 12769* (MO); Waspmam-Puerto Cabezas, Río Troncera, <200 m, ca. 14°43'N, 84°06'W, *Pipoly 4042* (MO); Río Wilike Grande, Wilike region, 100 m, 13°05'N, 84°57'W, *Moreno 23999* (MO). **PANAMA:** **Bocas del Toro:** Fortuna Dam-Chiriquí Grande, 3.2 mi. N of Continental Divide, 700 m, 8°45'N, 82°15'W, *Croat & Grayum 60281* (MO); Changuinola-Almirante, Milla 7.5, *Croat & Porter 16241* (MO); Valle del Silencio, Río Changuinola, ca. 1 km above mouth of Río Teribe, <100 m, 9°21'40"N, 82°31'40"W, *Croat & Zhu 76431* (MO). **Canal Area:** Frijoles, 25–30 m, *Pittier 3755* (US); Barro Colorado Island, *Shattuck 182* (F, GH, MO); *Ebinger 99* (MO); *Croat 11838* (MO); *Schmalzel 797* (MO); *Dwyer et al. 8474* (MO); *Croat 16535* (MO); 15067 (MO); 11006 (MO); 10741 (MO, UC); 10198 (MO, SCZ); 10085 (MO); 6214 (MO); 6294 (MO); 8605 (MO); 8040 (MO); 6861 (MO); 6495 (MO); 4001 (MO); Summit Garden, *Croat 17058* (MO); 10284 (MO, SCZ); 33015 (MO); 10915 (MO). **Chiriquí:** Volcán-Serrano, 7 mi. N of Volcán, 8°50'N, 82°38'W, *Croat 66228* (MO); vic. El Hato del Volcán at Las Lagunas, 3 mi. from Nueva California, 1380 m, 8°46'N, 82°40'W, *Croat 67707* (MO, PMA); vic. Santa Clara, 1520 m, *Croat 66288* (F, MO); Gualaca-Chiriquí Grande 2.5 mi. beyond Los Planes de Hornito, 1099 m, 8°40'N, 82°13'W, *Croat 67739* (CM, MO); 2.8 mi. beyond Los Planes, 1200 m, 48817 (MO); Quebrada La Mina, 2.3 mi. N of Los Planes, 800 m, 8°41'N, 82°13'W, *Croat & Zhu 76364* (CM, MO); along Río Colorado, 1200–1400 m, 82°43'W, 8°50'N, *Hamilton & Krager 3757* (MO, RSA); vic. Boquete, ca. 1450 m, *Croat 26744* (MO). **Coeló:** La Mesa, N of El Valle de Antón, 800–900 m, 8°38'N, 80°09'W, *Croat 67210* (CM, MO); *Croat 13358* (MO); ca. 1 mi. N of El Valle, *Gentry & Dwyer 3591* (MO); 900 m, *Gentry 7433* (F, GH, MO). **Coeló-Panamá:** trail beyond La Mesa towards Los Llano and the border, *Luteyn 3178* (MO); 850–900 m, *Luteyn 4066* (MO). **Colón:** Santa Rita Ridge, 26 km from Transisthmian Highway, 500 m, 9°26'N, 79°57'W, *Knapp et al. 1708* (MO); Mile ca. 12, ca. 500 m, 9°25'N, 79°40'W, *McPherson 11760* (MO). **Darién:** Parque Nacional del Darién, ca. 17 km E of Pucuro, Río Pucuro, 750–850 m, 8°03.5'N, 77°17'W, *Hammel et al. 16293* (MO); western slopes of Cerro Pirre, 17 km N of El Real, Río Perisenco, 100 m, 8°01'N, 77°40'W, *Croat & Zhu 77187* (MO); vic. Cana, 500–600 m, *Croat 37641* (MO); Alturas de Nique region, 900–1250 m, 7°45'N, 77°40'W, *McPherson 11603* (MO); Cerro Sapo, 800 ft., *Hammel 1330* (MO); E base of Cerro Sapo along stream (Río Celorio?), 1300 ft., 1131 (MO); 18 km SE of Jaque, 25 m, *Garwood et al. 98* (BM, CM, MO, PMA); N of Punta Guayabo Grande, NW of Ensenada El Guayabo ridges, 0–200 m, 7°24'N, 78°07'W, *Knapp & Mallet 3160* (MO, US); Río Coasi, Cerro Coasi, *Duke 15554* (US); Río Cecalito, SE of Jacque, 7°18'N, 77°58'W, *Whiteford & Eddy 223* (BM, MEXU, MO). **Panamá:** Juan Díaz, *Standley 30618* (US); Cerro Jefe region, 750–800 m, 9°14'N, 79°22'W, *Croat 67082* (F, MO); 67084 (MO); 4.6 km beyond peak on road to Altos de Pacora, ca. 600 m, *Croat 35900* (MO); at Altos de Pacora, 750 m, 9°15'N, 79°29'W, *Croat & Zhu 76606* (MO); 3–3.5 mi. NE of Altos de Pacora, 700–750 m, 9°15'N, 79°25'W, *Croat 68636* (CM, MO); Campo Tres, 5 km NE of Altos de Pacora, *Bussey 835* (MO); Lago Cerro Azul, 500 m, *Sullisán 82* (MO); Cerro Campana, 2500 ft., *Hammel 3767* (MO, SEL); Río Majé-Quebrada Brava, 60 m, *Croat 34630* (MO); El Llano-Cartí Road, 7–12 km from



Interamerican Highway, 360–400 m, *Croat 25122* (MO); Km ca. 16–18, 400 m, *Tyson & Nee 7355* (MO); El Llano Carti road, Mile 10.6, ca. 400 m, 9°17'N, 78°58'W, *Miller et al. 862* (MO); Mile 6–10, 200–250 m, 9°15–16'N, 78°59'W, *Thompson 4704* (CM, MO); Serranía de Majé, along Río Ipetí, near confluence with Río Ambrosio, 100 m, 8°57'N, 79°32'W, *Churchill & de Nevers 4490* (MO, PMA, RSA). **San Blas:** Nusagandi, along El Llano–Carti Road, 11.6 mi. N of highway, 450 m, 9°18'N, 79°59'W, *Croat 75135* (CM, MO, PMA). **Veraguas:** Santa Fe–Río San Luis, vic. Río Tercero Brazo, 700 m, 8°33'N, 81°08'W, *Croat 66925* (MO, PMA); Río Dos Bocas, ca. 5–8 km N of Alto Piedra, 730–770 m, *Croat 25915* (MO); ca. 500 m, *Croat 34242* (MO); vic. Escuela Agrícola–Calovéboa, ca. 1200 m, 81°07'W, 8°32'N, *Hamilton et al. 1279* (MO).

***Philodendron tysonii* Croat, sp. nov.** TYPE: Panama. Chiriquí: Gualaca–Chiriquí Grande, 7.2 mi. beyond Los Planes de Hornito, 1165–1200 m, 8°44'N, 82°14'W, *Croat 67844* (holotype, MO–3614677; isotypes, AAU, B, CAS, CM, CR, DUKE, F, GH, K, M, MEXU, NY, P, PMA, RSA, US, VEN, W). Figures 29, 425–430.

Planta plerumque hemiepiphytica, interdum terrestris; internodia 2–6 cm longa, 3.5–4 cm diam.; cataphylla (10)21–33 cm longa, incostata vel leniter 1-costata, saepe rubella, persistens semi-intacta, demum decidua; petiolus teres vel subteres, 39–90 cm longus, 8–15 mm diam.; lamina ovata, coriacea, nitida, 32–70 cm longa, 24–50 cm lata; costa postica plerumque vix aut haud nuda, raro nuda 3 cm; inflorescentia 4–5; pedunculus 3–12 cm longus, 3–8(11) mm diam.; spatia 11.5–19 cm longa, lamina spathae extus viridi, suffusa marronina aut rubella, intus alba vel pallide viridi; tubo spathae extus suffuso marronino aut rubello, intus atrimarronina suffuso in laminam; pistilla (5)6–8(9)-locularia; loculi (4)5–7-ovulati.

Usually hemiepiphytic, appressed-climber, sometimes terrestrial; internodes short, stout, semiglossy, smooth, 2–6 cm long, 3.5–4 cm diam., dark green; roots elongate, fine, branched; cataphylls (10)21–33 cm long, unribbed to weakly 1-ribbed, sometimes sharply 2-ribbed near base and unribbed near apex, frequently reddish, frequently striate, emarginate at apex, persisting semi-intact at upper nodes, eventually deciduous, emarginate at apex; **petioles** 39–90 cm long, 8–15 mm diam., terete to subterete, surface densely pale-striate, glossy; sheath 10.5–11 cm long; **blades** ovate, coriaceous to moderately coriaceous, glossy to semiglossy, conspicuously bicolorous, acuminate at apex (the acumen sometimes inrolled and twisted), cordate at base, 32–70 cm long, 24–50 cm wide (1–1.8 times longer than wide and averaging 1.4 times), (0.4–1.1 times longer than petiole), averaging about  $\frac{1}{3}$  as long as petiole, upper surface dark green, drying dark brown to blackish brown, lower surface slightly paler, drying dark yellow-brown to greenish brown; anterior lobe 25–52.5 cm

long, 20–50 cm wide (2–4.8 times longer than than posterior lobes); posterior lobes broadly rounded to broadly obtuse, 8–20 cm long, 11–22 cm wide; sinus parabolic, sometimes spatulate, (3)4–12 cm deep; midrib broadly convex to almost flat, conspicuously paler than surface above, convex to bluntly acute, darker than surface below; basal veins 5–9 per side, with (0)1(2) free to base, numbers 3–4 coalesced 1.5–3 cm; posterior rib usually scarcely or not at all naked, rarely naked for 3 cm; primary lateral veins 4–10 per side, departing midrib at a 60–70° angle, moderately straight to the margins, narrowly sunken above, convex and somewhat paler than surface below; interprimary veins sunken, usually concolorous above, raised, darker than surface below; minor veins moderately distinct below, arising from both the midrib and primary lateral veins. **INFLORESCENCES** 4–5 per axil; peduncle 3–12 cm long, 3–8(11) mm diam., green, finely striate; **spathe** 11.5–19 cm long (1.5–4.5 times longer than peduncle); spathe blade green to white, lightly to heavily tinged maroon or reddish, finely striate-speckled, margins clear outside, white to pale green inside; spathe tube tinged maroon or reddish, finely striate-speckled outside, 4.5–10 cm long, dark maroon, suffused into blade inside, weakly constricted above the tube; **spadix** white throughout, 10.7–15.6 cm long; pistillate portion 3.4–5.2 cm long in front, 2.4–3.1 cm long in back, broadest midway and gradually tapered toward both ends, 1.2 cm diam. at apex, 1.2–1.4 cm diam. at middle; staminate portion 7.7–9.4 cm long; fertile staminate portion white, barely constricted above the sterile staminate portion, barely tapered toward apex, 1–1.2 cm diam. at base, 1–1.2 cm diam. at middle, 6–7 mm diam. ca. 1 cm from apex; sterile staminate portion 9–10 mm diam.; pistils 1.5–2.7 mm long, 0.8–1.6 mm diam., white; ovary (5)6–8(9)-locular, with sub-basal placentation; locules (0.8)1.1–1.3 mm long, 0.3–0.6 mm diam.; ovules (4)5–7 per locule, (0.1)0.3–0.4 mm long, 2-seriate, contained within transparent ovule sac; style similar to style type B; stigma 0.8–1.2 mm diam., 0.2–0.3 mm high. **INFRUDESCENCE** with pistillate spadix 6.5 cm long, 4 cm diam.

Flowering in *Philodendron tysonii* occurs mostly in June and July but also in September. In reality, flowering probably occurs from January through September or perhaps October, since post-anthesis collections have been made nearly throughout the year except February. Immature fruiting collections have been made in March, September, and November.

*Philodendron tysonii* is known only from Panama, but is expected to be found in the Cordillera

de Talamanca of Costa Rica. It ranges from 600 to 1500 m elevation in *Premontane rain forest* and *Tropical Lower Montane rain forest* life zones.

*Philodendron tysonii* is a member of *P. sect. Calostigma* subsect. *Macrobilium ser. Reticulata*. This species is characterized by its stout, short (usually 3–5 cm diam.) internodes; unribbed or weakly one-ribbed, frequently reddish cataphylls persisting semi-intact (often becoming fibrous) at the upper nodes; terete to subterete petioles; and ovate-cordate blades averaging 1.4 times longer than wide and about three-fourths as long as the petioles, with 6–9 basal veins and a weakly developed posterior rib, which is rarely naked at the sinus.

*Philodendron tysonii* is perhaps most easily confused with *P. antonioanum*, with which it may occur. The latter species differs in having sharply 2-ribbed cataphylls, blades drying gray-green with prominent "cross-veins" and 4 to 6 pairs of primary lateral veins, and short-pedunculate inflorescences with spathe tubes dark magenta on both surfaces and barely constricted above. In contrast, *P. tysonii* has usually bluntly 1-ribbed cataphylls, blades drying ± blackened and smooth above, 6–10 primary lateral veins, more prominently pedunculate inflorescences, and prominently constricted spathes merely tinged red on the tube outside. In addition, the two species differ in the nature of their ovules, with *P. tysonii* having (5)6–8(9)-locular ovaries with (4)5–7 ovules per locule borne in an envelope with sub-basal placentation, while *P. antonioanum* has 5-locular ovaries with about 30 ovules per locule, with parietal placentation and not borne in an envelope.

This species may also be confused with *P. sagittifolium* (especially broader-leaved forms), which shares blades with mostly free basal veins with posterior ribs short or lacking and scarcely or not at all naked along the sinus. The latter species differs in having narrower stems (rarely more than 3 cm diam.), green, sharply 2-ribbed deciduous cataphylls (vs. persistent in *P. tysonii*), and blades usually somewhat reddish brown and proportionately longer (averaging about 2 times longer than broad) with the posterior lobes usually narrowly rounded. In addition, the juvenile blades of *P. sagittifolium* are elongated, usually oblong to narrowly triangular, and usually up to 2.5 times longer than wide (vs. ovate to narrowly ovate, less than 2.5 times longer than wide, and with broadly rounded posterior lobes in *P. tysonii*).

*Additional specimens examined.* PANAMA. **Bocas del Toro:** Fortuna Dam area, Gualaca-Chiriquí Grande, just N of the Continental Divide, 850–950 m, 8°46'N,

82°17'W, Hammel et al. 14700 (MO); ca. 8.5 mi. from bridge over Fortuna Lake, 600 m, 8°46'N, 82°16'W, Croat 67005 (CAS, CM, L, MO, PMA, TEX); Cerro Colorado, 9.2 mi. W of Chame, 1450–1480 m, 8°35'N, 81°50'W, Croat 69017 (DUKE, IBE, MEXU, MO, NY, PMA, QCA, RSA, WIS); 7 mi. from Chame, ca. 1500 m, 8°35'N, 81°45'W, McPherson 8810 (MO); 5.3 mi. N of center of Fortuna Dam, Continental Divide Trail, 8°44'N, 82°17'W, Croat & Zhu 76346 (MO); near Lago Fortuna, along trail to Río Hornito, 8°45'N, 82°18'W, Croat & Zhu 76374 (MO, PMA). **Chiriquí:** Cerro Colorado, above San Félix, 18–27 mi. N of Pan-American Highway, 1200–1500 m, Croat 33148 (MO); 24 mi. N of Río San Félix, 1430–1500 m, 48486 (MO); 34.1 km N of Río San Félix, 1390 m, 37258 (MO); 28 mi. above Río San Félix, 1200–1500 m, 33269 (MO); from Chame to ca. 9 mi. along road, 1100–1750 m, 8°35'N, 81°54'W, Hammel & Trainer 15008 (MO, NY); Fortuna Dam area, Fortuna-Chiriquí Grande, Continental Divide, 1170 m, 8°44'N, 81°17'W, Croat 66711 (B, BR, COL, CR, K, MO, NY, PMA, US); Hammel & Kress 13490 (DUKE); Continental Divide trail, 8°44'N, 82°17'W, Croat & Zhu 76330 (K, MO, NY, SCZ, US); trail to Río Hornito, 1100–1350 m, 8°45'N, 82°15'W, McPherson 11660 (MO); Croat & Zhu 76417 (AAU, CM, ENCB, GB, IBE, M, MO, NY, PMA, TEX, WIS); confluence of Río Hornito and Río Chiriquí, 1050–1100 m, ca. 8°44'N, 82°13'W, Stevens 18376 (MO); 18377 (MO); vic. IRHE facilities, 1100–1200 m, 8°45'N, 82°08'W, Croat 66589 (AAU, COL, DUKE, F, K, M, MEXU, MO, P, PMA, US, VDB); 1300 m, Croat 48760 (CM, MO); Quebrada Los Chorrros-Quebrada Frank, N of reservoir, 1100 m, 8°45'N, 82°13'W, Churchill & Churchill 6081 (MO); 8.3 mi. beyond Los Planes de Hornito, 1260 m, 8°44'N, 82°16'W, Croat 49943 (MO); 1300 m, 8°45'N, 82°17'W, 49821 (MO); 1400 m, 48704 (AAU, MEXU, MO, U, U, W). **Coclé:** El Copé region, Alto Calvario, Continental Divide, 9.4 km above El Copé, 750–900 m, Croat 44719 (MO); 710–800 m, 8°39'N, 80°36'W, 68766 (MO, PMA); 850 m, 67577 (AAU, CM, F, G, L, MEXU, MO, NY, PMA, TEX, US); 930 m, 49203 (MO); 650–850 m, Folsom 6217 (MO). **Panamá:** Cerro Jefe region, vic. of summit, 9°15'N, 79°30'W, Croat & Zhu 76210 (CM, MO); Witherspoon & Witherspoon 8531 (MO); ca. 1000 m, Croat 49084 (CAS, K, MO, PMA); 1000 m, 9°15'N, 79°30'W, Knapp 2240 (MO); La Enseida, 1000 m, Luteyn & Kennedy 1775 (DUKE); 0.8 mi. beyond turnoff to Altos de Pacora, 770 m, 9°15'N, 79°29'W, Croat & Zhu 76614 (CM, MO, WIS); 3–3.5 mi. NE of Altos de Pacora, 700–750 m, 9°15'N, 79°25'W, Croat 68688 (AAU, CM, F, G, MEXU, MO, PMA, TEX, US); 800–1000 m, 22671 (F, MO); 2400 ft., Antonio 4753 (MO); Croat 67059 (AAU, B, BR, C, CAS, CM, COL, CR, DUKE, EAP, ENCB, F, HNMM, HUA, IBE, JAUM, JBGP, K, L, M, MO, NY, PMA, QCA, US, USCG, WIS); 850–900 m, 9°15'N, 79°30'W, McDonagh et al. 36 (BM); ca. 1000 m, Mori et al. 3737 (MO); ca. 950 m, ca. 9°15'N, 79°30'W, McPherson 7109 (MO); Cerro Azul-Cerro Jefe, ca. 1000 m, Luteyn 3209 (DUKE); 3000 ft., Tyson et al. 44494 (MO, SCZ); Campo Tres, ca. 700 m, Croat 27047 (F, L, MO, US); El Llano-Carfí, 6.8 mi. from highway, 350 m, Croat 49120 (MO). **Veraguas:** Santa Fe region, Alto Piedra-Calorehara, 3200–5600 ft., Hammel 4604 (MO); Río Tercero Brazo, beyond Alto Piedra, Croat 27328 (MO); between Río Tercero Brazo and summit of hill leading down to the Atlantic lowlands, 700 m, 8°33'N, 81°08'W, Croat 66926 (B, CAS, F, K, MO, PMA, US); Parque Nacional Cerro Tute, near Alto Piedra,

0.5 mi. N of Alto Piedra, 1250–1350 m, *Croat 48955* (F, MO); 800–1030 m, *Croat & Zhu 76911* (CM, MO).

***Philodendron ubigantupense*** Croat, sp. nov.

TYPE: Panama. San Blas: on mainland opposite Isla Miria Ubigantupo, along trail to Digole, to 20 m, 9°26'N, 78°54'W, 19 July 1987, *H. Herrera 294* (holotype, MO-3585503; isotype, PMA). Figure 436.

Planta hemiepiphytica; internodia usque to 7.5 cm longa, 1 cm diam.; cataphylla decidua; petiolus subteres, 12–13 cm longus, in sicco 5–6 mm diam.; lamina oblonga, anguste cordulata basi, 27–32 cm longa, 8.5–9 cm lata, in sicco caniviridis; nervis lateralibus l obscuris; inflorescentia 1; pedunculus 8 cm longus, in sicco 5 mm diam.; spathe 9.3 cm longa, viridis; pistilla (7)8-locularia; loculi 1-ovulati.

Hemiepiphytic; stem scandent; internodes to 7.5 cm long, 1 cm diam., slender, drying 7 mm diam., gray-brown, irregularly and closely fissured, somewhat flaking; roots several per node, drying moderately smooth, reddish brown, semiglossy; cataphylls lanceolate, deciduous; **petioles** 12–13 cm long, drying 5–6 mm diam., subterete, drying dark gray-brown, minutely striate, sheathed to 1.5 cm; **blades** oblong, acuminate at apex (the acumen tightly inrolled, 5 mm long), narrowly cordulate at base, 27–32 cm long, 8.5–9 cm wide (3–3.8 times longer than wide), (2.2–2.5 times longer than petiole), about as long as petiole; upper surface dark green, drying weakly glossy, gray-green, lower surface slightly paler, semiglossy, drying greenish gray; sinus to 1.3 cm deep; midrib broadly convex above, convex and paler below, drying faintly striate; primary lateral veins almost totally obscure above and below; many minor veins, all free to base, arising from the midrib only, drying obscure above, close and weakly raised below, spreading at a 55–75° angle, ± straight to the margin. **INFLORESCENCES** solitary; peduncle 8 cm long, drying 5 mm diam.; **spathe** green, 9.3 cm long, (1.2 times longer than peduncle); spathe tube weakly constricted, ca. 4.5 cm long; **spadix** bluntly pointed at apex, 10.8 cm long; pistillate portion 4 cm long in front, 3.7 cm long in back, 8 mm diam. at apex, drying 1 cm diam. at middle; staminate portion 8.1 cm long; fertile staminate portion 8.5 mm diam. at middle, 6.5 mm diam. ca. 1 cm from apex; sterile staminate portion 8 mm diam. at apex, 9 mm diam. at base; pistils 1.5 mm long; ovary (7)8-locular, 1.2 mm diam., with sub-basal placentation; locules 1.1 mm long, 0.2–0.3 mm diam.; ovule sac ca. 1 mm long; ovules 1 per locule, contained within transparent ovule sac, ca. 0.3 mm long; funicle ca. 0.3 mm long (can be pulled free to base), style 0.9 mm

diam., similar to style type D; style apex sloping to weakly rounded; style boss moderately broad and fairly pronounced; the androecium truncate, sometimes prismatic and weakly clavate, margins irregularly 4–5-sided, 0.8–1.5 mm long; thecae oblong, 0.2–0.3 mm wide, ± parallel to one another; sterile staminate flowers clavate to prismatic, irregularly 4–5-sided, 1.3–1.7 mm long.

The flowering phenology of *Philodendron ubigantupense* is uncertain owing to its rarity, but since it has small inflorescences and was in flower in July, it probably both flowers and fruits in the wet season.

*Philodendron ubigantupense* is endemic to Panama, known only from the type locality along the Atlantic coast, at or near sea level in a *Premontane wet forest* life zone.

*Philodendron ubigantupense* is a member of *P. sect. Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is characterized by its scandent habit, long, slender internodes, subterete petioles and especially by its oblong, gray-green drying blades over twice as long as the petioles, weakly cordulate leaf bases, and almost totally obscure primary lateral veins.

*Philodendron ubigantupense* is perhaps most similar to *P. folsomii*, which has blades of similar size and color. The latter species differs in having short internodes and blades merely rounded at the base and with 8–10 distinct primary lateral veins. In addition, *P. folsomii* has 6–7 locules per ovary, whereas *P. ubigantupense* has mostly 8 locules per ovary. *Philodendron immixtum* is another species that might be confused with *P. ubigantupense*, but it differs in having proportionately broader blades (to 2.8 times longer than broad) with up to five, obvious primary lateral veins.

***Philodendron utleyanum***, Croat, sp. nov. TYPE:

Panama. Colón: Santa Rita Ridge, about 7 mi. from Transisthmian Highway, ca. 200 m, 21 Dec. 1971, *Wilbur et al. 15068* (holotype, DUKE 226389). Figures 431, 432.

Planta hemiepiphytica; internodia 1–3.5 cm longa, 1.5–2 cm diam., in sicco pallide brunnea; cataphylla usque 27 cm longa, leniter 2-costata, in sicco obscure rubri-brunnea, decidua; petiolus subteres, atribrunneus, obtuse subcomplanatus adaxialiter versus apicem, 19–21.5 cm longus, in sicco 6–7 cm diam.; lamina oblongo-oblanco-lata, 32–39 cm longa, 10–14 cm lata, aliquantum inequaliter et rotundata vel truncata aut leniter cordulata basi, atriviridis supra, in sicco canobrunnea, saepe aliquantum rubella infra, in sicco brunnea; nervis basalibus 2–3 utroque, omnibus liberis ad basim; inflorescentia 1; pedunculus usque 9 cm longus, in sicco atribrunneus; spathe usque 11 cm longa, pallide viridis, extus suffusa

subrosea, intus alba, tubo spathae 5 cm longa; pistilla 4-5-locularia; loculi 1-ovulati.

Hemiepiphytic; stem appressed-climbing; internodes drying moderately glossy, irregularly and acutely ribbed with a few flat smooth intervening areas, 1-3.5 cm long, 1.5-2 cm diam., drying light brown; roots several per node, short, drying brown, to 3 cm diam.; cataphylls to 27 cm long, weakly 2-ribbed, but sharply 2-ribbed toward the apex, red tinged toward the base, drying dull reddish brown, deciduous; petioles 19-21.5 cm long, drying 6-7 cm diam., subterete, dark brown, obtusely somewhat flattened toward the apex adaxially, surface smooth to weakly ribbed; blades somewhat inequilateral, oblong-ob lanceolate, subcoriaceous, semi-glossy, acuminate at apex, somewhat inequilateral and rounded to truncate to weakly cordulate at base, 32-39 cm long, 10-14 cm wide, broadest above the middle, 1.3-1.8 times longer than petioles, upper surface dark green, drying gray-brown, lower surface often somewhat reddish, drying medium brown; posterior lobes to ca. 2 cm long; midrib drying weakly and obtusely raised, slightly paler than surface above, drying convex, drying light brown with a distinct pale border along its edges and paler than surface below; basal veins 2-3 per side, and with all free to base; primary lateral veins 8-9 per side, departing midrib at a 65-80° angle in the lower one-half, to 45° angle toward apex, weakly arcuate to the margins, drying inconspicuous except near the midrib, paler than surface above, weakly convex and paler than surface, sometimes undulate below; minor veins drying moderately inconspicuous and close, weakly undulate, arising from the midrib only, the surface and minor veins minutely and densely granular at 10× magnification. INFLORESCENCES 1 per axil; peduncle to 9 cm long, drying 4.5 mm diam., dark brown; spathe moderately coriaceous, to 11 cm long, ca. 2 cm diam., pale green with pinkish tinge, drying dark brown throughout, white within; spathe tube 5 cm long, 1.8-2 cm diam.; spadix sessile, 10 cm long; pistillate portion 5 cm long, drying to 1.7 cm diam. near base, 1.5 cm diam. midway, 1 cm at apex; staminate portion 4.7 cm long, too deteriorated for detailed studies; pistils 2.5-3 mm long, 1.5-2 mm diam., drying light brown, smooth except warty near apex; style type not studied; stigma 0.4-0.6 m diam., button-like on drying with a medial nipple and 4-5 radiate arms, sometimes with fragments of the dried stigma apron-like still attached; locules 4-5; ovaries with sub-basal placentation, 1 ovule per locule.

Flowering in *Philodendron utleyanum* is poorly

known. The single collection with immature fruits was made in December. This species probably flowers in the rainy season.

*Philodendron utleyanum* is endemic to Panama, known only by the type specimen from Colón Province in a region of *Premontane wet forest* at 200 m elevation.

*Philodendron utleyanum* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is distinguished by its short tan internodes; sharply 2-ribbed, deciduous cataphylls; subterete petioles about three-fourths as long as the blades; and oblong-ob lanceolate, brown-drying blades broadest above the middle and rounded to weakly subcordate at the base.

In terms of coloration and texture, *P. utleyanum* appears closest to *P. sagittifolium*, but in terms of blade shape it is closest to *P. pseudauriculatum*. The former differs in having the blades broadest at the base and decidedly cordate, the latter in having dark gray-green-drying leaves, more or less spongy, proportionately shorter petioles, and the spadix clearly demarcated from the peduncles by a marked color contrast. *Philodendron utleyanum* lacks any clear limit between the peduncle and the spathe.

This species is named in honor of John Utley, a participant in the Duke University expedition that collected the type. Utley has been responsible for collecting many interesting and important Araceae during his trips to Central America and during his tenure as a Peace Corps volunteer in Costa Rica.

#### *Philodendron verapazense* Croat, sp. nov.

TYPE: Guatemala. Alta Verapaz: 7 mi. on road to Oxec, along road turning N off Hwy. 7E between Tzurú and El Estor, ca. 6 km NE of Panzós, 700 m, ca. 15°28'N, 89°04'W, Croat 41656 (holotype, MO-2743518; isotypes, GUAT, US). Figures 433-435.

Planta hemiepiphytica; internodia longiora quam lata, conspicue multisulcata; cataphylla 14-17 cm longa, acute 2-costata, decidua; petiolus acuto D-formatus coplanatus adaxialiter, 29-44 longus, in sicco 2-5 mm diam.; lamina anguste ovato-triangulari-cordata, longe acuminata apice, longe cordata vel sagittata basi, 26-34 cm longa, 11.5-14 cm lata; nervis lateralibus 13-4 utroque; inflorescentia 1; pedunculus 3-14 cm longus, 4-5 mm diam.; spathe 10-18 cm longa, extas rubriovaleacea; tubo spathae intus carmesino; pistilla 4-6-loculari; loculi 2-ovulati.

Hemiepiphytic; stem appressed-climbing or sometimes scandent; pre-adult internodes to 20 cm long, 1.5-3 cm long; adult internodes green becoming tinged purple, finally tan, conspicuously many-sulcate except above each petiole, longer than broad; roots few per node, 10-30 cm long,

drying dark brown, weakly glossy, 2.3 mm diam.; cataphylls 14–17 cm long, sharply 2-ribbed, green tinged red, deciduous; **petioles** 29–44 cm long, drying 2–5 mm diam., sharply D-shaped, flattened adaxially, with adaxial margins sharp; **blades** narrowly ovate-triangular-cordate, subcoriaceous, long-acuminate at apex (the acumen sometimes apiculate), long-cordate to sagittate at base, 26–34 cm long, 11.5–14 cm wide (2–3 times longer than wide), (0.7–1.1 times longer than petiole); anterior lobe (13)21–26 cm long, (5.4)13–14 cm wide (2.9–4.3 times longer than posterior lobes); posterior lobes 6–7.5 cm long, (2.3)6–7.1 cm wide; sinus hippocrepiform, sometimes spatulate, 7–7.5 cm deep; midrib broadly convex and paler above, convex and darker below, drying somewhat acute; basal veins 3(4) per side, with 0(1) free to base, the remainder coalesced 2–2.5 cm; posterior rib naked for most or all of its length; primary lateral veins 3–4 per side, departing midrib at a 55–65° angle, ± straight to weakly arcuate to the margins; minor veins arising from the midrib only. **INFLORESCENCES** 1 per axil; peduncle 3–14 cm long, 4–5 mm diam., green or tinged with reddish violet; **spathe** 10–18 cm long ((0.7)2–3.7 times longer than peduncle), reddish violet outside; spathe blade acuminate; spathe tube cylindrical, 8–9 cm long, 1–1.5 cm diam., crimson-red inside; **spadix** stipitate to 3.5 cm long; acute at apex, 15–16.5 cm long, constricted ca. 4 cm above base of sterile staminate portion; pistillate portion cream, 3.2 cm long in front, 1.7 cm long in back, 1.5–2.6 cm wide at base; staminate portion 8.3–9 cm long; fertile staminate portion cream, tapered to apex, 7–10 mm diam. at base and midway, 3–4 mm diam. ca. 1 cm from apex; sterile staminate portion 1.8–2.2 mm diam.; pistils 1.9–2.3 mm long; ovary 4–6-locular, 1 mm diam., with sub-basal placentation; locules 0.9 mm long, ca. 0.3 mm diam.; ovule sac 0.4–1 mm long; ovules 2 per locule, contained within transparent, gelatinous ovule sac, longer than funicle; funicle 0.2–0.3 mm long (can be pulled free to base), style 0.8 mm diam., similar to style type D; style apex somewhat rounded to flat, depressions surrounding stylar canal exits, style boss broad, very shallow; the androecium truncate, prismatic, ± oblong, margins irregularly 5–6-sided, 0.8–1.1 mm long; thecae oblong, 0.4–0.5 mm wide, ± parallel to one another, very nearly contiguous; sterile staminate flowers prismatic to weakly clavate, irregularly 4–6-sided, mostly 5-sided, 1.3–1.9 mm long.

Flowering in *Philodendron verapazense* occurs in April and May, with a single, post-anthesis collection known from July.

*Philodendron verapazense* ranges from southern Mexico (Chiapas) to Guatemala (Alta Verapaz), at 700 to 1525 m elevation in "Bosque Pino-Encino" in Mexico and *Tropical moist forest* in Guatemala.

*Philodendron verapazense* is a member of *P. sect. Calostigma* subsect. *Macrobelyum* ser. *Macrobelyum*. This species is recognized by its scandent habit with the internodes longer than broad and conspicuously pluri-sulcate, as well as by its sharply two-ribbed, deciduous cataphylls, sharply D-shaped petioles, narrowly ovate-triangular-cordate blades, solitary inflorescences, red to reddish violet spathe tubes, and greenish spathe blades.

*Philodendron verapazense* is probably most similar to *P. sagittifolium* especially in the shape and coloration of the blades, the sharply two-ribbed, deciduous cataphylls, and the D-shaped petioles. It differs from the latter species principally in its deeply sulcate stems, sharply D-shaped, shorter (less than 50 cm long) petioles, and leaf blades with 3–4 pairs of primary lateral veins (vs. subterete, longer petioles, which are greater than 50 cm long, and 4–8 pairs of primary lateral veins for *P. sagittifolium*).

*Additional specimens examined.* GUATEMALA. **Alta Verapaz:** near Chirriacté, on the Petén Highway, ca. 900 m. *Standley 91978* (F). **Izabal:** vic. EXMIBAL camp 2, NW of Lake Izabal, 400–500 m, *Jones & Facey 3354* (NY). MEXICO. **Chiapas:** Los Lagos, 3 mi. NW of Rancho San José, 5000 ft., *Carlson 1846* (F).

***Philodendron verrucosum* L. Mathieu ex Schott,**  
Syn. Aroid. 85. 1856. TYPE: Locality unknown (holotype, W? lost?). Schott ic. 2757–2759 (neotype, here designated). Figures 437–440, 445.

*Philodendron daguense* Linden & André, III. Hort. 18: 192, t. 79. 1871. TYPE: Colombia. Valle: Río Dagua, *Wallis s.n.* (holotype, K).

*Philodendron pilatonense* Engl., Bot. Jahrb. Syst. 37: 129. 1905. TYPE: Ecuador. Pichincha: Río Pilatón, May 1899, *Sodiro s.n.* (holotype, B; isotypes, G, MO).

*Philodendron discolor* K. Krause, Notizbl. Bot. Gard. Berlin-Dahlem 9: 273. 1925. TYPE: Peru. Junín: Prov. Jauja, in the valley of Río Masamerich (Río Pontachucla) confluent of Río Pangoa, above Rasthütte Calabaza, 1500–1600 m, 11°30'S, 7 May 1913, *Weberbauer 6663* (holotype, B; isotype, F).

Usually hemiepiphytic, rarely terrestrial or epiphytic; stem appressed-climbing, densely scaly or setose, trichomes green to greenish white; pre-adult internodes to 20 cm long, 1–2 cm diam.; adult internodes smooth, scurfy, minutely wrinkled, semi-glossy to matte, 3–10 cm long, 2–6 cm diam., about as long as broad or longer than broad, gray to brown, roots moderately numerous, mostly short,

covered at least on one side with fine root hairs, drying dark brown, semiglossy, faintly appressed-scaly; cataphylls 10–30 cm long, unribbed, sometimes bluntly 1-ribbed, green or reddish, densely scaly or setose, persisting as a tangled network of fibers. LEAVES erect-spreading; **petioles** 33–65(90) cm long, 3 cm diam. at base, 1.5 cm diam. at apex, subterete, purplish violet to brownish, surface densely scaly, the scales of two distinct types, short, broad, often lacerate scales interspersed with long acicular scales, the latter erect, or spreading then erect; **blades** broadly ovate-cordate, thinly coriaceous, moderately bicolorous, acuminate to narrow acuminate at apex (the acumen mostly inrolled, 1–2.5 mm long), cordate at base, 28–75 cm long, 19–60 cm wide (1–1.7 times longer than wide), (0.7–1.5 times longer than petiole), broadest below point of petiole attachment, upper surface dark green or bronze-green (blackish green on new leaves), velvety to matte, drying dark brown, yellow-green or gray-green, lower surface pale green tinged red-purple between secondary veins (weakly glossy and purple-violet on new leaves), matte, drying yellow-brown to reddish brown; anterior lobe 9.6–51 cm long, 10–59 cm wide (1.9–3 times longer than posterior lobes); posterior lobes 8–21 cm long, 5–28 cm wide, broadly rounded to obtuse; sinus hippocrepiform to obovate or closed; midrib flat to sunken, slightly paler than surface above, convex, matte, darker than surface below; basal veins 6–8 per side, with (0)1–2 free to base, numbers 3–4 coalesced 1–5 cm; posterior rib not at all naked on small blades or naked for 0.5–6 cm; primary lateral veins 3–6(8) per side, departing midrib at a 50–55° angle, sunken, paler than surface above, round-raised to convex, darker than surface below; minor veins distinct and darker than surface below, arising from both the midrib and primary lateral veins; “cross-veins” conspicuous, in part raised below. INFLORESCENCES 1–4 per axil; peduncles 5–25 cm long, 1–2 cm diam., medium green to purplish, usually 0.66–1.75 times longer than the spathe; **spathe** densely scaly or setose, 10–22 cm long, 4 cm diam.; spathe blade medium green, whitish or reddish green outside, white to pinkish inside; spathe tube reddish green, medium green, or dull purple-violet outside, red or pale reddish (dark green post-anthesis) inside; **spadix** 14.6 cm long; pistillate portion cylindrical to weakly clavate, 1.8–4.6 cm long, 1 cm diam. at apex, 8 mm wide at base; staminate portion 9.4 cm long; fertile staminate portion cream; pistils 2.6–3.3 mm long, 1.4–1.7 mm diam.; ovary 4–5-locular, 1.9–2.5 mm long, 1.4–1.7 mm diam., with axile placentation; locules 1.9–2.5 mm long, 0.5–0.7 mm diam.; ovule

sac 1.8–2.2 mm long; ovules 20–24(34) per locule, 2-seriate, contained within translucent, gelatinous ovule sac, 0.1–0.2 mm long, as long as or longer than funicle; funicle to 0.2 mm long, adnate to lower part of partition, style 0.5–0.6 mm long, 1.4–1.7 mm diam., similar to style type B; style apex  $\pm$  flat; stigma  $\pm$  lobed, 1.4 mm diam., 0.2–0.3 mm high, covering entire style apex; the androecium truncate, prismatic, oblong, margins  $\pm$  4–6-sided, 0.8–0.9 mm long, 2–2.5 mm diam. at apex; thecae oblong, 0.3–0.4 mm wide,  $\pm$  parallel to one another; sterile staminate flowers blunt, irregularly 4–5-sided, sometimes clavate or prismatic, 1.6–1.8 mm long, 1.4–1.7 and 1.9–3.4 mm wide. INFRUDESCENCE with ripe berries white.

Flowering in *Philodendron verrucosum* occurs from April through June (also September and November) in Central America. The species probably also flowers earlier in the dry season, perhaps as early as February, as suggested by the fact that a post-anthesis collection was made in March (and even in February in Colombia). Post-anthesis collections have also been made from April through July (but also in December). Immature fruits have been seen from July, October, and December. The flowering collections from September and October, as well as the immature fruits from October and December, appear to reflect bimodal flowering. On the other hand, no flowers were seen from July and no flowers or fruits from August.

*Philodendron verrucosum* ranges from Costa Rica to Peru at 200 to 1500 (mostly above 500) m elevation, mostly in *Premontane rain forest* and *Tropical Lower Montane rain forest* but also in *Tropical wet forest*. In Central America, it occurs principally on the Atlantic slope or near the Continental Divide in Costa Rica and Panama but also on the Pacific slope in southwestern Costa Rica. In South America, it ranges along the Pacific slope of the Andes to as far south as El Oro and Cotopaxi Provinces in Ecuador, but also occurs on the eastern slopes of the Andes in Napo and Morona-Santiago as well as in Peru in the departments of San Martín and Junín at 750 to 1850 m elevation.

*Philodendron verrucosum* is a member of *P. sect. Philodendron* subsect. *Achyropodium*. This species is characterized by its short internodes, subterete petioles (about as long as the blades), broadly ovate-cordate blades velvety on the upper surface, and especially by the densely scaly or setose vestiture of the stems, cataphylls, petioles, and inflorescences.

*Philodendron verrucosum* might be confused with a few other species that have scaly parts, including

*P. hammelii*, *P. malesevichiae*, *P. squamipetiolatum*, and especially *P. squamicaulis*. (See under the individual species for discussion of the differences.) *Philodendron squamipetiolatum* differs by its longer, more slender internodes and much smaller leaf blades (ca. 30 cm long on flowering plants) with a semi-glossy upper surface. It also differs in having solitary inflorescences (vs. usually more than one per axil in *P. verrucosum*). *Philodendron hammelii* differs in its much smaller size, petioles with foliaceous (rather than acicular) scales restricted to near the apex, and glabrous cataphylls, peduncles, and spathes. *Philodendron malesevichiae* differs by its terrestrial, creeping habit; persistent, semi-intact, glabrous cataphylls; narrowly sulcate petioles (rather than subterete as in *P. verrucosum*); and semi-glossy blades lacking the conspicuous "cross-veins" of *P. verrucosum*.

**Additional specimens examined.** COSTA RICA. **Alajuela:** Salto La Paz, 1350 m, *Ferreya 15788* (USM); San Ramón-Balsa, ca. 2.3 km N Río Balsa, ca. 1050–1150 m, 10°11'N, 84°30'W, *Stevens 14194* (MO); ca. 4.6 km N of Río Balsa, 900–1000 m, 10°12'N, 84°31'W, *Stevens 13762* (CR, MO); 4–7 km N of Balsa, ca. 750 m, 10°13'N, 84°32'W, *Liesner & Judziewicz 14763* (MO); 2.5 km N of Balsa, 1050–1100 m, 10°11'N, 84°30'W, *Liesner & Judziewicz 14726* (CR, MO); Finca Los Ensayos, ca. 11 mi. NW of Zarcero, ca. 850 m, *Croat 43636* (MO); 2 km N of Angeles Norte de San Ramón, 4000 ft., *Luteyn 3691* (MO); Zarcero-Quesada, 11.8 km past Zarcero, 1230 m, 10°12'N, 84°23'W, *Hoover 1352* (CR, MO); E of Río San Rafael, W of La Marina, 500 m, 10°23'N, 84°38'W, *Burger & Stolte 5041* (CR, US); Monteverde Biological Reserve, Río Peñas Blancas, 900 m, 10°18'N, 84°45'W, *Haber & Bello 8458* (CR, MO); 820 m, *Bello 2672* (CR); 850–900 m, 4478 (CR, INB); 950 m, *Bello & Méndez 2639* (CR, INB); Cuntón Alfaro Ruiz, La Peña de Zarcero, *Smith 961* (F); Los Chiles-Los Angeles, San Ramón, 500–1200 m, 10°10'N, 84°30'W, *Barringer et al. 2200* (CR, MO); *E. Schmidt 306* (CR). **Cartago:** 4000 ft., *J. D. Smith 391* (G); Moravia-Quebrada Plantainillo, Moravia, 3–5 km from Finca Racine, 1200–1300 m, *Croat 36623* (MO); 1200–1400 m, 36795 (MO); Turrialba-Limón, along Hwy. 32, ca. 11 mi. NE of Turrialba, 850 m, *Croat 43355* (CR, MO); 31 km S of Siquires on road to Turrialba (CR-101), 850 m, 9°57'N, 83°36'W, *Thompson & Rowlands 1163* (CM); El Mañeco, S of Río Navarro, 1400–1500 m, *Standley & Torres 51745* (CM, US); *Standley 33425* (US); Finca Navarro, 1350 m, *Maxon 636* (NY, US); 10 km S of Tapantí, 1600 m, 9°42'N, 83°46'W, *Burger & Burger 8424* (CR, F); 3.5 km SE of Tapantí, 1250 m, *Lent 851* (F, NY, US); Río Grande de Orovi, 2 km SE of Tapantí, 1350, *Lent 763* (F); 14 mi. from Turrialba, road to Pavones, *Carlson 3244* (F); Orovi, *Standley 39736* (US); La Estrella, *Standley 39547* (US); 4250 ft., *J. D. Smith 5963* (US); Río Gato, 4 km SE of Pejibaye, 700 m, 9°48'N, 83°42'W, *Liesner 14412* (MO); Tapantí Hydroelectric Reserve, 3–5 km from the gate, *D. Smith & Taylor 1009* (MO); 1300–1800 m, *Gómez 18889* (MO, RSA); Tapantí region, 1.5 km past Río Macho, along small stream along road to Humo, 1330 m, 9°17'N, 83°50'W, *Hoover 1345* (CR, MO); 4250 ft., *Cooper 5963* (CM, GH, US); Turcu-

rique, Las Vueltas, 635–700 m, *Tonduz 13321* (US); Reventazón Valley, Juan Viñas, *Cook & Doyle 288* (US). **Heredia:** 4 mi. N of Vara Blanca, 1350 m, *Croat 35669* (MO); 3 mi. S of Cariblanco, 760 m, *Croat 35794* (MO); La Selva Field Station, 280–300 m, 10°21'N, 84°03'W, *Schatz & Grayum 721* (DUKE). **Limón:** Cerro Chimú-Cerro Matama, 1200 m, *Gómez & Herrera 23533* (MO, US); Talamanca, Tsaki, 200 m, *Tonduz 9512* (CR, US); Fila Matama, 850 m, *Robles & Chacón 2724* (CR, MO). **Puntarenas:** Las Cruces Botanical Garden-Río Jaba, ca. 3 km SE of San Vito de Coto Brus, ca. 1050–1200 m, 8°47'N, 82°58'W, *Grayum 5621* (CR, MO); Monteverde Reserve, 1500 m, 10°17'N, 84°48'W, *Hammel & Zuchowski 13886* (MO). **San José:** San Isidro del General-Dominical, 4.8 mi. from Río Pacuare, 1000 m, *Croat 35251* (MO); 9 mi. SW of Río Pacuar, 680 m, *Croat 35349* (MO); Bijagua, 1300 m, *Gómez 20564* (M, MO, QCA); Alto San Juan, road to Dominical, 900 m, *Molina et al. 18096* (F, NY); La Honduras, 1300–1700 m, *Standley 37783* (US); Río Claro Valley (Bajo La Honduras), below La Palma, NE of San Jerónimo, 1000–1200 m, 10°3'N, 83°58'W, *Burger et al. 9433* (CR, F, U); below La Palma NE of San Jerónimo, 1000–1200 m, 10°3'N, 83°58'W, *Burger et al. 9286* (F, MO), *Burger & Genay 9077* (CR, F); 1500 m, 10°3'N, 83°58'W, *Burger & Stolte 5308* (CR, F), 8499 (F); Tarrazá, vic. Hormiguero, 1100–1200 m, *Croat 78935* (INB, MO). **PANAMA. Bocas del Toro:** Cerro Colorado, 9.2 mi. W of Chame, 1450–1480 m, 8°35'N, 81°50'W, *Croat 69010* (MO); Fortuna Dam area, Chiriquí Grande-Gualaca, 21.4 km past Gualaca, 8°32'N, 82°19'W, *Hoover 1327* (MO); Continental Divide, ca. 1200 m, ca. 8°45'N, 82°15'W, *McPherson 9033* (CAS, CM, K, MO); *Croat 60374* (MO); Old Bank Island, vic. of Chiriquí Lagoon, von Wedel 1923 (MO). **Chiriquí:** Volcán Cañas Gordas, 1 m E of Cañas Gordas, *Croat 22344* (MO); Cerro Colorado, 34.1 km beyond Río San Félix, 1390 m, *Croat 37254* (MO); Gualaca-Chiriquí Grande, vic. IRHE headquarters, 1200 m, 8°45'N, 82°18'W, *Croat 66719* (MO); 66558 (AAU, MEXU, MO, NY, PMA); road to Finca Landau, 1100 m, *Correa et al. 2141* (F, MO, PMA); NW of confluence of Río Hornito and Río Chiriquí, 1050–1100 m, 8°44'N, 82°13'W, *Sytma & Stevens 2234* (MO, PMA); Fortuna Dam, Cerro Fortuna, lower slopes, 1150 m, *Knapp & Vodička 5576* (MO); behind Vivero Forestal, 12 km E of Los Planes de Hornito, 1200–1300 m, 8°45'N, 82°12'W, *Knapp 4949* (MO); 7 km SE of Fortuna Dam, 1100 m, *O'Connor 91-512-004* (MICH); Continental Divide, 1200–1500 m, 8°47'N, 82°13'W, *Churchill 5298* (MO); 1200 m, *O'Connor 91-511-003* (MICH); Continental Divide trail, 1200–1300 m, 8°45'N, 82°15'W, *Thompson 4954* (CM, MO). **Coclé:** El Copé region, Alto Calvario, near Continental Divide, 5 mi. N of El Copé, 900–1000 m, 8°39'N, 80°36'W, *Croat 75058* (MO); 750–900 m, 44736 (MO); 8°39'N, 80°36'W, 68768 (CM, F, MO); El Valle region, La Mesa, *Gentry 5632* (MO, PMA); 860–900 m, *Croat 37339* (MO); *J3341* (MO); 900 m, *Sullivan 544* (MO); *Liesner 756* (MO); NE slopes of Cerro Caracoral, N rim of El Valle, 2700–3200 ft., *Sytma 3787* (MO); Cerro Pílon, 5 km NE of El Valle, 800–1045 m, *Mori et al. 6633* (MO); *Lallathin 5096* (MO); Cerro Gaital, 900–1000 m, 8°40'N, 80°07'W, *Knapp 5301* (MO); 800–900 m, *Knapp & Dressler 4913* (MO); 5.6 km S of El Valle, 870 m, 8°50'N, 80°07'W, *Hoover 1319* (MO, NY); ca. 3 km N of El Valle de Antón, 850 m, *Wilbur et al. 15656* (MO). **Darién:** Cerro Pirre region, 1000–1100 m, *Hartman 4829* (MO); 9–10 km N of Alto de Nique, 1520–1560 m, *Croat 37871* (MO); Río Tuquesa, vic. of Tyler

Kittredge gold mine, ca. 2 air km from Continental Divide, *Croat* 27215 (MO). **Panamá:** El Llano-Cartí, 12 mi. above Pan-American Highway, 200–500 m, *Liesner* 689 (MO); Campo Tres, 3 mi. NE of Altos de Pacora, 500–800 m, *Croat* 22716 (MO); Cerro Campana, 1000 m, *Nee & Stockwell* 11625 (MO); *Croat* 17200 (MO); ca. 850 m, 8°40'N, 79°50'W, *Thompson* 4576 (MO); 850 m, *Busey* 875 (MO); above Su Lin Motel, *Croat* 14728 (MO); near Florida State Univ. Building, *Croat* 12121 (F, MO, SCZ). **San Blas:** Cerro Habá, vic. of peak, 2500 ft., 9°23'N, 78°49'W, *Sytama et al.* 2699 (MO, US); Cerro Brewster, 850 m, 9°18'N, 79°16'W, *de Nevers et al.* 5546 (MO); Cerro Obu, 400–500 m, *de Nevers et al.* 8053 (MO); Nusagandi, 310 m, 9°20'N, 79°W, *Croat & Zhu* 76993A (MO). **Veraguas:** Santa Fe region, NW of Santa Fe, 11 km from Alto Piedra, Río Dos Bocas Valley, 450–550 m, *Mori et al.* 3815 (MO, PMA); NW of Santa Fe, along W fork of road beyond Alto Piedra, 0.6 mi. beyond fork in the road, 1300–1350 m, *Croat* 49060 (MO); Santa Fe–Río Calovébora, 0.6 mi. beyond Alto Piedra, 735 m, *Croat & Folsom* 33991 (MO); 1.7 mi. past Alto Piedra, 1.5 mi. beyond Quebrada Cosilla, 570 m, 8°33'N, 81°08'W, *Croat & Zhu* 76856 (MO).

***Philodendron warszewiczii*** K. Koch & Bouché, in A. Braun et al., *Append. gen. sp. Hort. berol.* 1855: 4. 1855–1856. TYPE: Guatemala. San José and Florida, *Warszewicz s.n.* (holotype, B? lost?). Guatemala. Santa Rosa: Volcán Jumaytepeque, 6000 ft., *Hyde & Lux* 4282 (neotype, here designated, K). Figures 16, 441–444, 446–449.

Usually terrestrial, or on rocks, sometimes epiphytic; stem appressed-climbing, thick, succulent, bare, leaf scars conspicuous, 1.5–2.6 cm long, 1.5–2 cm wide; internodes short near apex, semiglossy, glaucous, 3–5(15) cm long, 2.5–7(10) cm diam., often longer than broad lower down, dark green to gray-brown; roots pale green to dark brown, few per node, epidermis drying yellowish; cataphylls thin, soft, 18–33 cm long, weakly to sharply 2-ribbed, pale green to whitish, lightly and densely short-lineate, deciduous intact. LEAVES often deciduous in dry season; **petioles** 33–58(80) cm long, 1–2 cm diam., subterete to C-shaped, moderately soft, weakly flattened to rounded with thick, medial rib adaxially, with adaxial margins sharply raised, surface sharply striate, with a dark green ring around apex; **blades** triangular-sagittate in outline, deeply bipinnately or bipinnatisect to within 1–4 cm of the midrib, thin, semiglossy, moderately bicolorous, ± rounded at apex (the acumen 2 mm long), cordate at base, 31–78 cm long, 30–62 cm wide (0.9–1.3 times longer than wide), (1–1.3 times longer than petiole), upper surface moderately glossy, moderately paler, margins sinuate; median lobe 17–52 cm long, 23–63 cm wide, rounded to obtuse; lateral segments 1–30 cm long,

11–28 cm wide, pinnately lobed with 3 or more lobes per side, acute or acuminate; interlobal sinuses 0.78–0.94 the length of the blades; midrib flat, obscurely striate, slightly paler than surface above, raised, obscurely striate, slightly paler than surface below; basal veins 2–6 per side, with 0–1 free to base, second and third pair coalesced 2–5 cm, third and higher order veins coalesced 5–7 cm; posterior rib naked along most of its margin; primary lateral veins 3–6 per side, departing midrib at a 55–70° angle, straight to the margins, weakly raised above, raised, paler than surface below; tertiary veins raised, paler than surface above and below; minor veins visible, darker than surface below, distinctly visible on drying, arising from both the midrib and primary lateral veins. INFLORESCENCES 1(3) per axil; peduncle (2.5–3)4.5–11(16) cm long, 1–2 cm diam., dark green, dark short-lineate; **spathe** 13.5–30 cm long, 4–7.5 cm diam. (1.1–3.6(4.7) times longer than peduncle), medium to dark green throughout, blunt at apex, scarcely constricted; spathe blade short dark green lineate outside, 12 cm long, pale green to white inside; spathe tube weakly short dark green lineate outside, 14 cm long, pale green to white or purple or red inside; **spadix** sessile; white throughout, 14–24(30) cm long, constricted weakly above pistillate portion; pistillate portion 3 cm long in front, 2 cm long in back, 1.3 cm diam. at apex, 1.7 cm diam. at middle, 1.6 cm wide at base; staminate portion 8.3–15 cm long; fertile staminate portion 1.6 cm diam. at base, 1.4 cm diam. at middle, 9 mm diam. ca. 1 cm from apex; sterile staminate portion 1.3 cm diam.; pistils (1.1)4.1 mm long, 3.5–4.5 mm diam.; ovary 4–5-locular, 2.6–9.5 mm long, 3.5–4.5 mm diam., with axile placentation; locules 2.6 mm long, 0.8–1.1 mm diam.; ovule sac 2.4 mm long; ovules 3–4 per locule, 0.5 mm long, contained in thick, translucent matrix; style 1.2–1.6 mm long, 1.2–1.4 mm diam., similar to style type E; style funnel broad, moderately deep; style apex ± flat; stigma subdiscooid, unlobed, 0.7 mm diam., 0.3 mm high, lining entire upper surface of funnel; the androecium truncate, prismatic, oblong, margins irregularly 4–6-sided, 3.3 mm long, 1.3 mm diam. at apex; thecae oblong, 0.4 mm wide, ± parallel to one another and nearly contiguous; sterile staminate flowers rounded to somewhat clavate, irregularly 4–6-sided, 2.6 mm long, 1.3–1.8 mm wide. INFRACTESCENCE brownish outside, brownish inside, to 3 cm wide pistillate spadix 3–7.5 cm long; berries white, 8–11 mm long; seeds 3–4 per locule, pale yellow, oblong-ellipsoid, 2–2.5 mm long, 1–1.2 mm diam., with pale raphide cells.

Flowering collections of *Philodendron warszew-*



*iczi* are rare, known only from May. Post-anthesis collections, concentrated between April and June (but also January, February, March, July, August, September, and December), imply that the flowering period is relatively broad, perhaps throughout the entire dry season and the first half of the rainy season. Immature fruiting collections are scattered in a somewhat bimodal pattern, one group in the mid-rainy season from July through September, the other group primarily in the early dry season from December through March (especially December and January). The immature fruits collected in November might reflect an early dry season flowering. Mature fruits are known only from this same period between December and February at the beginning of the dry season.

*Philodendron warszewiczii* ranges from Mexico (Jalisco and Chiapas) to El Salvador on the Pacific slope, and to Honduras and Nicaragua at 300 to 1900 m elevation, in *Tropical dry forest* life zones. In Mexico, the species occurs in "Selva Mediana Subperennifolia" and "Selva Baja Caducifolia."

*Philodendron warszewiczii* is a member of *P.* sect. *Polytomium*. This species is distinguished by its moderately thin, deeply dissected, bipinnatifid leaf blades and thick, succulent stems often leafless during the dry season.

*Philodendron warszewiczii* is most similar to *P. radiatum* and *P. dressleri*. It differs from *P. radiatum* in having thinner, more highly divided blades and thicker stems. Although both species occur in some of the same forest types in Mexico, *P. radiatum* always occupies more mesic sites within these zones. In other parts of Central America for which there are Holdridge Life Zone maps, *P. radiatum* is found mostly in *Tropical moist forest* or *wetter life zones*, whereas *P. warszewiczii* is known from *Tropical dry forest* areas.

Although *Philodendron warszewiczii* and *P. dressleri* occupy equally dry sites, the latter is distinguished by occurring further north in Mexico and in having the leaves divided to no more than half-way to the midrib (vs. nearly all the way to the midrib in *P. warszewiczii*).

This species occurs principally on the Pacific slope except in Nicaragua (one Honduran collection is from the Atlantic slope). This is probably due to the fact that the Continental Divide runs very near the Pacific Ocean in Nicaragua, and because there is a second, generally much higher, more centrally located range of mountains and hills that creates a rain shadow throughout most of the Pacific side of the country. Because of this, *P. war-*

*szewiczii* inhabits many sites in the central part of the country well within the Atlantic watershed.

Common names for this species include: "Guacamayo," "Cupapayo," "Mano de León," "Copapayo," "Ocopayo," and "Papaya de Monte" (Standley & Steyermark, 1958b).

*Additional specimens examined.* EL SALVADOR. S of San Salvador, *Calderón 876* (GH, US). **Ahuachapán:** Ahuachapán vicinity, 800–1000 m, *Standley 19726* (GH, NY, US). **San Salvador:** *Calderón 416* (GH, NY, US); San Salvador vicinity, 650–850 m, *Standley 19624* (NY, US). GUATEMALA. **Jalapa:** Jalapa–San Pedro Pinula, 1400–1800 m, *Standley 77055* (F). **Jutiapa:** Monjas–El Progreso, above Ovejero, ca. 1400 m, *Standley 77660* (F); near Jutiapa, ca. 900 m, 60495 (F); *Agua Blanca–Amatillo*, 950–990 m, *Steyermark 30367* (F); *Jutiapa–La Calera*, SE of Jutiapa, ca. 850 m, *Standley 76087* (F). **Santa Rosa:** near Cuidapilla, ca. 900 m, *Standley 78051* (F). **Zacapa:** upper reaches of Río Sitio Nuevo, between Santa Rosalía and first waterfall, 1200–1500 m, *Steyermark 42281* (F); Río Lima, Sierra de las Minas, Finca Alejandria, 1500–1700 m, *Steyermark s.n.* (F). HONDURAS. Río de la Orilla, SE of El Zamorano, 900 m, *Anderson & Spoehr s.n.* (MO). **Comayagua:** vic. of Comayagua, ca. 600 m, *Standley & Chacón 6021* (F). **El Paraíso:** Danlí–Los Arcos, 740–850 m, *Standley 17007* (F); near Yuscarán, 960 m, *Standley et al. 1234* (F). **Morazán:** Tegucigalpa, Guaruabuzi, 2600 ft., *von Hagen & von Hagen 1272* (F, NY); S. Antonio de Occidente, 850 m, *Valerio 3654* (F); Escuela Agrícola Panamericana, 800 m, *Croat 34814* (cultivated) (MO); San Antonio de Oriente, 4 mi. N of Zamorano, *Croat 42765* (MO, US); Río de la Orilla region, SE of El Zamorano, 900–950 m, *Standley 23205* (F), 23213 (F), 23217 (F); near El Zamorano, 800 m, *Morton 7119* (US). **Olancho:** Río Olancho, San Francisco de la Paz–Gualaco, 1–4 mi. NE of San Francisco de la Paz, 690–900 m, 14°57'N, 86°13'W, *Croat & Hannon 64186* (B, MO); W of main 8.6 mi. SW of Catacamas, 400 m, 14°45'N, 86°W, *Croat & Hannon 64136* (CM, MO). MEXICO. **Chiapas:** 20–30 mi. SW of El Jorote, on road to Motozintla, 700–900 m, *Croat 40714* (MEXU, MO); Tuxtla Gutiérrez–San Fernando, Chacón canyon, 850 m, *Breedlove 69987* (CAS); 22 mi. SE of Comitán, *Carlson 1953* (F); 6–8 km E of Frontera Comalapa, road to Ciudad Cuauhtémoc, 1000 m, *Breedlove 23428* (MO); 10 km E of Motozintla, 1110–1150 m, *García et al. 1527* (MEXU, BM); vic. Tuxtla Gutiérrez, 830 m, *Breedlove & Bourel 68505*, (CAS); Mpio. Tzimol, 15 km S of Comitán, 1200 m, *Breedlove 51028* (CAS). **Jalisco:** Autlán–Barra de Navidad, at km 291.67, ca. 300 m, *Moore & Bunting 8744* (BH); 9 mi. N of road-junction at W end of Bahía de Navidad, 300 m, *McVaugh & Koelz 1758* (MICH); Hwy. 110, 2 mi. NE of Huizache, ca. 5 m, *Stevens & Fairhurst 1823* (US); 15–16 mi. NW of Barra de Navidad, *Dressler & Wirth 2671* (UC, US); Chamela Field Station, 100 km S of Puerto Vallarta, 50 m, 19°30'N, 105°03'W, *Gentry 63966* (MO), *Bullock 1101* (MO); El Tejón, 100 m, 19°30'N, 105°03'W, *Gentry 74426* (MO); 10 km E of Las Palmas, 340 m, 20°50'N, 105°02'W, *Cochrane et al. 12022* (MO); Reserva Biosfera Sierra de Manantlán, Lagunillas–Juluapan, Cerro Grande, ca. 1400–1500 m, 19°22'N, 103°56'30"W, *Ilitis et al. 29716* (MO); Rancho Cuixmalá, Cumbres I, Arroyo Cajones, 19°27'N, 104°58'30"W, *Latt et al. 3268* (MO). **Nayarit:** Mirador de Aguila, ca. 14 mi. N of Tepic, 450–600 m, *Feddesma 824* (MICH); 6–7 mi. S of Com-

postela, 1000 m, *McVaugh 18754* (MICH); 2 mi. SE of Las Varas, road to Mazatlán, 60–90 m, *McVaugh 15351* (MICH). NICARAGUA. **Boaco:** Cerro Mombachito, 8.5 km NW of Camoapa, 1059 m, *Araquistain & Moreno 931* (MO); **Chontales:** 3.9 km N of Comalapa, ca. 460 m, ca. 12°18'N, 85°30'W, *Stevens & Montiel 17206* (MO); Jitigalpa-La Libertad, ca. 17.4 km NE of Río Mayales, 350–400 m, 12°12'N, 85°17'W, *Stevens 4023* (MO). **Estelí:** 10 km S of Estelí, 1000 m, *Neill 1186* (MO); km 163 on Hwy. 1, ca. 11.2 km N of entrance to Estelí, ca. 920 m, ca. 13°13'N, 86°23'W, *Stevens 11213* (MO); Cerro Cucamonga, 800–850 m, 10°15'N, 86°22'W, *Moreno 14120* (MO); Salto de Estanzuela, ca. 5 km S of Estelí, 1000 m, 13°02'N, 86°20'W, *Guzmán et al. 1189* (MO); Río La Estanzuela, 6 km al S de Estelí, ca. 1000 m, 13°01'N, 86°21'W, *Moreno 9751* (MO). **Jinotega:** Jinotega-San Rafael, ca. 2 mi. NW of Jinotega, 1000 m, *Croat 43020* (MO); Lago de Abanas, 1–2 mi. E of San Rafael de Norte-Jinotega Road, 1000 m, *Croat 42970* (MO); ca. 1 mi. SE of Yalí, 850 m, *Croat 42891* (MO). **Madriz:** San Lucas-Hacienda El Volcán, 2 km NE de San Lucas, 800–900 m, 13°25'N, 86°35'W, *Araquistain & Moreno 2018* (K, MO); Cerro Volcán de Somoto (Tepesomoto), 900–1300 m, ca. 13°25'N, 86°35'W, *Stevens & Grijalva 16389* (MO); *Araquistain & Moreno 2046* (MO); ca. 7 mi. S of Ocotal, along Highway 15, 650 m, *Croat 42797* (MO). **Matagalpa:** 5–8 mi. E of Matagalpa, 800–1000 m, *Williams et al. 24048* (F); Cerro Apante, 1000–1400 m, *Moreno 155* (MO); 1 km SE of Matagalpa, 800 m, *Vincelli 297* (MO). **Nueva Segovia:** Cerro El Achote, 7 km N of El Jicaró, 700–1000 m, 13°46'N, 86°08'W, *Araquistain & Moreno 2316* (MO, NY); Quebrada El Nancital, ca. 6.2 km N of Ocotal, Quebrada El Nancital, 700–760 m, ca. 13°41'N, 86°24'W, *Stevens 3040* (BM, MO); 6.5 km W of Ocotal, road to Las Manos, ca. 700 m, 13°40'N, 86°29'W, *Moreno 14382* (MO); El Jicaró, "Casas Viejas," ca. 600 m, 13°44'N, 86°05'W, *Moreno 13504* (MO); El Jicaró, "El Cinchado," ca. 650–700 m, 13°46'N, 86°05'W, *Moreno 13589* (MO); El Jicaró, "Río Grande," road to Murra, 460–500 m, 13°43'N, 86°W, *Moreno 13670* (MO).

**Philodendron wendlandii** Schott, Prod. Syst. Aroid. 221. 1860. TYPE: Costa Rica. *Wendland s.n.* (holotype, W? lost). Schott ic. 2079 (neotype, here designated). Figures 15, 450, 451, 453.

Epiphytic, sometimes occurring high in the canopy, rarely terrestrial, perhaps by accident; stem short, scurfy, leaf scars inconspicuous, 1.3–2.1 cm long, 2–2.5 cm wide, obscured by root mass; internodes short, 0.5–3 cm long, 1.8–3.5 cm diam., glaucous, epidermis fissured transversely; roots reddish green becoming dark brown, blunt, branched near tip; cataphylls somewhat spongy or fleshy, 29–36 cm long, unribbed or sometimes sharply 2-ribbed, green to reddish green, persisting semi-intact at upper nodes, eventually fibrous. LEAVES rosulate; **petioles** 9–30 cm long, 1.5–4 cm diam., broader than thick, spongy, soft, sharply flattened adaxially, sharply convex abaxially, with adaxial margins erect on younger petioles, with dark green ring at apex; **blades** oblong-oblanco-

late, coriaceous, slightly bicolorous, semiglossy, acuminate, sometimes long-acuminate at apex (the acumen inrolled, 2–4 mm long),  $\pm$  narrowly rounded to obtuse at base, 32–67 cm long, 8.5–22 cm wide (2.6–4.7 times longer than wide), ((1.7)2.6–4.2(5.2) times longer than petiole), broadest  $\pm$  near the middle; midrib flat to convex, broadly convex near base, convex toward apex, concolorous above, round-raised to convex, slightly paler than surface below; basal veins lacking; primary lateral veins 7–12(16) per side, departing midrib at a (50)60–70° angle,  $\pm$  straight to the margins, weakly impressed to sunken above, convex to weakly raised, darker than surface below; minor veins obscure above, visible, darker than surface below, arising from midrib and paralleling primary lateral veins. INFLORESCENCES erect, 2 per axil; peduncle 3–10 cm long, 1–1.5 cm diam., pale green; **spathe** erect to erect-spreading, coriaceous, 12–18 cm long ((1.5)3.4–3.6(5.3) times longer than peduncle), constricted just above the tube to 2.1 cm diam. at constriction; spathe blade oblong-ovate, pale green to white, sometimes tinged pink, sometimes white-striate outside, 7–13 cm long, 2.6–7 cm diam. when open (opening broadly elliptic in face view, white (rarely rose-red)), pale-spotted with dark green central rib inside; spathe tube ellipsoid, pale to medium green, sometimes tinged with red, white-short-lineate in back outside, (3.5)5.5–7 cm long, 2.6–3.5 cm diam., pale green to white (sometimes creamy yellow or reddish with darker green central rib), minutely pale spotted inside; **spadix** stipitate to 1.3 mm long, light green; cylindrical, bluntly pointed at apex, (8.5)11–14.4(17.2) cm long, of nearly uniform width; pistillate portion pale green to yellowish green, cylindrical, or weakly tapered toward apex, 2.9–4.6(6.7) cm long, 1–1.2 cm diam. at apex, 1.1–1.4 cm diam. at middle, 1–1.5 cm wide at base; staminate portion 8.2–11.7 cm long; fertile staminate portion creamy white, with resin droplets, 9–13 mm diam. at base, 9–12 mm diam. at middle, 6–7(9) mm diam. ca. 1 cm from apex, broadest at the base or equally broad at base and middle, as broad as the pistillate portion, as broad as to slightly narrower than the sterile portion; sterile staminate portion slightly broader than the pistillate portion, white, 1–1.3 cm diam.; pistils 1.7–2.3 mm long, 1.4–1.7 mm diam.; ovary 6–7-locular, 0.7–1.3 mm long, 1.4–1.7 mm diam., with sub-basal placentation; locules 0.7–1.3 mm long, (0.2)0.4–0.5 mm diam.; ovule sac (0.7)1–1.1 mm long; ovules 2 per locule, translucent and contained within translucent or transparent ovule sac, 0.3–0.5 mm long, longer than funicle; funicle 0.2–0.4 mm long (can be pulled free to base), style 0.6–0.8 mm

long, 1.2–1.8 mm diam., similar to style type B; style apex sloping to weakly rounded, with depressions surrounding stylar canal exits; stigma subdiscoid, truncate, 0.7–1.1 mm diam., 0.2–0.3 mm high, covering center of style apex; the androecium truncate, prismatic, margins irregularly 3–6-sided, 1.2 mm long, 0.7–1.4 mm diam. at apex; thecae  $\pm$  ovate, 0.5 mm wide,  $\pm$  parallel to one another, sometimes nearly contiguous, sometimes distinctly contiguous; sterile staminate flowers blunt, irregularly 4–6-sided, sometimes clavate or prismatic, 1.4–1.6 mm long, 1.1–2(3.6) mm wide. Berries bright orange.

Flowering in *Philodendron wendlandii* occurs from the mid-dry season to the first part of the rainy season, with Costa Rican populations having been found in flower earlier (February and March) than those in Panama (May and June). Post-anthesis inflorescences have been collected between February and July, with an outlying collection from November. Immature fruits have been collected only in June and August, and mature fruits in September.

*Philodendron wendlandii* ranges from Nicaragua to Panama, from 10 to 1250 m elevation in *Tropical wet forest* and *Premontane rain forest* life zones. Most collections are from below 800 m elevation.

*Philodendron wendlandii* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Glossophyllum*. This species is distinguished by its true epiphytic habit, short stem, and rosulate, oblong-oblancheolate, short-petiole leaves but especially by the petioles, which are often broader than thick and sharply flattened adaxially. *Philodendron wendlandii* is one of relatively few truly epiphytic, rosulate "bird's nest" species in the genus.

*Philodendron wendlandii* is apparently related to several species of more or less rosulate habit and with elongate leaf blades and a green or purple line of demarcation between the petiole and the blade. These include *Philodendron annulatum*, *P. auriculatum*, and *P. ligulatum*. With the exception of *P. ligulatum* var. *heracleoanum*, all of these taxa differ from *P. wendlandii* in having more or less terete petioles. *Philodendron ligulatum* var. *heracleoanum* itself differs in being a long-stemmed vine (vs. rosulate in *P. wendlandii*), and in having D-shaped petioles scarcely or not at all wider than thick (vs. sharply flattened and conspicuously broader than thick for *P. wendlandii*) and undulate-winged margins.

*Additional specimens examined.* COSTA RICA. **Alajuela:** near La Laguna, 6–8 km S of Villa Quesada, 1200 m, *Molina et al.* 17553 (EAP, F, NY, US); Cañas-Upala, near Río Zapote, 1.8–2.7 km S of Río Canalete, 100 m,

*Croat* 36376 (MO); 8 km NE of Villa Quesada, 600 m, *Croat* 46925 (MO); Río Frío, near Los Chiles, 30–40 m, 11°02'N, 84°44'W, *Holm & Iltis* 826 (A, GH). **Cartago:** Moravia–Quebrada Platanillo, ca. 1250 m, *Croat* 36683 (MO); 4 km S of Tzurrique, Río Vueltas, 690 m, *Leut* 1096 (F). **Guacacaste:** El Silencio, near Tilarán, ca. 750 m, *Standley & Valerio* 44785 (US). **Heredia:** 3 mi. S of Cariblanco, 760 m, *Croat* 35833 (MO); S of Río Sarapiquí, opposite Chilamate, 60–100 m, 10°27'N, 84°04'W, *Grayum et al.* 5311 (MO); La Selva Field Station, Puerto Viejo de Sarapiquí, 100–150 m, *Croat* 44313 (MO); ca. 100 m, *Hammel* 9086 (F, MO); *McDowell* 943 (MO); *Folsom* 9596 (TEX); *Hammel* 8222 (DUKE). **Limón:** Finca La Suerte, *Davidson* 3265 (MO); Hacienda Tapasco–Hacienda La Suerte, 29 air km W of Tortuguero, 40 m, 10°30'N, 83°47'W, *Davidson & Donahue* 8599 (MO, RSA); *Croat* 69732 (CM, MO); Hitoy-Cerere, Quebrada Canahrit, Baja Talamanca, 500–800 m, *Gómez* 24081 (MO); Amubir, along Río Lari, 200 m, 9°29'40"N, 82°89'40"W, *Herrera* 2993 (AAU, CR, MO, US); Guápiles, 850 ft., *J. D. Smith* 4977 (US); Cerro Coronel, E of Laguna Danto, 20–170 m, 10°41'N, 83°38'W, *Stevens* 23863 (CR, MO); E of Laguna Danto, SW towards hills E of Río Zapote, 30–170 m, 10°40–41'N, 83°38–39'W, *Stevens et al.* 24988 (CR, MO); E of Río Zapote, 1 km from Río Colorado, 10–40 m, 10°40'N, 83°40'W, *Stevens & Montiel* 24313 (CR, MO); *Stevens et al.* 24683 (CR, MO); between Río Madre de Dios and Quebrada Cañahrit, 380–440 m, 10°02'N, 83°26'W, *Grayum et al.* 8712 (CR); Parque Nacional Tortuguero, NE of La Aurora, Guápiles, along Río Sierpe, 30 m, 10°22'N, 83°31'W, *Robles* 2254 (CR, MO). **San José:** Carrillo Station, 700 m, *Gómez et al.* 21192 (CR, MO). **NICARAGUA. Río San Juan:** Bocas de Sábalo, 70–100 m, 11°03'N, 84°27'W, *Morero* 26780 (MO). **PANAMA. Bocas del Toro:** Fortuna Dam Area, Gualaca–Chiriquí Grande, 9.4 mi. N of the Continental Divide, 175 m, 8°46'N, 82°16'W, *Croat* 66819 (K, MO); 2.8 mi. N of Continental Divide, 850–950 m, ca. 8°45'N, 82°15'W, *McPherson* 9653 (L, MO, PMA, US). **Colón:** Cerro Bruja from Río Escandaloso, 2000 ft., *Hammel* 3205 (MO); Santa Rita Ridge, Mile 10, *Porter et al.* 4740 (MO); Mile 6.5, 370 m, 9°21'15"N, 79°44'W, *Croat & Zhu* 76962 (MO). **Panamá:** Cerro Jefe region, Campo Tres, 3 mi. NE of Altos de Pacora, 500–800 m, *Croat* 22721 (MO); 3–3.5 mi. NE of Altos de Pacora, 7.8–8.2 mi. above Pan-American Highway, 700–750 m, 9°15'N, 79°25'W, *Croat* 68633 (MO); El Llano–Cartí, 10 mi. from main gate near El Llano, *Croat* 33715 (CM, MO); Mile 10.1, Nussagandi, 300 m, 9°20'N, 79°W, *Croat & Zhu* 76547 (MO). **Veraguas:** Santa Fe region, Santa Fe–Río Calovébora, 0.6 mi. N of Alto Piedra, 735 m, *Croat* 34141 (MO); Santa Fe–Calovébora, 1.7 mi. past Alto Piedra, 1.5 mi. beyond Quebrada Costilla, 570 m, 8°33'N, 81°08'W, *Croat & Zhu* 76820 (F, MO).

***Philodendron wilburii* Croat & Grayum, sp. nov.**

**TYPE:** Costa Rica. Alajuela: Cordillera de Tilarán, San Ramón–Bajo Rodríguez, vic. La Balsa, 8.9 mi. NW of center of San Ramón, 1100 m, 10°10'30"N, 84°30'W, 26 Sep. 1987, *Croat* 68083 (holotype, MO–3641132; isotypes, CR, K, US). Figures 452, 454–458.

Planta plerumque hemiepiphytica; caulis scandens; internodia 3–15 cm longa, 1.3–2 cm diam.; cataphylla 7–

26 cm longa, incostata vel obtuse 1-costata, decidua; petioli subteres vel teres, moderate spongiosus, obtuse complanatus adaxialiter, 16–55 cm longus, 5–12 mm diam.; lamina ovato-triangularis, 17–36 cm longa, 11–31 cm lata, cordata vel sagittata basi; inflorescentia 1–3(4); pedunculus 3.5–29 cm longus, 8–12 mm diam.; spathe 7.5–18 cm longa; lamina spathe extus alba vel pallide viridi aut cremeo-flava, intus alba (sub anthesi) vel cremeo-flava; tubo spathe extus virenti, intus rubella vel atrimarronino; pistilla (5)6–7-loculari; loculi 1–2-ovulati; baccæ pallide vel vivide aurantiacæ.

Usually hemiepiphytic, sometimes terrestrial; stem scandent, semiglossy, sap watery, spicy-scented, sometimes milky-white; internodes usually long, glossy to semiglossy, 3–15 cm long, 1.3–2 cm diam., usually longer than broad, dark green to gray-green, sometimes drying black, epidermis often cracking or minutely fissured, brittle, sometimes silver-gray; cataphylls soft, 7–26 cm long, unribbed to bluntly or sharply 1-ribbed, sometimes sharply D-shaped and sharply 2-ribbed, whitish or green, fleshy, glossy, deciduous; **petioles** 16–55 cm long, 5–12 mm diam., subterete to terete, moderately spongy or moderately firm, bluntly D-shaped or obtusely flattened adaxially, surface weakly glossy to semiglossy, smooth; sheathing 6–8 cm long; **blades** ovate-triangular, thinly coriaceous to subcoriaceous, matte to semiglossy, weakly to moderately bicolorous, acuminate to long-acuminate at apex (the acumen more or less inrolled), cordate to sagittate at base, 17–36 cm long (averaging 26 cm), 11–31 cm wide,  $\pm$  equal in length to or longer than petiole, margins hyaline, upper surface drying brownish, dark brown, or gray-brown to green, lower surface drying light brownish green to reddish brown or yellowish brown; anterior lobe 11.5–32 cm long, 7–24 cm wide; posterior lobes somewhat spreading, rounded to obtuse, 4–14 cm long, (2.7)4.2–13.5 cm wide, directed toward base or somewhat outward; sinus hippocrepiform, parabolic to arcuate or arcuate with petiole decurrent, 1–6(8) cm deep; midrib broadly convex to shallowly sunken to flat, paler than surface above, convex, paler than surface, drying darker than surface below; basal veins 3–5 per side, with 0–1 free to base, 1–3(4) coalesced (0.7)1.3–4(7) cm; posterior rib naked for 0.5–2.6 cm, sometimes not naked, directed to the tip of the posterior lobe 1–3.5 cm distant from the lower margin of the posterior lobe; primary lateral veins (2)3–4 per side, departing midrib at a 40–60(70°) angle,  $\pm$  straight to the margins, streaked reddish below, broadly impressed to shallowly and obtusely sunken above, weakly convex and darker than surface below; minor veins moderately distinct, arising from both the midrib and primary lateral veins. **INFLORESCENCES** erect,

1–3(4) per axil; peduncle 3.5–29 cm long, 8–12 mm diam., medium green, glossy, faintly white-striate near apex; **spathe** glossy, 7.5–18 cm long, 1.4–2.2 cm diam., weakly constricted midway above the tube, 1.9–2.2 mm diam. at constriction; spathe blade creamy-white to pale green or creamy-yellow, sometimes tinged faintly reddish outside, 6–7 cm long, 2.2–3.2 cm diam. (opening 6.7–9 cm long, 3.2 cm wide), whitish (at anthesis) to creamy-yellow inside, sometimes suffused with red; resin canals orange and appearing as continuous lines at least around the throat inside; spathe tube medium to dark green and light maroon band near base, sometimes weakly to heavily tinged reddish (B & K red-purple 5/7.5), white-striate outside, 5–7.5 cm long, 2–3.6 cm diam., reddish to dark maroon (B & K red-purple 2/10) inside; **spadix** sessile;  $\pm$  tapered toward apex, 6.7–11.1 cm long, broadest at or near the base, protruding strongly or weakly out of the spathe blade at anthesis; pistillate portion weakly tapered toward both ends, light green to yellowish green, (2.5)3.4–4.3 cm long in front, 2.9–3.8 cm long in back, 7–13 mm diam. at apex, 1–14 mm diam. at middle, 1–12 mm wide at base; staminate portion 5–8.4 cm long; fertile staminate portion white or light green, 10–11 mm diam. throughout, of nearly uniform diameter throughout or broadest in upper one-third, 9–13 mm diam. at base, 10–11 mm diam. at middle, 7–9 mm diam. ca. 1 cm from apex, broadest usually at the base, narrower than the pistillate portion, narrower than the sterile portion; sterile staminate portion as broad as the pistillate portion, (pre-anthesis) white to light gray, 9–14 mm diam.; pistils 1.1–2.3(3.5) mm long, 0.7–2 mm diam.; ovary (5)6–7-locular, 0.6–1.5(2) mm long, 0.8–1.3(1.6) mm diam., with sub-basal placentation; locules 0.7–1.1(2) mm long, 0.3–0.6 mm diam.; ovule sac 0.7–0.9 mm long; ovules 1–2 per locule, 2-seriate, contained within transparent, gelatinous ovule sac, 0.3–0.5 mm long, slightly longer than funicle; funicle 0.2–0.3 mm long (can be pulled free to base), style 0.3–1.2 mm long, 0.8–1.6 mm diam., similar to style type B; style apex flat to shallowly rounded or concave; stigma brushlike to discoid or subdiscoid, purplish violet or golden honey-colored, 1–1.5 mm diam., 0.2–0.4 mm high, covering  $\pm$  entire style apex; the androecium truncate, prismatic, margins irregularly 4–6-sided, 1–1.7 mm long, 0.7–1.5(2.7) mm diam. at apex; thecae oblong, 0.4–0.5 mm wide,  $\pm$  parallel to one another; sterile staminate flowers blunt, irregular, dense, 4–6-sided, prismatic, 1.5–2(2.5–3.3) mm long, (1.2)1.5–1.6(2) mm wide. Berries pale to bright orange (white, immature); seeds yellow-orange, narrowly ellipsoid.

*Philodendron wilburii* ranges from northern Costa Rica to the Canal Area of Panama from near sea level to 2000 m elevation. This species is highly versatile ecologically, ranging from *Premontane rain forest*, *Tropical Lower Montane wet forest*, and *Tropical Lower Montane rain forest* in the highlands to *Premontane wet forest* and *Tropical wet forest* life zones in the lowlands.

*Philodendron wilburii* is a member of *P.* sect. *Calostigma* subsect. *Glossophyllum* ser. *Ovata*. This species comprises somewhat scandent plants with elongate internodes drying grayish to yellowish brown and closely fissured; mostly 1-ribbed, deciduous cataphylls; subterete petioles mostly longer than the blades; ovate to ovate-triangular blades often with the lobes directed somewhat outward; and 1-3 prominently pedunculate white to greenish inflorescences per axil.

*Philodendron wilburii* is probably most closely related to *P. cotonense* from the mountains of eastern Costa Rica. That species is distinguished by its yellowish-brown-drying, more narrowly triangular blades averaging over two times longer than wide and thicker stems (2.5-3 cm diam. vs. up to 2 cm diam. in *P. wilburii*). It also has typically longer peduncles (mostly more than 10 cm) and more ovules per locule (4-5, vs. 1-2 in *P. wilburii*). The two species do not overlap in their distribution, but do occur in the same life zones.

*Philodendron wilburii* may be confused with *P. straminicaule* from Panama, but that species differs in having thicker and shorter internodes, 1-ribbed to sharply 2-ribbed cataphylls, and larger blades (rarely less than 35 cm long, vs. rarely more than 35 cm for *P. wilburii*), with frequently more than 6 basal veins and 6 primary lateral veins (vs. no more than 5 basal veins and mostly no more than 4 pairs of primary veins for *P. wilburii*). In addition, *P. straminicaule* frequently has the peduncle shorter than the spathe and has greenish white to lavender berries (vs. having the peduncle rarely shorter than the spathe and orangish berries in *P. wilburii*). *Philodendron wilburii* is also similar to *P. smithii*, which differs in having the dried stem epidermis conspicuously yellow-brown.

This species is named in honor of Robert Wilbur (DUKE), who has made major contributions to the study of Costa Rican plants and was one of the first to collect this species.

This species is here subdivided into two varieties.

KEY TO THE VARIETIES OF *P. WILBURII*

- 1a. Peduncles typically equal to or shorter than the spathe, usually less than 12 cm long; leaf blades

with sinus hippocrepiform to parabolic, usually drying greenish to yellowish green below; Costa Rica (Alajuela, Cartago, Guanacaste, Heredia, Puntarenas, San José), 400-2000 m — var. *wilburii*  
1b. Peduncles typically much longer than the spathe, usually more than 12 cm long; leaf blades with sinus usually arcuate or arcuate with leaf tissue decurrent on petiole, rarely hippocrepiform, usually drying reddish brown below; Costa Rica (Cartago and Puntarenas) and Panama (Veraguas), 900-1400 m, 40-450 m along the Pacific slope — var. *longipedunculatum* Croat & Grayum

*Philodendron wilburii* Croat & Grayum var. *wilburii*

Usually hemiepiphytic, sometimes terrestrial; cataphylls 13-26 cm long, unribbed to bluntly 1-ribbed; petioles 16-46 cm long, obtusely flattened adaxially; blades ovate-triangular, cordate to sagittate at base, 17-36 (averaging 26) cm long, 11-27 (averaging 16) cm wide (1.3-2 times longer than wide, averaging 1.5 times), (1.3-2 times longer than petiole); sinus usually hippocrepiform, sometimes shallowly parabolic. INFLORESCENCE 1-3(4) per axil; peduncle 3.5-14 cm long; spathe 7.5-13.5 cm long; spadix 9-11 cm long; pistillate portion (2.5)3.4-4.3 cm long.

Flowering in *Philodendron wilburii* var. *wilburii* appears to have a bimodal pattern, with the main event occurring in the late dry season to the mid-rainy season (March through September but especially April through August). The second flowering event occurs in the early dry season, especially in January. A flowering collection from November is unusual. Post-anthesis collections also occur in two clusters, June through September and November through February. Mature fruits have been collected only in March.

*Philodendron wilburii* var. *wilburii* is largely known from central Costa Rica to an area of the frontier with Panama but rarely also to central Panama (Veraguas), extending from 400 to 2000 m, mostly at over 1000 m elevation. It is largely known from central Costa Rica, mostly at elevations of higher than 1000 m in premontane rain, lower montane wet, and lower montane rain forests. It has been collected mostly in the Cordillera de Tilarán, especially in the San Ramón region and in the Monteverde Reserve at 1350 to 1700 m (averaging at more than 1500 m) but to a lesser extent in areas to the east and west of the Central Valley. It has been collected only rarely in the Cordilleras de Guanacaste and Talamanca.

One collection believed to be this species, Croat 76799, is from Veraguas Province near Santa Fe in *Premontane moist forest*. Because of the isolated na-

ture of this population it may prove to represent a distinct taxon. The plants are unusual in having unribbed to bluntly 1-ribbed cataphylls; thin, ovate to ovate-triangular, cordate, mostly greenish-drying blades; and peduncles 3.5 to 14 cm long (averaging 9.3 cm long), usually equal to or shorter than the spathe.

*Philodendron wilburii* var. *wilburii* might be confused with a lowland form of *P. wilburii* var. *longipedunculatum* from southeastern Costa Rica that has leaf blades more nearly as broad as long (e.g., *Grayum* 4759). Especially confusing are *Grayum* 4759 and 9167 from Puntarenas Province in the region of the Osa Peninsula at 300 to 400 m elevation. Both look much like *P. wilburii* var. *wilburii*, and if they prove to represent that taxon, they would be the only collections from such a low elevation and the only ones from southwestern Costa Rica.

Also noteworthy is *Croat* 61199, a sterile collection that had its locale and ovary number determined from a field-collected infructescence. This collection, from Monteverde in Puntarenas Province, Costa Rica, has ovaries with only 5 locules but 10–14 ovules per locule. It otherwise matches typical *P. wilburii*. Perhaps it represents a different species.

A collection from the Azuero Peninsula in Panama is somewhat intermediate between *P. wilburii* var. *wilburii* and *P. wilburii* var. *longipedunculatum*. Though it has very long peduncles like *P. wilburii* var. *longipedunculatum*, its blades dry somewhat blackened with the posterior lobes somewhat more rounded than normal for *P. wilburii* var. *longipedunculatum*.

*Additional specimens examined for P. wilburii* var. *wilburii*. COSTA RICA. **Alajuela:** San Ramón, N of Los Angeles, ca. 1200 m, *Primack et al.* 236 (DUKE); between Los Angeles and ca. 4 km beyond Río Balsa, 1100–1200 m, *Uley* 366 (DUKE); San Ramón–Bajo Rodríguez, 940 m, *Croat* 78833 (CR, INB, MO); 1025–1100 m, 78885 (CR, INB, MO); San Ramón–Cataritas, km 11, 850 m, *Pennington* 11543 (K); Cantón Alfaro Ruiz, Guadalupe, 1450 m, *A. Smith* 2844 (F); San Luis de Zarcero, 1575 m, *A. Smith* NY1023 (F, NY); 12 km NNW of San Ramón, 1100 m, 10°10'N, 84°29'W, *Liesner & Judziewicz* 14913 (CR, MO, WIS); 15 km NNW of San Ramón by road, 2.5 km N of Balsa on road to San Lorenzo, 1050–1100 m, 10°11'N, 84°30'W, *Liesner & Judziewicz* 14753 (MO); 17–20 km NNW of San Ramón, ca. 750 m, 10°13'N, 84°32'W, 14690 (MO, WIS); 1 km S of Balsa de San Ramón, 1200 m, 10°13'N, 84°31'W, *Lent* 3514 (F, MO); 9 km SE of Ramón, 1000 m, 10°16'N, 84°05'W, *Loiselle* 177 (MO); 11.2 km N of Zarcero, 1400 m, 10°12'N, 84°23'W, *Hooser* 1351 (CR, MO); 5.7 mi. N of San Ramón, 1200 m, *Croat* 46836 (MO); ca. 1.3 mi. N of Angeles Norte, ca. 1200 m, *Croat* 46874 (MO); Finca Los Ensayos, ca. 11 mi. NW of Zarcero, *Croat* 43634 (CM, MO); San Ra-

món–Balsa, ca. 5.7 km N of bridge over Quebrada Volio, 1100–1150 m, ca. 10°08'N, 84°29'W, *Stevens* 14182 (CAS, MO); 2.3 km N, Río Balsa, ca. 1050–1150 m, 10°01'N, 84°30'W, *Stevens* 14198 (K, MO); Reserva Forestal de San Ramón, ca. 10 km W of Laguitos, Río San Lorenzo, 850–1100 m, 10°18'N, 84°34'W, *Hammel et al.* 15252 (MO); 800–1000 m, 10°12'53"N, 84°36'28"W, *Herrera & Mora* 179 (MO); Río San Pedro, Cerro Azahar, 15 km NW of San Ramón by air, 1400–1500 m, 10°09'30"N, 84°34'–35'W, *Liesner et al.* 15486 (MO), 15487 (CR, MO, NY, WIS); UCR Reserva, Fila Volcán Muerte, 1000–1300 m, 84°32'W, 10°12'N, *Barringer & Gómez-Laurito* 2559 (F), *Barringer & Pérez* 3784 (CR, F); Monteverde Reserve, *Dryer* 794 (MO), 997 (F); 1500–1600 m, 10°17'N, 84°47'W, *Huber et al.* 10069 (CR, MO, MV); 10°18'N, 84°47'W, *Grayum & Sleeper* 3851 (CR, MO). **Cartago:** Moravia–Quebrada Platanillo, ca. 1250 m, *Croat* 36658 (MO); SE of Turrialba, ca. 3 km NE of La Suiza, ca. 1200 m, *Lellinger & White* 1403 (US); Tapantí Reserve, *Nilsson* 195 (CR), 215 (CR), 220 (CR), 238 (CR), 254 (CR), 266 (CR), 367 (CR), 647 (CR); 1250–1350 m, *Ferreira* 15702 (USM); 1500–1800 m, 9°43'N, 83°47'W, *Grayum & French* 5827 (INB, MO); ca. 1350 m, 9°47'N, 83°48'W, *Grayum & Sleeper* 3694 (MO, US); *Uley & Uley* 5613 (DUKE); 1500–1800 m, 9°43'N, 83°47'W, *Croat & Grayum* 68292 (CR, MO, US); *Croat* 36122 (MO); ca. 6 km S of Cartago by air, Quebrada Cangreja, 3 km S of Pan-American Highway, 1620–1650 m, 9°46'N, 83°57'W, *Liesner & Judziewicz* 14474 (MO); Alto Pañillos, 4 km S of Tapantí, 1480 m, *Lent* 1240 (F); Río Grande de Orosi, ca. 15 km S of Tapantí, 1500 m, 9°42'N, 83°47'W, *Burger & Liesner* 6752 (F, NY); La Sierra, 6 km NE of Empalme, 2000 m, *Gómez* 19762 (MO); Santa Teresa–Río Coliblanco, 6–7 km NE of Pacayas, ca. 1700 m, *Luteyn* 3246 (DUKE, MO); Quebrada Honda–Río Sombrero, ca. 1–2 km above El Muñeco, ca. 1400 m, 3236 (DUKE, MO); 31 km from San José, SE on CR-2, 1750 m, *Harmon* 6105 (UMO); Río Grande de Orosi valley, near Río Villegas, 1700 m, *Lent* 1860 (F); Río Macho, *Nilsson* 242 (CR), 274 (CR), 312 (CR). **Guanacaste:** Tilarán, Z.P. Tenorio, 1050 m, *G. Rodríguez* 24 (CR). **Heredia:** Río Peje–Río Sardinalito, Volcán Barva, 700–950 m, 10°17'30"N, 84°04'30"W, *Grayum* 6929 (MO); Río La Paz Grande, 7.5 km N of Vara Blanca, 1270–1350 m, *Croat* 36051 (MO); La Selva Field Station, *Valerio s.n.* (USJ). **Limón:** Kámká massif, between Río Tararria and NE Kámká páramo, 1900–2300 m, 9°14'–15'N, 82°59'W, *Davidse & Herrera* 29203 (CR, MO). **Puntarenas:** Monteverde Reserve, ca. 1350 m, *Croat* 47139 (MO); 2 km SW Station, 1500–1550 m, *Ingram & Ferrell-Ingram* 1734 (MO); ca. 3.5 km NE of Monteverde Reserve, ca. 1500 m, *Wilbur et al.* 15816 (MO); *Kennedy* 545 (F); *Huber* 2369 (MO); *Pounds* 321 (MO); 449 (MO); *Gentry et al.* 48860 (CR, MO, NY); *Huber* 3880 (MO); *Huber & Bello* 2458 (MO); *Bello & Cruz* 5237 (CR, INB); *Croat* 61199 (MO); *Feinsinger et al.* 83-66-1 (FLAS); *Burger & Baker* 9649 (F, MO, SEL); Palmar Norte–Jalisco, 300–400 m, 8°58'30"N, 83°28'W, *Grayum et al.* 9167 (MO). **San José:** Bajo La Hondura, Paracito–Río Claro, 1100–1400 m, *Croat* 44494 (F, MO); above Río Hondura, at Baja La Hondura, 1150 m, *Taylor* 17933 (NY); Cerro de la Muerte, along CA-2, N of turnoff for road 222, ca. 2000 m, *Croat* 32851 (MO); ca. 15 km S of San Isidro de Cartago, 1880 m, *Uley & Uley* 2948 (F); La Palma, ca. 1600 m, *Standley* 38300 (US); La Palma area, 1400 m, 10°12'N, 84°10'W, *Burger & Liesner* 6212 (CR, F, MO); ca. 6 km NE of San Jerónimo, *Primack et al.* 322 (DUKE); Braulio Carrillo National Park,

1000–1500 m, 10°05'N, 83°57'W, *Croat 61227* (CM, MO); Río Patria–Río Zurquí, Cerro Hondura, 1500–1600 m, 10°04'N, 84°01'W, *Grayum & Sleeper 6116* (CR, MO); Puriscal, Z.P. La Cangreja, 500 m, *Morales 924* (CR, INB). PANAMA. **Veraguas:** vic. Santa Fe, along road between Santa Fe and Alto Piedra, 1.7 mi. N of Hotel Santa Fe, 470 m, 8°31'N, 81°05'W, *Croat & Zhu 76799* (B, CAS, CR, F, HMNM, K, MO, NY, PMA, SCZ, US).

***Philodendron wilburii* Croat & Grayum var. *longipedunculatum* Croat & Grayum, var. nov.**

**TYPE:** Costa Rica. Puntarenas: vic. of San Vito de Java, ca. 1 km S of San Vito on road to Villa Neily, 8°48'N, 82°57'W, 1100 m, 13 June 1987, *Croat 66169* (holotype, MO–3610645; isotypes, CR, K, NY, US). Figures 459–464.

Planta plerumque terrestris, interdum hemiepiphytica; culmis scandens; internodia 3–15 cm longa, 1.3–1.7 cm diam.; cataphylla mollia, 7–15 cm longa, acute D-formata, plerumque acute 2-costata, decidua; petiolus subteretes vel obtuse complanatus adaxialiter, 16–55 cm longa, 5–10 mm diam.; lamina 22–35 cm longa, 16–31 cm lata, triangulari-ovata, lobis posticis angustis, patentibus; inflorescentia 1–2; pedunculus 11–29.5 cm longus, 5–10 mm diam., subteretes; spathe 8.7–18 cm longa, saepe viridis omnino; lamina spatheae extus viridi vel flaviviridi aut eburnea vel citrina, intus pallide viridi, cremae, albidula aut suffusa rubra; tubo spatheae extus virenti vel atriviridi, intus atrimarronino vel atripurpureo, rubro, rubriviolaceo aut violaceo; pistilla (5)6–7(8)-locularia; loculi 1–2-ovulati; baccae aurantiacae.

Usually terrestrial, sometimes hemiepiphytic; internodes 3–15 cm long, 1.3–1.7 cm diam.; cataphylls soft, 7–15 cm long, sharply D-shaped, usually sharply 2-ribbed, sometimes unribbed or sharply 1-ribbed; **petioles** 16–55 cm long, 5–10 mm diam., obtusely flattened or bluntly D-shaped adaxially; **blades** triangular-ovate with spreading lobes, 22–35 cm long (averaging 28 cm), 16–31 cm wide (averaging 21 cm; 1.03–1.4 times longer than wide, averaging 1.3 times), (0.6–1.3 times longer than petiole); sinus usually arcuate or arcuate with decurrent petiole. **INFLORESCENCES** 1 or 2 per axil; peduncle 11–29.5 cm long, 5–10 mm diam.; **spathe** 8.7–18 cm long; **spadix** 6.7–11.2 cm long; pistillate portion 2.8–5.5 cm long; ovary (5)6–7(8)-locular; ovules 1–2 per locule. Berries bright orange (mature) or white (immature); seeds yellow-orange, narrowly ellipsoid.

The flowering phenology of *Philodendron wilburii* var. *longipedunculatum* is poorly known. It appears to be bimodal or, in any event, unlike the pattern of any other species. All flowering collections have been made between September and April, in the late rainy season and throughout the dry season. However, post-anthesis flowering collections from July make it obvious that some flowering also takes

place during June or July, perhaps both months. Other post-anthesis collections are from January through April and also in September. The only mature infructescence was collected in March.

*Philodendron wilburii* var. *longipedunculatum* is known principally from eastern Costa Rica and western Panama with an outlying population at Tapantí in Cartago Province, at 900 to 1400 m elevation, especially on the Atlantic slope but also on the Pacific slope at lower elevations (40 to 450 m). The Pacific coastal populations range from Carara to the Osa and Burica Peninsulas in Puntarenas Province, with outlying populations on the Azuero Peninsula and in the Canal Area of Panama.

*Philodendron wilburii* var. *longipedunculatum* is distinguished by the sharply two-ribbed, deciduous cataphylls; subterete, moderately firm petioles; typically triangular-ovate blades often drying reddish to yellowish brown and usually with narrow, spreading lobes; and very long-pedunculate inflorescences.

The highland populations differ from those of the lowlands in having longer posterior lobes, which are narrower and more prominently directed outward, averaging about 11 cm long (vs. about 8 cm long for the lowland populations). The lowland populations have proportionately narrower blades (owing to their lack of long, outward-directed lobes), averaging 1.6 times longer than wide (vs. 1.3 times for highland populations).

Lowland populations in the vicinity of the Burica Peninsula have somewhat thicker blades drying with a minutely pustular or granular lower surface with the secretory ducts mostly not visible. In contrast, blades of highland populations dry darker brown to almost blackened on the upper surface and are smooth on the lower surface with the secretory ducts clearly visible and alternating with the minor veins.

The Azuero population, represented by *Croat 34476*, has leaf blades that dry somewhat blackened and have posterior lobes somewhat less narrowed and less directed outward than those of the highland Chiriquí populations. The Canal Area populations, represented by *Croat 12351* and *Croat & Zhu 77083*, also differ in having proportionately longer petioles than material elsewhere. The latter collection is also exceptional in having four inflorescences per axil.

*Grayum & Sleeper 3435* is a noteworthy collection from Tapantí, Cartago Province, Costa Rica. It is unusual both in being from well out of the normal range of the variety, but also in having narrower

leaf blades and shorter peduncles. Perhaps it represents another taxon.

*Additional specimens examined.* COSTA RICA. **Cartago:** Tapantí Region, N of Quebrada Casa Blanca, 1300 m, 9°47'N, 83°48'W, *Grayum & Sleeper 3435* (AAU, CR, MO). **Puntarenas:** San Vito-Ciudad Neily, Fila de Cal, ca. 500–620 m, 8°41'N, 82°56'30"W, *Grayum et al. 6048* (CR, INB, K, MO, PMA); Osa Peninsula, vic. Rincón de Osa, 250–540 m, 8°42'N, 83°31'W, *Croat & Grayum 59879* (CR, MO); Quebrada Banegas-Río Riyito, ca. 7 km W of Rincón de Osa, 100–300 m, 8°41'N, 83°33'W, *Grayum 4089* (CR, MO); Rincón-Aguabuena, road to Rancho Quemado, 100 m, 8°40'N, 83°31'W, *G. Herrera 4619* (CR, INB, MO); Finca Loma Linda, 1 mi. SW of Cañas Gordas, 1140 m, *Croat 22310* (F, MO); Río Piedras Blancas, Fila Costeña, Fila Cruces, Cerro Anguciana, 950–1150 m, 8°49'18"N, 83°11'15"W, *Grayum 10650* (CR, INB, MO); Quebrada Bonita, to ca. 1 km E of Costanera highway, Caraza Reserve, 30–40 m, 9°47'N, 84°37'W, *Grayum 4759* (CR, L, MEXU, MO). **San José:** San Isidro de El General-Dominical, 4.8 mi. S of Río Pacuare, 1000 m, *Croat 35249* (CR, K, MO, PMA, US); ca. 0.5 mi. above turnout to Canadán at Rivas, 900 m, *Croat 43429* (CR, MO); ca. 8 km SW of San Isidro de El General, 1000 m, *Williams et al. 28387* (F). PANAMA. **Bocas del Toro:** Fortuna-Chiriquí Grande, Continental Divide, 1200 m, 8°44'N, 82°17'W, *Croat & Grayum 60372* (COL, CR, DAV, IBE, K, MO, NY, PMA, S, U); 1.2 mi. N of Continental Divide, 910 m, 8°44'N, 82°17'W, *Croat 60454* (MO, PMA); 4.5–5 km N of dam over Fortuna Lake, 1100–1135 m, 8°43'N, 82°17'W, *Croat & Grayum 60026* (B, CAS, CR, F, L, KYO, MEXU, MO, NY, PMA, R, S); Cerro Colorado, ca. 7.5 mi. from Chame Camp, 1200–1250 m, ca. 8°35'N, 81°45'W, *McPherson 8868* (MO). **Canal Area:** Pipeline Road, near R. Agua Salud, *Croat 12351* (DUKE, F, MEXU, MO, SCZ); 6 mi. N of Gamboa, Río Mendoza, 9°11'N, 79°46'W, *Croat & Zhu 77083* (CM, CR, MO, NY). **Chiriquí:** Burica Peninsula, San Bartolo Límite, 18 km W of Puerto Armuéllens, 450 m, *Bussey 597* (CM, CR, MO); Rabo de Puerco, 8 km W of Puerto Armuéllens, 50–150 m, *Croat 21958* (F, MO); 200 m, *22482* (F, MO); Cerro Colorado, 34–35.6 km above Río San Félix, 1390–1410 m, *Croat 37250* (F, MO, NY, PMA); 28 mi. above San Félix, 1200–1500 m, *33202* (MO); Gualaca-Fortuna, 10 mi. NW of Los Planes de Hornito, 1260 m, 82°17'W, 8°45'N, *Croat 50093* (L, MO, NY); 5.9 mi. NW of Los Planes de Hornito, 1370 m, 8°43'N, 82°15'W, *Croat 49852* (MO); 7.2 mi. beyond Los Planes de Hornito, 1165–1200 m, 8°44'N, 82°14'W, *67836* (CM, F, MO, PMA); 11.8 mi. N of Los Planes de Hornito, 1400 m, *48696* (MO); near Lago Fortuna, 1200 m, 8°46'N, 82°17'W, *Croat 74907* (MO, PMA); 1100–1200 m, 8°45'N, 82°18'W, *66565* (CM, MO); 4.8 mi. beyond IRHE Facilities at Dam, 8°46'N, 82°16'W, *Croat 69032* (MO, NY, PMA, US); trail to Río Hornito, 1100–1300 m, 8°45'N, 82°15'W, *Thompson 5003* (CM, MO); 8°45'N, 82°18'W, *Croat & Zhu 76416* (AAU, CAS, GB, MEXU, MO, SCZ, US); *76378* (MO, SCZ); N edge of lake, ca. 1100 m, ca. 8°45'N, 82°15'W, *McPherson 9081* (MO, US); 3.4 km N of Quebrada Chorro, 1.6 mi. N of center of bridge over lake, 1205 m, 8°43'N, 82°14'W, *Croat 74955* (MO); Finca Loma Linda, 1 mi. E of Cañas Gordas near Costa Rican border on road to Volcán, *Croat 23310* (F, MO). **Darién:** Parque Nacional Darién, W side of Cerro Pirre, 800–1,050 m, 7°56'N, 77°45'W, *Croat 69001* (MO). **Los Santos:** Azuero Peninsula, Jobero-Río Pedregal

headwaters, 300–700 m, *Croat 34476* (MO, PMA). **Panamá:** Río Majé, ca. 4–5 mi. above waterfalls, near new Bayano Lake, <100 m, *Croat 34726* (MO).

**Philodendron zhanum** Croat, sp. nov. TYPE: Panama. Coclé: Alto Calvario, vic. El Copé, ca. 6 km N of El Copé, 770 m, 8°38'N, 80°35'W, 12 July 1994, *Croat & Zhu 76755* (holotype, MO–4619357–9; isotypes, AAU, B, CAS, COL, CR, F, GH, K, MEXU, NY, PMA, QCNE, SCZ, US, VEN). Figures 465–468.

Planta hemiepiphytica; internodia 2–4 cm longa, 3–5.5 cm diam.; cataphylla 37–50 cm longa, acute 2-costata, decidua; petiolus 40–63 cm longus, obtuse D-formatus, in sicco manifeste complanatus usque 3–6 cm latus; lamina anguste ovato-sagittata, 38–62 cm longa, 27–32 cm lata; inflorescentia 3; pedunculus 9–14 cm longus, 1–1.5 cm diam.; spatula 13–18.5 cm longa, lamina spathae alba, intus suffusa magenta in triente inferiore; tubo spathae extus viridi, intus atriviolaceo-purpureo; pistilla 8–9-locularia; loculi 1–3-ovulati.

Hemiepiphytic, loosely or appressed-climbing; leaf scars conspicuous, 3 cm long, 2 cm wide; internodes semiglossy, 2–4 cm long, 3–5.5 cm diam., dark green, drying dark brown, longitudinally wrinkled; roots drying light brown and weakly glossy, finely striate, epidermis peeling; cataphylls 37–50 cm long, sharply 2-ribbed, medium green, semiglossy, drying dark brown, deciduous; petioles 40–63 cm long (0.8–1.5 times longer than blades), bluntly D-shaped, faintly raised medially, moderately firm, medium green, surface dark green short-lineate, drying dark yellowish brown and glossy, prominently flattening to 3–6 cm wide and only a few mm thick, smooth to finely and closely striate; sheath 3–6 cm long, moderately closed; blades narrowly ovate-sagittate, moderately coriaceous, acuminate at apex, conspicuously sagittate at base, 38–62 cm long, 27–32 cm wide (1.8–2 times longer than wide), margins broadly undulate, much paler than surface, upper surface dark green, glossy, drying dark yellowish brown, lower surface weakly glossy, paler, drying reddish brown to yellowish green with conspicuous, dark intermittent secretory canals visible; anterior lobe 34–35 cm long, margins broadly convex, paler; posterior lobes 8.5–18.5 cm long, 7–13 cm wide, directed downward to slightly inward, broadly rounded; sinus narrow, spatulate to narrowly obovate, (7.5)15–19 cm deep, 1–3 cm wide; major veins drying with sharp ridges, usually darker than surface; midrib flat, paler than surface above, narrowly rounded, concolorous below; basal veins (3)5–6 per side, with the first free to base, numbers 3–5 coalesced (2)5–7 cm; posterior rib not at all naked or naked to 2.5 cm; primary lateral veins 6–7 per side, arising at an acute angle



then spreading to a 40–45° angle, weakly arcuate until near the margin, then turned up abruptly, weakly sunken above, convex, paler than surface below; minor veins clearly visible, arising from both the midrib and primary lateral veins, drying prominent above, faint and weakly wrinkled below. INFLORESCENCES 3 per axil; peduncle 9–14 cm long, 1–1.5 cm diam., somewhat flattened, medium green, faintly short-lineate; spathe 13–18.5 cm long, margins paler; spathe blade white, except yellow-green medially on back side outside, tinged magenta on lower one-third inside; spathe tube medium green outside, dark violet-purple inside, spadix 11.5–18 cm long; pistillate portion creamy green, 3.5–8 cm long, 3–6 cm long in back, 1.5 mm diam. at apex, 1.9 mm diam. at middle, 1.8 mm wide at base; staminate portion 8–10 cm long; fertile staminate portion slightly narrower than the pistillate portion; sterile staminate portion barely distinguishable when fresh, 1.2–1.3 cm diam. at base, tapered only slightly upward when fresh (drying to 10 mm diam. at apex); pistils 1.9–2.1 mm long, 1.2–1.6 mm diam.; ovary 8–9-locular, 0.8 mm long, with basal placentation; locules 0.8–1.2 mm long; ovules 1–3 per locule, translucent, contained within gelatinous matrix (no true envelope), 0.3–0.4 mm long, equal in length to funicle, style similar to style type B; stilar canals emerging directly onto surface; style apex truncate; stigma short, covering entire style apex; the androecium truncate, margins irregularly 4–6 sided, 0.8–1.2 mm diam. at apex; sterile staminate flowers irregularly rounded, 0.8–1 mm wide.

Flowering in *Philodendron zhuanium* is documented by a single collection from July, at the beginning of the rainy season. A cultivated plant of the same collection flowered in November at the Missouri Botanical Garden.

*Philodendron zhuanium* is endemic to Panama in Coclé Province in an area of Premontane rain forest at 700 to 900 m elevation.

*Philodendron zhuanium* is a member of *P.* sect. *Calostigma* subsect. *Macrobelyum* ser. *Macrobelyum*. This species is distinguished by its short, thick internodes; deciduous, sharply two-ribbed cataphylls; bluntly D-shaped petioles about as long as the blades or slightly shorter; narrowly ovate blades with prominent posterior lobes, a narrow sinus and prominently visible secretory canals; up to three inflorescences per axil; and spathes green outside and whitish inside on the blade, and dark purple-violet inside on the tube.

*Philodendron zhuanium* is apparently related to *P. sagittifolium* and *P. aromaticum*, differing from

both in having petioles which on drying become moderately spongy, markedly flattened (especially in the middle portions), dark yellow-brown to yellowish, and often glossy. It also differs in having conspicuous secretory canals visible in both living and dried conditions. While secretory canals are present in both *P. sagittifolium* and *P. aromaticum*, they are quite inconspicuous.

This species is perhaps most easily confused with *P. aromaticum*, which has similar leaf blades with a naked posterior rib. The latter species occurs at lower elevations and nearer the coast. In contrast, *P. zhuanium* rarely has the posterior rib naked. While *P. zhuanium* has the spathe tube dark purple-violet inside with the same color bleeding onto the lower one-third of the blade, the spathe tube of *P. aromaticum* is white to weakly maroon within and the color does not merge onto the lower part of the blade.

*Philodendron zhuanium* is named in honor of Guanghua Zhu, my student, field companion, and a monographer of the genus *Dracontium*, who helped to collect the type and many other aroids during our fieldwork in Panama.

*Additional specimen examined.* PANAMA. Coclé: Alto Calvario, vic. El Copé, ca. 5 mi. N of El Copé, 900–1000 m, Croat 75054 (CM, MO).

#### NOMEN INCERTAE SEDIS

*Philodendron auritum* Lindl., J. Hort. Soc. London 8: 60. 1853.

This is perhaps an older name for *Philodendron anisotomum* but was based on a cultivated plant collected by Skinner, perhaps in Guatemala, and no specimen or illustration is known to exist to confirm that it is even a *Philodendron*. Lindley stated that it had foliage very similar to a *Syngonium* illustrated by Vellozo (1825) in *Flora Fluminensis* (vol. 10, t. 113).

#### EXCLUDED NAMES

*Philodendron armigerum* Standl. & L. O. Williams, Ceiba 3: 107. 1952. = *Syngonium armigerum* (Standl. & L. O. Williams) Croat, Ann. Missouri Bot. Gard. 68: 585. 1981.

*Philodendron bresinodum* Standl. & L. O. Williams, Ceiba 1: 231. 1951. = *Monstera tuberculata* Lundell var. *bresinoda* (Standl. & L. O. Williams) Madison, Contr. Gray Herb. 207: 92. 1977.

*Philodendron hastiferum* Standl. & L. O. Williams, Ceiba 1: 232. 1951. = *Syngonium hastiferum*

(Standl. & L. O. Williams) Croat, *Ann. Missouri Bot. Gard.* 68: 595. 1981.

## Literature Cited

- Armbruster, W. S. 1984. The role of resin in angiosperm pollination: Ecological and chemical considerations. *Amer. J. Bot.* 71: 1149-1160.
- Bartlett, H. H. 1937. The vegetation of Petén. *Publ. Carnegie Inst. Wash.* 47b: 162.
- Bay, D. C. 1995[1996]. Thermogenesis in the aroids. *Aroidiana* 18: 32-39.
- Berlin, B. & P. Kay. 1969. *Basic Color Terms, Their Universality and Evolution*. Univ. California Press, Berkeley.
- Birdsey, M. 1951. *The Cultivated Aroids*. Gillick Press, Berkeley, California.
- Blanc, P. 1977a. Contribution à l'étude Aracées. I. Remarques sur la croissance monopodiale. *Rev. Gén. Bot.* 84: 115-126.
- . 1977b. Contribution à l'étude Aracées. II. Remarques sur la croissance sympodiale chez l'*Anthurium scandens* Engl., le *Philodendron fenzlii* Engl., et le *Philodendron speciosum* Schott. *Rev. Gén. Bot.* 84: 319-331.
- . 1977c. Contribution à l'étude Aracées. II. Remarques sur la croissance monopodiale. *Rev. Gén. Bot.* 84: 115-126.
- . 1978. Aspects de la ramification chez des Aracées tropicales. Thèse du Diplôme de Docteur 3<sup>ème</sup> Cycle, Université P. & M. Curie, Paris.
- . 1980. Observations sur les flagelles des Aracées. *Adansonia Ser.* 2, 20: 325-388.
- Bogner, H. & D. H. Nicolson. 1991. A revised classification of Araceae with dichotomous keys. *Willdenowia* 21: 35-50.
- Buggeln, R. G., B. J. D. Meeuse & J. R. Klima. 1971. The control of blooming in *Sauromatum gottatum* Schott by darkness. *Canad. J. Bot.* 49: 1025-1031.
- Bunting, G. S. 1963a. Studies in Araceae. *Ann. Missouri Bot. Gard.* 50: 23-28.
- . 1963b. A reconsideration of *Philodendron heteraceum*. *Baileya* 2: 62-67.
- . 1965. Commentary on Mexican Araceae. *Gentes Herb.* 9: 289-382.
- . 1968. Vegetative anatomy of the *Philodendron scandens* complex. *Gentes Herb.* 10: 136-168.
- . 1979. Sinopsis de las Araceae de Venezuela. *Rev. Fac. Agron. (Maracay)* 10: 139-290.
- . 1986. New taxa of Venezuelan Araceae. *Phytologia* 60: 293-344.
- . 1988. New taxa of Venezuelan Araceae—II. *Phytologia* 64: 459-486.
- . 1995. Araceae. Pp. 600-679. In: J. Steyermark, P. Berry & B. Holst (editors), *Flora of the Venezuelan Guayana*. Vol. 2: Acanthaceae—Araceae. Timber Press, Portland & Missouri Botanical Garden, St. Louis.
- Carvell, W. N. 1989. *Floral Anatomy of the Pothoideae and Monsteroideae (Araceae)*. Ph.D. Dissertation, Miami University, Oxford, Ohio.
- Croat, T. B. 1975. Phenological behavior of habit and habitat classes on Barro Colorado Island (Panama Canal Zone). *Biotropica* 7: 270-277.
- . 1978. *Flora of Barro Colorado Island (Panama Canal Zone)*. Stanford Univ. Press, Stanford.
- . 1979. The distribution of Araceae. Pp. 291-308 in K. Larsen & L. B. Holm-Nielsen (editors), *Tropical Botany*. Academic Press, London.
- . 1980. Flowering behavior of the neotropical genus *Anthurium* (Araceae). *Amer. J. Bot.* 67: 888-904.
- . 1981[1982]. A revision of *Syngonium* (Araceae). *Ann. Missouri Bot. Gard.* 68: 565-651.
- . 1983a. A revision of *Anthurium* (Araceae) of Mexico and Central America. Part I. Mexico and Middle America. *Ann. Missouri Bot. Gard.* 70: 211-420.
- . 1983b. *Dieffenbachia*. Pp. 234-236 in D. N. Janzen (editor), *Costa Rican Natural History*. Univ. Chicago Press, Chicago.
- . 1985a. Collecting and preparing specimens of Araceae. *Ann. Missouri Bot. Gard.* 72: 252-258.
- . 1986a. A revision of *Anthurium* (Araceae) for Mexico and Central America. Part II. Panama. *Monogr. Syst. Bot. Missouri Bot. Gard.* 14.
- . 1986b. The distribution of *Anthurium* (Araceae) in Mexico, Middle America and Panama. *Selbyana* 9: 94-99.
- . 1988[1990]. Ecology and life forms of Araceae. *Aroidiana* 11: 4-55.
- . 1990[1992]. A comparison of aroid classification systems. *Aroidiana* 13: 44-63.
- . 1991. A revision of *Anthurium* sect. *Pachynarium* (Araceae). *Ann. Missouri Bot. Gard.* 78: 539-855.
- . 1992. Species diversity of Araceae in Colombia: A preliminary survey. *Ann. Missouri Bot. Gard.* 79: 17-28.
- & G. S. Bunting. 1979. Standardization of *Anthurium* descriptions. *Aroidiana* 2: 15-25.
- & N. Lambert. 1986. The Araceae of Venezuela. *Aroidiana* 9(1-4): 3-213.
- Crosby, M. R. 1986. Index Muscorum—A computerized muscological database. *Bull. Brit. Bryol. Soc.* 48: 25-26.
- & R. Magill. 1986. TROPICOS: The botanical database at the Missouri Botanical Garden. *Missouri Botanical Garden*, St. Louis.
- Cullen, J. 1978. A preliminary survey of ptyxis (vernation) in the angiosperms. *Notes Roy. Bot. Gard. Edinburgh* 37: 161-214.
- Daghlian, C. P. 1981. A review of the fossil record of monocotyledons. *Bot. Rev. (Lancaster)* 47: 517-555.
- Dahlgren, R. M. T. & H. T. Clifford. 1982. *The Monocotyledons: A Comparative Study*. Academic Press, London.
- Davidse, G., M. Sousa S. & S. Knapp. 1995. In: G. Davidse, M. Sousa S. & A. O. Chater (editors), *Flora Mesoamericana*, Vol. 1, *Psilotaceae—Salvinaceae*. Universidad Nacional Autónoma de México, México D.F.; Missouri Botanical Garden, St. Louis; The Natural History Museum, London.
- Dilcher, D. L. & C. P. Daghljan. 1977. Investigations of angiosperms from the Eocene of southeastern North America: *Philodendron* leaf remains. *Amer. J. Bot.* 64: 526-534.
- Dodson, C. & A. Gentry. 1978. *Flora of the Río Palenque Science Center: Los Ríos Province, Ecuador*. *Selbyana* 41: 1-628.
- & F. Valverde. 1985. *Flora de Juaneche*. Ediciones del banco del Ecuador, Quito.
- Dugand, A. 1945. Revaluación de *Philodendron heteraceum* Schott como transferencia de *Arum heteraceum*. *Caldasia* 3: 445-452.
- Endlicher, S. 1837. *Genera Plantarum* 1(3). Vienna.
- Endress, P. K. 1982. Syncarpy and alternative nodes of

- escaping disadvantages of apocarpy in primitive angiosperms. *Taxon* 31: 48-52.
- Engler, A. 1876. Zur Morphologie der Araceae. *Bot. Zeitung* (Berlin) 34: 81-90, 95-105.
- . 1877. Vergleichende Untersuchungen über die morphologischen Verhältnisse der Araceae. II. Ueber Blattstellung und Sprossverhältnisse der Araceae. *Nova Acta Acad. Caes. Leop.-Carol. German. Nat. Cur.* 39: 157-233.
- . 1878. Araceae. Pp. 25-244 in: C. F. P. von Martius (editor), *Flora brasiliensis* 3(2A). F. Fleischer, Leipzig.
- . 1879. Araceae. In: A. & C. De Candolle (editors), *Monographie Phanerogamarum* 2: 1-681. Paris.
- . 1899. Beiträge zur Kenntnis der Araceae. IX. 16. Revision der Gattung *Philodendron* Schott. *Bot. Jahrb. Syst.* 26: 509-564.
- . 1905a. Beiträge zur Kenntnis der Araceae. X. 18. Araceae novae. *Bot. Jahrb. Syst.* 37: 110-143.
- . 1905b. Araceae-Pothoideae. In: A. Engler (editor), *Das Pflanzenreich IV.23B* (Heft 21): 1-330. W. Engelmann, Leipzig and Berlin.
- . 1908. Additamentum ad Araceae-Pothoideas. Pp. 1-3 in: A. Engler (editor), *Das Pflanzenreich IV.23B* (Heft 37). W. Engelmann, Leipzig and Berlin.
- . 1911. Araceae-Lasiodeae. In: A. Engler (editor), *Das Pflanzenreich IV.23B* (Heft 21): 1-310.
- . 1912. Araceae-Philodendroideae-Philodendreae-Homalomeninae und Schismatoglossitidinae. In: A. Engler (editor), *Das Pflanzenreich IV.23* (Heft 55): 1-134. W. Engelmann, Leipzig and Berlin.
- . 1915. Araceae-Philodendroideae-Amubiadae-Aglanometaceae-Dieffenbachieae-Zantedeschieae-Typhonodorea-Peltandreae. In: A. Engler (editor), *Das Pflanzenreich IV.23Dc* (Heft 64): 1-78.
- . 1920a. Araceae, pars generalis et Index familiae generalis. In: A. Engler (editor), *Das Pflanzenreich IV.23A* (Heft 74): 1-71.
- . 1920b. Araceae-Aroideae und Pistioidae. In: A. Engler (editor), *Das Pflanzenreich IV.23F* (Heft 73): 1-274.
- & K. Krause. 1908. Araceae-Monsteroideae. Pp. 4-138. In: A. Engler (editor), *Das Pflanzenreich IV.23* (Heft 37).
- & ———. 1920. Araceae-Colocasioidae. In: A. Engler (editor), *Das Pflanzenreich IV*. (Heft 71): 1-139.
- Eyde, R. H., D. H. Nicolson & P. Sherwin. 1967. A survey of floral anatomy in Araceae. *Amer. J. Bot.* 54: 478-497.
- Faegri, K. & L. van der Pijl. 1979. *The Principles of Pollination Ecology*. Pergamon Press, London.
- Flores, G., L. Jiménez, X. Madrigal, R. Moncayo & F. Taksaki. 1971. Mapa de Tipos de Vegetación de la República México. 1: 200,000. Secretaría de Recursos Hidráulicos, México.
- Foster, M. B. 1949. My flower has a temperature! *Natl. Hort. Mag.* 10-19 (Jan.).
- French, J. C. 1985. Patterns of endothelial wall thickenings in Araceae: Subfamilies Calloideae, Lasiodeae, and Philodendroideae. *Bot. Gaz.* 146: 521-533.
- . 1986a. Patterns of stamen vasculature in the Araceae. *Amer. J. Bot.* 73: 434-449.
- . 1986b. Ovarial vasculature of Araceae. *Bot. Gaz.* 147: 478-495.
- . 1987a. Systematic occurrence of sclerotic hypodermis in roots of Araceae. *Amer. J. Bot.* 74: 891-903.
- . 1987b. The structure of ovular and placental trichomes of Araceae. *Bot. Gaz.* 148: 198-208.
- . 1987c. Systematic survey of resin canals in roots of Araceae. *Bot. Gaz.* 148: 360-371.
- . 1988. Systematic occurrence of anastomosing laticifers in Araceae. *Bot. Gaz.* 149: 71-81.
- & P. B. Tomlinson. 1980. Preliminary observations on the vascular system in stems of certain Araceae. Pp. 105-116, pl. 1-9. In: C. D. Brickell, D. F. Cutler & M. Gregory (editors), *Petaloid Monocotyledons*. Academic Press, London.
- & ———. 1981. Vascular patterns in stems of Araceae: subfamily *Philodendroideae*. *Bot. Gaz.* 142: 550-563.
- & ———. 1984. Patterns of stem vasculature in *Philodendron*. *Amer. J. Bot.* 71: 1432-1443.
- , M. Chung & Y. Hsu. 1995. Chloroplast DNA phylogeny of Arifloae. Pp. 255-275 in P. J. Rudall, P. J. Cribb, D. F. Cutler & C. J. Humphries (editors), *Monocotyledons: Systematics & Evolution*. Royal Botanic Gardens, Kew.
- Gentry, A. H. 1982. Evidence for phylogeographic patterns as evidence for a Chocó refuge. Pp. 112-136 in G. T. Prance (editor), *Biological Diversification in the Tropics*. Columbia Univ. Press, New York.
- Goebel, K. & W. Sandt. 1930. Untersuchungen an Luftwurzeln. *Bot. Abb.* 17: 1-124.
- Gottsbarger, G. 1964. Pollination strategies in Brazilian *Philodendron* species. *Ber. Deutsch. Bot. Ges.* 97: 391-410.
- . 1986. Wärmeentwicklung von *Philodendron*-Blüten. *Naturwiss. Rundschau* (Stuttgart) 39: 350-351.
- . 1990. Flowers and beetles in South American tropics. *Ber. Deutsch. Bot. Ges.* 103: 360-365.
- & A. Amaral, Jr. 1984. Pollination strategies in Brazilian *Philodendron* species. *Ber. Deutsch. Bot. Ges.* 97: 391-410.
- & I. Silberbauer-Gottsbarger. 1991. Olfactory and visual attraction of *Eriocelis emarginata* (Cyclocephalini, Dynastinae) to the inflorescences of *Philodendron sellowii* (Araceae). *Biotropica* 23: 23-28.
- Grayum, M. H. 1984. *Palyonology and Phenology of the Araceae*. Ph.D. Dissertation, University of Massachusetts, Amherst.
- . 1985. Evolutionary and ecological significance of starch storage in pollen of the Araceae. *Amer. J. Bot.* 72: 1565-1577.
- . 1986. Correlations between pollination biology and pollen morphology in the Araceae, with some implications for angiosperm evolution. Pp. 313-327 in S. Blackmore & I. Ferguson, *Pollen and Spores—Form and Function*. Academic Press, London.
- . 1990. Evolution and phylogeny of Araceae. *Ann. Missouri Bot. Gard.* 77: 628-697.
- . 1991. Systematic embryology of the Araceae. *Bot. Rev.* 57: 167-203.
- . 1992a. Comparative external pollen ultrastructure of the Araceae and putatively related taxa. *Monogr. Syst. Bot. Missouri Bot. Gard.* 43: 1-167.
- . 1992b. New species of *Philodendron* subgenus *Pteromischum* (Araceae) from Mesoamerica and Pacific South America. *Phytologia* 73: 30-39.
- . 1996. Revision of *Philodendron* subgenus *Pteromischum* (Araceae) for Pacific and Caribbean Tropical America. *Syst. Bot. Monogr.* 47: 1-233.

- Gregor, H.-J. & J. Bogner. 1984. Fossile Araceen Mitteleuropas und ihre rezenten Vergleichsformen. *Documenta Naturae* 19: 1-12.
- Greuter, W., F. R. Barrie, H. M. Burdet, W. G. Chaloner, V. Demoulin, D. L. Hawksworth, P. M. Jørgensen, D. H. Nicolson, P. C. Silva, P. Treharne & J. McNeill. 1994. International Code of Botanical Nomenclature (Tokyo Code). Regnum Veg. 131.
- Hay, A. & D. J. Mabberley. 1991. Transference of function and the origin of aroids: Their significance in early angiosperm evolution. *Bot. Jahrb. Syst.* 113: 339-428.
- Henry, M. F. & E. J. Nyns. 1975. Cyanide insensitive respiration. An alternative mitochondrial pathway. *Subcell. Biochem.* 4: 1-65.
- Herk, A. W. H. van. 1937a. Die chemischen Vorgänge im Sauromatum-Kolben. I. *Rec. Trav. Bot. Néerl.* 34: 69-156.
- . 1937b. Die chemischen Vorgänge im Sauromatum-Kolben. II. *Proc. Kon. Ned. Akad. Wetensch.* 40: 607-617.
- . 1937c. Die chemischen Vorgänge im Sauromatum-Kolben. III. *Proc. Kon. Ned. Akad. Wetensch.* 40: 709-719.
- Hermann, P. 1698. Pp. 69-95 in *Paradisus batavus*. Leiden.
- Holmes, J. W. 1969. On the absolute fall of sea-level during the Quaternary. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 6: 237-239.
- Holdridge, L. R. 1967. Life zone ecology. Tropical Science Center, San José, Costa Rica.
- , W. C. Grenke, W. H. Hatheway, T. Liang & J. A. Tosi, Jr. 1971. *Forest Environments in Tropical Zones*. Pergamon Press, Oxford.
- Hotta, M. 1970. A system of the family Araceae in Japan and adjacent areas. *Mem. Fac. Sci. Kyoto Univ., Ser. Biol.* 4: 72-96.
- Jackson, B. D. 1971. *A Glossary of Botanic Terms*. Fourth Ed. Duckworth, London.
- Jacquin, N. J. 1760. P. 31. *Enumeratio Systematica Plantarum*. Leiden.
- . 1763. P. 240, Pl. 152. *Selectarum Stirpium Americanarum Historia. Ex officina Krausiana*. Vienna.
- . 1797. *Plantarum Rariorum Horti Caesarei Schoenbrunnensis*. Vienna.
- James, W. & H. Beever. 1950. The respiration of *Arum spadix*. *New Phytol.* 49: 353-374.
- Johnston, I. M. 1949. The botany of San José Island (Gulf of Panama). *Sargentia* 8: 1-306.
- Keating, R. C. In press. Vol. 10. Araceae. In: D. F. Cutler & M. Gregory (editors), *Anatomy of the Monocotyledons*. Clarendon Press, Oxford.
- Kelly, N. 1985. Epiphytes and climbers of a Jamaican rain forest: Vertical distribution, life forms and life histories. *J. Biogeogr.* 12: 223-241.
- Knutsen, R. M. 1974. Heat production and temperature regulations in eastern skunk cabbage. *Science* 186: 746-747.
- Koch, K. 1853. Araceae. P. 14 in *Index Seminum in horto botanico berlinensi* Colantur, Appendix. Berlin.
- Kraus, G. 1884. Über die Blütenwärme bei *Arum italicum*. *Abh. Naturf. Ges. Halle* 16: 746-747.
- . 1896. Physiologisches aus den Tropen. *Ann. Jard. Bot. Buitenzorg* 8: 217-275.
- Krause, K. 1913. Araceae-Philodendroidae-Philodendreae-Philodendrinae. In: A. Engler & K. Krause (editors), *Das Pflanzenreich IV.23Db (Heft 60)*: 1-143. W. Engelmann, Leipzig.
- Kunth, C. S. 1841. Araceae. Pp. 1-87 in *Enumeratio Plantarum*, Vol. III. J. G. Cotta, Stuttgart & Tubingen.
- Lelek, E. 1910. Untersuchungen über die Blütenwärme der Araceen. Verlag von Bruncken, Greifswald.
- . 1916. Die Erwärmungstypen der Araceen und ihre Blütenbiologische Deutung. *Ber. Deutsch. Bot. Ges.* 33: 518-537.
- Lierau, M. 1888. Über die Wurzeln der Araceen. *Bot. Jahrb. Syst.* 9: 1-38.
- Linnaeus, C. 1754. *Genera plantarum*, ed. 5. Stockholm.
- Madison, M. 1977. A revision of *Monstera* (Araceae). *Contr. Gray Herb.* 207: 3-100.
- . 1978. *Genera of Araceae in the northern Andes*. *Aroideana* 1: 31-53.
- Matuda, E. 1954. Las Araceae Mexicanas. *Anal. Inst. Biol. Univ. Nac. Auton. Mexico, Bot.* 25: 176.
- . 1962. Nuevas plantas de Mexico. *Anal. Inst. Biol. Univ. Nac. Auton. Mexico, Bot.* 32: 143-155.
- Mayo, S. J. 1986. *Systematics of Philodendron Schott (Araceae) with Special Reference to Inflorescence Characters*. Ph.D. Thesis, University of Reading, U.K.
- . 1988. Aspectos da evolução e da geografia do gênero *Philodendron* Schott (Araceae). *Acta Bot. Brasil.* 1(2) Supl.: 27-40.
- . 1989. Observations of gynoeceal structure in *Philodendron* (Araceae). *J. Linn. Soc. Bot.* 100: 139-172.
- . 1990. History and infrageneric nomenclature of *Philodendron* (Araceae). *Kew Bull.* 45: 37-71.
- . 1991. A revision of *Philodendron* subgenus *Mecanostigma* (Araceae). *Kew Bull.* 46: 601-681.
- , J. Bogner & P. C. Boyce. 1995. *The Arales*. Pp. 277-286 in P. J. Rudall, P. J. Cribb, D. F. Cutler & C. J. Humphries (editors), *Monocotyledons: Systematics & Evolution*. Royal Botanic Gardens, Kew.
- , & ———. 1997. *The Genera of Araceae*. Royal Botanic Gardens, Kew. In press.
- Meeuse, B. J. D. 1966. The voodoo lily. *Sci. Amer.* 215: 80-88.
- . 1975. Thermogenic respiration in aroids. *Annual Rev. Pl. Sci.* 26: 117-126.
- . 1978. The physiology of some sapromyophilous flowers. In: A. Richards (editor), *The Pollination of Flowers by Insects*. Academic Press, London.
- & R. G. Baggeln. 1969. Time, space, light and darkness in the metabolic flare-up of the spadix of *Sauromatum* appendix. *Acta Bot. Néerl.* 18: 159-171.
- Moodie, G. E. E. 1976. Heat production and pollination in Araceae. *Canad. J. Bot.* 54: 545-546.
- Nagy, K. A., D. K. Odell & R. S. Seymour. 1972. Temperature relation by the inflorescence of *Philodendron*. *Science* 178: 1195-1197.
- Nicolson, D. H. 1960. A brief review of classifications in the Araceae. *Baileya* 8: 62-67.
- . 1994. Report of the General Committee: 6. *Taxon* 43: 279-281.
- Plumier, C. 1756. *Plantarum americanarum fascicularis primus*. Fascicle 2. t. 51.
- Pohl, F. 1932. Anatomische und ökologische Untersuchungen am Blütenstande von *Philodendron sellowii* Schott, mit besonderer Berücksichtigung der Harzkanäle und der Beschaffenheit der Pollenkittstoffe. *Planta* 15: 506-529.
- Porsch, O. 1911. Die Anatomie der Nähr und Haftwurzeln von *Philodendron sellowii* C. Koch. *Denkschr. Kaiserl. Akad. Wiss., Math.-Naturwiss. Kl.* 79: 390-451, tt. 24-41.

- Putz, F. E. & N. M. Holbrook. 1986. Notes on the natural history of hemiepiphytes. *Selbyana* 9: 61-69.
- Raskin, I. 1992. Salicylate, a new plant hormone. *Pl. Physiol.* 99: 799-803.
- , I. M. Turner & W. R. Melander. 1989. Regulation of heat production in the inflorescences of an arum lily by endogenous salicylic acid. *Proc. Natl. Acad. Sci. U.S.A.* 86: 2214-2218.
- , A. Ehmann, W. R. Melander & B. J. D. Meese. 1987. Salicylic acid: A natural inducer of heat production in *Arum* lilies. *Science* 237: 1601-1602.
- Raven, P. H. & D. I. Axelrod. 1974. Angiosperm biogeography and past continental movements. *Ann. Missouri Bot. Gard.* 61: 539-673.
- Ray, T. S. 1986. Growth correlations within the segment in the Araceae. *Amer. J. Bot.* 73: 993-1001.
- . 1987a. Leaf types in the Araceae. *Amer. J. Bot.* 74: 359-1372.
- . 1987b. Diversity of shoot organization in the Araceae. *Amer. J. Bot.* 74: 1373-1387.
- . 1988. Survey of shoot organization in the Araceae. *Amer. J. Bot.* 75: 56-84.
- Ritterbusch, A. 1971. Morphologische Untersuchungen zur Wuchsform von *Philodendron*. *Bot. Jahrb. Syst.* 90: 527-649.
- Schutz, G. 1990. Chapter 7. Some aspects of pollination biology in Central American forests. Pp. 69-84 in K. S. Bawa & M. Hadley (editors), *Reproductive Ecology of Tropical Forest Plants*. Parthenon Publ. Group, Park Ridge, New Jersey.
- Schott, H. W. 1829. Für Liebhaber der Botanik. *Wiener Z. Kunst* 1829: 780.
- . 1832. Araceae. Pp. 16-22 in H. Schott & S. Endlicher, *Meletemata botanica*. Typis Caroli Gerold, Vienna.
- . 1856. Synopsis aroidearum complectens, Enumerationem systemicum generum et specierum hujus ordinis. Typis Congregationis Mechitharisticae, Vienna.
- . 1858. Genera Aroidearum. Typis Caroli Ueberreuter. Prostat Olomeui apud Ed. Hölzel, Vienna.
- . 1860. Prodromus systematis aroidearum. Typis Congregationis Mechitharisticae, Vienna.
- Seymour, R. S., G. A. Bartholomew & M. C. Barnhart. 1983. Respiration and heat production by the inflorescence of *Philodendron sellowii* Koch. *Planta* 157: 336-343.
- , M. C. Barnhart & G. A. Bartholomew. 1984. Respiratory gas exchange during thermogenesis in *Philodendron sellowii* K. Koch. *Planta* 161(3): 229-232.
- Sheridan, W. F. 1960. The Occurrence of a Temperature Fluctuation in the Spadix of *Philodendron sellowii*. Master's Thesis, University of Florida, Gainesville.
- Solereder, H. & F. J. Meyer. 1928. Systematische Anatomie der Monokotyledonen. Heft 13: 100-169.
- Sprengel, K. P. J. 1826. *In: Caroli Linnaei... Systema vegetabilium*. Ed. 16, 3. Göttingen.
- Standley, P. C. 1937. Flora of Costa Rica, Part 1. *Publ. Field Mus. Nat. Hist., Bot. Ser.* 18: 131-146.
- . 1944. Araceae. *In: R. E. Woodson, Jr. & R. Schery (editors), Flora of Panama*. *Ann. Missouri Bot. Gard.* 31: 1-60.
- & J. A. Steyermark. 1958a. Studies of Central American plants-III. *Publ. Field Mus. Nat. Hist., Bot. Ser.* 23: 1-28.
- & ———. 1958b. Araceae. Pp. 304-363 in: *Flora of Guatemala. Part 1. Fieldiana, Bot.* 24.
- Takhtajan, A. 1969. *Flowering Plants: Origin and Dispersal*. Smithsonian Institution Press, Washington, D.C.
- Tieghem, P. van. 1907. Recherches sur la structure des Aroidees. *Ann. Sci. Nat. Bot., ser. 9*, 5: 312-320.
- Torre, P. F. 1965. *Notas geológicas de la República de Panamá*. Comisión del Atlas de Panamá, Atlas de Panamá.
- Tournefort, J. P. 1700. *Institutiones rei herbariae*, editio altera. 1: 158-162. Paris.
- Urban, I. 1898. *Symbolae antillanae I. Bortraegero*, Berlin.
- Urdentlich, A., R. A. Linzer & I. Raskin. 1991. Alternative respiration and heat evolution in plants. *Pl. Physiol.* 97: 1545-1550.
- Usheer, G. 1966. *A Dictionary of Botany*. D. Van Nostrand, Princeton, New Jersey.
- Van der Pijl, L. 1937. Biological and physiological observations on the inflorescence of *Amorphophallus*. *Revue Trav. Bot. Néerl.* 34: 157-167.
- Vellozo, J. M. da Conceição. 1825 [1829]. *Flores Fluminenses*. Archivos do Museu Nacional Fluminense Janeiro. [Text corresponding to Vols. 1-8 of the plates.]
- Venenanat, E. P. 1800. Description des plantes nouvelles et peu connues. *Crapelet*, Paris.
- Walker, D. B., J. Gysi, L. Sternberg & M. J. DeNiro. 1983. Direct respiration of lipids during heat production in inflorescence of *Philodendron sellowii*. *Science* 220: 419-421.
- Warming, E. 1867. Nogle lagtagelser over Varmeudviklinger hos en Aroidee, *Philodendron lundii*. *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 1867(8-11): 127-144. t. 4.
- . 1883. *Tropische Fragmente. I. Die Bestäubung von Philodendron bipinnatifidum* Schott. *Bot. Jahrb. Syst.* 4: 328-340.
- Went, F. A. F. C. 1893. Ueber Haft- und Nahrwurzeln bei Kletterpflanzen und Epiphyten. *Ann. Jard. Bot. Buitenzorg* 12: 1-72.
- Willdenow, C. L. 1805. *In: Caroli a Linné Species Plantarum... Editio quarta* 4.
- Young, H. J. 1986. Beetle pollination of *Dieffenbachia longispatha* (Araceae). *Amer. J. Bot.* 73: 931-944.
- . 1987. Aroid observations: *Philodendron rothschuhianum*. *Aroideana* 10: 22.
- . 1988a. Differential importance of beetle species pollinating *Dieffenbachia longispatha* (Araceae). *Ecology* 69: 832-844.
- . 1988b. Neighborhood size in a beetle-pollinated tropical aroid: Effects of low density and asynchronous flowering. *Oecologia* 76: 461-466.
- . 1990. Chapter 11. Pollination and reproductive biology of an understory neotropical aroid. *In: K. S. Bawa & M. Hadley (editors), Reproductive Ecology of Tropical Forest Plants*. Parthenon Publ. Group, Park Ridge, New Jersey.
- Zavada, M. S. 1983. Comparative morphology of monocot pollen and evolutionary trends of apertures and wall structures. *Bot. Rev. (Lancaster)* 49: 331-379.











PHILODENDRON	SEED SHAPE	LEAF	LOCULES						PLACEN- TATION	OVULES/ LOCULE	Ovule/Locule Min	Ovule/Locule Max	OVULE		OVULE LENGTH (mm)	ENVELOPE CONDITION	
			L	P	E	M	Max	Min					Min	Max			AMPLING
grandipet	PPP cordate	B	5	5	1	1	1.00	1.50	axile	16-22	16	22	0.30	0.40	2-series	none	
granulata	CGG oblong	B	6	4	5	7	1.00	1.30	sub	1	1	0.30	0.50	seeds only	transp gel (leaves)		
grayana	OAM cordate	B	7	8	1	1	1.00	1.30	sub	(2)3-8	2	8	0.30	0.50	2-series	transp gel (leaves)	
hemmeli	PA cordate	B	6	6	1	1	1.00	1.30	axile	14	14	0.10	0.20	0.1-0.5	2-series	none	
herbertum	PPP cordate	B	5	8	1	1	1.30	1.50	axile	18-24	18	24	0.20	0.2	2-series	none	
herbertum var. hirsutum	PS cordate	B	4	6	8	8	8.00	8.00	axile	20-25	20	25	0.10	0.10	2-series	none	
hirsutum var. kirkbridei	PS cordate	B	5	5	6	6	6.00	6.00	axile	20-25	20	25	0.10	0.10	2-series	none	
hirsutum var. oycardium	PS cordate	B	5	5	6	6	6.00	6.00	axile	20-25	20	25	0.10	0.10	2-series	none	
helense	CO subcord	Cv	6	9	0.90	1.20	0.90	1.20	sub	(1)3-4	1	4	0.30	0.40	1-series	basal	
imbratum	CGG oblong	B	4	5	0.70	1.30	0.70	1.30	sub	1 (2)	1	2	0.40	0.70	transp	transp	
jacquini	MAC cordate	V	4	4	1.30	1.30	sub	sub	2	2	2	0.60	1.00	2-series	none-gel matrix		
obese	OMR cordate	C	7	8	0.32	3.20	axile	axile	6 seeds	6 seeds	6	6	0.30	0.30	2-series	none	
obovatum	PPP cordate	C	4	6	1.20	1.70	axile	axile	18-28	18	28	0.4	0.2	0.2	2-series	none?	
ocypete	OAM cordate	C	4	4	0.65	0.80	axile	axile	4	4	4	0.2	0.2	0.2	2-series	none	
obovatum	PPP cordate	C	4	4	1.10	1.60	axile	axile	18-14	10	14	0.30	0.30	0.50	0.3-0.5	1-2 series	trans
ovatum	OME subcord	Cv	6	7	1.20	2.00	sub	sub	(1-2)4-5	1	6	0.30	0.40	0.3-0.4	trans or gel		
liquidum var. liquidum	CGG oblong	Cv	6	8	0.65	2.40	sub	sub	1	1	1	0.30	0.40	0.4	trans		
liquidum var. hirsutum	CGG oblong	C	5	6	1.60	1.60	sub	sub	2	2	2	0.40	0.50	0.4-0.5	trans		
liquidum var. ovatum	CGG oblong	C	5	6	1.60	1.60	sub	sub	2 (seeds)	6	20	0.20	0.30	0.2-0.3	transp-basal		
ligense	PPP cordate	C	6	6	1.50	6.60	axile	axile	(6)12-14(20)	1	1	1	0.30	0.30	seeds only	transp	
madroaense	TRU 3-lobed	C	6	7	2.30	2.30	sub	sub	20	20	20	1	0.30	0.1-0.3	2-3 series	transp	
melanoceras	PA cordate	V	5	6	1.30	1.60	axile	axile	20	20	20	1	0.30	0.1-0.3	2-3 series	transp	
melanoceras var. melanoceras	OAM cordate	V	5	6	0.90	2.60	sub	sub	1-2(3)	1	3	0.30	1.00	0.3-0.6(1.0)	digitate		
microstichum	CGG subcord	V	6	7	0.80	0.80	sub	sub	1	1	1	0.4	0.4	0.4	basal		
mon	CGG subcord	C	4	5	1.30	1.30	sub	sub	1	1	1	1	0.4	0.4	seeds only	?	
nigricanum	OME cordate	C	?	?	?	?	?	?	?	?	?	?	?	?	?	?	
patersoniae	PPP cordate	C	4	6	1.00	3.00	axile	axile	20-31	20	31	0.30	0.30	0.30	2-series	none	
patersoniae	PPP cordate	C	4	6	1.70	1.70	axile	axile	20	20	20	0.20	0.20	0.2-0.3	2-series	none	
patersoniae	OMM cordate	V	7	8	0.70	1.50	sub	sub	3	3	3	0.20	0.50	(0.2)0.4(0.5)	1-series	transp-basal	
platyphallum	CGG oblong	C	6	6	1.40	1.10	sub	sub	1-2(4)	20	20	0.20	0.30	0.4-0.5	1-series	transp & gel	
pseudocucullatum	PPP cordate	C	6	6	1.50	1.90	axile	axile	20	20	20	0.20	0.30	0.2-0.3	2-series	none?	
pluratum	PS cordate	V	4	5	6.20	6.20	axile	axile	15-20	15	20	0.20	0.50	0.2-0.25	2-3 series	none	
purpurascens	PPP cordate	C	6	7	5.00	5.00	axile	axile	13-20	13	20	0.10	0.10	0.1	2-3 series	none	
purpurascens	POL incis-lob	C	5	7	1.60	6.00	axile	axile	8	8	8	0.10	0.30	0.1-0.3	2-series	transp-gel	
radicum var. radicum	POL incis-lob	C	5	7	1.60	6.00	axile	axile	3	3	3	0.30	0.30	0.3	1-series	transp	
radicum var. pseudoradicatum	PC oblong	C	5	7	1.40	1.10	axile	axile	14-18	14	18	0.30	0.50	0.3-0.5	2-series	none	
radicum var. roseospathum	PC oblong	C	4	4	1.10	1.10	axile	axile	6-8	6	8	0.30	0.30	0.3	1-series	none	
roseospathum var. angustilobum	TRU 3-lobed	C	5	3	1.00	6.30	sub	sub	2-4	2	4	0.10	0.50	0.1-0.5	1-2 series	transp to gel	
schubertianum	OMM cordate	C	6	2	1.00	6.30	sub	sub	(10)12-14	10	14	0.30	0.40	0.3-0.4	2-series	none	
seguinianum	PPP subcord	C	4	6	0.70	3.00	axile	axile	10-14(16)	10	15	0.10	0.30	0.1-0.3	2-series	transp or transp	
schubertianum	PPP cordate	C	4	6	1.90	2.50	axile	axile	10-14(16)	10	15	0.10	0.30	0.1-0.3	2-series	transp or transp	

Appendix 2. Continued.

PHILODENDRON	SECT	LEAF	H	S	A	TT	B	YY	I	L	P	LOCULE		PLACEN	TATION	OVULES/LOCULE	OVULE		ARRANG	ENVELOPE	CONDITION
												Min	Max				Min	Max			
smithii	CGO cordate	C	B	7	B	0	0	110	basal (sub)	1	1	1	1	0.30	0.40	0.3-0.4				transp.	
rossiae	OMM cordate	C	D	6	0	0.40	1.40	basal (sub)	2	2	2	2	0.10	0.20	0.1-0.2				transl transp gel		
squamocaula	PA cordate	C	D	4	4	1.00	1.20	axile	20-28	20	28	20	0.20	0.20	0.2				1-2 series		
squamipetiolum	PA subcord	V	B	4	5	1.00	1.20	axile	20-30	20	30	20	0.10	0.20	0.1-0.2				2-3 series		
straminea	CGO cordate	C	B	5	6	0.50	2.10	sub	1	1	1	1	0.20	0.30	0.2-0.3				ph-3y, transp		
stratum	PP1 cordate	C	B	5	6	1.60	1.70	axile	20-28	20	28	20	0.20	0.40	0.2-0.4				2-series		
subnigrum	OMM incisi-lob	C	B	4	4	0.40	0.90	axile	16	16	16	16	0.40	0.40	0.4				2-series		
subulocaula	CGO cordate	C	B	4	5	0.40	0.90	sub	1	1	1	1	0.20	0.30	0.2-0.3				transl or trans;		
tenue	PPF cordate	C	B	4	5	1.60	5.90	axile	12-14	12	14	12	0.40	0.40	0.4				2-series		
thalictrum	PP1 cordate	C	B	5	7	2.00	4.90	axile	16-22	16	22	16	0.40	0.40	0.2-0.4				none		
trichostemum	TR 3-lobed ov(B)	S	B	0.90	2.30	sub	1-2	sub	1-2	1	2	1	0.40	0.60	0.4-0.6				1-series		
tyocata	OMR cordate	C	B	5	6	0.80	1.30	sub	(4) 5-7	4	7	4	0.10	0.40	0.1, 0.3-0.4				2-series		
unguipetense	CGG oblong	V	D	8	0	1.10	1.10	sub	1	1	1	1	0.3	0.3	0.3				transp		
uruguayense	CGG oblong	C	7	4	5			sub	1	1	1	1									
venezuelense	OMM cordate	V	D	4	6	0.30	0.40	sub	2	2	2	2	0.4	0.4	0.4				transp & gel		
venezuelense	PA cordate	C	D	4	5	1.30	2.50	axile	20-24(32)	20	32	20	0.10	0.20	0.1-0.2				2-series		
venezuelense	POL incisi-lob	C	C	4	5	2.60	9.50	axile	3-4	3	4	3	0.50	0.50	0.5				1-2 series		
wendlandii	CGG oblong	C	B	6	7	0.70	1.30	sub	2	2	2	2	0.30	0.50	0.3-0.5				thick transl matine		
wilburii var wilburii	CGO cordate ov(B)	B	B	6	7	0.70	2.00	sub	1-2	1	2	1	0.30	0.50	0.3-0.5				1-series		
wilburii var longipedunculatum	CGO cordate ov(B)	B	B	6	7	0.40	1.50	sub	1-2	1	2	1	0.20	0.60	0.2-0.3(0.6)				transp-gel		
zuberium	OMM cordate	C	B	1	3	0.60	1.20	basal	1-3	1	3	1	0.30	0.40	0.3-0.4				1-series		

Code	Section	Subsection	Series	Code	Section	Subsection	Series
PP1	Philodendron	Philodendron	Impolita	sum	324	439	11
PP2	Philodendron	Philodendron	Impolita	count	46	46	44
PP3	Philodendron	Philodendron	Velutina	average	7.04	9.54	0.24
PP4	Philodendron	Philodendron	Ferrosa				
PPA	Philodendron	Philodendron	Atrorosea				
PM	Philodendron	Macrorrhynchium					
PC	Philodendron	Macrorrhynchium					
PL	Philodendron	Cerriophyllum					
PS	Philodendron	Solanostigma					
PA	Philodendron	Achyrocline					
OMM	Celestigma	Macrobellium					

Code	Section	Subsection	Series
OMR	Celestigma	Macrobellium	
OMX	Celestigma	Macrobellium	
OMY	Celestigma	Macrobellium	
OMP	Celestigma	Macrobellium	
OGG	Celestigma	Glossophyllum	
OGD	Celestigma	Glossophyllum	
OO	Celestigma	Chlocoepidium	
OB	Celestigma	Bulbovia	
OS	Celestigma	Evcardium	
TR	Tetramophyllum		
POL	Polytonium		
MAC	Macrogynium		

c = climber  
v = vine  
1 = terrestrial

Appendix 3. Sectional Composition of *Philodendron* subg. *Philodendron* in Central America with number of species in Central America.

## SECTION BAURISIA

Section 1. *Philodendron* sect. *Baurisia* (Rehb. ex Schott) Engl. in Mart. Fl. bras. 3(2): 134. 1878 ..... None

## SECTION PHILOPSAMMOS

Section 2. *Philodendron* sect. *Philopsamos* G. S. Bunting, Phytologia 60(5): 306. 1966 ..... None

## SECTION PHILODENDRON

Section 3. *Philodendron* (*Philodendrum*) Schott

Subsection 1. *Macrolonchium* (Schott) Engl., Martius, Fl. Bras. 3(2): 139. 1878 ..... 1 sp.

Subsection 2. *Canniphyllum* (Schott) Mayo, Bot. J. Linn. Soc. 100. 168. 1989 ..... 3 sp., 1 subsp.

Subsection 3. *Platyopodium* (Schott) Engl., Martius, Fl. Bras. 3(2): 137. 1878 ..... 5 sp.

Subsection 4. *Psoropodium* (Schott) Engl., Martius, Fl. Bras. 3(2): 138. 1878 ..... None

Subsection 5. *Solenasterigma* (Klotzsch ex Schott) Engl., Martius, Fl. Bras. 3(2): 139. 1878 ..... 3 sp., 3 subsp.

Subsection 6. *Philodendron* ..... 20 sp.

Series 1. *Philodendron* ..... None

Series 2. *Impolita* ser. nov ..... 3 sp.

Series 3. *Vesista* ser. nov ..... 1 sp.

Series 4. *Fibrosa* ser. nov ..... 15 sp.

Series 5. *Albisuccosa* ser. nov ..... 1 sp.

Subsection 7. *Achyropodium* (Schott) Engl., Martius, Fl. Bras. 3(2): 139. 1878 ..... 6 sp.

## SECTION CALOSTIGMA

Section 4. *Calostigma* (Schott) Engl., Martius, Fl. Bras. .... 48 sp., 4 subsp.

Subsection 1. *Macrobelyum* (Schott) Engl., Martius, Fl. Bras. 3(2): 143. 1878.

Series 1. *Macrobelyum* ..... 16 sp.

Series 2. *Ecordata* ser. nov ..... 5 sp., 2 subsp.

Series 3. *Reticulata* ser. nov ..... 2 sp.

Series 4. *Pachycaulia* ser. nov ..... 1 sp.

Subsection 2. *Glossophyllum* (Schott) Croat, comb. nov.

Series 1. *Glossophyllum* Schott ..... 15 spp., 3 subsp.

Series 2. *Osata* ser. nov ..... 7 spp., 2 subsp.

Subsection 3. *Oligocarpidium* Engl., Engler's Bot. Jahr. 26: 535. 1899 ..... 2 spp.

Subsection 4. *Bulaoanum* Mayo, Bot. J. Linn. Soc. 100: 168. 1989 ..... None

Subsection 5. *Eucardium* (Engl.) Mayo, Bot. J. Linn. Soc. 100: 168. 1989 ..... None

## SECTION TRITOMOPHYLLUM

Section 5. *Tritomophyllum* (Schott) Engl., Martius, Fl. Bras. 3(2): 144. 1878 ..... 6 spp.

## SECTION SCHIZOPHYLLUM

Section 6. *Schizophyllum* (Schott) Engl., Martius, Fl. Bras. 3(2): 144. 1878 ..... None

## SECTION POLYTOMIUM

Section 7. *Polytomium* (Schott) Engl., Martius, Fl. Bras. 3(2): 145. 1878 ..... 3 spp., 2 taxa

## SECTION MACROGYNIUM

Section 8. *Macrogygium* Engl., Bot. Jahrb. 26: 553. 1899 ..... 1 sp.

## SECTION CAMPTOGYNIUM

Section 9. *Camptogygium* K. Krause, in Engler, Das Pflanzenreich IV. 23Db (Heft 60): 3, 127. 1913 ..... None

Appendix 4. Phenological Patterns of Central American *Philodendron* subg. *Philodendron*.

PHILODENDRON	Flowering Pattern						
	All Year	Wet	Wet->Dry	Dry	Dry-> Wet	Bimodal	Unknown
advena	1						
albisuccus		1					
alticola					1		
angustilobum					1		
anisotomum			1				
annulatum					1		
antonianum		1					
aromaticum						1	
auniculatum					1		
bakeri				1			
basi		1					
breedlovei		1					
brenesii					1		
brevispalum					1		
brewsterense				1			
brunneicaule					1		
chiriquense		1					
chiripoense				1			
clewella					1		
coloradense		1					
copenae						1	
corrae		1					
colobrusense		1					
colonense					1		
crassispalum					1		
cretosum					1		
dauidsonii subsp. dauidsonii					1		
subsp. bocatoranum+A50		1					
dodsonii		1					
dolichophyllum					1		
dominicalense		1					
dressleri		1					
dwyeri				1			
edenudatum				1			
ferrugineum		1					
findens					1		
folsomii				1			
fortunense					1		
fragrantissimum					1		
gigas		1					
glanduliferum subsp. glanduliferum		1					
grandipes					1		
granulare		1					
grayumii					1		
harmeli		1					
hebetatum					1		
hederaceum var. hederaceum	1						
var. kirkbridei					1		
var. oxycardium							1
heleniae					1		
immedium					1		
jacquini		1					

## Appendix 4. Continued.

PHILODENDRON	Flowering Pattern						
	All Year	Wet	Wet->Dry	Dry	Dry-> Wet	BiModal	Unknown
efense		1					
jodavisanum					1		
knappiae				1			
lazoni					1		
lenti					1		
ligulatum var. ligulatum							1
var. heracleoanum							1
var. ovatum							1
llanense					1		
macronense		1					
malsevichiae					1		
mexicanum					1		
microbictum					1		
moii						1	
niqueanum		1					
panamense					1		
pirense		1					
platypetiolatum					1		
pseudauriculatum					1		
pterolum					1		
purpureovide	1						
purulense		1					
radiatum var. radiatum	1						
var. pseudoradiatum	1						
roseospathum var. roseospathum					1		
var. angustilaminatum		1					
rothschuhianum					1		
sagittifolium					1		
scaleriense					1		
schoisianum					1		
smithi					1		
sousae		1					
squamicaule		1					
squamipetiolatum					1		
stramineicaule					1		
strictum					1		
subincisum							1
sulcicaule					1		
tenue					1		
thalassicum					1		
tripartitum					1		
tysoni					1		
ubigantupense		1					
utleyanum		1					
verapazense					1		
verrucosum					1		
warszewiczii					1		
wendlandi					1		
wilburii var. wilburii							1
var. longipedunculatum				1			
zhuianum		1					
TOTAL	5	29	2	7	51	4	5

Appendix 4. Continued.

PHILODENDRON	Flowering Pattern						
	All Year	Wet	Wet->Dry	Dry	Dry->Wet	Bimodal	Unknown
% of Total	4.85%	28.16%	1.94%	6.00%	49.51%	3.88%	4.85%



Figures 1-4. —1 (top L). *Philodendron radiatum*, showing adventitious roots. Cultivated at Summit Gardens. —2 (top R). *P. schottianum* (Croat 66395), showing stems with transverse fissures and coarse striations at apex of node. —3 (bottom L). *P. malesreichiae*, showing trichome-like scales (Croat 76707). —4 (bottom R). *P. rothschuhianum*, showing stems that dry with minute cracks and scurfy condition (Croat 57199).





Figures 5-8. —5 (top L). *Philodendron conforme* (Venezuela), showing stems with transverse fissures at points of stress, petiole scar, and scar of inflorescence (Croat 59355). —6 (top R). *P. stramineaele*, dried stem showing conspicuous ribbing caused by drying (Croat & Grayum 60057). —7 (bottom L). *P. fendens*, dried stem showing interruption of conspicuous epidermis by petiole base (Croat 66811). —8 (bottom R). *P. radiatum* Schott var. *pseudoradiatum*, showing stems with cracking and protruding epidermis (Croat & Hannon 63381).



Figures 9–12. —9 (top L). *Philodendron rutachukianum*, showing inconspicuous cataphyll scars and petiole scars (Croat 57199) (photo: P. Malesevich). —10 (top R). *P. davidsonii* (Davidson 3956), showing stem with conspicuous cataphyll scars and petiole scars. —11 (bottom L). *P. megalophyllum* (Croat 54252), stem with intravaginal squamulae occurring just above the cataphyll scar. —12 (bottom R). *P. auriculatum* (Croat 59730), stem with spreading anchor roots showing spinelike root branch buds.




Figures 13-16. 13, 14, *Philodendron lazorii* (Croat 69833). —13 (top L). Showing clasp roots closely appressed to tree. —14 (top R). Showing clasp roots and a much larger feeder root. —15 (bottom L). *P. wendlandii*, showing persistent, intact cataphylls held in place by contiguous petioles (Croat 69732). —16 (bottom R). *P. warszewiczii*, showing weakly 2-ribbed, short-lineate cataphyll (Croat & Hannon 64186).




Figures 17-20. —17 (top L). *Philodendron schottianum*, showing bluntly 2-ribbed cataphyll (Croft & Zhu 76518). —18 (top R). *P. megalophyllum*, showing petioles which are narrowly sulcate with broadly rounded margins (Croft 542524). —19 (bottom L). *P. davidsonii* subsp. *bocatoranum*, showing sweet, viscid droplets and combination of short-lineate and lineate-streaked markings (Croft 38177). —20 (bottom R). *P. auriculatum*, showing dark green ring demarcating petiole and blade and free basal veins (Croft & Grayum 59730).



Figures 21-24. —21 (top L). *Philodendron malesichiae*, showing snapped petiole with slender strands of latex (Croat & Bay 75810). —22 (top R). *P. giganteum* (Croat 68568), showing midrib, primary lateral veins, and basal veins (the lowermost primarily fused to form posterior rib). —23 (bottom L). *P. auriculatum*, showing quilted primary lateral veins (Croat & Grayum 59730). —24 (bottom R). *P. megalophyllum*, showing arrangement of veins: midrib, primary lateral veins with lowermost branching (these secondary veins), and interprimary veins (Croat 53901).



Figures 25-28. —25 (top L). *Philodendron maleseichiae*, showing lowermost primary lateral veins and basal veins (these loosely coalesced toward the base) (Croft 74818) (photo: P. Malesevich). —26 (top R). *P. megalophyllum*, showing minor veins arising from the midrib and primary lateral veins (Croft 53901). 27, 28. *P. maleseichiae* (Croft & Zhu 76707). —27 (bottom L). Fresh spadix with pistillate flowers at base, sterile staminate flowers in middle, and a portion of the fertile staminate spadix at apex. —28 (bottom R). Older spadix showing the discolored staminate spadix and white sterile staminate portion.



Figures 29–32. —29 (top L). *Philodendron tysonii*, showing spadix with sterile male flowers having been consumed by beetle pollinators (Croat 67577). —30 (top R). *P. fragrantissimum*, pollen grain (Grayum 2436, DUKE). Photo: M. H. Grayum. —31 (bottom L). *P. rothschildianum*, showing strands of pollen emerging from anthers (Croat 35657). —32 (bottom R). Insect visitors to newly opening spadix of *P. anisotomum*. Large beetles shown (*Eriocelis colombica*) are legitimate pollinators; smaller insects (*Noella*, order Hemiptera) are frequent visitors to *Philodendron* but play no known role in their biology. Photo: H. Young.




Figures 33-36. —33 (top L). *Philodendron hebetatum*, showing spathe beginning to break up to expose mature berries (Croat & Monsalve 61396). —34 (top R). *P. findens*, showing scar left by fallen spathe and coarsely striate peduncle apex (Croat 38218). —35 (bottom L). *P. pseudauriculatum*, showing mature infructescences with spathe and staminate portion of spadix fallen (Croat & Zhu 76251). —36 (bottom R). *P. brenesii*, showing infructescence with exfoliating spathe and partially consumed berries (Croat 35519).





Figures 37-40. *Philodendron odivna*. —37 (top L). (Croat & Hannon 63107) Adult blade. —38 (top R). (Croat & Hannon 65549) Stem with two pre-anthesis inflorescences and semipersistent cataphylls and an elongated petiole sheath. —39 (bottom L). (Croat & Hannon 63107) Inflorescences (one showing spathe already fallen). —40 (bottom R). (Croat & Hannon 63430) Inflorescence nearing maturity (showing cut open view). Because of oblique attachment, adaxial side of spadix is longer than the abaxial side.



Figures 41-44. 41-43, *Philodendron albisuccus*. 41, 42 (top L & R). (Croat 3785J). —43 (bottom L). (Croat 68940).  
—44 (bottom R). *P. anisotomum* (Croat & Hannon 64364).



Figures 45-48. *P. alicicola* (Croat 74906). —45 (top L). Stem with post-anthesis inflorescence. —46 (top R). Juvenile and pre-adult leaves. —47, 48 (bottom L & R). Adult leaves.



Figures 49-52. *Philodendron angustilobum*. —49 (top L). (Croft 61162) Habit, in cultivation. —50 (top R). Plants displaced from trees, with inflorescences (Croft & Hannon 64522). 51, 52. (Croft 61162). —51 (bottom L). Inflorescence at anthesis. —52 (bottom R). Habit in cultivation.



Figures 53-56. *Philodendron anisotomum*. —53 (top L). Habit in cultivation (Berlin Botanical Garden (031-60-74-83)). —54 (top R). Stem with anchor roots spreading from stem. —55 (bottom L). Leaf in cultivation (Croat & Hanson 64364). —56 (bottom R). Plant with inflorescence in cultivation (Selby 80-1668, from Mexico, Chiapas, Ocosingo).



Figures 57-60. *Philodendron annulatum*. 57-59, (Croat 74803). —57 (top L). Habit. —58 (top R). Stem and inflorescences. —59 (bottom L). Leaf with petiole showing purple ring at apex. —60 (bottom R). (Croat 74855) Showing stem with cataphyll, unopened inflorescence, and bracteole.



Figures 61-64. *Philodendron antonioanum* (Croat & Zhu 76909). —61 (top L). Habit. —62 (top R). Leaf. —63 (bottom L). Inflorescences partially obscured by cataphylls. —64 (bottom R). Leaves, both lower and upper surfaces.



Figures 65-68. —65 (top L). *Philodendron auriculatum*, habit (cultivated by John Banta). 66-68. *P. aromaticum*. 66, 67. (Croft 68382). —66 (top R). Habit. —67 (bottom L). Stem and inflorescences. —68 (bottom R). (Croft 68423) Open inflorescence.





Figures 69-72. *Philodendron auriculatum*. —69 (top L). (Croat 59730) Habit in cultivation. —70 (top R). (Croat & Grayum 35205) Open inflorescence. 71, 72. (Croat 32956) (photo: P. Malesevich). —71 (bottom L). Open inflorescence showing protruded spadix. —72 (bottom R). Sharply 2-ribbed cataphyll.



Figures 73-76. —73 (top L), *Pileolodendron aderseni* (Croat 47777), habit. —74 (top R), *P. angustilobum*, post-anthesis inflorescences, showing spathe with spathe cut away and the discoloration of the staminate spathe (Croat 61162). —75 (bottom L), *P. antiochiensis* (Croat 78892), habit. —76 (bottom R), *P. ausii*, habit (cultivated by Conrad Fleming).



Figures 77-80. 77, 78. *Philodendron bakeri*. —77 (top L). Plant hand-held (Croat 37508). —78 (top R). Habit, showing abaxial surfaces of blades (Croat 67456). —79, 80 (bottom L & R). *P. breedlovei* (Breedlove 345181).



Figures 81-84. *Philodendron basii* (Croat 45442). —81 (top L). Habit (cultivated at MO; photo: P. Malesevich). —82, 83 (Cultivated at SEL). —82 (top R). Leaf, abaxial surface. —83 (bottom L). Stem with intact cataphylls and closed petiole sheaths. —84 (bottom R). Stem with inflorescence.



Figures 85-88. —85 (top L). *Philodendron breedlovei*, open inflorescence (Breedlove 35181) (photo: D. Breedlove). 86-88. *P. brenesii*, habit. —86 (top R). (Croat 67578). —87 (bottom L). (Croat 68084). —88 (bottom R). Apex of stem with bases of petioles, cataphyll, and unopened inflorescence (Croat 78806).



Figures 89-92. —89 (top L). *Philodendron brevesii*, habit (Croat 43521), 90, 91. *P. brevispanthum* (Croat 58536), —90 (top R). Habit. —91 (bottom L). Stem with entire cataphyll and unopened inflorescence. —92 (bottom R). *P. brevispanthum* (De Nevers et al. 5545).



Figures 93-96. *Philodendron brunneicaulis*. 93-95. (Croat & Zhu 76581). —93 (top L), Habit. —94 (top R), Plant showing stem attached to small tree with anchoring roots and inflorescence arising from elongated petiole sheath (plant displaced). —95 (bottom L), Habit (plant displaced). —96 (bottom R), Inflorescence cut open (Croat 68713).



Figures 97-100. —97 (top L). *Philodendron brunneicaule*, leaf blade adaxial surface (Croat & Zhu 76581). 98-100. *P. chiriquense* (Croat 69068). —98 (top R). Leaf blade adaxial surface. —99 (bottom L). Stem apex showing inflorescences emerging from cataphyll fibers. —100 (bottom R). Inflorescence with tube portion cut open.





Figures 101-104. —101 (top L). *Philodendron chirripoense* (Burger & Liesner 7139). —102 (top R). *P. clewellii* (Gentry & Clewett 7028). 103, 104. *P. coloradense*. —103 (bottom L). Habit (Croat 75039). —104 (bottom R). Plant hand-held with inflorescences (Croat 37168).



Figures 105-108. 105, 106. *Philodendron coloradense*. —105 (top L), Stem with inflorescences (Croat 37768). —106 (top R), Habit (Croat 75039). —107 (bottom L), *P. chariquense*, habit (Croat 69065). —108 (bottom R), *P. cotense* (plant held by T. K. Croat) (Croat 66262).



Figures 109-111. *Philodendron* coprosae. — 109 (top L). Habit (Croat 60159). — 110 (top R). Cluster of inflorescences (one open) (Croat 44729). — 111 (bottom L). Stem with D-shaped pedicels and unopened inflorescences (Croat 44729). — 112 (bottom R). *P. crassispatum*, habit (Croat 44444).



Figures 113–116. 113–115. *Philodendron copense*. —113 (top L). Leaf blade adaxial surface (Croat 44729). 114, 115. (Croat 68765). —114 (top R). Leaf blade abaxial surface. —115 (bottom L). Inflorescence emerging from cataphyll fibers. —116 (bottom R). *P. correae*, habit (Croat & Zhu 76395).



Figures 117–120. 117–119. *Philodendron correae* (Croat 66748). —117 (top L). Habit. —118 (top R). Habit with inflorescence and infructescence. —119 (bottom L). Plant with mature infructescence and fully sheathed petioles. —120 (bottom R). *P. cotobrusense* (Grayson & Hammel 5689).



Figures 121-124. 121-123. *Philodendron cotonense*. —121 (top L). Habit (Croat 66169). 122, 123. (Croat 66504). —122 (top R). Habit showing abaxial surfaces. —123 (bottom L). Stem with inflorescences. —124 (bottom R). *P. crassipathum*, habit (Croat 33150).



Figures 125-128. 125, 126. *Philodendron cretosum* (Croat & Zhu 76661). —125 (top L). Habit. —126 (top R). Apex of stem with persistent cataphyll fibers, obtusely sulcate petioles, and an unopened inflorescence. 127, 128. *P. crassispatum* (Croat 33150). —127 (bottom L). Habit. —128 (bottom R). Inflorescence with spathe tube cut open.



Figures 129-132. 129, 130. *Philodendron danielsonii* subsp. *danielsonii*, habit in cultivation. —129 (top L.), (Daxidson 70977). —130 (top R.), (Croat 522324). 131, 132. *P. danielsonii* subsp. *locatorum*. —131 (bottom L.), Habit in cultivation (Croat 39177). —132 (bottom R.), Apex of stem with unopened inflorescences (Croat 56655).





Figures 133-136. 133-135. *Philodendron davidsonii* subsp. *locatoranum* (Croat 38177). —133 (top L). Habit in cultivation. —134 (top R). Open inflorescence. —135 (bottom L). Leaf blade abaxial surface. —136 (bottom R). *P. davidsonii* subsp. *davidsonii*, open inflorescence (Davidson 7097).



Figures 137-140. *Philodendron dodsonii*. 137, 138, (Croat 72982). —137 (top L). Habit. —138 (top R). Apex of stem with petioles, cataphylls, and unopened inflorescence. —139 (bottom L). Unopened inflorescence (Croat & Hannon 79114). —140 (bottom R). Inflorescence on abaxial blade surface showing normal open nature.



Figures 141-144. 141-143. *Philodendron dodeonii*. 141, 142. Habit. —141 (top L). (Croat 72982). —142 (top R). (Croat & Hanson 79114). —143 (bottom L). Leaf blade abaxial surface (Croat 72982). —144 (bottom R). *P. crinitum*, habit (Croat 67723).



Figures 145-148. *Philodendron didymophyllum*. —145 (top L). Natural habit on road bank (Croat & Zha 77008), 146, 147. (Croat 67350). —146 (top R). Plant displaced from tree. —147 (bottom L). Apex of stem with unopened inflorescence. —148 (bottom R). Mature inflorescence. Not collected. Photo from El Llano-Carti Road, Panama Province (photo: D. Burne(6)).



Figures 149–152. —149 (top L). *Philodendron crassipathum*, stem with unopened inflorescence (Croat 33150). —150 (top R). *P. dodsonii*, habit with inflorescences arising from elongated petiole sheaths (Croat 72902). —151 (bottom L). *P. dolichophyllum*, habit (Croat 22908). —152 (bottom R). *P. dressleri*, leaf blade abaxial surface (Croat 45360).



Figures 153–156. 153, 154 (top L & R). *Philodendron dominicalense* (Croft 35268). —155 (bottom L). *P. dreyeri* (Dwyer & Lienser 12334). —156 (bottom R). *P. edenudatum*, habit (Croft & Zhu 77157).



Figures 157-160. *Philodendron dressleri*. 157, 158. (Croat 45360). —157 (top L). Habit. —158 (top R). Stem with unopened inflorescences. 159, 160. Cultivated at Joseph Fondeur's. —159 (bottom L). Habit. —160 (bottom R). Apex of stem showing persistent, intact cataphyll at uppermost node and petiole bases with closed sheath.



Figures 161-164. *Philodendron edenudatum*. 161, 162. (Croft & Zhu 77087). —161 (top L). Leaf blade adaxial surface. —162 (top R). Stem apex. 163, 164. (Croft 33988). —163 (bottom L). Blade abaxial surface, showing purplish spots on petiole and lower midrib (photo: P. Malesevich). —164 (bottom R). Inflorescence in cultivation in final stages of closure.





Figures 165–168. *Philodendron ferrugineum*. —165 (top L). Habit (Croat 33732). —166 (top R). Habit (Croat 75155). —167 (bottom L). Cluster of inflorescences (Croat & Zhu 77029), juvenile foliage. —168 (bottom R). Juvenile foliage (Croat 75116).



Figures 169–172. 169–171. *Philodendron findens*. —169 (top L). Showing young blade before shredding occurs (Croat 67153). —170 (top R). Showing divided blades which naturally become pinnate (Croat 67919). —171 (bottom L). Showing winged petiole (Croat & Zhu 76502). —172 (bottom R). *P. folsomii* (McPherson 13619).



Figures 173–176. 173, 174, *Philodendron ferrugineum*. —173 (top L), Habit (Croat 34349), 174, 175, Cluster of inflorescences with one open. —174 (top R), (Croat 25066), —175 (bottom L), *P. fulvum* (Croat 36218), —176 (bottom R), *P. fulvum* (McPherson 13619).



Figures 177-180. 177, 178. *Philodendron findens* (Croat 38218). —177 (top L). Apex of stem with unopened inflorescences. —178 (top R). Open inflorescence. —179 (bottom L). *P. fortuneense*, habit (Croat 67921). —180 (bottom R). *P. glanduliferum*, habit with open inflorescence (hand-held) (Croat 39753).



Figures 181-184. *Philodendron fortianense*. —181 (top L), Habit (hand-held by author) (Croat 50000). —182 (top R), Habit (Croat 66714). —183 (bottom L), Leaf showing abaxial surface and winged petiole (Croat 48727). —184 (bottom R), Stem apex with cluster of immature inflorescences (Croat 66714).



Figures 185-188. —185 (top L). *Philodendron ferrugineum*, cluster of inflorescences with one cut open (Croat 250908), 186-188. *P. fragrantissimum*, 186, 187 (top R & bottom L). (Gonselle 5975). Cultivated at the University of Utrecht Botanical Garden. —188 (bottom R). Habitat with leaves with D-shaped petioles and inflorescences emerging from cataphylls (Croat 11526).



Figures 189–192. 189–191. *Philodendron fragrantissimum*. —189 (top L). Habit (Croat 11526). —190 (top R). Apex of stem with persistent cataphyll fibers (Croat 53912). —191 (bottom L). Juvenile leaves (Croat 9003). —192 (bottom R). *P. gigas*, stem with petiole bases and intact cataphyll fibers (Croat 33680).




Figures 193-196. *Philodendron gigas* (Croat & Zhu 76988). —193 (top L). Plant on ground with loose inflorescences. —194 (top R). Plant held by author. —195, 196 (bottom L & R). Loose inflorescences showing unsopened spathe and prophylls.





Figures 197-200. *Philodendron glanduliferum*. 197-199. *P. glanduliferum* subsp. *camillosonum*, 197, 198 (top L. & R). Cultivated by Roberto Barthe-Marx. —199 (bottom L.). Open inflorescence (Croat 620659) (Venezuela). —200 (bottom R). *P. glanduliferum* subsp. *glanduliferum*. Cultivated at the Munich Botanical Garden (coll. (Forster s.n.), San Jose, Guatemala).



Figures 201-204. 201, 202. *Philodendron glanduliferum* var. *glanduliferum*. —201 (top L). (Croat 43909) Leaves showing petiolar glands. —202 (top R). Showing weathered cataphyll fibers (Croat 39753). 203, 204. *P. grandipet.* —203 (bottom L). Habit (displaced) (Croat & Watt 70200). —204 (bottom R). Apex of stem with intact cataphylls and unopened inflorescences (Croat 61323) (Colombia, Valle: Bujo Calima).



Figures 205–208. —205 (top L), *Philodendron gigas*, blade adaxial surface with unopened inflorescence (Croat & Zhu 76988). —206 (top R), *P. grandipes*, open inflorescence (Croat 33648) (photo: P. Malesevich). —207, 208 (bottom L & R), *P. granulare* (Croat & Porter 15543).



Figures 209–212. *Philodendron grayumii*. —209 (top L). Leaf blade adaxial surface (Croat 66814). —210 (top R). Cluster of unopened inflorescences (Croat & Zhu 76524). —211 (bottom L). Blade adaxial surfaces (Croat 66969). —212 (bottom R). Apex of stem with open inflorescence; petiole with obtusely sulcate base (Croat 74840).



Figures 213-216. 213-215. *Philodendron grousamii*, 213, 214 (Croat 66794), —213 (top L), Habit. —214 (top R), Habit. —215 (bottom L), (Croat & Zhu 77202) Open inflorescence. —216 (bottom R), *P. grandipes*, habit (Croat & Zhu 76664), Habit.



Figures 217-220. —217 (top L), *Philodendron hamamelii* (Hamamel 3551), 218-220, *P. acetosatum*. —218 (top R), Habit (Croat 56715). —219 (bottom L), Cluster of inflorescences (Croat 73150). —220 (bottom R), Open inflorescence in cultivation (Croat 33773).



Figures 221-224. *Philodendron hederaceum*. 221-223. *P. hederaceum* var. *hederaceum*. 221, 222. Habit. —221 (top L). (Croat 70891). —222 (top R). (Croat 69834). —223 (bottom L). Cultivated at Kiev Botanical Garden (juvenile form with velvety leaves). —224 (bottom R). *P. hederaceum* var. *kirkbridei*, close-up of stem showing minutely warty surface and anchor roots (Croat 75108).



Figures 225–228. 225, 226. *Philodendron hebetatum*. —225 (top L). Leaf blade adaxial surface (Croft 73150). —226 (top R). Open inflorescence (Croft 69231). 227, 228. *P. heleniae* (Croft & Zhu 76738). —227 (bottom L). Habit. —228 (bottom R). Habit with inflorescences (displaced from tree).





Figures 229-232. 229, 230, *Philodendron helensiae*. —229 (top L). Cluster of inflorescences (one open) (Croat & Zhu 76738). —230 (top R). Open inflorescence with protruding spadix (Croat 61275) (Colombia, Valle: Bajo Calima). 231, 232, *P. jacquinii*. —231 (bottom L). Habit (Croat 69835). —232 (bottom R). Habit with inflorescences (Croat 12458).



Figures 233-236. *Philodendron inornatum*. —233 (top L). Habit (hand-held) (Croat 16283). —234 (top R). Habit (Croat 34592), 235, 236. (Croat 33574). —235 (bottom L). Habit in cultivation. —236 (bottom R). Open inflorescence.



Figures 237-240. —237 (top L). *Philodendron jacquini*, showing infructescences with fallen spathe (Croat 1491). —238, 239 (top R & bottom L). *P. jefense* (McPherson 10038). —240 (bottom R). *P. knappiae*, habit (Croat 67982).



Figures 241-244. 241, 242. *Philodendron jacquinzii*, —241 (top L), Blade adaxial surface (Croat & Zhu 77072), —242 (top R), Inflorescences cut open showing prolonged styles (Croat 12450), —243 (bottom L), *P. jordanianum*, habit (Croat 60089), —244 (bottom R), *P. Annapurne*, blade abaxial surface (Croat 67902).



Figures 245-248. *Philodendron joazeiroense*. —245 (top L). Cataphyll fibers (Croat 60089). —246 (top R). Blade adaxial surface (Grayum 3216). —247 (bottom L). Habit (Croat 60450). —248 (bottom R). Open inflorescence (Croat 67213).



Figures 249-252. —249 (top L). *Philodendron knappiae*, habit (Croft 67982). 250-252. *P. lasozii* Croft. 250, 251. (Croft 68953). —250 (top R). Leaf blade adaxial surface. —251 (bottom L). Stem apex with cataphyll fibers and closed inflorescences. —252 (bottom R). (Croft 69833) Stem with cluster of inflorescences and persistent intact cataphylls.



Figures 253-256. —253 (top L), *Philodendron jordanianum*, stem apex with persistent cataphyll fibers and cluster of inflorescences (Croat 67129). —254 (top R), *P. knappianum*, habit (Croat 67982). 255, 256, *P. lasorii*. —255 (bottom L), Habit (Croat 69833). —256 (bottom R), Leaf blades showing adaxial and abaxial surfaces (Croat & Zhao 77126).



Figures 257-260. 257, 258, *Philodendron lazorii*. —257 (top L). Leaf blade adaxial surfaces (Croat & Zhu 77126). —258 (top R). Stem apex with partially intact cataphyll fibers (Croat 69833). 259, 260, *P. lentii*. —259 (bottom L). Habit, displaced from tree (Croat 50041). —260 (bottom R). Stem with open inflorescence (Croat 66647).





Figures 261-264. *Philodendron lenzii*. —261 (top L). Habit (Croat 66647). 262, 263. (Croat 67538). —262 (top R). Leaves showing abaxial surfaces and D-shaped petioles. —263 (bottom L). Habit. —264 (bottom R). Showing stem with sulcate petiole base (Croat 66391).





Figures 265-268. —265 (top L), *Philodendron lantifolium*, habit (Croat 67163). 266, 267, *P. lantifolium* var. *ligulatum* (Croat 49195). —266 (top R), Habit. —267 (bottom L), Inflorescence with protruding spathix. —268 (bottom R), *P. lantifolium*, open inflorescence with protruding spathix and cut-open spathix tube (Mori 46285).



Figures 269–272. 269–271. *Philodendron ligulatum* var. *ligulatum*. —269 (top L). Habit (Croat 67318). —270 (top R). Stem with open inflorescence (Croat 34302). —271 (bottom L). Habit (plant displaced from tree) (Croat & Zhu 76954); note dark ring separating petiole from blade. —272 (bottom R). *P. ligulatum* var. *ovatum*, habit (Croat & Zhu 76754).



Figures 273-276. —273 (top L). *Philodendron ligulatum* var. *osatum*, habit (Croat & Zhu 76888). 274-276. *P. ligulatum* var. *heraclioanum* (Croat & Zhu 77098). —274 (top R). Habit showing blades spotted on lower surface. —275 (bottom L). Leaf blade adaxial surface showing sharply D-shaped petiole with narrow wing and dark ring at petiole apex. —276 (bottom R). Unopened inflorescence, stem, and D-shaped petiole.




Figures 277-280. 277, 278. *Philodendron llanense*. —277 (top L). Leaf blade adaxial surface (Croat 60505). —278 (top R). Cluster of unopened inflorescences (Croat & Zhu 769934). —279, 280 (bottom L & R). *P. madronense* (Hammel & McPherson 14526).



Figures 281-284. *Philodendron malesevichiae* (Croat 74818). —281 (top L). Habit. —282 (top R). Stem with intact, persistent cataphylls. —283 (bottom L). Petiole with conspicuous, trichome-like glands. —284 (bottom R). Open inflorescence.




Figures 285-288. *Philodendron mexicanum*. 285, 286. (Croat & Bay 65778). —285 (top L). Habit, displaced from tree. —286 (top R). Stem close up. 287, 288. Cultivated by Monroe Birdsey (photo: K. Upton). —287 (bottom L). Inflorescence. —288 (bottom R). Open spathe.



Figures 289-292. 289, 290. *Philodendron microstictum* (Cross & Hannon 79210). —289 (top L). Stem, leaf blades, inflorescence. —290 (top R). Cataphyll, anchor roots. 291, 292. *P. morii*. —291 (bottom L). (Liesner 567). —292 (bottom R). (Hammel & McPherson 14530).






Figures 293–296. *Philodendron niqueanum*. 293–295. (Croat 37942). 293, 294 (top L & R). Adult leaves. —295 (bottom L). Juvenile leaves. —296 (bottom R). Juvenile leaves in cultivation (Croat 37886) (photo: P. Malesevich).



Figures 297-300. 297-299, *Philodendron panamense* (Croft 55184). —297 (top L). Leaf blade adaxial surface. —298, 299 (top R & bottom L). Open inflorescences and sharply 2-low-ribbed cataphylls. —300 (bottom R). *P. pseudoauriculatum*, habit (Croft 37595).



Figures 301-304. *Philodendron pittrense*. 301, 302. (Croat 37944). —301 (top L). Adult leaf. —302 (top R). Stem with detached fibrous cataphyll and inflorescence. 303, 304. (Croat 68952). —303 (bottom L). Apical half of blade with loose inflorescences. —304 (bottom R). Basal half of blade.



Figures 305-308. —305 (top L), *Philodendron jarroense*, habit (Croat 68952). 306-308, *P. platypteloidatum*. —306 (top R), Habit, displaced from tree (Croat 68136). 307, 308, (Croat 68205). —307 (bottom L), Habit, displaced from tree. —308 (bottom R), Plant with opened inflorescence (Croat 78878).



Figures 309–312. *Philodendron pseudauriculatum* (Croat 33526). —309 (top L), Habit. —310 (top R), Close-up of stem with petiole bases (photo: P. Malosевич), 311, 312. *P. pterotum*. —311 (bottom L), Habit (Croat 6561). —312 (bottom R), Apex of stems with cluster of inflorescences and persistent cataphylls (Croat 10792) (cultivated at Summit Gardens).



Figures 313-316. 313, 314. *Philodendron pseudauriculatum*. —313 (top L). Habit in cultivation with leaves, inflorescence, and sharply 2-ribbed cataphylls (Croat 33526). —314 (top R). (Croat 486964). 315, 316. *P. pterotum*. —315 (bottom L). Habit (Croat 6581). —316 (bottom R). Open inflorescence (Croat 10265).



Figures 317-320. *Philodendron purpureoviride*. 317, 318. (Croat 59857). —317 (top L). Leaf blade adaxial surface. —318 (top R). Habit. —319 (bottom L). Opened inflorescence. Cultivated at Wilson Botanical Garden (photo: D. Beath). —320 (bottom R). (Croat & Zhu 76677) Habit with cataphyll recurling and beginning to recur.



Figures 321-324. 321-323. *Philodendron parulhense*. —321 (top L). Habit (Croat 41752). —322 (top R). Leaf blade adaxial surface (Croat & Hannon 63765). —323 (bottom L). Stem apex with unopened inflorescence (Croat 41189). —324 (bottom R). *P. radiatum*, cultivated at Wilson Tropical Gardens.





Figures 325-328. 325, 326, *Philodendron parvifolium* (Croat & Hannon 63765). —325 (top L). Leaf blade adaxial surface. —326 (top R). Plant habit with inflorescence cut open. *Philodendron radiatum* (Croat & Hannon 63381). —327 (bottom L). Habit. —328 (bottom R). Plant on ground with inflorescence cut open.



Figures 329-332. 329, 330. *Philodendron radiatum* var. *pseudoradiatum* (Croat & Hannon 63381). —329 (top L). Closed inflorescence. —330 (top R). Leaf blade on ground with a displaced post-anthesis spadix. 331, 332. *P. radiatum* var. *radiatum*. —331 (bottom L). Habit (Croat 32767). —332 (bottom R). Pre-adult leaf blades (Croat 66075) (photo: P. Malesevich).



Figures 333-336. 333-335. *Philodendron radiatum* var. *radiatum*. —333 (top L). (Croat 66616). —334 (top R). Cultivated at the Missouri Botanical Garden trailer house in Curandá. —335 (bottom L). Habit (one leaf displaced, showing adaxial surface) (Croat 60660). —336 (bottom R). *P. radiatum* (Croat 74931).



Figures 337-340. *Philodendron rothschukianum*. —337 (top L). Adult leaves, Tortuguero National Park (not collected; photo: M. H. Grayum). —338 (top R). (Croat & Zhu 77224) Juvenile leaves. —339 (bottom L). Habit in cultivation at the Missouri Botanical Garden (Croat 35657). —340 (bottom R). Stem with inflorescences (Croat 27704).



Figures 341-344. *Philodendron roseospathum* var. *roseospathum*. —341 (top L). Habit in cultivation showing leaves and intact cataphylls (Croat 35931). 342-344. (Croat 57515). —342 (top R). Habit. —343 (bottom L). Stem with intact cataphylls and inflorescences. —344 (bottom R). Open inflorescence.



Figures 345-348. —345 (top L), *Philodendron roseospathum*, cultivated at Selby Gardens (SEL-92-353; photo: V. Sawyer). —346 (top R), *P. sagittifolium*, habit (Croft 67697), 347, 348, *P. rubrachydium*. —347 (bottom L), Habit (Croft 74931). —348 (bottom R), Stem with cluster of inflorescences at apex of stem (Croft & Grayum 60276).



Figures 349-352. —349 (top L). *Philodendron roseospathum* var. *angustilaminatum* (Hammel 3133). 350-352. *P. sagittifolium*. —350 (top R). Leaf blade adaxial surface (Croat 67918). —351 (bottom L). Apex of stem with petiole bases, cataphylls, and inflorescences (Croat & Grayum 60259). —352 (bottom R). Open inflorescence (Croat 69731).



Figures 353–356. *Philodendron sagittifolium*. —353 (top L). Leaf blade adaxial surface (Croat 74939). —354 (top R). Plant with open inflorescence (Croat 33179). —355 (bottom L). Inflorescence with open spathe and protruding spadix (Croat 69731). —356 (bottom R). Leaf blade adaxial surface (Croat 60506).





Figures 357-360. *Philodendron scalariserve*. —357 (top L). Habit, plant displaced on ground (Croat 67203). —358 (top R). Stem apex with petiole bases and persistent cataphyll fibers (Croat 68723). —359 (bottom L). Plant with inflorescence, displaced on ground (Croat 66959). —360 (bottom R). Open inflorescence (Croat 67203).



Figures 361-364. 361-363. *Philodendron scalariuense*. —361 (top L). Habit (Croat & Zhu 76419). —362 (top R). Habit (Croat 68722). —363 (bottom L). Cluster of inflorescences (Croat 68692). —364 (bottom R). *P. amabilis* Engl., displaced leaf blade with pre-anthesis and slightly post-anthesis inflorescences (Croat 45046).



Figures 365-368. *Philodendron schottianum*. —365 (top L). Leaf blade adaxial surface (Croat 74991). 366-368. (Croat & Zhu 76518). —366 (top R). Leaf blade adaxial surfaces. —367 (bottom L). Stem apex with partially intact cataphyll fibers and a cluster of inflorescences. —368 (bottom R). Cluster of inflorescences (one partially open).



Figures 369–372. 369–371. *Philodendron schottianum*. —369 (top L). Leaf blade adaxial surface (Croat 66505). —370 (top R). Stem apex with partially decomposed cataphyll fibers and immature inflorescences (Croat 74991). —371 (bottom L). Open inflorescence emerging from persistent cataphyll fibers (Croat 35620). —372 (bottom R). *P. smithii*, blade adaxial surfaces (Croat & Hannon 64588).



Figures 373-376. —373 (top L), *Philodendron anisikii*, habit (Croat & Honzon 64524). —374 (top R), *P. squamicaulis*, habit (Croat & Zhu 76798). —375 (bottom L), *P. squamipetiolatum*, juvenile plant (Croat 74857). —376 (bottom R), *P. stramineicaule*, habit (Croat 69013).



Figures 377-380. 377, 378. *Philodendron smithii*. —377 (top L). Stem apex, displaced on ground (Croat & Hannon 64588). —378 (top R). Open inflorescence (Croat 40079). 379, 380. *P. sousae*. —379 (bottom L). (Davidse et al. 20450). —380 (bottom R). (Breedlove 25142).



Figures 381-384. *Philodendron squamicale* (Croat & Zhu 76796). —381 (top L). Leaf blade adaxial surfaces. —382, 383 (top R & bottom L). Plant with clusters of inflorescences and persistent cataphyll fibers, displaced from original site on steep road bank. —384 (bottom R). Cluster of inflorescences, one open.



Figures 385-388. *Philodendron squamipetiolatum*. 385-387. (Croft 74857). —385 (top L). Habit. —386 (top R). Leaf blade adaxial surfaces. —387 (bottom L). Stem with scaly petioles. —388 (bottom R). Stem with leaf and inflorescence (Croft 68767).






Figures 389-392. *Philodendron stramineicaule*. —389 (top L). Habit (Croat 67920). —390, 391 (top R & bottom L). (Croat 69013) Open inflorescences. —392 (bottom R). Habit (Croat 66584).



Figures 393–396. 393–395. *Philodendron strictum* (Croat 59130). —393 (top L). Leaf blade adaxial surface. —394 (top R). Plant with internodes broader than long, persistent, intact cataphylls and inflorescences, one open. —395 (bottom L). Open inflorescence. —396 (bottom R). *P. subincisum*, flowering plant in cultivation, Cornell University (Moore & Bunting 8956) (photo: G. S. Bunting).



Figures 397-400. 397, 398. *Philodendron strictum* (Croat & Zhu 76311). —397 (top L), Habit. —398 (top R), Stem apex with cataphylls with thin, intact epidermis and cluster of inflorescences. 399, 400. *P. subseriale*. —399 (bottom L), Plant showing ribbed stem (Croat & Grayum 66249). —400 (bottom R), Habit (Croat 75167).



Figures 401-404. 401, 402 (top L & R). *Philodendron subincisum* (Moore & Bunting 8956). —403 (bottom L). *P. sulcicaule*, leaf blade adaxial surface (Croat 75167). —404 (bottom R). *P. tenue*, leaf blade adaxial surface (Neill 3608).



Figures 405-408. *Philodendron tenax*. —405 (top L), Stem apex with cluster of inflorescences, one open (Vell 36608), 406, 407. (Croat & Weat 70365), —406 (top R), Habai. —407 (bottom L), Stem apex showing cataphyll fibers and unopened inflorescence. —408 (bottom R), *P. thalassiacum*, abaxial leaf blade surface (Croat 66401).



Figures 409-412. *Philodendron thalassicon*. —409 (top L). Stem apex with cataphyll fibers and clusters of inflorescences (Croat 66407). —410 (top R). Leaf blade abaxial surface (Croat 61191). 411, 412. *P. tripartitum*. —411 (bottom L). Adult leaf blade adaxial surface and juvenile blades (Croat 8605). —412 (bottom R). Adult blades (Croat 56108).




Figures 413–416. 413, 414. *Philodendron tenue* (Croat 74634) (Venezuela, Zulia: La Fria). —413 (top L). Leaf blade adaxial surface. —414 (top R). Stem apex with cataphyll fibers and inflorescences. —415 (bottom L). *P. thalassicum*, leaf (Croat 79049). —416 (bottom R). *P. tripartitum*, habit (Croat 43438).



Figures 417-420. *Philodendron tripartitum*. 417, 418. (Croat 71903) Cultivated at Waimoa Arboretum. —417 (top L). Habit. —418 (top R). Open inflorescence. —419 (bottom L). Leaf blade adaxial surfaces (Croat 70136). —420 (bottom R). Leaf blade adaxial surfaces (Croat 56149).





Figures 421–424. *Philodendron tripartitum*. —421 (top L). Cultivated at the Lyon Arboretum (#80-890), habit. —422 (top R). Leaf blade abaxial surface (Croat 70136). —423 (bottom L). Stem with open spathe (Croat 59166). —424 (bottom R). Habit (Croat 56108).



Figures 425–428. *Philodendron tysonii*. —425 (top L). Habit (Crook 68688). —426 (top R). Leaf blade adaxial surfaces (Crook 67844). —427 (bottom L). Leaf blade adaxial surfaces (Crook 66711). —428 (bottom R). Stem apex with cluster of inflorescences, one spathe open (Crook & Zhu 76346).




Figures 429–432. 429, 430. *Philodendron tysonii*. —429 (top L). Leaf blade surfaces (Croat 66589). —430 (top R). Inflorescences emerging from cataphyll fibers (Croat 67577). —431, 432 (bottom L & R). *P. utleyanum* (Wilbur et al. 15068).




Figures 433-436. 433-435. *Philodendron verapazense*. —433 (top L). (Carlson 1846). —434, 435 (top R & bottom L). (Croat 41656). —436 (bottom R). *P. ubigantupense* (Herrera 294).



Figures 437-440. *Philodendron verrucosum*. —437 (top L). Habit, Panama, Panamá: Fortuna Road (not collected). —438 (top R). Leaf showing abaxial surface and petiolar scales, Costa Rica, Heredia: Braulio Carrillo Park (not collected). —439 (bottom L). Leaf showing abaxial surface with variegated coloration (Croat 71413). —440 (bottom R). Stem with inflorescences (Croat 66558).



Figures 441-444. *Philodendron warszewiczii*. 441, 442. (Croft 40714). —441 (top L). Leaf blade adaxial surface. —442 (top R). Habit. 443, 444. (Croft 34814) Cultivated at Escuela Agrícola Panamericana, Zamorano, Honduras. —443 (bottom L). Plant with young leaves at beginning of rainy season; stem shows short, thick internodes and deciduous cataphylls. —444 (bottom R). Same plant with full-sized leaves and inflorescences in the middle of rainy season.



Figures 445-448. —445 (top L). *Philodendron terracostum*, habit (Croat 66553). 446-448. *P. sourssewiczii*. —446 (top R). Habit, cultivated by L. Garnet, Homestead, Florida. —447 (bottom L). Habit (Croat 40714). —448 (bottom R). Stem with inflorescences (Croat & Hanson 64136).



Figures 449-452. —449 (top L). *Philodendron warszewiczii*, unopened inflorescences (Croft & Hannon 64136).  
450, 451. *P. wendlandii*, Costa Rica. —450 (top R). Habit. Heredia: La Selva (not collected). —451 (bottom L). Plant  
with open inflorescences and sharply flattened petioles (Croft 34141). —452 (bottom R). *P. wilburii* var. *wilburii*, habit  
(Croft 68083).





Figures 453-456. —453 (top L), *Philodendron serrulatum*, habit (Croat 34141), 454-456, *P. wilburii* var. *wilburii*, —454 (top R), habit, cultivated at the Munich Botanical Garden. —455 (bottom L), Plant displaced from tree with open inflorescences (Croat 69083), —456 (bottom R), habit (Croat 61199).



Figures 457-460. 457, 458. *Philodendron wilburii* var. *wilburii* (Croft 68083). —457 (top L). Plant with closed inflorescence. —458 (top R). Open inflorescence. 459, 460. *P. wilburii* var. *longipedunculatum* (Croft & Zhu 77083). —459 (bottom L). Habit. —460 (bottom R). Stem displaced with closed inflorescence.



Figures 461–464. *Philodendron wilburii* var. *longipedunculatum*. —461 (top L). Blade adaxial surface (Croat 66565). —462 (top R). Habit (Croat & Grayum 59879). 463, 464. (Croat & Grayum 60372). —463 (bottom L). Habit. —464 (bottom R). Stem with inflorescences.



Figures 465-468. *Philodendron zhuianum* (Croft & Zhu 76755). —465 (top L). Habit. —466, 467 (top R & bottom L). Leaf blade adaxial surfaces. —468 (bottom R). Open inflorescence.




Figure 469. Style types of Central American *Philodendron* subg. *Philodendron*. Semidiagrammatic longitudinal views (left) with with cut-away views (right) showing probable route of styler canals (based on Mayo, 1989); stigmatic area not shown. —Mayo Style Type B. *Philodendron adena*. —Mayo Style Type C. *Philodendron warzewiczii*. —Mayo Style Type D. *Philodendron ferrugineum*. —Mayo Style Type E. *Philodendron granulare*.

- Index
- Acorus* ..... 314
- Aglaonema* Schott ..... 315, 323
- Alocasia* (Schott) G. Don ..... 325
- Ambrosia* F. Bassi ..... 314
- Annona* L. .... 370
- Anthurium* Schott ..... 312, 315, 319, 344, 345, 346, 347, 349, 350, 369, 374, 375, 377
- sect. *Pachyneurium* Croat ..... 312
- surinamense* Miq. .... 380
- Anubias* Schott ..... 323, 360
- Arisaema* Mart. .... 314
- Arisarum* Mill. .... 314
- Arum* L. .... 314
- grandifolium* Jacq. .... 380
- hederaceum* Jacq. .... 314, 458, 459, 460
- lingulatum* L. .... 314
- tripartitum* Jacq. .... 546
- Asterostigma* F. E. L. Fischer & C. A. Mey. .... 323
- Baursia* [without rank] ..... 380
- subg. *Eubaurisia* T. Post & Kuntze ..... 380
- Bognera* Nicolson ..... 323
- Bucephalandra* Schott ..... 322
- Caladium* Vent. .... 314
- baurisia* Rehb. .... 380
- fragrantissimum* Hook. .... 444
- trifoliatum* Densf. .... 546
- tripartitum* (Jacq.) Willd. .... 543
- Calla* L. .... 314
- Calostigma* Decne. [without rank] ..... 324, 331
- Carex* Schott ..... 323, 360
- Colocasia* Schott ..... 314
- Culcasia* P. Beauv. .... 323, 360
- Cyclanthus* Poit. ex A. Rich. .... 374
- hipartitus* Poit. ex A. Rich. .... 370, 374
- Cyclophala* ..... 370, 508
- amblyopsis* Bates ..... 361
- amplata* Bates ..... 374
- conspicua* Sharp ..... 361
- gravis* Bates ..... 361
- kasabi* Endrödi ..... 361
- ligyrina* Bates ..... 361
- mafaffa* Burmeister ..... 361
- melanae* Bates ..... 361
- nigerrima* Bates ..... 361, 370
- rubescens* Bates ..... 361
- rustica* (Oliver) ..... 361
- saxipunctata* Casteinau ..... 361
- Cymbopetalum* Benth. .... 370
- Dieffenbachia* Schott ..... 315, 323, 353, 360, 370, 374
- longispatha* Engl. .... 374
- Dracontium* L. .... 314, 565
- Dracunculus* Mill. .... 314
- Elopium* H. W. Schott ..... 380
- surinamense* (Miq.) Engl. .... 380
- Eriocelis* ..... 370
- colombica* Endrödi ..... 361, 587
- proba* Sharp ..... 360, 361
- Euphiodendron* [without rank] ..... 324
- Furtadoa* M. Hotta ..... 323, 360
- Gamogyne* N. E. Br. .... 322
- Gearum* N. E. Br. .... 323
- Gorgonidium* Schott ..... 323
- Gymnostachys* R. Br. .... 314
- Homalomena* Schott ex Schott & Endl. .... 312, 322, 323, 324, 360, 370
- Mangonia* Schott ..... 323
- Meconostigma* [without rank] ..... 324
- Microcasia* Becc. .... 322
- Monstera* Adans. .... 324, 343, 351, 353
- tuberculata* Lundell var. *brevinoda* (Standl. & L. O. Williams) ..... 319, 565
- Montrichardia* Crueg. .... 323, 360
- Neelia* ..... 370, 587
- Nephtytis* Schott ..... 323, 360
- Orontium* L. .... 314
- Peltandra* Raf. .... 322, 323, 360
- Philodendron* Schott ..... 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 329, 336, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 356, 357, 358, 359, 360, 361, 362, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 377, 378, 379, 380, 394, 395, 436, 459, 461, 488, 546, 565, 576, 587
- subg. *Euphiodendron* Engl. .... 317, 380
- subg. *Macrolanichium* Engl. .... 326
- subg. *Meconostigma* (Schott) Engl. .... 315, 317, 322, 324, 325, 326, 344, 345, 346, 348, 351, 358, 360, 361, 363, 365, 367, 371, 372, 373, 375, 380
- subg. *Philodendron* Schott ..... 311, 312, 314, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 328, 338, 339, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 356, 358, 359, 360, 361, 362, 363, 364, 366, 367, 368, 369, 370, 371, 372, 376, 377, 378, 379, 380, 382, 395, 576, 697
- subg. *Pteromischum* Schott ..... 312, 314, 315, 318, 319, 322, 324, 325, 326, 328, 343, 343, 344, 346, 351, 356, 358, 359, 360, 361, 363, 364, 367, 368, 369, 370, 373, 374, 375, 377, 380, 481
- sect. *Baursia* (Rehb. ex Schott) Engl. .... 317, 318, 326, 327, 328, 334, 335, 338, 341, 366, 368, 576
- sect. *Calostigma* (Schott) Pfeiffer ..... 312, 317, 324, 326, 327, 328, 330, 331, 332, 335, 336, 338, 339, 347, 351, 364, 365, 366, 369, 370, 392, 397, 401, 404, 405, 407, 408, 410, 414, 419, 420, 422, 424, 426, 429, 433, 435, 437, 438, 439, 442, 452, 453, 466, 468, 472, 475, 478, 480, 490, 491, 492, 493, 497, 500, 517, 526, 528, 533, 537, 538, 550, 551, 552, 553, 559, 561, 565, 575, 576
- sect. *Campitognium* K. Krause ..... 312, 317, 318, 326, 337, 338, 576
- sect. *Cardiobelium* (Schott) Engl. .... 351
- sect. *Fraticosa* Grayum ..... 325
- sect. *Macrogynium* Engl. .... 312, 317, 318, 326, 332, 35, 337, 338, 347, 369, 470, 575, 576
- sect. *Macrolanichium* (Schott) Engl. .... 317, 318, 326, 329, 337
- sect. *Meconostigma* (Schott) Engl. .... 316, 348, 363
- sect. *Oligocarpidium* Engl. .... 332
- sect. *Oligospermium* Engl. .... 317, 318, 324, 326, 336, 364
- sect. *Philodendron* Schott ..... 312, 315, 316, 324, 326, 328, 330, 331, 332, 333, 335, 338, 342, 347, 351, 364, 365, 366, 370, 394, 395, 402, 409, 412, 415, 417, 418, 421, 428, 431, 441, 444, 445, 447, 448, 450, 455, 456, 459, 473, 477, 485, 488, 495, 496, 502, 503, 505, 512, 522, 524, 530, 532, 535, 540, 542, 554, 575
- sect. *Philosammus* G. S. Bunting ..... 312, 327, 328, 338, 365, 381, 576
- sect. *Platyopodium* (Schott) Engl. .... 329

sect. <i>Polyspermium</i> Engl. ...	317, 318, 324, 326, 328, 330, 364
sect. <i>Polytomium</i> (Schott) Engl. ....	312, 317, 318, 326, 328, 332, 337, 338, 347, 351, 436, 507, 557, 575, 576
sect. <i>Pteromischum</i> (Schott) Engl. ....	315, 316, 317, 326, 327, 348
sect. <i>Schizophyllum</i> (Schott) Engl. ....	312, 317, 318, 326, 337, 338, 576
sect. <i>Schizoplaclium</i> (Schott) Engl. ....	317, 351
sect. <i>Tritomophyllum</i> (Schott) Engl. ....	312, 317, 318, 326, 332, 335, 336, 337, 338, 347, 351, 365, 397, 398, 423, 487, 514, 549, 575, 576
subsect. <i>Achyropodium</i> (Schott) Engl. ....	331, 338, 448, 455, 488, 530, 532, 554, 575, 576
subsect. <i>Belocardium</i> (Schott) Engl. ....	332, 334
subsect. <i>Bulaoana</i> Mayo ...	335, 336, 337, 339, 351, 575, 576
subsect. <i>Canniphyllum</i> (Schott) Mayo ...	328, 331, 334, 339, 418, 428, 512, 575, 576
subsect. <i>Cardiobelium</i> (Schott) Engl. ....	330, 394
subsect. <i>Doratophyllum</i> ...	330, 338
subsect. <i>Eubelium</i> (Schott) Engl. ....	330
subsect. <i>Eucardium</i> (Engl.) Mayo ...	336, 339, 575, 576
subsect. <i>Glossophyllum</i> (Schott) Croat ...	327, 333, 334, 335, 339, 350, 369, 397, 406, 407, 414, 422, 424, 433, 435, 442, 452, 468, 480, 491, 492, 500, 526, 533, 538, 551, 552, 559, 561, 575, 576
subsect. <i>Macrobilium</i> (Schott) Engl. ...	332, 333, 334, 338, 339, 351, 392, 401, 404, 408, 410, 420, 426, 429, 437, 438, 439, 453, 472, 475, 478, 490, 493, 516, 529, 537, 550, 553, 565, 575, 576
subsect. <i>Macrolonchium</i> (Schott) Engl. ....	328, 329, 338, 445, 575, 576
subsect. <i>Oligocarpidium</i> (Engl.) Mayo ...	335, 339, 419, 466, 575, 576
subsect. <i>Philodendron</i> Schott ...	329, 330, 331, 339, 395, 402, 409, 431, 447, 450, 455, 473, 477, 485, 495, 496, 505, 522, 524, 535, 540, 542, 575, 576
subsect. <i>Platyepilatum</i> (Schott) Engl. ....	338
subsect. <i>Platypodium</i> (Schott) Engl. ...	329, 331, 415, 421, 441, 444, 502, 575, 576
subsect. <i>Psoropodium</i> (Schott) Engl. ...	329, 330, 338, 576
subsect. <i>Solenosterigma</i> (Klotzsch ex Schott) Engl. ....	329, 330, 331, 335, 338, 412, 459, 503, 575, 576
ser. <i>Albivaccosa</i> Croat ...	331, 339, 394, 575, 576
ser. <i>Ecordata</i> Croat ...	333, 339, 369, 410, 426, 429, 478, 493, 575, 576
ser. <i>Fibrosa</i> Croat ...	331, 339, 395, 402, 409, 417, 431, 450, 473, 477, 485, 495, 496, 505, 522, 524, 540, 575, 576
ser. <i>Glossophyllum</i> Croat ...	334, 339, 397, 405, 407, 415, 422, 433, 442, 452, 468, 480, 492, 500, 551, 552, 559, 575, 576
ser. <i>Impolita</i> Croat ...	330, 333, 339, 456, 535, 542, 575, 576
ser. <i>Macrobilium</i> (Schott) Croat ...	333, 339, 392, 401, 404, 420, 437, 438, 439, 453, 475, 490, 497, 517, 529, 537, 553, 565, 576
ser. <i>Ovata</i> Croat ...	334, 335, 339, 424, 435, 491, 526, 533, 538, 561, 575, 576
ser. <i>Pachycaulia</i> Croat ...	333, 339, 408, 575, 576
ser. <i>Philodendron</i> Schott ...	330, 339, 576
ser. <i>Reticulata</i> Croat ...	333, 339, 472, 550, 575, 576
ser. <i>Velutina</i> Croat ...	330, 336, 339, 447, 575, 576
grex <i>Achyropodium</i> Schott ...	316, 331
grex <i>Bauria</i> Rehb. ex Schott ...	315, 326
grex <i>Belocardium</i> Schott ...	316, 332, 334
grex <i>Canniphyllum</i> Schott ...	315, 328
grex <i>Cardiobelium</i> Schott ...	316, 330
grex <i>Cardiophylacium</i> Schott ...	316
grex <i>Doratophyllum</i> Schott ...	316
grex <i>Eubelium</i> Schott ...	316, 330
grex <i>Glossophyllum</i> Schott ...	315, 332, 334
grex <i>Imba</i> ...	316
grex <i>Macrobilium</i> Schott ...	316, 332, 459
grex <i>Macrolonchium</i> Schott ...	316, 328
grex <i>Meconostigma</i> Schott ...	316
grex <i>Oligophlebium</i> Schott ...	316, 327
grex <i>Platypodium</i> Schott ...	316, 329
grex <i>Polytomium</i> Schott ...	316
grex <i>Psoropodium</i> Schott ...	315, 329
grex <i>Pteromischum</i> Schott ...	315, 316
grex <i>Schizophyllum</i> Schott ...	316, 337
grex <i>Solenosterigma</i> Klotzsch ...	315, 329, 459
grex <i>Sphincterostigma</i> Schott ...	315, 316
grex <i>Tritomophyllum</i> Schott ...	316, 336
Gruppe <i>Achyropodium</i> Schott ...	317
Gruppe <i>Belocardium</i> Schott ...	317, 332
Gruppe <i>Cardiobelium</i> Schott ...	317, 330
Gruppe <i>Doratophyllum</i> Schott ...	335, 336, 351
Gruppe <i>Eucardium</i> Engl. ....	336
Gruppe <i>Macrobilium</i> Schott ...	317
Gruppe <i>Oligocarpidium</i> Engl. ....	317, 351
Gruppe <i>Platypodium</i> Schott ...	317
Gruppe <i>Solenosterigma</i> (Klotzsch) Schott ...	317, 335
<b>species:</b>	
<i>accescens</i> N. W. Simmonds ...	444
<i>acceanum</i> K. Krause ...	327
<i>acrocordium</i> Schott ...	316, 332, 461
<i>acuminatissimum</i> Engl. ....	330, 335, 339
<i>acutatum</i> Schott ...	330
<i>acutifolium</i> K. Krause ...	334
<i>adhatodifolium</i> Schott ...	334
<i>advena</i> Schott ...	317, 318, 319, 320, 332, 333, 334, 339, 344, 349, 353, 357, 358, 359, 363, 366, 368, 369, 370, 371, 376, 379, 390, 391-393, 505, 517, 529, 570, 573, 577, 589, 598, 697
<i>affine</i> Hemsl. ....	397, 543
<i>albivaccosa</i> Croat ...	311, 331, 339, 353, 355, 357, 376, 386, 393-394, 570, 573, 577, 590
<i>alticola</i> Croat & Grayum ...	311, 322, 331, 339, 355, 356, 358, 360, 370, 386, 394-395, 425, 524, 534, 570, 573, 577, 591
<i>amplectans</i> A. C. Sm. ....	413
<i>andreaum</i> Devansaye ...	331, 336, 447
<i>angustialatum</i> Engl. ....	334
<i>angustilobum</i> Croat & Grayum ...	311, 322, 336, 338, 345, 350, 354, 355, 357, 358, 359, 362, 379, 396-397, 490, 570, 573, 577, 592, 598
<i>angustum</i> K. Koch ...	318, 507
<i>angustisetum</i> Engl. ....	337
<i>anisotomum</i> Schott ...	316, 317, 318, 319, 320, 336, 338, 345, 351, 352, 353, 355, 357, 358, 361, 369, 370, 379, 383, 397-400, 544, 565, 570, 573, 577, 587, 590, 593, 598
<i>annulatum</i> Croat & Grayum ...	311, 322, 333, 339, 348, 349, 350, 351, 353, 355, 357, 370, 383, 384, 386, 387, 388, 400-402, 517, 559, 570, 573, 577, 594
<i>antonioanum</i> Croat ...	311, 331, 339, 347, 348, 351,

- 353, 356, 357, 358, 359, 360, 365, 376, 402-403,  
550, 570, 573, 577, 595
- apocarpum* Matuda ..... 318, 320, 469
- arcuatum* K. Krause ..... 331, 412
- armigerum* Standl. & L. O. Williams ..... 319, 565
- aromaticum* Croat & Grayum ..... 311, 322, 333, 339,  
353, 355, 357, 358, 359, 377, 390, 402-404, 453,  
565, 570, 573, 579, 596
- asperatum* K. Koch ..... 329
- augustinum* K. Koch ..... 317
- aurantifolium* Schott ..... 315, 316
- auriculatum* Standl. & L. O. Williams ..... 318, 319,  
334, 339, 346, 348, 350, 351, 352, 355, 358, 359,  
362, 369, 377, 385, 404-406, 434, 500, 559, 570,  
573, 577, 582, 584, 585, 596, 597
- auritum* Lindl. .... 565
- bahiense* Engl. .... 327
- bakeri* Croat & Grayum ..... 311, 321, 322, 334, 339,  
349, 350, 351, 355, 356, 357, 366, 369, 376, 383,  
406-407, 452, 518, 570, 575, 577, 599
- barrosoanum* G. S. Bunting ..... 335, 336, 337, 490
- basii* Matuda ..... 318, 319, 320, 333, 339, 342, 352,  
353, 355, 356, 357, 376, 377, 387, 407-408, 570,  
573, 577, 600
- bertae* K. Krause ..... 329
- billietiae* Croat ..... 330
- bipinnatifidum* Schott ex Endl. .... 361, 371, 372, 373
- Mancheianum* Schott ..... 328
- brandianum* K. Krause ..... 329
- breedlovei* Croat ..... 311, 312, 331, 339, 357, 358,  
360, 376, 377, 388, 389, 390, 409, 437, 529, 570,  
573, 577, 599, 601
- brenesii* Standl. .... 318, 319, 320, 321, 333, 339, 351,  
355, 358, 361, 365, 369, 370, 390, 409-412, 426,  
444, 570, 573, 577, 588, 601, 602
- brevilaminatum* Schott ..... 329, 444
- brevinodum* Standl. & L. O. Williams ..... 319, 565
- brevipathum* Schott ..... 316, 317, 319, 330, 331, 337,  
338, 343, 345, 357, 361, 369, 378, 379, 388,  
412-413, 470, 570, 573, 577, 602
- subsp. *holmquistii* (G. S. Bunting) G. S. Bunting  
..... 412, 413
- breustereae* Croat ..... 311, 312, 334, 339, 349, 352,  
354, 355, 357, 359, 376, 383, 413-414, 570, 573,  
577, 413-414, 602
- brunneicaule* Croat & Grayum ..... 311, 322, 329, 338,  
345, 350, 351, 352, 353, 354, 355, 357, 358, 378,  
388, 389, 414-416, 421, 570, 573, 577, 603, 604
- buloanum* Engl. .... 335, 337, 339
- buntingianum* Croat ..... 334
- burle-marxii* G. Barroso ..... 368
- callosum* K. Krause ..... 327, 328, 361
- subsp. *ptarianum* (Steyserm.) G. S. Bunting ..... 327
- canaimae* G. S. Bunting ..... 327
- cataniapoense* G. S. Bunting ..... 335, 336
- chimantae* G. S. Bunting ..... 445
- chimboanum* Engl. .... 327
- chiriquense* Croat ..... 311, 327, 331, 339, 344, 345,  
347, 349, 353, 356, 357, 376, 386, 416-417, 421,  
570, 573, 577, 604, 606
- chiriquense* Croat & Grayum ..... 311, 312, 321, 329,  
331, 344, 349, 352, 354, 355, 357, 359, 365, 376,  
377, 383, 417-418, 491, 570, 573, 577, 605
- clematis* Wright in Griseb. .... 329, 444
- clewellii* Croat ..... 311, 335, 339, 351, 355, 357, 359,  
389, 390, 418-419, 570, 573, 577, 605
- coarlescens* Engl. .... 318, 328
- colombianum* R. E. Schult. .... 455
- coloradense* Croat ..... 311, 333, 339, 353, 376, 388,  
390, 419-420, 570, 572, 577, 605, 606
- conforme* G. S. Bunting ..... 581
- consanguineum* Schott ..... 329, 332
- copense* Croat 311, 329, 338, 344, 349, 350, 353, 354,  
355, 356, 357, 358, 359, 365, 385, 386, 415, 417,  
420-421, 496, 570, 573, 577, 607, 608
- corocoense* Kunth ..... 344
- cordatum* hort. non Vell. .... 459
- corraeae* Croat ..... 311, 334, 339, 350, 351, 354, 355,  
357, 361, 364, 367, 376, 383, 401, 421-422, 443,  
480, 570, 573, 577, 608, 609
- cotabrusense* Croat & Grayum ..... 311, 312, 322, 336,  
338, 352, 353, 355, 357, 376, 377, 382, 422-423,  
487, 514, 544, 545, 570, 573, 577, 609
- cotonense* Croat & Grayum ..... 311, 335, 339, 345,  
350, 356, 357, 367, 369, 388, 390, 423-425, 561,  
570, 573, 577, 606, 610
- craspedotromum* R. E. Schult. .... 327
- crassinervium* Lindl. .... 326, 327
- crassipathum* Croat & Grayum ..... 311, 319, 321, 322,  
333, 339, 350, 353, 355, 357, 358, 359, 363, 369,  
370, 387, 410, 414, 425-427, 570, 573, 577, 607,  
610, 611, 617
- cretanum* Croat & Grayum ..... 311, 322, 328, 331, 339,  
349, 353, 355, 360, 365, 384, 394, 427-428, 570,  
573, 577, 611, 615
- cuneatum* Engl. .... 328, 512
- cuspidatum* K. Koch & Bouché ..... 459, 460
- cyrtocoleum* F. Diels ..... 334
- daemonum* Liebm. .... 316, 317, 320, 516
- dagilla* Schott ..... 316, 397, 399
- daguense* Linden & André ..... 553
- dauidsonii* Croat ..... 318, 319, 320, 333, 339, 345,  
349, 350, 351, 353, 357, 358, 359, 360, 362, 363,  
366, 384, 385, 428-429, 582
- subsp. *dauidsonii* Croat ..... 348, 355, 429-430, 570,  
573, 577, 612, 613
- subsp. *bocatoranum* Croat ..... 311, 351, 357, 365,  
385, 429, 430, 570, 573, 577, 584, 612, 613
- deltoidum* Poepp. & Endl. .... 327
- demevarae* Gleason ..... 444
- detansayeanum* L. Linden ..... 432
- devianum* Schott ..... 332, 335, 469
- discolor* K. Krause ..... 553
- distantilobum* K. Krause ..... 337
- dodsonii* Croat & Grayum ..... 311, 322, 331, 335, 339,  
345, 347, 352, 355, 357, 359, 360, 376, 378, 379,  
386, 387, 430-432, 435, 453, 524, 570, 573, 577,  
614, 615, 617
- dolichophyllum* Croat ..... 311, 322, 334, 339, 351,  
353, 355, 357, 360, 370, 384, 432-434, 570, 573,  
577, 616, 617
- dolosum* Schott ..... 329
- dominicalense* Croat & Grayum ..... 311, 335, 339, 355,  
376, 377, 387, 570, 573, 577, 431, 434-435, 618
- donnell-smithii* Engl. .... 527
- dressleri* G. S. Bunting ..... 318, 319, 320, 321, 338,  
344, 347, 351, 353, 354, 355, 357, 365, 376, 377,  
382, 435-436, 507, 510, 557, 570, 573, 577, 617,  
619
- dunstervilleorum* G. S. Bunting ..... 327
- dwyeri* Croat ..... 311, 312, 322, 333, 339, 352, 353,  
357, 359, 364, 376, 377, 390, 436-437, 570, 573,  
577, 618
- dyscarpium* R. E. Schult. var. *dyscarpium* ..... 327



<i>ecuadorensis</i> Engl.	539	<i>harlowii</i> I. M. Johnston	318, 320, 460
<i>edenatum</i> Croat	311, 333, 339, 345, 350, 354, 357, 360, 376, 388, 437-438, 453, 454, 570, 573, 577, 618, 620	<i>hastiferum</i> Standl. & L. O. Williams	319, 565
<i>effusilobum</i> Croat	335, 336	<i>hebetatum</i> Croat	311, 330, 339, 347, 351, 352, 353, 355, 357, 359, 360, 362, 368, 378, 379, 455, 536, 542, 571, 574, 577, 588, 634, 636
<i>eichleri</i> Engl.	372	<i>hederaceum</i> (Jacq.) Schott	314, 315, 316, 317, 318, 319, 320, 329, 330, 332, 335, 338, 341, 343, 345, 353, 357, 358, 360, 363, 364, 365, 369, 370, 376, 377, 378, 379, 385, 388, 389, 458-460, 470, 504, 635
<i>elaphoglossoides</i> Schott	334	var. <i>hederaceum</i>	317, 343, 344, 378, 460, 460-463, 464, 571, 574, 577, 635
<i>elegans</i> K. Krause	337	var. <i>hederaceum</i> f. <i>micans</i>	317, 318
<i>ellipseicum</i> Engl.	327	var. <i>kirkbridei</i> Croat	311, 321, 348, 378, 388, 460, 463-464, 571, 574, 577, 635
<i>englerianum</i> Steyerf.	328	var. <i>oxycardium</i> (Schott) Croat	311, 316, 317, 318, 377, 460, 464-465, 571, 574, 578
<i>erlansonii</i> I. M. Johnston	318, 320, 469	<i>hederaceum sensu Kunth</i>	340
<i>ernestii</i> Engl.	329, 415	<i>heleniae</i> Croat	311, 335, 339, 345, 349, 350, 353, 355, 356, 357, 359, 360, 362, 370, 377, 378, 380, 384, 465-467, 478, 570, 574, 578, 636, 637
<i>crubescens</i> Linden	332	<i>heterophyllum</i> Poepp.	334
<i>fenderli</i> K. Krause	328, 337	<i>hoffmannii</i> Schott	316, 317, 319, 320, 332, 337
<i>fenalii</i> Engl.	317, 318, 543, 545	<i>hoffmannii</i> Schott sensu Engl.	318, 459, 469
<i>ferrugineum</i> Croat	311, 333, 339, 344, 350, 351, 353, 355, 359, 360, 364, 369, 370, 376, 378, 390, 438-440, 453, 486, 570, 573, 577, 621, 623, 626, 697	<i>holtonianum</i> Schott	335, 336, 546
<i>fibrillosum</i> Poepp.	328, 334, 345	<i>holmuquianii</i> G. S. Bunting	412
<i>findens</i> Croat & Grayum	311, 319, 322, 329, 338, 345, 349, 351, 352, 354, 355, 356, 357, 358, 359, 360, 369, 370, 382, 385, 440-442, 502, 524, 570, 573, 577, 581, 588, 622, 623, 624	<i>houlettiana</i> Engl.	337
<i>foliosum</i> Croat	311, 312, 334, 339, 353, 355, 357, 376, 384, 442-443, 551, 570, 573, 577, 620, 623	<i>hylanse</i> G. S. Bunting	335, 336
<i>fortunense</i> Croat	311, 329, 338, 345, 349, 350, 352, 353, 357, 389, 443-444, 571, 573, 577, 624, 625	<i>imbe</i> Schott	331
<i>fragrantissimum</i> (Hook.) G. Don	320, 328, 329, 338, 341, 344, 349, 357, 359, 360, 363, 365, 369, 376, 378, 379, 385, 444-446, 571, 573, 577, 587, 626, 627	<i>imixatum</i> Croat	311, 334, 339, 343, 345, 349, 350, 351, 352, 353, 355, 357, 360, 362, 364, 365, 378, 379, 383, 384, 407, 467-469, 481, 551, 570, 574, 638, 638
<i>fragrantissimum</i> (Hook.) Kunth	316, 317, 318	<i>impolium</i> Schott	316, 507
<i>fraternum</i> Schott	330	<i>inacuilaterum</i> Liebm.	315, 317, 328, 369
<i>fuertesii</i> K. Krause	329	<i>inconcinnum</i> Schott	364, 468
<i>giganteum</i> Schott	330, 339, 585	<i>jacquinii</i> Schott	316, 317, 318, 319, 320, 332, 335, 337, 338, 341, 343, 345, 349, 357, 363, 364, 365, 369, 370, 376, 378, 379, 388, 389, 413, 459, 460, 469-471, 571, 574, 578, 637, 639, 640
<i>gigas</i> Croat	311, 330, 331, 336, 339, 344, 346, 348, 349, 351, 352, 353, 355, 356, 357, 359, 360, 376, 386, 446-447, 571, 573, 577, 627, 628, 631	<i>jamapanum</i> G. S. Bunting	318, 320, 391
<i>glanduliferum</i> Matuda	317, 318, 319, 320, 321, 338, 342, 347, 348, 350, 351, 352, 355, 357, 371, 379, 447-449, 488, 573, 624	<i>jeffense</i>	Croat 311, 312, 333, 339, 353, 357, 366, 370, 376, 387, 471-472, 477, 571, 574, 578, 639
subsp. <i>glanduliferum</i>	331, 379, 387, 448, 571, 577, 629, 630	<i>jenmanii</i> K. Krause	330, 335
subsp. <i>camiloanum</i> Croat	448, 629	<i>jodasianum</i> G. S. Bunting	318, 319, 320, 321, 331, 339, 340, 347, 349, 350, 353, 357, 359, 360, 361, 362, 363, 378, 379, 386, 450, 472-475, 495, 542, 571, 574, 578, 640, 641, 643
<i>gloriosum</i> André	336, 444	<i>juninense</i> Engl.	328
<i>goeldii</i> G. M. Barroso	342, 351	<i>karstedianum</i> Schott	319
<i>gracile</i> Schott	315, 316, 317, 318, 320, 539	<i>knappiae</i> Croat	311, 333, 339, 342, 349, 355, 357, 359, 365, 376, 390, 411, 475, 571, 574, 578, 609, 640, 642, 643
<i>grandifolium</i> (Jacq.) Schott	315, 330, 380	<i>krebii</i> Schott	329
<i>grandipes</i> K. Krause	317, 318, 319, 329, 331, 338, 339, 342, 345, 347, 348, 349, 350, 353, 354, 357, 359, 360, 361, 363, 369, 374, 376, 378, 379, 380, 387, 449-451, 473, 571, 574, 577, 630, 631, 633	<i>lacerum</i> (Jacq.) Schott	315, 337
<i>granulare</i> Croat	311, 334, 339, 353, 355, 357, 364, 368, 369, 370, 384, 451-452, 571, 574, 577, 631, 697	<i>laciniosum</i> Schott	315
<i>graveolens</i> Engl.	327	<i>lanceigerum</i> Standl. & L. O. Williams	319, 516, 518
<i>grayumii</i> Croat	311, 322, 333, 339, 345, 351, 353, 354, 355, 356, 357, 360, 361, 370, 388, 419, 420, 431, 438, 440, 452-454, 571, 574, 577, 632, 633	<i>latilobum</i> Schott	336
<i>guatemalense</i> Engl.	331	<i>latipes</i> K. Koch & Augustin	444
<i>guatemalense</i> Engl.	317	<i>latisagittatum</i> Matuda	319, 320, 489
<i>guttiferum</i> Kunth	317, 319	<i>lauroi</i> Croat	311, 322, 331, 339, 347, 352, 353, 360, 387, 472, 476-477, 495, 571, 574, 578, 583, 642, 643, 644
<i>hammellii</i> Croat	311, 312, 331, 338, 342, 345, 350, 357, 376, 387, 454-455, 488, 532, 555, 571, 574, 577, 634	<i>leal-costae</i> Mayo & G. M. Barroso	344
		<i>lechlerianum</i> Schott	329, 498, 504
		<i>lehmannii</i> Engl.	327
		<i>lentii</i> Croat & Grayum	311, 321, 333, 339, 342,

- 349, 352, 354, 355, 457, 369, 383, 384, 466,  
477-479, 484, 493, 571, 574, 578, 645, 646
- levelii* G. S. Bunting ..... 336, 337
- liesseri* G. S. Bunting ..... 334
- ligulatum* Schott ..... 316, 317, 318, 320, 334, 339,  
345, 349, 350, 351, 355, 357, 359, 360, 362, 397,  
401, 422, 468, 479-481, 500, 559
- var. *ligulatum* ..... 354, 357, 378, 379, 380, 384,  
480, 481, 481-482, 484, 500, 571, 574, 578,  
646, 647
- var. *heracleianum* Croat ..... 311, 334, 349, 353,  
357, 364, 367, 384, 480, 481, 482, 482-483,  
500, 559, 571, 574, 578, 648
- var. *osatum* Croat ..... 311, 334, 349, 357, 360, 383,  
480, 481, 482, 483-484, 571, 574, 578, 647,  
648
- linnensis* Dilcher & Daphlian ..... 322
- lindenii* Schott ..... 332, 334, 336
- lingulatum* (L.) K. Koch ..... 315, 481
- linnaei* Kunth ..... 327, 344
- llanense* Croat ..... 311, 322, 331, 339, 349, 350, 356,  
360, 362, 370, 386, 387, 439, 484-486, 524, 571,  
574, 578, 646, 649
- longilaminatum* Schott ..... 327
- longipes* Engl. .... 327
- longipetiolatum* Engl. .... 334
- longistilum* K. Krause ..... 338
- lundellii* Bartlett ex Lundell ..... 319, 470
- madronense* Croat ..... 311, 312, 336, 338, 351, 352,  
354, 357, 376, 382, 423, 486-487, 545, 571, 574,  
578, 649
- maguirei* G. S. Bunting ..... 540
- malesovichiae* Croat ..... 311, 331, 338, 342, 345, 350,  
351, 357, 360, 378, 380, 385, 487-489, 555, 571,  
574, 578, 580, 585, 586, 650
- mamei* André ..... 336, 447
- marginatum* Urb. .... 329
- maximum* K. Krause ..... 329
- megalophyllum* Schott ..... 330, 342, 351, 369, 370,  
582, 584, 585, 586
- melanochrysum* Linden & André ..... 329
- melinoii* Brongn. ex Regel ..... 328, 337
- mexicanum* Engl. .... 317, 318, 319, 320, 333, 339,  
342, 345, 352, 354, 357, 358, 359, 360, 362, 363,  
370, 378, 379, 383, 388, 389, 397, 489-490, 571,  
574, 578, 651
- micans* K. Koch ..... 315, 316, 317, 318, 329, 459, 460
- micranthum* Poepp. ex Schott ..... 327
- microphyllum* K. Koch ..... 461
- microstictum* Standl. & L. O. Williams ..... 319, 335,  
339, 349, 351, 352, 353, 354, 355, 357, 358, 369,  
377, 383, 418, 490, 538, 571, 574, 578, 652
- midahoi* Matuda ..... 319, 320, 459, 460
- mirificum* Standl. & L. O. Williams ..... 319, 501
- monticola* Matuda ..... 319, 320, 391, 392
- morii* Croat ..... 311, 334, 339, 352, 353, 365, 370,  
377, 384, 492-493, 571, 574, 578, 652
- multispadicum* Engl. .... 335
- muricatum* Willd. ex Schott ..... 330, 335, 413
- maschlerianum* K. Krause ..... 329
- myrmecophyllum* Engl. & Schenck ..... 334
- nanegalense* Engl. .... 331
- nigraeanum* Croat ..... 311, 333, 339, 348, 353, 355,  
365, 367, 376, 377, 385, 493, 571, 574, 578, 653
- obtusilobum* Miq. .... 336
- ornatum* Schott ..... 329, 330, 338, 350, 448
- oxycardium* Schott ..... 315, 316, 317, 318, 329, 459,  
460, 464
- oxyprorum* Schott ..... 461
- pachycaule* K. Krause ..... 327
- pachyphyllum* K. Krause ..... 334
- paranense* K. Krause ..... 317, 318, 319, 320, 330,  
331, 339, 348, 351, 352, 353, 355, 357, 358, 360,  
362, 365, 376, 386, 387, 473, 477, 493-495, 526,  
571, 574, 578, 654
- panduriforme* (Kunth) Kunth ..... 327, 336
- pastazanum* K. Krause ..... 336
- paxianum* K. Krause ..... 327
- pedatum* (Hook.) Kunth ..... 337
- peraiense* G. S. Bunting ..... 327
- phlebodes* G. S. Bunting ..... 327
- pilatense* Engl. .... 331, 553
- pinichinense* G. S. Bunting ..... 327
- pinnatifidum* (Jacq.) Schott ..... 315, 328, 337
- pirrense* Croat ..... 311, 331, 339, 347, 348, 350, 353,  
354, 357, 376, 387, 495-496, 571, 574, 578, 655,  
656
- pitieri* Engl. .... 317, 318, 320, 335, 377, 461
- platypetiolatum* Madison ..... 319, 320, 333, 339, 350,  
352, 353, 355, 357, 378, 379, 388, 496, 498, 571,  
574, 578, 656
- pleistoneurum* Standl. & L. O. Williams ..... 319, 449
- polytomum* Schott ..... 316, 318, 507
- poepigii* Schott ..... 444
- populneum* K. Koch ..... 332
- pruriatum* Schott ..... 329, 459
- pseudauriculatum* Croat ..... 311, 322, 334, 339, 348,  
351, 352, 355, 358, 359, 369, 378, 380, 384, 405,  
433, 434, 483, 498-501, 552, 571, 574, 578, 588,  
654, 657, 658
- pseudoradiatum* Matuda ..... 319, 320, 510
- ptarianum* Steyererm. .... 327, 328, 361
- pterosum* K. Koch & Augustin ..... 316, 317, 318, 319,  
320, 329, 331, 338, 345, 349, 352, 357, 359, 360,  
361, 363, 376, 379, 385, 441, 501-503, 571, 574,  
578, 657, 658
- pulchrum* G. M. Barroso ..... 327
- purpureoviride* Engl. .... 317, 318, 330, 338, 345, 347,  
353, 357, 369, 370, 376, 378, 389, 459, 503-504,  
538, 571, 574, 578, 659
- purpureoviridis* ..... 317, 329
- purulhense* Croat ..... 311, 319, 320, 331, 339, 350,  
352, 353, 355, 357, 358, 359, 370, 376, 379, 387,  
392, 504-506, 517, 572, 574, 578, 660, 661
- quiense* Engl. .... 330
- radiatum* Schott ..... 316, 317, 318, 319, 320, 321,  
337, 338, 342, 349, 352, 354, 355, 357, 360, 361,  
363, 364, 367, 370, 374, 376, 377, 379, 436, 506,  
507, 537, 557, 580, 660
- var. *radiatum* ..... 349, 378, 382, 507, 507-510,  
572, 574, 578, 662, 663
- var. *pseudoradiatum* (Matuda) Croat ..... 311, 319,  
320, 337, 357, 377, 382, 436, 507, 510, 537,  
572, 574, 578, 581, 661, 662
- reichenbachianum* Schott ..... 327
- remifolium* R. E. Schult. .... 327
- robustum* Schott ..... 328
- robustum* Sodiro ined. .... 452
- roraimae* K. Krause ..... 328
- rosospathum* Croat ..... 311, 328, 331, 341, 347, 349,  
350, 353, 355, 360, 366, 370, 384, 511-512, 660
- var. *rosospathum* ..... 342, 512, 513, 572, 574, 578,  
665

- var. *angustilaminatum* Croat 311, 312, 359, 512, 513, 572, 574, 578, 667
- rothschuhianum* (Engl. & K. Krause) Croat & Grayum 336, 338, 345, 348, 351, 352, 357, 358, 359, 360, 361, 370, 379, 382, 513-516, 572, 574, 578, 580, 582, 663, 664, 666
- rubens* Schott 329, 330, 336
- ruizi* Schott 334
- sabulosum* G. S. Bunting 327
- sagittifolium* Liebm. 316, 317, 318, 319, 320, 332, 333, 342, 349, 350, 353, 353, 354, 360, 361, 362, 364, 365, 369, 370, 376, 378, 379, 390, 392, 404, 407, 516-521, 537, 545, 550, 552, 553, 565, 572, 574, 578, 666, 667, 668
- sagittifolium* Schott 316, 339, 401
- cf. *sagittifolium* 518
- sanguineum* Regel 317, 318, 320, 516, 518
- scabrum* K. Krause 330, 335
- scalarinerve* Croat & Grayum 311, 322, 331, 339, 347, 349, 350, 353, 355, 356, 358, 360, 378, 380, 384, 521-523, 572, 574, 578, 669, 670
- scandens* K. Koch & Sello 315, 316, 317, 318, 329, 377, 459, 460, 491
- f. *micans* (Schott) G. S. Bunting 460
- subsp. *oxycardium* (Schott) G. S. Bunting 464
- var. *cuspidatum* (K. Koch & Bouché) Engl. 460
- schottianum* H. Wendl. ex Schott 317, 318, 320, 330, 331, 339, 345, 347, 348, 350, 351, 352, 353, 356, 357, 358, 359, 360, 361, 385, 386, 387, 396, 435, 485, 486, 523-525, 524, 542, 572, 574, 578, 580, 584, 671, 672
- seguine* Schott 315, 317
- sellowii* K. Koch 372, 373
- serpens* Hook. f. 331, 530
- simsii* (Hook.) Don 328
- smithii* Engl. 317, 318, 320, 334, 335, 339, 345, 350, 351, 353, 359, 360, 361, 364, 368, 369, 370, 379, 389, 390, 525-528, 561, 572, 575, 578, 670, 672, 673, 674
- sodioreanum* Engl. 539
- sodiroi* hort. 336
- solimoense* A. C. Sm. 342, 346
- sonderianum* Schott 361
- sousae* Croat 311, 312, 333, 339, 352, 357, 359, 364, 366, 376, 377, 389, 409, 528-529, 572, 575, 578, 674
- speciosum* Schott 372
- sphalerum* Schott 442, 468, 492
- spligerberi* Schott 329
- squamicaule* Croat & Grayum 311, 331, 338, 342, 345, 347, 350, 352, 353, 354, 355, 357, 358, 365, 366, 378, 380, 385, 529-531, 555, 572, 575, 578, 673, 675
- squamipetalatum* Croat 311, 331, 338, 345, 350, 352, 354, 357, 365, 378, 380, 383, 389, 447, 455, 488, 531-532, 555, 572, 575, 578, 673, 676
- standleyi* Grayum 370
- steyermarkii* G. S. Bunting 327
- stramineaule* Croat 311, 322, 335, 339, 344, 345, 351, 353, 355, 358, 359, 364, 386, 390, 394, 395, 410, 526, 532-534, 561, 572, 575, 578, 581, 673, 677
- strictum* G. S. Bunting 319, 321, 330, 339, 347, 353, 355, 359, 360, 370, 378, 379, 385, 456, 457, 534-536, 542, 572, 575, 578, 678, 679
- subbasatum* Engl. & K. Krause 334
- subincisum* Schott 316, 333, 339, 345, 350, 352, 353, 360, 365, 377, 388, 536-537, 572, 575, 578, 678, 680
- subrotundum* Schott 317, 318, 332, 334, 391
- sulcicaule* Croat & Grayum 311, 321, 335, 339, 343, 345, 352, 353, 354, 355, 357, 362, 388, 389, 537-539, 572, 575, 578, 679, 680
- surinamense* (Miq. ex Schott) Engl. 368
- talamancae* Engl. 317
- tanyphyllum* Schott 316, 516
- tatei* K. Krause 327
- tenae* K. Koch & Augustin 315, 316, 317, 318, 320, 330, 331, 339, 348, 349, 350, 351, 352, 357, 360, 378, 379, 388, 539-541, 572, 575, 578, 680, 681, 683
- tenuipes* Engl. 334, 428
- thalassicum* Croat & Grayum 311, 321, 322, 330, 339, 349, 350, 351, 354, 355, 385, 386, 456, 524, 536, 540, 541-543, 572, 575, 578, 681, 682, 683
- thalifolium* Schott 329
- tobagoense* Engl. 329
- trauzii* Engl. 329
- tripartitum* (Jacq.) Schott 315, 316, 317, 318, 319, 320, 326, 336, 338, 345, 350, 352, 355, 359, 361, 362, 364, 367, 368, 378, 379, 382, 398, 399, 423, 487, 514, 543-549, 572, 575, 578, 683, 684, 685
- var. *tricuspidatum* Engl. 543
- trisetrum* Standl. 319, 320, 397
- tuxtilanum* G. S. Bunting 320, 516
- tysonii* Croat 311, 322, 333, 339, 348, 352, 353, 356, 357, 359, 361, 370, 387, 388, 402, 403, 419, 549-551, 572, 575, 578, 587, 686, 687
- ubiganapense* Croat 311, 312, 334, 339, 349, 355, 359, 376, 377, 383, 551, 572, 575, 578, 688
- uleanum* Engl. 334
- urbanianum* K. Krause 329
- utleyanum* Croat 311, 312, 334, 339, 355, 357, 367, 384, 551-552, 572, 575, 579, 687
- validinervium* Engl. 410
- verapazense* Croat 311, 312, 319, 320, 333, 339, 345, 352, 355, 357, 379, 389, 552-553, 572, 575, 579, 688
- verrucosum* L. Mathieu ex Schott 316, 317, 318, 320, 331, 338, 345, 350, 353, 357, 358, 365, 377, 378, 379, 385, 387, 447, 488, 530, 531, 553-556, 572, 575, 579, 689, 691
- victoriae* G. S. Bunting 336, 337
- viride* Engl. 334
- wallisi* Regel ex Engl. 336
- warzeniczii* K. Koch & Bouché 316, 317, 320, 321, 325, 337, 338, 342, 345, 347, 348, 349, 350, 351, 352, 353, 357, 359, 364, 365, 366, 367, 369, 370, 379, 382, 408, 436, 507, 556-558, 572, 575, 579, 583, 690, 691, 692, 697
- weberbaueri* Engl. 334
- wendlandii* Schott 315, 317, 318, 319, 320, 327, 334, 338, 339, 345, 351, 355, 358, 360, 363, 369, 377, 379, 385, 481, 500, 558-559, 572, 575, 579, 583, 692, 693
- wilburii* Croat & Grayum 311, 321, 322, 335, 339, 345, 348, 362, 369, 370, 389, 377, 388, 475, 492, 504, 526, 559-561, 562
- var. *wilburii* 319, 321, 357, 377, 561-563, 572, 575, 579, 692, 693, 694
- var. *longipedunculatum* Croat & Grayum 311, 321, 352, 353, 358, 366, 389, 525, 534, 561, 562, 563-564, 572, 575, 579, 694, 695
- williamsii* Hook. f. 372

<i>uitrianum</i> Engl. ....	334	<i>Synandropodix</i> Engl. ....	323
<i>wardackii</i> G. S. Bunting .....	334	<i>Syngonium</i> (Engl.) Croat .....	314, 315, 343, 370, 565
<i>zhuangum</i> Croat 311, 333, 339, 350, 353, 356, 357, 360, 364, 388, 390, 564-565, 572, 575, 579, 696		<i>armigerum</i> (Standl. & L. O. Williams) Croat .....	319, 565
<i>Philodendrum</i> Schott .....	315, 380, 576	<i>hastiferum</i> (Standl. & L. O. Williams) Croat .....	319, 565
<i>Piptospatha</i> N. E. Br. ....	322	<i>rothschuhianum</i> Engl. ....	513
<i>Pistia</i> L. ....	314	<i>Taccarum</i> Brongn. ex Schott .....	323
<i>Pathos</i> L. ....	314	<i>Telipodus</i> Raf. ....	380
<i>Rhodospatha</i> Poepp. ....	326, 353	<i>grandifolius</i> (Jacq.) Raf. ....	380
<i>Sauromatum venosum</i> Kunth .....	373	<i>Thaumatophyllum</i> Schott .....	380
<i>Schismatoglottis</i> Zoll. & Moritzii .....	324	<i>spruceanum</i> Schott .....	380
<i>Spathantheum</i> Schott .....	323	<i>Typhonodorum</i> Schott .....	322, 360
<i>Spathicarpa</i> Hook. ....	323	<i>Xanthosoma</i> Schott .....	370
<i>Spathiphyllum</i> Schott .....	315, 324, 353	<i>andipes</i> (K. Koch & Bouché) K. Koch .....	374
<i>Sphinctrostigma</i> [unranked] .....	324	<i>Zantedeschia</i> Spreng. ....	314, 360
<i>Stenospermatum</i> Schott .....	353		

Petioles .....	349
Blades .....	351
Blade Shape .....	351
Posterior Lobes .....	352
Sinus Shape .....	352
Blade Size .....	352
Venation .....	353
Midribs .....	353
Primary Lateral Veins .....	354
Basal Veins .....	354
Posterior Ribs .....	354
Lesser Order Veins .....	355
Morphology of Reproductive Structures .....	356
Inflorescences .....	356
Peduncles .....	357
Spathes .....	358
Spathe Color .....	359
Spadix .....	362
Male Flowers .....	363
Pollen .....	363
Female Flowers .....	363
Placentation .....	364
Ovary Locule Number .....	365
Ovules per Locule .....	365
Style and Stigma Morphology .....	366
Style Types .....	367
Fruits and Seeds .....	368
Pollination Biology .....	370
Breeding Behavior .....	374
Phenology .....	375
A. Flowering in Dry Season and Wet Season .....	376
B. Flowering only in Wet Season .....	376
C. Flowering only in Dry Season .....	376
D. Flowering All Year .....	376
E. Flowering Bimodally .....	377
Cytology .....	377
Geographical Distribution and Endemism .....	377
Taxonomic Treatment .....	380
Generic Description .....	380
Key to Subgenera of <i>Philodendron</i> .....	380
Subgeneric Description .....	380
Keys to Species of <i>Philodendron</i> Subg. <i>Philodendron</i> .....	382
Species Descriptions .....	391
Excluded Names .....	565
Literature Cited .....	566
Appendix 1: Geographic Distribution of Central American <i>Philodendron</i> Subg. <i>Philodendron</i> .....	570
Appendix 2: Technical Data on Pistils .....	573
Appendix 3: Composition of <i>Philodendron</i> Subg. <i>Philodendron</i> in Central America .....	576
Appendix 4: Phenological Patterns of Central American <i>Philodendron</i> .....	577
Figures .....	580
Index .....	698

## CONTENTS

A Revision of <i>Philodendron</i> Subgenus <i>Philodendron</i> (Araceae) for Mexico and Central America Thomas B. Croat	311
Abstract	311
Introduction	312
Materials and Methods	312
Acknowledgments	313
History of the Genus <i>Philodendron</i>	314
Heinrich Wilhelm Schott	314
Adolf Engler	316
Kurt Krause	317
Modern Work	318
Collecting History	321
Fossil History	322
Intrageneric Relationships	322
Infrageneric Relationships	324
Key to Subgenera of <i>Philodendron</i>	325
Relationships Within <i>P.</i> Subg. <i>Philodendron</i>	326
Discussion of Subgeneric Classification	326
1. <i>P.</i> Sect. <i>Baurisia</i>	326
2. <i>P.</i> Sect. <i>Philopsammis</i>	327
3. <i>P.</i> Sect. <i>Philodendron</i>	328
1. <i>P.</i> Subsect. <i>Macrolonchium</i> (328); 2. <i>P.</i> Subsect. <i>Canniphyllum</i> (328); 3. <i>P.</i> Subsect. <i>Platypodium</i> (329); 4. <i>P.</i> Subsect. <i>Psoropodium</i> (329); 5. <i>P.</i> Subsect. <i>Solenosterigma</i> (329); 6. <i>P.</i> Subsect. <i>Philodendron</i> (330); 7. <i>P.</i> Subsect. <i>Achyropodium</i> (331)	328
4. <i>P.</i> Sect. <i>Calostigma</i>	331
1. <i>P.</i> Subsect. <i>Macrobelium</i> (332); 2. <i>P.</i> Subsect. <i>Glossophyllum</i> (334); 3. <i>P.</i> Subsect. <i>Oligocarpidium</i> (335); 4. <i>P.</i> Subsect. <i>Bulaoana</i> (335); 5. <i>P.</i> Subsect. <i>Eucardium</i> (336)	332
5. <i>P.</i> Sect. <i>Tritomophyllum</i>	336
6. <i>P.</i> Sect. <i>Schizophyllum</i>	337
7. <i>P.</i> Sect. <i>Polytomium</i>	337
8. <i>P.</i> Sect. <i>Macrogynium</i>	337
9. <i>P.</i> Sect. <i>Camptogynium</i>	338
Key to Sections and Subsections of <i>P.</i> Subg. <i>Philodendron</i>	338
Morphology of Vegetative Structures	340
Anatomy	340
Vegetative Anatomy (by R. C. Keating)	340
Vascular Anatomy	341
Habit and Growth Patterns	342
Stems	344
Petiolar Scars	345
Roots	346
Cataphylls	347
Leaves	348

Contents continued on inside back cover