

NOTAE MYCOLOGICAE, SER. XXIX

Micromycetes Dakotenses et Utahenses a Doct. J. F. Brenckle Lecti et Communicati

P. A. SACCARDO

A. TELEOMYCETAE

1. ROSELLINIA AMPHISPHEROIDES Sacc. et Speg., Syll. I: 262, F. Ital. fig. 588

Hab. in ramis putrescentibus *Ribis floridani*, Kulm, N. D., Oct., 1916 (1205).

Asci $100 \times 7-8$, stip. $20-26 \mu$ longis; sporidia 1-sticha $17-21 \times 7-8$, fuliginea, senio longitrosum rimosa ut in *Anthostomate taeniosporo*. Esse potest forma, a loco sub cortice relaxato, deminuta et deformata, *R. mastoideae* Sacc.

2. *Rosellinia subsimilis* Sacc., sp. nov.

Peritheciis gregariis, subglobosis, minutis, nigris, glabris, $100-180 \mu$ diam. brevissime papillatis; ascis filiformi-paraphysatis, cylindricis, subsessilibus, $89-90 \times 5.5-6$, 8-sporis; sporidiis oblique monostichis e cylindraceo ellipsoideo-oblongis, saepius leviter inaequilateris, utrinque rotundatis, $12-14 \times 5-6$, fuligineis.

Hab. in ramis decorticates emortuis *Crataegi* sp. Whitestone Gully,¹ N. D., Nov., 1916 (1188). Affinis *R. rimincolae* differt sporidiis angustioribus subcylindraceis, etc.

3. MELANOPSAMMA POMIFORMIS (Pers.) Sacc., Syll. I: 575

Hab. in ramis putrescentibus *Corni stoloniferae*. Whitestone Gully,¹ N. D., Nov., 1916 (1185). Specimina immatura et non omnino certa.

4. DIDYMELLA EUPYRENE Sacc., Syll. I: 554

Hab. in caulibus putrescentibus *Urticae gracilis*, Ft. Douglas, Utah, Majo, 1918 (53).

¹ Whitestone Gully extends from and lies east of the U. S. Monument of "Whitestone Battlefield" in Dickey County.

5. *DIDYMELLA EFFUSA* (Niessl) Sacc., Syll. **I**: 552

Hab. in caule emortuo *Artemisiae biennis*, Kulm, N. D., Sept., 1914 (632). Forma ascis $60-70 \times 7$, sporidiis breviter fusoides, $12-15 \times 4$, plasmate bipartito 1-septatis, vix constrictis.

6. *MELANOMMA MEDIUM* Sacc. et Speg., Syll. **II**: 104

Hab. in ramis decorticatis putrescentibus *Sambuci glaucae*, Ft. Douglas, Utah, Apr., 1918 (70).

Asci $90-100 \times 14$; sporidia disticha v. oblique 1-sticha, teretibus, utrinque rotundata, $22-25 \times 7-8$, 3-septata, leviter constricta, dilute brunnea.

7. *TEICHOSPORA SOLITARIA* Ellis, Pyr. N. Amer., p. 214, *Cucurbitaria* Ell. Syll. **II**: 321.

Hab. in trunco decorticato et intemperie indurato *Atriplicis confertifoliae*, Grantville, Utah, Apr., 1918 (32).

Valde affinis est *T. oxythela* Sacc. et Briand. Syll. **IX**: 903. Berl. Jc. II, t. 74.

8. *PLEOSPORA SHEPHERDIAE* Peck., Syll. **IX**: 876

Hab. in ramis corticatis emortuis *Shepherdiae argenteae*, Whitestone Gully, N. D., Nov., 1918 (1192).

9. *HYPOXYLON EFFUSUM* Nits., Syll. **I**: 379

Hab. ad ligna putrescentia, Fort Riley, Kans., Oct., 1917 (101).

Phaeotrype Sacc., gen. nov. (Etym. quasi *Diatrype phaeospora*)

Stromata pulvinata, erumpenti superficialia, nigra, ostiolis vix exantibus punctulata; perithecia pauca monosticha, substantia discolori excepta. Asci breviter stipitati, octospori, apophysati (?). Sporidia allantoides, majuscula, atro-olivacea. Est omnino *Diatrype* sed vere phaeospora.

10. **Phaeotrype Brencklei** Sacc., sp. nov.

Stromatibus gregariis, subsuperficialibus, peridermii laciniis basi cinctis, depresso pulvinatis, atro-nitidulis, 0.5×0.7 mm.

diam., superficie levissime colliculosa et punctata; intus sordide albido, carnosulo; peritheciis 8-10 monostichis, globulosis, 0.3 mm. diam., ostiolis punctiformis, obtusis, integris; ascis fusoideis breviter stipitatis, $80-85 \times 8-9$, octosporis; sporodiis distichis allantoideis curvis, utrinque rotundatis; $16-17 \times 3.5-4$, atro-olivaceis, subnigricantibus.

Hab. in ramis corticatis emortuis *Rosae* sp., Whitestone Gully, N. Dak., Nov., 1916 (1198).

11. **Diatrype paurospora** Sacc., sp. nov.

Stromatibus gregariis, erumpenti-superficialibus, peridermo cinctis, pulvinatis, nigris, duris, fragilibus, superficie colliculosis v. rimosis, 1-2 mm. diam.; peritheciis paucis, globosis, 0.3 mm diam., substantia sorde grisea exceptis; ostiolis vix excedentibus obtusis; ascis fusoideis, apice rotundatis, $35-40 \times 7$, subsessilibus, aparaphysatis (?), 2-4-sporis; sporidiis allantoideis, curvulis, utrinque rotundatis, $16 \times 4-5$, hyalinis, vix chlorinis, tunica crassiuscula.

Hab. in ramis corticatis emortuis *Quercus utahensis* pr. Ft. Douglas, Utah, Jan., 1918 (12). Imprimis ascis tantum 2-4-sporis distinguenda species.

12. **EUTYPA SCABROSA** (Bull.) Fuck., Syll. I: 171

Hab. in ramis emortuis corticatis et decorticatis *Amelanchieris alnifoliae*, Whitestone Gully, Nov., 1916 (1199).

13. **EUTYPA LEIOPLACA** (Fr.) Cooke, Syll. I: 170

Hab. in ramis decorticatis emortuis *Amelanchieris alnifoliae*, Whitestone Gully, Nov., 1916 (1182). Socia adest forma diminuta *Calosphaeriae Principis*. Specimina *Eutypae* juvenilia et tunc dubium an potius *Eutypam latam* spectent.

14. **EUTYPELLA SARCOBATI** Ellis et Ev., Syll. XIV: 484

Hab. in ramis corticatis *Sarcobati vermiculati*, Salt Lake pr. Grantville, Utah, Apr., 1918 (29). Superficiis ligni late mycelio atrata, hinc speciis ad *Eutypam* nutat.

15. *VALSA CORNICOLA* Cooke, Syll. I: 122

Hab. in ramis corticatis emortuis *Corni stoloniferae*, Whitestone Gully, N. D., Nov., 1916 (1184). Est forma minor.

16. *VALSA CINCTA* Fr., Syll. I: 142

Hab. in ramis corticatis *Amelanchieris*?, Whitestone Gully, Nov., 1916 (1183a).

17. *VALSA LEUCOSTOMA* (Pers.) Fr., Syll. I: 139

Hab. in ramis corticatis *Pruni melanocarpi*, Ft. Douglas, Utah, Jul., 1918 (85). Specimina non bene evoluta.

18. *Chorostate utahensis* Sacc., sp. nov.

Stromatibus erumpentibus, prominulis pulvinatis, 1.5 mm. diam., nigricantibus; peritheciis numerosis, monostichis, 0.3 mm. diam., globosis, ostiis punctiformibus, vix extantibus; ascis cylindraceutis apice rotundatis, ibique non v. vix lumine bifoveolatis, breviter stipitatis, $55-60 \times 7.3-8$, aparthysatis, octosporis; sporidiis oblique monostichis, oblongo-clavatis, curvatis, $12-14 \times 3-6$, eguttulatis, hyalinis, constricto-1-septatis.

Hab. in ramis corticatis emortuis *Quercus utahensis*, Ft. Douglas, Utah, Majo, 1918 (94). Affinis *C. teiphaemae*, differt sporidiis curvis, loculo super crassiore rotundato, ascis angustioribus, ostiis abbreviatis.

19. *Diaporthe (Euporthe) Brenckleana* Sacc., sp. nov.

Peritheciis laxe gregariis, ligno immutato sed linea stromatica percurso immersis, globosis, $350-500 \mu$ diam., contextu minute celluloso atro-olivaceo; ostiis erumpentibus tereti-conicis parum extantibus; ascis fusoides, apice rotundatis, $45-55 \times 8-9$, aparthysatis, octosporis; sporidiis oblique 1-stichis, cylindraceuto-fusoides, utrinque rotundatis, 14×3.6 , inaequaliter 4-guttulatis, hyalinis, leviter constrictis.

Hab. in ramis corticatis, interdum superficie atratis, *Corni stoloniferae*, Whitestone Gully, N. D., Nov., 1916 (1186). A *D. crassicoli* ostiolo omino diverso, etc. secedit.

20. FENESTELLA MINOR Tul., Syll. II: 326

Hab. in ramis corticatis emortuis *Corni stoloniferae*, White-stone Gully, N. D., Nov., 1916 (1194). Asci 100–120 $\times$ 10–12, p. spor. 80–85 μ longa, plerumque 4-spori; sporidia 6–10-septato-muralia, 25–30 $\times$ 9–11, fuliginea; stromata valsea.

21. *Lachnum crystalligerum* Sacc., sp. nov. (mox endenda)

Hab. in ramis corticatis emortuis *Rubi parviflori*, Ft. Douglas, Utah, Jul., 1918 (81). Specimina non omnino matura.

22. PYRENOPEZIZA RUBI (Fr.) Rehm., Syll. VIII: 361

Hab. in ramis emortuis *Rubi strigosi*, Kulm, N. D., Sept., 1916 (1190).

23. ORBILIA VINOSA (A. et S.) P. Karst., Syll. VIII: 622

Hab. in lignis putridis pr. Anselm, N. D., Aug., 1916 (1072).

24. CENANGIUM FURFURACEUM (Roth.) De Not., Syll. VIII: 565

Hab. in ramis corticatis emortuis *Ceanothi velutini*, Ft. Douglas, Utah, Apr., 1918 (39). Probabiliter haec species, sed specimina imperfecta et tunc paullulum dubia.

25. *Patinella Brenckleana* Sacc., sp. nov.

Ascomatibus gregariis, subsuperficialibus, scutellatis, applanatis, 0.7–1 mm. diam., nigris, glabris, margine rectiusculo, acuto, disco plano intus flavido, epithecio vero grumoso, atro-fulvo; ascis tereti-clavatis, apice rotundatis, deorsum sensim tenuato-stipitatis, 70 $\times$ 11–12, octosporis; sporidis distichis, fusoideis, saepe leniter curvis, utrinque obtusatis, dilutissime flavidis, fartis; paraphysibus filiformibus, hyalinis.

Hab. in cortice *Amelanchieris alnifoliae*, Whitestone Gully, N. D., Nov., 1916 (1196). Affinis *P. inquinanti* a qua differt ascomate 1 mm. lato, disco plano, sporidiis paullo minoribus, paraphysibus aequalibus.

B. DEUTEROMYCETAE

26. *SPHAERONAEMA POLYMORPHUM* Auersw., Syll. **III**: 185

Hab. in ramis corticatis *Pruni melanocarpae*, Ft. Douglas, Utah, Nov., 1917 (4). Dicitur status pycnid. *Cenangellae vernicosae* (Fuck.) Sacc.

27. *DIPLODIA ABROTANI* Fuck., Syll. Syll. **III**: 369

Hab. in caulibus emortuis *Artemisiae abrotani*, cultae, Kilm., N. D., Jan., 1917 (1207). Sporulae ovato-ellipsoideae, $20-24 \times 11$, diu continuae (ut *Sphaeropsis*) denique obsolete 1-septatae, sporophora interdum in pseudoparaphyses $20-30 \times 1.5-2$ hyalinas abeunt.

28. *Septoria Lunelliana* Sacc., sp. nov.

Pycnidiis gregariis globoso-depressis, epidermide initio velatis, 180μ diam. subastomis, dein late dehiscentibus et collabescentibus, extus nigricantibus; contextu fulvo-brunneo, celluloso; sporulis filiformibus, rectis v. curvulis, utrinque acutulis, $65-75 \times 2.3-2.5$, hyalinis, obsolete 3-septatis, guttulatisque.

Hab. in foliis languidis expallentibus *Caricis atrostaticae* pr. Leeds, N. D., Jul., 1915 (J. Lunell, n. 1206). A *Septoriis* cariciolis praecipue pycnidiis contextu fulvescentibus, collabescentibus, sporulis aequalibus, obsolete 3-septatis dignoscitur.

29. *Melanconium botryosum* Sacc., sp. nov.

Acervulis erumpentibus, gregariis, nigris, peridermio laciniato cinctis, ob conidia expulsa et compacta superficie colliculoso-botryosis (sphaeriae-formibus); conidiis copiosissimis, globosis, levibus, $7-8 \mu$ diam., ochraceo-fuliginosis; conidiophoris papilliformibus, exiguis ex hypostromate celluloso, ochraeo-fusco orientibus.

Hab. in ramis corticatis emortuis *Pruni melanocarpae*, Ft. Douglas, Utah, Nov., 1917 (3). Typus singularis et vix *Melanconio* proximus, probibilter gen. nov.

30. *Steganosporium utahense* Sacc., sp. nov.

Acervulis gregariis, subcutaneo-erumpentibus, sed non praeminentibus, 1 mm. diam., primo peridermio bullato tectis, dein ab



Saccardo, P. A. 1920. "Notae mycologicae, ser. XXIX. Micromycetes Dakotenses et Utahenses a Doct. J.F. Brenckle lecti et communicati." *Mycologia* 12(4), 199–205.

View This Item Online: <https://www.biodiversitylibrary.org/item/173509>

Permalink: <https://www.biodiversitylibrary.org/partpdf/246278>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Not in copyright. The BHL knows of no copyright restrictions on this item.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.