

NOMENCLATURAL CONTRIBUTIONS TO A REVISION OF THE GENUS WISTERIA

Lawrence R. Stritch  
Department of Botany  
Southern Illinois University  
Carbondale, IL 62901

In a forthcoming revision of the genus Wisteria Nuttall, the genus will be divided into two separate but closely related genera. In order to facilitate timely annotations of specimens borrowed from various herbaria, this nomenclatural contribution is presented.

Rehsonia Stritch, gen. nov. (Fabaceae)

Perennis; rhizomata longa repentina. Caules lignei, sinistrorse aut dextre volubiles. Folia alterna, imparipinnata; foliola 7-17, petiolulata, alterna, oblongo-elliptica, cuneata ad rotundatas ad basim, acuminata ad apicem, integra, glabra ad villosas, ad 8.0 (10.0) cm longa, ad 3.0 cm lata; stipulae et stipellae pusillae, caducae.

Inflorescentia racemus terminalis, 15.0 ad 90.0 cm longus, axis pubescens. Flores pedicellati, pedicellus magnior quam 1.5 cm longus; azurei, purpurei, rosei v. candidi, ostende, magni; bracteae caducae; bracteolae absentes; calyx campanulatus, bilabiatus, pubescens, lobi 5, labium superum bilobus, labium infernum trilobus lobo medio longescenti acuminato; petala subequalia, carina incurva, petala carinae unita ad apicem, auriculata ad bases, alae oblongo-falcatae, monounguiculatae, projecturis cornuatis angulatis varie, vexillum magnum, ad 2.7 (3.0) cm longum, 2.5 (2.8) cm latum, reflexum callis binis auritis ad basim; stamina decem, diadelphi, stamen unus liber vel connatus ad tubum stamineum; ovarium super gynophorum, pubescens, discus cingens gynophorum; multiovula.

Fructus legumen; legumen oblongo-ellipticum ad oblongo-lanceolatum, complanatum, elongatum, torulosum, constrictum leviter inter semina interne, pubescens, bivalve, dehiscens tarde. Semina aliquot, complanata, lenticularia, anthracina.

Type: Rehsonia floribunda (Willd.) Stritch, comb. nov. Glycine floribunda Willd., Sp. Pl. 3,2:1066. 1819.

NEW COMBINATIONS IN THE GENUS REHSONIA

Rehsonia sinensis (Sims) Stritch, comb. nov. Glycine sinensis Sims, Bot. Mag. 46:t2083. 1819.

Rehsonia venusta (Rehd. & Wils.) Stritch, comb. nov.  
Wisteria venusta Rehd. & Wils. in Sargent, Plantae Wilsoniae 2:514. 1916.

Rehsonia villosa (Rehd.) Stritch, comb. nov. Wisteria  
villosa Rehd., J. Arnold Arb. 7:162. 1926.

Rehsonia brevidentata (Rehd.) Stritch, comb. nov. Wisteria  
brevidentata Rehd. J. Arnold Arb. 7:163. 1926.

Rehsonia brachybotrys (Sieb. & Zucc.) Stritch, comb. nov.  
Wisteria brachybotrys Sieb. & Zucc., Fl. Jap. 1:92, t.45. 1839.

Rehsonia Xformosa (Rehd.) Stritch, comb. nov. Wisteria  
Xformosa Rehd., J. Arnold Arb. 3:36. 1921.

#### A NEW SPECIES OF REHSONIA FROM JAPAN

Rehsonia nipponica Stritch, sp. nov. (Fabaceae)

Perennis; rhizomata longa repentina. Caules lignei,  
sinistrorse volubiles. Folia alterna, imparipinnata; foliola 9 -  
11 (13), petiolulata, alterna ad suboppositas, oblongo-elliptica,  
cuneata ad rotundatas ad basim, acuminata ad apicem, integra,  
glabra ad maturitatibus, ad 6.0 cm longa, ad 3.0 cm lata; stipulae  
et stipellae pusillae, caducae.

Inflorescentia racemus terminalis, 15.0 ad 22.0 cm longus,  
axis pubescens. Flores pedicellati, pedicellus 1.5 ad 2.3 cm  
longus; azurei, purpurei, v. candidi, ostende, magni; bracteae  
caducae; bracteolae absentes; calyx campanulatus, bilabiatus,  
pubescens, lobi 5, labium superum bilobus, labium infernum  
trilobus lobo medio longescenti acuminato, ad 6.5 mm longo; petala  
subequalia, carina incurva, petala carinae unita ad apicem,  
auriculata ad bases, alae oblongo-falcatae, monounguiculatae,  
projecturis cornutis perpendicularibus aut propemodo ad unquem,  
vexillum magnum, ad 2.1 - 2.2 cm longum, 2.1 - 2.2 cm latum,  
reflexum callis binis auritis ad basim, auriculae 1.7 - 2.0 mm  
longae; stamna decem, diadelphi, stamen unus liber; ovarium super  
gynophorum, pubescens, discus cingens gynophorum; multiovula.

Fructus legumen; legumen oblongo-ellipticum ad oblongo-  
lanceolatum, complanatum, elongatum, torulosum, constrictum  
leviter inter semina interne, pubescens, bivalve, dehiscens tarde.  
Semina aliquot, complanata, lenticularia, anthracina.

Type: E.H. Wilson 6579. (A). April 27, 1914. Near Tokyo.



BHL

# Biodiversity Heritage Library

Stritch, L R. 1984. "Nomenclatural contributions to a revision of the genus Wisteria." *Phytologia* 56, 183–184.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/47412>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/31749>

## Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

## Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

## Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.