

1 Drought frequency predicts life history strategies in *Heliophila*

2 J. Grey Monroe^{1,2}, Brian Gill³, Kathryn G. Turner⁴, & John K. McKay²

3 ¹ Graduate Degree Program in Ecology, Colorado State University, Fort Collins, CO 80521,
4 USA

5 ² College of Agriculture, Colorado State University, Fort Collins, CO 80521, USA

6 ³ Institute for Environment and Society, Brown University, Providence, RI 02912, USA

7 ⁴ Biology Department, Pennsylvania State University, State College, PA 16802, USA

8 Author Note

9 Correspondence concerning this article should be addressed to J. Grey Monroe, 307
10 University Ave, Fort Collins, CO 80521. E-mail: monroejg@colostate.edu

Abstract

11

12 Explaining variation in life history strategies is a long-standing goal of evolutionary biology.
13 For plants, annual and perennial life histories are thought to reflect adaptation to
14 environments that differ in the frequency of stress events such as drought. Here we test this
15 hypothesis in *Heliophila* (Brassicaceae), a diverse genus of flowering plants native to Africa,
16 by integrating 34 years of satellite-based drought measurements with 2192 herbaria
17 occurrence records. Consistent with predictions from classic life history theory, we find that
18 perennial *Heliophila* species occur in environments where droughts are significantly less
19 frequent compared to annuals. These associations are predictive while controlling for
20 phylogeny, lending support to the hypothesis that drought related natural selection has
21 influenced the distributions of these strategies. Additionally, the collection dates of annual
22 and perennial species indicate that annuals escape drought prone seasons during the seed
23 phase of their life cycle. Together, these findings provide empirical support for classic
24 hypotheses about the drivers of life history strategy in plants - that perennials out compete
25 annuals in environments with less frequent drought and that annuals are adapted to
26 environments with more frequent drought by escaping drought prone seasons as seeds.

27 *Keywords:* drought adaptation, life history evolution, remote sensing, phylogeography,
28 herbaria records

29 Drought frequency predicts life history strategies in *Heliophila*

30 **Introduction**

31 Understanding the causes and consequences of life history variation is a longstanding
32 goal of ecology and evolutionary biology (Cole, 1954). In plants, life histories are especially
33 diverse, with herbaceous species that complete their life cycle in a number of weeks to trees
34 that live for thousands of years (Brown, 1996). Along this continuum in angiosperms an
35 important division exists, distinguishing annuals which complete their seed to seed life cycle
36 within a single calendar year from perennials which can persist over multiple years. Annual
37 plants flower once, set seed, senesce, and then die, spending at least some portion of the year
38 as a seed, where they are relatively protected from environmental stress. In contrast,
39 perennial plants can continue vegetative growth after reproduction and must survive
40 conditions experienced during all seasons. These represent fundamentally different life
41 history strategies, but the ecological factors that explain their evolution and distributions
42 remain empirically unresolved (Friedman & Rubin, 2015).

43 Classical theory predicts shorter life spans in environments where adult mortality is
44 high (Charnov & Schaffer, 1973; Stearns, 1992; Franco & Silvertown, 1996). In plants, this
45 has been extended to the hypothesis that annuality is adaptive when it allows plants to
46 escape drought (Schaffer & Gadgil, 1975). Lack of water is perhaps the greatest threat to
47 survival during vegetative or reproductive growth and annuals can remain dormant (and
48 protected as a seed) during drought. Thus, environments with greater seasonal drought
49 frequency may select for annual life histories that complete reproduction prior to drought
50 prone seasons. Conversely, environments with less frequent drought may select for perennial
51 species, which benefit from multiple bouts of reproduction and competitive advantage by
52 preventing recruitment of annual species (Corbin & D'Antonio, 2004). These predictions
53 have been supported by the observation of annuals in arid environments in *Oryza perennis*
54 (Morishima *et al.*, 1984) and *Oenothera* (Evans *et al.*, 2005). Additionally, annual and

55 perennial species of *Nemesia* were qualitatively associated with winter rather and summer
56 rainfall environments respectively (Datson *et al.*, 2008) and annual species of *Scorzoneroideis*
57 were associated with environments classified as unpredictable (Cruz-Mazo *et al.*, 2009).
58 However, whether the history frequency of drought events indeed predicts the distributions
59 annual or perennial life history strategies has yet to be tested.

60 Here we combine a long-term global dataset of satellite detected drought events with
61 metadata from natural history collections to test these classic hypotheses within the African
62 endemic mustard genus, *Heliophila* L. (Brassicaceae). If annuality is an adaptive strategy
63 allowing plants to escape drought prone seasons, then drought frequency should predict the
64 distribution of life history strategies across landscapes, and annual species should be more
65 commonly associated with drought prone regions than perennial species. Furthermore, if
66 annual species have adapted to escape drought prone seasons, observations of growing annual
67 species (i.e. occurring in forms other than seed) should be rare during drought prone seasons.
68 Phylogenetic relatedness can influence tests of associations between species' traits and their
69 environments (Felsenstein, 1985; Barrett *et al.*, 1996), and therefore we assessed the
70 relationship between life history distribution and drought frequency in a phylogenetic
71 context.

72 Materials and Methods

73 Data

74 **Availability.** All analyses were performed using R. All data and the source code to
75 produce this manuscript are available at <https://github.com/greymonroe/heliophila>.
76 Software used is listed in the supplement.

77 **Satellite-detected drought data.** Remotely sensed data is a powerful tool for
78 characterizing seasonal patterns in drought because it is less limited in spatial and temporal

79 scope and resolution than weather stations or field observations (AghaKouchak *et al.*, 2015).
80 To quantify the frequency of drought during different seasons across landscapes, we used the
81 remotely sensed Vegetative Health Index (VHI), which measures landscape scale reductions
82 in plant cover and temperature conditions characteristic of drought (Kogan, 2001).
83 Generated from data collected by NOAA AVHRR satellites since 1981, the VHI combines
84 Normalized Difference Vegetation Index (NDVI) derived measures of vegetative stress
85 (Vegetative Condition Index - VCI) with temperature stress indicated by anomalies in
86 thermal spectra (Temperature Condition Index - TCI). The VHI of year y during week w of
87 [1, 52] at pixel i is derived from the following equations, where n is the number of years
88 observed.

$$VCI_{y,w,i} = 100 \frac{NDVI_{y,w,i} - NDVI_{min}}{NDVI_{max} - NDVI_{min}}$$

$$TCI_{y,w,i} = 100 \frac{T_{y,w,i} - T_{min}}{T_{max} - T_{min}}$$

$$VHI_{y,w,i} = 0.5(VCI_{y,w,i}) + 0.5(TCI_{y,w,i})$$

89 where $NDVI_{min} = \min(NDVI_{1981,w,i} \dots NDVI_{1981+n,w,i})$ and
90 $NDVI_{max} = \max(NDVI_{1981,w,i} \dots NDVI_{1981+n,w,i})$ and $T_{min} = \min(T_{1981,w,i} \dots T_{1981+n,w,i})$
91 and $T_{max} = \max(T_{1981,w,i} \dots T_{1981+n,w,i})$

92 Thus, VHI measurements are standardized according to conditions historically
93 observed at each locations. These measurements have been validated and generally used for
94 evaluating drought risk and predicting crop yields in agriculture (e.g., Rojas *et al.*, 2011;
95 Kogan *et al.*, 2016). But they also present a new tool to study seasonal patterns in the
96 frequency of drought across environments and to test hypotheses about the effect of drought
97 on ecological and evolutionary processes (Kerr & Ostrovsky, 2003). As such, the VHI has

98 been applied recently to study drought related ecology of natural species and proven useful
99 for predicting intraspecific variation in drought tolerance traits and genes (Mojica *et al.*,
100 2016; Dittberner *et al.*, 2018; Monroe *et al.*, 2018b). Here, we accessed VHI data at 16km²
101 resolution from 1981 to 2015
102 (https://www.star.nesdis.noaa.gov/smcd/emb/vci/VH/vh_ftp.php) to characterize the
103 seasonal drought frequencies experienced by annual and perennial *Heliophila* species.

104 **Life history data for *Heliophila*.** *Heliophila* is a genus of flowering plants
105 endemic to the southern portion of Africa including the Cape Floristic and Succulent Karoo
106 Regions. These are among the most botanically diverse environments on Earth and the
107 *Heliophila* species occurring there are considered to make up the most diverse genus of the
108 family Brassicaceae (Mummenhoff *et al.*, 2005; Mandáková *et al.*, 2012). This genus includes
109 both perennial and annual species and this change in life history strategy has likely arisen
110 multiple independent times (Appel & Al-Shehbaz, 1997; Mummenhoff *et al.*, 2005).
111 Furthermore, the fine scale climatic heterogeneity of Southern Africa is ideal for studying the
112 distribution of traits in relation to environmental parameters (Sayre *et al.*, 2013). We used
113 life histories reported by Mummenhoff *et al.* (2005), grouping species with annual or
114 perennial life histories. Perenniality was defined based any form of perennial life history (e.g.,
115 herbs, shrubs, mixed, etc). Because the nature of species reported with mixed traits were
116 unknown (i.e. plasticity vs. genetic variation), we classified these species here as perennial
117 since they can maintain vegetative growth after reproduction at least to some capacity.

118 ***Heliophila* occurrence records.** Botanists have collected and maintained over 350
119 million botanical specimens worldwide over the past 300 years (Thiers, 2016). Herbarium
120 specimens and their associated metadata have been used since the 1960s to study species'
121 geographical distributions (reviewed by Willis *et al.* (2017) and Lang *et al.* (2018)). And as
122 they become digitized (Soltis, 2017), these collections have been used to study relationships
123 between trait distributions, geography, and climate (Davis *et al.*, 2015; Stropp *et al.*, 2016;

124 Wolf *et al.*, 2016; Václavík *et al.*, 2017). To characterize the distributions of annual and
125 perennial *Heliophila* species, all records for the genus *Heliophila* were downloaded from the
126 Global Biodiversity Information Facility (gbif.org) on July 21, 2018 (GBIF, 2018).

127 **Sequence data for phylogeny.** An alignment of ITS I and II sequences for
128 *Heliophila* species was obtained from the authors of Mandáková *et al.* (2012). Individual ITS
129 I and II sequences for *Aethionema grandiflorum*, *Alliaria petiolata*, *Cardamine matthioli*,
130 *Chamira circaeoides*, and *Rorippa amphibia* were downloaded from Genbank.

131 **Analyses**

132 **Drought frequency calculations.** To characterize drought regimens across the
133 distributions of annual and perennial species of *Heliophila*, we calculated drought during
134 different seasons at the location of observations for *Heliophila* records using the VHI.
135 Specifically, we created global maps of the frequencies of observing drought conditions
136 (VHI<40, NOAA) during the winter (quarter surrounding winter solstice), spring (quarter
137 surrounding spring equinox), summer (quarter surrounding summer solstice) and fall (quarter
138 surrounding fall equinox) from 1981 to 2015. From these maps, the drought frequency during
139 the winter, spring, summer, and fall were extracted for the locations of all GBIF records.

140 **Filtering of occurrence records.** To avoid instances with spurious location data,
141 we filtered raw GBIF by restricting our analyses to include only:

- 142 • records for species with reported life history
- 143 • records with geospatial data
- 144 • records without known geospatial coordinate issues (i.e., coordinates reported are those
145 of herbarium)
- 146 • records from collection sites classified as land pixels in the VHI dataset
- 147 • records from Africa (to exclude locations of cultivation)

- 148 • records without duplicates (i.e., identical species, location, collection date)

149 **Phylogeny construction.** Out group (*Aethionema grandiflorum*, *Alliaria petiolata*,
150 *Cardamine matthioli*, *Chamira circaeoides*, and *Rorippa amphibia*) and ingroup *Heliophila*
151 ITS I and II sequences were aligned using MAFFT (Kato *et al.*, 2002) with strategy
152 G-INS-I, offset value 0.1, and all other options set as default. The *GTR* + Γ model of
153 nucleotide substitution was determined to best fit the data based on AIC using jModelTest2
154 (Guindon & Gascuel, 2003; Darriba *et al.*, 2012). A maximum clade credibility tree with
155 branch lengths as relative time was estimated by summarizing data from six runs of
156 100,000,000 generations of Bayesian Markov chain Monte Carlo conducted in BEAST 2
157 (Bouckaert *et al.*, 2014). Model selection and phylogenetic analyses were conducted through
158 the CIPRES Science Gateway (Miller *et al.*, 2010).

159 **Comparison of drought frequency between annual and perennial species.**

160 To evaluate the hypothesis that annual and perennial life history strategies reflect
161 adaptations to alternative drought regimes, we tested the corresponding prediction that the
162 observed distributions of annual and perennial *Heliophila* species would be significantly
163 associated with historic drought frequency. First, we compared the frequency of drought
164 during the winter, spring, summer, and fall between total occurrence records of annual and
165 perennial species by t-tests. To account for variation in the number of occurrence records per
166 species, we next calculated the mean drought frequency during the winter, spring, summer
167 and fall for each species. Because shared evolutionary history of closely related species can
168 lead to spurious associations between traits and environments (Felsenstein, 1985), we tested
169 for a relationship between life history strategy and drought frequency while controlling for
170 phylogeny using phylogenetic logistic regression (Ives & Garland, 2010).

171 **Collection dates.** To test the hypothesis that annual species have adapted to
172 escape drought prone seasons as seeds, collection dates for herbarium specimens were
173 compared between annual and perennial species. Comparisons of distributions were made by

174 Two-sample Kolmogorov-Smirnov test and Barlett variance test.

175

Results

176 Out of 8670 *Heliophila* GBIF records, 6634 were for species with reported life history
177 (Mummenhoff *et al.*, 2005), 2856 had geospatial data, 2833 did not have geospatial issues,
178 2684 were located on pixels classified as land having drought measurements, 2543 were
179 located in Africa, 2192 were not duplicated. Thus, after all filtering steps, 2192 records for
180 42 species (Figure 1, Table S1) passed for further analyses. The number of samples varied
181 between species, with a mean of 52.19 samples per species. *H. rigidiuscula* had the most
182 records, 201, and *H. cornellsbergia* the fewest, 2 (Table S1).

183 There were clear visual differences between the distributions of the 960 annual and the
184 1232 perennial *Heliophila* observation records (see Figure S1 for maps of individual species).
185 While annual species were generally found in the western regions of South Africa and
186 Namibia, primarily in the Cape Floristic Region and Succulent Karoo (Figure 2a), the
187 occurrence of perennials extended to the east coast of South Africa (Figure 2b).

188 The frequency of drought varied considerably across the ranges of *Heliophila* species
189 (Figure 2c-f). This heterogeneity is expected, given that this is one of the most climatically
190 diverse regions of the Earth (Sayre *et al.*, 2013). It is worth noting the east to west cline in
191 drought frequency observed during the summer, which distinguishes the high drought
192 frequency of the Kalahari Sands and Namid Desert phytogeographic regions from the low
193 drought frequency of the Drakensberg Mountains and Coastal Zambesian phytogeographic
194 regions. In the Cape phytogeographic region there was finer scale heterogeneity in drought
195 frequency during the summer.

196 Theory predicts that annuality should be adaptive in places where stresses such as
197 drought are more common. Conversely, perenniality should be adaptive in places where such




Figure 1. Species and examples of herbaria specimens of *Heliophila* (a) Phylogeny and life history strategies of species studied. Orange circles at branch tips mark annual species and blue circles mark perennial species. Example herbaria specimens accessed via GBIF of (a) *H. minima*, (b) *H. deserticola*, (c) *H. coronopifolia* and (d) *H. ephemera*. Images (a,c,d) courtesy of The Bavarian Natural History Collections (CC BY-SA 4.0) and (b) The London Natural History Museum (CC BY 4.0). Links to images are found in the supplement.

198 stresses are less frequent. We found that the frequency of drought was significantly higher at
199 the locations of occurrence records for annual species. When comparing across all occurrence

DROUGHT AND LIFE HISTORY

11




Figure 2. Locations of occurrence records of (a) annual and (b) perennial *Heliophila*. Drought frequency during the (c) winter, (d) spring, (e) summer and (f) fall measured using the VHI. (g) Drought frequencies during each season at the observation locations of annual and perennial *Heliophila* (t tests, ** = $p < 0.01$).

200 records (all records rather than species means, Figure 2g), the frequency of drought was
201 significantly higher at the location of annuals during the winter ($t = 10.65$, $p = 0.00$), spring
202 ($t = 10.73$, $p = 0.00$), summer ($t = 12.67$, $p = 0.00$), and fall ($t = 15.26$, $p = 0.00$). Because
203 a comparison across all occurrence records does not account for variation in the number of
204 records per species (Table S1) or species relatedness (Figure 1a), we also tested whether
205 mean drought frequency values of each species were significantly different between annuals
206 and perennials using phylogenetic logistic regression. We found that the mean drought
207 frequencies were significantly higher ($\alpha = 0.05$) in annual species during the spring, summer,
208 and fall (Table 1, Figure 3a). These findings indicate that common ancestry alone does not
209 explain differences the drought frequencies experienced between the environments of annual
210 and perennial *Heliophila*.

211 The preceding results indicate that annual species are found in environments where
212 droughts are significantly more frequent, especially in the summer and fall. Classic life
213 history theory hypothesizes that annuality reflects adaptation to such environments because
214 it allows species to escape stressful conditions. If this is the case, we would expect that
215 annuals spend the drought prone seasons of summer and fall as seeds. To test this
216 hypothesis, we compared the dates of occurrence records between annual and perennial
217 *Heliophila* species. The distributions reveal a considerable difference in the timing of
218 observation of these two life histories. In comparison to perennials, which appear to be
219 collected throughout the year, annuals are almost exclusively observed during the winter and
220 spring (Figure 3b). The differences between the distribution of collection dates were
221 significant by all tests (ks.test $D = 0.25$, $p = 0$; bartlett.test $K2 = 503.18$, $p = 0.00$) This is
222 consistent with a model of life history in which annual species flower in the spring, set seed,
223 senesce, and die before the summer. Thus, these annual species are likely to remain dormant
224 during the summer and fall, when drought is the strongest predictor of the distributions of
225 annual and perennial life histories (Figure 3a).

Table 1

*Phylogenetic logistic regressions between life history, and the mean drought frequency observed at specimen sites of *Heliophila* species the winter, spring, summer, and fall.*

Predictor	Estimate	P
Intercept	0.7231	0.6636
Winter drought freq.	-1.5452	0.7274
Intercept	5.0107	0.0534
Spring drought freq.	-12.9014	0.0464
Intercept	7.7093	0.0054
Summer drought freq.	-19.9056	0.0042
Intercept	7.0162	0.0082
Fall drought freq.	-20.8174	0.0067

Note. Annual species were scored as 0 and perennial species as 1.

226

Discussion

227 To test the hypothesis that annual and perennial plants reflect adaptation to
228 alternative drought environments we examined the landscape distribution of life history
229 strategies in the large and diverse mustard genus, *Heliophila*. Using metadata of 2192
230 occurrence records and a 34 year dataset of satellite-detected droughts, we tested the
231 prediction that annual species are more often observed in drought-prone locations than
232 perennial species, when controlling for phylogenetic relatedness. We found that drought
233 frequency is significantly different between the distributions of annual and perennial species,




Figure 3. (a) Comparison (mean \pm SE) of drought frequency across seasons measured at the GBIF records of annual and perennial species of *Heliophila*. (phylogenetic logistic regression, * = $p < 0.05$, ** = $p < 0.01$) (b) Collection dates of GBIF records of annual and perennial species of *Heliophila*.

234 with annuals being found in environments with more frequent drought, and that this signal
235 is strongest during the seasons when annuals are likely escaping via seed dormancy. These
236 results remain significant while controlling for the phylogenetic relationships of *Heliophila*
237 species, yielding support for the role that natural selection has played in driving
238 contemporary distributions of these alternatives strategies in relation to drought regimens.

239 We cannot eliminate the possibility that confounding traits or environmental variables

240 are the causative factors explaining variation in the distributions of annual and perennial
241 species. Nevertheless, these results provide quantitative support for the classic prediction
242 that annual species are found in environments that experience more frequent drought than
243 perennial species. These findings complement previous reports of qualitative associations
244 between annuality with environments characterized as having increased aridity (Evans *et al.*,
245 2005), alternative precipitation defined habitats (Morishima *et al.*, 1984; Datson *et al.*, 2008),
246 or greater unpredictability (Cruz-Mazo *et al.*, 2009). However, to our knowledge this is the
247 first study to demonstrate a significant association between life history and drought in a
248 phylogenetic context informed by large scale species distribution data and long term drought
249 measures.

250 Unfortunately, herbarium collections and their associated data do not represent
251 systematic or random sampling of a species distribution. Significant biases in collecting exist,
252 which we have not necessarily controlled for here, and may have some effect on our findings,
253 such as a bias toward collecting near roads or near the locations of natural history collections
254 (Daru *et al.*, 2018). Future research will benefit from systematic sampling efforts to avoid
255 these noted biases. However, the ecosystems of southern Africa include several biodiversity
256 hotspots and are among the most botanically well sampled regions on Earth (Daru *et al.*,
257 2018), suggesting that this may currently be the optimal region for our analyses of life
258 history distribution. Indeed, we were able to use 2192 occurrence records to study 42 species,
259 which represents a significant advance over relying on personal observations to characterize
260 species distributions.

261 These findings support classical theoretical predictions about the adaptive value of
262 annual and perennial life history strategies. Taken together, they suggest that in *Heliophila*,
263 annual species are adapted to environments with increased summer droughts by avoiding
264 these seasons in a dormant seed phase of their life cycle. They also suggest that perenniality
265 is adaptive in environments where droughts are less frequent. While most previous work has

266 focused on describing the evolutionary origins of annuality (Barrett *et al.*, 1996; Conti *et al.*,
267 1999; Andreasen & Baldwin, 2001; Verboom *et al.*, 2004; Friedman & Rubin, 2015) there are
268 at least a few other cases where perenniality appears to have arisen from an annual ancestor
269 (Bena *et al.*, 1998; Tank & Olmstead, 2008). And while early theory predicted selection for
270 annuality when adult mortality is high (Stearns, 1992), we also find evidence that perenniality
271 could be explained by reduced frequency of drought. The phylogeny reveals several
272 transitions from annual to perennial life history (Figure 1a) and the distributions of
273 perennial *Heliophila* extend into regions where drought frequency is low (Figure 2b, Figure
274 S1). Perennials may be able to out compete annual relatives in environments where the
275 infrequency of drought favors strategies that allow plants to benefit from growth over many
276 seasons. This also suggests that annuals rely on drought as a source of disturbance for
277 seedling recruitment when competing with perennials (Corbin & D'Antonio, 2004). Indeed,
278 no annual species were observed in the low drought regions of eastern South Africa (Figure 2,
279 Figure S1).

280 These findings suggest that species with locally adaptive life history strategies could be
281 threatened by rapidly changing drought regimens (Dai, 2011). This could have impacts on
282 ecosystem functioning and processes such as carbon cycling if life history traits evolve or the
283 composition of annual and perennial species changes in response (Garnier *et al.*, 1997;
284 Roumet *et al.*, 2006; Monroe *et al.*, 2018a). Furthermore, the frequency of drought may be
285 an important factor when considering the use of perennial cropping systems (Parry *et al.*,
286 2005; Lelièvre & Volaire, 2009).

287 In conclusion, we find strong support for classic life history theory which predicts that
288 annuality is adaptive in environments where droughts occur more frequently. Additionally,
289 we report evidence consistent with a life history model in annuals in which they escape
290 drought prone seasons during the seed phase of their life cycle. Finally, we find evidence that
291 the distributions of perennial lineages may indicate a competitive advantage in areas where

292 droughts are infrequent. More broadly, this work highlights the irreplaceable value of natural
293 history collections and demonstrates the power of combining such information with large
294 scale remote sensing data to address outstanding classic hypotheses in ecology and evolution.

295

Acknowledgments

296 We thank members of the Sloan lab for generous feedback that improved the quality of
297 this work. This research was supported by NSF Award 1701918 and USDA-NIFA Award
298 2014-38420-21801 to JGM.

299

Author contributions

300 JGM, BG, KGT and JKM contributed to the design of the research, interpretation,
301 and writing the manuscript. JGM, BG, and KGT contributed to the performance of the
302 research and data analysis.

303

References

- 304 AghaKouchak A, Farahmand A, Melton F, Teixeira J, Anderson M, Wardlow BD, Hain C.
305 **2015**. Remote sensing of drought: Progress, challenges and opportunities. *Reviews of*
306 *Geophysics* **53**: 452–480.
- 307 Alfaro M, Santini F, Brock C, Alamillo H, Dornburg A, Rabosky D, Carnevale G, Harmon
308 L. **2009**. Nine exceptional radiations plus high turnover explain species diversity in
309 jawed vertebrates. *Proceedings of the National Academy of Sciences of the United*
310 *States of America* **106**: 13410–13414.
- 311 Andreassen K, Baldwin BG. **2001**. Unequal evolutionary rates between annual and perennial
312 lineages of checker mallows (sidalcea, malvaceae): Evidence from 18S–26S rDNA
313 internal and external transcribed spacers. *Molecular Biology and Evolution* **18**:
314 936–944.
- 315 Appel O, Al-Shehbaz IA. **1997**. Generic limits and taxonomy of hornungia, pritzelago, and
316 hymenolobus (brassicaceae). *Novon*: 338–340.
- 317 Aust F, Barth M. **2018**. *papaja: Create APA manuscripts with R Markdown*.
- 318 Barrett SCH, Harder LD, Worley AC. **1996**. The comparative biology of pollination and
319 mating in flowering plants. *Phil. Trans. R. Soc. Lond. B* **351**: 1271–1280.
- 320 Bates D, Maechler M. **2018**. *Matrix: Sparse and dense matrix classes and methods*.
- 321 Bena G, Lejeune B, Prosperi J-M, Olivieri I. **1998**. Molecular phylogenetic approach for
322 studying life-history evolution: The ambiguous example of the genus medicago l.
323 *Proceedings of the Royal Society of London B: Biological Sciences* **265**: 1141–1151.
- 324 Bouckaert R, Heled J, Kühnert D, Vaughan T, Wu C-H, Xie D, Suchard M, Rambaut A,
325 Drummond A. **2014**. BEAST 2: A software platform for bayesian evolutionary
326 analysis. *PLoS Computational Biology* **10**: doi:10.1371/journal.pcbi.1003537.

- 327 Brown PM. **1996**. OLDLIST: A database of maximum tree ages. *Tree rings, environment,*
328 *and humanity. Radiocarbon* **1996**: 727–731.
- 329 Charnov EL, Schaffer WM. **1973**. Life-history consequences of natural selection: Cole’s
330 result revisited. *The American Naturalist* **107**: 791–793.
- 331 Cole LC. **1954**. The population consequences of life history phenomena. *The Quarterly*
332 *Review of Biology* **29**: 103–137.
- 333 Conti E, Soltis DE, Hardig TM, Schneider J. **1999**. Phylogenetic relationships of the silver
334 saxifrages (saxifraga, sect. Ligulatae haworth): Implications for the evolution of
335 substrate specificity, life histories, and biogeography. *Molecular Phylogenetics and*
336 *Evolution* **13**: 536–555.
- 337 Corbin JD, D’Antonio CM. **2004**. Competition between native perennial and exotic annual
338 grasses: Implications for an historical invasion. *Ecology* **85**: 1273–1283.
- 339 Cruz-Mazo G, Buide M, Samuel R, Narbona E. **2009**. Molecular phylogeny of
340 scorzoneroïdes (asteraceae): Evolution of heterocarpy and annual habit in
341 unpredictable environments. *Molecular phylogenetics and evolution* **53**: 835–847.
- 342 Dai A. **2011**. Drought under global warming: A review. *Wiley Interdisciplinary Reviews:*
343 *Climate Change* **2**: 45–65.
- 344 Darriba D, Taboada G, Doallo R, Posada D. **2012**. JModelTest 2: More models, new
345 heuristics and parallel computing. *Nature Methods* **9**: 772.
- 346 Daru BH, Park DS, Primack RB, Willis CG, Barrington DS, Whitfeld TJ, Seidler TG,
347 Sweeney PW, Foster DR, Ellison AM *et al.* **2018**. Widespread sampling biases in
348 herbaria revealed from large-scale digitization. *New Phytologist* **217**: 939–955.

- 349 Datson P, Murray B, Steiner K. **2008**. Climate and the evolution of annual/perennial
350 life-histories in nemesia (scrophulariaceae). *Plant Systematics and Evolution* **270**:
351 39–57.
- 352 Davis CC, Willis CG, Connolly B, Kelly C, Ellison AM. **2015**. Herbarium records are
353 reliable sources of phenological change driven by climate and provide novel insights
354 into species' phenological cueing mechanisms. *American Journal of Botany* **102**:
355 1599–1609.
- 356 Dittberner H, Korte A, Mettler-Altmann T, Weber A, Monroe G, Meaux J de. **2018**.
357 Natural variation in stomata size contributes to the local adaptation of water-use
358 efficiency in arabidopsis thaliana. *bioRxiv*: 253021.
- 359 Eastman J, Alfaro M, Joyce P, Hipp A, Harmon L. **2011**. A novel comparative method for
360 identifying shifts in the rate of character evolution on trees. *Evolution* **65**: 3578–3589.
- 361 Evans ME, Hearn DJ, Hahn WJ, Spangle JM, Venable DL. **2005**. Climate and life-history
362 evolution in evening primroses (oenothera, onagraceae): A phylogenetic comparative
363 analysis. *Evolution* **59**: 1914–1927.
- 364 Felsenstein J. **1985**. Phylogenies and the comparative method. *American Naturalist* **125**:
365 1–15.
- 366 Franco M, Silvertown J. **1996**. Life history variation in plants: An exploration of the
367 fast-slow continuum hypothesis. *Phil. Trans. R. Soc. Lond. B* **351**: 1341–1348.
- 368 Friedman J, Rubin MJ. **2015**. All in good time: Understanding annual and perennial
369 strategies in plants. *American journal of botany* **102**: 497–499.
- 370 Garnier E, Cordonnier P, Guillerm J-L, Sonié L. **1997**. Specific leaf area and leaf nitrogen
371 concentration in annual and perennial grass species growing in mediterranean

372 old-fields. *Oecologia* **111**: 490–498.

373 GBIF. **2018**. GBIF occurrence download.

374 Genz A, Bretz F. **2009**. *Computation of multivariate normal and t probabilities*. Heidelberg:
375 Springer-Verlag.

376 Guindon S, Gascuel O. **2003**. A simple, fast and accurate method to estimate large
377 phylogenies by maximum-likelihood. *Systematic Biology* **52**: 696–704.

378 Hadfield JD. **2010**. MCMC methods for multi-response generalized linear mixed models:
379 The MCMCglmm R package. *Journal of Statistical Software* **33**: 1–22.

380 Harmon L, Weir J, Brock C, Glor R, Challenger W. **2008**. GEIGER: Investigating
381 evolutionary radiations. *Bioinformatics* **24**: 129–131.

382 Heinze G, Ploner M. **2018**. *Logistf: Firth's bias-reduced logistic regression*.

383 Henry L, Wickham H. **2018**. *Purrr: Functional programming tools*.

384 Hijmans RJ. **2018**. *Raster: Geographic data analysis and modeling*.

385 Ho LST, Ane C. **2014**. A linear-time algorithm for gaussian and non-gaussian trait
386 evolution models. *Systematic Biology* **63**: 397–408.

387 Ives A, Garland T. **2010**. CPhylogenetic logistic regression for binary dependent variables.
388 *Systematic Biology* **59**: 9–26.

389 Katoh, Misawa, Kuma, Miyata. **2002**. MAFFT: A novel method for rapid multiple
390 sequence alignment based on fast fourier transform. *Nucleic Acids Research* **30**:
391 3059–3066.

- 392 Kerr JT, Ostrovsky M. **2003**. From space to species: Ecological applications for remote
393 sensing. *Trends in ecology & evolution* **18**: 299–305.
- 394 Kogan FN. **2001**. Operational space technology for global vegetation assessment. *Bulletin*
395 *of the American Meteorological Society* **82**: 1949–1964.
- 396 Kogan F, Guo W, Strashnaia A, Kleshchenko A, Chub O, Virchenko O. **2016**. Modelling and
397 prediction of crop losses from noaa polar-orbiting operational satellites. *Geomatics,*
398 *Natural Hazards and Risk* **7**: 886–900.
- 399 Lang PL, Willems FM, Scheepens J, Burbano HA, Bossdorf O. **2018**. *Using herbaria to*
400 *study global environmental change*. PeerJ Preprints.
- 401 Lelièvre F, Volaire F. **2009**. Current and potential development of perennial grasses in
402 rainfed mediterranean farming systems. *Crop Science* **49**: 2371–2378.
- 403 Mandáková T, Mummenhoff K, Al-Shehbaz IA, Mucina L, Mühlhausen A, Lysak MA.
404 **2012**. Whole-genome triplication and species radiation in the southern african tribe
405 heliophileae (brassicaceae). *Taxon* **61**: 989–1000.
- 406 Miller M, Pfeiffer W, Schwartz T and. **2010**. Creating the cipres science gateway for
407 inference of large phylogenetic trees. *Proceedings of the Gateway Computing*
408 *Environments Workshop*: 1–8.
- 409 Mojica JP, Mullen J, Lovell JT, Monroe JG, Paul JR, Oakley CG, McKay JK. **2016**.
410 Genetics of water use physiology in locally adapted *Arabidopsis thaliana*. *Plant*
411 *Science*.
- 412 Monroe J, Markman D, Beck W, Felton A, Vahsen M, Pressler Y. **2018a**. Ecoevolutionary
413 dynamics of carbon cycling in the anthropocene. *Trends in ecology & evolution* **33**:
414 213–225.

- 415 Monroe J, Powell T, Price N, Mullen J, Howard A, Evans K, Lovell J, McKay J. **2018b**.
416 Drought adaptation in nature by extensive genetic loss-of-function. *eLife*: DOI:
417 10.7554/eLife.41038.
- 418 Morishima H, Sano Y, Oka H. **1984**. Differentiation of perennial and annual types due to
419 habitat conditions in the wild rice *oryza perennis*. *Plant Systematics and Evolution*
420 **144**: 119–135.
- 421 Mummenhoff K, Al-Shehbaz IA, Bakker FT, Linder HP, Mühlhausen A. **2005**. Phylogeny,
422 morphological evolution, and speciation of endemic brassicaceae genera in the cape
423 flora of southern africa. *Annals of the Missouri Botanical Garden*: 400–424.
- 424 Müller K. **2018**. *Bindrcpp: An 'rcpp' interface to active bindings*.
- 425 Müller K, Wickham H. **2018**. *Tibble: Simple data frames*.
- 426 Orme D, Freckleton R, Thomas G, Petzoldt T, Fritz S, Isaac N, Pearse W. **2018**. *Caper:*
427 *Comparative analyses of phylogenetics and evolution in r*.
- 428 Paradis E, Schliep K. **2018**. Ape 5.0: An environment for modern phylogenetics and
429 evolutionary analyses in R. *Bioinformatics* **xx**: xxx–xxx.
- 430 Parry M, Flexas J, Medrano H. **2005**. Prospects for crop production under drought:
431 Research priorities and future directions. *Annals of Applied Biology* **147**: 211–226.
- 432 Pebesma EJ, Bivand RS. **2005**. Classes and methods for spatial data in R. *R News* **5**: 9–13.
- 433 Plummer M, Best N, Cowles K, Vines K. **2006**. CODA: Convergence diagnosis and output
434 analysis for mcmc. *R News* **6**: 7–11.
- 435 R by Thomas Lumley VJCP to, author. BRN that maintainers are not available to give
436 advice on using a package they did not. **2015**. *Gee: Generalized estimation equation*

437 *solver*.

438 R Core Team. **2018**. *R: A language and environment for statistical computing*. Vienna,
439 Austria: R Foundation for Statistical Computing.

440 Revell LJ. **2012**. Phytools: An r package for phylogenetic comparative biology (and other
441 things). *Methods in Ecology and Evolution* **3**: 217–223.

442 Richard A. Becker OS code by, Ray Brownrigg. Enhancements by Thomas P Minka ARWR
443 version by, Deckmyn. A. **2018**. *Maps: Draw geographical maps*.

444 Rojas O, Vrieling A, Rembold F. **2011**. Assessing drought probability for agricultural areas
445 in africa with coarse resolution remote sensing imagery. *Remote sensing of*
446 *Environment* **115**: 343–352.

447 Roumet C, Urcelay C, Di'az S. **2006**. Suites of root traits differ between annual and
448 perennial species growing in the field. *New phytologist* **170**: 357–368.

449 Sayre RG, Comer P, Hak J, Josse C, Bow J, Warner H, Larwanou M, Kelbessa E, Bekele
450 T, Kehl H *et al.* **2013**. A new map of standardized terrestrial ecosystems of africa.
451 *African Geographical Review*.

452 Schaffer W, Gadgil M. **1975**. Selection for optimal life histories in plants. *Ecology and*
453 *evolution of communities*.: 142–157.

454 Slater G, Harmon L, Wegmann D, Joyce P, Revell L, Alfaro M. **2012**. Fitting models of
455 continuous trait evolution to incompletely sampled comparative data using
456 approximate bayesian computation. *Evolution* **66**: 752–762.

457 Soetaert K. **2017**. *Diagram: Functions for visualising simple graphs (networks), plotting*
458 *flow diagrams*.

- 459 Soetaert K. **2018**. *Shape: Functions for plotting graphical shapes, colors*.
- 460 Soltis PS. **2017**. Digitization of herbaria enables novel research. *American journal of botany*
461 **104**: 1281–1284.
- 462 Stearns SC. **1992**. *The evolution of life histories*.
- 463 Stropp J, Ladle RJ, M. Malhado AC, Hortal J, Gaffuri J, H. Temperley W, Olav Skøien J,
464 Mayaux P. **2016**. Mapping ignorance: 300 years of collecting flowering plants in
465 africa. *Global Ecology and Biogeography* **25**: 1085–1096.
- 466 Tank DC, Olmstead RG. **2008**. From annuals to perennials: Phylogeny of subtribe
467 castillejinae (orobanchaceae). *American Journal of Botany* **95**: 608–625.
- 468 Thiers B. **2016**. Index herbariorum: A global directory of public herbaria and associated
469 staff. New York Botanical Garden's virtual herbarium. <http://sweetgum.nybg.org/ih>.
- 470 Václavík T, Beckmann M, Cord AF, Bindewald AM. **2017**. Effects of uv-b radiation on
471 leaf hair traits of invasive plants—combining historical herbarium records with novel
472 remote sensing data. *PloS one* **12**: e0175671.
- 473 Venables WN, Ripley BD. **2002**. *Modern applied statistics with s*. New York: Springer.
- 474 Verboom GA, Linder HP, Stock WD. **2004**. Testing the adaptive nature of radiation:
475 Growth form and life history divergence in the African grass genus *Ehrharta* (Poaceae:
476 Ehrhartoideae). *American Journal of Botany* **91**: 1364–1370.
- 477 Wickham H. **2016**. *Ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York.
- 478 Wickham H. **2017**. *Tidyverse: Easily install and load the 'tidyverse'*.
- 479 Wickham H. **2018a**. *Forcats: Tools for working with categorical variables (factors)*.

480 Wickham H. **2018b**. *Stringr: Simple, consistent wrappers for common string operations*.

481 Wickham H, François R, Henry L, Müller K. **2018**. *Dplyr: A grammar of data*
482 *manipulation*.

483 Wickham H, Henry L. **2018**. *Tidyr: Easily tidy data with 'spread()' and 'gather()' functions*.

484 Wickham H, Hester J, François R. **2017**. *Readr: Read rectangular text data*.

485 Willis CG, Ellwood ER, Primack RB, Davis CC, Pearson KD, Gallinat AS, Yost JM,
486 Nelson G, Mazer SJ, Rossington NL *et al.* **2017**. Old plants, new tricks:
487 Phenological research using herbarium specimens. *Trends in ecology & evolution* **32**:
488 531–546.

489 Wolf A, Zimmerman NB, Anderegg WR, Busby PE, Christensen J. **2016**. Altitudinal shifts
490 of the native and introduced flora of California in the context of 20th-century
491 warming. *Global ecology and biogeography* **25**: 418–429.

Supplement

492

493 **Images used.** <https://www.gbif.org/occurrence/1099023487>
494 <https://www.gbif.org/occurrence/1057389408> <https://www.gbif.org/occurrence/1099023562>
495 <https://www.gbif.org/occurrence/1099023490>

496 **Software used.** We used R (Version 3.5.1; R Core Team, 2018) and the R-packages
497 *ape* (Version 5.2; Paradis & Schliep, 2018; Orme *et al.*, 2018; Soetaert, 2018), *bindrcpp*
498 (Version 0.2.2; Müller, 2018), *caper* (Version 1.0.1; Orme *et al.*, 2018), *coda* (Version 0.19.2;
499 Plummer *et al.*, 2006), *diagram* (Version 1.6.4; Soetaert, 2017), *dplyr* (Version 0.7.8;
500 Wickham *et al.*, 2018), *forcats* (Version 0.3.0; Wickham, 2018a), *gee* (Version 4.13.19; R by
501 Thomas Lumley & author., 2015), *geiger* (Version 2.0.6; Alfaro *et al.*, 2009; Harmon *et al.*,
502 2008; Eastman *et al.*, 2011; Slater *et al.*, 2012), *ggplot2* (Version 3.1.0; Wickham, 2016),
503 *logistf* (Version 1.23; Heinze & Ploner, 2018), *maps* (Version 3.3.0; Richard A. Becker *et al.*,
504 2018), *MASS* (Version 7.3.51.1; Venables & Ripley, 2002), *Matrix* (Version 1.2.15; Bates &
505 Maechler, 2018), *MCMCglmm* (Version 2.26; Hadfield, 2010), *mvtnorm* (Version 1.0.8; Genz
506 & Bretz, 2009), *papaja* (Version 0.1.0.9842; Aust & Barth, 2018), *phylolm* (Version 2.6; Ho &
507 Ane, 2014), *phytools* (Version 0.6.60; Revell, 2012), *purrr* (Version 0.2.5; Henry & Wickham,
508 2018), *raster* (Version 2.8.4; Hijmans, 2018), *readr* (Version 1.2.1; Wickham *et al.*, 2017),
509 *shape* (Version 1.4.4; Soetaert, 2018), *sp* (Version 1.3.1; Pebesma & Bivand, 2005), *stringr*
510 (Version 1.3.1; Wickham, 2018b), *tibble* (Version 1.4.2; Müller & Wickham, 2018), *tidyr*
511 (Version 0.8.2; Wickham & Henry, 2018), and *tidyverse* (Version 1.2.1; Wickham, 2017) for
512 all our analyses.

513

Supplementary tables and figures.

Table S1

Heliophila species records and the mean drought frequencies during different seasons at the location of records

Species	LH	n	Winter	Spring	Summer	Fall
<i>Heliophila acuminata</i>	a	28	0.32	0.38	0.41	0.36
<i>Heliophila africana</i>	a	91	0.33	0.35	0.34	0.34
<i>Heliophila amplexicaulis</i>	a	60	0.32	0.36	0.39	0.33
<i>Heliophila arenaria</i>	a	65	0.34	0.37	0.38	0.34
<i>Heliophila carnososa</i>	p	129	0.33	0.37	0.39	0.31
<i>Heliophila cedarbergensis</i>	p	3	0.40	0.43	0.32	0.27
<i>Heliophila collina</i>	a	16	0.35	0.47	0.48	0.45
<i>Heliophila cornellsbergia</i>	a	2	0.33	0.42	0.35	0.21
<i>Heliophila cornuta</i>	p	101	0.35	0.40	0.40	0.34
<i>Heliophila coronopifolia</i>	a	40	0.37	0.42	0.40	0.37
<i>Heliophila crithmifolia</i>	a	97	0.35	0.42	0.45	0.38
<i>Heliophila descurva</i>	a	12	0.36	0.38	0.38	0.29
<i>Heliophila deserticola</i>	a	133	0.48	0.48	0.46	0.45
<i>Heliophila digitata</i>	a	30	0.33	0.38	0.44	0.38
<i>Heliophila dregeana</i>	p	17	0.33	0.37	0.33	0.32
<i>Heliophila elongata</i>	p	82	0.26	0.32	0.30	0.25
<i>Heliophila ephemera</i>	a	3	0.14	0.27	0.31	0.26
<i>Heliophila esterhuyseniae</i>	p	3	0.21	0.30	0.37	0.27
<i>Heliophila eximia</i>	p	12	0.42	0.41	0.32	0.34
<i>Heliophila gariepina</i>	a	12	0.50	0.53	0.48	0.41
<i>Heliophila glauca</i>	p	35	0.29	0.35	0.34	0.33
<i>Heliophila juncea</i>	p	150	0.32	0.37	0.39	0.35
<i>Heliophila linearis</i>	p	94	0.32	0.33	0.28	0.30

DROUGHT AND LIFE HISTORY

29

<i>Heliophila macowaniana</i>	a	31	0.33	0.38	0.44	0.39
<i>Heliophila macra</i>	p	22	0.30	0.30	0.32	0.29
<i>Heliophila macrosperma</i>	p	5	0.28	0.36	0.35	0.25
<i>Heliophila minima</i>	p	35	0.36	0.45	0.51	0.39
<i>Heliophila namaquana</i>	a	16	0.39	0.46	0.48	0.39
<i>Heliophila nubigena</i>	p	19	0.31	0.36	0.43	0.38
<i>Heliophila pectinata</i>	a	16	0.27	0.34	0.50	0.34
<i>Heliophila polygaloides</i>	p	12	0.40	0.48	0.42	0.34
<i>Heliophila pubescens</i>	a	9	0.31	0.40	0.48	0.39
<i>Heliophila pusilla</i>	a	45	0.32	0.38	0.38	0.34
<i>Heliophila rigidiuscula</i>	p	201	0.30	0.33	0.28	0.24
<i>Heliophila scoparia</i>	p	106	0.31	0.37	0.36	0.31
<i>Heliophila seselifolia</i>	a	80	0.36	0.42	0.45	0.40
<i>Heliophila suavissima</i>	p	92	0.30	0.39	0.42	0.31
<i>Heliophila subulata</i>	p	103	0.29	0.33	0.31	0.29
<i>Heliophila tricuspidata</i>	p	8	0.28	0.33	0.38	0.30
<i>Heliophila trifurca</i>	a	77	0.45	0.48	0.48	0.43
<i>Heliophila tulbaghensis</i>	p	3	0.36	0.41	0.36	0.35
<i>Heliophila variabilis</i>	a	97	0.35	0.41	0.40	0.37

Note. LH = Life history (a = annual, p = perennial). n=sample size of GBIF records. Seasons are mean drought frequencies observed at locations of records.




Figure S1. Maps of occurrence records for individual species. Orange points indicate annual species. Blue points indicate perennial species.