



Review article

Global species diversity and distribution of the psychedelic fungal genus *Panaeolus*

Dominique Strauss, Soumya Ghosh, Zurika Murray, Marieka Gryzenhout^{*}

Department of Genetics, Natural and Agricultural Sciences, University of the Free State, Bloemfontein, South Africa

ARTICLE INFO

Keywords:

DNA sequence
Geography
Hallucinogenic
Panaeolus
Psilocin
Psilocybin
Taxonomy

ABSTRACT

Psychedelic fungi have received considerable attention recently due to their promising treatment potential of several psychiatric disorders and medical conditions, both in clinical settings but also as a nutraceutical. Besides research, a growing number of companies are developing capacity to conduct research and clinical trials where these fungi and their products can be used, and to provide these fungi to the public market that are rapidly becoming legal across the world. Whereas *Psilocybe* species are better known as psychedelic fungi, species in *Panaeolus* are also reputed to contain the psychedelic compound psilocybin and used recreationally. For the novice, there is no contemporary scientific summary of all the species in this genus that are known to be psychedelic, compared to those that are not. The global distribution and species diversity of these brown to white, often inconspicuous mushrooms are also not summarised, nor is it known to what extent DNA sequence data that are needed for identification have been generated for all of the species in this genus. However, psychedelic *Panaeolus* species are used and moved across the world. This lack of data makes it difficult to regulate bioexploitation and apply law enforcement of these fungi and the compounds they contain, especially seen in the light of the rapid development of the related markets. The aim of this review is to summarise current scientific data and knowledge on the species biodiversity, geographical distribution, extent of sequence data for identification purposes, and the psychedelic potential of species, based on published results. The review revealed where species are mostly known from, while also indicating areas seriously lacking such biodiversity data. A significant degree of study across the world is still needed to confirm which of these species are truly psychedelic and exactly what compounds they can produce.

1. Introduction

The use of psychedelic fungi has its roots in the histories of various cultures across the world [1]. Of these, psilocybin-containing mushrooms have mostly been used in the Americas by shamans, and were consequently used recreationally by Western societies after their discovery [2–4]. Psilocybin and its related metabolite psilocin are two serotonin derivatives that are increasingly demanded in modern medicine [5–11]. When psilocybin and psilocin are consumed by an individual, these substances can cross the blood-brain barrier and act as a serotonin inhibitor, creating hallucinations and long lasting beneficial changes in the brain [12,13]. Psilocybin and psilocin are used as an alternative therapeutic tool for the treatment of major depression disorder, obsessive-compulsive disorder,

^{*} Corresponding author.

E-mail address: GryzenhoutM@ufs.ac.za (M. Gryzenhout).

<https://doi.org/10.1016/j.heliyon.2023.e16338>

Received 6 September 2022; Received in revised form 9 May 2023; Accepted 12 May 2023

Available online 18 May 2023

2405-8440/© 2024 Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

anorexia nervosa, post-traumatic stress disorder, cluster headaches, chronic pain, and substance abuse (e.g. alcohol, cocaine, nicotine) [14–18].

Upon legalization in several countries research into medicinal aspects are exploding with small to large pharmaceutical companies starting to invest significant resources to harness the therapeutic powers of psilocybin analogues and the fungi that produce them [19–21]. Whereas these compounds can be synthesized, it is the organic nature of consuming the mushrooms that naturally produce these compounds, which appeal most to the public. Furthermore, these mushrooms are also a source of self help, or assisted help, for individuals that does not require prescriptions. Several fungal genera are known to include psychedelic species that can produce psilocybin, with *Psilocybe* being the best known [4,22].

A number of powerful psilocybin-producing species are also known in *Panaeolus* [22] with species shown to produce psilocybin, its psychoactive metabolite psilocin, and other related alkaloids [19]. Hence species of *Panaeolus*, typified by *Panaeolus papilionaceus* [23, 24], are known for their notable health benefits and hallucinogenic properties [25–35]. *P. antillarum* and *P. cyanescens* have significant antioxidant, antiviral, antibacterial and antifungal activities [33,35], while *P. cyanescens* can also be anti-inflammatory [36]. Some species that are hallucinogenic include *P. subbalteatus*, *P. cyanescens* and *P. cambodginiensis* [5,19,37].

Panaeolus mushrooms [Fig. 1(A–D)] are small, brown to blackish to cream to white, with conical- or bell-shaped caps, elongated slender stalks, attached gills and a purple-brown to black spore print [5,38–40]. Microscopic characteristics include spores with an apical germ pore and a cellular pileipellis [23,41]. Most *Panaeolus* species are coprophilic [42–44] but non-coprophilic species, such as *P. bisporus*, *P. subbalteatus*, and *P. cyanescens*, are also known from substrates such as wood [45,46]. They can be easily grown on a number of substrates [47]. Family and subfamily relationships of *Panaeolus* remain complex, as well as the total number of species after previous taxonomic changes [22]. The genus was classified under *incertae sedis* due to its uncertain taxonomic level and unknown broader relationships [40,48,49]. However, recently the genus was placed in the Galeropsidaceae [24], sometimes referred to as tribe Panaeoleae. This family includes *Copelandia*, *Panaeolina*, *Panaeolopsis*, and *Panaeolus* [24]. Of these genera, *Panaeolina* and *Panaeolopsis* are currently valid but not *Copelandia* [50,51], even if these names are inconsistently used in literature [52–55].

The total number of currently legitimate species in *Panaeolus* is confusing for individuals that lack taxonomic expertise and knowledge of this group. There are no up to date monographs of *Panaeolus*, but only previous studies on groups of 15 and 29 *Panaeolus* species [39,56,57]. Of these, only 16 species were linked to DNA sequence data of the ITS region at the National Library of Medicine [58]. In comparison, online taxonomic databases such as Index Fungorum [50] and Mycobank [51] show that *Panaeolus* has 179 and 202 names, respectively. Other online biodiversity databases, such as the Catalogue of Life: 2019 Annual Checklist [59], has 162 names of *Panaeolus*, while the Global Biodiversity Information Facility [60] recognises 90 names from 23,000 observations and photographs. These still include synonyms and species names moved to other genera.

A publication capturing only currently accepted species names of *Panaeolus* to know how diverse the genus is will be of substantial value to researchers, students, taxonomists, forensic investigators and amateur mycologists. Furthermore, it is not well-known and




Fig. 1. (A) *Panaeolus antillarum* associated with dung (photograph by L. Popich). (B) *Panaeolus cyanescens* associated with dung, one of the most potent hallucinogenic species (photograph by J. MacGillivray). (C) *Panaeolus cinctulus* (photograph by L. Popich). (D) Mottled gills characteristic of *Panaeolus* species (photograph by L. Popich).

summarised how widely these species occur across the world. This systematic review aimed to compile only the currently accepted species, along with valuable information on each species, including type locations, global distributions as well as DNA sequence availability. The data were used to summarise the diversity and occurrences of *Panaeolus* species in each continent, region and country to show areas of the world where the genus *Panaeolus* is well studied compared to areas where there are missing data. Lastly, the extent of the state of knowledge if each species contains psychoactive compounds, or an indication thereof (i.e. bruising blue), is not recorded consistently in published literature and the scientific proof for the existence of psilocybin and psilocin in some species is thus unclear. Hallucinogenic properties captured in scientific literature were thus included in this review for each species, which differs from on the ground knowledge by those growing and using these fungi. Similarly it summarised what is known about the occurrence of psilocybin and psilocin across species only based only on published scientific proof.

2. Materials and methods

A list only of species that are currently legitimate was obtained from Dr Paul Kirk (Kew Botanical Gardens, UK) and formed the basis of the review. The approach used to obtain information is based on literature and data from Google Scholar (61) and Scopus (62) (Fig. 2) and results were summarised in Tables 1–3. Literature for *Panaeolus* each year were found using the search terms “*Panaeolus*” or “*Panaeolus* species”, as well as “*Panaeolus* species”, “novel species” or “sp. nov.”, “first record”, “first finding”, “species description”, “check lists”, “field guides”, “species distribution”, “hallucinogenic”, “psychedelic”, “psychoactive”, “bluing”, “psilocybin” and “psilocin”, with titles, abstracts and text searches from 1922 to 2022 (Fig. 2). Sequences found on NCBI representing ITS [63], the most variable gene region currently sequenced, was representatively sequenced across species and added to Table 1 for easy referencing. Separate datasets were compiled using Microsoft Excel to determine the diversity of species per country, region and continent. Regions were defined according to the United Nations Geoscheme (Table 4). A map was created to visualise the distribution of *Panaeolus* species using MapChart [64].




Fig. 2. A pipeline for creating the referral datasets of *Panaeolus*, including characteristics of species in the genus such as the diversity, distribution, hallucinogenic properties and available DNA data.

Table 1
Published records of *Panaeolus* species distribution across the world.

No.	Species	Type location	Distribution	References	ITS Accession Number (NCBI)
1	<i>Panaeolus acidus</i> Sumst.1905	USA	USA.	[115]	–
2	<i>Panaeolus acuminatus</i> (P. Kumm.) Quél. 1872	France	Italy; British Isles; Brazil; India; Argentina; Iran; China; Morocco; Spain; Turkey; Macedonia; Poland; Australia; Austria; Slovakia; Scotland; Serbia; USA; Czech Republic; Croatia; France.	[23,116–125]	MH856251.1; MH856250.1; JF908518.1; MW376698.1; MW352021.1.
3	<i>Panaeolus affinis</i> (E. Horak) Ew. Gerhardt 1996	Papua New Guinea	Papua New Guinea.	[39]	–
4	<i>Panaeolus africanus</i> Ola'h 1969	Central African Republic	India; Chad; Republic of Central Africa; Sudan.	[25,49,57,126]	–
5	<i>Panaeolus albellus</i> Masee 1902	Thailand	Thailand.	[74,127,128]	–
6	<i>Panaeolus albovelutinus</i> (Rick) Raitelh. 1995	Brazil	Brazil	[129]	–
7	<i>Panaeolus alcis</i> M.M. Moser 1984	Sweden	Italy; Poland; India; Sweden.	[75,126,130,131]	KM982723.1; MW597122.1.
8	<i>Panaeolus alveolatus</i> Peck 1902	USA	USA.	[132,133]	–
9	<i>Panaeolus annulatus</i> Natarajan & Raman 1983	India	India.	[126,134]	–
10	<i>Panaeolus anomalus</i> (Murrill) Sacc. & Trotter 1925	Jamaica	Jamaica	[135]	–
11	<i>Panaeolus antillarum</i> (Fr.) Dennis 1961	Venezuela	India; Kenya; Uganda; Tanzania; British Isles; China; South Africa; Australia, Italy, East Falkland; Ukraine; Brazil; Poland; Thailand; Philippines; United States of America; Mexico; Iceland; Taiwan; Thailand; Indonesia; Sierra Leone; Netherlands; Austria; Argentina; Spain; Panama; Venezuela.	[49,73,74,78,83,95,96,119,123,126,130,136–147]	MF497586.1; MF497585.1; JF908515.1; KR998382.1.
12	<i>Panaeolus ater</i> (J.E. Lange) Kühner & Romagn. ex Bon 1985	Denmark	India; British Isles; USA; Denmark; Australia; Russia; Israel; Japan; Sri Lanka; Germany; Central Russia; Morocco; Macedonia; Turkey; Italy; Indonesia; Jordan; Greece; Scotland. British Isles.	[25,45,49,78,100,117,119,123,126,148–154]	–
13	<i>Panaeolus atrolbateus</i> Pegler & A. Henrici 1998	British Isles	British Isles.	[155]	–
14	<i>Panaeolus axfordii</i> Y. Hu, S.C. Karunarathna, P.E. Mortimer & J.C. Xu 2020	China	China.	[7]	NR169700.1; MN482689.1.
15	<i>Panaeolus bernicis</i> A.M. Young 1989	Australia	Australia.	[119]	–
16	<i>Panaeolus bisporus</i> (Malençon & Bertault) Ew. Gerhardt 1996	Morocco	USA; Morocco; Spain, South Africa.	[39,46,156]	MT110229.1; MG966283.1.
17	<i>Panaeolus bolombensis</i> Beeli 1928	Congo	Congo.	[157]	–
18	<i>Panaeolus cambodginiensis</i> Ola'h & R. Heim 1969	Colombia	Brazil; Mexico; Colombia; Hawaii.	[57,139,158,159]	AB158633.1.
19	<i>Panaeolus campanuloides</i> Guzmán & K. Yokoy. 1979	Papua New Guinea	Papua New Guinea.	[160]	–
20	<i>Panaeolus chlorocystis</i> (Singer & R.A. Weeks) Ew. Gerhardt 1987	USA	USA.	[39,46,161]	–
21	<i>Panaeolus cinctulus</i> (Bolton) Sacc. 1887	Brazil	Italy; Brazil; Poland; Mexico; Morocco.	[130,131,139,162,163]	MH590045.1; MW241166.1; MW352022.1.
22	<i>Panaeolus conicodiffractus</i> (Rick) Raitelh. 1995	Brazil	Brazil	[129]	–
23	<i>Panaeolus convexulus</i> Singer 1965	Chile	Argentina; Chile.	[164,165]	–
24	<i>Panaeolus cyanescens</i> Sacc. 1887		Brazil; India; Hawaii; Indonesia; Italy; Phillippean; Australia; Germany; Japan; Mexico; USA; Hungary (single occurrence); Thailand; Sri Lanka; Venezuela, South Africa.	[45,93,119,126,139,140,158,159,163,166–170]	MW452990.1; MW018891.1; MK855518.1; MK855517.1; MK855516.1; MK855515.1; MH547103.1; KU640172.1; KU640168.1; KT002152.1; HM035084.1; EU834287.1.

(continued on next page)

Table 1 (continued)

No.	Species	Type location	Distribution	References	ITS Accession Number (NCBI)
25	<i>Panaeolus cyanoannulatus</i> Atri, M. Kaur & Amand. Kaur 2014	India	India.	[49,171]	–
26	<i>Panaeolus cylindrosporus</i> E. Ludwig 2001	Germany	Germany	[172]	–
27	<i>Panaeolus desertorum</i> (Velen. & Dvořák) E.F. Malysheva, G. Moreno, Svetash. & M. Villarreal 2019	Spain	Spain.	[86]	–
28	<i>Panaeolus diffractus</i> (Rick) Raithelh. 1995	Brazil	Brazil	[129]	–
29	<i>Panaeolus digressus</i> Peck 1895	USA	USA.	[173]	–
30	<i>Panaeolus fimicola</i> (Pers.) Gillet 1878	France	Italy; Australia; USA; Jamaica; Puerto Rico; Brazil; Austria; Denmark; British Isles; Ireland; Spain; Switzerland; Morocco; Japan; Korea; Mongolia; Germany; Brazil; India; Macedonia; Argentina; Norway; Slovak; Turkey; Mexico; Iran; Thailand; Panama; Nigeria; Finland; Egypt; France; Poland.	[25,74,96,119,123,126,130,139,149,165,174–185]	MT451924.1; MN894012.1; JF908519.1; JF908514.1; MK394183.1; MT347601.1.
31	<i>Panaeolus fimicoloides</i> A. Pearson 1950	South Africa	South Africa.	[186,187]	–
32	<i>Panaeolus fontinalis</i> A.H. Sm. 1948	USA	USA; Mexico.	[188]	–
33	<i>Panaeolus fraxinophilus</i> A.H. Sm. 1948	USA	USA.	[116,189]	–
34	<i>Panaeolus georgii</i> Szem. 1944	Hungary	Hungary.	–	–
35	<i>Panaeolus goossensiae</i> Beeli 1928	Congo	Hawaii; Congo.	[105,157,190]	–
36	<i>Panaeolus griseofibrillosus</i> (Rick) Raithelh. 1995	Brazil	Brazil	[129]	–
37	<i>Panaeolus guttulatus</i> Bres. 1893	Italy	Italy, Iran, Greece, Turkey	[191–194]	MH592651.1, LC458688.1, KU725994.1, KU725993.1
38	<i>Panaeolus hippophilus</i> E.H.L. Krause 1928	Germany	Germany.	–	–
39	<i>Panaeolus hygrophanus</i> Velen. 1921	Czech Republic	Czech Republic.	[195]	–
40	<i>Panaeolus indicus</i> Sathe & J.T. Daniel 1979	India	India.	[196]	–
41	<i>Panaeolus lentisporus</i> Ew. Gerhardt 1996	Papua New Guinea	Papua New Guinea.	[39]	–
42	<i>Panaeolus lepistercoris</i> Atri, M. Kaur & Amand. Kaur 2014	India	India.	[171]	–
43	<i>Panaeolus lepus-stercus</i> Atri, M. Kaur & A. Kaurin 2014	India	India.	[49,126,171]	–
44	<i>Panaeolus lignicola</i> Rick 1930	Brazil	Brazil.	[139]	–
45	<i>Panaeolus linnaeanus</i> S. Imai 1938	Japan	Japan.	[197]	–
46	<i>Panaeolus microsporus</i> Ola'h & Cailleux 1969	Central African Republic	Republic of Central Africa.	[25,57]	–
47	<i>Panaeolus moellerianus</i> Singer 1960	Macquarie Is.	Faeroes Islands; Antarctic; Argentina; Macquarie Island.	[25,165,174,198]	–
48	<i>Panaeolus niveus</i> Velen. 1921	Czech Republic	Czech Republic.	[195]	–
49	<i>Panaeolus olivaceofuscus</i> Raithelh. 1977	Argentina	Argentina.	[165,174,199]	–
50	<i>Panaeolus olivaceus</i> F.H. Møller 1945	Føroyar (Faeroe Islands)	Czechoslovakia; Denmark; Finland; British Isles; Sweden; Switzerland; Australia; Morocco; Macedonia; Iran; Poland; Faeroe Islands; Turkey; Italy; Austria; France; India.	[25,95,123,149,179,200–206]	MH593015.1; MH285992.1; MF955153.1.
51	<i>Panaeolus paludosus</i> Cleland 1933	Australia	China; Australia.	[207,208]	MK278434.1
52	<i>Panaeolus panaiensis</i> Copel. 1905	Philippines	Philippines.	[209]	–
53	<i>Panaeolus papilionaceus</i> (Bull.) Quél. 1872	France	Italy; USA; Bahamas; Cuba; Puerto Rico; San Vincent Island; Brazil; Chile;	[23,25,44,68,74,122,123,125,126,	MH632116.1; MW915589.1; MW633031.1; MK397571.1;

(continued on next page)

Table 1 (continued)

No.	Species	Type location	Distribution	References	ITS Accession Number (NCBI)
			Colombia; Uruguay; Venezuela; Czechoslovakia; Finland; British Isles; Ireland; Macedonia; Norway; Russia; Spain; Sweden; Switzerland; Morocco; Ukraine; South Africa; Uganda; China; Hong Kong; India; Israel; Japan; Korea; Kuwait; Philippines; Australia; Sri Lanka; Germany; Netherlands; Argentina; Turkey; Taiwan; Iran; Egypt; Poland; Austria; Thailand; Croatia; Paraguay; Iceland; France.	[130,148,174,179, 181,183,200,203, 210]	MK500858.1; MN258670.1; LC481956.1; MK028487.1; MK439503.1; LC458685.1; MH979305.1; MF156263.1; MH169580.1; MH100727.1; MH100681.1; MF628989.1; LT716041.1; KF830093.1; KC414234.1.
54	<i>Panaeolus plantaginiformis</i> (Lebedeva) E.F. Malysheva 2019	Russia	Argentina; Peru; Russia; Asia	[86]	–
55	<i>Panaeolus pseudoguttulatus</i> Hauskn. & Krisai 2009	Austria	Austria.	[146]	–
56	<i>Panaeolus pseudopapilionaceus</i> Copel. 1905	Philippines	Philippines.	[209,211]	–
57	<i>Panaeolus regis</i> De Seynes 1901	France	France.	[212]	–
58	<i>Panaeolus reticulatus</i> Overh. 1916	USA	Macedonia; USA; Slovakia.	[149,213]	–
59	<i>Panaeolus rickenii</i> Hora 1960	British Isles	Australia; Brazil; India; Spain; Iran; Poland; Siberia; Korea; Argentina; Turkey; Greece; Austria; British Isles.	[100,165,174,179, 181,214–219]	JF908516.1; KY559329.1; MK966650.1; MK966649.1; MK966648.1; JF908523.1; MK351680.1.
60	<i>Panaeolus rubricaulis</i> Petch 1925	Sri Lanka	Papua New Guinea; Sri Lanka; Vietnam.	[25,44,220]	–
61	<i>Panaeolus rufus</i> Overh. 1916	USA	USA.	[213]	–
62	<i>Panaeolus semiglobatus</i> (Murrill) Sacc. & Trotter 1925	USA	USA; Mongolia	[135,221]	–
63	<i>Panaeolus semilanceatus</i> Peck 1909	Canada	Ireland; Australia.	[25,119,222]	–
64	<i>Panaeolus semiovatus</i> (Sowerby) S. Lundell & Nannf. 1938	Germany	Italy; British Isles; Brazil; India; Indonesia; Morocco; Argentina; Macedonia; Spain; Macedonia; Colombia; Mongolia; Falkland Island; Hawaii; Canada; USA; Turkey; Poland; Thailand; Croatia; Mexico; Bulgaria; Kazakhstan; Iceland.	[66,68,73,74,105, 117,123,125,126, 130,131,139,143, 149,165,174,200, 223–232]	MH856012.1; MT712776.1; MK386836.1; JF908517.1; MH856675.1; MK386822.1; MF955154.1.
65	<i>Panaeolus squamulosus</i> Velen. 1921	Czech Republic	Czech Republic.	[195]	–
66	<i>Panaeolus subbalteatus</i> (Berk. & Broome) Sacc. 1887	British Isles	USA; Guadalupe; Martinique; Argentina; Brazil; British Isles; Iceland; Italy; Russia (including Siberia); South Africa; India; Japan; Papua New Guinea; Philippines; Australia; New Zealand; Hawaii; Germany; Brussels; Belgium; India; Iran; Bulgaria; Canada.	[25,126,158,163, 174,179,230,233]	MW192454.1; MH855554.1; MH855553.1; MH855551.1; MH855550.1; MN960188.1; MN622762.1; MF955157.1; MF955156.1; MF955155.1; JF961370.1; AB092794.1; MH855552.1; MZ197976.1.
67	<i>Panaeolus subfirmus</i> P. Karst. 1889	Finland	Italy; Poland; Argentina; Iran; Greece; Falkland Islands; Turkey; Finland.	[73,123,130,131, 174,179,234,235]	–
68	<i>Panaeolus sylvaticus</i> Silva-Filho & Cortez 2019	Brazil	Brazil.	[88,139]	–
69	<i>Panaeolus texensis</i> V.E. Tyler & A.H. Sm. 1963	USA	USA.		–
70	<i>Panaeolus tirunelveliense</i> (Natarajan & Raman) Ew. Gerhardt 1996	India	India; Hawaii; Cambodia.	[39,45,49]	–
71	<i>Panaeolus tropicalis</i> Ola'h 1969	Cambodia	Brazil; Cambodia; Republic of Central Africa; Mexico; Hawaii; India.	[126,139,158,159, 236]	JF961377.1.
72	<i>Panaeolus uliginicola</i> (Speg.) Sacc. 1891	Argentina	Argentina.	[174,237]	–
73	<i>Panaeolus uliginosus</i> Jul. Schäff. 1947	Germany	Slovakia; France; Germany; Canada.	[212,238–240]	AY129363.1.
74	<i>Panaeolus variabilis</i> Overh. 1916	USA	USA.	[213]	–

(continued on next page)

Table 1 (continued)

No.	Species	Type location	Distribution	References	ITS Accession Number (NCBI)
75	<i>Panaeolus venenosus</i> Murrill 1916	USA	British Isles; USA.	[201,241]	–
76	<i>Panaeolus venezolanus</i> Guzmán 1978	Venezuela	Mexico; Venezuela; India.	[25,126,242]	–
77	<i>Panaeolus westii</i> (Murrill) Murrill 1942	USA	USA	[243]	–

3. Results

3.1. Literature

In this review, 258 articles were studied for information on the 77 *Panaeolus* species that are currently legitimate (Table 1). Publications on *Panaeolus* mushrooms have up to now slightly increased, but especially in popular literature, as indicated by Google Scholar searches (Fig. 3). The number of citations of published literature on *Panaeolus* has also increased. This trend that is most evident over the last decade, probably is a result of the increase in the interest in psilocybin and psilocin containing species of the genus.

3.2. Type locations

In North America, the United States of America (USA) has 13 type locations for species from out of the 77 currently legit species, and one type species from Canada (Table 1). In South America (9 species), three species have been described from type locations in Brazil, two more from Venezuela, two from Argentina, one from Chile, and one from Colombia. In Asia (12 species), India has six species described from type locations, one species in Japan, one species from China, two species from the Philippines, one species from Thailand and another from Cambodia. In Europe (19 species), species were described from type locations all over the continent, including three species from the United Kingdom, one from Austria, three from Czech Republic, one from Denmark, one from Finland, four from France, three from Germany, one from Hungary, one from Spain and one from Sweden. The Oceania region (five species) has two species described from type locations in Australia, and three from Papua New Guinea. Six species have been described from type locations in Africa and include two species from Central African Republic, two more from Congo, one species from Morocco, and one from South Africa.

3.3. Distribution and diversity

The highest known species diversity of *Panaeolus* occurs in Asia (32 species) (Fig. 4). This is followed by South America (27 species) and Europe (26 species). North America had 21 species, while Oceania, with Australia, New Zealand and Melanesia, had 19 species of *Panaeolus*, and Africa 16 species.

Some species have a wide geographic distribution from multiple continents whereas others appear to have a restricted distribution and are currently known from one or a few countries (Table 1). Twenty eight species of *Panaeolus* are intercontinental and occur on more than two continents. Some examples include *P. cinculus*, *P. rickenii*, *P. subbalteatus*, *P. tropicalis*, *P. africanus*, *P. ater*, *P. subfirmis*, *P. olivaceus*, *P. fimicola*, *P. papilionaceus*, *P. semiovatus*, *P. uliginosus*, *P. antillarum* and *P. cyanescens*. Species that are only known intra-continental but occur in more than one location include *P. cambodginiensis*, *P. convexulus*, *P. moellerianus*, *P. bisporus*, *P. alcis* and *P. fontinalis*. However, the majority of species have only been recorded at their type locations (*P. olivaceofuscus*, *P. uliginosa*, *P. benicis*, *P. pseudoguttulatus*, *P. atrobalteatus*, *P. axfordii*, *P. bolombensis*, *P. hygrophanus*, *P. niveus*, *P. squamulosus*, *P. georgii*, *P. desertorum*, *P. lignicola*, *Pa. sylvaticus*, *P. annulatus*, *P. cyanoannulatus*, *P. indicus*, *P. lepistercoris*, *P. lepus-stercus*, *P. affinis*, *P. campanuloides*, *P. lentisporus*, *P. panaiensis*, *P. pseudopapilionaceus*, *P. fimicoloides*, *P. albellus*, *P. acidus*, *P. alveolatus*, *P. chlorocystis*, *P. digressus*, *P. fraxinophilus*, *P. rufus*, *P. texensis*, and *P. variabilis*).

In Asia with the highest diversity, division of data from published records into regions according to the United Nations Geoscheme shows that most *Panaeolus* species are recorded in southern Asia, south-eastern Asia, and eastern Asia with 23, 15 and 13 species, respectively (Table 2, Figs. 5 and 6). Western Asia has ten known species and central Asia only has a single species from Kazakhstan, namely *P. semiovatus*, which is also a widely distributed species.

In Europe, most *Panaeolus* species are recorded in eastern Europe (20 species) and southern Europe (17 species), while western Europe has 15 species and northern Europe records 13 species. In South America, *Panaeolus* has a diversity of 21 species, while in North America, 21 species have also been recorded. Central America and the Caribbean have nine and three species recorded, respectively. In Australia and New Zealand, 12 and two species have been recorded, while Papua New Guinea in the Melanesia region, have five species recorded. Africa has the lowest diversity of *Panaeolus* compared to its size (Fig. 4). Northern Africa has the highest diversity on the continent with eight species of *Panaeolus*, while Central Africa only has five species, southern Africa four species and eastern and western Africa each having two recorded species.

Table 2
Global distribution of *Panaeolus* according to continent, region and country.

Continent	Species per continent	Regions	Species per region	Country	Species per country		
Asia	<i>Panaeolus acuminatus</i>	Central Asia	<i>Panaeolus semiovatus</i>	Kazakhstan	<i>Panaeolus semiovatus</i>		
	<i>P. africanus</i>	Eastern Asia	<i>P. acuminatus</i>	China	<i>P. acuminatus</i>		
	<i>P. albellus</i>		<i>P. antillarum</i>		<i>P. antillarum</i>		
	<i>P. alcis</i>		<i>P. ater</i>	Japan	<i>P. axfordii</i>		
	<i>P. annulatus</i>		<i>P. axfordii</i>		<i>P. paludosus</i>		
	<i>P. antillarum</i>		<i>P. cyanescens</i>		<i>P. papilionaceus</i>		
	<i>P. ater</i>		<i>P. fimicola</i>		<i>P. ater</i>		
	<i>P. axfordii</i>		<i>P. linnaeanus</i>		<i>P. cyanescens</i>		
	<i>P. cyanescens</i>		<i>P. paludosus</i>		<i>P. fimicola</i>		
	<i>P. cyanoannulatus</i>		<i>P. papilionaceus</i>		<i>P. linnaeanus</i>		
	<i>P. fimicola</i>		<i>P. rickenii</i>		<i>P. papilionaceus</i>		
	<i>P. guttulatus</i>		<i>P. semiglobatus</i>		<i>P. subbalteatus</i>		
	<i>P. indicus</i>		<i>P. semiovatus</i>		<i>P. fimicola</i>		
	<i>P. lepistercoris</i>		<i>P. subbalteatus</i>		<i>P. papilionaceus</i>		
	<i>P. lepus-stercus</i>		South-Eastern Asia		<i>P. albellus</i>	Mongolia	<i>P. semiglobatus</i>
	<i>P. linnaeanus</i>						<i>P. semiovatus</i>
	<i>P. olivaceus</i>						<i>P. fimicola</i>
	<i>P. paludosus</i>						<i>P. papilionaceus</i>
	<i>P. panaiensis</i>						<i>P. rickenii</i>
	<i>P. papilionaceus</i>						<i>P. antillarum</i>
	<i>P. pseudopapilionaceus</i>	<i>P. papilionaceus</i>					
	<i>P. rickenii</i>	<i>P. tirunelveliense</i>					
	<i>P. rubricaulis</i>	<i>P. tropicalis</i>					
	<i>P. semiglobatus</i>	<i>P. albellus</i>					
	<i>P. semiovatus</i>	<i>P. antillarum</i>	Thailand				
	<i>P. sepulchralis</i>	<i>P. ater</i>					
	<i>P. solidipes</i>	<i>P. cyanescens</i>	Vietnam				
	<i>P. subbalteatus</i>	<i>P. fimicola</i>					
	<i>P. subfirmus</i>	<i>P. panaiensis</i>	Indonesia				
	<i>P. tirunelveliense</i>	<i>P. papilionaceus</i>					
	<i>P. tropicalis</i>	<i>P. pseudopapilionaceus</i>	Philippines				
	<i>P. venezolanus</i>	<i>P. rubricaulis</i>					
					<i>P. antillarum</i>		
					<i>P. ater</i>		
					<i>P. cyanescens</i>		
					<i>P. semiovatus</i>		
					<i>P. panaiensis</i>		
					<i>P. papilionaceus</i>		
					<i>P. pseudopapilionaceus</i>		
					<i>P. subbalteatus</i>		
					<i>P. acuminatus</i>		
					<i>P. africanus</i>		
					<i>P. alcis</i>		
					<i>P. annulatus</i>		
					<i>P. antillarum</i>		
					<i>P. ater</i>		
					<i>P. cyanescens</i>		
				<i>P. cyanoannulatus</i>			
				<i>P. fimicola</i>			
				<i>P. guttulatus</i>			
				<i>P. indicus</i>			
				<i>P. lepistercoris</i>			
				<i>P. lepus-stercus</i>			
				<i>P. olivaceus</i>			
				<i>P. papilionaceus</i>			
				<i>P. rickenii</i>			
				<i>P. semiovatus</i>			
				<i>P. sepulchralis</i>			
				<i>P. solidipes</i>			
				<i>P. subbalteatus</i>			
				<i>P. tirunelveliense</i>			
				<i>P. tropicalis</i>			
				<i>P. venezolanus</i>			
				<i>P. ater</i>			
				<i>P. cyanescens</i>			
				<i>P. papilionaceus</i>			
				<i>P. rubricaulis</i>			
				<i>P. acuminatus</i>			
				<i>P. fimicola</i>			
				<i>P. guttulatus</i>			
				<i>P. olivaceus</i>			
				<i>P. papilionaceus</i>			
				<i>P. acuminatus</i>			
				<i>P. fimicola</i>			
				<i>P. guttulatus</i>			
				<i>P. olivaceus</i>			
				<i>P. papilionaceus</i>			

(continued on next page)

Table 2 (continued)

Continent	Species per continent	Regions	Species per region	Country	Species per country
			<i>P. subfirmus</i> <i>P. uliginicola</i> <i>P. cambodginiensis</i> <i>P. cinctulus</i> <i>P. cyanescens</i> <i>P. sylvaticus</i> <i>P. tropicalis</i> <i>P. lignicola</i> <i>P. convexulus</i> <i>P. subfirmus</i> <i>P. venezolanus</i>	Brazil	<i>P. subfirmus</i> <i>P. uliginicola</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. cambodginiensis</i> <i>P. cinctulus</i> <i>P. cyanescens</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. sylvaticus</i> <i>P. tropicalis</i> <i>P. lignicola</i> <i>P. convexulus</i> <i>P. papilionaceus</i> <i>P. cambodginiensis</i>
				Chile	<i>P. papilionaceus</i> <i>P. semiovatus</i> <i>P. antillarum</i> <i>P. subfirmus</i> <i>P. papilionaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subfirmus</i> <i>P. papilionaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. cyanescens</i> <i>P. papilionaceus</i> <i>P. venezolanus</i>
				Colombia	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				Falkland Island	<i>P. subfirmus</i> <i>P. papilionaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. semiovatus</i> <i>P. subfirmus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. cyanescens</i> <i>P. papilionaceus</i> <i>P. venezolanus</i>
				Paraguay	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				Uruguay	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				Venezuela	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
Europe	<i>P. acuminatus</i> <i>P. alcis</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. atrolbalteatus</i> <i>P. bisporus</i> <i>P. cinctulus</i> <i>P. cyanescens</i> <i>P. cylindrosporus</i> <i>P. fimicola</i> <i>P. georgii</i> <i>P. hippophilus</i> <i>P. niveus</i> <i>P. olivaceus</i> <i>P. panaiensis</i> <i>P. plantaginiformis</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. regis</i> <i>P. reticulatus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. semiovatus</i> <i>P. squamulosus</i> <i>P. subbalteatus</i> <i>P. subfirmus</i> <i>P. uliginosus</i> <i>P. venenosus</i>	Northern Europe	<i>P. acuminatus</i> <i>P. alcis</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. atrolbalteatus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i>	Denmark	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. alcis</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				Faeroe Islands	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. alcis</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				Finland	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i>
				Iceland	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. alcis</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				Norway	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. alcis</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				Sweden	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. moellerianus</i> <i>P. olivaceus</i> <i>P. subfirmus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. alcis</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i>
				British Isles	<i>P. ater</i> <i>P. atrolbalteatus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. olivaceus</i> <i>P. venenosus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i>
		Western Europe	<i>P. acuminatus</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. cyanescens</i> <i>P. cylindrosporus</i> <i>P. fimicola</i> <i>P. hippophilus</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. regis</i> <i>P. rickenii</i>	Austria	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. venenosus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i>
				Belgium	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. venenosus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i>
				France	<i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. subbalteatus</i> <i>P. venenosus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. pseudoguttulatus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. venenosus</i>

(continued on next page)

Table 2 (continued)

Continent	Species per continent	Regions	Species per region	Country	Species per country
			<i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. uliginosus</i>	Germany	<i>P. papilionaceus</i> <i>P. regis</i> <i>P. uliginosus</i> <i>P. ater</i> <i>P. cyanescens</i> <i>P. cylindrosporus</i> <i>P. fimicola</i> <i>P. hippophilus</i> <i>P. papilionaceus</i> <i>P. subbalteatus</i> <i>P. uliginosus</i>
				Netherlands	<i>P. antillarum</i> <i>P. papilionaceus</i>
				Switzerland	<i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i>
		Eastern Europe	<i>P. acuminatus</i> <i>P. alcis</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. cinctulus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. panaiensi</i> <i>P. plantaginiformis</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subfirmus</i> <i>P. subbalteatus</i> <i>P. niveus</i> <i>P. squamulosus</i> <i>P. cyanescens</i> <i>P. georgii</i> <i>P. reticulatus</i> <i>P. uliginosus</i>	Bulgaria	<i>P. papilionaceus</i> <i>P. semiovatus</i> <i>P. subbalteatus</i>
				Czech Republic	<i>P. hygrophanus</i> <i>P. acuminatus</i> <i>P. niveus</i>
				Hungary	<i>P. squamulosus</i> <i>P. cyanescens</i> <i>P. georgii</i>
				Poland	<i>P. acuminatus</i> <i>P. alcis</i> <i>P. antillarum</i> <i>P. cinctulus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subfirmus</i> <i>P. ater</i>
				Russia (based on all name variations)	<i>P. panaiensis</i> <i>P. plantaginiformis</i> <i>P. subbalteatus</i> <i>P. acuminatus</i> <i>P. reticulatus</i> <i>P. uliginosus</i> <i>P. antillarum</i> <i>P. papilionaceus</i>
				Slovakia	<i>P. acuminatus</i> <i>P. reticulatus</i> <i>P. uliginosus</i> <i>P. antillarum</i> <i>P. papilionaceus</i>
				Ukraine	<i>P. acuminatus</i> <i>P. acuminatus</i> <i>P. papilionaceus</i> <i>P. semiovatus</i> <i>P. ater</i> <i>P. guttulatus</i> <i>P. rickenii</i> <i>P. subfirmus</i>
		Southern Europe	<i>P. acuminatus</i> <i>P. alcis</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. bisporus</i> <i>P. cinctulus</i> <i>P. cyanescens</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. reticulatus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. subfirmus</i>	Croatia	<i>P. acuminatus</i> <i>P. papilionaceus</i> <i>P. semiovatus</i>
				Greece	<i>P. ater</i> <i>P. guttulatus</i> <i>P. rickenii</i> <i>P. subfirmus</i>
				Italy	<i>P. acuminatus</i> <i>P. alcis</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. cinctulus</i> <i>P. cyanescens</i> <i>P. fimicola</i> <i>P. guttulatus</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. subfirmus</i> <i>P. acuminatus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. bisporus</i> <i>P. desertorum</i> <i>P. fimicola</i>
				Serbia	<i>P. acuminatus</i> <i>P. acuminatus</i>
				Spain	<i>P. antillarum</i> <i>P. bisporus</i> <i>P. desertorum</i> <i>P. fimicola</i>

(continued on next page)

Table 2 (continued)

Continent	Species per continent	Regions	Species per region	Country	Species per country
					<i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semiovatus</i> <i>P. acuminatus</i> <i>P. ater</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. reticulatus</i> <i>P. semiovatus</i> <i>P. acuminatus</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. bernicis</i> <i>P. cyanescens</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. paludosus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. subbalteatus</i> <i>P. albovelutinus</i> <i>P. subbalteatus</i> <i>P. affinis</i> <i>P. campanuloides</i> <i>P. lentisporus</i> <i>P. rubricaulis</i> <i>P. semilanceatus</i> <i>P. subbalteatus</i>
Oceania	<i>P. acuminatus</i> <i>P. affinis</i> <i>P. albovelutinus</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. bernicis</i> <i>P. campanuloides</i> <i>P. cyanescens</i> <i>P. fimicola</i> <i>P. lentisporus</i> <i>P. olivaceus</i> <i>P. paludosus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. rubricaulis</i> <i>P. semilanceatus</i> <i>P. subbalteatus</i>	Australia and New Zealand	<i>P. acuminatus</i> <i>P. albovelutinus</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. bernicis</i> <i>P. cyanescens</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. paludosus</i> <i>P. papilionaceus</i> <i>P. rickenii</i> <i>P. semilanceatus</i> <i>P. subbalteatus</i>	Australia	
		Melanesia	<i>P. affinis</i> <i>P. campanuloides</i> <i>P. lentisporus</i> <i>P. rubricaulis</i> <i>P. subbalteatus</i>	Papua New Guinea	<i>P. affinis</i> <i>P. campanuloides</i> <i>P. lentisporus</i> <i>P. rubricaulis</i> <i>P. subbalteatus</i> <i>P. fimicola</i> <i>P. papilionaceus</i> <i>P. acuminatus</i> <i>P. ater</i> <i>P. bisporus</i> <i>P. cinctulus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. semiovatus</i> <i>P. fimicola</i> <i>P. antillarum</i> <i>P. africanus</i> <i>P. microsporius</i> <i>P. tropicalis</i> <i>P. africanus</i> <i>P. bolombensis</i> <i>P. goossensiae</i> <i>P. fimicola</i> <i>P. antillarum</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. papilionaceus</i> <i>P. antillarum</i> <i>P. antillarum</i> <i>P. fimicoloides</i> <i>P. papilionaceus</i> <i>P. subbalteatus</i>
Africa	<i>P. acuminatus</i> <i>P. africanus</i> <i>P. antillarum</i> <i>P. ater</i> <i>P. bisporus</i> <i>P. bolombensis</i> <i>P. cinctulus</i> <i>P. fimicola</i> <i>P. fimicoloides</i> <i>P. goossensiae</i> <i>P. microsporius</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. semiovatus</i> <i>P. subbalteatus</i> <i>P. tropicalis</i>	Northern Africa	<i>P. acuminatus</i> <i>P. ater</i> <i>P. bisporus</i> <i>P. cinctulus</i> <i>P. fimicola</i> <i>P. olivaceus</i> <i>P. papilionaceus</i> <i>P. semiovatus</i>	Egypt	
				Morocco	
		Western Africa	<i>P. fimicola</i> <i>P. antillarum</i>	Nigeria	
		Middle/Central Africa	<i>P. africanus</i> <i>P. microsporius</i> <i>P. tropicalis</i> <i>P. bolombensis</i> <i>P. goossensiae</i>	Sierre Leone	
				Central African Republic	
				Chad	
				Congo	
		Eastern Africa	<i>P. antillarum</i> <i>P. papilionaceus</i>	Kenya	
				Uganda	
				Tanzania	
		Southern Africa	<i>P. antillarum</i> <i>P. fimicoloides</i> <i>P. papilionaceus</i> <i>P. subbalteatus</i>	South Africa	

3.4. Distribution of *Panaeolus* species per country

Using published literature [63,64] (Tables 1 and 2), a constructed map shows that 78 countries have records of *Panaeolus* (Fig. 6). However, many others lacked records (light grey). On average, countries with records had between one to six species (blue). Below follows a more detailed summary of trends per country for each continental region (Figs. 5 and 6, Table 2).

3.4.1. Asia

South eastern Asia has species reported from Cambodia, Thailand, Vietnam, Indonesia and the Philippines, while Malaysia, Myanmar and Singapore lack records. In the southern Asia region, India has the highest diversity of species, with 22 species. Countries surrounding India and Iran such as Afghanistan, Bangladesh, Bhutan, Maldives, Nepal and Pakistan lack records of *Panaeolus* species. Central Asia only has a report from Kazakhstan, namely of the commonly occurring *P. semiovatus*, and lacks reports from Kyrgyzstan,

Table 3
Reported hallucinogenic characteristics of *Panaeolus* species.

	Species	Hallucinogenic	Bluing	Psilocybin	Psilocin	References
1	<i>Panaeolus acuminatus</i>	N	–	N	N	[104,158,244]
2	<i>P. affinis</i>	Y	–	–	–	[190]
3	<i>P. africanus</i>	Y	Y	Y	Y	[25,158,190,236,245,246]
4	<i>P. antillarum</i>	Y?	N	Y?	Y?	[25,190,246]
5	<i>P. ater</i>	N	–	N	N	[25,190,246]
6	<i>P. bisporus</i>	Y	Y	–	–	[46,247]
7	<i>P. cambodginiensis</i>	Y	Y	Y	N	[9,98,158,236,245,248]
8	<i>P. campanuloides</i>	Y	–	Y	N	[9,158,246]
9	<i>P. chlorocystis</i>	Y	Y	–	–	[25]
10	<i>P. cinctulus</i>	Y	Y	–	–	[249]
11	<i>P. cyanescens</i>	Y	Y	Y	Y	[9,97,98,104,158]
12	<i>P. cyanoannulatus</i>	Y	Y	–	–	[87]
13	<i>P. fimicola</i>	Y	Y	Y	Y	[9,25,98,158]
14	<i>P. fontinalis</i>	N	–	N	N	[9,158,250]
15	<i>P. fraxinophilus</i>	–	–	N	–	[9,158]
16	<i>P. goossensiae</i>	N	–	N	N	[9,190]
17	<i>P. microsporus</i>	Y	Y	N	Y	[9,25,158,236]
18	<i>P. moellerianus</i>	Y	–	–	–	[25,198]
19	<i>P. olivaceofuscus</i>	Y	Y	–	–	[251]
20	<i>P. olivaceus</i>	Y	Y	Y	N	[25,98,252]
21	<i>P. papilionaceus</i>	N	N	N	N	[9,25]
22	<i>P. reticulatus</i>	Y	–	–	–	[25]
23	<i>P. rickenii</i>	N	–	N	N	[9,25,253]
24	<i>P. rubricaulis</i>	Y	–	–	–	[220]
25	<i>P. semilanceatus</i>	Y	–	–	–	[254]
26	<i>P. semiovatus</i>	N	–	N	N	[158,190]
27	<i>P. subbalteatus</i>	Y	Y	Y	N	[9,25,98,158,244,245,255,256]
28	<i>P. tropicalis</i>	Y	Y	Y	Y	[9,105,158]
29	<i>P. uliginosus</i>	N	–	N	N	[9]
30	<i>P. venenosus</i>	Y	–	–	–	[244,245,252,257]
31	<i>P. venezolanus</i>	Y	–	–	–	[25,258]
	TOTAL Yes	20	13	8	5	
	TOTAL No	8	2	10	12	
	TOTAL uncertain from publications	1	0	1	1	
	TOTAL unknown from publications	1	16	12	13	

Tajikistan, Turkmenistan, Uzbekistan. Western Asia has reports of *Panaeolus* species from Israel, Jordan, Kuwait and Turkey, but no reports from Armenia, Azerbaijan, Bahrain, Cyprus, Georgia, Iraq, Lebanon, Oman, Qatar, Saudi Arabia, State of Palestine, Syrian Arab Republic, United Arab Emirates, and Yemen. Of the reported species some species only have records from their type locations, such as *P. albellus* from Thailand, *P. annulatus*, *P. cyanoannulatus*, *P. indicus*, *P. lepistercoris*, *P. lepus-stercus* from India, *P. axfordii* and *P. paludosus* from China, and *P. linnaeanus* from Japan.

3.4.2. North America, Central America, the Caribbean and South America

In North America, the United States of America (USA) had 20 species of which two also occurred in Canada, namely *P. semiovatus* and *P. subbalteatus*. Canada had reports of a third species, namely *P. uliginosus*, which does not occur in the USA. Given that two species found in Canada are also found in the United States, there are possibly more species. Checklists of North American fungi list detailed descriptions of many of these species including *P. campanulatus*, *P. cinctulus*, *P. cyanescens*, *P. semiovatus*, *P. solidipes* and *P. subbalteatus* [65–67].

Central America and the Caribbean have nine recorded species, all of which were also found in Mexico. Three species appear to be widely distributed in Central America and the Caribbean, namely *P. papilionaceus*, *P. subbalteatus* and *P. fimicola*. In the Caribbean islands, *P. papilionaceus* occurs in the Bahamas, Cuba, Puerto Rico and San Vincent Island, *P. subbalteatus* occurs in Guadalupe and Martinique, and *P. fimicola* in Jamaica and Puerto Rico.

South America has 21 recorded species of *Panaeolus*, mostly observed from Argentina, Brazil and Colombia [68]. *P. papilionaceus* and *P. subbalteatus* occurs all over South America [69]. *P. cyanescens* and *P. antillarum* are reported from Brazil [68,70,71], *P. venezolanus* from Venezuela [72], and *P. cubensis* from Colombia. *P. antillarum*, *P. papilionaceus*, *P. subfirmis* and *P. semiovatus* were reported on the Falkland Islands from horse and cattle dung [73]. Across both American continents *P. antillarum* is thus widely reported [74]. Furthermore, in South America species such as *P. papilionaceus* and *P. subbalteatus* are widespread across the continent.

Species that are restricted to the USA include *P. acidus*, *P. alveolatus*, *P. chlorocystis*, *P. digressus*, *P. faxinophilus*, *P. rufus*, *P. texensis*, and *P. variabilis*. Species from the USA but with restricted global distributions include *P. fontinalis* (USA and Mexico), *P. goossensiae* (USA and Congo), *P. bisporus* (USA, Morocco and Spain) and *P. venenosus* (USA and British Isles). In South America, *P. convexus* is only recorded in Argentina and Chile. *P. cambodginiensis* is only reported from Brazil, Mexico, and Colombia, but then also recorded from the USA. *P. venezolanus* is found in Mexico and Venezuela, also with a report from India. *P. moellerianus* appears to occur far south in the Faeroes Islands, Antarctic, Argentina and Macquarie Island. *P. lignicola* and *P. sylvaticus* are only located from their type locations in

Table 4

United Nations Geoscheme (data from <https://unstats.un.org/unsd/methodology/m49/>). The United Nations Geoscheme divides the world into regions and sub-regions. This assignment is for statistical convenience and does not imply any assumption regarding political or other affiliation of countries or territories.

Region	Sub-region	Countries and territories
Africa	Northern Africa	Algeria, Egypt, Libya, Morocco, Sudan, Tunisia, Western Sahara.
	Eastern Africa	British Indian Ocean Territory, Burundi, Comoros, Djibouti, Eritrea, Ethiopia, French Southern Territories, Kenya, Madagascar, Malawi, Mauritius, Mayotte, Mozambique, Réunion, Rwanda, Seychelles, Somalia, South Sudan, Uganda, United Republic of Tanzania, Zambia, Zimbabwe.
	Middle/Central Africa	Angola, Cameroon, Central African Republic, Chad, Congo, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Sao Tome and Principe.
	Southern Africa	Botswana, Eswatini, Lesotho, Namibia, South Africa.
Americas	Latin America and the Caribbean	Benin, Burkina Faso, Cabo Verde, Côte d'Ivoire, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Mali, Mauritania, Niger, Nigeria, Saint Helena, Senegal, Sierra Leone, Togo. Caribbean: Anguilla, Antigua and Barbuda, Aruba, Bahamas, Barbados, Bonaire, Sint Eustatius and Saba, British Virgin Islands, Cayman Islands, Cuba, Curaçao, Dominica, Dominican Republic, Grenada, Guadeloupe, Haiti, Jamaica, Martinique, Montserrat, Puerto Rico, Saint Barthélemy, Saint Kitts and Nevis, Saint Lucia, Saint Martin (French Part), Saint Vincent and the Grenadines, Sint Maarten (Dutch part), Trinidad and Tobago, Turks and Caicos Islands, United States Virgin Islands Central America: Belize, Costa Rica, El Salvador, Guatemala, Honduras, Mexico, Nicaragua, Panama South America: Argentina, Bolivia (Plurinational State of), Bouvet Island, Brazil, Chile, Colombia, Ecuador, Falkland Islands (Malvinas), French Guiana, Guyana, Paraguay, Peru, South Georgia and the South Sandwich Islands, Suriname, Uruguay, Venezuela (Bolivarian Republic of).
	Northern America	Bermuda, Canada, Greenland, Saint Pierre and Miquelon, United States of America.
Asia	Central Asia	Kazakhstan, Kyrgyzstan, Tajikistan, Turkmenistan, Uzbekistan.
	Eastern Asia	China, Democratic People's Republic of Korea, Japan, Mongolia, Republic of Korea.
	South - Eastern Asia	Brunei Darussalam, Cambodia, Indonesia, Lao People's Democratic Republic, Malaysia, Myanmar, Philippines, Singapore, Thailand, Timor-Leste, Vietnam.
Europe	Southern Asia	Afghanistan, Bangladesh, Bhutan, India, Iran, (Islamic Republic of), Maldives, Nepal, Pakistan, Sri Lanka.
	Western Asia	Armenia, Azerbaijan, Bahrain, Cyprus, Georgia, Iraq, Israel, Jordan, Kuwait, Lebanon, Oman, Qatar, Saudi Arabia, State of Palestine, Syrian Arab Republic, Turkey, United Arab Emirates, Yemen.
	Eastern Europe	Belarus, Bulgaria, Czech Republic, Hungary, Poland, Republic of Moldova, Romania, Russian Federation, Slovakia, Ukraine.
Oceania	Northern Europe	Åland Islands, Channel Islands (Guernsey, Jersey, Sark), Denmark, Estonia, Faroe Islands, Finland, Iceland, Ireland, Isle of Man, Latvia, Lithuania, Norway, Svalbard and Jan Mayen Islands, Sweden, United Kingdom of Great Britain and Northern Ireland.
	Southern Europe	Albania, Andorra, Bosnia and Herzegovina, Croatia, Gibraltar, Greece, Holy See, Italy, Malta, Montenegro, Portugal, San Marino, Serbia, Slovenia, Spain, Republic of Macedonia.
	Western Europe	Austria, Belgium, France, Germany, Liechtenstein, Luxembourg, Monaco, Netherlands, Switzerland.
Oceania	Australia and New Zealand	Australia, Christmas Island, Cocos (Keeling) Islands, Heard Island and McDonald Islands, New Zealand, Norfolk Island.
	Melanesia	Fiji, New Caledonia, Papua New Guinea, Solomon Islands, Vanuatu.
	Micronesia	Guam, Kiribati, Marshall Islands, Micronesia (Federated States of), Nauru, Northern Mariana Islands, Palau, United States Minor Outlying Islands.

Brazil, and *P. olivaceofuscus* and *P. uliginicola* in Argentina.

Areas that lack published records of *Panaeolus* species in North America include Greenland, Saint Pierre and Miquelon, according to the names of the United Nations Bioschemes. Countries in the Caribbean and Central America still lacking records of *Panaeolus* species include British Virgin Islands, Cayman Islands, Curaçao, Dominica, Grenada, Montserrat, Saint Barthélemy, Saint Lucia, Saint Martin (French Part), Sint Maarten (Dutch part), Trinidad and Tobago, Turks and Caicos Islands, United States Virgin Islands, and Bouvet Island. South American countries that do not have any reported *Panaeolus* species are Anguilla, Antigua and Barbuda, Aruba, Barbados, Bonaire, Bolivia, Ecuador, French Guiana, Guyana, Peru, Sint Eustatius and Saba, South Georgia, Suriname, and the South Sandwich Islands.

3.4.3. Europe

A significant amount of literature is available on the coprophilous agarics of Europe [68], including records of *Panaeolus* from 26 different countries. The continent is well studied and *Panaeolus* species are widespread. Across Europe *P. subbalteatus* and *P. acuminatus* are the most widely distributed species [5,69].

In Northern Europe *P. alcidis* was recorded in Sweden [68,75] and *P. antillarum*, *P. papilionaceus* and *P. semiovatus* in Iceland [68,76,77]. These and eight other species were recorded in the British Isles [68,78], which were also the country in the region with the most records. In southern Europe, regions around Italy were quite diverse after 13 species were recorded [79]. Other countries having numerous species included Austria and Germany in Western Europe (eight each) and Poland (10 species) in Eastern Europe.

The Czech Republic had the rare species *P. hygrophanus*, *P. niveus* and *P. squamulosus* reported from only their type locations. In Austria *P. pseudoguttulatum* and in France *P. regis*, were recorded only from their type locations. *P. semilanceata* has only been described from Ireland and Australia, and *P. atrobalteatus* from the British Isles. *P. desertorum* is only recorded at its type location in Spain.

Although most of Europe has been well investigated, there are countries with no *Panaeolus* species recorded. In northern Europe




Fig. 3. Number of publications and citations per year for *Panaeolus* based on Microsoft Academic [114] and Scopus (2022) [58] (accessed on 15 January 2023).




Fig. 4. An overview of the distribution of *Panaeolus* in different continents of the world according to published literature.

many islands had no records, including Åland Islands, Channel Islands (Guernsey, Jersey, Sark), Svalbard and Jan Mayen Islands, as well as the countries Estonia, Isle of Man, Latvia, and Lithuania. In eastern Europe countries such as Belarus, Republic of Moldova, and Romania do not have records of *Panaeolus* species. In Southern Europe, Albania, Andorra, Bosnia and Herzegovina, Gibraltar, Holy See, Malta, Montenegro, Portugal, San Marino and Slovenia lack records of *Panaeolus* species. In Western Europe, only a small region remains void of records, including Liechtenstein, Luxembourg and Monaco.

3.4.4. Oceania

The Oceania region consists out of Australia and New Zealand, Melanesia (containing Fiji, New Caledonia, Papua New Guinea, Solomon Islands, Vanuatu) and Micronesia (Guam, Kiribati, Marshall Islands, Federated States of Micronesia, Nauru, Northern Mariana Islands, Palau, United States Minor Outlying Islands). Only Australia, New Zealand and Papua New Guinea have recorded records. No records were found of *Panaeolus* species in Micronesia, inclusive of Guam, Kiribati, Marshall Islands, Nauru, Northern Mariana Islands, Palau and United States Minor Outlying Islands. In the Melanesia region countries including Fiji, New Caledonia,




Fig. 5. An overview of *Panaeolus* distribution in different regions across the world according to publications.




Fig. 6. A distribution map of *Panaeolus* across the world. Zero records are presented as grey, countries with one to six species are presented as blue, which fades to a red in countries that have 13–24 records. Created from published literature (Table 1) and illustrated using Mapchart [62]. (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

Solomon Islands, and Vanuatu lacked species.

Australia has 12 species including some of the most widely spread global species such as *P. fimicola*, *P. papilionaceus*, *P. semilanceata* and *P. subbalteatus*. New Zealand only has two species, namely *P. alboveletinus* and *P. subbalteatus*. The Melanesia region has five species, all from Papua New Guinea, namely *P. affinis*, *P. campanuloides*, *P. lentisporus*, *P. rubricaulis* and *P. subbalteatus*, with *P. affinis*, *P. campanuloides* and *P. lentisporus* restricted from this country. Australia has an indigenous species *P. bernicis*, only known from the type location, as well as a rare species, *P. paludosus*, only found in Australia and China.

3.4.5. Africa

Northern Africa has the highest known diversity of *Panaeolus* but only from two countries, with eight species from Morocco, while *P. fimicola* and *P. papilionaceus* also occur in Egypt. Southern Africa has four species recorded from South Africa only, including *P. antillarum*, *P. fimicoloides*, *P. papilionaceus* and *P. subbalteatus* [80–82]. Eastern Africa has two species, namely *P. antillarum* and *P. papilionaceus*, recorded in Kenya, Uganda and Tanzania [68,74,83]. Western Africa recorded *P. fimicola* and *P. antillarum* from Nigeria and Sierra Leone, respectively, and Middle Africa recorded five species, namely *P. africanus*, *P. microsporus*, *P. tropicalis*, *P. bolombensis*, and *P. goossensiae*, from Central African Republic, Chad and the Congo [5]. Africa has three species that have only been recorded from their type locations including *P. bolombensis* from the Congo, *P. fimicoloides* from South Africa, and *P. microsporus*

described in the Republic of Central Africa.

A large number of countries lack published records of *Panaeolus*, and Africa appears to be the most poorly studied with regards to the biodiversity of *Panaeolus*. These include Algeria, Libya, Sudan, Tunisia, and Western Sahara in northern Africa. In eastern Africa, no *Panaeolus* records exist from British Indian Ocean Territory, Burundi, Comoros, Djibouti, Eritrea, Ethiopia, French Southern Territories, Madagascar, Malawi, Mauritius, Mayotte, Mozambique, Réunion, Rwanda, Somalia, South Sudan, Zambia, and Zimbabwe. Angola, Democratic Republic of the Congo, Equatorial Guinea, Gabon, Sao Tome and Principe in central Africa, and Lesotho, and Namibia in Southern Africa, and Benin, Burkina Faso, Cabo Verde, Côte d'Ivoire, Gambia, Ghana, Guinea, Guinea-Bissau, Liberia, Mali, Mauritania, Niger, Saint Helena, Senegal, Togo in Western Africa lack records.

3.5. DNA sequences of *Panaeolus*

Of the 77 legitimate *Panaeolus* species, sequences of the ITS region are available for 18 species on Genbank [58]. These include recent novel species and species that are widely occurring such as *P. antillarum*, *P. sphinctrinus*, *P. acuminatus*, *P. antillarum*, *P. campanulatus*, *P. retirugis*, *P. rickenii*, *P. semiovatus*, *P. alcis*, *P. bisporus*, *P. foeniseccii* and *P. papilionaceus*. *P. cambodiginensis*, *P. subbalteatus*, *P. campanulatus* and *P. retirugis* (Table 1). Only one ex-type sequence, namely that of the ITS sequence for *P. axfordii*, exist. Whereas Large and Small Subunit sequences for the ribosomal operon are available these have low resolution power at the species level, but will be helpful for the resolution of deeper relationships of the genus. Only one Translation Elongation Factor sequence for a *P. papilionaceus* strain is available, and no other genes that are normally used for multi-gene phylogenies at the species level have been sequenced. Genome sequences are available but thus far only for *P. papilionaceus* and *P. cyanescens* [84].

3.6. *Panaeolus* hallucinogenic species

Indications of hallucinogenic properties have been recorded for only 22 *Panaeolus* species (Table 3) out of the 77 total number of species (29%). As many as 20 species were recorded to be hallucinogenic in published literature, and eight species were reported as non-hallucinogenic, including *P. acuminatus*, *P. ater*, *P. fontinalis*, *P. fraxinophilus*, *P. goossensiae*, *P. rickenii*, *P. papilionaceus*, *P. semiovatus* and *P. uliginosus*. Bruising was recorded for 13 species (17%), leaving 64 species lacking descriptions of presence of absence of bruising. Psilocybin was recorded in eight species (10% of total number) and psilocin in five species (6%). Psilocybin was not found in any trace amounts in 10 species and similarly, psilocin was not found in 12 species. Species that are shown to contain both psilocybin and psilocin include only four species, namely *P. africanus*, *P. cyanescens*, *P. fimicola*, and *P. tropicalis*. Psilocybin can be found in four species, which lack psilocin, namely *P. campanuloides*, *P. cambodiginensis*, *P. olivaceus*, and *P. subbalteatus*. One species, namely *P. microsporus*, has concentrations of psilocin but lacks psilocybin. Hallucinogenic species that remain unexplored for psilocybin and psilocin are 12 and 13 species, respectively, including *P. affinis*, *P. bisporus*, *P. chlorocystis*, *P. cinctulus*, *P. cyanoannulatus* and *P. olivaceofuscus*, to name a few (Table 4).

A third of the known hallucinogenic species are widely distributed and occur on three or more continents, including *P. acuminatus*, *P. cinctulus*, *P. cyanescens*, *P. fimicola*, *P. olivaceus*, *P. reticulatus*, *P. semilanceatus* and *P. rickenii*. However, two thirds of the known hallucinogenic species have limited distributions, including *P. rubricaulis* (from Papua New Guinea, Sri Lanka and Vietnam), *P. moellerianus* (Faeroes Islands, Antarctic, Argentina and Macquire Island), *P. microsporus* (Republic of Central Africa), *P. chlorocystis* (USA), *P. campanuloides* (Papua New Guinea), *P. africanus* (India, Chad, Republic of Central Africa and Sudan) and *P. affinis* (Papua New Guinea) (Tables 1 and 2). The extent to which this distribution has been aided by human movement and use of these mushrooms is unrecorded.

4. Discussion

The systematic review of 258 articles, and the records of Index Fungorum, showed that 77 species are currently in *Panaeolus* [50, 51]. More species have thus been added since those listed in previous monographs [23,39,85]. The increase is a result of the discovery of novel species from across the world and subspecies being brought to species rank, while other species were also transferred to other genera such as to *Galeropsis* [86]. Six novel species have been described in the last decade, including *P. axfordii* [7], *P. cyanoannulatus* [87], *P. desertorum* [86], *P. lepus-stercus* [87], *P. lignicola* [87] and *P. sylvaticus* [88].

The distribution of *Panaeolus* species was found to span across 78 countries. The highest diversity was in Asia with reports from numerous countries such as India, Sri Lanka, Iran, Japan, China, and Thailand. South America and North America had the highest diversity observed from Brazil, Argentina, the USA, and Mexico. Europe had species reported from each region but the most species were from the United Kingdom, Italy, Poland, Austria and Germany. Australia has a high diversity of *Panaeolus* with 12 species. Africa is known to have *Panaeolus* biodiversity but has the least number of records of species across countries. Numerous other countries across the world also still lack reports of any *Panaeolus* species. Evidence of rare species existed, since numerous species are only known from their type locations. In contrast, some species occurred commonly across the world.

The geographic trends reported in this study most likely reflect the availability of mycological skills and access to funding, rather than true biodiversity [89,90]. The global distributions could also be attributed to the recreational use of hallucinogenic species and their often illegal spread through trade, cultivation and human consumption. This is because species that are notably hallucinogenic usually also occur widely, such as *P. cyanescens*, *P. fimicola*, *P. olivaceus*, *P. papilionaceus*, *P. subbalteatus*, and *P. tropicalis*. These better known species may also be confused with morphologically slightly different species, which could actually represent novel species.

As many as 20 *Panaeolus* species are hallucinogenic. Previously nine [91] and 13 [25] hallucinogenic *Panaeolus* species were

known. An additional seven species have now been recognised, with *P. axfordii* the most recently described [7]. Psilocybin-containing species are often, but not always, identified by a strong blue staining reaction described when bruised on the cap or stem due to the oxidative reaction between the air and psilocybin [92,93]. *P. fimicola*, *P. cyanescens* and *P. rickenii* are a few bluing species that are widely distributed [94–97]. However, presence or absence of bruising are often not reported in publications.

Blueing and hallucinogenic properties of species have been reported by end-users, and the presence of psilocin and psilocybin could perhaps be assumed from these reactions. However, relatively few scientific studies focused on establishing whether both psilocin and psilocybin are actually present, including across all of the known species. The presence of psilocybin and psilocin was confirmed in fruiting bodies for *P. africanus*, *P. ater*, *P. cambodginiensis*, *P. campanulatus*, *P. castaneifolius*, *P. cyanescens*, *P. fimicola*, *P. microspores*, *P. sphinctrinus*, *P. subbalteatus* and *P. tropicalis*. The highest levels of psilocybin was regularly found to be in *P. cyanescens* fruiting bodies [8,93,106]. Furthermore, alkaloid detection studies for all of the other known related alkaloids [19] have not yet been published. Chemical methods to detect psilocybin and psilocin include gravity flow liquid chromatography, liquid chromatography-mass spectrometry, thin layer chromatography, gas chromatography, high-performance liquid chromatography, and capillary zone electrophoresis [37,57,93,98–105].

The concentration of psilocybin and psilocin can vary appreciably within a collection of mushrooms and between species that are widely distributed [97]. For example, collections of *P. cyanescens* from Hawaii contained significantly higher levels of psilocybin compared to collections from Australia and Thailand, which barely contained traces [8,97]. Variation in concentrations of psilocybin and psilocin between specimens of the same species was also found in *P. campanulatus*, *P. castaneifolius*, *P. foenicicii*, *P. microspores*, *P. sphinctrinus* and *P. subbalteatus* [8,93,106]. There have also been instances where a species is reported both with and without psilocybin and psilocin [25,95], for example samples of *P. antillarum* from Poland contained no psilocybin or psilocin [95], but sampled did contain psilocybin and psilocin in Taiwan [107]. Misidentification of species could be the largest problem contributing to the confusion concerning published chemical studies of psychoactive fungi [25]. Psilocybin and psilocin amounts varied even in the same mushroom specimen (e.g. cap, stem, gills) [9]. This could be due to several factors including age, growing and drying conditions, and specific parts of mushrooms used for psilocybin testing [98].

For any new species, it is common practice that novel species of *Panaeolus* are also characterized using at least the internal transcribed spacer (ITS) region as part of their taxonomical description, as has been done for the recently discovered species *P. axfordii*, *P. plantaginiformis* and *P. desertorum* [7,108]. Numerous DNA extraction protocols exist, while in some studies DNA extraction methods were modified for more efficient DNA extractions to aid identifying mushrooms of hallucinogenic genera based on DNA sequence data [61,109]. Of the 77 known *Panaeolus* species, sequences of the ITS region are available for only 18 species on Genbank [58]. Because the remaining species still need to be sequenced, this results in poor species coverage for the genus that may skew and bias DNA sequence comparison results. Moreover, only one type specimens has representative sequence, namely that of *P. axfordii*. Obtaining more ex-type sequences will be vital to fully ascertain species affiliations of samples since the often indistinct, highly variable morphologies of these fungi can lead to misidentifications. Additional regions, such as the translation elongation factor, beta-tubulin, large subunit and small subunit of rRNA [110–112], should also be tested for species resolution and consistently be sequenced with the ITS region to confirm results through multi-gene analyses, which would solidify understanding of species relationships.

5. Conclusion

Panaeolus species occur across the globe. Since the first record of *P. papilionaceus* in 1872, a total of 77 species have been recorded and many more will be discovered. This is especially evident from the fact that numerous countries still have no records of *Panaeolus*, and that the true biodiversity of the genus is poorly represented in sequence data. There are at least 20 psychoactive species that belong to the genus and at least 10 species that can be considered medically valuable because they are proven to contain psilocybin. However, analysis showed that this aspect across all of the species in *Panaeolus* is still poorly studied based on sound experimental data, and not just invaluable, but unpublished, experiences by the community that uses these mushrooms recreationally or as self-medication.

Current and future regulation of these fungi and their products for psycho-assisted therapy, in clinical settings, and trade and product development by pharmaceutical and nutraceutical companies (nutraceutical companies being companies obtaining additional value from natural products other than nutrition) [19], will not be possible without up-to-date databases from various parts of the world. Such species databases should include geographical distributions, flagged geographical areas still lacking any data, whether species are rare or commonly occurring, threatened and in need of conservation, what their hallucinogenic properties are, and the extent of study as well as of use. Such information should aid studies and regulation of species of interest for bioexploitation. However, more complete surveys on the biogeography of *Panaeolus* spp. will be needed to service the various levels of regulation (biodiversity, ethics and knowledge sharing with traditional communities, conservation, commercialization, trade, use and movement of biological material, strain development, and clinical aspects) required by individual countries, and to protect their natural resources and cultural heritage against illegal trade and movement [4,19,22,113].

This review only assimilated facts that have been published in scientific literature and not word-of-mouth knowledge from the recreational community, or the more confidential data on species and strains used by industry. Whereas all effort was placed to have the review as complete and correct as possible, there are most likely areas to be improved or corrected, records that have been missed, and new records that will continuously have to be added. Furthermore, considerable more knowledge exist about the psychoactive properties of *Panaeolus* species by the recreational community and industry, and strains and species with different properties are being collected from across the world to be developed and refined with differing psilocybin content and properties. This knowledge also needs to be incorporated more actively in scientific literature and regulation schemes to better protect the important resource that presents *Panaeolus* biodiversity.

Author contribution statement

All authors listed have significantly contributed to the development and the writing of this article.

Data availability statement

Data included in article/supp. material/referenced in article.

Additional information

No additional information is available for this paper.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

References

- [1] P. Stafford, *Psychedelics Encyclopedia*, Ronin Publishing, 2013.
- [2] G. Guzmán, New studies on hallucinogenic mushrooms: history, diversity, and applications in psychiatry, *Int. J. Med. Mushrooms* 17 (2015) 1019–1029, <https://doi.org/10.1615/intjmedmushrooms.v17.i1.1.10>.
- [3] R.R. Griffiths, W.A. Richards, U. McCann, R. Jesse, Psilocybin can occasion mystical-type experiences having substantial and sustained personal meaning and spiritual significance, *Psychopharmacology (Berl.)* 187 (2006) 268–283, <https://doi.org/10.1007/s00213-006-0457-5>.
- [4] R.C. Van Court, M.S. Wiseman, K.W. Meyer, D.J. Ballhorn, K.R. Amses, J.C. Slot, B.T.M. Dentinger, R. Garibay-Orijel, J.K. Uehling, Diversity, biology, and history of psilocybin-containing fungi: suggestions for research and technological development, *Fungal Biol* 126 (2022) 308–319, <https://doi.org/10.1016/j.funbio.2022.01.003>.
- [5] P. Stamets, *Psilocybin Mushrooms of the World*, 1996.
- [6] S. Hartman, *Psilocybin could Be legal for therapy by 2021*, *Roll. Stone* (2018) 1.
- [7] Y. Hu, P. Mortimer, S. Karunarathna, O. Raspé, I. Promputtha, K. Yan, J. Xu, K. Hyde, A new species of *Panaeolus* (Agaricales, Basidiomycota) from Yunnan, Southwest China, *Phytotaxa* 434 (2020) 22–34, <https://doi.org/10.11646/phytotaxa.434.1.3>.
- [8] T. Stijve, Worldwide occurrence of psychoactive mushrooms - an update, *Czech Mycol.* 48 (1995) 11–19.
- [9] C. Andersson, J. Kristinsson, J. Gry, Occurrence and Use of Hallucinogenic Mushrooms Containing Psilocybin Alkaloids, Nordic Council of Ministers., Denmark., 2009.
- [10] COMPASS Pathways, COMPASS pathways receives FDA Breakthrough therapy designation for psilocybin therapy for treatment-resistant depression, PR Newswire. <https://compasspathways.com/compass-pathways-receives-fda-breakthrough-therapy-designation-for-psilocybin-therapy-for-treatment-resistant-depression/>, 2018 (accessed September 28, 2020).
- [11] R.E. Doblin, M. Christiansen, L. Jerome, B. Burge, The past and future of psychedelic science: an introduction to this issue, *J. Psychoact. Drugs* 51 (2019) 93–97, <https://doi.org/10.1080/02791072.2019.1606472>.
- [12] R.L. Carhart-Harris, M. Bolstridge, C.M.J. Day, J. Rucker, R. Watts, D.E. Erritzoe, M. Kaelen, B. Giribaldi, M. Bloomfield, S. Pilling, J.A. Rickard, B. Forbes, A. Feilding, D. Taylor, H.V. Curran, D.J. Nutt, Psilocybin with psychological support for treatment-resistant depression: six-month follow-up, *Psychopharmacology (Berl.)* 235 (2018) 399–408, <https://doi.org/10.1007/s00213-017-4771-x>.
- [13] G. Guzman, in: J. Misra, S. Deshmukh (Eds.), *The Hallucinogenic Mushrooms: Diversity, Traditions, Use and Abuse with Special Reference to the Genus Psilocybe, Fungi Differ. Environ.*, Taylor & Francis, New York, 2009, pp. 256–276.
- [14] F. Tyš, T. Páleníček, J. Horáček, Psilocybin - summary of knowledge and new perspectives, *Eur. Neuropsychopharmacol* 24 (2014) 342–356, <https://doi.org/10.1016/j.euroneuro.2013.12.006>.
- [15] T.S. Krebs, P.Ø. Johansen, Psychedelics and mental health: a population study, *PLoS One* 8 (2013), <https://doi.org/10.1371/journal.pone.0063972>.
- [16] S. Reiche, L. Hermle, S. Gutwinski, H. Jungaberle, P. Gasser, T. Majjić, Serotonergic hallucinogens in the treatment of anxiety and depression in patients suffering from a life-threatening disease: a systematic review, *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 81 (2018) 1–10, <https://doi.org/10.1016/j.pnpbp.2017.09.012>.
- [17] R.L. Carhart-Harris, G.M. Goodwin, The therapeutic potential of psychedelic drugs: past, present, and future, *Neuropsychopharmacology* 42 (2017) 2105–2113, <https://doi.org/10.1038/npp.2017.84>.
- [18] F.A. Moreno, C.B. Wiegand, E.K. Taitano, P.L. Delgado, Safety, tolerability, and efficacy of psilocybin in 9 patients with obsessive-compulsive disorder, *J. Clin. Psychiatry* 67 (2006) 1735–1740, <https://doi.org/10.4088/JCP.v67n1110>.
- [19] D. Strauss, S. Ghosh, Z. Murray, M. Gryzenhout, Psilocybin containing mushrooms: a rapidly developing biotechnology industry in the psychiatry, biomedical and nutraceutical fields, *3 Biotech* 12 (2022) 339, <https://doi.org/10.1007/s13205-022-03355-4>.
- [20] J. Phelps, R.N. Shah, J.A. Lieberman, The rapid rise in investment in psychedelics - cart before the horse, *JAMA Psychiatr.* 79 (2022) 189–190, <https://doi.org/10.1001/jamapsychiatry.2021.3972>.
- [21] P. Tullis, How ecstasy and psilocybin are shaking up psychiatry, *Nature* 589 (2021) 506–509, <https://doi.org/10.1038/d41586-021-00187-9>.
- [22] D. Strauss, S. Ghosh, Z. Murray, M. Gryzenhout, An overview on the taxonomy, phylogenetics and ecology of the psychedelic genera *Psilocybe*, *Panaeolus*, *Pluteus* and *Gymnopilus*, *Front. For. Glob. Change.* 5 (2022) 1–9, <https://doi.org/10.3389/fgc.2022.813998>.
- [23] L. Quelet, *Les Champignons du Jura et des Vosges*, *Mém. Société D'Emulation Montbél.* 2 (5) (1872) 43–332.
- [24] J. Kalichman, P.M. Kirk, P.B. Matheny, A compendium of generic names of agarics and Agaricales, *Taxon* 69 (2020) 425–447, <https://doi.org/10.1002/tax.12240>.
- [25] G. Guzmán, J.W. Allen, J. Gartz, A Worldwide geographical distribution of the Neurotropic fungi, an analysis and discussion, *Africa* 14 (1998) 1–107.
- [26] D. Reingardiene, J. Vilcinskaitė, R. Lazauskas, [Hallucinogenic mushrooms], *Med. Kaunas Lith.* 41 (2005) 1067–1070.
- [27] S. Ling, F. Ceban, L.M.W. Lui, Y. Lee, K.M. Teopiz, N.B. Rodrigues, O. Lipsitz, H. Gill, M. Subramaniapillai, R.B. Mansur, K. Lin, R. Ho, J.D. Rosenblat, D. Castle, R.S. McIntyre, Molecular Mechanisms of Psilocybin and Implications for the Treatment of Depression, *CNS Drugs*, 2021, <https://doi.org/10.1007/s40263-021-00877-Y>.
- [28] R. Carhart-Harris, B. Giribaldi, R. Watts, M. Baker-Jones, A. Murphy-Beiner, R. Murphy, J. Martell, A. Blemings, D. Erritzoe, D.J. Nutt, Trial of psilocybin versus escitalopram for depression, *N. Engl. J. Med.* 384 (2021) 1402–1411, <https://doi.org/10.1056/NEJMoa2032994>.
- [29] S.J. Tai, E.M. Nielson, M. Lennard-Jones, R.L. Johanna Ajantaival, R. Winzer, W.A. Richards, F. Reinholdt, B.D. Richards, P. Gasser, E. Malievskaja, Development and evaluation of a therapist training program for psilocybin therapy for treatment-resistant depression in clinical research, *Front. Psychiatr.* 12 (2021), <https://doi.org/10.3389/FPSYT.2021.586682/FULL>.

- [30] J. Daniel, M. Haberman, Clinical potential of psilocybin as a treatment for mental health conditions, *Ment. Health Clin.* 7 (2017) 24–28, <https://doi.org/10.9740/mhc.2017.01.024>.
- [31] T. Chi, J.A. Gold, A review of emerging therapeutic potential of psychedelic drugs in the treatment of psychiatric illnesses, *J. Neurol. Sci.* 411 (2020) 116–175, <https://doi.org/10.1016/j.jns.2020.116715>.
- [32] F. Moreno, C. Wiegand, E. Taitano, Safety, tolerability, and efficacy of psilocybin in 9 patients with obsessive-compulsive disorder, *J. Clin. Psychiatry* 67 (2006) 1735–1740.
- [33] R.G. Bustillos, R.R. Milton Dulay, J. John Bauto, F. Pascual, K. Baltazar, H. Weisel Bunag, A. Macatula, M. Ann Nicolas, M.M. Ann Torres, J.C. Nillosa, J. Cesar Dela Cruz, S.P. Kalaw, R.G. Reyes, Mycochemical profile of mycelia and fruiting body of *Panaeolus cyanescens* and its optimal submerged culture conditions for antioxidant properties, *Int. J. PURE Appl. Biosci.* 2 (2014) 175–181.
- [34] S. Wasser, Medicinal mushrooms as a source of antitumor and immunomodulating polysaccharides, *Appl. Microbiol. Biotechnol.* 60 (2002) 258–274, <https://doi.org/10.1007/s00253-002-1076-7>.
- [35] R.R. Milton Dulay, D. Rmr, D. Roxas Mjb, C. Jmp, D.E. Guzman, Proximate Composition and Antioxidant Activity of *Panaeolus antillarum*, a Wild Coprophilous Mushroom, 2015, <https://doi.org/10.5943/cream/5/1/7>.
- [36] S.M. Nkadameng, C.M.L. Steinmann, J.N. Eloff, Anti-inflammatory effects of four psilocybin-containing magic mushroom water extracts in vitro on 15-lipoxygenase activity and on lipopolysaccharide-induced cyclooxygenase-2 and inflammatory cytokines in human U937 macrophage cells, *J. Inflamm. Res.* 14 (2021) 3729–3738, <https://doi.org/10.2147/JIR.S317182>.
- [37] J. Kristinsson, Occurrence and Use of Hallucinogenic Mushrooms Containing Psilocybin Alkaloids, Nordic Council of Ministers, 2008.
- [38] B.H. Rumack, D.G. Spoerke, *Handbook of Mushroom Poisoning: Diagnosis and Treatment* Barry H. Rumack, David G. Spoerke - Google Books, CRC Press, 1994.
- [39] E. Gerhardt, Taxonomische Revision der Gattungen *Panaeolus* und *Panaeolina* (Fungi, Agaricales, Coprinaceae), *Bibl. Bot.* 147 (1996) 1–149.
- [40] M.Q. He, R.L. Zhao, K.D. Hyde, D. Begerow, M. Kemler, A. Yurkov, E.H.C. McKenzie, O. Raspé, M. Kakishima, S. Sánchez-Ramírez, E.C. Vellinga, R. Halling, V. Papp, I.V. Zmitrovich, B. Buyck, D. Ertz, N.N. Wijayawardene, B.K. Cui, N. Schoutteten, X.Z. Liu, T.H. Li, Y.J. Yao, X.Y. Zhu, A.Q. Liu, G.J. Li, M.Z. Zhang, Z. L. Ling, B. Cao, V. Antonín, T. Boekhout, B.D.B. da Silva, E. De Crop, C. Decock, B. Dima, A.K. Dutta, J.W. Fell, J. Geml, M. Ghobad-Nejhad, A.J. Giachini, T. B. Gibertoni, S.P. Gorjón, D. Haelewaters, S.H. He, B.P. Hodkinson, E. Horak, T. Hoshino, A. Justo, Y.W. Lim, N. Menolli, A. Mešić, J.M. Moncalvo, G. M. Mueller, L.G. Nagy, R.H. Nilsson, M. Noordeloos, J. Nuytinck, T. Orihara, C. Ratchadawan, M. Rajchenberg, A.G.S. Silva-Filho, M.A. Sulzbacher, Z. Tkalčec, R. Valenzuela, A. Verbeken, A. Vizzini, F. Wartchow, T.Z. Wei, M. Weiß, C.L. Zhao, P.M. Kirk, Notes, outline and divergence times of Basidiomycota, *Fungal Divers.* 99 (2019) 105–367, <https://doi.org/10.1007/s13225-019-00435-4>.
- [41] L. Quélet, *Panaeolus* (Fr.) Quélet, *Mém Soc Émul Montbél.* 2 (1872) 151.
- [42] Y.-W. Wang, S.-S. Tzean, Dung-associated, potentially hallucinogenic mushrooms from taiwan, *Taiwania* 60 (2015) 160–168, <https://doi.org/10.6165/tai.2015.60.160>.
- [43] R. O' Hanlon, in: K. Kavanagh (Ed.), *Fungi in the environment*, third ed. *Fungi Biol. Appl.*, 2018, pp. 333–355, <https://doi.org/10.1002/9781119374312>.
- [44] S. Ediriweera, R.L.C. Wijesundera, C. Nanayakkara, J. Weerasena, First report of *Panaeolus sphinctrinus* and *Panaeolus foenicicii* (Psathyrellaceae, Agaricales) on elephant dung from Sri Lanka, *Front. Environ. Microbiol.* 1 (2015) 19–23, <https://doi.org/10.11648/j.fem.20150102.12>.
- [45] Paul Stamets, *Psilocybin Mushrooms of the World*, Ten Speed Press, Berkeley California, 1999, p. 245.
- [46] B. Senn-Irlet, A. Nyffenegger, R. Brenneisen, *Panaeolus bisporus* - an adventitious fungus in central Europe, rich in psilocin, *Mycologist* 13 (1999) 176–179, [https://doi.org/10.1016/S0269-915X\(99\)80107-4](https://doi.org/10.1016/S0269-915X(99)80107-4).
- [47] P. Stamets, J.S. Chilton, *The Mushroom Cultivator: A Practical Guide to Growing Mushrooms at Home*, first ed., Agarikon Press, 1983.
- [48] N. Wijayawardene, K. Hyde, L.K.T. Al-Ani, L. Tedersoo, D. Haelewaters, K.C. Rajeshkumar, R.L. Zhao, A. Aptroot, D.V. Leontyev, R.K. Saxena, Outline of Fungi and fungus-like taxa, *Mycosphere* 11 (2020) 1060–1456.
- [49] A. Kaur, N.S. Atri, M. Kaur, Diversity of coprophilous species of *Panaeolus* (Psathyrellaceae, Agaricales) from Punjab, India, *Biodiversitas* 15 (2014) 115–130, <https://doi.org/10.13057/biodiv/d150202>.
- [50] Index Fungorum, <http://www.indexfungorum.org/>, 2021. (Accessed 7 October 2021) accessed.
- [51] Mycobank, <https://www.mycobank.org/>, 2021. (Accessed 7 October 2021) accessed.
- [52] S. Ediriweera, R.L.C. Wijesundera, C. Nanayakkara, J. Weerasena, First report of *Panaeolus sphinctrinus* and *Panaeolus foenicicii* (Psathyrellaceae, Agaricales) on elephant dung from Sri Lanka, *Front. Environ. Microbiol.* 1 (2015) 19–23, <https://doi.org/10.11648/j.fem.20150102.12>.
- [53] A. Tóth, A. Hausknecht, I. Krisai-Greilhuber, T. Papp, C. Vágvölgyi, L.G. Nagy, Iteratively refined guide trees help improving alignment and phylogenetic inference in the mushroom family Bolbitiaceae, *PLoS One* 8 (2013), <https://doi.org/10.1371/journal.pone.0056143>.
- [54] F. Wartchow, A.S. Carvalho, M.C.A. Sousa, First Record of the Psychotropic Mushroom *Copelandia cyanescens* (Agaricales) from Pernambuco State, Northeast Brazil, 2009, pp. 1679–2343.
- [55] J. Kalichman, P.M. Kirk, P.B. Matheny, A compendium of generic names of agarics and Agaricales, *Taxon* 69 (2020) 425–447, <https://doi.org/10.1002/TAX.12240>.
- [56] M.Q. He, R.L. Zhao, K.D. Hyde, D. Begerow, M. Kemler, A. Yurkov, E.H.C. McKenzie, O. Raspé, M. Kakishima, S. Sánchez-Ramírez, E.C. Vellinga, R. Halling, V. Papp, I.V. Zmitrovich, B. Buyck, D. Ertz, N.N. Wijayawardene, B.K. Cui, N. Schoutteten, X.Z. Liu, T.H. Li, Y.J. Yao, X.Y. Zhu, A.Q. Liu, G.J. Li, M.Z. Zhang, Z. L. Ling, B. Cao, V. Antonín, T. Boekhout, B.D.B. da Silva, E. De Crop, C. Decock, B. Dima, A.K. Dutta, J.W. Fell, J. Geml, M. Ghobad-Nejhad, A.J. Giachini, T. B. Gibertoni, S.P. Gorjón, D. Haelewaters, S.H. He, B.P. Hodkinson, E. Horak, T. Hoshino, A. Justo, Y.W. Lim, N. Menolli, A. Mešić, J.M. Moncalvo, G. M. Mueller, L.G. Nagy, R.H. Nilsson, M. Noordeloos, J. Nuytinck, T. Orihara, C. Ratchadawan, M. Rajchenberg, A.G.S. Silva-Filho, M.A. Sulzbacher, Z. Tkalčec, R. Valenzuela, A. Verbeken, A. Vizzini, F. Wartchow, T.Z. Wei, M. Weiß, C.L. Zhao, P.M. Kirk, Notes, outline and divergence times of Basidiomycota, *Fungal Divers.* 99 (2019) 105–367, <https://doi.org/10.1007/s13225-019-00435-4>.
- [57] G. Ola'h, Le Genre *Panaeolus*, *Essai Taxinomique et Physiologique*, *Rev. Mycol. Paris.* 10 (1969) 1–273.
- [58] NCBI, The NCBI taxonomy database (Nucleotide), <https://www.ncbi.nlm.nih.gov/nucleotide>, 2021. (Accessed 7 October 2021) accessed.
- [59] COL, COL | The Catalogue of Life, <https://www.catalogueoflife.org/>, 2021. (Accessed 7 October 2021) accessed.
- [60] GBIF, GBIF | Global Biodiversity Information Facility, <https://www.gbif.org/>, 2021. (Accessed 7 October 2021) accessed.
- [61] Google Scholar, <https://scholar.google.com/>, 2021. (Accessed 7 October 2021) accessed.
- [62] Scopus, <https://www.scopus.com/search/form.uri?zone=TopNavBar&origin=searchbasic&display=basic#basic>, 2021. (Accessed 7 October 2021) accessed.
- [63] A. Wesselink, DNA Markers for Forensic Identification of Non-human Biological Traces, Thesis, University of Amsterdam, 2018. Retrieved from, <https://hdl.handle.net/11245>.
- [64] MapChart, <https://mapchart.net/world.html>, 2021. (Accessed 7 October 2021) accessed.
- [65] D. Arora, *Mushrooms Demystified: A Comprehensive Guide to the Fleshy Fungi*, Ten Speed Press, 1986.
- [66] O.K. Miller, Interesting fungi of the St. Elias Mountains, Yukon Territory, and adjacent Alaska, *Mycologia* 60 (1968) 1190–1203, <https://doi.org/10.1080/00275514.1968.12018686>.
- [67] F.C. Ghouled, *Field Guide to the Psilocybin Mushroom: Species Common to North America*, Guidance Publications, 1972.
- [68] A. Kaur, N. Atri, M. Kaur, Studies on coprophilous agaricoid mushrooms: an appraisal, *Kavaka* 52 (2019) 66–84.
- [69] A. Kaur, N.S. Atri, M. Kaur, Studies on coprophilous agaricoid mushrooms: an appraisal, *Kavaka* 52 (2019) 66–84.
- [70] R.F.R. Melo, R.D.S. Chikowski, A.N. Miller, L.C. Maia, Coprophilous agaricales (Agaricomycetes, Basidiomycota) from Brazil, *Phytotaxa* 266 (2016) 1–14, <https://doi.org/10.11646/phytotaxa.266.1.1>.
- [71] F.J.S. Calaça, N.C. Silva, S. Xavier-Santos, A checklist of coprophilous fungi and other fungi recorded on dung from Brazil, *Mycotaxon* 128 (2014) 205, <https://doi.org/10.5943/cream/5/1/8>.
- [72] G. Guzmán, Index of Taxa in the genus *Psilocybe*, *Mycotaxon* 6 (1978) 464–476.
- [73] R. Watling, M.J. Richardson, Coprophilous fungi of the Falkland Islands, *Edinb. J. Bot.* 67 (2010) 399–423, <https://doi.org/10.1017/S0960428610000156>.

- [74] D.E. Desjardin, B.A. Perry, *Panaeolus antillarum* (Basidiomycota, Psathyrellaceae) from wild elephant dung in Thailand the species has been reported from elephant dung, from Uganda, *Hausknecht Krisai-Greilhuber* 7 (2017) 275–281, <https://doi.org/10.5943/cream/7/4/4>.
- [75] M. Moser, *Panaeolus acidis*, a new species from Scandinavia and Canada, *Mycologia* 76 (1984) 551–554.
- [76] M.J. Richardson, Coprophilous fungi from Iceland, *Acta Bot. Isl.* 14 (2004) 77–102.
- [77] M.J. Richardson, Additions to the coprophilous mycota of Iceland, *Acta Bot. Isl.* 15 (2011) 23–49.
- [78] R. Watling, N.M. Gregory, *British Fungus Flora-Agaric and Boleti* 5., Strophariaceae Coprinaceae, Royal Bot. Gard. Edinb., 1987.
- [79] F. Doveri, Occurrence of coprophilous Agaricales in Italy, new records, and comparisons with their European and extra-European distribution, *Mycosphere* 1 (2010) 103–140.
- [80] G.B. Goldman, M. Gryzenhout, *Field Guide to Mushrooms and Other Fungi of South Africa*, first ed., Penguin Random House South Africa, 2019.
- [81] R. van der Walt, J. Dames, G. Hawley-McMaster, *Fungi and Lichens of the Limpopo Valley and Mapungubwe National Park, Ludwigslust Game Farms*, 2020.
- [82] M. Gryzenhout, *Pocket Guide Mushrooms of South Africa*, first ed., Struik Nature, 2021.
- [83] D.N. Pegler, A preliminary agaric flora of East Africa, *Kew Bull.* 6 (1977) 1–615.
- [84] National Center for Biotechnology Information (NCBI), No Title, Bethesda MD Natl. Libr. Med. US Natl. Cent. Biotechnol. Inf, 1988.
- [85] G.M. Ola'h, Le genre *Panaeolus*: Essai taxinomique et physiologique, Louis Marie, Paris Herb, 1969, p. 273.
- [86] E. Malysheva, G. Moreno, M. Villarreal, V. Malysheva, T. Svetasheva, The secotioid genus *Galeropsis* (Agaricomycetes, Basidiomycota): a real taxonomic unit or ecological phenomenon? *Mycol. Prog.* 18 (2019) 805–831, <https://doi.org/10.1007/s11557-019-01490-6>.
- [87] A. Kaur, N.S. Atri, M. Kaur, Two New Species of *Panaeolus* (Psathyrellaceae, Agaricales) from Coprophilous Habitats of Punjab, India, 2014.
- [88] A.G.S. Silva-Filho, C. Seger, V.G. Cortez, *Panaeolus* (Agaricales) from western Paraná state, south Brazil, with a description of a new species, *Panaeolus sylvaticus*, *Edinb. J. Bot.* 76 (2019) 297–309, <https://doi.org/10.1017/S0960428619000064>.
- [89] M. Gryzenhout, J.M. Jefwa, N.S. Yorou, The status of mycology in Africa: a document to promote awareness, *IMA Fungus* 3 (2012) 99–102, <https://doi.org/10.5598/imafungus.2012.03.01.11>.
- [90] M. Gryzenhout, The need to engage with citizen scientists to study the rich fungal biodiversity in South Africa, *IMA Fungus* 6 (2015) 58–68, <https://doi.org/10.1007/bf03449355>.
- [91] J. Ott, *Pharmacothoeon: etheogenic drugs, their plant sources and history*, *Nat. Prod.* (1993).
- [92] C. Lenz, J. Wick, D. Braga, M. García-Altares, G. Lackner, C. Hertweck, M. Gressler, D. Hoffmeister, Injury-triggered blueing reactions of *Psilocybe* “magic” mushrooms, *Angew. Chem. Int. Ed.* 59 (2020) 1450–1454, <https://doi.org/10.1002/anie.201910175>.
- [93] F. Müsshoff, B. Madea, J. Beike, Hallucinogenic Mushrooms on the German Market - Simple Instructions for Examination and Identification, in: *Forensic Sci. Int.*, 2000, pp. 389–395, [https://doi.org/10.1016/S0379-0738\(00\)00211-5](https://doi.org/10.1016/S0379-0738(00)00211-5).
- [94] N.C. Karun, K. Sridhar, Elephant dung-inhabiting macrofungi in the Western Ghats, *Curr. Res. Environ. Appl. Mycol.* 5 (2015) 60–69, <https://doi.org/10.5943/cream/5/1/8>.
- [95] M. Halama, D. Witkowska, I. Jasicka-Misiak, A. Poliwoła, An adventive *Panaeolus antillarum* in Poland (Basidiomycota, Agaricales) with notes on its taxonomy, geographical distribution, and ecology, *Cryptogam. Mycol.* 35 (2014) 3–22, <https://doi.org/10.7872/crym.v35.iss1.2014.3>.
- [96] M. Piepenbring, *Reportes Nuevos De Agaricales Para Panamá*, Panama (2008).
- [97] T. Stijve, Psilocin, psilocybin, serotonin and urea in *Panaeolus cyanescens* from various origin, *Persoonia-Mol. Phylogeny Evol. Fungi.* 15 (1992) 117–121.
- [98] Z. Alam Mahmood, Bioactive alkaloids from fungi: Psilocybin, in: *Nat. Prod. Phytochem. Bot. Metab. Alkaloids Phenolics Terpenes*, 2013, pp. 523–552, https://doi.org/10.1007/978-3-642-22144-6_19.
- [99] M.W. Beug, J. Bigwood, Psilocybin and psilocin levels in twenty species from seven genera of wild mushrooms in the Pacific Northwest, U.S.A., *J. Ethnopharmacol.* 5 (1982) 271–285, [https://doi.org/10.1016/0378-8741\(82\)90013-7](https://doi.org/10.1016/0378-8741(82)90013-7).
- [100] L.S. Gurevich, Indole derivatives in certain *Panaeolus* species from east Europe and siberia, *Mycol. Res.* 97 (1993) 251–254, [https://doi.org/10.1016/S0953-7562\(09\)80249-9](https://doi.org/10.1016/S0953-7562(09)80249-9).
- [101] J.C.I. Lee, M. Cole, A. Linacre, Identification of members of the genera *Panaeolus* and *Psilocybe* by a DNA test: a preliminary test for hallucinogenic fungi, *Forensic Sci. Int.* 112 (2000) 123–133, [https://doi.org/10.1016/S0379-0738\(00\)00181-X](https://doi.org/10.1016/S0379-0738(00)00181-X).
- [102] J. Borovicka, M. Oborník, J. Štrifbrný, M.E. Noordeloos, L.A. Parra Sánchez, M. Gryndler, Phylogenetic and chemical studies in the potential psychotropic species complex of *Psilocybe atrobrunnea* with taxonomic and nomenclatural notes, *Persoonia Mol. Phylogeny Evol. Fungi.* 34 (2015) 1–9, <https://doi.org/10.3767/003158515X685283>.
- [103] A. Adamczyk, A. Sadakierska-Chudy, J. Janoszka, A. Rymkiewicz, T. Dobosz, Hallucinogenic fungi (*Psilocybe*). Part II. Identification of *Psilocybe semilanceata* by PCR, *Arch. Med. Sadowej Kryminol.* 57 (2007) 285–288.
- [104] J. Štrifbrný, J. Borovicka, M.O. Sokol, [Levels of psilocybin and psilocin in various types of mushrooms], *Soud Lek.* 48 (2003) 45–49.
- [105] M.D. Merlin, J.W. Allen, Species identification and chemical analysis of psychoactive fungi in the Hawaiian islands, *J. Ethnopharmacol.* 40 (1993) 21–40, [https://doi.org/10.1016/0378-8741\(93\)90086-K](https://doi.org/10.1016/0378-8741(93)90086-K).
- [106] T. Maruyama, O. Shirota, N. Kawahara, K. Yokoyama, Y. Makino, Y. Goda, Discrimination of psychoactive fungi (commonly called “magic mushrooms”) based on the DNA sequence of the internal transcribed spacer region, *J. Food Hyg. Soc. Jpn.* 44 (2003) 44–48, <https://doi.org/10.3358/shokueishi.44.44>.
- [107] Y.W. Wang, S.S. Tzean, Dung-associated, potentially hallucinogenic mushrooms from Taiwan, *Taiwania* 60 (2015) 160–168, <https://doi.org/10.6165/tai.2015.60.160>.
- [108] E. Malysheva, G. Moreno, M. Villarreal, V. Malysheva, T. Svetasheva, The secotioid genus *Galeropsis* (Agaricomycetes, Basidiomycota): a real taxonomic unit or ecological phenomenon? *Mycol. Prog.* 18 (2019) 805–831, <https://doi.org/10.1007/s11557-019-01490-6>.
- [109] D. Strauss, S. Ghosh, Z. Murray, M. Gryzenhout, Genomic DNA extraction from minimal amount of dried mushroom samples, *Microb. Biosyst.* 6 (2021) 49–54, <https://doi.org/10.21608/MB.2021.91637.1036>.
- [110] D. Begerow, H. Nilsson, M. Unterseher, W. Maier, Current state and perspectives of fungal DNA barcoding and rapid identification procedures, *Appl. Microbiol. Biotechnol.* 87 (2010) 99–108, <https://doi.org/10.1007/s00253-010-2585-4>.
- [111] A.D. Tekpinar, A. Kalmer, Utility of various molecular markers in fungal identification and phylogeny, *Nova Hedwigia* 109 (2019) 187–224, https://doi.org/10.1127/nova_hedwigia/2019/0528.
- [112] W. Meyer, L. Irinyi, M.T.V. Hoang, V. Robert, D. Garcia-Hermoso, M. Desnos-Ollivier, C. Yurayart, C.C. Tsang, C.Y. Lee, P.C.Y. Woo, I.M. Pchelin, S. Uhrlaß, P. Nenoff, A. Chindamporn, S. Chen, P.D.N. Hebert, T.C. Sorrell, Database establishment for the secondary fungal DNA barcode translational elongation factor 1 α (TEF1 α), *Genome* 62 (2019) 160–169, <https://doi.org/10.1139/gen-2018-0083>.
- [113] K. Gerber, I.G. Flores, A.C. Ruiz, I. Ali, N.L. Ginsberg, E.E. Schenberg, Ethical concerns about psilocybin intellectual property, *ACS Pharmacol. Transl. Sci.* 4 (2021) 573–577, <https://doi.org/10.1021/acspsci.0c00171>.
- [114] Microsoft Academic. <https://academic.microsoft.com/home>, 2021. (Accessed 7 October 2021) accessed.
- [115] D.R. Sumstine, *Panaeolus acidis* sp. nov., *Torreya* 5 (1905) 34.
- [116] M. Contu, A. Hausknecht, A lignicolous variety of *Panaeolus acuminatus* from Sardinia, *Österr Z Pilzk* 12 (2003) 85–88.
- [117] R. Watling, R. King, N. Riddiford, R. Kingt, Botanical Journal of Scotland New and interesting records of fungi from shetland new and interesting records of Fungi from Shetland, *Bot. J. Scotl.* 53 (2009) 57–64, <https://doi.org/10.1080/03746600108684955>.
- [118] M. Xavier, Morphological description and new record of *Panaeolus acuminatus* (Agaricales) in Brazil, *Stud. Fungi.* 4 (2019) 135–141, <https://doi.org/10.5943/sif/4/1/16>.
- [119] A.M. Young, The panaeoloideae (Fungi, Basidiomycetes) of Australia, *Aust. Syst. Bot.* 2 (1989) 75–97, <https://doi.org/10.1071/SB9890075>.
- [120] V. Kučerová, I. Kautmanová, Contribution to the knowledge of macrofungi of the Muránska Planina National Park and adjacent areas Príspevok k poznaniu húb Národného parku Muránska planina a priľahlých oblastí, *Reussia* 6 (2011) 87–96.
- [121] E. Čapelja, Basidiomycetes of Temska village area (Eastern Serbia, Mt Stara Planina), *Biol. Nyssana.* 3 (2012) 91–96.
- [122] I. Akata, B. Çetin, M. Işiloğlu, Macrofungi diversity of ilgaz mountain national park and its environs (Turkey), *Mycotaxon* 113 (2010) 287–290.

- [123] V.M. Granito, D. Lunghini, Biodiversity of macrofungi in the beech forests and calcareous grasslands of the simbruini mountains regional park (central Apennines, Italy), *Plant Biosyst.* 145 (2011) 381–396, <https://doi.org/10.1080/11263504.2011.563523>.
- [124] M. Wurst, R. Kysilka, M. Flieger, Psychoactive tryptamines from basidiomycetes, *Folia Microbiol. (Praha)* 47 (2002) 3–27, <https://doi.org/10.1007/BF02818560>.
- [125] A. Mešić, Z. Tkalčec, Preliminary checklist of agaricales from Croatia Iv: Families Bolbitiaceae, coprinaceae, entolomataceae and pluteaceae, *Mycotaxon LXXXVII* (2003) 283–309.
- [126] K. Amandeep, N.S. Atri, K. Munruchi, A checklist of Coprophilous Agarics of India, *Curr. Res. Environ. Appl. Mycol.* 5 (2015) 322–348, <https://doi.org/10.5943/cream/5/4/3>.
- [127] S. Somrithipol, E. Jones, M. Tantichareon, K. Hyde, Coprophilous fungi, in: *Thai Fungal Diversity, Natl. Cent. Genet. Eng. Biotechnol. Pathum Thani* Thailand, 2004, pp. 119–128.
- [128] G.E. Massee, in: J. Schmidt (Ed.), *Flora of Koh Chang. Contributions to the knowledge of the vegetation of the gulf of Siam, Agaricineae, Bot. Tidsskr., vol. 24*, 1902, pp. 363–367.
- [129] J. Raitelhuber, *Species et combin. nov. Metrodiana* 23 (1995) 3–10.
- [130] F. Doveri, Occurrence of coprophilous Agaricales in Italy, new records, and comparisons with their European and extra-European distribution, *Mycosphere* 1 (2010) 103–140.
- [131] M. Halama, *Panaeolus subfirmus* (Agaricales, Basidiomycota), a species new for Poland, *Pol. Bot. J.* 59 (2014) 271–277, <https://doi.org/10.2478/pbj-2014-0029>.
- [132] A.P. Morgan, North American species of agaricaceae. The melanosporae, *J. Mycol.* 13 (1907) 53, <https://doi.org/10.2307/3752667>.
- [133] C.H. Peck, Report of the state Botanist (1900), *Annu. Rep. N. Y. State Mus. Nat. Hist.* 54 (1902) 131–199.
- [134] K. Natarajan, N. Raman, South Indian agaricales, *Bibl. Mycol.* 89 (1983) 1–203.
- [135] P.A. Saccardo, G.B. Traverso, A. Trotter, *Sylloge Fungorum Omnium Hucusque Cognitorum, Sumptibus Auctoris, Patavii*, 1925. <https://www.biodiversitylibrary.org/item/25489>.
- [136] B. Zhishu, Z. Guoyang, L. Taihui, *The Macrofungus Flora of China's Guangdong Province*, 1993, p. 734.
- [137] D. Reid, A. Eicker, South African Fungi 10: new species, new records and some new observations, *Mycotaxon* 73 (1999) 169–197.
- [138] A. Hausknecht, I. Krisai-Greilhuber, Pilzbeobachtungen in einem neugeschaffenen Weidegebiet, *Österr. Z. Pilzk.* 12 (2003) 101–123.
- [139] K.D. Hyde, D.S. Tennakoon, R. Jeewon, D.J. Bhat, S.S.N. Maharachchikumbura, W. Rossi, M. Leonardi, H.B. Lee, H.Y. Mun, J. Houbraeken, T.T.T. Nguyen, S. J. Jeon, J.C. Frisvad, D.N. Wanasinghe, R. Lücking, A. Aptroot, M.E.S. Cáceres, S.C. Karunarathna, S. Hongsanan, R. Phookamsak, N.I. de Silva, K. M. Thambugala, R.S. Jayawardena, I.C. Senanayake, S. Boonmee, J. Chen, Z.L. Luo, C. Phukhamsakda, O.L. Pereira, V.P. Abreu, A.W.C. Rosado, B. Bart, E. Randrianjohany, V. Hofstetter, T.B. Gibertoni, A.M. da S. Soares, H.L. Plautz, H.M.P. Sotão, W.K.S. Xavier, J.D.P. Bezerra, T.G.L. de Oliveira, C.M. de Souza-Motta, O.M.C. Magalhães, D. Bundhun, D. Harishchandra, I.S. Manawasinghe, W. Dong, S.N. Zhang, D.F. Bao, M.C. Samarakoon, D. Pem, A. Karunarathna, C. G. Lin, J. Yang, R.H. Perera, V. Kumar, S.K. Huang, M.C. Dayaratne, A.H. Ekanayaka, S.C. Jayasiri, Y. Xiao, S. Konta, T. Niskanen, K. Liimatainen, Y.C. Dai, X. H. Ji, X.M. Tian, A. Mešić, S.K. Singh, K. Phuththacharoen, L. Cai, T. Sorvongxay, V. Thiyagaraja, C. Norphanphoun, N. Chaiwan, Y.Z. Lu, H.B. Jiang, J.F. Zhang, P.D. Abeywickrama, J.V.S. Aluthmuhandiram, R.S. Brahmanage, M. Zeng, T. Chethana, D. Wei, M. Réblová, J. Fournier, J. Nekvindová, R. do Nascimento Barbosa, J.E.F. dos Santos, N.T. de Oliveira, G.J. Li, D. Ertz, Q.J. Shang, A.J.L. Phillips, C.H. Kuo, E. Camporesi, T.S. Bulgakov, S. Lumyong, E.B.G. Jones, P. Chomnunti, E. Gentekaki, F. Bungartz, X.Y. Zeng, S. Fryar, Z. Tkalčec, J. Liang, G. Li, T.C. Wen, P.N. Singh, Y. Gafforov, I. Promputtha, E. Yasanthika, I. D. Goonasekara, R.L. Zhao, Q. Zhao, P.M. Kirk, J.K. Liu, J.Y. Yan, P.E. Mortimer, J. Xu, M. Doilom, Fungal diversity notes 1036–1150: taxonomic and phylogenetic contributions on genera and species of fungal taxa, *Fungal Divers.* 96 (2019) 1–242, <https://doi.org/10.1007/s13225-019-00429-2>.
- [140] R.G. Bustillos, B. Rg, D. Rmr, K. Sp, R. Rg, Optimization of culture conditions for mycelial growth and basidiocarp production of Philippine strains of *Panaeolus antillarum* and *Panaeolus cyanescens*, *Mycosphere* 5 (2014) 398–404, <https://doi.org/10.5943/mycospere/5/3/1>.
- [141] G. Guzmán, Some distributional relationships between Mexican and United States mycofloras, *Mycologia* 65 (1973) 1319–1330, <https://doi.org/10.1080/00275514.1973.12019555>.
- [142] J. Gartz, J.W. Allen, M.D. Merlin, Ethnomycology, biochemistry, and cultivation of *Psilocybe samuiensis* Guzmán, Bandala and Allen, a new psychoactive fungus from Koh Samui, Thailand, *J. Ethnopharmacol.* 43 (1994) 73–80, [https://doi.org/10.1016/0378-8741\(94\)90006-X](https://doi.org/10.1016/0378-8741(94)90006-X).
- [143] A. Marisa, A. Sara, M. Mendes, José, P. Sousa, H. Freitas, A.M. Azul, A.H. Freitas, S.M. Mendes, A.J.P. Sousa, Fungal fruitbodies and soil macrofauna as indicators of land use practices on soil biodiversity in Montado, *Agrofor. Syst.* 82 (2011) 121–138, <https://doi.org/10.1007/s10457-010-9359-y>.
- [144] D.N. Pegler, *Studies on african agaricales: I*, Springer Behalf R, *Bot. Gard. Kew.* 21 (1968) 499–533.
- [145] T. Kuyper, A. Van Peer, J. Baars, Coprophilous Fungi Closing the Loop: Improving Circularity with Manure-Loving Mushrooms, 2021, <https://doi.org/10.18174/539315>.
- [146] A. Hausknecht, I. Krisai-Greilhuber, Die Gattungen *Panaeolina* und *Panaeolus* in Österreich und Bemerkungen zu einigen sonstigen, interessanten *Panaeolus*-Funden, *Österr. Z. Pilzk.* 18 (2009).
- [147] R.W.G. Dennis, Fungi venezuelani: IV, agaricales, *Kew Bull.* 15 (1961) 67–156.
- [148] E.S. Kholfy, O.A. Touhami, Study of *Panaeolus semiovatus* (sowerby) S. Lundell and Nannf., (1938), A coprophilous fungus, *Plant Arch* 20 (2020) 1856–1858.
- [149] M. Karadelev, S. Spasikova, Second contribution to hallucinogenic fungi in the Republic of Macedonia, *Fungal Divers.* (2006) 441–449.
- [150] A. Kaya, K. Demirel, Y. Uzun, Macrofungi diversity of Araban (Gaziantep/Turkey) district, *Biol. Divers. Conserv.* 5 (2012) 162–166.
- [151] A. Mumpuni, A. Amurwanto, D.J. Wahyono, Molecular identification of coprophilous microfungi from Banyumas District, Central Java, Indonesia, *Biodiversitas* 22 (2021) 1550–1557.
- [152] A.S. Salhab, A minireview on mushroom: emphasis on the wild mushroom of, *Jpn. Med. J.* 41 (2007) 170–178.
- [153] S. Diamandis, C. Perlerou, *The Mycoflora of the Chestnut Ecosystems in Greece*, 2001.
- [154] R. Watling, R. King, N. Riddiford, New and interesting records of fungi from Shetland, *Bot. J. Scotl.* 53 (2001) 57–64, <https://doi.org/10.1080/03746600108684955>.
- [155] A. Pegler, D.N. Henrici, *Panaeolus atrolateatus* sp. nov., a member of *Panaeolus* stirps subbalteatus (Agaricales, Strophariaceae), *Folia Cryptogam. Est.* 33 (1998) 105–108.
- [156] C. Brown, Warm Temperatures Chase the Thin Alpine Snowpack, Initiating an Early Flush of Spring Mushrooms in Washington, *MushRumors Newsl. Northwest Mushroomers Assoc.*, 2015, pp. 10–15.
- [157] M. Beeli, Contribution à l'étude de la flore mycologique du Congo. VI Fungi Goossensiani. Agaricacées rhodosporeées, *Bull. Société R. Bot. Belg.* 61 (1928) 78–103.
- [158] S.H. Pollock, *Psilocybian mycetismus* with special reference to *Panaeolus*, *J. Psychoact. Drugs* 8 (1976) 43–57, <https://doi.org/10.1080/02791072.1976.10472007>.
- [159] J.W. Allen, M.D. Merlin, Psychoactive mushroom use in Koh Samui and Koh Pha-Ngan, Thailand, *J. Ethnopharmacol.* 35 (1992) 205–228, [https://doi.org/10.1016/0378-8741\(92\)90020-R](https://doi.org/10.1016/0378-8741(92)90020-R).
- [160] Yokoyama Guzmán, *Studies on Cryptogams of Papua New Guinea*, 1979.
- [161] E. Gerhardt, *Panaeolus cyanescens* (Bk. & Br.) Sacc. und *Panaeolus antillarum* (Fr.) Dennis, zwei Adventivarten in Mitteleuropa, *Beitr. Zur Kennt. Pilze Mitteleur.* 3 (1987) 223–227.
- [162] F. Garza Ocañas, M. Quiñónez Martínez, L.G. Ocañas, A.C. Parra, H.V. Mendoza, H. Gonzalez Rodríguez, R.V. Garza, G. Guevara Guerrero, J. García Jiménez, M.G. Aranda, Some edible, toxic and medicinal mushrooms from temperate forests in the North of Mexico, *Adv. Macrofungi Divers. Ecol. Biotechnol.* (2019).
- [163] P.A. Saccardo, *Sylloge Hymenomycetum, Vol. I, Sylloge Fungorum* 5 (1887) 1–1146.
- [164] R. Singer, M. Moser, Forest mycology and forest communities in South America. 1. The early fall aspect of the mycoflora of the Cordillera Pelada (Chile), *Mycopathol. Mycol. Appl.* 26 (1965) 129–191.

- [165] N. Niveiro, E. Albertó, Checklist of the Argentine agaricales 7. Cortinariaceae and entolomataceae, Check List. 10 (2014) 72–96, <https://doi.org/10.15560/10.1.72>.
- [166] T. Stijve, A.A.R. de Meijer, Macromycetes from the state of Parana, Brazil. The psychoactive species, ArqBiol. Tecnol. 36 (1993) 313–329.
- [167] F. Wartchow, A.S. Carvalho, M.C.A. Sousa, First Record of the Psychotropic Mushroom *Copelandia cyanescens* (Agaricales) from Pernambuco State, Northeast Brazil, 2009, pp. 1679–2343.
- [168] K. Gonmori, K. Yokoyama, Acute encephalopathy caused by cyanogenic fungi in 2004, and magic mushroom regulation in Japan, Chūdoku Kenkyū Chūdoku Kenkyūkai Jun Kikanshi Jpn. J. Toxicol. 22 (2009) 61–69.
- [169] I. Siller, G. Vasas, Red list of macrofungi of Hungary (revised edition), Stud. Bot. Hung. 26 (1995) 7–14.
- [170] A.E. Delgado, G. Urdaneta, Basidiomycota fungi of the Agaricales order, in five municipalities in Zulia State, Venezuela, Rev. Fac. Agron. 19 (2002) 56–70.
- [171] A. Kaur, N.S. Atri, M. Kaur, Two new species of *Panaeolus* (Psathyrellaceae, Agaricales) from coprophilous habitats of Punjab, India, J. N. Biol. Rep. 3 (2014) 125–132.
- [172] E. Ludwig, Pilzkompendium, 2001.
- [173] ChasH. Peck, New species of fungi, Bull. Torrey Bot. Club 22 (1895) 198, <https://doi.org/10.2307/2478162>.
- [174] N. Niveiro, E. Albertó, Checklist of the Argentine agaricales 4. Tricholomataceae and polyporaceae, Mycotaxon 121 (2012) 499–500.
- [175] A.F. Parker-Rhodes, The basidiomycetes of Skokholm Island, Trans. Br. Mycol. Soc. 34 (1951) 360–367, [https://doi.org/10.1016/s0007-1536\(51\)80063-9](https://doi.org/10.1016/s0007-1536(51)80063-9).
- [176] M. Krishnaappa, S. Swapna, S. Abrar, Diversity of macrofungal communities in Chikmagalur District Of Western Ghats, India, in: Proc. 8th Int. Conf. Mushroom Biol. Mushroom Prod. ICMBMP8, 2014, pp. 1–380.
- [177] I. Mihal, Production of fruiting bodies of saprophytic fungi in spruce monocultures planted on former arable land, Ekol6gia Bratisl. 7 (1998) 152–161.
- [178] A. Kaya, Macrofungal diversity of Adiyaman province (Turkey) view project macrofungal diversity of adiyaman province (Turkey), Mycotaxon 110 (2010) 43–46, <https://doi.org/10.5248/110.43>.
- [179] E. Seidmohammadi, S. Abbasi, M.R. Asef, The first report of *Panaeolus olivaceus* and *Panaeolus guttulatus* from Iran, taxon, Biosystems 11 (2019) 2–10, <https://doi.org/10.22108/tbj.2020.118170.1093>.
- [180] J.L. Siquier, J.C. Salom, J. Espinosa, F. Esteve-Raventós, J. Llistosella, I.S. Gomes, Contribució al coneixement micològic de les illes balears (espanya). XXI, Rev. Catalana Micol. 36 (2015) 59–88.
- [181] J. Adamczyk, L. Kucharski, Macrofungi in different habitats of small postglacial ponds margins in the kujawy region (central Poland), Lejeunia Rev. Bot. 177 (2005) 1–18.
- [182] K. Kokkonen, Macrofungi in fire alleys in old quarters of Nurmes, Eastern Finland, Karstenia 45 (2005) 51–61.
- [183] S. Esraa, S. Ea, A.-A. Am, S. Fm, N. Na, M. Mt, A. Ais, I. Me, A.-A. Ma, H. Ss, A taxonomic analysis of fungi collected and described from Egypt up to 1931, Stud. Fungi 1 (2016) 11–33, <https://doi.org/10.5943/sif/1/1/2>.
- [184] M.J. Richardson, Records of French coprophilous fungi, Cryptogam. Mycol. 29 (2008) 157–177.
- [185] C.C. Gillet, Les hyménomycètes: ou, Description de tous les champignons (fungi) qui croissent en France, avec l'indication de leurs propriétés utiles ou vénéneuses, C Thomas., 1878, pp. 561–828.
- [186] T.R. Kinge, G. Goldman, A. Jacobs, G.G. Ndiritu, M. Gryzenhout, A first checklist of macrofungi for South Africa, MycoKeys 63 (2020) 1–48, <https://doi.org/10.3897/mycokeys.63.36566>.
- [187] A.A. Pearson, Cape agarics and boleti, Trans. Br. Mycol. Soc. 33 (1950), [https://doi.org/10.1016/s0007-1536\(50\)80080-3](https://doi.org/10.1016/s0007-1536(50)80080-3), 276–IN8.
- [188] A.H. Smith, Studies in the dark-spored agarics, Mycologia 40 (1948) 669–707, <https://doi.org/10.1080/00275514.1948.12017737>.
- [189] A.H. Smith, Studies in the dark-spored agarics, Mycologia 40 (1948) 669–707.
- [190] J.W. Allen, A Chemical Referral and Reference Guide to the Known Species of Psilocin and/or Psilocybin-Containing Mushrooms and Their Published Analysis and Bluing Reactions: an Updated and Revised List, 2011.
- [191] G. Bresadola, Fungi Tridentini Novi Vel Nondum Delineati, Descripti Et Iconibus Illustrati, vol. 1, Forgotten Books, 1883.
- [192] A. Kaya, Macrofungal diversity of Adiyaman Province (Turkey), Mycotaxon 110 (2010) 43–46, <https://doi.org/10.5248/110.43>.
- [193] V. Kaounas, G. Konstantinidis, D. Sofronis, M. Gkilas, L. Kottis, Seven interesting Basidiomycetes recorded in Greece for the first time, Mycol Monten 19 (2016) 97–112.
- [194] E. Seidmohammadi, S. Abbasi, M.R. Asef, The first report of *Panaeolus olivaceus* and *Panaeolus guttulatus* from Iran, Taxon, Biosystems 11 (2019) 2–10, <https://doi.org/10.22108/tbj.2020.118170.1093>.
- [195] J. Velenovský, Ceske Houby, 1921, pp. 425–632.
- [196] A.V. Sathe, J.T. Daniel, A new species of *Panaeolus* from India, Curr. Sci. 48 (1979) 905–906.
- [197] S. Imai, Studies on the agaricaceae of hokkaido. II, J. Fac. Agric. Hokkaido Imp. Univ. 43 (1938) 179–378.
- [198] R. Singer, Sobre algunas especies de hongos presumiblemente psicotropicos, Lilloa 30 (1960) 117–127.
- [199] J. Raithelhuber, Hongos Argentinos. II, 1977.
- [200] A. Pauline N'douba, A.O. Touhami, R. Benkirane, L. Zidane, A. Douira, Etude de cinq espèces coprophiles du genre *Panaeolus* (Fr.) Quélet dont trois nouvelles pour le Maroc : *Panaeolus ater*, *P. olivaceus* et *P. semiovatus*, Bull. L'Institut Sci. Rabat Sect. Sci. Vie. 34 (2012) 71–74.
- [201] R. Watling, A *Panaeolus* poisoning in scotland, Mycopathologia 61 (1977) 187–190.
- [202] H. Hüseyin Doğan, C. Öztürk, G. Kaşık, S. Aktaş, New records for Turkish mycoflora from aAlanya (Antalya) District, fen-edeb, Fakültesi Fen Derg. Sayı. 21 (2003) 21–41.
- [203] A. Hausknecht, I. Krisai-Greilhuber, W. Jaklitsch, Ergebnisse des Mykologischen Arbeitstreffens in Jerischach (Sülkärn-ten) im August/September 1998, Osterr Z Pilzk 9 (2000) 192.
- [204] P.-A. Moreau, G. Corriol, D. Borgarino, P. Aubel, C. Lavoise, F. Richard, M.-A. Selosse, Contribution À La Connaissance Des Champignons De L'étagé Thermoméditerranéen Corse II, Bull Famm N S 31 (2007) 9–33.
- [205] G. Gogoi, V. Parkash, A checklist of gilled mushrooms (Basidiomycota: agaricomycetes) with diversity analysis in Hollongapar Gibbon, J. Threat. Taxa 7 (2015) 8272–8287, <https://doi.org/10.11609/jott.1770.7.15.8272-8287>.
- [206] F.H. Møller, Fungi of the Farøes, Part I, Basidiomyceten, 1945.
- [207] Z. Ping, H. Hongying, C. Zuohong, A new record of *Panaeolus* in China, Mycosystema 23 (2004) 308–309.
- [208] J.B. Cleland, Australian fungi: notes and descriptions, Trans. Proc. R. Soc. S. Aust. No 9 (57) (1933) 187–194.
- [209] E.B. Copeland, Fungi esculentes philippinenses, Ann. Mycol. 3 (1905) 25–29.
- [210] A. Flecha Rivas, N. Niveiro, Checklist of Agaricoid Fungi from Paraguay, 2019, <https://doi.org/10.5248/133.729>.
- [211] P.W. Graff, Bibliography and new species of Philippine Fungi, Mycologia 8 (1916) 253–288, <https://doi.org/10.1080/00275514.1916.12018891>.
- [212] P.-A. Moreau, P. Hériveau, V. Bourgade, J.-M. Bellanger, R. Courteuisse, F. Fons, S. Rapior, Redécouverte et typification des champignons de la région de Montpellier illustrés par Michel-Félix Dunal et Alire Raffeneau-Delile, Cryptogam. Mycol. 32 (2011) 255–276.
- [213] L.O. Overholts, New or interesting species of gill fungi from Missouri, Ann. Mo. Bot. Gard. 3 (1916) 195, <https://doi.org/10.2307/2990074>.
- [214] S. Eun Cho, J. Won Jo, N. Kyu Kim, Y.-N. Kwag, S.-K. Han, K. Sun Chang, S. Hwan Oh, C. Sun Kim, Macrofungal survey of the Tian Shan mountains, Kyrgyzstan, Mycobiology. 47 (2019) 378–390, <https://doi.org/10.1080/12298093.2019.1661565>.
- [215] H. Alii, B. Çöl, İ. Şen, Macrofungi biodiversity of Kütahya (Turkey) province, Biol. Divers. Conserv. 10 (2017) 133–143.
- [216] V. Lucia, R. Bononi, A.K. Morbeck De Oliveira, A. De Melo Gugliotta, J. Ratier De Quevedo, Agaricomycetes (Basidiomycota, Fungi) diversity in a protected area in the Maracaju Mountains, in the Brazilian central region, Hoehnea 44 (2017) 361–377, <https://doi.org/10.1590/2236-8906-70/2016>.
- [217] S. Diamandis, A contribution to the mycoflora of the Greek forest ecosystems, Minist. Agric. Thessalon. Greece For. Res. Inst. 4 (1990) 59–66.
- [218] T. Rüdcker, Beiträge zur Pilzflora von Salzburg II-ein Zwischenbericht, Jb Haus Nat 10 (1987) 93–99.
- [219] F.B. Hora, New check list of British agarics and boleti part IV. Validations, new species and critical notes, Trans. Br. Mycol. Soc. 43 (1960) 440–459.
- [220] T. Petch, Additions to ceylon fungi. III, Ann. R. Bot. Gard. Perad. 9 (1925) 313–328.

- [221] N. Kherlenchimeg, D. Otgontsetseg, G. Kherlenchimeg, M. Urgamal, Results of the research of higher fungi in Khan Khentey, Mongolia, *Проблемы Ботаники Южной Сибири И Монголии* 21 (2022) 196–203, <https://doi.org/10.14258/pbssm.2022085>.
- [222] C.H. Peck, Report of the state Botanist. 1908, *Bull. N. Y. State Mus.* 131 (1909) 1–202.
- [223] M. Karadelev, S. Spasikova, Hallucinogenic fungi in the Republic of Macedonia, in: *Proc. 2nd Congr. Ecol. Repub. Maced. Int. Particip.* 25-29102003 Ohrid, Special Issues of Macedonian, Ecological Society, 2004, pp. 479–483.
- [224] A.F. Montoya, D.F. Arias, M.B. Agudelo, Contribución Al Conocimiento De Los Hongos Macromicetos Del Resguardo Indígena Nuestra Señora De La Candelaria De La Montaña Riosucio-Caldas, *Mus. Hist. Nat.* 9 (2005) 21–32.
- [225] K. Nyamsuren, U. Magsar, S. Batsumber, T. Myagmardorj, E. Tsogtbaatar, Y. Cho, O. Kwon, Survey on the distribution of macrofungi in Mongolia, *J Ecol Evn* 39 (2016) 91–97, <https://doi.org/10.5141/ecoenv.2016.010>.
- [226] M.W. Beug, J. Bigwood, Psilocybin and psilocin levels in twenty species from seven genera of wild mushrooms in the Pacific Northwest, U.S.A., *J. Ethnopharmacol.* 5 (1982) 271–285, [https://doi.org/10.1016/0378-8741\(82\)90013-7](https://doi.org/10.1016/0378-8741(82)90013-7).
- [227] H. Alli, M. Isiloglu, M.H. Solak, Mehmet Halil Solak Mugla Üniversitesi Macrofungi of Aydin Province, Turkey, *Mycotaxon* 99 (2007) 163–165.
- [228] A.E. Franco-Molano, E. Uribe-Calle, Agaricals and Boletals Fungi of Colombia Hongos Agaricales y Boletales de Colombia, *Biota Colomb.* 1 (2000) 25–43.
- [229] F. Garza-Ocañas, Sustainable management of edible forest mushrooms in Chihuahua, Mexico, *Actes Colloq.* in: *Int. Champignons for. Comest. Potentiel Commer. Qué. Canada Univ. Laval.*, 2009, pp. 1–14.
- [230] V.I. Fakirova, M.M. Gyosheva, C.M. Denchev, Checklist of the macromycetes of central balkan mountain (Bulgaria), in: *Proc. 6th Symp. Flora Southeast. Serbia Adjac. Territ. Sokobanja Yugosl.*, 2000, pp. 25–38.
- [231] Y. V. Rakhimova, G.A. Nam, B.D. Yermekova, U.K. Jetigenova, B. Zh Yessengulova, Ecological and Trophic Differentiation of Fungal Diversity in Aksu-Zhabagly Nature Reserve (Kazakhstan), *Sib. Ekol. Zhurnal.* 10 (2017) 595–608, <https://doi.org/10.1134/S1995425517050110>.
- [232] S. Lundell, J.A. Nannfeldt, *Panaeolus semiovatus*, *Fungi Exsicc. Succ. Fasc.* 11 (1) (1938) 501–600.
- [233] H.J. Brodie, The Heterothallism Of *Panaeolus Subbalteatus* Berk., A Sclerotium-Producing Agaric, *Can. J. Res.* 12 (1935) 657–660, <https://doi.org/10.1139/cjr35-052>.
- [234] D. Yağız, A. Afyon, M. Konuk, S. Helfer, Contributions to the Macrofungi of Kastamonu Province, Turkey, 2006.
- [235] P. Karsten, *Fragmenta mycologica XXVIII*, *Hedwigia* 28 (1889) 363–367.
- [236] G.M. Ols'ha, Le genre *Panaeolus*: Essai taxinomique et physiologique, *Louis Marie, Paris Herb.* 1969, pp. 116–119.
- [237] P.A. Saccardo, *Supplementum Universale, Pars I. Agaricaceae-Laboulbeniaceae, Sylloge Fungorum* 9 (1891) 1–1141.
- [238] I. Kautmanová, Redlist species of fungi held in the collections of slovak national museum-natural history museum (bra). ii. Endangered speci redlist species of fungi held in the collections of slovak national museum-natural history museum (bra). ii. Endangered species (en), *Acta Rer Nat. Mus Nat Slov. LI* (2005) 1–12.
- [239] A. Bresinsky, “*Galerina beinrothii*” nov. spec., “*Panaeolus uliginosus*” J. Schaeff. and other agaricales from flat moors in Upper Bavaria, *J. Mushroom Sci.* 1 (1966) 8–17.
- [240] K.S. Han, T.J. Volk, H.K. Kim, Identification of *Lacrymaria velutina* (Pers. Ex Fr.) Konrad & Maubl. from Micheon-myeon, Jinju-City, *Mycobiology* 38 (2010) 249–255, <https://doi.org/10.4489/MYCO.2010.38.4.249>.
- [241] W.A. Murrill, *Panaeolus venenosus*, *Mycologia* 8 (1916) 186–187.
- [242] G. Guzmán, *Panaeolus venezolanus*, *Mycotaxon* 7 (1978) 221–224.
- [243] W.A. Murrill, New fungi from Florida, *Lloydia Cincinnati* 5 (1942) 136–157.
- [244] M.W. Beug, J. Bigwood, Psilocybin and psilocin levels in twenty species from seven genera of wild mushrooms in the Pacific Northwest, U.S.A., *J. Ethnopharmacol.* 5 (1982) 271–285, [https://doi.org/10.1016/0378-8741\(82\)90013-7](https://doi.org/10.1016/0378-8741(82)90013-7).
- [245] M. Wurst, R. Kysilka, M. Flieger, Psychoactive Tryptamines from Basidiomycetes, *Folia Microbiol.* 47 (2002) 3–27.
- [246] Z. Alam Mahmood, in: K.G. Ramawat, J.M. Merillon (Eds.), *Bioactive alkaloids from fungi: Psilocybin, Nat. Prod. Phytochem. Bot. Metab. Alkaloids Phenolics Terpenes*, Springer-Verlag Berlin Heidelberg, 2013, pp. 523–552, https://doi.org/10.1007/978-3-642-22144-6_19.
- [247] S. Ediriweera, R.L.C. Wijesundera, C. Nanayakkara, J. Weerasena, First Report of *Panaeolus sphinctrinus* and *Panaeolus foenicicii* (Psathyrellaceae, Agaricales) on Elephant Dung from Sri Lanka, *Front. Environ. Microbiol.* 1 (2015) 19–23, <https://doi.org/10.11648/j.fem.20150102.12>.
- [248] G. Guzmán, J. Ott, Description and Chemical Analysis of a New Species of Hallucinogenic *Psilocybe* from the Pacific Northwest, *Mycologia* 68 (1976) 1261–1267, <https://doi.org/10.1080/00275514.1976.12020019>.
- [249] L.-A. Cardon, *Panaeolus cinctulus* [Bolt.] Saccardo et *Panaeolus cyanescens* [Berk & Br.] Saccardo, deux champignons hallucinogènes présents en Normandie, 2014.
- [250] R.G. Benedict, Chemotaxonomic relationships among the basidiomycetes, *Adv. Appl. Microbiol.* 13 (1970) 1–23, [https://doi.org/10.1016/S0065-2164\(08\)70399-X](https://doi.org/10.1016/S0065-2164(08)70399-X).
- [251] J. Raitelhuber, *Panaeolus olivaceofuscus*, *Cat. Life Checkl* 2 (1977).
- [252] E. Ohenoja, J. Jokiranta, T. Marinen, A. Kaikkonen, M.M. Alraksinen, The occurrence of psilocybin and psilocin in Finnish fungi, *J. Nat. Prod.* 50 (1987) 741–744.
- [253] A.L. Christiansen, K.E. Rasmussen, Screening of hallucinogenic mushrooms with high-performance liquid chromatography and multiple detection, *J. Chromatogr. A* 270 (1983) 293–299, [https://doi.org/10.1016/S0021-9673\(01\)96375-7](https://doi.org/10.1016/S0021-9673(01)96375-7).
- [254] D.B. Repke, D.T. Leslie, G. Guzmán, Baecocystin in *Psilocybe*, *Conocybe* and *Panaeolus*, *Lloydia* 40 (1977) 566–578.
- [255] T. Stijve, A.A.R. de Meijer, Macromycetes from the state of Parama, Brazil - The psychoactive species, *Arq. Biol. Tecnol.* 36 (1993) 313–329.
- [256] Jochen Gartz, Magic mushrooms around the world: a scientific journey across cultures and time - the case for challenging research and value systems, *Aware J.* (1996) 1–136.
- [257] J. Gartz, G.K. Moller, Analysis and Cultivation of Fruit Bodies and Mycelia of *Psilocybe bohemica*, *Biochem. Physiol. Pflanz.* 184 (1989) 337–341, [https://doi.org/10.1016/S0015-3796\(89\)80023-x](https://doi.org/10.1016/S0015-3796(89)80023-x).
- [258] G. Guzmán, Index of taxa in the genus *Psilocybe*. Taxonomic studies on dark-spored agarics, *Mycotaxon* 6 (1978) 464–476.