Species

23(72), 2022

To Cite:

Mathew J, Pichan S. *Ixora lavanya* (Rubiaceae), a new species from south Western Ghats, India. *Species*, 2022, 23(72), 528-534

Author Affiliation:

¹Department of Botany, Sanatana Dharma College, Alappuzha, IN - 688003, Kerala, India

²Senior Technical Assistant, MS Swaminathan Research Foundation, Puthurvayal, Kalpetta, Wayanad, IN - 673577, Kerala, India

*Corresponding author

Department of Botany, Sanatana Dharma College, Alappuzha, IN - 688003, Kerala,

India

Email: polachirayan@yahoo.co.in

Peer-Review History

Received: 18 August 2022

Reviewed & Revised: 25/August/2022 to 18/October/2022

Accepted: 20 October 2022 Published: 23 October 2022

Peer-Review Model

External peer-review was done through double-blind method



© The Author(s) 2022. Open Access. This article is licensed under a Creative Commons Attribution License 4.0 (CC BY 4.0)., which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/.

Ixora lavanya (Rubiaceae), a new species from south Western Ghats, India

Jose Mathew^{1*}, Salim Pichan²

ABSTRACT

Ixora lavanya is described as a new species from Agasthyamala Hills in the Kerala part of south Western Ghats. The detailed descriptions and images of the new taxa along with the morphological keys and photographs of the indigenous *Ixora* species of Kerala are presented.

Keywords: Ixora species; Agasthyamala Hills; morphology

INTRODUCTION

Ixora Linnaeus is a large pantropical genus of Rubiaceae with more than 575 species worldwide (Mouly et al. 2009; POWO 2022), growing mostly in humid forests. It is a genus of shrubs and small trees with white, pink, orange or red entomophilous flowers, contorted corolla lobe aestivation, bilocular ovary, each locule with a single ovule attached peltately to the septum, and red or purple drupaceous fruits with usually two plano-convex pyrenes (Smith and Darwin, 1988; Burger & Taylor, 1993; De Block 1998; Moulyet al. 2009). In India, the genus is represented by 47 species (Husain & Paul, 1989), of which 22 native species, excluding ornamentals were reported from Kerala state (Sasidharan, 2013; Nayar et al., 2014; Balan & Shanmugham, 2016). Among them, two names viz., I. predeepii Balan & S.Harikr. and I. sivarajiana Pradeep. were reduced as synonyms of I. cuneifolia Roxb and I. notoniana Wall. ex G.Don respectively (Murugan et al. 2021).

Botanical exploration in the Agasthyamala Biosphere Reserve, during 2018-2021 has yielded some interesting specimens of Ixora. Critical analysis of the literature (Hooker, 1880; Gamble, 1921; Husain & Paul, 1984; 1989; Pradeep, 1997; Sasidharan 2013; Nayar et al., 2014; Balan & Shanmugham, 2016; Kottaimuthu, 2016; Gangopadhyay, 2017; Murugan et al. 2021), as well as from the scrutiny of vouchers deposited in BSI, CAL, CALI, DD, DEV, E, FRC, ICB, KFRI, MH, SKU, **TBGT** online (https://plants.jstor.org; https://www.ipni.org; https://www.tropicos.org; https://www.biodiversitylibrary.org and https://www.wcsp.science.kew.org) showed that this specimen is hitherto unknown to science and warrant to be a new taxa. This has resulted in the recognition of Ixora lavanya sp. nov.

Ixora lavanya J.Mathew & P.M.Salim, sp. nov. (Fig. 1)





Figure 1 *Ixora lavanya* J.Mathew & P.M.Salim; A–E. Flowering twigs from type locality; F. Corolla; G–H. Close-up images of stamen and stigma; I. Stipule

DIAGNOSIS

Ixora lavanya is morphologically similar to *Ixora lawsonii* Gamble in its terminal, congested, peduncled corymbose inflorescence with pink colour flowers, but differs mainly in having: lanceolate coriaceous leaves with redcolour margins in apical portions (vs. elliptic-ovate to obovate chartaceous leaves and without any markings in apical regions in *I. lawsonii*), glabrous calyx (vs. ciliate calyx in *I. lawsonii*); oblanceolate glabrascent corolla lobes (lanceolate pubescent corolla lobes in *I. lawsonii*) and beaked anthers (attenuate anther without any beak in *I. lawsonii*).




Figure 2 Diversity of indigenous *Ixora* species of Kerala; A. *I. acuminata*; B. *I. agasthyamalayana*; C. *I. brachiata*; D. *I. beddomei*; E. *I. coccinea*; F. *I. cuneifolia*; G. *I. elongata*; H. *I. finlaysoniana*; I. *I. gamblei*; J. *I. Johnsoni*; K. *I. pavetta*; L–N. *I. notoniana*

Type: India. Kerala, Thiruvananthapuram District, Peppara Wildlife Sanctuary, Way to Agasthyarkoodam, altitude 1720 m a. s. l., 14 April 2012, *J.Mathew 2127* (holotype: MH!; isotype: KUBH! (Kerala University Herbarium). – Paratype: Same locality, 27 April 2022, *J.Mathew 6033-6034* (KUBH!).

Since the herbarium of the Kerala University Herbarium, Thiruvanathapuram has not been formally registered with Index Herbariorum (Thiers [continuously updated]), it is here referred to as "KUBH".

Shrub, 1–1.5 m tall, profusely branched, glabrous; young internodes purple, older internodes grayish to brown. Stipule sheath subtruncate,9 mm long, awn 5–6 mm long. Leaves opposite with petioles 0.5–1 mm long, leaves lanceolate, 4–6× 1.5–2.3 cm, coriaceous, glabrous in both sides, apical portions with red colour margins in young leaves, apex narrowly acute to acuminate; base acute; secondary veins 6–7 pairs on each side. Inflorescence terminal, peduncle 2– 2.5 cm long, glabrous. Flowers pink, 4-merous; flower buds ellipsoid, pedicel bracteoles 3 – 3.5 mm long, lanceolate, acute, glabrous; calyx lobes glabrous, linear lanceolate, lobes 2.5 – 3 mm long; corolla tube 11-13.5 mm long, lobes oblanceolate, glabra scent, acute; Stamens inserted at corolla mouth, filaments 0.5–1 mm long, glabrous; anthers beaked, style 13-13.5 mm long, glabrous; stigmatic lobes 1–1.5 mm long. Fruits unknown.



Figure 3 Diversity of indigenous *Ixora* species of Kerala; A. *I. nigricans*; B. *I. lawsonii*; C. *I. thwaitesii*; D. *I. lanceolaria*; E. *I. polyantha*; F– G. *I. malabarica*; G. *I. undulata*; H. *I. ravikumarii*; I. *I. lavanya*

Flowering: April

Distribution and Habitat: Found in montane shola forests at altitudes of 1720 m, in Pongalappara regions of Agasthyarkoodam forests of Western Ghats, India. Common associates are *Canthium neilgherrense* and *Ochalandra travancorica*.

Etymology: The specific epithet 'lavanya' (beauty and grace in Indian languages) refers the beautiful flowers of the new species.

Conservation status: Only a single population with 4 plants of this species is known from which the type collection was made. The conservation status of this variety is proposed as Data Deficient. Further survey for this variety is suggested, which would need to be conducted in March to April, when the plants are flowering.

Key to native Ixora species of Kerala part of south Western Ghats

1. Calyx lobes longer than the tube
1: Calyx lobes equal or smaller than the tube
2: Calyx lobes lanceolate or linear lanceolate
3: Inflorescence glabrous
4: Calyx lobe linear, ciliate.125. Corymbs lax.6
5: Corymbs dense
6: Leaves elliptic-ovate or oblong
7: Flowers green to yellow
9: Corolla white
10: Leaves lanceolate
11: Leaves sub-membranous, corymbs sessile
12: Flowers pink, corolla tube 8 -12 mm long
13: Calyx lobe and tube are glabrous
14: Peduncle short or absent.1515. Corolla lobes glabrous.I. cuneifolia
15: Corolla lobes puberulous
16: Leaves flat on the margins.1817. Cymes sessileI. thwaitesii
17: Cymes peduncled

SPECIES I REPORT

18. Calyx-lobes triangular, acute	19
18: Calyx lobes much reduced, almost 0	20
19. Flowers white	I. nigricans
19: Flowers scarlet	I. coccinea
20. Leaves narrowed at base; flower buds globose	I. brachiata
20: Leaves round or cordate at base; flower buds ellipsoid	I. pavetta

ACKNOWLEDGEMENT

The authors are thankful to the Kerala Forest Department for permission to conduct research. The corresponding author is grateful to the Principal and Management of the Sanatana Dharma College, Alappuzha, Kerala for support; to Dr. K.N. Gandhi, Harvard University, USA for nomenclatural corrections; and to the Mohamed bin Zayed Species Conservation Fund (project 190521457); SERB (DST – INDIA) Project No. NPDF-PDF/2017/000135; DOEACC, Kerala (head no. 3435–04–104–98(P) –1) and NMPB (Project No. HG/KE–01/2019–2020) for financial assistance.

Funding

This study has been funded by Mohamed bin Zayed Species Conservation Fund (project 190521457); SERB (DST – INDIA) Project No. NPDF-PDF/2017/000135; DOEACC, Kerala (head no. 3435–04–104–98(P) –1) and NMPB (Project No. HG/KE–01/2019–2020).

Ethical issues

Ixora lavanya plants were used in this research and studied with the permission of Kerala Forest Department; Sanatana Dharma College, Alappuzha, Kerala for support; with help of Dr. K.N. Gandhi, Harvard University, USA for nomenclatural corrections.

Conflicts of interests

The authors declare that there are no conflicts of interest.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES AND NOTES

- 1. Balan AP and Shanmugham H. *Ixora pradeepii*, a new species of Rubiaceae from southern Western Ghats, India. Bangladesh Journal of Plant Taxonomy 2016; 23(1):65 69.
- 2. Burger WC and Taylor CM. Family 202 Rubiaceae. In: Burger WC ed. Flora costaricensis. *Fieldiana*: Botany, n.s. 1993; 33:1–333.
- 3. De Block P. The African species of *Ixora* (Rubiaceae Pavetteae). *Opera Botanica Belgica* 1998; 9:1–218.
- 4. Gamble JS. *The Flora of the Presidency of Madras, Vol.* 2. Adlard and Son Ltd., London 1921; 445.
- 5. Gangopadhyay M. Notes on Rubiaceae from India. *Nelumbo* 2017; 59(1):(23-24).
- 6. Hooker JD. *The Flora of British India, vol.* 3. Reeve and Co., London 1880; 137-149.
- 7. Husain T and Paul SR. A new species of *Ixora* (Rubiaceae) from the Andaman & Nicobar Islands *Blumea* 1984; 30:153–156.
- 8. Husain T and Paul SR. Taxonomic studies on Indian species of genus *Ixora* L. (Rubiaceae). *J Econ Tax Bot Additional series* 6. Scientific publishers, Jodhpur, India 1989.

- 9. Kottaimuthu R. *Ixora ravikumarii*, a new name for *Ixora monticola* Gamble (Rubiaceae: Ixoroideae). *Phytotaxa* 2016; 263(1):079–080.
- 10. Mouly A, Razafimandimbison S, Florence J, Jérémie J and Bremer B. Paraphyly of *Ixora* and new tribal delimitation of Ixoreae (Rubiaceae): inference from combined chloroplast (rps16, rbcL, and trnT-F) sequence data. Annals of Missouri Botanical Garden 2009; 96: 146–160. doi: 10.3417/2006194
- 11. Murugan P, Murugan C and Karthigeyan K. Notes on the identity and taxonomy of *Ixora cuneifolia* and *I. notoniana* and typification of three names of *Ixora*. *Gardens' Bulletin Singapore* 2021; 73(2):481-487.
- 12. Nayar TS, Rasiya Beegam A and Sibi M. Flowering *Plants of the Western Ghats, India.* Jawaharlal Nehru Tropical Botanic Garden and Research Institute, Palode, Thiruvananthapuram, Kerala. 2014.
- 13. POWO. Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew 2022.

- 14. Pradeep AK. *Ixora sivarajiana*, a new species of Rubiaceae from India. *Nordic Journal of Botany* 1997; 17:315–317. doi: 10.1111/j.1756-1051.1997.tb00319.x
- 15. Sasidharan N. *Flowering plants of Kerala: CD-ROM ver.* 2.0. (Kerala Forest Research Institute, Peechi, Kerala). 2013.
- 16. Smith AC and Darwin SP. *Ixora*. In: Smith AC, ed. *Flora Vitiensis Nova*. *Rubiaceae*, Vol. 4. Lawai: Pacific Tropical Botanical Garden 1988; 202–222.
- 17. Thiers B. continuously updated. Index *Herbariorum: A global directory of public herbaria and associated staff.* New York Botanical Garden's Virtual Herbarium. http://sweetgum.nybg.org/science/ih/ (accessed Jun 2018)