

ENTOLOMA SUBGENUS NOLANEA IN THE NETHERLANDS  
AND ADJACENT REGIONS  
WITH A RECONNAISSANCE OF ITS  
REMAINING TAXA IN EUROPE

M. E. NOORDELOOS

*Rijksherbarium, Leiden*

(With 49 Text-figures)

*Entoloma* subgenus *Nolanea* (emend. Romagnesi 1978) is revised on account of personal observations and studies on collections from various herbaria. The types of European taxa, as far as they could be recovered, have been examined. Observations on extralimital taxa are included. The infrageneric classification of Romagnesi (1978) is followed in broad outline with some slight alterations: sect. *Papillata* is emended by including sections *Minuta* Romagn. and *Cosmeoxonema* Largent & Thiers. One new section, viz. sect. *Fernandae*, and four new subsections, viz. subsect. *Tristia*, *Icterina*, *Infularia*, and *Cheilocystidiata* are introduced. 54 Taxa are recognized, five of which are new: *Entoloma sericeum* var. *cinereo-opacum*, *E. fernandae* f. *eccilioides*, *E. chlorophyllum*, *E. globulifer*, and *E. sphaerocystis*. Several new combinations are proposed, viz. *Entoloma ambrosium*, *E. clandestinum*, *E. conferendum*, *E. conferendum* var. *pusillum*, *E. cuspidifer*, *E. hirtipes* var. *sericoides*, *E. inutile*, *E. infula*, *E. sacchariolum*, *E. sericeoides*, *E. sericeonitens*, *E. sericeum* f. *nolaniforme*, *E. solstitiale*, *E. verecundum*, and *Entoloma* sect. *Staurospora*, sect. *Turfosa*, subsect. *Cosmeoxonema*, subsect. *Fibulata*, and subsect. *Minuta*. For nomenclatural reasons two new names are introduced, viz. *Entoloma leptopus* to replace *Nolanea tenuipes* P. D. Orton and *E. foetulentum* to replace *Nolanea foetida* Killermann. Keys, descriptions and illustrations are given to all species accepted. In an appendix doubtful and excluded species are briefly discussed.

CONTENTS

Introduction . . . . .	428
Material, methods and presentation. . . . .	428
Notes on the characters used in the keys . . . . .	429
Clamp-connections . . . . .	429
Pigmentation . . . . .	429
Acknowledgments . . . . .	430
Taxonomic part . . . . .	431
<i>Entoloma</i> subgenus <i>Nolanea</i> . . . . .	431
Key to the sections and subsections of subgenus <i>Nolanea</i> . . . . .	431
Key to the species of subgenus <i>Nolanea</i> . . . . .	432
Synoptical key to the species of subgenus <i>Nolanea</i> . . . . .	436
Section <i>Nolanea</i> . . . . .	438
Section <i>Staurospora</i> . . . . .	445
Section <i>Papillata</i> . . . . .	453
Subsections: <i>Papillata</i> , 453; <i>Fibulata</i> , 463; <i>Minuta</i> , 468; <i>Cosmeoxonema</i> , 472	
Section <i>Fernandae</i> . . . . .	486

Section <i>Endochromonema</i> . . . . .	494
Subsections: <i>Endochromonema</i> , 495; <i>Infularia</i> , 503; <i>Tristia</i> , 508; <i>Cheilocystidiata</i> , 510; <i>Icterina</i> , . . . . .	514
Extralimital species . . . . .	520
Insufficiently known and excluded taxa . . . . .	524
References . . . . .	530
Index . . . . .	531

## INTRODUCTION

Fries (1821: 10) erected *Agaricus* tribus *Nolanea* for those pink-spored agarics with a mycenoid habit. The concept of this taxon was generally accepted by later mycologists, though on different taxonomical levels: as a subgenus by Quélet (1886: 63) in *Rhodophyllus*, and by Schroeter (1889: 63) in *Hyporrhodius*, or as a genus by Kummer (1871: 24) and Saccardo (1887: 716).

The mycenoid habit appeared to be a morphological criterion difficult to maintain for modern mycologists. Romagnesi (1978) emended the concept of *Nolanea* on the base of microscopical characters such as shape of spores, type and topography of pigments, and presence and topography of clamp-connections. Kühner (1977: 450) suggested the use of size and shape of tramal elements as a distinctive infrageneric character, a suggestion worked out by me in the present study. On account of these characters a more natural classification appears to be possible. In the concept of Romagnesi followed by me, subgenus *Nolanea* now comprises 'classical' species such as *Entoloma cetratum* (Fr.) Moser, *E. conferendum* (Britz.) Noordeloos (= *Nolanea staurospora* Bres.), and *E. papillatum* (Bres.) Dennis as well as species with a more tricholomatoid habit formerly placed in subgenus *Entoloma*, such as *E. sericeum* (Bull. ex Mérat) Qué. and *E. ameiides* Berk. & Br., or with a collybioid habit, such as *E. icterinum*.

*Nolanea* is treated here as a subgenus in agreement with the classification of Romagnesi, as in my experience it is impossible to divide *Entoloma* sensu lato (or *Rhodophyllus*) into clear-cut genera. This is confirmed implicitly by the recent work on mostly extra-European taxa of *Entoloma* by Horak (1973, 1976 & 1978) and by Romagnesi & Gilles (1979). The criteria used by Largent & Benedict (1971: 35), such as the diameter of hyphae of the pileipellis, to distinguish genera cannot be maintained.

Subgenus *Nolanea* is closely related to subgenus *Entoloma* via sect. *Endochromonema* on one hand, and sect. *Turfosa* on the other, and with subgenus *Pouzaromyces* via sect. *Fernandae*. For further details and schemes of classification the reader is referred to a future paper (Noordeloos, 1980a; in preparation).

## MATERIAL, METHODS AND PRESENTATION

Most species treated here have been studied by me in fresh as well as in dried condition. Macroscopic descriptions are thus in most cases based on personal observations completed by notes in descriptions of others. Many of the latter are supplied by Dr. C. Bas and Dr. R. A. Maas Geesteranus, who laid the base of a rather well documented set of Agaricales collections in the Rijksherbarium during the past 25 years. Unless otherwise stated, the colours of fresh specimens are compared with Munsell Soil Colour Charts, Baltimore.

Microscopical structures were observed and measured in water (fresh carpophores) or in a 10% ammonia solution, or in an ammoniac 1% Congo Red solution (dried specimens), usually under oil-immersion.

Spores, basidia and cystidia were observed and measured in squash preparations of minute parts of the lamellae. Hymenophoral trama has been examined in transversal sections as well as in squash preparations. The pigmentation of the upper layers of the pileus has been observed preferably after plasmolysis in radial sections of fresh carpophores. (For obtaining plasmolysis a saturated salt solution has been used.) In dried specimens it has been observed after boiling sections in ammonia solution or with the method described by Kühner & Romagnesi (1953: 177) after treatment with picroformol de Hollande observed in boiling Chloralhydrate. Part of the collections have been fixed in Carnoy solution and conserved in 70% alcohol. This material appeared to be very suitable for the study of pigmentation patterns too. However, observations on fresh material are preferable because they give the most reliable results, particularly in cases of pale, diffusely intracellular pigments.

Drawings were made with the aid of a drawing prisma. The magnifications of the figures are: carpophores, natural size; spores,  $\times 1000$ ; all other microscopical details,  $\times 670$ .

Unless otherwise stated all collections studied are deposited in the Rijksherbarium, Leiden (L).

#### NOTES ON THE CHARACTERS USED IN THE KEYS

**CLAMP-CONNECTIONS.**—To have a criterion for the presence or absence of clamp-connections, the hymenium was selected. In most species of *Nolanea* clamp-connections are abundant in the hymenium, but only very sparse in other tissues. The best way to study clamp-connections is to stain them in ammoniac 1% Congo Red. One should be careful of proliferating clamps (see Bas, 1965: 355) and look preferably at the base of young basidia.

**PIGMENTATION.**—The type of pigmentation and its topography is considered a character of great importance. Three main types of pigmentation are to be distinguished, viz.:

(i). **Intracellular.**—Pigment present in plasma and/or vacuoles inside the hyphal elements. This pigment can be diffuse or present itself as smaller or larger granules or clots, which sometimes agglutinate. In plasmolysis this type of pigmentation becomes more distinct at the moment the plasma retires from the hyphal walls in one or more coloured globules, leaving a non-coloured zone between them and the hyphal wall.

(ii). **Membranal.**—Pigment present in the hyphal wall. In plasmolysis there is no difference to be seen in the location of the pigmentation in comparison to observations on the pigments in water.

(iii). **Encrusting.**—Pigments present on the outer wall of the hyphal elements in the form of minute to rather large crust-like patches, which sometimes form distinct patterns (e.g. a 'zebra' pattern). This type of pigmentation is usually easy to recognize also in dried collections. In some cases the encrustations are very minute and inconspicuous and should be looked for very carefully, particularly on the narrowest hyphae of pileipellis and pileitrama, and near the septa.

Combinations of different types of pigmentation frequently occur, particularly (i) and (ii), or (i) and (iii).

The following abbreviations are used:

Q.—Length-width ratio, usually given as follows: Q = 1.2–1.3–1.4, which means Q = between 1.2 and 1.4, with an average of 1.3. (The length of a spore is measured without the apiculus.)

L–D = 1–2–3  $\mu\text{m}$ .—Length minus width between 1 and 3, with an average of 2  $\mu\text{m}$ .

Lamellae L = 20–25, l = 1–3(–5).—20 to 25 entire lamellae per carpophore with 1–3, sometimes 5 short lamellae ('lamellulae') between each pair.

#### ACKNOWLEDGMENTS

I am deeply indebted to Dr. C. Bas for his guidance and encouragement during my studies in *Entoloma* which resulted in the present paper. Sincere thanks are due to Prof. H. Romagnesi, Paris, for valuable discussions, and for making available his rich herbarium and collection of unpublished notes. Dr. E. Kits van Waveren, Amsterdam, Dr. F. Tjallingii & Mrs. G. J. M. G. Tjallingii-Beukers, Wageningen, and Mr. P. B. Jansen, Breda, kindly put their herbaria at my disposal, for which I am very grateful. Many members of the Netherlands' Mycological Society (N.M.V.) supplied me with fresh and dried material. Without their help the present revision would have been very incomplete. Grateful acknowledgments are made to the directors and/or keepers of the following herbaria for the loan of types and other critical collections: Bündner Naturhistorisches und Nationalpark Museum (CHUR); Nationale Plantentuin van België, Meise (BR); Botanical Museum and Herbarium, Copenhagen (C); Botanical Museum, Helsinki (H); Royal Botanic Garden, Edinburgh (E); Royal Botanic Gardens, Kew (K); Botanische Staatssammlung, München (M); Botanical Museum, Oslo (O); Laboratoire de Cryptogamie, Muséum de l'Histoire Naturelle, Paris (PC); National Museum, Prague (PRM); Botanical Department of Charles University, Prague (PRC); Botanical Museum, Stockholm (S); Institute for Systematic Botany, Uppsala (UPS), and the herbarium of the Biological Station Dr. W. Beyerinck, Wijster, Netherlands (WBS). I am indebted to the directors of the following institutions for providing working facilities: The Herbarium, Royal Botanic Gardens, Kew; Laboratoire de Cryptogamie, Paris; Botanische Staatssammlung, München; Botanical Museum, Oslo; National Museum, Prague, and Department of Botany of Charles University, Prague. Special help was received from Prof. Dr. J. J. Barkman and Eef Arnolds, Wijster; Dr. H. Knudsen, Copenhagen; Prof. Dr. H. Hertel, München; Dr. D. N. Pegler, Kew; Dr. R. Watling, Edinburgh; M<sup>me</sup> Dr. J. Perreau, Paris; Konservator A. E. Torkelsen, Oslo; Dr. J. Klán, Dr. Z. Pouzar and Dr. M. Svrček, Prague. I am indebted to Dr. R. A. Maas Geesteranus, Oegstgeest, for correcting the Latin diagnoses. Ruth van Crevel, Leiden, spent a lot of her time and energy arranging the plates and preparing them for printing, for which I am very grateful. Eva van Santen, Leiden, is thanked for her improvements of the English text. Mrs. A. Pots-Hageman is gratefully thanked for the accurate way she typed out the manuscript and searched for errors in the text.

## TAXONOMIC PART

**Entoloma** (Fr.) Kumm. *emend.* Donk  
subgenus **Nolanea** (Fr.) Noordeloos, *comb. nov.*

*Agaricus* L. ex Fr. trib. *Nolanea* Fr., Syst. mycol. 1: 10. 1821 (basionym). — *Agaricus* L. ex Fr. subgen. *Nolanea* (Fr.) Loud., Encl. Pl.: 998. 1829. — *Nolanea* (Fr.) Kumm., Führ. Pilzk.: 24. 1871. — *Rhodophyllus* Quéf. subgen. *Nolanea* (Fr.) Quéf., Enchir.: 63. 1886. — *Hyporrhodius* (Fr.) Schroet. subgen. *Nolanea* (Fr.) Schroet. in Cohn, KryptogFl. Schles. 3(1): 613. 1889. — Lectotype (Largent, 1974): *Agaricus hirtipes* Schum. ex Fr.

*Latzinea* O.K., Rev. Gen. Pl. 2: 857. 1891. — Holotype: *Agaricus pasceus* Pers. ex Fr.

*Lanolea* Nieuwl. in Amer. Midl. Nat. 4: 381. 1916. — Holotype: *Agaricus pasceus* Pers. ex Fr.

*Arenicola* Velen., Novitates mycologicae novissimae: 62. 1947. — Holotype: *A. flavispora* Velen.

Carpophores mycenoid, rarely tricholomatoid or omphalioid. Pileus conical to campanulate or semiglobose, generally finally expanding to convex rarely plano-convex, usually papillate or umbonate, rarely truncate or umbilicate, hygrophanous (but sometimes only very weakly), when moist usually translucently striate at least at margin, surface glabrous and smooth or minutely radially lustrously fibrillous with aeriferous fibrils or sometimes rugulose to minutely fluffy-scaly at centre. Lamellae narrowly adnate to almost free or emarginate, rarely broadly adnexed to subdecurrent, frequently with distinct grey or brown tinge when mature. Stipe slender, often more or less cartilaginous and easily snapping across but in some taxa fibrous and easily splitting lengthwise. Smell and taste often farinaceous-rancid.

Spores iso- to heterodiametrical, with basal facet or dihedral base, in extra-European (tropical) taxa often prismatical or cuboid. Cheilocystidia present or not. Pleurocystidia usually lacking. Hymenophoral trama regular, usually composed of long cylindrical cells attenuated towards septa ('fusiform'), often up to 450  $\mu\text{m}$  long or more. Pileipellis two-layered; suprapellis a simple cutis made up of radially arranged cylindrical hyphae, often with (clavate) terminal and sometimes ascending cells, thus forming transitions to a trichodermium; subpellis of short inflated cells well developed or not. Pileitrampa regular, in general composed of the same type of cells as the hymenophoral trama. Pigmentation variable: membranous, encrusting or intracellular or combinations of different types. Clamp-connections present or not, if present then usually abundant in hymenium and elsewhere rare or lacking.

KEY TO THE SECTIONS AND SUBSECTIONS  
OF SUBGENUS NOLANEA

- |   |   |
|---|---|
| 1a. Spores cuboid or prismatical (cruciform-stellate).  | Sect. <i>Staurospora</i> , p. 445         |
| b. Spores otherwise: (sub-)isodiametrical and 5-6-angled in side-view or distinctly heterodiametrical.  | 2   |
| 2a. Cheilocystidia present.   | 3   |
| b. Cheilocystidia absent.   | 7   |
| 3a. Lamellae olivaceous-greenish.   | Subject. <i>Icterina</i> , p. 514         |
| b. Lamellae never olivaceous-greenish.  | 4   |
| 4a. Pigment minutely encrusting hyphae of pileipellis and pileitrampa and simultaneously intracellular, particularly in hypoderm; cheilocystidia cylindrical-subcapitate, spores heterodiametrical. | Sect. <i>Nolanea</i> , p. 438             |
| b. Pigment either exclusively intracellular or exclusively membranous and encrusting.   | 5   |
| 5a. Pigment intracellular.  | Subject. <i>Cheilocystidiata</i> , p. 510 |
| b. Pigment membranous and encrusting.   | 6   |

- 6a. Cheilocystidia tibiiform or spheropedunculate; spores with dihedral base. . . . . Subject. *Cosmeoxonema*, p. 472
- b. Cheilocystidia (sub-)cylindrical, sometimes subcapitate; spores with basal facet. . . . . Subject. *Papillata*, p. 453
- 7a. Clamp-connections present, rarely absent but then pigment never encrusting. . . . . 8
- b. Clamp-connections absent; pigment of two types: coarsely encrusting and granular-intracellular. . . . . Sect. *Fernandae*, p. 486
- 8a. Pigment minutely to coarsely encrusting hyphae of pileipellis and pileitrama, sometimes in addition intracellular, particularly in hypoderm. . . . . Sect. *Papillata* 9
- b. Pigment diffuse and intracellular, sometimes in addition membranal, never encrusting. . . . . Sect. *Endochromonema* 12
- 9a. Spores isodiametrical,  $Q = 1.0-1.1-1.2$ . . . . . 10
- b. Spores heterodiametrical,  $Q \geq 1.2$  on the average per collection. . . . . 11
- 10a. Pigment exclusively and coarsely encrusting hyphae of pileipellis and pileitrama. . . . . Subject. *Cosmeoxonema*, p. 472
- b. Pigment minutely encrusting as well as intracellular, particularly in hypoderm. . . . . Subject. *Fibulata*, p. 463
- 11a. Carpophores pale; lamellae white then pale pink; pileus often variable in shape from conical to plano-convex, with papilla or central depression; spores with distinct dihedral base. . . . . Subject. *Minuta*, p. 468
- b. Carpophores moderately to strongly pigmented; lamellae always with distinct grey or brown tinge; pileus usually papillate; spores usually with basal facet. . . . . Subject. *Papillata*, p. 453
- 12a. Stipe smooth, glabrous, as if polished. . . . . 13
- b. Stipe fibrillose, often striate. . . . . 14
- 13a. Pileus blackish brown; spores thin-walled, multiangular-gibbose in outline. Subject. *Tristia*, p. 508
- b. Pileus moderately dark brown; spores regularly 5-6-7-angled in outline with pronounced angles. . . . . Subject. *Infularia*, p. 503
- 14a. Pileus and stipe with lemon-yellow tinges and/or flesh with strong aromatic smell like orange blossom. . . . . Subject. *Icterina*, p. 514
- b. Pileus and stipe never with lemon-yellow tinges, only sometimes yellow-ochraceous; smell never aromatic. . . . . Subject. *Endochromonema*, p. 495

## KEY TO THE SPECIES OF SUBGENUS NOLANEA

- 1a. Basidia in majority 2-spored. . . . . 2
- b. Basidia in majority 4-spored. . . . . 4
- 2a. Clamp-connections absent; pigment membranal and intracellular; stipe without differentiated hairs. . . . . 3
- b. Clamp-connections present; pigment minutely encrusting; stipe with capitate hairs, particularly at apex. . . . . *E. cuspidifer*, p. 461
- 3a. Carpophores mycenoid with conical, then slightly expanding pileus, usually with distinct umbo; stipe silvery striate; taste absent or slightly farinaceous. . . . . *E. cetratum*, p. 496
- b. Carpophores small and thick-set; pileus soon expanding, with or without a slight umbo, sometimes even slightly depressed; stipe polished; taste strongly rancid-fishy. . . . . *E. farinogustus*, p. 497
- 4a. Spores cuboid or cruciform-stellate (see Figs. 4, 5, 6). . . . . 5
- b. Spores otherwise: (sub-)isodiametrically 5-6-angled or distinctly heterodiametrical in side-view. . . . . 9
- 5a. Spores cruciform-stellate. . . . . 6
- b. Spores cuboid. . . . . 7
- 6a. Carpophores small and slender; pileus up to 15 mm broad; stipe 1-1.5 mm thick; on decaying trunks of frondose trees. . . . . *E. conferendum* var. *pusillum*, p. 450
- b. Carpophores more robust and terrestrial. . . . . *E. conferendum* var. *conferendum*, p. 446

- 7a. Spores 10.2–12.4  $\mu\text{m}$  long; pileus blackish brown, strongly contrasting with pale stipe. . . . . *E. inodorum*, p. 520
- b. Spores up to 10  $\mu\text{m}$  long; pileus and stipe concolorous and paler. . . . . 8
- 8a. Stipe minutely pruinose at least at upper half, with large fusiform caulocystidia; pigment membranal, not encrusting. . . . . *E. prismatospermum*, p. 523
- b. Stipe smooth, polished, without caulocystidia; pigment intracellular, sometimes in addition encrusting. . . . . *E. rhombisporum*, p. 450
- 9a. Spores isodiametrical, 5–6-angled in side-view,  $Q=1.0-1.1-1.2$ . . . . . 10
- b. Spores heterodiametrical,  $Q \geq 1.2$  on the average per collection. . . . . 18
- 10a. Cheilocystidia present, subcylindrical to tibiiform. . . . . *E. tibilocystidiatum*, p. 473
- b. Cheilocystidia absent. . . . . 11
- 11a. Pigment membranal, not encrusting. . . . . *E. occultopigmentatum*, p. 501
- b. Pigment minutely to coarsely encrusting, sometimes in addition intracellular. . . . . 12
- 12a. Pigment minutely encrusting and in addition also intracellular, particularly in subpellis; in forests, rarely also in grassland. . . . . 13
- b. Pigment moderately to coarsely encrusting; no trace of intracellular pigment; in grassland. . . . . 14
- 13a. Pileus moderately dark grey-brown, often with ochraceous tinge, already when moist covered with loose aëriferous fibrils, strongly lustrous when dry; stipe strongly silvery striate; smell weak, not farinaceous. . . . . *E. nitens*, p. 466
- b. Pileus fairly dark grey-brown, smooth; stipe smooth or sparsely striate; smell strongly farinaceous. . . . . *E. juncinum*, p. 464
- 14a. Pileus very obscurely striate when moist, very dark grey-brown, surface hoary with scattered silvery-fibrillous patches; lamellae dark grey (species reminding of *Tephrocycbe* spp.). . . . . *E. sericeum* var. *cinereo-opacum*, p. 482
- b. Pileus distinctly striate when moist, dark sepia or reddish brown. . . . . 15
- 15a. Pileus relatively thick-fleshed with faint umbo or central depression; stipe usually as long as diameter of pileus. . . . . *E. sericeum* var. *sericeum* f. *sericeum*, p. 479
- b. Pileus relatively thin-membranaceous; stipe considerably longer than diameter of pileus. . . . . 16
- 16a. Pileus distinctly umbilicate. . . . . *E. sericeoides*, p. 483
- b. Pileus usually distinctly papillate. . . . . 17
- 17a. Spores 7.4–8.3(–9.3)  $\times$  5.7–7.4  $\mu\text{m}$ ,  $Q=1.1-1.2-1.3$ , only very weakly angled, subglobose-ellipsoid in outline. . . . . *E. ortonii*, p. 459
- b. Spores 7.4–10.4  $\times$  (6.4–)6.7–8.7(–9.3)  $\mu\text{m}$ ,  $Q=1.0-1.1-1.2$ , distinctly angled in side-view. . . . . *E. sericeum* var. *sericeum* f. *nolaniforme*, p. 480
- 18a. Cheilocystidia present. . . . . 19
- b. Cheilocystidia absent. . . . . 30
- 19a. Lamellae olivaceous-greenish. . . . . *E. chlorophyllum*, p. 518
- b. Lamellae without any olivaceous or greenish tinge. . . . . 20
- 20a. Pileus very pale grey when moist, on drying pallescent to white, no pigment present; stipe white. . . . . *E. nivescens*, p. 524
- b. Pileus and stipe distinctly pigmented. . . . . 21
- 21a. Cheilocystidia globuliform or spheropedunculate. . . . . 22
- b. Cheilocystidia differently shaped. . . . . 23
- 22a. Clamp-connections absent; pigment intracellular. . . . . *E. globulifer*, p. 513
- b. Clamp-connections present; pigment minutely encrusting. . . . . *E. sphaerocystis*, p. 485
- 23a. Pigment intracellular. . . . . 24
- b. Pigment minutely to coarsely encrusting, sometimes particularly in subpellis in addition also intracellular. . . . . 27
- 24a. Lamellae very dark brown; cheilocystidia in majority lecitiform to tibiiform. . . . . *E. inutile*, p. 512
- b. Lamellae never dark brown; cheilocystidia differently shaped. . . . . 25
- 25a. Cheilocystidia cylindrical, sometimes subcapitate, up to 40  $\mu\text{m}$  long. . . . . *E. cryptocystidiatum*, p. 511
- b. Cheilocystidia fusiform to slenderly lecitiform, usually considerably longer than 40  $\mu\text{m}$ . . . . . 26

- 26a. Spores (10.0-)10.4-12.0  $\mu\text{m}$  long; cheilocystidia 42-90(-120)  $\times$  7-14(-20)  $\mu\text{m}$ .  
*E. velenovskyi* var. *velenovskiyi*, p. 510
- b. Spores (10.1-)11-15.9(-16.3)  $\mu\text{m}$  long; cheilocystidia (50-)70-149  $\times$  (6.2-)7.4-15.9  $\mu\text{m}$ .  
*E. velenovskyi* var. *longicystidium*, p. 511
- 27a. Pigment minutely encrusting; smell aromatic-fruity or like that of orange blossom; in grasslands.  
*E. sacchariolenis*, p. 474
- b. Pigment minutely encrusting and in addition particularly in subpellis intracellular; smell farinaceous, rarely absent, never aromatic-fruity; in forests or grasslands. . . . . 28
- 28a. Spores (8.1-)8.5-12(-13.9)  $\times$  (5.2-)5.8-7.5(-8.1)  $\mu\text{m}$ ; stipe long and slender, much longer than diameter of pileus, 1-3(-5) mm thick; in forests, autumn. . . . . *E. leptopus*, p. 442
- b. Spores 8.0-9.3(-9.5)  $\mu\text{m}$  wide; stipe considerably thicker, 3-7(-10) mm. . . . . 29
- 29a. Stipe much longer than diameter of pileus; pileus conical, only slightly expanding; in coniferous forests in spring, also in dune valleys and on grassy, open places in mixed forest in late autumn.  
*E. hirtipes* var. *hirtipes*, p. 438
- b. Stipe about as long as diameter of pileus; pileus more or less flattened; in grasslands.  
*E. hirtipes* var. *sericoides*, p. 442
- 30a. Pigment intracellular, sometimes in addition membranous but never encrusting. . . . . 31
- b. Pigment minutely to coarsely encrusting, sometimes in addition diffusely or granularly intracellular. . . . . 43
- 31a. Pileus with yellowish or brownish olivaceous tinge; often with strong, aromatic smell like that of *Hebeloma sacchariolenis* or *Entoloma ameides*, like fruit, etc. . . . . 32
- b. Pileus never with yellow-olivaceous or brown-olivaceous tinges; smell never aromatic like that of *Hebeloma sacchariolenis*, etc. . . . . 33
- 32a. Pileus with yellow-olivaceous tinge. . . . . *E. icterinum*, p. 516
- b. Pileus with brown-olivaceous tinge. . . . . *E. ambrosium*, p. 520
- 33a. Spores 5-7  $\mu\text{m}$  wide. . . . . 34
- b. Spores 7-9.5  $\mu\text{m}$  wide. . . . . 37
- 34a. Spores 6-8  $\times$  5-7  $\mu\text{m}$ , Q = 1.15-1.2 on the average per collection, rather thin-walled, multiangular in side-view, slightly cyanophilous. . . . . *E. vinaceum* and varieties, see excluded species, p. 529
- b. Spores 7-10  $\times$  5-7  $\mu\text{m}$ , Q = 1.3 on the average per collection, 5-7(-9)-angled in side-view, with relatively thick, non-cyanophilous walls. . . . . 35
- 35a. Pileus blackish brown; spores irregularly 6-9-angled-nodulose in side-view.  
*E. undulatosporum*, p. 509
- b. Pileus moderately dark brown, reddish brown or yellowish brown; spores rather regularly 5-6(-7)-angled in side-view. . . . . 36
- 36a. Stipe grey-brown to blackish brown; pileus moderately dark brown, often with ochraceous tinge; smell absent.  
*E. solstitiale*, p. 505
- b. Stipe and pileus yellowish to reddish brown; smell strongly like that of chlorine.  
*E. chlorinosum*, p. 507
- 37a. Stipe smooth, polished. . . . . 38
- b. Stipe striate, with (aeriferous) silvery white fibrils. . . . . 39
- 38a. Stipe yellow; pileus pinkish brown or with reddish tinge. . . . . *E. verecundum*, p. 507
- b. Stipe (dark) horn brown; pileus blackish brown. . . . . *E. triste*, p. 509
- 39a. Spores 8.3-11  $\mu\text{m}$  long, Q = 1.1-1.25-1.4; in grasslands. . . . . 40
- b. Spores 10-12  $\mu\text{m}$  long, Q = 1.25-1.4-1.5; in forests. . . . . 41
- 40a. Pileus pale grey-brown with slight ochraceous tinge, with dark brown striae up to centre; lamellae grey, then brown-pink, thickish, broadly ventricose, exceeding the pileus. . . . . *E. ventricosum*, p. 502
- b. Pileus pale orange-brown with grey-brown striae up to centre; lamellae pale, then pink without any grey or brown tinge, of normal thickness, not exceeding the pileus. . . . . *E. cathionis*, p. 502
- 41a. Stipe entirely downy pruinose with cylindrical, sometimes subcapitate hairs; pileus and stipe very pale brown with yellow or grey tinge. . . . . *E. lanuginosipes*, p. 499
- b. Stipe smooth or pruinose at apex or upper half only, rarely entirely pruinose, then pileus dark brown or date brown at least at limb. . . . . 42



- 42a. Pileus bicolorous; limb sepiceous or date brown, centre yellowish; lamellae with ochraceous tinge. *E. cuneatum*, p. 498  
 b. Pileus unicolorous, reddish brown; lamellae pale when young, becoming pinkish brown with age. *E. pallescens*, p. 521
- 43a. Pigment of two types: (coarsely) encrusting the hyphae of pileipellis and pileitrama, in addition intracellular in form of granules or clots in hyphae of pileipellis and upper pileitrama; pileipellis a cutis, often with numerous, ascending terminal cells, forming a transition to a trichodermium, particularly at centre of pileus; clamps absent. . . . . 44  
 b. Pigment encrusting, exceptionally in addition with pale, diffusely intracellular pigment; pileipellis usually a simple cutis, occasionally with tufts of ascending hyphae at centre of pileus; clamps present. . . . . 51
- 44a. Stipe strongly silvery striate; pileus smooth, when moist translucently striate up to centre. . . . . 45  
 b. Stipe not striate but smooth or downy-pruinose, and then pileus rugulose-felted to subsquamulose, at least at centre, and striate at margin only. . . . . 46
- 45a. Pileus grey-brown with yellow or russet tinge, pale greyish ochraceous when dry; lamellae white, then pink without any brown or grey tinge; flesh rather brittle in pileus; spores  $7.9-10.3(-10.8) \times (6.3-6.7-7.7(-8.0)) \mu\text{m}$ ; 2- and 4-spored basidia equally distributed in hymenium. . . . . *E. acidophilum*, p. 489  
 b. Pileus rather dark grey-brown, brown-grey on drying; lamellae pale brown-grey, then flesh-coloured pink; flesh in pileus more or less firm; spores  $7.4-9.5(-10.6) \times 5.8-6.9(-7.4) \mu\text{m}$ ; basidia 4-spored. *E. argenteostriatum*, p. 490
- 46a. Spores rather strongly rounded and weakly angular in side-view,  $(6.5-6.7-7.9(-8.1)) \times (5.3-5.8-6.9(-7.4)) \mu\text{m}$ ,  $Q = 1.1-1.2-1.3$ . . . . . *E. defibulatum*, p. 492  
 b. Spores larger and more pronouncedly angular in side-view,  $Q \geq 1.25$  on the average per collection. . . . . 47
- 47a. Stipe with distinct yellow tinge, quite different from colour of pileus. . . . . *E. xanthocaulon*, p. 493  
 b. Stipe greyish-brownish, more or less with the same colour as the pileus, but usually paler. . . . . 48
- 48a. Spores  $(8.4-9.0-11.4(-12.5)) \times (6.2-6.9-7.7(-8.2)) \mu\text{m}$ ,  $Q = 1.3-1.45-1.7(-1.8)$ . *E. cuniculorum*, p. 490  
 b. Spores not longer than  $10.5 \mu\text{m}$ , and  $Q = 1.1-1.3-1.4$ . . . . . 49
- 49a. Pileus when young more or less conical, quickly expanding to convex or flattened, sometimes slightly depressed at centre, sometimes with small umbo, pale to moderately dark grey-brown; in grasslands. . . . . 50  
 b. Pileus acutely conical, not or only slightly expanding; blackish brown; gregarious in coniferous forests. *E. fractum*, p. 492
- 50a. Pileus strongly hygrophanous, when moist translucently striate up to centre, absolutely smooth and glabrous; stipe brilliant, smooth and glabrous. . . . . *E. psilopus*, p. 489  
 b. Pileus only weakly hygrophanous, when moist translucently striate at margin only, rugulose-fluffy or subsquamulose particularly at centre; stipe smooth, glabrous or minutely downy-woolly all over. *E. fernandae*, p. 487
- 51a. Smell strong, aromatic, like that of amylicetate or fruit, also reminding of *Hebeloma sacchariolenis*; pileus and stipe pale ochraceous grey or brownish grey. . . . . *E. ameides*, p. 473  
 b. Smell absent or farinaceous-rancid. . . . . 52
- 52a. Pileus and stipe pale beige to pale brown; carpophores tiny and brittle; smell absent; on marshy places. . . . . 53  
 b. Pileus and stipe moderately to fairly dark brown, sepia, reddish brown or yellowish brown; in grasslands and smell farinaceous, or in forests and smell absent. . . . . 54
- 53a. Pileus usually (slightly) depressed at centre and with conspicuous dark spot, rarely with small papilla; lamellae pale pink; stipe glabrous, without differentiated hairs; spores  $(7.9-8.0-10.4(-11.5)) \times (6.4-6.8-8.1(-8.7)) \mu\text{m}$ . . . . . *E. minutum*, p. 468  
 b. Pileus convex with or without small papilla, never depressed; lamellae with distinct grey-brown colour; apex of stipe with well differentiated, subcapitate hairs; spores  $9.3-12(-14) \times 7.2-9.3 \mu\text{m}$ . *E. tenellum*, p. 470

- 54a. Spores very obtusely angled, more or less rounded in outline,  $7.4-8.3(-9.4) \times 5.7-7.4(-8.0) \mu\text{m}$ ,  $Q = 1.1-1.2-1.3$ ; stipe strongly silvery striate; smell farinaceous. . . . . *E. ortonii*, p. 459
- b. Spores more pronouncedly angled and slightly to considerably larger; stipe smooth or sparsely silvery striate. . . . . 55
- 55a. Spores  $7.9-10.4 \times 6.6-8.6(-9.0) \mu\text{m}$ ,  $Q = 1.1-1.2-1.3$ ; lamellae never dark brown; stipe sparsely but distinctly silvery striate; smell farinaceous. . . . . *E. lucidum*, p. 458
- b. Spores with average  $Q$  per collection at least 1.25 and usually larger; if spores of same size then lamellae fairly dark brown, stipe never striate and smell absent. . . . . 56
- 56a. Pileus moderately dark reddish brown or ochraceous brown; lamellae pale, then pink, sometimes with brown or grey tinge; hyphae of hymenophoral trama never encrusted; smell absent. . . . . 57
- b. Pileus very dark sepia, reddish brown or blackish brown; lamellae very dark brown when mature; usually the narrowest hyphae of the hymenophoral trama encrusted; smell farinaceous or absent. . . . . 58
- 57a. Lamellae with brown tinge when mature; spores  $7.0-8.7 \mu\text{m}$  wide; stipe sparsely silvery striate. . . . . *E. sericeonitens*, p. 459
- b. Lamellae white, then pink without brown or grey tinges; spores  $5.8-7 \mu\text{m}$  wide; stipe smooth, polished. . . . . *E. infula*, p. 503
- 58a. Smell farinaceous-rancid; lamellae pale when young but soon very dark brown with pink tinge; stipe often paler than pileus and with some scattered silvery fibrils or smooth, polished; spores  $(9.0-9.3-11.3(-13) \times (6.7-7.1-8.0(-8.5) \mu\text{m}$ ,  $Q = 1.2-1.3-1.5(-1.6)$ , with basal facet; in grasslands. . . . . *E. papillatum*, p. 454
- b. Smell absent. . . . . 59
- 59a. Strictly vernal, appearing from March to the end of May; pileus moderately dark brown to very dark sepia; lamellae of normal thickness, not very much distant, paler than pileus; apex of stipe with well differentiated, cylindrical, sometimes subcapitate hairs; spores  $(8.1-9.0-11.5(-12.0) \times 7.0-9.3 \mu\text{m}$ ,  $Q = 1.1-1.25-1.4$ , with dihedral base; in or near coniferous forests. . . . . *E. vernum*, p. 476
- b. Appearing in late summer and autumn; pileus and stipe concolorous, very dark, almost blackish brown, often with reddish tinge; lamellae very distant and thickish, brown already when young, when mature almost as dark as pileus; stipe smooth, polished, without differentiated hairs; spores  $(7.8-8.2-10.2(-10.6) \times (5.7-6.2-7.5(-7.9) \mu\text{m}$ ,  $Q = 1.2-1.3-1.4(-1.5)$ , with basal facet; in clearings in frondose forests, rarely also in grasslands. . . . . *E. clandestinum*, p. 456

## SYNOPTICAL KEY TO THE SPECIES OF SUBGENUS NOLANEA

(numbers refer to those in the Synopsis on p. 439;  
numbers in brackets means: characters present or not)

## Pileus

## colour

- when moist very pale grey, becoming (almost) white on drying: (15), (18), (38), 59  
with distinct olivaceous yellow or olivaceous brown tinge: 53, 54, 55  
date brown with yellow papilla: 37

## surface

- hardly hygrophanous, not striate: 26, (27)  
with slightly rugulose or fluffy centre: (3), (4), (5), (13), (14), (19), (21), (23), (49), (51), (53), (54)  
centre minutely squamulose: 26, 27

## shape

- (acutely) conical, not or only slightly expanding: (1), 7, (10), 12, (16), 18, 20, 34, 35, (37), 43, (44), (45), 48, 57  
with (slight) central depression or umbilicate: (6), (15), (21), 24, (26), 27, (36), (38), (39), (51), (53), (54)

**Lamellae**

## colour

with greenish or olivaceous tinge: (53), 54

white, then pink without any brown or grey tinge: (6), 15, 29, (34), (36), 40, 42, 43, 44, 45, (53), 56, 59

very dark brown or grey-brown, only slightly tinged pink: 7, 8, (13), (21), (22), 23, 51, (57)

## shape

with decurrent tooth: (15), 24, (26), 27, (53), (54)

thickish: 8, 41, 54

broadly ventricose, exceeding pileus: (4), (6), 12, (13), 15, (16), (22), (39), 41

**Stipe**

## colour

white: 56, 59

yellow or at least predominantly yellow: 5, 6, (15), (16), (18), 34, (35), (36), 38, (40), 45, (53)

## surface

smooth, as if polished: 6, (7), 8, (26), 27, 28, 31, (34), 36, 42, 43, 44, 45, 47

**Smell**

like chlorine: 44

aromatic, like orange blossom, fruit, amylacetate, etc.: 18, 19, 54, 55

raphanoid: 12

absent: (4), (5), (7), 8, 15, 16, 20, (24), (25), 35, 37, (38), 41, 42, 43, 45, 46, 50, 51, 54, 56

fetid, like rotten meat: 60

farinaceous or farinaceous-rancid, like fish or cucumber: all other numbers

**Spores**

## shape

cruciform-stellate: 4, 5

cuboid: 6, 56, 58

(sub-)isodiametrical,  $Q=1.0-1.1-1.2$ : (9), (11), 13, 14, 17, 21, 22, 23, 24, 32, 39

heterodiametrical: all other numbers

## size

width not exceeding  $7-7.5 \mu\text{m}$ : 3, 8, 11, 26, 27, 28, (29), 30, 32, 33, 34, 42, 43, 44, 47, (55)

length not exceeding  $8.5 \mu\text{m}$ : (11), 32, 34

length not exceeding  $10.5 \mu\text{m}$ : 6, 8, 9, 13, 14, (15), (17), (19), 21, 22, 23, 24, 26, 27, 28, 29, 30, 31, 32, 33, 39, (40), 42, 43, 44, (47), 50, 54, 58

length always  $\geq 9.5 \mu\text{m}$ : 1, 2, (5), (7), (10), 12, 16, 18, 19, (20), (31), 35, 37, 38, 45, 46, 48, 49, 51, (52), (54), 56, 57, 59

**Basidia**

in majority 2-spored: 12, 35, 36

clampless: 4, 5, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 52

**Cheilocystidia**

present: 1, 2, 3, 6, 12, (16), 17, 19, 25, 48, 49, 50, 51, 52, 54, 59

## shape

semiglobose, globuliform or spheropedunculate: 25, 52

tibiiform: 17, 51

irregularly cylindrical-flexuous, with attenuate apex or subcapitate,  $50-90(-140) \mu\text{m}$  long, protruding: 48, 49, 54

**Pigment**

- exclusively encrusting: 7, 8, 9, 10, 11, 12, (15), 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, (42)  
 exclusively intracellular: 4, 5, (6), 40, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 56  
 encrusting and intracellular: 1, 2, 3, (6), 13, 14, (15), 26, 27, 28, 29, 30, 31, 32, 33, 34, (42)  
 membranal and intracellular: 35, 36, 37, 38, (39), 53, 54, 55, 57  
 exclusively membranal: (39), 58.

**Stipitepellis with well-developed hairs**

- (sub-)capitate: 12, 16, 20  
 (sub-)cylindrical: (37), 38

## SECTION NOLANEA

*Rhodophyllus* Quél. subsect. *Mammosi* Romagn. in Bull. Soc. mycol. Fr. 53: 331. 1937. — *Rhodophyllus* Quél. sect. *Mammosi* (Romagn.) Romagn. in Bull. mens. Soc. linn. Lyon 43: 331. 1974.

Characteristics.—Pigment encrusting the hyphae of pileipellis and pileitrama as well as diffusely intracellular, particularly in hypoderm; cheilocystidia present, cylindrical-subcapitate; spores heterodiametrical with basal facet; clamps at least in hymenium always present.

## KEY TO THE SPECIES OF SECTION NOLANEA

- 1a. Spores on the average considerably less than 8  $\mu$ m wide; stipe long and slender, 1–3(–5) mm thick; general habit *Mycena*-like; in forests. . . . . *E. leptopus*, p. 442  
 b. Spores on the average more than 8  $\mu$ m wide; stipe if long then considerably thicker (3–7(–10) mm). . . . . 2  
 2a. Stipe much longer than diameter of pileus; pileus conical, only slightly expanding; in coniferous forests, frequent in spring but also in autumn, then also in dune valleys. . . . . *E. hirtipes* var. *hirtipes*, p. 438  
 b. Stipe about as long as diameter of pileus; pileus dark sepia and more or less flattened (resembling *E. sericeum*); in grasslands. . . . . *E. hirtipes* var. *sericoides*, p. 442

## I. ENTOLOMA HIRTIPES (Schum. ex Fr.) Moser

var. HIRTIPES—Figs. 1a–d

*Agaricus hirtipes* Schum. ex Fr., Syst. mycol. 1: 206. 1821. — *Nolanea hirtipes* (Schum. ex Fr.) Kumm., Führ. Pilzk.: 95. 1871. — *Rhodophyllus hirtipes* (Schum. ex Fr.) Quél., Enchir.: 64. 1886. — *Entoloma hirtipes* (Schum. ex Fr.) Moser in Gams, Kl. KryptogFl., 4. Aufl. 2(b/2): 206. 1978.

*Agaricus acceptandus* Britz. in Ber. naturw. Ver. Schwaben, Augsburg 26: 140. 1881. — *Nolanea acceptanda* (Britz.) Sacc., Syll. Fung. 5: 724. 1887.

MISAPPLIED NAMES.—*Nolanea mammosa* (L. ex Fr.) Quél. sensu Boud., Ricken, non Konr. & Maubl. nec P. D. Orton. ? — *Rhodophyllus mammosus* (L. ex Fr.) Quél. sensu Kühn. & Romagn.

*Nolanea pascua* (Pers. ex Fr.) Kumm. sensu Quél., Rea, Bres.

*Nolanea majalis* (Fr. ex Fr.) Konr. sensu Konr., Rea non Fr.

SELECTED ICONES AND DESCRIPTIONS.—Boud., Icones mycol. 1: 49, pl. 97. 1906 (as *Nolanea mammosa*). — Cetto, Funghi Vero 1: 247 (No. 100). 1975. — Konr. in Bull. Soc. mycol. Fr. 39: 36, pl. 2, figs. 1–4. 1923 (as *N. majalis* Fr.). — Konr. in Bull. Soc. mycol. Fr. 45: 47–49. 1929. — Konr. & Maubl., Icones sel. Fung., pl. 177. 1930. — J. Lange, Fl. agar. dan. 2: 101, pl. 78 G. 1936. — P. D. Orton in Trans. Br. mycol. Soc. 43: 329. 1960 (in key only). — Trimbach in Doc. mycol. 8(29): 39–41. 1978.

TABLE I

Synopsis of the taxa of *Entoloma* subgenus *Nolanea* treated in the present work

Section <i>Nolanea</i>		27. <i>f. eccitoides</i>	p. 488
1. <i>E. hirtipes</i>		28. <i>E. psilopus</i>	p. 489
var. <i>hirtipes</i>	p. 438	29. <i>E. acidophilum</i>	p. 489
2. <i>var. sericoides</i>	p. 442	30. <i>E. argenteostriatum</i>	p. 490
3. <i>E. leptopus</i>	p. 442	31. <i>E. cuniculorum</i>	p. 490
59. <i>E. nivescens</i>	p. 524	32. <i>E. defibulatum</i>	p. 492
Section <i>Staurospora</i>		33. <i>E. fractum</i>	p. 492
4. <i>E. conferendum</i>		34. <i>E. xanthocaulon</i>	p. 493
var. <i>conferendum</i>	p. 446	Section <i>Endochromonema</i>	
5. <i>var. pusillum</i>	p. 450	Subsection <i>Endochromonema</i>	
6. <i>E. rhombisporum</i>	p. 450	35. <i>E. cetratum</i>	p. 496
56. <i>E. inodorum</i>	p. 520	36. <i>E. farinogustus</i>	p. 497
58. <i>E. prismatospermum</i>	p. 523	37. <i>E. cuneatum</i>	p. 498
Section <i>Papillata</i>		38. <i>E. lanuginosipes</i>	p. 499
Subsection <i>Papillata</i>		39. <i>E. occultopigmentatum</i>	p. 501
7. <i>E. papillatum</i>	p. 454	40. <i>E. calthionis</i>	p. 502
8. <i>E. clandestinum</i>	p. 456	41. <i>E. ventricosum</i>	p. 502
9. <i>E. lucidum</i>	p. 458	57. <i>E. pallescens</i>	p. 521
10. <i>E. sericeontitens</i>	p. 459	60. <i>E. foetulentum</i>	p. 525
11. <i>E. ortonii</i>	p. 459	Subsection <i>Infularia</i>	
12. <i>E. cuspidifer</i>	p. 461	42. <i>E. infula</i>	p. 503
Subsection <i>Fibulata</i>		43. <i>E. solstitiale</i>	p. 505
13. <i>E. juncinum</i>	p. 464	44. <i>E. chlorinosum</i>	p. 507
14. <i>E. nitens</i>	p. 466	45. <i>E. verecundum</i>	p. 507
Subsection <i>Minuta</i>		Subsection <i>Tristes</i>	
15. <i>E. minutum</i>	p. 468	46. <i>E. triste</i>	p. 509
16. <i>E. tenellum</i>	p. 470	47. <i>E. undulatosporum</i>	p. 509
Subsection <i>Cosmeoxonema</i>		Subsection <i>Cheilocystidiata</i>	
17. <i>E. tibiicystidiatum</i>	p. 473	48. <i>E. velenovskyi</i>	
18. <i>E. ameides</i>	p. 473	var. <i>velenovskyyi</i>	p. 510
19. <i>E. saccharioliens</i>	p. 474	49. <i>var. longicystidiatum</i>	p. 511
20. <i>E. vernum</i>	p. 476	50. <i>E. cryptocystidiatum</i>	p. 511
21. <i>E. sericeum</i>		51. <i>E. inutile</i>	p. 512
f. <i>sericeum</i>	p. 479	52. <i>E. globulifer</i>	p. 513
22. <i>f. nolaniforme</i>	p. 480	Subsection <i>Icterina</i>	
23. <i>var. cinereo-opacum</i>	p. 482	53. <i>E. icterinum</i>	p. 516
24. <i>E. sericoides</i>	p. 483	54. <i>E. chlorophyllum</i>	p. 518
25. <i>E. sphaerocystis</i>	p. 485	55. <i>E. ambrosium</i>	p. 520
Section <i>Fernandae</i>			
26. <i>E. fernandae</i>			
f. <i>fernandae</i>	p. 487		

**CHARACTERISTICS.**—Carpophores robust, pileus 30–70 mm broad, dark sepia or reddish brown; stipe 90–160 × 3–7(–10) mm, striate; smell strong, rancid-farinaceous; spores 10.4–13.4 (–14.5) × 8–9.3 μm; subcylindrical-subcapitate cheilocystidia present; on calcareous soils in coniferous forests and near coastal dunes, frequent in spring.

Pileus 30–70 mm broad, acutely conical then expanding to conico-campanulate, hemispherical or conico-convex with or without small umbo, with margin involute later straight, strongly hygrophanous, when moist rather dark sepia, date brown or reddish brown (centre e.g. 10 YR 3/2; 7.5 YR 3/2), towards margin slightly paler (7.5 YR 5/2, 5/4, 5/6), translucently striate up to one half of the radius, on drying pallescent to grey-brown (10 YR 4/2, margin to 10 YR 7/3), smooth, innately fibrillous, shining. Lamellae L = about 30, l = 3–5, adnate to almost free, broadly ventricose, pale then brownish pink (10 YR 7/3, 8/3 then 7.5 YR 8/4, 7/4, finally 5 YR 5/4), with eroded concolorous edge. Stipe 90–160 × 3–7(–10) mm, straight, usually slightly broadened at base, yellowish brown to sepia, with paler apex (apex 10 YR 7/2, 6/3; 2.5 Y 7/2, base 10 YR 4/3, 4/2), innately silvery striate, twisted, pruinose at apex, tomentose-hairy at base. Smell and taste strongly farinaceous-rancid or like cod-liver oil.

Spores 10.4–13.4(–14.5) × 8.1–9.3(–9.5) μm, Q = (1.25–)1.3–1.5(–1.6); L-D = (1.7–)2.3–3.7–5.0 μm; irregularly (5–)6(–7)-angled in side-view, with basal facet. Basidia 32–48 × 9–14 μm, 4-spored. Cheilocystidia 35–70 × 5.5–14 μm, subcylindrical to sublageniform, often subcapitate, numerous, sometimes mixed with basidia. Hymenophoral trama regular; cells subcylindrical to inflated, up to 350 μm long and 10–19(–25) μm wide. Pileipellis a cutis made up of radially arranged 4–11 μm wide cylindrical hyphae with intracellular pigment and minutely encrusted walls. Pileitrama regular; cells subcylindrical to inflated, 170–320 × 7.5–19 μm, with pale brown, often minutely encrusted walls. Clamp-connections frequent in hymenium, rare in other tissues.

**HABITAT & DISTRIBUTION.**—In coniferous forests on calcareous soils and (in the Netherlands) in coastal dunes near frondose and coniferous trees. Common in central and northern Europe in spring, also recorded in autumn (Netherlands, Denmark).


**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Noord-Holland, IJmuiden, Midden Heerenduin, 6 Nov. 1976, *M. E. Noordeloos 251*; Overveen, 'Koningshof', 21 Nov. 1976, *J. van Brummelen*; prov. Zuid-Holland, Warmond, 'Huys te Warmont', 8 Nov. 1953, *R. A. Maas Geesteranus 9574*.

NORWAY, Oppland, Jevnaker, Vangsaasen, 1 June 1969, *A. Efstøl & G. Gulden* (O). — DENMARK, Isl. of Mön, 16 Oct. 1977, *M. E. Noordeloos 524*. — BELGIUM, prov. Namur, Rochefort, 3 May 1965, *H. S. C. Huijsman*. — GERMAN FEDERAL REPUBLIC, Bayern, Geltendorf, 3 May 1970, *A. Bresinsky* (M); Gnadewald, 14 May 1951, *M. Moser 51/25* (M).

Orton (1960: 329) gives a key to *Entoloma hirtipes* and related species, mainly based on differences in spore size. In my experience these differences are not as clear-cut as Orton suggests. Particularly Orton's *N. hirtipes* and *N. mammosa* are difficult to distinguish. In the collections of *E. hirtipes* var. *hirtipes* studied I found a variation in spore dimensions covering the measurements given by Orton for both species, without correlations with such characters as size of the carpophore and striation-intensity of the stipe.

I found the size of the spores in typical *Nolanea tenuipes* to differ from the measurements given by Orton, though the difference between the latter and *E. hirtipes* var. *hirtipes* remains clear-cut indeed (see also below, p. 443).

The main distribution of var. *hirtipes* is in central and northern Europe particularly in hilly or mountainous areas in coniferous forests on calcareous soils, frequently in spring, but occasionally in autumn. Orton records *E. hirtipes* var. *hirtipes* from the same habitat in Scotland in late autumn. The Netherlands' collections cited were gathered in or near the coastal dunes also in late autumn. The habitat was not strictly coniferous forest but grassy spots near frondose



Figs. 1a-d. *Entoloma hirtipes* var. *hirtipes*. — Habit, spores and cheilocystidia (1a from Noordeloos 251; 1b from van Brummelen, 21 Nov. 1976; 1c-d from Huijsman, 3 May 1965).

Figs. 2a-b. *Entoloma hirtipes* var. *sericoides*. — Spores and cheilocystidia (2a-b from Bas 5514).

and/or coniferous trees. The size of the carpophores of these collections was somewhat small, but the large spores made me consider them to belong to var. *hirtipes*.

2. *ENTOLOMA HIRTIPES* (Schum. ex Fr.) Moser  
var. *sericoides* (Kühn.) Noordeloos, *comb. nov.*—Figs. 2a–b

*Rhodophyllus mammosus* (Fr.) var. *sericoides* Kühn. apud Kühn. & Romagn. in *Rev. Mycol.* 19: 10. 1954. (Fl. anal.: 187. 1953, nom. nud.)

**CHARACTERISTICS.**—Carpophores thick-set, resembling *Entoloma sericeum* var. *sericeum* and occurring in the same habitat with that species; pileus and stipe rather dark grey-brown; smell farinaceous-rancid; spores  $10\text{--}12.5 \times 8\text{--}9.3 \mu\text{m}$ ; cheilocystidia present cylindrical-sublageniform, often subcapitate; in poorly manured and extensively grazed grasslands in coastal dunes; autumn.

Pileus 15–38 mm broad, conico-convex soon flattening, with or without small umbo, with margin slightly involute, strongly hygrophanous, rather dark brown when moist (7.5 YR 2/2 to 5 YR 2/2), paler at margin, translucently striate up to one-third of the radius, on drying pallescent to grey-brown along radial streaks, glabrous, radially innately fibrillose, shining. Lamellae subdistant, deeply emarginate, pale grey then moderately dark grey-brown with pink tinge, paler towards slightly eroded edge. Stipe 25–45  $\times$  (1.5–)2–4 mm, grey-brown to dark brown-grey, pruinose-flocculose at apex, downwards silvery striate. Flesh in pileus and stipe concolorous with surface, on drying pallescent. Smell strong, like cod-liver oil. Taste rancid.

Spores  $10\text{--}12.5 \times 8.0\text{--}9.3 \mu\text{m}$ ,  $Q = 1.2\text{--}1.3\text{--}1.5$ ,  $L\text{--}D = 2\text{--}4 \mu\text{m}$ , 6–7-angled in side-view, with basal facet. Basidia  $32\text{--}46 \times 8.2\text{--}12 \mu\text{m}$ , 4-spored. Cheilocystidia  $25\text{--}52 \times 2.5\text{--}7 \mu\text{m}$ , (sub-)cylindrical to sublageniform, with rounded, sometimes subcapitate or attenuate tip. Hymenophoral trama regular, with cylindrical to inflated cells up to  $270 \times 9\text{--}21 \mu\text{m}$ , with hyaline, colourless walls. Pileipellis a simple cutis made up of radially arranged,  $2.7\text{--}8 \mu\text{m}$  wide, cylindrical hyphae with intracellular pigment and sometimes minutely encrusted walls, gradually passing into pileitrama. Pileitrama regular with long cylindrical to inflated cells up to  $290 \times 6\text{--}19\text{--}(26) \mu\text{m}$ , with brown, minutely encrusted walls. Clamp-connections frequent in hymenium, rare elsewhere.

**HABITAT & DISTRIBUTION.**—In grasslands, in the Netherlands in extensively grazed meadows in old coastal dunes, rare. Also known from France (type-locality). September–November.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Zuid-Holland, Isl. Goeree, Westduinen, 11 Nov. 1976, *F. Tjallingii* & *G. J. M. G. Tjallingii-Beukers*; prov. Zeeland, Schouwen, Renesse, 'Zouten Haard', 10 Sept. 1970, *C. Bas* 5514.

*Entoloma hirtipes* var. *sericoides* differs from the type by habit, and from *E. leptopus* by the large spores. In the field it is easily confused with *E. sericeum*, with which it grows in the same habitat. Microscopically it is easily distinguished, however, by the large spores, the presence of cheilocystidia and two types of pigment.

3. *Entoloma leptopus* Noordeloos, *nom. nov.*—Figs. 3a–f

*Nolanea tenuipes* Orton in *Trans. Br. mycol. Soc.* 43: 334. 1960 (basionym), non *Entoloma tenuipes* Murr. 1917.

*Rhodophyllus mammosus* (L. ex Fr.) Quél. var. *obsoletus* Romagn. in *Rev. Mycol.* 19: 7. 1954. (Fl. anal.: 188. 1953, nom. nud.)



MISAPPLICATIONS.—*Rhodophyllus mammosus* var. *tenuis* (Fr.) Kühn. & Romagn. sensu Kühn. & Romagn. non Fr. — *Entoloma mammosum* (Fr.) Hesler sensu Moser.

SELECTED ICONES & DESCRIPTIONS.—Dähne & Dähne, 700 Pilze: 261. 1979 [as *E. mammosum*].

CHARACTERISTICS.—Carpophores slender, mycenoid; pileus 10–40(–45) mm broad, yellowish to reddish brown, often with discolor, darker papilla; stipe up to 110 × 1–3(–5) mm; smell usually strongly farinaceous-rancid; spores 8.5–12 × 5.8–7.5  $\mu$ m; cylindrical-subcapitate cheilocystidia present; in deciduous forests; summer-autumn.

Pileus 9–40(–45) mm broad, conico-convex, campanulate or hemispherical then expanding to plano-convex, usually with distinct papilla, rarely more or less flattened, with margin slightly involute when young only, mostly straight, sometimes exceeding the lamellae, often splitting with age and more or less crenulate, strongly hygrophanous, when moist yellow-brown, grey-brown, leather-coloured or reddish brown, centre (papilla) frequently distinctly darker than rest (centre 10 YR 6/3, 5/3, 4/3, 4/2, 3/2, even 2/2; 7.5 YR 5/4, 4/2, 3/2 towards margin and interstriation distinctly paler, 10 YR 6/3, 6/4, 7/3, 7/4; 7.5 YR 6/4, 7/4), translucently striate up to centre, strongly pallescent on drying towards yellowish brown or greyish isabella (10 YR 7/2, 7/3, 8/3, 8/4), glabrous, strongly radially fibrillous-lustrous, at centre sometimes more or less fluffy-rugulose. Lamellae L = 20–30(–35), l = 1–3.5(–7), almost free or adnate-emarginate, ventricose, pale (brown) then pink finally brownish pink (10 YR 8/3; 7.5 YR 8/3, 8/4, 7/4, 6/4 to 6/6), with almost entire concolorous edge. Stipe 20–85(–110) × 1–3(–5) mm, straight-cylindrical, usually slightly broadened at base, solid then fistulose, pale brown, reddish brown or grey-brown, usually slightly paler than pileus (10 YR 7/4, 6/4, 6/3, 5/4, 5/6, 5/8; 7.5 YR 5/4), darker at base (10 YR 3/2; 7.5 YR 3/2), pruinose-flocculose at apex, downwards sparsely to abundantly fibrillously striate, never absolutely smooth, sometimes twisted, white-tomentose at base. Flesh thin-membranaceous, rather brittle in pileus, firmer to subcartilagineous in stipe, almost concolorous with surface, slightly pallescent on drying. Smell usually strongly like cucumber; rancid-farinaceous, particularly when cut, rarely absent. Taste strongly rancid, rarely indistinct.

Spores (8.1–)8.5–12.0(–13.9) × (5.2–)5.8–7.5(–8.1)  $\mu$ m, Q = 1.2–1.45–1.6(–1.8), L–D = 2.5–4–6  $\mu$ m, 7–8-angled in side-view, with basal facet, often with strong suprahilar depression. Basidia 24–40 × 7.5–13  $\mu$ m, 4-(rarely 2-)spored. Cheilocystidia 25–52 × 4–10  $\mu$ m, numerous, usually mixed with basidia, versiform, mostly cylindrical-subcapitate to flexuose or clavate, more rarely cylindrico-attenuate. Hymenophoral trama regular, composed of cylindrical to inflated cells up to 380  $\mu$ m long and 14–34(–50)  $\mu$ m wide. Pileipellis a thin cutis made up of 3–9.5(–13.5)  $\mu$ m wide cylindrical hyphae with pale, often minutely encrusted walls, gradually passing into pileitrama or subpellis, composed of short, inflated cells 32–80 × 11.5–32  $\mu$ m, with scarce to abundant intracellular pigment and with slightly coloured, sometimes minutely encrusted walls. Pileitrama regular, composed of cylindrical to inflated cells up to 470  $\mu$ m long and up to 27  $\mu$ m wide, with in upper layer usually pale, minutely encrusted hyphae (particularly the narrowest ones). Clamp-connections abundant in hymenium, elsewhere rare or lacking.


HABITAT & DISTRIBUTION.—Terrestrial on damp, humus-rich soils in frondose forests, common in the Netherlands, particularly in Alder and Ash-woods (*Alno-Padion* and *Ulmion-carpinifoliae* associations). Also known to occur in Belgium, Great Britain, France and Germany. July–November.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Roden E. of Vagevuur, 14 Sept. 1976, M. E. Noordeloos 169; prov. Gelderland, Winterswijk, Bek-en-Delle, 3 Sept. 1979, J. Schreurs; ditto, Willinks Weust, 20 Oct. 1977, M. E. Noordeloos 553; Daarlerveen, 1 Sept. 1979, C. M. den Held-Jager; Hierden, Hulshorst, Grote Water, 14 Oct. 1956, A. Groeneweg; prov. Utrecht, Leusden, Schoolsteegbosjes, 6 Aug. 1979, J. Wisman 71; Bunnik, Oud Amelisweerd, 22 July 1954, R. A. Maas Geesteranus 9997; prov. Noord-Holland, Kortenhoef, 11 July 1970, J. Daams; Noordwijkerhout, Vogelenzang, 19 Oct. 1958, C. Bas 1618; prov. Zuid-Holland, Voorschoten, estate 'ter Horst', 2 July 1974, C. Bas 6313/14/15; 7 Aug. 1974, M. E. Noordeloos 25; 3 Sept. 1974, M. E. Noordeloos 42; 28 Sept. 1977, M. E.

*Noordeloos* 472; 15 Aug. 1979, *M. E. Noordeloos* 972; Isl. of Voorne, Quackjeswater, 27 Oct. 1976, *M. E. Noordeloos* 234; Katwijk, Pan van Persijn, 17 Oct. 1969, *E. Arnolds* 411; prov. Zeeland, Domburg near Castle, 30 Oct. 1976, *M. E. Noordeloos* 239; prov. Noord-Brabant, Nuenen, Nuenense Broek, 15 Sept. 1979, *M. E. Noordeloos* 1021; Terheyden, Eendekooi, 28 Oct. 1978, *P. B. Jansen* 78-453; Drunen, along Channel, 23 Oct. 1969, *E. Arnolds* 469; prov. Limburg, Wijlre, 12 Oct. 1968 and 11 Nov. 1970, *P. B. Jansen*; Schelsberg, 8 Nov. 1974, *P. B. Jansen*; Gronsveld, Savelsbos, 28 Sept. 1976, *M. E. Noordeloos* 195/6.

GREAT BRITAIN, Sussex, Friston, 17 Nov. 1957, *P. D. Orton* (holotype, K). — BELGIUM, prov. Namur, Han-sur-Lesse, Fond d'Auffe, 2 Oct. 1977, *M. E. Noordeloos* 474/5. — FRANCE, dept. Oise, Orry-la-Ville, 3 Nov. 1976, *H. Romagnesi* 76.170 (Herb. Romagn., PC).

NOTES ON THE HOLOTYPE.—The holotype consists of fragments of one specimen with the following microscopical characters. Spores  $(8.7-9.3-11(-11.4) \times 6.4-7.5(-8.1) \mu\text{m}$  (average spore  $10.1 \times 7 \mu\text{m}$ ),  $Q=(1.2-1.3-1.4-1.55$ ,  $L-D=2.4-3.0-4.0 \mu\text{m}$ , 6-8-angled in side-view, with blunt base (basal facet). Basidia  $25-40 \times 11.5-14 \mu\text{m}$ , 4-spored. Cheilocystidia  $21-43 \times 5-9(-11) \mu\text{m}$ , cylindrical sometimes subcapitate, frequent, mixed with basidia. Hymenophoral trama with cells up to  $400 \mu\text{m}$  long and up to  $26 \mu\text{m}$  wide, cylindrical or inflated. Pileipellis a simple cutis



Figs. 3a-f. *Entoloma leptopus*. — Habit, spores and cheilocystidia (3a from *Noordeloos* 195; 3b-c from *Scheurs*, 20 Oct. 1977; 3d, 3f from *Jansen*, 8 Nov. 1974; 3e from *Jansen*, 11 Nov. 1970).

made up of 3–11  $\mu\text{m}$  wide, cylindrical hyphae with minutely encrusted walls. Pileitrama regular, with cells of similar size and shape as in hymenophoral trama. Clamp-connections seen in meniumium.

*Entoloma leptopus* is easy to distinguish from *E. hirtipes* and varieties by its different habitat, slender habit and small and narrow spores. Macroscopically the slender habit, discolor pileus with dark papilla and the habitat make it often easy to recognize *E. leptopus* in the field. The odourless form, described by Romagnesi (1954, l.c.) as *R. mammosus* var. *obsoletus* is, besides the lack of smell, not sufficiently different from the typical *leptopus* to justify a distinction on varietal level. Usually, however, the smell is quite strong and can be noticed from a distance.

*Agaricus mammosus* Fr. var. *tenuior* Fr., Icon. sel. pl. 98 fig. 4 (1874), is most probably the same as *Nolanea papillata* Bres. The interpretation of this taxon by Kühn. & Romagn. (1953, l.c.) may be considered a misapplication. (See also notes on *Agaricus mammosus* under the excluded names, p. 526).

*Rhodophyllus hebes* Romagn. is said to differ from *E. leptopus* by the different smell (farinaceous and not like cucumber or fish), the more flattened, subpapillate pileus and the more irregularly shaped spores. In a note following the diagnosis Romagnesi mentions, however, that there is a chain of forms connecting typical '*R. mammosus*' (= our *E. hirtipes* var. *hirtipes*) with *R. hebes*. Indeed I found the distinguishing characters mentioned above to be rather variable. Except in the type of *R. hebes* and in a few additional collections in Herb. Romagnesi I failed to find the extremely elongate spores as depicted by Romagnesi (1954: fig. 14). In *E. leptopus* in general, however, the shape of the spores is rather variable (see figs. 3c–d). There seems to be no reason either to consider the shape of the pileus in this case as a taxonomically valuable character. Moreover the difference in smell seems to be too subtle. Future studies are necessary to prove the difference between *E. leptopus* and *R. hebes*.

In the field *Entoloma leptopus* may be confused with *E. juncinum*, which has about the same colours particularly when dried up, and the same strong smell, but *E. juncinum* is easily distinguished microscopically by the lack of cheilocystidia and by the totally different spores.

**Entoloma section *Staurospora* (Largent & Thiers) Noordeloos,  
comb. nov.**

*Nolanea* (Fr.) Kumm. emend. Largent & Benedict sect. *Staurospori* Largent & Thiers in Northwest Sci. 46: 37. 1972. — *Rhodophyllus* Quél. sect. *Staurospori* (Largent & Thiers) Romagn. in Bull. mens. Soc. linn. Lyon 43: 330. 1974. — Holotype: *Entoloma conferendum* (Britz.) Noordeloos (= *Nolanea staurospora* Bres.).  
*Rhodophyllus* Quél. subsect. *Staurospori* Romagn. in Bull. Soc. mycol. Fr. 53: 331. 1937 (nom. nud.).

CHARACTERISTICS.—Spores cuboid or prismatical. Pigment membranal, encrusting and/or intracellular.

KEY TO THE SPECIES OF SECTION STAUROSPORA

- 1a. Spores prismatical-stellate; pigment intracellular. . . . . 2  
b. Spores cuboid; pigment encrusting and/or intracellular. . . . . 3

- 2a. Carpophores small and slender; pileus up to 15 mm broad, stipe up to 1.5 mm wide; on rotten trunks of frondose trees. . . . . *E. conferendum* var. *pusillum*, p. 450  
 b. Carpophores more robust, terrestrial. . . . . *E. conferendum* var. *conferendum*, p. 446  
 3a. Spores large, 10.2–12.4  $\mu\text{m}$  long; pileus blackish brown, strongly contrasting with pale, whitish stipe. . . . . *E. inodorum*, p. 520  
 b. Spores smaller, on the average less than 10  $\mu\text{m}$  long; pileus and stipe paler and of same colour. . . . . 4  
 4a. Stipe minutely pruinose at least at upper half, with large fusiform caulocystidia 60–75  $\mu\text{m}$  long, pigment membranous (not encrusting). . . . . *E. prismatospermum*, p. 523  
 b. Stipe smooth, pigment intracellular, sometimes also membranous and/or encrusting. . . . . *E. rhombisporum*, p. 450

4. *Entoloma conferendum* (Britz.) Noordeloos, *comb. nov.*  
 var. *conferendum*—Figs. 4a–d

*Agaricus conferendus* Britz. in Ber. naturw. Ver. Schwaben, Augsburg 26: 140. 1881. — *Nolanea conferenda* (Britz.) Sacc., Syll. Fung. 5: 723. 1887.

*Nolanea staurospora* Bres., Fungi trident. 1: 18, pl. 20. 1882. — *Rhodophyllus staurosporus* (Bres.) J. Lange, Fl. agar. dan. 2: 99, pl. 99A. 1936. — *Entoloma staurosporum* (Bres.) Horak in Sydowia 28: 222. 1976 "1974/75".

*Rhodophyllus staurosporus* var. *typicus* Kühn. & Romagn., Fl. anal.: 187. 1953.

*Rhodophyllus staurosporus* var. *platyphyllus* Romagn. & Favre in Rev. Mycol. 3: 77. 1938.

*Nolanea staurospora* var. *farinacea* Largent & Thiers in Northwest Sci. 46: 37. 1972.

*Nolanea staurospora* var. *incrustedata* Largent & Thiers in Northwest Sci. 46: 38. 1972.

*Agaricus dissidens* Britz. in Ber. naturw. Ver. Schwaben, Augsburg 26: 140. 1881, pl. 176, No. 27; pl. 188, No. 109. 1881. — *Nolanea dissidens* (Britz.) Sacc., Syll. Fung. 5: 723. 1887.

*Agaricus subpostumus* Britz. in Ber. naturw. Ver. Schwaben, Augsburg 26: 143, pl. 173, No. 38. 1881. — *Nolanea subpostuma* (Britz.) Sacc., Syll. Fung. 5: 725. 1887.

*Rhodophyllus rickenii* Romagn. in Bull. Soc. mycol. Fr. 48: 320. 1932. — *Rhodophyllus staurosporus* subsp. *rickenii* (Romagn.) Romagn. in Rev. Mycol. 2: 86. 1937. — *Rhodophyllus staurosporus* var. *rickenii* (Romagn.) Kühn. & Romagn., Fl. anal.: 187. 1953. — *Nolanea rickenii* (Romagn.) Konr. & Maubl., Agaricales: 264. 1948.

*Rhodophyllus rickenii* var. *obscurior* Romagn. in Bull. Soc. mycol. Fr. 48: 320. 1932. — *Rhodophyllus staurosporus* var. *obscurior* (Romagn.) Romagn. in Rev. Mycol. 2: 86. 1937.

*Rhodophyllus rickenii* var. *subrugosus* Romagn. in Bull. Soc. mycol. Fr. 48: 321. 1932. — *Rhodophyllus staurosporus* var. *subrugosus* (Romagn.) Romagn. in Rev. Mycol. 2: 86. 1937.

*Entoloma nothofagi* Stevenson in Kew Bull. 16: 234. 1962.


*Entoloma botanicum* Stevenson in Kew Bull. 16: 236. 1962.

MISAPPLIED NAMES.—*Nolanea pascua* (Pers. ex Fr.) Kumm. sensu Britz., Cooke, Ricken non Bres., Boud., Rea. — *Nolanea proletaria* (Fr.) Gill. sensu Boud., Rea non Fr., Ricken nec Bres. — *Agaricus cetratus* Fr. sensu Britz. pro parte.

SELECTED ICONS AND DESCRIPTIONS.—Bresadola, Iconogr. mycol.: 12: 584. 1929. — Cetto, Funghi Vero 2: 248. 1976. — Horak, l.c. 1976. — Konrad in Bull. Soc. mycol. Fr. 45: 49–50. 1929. — Konrad & Maublanc, Icon. sel. Fung., pl. 178 fig. 1. 1930. — J. Lange, Fl. agar. dan., pl. 77 fig. a. 1936. — F. H. Möller, Fungi Faerøes 1: 239, fig. 111. 1945.

CHARACTERISTICS.—Pileus grey-brown, horn or reddish brown, strongly lustrous particularly when dry, usually with rugulose-felted centre; stipe striate with more or less aeriferous fibrils, often twisted; spores 4–5-angled in side-view, cruciform-stellate.

Pileus 23–50 mm broad, hemispherical or conical, then conico-campanulate to conico-convex, finally convex with or (mostly) without weak umbo, rarely acutely papillate, sometimes truncate,



Figs. 4a-d. *Entoloma conferendum* var. *conferendum*. — Habit and spores (4a from Brand, 11 Oct. 1979; 4b, 4d from Bas 7605; 4c from Noordeloos 181).

Figs. 5a-b. *Entoloma conferendum* var. *pusillum*. — Habit and spores (5a-b from Noordeloos 178).

with margin slightly involute when young, usually straight with age, sometimes splitting; strongly hygrophanous, when moist translucently striate often up to centre, at least up to one half of the radius of the pileus, dark grey-brown, sepia, also horn brown or more reddish brown, usually fairly dark at centre, becoming paler towards margin (centre 10 YR 2/2, 3/2, 3/3, 3/4; 7.5 YR 3/2 or 5 YR 2/2, 3/2, 4/2, towards margin 10 YR 5/4, 6/4, 6/3; 7.5 YR 5/2, 4/2 or 5 YR 5/4, 5/6, outermost margin 10 YR 7/4, 7.5 YR 7/4), on drying strongly pallescent to greyish brown or ochraceous-greyish with centre long remaining darker (centre 10 YR 5/2, 5/3, 5/4; 2.5 Y 5/4, rest 10 YR 6/3, 7/3, 8/3), strongly lustrous at centre often with aeriferous fibrils fluffy-rugulose to subsquamulose, sometimes on limb radially wrinkled. Lamellae  $L=20-50$ ,  $l=1-3-5(-7)$ , (almost) free, moderately to broadly ventricose, up to 10 mm broad, pale then pink often tinged grey or brown, becoming reddish brown with age (7.5 YR 7/4, 6/4; 5 YR 7/6, 7/4 towards 5 YR 6/4, 6/6), with entire or serrulate concolorous edge. Stipe  $25-80 \times 2-6.5$  mm, cylindrical or flattened with longitudinal groove, with broadened to subbulbous base (up to 12 mm wide) or not, strongly silvery-fibrilously striate on yellowish brown to greyish brown background (upper half e.g. 2.5 YR 7/6 or 7/2, basal part 10 YR 4/2, 3/2 or 5/4 towards 2.5 Y 4/4), often twisted, apex sometimes minutely pruinose, base usually with white tomentum, rest glabrous, solid then fistulose. Flesh concolorous with surface but in fleshy specimens context distinctly paler, fibrillose, fairly brittle particularly in pileus. Smell spontaneously often weak or absent but usually distinctly farinaceous-rancid on cutting, rarely absent. Taste strong, disagreeably rancid-farinaceous, rarely more or less mild. Spore print (yellowish) red, 2.5 YR 6 or 5 YR 5/6.

Spores  $(7.9-8.0-12.7(-13.9) \times (7.0-7.4-11.5(-12.7)) \mu\text{m}$ ,  $Q=1.0-1.3-1.5(-1.6)$ ,  $L-D=0-2.5-4 \mu\text{m}$ , rather variable in shape, mostly irregularly 4-angled in side-view with pointed, sharp angles causing a cruciform or stellate appearance, often the axe distorted or twisted, also transitions to 5-6-angled spore type occurring. Basidia  $27-37 \times 10.5-16.3 \mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular; cells cylindrical or very slightly inflated, never fusoid,  $70-324 \times 13.5-32(-35) \mu\text{m}$ , with hyalinous, colourless walls. Pileipellis a cutis with transitions to a trichodermium, particularly at centre of pileus, made up of repent to ascending cylindrical hyphae  $6-12(-14) \mu\text{m}$  wide, with brown intracellular pigment, subpellis usually well developed, composed of chains of short inflated cells  $40-100 \times 18-40(-54) \mu\text{m}$ , with abundant intracellular pigment; pileitrama regular, hyphae cylindrical with cells  $70-220 \times 18-32 \mu\text{m}$ , usually colourless. Clamp-connections none.

**HABITAT & DISTRIBUTION.**—Euryoecious, viz. occurring in all sorts of habitats: grasslands, marshy places, in *Sphagnum*, in humus in both coniferous and frondose forests on different types of soil. Widespread, most probably cosmopolitan (Horak 1976: 222-224). In the Netherlands fairly common from spring to late fall.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Drenthe, Roden, Moltmakersstuk, 15 Sept. 1976, *M. E. Noordeloos* 181/182; Norg, along Peizerdiep, 16 Sept. 1976, *J. Schreurs*; prov. Overijssel, Gorsseel, 't Joppe, 31 May 1954, *C. Bas* 473; Ommen, Eerder Achterbroek, 14 Oct. 1962, *C. Bas* 2875; Delden, estate 'Twickel', 9 Sept. 1972 and 7 Oct. 1977, *E. Kits van Waveren*; Oldenzaal, estate 'het Snippert', 19 Oct. 1972, *E. Kits van Waveren*; Denekamp, estate 'Singraven', 3 Oct. 1965, *E. Kits van Waveren*; prov. Gelderland, Winterswijk, Korenburgerveen, 29 Sept. 1973, *C. Bas* 6173; idem, Willinks Weust, 20 Oct. 1977, *M. E. Noordeloos* 552; idem, Wooldse Veen, 16 Oct. 1972, *J. Schreurs*; Overassel, near 'de Diervoort', 26 Oct. 1960, *R. A. Maas Geesteranus* 13491 & *C. Bas* 2315; idem, 21 Oct. 1969, Hatertsebroek, *E. Arnolds*; Gietelo, estate 'de Poll', 6 May 1979, *G. Piepenbroek-Grooters* & *H. Piepenbroek* 1168; Brummen, Voorstonden, 17 Sept. 1978, *E. Batten*; Vorden, Dennendijk, 4/5 June 1960, *E. Kits van Waveren*; Voorst, estate 'het Hartelaar', 8 May 1978, *H. Piepenbroek* & *G. Piepenbroek-Grooters* 1079; Niimegen, Hatert, 26 Sept. 1973, *C. Bas* 6175; prov. Utrecht, Amersfoort, Woudenberg, estate 'de Boom', 12 Sept. 1953, *A. F. M. Reijnders*; prov. Noord-Holland, Isl. Texel, 'de Geul', 25 Oct. 1977, *M. E. Noordeloos* 561; idem, near Den Hoorn, 28 Oct. 1978, *M. E. Noordeloos* 580; Bergen, near 'Klompduin', 5 Oct. 1969, *E. Arnolds* 391; Kortenhoef, 'Oppad', 8 Sept. 1976, *M. E. Noordeloos* 164; Noordwijkerhout, estate 'Vogelzang', 20 Oct. 1951, *R. A. Maas Geesteranus* 8101; prov. Zuid-Holland, Warmond, 'Huys te Warmont', 7 Nov. 1950, 13 Oct. 1951, 26 Oct. 1951, 23 Nov. 1951, and 3 Nov. 1953, *R. A. Maas Geesteranus* 7623, 8082, 8121, 8179.

9580; and 10 Oct. 1969, *E. Arnolds* 397; Isl. Voorne, Oostvoorne, Heveringen, 17 May 1968, *C. Bas* 4979A; Isl. Goeree, Westduinen, 11 Oct. 1979, *M. Brand*; prov. Noord-Brabant, Chaam, estate 'de Honsdonk', 19 Sept. 1956, *R. A. Maas Geesteranus* 11717; Zundert, 'de Krochten', 10 July 1956, *C. Bas* 1049; Drunen, along 'Drongelens kanaal', 20 Aug. 1972, *P. B. Jansen* 72-227; Bergen-op-Zoom, estate 'de Mattenburg', 25 Sept. 1971, *P. B. Jansen* 71-193; Baarle-Nassau, 2 May 1965, *P. B. Jansen* 65-175; Nw Ginneken, Goudbergven, 25 Aug. 1963, *P. B. Jansen* 63-168; prov. Limburg, Wijre, 28 Oct. 1979, *C. Bas* 7605; Mook, 10 Oct. 1964, *E. Kits van Waveren*; Heerlen, Schelsberg, 8 Nov. 1974, *P. B. Jansen* 74-280. NORWAY, Oppland, Dombås-region near Fokstua, 2 Aug. 1977, *M. E. Noordeloos* 370 and 5 Aug. 1977, *M. E. Noordeloos* 388. — GREAT BRITAIN: Wales, Lake Yrnwy, 16 Sept. 1976, *E. Kits van Waveren*; Devonshire, New Forest, 14 Sept. 1971, *E. Kits van Waveren*. — BELGIUM: prov. Limburg, Genk, 21 May 1923, *Beeli* 1140 (BR); Bévercé, 24 Sept. 1976, *P. B. Jansen*; prov. Luxemburg, 13 Oct. 1930, *Beeli* 1880 (BR).

*Entoloma conferendum* is a very common and a very polymorphic species: it occurs in all kinds of habitats and has most probably a world-wide distribution (Horak 1976: 222).

It is well-known to all European mycologists. In Fries's time it was generally called '*Agaricus pascuus* Pers.', a collective species which comprised *E. conferendum* as well as *E. sericeum* and *E. vernum*. At present *A. pascuus* is generally considered a nomen dubium. No type collection is available in Herb. Persoon (Singer 1961: 36).

Bresadola (1882, l.c.) recognized *N. staurospora* as a species in its own right in the *A. pascuus*-complex, a view generally accepted by his contemporaries as well as by most modern mycologists. Looking through the plates of Britzelmayr I found on plate 179 as No. 26A a picture of *Agaricus conferendus* Britz. which is unmistakably identical with *N. staurospora* Bres. Unfortunately Britzelmayr's name for the species has been published one year before that of Bresadola, and therefore has priority. Consequently in the present work the well-known and widely used epithet *staurosporum* unfortunately has to be changed in *conferendum* Britz. In the same work of Britzelmayr two more older synonyms of *Nolanea staurospora* Bres. were encountered, viz. *A. dissidens* Britz. and *A. subpostumus* Britz.

*Entoloma conferendum* is a very variable species, in size and shape as well as in the colour of the carpophores. The pileus shows tinges varying from grey-brown to sepia, leather-brown or reddish brown, its surface may be smooth, but has more often a covering of loose aceriferous fibrils, particularly when dry, the centre is sometimes even fluffy-subscamulose. The lamellae are usually tinged brown or grey and narrowly to rather broadly ventricose. The smell may be absent or farinaceous-rancid. This variability is reflected in the large number of varieties published. They are based upon (i) lack of smell and pale lamellae (var. *rickenii* Romagn.); (ii) a slightly darker pileus (var. *obscurior* Romagn.); (iii) the subfelted-subrugulose aspect of the pileus (var. *subrugosus* Romagn.); (iv) the broad lamellae (var. *platyphyllus* Romagn.-Favre); (v) the farinaceous smell (var. *farinacea* Largent). In the course of my investigations I found these differences not correlated with other taxonomically important characters and therefore consider these varieties as mere forms of var. *conferendum*.

A taxonomically more important case is that of *Nolanea staurospora* var. *incrustedata* Largent & Thiers from California, described as having encrusting pigments only. I did not study the type of this variety but if it really belongs to the complex of *E. conferendum* this throws a new light upon the constancy of intracellular pigment as a character for that complex. Compare also the discussion under *E. rhombisporum*, p. 452.

*Entoloma conferendum* var. *pusillum* (= *Rhodophyllum xylophilus* J. Lange) is considered here a true variety on account of its habitat and slender habit.

The synonymy of *E. botanicum* and *E. nothofagi* is accepted on the strength of the evidence supplied by Horak (l.c.).

5. *ENTOLOMA CONFERENDUM* var. ***pusillum*** (Velen.)

Noordeloos, *comb. nov.*—Figs. 5a–b

*Nolanea pusilla* Velen., České Houby: 626. 1921. — *Rhodophyllus pusillus* (Velen.) Romagn. in Bull. Soc. mycol. Fr. 53: 332. 1937. — *Entoloma staurosporium* (Bres.) Horak var. *pusillum* (Velen.) Noordeloos in Persoonia 10: 252. 1979.

*Rhodophyllus xylophilus* J. Lange in Dansk bot. Ark. 2(11): 35. 1921 (non *Entoloma xylophilum* Speg.). — *Nolanea xylophila* (J. Lange) P. D. Orton in Trans. Br. mycol. Soc. 43: 179. 1960.

**CHARACTERISTICS.**—Fruitbodies rather slenderly mycenoid, pileus up to 15 mm broad; stipe 21–50 × 1–1.5 mm, on rotten stumps of frondose trees; spores cruciform-stellate.

Pileus 10–15 mm broad, (conico-)convex to flattened with involute margin, later straight, hygrophanous, when moist translucently striate up to centre, yellowish brown or slightly reddish tinged (10 YR 6/4 to 7.5 YR 6/4), with darker centre (10 YR 4/4, 7.5 YR 4/4), on drying pallescent to ochraceous grey or pale brown (10 YR 7/3), smooth, dull or subsquamulose-rugulose at centre. Lamellae L=24–25, l=1(–3), absolutely free, ventricose, exceeding the pileus, pink (7.5 YR 7/4, 6/4 or 5 YR 7/4, 6/4) without any brown tinge, with entire concolorous edge. Stipe 21–50 × 1–1.5 mm, straight, cylindrical with slightly swollen base, pale yellow (2.5 Y 7/4), minutely pruinose at apex, rest silvery striate lengthwise. Smell and taste indistinct to farinaceous.

Spores (8.0–)9–12(–12.5) × 7–10.4(–11) μm, Q=1.0–1.2–1.4(–1.5), L–D=0–1.8–3.5(–5) μm, cruciform-stellate, rather irregular in shape, mostly with 4 angles, also transitions to asymmetric 6-angled type. Basidia 22–45 × 10–15 μm, 4-(rarely 2-)spored. Cystidia absent. Hymenophoral trama regular; cells cylindrical, up to 150 × 4.5–12 μm. Pileipellis a cutis made up of cylindrical, 4.8–12 μm wide hyphae with cylindrical to clavate endcells up to 12(–15) μm wide with at centre of pileus transitions to a trichodermium with ascending terminal cells; intracellular pigment abundant, subpellis well-developed, composed of short inflated cells up to 25 μm wide. Pileitrama regular; cells up to 150 μm long, 12.5–24 μm wide. Clamp-connections absent.

**HABITAT & DISTRIBUTION.**—On rotten trunks of frondose trees such as *Betula*, *Alnus*, *Corylus* and *Fagus*. Known from Denmark (J. Lange, l.c.), Great Britain (Orton & al., 1960), the Netherlands, and Czechoslovakia.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Drenthe, Roden, Mensingebos, 15 Sept. 1976, M. E. Noordeloos 178; prov. Noord-Holland, Bergen, Eeuwige Laan, 8 May 1958, C. Bas 1420.


DENMARK, Jylland, Vellin-plantage, 10 Nov. 1969, leg. Mykol. Congress (C). — CZECHOSLOVAKIA, Bohemia, Prague, Mnichovice, July 1919, J. Velenovský (holotype of *Nolanea pusilla*, PRC); idem, 27 July 1946, V. Vacek (PRM).

The microscopical characters of *Entoloma conferendum* var. *pusillum*, particularly the size and the shape of the spores, are similar to those of the type-variety. It is considered a variety, however, on account of the rather slender appearance and the habitat.

6. *ENTOLOMA RHOMBISPORUM* (Kühn. & Bours.) Horak—Figs. 6a–d

*Leptonia rhombispora* Kühn. & Bours. in Bull. Soc. mycol. Fr. 45: 276. 1929. — *Rhodophyllus rhombisporus* (Kühn. & Bours.) Romagn., Rhodoph. Madag.: 36. 1941. — *Entoloma rhombisporum* (Kühn. & Bours.) Horak in Sydowia 28: 228. "1974/1975" 1976.





Figs. 6a-d. *Entoloma rhombisporum*. — Habit, spores and cheilocystidia (6a, 6c-d from Noordeloos 1070; 6b from Romagnesi, 26 Sept. 1974).

**CHARACTERISTICS.**—Pileus conical or conico-campanulate with papillate or truncate-subdepressed centre, yellowish brown; stipe pale, yellowish, polished; spores cuboid; cheilocystidia lageniform-tibiiform.

Pileus 20–34 mm broad, truncate-conical or conico-campanulate with centre blunt or papillate or sometimes slightly depressed, later expanding to campanulate-convex or even flattened, then with large conical umbo, with straight margin, strongly hygrophanous, when moist dark yellowish brown (10 YR 4/4, 4/6) at centre and striae, interstriation and margin paler (10 YR 5/4), translucently striate up to three fourths of the radius, on drying pallescent to pale yellowish brown (2.5 Y 6/4, 7/4), but at centre remaining slightly darker, glabrous, smooth when moist, becoming coarsely radially fibrillose and shining on drying. Lamellae  $L = 20-25$ ,  $l = 3-5$ , distant, almost free, triangular then ventricose, pink with yellow tinge (10 YR 7/6 or 6/6 to 7.5 YR 7/6 or 6/6), slightly pallescent on drying, sometimes very slightly transversely veined, with entire, concolorous edge sometimes exceeding the pileus. Stipe  $30-60 \times 2-4$  mm, cylindrical, straight, sometimes somewhat broader at base, yellow to yellowish brown, with apex and sometimes also base paler (10 YR 5/4, 6/4 to 2.5 Y 5/4, apex and base 2.5 Y 6/4, 7/4 or entire stipe 5 Y 7/4), not fibrillose, smooth, glabrous, more or less shining, solid, then narrowly fistulose. Flesh slightly paler than surface, subcartilagineous in stipe, brittle in pileus. Smell absent or faint (in one specimen slightly fruity), distinctly farinaceous when cut. Taste farinaceous.

Spores  $8.1-10.5(-10.6) \times (7.0-7.6-10.4 \mu\text{m})$ ,  $Q = 1.0-1.1-1.2$ ,  $L-D = 0-1-1.5 \mu\text{m}$ , cuboid, 4- (rarely 5-)angled in side-view. Basidia  $35-52 \times 10-17 \mu\text{m}$ , broadly clavate, 4-spored. Cheilocystidia  $40-70 \times 10.5-21 \times 3.5-8.1 \mu\text{m}$ , lageniform to tibiiform with broadly swollen basal half and usually long, tapering, moniliform or capitate neck, thin-walled, colourless, numerous; edge usually homomorph. Hymenophoral trama regular; cells fusoid to cylindrical,  $(70-135-297 \times (3.5-8-27(-30) \mu\text{m})$ , colourless, thin-walled. Pileipellis a thin cutis made up of  $2-8 \mu\text{m}$  wide cylindrical hyphae with uniformly coloured, sometimes minutely encrusted walls and intracellular pigment, gradually passing into pileitrama or subpellis differentiated into a distinct layer of short inflated cells  $70-120 \times 10-32 \mu\text{m}$ . Pileitrama regular, cells cylindrical to inflated,  $162-324 \times 8-27 \mu\text{m}$ , intracellular pigment in upper layer, sometimes in addition minutely encrusted. Clamp-connections numerous in hymenium, less abundant to absent in other tissues.

**HABITAT & DISTRIBUTION.**—The Netherlands' collections were made in a moist valley in old coastal dunes on sandy soil with relatively low pH between *Salix repens*, *Menyanthes trifoliata* and *Sphagnum* species. It was originally described from a meadow without indication of type of vegetation. Favre (1948: 53) collected it in a *Sphagnum*-bog. Rare; known from France (type-locality), Switzerland, and the Netherlands.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Noord-Holland, Callantssoog, Nature-reserve 'Zwanewater', S. of the lakes, three different localities, 21 Oct. 1979, *M. E. Noordeloos 1070, 1071 and 1075*.

FRANCE, Dordogne, Ch. de Rossignol, 26 Sept. 1974, *H. Romagnesi 74.263* (Herb. Romagn., PC).

*Entoloma rhombisporum* is fairly easily distinguished by its cuboid spores and the type of cheilocystidia. Macroscopically the species is characterized by the pale, polished and smooth stipe and the fibrillose pileus. The pigmentation of the pileus is remarkably variable: in all specimens examined the intracellular pigment in the pileipellis and particularly in the upper part of the trama is rather distinct. Some specimens of the Netherlands' collection cited above, however, in addition show distinct encrustations on some hyphae of the pileipellis and the pileitrama. Horak (1976: 230) studied the Favre collection and found only encrusting pigments. Unfortunately I failed to locate the type, nor did I study the Favre collection. In consequence of my own observations I am inclined to consider the pigmentation in *E. rhombisporum* very variable. Largent's (1977: 147) concept of *Leptonia rhombispora* is according to the description a

misinterpretation. His Californian collection differs a.o. in colour, smell, and type of pileipellis (the last mentioned character proves it to be a true *Leptonia*).

ENTOLOMA section PAPILLATA (Romagn.) Noordeloos  
emend. Noordeloos

*Rhodophyllus* subgen. *Nolanea* sect. *Papillati* Romagn. in Bull. mens. Soc. linn. Lyon 43: 330. 1974. —  
*Entoloma* subgen. *Nolanea* sect. *Papillata* (Romagn.) Noordeloos in Persoonia 10: 246. 1979. — Holotype:  
*Entoloma papillatum* (Bres.) Dennis.

CHARACTERISTICS.—Pigment encrusting at least the narrowest hyphae of pileipellis and pileitrama; diffusely intracellular pigment sometimes present too but then never dominant; spores iso- to heterodiametrical with variable base; clamp-connections present; cystidia usually absent.

Section *Papillata* Romagn. is emended by including both the sections *Minuta* Romagn. and *Cosmeoxonema* Largent & Thiers on the level of subsections. In the present concept sect. *Papillata* covers about the same species as sect. *Nolanea* sensu Largent (1974: 1003), except for the exclusion of *Entoloma hirtipes* and related species (= sect. *Nolanea* s. restr. mihi). For details the reader is referred to Noordeloos (1980b).

Subsection PAPILLATA

Spores heterodiametrical with average Q per collection at least 1.2, with basal facet; lamellae usually with distinct grey and/or brown tinge, at least in mature specimens.

KEY TO THE SPECIES OF SUBSECTION PAPILLATA

- 1a. Basidia 2-spored. . . . . *E. cuspidifer*, p. 461
- b. Basidia 4-spored. . . . . 2
- 2a. Spores very obtusely rounded-angular,  $7.4-8.3(-9.4) \times 5.7-7.4(-8.0) \mu\text{m}$ ,  $Q = 1.07-1.2-1.3$ ; stipe strongly silvery striate. . . . . *E. ortonii*, p. 459
- b. Spores more pronouncedly angular and slightly to considerably larger; stipe smooth or only sparsely silvery striate. . . . . 3
- 3a. Spores  $7.9-10.4 \times 6.6-8.6(-9.0) \mu\text{m}$ ,  $Q = 1.1-1.2-1.3$ ; lamellae never dark reddish brown; stipe innately but distinctly silvery striate; smell farinaceous. . . . . *E. lucidum*, p. 458
- b. Spores with average Q per collection at least 1.3 and usually larger, if not so then lamellae fairly dark brown when mature, and stipe never striate and smell absent. . . . . 4
- 4a. Pileus moderately dark reddish brown or ochraceous- or greyish brown; lamellae pale then pink, sometimes with brown shade but hyphae of hymenophoral trama never encrusted; smell absent. . . . . 5
- b. Pileus fairly dark reddish or blackish brown; lamellae rather dark brown when mature; at least the narrowest hyphae of hymenophoral trama encrusted; smell farinaceous or absent. . . . . 6
- 5a. Lamellae tinged brown when mature; spores  $(8.7-9.0-10.4(-11.5)) \times (7.0-7.6-8.7) \mu\text{m}$ ; stipe sparsely silvery striate. . . . . *E. sericeonitens*, p. 459
- b. Lamellae white then pink, never with brown tinge; spores  $7-9.3(-10) \times 5.8-7 \mu\text{m}$ ; stipe smooth as if polished. . . . . (see subsection *Infularia*: *E. infula*), p. 503

6a. Lamellae pale when young, becoming dark reddish brown with age, of normal thickness, moderately distant; stipe usually without the reddish tinge of the pileus and usually paler, smooth or with spread silvery fibrils lengthwise; spores (9.0–)9.3–11.3(–13) × (6.7–) 7.1–8.0(–8.5)  $\mu\text{m}$ ; smell farinaceous.

*E. papillatum*, p. 454

b. Lamellae brown already when young, thickish and fairly distant; stipe concolorous with pileus, not striate, smooth; spores (7.8–)8.2–10.2(–10.6) × (5.7–)6.2–7.5(–7.9)  $\mu\text{m}$ ; smell absent.

*E. clandestinum*, p. 456

## 7. ENTOLOMA PAPILLATUM (Bres.) Dennis—Figs. 8a–d

*Nolanea papillata* Bres., *Fungi trident.* 1: 75, pl. 82. 1887. — *Rhodophyllum papillatus* (Bres.) J. Lange, *Fl. agar. dan.* 2: 101. 1936. — *Entoloma papillatum* (Bres.) Dennis in *Bull. Soc. mycol. Fr.* 69: 162. 1953. — *Nolanea mammosa* (L. ex Fr.) Quél. subsp. *papillata* (Bres.) Konr. & Maubl., *Icon. sel. Fung. pl.* 180 fig. 2. 1932.

*Agaricus mammosus* Fr. var. *tenuior* Fr., *Icon. sel.* 1: 113, pl. 98 fig. 4. 1867.

*Leptonia papillata* Velen., *České Houby:* 622. 1921.

*Leptonia mamillata* Velen., *České Houby:* 622. 1921.

MISAPPLIED NAMES.—*Agaricus mammosus* L. ex Fr. sensu Cooke, *Ill. Br. Fungi* 363(377). 1876. — *Rhodophyllum mammosus* (L. ex Fr.) Quél. sensu J. Lange in *Dansk bot. Ark.* 2(11): 36. 1921. — *Nolanea clandestina* (Fr.) Kummer sensu Bres. 1929, pl. 585 fig. 2; sensu Konr. & Maubl. 1936, pl. 178 fig. 2, non sensu Fr., J. Lange.

SELECTED ICONES & DESCRIPTIONS.—Arnolds & Noordeloos in *Fung. rar. Icon. col.* 12, pl. 91 fig. d. 1980. — Bresadola, *Iconogr. mycol.*, pl. 538. 1929. — Konr. & Maubl., *Icon. sel. Fung.*, pl. 180 fig. 2. 1932. — J. Lange, *Fl. agar. dan.* 2: 101, pl. 78D. 1936. — F. H. Möller, *Fungi Faeröes I:* 239. 1945. — Nathorst-Windahl in *Friesia* 8: 14, fig. 1. 1966. — Einhellinger in *Ber. bayer. bot. Ges.* 41: 106, figs. 19, 22, 23, and pl. 10f. 1969. — Gulden & M. Lange in *Norw. J. Bot.* 18: 13. 1971. — Höyland in *Blyttia* 35: 149. 1977.

CHARACTERISTICS.—Pileus dark sepia brown to almost black or fairly dark reddish brown; lamellae white or pallid at first, then grey-pink, finally (dark) reddish brown, moderately distant, thin; stipe usually (slightly) paler than pileus and less reddish, pruinose at apex, downwards with scattered innate silvery fibrils; smell and taste often distinctly farinaceous; spores 9–12.7  $\mu\text{m}$  long.

Pileus 6–28(–35) mm broad, acutely conical to conico-campanulate, rarely hemispherical or truncate-campanulate, expanding to conico-convex rarely convex, almost always with distinct papilla, with margin involute when young later straight, strongly hygrophanous, when moist dark sepia, blackish brown or reddish brown (10 YR 2/2, 3/2, 3/3, 4/4; 7.5 YR 3/2, 4/4; 5 YR 3/2 even 2.5 YR 3/2), not or only slightly paler at margin (10 YR 5/4, 6/4; 7.5 YR 4/4; 5 YR 5/4, 5/6), translucently striate often up to one half or two-thirds of the radius, on drying pallescent to (pale) sepia brown, often considerably less reddish than in moist state (7.5 YR 4/4, 4/2 or 6/4 at centre, rest 10 YR 5/3, 5/4, 6/3, 7/4, 7/6), smooth, shining, rarely minutely radially wrinkled. Lamellae L = 15–24, l = (1–)3–7(–9), moderately distant, thin, deeply emarginate to almost free, rarely adnate, triangular to broadly ventricose, pale grey or whitish when young, soon greyish pink, finally reddish brown (young 10 YR 8/3, 7/3, then 10 YR 4/4; 7.5 YR 6/4, 5/4, 4.4, 4/2 or 5 YR 7/3, 6/3, 5/3, 5/2, 4/3) with entire concolorous or slightly paler edge. Stipe 20–50 × 1–3 mm, cylindrical with base slightly swollen or not, straight or flexuose, sepia brown with paler apex, often missing red tinge of pileus (2.5 Y 7/6, 7/4; 10 YR 6/3, 5/3, 5/4, 5/6, 4/3, 4/2; 7.5 YR 3/2), pruinose-flocculose at apex, downwards with scattered, innate, silvery fibrils (hand lens), usually with white tomentum at base. Flesh thin-cartilagineous in pileus and stipe, concolorous with surface in pileus and cortex of stipe, paler in context of stipe. Smell usually strongly farinaceous, rarely weak. Taste farinaceous.

Spores (9.0–)9.3–11.3(–13) × (6.7–)7.1–8(–8.5)  $\mu\text{m}$ , Q = (1.15–)1.2–1.3–1.5(–1.6), L–D = 1.2–

2.5–5  $\mu\text{m}$ , variable, regularly 6- to irregularly 6–8-angled in side-view, with basal facet. Basidia 32–50  $\times$  8.5–13  $\mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular, composed of long cylindrico-fusiform cells 170–400  $\times$  10–28  $\mu\text{m}$ , mixed with narrow cylindrical hyphae with minutely encrusted walls. Pileipellis a simple cutis made up of radially arranged cylindrical 4–9 (–12)  $\mu\text{m}$  wide hyphae with minutely to coarsely encrusted walls gradually passing into pileitrama or with weakly developed subpellis of relatively short inflated cells 60–100  $\times$  12–30  $\mu\text{m}$ , lower trama regular, with long cylindrical-fusiform cells up to 450  $\mu\text{m}$  long and up to 29  $\mu\text{m}$  wide with walls encrusted by brown pigment, and this particularly abundant and dark in hypodermal layer. Clamp-connections abundant in hymenium, rare elsewhere.

**HABITAT & DISTRIBUTION.**—In poorly manured, extensively grazed grasslands, more rarely in open places in forest, at roadsides etc., not uncommon and widespread in the temperate and boreal and (sub-)alpine zones of the northern (and probably also the southern, see below) hemisphere. May to November.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Friesland, Isl. of Vlieland, Kroonpolders, 1 Nov. 1976, *C. Bas* 7147 and 2 Nov. 1976, *C. Bas* 7151; Isl. of Terschelling, Oosterend, Boschplaat, 30 July 1958, *C. Bas* 1514, Oosterend, Dazenplak, 3 Nov. 1978, *M. E. Noordeloos* 845 and 5 Nov. 1978, *M. E. Noordeloos* 862; prov. Drenthe, Beilen, 17 Oct. 1974, *E. Arnolds* 3245 (WBS); Gees, state forest, 5 June 1974, *K. Booy*; prov. Gelderland, Mook-Nijmegen, Heumensoord, 27 Aug. 1977, *T. Kuypers*.

NORWAY, Oppland, Dombås, 5 Aug. 1977, *M. E. Noordeloos* 393. — GREAT BRITAIN, Wales, Lake Vyrnwy, 30 Aug. 1979, *E. Kits van Waveren*. — GERMAN FEDERAL REPUBLIC, Westphalen, Detmold, Merlsheim, 6 Oct. 1976, *C. Bas*. — FRANCE, dept. Oise, Coye-la-Forêt, 11 Sept. 1945, *H. Romagnesi* 116 (Herb. Romagn., PC). — CZECHOSLOVAKIA, Bohemia, Mnichovice, July 1919, *J. Velenovský* (PRC, type of *L. mamillata*), ibidem, July 1919, *J. Velenovský* (PRC, type of *L. papillata*).

*Entoloma papillatum* is a species very variable in colour and habit as well as in size and shape of the spores. The presence of red tinges in pileus, stipe and lamellae makes it sometimes difficult to distinguish it from *E. clandestinum*. In the present work *E. clandestinum* is defined as a species with the following characters: fairly dark (red-)brown tinges in all parts of the carpophore, distant, thickish lamellae, a smooth stipe without innate silvery fibrils, and relatively small spores. As a consequence dark pigmented specimens of *E. papillatum* can be distinguished from *E. clandestinum* in having 'normally' thin lamellae, a minutely silvery striate stipe (hand lens) and (considerably) larger spores.

The strong resemblance between *E. papillatum* and *E. clandestinum* is reflected in the confusion in literature. As suggested by Romagnesi (Kühn. & Romagn., 1953: 186), Bresadola's (1929, pl. 585 fig. 2) plate of *Nolanea clandestina* most probably represents *Rhodophyllus papillatus* sensu J. Lange and Kühn. & Romagn., which interpretation comes very close to my own concept. In the Bresadola Herb. (S) 2 collections were found labelled as *E. clandestinum*, which appeared to fit in perfectly with our present concept of *E. papillatum*.

The plates and descriptions of Konrad & Maublanc (1936, pl. 178 fig. 2, as *Nolanea clandestinum*, and pl. 180 fig. 2, as *N. mammosa* var. *papillata*) illustrate the variability of *E. papillata*. In my opinion the first plate represents a dark, reddish tinged form of *E. papillatum* rather than *E. clandestinum*.

The only plates very well illustrating *E. clandestinum* in our concept can be found in J. Lange (1936, pl. 78C) and in Arnolds & Noordeloos (1980, pl. 91 fig. e).

The size and shape of the spores vary considerably in *E. papillatum*, often within one collection. This is illustrated in Fig. 8d and in earlier published figures (Noordeloos 1979a: 259, fig. 22 and 262, fig. 28). In the type of *Leptonia papillata* Velen. only the simplest type of spore is found, whereas in *L. mamillata* Velen. the more elongate-complex type is predominant.

Outside Europe *E. papillatum* has been recorded from the U.S.A. (Hesler, 1967: 100) and South America (Dennis, 1953: 162; Horak, 1978: 75). I did not study material from these localities, but judging from these descriptions it seems probable that *E. papillatum* also occurs outside the temperate-boreal regions of Europe.

8. *Entoloma clandestinum* (Fr.) Noordeloos,  
comb. nov.—Figs. 7a–e

*Agaricus clandestinus* Fr., Syst. mycol. 1: 206. 1821 (basionym). — *Nolanea clandestina* (Fr.) Kumm., Führ. Pilzk.: 95. 1871. — *Rhodophyllus clandestinus* (Fr.) Quél., Enchir.: 64. 1886.

SELECTED ICONES AND DESCRIPTIONS.—Arnolds & Noordeloos in Fung. rar. Icon. col. 12: pl. 91 fig. e. 1980.  
— J. Lange, Fl. agar. dan. 2: 101, pl. 78C. 1936.


CHARACTERISTICS.—Pileus rather dark blackish brown often with distinct red tinge; lamellae distant, thickish, fairly dark brown already when young; stipe (sub-)concolorous with pileus; smell and taste indistinct; spores 7.8–10.2  $\mu\text{m}$  long.

Pileus 6–22(–30) mm broad, conical or hemispherical, then slightly expanding to conico-convex, finally convex, with or without small papilla, with margin involute then expanding, hygrophanous, when moist rather dark blackish brown often with red tinge (2.5 YR 2.5/2; 5 YR 2/2–3/2; 7.5 YR 2/2–3/2; Meth. 6F4), not or only slightly paler at margin (e.g. 7.5 YR 4/4), mostly more or less unicolorous, translucently striate at least at margin, on drying pallescent to grey-brown (e.g. Meth. 6E6/5), glabrous, often ornated with minute warts or minutely radially wrinkled, shining. Lamellae L = 12–20, l = 1–3(–7), distant, thickish, variably inserted, broadly adnate or emarginate, uncinuate to almost free, narrowly ventricose, up to 5 mm broad, dark grey-brown already when young, darkening with age, mostly without real pink tinge (7.5 YR 5/4, 5 YR 4/3; Meth. 6D5/E5, 6E5), sometimes veined, sometimes anastomosing, with concolorous or slightly paler edge. Stipe 18–40  $\times$  1–2.3 mm, cylindrical, often flexuous, broadened towards base or not, concolorous with pileus or paler (Meth. 6E6; 5 YR 2.5/2, 3/3, 3/2, 3/1), not striate, not fibrillous, glabrous or at apex minutely pruinose, rarely entirely pruinose (*Noordeloos 187*), at base often with white tomentum, solid, then narrowly fistulose. Flesh thin-membranaceous but firm in pileus, concolorous with surface, in stipe firm and cartilaginous; cortex with same colour as surface, context paler. Smell and taste weak, inconspicuous, mild.

Spores (7.8–)8.2–10.2(–10.6)  $\times$  (5.7–)6.2–7.5(–7.9)  $\mu\text{m}$ , Q = 1.2–1.3–1.4(–1.5), L–D = 1.5–2.5–3.5  $\mu\text{m}$ , mostly rather regularly 6-angled in side-view, with basal facet. Basidia 30–50  $\times$  8–15  $\mu\text{m}$ , 2–4-spored. Cystidia absent. Hymenophoral trama regular, composed of fusiform cells (170–)220–450(–500)  $\times$  10–29  $\mu\text{m}$ , with brown walls, sometimes with additional intracellular pigment. Pileipellis a simple cutis made up of 4.5–11  $\mu\text{m}$  wide, cylindrical hyphae with coarsely encrusted walls. Pileitrama regular, composed of long, cylindrical to fusiform cells as in hymenophoral trama. Clamp-connections abundant in hymenium, rare or absent elsewhere.

HABITAT & DISTRIBUTION.—In frondose forest, often on clear spots, roadsides etc., but also in poorly manured grasslands. Rare but widespread. Recorded from the Netherlands, Denmark, Great Britain, German Federal Republic, France, Italy and Czechoslovakia. June–November.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Anloo, Gasterse duinen, 28 Oct. 1973, *E. Arnolds 3288* (WBS); prov. Gelderland, Winterswijk, Bek-en-Delle, 20 Sept. 1976, *M. E. Noordeloos 187*; prov. Noord-Holland, IJmuiden, Duin & Kruidberg, 29 Sept. 1975, *E. Kits van Waveren*; Aerdenhout, Naaldenveld, 2 Oct. 1975, *E. Kits van Waveren*; Castricum, 4 Oct. 1975, *E. Kits van Waveren*.



Figs. 7a-c. *Entoloma clandestinum*. — Habit and spores (7a from Kits van Waveren, 2 Oct. 1975; 7b, 7c from Noordeloos 187; 7c-d from Kits van Waveren, 4 Oct. 1975).

Fig. 8a-d. *Entoloma papillatum*. — Habit and spores (8a from Noordeloos 421; 8b from Noordeloos 418; 8c-d from Noordeloos 420).

GERMAN FEDERAL REPUBLIC, Bayern, Sugenburg bei Abensberg i.d. Hallertau, 10 Oct. 1968, A. Einhellinger (M). — FRANCE, dept. Oise, Lamorlaye, 24 Aug. 1950, H. Romagnesi (Herb. Romagn. 50.195, PC); loc. unknown, 14 Nov. 1977, anonym (Herb. Romagn. 77.378, PC). — ITALY (Trento?), Herb. Bres., June 1900 and July 1901 (S).

*Entoloma clandestinum* is closely related to *E. papillatum* but can be distinguished by the distant, thickish lamellae, smooth stipe, and smaller spores (see also the discussion under *E. papillatum*, p. 455).

9. *ENTOLOMA LUCIDUM* (P. D. Orton) Moser—Figs. 9a–e

*Nolanea lucida* P. D. Orton in Trans. Br. mycol. Soc. 43: 331. 1960. — *Rhodophyllus lucidus* (P. D. Orton) Moser in Gams, Kl. KryptogFl. 3. Aufl., 2(b/2): 166. 1967. — *Entoloma lucidum* (P. D. Orton) Moser in Gams, Kl. KryptogFl. 4. Aufl., 2(b/2): 206. 1978.

CHARACTERISTICS.—Pileus and stipe dark sepia or horn brown; stipe striate; spores subisodiametrical,  $Q=1.2$  on the average per collection.

Pileus 18–32 mm broad, conico-convex soon expanding to convex, finally flattened, with faint papilla, with straight margin, strongly hygrophanous, when moist rather dark sepia or umber brown (10 YR 4/3, 3/3), paler towards margin (10 YR 5/3, 6/3), translucently striate at least up to one half of the radius, pallescent on drying to pallid grey-brown or ochraceous-grey (10 YR 7/3, 7/2), innately radially fibrillose, dull or shining. Lamellae  $L=$  up to 36,  $l=1-3$ , adnate to almost free, segmentiform to ventricose, pallid grey then pinkish grey, finally brownish pink (7.5 YR 8/2, 8/4 then 7/4, 6/4, finally 5 YR 6/4), with concolorous, almost entire edge. Stipe 36–58  $\times$  1.5–3 mm, cylindrical, mostly slightly curved and broadened at base, solid then narrowly fistulose, concolorous with or, particularly towards base, paler than pileus, strong brown to grey- or yellowish brown (10 YR 5/4, base 7/6), minutely silvery striate lengthwise. Flesh concolorous with surface in cortical layer, in context paler, brittle in pileus, more rigid in stipe but never cartilaginous. Smell and taste strong, rancid-farinaceous.

Spores 7.9–10.4  $\times$  6.6–8.6(–9.0)  $\mu\text{m}$ ,  $Q=1.1-1.2-1.3$ ,  $L-D=(0.6-1.0-1.5-2 \mu\text{m})$ , (5-)6-7(–8)-angled in side-view, probably with basal facet. Basidia 24–40  $\times$  9–15  $\mu\text{m}$ , 4-, a few 2-spored. Cystidia absent. Hymenophoral trama regular; cells (140-)158–260  $\times$  14–26  $\mu\text{m}$ , mixed up with narrow, cylindrical hyphae. Pileipellis a cutis made up of 3–11(–20)  $\mu\text{m}$  wide cylindrical hyphae with brownish, sometimes minutely encrusted walls and sometimes with a pale, diffuse, intracellular pigment, gradually passing into pileitrama. Pileitrama regular, composed of inflated hyphae up to 32  $\mu\text{m}$  wide with minutely encrusted walls. Clamp-connections abundant in hymenium, rare in other tissues.

HABITAT & DISTRIBUTION.—In grasslands, rare. Known from the Netherlands and Great Britain.

COLLECTIONS EXAMINED.—NETHERLANDS: IJsselmeerpolders, Oostelijk Flevoland, Roggebootszand, 1 July 1978, J. Hoogschagen; idem, Revebos, 14 June 1979, G. J. M. G. Tjallingii-Beukers; prov. Gelderland, Winterswijk, Korenburgerveen, 29 Sept. 1973, C. Bas 6164.

GREAT BRITAIN, Devon, Membury, 1 Nov. 1957, P. D. Orton 1302 (K, holotype).

NOTES ON THE TYPE.—The type collection consists of two fragments of pilei, obviously belonging to two different carpophores with the following microscopical characters.

Spores of first specimen: (7.3-)7.9–10.2  $\times$  (6.6-)6.8–7.3(–8.6)  $\mu\text{m}$ ,  $Q=1.08-1.11-1.25$ ,  $L-D=0.6-1.1-1.7 \mu\text{m}$ ; spores of second specimen: 7.9–10.2  $\times$  (6.3-)6.8–7.9(–8.5)  $\mu\text{m}$ ,  $Q=1.12-1.2-1.27$ ,  $L-D=1.0-1.8-2.7 \mu\text{m}$ . In both specimens the spores are irregularly 5–6–7-sided in side-view. Basidia 29–38  $\times$  9–11, 3  $\mu\text{m}$ , 4-spored. Pileipellis a cutis made up of narrow cylindrical hyphae with minutely encrusted walls. Pileitrama regular, with long, subcylindrical to inflated cells with minutely encrusted walls, particularly those of the narrowest hyphae. Clamp-connections seen in hymenium.

Owing to the scarcity of material (most of the collections studied were poor in specimens) it is difficult to get a clear picture of the variability of *E. lucidum*. The dark colours and slightly more isodiametrical spores distinguish it from *E. minutum*.

*Entoloma sericeonitens* is very similar, but differs by the slightly more elongate spores and the aspect of pileal surface (see below).



10. *Entoloma sericeonitens* (P. D. Orton) Noordeloos,  
*comb. nov.*—Fig. 12

*Nolanea sericeonitens* P. D. Orton in Trans. Br. mycol. Soc. 43: 333. 1960 (basionym).

CHARACTERISTICS.—Pileus reddish brown, strongly pallescent and fibrillous-aeriferous on drying; spores  $9-11 \times 7-8 \mu\text{m}$ ,  $Q = 1.2-1.3-1.4$ .

Pileus up to 30 mm broad, conical to conico-campanulate, with margin straight at first but reflexed with age, hygrophanous, when moist reddish brown, slightly paler at margin, translucently striate up to centre, on drying strongly pallescent to pale grey-brown. Lamellae  $L =$  about 28,  $l = 1-3$ , almost free, ventricose, brownish pink, with concolorous edge. Stipe  $50 \times 2$  mm, cylindrical, concolorous with pileus, slightly silky striate lengthwise, white tomentose at base. Flesh concolorous with surface, membranaceous-brittle. Smell and taste absent.

Spores  $(8.7-9.0-10.4(-11.5) \times (7.0-7.6-8.7 \mu\text{m})$ ,  $Q = 1.2-1.3-1.4$ ,  $L-D = 1.0-2.0-3.0 \mu\text{m}$ , irregularly 6-angled in side-view, probably with basal bacet. Basidia  $22-37 \times 7.5-11.5 \mu\text{m}$ , 4-spored. Hymenophoral trama regular, with cells  $180-300 \times 15-23 \mu\text{m}$ , cylindrical to inflated. Pileipellis a simple cutis made up of radially arranged, cylindrical hyphae  $4-8.7 \mu\text{m}$  wide with pale, minutely encrusted walls. Pileitrama regular with long, inflated cells  $175-340 \times 12-24 \mu\text{m}$ . Clamp-connections frequent in hymenium, scattered in trama of lamellae and pileus, rare in pileipellis.

HABITAT & DISTRIBUTION.—In grasslands (type); the Netherlands' collection was made on peaty soil. Rare; up till now known from the type locality and one locality in the Netherlands.

COLLECTIONS EXAMINED.—NETHERLANDS, prov. Gelderland, Winterswijk, Korenburgerveen, 29 Sept. 1973, G. J. M. G. Tjallingii-Beukers.

GREAT BRITAIN, Devon, Dawlish Warren, 12 Dec. 1956, T. J. Wallace (holotype, K).

NOTE ON THE TYPE.—The type collection consists of fragments of one specimen with the following characters.

Spores  $9.0-11.0(-11.2) \times 7.0-8.3(-8.7) \mu\text{m}$ ,  $Q = 1.2-1.3-1.36$ ,  $L-D = 1.7-2.3-3.0$ . Basidia  $27-34 \times 10.4-12.7 \mu\text{m}$ , 4-spored. Hymenophoral trama regular with cylindrical cells  $190-370 \times 11-24 \mu\text{m}$ . Pileipellis a thin cutis made up of parallel,  $3-8(-11) \mu\text{m}$  wide, cylindrical hyphae with minutely encrusted walls, gradually passing into trama. Pileitrama regular, with the same long cylindrical-inflated cells as in hymenophoral trama. Clamp-connections present in hymenium.

*Entoloma sericeonitens* resembles *E. lucidum* very strongly. It differs by the slightly more reddish tinged pileus, the aeriferous aspect of dry pileus, the lack of smell and the slightly larger and more heterodiametrical spores. The shape of the spores, however, is shown to be rather variable in *E. lucidum*. Unfortunately the observations above are based on very poor material, the Netherlands' collection consisting of one specimen. Further studies are necessary to prove that both species are not conspecific.

*Entoloma minutum* differs from *E. sericeonitens* by the much paler pileus and lamellae and the distinctly dihedral base of the spores.

11. *Entoloma ortonii* Arnolds & Noordeloos—Fig. 10


*Entoloma ortonii* Arnolds & Noordeloos in Persoonia 10: 292. 1979 (name change). — *Nolanea farinolens* P. D. Orton in Trans. Br. mycol. Soc. 43: 330. 1960. — *Rhodophyllus farinolens* (P. D. Orton) Moser in Gams, Kl. KryptogFl. 3. Aufl., 2(b/2): 166. 1967. — *Entoloma farinolens* (P. D. Orton) Moser in Gams, Kl.

KryptogFl. 4. Aufl., 2(b/2): 206. 1978 (non *Entoloma farinolens* Horak, 1973).

SELECTED ICONES AND DESCRIPTION.—Arnold & Noordeloos in Fung. rar. Icon. col. 12, pl. 92 fig. b. 1980.

CHARACTERISTICS.—Pileus flattened, never truly papillate; stipe strongly silky striate lengthwise; spores rather rounded-angular and thin-walled.

Pileus 24–35 mm broad, convex to plano-convex, sometimes with very weak umbo or slightly depressed at centre, with margin slightly involute at first then straight, hygrophanous, when moist dark grey-brown at centre (Meth. 5E6; 10 YR 3/2–4/2), paler towards margin (Meth. 5D5; 10 YR 4/3–5/3), translucently striate up to centre, when dry brown-grey to ochraceous grey, dull. Lamellae L = 18–23, l = 3–7, adnate or emarginate, segmentiform to subventricose, brown-grey with pink tinge (Meth. 5C4; 7.5 YR 8/4–6/4), paler towards edge. Stipe 50–95 × (2–)2.5–5 mm, cylindrical, sometimes flexuous, with or without broadened base, grey-brown (10 YR 6/4, 5/4, 4/4), strongly silvery striate lengthwise, sometimes twisted, pruinose at apex, white tomentose at



Figs. 9a–e. *Entoloma lucidum*. — Habit and spores (9a from Bas 6164; 9b, 9e from Tjallingii-Beukers, 14 June 1979; 9c from Hoogschagen, 1 July 1978; 9d from holotype).

Fig. 10. *Entoloma ortonii*. — Spores (from holotype).

Figs. 11a–b. *Entoloma sphaerocystis*. — Habit and spores (11a–b from holotype).

Fig. 12. *Entoloma sericeonitens*. — Spores (from holotype).

base. Flesh in pileus rather firm, tough, watery grey-brown, in stipe more fibrillous, rigid, grey- to ochraceous brown. Smell and taste strongly rancid-farinaceous.

Spores  $7.4-8.5(-9.4) \times 5.7-7.4(-8.0) \mu\text{m}$ ,  $Q = 1.07-1.2-1.3(-1.4)$ ,  $L-D = (0.6-)1.5-1.7-2.3 \mu\text{m}$ , 5-6-angled in side-view, with very much rounded angles, probably with basal facet. Basidia  $25-40 \times 10.5-14 \mu\text{m}$ , 4- (rarely 2-)spored. Cystidia absent. Hymenophoral trama regular, with cylindrical to inflated cells up to  $450 \mu\text{m}$  long and  $32 \mu\text{m}$  wide. Pileipellis a thin cutis made up of 4-12  $\mu\text{m}$  wide cylindrical hyphae with minutely encrusted walls and diffusely intracellular pigment. Pileitrama regular, with cells up to  $500 \mu\text{m}$  long and  $10-23 \mu\text{m}$  wide, in upper layers with pale brown, minutely encrusted walls. Clamp-connections present in hymenium.

**HABITAT & DISTRIBUTION.**—In grasslands or grassy spots in frondose forest, rare. Known from the Netherlands and Great Britain.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Drenthe, Anloo, CRM-reserve 'de Burgwalen', 4 July 1974, *E. Arnolds 3388* (WBS); prov. Zuid-Holland, Warmond, estate 'Huys te Warmond', 12 Oct. 1976, *C. Bas 7094*.

GREAT BRITAIN, Inverness-shire, Plodda Falls, 22 Aug. 1955, *P. D. Orton* (K, holotype).

**NOTES ON THE TYPE.**—The holotype collection consists of 5 specimens with the following microscopical characters.

Spores  $7.3-8.5(-9.0) \times (5.7-6.2-6.8(-7.3)) \mu\text{m}$ ,  $Q = (1.08-)1.1-1.2-1.3(-1.4)$ ,  $L-D = (0.6-)0.9-1.3-2.3 \mu\text{m}$ , (5-)6-angled in side-view with very much rounded angles. Basidia  $(28.5-)31-40 \times 8.1-10.5 \mu\text{m}$ , 4-spored. Cystidia absent. Pileipellis a thin cutis made up of 4-9  $\mu\text{m}$  wide cylindrical hyphae with minutely encrusted walls. Clamp-connections frequent in hymenium.

*Entoloma ortonii* has very distinctive rounded-angular, thin-walled spores, which are also smaller than in *E. lucidum* and *E. sericeonitens*. *Entoloma juncinum* has more isodiametrical, more pronouncedly angled spores.

## 12. *Entoloma cuspidifer* (Kühn. & Romagn.) Noordeloos, *comb. nov.*

—Figs. 13a-c

*Rhodophyllum cuspidifer* Kühn. & Romagn., *Fl. anal.*: 189. 1953. — *Nolanea cuspidifer* (Kühn. & Romagn.) P. D. Orton in *Trans. Br. mycol. Soc.* **43**: 179. 1960.

*Agaricus junceus* Fr. var. *cuspidatus* Fr., *Icon. sel.* 1: pl. 99 fig. 2. 1875. — *Nolanea juncea* Fr. var. *cuspidata* (Fr.) Favre in *Bull. Soc. mycol. Fr.* **52**: 137. 1936. — *Rhodophyllum cuspidatus* (Fr.) Favre, *Assoc. fung. hautes-marais*: 44. 1948, non *R. cuspidatus* Pat. (1924) nec *Entoloma cuspidatus* (Peck, 1870) Sacc. 1887.


**SELECTED DESCRIPTIONS.**—Favre, l.c. 1948. — Einhellinger in *Ber. bayer. bot. Ges.* **47**: 125, fig. 4. 1976.

**CHARACTERISTICS.**—Pileus pointedly conical, only slightly expanding, sepia, strongly pallescent on drying; stipe paler with pruinose-subflocculose apex; basidia 2-spored; capitate hairs at apex of stipe.

Pileus 17-35 mm broad, up to 18 mm high, acutely conical with small papilla, only slightly expanding to conico-campanulate with age, with margin slightly inflexed but not involute when young, later straight, strongly hygrophanous, when moist translucently striate up to one half of the radius, (greyish) sepia to greyish bistre, only slightly paler at margin (10 YR 4/4, towards margin 10 YR 5/4-6/4, on drying pallescent to very pale brown (10 YR 7/4-7/3), with radial streaks from centre outward, when moist already with some radially arranged silvery fibrils, becoming strongly silvery fibrillous-silky on drying. Lamellae  $L =$  about 30,  $l = 3-5$ , (almost) free, triangular-ventricose, up to 7 mm broad, exceeding mature pilei, grey then grey-pink, finally with brown tinge (10 YR 7/2, 7/3, 7/4 then 7.5 YR 7/4 (6/4), with eroded concolorous

edge. Stipe 72–80 × 2.5–4 mm, cylindrical, straight, with equal base, paler and more yellow than pileus (10 YR 8/4, 7/4, 8/6), browner at base (10 YR 6/4, 5/4), upper 15 mm minutely pruinose-downy, downwards with a few longitudinal silvery fibrils, base white tomentose, solid, then narrowly fistulose. Flesh relatively firm in pileus and concolorous with surface, fibrillous in stipe, concolorous with surface in cortex, pale in context. Smell strongly raphanoid. Taste rancid-raphanoid.

Spores 10–13 × 9.3–11 μm, Q = 1.05–1.2–1.4, L–D = 1.2–2.5–4 μm, 5–6-sided in side-view, structure of base difficult to interpret. Basidia 25–41 × 9–15 μm, mostly broadly clavate, 2-spored, rarely 4-spored. Cheilocystidia usually absent, in one specimen scattered, 42–60 × 12–17 μm, clavate-subcapitate. Hymenophoral trama with long, only slightly inflated cells 280–380 × 14–32 μm. Pileipellis a poorly differentiated cutis made up of 2.5–10(–14) μm wide cylindrical hyphae with pale, very minutely encrusted walls. Subpellis composed of broad, relatively short cells about 135–170 × 32–42 μm, gradually passing into pileitrama. Pileitrama regular with much



Figs. 13a–e. *Entoloma cuspidifer*. — Habit, spores, basidia, cheilocystidia and stipitepellis. (All figs. from Noordeloos 993.)

longer cells, up to 450  $\mu\text{m}$  long and 15–36  $\mu\text{m}$  wide, with pale, sometimes slightly encrusted walls particularly in the narrowest hyphae near septa, in some superficial hyphae with additional diffuse, pale intracellular pigment. Stipitipellis with an arachnoid covering of 2–5  $\mu\text{m}$  wide cylindrical branching hyphae with long terminal cells with a distinct capitulum 4–8  $\mu\text{m}$  wide. Clamp-connections frequent in all tissues studied.

**HABITAT & DISTRIBUTION.**—In acid, peaty habitats, in *Sphagnum*, but in the Netherlands found in a marshy vegetation with *Salix repens*, *S. aurita*, and *Menyanthes trifoliata* in a dune valley near the coast; known to occur in Norway, Great Britain, France, Switzerland, and German Federal Republic.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Noord-Holland, Callantssoog, Nature-reserve 'Zwanewater', 25 Aug. 1979, M. E. Noordeloos 993.

NORWAY, prov. Telemark, Nesland near Rauland, 26 July 1975, F. Tjallingii & G. J. M. G. Tjallingii-Beukers. — GERMAN FEDERAL REPUBLIC, Bayern, Ascholding, Eglinger Moor, 3 Aug. 1976, A. Einhellinger (M).

*Entoloma cuspidifer* has been excellently delineated by Favre (1936, l.c.) in a description fitting perfectly our material. Also Einhellinger's (1976, l.c.) illustrations are well qualified. Both authors collected *E. cuspidifer* in Sphagneta. The Netherlands' locality has a different vegetation: marshy ground with *Salix repens*, *S. aurita*, *Menyanthes trifoliata*, and mosses, but without *Sphagnum*, in a dune area with a relatively low pH. Obviously *E. cuspidifer* is not restricted to *Sphagnum*, in contrast with many other agarics found on species of that genus of mosses.

The pale colours and two-spored basidia may cause confusion with *E. cetratum*, but the raphanoid smell and completely different spores make it easy to recognize *E. cuspidifer*. *Entoloma tenellum* Favre is closely related, but differs by the mostly 4-spored basidia and slightly different spores.

**Entoloma** subsection **Fibulata** (Largent) Noordeloos, *comb. nov.*  
emend. Noordeloos

*Nolanea* (Fr.) Kumm. subsect. *Fibulatae* Largent in *Mycologia* 66: 1008. 1974. — Type: *Nolanea juncea* (Fr.) Quél.

Pigment minutely encrusting hyphae of pileipellis and pileitrama, and at the same time diffusely intracellular pigment present particularly in subpellis of pileus; spores isodiametrical to subglobose,  $Q = 1.0\text{--}1.1\text{--}1.2$ .

*Entoloma* subsection *Fibulata* comprised in its original concept species which are treated here in subsection *Papillata* and section *Nolanea*. In the present work subsection *Fibulata* is emended by restricting it to those species in section *Papillata* which have isodiametrical spores and two types of pigment.

KEY TO THE SPECIES OF SUBSECTION FIBULATA

- 1a. Pileus moderately dark grey-brown or slightly ochraceous tinged grey-brown, not entirely smooth but already when moist with aciferous silvery fibrils, becoming strongly lustrous fibrillose on drying; stipe strongly silvery striate lengthwise; smell weak, spermatial, not farinaceous. . . . *E. nitens*, p. 466

- b. Pileus fairly dark grey-brown, smooth and glabrous, when moist becoming silvery shining, innately fibrillous on drying; stipe smooth or with some scattered silvery longitudinal fibrils; smell farinaceous-rancid. . . . . *E. juncinum*, p. 464

### 13. ENTOLOMA JUNCINUM (Kühn. & Romagn.) Noordeloos—Figs. 14a–d

*Rhodophyllus juncinus* Kühn. & Romagn. in Rev. Mycol. 19: 5. 1954. — *Nolanea juncina* (Kühn. & Romagn.) P. D. Orton in Trans. Br. mycol. Soc. 43: 179. 1960. — *Entoloma juncinum* (Kühn. & Romagn.) Noordeloos in Persoonia 10: 255. 1979.

*Nolanea globispora* Velen., České Houby: 628. 1921; non *Entoloma globisporum* Morg.-Jones, 1971.

*Nolanea zonata* Velen., Novitates mycologicae novissimae: 78. 1947; non *Entoloma zonatum* Hesler, 1967.

MISAPPLIED NAMES.—*Rhodophyllus juncus* sensu J. Lange in Dansk bot. Ark. 2: 37–38. 1921. — *Nolanea proleteria* sensu Ricken, Blätterpilze: 299, pl. 74 fig. 4. 1910.

CHARACTERISTICS.—Pileus rather dark brown, convex, weakly papillate or not, translucently striate when moist; lamellae rather dark brown with pink tinge; stipe concolorous with pileus, smooth, not or only weakly striate; spores isodiametrical.

Pileus (10–)15–35 mm broad, conico-convex soon expanding to plano-convex, obtuse or papillate, with margin slightly involute when young but soon straight, slightly undulating and/or splitting with age, strongly hygrophanous, when moist rather dark reddish or greyish brown at centre and on striae, slightly paler towards margin (centre 10 YR 3/2, 4/2, 3/3; 7.5 YR 3/2, 4/2, towards margin 10 YR 4/4, 5/3, 5/4, 6/3, 7/2; 7.5 YR 6/3, 5/3), translucently striate from margin up to one half of the radius at least, often up to centre; on drying strongly pallescent towards pale yellowish-greyish brown (10 YR 7/2, 7/3, 8/3, 8/4), glabrous, smooth or minutely fluffy-rugulose at centre when dry, sometimes with concentric rings when drying. Lamellae L = 15–30, l = (1–)3–7, deeply emarginate to almost free, ventricose, up to 7 mm broad, often exceeding the pileus, rather dark grey-brown with pink tinge with age (10 YR 5/3, 5/4; 7.5 YR 5/4, 6/4; 5 YR 5/3, 5/4, 6/3), with entire concolorous edge, on drying distinctly pallescent from edge upwards; rarely veined and/or anastomosing. Stipe 45–80 × 1–4 mm, cylindrical or flattened with longitudinal groove, sometimes flexuous, slightly broadened at base or not, concolorous or slightly paler than pileus, dark grey-brown or golden brown, sometimes with reddish tinge (10 YR 4/3, 3/4, 4/4; 7.5 YR 4/2, 3/2; 5 YR 4/2, 3/3), glabrous, sometimes minutely pruinose at apex, smooth and shining or innately silvery fibrillous under lens, often with white tomentum at base, solid then narrowly fistulose. Flesh when moist concolorous with surface, pallescent on drying, brittle in pileus, firm-subcartilaginous in stipe. Smell spontaneously farinaceous, sometimes weak but then distinct after cutting. Taste strongly farinaceous-rancid.

Spores 8.0–10.4 × (7.0–)7.2–9.0(–9.3)  $\mu\text{m}$ , Q = 1.0–1.1–1.2, L–D = 0.5–1.5  $\mu\text{m}$ , 5–6-sided isodiametrical in side-view with pronounced angles and protruding apiculus; basal diedron sometimes well developed. Basidia 28–45(–56) × 10–16  $\mu\text{m}$ , Q = 2.3–4.0, 4-spored. Cystidia absent. Hymenophoral trama regular, composed of cylindrical or inflated cells (170–)230–540 × 12.5–42  $\mu\text{m}$ , mixed with narrow cylindrical hyphae. Pileipellis a thin cutis made up of 2–4–14  $\mu\text{m}$  wide cylindrical hyphae with scattered fusiform to clavate terminal cells, up to 20  $\mu\text{m}$  broad; with pale brown and minutely encrusted walls, sometimes with additional pale, diffuse intracellular pigment. Pileitrama regular, composed of long cylindrical to slightly inflated hyphae with cells up to 450  $\mu\text{m}$  long and 12–32  $\mu\text{m}$  wide, with pale, sometimes minutely encrusted walls. Clamp-connections frequent in hymenium, rare elsewhere.

HABITAT & DISTRIBUTION.—In humus-rich frondose forests, solitary; rarely also in grasslands. Widespread and locally constant and common. July–November.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Westerbork, 'de Reitema', E. of Elp, 13 Oct. 1976, *E. Arnolds 3648* (WBS); prov. Overijssel, Delden, along Springendaalse beek, 11 Sept.

1976, *J. Schreurs*; idem, estate 'Singraven', 18 June 1960 and 19 Sept. 1969, *E. Kits van Waveren*; idem, estate 'Twickel', 28 Sept. 1974, *E. Kits van Waveren*; Ommen, estate 'Stekkenkamp', 13 Sept. 1979, *J. Schreurs & T. Boekhout*; idem, near Castle, 18 Oct. 1969, *E. Kits van Waveren*; idem, near 'Ada's Hoeve', 26 Sept. 1964 and 19 Oct. 1969, *E. Kits van Waveren*; prov. N o o r d - H o l l a n d, Noordwijkerhout, Vogelenzang, 13 Nov. 1962, *C. Bas* 2898; idem, State Forest, 24 Nov. 1977, *M. E. Noordeloos* 575; Velzen, 'Duin en Kruidberg', 13 Nov. 1962, *C. Bas* 2913; Castricum, 'Naaldenveld', 19 Oct. 1964, *E. Kits van Waveren*; Overveen, 'Koningshof', 16 Nov. 1976, *M. E. Noordeloos* 266; Callantsoog, 'Zwanewater', 13 Nov. 1977, *C. Bas*; prov. Z u i d - H o l l a n d, Isl. Goeree, Westduinen, Oct. 1979, *M. Brand*; prov. N o o r d - B r a b a n t, Nuenen, Broeklanden, 15 Sept. 1979, *M. E. Noordeloos* 1020; Drunen, along Drongelens kanaal, 1 Nov. 1970, *P. B. Jansen* 70-182.

BELGIUM, prov. Namur, Rochefort, near Resteigne, 1 Oct. 1977, *M. E. Noordeloos* 476; St. Hubert, 'Poix', 24 Sept. 1975, *P. B. Jansen* 75-243. — GERMAN FEDERAL REPUBLIC, Teutoburgerwald, Detmold, Sandebeck, 5 Oct. 1976, *M. E. Noordeloos* 214; idem, Externsteinerwald, 7 Oct. 1976, *C. Bas*; idem, Beller Holz, 4 Oct. 1976, *M. E. Noordeloos* 217, 218, 219; idem, Merlsheimerwald, 6 Oct. 1976, *C. Bas*; Hörter, Elschenbergerwald, 8 Oct. 1976, *C. Bas*. — FRANCE, dept. Oise, Chantilly, 16 Aug. 1954, *H. Romagnesi* 54.136 (lectotype, Herb. Romagn., PC); Chamont, 19 July 1953, *H. Romagnesi* 53.95 (Herb. Romagn., PC); Montlognon, 18 Aug. 1954, *H. Romagnesi* 54.145 (Herb. Romagn., PC); Luzarches, 24 Aug. 1946, *H. Romagnesi* 46.271 (Herb. Romagn., PC). — CZECHOSLOVAKIA, Bohemia, Mnichovice, July 1919, *J. Velenovský* (lectotype of *Nolanea globispora* Velen., PRC); idem, July 1940, *J. Velenovský* (holotype of *Nolanea zonata* Velen., PRM 154454).

*Entoloma juncinum* is very well depicted by J. Lange (1936, pl. 79F) as *Rhodophyllus junceus* (Fr.) Quél. *Agaricus junceus* Fr., however, is a sphagnicolous species of subgenus *Leptonia*, close to *A. (Leptonia) asprellus*, characterized by a subumbilicate pileus with a villous to subsquamulose centre. *Nolanea juncea* sensu Ricken (1913: 300, No. 896) is probably the same taxon as J. Lange, l.c., had in mind. Ricken's description is somewhat hybrid; obviously he combined his own observations with those of Fries: 'Hut kegelig-ausgebreitet, warzig-zugespitzt aber auch mit genabeltem und fast schuppigem Scheitel.' In a note Ricken admits: 'Ich habe ihn nur streng kegelig, mit fast abgesetzter Warze beobachtet', and this suggests that Ricken's own observations refer to the species described here as *E. juncinum*.

Another description of Ricken (1913: 299, No. 892, pl. 74 fig. 4), viz. that of *Nolanea proletaria* Fr. likewise seems to refer to *E. juncinum*. Again it is not quite clear whether Ricken's interpretation is the same as Fries's, or not. Fries's descriptions (1836: 8; 1857: 292; 1874: 206) give a picture of a species with a grey campanulate pileus, with a blunt, villous, umber brown centre and with a smooth, fuliginous stipe. Fries refers to a drawing of Battarra (1755: pl. 18D) which may figure as type plate for his species. To me the Friesian concept remains obscure and does not agree with my view of *E. juncinum*, nor with any other species known by me.

Considering all this I think Kühner & Romagnesi (1954: 5) were right in giving *Rhodophyllus junceus* sensu J. Lange a new name, viz. *Rhodophyllus juncinus*.

Earlier synonyms of *E. juncinum*, viz. *Nolanea globispora* Velen. and *N. zonata* Velen. cannot be recombined in *Entoloma* for nomenclatural reasons (see Noordeloos, 1979b: 255).

In the field *E. juncinum* can be distinguished by the dark grey-brown, strongly striate, smooth pileus, the almost glabrous or minutely striate stipe, and often also by the strong mealy smell. *E. nitens* differs in the lustrous-fibrillous pileal surface, the strongly striate stipe, and in the colour of the pileus.

In forests, a habitat which is preferred by *E. juncinum*, such as *Alno-Padion* and *Ulmion*

*carpinifoliae* vegetations, this species can be found growing together with *E. leptopus*. The latter is easily distinguished by its heterodiametrical spores and the presence of cheilocystidia.

In grasslands *E. juncinum* frequently grows together with *E. papillatum*, which differs by having heterodiametrical spores.

*Entoloma juncinum* can be distinguished from small specimens of *E. sericeum* by slightly larger spores, less intensive membranal pigments and additional intracellular pigment.

#### 14. ENTOLOMA NITENS (Velen.) Noordeloos—Figs. 15a–c

*Nolanea nitens* Velen., České Houby: 627. 1921. — *Rhodophyllus nitens* (Velen.) Kühn. & Romagn., Fl. anal.: 190. 1953. — *Entoloma nitens* (Velen.) Noordeloos in *Persoonia* 10: 252. 1979.

SELECTED ICONES AND DESCRIPTIONS.—Romagnesi in (Trav. mycol. d'éd. R. Kühner) Bull. mens. Soc. linn. Lyon 43 (Num. spéc.): 372–374. 1974. — Arnolds & Noordeloos in *Fung. rar. Icon. col.* 12: pl. 93 fig. d. 1980.

CHARACTERISTICS.—Pileus and stipe grey-brown with silvery fibrillous lustre, particularly when dry; spores isodiametrical.


Pileus 11–42 mm broad, at first conical, conico-campanulate or hemispherical, then expanding to convex or plano-convex, usually with small indistinct papilla, sometimes truncate, sometimes with small abrupt papilla, with straight margin, often reflexed and splitting with age, at first regularly shaped, later with undulating limb, hygrophanous, when moist grey-brown (10 YR 4/3, 4/4, 5/4, 5/3), slightly paler towards margin, translucently striate often up to one half of the radius or more, pallescent on drying to pale greyish brown or pale beige, sometimes slightly tinged ochre at centre (10 YR 6/2, 7/2, 6/3, 7/3); surface when moist with some minute aeriferous fibrils, when dry rather strongly radially satiny-lustrous, at centre sometimes becoming velvety-fluffy. Lamellae L=20–30, l=1–3(–5), moderately distant, narrowly to broadly adnate, sometimes deeply emarginate to almost free, segmentiform then ventricose, sometimes transversely veined, grey-brown, only in later stages tinged pink (10 YR 6/3, 6/4; 7.5 YR 6/4, 5/4), with concolorous entire edge. Stipe 40–70 × 2–4 mm, cylindrical or flattened with longitudinal groove, with or without broadened base, concolorous with moist pileus or paler, strongly silvery-aeriferous striate lengthwise, often twisted, minutely pruinose at apex, white tomentose at base, tomentum often covering basal third of stipe. Flesh in cortex of pileus and stipe concolorous with surface, pale and fibrous inside stipe, brittle in pileus, brittle-fibrous, easily splitting lengthwise in stipe. Smell weak, somewhat earthlike or subraphanoid (raphanoid-subspERMATICAL according to Romagnesi, l.c.). Taste weak, more or less raphanoid, never farinaceous.

Spores 8.0–10.4 × (7.3–)7.5–9.3(–10.4)  $\mu\text{m}$ , Q=1.0–1.1–1.2, L–D=0–0.6–1.5  $\mu\text{m}$ , 5–6-angled in side-view, with distinct basal diedron. Basidia 32–52(–60) × 9–14  $\mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular, cells cylindrical to inflated, 200–340 × 12.5–27  $\mu\text{m}$ , with hyaline, colourless walls. Pileipellis a cutis made up of 3.4–9  $\mu\text{m}$  wide cylindrical hyphae with pale brown minutely encrusted walls. Subpellis sometimes more or less well differentiated, consisting of broad cylindrical cells 38–87 × 10–19  $\mu\text{m}$ , with pale or moderately dark brown intracellular pigment. Pileitrama regular with long cylindrical or inflated cells up to 450  $\mu\text{m}$  long and 10–24  $\mu\text{m}$  wide, with pale, sometimes minutely encrusted walls, intermixed with narrow 3–7.5  $\mu\text{m}$  wide cylindrical hyphae with minutely encrusted walls. Clamp-connections present in hymenium, rare or absent elsewhere.

HABITAT & DISTRIBUTION.—In frondose forests on rich loamy soils or in grassy vegetations, once found in juniper-heath (*Barkman 10071*). Rare but widespread, known from the Netherlands, France, and Czechoslovakia. April–November.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Overijssel, Ommen, estate 'Stekkenkamp', 13 Sept. 1979, *J. Schreurs* & *T. Boekhout*; Ommen, Junne, Junner Koeland, 5 Nov. 1976, *J. J. Barkman 10071*





Figs. 14a-d. *Entoloma juncinum*. — Habit and spores (14a, 14c from *Kits van Waveren*, 18/19 Oct. 1969; 14b, 14d from *Noordeloos* 203).

Figs. 15a-c. *Entoloma nitens*. — Habit and spores (15a from *de Bernon*, April 1977; 15b-c from *Noordeloos* 316).

(WBS); Delden, estate 'Twickel', 20 Aug. 1977, *M. E. Noordeloos 401*; idem, near Oldenzaal, Roderveld, 12 Oct. 1976, *E. Kits van Waveren*; prov. Noord-Brabant, Strijbeek, Goudbergven, 15 Aug. 1964, *P. B. Jansen*.

FRANCE, dept. Manche, Tallepied, 23 Aug. 1971, *H. Romagnesi 71.179* (Herb. Romagn., PC); idem, April 1977, *M. de Bernon*.

*Entoloma nitens* is close to *E. juncinum* from which it differs mainly by the lustrous aspect of the covering layers and the slightly paler, sometimes more ochraceous colour. Romagnesi (1974: 374) also emphasizes the difference in smell. In *E. nitens* the raphanoid smell can be strong, as I personally recorded (coll. *M. de Bernon*, April 1977), but is found to be weak, however, in some collections of both species and is therefore considered less indicative than the striking macroscopical differences.

In the field *E. nitens* can be confused with *E. lucidum* by the lustrous-fibrillose surface of the pileus, but the spores are distinctly isodiametrical.

*Nolanea sericeonitens* P. D. Orton, according to its author closely related to, if not identical with, *Rhodophyllus nitens* (Velen.) Kühn. & Romagn., has differently shaped spores and is considered here to be related to *E. lucidum* (see p. 458).

**Entoloma subsection Minuta (Romagn.) Noordeloos,  
comb. & stat. nov.**

*Rhodophyllus* Quél. section *Minuti* Romagn. in Bull. mens. Soc. linn. Lyon **43**: 331. 1974 (basonym). (In Bull. Soc. mycol. Fr. **53**: 330. 1937, nom. nud.) Typus: *Entoloma minutum* (P. Karst.) Noordeloos.

Carpophores small, pale grey-brown or flesh-coloured beige, pigments membranous and at least on some hyphae of pileipellis and pileitrama minutely encrusting; spores subisodiametrical to heterodiametrical with distinctly dihedral base.

KEY TO THE SPECIES OF SUBSECTION MINUTA

- 1a. Pileus usually distinctly depressed at centre; lamellae pale pink with or without slight grey shade; spores (7.9–)8.0–10.4(–11.5) × (6.4–)6.8–8.1(–8.7) μm; surface of stipe without differentiated hairs.  
*E. minutum* var. *minutum*, p. 468
- b. Pileus usually convex with or without small papilla, never depressed; lamellae distinctly grey-brown with pink tinge; spores 9.3–12(–14) × 7.2–9.3 μm; apex of stipe with well-differentiated, subcapitate hairs.  
*E. tenellum*, p. 470

15. ENTOLOMA MINUTUM (P. Karst.) Noordeloos  
var. MINUTUM—Figs. 16a–c

*Nolanea minuta* P. Karst. in Meddn Soc. Fauna Fl. fenn. **5**: 10. 1879 (= Symb. ad Mycol. fenn. 6). — *Rhodophyllus minutus* (P. Karst.) J. Lange in Dansk bot. Ark. **2**: 37. 1921. — *Entoloma minutum* (P. Karst.) Noordeloos in Persoonia **10**: 248. 1979 (non *Ecclilia minuta* Velen. 1921 nec *Claudopus minutus* Velen. 1929).

SELECTED ICONS AND DESCRIPTIONS.—Arnolds & Noordeloos in Fung. rar. Icon. col. **12**: pl. 93, figs. a, b. 1980. — Jossierand in Bull. Soc. mycol. Fr. **53**: 213–215. 1937. — J. Lange, Fl. agar. dan. **2**: 102, pl. 79H. 1936. — Nathorst-Windahl in Friesia **8**: 13. 1966.

**CHARACTERISTICS.**—Carpophores rather small and thin-fleshed, brittle; pileus small and flattened, usually with central depression; pale flesh-coloured to pale grey-brown, remarkably translucently striate; stipe concolorous with pileus.

Pileus 6–25 mm broad, conical, soon expanding to (plano-)convex usually with slightly depressed, rarely minutely papillate centre, with straight margin, strongly hygrophanous, when moist pale flesh-colour beige to pale grey-brown with remarkably contrasting grey-brown striation and centre (centre and striae 10 YR 6/3, 6/4, 5/3; centre sometimes 7.5 YR 5/4; interstriation and margin 10 YR 7/6, 7/4, 8/4, 8/2), strongly pallescent on drying to pale grey-brown or sordid white (10 YR 8/3, 8/4), innately to aeriferously fibrillous-lustrous, shining, smooth or at centre, particularly in young specimens, slightly rugulose to fluffy-subsquamous. Lamellae  $L=13-22$ ,  $l=1-3$ , moderately distant, adnate to almost free, ventricose, often exceeding the pileus, white to pale grey, then pink with or without grey tinge (10 YR 8/4, 7/4 then 7.5 YR 7/2, 7/4), with concolorous, entire or slightly eroded edge. Stipe 25–65 × 1–1.5(–2) mm, cylindrical, straight, somewhat glassy hyaline pale grey-brown or ochraceous-orange to greyish yellow, sometimes innately fibrillous to very slightly silvery striate lengthwise (lens), glabrous apex sometimes minutely pruinose, solid then narrowly fistulose. Flesh very thin-membranaceous, brittle, glassy, pale grey-brown or flesh-colour, concolorous with surface. Smell weak, never farinaceous. Taste none or slightly bitterish to subfarinaceous.

Spores (7.9–)8.0–10.4(–11.5) × (6.4–)6.8–8.1(–8.7)  $\mu\text{m}$ ,  $Q=1.1-1.25-1.4(-1.5)$ ,  $L-D=(1.2-)$  1.7–2.5(–4)  $\mu\text{m}$ , irregularly 5–6(–7)-angled in side-view, with distinctly dihedral base. Basidia (26–)27–40 × 9–12.5  $\mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular, composed of long inflated-subcylindrical cells (198–)208–340(–511) × (10.2–) 11.4–30.5(–37.5)  $\mu\text{m}$ , with pale, hyaline walls. Pileipellis a cutis made up of radially arranged repent cylindrical hyphae (2.7–)4–13.2(–15)  $\mu\text{m}$  wide, gradually passing into trama, scattered, with—particularly at centre—repent to ascending (clavate) terminal cells up to 17  $\mu\text{m}$  wide. Pileitrarma regular; hyphae strongly inflated, up to 34  $\mu\text{m}$  wide; cells up to 340(–429)  $\mu\text{m}$  long, with pale, sometimes minutely encrusted walls and rarely also diffusely or granularly intracellular pigment. Clamp-connections frequent in hymenium, rare in other tissues.

**HABITAT & DISTRIBUTION.**—On damp soil, usually in forests of *Alnus*, *Betula*, *Salix*, more rarely in moist grasslands, widespread. Recorded from the Netherlands, Scandinavia, Great Britain, German Federal Republic, France, and Czechoslovakia.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Drenthe, Beilen, Holtherzand, 26 Sept. 1979, *E. Arnolds* (WBS); Vries, Taarlo, along Taarlose Diep, 4 July 1974, *E. Arnolds* 3126 (WBS); Emmen, 12 Sept. 1974, *E. Kits van Waveren*; prov. Gelderland, Delden, 9 Oct. 1974, *E. Kits van Waveren*; Winterswijk, Korenburgerveen, 29 Sept. 1973, *C. Bas* 6158 & 6159; prov. Noord-Brabant, Bergen op Zoom, Zoomland, 5 July 1974, *P. B. Jansen* 74–274; Nieuw Ginneken, Goudbergven, 13 Sept. 1964, *P. B. Jansen* 64–202; Zundert, Lange Goren, 6 July 1963, *P. B. Jansen* 63–175.

FINLAND, Inari Lappland, Kevo, Kevoosu, 15 Aug. 1978, *M. E. Noordeloos* 683. — GERMAN FEDERAL REPUBLIC, Bayern, Rosenau near Dingolfing, 14 Aug. 1967, *A. Einhellinger* (M).

The pigmentation of *Entoloma minutum* is difficult to study because of the pale colours of the pileus. Careful observations, however, reveal that the membranes of the hyphae in pileipellis and -trama are uniformly coloured (membranal pigmentation) and minutely encrusted at least in part of the hyphae, often the narrowest ones. Two collections studied (*Bas* 6159 and *Arnolds*, 26 Sept. 1979) also showed an intracellular pigmentation, diffuse or in the form of granules, in some hyphae of the surface of the pileus.

Josserand (1937: 213–215, fig. 13) gave an excellent description of *E. minutum* but found the pigmentation to be 'vacuolaire', and this statement was repeated by Kühner & Romagnesi (1953: 183). Later Romagnesi changed his mind on this character (1954: 199 and 1974b: 331) and described it as 'membranaire parfois un peu incrustante.'

Einhellinger (1969: 106) described a collection from Bavaria with intracellular pigment. A study of this material revealed at least some encrusted hyphae in the upper layer of the pileus.<sup>1</sup>

The spores are found to be rather variable, e.g. within one collection from more or less isodiametrical to distinctly heterodiametrical. The first type reminds of the spores of *E. juncinum* and *E. nitens*, which species are easily distinguished by their different habit and dark pigmentation. The latter heterodiametrical type of spores resemble those of *E. tenellum* (see below).

*Entoloma minutum* var. *polymorphum* (Romagn.) Noordeloos (1979b: 248) differs from the type among others by the rather variable shape of the pileus and insertion of the lamellae. On account of the presence of the same characters I have placed *Nolanea depressa* Velen. in the synonymy of this variety. At present this variety is known only from France and Czechoslovakia. For a detailed description I refer to Romagnesi (1954: 197–199).

#### 16. ENTOLOMA TENELLUM (Favre) Noordeloos—Figs. 18a–d

*Rhodophyllus tenellus* Favre, Assoc. fong. hauts-marais: 212. 1948. — *Nolanea tenella* (Favre) P. D. Orton in Trans. Br. mycol. Soc. 43: 179. 1960. — *Entoloma tenellum* (Favre) Noordeloos in Persoonia 10: 256. 1979. *Leptonia cinerascens* Velen., České Houby: 623. 1921 (non *Entoloma cinerascens* Hesler, 1967).

CHARACTERISTICS.—Carpophores slender; pileus 5–13 mm broad, pale grey-brown; stipe concolorous, polished, apex pruinose with (sub-)capitate hairs; smell indistinct.

Pileus 5–13 mm broad, conico-convex, often with small papilla, with margin slightly involute, exceeding the lamellae, hygrophanous, when moist brown-grey to beige-grey (e.g. 10 YR 5/3), translucently striate up to two-thirds of the radius, strongly pallescent on drying to sordid white (10 YR 7/2), brilliantly shining, smooth, glabrous. Lamellae moderately distant, narrowly adnate to almost free, broadly ventricose, 2.5–3 mm broad, greyish brown with pink tinge (10 YR 5/2, 5/3), with entire, concolorous edge. Stipe 30 × 1.5–2 mm, straight, cylindrical, hyaline, ochraceous-yellow to grey-brown, minutely pruinose at apex, downwards smooth and shining. Flesh rather thin, firm-subcartilagineous, concolorous with surface. Smell and taste indistinct.

Spores (8.1–)9.3–12(–14) × 7.2–9.3 μm, Q = 1.1–1.25–1.5, L–D = 0.6–4(–5) μm, (5–)6–7(–8)-angled in side view, with dihedral base. Basidia 32–48 × 9.4–12 μm, 4-(rarely 2-)spored. Cystidia absent.<sup>2</sup> Hymenophoral trama regular, composed of cylindrical to inflated cells 180–270 × 12–22 μm. Pileipellis a cutis made up of cylindrical 2.7–14 μm wide hyphae with minutely encrusted walls. Stipitepellis a cutis with at apex of stipe cylindric-capitate hairs 50–150 × 6.5–10 × 3.5–9 μm. Clamp-connections frequent in hymenium, rare elsewhere.

HABITAT & DISTRIBUTION.—In marshy vegetation with *Filipendula ulmaria* (Favre, l.c.). In the Netherlands found in a mossy vegetation on acid, sandy soil with *Juniperus communis* and in *Quercus* forest.

<sup>1</sup> I also failed to find any cheilo- and/or pleurocystidia in this collection from Bayern, Rosenau near Dingolfing, 14 Aug. 1967, A. Einhellinger (M).

<sup>2</sup> Cheilocystidia were said to be present but rare in the holotype, 36–58 × 7.5–11.5 μm, cylindrical-flexuose ('vermiciform'), scattered among basidia, rare. See discussion below.


COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Sleen, Slenerzand, 27 Oct. 1965, A. K. Masselink (WBS); prov. Limburg, Mook, 12 Oct. 1965, E. Kits van Waveren.

SWITZERLAND, Grisons, Val Sesvenna, 14 Aug. 1943, J. Favre (lectotype, design. mihi, CHUR). — CZECHOSLOVAKIA, Bohemia, Bozkov, July 1919, J. Velenovský (holotype of *L. cinerascens*, PRC).

STUDY OF LECTOTYPE.—The lectotype consists of three specimens with the following microscopical characters.

Spores  $9.3\text{--}11.5\text{--}(12) \times (7.6\text{--})8.1\text{--}9.3 \mu\text{m}$ ,  $Q = 1.13\text{--}1.23\text{--}1.43$ ,  $L\text{--}D = 1.2\text{--}2\text{--}3.4 \mu\text{m}$ , 6–8-angled in side-view, with distinctly dihedral base. Basidia  $31\text{--}53 \times 8\text{--}12 \mu\text{m}$ , 4-(rarely 2-)spored. Cheilocystidia not found. Hymenophoral trama regular, composed of long cylindrical to slightly inflated cells e.g.  $230 \times 17 \mu\text{m}$ . Pileipellis a thin cutis made up of 2.6–8  $\mu\text{m}$  wide, cylindrical hyphae with minutely encrusted walls. Stipitopellis with some cylindrical-capitate hairs,  $35\text{--}60 \times 4\text{--}7.5 \times 5\text{--}10 \mu\text{m}$ . Clamp-connections seen at base of basidia.

As no holotype is left at CHUR nor at G, I select the authentic collection of Val Sesvenna, 14 Aug. 1943, collected and named by J. Favre, as lectotype. In the lectotype the spore dimensions are slightly different from those given by Favre in the protologue, viz.  $11\text{--}12.5 \times 6\text{--}7.5 \mu\text{m}$ , thus



Figs. 16a–c. *Entoloma minutum*. — Habit and spores (16a–b from Noordeloos 683; 16c from Jansen 74–274).

Figs. 17a–b. *Entoloma tibilocystidiatum*. — Spores and cheilocystidia (17a–b from Arnolds 456).

Figs. 18a–d. *Entoloma tenellum*. — Habit, spores and hairs of stipe (18a–b, 18d from Kits van Waveren, 12 Oct. 1965; 18c from lectotype).

distinctly narrower. In the discussion following the description Favre (1948: 57) mentioned this collection. He said that he was unable to locate any cystidia in this collection. (I could not find any cystidia either, nor in the lectotype nor in the other collections studied.) The difference in spore size apparently did not impress him, as he did not mention it at all. He explicitly considered this collection to belong to *Nolanea tenella*.

*Entoloma cuspidifer* resembles *E. tenellum* in spore shape and covering of the stipe. The spores, however, are somewhat larger and born on bisporous basidia. Besides, the habit and colour of the carpophore are rather different.

*Entoloma vernum* has the same type of hairs on the stipe as *E. tenellum*, but differs among other things in the dark grey-brown colours and strictly vernal appearance.

**Entoloma subsection Cosmeoxonema (Largent & Thiers) Noordeloos**  
*comb. & stat. nov.*

*Nolanea* (Fr.) Kummer sect. *Cosmeoxonema* Largent & Thiers in Northwest Sci. 46: 35. 1972. —  
*Rhodophyllus* Quél. sect. *Cosmeoxonema* (Largent & Thiers) Romagn. in Bull. mens. Soc. linn. Lyon 43: 331.  
1974.

Pigment exclusively encrusting, often coarsely so, the hyphae of pileipellis and pileitrama; spores isodiametrical or heterodiametrical and then with distinctly dihedral base. Type: *Entoloma sericeum* (Bull. ex Mérat) Quél.

KEY TO THE SPECIES OF SUBSECTION COSMEOXONEMA

- |  |  |
|--|--|
| 1a. Cheilocystidia present. . . . .  | 2  |
| b. Cheilocystidia absent. . . . .  | 3  |
| 2a. Cheilocystidia tibiiform; spores isodiametrical; smell farinaceous. . . . .  | <i>E. tibilocystidium</i> , p. 473                                     |
| b. Cheilocystidia cylindrical-subcapitate; spores heterodiametrical; smell sweet, like in <i>Hebeloma sacchariolens</i> . . . . .  | <i>E. sacchariolens</i> , p. 474                                       |
| c. Cheilocystidia sphaeropedunculate, subglobose or broadly clavate; spores heterodiametrical; smell absent. . . . .   | <i>E. sphaerocystis</i> , p. 485                                       |
| 3a. Spores isodiametrical, $Q = 1.1$ on the average per collection. . . . .  | 4  |
| b. Spores heterodiametrical, $Q \geq 1.25$ on the average per collection. . . . .  | 7  |
| 4a. Pileus when moist very obscurely striate at margin only, very dark grey-brown, with scattered silvery fibrillous patches producing a hoary impression; lamellae predominantly with grey tinge (like in <i>Tephroclype</i> spp.). . . . . | <i>E. sericeum</i> var. <i>cinereo-opacum</i> , p. 482                 |
| b. Pileus distinctly striate when moist, sepia or reddish brown, smooth; lamellae reddish brown. . . . .   | 5  |
| 5a. Pileus relatively thick-fleshed, with faint umbo or flattened or slightly depressed; stipe usually as long as diameter of pileus, or shorter. . . . .  | <i>E. sericeum</i> var. <i>sericeum</i> f. <i>sericeum</i> , p. 479    |
| b. Pileus thin-fleshed, stipe often distinctly longer than diameter of pileus. . . . .   | 6  |
| 6a. Pileus distinctly papillate. . . . .   | <i>E. sericeum</i> var. <i>sericeum</i> f. <i>nolaniforme</i> , p. 480 |
| b. Pileus distinctly umbilicate. . . . .   | <i>E. sericeoides</i> , p. 483   |
| 7a. Pileus fairly dark sepia brown ('bistre'); hyphae of pileipellis coarsely encrusted; spores bluntly angled; smell none; strictly vernal in or near coniferous forests. . . . .   | <i>E. vernum</i> , p. 476  |
| b. Pileus pale to moderately dark grey-brown, hyphae of pileipellis minutely encrusted; spores pronouncedly angled; smell strong, sweet, like in <i>Hebeloma sacchariolens</i> ; summer-autumn in grasslands. . . . .                        | <i>E. amecides</i> , p. 473  |

17. *ENTOLOMA TIBIICYSTIDIATUM* Arnolds & Noordeloos—Figs. 17a–b

*Entoloma tibiicystidium* Arnolds & Noordeloos in *Persoonia* 10: 294–295, figs. 23–26. 1979.

Pileus 10–15 mm broad, irregularly plano-convex without papilla, with straight margin, hygrophanous, when moist rather pale grey-brown, translucently striate, pallescent on drying. Lamellae fairly distant, almost free, pale salmon-pink. Stipe up to 38 × 2 mm, slender, cylindrical, pale grey-brown, silvery striate. Smell farinaceous. Spores (7.0–)7.9–10.5(–11) × (6.4–)6.8–9.3 μm, Q = 1.0–1.1–1.2(–1.3), isodiametrical. Basidia 4-(rarely 2-)spored. Cheilocystidia subcylindrical-subcapitate to tibiiform, (17–)20–35 × 6–8 × 2.5–5 μm, often with hyaline mucous cap covering the apex. Pileipellis a poorly differentiated cutis made up of 2.5–8 μm wide cylindrical hyphae gradually passing into trama, with membranal, often encrusting pigment. Clamp-connections present.

HABITAT.—In poorly fertilized hayfield (*Calthion palustris*) on wet, in winter inundated, peaty soil.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, de Wijk, 'Reest valley', 15 Oct. 1979, *E. Arnolds* 3663 (holotype, WBS; isotype, L); prov. Noord-Brabant, Udenhout, loam pits, 24 Oct. 1969, *E. Arnolds* 456.

The type of pigmentation and the spores of *E. tibiicystidium* suggest close relationship with *E. sericeum*, in particular with var. *sericeum* f. *nolaniforme*. It differs, however, so strikingly by the remarkably shaped cheilocystidia, that I consider it a distinct species.

18. *ENTOLOMA AMEIDES* (Berk. & Br.) Sacc.—Figs. 19a–b

*Agaricus ameides* Berk. & Br. in *Ann. Mag. nat. Hist.*, Ser. III, 15: 315. 1865 (Notices on Br. Fungi no. 999). — *Rhodophyllus ameides* (Berk. & Br.) Quél., *Enchir.*: 58. 1886. — *Entoloma ameides* (Berk. & Br.) Sacc., *Syll. Fung.* 5: 686. 1887.

SELECTED ICONES.—Cooke, III. *Br. Fungi* pl. 329 (341). 1886. — Konr. & Maubl., *Icon. sel. Fung.* pl. 192. 1925.

SELECTED DESCRIPTIONS.—Einhelliger in *Ber. bayer. bot. Ges.* 41: 102–103 figs. 17, 21; pl. 9a, 11c. 1969. — Konr. in *Bull. Soc. mycol. Fr.* 41: 45–46. 1925; *ibid.* 45: 52. 1929. — F. H. Möller, *Fungi Faerøes* 1: 248. 1945.

CHARACTERISTICS.—Habit rather variable, entolomatoid to nolaneoid, rarely with umbilicate pileus and subdecurrent lamellae; pileus pale grey to grey-brown, at centre sometimes with ochre tinge; stipe pale grey-brown, silvery striate; smell strong and persistent, like that of *Hebeloma sacchariolens*.

Pileus 12–35 mm broad, 7–30 mm high, conical to conical-campanulate, only slightly expanding to conico-convex with pointed papilla, rarely plano-convex with pointed papilla, with margin involute when young, then straight, strongly hygrophanous, when moist pale grey-ochre to grey-brown (10 YR 6/2, 6/3, 5/2, 5/3, sometimes more like 10 YR 5/4), translucently striate up to one half of the radius, on drying pallescent to greyish, very pale brown, ochraceous, remaining darker at centre (centre 10 YR 5/4, margin 10 YR 7/4, 8/3), glabrous, lustrous. Lamellae L = 22–35, l = (1–)3, narrowly adnate-emarginate, almost free, ventricose, thickish, grey then pink, finally dark grey-brown with pink tinge (10 YR 7/2, 7/3, then between 10 YR 7/4 and 7.5 YR 7/4, finally more like 10 YR 6/3, 5/2 with pink tinge), slightly hygrophanous, pallescent on drying from edge towards back, with subserrulate, concolorous edge. Stipe 30–80 × 3–7 mm, cylindrical, often flexuous-twisted, solid, then fistulose, pale isabella-ochraceous as dry pileus (10 YR 8/2,

8/3 to 7/3, sometimes 8/4), brilliantly and innately fibrillosly striate lengthwise, slightly pruinose at apex, smooth downwards, at base more or less white tomentose. Flesh concolorous with surfaces, relatively thin in pileus and brittle-fibrillose, easily splitting lengthwise in stipe. Smell rather strong like acetylaldehyde, *Hebeloma sacchariolens*. Taste sweet, like caramel.

Spores  $9.3-11.5(-12) \times 7.6-9.3 \mu\text{m}$ ,  $Q = 1.1-1.25-1.4$ ,  $L-D = 0.6-2-3.5 \mu\text{m}$ , 5-6-angled in side-view and with rather pronounced suprahilar depression, most probably with bluntly dihedral base. Basidia  $35-42 \times 10.5-16.3 \mu\text{m}$ , 4-spored, slenderly clavate. Cystidia absent. Hymenophoral trama regular, composed of cylindrical to slightly inflated cells up to  $270 \mu\text{m}$  long and  $8.5-26 \mu\text{m}$  wide. Pileipellis a thin cutis made up of  $2.7-9 \mu\text{m}$  wide cylindrical hyphae with minutely encrusted walls. Pileitrama regular, composed of cylindrical or inflated hyphae with particularly in upper layers pale coloured, sometimes encrusted walls; cells  $190-295 \times 12-28 \mu\text{m}$ . Clamp-connections abundant in hymenium, also rarely seen in pileipellis.

**HABITAT & DISTRIBUTION.**—In hayfields and extensively grazed meadows, widespread but apparently rare. Known from the Netherlands, Scandinavia, Great Britain, Germany, and France.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Limburg, Wijlre, Dolsberg, 20 Sept. 1978, *M. E. Noordeloos* 724.

GREAT BRITAIN, Flintshire, Bodelwyddan, Sept. 1863, *Berkeley & Broome* (holotype, K). — FRANCE, Jura, Curtil, 14 Aug. 1941, *Métrod* 1222 (PC).

The description given above is based on the Netherlands' collection which represents a slender, nolaneoid form of *Entoloma ameides*. The more typical form as depicted by Cooke (l.c.) has a more entolomatoid habit. Konrad (1925, 1929; l.c.) clearly illustrated the variability of this species. The plate of Konrad & Maublanc (1925, l.c.) gives an excellent impression of the colours of the Netherlands' collection; the plate of Cooke, however, is too much yellow. I was able to compare Cooke's plate with the original at Kew, which seems to be more 'natural' in colour. Comparing other plates in the same work, I came to the conclusion that, generally speaking, the *Entoloma* plates are printed in a somewhat too yellow tone.

The general appearance of *E. ameides* may be that of a member of subgenus *Entoloma*, but the type of pigmentation and structure of trama definitely place it in subgenus *Nolanea*, close to *E. sericeum* and *E. papillatum*.

*Entoloma sacchariolens* may be considered a slender form of *E. ameides*, but the slight difference in colour and the presence of cheilocystidia suggest that the former is a species in its own right. The spores, however, are remarkably similar. In this case more material is needed for a final decision about the value of cheilocystidia as a distinctive character.

19. *Entoloma sacchariolens* (Romagn.) Noordeloos,  
*comb. nov.*—Figs. 20a-c

*Rhodophyllus sacchariolens* Romagn. in Bull. mens. Soc. linn. Lyon 43, No. spéc. (Trav. déd. R. Kühner): 385. 1974 (basionym).

**CHARACTERISTICS.**—Pileus grey-brown with ochraceous tinge, rather strongly acriferosly lustrous when dry; stipe silvery striate lengthwise; smell strong, sweet, like in *Hebeloma sacchariolens*; cheilocystidia present, cylindrical-subcapitate.

Pileus 20-24 mm broad, conico-convex, then expanding to plano-convex with blunt or subumbonate centre and marginal zone irregularly undulating with age, with outermost margin






Fig. 19a–b. *Entoloma ameides*. — Habit and spores (19a–b from Noordeloos 724).

Figs. 20a–c. *Entoloma sacchariolens*. — Habit, spores and cheilocystidia (20a–c from Bas 7623).

involute, weakly hygrophanous, when moist brown with grey or sometimes ochraceous tinge (10 YR 4/2, 4/3, rarely 3/2), paler towards margin (10 YR 7/4, 7/6), translucently striate at margin only, pallescent on drying to greyish ochraceous (10 YR 8/3, 8/6), with surface already when moist with some aeriferous fibrils, becoming strongly lustrous-shining on drying; at centre often somewhat fluffy. Lamellae L = 21–30, l = 3–7, moderately distant, deeply emarginate to almost free, or adnate, segmentiform then narrowly ventricose, brown (10 YR 6/3), later with pink shade, slightly paler towards serrulate edge. Stipe 40–75 × 2–5 mm, cylindrical, often bent at base, concolorous with pileus, pruinose at apex, downwards glabrous, strongly silvery striate lengthwise, white tomentose at base, solid or narrowly fistulose. Flesh in cortex concolorous with surface, inner part paler, firm in pileus, more fibrous in stipe. Smell strong, sweet, like that of orange blossom or *Hebeloma sacchariolens*, sometimes weak, but then becoming more distinct after being kept for some time in a closed box. Taste unpleasant, soap-like.

Spores 9.3–10.5(–11.0) × 7.0–8.1 μm, Q = 1.15–1.3–1.4, L–D = 1.2–3.0 μm, 5–6-angled in side-view, with distinctly dihedral base. Basidia 28.5–35.0 × 10.5–12 μm, 4-(rarely 2-)spored. Cheilocystidia 25–45 × 3.5–8(–12.0) μm, cylindrical to clavate, sometimes subcapitate, scattered among basidia. Hymenophoral trama regular, with cylindrical to slightly inflated cells 140–325 × 5.7–18.0(–23) μm. Pileipellis a simple cutis made up of 1.5–3(–6) μm wide, cylindrical hyphae

with pale brown, sometimes minutely encrusted walls. Pileitrama regular, with cylindrical to inflated hyphae with cells up to 350  $\mu\text{m}$  long and 7–28  $\mu\text{m}$  wide, with pale, sometimes minutely encrusted walls. Clamp-connections frequent in hymenium, but very rare in trama.

**HABITAT & DISTRIBUTION.**—In *Salix repens* dominated dune meadow in old dune valley on slightly acid sand. Very rare; only known from one locality in the Netherlands and from the type-locality in France.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Noord-Holland, Callantsoog, nature-reserve 'Zwanewater', 5 Nov. 1977, C. Bas 7623.

FRANCE, dept. Oise, Luzarches, 18 July 1951, H. Romagnesi 51.71 (holotype, Herb. Romagn., PC).

*Entoloma sacchariolens* is very closely related to *E. ameides*, but differs from the latter by the more ochraceous tinged colour of the pileus, the remarkably lustrous pileal surface, and by the cylindrico-subcapitate cheilocystidia. Normally the habit of *E. ameides* is more entolomatoid (=tricholomatoid) than that of *E. sacchariolens*, but slender specimens of *E. ameides* do also occur (see above). In addition to the characteristic smell, *E. sacchariolens* differs from other cystidiate members of subgenus *Nolanea* in the type of pigmentation (sect. *Nolanea* has two types of pigment, subsect *Cheilocystidiata* has intracellular pigment) and/or in size and shape of spores (compare *E. cryptocystidiatum* and *E. tibiicystidiatum*).

## 20. ENTOLOMA VERNUM Lundell—Figs. 21a–h

*Entoloma vernum* Lund. in Svensk bot. Tidskr. 31: 193. 1937. — *Rhodophyllus vernus* (Lund.) Romagn. in Bull. Soc. mycol. Fr. 63: 195. 1947. — *Nolanea verna* (Lund.) Kotl. & Pouz. in Česká Mykol. 26: 221. 1972. *Rhodophyllus cucullatus* Favre, Champ. supér. Zone alp.: 62. 1955. — *Nolanea cucullata* (Favre) P. D. Orton in Trans. Br. mycol. Soc. 43: 179. 1960.

**MISAPPLICATIONS.**—*Nolanea pascua* sensu Bres., Iconogr. mycol. pl. 580. 1929. — *Nolanea erophila* sensu Velen., České Houby: 630. 1921.


**SELECTED ICONES AND DESCRIPTIONS.**—Lund., l.c.: 191 fig. 3. 1937. — Herink in Česká Mykol. 10: 1–9, pl. 21. 1956. — Ayer in Schweiz. Z. Pilzk. 52: 17–19, pl. 1. 1974. — Mazzer in Mich. Bot. 16: 195–200, figs. 1–6. 1977. — Cetto, Funghi Vero 2: 247, pl. 532. 1976.

**CHARACTERISTICS.**—Strictly vernal; pileus rather dark sepia brown, reminding of *E. sericeum*; lamellae with brown tinge; smell absent.

Pileus 11–60 mm broad, acutely conical to conico-campanulate, only slowly expanding to convex, finally plano-convex, then usually with conical papilla, with margin slightly involute when young, later straight, exceeding lamellae and becoming lobed and undulate with age, strongly hygrophanous, when moist rather dark sepia, at centre even blackish brown, paler at margin (2.5 Y 3/2, 4/3, rarely 6/4; 10 YR 3/3, 3/2; Expo H62; Ség. 131, at margin 10 YR 5/4, 6/4; 2.5 Y 6/6; Ség. 134; Expo D62), translucently striate in fleshy specimens at least at margin, but in small membranaceous pilei frequently up to centre, on drying pallescent to golden or grey-brown, ochraceous grey or isabella (2.5 Y 6/4 to 6/6; 10 YR 6/4, 5/4; Expo B 61), shining, glabrous. Lamellae L = 19–40, l = 1–3(–5), adnexed to almost free, ventricose, up to 7 mm broad, pale grey-brown when young, later rather dark brown-pink or rusty brown (10 YR 6/4; 7.5 YR 7/4, 6/4; Expo E42), with concolorous entire or crenulate edge. Stipe 20–67  $\times$  2–7 (base –10) mm, cylindrical, sometimes distinctly but gradually broadening towards base, sometimes flattened with longitudinal groove, solid to narrowly fistulose, concolorous with pileus or paler, then more or less ochraceous tinged, pruinose at apex (rarely pruinose all over in young specimens), downwards glabrous, innately fibrillose, not distinctly striate, white tomentose at base. Flesh

pale brown in pileus and stipe, in cortex more or less concolorous with surface, brittle. Smell none or faint and herbaceous. Taste mild.

Spores (8.1–)9.0–11.5(–12) × 7.0–8.7(–9.3)  $\mu\text{m}$ , Q = 1.1–1.25–1.4, L–D = 1–2.5  $\mu\text{m}$ . 5–6–7-angled in side-view, with more or less distinctly dihedral base. Basidia 27–50 × 8.5–12.5(–16)  $\mu\text{m}$ , 4-spored, in some collections frequently with hyaline, thickened walls ('sklerobasidia'). Cystidia absent. Hymenophoral trama regular, composed of cylindrical to inflated cells up to 450  $\mu\text{m}$  long and 7.5–28  $\mu\text{m}$  wide, frequently with brown walls. Pileipellis a cutis made up of radially arranged



Figs. 21a–h. *Entoloma vernum*. — Habit, spores and hairs of stipe (21a, 21f, 21h from *Bas 1700*; 21b from *Noordeloos 317*; 21c from *Tjallingii*, 8 April 1978; 21d from type of *Rhodophyllus cucullatus*; 21e, 21g from holotype).

cylindrical hyphae 4–10(–11)  $\mu\text{m}$  wide, with minutely to coarsely encrusted walls. Pileitrama regular, with inflated hyphae, composed of up to 370  $\mu\text{m}$  long and 6.5–29  $\mu\text{m}$  wide cells with brown, encrusted walls, encrustations often coarse, particularly on narrowest hyphae. Apex of stipe with clusters of long, subcylindrical to clavate, frequently subcapitate hairs 23–120  $\times$  4.5–15  $\times$  5–11  $\mu\text{m}$ . Clamp-connections frequent in hymenium, rare elsewhere.

**HABITAT & DISTRIBUTION.**—Terrestrial on grassy spots or in grasslands on sandy soil, preferably near coniferous trees (*Pinus*, *Juniperus*). Widespread and frequent in boreal and mountainous habitats; rare in the lowlands of western Europe. March–May.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Drenthe, Sleen, Sleener Zand, 13 April 1967, *J. J. Barkman* 8372 (WBS); prov. Gelderland, Apeldoorn, 11 April 1966, *A. N. Koopmans*; Vierhouten, 5 April 1968, *J. van Brummelen* 658; prov. Zuid-Holland, Meydel near entrance to 'Kijfhoek', 27 March 1959, *C. Bas* 1700 and 19 April 1975, *C. Bas* 6555; IJsselmeerpolders, Zuidelijk Flevoland, beach area near Nykerkersluis, 8 April and 6 May 1978, *G. J. M. G. Tjallingii-Beukers*.

SWEDEN, prov. Upland, Årentuna parish, 19 May 1933, *S. Lundell* (holotype, U); Fungi exsicc. succ. 402; idem, 22 May 1955, *H. Belin* and *L. Holm* (PC); Bålinge parish, Märsta skog, 4 and 8 May 1952, *H. Smith* (PC, Fungi exsicc. succ. 2010). — GERMAN FEDERAL REPUBLIC, Kreis Meppen, Haselünne, 4 May 1977, *B. de Vries* (WBS). — FRANCE, dept. Oise, Forêt de Fontainebleau, 1 May 1977, *M. E. Noordeloos* 317. — SWITZERLAND, Parc nat. Suisse, Col dal Fuorn, 19 May 1958, *Dr. Zimmerli* (holotype of *R. cucullatus* Favre, CHUR).

As has been observed in other species of subject, *Cosmeoxonema*, *E. vernum* appears to have a rather variable habit. This is reflected in literature. Smotlacha (1945: 67) and Herink (1956, l.c.) distinguish a robust, 'entolomatoid', and a slender, 'nolaneoid' form. The typical form of *E. vernum* as characterized by Lundell (1937, l.c.) is a rather robust fungus with a fleshy pileus and firm stipe. As a consequence the pileus is striate at the margin only. This form has its main distribution in the boreal and mountainous habitats of northern and central Europe. It is not uncommon in or near coniferous forests, for example in Sweden and in Czechoslovakia. According to Mazzer (1977, l.c.) this form is also frequently met with in the Great Lakes area in the northern United States.

In western Europe a slender form of *E. vernum* is observed (Fig. 21c) with a membranaceous, deeply striate pileus. The dimension of the pileus falls within the range of 11–35 mm with 19–25 entire lamellae. The apex of the stipe is frequently abundantly pruinose, which is microscopically reflected in the relative abundance of hairs. As there are no strict limits between the robust 'typical' form and the slender one, a distinction on varietal level does not seem justified.

*Rhodophyllus cucullatus* Favre comes very close to the slender form of *E. vernum* as described above. After microscopical investigation of the holotype, I consider it a synonym.

On account of the abundantly encrusted pileal hyphae and the dark colours of pileus and stipe *E. vernum* is placed in subsection *Cosmeoxonema* close to *E. sericeum*, from which it differs among other things in the heterodiametrical spores. *Entoloma papillatum* has a different spore shape, a farinaceous smell, and occurs in summer and autumn.

## 21. ENTOLOMA SERICEUM (Bull. ex Mérat) Quél.

*Agaricus sericeus* Bull. ex Mérat, Fl. Envir. Paris, 2nd Ed.: 78. 1821, non Schaeff. 1762, nec Pers. 1798. — *Entoloma sericeum* (Bull. ex Mérat) Quél. in Mém. Soc. ém. Montbéliard. Sér. II, 5: 119. 1872. — *Rhodophyllus sericeus* (Bull. ex Mérat) Quél., Enchir.: 59. 1886. — *Nolanea sericea* (Bull. ex Mérat) P. D. Orton in Trans. Br. mycol. Soc. 43: 179. 1960.

## var. SERICEUM f. SERICEUM—Figs. 23a-f

*Rhodophyllus sericeus* var. *typicus* Kühn. & Romagn., Fl. anal.: 193. 1953.

SELECTED ICONES AND DESCRIPTIONS.—Ricken, Blätterpilze, pl. 72 fig. 5. 1913. — J. Lange, Fl. agar. dan. 2: 97, pl. 76G, Gl. 1936. — Einhellinger in Ber. bayer. bot. Ges. 41: 107, fig. 19. 1969.

CHARACTERISTICS.—General habit thick-set: length of stipe about equal to diameter of pileus; pileus soon more or less flattened without a pronounced papilla, moderately to dark brown, often tinged red, deeply striate when moist, strongly pallescent on drying; lamellae pale then brown, finally pinkish brown; stipe concolorous with or slightly paler than pileus, usually covered with aeriferous fibrils; smell strongly farinaceous-rancid; spores isodiametrical; pigment coarsely encrusting at least the narrow hyphae of pileipellis and -trama; clamp-connections present.

Pileus 12.5–70 mm broad, from conico-convex or hemispherical expanding to convex, with centre usually flattened, rarely papillate or slightly depressed, with at first involute but later on straight margin, usually undulating, and sometimes splitting with age, strongly hygrophanous, when moist moderately to rather dark sepia or date brown, sometimes with reddish tinge (10 YR 2/2, 3/2; 7.5 YR 3/2; 5 YR 3/2, 3/3), paler towards margin (10 YR 5/2, 5/4, 6/4; 7.5 YR 4/4, 5/2, 5/4, 6/4), translucently striate up to two-thirds of the radius, on drying strongly pallescent along radial streaks to pale sepia brown or ochraceous brown (10 YR 6/4, 7/3; 2.5 Y 7/2; 5 Y 7/3), smooth or minutely radially wrinkled or grooved, radially fibrillous, at centre sometimes slightly rugulose, shining. Lamellae L = 22–36, l = 3–5–7(–9), crowded to moderately distant, at first segmentiform but later usually (broadly) ventricose, rarely veined, adnate-emarginate, sometimes almost free, pale grey-brown at first, later with pink tinge, finally red-brown (10 YR 6/3 or 7.5 YR 6/4, then 7.5 YR 6/4, 5/4, 5/2; 5 YR 5/3), with irregular, concolorous edge. Stipe 13–55(–75) × (2–)2.5–6 mm, cylindrical or flattened, rarely slightly broadened at base, sometimes twisted, concolorous with or slightly paler than pileus (10 YR 6/3, 6/4, 5/3, 5/2; 7.5 YR 6/2, 5/2, 4/2), usually longitudinally striate with adpressed or loose silvery fibrils, rarely glabrous and more or less polished, sometimes pruinose at apex, often white tomentose at base. Flesh concolorous with surface or (in fleshy specimens) paler, fibrous, brittle in pileus, brittle to subcartilaginous in stipe. Smell and taste strongly farinaceous-rancid.

Spores (7.0–)7.4–10.4 × (6.4–)6.7–8.7(–9.3)  $\mu\text{m}$ , Q = 1.0–1.1–1.2(–1.25), L–D = 0–1–1.7  $\mu\text{m}$ , variable in shape, from rather regularly 5-angled isodiametrical to irregularly 6–7(–8)-angled isodiametrical in side-view, with basal facet. Basidia 24–45 × 10.5–12.7(–17.5)  $\mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular, composed of cylindrical to inflated cells at least up to 240 but often up to 400  $\mu\text{m}$  long and 14–30(–50)  $\mu\text{m}$  wide, intermixed with narrow, cylindrical, minutely encrusted hyphae. Pileipellis a thin cutis made up of 2.7–12  $\mu\text{m}$  wide, cylindrical hyphae with minutely to coarsely encrusted walls; subpellis usually weakly developed and more intensively pigmented than suprapellis. Pileitrarma regular with cylindrical to inflated cells up to 350(–400)  $\mu\text{m}$  long and 12–27(–33)  $\mu\text{m}$  wide. Clamp-connections present in hymenium, often difficult to find because of their proliferation. Vascular hyphae sometimes rather abundant in trama.

HABITAT & DISTRIBUTION.—In natural grasslands extensively grazed by cattle, as well as in lawns, meadows, and hayfields. Fairly common in western and northern Europe.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Friesland, Isl. of Terschelling, 30 Aug. 1958, *C. Bas* 1513; prov. Overijssel, Hengforderwaarden between Deventer and Olst, 12 Nov. 1977, *H. Piepenbroek* & *G. Piepenbroek-Grooters* 1066; prov. Gelderland, Doetinchem, in garden, 10 Jan. 1948, *H. S. C. Huijsman*; Wilp, on dyke, Nov. 1977, *H. Piepenbroek* & *G. Piepenbroek-Grooters* 1053; prov. Utrecht, Rhenen, along Oude Veenendaalseweg, 16 Sept. 1955, *C. Bas* 878; prov. Noord-Holland, Isl. of Texel, near Slufter, 29 Oct. 1971, *P. B. Jansen* 71.194; Wieringermeer, Robbenoordbos, 23 Oct. 1976, *C. Bas* 7108A; Castricum, Geversduin, 13 Oct. 1954, *G. D. Swanenburg de Veye* and 21 Oct.

1956, *R. A. Maas Geesteranus 11856*; idem, Waterleidingduin, 10 Aug. 1952, *R. A. Maas Geesteranus 8902*; Velzen, Duin en Kruidberg, 3 Nov. 1965, *E. Kits van Waveren*; IJmuiden, Midden Heerenduin, 6 Nov. 1976, *M. E. Noordeloos 253*; Noordwijkerhout, dunes of Amsterdam Water Supply, 3 Oct. 1953, *C. Bas 667*; idem 29 Nov. 1953, *R. A. Maas Geesteranus 9610*; idem 4 Nov. 1976, *M. E. Noordeloos 250*; idem, State forest, 8 Dec. 1976, *M. E. Noordeloos 275*; Huizen, 18 Oct. 1976, *J. Frencken*; prov. Zuid-Holland: Katwijk a/Zee, 9 Nov. 1952, *R. A. Maas Geesteranus 9294*; Wassenaar, Meyendel, 17 Nov. 1974, *C. Bas 6511*; idem 16 Oct. 1979, *C. Bas 7572*; Wassenaar, estate 'Den Deyl', Maaldrift, 18 Sept. 1955, *R. A. Maas Geesteranus 10706*; prov. Zeeland: Walcheren, estate 'Zeeduin', 10 June 1937, *H. S. C. Huijsman 1262*; idem Vrouwenpolder, 31 Oct. 1976, *M. E. Noordeloos 244*; prov. Noord-Brabant: Drunen, Drunense Duinen, 15 Aug. 1976, *N. Hanegraaff-Lissenberg*; idem, along Drongelens Kanaal, 19 Dec. 1971, *P. B. Jansen 71-196*.

BELGIUM, prov. Namur, Ave-et-Auffe, near 'le Roptai', 6 Oct. 1977, *M. E. Noordeloos 495*. — GERMAN FEDERAL REPUBLIC, Westphalen, Detmold, Externsteine, 7 Oct. 1976, *M. E. Noordeloos 215*; idem, Merlshiem, 6 Oct. 1976, *G. J. M. G. Tjallingii-Beukers*.

For a discussion, see below under *E. sericeum* var. *sericeum* f. *nolaniforme*.

22. var. *SERICEUM* f. **nolaniforme** (Kühn. & Romagn.)

Noordeloos, *comb. & stat. nov.*—Figs. 22a-d

*Rhodophyllus sericeus* var. *nolaniformis* Kühn. & Romagn. in *Rev. Mycol.* 19: 9. 1954.

SELECTED ICONS AND DESCRIPTION.—Arnolds & Noordeloos, in *Fung. rar. Icon. col.* 12: pl. 92A. 1980.


CHARACTERISTICS.—Carpophores slender, mycenoid; pileus 15–37(–45) mm broad, usually pronouncedly papillate and with strongly involute margin; lamellae deeply emarginate to almost free; stipe 20–70 × 2–5 mm, paler than pileus, strongly silvery striate.

Pileus 15–37(–45) mm broad, conical then expanding to conico-convex, finally convex, usually pronouncedly papillate, sometimes with abrupt acutely conical papilla, rarely more or less flattened, with margin strongly involute when young and expanding only in later stages, strongly hygrophorous, when moist dark sepia or reddish brown (10 YR 2/1, 2/2, 3/2, 4/2, 4/3; 7.5 YR 3/2, 4/2), slightly paler towards margin (10 YR 5/3, 6/4, 7/4), translucently striate up to two-thirds of the radius, on drying strongly pallescent to greyish brown (10 YR 4/3, 5/4, 6/4, 7/4), lustrous-fibrillose. Lamellae L = 25–50, l = 3–5, moderately distant to fairly crowded, deeply emarginate to almost free, segmentiform to ventricose, pale, then pink with distinct grey or brown tinge, finally brown-pink (10 YR 7/3, 6/3 then 7.5 YR 4/2, 5/4, 6/4), with concolorous or slightly paler, slightly irregular to distinctly serrulate edge. Stipe 20–70 × 2–5 mm, cylindrical, grey-brown or yellowish brown (10 YR 3/2, 3/3, 4/3, 5/3, 6/4, 7/4), strongly silvery striate with aeriferous longitudinal fibrils, often white tomentose at base. Flesh concolorous with surface or slightly paler, brittle. Smell and taste strongly farinaceous-rancid.

Microscopical characters and habitat as in type form.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Friesland, Isl. of Vlieland, in meadow S. of Vianenslid, N. of Postweg, 2 Nov. 1976, *C. Bas 7148, 7149*; prov. Drenthe, Beilen, 8 Nov. 1977, *E. Arnolds*; prov. Gelderland: Arnhem, 25 Sept. 1978, *J. Hoogschagen*; Wilp, 9 Oct. 1977, *H. Piepenbroek & G. Piepenbroek-Grooters 1051, 1052*; prov. Noord-Holland: Callantssoog, nature reserve 'Zwanewater', 5 Nov. 1977, *C. Bas*; Velzen, Duin-en Kruidberg, 13 Nov. 1962, *C. Bas 2909*; prov. Zuid-Holland, Wassenaar, Den Deyl-Maaldrift, 18 Sept. 1955, *R. A. Maas Geesteranus 10707*.

*Entoloma sericeum* is rather variable species both in habit and in microscopical characters. The typical habit of *E. sericeum* is somewhat entolomatoid. Therefore many mycologists placed the species in subgenus *Entoloma*. In my opinion the type of pigmentation and the size and shape of



Figs. 22a-d. *Entoloma sericeum* f. *nolaniforme*. — Habit and spores (22a, 22c from Bas 7149; 22b, 22d from Bas, 5 Nov. 1977).

Figs. 23a-f. *Entoloma sericeum* f. *sericeum*. — Habit and spores (23a, 23d from Noordeloos 244; 23b, 23c from Maas Geesteranus 10706; 23c, 23f from Noordeloos 259).

the tramal elements make *E. sericeum* a true *Nolanea*. Nolaneoid forms of *E. sericeum* occur, and are distinguished here as forma *nolaniforme*. This form is frequently met with in the same habitat as *E. sericeum*, sometimes in separate groups and easily to distinguish from the typical form; sometimes even it is evident that both forms occur on the same mycelium, e.g. in one fairy ring. For the sake of convenience I decided to give the nolaneoid forms the rank of forma but I do not attach taxonomic value to it. There are no differences in microscopy between f. *sericeum* and f. *nolaniforme*.

A dark grey '*Entoloma sericeum*' with a non-striate pileus is distinguished as a new variety, viz. *E. sericeum* var. *cinereo-opacum* (see below).

The stipe of *E. sericeum* is usually covered with loose (aeriferous) fibrils which give the surface a striate appearance. Sometimes, however, specimens were found with a more or less glabrous and smooth stipe without these fibrils. Perhaps this is *Nolanea radiata* sensu Orton. I do not think that at least my collections of this form deserve a taxonomic rank, as there are no significant other differences with the typical form.<sup>3</sup>

The size and shape of the spores are rather variable, as indicated in the description above. The intensity of the pigmentation, particularly that of the pileipellis, is rather variable as well. Clamp-connections are often difficult to find due to proliferation (see Bas 1965: 355, 1969: 319), but careful observation always revealed clamp-connections at least at the base of the youngest basidia.

In the field *E. sericeum* can be confused with other grassland species such as *E. papillatum*, *E. hirtipes* var. *sericeoides* and *E. vernum*, but all these species have distinctly heterodiametrical spores. *Entoloma juncinum*, occurring sometimes in the same habitat, differs by the slender mycenoid habit, slightly different colours and the type of pigmentation.

### 23. ENTOLOMA SERICEUM var. *cinereo-opacum* Noordeloos,

var. nov.—Figs. 25a–d

A var. *sericeo* differt pileo haud striato, obscure cinereo, opaco, superficie incana lamellisque obscure griseis. Typus: *M. E. Noordeloos* 566, 26 X 1977 'Westerslag, Isl. Texel, prov. Noord-Holland, Netherlands' (L).

CHARACTERISTICS.—Differs from *E. sericeum* var. *sericeum* by the non-striate, dark grey-brown to blackish brown pileus with minute white pruinum causing a hoary impression (lens), and by the grey lamellae.

Pileus 13–44 mm broad, obtusely rounded conical or conico-convex with flattened or slightly depressed, rarely subpapillate centre, with margin inflexed then straight, often undulating and slightly reflexed with age, strongly hygrophanous, when moist blackish brown (10 YR 2/1, 2/2, 3/1, 3/4; 7.5 YR 2/1, 3/1, 3/2; 5 YR 3/2, 3/3), non-striate or striatulate at margin only, with minute whitish pruinum causing a hoary impression (lens), sometimes zonate, on drying strongly pallescent to brown with ochre tinge particularly at centre (10 YR 5/2, 5/3; 7.5 YR 6/4, 5/4), dull.

<sup>3</sup> *Rhodophyllus radiatus* J. Lange in its original concept is another taxon, probably belonging to subgenus *Entoloma*, and is perhaps close to *Entoloma sordidulum* (Kühn. & Romagn.) P. D. Orton (see also under the excluded taxa, p. 528).



Lamellae L = 20–30, l = 3–9, distant, adnexed, broadly adnate or deeply emarginate to almost free, grey (10 YR 7/2, 6/2, 6/3), hardly tinged pink when mature, with concolorous, subserulate edge. Stipe 25–70 × 2–5 mm, cylindrical or flattened with longitudinal groove, grey-brown, almost concolorous with pileus (10 YR 3/2, 4/2, 4/3; 7.5 YR 6/4, 5/4), scarcely to abundantly silvery striate with aeriferous fibrils, not or minutely pruinose at apex, abundantly white tomentose at base. Flesh concolorous with surface, subcartilagineous. Smell strong, rancid-farinaceous. Taste unpleasant, rancid, farinaceous.

Spores (7.1–)8.0–10.0(–11) × 7.1–8.7(–10.0)  $\mu\text{m}$ , Q = 1.0–1.1–1.2(–1.3), L–D = 0.6–1.2–1.8  $\mu\text{m}$ , obtusely 5–8-angled in side-view, probably with a basal facet. Basidia (27–)30–45(–52) × 10.5–15  $\mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular, cells cylindrical, (75–)125–290 × 5.0–19  $\mu\text{m}$ , brown, sometimes minutely encrusted. Pileipellis a thin cutis made up of cylindrical, 3.5–10  $\mu\text{m}$  wide hyphae with brown encrusted walls. Subpellis usually well developed, composed of short, inflated cells 27–78 × 10.8–16  $\mu\text{m}$ , with coarsely brown encrusted walls. Pileitrama regular; cells in deeper layers cylindrical, 70–300 × 8–14  $\mu\text{m}$ , with brown encrusted walls. Vascular hyphae numerous in some specimens. Clamp-connections numerous in hymenium and in trama of pileus, rare elsewhere.

HABITAT & DISTRIBUTION.—On poorly grazed, not artificially fertilized dune meadows on the West-Frisian islands in the North of the Netherlands and the German Federal Republic. October–November.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Friesland, Isl. of Terschelling, Oosterend, Dazenplak, 3 Nov. 1978, *M. E. Noordeloos 846*; Isl. of Ameland, Nes, 8 Nov. 1979, *T. Kuijper 1387*; prov. Noord-Holland: Isl. of Texel, Westerslag, 26 Oct. 1977, *M. E. Noordeloos 566* (holotype); idem, nature reserve 'de Geul', 28 Oct. 1977, *M. E. Noordeloos 579*.

GERMAN FEDERAL REPUBLIC, Ost Friesland, Isl. of Borkum, in dunes between Südstrand and village, 29 Oct. 1977, *C. Bas 7251*.

*Entoloma sericeum* var. *cinereo-opacum* is easily distinguished from the type form by the predominantly grey colour of the entire carpophore and by the non-striate, more or less hoary pileus. So far it has only been found in dune meadows on the West-Frisian islands. It is remarkable that I failed to find this characteristic form in dune meadows in the western and south-western parts of the Netherlands. One collection, made by the late Mr. H. F. van der Laan in the new polders of the former Southern Sea (Oostelijk Flevoland, recreation grounds in 'de Abbert', 10 Nov. 1976), probably belongs to var. *cinereo-opacum* but there are some slight differences in colour of lamellae and stipe, more typical for var. *sericeum*.

The habit of *E. sericeum* var. *cinereo-opacum* shows the same variation as that of the type variety, and also includes entolomatoid as well as nolaneoid forms.

#### 24. *Entoloma sericeoides* (J. Lange) Noordeloos,

*comb. nov.*—Figs. 24a–c

*Rhodophyllum sericeoides* J. Lange, Fl. agar. dan. 5: 99, pl. 198E. 1940 (basionym).


CHARACTERISTICS.—Pileus convex, moderately to deeply umbilicate, dark sepia; lamellae dark brown; stipe slightly paler than pileus, fibrillosly striate; smell weak; spores isodiametrical; pigment encrusting.

Pileus 15–30 mm broad, convex with rather shallow to deep umbilicus, with involute margin expanding only in later stage, hygrophanous, when moist dark sepia brown in centre (10 YR 3/2

to 5 YR 3/2), towards margin slightly paler (about 7.5 YR 3/2–4/2), translucently striate up to half of the radius, on drying pallescent to moderately dark brown (10 YR 6/3), brilliantly shining. Lamellae L about 40, 1=3–5, crowded, broadly adnate with decurrent tooth to slightly ascending-emarginate, segmentiform to narrowly ventricose, brown, later with pink tinge (7.5 YR 6/4 to 5/4), with slightly irregular, concolorous edge. Stipe 35–40 × (2–)3–5 mm, cylindrical or tapering upwards, sometimes twisted, concolorous with dry pileus, shingly fibrilously striate. Flesh cartilagineous; white. Smell indistinct or weakly farinaceous. Taste not tried.

Spores 7.3–9.3 × (5.8–)6.4–8.1 μm. Q = 1.0–1.1–1.2(–1.25), L–D = 0–1.0–1.7 μm, isodiametrically 5–6-angled in side-view, with basal facet. Basidia 30–36 × 10.4–11.5 μm, 4-spored. Cystidia absent. Hymenophoral trama regular; cells cylindrical, 120–200(–240) × 9.5–15 μm, with brown, sometimes minutely encrusted walls. Pileipellis a poorly differentiated cutis made up of radially arranged, cylindrical, 6–12 μm wide hyphae with (coarsely) encrusted walls, gradually passing into trama. Pileitrama regular with inflated cells up to 300 μm long and 6.5–20 μm wide. Clamp-connections not seen in hymenium, nor in other tissues.

**HABITAT & DISTRIBUTION.**—In extensively grazed grassland on dyke along river and in probably grassy clearing between *Fagus* forest and *Pinus* plantation. Recorded from the Netherlands and Denmark. Rare.



Figs. 24a–c. *Entoloma sericeoides*. — Habit and spores (24a–b from *Piepenbroek 1007*; 24c from holotype).  
 Figs. 25a–d. *Entoloma sericeum* var. *cinereo-opacum*. — Habit and spores (25a, 25d from holotype; 25b–c from *Bas 7251*).

COLLECTIONS EXAMINED.—NETHERLANDS, prov. Overijssel, Hengforder waarden between Olst and Deventer, 10 Oct. 1976, H. Piepenbroek & G. Piepenbroek-Grooters 1007.

DENMARK, Isl. of Fyn, Sønderskov near Kvaerndrup, 12 July 1938, J. Lange (holotype, C).

*Entoloma sericeoides* resembles *E. sericeum* very much in colour of the carpophore, type of pigmentation, and in the isodiametrical spores. The main differences are the deeply umbilicate pileus, the lack of clamp-connections in the hymenium, and the lack of a distinct farinaceous smell. Another collection from the Netherlands—prov. Zuid-Holland, Isl. of Goeree, 9 Nov. 1977, G. J. M. G. Tjallingii-Beukers—is quite similar to *E. sericeoides*, but differs by the strong farinaceous-rancid smell and the presence of clamp-connections in the hymenium, which makes it a form of *E. sericeum*. It is clear that more material is required for study before a more satisfactory decision can be made on the true status of *E. sericeoides*.

## 25. *Entoloma sphaerocystis* Noordeloos, *spec. nov.*—Figs. 11a–b, 49

Pileus 16 mm latus, conicus, hygrophanus, striatus, badius, expallens, fibrilloso-sericeus. Lamellae subliberae, ventricosae, roseae brunneo-tinctae. Stipes 32 × 1.5 mm, cylindraceus, fulvus. Odore nullo. Sporae 8.5–11 × 7–8.1 μm, Q = 1.15–1.25–1.4, 6–7-angulatae basi nominatur 'dihedral base'. Cheilocystidia (sub-)globosa, 25–38(–45) × 13–35 μm. Pileipellis cutis, hyphae 3.5–7.5 μm latae. Pigmentis incrustatus. Fibulae ad basim basidiorum frequentes. Typus: J. Schreurs, 19 VIII 1979, 'Willinks Weust, Winterswijk, prov. Gelderland, Netherlands' (L).

CHARACTERISTICS.—Carpophores mycenoid; pileus conical, reddish brown, pallescent and lustrous on drying; lamellae brownish pink; stipe minutely striate with some scattered silvery fibrils; spores 8.5–11 × 7–8.1 μm; cheilocystidia present, subglobose; pigment minutely encrusting; clamp-connections present.

Pileus 16 mm broad, conical with straight margin, hygrophanous, when moist reddish brown (7.5 YR 4/4, 5/4; 5 YR 4/4), paler at margin, translucently striate up to three fourths of the radius, on drying strongly pallescent to pale grey, strongly fibrillous with loose-lying fibrils, shining. Lamellae L = 24, l = 1–3, almost free, ventricose, brownish pink (7.5 YR 6/4, 5/4). Stipe 32 × 1.5 mm, cylindrical, subcartilagineous, pale yellowish brown (10 YR 5/4, 4/4), contrasting with pileus, almost smooth, with only very few silvery fibrils lengthwise. Flesh concolorous with surface. Smell absent.

Spores 8.5–11 × 7–8.1 μm, Q = 1.15–1.25–1.4, L–D = 1.7–2.0–2.3 μm, 6–7-angled in side-view, with dihedral base. Basidia 26–35 × 8–15 μm, 4-spored. Cheilocystidia 25–38(–45) × 13–35 μm, (sub-)globose, rarely broadly clavate, sometimes with thickened, colourless walls. Hymenophoral trama regular, cells 170–360 × 14–23 μm. Pileipellis a cutis made up of radially arranged, cylindrical hyphae with brown, sometimes minutely encrusted walls. Pileitrama regular, cells slightly inflated, up to 270 μm long, 12–31 μm wide, with brown walls particularly in upper layers. Clamp-connections abundant in hymenium.

HABITAT & DISTRIBUTION.—In hayfield on rich, loamy soil (type) and on branchlet of *Fraxinus excelsior* in frondose forest, up till now only known from two localities in the Netherlands.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Gelderland, Winterswijk, Willinks Weust, 19 Aug. 1979, J. Schreurs (holotype); prov. Noord-Brabant, Nieuw Ginneken, Ulvenhoutse bos, 27 Aug. 1979, P. B. Jansen.

*Entoloma sphaerocystis* differs from *E. globulifer*, which has about the same type of cheilocystidia, by the differently shaped spores, the type of pigmentation, and the presence of

clamp-connections. On account of the presence of membranal and minutely encrusting pigments *E. sphaerocystis* is placed in subsection *Cosmeoxonema*. Macroscopically there is some resemblance with *E. sericeonitens*, which differs by the spores with basal facet and the absence of cheilocystidia.

### Entoloma section *Fernandae* Noordeloos, *sect. nov.*

Pigmentis duobus, pileipellis pileitramaeque hypharum tunicam valde incrustantibus, pariter in pileipelli intracellulosis-granulatis; sporis 5-6-7-angulatis cum structura nominatur 'dihedral base'; fibulae nullae. — Typus: *Entoloma fernandae* (Romagn.) Noordeloos.

Two types of pigment present: one encrusting the hyphae of pileipellis and pileitrama, the other intracellular in the form of (agglutinated) clots or granulae, particularly in the pileipellis; spores 5-6-7-angled in side view with distinctly dihedral base; clamp-connections absent. — Type: *Entoloma fernandae* (Romagn.) Noordeloos.

#### KEY TO THE SPECIES OF SECTION FERNANDAE

- 1a. Stipe strongly striate lengthwise (as in *E. conferendum*); pileus smooth, in moist state translucently striate up to centre. . . . . 2
- b. Stipe not striate but smooth or downy-woolly and then pileus rugulose-fluffy to subsquamulose and striate at margin only. . . . . 3
- 2a. Pileus pale grey-brown with yellow or russet tinge, pale greyish-ochraceous when dry; lamellae white then pink without any brown or grey tinge; flesh in pileus rather brittle; spores 7.9-10.3(-10.8) × (6.3-) 6.7-7.7(-8.0) μm; basidia 2- and 4-spored. . . . . *E. acidophilum*, p. 489
- b. Pileus rather dark grey-brown, brown-grey when dry; lamellae pale brown-grey, then sordid flesh colour; flesh in pileus more or less firm; spores 7.4-9.5(-10.6) × 5.8-6.9(-7.4) μm; basidia 4-spored. . . . . *E. argenteostriatum*, p. 490
- 3a. Spores rather strongly rounded-angular in side-view, (6.5-)6.7-7.9(-8.1) × (5.3-)5.8-6.9(-7.4) μm, Q = 1.1-1.2-1.3. . . . . *E. defibulatum*, p. 492
- b. Spores slightly larger and more pronouncedly angular in side-view, Q ≥ 1.3 on the average per collection. . . . . 4
- 4a. Stipe with distinct yellow tinge quite different from colour of pileus. . . . . *E. xanthocaulon*, p. 493
- b. Stipe greyish-brownish, more or less with the same colour as pileus, though usually paler. . . . . 5
- 5a. Spores (8.4-)9.0-11.4(-12.5) × (6.2-)6.9-7.7(-8.2) μm, Q = 1.3-1.45-1.7(-1.8). . . . . *E. cuniculorum*, p. 490
- b. Spores not longer than 10.5 μm and Q = 1.1-1.3-1.4. . . . . 6
- 6a. Pileus when young more or less conical, quickly expanding to convex or flattened, sometimes slightly depressed at centre, sometimes with small umbo, pale to moderately dark grey-brown; in grasslands. . . . . 7
- b. Pileus acutely conical, not or only slightly expanding, blackish brown; gregarious in coniferous forests. . . . . *E. fractum*, p. 492
- 7a. Pileus strongly hygrophanous, when moist translucently striate up to centre, absolutely smooth and glabrous; stipe brilliant, smooth and glabrous. . . . . *E. psilopus*, p. 489
- b. Pileus weakly hygrophanous, when moist translucently striate at margin only, rugulose-fluffy particularly at centre, sometimes even subsquamulose; stipe mostly smooth and glabrous, rarely minutely downy-woolly all over. . . . . *E. fernandae*, p. 487

26. *ENTOLOMA FERNANDAE* (Romagn.) Noordeloos—Figs. 29a–b

*Rhodophyllus fernandae* Romagn. in Rev. Mycol. 1: 162. 1936. — *Nolanea fernandae* (Romagn.) P. D. Orton in Trans. Br. mycol. Soc. 43: 179. 1960. — *Entoloma fernandae* (Romagn.) Noordeloos in Persoonia 10: 250. 1979.

SELECTED ICONES AND DESCRIPTIONS.—Romagn. l.c., pl. 9, fig. 3. 1936. — Arnolds & Noordeloos in Fung. rar. Icon. col. 12, pl. 91 fig. b. 1980.

CHARACTERISTICS.—Pileus and stipe grey-brown; pileus only weakly hygrophanous, translucently striate at margin or not; surface of pileus subfelted-rugulose to minutely squamulose, particularly at centre.

Pileus 12–35 mm broad, conical or campanulate-convex, soon expanding to plano-convex, with or without a faint central depression, rarely with small papilla, with margin straight when young, but often irregularly undulating and/or splitting with age, weakly hygrophanous, when moist translucently striate at margin only or not, pale to moderately dark grey-brown sometimes with reddish tinge (10 YR 3/3–3/4, 7.5 YR 4/4, 10 YR 4/3, 5/3, 6/3 at centre), paler towards margin, slightly pallescent when dry, radially fibrillous-satyne becoming filthy rugulose or minutely squamulose, particularly at centre, often brilliantly shining. Lamellae L = 22–30, l = 1–3(–5), almost free, (sub-) ventricose, white, then (sordid) pink, finally brownish pink (7.5 YR 6/4, 5/4), with concolorous, slightly irregular edge. Stipe 20–45 × 2–4 mm, cylindrical or flattened, with longitudinal groove(s), tapering or slightly broadening downwards, concolorous with pileus or slightly paler, sometimes pruinose at apex and white tomentose at base, often tomentum covering basal half of stipe, remaining surface shining, sometimes very minutely downy-pubescent, rarely faintly striate near basal tomentum, solid, then fistulose. Flesh concolorous with surface, inner parts slightly paler and fibrous in stipe. Smell and taste farinaceous.

Spores (7.0–)7.4–9.0(–9.6) × (5.4–)5.7–7.0(–7.3) μm, Q = 1.15–1.3–1.4(–1.5), L–D = 1.0–1.7–2.3(–2.8) μm, mostly 6-angled in side-view, with distinctly dihedral base. Basidia 23–33(–34) × 8–14 μm, 4(–rarely 2-)spored. Cystidia absent. Hymenophoral trama regular, composed of inflated hyphae with cells (110–)210–320(–600) × 12–32(–37) μm. Pileipellis a cutis made up of repent, 3–20 μm wide cylindrical hyphae with numerous repent or slightly ascending terminal cells sometimes in chains of 2–3–4 cells, with often deep brown, coarsely encrusted walls and with intracellular clustered clots of brown pigment. Pileitrama regular, composed of cylindrical to slightly inflated cells 190–320 × 16–34 μm, mixed with narrow 2.3–7 μm wide cylindrical hyphae, with brown encrusted walls, in upper layer often also intracellular pigment as in pileipellis. Clamp-connections absent.

HABITAT & DISTRIBUTION.—Terrestrial on poor acid soils, frequently met with in *Calluna*-heaths on the pleistocene soils in central and eastern parts of the Netherlands; also found on peat but never been observed growing in or on living *Sphagnum*. Known to occur in Great Britain (P. D. Orton & al., 1960: 125), the Netherlands, France and in the German Federal Republic.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Roden, Moltmakersstuk, 15 Sept. 1976, *M. E. Noordeloos* 177; prov. Utrecht, Leusden, near Pyramide of Austerlitz, 25 Aug. 1979, *A. F. M. Reijnders*; idem 30 Aug. 1979, *M. E. Noordeloos* 994; prov. Noord-Holland, Callantssoog, Eendekooi, 10 Oct. 1979, *M. E. Noordeloos* 1067; Hilversum, 13 Aug. 1979, *J. Frencken*; Vogelenzang, A. D. W.-dunes, 15 Sept. 1979, *C. Bas* 7477; prov. Noord-Brabant, Nieuw Ginneken, Goudbergven, 23 Aug. 1954, *R. A. Maas Geesteranus* 10120; Zundert, 'Krochten', 3 Sept. 1970, *C. Bas* 5324; Deurne, Peel, 23 Sept. 1961, *C. Bas* 2446b.

GERMAN FEDERAL REPUBLIC, Ost Friesland, Aurich, Ewiges Moor, 22 Aug. 1962, *C. Bas* 2620. — FRANCE, dept. Seine & Oise, Yerres, bois de l'Etoile, Aug. 1932, *H. Romagnesi* 209 (holotype, Herb. Romagn., PC).

*Entoloma fernandae* belongs to a group of closely related species, in the present work all placed together in the new section *Fernandae*. Romagnesi (in Kühn. & Romagn. 1953: 193) placed *Rhodophyllus fernandae* in the subgenus *Entoloma* because of the habit of the type specimens. In my opinion, however, *E. fernandae* and related species are far better placed in subgenus *Nolanea* on account of the type of pigmentation and tramal structures. Within subgenus *Nolanea* section *Fernandae* seems to form a bridge to subgenus *Pouzaromyces* because of the tendency to form a trichoderm-like pileipellis, the type of pigmentation, and the lack of clamp-connections. For more details the reader is referred to a publication (in preparation) on the general taxonomy in *Entoloma* (Noordeloos, 1980b).

Almost in complete accordance with the original diagnosis *E. fernandae* is defined here as a species with a weakly hygrophanous, hardly striate pileus with fluffy to subsquamulose surface and a smooth, rarely pruinose-downy stipe. *Entoloma psilopus* differs by having a smooth, deeply striate pileus, whereas *E. acidophilum* and *E. argenteostriatum* differ by having a strongly silvery-fibrillous surface of the stipe.

27. ENTOLOMA FERNANDAE f. **eccilioides** Noordeloos,  
f. nov.—Figs. 30a–b

A forma typica differt lamellis decurrentibus atque statura Ecciliae. Typus: *M. E. Noordeloos 1073*, 20 X 1979, 'Eendekooi, Callantsoog, prov. Noord-Holland, Netherlands' (L).

Pileus 20 mm broad, convex with umbilicus, with margin slightly involute, faintly hygrophanous, when moist grey-brown (10 YR 4/4, 4/3) and translucently striate at margin only, on drying pallescent to pale grey-brown (10 YR 5/3, 5/4) with marginal zone grooved on back of lamellae, entirely minutely fibrillous-squamulose, particularly at centre. Lamellae L = 15, l = 7–9, distant, deeply decurrent, sordid flesh colour (7.5 YR 6/4, 5/4), with slightly thickened, concolorous edge. Stipe 20 × 2–3 mm, cylindrical, horn brown (10 YR 6/4, 7/4), pruinose at apex, downwards smooth, glabrous, shining. Flesh firm in stipe, brittle in pileus, slightly paler than surface. Smell strong, farinaceous.

Spores 8.1–9.0 × 5.8–7.0 μm, Q = 1.1–1.25–1.4, L–D = 1.2–1.8 μm, 6-angled in side-view, with distinctly dihedral base. Basidia 27–42 × 8.1–11 μm, 4-spored. Cystidia absent. Hymenophoral trama regular with cells up to 250 μm long and 9–17 μm wide. Pileipellis a cutis with transitions to a trichodermium, composed of 4–12 μm wide cylindrical hyphae with repent or slightly ascending, terminal, up to 17 μm wide cells with brown-encrusted and/or deeply brown coloured walls and intracellular clots of pigment. Pileitrama regular, with coarsely encrusted hyphae. Clamp-connections absent.

HABITAT & DISTRIBUTION.—Terrestrial in *Calluna*-heath on poor, acid sandy soil. So far only found in type locality near typical *Entoloma fernandae*.

COLLECTIONS EXAMINED.—NETHERLANDS, prov. Noord-Holland, Callantsoog, Eendekooi, 20 Oct. 1979, *M. E. Noordeloos 1073* (holotypus, L.).

The eccilioid habit with long decurrent lamellae is considered so remarkable that the collection concerned is described as a new forma to draw attention to the fact that a species in subgenus *Nolanea* may have a strictly eccilioid appearance.

28. *ENTOLOMA PSILOPUS* Arnolds & Noordeloos

*Entoloma psilopus* Arnolds & Noordeloos in Persoonia 10: 293–294, Figs. 19–22. 1979.

**CHARACTERISTICS.**—Pileus (dark) grey-brown and translucently striate up to centre; when moist smooth, glabrous. Stipe paler than pileus, grey-brown, smooth and not striate.

Pileus 8–23 mm broad, conical when young, then hemispherical, expanding, finally flattened, mostly not papillate, sometimes with weak papilla, often slightly depressed at centre when old, with margin straight, sometimes exceeding the lamellae, hygrophanous, when moist dark grey-brown at centre, paler towards margin (10 YR 3/2; M 6E6, 6E6/6F6, 6E6/7E6), translucently striate up to centre, on drying pallescent to pale grey-brown (10 YR 6/4, 7/4; M 6C4, 5C4), smooth, mostly dull but sometimes lustrous, glabrous. Lamellae L=about 20, l=3–7, moderately crowded to rather distant, narrowly to broadly adnate or deeply emarginate, ventricose, white, then salmon pink or incarnate, sometimes slightly tinged brown (7.5 YR 7/4, 6/4; M 7A4/7B4, 7B4, 6B4), sometimes transversely veined, with entire, concolorous edge. Stipe 15–50 × 1–2.5(–3) mm, cylindrical, straight or flexuous, with base slightly broadened or not, solid, then fistulose, pale to rather dark grey-brown (10 YR 7/4, 6/4, base 5/4; M 4C4, 6C4, 6C4/D4, 6D5/E5), non-striate, smooth, with moderately to strongly developed tomentum at base. Flesh thin-membranaceous in pileus, rather firm or fragile in stipe but mostly more or less cartilaginous, concolorous with surfaces, in fleshy specimens inner part slightly paler and fibrous. Smell mostly distinctly farinaceous, sometimes spontaneously rather weak but then becoming distinct when cut. Taste weakly to distinctly farinaceous-rancid.

Spores (6.5–)7.0–9.1(–10.1) × 5.7–7.0(–7.5)  $\mu$ m. Q = 1.15–1.4, on the average between 1.25 and 1.3 per collection, L:D = 1.0–1.8–2.5  $\mu$ m, 5–6-angled in side-view, with dihedral base. Basidia (20–)23–40(–44) × 8.5–12.5(–14)  $\mu$ m, 4-(rarely 2)-spored. Hymenophoral trama regular, composed of cylindrical to distinctly inflated cells (55–)176–450 × 7.6–32  $\mu$ m, mixed up with 3–7  $\mu$ m wide cylindrical hyphae. Pileipellis a poorly differentiated cutis made up of 2.9–8  $\mu$ m wide cylindrical hyphae, at centre of pileus sometimes with some clavate cells 20–85 × 6–14(–18)  $\mu$ m, forming transitions to a trichodermium with brown encrusted walls and scattered to abundant, intracellular, mostly clustered clots of pigment. Pileipellis sometimes with subpellis of inflated cells 46–110 × 18–25  $\mu$ m, passing into regular trama with cylindrical to inflated cells up to 250  $\mu$ m long and up to 30  $\mu$ m wide. Clamp-connections absent.

**HABITAT & DISTRIBUTION.**—Terrestrial on oligotrophic, pleistocene sands and holocene peat in the northeastern and southeastern parts of the Netherlands. July–November.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Friesland, Appelscha, 3 Oct. 1968, R. Kramer (WBS); prov. Drenthe, Westerbork, Mantingerzand, 3 Nov. 1976, E. Arnolds 3734 (holotype); idem, Hullenzand, 3 Nov. 1976, E. Arnolds 3732 & 3733 and 15 Oct. 1974, E. Arnolds 3237 (WBS); Rolde, Eexterveld, Westerholt, N. E. of Anderen, 22 Sept. 1976, E. Arnolds 3595 (WBS); prov. Overijssel, Rijssen-Markelo, 11 July 1972, B. de Vries (WBS); prov. Gelderland, Kootwijkerveld, Sept. 1970, G. S. de Hoog; prov. Noord-Brabant, Bergen op Zoom, Zoomland, 7 July 1974, P. B. Jansen 74–272; Alphen, 'Ooyevaarsnest', 26 July 1979, P. B. Jansen 79–148; prov. Limburg, Grote Peel reserve near Ospel, 22 Aug. 1972, P. B. Jansen 72–213.

*Entoloma psilopus* is easily distinguished from *E. fernandae* by the more hygrophanous, strongly striate, absolutely smooth pileus. *Entoloma xanthocaulon* differs by the yellow-tinged stipe.

29. *ENTOLOMA ACIDOPHILUM* Arnolds & Noordeloos

*Entoloma acidophilum* Arnolds & Noordeloos in Persoonia 10: 285. 1979.

ICON AND DESCRIPTION.—Arnolds & Noordeloos in Fung. rar. Icon. col. 12: pl. 91 fig. a. 1980.

**CHARACTERISTICS.**—Pileus 8–16 mm broad, conical or hemispherical, expanding to convex with or without small papilla, finally flattened, slightly depressed at centre, hygrophanous, pale grey-brown when moist, deeply translucently striate, on drying pallescent, smooth; lamellae  $L = 12-17$ ,  $l = (1-3)-4$ , moderately distant, narrowly adnate to almost free, pale then pink; stipe 10–32(–53)  $\times$  1–2.5(–3) mm, pale greyish brown, silvery striate, apex sometimes pruinose; smell strongly farinaceous; spores 7.9–10.2(–10.8)  $\times$  6.3–7.5(–8.0)  $\mu\text{m}$ ,  $Q = 1.15-1.3-1.4$ ; basidia 2- and 4-spored; cystidia absent; pileipellis a poorly differentiated cutis made up of 4–10  $\mu\text{m}$  wide hyphae gradually passing into pileitrama with brown, granularly intracellular pigment and minutely encrusted walls; clamp-connections absent.

**HABITAT & DISTRIBUTION.**—In grass on acid sandy soil near entrance of a rabbit-hole. Known from five localities in the provinces of Drenthe, Friesland and Noord-Brabant, Netherlands.

The pale, strongly hygrophanous pileus, distant lamellae and silvery striate stipe distinguish *Entoloma acidophilum* from *E. fernandae*. *Entoloma psilopus* has a more intensively pigmented pileus and a smooth, non-striate stipe. *Entoloma argenteostriatum* differs by the darker pileus, firm flesh and brown-tinged lamellae. So far the 2- and 4-spored basidia equally distributed in the hymenium of *E. acidophilum* are unique in section *Fernandae*.

Up to now the species has been found in three different localities in the province of Drenthe (Arnolds & Noordeloos, 1980, l.c.) and in addition in the province of Friesland, Heerenveen, Oranjewoud, 31 July 1979, *J. Wisman*, and in the province of Noord-Brabant, Bergen op Zoom, Zoomland, 5 July 1974, *P. B. Jansen* 74–271.

### 30. ENTOLOMA ARGENTEOSTRIATUM Arnolds & Noordeloos

*Entoloma argenteostriatum* Arnolds & Noordeloos in *Persoonia* 10: 285–287, figs. 1–4. 1979.

**CHARACTERISTICS.**—Pileus 9–25 mm broad, conico-convex or convex, then expanding, with straight, often undulating margin, strongly hygrophanous, when moist dark grey-brown, translucently striate up to centre, on drying pallescent, smooth; lamellae moderately distant, variably inserted, pale brown-grey, then dingy flesh-coloured; stipe 23–32  $\times$  1.8–4 mm, cylindrical or flattened, grey-brown paler than pileus, silvery and aeriferously striate lengthwise; smell strongly farinaceous-cucumberlike; spores 7.4–9.5(–10.6)  $\times$  5.8–6.9(–7.4)  $\mu\text{m}$ ,  $Q = 1.2-1.35-1.5$ ; basidia 4-spored; cystidia absent; pileipellis a cutis made up of 4–12(–13)  $\mu\text{m}$  wide hyphae with coarsely encrusted walls and granularly intracellular pigment; clamp-connections absent.

**HABITAT & DISTRIBUTION.**—In poor vegetation of short grass and moss (*Polytrichum piliferum*) on dry, acid sandy soil. Known only from the type locality in the province of Drenthe, Netherlands.

*Entoloma argenteostriatum* differs from *E. acidophilum* by the more intensively pigmented pileus, firm flesh and 4-spored basidia; from *E. fernandae* mainly by the entirely smooth, translucently striate pileus and strongly striate stipe. *Entoloma psilopus* differs by the smooth stipe.

So far *E. argenteostriatum* has only been found in the type locality and in a nearby locality: province of Drenthe, Spier, Reigersplas, 22 Sept. 1964, *J. J. Barkman* (WBS).

### 31. ENTOLOMA CUNICULORUM Arnolds & Noordeloos—Fig. 27

*Entoloma cuniculorum* Arnolds & Noordeloos in *Persoonia* 10: 289–290, figs. 11–14. 1979.



CHARACTERISTICS.—Pileus 12–16 mm broad, bluntly conical or hemispherical with small papilla, with straight margin sometimes exceeding the lamellae, hygrophanous, when moist rather pale grey-brown, with dark brown striae up to centre, on drying pallescent; lamellae moderately distant, narrowly adnate, pale grey-brown then with pink tinge; stipe 20–32 × 1.5–2 mm, rather pale grey-brown, not striate, smooth; smell rather strongly farinaceous; spores (8.4–)



Figs. 26a–c. *Entoloma xanthocaulon*. — Habit and spores (26a from Noordeloos 407; 26b–c from Noordeloos 400).

Fig. 27. *Entoloma cuniculorum*. — Spores (from holotype).

Fig. 28. *Entoloma defibulatum*. — Spores (from holotype).

Figs. 29a–b. *Entoloma fernandae* f. *fernandae*. — Habit and spores (29a–b from Noordeloos 177).

Figs. 30a–b. *Entoloma fernandae* f. *eccilioides*. — Habit and spores (30a–b from holotype).

Figs. 31a–b. *Entoloma fractum*. — Habit and spores (31a–b from Barkman 7164).

9.0–11.4(–12.5) × (6.0–)6.2–7.7(–8.2)  $\mu\text{m}$ ,  $Q = 1.3–1.45–1.7(–1.8)$ ; basidia 4-spored; cystidia absent; pileipellis a dry cutis made up of (4.5–) 6–14  $\mu\text{m}$  wide cylindrical hyphae with brown encrusted walls and diffusely and/or granularly intracellular pigment; clamp-connections absent.

**HABITAT & DISTRIBUTION.**—In poor vegetation of moss-interspersed short grass, grazed by rabbits, on dry acid, sandy soil. Known only from two different localities in the province of Drenthe, Netherlands.

*Entoloma cucicularum* is quite distinct in section *Fernandae* by its large and irregularly shaped spores. The smooth and non-striate stipe and the colours of pileus and stipe remind of *E. psilopus*.

The macroscopic characters seem to point to a relationship with some members of section *Papillata* but they are contradictory to the microscopic characters such as type of pigmentation and lack of clamps, which make it necessary to place *E. cucicularum* in section *Fernandae*.

### 32. ENTOLOMA DEFIBULATUM Arnolds & Noordeloos—Fig. 28

*Entoloma defibulatum* Arnolds & Noordeloos in *Persoonia* 10: 290–291, figs. 15–18. 1979.

**CHARACTERISTICS.**—Pileus 19–21 mm broad, conico-campanulate, then expanding, not papillate, hygrophanous, when moist dark grey-brown, paler towards margin, translucently striate up to centre, on drying pallescent, smooth; lamellae  $L = 16–21$ ,  $l = 3–7$ , moderately distant, narrowly adnate to almost free, pale grey-brown, then with pink tinge; stipe 28–40 × 2 mm, cylindrical, rather dark greyish brown, finely striate under lens, apex minutely pruinose; smell and taste cucumber-like or farinaceous-rancid; spores (6.5–)6.7–7.9(–8.1) × (5.3–)5.6–6.8(–7.0)  $\mu\text{m}$ ,  $Q = 1.1–1.2–1.3(–1.4)$ ; basidia 4-spored; cystidia absent; pileipellis a poorly differentiated cutis made up of (2.8–)5–12(–15)  $\mu\text{m}$  wide cylindrical hyphae with encrusted walls and granularly intracellular pigment; clamp-connections absent.

**HABITAT & DISTRIBUTION.**—In poor vegetation of moss and short grass on rather moist to rather dry acid peat or humus-rich sand. Known only from two different localities in the province of Drenthe, Netherlands.

*Entoloma defibulatum* differs from all other species in section *Fernandae* by the rather strongly rounded, only weakly angled spores. Macroscopically it resembles *E. psilopus* but the stipe appears to be minutely striate under a lens, although not so silvery striate as in *E. argenteostriatum* and *E. acidophilum*. The type of spores of *E. defibulatum* resembles that of *E. ortonii* but the latter species has clamped basidia, lacks intracellular pigment and has a quite different habit.

### 33. ENTOLOMA FRACTUM (Velen.) Noordeloos—Figs. 31a–b

*Nolanea fracta* Velen., *Novitates mycologicae*: 146. 1939. — *Entoloma fractum* (Velen.) Noordeloos in *Persoonia* 10: 248. 1979.

Pileus 10–20 mm broad, acutely conical, hygrophanous, blackish at centre, blackish brown towards margin, translucently striate when moist, glabrous, subrugose at centre, margin acutely deflected. Lamellae crowded, white, then greyish pink, broad. Stipe 20–40 × 1–2 mm, concolorous with pileus, glabrous, fistulose.

Spores  $6.8-7.9 \times (4.5-5.7-6.7 \mu\text{m})$ ,  $Q = 1.1-1.2-1.4$ ,  $L-D = 1.2-1.8-2.4 \mu\text{m}$ , 5-6-angled in side-view. Basidia  $25-32 \times 7.6-10.2 \mu\text{m}$ , 4-spored. Hymenophoral trama regular; cells cylindrical to inflated, up to  $270 \mu\text{m}$  long and  $10-24 \mu\text{m}$  wide, hyaline, thin-walled, colourless. Pileipellis a cutis with transitions to a trichodermium, composed of cylindrical to slightly inflated hyphae  $7-17(-22) \mu\text{m}$  wide, at centre of pileus with tufts of clavate, terminal cells  $32-60 \times 8-22 \mu\text{m}$ , with brown encrusted walls and olivaceous-brown intracellular clots of pigment. Pileitrama regular with cylindrical to slightly inflated, brown encrusted hyphae. Clamp-connections absent.

HABITAT.—Terrestrial in *Pinus* forest on clayey soil.

COLLECTION EXAMINED.—CZECHOSLOVAKIA, Bohemia, Mnichovice, Struhařov, July 1936, J. Velenovský (holotype, PRM 154483).

A microscopical study of the type collection of this species revealed close relationship with *E. fernandae*. It differs from all species described above by the acutely conical, rather darkly pigmented pileus and by the habitat. In WBS one collection is present with the following characters suiting several important characters in Velenovský's description:

Pileus 14 mm broad, convex with conical papilla, dark grey-brown, not translucently striate, weakly hygrophanous, glabrous, smooth, strongly silky-shining. Lamellae almost free, ascending, ventricose, greyish pink with entire, concolorous edge. Stipe  $40 \times 1.5$  mm, cylindrical, fistulose, glabrous, only minutely pruinose at apex, dark grey-brown, shining.

Spores  $7.9-9.0 \times 5.7-6.9 \mu\text{m}$ ,  $Q = 1.25-1.3-1.4$ ,  $L-D = 1.7-2.3 \mu\text{m}$ , 5-6-angled in side-view, with distinctly dihedral base. The material is in poorly dried state, therefore a critical study of the tissues appeared to be impossible. In the surface layer of the pileus, however, the pigmentation appeared to be clearly of two kinds: coarsely encrusting and granularly intracellular.

HABITAT.—In old *Pinus* forest on poor, acid sandy soil in open spot in dense moss layer of *Pleurozium schreberi*, *Dicranum scoparium*, *D. undulatum*, etc.

COLLECTION EXAMINED.—NETHERLANDS, prov. Drenthe, Lhee, state forest, 11 Oct. 1961, J. J. Barkman 7164 (WBS).

On account of the shape of the pileus, the dark pigmentation as well as the habitat this collection, certainly representing a member of section *Fernandae*, might be considered to belong to *Entoloma fractum*. As the collection consists of only one specimen in a poor state, the occurrence of Velenovský's species in the Netherlands remains somewhat doubtful also because of the slight differences in macroscopy such as the non-striate pileus and the slightly larger and more elongate spores of the Netherlands' collection.

#### 34. ENTOLOMA XANTHOCAULON Arnolds & Noordeloos—Figs. 26a-c

*Entoloma xanthocaulon* Arnolds & Noordeloos in Persoonia 10: 299. 1979.

ICON AND DESCRIPTION.—Arnolds & Noordeloos in Fung. rar. Icon. col. 12: pl. 91 fig. c. 1980.

CHARACTERISTICS.—Pileus brown with pinkish or reddish tinge; stipe with distinct yellow tinge and therefore contrasting with the pileus.

Pileus 13-32 mm broad, conical, then expanding to convex, finally flattened, rarely with small papilla, sometimes with slight depression, with margin often slightly to distinctly involute and exceeding the lamellae, strongly hygrophanous, when moist pale to rather dark reddish or

pinkish brown (centre 10 YR 6/3, 5/3; 7.5 YR 4/2, 4/4, 3/2; 5 YR 3/3, paler towards margin, 10 YR 6/3, 6/4, rarely 7/3), translucently striate up to centre, on drying strongly pallescent to ochraceous-greyish ( $\pm$  10 YR 7/2, 7/3), smooth, glabrous, shining. Lamellae  $L = (15-20-30, 1 = (1-3)-5$ , adnate or emarginate, sometimes almost free, pale then pink, often without any grey or brown shade (7.5 YR 7/4), with entire concolorous edge. Stipe  $20-50 \times 1-3$  mm, cylindrical, sometimes slightly broadened at base, with predominant yellow tinge (10 YR 7/4, 5/4; 2.5 Y 7/4, 6/4, 5/4), in colour remarkably different from pileus, smooth, polished, sometimes slightly pruinose at apex, white tomentose at base, solid then fistulose. Flesh thin-membranaceous, fragile, concolorous with surface. Smell and taste strongly farinaceous.

Spores  $7.2-9(-9.3) \times 5.7-7.4(-7.6) \mu\text{m}$ ,  $Q = (1.1-1.15-1.3-1.4, L-D = 1.2-1.9-3 \mu\text{m}, 5-6$ -angled in side-view, with distinctly dihedral base. Basidia  $24-42 \times 8-11.5(-12.6) \mu\text{m}$ , 4-(rarely 2-) spored. Hymenophoral trama regular with cylindrical to slightly inflated cells  $159-370 \times 11-32 \mu\text{m}$ . Pileipellis a thin cutis made up of  $2.5-10(-11) \mu\text{m}$  wide cylindrical hyphae rarely with clavate terminal cells up to  $12(-15) \mu\text{m}$  wide, with pale intracellular pigment, and that sometimes granular or even in large clots sticking to inner walls of hyphae and with additional brown, usually (coarsely) encrusted walls, subpellis well differentiated of short, broad cells  $50-110 \times 14-27 \mu\text{m}$ , gradually passing into pileitrama with cells up to  $320 \mu\text{m}$  long and  $12-31 \mu\text{m}$  wide, with brown encrusted walls. Clamp-connections absent.

HABITAT & DISTRIBUTION.—In mossy grasslands, usually on acid soils, not uncommon on the pleistocene sands and old coastal dunes. Known from several localities in the Netherlands. August-October.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Sleen, Sleenerzand, 9 Aug. 1971, *B. de Vries* 1008 (WBS); Westerbork, E. of Eursinge, 27 Sept. 1974, *E. Arnolds* 3182 (WBS); Beilen, along Oude Westerborkseweg near Holtherzand, 13 Oct. 1976, *E. Arnolds* 3645 (holotype); idem, Terhorsterzand, 15 Oct. 1973, *P. Ypelaar* (WBS); Balinge, Balingierzand, 4 Sept. 1968, *B. de Vries* 40 (WBS); prov. Overijssel, Delden, estate 'Twickel', 20 Aug. 1977, *M. E. Noordeloos* 400; Staphorst, 16 Oct. 1976, *M. E. Noordeloos* 225; prov. Utrecht, Baarn, Paleis Soestdijk, 8 Sept. 1979, *M. E. Noordeloos* 1000; prov. Zuid-Holland, Wassenaar, Meyendel, 24 Aug. 1977, *M. E. Noordeloos* 408.

The pigmentation in the pileus varies considerably. In pale forms the encrusting pigments can be entirely lacking from the pileipellis but then it is always distinct in the pileitrama. The intracellular pigments are always manifest.

The yellowish stipe distinguishes *E. xanthocaulon* from all other species in section *Fernandae*. There are, however, some other *Entoloma* species with yellowish stipes with which our species may be confused. *Entoloma verecundum* (Fr.) Noordeloos has absolutely no encrusting pigments in the pileus and larger spores; *E. vinaceum* var. *vinaceum* differs among other things by the spores that are smaller, thin-walled and polyangular in side-view, and also by a different pigmentation, viz. intracellular. It should be noted also that in contrast to *E. xanthocaulon* all these species have clamped basidia. *Entoloma rhombisporum* (Bres.) Horak has sometimes also two types of pigmentation (see p. 452), but its cuboid spores make confusion unlikely.

#### ENTOLOMA section ENDOCHROMONEMA (Largent & Thiers) Noordeloos

*Nolanea* (Fr.) Kumm. sect. *Endochromonema* Largent & Thiers in Northwest Sci. 46: 36. 1972. — *Rhodophyllus* Quél. sect. *Endochromonema* (Largent & Thiers) Romagn. in Bull. mens. Soc. linn. Lyon 43: 331. 1974. — *Entoloma* (Fr.) Kumm. sect. *Endochromonema* (Largent & Thiers) Noordeloos in Persoonia 10: 246. 1979. — Holotype: *Entoloma cetratum* (Fr.) Moser.

CHARACTERISTICS.—With diffusely intracellular, sometimes very pale pigment and in addition pale membranal pigment<sup>4</sup>; spores heterodiametrical with dihedral base, rarely subisodiametrical.

#### Subsection ENDOCHROMONEMA

Pigment usually pale and inconspicuous, diffusely intracellular and in addition sometimes membranal; spores usually simple, never polyangular-nodulose; pileus grey-brown or ochraceous-fulvous, never blackish brown; stipe fibrillose, often striate, never with lemon yellow tinge; smell usually absent or faint, rarely farinaceous, never strongly aromatic like blossom or fruit.

Largent (1974: 1009) characterized this subsection by the presence of clamp-connections, and consequently erected a new subsection for those species in section *Endochromonema* without clamps, viz. subsect. *Efibulatae*. The type species of sect. *Endochromonema*, *E. cetratum* is, in the probably correct sense of European authors, a clampless species. This suggests that Largent's observations were based upon a misinterpretation. In the present work subsect. *Endochromonema* contains both clamped and non-clamped species. I did not study *Nolanea californica* (Murrill) Largent yet, the type species of subsect. *Efibulatae*, therefore I hesitate to put the name *Efibulatae* into the synonymy of subsect. *Endochromonema*.

#### KEY TO THE SPECIES OF SUBSECTION ENDOCHROMONEMA

- 1a. Basidia in majority 2-spored without clamps. . . . . 2
- b. Basidia in majority 4-spored with clamps. . . . . 3
- 2a. Stipe silvery striate; fruitbody slender, mycenoid with conical then slightly expanding pileus, usually with distinct umbo; taste absent or faintly farinaceous; smell absent. . . . . *E. cetratum*, p. 496
- b. Stipe smooth, glabrous; fruitbody thick-set, small; pileus quickly expanding, with or without a slight umbo, sometimes even slightly depressed; taste strongly farinaceous-rancid or fishy, smell weakly to distinctly farinaceous. . . . . *E. farinogustus*, p. 497
- 3a. Spores subisodiametrical,  $(7.4-7.6-9.4(-9.6) \times (6.4-6.9-8.1(-8.6) \mu\text{m})$ ,  $Q = 1.05-1.15-1.25$ . . . . . *E. occultpigmentatum*, p. 501
- b. Spores distinctly heterodiametrical,  $Q$  on the average per collection  $\geq 1.25$ . . . . . 4
- 4a. Spores  $8.3-11 \mu\text{m}$  long,  $Q = 1.1-1.25-1.4$ ; in grasslands. . . . . 5
- b. Spores  $10-12 \mu\text{m}$  long,  $Q = 1.25-1.4-1.5$ ; in forests. . . . . 6
- 5a. Pileus pale orange-brown with grey-brown striation; lamellae pale, then pink without any grey or brown tinge, of normal thickness, not exceeding the pileus. . . . . *E. calthionis*, p. 502
- b. Pileus pale grey-brown with slight ochre tinge, with dark brown striae; lamellae grey, then brown-pink, thickish, broadly ventricose, exceeding the pileus. . . . . *E. ventricosum*, p. 502
- 6a. Stipe downy-pruinose all over, with cylindrical often subcapitate hairs; pileus and stipe concolorous and pale. . . . . *E. lanuginosipes*, p. 499
- b. Stipe glabrous or pruinose at apex only; pileus often dark brown; stipe paler. . . . . 7
- 7a. Pileus bicolorous: limb sepiaecous or date brown, centre paler, yellowish, lamellae ochraceous. . . . . *E. cuneatum*, p. 498
- b. Pileus unicolorous, reddish brown, lamellae flesh pink, then brownish pink. . . . . *E. pallescens*, p. 521

<sup>4</sup> Except in *E. infula*, a species of uncertain taxonomical position, intermediate in this character between sect. *Endochromonema* and sect. *Papillata*.

35. *ENTOLOMA CETRATUM* (Fr.) Moser—Figs. 32a–c

*Agaricus cetratus* Fr., Syst. mycol. 1: 207. 1821. — *Nolanea cetrata* (Fr.) Kumm., Führ. Pilzk.: 95. 1871. — *Rhodophyllus cetratus* (Fr.) Quél., Enchir.: 64. 1886. — *Hyporrhodius cetratus* (Fr.) Schroet. in Cohn, KryptogFl. Schles. 3(1): 613. 1889. — *Entoloma cetratum* (Fr.) Moser in Gams, Kl. KryptogFl. 4. Aufl. 2(b/2): 206. 1978.

SELECTED ICONES AND DESCRIPTIONS.—Arnolds & Noordeloos in Fung. rar. Icon. col. 12: pl. 89 fig. a. 1980. — Konrad in Bull. Soc. mycol. Fr. 45: 51–52. 1929. — Konrad & Maublanc, Icon. sel. Fung., pl. 179 fig. 1. 1930. — J. Lange, Fl. agar. dan., pl. 78F. 1936. — Rea in Trans. Br. mycol. Soc. 12: 217. 1927. — Ricken, Blätterpilze: 297, pl. 74 fig. 1. 1913.

CHARACTERISTICS.—Habit mycenoid; pileus conical, only slightly expanding, ochraceous brown often with reddish tinge; lamellae ochraceous, then pink with brown tinge; stipe concolorous with or paler than pileus, silvery striate; smell weak, rarely slightly farinaceous; basidia 2-spored; clamp-connections absent.

Pileus 9–40(–50) mm broad, conical, then hemispherical to conico-campanulate, finally conico-convex, with broad rounded umbo, rarely truncate, with margin slightly inflexed when young, but straight later on, hygrophanous, pale ochraceous, horn brown or sepia often with reddish tinge, distinctly paler and more yellow towards margin (7.5 YR 4/4, 4/2, 5/4, rarely 3/2 at centre; margin 7.5 YR 7/4; 10 7/4, 8/4), translucently striate up to two thirds of the radius, on drying strongly pallescent to pale yellowish or ochraceous brown (10 YR 7/4, 7/6, 8/3, 8/4), when moist smooth, when dry with aeriferous, silvery lustre. Lamellae L = (15–)20–30, l = (1–)3–5(–7), narrowly adnate to almost free, triangular, then ventricose, up to 7 mm broad, yellowish-ochraceous, then pink, finally brownish pink (10 YR 7/4, then 7.5 YR 8/4, 7/4, 6/4, rarely 5/4), with irregularly crenulate concolorous edge. Stipe (20–)35–85 × (1.5–)2–4(–5) mm, cylindrical, gradually broadening towards base, sometimes flattened and/or twisted, pale yellow to yellowish red (2.5 Y 7/4, 7/6, 6/4, 5/4, or 10 YR 7/3, 7/4), strongly silvery striate lengthwise with aeriferous fibrils, white tomentose at base. Flesh concolorous, inner parts paler. Smell usually absent, rarely very slightly farinaceous. Taste indistinct or rarely with farinaceous or rancid aftertaste.

Spores (9.6–)10.2–13.4(–14) × 7.2–8.0(–9.3)  $\mu\text{m}$ , Q = 1.2–1.45–1.8, L–D = 2.4–4  $\mu\text{m}$ , irregularly (5–)6–7(–8)-angled in side-view, with dihedral base. Basidia 27–45 × 8–12.5  $\mu\text{m}$ , 2-(rarely a few 1- or 4-)spored. Cystidia absent. Hymenophoral trama regular with cylindrical cells 50–140 (–190) × 17–32  $\mu\text{m}$ , with colourless walls. Pileipellis a thin cutis made up of radially arranged, cylindrical, 4–7(–10)  $\mu\text{m}$  wide hyphae, subpellis well developed, composed of inflated cells 52–90(–140) × 16–30  $\mu\text{m}$ , with pale brown walls and in addition a very pale intracellular pigment. Pileitrama regular, composed of cylindrical cells up to 230  $\mu\text{m}$  long and 7.5–21(–27)  $\mu\text{m}$  wide. Clamp-connections absent in all tissues.

HABITAT & DISTRIBUTION.—Terrestrial in humus in or near coniferous forests (*Picea abies*, *Pinus sylvestris*, *P. nigra*) and on heaths (*Ericetum*, *Empetretum*) on pleistocene sandy soils and in coastal dunes; fairly common throughout the year when weather conditions are favourable for the development of carpophores. Common also in entire western and northern Europe in comparable habitats.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Friesland, Isl. of Vlieland, state forest near village, 1 Sept. 1976, *C. Bas* 7146; prov. Drenthe: Gees, state forest, 9 May 1973, *K. Booy*; Spier, state forest, 18 Aug. 1960, *J. J. Barkman* 6734 (WBS); Schoonloo, 19 Oct. 1977, *P. B. Jansen* 77–312; Berkenheuvel, 6 Nov. 1978, *A. E. Jansen*; prov. Overijssel: Rijssen, Friezenberg, 31 Oct. 1957, *R. A. Maas Geesteranus* 12435; prov. Gelderland: Nijmegen, Hatert, 26 Sept. 1954, *C. Bas* 658; Arnhem, state forest N. of town, 10 Nov. 1957, *E. Kits van Waveren*; Nieuw Milligen near Apeldoorn, state forest, 12 Nov. 1954, *C. Bas* 737; Otterloo, *Juniperus* reserve, 24/27 Oct. 1963, *J. J. Barkman* 7664 (WBS); prov. Utrecht: Soestdijk, Paleis Soestdijk, 12 Oct. 1979, *E. Kits van Waveren*; prov. Noord-Holland: Isl. of Texel, de Koog, 27 Oct. 1977, *M. E. Noordeloos* 567; Bergen, 23 Oct. 1965, *J. van Brummelen* 1956;

Overveen, 'Koningshof', 16 Nov. 1976, *M. E. Noordeloos* 269 & 270; Vogelenzang, dunes of Amsterdam Water Supply, 4 Nov. 1976, *M. E. Noordeloos* 246; idem, estate 'Vogelenzang', 11 May 1977, *C. Bas*; prov. Noord-Brabant: Breda, Mastbos, 5 May 1965, *P. B. Jansen* 65-178; Dorst, 2 April 1957, *C. Bas* 1199; Nieuw Ginneken, Goudbergven, 28 Nov. 1964, *P. B. Jansen* 64-204; prov. Limburg, Schinveld, Schinvelderbos, 8 Sept. 1974, *P. B. Jansen* 74-279.

*Entoloma cetratum* is easily recognized in the field by the following combination of characters: pileus and stipe with yellow-ochraceous tinges, silvery striate stipe, and habitat. The smell is usually indistinct; only in three collections examined the smell was more or less farinaceous. The taste is usually weak and mild but in the same three collections mentioned above it was distinctly farinaceous. *Entoloma farinogustus* differs from *E. cetratum* by the dwarfish habit, polished stipe and persistent and strong farinaceous-rancid taste. *Entoloma cuspidifer* resembles *E. cetratum* somewhat in colour of the carpophore but differs by many microscopic characters such as size and shape of spores, pigmentation, and presence of clamp-connections.

*Nolanea testacea* Bres., *Rhodophyllus putus* Romagn., and *R. fulviceps* Romagn. are insufficiently known extralimital species which seem to be closely related to, if not identical with, *E. cetratum*. More information is needed on the variability of *E. cetratum* in entire Europe to take a final decision on the taxonomical rank of these species (see p. 529).

*Entoloma cetratum* has been very well depicted by J. Lange (1936, l.c.), whose plate is representative for the Netherlands' collections; the specimens depicted by Konrad & Maublanc, l.c., are much less characteristic, but nevertheless their description suits my concept of *E. cetratum* very well, just as the description given by Ricken (1913, l.c.).

### 36. ENTOLOMA FARINOGUSTUS Arnolds & Noordeloos—Figs. 34a-b

*Entoloma farinogustus* Arnolds & Noordeloos in *Persoonia* 10: 292. 1979.

ICON AND DESCRIPTION.—Arnolds & Noordeloos in *Fung. rar. Icon. col.* 12: pl. 89 fig. b. 1980.

CHARACTERISTICS.—Carpophores dwarfish; pileus 6-20 mm broad, pale ochraceous brown, flattened; lamellae pale, then pink without any grey or brown; stipe 11-28 × 1.5-3 mm, yellowish, polished; taste farinaceous-rancid; basidia 2-spored; clamp-connections absent.

Pileus 6-20 mm broad, when young bluntly conical or hemispherical, soon flattened with or without weak umbo, rarely slightly depressed, with straight margin, strongly hygrophanous, when moist pale ochraceous brown, at centre slightly darker orange-brown, translucently striate up to centre, on drying strongly pallescent to cream-colour or pinkish orange, smooth. Lamellae L = 20-25, l = 1-5, moderately distant, narrowly adnate, narrowly ventricose, pale, then pink without any grey or brown tinge, with concolorous, entire edge. Stipe 11-28 × (1.5-2-3) mm, cylindrical, slightly broadened at base, yellowish or orange-brown, smooth as if polished, apex sometimes minutely pruinose. Smell weak, slightly farinaceous. Taste strongly farinaceous-rancid or like cod-liver oil.

Spores (8.7-9.0-12(-12.4) × (6.4-6.9-8.2(-9.5)  $\mu\text{m}$ , Q = (1.2-1.25-1.35-1.5(-1.6), L-D = 1.7-2.5-4  $\mu\text{m}$ , irregularly 6-8-angled in side-view, with dihedral base. Basidia 25-40 × 8-12  $\mu\text{m}$ , 2-spored. Cystidia absent. Hymenophoral trama regular with cylindrical or slightly inflated cells (54-64-200(-240) × 7.5-27  $\mu\text{m}$ . Pileipellis a cutis made up of radially arranged cylindrical hyphae 4-12  $\mu\text{m}$  wide, with pale membranous and intracellular pigment; subpellis well developed, composed of short and broad cells 24-85 × 13-28  $\mu\text{m}$ . Pileitrama regular, with broad cylindrical cells up to 200  $\mu\text{m}$  long and 8-25  $\mu\text{m}$  wide. Clamp-connections absent.

**HABITAT & DISTRIBUTION.**—On the ground and on litter of grasses and *Calluna vulgaris* on dry sandy or slightly loamy soils. Known only from the province of Drenthe, Netherlands.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Drenthe, Dwingeloo, Dwingelose hei S. of Smitsveen, 17 Nov. 1976, *E. Arnolds* 3775 (WBS; holotype, L); idem, 15 Oct. 1979, *B. de Vries* (WBS); Schoonloo, Schoonloër Strubben, 11 Oct. 1979, *A. E. Jansen* 582 (WBS).

*Entoloma farinogustus* is closely related to *E. cetratum* by its bisporous basidia, size and shape of spores, and clampless hyphae. It is considered a distinct species on account of dwarfish habit, shape of pileus and polished stipe.

*Eccilia bisporigera* Orton differs by the dark grey-brown pileus, the usually slightly decurrent lamellae, and by the slightly broader and more irregularly shaped spores and abundant clamp-connections, which place it in subgenus *Entoloma*, close to *E. politus*.

### 37. *ENTOLOMA cuneatum* (Bres.) Moser—Figs. 35 a-c

*Nolanea cuneata* Bres., *Fungi trident.* 1: 77, pl. 82 fig. 2. 1887. — *Entoloma cuneatum* (Bres.) Moser in *Gams, Kl. KryptogFl.* 4. Aufl. 2 (b/2): 205. 1978.

**SELECTED DESCRIPTION.**—P. D. Orton in *Trans. Br. mycol. Soc.* 43: 330. 1960 (as *Nolanea cuneata*).

**CHARACTERISTICS.**—Habit mycenoid; pileus dark brown with contrasting yellow papilla; stipe yellowish, paler than pileus, entirely pruinose or at apex only, silvery striate; smell absent; spores (10.2–)10.6–12.4 × 7.4–8.4(–9.0)  $\mu\text{m}$ , Q = 1.25–1.4–1.5(–1.6); basidia 4-spored; clamp-connections present.

Pileus 23–56 mm broad, conical, expanding to conico-convex, then with small papilla, with straight margin, strongly hygrophanous, when moist rather dark (reddish) brown (5 YR 4/3, 4/4; Meth. 5E8), with contrasting yellowish-glassy, apex or papilla, translucently striate up to centre, on drying strongly pallescent to sordid ochraceous grey (10 YR 7/2, 8/4), smooth, lustrous. Lamellae L = up to 32, l = 1–3, moderately distant, almost free, narrowly ventricose, sordid ochraceous pink, finally more brownish pink (one collection: 5 YR 6/6), with concolorous or slightly paler, entire edge. Stipe 25–70 × 2–4 mm, cylindrical, sometimes slightly broadened towards base, yellowish to yellowish brown, paler than pileus, silvery striate lengthwise, at apex or at upper half, sometimes even entirely white pruinose. Flesh concolorous with or paler than surface. Smell and taste indistinct.

Spores (10.2–)10.6–12.4 × 7.4–8.4(–9.0)  $\mu\text{m}$ , Q = 1.25–1.4–1.5(–1.6), L–D = 2.5–4  $\mu\text{m}$ , 5–6–7-angled in side-view, with dihedral base. Basidia 36–50 × 10.8–14.5  $\mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular with cylindrical to strongly inflated cells (160–)220–514(–550) × 10.5–27  $\mu\text{m}$ , with colourless walls. Pileipellis a cutis made up of 2.5–7(–9)  $\mu\text{m}$  wide cylindrical hyphae with pale diffusely intracellular and pale membranal pigment; subpellis not or only weakly developed. Pileitrama regular with cylindrical to inflated cells 270–650 × 12–35  $\mu\text{m}$ , with pale intracellular pigment particularly in upper layers, rarely in addition with pale membranal, never with encrusting pigment. Vascular hyphae numerous in pileitrama. Stipitepellis at apex, rarely also downwards with numerous cylindrical or flexuous hairs with rounded, rarely subcapitate apex, up to 110  $\mu\text{m}$  long and 4.5–6(–8)  $\mu\text{m}$  wide. Clamp-connections abundant in hymenium, rare or even seemingly absent elsewhere.

**HABITAT & DISTRIBUTION.**—In coniferous forests, probably with a preference to rich, calcareous soils. Rare. Known to occur in the Netherlands, Great Britain, and Italy.

**COLLECTIONS EXAMINED.**—NETHERLANDS, IJsselmeerpolders, Oostelijk Flevoland, Abbertsbos, 16 Oct. 1978 and 14 June 1979, *G. J. M. G. Tjallingii-Beukers*.



GREAT BRITAIN: Scotland, Isl. of Rhum, Spruce Woods, north shore of Lock, 30 Aug. 1969, R. W. G. Dennis (K); Wales, Lake Vyrnwy, 8 Sept. 1977, E. Kits van Waveren. — ITALY, Trentino, Dagario, 6 Aug. 1924, G. Bresadola (neotype: design. mihi, M).

DESCRIPTION OF THE NEOTYPE.—Ex Herb. Bres. no. 1198, Dagario, in herbidis, 6 Aug. 1924, G. Bresadola (M).

The neotype consists of one specimen in relatively good state with the following microscopical characters.

Spores  $10.6\text{--}12.4 \times (7.4\text{--})7.9\text{--}8.4\text{--}(9.0) \mu\text{m}$ ,  $Q = 1.3\text{--}1.4\text{--}1.55$ ,  $L\text{--}D = 3.5\text{--}4 \mu\text{m}$ , mostly 6-angled in side-view, with dihedral base. Basidia  $36\text{--}42 \times 12\text{--}14.5 \mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular with long inflated cells  $324\text{--}514\text{--}(540) \times 16\text{--}25 \mu\text{m}$ . Pileipellis a dry cutis made up of  $4.5\text{--}8 \mu\text{m}$  wide cylindrical hyphae with intracellular pigment. Pileitrama regular with strongly inflated cells  $270\text{--}456 \times 10.8\text{--}35 \mu\text{m}$ , some with intracellular pigment particularly in upper layer. Clamp-connections abundant, at least in hymenium.

In the Bresadola herbarium at Stockholm no authentic material of *Entoloma cuneatum* is left. Fortunately I found in the exsiccatae collection of the Bavarian mycologist Killermann (1870–1956) at Munich duplicates from the Bresadola herbarium, among other things the collection cited above, which I selected as a neotype. The microscopical characters fit the recent collections of the same species very well.


*Entoloma cuneatum* is very well characterized by its bicoloured pileus, by which it is easy to distinguish from *E. pallescens*, which in addition is much larger and has a different habitat and distribution. *Entoloma lanuginosipes* has a rather pale, unicoloured pileus and a constantly downy-pruinose stipe. The latter character, which I thought to be very characteristic when creating this new name for *Nolanea crassipes* Velen., is less unique than I thought as the collection from Wales, Lake Vyrnwy, leg E. Kits van Waveren, shows that typical *E. cuneatum* may possess an entirely pruinose stipe as well. For the time being I consider the difference in colour of the pileus sufficiently distinctive to maintain *E. lanuginosipes* as a species in its own right. Further examinations must prove the constancy of that character.

### 38. ENTOLOMA LANUGINOSIPES Noordeloos—Figs. 33a–b

*Nolanea crassipes* Velen., Česká Houby: 627. 1921; non *Entoloma crassipes* Petch, 1924. — *Entoloma lanuginosipes* Noordeloos in Persoonia 10: 248. 1979 [change of name].

CHARACTERISTICS.—Pileus and stipe rather pale yellow(-brown); stipe minutely woolly-downy all over, particularly when dry; hairs on stipe rather long, cylindrical with rounded, attenuate or capitate apex.

Pileus 20–27 mm broad, conical, expanding to plano-convex with or without weak central depression, with straight margin, strongly hygrophanous, when moist pallid honey to yellow(-brown) (Expo C66–C74; 10 YR 7/6–6/4), translucently striate up to one half of the radius, on drying pallescent to pallid isabella-buff or yellow (2.5 Y 8/2, 8/4), strongly silky-shining. Lamellae  $L = \text{about } 30$ ,  $l = 3\text{--}(5)$ , fairly crowded, flesh-coloured pink (7.5 YR 7/4–6/4), with more or less entire, concolorous edge. Stipe 85–90  $\times$  2–3 mm (in type up to  $13 \times 4\text{--}5$  mm, base up to 13 mm), slightly broadened towards base, cylindrical or flattened, pale yellow (10 YR 7/8 to 2.5 Y 8/6), on drying pallescent, silvery striate and minutely downy-pruinose all over, with white



Figs. 32a-c. *Entoloma cetratum*. Habit and spores (32a from Noordeloos 246; 32b-c from Kits van Waveren, 22 Oct. 1979).

Figs. 33a-b. *Entoloma lanuginosipes*. — Habit and spores (33a-b from Noordeloos 1032).

Figs. 34a-b. *Entoloma farinogustus*. — Habit and spores (34a-b from de Vries, 15 Oct. 1979).

Figs. 35a-c. *Entoloma cuneatum*. — Habit and spores (35a-b from Tjallingii, 16 Oct. 1978; 35c from neotype).

toementum at base. Flesh pale, concolorous with surface, inner part paler, subcartilaginous in stipe, brittle in pileus. Smell spontaneously weak, subfarinaceous when cut (in type: 'penetrantly acid'). Taste farinaceous or absent.

Spores 10.2–12.4(–12.7) × 7.4–9.0(–10.2), Q = (1.1)–1.2–1.4–1.5(–1.7), L–D = 2–4.5 μm, 6–7 angled in side-view, with distinctly dihedral base. Basidia 30–58 × 9–17.5 μm, 4-spored. Cystidia absent. Hymenophoral trama regular; cells fusoid, 200–435 × 12–21 μm. Pileipellis a simple cutis made up of 2.5–8(–10) μm wide cylindrical hyphae with pale coloured walls and rather pale diffusely intracellular pigment. Pileitrama regular, composed of cylindrical to inflated cells 280–520 × 12–32 μm, with colourless walls. Stipitepellis a cutis bearing numerous cylindrical hairs 43–163 × 4–10 μm, with rounded, attenuated or capitate apex. Clamp-connections abundant in hymenium, rare elsewhere.

HABITAT & DISTRIBUTION.—In frondose and coniferous forests, rare. Recorded from the Netherlands, German Federal Republic and Czechoslovakia.

COLLECTIONS EXAMINED.—NETHERLANDS, prov. Gelderland, Gietelo, Gietelse bos, 9 Oct. 1966, C. Bas 4804.

GERMAN FEDERAL REPUBLIC, Eifel, Gerolstein, 1 Oct. 1979, M. E. Noordeloos 1032. — CZECHOSLOVAKIA, Bohemia, Hvězda near Prague, April 1920, O. Zvěřina (holotype, PRC).

*Entoloma lanuginosipes* is quite a distinct species with its pale colours and entirely downy-pubescent stipe. Macroscopically it is very similar to *E. cetratum*, which differs by the glabrous stipe, 2-spored basidia and lack of clamp-connections. *Entoloma cuneatum* and *E. pallescens* have a dark brown pileus. See also the discussion under *E. cuneatum*, p. 499.

### 39. *ENTOLOMA OCCULTOPIGMENTATUM* Arnolds & Noordeloos

*Entoloma occultopigmentatum* Arnolds & Noordeloos in *Persoonia* 10: 292. 1979.

ICON AND DESCRIPTION.—Arnolds & Noordeloos in *Fung. rar. Icon. col.* 12: pl. 92 fig. c. 1980.

CHARACTERISTICS.—Robust species, pileus 25–56 mm broad, only weakly umbonate or slightly depressed, rather dark brown; stipe 30–100 × 2.5–8(–9) mm, grey-brown, silvery striate; smell farinaceous; spores isodiametrical; pigment membranous.

Pileus 25–56 mm broad, conico-convex to convex, finally flattened, with or without blunt umbo, rarely slightly depressed at centre, with slightly involute or straight margin, strongly hygrophanous, when moist rather dark brown (10 YR 2/2, 3/2; 5 YR 2/2), paler towards margin and translucently striate at least up to one half of the radius, on drying strongly pallescent to ochraceous-greyish brown (10 YR 6/4, 6/3, 7/2, 7/3), innately fibrillous, lustrous. Lamellae L = 24–60, l = 1–14, moderately distant to rather crowded, broadly adnate to deeply emarginate, rarely almost free, segmentiform to rather broadly ventricose, up to 9 mm broad, pale, sordid grey, then pink with brown tinge, finally flesh-coloured pink (7.5 YR 6/4, 5/4, finally 4/4), with concolorous, entire or slightly irregular edge. Stipe 30–100 × 2.5–8(–9) mm, cylindrical or flattened, straight or flexuous, sometimes gradually broadening towards base, pale to moderately dark grey-brown (10 YR 7/4, 6/4, 5/2, 4/3, 4/2), strongly silvery striate lengthwise, white tomentose at base. Flesh thin and relatively firm in pileus, brittle and fibrous in stipe, concolorous with surface. Smell and taste strongly farinaceous.

Spores (7.4)–7.6–9.4(–9.6) × (6.4)–7.0–8.1(–8.7) μm, Q = 1.0–1.1–1.2(–1.25), L–D = 0–1–1.7 μm, 4–5–6 angled in side-view, base difficult to interpret. Basidia 28.5–41 × 8.5–14 μm, 4-spored. Cystidia absent. Hymenophoral trama regular, composed of rather long, inflated cells up to 500(–700) μm long and up to 35 μm wide. Pileipellis a cutis made up of 3–7(–11) μm wide, cylindrical hyphae with pale membranous and rather diffusely intracellular pigment. Pileitrama

regular, composed of inflated cells up to 350  $\mu\text{m}$  long and up to 27  $\mu\text{m}$  wide, with pale brown walls and, particularly in upper layers pale, diffusely intracellular pigment. Clamp-connections abundant in hymenium, rare in other tissues.

**HABITAT & DISTRIBUTION.**—In grasslands (*Scirpetum sylvatici*); once found on poorly grazed dike); rare. So far known from three different localities in the Netherlands.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Drenthe, Vries, along Taarlose Diep, E. of Taarlo, 15 Sept. 1976, *E. Arnolds 3588* (holotype); prov. Noord-Holland, Schoorl, old sea dike near 'Schotsbrugsluis', 21 Oct. 1979, *M. E. Noordeloos 1074*; IJsselmeerpolders, Oostelijk Flevoland, Roggebotszand, 17 Oct. 1976, *G. Boezwinkel & H. J. W. Langevoord-Dull*.

*Entoloma occultopigmentatum* is a fairly robust grassland species. The dark brown pileus, strong farinaceous smell and isodiametrical spores may cause confusion with *E. sericeum*, but the type of pigmentation is quite different. Because of the latter character *E. occultopigmentatum* is placed in section *Endochromonema*. It has some resemblance with *E. pallescens*, particularly in habit, but the isodiametrical spores prevent confusion.

#### 40. ENTOLOMA CALTHIONIS Arnolds & Noordeloos

*Entoloma calthionis* Arnolds & Noordeloos in *Persoonia* 10: 287. 1979.

ICON AND DESCRIPTION.—Arnolds & Noordeloos in *Fung. rar. Icon. col.* 12: pl. 89 fig. c. 1980.

**CHARACTERISTICS.**—Pileus (10–)12–28 mm broad, obtusely conical to convex or flattened, with or without small umbo, with straight margin, hygrophanous, when moist pale brown or with orange tinge, with darker brown striation up to centre, smooth but innately radially fibrillous, lustrous; lamellae  $L = 22\text{--}27$ ,  $l = 1\text{--}3\text{--}(5)$ , crowded, free, ventricose, pink without any brown or grey tinge; stipe 22–47  $\times$  1.5–3.5 mm, cylindrical, slightly broadened towards base, greyish to brownish yellow, silvery striate. Flesh concolorous, brittle; smell farinaceous; spores (8.5–)9–10.6(–11.5)  $\times$  7.2–8.5(–9.0)  $\mu\text{m}$ ,  $Q = 1.1\text{--}1.25\text{--}1.4$ , 5–6-angled in side-view, with dihedral base; basidia 4-spored; cystidia absent; hymenophoral trama regular with cells 200–350  $\times$  11–15  $\mu\text{m}$ ; pileipellis a dry cutis made up of 2–5  $\mu\text{m}$  wide hyphae with pale intracellular pigment. Pileitrama regular; hyphae 7–17.5  $\mu\text{m}$  wide. Clamp-connections abundant in hymenium.

**HABITAT & DISTRIBUTION.**—In *Calthion palustris* on moist soil. Known only from the type locality in the province of Drenthe in the Netherlands.

*Entoloma calthionis* resembles *E. cetratum* very much in colour, but differs by the habitat, slightly smaller and less elongate spores, the 4-spored basidia and the clamp-connections in the hymenium. *Entoloma ventricosum* differs by the grey tinges in all parts of the carpophore, veined, thickish lamellae and different habit.

#### 41. ENTOLOMA VENTRICOSUM Arnolds & Noordeloos

*Entoloma ventricosum* Arnolds & Noordeloos in *Persoonia* 10: 298. 1979.

ICON AND DESCRIPTION.—Arnolds & Noordeloos in *Fung. rar. Icon. col.* 12: pl. 93 fig. c. 1980.

**CHARACTERISTICS.**—Pileus 13–24 mm broad, bluntly conical to convex, with slightly involute margin, hygrophanous, when moist grey-brown, on drying strongly pallescent to sordid

ochraceous-grey; lamellae  $L=15-18$ ,  $l=1-3$ , narrowly adnate, rather strongly ventricose, exceeding the pileus, thickish, veined, grey, then brown-pink with flesh-coloured edge; stipe  $30-48 \times 2-3.7$  mm, pale ochraceous grey-brown to dark grey-brown, smooth but very minutely longitudinally striate, white tomentose at base; flesh thin-membranaceous, concolorous with surface; smell and taste absent; spores  $(8.3-9.1-11(-11.5) \times 7-8.3 \mu\text{m})$ .  $Q=1.15-1.25-1.4$ ; basidia 4-spored; cystidia absent; hymenophoral trama regular with cells up to  $650 \mu\text{m}$  long and up to  $30 \mu\text{m}$  wide; pileipellis a thin cutis made up of  $4-8 \mu\text{m}$  wide, cylindrical hyphae with pale diffusely intracellular pigment; pileitrarna regular with cells up to  $500 \mu\text{m}$  long and up to  $32 \mu\text{m}$  wide. Clamp-connections abundant in hymenium.

**HABITAT & DISTRIBUTION.**—In poor grassland with much moss, grazed by sheep. Known only from the type locality in the province of Drenthe in the Netherlands.

**Entoloma** subsection **Infularia** (Romagn.) ex Noordeloos emend.

*Rhodophyllus* sect. *Infularii* Romagn. in Bull. Soc. mycol. Fr. 53: 332. 1937 (nom. nud.).

Lamellis albidis demum roseis; stipite polito; sporis 5-6-angulatis cum structura nominatur 'dihedral base'. — Typus: *Entoloma infula* (Fr.) Noordeloos.

Lamellae white, then salmon pink without any grey or brown tinge; spores heterodiametrical, 5-6-angled in side-view, with distinctly dihedral base. — Type: *Entoloma infula* (Fr.) Noordeloos.

The original concept of sect. *Infularia* is emended by restricting it to the species in sect. *Endochromonema* with lamellae first white, then pink without any brown or grey tinge, with a polished stipe, and with heterodiametrical spores with a dihedral base.

KEY TO THE SPECIES OF SUBSECTION INFULARIA

- |  |                                |
|--|--------------------------------|
| 1a. Pigment distinctly membranal-encrusting in hyphae of pileipellis and (particularly) of pileitrarna, sometimes also diffusely intracellular . . . . . | <i>E. infula</i> , p. 503      |
| b. Pigment exclusively intracellular . . . . .   | 2                              |
| 2a. Stipe yellow; spores $9.3-11(-11.5) \times 7-8.1 \mu\text{m}$ . . . . .  | <i>E. verecundum</i> , p. 507  |
| b. Stipe brown; spores $7.5-9.3 \times 5.4-7 \mu\text{m}$ . . . . .  | 3                              |
| 3a. Stipe grey-brown to blackish brown; smell absent. . . . .  | <i>E. solstitiale</i> , p. 505 |
| b. Stipe yellowish brown to reddish brown; smell strong, like chlorine . . . . .   | <i>E. chlorinosum</i> , p. 507 |

42. **Entoloma infula** (Fr.) Noordeloos, *comb. nov.*—Figs. 37a-b

*Agaricus infula* Fr., Spicilegium: 8. 1836. — *Nolanea infula* (Fr.) Gill., Hymen. Fr.: 421. 1876. — *Rhodophyllus infula* (Fr.) Quél., Enchir.: 64. 1886.

SELECTED ICONS.—Arnolds & Noordeloos in Fung. rar. Icon. col. 12: pl. 90 figs. a, d. 1980. — Ricken, Blätterpilze, pl. 74 fig. 9. 1913.

**CHARACTERISTICS.**—Pileus and stipe moderately dark ochraceous brown or greyish brown; stipe smooth; minutely encrusting pigment, sometimes combined with diffusely intracellular pigment.

Pileus  $10-43$  mm broad, at first conical, then conico-campanulate, later expanding to convex, finally plano-convex or more or less concave with strongly undulating margin, usually with

pronounced papilla, hygrophanous, when moist ochraceous- or greyish brown to sepia (Meth. 6D6, 6C5 to 6C6 or 6E6), translucently striate at least up to one half of the radius, on drying pallescent to ochre-grey (Meth. 5C5/C4), glabrous, smooth or minutely radially wrinkled ('chagriné'); shining. Lamellae  $L=18-29$ ,  $l=(1-3)(-7)$ , crowded, narrowly adnate to free, ventricose, white, then salmon pink without any trace of grey or brown, with entire, concolorous edge. Stipe  $17-78 \times 1.5-3$  mm, cylindrical, sometimes slightly broadened at base, concolorous with or slightly paler than pileus (Meth. 6C3, 6C5, 6D5, 6D4/C4), pruinose at apex, glabrous, polished or innately fibrillose downwards, not striate, white tomentose at base. Flesh membranaceous, usually rather firm, concolorous with surface or inner part paler. Smell absent or faint. Taste inconspicuous, mild.

Spores  $7-9.3(-10) \times 5.8-7 \mu\text{m}$ ,  $Q=1.15-1.3-1.4(-1.5)$ ,  $L-D=(0.6-)1.0-1.7-2.4 \mu\text{m}$ , (4-)5-6(-7)-angled in side-view, with dihedral base. Basidia  $26-39 \times 7.5-14 \mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular, cells cylindrical or slightly inflated,  $220-520 \times 5.6-20(-32) \mu\text{m}$ , with hyaline, colourless walls. Pileipellis a cutis made up of  $(2.7-)4.0-11 \mu\text{m}$  wide cylindrical hyphae with (minutely) encrusted walls and sometimes in addition diffusely intracellular pigment, with some scattered inflated terminal cells up to  $17 \mu\text{m}$  wide. Subpellis sometimes more or less distinct, composed of cylindrical to inflated cells  $32-89 \times 12-17 \mu\text{m}$ , gradually passing into trama. Pileitrama regular with up to  $250 \mu\text{m}$  long and  $7-19(-23) \mu\text{m}$  wide cells with encrusted walls particularly in upper zones. Clamp-connections frequent in hymenium, rare elsewhere.

HABITAT & DISTRIBUTION.—In grasslands; rare, known to occur in Sweden, the Netherlands and France; probably widespread (lit.).

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Vries, Taarlose Diep, 13 Oct. 1975, *E. Arnolds 3443* (WBS); Westerbork, CRM-reserve 'de Reitsma', 13 Oct. 1976, *E. Arnolds* (WBS); prov. Limburg, Canne, Cannerberg, 30 Oct. 1978, *E. Arnolds 4141* (WBS).

SWEDEN, Småland, Femsjö, S. of Källebo, 22 Sept. 1949, *S. Lundell & J. Stordahl* (Fungi exsicc. suec. 2013, PC; neotype, design. mihi). —FRANCE, dept. Oise, Chantilly, 28 Aug. 1951, *H. Romagnesi 51.210* (Herb. Romagn., PC).

OBSERVATIONS ON THE NEOTYPE.—Spores  $(7.6-)8.1-9.3(-10) \mu\text{m}$ ,  $Q=1.3-1.35-1.5$ ,  $L-D=1.7-2-2.7(-3) \mu\text{m}$ , 5-6-angled in side-view, with dihedral base. Basidia  $25-36 \times 8-11.5 \mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama with cylindrical to inflated cells up to at least  $250 \mu\text{m}$  long and  $7-15(-19) \mu\text{m}$  wide. Pileipellis a cutis made up of  $3.5-6(-8) \mu\text{m}$  wide cylindrical hyphae with uniformly coloured, sometimes minutely encrusted walls. Subpellis more or less differentiated, composed of short cells  $27-50 \times 8-17 \mu\text{m}$  with brown encrusted walls. Pileitrama regular, composed of cells much longer and more cylindrical,  $85-320 \times 7-21 \mu\text{m}$ . Clamp-connections frequent in hymenium, rare in other tissues.

The macroscopical characters as indicated by Lundell agree in a satisfactory way with Fries's diagnosis and with my own observations. As the collection of Lundell & Stordahl was made at what may be considered the type locality, I selected this collection for neotype. Another collection in the exsiccatae series of Lundell & Nannfeldt, viz. no. 1133, studied by me, appeared to be quite close to *E. solstitialis* in my concept, as it showed intracellular pigment and similar spores. Unfortunately, detailed macroscopical characters on this collection are lacking. It is said to resemble the picture of *J. Lange* (1936, pl. 79G).

*Leptonia pilatii* Svrček is very likely a synonym of *Entoloma infula*. The macroscopical characters are similar. The dimensions of the spores as given by Svrček are slightly too great, probably because of including the measurements of the apiculus. I measured  $7.0-9.3 \times 5.8-7.2 \mu\text{m}$  (see Fig. 37a). Svrček placed the species in *Leptonia* on account of the (classical Friesian) characters as involute margin of pileus and rigid, polished stipe. I consider these characters subordinate to characters such as pigmentation, size and shape of tramal elements, and clamped

basidia. Concerning the latter characters *L. pilatii* is a typical *Nolanea*. The colour of pileus and stipe of *L. pilatii* is said to be 'rubro-fusco', which may range within the variation of *E. infula*.

*Entoloma infula* appears to be rather rare in the Netherlands, and probably outside this area as well. It seems to be restricted to a type of grassland of which, due to artificial fertilization, little is left. It may be confused with other species in subgenus *Nolanea* with pale lamellae, such as *E. minutum* and members of sect. *Fernandae*. The former differs by the pale carpophores and different size and shape of the spores. The latter have, among other things, quite an other type of pileipellis and pigmentation, though the spores are very similar.

#### 43. *Entoloma solstitiale* (Fr.) Noordeloos,

*comb. nov.*—Figs. 38a–b

*Agaricus solstitialis* Fr., Epicr.: 152, 1838. — *Leptonia solstitialis* (Fr.) Gill., Hymen. Fr.: 416, 1876. — *Rhodophyllus solstitialis* (Fr.) Quél., Enchir.: 60, 1886. — *Nolanea solstitialis* (Fr.) P. D. Orton in Trans. Br. mycol. Soc. 43: 179, 1960.

EXCLUDED NAME.—*Leptonia solstitialis* sensu Ricken 1913 (? = *E. rhombisporum*).

SELECTED ICON.—Arnolds & Noordeloos in Fung. rar. Icon. col. 12: pl. 90 fig. c. 1980.

CHARACTERISTICS.—Pileus acutely conical, only slightly expanding, brown; lamellae white, then pink; stipe rigid, polished, with grey tinge; pigment intracellular.

Pileus 10–24 mm broad, (acutely) conical, only slightly expanding, with small, often abrupt papilla, with margin first involute but later more straight, hygrophanous, when moist moderately dark brown, sometimes more ochraceous tinged (Meth. 6D6 to 6E6; 10 YR 5/4, 5/6, at centre towards 10 YR 3/4), slightly paler at margin, translucently striate up to three fourths of the radius, on drying pallescent to greyish brown or greyish yellow (Meth. 6D5; 10 YR 7/3 or 2.5 Y 7/2[8/2]), strongly radially fibrillose-canescens with some aeriferous fibrils, shining. Lamellae L = 30–35, l = (1)–3–9, fairly crowded, deeply emarginate to almost free, ventricose, first pale, then salmon pink without any grey or brown (Meth. 6B5, 7B5; 7.5 YR 8/4, 7/4), with entire, concolorous edge. Stipe 27–60 × 1–2 mm, slender, cylindrical, sometimes slightly broadened at base, solid, then narrowly fistulose, greyish or blackish brown — distinctly more grey than pileus — (Meth. 6E4; 10 YR 2/2, 3/2), paler at apex, glabrous, polished or with some innate fibrils, not striate, white tomentose at base. Flesh firm-subcartilagineous, concolorous with surface, in inner parts paler. Smell and taste mild, indistinct.

Spores 7.5–9.3 × (5.2)–5.4–7 μm, Q = (1.0)–1.2–1.35–1.6, L–D = (0)–1–2–3.5 μm, (4)–5–6 angled in side-view, with very distinctly dihedral base. Basidia 26–34(–35) × (6.2)–6.6–12 μm, 4-spored. Cystidia absent. Hymenophoral trama regular with cylindrical to inflated cells 90–220 × 7–15 μm. Pileipellis a cutis made up of cylindrical to slightly inflated hyphae 3–9.5 μm wide, with hardly any pigment. Subpellis only weakly developed, with cylindrical to inflated cells 27–72 × 6.5–15(–19) μm, with brown intracellular pigment. Pileitrama regular with cylindrical or more or less inflated hyphae composed of relatively short cells 75–190(–230) × 6–17 μm, with intracellular pigment in upper layers. Clamp-connections only seen in hymenium.

HABITAT & DISTRIBUTION.—In grasslands; rare. Recorded from the Netherlands, Sweden, Great Britain, and France.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Overijssel, Diepenveen, Hengforder waarden, 12 Nov. 1977, H. Piepenbroek & G. Piepenbroek-Grooters 1068; prov. Gelderland, Voorst, on dike 500 m S. of Wilp, 6 Oct. 1975, E. Arnolds 3425 (WBS).

FRANCE, dept. Oise, Coye-la-Forêt, 27 Aug. 1945, H. Romagnesi 208 (Herb. Romagn., PC).




Fig. 36-b. *Entoloma chlorinosum*. — Habit and spores (36a-b from Noordeloos 815).

Figs. 37a-b. *Entoloma infula*. — Spores (37a from type of *Leptonia pilatii*; 37b from neotype).

Figs. 38a-b. *Entoloma solstitiale*. — Habit and spores (38a-b from Piepenbroek 1068).

Figs. 39a-b. *Entoloma verecundum*. — Habit and spores (39a-b from Noordeloos 844).

In *Entoloma solstitiale* the spore shape is slightly more variable than in *E. infula*. A certain amount of spores (which remains small, however, and may lack in some carpophores) appears to be more or less rhomboid or 4-angled in side-view, whereas the majority of the spores is 5-6-angled. These spores most probably led Kühn. & Romagn. (1953: 189) and Orton & al. (1960: 127) to identify the species with Ricken's plate of *Leptonia solstitialis* (1913, pl. 73 fig. 3). This plate, however, does not agree in a satisfactory way with Fries's diagnosis, particularly by the colours of pileus and stipe. Ricken's fungus is far more likely to be identical with *E. rhombisporum* (see p. 450).

The fungus as described above may be the same as the original *Agaricus solstitialis* Fr. But I never found the pileus to be slightly depressed as mentioned by Fries (e.g. in Kalchbrenner 1873, pl. 12 fig. 3). According to Fries there is always a papilla, even in the depression of the pileus. In other macroscopical characters my fungus is quite similar to that of Fries. For the same reasons as described above for *Leptonia pilatii*, Fries and many of his followers placed *A. solstitialis* in the (sub-) genus *Leptonia*.



*Entoloma solstitiale* is distinguished from *E. chlorinosum* by the different colours of pileus and stipe and the lack of a smell. *Entoloma infula* differs among other things by the encrusting pigments.

#### 44. *ENTOLOMA CHLORINOSUM* Arnolds & Noordeloos—Figs. 36a–b

*Entoloma chlorinosum* Arnolds & Noordeloos in *Persoonia* 10: 287. 1979.

ICON AND DESCRIPTION.—Arnolds & Noordeloos in *Fung. rar. Icon. col.* 12: pl. 90 fig. b. 1980.

CHARACTERISTICS.—Pileus small, acutely conical or hemispherical, with papilla, yellowish brown or reddish brown; lamellae white, then pink; stipe 28–40 × 0.6–1.3 mm, concolorous with pileus, rigid, polished; smell strong, like chlorine; pigment intracellular.

Pileus 7–14 mm broad, acutely conical or hemispherical, with small, abrupt papilla, expanding to conico-campanulate, finally convex with small conical papilla, with margin straight but reflexed with age, hygrophanous, when moist (dark) yellowish brown or reddish brown (Meth. 5C7; 10 YR 3/3, 4/3, 4/4), slightly paler towards margin, translucently striate up to three fourths of the radius, on drying pallescent to pale yellowish brown (Meth. 5B5; 10 YR 5/4, 6/4), shining, sometimes with aeriferous fibrils, slightly fluffy particularly at centre. Lamellae L = 12–24, l = 0–1(–3), moderately distant, almost free, thin, broadly ventricose, white, then salmon (7.5 YR 7/4, becoming 6/4 with age), with entire concolorous edge. Stipe 28–40 × 0.6–1.3 mm, cylindrical, pale to moderately dark yellowish brown (Meth. 5B5, 5D5; 10 YR 4/3, 4/4), glabrous, smooth as if polished, white tomentose at base. Flesh thin, membranaceous-brittle in pileus, firm, rigid in stipe, concolorous with surface. Smell strong, like chlorine. Taste mild or somewhat soapy, not farinaceous.

Spores (6.8–)7.4–9.0 × (5.1–)5.2–6.8 μm, Q = 1.2–1.3–1.5, L–D = 1–2.5–3.6 μm, 5–6(–7)-angled in side-view, with distinctly dihedral base. Basidia 24–35 × 7.5–10 μm, 4-spored. Cystidia absent. Hymenophoral trama regular, with cylindrical to inflated cells 100–200 × (5–)8–15 μm. Pileipellis a simple cutis made up of repent 2–8 μm wide, cylindrical hyphae. Pileitrama regular, hyphae cylindrical with cells up to 260 μm long and up to 21 μm wide, with abundant intracellular pigment in upper layer. Clamp-connections abundant in hymenium, rare in other tissues.

HABITAT & distribution.—In grasslands; rare. The Netherlands.

COLLECTIONS EXAMINED.—Netherlands: prov. Groningen, Vlagtwedde, reserve 'Metbroek', 29 Oct. 1978. *M. E. Noordeloos* 815; prov. Drenthe, Anloo, Burgwallen, N. of Nieuwlanden, 24 Sept. 1974. *E. Arnolds* 3168 (holotype, L; isotype, WBS).

*Entoloma chlorinosum* is closely related to *E. solstitiale* from which it differs by the colour of the stipe and the strong smell. It can easily be distinguished from other *Entoloma* species with similar smell (viz. *E. nidorosum*, '*Leptonia*' *pernitrosa* Orton, '*Rhodophyllus*' *nitriolens* Kühn.) by the nolaneoid habit, the size and the shape of the spores and of the tramal elements. Those species all belong to the subgenus *Entoloma* and will be treated in a future paper (Noordeloos, 1980a; in prep.).

#### 45. *Entoloma verecundum* (Fr.) Noordeloos, *comb. nov.*—Figs. 39a–b

*Agaricus rubellus* Scop. subsp. *verecundus* Fr., *Spicilegium*: 6. 1836. — *Agaricus verecundus* (Fr.) Fr., *Epicr.*: 158. 1838. — *Nolanea verecunda* (Fr.) Gill., *Hymen. Fr.*: 422. 1876. — *Rhodophyllus verecundus* (Fr.) Quel., *Enchir.*: 64. 1886.

**CHARACTERISTICS.**—Pileus conical to conico-campanulate, pinkish brown, translucently striate up to centre; lamellae white, then pink; stipe yellowish, rigid, polished; pigment intracellular.

Pileus 17–22 mm broad, conical, then expanding to conico-campanulate, with straight margin, hygrophanous, pinkish brown (7.5 YR 5/4 at centre, towards margin 7.5 YR 6/4, 7/4), translucently striate up to centre, on drying strongly pallescent to ochraceous-pink, smooth, glabrous, shining. Lamellae  $L=25$ ,  $l=(1-3)(-5)$ , almost free, ventricose, white, then pink (7.5 YR 7/2), with entire concolorous edge. Stipe  $25-30 \times 1.5-2$  mm, cylindrical, straight, yellow (2.5 Y 7/4), smooth, polished, rigid. Flesh thin-membranaceous, rigid. Smell absent. Taste not tried.

Spores  $9.3-11(-11.5) \times 7.0-8.1 \mu\text{m}$ ,  $Q=1.1-1.3-1.5$ ,  $L-D=1.2-3.5 \mu\text{m}$ , 5–6–7-angled in side-view, with dihedral base. Basidia  $28-35 \times 11-14 \mu\text{m}$ , 4-spored. Cystidia absent. Hymenophoral trama regular with slightly inflated cells at least up to  $240 \mu\text{m}$  long and  $7-18 \mu\text{m}$  wide. Pileipellis a cutis made up of repent, cylindrical,  $2.5-11 \mu\text{m}$  wide hyphae with some clavate terminal cells up to  $16 \mu\text{m}$  wide, with pale, diffusely intracellular pigment, gradually passing into trama. Pileitrama regular with cells similar to those in hymenophoral trama. Clamp-connections abundant in hymenium, not seen in other tissues.

**HABITAT & DISTRIBUTION.**—In extensively grazed dune meadow with *Salix reprens*; known from one locality in the Netherlands.

**COLLECTION EXAMINED.**—NETHERLANDS, prov. Friesland, Isl. of Terschelling, Oosterend, Dazenplak, 3 Nov. 1978, *M. E. Noordeloos 844*.

*Entoloma verecundum* with its remarkably pinkish brown pileus and yellowish stipe has disappeared from mycological literature since the beginning of this century. The plates of Fries (1874, pl. 99 fig. 5) and Cooke (1886, pl. 340 fig. a) are representative for my collection and made me identify it as *E. verecundum*. Konrad & Maublanc (1930: pl. 179) and Ricken (1913: 297) consider *E. verecundum* a synonym of *Rhodophyllus icterinus* and *Nolanea pleopodia* respectively, two taxa considered conspecific by me (see p. 516). *Entoloma verecundum* is distinguished from *E. icterinum* by the differently coloured pileus and slightly different spores. Other species of *Entoloma* with a yellow stipe, viz. *E. vinaceum* var. *vinaceum* and *E. xanthocaulon*, differ by smaller spores and an other type of pigmentation respectively.

#### **Entoloma** subsection **Tristia** Noordeloos, *subsect. nov.*

Pileo atrobrunneo; stipite polito; sporis irregulariter noduloso-angulatis; fibulis nullis. — Typus: *Entoloma triste* (Velen.) Noordeloos.

Pileus blackish brown; stipe smooth and glabrous; spores irregularly angular-gibbose in side-view; clamp-connections present. — Holotype: *Entoloma triste* (Velen.) Noordeloos.

#### KEY TO THE SPECIES OF SUBSECTION TRISTIA

- 1a. Spores  $10.4-12(-14) \times (7.0-7.6-8.7(-9.0) \mu\text{m}$ . . . . . *E. triste*, p. 509  
 b. Spores  $(7.7-7.9-10.8(-12) \times (5.7-6.0-6.8(-7.5) \mu\text{m}$ . . . . . *E. undulatosporum*, p. 509

46. *ENTOLOMA TRISTE* (Velen.) Noordeloos

*Nolanea tristis* Velen., České Houby: 630. 1921. — *Entoloma triste* (Velen.) Noordeloos in Persoonia 10: 254. 1979.<sup>2</sup>

**CHARACTERISTICS.**—Pileus rather dark blackish brown; stipe paler; spores on the average 7.6–8.7  $\mu\text{m}$  wide.

Pileus 10–20 mm broad, conical, finally expanding to convex, with umbo, hygrophanous, when moist blackish brown, not striate or only very obscurely striate at margin only, on drying pallescent, radially fibrillous, shining. Lamellae distant, greyish-ochraceous (with pink tinge?). Stipe  $20 \times 1.5$  mm, horn brown at apex, darker towards base, paler than pileus, white tomentose at base. Smell absent.

Spores  $10.4\text{--}12.0\text{--}(14.0) \times (7.0\text{--})7.6\text{--}8.7\text{--}(9.0)$   $\mu\text{m}$ ,  $Q = 1.2\text{--}1.35\text{--}1.5\text{--}(1.6)$ ,  $L\text{--}D = 1.7\text{--}2.6\text{--}3.4$   $\mu\text{m}$ , 6–8-angled in side view, with dihedral base. Basidia  $27\text{--}38 \times 12.5\text{--}14.5$   $\mu\text{m}$ , 4-spored. Hymenophoral trama regular; cells cylindrical to strongly inflated, up to 250  $\mu\text{m}$  long and 10–23  $\mu\text{m}$  wide. Pileipellis a cutis made up of 4–10  $\mu\text{m}$  wide cylindrical hyphae, sometimes with clavate terminal cells up to 15  $\mu\text{m}$  wide, with abundant intracellular pigment. Clamp-connections frequent in hymenium.

**HABITAT & DISTRIBUTION.**—The Netherlands' collection was made in a dried up fen, growing on dead *Sphagnum* in *Calluna* heath on acid, sandy soil. The type collection came from a grassy clearing in a coniferous forest. Rare; known only from the type locality in Czechoslovakia and one locality in the Netherlands.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Noord-Brabant, Nieuw Ginneken, Strijbeekse heide, 27 July 1960, P. B. Jansen.

CZECHOSLOVAKIA, Bohemia, Krč, May 1920, J. Velenovský (holotype, PRC).

The description given above is based on the Netherlands' collection. Unfortunately some important macroscopical notes are lacking, such as the colour of the lamellae and the aspect of the stipe-surface. The rest of the macroscopy and the microscopical characters lead me to consider this collection, made by the eminent amateur mycologist P. B. Jansen of Breda, conspecific with *Entoloma triste* (Velen.) Noordeloos.

The broad spores differentiate this species from *E. undulatosporum*. Both species are to be placed in sect. *Endochromonema* on account of their pigmentation pattern. The characteristic shape of the spores and the fairly dark colour of the pileus made me place them in a new subsection.

47. *ENTOLOMA UNDULATOSPORUM* Arnolds & Noordeloos

*Entoloma undulatosporum* Arnolds & Noordeloos in Persoonia 10: 295–296, figs. 27–29. 1979.

**CHARACTERISTICS.**—Pileus 15–23 mm broad, convex, then flattened, not papillate, rather dark blackish brown, striate at margin only, on drying slightly pallescent, minutely radially rugulose. Lamellae moderately distant, grey-brown, then flesh-coloured grey. Stipe up to  $27 \times 1.8\text{--}2.8$  mm,

<sup>2</sup> In combination with *Entoloma* the correct spelling of the epithet is 'triste', and not 'tristis'.

paler and more brown than pileus. Smell and taste farinaceous. Spores (7.7–)7.9–10.8(–12.0) × (5.7–)6.0–6.8(–7.5) μm, irregularly 6–9-angled-nodulose in side-view. Pileipellis a cutis made up of 3.8–15 μm wide cylindrical hyphae with abundant intracellular pigment. Clamp-connections frequent in hymenium.

**HABITAT & DISTRIBUTION.**—In poor non-fertilized meadow (*Lolio-Cynosuretum*) on dry, humus-rich sandy soil. Known only from type locality.

The narrow, slightly more irregularly angular spores and perhaps also the farinaceous smell and the habitat distinguish *E. undulatosporum* from *E. triste*.

#### **Entoloma** subsection **Cheilocystidiata** Noordeloos, *subsect. nov.*

Cheilocystidia adsunt, variabilibus; pigmentis intracellulosis. — Typus: *Entoloma velenovskyi* Noordeloos.

Cheilocystidia present, variable in shape; pigment intracellular. — Type: *Entoloma velenovskyi* Noordeloos.

#### KEY TO THE SPECIES OF SUBSECTION CHEILOCYSTIDIATA

- 1a. Cheilo- and pleurocystidia present, globuliform or sphaeropedunculate; clamp-connections absent. *E. globulifer*, p. 513
- b. Cheilocystidia different; pleurocystidia absent; clampconnections present. . . . . 2
- 2a. Cheilocystidia in majority distinctly lecithiform with abrupt capitulum; lamellae very dark brown. *E. inutile*, p. 512
- b. Cheilocystidia different; if (sub-)capitate, then lamellae never very dark brown. . . . . 3
- 3a. Cheilocystidia cylindrical, sometimes subcapitate, (8.1–)15–38(–42) μm long *E. cryptocystidiatum*, p. 511
- b. Cheilocystidia fusiform-lageniform, usually much longer than 40 μm . . . . . 4
- 4a. Spores (10–)10.4–12.0 μm long; cheilocystidia 42–90(–120) × 7–14(–20) μm *E. velenovskyi* var. *velenovskyi*, p. 510
- b. Spores (10.1–)11–15.9(–16.3) μm long; cheilocystidia (50–)70–149 × (6.2–)7.4–15.9 μm *E. velenovskyi* var. *longicystidiatum*, p. 511

#### 48. **ENTOLOMA VELENOVSKYI** Noordeloos var. **VELENOVSKYI** Figs. 41a–c

*Nolanea conica* Velen., České Houby: 623. 1921 (non *Entoloma conicum* (Peck 1872) Hesler 1967). — *Entoloma velenovskyi* Noordeloos in *Persoonia* 10: 258–259. 1979 (change of name).

**CHARACTERISTICS.**—Pileus conical, only slightly expanding, with acute papilla, moderately dark grey-brown; stipe concolorous or slightly paler, smooth; spores 10–12.5 μm long; cheilocystidia lageniform, 42–90(–120) μm long.

Pileus 25 mm broad, conico-convex with acute papilla, with margin slightly involute, then straight, hygrophanous, when moist clay buff (7.5 YR 6/2, 5/2), translucently striate, on drying pallescent, silvery fibrillous, shining. Lamellae moderately distant, almost free, ventricose, clay pink (5 YR 7/3, 6/3). Stipe 60 × 3 mm, cylindrical, straight, buff, smooth, glabrous, white tomentose at base. Smell absent.

Spores (10–)10.4–12.0(–12.7) × 7.6–8.7(–9.3) μm, Q = 1.2–1.3–1.4(–1.5), L–D = 1.7–2.4–4.0 μm, 5–6–7-angled in side-view, probably with dihedral base. Basidia 34–42 × 10.5–12.7 μm, 4-

(rarely 2-)spored. Cheilocystidia  $42-93(-120) \times 7-14 \mu\text{m}$ , fusiform-lageniform, scattered among basidia. Hymenophoral trama regular, with elements  $230-300 \times 11-27 \mu\text{m}$ . Pileipellis a cutis made up of radially arranged,  $4-10 \mu\text{m}$  wide, cylindrical hyphae with diffusely intracellular pigment. Pileitrama regular, composed of cylindrical to slightly inflated cells  $210-320 \times 12-32 \mu\text{m}$ . Clamp-connections frequent in hymenium, rare in trama and covering layers.

**HABITAT & DISTRIBUTION.**—In grasslands, rare. Known from the type locality and from one locality in the Netherlands.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Noord-Brabant, Drunen, along Drongelen canal, 20 Aug. 1972, P. B. Jansen 72-226.

CZECHOSLOVAKIA, Bohemia, Prague, Mnichovice, July 1919, J. Velenovský (lectotype, PRC).

*Entoloma velenovskyi* is fairly well characterized by the long attenuate cheilocystidia and the intracellular pigment.

#### 49. ENTOLOMA VELENOVSKYI Noordeloos var. LONGICYSTIDIATUM Arnolds & Noordeloos

*Entoloma velenovskyi* Noordeloos var. *longicystidium* Arnolds & Noordeloos in *Persoonia* 10: 296-298, figs. 30-33, 1979.

**CHARACTERISTICS.**—Differs from the type variety by the large, more fusiform and slender cheilocystidia and by the larger spores. Known only from the type locality in the province of Drenthe in the Netherlands.

#### 50. ENTOLOMA CRYPTOCYSTIDIATUM Arnolds & Noordeloos

*Entoloma cryptocystidium* Arnolds & Noordeloos in *Persoonia* 10: 287-288, 1979.

**CHARACTERISTICS.**—Pileus 20-32 mm broad, conico-convex with slightly involute margin, strongly hygrophanous, pale grey-brown, strongly striate, pallescent on drying, smooth; lamellae crowded, free, ventricose, pale brown-grey; stipe  $47-53 \times 1.5-2.5$  mm, pale grey-brown, silvery striate; smell and taste indistinct; spores  $(8.5-10.2) \times (7.5-8.3) \mu\text{m}$ ; cheilocystidia  $(8-15) \times 4.8-7.0(-7.6) \mu\text{m}$ , cylindrical, sometimes flexuous, often subcapitate, sometimes with thickened, refringent apex, scattered among basidia, not protruding from hymenium; pileipellis a cutis made up of cylindrical, repent,  $3.3-5.8 \mu\text{m}$  wide hyphae with pale diffusely intracellular pigment; clamp-connections frequent in hymenium.

**HABITAT & DISTRIBUTION.**—In non-manured hayfield (*Calthion palustris*) on rather eutrophic wet peaty soil. Known only from the type locality in the province of Drenthe, Netherlands.

**COLLECTIONS EXAMINED.**—NETHERLANDS, prov. Drenthe, Vries, Taarlo, Taarlose Diep, 3 Sept. 1975, E. Arnolds 3408 (holotype WBS, isotype L).


*Entoloma cryptocystidium* has been placed in subsect. *Cheilocystidiata* on account of the intracellular pigment and the dihedral base of the spores. It is better placed here than in sect. *Paramammosi* Romagn. as that section unites the species with cheilocystidia that are not cylindrical-subcapitate, spores with basal facet, and clampless basidia. *Entoloma hirtipes* and varieties differ among other things by the presence of two types of pigment and by differently shaped spores.

51. *Entoloma inutile* (Britz.) Noordeloos, *comb. nov.*—Figs. 40a–e

*Agaricus inutilis* Britz. in Ber. naturhist. Ver. Augsburg 30: 16. 1890. — *Nolanea inutilis* (Britz.) Sacc. & Trav., Syll. Fung. 20: 199. 1911. — *Rhodophyllus inutilis* (Britz.) Romagn. in Rev. Mycol. 1: 160. 1937.

SELECTED ICONES AND DESCRIPTION.—Arnolds & Noordeloos in Fung. rar. Icon. col. 12, pl. 93 fig. e. 1980. — Romagn. in Rev. Mycol. 1: 160, pl. 12 fig. 2. 1937.

CHARACTERISTICS.—Pileus blackish brown; lamellae dark brown with pink tinge; stipe silvery striate; cheilocystidia numerous, lecitiform.



Figs. 40a–e. *Entoloma inutile*. — Habit, spores and cheilocystidia (40a, 40d–e from Kits van Waveren, 26 Nov. 1966; 40b–c from Kramer, 3 Oct. 1968).

Figs. 41a–c. *Entoloma velenovskiyi* var. *velenovskiyi*. — Habit, spores and cheilocystidia (41a–c from Jansen 72–226).

Pileus 12–30 mm broad, convex, then flattened with or without weak umbo or slightly depressed, with margin straight but often undulating with age, hygrophanous, when moist blackish brown (Meth. 6F6), slightly paler towards margin (Meth. 6E6), translucently striate up to two thirds of the radius, on drying pallescent, smooth or at centre slightly rugulose, shining. Lamellae crowded, L = 28–32, I = 1–3, almost free or narrowly adnate, ventricose, up to 5 mm broad, dark brown with pink shade (5 YR 5/4; Meth. 7D4/E4), with serrulate, concolorous or slightly paler edge. Stipe 25–40 × 2.5–4 mm, straight, pale to moderately dark brown, moderately to strongly striate with silvery fibrils, pruinose at apex, solid, then fistulose. Flesh in pileus concolorous with surface, in stipe paler, brittle. Smell and taste indistinct.

Spores (8.7–)9.3–11.5(–12.0) × 7–8.1  $\mu\text{m}$ , Q = (1.15–)1.3–1.4–1.5, L–D = 1.7–3.5–5  $\mu\text{m}$ , pronouncedly nodulose-angular in side-view, with dihedral base. Basidia 24–32 × 9.5–11.5  $\mu\text{m}$ , 4-spored. Cheilocystidia 25–52(–80) × 6–14.5(–17) × 2.5–7  $\mu\text{m}$ , versiform, mostly lecithiform but also subcylindrical-subcapitate or even clavate with conical apex, numerous. Hymenophoral trama regular, composed of fusiform cells 220–350 × 7.5–15(–22)  $\mu\text{m}$ , with intracellular pigment. Pileipellis a simple cutis made up of cylindrical, 3.5–14  $\mu\text{m}$  wide hyphae, at centre of pileus sometimes with cylindrical to clavate, up to 18  $\mu\text{m}$  wide, repent or ascending terminal cells, with intracellular pigment. Pileitrama regular with cylindrical to inflated hyphae, composed of cells up to 370  $\mu\text{m}$  long and 7.5–16(–22)  $\mu\text{m}$  wide, with intracellular pigment particularly in upper layer, and sometimes additional membranal pigments; encrustations never seen. Clamp-connections abundant in hymenium, also observed in covering layers.

**HABITAT & DISTRIBUTION.**—In the Netherlands found on *Calluna* heaths on poor, acid sandy soils and in coastal dunes in *Quercus* forest; rare. October–November.

**COLLECTIONS EXAMINED.**—NETHERLANDS: prov. Friesland, Appelscha, 3 Oct. 1968, R. Kramer (WBS); prov. Drenthe, Beilen, E. of Holthe, 13 Oct. 1976, E. Arnolds 3644 (WBS); prov. Noord-Holland, near Heemstede, Leyduin, 12 Nov. 1966, E. Kits van Waveren.

*Entoloma inutile* has rather distinctive dark lamellae and lecithiform cheilocystidia. The latter character and the striate stipe make it easy to distinguish it from *E. clandestinum*. Its taxonomical situation seems to be somewhat isolated. Orton (1960: 229–230) suggested a relationship with *E. jubatum* and allied species, probably on account of the shape of the cheilocystidia. *Entoloma jubatum* and related species are placed by me in subgenus *Trichopilus* Romagn., characterized among other things by the non-hygrophanous, non-striate, fibrillose-subsquamulose pileus. The Netherlands' collections of *E. inutile*, however, have a distinctly hygrophanous striate pileus which is smooth or only slightly rugulose at centre. This character, together with the nolaneoid habit, almost free lamellae, and type of pigmentation, made me decide to place *E. inutile* in *Nolanea* subsect. *Cheilocystidiata*.

*Rhodophyllus* sect. *Paramammosi* Romagn. does not seem to be suited to accommodate *E. inutile*, as this section is characterized by the combination of non-capitate cheilocystidia and clampless basidia. The species in *Rhodophyllus* sect. *Lecithipori* Romagn. have a totally different habit, typical of subgenus *Eccilia*.

## 52. *Entoloma globulifer* Noordeloos, *spec. nov.*—Figs. 42a–c

Pileus 17–45 mm latus, conicus vel conico-campanulatus demum explanatus, papillatus, hygrophanus, in udo sepiaceus, in sicco pallescens, glaber. Lamellae confertae, subliberae, ventricosae, pallide brunneae demum incarnatae-brunneae. Stipes 40–85 × 2.5–6 mm, cylindraceutus vel compressus, solidus, firmus, pileo concolor, innato fibrilloso-striatus. Caro tenax. Odore saporeque farinaceis.

Spores 9–11.5(–12.4) × 6.8–7.9  $\mu\text{m}$ , Q = 1.3–1.4–1.6. Basidia 2- vel 4-sporigera. Cheilo- et pleurocystidia

17–40 × 12–24  $\mu\text{m}$ , globuliformia vel sphaeropedunculata. Trama lamellarum e elementis subfusiformibus 80–270 × 10–17  $\mu\text{m}$  constantes. Pileipellis cutis differentiatia, hyphae cylindraceae, 2.5–6  $\mu\text{m}$  latae cum cellulis terminalibus clavatis intermixtae. Pigmentis intracellularibus in pileipelle dispersis. Fibulae nullae.

HABITAT: In graminosis inter *Salicem repentem* in dunis. — Typus: *M. E. Noordeloos 811*, 18 X 1978, 'Prov. Zuid-Holland, Isl. Goeree, West-duinen', Netherlands.

CHARACTERISTICS.—Carpophores robust; pileus 17–45 mm broad, conical then expanding, very dark sepia, pallescent; lamellae pale brown with pink tinge; stipe 40–85 × 2.5–6 mm; sepia or horn-brown, silvery striate; smell farinaceous; spores 9.0–11.3(–12.4) × 6.8–7.9  $\mu\text{m}$ ; cheilo- and pleurocystidia present, subglobose; pigment intracellular; clamp-connections absent.

Pileus 17–45 mm broad, conical to conico-campanulate, expanding to convex, finally flattened with broad, obtuse umbo, with outermost margin involute, with marginal zone irregularly undulating with age, hygrophanous, very dark sepia brown when moist, translucently striate at margin, often obscurely so (10 YR 3/3, 4/3, between striae and at margin paler: 10 YR 5/4, rarely 6/4), drying pale brown (10 YR 6/3 to 7/3), brilliantly shining when moist, when dry innately radially satiny-fibrillose, in exposed pilei often radially splitting and pileitrama showing. Lamellae L = 40–50, l = 1–3, fairly crowded, almost free, ventricose, pale brown, then pink with brown tinge (10 YR 6/3, then 7.5 YR 6/4 to 5/4), with more or less serrulate edge, concolorous with or slightly paler than sides. Stipe 40–85 × 2.5–6 mm, cylindrical or flattened, with rounded only slightly swollen base, firm, solid, subfistulose with age, sepia or horn brown (10 YR 6/4, 5/4), sometimes slightly pruinose at apex, downwards glabrous, innately silvery striate lengthwise, with white tomentum at base. Flesh rather firm-subcartilagineous particularly in pileus, but thin, except under papilla in largest specimens. Smell farinaceous. Taste strong, rancid, very unpleasant.

Spores 9.0–11.3(–12.4) × 6.8–7.9  $\mu\text{m}$ , Q = 1.3–1.4–1.6, L–D = 2.4–4.8  $\mu\text{m}$ , rather irregularly asymmetrical in side-view, with 5–7 pronounced angles, with weakly to distinctly dihedral base. Basidia 28–44 × 10.5–14  $\mu\text{m}$ , Q = 2.3–2.9–3.7, 2- and 4-spored. Cheilo- and pleurocystidia broadly globuliform to sphaeropedunculate, 17–40 × 12–24  $\mu\text{m}$ , sometimes slightly thick-walled, hyaline, colourless, scattered in hymenium, particularly abundant near edge of older specimens. Hymenophoral trama regular, composed of subfusiform to cylindrical cells 80–270 × 10–17  $\mu\text{m}$ , with hyaline, colourless walls. Pileipellis a differentiated cutis made up of 2.5–6  $\mu\text{m}$  wide cylindrical hyphae, intermixed with more or less ascending up to 12  $\mu\text{m}$  wide terminal cells of underlying hyphae, with abundant, brown intracellular pigment. Pileitrama regular, composed of cylindrical cells 120–315 × 10–23  $\mu\text{m}$ , with intracellular pigment most abundant in upper layers. Clamp-connections absent.

HABITAT.—In *Salix repens* dominated grassland in old dune area, with *E. papillatum*, *E. staurosporum*, *Hygrophorus* spp. div., *Dermocybe uliginosa*, etc. Known only from the type locality.

COLLECTION EXAMINED.—NETHERLANDS, prov. Zuid-Holland, Isl. of Goeree, Westduinen, 18 Oct. 1978. *M. E. Noordeloos 811* (holotype, L).

The type of pigmentation and the shape of the spores place *Entoloma globulifer* in sect *Endochromonema*. The absence of clamp-connections suggests relationship with *E. cetratum*. The unique cystidia, however, make *E. globulifer* a species in its own right in a somewhat isolated position. *Entoloma sphaerocystis* differs by the slender habit, encrusting pigments and differently shaped spores.

#### **Entoloma** subsection **Icterina** Noordeloos, *subsect. nov.*

*Nolanea* sect. *Icterinae* Konr. & Maubl., Agaricales: 266. 1948 [not validly published; no Latin diagnosis].




Pileus tinctus luteolus vel olivaceus, statura variabilis, truncatus vel planus, subpapillatus vel subumbilicatus. Odor suavissimus frequens. — Typus: *Entoloma icterinum* (Fr.) Moser.

Pileus with yellow and/or olivaceous tinges, variable in shape: truncate-campanulate to flattened, subpapillate to subumbilicate; frequently with strong aromatic smell. — Type: *Entoloma icterinum* (Fr.) Moser.

KEY TO THE SPECIES OF SUBSECTION ICTERINA

- 1a. Pileus yellow, lemon yellow or olivaceous yellow, often with reddish brown flush at centre. . . . .  
 b. Pileus dark brown ('bistre') with olivaceous tinge (as in the pileus of *Naucoria centunculus*).  
*E. ambrosius*, p. 520
- 2a. Cheilocystidia scattered among basidia on edge of lamellae, long subcylindrical-subcapitate to slenderly lageniform-tibiiform; lamellae pale green-olivaceous. . . . . *E. chlorophyllum*, p. 518  
 b. Cheilocystidia absent; lamellae white or pale yellow, then pink . . . . . *E. icterinum*, p. 516



Figs. 42a-c. *Entoloma globulifer*. — Habit, spores and cheilocystidia (42a-c from holotype).

## 53. ENTOLOMA ICTERINUM (Fr.) Moser—Figs. 44a–c

*Agaricus icterinus* Fr., Syst. mycol. 1: 207. 1821. — *Nolanea icterina* (Fr.) Kumm., Führ. Pilzk.: 95. 1871. — *Rhodophyllus icterinus* (Fr.) Quél., Enchir.: 64. 1886. — *Hyporrhodius icterinus* (Fr.) Schroet. in Cohn, KryptogFl. Schles. 3(1): 613. 1889. — *Entoloma icterinum* (Fr.) Moser in Gams, Kl. KryptogFl., 4. Aufl. 2(b/2): 205. 1978.

*Rhodophyllus icterinus* f. *gracillimus* J. Lange in Dansk bot. Ark. 2(11): 37. 1921.

*Agaricus pleopodius* Fr., Syst. mycol. 1: 207. 1821. — *Nolanea pleopodia* (Fr.) Gill., Hymen. Fr.: 420. 1876. — *Rhodophyllus pleopodius* (Fr.) Quél., Enchir.: 64. 1886.

SELECTED ICONES.—Fries. Icon. sel., pl. 99, fig. 4. 1874. — Ricken, Blätterpilze pl. 74, figs. 2, 5. 1913. — J. Lange, Fl. agar. dan. 2: pl. 78A. 1936. — Konr. & Maubl., Icon. sel. Fung., pl. 179 fig. 3. 1930.

CHARACTERISTICS.—Pileus with yellow and/or olivaceous tinge; stipe usually more reddish brown and contrasting with pileus; smell usually strong, aromatical.

Pileus 10–35 mm broad, campanulate or paraboloid, often truncate, then expanding to convex, finally flattened to concave, often with subumbilicate or subpapillate centre, with margin slightly involute or straight, rarely exceeding the lamellae, at marginal zone often slightly sulcate and/or undulating with age, hygrophanous, when moist pale to moderately dark yellow, olivaceous yellow or lemon yellow, at centre often darker and more brown-tinged, paler towards margin (5 Y 7/6, 6/6, 6/4; 2.5 Y 7/6, 6/6, 6/4, 5/4, at centre 5 Y 5/6, 10 YR 5/4, margin 2.5 Y 8/6), translucently striate up to one half of the radius, on drying pallescent to pale yellow or olivaceous yellow (5 Y 8/6, 8/4; Meth. 2A5/6), smooth when moist, on drying with radially arranged aiferous fibrils, sometimes becoming subsquamulose at centre. Lamellae moderately distant, L = 20–30, l = 0–3, usually adnate or emarginate, sometimes broadly adnate with decurrent tooth, segmentiform, then ventricose, up to 6.5 mm broad, white, then pale yellow, finally pink, often with slight brown tinge when old (5 Y 8/2; 2.5 Y 8/6, 8/4; 10 YR 8/6, 8/4, then 7.5 YR 8/4, 7/6), with entire or subserrulate, concolorous edge. Stipe 20–80 × 2–4(–4.5) μm, cylindrical, often slightly broadened at base, straight or flexuous, pale yellowish or greyish brown at apex, downwards more flesh-coloured or pinkish brown, at base often with purplish tinge, more rarely entirely yellowish olivaceous and concolorous with pileus (10 YR 8/4, 8/6, 7/4, 7/6, base 7.5 YR 7/6, 6/6, 5/6, rarely 5 Y 6/4 to 2.5 Y 6/4), minutely pruinose to flocculose at apex, downwards smooth or minutely striate with scattered silvery white fibrils, (often abundantly) white tomentose at base. Flesh subcartilagineous or brittle, yellow-olivaceous in pileus, pale in centre of stipe. Smell strongly aromatical (like amylicetate, fruity, often compared with that of *Entoloma ameides*, *Hebeloma sacchariolens* or *Inocybe pyriodora*), rarely indistinct. Taste weak or nasty-unpleasant.

Spores (8.1–)8.7–11(–11.5) × (6.3–)7.0–8.1(–9.0) μm, Q = (1.15–)1.2–1.35–1.5, L–D = (1.4–)1.7–2.4–4 μm, 5–6-angled in side-view, with pronounced angles, with dihedral base. Basidia 32–52 × 9.5–16 μm, 2–4-spored. Cystidia absent. Hymenophoral trama regular with cylindrical to inflated cells 120–300 × 11.5–17(–21) μm, often with pale intracellular pigment. Pileipellis a cutis made up of radially arranged, 4–17 μm wide cylindrical hyphae with scattered clavate terminal cells 40–92 × 6–25 μm; subpellis with strongly inflated, up to 50 μm wide cells with intracellular pigment. Pileitrama regular with cylindrical to inflated hyphae with cells up to 320 μm long and 11–30 μm wide, often with pale yellowish walls and, particularly in upper layer, intracellular pigment. Clamp-connections abundant in hymenium, more rare and scattered in other tissues.

HABITAT & DISTRIBUTION.—Terrestrial, on rich humus, particularly at places with much nitrogen such as gardens, waste-places etc., often near *Urtica dioica*; common; recorded from Western Europe. July–November.

COLLECTIONS EXAMINED.—NETHERLANDS: prov. Drenthe, Beilen, in garden. Aug. 1977, H. S. C. Huijsman; prov. Overijssel: Hengelo, 24 Oct. 1956, M. G. J. Meijer; Delden, near Hotel Carelshaven, 12 Oct. 1975, E. Kits van Waveren; Twello, Noordijk, 13 Oct. 1957, E. Kits van Waveren; prov. Gelderland: Rheden, Nov. 1953, Mrs. Schutte; Valburg near Nijmegen, estate Oosterhout, 25 Sept. 1954, C. Bas 644; prov. Utrecht: Leusden, 16 Sept. 1978, J. Wisman 60; Amersfoort, estate 'de Treek', 11

Oct. 1958, *E. Kits van Waveren*; Breukelen, estate 'Over Holland', 28 July 1962, *E. Kits van Waveren*; prov. Noord-Holland, Vogelenzang, estate 'Vogelenzang', 11 Nov. 1953, *R. A. Maas Geesteranus* 9533; Castricum, 30 Sept. 1975, *E. Kits van Waveren*; Amsterdam, Amsterdamse bos, 21 July 1960, *E. Kits van Waveren*; Laren, Gooiersgracht, 1 Oct. 1975, *P. A. A. van Winden*; Kortenhoeft, near Kerklaan, 21 Nov. 1970, and near 'de Zuwe', 1 Nov. 1970, *J. Daams*; 's-Gravenland, estate 'Boekesteyn', 28 Aug. 1956, *J. Daams*; prov. Zuid-Holland: Leidschendam, Veenzijdse Polder, 29 Oct. 1969, *C. Bas* 5163; Voorschoten, estate 'Ter Horst', 12 Sept. 1974, *M. E. Noordeloos* 48, and 1 Oct. 1974, *M. E. Noordeloos & A. E. Jansen*; Leiden, in garden, 17 Nov. 1963, *C. Bas* 4033; idem, 16 Oct. 1966 and 22 Nov. 1966, *C. Bas* 4819 a & b; Oegstgeest, estate 'Oud Poelgeest', 26 Aug. 1956, *C. Bas* 1064; prov. Noord-Brabant: Breda, in garden Meerten Verhoff str., 14 Oct. 1967, *P. B. Jansen* 67-206; in garden Paul Windhausenweg 26, 10 Nov. 1974, *P. B. Jansen* 74-275; 11 Oct. 1978, *P. B. Jansen* 78-276 and 30 Nov. 1974, *A. E. Jansen* 57; idem, Liesbos, 19 Aug. 1967, *P. B. Jansen* 67-208; Dorst, loam pits, 24 Aug. 1968, *R. A. Maas Geesteranus* 15266; Nuenen, Nuenense Broek, 15 Sept. 1979, *M. E. Noordeloos* 1019; prov. Limburg: Echt, estate 'De Doordt', 6 Oct. 1962, *C. Bas* 2820; Gronsveld, Savelsbos, 27 Aug. 1979, *J. Schreurs*; Gulpen, 14/20 Sept. 1970, *E. Kits van Waveren*.

BELGIUM, prov. Brabant, Oppem, S. E. of Brussels, Wegembeek, 1 Nov. 1960, *P. Heinemann* (BR).

*Entoloma icterinum* is a rather variable species (Kits van Waveren, 1977: 54-59). Distinctive characters are the yellowish olivaceous tinges in the pileus and the aromatic smell. In literature this smell has been described in various ways: 'burnt sugar' (Quél., 1872: 248); fruity: 'ananas' (J. Lange, 1936: 102); amylicetate or 'bonbons anglais' (Kühn. & Romagn., 1953: 184). Others compare the smell with that of *Hebeloma sacchariolens*. Kits van Waveren (pers. comm.) suggests that it is identical with that of *Entoloma ameides* (Berk. & Br.) Sacc., which I can confirm. In my experience the intensity of the smell is rather variable as well: some collections were smelled before they actually were found, others showed no distinct smell when collected, but after some hours in a closed box they emitted their delicate perfume quite distinctly. Temperature is also of great importance with regard to the intensity of the smell: low temperatures prevent the perception of the smell which becomes distinct when the carpophores are warmed up. This can easily be demonstrated when fresh collections have been kept in the refrigerator for some time.

Fries (1821: 207, 1857: 296, 1874: 207), nor Gillet (1876: 421-422), Cooke (1886, pl. 338), Quélet (1888: 171) or Rea (1922: 404, 405) do mention the smell of *Entoloma icterinum*. This may suggest that either the smell was absent or it was ignored or considered of minor importance. Only Ricken (1913: 297) positively stated that '*Nolanea*' *icterina* is inodorous.

Some authors like Moser (1967: 167, and 1978: 208) consider *Entoloma icterinum* to be distinct from *Agaricus pleopodius* Fr. The differences between *Agaricus icterinus* and *A. pleopodius* are according to Fries:

*A. icterinus*

Pileus campanulate, then convex, not umbilicate, centre sometimes blunt, sometimes umbonate.

Lamellae pale, then yellow, finally pink.

Stipe concolorous with pileus or darker and more brown, entirely pruinose or at apex only.

*A. pleopodius*

Pileus conical, then expanding, subumbonate.

Lamellae white, then pink.

Stipe pale, yellowish, pruinose at apex only.

Moser adds to these differences the lack of an aromatic smell in '*Rhodophyllus pleopodius*' and a difference in spore size, viz.  $10\text{--}12 \times 6\text{--}8 \mu\text{m}$  in *E. icterinum* and  $9\text{--}11 \times 7\text{--}8 \mu\text{m}$  in *R. pleopodius*. As I tried to point out in the discussion above, the lack of a distinct smell cannot be used as a distinctive character. My own observations and those of Kits van Waveren (1977, l.c.) and P. van Winden (1978: 9–10) do not permit a distinction between *E. icterinum* and *Agaricus pleopodius* on specific level.

The shape of the pileus and the sizes of the carpophores are rather variable in *Entoloma icterinum* often within one collection, and show 'typical' *E. icterinum* as well as *A. pleopodius* characters. In addition the lamellae in *E. icterinum* are usually purely white when young; the yellow tinge may appear to be rather pale and is sometimes only suggested when the colour of the pileus shines through the flesh; finally the smell may be present or not in both forms. Spore size in *Entoloma icterinum* is found to cover a range which comprizes both sizes given by Moser, and cannot be used to distinguish the two forms.

#### 54. *Entoloma chlorophyllum* Noordeloos, *spec. nov.* — Figs. 43a–c

Pileus 15–27 mm latus, campanulato-truncatus, centro leviter depressus interdum papillatus, hygrophanus, in udo olivaceus, in sicco pallescens. Lamellae distantes, crassiusculeae, chlorino-olivaceae. Stipes  $50 \times 2\text{--}4$  mm, deorsum bulbosus, olivaceo vel olivaceo-brunneus, argenteo-striatus. Odore nulla.

Sporea ( $8.6\text{--}9.3\text{--}10.4 \times 7.0\text{--}8.1\text{--}(8.7) \mu\text{m}$ ). Basidia 4-sporigera. Cheilocystidia cylindracea vel flexuosa vel lageniformes, interdum capitata, Pileipellis cutis; hyphae cylindraceae,  $6\text{--}21 \mu\text{m}$  latae. Pigmentis intracellulosi pariter in hypharum tunicam, haud incrustantibus. Fibulae presentes. Habitat ad humum in hortum. — Holotypus: *J. J. Barkman 6888*, 27 X 1960, 'Wijster, prov. Drenthe, Netherlands' (WBS).


CHARACTERISTICS.—Pileus olivaceous, weakly hygrophanous, minutely radially fibrillous; lamellae pale olivaceous; flesh inodorous; cheilocystidia present.

Pileus 15–27 mm broad, truncate-campanulate with slight central depression and sometimes with small papilla, with slightly involute margin, weakly hygrophanous, when moist olivaceous-greenish (5 Y 6/6) with slightly darker centre, slightly pallescent on drying, non-striate, minutely radially fibrillous with whitish pruinum at centre, somewhat dull or shining. Lamellae distant, thickish, ventricose, 3–5 mm broad, adnate with small decurrent tooth, pale olivaceous-green. Stipe  $50 \times 2\text{--}4$  mm, cylindrical with abruptly swollen base up to 6 mm wide, olivaceous brown at apex, downwards more brown, flocculose at apex, downwards densely covered with silvery fibrils, often twisted, white tomentose at base. Flesh dark olivaceous when moist, pallescent on drying. Smell absent. Taste not recorded.

Spores ( $8.6\text{--}9.3\text{--}10.4 \times 7.0\text{--}8.1\text{--}(8.7) \mu\text{m}$ ,  $Q = (1.07\text{--})1.2\text{--}1.3\text{--}1.4$ ,  $L\text{--}D = (0.6\text{--})1.2\text{--}3.5 \mu\text{m}$ , 5–6-angled in side-view, with pronounced angles, with dihedral base. Basidia  $32\text{--}50 \times 10\text{--}13.5 \mu\text{m}$ , 4-spored. Cheilocystidia  $52\text{--}85\text{--}(140) \times 10.5\text{--}17.5 \mu\text{m}$ , cylindrical-flexuous with (sub-)capitate apex to slenderly lageniform-tibiiform, scattered among basidia. Hymenophoral trama regular, composed of cylindrical to inflated cells up to  $270 \mu\text{m}$  long and  $12\text{--}18\text{--}(23) \mu\text{m}$  wide, probably with pale intracellular pigment and pale yellow, non-encrusted walls. Pileipellis a cutis made up of radially arranged,  $6\text{--}21 \mu\text{m}$  wide cylindrical hyphae with repent or ascending terminal cells with yellowish brown intracellular pigment. Pileitrama regular, composed of cylindrical or slightly inflated, up to  $28 \mu\text{m}$  wide hyphae with yellow-brown walls. Vascular hyphae numerous. Clamp-connections frequent in hymenium, rare in other tissues.

HABITAT & DISTRIBUTION.—On humus in garden; known only from type locality.

COLLECTION EXAMINED.—NETHERLANDS, prov. Drenthe, Wijster, in lawn on humus, 27 Oct. 1960, *J. J. Barkman 6888* (holotype, WBS).



Figs. 43a-c. *Entoloma chlorophyllum*. — Habit, spores and cheilocystidia (43a-c from holotype).

Figs. 44a-c. *Entoloma icterinum*. — Habit and spores (44a from Kits van Waveren, 20 Sept. 1970; 44b-c from Kits van Waveren, 26 July 1962).

*Entoloma chlorophyllum* is very well characterized by the olivaceous tinges in the lamellae and the enormous protruding cheilocystidia. In many characters such as the habit, the type of pigmentation, the predominantly olivaceous colour of the pileus and particularly the size and the shape of the spores, *E. chlorophyllum* resembles *E. icterinum* very much. For this reason I place *E. chlorophyllum* in subsection *Icterina*. It appears to be better in its place here than in subgenus *Leptonia*, also with regard to the strongly fibrillous-striate stipe and the clamped basidia.

In the field, *E. chlorophyllum* was taken by the collector for *E. versatilis*. The latter, however, has differently shaped cystidia and spores, and lacks clamp-connections in the hymenium.

### EXTRALIMITAL SPECIES

#### 55. *Entoloma ambrosium* (Quél.) Noordeloos, *comb. nov.*

*Rhodophyllum ambrosius* Quél. in C. r. Ass. franç. Av. Sci. 24 (2): 618. 1896 (basionym). — *Pouzarella ambrosia* (Quél.) Mazzet in *Biblca mycol.* 46: 83. 1976.

**CHARACTERISTICS.**—Pileus campanulate-convex, dark olivaceous brown, hygrophanous; lamellae pale grey-brown, then pink; stipe paler than pileus and more reddish brown, not really striate; smell strong, fragrant, like orange blossom.

Pileus about 14 mm broad, campanulate-convex without papilla, with involute margin, slightly undulating, rather dark olive brown ('bistre olive'), translucently striate, strongly pallescent on drying, when moist surface minutely granular (under lens), becoming strongly lustrous on drying. Lamellae distant, thickish, free, slightly ventricose, pale grey-brown, then pink with entire or slightly irregular, concolorous edge. Stipe 60 × 3 mm, cylindrical, somewhat flexuous, distinctly broadened towards base, narrowly fistulose, brown, slightly paler than pileus and more reddish, not striate but with scattered minutely silvery fibrils under lens, in some places nearly pruinose. Flesh thickish in pileus and concolorous with surface, in stipe paler, whitish. Smell strong, like orange blossom.

Spores 7.5–10(11.5) × 6–8 μm, 5–7-angled in side-view, with dihedral base. Basidia 35–50 × 9–11.7 μm, 4-spored. Cystidia absent. Hymenophoral trama regular; hyphae 7.5–18 μm wide. Pileipellis a thin cutis with narrow hyphae; subpellis with inflated, sometimes up to 43 μm wide hyphae. Pigment uniformly membranous, not encrusting. Clamp-connections present in hymenium, rare in other tissues.

**HABITAT.**—Among mosses, near *Pinus sylvestris*; rare, known only from France.

**COLLECTION EXAMINED.**—FRANCE, dept. Oise, Coye-la-Forêt, 22 Nov. 1951, G. Robert & P. Causse (Herb. Romagn., PC).

*Entoloma ambrosium* is easily recognized by its dark pileus, coloured as in *Naucoria centunculus*, and by its aromatic smell. It seems to be rather rare. The description given above is a translation of that of Romagnesi (1974a: 365–368) who found only one, probably juvenile specimen. For line drawings I refer to Romagnesi (1974a, l.c.). The microscopic characters are very similar to those of *E. icterinum*, which has a more yellow tinged pileus.

#### 56. *Entoloma inodorum* (Velen.) Noordeloos

*Nolanea inodora* Velen., České Houby: 629. 1921. — *Entoloma inodorum* (Velen.) Noordeloos in *Persoonia* 10: 251. 1979.

Pileus 10–30 mm broad, campanulate-convex with abrupt papilla, strongly hygrophanous, blackish brown, translucently striate when moist, on drying pallescent to grey, strongly silky

shining, smooth. Lamellae crowded, emarginate, white, then reddish. Stipe long, 1–2 mm broad, transparent, white, not shining, distinctly powdered at apex, hollow, fragile. Smell absent or weakly unpleasant-sourish.

Spores  $10.2\text{--}12.4 \times 8.3\text{--}9.7 \mu\text{m}$ ,  $Q = 1.1\text{--}1.15\text{--}1.25$ ,  $L\text{--}D = 1\text{--}2 \mu\text{m}$ , irregularly cuboid-4-angled in side-view. Clamp-connections present. Pigment intracellular in upper layers of pileus.

HABITAT & DISTRIBUTION.—In grass on sunlit hill, Czechoslovakia. Known only from type locality.

COLLECTION EXAMINED.—CZECHOSLOVAKIA, Bohemia, Myslin, July 1919, *J. Velenovský* (holotype, PRC).

The rather dark pileus strongly contrasting with the whitish stipe as well as the large spores make *E. inodorum* easy to distinguish from all other cuboid-spored *Entoloma*'s in Europe. For details on the type collection of this rather remarkable species the reader is referred to Noordeloos, 1979b: 251.

#### 57. ENTOLOMA PALLESCENS (P. Karst.) Noordeloos—Figs. 45a–d

*Nolanea pascua* var. *pallescens* P. Karst., Hattusv. I: 280. 1879. — *Nolanea pallescens* (P. Karst.) P. Karst. in Meddn Soc. Fauna Flora fenn. 16: 94. 1890 (Symb. Myc. fenn. 29). — *Entoloma pallescens* (P. Karst.) Noordeloos in Persoonia 10: 251. 1979.

*Nolanea pallescens* var. *procera* P. Karst. in Meddn Soc. Fauna Flora fenn. 16: 64. 1890.

*Nolanea majalis* Velen., Novitates mycologicae novissimae: 147. 1939.

CHARACTERISTICS.—Carpophores medium-sized to rather large; pileus up to 75 mm broad, conical to conico-campanulate, only slightly expanding, yellowish brown to reddish brown, strongly pallescent on drying; lamellae pale, then brown, finally reddish brown; stipe up to 150 mm long and 4–11(–14) mm broad, concolorous with pileus or distinctly paler, strongly silvery striate lengthwise; pigment membranal and intracellular; basidia 4-spored, with clamp.

Pileus 25–75 mm broad, acutely conical or conico-campanulate, only slightly expanding, rarely flattened, with conical papilla, with margin slightly involute when young, finally straight, often with plicate marginal zone particularly when young and not expanded; strongly hygrophanous, when moist yellowish brown to reddish brown, darkening with age and when water-soaked (10 YR 6/6; 7.5 YR 5/6, 4/4, finally 7.5 YR 3/2), slightly paler towards margin (10 YR 6/6; 7.5 YR 6/8), translucently striate up to three fourths of radius; on drying strongly pallescent to pale brown, smooth, glabrous, innately fibrillose, shining. Lamellae  $L =$  up to 50,  $l = 3\text{--}5$ , almost free, ventricose, pale, then brown, finally reddish brown (10 YR 7/6, 6/6, then 7.5 YR 5/6, 4/4, 3/2), with slightly eroded, concolorous edge. Stipe up to  $150 \times 4\text{--}11(14)$  mm, cylindrical or gradually broadening towards base, pale to moderately dark brown, usually paler than pileus (10 YR 6/5, 5/4, 3/4; 7.5 YR 3/2), strongly silvery striate lengthwise, often twisted, white-tomentose at base. Flesh fairly brittle, concolorous with surfaces or slightly paler in inner parts. Smell faint. Taste mild.

Spores  $(9.3\text{--})10.0\text{--}12.0 \times (7.0\text{--})7.4\text{--}8.1(9.0) \mu\text{m}$ ,  $Q = 1.2\text{--}1.4\text{--}1.5$ ,  $L\text{--}D = 1.7\text{--}2.5\text{--}4.0 \mu\text{m}$ , 5–6–7-angled in side-view, with distinctly dihedral base. Basidia  $32\text{--}52 \times 10.5\text{--}14 \mu\text{m}$ , 2–4-spored. Cystidia absent. Hymenophoral trama regular, with cells  $160\text{--}340(460) \times (3.5\text{--})7\text{--}16(27) \mu\text{m}$ . Pileipellis a cutis made up of  $3.5\text{--}10 \mu\text{m}$  wide cylindrical hyphae with brown walls and pale diffusely intracellular pigment. Pileitrama regular, composed of strongly inflated cells  $230\text{--}450 \times 12\text{--}32 \mu\text{m}$ . Clamp-connections abundant in hymenium, rare elsewhere.

HABITAT & DISTRIBUTION.—In subboreal and submountainous coniferous forest. In Norway found in mixed forest of *Picea abies*, *Pinus sylvestris* and scattered *Betula pubescens* with





understory of *Vaccinium vitis-idaea* and *Linna borealis*; May–October. Known to occur in Finland, Norway and Czechoslovakia.

COLLECTIONS EXAMINED.—FINLAND, Tavestia australis, Tammela, Syrjä, Mustiala-W, 6 Oct. 1889, P. A. Karsten (neotype, design. mihi, Herb. Karsten, H). —NORWAY: Östfold, Moss, Jelöya, 28 May 1972, Gro Gulden 181/72 (O); Hedmark, Tynset, Bygdemuseum, 21 July 1979, M. E. Noordeloos 975. —CZECHOSLOVAKIA, Bohemia, Mnichovice near Prague, 12 May 1937, J. Velenovský (holotype of *Nolanea majalis* Velen., PRM).

In the Karsten Herbarium at Helsinki only one collection is available of *Nolanea pallescens* P. Karst., collected and named by the author. It consists of one specimen in good state and was selected by me as a neotype. The neotype collection has the following microscopical characters:

Spores (10.0–)10.4–12.1(–12.7) × (6.9–)7.4–8.5(–9) μm, Q = (1.25–)1.3–1.4–1.5, L–D = 2.3–4.0 μm, 5–6(–7)-angled in side-view, with dihedral base. Basidia 35–42 × 10.5–14.0 μm, 4-spored. Hymenophoral trama regular with strongly inflated cells 226–340(–460) × 17–27 μm, with thin, colourless walls. Pileipellis a cutis made up of cylindrical, up to 10 μm wide hyphae with pale intracellular pigment and pale brown coloured walls; no encrustations seen. Pileitrama regular, with strongly inflated cells 230–450 × 12–32 μm. Clamp-connections abundant in hymenium; elsewhere not seen.

*Entoloma pallescens* is closely related to *E. cuneatum* and *E. cetratum*. From *E. cuneatum* it differs by the unicoloured pileus, different habitat and probably also by the more robust habit. *Entoloma cetratum* is usually smaller, and in addition has bisporous, clampless basidia.

*Entoloma pallescens* occurs in the same habitat and seems to have more or less the same distribution as *E. hirtipes* var. *hirtipes*. The latter species, however, is strictly vernal, has cheilocystidia, larger spores, and a strong rancid-farinaceous smell.

## 58. ENTOLOMA PRISMATOSPERMUM (Romagn.) Horak

*Rhodophyllum prismatospermum* Romagn. in (Trav. mycol. död. R. Kühner) Bull. Soc. linn. Lyon 43 (No. spéc.): 386. 1974. — *Entoloma prismatospermum* (Romagn.) Horak in Sydowia 28: 216. 1976 ("1974/75").

Pileus 5–9 mm broad, conical, then expanding to convex with or without distinct papilla, with margin involute when young, then straight, hygrophanous, hardly striate when moist, pale grey-brown with darker papilla and paler margin, on drying pallescent, glabrous, satiny lustrous. Lamellae fairly distant, thickish, with 1–3 lamellulae between each pair, slightly ascending, broadly adnate to uncinately, narrow, up to 1.5 mm broad, greyish to grey-brown without any pink tinge, with whitish edge. Stipe (10–)20–30 × 1–2 mm, cylindrical, thickened to subbulbous at base, fairly pale grey-brown to pearl grey, becoming browner with age, paler towards base or with white mycelium, minutely flocculose at apex, more sparsely so towards base. Flesh thin, grey-brown in pileus, pale in stipe. Smell and taste farinaceous-rancid.

Spores 7.5–9 × (6–)7–8.5 μm, 3–4-angled in side-view, prismatical-cuboid with fairly blunt angles. Basidia 50–60 × 12–14 μm, 4-spored. Cheilocystidia numerous, fusiform often with broadly swollen basal part, 60–75 × 9–12 μm. Pileipellis a simple cutis made up of 3.5–7 μm wide cylindrical hyphae with pale yellow, non-encrusted walls without any intracellular pigment. Stipiteipellis with caulocystidia similar to cheilocystidia. Clamp-connections found in hymenium.

COLLECTION EXAMINED.—FRANCE, dept. Oise, Neuville-en-Hez, 5 Sept. 1956, *H. Romagnesi* 56.146 holotype, (Herb. Romagn., PC).

The description given above is a concise translation of the original diagnosis. For full details and line drawings the reader is referred to Romagnesi, l.c.: 369–372. *Entoloma prismatospermum* is known only from its type locality. It differs from all other cuboid-spored species of *Entoloma* in Europe by its characteristic cheilo- and caulocystidia and simple pigmentation. *Entoloma inodorum* differs among other things by the larger and more cuboid spores, and *E. rhombisporum* by differently shaped cheilocystidia, a smooth stipe and a different pigmentation. Romagnesi suggested that the small size of the carpophores in *E. prismatospermum* might be due to the young state of the type collection.

### INSUFFICIENTLY KNOWN AND EXCLUDED TAXA

**alba.**—*Nolanea alba* Velen., České Houby: 629. 1921. — *Entoloma nivescens* Noordeloos in *Persoonia* 10: 246. 1979 (name change).

CHARACTERISTICS.—Pileus 20–40 mm broad, obtusely convex, sometimes weakly umbonate with undulating, crenulate margin, distinctly hygrophanous, when moist pale grey and translucently striate, on drying pallescent to white, smooth, shining; lamellae broadly ventricose, white, then pink, with denticulate edge; stipe about 40–80 × 2–4 mm, white, smooth, furfuraceous at apex only; flesh inodorous; spores (9.0–)9.6–10.8(–11.3) × 6.8–7.9 μm; basidia 4-spored; cheilocystidia 40–55 × 7–9 μm, cylindrical-subcapitate; pileipellis a cutis made up of 4–11 μm wide cylindrical hyphae without pigment; clamp-connections abundant in hymenium.

HABITAT & DISTRIBUTION.—Under *Corylus*; Czechoslovakia; rare.

COLLECTION EXAMINED.—CZECHOSLOVAKIA, Bohemia, Mnichovice, collis Rlechač, July 1940, *J. Velenovský* (neotype, PRM).

*Entoloma nivescens* seems to be closely related to *E. hirtipes* var. *hirtipes* and *E. leptopus* by size and shape of the spores and the cheilocystidia. The very pale colour makes it a species of its own kind, which needs rediscovery.

**araneosa.**—*Nolanea araneosa* Quél. in Bull. Soc. bot. Fr. 23: 327. 1876. — Belongs to subgenus *Pouzaromyces* (see Noordeloos 1979a: 234).

**babingtonii.**—*Nolanea babingtonii* (Blox. apud Berk. & Br.) Sacc., Syll. Fung. 5: 717. 1887. — Nomen dubium (see Noordeloos 1979a: 241).

**brassicolens.**—*Nolanea brassicolens* Reid in Trans. Br. mycol. Soc. 48: 518. 1965.

This fungus has to be transferred to subgenus *Entoloma* on account of size and shape of the 'barrel'-shaped cells forming the hymenophoral and pileal trama. In addition the habit suggests relationship with members of the *Entoloma nidorosum* complex. It will be treated in a future paper (Noordeloos, 1980a).

**dysthales.**—*Nolanea dysthales* (Peck) Murrill in North Am. Fl. 10: 101. 1917. — Belongs to the subgenus *Pouzaromyces* (see Noordeloos, 1979a: 215).

**foetida.**—*Nolanea foetida* Killermann, Pilze Bayern 2: 55. 1925. — Figs. 47a–b.

Characteristics.—Pileus 20–30 mm broad, campanulate or hemispherical, then expanding to convex with weak umbo, thin-fleshed, hygrophanous, when moist fulvous, striate, on drying pallescent; lamellae almost free, rusty brown when mature (ferrugineous); stipe about 60 × 2–4 mm, cylindrical, straight or slightly broadening towards base, concolorous with pileus; smell strong, foetid, like that of a putrefying cadaver; spores' 11–13.9(–15.0) × (7.1–)7.6–9.3  $\mu\text{m}$ , Q = 1.25–1.5–1.7(–2.0), L–D = 2.5–4.0–6.5  $\mu\text{m}$ , irregularly 6–8-angled in side-view, with dihedral base; basidia 24–37(–42) × 8.0–12.7  $\mu\text{m}$ , 4-spored. Cystidia absent. Pileipellis a cutis made up of 4.7–10.4  $\mu\text{m}$  wide cylindrical hyphae with membranal pigment. Clamp-connections seen in hymenium. Vascular hyphae numerous in pileitrama.

HABITAT & DISTRIBUTION.—In forest between decaying needles of *Pinus*; known only from Bavaria (German Federal Republic).

COLLECTION EXAMINED.—GERMAN FEDERAL REPUBLIC, Bavaria, near Regensburg, 1 June 1913, S. Killermann (holotype, M).

*Nolanea foetida* belongs undoubtedly to subsection *Endochromonema*. Killermann considered it to be closely related to *Entoloma cetratum*. Though the macroscopical characters and the habitat are similar, *Nolanea foetida* cannot be reduced to a form or variety of *Entoloma cetratum* as there are in addition to the foetid smell some striking microscopical differences, respectively (i) the larger and more irregularly shaped spores; (ii) the 4-spored basidia; (iii) the presence of clamp-connections in the hymenium. The last two characters indicate relationship with *E. cuneatum* and *E. pallescens*. Recombination of Killermann's epithet would result in a later homonym of *Entoloma foetidum* Hesl. 1967. Therefore the following new name is introduced: **Entoloma foetulentum** Noordeloos, *nom. nov.* (basionym *Nolanea foetida* Killermann, Pilze Bayern 2: 55. 1925).

**fulviceps.**—*Rhodophyllus fulviceps* Romagn. in Bull. Soc. mycol. Fr. 48: 321. 1932. — Fig. 46.

CHARACTERISTICS.—Pileus 13–15 mm broad, hemispherical or convex, expanding, not umbonate, hygrophanous, when moist brown with reddish ochraceous tinge, with paler margin, translucently striate, pallescent on drying; lamellae rather distant, free, very strongly ventricose, pinkish brown; stipe 30–45 × 1.8 mm, cylindrical, reddish or yellowish, shining, not striate, white tomentose at base; flesh reddish, inodorous; spores (9.3–)10.0–11.5 × 7.6–8.1(–8.7)  $\mu\text{m}$ , Q = 1.2–1.4–1.5, L–D = 1.5–2.5–4.0  $\mu\text{m}$ , (5–)6–7(–8)-angled in side-view, with dihedral base; basidia 2-spored; clamp-connections not seen.

**HABITAT & DISTRIBUTION.**—In (frondose?) forest between mosses. Known only from the type locality.

**COLLECTION EXAMINED.**—France, dept. Seine & Oise, Yerres, April 1932, *H. Romagnesi 129* (holotype, herb. Romagn., PC).

The type collection is in a very poor state; the tissues are difficult to study. I failed to find intact basidia, but they seem to be mostly 2-spored and clampless. *Rhodophyllus fulviceps* most probably represents a form of *Entoloma cetratum*.

**hirta.**—*Nolanea hirta* Velen. in *Mykologia* 6: 28. 1929. — Belongs to subgenus *Pouzaromyces* (see Noordeloos, 1979a: 223).

**limosella.**—*Nolanea limosella* P. D. Orton in *Notes R. bot. Gdn Edinb.* 29: 108. 1969.

**CHARACTERISTICS.**—Cap 12–24 mm, conico-convex then expanded rather acutely umbonate, later often depressed around umbo, dark sepia-horn, paler when dry, smooth, when moist striate at margin, when dry silky shiny and sometimes rugulose at centre sub lente. Gills adnate often slightly emarginate, whitish then pale brownish pink, not crowded, L = 18–22, l = 1–(3), rather narrow at first then ventricose near stem, edge even, fairly thick. Stem 15–30/2–3 mm, equal or +/– clavate, pale grey-horn, apex almost white and white pruinose, remainder whitish silky striate, stuffed, base white tomentose. Flesh concolorous, white in centre of cap and stem. Smell fairly strong, mealy, especially when cut. Spores nodulose oblong-angular, (10)11–13/7–9  $\mu$ m. Basidia 2-spored. Cystidia absent.

**HABITAT.**—On mud at edge of pond.

Judging from the description *Nolanea limosella* is very close to *Entoloma cetratum* but has the colours of *E. cuspidifer*. That species, however, has differently shaped spores, viz. more broadly ellipsoid to subsodiametrical. The type collections at K and E contain fragments of a brown-spored agaric which evidently has been put erroneously under this label. I was therefore unable to study critically the microscopical characters of *Nolanea limosella*, and for the moment consider it a nomen dubium.

**mamosus.**—*Agaricus mamosus* L. ex Fr., *Spicilegium*: 7. 1836. — *Nolanea mamosus* (L. ex Fr.) Quél. in *Mém. Soc. Emul. Montbéliard, Sér. II*, 5: 122. 1872 (Champ. Jura Vosges 1). — *Rhodophyllus mamosus* (L. ex Fr.) Quél., *Enchir.*: 64. 1886. — *Hyporrhodius mamosus* (L. ex Fr.) Schroet. in *Cohn, KryptogFl. Schles.* 3 (1): 614. 1889. — *Entoloma mamosum* (L. ex Fr.) Hesler in *Beih. Nova Hedwigia* 23: 185. 1967.

Fries (1836: 7) placed *Agaricus mamosus* in tribus *Nolanea* close to *A. pascuus* Pers. ex Fr. and *A. clandestinus* Fr., apparently considering the polished, smooth stipe the most important difference with *A. pascuus*; Fries had this character printed in italics. His *A. clandestinus* differs from *A. mamosus* by the dark coloured lamellae. Furthermore Fries referred to the plates of Schaeffer (1800, pl. 254) and Batsch (1783, pl. 5), both depicting a small *Nolanea* with a moderately dark brown pileus and a smooth, shining stipe.

Another character mentioned by Fries a few years later (1838: 156) is the peculiar smell. This may be considered important, as Fries seemingly had no good sense for odours. Unfortunately in this case he did not indicate the nature of it.

Considering this, I came to the conclusion that among the different interpretations of *Agaricus mammosus* Fr. in literature, that of Konrad (1923: 36 and 1929: 50) comes closest to that of Fries. This interpretation might be the same as P. D. Orton's (1960: 329). I did not study authentic material neither from Konrad nor from P. D. Orton, and I was unable to establish the occurrence of this taxon in the Netherlands.

*Rhodophyllus mammosus* sensu Kühn. & Romagn. (1953: 188) is conspecific with *Entoloma hirtipes* (Schum. ex Fr.) Moser.

*Agaricus mammosus* var. *temuioir* Fr., Icon. sel. 1: 113, pl. 98 fig. 4. 1867 is the same as *Entoloma papillatum* (Bres.) Dennis (see p. 454).

**nivescens.**—*Entoloma nivescens* Noordeloos in Persoonia 10: 246. 1979. See under *Nolanea alba* Velen. on p. 524.

**pascuus.**—*Agaricus pascuus* Pers., Synopsis: 427. 1801. — *Agaricus pascuus* Pers. ex Fr., Syst. mycol. 1: 205. 1821. — *Nolanea pascua* (Pers. ex Fr.) Kumm. Führ. Pilzk.: 95. 1871. — *Rhodophyllus pascuus* (Pers. ex Fr.) Quél., Enchir.: 63. 1886. — *Hyporrhodius pascuus* (Pers. ex Fr.) Schroet. in Cohn, KryptogFl. Schles. 3 (1): 614. 1889. — *Entoloma pascuum* (Pers. ex Fr.) Donk in Bull. bot. Gdns Buitenzorg, ser. 3, 18: 158. 1948.

*Agaricus pascuus* is a collective species, comprizing at least three common grassland-species, viz. *E. conferendum*, *E. sericeum*, and *E. vernum*. As the protologue does not permit an exact definition it is considered a nomen dubium (see p. 449).

**pseudotelamonia.**—*Entoloma pseudotelamonia* Noordeloos in Persoonia 10: 263. 1979. — *Telamonia brevipes* Velen., České Houby: 458. 1921, non *Entoloma brevipes* Murrill 1917.

**CHARACTERISTICS.**—General habit that of a species of *Telamonia* (Cortinariaceae); pileus 20–40 mm broad, more or less flattened or with weakly depressed centre, hygrophanous, when moist very dark chestnut brown, centre almost black, not striate, pallescent on drying; lamellae somewhat crowded, deeply emarginate, sordid white, then ochraceous, finally reddish brown; stipe about as long as diameter of pileus, 10 mm thick, brown, with white striation, base with ochraceous 'veil'; flesh brownish, inodorous; spores 8.2–9.7(–10.3) × (7.7–)8.2–8.7(–9.3) μm, Q=1.0–1.05–1.1(–1.15), isodiametrically 5(–6)-angled in side-view; basidia 4-spored, with cystidia absent; pigment encrusting.

**HABITAT.**—In coniferous forest; known only from type locality.

**COLLECTION EXAMINED.**—CZECHOSLOVAKIA, Bohemia, Mnichovice, August 1915, J. Velenovský (holotype; PRC, bottle 183).

*Entoloma pseudotelamonia* is microscopically very much like *E. juncinum* but differs from the latter by the different habit, the dark pigmented, flattened, often depressed pileus and by the habitat.

**putus.**—*Rhodophyllus putus* Romagn. in Bull. Soc. mycol. Fr. 48: 322. 1932. — Fig. 48.

CHARACTERISTICS.—Pileus 15–25 mm broad, variable in shape, campanulate or convex, sometimes flattened or with reflexed margin, with or without umbo, hygrophanous, when moist reddish bistre, sometimes with ochraceous tinge, translucently striate, on drying pallescent to ochraceous or pale reddish brown; lamellae relatively distant, not ventricose, white, then pale pink with slight brown tinge; stipe 36–60 × 2–3 mm, cylindrical, smooth, not striate, shining, whitish or pale reddish brown, much paler at apex; flesh pale reddish, fragile; smell absent; spores (9.3–)10–12(–12.7) × 7.6–8.1(–9.3)  $\mu\text{m}$ ,  $Q = 1.25\text{--}1.3\text{--}1.6$ ,  $L\text{--}D = 2\text{--}3.5\text{--}4.5$   $\mu\text{m}$ , irregularly 6–8-angled in side-view, with dihedral base; basidia 2-spored, without clamp-connections; cystidia absent; pigment probably membranous, no encrustations seen.

HABITAT & DISTRIBUTION.—In forest; known only from type locality.

COLLECTION EXAMINED.—FRANCE, dept. Seine & Oise, Yerres, May 1937, *H. Romagnesi* 127 (holotype; herb. Romagn., PC).

*Rhodophyllus putus* undoubtedly belongs to the *Entoloma cetratum* complex. The smooth, shining stipe resembles that of *E. farinogustus* which differs, however, by colour, habit, strongly farinaceous-rancid taste, and by habitat.

**radiatus.**—*Rhodophyllus elaphinus* var. *radiatus* J. Lange in Dansk bot. Ark. 2 (11): 31. 1921. — *Rhodophyllus radiatus* (J. Lange) J. Lange, Fl. agar. dan. 2: 96. 1939. — *Nolanea radiata* (J. Lange) P. D. Orton in Trans. Br. mycol. Soc. 43: 179. 1960.

*Rhodophyllus radiatus* in its original sense represents a species that belongs to subgenus *Entoloma*, and is close to *R. sordidulus* Kühn. & Romagn. *Nolanea radiata* sensu P. D. Orton seems to be another taxon, close to *E. sericeum* (see p. 482).

**robiniae.**—*Nolanea robiniae* Velen., Novitates mycologicae novissimae: 78. 1947. — *Entoloma robiniae* (Velen.) Noordeloos in Persoonia 10: 252. 1979.

CHARACTERISTICS.—Carpophores slender, mycenoid; pileus 6–8 mm broad, hemispherical to convex, with sulcate margin, not umbonate, pale greyish ochraceous when moist; lamellae crowded, almost free, pale, then pink; stipe long, 1 mm thick, shining, striate, twisted, with basal tomentum; spores 9.0–10.2(–11.0) × 7.4–7.9  $\mu\text{m}$ ,  $Q = 1.2\text{--}1.3\text{--}1.4$ , 5–6-angled in side-view, with blunt base (basal facet?); basidia 4-spored, with clamp-connection; cystidia absent; pigment membranous (?); subpellis cellular, strongly developed.

HABITAT & DISTRIBUTION.—In *Robinia pseudoacacia* forest; known only from type locality.

COLLECTION EXAMINED.—CZECHOSLOVAKIA, Bohemia, Mnichovice, July 1940, J. Velenovský (holotype, PRM).

The slender mycenoid habit and pale colours of *E. robiniae* are very distinctive. Furthermore the habitat and the strongly developed subpellicular layer in the pileus are characteristic. This species belongs to subsection *Endochromonema*, close to *E. cuneatum*, *E. lanuginosipes* and *E. pallescens*.

**setulosa.**—*Nolanea setulosa* Velen., *Novitates mycologicae*: 147. 1939.

Identical with *Entoloma hirtum* (Velen.) Noordeloos in subgenus *Pouzaromyces* (see Noordeloos, 1979a: 223).

**strigosissima.**—*Nolanea strigosissima* Rea in *Trans. Br. mycol. Soc.* 6: 325. 1920.

Belongs to subgenus *Pouzaromyces* (see Noordeloos, 1979a: 211).

**testacea.**—*Nolanea cetrata* var. *testacea* Bres., *Fungi trident.* 1: 77, pl. 83 fig. 1. 1887. — *Nolanea testacea* (Bres.) P. D. Orton in *Trans. Br. mycol. Soc.* 43: 179. 1960.

The 2-spored taxon to which P. D. Orton applied this epitheton represents a form or variety of *Entoloma cetratum* with reddish date brown pileus, gathered under *Betula* and *Corylus* and on a *Calluna* heath, near *Betula*. There is some doubt whether this is the same as the original 'testacea' depicted by Bresadola, with 4-spored basidia. I failed to locate the type or other authentic material.

This case again illustrates the need of careful collecting and comparison of *E. cetratum*-like fungi in all sorts of habitats in western and northern Europe, to get a clear picture of the variability of this species. It might then appear to be a very polymorphic one, including such forms as *Nolanea testacea*, *Rhodophyllus fulviceps* Romagn., *R. putus* Romagn., and perhaps even *E. farinogustus* Arnolds & Noordeloos.

**versatilis.**—*Nolanea versatilis* (Fr.) Gill., *Hymenom. Fr.*: 418. 1874.

Belongs to subgenus *Pouzaromyces* (see Noordeloos, 1979a: 230).

**vinaceus.**—*Agaricus vinaceus* Scop., *Fl. carn.* 1: 444. 1772. — *Agaricus vinaceus* Scop. ex Fr., *Epicr.*: 157. 1838. — *Nolanea vinacea* (Scop. ex Fr.) Kumm., *Führ. Pilzk.*: 95. 1871. — *Rhodophyllus vinaceus* (Scop. ex Fr.) Quél., *Enchir.*: 64. 1886. — *Entoloma vinaceum* (Scop. ex Fr.) Arnolds & Noordeloos in *Persoonia* 10: 298. 1979.

*Entoloma vinaceum* and its varieties *fumosipes* Arnolds & Noordeloos and *violeipes* Arnolds & Noordeloos are considered to belong to subgenus **Entoloma** section **Turfosa** (Kühn. & Romagn. ex Romagn.) Noordeloos, *comb. nov.* [basionym *Rhodophyllus* sect. *Turfosi* Kühn. & Romagn. ex Romagn. in (*Trav. mycol. d'éd. R. Kühner*) *Bull. mens. Soc. linn. Lyon* 43 (Num. spéc.): 387. 1974].

## REFERENCES

- ARNOLDS, E. J. M. & NOORDELOOS, M. E. (1979). New species of *Entoloma* from grasslands in the prov. of Drenthe, the Netherlands. *In Persoonia* 10: 283–300.
- & — (1980). New, rare and interesting species of *Entoloma*. *In Fungorum rariorum icones coloratae* 12: pl. 89–96. J. Cramer, Vaduz.
- BAS, C. (1965). The genus *Squamanita*. *In Persoonia* 3: 331–364.
- (1969). Morphology and subdivision of *Amanita* and a monograph of its section *Lepidella*. *In Persoonia* 5: 285–579.
- BATSCH, J. (1783). *Elenchus Fungorum. Halae Magdeburgicae.*
- BATTARRA, A. J. A. (1755). *Fungorum agri Ariminensis Historia. Faventiae.*
- BRESADOLA, G. (1929). *Iconographia Mycologica* 12: pl. 551–600. Mediolani.
- COOKE, M. C. (1886). *Illustrations of British Fungi* 3. London.
- DENNIS, R. W. G. (1953). Les Agaricales de l'île de la Trinité: Rhodosporeae-Ochrosporeae. *In Bull. Soc. mycol. Fr.* 69: 145–198.
- EINHELLINGER, A. (1969). Die Pilze der Garginger Heide. *In Ber. bayer. bot. Ges.* 41: 79–130.
- (1976). Die Pilze in primären und sekundären Pflanzengesellschaften oberbayerischer Moore. *In Ber. bayer. bot. Ges.* 47: 75–149.
- FAVRE, J. (1936). Champignons rares ou peu connus des hauts-marais jurassiens. *In Bull. Soc. mycol. Fr.* 52: 129–146.
- (1948). Les associations fongiques des Hauts Marais jurassiens et de quelques régions voisines. *In Matér. Fl. Crypt. Suisse* 10 (3): 43–59.
- FRIES, E. (1821). *Systema mycologicum* 1. Lundae.
- (1836). *Spicilegium plantarum neglectarum. Decadem primam, Agaricos Hyperhodios sistentem. Upsaliae.*
- (1838). *Epicrisis Systematis Mycologici seu Synopsis Hymenomycetum. Upsaliae.*
- (1857). *Monographia Hymenomycetum Sueciae* 2. (Reprint, 1963, Amsterdam.)
- (1874). *Hymenomycetes europaei. Upsaliae.*
- GILLET, C. C. (1876). Les Hyménomycètes. Alençon 1874–1878.
- HESLER, L. R. (1967). *Entoloma* in southeastern North America. *In Beih. Nova Hedwigia* 23.
- HORAK, E. (1973). Fungi Agaricini Novaezelandiae. *Entoloma* (Fr.) and related genera. *In Beih. Nova Hedwigia* 43.
- (1976). On cuboid-spored species of *Entoloma* (Agaricales). *In Sydowia* 28: 171–236 '1974/75'.
- (1978). *Entoloma* in South America. *In Sydowia* 30: 40–111 '1977'.
- JOSSERAND, M. (1937). Champignons de la région lyonnaise. *In Bull. Soc. mycol. Fr.* 53: 178–230.
- KALCHBRENNER, K. (1873). *Icones selectae hymenomycetum Hungariae. Pest.*
- KITS VAN WAVEREN, E. (1977). *Rhodophyllus icterinus* versus *Rhodophyllus pleopodius*. *In Coolia* 20: 54–58.
- KONRAD, P. (1923). Notes critiques sur quelques Champignons du Jura. *In Bull. Soc. mycol. Fr.* 39: 27–45.
- (1925). Notes critiques sur quelques Champignons du Jura 2. *In Bull. Soc. mycol. Fr.* 41: 33–70.
- (1929). Notes critiques sur quelques Champignons du Jura 4. *In Bull. Soc. mycol. Fr.* 45: 35–77.
- KONRAD, P. & MAUBLANC, A. (1930). *Icones selectae fungorum* 2. Paris.
- KÜHNER, R. (1977). Agaricales de la zone alpine. Genre *Rhodophyllus* Quél. *In Bull. Soc. mycol. Fr.* 93: 445–502.
- KÜHNER, R. & ROMAGNESI, H. (1953). *Flore analytique des champignons supérieurs. Paris.*
- KUMMER, P. (1871). *Der Führer in die Pilzkunde. Zerbst.*
- LANGE, J. (1936). *Flora agaricina danica* 2. Copenhagen.
- (1940). *Flora agaricina danica* 5. Copenhagen.
- LARGENT, D. L. (1974). Rhodophylloid Fungi of the Pacific Coast (United States) IV: Infrageneric concepts in *Entoloma*, *Nolanea* and *Leptonia*. *In Mycologia* 66: 987–1021.
- (1977). The genus *Leptonia* on the Pacific Coast of the United States, including a study of North American types. *In Bibliotheca mycol.* 55. J. Cramer, Vaduz.
- LARGENT, D. L. & BENEDICT, R. G. (1971). Studies in the rhodophylloid fungi I. Generic concepts. *In Macroño* 21: 32–39.



- MAZZER, S. J. (1977). *Nolanea verna* in North America. *In* Mich. Bot. **16**: 195–200.
- MOSER, M. (1967). Die Röhrlinge und Blätterpilze. *In* Gams, Kl. KryptogFl. 3. Aufl., 2 (b/2). Stuttgart.
- (1978). Die Röhrlinge und Blätterpilze. *In* Gams, Kl. KryptogFl. 4. Aufl., 2(b/2). Stuttgart.
- NOORDELOOS, M. E. (1979a). *Entoloma* subgenus *Pouzaromyces* emend. in Europe. *In* Persoonia **10**: 207–243.
- (1979b). Type studies on entolomatoid species in the Velenovský Herbarium I. Species described in the genera *Nolanea*, *Leptonia* and *Telamonia*. *In* Persoonia **10**: 245–265.
- (1980a). *Entoloma* subgenus *Entoloma* in the Netherlands and adjacent regions, with a reconnaissance of its remaining taxa in Europe. *In* Persoonia **11** (1) (in preparation).
- (1980b). Introduction to the taxonomy and characters of *Entoloma*. *In* Persoonia **11** (1) (in preparation).
- ORTON, P. D. (1960). New Check List of British Agarics and Boleti 3. Notes on genera and species in the List. *In* Trans. Br. mycol. Soc. **43**: 159–439.
- ORTON, P. D., DENNIS, R. W. G. & HORA, F. B. (1960). New Check List of British Agarics and Boleti. *In* Trans. Br. mycol. Soc. **43**. Suppl. 1–225.
- QUÉLET, L. (1872). Les Champignons du Jura et des Vosges. *In* Mém. Soc. Emul. Montbéliard, Sér. II, **5**: 43–332.
- (1886). Enchiridion fungorum in Europa media et praesertim in Gallia vigentium. Lutetiae.
- (1888). Flore mycologique de la France. Paris.
- REA, C. (1922). British Basidiomycetae. Cambridge.
- RICKEN, A. (1913). Die Blätterpilze: 257–320, pl. 65–80. Leipzig.
- ROMAGNESI, H. (1954). In Kühner & Romagnesi, Espèces nouvelles ou critiques de *Rhodophyllus*. *In* Rev. Mycol. **19**: 3–46.
- (1974a). Etude de quelques Rhodophylles. *In* (Trav. mycol. déd. R. Kühner) Bull. mens. Soc. linn. Lyon **43** (No. spéc.): 365–387.
- (1974b). Essai d'une classification des Rhodophylles. *In* Bull. mens. Soc. linn. Lyon **43**: 325–332.
- (1978). Les fondements de la taxinomie des Rhodophylles et leur classification. *In* Beih. Nova Hedwigia **59**: 1–80. (Prepublication.)
- ROMAGNESI, H. & GILLES, G. (1979). Les Rhodophylles des forêts côtières du Gabon et de la Côte d'Ivoire, avec une introduction générale sur la taxinomie du genre. *In* Beih. Nova Hedwigia **59**: 1–649.
- SACCARDO, P. (1887). Sylloge Fungorum **5**. Patavii.
- SCHAEFFER, J. C. (1800). Fungorum Bavariae Icones. Ed. 2, 3. Erlangae.
- SCHROETER, J. (1889). *Hyporrhodius*. *In* Cohn, KryptogFl. Schles. **3** (1): 613–617. Breslau.
- SINGER, R. (1961). Type-studies on Basidiomycetes. X. *In* Persoonia **2**: 1–62.
- SMOTLACHA, F. (1945). *In* Čas. Česká Houby **23**: 67.
- WINDEN, P. van (1978). Dag, *Rhodophyllus pleopodius*! *In* Coolia **21**: 9–10.

## INDEX

New names are in bold-face type. Subdivisions of genera are indicated by the sign §, illustrations by an asterisk (\*)

## Agaricus

- § *Nolanea* 428, 431  
 acceptandus 438  
 ameides 473  
 asprellus 465  
 cetratus 446, 496  
 clandestinus 456, 526  
 conferendus 446, 449  
 dissidens 446, 449

- hirtipes 431, 438  
 icterinus 516, 517  
 infula 503  
 inutilis 512  
 junceus 465  
   var. *cuspidatus* 461  
 mammosus 445, 454, 526, 527  
   var. *tenuior* 445, 454, 527  
 pascuus 431, 449, 526, 527

*(Agaricus)*

- pleopodius 516, 517, 518  
 rubellus  
   subsp. *verecundus* 507  
 sericeus 478  
 solstitialis 505, 506  
 subpostumus 446, 449  
 verecundus 507  
 vinaceus 529  
**Arenicola** 431  
   *flavispora* 431  
**Claudopus**  
   *minutus* 468  
**Eccilia**  
   *bisporigera* 498  
   *minuta* 468  
**Entoloma** 465, 488, 494, 507, 508, 509, 521, 524, 525  
   § **Cheilocystidiata** 476, 510, 511, 513  
   § *Cosmoxonema* 453, 472, 478, 486  
   § *Eccilia* 513  
   § *Efibulatae* 495  
   § *Endochromonema* 494, 495, 502, 503, 509, 514, 525, 528  
   § *Entoloma* 428, 474, 480, 498, 507, 528  
   § **Fernandae** 428, 486, 488, 490, 492, 493, 494, 505  
   § **Fibulata** 463  
   § **Icterina** 514, 519  
   § **Infularia** 503  
   § *Leptonia* 519  
   § **Minuta** 453, 468  
   § **Nolanea** 427, 428, 429, 431, 438, 463, 474, 476, 488, 505, 526  
   § *Papillata* 453, 463, 492  
   § *Pouzaromyces* 428, 488, 524, 525, 526, 529  
   § **Staurospora** 445  
   § *Trichopilus* 513  
   § **Tristia** 508  
   § **Turfosa** 529  
   *acidophilum* 488, 489, 490, 492  
   **ambrosium** 520  
   *ameides* 428, 473, 474, 475\*, 476, 517  
   *argenteostriatum* 488, 490, 492  
   *botanicum* 446, 450  
   *brevipes* 527  
   *calthionis* 520  
   *cetratum* 428, 463, 494, 495, 496, 497, 498, 500\*, 501, 502, 514, 523, 525, 526, 528, 529  
   *chlorinosum* 506\*, 507  
   **chlorophyllum** 518, 519\*, 520  
   *cinerascens* 470  
   **clandestinum** 455, 456, 457\*, 513  
   **conferendum** 428, 445, 446, 449, 527  
     var. *conferendum* 447\*, 449  
     var. **pusillum** 447\*, 449, 450  
   *conicum* 510  
   *crassipes* 499  
   *cryptocystidiatum* 476, 511  
   *cuneatum* 498, 499, 500\*, 501, 523, 525, 528  
   *cuniculorum* 490, 491\*, 492  
   *cuspidatus* 461  
   **cuspidifer** 461, 462\*, 463, 526  
   *defibulatum* 491, 492  
   *farinogustus* 497, 498, 500\*, 528, 529  
   *farinolens* 459, 460  
   *fernandae* 486, 487, 488, 489, 490, 493  
     f. **eccilioides** 488, 491\*  
     f. *fernandae* 491\*  
   *foetidum* 525  
   **foetulentum** 522\*, 525  
   *fractum* 491\*, 492, 493  
   *globisporum* 464  
   **globulifer** 485, 513, 514, 515\*  
   *hirtipes* 438, 440, 445, 453, 527  
     var. *hirtipes* 438, 440, 441\*, 445, 523, 524  
     var. **sericoides** 441\*, 442, 482, 511  
   *hirtum* 529  
   *icterinum* 428, 508, 515, 516, 517, 518, 519\*, 520  
   *ifula* 503, 504, 505, 506\*, 507  
   *inodorum* 520, 521, 524  
   **inutile** 512\*, 513  
   *jubatum* 513  
   *juncinum* 445, 464, 465, 466, 467\*, 468, 470, 482, 527  
   *lanuginosipes* 499, 500\*, 501, 528  
   **leptopus** 442, 444\*, 445, 466, 524  
   *lucidum* 458, 459, 460\*, 461, 468  
   *mammosum* 526  
   *minutum* 458, 459, 468, 469, 471\*  
     var. *minutum* 468  
     var. *polymorphum* 470  
   *nidosorum* 507  
   *nitens* 465, 466, 467\*, 468, 470  
   *nivescens* 524, 527  
   *nothofagi* 446, 450  
   *occultopigmentatum* 501, 502  
   *ortonii* 459, 460\*, 461, 492  
   *pallescens* 499, 501, 502, 521, 522\*, 523, 525  
   *papillatum* 428, 454, 455, 456, 457\*, 466, 474, 478, 482, 527  
   *pascuum* 527  
   *politum* 498  
   *prismatospermum* 523, 524  
   *pseudotelamonia* 527  
   *psilopus* 488, 489, 490, 492  
   *rhombisporum* 449, 450, 451\*, 452, 505, 506, 524

*(Entoloma)*

- robiniae 528  
**sacchariolens** 474, 475\*, 476  
**sericeoides** 483, 484\*, 485  
**sericeonitens** 458, 459, 460\*, 461  
 sericeum 428, 442, 449, 466, 473, 474, 476, 478,  
 480, 482, 485, 502, 527, 528  
   var. **cinereo-opacum** 482, 483, 484\*  
   var. sericeum  
     f. **nolaniforme** 473, 480, 481\*, 482  
     f. sericeum 479, 481\*, 482  
**solstitiale** 504, 505, 506\*, 507  
 sordidulum 482  
**sphaerocystis** 460\*, 485, 486, 514, 522\*  
 staurosporium 446  
   var. pusillum 450  
 tenellum 463, 470, 471\*  
 tenuipes 442  
 tibicycstidiatum 471\*, 473, 476  
 triste 508, 509, 510  
 undulatosporium 509, 510  
 velenovskyi 510, 511  
   var. longicycstidiatum 511  
   var. velenovskyi 510, 512\*  
 ventricosum 502  
**verecundum** 494, 506\*, 507, 508  
 vernum 449, 476, 477\*, 478, 482, 527  
 versatilis 520  
 vinaceum 529  
   var. fumosipes 529  
   var. vinaceum 494, 508  
   var. violeipes 529  
 xanthocaulon 489, 491\*, 493, 494, 508  
 xylophilum 450  
 zonatum 464
- Hebeloma**  
 sacchariolens 475, 517
- Hyporrhodius** 428  
   § *Nolanea* 431  
   cetratus 496  
   icterinus 516  
   mammosus 526  
   pascuus 527
- Lanolea** 431  
**Latzinea** 431
- Leptonia** 453, 465, 504, 506  
   cinerascens 470  
   mamillata 454, 455  
   papillata 454, 455  
   pernitrosa 507  
   pilatii 504, 505  
   rhombispora 450, 452  
   solstitialis 505, 506
- Naucoria**  
 centunculus 520
- Nolanea** 431, 482, 505  
   § *Cosmeoxonema* 472  
   § *Efibulatae* 495  
   § *Endochromonema* 494  
   § *Fibulatae* 463  
   § *Icterinae*  
   § *Staurospori* 445  
   acceptanda 438  
   alba 524  
   araneosa 524  
   babingtonii 524  
   brassicolens 525  
   californica 495  
   cetrata  
     var. testacea 529  
   clandestina 454, 455, 456  
   conferenda 446  
   conica 510  
   crassipes 499  
   cucullata 476  
   cuneata 498  
   cuspidifer 461  
   depressa 470  
   dysthales 525  
   erophila 476  
   farinolens 459  
   fernandae 487  
   foetida 525  
   fracta 492  
   globispora 464, 465  
   hirta 526  
   hirtipes 438, 440  
   icterina 516, 517  
   infula 503  
   inodora 520  
   inutilis 512  
   junceae 465  
     var. cuspidata 461  
   juncina 464  
   limosella 526  
   lucida 458  
   majalis 438, 521  
   mammosa 438, 440, 454, 526  
     var. papillata 455  
   minuta 468  
   nitens 466  
   palescens 521, 523  
     var. procera 521  
   papillata 445, 454  
   pascua 427, 438, 446, 527  
     var. palescens 521

*(Nolanea)*

- pleopodia 508, 516  
 proletaria 446, 464, 465  
 pusilla 450  
 radiata 482, 528  
 rickenii 446  
 robiniae 528  
 sericea 478  
 sericeonitens 459, 468  
 setulosa 529  
 solstitialis 505  
 staurospora 428, 445, 446  
   var. *farinacea* 446, 449  
   var. *incrustedata* 446, 449  
 strigosissima 529  
 subpostuma 446  
 tenella 470  
 tenuipes 440, 442  
 testacea 497, 529  
 tristis 509  
 verecunda 507  
 verna 476  
 versatilis 529  
 vinacea 529  
 xylophila 450  
 zonata 464, 465
- Pouzarella**  
 ambrosia 520
- Rhodophyllus** 428  
   § *Cosmeoxonema* 472  
   § *Endochromonema* 494  
   § *Infularii* 503  
   § *Lecithiphori* 513  
   § *Mammosi* 438  
   § *Minuti* 468  
   § *Nolanea* 431  
   § *Papillati* 453  
   § *Paramammosi* 511, 513  
   § *Staurospori* 445  
   § *Turfosi* 529  
 ambrosius 520  
 ameides 473  
 cetratus 496  
 cucullatus 476, 477  
 cuspidatus 461  
 cuspidifer 461  
 elaphinus  
   var. *radiatus* 528  
 farinolens 459  
 fernandae 487, 488
- fulviceps 497, 522\*, 525, 526, 529  
 hebes 445  
 hirtipes 438  
 icterinus 508  
   f. *gracillimus* 516  
 infula 503  
 inutilis 512  
 junceus 464, 465  
 juncinus 464, 465  
 lucidus 458  
 mammosus 438, 445, 454, 526, 527  
   var. *obsoletus* 442, 445  
   var. *sericoides* 442  
   var. *tenuis* 443  
 minutus 468  
 nitens 466, 468  
 nitriolens 507  
 papillatus 454, 455  
 pascuus 527  
 pleopodius 516, 518  
 prismatospermus 523  
 pusillus 450  
 putus 497, 522\*, 528, 529  
 radiatus 482, 528  
 rhombisporus 450  
 rickenii 446  
   var. *obscurior* 446  
   var. *subrugosus* 446, 449  
 sacchariolenis 474  
 sericeoides 483  
 sericeus 478  
   var. *nolaniformis* 480  
   var. *typicus* 479  
 solstitialis 505  
 sordidulus 528  
 staurosporus 446  
   subsp. *rickenii* 446  
   var. *obscurior* 446, 449  
   var. *platyphyllus* 446, 449  
   var. *rickenii* 446, 449  
   var. *subrugosus* 446, 449  
   var. *typicus* 446  
 tenellus 470  
 verecundus 507  
 vernus 476  
 vinaceus 529  
 xylophilus 449
- Telamonina** 527  
 brevipex 527

NOTES AND BRIEF ARTICLES

ON MUCRONELLA RICKII AND PTERULA GRACILIS

W. JÜLICH

*Rijksherbarium, Leiden*

Among the minute clavarioid fungi, my attention was drawn to *Ceratellopsis* and the tiny *Pterula gracilis* of which the last one has been collected several times in the Netherlands. *Pterula gracilis* is a species not always easily recognized as a *Pterula* and more often identified as a *Typhula*, although the hyphae are too thick-walled for that genus.

*Typhula*, inclusive of *Cnazonaria*, *Pistillaria* and several other closely related genera, is characterized by a monomitic hyphal system, and so is the genus *Ceratellopsis*, which differs from *Typhula* mainly in the distinct sterile, apex of the basidiocarp and the absence of sclerotia.

*Pterula*, on the other hand, is characterized by a dimitic hyphal system with unbranched, hyaline to pale yellowish-brown skeletal hyphae. The latter are rather obvious in the large, much-branched, dark coloured tropical species, but are less easily observed in the small and unbranched *Pterula gracilis*. This not because there are so few of them, in fact they constitute the main part of the basidiocarp, but because they are not very thick-walled, densely packed and longitudinally arranged, while they show several secondary septa. A closer study of *P. gracilis* reveals without any doubt that the conspicuous hyphae are all skeletal hyphae which are closely interwoven with a few, mostly thin-walled and easily collapsed generative hyphae bearing clamp-connexions.

After having studied the Netherlands' material of *P. gracilis*, I came across the description of *Ceratellopsis rickii* which, judged by the description, seemed to be a very similar species.

*Ceratellopsis rickii* has been described by Oudemans (as *Mucronella rickii*) in 1902 and is based on a specimen collected by Rick on *Asparagus* in the Netherlands. The basidiocarp was described as measuring 3-5 mm  $\times$  100-125  $\mu$ m, and as having ellipsoid spores (7-9.3  $\times$  3.5-4.7  $\mu$ m) and probably monosporous basidia.

Later Donk (1933) gave a more detailed description of the species and placed it in *Cnazonaria*, a genus closely allied to or probably identical with *Typhula*. He found the basidiocarps to be fertile except for the sterile apex, the hyphae slightly thick-walled and with rare septa lacking clamps, the basidia two-spored and the spores slightly larger than given by Oudemans, viz. 9.5-10-(12)  $\times$  4.5-5.5  $\mu$ m. The character 'sterile apex' leads to the genus *Ceratellopsis* and it was Corner (1950) who transferred the species to that genus. In a recent key to the Netherlands' species of clavarioid fungi, Maas Geesteranus (1976) furnished a description of that species under the genus *Ceratellopsis*.

The holotype, restudied only by Donk, is preserved in Leiden and has been re-examined by me. It consists of several basidiocarps arranged singly or in small groups, without a subiculum, acicular or narrowly subulate, straight or in most cases strongly curved, cream-coloured to ochraceous (white when fresh), gradually tapering towards the apex, without a distinct sterile apex in mature specimens (as in *Ceratellopsis*), stipe indistinct under a hand lens (but because of the absence of basidia very distinct under a microscope), the whole basidiocarp glabrous, 1–3 mm long and 50–120  $\mu\text{m}$  thick. Hyphal system dimitic with skeletal hyphae. Generative hyphae difficult to observe, cylindrical, not inflated, thin- to slightly thick-walled (0.2–0.4  $\mu\text{m}$ ), 3–3.5  $\mu\text{m}$  wide, with clamps. Skeletal hyphae hyaline to very pale yellowish-brownish, 2.5–4  $\mu\text{m}$  wide, with thick walls (0.5–1  $\mu\text{m}$ ) but never solid, cylindrical, not branched. Leptocystidia few or absent, clavate to irregularly subulate, 30–40  $\times$  5.5–7  $\mu\text{m}$ , slightly projecting (10–20  $\mu\text{m}$ ). Basidia 22–28  $\times$  6–8  $\mu\text{m}$ , always two-spored. Spores hyaline, ellipsoid, thin-walled, smooth, 10–12  $\times$  4.8–5.6  $\mu\text{m}$ , with distinct apiculus, not amyloid.




Fig. 1. *Mucronella rickii* Oud. — Holotype: spores, basidia, generative and skeletal hyphae.

Holotype: Netherlands, prov. Limburg, 'Valkenburg (L.) Majo 1901. In caulibus herbarum variarum praesertim Asparagi off. *J. Rick*' (L.: Herb. Oudemans).

The description leaves no doubt that this is typical *Pterula gracilis*, which among the European *Pterulas* is easily recognized because of its unbranched basidiocarps lacking a basal sclerotium, and its constantly two-spored basidia. The only similar species is *Pterula caricis-pendula* Corner which differs in having larger, normally four-spored basidia and larger spores.

#### PTERULA GRACILIS (Desm. & Berk. apud Berk.) Corner

- Typhula gracilis* Desm. & Berk. apud Berk. in Ann. Mag. nat. Hist. 1: 202. pl. 7, fig. 1. 1838. — *Clavaria gracilis* (Desm. & Berk. apud Berk.) P. Karst. in Bidr. Känn. Finl. Nat. Folk 37: 181. 1882; non ~ Pers. ex Fr. 1821; non ~ Bolt. ex Purt. 1821. — *Pistillaria gracilis* (Desm. & Berk. apud Berk.) Pat., Tab. anal. Fung. 2: 30. 1887. — *Ceratella gracilis* (Desm. & Berk. apud Berk.) Pat. in J. de Bot. 3: 36. 1889. — *Hirsutella gracilis* (Desm. & Berk. apud Berk.) Pat. Essai taxon, Hym.: 50. 1900. — *Pterula gracilis* (Desm. & Berk. apud Berk.) Corner, Monogr. *Clavaria*: 514. 1950.
- Clavaria aculina* Quél. in Compt.-Rend. Ass. franç. Avanç. Sci. 9: 670, pl. 8, fig. 11. 1881. — *Pistillaria aculina* (Quél.) Pat., Tab. anal. Fung.: 29. 1887. — *Ceratella aculina* (Quél.) Pat., Hym. Eur.: 157. 1887. — *Cnazonaria aculina* (Quél.) Donk in Meded. Nederl. mycol. Ver. 22: 97. 1933.
- Typhula brunaudii* Quél. in Compt.-Rend. Ass. franç. Avanç. Sci. 13: 283. 1884.
- Ceratella ferryi* Quél. & Fantrey apud Roum. in Revue mycol. 15 (57): 15. 1893. — *Pistillaria ferryi* (Quél. & Fantrey apud Roum.) Sacc., Syll. Fung. 11: 141. 1895.
- Mucronella rickii* Oud. in Beih. bot. Zbl. 11: 525. 1902; in Ned. kruidk. Arch., ser. III, 2: 667. 1902. — *Cnazonaria rickii* (Oud.) Donk in Meded. Nederl. mycol. Ver. 22: 99. 1933. — *Ceratellopsis rickii* (Oud.) Corner, Monogr. *Clavaria*: 205. 1950.
- Pistillaria equiseticola* Boud. in Bull. Soc. mycol. France 33: 13, pl. 4, fig. 5. 1917. — *Pistillaria aculina* subsp. *equiseticola* (Boud.) Bourd. & Galz., Hym. France: 138. 1928. — *Ceratellopsis equiseticola* (Boud.) Corner, Monogr. *Clavaria*: 204. 1950.
- Pistillaria aculina* subsp. *acicula* Bourd. & Galz., Hym. France: 139. 1928.
- Pistillaria aculina* subsp. *juncicola* Bourd. & Galz., Hym. France: 138. 1928.

Basidiocarp single or often more or less gregarious, seldom slightly fasciculate, normally without a mycelial patch on the substrate, erect, acicular to narrowly subulate, attenuate towards the obtuse or slightly filiform apex, often fertile at the top, 1–5–8 mm high and 80–250  $\mu\text{m}$  thick, white when fresh, cream-coloured to ochraceous when dry, stipe indistinct (under a hand lens), visible under the microscope as a sterile basal zone 100–300  $\times$  30–60  $\mu\text{m}$ .

Hyphal system dimittic with skeletal hyphae. Generative hyphae hyaline, cylindrical in the trama, often torulose and short-celled in the thin subhymenium where they are densely arranged, in the trama closely interwoven with the more abundant skeletal hyphae, branching from or near the clamps, thin- to slightly thick-walled (0.2–0.4  $\mu\text{m}$ ), 2–4(–5)  $\mu\text{m}$  wide, with smooth surface; clamps at all primary septa; secondary septa not rare; contents homogeneous or slightly guttulate. Skeletal hyphae pale yellowish brown, often becoming hyaline in KOH but remaining slightly coloured in lactophenol and Melzer's, cylindrical, parallel and densely arranged throughout the basidiocarp, unbranched, somewhat thick-walled (0.4–0.8  $\mu\text{m}$ ), but never solid, 2–3  $\mu\text{m}$  wide, with smooth surface; contents homogeneous. Cystidia (leptocystidia) lacking or rare, of hymenial origin, hyaline, subulate or with slightly moniliform apex, 25–50  $\times$  6–8  $\mu\text{m}$ , thin-walled, smooth, with basal clamp, not or only slightly projecting. Basidia hyaline, slightly suburniform to clavate when mature, with or without a lateral outgrowth at the base, somewhat stalked or sessile; young basidia narrowly clavate, 25–40  $\times$  6–8  $\mu\text{m}$ , thin-walled, smooth, always with basal clamp, with homogeneous contents, and with two subulate sterigmata. Spores




Fig. 2. *Pterula gracilis*. — Bas 2446a: spores, germinating spores, basidia, generative and skeletal hyphae.



hyaline, ellipsoid, thin-walled, smooth,  $10-14 \times 5.5-6.5 \mu\text{m}$ , with distinct apiculus, with contents homogeneous or slightly guttulate, neither amyloid, nor dextrinoid or cyanophilous.

**SUBSTRATE.**—Saprophytic on old stems and leaves of e.g. *Asparagus*, *Carex*, *Cladium*, *Equisetum*, *Eupatorium*, *Iris*, *Juncus*, *Typha*, also on straw in greenhouses, mostly found at wet or swampy places.

**CHEMICAL REACTIONS.**—No part of the basidiocarp is amyloid, dextrinoid, or cyanophilous.

**GERMINATION.**—Spores often germinating with only one cylindrical germ-tube opposite to the apiculus, but in some cases with two to three from other places inclusive of the apiculus area. Germ-tube mostly unbranched for a considerable length, occasionally branched already near the spore surface.

**DISTRIBUTION.**—Recorded from Czechoslovakia, France, Great Britain and the Netherlands.

**MATERIAL STUDIED.**—NETHERLANDS: prov. Noord-Brabant, Deurne, Peel W. of Hele-naeven, 23.IX.1961, *C. Bas 2445a* (L); prov. Utrecht, Kortenhoeft, 14.III.1968, *J. Daams* (L); prov. Noord-Holland, 's-Graveland, 24.X.1966 and 7.II.1968, *J. Daams* (L).

#### REFERENCES

- BERTHIER, J. (1976). Monographie des *Typhula* Fr., *Pistillaria* Fr. et genres voisins. In Bull. Soc. linn. Lyon 45 (Num. Spéc.): 213 pp.
- CORNER, E. J. H. (1950). A monograph of *Clavaria* and allied genera. Oxford, 740 pp.
- DONK, M. A. (1933). Revision der niederländischen Homobasidiomycetae — Aphyllophoraceae II. In Meded. Nederl. mycol. Ver. 22: 278 pp.
- MAAS GEESTERANUS, R. A. (1976). De clavarioïde fungi. In Wetensch. Meded. Kon. Ned. nat. Ver. 113: 92 pp.

## A NEW SPECIES OF PARAPHELARIA (AURICULARIALES) FROM BORNEO.

W. JÜLICH

*Rijksherbarium, Leiden*

### *Paraphelaria borneensis* Jülich, *spec. nov.*

Fructificatio erecta, cylindracea, apice simplici vel plus minusve ramoso, usque ad  $15 \times 1-2 \text{ mm}$ , brunnea. Systema hypharum monomiticum. Hyphae luteae vel brunneae, leviter tenui-tunicatae,  $2.5-4.5 \mu\text{m}$  in diam., efibulatae. Cystidia desunt. Basidia (phragmobasidia) hyalina,  $30-36 \times 4-5 \mu\text{m}$ , tetraspora. Sporae hyalinae, plus minusve allantoideae, tenui-tunicatae, laeves,  $12-14 \times 4-4.5 \mu\text{m}$ , inamyloideae.

Typus: 'W. Jülich 78/2541, Borneo, Sarawak, Gunong Mulu National Park, 19 III 1978 (L).

Basidiocarp annual, erect, cylindrical, the upper part simple or slightly branched, up to 15 mm high and 1-2 mm thick; consistency tough coriaceous; context homogeneous. Hymenial surface even, dark brown when dry. Hyphal system monomitic. Hyphae yellowish brown when young, later darker brown, cylindrical, densely packed, somewhat thick-walled and  $2.5-4.5 \mu\text{m}$  wide, with smooth surface; clamps absent from all septa. Cystidia lacking. Basidia (phragmobasidia)

scattered in the context, hyaline, sinuous-cylindrical when mature,  $30\text{--}36 \times 4\text{--}5 \mu\text{m}$ , with four large sterigmata; a basal calmp absent. Spores hyaline, curved-cylindrical to allantoid, thin-walled, smooth,  $12\text{--}14 \times 4\text{--}4.5 \mu\text{m}$ , with rather small apiculus and homogeneous or somewhat guttulate contents, not amyloid. Substrate: on rotten wood (?) on the ground.

Type specimen: Borneo, Sarawak, Gunong Mulu National Park, around Camp 1, 19. III. 1978, W. Jülich 78/2541 (L).

The species differs from *Paraphelaria amboinensis* (Lév.) Corner — the only other species known from Southeast Asia — in its much smaller, only slightly branched and distinctly darker coloured basidiocarps and the different sizes of basidia and spores.

#### REFERENCES

- CORNER, E. J. H. (1966). *Paraphelaria*, a new genus of Auriculariaceae (Basidiomycetes). In *Persoonia* 4: 345–350.

## AN UNUSUAL FORM OF TRECHISPORA VAGA

W. JÜLICH

*Rijksherbarium, Leiden*

During an excursion of the Dutch Mycological Society to the Bois de Resteigne (prov. Namur, Belgium) in 1977, a curious lignicolous clavate fungus was found growing out of a resupinate basidiocarp. Since the clavate body had the same colour as the resupinate part, a possible relationship between the two parts was supposed but seemed highly improbable since no such large structures connected with a thin resupinate basidiocarp are known up to now. The resupinate part showed a yellowish-brownish colour, stained slightly reddish in a diluted solution of KOH and was easily identified as *Trechispora vaga* (Fr.) Liberta, owing to its monomitic hyphal system, inflated and clamped hyphae, small four-spored basidia which often develop a lateral, basal outgrowth (pleurobasidia), and small, hyaline, inamyloid spores densely covered with warts.

The clavate body was hanging out of the centre of the resupinate part and was up to 12 cm long and 9 cm wide. The point of attachment was rather narrow, ranging from a few millimeters in the small body to a few centimeters in the larger one. The surface was yellowish and showed at the marginal parts small, cup-shaped depressions caused by guttation drops. A longitudinal section reveals a yellowish context with small, darker spots (much more pronounced in dried condition) and a distinct concentric zonation, probably demonstrating the successive growth of the fungus. The whole clavate body is sterile and shows no traces of hymenial formation or conidium production. The hyphal structure is monomitic and shows the same cylindrical, partly inflated hyphae which deviate in no way from the hyphae of the underlying perfect state. There is no doubt that the sterile, clavate structures belong to *Trechispora vaga*, with the resupinate basidiocarp of which they were intimately connected.

There are some other short sterile outgrowths known from different resupinate species, mostly caused by a cover of the fungus with leaves etc., but these large clavate bodies growing out of a




Fig. 1. Upper surface of two sterile, clavate bodies growing out of the resupinate, fertile part of *Trechispora vaga* (Fr.). Half natural size.

Fig. 2. Longitudinal section through the large sterile body. Half natural size.

thin resupinate fungus and hanging freely down are a 'novum' for Aphyllophorales. There are no suggestions about what could have caused such a structure, since the underlying resupinate part seemed to be excellently developed and showed no signs of deformation or abnormality. Another fact makes an explanation equally difficult, viz. the presence of other resupinate basidiocarps of the same species in the immediate vicinity, supposedly growing under the same ecological conditions.

This collection shows that resupinate species may be able to produce quite differently shaped large structures and it would be very interesting to look for similar structures in other species, in the hope that eventually the causal factors may be found.

## PHOMA HETEROMORPHOSPORA NOM. NOV.

H. A. VAN DER AA

*Centraalbureau voor Schimmelcultures, Baarn.*

H. A. VAN KESTEREN

*Plantenziektenkundige Dienst, Wageningen*

Van der Aa & van Kesteren (1979) redescribed a *Phoma* species commonly occurring on species of *Chenopodium* in Europe, as *Phoma variospora* v. d. Aa & v. Kest. After the appearance of this publication, in November 1979, descriptions of some new species of *Phoma* came to their attention amongst which was also *Phoma variospora* Shreemali (1978, issued in July 1979) based on a different fungus. According to the rules of priority *Phoma variospora* v. d. Aa & v. Kest. has to be renamed. The following name change is proposed:

***Phoma heteromorphospora* v. d. Aa & v. Kest., nom. nov.**

*Phyllosticta chenopodii* Westend. in Bull. Acad. r. Belg. Cl. Sci. II. 2: 567. 1857. — *Phoma variospora* v. d. Aa & v. Kest. in Persoonia 10: 268. 1979. — non *Phoma variospora* Shreemali in Indian J. Mycol. Pl. Path. 8: 221. '1978' (in 1979).

For further synonyms and discussion see van der Aa & van Kesteren (1979).

## REFERENCES

- AA, H. A. VAN DER & KESTEREN, H. A. VAN (1979). Some pycnidial fungi occurring on *Atriplex* and *Chenopodium*. In *Persoonia* 10: 267–276.
- SHREEMALI, J. L. ('1978', in 1979). Two new species of *Phoma* from India. In *Indian J. Mycol. Pl. Path.* 8: 220–221.

VALIDATION OF SYMBIOTAPHRINA  
(IMPERFECT YEASTS)

W. GAMS &amp; J. A. VON ARX

*Centraalbureau voor Schimmelcultures, Baarn*

***Symbiotaphrina* Kühlwein & Jurzitza ex W. Gams & v. Arx, gen. nov.**

*Symbiotaphrina* Kühlwein & Jurzitza in *Arch. Mikrobiol.* 40: 258. 1961 (nom. inval., Art. 36).

Crescit in mycetomatibus intestinalibus coleopterorum Anobiidarum. Coloniae zymoideae, restrictae, cremeae vel luteae vel rubrae, mucidae; cellulae dacryoideae vel clavatae vel pyriformes, apicem attenuatum versus enteroblastic successionem basipetali gemmantes.

Species typica: *Symbiotaphrina buchneri* Graebner ex W. Gams & v. Arx

Isolated from intestinal mycetomata of Anobiid beetles. Colonies yeast-like, growing restrictedly, cream, yellow to red, mucoid; cells dacryoid, clavate or pyriform, budding enteroblastically in basipetal succession at the attenuated end. The colonies resemble *Rhodotorula*. A full description with physiological data is given in the papers by Kühlwein & Jurzitza (1961) and Jurzitza (1964). *Symbiotaphrina* was placed there in the family Taphrinaceae but differs from *Taphrina* by its symbiotic mode of life.

***Symbiotaphrina buchneri* Graebner ex W. Gams & v. Arx, spec. nov.**

*Torulopsis buchneri* Graebner in Z. Morphol. Oekol. Tiere **42**: 513. 1954 (nom. inval., Art. 36) — *Symbiotaphrina buchneri* (Graebner) Kühlwein & Jurzitza in Arch. Mikrobiol. **40**: 258. 1961 (nom. inval., Art. 43).

Crescit in mycetomatibus intestinalibus *Sitodrepae paniceae* L. (Coleoptera, Anobiidae). Coloniae sub luce vel obscuritate rubidae, mucidae, leves, nitentes. In musto cerevisiae cellulae dacryoideae ad clavatae,  $5-8 \times 2-6 \mu\text{m}$ , haud aggregatae, in parte angustata gemmantes. Biotinum et aneurinum et acidum asparaginicum et glutaminicum necessaria.

Typus: CBS 420.63, isolatus e mycetomate *Sitodrepae paniceae* in pulvere fructus *Capsici* in Instituto botanico Karlsruhe in Germania, a G. Jurzitza.

Observed in and isolated from intestinal mycetomata of *Sitodrepa paniceae* L. (Anobiidae). Colonies in light and in darkness reddish, mucoid, smooth, shiny. In beer wort cells dacryoid to clavate,  $5-8 \times 2-6 \mu\text{m}$ , not forming aggregations, budding from the attenuated pole. Biotin and aneurin are necessary for growth, aspartic and glutamic acids are partially required.

***Symbiotaphrina kochii* Jurzitza ex W. Gams & v. Arx, spec. nov.**

*Symbiotaphrina kochii* Jurzitza in Arch. Mikrobiol. **49**: 338. 1964 (nom. inval., Art. 36, 43).

Crescit in mycetomatibus intestinalibus *Lasiodermais serricornis* F. (Coleoptera, Anobiidae). Coloniae sub luce rubrae, obscuritate albae, mucidae, leves, nitentes. Cellulae in musto cerevisiae rotundatae ad ovaes, raro pyriformes,  $9-11 \times 4-7 \mu\text{m}$ , haud aggregatae; nonnumquam ad  $18-42 \times 6.5-13 \mu\text{m}$  inflatae et uni- ad triseptatae. Vitamina haud necessaria.

Typus: CBS 250.77, isolatus ex mycetomate *Lasiodermais serricornis* cigarros infestantis in Germania a G. Jurzitza.

Observed in and isolated from intestinal mycetomata of *Lasioderma serricorne* F. (Anobiidae). Colonies in light red, in darkness white, mucoid, smooth, shiny. Cells in beer wort rounded to oval, rarely pyriform,  $9-11 \times 4-7 \mu\text{m}$ , not forming aggregations; sometimes swollen to  $18-42 \times 6.5-13 \mu\text{m}$  and one- to three-septate. Vitamines are not required for growth.

REFERENCES

- JURZITZA, G. (1964). Studien an der Symbiose der Anobiiden. II. Physiologische Studien am Symbionten von *Lasioderma serricorne* F. In Arch. Mikrobiol. **49**: 331-340.  
KÜHLWEIN, H. & JURZITZA, G. (1961). Studien an der Symbiose der Anobiiden. I. Die Kultur des Symbionten von *Sitodrepa panicea* L. In Arch. Mikrobiol. **40**: 247-260.

### BOOKS RECEIVED BY THE RIJKSHERBARIUM LIBRARY

M. V. LOCQUIN, *Flore mycologique 3 and 4, Cortinariales-A*. (Ed. J.-F. Guyot, Paris, 1977).  
Respect. 160 pp. and 156 pp., including 75 coloured Plates. Price: Frs. 160.-, bound Frs. 200.-.

These two volumes in which *Hebeloma* s.l. and the viscid species of *Cortinarius* s.l. are treated, are part of a work on higher fungi (with emphasis on Agaricales s.l.) of the northern hemisphere, planned in 24 volumes.

Volume three contains an introductory text of about 50 pages on technics, terminology, classification schemes, etc., followed by about 80 pages with tabulated keys to the species of the many genera into which the classical genera *Hebeloma* and *Cortinarius* are subdivided and very brief descriptions of the illustrated species of '*Hebeloma*'. The descriptions of the '*Cortinarius*' species will follow in volume five. Microscopic characters are hardly used in the keys and only very briefly mentioned in the descriptions. A few pages are added with Latin diagnoses of '*Hebeloma*' and '*Cortinarius*' species in Fries's '*Hymenomyces europaei*'.

In volume four about 180 species of '*Hebeloma*' and '*Cortinarius*' are illustrated in colour on 75 plates. Facing each plate is a black and white copy on which the most important characters and data are indicated in symbols. The coloured illustrations are redrawn copies of plates of prominent mycologists, but the origin of the plates is not indicated.

C. H. PECK, *Annual reports of the state botanist 1868-1912*. (Reprint edition, Boerhaave Press, p.o. box 1051, 2302 BB Leiden; L. Vogelenzang Ed.) Vol. 1, reports over the period 1868-1877, about 650 pp., 10 col. pls. on microfiche, biographical and bibliographical introduction (25 pp.) by R. H. Petersen and portrait of Peck added. Price Hfl. 320,- (excl. of postage).

The first volume of a set of five, containing reprints of Peck's important yearly reports of the state botanist of New York in which the great majority of Peck's about 2700 new taxa of fungi are published. Great trouble is taken to establish the correct dates of publication.

## INDEX

New names are in **bold-face** type. Subdivisions of genera are indicated by the sign §, illustrations by an asterisk (\*) added to the page number.

N. B. — The paper by Noordeloos on *Entoloma* subgenus *Nolanea* in Part 4 is separately indexed on pp. 531–534.

- Acanthobasidium* 321–323; **delicatum** 335  
*Acanthostereum* *acanthophysalum* 320; *illudens* 320; *peculiare* 320; *reflexulum* 320; *spectabile* 320  
*Achaetomium* 144, 146; *globosum* 146; **thielavioides** 144, 145\*  
*Acia* *subfascicularis* 327  
*Aciculariella* *foliicola* 70  
*Acremonium* 78, 97–100, 102, 111, 143; *diversisporum* 97, 107; *griseoviride* 102; *humicola* 102; *longisporum* 97; *luzulae* 108\*, 111; *meliolae* 60; *murorum* 98; *persicinum* 97, 98; *sagenomatis* 100, 105; *striatisporum* 100, 101; *strictum* 67, 78; *verticillatum* 47  
*Acrocylindrium* *cordae* 72; *roseum* 78  
*Acrodontium* 64; *album* 64  
*Acrophialophora* 101  
*Aculeatostereum* 321; *australe* 321; *insignitum* 321  
***Adustomyces*** 325, 326; ***lusitanicus*** 326  
*Agaricus* 373, 407; *babingtonii* 240, 241; *brunnescens* 44; *clavularis* 280; *corticola* 280, 281; *corticalis* 280; *dysthales* 215, 241; *fulvostrigosus* 236; *fumosellus* 241; *fuscoalbus* 364; *geophyllus* 157; *limacinus* 364, 365, 370, 372, 373; *olivaceoalbus* 364, 373, 376, 380; *parkensis* 262; *phylogena* 279; *piceus* 254; *umbellifera* 280; *versatilis* 229, 230; *vinaceum* 298  
*Agmocybe* 157  
*Agrocybe* *aegerita* 408; *praecox* 86, 92, 94, 96  
*Aleuria* 125, 126, 128; *aurantia* 119\*, 128  
*Aleurodiscus* 321, 323; § *Acanthophysium* 311, 321, 322; § *Aleurobolus* 311, 321, 322; § *Aleurodiscus* 311, 321, 322; *apricans* 322; *delicatus* 321, 335; *mirabilis* 321; *penicillatus* 321  
*Aleyrodes* 54  
*Allescheria* *boydii* 25  
*Alnicola* *melinoides* 410  
*Alnus* 2, 218, 222; *glutinosa* 218; *viridis* 218  
*Alternaria* 275  
*Alytosporium* *roseum* 78  
*Amanita* 1–3, 12, 301, 307, 308, 386, 410; § *Amanita* 3, 9, 12, 14; § *Lepidella* 1–3; § *Vaginatae* 3, 14, 17, 20, 22; § *Validae* 5, 7; *antillana* 1, 2, 22; *austro-olivacea* 1; ***campinaranae*** 1–4, 5\*; *cinctipes* 22; *coacta* 1, 2, 4, 15, 16\*–18, 20; *craseoderma* 1–3, 17, 20, 21\*, 22; ***crebresulcata*** 1–3, 17, 18, 19\*, 20; *diemii* 1; *farinosa* 9, 12; *fritillaria* 7; *gayana* 1; *humboldtii* 1; *inaurata* 22; ***lanivolvula*** 1–3, 12, 13\*–15\*, 17; *morrisii* 7; *muscaria* 1; *obsita* 9, 12; *phaea* 1–3, 6\*, 7; *phalloides* 1; *pseudospretta* 1; *spissaceae* 7; *subvaginata* 9, 12; ***sulcatissima*** 1–3, 9, 12; *tristis* 7; *umbrinella* 1; *ushuaiensis* 1; ***xerochybe*** 1–3, 7, 8\*, 9, 12  
*Amanitopsis* *plumbea* 1, 2, 17, 18, 20  
Anisoptera 191  
*Anthracobia* 126; *maurilabra* 119\*; *melaloma* 128  
*Aphanosaceus* 101  
***Aphanobasidium*** 326; ***allantosporum*** 326; ***filicinum*** 326; ***gaspesicum*** 326; ***grisellum*** 326; ***lloydii*** 326; ***ralla*** 326; ***subnitens*** 326  
*Aphylophorales* 335  
*Aporoethelavia* 30  
*Arachis* *hypogaea* 51  
*Arbutus* 218; *unedo* 318  
*Arenicola* 245  
*Arthrographis* 23, 28; *kalrai* 25; *langeronii* 23, 25  
*Arthyrium* 330  
*Ascobolus* 113, 126; *crenulatus* 128; *furfuraceus* 117\*; *immersus* 115; *sacchariferus* 117; *stercorarius* 115; *viridulus* 115  
*Ascochyta* 267, 268, 270, 275; § ***Chaetodiopodia*** 270; *atriplicis* 271, 273, 275; *caulina* 267–271, 272\*–276; *chenopodii* 271, 273, 275; *chenopodiicola* 271; *hyalospora* 275; *nebulosa* 271, 273, 275  
*Ascochyttula* 267; *atriplicis* 271  
*Ascodesmis* 114, 126; *microscopica* 118\*; *nigricans* 115, 118\*; *sphaerosporus* 115  
*Ascophanus* *coemansii* 122, 125\*  
*Ascozonus* 113, 126; *woolhopensis* 122, 125\*, 128  
*Asparagus* 535, 539  
*Aspergillus* 99, 101, 309, 328; *niger* 97, 98  
*Asperisporium* 58

- Aspidiotus 54  
 Asterotheca 87; lycoperdoides 85, 86, 94  
 Astrosporina 157, 158, 179, 181, 185, 187, 193,  
 196, 203, 204; § Calosporae 164, 172; § Corti-  
 natae 175; § Marginatae 191; § Petiginosae 163;  
 § Rubellae 162; **aberrans** 161, 187, 188\*; **albovic-  
 cida** 159, 191, 192\*; **alienospora** 161, 173, 174\*;  
**angustifolia** 159, 195, 196\*; asterospora 159,  
 190, 202\*; avellana 159, 200, 201\*; **bresadolae**  
 159, 162\*; bucknallii 215; brevispora 182; ceras-  
 phora 182; **cingulatipes** 161, 177, 178\*; com-  
 mixta 191; **corneri** 159, 184, 185\*; discissa 189\*,  
 204; **echinosimilis** 160, 168, 169\*, 173; emergens  
 203; exigua 189\*, 203; **gemina** 158, 166, 167\*;  
**granulosiceps** 160, 191, 193\*; **hydrocymbiformis**  
 160, 170, 171\*; umbricata 157, 160, 190\*; **in-  
 cognita** 161, 179, 180\*, 181, 184; lanuginosa  
 177, 182; lasseri 173; **lasseroides** 161, 172\*, 173;  
 longicystis 187; lutea 159, 176, 197\*; **luteifolia**  
 160, 176\*, 177; **magnifica** 160, 185, 186\*, 187;  
**mediocris** 158, 198\*; napipes 199, 204; **neoum-  
 brina** 158, 160, 161, 182, 183\*; **olivaceonigra**  
 160, 194\*, 195; olivaceonigra f. **volvata** 158, 159,  
 160, 195; **pahangi** 160, 168, 170\*; **papillata** 161,  
 183, 184\*; paracerasphora 161, 181\*, 182, 184;  
 petchii 159, 164, 165\*, 166; petiginosa 164;  
**punctatosquamosa** 161, 178, 179\*, 184; pusil-  
 lina 160, 163, 164\*; rennyi 191; scissa 199;  
**squarrosolutea** 158, 175\*, 176; straminea 191;  
 subclavata 182; titibuensis 159, 199, 200\*; um-  
 brina 182, 191  
 Athelia 79, 149, 150; andina 147, 150\*, 151;  
 epiphylla 149, 150\*, 151; phycophila 149, 150\*,  
 151; **poeltii** 149, 150\*, 151  
 Athelopsis 331  
 Atriplex 267, 273, 274; hastata 273, 274; hortensis  
 274; lacinata 274; littoralis 273; nitens 273, 274;  
 oblongifolium 274; patula 274; spec. 274  
**Australohydnum 137; griseo-fuscescens 138**  
  
 Basifimbria peruviana 34  
 Basipetospora rubra 34  
 Beauveria 53; alba 53  
 Blastotrichum araneanum 57; puccinioides 37  
 Bolbitius vitellinus 408  
 Boothiella tetraspora 146  
 Botryobasidium 328; alutaceum 334; **cystidium**  
**334; scabridum 334**  
 Botryohypochnus 328; **alutaceum 334**  
 Botrytis agaricina 67; dendroides 41; macrospora  
 41; varia 46; variosperma 46  
 Boudiera 126; echinulata 117\*  
  
 Briosia 26  
 Burseraceae 11  
 Byssochlamys 101  
  
 Calcarisporium 55, 77; arbuscula 72; **ovalisporum**  
**77**  
 Calocybe 83, 87; gambosa 86, 94; ionides 86, 94  
 Calocectria 101  
 Calthion palustris 288, 289, 295  
 Carex 539  
 Castanopsis 158–161, 163, 166, 168, 173, 178–180,  
 182, 183, 193–195, 197, 199  
 Catenularia 101  
 Cephalotheca pulcherrima 28  
**Ceraceohydnum 138; brunneum 137, 138**  
 Ceratella aculina 537; ferryi 537; gracilis 537  
 Ceratellopsis 535, 536; equiseticola 537; rickii 535,  
 537  
 Cercospora dubia 268, 275; tertia 53  
 Certocystis 101  
 Chaetocypha libertiana 339  
 Chaetodiplodia 267, 270, 273; caulina 271, 273,  
 275  
 Chaetomiaceae 23, 99, 146  
 Chaetomium 101  
 Chaetosphaeria 101  
 Chalara 101  
 Chamaezyces fracidus 410  
 Cheilymenia 125, 126; pulcherrima 119; vitellina  
 119, 121\*  
 Chenopodium 267, 269, 270, 274, 542; album 269,  
 270, 273–276; bonus-henricus 274; glaucum  
 273; hircinum 270; hydridum 274, murale 274;  
 quinoa 270, 275, 276; sueticum 268, 273, 274;  
 urbicum 268; viride 274; vulvaria 274; spec. 273  
 Chloridium virescens 43  
 Chrysosporium merdarium var. **roseum 78**  
 Cistus monsepliensis 318  
 Cladium 539  
 Cladobotryum 33–35, 37, 41, 44, 45, 47, 57, 68, 76,  
 80; agaricina 42; agaricinum 34, 42, 47, 68;  
 aleuritidis 69; apiculatum 34–36\*, 37, 67–69,  
 79; arbuscula 77; **asterophorum 34, 35, 37, 38\***,  
 44, 70; australe 34, 70; binatum 34, 41; bole-  
 torum 42; candidum 42; capitatum 34, 72;  
**curvatum 34, 36, 39\***, 72; cylindrosporium 47;  
 dendroides 34, 35, 41, 42\*, 55, 68, 70–72, 76, 78,  
 79; elegans 34, 45, 72; gelatinosum 34, 73;  
 heterocladium 34, 75; leptosporium 34, 55, 56;  
 macrosporium 34, 41; majus 45; minus 47;  
**multiseptatum 34, 36, 43, 44\***, 76; mycophilum  
 34, 36, 41, 45, 73, 76; ovalisporum 34, 76, 77\*;



*(Cladobotryum)*

roseum 47; tenellum 42; tenera 42, 67; ternatum 34, 41, 45, 47; ternatum var. binatum 41; terrigenum 34, 42; thumenii 34, 79; variosperrum 34, 46; varium 34, 36, 45-47, 76-79; verticillatum 34, 35, 38, 46\*, 47, 68, 75, 79

## Cladonia 293

Cladotrichum ternatum 79

Cladosporium modestum 51, 53; varium 78

## Claudopus 245

Clavaria aculina 537; gracilis 537

Clitocybe 245, 406

Cnazonaria 535; aculina 537; rickii 537

**Colleticoconis 33, 48; aecidiophila 48, 49\*, 50\*, 67**

Colletotrichum aeciicola 51; uredinophilum 51

Collybia sulcatissima 1, 12

**Columnodontia 326, 327; columellifera 327; lutea 327; resupinata 327; subfascicularia 327**

Coniophora 335

Conocybe 418

Coprinaceae 425

Coprinus 383, 384, 386, 390, 394, 400, 404-406, 408, 409, 412, 418, 419\*, 422, 423, 425; § Alachuanii 386, 410, 418; § Auricomi 421; § Cycloidei 390, 419; § Domestici 421; § Exanulati 421; § Farinosi 422; § Glabri 422, 423; § Hemerobii 421, 423; § Impexi 386, 418, 423; § Lanatuli 421; § Micacei 419, 421; § Nudi 422; § Setulosi 409, 421-423; § Vestiti 390, 419, 421; angulatus 403\*, 405, 416; atramentarius 415; auricomus 415, 422; bisporus 417; boudieri 405, 416; brassicae 386, 387\*, 388, 389\*, 390, 408, 409, 412, 418; bulbillosus 413; comatus 415; cortinatus 413, 419; cubensis 413; curtus 399\*-401\*, 402, 404, 416, 421, 422; disseminatus 404, 407, 416, 421, 422; ephemerus 407, 417; fimetarius 383, 396; flocculosus 392, 393\*, 395\*, 407, 414, 421; heptemerus 404, 422; hexagonosporus 401\*, 403\*, 404, 407, 409, 416, 422; lagopus 415; macrocephalus 388, 396, 397\*-399\*, 407, 408, 415, 421; macrorrhizus 388, 390, 395\*-397\*, 398, 407, 415, 421; micaceus 407, 414, 421; miser 407, 417; narcoticus 413, 419; niveus 412, 419; patouillardii 389\*, 391\*, 413, 419, 421; phaeosporus 385\*-387\*, 388, 390, 407, 408, 412, 418, 419; plicatilis 409, 417, 423; poliommallus 390, 391\*, 393\*, 408, 414, 418, 419, 421; radians 408, 414, 419; radiatus 396; stellatus 407, 417; stercorarius 406, 409, 412, 419

Coprobria 125, 126; granulata 119, 120\*

Coprotus 125, 126

Cremiella 26; cuboidea 26

Corneromyces 335, 336; kinabalui 335

**Corneromycetaceae 325, 336**

Cornus mas 339; mascula 343

Corticiciae 325

Corticium fallax 137, 139; filicinum 326; grisellum 326; insperatum 334; leucoxanthum 137, 139, 140; luridum 140; niveocremente 335; **rallum 326**; subflavidogriseum 335; subnitens 326; sulfureo-isabellinum 334; tulasnellodeum 335; vellereum 328

Cortinariu 172, 410, 545; acutus 172

Costantinella 72

**Crustomyces 137, 140; pini-canadensis 140; sub-abruptus 140**

Cryptodoxyla 101

Cryptocline 73

Cryptosporiopsis 48

**Cyanobasidium 327; asperulum 328; chordulatum 328**

Cylindrocarpon 71, 72, 76

Cylindrocladium 74; **gregarium 74\*, 75**; parvum 75; peruvianum 75; scoparium 72

Cylindrodendrum 70; album 70

Cylindrophora 67; alba 67, 68; albedinis 67-69; apiculata 36, 67; aranearum 67, 69; candida 67; epitrichia 67, 73; fagi 67, 73; fagi var. candida 70, 71; hoffmannii 75; lindaviana 67, 76; tenera 67, 78; verticillata 67, 72; virgata 67, 79

Cyphella applanata 336; libertiana 339

Cystolepiota 419

Cystospora 73

Dactylaria mycophila 60

Dactylium sect. Helminthophora 55; agaricium 68; boletorum 70; dendroides 41, 68; dendroides var. leptosporum 56; leptosporum 56; macrosporium 41; melleum 46, 47; mycophilum 45; rennyi 46, 47; schlechtendalianum 42; sphaerocephalum 55, 56; tenellum 78; tenerum 56

Debaryomyces 146, 147; castelii 147; coudertii 148; globus 146, 147; hansenii 147; tamarii 148

**Debaryozyma 146, 147; castelii 147; coudertii 148; hansenii 147; polymorpha 148; pseudopolymorpha 148; tamarii 148; vanriji 148**

Delicatula 130

Dendryphon 275

Dennisiopsis 114

Denticularia 33, 51, 53, 62, 66; fici 51, 73; **limoniiformis 51, 52\***, 60, 76; modesta 51, 53, 76; tertia 51, 53

- (Depazea) see *Sphaeria vagans*  
*Desmazierella* 126; *acicula* 120  
*Dichotomyces* 101  
*Didymocladium* 34; *ternatum* 34, 79  
*Diplocladium* 34; *clavarium* 34; *cylindrosporum* 34, 72; *elegans* 34, 72; *gregarium* 34, 73, 74; *hydrangae* 34, 75; *majus* 34, 76; *melleum* 34, 46; *minus* 34, 76; *penicilloides* 34, 46, 47; *penicilloides* var. *clavarium* 72; *preussii* 34, 79; *rennyi* 34, 46; *tenuis* 34, 78; *theobromae* 34, 79  
*Diplodia* 267; *chenopodii* 271, 275; *hyalospora* 275, 276  
*Diplodina* 267; *ellisi* 275, 276; *atriplicis* 271, 273; *chenopodii* 271, 274  
*Dryopteris* 331
- Eccila* 245  
*Ellecevera parasitica* 58  
*Emericella* 101  
*Emericellopsis* 143  
*Engyodontium* 33, 53; *album* 53, 55, 69; *parvisporum* 53, 54\*, 55, 77  
*Enterocarpus* 23  
*Entoloma* 283, 284; § *Alboleptonia* 246, 247; § *Dysthales* 215; § *Endochromonema* 246, 254; § *Entoloma* 247, 284; § *Inopilus* 209, 229; § *Nolana* 284, 286, 289; § *Papillati* 246, 261, 263, 286, 296; § *Paramammosi* 289; § *Pouzarella* 215; § *Pourzaromyces* 207–211, 215, 226, 227; § *Versatilis* 210, 229, 234, 240; *acidophilum* 283, 285; *album* 247; *araneosum* 234, 238–240; *araneosum* var. *rubrolanatus* 238; *araneosum* f. *araneosum* 229, 234, 235\*, 236, 239; *araneosum* f. *fulvostrigosum* 207, 229, 236, 237\*, 239; *ardosiacum* 259; *argenteostriatum* 283, 285, 286\* *babingtonii* 241; *brevipes* 263; *brunneum* 256; *cathionis* 283, 287; *cephalotrichum* 260; *chlorinosum* 283, 287; *cinerascens* 247, 256; *cinereum* 258; *elandestinum* 223; *conicum* 258, 259; *crasipes* 248; *cryptocystidium* 283, 287, 288\*, 289; *cuneatum* 247, 248; *cuniculorum* 283, 289, 291\*; *defibulatum* 283, 290, 291\*; *dysthales* 211, 214–216\*, 217\*–219, 222, 223, 241, 242; *dysthales* f. *acystidiosum* 207, 211, 219, 231\*; *dysthales* f. *dysthales* 215; *dysthaloides* 207, 211, 215, 218, 219, 220\*, 221\*, 223, 225, 227, 238, 239, 241; *farinogustus* 283, 292; *farinolens* 292; *fernandae* 250, 286, 287, 290, 291, 294; *fractum* 250, 287; *fulvostrigosa* 236; *fulgostrigosum* 238; *globisporum* 255; *griseocyaneum* 259; *hirtum* 207, 211, 223, 224\*, 225\*, 240, 241, 250, 251, 254, 258; *incertum* 251; *indutum* 229, 239\*, 240; *inodorum* 251; *juncinum* 235; *kervernii* 255; *lanuginosipes* 248; *leptonipes* 262; *longipes* 204; *mammosum* 298; *minimum* 260; *minutum* var. *polymorphum* 248; *nitens* 252; *nivescens* 247; *nodospora* 241; *nodosporum* 207, 215; *occultopigmentatum* 283, 292; *olorinum* 260; *omphaliaeformis* 262; *ortonii* 283, 291, 292; *pallescens* 251, 254; *papillatum* 260, 262 290; *parkensis* 262; *poliopus* 262; *prismatospermum* 251; *pseudotelamonia* 263; *psilopus* 250, 283, 287, 293, 294\*; *pulverum* 211, 223, 225, 226, 228\*, 229\*; *rhombsporum* 251; *robiniae* 254; *romagnesii* 207, 211, 223, 225, 226, 231\*; *sericellum* 258, 260; *stauosporum* 252, 288; *stauosporum* var. *pusillum* 252; *strigosissimum* 207, 209, 211, 212\*, 213\*–215, 241, 242; *subsericellum* 260; *tenellum* 256; *tibiicystidium* 283, 294\*, 295; *tristis* 254; *undulatosporum* 283, 295–297\*; *velenovskiyi* 258, 298; *velenovskiyi* var. *longicystidium* 259, 283, 296, 297\*; *ventricosum* 283, 298; *versatilis* 229, 230, 232\*, 233\*, 234; *vinaceum* 298, *vinaceum* var. *fumosipes* 283, 297\*, 298, 299; *vinaceum* var. *vinaceum* 299; *vinaceum* var. *violeipes* 283, 299; *xanthocaulon* 283, 299; *xylophilum* 252; *zonatum* 255
- Epithela* 331; *galzinii* 331; *nikau* 331; *typhae* 331  
*Equisetum* 539  
*Erica arborea* 318  
*Ericum* 293  
*Eriomycopsis tenuis* 61  
*Eupatorium* 539  
*Eupenicillium* 101  
*Euphorbiaceae* 14, 20  
*Eurasia* 37, 55; *bondarzewiae* 56  
*Eurotium* 101; *pulcherrimum* 28  
*Exochalara* 101
- Fagus* 218, 374; *sylvatica* 214, 344  
*Faurelina* 23, 24, 28; *elongata* 23, 27\*, 28\*; *firmigenes* 28  
*Flacourtiaceae* 14, 20  
*Flagelloscypha* 337, 339, 344–346; § *Lachnelloscypha* 337, 345, 356; *dextrinoidea* 344; *donkii* 344; *faginea* 344; *libertiana* 337, 339, 340\*, 342–345; *minutissima* 344; *montis-anagae* 337, 339, 342, 344, 345; *spec.* 342, 343\*–345  
*Fusarium* 68, 75, 101; § *Arthrosporiella* 57; *albedinis* 68; *decemcellulare* 79; *longum* 78; *oxysporum* 75; *oxysporum* var. *albedinis* 68; oxy-

*(Fusarium)*

- sporum f. spec. albedinis 68; solani 79; theobromae 79  
 Fusichalara 101  
 Gabarnaudia 101, 143  
 Gamsia 101  
 Geopyxis 126; carbonaria 120, 122\*; majalis 128  
 Geotrichum microsporum 26  
 Gibberella 101  
 Gibellula 54, 70  
 Gliomastix 97-99, 111; protea 111  
 Gloecystidiellum humile 334  
 Gloeosporidiella 48  
 Gloeosporium 33; aecidiicola 51; aecidiophilum 33, 48, 51; fructigenum 79  
 Gloiosphaera 74; globuligera 74  
 Glomerella cingulata 79  
 Gomphideus roseus 410  
 Gonatorrhodiella coccorum 54  
 Gonytrichum 43  
**Granulobasidium 328; vellereum 328**  
**Granulomanus 70; araneorum 69, 70**  
 Graphium 23, 27  
**Gyrophanopsis 329; zealandica 329**  
 Hadrotrichum 58  
 Haematostereum 320  
 Hamigera 101  
 Hansfordia 75; pulvinata 69, 70, 75  
 Harposporium 71; siccyodes 71  
 Hebeloma 545  
 Hedera spec. 71  
 Helminthophora 33, 35, 55; agaricinum 56; fusariorum 56; **sphaerocephala 56\***, 68, 70, 73, 75, 78; tenera 55, 56  
 Helvella 126; crispa 120, 122\*  
 Heterosecypha 336  
**Heterosecyphaceae 325, 336**  
 Hevea brasiliensis 54  
 Hirsutella gracilis 537  
 Humaria hemisphaerica 128  
 Huminiaceae 11  
 Hyalostachybotrys 97  
 Hydnium griseo-fuscescens 137, 138; pseudo-mucidum 334  
 Hydrasidium 334  
 Hydrocybe 372  
 Hygrophorus 357, 358, 359, 373, 376, 406; § Colorati 358; § Olivaceoubrini 357, 359, 363; § Tephroleuci 364; agathosmus 364; dichrous 357, 358, 363, 365, 371-374, 376; dichrous var. fuscovinosus 371; fuscoalbus 357, 358, 360, 363-365; latitabundus 357-361\*, 362\*-364\*, 365, 372, 379\*, 380\*; leucophaeus 374 limacinus 357, 360, 364, 365, 372, 373; mesotephrus 358, 359, 364\*, 365, 369\*, 372, 374, 375\*, 376, 379\*, 380\*; olivaceoalbus 357-359, 363, 373, 376, 378, 379\*, 380\*; olivaceoalbus var. gracilis 380; olivaceoalbus f. gracilis 376; olivaceoalbus f. obesa 371, 380; olivaceoalbus f. obesus 360, 372; **persoonii 357, 358, 360, 363, 365, 366\***, 368, 369\*, 370\*-373, 376, 379\*, 380\*; pustulatus 364  
 Hyphodontia breviseta 33  
 Hypholoma 410; elongatipes 410  
 Hypochnium 328  
 Hypochnium dussii 335; subviolaceus 334  
 Hypocreaceae 99  
 Hypocreales 143  
 Hypomyces aurantius 45-47; dactylarioides 34, 35, 40\*, 41; odoratus 45; petchii 35; rosellus 42; trichothecioides 36  
 Hypsizygus 89; tessulatus 89  
 Hyptis capitata 51  
 Hyphodontia breviseta 52  
 Incrustoporia 347, 348, 353, 354; alutacea 347, 348; carneola 347, 348, 352\*-354; nivea 347, 348; percandida 347, 348; stellae 347, 348; subincarnata 347, 348; tschulymica 347, 348;  
 Inocybe 157, 158, 218, 239; § Clypeus 157, 158; § Inocybe 157; § Inocybium 157; asterospora 202; avellana 200; bresadolae 162; bucknallii 215, 218; descissa 204; longipes 204; murrayana 190; neoumbrina 182; petchii 164; rhombospora 204; umbonata 164, 166  
 Inopilus 207  
 Intsia 191  
 Iodophanus 126; carneus 117\*; granulipolaris 128  
 Iris 539  
**Jacksonomyces 329; phlebioides 329**  
 Jafnea fusicarpa 128  
 Juncus 539  
 Juniperus communis 224  
 Kernia 23, 24, 101  
 Kneiffia coriacea 321  
 Lachnella 337, 339, 344-346; libertiana 339; rosae 337, 339, 341\*, 342, 345; subfalcispora 337; tiliiae 337, 338\*  
 Lactarius 33, 92, 307; griseus 85, 86, 96; maculatus 86, 96

- Lagarobasidium detriticum* 334  
*Lasiobolus* 125, 126; *monascus* 114, 122, 125\*, 128; *pilosus* 122, 125\*  
*Lasiodermatis serricornis* 543  
*Lasiosphæria* 101  
*Lecanium* 54  
 Lecidithaceae 22  
 Leguminosae 5, 11, 14, 20, 22  
*Lepidomyces* 329, 330; *subcalceus* 330  
*Lepiota* 410; *clypeolaria* 386  
*Leptodontium* 43  
*Leptographium* 74  
*Leptokalpion* 114, 122  
*Leptonia* 240, 245, 256, 259; § *Alboleptonia* 246, 255, 258, 260; § *Eccilia* 262; § *Leptonia* 259; *aurea* 253\*, 255; *babingtonii* 215, 241; *brevipes* 263; *brunnea* 253\*, 255, 256; *cephalotricha* 260; *cinerascens* 256, 257\*; *cinerea* 225, 241, 256, 257\*, 258; *citrina* 257\*, 258; *conica* 253, 258, 259; *decurrens* 259; *densifolia* 257\*, 259 *fulvostrigosa* 236, 238; *induta* 239; *involuta* 257\*, 259; *kervernii* 255; *mammillata* 257\*, 259, 260; *minima* 260, 261\*; *mycenoides* 260, 261\*; *nigrella* 226, 242, 260, 261\*; *nivea* 260; *omphaliaeformis* 249\*, 261, 262; *papillata* 261, 262; *pulverea* 226; *strigossima* 211; *viridula* 230; *xanthopa* 262  
*Leptoporus chioneus* 347; *semipileatus* 347  
*Leptotrimitis* 347  
*Leucogyrophana* 330  
*Leuconeuospora* 23, 24; *elongata* 23, 27, 28; *pulcherrima* 23, 28\*, 30  
*Leucosphaera* 141, 143; *indica* 141, 142\*  
*Limacella* 301; *glioderma* 301, 303\*, 305\*–308, 410; *guttata* 301, 306–308; *lenticulare* 301  
*Limacium fuscoalbum* 360; *mesotephrus* 374; *olivaceoalbum* 376  
*Lindtneria trachyspora* 86, 89, 94  
*Lithocarpus* 158–161, 163, 166, 168, 173, 178, 179, 182, 183, 193, 195, 197, 199  
*Litschauerella abietis* 335  
*Lolium perenne* 66  
*Lophotrichus* 23, 24; *ampullus* 24; *bartletii* 24; *brevirostris* 24; *martinii* 24  
*Luzula* 374  
 Lyophylleae 83  
*Lyophyllum* 83, 85, 87, 89; § *Difformia* 87; *anthracophilum* 86, 94; *connatum* 86, 87, 89; *favrei* 86, 94; *fumosum* 86, 94; *leucophaeatum* 86, 94; *ulmarium* 86, 87, 89  
*Macrocyrtidia cucumis* 254  
*Macrophoma* 267; *chenopodii* 271, 275  
*Mariannaea* 101  
*Megalobasidium humili* 334  
*Megalocystidium* 137, 139; *lactescens* 140; *leucoxanthum* 140; *luridum* 140  
*Melonoleuca alboflavida* 86, 90, 92, 96  
*Melanopsamma* 99, 101  
*Memnoniella* 98, 99, 101  
*Merulius aureus* 330; *elliottii* 330  
*Metarrhizium* 99, 101  
*Michelia champaka* 314  
 Microasceae 23, 28, 30  
*Mucroascus* 23, 24, 101  
*Microplodia* 267; *henningsii* 271, 274, 275  
*Molinia* 363  
*Mollisia microcarpa* 73  
*Monilochaetes* 101  
*Monocillium indicum* 101  
*Monosporium agaricinum* 68  
*Mucor* 309; *dendroides* 41  
*Mucronella rickii* 535, 536\*, 537  
*Mucrosporium* 34; *dendroides* 34; *fusarisporum* 34, 73; *leptosporum* 34, 75; *schlechtendalianum* 34; *sphaerocephalum* 34, 56; *tenellum* 34, 78; *unisetatum* 34, 79; *verticillatum* 34, 42  
*Mycena* 129, 130, 132, 277, 278, 411; § *Adonidae* 130; § *Amabilissimae* 130; § *Filipedes* 130; § *Fragilipedes* 134; § *Hiemalis* 130; § *Lacteeae* 130; § *Rigidipedes* 130, 134; *abramsii* 132; *aetites* 132; *alba* 280; *alcalina* 132; *atromarginata* 134; *capillaripes* 134; *chlorinella* 132; *citrinomarginata* 132, 134; *corticola* 277, 280; *delectabilis* 130; *filopes* 130, 131\*, 134; *flavoalba* 130, 133\*; *flocculentipes* 277, 278, 279\*, *galericulata* 130, 131\*, 132; *gracilis* 129; *hiemalis* 130, 133\*, 277–279\*, 280, 282; *jacobi* 129, 132; *lactea* 130, 133\*, *leptocephala* 132, 134; *melligena* 277, 280; *metata* 130, 131\*, 134, 277, 279, 280; *niveipes* 129, 132; *olida* 130, 133\*; *olivaceomarginata* 134; *phylligena* 277, 279, 280; *polygramma* 134; *polygramma* f. *ambigua* 134; *praecox* 132; *purpureofusca* 134; *rubromarginata* 134; *sepia* 130, 131\*, 280; *seynii* 134; *speirea* 280; *strobilicola* 132; *supina* 280; *venustula* 280; *vitilis* 134; *vitrea* 280; *vitrea* var. *tenella* 277, 279, 280; *zephyrus* 132  
*Mycogone puccinioides* 37; *verticillata* 47  
*Myrtus communis* 318  
  
*Naucoria* 410  
*Neea* 2, 14, 17, 20  
*Nematoloma polytrichi* 410

- Niesslia 101  
 Niptera microcarpa 73  
 Nodulisporium didymosporium 57; hinnuleum 34  
 Nolanea 207, 210, 245, 258, 283, 296; § Endochromonema 246; alba 246, 247, 249\*; araneosa 234; autumnalis 240, 247; babingtonii 215, 241; betulina 247, 257\*; crassipes 247, 248, 257\*; depressa 248, 249\*; dysthales 215, 219; farinolens 291, 292; fibrillosipes 218, 219; fracta 248, 249\*, 250; fulvostrigosa 222, 238; fumosella 207–209; galeraeformis 250; globispora 249\*, 250, 255; hirta 223, 250; holoconiota 248; inodora 249\*, 251; majalis 251, 253\*; mammosa 247; minuta 248; nitens 252; nodospora 215; pallescens 251; paludosa 252; pusilla 249\*, 252; rhodoura 238; robiniae 249\*, 252, 254; setulosa 223, 253\*, 254; strigosissima 207, 208, 211; tristis 249\*, 254; undulata 255; variegata 255; versatilis 230, 233; zonata 249\*, 250, 255  
 Nothofagus 1, 2, 160, 161, 163, 177, 180–183, 187, 201, 202; carrii 177, 187; grandis 177, 187  
 Nyctaginaceae 2, 14, 20
- Ochnaceae 11  
 Octospora 125, 126; musci-muralis 119, 121\*  
 Odontia columellifera 327; lutea 327; subabrupta 137, 140  
 Odiiodendron kalrai 25  
 Ophalia 129; corticola 282  
 Omphalina 245  
 Ophiostoma epichloes 66; stenoceras 63  
 Ophiostomataceae 23  
 Otidea 128; leporina 128; onotica 119
- Paecilomyces 98, 101, 107, 111; griseoviridis 102; humicola 102; striatisporus 102; variabilis 102  
 Palmea 14, 20  
 Paraphelaria 539; amboinesis 540; **borneensis** 539  
 Parasitella 309  
 Parvobasidium **lundellii** 334  
 Paullicorticium **niveocremeum** 335  
 Pelargonium 178, 179  
 Pellicularia asperula 328; chordulata 327, 328; cystidiata 334; scabrida 334; zealandica 329  
 Penicillifer 101  
 Penicillium 99, 101, 328; claviforme 98; *co-ryophilum* 97  
 Peniophora aegerita subsp. abietis 335; arenata 334; detritica 334; incarnata 318; pelliculosa 334; phlebioides 329; radicata subsp. leprosa 334; subcalcea 330; umbracula 332; vermifera 335
- Petriella 23, 24  
 Petriellidium 23, 24; boydii 25; desertorum 26, 27; *fimeti* 23, 24, 25, 26\*  
 Peziza 126; ammophila 118\*; badia 118; cerea 118; cunicularia 113; depressa 118; succosa 128; succosella 118\*  
 Phaeolepiota 410, 419  
 Phaeomarasmium 419  
 Phanerochaete **arenata** 334; **leprosa** 334; **pelluculosa** 334  
 Phialocephala 74, 75  
 Phialomyces 98, 101  
 Phialophora 101  
 Phialotubus 101  
 Phlebia 329, 330  
**Phlebiopsis** 137; **gigantea** 137  
 Phlebospora 267; chenopodii 275, 276  
 Phloenix dactylifera 68  
 Phoma 267, 268, 275, 542; chenopodii 268, 270; **dimorphospora** 267, **269**, 270, 272\*, 273; exigua 268; **heteromorphospora** 542; **variospora** 267, **268**–270, 272\*, 273, 542; westendorpii 268  
 Phyllosticta 267; atriplicis 271, 274; chenopodii 268, 270, 542; dimorphospora 269, 270  
 Physodentia lundellii 334  
 Picea 363, 378; abies 52, 110, 378  
 Pichia polymorpha 148; pseudopolymorpha 148; vanriji 148  
 Pidoplickoviella 30  
 Pinus 363; sylvestris 70, 73, 107  
 Pirola 180  
 Pistillaria 535; aculina 537; aculina subsp. acicula 537; aculina subsp. equiseticola 537; aculina subsp. juncicola 537; equiseticola Boud. 537; ferryi 537; gracilis 537  
 Pithoascus 23–25; **langeronii** 23, **24**, 25\*, 26, 28\*  
 Pithya 126; cupressina 120, 123\*  
 Pleospora 275; calvescens 275  
 Pleurotus ulmarius 89  
 Pluteus 245, 247; **betulinus** 247  
 Polypaecilium 101  
 Polyporus chioneus 347; niveus 347, 353; semi-pileatus 347  
 Polytrichum piliferum 286, 299  
 Populus 73  
 Poria 347; alutacea 353; attenuata var. subincarnata 353; carneola 352\*; percandida 353; stellae 347, 353; tschulymica 252  
 Pouzarella 208, 209; § Dysthales 210; § Pouzarella 210; § Versatilis 229; ambrosia 240; araneosa 234; autumnalis 240; babingtonii 215, 241; dysthales 215; hirta 223; induta 239; nigrella

- (*Pouzarella*)  
 225, 226; nodospora 209, 210; pulverea 226; setulosa 223, 254; strigosissima 211, 241; versatilis 230  
*Pouzaromyces* 207–209, 240, 261; fumosellus 211, 242; strigosissimus 211  
*Prunus spinosa* 224  
*Psathyrella* 242; multipedata 409; pyrotircha 407  
*Pseudeurotiaceae* 141, 143  
*Pseudeurotium* 143  
*Pseudocercospora* 58  
*Pseudoepicoccum* 58  
*Pseudofusarium* 57  
*Pseudohansfordia* 33, 35, 41, 53, 55, 57, 58, 62; **didyma** 58, 72; hughii 58; irregularis 35, 57–59\*, 60, 75; **meliolae** 52 58, 60, 76; **mycophila** 58, 60, 61\*, 76; **pulchella** 58, 60, 77; **stereicola** 58, 61, 78; **tenuis** 58, 61, 78; **varanasiensis** 58, 61, 79; **venezuelensis** 58, 62, 79  
***Pseudomerulius* 330; aureus 330; elliottii 330**  
*Pseudomicrodochium* 71; aciculare 70; **candidum 71**  
*Pseudoplectania* 126; nigrella 120, 123\*  
*Pseudeurotiaceae* 30  
*Psilocybe* 204  
*Psychotria* 2, 17, 20  
***Pteridomyces* 331; galzinii 331**  
*Pterula* 535; carcis-pendula 537; gracilis 535, 537, 538\*  
*Puccinia* 51  
*Pustularia cupularis* 115  
*Pyronema* 125, 126; omphalodes 119\*  
*Quercus* 1, 2, 227, 367, 368, 374; ilex 318; robur 368  
*Radulomyces* 326; **pseudomucidus 334; sulfureo-isabellinus 334**  
*Radulum pini-canadense* 140  
*Ramularia* 43, 71; tenuis 61  
*Rhinotrichum parvisporum* 33, 53, 54  
*Rhizomucor* 309  
*Rhodocybe* 87; alutacea 87; leucopaxilloides 86, 87, 94; nitellinia 86, 89; parilis 86, 92, 94, 96; suburens 86, 87, 94; spec. 94  
*Rhodophyllus* 85, 89, 94, 130, 209, 245, 246, 283; § **Induti** 207, 210; § **Inopilus** 209, 229; § **Luctuarii** 210; § **Papillati** 246; § **Pouzaromyces** 209, 210; § **Versatilis** 207, 208, 229; ambrosius 240; araneosus 222, 234, 236, 238, 239; asprellus 256; babingtonii 207, 210, 211, 223, 231\*, 240, 241; bicolor 86, 89, 94; dysthales 215, 219, 238; dysthales f. **subnigrellus** 225; fernandae 250; fumosellus 215, 242; fumosellus var. **homomorphus** 219; hirtipes 86, 94; incanus 86, 90, 94; indutus 210, 239; inocephalus 207; juncinus 250, 255; minutus var. **polymorphus** 248; mougeotii 259; nitens 252; olorinus 260; omphaliformis 260; poliopus 262; prunuloides 86, 94; strigosissimus 211 **subnigrellus** 225, 226; tenellus 256; turci 86, 94; versatilis 207, 209, 230; viridulus 86, 94, 230, 233, 234; xylophilus 252; spec. 222  
*Rhodotorula* 542  
*Rhyarobius* 126; caninus 122, 124\*; crustaceus 122, 124\*; myriosporus 122, 124\*  
*Riclaretia* 101  
*Rosa spauldingii* 339  
*Rozites* 410  
*Rubiaceae* 2, 14, 20  
*Russula* 33, 92, 307; adusta 86, 89, 92, 96; fragilis 86, 92, 96; sardonias 86, 90, 96  
*Saccharomyces hansenii* 147  
*Saccobolus* 114, 126\*; depauperatus 128; glaber 117\*; kernvernii 115  
*Sagenoma* 100; viride 97, 105  
***Sagenomella* 97, 100, 107, 111; alba 102, 104\*, 110\*; diversispora 100, 102, 103\*; griseoviridis 102, 104\*, 107; humicola 102; oligospora 102, 103\*, 105\*, 111; sagenomatis 102, 104\*, 105, 111; sclerotialis 102, 108\*, 109\*, 111; striatospora 102, 103\*; verticillata 102, 106\*, 107, 108\***  
*Sagrahamala* 98, 99, 102, 111; griseoviridis 102; luzulae 98; murorum 98; striatospora 98, 102; variabilis 102  
*Salix repens* 214, 218  
*Sapindaceae* 11, 14, 20  
*Sapotaceae* 5, 11, 20  
*Sarcoscypha* 114, 126; coccinea 120, 123\*  
*Sarothamnus* 261  
*Seedosporium* 23  
*Scopulariopsis* 23, 101; diversispora 102  
***Scotomyces* 137, 139; fallax 139; subviolaceus 334**  
*Scutellinia* 125, 126; armatospora 119, 121\*; scutellata 128  
*Sepedonium meliolae* 60  
*Septofusidium* 101; herbarum 101  
*Septoria* 267; atriplicis 271, 274; chenopodii 274, 275; westendorpii 268–270  
*Serpula* 330  
*Sibirina* 41, 55; fungicola 41, 73; orthospora 41  
*Sitodrepa paniccae* 543  
*Skeletocutis* 347, 348, 353, 354; **alutacea** 347, 350\*,

*(Skeletocutis)*

**353**, 354; amorphe 347, 348, 353–355\*; **nivea** 347, 349\*, **353**, 354; **percandida** 347, **353**, 354; **stellae** 347, 351\*, **353**, 354; **subincarnata** 347, 351\*, **353**, 354; **tschulymica** 347, 352\*, **353**

**Skeletohydnum 331; nikau 331**

Sordariaceae 23, 30, 146

Sowerbyella 125, 126; imperialis 128; unicolor 119, 121\*

Sphaeria vagans 275; vagans f. atriplicicola 271, 275

Sphaeriaceae 99

Sphaeriales 143

**Sphaerobasidium minutum 335**

Sphaeronaemella 101, 143; fimicola 143

Sphaerosporella brunnea 128

Sporendocladia 101

Sporoschisma 101

Sporothrix 33, 62, 64, 66; fungorum 63; ghanensis 62, 66; **guttuliformis 62**, 63\*, 75; pallida 63; ramosissima 63, 66; **rectidentata 64**, 69, 77; schenckii 63, 66; **sclerotialis 64**, 65\*, 78; setiphila 77; **vizei 66**, 67\*, 79

Sprottrichum agaricinum 68; boletorum 70; candidum 71; cylindrosporium 78; merdarium var. roseum 78; roseum 78

Stachybotryna 101

Stachybotrys 97, 99

Stagnospora 267, 271; atriplicis 271; chenopodii 269, 271, 275

Stagonosporopsis 267; atriplicis 271, 274

Stemphylium 275

Stereum 311, 319, 320, 321, 323; § **Acanthostereum**

311, **320–323**; § **Aculeatostereum 311**, **320**, 322, 323; § **Cruentata 311**, 320; § **Luteola 311**, 320; § **Stereum 321–323**; acanthophysatum 314, 320; australe 321; complicatum 320, 321; durbanense 321; fasciatum 319; hirsutum 319, 321; hirsutum subsp. sulphuratum 321; illudens 320, 321; insignitum 318–321; kurilense 314; lobatum 321; macrocystidiatum 321; obscurans 321; ochraceo-flavum 321; ochroleucum 319, 321; ostrea 320, 321; **peculiare 311**, 313\*–315\*, 317 319, 320, 323; radiato-fissum 320; rameale 319–321; reflexulum 311, 317, 318\*–321, 323; repandum var. lusitanicum 325, 326 rimosum var. africanum 321; rugosum 320, 321; sanguinolentum 321; scutellatum 321; sericeum 321; spectabile 320; striatum 320, 321; styracifluum 321; subtomentosum 319–321; traplanum 321; vellereum 321; versicolor 321; zebra 321; zonarium 321

Strobilomyces floccopus 410

**Symbiotaphrina 542**, 543; **buchneri 542**, **543**; **kochii 543**

Sympodiophora 33, 57; didyma 57, 58; pulchella 60; stercicola 57, 58, 61; tenuis 61; varanasiensis 61; venezuelensis 58, 62

Talaromyces 101

Taphrina 542

Telamonia 158, 245; brevipes 261, 263

Tephrocye 83

Testudina terrestris 30

Thecotheus 126; pelletieri 128; spec. 117\*

Thelebolus 113, 114, 122, 125, 126; stercoreus 114, 122, 124\*

Thelephora concolor 321; gigantea 137; lactescens 140

Thermoascus 101

Thielavia 145, 146; hyalocarpa 144; octospora 146; terrestris 101; tetrasperma 146

Thysanophora 101

Torrubiella albolanata 69

Torulaspora 147

Torulomyces 101, 111; lagena 108\*

Torulopsis buchneri 543

Trechispora vaga 540, 541\*

Tremellales 335

Tremelloscypha 336

Trichobolus zukalii 114

Tricholoma 406; focale 408

Tricholomopsis rutilans 408

Trichothecium 35, 71; agaricinum 68; candidum 71; luteum 71; roseum 34, 35

Tritirachium 53, 64; album 53; rectidentatum 64 Tuber 68

Tuberculina persicina 51

Tubulicium clematidis 335; **dussii 335**; **vermiferum 335**

Tubulicrinis 332; **corneri 332**; hamata 332

Typhula 535, 539; brunaudii 537; gracilis 537

Tylopilus felleus 86, 89, 92, 94

Tyromyces semipeleatus 347

Uncigera 70, 71

Urnula 126; platensis 120, 123\*

Verticillium 72, 75, 78; agaricinum 68; cinnamomeum 75; cylindrophorum 72; cylindrosporium 67, 72; falcatum 78; fungicola 67, 72; heterocladum 75, 77; lactarii 47; lecanii 75; vizei 66

Violaceae 20

Wallrothiella subiculosa 111

Wardomyces 101

Xenasma 326, 330; gaspeticum 326; lloydii 326; minutum 335

Xenasmatella allantospora 326; aurora 335; insperata 334; subflavidogrisea 335; tulasnelloidea 335

**Xenosperma ludibundum 335**

Xerocomus 33

Xylariaceae 23

Xylobolus 311, 317, 321-323; frustulatus 317; princeps 317; sepium 317; subpileatus 317, 319

Zopfiella 30; curvata 30