

**Rapid Communication****First records and molecular confirmation of invasive species  
*Elodea nuttallii* (Planch.) H.St.John, 1920 in Lithuania**Jurgita Butkuvienė<sup>1\*</sup>, Liucija Kamaitytė-Bukelskienė<sup>2</sup>, Donatas Naugžemys<sup>3</sup>, Jolanta Patamsytė<sup>1</sup> and Zofija Sinkevičienė<sup>2</sup><sup>1</sup>Life Sciences Center, Vilnius University, Saulėtekio av. 7, LT-10222 Vilnius, Lithuania<sup>2</sup>Nature Research Centre, Institute of Botany, Akademijos Str. 2, LT-08412 Vilnius, Lithuania<sup>3</sup>Botanical Garden, Vilnius University, Kairėnų Str. 43, 01100 Vilnius, Lithuania

\*Corresponding author

E-mail: [jurgita.makaviciute@gmail.com](mailto:jurgita.makaviciute@gmail.com)

**Citation:** Butkuvienė J, Kamaitytė-Bukelskienė L, Naugžemys D, Patamsytė J, Sinkevičienė Z (2022) First records and molecular confirmation of invasive species *Elodea nuttallii* (Planch.) H.St.John, 1920 in Lithuania. *BioInvasions Records* 11(4): 1019–1030, <https://doi.org/10.3391/bir.2022.11.4.20>

**Received:** 7 February 2022**Accepted:** 6 August 2022**Published:** 24 October 2022**Handling editor:** Laura Garzoli**Thematic editor:** Karolina Bączela-Psychalska**Copyright:** © Butkuvienė et al.This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International - CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).**OPEN ACCESS****Abstract**

Alien aquatic plant species have significant ecological and economic impacts on freshwater ecosystems and are mentioned amongst the major threats to aquatic biodiversity. Detection of alien species at the beginning of invasion can help avoid their spread and prevent negative impacts on aquatic ecosystems. This study presents the first records of invasive species *Elodea nuttallii* in Lithuania. This species was found in three localities of the Nemunas River (southern Lithuania) in 2020 and 2021. The localities were mapped using system grid cells arranged based on geographical coordinates with sides of 6' latitude and 10' longitude. The habitat characteristic and associated species were presented. The identity of *Elodea nuttallii* was confirmed using a molecular method (sequencing of two regions *ITS* and *psbE-petL*). The molecular analysis was effective in supporting the identification of individuals collected in the early stages of growth. We suppose that *Elodea nuttallii* may be more widespread in the region of north-eastern Poland, north-eastern Belarus and southern Lithuania and will continue to spread north and northeast directions. The expansion of *Elodea nuttallii* can have a negative impact on valuable natural habitats, particularly on protected throughout Europe water courses with the *Ranunculion* vegetation.

**Key words:** aquatic plants, molecular identification, rivers, alien species**Introduction**

Increasing human populations and expanded global commercial relations together with human activities and climate change-induced environmental changes cause spreading and often the establishment of species worldwide (Rodríguez-Merino et al. 2018; Pyšek et al. 2020). Despite a relatively low number of alien aquatic plant species reported for Europe (almost 100) (Hussner 2012), they make a significant ecological and economic impact on freshwater ecosystems and are mentioned amongst the major threats to aquatic biodiversity (Havel et al. 2015). Early detection of alien species at the beginning of invasion and rapid response can help avoid their widespread and prevent negative impact on aquatic ecosystems (Reaser et al. 2020). Most alien aquatic plants naturalised in Europe are native to

Northern America. The genus *Elodea* (Hydrocharitaceae) is presented by six species: *E. bifoliata* H.St.John 1920, *E. callitrichoides* (Rich.) Casp., 1857, *E. canadensis* Michx., 1803, *E. granatensis* Humb. & Bonpl. 1810, *E. nuttallii* (Planch.) H.St.John, 1920, *E. potamogeton* (Bertero) Espinosa 1928 (Cook and Urmi-König 1985; WFO 2021). Three species of *Elodea* (*E. canadensis*, *E. nuttallii*, *E. callitrichoides*) are still known in Europe (Josefsson 2011). *Elodea canadensis* is the most widespread alien aquatic plant in Europe. It has been reported in 41 countries (Hussner 2012), however, *E. nuttallii* is becoming more frequent and is replacing *E. canadensis* in many regions (e.g., Cook and Urmi-König 1985; Simpson 1990; Barrat-Segretain 2001). Like other *Elodea* species, *E. nuttallii* is a perennial submerged aquatic plant. It has been found growing in very different water bodies, but usually in quiet water at shorelines of lakes, reservoirs, and ponds or in slowly flowing water along rivers and streams, in canals and ditches and wetlands such as cranberry plantations (Uotila 2009; Josefsson 2011; Dzhus 2014). Although both male and female plants have been introduced into European botanical gardens, only female plants appear to be spreading in nature (Cook and Urmi-König 1985). Since the first wild record in 1914 (Cook and Urmi-König 1985), *E. nuttallii* has spread to many European countries. According to the latest summaries (Steen et al. 2019; Dembowska et al. 2021), *E. nuttallii* (with an indication of the first record) occurs in 26 European countries: England (1914), Belgium (1939), the Netherlands (1941), France (1950), Germany (1953), Ireland (1960), Belarus (1964), Austria (1970), Denmark (1974), Luxemburg (1980), Switzerland (1980), Slovakia (1986), Czech Republic (1988), Hungary and Sweden (1991), Italy and Poland (1994), Romania (1997), Ukraine (2001), Bulgaria (2002), Serbia (2005), Croatia and Norway (2006), Slovenia (2007), Albania and Russia (2017). The last record of occurrence in Russia (Panasenko and Shcherbakov 2018) has indicted the easternmost distribution point. As the plant species widely distributed in many European freshwaters, *E. nuttallii* has been included into the List of invasive alien species of Union concern (European Union (EU) Regulation 1143/2014).

In many cases, the spread of *E. nuttallii* is related to river basins, for example, the Rhône River (Barrat-Segretain 2001) and the Alsatian Rhine (Greulich and Trémolières 2006) in France, the Danube River crossing Germany, Austria (Janauer et al. 2007), Slovakia (Hrivnák et al. 2019; Bubíková et al. 2021), Serbia (Vukov et al. 2008), Bulgaria (Georgiev et al. 2019), Romania (Ciocârlan et al. 1998; Sârbu et al. 2006), Croatia (Kočić et al. 2014) and its tributary Drava (Grudnik et al. 2014) in Slovenia, the Dniپر in Ukraine (Prokopuk and Zub 2019; Davydova et al. 2021), the Pripet River in Belarus (Dubovik 2013), the Vistula in Poland (Kamiński 2010).

According to GBIF data (GBIF Secretariat 2021) and literature sources cited above *E. nuttallii* is rare in Eastern European and Scandinavian Peninsula countries. Of the Eastern Baltic countries, this species has so far

**Table 1.** Physico-chemical characteristics (average annual values) of the Nemunas River near Druskininkai.

pH	8.2
Flow velocity m/s	0.64
Temperature °C	10.6
Conductivity µS/cm	447.8
Dissolved oxygen (O <sub>2</sub> mg/l)	10.6
Biochemical oxygen demand BOD <sub>7</sub> (O <sub>2</sub> mg/l)	3.91
Ca <sup>2+</sup> mg/l	225.8
NH <sub>4</sub> <sup>+</sup> (N mg/l)	0.128
NO <sub>2</sub> (N mg/l)	0.015
NO <sub>3</sub> (N mg/l)	1.014
TN mg/l	1.43
PO <sub>4</sub> <sup>3-</sup> (P mg/l)	0.051
TP mg/l	0.102

only been recorded in Estonia (Kukk et al. 2020), however it has never been found in Latvia and Lithuania. In this paper we described the first sites of invasive *E. nuttallii* in Lithuanian fresh waters (Nemunas River). Identification of the specimens collected in the early stage of growth was supported by molecular analysis.

## Materials and methods

### *Study area*

Field surveys were conducted in southern Lithuania in June 2020 and August–September 2021. The approximately 60 km long section of the middle Nemunas River from the state border of Belarus to Merkinė village in Lithuania was investigated. Upstream of the state border, the Nemunas River via the Čarna Hančia River and Augustow Canal is connected with different water bodies in the territory of Belarus and further in Poland. The studied part of the Nemunas River crosses the Dainava plain characterized by a low precipitation (600–650 mm per year) and weak density of the hydrographic network (0.7 km/km<sup>2</sup>) (Česnulevičius 2008). The data on water physicochemical parameters of the monitored Nemunas River section were acquired from the Environmental Protection Agency (AAA 2020) and presented in Table 1.

### *Species identification and distribution*

During the field surveys of the Nemunas River, *Elodea* sp., having a different appearance from *E. canadensis*, was found in three locations. In the field these plants differed from the known *E. canadensis* by relatively long, almost linear leaves and incompact flat apices, however without the obvious characteristic to *E. nuttallii* leaves recurving and twisting. Phytosociological relevés of the plant communities with *Elodea* spp. were conducted applying the Braun-Blanquet (1964) approach.

Identification and morphological measurement of dried specimens tentatively recognised as *E. nuttallii* were performed in the laboratory.

Measurement of the stems and internodes length, counting of the branches was made for 4 well-developed and preserved plants collected in August 2021. Morphology of 30 moistened verticillate leaves from main stem of plants collected in August and September 2021 was studied. The most questionable specimens collected at the beginning of the vegetation period (April 2020) were used for genetic testing.

Herbarium specimens were deposited at the Herbarium of the Institute of Botany of the Nature Research Centre (BILAS 92710, 92711 leg./det. Liucija Kamaitytė-Bukelskienė and Zofija Sinkevičienė) and the Herbarium of Vilnius University (WI P33612, leg./det. Jurgita Butkuvienė). Nomenclature and taxonomy follow WFO (2021). The map with the first localities of *E. nuttallii* in Lithuania was compiled by applying a system of grid cells arranged based on geographical coordinates with sides of 6' latitude and 10' longitude.

### *Molecular analyses*

Six specimens collected in April 2020 that tentatively identified as *E. nuttallii* and six specimens of *E. canadensis* from the Nemunas River above Merkinė were used for genetic testing. The 10–15 mg of green plant material was used for DNA isolation. The total genomic DNA was extracted using Plant Mini Kit (Qiagen), following the manufacturer's protocol. The quality of DNA extractions was verified by electrophoresis on 1% agarose gel.


The *psbE-petL* region (primers *psbE* (5'-AGTAGAAAACCGAAATAA CTAGTTA-3') and *petL* (5'-AGTAGAAAACCGAAATAACTAGTTA-3')) of the *cpDNA* and additional analysis using *ITS* sequences of nuclear DNA (primers *ITS1* (5'-TCCGTAGGTGAACCTGCGG-3') and *ITS4* (5'-TCCT CCGCTTATTGATATGC-3')) were analysed in 12 *Elodea* plants presumably representing different species – *E. canadensis* and *E. nuttallii*. The amplification reactions of *psbE-petL* region were performed as described by Bobrov et al. (2018), and the amplification of the *ITS1-ITS2* region was performed according to Butkuvienė et al. (2020). All polymorphic variants of the *ITS* and *cpDNA* sequences were submitted to GenBank (ON584397–ON584398, ON600780–ON600781).

The MEGA X ver.10.1.8, programme (Tamura et al. 2021) was employed to align sequencing results with *Elodea ITS* region and *psbE-petL* sequences from the NCBI GenBank database, which was extracted using the BLAST® (Madden et al. 1996) tool.

## **Results**

### *Habitats and associated species*

*Elodea nuttallii* was recorded in three localities of the Nemunas River (southern Lithuania) in 2020 and 2021. The species was mapped in two grid cells (Figure 1). The first time *E. nuttallii* was found in the Nemunas River upstream of Merkinė village (Y 54.149192, X 24.180477) on 22 June



**Figure 1.** Distribution of *Elodea nuttallii* in Lithuania (upper point includes one location, lower – two locations).

2020 (Figure 1, upper point). The growing site is on the edge of a rapid covered with boulders of various sizes and patches of silty sand deposits and with typical vegetation of the *Ranunculon fluitantis*. Several young shoots grew at a depth of 0.65 m, on a sandy and gravelly bottom with dominant *Myriophyllum spicatum* L. and other submerged plants (Table 2, 1<sup>st</sup> relevé).

The second location of *E. nuttallii* was discovered in the Nemunas River downstream of the Baltoji Ančia River mouth situated along the left shore (Y 53.954161; X 23.835154) on 31 August 2021 (Figure 1, lower point). Both species *E. nuttallii* and *E. canadensis* were found growing together in this location. The plants of both species were well-developed and had clearly different appearance (Figure 2).

The third location was detected about 10 km downstream from the second (Y 53.987915; X 23.914747) on 16 September 2021 (Figure 1, lower point). In these two locations, *E. nuttallii* was found growing under similar conditions. Shallow depressions (0.1–0.5 m) formed on the edges of the riverbed with abundant boulders were typical habitats of *E. nuttallii*. The bottom of the depressions mainly consisted of nutrient-rich silt or silty sand. *Elodea nuttallii* was not very abundant because the water surface was covered by *Lemnaceae* species (Table 2, 2<sup>nd</sup>–4<sup>th</sup> relevés, Figure 3). Plants in the third location were very fragile with many formed turions. The species composition of relevés 3<sup>rd</sup> and 4<sup>th</sup> with dominant helophytes can be attributed to the helophyte communities.



**Table 2.** Phytosociological relevés with occurrences of *Elodea nuttallii* in the Nemunas River (South Lithuania).

Species	Relevé			
	1	2	3	4
<b><i>Elodea nuttallii</i> (Planch.) H.St.John</b>	+	1	1	+
<i>Elodea canadensis</i> Michx.		1		
<b>Ch. Potamogetonetea</b>				
<i>Ceratophyllum demersum</i> L.	+			
<i>Najas major</i> All.			+	+
<i>Myriophyllum spicatum</i> L.	4	1		
<i>Potamogeton nodosus</i> Poir.	1	1	1	
<i>Potamogeton perfoliatus</i> L.		1		
<i>Ranunculus</i> cf. <i>fluitans</i> Lam.	1		+	
<i>Stuckenia pectinata</i> (L.) Börner	2	+		
<b>Ch. Lemnetea</b>				
<i>Lemna turionifera</i> Landolt		1	3	
<i>Lemna gibba</i> L.			2	3
<i>Spirodela polyrhiza</i> (L.) Schleid.	+	3	2	3
<b>Ch. Phragmito-Magno-Caricetea</b>				
<i>Agrostis stolonifera</i> L.		1		1
<i>Alisma plantago-aquatica</i> L.		+		1
<i>Butomus umbellatus</i> L.	1	1	+	3
<i>Carex</i> sp.		+		1
<i>Glyceria maxima</i> (Hartm.) Holmb.		1	+	+
<i>Leersia oryzoides</i> (L.) Sw.		+		+
<i>Mentha aquatica</i> L.		1		+
<i>Phalaris arundinacea</i> L.				1
<i>Rorippa amphibia</i> (L.) Besser		+		
<i>Sparganium erectum</i> L.		+	4	1
Depth m	0.65	0.1–0.3	0.1–0.2	0.2
Substrates	Sand/gravel	Silt/sand/boulders	Silt/sand/boulders	Silt/sand/boulders

### Morphological features of collected specimens

Suspected *E. nuttallii* collected in the Nemunas River downstream of the Baltoji Ančia River mouth was non-flowering but well-developed and suitable for morphological examination and final identification. The length of 4 measured stems varied from 34.7–42.1 cm. They have 3–8 lateral branches, several of them were repeatedly branched. The length of internodes ( $N = 20$ ) was 3.2–22 cm, with an average  $6.5 \pm 4.1$  cm.

Leaves morphology was studied from plants of locality near Baltoji Ančia as well as from the site located downstream. The leaves were in whorls of three, linear to linear lanceolate, straight, or slightly recurved, with weakly undulate margins and acuminate apices. Lamina of some leaves was only slightly twisted. According to Simpson (1986, 1988) morphological differences of *E. nuttallii* and *E. canadensis* are based on a combination of leaf shape, leaf apex shape and leaf width 0.5 mm below the apex-tip. Leaf shape and character of apex in collected plants fully corresponded with features of *E. nuttallii* indicated by Simpson (1986, 1988). Moreover, leaf width 0.5 mm below the apex in both populations ranged from 0.5 to 0.7 mm, with an average  $0.6 \pm 0.07$  mm and not exceeded maximal value 0.7(0.8) mm recorded by Simpson (1986, 1988) for *E. nuttallii*. Plant leaves from first site were longer ( $16.49 \pm 1.4$ ) and wider ( $2.4 \pm 0.3$ ) than in plant collected



**Figure 2.** Habitus of *Elodea canadensis* and *E. nuttallii* from a location downstream of the Baltoji Ančia River mouth. Photograph by dr. Z. Sinkevičienė.

downstream (respectively  $15.5 \pm 1.8$ ,  $1.7 \pm 0.3$ ). In general *E. nuttallii* from the Nemunas River had significantly longer leaves than indicated in many literature sources (Table 3). Similar long-leaved morphotypes have been also described by Simpson (1988).

#### *Molecular analysis*

The sequences of nuclear DNA *ITS* region obtained from the samples by direct sequencing were 806 bp (aligned sequences). Analysis of *ITS* sequencing data revealed 26 variables and 780 conserved sites. Only one site was identified as a six bases insertion/deletion (Table 4 – from 705 bp position to 710 bp position). In addition, single nucleotide polymorphism (SNP) was detected in twenty positions (Table 4).





**Figure 3.** Typical habitat of *Elodea nuttallii* in the Nemunas River – shallow depression along the helophyte belt at the shore covered by *Lemnaceae* species. Photograph by dr. Z. Sinkevičienė.

The alignment of the sequences of *psbE-petL* region included 1109 base pairs. However, the *psbE-petL* region turned out to be less variable. Here, in total 12 variables and 1097 conserved sites. Predominantly single nucleotide substitutions (SNP) (eight positions) and one four base pairs insertion/deletion (from 355 nt position to 358 nt position) (Table 5).

### Discussion

*Elodea nuttallii* has been detected in Lithuania accidentally, however this finding was not a complete surprise considering the rapid spread of *E. nuttallii* in the last two decades. The closest countries where *E. nuttallii* was recorded are Belarus (Dubovik 2013; Dzhus 2014) and Poland (Kamiński 2010), located south of Lithuania. *Elodea nuttallii* has not been found in Latvia (Grinberga



**Table 3.** Morphological characters of *Elodea nuttallii* leaves. Mean  $\pm$  standard deviation and range of parameter values for the Nemunas River plants (N = 30) comparing with literature (Thiébaud and Di Nino 2009; Prokopuk and Zub 2019).

	Nemunas	West Europe	Ukraina	Croatia
Leaf length, mm	15.9 $\pm$ 1.6 12.0–18.5	11.7 $\pm$ 3.1	10.2 $\pm$ 3.0	10.45 $\pm$ 2.11 7.62–14.01
Leaf width at the mid-point, mm	1.9 $\pm$ 0.3 1.1–2.4	2.2 $\pm$ 0.9	1.9 $\pm$ 0.4	1.14 $\pm$ 0.23 0.63–1.64

**Table 4.** Sequences variation in the *ITS* region of *Elodea nuttallii* and *Elodea canadensis*.

<i>ITS</i>	139	140	178	253	255	256	257	258	433	481	485	517	520	591	621	690	705	706	707	708	709	710	712	714	719	723
<i>Elodea nuttallii</i>	G	T	T	C	G	C	T	C	G	C	A	T	A	A	T	T	G	T	G	T	A	A	C	T	C	C
<i>Elodea canadensis</i>	C	C	C	T	T	G	C	T	A	T	T	C	G	G	C	C	-	-	-	-	-	-	T	C	T	T

**Table 5.** Sequences variation in the *psbE-petL* region of *Elodea nuttallii* and *Elodea canadensis*.

<i>psbE-petL</i>	324	346	355	356	357	358	424	480	615	722	725	816
<i>Elodea nuttallii</i>	G	T	T	T	T	G	G	C	C	G	T	G
<i>Elodea canadensis</i>	T	G	-	-	-	-	T	T	T	A	G	C

and Priede 2010), situated to the north of Lithuania. The findings in the southern part of Lithuania suggest that *E. nuttallii* has spread from Belarus. The Nemunas River in the west northern part of Belarus via the Čarna Hanča River and Augustow Canal connects with many water bodies in Poland and the Vistula River Basin, where *E. nuttallii* occurs (Kamiński 2010). The first possible finding of *E. nuttallii* in Poland mentioned by Kamiński 2010) is in the Biebrza River, having a connection with Augustow Canal. The location is about 80 km away from the Lithuanian border. The other close to Lithuania locality of *E. nuttallii* known since 1964 in Belarus is Lake Svityaz (Dubovik 2013). It is possible that *E. nuttallii* has been neglected in north-eastern Poland and north-eastern Belarus as well as in Lithuania. We suppose that *E. nuttallii* may be more widespread in this region. The species has all the typical traits of successful invaders: rapid growth, vegetative reproduction and easy dispersion by waterfowl and currents (Cook and Urmi-König 1985). The rapid spread and climate change with higher temperatures during the vegetation season suggest that *E. nuttallii* will continue to spread to north and northeast directions. The species has also been recorded in Northern countries such as Sweden, Finland and Norway, however, it has not been recorded in Latvia (GBIF Secretariat 2021). Moreover, the specimens identified as *E. nuttallii* are already included in the Estonian herbarium base (<https://elurikkus.ee/generic-hub/occurrences>; <https://elurikkus.ee/plant-atlas/taxon>) and its distribution is mapped (GBIF Secretariat 2021; Kukk et al. 2020). The detection of this species in Latvia – the last Eastern Baltic country is very likely in the nearest future. The occurrence of *E. nuttallii* is highly undesirable in valuable protected areas, especially in such as Natura 2000 (Steen et al. 2019). Bearing in mind the spread of *E. nuttallii* through the river systems this expansion can have a negative impact on river biota and especially on protected throughout Europe habitat of water courses with *Ranunculion* vegetation.

Both *Elodea* species (*E. canadensis* and *E. nuttallii*) exhibit wide morphological variation and morphological features may overlap, especially when plants grow under extreme conditions (Simpson 1988; Kočić et al. 2014; Prokopuk and Zub 2019; Thiébaud and Di Nino 2009). That can be a significant problem of species identification. Typical *E. nuttallii* have narrower and more elongated linear or linear-lanceolate leaves, while leaves of *E. canadensis* are linear oblong to ovate. Other distinguishing features of *E. nuttallii* are strongly recurved and twisted leaves (Simpson 1986). *Elodea nuttallii* from the Nemunas River had typical elongated linear to linear-lanceolate leaves, however without clear expressed recurving and twisting. Absence of these features was observed also in plants from the Iberian Peninsula (Simpson 1986). Leaf shape, leaf apex shape and leaf width 0.5 mm below the apex-tip in our collected plants fully corresponded with features combination indicated by Simpson (1986, 1988) as typical for *E. nuttallii*. This species from the Nemunas River distinguished by significantly longer leaves than indicated in literature cited. Thiébaud and Di Nino (2009) found that the shorter and broader-leaved phenotype of *E. nuttallii* typically occurs in shallow streams, whereas the longer and narrower-leaved phenotype occurs in deeper waters. The plants we observed were growing in shallows, so they may have been affected by several factors such as water level fluctuations, current, shading, abundance of free-floating plants and other.

We had no major problem identifying well-developed plants collected in August and even these collected in September. The specimens collected in June at an early stage of development were questionable. In this case, molecular identification was helpful in distinguishing two morphologically similar species. The regions of nuclear and chloroplast sequencing confirmed morphological species identification. The identical genotypes of *E. canadensis* and *E. nuttallii* have been described in other countries (Huotari and Korpelainen 2012; Rybicky and Voytek 2013).

## Conclusions

The occurrence of invasive species *E. nuttallii* in the Nemunas River at the border crossing of Lithuania, Poland and Belarus suggest that this species could be more widely distributed in this region. Extensive research is needed in the nearest future to assess the true extent of the invasion and the potential impact on native species communities and habitats of water bodies in the region. Ongoing climate change is favourable for the species expansion to the northeast.

## Acknowledgements

The authors are thankful to Lukas Petrulaitis for compiling the distribution map and Violeta Ptašekienė for kind linguistic assistance. We owe gratitude to the subject editor and all anonymous reviewers for their valuable comments to previous versions of the manuscript.

## Funding declaration

The research was carried out within the framework of a project (No. S-SIT-20-1) funded by the Research Council of Lithuania.

## Authors' contribution

All authors performed the field research, analyzed the data, and contributed to editorial changes in the manuscript. JB, DN, JP carried out genetic testing. ZS and JB wrote the manuscript.

## References

- AAA (2020) Aplinkos apsaugos agentūra. Upių monitoringo rezultatai. Vidutiniai hidrocheminiai duomenys. [Results of monitoring of rivers monitoring. Average hydrochemical data]/ [https://aaa.lrv.lt/uploads/aaa/documents/files/Vidurkiai\\_up%C4%97s\\_2020.xlsx/](https://aaa.lrv.lt/uploads/aaa/documents/files/Vidurkiai_up%C4%97s_2020.xlsx/) (accessed 24 May 2020)
- Barrat-Segretain MH (2001) Invasive species in the Rhône River floodplain (France): replacement of *Elodea canadensis* Michaux by *E. nuttallii* St. John in two former river channels. *Archiv für Hydrobiologie* 152: 237–251, <https://doi.org/10.1127/archiv-hydrobiol/152/2001/237>
- Bobrov AA, Chemeris EV, Filipova VA, Maltseva SY (2018) European pondweed in East Siberia: evidence of *Potamogeton rutilus* (Potamogetonaceae) in Yakutia (Asian Russia) with evaluation of current distribution and conservation status. *Phytotaxa* 333: 58–72, <https://doi.org/10.11646/phytotaxa.333.1.4>
- Braun-Blanquet J (1964) Pflanzensoziologie. Grundzüge der Vegetationskunde [Plant sociology. Fundamentals of vegetation science]. 3. Auflage. Springer, Vienna and New York, 634 pp
- Bubíková K, Svitková I, Svitok M, Hrivnák R (2021) Invasive elodeas in Slovakia (Central Europe): distribution, ecology and effect on native macrophyte assemblages. *Aquatic Invasions* 16: 617–636, <https://doi.org/10.3391/ai.2021.16.4.03>
- Butkuvienė J, Sinkevičienė Z, Naugžemys D, Žvingila D, Skridaila A, Bobrov AA (2020) Genetic diversity of aquatic *Ranunculus* (Batrachium, Ranunculaceae) in one river basin caused by hybridization. *Plants* 9: 1–17, <https://doi.org/10.3390/plants9111455>
- Ciocârlan V, Sarbu I, Stefan N, Marian T (1998) *Elodea nuttallii* (Planchon) H. St. John a new species in Romanian flora. *Buletinul Grdinii Botanice Iași* 6(1): 215–231
- Cook CDK, Urmí-König K (1985) A revision of the genus *Elodea* (Hydrocharitaceae). *Aquatic Botany* 21: 111–156, [https://doi.org/10.1016/0304-3770\(85\)90084-1](https://doi.org/10.1016/0304-3770(85)90084-1)
- Česnulevičius A (2008) Dainavos lyguma. Visuotinė lietuvių enciklopedija. [Dainava plain. Universal Encyclopaedia of Lithuania. <https://www.vle.lt/straipsnis/dainavos-lyguma/#:~:text=Atgal,Dainavos%20lyguma,-Algimantas%20%C4%8Cesnulevi%C4%8Dius/> (accessed 20 January 2022)
- Davydova AO, Orlova-Hudim KS, Shevchenko IV, Davydov DA, Dzerkal VM (2021) The first record of *Elodea nuttallii* (Hydrocharitaceae) in the Lower Dnipro River. *Ukrainian Botanical Journal* 78: 303–307, <https://doi.org/10.15407/ukrbotj78.04.303>
- Dembowska EA, Kamiński D, Wojciechowska A (2021) Phytoplankton response to the massive expansion of *Elodea nuttallii* (Planch.) H.St.John, 1920 in a floodplain lake of the Vistula River (Poland). *Aquatic Invasions* 16: 601–616, <https://doi.org/10.3391/ai.2021.16.4.02>
- Dubovik DV (2013) Hydrocharitaceae Juss. In: Parfenov VI (ed.), Flora Belarus, vascular plants, Vol.2, pp 94–102
- Dzhus MA (2014) Alien species of American origin on cranberry plantation in Belarus. *Botanicheski Zhurnal* 99: 540–554
- EU (2014) Regulation (EU) No 1143/2014 of the European Parliament and of the Council on the prevention and management of the introduction and spread of invasive alien species. *Official Journal of the European Union* L 317/35
- GBIF Secretariat (2021) GBIF Backbone Taxonomy. Checklist dataset <https://doi.org/10.15468/39omei> (accessed 16 May 2022)
- Georgiev V, Tsoneva S, Kenderov L, Trichkova T, Todorov M, Vladimirov V (2019) Distribution of *Elodea nuttallii*, an invasive alien species of EU concern, in Bulgaria. *Phytologia Balcanica* 25(3): 417–423
- Greulich S, Trémolières M (2006) Present distribution of the genus *Elodea* in the Alsatian Upper Rhine floodplain (France) with a special focus on the expansion of *Elodea nuttallii* St. John during recent decades. *Hydrobiologia* 570: 249–255, <https://doi.org/10.1007/s10750-006-0188-y>
- Grinberga L, Priede A (2010) *Elodea canadensis* Michx. in Latvia. *Acta Biologica Universitatis Daugavpiliensis* 10: 43–50
- Grudnik ZM, Jelenko I, Germ M (2014) Influence of abiotic factors on invasive behaviour of alien species *Elodea nuttallii* in the Drava River (Slovenia). *Annales de Limnologie - International Journal of Limnology* 50: 1–8, <https://doi.org/10.1051/limn/2013065>
- Havel JE, Kovalenko KE, Tomaz SM, Amalfitano S, Kats LB (2015) Aquatic invasive species: challenges for the future. *Hydrobiologia* 750: 147–170, <https://doi.org/10.1007/s10750-014-2166-0>



- Hrivnák R, Medvecká J, Baláži P, Bubíková K, Ořaheřová H, Svitok M (2019) Alien aquatic plants in Slovakia over 130 years: historical overview, current distribution and future perspectives. *NeoBiota* 49: 37–56, <https://doi.org/10.3897/neobiota.49.34318>
- Huotari T, Korpelainen H (2012) Complete chloroplast genome sequence of *Elodea canadensis* and comparative analyses with other monocot plastid genomes. *Gene* 508: 96–105, <https://doi.org/10.1016/j.gene.2012.07.020>
- Hussner A (2012) Alien aquatic plant species in European countries. *Weed Research* 52: 297–306, <https://doi.org/10.1111/j.1365-3180.2012.00926.x>
- Janauer GA, Gaberščik A, Vukov D (2007) Macrophytes and habitat characteristics of the Danube. Danube News, Bulletin of the International Association for Danube Research IAD 13/14, Wilhering, 15-18
- Josefsson M (2011) NOBANIS - Invasive Species Fact Sheet - *Elodea canadensis*, *Elodea nuttallii* and *Elodea callitrichoides* - Online Database of the European Network on Invasive Alien Species – NOBANIS. [www.nobanis.org](http://www.nobanis.org) (accessed 05 January 2021)
- Kamiński D (2010) *Elodea nuttallii* (Hydrocharitaceae) nowy gatunek w wodach Wisły. *Fragmenta Floristica and Geobotanica Polonica* 17(1): 182–184
- Kočić A, Horvatić J, Jelaska SD (2014) Distribution and morphological variations of invasive macrophytes *Elodea nuttallii* (Planch.) H. St. John and *Elodea canadensis* Michx in Croatia. *Acta Botanica Croatica* 73: 437–446, <https://doi.org/10.2478/botcro-2014-0011>
- Kukk T, Kull T, Luuk O, Mesipuu M, Saar P (2020) Atlas of the Estonian flora. Tartu: Estonian Seminatural Community Conservation Association, Institute of Agricultural and Environmental Sciences of the Estonian University of Life Sciences, 643 pp
- Madden TL, Tatusov RL, Zhang J (1996) Applications of network BLAST server. *Methodology of Enzymology* 266: 131–141, [https://doi.org/10.1016/S0076-6879\(96\)66011-X](https://doi.org/10.1016/S0076-6879(96)66011-X)
- Prokopuk M, Zub L (2019) Ecology of *Elodea* spp. in Ukraine. *Phytologia Balcanica* 25(3): 381–386
- Panasenko NN, Shcherbakov AV (2018) *Elodea nuttallii* (Planch.) H. St. John (Hydrocharitaceae), a new potentially invasive species for the Russian flora. *Bulletin of the Moscow Society of Naturalists. Department of Biology* 123(6): 58–59
- Pyšek P, Hulme PE, Simberloff D, Bacher S, Blackburn TM, Carlton JT, Dawson W, Essl F, Foxcroft LC, Genovesi P, Jeschke JM, Kühn I, Liebhold AM, Mandrak NE, Meyerson LA, Pauchard A, Pergl J, Roy HE, Seebens H, van Kleunen M, Vilf M, Wingfield MJ, Richardson D M (2020) Scientists' warning on invasive alien species. *Biological Reviews* 95: 1511–1534, <https://doi.org/10.1111/brv.12627>
- Reaser JK, Burgiel SW, Kirkey J, Brantley KA, Veatch SD, Rodríguez JB (2020) The early detection of and rapid response (EDRR) to invasive species: a conceptual framework and federal capacities assessment. *Biological Invasions* 22: 1–19, <https://doi.org/10.1007/s10530-019-02156-w>
- Rodríguez-Merino A, García-Murillo P, Cirujano S, Fernández-Zamudio R (2018) Predicting the risk of aquatic plant invasions in Europe: How climatic factors and anthropogenic activity influence potential species distributions. *Journal for Nature Conservation* 45: 58–71, <https://doi.org/10.1016/j.jnc.2018.08.007>
- Rybicky N, Voytek M (2013) Molecular techniques to distinguish morphologically similar *Hydrilla verticillata*, *Egeria densa*, *Elodea nuttallii*, and *Elodea canadensis*. *Journal of Aquatic Plants Management* 51: 94–102
- Sârbu A, Smarandache D, Janauer G, Pascale G (2006) *Elodea nuttallii* (Planchon) St. John - a competitive hydrophyte in the Romanian Danube River corridors. 36<sup>th</sup> International Conference of International Association for Danube Research. 4-8 September 2006, pp 107–111
- Simpson DA (1986) Taxonomy of *Elodea* Michx in the British Isles. *Watsonia* 16: 1–14
- Simpson DA (1988) Phenotypic plasticity of *Elodea nuttallii* (Planch.) H. St John and *Elodea canadensis* Michx in the British Isles. *Watsonia* 17: 121–132
- Simpson DA (1990) Displacement of *Elodea canadensis* Michx. by *Elodea nuttallii* (Planch.) H. St John in the British Isles. *Watsonia* 18(2): 173–177
- Steen B, Cardoso AC, Tsiamis K, Nieto K, Engel J, Gervasini E (2019) Modelling hot spot areas for the invasive alien plant *Elodea nuttallii* in the EU. *Management of Biological Invasions* 10: 151–170, <https://doi.org/10.3391/mbi.2019.10.1.10>
- Tamura K, Stecher G, Kumar S (2021) MEGA11: Molecular Evolutionary Genetics Analysis Version 11. *Molecular Biology and Evolution* 38: 3022–3027, <https://doi.org/10.1093/molbev/msab120>
- Thiébaud G, Di Nino F (2009) Morphological variations of natural populations of an aquatic macrophyte *Elodea nuttallii* in their native and in their introduced ranges. *Aquatic Invasions* 4: 311–320, <https://doi.org/10.3391/ai.2009.4.2.2>
- Uotila P (2009) Hydrocharitaceae. In: Euro+Med Plantbase - the information resource for Euro-Mediterranean plant diversity, <http://ww2.bgbm.org/EuroPlusMed> (accessed 6 January 2022)
- Vukov D, Boža P, Igić R, Anačkov G (2008) The distribution and the abundance of hydrophytes along the Danube River in Serbia. *Central European Journal of Biology* 3: 177–187, <https://doi.org/10.2478/s11535-007-0048-4>
- WFO (2021) *Elodea* Michx. <http://www.worldfloraonline.org/taxon/wfo-4000013242> (accessed 30 December 2021)