

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY INSTRUCTIONS BEFORE OPENING OR USING

EPOXICONAZOLE

500 FUNGICIDE

ACTIVE CONSTITUENT:
500 g/L EPOXICONAZOLE

GROUP 3 FUNGICIDE

Crops: Barley, Oats, Wheat
Controls/Suppresses: Eyespot, Leaf Rust, Leaf Scald, Net form of Net Blotch, Powdery Mildew, Septoria Nodorum Blotch, Septoria Leaf Blotch, Septoria Tritici Blotch, Spot form of Net Blotch, Stem Rust, Stripe Rust and Yellow Spot as per the Directions for Use

IMPORTANT: READ THIS LEAFLET BEFORE USE

RELYON



DIRECTIONS FOR USE

RESTRAINTS

DO NOT exceed a total rate of 250 mL/ha of Relyon EPOXICONAZOLE 500 FUNGICIDE in wheat and barley per growing season.

DO NOT apply more than two applications of Relyon EPOXICONAZOLE 500 FUNGICIDE in oats per growing season.

SPRAY DRIFT RESTRAINTS

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour, as measured at the application site.

DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product **MUST make an accurate written record** of the details of each spray application within 24 hours following application, and must **KEEP** this record for at least 2 years. The spray application details that must be recorded are:

1. Date with start and finish times of application;
2. Location address and paddock(s) sprayed;
3. Full name of this product;
4. Amount of product used per hectare and number of hectares treated;
5. Crop or situation and weed or pest;
6. Wind speed and direction during application;
7. Air temperature and relative humidity during application;
8. Nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application;
9. Name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in Table 1 below.

Table 1 – No-Spray Zones for Protection of the Aquatic Environment	
FOR GROUND and AERIAL APPLICATION	
From 3 to 20 kilometres per hour	30 metres

CROP	DISEASE	RATE/HA	CRITICAL COMMENTS
Barley	Eyespot (<i>Oculimacula yallundae</i>) Suppression only	125 mL	Apply to at risk crops when conditions favour disease development and prior to establishment of disease in the crop. Aim to apply from mid-tillering (Z25) to flag leaf emergence (Z39). Application prior to canopy closure may improve coverage, enhance efficacy and reduce lodging from eyespot infections. Applications beyond flag leaf emergence (Z39) may not provide any yield benefit. DO NOT apply later than Z59.
	Leaf Rust (<i>Puccinia hordei</i>) Leaf Scald (<i>Rhynchosporium secalis</i>), Net form of Net Blotch (<i>Pyrenophora teres</i> f.sp. <i>teres</i>), Spot form of Net Blotch (<i>Pyrenophora teres</i> f. sp. <i>maculata</i>)	65 to 125 mL	Apply when conditions favour disease development and prior to development of high levels of infection in the crop. Aim to apply from jointing (Z30) if disease is present and/or if conditions favour infection. Use the higher rate when disease is present on the top leaf or conditions are favourable for infection during these stages of crop development. Regularly monitor crops from 3-4 weeks post application. Repeat spraying may be required, particularly if infection occurs early. DO NOT apply after Z59 as yield responses are unlikely.
	Powdery Mildew (<i>Blumeria graminis</i> f.sp. <i>tritici</i>)	Apply at the first sign of the disease during the tillering stage (Z21-22). Regularly monitor the crop 3-4 weeks post-application. Apply a second spray if infection pressure persists. DO NOT apply after Z59 as yield responses are unlikely.	
Oats and Oaten Hay	Septoria Leaf Blotch (<i>Phaeosphaeria avenaria</i>)	65 mL	Apply when conditions favour disease development and preferably prior to development of high levels of disease in the crop. Aim to apply between stem elongation and completion of flag leaf emergence (Z31-41) if disease is present or conditions favour disease development. Repeat spraying may be required, particularly if infection occurs early. Regularly monitor the crop from 3-4 weeks after the first application for signs of re-infection. DO NOT apply later than Z59 as yield responses are unlikely.

CROP	DISEASE	RATE/HA	CRITICAL COMMENTS
Wheat	Eyespot (<i>Oculimacula yallundae</i>) Suppression only	125 mL	Apply to at risk crops when conditions favour disease development and prior to establishment of disease in the crop. Aim to apply from mid-tillering to flag leaf emergence (Z24-39). Application prior to canopy closure may improve coverage, enhance efficacy and reduce lodging from eyespot infections. Applications beyond flag leaf emergence (Z39) may not provide any yield benefit. A second application after a 28-day spray interval may be required, particularly if re-infection occurs. DO NOT apply later than Z59
	Leaf Rust (<i>Puccinia recondita</i>), Septoria Nodorum Blotch (<i>Phaeosphaeria nodorum</i>), Septoria Tritici Blotch (<i>Mycosphaerella graminicola</i>), Stem Rust (<i>Puccinia graminis</i>), Stripe Rust (<i>Puccinia striiformis</i>), Yellow Spot (<i>Pyrenophora tritici-repentis</i>)	65 to 125 mL	Apply when conditions favour disease development and prior to development of high levels of disease in the crop. Aim to apply during between stem elongation and complete ear emergence (Z31-59). Use the higher rate when disease is present on the top leaf or conditions are favourable for infection during these stages of crop development. Regularly monitor crops from 3-4 weeks post application. Repeat spraying may be required, particularly if infection occurs early. DO NOT apply later than Z59.
	Powdery Mildew (<i>Blumeria graminis</i> f.sp. <i>tritici</i>)		Apply at the first sign of the disease during the tillering stage (Z21-22). Regularly monitor the crop 3-4 weeks post-application. Apply a second spray if infection pressure persists. DO NOT apply later than Z59 as yield responses are unlikely.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 3 WEEKS AFTER APPLICATION.

TRADE ADVICE

Export of treated produce

Growers should note that Maximum Residue Limits (MRLs) or import tolerances do not exist in all markets for produce treated with Relyon Epoxiconazole 500 Fungicide. Additionally, some export markets have established MRLs different to those in Australia. If you are growing produce for export, please check with Relyon (Australia) Pty Ltd. or your produce exporter for the latest information on MRLs and import tolerances BEFORE using Relyon Epoxiconazole 500 Fungicide.

LIVESTOCK DESTINED FOR EXPORT MARKETS

The grazing withholding period only applies to stock slaughtered for the domestic market. Some export markets apply different standards. To meet these standards, ensure that in addition to complying with the grazing withholding period, the Export Slaughter Interval is observed before stock are sold or slaughtered.

EXPORT SLAUGHTER INTERVAL (ESI)

AFTER OBSERVING THE WITHHOLDING PERIOD FOR GRAZING OR CUTTING FOR STOCKFOOD, LIVESTOCK THAT HAVE BEEN GRAZED ON OR FED TREATED CROPS SHOULD BE PLACED ON CLEAN FEED FOR 3 DAYS PRIOR TO SLAUGHTER.

GENERAL INSTRUCTIONS:

Relyon Epoxiconazole 500 Fungicide is a broad spectrum contact and systemic fungicide. It is taken up via the stem and foliage and is translocated upwards and outwards to provide protection to new growth. Thorough coverage of the crop is necessary for best results.

Check the crop regularly for any symptoms of disease infection. Apply as soon as disease is present, and preferably prior to disease establishment. For effective management of eyespot disease, avoid tank mixtures with herbicides and apply prior to canopy closure. Application beyond flag leaf emergence (Z39) may not provide any yield benefit, particularly under dry seasonal conditions.

MIXING

Relyon Epoxiconazole 500 Fungicide is a suspension concentrate formulation. Add the product to the half-filled spray tank while agitating. Continue to agitate while topping up the tank and during spraying.

APPLICATION

Application of Relyon Epoxiconazole 500 Fungicide with Wetspray® 1000 (or equivalent formulation) at 200 mL/100 L may improve disease control in cereals, particularly at the lower application rate.

Ground Application: Apply in a water volume of between 50 and 100 L/ha, using flat fan nozzles operating at around 50 cm above the top of the crop. Use the higher water volume in crops with heavier canopies.

Aerial Application: Apply with suitable aircraft, set up and operated to apply fungicides to cereal crops in a minimum water volume of 20 L/ha.

FUNGICIDE RESISTANCE WARNINGS

GROUP 3 FUNGICIDE

Relyon Epoxiconazole 500 Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Relyon Epoxiconazole 500 Fungicide is a Group 3 fungicide. Some naturally occurring individual fungi resistant to Relyon Epoxiconazole 500 Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product or

other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of this product to control resistant fungi. Uses of Relyon Epoxiconazole 500 Fungicide are subject to CropLife Australia Fungicide Resistance Management guidelines. Refer to the relevant guidelines prior to using Relyon Epoxiconazole 500 Fungicide.

PRECAUTIONS

DO NOT use human flaggers/markers unless they are protected by engineering controls such as controlled cabs.

RE-ENTRY PERIOD

DO NOT allow entry into treated areas until the spray has dried. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist and a washable hat and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift over nearby susceptible plants, crops, cropping lands or pastures.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Highly toxic to aquatic organisms. DO NOT apply Relyon Epoxiconazole 500 Fungicide aerially to waterbodies, watercourses or wetlands. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL

1 L: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

2 L – 100 L: Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple-rinse container for disposal. Dispose of rinsate by adding it to the spray tank. Do not dispose of undiluted chemical on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be taken separately.

110 L: Store the original sealed container in a cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the container with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the container have been used, please return the container to the point of purchase.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

RELYON®

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW 2113.
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92
relyonaustralia.com.au

®RELYON is a Registered Trademark of Ruralco Holdings Limited