

Notes for genera: *Ascomycota*

Nalin N. Wijayawardene^{1,2,3,4} · Kevin D. Hyde^{1,2,3,4} · Kunhiraman C. Rajeshkumar⁵ · David L. Hawksworth^{6,7,8} · Hugo Madrid⁹ · Paul M. Kirk⁶ · Uwe Braun¹⁰ · Rajshree V. Singh⁵ · Pedro W. Crous¹¹ · Martin Kukwa¹² · Robert Lücking¹³ · Cletus P. Kurtzman¹⁴ · Andrey Yurkov¹⁵ · Danny Haelewaters^{16,17} · André Aptroot¹⁸ · H. Thorsten Lumbsch¹⁹ · Einar Timdal²⁰ · Damien Ertz²¹ · Javier Etayo²² · Alan J. L. Phillips²³ · Johannes Z. Groenewald¹¹ · Moslem Papizadeh²⁴ · Laura Selbmann²⁵ · Monika C. Dayarathne^{1,2} · Gothamie Weerakoon^{26,27} · E. B. Gareth Jones²⁸ · Satinee Suetrong²⁹ · Qing Tian^{2,3} · Rafael F. Castañeda-Ruiz³⁰ · Ali H. Bahkali³¹ · Ka-Lai Pang³² · Kazuaki Tanaka³³ · Dong Qin Dai³⁴ · Jariya Sakayaroj²⁹ · Martina Hujslová³⁵ · Lorenzo Lombard¹¹ · Belle D. Shenoy³⁶ · Ave Suija³⁷ · Sajeewa S. N. Maharachchikumbura³⁸ · Kasun M. Thambugala^{2,4,39} · Dhanushka N. Wanasinghe^{2,3,4} · Bharati O. Sharma⁴⁰ · Subhash Gaikwad⁴⁰ · Gargee Pandit⁴⁰ · Laura Zucconi²⁵ · Silvano Onofri²⁵ · Eleonora Egidi⁴¹ · Huzefa A. Raja⁴² · Rampai Kodsub⁴³ · Marcela E. S. Cáceres⁴⁴ · Sergio Pérez-Ortega⁴⁵ · Patrícia O. Fiúza⁴⁶ · Josiane Santana Monteiro⁴⁶ · Larissa N. Vasilyeva⁴⁷ · Roger G. Shivas⁴⁸ · Maria Prieto^{49,50} · Mats Wedin⁵⁰ · Ibai Olariaga⁵⁰ · Adebola Azeez Lateef⁵¹ · Yamini Agrawal⁵² · Seyed Abolhassan Shahzadeh Fazeli^{24,53} · Mohammad Ali Amoozegar⁵⁴ · Guo Zhu Zhao⁵⁵ · Walter P. Pfleigler⁵⁶ · Gunjan Sharma⁵⁷ · Magdalena Oset¹² · Mohamed A. Abdel-Wahab⁵⁸ · Susumu Takamatsu⁵⁹ · Konstanze Bensch^{11,60} · Nimali Indeewari de Silva^{1,2,3,61} · André De Kesel²¹ · Anuruddha Karunarathna^{2,3,4,62} · Saranyaphat Boonmee^{2,3} · Donald H. Pfister^{16,17} · Yong-Zhong Lu^{2,3} · Zong-Long Luo^{2,3,63} · Nattawut Boonyuen²⁹ · Dinushani A. Daranagama^{2,4} · Indunil C. Senanayake^{1,2,3} · Subashini C. Jayasiri^{2,64} · Milan C. Samarakoon^{2,61} · Xiang-Yu Zeng^{2,3,64} · Mingkwan Doilom² · Luis Quijada⁶⁵ · Sillma Rampadarath⁶⁶ · Gabriela Heredia⁶⁷ · Asha J. Dissanayake² · Ruvishika S. Jayawardana² · Rekhani H. Perera^{2,4,39} · Li Zhou Tang³⁴ · Chayanard Phukhamsakda^{1,2,3} · Margarita Hernández-Restrepo¹¹ · Xiaoya Ma^{2,3} · Saowaluck Tibpromma^{2,3,4} · Luis F. P. Gusmao⁴⁶ · Darshani Weerahewa⁶² · Samantha C. Karunarathna^{1,2,3,4}

Received: 29 June 2017/Accepted: 14 July 2017

© School of Science 2017

Abstract Knowledge of the relationships and thus the classification of fungi, has developed rapidly with increasingly widespread use of molecular techniques, over the past 10–15 years, and continues to accelerate.

This paper is dedicated to the memory of K. Walter Gams, Hubertus Antonius van der Aa, Larissa N. Vasilyeva, E. Punithalingam and Vadim A. Mel'nik whose efforts contributed greatly to mycology.

 Samantha C. Karunarathna
samanthakarunarathna@gmail.com

¹ Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Science, Kunming 650201, Yunnan, People's Republic of China

² Center of Excellence in Fungal Research, Mae Fah Luang University, Chiang Rai 57100, Thailand

³ World Agroforestry Centre, East and Central Asia, Kunming 650201, Yunnan, People's Republic of China

Several genera have been found to be polyphyletic, and their generic concepts have subsequently been emended. New names have thus been introduced for species which are phylogenetically distinct from the type species of particular genera. The ending of the separate naming of morphs of the same species in 2011, has also caused

⁴ Mushroom Research Foundation, 128 M.3 Ban Pa Deng T. Pa Pae, A. Mae Taeng, Chiang Mai 50150, Thailand

⁵ National Fungal Culture Collection of India (NFCCI), Biodiversity and Palaeobiology (Fungi) Group, Agharkar Research Institute, Pune, Maharashtra 411 004, India

⁶ Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3DS, UK

⁷ Department of Life Sciences, The Natural History Museum, Cromwell Road, London SW7 5BD, UK

changes in fungal generic names. In order to facilitate access to all important changes, it was desirable to compile these in a single document. The present article provides a list of generic names of *Ascomycota* (approximately 6500 accepted names published to the end of 2016), including those which are lichen-forming. Notes and summaries of the changes since the last edition of ‘Ainsworth & Bisby’s Dictionary of the Fungi’ in 2008 are provided. The notes include the number of accepted species, classification, type species (with location of the type material), culture availability, life-styles, distribution, and selected publications that have appeared since 2008. This work is intended to provide the foundation for updating the ascomycete component of the “Without prejudice list of generic names of *Fungi*” published in 2013, which will be developed into a list of protected generic names. This will be subjected to the XIXth International Botanical Congress in Shenzhen in July 2017 agreeing to a modification in the rules relating to protected lists, and scrutiny by procedures determined by the Nomenclature Committee for Fungi (NCF). The previously invalidly published generic names *Barriopsis*, *Collophora* (as *Collophorina*), *Cryomyces*, *Dematiopleospora*, *Heterospora* (as *Heterosporicola*), *Lithophila*, *Palmomyces* (as *Palmaria*) and *Saxomyces* are validated, as are two previously invalid family names, *Bartaliniaceae* and *Wiesneriomycetaceae*. Four species of *Lalaria*, which were invalidly published are transferred to *Taphrina* and validated as new combinations.

⁸ Jilin Agricultural University, Changchun 130118, Jilin, People’s Republic of China

⁹ Centro de Genómica y Bioinformática, Facultad de Ciencias, Universidad Mayor de Chile, Camino La Pirámide 5750, Huechuraba, Santiago, Chile

¹⁰ Martin-Luther-Universität, Institut für Biologie, Bereich Geobotanik und Botanischer Garten, Herbarium, Neuwerk 21, 06099 Halle (Saale), Germany

¹¹ Westerdijk Fungal Biodiversity Institute, Uppsalaalaan 8, 3584 CT Utrecht, The Netherlands

¹² Department of Plant Taxonomy and Nature Conservation, Faculty of Biology, University of Gdańsk, Wita Stwosza 59, 80-308 Gdańsk, Poland

¹³ Botanic Garden and Botanical Museum, Königin -Luise-Str. 6–8, 14195 Berlin, Germany

¹⁴ Mycotoxin Prevention and Applied Microbiology Research Unit, National Center for Agricultural Utilization Research, Agricultural Research Service, U.S. Department of Agriculture, Peoria, IL 61604, USA

¹⁵ Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, Inhoffenstraße 7B, 38124 Brunswick, Germany

¹⁶ Farlow Reference Library and Herbarium of Cryptogamic Botany, Harvard University, 22 Divinity Avenue, Cambridge, MA 02138, USA

Catenomycopsis Tibell & Constant. is reduced under *Chaenothecopsis* Vain., while *Dichomera* Cooke is reduced under *Botryosphaeria* Ces. & De Not. (Art. 59).

Keywords Article 59 · Classification · One name · Pleomorphic fungi · Taxonomy

Introduction

Knowledge of the taxonomy of fungi has been changing rapidly since the last edition of *Ainsworth & Bisby’s Dictionary of the Fungi* (Kirk et al. 2008), as a consequence of new information from DNA and protein sequence studies, and the integration of asexual and sexual morphs into a unified system. Phylogenetic affiliations and taxonomic placements of a large number of genera are being clarified, and at the same time species complexes and generic circumscriptions are being resolved (Damm et al. 2012a, b; 2013; Choi et al. 2013a; Giraud et al. 2013; Nilsson et al. 2014; Udayanga et al. 2014a). Sequences are also increasingly being obtained from genera representing fungi with ecologies not previously well-represented in the system, such as rock-inhabiting (Egidi et al. 2014; Ertz et al. 2014b; Pérez-Ortega et al. 2015) and lichenicolous fungi (Trakunyingcharoen et al. 2014; Sipman and Raus 2015; Suija et al. 2015b, c), or from many which are

¹⁷ Department of Organismic and Evolutionary Biology, Harvard University, 22 Divinity Avenue, Cambridge, MA 02138, USA

¹⁸ ABL Herbarium, G.v.d. Veenstraat 107, 3762 XK Soest, The Netherlands

¹⁹ Science & Education, The Field Museum, 1400 S. Lake Shore Drive, Chicago, IL 60605, USA

²⁰ Natural History Museum, University of Oslo, Blindern, P.O. Box 1172, 0318 Oslo, Norway

²¹ Botanic Garden Meise - Research Department, Nieuwelaan 38, 1860 Meise, Belgium

²² Department of Biology, IES Zizur, Ronda S. Cristóbal 196, 31180 Zizur Mayor, Navarra, Spain

²³ Faculty of Sciences, Biosystems and Integrative Sciences Institute (BioISI), University of Lisbon, Campo Grande, 1749-016 Lisbon, Portugal

²⁴ Microorganisms Bank, Iranian Biological Resource Center (IBRC), Academic Center for Education, Culture and Research (ACECR), Tehran, Iran

²⁵ DEB, University of Tuscia, Largo dell’Università, Viterbo, Italy

²⁶ Scientific Affiliate, Integrative Research Center, Science & Education, The Field Museum, Chicago, IL, USA

²⁷ National Institute of Fundamental Studies, Kandy, Sri Lanka

uncultured or unculturable, the latter through new generation sequencing (Hodkinson and Lendemer 2013). In addition, neo- and epitypifications are enabling the application of historic species and genera to be accommodated into molecularly based systems (Zhang et al. 2008a, b; Ariyawansa et al. 2014b). Thus, major progress towards a natural classification system is being made (Hyde et al. 2017).

The ending of dual nomenclature for pleomorphic non-lichenized ascomycetes and basidiomycetes in 2011 (McNeill et al. 2011) has also led to a need to select the name to be adopted for pleomorphic genera. Lists of names are now being proposed for protection or suppression by working groups established under the auspices of the International Commission on the Taxonomy of Fungi (ICTF). The proposals from the working groups are being published and will then be scrutinized by the Nomenclature Committee for Fungi (NCF) and the General Committee (Art. 14.13) established by the last International Botanical Congress.

The problem is especially acute in the phylum *Ascomycota*, the largest in kingdom *Fungi*, with some 15 classes, 68 orders, 327 families, ca. 6300 genera, and over 64 000 species (Kirk et al. 2008). However, 1000–1500 new taxa are introduced each year, and our understanding of the *Ascomycota* has been developing rapidly. Keeping track of all the changes and the incorporation of so much new knowledge is a major challenge, though some

important compilations have been prepared, for example those of Hyde et al. (2013) and Wijayawardene et al. (2014c) for *Dothideomycetes*, and Maharachchikumbura et al. (2015b, 2016b) for *Sordariomycetes*, which considered both asexually and sexually typified genera. Molecular clock analyses has also provided additional evidence for the placement of genera at the family, order and class levels (Phukhamsakda et al. 2016b; Samarakoon et al. 2016; Divakar et al. 2017; Hongsanan et al. 2016c, 2017; Hyde et al. 2017; Liu et al. 2017). There has also been a major attempt to produce an overall system for *Ascomycota* referring to accepted families down to genus level (Jaklitsch et al. 2016a).

The long-running series *Outline of the Ascomycota* (1982–1986) and *Notes on Ascomycete Systematics* (1986–2010) aimed to provide a current awareness system of novel taxa and changes proposed, and provided 5113 notes. This has not, however, been continued to include asexually typified genera. Here we start to redress this problem, taking the last edition of the *Dictionary* (Kirk et al. 2008) as our starting point. In addition to recent nomenclature changes, we include typification details of the type species of genera (including lecto-, neo- and epitypifications), information on the availability of cultures, and references to accessible sequence data. References to recently established DNA barcodes for plant pathogenic genera are also included where available.

²⁸ 33 B St. Edwards Road, Southsea PO5 3DH, UK

²⁹ National Center for Genetic Engineering and Biotechnology (BIOTEC), 113 Thailand Science Park, Phahonyothin Road, Khlong Nueng, Khlong Luang, Pathum Thani 12120, Thailand

³⁰ Instituto de Investigaciones Fundamentales en Agricultura Tropical, ‘Alejandro de Humboldt’ (INIFAT), Santiago de Las Vegas, Habana, Cuba

³¹ Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box: 2455, Riyadh 1145, Saudi Arabia

³² Institute of Marine Biology and Center of Excellence for the Oceans, National Taiwan Ocean University, 2 Pei-Ning Road, Keelung 20224, Taiwan

³³ Faculty of Agriculture and Life Science, Hirosaki University, 3 Bunkyo-cho, Hirosaki, Aomori 036-8561, Japan

³⁴ Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qijing Normal University, Qijing 655011, Yunnan, China

³⁵ Institute of Microbiology ASCR, Videnska 1083, 142 20 Prague 4, Czech Republic

³⁶ CSIR-National Institute of Oceanography Regional Centre, 176, Lawsons Bay Colony, Visakhapatnam – 17, Andhra Pradesh, India

³⁷ Institute of Ecology and Earth Sciences, University of Tartu, Tartu, Estonia

³⁸ Department of Crop Sciences, College of Agricultural and Marine Sciences, Sultan Qaboos University, PO Box 34, 123 Alkhoud, Oman

³⁹ Guizhou Key Laboratory of Agricultural Biotechnology, Guizhou Academy of Agricultural Sciences, Guiyang 550006, Guizhou, People’s Republic of China

⁴⁰ Biodiversity and Palaeobiology (Lichens) Group, Agharkar Research Institute, Pune, Maharashtra 411 004, India

⁴¹ Department of Physiology, Anatomy and Microbiology, La Trobe University, Bundoora, VIC 3086, Australia

⁴² Department of Chemistry and Biochemistry, University of North Carolina at Greensboro, Greensboro, NC 27402, USA

⁴³ Microbiology Program, Faculty of Science and Technology, Pibulsongkram Rajabhat University, Phitsanulok 65000, Thailand

⁴⁴ Departamento de Biociências, Universidade Federal de Sergipe, Itabaiana, Sergipe CEP: 49500-000, Brazil

⁴⁵ Real Jardín Botánico (CSIC), Plaza Murillo 2, 28014 Madrid, Spain

⁴⁶ Departamento de Ciências Biológicas, Laboratório de Micologia, Universidade Estadual de Feira de Santana, Av. Transnordestina, s/n - Bairro: Novo Horizonte, Feira de Santana, BA CEP: 44036-900, Brazil

In addition to providing an overview of current knowledge on accepted genera of ascomycetes, this work is also intended to provide the foundation for updating the ascomycete component of the “Without prejudice list of generic names of *Fungi*” published in 2013 (Kirk et al. 2013), which is anticipated will be developed into a list of protected generic names. This will, however, be subjected to the XIXth International Botanical Congress in Shenzhen in July 2017 agreeing to a modification in the rules relating to protected lists of names supported by the 10th International Mycological Congress in 2014 (Redhead et al. 2014; Hawksworth 2015), and scrutiny by procedures determined by the Nomenclature Committee for Fungi (NCF).

In this study, we list 6540 accepted genera of ascomycetes with an estimate of over 80,000 species. They belong in 485 families, 115 orders and 17 classes. We have attempted to include all genera and place them in a natural classification using the most recent evidence and publications. We however, have surely missed some entries and may not have found all of the latest data. We therefore regard this as a working paper and will continually be updated it in our webpage www.fungalgenera.org. Suggestions for changes or omissions will be welcome and readers can contact the corresponding author.

Materials and Methods

Data collection

Collecting data on existing names of taxa in use was based on Kirk et al. (2008, 2013) and Species Fungorum (2016). The works of Sutton (1977, 1980) and Seifert et al. (2011) were also used for further clarification of nomenclatural issues for asexually typified genera. Previous outlines of sexually typified genera (i.e. Lumbsch and Huhndorf 2010) and asexually typified genera (i.e. Hyde et al. 2011; Wijayawardene et al. 2012, 2017) have been used to extract taxonomic placements. For the taxonomic placement of lichen-forming genera, the compilation of Lücking et al. (2017a, b) was used. The citation of the articles given in each entry was decided by the author who provided the entry and is not standardized. Synonyms of generic names (if less than 10) are also provided based on recent publications and Species Fungorum (2017).

One fungus: one name

Recent proposals on names to be selected for use in non-lichenized pleomorphic ascomycete genera were gathered from the reports of ICTF working groups (Table 1).

⁴⁷ Institute of Biology & Soil Science, Far East Branch of the Russian Academy of Sciences, Vladivostok, Russia 690022

⁴⁸ Centre for Crop Health, Institute for Agriculture and the Environment, University of Southern Queensland, Toowoomba, QLD 4350, Australia

⁴⁹ Departamento de Biología y Geología, Física y Química Inorgánica, Universidad Rey Juan Carlos, C/Tulipan s/n, 28933 Mostoles, Madrid, Spain

⁵⁰ Department of Botany, Swedish Museum of Natural History, PO Box 50007, 10405 Stockholm, Sweden

⁵¹ Department of Plant Biology, Faculty of Life Sciences, University of Ilorin, Ilorin, Nigeria

⁵² Shahid Rajguru College of Applied Sciences for Women, Vasundhara Enclave, East Delhi 110096, India

⁵³ Department of Molecular and Cellular Biology, Faculty of Basic Sciences and Advanced Technologies in Biology, University of Science and Culture, Tehran, Iran

⁵⁴ Extremophiles Laboratory, Department of Microbiology, Faculty of Biology and Center of Excellence in Phylogeny of Living Organisms, College of Science, University of Tehran, Tehran, Iran

⁵⁵ College of Biological Sciences and Biotechnology, Beijing Forestry University, Beijing 100083, People's Republic of China

⁵⁶ Department of Biotechnology and Microbiology, University of Debrecen, Debrecen, Hungary

⁵⁷ Regional Centre for Biotechnology, NCR Biotech Science Cluster, Faridabad, Haryana 121001, India

⁵⁸ Department of Botany and Microbiology, Faculty of Science, Sohag University, Sohag, Egypt

⁵⁹ Graduate School of Bioresources, Mie University, 1577 Kurima-Machiya, Tsu 514-8507, Japan

⁶⁰ Botanische Staatssammlung München, Menzinger Straße 67, 80638 Munich, Germany

⁶¹ Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

⁶² Department of Botany, The Open University of Sri Lanka, Nawala, Nugegoda, Sri Lanka

⁶³ College of Agriculture and Biological Sciences, Dali University, Dali 671003, Yunnan, People's Republic of China

⁶⁴ Engineering Research Center of Southwest Bio-Pharmaceutical Resources, Ministry of Education, Guizhou University, Guiyang 550025, Guizhou, People's Republic of China

⁶⁵ Departamento de Botánica, Ecología y Fisiología Vegetal, Universidad de La Laguna, 38200 apdo. 456 Avda. Francisco Sánchez s/n, Tenerife, Canary Islands, Spain

⁶⁶ Department of Agriculture and Food Science, Faculty of Agriculture, University of Mauritius, Réduit, Mauritius

Table 1 ICTF working groups and reports

Taxonomic group	Reference
<i>Diaporthales</i>	Rossman et al. (2015a)
<i>Dothideomycetes</i>	Hyde et al. (2013), Wijayawardene et al. (2014c), Rossman et al. (2015b, 2016a)
<i>Hypocreales</i>	Rossman et al. (2013b)
<i>Leotiomycetes</i>	Johnston et al. (2014b)
<i>Magnaportheales</i>	Zhang et al. (2016d)
Miscellaneous genera	Rossman et al. (2016a)
<i>Pezizomycetes</i>	Healy et al. (2016)
<i>Sordariomycetes</i> – (except <i>Diaporthales</i> , <i>Hypocreales</i> , and <i>Magnaportheales</i>)	Maharachchikumbura et al. (2015b, 2016a), Réblová et al. (2016c)
<i>Xylariales</i>	Stadler et al. (2014), Daranagama et al. (2017), Wendt et al. (2017)

Results

Validating of invalid families

Bartaliniaceae Wijyaw., Maharachch., P.M. Kirk & K.D. Hyde

Index Fungorum number: 551079

= *Bartaliniaceae* Wijyaw., Maharachch. & K.D. Hyde,

Fungal Diversity **73**: 85. 2015, Nom. inval., Art. 42.1 (Melbourne)

Type: *Bartalinia* Tassi, Bulletin Labor. Orto Bot. de R. Univ. Siena 3: 4 (1900)

Description: see Senanayake et al. (2015)

Notes: Senanayake et al. (2015) introduced *Bartaliniaceae* but it is invalid (Art. 42.1) as the identifier cited in the protologue, (i.e. IF511183), was not issued for the name published (i.e. *Bartaliniaceae*). This identifier was issued for *Boletellus exiguus* T.W. Henkel & Fulgenzi 2008. A new identifier was obtained (i.e. 551079), thus *Bartaliniaceae* is herein validly published. However, Hongsanan et al. (2017a) treated this family as a synonym of *Sporocadaceae*.

Wiesneriomycetaceae Suetrong, Rungjindamai, Somrithipol. & E.B.G. Jones

Index Fungorum number: IF550306

= *Wiesneriomycetaceae* Suetrong, Rungjindamai, Somrithipol. & E.B.G. Jones, *Phytotaxa* 176(1): 288. 2014; Nom. inval., Art. 42.1 (Melbourne)

Type: *Wiesneriomycetes* Koord., Verh. K. Akad. Wet., tweede sect. 13(4): 246 (1907) IF 10438

Description: see Suetrong et al. (2014)

Notes: Suetrong et al. (2014) introduced *Wiesneriomycetaceae* but it is invalid (Art. 42.1) as the identifier cited in the protologue, (i.e. MB809097), was not issued for the name published (i.e. *Wiesneriomycetaceae*). This identifier was issued for *Hypogymnia papilliformis* McCune, Tchabanenko & X.L. Wei 2015. A new identifier was obtained (i.e. IF550306), thus *Wiesneriomycetaceae* is herein validly published.

Notes for genera

Aabaarnia Diederich 2014, *Ostropales* genera incertae sedis, *Lecanoromycetes*, one species, type: *A. siphulicola* Diederich, asexual morph is unknown, lichenicolous, terrestrial, New Zealand and Tasmania, see Diederich (2014), holotype: UPS, James 992/1, sequences are unavailable.

Aaosphaeria Aptroot 1995, *Dacampiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. arxii* (van der Aa) Aptroot, asexual morph coelomycetous, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Dacampiaceae*), Kirk et al. (2013; genus accepted), sequence available for the type: KU158145 (ITS) (unpublished).

Abropelta B. Sutton 1986, *Ascomycota* genera incertae sedis, one species, type: *A. fusariooides* B. Sutton, coelomycetous, sexual morph unknown, caulicolous, terrestrial, India, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: IMI 236263.

Abrothallus De Not. 1845 (= *Abrothallomyces* Cif. & Tomas. 1953; = *Phymatopsis* Tul. ex Trevis. 1857; = *Pseudolecidea* Marchand 1896; = *Vouauxiomycetes* Dyko & D. Hawksw. 1979 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Abrothallaceae*, *Abrothallales*,

⁶⁷ Laboratory of Microfungi, Instituto de Ecología A. C., Xalapa, Ver., Mexico

Dothideomycetes, c. 40 species, type: *A. bertianus* De Not., obligately lichenicolous, parasymbiotic or parasitic on macrolichens from several families of lichen-forming fungi, sexual morph with apothecioïd ascocarps, asexual morph coelomycetous, some species are known only either by their asexual or sexual morph, terrestrial, worldwide, see Pérez-Ortega et al. (2011b, 2014; sexual-aseexual connection, phylogeny), Suija et al. (2015a; phylogeny, key for species on *Peltigerales*), Rossman et al. (2016a; nomenclature), holotype of the type: likely in VER or RO. the species growing on *Parmeliaceae* are in urgent need of revision, preliminary molecular data suggest that current species delimitation, largely based on the host identity is inaccurate (Pérez-Ortega and Suija, data not published).

Asconditella Vězda 1965, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, twelve species, type: *A. sphagnorum* Vězda & Poelt, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Abyssomyces Kohlm. 1970, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. hydrozoicus* Kohlm., saprobes, asexual morph unknown, marine, Scotland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Acallomyces Thaxt. 1903, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *A. homalotae* Thaxt., asexual morph unknown, biotrophic on *Coleoptera* in decaying agarics (Staphylinidae), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), morphologically related to *Polyascomyces* Thaxter, cultures and sequences are unavailable.

Acanthogymnomyces Udagawa & Uchiy. 2000, *Eoterfeziaceae*, *Ascomycota* families *incertae sedis*, one species, type: *A. terrestris* Udagawa & Uchiy., asexual morph unknown, from soil, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), sequences are unavailable, holotype of the type: CBM SUM3162.

Acanthohelicospora Boonmee & K.D. Hyde 2014, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, four species, type: *A. pinicola* Boonmee & K.D. Hyde, saprobes, worldwide, asexual morph hyphomycetous, see Boonmee et al. (2014b; taxonomy, phylogeny), Rossman et al. (2016b; plant pathogens, new combinations), holotype and ex-type: MFLU 10-0049, MFLUCC 10-0116 =BCC 52036.

Acanthonitschkea Speg. 1908 (= *Euacanthe* Theiss. 1917; = *Fitzpatrickia* Cif. 1928; = *Hystrix* Alstrup & Olech 1993; = *Petelotia* Pat. 1924; = *Sydowninula* Petr. 1923 *fide* Species Fungorum 2017), *Nitschkiaceae*, *Coronophorales*, *Sordariomycetes*, nine species, type: *A. argentinensis* Speg., asexual morph acermonium-like, saprobes and on

lichens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Mugambi and Huhndorf (2010; DNA), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Acanthophiobolus Berl. 1893 (= *Ophiobolus* subgen. *Ophiochaeta* Sacc. 1883; = *Ophiochaeta* (Sacc.) Sacc. 1895; = *Ophiophaeria* Kirschst. 1907 [1906]; = *Ophiotrichia* Berl. 1893 *fide* Species Fungorum 2017), *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, three species, type: *A. helminthosporus* (Rehm) Berl., asexual morph unknown, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline), Boonmee et al. (2011, 2014b; accepted as a genus in *Tubeufiaceae*), Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Acanthorus Bat. & Cavalc. 1967, *Ascomycota* genera *incertae sedis*, one species, type: *A. maranhensis* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as a genus in *Dothideomycetes*), Wijayawardene et al. (2017) and in here we treat this genus as *Ascomycota* genera *incertae sedis* since cultures and sequences are unavailable.

Acanthosphaeria Kirschst. 1939, *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, two species, type: need typification, asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Acanthostigma De Not., 1863, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, 64 species, type: *A. perpusillum* de Not., saprobes on dead wood, widespread, asexual morph hyphomycetous, see Lumbsch and Huhndorf (2010; outline), Boonmee et al. (2011, 2014b; accepted as a genus in *Tubeufiaceae*), Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available for three species, strains available for type: UAMH 7237.

Acanthostigmella Höhn., 1905, *Dothideomycetes* genera *incertae sedis*, six species, type: *A. genuflexa* Höhn., asexual morph unknown, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Tubeufiaceae*), Boonmee et al. (2011; accepted as a genus in *Tubeufiaceae*), Hyde et al. (2013) and Boonmee et al. (2014b) did not accept as a genus in *Tubeufiaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Acanthostigmmina Höhn. 1909, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, three species, type: *A. minuta* (Fuckel) Clem. & Shear, saprobes on decaying

wood, cosmopolitan, see Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; taxonomy), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Acanthotheciella Höhn. 1911 (= *Acanthotheca* Clem. & Shear 1931 *fide Species Fungorum* 2017), *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. barbata* (Pat.) Höhn., asexual morph unknown, on dead scale insects, terrestrial, Asia, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Acanthothecis Clem. 1909 (= *Acanthographina* (Vain.) Walt. Watson 1929; = *Acanthographis* (Vain.) Walt. Watson 1929; = *Acanthotheciopsis* Zahlbr. 1923 [1924]; = *Acanthothecium* Vain. 1890; = *Acanthothecium* sect. *Acanthographina* Vain. 1890; = *Acanthothecium* sect. *Acanthographis* Vain. 1890; *Acanthothecomyces* Cif. & Tomas. 1953; = *Psorographis* Clem. 1909 *fide Species Fungorum* 2017), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, c. 56 species, type: *A. hololeuroides* (Nyl.) Staiger & Kalb, all with ascocarps, lichenized (*Trenthepohlia*), terrestrial (mainly corticolous), pantropical-subtropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rivas Plata et al. (2013; taxonomy), Lumbsch et al. (2014a, b; taxonomy), Lücking et al. (2017a; classification), polyphyletic, sequences are available.

Acanthotrema Frisch 2006, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *A. brasiliandum* (Hale) Frisch, lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rivas Plata et al. (2010a, 2012a; key), Sipman et al. (2012; key), sequences are available.

Acarella Syd. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *A. costaricensis* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Central America, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Acarellina Bat. & H. Maia 1960, *Ascomycota* genera *incertae sedis*, one species, type: *A. psidii* Bat. & H. Maia, coelomycetous, sexual morph unknown, epiphytes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acaroconium Kocourk. & D. Hawksw. 2008, *Ascomycota* genera *incertae sedis*, one species, type: *A. punctiforme* Kocourk. & D. Hawksw., coelomycetous, sexual morph unknown, lichenicolous, terrestrial, Europe, North America, see Kocourková and Hawksworth (2008; taxonomy), Wijayawardene et al. (2016b, 2017; taxonomy),

outline), holotype of the type: PRM 8926543, cultures and sequences are unavailable, some lichenicolous *Microsphaeropsis* species may belong to this genus.

Acarocybe Syd. 1937, *Ascomycota* genera *incertae sedis*, three species, type: *A. hansfordii* Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Central America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acarocybellina Subram. 1992, *Ascomycota* genera *incertae sedis*, one species, type: *A. arengae* (Matsush.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acarocybiopsis J. Mena, A. Hern.-Gut. & Mercado 1999, *Ascomycota* genera *incertae sedis*, one species, type: *A. cubensis* J. Mena, A. Hern. Gut. & Mercado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acaropeltis Petr. 1937, *Ascomycota* genera *incertae sedis*, one species, type: *A. costaricensis* Petr., coelomycetous, sexual morph unknown, on living leaves, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acarospora A. Massal. 1852, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, c. 200 species, type: *A. schleicheri* (Ach.) A. Massal. all with ascocarps, lichenized (green algae), terrestrial (mostly saxicolous or terricolous), cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Westberg et al. (2015; phylogeny), Lücking et al. (2017a; classification), polyphyletic, sequences are available.

Acarosporina Sherwood 1977, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *A. monilifera* (Ellis & Everh.) Sherwood, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Acaulium Sopp 1912 (= *Acaulium* Sopp 1912; = *Masonry* G. Sm. 1952; = *Masoniella* G. Sm. 1952; = *Phaeoscopulariopsis* M. Ota 1928 *fide Species Fungorum* 2017), *Microascaceae*, *Microascales*, *Sordariomycetes*, three species, type: *A. albonigrescens* Sopp, hyphomycetous, the genus was treated as a synonym of *Scopulariopsis* Bainier 1907 (*fide* Seifert et al. 2011), Sandoval-Denis et al. (2016b; phylogeny, reinstated the genus, new species), cultures and sequences are available.

Acephala Grünig & T.N. Sieber 2005, *Vibrisseaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *A. applanata* Grünig & T.N. Sieber, hyphomycetous, sexual morph unknown, from fine roots of *Pinus*, terrestrial, Germany, Switzerland, see Munzenberger et al. (2009; morphology, new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), ex-type strain of the type: CBS 109321.

Acerbiella Sacc. & D. Sacc. 1905, *Sordariomycetes* genera *incertae sedis*, two species, type: *A. macrospora* (Rick) Sacc. & D. Sacc., asexual morph unknown, saprobes, terrestrial, South America, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Acerviclypeatus Hanlin 1990, *Ascomycota* genera *incertae sedis*, one species, type: *A. poriformans* Hanlin, coelomycetous, sexual morph unknown, on living leaves of *Vaccinium*, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acervuloseptoria Crous & Jol. Roux 2014, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *A. ziziphicola* Crous & Jol. Roux, coelomycetous, sexual morph unknown, associated with leaf spots, terrestrial, South Africa, see Crous et al. (2014c, 2015c; morphology and phylogeny), holotype and ex-type culture of type: CBS H-21723, CPC 23707.

Acervus Kanouse 1938 (= *Phaedropezia* Le Gal 1953 *fide* Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, eight species, type: *A. aurantiacus* Kanouse, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhuang et al. (2011; morphology, taxonomy), Kirk et al. (2013; genus accepted), Ekanayaka et al. (2016d; new species, China), Jaklitsch et al. (2016a; classification), cultures and sequences are available but lacking for the type species.

Achaetobotrys Bat. & Cif. 1963, *Antennulariellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *A. affinis* (L.R. Fraser) Bat. & Cif., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Antennulariellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Achaetomium J.N. Rai, J.P. Tewari & Mukerji 1964, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, 13 species, asexual morph unknown, type: *A. globosum* J.N. Rai & J.P. Tewari, saprobes, terrestrial, Asia, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequence are available.

Achorella Theiss. & Syd. 1915, *Dothideomycetes* genera *incertae sedis*, ten species, type: *A. ametableta* (Rehm) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Achorodothis Syd. 1926, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *A. poasensis* Syd., asexual morph unknown, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mycosphaerellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Achoropeltis Syd. 1929, *Ascomycota* genera *incertae sedis*, one species, type: *A. modesta* Syd., coelomycetous, saprobes, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acroceratosphaeria Réblová, J. Fourn., K.D. Hyde & Ranghoor 2010, *Pisorisporiaceae*, *Pisorisporiales*, *Sordariomycetes*, two species, type: *A. potamia* Réblová, Fournier & K.D. Hyde, asexual morph unknown, saprobes, aquatic, Europe, see Réblová et al. (2010; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype and ex-type of the type: PRM 915719, CBS 125414.

Achroistachys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, six species, type: *A. humicola* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, from soil, terrestrial, cosmopolitan, see Lombard et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: CBS H-22420, CBS 868.73.

Aciascus Doweld 2013, *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *A. purpureus* (Thaxt. ex Currah) Doweld, asexual morph unknown, saprobes, USA, see Index Fungorum (2013; introduced to replace *Acitheca* Currah 1985), Wijayawardene et al. (2017; outline).

Aciculonidium D.S. King & S.C. Jong 1976, *Metschnikowiaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *A. aculeatum* (Phaff, M.W. Mill. & Shifrine) D.S. King & S.C. Jong, asexual morph unknown (budding of cells and formation of true hyphae with blastoconidia), saprophytes, isolated from fruit flies (*Drosophila* spp.), terrestrial, California, USA, see Kirk et al. (2013; genus accepted), cultures and sequences are available.

Aciculopsora Aptroot & Trest 2006, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *A. salmonaea* Aptroot & Trest, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf

(2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), holotype of the type: CR, Lücking 17543, cultures and sequences are unavailable.

Aciculosporium I. Miyake 1908 (= *Albomyces* I. Miyake ex I. Hino 1908; = *Mitosporium* Clem. & Shear 1931 *fide Species Fungorum* 2017), *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, seven species, type: *A. take* I. Miyake, asexual morph unknown, saprobes, terrestrial, Asia, see Tanaka (2009; in situ visualization), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Acicuseptoria Quaedv., Verkley & Crous 2013, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. rumicis* Quaedv., Verkley & Crous, coelomycetous, sexual morph unknown, saprobes, one species, France, see Quaedvlieg et al. (2013; morphology and phylogeny), holotype and ex-type culture of type: CBS H-18163, CPC 522.78.

Acidea Hujsová & M. Kolařík 2014, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. extrema* Hujsová & M. Kolařík, hyphomycetous, sexual morph unknown, saprobes, terrestrial, aquatic, Europe, Antarctica, see Hujsová et al. (2014; morphology, phylogeny), holotype and ex-type of the type: PRM 922617, CBS 136258.

Acidiella Hujsová & M. Kolařík 2013, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *A. bohemica* Hujsová & M. Kolařík, hyphomycetous, sexual morph unknown, saprobes, terrestrial and aquatic, Europe, Australia, see Hujsová et al. (2013; morphology, phylogeny), Kolařík et al. (2015a; new species), holotype and ex-type of cultures of the type: PRM 859933, CBS 132721.

Acidomelania E. Walsh & N. Zhang 2014, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. panicicola* E. Walsh & N. Zhang, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Walsh et al. (2014; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: RUTPP 61R8.

Acidomyces B.J. Baker, M.A. Lutz, S.C. Dawson, P.L. Bond & Banfield ex Selbmann, de Hoog & De Leo 2008 (= *Acidomyces* B.J. Baker, M.A. Lutz, S.C. Dawson, P.L. Bond & Banfield, Appl. 2004 *fide Species Fungorum* 2017), *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *A. acidophilus* (Sigler & J.W. Carmich.) Selbmann, de Hoog & De Leo, hyphomycetous, sexual morph unknown, from soil, terrestrial, aquatic, Worldwide, see Selbmann et al. (2008; morphology; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017;

outline), Hujsová et al. (2013; taxonomy), holotype and ex-type of the type: UAMH 3460, CBS 270.74.

Acidothrix Hujsová & M. Kolařík 2014, *Amplistromataceae*, *Amplistromatales*, *Sordariomycetes*, one species, type: *A. acidophila* Hujsová & M. Kolařík, hyphomycetous, sexual morph unknown, in highly acidic soil, terrestrial, Czech Republic, see Hujsová et al. (2014; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type species: PRM 922615, CBS 136259.

Acleistia Bayl. Ell. 1917, *Ascomycota* genera *incertae sedis*, one species, type: *A. alniella* Bayl. Ell., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Acolium (Ach.) Gray 1821 (= *Acolium* Trevis. 1862; = *Calicium* subdiv. *Acolium* Ach. 1808; = *Crateridium* Trevis. 1862; = *Cypheliomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Mycacolium* Reinke 1895; = *Pseudacoliumyces* Cif. & Tomas. 1953; = *Pseudacolium* Stizenb. ex Clem. 1909 *fide Species Fungorum* 2017), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, five species, type: *Calicium tympanellum* Ach. (lectotype, designated by) (= *Acolium inquinans* (Sm.) A. Massal.), lichenized, see Gaya et al. (2012; taxonomy), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Acompsomyces Thaxt. 1901, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, six species, type: *A. corticariae* Thaxt., asexual morph unknown, biotrophic on *Coleoptera* (Cryptophagidae and Lathridiidae), America, Europe, probably worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acontium Morgan 1902, *Ascomycota* genera *incertae sedis*, four species, type: *A. album* Morgan, see Seifert et al. (2011; mentioned as a doubtful genus), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Acremoniopsis Giraldo, Gené & Guarro 2014, *Hypocreales*, genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. suttonii* Giraldo, Gené & Guarro, hyphomycetous, sexual morph unknown, from soil, terrestrial, see Crous et al. (2014d; taxonomy), Maharachchikumbura et al. (2015b, 2016b), holotype and ex-type: CBS H-21936, CBS 138708.

Acremoniula G. Arnaud ex Cif. 1962 (= *Acremoniula* G. Arnaud 1954 [1953] nom. inval. *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, five species, type: *A. sarcinellae* (Pat. & Har.) G. Arnaud ex Deighton, hyphomycetous, sexual morph unknown, saprobes,

terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acremonium Link 1809, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, c. 150 species, type: *A. alternatum* Link, hyphomycetous, sexual morph unknown, saprobes, lichenicolous, cosmopolitan, see Weisenborn et al. (2010; new species), Gräfenhan et al. (2011; overview), Kiyuna et al. (2011; new species), Seifert et al. (2011; morphology), Summerbell et al. (2011; overview), Giraldo et al. (2012, 2014c; new species, phylogeny), von Brackel et al. (2012; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Lynch et al. (2016; new species), cultures and sequences are available.

Acroclymma Alcorn & J.A.G. Irwin 1987, *Acroclymmaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *A. medicaginis* Alcorn & J.A.G. Irwin, coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Zhang et al. (2012a; new species, phylogeny), Kirk et al. (2013; genus accepted), Trakunyingcharoen et al. (2014; phylogeny, introduced Acroclymmaceae), Wijayawardene et al. (2014c, 2017; outline), cultures and sequences are available for the type species.

Acroconidiella J.C. Lindq. & Alippi 1964, *Cladosporiaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *A. tropaeoli* (T.E.T. Bond) J.C. Lindq. & Alippi, hyphomycetous, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Acroconidiellina M.B. Ellis 1971 (= *Zeuctomorpha* Sivan. et al. 1984 *fide* Rossman et al. 2016a), *Sympoventuriaceae*, *Venturiales*, *Dothideomycetes*, four species, type: *A. loudeiae* M.B. Ellis, hyphomycetous, sexual morph formerly known in *Zeuctomorpha*, widespread (especially in tropical regions), see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Acrocordia A. Massal. 1854 (= *Acrocordiomycetes* Cif. & Tomas. 1953; = *Amphididymella* Petr. 1928 *fide* Species Fungorum 2017), *Monoblastiaceae*, *Monoblastiales*, *Dothideomycetes*, ten species, type: *A. gemmata* (Ach.) A. Massal., asexual morph unknown, lichenized, terrestrial, temperate, placement in *Monoblastiaceae* supported by molecular data, see Nelsen et al. (2009, 2011b; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013;

genus accepted), cultures available but lacking for the type species.

Acrocordiella O.E. Erikss 1982, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, one species, type: *A. occulta* (Romell) O.E. Erikss, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acrocordiopsis Borse & K.D. Hyde 1989, *Salsugineaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. patillii* Borse & K.D. Hyde, asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Melanommataceae*), Suetrong et al. (2009; phylogeny, morphology), Jones et al. (2009b, 2015; classification), Alias and Jones (2009, monograph), Besitulo et al. (2010; diversity), Pang et al. (2010b; diversity, Malaysia), Zainuddin et al. (2010; antimicrobial activities), Sakayaroj et al. (2011b; diversity, Thailand), Lee et al. (2012; checklist, Malaysia), Zhang et al. (2012e; phylogeny, morphology), Hyde et al. (2013; phylogeny), Samón-Legrá et al. (2014; diversity), Tian et al. (2015; morphology, phylogeny), holotype of type: IMI 297769, available cultures of type: BCC 28167, BCC 28166.

Acrodictyella W.A. Baker & Partr. 2001, *Ascomycota* genera *incertae sedis*, one species, type: *A. obovata* W.A. Baker & Partr., hyphomycetous, sexual morph unknown, wood decaying, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Acrodictyopsis P.M. Kirk 1983, *Ascomycota* genera *incertae sedis*, one species, type: *A. lauri* P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Acrodictys M.B. Ellis 1961, *Sordariomycetidae* genera *incertae sedis*, 22 species, type: *A. bambusicola* M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, aquatic, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Zhao et al. (2011a; descriptions of *Acrodictys* and related genera from China), Ma et al. (2014b; new species), sequences are available.

Acrodontiella U. Braun & Scheuer 1995, *Ascomycota* genera *incertae sedis*, one species, type: *A. fallopiae* U. Braun & Scheuer, hyphomycetous, sexual morph unknown, terrestrial, folicolous, associated with leaf-spots, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of type: GZU 294903, cultures and sequences are unavailable.

Acrodontium de Hoog 1972, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, 13 species, type: *A. crateriforme* (J.F.H. Beyma) de Hoog, hyphomycetous, sexual morph unknown, saprobes, cosmopolitan, see Seifert et al. (2011; morphology), Crous et al. (2014c; new species), Videira et al. (2016; phylogeny), Wijayawardene et al. (2017; outline), ex-type of the type: CBS 144.33 =ATCC 15679.

Acrogenospora M.B. Ellis 1971 (= *Farlowiella* Sacc. 1891 *fide* Rossman et al. 2015b), *Pleosporomycetidae*, genera *incertae sedis*, *Dothideomycetes*, nine species, type: *A. sphaerocephala* (Berk. & Broome) M.B. Ellis, hyphomycetous, sexual morph formerly known in *Farlowiella*, saprobes, cosmopolitan, see Hu et al. (2010b; new species), Seifert et al. (2011; morphology), Rossman et al. (2015b; nomenclature), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Acrogenotheca Cif. & Bat. 1963 (= *Laterotheca* Bat. 1963 *fide* Species Fungorum 2017), *Trichopeltinaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *A. pulcherrima* Bat. & Cif., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Acrogynomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, six species, type specie: *A. eumicri* Thaxt., parasitic on African scydmaenid genus *Eumicrus*, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acrophialophora Edward 1961 (= *Taifanglania* Z.Q. Liang, Y.F. Han, H.L. Chu & R.T.V. Fox 2009 *fide* Zhang et al. 2015b), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, six species, type: *A. nainiana* Edward, sexual morph unknown, pathogen, terrestrial, worldwide, thermo-tolerant, see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2016b; outline), Kirk et al. (2013; genus accepted), Sandoval-Denis et al. (2015a; clinical aspects), Zhang et al. (2015b; DNA), ex-isotype of type: CBS 100.60, ex-type culture of type: ATCC 22555, IMI 076567.

Acrophragmis Kiffer & Reisinger 1970, *Ascomycota* genera *incertae sedis*, three species, type: *A. coronata* Kiffer & Reisinger, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acrosyphus Lév. 1846, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *A. sphaerophoroides* Lév., lichenized, terrestrial (saxicolous), Americas, Asia, Africa, see Gaya et al. (2012; phylogeny), Prieto et al.

(2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Acrospeira Berk. & Broome 1857 (= *Spirospora* L. Mangin & Vincens 1920 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *A. mirabilis* Berk. & Broome, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acrospermoïdes J.H. Mill. & G.E. Thomps. 1940, *Sordariomycetes* genera *incertae sedis*, two species, type: *A. subulata* J.H. Mill. & G.E. Thomps., asexual morph unknown, saprobes, terrestrial, Greece, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Acrospermum Tode 1790 (= *Scleroglossum* Fr. 1874; = *Xyloglossum* Pers. 1818 *fide* Species Fungorum 2017), *Acrospermaceae*, *Acrospermales*, *Dothideomycetes*, eleven species, type: *A. compressum* Tode, asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; phylogeny), Hyde et al. (2013; morphology, phylogeny), Kirk et al. (2013 genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available for type species.

Acrostalagmus Corda 1838, *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*, three species, type: *A. cinnabarinus* Corda, hyphomycetous, sexual morph unknown, widespread, see Wijayawardene et al. (2012, 2017; outline), Réblová et al. (2011; DNA, phylogeny), cultures and sequences are available.

Acrostaurus Deighton & Piroz. 1972, *Ascomycota* genera *incertae sedis*, one species, type: *A. turneri* Deighton & Piroz., asexual morph unknown, fungicolous, saprobes, widespread especially in tropical regions, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Actidiographium Lar. N. Vassiljeva 2000, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, one species, type: *A. orientale* Lar & Vassiljeva, asexual morphs unknown, saprobes, terrestrial, north temperate, see Boehm et al. (2009a; accepted as in *Hysteriaceae*), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mytilinidiaceae*), Wijayawardene et al. (2014c; outline), holotype: VLA P-2938, cultures and sequences are unavailable.

Actidium Fr. 1815 (= *Bulliardella* (Sacc.) Paoli 1905; = *Mytilinidion* subgen. *Bulliardella* Sacc. 1883; = *Ostreionella* Seaver 1926 *fide* Species Fungorum 2017),

Mytilinidiaceae, *Mytilinidiales*, *Dothideomycetes*, nine species, type: *A. hysteroides* Fr., asexual morphs unknown, saprobes, terrestrial, Europe, North America, see Boehm et al. (2009a, b; accepted as a genus in *Mytilinidiaceae*), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as in *Mytilinidiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Actinocladium Ehrenberg 1819, *Ascomycota* genera *incertae sedis*, eight species, type: *A. rhodosporum* Ehrenb., hyphomycetous, sexual morph unknown, pathogen of wood and bark of fallen twigs and trunk, worldwide, mainly Europe, Asia, Africa, Caribbean, see Manoharachary et al. (2010; two new species), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Actinocymbé Höhn. 1911 (= *Limaciniella* J.M. Mend. 1925 *fide* Species Fungorum 2017), *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, three species, type: *A. separato-setosa* (Henn.) Höhn., asexual morph unknown, parasitic, epiphyllous to hypophyllous, terrestrial, temperate to tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Actinomyxa Syd. & P. Syd. 1917, *Dothideomycetes* genera *incertae sedis*, one species, type: *A. australiensis* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013) did not accept as a genus in *Microthyriaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Actinoplaeca Müll. Arg. 1891 (= *Actinoplacomycetes* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *A. strigulacea* Müll. Arg., lichenized, terrestrial (folicolous), tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Actinotexis Arx 1960, *Ascomycota* genera *incertae sedis*, one species, type: *A. mirabilis* Berk. & Broome, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Actinothecium Flot. 1855 (see Species Fungorum 2017 for synonyms), *Ascomycota* genera *incertae sedis*, five species, type: *A. caricicola* Ces., coelomycetous, sexual

morph unknown, widespread, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Actinothyrium Kunze 1823, *Ascomycota* genera *incertae sedis*, ten species, type: *A. graminis* Kunze ex Fr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Acumispora Matsush. 1980, *Ascomycota* genera *incertae sedis*, four species, type: *A. uniseptata* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Adautomilanezia Gusmão, S.S. Silva, Fiúza, L.A. Costa, T.A.B. Santos 2016, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *A. caesalpiniae* Gusmão, S.S. Silva, Fiúza, L.A. Costa, T.A.B. Santos, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Crous et al. (2016a; taxonomy, phylogeny), holotype and ex-type: HUEFS216632, CCLAMIC 102/12.

Adelococcus Theiss. & Syd. 1918, *Adelococcaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *A. alpestris* (Zopf) Theiss. & Syd., asexual morph unknown, lichenicolous, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Adelodiscus Syd. 1931, *Leotiomycetes* genera *incertae sedis*, three species, type: *A. philippinensis* Syd., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, need generic revision.

Adelolecia Hertel & Hafellner in Hafellner 1984 (= *Leclideomyces* E.A. Thomas ex Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, four species, type: *A. pilati* (Hepp) Hertel & Hafellner, all with ascospores, lichenized (green algae), terrestrial (saxicolous, rarely corticolous), temperate regions of both Hemispheres, Ekman et al. (2008; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Adelosphaeria Réblová 2016 [2017], *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, one species, type: *A. catenata* Réblová, asexual morph hyphomycetous, saprobes, terrestrial, Czech Republic, see

Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2017; taxonomy), holotype and ex-type: PRM 933853, CBS 138679.

Aderkomycetes Bat. 1961, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, c. 30 species, type: *A. couepiae* Bat, with ascomata, lichenized (green algae), terrestrial (foliicolous or corticolous), tropical, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type.

Adomia S. Schatz 1985, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. avicenniae* S. Schatz, saprobes, on mangrove wood, saprobe, Australia, Egypt, Red Sea, Thailand, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; morphology, outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Aequabiliella Crous 2015, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, coelomycetous, sexual morph unknown, one species, type: *A. effusa* (Damm & Crous) Crous in necrotic wood of *Prunus persica*, pathogens, terrestrial, South Africa, see Damm et al. (2010b as *Phaeomoniella effusa* Damm & Crous, morphology), Crous et al. (2015d; phylogeny), holotype and ex-type culture of type: CBS H-19998, CBS 120883.

Aeruginoscyphus Dougoud 2012, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *A. sericeus* (Alb. & Schwein.) Dougoud, asexual morph unknown, saprobes, terrestrial, Europe, see Dougoud (2012; taxonomy), cultures and sequences are unavailable.

Agaricodochium X.J. Liu 1981, *Ascomycota* genera *incertae sedis*, one species, type: *A. camelliae* X.J. Liu, A.J. Wei & S.G. Fan, hyphomycetous, on leaves of *Camellia oleosa*, pathogen of *Camellia oleiferra*, terrestrial, Zhejiang, China, see Guang-yu et al. (2008; control), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Fangyan et al. (2013; control), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Agarwalyomyces R.K. Verma & Kamal 1987, *Ascomycota* genera *incertae sedis*, one species, type: *A. indicus* R.K. Verma & Kamal, hyphomycetous, on fruit of *Lythraceae*, terrestrial, India, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Aggregatorygma M. Cáceres, Aptroot & Lücking 2014, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *A. triseptatum* M. Cáceres, Aptroot & Lücking, lichenized, see Cáceres et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Agonimia Zahlbr. 1909 (= *Agonimiella* H. Harada 1993 *fide Species Fungorum* 2017), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, c. 17 species, type: *A. tristicula* (Nyl.) Zahlbr., asexual morph unknown, lichenized, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Muggia et al. (2010a; DNA), Dymytrova et al. (2011; new species), Guzow-Krzemińska et al. (2012; new species), Aptroot and Cáceres (2013; new species), Harada (2013; new species, Japan), Kirk et al. (2013; genus accepted), Kondratyuk (2015; new species), Lücking and Moncada (2017; taxonomy and phylogeny), cultures available for the type: L469 (*fide* Muggia et al. 2010a).

Agrabeja Subram. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *A. kavakapriya* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Singapore, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype MUBL 3140.

Agyriella Sacc. 1884, *Ascomycota* genera *incertae sedis*, two species, type: *A. nitida* (Lib.) Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Agyriellopsis Höhn. 1903, *Ascomycota* genera *incertae sedis*, three species, type: *A. caeruleoatra* Höhn., coelomycetous, sexual morph unknown, Europe see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Agyrium Fr. ex Fr. 1822 (= *Agyrium* Fr. 1821 nom. inval.; = *Exogone* Henn. 1908; = *Myxomphalos* Wallr. 1833 *fide Species Fungorum* 2017), *Pertusariaceae*, *Pertusariales*, *Lecanoromycetes*, three species, type: *A. rufum* (Pers.) Fr., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; taxonomy), Kirk et al. (2013; genus accepted), Resl et al. (2015; DNA), cultures and sequences are available.

Ahmadia Syd. 1939, *Ascomycota* genera *incertae sedis*, one species, type: *A. pentatropidis* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Pakistan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ahtiaria Goward 1986, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *A. sphaerospora* (Müll. Arg.) Goward, all with ascomata, lichenized (green algae), terrestrial (corticulous, epixylic), western North America, see Thell et al. (2009; phylogeny), Nelsen et al.

(2011a; phylogeny), Miadlikowska et al. (2014a; phylogeny), Divakar et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Aigialus Kohlm. & S. Schatz 1986, *Aigialaceae*, *Pleosporales*, *Dothideomycetes*, asexual morph unknown, five species, type: *A. grandis* Kohlm. & Schatz, saprobes, marine, worldwide, see Alias and Jones (2009; monograph), Jones et al. (2009b, 2012, 2015; classification, outline, review), Schoch et al. (2009; phylogeny), Suetrong et al. (2009, 2011a, b, 2014; phylogeny, morphology), Zhang et al. (2009a, 2012e; phylogeny), Isaka et al. (2009; secondary metabolites), Nambiar (2009; taxonomy), Nambiar and Raveendran (2009; diversity), Nambiar et al. (2009; diversity), Zainuddin et al. (2010; antimicrobial activities), Besitudo et al. (2010; diversity), Lumbsch and Huhndorf (2010; outline), Pang et al. (2010b, 2011b, 2013a; diversity, phylogeny), Sakayaroj et al. (2011b; diversity), Josphine et al. (2011; diversity), Liu et al. (2011b, 2015a; phylogeny, morphology), Boonmee et al. (2012; phylogeny), Dai et al. (2012; phylogeny), Lee et al. (2012; checklist), Pang and Jheng (2012a; checklist), Hyde et al. (2013; phylogeny, morphology, outline), Ariyawansa et al. (2014a, c; phylogeny), Kocharin et al. (2013; optimization), Sivakumar (2013; review), Sarma and Raghukumar (2013; diversity), Gnavi et al. (2014, phylogeny), Phookamsak et al. (2014c, 2015a; phylogeny), Wijayawardene et al. (2014c; outline), Shearer et al. (2014; phylogeny), Pratibha and Prabhugaonkar (2015; phylogeny), Shen et al. (2015; secondary metabolites), Doilom et al. (2017; phylogeny), holotype: J.K. 4332a (IMS), ex-isotype strain J.K.4332b (NY).

Ainoa Lumbsch & I. Schmitt 2001, *Baeomycetaceae*, *Baeomycetales*, *Lecanoromycetes*, three species, type: *A. geochroa* (Körb.) Lumbsch & Schmitt, ascomata apotheciate, conidiomata pycnidial, lichen-forming, saxicolous or terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Resl et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Aithaloderma Syd. & P. Syd. 1913 (= *Asbolisia* Bat. & Cif. 1963; = *Blastocapnia* Cif. & Bat. 1963; *Chaetopotius* Bat. 1951; = *Hypocapnodium* Speg. 1918; = *Phaeochaetia* Bat. & Cif. 1962; = *Vertixore* V.A.M. Mill. & Bonar 1941 *fide* Species Fungorum 2017), *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, 15 species, type: *A. clavatisporum* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2011), Hyde et al. (2013) and Wijayawardene et al. (2014c) did not accept as a genus

in *Capnodiaceae*, Kirk et al. (2013; outline, genus accepted), cultures and sequences are unavailable.

Aivenia Svrček 1977, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *A. tantula* Svrček, asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ajrekarella Kamat & Kalani 1964, *Ascomycota* genera *incertae sedis*, one species, type: *A. polychaetiae* Kamat & Kalani, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Akanthomyces Lebert 1858, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, c. 13 species, type: *A. aculeatus* Lebert, hyphomycetous, sexual morph unknown, saprobes, on insects and spiders, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), cultures and sequences are available.

Alanphillipsia Crous & M.J. Wingf. 2013, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, five species, type: *A. aloes* Crous & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes or possibly pathogenic, associated with leaf lesions, South Africa, see Crous et al. (2013b, 2014d; morphology), Wijayawardene et al. (2016b, 2017; outline, morphology, phylogeny), Dissanayake et al. (2016b; phylogeny), holotype and ex-type: CBS H-21418, CPC 21298.

Alascospora Raja, Violi & Shearer 2010, *Dothideomycetes*, genera *incertae sedis*, one species, type: *A. evergladensis* Raja, Violi & Shearer, asexual morph unknown, saprobes, aquatic, USA, see Raja et al. (2010c; morphology), Wijayawardene et al. (2014c; outline), holotype of the type: ILL 40789, cultures and sequences are unavailable.

Alatosessilispora K. Ando & Tubaki 1984, *Ascomycota* genera *incertae sedis*, one species, type: *A. bibrachiata* K. Ando & Tubaki, hyphomycetous, sexual morph unknown, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available for the type, need generic revision.

Alatospora Ingold 1942, *Leotiaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *A. acuminata* Ingold, hyphomycetous, sexual morph unknown, saprobes, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Baschien et al. (2013; DNA), Kirk et al. (2013; genus accepted),

Duarte et al. (2015; DNA), neotype of type: IMI 222997, available cultures of the type: CCM FI2186.

Albertiniella Kirschst. 1936, *Cephalothecaceae*, *Sordariomycetes*, families *incertae sedis*, one species, type: *A. reticulata* Kirschst., asexual morph acremonium-like, saprobes, terrestrial, Europe, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures available for the type: NBRC 9424.

Albifimbria L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *A. verrucaria* (Alb. & Schwein.) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, from soil, terrestrial, cosmopolitan, see Lombard et al. (2016; taxonomy, phylogeny), cultures and sequences are available.

Albonectria Rossman & Samuels 1999, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, type: *A. rigidiuscula* (Berk. & Broome) Rossman & Samuels, one species, asexual morph hyphomycetous, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Zhao et al. (2011b; phylogeny), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Alborbis Senan. & K.D. Hyde 2017, *Sydiomellaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. galericulata* (Tul. & C. Tul.) Senan. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Europe, see Senanayake et al. (2017; taxonomy, phylogeny), holotype of the type: NYBG 02929068, cultures and sequences are available.

Albosynnema E.F. Morris 1967, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *A. elegans* E.F. Morris 1967, asexual morph hyphomycetous, America, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2016; phylogeny), cultures and sequence are available.

Albotricha Raity. 1970, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, c. 19 species, type: *A. acutipila* (P. Karst.) Raity., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Alciphila Harmaja 2002, *Ascomycota* genera *incertae sedis*, one species, type: *A. vulgaris* Harmaja, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Aldona Racib. 1900, *Parmulariaceae*, *Dothideomycetes*, families *incertae sedis*, three species, type: *A. stella-nigra*

Racib., asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), Tian et al. (2016a; taxonomy), cultures and sequences are unavailable.

Aldonata Sivan. & A.R.P. Sinha 1989, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *A. pterocarpi* Sivan. & A.R.P. Sinha, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), Tian et al. (2016a; taxonomy), cultures and sequences are unavailable.

Alectoria Ach. 1809 (= *Alectoria* sect. *Eualectoria* Th. Fr. 1880; = *Alectoriomyces* Cif. & Tomas. 1953; = *Bryopogon* Link 1833; = *Ceratocladia* Schwend. 1860; = *Eualectoria* (Th. Fr.) Gyeln. fide Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, nine species, type: *A. sarmentosa* (Ach.) Ach., with ascomata, but often absent, lichenized (green algae), terrestrial (corticulous, terricolous), widespread, see Halonen et al. (2009; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Divakar et al. (2015; phylogeny), Wang et al. (2015f; new species), Lücking et al. (2017a; classification), McMullin et al. (2016; phylogeny), sequences are available.

Aleuria Fuckel 1870 (= *Peziza* Fuckel 1870; = *Peziza* subgen. *Aleuria* (Fuckel) Quél. 1886 fide Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 17 species, type: *A. aurantia* (Pers.) Fuckel, asexual morph unknown, on soil, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for the type species.

Aleurina Massee 1898 (= *Jafneadelphus* Rifai 1968 fide Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, eleven species, type: *A. tasmanica* Massee, asexual morph unknown, on soil, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Alfaria Crous, Montano-Mata & Garcia-Jim. 2014, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, eight species, type: *A. cyperi-esculentii* Crous, N.J. Montaño-Mata & García-Jim., asexual morph coelomycetous, pathogenic, terrestrial, Spain, Italy, see Crous et al. (2014c; morphology of sexual morph), Lombard et al. (2016; phylogeny, new species), Wijayawardene et al. (2016b, 2017; morphology of asexual morph, phylogeny), ex-type strains of type: CPC 23153; holotype CBS H-21703.

Alfariacladiella Crous & R.K. Schumach. 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *A. spartii* Crous & R.K. Schumach., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Spain, see Hernández-Restrepo et al. (2016a; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and culture ex-type of the type: CBS H-22655, CPC 24966.

Algincola Velen. 1939, *Helotiales* genera *incertae sedis*, three species, type: *A. quercina* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Algonquinia R.F. Castañeda & W.B. Kendr. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *A. inflata* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Canada, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), holotype of the type: INIFAT C89/401, cultures and sequences are unavailable, compare with *Catenocuneiphora*, *Hormomyces* and *Sphaerosporium*.

Alina Racib. 1909, *Perisporiopsidaceae*, *Dothideomycetes*, families *incertae sedis*, one species, type: *A. jasmini* Racib., asexual morph hyphomycetous, parasitic on living leaves of *Jasminum* spp., Buitenzorg, Indonesia, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, re-collection and revision required.

Aliquandostipite Inderb. 2001, *Aliquandostipitaceae*, *Jahnulales*, *Dothideomycetes*, seven species, type: *A. khaoyaiensis* Inderb., asexual morph unknown, saprobes, freshwater, Australia, China, Costa Rica, Egypt, Panama, Thailand, Venezuela, USA, see Schoch et al. (2009; phylogeny), Shearer et al. (2009; phylogeny), Suetrong et al. (2010, 2011a; outline, phylogeny), Raja et al. (2011b; phylogeny), Liu et al. (2015a; morphology), holotype and ex-type of the type: UBC F13875, CBS 118232.

Alisea Dupont & E. B. G. Jones 2009, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *A. longicolla* Dupont & Jones, asexual morph unknown, saprobes, marine, ascomata on deep sea sunken wood, south Pacific Ocean and South China Sea, see Dupont et al. (2009; taxonomy), Jones et al. (2015; outline), Maharachchikumbura et al. (2016b; outline), holotype of type: PC 0115840.

Allantonectria Earle 1901 (see Species Fungorum 2017 for synonyms), *Nectriaceae*, *Hypocreales*,

Sordariomycetes, type: *A. yuccae* Earle, four species, asexual morph hyphomycetous, terrestrial, saprobes, cosmopolitan, see Hirooka et al. (2012; monograph), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Allantoparmelia (Vain.) Essl. 1978 (= *Parmelia* subgen. *Allantoparmelia* Vain. 1909 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *A. alpicola* (Th. Fr.) Essl., all with ascomata (but often absent), lichenized (green algae), terrestrial (saxicolous, rarely terricolous), boreal or arctic alpine in Northern Hemisphere, see Crespo et al. (2010b; phylogeny), Lumbsch and Huhndorf (2010; outline), Spribille et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; DNA, phylogeny), Divakar et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Allantophomopsiella Crous 2014, *Phacidiaceae*, *Phacidiales*, *Leotiomycetes*, one species, type: *A. pseudotsugae* (M. Wilson) Crous, saprobes, Europe, see Crous et al. (2014b; morphology, phylogeny), reference specimen and culture of type: CBS H-15946, culture CBS 562.63.

Allantophomopsis Petr. 1925 (= *Apostrasseria* Nag Raj 1983 *fide* Jaklitsch et al. 2016a), *Phacidiaceae*, *Phacidiales*, *Leotiomycetes*, eight species, type: *A. cytisporea* (Fr.) Petr., asexual morph unknown, saprobes, terrestrial, worldwide, see Crous et al. (2014b, 2015a; phylogeny), Wijayawardene et al. (2017; outline), ex-epitype of the type: CPC 24977.

Allantoporthe Petr. 1921 (= *Allantoporthe* Petr. 1921; = *Aporhytisma* Höhn. 1917; = *Chorostate* (Nitschke ex Sacc.) Traverso 1906; = *Clypeorhynchus* Kirschst. 1936; = *Dialytes* Nitschke 1867; = *Diaporthe* subgen. *Chorostate* Nitschke ex Sacc. 1882; = *Melanoporthe* Wehm. 1938; = *Septomazzantia* Theiss. & Syd. 1915 *fide* Species (Fungorum 2017), *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. leucothoës* Lar.N. Vassiljeva, asexual morph unknown, saprobes, terrestrial, North America, see Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Alloanthostomella Daranag., Camporesi & K.D. Hyde 2016, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. rubicola* (Speg. ex Sacc. & Trotter) Daranag., Camporesi & K.D. Hyde, asexual morph unknown, terrestrial, saprobes, see Daranagama et al. (2016a; taxonomy), Wendt et al. (2017; placed in *Xylariales*), cultures and sequences are available.

Alloarthopyrenia Phukhams., Lücking & K.D. Hyde 2016, *Tryptethiaceae*, *Tryptethiales*, *Dothideomycetes*,

one species, type: *A. italicica* Phukhams., Camporesi, Ariyaw. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Hyde et al. (2016; taxonomy, phylogeny), holotype of the type: MFLU 15-0399.

Alloascoidea Kurtzman & Robnett 2013, *Alloascoideaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *A. africana* (L.R. Batra & Francke-Grosm.) C.P. Kurtzman & C.J. Robnett, asexual morph unknown (budding of cells and formation of true hyphae), saprophytes, from trees and associated insects, terrestrial, Europe, Africa, see Kurtzman and Robnett (2013a; taxonomy), cultures and sequences are available.

Allocallicium M. Prieto & Wedin 2016, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *A. adaequatum* (Nyl.) M. Prieto and Wedin., lichenized, see Prieto and Wedin (2016; taxonomy, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Allocetraria Kurok. & M.Y. Lai 1991 (= *Usnocetraria* M.J. Lai & J.C. Wei 2007 *fide Species Fungorum* 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, nine species, type: *A. stracheyi* (C. Bab.) Kurok. & M.J. Lai; lichenized; see Thell et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2011a; DNA, phylogeny), Miadlikowska et al. (2014a; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Alloconiothyrium Verkley, Göker & Stielow 2014, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. aptrootii* Verkley & Stielow, coelomycetous, sexual morph unknown, from soil, Papua New Guinea, see Verkley et al. (2014; morphology, phylogeny); Wijayawardene et al. (2016b; morphology, phylogeny), holotype and ex-type culture of type species CBS H-21035, CBS 980.95.

Alloleptosphaeria Ariyaw., Wanaw. & K.D. Hyde 2015, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. italicica* Wanaw., Camporesi, Ariyaw. & K.D. Hyde, asexual morph unknown, terrestrial, Europe, see Ariyawansa et al. (2015b; taxonomy, phylogeny), holotype of the type: MFLU 15-1517, cultures and sequences are available.

Allonecte (Pat.) Syd. 1939, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. lagerheimii* (Pat.) Syd. (= *Broomella lagerheimii* Pat. 1895), asexual morph coelomycetous, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Boonmee et al. (2011; transferred to *Pleosporaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Allonectella Petr. 1950, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *A. rubescens* Petr.,

sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b), cultures and sequences are available.

Alloneottiosporina Nag Raj 1993, *Ascomycota* genera *incertae sedis*, two species, type: *A. carolinensis* Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: BPI 1110181.

Allophaeosphaeria Ariyaw., Camporesi & K.D. Hyde 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *A. muriformia* Ariyawansa, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Europe, see Ariyawansa et al. (2015a; taxonomy, phylogeny), Liu et al. (2015a; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type cultures of the type: MFLU 15-0067, MFLUCC 13-0349.

Allophoma Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *A. tropica* (R. Schneid. & Boerema) Q. Chen & L. Cai, coelomycetous, saprobes, pathogens, terrestrial, cosmopolitan, see Chen et al. (2015b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), isotype and ex-isotype of the type: CBS H-7629, CBS 436.75.

Allophoron Nádv. 1942, *Pezizomycotina* genera *incertae sedis*, one species, type: *A. farinosum* Nádv., asexual morph unknown, lichenized, terrestrial, Colombia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017b; classification), cultures and sequences are unavailable.

Allophylaria (P. Karst.) P. Karst. 1870 (= *Helotidium* Sacc. 1884; = *Peziza* sect. *Allophylaria* P. Karst. 1869 *fide Species Fungorum* 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *X. sticticum* (Fr.) Wallr., asexual morph unknown, saprobes, terrestrial, China, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Allosoma Syd. 1926, *Englerulaceae*, *Dothideomycetes* families *incertae sedis*, four species, type: *A. cestri* Syd., asexual morph hyphomycetous, epiphytic on leaves or saprobes on dead stems, North and South America, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014e), cultures and sequences are unavailable, holotype of type: GZU157-80.

Allothyrina Bat. & J.L. Bezerra 1964, *Ascomycota* genera *incertae sedis*, one species, type: *A. serjaniae* Bat. & J.L. Bezerra, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Allothyriopsis Bat., Cif. & H. Maia 1959, *Ascomycota* genera *incertae sedis*, one species, type: *A. landolphiae* Bat., Cif. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, Ghana, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Alnecium Voglmayr & Jaklitsch 2014, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *A. auctum* (Berk. & Broome) Voglmayr & Jaklitsch, asexual morph unknown, saprobes, terrestrial, Europe, see Voglmayr and Jaklitsch (2014; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Alpakesa Subram. & K. Ramakr. 1954, *Ascomycota* genera *incertae sedis*, four species, type: *A. yuccifolia* (J.G. Hall) Subram. & K. Ramakr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, Thiwan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), available specimens of type: IMI 177975, cultures are unavailable.

Alpakesiopsis Abbas, B. Sutton, Ghaffar & A. Abbas 2003, *Ascomycota* genera *incertae sedis*, one species, type: *A. ghaffarii* Abbas, B. Sutton & Al. Abbas, coelomycetous, sexual morph unknown, saprobes, terrestrial, Pakistan, see Wijayawardene et al. (2012, 2017; outline), holotype of the type: IMI 202284, cultures are unavailable.

Alpestrisphaeria Thambug. & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. terricola* (G.S. Gong) Thambug. & K.D. Hyde, asexual morph unknown, associated with soil, China, see Thambugala et al. (2015b; phylogeny), Ariyawansa et al. (2016c; new species), ex-type culture of type: SC-12.

Alternaria Nees 1817 (= *Allewia* E.G. Simmons 1990 *fide* Woudenberg et al. 2013; Wijayawardene et al. 2014c; Rossman et al. 2015b; see Woudenberg et al. 2013 for all synonyms), *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, 663 species, type: *A. alternata* (Fr.) Keissl., sexual morph formerly known in *Lewia*, saprobes, plant and animal pathogens, terrestrial, worldwide, see Aly et al. (2008; cytotoxic metabolites), Andersen et al. (2008; metabolite profiling), Avenot et al. (2008; succinate dehydrogenase mutation), Azcarate et al. (2008; toxins), Bartra et al. (2008; respiratory allergy), Cota et al. (2008; Altenuasin), Craven et al. (2008; Anastomosis), Flors et al. (2008; cell signalling), Grinn-Gofroń and Strzelczak (2008; meteorological factors), Harimoto et al. (2008; AMT2 gene), Kumar et al. (2008a, b; variability, India, host resistance), Lawrence et al. (2008; pathogenicity review), Miyamoto et al. (2008, 2009; ACT-toxin biosynthesis,

phylogeny), Ostry (2008; *Alternaria* mycotoxins), Park et al. (2008; phylogeny), Parada et al. (2008; toxin from germinating spores), Pasche and Gudmestad (2008; F129L mutation), Pastor and Guarro (2008; infections), Perello et al. (2008; Argentina, black point of wheat), Rosenzweig et al. (2008; sensitivity to azoxystrobin), Sabirov et al. (2008; nasal polyposis), Saravanakumar et al. (2008; competition for host), Schmechel et al. (2008; environmental immunoassays), Tabar et al. (2008; immunotherapy), Véliz et al. (2008; Mannitol biosynthesis), Wang et al. (2008c; biocontrol), Yuan et al. (2008; Tricycloalternareses), Logrieco et al. (2009; pathogenicity review), Latha et al. (2009; host resistance induction), Lee et al. (2009b; protein disruption study), Pogány et al. (2009; biochemistry), Kjer et al. (2009; phenolic extrolites), Mishra et al. (2009; inhibition by Cinnamomum), Kobayashi et al. (2009; hypersensitivity), Hadizadeh et al. (2009; antifungal activity), Andrew et al. (2009; phylogeny), Lin et al. (2009; pathological mechanism), Tu (2009; keratitis), Avenot et al. (2009; protein mutation), Ali et al. (2009; malachite green degradation), Pedras et al. (2009; review), Siegel et al. (2009; HPLC, MS), Grinn-Gofroń and Rapiejko (2009; Poland), Aoki et al. (2009; mycovirus), Wight et al. (2009; biochemistry), Fawzi et al. (2009; antifungal effects), Wang et al. (2009c, d; cerebrosides, postharvest biocontrol), Gu (2009, Bioactive metabolites), Andersen et al. (2009; polyphasic taxonomy), Phuwapraisirisan et al. (2009; antitumour metabolites), Yang et al. (2009; AP1 pathway study), Cho et al. (2009; virulence factors), Fernandes et al. (2009; fermentation extract activity), Stępalska and Wołek (2009; Intradiurnal periodicity), Hedayati et al. (2009; IgE analysis), Nallathambi et al. (2009; *Trichoderma* effects), Gao et al. (2009; Perylene derivatives), Anand et al. (2009; host resistance), Pose et al. (2009; growth factors), Kim et al. (2009b; Alternaramide), Evidente et al. (2009; Alternethanoxins), Hipolito et al. (2009; brain abscess), Kasahara et al. (2010; Solanapyrone synthase), Mukherjee et al. (2010; infection proteomics), Wolf et al. (2010; CO₂ effects), Imazaki et al. (2010; peroxisomes), Lin and Chung (2010; signalling pathways), Ntui et al. (2010; wasabi defensin gene), Horsfield et al. (2010; fungicides), Randriamanantany et al. (2010; allergic rhinitis), Siegel et al. (2010; tenuazonic acid), Boyce et al. (2010; Phaeohyphomycosis), Kilic et al. (2010; effect on children), Lin et al. (2010, 2011; FUS3 MAPK pathway, pathogenicity), Dellavalle et al. (2011; antifungal activity), Kuna et al. (2011; immunotherapy), Boitano et al. (2011; molecular pathogenesis), Dey et al. (2011; lipid profiling), Song et al. (2011; Abscisic acid resistance), Huang et al. (2011; bianthraquinone derivatives), Escuredo et al. (2011; meteorological factors), Joubert et al. (2011a, b; stress

adaptation, pathogenicity), Sinha and Chattopadhyay (2011; host proteome profile), Yago et al. (2011; cell signalling), Noser et al. (2011; UPLC-MS/MS), Karaoglanidis et al. (2011; QoI fungicides resistance), Zheng et al. (2012; bioactive metabolites), Doherty et al. (2012; immunology), Fleck et al. (2012; toxins in mammals), Nowicki et al. (2012; crucifers host), Chruszcz et al. (2012; allergen), Pane et al. (2012; whey compost-tea control), Matsuwaki et al. (2012; immunological pathway), Recio et al. (2012; spore fluctuation), Skjøth et al. (2012; Copenhagen), Yang and Chung (2012; stress resistance), Hu et al. (2012d; Genomic characterization), Cho et al. (2012; Amr1), Fuiano et al. (2012; nasal IgE measurement), Asam et al. (2012; mycotoxin in food), Sarkar et al. (2012; Au nanoparticles), Twaroch et al. (2012; vaccine candidates), Chhikara et al. (2012; host resistance), Saha et al. (2012; biochemistry), Woudenberg et al. (2013, 2014; review), Tsuge et al. (2013; toxins), Lawrence et al. (2013; phylogeny), Lou et al. (2013; metabolites review), Jia et al. (2013; host pathogen interaction), Prelle et al. (2013; toxin detection in food), Müller and Korn (2013; mycotoxins in wheat), Chakraborty et al. (2013; congo red dye degradation), Mamgain et al. (2013; pathogenicity), Snelgrove et al. (2014; immunology), Wijayawardene et al. (2014c; nomenclature), Ouda (2014; nanoparticle effect), Ariyawansa et al. (2015a, c; new species, phylogeny), Rossman et al. (2015b; nomenclature), Templeton (2016, pathogenesis), Meena et al. (2016; mustard disease), Thambugala et al. (2017b, new species, phylogeny), Zhang et al. (2017a; LAMP), molecular data available for most of the species, including type species; GenBank Number NR_131316, epitype and ex-epitype of type: IMI 254138; CBS 916.96.

Alternariaster E.G. Simmons 2007, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. helianthi* (Hansf.) E.G. Simmons, hyphomycetous, sexual morph unknown, pathogens, Brazil, see Alves et al. (2013; new species, morphology, phylogeny), Hyde et al. (2013; accepted as a genus in *Pleosporaceae*), Wijayawardene et al. (2014c; outline), Woudenberg et al. (2014; phylogeny), cultures and sequences are available.

Alveariospora Meir. Silva, R.F. Castañeda, O.L. Pereira & R.W. Barreto 2012, *Ascomycota* genera *incertae sedis*, one species, type: *A. distoseptata* Meir. Silva, R.F. Castañeda, O.L. Pereira & R.W. Barreto, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Silva et al. (2012; taxonomy), Wijayawardene et al. (2017; outline), holotype of the type: VIC 31399, cultures and sequences are unavailable, compare with *Annellophragmia*, *Annellosympodia* and *Spiropes*.

Alveophoma Alcalde 1952, *Ascomycota* genera *incertae sedis*, one species, type: *A. caballeroi* Alcalde,

coelomycetous, sexual morph unknown, saprobes, Spain, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need further revisions.

Alysidiella Crous 2006, *Dothideomycetes* genera *incertae sedis*, four species, type: *A. parasitica* Crous, hyphomycetous, sexual morph unknown, pathogens, Cyprus, South Africa, Uruguay, see Seifert et al. (2011; morphology), Cheewangkoon et al. (2012; morphology, phylogeny), Wijayawardene et al. (2012, 2017; outline), holotype and ex-type culture of type: CBS H-19742, CPC 12835.

Alysidiopsis B. Sutton 1973, *Ascomycota* genera *incertae sedis*, five species, type: *A. pipsissevae* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need further revisions.

Alyxoria Ach. 1821, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, twelve species (many additional species in the tropics; revision needed), type: *A. varia* (Pers.) Ertz & Tehler, lichenized, see Ertz and Tehler (2011; taxonomy, phylogeny), Diederich et al. (2012a; taxonomy, checklist), Lücking et al. (2017a; classification), sequences are available.

Amallospora Penz. 1897, *Ascomycota* genera *incertae sedis*, one species, type: *A. dacrydion* Penz., hyphomycetous, sexual morph unknown, saprobes, Java, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need further revisions.

Amandinea M. Choisy ex Scheid. & M. Mayrhofer 1993 (= *Amandinea* M. Choisy, Bull. 1950 *fide* Species Fungorum 2017), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 35 species, type: *A. coniops* (Wahlenb.) M. Choisy ex Scheid. & H. Mayrhofer, lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny), Kirk et al. (2013; outline, genus accepted), Kondratyuk et al. (2015d; notes), Prieto et al. (2013; phylogeny), Prieto and Wedin (2016; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Amarenographium O.E. Erikss. 1982 (= *Amarenomyces* O.E. Erikss. 1981 *fide* Rossman et al. 2016a), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *A. metableticum* (Trail) Erikss., asexual morph coelomycetous, sexual morph unknown, endophytic and saprobes, China, Saudi Arabia, Sweden, Italy, see Hodhod et al. (2012; morphology, phylogeny); Wijayawardene et al. (2016b; morphology, phylogeny),

Rossman et al. (2016a; nomenclature), cultures and sequences are available but lacking for the type.

Amaurascopsis Guarro, Gené & De Vroey 1992, *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, asexual morph unknown, two species, type: *A. perforata* Guarro, Gené & De Vroey, saprobes, from chicken yard soil, Burundi, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Amauroascus J. Schröt. 1893, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, 14 species, type: *A. verrucosus* (Eidam) J. Schröt., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Sigler et al. (2013; DNA), cultures and sequences are available but lacking for the type.

Amazonia Theiss. 1913 (= *Actinodothis* Syd. & P. Syd. 1914; = *Amazoniella* Bat. & H. Maia 1960; = *Meliolaster* Doidge 1920 *fide* Species Fungorum 2017), *Meliolaceae*, *Meliolales*, *Sordariomycetes*, more than 70 species, type specie: *A. psychotriae* (Henn.) Theiss., asexual morph unknown, pathogens causing black mildew disease, terrestrial, tropical and subtropical distribution, see Hosagoudar and Archana (2009; host distribution in India), Lumbsch and Huhndorf (2010; outline), Hosagoudar et al. (2010c; new species, morphology), Hosagoudar and Robin (2011; new species, morphology), Hosagoudar (2013; new species, morphology), LSU sequence are available for *A. acacia*.

Amazonomyces Bat. 1964, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *A. palmae* Bat. & Cavalc., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Amazonotheca Bat. & H. Maia 1959, *Schizothyriaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *A. sanitiae* Bat. & H. Maia, asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Phookamsak et al. (2016; accepted as *Schizothyriaceae*), cultures and sequences are unavailable.

Amazonotrema Kalb & Lücking 2009, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *A. nigrum* Kalb & Lücking, asexual morph unknown, lichenized, terrestrial, tropical, see Kalb (2009; new records), Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; classification), Lücking et al. (2017a; classification), sequences unavailable.

Ambarignomonia Sogonov 2008, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. petiolorum* (Schwein.) Sogonov, asexual morph

unknown, saprobes, USA, see Sogonov et al. (2008; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Amblyosporium Fresen. 1863 (= *Gueguenia* Bainier 1907; = *Searchomyces* B.S. Mehrotra & M.D. Mehrotra 1963 *fide* Seifert et al. 2011; Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, four species, type: *A. botrytis* Fresen., hyphomycetous, sexual morph unknown, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Ambrodiscus S.E. Carp. 1988, *Hyaloscrophaceae*, *Helotiidales*, *Leotiomycetes*, one species, type: *A. pseudotsugae* S.E. Carp., asexual morph unknown, on bark beetle galleries, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ambrosiella Brader ex Arx & Hennebert 1965, *Ceratocystidaceae*, *Microascales*, *Sordariomycetes*, seven species, type: *A. xylebori* Brader, hyphomycetous, obligate, mutualistic symbionts of ambrosia beetles, from bark beetle galleries/ mycangia in Ivory Coast, USA, Africa, Asia, Europe, USA, Japan, Taiwan, terrestrial, polyphyletic, some species belong to *Ophiostomataceae*, see Six et al. (2009; new species), Alamouti et al. (2009; phylogeny), Harrington et al. (2010, 2014; taxonomic revisions and new combinations), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Mayers et al. (2015; phylogeny), Maharachchikumbura et al. (2016b; outline), isotype: CBS 110.61, sequences are available at GenBank from Alamouti et al. (2009).

Ambrosiozyma Van der Walt 1972 (= *Hormoascus* Arx 1972 *fide* Species Fungorum 2017), *Saccharomycopsideaceae*, *Saccharomycetales*, *Saccharomycetes*, six species, type: *A. monospora* (Saito) Van der Walt, yeast or yeast-like, fungal symbionts of ambrosia beetles, widespread, see Endoh et al. (2008; phylogeny), Kirk et al. (2013; genus accepted), Kurtzman and Robnett (2013a; phylogeny), ex-type strains of type: CBS 2554.

Ameghiniella Speg. 1887 (= *Ionomidotis* E.J. Durand ex Thaxt. 1923; = *Polonioidiscus* Svrček & Kubička 1967 *fide* Species Fungorum 2017), *Cordieritidaceae*, *Helotiidales*, *Leotiomycetes*, two species, type: *A. australis* Speg., asexual morph unknown, bark beetle galleries, terrestrial, North and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny), cultures and sequences are available.

Ameliella Fryday & Coppins 2008, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *A. andreaeicola* Fryday & Coppins, lichenized, Europe and North America, see Coppins (2008; taxonomy), Fryday and Tønsberg (2015; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Ameroconium U. Braun & Zhurb. 2013, *Ascomycota* genera *incertae sedis*, one species, type: *A. hansfordii* Syd., hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, cosmopolitan, see Zhurbenko and Braun (2013; taxonomy), Wijayawardene et al. (2017; outline), holotype: LE 260838, cultures and sequences are unavailable.

Amerodiscosia Farr 1961, *Ascomycota* genera *incertae sedis*, one species, type: *A. renispora* M.L. Farr 1961, coelomycetous, sexual morph unavailable, saprobes, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Amerodiscosielina Bat. & Cavalc. 1966, *Ascomycota* genera *incertae sedis*, one species, type: *A. annonacearum* Bat. & Cavalc., coelomycetous, sexual morph unavailable, saprobes, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ameromassaria Hara 1918, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. japonica* Hara, asexual morph unknown, pathogens, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Amerosporiopsis Petr. 1941, *Ascomycota* genera *incertae sedis*, one species, type: *A. gaubae* Petr., coelomycetous, sexual morph unavailable, saprobes, Iran, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Amerosympodula Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *A. malaysiana* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Amicodisca Svrček 1987, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *A. brdensis* (Velen.) Svrček, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Han et al. (2012, 2014; new species, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Amniculicola Y. Zhang ter & K.D. Hyde 2008 (= *Anguillospora* Ingold 1942 *fide* Rossman et al. 2016a), *Amniculicolaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *A. lignicola* Y. Zhang ter & K.D. Hyde, asexual morph hyphomycetous, saprobes in freshwater habitats, Europe, see Zhang et al. (2008b, 2009c, 2012e; taxonomy, phylogeny), Wijayawardene et al. (2012, 2014c), Wanasinghe et al. (2015; phylogeny), Rossman et al. (2016a; nomenclature), sequence are available for all species, holotype of type: PC 0092661.

Amoenodochium Peláez & R.F. Castañeda 1996, *Ascomycota* genera *incertae sedis*, one species, type: *A. humicola* Peláez & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: MA-33881, cultures and sequences are unavailable, compare with *Acanthodochium*, *Asperisporium*, *Megalodochium* and *Polythrincium*.

Amoenomyces R.F. Castañeda, Saikawa & Hennebert 1996, *Ascomycota* genera *incertae sedis*, one species, type: *A. catenosporus* R.F. Castañeda, Saikawa & Hennebert, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: INIFAT C94/192, cultures and sequences are unavailable, compare with *Alternaria*, *Embellisia* and *Parapithomyces*.

Amorosia Mantle & D. Hawksw. 2006, *Amorosiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. littoralis* Mantle & D. Hawksw., hyphomycetous, sexual morph unknown, recovered from intertidal sediments, The Bahamas, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Jones et al. (2015; outline of marine genera), Thambugala et al. (2015b; phylogeny), holotype of the type: K(M) 139387, cultures and sequences are available, AM292047 (ITS).

Amorphomyces Thaxt. 1893, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, ten species, type: *A. falagriae* Thaxt., asexual morph unknown, dioecious, biotrophic on *Coleoptera* (Staphylinidae), terrestrial, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Amorphotheca Parbery 1969, *Amorphothecaceae*, *Leotiomycetes*, families *incertae sedis*, one species, type: *A. resinae* Parbery, asexual morph unknown, saprobes on resins, hydrocarbons, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wang et al. (2015b; applications), cultures available for the type: CBS 184.54.

Ampelomyces Ces. ex Schltl. 1852 (= *Byssocystis* Riess 1853; = *Cicinobolus* Ehrenb. 1853 *fide* Species Fungorum 2017), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *A. quisqualis* Ces., coelomycetous, sexual morph unknown, fungicolous, terrestrial, worldwide, see de Gruyter et al. (2009; phylogeny), Park et al. (2010; genetic variability), Wijayawardene et al. (2012, 2014c, 2017; taxonomy, phylogeny, outline), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*), Verkley et al. (2014; phylogeny), cultures of the type: CBS 131.31, CBS 129.79.

Amphibambusa D.Q. Dai & K.D. Hyde 2015, *Cainiaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *A. bambusicola* Dai & Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Liu et al. (2015a; phylogeny, taxonomy), Senanayake et al. (2015; phylogeny), holotype and ex-type culture of type: MFLUCC 11-0617, MFLU 14-0825.

Amphichaetella Höhn. 1916 (= *Amphichaete* Kleb. 1914 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *A. echinata* (Kleb.) Höhn., hyphomycetous, sexual morph unknown, Europe, Australia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Amphilogia Gryzenh., H.F. Glen & M.J. Wingf. 2005, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *A. gyrosa* (Berk. & Broome) Gryzenh., H.F. Glen & M.J. Wingf., asexual morph coelomycetous, pathogenic, terrestrial, temperate, Sri Lanka, New Zealand, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-epitype strain of the type: CBS H-20178, CMW 10470.

Amphimyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *A. cerylonis* Thaxt., asexual morph unknown, biotrophic on *Coleoptera* from the genus *Cerylon* (Cerylonidae), West Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Amphiporthe Petr. 1971 (= *Amphicytostroma* Petr. 1921 *fide* Rossman et al. 2015a), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *A. tiliae* (Sacc.) Petr., asexual morph hyphomycetous, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b), Rossman et al. (2015a; nomenclature).

Amphirosellinia Y.M. Ju, J.D. Rogers, H.M. Hsieh & Lar.N. Vassiljeva 2004, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, five species, type: *A. nigrospora* Y.M. Ju, J.D. Rogers & H.M. Hsieh, asexual morph unknown, saprobes, terrestrial, north temperate, see Hsieh et al. (2010; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: HAST Ju & Hsieh 91092308, sequences are available for the type species.

Amphisphaerellula Gucevič 1952, *Sordariomycetes*, genera *incertae sedis*, two species, type: *A. fagi* Gucevič, sexual morph unknown, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Amphisphaeria Ces. & De Not. 1863 (= *Macrothelia* M. Choisy 1949; = *Massariella* Speg. 1880; = *Massariopsis* Niessl 1876; = *Phorcys* Niessl 1876; = *Poikiloderma* Füsting 1868; = *Rhynchostomopsis* Petr. & Syd. 1923 *fide* Species Fungorum 2017), *Amphisphaeriaceae*, *Amphisphaerales*, *Sordariomycetes*, 196 species, type: *A. umbrina* (Fr.) De Not., asexual morph coelomycetous, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Senanayake et al. (2015, morphology and phylogeny), Maharachchikumbura (2015b, 2016b; morphology, phylogeny), sequences are available.

Amphisphaerina (Cooke) Höhn 1919, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. texensis* (Cooke) Höhn, asexual morph unknown, saprobes, terrestrial, USA, See Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Amphobotrys Hennebert 1973, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *A. ricini* (N.F. Buchw.) Hennebert, hyphomycetous, sexual morph unknown, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, need generic revision.

Amphoropycnum Bat. 1963, *Ascomycota* genera *incertae sedis*, one species, type: *A. philippinense* Bat., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Amphorothecium P.M. McCarthy, Kantvilas & Elix 2001, *Ostropales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *A. occultum* P.M. McCarthy, Kantvilas & Elix, lichenized, terrestrial, Australia, see Lumbsch and

Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences unavailable.

Amphorulopsis Petr. 1959, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. polygonacearum* Petr., asexual morph unknown, terrestrial, former Yugoslavia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Index Fungorum (2017) listed the type species as a synonym of *Ceriospora polygonacearum*.

Ampliotrema Kalb ex Kalb 2006 (= *Ampliotrema* Kalb 2004 *fide* Species Fungorum 2017), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, c. 20 species, type: *A. amplius* (Nyl.) Kalb, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Sipman et al. (2012; key Costa Rica), Rivas Plata et al. (2012b; phylogeny), Cáceres et al. (2012b), Rivas Plata and Lücking (2012; classification), Mercado-Díaz et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Amplistroma Huhndorf, A.N. Mill., Greif & Samuels 2009, *Amplistromataceae*, *Amplistromatales*, *Sordariomycetes*, nine species, type: *A. carolinianum* Huhndorf, A.N. Mill., Greif & Samuels, asexual morph hyphomycetous, acrodontium-like in culture, sexual morph ascomycetes, saprobes, French Guiana, Costa Rica, Spain, Singapore, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura (2015b, 2016b; morphology, phylogeny), holotype: BPI 878925, sequences are available.

Ampullicephala R.F. Castañeda, Minter & M. Stadler 2009, *Ascomycota* genera *incertae sedis*, one species, type: *A. setiformis* (R.F. Castañeda) R.F. Castañeda, Minter & M. Stadler, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Castañeda-Ruiz et al. (2009b; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: INIFAT C85/79, cultures and sequences are unavailable, compare with *Gonatobotryum* and *Gonatobotrys*.

Ampulliferina B. Sutton 1969, *Ascomycota* genera *incertae sedis*, three species, type: *A. persimplex* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: IMI 134269.

Amundsenia Søchting, Garrido-Benavent, Arup & Frödén 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *A. austrocontinentalis* Garrido-Benavent, Søchting, Pérez-Ortega & Seppelt, lichenized, see Søchting et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Amicosphaerella Quaedvl. & Crous 2014, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one

species, type: *A. africana* (Crous & M.J. Wingf.) Quaedvl. & Crous, asexual morph unknown, pathogens, terrestrial, Australia, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), holotype and ex-type of the type: PERTH 05849543, CBS 110500.

Amygdalaria Norman 1853, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, 14 species, type: *A. pelobotryon* (Wahlenb.) Norman, lichenized, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Amylascus Trappe 1971, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, five species, type: *A. herbertianus* (J.W. Cribb) Trappe, mycorrhizal, Australia, the Americas, see Lumbsch and Huhndorf (2010; outline), Healy et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are for several undescribed species.

Amylis Speg. 1922, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. memorabilis* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Amylocarpus Cur. 1859 (= *Plectolitus* Kohlm. 1960 *fide* Species Fungorum 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *A. encephaloides* Curr., asexual morph unknown, drift wood, temperate marine waters, saprobe, sequence are available, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rämä et al. (2014; DNA, records in Norway), Jones et al. (2015; outline), cultures and sequences are available.

Amylopora Rambold 1994, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, one species, type: *A. cervinocuprea* (Arnold) Rambold, lichenized, see Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Anabahusakala Carmo, J.S. Monteiro, Gusmão & R.F. Castañeda 2014, *Ascomycota* genera *incertae sedis*, one species, type: *A. amapensis* L.T. Carmo, J.S. Monteiro, Gusmão & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, Brazil, Carmo et al. (2014; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Anacacumisporium Y.R. Ma & X.G. Zhang 2016, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *A. appendiculatum* Y.R. Ma & X.G. Zhang, hyphomycetous, sexual morph unknown, saprobic, terrestrial, China, see Ma et al. (2016c; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Anacoronospora J.S. Monteiro, Gusmão & R.F. Castañeda 2016, *Ascomycota genera incertae sedis*, one species, type species: *A. diversiseptata* J.S. Monteiro, Gusmão & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobic, Brazil, see Monteiro et al. (2016b; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Anacraspedodidymum C.R. Silva, R.F. Castañeda & Gusmão 2014, *Ascomycota genera incertae sedis*, two species, type: *A. aquaticum* C.R. Silva, R.F. Castañeda & Gusmão, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Silva et al. (2014a; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Anaexserticlava T.S. Santa Izabel, R.F. Castañeda & Gusmão 2015, *Ascomycota genera incertae sedis*, one species, type: *A. caatingae* T.S. Santa Izabel, R.F. Castañeda & Gusmão, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Santa Izabel et al. (2015; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Anamylopsora Timdal 1991, *Baeomycetaceae*, *Baeomycetales*, *Lecanoromycetes*, one species, type: *A. pulcherrima* (Vain.) Timdal (*Lecidea pulcherrima* Vain.), lichenized, saxicolous, terricolous, temperate northern hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Resl et al. (2015; phylogeny), holotype: TUR-V-22644 (holotype), sequences are available.

Anaphysmene Bubák 1906 (= *Wakefieldia* G. Arnaud, Bull 1954 [1953] *fide* Species Fungorum 2017), *Ascomycota genera incertae sedis*, two species, type: *A. heraclei* (Lib.) Bubák, coelomycetous, sexual morph unknown, Europe, Gautamala see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Anapleurothecium Hern.-Restr. et al. 2017, *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, one species, type: *A. botulisporum* Hern.-Restr., J. Mena & Gene, hyphomycetous, sexual morph unknown, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Anaptychia Körb. 1848 (= *Anaptychiomyces* E.A. Thomas 1939; = *Hagenia* Eschw. 1824; = *Imbricaria* (Schreb.) Michx. 1803; = *Lichen* sect. *Imbricaria* Schreb. 1791; *Lichenoides* Hoffm. 1789 [1790]; = *Parmelia* sect. *Imbricaria* (Schreb.) Fr. 1825; = *Tornabenia* A. Massal. 1853 *fide* Species Fungorum 2017), *Physciaceae*, *Caliciales*, *Lecanoromycetes*, 15 species, type: *A. ciliaris* (L.) Körb. ex A. Massal., lichenized, see Gaya et al. (2012; phylogeny), Prieto et al. (2013; phylogeny), Miadlikowska et al. (2014a; phylogeny), Prieto and Wedin (2016;

classification), Lücking et al. (2017a; classification), sequences are available.

Anarhyma M.H. Pei & Z.W. Yuan 1986, *Ascomycota genera incertae sedis*, one species, type: *A. rhododendri* M.H. Pei & Z.W. Yuan, coelomycetous, sexual morph unknown, saprobes, terrestrial, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Anariste Syd. 1927, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *A. poliothea* Syd., asexual morph unknown, epiphytes, terrestrial, Costa Rica, see Hofmann (2009; review, accepted as a genus in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, keys, transferred to *Micropeltidaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Micropeltidaceae*), placement in *Micropeltidaceae* supported by morphology, syntype of type: BPI 690985, cultures and sequences are unavailable.

Anaselenosporella Heredia, R.F. Castañeda & R.M. Arias 2010, *Ascomycota genera incertae sedis*, two species, type: *A. sylvatica* Heredia, R.F. Castañeda & R.M. Arias, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Mexico, see Castañeda-Ruiz et al. (2010a; morphology), Pratibha et al. (2011; new species), Wijayawardene et al. (2012, 2017; outline), holotype of the type: MUCL 45630, cultures and sequences are unavailable, compare with *Selenosporella* and *Selenosporopsis*.

Anaseptoidium R.F. Castañeda, Heredia & R.M. Arias 2012, *Ascomycota genera incertae sedis*, one species, type: *A. mycophilum* R.F. Castañeda, Heredia & R.M. Arias, hyphomycetous, sexual morph unknown, mycophilous, on synnemata of *Phaeoisaria clavulata*, terrestrial, Mexico, see Castañeda-Ruiz et al. (2010a; taxonomy), Wijayawardene et al. (2017; outline), holotype of type: XAL CB1696, cultures and sequences are unavailable, compare with *Septoidium*.

Ancistrospora G. Thor. 1995 (= *Ancistrospora* G. Thor 1991 *fide* Species Fungorum 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *A. australiensis* (G. Thor) G. Thor, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ertz et al. (2015c; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Ancorasporella Mig. Rodr. 1982, *Ascomycota genera incertae sedis*, one species, type: *A. cubensis* Mig. Rodr., hyphomycetous, sexual morph unknown, saprobes, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Andamanomyces Hosag. 2014, *Asterinales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *A. fragariae* Hosag., S.P. Mathew & D. Babu, asexual morph unknown, epiphytes, biotrophic, terrestrial, India, see Hosagoudar et al. (2014; morphology, accepted as a genus in *Asterinales*), placement in *Asterinales* supported by morphology, holotype: TBGT 6666, cultures and sequences are unavailable.

Andreomyces B.P. Hodk. & Lendemer 2013, *Andreomycetaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *A. morozianus* (Lendemer) B.P. Hodk. & Lendemer, lichenized, see Hodkinson and Lendemer (2013; taxonomy), Lücking et al. (2017a; classification), cultures available for the type: NY1377.

Anema Nyl. ex Forssell 1885 (= *Omphalaria* A. Massal. 1855 *fide* Species Fungorum 2017), *Lichinaceae*, *Lichinales*, *Lichinomycetes*, ten species, type: *A. decipiens* (A. Massal.) Forssell, with ascocarps, lichenized (cyanobacteria), terrestrial (saxicolous), arid, semiarid to warm temperate regions, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), see sequences are available.

Angatia Syd. & P. Syd. 1914, *Saccardiaceae*, *Ascomycota* families *incertae sedis*, five species, type: *A. eugeniae* Syd. & P. Syd., asexual morph unknown, saprobes, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Angelina Fr. 1849, *Helotiaceae* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. rufescens* (Schwein.) Duby, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dermateaceae*), Kirk et al. (2013; genus accepted), Karakehian et al. (2014; DNA, phylogeny), Jaklitsch et al. (2016a; classification), culture available for the type: JK 12040101.

Angiactis Aptroot & Sparrius 2008, *Arthoniales* genus *incertae sedis*, *Arthoniomycetes*, three species, type: *A. littoralis* (Kantvilas) Aptroot & Sparrius, lichenized, terrestrial, tropical, see Aptroot et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Angiopomopsis Höhn. 1912, *Ascomycota* genera *incertae sedis*, one species, type: *A. lophostoma* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Java, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Anguillosporella U. Braun 1995, *Ascomycota* genera *incertae sedis*, two species, type: *A. vermiformis* (Davis) U. Braun., hyphomycetous, terrestrial, foliicolous, leaf-

spotting, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), lectotype of type: BPI 442755, cultures and sequences are unavailable.

Angulimaya Subram. & Lodha 1964, *Ascomycota* genera *incertae sedis*, one species, type: *A. sundara* Subram. & Lodha, hyphomycetous, sexual morph unknown, on dung, terrestrial, Asia, see Seifert et al. (2011; morphology, accepted as a genus in *Lasiosphaeriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline, treated as *Ascomycota* genera *incertae sedis*), cultures and sequences are unavailable.

Angulospora Sv. Nilsson., 1962, *Ascomycota* genera *incertae sedis*, one species, type: *A. aquatica* Sv. Nilsson, hyphomycetous, sexual morph unknown, aquatic, on submerged decaying wood, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Angustumassarina Thambug., Kaz. Tanaka & K.D. Hyde 2015, *Amorosiaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *A. populi* Thambug. & K.D. Hyde, asexual morph hyphomycetous, fungicolous or saprobes, terrestrial, Germany and Italy, three species, see Thambugala et al. (2015b; morphology, phylogeny), Tibpromma et al. (2017b; new species), holotype and ex-type strain of type: MFLU 14-0588, MFLUCC 13-0034.

Angustospora Abdel-Aziz 2016, *Testudinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. nilensis* Abdel-Aziz, asexual morph unknown, saprobes, aquatic, Egypt, see Li et al. (2016a; taxonomy, phylogeny), holotype and ex-isotype of the type: MFLU 15-1511, MF804.

Anhelia Racib. 1900 (= *Agostaea* (Sacc.) Theiss. & Syd. 1915; = *Ramosiella* Syd. & P. Syd. 1917; = *Trabutia* subgen. *Agostaea* Sacc. 1895; = *Whetzelomyces* Viégas 1945 *fide* Species Fungorum 2017), *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, nine species, type: *A. tristis* Racib., asexual morph unknown, parasitic on leaves, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Pinho et al. (2012; morphology, phylogeny), Kirk et al. (2013; genus accepted), Dissanayake et al. (2014; morphology), holotype of the type: F47931.

Aniptodera Shearer & M. A. Mill. 1977, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, c. 16 species, type: *A. chesapeakensis* Shearer & Mill., asexual morph unknown, saprobes, freshwater, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline),

genus polyphyletic, ex-type strains of type: CS-97-26 =ATCC 32818, holotype: ILLS 36523.

Anisogramma Theiss. & Syd. 1917 (= *Apioporth* Höhn. 1917 *fide* Species Fungorum 2017), *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, three species, type: *A. virgultorum* (Fr.) Theiss. & Syd., asexual morph unknown, terrestrial, saprobes, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Anisomeridium (Müll. Arg.) M. Choisy 1928 (= *Arthopyrenia* sect. *Anisomeridium* Müll. Arg., Flora 1883; = *Compsosporiella* Sankaran & B. Sutton 1991; = *Ditremis* Clem. 1909; = *Lembidium* Körb. 1855; = *Microthelia* Körb. 1855; = *Sarcinulella* B. Sutton & Alcorn 1983 *fide* Species Fungorum 2017), *Monoblastiaceae*, *Monoblastiales*, *Dothideomycetes*, c. 200 species, type: *A. xylogena* Müll. Arg., asexual morph unknown, lichenized, terrestrial, mostly tropical, see Nelsen et al. (2009, 2011b; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Anisomyces Theiss. & Syd. 1914 (= *Apiospora* subgen. *Phaeoapiospora* Sacc. & P. Syd. 1902; *Phaeoapiospora* (Sacc. & P. Syd.) Theiss. & Syd. 1915; = *Stegastroma* Syd. & P. Syd. 1916 *fide* Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, seven species, type: *A. papilloideoseptatus* (Henn.) Theiss. & Syd., asexual morph unknown, saprobes?, terrestrial, tropical America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Anisomycopsis I. Hino & Katum. 1964, *Diaporthales*, genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. rosae* I. Hino & Katum., asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Anisostagma K.R.L. Petersen & Jørg. Koch 1996, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *A. rotundatum* K.R.L. Petersen & Jørg. Koch, asexual morph unknown, saprobes, marine, Denmark, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: K. R. L. Petersen 9 slides numbered 1-9, Herbarium C, Copenhagen, cultures and sequences are unavailable.

Annelloidentimyces Matsush. 1985, *Ascomycota* genera *incertae sedis*, one species, type: *A. antennatuloideus* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011;

morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Annellodochium Deighton 1969, *Ascomycota* genera *incertae sedis*, one species, type: *A. ramulisperum* Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Annelolacinia B. Sutton 1964, *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, two species, type: *A. dinemasporioides* B. Sutton, coelomycetous, sexual morph unknown, pathogens, Ghana, Sierra Leone, Trinidad, India, Australia, cultures and sequences are unavailable, see Wijayawardene et al. (2012, 2016b, 2017; taxonomy, outline), Kirk et al. (2013; genus accepted), holotype: IMI37897c.

Anellophora S. Hughes 1952 (= *Chaetotrichum* Syd. 1927 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, 13 species, type: *A. solani* (Syd.) S. Hughes, hyphomycetous, sexual morph unknown, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Anellophorella Subram. 1962, *Ascomycota* genera *incertae sedis*, five species, type: *A. densa* (Syd. & P. Syd.) Subram., hyphomycetous, sexual morph unknown, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequence are unavailable.

Anellophragmia Subram. 1963, *Ascomycota* genera *incertae sedis*, one species, type: *A. coonoorensis* (Subram.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Annelospermosporella P.R. Johnst. 1999, *Ascomycota* genera *incertae sedis*, one species, type: *A. meliolinae* P.R. Johnst., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Annelosympodia McTaggart, R.G. Shivas & U. Braun 2007, *Ascomycota* genera *incertae sedis*, one species, type: *A. orbiculata* McTaggart, Shivas & Braun, hyphomycetous, sexual morph unknown, pathogens, terrestrial, Australia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: PERTH 03270173, isotype: BRIP 49805, paratype BRIP 19908, living cultures unavailable.

Annellosympodiella Crous & Assefa 2014, *Mycosphaerellaceae, Capnodiales, Dothideomycetes*, one species, type: *A. juniperi* Crous & Assefa, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Ethiopia, see Crous et al. (2014c; morphology, phylogeny), Wijayawardene et al. (2014c; outline), holotype of the type: CBS H-21706, ex-type strain of type: CPC 23276.

Annulatascus Ces. & De Not. 1863, *Annulatascaeae, Annulatascales, Sordariomycetes*, c. 19 species, type: *A. velatispora* K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline), Boonyuen et al. (2012; new species; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Luo et al. (2015b; taxonomy), Dayarathne et al. (2016a; epitypification of *A. velatisporus*), holotype of type: BRIP 17373; epitype MFLU 16-2204.

Annulohypoxylon Y.M. Ju, J.D. Rogers & H.M. Hsieh 2005, *Hypoxylaceae, Xylariales, Sordariomycetes*, three species, type: *A. truncatum* (Schwein.) Y.M. Ju, J.D. Rogers & H.M. Hsieh, four species, asexual morph hyphomycetous, terrestrial, saprobes, Europe, North America, see Marincowitz et al. (2008; new species, South Africa), Hladki and Romero (2009; new species), Lumbsch and Huhndorf (2010; outline), Pereira et al. (2010; new species, Brazil), Liu et al. (2015a; new species), Li et al. (2016a; new species, Thailand), Maharachchikumbura et al. (2015b, 2016b; outline), Vasilyeva et al. (2016; new species), Kuhnert et al. (2017; new species, revision), Wendt et al. (2017; phylogeny), cultures and sequences are available.

Annulusmagnus J. Campb. & Shearer 2004, *Annulatascaeae, Annulatascales, Sordariomycetes*, one species, type: *A. triseptatus* (Wong, Hyde & Jones) Campb. & Shearer, asexual morph unknown, saprobes, aquatic, Australia, Canada, USA, see Réblová et al. (2012; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), neotype of the type: HKU889, cultures available CBS 128831.

Anomalographis Kalb 1992, *Graphidaceae, Ostropales, Lecanoromycetes*, two species, type: *A. madeirensis* (Tav.) Kalb, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Anomomorpha Nyl. ex Hue 1891 (= *Anomomorpha* Nyl. ex Hue, Nouv 1891; = *Anomomorpha* Nyl. 1889; = *Aulacographa* Leight. 1854; = *Ctesium* Pers. 1827 [1826-1830]; = *Digraphis* Clem. 1909; = *Diplographis* Kremp. ex A. Massal. 1860 [1859-1860]; = *Dyplolabia* A. Massal. 1854; = *Graphidomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Opegrapha* Humb. 1793; = *Oxystoma* Eschw.

1824; = *Scaphis* Eschw. 1824; = *Thelographis* Nyl. 1857 *fide* Species Fungorum (2017), *Graphidaceae, Ostropales, Lecanoromycetes*, eight species, type: *A. turbulentula* (Nyl) Hue, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; phylogeny), Lücking et al. (2017a; classification), sequences are unavailable.

Anopeltis Bat. & Peres 1960, *Dothideomycetes* genera *incertae sedis*, one species, type: *A. venezuelensis* Bat. & Peres, asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Capnodiaceae*), Hyde et al. (2013) did not accept as a genus in *Capnodiaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Anopodium N. Lundq. 1964, *Lasiostphaeriaceae, Sordariales, Sordariomycetes*, two species, type: *A. ampullaceum* N. Lundq., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kruys et al. (2015; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Antarctomyces Stchigel & Guarro 2001, *Thelebolaceae, Thelebolales, Leotiomyces*, one species, type: *A. psychrotrophicus* Stchigel & Guarro, asexual morph sporothrix-like, saprobes, terrestrial, Antarctica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Bansal et al. (2014; antifreeze proteins), cultures and sequences are available.

Anteaglonium Mugambi & Huhndorf 2009, *Anteagloniaceae, Pleosporales, Dothideomycetes*, six species, type: *A. abbreviatum* (Schwein.) Mugambi & Huhndorf, asexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Mugambi and Huhndorf (2009a, 2010, taxonomy), Almeida et al. (2014b, new species), Hyde et al. (2013, included in to new family, *Anteagloniaceae*), Wijayawardene et al. (2014c; outline), Jayasiri et al. (2016c; new species, asexual morph of this genus and reference specimens), cultures available for the type: ANM 925aT.

Antealophiotrema A. Hashim. & Kaz. Tanaka 2017, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *A. brunneosporum* (Ying Zhang, J. Fourn. & K.D. Hyde) A. Hashim. & Kaz. Tanaka, asexual morph hyphomycetous, saprobes, Europe, see Hashimoto et al. (2017; taxonomy, phylogeny), sequences are available.

Antennatula Fr. ex F. Strauss 1850 (= *Hormisciella* Bat. 1956 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, ten species, type: *A. pinophila* Fr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011;

morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Antennopsis R. Heim 1952, *Ascomycota* genera *incertae sedis*, three species, type: *A. gallica* R. Heim & Buchli ex R. Heim, hyphomycetous, sexual morph unknown, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Antennospora Meyers 1957, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *A. caribbea* Meyers, asexual morph unknown, saprobes, marine, worldwide, see Pang et al. (2008b; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Antennulariella Woron. 1915 (= *Antennariella* Bat. & Cif.; = *Capnodendron* S. Hughes 1963; = *Capnociferria* Bat. 1963; = *Capnocrinum* Bat. & Cif. 1963; = *Capnodina* (Sacc.) Sacc. 1926; = *Capnodium* subgen. *Capnodina* Sacc. 1913 *fide* Wijayawardene et al. 2014c; Species Fungorum 2017), *Antennulariellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *A. fuliginosa* Woron., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Antennulariellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, treated *Antennariella* Bat. & Cif. and *Capnodendron* S. Hughes as synonyms), Rossman et al. (2015b) did not agree with Wijayawardene et al. (2014c), Hongsanan et al. (2016c; evolution), cultures and sequences are unavailable.

Anthocanalis Daranag., Camporesi & K.D. Hyde 2015, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *A. sparti* Daranag., Camporesi & K.D. Hyde, asexual morph nodulisporium-like, saprobes, terrestrial, Europe, see Daranagama et al. (2015a), Maharachchikumbura et al. (2015b, 2016b), Wendt et al. (2017; phylogeny), holotype and ex-type strains of type: MFLU 14–0225, MFLUCC 14–0010.

Anthonectria Döbbeler 2010, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, eight species, type: *A. mammispora* Döbbeler, asexual morph unknown, saprobes, epiphytes, terrestrial, cosmopolitan, see Döbbeler (2010; taxonomy), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Anthopsis Fil. March., A. Fontana & Luppi Mosca 1977, *Ascomycota* genera *incertae sedis*, three species, type: *A. deltoidea* Fil. March., A. Fontana & Luppi Mosca, asexual morph hyphomycetous, Europe, Japan, see Seifert

et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequence are unavailable.

Anthostoma Nitschke 1867, *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *N. stictarum* Nannf. & R. Sant., asexual morph unknown, saprobes, see Kirk et al. (2008; treated as a synonym of *Cryptosphaeria* Ces. & De Not.), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a distinct genus), cultures and sequences are available.

Anthostomaria (Sacc.) Theiss. & Syd. 1918 (= *Anthostomella* subgen. *Anthostomaria* Sacc. 1905 *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, one species, type: *A. apogyla* (Nyl.) Theiss. & Syd., asexual morph unknown, lichenicolous, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline).

Anthostomella Sacc. 1875 (= *Cryptosordaria* De Not. ex Sacc. 1891; = *Entosordaria* Speg. 1910, nom. inval.; = *Maurinia* Niessl 1876; = *Myconenesia* Kirschst. 1936; ≡ *Neesiella* Kirschst. 1935; = *Paranthostomella* Speg. 1910; = *Phaeophomatospora* Speg. 1909 *fide* Wendt et al. 2017), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, more than 100 species, type: *A. tomentoides* Sacc., asexual morph hyphomycetous, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Daranagama et al. (2015a, 2016a; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Dai et al. (2017; new species), cultures and sequences are available.

Anthostomellina L.A. Kantsch. 1928, *Sordariomycetes* genera *incertae sedis*, two species, type: *A. carpinea* L.A. Kantsch., asexual morph unknown, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline).

Anthostomelloides Tibpromma & K.D. Hyde 2016, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, five species, type: *A. krabiensis* Tibpromma & K.D. Hyde, asexual morph unknown, terrestrial, saprobes, Asia, see Tibpromma et al. (2017a; taxonomy), Wendt et al. (2017; placed in *Hypoxylaceae*), cultures and sequences are available.

Anthracobia Boud. 1885, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 20 species, type: *A. melaloma* (Alb. & Schwein.) Boud., scyphalidium-like asexual morph has been reported, saprobes, terrestrial, on burnt soil, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hansen et al. (2013; phylogeny), cultures and sequences are available.

Anthracocarpon Breuss 1996, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, three species, type: *A. virescens* (Zahlbr.) Breuss, lichenized, see Lumbsch and Huhndorf (2010; outline), Prieto et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Anthracoderma Speg. 1888, *Ascomycota* genera *incertae sedis*, three species, type: *A. hookeri* Speg., coelomycetous, sexual morph unknown, South America see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Anthracotheicum Hampe ex A. Massal. 1860 (= *Phaeodictyon* M. Choisy 1929 *fide Species Fungorum* 2017), *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, five species, type: *A. doleschallii* A. Massal., asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Aptroot (2012; key), Weerakoon et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Gueidan et al. (2016; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type species.

Antimanoa Syd. 1930, *Ascomycota* genera *incertae sedis*, one species, type: *A. grislea* Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Antoniomyces Inácio 2012, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, one species, type: *A. loranthicola* Inácio, Araúz & M. Piepenbr., asexual morph unknown, epiphytes, terrestrial, Panama, see Inácio et al. (2012; morphology, accepted as a genus in *Parmulariaceae*), Hyde et al. (2013; review, keys), Wijayawardene et al. (2014c; outline), placement in *Parmulariaceae* supported by morphology, holotype of type: PMA, cultures and sequences are unavailable.

Antromyces Fresen. 1850, *Ascomycota* genera *incertae sedis*, four species, type: *A. copridis* Fresen., hyphomycetous, sexual morph unknown, Europe, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Anulohypha Cif. 1962, *Ascomycota* genera *incertae sedis*, one species, type: *A. serpens* Cif., hyphomycetous, sexual morph unknown, Dominican Republic, see Seifert et al. (2011; did not accept), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Anungitea B. Sutton 1973, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, 17 species, type: *A. fragilis* B.

Sutton, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014c, 2016a; new species), Xia et al. (2015b; new species), cultures and sequences are available but lacking for the type species.

Anungitopsis R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, nine species, type: *A. speciosa* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type species: INIFAT C88/275, sequences are available, compare with *Anungitea* and *Iyengarina*.

Anaverticicladus P.O. Costa, Malosso & R.F. Castañeda 2016, *Ascomycota* genera *incertae sedis*, one species, type: *A. uncinatus* P.O. Costa, Malosso & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobic, terrestrial Brazil, see Costa et al. (2016; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Anzia Stizenb. 1861 (= *Chondrospora* A. Massal. 1860 [1859-1860] *fide Species Fungorum* 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, c. 25 species, type: *A. colpodes* (Ach.) Stizenb., with ascocata, lichenized (green algae), terrestrial, tropical to subtropical, see Crespo et al. (2010b; phylogeny), Lumbsch and Huhndorf (2010; outline), Jayalal et al. (2012b; new species), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; phylogeny), Divakar et al. (2015; phylogeny), Wang et al. (2015d; new species), Lücking et al. (2017a; classification), sequences are available.

Anzina Scheid. 1982, *Ostropales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *A. carneonivea* (Anzi) Scheid., lichenized, terrestrial, Eurasia, see Lumbsch and Huhndorf (2010; outline), Wang et al. (2012; China), Kirk et al. (2013; genus accepted), Resl et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Aoria Cif. 1962, *Ascomycota* genera *incertae sedis*, one species, type: *A. amphistroma* Cif., coelomycetous, sexual morph unknown, Dominican Republic, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Aparaphysaria Speg. 1922, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, two species, type: *A. doelloi* Speg., asexual morph unknown, terrestrial, South America, Tierra del Fuego, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Apatelomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *A. ogmoceri* Thaxt., asexual morph unknown, biotrophic only host known is *Ogmocerus* sp. (Coleoptera, Pselaphidae), Afrotropical, Liberia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Apatomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *A. laboulbenoides* Thaxt., asexual morph unknown, dioecious, biotrophic only host known is *Tachys ceylonicus* Nietn. (Coleoptera, Carabidae), Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Apatopla Poelt & Hafellner 1980 (see Species Fungorum 2017 for synonyms), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *A. oblongula* (H. Magn.) Poelt & Hafellner, lichenized, see Vondrák et al. (2012; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Apendiella Quaedvly. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *A. strumelloidea* (Milko & Dunaev) Quaedvly. & Crous, hyphomycetous, sexual morph unknown, pathogens, terrestrial, Russia, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; outline), holotype of the type: BKMF-2534, ex-type strain of type: CBS 114484.

Aphanandromyces W. Rossi 1982, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *A. audisioi* W. Rossi, asexual morph unknown, biotrophic only host known is *Brachypterus urticae* (Coleoptera, Kateretidae), Europe, temperate to cool, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Aphanoascus Zukal 1890, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, c. 10 species, type: *A. cinnabarinus* Zukal, asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), sequences and culture available.

Aphanocladium W. Gams 1971, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *A. album* (Preuss) W. Gams, asexual morphs unknown, on myctozoans, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Aphanofalx B. Sutton 1986, *Ascomycota* genera *incertae sedis*, two species, type: *A. mali* B. Sutton, coelomycetous, sexual morph unknown, saprobes, Pakistan, Zambia, see Wijayawardene et al. (2012, 2017;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: IMI293036.

Aphanophora Réblová & Unter. 2013, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, three species, type: *A. eugeniae* (Crous & Alfenas) Réblová & Unter., hyphomycetous, sexual morph unknown, from living leaves of *Eugenia uniflora*, terrestrial, Rio Grande do Sul, see Réblová et al. (2013; taxonomy), ex-type cultures of the type: CBS 124105.

Aphanopsis Nyl. ex Sydow 1882 (= *Aphanopsis* Nyl. ex P. Syd. 1887 *fide* Species Fungorum 2017), *Aphanopsidaceae*, *Ascomycota* families *incertae sedis*, two species, type: *A. terrigena* (Ach.) Nyl. ex P. Syd., lichenized, terrestrial (terricolous), Europe, see Lumbsch and Huhndorf (2010; outline), Printzen et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Aphanostigme Syd. 1926 (= *Gomezina* Chardón & Toro 1934; = *Somatexis* Toro, in Chardon & Toro 1934 *fide* Species Fungorum 2017), *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, 21 species, type: *A. solani* Syd., asexual morph unknown, ascocata occurring as a saprophyte or a leaf hair parasite, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Aphanotria Döbbeler 2007, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *A. paradoxa* Döbbeler, asexual morph unknown, saprobes, Colombia, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Apharia Bonord. 1864, *Sordariomycetes* genera *incertae sedis*, one species, type: *H. nigrescens* Bonord., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Aphidomyces Brain 1923, *Saccharomycetales* genera *incertae sedis*, *Saccharomycetes*, three species, type: no typification data, asexual morph unknown, in insects, terrestrial, worldwide, see Lumbsch and Huhndorf (2010) did not include, Kirk et al. (2013; genus accepted).

Apiculospora Wijayaw., Camporesi, A.J.L. Phillips & K.D. Hyde 2016, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. spartii* Wijayaw., W.J. Li, Camporesi, A.J.L. Phillips & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2016b, 2017;

taxonomy, phylogeny, outline), holotype and ex-type of the type: MFLU 15-3556, MFLUCC 13-0400.

Apinisia La Touche 1968, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *A. graminicola* La Touche, asexual morph unknown, saprobes, terrestrial, Europe, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), culture available for the type: CBS 721.68.

Apiocamarops Samuels & J.D. Rogers 1987, *Boliniaceae*, *Boliniales*, *Sordariomycetes*, three species, type: *A. alba* Samuels & J.D. Rogers, asexual morphs unknown, saprobes, terrestrial, Central and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Apiocarpella Syd. & P. Syd. 1919 (= *Apiosporella* Speg. 1910 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, eight species, type: *A. macrospora* (Speg.) Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but unpublished.

Apioclypea K.D. Hyde 1994, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, six species, type: *A. livistonae* K.D. Hyde, asexual morphs unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, treated as *Clypeosphaeriaceae*), Hongsan et al. (2017a; phylogeny, evolution, treated *Clypeosphaeriaceae* as a synonym of *Xylariaceae*), cultures and sequences are available for unidentified species.

Apiodiscus Petr. 1940, *Rhytismatales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. gillii* Petr., asexual morph unknown, saprobes, terrestrial, Iran, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Apiognomonia Höhn. 1917, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, c. 15 species, type: *A. veneta* (Sacc. & Speg.) Höhn., asexual morph unknown, saprobes, terrestrial, North America, Europe, see Lumbsch and Huhndorf (2010; outline), Vasilyeva and Stephenson (2010; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Crous et al. (2016a; new species), cultures and sequences are available.

Apioplaciostoma M.E. Barr 1978, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *A. populi* (E.K. Cash & Waterman) M.E. Barr, asexual morph discula-like, weak pathogens, terrestrial, temperate, Europe, North America, see Lumbsch and Huhndorf (2010;

outline, accepted genus in *Valsaceae*), Sogonov et al. (2008; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Apioporthella Petr. 1929 (= *Apiothecium* Lar.N. Vassiljeva 1987 *fide* Species Fungorum 2017), *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. bavarica* Petr., asexual morph unknown, saprobe, terrestrial, temperate, Europe, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Apiorhynchostoma Petr. 1923, *Boliniaceae*, *Boliniales*, *Sordariomycetes*, five species, type: *A. apiculatum* (Sacc.) Petr., asexual morphs unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Apiosordaria Arx & W. Gams. 1967 (= *Cainea* S. Hughes 1951; = *Echinopodospora* B.M. Robison 1970; = *Echinospora* Mirza 1963; = *Lacunospora* Cailleux 1969 [1968] *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, 23 species; type: *A. verruculosa* Arx & W. Gams, asexual morph unknown, saprobes, endophytes, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; genus accepted), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences and culture are available for type: CBS 244.71.

Apiosphaeria (Spegazzini) Höhn 1909 (= *Anisochora* Theiss. & Syd. 1915; = *Oswaldia* Rangel 1921 *fide* Species Fungorum 2017), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, four species, type: *A. guaranitica* (Speg.) Höhn., asexual morph unknown, terrestrial, South America, Lumbsch and Huhndorf (2010; genus accepted), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), culture is available for type: CCMF 766.

Apiosporina Höhn. 1910 (= *Dibotryon* Theiss. & Syd. 1915 *fide* Zhang et al. 2011d), *Venturiaceae*, *Venturiales*, *Dothideomycetes*, c. two species, type: *A. collinsii* (Schwein.) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Raja et al. (2011a; DNA), Zhang et al. (2011d; morphology, phylogeny), Hyde et al. (2013; accepted as in *Venturiaceae*, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Apiosporopsis (Traverso) Mariani 1911 (= *Guignardia* subgen. *Apiosporopsis* Traverso 1907; = *Sphaerognomonia* Potebnia ex Höhn. 1917 *fide* Species Fungorum 2017), *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, three

species, type: *A. saccardoana* Mariani (as ‘*saccardiana*’), asexual morph unknown, saprobes, endophytes, terrestrial, Dominican Republic, Europe, India, Japan, see Lumbsch and Huhndorf (2010; outline), Osono and Masuya (2012; diversity on Betulaceae), Kirk et al. (2013; genus accepted), Keča et al. (2015; diversity on Poplars), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available for *Apiosporopsis carpinea*.

Apiothyrium Petr. 1947, *Hyponectriaceae*, *Amphisphaeriales*, *Sordariomycetes*, two species, type: *A. arcticum* Petr., asexual morph unknown, saprobes, terrestrial, Finland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Apiotypa Petr. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *A. philippinensis* Petr., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, needs generic revision as Kirk et al. (2008) mentioned that type material is missing.

Aplanocalenia Lücking, Sérus. & Vězda 2005, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *A. inconspicua* (Müll. Arg.) Lücking, Sérus. & Vězda, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Apłosporella Speg. 1880 (= *Apłosporella* subgen. *Apłosporella* Speg. 1880; = *Apłosporella* subgen. *Euapłosporella* Petr. & Syd. 1926; = *Apłosporella* subgen. *Pleosphaeropsis* (Died.) Petr. & Syd. 1880; = *Epicyta* Syd. 1926; = *Microhapłosporella* Sousa da Câmara 1949; = *Pleosphaeropsis* Died. 1916; = *Podosporium* Bonord. 1851; = *Podosporium* Sacc. & Schulzer 1884 *fide* Species Fungorum 2017), *Apłosporellaceae*, *Botryosphaeriales*, *Dothideomycetes*, c. 70 species, type: *A. chlorostroma* Speg., asexual morph unknown coelomycetous, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c, 2016b, 2017; outline, taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Kirk et al. (2013; genus accepted), Slippers et al. (2013; phylogeny), Trakunyingcharoen et al. (2015; new species, phylogeny), Crous et al. (2016a; new species, phylogeny), Ekanayaka et al. (2016b; new species), cultures and sequences are available.

Apodospora Cain & J.H. Mirza 1970, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, four species, type: *A. simulans* Cain & J.H. Mirza, asexual morph unknown, coprophilous, North America, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al.

(2015b, 2016b; outline), culture and sequences are available.

Apodothina Petr. 1970, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. pringlei* (Peck) Petr., asexual morph unknown, saprobes, from living leaves, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Apodus Malloch & Cain 1971, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, two species, type: *A. deciduus* Malloch & Cain, asexual morph unknown, coprophilous, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 506.70.

Apogaeumannomyces Matsush. 2003, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. perplexus* Matsush., asexual morph unknown, saprobes, from living leaves, terrestrial, Peru, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: MFC 1P446, cultures and sequences are unavailable.

Apogloeum Petr. 1954, *Ascomycota* genera *incertae sedis*, one species, type: *A. concinnum* Petr., coelomycetous, sexual morph unknown, rarely reported and in case on leaves of *Lomatia polymorpha* in Tasmania, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype: BPI 1110296.

Apoharknessia Crous & S.J. Lee 2004, *Apoharknessiaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. insueta* (B. Sutton) Crous & S.J. Lee, sexual morph unknown, pathogens, Mauritius, Brazil, Colombia, see Crous et al. (2012f; phylogeny), Wijayawardene et al. (2012, 2016b, 2017; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline) epitype and ex-epitype culture of type: herb. CBS 9913, CBS 111377.

Apomelasmia Grove 1937 (= *Cheilarlopsis* Petr. 1959 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, eight species, type: *A. urticae* (Fr.) Grove, coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Wijayawardene et al. (2012), Maharachchikumbura et al. (2015b, 2016b) treated as a genus in *Diaporthales*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Aporellula B. Sutton 1986 [1985] (= *Aporella* Syd. 1939 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *A. erigerontis* (Syd.) B. Sutton, coelomycetous, sexual morph unknown, saprobes, Ecuador, see Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, isotype: IMI 288252.

Aporomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eleven species, type: *A. uniflagellatus* Thaxt., asexual morph unknown, parasitic on Limnichidae, Histeridae, and Staphylinidae beetles, terrestrial, Asia, Europe, North and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossi and Santamaria (2015; Ecuador), cultures and sequences are unavailable.

Aposphaeria Sacc. 1880, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, 189 species, type: *A. pulviscula* (Sacc.) Sacc., coelomycetous, sexual morph ?*Melanomma* Nitschke ex Fuckel, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Tian et al. (2015; morphology, phylogeny), Li et al. (2016a; new species), Tibpromma et al. (2017b; sexual report), cultures and sequences are available.

Aposporella Thaxt. 1920, *Ascomycota* genera *incertae sedis*, three species, type: *A. elegans* Thaxt., hyphomycetous, sexual morph unknown, on insects, terrestrial, Africa, see Seifert et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Apostrasseria Nag Raj 1983, *Ascomycota* genera *incertae sedis*, two species, type: *A. lunata* (Shear) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, New Zealand, North America, see Wijayawardene et al. (2012) treated as a genus in *Phaciidaeae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Appendichordella R.G. Johnson, E.B.G. Jones & S.T. Moss 1987, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *A. amicta* (Kohlm.) R.G. Johnson, E.B.G. Jones & S.T. Moss, asexual morph unknown, saprobes, marine, Denmark, France, Germany, Yugoslavia, USA, see Jones et al. (2009b, 2015; outline, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Appendicospora K.D. Hyde 1995, *Apiosporaceae*, *Amphisphaerales*, *Sordariomycetes*, two species, type: *A. coryphae* (Rehm) K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanan et al. (2017a; phylogeny, evolution), sequences are available.

Appendiculella Höhn. 1919 (= *Irenina* F. Stevens 1927; = *Meliola* subgen. *Irenina* (F. Stevens) Cif. 1938 *fide*

Species *Fungorum* (2017), *Meliolaceae*, *Meliolales*, *Sordariomycetes*, c. 110 species, type: *A. calostroma* (Desm.) Höhn., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type species.

Appendispora K.D. Hyde 1994, *Roussellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. frondicola* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Brunei, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Didymosphaeriaceae*), Hyde et al. (2013; accepted as a genus in *Didymosphaeriaceae*), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014a; transferred to *Roussellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Appendixia B.S. Lu & K.D. Hyde 2000, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, five species, type: *A. closterium* (Berk. & M.A. Curtis) B.S. Lu & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; placed in *Xylariales*), cultures and sequences are unavailable.

Apseudocercospora Videira & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *A. trigonotidis* Videira, H.D. Shin & Crous, hyphomycetous, sexual morph unknown, plant pathogenic, see Videira et al. (2016; morphology, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: KUS-F20054, CBS 131890 =CPC 10864.

Aptrootia Lücking & Sipman 2007, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, three species, type: *A. terricola* Lücking, Umaña & Sipman, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Trypetheliaceae*), Aptroot and Lücking (2016; monograph), Lücking et al. (2016a, 2017a; classification), sequences are available.

Aquadiscula Shearer & J.L. Crane 1985, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *A. appendiculata* Shearer & J.L. Crane, asexual morph unknown, saprobes on submerged leaf, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Aquadulciospora Fallah & Shearer 2001, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. rhomboidia* Fallah & Shearer, asexual morph unknown, saprobes, aquatic, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b;

outline), holotype: ILLS 53999, cultures and sequences are unavailable.

Aqualignicola Ranghoo, K.M. Tsui & K.D. Hyde 2001, *Annulatascaceae*, *Annulatascales*, *Sordariomycetes*, two species, type: *A. hyalina* Ranghoo, K.M. Tsui & K.D. Hyde, asexual morph unknown, saprobes, fresh water, terrestrial, China, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Hu et al. (2012a; new species, China), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: HKU (M) 12178, cultures and sequences are unavailable.

Aquamarina Kohlm., Volk.-Kohlm. & O.E. Erikss. 1996, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. speciosa* Kohlm., Volk.-Kohlm. & O.E. Erikss., asexual morph unknown, saprobes, marine, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Aquanectria L. Lombard & Crous 2015, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *A. penicilliodes* (Ingold) L. Lombard & Crous, asexual morph hyphomycetous, saprobes, aquatic, Jamaica, USA, see Lombard et al. (2015b; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type of type: CBS 257.54.

Aquapeziza D.M. Hu, L. Cai & K.D. Hyde 2012, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *A. globispora* D.M. Hu, L. Cai & K.D. Hyde, asexual morph unknown, saprobes, aquatic, China, see Hu et al. (2012c; taxonomy, phylogeny), holotype and cultures available for the type: IFRD 8978, DMH-2010.

Aquaphila Goh, K.D. Hyde & W.H. Ho 1998, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, two species, type: *A. albicans* Goh, K.D. Hyde & W.H. Ho, sexual morph tubeufia-like, saprobes, submerged wood in aquatic habitats, see Boonmee et al. (2011, 2014b; accepted as *Tubeufiaceae*), Seifert et al. (2011; morphology), Wijayawardene et al. (2014c; outline), Hyde et al. (2016; new species, sexual morph), cultures and sequences are available for authentic strains BCC 3520, BCC 3543 and reference strains MFLUCC16-0010, MFLUCC16-0020.

Aquapoterium Raja & Shearer 2008, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. pinicola* Raja & Shearer, saprobes, aquatic, see Koukol and Kolárová (2010; demonstrated that *Bactrodosmium gabretiae* Koukol & Kolárová, which was isolated from spruce needles is closely related to *A. pinicola* Raja & Shearer, in the *Helotiales*), Lumbsch and Huhndorf (2010; outline), ex-type strain F47-1 =MYA 4213 (American Type culture Collection), holotype: F47-1, ILL 40117

Aquapteridospora J. Yang, K.D. Hyde & Maharachch. 2015, *Diaporthomycetidae* genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. lignicola* Yang

Jing, K.D. Hyde & Maharachch., hyphomycetous, sexual morph unknown, saprobes, aquatic, Thailand, see Yang et al. (2015a; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: MFLU 15-1172.

Aquasphaeria K.D. Hyde 1995, *Sordariomycetes*, genera *incertae sedis*, one species, type: *A. dimorphospora* K.D. Hyde, asexual morph unknown, saprobes, aquatic, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: BRIP 21482, cultures and sequences are unavailable.

Aquastroma Kaz. Tanaka & K. Hiray. 2015, *Parabambusicolaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. magniostiolata* Kaz. Tanaka & K. Hiray., asexual morph unknown, saprobes, aquatic, Japan, see Tanaka et al. (2015; taxonomy, phylogeny), holotype and ex-type of the type: HHUF 30122, CBS 139680.

Aquasubmersa K.D. Hyde & Huang Zhang 2012, *Aquasubmersaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. mircensis* Huang Zhang & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, aquatic, Asia, see Wijayawardene et al. (2012, 2017; outline), Zhang et al. (2012a; taxonomy, phylogeny), Ariyawansa et al. (2015a; new species), Hashimoto et al. (2017; phylogeny), cultures and sequences are available.

Aquaticola W.H. Ho, K.M. Tsui, Hodgkiss & K. D. Hyde 1999, *Annulatascaceae*, *Incertae sedis*, *Sordariomycetes*, six species, type: *A. triseptata* K.M. Tsui, Hodgkiss & K.D. Hyde, asexual morph unknown, saprobes, aquatic, Hong Kong, Australia, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: HKU(M) 2969.

Aquilomyces D.G. Knapp, Kovács, J.Z. Groenew. & Crous 2015, *Morosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. patris* D.G. Knapp, Kovács, Groenewald & Crous, endophytes, see Knapp et al. (2015; phylogeny), cultures and sequences are available.

Arachniotus J. Schröt. 1893 (= *Disarticulatus* G.F. Orr 1977; = *Petalosporus* G.R. Ghosh, G.F. Orr & Kuehn 1963; = *Plunkettomyces* G.F. Orr 1977; = *Rollandina* Pat. 1905; = *Waldemaria* Bat., H. Maia & Cavalc. 1960 *fide* Species Fungorum 2017), *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, three species, type: *A. ruber* (Tiegh.) J. Schröt., asexual morph unknown, saprobes, from soil, Poland, see Lumbsch and Huhndorf (2010; outline), Ropars et al. (2012; DNA), Kirk et al. (2013; genus accepted), cultures available for the type: MUCL 39748.

Arachnocrea Z. Moravec 1956, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *A. stipata* (Fuckel) Z. Moravec, asexual morphs

?verticillium-like, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Arachnomyces Massee & E.S. Salmon 1902, *Arachnomycetaceae*, *Onygenales*, *Eurotiomycetes*, ten species, type: *A. nitidus* Massee & E.S. Salmon, asexual morph unknown, saprobes, Europe, see Gueidan et al. (2008; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence available for the type: AB075351.

Arachnopeziza Fuckel 1870 (= *Arachnopezizella* Kirschst. 1938; = *Arachnoscypha* Boud. 1885 *fide* Species Fungorum 2017), *Arachnopezizaceae*, *Helotiales*, *Leotiomycetes*, c. 15 species, type: *A. aurelia* (Pers.) Fuckel., asexual morph hyphomycetous, saprobes worldwide, common in temperate and subtropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; phylogeny), type in L herbarium, cultures and sequence are available but lacking for the type species.

Arachnophora Hennebert 1963 (= *Cacahualia* Mercado & R.F. Castañeda 1984; = *Sopagraha* Subram. & Sudha 1979 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, three species, type: *A. fagicola* Hennebert, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Pratibha et al. (2011; new species), Seifert et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Arachnoscypha Boud. 1885, *Arachnopezizaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *A. aranea* (De Not.) Boud., sexual morph unknown, saprobes, UK, see Jaklitsch et al. (2016a; classification), Index Fungorum (2017; treated as a synonym of *Arachnopeziza* Fuckel), cultures and sequences are unavailable.

Arachnospora R.F. Castañeda, Minter & Camino 2003, *Ascomycota* genera *incertae sedis*, one species, type: *A. insolita* R.F. Castañeda, Minter & Camino, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: MUCL 42859, cultures and sequences are unavailable, compare with *Arachnophora* and *Iyengarina*.

Arachnotheca Arx 1971, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, three species, type: *A. glomerata* (E. Müll. & Pacha-Aue) Arx, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Sigler et al. (2013; DNA), culture available for the type: UAMH 3551.

Araneomyces Höhn. 1909 (= *Paranectriella* (Henn. ex Sacc. & D. Sacc.) Höhn. *fide* Wijayawardene et al. 2014c;

Rossman et al. 2015b), *Paranectriellaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *A. acarifer* Höhn., sexual morph formerly known in *Paranectriella*, South America, see Seifert et al. (2011; taxonomy), Wijayawardene et al. (2012, 2014c, 2017; outline, nomenclature), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), cultures and sequences are unavailable.

Arborillus Munt.-Cvetk. & Gómez-Bolea 1998, *Ascomycota* genera *incertae sedis*, one species, type: *A. llimonae* Munt.-Cvetk. & Gómez-Bolea, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Arborispora K. Ando., 1986, *Ascomycota* genera *incertae sedis*, four species, type: *A. palma* K. Ando, in rainwater draining from leaves of *Quercus*: Honshu, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sexual morph unknown, cultures and sequences are unavailable.

Arbusculina Marvanová & Descals 1987, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *A. irregularis* (R.H. Petersen) Marvanová & Descals, asexual morph unknown, saprobes, aquatic, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Funck et al. (2015; DNA), cultures and sequences are available but lacking for the type species.

Archaeorhizomyces Rosling & T.Y. James 2011, *Archaeorhizomycetaceae*, *Archaeorhizomycetales*, *Archaeorhizomycetes*, two species (but many others are predicted from the analysis of environmental sequences), type: *A. finlayi* Rosling & T.Y. James, asexual morph unknown, saprobes, from soil, terrestrial, Sweden, Lithuania, see Rosling et al. (2011; taxonomy, phylogeny), Menkis et al. (2014; new species, Lithuania, phylogeny), holotype and ex-type cultures: MICH 72671, CBS 128710.

Architrypethelium Aptroot 1991, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, seven species, type: *A. seminudum* (Mont.) Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Aptroot and Lücking (2016; monograph), Lücking et al. (2017a; classification), sequences are available.

Arctocetraria Kärnefelt & Thell 1993, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *A. andrejevii* (Oxner) Kärnefelt & A. Thell, lichenized, terrestrial (corticicolous or terricolous), Northern

Hemisphere (arctic), see Thell et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2011a; DNA, phylogeny), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; DNA, phylogeny), Divakar et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Arctomia Th. Fr. 1860, *Arctomiaceae*, *Arcomiales*, *Lecanoromycetes*, ten species, type: *A. delicatula* Th. Fr. lichenized (cyanobacteria), terrestrial (saxicolous, terricolous or corticolous), cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Magain and Sérusiaux (2012; new species), Kirk et al. (2013; genus accepted), Otálora and Wedin (2013; DNA, phylogeny), Miądlikowska et al. (2014a; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Arctomollisia Raity. 2008, *Dermateaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *A. oxyparaphysata* (Rehm) Raity., asexual morph unknown, saprobes, terrestrial, Magadan, see Raityir (2008; taxonomy), cultures and sequences are unavailable.

Arctoparmelia Hale 1986, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *A. centrifuga* (L.) Hale, lichenized (green algae), terrestrial (mainly saxicolous), arctic and alpine regions of Northern Hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; DNA, phylogeny), Divakar et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Arcuadendron Sigler & J.W. Carmich. 1976, *Ascomycota* genera *incertae sedis*, three species, type: *A. ovatum* Sigler & J.W. Carmich, hyphomycetous, sexual morph unknown, soil, India, former Yugoslavia, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), ex-type strain of type: ATCC 34519, CBS 122.77, sequences are unavailable.

Arecacicola Joanne E. Taylor, J. Fröhl. & K.D. Hyde 2001, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. calami* Joanne E. Taylor, J. Fröhl. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: HKU (M) 1126a, cultures and sequences are unavailable.

Arecomyces K.D. Hyde 1996, *Hypocretriaceae*, *Amphisphaeriales*, *Sordariomycetes*, eleven species, type: *A. frondicola* K.D. Hyde, ascomycetous, asexual morph unknown, saprobes, reported only from palms, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Vitoria et al. (2011, new species), Kirk et al. (2013; genus

accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Arecophila K.D. Hyde 1996, *Cainiaceae*, *Xylariales*, *Sordariomycetes*, 15 species, type: *A. gulubiicola* K.D. Hyde, asexual morph unknown, saprobes, South East Asia and Papua New Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lee et al. (2013a; sesquiterpenes from *A. saccharicola*), Maharachchikumbura et al. (2015b, 2016b; outline), sequence data available for two species but lacking for the type.

Arenariomyces Höhnk 1954, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, five species, type: *A. trifurcatus* Höhnk, aquatic, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Areolospora S.C. Jong & E.E. Davis 1974, *Sordariomycetes* genera *incertae sedis*, two species, type: *A. terrophila* S.C. Jong & E.E. Davis, asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Sordariomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Argentinomyces N.I. Peña & Aramb. 1997, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. naviculisperus* Peña & Aramb., asexual morph unknown, saprobes, marine, Argentina, Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Ascomycota* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: LPS 45469.

Argopericonia B. Sutton & Pascoe 1987, *Ascomycota* genera *incertae sedis*, two species, type: *A. elegans* B. Sutton & Pascoe, hyphomycetous, sexual morph unknown, saprobes, saprophytes, Australia, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need further revisions, holotype: VPRI13846.

Argopsis Th.Fr. 1857, *Lecanoromycetes* genera *incertae sedis*, three species, type: *A. megalospora* Th. Fr., lichenized, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Brigantiaeaceae*), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Argynna Morgan 1895, *Argynnaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *A. polyhedron* (Schwein.) Morgan, asexual morph unknown, saprobes,

terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Argynnaceae*, taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Aristostoma Tehon 1933, *Ascomycota* genera *incertae sedis*, one species, type: *A. concentrica* Tehon, coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Zhu et al. (2012; DNA, pathogens), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Arkoola J. Walker & Stovold 1986, *Dothideomycetes* genera *incertae sedis*, one species, type: *A. nigra* J. Walker & Stovold, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Zhang et al. (2011d; provisionally accepted as a genus in *Venturiaceae*), Hyde et al. (2013; did not accept as *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Armata W. Yamam. 1958, *Dothideomycetes* genera *incertae sedis*, one species, type: *A. macrospora* (W. Yamam.) W. Yamam., asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Micropeltidaceae*), Hyde et al. (2013; did not accept as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Arnaudiella Petr. 1927 (= *Seynesiola* Speg. 1918 *fide* Wu et al. 2011), *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, 13 species, type: *A. caronae* (Pass.) Petr., asexual morph hyphomycetous, saprophyte, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Armatella Theiss. & Syd. 1915 (= *Artallendea* Bat. & H. Maia 1960 *fide* Species Fungorum 2017), *Armatellaceae*, *Meliolales*, *Sordariomycetes*, 20 species, type: *A. litseae* (Henn.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, worldwide distribution, see Hosagoudar and Archana (2009; distribution on Lauraceae), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hosagoudar et al. (2013; new species), Hongsanan et al. (2015d, 2016c; family placement, evolution), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Arnium Nitschke ex G. Winter 1873, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, twelve species, type: *A. lanuginosum* Nitschke, asexual morph unknown, saprobes and coprophilous, worldwide, see

Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Aropsiclus Kohlm. & Volk.-Kohlm. 1994 (= *Sulcospora* Kohlm. & Volk.-Kohlm. 1993 *fide* Species Fungorum 2017), *Sordariomycetes*, genera *incertae sedis*, one species, type: *A. junci* (Kohlm. & Volk.-Kohlm.) Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Arpinia Berthet 1974, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, four species, type: *A. inops* Berthet, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Arrhenosphaera Stejskal 1974, *Ascospaeraceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *A. craneae* Stejskal, asexual morph unknown, from bee hive, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), sequences and culture available.

Arthonia Ach. 1806 (see Species Fungorum 2017 for synonyms), *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, c. 500 species, type: *A. radiata* (Pers.) Ach. 1808, sexual morph with mainly apothecoid ascomata, asexual morph coelomycetous, lichenized, lichenicolous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010, 2014; outline), Ertz and Tehler (2011; phylogeny), Kirk et al. (2013; genus accepted), Frisch et al. (2014; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Arthophaeopsis Hafellner 1998, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *A. parmeliarum* Hafellner, lichenicolous, terrestrial, North America, Europe (including Canary Islands), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype: Hafellner 41023 (GZU), cultures and sequences are unavailable.

Arthopyrenia A. Massal. [nom. cons.] 1852, *Arthopyreniaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *A. cerasi* (Schrad.) A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Arthopyreniaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Lücking et al. (2017a; classification), cultures and sequences are available.

Arthotheliopsis Vain. 1896, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *A. hymenocarpoides*

Vain., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Arthothelium A. Massal. 1852, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, c. 130 species, type: *A. spectabile* Flot. ex A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), polyphyletic, sequences are available.

Arthrinium Kunze 1817 (see Species Fungorum 2017 for synonyms), *Apiosporaceae*, *Amphisphaeriales*, *Sordariomycetes*, 64 species, type: *A. caricicola* Kunze & J.C. Schmidt, saprobes, pathogenic, endophytic, terrestrial, widely distribute, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Crous and Groenewald (2013; phylogeny), Crous and Groenewald (2013; asexual morph, morphology, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; morphology, phylogeny), Dai et al. (2016, 2017; new species, sexual and asexual morph, morphology, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are unavailable for type species.

Arthrobotrys Corda 1839 (= *Dactylariopsis* Mekht. 1967; = *Dactylium* Nees 1816 [1816-17]; = *Didymozoophaga* Soprakov & Galiulina 1951; = *Duddingtonia* R.C. Cooke 1969; = *Monacrosporium* Oudem. 1885; = *Orbiliella* Kirschst. 1938; = *Woroninula* Mekht. 1979 *fide* Species Fungorum 2017; see Baral et al. 2015 for additional synonyms), *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, 112 species, type: *A. superba* Corda, sexual morph unknown, nematophagous, terrestrial, marine, worldwide see Han et al. (2008; extrolite profiling), Li et al. (2008c, 2009a; taxonomic revision, new species), Swe et al. (2008; new species, marine), Wang et al. (2008b, 2013a, e; amino acid effect, mutant predatory activity, immunological effect of nanoparticles), Braga et al. (2009, 2010, 2014; predatory activity), Ranjbar-Bahadori et al. (2010; in vivo study), Silva et al. (2010; control sheep gastroenteritis), Zhang et al. (2011f, 2012b, 2013e; phylogeny, metabolites study, genetic diversity), Carvalho et al. (2011; predatory activity after freeze dry), Niu and Zhang (2011; review), Su et al. (2011a; new species), Wei et al. (2011; Arthrobotrisins, chitinase encoding gene analysis), Yang et al. (2011b, 2013; genomics, proteomics), Freire et al. (2012; effect of Fusarium extrolites), Qiao et al. (2012; description), Singh et al. (2012, 2013c; role in plant immunity, biocontrol potential), Wu et al. (2013a; new species, metabolite profiling), Arias et al. (2013; developmental biology), Chen et al. (2013c; trap formation), Falbo et al. (2013; characterization),

Jamshidnejad et al. (2013; biocontrol activity), Liang et al. (2013, 2015; cell signalling, proteomic and transcriptional analyses), Wang et al. (2013a; nematophagous), Zhang et al. (2013e; nematode trapping), Braga et al. (2014; predatory activity after preservation), Liu et al. (2014c; new species), Mostafanezhad et al. (2014; nematode control), Pandit et al. (2014; protease study), Zhao et al. (2014b; malate synthase gene), Baral et al. (2015; nomenclature), sequences are available for few species, including type: KT215210.

Arthrobotryum Ces. 1854 (= *Arthrographium* Ces. 1854; = *Lindauomyces* Koord. 1907; = *Wettsteinella* Kuntze 1891 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, five species, type: *A. stilboideum* Ces., coelomycetous, sexual morph unknown, widespread, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Arthromoniliphora S.S. Silva, Gusmão & R.F. Castañeda, *Ascomycota* genera *incertae sedis*, one species, type: *A. araucariae* S.S. Silva, Gusmão & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Silva et al. (2016; taxonomy), cultures and sequences are unavailable, compare with *Arthrographis*, *Geomycetes*, *Geotrichum*, *Thedgonia*.

Arthrocataena Egidi & Selbmann 2014, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *A. tenebrio* Egidi & Selbmann, rock-inhabiting fungus, conidia resulting from arthric secession, sexual morph unknown, see Egidi et al. (2014; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), ex-type cultures available for the type: CBS 136100 = CCFEE 5413.

Arthrocladiella Vassilkov 1960, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, one species, type species: *A. mougeotii* (Lév.) Vassilkov (≡ *Microsphaera mougeotii* Lév.), asexual morph hyphomycetous, obligate plant pathogen, terrestrial, almost worldwide, see Braun and Cook (2012; taxonomy), lectotype: France, Vaugirard, on *Lycium europaeum*, ex herb. Léveillé (PC), sequences available.

Arthrocladium Papendorf 1969, *Trichocomiaceae*, *Chaetothyriales*, *Eurotiomycetes*, four species, type: *A. caudatum* Papendorf, asexual morph hyphomycetous, saprobes, terrestrial, Africa and Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Nascimento et al. (2016; phylogeny, human opportunist), ex-type strains of type: CBS 457.67.

Arthrocristula Sigler, M.T. Dunn & J.W. Carmich. 1982, *Ascomycota* genera *incertae sedis*, one species, type: *A. hyphenata* Sigler, M.T. Dunn & J.W. Carmich.,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Arthroderma Curr. 1860 (= *Nannizzia* Stockdale 1961 *fide* Species Fungorum 2017), *Arthrodermataceae*, *Onygenales*, *Eurotiomycetes*, 41 species, type: *A. curreyi* Berk, asexual morph unknown, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), sequences and culture are available for type: NCPF 0797.

Arthrographis G. Cochet ex Sigler & J.W. Carmich. 1976, *Eremomycetaceae*, *Dothideomycetes* families *incertae sedis*, twelve species, type: *A. kalrae* (R.P. Tewari & Macph.) Sigler & J.W. Carmich., hyphomycetous, sexual morph unknown, saprobes, on human, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Giraldo et al. (2014a; phylogeny, new species), cultures available for type: CBS 693.77.

Arthropsis Sigler, M.T. Dunn & J.W. Carmich. 1982, *Onygenales*, *Eurotiomycetes*, four species, type: *A. truncata* Sigler, M.T. Dunn & J.W. Carmich., hyphomycetous, sexual morph unknown, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, need generic revision.

Arthrorhaphis Th. Fr. [nom. cons.] 1860 (= *Gongylia* Körb. 1855; = *Mycobacidia* Rehm 1890 [1896]; = *Parathalle* Clem. 1909; = *Raphiospora* A. Massal. 1853 *fide* Species Fungorum 2017), *Arthrorhaphidaceae*, order *incertae sedis*, *Lecanoromycetes*, 13 species, type: *A. flavovirescens* (A. Massal.) Th. Fr.; lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Arthrorhynchus Kolen. 1857 (= *Helminthophhana* Peyr. 1873 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, one additional species (*A. acrandros* Merola) is doubtful, type: *A. nycteribiae* (Peyr.) Thaxt., asexual morph unknown, biotrophic, hosts are Nycteriidae (Diptera: Hippoboscidae), Eastern Hemisphere distribution, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted) Haelewaters et al. (2017a; review, DNA), sequences are available for type species.

Arthrosporium Sacc. 1880 (= *Phragmostilbe* Subram. 1959 *fide* Seifert et al. 2011), *Ascomycota* genera *incertae sedis*, two species, type: *A. albicans* (Sacc.) Sacc., hyphomycetous, sexual morph unknown, Europe, North

America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Arthrosporum A. Massal. 1853 (= *Arthrospora* Th. Fr. 1861 *fide* Species Fungorum 2017), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *A. populorum* Massal., with ascomata, lichenized (green algae), terrestrial (corticolous), Northern Hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Arthrowallemia R.F. Castañeda, Dania García & Guarro 1998, *Ascomycota* genera *incertae sedis*, two species, type: *A. formosa* R.F. Castañeda, Dania García & Guarro, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: INIFAT C95/88, sequences are unavailable, compare with *Arthrocristala*, *Bahusakala* and *Parasymodiella*.

Arthroxylaria Seifert & W. Gams 2002, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *A. elegans* Seifert & W. Gams, hyphomycetous, sexual morph unknown, terrestrial, saprobes, North America, see Seifert et al. (2011; morphology), Wendt et al. (2017; placed in *Hypoxylaceae*), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Articulophora C.J.K. Wang & B. Sutton 1982, *Ascomycota* genera *incertae sedis*, one species, type: *A. variispora* C.J.K. Wang & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype: Wang 11021.

Articulospora Ingold 1942, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *A. tetracladia* Ingold, hyphomycetous, saprobes, endophytic, aquatic, worldwide, see Seifert et al. (2011; morphology), Seena et al. (2012; phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Baschien et al. (2013; phylogeny), compare with *Geniculospora* and *Isthmotrichiacladia*.

Artocarpomyces Subram. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *A. paradoxa* Subram., saprobes, terrestrial, Singapore, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: MUBL 3147, cultures and sequences are unavailable.

Arundellina Wanas., E.B.G. Jones & K.D. Hyde 2016, *Lindgomycetaceae*, *Pleosporales*, *Dothideomycetes*, one

species, type: *A. typhae* Wanas., E.B.G. Jones & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, UK, see Hyde et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 16-1276, MFLUCC 16-0310.

Arwidssonia B. Erikss. 1974, *Hyponectriaceae*, *Amphisphaeriales*, *Sordariomycetes*, two species, type: *A. empetri* (Rehm) B. Erikss., asexual morph unknown, plant pathogen, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Arxiella Papendorf 1967, *Dothideomycetes*, genera *incertae sedis*, one species, type: *A. terrestris* Papendorf, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014d; DNA, new species), cultures and sequences are available but lacking for the type.

Arxiomyces P.F. Cannon & D. Hawksw. 1983 (= *Ceratostoma* Pers. 1818; = *Phaeostoma* Arx & E. Müll. 1954 *fide Species Fungorum* 2017), *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *A. vitis* (Fuckel) P.F. Cannon & D. Hawksw., asexual morph unknown, on bark, Europe, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Asahinea W.L. Culb. & C.F. Culb. 1965, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, four species; type: *A. chrysantha* (Tuck.) W.L. Culb. & C.F. Culb., lichenized, terrestrial (saxicolous or on plant debris), arctic-montane, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Asaphomyces Thaxt. 1931 (= *Barbariella* Middelh. 1949 *fide Species Fungorum* 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *A. cholevae* Thaxt., asexual morph unknown, parasitic on Leiodidae beetles, terrestrial, widespread, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascitendus J. Campb. & Shearer 2004, *Annulatascae*, *Annulatascales*, *Sordariomycetes*, one species, type: *A. austriacus* (Rébllová, Winka & Jaklitsch) J. Campb. & Shearer, asexual morph unknown, saprobes, fresh water, terrestrial, Austria, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: PRM 842991, cultures available for the type: CBS 131685.

Ascluella DiCosmo, Nag Raj & W.B. Kendr. 1983, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. symplocina* (P. Syd.) DiCosmo, Nag Raj & W.B. Kendr., asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascobolus Pers. 1792 (= *Anserina* Velen. 1934; = *Ascobolus* sect. *Sphaeridiobolus* (Boud.) Brumm. 1967; = *Ascobolus* subgen. *Dasyobolus* Sacc. 1889; = *Crouaniella* (Sacc.) Lambotte 1887 [1888]; = *Dasyobolus* (Sacc.) Sacc. 1895; = *Phaeopezia* subgen. *Crouaniella* Sacc. 1884; = *Selianniella* Arx & E. Müll. 1955; = *Sphaeridiobolus* Boud. 1885 *fide Species Fungorum* 2017), *Ascobolaceae*, *Pezizales*, *Pezizomycetes*, c. 70 species, type: *A. pezizoides* Pers., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Doveri (2011, 2014; worldwide key), Mungai et al. (2012e; from dung), Aluoch et al. (2015; fungi on wildlife dung), cultures and sequences are available but lacking for the type species.

Ascocalathium Eidam ex J. Schröt. (1893) [1908], *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *A. stipitatum* Eidam ex J. Schröt., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, treated as *Pyronemataceae*), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Ascocalvatia Malloch & Cain 1971, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *A. alveolata* Malloch & Cain, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), culture available for the type: ATCC 22147.

Ascocalyx Naumov 1926 (= *Bothrodiscus* Shear 1907; = *Lagerbergia* J. Reid 1971 *fide* Johnston et al. 2014b; *Species Fungorum* 2017), *Godroniaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *A. abietis* Naumov, asexual morph *Bothrodiscus*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available.

Ascocephalophora K. Matsush. & Matsush. 1995, *Saccharomycetales* genera *incertae sedis*, *Saccharomycetes*, one species, type: *A. petasiformis* K. Matsush. & Matsush., Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascochalara Rébllová 1999, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type:

A. gabretiae Réblová, asexual morph chalara-like, saprobes, Czech Republic, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequence available but lack for the type species, generic placement is based on non-type species.

Ascochyta Lib. 1830 (= *Ascochyta* subgen. *Ascochytyula* Potebnia 1907; = *Ascochytyella* Tassi 1902; = *Ascochytyula* (Potebnia) Died. 1912; = *Macrodiplodina* Petr. 1962 [1961]; = *Stagonosporopsis* Died. 1912 *fide* Species Fungorum 2017), *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, c. 400 species, type: *A. pisi* Lib., coelomycetous, saprobes, pathogens, terrestrial, worldwide, see Chilvers et al. (2009; pathogens), Aveskamp et al. (2010; phylogeny), de Gruyter et al. (2009; phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), Kirk et al. (2013; genus accepted), Chen et al. (2015b; taxonomy, phylogeny), Tibpromma et al. (2017b; new species), cultures and sequences are available, needs generic revision as more than 1000 records are available.

Ascochytopsis Henn. 1905 (= *Cryptogene* Syd. 1939; = *Cryptogenella* Syd. 1939; = *Scleroparodia* Petr. 1934 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, five species, type: *A. vignae* Henn., coelomycetous, saprobes, endophytic, pathogenic, terrestrial and aquatic, cosmopolitan, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascochytulina Petr. 1922 (= *Clypeodiplodina* F. Stevens 1927 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, three species, type: *A. deflectens* (P. Karst.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascoclavulina Otani 1974, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, eight species, type: *A. sakaii* Y. Otani, asexual morph gliomastix-like, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascodiodinaea Samuels, Cand. & Magni 1997, *Glomerellales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *A. stereicola* Samuels, Cand. & Magni, asexual morph dictyochaeta-like, saprobes, on old polypore, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ascoconidium Seaver 1942 (= *Sageria* A. Funk 1975 *fide* Johnston et al. 2014b), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *A. castaneae* Seaver,

hyphomycetous, sexual morph formerly known in *Sageria*, saprobes, Asia, North America, see Seifert et al. (2011; morphology), Johnston et al. (2014b; nomenclature), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascocorticiellum Jülich & B. de Vries 1982, *Ascocorticiaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *A. vermisporum* (Hauerslev) Jülich & B. de Vries, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Ascocorticium Bref. 1891, *Ascocorticiaceae*, *Helotiales*, *Leotiomycetes*, eight species, type: *A. albidum* Bref. & Tavel, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Ascocoryne J.W. Groves & D.E. Wilson 1967 (= *Coryne* Nees 1816; = *Pirobasidium* Höhn. 1902; *Pleurocolla* Petr. 1924; = *Endostilbum* Malençon 1964 *fide* Johnston et al. 2014b), *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, eight species, type: *A. sarcooides* (Jacq.) J.W. Groves & D.E. Wilson, asexual morph *Monilia*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Johnston et al. (2014b; nomenclature), Zhu et al. (2016; pathogens, phylogeny), cultures and sequences are available.

Ascocratera Kohlm. 1986, *Aigialaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. manglicola* Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Schoch et al. (2009; phylogeny), Suetrong et al. (2009; phylogeny, morphology), Besitulo et al. (2010; diversity), Zainuddin et al. (2010; antimicrobial activities), Pang et al. (2010b; diversity), Sakayaroj et al. (2011b; diversity), Zhang et al. (2012e, 2016a; phylogeny), Boonmee et al. (2012; phylogeny), Jones et al. (2012; phylogeny, review), Lee et al. (2012; checklist), Ariyawansa et al. (2014a; phylogeny), Wanasinghe et al. (2015; phylogeny), holotype and ex-isotype of type: J.K. 4812a (FH), J.K.4812b (IMS).

Ascocylindrica Abdel-Wahab, Bahkali & E.B.G. Jones 2015, *Ascocylindricaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. marina* Abdel-Wahab, Bahkali & E.B.G. Jones, asexual morph unknown, saprobes, marine, Saudi Arabia, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype of the type: MFLU 15-0750

Ascodesmis Tiegh. 1876, *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, six species, type: *A. aurea* Tiegh., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Doveri (2011;

new record from Italy), Hansen et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for type species.

Ascodichaena Butin 1977 (= *Polymorphum* Chevall. 1822 *fide* Johnston et al. 2014b), *Ascodichaenaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *A. rugosa* Butin, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), culture available for type: DSM 4703.

Ascofascicula Matsush. 2003, *Ascomycota* genera *incertae sedis*, six species, type: *A. talaroluteoides* Matsush., asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available but lacking for type species.

Ascographa Velen. 1934, *Lecanoromycetes* genera *incertae sedis*, *Leotiomycetes*, one species, type: *A. hederae* Velen., asexual morph unknown, former Czechoslovakia, on decaying trunk of *Hederae helicis*, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Ascoidea Bref. 1891, *Ascoideaceae*, *Saccharomycetales*, *Saccharomycetes*, three species, type: *A. rubescens* Bref., asexual morph unknown (by budding of cells and formation of hyphae), saprophyte, terrestrial, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Ascolacicola Ranghoo & K.D. Hyde 1998, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. aquatica* Ranghoo & K.D. Hyde, asexual morph trichocladium-like, saprobes, aquatic, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Ascolectus Samuels & Rogerson 1990, *Saccardiaceae*, *Ascomycota* families *incertae sedis*, one species, type: *A. albus* Samuels & Rogerson, asexual morph unknown, saprobes, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascomauritiana Ranghoo & K.D. Hyde 1999, *Ascomycota* genera *incertae sedis*, one species, type: *A. lignicola* Ranghoo & K.D. Hyde, asexual morph unknown, saprobes, aquatic, Mauritius, see Kirk et al. (2013; genus accepted), holotype of type species HKU (M) 5255.

Ascominuta Ranghoo & K.D. Hyde 2000, *Dothideomycetes* genera *incertae sedis*, two species, type: *A. lignicola* Ranghoo & K.D. Hyde, asexual morph unknown, saprobes, aquatic, Thailand, Hong Kong, see

Wijayawardene et al. (2014c; outline), holotype: HKU (M) 5246.

Ascophanus Boud. 1869, *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, one species, type: *A. granuliformis* (P. Crouan & H. Crouan) Boud., asexual morph unknown, saprobes, terrestrial, Antarctica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Bansal et al. (2014; antifreeze proteins), cultures and sequences are available.

Ascoplyporus Möller 1901, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *A. polychrous* Möller, asexual morph unknown, saprobes, terrestrial, Central and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ascoporia Samuels & A.I. Romero 1993 (= *Pseudosolidum* Lloyd (as McGinty) 1923 *fide* Hyde et al. 2013), *Ascoporiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *A. lateritia* Samuels & A.I. Romero, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures are unavailable.

Ascorhiza Lecht.-Trinka 1931, *Sordariomycetes*, genera *incertae sedis*, one species, type: *A. leguminosarum* Lecht.-Trinka, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ascorhizoctonia Chin S. Yang & Korf 1985, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, seven species, type: *A. praecox* Chin S. Yang & Korf, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Ascorhombispora L. Cai & K.D. Hyde 2007, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *A. aquatica* L. Cai & K.D. Hyde, asexual morph unknown, China, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), holotype: HKU (M) 10859, cultures and sequences are available.

Ascoronospora Matsush. 2003, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *A. yakuensis* Matsush., asexual morph coronospora-like, saprobes, terrestrial, Japan, see Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Ascosacculus J. Campb., J.L. Anderson & Shearer 2003, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two

species, type: *A. aquaticus* (K.D. Hyde) J. Campb., J.L. Anderson & Shearer, saprobes, aquatic, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Ascisorus Henn. & Ruhland 1900 (= *Gymnoascopsis* C. Moreau & M. Moreau 1959 *fide* Species Fungorum 2017), *Ascocorticiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *A. floridanus* (Ellis & G. Martin) Henn. & Ruhland, asexual morph unknown, saprobes, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Ascosparsassis Kobayasi 1960, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *A. shimizuensis* Kobayasi, saprobes, terrestrial, tropics, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Ascospheara L.S. Olive & Spiltoir 1955 (= *Pericystis* Betts 1912 *fide* Species Fungorum 2017), *Ascospheeraceae*, *Onygenales*, *Eurotiomycetes*, c. 30 species, type: *A. apis* (Maassen ex Claussen) L.S. Olive & Spiltoir, pathogens, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences and culture available for type: NBRC 31608.

Ascostratum Syd. & P. Syd. 1912, *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, two species, type: *A. insigne* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, South Africa, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Tian et al. (2014; morphology, nomenclature, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Ascotaiwania Sivan. & H.S. Chang 1992, *Savoryllaceae*, *Savoryllales*, *Sordariomycetes*, 13 species, type: *A. lignicola* Sivan. & H.S. Chang, asexual morph hyphomycetous, saprobes, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Boonyuen et al. (2011; review, phylogeny), Maharachchikumbura et al. (2015b, 2016b), holotype: IMI 348977 sequences available.

Ascotremella Seaver 1930, *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *A. faginea* (Peck) Seaver, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ascotricha Berk. 1838 (= *Dicyma* Boulanger 1897; = *Gonytrichella* Emoto & Tubaki 1971; = *Puciola* De Bert. 1976 *fide* Wendt et al. 2017), *Xylariaceae*, *Xylariales*,

Sordariomycetes, one species, type: *A. chartarum* Berk., asexual morph unknown, saprobes, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Cheng et al. (2015; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ascotrichella Valldos. & Guarro 1988, *Xylariales*, genera *incertae sedis*, *Sordariomycetes*, one species, type: *A. hawksworthii* Valldos. & Guarro, asexual morph hyphomycetous, saprobes, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ascovaginomysora Fallah, Shearer & W.D. Chen 1997, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *A. stellipala* Fallah, Shearer & W.D. Chen, saprobes, aquatic, USA, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariomycetidae*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: ILLS 52319, sequences are available.

Ascoyunnania L. Cai & K.D. Hyde 2005, *Sordariomycetes* genera *incertae sedis*, two species, type: *A. aquatica* L. Cai & K.D. Hyde, saprobes, aquatic, China, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ascozonus (Renny) E.C. Hansen 1877 (= *Ascobolus* sect. *Ascozonus* Renny, Trans. 1873 [1871-1873]; = *Gymnodiscus* Zukal 1887; = *Streptotheca* Vuill. 1887; = *Zukalina* Kuntze 1891 *fide* Species Fungorum 2017), *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, c. six species, type: *A. cunicularius* (Boud.) Marchal, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacks for the type species.

Ashtaangam Subram. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *A. sundaram* Subram., hyphomycete, sexual morph unknown, saprobe, terrestrial, Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: MUBL 3127.

Aspergillus P. Micheli ex Haller 1768 (= *Chaetosartorya* Subram. 1972; = *Cristaspora* Fort & Guarro 1984; = *Dichotomomyces* Saito ex D.B. Scott 1970; = *Emericella* Berk. 1857; = *Eurotium* Link 1809; = *Fennellia* B.J. Wiley & E.G. Simmons 1973; = *Neocarpenteles* Udagawa & Uchiy. 2002; = *Neopetromyces* Frisvad & Samson 2000; = *Neosartorya* Malloch & Cain 1973; = *Petromyces* Malloch & Cain 1973; = *Phialosimplex* Sigler, Deanna A. Sutton, Gibas, Summerb. & Iwen 2010; = *Polypaecilum* G. Sm. 1961),

Aspergillaceae, *Eurotiales*, *Eurotiomycetes*, c. 360 species, type: *A. glaucus* (L.) Link, mostly terrestrial, but also reported from aquatic environments, mainly saprobes associated with soil, plant material and foodstuffs, but also including phytopathogenic and clinically-relevant fungi causing opportunistic mycoses (aspergillosis) and mycotoxicoses in various vertebrates including humans, cosmopolitan, monophyletic but morphologically diverse both in its sexual and asexual morphs, see Peterson (2008; phylogeny), Sigler et al. (2010b; morphology as “*Phialosimplex*”), Varga et al. (2010a, b, 2011; phylogeny, new species, mycotoxins), Houbraken and Samson (2011; phylogeny), Guarro et al. (2012; selected soil borne species), Hubka et al. (2013a, b, c, 2015, 2016a, 2016b; phylogeny, new combinations, new species), Asgari et al. (2014; new species, sexual morph), Samson et al. (2014; polyphasic taxonomy, new species, outline), Sugui et al. (2014; new species), Crous et al. (2015e; new species), Guinea et al. (2015; new species), Jurjević et al. (2015; new section and new species), Arzanlou et al. (2016b; new species), Tsang et al. (2016; new species), Visagie et al. (2017; new species), ex-neotype: CBS 516.65, sequences are available.

Asperisporium Maubl. 1913, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, twelve species, type: *A. caricae* (Speg.) Maubl., hyphomycetous, saprobes, terrestrial, America, see Seifert et al. (2011; morphology), Minnis et al. (2011; phylogeny, typification), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), ex-epitype strains of type: CBS 130298.

Asperopilum Spooner 1987, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *A. juncicola* (Dennis) Spooner, asexual morph unknown, saprobes, terrestrial, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Aspicilia A. Massal. [nom. cons.] 1852 (= *Agrestia* J.W. Thomson 1961; = *Athecaria* Nyl. 1897; = *Chlorangium* Link 1849; *Manzonia* Garov. 1866; *Paraplacodium* Motyka 1996; *Sagedia* Ach. 1809; *Sphaerothallia* Nees ex Eversm. 1831; = *Urceolaria* Hook. 1821 *fide* Species Fungorum 2017), *Megasporaceae*, *Pertusariales*, *Lecanoromycetes*, 200 species, type: *A. cinerea* (L.) Körb., lichenized (green algae), worldwide, see Lumbsch and Huhndorf (2010; outline), Nordin et al. (2010; phylogeny), Sohrabi et al. (2010a; DNA), Raggio et al. (2011; survive exposure to space), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Aspiciliopsis (Müll. Arg.) M. Choisy 1929, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, one species;

type: *A. macrophthalmia* (Hook. f. & Taylor) B. de Lesd., lichenized, see Hodkinson and Lendemer (2011b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Aspidothelium Vain. 1890 (= *Aspidopyrenis* Clem. & Shear 1931; = *Aspidopyrenium* Vain. 1890; = *Aspidotheliomyces* Cif. & Tomas. 1953; = *Lecania* sect. *Secoligella* Müll. Arg. 1919 [1918]; = *Porinopsis* Malme 1928; = *Secoligella* (Müll. Arg.) Vain. 1896 *fide* Species Fungorum 2017), *Thelenellaceae*, *Ostropales*, *Lecanoromycetes*, 17 species, type: *A. cinerascens* Vain., lichenized, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Aspidotheliaceae*), Kirk et al. (2013; genus accepted), Aptroot et al. (2014d; new species), Nelsen et al. (2016; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Aspilaima Bat. & H. Maia 1961, *Ascomycota* genera *incertae sedis*, one species, type: *A. platoniae* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Aspilidea Hafellner 2001, *Ostropomycetidae* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *A. myrinii* (Fr.) Hafellner, lichenized (green algae), terrestrial (saxicolous), restricted to the alpine zone, see Jørgensen and Nordin (2009, DNA), Lumbsch and Huhndorf (2010; outline), Nordin et al. (2010; taxonomy), Lücking et al. (2017a; classification), sequences unavailable.

Asteridiella McAlpine 1897 (= *Irene* Theiss. 1917; = *Meliola* subgen. *Irene* (Theiss., Syd. & P. Syd.) 1938; = *Parasteridiella* H. Maia 1960 *fide* Species Fungorum 2017), *Meliolaceae*, *Meliolales*, *Sordariomycetes*, more than 300 species, type species: *A. solani* McAlpine, asexual morph unknown, pathogens saprobes, terrestrial, worldwide, see Hosagoudar and Archana (2009; host distribution in India), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013 genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Asterina Lév. 1845 (see Hongsanan et al. 2014c for synonyms), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, more than 700 species, type: *A. melastomatis* Lév., asexual morph coelomycetous or hyphomycetous (=asterostomella-like, clasterosporium-like and mahanteshomyces-like, *fide* Hofmann 2009), epiphytes, biotrophic, terrestrial, worldwide, most species do not have molecular data, see Hofmann (2009; review, DNA), Hosagoudar et al. (2010a; morphology of *Bheemamyces*), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Hongsanan et al. (2014c, 2016c; taxonomy, DNA, keys, nomenclature, evolution), Wijayawardene et al. (2014c;

outline), Guatimosim et al. (2015; phylogeny, taxonomy, DNA), Liu et al. (2017, phylogeny), cultures and sequences are available.

Asterinella Theiss. 1912 (= *Calothyrium* Theiss. 1912; = *Opasterinella* Speg. 1919 [1918] fide Species Fungorum 2017), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, c 40 species, type: *A. puiggarii* (Speg.) Theiss., asexual morph coelomycetous (=asterostomula-like), epiphytes, biotrophic, terrestrial, worldwide, no molecular data, see Hofmann (2009; review, accepted as a genus in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Wijayawardene et al. (2014c; outline), Wu et al. (2014b; review, transferred to *Asterinaceae*), holotype: LPS 1377, cultures and sequences unavailable.

Asterinema Bat. & Gayão 1953, *Dothideomycetes* genera *incertae sedis*, one species, type: *A. puiggarii* (Speg.) Theiss., asexual morph unknown, foliar epiphytes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; accepted as a genus in *Dothideomycetes* genera *incertae sedis*), Wu et al. (2014b; accepted as a genus in *Asterinaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Asterinothyriella Bat. & Cif. 1959, *Ascomycota* genera *incertae sedis*, three species, type: *A. landolphiae* Bat. & Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, Uganda, see Wijayawardene et al. (2012, 2014c, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Asterinothyrium Bat., Cif. & H. Maia 1959, *Ascomycota* genera *incertae sedis*, one species, type: *A. singulatum* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Asterition Leight. 1870, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *A. erumpens* Leight., asexual morph unknown, lichenized fungi, terrestrial, tropical, placement in *Graphidaceae* supported by molecular data, new synonym *Phaeotrema* Müll.Arg., see Lücking et al. (2014, 2017a; classification), sequences are available.

Astroconium Syd. & P. Syd. 1903, *Ascomycota* genera *incertae sedis*, two species, type: *A. saccardoi* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Central America, China, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013 genus accepted), cultures and sequences are unavailable.

Asterodiscus Voglmayr, Gardiennet & Jaklitsch 2016, *Stigmatodiscaceae*, *Stigmatodiscales*, *Dothideomycetes*, two species, type: *A. tamaricis* Voglmayr, Gardiennet & Jaklitsch, asexual morph coelomycetous, saprobes, Europe, see Voglmayr et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: WU 35906, cultures and sequences are available.

Asterodothis Theiss. 1912, *Dothideomycetes* genera *incertae sedis*, one species, type: *A. solaris* (Kalchbr. & Cooke) Theiss., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; transferred this genus from *Asterinaceae* to *Dothideomycetes* genera *incertae sedis*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Asterooglobulus Brackel 2011, *Ascomycota* genera *incertae sedis*, one species, type: *A. giselae* Brackel, coelomycetous, sexual morph unknown, lichenicolous, terrestrial, Europe, see Brackel (2011a; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Asteroma DC. 1815 (= *Actinonemella* Höhn. 1916; = *Depazea* Fr. 1818; = *Gloeosporina* Höhn. 1916; = *Ribaldia* Cif. 1954; = *Titaeosporina* Luijk 1920 fide Sutton 1977; Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, more than 200 species, type: *M. stellaris* (Pers.) Theiss. & Syd., sexual morph *Gnomoniella*, *Plagiostoma*, pathogens associated with leaf spots, saprobes, terrestrial, worldwide distribution, see Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available for *A. alneum*.

Asteromassaria Höhn. 1917, *Dothideomycetes* genera *incertae sedis*, type: *A. macrospora* (Desm.) Höhn., twelve species but could be higher, asexual morph coelomycetous, saprobes, parasites, terrestrial, Europe, India, Japan, USA, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Pleomassariaceae*), Zhang et al. (2012e; accepted as a genus in *Morosphaeriaceae*), Hyde et al. (2013; did not list in *Pleomassariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline as *Dothideomycetes* *incertae sedis*), holotype of type: CBS H-15119, cultures and sequences are available but lacking for the type species.

Asteromella Pass. & Thüm. 1880 (= *Apiospora* Speg. 1912; = *Aplosporidium* Speg. 1912; = *Plectophoma* Höhn. 1907; = *Porterula* Speg. 1920; = *Stictochorella* Höhn. 1917; = *Stictochorellina* Petr. 1922 fide Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, c. 265 species, type: *A. ovata* Thüm., polyphyletic, coelomycetous, sexual morph unknown, widespread, see Hyde et al. (2011; outline), Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available.

Asteromelopsis H.E. Hess & E. Müll. 1951, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, one species, type: *A. insculpta* H.E. Hess & E. Müll., saprobes, Switzerland, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Asteromidium Speg. 1888 (= *Asterostomidium* Lindau 1900 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, three species, type: *A. imperspicuum* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Asteromyces Moreau & M. Moreau ex Hennebert 1962, *Ascomycota* genera *incertae sedis*, one species, type: *A. cruciatus* Moreau & M. Moreau ex Hennebert, hyphomycetous, on seaweeds, sand grains, drift wood, saprobe, mostly marine temperate, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Gulder et al. (2012; biochemistry), Jones et al. (2015; outline), cultures and sequences are unavailable.

Asteronectrioidae Cant. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *A. heveicola* Cant., possibly hyphomycetous, sexual morph unknown, saprobes, Africa, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Asteronia (Sacc.) Henn. 1895 (= *Asterina* subgen. *Asteronia* Sacc. 1882 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, two species, type: *A. sweetiae* Henn., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Asteroporum Müll. Arg. 1884 (= *Asteroporomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, seven species, type: *A. perminutum* Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017b; classification), sequences are unavailable.

Asteroscutula Petr. 1948, *Ascomycota* genera *incertae sedis*, one species, type: *A. sydowii* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Ecuador, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013 genus accepted), cultures and sequences are unavailable.

Asterosporium Kunze 1819, *Asterosporiaceae*, *Diaporthales*, *Sordariomycetes*, seven species, type: *A. asterospermum* (Pers.) Hughes, coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see

Tanaka et al. (2010; taxonomy, phylogeny), Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 112404.

Asterostomopora Bat. & H. Maia 1960, *Ascomycota* genera *incertae sedis*, one species, type: *A. jamaicensis* Bat. & H. Maia, coelomycetous, Jamaica, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Asterostomopsis Bat., Cif. & H. Maia 1959, *Ascomycota* genera *incertae sedis*, one species, *A. ghanaensis* Bat., Cif. & H. Maia, coelomycetous, rarely reported and in case on leaves of *Lawsonia inermis* in Ghana and India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), and sequences are unavailable.

Asterostomula Theiss. 1916, *Ascomycota* genera *incertae sedis*, four species, type: *A. loranthi* Theiss., coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Asterostomulina Bat., J.L. Bezerra & H. Maia 1964, *Ascomycota* genera *incertae sedis*, one species, type: *A. phoebeicola* Bat., J.L. Bezerra & Cavalc., coelomycetous, sexual morph unknown, rarely reported and in case on *Phoebe rivularis*, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Asterotexis Arx 1958, *Asterotexaceae*, *Asterinales*, *Dothideomycetes*, two species, type: *A. cucurbitacearum* (Rehm) Arx, asexual morph unknown, epiphytes, pathogens, biotrophic, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hyde et al. (2013; review, keys, accepted as a genus in *Asterinaceae*), Hongsan et al. (2014c; taxonomy, keys, accepted as a genus in *Asterinaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Asterinaceae*), Guatimosim et al. (2015; phylogeny, DNA, transferred to a newly described family *Asterotexaceae*), Liu et al. (2017, phylogeny, placement), placement in *Asterotexaceae* supported by morphology and phylogeny, syntype: S-F7565, type sequence KP143734 (voucher VIC 24814), HQ610510 (voucher PMA M-0141224).

Astrothyrium Müll. Arg. 1890 (= *Actinoteichus* Cavalc. & Poroca 1971; = *Astrothyriomyces* Cif. & Tomas. 1953; = *Diploeltopsis* Henn. ex Höhn. 1911; = *Psorotheciella* Sacc. & P. Syd. 1905 *fide* Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, 33 species; type: *A. argenteum* Müll. Arg., lichenized, see Flakus and Lücking (2008; new

species), Lücking et al. (2017a; classification), sequences are available.

Astomella Thirum. 1947, *Ascomycota* genera *incertae sedis*, one species, type: *A. neolitseaie* Thirum., saprobes, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Astrochapsa Parnmen, Lücking & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 26 species, type: *A. astroidea* (Berk. & Broome) Parnmen, Lücking & Lumbsch, asexual morph unknown, ascomata apotheciate, conidiomata pycnidial, lichen-forming, corticolous, pantropical, see Parnmen et al. (2012; taxonomy, phylogeny), Rivas Plata et al. (2013; phylogeny), Lumbsch et al. (2014a, b; classification).

Astrocytis Berk. & Broome 1873, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, c. 20 species, type: *P. invocans* Morgan, asexual morph unknown, saprobes, North America, Asia, see Lumbsch and Huhndorf (2010; outline), Pinnoi et al. (2010; new species), Kirk et al. (2013; genus accepted), Daranagama et al. (2015a; new species, DNA), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Astronatelia Bat. & H. Maia 1962, *Ascomycota* genera *incertae sedis*, one species, type: *A. andromedae* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Astrosphaeriella Syd. & P. Syd. 1913 (= *Asterella* Hara 1936; = *Astrotheca* I. Hino 1938; = *Astrosphaeria* (Höhn.) Syd. 1913; = *Astrotheca* I. Hino 1938; = *Leptosphaeria* sect. *Astrosphaeria* Höhn. 1909 *fide* Species Fungorum 2017), *Astrosphaeriellaceae*, *Pleosporales*, *Dothideomycetes*, c. 10 species, type: *A. fusispora* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, widespread, see Tanaka et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Liu et al. (2011b; phylogeny), Zhang et al. (2012e; morphology, phylogeny), Kirk et al. (2013; genus accepted), Ren et al. (2013b; phylogeny), Phookamsak et al. (2015b; taxonomy, phylogeny), cultures available for type: MFLUCC 10-0555.

Astrosphaeriellopsis Phook., J.K. Liu & K.D. Hyde 2015, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *P. thailandensis* Phook., Z.L. Luo & K.D. Hyde, asexual morph unknown, saprobes, Asia, see Phookamsak et al. (2015b; taxonomy, phylogeny), cultures and sequences are available.

Astrothelium Eschw. 1824, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, c. 250 species, type: *A. conicum* Eschw., asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010;

outline), Kirk et al. (2013; genus accepted), Nelsen et al. (2014a; phylogeny) Aptroot and Lücking (2016; monograph), Lücking et al. (2016a, 2017a; classification).

Asymmetrispora Thambug. & K.D. Hyde 2015, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. tennesseensis* (Mugambi, A.N. Mill. & Huhndorf) Thambugala & K.D. Hyde, asexual morph unknown, saprobes, cosmopolitan, see Thambugala et al. (2015b; taxonomy), cultures and sequences are available

Asymmetricospora J. Fröh. & K.D. Hyde 1998, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *A. calamicola* J. Fröh. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Tian et al. (2015; morphology, phylogeny), cultures and sequences are unavailable.

Athallia Arup, Frödén & Söchting 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 13 species, type: *A. holocarpa* (Hoffm.) Arup, Frödén & Söchting, with ascocata, lichenized ('*Pseudotrebouxia*'), (corticulous, epixylic, terricolous, saxicolous), cosmopolitan, see Arup et al. (2013, DNA), Lücking et al. (2017a; classification), Vondrák et al. (2016, geographical range), sequences are available for voucher JV14428 of the type (*fide* Arup et al. 2013).

Atichia Flot. 1850 (= *Actinomma* Sacc. 1884; = *Hyphodictyon* Millardet 1866; = *Myriophysa* Fr. 1849; = *Seuratia* Pat. 1904 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Ascomycota* families *incertae sedis*, four species, type: *A. mosigii* Flot., hyphomycetous, sexual morph formerly known in *Seuratia* Pat., widespread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Atkinsonella Diehl 1950, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *A. hypoxylon* (Peck) Diehl, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Atla Savić & Tibell 2008, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, nine species, type: *A. alpina* Savić & Tibell, lichenized, terrestrial, Europe, see Savić and Tibell (2008a; taxonomy), Lumbsch and Huhndorf (2010; outline), Tibell and Tibell (2015; new species, Alaska), Lücking et al. (2017a; classification), Pykälä and Myllys (2016; new species), holotype and ex-type of the type: UPS, Savić 3129, SS193.

Atopospora Petr. 1925, *Venturiaceae*, *Venturiales*, *Dothideomycetes*, two species, type: *A. betulina* (Fr.)

Petr., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011; accepted as a genus in *Venturiaceae*), Hyde et al. (2013; accepted as a genus in *Venturiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Atractilina Dearn. & Barthol. 1924, *Ascomycota* genera *incertae sedis*, two species, type: *A. callicarpae* Dearn. & Barthol., hyphomycetous, sexual morph unknown, fungicolous, terrestrial, tropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Atractium Link 1809, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *A. stilbaster* Link, hyphomycetous, sexual morph unknown, saprobes or isolated from drinking water, terrestrial, aquatic, worldwide, see Gräfenhan et al. (2011; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), epitype and ex-epitype: CBS 410.67.

Atractobolus Tode 1790, *Ascomycota* genera *incertae sedis*, one species, type: *A. ubiquitarius* Tode, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Atradidymella M.L. Davey & Currah 2009, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *A. muscivora* M.L. Davey & Currah, asexual morph unknown, saprobes, Canada, see Davey and Currah (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), holotype: UAMH 10909, cultures and sequences are available.

Atrocalyx A. Hashim. & Kaz. Tanaka 2017, *Lophiotremataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. acutisporus* A. Hashim. & Kaz. Tanaka, asexual morph coelomycetous, saprobes, Asia, Europe, see Hashimoto et al. (2017; taxonomy, phylogeny), sequences are available.

Atrogeniculata J.S. Monteiro, Gusmão & R.F. Castañeda 2014, *Ascomycota* genera *incertae sedis*, one species, type: *A. submersa* J.S. Monteiro, Gusmão & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes of submerged leaves, Brazil, see Monteiro et al. (2014c; taxonomy), Santa Izabel et al. (2015; taxonomy), Wijayawardene et al. 2017; outline), cultures and sequences are unavailable.

Atrokylindriopsis Y.R. Ma & X.G. Zhang 2015, *Chaetothyriales* genera *incertae sedis*, *Eurotiomycetes*, one species, type: *A. setulosa* Y.R. Ma & X.G. Zhang, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Ma et al. (2015; taxonomy, DNA),

Wijayawardene et al. (2017; outline), holotype and ex-type strain of type: HSAUP H4560; HSAUPmyr4560, HMAS 245592.

Atronectria Etayo 2008, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *A. magellanica* Etayo, asexual morph unknown, parasymbiont on macrolichens, aquatic, South America, see Etayo and Sancho (2008; morphology, distinction of similar lichenicolous genera), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), need revision, holotype: MAF, Etayo 22938, cultures and sequences are unavailable.

Atropellis Zeller & Goodd. 1930, *Godroniaceae*, *Helotiidae*, *Leotiomycetes*, four species, type: *A. pinicola* Zeller & Goodd., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Atrosetaphiale Matsush. 1995, *Ascomycota* genera *incertae sedis*, two species, type: *A. flagelliformis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Atrosynnema J.W. Xia, X.G. Zhang & Z. Li 2016, *Ascomycota* genera *incertae sedis*, one species, type: *A. digitosporum* J.W. Xia, X.G. Zhang & Z. Li, hyphomycetous, sexual morph unknown, saprobic, terrestrial, China, see Xia et al. (2016b; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Atrotorquata Kohlm. & Volk.-Kohlm. 1993, *Cainiaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *A. lineata* Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, Italy, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Liu et al. (2015a; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), only *Atrotorquata spartii* has sequence: MFLU 14-0738, MFLUCC 13-0444, further sequences are necessary for confirm its placement in *Cainiaceae*.

Auerswaldia Sacc. 1883, *Dothideomycetes* genera *incertae sedis*, c. 20 species, type: *A. examinans* (Mont. & Berk.) Sacc., asexual morph coelomycetous, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Liu et al. (2012b; phylogeny, treated as a genus in *Botryosphaeriaceae*), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Botryosphaeriaceae*), Phillips et al. (2013; phylogeny, excluded from *Botryosphaeriaceae*), cultures and sequences are unavailable for type species.

Auerswaldiella Theiss. & Syd. 1914 (= *Dimeriellina* Chardón 1939; = *Stichodothis* Petr. 1927 *fide* Species Fungorum 2017), *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, four species, type: *A. puccinoides* (Speg.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Liu et al. (2012b; accepted as a genus in *Botryosphaeriaceae*), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Aulacostroma Syd. & P. Syd. 1914, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, five species, type: *A. palawanense* Syd. & P. Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Aulaxina Fée 1825, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 13 species, type: *A. opegraphina* Fée, with ascocarps, lichenized (*Trebouxia*), terrestrial (foliicolous and corticolous), tropical, see Lücking (2008; outline), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Sanders and de los Ríos (2016; outline), sequences are available.

Aulographina Arx & E. Müll. 1960, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *A. pinorum* (Desm.) Arx & E. Müll., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as genus in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted as genus in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, transferred to *Teratosphaeriaceae*), Wijayawardene et al. (2014c; outline), holotype: PC 0084689.

Aulographum Lib. 1834, *Aulographaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, c. 30 species, type: *A. hederae* Lib., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures available for type species MFLUCC 13-0001, CPC21373.

Aulospora Speg. 1909, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. epimyces* (Speg.) Speg., asexual morph unknown, saprobes, terrestrial, Argentina, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Aurantiascoma Thambugala & K.D. Hyde 2014, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, one

species, type: *A. minimum* (Mugambi, A.N. Mill. & Huhndorf) Thambugala & K.D. Hyde, asexual morph unknown, saprobes, cosmopolitan, see Mugambi and Huhndorf (2009a; taxonomy, as *Misturatospaeria minima* Mugambi, A.N. Mill. & Huhndorf), Thambugala et al. (2015b; taxonomy), Jaklitsch et al. (2016c; treated as a synonym of *Teichospora* but in here we do not agree with them), cultures and sequences are available.

Aurantiosacculus Dyko & B. Sutton 1979, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *A. eucalypti* (Cooke & Massee) Dyko & B. Sutton, coelomycetous, sexual morph unknown, foliicolous, pathogens, saprobes, three species, Australia, see Crous et al. (2012d; morphology, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of the type: Reader No. 24 K.

Arapex Gryzenh. & M.J. Wingf. 2006, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. penicillata* Gryzenh. & M.J. Wingf., coelomycetous, sexual morph unknown, ?pathogens, saprobes, terrestrial, cosmopolitan, see Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type cultures of type: PREM 57520, CMW 10030 = CBS115740.

Auratiopycnidia Crous & Summerell 2012, *Auratiopycnidellaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. tristaniopsis* Crous & Summerell, coelomycetous, sexual morph unknown, pathogens, terrestrial, Australia, see Crous et al. (2012d; morphology, phylogeny), Maharachchikumbura et al. (2015b; classification, phylogeny), holotype and ex-type: CBS H-20932, CPC 16371.

Aureobasidium Viala & G. Boyer 1891 (= *Aureobasis* Clem. & Shear 1931; = *Dematoidium* Stautz 1931; = *Pachybasidiella* Bubák & Syd. 1915; = *Polyspora* Laff. 1921 [1920-22]; = *Protocoronis* Clem. & Shear 1931; = *Protocoronospora* G.F. Atk. & Edgerton 1907; = *Pullularia* Berkhouw 1923 *fide* Species Fungorum 2017), *Saccotheciaceae*, *Dothideales*, *Dothideomycetes*, 36 species, type: *A. vitis* Viala & G. Boyer, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Peterson et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; taxonomy), ex-epitype: CBS 130298.

Auricularora Kalb 1988, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *A. byssomorpha* (Nyl.) Kalb, lichenized, Neotropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), Lücking et al. (2017a; classification), sequences are unavailable.

Aurifilum Begoude, Gryzenh. & Jol. Roux 2010, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *A. marmelostoma*, asexual morph coelomycetous, pathogens, terrestrial, Cameroon, Taiwan, see Begoude et al. (2010; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type: CBS 124928 =CMW28290; ex-paratypes of type: CBS 124929, CBS 124930; holotype: PREM 60256.

Aurophora Rifai 1968, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *A. dochmia* (Berk. & M.A. Curtis) Rifai, asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Aurophaeria Sun J. Lee, Strobel, Eisenman, Geary, Vargas & S.A. Strobel 2009, *Ascomycota* genera *incertae sedis*, one species, type: *A. flaviradians* Sun J. Lee, Strobel, Eisenman, Geary, Vargas & S.A. Strobel, coelomycetous, sexual morph unknown, saprobes, terrestrial, Bolivia, see Lee et al. (2009c; taxonomy), Wijayawardene et al. (2012, 2017; outline), holotype: MONT 2349, cultures and sequences are unavailable.

Australiaena Matzer, H. Mayrhofer & Elix 1997, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *A. streimannii* Matzer, H. Mayrhofer & Elix, with ascomata, lichenized (green algae), terrestrial (saxicolous), Australia, Papua-New Guinea see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Austrella P.M. Jørg. 2004, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *A. arachnoidea* P.M. Jørg., with ascomata, lichenized (*Nostoc*), terrestrial (corticulous), Australia, New Zealand, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available for the type species.

Astroafricana Quaedvl. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *A. associata* (Crous & Carnegie) Quaedvl. & Crous, asexual morph unknown, pathogens, terrestrial, Australia, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; outline), holotype: CBS-H 19833, ex-type: CPC 13119.

Astrocenangium Gamundí 1997, *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *A. australe* (Speg.) Gamundí, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Astrographa Sparrius, Elix & A.W. Archer 2010, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, three species,

type: *A. kurriminensis* Sparrius, Elix & A.W. Archer, lichenized, see Sparrius et al. (2010, 2013; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Austrolecia Hertel 1984, *Catillariaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *A. antarctica* Hertel, lichenized (green algae), terrestrial (saxicolous), Antarctica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Austroparmeliella (P.M. Jørg.) P.M. Jørg. 2014, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *A. arachnoidea* P.M. Jørg., with ascomata, lichenized (*Nostoc*), terrestrial (corticulous), Australia, New Zealand, see Lücking et al. (2017a; classification), sequences are available.

Austroparmelia A. Crespo, Divakar & Elix 2010, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 15 species, type: *A. endoleuca* (Taylor) A. Crespo, Divakar & Elix, with ascomata, lichenized (green algae), terrestrial (corticulous, saxicolous), Australasia, South Africa, see Crespo et al. (2010a, b; taxonomy), Kantvilas and Divakar (2013; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Austropeltum Henssen, Döring & Kantvilas 1992, *Sphaerophoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *A. glareosum* Henssen, Döring & Kantvilas, with ascomata, lichenized (green algae), terrestrial (terricolous), Tasmania and New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification).

Austropozia Spooner 1987, *Arachnopezizaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *A. samuelsii* (Korf) Spooner, sexual morph unknown, saprobes, New Zealand, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Hyaloscyphaceae*), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Austroplaca Søchting, Frödén & Arup 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, ten species, type: *A. ambitiosa* (Darb.) Søchting, Frödén & Arup, lichenized, terrestrial, widespread, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification); sequences are available.

Astropleospora R.G. Shivas & L. Morin 2010, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *A. osteospermi* R.G. Shivas & L. Morin, asexual morph coelomycetous, pathogens, terrestrial, Australia, South Africa, see Morin et al. (2010), Thambugala et al. (2014e; taxonomy), Ariyawansa et al. (2015a; phylogeny), holotype of the type: BRIP 52234, isotype BRIP 51628 (includes ex-isotype culture).

Austroroccella Tehler & Ertz 2013, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *A. gayana* (Mont.) Tehler & Ertz, sexual morph with apothecioid ascomata, asexual morph coelomycetous, lichenized, terrestrial, Chile, see Tehler et al. (2013b; morphology, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Austrostigmidium Pérez-Ortega & Garrido-Benavent 2015, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *A. mastodiae* Pérez-Ortega & Garrido-Benavent, lichenicolous, terrestrial, Antarctica, Tierra del Fuego, asexual morph coelomycetous, see Pérez-Ortega et al. (2015, phylogeny, taxonomy), closely related to the lichenicolous genus *Xanthoriicola* and the rock-inhabiting fungal genus *Friedmanniomycetes*, holotype: MA-Lich 18215.

Austrotrema I. Medeiros, Lücking & Lumbsch 2017, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *A. bicinctulum* (Nyl.) I. Medeiros, Lücking & Lumbsch, lichenized, see Lücking et al. (2017a; classification), Medeiros et al. (2017; taxonomy), sequences are available.

Autoicomycetes Thaxt. 1908, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 28 species, type: *A. acuminatus* (Thaxt.) Thaxt., asexual morph unknown, parasitic on Hydrophilidae beetles, aquatic, Asia, Europe, North America, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Autophagomyces Thaxt. 1912 (= *Bordea* Maire 1916 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 20 species, type: *A. platensis* Thaxt., asexual morph unknown, parasitic on Anthicidae, Phalacridae, and Staphylinidae beetles, terrestrial, Africa, Asia, North America, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Auxarthron G.F. Orr & Kuehn 1963, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, 17 species, type: *A. californiense* G.F. Orr & Kuehn, asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Crous et al. (2013b; new species, Portugal), Hubka et al. (2013a; new species), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Auxarthronopsis Rahul Sharma, Gräser & S.K. Singh 2011, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, 18 species, type: *A. californiense* G.F. Orr & Kuehn, asexual morph hyphomycetous, saprobes, terrestrial, India, see Sharma et al. (2011; taxonomy), cultures and sequences are available.

Avesicidiella W.P. Wu, B. Sutton & Gange 1997, *Ascomycota* genera *incertae sedis*, two species, type: *A. britannica* Wu, Sutton & Gange, saprobes, hyphomycetous, sexual morph unknown, UK, China, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: IMI 372453, sequence are unavailable.

Avettaea Petr. & Syd. 1927, *Ascomycota* genera *incertae sedis*, three species, type: *A. philippinensis* Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Awasthiella Kr. P. Singh 1980, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *A. indica* Kr. P. Singh, lichenized, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Ayria Fryar & K.D. Hyde 2004, *Annulatascaeae*, *Annulatascales*, *Sordariomycetes*, two species, type: *A. appendiculata* Fryar & K.D. Hyde, saprobes, aquatic, Costa Rica, USA, Brunei, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2016b; outline), holotype: HKU (M) 15553.

Azbuskinia Lar. N. Vassiljeva 1989, *Sordariomycetes* genera *incertae sedis*, one species, type: *A. ferruginea* (Fuckel) Lar.N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Azureothecium Matsush. 1989, *Eurotiomycetidae* genera *incertae sedis*, *Eurotiomycetes*, two species, type: *A. australiense* Matsush., asexual morph unknown, saprobes, from soil, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Babjeviella Kurtzman & M. Suzuki 2010, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *B. inositovora* (Golubev & Blagod.) Kurtzman & M. Suzuki, asexual reproduction by multilateral budding and cells are short-ellipsoid to elongate, saprophytes, known from peat, rotted balsam fir (*Abies balsamea*), insect frass, unidentified tree, Russia, Canada, USA, see Kurtzman and Suzuki (2010; taxonomy), Kurtzman and Robnett (2013b; phylogeny), cultures and sequences are available.

Bachmanniomycetes D. Hawksw. 1981, *Ascomycota* genera *incertae sedis*, type: *B. uncialicola* (Zopf) D. Hawksw. (syn. *Phyllosticta uncialicola* Zopf), three species, coelomycetous, sexual morph unknown, gall-

forming on lichens, Europe, North and South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), lectotype: E, cultures and sequences are unavailable.

Bacidia De Not. 1846 (= *Bacidiomyces* Cif. & Tomas. 1953; = *Byssopsora* A. Massal. 1861; = *Psorella* Müll. Arg. 1894; = *Sporacestra* A. Massal. 1860 [1859–1860]; = *Urophora* Sommerf. ex Arnold 1899 *fide* Species Fungorum 2017), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, 230 species, type: *B. rosella* (Pers.) De Not., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Bacidina Vězda [nom. cons.] 1991, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, twelve species, type: *B. phacodes* (Körb.) Vězda, lichenized, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Czarnota and Guzow-Krzeminska (2012; DNA), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Baciadiopsora Kalb 1988, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *B. squamulosa* (Nyl.) Kalb (*Lecidea squamulosa* Nyl.), lichenized, corticolous, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Bacillopeltis Bat. 1957, *Ascomycota* genera *incertae sedis*, one species, type: *B. paypayrolae* Bat., coelomycetous, sexual morph unknown, saprobes, on soil, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bactridium Kunze 1817 (= *Damnosporium* Corda 1842; = *Ericianella* Brond. 1828 [1828–1830]; = *Podobactridium* Penz. & Sacc. ex Petch 1916 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, 15 species, type: *B. flavum* Kunze, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bactrodesmiastrum Hol.-Jech. 1984, *Sordariomycetes* genera *incertae sedis*, two species, type: *B. obscurum* Hol.-Jech., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Bactrodesmiella M.B. Ellis 1959, *Ascomycota* genera *incertae sedis*, two species, type: *B. masonii* (S. Hughes) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bactrodesmium Cooke 1883, *Dothideomycetes* genera *incertae sedis*, more than 48 species, type: *B. abruptum* (Berk. & Broome) Mason & Hughes, sexual morph *Stuartella* spp., saprobes, terrestrial and aquatic, worldwide, see Zhao et al. (2009; outline, diversity), da Cruz and Gusmão (2009; diversity), Koukol and Kolářová (2010; morphology, phylogeny), Kalko (2011; diversity), Seephueak et al. (2011a, 2011b; diversity), Hernández-Restrepo et al. (2013; morphology), Tanaka et al. (2015; phylogeny), sequences are available.

Bactrosphaeria Penz. & Sacc. 1897, *Sordariomycetes* genera *incertae sedis*, one species, type: *B. asterostoma* Penz. & Sacc., asexual morph unknown, saprobe, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bactrospora A. Massal. 1852 (= *Bactrosporomyces* Cif. & Tomas. 1953; = *Melampydiomyces* Cif. & Tomas. 1953; = *Melampyllum* Stirt. ex Müll. Arg. 1894; = *Scolecactis* Clem. 1909 *fide* Species Fungorum 2017), *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, 35 species, type: *B. dryina* (Ach.) A. Massal., lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Knudsen et al. (2011; new species), Kirk et al. (2013; genus accepted), Sobreira et al. (2015; new species), Lücking et al. (2017a; classification), sequences are unavailable.

Baculifera Marbach & Kalb 2000, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 14 species, type: *B. orosa* Marbach., with ascocata, lichenized (green algae), terrestrial (mostly corticolous and lignicolous), tropical, see Elix and Kantvilas (2014; new records), Elix and Mayrhofer (2016; outline), Lücking et al. (2017a; classification), Prieto and Wedin (2016; DNA, genus accepted), sequences are available but lacking for the type.

Baculospora Zukal 1887, *Ascomycota* genera *incertae sedis*, one species, type: *B. pellucida* Zukal, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bacusphaeria Norlailatul, Alias & Suetrong 2017, *Tirisporellaceae*, *Tirisporellales*, *Sordariomycetes*, one species, type: *B. nypae* Norlailatul, Alias & Suetrong, asexual morph unknown, saprobes, marine, Malaysia, see Abdel-Wahab et al. (2017; phylogeny), holotype and ex-type: MFLU 13-0617, BCC51275, sequences are available.

Badarisama Kunwar, J.B. Manandhar & J.B. Sinclair 1986, *Ascomycota* genera *incertae sedis*, one species, type: *B. sojae* Kunwar, J.B. Manandhar & J.B. Sinclair, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Badimia Vězda [nom. cons.] 1986 (= *Pseudogyalecta* Vězda 1975 fide Species Fungorum 2017), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, 18 species, type: *B. dimidiata* (Bab. ex Leight.) Vězda, lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; DNA), Lücking et al. (2017a; classification), sequences are available.

Badimiella Malcolm & Vězda 1994, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *B. serusiauxii* Malcolm & Vězda, lichenized, south temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Baeomyces Pers. 1794 (= *Baeomycetomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Cladoniopsis* Zahlbr. 1941; = *Cyanobaeis* Clem. 1909; = *Ludovicia* Trevis. 1857; = *Sphyridiomycetes* E.A. Thomas ex Cif. & Tomas. 1953; = *Sphyridium* Flot. 1843 fide Species Fungorum 2017), *Baeomycetaceae*, *Baeomycetales*, *Lecanoromycetes*, nine species, type: *B. byssoides* (L.) P. Gaertn., G. Mey. & Scherb., lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; DNA), Lücking et al. (2017a; classification), sequences are available.

Baflavia Lücking 2008, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *B. flavescens* (Lücking) Lücking, lichenized, see Lücking et al. (2017a; classification), sequences are unavailable.

Bagadiella Cheew. & Crous 2009, *Clypeophysalosporaceae*, *Amphisphaeriales*, *Sordariomycetes*, two species, type: *B. lunata* Cheew. & Crous, hyphomycetous, sexual morph unknown, saprobes, ?pathogens, terrestrial, Australasia, see Cheewangkoon et al. (2009; taxonomy, phylogeny), Crous et al. (2011a; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Giraldo et al. (2017; phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), holotype and ex-type: CBS H-20281, CBS 124762.

Bagcheea E. Müll. & R. Menon 1954, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *B. albomaculans* (Fukui) Hino & Katumoto asexual morph unknown, mild pathogens, terrestrial, temperate and tropical, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Baggea Auersw. 1866, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *B. pachyascus*

Auersw., asexual morph unknown, saprobes, terrestrial, Germany, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), neotype: K herbarium.

Bagliettoa A. Massal. 1853, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 20 species, type: *B. limborioides* A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Bagnisiella Speg. 1880 (= *Robertomyces* Starbäck 1880 fide Species Fungorum 2017), *Aplosporellaceae*, *Botryosphaeriales*, *Dothideomycetes*, c. 20 species, type: *B. australis* Speg., asexual morph unknown, saprobes, South America, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; transferred to *Aplosporellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Bagnisimitrula S. Imai 1942, *Leotiomycetes* genera incertae sedis, one species, type: *B. saccardoa* (Bagnis) S. Imai, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bahianora Kalb 1984, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *B. poeltii* Kalb, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Bahuchashaka Subram. 1978, *Ascomycota* genera incertae sedis, one species, type: *B. graminis* (Matsush.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bahukalasa Subram. & Chand rash. 1979, *Ascomycota* genera incertae sedis, one species, type: *B. samala* Subram. & Chand rash., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bahusakala Subram. 1958, *Dothideomycetes* genera incertae sedis, four species, type: *B. olivaceonigra* (Berk. & Broome) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, Caribbean, North America, see Kang et al. (2010; DNA), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), cultures and sequences are available.

Bahusandhika Subram. 1956, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, four species, type: *B. indica* (Subram.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Pratibha et al. (2014; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Bahusutrabeja Subram. & Bhat 1977, *Botryosphaeriales* genera *incertae sedis*, *Dothideomycetes*, six species, type: *B. dwaya* Subram. & Bhat, hyphomycetous, sexual morph unknown, saprobes, terrestrial, sometimes aquatic, cosmopolitan see Shenoy et al. (2010, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Li et al. (2013; new species, key), Gao et al. (2015a; new record for *B. angularis* from China), sequences are available.

Baipadisphaeria Pinruan 2011, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *B. spathulospora* Pinruan, asexual morph unknown, saprobes, aquatic, on decaying leaves of peat swamp palm *Licuala longicalycata*, Thailand, see Pinruan et al. (2010; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type culture of type: BBH Pinruan Wah32A, BCC16119.

Bakerophoma Died. 1916, *Ascomycota* genera *incertae sedis*, three species, type: *B. sacchari* Died., lichenicolous, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Balaniopsis P.M. Kirk 1985, *Ascomycota* genera *incertae sedis*, four species, type: *B. africana* (Kiffer) P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, on soil, terrestrial, Africa, Australasia, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Balanium Wallr. 1833, *Ascomycota* genera *incertae sedis*, one species, type: *B. stygium* Wallr., hyphomycetous, sexual morph unknown, saprobes, on soil, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Balsania Speg. 1885 (= *Dothichloë* G.F. Atk. 1894; = *Ephelis* Fr. 1849; = *Ophiodothis* Sacc. 1883 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, 27 species, type: *B. claviceps* Speg., asexual morph unknown, saprobes, terrestrial, subtropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are available.

Balazucia R.K. Benj. 1968, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *B. bilateralis* R.K. Benj., asexual morph unknown, parasitic on Staphylinidae beetles, terrestrial, Central America and Japan, see Rossi and Weir (2008; treated as a synonym of *Cucujomyces*), Lumbsch and Huhndorf (2010; outline, misspelled as *Balazusia*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Balladyna Racib. 1900 (= *Wageria* Stevens & Dalbey. 1918), *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, c. 17 species, type: *B. gardeniae* Racib., asexual morph unknown, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Balladynocallia Bat. 1965, *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, three species, type: *B. glabra* (Hansf.) Bat., asexual morph unknown, saprophytes on leaves, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014; outline), cultures and sequences are unavailable.

Balladynopsis Theiss. & Syd., 1917 (= *Balladynastrum* Hansf. 1941 [1940-41]; = *Xenostigmella* Petr. 1950 *fide* Species Fungorum 2017), *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, 17 species, type: *B. philippinensis* Syd. & P. Syd., asexual morph hyphomycetous, mostly saprobes on leave, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences unavailable.

Balsamia Vittad. 1831 (= *Pseudobalsamia* E. Fisch. 1907 *fide* Species Fungorum 2017), *Helvellaceae*, *Pezizales*, *Pezizomycetes*, six species, type: *B. vulgaris* Vittad., asexual morph unknown, saprobes, ?from soil, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014d; DNA, phylogeny), sequences are available.

Bambusaria Jaklitsch, D.Q. Dai, K.D. Hyde & Voglmayr 2015, *Valsariaceae*, *Valsariales*, *Dothideomycetes*, one species, type: *B. bambusae* (J.N. Kapoor & H.S. Gill) Jaklitsch, D.Q. Dai, K.D. Hyde & Voglmayr, asexual morph coelomycetous, saprobes, terrestrial, India and Thailand, see Jaklitsch et al. (2015; phylogeny, morphology, sexual and asexual morphs), Dai et al. (2017; description), reference specimen culture of type: MFLUCC 12-0851 = CBS 139763 = MUCL 55880.

Bambusicola D.Q. Dai & K.D. Hyde 2012, *Bambusicolaceae*, *Pleosporales*, *Dothideomycetes*, nine species, type: *B. massarinia* D.Q. Dai & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Thailand, see Dai et al. (2012, 2015b, 2017; phylogeny,

morphology, sexual morph, key), Wijayawardene et al. (2014c; outline), Thambugala et al. 2017b, new species, phylogeny), ex-type: MFLUCC 11–0389 =CBS 139985.

Bambusicularia Klaubauf, Lebrun & Crous 2014, Pyriculariaceae, Magnaportheales, Sordariomycetes, one species, type: *B. brunnea* Klaubauf, Lebrun & Crous, hyphomycetous, sexual morph unknown, pathogens, terrestrial, Japan, see Klaubauf et al. (2014; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: CBS H-21839, ex-type strain of type: CBS 133599.

Bambusistroma D.Q. Dai & K.D. Hyde 2015, Massarinaceae, Massarineae, Pleosporales, Dothideomycetes, one species, type: *B. didymosporum* D.Q. Dai & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Adamčík et al. (2015; phylogeny, morphology, sexual morph), holotype and ex-type strain of type: MFLU 15–0057, MFLUCC 13–0862.

Banhegyia Zeller & Tóth 1960, Dothideomycetes genera incertae sedis, two species, type: *B. setispora* L. Zeller & Tóth, asexual morph unknown, saprobes, terrestrial, Europe, see Sanderson et al. (2009; excluded the genus *Banhegyia* due to differing morphology and phylogeny and synonymized *Banhegyia* with *Mycoelaspilea*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; did not accept as a genus in Patellariaceae but included in key due to historical reasons), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; did not accept as a genus in Patellariaceae), holotype: BP 2835.

Banksiamyces G.W. Beaton 1982, Helotiales genera incertae sedis, Leotiomycetes, four species, type: *B. macrocarpus* G.W. Beaton, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence (ITS) is available for an unidentified species.

Banksiophoma Crous 2017, Phaeosphaeriaceae, Pleosporales, Dothideomycetes, one species, type: *B. australiensis* Crous, coelomycetous, sexual morph unknown, saprobes, Australia, see Crous et al. (2017; taxonomy), holotype: CBS H-23121, cultures and sequences are available.

Bapalmuia Sérus. 1993, Pilocarpaceae, Lecanorales, Lecanoromycetes, 23 species, type: *B. palmularis* (Müll. Arg.) Sérus., lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), van den Broeck et al. (2014; new species, Congo), Lücking et al. (2017a; classification), sequences are available.

Barbarosporina Kirulis 1942, Ascomycota genera incertae sedis, one species, type: *B. rhytismatis* Kirulis, coelomycetous, sexual morph unknown, former USSR see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Barbatosphaeria Réblová 2008 (= *Tectonidula* Réblová 2009 fide Réblová et al. 2015c), Sordariomycetes genera incertae sedis, eight species, type: *B. barbirostris* (Dufour) Réblová, asexual morph unknown, saprobes, tropical, see Lumbsch and Huhndorf (2010; outline), Réblová et al. (2015c; new species), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures available for the type: CBS 121149.

Barnettella D. Rao & P. Rag. Rao 1964 (= *Vinculum* R.Y. Roy, R.S. Dwivedi & P.K. Khanna 1965 fide Species Fungorum 2017), Ascomycota genera incertae sedis, one species, type: *B. speciosa* D. Rao & P. Rag. Rao, coelomycetous, sexual morph unknown, saprobes, on soil, terrestrial, India, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), cultures and sequences are unavailable.

Barnettozyma Kurtzman, Robnett & Bas.-Powers 2008, Phaffomycetaceae, Saccharomycetales, Saccharomycetes, ten species, type: *B. populi* (Phaff, Y. Yamada, Tredick & M. Miranda) Kurtzman, Robnett & Basehoar-Powers, asexual reproduction by budding, saprophytic, on soil, water, tree fluxes, insect frass in trees, animal dung, one clinical isolate, worldwide, see Kurtzman et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Barrenia E. Walsh & N. Zhang 2015, Helotiales genera incertae sedis, Leotiomycetes, two species, type: *B. panicia* E. Walsh & N. Zhang, hyphomycetous, sexual morph unknown, root endophytes, terrestrial, USA, see Walsh et al. (2015; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: RUTPP WSF1R37.

Barretomyces Klaubauf, Lebrun & Crous 2014, Pyriculariaceae, Magnaportheales, Sordariomycetes, hyphomycetous, one species, type: *B. calathea* (Soares, Rocha & Barreto) Klaubauf, Lebrun & Crous, sexual morph unknown, pathogens, terrestrial, Brazil, see Klaubauf et al. (2014; morphology, phylogeny), Maharachchikumbura et al. (2015b; outline), holotype of the type: VIC 30699, ex-type strain of type: CBMAI 1060.

Barria Z.Q. Yuan 1994, Didymosphaeriaceae, Pleosporales, Dothideomycetes, one species, type: *B. piceae* Z.Q. Yuan, asexual morph unknown, pathogens, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014e; transferred the genus *Barria* from Phaeosphaeriaceae to Didymosphaeriaceae), Phookamsak et al. (2014c; accepted as a genus in Didymosphaeriaceae), Wijayawardene et al. (2014c; outline), sequences are unavailable, isotype of type species NY 00164027.

Barrina A.W. Ramaley 1997, *Coniochaetaceae*, *Coniochaetales*, *Sordariomycetes*, one species (type): *B. polyspora* A.W. Ramaley, asexual morph *phialophora*-like, saprobes, USA see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b; outline), sequences are available.

Barriopsis A.J.L. Phillips, A. Alves & Crous, Persoonia 21: 39. 2008, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, four species, type: *P. fusca* N.E. Stevens [\equiv *Barriopsis fusca* nom. illeg.; \equiv *Barriopsis stevensiana* A.J.L. Phillips & Pennycook], coelomycetous, saprobes, weak pathogens, terrestrial, distribution not known, see Abdollahzadeh et al. (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Liu et al. (2012b; phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Phillips et al. (2013; phylogeny), Doilom et al. (2014; new species), Wijayawardene et al. (2014c, 2016b; outline, taxonomy), Konta et al. (2016c; new species), Tibpromma et al. (2017b; new species, phylogeny), ex-type: CBS 174.26.

Barriopsis stevensiana A.J.L. Phillips & Pennycook, nom. nov. MycoBank MB819573

\equiv *Barriopsis fusca* (N.E. Stevens) A.J.L. Phillips, A. Alves & Crous, Persoonia 21: 39. 2008, nom illeg. [ICN(Melbourne) Art. 52.1], replaced synonym.

\equiv *Phaeobotryosphaeria fusca* (N.E. Stevens) Petr., Sydowia 6: 317. 1952.

\equiv *Physalospora fusca* N.E. Stevens, Mycologia 18: 210. 1926.

Etymology: Named in honour of N.E. Stevens, the American mycologist and plant pathologist.

See Phillips et al. (2008) for illustrations, description and diagnosis.

Holotype: Cuba, Herradura, on twigs of *Citrus* sp., 15 Jan. 1925, N.E. Stevens, holotype BPI 599052, culture ex-type CBS 174.26.

Notes: Phillips et al. (2008) introduced *Barriopsis fusca* as a new combination based on *Physalospora fusca*, but inadvertently rendered it illegitimate (superfluous) by listing *Botryosphaeria disrupta* (Berk. & M.A. Curtis) Arx & E. Müll., as a synonym. However, this synonymy is erroneous; the dark brown, aseptate ascospores of *B. stevensiana* are distinct from the light brown (“hellbraun”) ascospores of *B. disrupta* sensu von Arx and Müller (1954), and clearly distinct from the hyaline ascospores of *Sphaeria disrupta*, the basionym of *B. disrupta* (Berkeley 1876).

Barraelia Rappaz 1995, *Xylariales* genera incertae sedis, *Sordariomycetes*, three species, type: *B. rhamnicola* Rappaz, six species, asexual morph libertella-like, terrestrial, saprobes, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al.

(2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; placed in *Xylariales* genera incertae sedis), cultures and sequences are available.

Barssia Gilkey 1925 (= *Phymatomyces* Kobayasi 1937 fide Species Fungorum 2017), *Helvellaceae*, *Pezizales*, *Pezizomycetes*, four species, type: *B. oregonensis* Gilkey, asexual morph unknown, saprobes, ?from soil, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014d; new species), Hongsanan et al. (2017a; evolution), sequences are available for the type species.

Bartalinia Tassi 1900 (= *Hyalotia* Guba 1961; = *Pestalozzina* P. Karst. & Roum. 1890 fide Species Fungorum 2017), *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, 17 species, type: *B. robillardoides* Tassi, saprobes, associated with leaf spots, cosmopolitan, see Marincowitz et al. (2010; new species), Wijayawardene et al. (2012, 2016b, 2017; phylogeny, taxonomy, outline), Maharachchikumbura et al. (2015b, 2016b; phylogeny, outline), Senanayake et al. (2015; phylogeny, morphology), cultures and sequences are available.

Bartlettiella D.J. Galloway & P.M. Jørg. 1990, *Lecanoromycetes* genus incertae sedis one species, type: *B. fragilis* D.J. Galloway & P.M. Jørg., with ascomata, lichenized (green algae), lichenized, terrestrial (saxicolous), New Zealand, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a, classification), cultures and sequences are unavailable.

Barubria Vezda 1986, *Pilocarpacae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *B. fuscorubra* (Vezda) Vezda, with ascomata, lichenized (green algae), terrestrial (foliicolous), Neotropics and tropical Africa, rare see Lücking (2008; outline), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Farkas (2015; nomenclature), Lücking et al. (2017a, classification), cultures and sequences are unavailable.

Basauxia Subram. 1995 (= *Basauxia* Subram. 1992 fide Species Fungorum 2017), *Ascomycota* genera incertae sedis, one species, type: *B. pulchra* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: MUBL 3126, cultures and sequences are unavailable.

Basavamyces Hosag. 2005, *Meliolaceae*, *Meliolales*, *Sordariomycetes*, two species, type: *B. litseae* Hosag., C.K. Biju & T.K. Abraham, asexual morph unknown, obligate parasites, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Hosagoudar and Archana (2009; key), holotype (HCIO 43453), cultures and sequences are unavailable.

Basidiobotrys Höhn. 1909, *Sordariomycetes* genera *incertae sedis*, one species, type: *B. clautriavii* (Pat.) Höhn., hyphomycetous, sexual morph unknown, terrestrial, saprobes, see Wendt et al. (2017; resurrected the genus), cultures and sequences are available.

Basifimbria Subram. & Lodha 1968, *Ascomycota* genera *incertae sedis*, three species, type: *B. aurea* Subram. & Lodha, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology, accepted as genus in *Pyronemataceae*), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Batcheloromyces Marasas, P.S. van Wyk & Knox-Dav. 1975, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *B. proteae* Marasas, P.S. van Wyk & Knox-Dav., hyphomycetous, sexual morph unknown, pathogens, saprobes, terrestrial, Australia, Thailand, see Crous et al. (2009e; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017, phylogeny, outline), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2014; phylogeny), cultures available for type: CBS 110696.

Bathelium Ach. 1803, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, 16 species, type: *B. mastoideum* Afzel. ex Ach., lichenized, see Nelsen et al. (2014a; phylogeny), Aptroot and Lücking (2016; monograph), Lücking et al. (2017a; phylogeny; classification), sequences are available.

Bathyascus Kohlm. 1977, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, five species, type: *B. vermisporus* Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: J.K. 1947 =00966736 NYBG, cultures and sequences are unavailable.

Batistia Cif. 1958 (= *Acrostroma* Seifert 1987 *fide* Rossman et al. 2016a), *Batistiaceae*, *Ascomycota* families *incertae sedis*, one species, type: *B. annulipes* (Mont.) Cif., asexual morph formerly known in *Acrostroma*, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Batistina Peres 1961, *Ascomycota* genera *incertae sedis*, one species, type: *B. splendida* Peres, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Batistinula Arx 1960, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, one species, type: *B. gallesiae* Arx, asexual morph triposporium-like, epiphytes, biotrophic, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al.

(2013; review, keys), Hongsanan et al. (2014c; taxonomy, keys), Wijayawardene et al. (2014c; outline), Guatimosim et al. (2015; phylogeny, DNA), holotype: URM 19988, cultures and sequences are unavailable.

Batistospora J.L. Bezerra & M.P. Herrera 1964, *Ascomycota* genera *incertae sedis*, one species, type: *B. septentrionalis* J.L. Bezerra & M.P. Herrera, asexual morph unknown, saprobes, Neotropics, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Battarrina (Sacc.) Clem. & Shear 1931 (= *Hypocrea* subgen. *Battarrina* Sacc. 1883 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *B. inclusa* (Berk. & Broome) Clem. & Shear, asexual morph unknown, parasitic (fungicolous), terrestrial, UK, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: K.

Baudoinia J.A. Scott & Unter. 2007, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *B. compniacensis* (Richon) J.A. Scott & Unter., hyphomycetous, sexual morph unknown, saprobes, terrestrial and aquatic, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Scott et al. (2016; new species, phylogeny), epitype and ex-epitype: DAOM 238773, UAMH 10808.

Beauveria Vuill. 1912, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *B. bassiana* (Bals.-Criv.) Vuill., asexual morph unknown, saprobes, terrestrial, Central and South America, see Lumbsch and Huhndorf (2010; outline), Rehner et al. (2011; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Robène-Soustrade et al. (2015; new species), Imoulan et al. (2016; new species, entomopathogens), cultures and sequences are available.

Beauveriphora Matsush. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *B. sporotrichis* Matsush., hyphomycetous, sexual morph unknown, saprobes, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017, outline), cultures and sequences are unavailable.

Beccopycnidium F. Stevens 1930, *Ascomycota* genera *incertae sedis*, one species, type: *B. palmicola* F. Stevens, coelomycetous, sexual morph unknown, see Wijayawardene et al. (2012, 2017, outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bejjadwya Subram. 1977, *Ascomycota* genera *incertae sedis*, one species, type: *B. bispora* (Matsu.) Subram., hyphomycetous, sexual morph unknown, saprobe, terrestrial, Japan, see Seifert et al. (2011; morphology),

Wijayawardene et al. (2012, 2017, outline), cultures and sequences are unavailable, holotype: Matsushima herb. 4861.

Beelia F. Stevens & R.W. Ryan 1925, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *B. suttoniae* F. Stevens & R.W. Ryan, asexual morph unknown, saprobes, Hawaii, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Elsinoaceae*), Kirk et al. (2013; genus accepted), Jayawardene et al. (2014; transferred to *Chaetothyriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Belemnospora P.M. Kirk 1981, *Ascomycota* genera *incertae sedis*, seven species, type: *B. epiphylla* P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, Asia, Australasia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Belizeana Kohlm. & Volk.-Kohlm. 1987, *Dothideomycetes* genera *incertae sedis*, one species, type: *B. tuberculata* Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, marine, Belize, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Elsinoaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Jones et al. (2015; outline, accepted as a genus in *Dothideales* genera *incertae sedis*), cultures and sequences are unavailable.

Bellemerea Hafellner & Cl. Roux 1984, *Lecanoromycetes* genus *incertae sedis*, nine species, type: *B. alpina* (Sommerf.) Clauzade & Cl. Roux, with ascocarps, lichenized (green algae), terrestrial (saxicolous), Eurasia, North America, Australia, New Zealand and Antarctica, montane and boreal, arctic-alpine/Antarctic regions, see Ekman et al. (2008, DNA), Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Miadlikowska et al. (2014a; DNA, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available for the type.

Bellemerella Nav.-Ros. & Cl. Roux 1997, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, four species, type: *B. trapeliae* Nav.-Ros. & Cl. Roux, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bellojisia Réblová 2008, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *B. rhynchostoma* (Höhn.) Réblová, asexual morph unknown, saprobes, terrestrial, Austria, see Réblová (2008; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), culture of the type: CBS 118484.

Bellulicauda B. Sutton 1967, *Ascomycota* genera *incertae sedis*, two species, type: *B. dialii* (Syd.) B.

Sutton, coelomycetous, sexual morph unknown, saprobes, Italy, Sierra Leone, Zambia, see Wijayawardene et al. (2012, 2016b, 2017; taxonomy, new species, outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of the type: IMI 21609.

Beloniocyphella Höhn. 1918, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, four species, type: *B. hypnorum* (Syd. & P. Syd.) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Huhtinen et al. (2010; new records), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequence are available.

Beltramono Rashmi Dubey, A.K. Pandey bis & Manohar. 2011, *Ascomycota* genera *incertae sedis*, one species, type: *B. costi* Dubey, Pandey & Manohar., hyphomycetous, sexual morph unknown, pathogens, terrestrial, India, see Dubey et al. (2011; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Beltrania Penz. 1882, *Beltraniaceae*, *Amphisphaeriales*, *Sordariomycetes*, 14 species, type: *B. rhombica* Penz., saprobes, worldwide, see Kirk et al. (2013; genus accepted), Crous et al. (2015e; emended the family), Maharachchikumbura et al. (2015b, 2016b; outline), Rajeshkumar et al. (2016; taxonomy, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Beltraniella Subram. 1952 (= *Ellisiopsis* Bat. 1956 *fide* Species Fungorum 2017), *Beltraniaceae*, *Amphisphaeriales*, *Sordariomycetes*, 21 species, type: *B. odinae* Subram., saprobes, worldwide, see Shirouzu et al. (2010; new species, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2014c, 2016a; new species, New South Wales), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; new species, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Beltraniomyces Manohar., D.K. Agarwal & Rao 2003, *Ascomycota* genera *incertae sedis*, one species, type: *B. lignicola* Manohar., D.K. Agarwal & Rao, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Beltraniopsis Bat. & J.L. Bezerra 1960, *Beltraniaceae*, *Xylariales*, *Sordariomycetes*, nine species, type: *B. odinae* Subram., saprobes, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2014c; new species, New South Wales), Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Benguetia Syd. & P. Syd. 1917, *Helotiales* genera *incertae sedis*, one species, type: *B. omphalodes* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Beniowskia Racib. 1900 (= *Clathrotrichum* Pat. 1921 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, four species, type: *B. graminis* Racib., hyphomycetous, sexual morph unknown, pantropical, saprobes, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Benjaminiomycetes I.I. Tav. 2000 (= *Benjaminella* I.I. Tav. 1981 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *B. melanophthalmae* (Thaxt.) I.I. Tav., asexual morph unknown, parasitic on Cryptophagidae, Latridiidae, Monotomidae, and Phalacridae beetles, terrestrial, North and South America, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Benjalia Subram. & Bhat 1989, *Ascomycota* genera *incertae sedis*, one species, type: *B. sundara* Subram. & Bhat, hyphomycetous, sexual morph unknown, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Berkelella (Sacc.) Sacc. 1891 (= *Byssostilbe* Petch 1912; = *Hypomyces* subgen. *Berkelella* Sacc. 1883 *fide* Species Fungorum 2017)), *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *B. caledonica* (Pat.) Sacc., asexual morph unknown, on Myxomycetes, worldwide, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Berkleasmium Zobel 1854, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, 36 species, type: *B. concinnum* (Berk.) S. Hughes, sexual morph unknown, saprobes, terrestrial and aquatic, see McKenzie (2008; new species), Zhang et al. (2009a; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Whitton et al. (2012a; key to 34 *Berkleasmium* species), Hüseyin et al. (2014; new species), Qu et al. (2014; new species), Tibpromma et al. (2017b; new species), four species have sequences.

Bertia De Not. 1844 (= *Sphaeria a Bertiae* (De Not.) Fr. 1849 *fide* Species Fungorum 2017), *Bertiaceae*, *Coronophorales*, *Sordariomycetes*, 30 species, type: *B. moriformis*, worldwide, see Lumbsch and Huhndorf (2010; outline), Mugambi and Huhndorf (2010; new species), Kirk et al. (2013; genus accepted),

Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Bertiella (Sacc.) Sacc. & P. Syd. 1899, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *B. polyspora* Kirschst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013, 2016; accepted as a genus in *Melanommataceae*, new species), Wijayawardene et al. (2014c; outline), Tian et al. (2015; accepted as a genus in *Melanommataceae*), Almeida et al. (2017; new species), cultures and sequences are available.

Bettisia Skou 1972, *Ascosphaeraceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *B. alvei* (Betts) Skou, asexual morph hyphomycetous, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pitt et al. (2013; new species), sequences and culture available.

Beverwykella Tubaki., 1975, *Pleomassariaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *B. pulmonaria* (Beverw.) Tubaki, sexual morph unknown, aquatic, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c; outline, accepted as a genus in *Melanommataceae*), Hyde et al. (2013; accepted as a genus in *Melanommataceae*), Kirk et al. (2013; genus accepted), Tian et al. (2015; phylogeny, transferred to *Pleomassariaceae*), Wijayawardene et al. (2017; outline), cultures and sequences are available for authentic strains CBS 283.53, MFUCC 15–0183, reference specimen MFLU 15–1398.

Bezerrromyces J.D.P. Bezerra, C.M. Souza-Motta & Crous 2016, *Bezerrromycetaceae*, *Tubeufiales*, two species, type: *B. brasiliensis* J.D.P. Bezerra, C.M. Souza-Motta & Crous, Bezerra et al. (2017a; taxonomy, phylogeny), Liu et al. (2017; treated as *Tubeufiales*), cultures and sequences are available.

Bhadradriella Nagaraju, Kunwar & Manohar. 2011, *Ascomycota* genera *incertae sedis*, one species, type: *B. hyalina* Nagaraju, Kunwar & Manohar., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Nagaraju et al. (2011b; taxonomy), Wijayawardene et al. (2017; outline), holotype: HCIO 50143, cultures and sequences are unavailable.

Bhadradriomyces Sureshk., Manohar. & Kunwar 2009, *Ascomycota* genera *incertae sedis*, one species, type: *B. elegans* Sureshk., Manohar. & Kunwar, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Suresh Kumar et al. (2009; taxonomy), Wijayawardene et al. (2017; outline), holotype: HCIO 43002, cultures and sequences are unavailable.

Bharatheeya D’Souza & Bhat 2002, *Ascomycota* genera *incertae sedis*, one species, type: *B. goanensis* (Bhat & W.B. Kendr.) D’Souza & Bhat, hyphomycetous, sexual

morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: DAOM 214612, cultures and sequences are unavailable.

Bhatia W.A. Baker & Morgan-Jones 2009, *Ascomycota* genera *incertae sedis*, one species, type: *B. malabarica* (Subram. & Bhat) W.A. Baker & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Biappendiculispora Thambug., Kaz. Tanaka & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *B. japonica* Thambug., Wanas., Kaz. Tanaka & K.D. Hyde, asexual morph unknown, saprobes on *Poaceae* spp., terrestrial, river bank, Japan, see Thambugala et al. (2015b; taxonomy, phylogeny), holotype of type: HHUF 27313, ex-type living culture of type: MAFF 239452.

Biatora Fr. 1817, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, 42 species, type: *B. vernalis* (L.) Fr., usually with ascocarps and often pycnidia, some species sorediate, lichenized (green algae), terrestrial (various substrata), probably cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Printzen (2014; DNA), Rodriguez Flakus and Printzen (2014; DNA), sequences are available.

Biatorella De Not. 1846, *Biatorellaceae*, *Lecanorales*, *Lecanoromycetes*, 30 species, type: *B. rousselii* (Durieu & Mont.) De Not., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Biatoridium J. Lahm ex Körb. 1860 (= *Chilospora* A. Massal. 1860 [1859-1860] *fide* Species Fungorum 2017), *Lichenomyces* genera *incertae sedis*, two species, type: *B. monasteriense* J. Lahm ex Körb., with ascocarps only, lichenized (green algae), terrestrial (corticicolous), Northern Hemisphere, see Lumbsch and Huhndorf (2010; outline), Miądlikowska et al. (2014a; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Biatriospore K.D. Hyde & Borse 1986, *Biatriosporaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *B. marina* K.D. Hyde & Borse, asexual morph coelomycetous, on mangroves, human mycetoma, Seychelles, Venezuela, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; introduced *Biatriosporaceae*), Kirk et al. (2013; genus accepted), Ahmed et al. (2014a; new species, asexual morph), Wijayawardene et al. (2014c; outline), Tibpromma et al. (2017b; placement), sequences and culture are suspect.

Bibanasiella R.F. Castañeda & W.B. Kendr. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *B. ramispora* R.F. Castañeda & W.B. Kendr.,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: INIFAT C91/137, cultures and sequences are unavailable, compare with *Catenosynnema* and *Speiropsis*.

Biciliopsis Diederich 1997, *Dothideomycetes* genera *incertae sedis*, one species, type: *B. leptogiicola* Diederich, asexual morph unknown, lichenicolous, non-lichenized, terrestrial, Papua New Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Biciliiospora Petr. 1952, *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *B. velutina* Petr., asexual morph unknown, saprobes, see Kirk et al. (2008; treated as a synonym of *Nitschkiella*), Mugambi and Huhndorf (2010; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Biciliiosporina Subram. & Sekar 1993, *Nitschkiaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *B. kwarensis* Subram. & Sekar, asexual unknown, saprobe, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: herb. FSI [sic] 3189; isotype IMI 32468, cultures and sequences are unavailable.

Bicoloromyces Heuchert, U. Braun & D. Hawksw. 2015, *Ascomycota* genera *incertae sedis*, one species, type: *B. kyffinensis* Heuchert, U. Braun & D. Hawksw., sexual morph unknown, on sterile lichen on rock, Antarctica, see Heuchert et al. (2015; taxonomy), holotype: MAF 19121, cultures and sequences are unavailable.

Biconiosporella Schaumann 1972, *Sordariomycetes* genera *incertae sedis*, one species, type: *B. corniculata* Schaumann, asexual morph unknown, saprobes, marine, Germany, see Jones et al. (2009b, 2015; classification), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Bicornispore Checa, Barrasa, M.N. Blanco & A.T. Martínez 1996, *Rutstroemiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *B. exophiala* Checa, Barrasa, M.N. Blanco & A.T. Martínez, saprobes, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Coryneliaceae*), Kirk et al. (2013; genus accepted), Galán et al. (2015; phylogeny, new species), cultures and sequences are available but lacking for the type.

Bicrouania Kohlm. & Volkmar-Kohlm. 1990, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, one

species, type: *B. maritima* (P. Crouan & H. Crouan) Kohlm. & Volkmar-Kohlm., asexual morph unknown, saprobes, marine aquatic, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Tian et al. (2015; morphology, phylogeny), cultures and sequences are unavailable.

Bifidocarpus Cano, Guarro & R.F. Castañeda 1994, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *B. cubensis* Cano, Guarro & R.F. Castañeda, saprobes, terrestrial, Cuba, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype: IMI 356298, cultures and sequences are unavailable, compare with *Renispora* and *Onygena*.

Biflagellospora Matsush. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *B. japonica* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Biflagellosporella Matsush. 1993, *Ascomycota* genera *incertae sedis*, one species, type: *B. amazonensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Biflua Jørgen Koch & E.B.G. Jones 1989, *Ascomycota* genera *incertae sedis*, one species, type: *B. physasca* Jørg. Koch & Jones, asexual morph unknown, on wood associated with sand, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), cultures and sequences are unavailable.

Bifrontia Norman 1872, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, three species, type: *B. compactior* Norman, saprobes on decaying wood, Europe, see Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; taxonomy), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Bifusella Höhn. 1917, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, eleven species, type: *B. linearis* (Peck) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016b; new combination), cultures and sequences are available (unpublished).

Bifusepta Darker 1963, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *B. tehonii* Darker, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bilimbia De Not. 1846, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *B. hexamera* De Not., lichenized, see Ekman et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Printzen (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Bimeris Petr. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *B. aperta* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Ecuador, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bimuria D. Hawksw., Chea & Sheridan 1979, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, type: *B. novae-zelandiae* D. Hawksw., Chea & Sheridan, one species, asexual morph unknown, isolated from soil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014e; taxonomy), Wijayawardene et al. (2014c; outline), holotype: K(M)-IMI 229814, cultures and sequences are available.

Bioconiosporium Bat. & J.L. Bezerra 1964, *Ascomycota* genera *incertae sedis*, two species, type: *B. baccharidis* Bat. & J.L. Bezerra, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Biophomopsis Petr. 1931, *Ascomycota* genera *incertae sedis*, three species, type: *B. brenesii* Petr., coelomycetous, sexual morph unknown, West Indies, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Bioscypha Syd. 1927, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *B. cyatheae* Syd., asexual morph hyphomycetous, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequence are unavailable.

Biostictis Petr. 1950, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *B. rubiacearum* (Pat.) Petr., with ascospores only, terrestrial (parasitic on leaves), tropical America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Bipolaris Shoemaker 1959, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, 121 species, type: *B. maydis* (Y. Nisik. & C. Miyake) Shoemaker, sexual morph *Cochliobolus*, plant and animal pathogen, saprobes, terrestrial, worldwide, see Aggarwal et al.

(2008, 2009, 2010, 2011a, b; intraspecific variations, DNA fingerprinting, pathogen histopathology), Antoniazzi et al. (2008; xanthan and allicin as elicitors), Chand et al. (2008, 2014; cell biology, disease resistance screening), de Moura Nascimento and Van Der Sand (2008; restriction analysis), Durkin et al. (2008; phaeohyphomycotic keratitis), Dyer et al. (2008; bronchopulmonary mycosis), Felle et al. (2008; host pathogen interaction), Geng et al. (2008; *Echinochloa* control), Ghazvini and Tekauz (2008; Host-pathogen interaction), Janni et al. (2008; resistance in wheat), Jiang and Zhang (2008; new species), Kihara et al. (2008, 2009; BMR1 gene characterization), Kobayashi et al. (2008; Disseminated infection), Motlagh and Kaviani (2008; Iran), Pandey et al. (2008; clonal variability), Persson et al. (2008, 2009; mutant host resistance), Shabana et al. (2008; pathogen control), Asad et al. (2009; Pakistan), Bashir et al. (2009; keratomycosis, endophthalmitis), El Mhadri et al. (2009; *Citrullus lanatus* host), Iftikhar et al. (2009a, b; synthetic host screening), Kamal and Mia (2009; DNA fingerprinting), Kumar et al. (2009; host resistance), Naumann et al. (2009; host enzyme modification), Poloni et al. (2009a, b; morphology, physiology, virulence, vegetative incompatibility), Zhang and Li (2009; new species), Bashyal et al. (2010, 2011; conidiogenesis study, host resistance components), Carissimi et al. (2010; *Bacillus* sp. E164 activity), Dong et al. (2010; TaPIEP1 study), Gomathinayagam et al. (2010; *Trichoderma* effect), Han et al. (2010b; cytology), Kleczewski and Flory (2010; *Microstegium vimineum* host), Knight et al. (2010; genetic variation), Kumar et al. (2010; host resistance), Lapmak et al. (2010; L-asparaginase production), Martinez et al. (2010; Tanzania), Matusinsky et al. (2010; species specific detection), Motlagh and Anvari (2010; genetic variation), Saha et al. (2010; Ag nanoparticle production), Shali et al. (2010; SG2 chitinases inhibition), Tomaso-Peterson and Balbalian (2010; Mississippi), Acharya et al. (2011; review), Ahmadpour et al. (2011, 2012; new species), Baturo-Ciesniewska (2011; Poland), Ben-Ze'ev et al. (2011; Israel), Chowdhary et al. (2011; bronchopulmonary mycosis), Dallagnol et al. (2011a, b, 2013; host pathogen interaction), Giri et al. (2011; phaeohyphomycosis in dog), Ishihara et al. (2011; defense response), Jiao et al. (2011; MYB73 analysis), Kawahigashi et al. (2011; resistance gene cloning), Kumar et al. (2011; molecular and pathological characterization), Moya-Elizondo et al. (2011; population dynamics), Rosow et al. (2011; Cerebral phaeohyphomycosis), Vu et al. (2011, 2015; on switchgrass), Arunpanichlert et al. (2012; metabolite profiling), Da Cunha et al. (2012; clinical diversity), Manamgoda et al. (2012; phylogeny), Mizuno et al. (2012; host pathogen interaction), Nizam et al. (2012; transformation), Zhang et al. (2012d, k; host resistance,

from soil), Almaguer et al. (2013; conidial germination), Burgos et al. (2013; Philippines), Eisa et al. (2013; resistance screening), Gupta et al. (2013; fungicidal management), Rossman et al. (2013a; name conservation), Tarnowski et al. (2013; South Florida), Wang et al. (2013c; metabolite as drug), Yazawa et al. (2013; transcriptome analysis), Amaradasa and Amundsen (2014; Nebraska), Bencsik et al. (2014; Ophiobolin A), Manamgoda et al. (2014; review), Meena et al. (2014; genetic evaluation of host), Minckler et al. (2014; *Bipolaris endophthalmitis*), Mishra et al. (2014b; Ag nanoparticle fungicide), Small et al. (2014; endophthalmitis), Teran et al. (2014; fatal infection), Ariyawansa et al. (2015c; phylogeny), Eghtedari (2015; conjunctiva cyst), Fatima et al. (2015; nanoparticle synthesis), Kuan et al. (2015; phylogeny, genomics), Waxman and Bergstrom (2015; New York), Kekuda et al. (2016; antifungal activity), sequences are available, paratype and ex-paratype of type species CBS 136.29.

Biporispora J.D. Rogers, Y.M. Ju & Cand. 1999, *Xylariales* genera *incertae sedis*, one species, type: *B. europaea* J.D. Rogers, Y.M. Ju & Cand., asexual morph unknown, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; placed in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Bisbyopeltis Bat. & A.F. Vital 1957, *Ascomycota* genera *incertae sedis*, one species, type: *B. phoebe* Bat. & A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Biscogniauxia Kuntze 1891 (= *Albocrustum* Lloyd 1925; = *Kommamyce* Nieuwl., Am. Midl. 1916; = *Nummularia* Tul. & C. Tul 1863; = *Nummulariella* Eckblad & Granmo 1978; = *Numulariola* House 1923 *fide* Wendt et al. 2017), *Graphostromataceae*, *Xylariales*, *Sordariomycetes*, 33 species, type: *B. nummularia* (Bull.) Kuntze, asexual morph unknown, saprobes, terrestrial, France, see Mugambi et al. (2009; new species, Kenya), Lumbsch and Huhndorf (2010; outline), Vasilyeva and Stephenson (2010; new species), Vasilyeva et al. (2012; new species, Thailand), Kirk et al. (2013; genus accepted), Liu et al. (2015a; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; transferred to *Graphostromataceae*), cultures and sequences are available.

Bisifusarium L. Lombard, Crous & W. Gams 2015, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, seven species, type: *B. dimerum* (Penz.) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lombard et al. (2015b;

taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type of the type: CBS 108944.

Bispora Corda 1837, *Ascomycota* genera *incertae sedis*, 31 species, type: *B. monilioides* Corda, hyphomycetous, phialophora-like synanamorph, sexual morph unknown, meristematic black fungi, saprobes on wood and leaves, widespread (temperate), producer of various enzymes possessing industrial applications, see Luo et al. (2009a, b; enzymes), Wang et al. (2010a; enzyme), Dubey and Pandey (2011; new species), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Xu et al. (2013; enzymes), Hua et al. (2014; enzyme), Liu et al. (2015b; enzyme), molecular data not available from the type materials.

Bisporella Sacc. 1884 (= *Bispora* Fuckel 1870 [1869-70]; = *Calycella* Quél. 1886 *fide* Species Fungorum 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, 19 species, type: *B. monilifera* (Fuckel) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequence are available but lacking for the type species.

Bisporostibella Brandsb. & E.F. Morris 1971, *Ascomycota* genera *incertae sedis*, one species, type: *B. fusca* Brandsb. & E.F. Morris, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bisseomyces R.F. Castañeda, 1985, *Ascomycota* genera *incertae sedis*, one species, type: *B. havanensis* R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: C84/90-2, cultures and sequences are unavailable, compare with *Neojohnstonia* and *Parapithomyces*.

Bivallum P.R. Johnst. 1991, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, seven species, type: *B. zelandicum* P.R. Johnst., asexual morph unknown, saprobes, terrestrial, Australasia, Chile, Panama, see Hou and Piepenbring (2009a; new species, Panama), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blasdalea Sacc. & P. Syd., 1902 (= *Singeriella* Petr. 1959; = *Chrysogloeum* Petr. 1959 *fide* Wijayawardene et al. 2014c; Species Fungorum 2017), *Vizellaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *B. disciformis* (Rehm) Sacc. & P. Syd., asexual morph coelomycetous, ring-like or linear loculus arrangement, associated with leaf spots on leaves of *Escallonia vaccinoides*, Brazil, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013;

morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Rossman et al. (2016a; accept the genus status), cultures and sequences are unavailable.

Blastacervulus H.J. Swart 1988, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, one species, type: *B. eucalypti* H.J. Swart, coelomycetous, sexual morph unknown, pathogens, terrestrial, Australia, see Cheewangkoon et al. (2009; taxonomy, phylogeny), Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy, accepted as a genus in *Planistromellaceae*), Kirk et al. (2013; genus accepted), Giraldo et al. (2017; phylogeny), cultures available for the type: CBS124759.

Blastenia A. Massal. 1852, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, eleven species, type: *B. ferruginea* (Huds.) A. Massal., lichenized, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Blasticomyces I.I. Tav. 1985, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *B. lispi* (Thaxt.) I.I. Tav., asexual morph unknown, parasitic on Hydrophilidae: Sphaeridiinae and Staphylinidae beetles, asexual morph unknown, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters and Yaakop (2014; Borneo), cultures and sequences are unavailable.

Blastobotrys Klopotek 1967 (= *Sympodiomyces* Fell & Statzell 1971; = *Trichomonascus* H.S. Jackson emend Kurtzman & Robnett 2007; = *Stephanoascus* M.T. Sm., Van der Walt & Johannsen 1976 *fide* Species Fungorum 2017), *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, 30 species, type: *B. nivea* Klopotek, asexual reproduction by budding (occasionally) or hyphomycetous, primarily saprophytic, some species cause human infections, insect frass from trees, growth on mushrooms, moss, ensiled maize, soil, air, sea water, bat liver, human infections, worldwide, Kirk et al. (2013; genus accepted), Crous et al. (2016b; new species), cultures and sequences are available.

Blastocatena Subram. & Bhat 1989, *Ascomycota* genera *incertae sedis*, one species, type: *B. pulneyensis* Subram. & Bhat, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blastodesmia A. Massal. 1852, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, one species, type: *B. nitida* A. Massal., asexual morph unknown, lichenized, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), holotype US National Herbarium barcode

01138889 – on bark of *Fraxini*, Austria (Schuler J), cultures and sequences are unavailable.

Blastodictys M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, one species, type: *B. hibisci* (Hansf.) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blastofusariooides Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *B. fusca* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blastomyces Gilchrist & W.R. Stokes 1898 (= *Gilchristia* Cif. & Redaelli 1934; = *Ajellomyces* McDonough & A.L Lewis 1968; = *Zymonema* Beurm. & Gougerot 1909) *Ajellomycetaceae*, *Onygenales*, *Eurotiomycetes*, two species, (but phylogenetic studies suggest the existence of more taxa *fide* de Hoog et al. 2016b), type: *B. dermatitidis* Gilchrist & W.R. Stokes, terrestrial, pathogenic to humans, canids, felines and other animals, also reported from soil, Africa, India, North America, *Blastomyces* Gilchrist & W.R. Stokes is an illegitimate genus (Art. 53.1) but, due to its wide use in medical literature, most modern authors prefer this name instead of any of its synonyms (Seifert et al. 2011), and a proposal of conservation against an earlier homonym, *Blastomyces* Costantin & Rolland, has been published (de Hoog et al. 2016b), see Saccante and Woods (2010; clinical and laboratory aspects), Brown et al. (2013; phylogeny, new species), Kirk et al. (2013; genus accepted), ex-epitype strain of the type species =CBS 674.68, sequences are available.

Blastophorella Boedijn 1937, *Ascomycota* genera *incertae sedis*, one species, type: *B. smithii* Boedijn, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South East Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blastophorum Matsush. 1971, *Reticulascaceae*, *Glomerellales*, *Sordariomycetes*, five species, type: *B. truncatum* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, China, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2016; new species, China), cultures and sequences are unavailable.

Blastophragma Subram. 1995, *Ascomycota* genera *incertae sedis*, four species, type: *B. subulatum* Subram.,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Gao et al. (2015a; new species), Xia et al. (2016a; new species), holotype: MUBL 3128, cultures and sequences are unavailable.

Blennoria Moug. & Fr. 1825, *Ascomycota* genera *incertae sedis*, four species, type: *B. buxi* Fr., coelomycetous, sexual morph unknown, saprobes and plant endophytes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blennoriopsis Petr. 1920, *Ascomycota* genera *incertae sedis*, one species, type: *B. moravica* Petr., coelomycetous, sexual morph unknown, rarely reported and in case as saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blennothallia Trevis. 1853, *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, four species, type: need typification, lichenized, see Otálora et al. (2013, 2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Bleptosporium Steyaert 1961 (= *Monochaetina* Subram. 1961 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, four species, type: *B. montteae* Speg. ex Steyaert, coelomycetous, sexual morph unknown, saprobes, terrestrial, Argentina, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Blodgettia Harv. 1858 (= *Blodgettia* E.P. Wright 1881; = *Blodgettomyces* Feldmann 1939 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *B. confervoides* Harv., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bloxamia Berk. & Broome 1854 (= *Endosporostilbe* Subram. 1958; = *Thecostroma* Clem. 1909; = *Trichotheca* P. Karst. 1887 *fide* Species Fungorum 2017), *Bloxamiaceae*, *Helotiales*, *Leotiomycetes*, 19 species, type: *B. truncata* Berk. & Broome, hyphomycetous, sexual morph unknown, saprobes, terrestrial, fresh water, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Spooren (2014; new species), Guatimosim et al. (2016b; new species, DNA, phylogeny), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), cultures and sequences are available but lacking for the type species.

Blumeria Golovin ex Speer 1974 (= *Oidium* Link 1824 *fide* Braun and Cook 2012), *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, one species, type: *B. graminis* (DC.) Speer, asexual morphs hyphomycetous, biotrophic, obligate plant pathogens, terrestrial, worldwide, see Braun and Cook (2012; taxonomy), neotype: Germany, Bonn, Botanical garden Poppelsdorf, on *Triticum aestivum*, 25 June 1869, Körnicke, in herb. de Candolle (G 00122110), sequences are available.

Blumeriella Arx 1961 (= *Higginsia* Nannf. 1932; = *Microgloeum* Petr. 1922; = *Phloeosporella* Höhn. 1924 *fide* Johnston et al. (2014b); Species Fungorum 2017), *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *B. jaapii* (Rehm) Arx, asexual morphs *Microgloeum*, *Phloeosporella*, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available.

Boeremia Aveskamp, Gruyter & Verkley 2010, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, eleven species, type: *B. exigua* (Desm.) Aveskamp, Gruyter & Verkley, asexual morph coelomycetous, saprobes or plant pathogenic, widespread, see Aveskamp et al. (2010; taxonomy, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Chen et al. (2015b; phylogeny), ex-type: CBS 101207 (*Boeremia exigua* var. *pseudolilacis*), CBS 443.94 (*Boeremia exigua* var. *heteromorpha*), CBS 100167 (*Boeremia exigua* var. *populi*).

Boerlagiomyces Butzin 1977 (= *Boerlagella* Penz. & Sacc. 1897; = *Garethjonesia* K.D. Hyde 1992 *fide* Species Fungorum 2017), *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, c. nine species, type: *B. velutinus* (Penz. & Sacc.) Butzin, asexual morph coelomycetous, saprobes, aquatic, Australia, India, Indonesia, Philippines and USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Boonmee et al. (2011, 2014b); Wijayawardene et al. (2014c; outline), Doilom et al. (2017, description), sequences are available.

Bogoriella Zahlbr. 1928 (= *Bogoriellomyces* Cif. & Tomas. 1954 *fide* Species Fungorum 2017), *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, 29 species, type: *B. subpersicina* Zahlbr., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Nelsen et al. (2014b; phylogeny), Aptroot and Lücking (2016; taxonomy), Lücking et al. (2017a; phylogeny), sequences are available.

Boliniella (Nitschke) Sacc. 1882 (= *Naemaspora* Willd. 1849 *fide* Species Fungorum 2017), *Boliniales*, *Sordariomycetes*, three species, type: *B. tubulina* (Alb. & Schwein.) Sacc., asexual morph unknown, saprobes, cosmopolitan, see Kirk et al. (2008; treated as a synonym of *Camarops* P. Karst.), Lumbsch and

Huhndorf (2010; did not list), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a distinct genus), cultures and sequences are unavailable.

Bombardia (Fr.) P. Karst. 1873, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *B. bombarda* (Batsch) J. Schröt., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariomycetes*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Bombardiastrum Pat. 1893, *Sordariomycetes* genera *incertae sedis*, one species, type: *B. andinum* Pat., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Bombardiella Höhn. 1909, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *B. caespitosa* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Bombardioidea C. Moreau ex N. Lundq. 1972 (= *Bombardioidea* C. Moreau 1954 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, four species, type: *B. bombardoides* (Auersw.) C. Moreau ex C. Moreau, asexual morph unknown, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Bommerella Marchal 1885, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *B. trigonospora* Marchal, asexual morph scopulariopsis-like, saprobes, from soil, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), type species have been transferred to *Chaetomium* (*fide* Index Fungorum 2017).

Bonansea Sacc. 1906, *Rhytismatales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *B. mexicana* Sacc., asexual morph unknown, saprobes, terrestrial, Mexico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bonaria Bat. 1959, *Dothideomycetes* genera *incertae sedis*, one species, type: *B. lithocarpi* (V.A.M. Mill. & Bonar) Bat., asexual morph unknown, saprobes, terrestrial, Mexico, see Lumbsch and Huhndorf (2010; outline,

accepted as a genus in *Micropeltidaceae*), Hyde et al. (2013) did not accept as a genus in *Micropeltidaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Bondiella Piroz. 1972, *Mesnieraceae, Dothideomycetes* families *incertae sedis*, one species, type: *B. palmicola* Piroz., asexual morph unknown, saprobes, terrestrial, Tanzania, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Boothiella Lodhi & Mirza 1962 (= *Thielaviella* Arx & T. Mahmood 1968 *fide Species Fungorum* 2017), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *B. tetraspora* Lodhi & J.H. Mirza, asexual morph unknown, saprobes, from soil, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Boreoplaca Timdal 1994, *Ophioparmaceae*, *Umbilicariales*, *Lecanoromycetes*, one species; type: *B. ultrafrigida* Timdal, lichenized, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; phylogeny), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; taxonomy), Lücking et al. (2017a; classification), holotype: O-L-127, sequences are available.

Borinquenotrema Mercado-Díaz, Lücking & Parnmen 2014, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *B. sorediocarpum* Mercado-Díaz, Lücking & Parnmen, lichenized, terrestrial, see Lumbsch et al. (2014a, b; taxonomy), Lücking et al. (2017a; classification), holotype: UPR, Mercado-Díaz 883, sequences are available.

Botanamphora Nograsek & Scheuer 1990, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *B. pachycarpa* (Sacc. & Marchal) Nograsek & Scheuer, asexual morph unknown, saprobes, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Botryandomyces I.I. Tav. & T. Majewski 1976, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *B. heteroceri* (Maire) I.I. Tav. & T. Majewski, asexual morph unknown, parasitic on Heteroceridae beetles, semi-aquatic, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence is available but lacking for the type species.

Botryobambusa Phook., J.K. Liu & K.D. Hyde 2012, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, one species, type: *B. fusicoccum* Phook., J.K. Liu & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Thailand, see Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Phillips et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline), holotype and ex-type: MFLU 11-0179, MFLUCC 11-0143.

Botryochora Torrend 1914, *Dothideomycetes* genera *incertae sedis*, one species, type: *B. nigra* (Torrend) Torrend, asexual morph unknown, saprobes, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothioraceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Botryoderma Papendorf & H.P. Upadhyay 1969, *Ascomycota* genera *incertae sedis*, four species, type: *B. lateritium* Papendorf & H.P. Upadhyay, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Botryodiplodina Dias & Sousa da Câmara 1954, *Ascomycota* genera *incertae sedis*, one species, type: *B. mori* Dias & Sousa da Câmara, coelomycetous, sexual morph unknown, saprobes, terrestrial, Portugal, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Botryohypoxylon Samuels & J.D. Rogers 1986 (= *Iledon* Samuels & J.D. Rogers 1986 *fide* Wijayawardene et al. 2014c), *Dothideomycetes* genera *incertae sedis*, one species, type: *B. nigra* (Torrend) Torrend, asexual morph coelomycetous, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Rossman et al. (2015b; nomenclature), cultures and sequences are unavailable.

Botryola Bat. & J.L. Bezerra 1964, *Nitschiaceae*, *Coronophorales*, *Sordariomycetes*, six species, type: *B. tetrasperma* Bat. & J.L. Bezerra, asexual morph unknown, saprobes, lichenicolous, terrestrial, saprobes, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Botryolepraria Canals, Hern.-Mar., Gómez-Bolea & Llimona 1997, *Verrucariales* genera *incertae sedis*, *Eurotiomycetes*, two species, type: *B. lesdainii* (Hue) Canals, Hern.-Mar., Gómez-Bolea & Llimona, lichenized, worldwide, see Kukwa and Pérez-Ortega (2010; taxonomy), Lumbsch and Huhndorf (2010; outline,

accepted as a genus in *Lecanoromycetes*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Botryomonilia Goos & Piroz. 1975, *Ascomycota* genera *incertae sedis*, four species, type: *B. scheeleae* Goos & Piroz., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Central America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Botryosphaeria Ces. & De Not. 1863 (= *Fusicoccum* Corda 1829; = *Dichomera* Cooke 1878 *fide* Wijayawardene et al. 2014c, 2016b; Rossman et al. 2015b; Art. 59), *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, eleven species, type: *B. dothidea* (Moug. Fr.) Ces. & De Not., asexual morph coelomycetous, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Liu et al. (2012b; accepted as a genus in *Botryosphaeriaceae*), Abdollahzadeh et al. (2013; species in Iran), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Phillips et al. (2013; taxonomy, phylogeny), Slippers et al. (2013; phylogeny), Yan et al. (2013; species associated with grape vine), Wijayawardene et al. (2014c, 2016b; outline, taxonomy), Xu et al. (2015, apple ring rot), Ariyawansa et al. (2016a; new species), Dissanayake et al. (2016b; taxonomy), cultures and sequences are available.

Botryosporium Corda 1831 (= *Peylia* Opiz 1857; *Radaiella* Bainier 1910 *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, seven species, type: *B. diffusum* (Grev.) Corda, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, cosmopolitan, see Campbell et al. (2007; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Choi et al. (2014; pathogens, DNA), cultures and sequences are available but lacking for the type.

Botryostroma Höhn. 1911 (= *Parodiellina* Viégas 1944; = *Robledia* Chardón 1929 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *B. inaequale* (G. Winter) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, America (tropical) see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Botryozyma Shann & M.Th. Smith emend. Lachance & Kurtzman 2013, *Trigonopsidaceae*, *Saccharomycetales*, *Saccharomycetes*, four species, type: *B. nematodophila* Shann & M.T. Smith, asexual reproduction by budding (multilateral), ?saprophytic, from Panagrellus nematodes or associated with *Drosophila* flies, north western USA, Italy, see Kirk et al. (2013; genus accepted), Lachance and

Kurtzman (2013; re-visited, generic revision), cultures and sequences are available.

Botryotis P. Micheli ex Pers. 1794 (= *Botryotinia* Whetzel 1945 *fide* Johnston et al. 2014b; see Species Fungorum 2017 for synonyms), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *B. cinerea* Pers., asexual morph hyphomycetous, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available.

Boubovia Svrček 1977, *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, five species, type: *B. luteola* (Velen.) Svrček, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Boudiera Cooke 1877 (= *Svrcekia* Kubička 1960 *fide* Species Fungorum 2017), *Pezizaceae*, *Pezizales*, *Pezizomycetes*, ten species, type: *B. areolata* Cooke & W. Phillips, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Boudierella Sacc. 1895, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *B. cana* Marchal & É.J. Marchal, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Bouvetiella Øvstedal 1986, *Lecanoromycetes* genera *incertae sedis*, one species; type: *B. pallida* Øvstedal, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Brachiosphaera Nawawi 1976, *Aliquandostipitaceae*, *Jahnulales*, *Dothideomycetes*, two species, type: *B. tropicalis* Nawawi, isolated from river foam: Peninsular Malaysia, see Seifert et al. (2011; morphology), Suetrong et al. (2011a; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sexual morph unknown, cultures and sequences are available for authentic strains SS2523, E192–1.

Brachyalara Réblová & W. Gams 2011, *Helotiales*, genera *incertae sedis*, *Leotiomycetes*, one species, type: *B. straminea* Réblová & W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Réblová and Gams (2011; morphology, phylogeny), Wijayawardene et al. (2012, 2017; outline), ex-type of the type: CBS 175.74.

Brachycephala J.S. Monteiro, Gusmão & R.F. Castañeda 2015, *Ascomycota* genera *incertae sedis*, one

species, type: *B. exotica* J.S. Monteiro, Gusmaño & Castañeda-Ruiz, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Monteiro et al. (2015b; morphology), holotype of the type: HUEFS 210446, cultures and sequences are unavailable, compare with *Acarocybellina*, *Brachysporiella* and *Kramabeeja*.

Brachyconidiella R.F. Castañeda & W.B. Kendr. 1990 *Ascomycota* genera *incertae sedis*, *Dothidiomycetes*, one species, type: *B. monilispora* R.F. Castañeda & W.B. Kendr., sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: INIFAT C88/300, sequences are available, compare with *Brachyconidiellopsis* and *Matsushimaea*.

Brachyconidiellopsis Decock, R.F. Castañeda & Adhikari 2004, *Microascaceae*, *Microascales*, *Sordariomycetes*, two species, type: *B. fimicola* Decock, R.F. Castañeda & Adhikari, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharanachchikumbura et al. (2015a, b; outline, phylogeny), holotype: MUCL 44677, cultures and sequences are available.

Brachydesmiella G. Arnaud ex S. Hughes 1961, *Ascomycota* genera *incertae sedis*, eight species, type: *B. biseptata* G. Arnaud ex S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Brachyphoris Juan Chen, L.L. Xu, B. Liu & Xing Z. Liu 2007, *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, five species, type: *B. oviparasitica* (G.R. Stirling & Mankau) Juan Chen, L.L. Xu, B. Liu & Xing Z. Liu, saprobes, on nematode eggs, on soil, terrestrial, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Brachysporiella Bat. 1952 (= *Edmundmasonia* Subram. 1958; = *Monosporella* S. Hughes 1953; = *Monotosporella* S. Hughes 1958; = *Sporidesmiopsis* Subram. & Bhat 1989 [1987] *fide Species Fungorum* 2017), *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, 16 species, type: *B. gayana* Bat., hyphomycetous, sexual morph unknown, saprobes, widespread, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2017; introduced *B. navarrica* Hern.-Restr. et al. and accepted as *Kirschsteiniotheliales*, *incertae sedis*), sequences are available.

Brachysporiellina Subram. & Bhat 1989, *Ascomycota* genera *incertae sedis*, two species, type: *B. pulneyensis* Subram. & Bhat, hyphomycetous, sexual morph unknown,

saprobes, terrestrial, Asia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Brachysporiopsis Yanna, W.H. Ho & K.D. Hyde 2004, *Ascomycota* genera *incertae sedis*, one species, type: *B. chinensis* Yanna, W.H. Ho & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Brachysporium Sacc. 1886 (= *Cryptadelphia* Réblová & Seifert 2004 *fide* Réblová et al. 2016c), *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, twelve species, type: *B. obovatum* Keissl., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharanachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures and sequences are available.

Bradymyces Hubka, Réblová, Selbmann & M. Kolařík 2014, *Trichomeriaceae*, *Chaetothyriales*, *Eurotiomycetes*, three species, type: *B. oncorhynchi* Hubka, Řehulková, & M. Kolařík, sexual morph unknown, cold-blooded, water vertebrates opportunist, rock-inhabiting, see Hubka et al. (2014b; taxonomy, phylogeny), Réblová et al. (2016b; new species), holotype and ex-type of the type: PRM 861507 CBS 133066 = CCF 4369 = CCFFEE 6134.

Bramhamyces Hosag. 2009, *Dothideomycetes* genera *incertae sedis*, one species, type: *B. ilicis* Hosag. & Chandrapr, sexual morph unknown, epiphytes, terrestrial, India, see Hosagoudar and Chandraprabha (2009a; protologue, accepted as an anamorphic genus in *Asterinaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes* genera *incertae sedis*), holotype of the type: HCIO 48300, cultures and sequences are unavailable.

Brasilicia Lücking, Kalb & Sérus. 2008, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *B. brasiliensis* (Müll. Arg.) Lücking, Kalb & Sérus., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences unavailable.

Braunomyces V.A. Melnik & Crous 2014, *Ascomycota* genera *incertae sedis*, one species, type: *B. dictyosporus* V.A. Melnik & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Vietnam, see Mel'nik and Crous (2014; taxonomy), Wijayawardene et al. (2017; outline), holotype: HAL 2606 F, cultures and sequences are unavailable.

Brefeldiella Speg. 1889, *Trichopeltinaceae*, *Dothideomycetes* families *incertae sedis*, four species, type: *B. brasiliensis* Speg., asexual morph unknown, epiphytes, appearance as thallus on leave, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), Hongasan et al. (2014a; accepted as a genus in *Trichopeltinaceae*), Wijayawardene et al. (2014c; outline), need recollecting and sequencing to verify generic placement.

Brefeldiopycnis Petr. & Cif. 1932, *Ascomycota* genera *incertae sedis*, one species, type: *B. membranacea* Petr. & Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Brefeldochium Verkley 2005, *Helotiales* genera *incertae sedis* *Leotiomycetes*, one species, type: *B. pruinatum* Verkley, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline, taxonomy), cultures available for the type: CBS 160.92.

Brenesiella Syd. 1929, *Sordariomycetes*, genera *incertae sedis*, one species, type: *B. erythroxyli* Syd., asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Brettanomyces Kufferath & van Laer 1921, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, six species, type: *B. bruxellensis* Kufferath & van Laer, asexual reproduction is either by multilateral budding or, rarely, by bipolar budding on a narrow base, saprophytes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), cultures and sequences are available.

Breviappendix Senan. & K.D. Hyde 2017, *Sydiowilliaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *B. rubi* (Rehm) Senan., Maharachch. & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Europe, see Senanayake et al. (2017; taxonomy, phylogeny), epitype: MFLU 16–2861, cultures and sequences are available.

Brevicatenospora R.F. Castañeda, Minter & Saikawa 2006, *Ascomycota* genera *incertae sedis*, one species, type: *B. enteroproliferata* R.F. Castañeda, Minter & Saikawa, hyphomycetous, sexual morph unknown, saprobes, aquatic (fresh water), Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: INIFAT C04/111, cultures and sequences are unavailable, compare with *Annellophora*, *Acrogenospora*, *Belemnospora*, *Sporidesmiella* and *Repetophragma*.

Brevistachys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, five

species, type: *B. variabilis* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lombard et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: CBS H-22431, CBS 141057.

Brianaria S. Ekman & M. Svensson 2014, *Psoraceae*, *Lecanorales*, *Lecanoromycetes*, four species, type: *B. sylvicola* (Flot. ex Körb.) S. Ekman & M. Svensson, lichenized, terrestrial, see Ekman and Svensson (2014; taxonomy), Lücking et al. (2017a; classification), type specimen UPS-L-133138 (lectotype), sequences are available.

Briancoppinsia Diederich, Ertz, Lawrey & van den Boom 2012, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *B. cytospora* (Vouaux) Diederich, Ertz, Lawrey & van den Boom, coelomycetous, sexual morph unknown, lichenicolous, non-lichenized, terrestrial, Europe, North America, see Diederich et al. (2012b; morphology, phylogeny), Wijayawardene et al. (2017; outline), sequences are available.

Briania D.R. Reynolds 1989, *Meliolinaceae*, *Dothideomycetes*, families *incertae sedis*, one species, type: *B. fruticetum* D.R. Reynolds, sexual morph unknown, saprobes, terrestrial, Hawaiian Island, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bricookea M.E. Barr 1982, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *B. sepalorum* (Vleugel) M.E. Barr, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Kirk et al. (2013; treated as a synonym of *Lophiostoma*), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Brigantiae Trevis. 1853 (= *Myxodictyon* A. Massal. 1860 [1859–1860]; = *Myxodictyonomyces* Cif. & Tomas. 1953 fide Species Fungorum 2017), *Brigantieaceae*, *Teloschistales*, *Lecanoromycetes*, 29 species; type: *B. tricolor* (Mont.) Trevis., lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Briosia Cavara 1888, *Ascomycota* genera *incertae sedis*, six species, type: *B. ampelophaga* Cavara, hyphomycetous, plant pathogen, widespread, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Brodingnagia K.D. Hyde & P.F. Cannon 1999, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two

species, type: *B. nigeriensis* (Sivan. & Okpala) Hyde & Cannon, asexual morph unknown, pathogens, saprobes, terrestrial, Australia, Sierra Leone, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of *Brodingnagia nigeriensis* subsp. *corneri* IMI 390a.

Brodoa Goward 1987, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *B. oroorctica* (Krog) Goward, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Brookzia Hansf. 1956 (= *Capnogonium* Bat. & Peres 1961; = *Hiospira* R.T. Moore 1962; = *Overeemia* G. Arnaud 1954 [1953] *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b; Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, one species, three variety, type: *B. tropicalis* Hansf., hyphomycetous, on leaf as parasite, common in tropical regions, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; nomenclature), Rossman et al. (2015b; nomenclature), sequences are unavailable, need recollecting to verify generic placement.

Broomella Sacc. 1883 (= *Keissleria* Höhn. 1918 *fide* Species Fungorum 2017), *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, two species, type: *B. vitaliae* (Berk. & Broome) Sacc., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Li et al. (2015d; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; outline), cultures available for the type: MFLUCC 15-0023 (*fide* Li et al. 2015).

Brownliella S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, four species, lichenized, terrestrial, see Kondratyuk et al. (2013a, 2015b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Bruceomyces Rikkinen 2012 (= *Brucea* Rikkinen 2003 *fide* Species Fungorum 2017), *Lecanorales* genera *incertae sedis*, *Lecanoromycetes*, four species, type: *B. castoris* (Rikkinen) Rikkinen, lichenized, terrestrial, see Tuovila et al. (2012; taxonomy), cultures and sequences are unavailable.

Brunneiapiospora K.D. Hyde, J. Fröhl. & Joanne E. Taylor 1998, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, six species, type: *B. javensis* K.D. Hyde, J. Fröhl. & Joanne E. Taylor, asexual morphs unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Vitoria et al. (2012; new species), Crous et al. (2012c; new species, New Zealand), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, treated as *Clypeosphaeriaceae*), Hongasan et al. (2017a;

phylogeny, evolution, treated *Clypeosphaeriaceae* as a synonym of *Xylariaceae*), cultures and sequences are available for unidentified species.

Brunneiperidium Daranag., Camporesi & K.D. Hyde 2015, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *B. gracilemum* Daranag., Camporesi & K.D. Hyde, asexual morph hyphomycetous, only known for *B. gracilemum*, saprobes, terrestrial, Italy, Europe, two species, all the species have molecular data, see Daranagama et al. (2015a; taxonomy, phylogeny), Maharachchikumbura et al. (2016b; outline), ex-type: MFLUCC 14-0011; holotype MFLU 14-0229.

Brunneocarpos Giraldo & Crous 2016, *Mycocaliciaceae*, *Mycocaliciales*, *Eurotiomycetes*, one species, type: *B. banksiae* Giraldo & Crous, asexual morph unknown, saprobes, Australia, see Crous et al. (2016b; taxonomy, phylogeny), holotype and ex-type: CBS H-22633, CPC 29841.

Brunneoclavispora Phook. & K.D. Hyde 2015, *Halothiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *B. bambusae* Phook. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 11-0213, MFLUCC 11-0177.

Brunneodinemasporium Crous & R.F. Castañeda 2012, *Chaetosphaeriaceae*, *Chaetosphaerales*, *Sordariomycetes*, two species, type: *B. brasiliense* Crous & Castañeda, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Crous et al. (2012g; morphology, phylogeny), Maharachchikumbura et al. (2015a, b; classification, phylogeny), Lu et al. (2016; new species), Wijayawardene et al. (2016b; morphology, phylogeny), ex-type: CPC 112007; holotype CBS H-20948.

Brunneosphaerella Crous 2009, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *B. protearum* (Syd. & P. Syd.) Crous., asexual morph coelomycetous, saprobes, terrestrial, South Africa, see Crous et al. (2009d, 2011d; morphology; phylogeny), Schoch et al. (2009; morphology, phylogeny), Hyde et al. (2013; accepted as in *Mycosphaerellaceae*), Wijayawardene et al. (2014c; accepted as in *Mycosphaerellaceae*), epitype: CBS H-20335, ex-epitype culture CBS 130597 = cCPC 16368.

Brunneosporella Ranghoo & K.D. Hyde 2001, *Papulosaceae*, *Diaportheomycetidae* families *incertae sedis*, *Sordariomycetes*, one species, type: *B. aquatica* Ranghoo & K.D. Hyde, one species, asexual morph unknown, saprobes, aquatic, Hong Kong, see Lumbsch and Huhndorf (2010; outline, as a genus in *Annulatasaceae*), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Papulosaceae*), holotype: HKU (M) 5251.

Brunnipila Baral 1985, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, c. ten species, type: *B. clandestina* (Bull.) Baral, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), Jaklitsch et al. (2016a; classification), cultures and sequences are available but lacking for the type species.

Brycekendrickia Nag Raj 1973, *Ascomycota* genera *incertae sedis*, one species, type: *B. indica* Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Brycekendrickomyces Crous & M.J. Wingf. 2009, *Trichomeriaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *B. acaciae* Crous & Wingf, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Indonesia, see Crous et al. (2009a; morphology, phylogeny), holotype and ex-type: CBS H-20198, CPC 15078 = CBS 124104.

Bryobilimbia Fryday, Printzen & S. Ekman 2014, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, six species, type: *B. hypnorum* (Lib.) Fryday, Printzen & S. Ekman, lichenized, see Fryday et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Bryocaulon Kärnefelt 1986, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, four species, type: *B. divergens* (Ach.) Kärnefelt, lichenized, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Bryocentria Döbbeler 2004, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, eight species, type: *B. brongniartii* (P. Crouan & H. Crouan) Döbbeler, sexual morph unknown, saprobes, epiphytes, terrestrial, cosmopolitan, see Döbbeler (2010; new species), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Bryochiton Döbbeler & Poelt 1978, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, four species, type: *B. monascus* Döbbeler & Poelt, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available for unidentified species.

Bryoclaviculus L. Ludw., P.R. Johnst. & Steel 2013, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *B. campylopi* P.R. Johnst., asexual morph

unknown, saprobes, terrestrial, New Zealand, see Johnston et al. (2013; taxonomy, DNA), holotype of the type: OTA 062197, cultures and sequences are available.

Bryodina Hafellner 2001, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *B. rhypariza* (Nyl.) Hafellner, with ascomata, terrestrial (bryophilous), Northern Hemisphere, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences unavailable.

Bryoglossum Redhead 1977, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *B. gracile* (P. Karst.) Redhead, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available for the type: DAOM178087.

Bryogomphus Lücking, W.R. Buck, Sérus. & L.I. Ferraro 2005, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *B. caribaeus* (W.R. Buck) Lücking, W.R. Buck, Sérus. & L.I. Ferraro, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Bryonora Poelt 1983, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, eleven species, type: *B. castanea* (Hepp) Poelt, with ascomata, terrestrial (bryophilous), bipolar, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available only for the type species.

Bryopelta Döbbeler & Poelt 1978, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *B. variabilis* Döbbeler & Poelt, asexual morph unknown, saprobes, terrestrial, Norway, Sweden, see Lumbsch and Huhndorf (2010; outline; accepted as a genus in *Dothideomycetes*), Kirk et al. (2013; genus accepted), Li et al. (2014b; accepted as a genus in *Mycosphaerellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Bryoplaca Søchting, Frödén & Arup 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *B. sinapisperma* (DC.) Søchting, Frödén & Arup, lichenized, Northern Hemisphere and Antarctica, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Bryrella Döbbeler 1978, *Dothideomycetes* genera *incertae sedis*, ten species, type: *B. acrogena* Döbbeler, asexual morph unknown, saprobes or parasitic, terrestrial, Europe (Italy, Austria, Poland, Sweden, Switzerland), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Li et al. (2014b; taxonomy), Wijayawardene et al. (2014c; outline), sequences are available.

Bryoria Brodo & D. Hawksw. 1977 (= *Bryopogon* Th. Fr. 1860; *Setaria* Ach. ex Michx. 1803 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, c. 60 species, type: *B. trichodes* (Michx) Brodo & D. Hawksw., widespread but mainly N Hemisphere, see Lumbsch and Huhndorf (2010; outline), Myllys et al. (2011, 2014; phylogeny), Kirk et al. (2013; genus accepted), Velmala et al. (2014; phylogeny), Jaklitsch et al. (2016a; position), Lücking et al. (2017a; classification), holotype: PC, sequences are available.

Bryoscyphus Spooner 1984 (= *Calyptellopsis* Svrček 1986 *fide* Jaklitsch et al. 2016a), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, 19 species, type: *B. conocephali* (Boyd) Spooner, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; DNA), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available but lacking for the type.

Bryosphaeria Döbbeler 1978, *Trematosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, nine species, type: *B. cinclidoti* (Racov.) Döbbeler, asexual morph unknown, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Li et al. (2014b; transferred to *Trematosphaeriaceae*), sequences are unavailable.

Bryostigma Poelt & Döbbeler 1979, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, two species, type: *B. leucodontis* Poelt & Döbbeler, lichenized, see Sundin et al. (2012; taxonomy), Frisch et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Bryostroma Döbbeler 1978, *Dothideomycetes* genera *incertae sedis*, seven species, type: *B. rhacomitrii* Döbbeler, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Bryothele Döbbeler 1998, *Dothideomycetes* genera *incertae sedis*, one species, type: *B. mira* Döbbeler & Melnik, asexual morph unknown, saprobes or parasitic, terrestrial, America and Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Budhangurabania P. Wong, Khemmuk & R.G. Shivas 2015, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *B. cynodonticola* P. Wong, Khemmuk & R.G. Shivas, coelomycetous, saprobes, Australia, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2017; outline) holotype of the type: BRIP 59305, includes cultures.

Buellia De Not. [nom. cons.] 1846, *Caliciaceae*, *Calicales*, *Lecanoromycetes*, 300 species, type:

B. disciformis (Fr.) Mudd, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Prieto and Wedin (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Buelliastrum Zahlbr. 1930, *Lecanoromycetes* genera *incertae sedis*, two species, type: need typification data, lichenized, China, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Buellia Fink 1935, *Asterinales* genera *incertae sedis*, *Dothideomycetes*, eleven species, type: *B. minimula* (Tuck.) Fink, lichenicolous, Asia, see Hafellner et al. (2008; new species), Lumbsch and Huhndorf (2010; outline), Pérez-Ortega and Etayo (2010; new species), van den Boom (2010; new species), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Ertz and Diederich (2015; phylogeny, accepted in *Asterinales*), sequences are available.

Buerenia M.S. Reddy & C.L. Kramer 1975, *Protomycetaceae*, *Taphrinales*, *Taphrinomycetes*, three species, type: *B. cicutae* (Lindr.) M.S. Reddy & C.L. Kramer, asexual morph unknown, ?saprobes or parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Buergenerula Syd. 1936 (= *Yukonia* R. Sprague 1962 *fide* Species Fungorum 2017), *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, six species, type: *B. biseptata* (Rostr.) Syd., asexual morph harpophora-like and pyricularia-like, pathogen, marine and freshwater habitats, worldwide, see Jones et al. (2009b, 2015; outline), Lumbsch and Huhndorf (2010; outline), Yuan et al. (2010; phylogeny), Al-Nasrawi and Hughes (2012; diversity), Kirk et al. (2013; genus accepted), Luo and Zhang (2013; phylogeny), Chen et al. (2014; mitoviruses), Hujslová et al. (2014; phylogeny), Klaubauf et al. (2014; phylogeny), Calado et al. (2015; diversity, ecology), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Bulbilopycnis Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *B. caffra* Matsush., coelomycetous, sexual morph unknown, saprobes, terrestrial, Transvaal, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bulbithecum Udagawa & T. Muroi 1990, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *B. hyalosporum* Udagawa & T. Muroi, asexual morph acremonium-like, saprobes, India, see Lumbsch and Huhndorf (2010; outline), Giraldo et al. (2012; DNA, phylogeny), Kirk et al. (2013; genus accepted),

Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 318.91.

Bulbocatenospora R.F. Castañeda & Iturr. 2000, *Ascomycota* genera *incertae sedis*, one species, type: *B. complanata* R.F. Castañeda, Iturr. & Decock, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Venezuela, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: MUCL 40705, cultures and sequences are unavailable, compare with *Piricauda*, *Chuppia*, *Neoarbuscula* and *Peyronelia*.

Bulbomollisia Graddon 1984, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *B. radiata* Graddon, sexual morph unknown, saprobes, UK, see Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Bulbothrix Hale 1974, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 60 species, type: *B. semilunata* (Lynge) Hale, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Benatti (2013; review), Kirika et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Bulgaria Fr. 1822, *Phaciidaeae*, *Helotiales*, *Leotiomycetes*, two species, type: *B. inquinans* (Pers.) Fr., asexual morph unknown, saprobes, terrestrial, northern and southern temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014b; phylogeny), cultures available for the type: CBS 315.71.

Bulgariella P. Karst. 1885, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, four species, type: *B. pulla* (Fr.) P. Karst., asexual morph unknown, saprobes, terrestrial, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available (unpublished).

Bulgariopsis Henn. 1902, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *B. moelleriana* Henn., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bullanockia Crous 2016, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *B. australis* Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type of the type: CBS H-22879, CPC 28976.

Bullaserpens Bat., J.L. Bezerra & Cavalc. 1965, *Ascomycota* genera *incertae sedis*, one species, type: *B. bignoniacearum* Bat., J.L. Bezerra & Cavalc., coelomycetous, sexual morph unknown, saprobes,

terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Bullimyces A. Ferrer, A.N. Mill., C. Sarmiento & Shearer 2012, *Sordariomycetidae* *incertae sedis*, *Sordariomycetes*, three species, type: *B. communis* A. Ferrer, A.N. Mill., Sarmiento & Shearer, Costa Rica, see Ferrer et al. (2012; taxonomy), Cai et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: AF281-1, ILL, cultures and sequences are available.

Bunodophoron A. Massal. 1861, *Sphaerophoraceae*, *Lecanorales*, *Lecanoromycetes*, 24 species, type: *B. australe* (Laurer) A. Massal., asexual morph unknown, lichenized, terrestrial, mainly in the temperate rainforests of the Southern Hemispheres, rarely in oceanic areas of the Northern Hemisphere or in high-altitude rainforests in tropical or subtropical areas, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Burenia M.S. Reddy & C.L. Kramer 1975, *Protomycetaceae*, *Taphriniales*, *Taphrinomycetes*, three species, type: *B. cicutae* (Lindr.) M.S. Reddy & C.L. Kramer, asexual reproduction is unknown, plant parasitic, parasitic on Apiaceae, Europe, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Bussabanomyces Klaubauf, Lebrun & Crous 2014, *Magnaportheaceae*, *Magnaporthales*, *Sordariomycetes*, one species, type: *B. longisporus* (Bussaban) Klaubauf, Lebrun & Crous, hyphomycetous, sexual morph unknown, endophyte, terrestrial, Thailand, see Klaubauf et al. (2014; morphology, phylogeny), Maharachchikumbura et al. (2015; outline), Wijayawardene et al. (2017; outline), holotype and ex-type: BCC11377, CBS 125232.

Butleria Sacc. 1914, *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, one species, type: *B. inaghatahani* Sacc., asexual morph unknown, parasitic, terrestrial, tropical, holotype of type: PAD 1677, see Lumbsch and Huhndorf (2010; outline), Li et al. (2011; morphology), Kirk et al. (2013; genus accepted), Dissanayake et al. (2014; morphology), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Byrsomyces Cavalc. 1972, *Sordariomycetes* genera *incertae sedis*, one species, type: *B. olivaceus* Cavalc., asexual morph unknown, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), type material is not in good condition hence need to recollect and subject them to molecular analysis, holotype: URM 27661 /Exs. 20057.

Byssosascus Arx 1971, *Myxotrichaceae*, *Leotiomycetes* families *incertae sedis*, two species, type: *B. striatosporus* (G.L. Barron & C. Booth) Arx, asexual morph unknown,

saprobes, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype: BF 24290, cultures available for type: UAMH 3572.

Byssocallis Syd. 1927, *Dothideomycetes* genera *incertae sedis*, two species, type: *B. phoebe* Syd., asexual morph unknown, parasitic on living leaves (*Phoebe tonduzii* Mez. and *Podocarpus elongate* L'Herit.), see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Tubeufiaceae*), Hyde et al. (2013) and Boonmee et al. (2014b) did not accept as a genus in *Tubeufiaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Byssocaulon Mont. 1835, *Chrysotrichaceae*, *Arthoniales*, *Arthoniomycetes*, six species, type: *B. nivale* Mont., lichenized, saxicolous or epiphytic, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Byssogene Syd. 1922, *Dothideomycetes* genera *incertae sedis*, two species, type: *B. amboinensis* Syd., asexual morph unknown, saprobes, terrestrial, Brazil and Indonesia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), da Silva (2016; taxonomy, phylogeny, accepted as a genus in *Saccardiaceae*), cultures and sequences are unavailable.

Byssolecania Vain. 1921 (= *Dothiomycetes* Bat. & J.L. Bezerra 1961; = *Gonolecania* Zahlbr. 1924; = *Lecaniella* Vain. 1896 *fide Species Fungorum* 2017), *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, seven species, type: *B. fuscolivida* Vain., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Byssoloma Trevis. 1853 (= *Calidia* Stir. 1876; = *Limbalba* Nieuwl. 1916; = *Pilocarpon* Vain. 1890; = *Tricholechia* A. Massal. 1853; = *Wainioa* Nieuwl. 1916 *fide Species Fungorum* 2017), *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, possibly heterogeneous, c. 50 species, type: *B. leprieurii* Trevis., sexual morph with apothecoid ascocarps usually having a byssoid margin, asexual morph coelomycetous, rarely campylidial, lichenized, terrestrial, mainly foliicolous in tropical rainforests, see Lücking (2008; morphology), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), van den Boom (2016; new species), holotype: Leprieur 1366 (PC), sequences are available but lacking for the type species.

Byssonectria P. Karst. 1881 (= *Inermisia* Rifai 1968; = *Pseudocollema* Kanouse & A.H. Sm. 1940 *fide Species Fungorum* 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, eleven species, type: *B. obducens* P.

Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lindemann et al. (2015; new species, DNA), cultures and sequences are available but lacking for the type.

Byssonyxena Guarro, Punsola & Cano 1987, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *B. ceratinophila* Guarro, Punsola & Cano, asexual morph unknown, saprobes, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), culture available for the type: IMI 370021.

Byssophytum Mont. 1848, *Ascomycota* genera *incertae sedis*, one species, type: *B. sulphureum* Mont., asexual morph unknown, lichenized, terrestrial, Java, Tahiti, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Byssosphaeria Cooke 1879 (= *Macbriddella* Seaver 1909; = *Xenonectria* Höhn. 1920 *fide Species Fungorum* 2017), *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, 25 species, type: *B. keithii* (Berk. & Broome) Cooke, chaetophoma-like, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Tian et al. (2015; morphology, phylogeny), cultures and sequences of type species are unavailable.

Byssotheciella Petr. 1923, *Sordariomycetes*, genera *incertae sedis*, two species, type: *B. tiliae* Petr., asexual morph unknown, saprobes, terrestrial, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Byssothecium Fuckel 1861, *Dothideomycetes* genera *incertae sedis*, eight species, type: *B. circinans* Fuckel, asexual morph coelomycetous, saprobes or parasites, terrestrial, freshwater or marine, worldwide, see Zhang et al. (2009e) accepted as *Massarinaceae* but Zhang et al. (2012e) showed it distinct from *Massarinaceae*, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Byssotrema M. Cáceres, Aptroot & Lücking 2014, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *B. mirabile* M., Aptroot & Lücking, lichenized, see Cáceres et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Cacobius Kimbr. 1967, *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, c. six species, type: *C. minusculus* Kimbr., asexual morph unknown, coprophilous, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cacumisporium Preuss 1851, *Ascomycota* genera *incertae sedis*, seven species, type: *C. tenebrosum* Preuss, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, Europe, North America, see Wongsawas et al. (2009; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cadophora Lagerb. & Melin 1927, *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, 15 species, type: *C. fastigiata* Lagerb. & Melin, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Navarrete et al. (2011; associated with grapevine trunk diseases), Seifert et al. (2011; morphology), Day et al. (2012; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Travadon et al. (2015; new species), cultures and sequences are available.

Caeruleum Arcadia 2012, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, two species; type: *C. heppii* (Nägeli ex Körb.) K. Knudsen & Arcadia, lichenized, see Arcadia and Knudsen (2012; taxonomy), Lücking et al. (2017a; classification), sequences unavailable.

Caespitotheca S. Takam. & U. Braun 2005, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, one species, type: *C. forestalis* (Mena) S. Takam. & U. Braun, asexual morphs hyphomycetous, biotrophic, obligate plant pathogens, terrestrial, South America (Argentina), see Lumbsch and Huhndorf (2010; outline), Braun and Cook (2012; taxonomy), sequences are available.

Cainia Arx & E. Müll. 1955, *Cainiaceae*, *Xylariales*, *Sordariomycetes*, seven species, type: *C. graminis* (Niessl) Arx & E. Müll., asexual morph coelomycetous, saprobes, pathogenic, terrestrial, temperate, see Kirk et al. (2008, 2013; genus accepted), Lumbsch & Huhndorf (2010; outline), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; phylogeny), cultures and sequences are available, reference specimens: CBS 136.62, MFLUCC 15-0540 = It 1462.

Cainiella E. Müll. 1957, *Sydowiellaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *C. johansonii* (Rehm) E. Müll., asexual morphs unknown, saprobes, terrestrial, Canada, Europe, see Lumbsch and Huhndorf (2010; outline), Kruys and Castlebury (2012; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Senanayake et al. (2017, placement), cultures and sequences are available.

Cainomyces Thaxt. 1901, *Laboulbeniales* genera *incertae sedis*, *Laboulbeniomycetes*, one species, type: *C. isomali* Thaxt., asexual morph unknown, saprobes, terrestrial, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Calathaspis I.M. Lamb & W.A. Weber 1972, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *C. devexa* I.M. Lamb & W.A. Weber, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Calcarispora Marvanová & Marvan 1963, *Ascomycota* genera *incertae sedis*, one species, type: *C. hiemalis* Marvanová & Marvan, hyphomycetous, sexual morph unknown, aquatic, Europe and Canada, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Calcarisporium Preuss 1851, *Calcarisporiaceae*, *Hypocreales*, *Sordariomycetes*, 15 species, type: *C. arbuscula* Preuss, hyphomycetous, sexual morph unknown, saprobes, widespread, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Sun et al. (2016, 2017; new species, placement), cultures and sequences Anthostomelloides available.

Calceispora Matsush. 1975, *Ascomycota* genera *incertae sedis*, two species, type: *C. hachijoensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Calceomyces Udagawa & S. Ueda 1988, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. lacunosus* Udagawa & S. Ueda, asexual morph hyphomycetous, saprobes, terrestrial (collected from activated sludge), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as *Xylariales* genera *incertae sedis*), cultures and sequences are available, type: CBS 633.88 = NHL 2983.

Calenia Müll. Arg. 1890 (= *Phlyctidium* Müll. Arg. 1888 fide Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, 30 species, type: *C. depressa* Müll. Arg., lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Caleniopsis Vězda & Poelt 1987 (= *Caleniomyces* Cif. & Tomas. 1953 fide Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *C. laevigata* (Müll. Arg.) Vězda & Poelt, lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Caleutypa Petr. 1934, *Sordariomycetes*, genera *incertae sedis*, two species, type: *C. maculans* Petr., saprobes, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Caliciopsis Peck 1883 (= *Capnodiella* (Sacc.) Sacc. & D. Sacc. 1905; = *Capnodium* subgen. *Capnodiella* Sacc. 1882; = *Hypotheeca* Ellis & Everh. 1885; = *Lagenula* G. Arnaud 1930; = *Sorica* Giesenh. 1904 *fide* Species Fungorum 2017), *Coryneliaceae*, *Coryneliales*, *Eurotiomycetes*, 31 species, type: *C. pinea* Peck, asexual morph coelomycetous (phoma-like), parasites, pathogens, see Lumbsch and Huhndorf (2010; outline), Pratibha et al. (2010; new species), Garrido-Benavent and Pérez-Ortega (2015; new species), Crous et al. (2016a; new species, phylogeny), Wood et al. (2016; DNA, phylogeny), cultures available for the type: CBS 139.64.

Calicium Pers. 1794 (= *Cyphelium* Ach. 1815; see Species Fungorum 2017 for additional synonyms), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 30 species, type: *C. viride* Pers., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Gaya et al. (2012; phylogeny), Prieto et al. (2013; phylogeny), Prieto and Wedin (2016; classification), Lücking et al. (2017a; classification), sequences are available.

Callebaea Bat. 1962, *Dothideomycetes*, genera *incertae sedis*, one species, type: *C. rutideae* (Hansf.) Bat., saprobes, terrestrial, Uganda, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Callerascus Whitton, K.D. Hyde & McKenzie 2012, *Leotiomycetes* genera *incertae sedis*, one species, type: *C. tunicaerulus* Whitton, K.D. Hyde & McKenzie, asexual morph unknown, saprobes, terrestrial, Australia, see Whitton et al. (2012b; taxonomy holotype of the type: IFRD 9021, cultures and sequences are unavailable.

Callistospora Petr. 1955, *Ascomycota* genera *incertae sedis*, one species, type: *C. gaubae* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Callome Otálora & Wedin 2014, *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, one species, type: *C. multipartita* (Sm.) Otálora, P.M. Jørg. & Wedin, lichenized, see Otálora et al. (2013; 2014; taxonomy, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Calloria Fr. 1836 (= *Beloniella* Rehm 1892; = *Belonopeziza* Höhn. 1917; = *Cylindrocolla* Bonord. 1851;

Callorina Korf 1971 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Calloriaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *C. fusariooides* (Berk.) Fr., asexual morph *Cylindrocolla*, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Seifert et al. (2011; morphology of asexual morph), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are available but lacking for the type species.

Calloriopsis Syd. & P. Syd. 1917 (= *Diplothrix* Vain. 1921 *fide* Jaklitsch et al. 2016a), *Helicogoniaceae*, *Phacidiales*, *Leotiomycetes*, five species, type: *C. gelatinosa* (Sacc.) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Calogaya Arup, Frödén & Søchting 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 14 species, type: *C. biatorina* (A. Massal.) Arup, Frödén & Søchting, lichenized, see Arup et al. (2013; taxonomy, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Calonectria De Not. 1867 (= *Cylindrocladium* Morgan 1892; = *Candelospora* Rea & Hawley 1912 *fide* Rossman et al. 2016a), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, type: *pyrochroa* (Desm.) Sacc., 248 species are listed in Index Fungorum 2016, type: *C. daldiniana* De Not., pathogenic to plants or isolated from soil, terrestrial, worldwide, see Lombard et al. (2010a, b, 2015a, b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), ex-epitype: CBS 749.70; epitype: of the type: CBS H-15110.

Calongea Healy, Bonito & Trappe 2009, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *C. prieuensis* (Mor.-Arr., J. Gómez & Calonge) Healy, Bonito & Trappe, asexual morph unknown, saprobes, from soil, terrestrial, Europe, see Healy et al. (2009b; taxonomy), holotype of the type: MA (Fungi) 33502, cultures and sequences are available.

Calongeomyces Hawksworth & Etayo 2011 (= *Calongia* D. Hawksw. & Etayo 2010 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *C. gibelluloides* (D. Hawksw. & Etayo) D. Hawksw. & Etayo, coelomycetous, sexual morph unknown, parasymbiont on *Cladonia*, *Flavoparmelia*, *Pertusaria*, terrestrial, southern Europe, see Hawksworth and Etayo (2011; morphology, distinction of similar lichenicolous genera), holotype: (K, (M) 164014, J. Etayo & A. Etayo), cultures and sequences are unavailable.

Calopadia Vězda 1986 (= *Crocicreomyces* Bat. & Peres 1964 [1963]; = *Cyrtula* Bat. & H. Maia 1961 *fide* Species Fungorum 2017), *Pilocarpaceae*, *Lecanorales*,

Lecanoromycetes, 27 species, type: *C. fusca* (Müll. Arg.) Vězda, lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Calopadiopsis Lücking & R. Sant. 2002, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *C. tayabasensis* (Vain.) Lücking & R. Sant., lichenized, Argentina, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Calophoma Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *C. clematidina* (Thüm.) Q. Chen & L. Cai, coelomycetous, saprobes, ?pathogens, terrestrial, The Netherlands, New Zealand, Switzerland, see Chen et al. (2015b; taxonomy, phylogeny), Tibpromma et al. (2017b; new species), Wijayawardene et al. (2017; outline), epitype and ex-epitype of the type: CBS H-16193, CBS 108.79.

Caloplaca Th. Fr. [nom. cons.] 1860 (see Species Fungorum 2017 for synonyms), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 350 species, type: *C. cerina* (Hedw.) Th. Fr., lichenized, see Lumbsch and Huhndorf (2010; outline), Arup et al. (2013; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Caloscypha Boud. 1885 (= *Geniculodendron* G.A. Salt 1974; = *Otidella* Sacc. 1889 fide Healy et al. 2016; Species Fungorum 2017), *Caloscyphaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *C. fulgens* (Pers.) Boud., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures available for the type: AFTOL-ID 152.

Calosphaeria Tul. & C. Tul. 1863 (= *Calosphaeriophora* Réblová, L. Mostert, W. Gams & Crous 2004 fide Réblová et al. 2016c), *Calosphaeriaceae*, *Calosphaeriales*, *Sordariomycetes*, c. 30 species, type: *C. princeps* Tul. & C. Tul., asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Damm et al. (2008b; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2015a, 2016c; phylogeny, nomenclature), cultures available but lacking for the type species.

Calosphaeriopsis Petr. 1941, *Sordariomycetes*, genera *incertae sedis*, one species, type: *C. huberiana* (Kirschst.) Petr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Calosporella J. Schröt. 1897, *Sydiomycetidae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. innesii* (Carr.) J. Schröt., asexual morph unknown, saprobes, terrestrial, Europe, see Voglmayr and Jaklitsch (2014; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2017; placement), cultures and sequences are unavailable, ex-type and epitype: CBS 123810, WU 32161.

Calostilbe Sacc. & P. Syd. 1902 (= *Calostilbella* Höhn. 1919; = *Nectria* subgen. *Phaeonectria* Sacc. 1895; = *Phaeonectria* (Sacc.) Sacc. & Trotter 1913; = *Xenostilbum* Petr. fide Rossman et al. 2016a; Species Fungorum 2017), *Nectriaceae*, *Hypoocreales*, *Sordariomycetes*, two species, type: *C. striispora* (Ellis & Everh.) Seaver, asexual morph hyphomycetous, saprobes on wood and bark, commonly in tropical regions, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2015a; phylogeny); Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), lectotype: NY, iso-lectotypes BPI 553210, 553211.

Calothyriopsis Höhn. 1919 (= *Ptychopeltis* Syd. 1927 fide Species Fungorum 2017), *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, two species, type: *C. conferta* (Theiss.) Höhn., saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Microthyriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Calotrichopsis Vain. 1890, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, three species, type: *C. insignis* Vain., lichenized, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Calvitimela Hafellner 2001, *Tephromelataceae*, *Lecanorales*, *Lecanoromycetes*, ten species, type: *C. armeniaca* (DC.) Hafellner, lichenized, see Fryday (2011; taxonomy), Bendiksby et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Calvolachnella Marinc., T.A. Duong & M.J. Wingf. 2016, *Chaetosphaeriales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. guaviyuensis* (Marinc. T.A. Duong, M.J. Wingf. & C.A. Perez) Marinc., T.A. Duong & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, terrestrial, Uruguay, see Hernández-Restrepo et al. (2016a; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type: PREM 60964, CBS 134695 = CMW 39055.

Calycella (Sacc.) Sacc. 1899 (= *Helotium* subgen. *Calycella* Sacc. 1889 *fide* Species Fungorum 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *C. gelatinosa* (Sacc.) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Europe, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Calycellina Höhn. 1918 (= *Chaetochalara* B. Sutton & Piroz. 1965; *Phialina* Höhn. 1926; = *Phialoscypha* Raity. 1977; = *Scutoscypha* Graddon 1980; = *Setoscypha* Velen. 1934 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, c. 45 species, type: *C. punctiformis* (Grev.) Höhn., asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Seifert et al. (2011; morphology of asexual morph), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available but lacking for the type.

Calycellinopsis W.Y. Zhuang 1990, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. xishuangbanna* W.Y. Zhuang, asexual morph unknown, saprobes, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Calycidium Stirt. 1877 (= *Calycidiomyces* Cif. & Tomas. 1953; = *Coniophyllum* Müll. Arg. 1892 *fide* Species Fungorum 2017), *Sphaerophoraceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *C. cuneatum* Stirt., lichenized, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Calycidiaceae*), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Calycina Nees ex Gray 1821 (= *Antinoa* Velen. 1934; = *Calycella* (Fr.) Boud., 1885; = *Cystopezizella* Svrček 1983; *Eubelonis* Clem. 1909; = *Gemmina* Raity. 2004; = *Helotium* subgen. *Calycella* Fr. 1849; = *Malotium* Velen. 1934, nom. Nom. illegit., Art. 53.1; = *Parthenope* Velen. 1934; = *Pezizella* Fuckel 1870 [1869-70]; = *Septopezizella* Svrček 1987; = *Weinmannioscyphus* Svrček *fide* Jaklitsch et al. 2016a; Species Fungorum 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, c. 30 species, type: *C. herbarum* (Pers.) Gray, asexual morph hyphomycetous, saprobes, cones scales, herbaceous stems and leaves, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Baral et al. (2013; genus accepted), Han et al. (2014; genus accepted), Baral and Rämä (2015; genus accepted), type in S herbarium, cultures and sequences are unavailable for the type of the type species.

Calyptra Theiss. & Syd. 1918, *Dothideomycetes* genera *incertae sedis*, five species, type: *C. cordobensis* (Speg.)

Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Calyptronectria Speg. 1909, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *C. platensis* Speg., asexual morph unknown, saprobes, terrestrial, Argentina, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Tian et al. (2015; taxonomy, phylogeny), cultures and sequences of type species are unavailable.

Calyptrozyma Boekhout & Spaay 1995, *Eurotiomycetidae*, genera *incertae sedis*, one species, type: *C. arxii* Boekhout & Spaay, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 354.92.

Camaroglobulus Speer 1986, *Ascomycota* genera *incertae sedis*, one species, type: *M. caespitosa* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Camarographium Bubák 1916, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, seven species, type: *C. stephensii* (Berk. & Broome) Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Crous et al. (2011c; new species, key), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2016b, 2017; outline, taxonomy, new species), cultures and sequences are available but lacking for the type species.

Camaropella Lar. N. Vasilyeva 1997, *Boliniaceae*, *Boliniales*, *Sordariomycetes*, two species, type: *C. pugillus* (Schwein.) Lar. N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, north temperate zone, see Huhndorf and Miller (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Untereiner et al. (2013; phylogeny), Friebes and Wendelin (2014), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Camarops P. Karst. 1873 (= *Biscogniauxia* sect. *Camarops* (P. Karst.) Lar.N. Vassiljeva et al. 1998; = *Bolinia* (Nitschke) Sacc. 1882; = *Chromendothia* Lar.N. Vassiljeva 1993; = *Hypoxylon* sect. *Bolinia* Nitschke 1867; = *Peridoxyylon* Shear 1923; = *Phaeosperma* Nitschke ex Fuckel 1870 [1869-70]; = *Sarcostromella* Boedijn 1959; = *Solenoplea* Starbäck 1901 *fide* Species Fungorum 2017), *Boliniaceae*, *Boliniales*, *Sordariomycetes*, 20 species, type: *C. hypoxylonoides* P. Karst., asexual morphs unknown, saprobes, terrestrial, worldwide, see Huhndorf and Miller (2008; new species), Lumbsch and Huhndorf (2010;

outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Camaropycnis E.K. Cash 1945, Ascomycota genera *incertae sedis*, one species, type: *C. libocedri* E.K. Cash, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Camarosporellum Tassi 1902, Ascomycota genera *incertae sedis*, one species, type: *C. nervisequum* (Tassi) Tassi, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2016b, 2017; taxonomy, outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Camarosporoides W.J. Li & K.D. Hyde 2016, Phaeosphaeriaceae, Pleosporales, Dothideomycetes, one species, type: *C. phragmitis* W.J. Li & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Germany, see Hyde et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 16-1488, MFLUCC 13-0365.

Camarosporiopsis Abbas, B. Sutton & Ghaffar 2000, Ascomycota genera *incertae sedis*, one species, type: *C. capparis* (S. Ahmad) Abbas, Sutton & Ghaffar, coelomycetous, sexual morph unknown, one species, saprobes, Pakistan, see Wijayawardene et al. (2012, 2016b, 2017; taxonomy, outline), sequences are unavailable.

Camarosporium Schulzer 1870 (= *Piringa* Speg. 1911; = *Sclerotheca* Bubák & Vleugel 1917 *fide* Species Fungorum 2017), Camarosporaceae, Pleosporales, Dothideomycetes, more than 100 species, type: *C. quaternatum* Schulzer, coelomycetous, sexual morph cucurbitaria-like, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2012, 2014a, c, 2016b, 2017; taxonomy, phylogeny, outline), Liu et al. (2015a, taxonomy, phylogeny), Tibpromma et al. (2016e, 2017b; taxonomy, phylogeny), sequences are available for 11 species.

Camarosporula Petr. 1954 (= *Anthracostroma* Petr. 1954 *fide* Rossman et al. 2015b), Teratosphaeriaceae, Capnodiales, Dothideomycetes, one species, type: *C. persoonii* (Henn.) Petr., asexual morph coelomycetous, saprobes, terrestrial, Australia, see Crous et al. (2011e; phylogeny), Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), cultures available for the type: CBS 116258.

Camarosporulum Tassi 1902, Ascomycota genera *incertae sedis*, one species, type: *C. ampelopsidis* Tassi, coelomycetous, sexual morph unknown, saprobes,

terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Camarotella Theiss. & Syd. 1915, Phyllachoraceae, Phyllachorales, Sordariomycetes, three species, type: *C. astrocaryae* (Rehm) Theiss. & Syd., asexual morph unknown, saprobes, see Kirk et al. (2008; treated as a synonym of *Coccodiella* Hara), Lumbsch and Huhndorf (2010; did not list), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a distinct genus), cultures and sequences are unavailable.

Cameronia Kantvilas 2012, Cameroniaceae, Baeomycetales, Lecanoromycetes, two species, type: *C. pertusariooides* Kantvilas, lichenized, see Kantvilas (2012; taxonomy), Lumbsch et al. (2012; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Camillea Fr. 1849 (= *Bacillaria* Mont. 1840; *Diatrype* subgen. *Nummularoidea* Cooke & Massee 1892; = *Diatrypeopsis* Speg. 1884; = *Nummularoidea* (Cooke & Massee) Lloyd 1924; = *Xylocladium* P. Syd. ex Lindau 1900; = *Jongiella* M. Morelet 1971; = *Nummularoidea* (Cooke & Massee) Lloyd 1924 *fide* Species Fungorum 2017; Wendt et al. 2017), Graphostromataceae, Xylariales, Sordariomycetes, more than 48 species, type: *C. leprieurii* (Mont.) Mont., endophytes-saprobes, widespread and high in tropics, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2013), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; transferred to Graphostromataceae), cultures and sequences are available.

Camposia Speg. 1921 (= *Myxostomella* Syd. 1931 *fide* Species Fungorum 2017), Parmulariaceae, Dothideomycetes families *incertae sedis*, two species, type: *C. pulcherrima* Speg., asexual morph unknown, saprobes, terrestrial, South America, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Camporesia W.J. Li & K.D. Hyde 2016, Xylariales genera *incertae sedis*, Sordariomycetes, two species, type: *C. sambuci* W.J. Li & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Li et al. (2016a; taxonomy, phylogeny), Wendt et al. (2017; transferred to *Xylariales* genera *incertae sedis*), holotype and ex-type: MFLU 15-3905, MFLUCC 13-0203.

Camposporidium Nawawi & Kuthub. 1988, Ascomycota genera *incertae sedis*, three species, type: *C. cristatum* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, aquatic, terrestrial, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Camposporium Harkn. 1884, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 18 species, type: *C. antennatum* Harkn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology, accepted as a genus in *Leptosphaeriaceae*), Wijayawardene et al. (2012, 2017; outline), Adamčík et al. (2015; new species), cultures available for the type: ICMP15321.

Camptomeriphila Crous & M.J. Wingf. 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *C. leucaenae* Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type: CBS H-22884, CPC 27608.

Camptomeris Syd. 1927, *Ascomycota genera incertae sedis*, seven species, type: *C. calliandrae* Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Campomyces Thaxt. 1894, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eight species, type: *C. melanopus* Thaxt., asexual morph unknown, parasitic on Staphylinidae beetles, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Campophora Réblová & Unter. 2013, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *C. hylomeconis* (Crous, de Hoog & H.D. *Chaetothyriales* Shin) Réblová & Unter., asexual morph unknown, from living leaves of *Hylomecon vernalis*, terrestrial, South Korea, see Réblová et al. (2013; genus accepted), Madrid et al. (2016; phylogeny), holotype and ex-type of the type: CBS H-19907, CBS 113311.

Campsphaeria Fuckel 1870, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, four species, type: *C. sulphurea* Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, need generic revision.

Campylocarpon Haldeen, Schroers & Crous 2004, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *C. fasciculare* Schroers, Haldeen & Crous, hyphomycetous, sexual morph unknown, pathogenic on *Vitis vinifera*, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline),

holotype and ex-type of the type: CBS 6590, CBS 112613 = CPC 3970.

Campylospora Ranzoni. 1953, *Hypocreomycetidae* genera *incertae sedis*, five species, type: *C. chaetocladia* Ranzoni, hyphomycetous, sexual morph unknown, saprobes, endophytes, aquatic, Asia, Europe, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Marvanová and Laichmanová (2014; new species), cultures and sequences are available for authentic strains CCM F-17708, CCM F-10610, CCM F-10710, CCM F-10810.

Campylothelium Müll. Arg. 1883, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, six species, type: *C. puiggarii* Müll. Arg., lichenized, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Nelsen et al. (2014b; taxonomy), cultures available for the type: MPN268.

Canalisporium Nawawi & Kuthub 1989 (= *Ascothailandia* Sri-indr. et al. 2010 *fide* Réblová et al. 2016c), *Savoryellaceae*, *Savoryellales*, *Sordariomycetes*, twelve species, type: *C. caribense* (Hol. Jech. & Mercado) Nawawi & Kuthub., hyphomycetous, freshwater habitats except *C. microsporum* and *C. nigrum*, Africa, see Seifert et al. (2011; morphology), Sri-Indrasutdh et al. (2010; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), Réblová et al. (2016c; nomenclature) cultures and sequences available.

Canariomyces Arx 1984, *Microascaceae*, *Microascales*, *Sordariomycetes*, one species, type: *C. notabilis* Arx, asexual morph unknown, saprobes, terrestrial, Canary Islands, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype of the type: MUCL 44677, a sequence available but unpublished.

Canavirgella W. Merr., N.G. Wenner & Dreisbach 1996, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *C. banfieldii* W. Merr., N.G. Wenner & Dreisbach, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cancellidium Tubaki 1975, *Sordariales* genera *incertae sedis*, two species, type: *C. applanatum* Tubaki, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, Europe, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Zhao et al. (2013a; China), Pratibha et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Candelabrum Beverw. 1951 (= *Dionysia* G. Arnaud *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, seven species, type: *C. spinulosum* Beverw., hyphomycetous, sexual morph unknown, on decaying leaves in water, aquatic, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Candelaria A. Massal. 1852, *Candelariaceae*, *Candelariales*, *Lecanoromycetes*, seven species, type: *C. concolor* (Dicks.) Arnold, lichenized, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Candelariella Müll. Arg. 1894 (= *Caloplacopsis* (Zahlbr.) B. de Lesd. 1932; = *Candelariella* sect. *Caloplacopsis* Zahlbr. 1926; = *Candelariellomyces* E.A. Thomas ex Cif. & Tomas. 1953 [1952], = *Candelariellopsis* Werner 1936; = *Eklundia* C.W. Dodge 1968; = *Pleochroma* Clem. 1909 *fide Species Fungorum* 2017), *Candelariaceae*, *Candelariales*, *Lecanoromycetes*, 50 species; type: *C. vitellina* (Hoffm.) Müll. Arg., lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Candelina Poelt 1974, *Candelariaceae*, *Candelariales*, *Lecanoromycetes*, three species, type: *C. mexicana* (B. de Lesd.) Poelt, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Candelosynnema K.D. Hyde & Seifert 1992, *Ascomycota* genera *incertae sedis*, one species, type: *C. ranunculosporum* K.D. Hyde & Seifert, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Candida Berkout 1923 (= *Lodderomyces* Van der Walt 1966), *Saccharomycetales*, genera *incertae sedis* *Saccharomycetes*, 40 species, type: *C. tropicalis* var. *tropicalis* (Castell.) Berkout, asexual reproduction by multilateral budding and formation of pseudo-hyphae and true hyphae, saprophytic, others pathogens or opportunistic pathogens of humans and animals, terrestrial, worldwide, phylogenetic analysis of gene sequences shows that there are about 40 species in the *Candida tropicalis* clade, the type species of the genus *Candida*, which also includes *C. albicans* and *Lodderomyces elongisporus*. Prior to DNA sequence analysis and changes in the Code of

nomenclature for fungi (Melbourne Code), approximately 400 described anamorphic species were included in the genus *Candida*.

Cannonia Joanne E. Taylor & K.D. Hyde 1999, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. australis* (Speg.) Joanne E. Taylor & K.D. Hyde, asexual morph unknown, saprobes, Argentina, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wendt et al. (2017; transferred to *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Canoparmelia Elix & Hale 1986, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 40 species; type: *C. texana* (Tuck.) Elix & Hale, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Leonardo et al. (2011; indicator for environmental pollution), Kirk et al. (2013; genus accepted), Kirika et al. (2016; phylogeny and circumscription), Lücking et al. (2017a; classification), sequences are available.

Cantharomyces Thaxt. 1890, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 29 species, type: *C. bledii* Thaxt., asexual morph unknown, biotrophic, hosts are Dryopidae, Hydrophilidae, Limnichidae, and Staphylinidae (Coleoptera), widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters and De Kesel (2013; morphology), cultures and sequences are unavailable.

Capillataspora K.D. Hyde 1989, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. corticola* K.D. Hyde, asexual morph unknown, saprobes, collected from mangroves, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Capillipes R. Sant. 1956, *Helotiaceae* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. cavorum* R. Sant., asexual morph unknown, saprobes, terrestrial, Lapland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Capillistichus Santam. 2004, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *C. tenellus* Santam., asexual morph unknown, parasitic on Hydrophilidae beetles, aquatic, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Capitofimbria L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. compacta* (R.F. Castañeda et al.) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Lombard et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: CBS-H6584a, CBS 111739.

Capitorostrum Bat. 1957, *Ascomycota genera incertae sedis*, one species, type: *C. asteridiellae* Bat., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, Papua New Guinea, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Capitotricha (Raity.) Baral 1985, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, ten species, type: *C. bicolor* (Bull.) Baral, sexual morph unknown, saprobes, UK, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Hyaloscypheaceae*), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), Index Fungorum (2017; treated as a synonym of *Lachnum* Retz.), cultures and sequences are unavailable.

Capnobotryella Sugiy. 1987, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *C. renispora* Sugiy., hyphomycetous, sexual morph unknown, saprobes, Asia, Europe, North America, see Seifert et al. (2011; morphology), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 214.90.

Capnobotrys S. Hughes 1970, *Metacapnodiaeae*, *Capnodiales*, *Dothideomycetes*, c. 20 species, type: *C. neesii* S. Hughes, hyphomycetous, sexual morph unknown, sooty mould on living plants, terrestrial, Asia, Australasia, Europe, North America, South America, North America, see Seifert et al. (2011; morphology), Hughes and Seifert (2012; new species), Wijayawardene et al. (2012, 2017; outline), Hyde et al. (2013; accepted as a genus in *Metacapnodiaeae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Capnocheirides J.L. Crane & S. Hughes 1982, *Ascomycota genera incertae sedis*, one species, type: *C. rhododendri* (Kunze) J.L. Crane & S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Capnodiastrum Speg. 1886, *Ascomycota genera incertae sedis*, one species, type: *C. guaraniticum* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable thus placed in *Ascomycota incertae sedis* in here.

Capnodium Mont. 1849 (= *Polychaeton* (Pers.) Lév. 1846; = *Fumagospora* G. Arnaud 1911 *fide* Chomnunti et al. 2011; Hyde et al. 2013; Wijayawardene et al. 2014c), *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, one

species, type: *C. citri* Berk. & Desm., asexual morph coelomycetous, sooty moulds, worldwide, see Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2011, 2014; taxonomy, nomenclature), Hyde et al. (2013; accepted as a genus in *Capnodiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Hongsanan et al. (2015c, 2016c; new species, evolution), cultures and sequences are available.

Capnofrasera S. Hughes 2003, *Antennulariellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *C. dendryphioides* S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Capnokyma S. Hughes 1975, *Euantennariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *C. corticola* S. Hughes, asexual morph unknown, saprobes, terrestrial, Australasia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Euantennariaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Caprettia Bat. & H. Maia 1965 (= *Porinula* Vězda 1975 *fide* Species Fungorum 2017), *Monoblastiaceae*, *Monoblastiales*, *Dothideomycetes*, eight species, type: *C. amazonensis* Bat. & H. Maia, asexual morph coelomycetous, saprobes, thallus epiphyllous, worldwide, see Nelsen et al. (2009, 2011b; taxonomy), Yeshitela et al. (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Capricola Velen. 1947, *Helotiales* genera *incertae sedis*, one species, type: *C. fuscidula* Velen., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Capronia Sacc. 1883 (= *Berlesiella* Sacc. 1888; = *Caproniella* Berl. 1899; = *Dictyotrichiella* Munk 1953; = *Didymotrichiella* Munk 1953; = *Herpotrichiella* Petr. 1914; = *Melanostigma* Kirschst. 1939; = *Polytrichiella* M.E. Barr 1972 *fide* Species Fungorum 2017), *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, c. 60 species, type: *C. sexdecimspora* (Cooke) Sacc., asexual morphs hyphomycetous, parasitic (on fungi or lichens) or saprobes, terrestrial, worldwide, see Badali et al. (2008; taxonomy), Halici et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Untereiner et al. (2011; phylogeny), Friebes (2012; morphology), Diederich et al.

(2013; phylogeny), Kirk et al. (2013; genus accepted), Gueidan et al. (2014a; phylogeny), sequences are available.

Caproniella Berl. 1896, *Sordariomycetes*, genera *incertae sedis*, two species, type: *C. pleiospora* (Mouton) Berl., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Caproventuria U. Braun 1998, *Venturiaceae*, *Venturiales*, *Dothideomycetes*, two species, type: *C. hanliniana* (U. Braun & Feiler) U. Braun, asexual morph hyphomycetous, saprobes, soil-borne, Africa, Asia, Europe, New Zealand, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011d; phylogeny, accepted as a genus in *Venturiaceae*), Hyde et al. (2013; accepted as a genus in *Venturiaceae*), Wijayawardene et al. (2014c; outline), strains of the type: ATCC 9601, IFM 52023, holotype of type: HAL 7623 F.

Capsicumyces Gamundí, Aramb. & Giaiotti 1979, *Ascomycota* genera *incertae sedis*, one species, type: *C. delicatus* Gamundí, Aramb. & Giaiotti, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Capsulospora K.D. Hyde 1996, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, nine species, type: *C. frondicola* K.D. Hyde, asexual morph unknown, sexual morph ascomycetes, saprobes on dead branches, Vanuatu, Brunei, Indonesia, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Capulatispora Thambug., Kaz. Tanaka & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *C. sagittiformis* (Kaz. Tanaka & Hosoya) Thambug. & Hyde, asexual morph unknown, saprobes, terrestrial, Japan, see Thambugala et al. (2015b; morphology, phylogeny), holotype of type: TNS-F-12453, ex-type strains of type: JCM 15100.

Carassea S. Stenroos 2002, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, one species; type: *C. connexa* (Vain.) S. Stenroos, lichenized, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Carbacanthographis Staiger & Kalb 2002, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 28 species, type: *C. chionophora* (Redinger) Staiger & Kalb, lichenized (*Trenthepohlia*), terrestrial (mainly corticolous), tropical, see Tripp et al. (2010; new species), Sharma et al. (2010; key), Rivas Plata et al. (2013; DNA, phylogeny), Lumbsch et al. (2014a, b; tribe

Wirthiotremateae), Sharma and Khadilkar (2011; new species), Kukwa et al. (2013; new species), sequences are available.

Carbomyces Gilkey 1954, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, three species; type: *C. emergens* Gilkey, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Moreno et al. (2012b; re-visited, DNA, phylogeny), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as a genus in *Pezizaceae* and *Carbomycetaceae* as a synonym of *Pezizaceae*), sequences are available for the voucher AH 39344 of the type species.

Carbonicola Bendiksby & Timdal 2013, *Carbonicolaceae*, *Lecanorales*, *Lecanoromycetes*, *Ascomycota*; three species; type: *C. anthracophila* (Nyl.) Bendiksby & Timdal; lichenized; temperate northern hemisphere, Australia, see Bendiksby and Timdal (2013; taxonomy), Lücking et al. (2017a; classification), holotype: H-NYL-20375 p.p. (lectotype by Timdal 1984), sequences are available.

Carbosphaerella I. Schmidt 1969, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *C. pleosporoides* Schmidt, asexual morph unknown, saprobes, marine, Denmark, Germany, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), cultures and sequences are available.

Carestiella Bres. 1897, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *C. socia* Bres., coelomycetous, sexual morph unknown, saprobes, cosmopolitan, see Baloch et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Caribaeomyces Cif. 1962, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *C. tetrasporus* (Toro) Cif., saprobes, terrestrial, Dominican Republic, see Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Microthyriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Carinispora K.D. Hyde 1992, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *C. nypae* K.D. Hyde, saprobes, marine habitat (collected from palm *Nypa fruticans*), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available, holotype: BCC 36316.

Carnegieispora Etayo 2009, *Ascomycota* genera *incertae sedis*, one species, type: *C. rimeliae* Etayo & Berger, coelomycetous, sexual morph unknown, parasymbiont on *Parmotrema*, terrestrial, Azores, see Etayo and Berger (2009; morphology, distinction of

similar lichenicolous genera), Wijayawardene et al. (2016b; morphology), holotype: LI, F. Berger 18380, cultures and sequences are unavailable.

Carnia Bat. 1960, *Ascomycota* genera *incertae sedis*, one species, type: *C. tabebuiae* Bat. & Peres, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Carpophoromyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *C. cybocephali* Thaxt., asexual morph unknown, parasitic on Cybocephalidae beetles, terrestrial, Asia, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Carrismyces R.F. Castañeda & Heredia 2000, *Ascomycota* genera *incertae sedis*, one species, type: *C. proliferatus* R.F. Castañeda & Heredia, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Mexico, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: XAL CB736, cultures and sequences are unavailable, compare with *Bioconosporium*, *Oncopodiella*, *Pseudopetrakia* and *Septosporium*.

Caryophylloseptoria Verkley, Quaedvl. & Crous 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *C. lychnidis* (Desm.) Verkley, Quaedvl. & Crous, coelomycetous, sexual morph unknown, plant pathogenic, see Verkley et al. (2013; morphology, phylogeny), Wijayawardene et al. (2014c, 2017; outline), reference material CBS H-21161, living cultures CBS 109098, 109102.

Caryospora De Not. 1855, *Caryosporaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *C. putaminum* (Schwein.) De Not., saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Zopfiaceae*), Hyde et al. (2013; accepted as a genus in *Zopfiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as a genus in *Zopfiaceae*), Zhao and Zhao (2012; new species, China), Ariyawansa et al. (2015a; new species, introduced *Caryosporaceae*), cultures and sequences are available but lacking for the type species.

Caryosporella Kohlm. 1985, *Dothideomycetes*, genera *incertae sedis*, one species, type: *C. rhizophorae* Kohlm., asexual morph unknown, saprobes, marine aquatic, America, see Jones et al. (2009b, 2015; classification), Suetrong et al. (2009; phylogeny, morphology), Besitulo et al. (2010; diversity), Lumbsch and Huhndorf (2010; outline), Zainuddin et al. (2010; antimicrobial activities), Lee et al. (2012; checklist), Zhang et al. (2012e; morphology), Hyde et al. (2013; morphology, outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Samón-Legrá et al. (2014; diversity), Tian

et al. (2015; morphology; nomenclature; phylogeny), cultures available for the type: JK 5302A.

Casaresia Gonz. Frag. 1920 (= *Ankistrocladium* Perrott 1960 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, one species, type: *C. sphagnorum* Gonz. Frag, hyphomycetous, sexual morph unknown, aquatic, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cashiella Petr. 1951, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *C. atra* Petr., asexual morph unknown, saprobes, terrestrial, North America, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Castanedaea W.A. Baker & Partr. 2001, *Ascomycota* genera *incertae sedis*, one species, type: *C. minor* (R.F. Castañeda) W.A. Baker & Partr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Castanediella M. Hern.-Restr., Crous & M.J. Wingf. 2015, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, nine species, type: *C. acaciae* Crous, M. Hern.-Restr. & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Crous et al. (2015d, 2016a, b; taxonomy), Hernández-Restrepo et al. (2016a; new species; phylogeny), holotype and ex-type: CBS H-22225, CPC 24869 = CBS 139896.

Castanedomyces Cano, L.B. Pitarch & Guarro 2002, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *C. australiensis* Cano, L.B. Pitarch & Guarro., asexual morph unknown, saprobes, terrestrial, Australia, see Liang et al. (2009a; phylogeny), cultures available for the type: IMI 370015.

Catabotrys Theiss. & Syd. 1915 (= *Pseudonectriella* Petr. 1959 *fide Species Fungorum* 2017), *Catabotrydaceae*, *Amplistromatales*, *Sordariomycetes*, one species, type: *C. palmarum* (Pat.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Catapyrenium Flot. 1850 (= *Catapyrenium* Flot. 1849; = *Endopyrenium* Flot. 1855; = *Involucrocarpone* Servít 1953 *fide Species Fungorum* 2017), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, six species, type: *C. cinereum* (Pers.) Körb., lichenized, see Lumbsch and Huhndorf (2010; outline), Prieto et al. (2010c; species in the Iberian Peninsula and the Balearic Islands), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Cataractispora K.D. Hyde, S.W. Wong & E.B.G. Jones 1999, *Annulatasaceae*, *Annulatascales*, *Sordariomycetes*, five species, type: *C. aquatica* K.D. Hyde, S.W. Wong & E.B.G. Jones, saprobes, aquatic, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: HKU (M) 5239, cultures and sequences are available.

Catarraphia A. Massal. 1860 (= *Cyclographa* Vain. 1921 *fide* Species Fungorum 2017), *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *C. dictyoplaca* (Mont. & Bosch) A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Catarrhospora Brusse 1994, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, two species, type: *C. mira* Brusse, lichenized, South Africa, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Catenarina Søchting, Søgaard, Arup, Elvebakk & Elix 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *C. desolata* Søchting, Søgaard & Elvebakk, lichenized, see Søchting et al. (2014b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Catenocuneiphora Matsush. 2003, *Ascomycota* genera *incertae sedis*, one species, type: *C. mucosa* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Catenophoropsis Nag Raj & W.B. Kendr. 1988, *Ascomycota* genera *incertae sedis*, one species, type: *C. eucalypticola* Nag Raj & W.B. Kendr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Catenosubulispora Matsush. 1971, *Ascomycota* genera *incertae sedis*, one species, type: *C. honiaraensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South East Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Catenosynnema Kodsueb, K.D. Hyde & W.H. Ho 2007, *Ascomycota* genera *incertae sedis*, one species, type: *C. micheliae* Kodsueb, K.D. Hyde & W.H. Ho, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Thailand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

holotype: PDD 87914, cultures and sequences are unavailable.

Catenularia Grove 1886, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, type: *C. simplex* Grove, hyphomycetous, sexual morph unknown, saprobes, terrestrial (wood), worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Catenulomyces Egidi & de Hoog 2014, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *C. convolutus* Egidi & Quaedvly., hyphomycetous, sexual morph unknown, rock-inhabiting, Spain, see Egidi et al. (2014; taxonomy, phylogeny), Wijayawardene et al. (2014c, 2017; outline, accepted as *Capnodiales* genera *incertae sedis*), ex-type cultures of the type species CBS 118609 = TRN9.

Catenulostroma Crous & U. Braun 2007, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *C. protearum* (Crous & M.E. Palm) Crous & U. Braun, hyphomycetous, sexual morph ?teratosphaeria-like, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Teratosphaeriaceae*), Quaedvlieg et al. (2014; phylogeny, introduced new genus *Neocatenulostroma*), Hernández-Restrepo et al. (2017; new species, phylogeny), holotype and ex-type of the type: BPI 1107938, CPC 18276.

Catillaria A. Massal. 1852 (= *Catillaria* A. Massal. ex R. Sant. 1952; = *Microlecia* M. Choisy 1949; = *Ulocodium* A. Massal. 1855 *fide* Species Fungorum 2017), *Catillariaceae*, *Lecanorales*, *Lecanoromycetes*, 150 species, type: *C. chalybeia* (Borrer) A. Massal., lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Catilochroma Kalb 2007, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, seven species, type: *C. endochroma* (Fée) Kalb, lichenized, see Lendemer (2010c; re-visit), Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Miadlikowska et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Catinaria Vain. [nom. cons.] 1922 (= *Biatorina* A. Massal. 1852 *fide* Species Fungorum 2017), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *C. atropurpurea* (Schaer.) Vězda & Poelt, lichenized, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Catinella Boud. 1907, *Dothideomycetes* genera *incertae sedis*, one or two species, type: *C. olivacea* (Batsch) Boud., asexual morph unknown, saprobes or parasitic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequence are available only for the type specie.

Catinopeltis Bat. & C.A.A. Costa 1957, *Ascomycota* genera *incertae sedis*, one species, type: *C. recifensis* Bat. & C.A.A. Costa, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Catolechia Flot. 1850 (= *Dimaura* Norman 1853 [1852]; = *Xanthopsis* Acloque 1893 *fide Species Fungorum* 2017), *Rhizocarpaceae*, *Rhizocarpales*, *Lecanoromycetes*, one species, type: *C. wahlenbergii* (Ach.) Flot., lichenized, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), type specimen UPS-ACH-L-244 (isotype), sequences are available.

Catulus Malloch & Rogerson 1978, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. aquilonius* Malloch & Rogerson, asexual morph unknown, saprobes, fungicolous, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Caudatispora J. Fröhl. & K.D. Hyde 1995, *Chaetosphaeriales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *C. palmicola* J. Fröhl. & K.D. Hyde, asexual morph unknown, terrestrial, saprobes, America (tropical), see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariomycetidae*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available but lacking for the type species.

Caudella Syd. & P. Syd. 1916, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, two species, type: *C. oligotricha* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline) cultures and sequences are unavailable.

Caudospora Starbäck 1889, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. taleola* (Fr.) Starbäck, asexual morph unknown, terrestrial, saprobes, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cavaraella Spieg. 1921, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *C. micraspis* (Berk. & M.A. Curtis) Spieg., asexual morph unknown, saprobes, terrestrial, Cuba, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cavimalum Yoshim. Doi, Dargan & K.S. Thind 1977, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *C. indicum* Yoshim. Doi, Dargan & K.S. Thind, asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cazia Trappe 1989, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *C. flexiascus* Trappe, asexual morph unknown, from soil, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available for voucher SOC 469 of the type species.

Cecidiomyces U. Braun & Zhurb. 2012, *Ascomycota* genera *incertae sedis*, one species, type: *C. santessonii* U. Braun & Zhurb., hyphomycetous, sexual morph unknown, terrestrial, lichenicolous, Australia, see Zhurbenko and Braun (2012; taxonomy), Wijayawardene et al. (2017; outline), holotype of type: UPS F-552770, cultures and sequences are unavailable.

Cecidioskyttea Etayo 2004, ?*Lecanoromycetes* genera *incertae sedis*, one species, type: *C. osorioi* Etayo, asexual morph coelomycetous, sexual morph stromatic and hairy and spores simples, parasymbiont on macrolichens belonging to *Parmeliaceae*, South America, see Suija et al. (2015c; taxonomy), cultures and sequences are unavailable, holotype of the type: Hb. Osorio, H.S. Osorio & G. Geymonat 9639a.

Cecidonia Triebel & Rambold 1988, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, two species, type: *C. umbonella* (Nyl.) Triebel & Rambold, asexual morph unknown, lichenicolous, Europe, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Ceeveesubramaniomyces Pratibha, K.D. Hyde & Bhat 2005, *Ascomycota* genera *incertae sedis*, one species, type: *C. litsea* Pratibha, K.D. Hyde & Bhat 2005, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: GUBH P 198, cultures and sequences are unavailable.

Cepia Velen. 1934, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *C. coerulea* Velen., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013;

genus accepted), cultures and sequences are available but lacking for the type.

Celerioriella Crous 2015, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, three species, type: *C. dura* (Damm & Crous) Crous, coelomycetous, sexual morph unknown, in necrotic wood of *Prunus* spp., pathogens, terrestrial, South Africa, see Damm et al. (2010b) as *Phaeomoniella dura* Damm & Crous, morphology), Crous et al. (2015d, 2016a; phylogeny), holotype and ex-type: CBS H-19999, CBS 120882.

Celoporthes Nakab., Gryzenh., Jol. Roux & M.J. Wingf. 2006, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *C. dispersa* Nakab., Gryzenh., Jol. Roux & M.J. Wingf., asexual morph coelomycetous, pathogenic, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Chen et al. (2011; species from *Eucalyptus* and *Syzygium* trees in China and Indonesia), Crous et al. (2012d; phylogeny), Vermeulen et al. (2013; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), holotype and ex-epitype strain of the type: PREM 58896, CMW 9976.

Celothelium A. Massal. 1860, *Celotheliaceae*, *Phaeomoniellales*, *Eurotiomycetes*, eight species, type: *C. socialis* (Zenker) A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Gueidan et al. (2014b; taxonomy, DNA), Chen et al. (2015a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Celtidia J.D. Janse 1897, *Zopfiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *C. duplicitispora* J.D. Janse, asexual morph unknown, marine habitat (root nodules of *Celtis* in Java), see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Zopfiaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cenangiopsis Rehm 1912 (= *Pyrenopezizopsis* Höhn. 1917 *fide* Species Fungorum 2017), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *C. quercicola* (Romell) Rehm, asexual morph unknown, saprobes, terrestrial, China, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärte et al. (2017; DNA, phylogeny), cultures available for the type: KL174.

Cenangium Fr. 1818 (= *Cenangina* Höhn. 1909; = *Cenangium* trib. *Clithris* Fr. 1822; = *Clithris* (Fr.) Bonord. 1851 *fide* Species Fungorum 2017), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, c. 30 species, type: *C. ferruginosum* Fr., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärte et al. (2017; DNA, phylogeny), cultures available for the type: KL390.

Cenangiumella J. Fröhl. & K.D. Hyde 2000, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. rattanicola* J. Fröhl. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Brunei, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Cenococcum Moug. & Fr. 1829, *Gloniaceae*, *Mytilinidiales*, *Dothideomycetes*, nine species, type: *C. geophilum* Fr., coelomycetous, sexual morphs unknown, ectomycorrhizal, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as in *Gloniaceae*), Kirk et al. (2013; genus accepted), Peter et al. (2016; DNA, ectomycorrhizal), cultures and sequences are available.

Cenozosia A. Massal. 1854, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *C. inanis* (Mont.) A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences unavailable.

Cephaliophora Thaxt. 1903 (= *Cephalomyces* Bainier 1907 *fide* Species Fungorum 2017), *Pezizales* genera *incertae sedis*, four species, type: *C. tropica* Thaxt., hyphomycetous, sexual morph unknown, saprobes, on dung, soil, terrestrial, pantropical, temperate, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Cephaloascus Hanawa 1920 (= *Ascocybe* D.E. Wells 1954; = *Aureomyces* Ruokola & Salonen 1970 *fide* Species Fungorum 2017), *Cephaloascaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *C. dispersa* Nakab., Gryzenh., Jol. Roux & M.J. Wingf., asexual reproduction by budding and blastoconidia, saprophytic, although one strain of *C. fragrans* was isolated from a human ear, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kurtzman and Robnett (2013b; DNA, phylogeny), cultures and sequences are available.

Cephalophysis (Hertel) H. Kilias 1985 (= *Lecidea* subgen. *Cephalophysis* Hertel 1967 *fide* Species Fungorum 2017), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *C. leucospila* (Anzi) H. Kilias & Scheid., asexual morph unknown, lichenized, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Arup et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Cephalotheca Fuckel 1871 (= *Aposphaeriopsis* Died. 1913; = *Carothecis* Clem. 1931; = *Crepinula* Kuntze 1891; *Erythrophaera* Sorokin 1871 *fide* Species Fungorum 2017), *Cephalothecaceae*, *Sordariomycetes* families *incertae sedis*, six species, type: *C. sulfurea* Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide,

see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Cephalotrichiella Crous 2014, *Microascales*, genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. penicillata* Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Netherlands, see Crous et al. (2014c; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), holotype and ex-type: CBS H-21725, CPC 23865.

Cephalotrichum Link 1809 (= *Berkeleya* Kuntze 1898; = *Capnostysanus* Speg. 1918; = *Doratomyces* Corda 1829; = *Stelechotrichum* Ritgen 1831; = *Stysanopsis* Ferraris 1909; = *Stysanus* Corda 1837; = *Synpenicillium* Costantin 1888; = *Trichurus* Clem. 1896 *fide* Seifert et al. 2011; Species Fungorum 2017), *Microascaceae*, *Microascales*, *Sordariomycetes*, 31 species, type: *C. stemonitis* (Pers.) Nees, hyphomycetous, sexual morph unknown, saprobes, soil, dung terrestrial, worldwide, see Jiang et al. (2011; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), de Beer et al. (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Sandoval-Denis et al. (2016a; new species), cultures and sequences are available.

Ceramoclasteropsis Bat. & Cavalc. 1962, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. coumae* Bat. & Cavalc., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Capnodiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Ceramothyrium Bat. & H. Maia 1956, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, 34 species, type: *C. paiveae* Bat. & H. Maia, asexual morph hyphomycetous, *Stanughesia* Constant., epiphytic, epiphyllous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2012b, 2014; morphology, phylogeny), Kirk et al. (2013; genus accepted), Zeng et al. (2016; morphology, phylogeny), cultures and sequences of type species are unavailable.

Ceratocarpia Rolland 1896, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, two species, type: *C. cactorum* Rolland, asexual morph unknown, saprobes, terrestrial, America, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Tian et al. (2014; morphology; nomenclature), cultures and sequences are unavailable.

Ceratocladium Corda 1839, *Ascomycota* genera *incertae sedis*, two species, type: *C. microspermum* Corda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ceratocystiopsis H.P. Upadhyay & W.B. Kendr. 1975, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, eleven species, type: *C. minuta* (Siemaszko) H.P. Upadhyay & W.B. Kendr., sexual morph unknown, saprobes, terrestrial, worldwide, see Plattner et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Ceratocystis Ellis & Halst. 1890 (= *Chaetonaemosphaera* Schwarzman 1968); = *Milowia* Massee 1884; = *Mycorrhynchella* Höhn. 1918; = *Rostrella* Fabre 1879 [1878]; = *Rostrella* Zimm. 1900 *fide* Species Fungorum 2017), *Ceratocystidaceae*, *Microascales*, *Sordariomycetes*, c. 70 species, type: *C. fimbriata* Ellis & Halst., asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), van Wyk et al. (2009, 2010, 2011; new species), Kajitani and Masuya (2011; new species, pathogens), Tarigan et al. (2011; new species, Indonesia), Nkuekam et al. (2012; new species), Chen et al. (2013a; species associated with Eucalyptus, China), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Crous et al. (2015d; new species, phylogeny), cultures and sequences are available.

Ceratolenta Réblová 2013, *Sordariomycetidae* genera *incertae sedis*, *Sordariomycetes*, eight species, type: *C. caudata* Réblová, asexual morph unknown, saprobes, Europe, see Réblová et al. (2013; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype and ex-type cultures available for the type: PRM 899854, CBS 125234.

Ceratomyces Thaxt., 1892, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 32 species, type: *C. mirabilis* Thaxt., parasitic on Hydrophilidae and Staphylinidae beetles, terrestrial and aquatic, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence is available for the type: AF431764 (SSU).

Ceratopachnidium J. Reid & Piroz. 1966, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *C. aristosporum* (Bonar) J. Reid & Piroz., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ceratophoma Höhn. 1917, *Ascomycota* genera *incertae sedis*, one species, type: *C. rostrata* (Fuckel) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ceratophorum Sacc. 1880, *Ascomycota* genera *incertae sedis*, two species, type: *C. helicosporium* (Sacc.) Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ceratoportha Petr. 1925, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. didymospora* Petr., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Ceratopycnis Höhn. 1915 (= *Rhynchophorus* Hollós 1926 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *C. clematidis* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy, new species), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ceratospermopsis Bat. 1951, *Ascomycota* genera *incertae sedis*, two species, type: need typification data, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded from *Meliolaceae*), cultures and sequences are available but lacking for the type species.

Ceratosphaerella Huhndorf, Greif, Mugambi & A.N. Mill. 2008, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *C. castillensis* (C.L. Sm.) Huhndorf, Greif, Mugambi & A.N. Mill., asexual morph hyphomycetous, saprobes, terrestrial, Costa Rica, Nicaragua, Puerto Rico, see Huhndorf et al. (2008; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type culture of type: SMH1865.

Ceratosphaeria Niessl 1876, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, twelve species, type: *C. lampadophora* (Berk. & Broome), asexual morph harpophora-like, saprobes, terrestrial, worldwide, see Huhndorf et al. (2008; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), culture available for type: CBS 117555.

Ceratospora Höhn. 1923, *Ascomycota* genera *incertae sedis*, 20 species (14 valid taxa as per Monteiro and Gusmao 2014), type: *C. bicornis* Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), Monteiro and Gusmao (2014; key to species), sequence available but lacking for the type species.

Ceratosporium Schwein. 1832, *Ascomycota* genera *incertae sedis*, c. 14 species, type: *C. fuscescens* Schwein, widespread, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus not accepted), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Ceratostomella Sacc. 1878 (= *Cerastomis* Clem. 1931; = *Rhynchophoma* P. Karst. 1884 *fide* Species Fungorum 2017), *Sordariomycetidae* genera *incertae sedis*, *Sordariomycetes*, 18 species, type: need typification, asexual morph unknown, saprobes, terrestrial, temperate, see Réblová and Štěpánek (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Cercidospora Körb. 1865 (= *Neonorrlinia* Syd. 1923; = *Norrlinia* Vain. 1921; = *Pleosphaeropsis* Vain. 1921; = *Prolisea* Clem. 1931; = *Vainiona* Werner 1943 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, more than 100 species, type: *C. ulothii* Körb., on lichens, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Cercophora Fuckel 1870 (= *Bovilla* Sacc. 1883; = *Heteronectria* Penz. & Sacc. 1898 [1897]; = *Lasiosordaria* Chenant. 1919; = *Lasiosordariopsis* Chenant. 1919 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, 51 species, type: *C. mirabilis* Fuckel, asexual morph unknown, saprobes, terrestrial, Europe, see Chang et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline), Catania et al. (2011; new species), Kirk et al. (2013; genus accepted), Kruys et al. (2015; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Crous et al. (2016a, b; new species), Dai et al. (2017; new species, phylogeny), cultures and sequences are available.

Cercosperma G. Arnaud ex B. Sutton & Hodges 1983, *Ascomycota* genera *incertae sedis*, two species, type: *C. arnaudii* B. Sutton & Hodges, hyphomycetous, sexual morph unknown, saprobes, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Cercospora Fresen. 1863 (= *Cercosporina* Speg. 1911; = *Virgasporium* Cooke 1875 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, about 3000 described species (c. 700 accepted species), and 357 orthographic variants (MycoBank), type: *C. apii* Fresen., hyphomycetous, saprobes, plant pathogens,

terrestrial, worldwide, see Groenewald et al. (2008, 2013; sexual reproduction, phylogeny), To-Anun et al. (2010, 2011; phylogeny, key), Hesami et al. (2011; phylogeny, new species), Seifert et al. (2011; morphology), Phengsintham et al. (2013; species in Laos), Hyde et al. (2013; accepted as a genus in *Mycosphaerellaceae*), Wijayawardene et al. (2014c, 2017; outline), Bakhshi et al. (2015a, b; phylogeny), Nguanhom et al. (2015; phylogeny), Braun and Crous (2016; nomenclature), sequences are available for most of the species, holotype: *Fungi rhen.* 117.

Cercospora Sacc. 1880, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 40 species, type: *C. cana* (Sacc.) Sacc., hyphomyceteous, sexual morph mycosphaerella-like, plant pathogen, terrestrial, worldwide, see Kirschner (2009; differentiation from *Ramularia*), Seifert et al. (2011; morphology), Braun (2013a; taxonomy), Braun and Urtiaga (2013; new species), Braun et al. (2015; new species, key), Crous et al. (2014c; new species), Machado et al. (2012; pathology), Zhang et al. (2013b; secondary metabolites), Videira et al. (2016; morphology, phylogeny), lectotype of type: Sacc., Mycoth. Ven. 593 (K), non-European reference cultures CPC 19492, CPC 11461, CBS 113304.

Cercosporidium Earle 1901, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 70 species, of which five are confirmed based on DNA, type: *C. euphorbiae* Tracy & Earle, hyphomyceteous, sexual morph unknown, plant pathogenic, see Braun et al. (2013; morphology), Videira et al. (2017; morphology, phylogeny), resurrect the genus, and designate an epitype.

Cerillum Clem. 1931, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. paradoxa* (Har. & Pat.) Clem., terrestrial, saprobes, see Wendt et al. (2017; resurrected the genus in *Xylariales*), cultures and sequences are available.

Cerion Massee 1901, *Rhytidomycetaceae*, *Rhytidomycetales*, *Leotiomycetes*, two species, type: *C. coccineum* Massee & Rodway, asexual morph unknown, saprobes, terrestrial, Australia, America (tropical), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ceriophora Höhn. 1919, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type species: *C. palustris* (Sacc.) Höhn., Europe, doubtful genus, see Kirk et al. (2008; genus accepted), Lumbsch and Huhndorf (2010; outline), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ceriospora Niessl 1876 (= *Ceriophora* Höhn. 1919; = *Hindersonia* Moug. & Nestl. ex J. Schröt. 1897 [1908] *fide* Species Fungorum 2017), *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, 18 species, type species: *C. dubyi*

Niessl., doubtful genus, see Kirk et al. (2008; 2013), Lumbsch and Huhndorf (2010; outline), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b; outline), cultures and sequences are unavailable.

Ceriosporopsis Barghoon & Linder 1944, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, four species, type: *C. halima* Linder, asexual morph not unknown, saprobes, marine, worldwide, see Jones et al. (2009b, 2015; outline, phylogeny), Sakayaroj et al. (2011a; new species, phylogeny, however this species is invalid *fide* Index Fungorum 2017), Abdel-Wahab et al. 2017; new species), sequence data available for *C. halima* (BCC15553).

Cerodothis Muthappa 1969, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. aurea* Muthappa, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cerothallia Arup, Frödén & Søchting 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, four species, type: *C. luteoalba* (Turner) Arup, Frödén & Søchting, lichenized, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Cesariella W. Rossi & Santam. 2008, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *C. graeca* W. Rossi & Santam., asexual morph unknown, parasitic on Carabidae beetles, terrestrial, Europe, Rossi and Santamaría (2008; taxonomy, did not explicitly state family level placement, but suggested very close relationship to *Laboulbenia*), Lumbsch and Huhndorf (2010; outline, accepted as *Laboulbeniales incertae sedis*), cultures and sequences are unavailable.

Cesatiella Sacc. 1878, *Hypocreaceae*, *Amphisphaerales*, *Sordariomycetes*, four species, type species: *C. australis* Sacc. & Speg., see Kirk et al. (2008, 2013; genus accepted), Lumbsch and Huhndorf (2010; outline), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cetradonia J.C. Wei & Ahti. 2002, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *C. linearis* (A. Evans.) J.C. Wei & Ahti., asexual morph unknown, lichenized, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Cetraria Ach. 1803 nom. cons. (= *Cetrariomyces* E.A. Thomas 1939; = *Cetrariella* Kärnefelt & A. Thell, 1993; = *Coelocaulon* Link 1833; = *Geissodea* Vent. 1799; = *Platism* Hoffm. 1796 [1795]; = *Platiphyllum* Vent. 1799; = *Pseudocornicularia* Gyeln. 1933; = *Squamaria* Hoffm. 1789 [1790] = *Usnocetraria* M.J. Lai & J.C. Wei 2007; = *Vulpicida* J.-E. Mattsson & M.J. Lai 1993 *fide*

Divakar et al. 2017; Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 21 species, type: *C. islandica* (L.) Ach., asexual morph unknown, epiphytic or saxicolous, widespread, see Thell et al. (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Cetrelia W.L. Culb. & C.F. Culb. 1968, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, c. 18 species, type: *C. cetrariooides* (Delise) W.L. Culb. & C.F. Culb., asexual morph unknown, lichenized, terrestrial or on lignum, widespread, see Lumbsch and Huhndorf (2010; outline), Crespo et al. (2010b; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Cetreliopsis M.J. Lai. 1980, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, seven species, type: *C. rhytidocarpa* (Mont. & Bosch) Lai, asexual morph unknown, lichenized, epiphytic, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Ceuthodiplospora Died. 1912, *Ascomycota* genera *incertae sedis*, one species, type: *C. robiniae* (Roberge) Died., asexual morph coelomycetous, terrestrial, Czech Republic, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Ceuthosira Petr. 1924, *Ascomycota* genera *incertae sedis*, one species, type: *C. aesculicarpa* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ceuthosporella Petr. & Syd. 1923 (= *Microcyta* Petr. & Syd. 1927 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *C. sambuci* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chadefaudia Feldm.-Maz. 1957 (= *Mycophycophila* Cribb & J.W. Cribb 1960 *fide* Species Fungorum 2017), *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, six species, type: *C. marina* Feldm.-Maz., saprobes, ?marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline of marine genera), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Chadefaudiella Faurel & Schotter 1959, *Chadefaudiellaceae*, *Microascales*, *Sordariomycetes*, two species, type: *C. quezelii* Faurel & Schotter, saprobes,

coprophilous, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Chadefaudiomyces Kamat, V.G. Rao, A.S. Patil & Ullasa 1974, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. indicus* Kamat, V.G. Rao, A.S. Patil & Ullasa, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Chaenocarpus Rebent. 1804 (= *Chaenocarpus* Spreng. 1806; = *Chaenocarpus* Lév. 1843; = *Coenocarpus* Fr. 1825; = *Cryptothamnium* Wallr. 1842; = *Simoninus* Roum. 1879 *fide* Species Fungorum 2017; Wendt et al. 2017), *Xylariales* genera *incertae sedis*, *Sordariomycetes*, three species, type: *C. setosus* (Roth) Rebent., asexual morph unknown, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), Wendt et al. (2017; accepted as *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Chaenotheca (Th. Fr.) Th. Fr. 1860 (= *Allodium* Nyl. 1896 [1895]; = *Calicum* b *Chaenotheca* Th. Fr. 1856; = *Chaenothecomyces* Cif. & Tomas. 1953; = *Cybebe* Tibell 1984; = *Cyphelium* Chevall. 1826; = *Embolus* Batsch 1786; = *Eucyphelis* Clem. 1909; = *Heydeniopsis* Naumov 1915 *fide* Species Fungorum 2017), *Coniothybaceae*, *Coniothybales*, *Coniothybomycetes*, c. 25 species, type: *C. trichialis* (Ach.) Th. Fr., asexual morph unknown, lichenized, epiphytic, widespread but mostly Northern Boreal, see Lumbsch and Huhndorf (2010, outline), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Chaenothecopsis Vain. 1927 (= *Catenomycopsis* Tibell & Constant. 1991 = *Chaenotheciella* Räsänen 1943; = *Pseudocalicium* Marchand 1896; = *Strongyleuma* Vain. 1927 *fide* Species Fungorum 2017; Art. 59), *Mycocaliciaceae*, *Mycocaliciales*, *Eurotiomycetes*, 27 species, type: *C. rubescens* Vain., asexual morph coelomycetous or hyphomycetous, lichenicolous, fungicolous, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Seifert et al. (2011; morphology of *Catenomycopsis*), Tuovila et al. (2011; new species, Oregon), Kirk et al. (2013; genus accepted), Selva (2013; new species), Hawksworth et al. (2014; new species), Rikkinen et al. (2014; new species), cultures and sequences are available but lacking for the type.

Chaetarthriomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *C. flexatus* Thaxt., asexual morph unknown, parasitic on

Hydrophilidae beetles, aquatic, widespread, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2014b; Netherlands), cultures and sequences are unavailable.

Chaetasbolisia Speg. 1918, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *C. erysiphoides* Griffon & Maubl., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see de Gruyter et al. (2009; DNA, phylogeny, Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 148.94.

Chaetendophragmia Matsush. 1971, *Ascomycota* genera *incertae sedis*, six species, type: *C. triangularis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetendophragmiopsis B. Sutton & Hodges 1978, *Ascomycota* genera *incertae sedis*, two species, type: *C. pulchra* Sutton & Hodges, hyphomycetous, sexual morph unknown, two species, saprobes, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Chaetoamphispshaeria Hara 1918, *Sordariomycetes*, genera *incertae sedis*, two species, type: *C. japonica* Hara, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Chaetoblastophorum Morgan-Jones 1977, *Ascomycota* genera *incertae sedis*, one species, type: *C. ingramii* Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetocapnodium Hongsanan & K.D. Hyde 2015, *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *C. siamensis* Hongsanan & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Liu et al. (2015a; taxonomy, phylogeny), holotype and ex-type: MFLU 14-0747, MFLUCC 13-0778.

Chaetoconis Clem. 1909 (= *Amphorula* Grove 1922 *fide* Species Fungorum 2017), *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. polygoni* (Ellis & Everh.) Clem., coelomycetous, sexual morph unknown, saprobes, terrestrial, north temperate, see de Gruyter et al. (2009; DNA, phylogeny), Kirk et al. (2013;

genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures available for the type: CBS 405.95.

Chaetocrea Syd. 1927, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. parasitica* Syd., asexual morph unknown, fungicolous associated with black areas of fungi on leaves, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; excluded from *Tubeufiaceae*), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Chaetocylostroma Petr. 1920, *Ascomycota* genera *incertae sedis*, one species, type: *C. arundinacea* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetodiplis Clem. 1931, *Ascomycota* genera *incertae sedis*, one species, type: *C. hirta* (Sacc.) Clem., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetodiplodia P. Karst. 1884, *Ascomycota* genera *incertae sedis*, nine species, type: *C. caulinia* P. Karst., coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Chaetodiplodina Speg. 1911, *Ascomycota* genera *incertae sedis*, two species, type: *C. graminicola* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetomastia (Sacc.) Berl. 1890 (= *Herpothrix* Clem. 1909; = *Melanomma* subgen. *Chaetomastia* Sacc. 1883 *fide* Species Fungorum 2017), *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, ten species, type: *C. hirtula* (P. Karst.) Berl., asexual morph unknown, terrestrial, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Teichosporaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Chaetomella Fuckel 1870 (= *Harikrishnaella* D.V. Singh & A.K. Sarbhoy 1972 [1971]; = *Volutellospora* Thirum. & P.N. Mathur 1965 [1964]; = *Zoellneria* Velen. 1934 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Chaetomellaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *C. oblonga* Fuckel, asexual morph coelomycetous,

saprobes, in soil, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species, Zambia), Johnston et al. (2014b; nomenclature), cultures available for the type: BPI 843552.

Chaetomidium (Zopf) Sacc. 1882 (= *Chaetomium* subgen. *Chaetomidium* Zopf 1881 *fide* Species Fungorum 2017), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, 16 species, type: *C. fimetii* (Fuckel) Zopf, asexual morph phialophora-like, saprobes, in soil, coprophilous, worldwide, see Greif et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 168.71.

Chaetomiopsis Moustafa & Abdul-Wahid 1990, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *C. dinae* Moustafa & Abdul-Wahid, asexual morph unknown, saprobes, Egypt, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a distinct genus), cultures and sequences are unavailable.

Chaetomium Kunze, in Kunze & Schmidt 1817 (= *Achaetomiella* Arx 1970; = *Bolacotricha* Berk. & Broome 1851; = *Bommerella* Marchal, 1885; = *Botryotrichum* Sacc. & Marchal 1885; *Chaetomiotricha* Peyronel 1914; = *Cladochaete* Sacc. 1912; = *Trichocladium* Harz 1871; = *Vanhallia* L. Marchand 1828 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, more than 150 species., type: *C. globosum* Kunze, asexual morph hyphomycetous, saprophytic fungus that primarily resides on plants, soil, straw, and dung, potential biological control agents, worldwide, see Ahmed et al. (2012; enzymes), Mungai et al. (2012a; from dung), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), Wang et al. (2016d; taxonomy), cultures and sequences are available.

Chaetomyces Thaxt. 1938, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *C. pinophilii* Thaxt., asexual morph unknown, parasitic on Staphylinidae beetles, terrestrial, North and South America, Rossi and Bergonzo (2008; Brazil), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetonaevia Arx 1951, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *C. nannfeldtii* Arx, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetonectrioides Matsush. 1996, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. malaysiana* Matsush., saprobes, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: 5T121 (Matsushima Fungus Collection, Kobe), cultures and sequences are unavailable.

Chaetopeltaster Katum. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *C. japonicus* Katum., coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetophiophoma Speg. 1911, *Ascomycota* genera *incertae sedis*, one species, type: *C. trematis* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetophoma Cooke 1878, *Ascomycota* genera *incertae sedis*, c. 30 species, type: *C. quercifolia* Cooke, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetoplaca Syd. & P. Syd. 1917, *Ascomycota* genera *incertae sedis*, one species, type: *C. memecyli* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Schizophyriaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013, accepted as a genus in *Schizophyriaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Schizophyriaceae*), Phookamsak et al. (2016; taxonomy, transferred to *Ascomycota* genera *incertae sedis*), cultures and sequences are unavailable.

Chaetoplea (Sacc.) Clem. 1931 (= *Parahendersonia* A.W. Ramaley 1995; = *Pyrenophora* subgen. *Chaetoplea* Sacc. 1883 *fide* Wijayawardene et al. 2016b; Species Fungorum 2017), *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, c. 17 species, type: *C. calvescens* (Fr.) Clem., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as *Phaeosphaeriaceae*), Phookamsak et al. (2014c; morphology, accepted as *Leptosphaeriaceae*), Wijayawardene et al. (2014c, 2016b; outline, morphology of asexual morph), sequence data and cultures are unavailable.

Chaetopreussia Locq.-Lin. 1977, *Sporormiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *C. chadefaudii* Locq.-Lin., asexual morph unknown, saprobes, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as *Sporormiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Chaetopsina Rambelli 1956 (= *Chaetopsinectria* J. Luo & W.Y. Zhuang 2010 *fide* Rossman et al. 2016a), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, 18 species, type: *C. fulva* Rambelli, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Nagaraju et al. (2011a; new species), Crous et al. (2013b, 2014c; morphology, new species), Lombard et al. (2015b; DNA), Maharachchikumbura et al. (2016b; outline), Rossman et al. (2016a; nomenclature), Wijayawardene et al. (2017; outline), available cultures of type: CBS 142.56.

Chaetopsis Grev. 1825 (= *Chaetopsella* Höhn. 1930 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, seven species, type: *C. wauchii* Grev., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetopyrena Pass. 1881 (= *Sclerochaeta* Höhn. 1917 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *C. poae* (Niessl) Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetorostrum Zelski, Raja, A.N. Mill. & Shearer 2011, *Annulatasaceae*, *Annulatascales*, *Sordariomycetes*, one species, type: *C. quincemilense* Zelski, Raja, A.N. Mill. & Shearer, asexual morph unknown, saprobes, aquatic, Peru, see Zelski et al. (2011b; morphology), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: ILL 40822, cultures and cultures and sequences are unavailable.

Chaetosclerophoma Petr. 1924, *Ascomycota* genera *incertae sedis*, two species, type: *C. coluteae* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Pakistan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetoscutula E. Müll. 1958, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *C. juniperi* E. Müll., asexual morph unknown, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Chaetoseptoria Tehon 1937, *Ascomycota* genera *incertae sedis*, one species, type: *C. vignae* Tehon, coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences and cultures are unavailable.

Chaetosphaerella E. Müll. & C. Booth 1972 (= *Cladotrichum* Corda 1831; = *Dimera* Fr. 1825; = *Diplosporium* Link 1824; = *Gongylocladium* Wallr. 1833; = *Jacobaschella* Kuntze 1891; = *Oedemium* Link 1824; = *Sphinctrosporium* Kunze 1849; = *Veramyces* Subram. 1994; = *Veramycina* Subram. 1995 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Chaetosphaerellaceae*, *Coronophorales*, *Sordariomycetes*, 14 species, type: *C. phaeostroma* (Durieu & Mont.) E. Müll. & C. Booth, hyphomycetous, *Chaetosphaerella* sexual morph reported for *E. didymum* (= *E. atrum*) and *E. minus*, while *Acrogenotheca* sexual morph reported for *E. robustum*, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Réblová et al. (2016c; nomenclature).

Chaetosphaeria Tul. & C. Tul. 1863 (= *Aposphaeriella* Died. 1912; = *Chaetolentomita* Maubl. 1915; = *Didymopsamma* Petr. 1925; = *Lentomita* Niessl 1876; = *Melanopsamma* Höhn. 1919; = *Miyoshia* Kawam. 1907; = *Miyoshiella* Kawam. 1929; = *Montemartinia* Curzi 1927; = *Trichocollonema* Höhn. 1902; = *Urnularia* P. Karst. 1866 *fide* Species Fungorum 2017), *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, more than 100 species, type: *C. innumera* Berk. & Broome ex Tul. & C. Tul., asexual morph complex, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Perera et al. (2016b; new species), cultures and sequences are available.

Chaetosphaerides Matsush. 2003, *Sordariomycetidae* genera *incertae sedis*, one species, type species: *C. ramichloridifera* Matsush., saprobes, terrestrial, Japan, see Maharachchikumbura et al. (2015b, 2016b; outline, accepted as *Sordariomycetidae* genera *incertae sedis*), holotype: MFC 210, sequences and cultures are unavailable.

Chaetosphaeronema Moesz 1915 (= *Pseudophoma* Höhn. 1916 *fide* Species Fungorum 2017), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, eleven species, type: *C. hispidulum* (Corda) Moesz, asexual morph coelomycetous, saprobes or plant pathogenic, widespread, see de Gruyter et al. (2010; phylogeny), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Quaedvlieg et al. (2013; phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequence are available.

Chaetosphaerulina I. Hino 1938 (= *Thaxteriellopsis* Sivan. 1977 [1976] *fide* Species Fungorum 2017), ?*Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, six species, type: *C. yasudai* I. Hino, asexual morph hyphomycetous, saprobes on dead wood, see Lumbsch and Huhndorf (2010; phylogeny).

outline), Kirk et al. (2013; genus accepted), Boonmee et al. (2011, 2014b; treated as a doubtful genus but listed under *Tubeufiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Chaetosticta Petr. & Syd. 1925 (= *Pyrenochaeta* subgen. *Trichocicinnus* Sacc. 1906 [1905]; = *Trichocicinnus* (Sacc.) Höhn. 1919 *fide* Species Fungorum 2017), *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, three species, type: *C. perforata* (Ellis & Everh.) Petr. & Syd., asexual morph unknown, saprobes or parasitic, terrestrial, Europe, Japan, North America and South America, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequence data unavailable, need adequate sequence data.

Chaetotheca Zukal 1890, *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, one species, type: *C. fragilis* Zukal, asexual morph unknown, terrestrial, on bile concretion, Austria, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, needs phylogenetic assessment.

Chaethiersia B.A. Perry & Pfister 2008, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *C. vernalis* B.A. Perry & Pfister, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), holotype of the type: FH, Perry 492, cultures and sequences are available but lacking for the type species.

Chaetothyrina Theiss. 1913 (= *Ceratochaetopsis* F. Stevens & Weedon 1927; = *Chaetopeltopsis* Theiss. 1913; = *Plochmopeltidella* J.M. Mend. 1925; = *Setopeltis* Bat. & A.F. Vital 1959 *fide* Species Fungorum 2017), *Phaeothecoidiellaceae*, *Dothideomycetes* families *incertae sedis*, six species, type: *C. musarum* (Speg.) Theiss., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Singtripop et al. (2016; taxonomy), Hongsanan et al. (2016c, 2017b; new species, phylogeny, new family placement, evolution), cultures available for the type: MFLUCC 15-0383.

Chaetothyriomyces Pereira-Carv., Inácio & Dianese 2009, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *C. brasiliensis* Pereira-Carv., Inácio & Dianese, asexual morph unknown, epiphytic, terrestrial, Brazil, see Pereira-Carvalho et al. (2009a; morphology), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chaetothyriothecium Hongsanan & K.D. Hyde 2014, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *C. elegans* Hongsanan & K.D. Hyde, asexual morph unknown, epiphytes, Thailand, see Hongsanan et al.

(2014b, 2016c; taxonomy, phylogeny, evolution), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Chaetothyrium Speg. 1888 (= *Almeidaea* Cif. & Bat. 1962; = *Chaetasterina* Bubák 1909; = *Gilmania* Bat. & Cif. 1962; = *Sphaerochaetia* Bat. & Cif. 1962; = *Trotterula* Speg. 1921; = *Uloseia* Bat. 1963; = *Zukalia* Sacc. 1891 *fide* Species Fungorum 2017), *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, 66 species, type: *C. guaraniticum* Speg., asexual morph coelomycetous, epiphytic, sooty moulds, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2012b; morphology; phylogeny), Wijayawardene et al. (2012; 2017; outline), Kirk et al. (2013; genus accepted), Liu et al. (2015a; morphology; phylogeny), cultures and sequences are available for some species.

Chalara (Corda) Rabenh. 1844 (= *Chaetochalara* B. Sutton & Piroz. 1965; = *Cylindrocephalum* Bonord. 1851; = *Excioconidium* Plunkett 1925; = *Excioconis* Clem. & Shear 1931; = *Hughesiella* Bat. & A.F. Vital 1956; = *Stilbochalara* Ferd. & Winge 1910; = *Torula* subgen. *Chalara* Corda 1838 *fide* Species Fungorum 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, c. 100 species, type: *C. fusidioides* (Corda) Rabenh., hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, cosmopolitan, see Kowalski and Holdenrieder (2009; pathogens), Koukol (2011; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2016a, b; taxonomy, phylogeny), cultures and sequences are available.

Chalarodendron C.J.K. Wang & B. Sutton 1984, *Ascomycota* genera *incertae sedis*, one species, type: *C. fuscum* C.J.K. Wang & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chalarodes McKenzie 1991, *Ascomycota* genera *incertae sedis*, two species, type: *C. bisetis* McKenzie, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chalaropsis Peyronel 1916, *Ceratocystidaceae*, *Microascales*, *Sordariomycetes*, c. five species, type: *C. thielavioides* Peyronel, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see de Beer et al. (2014; revision, new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures are available.

Chalazion Dissing & Sivertsen 1975, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, three species, type: *C. sociabile* Dissing & Sivertsen, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype: FH, Perry 492, cultures and sequences are available but lacking for the type.

Chamaeascus L. Holm, K. Holm & M.E. Barr 1993, *Hypnoriaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *C. arcticus* L. Holm, K. Holm & M.E. Barr, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Chamaeleomyces Sigler 2010, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *C. granulomatis* Sigler, asexual morph unknown, reptile pathogen, terrestrial, America, see Sigler et al. (2010a; taxonomy), Schmidt et al. (2012; outline), Schmidt (2015; reptile pathogens), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Chantransiopsis Thaxt. 1914, *Ascomycota* genera *incertae sedis*, three species, type: *C. decumbens* Thaxt., coelomycetous, sexual morph unknown, entomogenous, terrestrial, Java, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chapeckia M.E. Barr 1978, *Sydiomycetaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *C. nigrospora* (Peck) M.E. Barr, asexual morphs unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kruys and Castlebury (2012; DNA, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Senanayake et al. (2017, placement), cultures available for the type: CBS 125532.

Chapsa A. Massal. 1860, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 62 species, type: *C. indica* A. Massal., asexual morph unknown, lichenized fungi, terrestrial, tropical, see Mangold et al. (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2010a, 2012a; key), Sipman et al. (2012; key, Costa Rica), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Characonidia Bat. & Cavalc. 1965, *Ascomycota* genera *incertae sedis*, one species, type: *C. ingae* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Charcotiana Søchting, Garrido-Benavent & Arup in Søchting et al. 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *C. antarctica* Søchting et al., asexual morph unknown, lichenized, saxicolous, Antarctica, see Søchting et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Charomyces Seifert 1987, *Ascomycota* genera *incertae sedis*, one species, type: *C. amphimelas* Seifert, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Hawaiian Island, Cuba, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences and cultures are unavailable.

Charonectria Sacc. 1880, *Hypnoriaceae*, *Amphisphaeriales*, *Sordariomycetes*, two species, type: *C. sceptrum* (P. Karst.) Lowen, temperate, dead leaves or lichenicolous, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013), Maharachchikumbura et al. (2015b, 2016b; outline), sequences and cultures are unavailable.

Chasakopama Manohar., Bagyanaryana, N.K. Rao & Kunwar 2009, *Ascomycota* genera *incertae sedis*, one species, type: *C. velgodensis* Manohar., Bagyanaryana, N.K. Rao & Kunwar, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Manoharacharya et al. (2009; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: IMI 296874, sequences are unavailable.

Cheilaria Lib. 1830, *Ascomycota* genera *incertae sedis*, one species, type: *C. agrostidis* Lib., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), isotype: IMI 152241, cultures and sequences are unavailable.

Cheilymenia Boud. 1885 (= *Coprobia* Boud. 1885 *fide* Species Fungorum 2017, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, c. 66 species, type: *C. stercorea* (Pers.) Boud., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Cheiroidea W.A. Baker & Morgan-Jones 2009, *Ascomycota* genera *incertae sedis*, one species, type: *C. triarmata* (Whitton, McKenzie & K.D. Hyde) W.A. Baker & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial (on *Pandanus*), Mauritius, see Gams et al. (2009; morphology), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: HKU(M) 13034, cultures and sequence unavailable.

Cheiromycella Höhn. 1910, *Ascomycota* genera *incertae sedis*, three species, type: *C. speiroidea* (Höhn.) Höhn., hyphomycetous, sexual morph unknown, on cheese and fungi, terrestrial, Europe, North America, see Braun et al.

(2009; new species), Seifert et al. (2011; morphology, treated as a genus in *Hyaloscypheaceae*), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cheiromyceopsis Mercado & J. Mena 1988, *Ascomycota* genera *incertae sedis*, one species, type: *C. echinulata* Mercado & J. Mena, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), sequences and cultures are unavailable.

Cheiromycinia B. Sutton 1986, *Ascomycota* genera *incertae sedis*, type: *C. flabelliformis* B. Sutton, five species, hyphomycetous, sexual morph unknown, saprobes, lichenized, Austria, Russia, Sweden, see Diederich et al. (2008; treated *C. ananas* Aptroot & Schiebelb. as a synonym of *Dictyocatenulata alba* Finley & E.F. Morris), An et al. (2012; morphology), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype: IMI288326.

Cheiropolyschema Matsush. 1980, *Ascomycota* genera *incertae sedis*, two species, type: *C. formosana* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, Taiwan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: MFC-6717, sequences and cultures are unavailable.

Cheirospora Moug. & Fr. 1825 (= *Hyperomyxa* Corda 1839; = *Moronopsis* Delacr. 1891; = *Myriocephalum* De Not. ex Corda 1842; = *Rabdosporium* Chevall. 1826; = *Thyrsidium* Mont. 1848 [1846-49] fide Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *C. botryospora* (Mont.) Berk. & Broome, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Crous et al. (2015c; morphology, phylogeny, DNA), Wijayawardene et al. (2016b; morphology, phylogeny, outline), reference specimens available for type: MFLU 15-3550, cultures available: CPC 24603 = CBS 139999, CPC 24605, CPC 24607, CPC 24611.

Cheiosporium L. Cai & K.D. Hyde 2008, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *C. triseriale* L. Cai & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, Asia, Egypt, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Abdel-Aziz (2016b; new species), Boonmee et al. (2016a; phylogeny), cultures and sequences are available.

Chevalieropsis Arnaud. 1923 (= *Chevalieria* Arnaud 1920; = *Parodiellinopsis* Hansf. 1946 [1944-45] fide Species Fungorum 2017), *Perisporiopsidaceae*, *Dothideomycetes* genera *incertae sedis*, two species, type: *C. ctenotricha* (Pat. & Har.) G. Arnaud, asexual morph

hyphomycetous, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences are unavailable, need recollecting to verify generic placement.

Chiastospora Riess 1852, *Ascomycota* genera *incertae sedis*, one species, type: *C. parasitica* Riess, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chikaneea B. Sutton 1973, *Ascomycota* genera *incertae sedis*, one species, type: *C. holleriae* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, Canada, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), sequences are unavailable.

Chiodecton Ach. 1814 (= *Melanodecton* A. Massal. 1860 [1859-1860] fide Species Fungorum 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, c. 24 species, type: *C. sphaerale* Ach., lichenized, corticolous or saxicolous, worldwide, see Ertz et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Chionomyces Deighton & Piroz. 1972, *Ascomycota* genera *incertae sedis*, seven species, type: *C. meliolicola* (Cif.) Deighton & Piroz., hyphomycetous, sexual morph unknown, saprobes, worldwide (tropical), see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), sequences and cultures are unavailable.

Chirleja Lendemer & B.P. Hodk. 2012, *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type: *C. buckii* Lendemer & B.P. Hodk., asexual morph unknown, lichenized, terrestrial, South America, see Lendemer and Hodkinson (2012a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Chithramia Nag Raj 1988, *Ascomycota* genera *incertae sedis*, one species, type: *C. elegantissima* Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2016b, 2017; morphology, outline), Kirk et al. (2013; genus accepted), holotype: BPI 71492, sequences and cultures are unavailable.

Chitonomyces Peyr. 1873 (= *Heimatomyces* Peyr. 1873 fide Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 98 species, type: *C. melanurus* Peyr., asexual morph unknown, biotrophic, worldwide, see Lumbsch and Huhndorf (2010; outline), Goldmann and Weir (2012; phylogeny), Kirk et al. (2013; genus accepted), sequences are available for type: (SSU, LSU).

Chitonospora E. Bommer, M. Rousseau & Sacc. 1891, *Phlogi cylindriaceae*, *Amphisphaeriales*, *Xylariomycetidae*, *Sordariomycetes*, one species, type: *C. ammophila* E.

Bommer, M. Rousseau & Sacc., from dead grasses, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015; outline), sequences and cultures are unavailable.

Chlamydomyces Bainier 1907, *Ascomycota* genera *incertae sedis*, two species, type: *C. diffusus* Bainier, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences and cultures are unavailable.

Chlamydopsis Hol.-Jech. & R.F. Castañeda 1986, *Ascomycota* genera *incertae sedis*, one species, type: *C. proliferans* Hol.-Jech. & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: INIFAT C84/140, cultures and sequences are unavailable, compare with *Acrodictys*, *Ashtaangam* and *Petrakia*.

Chlamydorubra K.B. Deshp. & K.S. Deshp. 1966, *Ascomycota* genera *incertae sedis*, one species, type: *C. verrucosa* K.B. Deshp. & K.S. Deshp., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences and cultures are unavailable.

Chlamydosauromyces Sigler, Hambl. & Paré 2002, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *C. punctatus* Sigler, Hambl. & Paré, asexual morph unknown, from bee hive, USA, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), sequences and culture available.

Chlamydotubeufia Boonmee & K.D. Hyde 2011, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, five species, type: *C. huaijangplaensis* Boonmee & K.D. Hyde, saprobes on decaying wood, Thailand, asexual morph hyphomycetous, see Boonmee et al. (2011, 2012; taxonomy), Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), Wijayawardene et al. (2014c; outline), Brahamanage et al. (2017, new species, phylogeny), cultures and sequences are available, holotype and ex-type strains of the type: MFLUCC 10-0926, MFLU 10-0972.

Chlorencoelia J.R. Dixon 1975, *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *C. versiformis* (Pers.) J.R. Dixon, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ren and Zhuang (2014; new species, China), Pärtel et al. (2017; DNA, phylogeny), cultures and sequences are available.

Chloridium Link 1809 (= *Gonytrichum* Nees & T. Nees 1818; = *Melanopsammella* Höhn. 1920 *fide* Réblová et al. 2016c), *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, c. five new species, type: *C. viride* Link, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable for type species.

Chlorociboria Seaver ex C.S. Ramamurthi, Korf & L.R. Batra 1958, *Chlorociboriaceae*, *Helotiales*, *Leotiomycetes*, c. 22 species, saprobes, worldwide, see Huhtinen et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; new species, phylogeny), cultures and sequences are available.

Chloroepilichen Etayo 2010, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. rolfii* Etayo, asexual morph unknown, lichenicolous, terrestrial, Southern America, see Etayo (2010; description and distinction of similar lichenicolous genera), cultures and sequences are unavailable, holotype: UPS, R. Santesson & G. Thor P75:5.

Chloroscypha Seaver 1931 (= *Parksia* E.K. Cash 1945 *fide* Species Fungorum 2017), *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, 14 species, type: *C. seaveri* Rehm ex Seaver, asexual morph sporodochia-like, endophytic, weak pathogen and saprobes, on leaves of *Cupressaceae* (rarely *Taxodiaceae*), widespread in temperate areas, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype: NY herbarium (NY00915009), living cultures and sequences are unavailable for the type of the type species.

Chlorospleniella P. Karst. 1885, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. fennica* (P. Karst.) Sacc. ex Clem. & Shear, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chlorosplenium Fr. 1849 (= *Chrysosplenium* Allesch. 1898 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, five species, type: *C. chlora* (Schwein.) M.A. Curtis, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chlorostroma A.N. Mill., Lar. N. Vassiljeva & J. D. Rogers. 2007, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, three species, type: *C. subcubisporum* A.N. Mill., Lar.N. Vassiljeva & J.D. Rogers, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline),

Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are unavailable.

Chlorovibrissa L.M. Kohn 1989, ?*Vibrisseaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *C. bicolor* (G.W. Beaton & Weste) L.M. Kohn, asexual morph phialophora-like, aquatic or semi-aquatic on wood and twigs, Australasia, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Sandoval-Leiva et al. (2014; DNA), holotype in MELU herbarium, cultures and sequences are available.

Choanatiara DiCosmo 1984, *Ascomycota* genera *incertae sedis*, two species, type: *C. lunata* DiCosmo & Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, Canada, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Choiromyces Vittad. 1831 (= *Choeromyces* Tul. & C. Tul. 1862; = *Piersonia* Harkn. 1899; = *Tartufa* (Gray) Kuntze 1898; = *Tuber* sect. *Tartufa* Gray 1821; = *Zobelia* Opiz 1855 *fide* Species Fungorum 2017), *Tuberaceae*, *Pezizales*, *Pezizomycetes*, type: *C. meandriformis* Vittad., asexual morph unknown, saprobes, hypogeous, terrestrial, worldwide, see Wedén et al. (2009; edible truffle *Choiromyces venosus* in Sweden), Lumbsch and Huhndorf (2010; outline), Moreno et al. (2012b; phylogeny), Bonito et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Chondroderris Maire 1937, *Helotiales*, genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. nevadensis* Maire, asexual morph unknown, saprobes, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chordomyces Bilanenko, M.L. Georgieva & Grum-Grzhimaylo 2016, *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*, one species, type: *C. antarcticus* E.N. Bilanenko, M.L. Georgieva, A.A. Grum-Grzhimaylo, saprobes, from soil, see Grum-Grzhimaylo et al. (2016; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Choreospora Constant. & R. Sant. 1987, *Ascomycota* genera *incertae sedis*, two species, type: *C. tibellii* Constant. & R. Sant., hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, Australia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chorioactis Kupfer ex Eckblad 1968 (= *Chorioactis* Kupfer 1902 nom. inval. Art. 38.5, Ex. 1 (Melbourne), *Chorioactidaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *C. geaster* (Peck) Kupfer ex Eckblad, asexual morph unknown, saprobes, terrestrial, USA, China, see Lumbsch

and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), sequences are available.

Chrismofulvea Marbach 2000, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, four species, type: *C. dialyta* (Nyl.) Marbach, asexual morph unknown, lichenized, epiphytic, North and South America, see Lücking et al. (2017a; classification), sequences are available.

Chromatochlamys Trevis. 1860 (= *Microglaenomyces* Cif. & Tomas. 1953; = *Weitenwebera* Körb. 1863 *fide* Species Fungorum 2017), *Thelenellaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *C. muscicola* (Ach.) Trevis., lichenized, see Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Chromendothia Lar. N. Vassiljeva 1993, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *C. appendiculata* Lar. N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, temperate north-eastern Asia, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2016b; phylogeny, outline), holotype of type: VLA P-2939, sequences are available for *C. citrina*.

Chroodiscus (Müll. Arg.) Müll. Arg. 1890 (= *Ocellularia* sect. *Chroodiscus* Müll. Arg. 1883; = *Phyllobrassia* Vain. 1921 *fide* Species Fungorum 2017), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 14 species, type: *C. coccineus* (Leight.) Müll. Arg., lichenized (*Trenthepohlia* or *Phycopeltis*), terrestrial (foliicolous), tropical, see Papong et al. (2009a, b; DNA), Parnmen et al. (2012; DNA), Sipman et al. (2012; key), Rivas Plata et al. (2013; DNA), Lumbsch et al. (2014a, b; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Chrysachne Cif. 1938, *Ascomycota* genera *incertae sedis*, two species, type: *C. hyphochoria* Cif., hyphomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Chrysocrypta Crous & Summerell 2012, *Erythrogloeaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. corymbiae* Crous & Summerell, coelomycetous, sexual morph unknown, terrestrial, Australia, see Crous et al. (2012e; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), holotype: CBS H-20963, cultures and sequences are available.

Chrysosolia Crous & M.J. Wingf. 2015, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. colombiana* Crous, Rodas & M.J. Wingf., coelomycetous, saprobes, Colombia, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al.

(2017; outline, phylogeny), holotype and ex-type: CBS H-22238, CPC 24986.

Chrysoporthe Gryzenh. & M.J. Wingf. 2004 (= *Chrysoporthella* Gryzenh. & M.J. Wingf. 2004 *fide* Maharachchikumbura et al. 2015b, 2016b), *Cryphomycetaceae*, *Diaporthales*, *Sordariomycetes*, eight species, type: *C. cubensis* (Bruner) Gryzenh. & M.J. Wingf., asexual morph coelomycetous, one species, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Chungu et al. (2010; new species), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: BPI 631857, cultures and sequences are available.

Chrysosporium Corda 1833 (= *Aleurophora* O. Magalh. 1916; = *Atrichophyton* Castell. & Chalm. 1919; = *Blastomyces* Costantin & Rolland 1889 [1888]; = *Emmonsia* Cif. & Montemart. 1959; = *Rhinocladiella* Kamyschko 1960; = *Rhinocladiopsis* Kamyschko 1961 *fide* Species Fungorum 2017), *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, c. 70 species, type: *C. corii* Corda, hyphomycetous, terrestrial, saprobes on keratinous substrates, but also reported from other sources, including dung, soil and foodstuffs, several species are opportunistic pathogens of vertebrates including humans, cosmopolitan, highly polyphyletic, see Liang et al. (2009a; taxonomy), Pitt et al. (2013; reallocation of some xerophilic species, with comments on the polyphyly of the genus), Crous et al. (2016b; new species, phylogeny), cultures and sequences properly representing the type species are unavailable, needs phylogenetic assessment, *Chrysosporium* is tentatively considered herein as a member of *Onygenaceae* following de Hoog et al. (2017).

Chrysotrichix Mont. 1852 (= *Allarthonia* (Nyl.) Zahlbr. 1902; = *Plearthonis* Clem. 1909 *fide* Jaklitsch et al. 2016a), *Chrysotrichaceae*, *Arthoniales*, *Arthoniomycetes*, 14 species, type: *C. noli-tangere* (Mont.) Mont., asexual morph unknown, lichenized, saxicolous or epiphytic, widespread, see Nelsen et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Laundon (2012, conserved against *Alysphaeria*), Sundin et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Chuppia Deighton 1965, *Ascomycota* genera *incertae sedis*, one species, type: *C. sarcinifera* Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), sequences and cultures unavailable.

Ciboria Fuckel 1870 (= *Ciboriopsis* Dennis 1962; = *Moellerodiscus* Henn. 1902 *fide* Species Fungorum 2017), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, c. 21 species, type: *C. caucus* (Rebent.) Fuckel, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf

(2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny), cultures and sequences are available.

Ciborinia Whetzel 1945, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, 16 species, type: *C. bifrons* (Whetzel) Whetzel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Cicadocola Brain 1923, *Ascomycota* genera *incertae sedis*, one species, type: *C. cicadarum* (Šulc) Bain, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), sequences and cultures unavailable.

Cidaris Fr. 1849, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, three species, type: *C. caroliniana* (Schwein.) Fr., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Cidaris Fr. 1849, *Pezizales*, genera *incertae sedis*, *Pezizomycetes*, one species, type: *C. caroliniana* (Schwein.) Fr., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, treated as *Helvellaceae*), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Ciferria Gonz. Frag. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *C. coccothrinacis* Gonz. Frag., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences and cultures unavailable.

Ciferriascosea Senan., Bhat, Camporesi & K.D. Hyde 2015, *Phlogi cylindriaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *C. rectamurum* Senanayake, Bhat, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, Italy, see Senanayake et al. (2015; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ciferrina Petr. 1932, *Ascomycota* genera *incertae sedis*, one species, type: *C. pulchella* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequence data and cultures are unavailable.

Ciferriopeltis Bat. & H. Maia 1965, *Ascomycota* genera *incertae sedis*, one species, type: *C. domingensis* (Petr. & Cif.) Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Ciliella Sacc. & P. Syd. 1902, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. epidendri* (Rehm) Sacc. & P. Syd., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ciliochora Höhn. 1919 (= *Petrakomyces* Subram. & K. Ramakr. 1953 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, two species, type: *C. longiseta* (Racib.) Höhn. coelomycetous, sexual morph unknown, saprobes, terrestrial, Dominican Republic, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ciliochorella Syd. 1935 (= *Phaeociliospora* Bat. & Peres 1967; = *Plagionema* Subram. & K. Ramakr. 1927 *fide Species Fungorum* 2017), *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, four species, type: *C. mangiferae* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Allegrucci et al. (2011; morphology, new species), Wijayawardene et al. (2012, 2016b, 2017; morphology, phylogeny, outline), Kirk et al. (2013; genus accepted), Tangthirasunun et al. (2015; morphology, DNA, phylogeny), Hyde et al. (2016; new species, phylogeny), available specimens and cultures of type: MFLU 13-0264, MFLUCC 12-0310 =ICMP 20052.

Ciliofusospora Bat. & J.L. Bezerra 1963, *Sordariomycetes*, genera *incertae sedis*, one species, type: *C. oenocarpi* Bat. & J.L. Bezerra, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ciliolarina Svrček 1977, *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *C. laricina* (Raitv.) Svrček, asexual morph hyphomycetous (septonema-like), saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Filippova (2012; records from west Siberia), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ciliophora Petr. 1929, *Ascomycota* genera *incertae sedis*, two species, type: *C. cryptica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Central America, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ciliophorella Petr. 1940, *Ascomycota* genera *incertae sedis*, two species, type: *C. insignis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Austria, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Cilioplea Munk 1953, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. coronata* (Niessl) Munk, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Lophiostomataceae*), Hyde et al. (2013) did not accept as a genus in *Lophiostomataceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Ciliosporella Petr. 1927, *Ascomycota* genera *incertae sedis*, two species, type: *C. selenospora* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Austria, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cipisia Marbach 2000, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *C. wheeleri* (R.C. Harris) Marbach, asexual morph unknown, lichenized, epiphytic, North America, see Lücking et al. (2017a; classification), sequences are unavailable.

Circinaria Link 1809, *Megasporaceae*, *Pertusariales*, *Lecanoromycetes*, 25 species, type: *C. contorta* (Hoffm.) A. Nordin, Savić & Tibell, lichenized, see Nordin et al. (2010; phylogeny), Sohrabi et al. (2013a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Circinoconiopsis A. Hern.-Gut. 2013, *Ascomycota* genera *incertae sedis*, one species, type: *C. amazonica* A. Hern. Gut., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Hernández-Gutiérrez (2013; morphology), holotype: MG 203121, sequences and cultures unavailable.

Circinoconis Boedijn 1942, *Ascomycota* genera *incertae sedis*, one species, type: *C. paradoxa* Boedijn, hyphomycetous, sexual morph unknown, saprobes, terrestrial, East Indies, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Circinoniesslia Samuels & M.E. Barr 1997, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. neciae* Samuels & M.E. Barr, asexual morph unknown, saprobes or fungicolous, terrestrial, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: BPI, Samuels 8028, cultures and sequences are unavailable.

Circinotrichum Nees 1816 (= *Dephilippia* Rambelli 1959; = *Gyrocerus* Corda, Icon. fung. (Prague) 1: 9 1837; = *Gyrotrichum* Spreng. 1827 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, c. 20 species, type:

C. maculiforme Nees, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014c; morphology), cultures available for type: CBS 122758, CPC 24566.

Cirrenalia Meyers & R.T. Moore 1960, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, 14 species, type: *C. macrocephala* (Kohlm.) Meyers & R.T. Moore, hyphomycetous, sexual morph unknown, saprobes, aquatic, terrestrial, worldwide, see Abdel-Wahab et al. (2010; phylogeny, key), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available MUCL 15736.

Cirrosporium S. Hughes 1980, *Eurotiomycetes*, genera *incertae sedis*, one species, type: *C. novae-zelandiae* S. Hughes, coelomycetous, sexual morph unknown, saprobes, terrestrial, New Zealand, see Réblová and Seifert (2012; morphology, phylogeny), Wijayawardene et al. (2012, 2016b, 2017; morphology, phylogeny, outline), Kirk et al. (2013; genus accepted), holotype: PDD 36101, cultures available CBS 123236, ICMP 18891.

Cirsosia G. Arnaud 1918 (= *Cirsosiella* G. Arnaud 1918 [1917]; = *Halbanina* G. Arnaud 1918 [1917]; = *Lembopodia* Bat. 1960; = *Morqueria* Bat. & H. Maia 1963 *fide* Species Fungorum 2017), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, 15 species, type: *C. manaosensis* (Henn.) G. Arnaud, asexual morph coelomycetous (=homalopeltis-like *fide* Hofmann 2009), epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Asterinaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; taxonomy, keys), Wijayawardene et al. (2014c; outline), placement in *Asterinaceae* supported by morphology, isotype of type: K(M) 176510, cultures and sequences are unavailable.

Cirsosina Bat. & J.L. Bezerra 1960, *Dothideomycetes* genera *incertae sedis*, two species, type: *C. rhododendri* Bat. & J.L. Bezerra, asexual morph unknown, saprobes, terrestrial, Europe, Borneo, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Cirsosiosis Butin & Speer 1979, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *C. violaceascens* Butin & Speer, asexual morph unknown, epiphytes, terrestrial, Brazil, see Hofmann (2009; review, accepted as a genus in *Microthyriaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; did not accept as a genus in *Microthyriaceae*), Wijayawardene et al. (2014c;

outline, accepted as a genus in *Dothideomycetes, genera incertae sedis*), Wu et al. (2014b; review, transferred to *Parmulariaceae*) placement in *Parmulariaceae* supported by morphology, holotype: ZT Myc 4325, cultures and sequences are unavailable.

Cissococomyces Brain 1923, *Ascomycota* genera *incertae sedis*, one species, type: *C. natalensis* Brain, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Cistella Quél. 1886 (= *Chimaeroscypha* Raity. 2004; *Clavidisculum* Kirschst. 1938 *fide* Jaklitsch et al. 2016a), *Hyaloscypheaceae*, *Helotiales*, *Leotiomycetes*, c. 45 species, type: *C. dentata* (Pers.) Quél., asexual morph unknown, erroneously linked with phialophora-like anamorph (*Cistella rubescens* = *Hyphodiscus hymenophilus*, saprobes on wood, dead herbaceous stems, fallen leaves, needles and cones, widespread, mainly in temperate zones, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; taxonomy), Baral et al. (2015; taxonomy), cultures and sequences are available.

Citeromyces Santa María 1957, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, five species, type: *C. matritensis* (Santa María) Santa María, saprobes, terrestrial, worldwide, see Kurtzman (2012; new species), Casaregola et al. (2013; new species), cultures and sequences are available.

Civisubramaniana Vittal & Dorai 1986, *Ascomycota* genera *incertae sedis*, two species, type: *C. eucalypti* Vittal & Dorai hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: IMI 298300, cultures and sequences are unavailable.

Cladia Nyl. 1870 (= *Clathrina* Müll. Arg. 1883 *fide* Species Fungorum 2017), *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, c. 25 species, type: *C. aggregata* (Sw.) Nyl., asexual morph unknown, lichenized, terrestrial, austral or widespread, see Parnmen et al. (2010; generic delimitation), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Lecanoraceae*), Lumbsch et al. (2010b; generic delimitation), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Cladidium Hafellner 1984 (= *Cladodium* (Tuck.) Gyeln. 1934; = *Lecanora* sect. *Cladodium* Tuck. 1882 *fide* Species Fungorum 2017), *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *C. thamnitis* (Tuck.) Hafellner (= *C. bolanderi* (Tuck.) B.D. Ryan), asexual morph unknown, lichenized, saxicolous, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al.

(2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Cladochasiella Marvanová 1997, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *C. divergens* Marvanová, hyphomycetous, sexual morph unknown, saprobes, aquatic, Czech Republic, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Baschien et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), holotype and ex-type of the type: PRM 842967, CCM F-13489.

Cladoconidium Bandoni & Tubaki 1985, *Ascomycota* genera *incertae sedis*, one species, type: *C. articulatum* Bandoni & Tubaki, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, Canada, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cladographium Peyronel 1918, *Ascomycota* genera *incertae sedis*, one species, type: *C. rivulorum* Peyronel, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Cladonia P. Browne 1756 (see Species Fungorum 2017 for synonyms), *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, c. 550 species, type: *C. subulata* (L.) F.H. Wigg., asexual morph unknown, lichenized, terricolous or on lignum, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Lecanoraceae*), Pino-Bodas et al. (2012; polymorphic species), Ahti and Stenroos (2013; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Cladoniicola Diederich, van den Boom & Aptroot 2001, *Pezizomycotina*, genera *incertae sedis*, one species, type: *C. staurospora* Diederich, van den Boom & Aptroot, asexual morph unknown, lichenicolous, terrestrial, temperate, see Etayo (2010b; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cladophialophora Borelli 1980, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, 35 species, type: *C. carriionii* (Trejos) de Hoog, Kwon-Chung & McGinnis, hyphomycetous, mostly terrestrial, but also reported from mangrove environments, mainly saprobes associated with soil and plant material but a few species are lichenicolous, endophytic or plant-pathogenic, several species cause infections in animals, including humans, cosmopolitan, although the type species belongs in *Herpotrichiellaceae*, some species are related to other families of *Chaetothyriales*, see Crous et al. (2011c, 2013a; new species, pathogenicity to plants), Badali et al. (2008, 2009, 2011; outline, new species), Diederich et al.

(2013; lichenicolous species), Feng et al. (2013; new species), Seyedmousavi et al. (2013; pathogenicity to animals), Gueidan et al. (2014a; phylogeny), Madrid et al. (2016; new species, comments about the polyphyly of the genus), Usui et al. (2016; new species), ex-type strain of the type: =CBS 160.54, sequences are available.

Cladoriella Crous 2006, *Dothideomycetes* genera *incertae sedis*, three species, type: *C. eucalypti* Crous, hyphomycetous, sexual morph unknown, saprobes or ?pathogens, terrestrial, Australia, South Africa, see Cheewangkoon et al. (2009; morphology, new species, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), holotype and ex-type: CBS H-18043, CPC 10953–10955 =CBS 115898–115890.

Cladorrhinum Sacc. & Marchal 1885 (= *Bahupaathra* Subram. & Lodha 1964 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, eight species, type: *C. foecundissimum* Sacc. & Marchal, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Madrid et al. (2011a; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Carmaran et al. (2015; phylogeny), Maharachchikumbura et al. (2015b, 2016b; phylogeny, outline), available cultures of type: BCCM 6980.

Cladosphaera Dumort. 1822, *Ascomycota* genera *incertae sedis*, one species, type: *C. cespitosa* (Tode) Dumort., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cladosporiella Deighton 1965, *Ascomycota* genera *incertae sedis*, five species, type: *C. cercosporicola* Deighton, hyphomycetous, sexual morph unknown, saprobes, fungicolous, terrestrial, Africa, Asia, Caribbean, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cladosporiopsis S.C. Ren & X.G. Zhang 2012, *Ascomycota* genera *incertae sedis*, one species, type: *C. ovata* S.C. Ren & X.G. Zhang, hyphomycetous, sexual morph unknown, saprobes, fungicolous, terrestrial, China, see Ren et al. (2012a; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Cladosporium Link 1816 (= *Acrosperella* Riedl & Ershad 1977 [1976]; = *Azosma* Corda 1831; = *Davidiella* Crous & U. Braun 2003; = *Didymotrichum* Bonord. 1851; = *Heterosporium* Klotsch ex Cooke 1877; = *Hormodendrum* Bonord. 1851; = *Mydonosporium* Corda 1833; = *Myxocladium* Corda 1837; = *Polyrhizium* Giard 1889; = *Spadicesporium* V.N. Boriss. & Dvoříns 1982; =

Sporocladium Chevall. 1826 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b; Species Fungorum 2017), *Cladosporiaceae*, *Capnodiales*, *Dothideomycetes*, c. 300 species, type: *C. herbarum* (Pers.) Link, hyphomycetous, sexual morph formerly known in *Davidiella*, saprobes, endophytes, animal, human and plant pathogens, worldwide, see Seifert et al. (2011; morphology), Bensch et al. (2012, 2015; phylogeny, taxonomy), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Cladosporiaceae*), Sandoval-Denis et al. (2015b; clinical), lectotype of type: L 910.225-733, epitype: CBS-H 19853; ex epitype culture, CBS 121621.

Clandestinotrema Rivas Plata, Lücking & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 17 species, type: *C. clandestinum* (Ach.) Rivas Plata, Lücking & Lumbsch, lichenized, see Rivas Plata et al. (2012a; taxonomy), Lumbsch et al. (2014a, b; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Clasteropycnis Bat. & Cavalc. 1963, *Ascomycota* genera *incertae sedis*, one species, type: *C. coumae* Bat. & Cavalc., sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clasterosphaeria Sivan. 1984, *Magnaportheaceae*, *Magnaporthales*, *Sordariomycetes*, two species, type: *C. cyperi* Sivan., asexual morph clasterosporium-like, saprobes, terrestrial, India, Malaysia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: IMI 56878, cultures and sequences are unavailable.

Clasterosporium Schwein. 1832 (= *Apotemnoum* Corda 1833; = *Brachydesmium* (Sacc.) Costantin 1888; = *Cheiropodium* Syd. & P. Syd. 1915; = *Clasterosporium* sect. *Brachydesmium* Sacc. 1886; = *Cometella* Schwein. ex Fr. 1849; = *Hymenopodium* Corda 1837; = *Phanerocoryneum* Höhn. 1923; = *Sporhelminthium* Speg. 1918 *fide* Species Fungorum 2017), *Magnaportheaceae*, *Magnaporthales*, *Sordariomycetes*, ten species, type: *C. caricum* Schwein., sexual morph ?clasterosphaeria-like, saprobes, terrestrial, worldwide, see Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, sexual-aseexual link was not confirmed.

Clathroconium Samson & H.C. Evans 1982, *Ascomycota* genera *incertae sedis*, one species, type: *C. arachnicola* Samson & H.C. Evans, hyphomycetous, sexual morph unknown, entomopathogens, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Clathroporina Müll. Arg. 1882, *Porinaceae*, *Ostropales*, *Lecanoromycetes*, 25 species, type: *C. endochrysea* (C. Bab.) Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Clathrosphaerina Beverw. 1951, *Hyaloscypheaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *C. zalewskii* Beverw., sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clathrospora Rabenh. 1857, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, more than 40 species, type: *C. elynae* Rabenh., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2015c; morphology, accepted as *Pleosporaceae*), sequences are available, type sequence data are available: CBS 161.51; CBS 196.54.

Clathrosporium Nawawi & Kuthub. 1987, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. intricatum* Nawawi & Kuthub., sexual morph unknown, saprobes, aero-aquatic, Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Clauroxia D. Hawksw. 1988 (= *Pseudolecidea* Clauzade & Cl. Roux 1984 [1983] *fide* Species Fungorum 2017), *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *C. chalybeoides* (Nyl.) D. Hawksw., asexual morph unknown, lichenized, saxicolous, Europe, North America, see Gilbert and Hawksworth (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Claussenomyces Kirschst. 1923 (= *Corynella* Boud. 1885; = *Dendrostilbella* Höhn. 1905 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Tympanidaceae*, *Phacidiales*, *Leotiomycetes*, 19 species, type: *C. jahnianus* Kirschst., asexual morph formerly known in *Dendrostilbella*, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2012; outline and asexual morph), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), sequence data are available for unidentified *Claussenomyces* species.

Clauzadea Hafellner & Bellem. 1984, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, five species, type: *C. monticola* (Ach.) Hafellner & Bellem., asexual morph unknown, lichenized, saxicolous, widespread, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Clauzadeana Cl. Roux 1984, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *C. instratula* (Nyl.) Cl. Roux, asexual morph unknown, lichenized, saxicolous, Asia, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Clauzadella Nav.-Ros. & Cl. Roux 1996, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *C. gordensis* Nav.-Ros. & Cl. Roux, asexual morph unknown, terrestrial, lichenized, on *Verrucaria viridula* (Schrad.) Ach., France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Clauzadeomyces Diederich 1994, *Ascomycota* genera *incertae sedis*, one species, type: *C. verrucosus* Diederich, hyphomycetous, sexual morph unknown, lichenicolous, non-lichenized, terrestrial, Belgium (Ardennes), see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: LG, Lambinon, 22 Aug. 1965, cultures and sequences are unavailable.

Clavariana Nawawi 1976, *Ascomycota* genera *incertae sedis*, one species, type: *C. aquatica* Nawawi, hyphomycetous, sexual morph unknown, in river foam, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Clavascidium Breuss 1996, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, eight species, type species: *C. umbrinum* (Breuss) Breuss, asexual morph coelomycetous, terrestrial, lichenized, occurring on soil, rocks and mosses, America, Asia, Europe, see Gueidan et al. (2009; phylogeny, proposal to synonymize *Clavascidium* with *Placidium* A. Massal.), Prieto et al. (2010c, 2012; morphology, reinstatement of *Clavascidium* as a distinct genus), sequences of the type material of the type species are unavailable, but sequences of non-type material are available.

Clavatispora Boonmee & K.D. Hyde 2014, *Venturiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *C. thailandica* Boonmee & K.D. Hyde, asexual morph hyphomycetous, saprobes, terrestrial, Thailand, see Boonmee et al. (2014a; taxonomy), sequence data available, holotype and ex-type: MFLU 10-0038, MFLUCC 10-0107 = BCC 38863 = IFRD 2170.

Clavatisporella K.D. Hyde 1995, *Magnaportheaceae*, *Magnaporthales*, *Sordariomycetes*, one species, type: *C. musicola* K.D. Hyde, asexual morph unknown, saprobes, Irian Jaya (Papua), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted),

Maharachchikumbura et al. (2016b; outline), holotype: BRIP 22403, cultures and sequences are unavailable.

Clavatospora Sv. Nilsson ex Marvanová & Sv Nilsson 1971 (= *Clavatospora* Sv. Nilsson 1933 *fide* Species Fungorum 2017), *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, three species, type: *C. longibrachiata* (Ingold) Sv. Nilsson ex Marvanová & Sv. Nilsson, hyphomycetous, sexual morph corollospora-like, saprobes, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016a; outline), needs revision since genus shown to be polyphyletic, reference cultures of type: UMB-1109.13.

Claviceps Tul. 1853 (= *Balansiella* Henn. 1904; = *Kentrosporium* Wallr. 1844; = *Mothesia* Oddo & Tonolo 1967; = *Spermoedia* Fr. 1822; = *Sphacelia* Lév. 1827; = *Ustilagopsis* Speg. 1880 *fide* Rossman et al. 2016a), *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, 56 species, type: *C. purpurea* (Fr.) Tul., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Pažoutová et al. (2008, 2011, 2015; new species), Kirk et al. (2013; genus accepted), Kolařík (2015a, 2015b; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are available.

Claviradulomyces P.R. Johnst., D.C. Park, H.C. Evans, R.W. Barreto & D.J. Soares 2010, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *C. dabeicola* P.R. Johnst., D.C. Park, H.C. Evans, R.W. Barreto & D.J. Soares, asexual morph (putative) pycnidial on branches of vascular plants, Africa, South America, see Evans et al. (2010; taxonomy), Barreto et al. (2012; phylogeny), cultures and sequences are available.

Clavisporea Rodrigues de Miranda 1979, *Metschnikowiaeae*, *Saccharomycetales*, *Saccharomycetes*, 20 species, type: *C. lusitaniae* Rodrigues de Miranda, asexual reproduction by budding, saprophytic, from cacti, agave, processed dates and other high osmotic environments, occasionally an opportunistic human pathogen, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Cleistocystis Sousa da Câmara 1931, *Ascomycota* genera *incertae sedis*, one species, type: *C. rosarum* Sousa da Câmara, coelomycetous, sexual morph unknown, saprobes, terrestrial, Portugal, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cleistoiodophanus J.L. Bezerra & Kimbr. 1976, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type:

C. conglutinatus J.L. Bezerra & Kimbr., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Cleistonium Speer 1986, *Ascomycota genera incertae sedis*, one species, type: *C. parasiticum* Speer, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2014c) accepted as a genus in *Gloniaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline, accepted as *Ascomycota genera incertae sedis*), cultures and sequences are unavailable.

Cleistophoma Petr. & Syd. 1927, *Ascomycota genera incertae sedis*, two species, type: *C. suberis* (Prill. & Delacr.) Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cleistosphaera Syd. & P. Syd. 1916, *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *C. macrostegia* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, S. America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as in *Perisporiopsidaceae*, key), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Cleistothelebolus Malloch & Cain 1971, *Thelebolaceae*, *Thelebolales*, *Pezizomycetes*, one species, type: *C. nippigenensis* Malloch & Cain, asexual morph unknown, coprophilous, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Clematidis Tibpromma, Camporesi & K.D. Hyde 2016, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *C. italicica* Tibpromma, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Li et al. (2016a; taxonomy, phylogeny), holotype and ex-type: MFLU 14-0669, MFLUCC 15-0084.

Clematomyces Thaxt. 1900 (= *Peziza* sect. *Clibanites* P. Karst. 1869 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, five species, type: *C. pinophili* Thaxt., asexual morph unknown, biotrophic, Asia, West Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clibanites (P. Karst.) P. Karst. 1871, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. paradoxa* (P. Karst.) P. Karst., asexual morph unknown, saprobes on wood, Finland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b;

outline), cultures and sequences are unavailable, holotype in H.

Cliostomum Fr. 1825 (= *Catillariomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Rhytismella* P. Karst. 1884; = *Sporoblastia* Trevis. 1851 *fide* Species Fungorum 2017), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, c. 25 species, type: *C. corrugatum* (Ach.) Fr., asexual morph unknown, lichenized, epiphytic, world-wide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Printzen (2014; phylogeny and relationships), Lücking et al. (2017a; classification), sequences are available.

Clohesomyces K.D. Hyde 1993, *Lindgomycetaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *C. aquaticus* K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, aquatic, Australia, see Zhang et al. (2012a; new species, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), holotype: BRIP 20092, cultures and sequences are available.

Clohesisia K.D. Hyde 1995, *Annulatascaeae*, *Annulatascales*, *Sordariomycetes*, three species, type: *C. corticola* K.D. Hyde, sexual morph unknown, saprobes, terrestrial, China, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: BRIP 21485, cultures available for *Clohesisia corticola* HKUCC 3712.

Clonophoromyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *C. grenadinus* Thaxt., asexual morph unknown, biotrophic, West Indies, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2015c; notes), cultures and sequences are unavailable.

Clonostachys Corda 1839 (= *Bionectria* Speg. 1919 *fide* Rossman et al. 2013b), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, c. 50 species, type: *C. araucaria* Corda, hyphomycetous, sexual morph formerly known in *Bionectria*, on soil, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; phylogeny, new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Clypeispora A.W. Ramaley 1991, *Ascomycota genera incertae sedis*, one species, type: *C. angustifoliorum* A.W. Ramaley, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), holotype: BPI 1102631, cultures and sequences are unavailable.

Clypeoceriospora Sousa da Câmara 1946, *Sordariomycetes* genera *incertae sedis*, one species, type: *C. rubi* Sousa da Câmara, asexual morph unknown,

saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Clypeochorella Petr. 1923, *Ascomycota genera incertae sedis*, one species, type: *C. orientalis* (Sacc. & Penz.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clypeococcum D. Hawksw. 1977, *Polycoccaceae*, *Trypetheliales*, *Dothideomycetes*, eleven species, type: *C. cladonema* (Wedd.) D. Hawksw., on lichens, widespread, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Dacampiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Ertz et al. (2015a; sequences two species, but not the type species, placed in *Polycoccaceae*), Jaklitsch et al. (2016a; accepted as a genus in *Dacampiaceae*), holotype of type: PC.

Clypeolculus Kaz. Tanaka & K. Hiray. 2015, *Morosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *C. akitaensis* Kaz. Tanaka & K. Hiray., asexual morph unknown, saprobes, aquatic, Japan, see Tanaka et al. (2015; morphology, phylogeny), holotype and ex-type: HHUF 27557, CBS 139681.

Clypeolum Speg. 1881, *Ascomycota genera incertae sedis*, eight species, type: *C. atroareolatum* Speg., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clypeopatella Petr. 1942, *Ascomycota genera incertae sedis*, one species, type: *C. cylindrospora* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Rhodes, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clypeophialophora Bat. & Peres 1962, *Ascomycota genera incertae sedis*, one species, type: *C. erythroxyli* Bat. & Peres, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clypeophysalospora H.J. Swart 1981, *Clypeophysalosporaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *C. latitans* (Sacc.) H.J. Swart, asexual morph unknown, cosmopolitan, see Giraldo et al. (2017; phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Clypeoporthea Höhn. 1919, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *C. monocarpa* Höhn., asexual morphs unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Clypeoporthella Petr. 1924 (= *Batschiella* Kirschst. 1938 *fide* Species Fungorum 2017), *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *C. brencklei* Petr., asexual morphs unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Clypeopycnis Petr. 1925, *Ascomycota genera incertae sedis*, three species, type: *C. aeruginascens* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clypeopyrenis Aptroot 1991, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, two species, type: *C. microsperma* (Müll. Arg.) Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Clypeoseptoria F. Stevens & P.A. Young 1925, *Ascomycota genera incertae sedis*, three species, type: *C. rockii* F. Stevens & P.A. Young, coelomycetous, sexual morph unknown, saprobes, terrestrial, Hawaii, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clypeosphaerella Guatimosim, R.W. Barreto & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *C. quasiparkii* (Cheew., K.D. Hyde & Crous) Guatimosim, R.W. Barreto & Crous, asexual morph unknown, plant pathogenic, see Guatimosim et al. (2016a; morphology, phylogeny), holotype CBS H-22088, isotype VIC 42607, culture ex-type CPC 24705.

Clypeosphaeria Fuckel 1870 (= *Anthostoma* subgen. *Entosordaria* (Sacc.) Cooke 1889; = *Anthostomella* subgen. *Entosordaria* Sacc. 1882; = *Coccochorina* Hara 1927; = *Entosordaria* (Sacc.) Höhn. 1920; = *Stereosphaeria* Kirschst. 1939 *fide* Species Fungorum 2017), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, c. 30 species, type: *C. notarisii* Fuckel, asexual morph unknown, saprobes (dead wood and bark), terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al.

(2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, treated as *Clypeosphaeriaceae*), Hongsanan et al. (2017a; phylogeny, evolution, treated *Clypeosphaeriaceae* as a synonym of *Xylariaceae*), cultures and sequences are unavailable for type species.

Clypeosphaerulina Sousa da Câmara 1939, *Sordariomycetes* genera *incertae sedis*, one species, type: *C. vincae* (Sousa da Câmara) Sousa da Câmara, asexual morph unknown, saprobes, terrestrial, Portugal, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Clypeostagonospora Punith. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *C. muhlenbergiae* Punith., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Clypeostroma Theiss. & Syd. 1914, *Dothideomycetes* genera *incertae sedis*, two species, type: *C. spilomeum* (Berk.) Theiss. & Syd., asexual morph unknown, saprobes, Australasia, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Coccidiascus Chatton emend. Lushbaugh, Rowton & McGhee 1976, *Saccharomycetales* genera *incertae sedis*, *Saccharomycetes*, one species, type: *C. legeri* Chatton, asexual reproduction by budding, perhaps parasitic on *Drosophila* spp., found within the intestinal epithelial cells of *Drosophila*, possibly worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coccidioides G.W. Stiles 1896 (= *Blastosporidium* M. Hartmann 1912; = *Megalocitospoides* Wernicke 1892; = *Posadasia* Cantón 1898; = *Pseudococcidioides* Fonseca 1928 *fide* Species Fungorum 2017), *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *C. immitis* G.W. Stiles, sexual morph unknown, saprobes, terrestrial, southern USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Coccidomyces Buchner 1912, *Saccharomycetidae* genera *incertae sedis*, *Saccharomycetes*, one species, type: *C. pierantonii* Buchner, asexual morphs unknown, in insects, terrestrial, Germany, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coccinonectria L. Lombard & Crous 2015, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *C. pachysandricola* (B.O. Dodge) L. Lombard & Crous, asexual morph hyphomycetous, saprobes, France, USA, see Lombard et al. (2015b; morphology, phylogeny),

Maharachchikumbura et al. (2015b, 2016b; outline), available cultures of the type: CBS 501.63.

Coccisia Norman 1870, *Dothideomycetes* genera *incertae sedis*, two species, type: *C. hamperi* Norman, asexual morph unknown, saprobes, Scandinavia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dacampiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Coccocarpia Pers. 1827 (= *Circinaria* Fée 1825 [1824]; = *Vischia* C.W. Dodge 1971 [1970] *fide* Species Fungorum 2017), *Coccocarpiaceae*, *Peltigerales*, *Lecanoromycetes*, c. 25 species, type: *C. molybdaea* Pers., asexual morph unknown, lichenized, epiphytic or epiphyllous, worldwide tropical-subtropical, see Wedin et al. (2009b; systematics), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Coccochora Höhn. 1909 (= *Coccochorella* Höhn. 1910 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *C. quercicola* (Henn.) Höhn., asexual morph unknown, saprobes, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Coccochorina Hara 1927, *Dothideomycetes* genera *incertae sedis*, two species, type: need typification, asexual morph unknown, saprobes, terrestrial, see Kirk et al. (2008; treated as a synonym of *Clypeosphaeria*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Coccidiella Hara 1910 (see Species Fungorum 2017 for synonyms), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, 21 species, type: *C. arundinariae* Hara, asexual morph unknown, biotrophic, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures available for *C. melastomatum* CMU78543, *C. miconiae* ppMP 1342.

Coccodinium A. Massal. 1860, *Coccodiniaceae*, *Chaetothyriales*, *Eurotiomycetes*, two species, type: *C. bartschii* A. Massal., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Coccodiniaceae*; excluded family from *Dothideomycetes*), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Coccodothis Theiss. & Syd. 1914, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. sphaeroidea* (Cooke) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, Georgia, see Hofmann (2009;

review, accepted as a genus in *Parmulariaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Parmulariaceae*), Hyde et al. (2013; excluded from *Parmulariaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Coccogloeum Petr. 1955, *Ascomycota* genera *incertae sedis*, one species, type: *C. microsporum* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coccoidea Henn. 1900 (= *Apiodothina* Petr. & Cif. 1932; = *Coccodiscus* Henn. 1904 *fide* Species Fungorum 2017), *Coccoideaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *C. quercicola* Henn. & Shirai, asexual morph unknown, saprobes, Japan, India, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Coccoideaceae*, morphology), cultures and sequences are unavailable.

Coccoidella Höhn. 1909 (= *Coccodothella* Theiss. & Syd. 1915; = *Eumycrocyclus* Hara 1915 *fide* Species Fungorum 2017), *Coccoideaceae*, *Dothideomycetes* families *incertae sedis*, seven species, type: *C. scutula* (Berk. & M.A. Curtis) Höhn., asexual morph unknown, saprobes, America, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Coccoideaceae*), cultures and sequences are unavailable.

Cocomyces De Not. 1847 (= *Cocomycella* Höhn. 1917; = *Malenconia* Bat. & H. Maia 1960 *fide* Species Fungorum 2017), *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, 118 species, type: *C. coronatus* (Schumach.) De Not., asexual morph unknown, pathogens, worldwide, see Marincowitz et al. (2008; new species, morphology), Jia et al. (2011; new species, morphology), Li et al. (2014d, 2015b; new species, morphology), Moreno-Rico et al. (2014; new species, morphology), Wang et al. (2014c; new species, morphology), Li et al. (2015b; new species), sequences are available for *C. dentatus* and *C. strob.*

Cocomycetella Höhn 1917, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *C. belonospora* (Nyl.) Höhn, asexual morph unknown, epixylic, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Baloch et al. (2013; systematics), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), sequences are available.

Cocconia Sacc. 1889, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, c 25 species, type: *C. placenta* (Berk.

& Broome) Sacc., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), Jayasiri et al. (2016a; re-visited), placement in *Parmulariaceae* supported by morphology, unknown, cultures are sequences unavailable.

Coccostromella Petr. 1968, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, one species, type: *C. puttemansi* (Henn.) Petr., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline, accepted as *Dothideaceae*), Hyde et al. (2013; accepted as a genus in *Dothideaceae*), Thambugala et al. (2014a; excluded from *Dothideaceae* and placed *Botryosphaeriaceae*), Wijayawardene et al. (2014c; outline, taxonomy), cultures and sequences are unavailable.

Coccotrema Müll. Arg. 1888 (= *Perforaria* Müll. Arg. 1891; = *Perforariomyces* Cif. & Tomas. 1953; = *Verrucaria* sect. *Verrucaria* Hue 1914; = *Verrucarina* (Hue) Zahlbr. 1931 [1932] *fide* Species Fungorum 2017), *Coccotremataceae*, *Pertusariales*, *Lecanoromycetes*, 16 species, type: *C. antarcticum* Müll. Arg., asexual morph unknown, lichenized, terricolous, saxicolous or epiphytic, world-wide in arctic and sub Antarctic areas, see Lumbsch and Huhndorf (2010), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available

Cochliomyces Speg. 1912, *Euceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *C. argentinensis* Speg., asexual morph unknown, biotrophic, Central and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cocoicola K.D. Hyde 1995, *Phaeochoraceae*, *Phyllachorales*, *Sordariomycetes*, five species, type: *C. cylindrospora* (C. Booth & D.E. Shaw) K.D. Hyde, saprobes, worldwide, see Lumbsch and Huhndorf (2010), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Codinaea Maire 1937 (= *Menisporella* Agnihothr. 1962 *fide* Species Fungorum 2017), *Chaetosphaeriaceae*, *Chaetosphaerales*, *Sordariomycetes*, c. 80 species, type: *C. aristata* Maire, hyphomycetous, sexual morphs unknown, saprobes, terrestrial, tropical, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Almeida et al. (2014c; new species, Brazil), Crous et al. (2014d, 2015d; new species), Granados et al. (2014; new species, Costa Rica), Maharachchikumbura et al. (2015b, 2016b; outline), Xia et al. (2015a; new species, China), Barbosa et al. (2016; new species, Brazil), Oliveira et al. (2016b; new

species, Brazil, aquatic), cultures and sequences are available.

Codonmyces Calatayud & Etayo 1999, Ascomycota genera *incertae sedis*, one species, type: *C. lecanorae* Calat. & Etayo, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, Spain, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: VAB (lich.) 15404, cultures and sequences are unavailable.

Coelodictyosporium Thambug. & K.D. Hyde 2015, Lophiostomataceae, Pleosporales, Dothideomycetes, two species, type: *C. pseudodictyosporium* (Qing Tian, Camporesi & K.D. Hyde) Thambug. & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Italy, all species has sequence data, see Thambugala et al. (2015b; phylogeny), Wijayawardene et al. (2016b; morphology, phylogeny), holotype: MFLU 14-0737, ex-type strains of type: MFLUCC 13-0451.

Coelopogon Brusse & Kärnefelt 1991, Parmeliaceae, Lecanorales, Lecanoromycetes, two species, type: *C. abraxas* Brusse, lichenized, south temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Coenogonium Ehrenb. ex Nees 1820 (see Species Fungorum 2017 for synonyms), Coenogoniaceae, Ostropales, Lecanoromycetes, 90 species, type: *C. linkii* Ehrenb., lichenized, see Baloch et al. (2010; taxonomy), Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; DNA), Kirk et al. (2013; genus accepted), Joshi et al. (2015b; new species, key), Lücking et al. (2017a; classification), sequences are available.

Colemaniella Agnihothr. 1974, Ascomycota genera *incertae sedis*, one species, type: *C. ossoori* Agnihothr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Colensonella Hafellner 1979, Patellariaceae, Patellariales, Dothideomycetes, one species, type: *C. torulispora* (W. Phillips) Hafellner, asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Tian et al. (2014; morphology; nomenclature), cultures and sequences are unavailable.

Coleodictyospora Charles 1929 (= Coleodictys Clem. & Shear 1931 *fide* Species Fungorum 2017), Ascomycota genera *incertae sedis*, one species, type: *C. cubensis* Charles, hyphomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coleophoma Höhn. 1907 (= *Bactropycnis* Höhn. 1920; = *Coleonaema* Höhn. 1924; = *Rhabdostromellina* Höhn. 1917; = *Xenodomus* Petr. 1922 *fide* Species Fungorum 2017), Dermateaceae, Helotiales, Leotiomycetes, more than 35 species, type: *C. crateriformis* (Durieu & Mont.) Höhn., coelomycetous, sexual morph unknown, pathogenic, saprobes or endophytic on a wide host range, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2016a; new species), holotype of type: FH 00304449.

Coleoseptoria Petr. 1940, Ascomycota genera *incertae sedis*, one species, type: *C. ephedrae* (Auersw.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Germany, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coleosperma Ingold 1954, Helotiales genera *incertae sedis*, Leotiomycetes, one species, type: *C. lacustre* Ingold, asexual morph unknown, saprobes, aquatic, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coleroa Rabenh. 1850 (= *Aphysa* Theiss. & Syd. 1917; = *Cyphospilea* Syd. 1926; = *Hormotheca* Bonord. 1864 *fide* Species Fungorum 2017), Venturiaceae, Venturiales, Dothideomycetes, eight species, type: *C. chaetomium* (Kunze) Rabenh., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011d; morphology, phylogeny), Hyde et al. (2013; accepted as in Venturiaceae, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequence data available for *Coleroa robertiani* CBS 458.64.

Colipila Baral & Guy García 2012, Helotiales genera *incertae sedis*, Leotiomycetes, two species, type: *C. masduguana* Baral & Guy García, asexual morph unknown, saprobes, cosmopolitan, see Baral et al. (2012; taxonomy, phylogeny), holotype: M-014088, cultures and sequences are available.

Colispora Marvanová 1988, Sarcosomataceae, Pezizales, Pezizomycetes, three species, type: *C. elongata* Marvanová, hyphomycetous, sexual morph unknown, saprobes, aquatic, worldwide, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences and cultures available for *Colispora elongata* (F-08382 *fide* Baschien et al. 2013).

Collarina Giraldo, Gené & Guarro 2012, Clavicipitaceae, Hypocreales, Sordariomycetes, two species, type: *C. aurantiaca* Giraldo, Gené & Guarro, hyphomycetous, sexual morph unknown, from water

sediment, aquatic, Spain, see Crous et al. (2014d; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Collema F.H. Wigg. [nom. cons.] 1780 (see Species Fungorum 2017 for synonyms), *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, 40 species, type: *C. lactuca* (Weber) Weber ex F.H. Wigg., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Collematospora Jeng & Cain 1976, *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, one species, type: *C. venezuelensis* Jeng & Cain, asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Collembolispora Marvanová & Pascoal 2003, *Helotiales* genera *incertae sedis*, *Leotiomyces*, two species, type: *C. barbata* Marvanová, Pascoal & Cássio, hyphomycetous, sexual morph unknown, saprobes, aquatic, Portugal, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2012c; new species, DNA, morphology), holotype and ex-type: IMI 389661, UMB-088.01.

Collemopsidium Nyl. 1881 (= *Collemopsidiomyces* Cif. & Tomas. 1953; = *Xanthoporina* C.W. Dodge 1948; = *Xanthopyrenia* Bachm. 1919 *fide* Species Fungorum 2017), *Xanthopyreniaceae*, *Collemopsidiales*, *Dothideomycetes*, c. 40 species (but the number could be higher), type: *C. iocarpum* (Nyl.) Nyl., molecular data suggest that the number of marine species has been largely underestimated (Pérez-Ortega et al. 2016), asexual morph coelomycetous, lichen-forming fungi, associated with cyanobacteria, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a, classification), Pérez-Ortega et al. (2016, phylogeny), lectotype of type: H-NYL 42858.

Colletococonis de Hoog & Aa 1978, *Ascomycota* genera *incertae sedis*, one species, type: *C. aecidiophila* (Speg.) de Hoog, Aa & U.P. Singh, coelomycetous, sexual morph unknown, saprobes, fungicolous, Argentina, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Colletogloeum Petr. 1953, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *C. sissoo* (Syd.) B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Crous et al. (2009f; taxonomy, phylogeny), Wijayawardene et al. (2012, 2016b, 2017; outline), Zhao and Zhao (2012; new

species, China), cultures and sequences are available for unidentified species.

Colletotrichum Corda. 1831 (= *Glomerella* Spauld. & H. Schrenk 1903 *fide* Réblová et al. 2016c; see Species Fungorum 2017 for additional synonyms), *Glomerellaceae*, *Glomerellales*, *Sordariomycetes*, more than 800 species, type: *C. lineola* Corda, coelomycetous, sexual morph formerly known in *Glomerella*, pathogens, endophytes, saprobes, terrestrial and aquatic, worldwide, many species have molecular data including, the type species, sequence data available for accepted species, see Hyde et al. (2009, 2014; phylogeny, species number), Damm et al. (2010a, 2012a, b, 2013, 2014; species complexes, phylogeny), Cannon et al. (2012; phylogeny), O'Connell et al. (2012; pathogenic species), Peng et al. (2012; species in Citrus), Weir et al. (2012; phylogeny, species complex), Lima et al. (2013; pathogens), Manamgoda et al. (2013; endophytic species), Udayanga et al. (2013; pathogens), Vieira et al. (2014; endophytic species), Liu et al. (2015a; new species), Sharma et al. (2015b; *C. siamense* complex), Yan et al. (2015; species associated with grape wine diseases), Hou et al. (2016; new species), Huang et al. (2016b; genomics), Jayawardena et al. (2016b, c, d, e; new species, biocontrol, current status, secondary metabolites, on strawberry), Sharma and Shenoy (2016; methods), Marin-Felix et al. (2017; accepted species, phylogeny, new species), Tibpromma et al. (2017b; new species), epitype and culture ex-epitype of type species: CBS H-20362, CBS 1253376.

Collodiscula I. Hino & Katum. 1955 (= *Acanthodochium* Samuels et al. 1987 *fide* Réblová et al. 2016c), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *C. japonica* I. Hino & Katum., asexual morph hyphomycetous (*Acanthodochium*), saprobes, Japan, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch and Voglmayr (2012; phylogeny), Wijayawardene et al. (2012, 2017; outline), Stadler et al. (2013; one name), Li et al. (2015d; new species), Maharachchikumbura et al. (2015b, 2016b; outline, one name), Réblová et al. (2016c; nomenclature), cultures and sequences are available.

Collophorina Damm & Crous, *nom. nov.* MycoBank MB821422

≡ *Collophora* Damm & Crous, Persoonia 24: 64. 2010. nom. illeg., [ICN (Melbourne) Art. 53.1], non *Collophora* Mart. 1830 (*Apocynaceae*).

Leotiomyces genera *incertae sedis*, five species, coelomycetous, sexual morph unknown, endophytes, saprobes, plant pathogens, Baleares, Pacific Northwest regions (USA), South Africa, Germany, Iran, see Damm et al. (2010a; morphology, phylogeny), Gramaje et al. (2012; new species, phylogeny), Wijayawardene et al. (2012; outline), Xie et al. (2013; new species).

See Damm et al. (2010a) for illustrations, description and diagnoses.

Holotype: South Africa, Western Cape Province, Paarl, from wood of *Prunus persica*, 10 Jun. 2004, U. Damm, holotype CBS H-19992, ex-type strain of the type species *C. rubra*: CBS 120873 = STE-U 6109.

Collophorina aceris (J. Xie, Strobel & Geary) Damm & Crous, comb. nov. MycoBank MB 821434

≡ *Collophora aceris* J. Xie, Strobel & Geary, Microbial Ecology 66: 787. 2013.

Collophorina africana (Damm & Crous) Damm & Crous, comb. nov. MycoBank MB821423

≡ *Collophora africana* Damm & Crous, Persoonia 24: 65. 2010.

≡ *Collophora capensis* Damm & Crous, Persoonia 24: 67. 2010.

Collophorina hispanica (Gramaje, Armengol & Damm) Damm & Crous, comb. nov. MycoBank MB 821430

≡ *Collophora hispanica* Gramaje, Armengol & Damm, Persoonia 28: 9. 2012.

Collophorina paarla (Damm & Crous) Damm & Crous, comb. nov. MycoBank MB821427

≡ *Collophora paarla* Damm & Crous, Persoonia 24: 67. 2010.

≡ *Collophora pallida* Damm & Crous, Persoonia 24: 69. 2010.

Collophorina rubra (Damm & Crous) Damm & Crous, comb. nov. MycoBank MB821428

≡ *Collophora rubra* Damm & Crous, Persoonia 24: 69. 2010.

Notes: The genus *Collophora* Damm & Crous 2010 is illegitimate, being a later homonym of the plant genus *Collophora* Mart. 1830. In order to replace this name, the new name *Collophorina* nom. nov. is provided for this genus. Species of the genus are associated with black spots and dark brown to black streaking of the xylem tissues of woody plant hosts, and can cause wood necrosis (Gramaje et al. 2012; Arzanlou et al. 2016a); it was also collected from air samples (Fischer et al. 2017).

Collostroma Petr. 1947, Ascomycota genera incertae sedis, one species, type: *C. gelatinosum* Petr., coelomycetous, sexual morph unknown, saprobes, Austria, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Colonomyces R.K. Benj. 1955, Euceratomycetaceae, Laboulbeniales, Laboulbeniomycetes, two species, type: *C. appendiculatus* R.K. Benj., asexual morph unknown, biotrophic, USA, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Colpoma Wallr. 1833 (= *Hysterium* Tode 1791; = *Pragmoparopsis* Höhn. 1917 fide Species Fungorum 2017),

Rhytismataceae, Rhytismatales, Leotiomycetes, 14 species, type: *C. quercinum* (Pers.) Wallr., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lantz et al. (2011; phylogeny), Lee et al. (2016; antifungal properties), cultures and sequences are unavailable, available cultures for type: Lantz 368 (UPS).

Columnnodomus Petr. 1941, Ascomycota genera incertae sedis, one species, type: *C. microsporus* Petr. coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Columnnomyces R.K. Benj. 1955, Laboulbeniaceae, Laboulbeniales, Laboulbeniomycetes, two species, type: *C. ptomaphagi* R.K. Benj., asexual morph unknown, biotrophic, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Columnosphaeria Munk 1953, Saccotheciaceae, Dothideales, Dothideomycetes, c. four species, type: *C. sarothamni* Munk, asexual morph unknown, saprobes, terrestrial, worldwide, see Thambugala et al. (2015b; phylogeny), cultures and sequences are available.

Columnnothyrium Bubák 1916, Ascomycota genera incertae sedis, one species, type: *C. myriospermum* (C. Massal.) Bubák, coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coma Nag Raj & W.B. Kendr. 1972 (= *Ascocoma* H.J. Swart 1987 fide Johnston et al. 2014b; Wijayawardene et al. 2016b), Phacidiales genera incertae sedis, Leotiomycetes, two species, type: *C. circularis* (Cooke & Massee) Nag Raj & W.B. Kendr., coelomycetous, sexual morph *Ascocoma*, saprobes, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Wijayawardene et al. (2016b; outline, one name, morphology), available specimens for type: BRIP 14753, cultures and sequences are unavailable.

Comatospora Piroz. & Shoemaker 1971, Ascomycota genera incertae sedis, one species, type: *C. suttonii* Piroz. & Shoemaker, coelomycetous, sexual morph unknown, saprobes, Canada, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Combea De Not. 1846, Opegraphaceae, Arthoniales, Arthoniomycetes, one species, type: *C. mollusca* (Ach.) Nyl., lichenized, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Roccellaceae), Kirk et al. (2013; genus accepted), Ertz and Tehler (2011; taxonomy,

placed in *Opegraphaceae*), Lücking et al. (2017a; classification), sequences are available.

Comesella Speg. 1923, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. anomala* (Berk. & M.A. Curtis) Speg., asexual morph unknown, saprobes, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Comesia Sacc. 1884, *Helotiales* genera *incertae sedis*, three species, type: *C. felicitatis* (P. Crouan & H. Crouan) Sacc., asexual morph unknown, saprobes, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Comminutispora A.W. Ramaley 1996 (= *Hyphospora* A.W. Ramaley 1996 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b), *Dothideomycetes* genera *incertae sedis*, one species, type: *C. agavacearum* A.W. Ramaley, asexual morph hyphomycetous (*Hyphospora*), saprobes, USA, see Lumbsch and Huhndorf (2010; outline; accepted as *Planistromellaceae*), Hyde et al. (2013; did not accept as *Planistromellaceae*); Wijayawardene et al. (2014c; outline, nomenclature), Rossman et al. (2015b; nomenclature), holotype: BPI 802958, cultures and sequences are unavailable.

Comocephalum Syd. 1939, *Ascomycota* genera *incertae sedis*, one species, type: *C. visiae* Syd., coelomycetous, sexual morph unknown, saprobes, Ecuador, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Comoclathris Clem. 1909 (= *Platyspora* Wehm. 1961 *fide* Species Fungorum 2017), *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, 26 species, type: *C. lanata* Clem., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline; accepted as a genus in *Diademaceae*), Hyde et al. (2013; accepted as *Diademaceae*) Wijayawardene et al. (2014c; outline; accepted as a genus in *Diademaceae*), Ariyawansa et al. (2015c; morphology, accepted as *Pleosporaceae*), Li et al. (2016a; new species, Russia), Thambugala et al. (2017b, new species, phylogeny), holotype of the type: COLO 62872, sequences are available.

Complexipes C. Walker 1979, *Ascomycota* genera *incertae sedis*, two species, type: *C. moniliformis* C. Walker, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Seifert et al. (2011 morphology, accepted as a genus in *Otideaceae*, *Pezizales*), Wijayawardene et al. (2012; outline, accepted as a genus in *Pyronemataceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; transferred to *Ascomycota* genera *incertae sedis*), cultures and sequences are unavailable.

Compositrema Rivas Plata, Lücking & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *C. cerebriforme* J.E. Hern. & Lücking, lichenized, see Rivas Plata et al. (2012b), Kraichak et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Compsocladium I.M. Lamb 1956, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *C. archboldianum* I.M. Lamb, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Compsomyces Thaxt. 1894 (= *Moschomyces* Thaxt. 1894 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, seven species, type: *C. verticillatus* (Thaxt.) Thaxt., asexual morph unknown, biotrophic, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Condylospora Nawawi 1976, *Ascomycota* genera *incertae sedis*, four species, type: *C. spumigena* Nawawi, hyphomycetous, saprobes, on decaying leaves, aquatic, Brazil, Malaysia, Poland, Puerto Rico, Republic of Cameroon, Venezuela, Vietnam, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Fiúza et al. (2015c; first record from Amazon), sequence data available for one species, compare with *Angulospora*, isotype of type: IMI 193709.

Coniambigua Etayo & Diederich 1995, *Ascomycota* genera *incertae sedis*, one species, type: *C. phaeographidis* Etayo & Diederich, coelomycetous, sexual morph unknown, lichenicolous, non-lichenized, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2016b, 2017; morphology, outline), Kirk et al. (2013; genus accepted), holotype: MA (Lich.), Etayo 4215, sequence data of type species is unavailable.

Coniangium Fr. 1821 (see Species Fungorum 2017 for synonyms), *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, four species, type: *C. vulgare* Fr., lichenized, see Lumbsch and Huhndorf (2010; outline), Sundin et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Coniarthronia Grube 2001, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, seven species, type: *C. pyrrhula* (Nyl.) Grube, lichenized, see Lumbsch and Huhndorf (2010; outline), Aptroot et al. (2015a; phylogeny), Lücking et al. (2017a; classification), sequences are available but lacking for the type.

Conicomycetes R.C. Sinclair, Eicker & Morgan-Jones 1983, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, four species, type: *C. transvaalensis* R.C. Sinclair, Eicker & Morgan-Jones, hyphomycetous,

worldwide, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Liu et al. (2015a; new species, morphology, DNA), sequences are available for *C. pseudotransvaalensis* (HHUF 29956 *fide* Liu et al. 2015a).

Conidiascus Holterm. 1898, *Saccharomycetes* genera *incertae sedis*, one species, type: *C. paradoxus* Holterm., asexual morphs unknown, terrestrial, Java, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Conidiotheca Réblová & L. Mostert 2007, *Calosphaeriaceae*, *Calosphaeriales*, *Sordariomycetes*, one species, type: *C. tympanoides* (M.E. Barr) Réblová & L. Mostert, asexual morph unknown, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; phylogeny, outline), holotype: holotype of *Romellia tympanoides*, NY.

Coniella Höhn. 1918 (= *Anthasthoopa* Subram. & K. Ramakr. 1956; = *Baeumleria* Petr. & Syd. 1927; = *Cyclodomella* P.N. Mathur, V.V. Bhatt & Thirum. 1959; = *Emboldidium* Bat. 1964; = *Pilidiella* Petr. & Syd. 1927; = *Schizoparme* Shear 1923 *fide* Alvarez et al. 2016; Species Fungorum 2017), *Schizopharmaceae*, *Diaporthales*, *Sordariomycetes*, 29 species, type: *C. pulchella* Höhn., coelomycetous, sexual morph formerly known in *Schizopharme*, saprobes, pathogens, terrestrial, widespread, see Crous et al. (2015c; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; phylogeny, outline), Alvarez et al. (2016; revision, nomenclature, treated *Schizopharme* and *Pilidiella* as synonyms), Wijayawardene et al. (2016b; morphology, phylogeny), cultures and sequences are available.

Coniobrevicolla Réblová 1999, *Trichosphaeriaceae*, *Trichosphaeriales*, *Sordariomycetes*, one species, type: *C. larsenii* Réblová, asexual morph unknown, terrestrial, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Coniocarpon DC. 1805 (see Species Fungorum 2017 for synonyms), *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, five species, type: *C. cinnabarinum* DC., lichenized, see Sundin et al. (2012; taxonomy), Frisch et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Coniocessia Dania García, Stchigel, D. Hawksw. & Guarro 2006, *Coniocessiaceae*, *Xylariales*, *Sordariomycetes*, five species, type: *C. nodulisporioides* (D. Hawksw.) Dania García, Stchigel, D. Hawksw. & Guarro, asexual morph hyphomycetous (nodulisporioides-like), terrestrial, saprobes, coprophilous, from soil, Jordan,

Iran, Spain, see Lumbsch and Huhndorf (2010; outline), Asgari and Zare (2011; new species, DNA), Senanayake et al. (2015; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: IMI 204247 (as *Coniochaeta nodulisporioides* D. Hawksw.), sequence data and cultures available.

Coniochaeta (Sacc.) Cooke 1887 (= *Lecythophora* Nannf. 1934 *fide* Réblová et al. 2016c; see Species Fungorum 2017 for additional synonyms), *Coniochaetaceae*, *Coniochaetales*, *Sordariomycetes*, 78 species, type: *C. lignaria* (Grev.) Cooke, asexual morph formerly known in *Lecythophora*, terrestrial, aquatic, saprobes, coprophilous, widespread, see Damm et al. (2010b; South Africa, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; phylogeny), Réblová et al. (2016c; nomenclature), no ex-type but none-type cultures of the type species are available.

Coniolariella Dania García, Stchigel & Guarro 2006, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, eight species, type: *C. limonispora* (Ellis & Everh.) Checa, Arenal & J.D. Rogers, asexual morph rhinocladiella-like, saprobes, worldwide, see Checa et al. (2008; new species), Lumbsch and Huhndorf (2010; outline), Zare et al. (2010; new species, morphology), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype: IRAN 12363F (as *Coniochaeta gamsii* Asgari & Zare), sequence data available.

Conioscyphula Höhn. 1904 (= *Conioscyphascus* Réblová & Seifert 2004; = *Cylicogone* Emden & Veenb.-Rijks 1974 [1973] *fide* Maharachchikumbura et al. 2016b; Réblová et al. 2016c; Species Fungorum 2017), *Conioscyphaceae*, *Conioscyphales*, *Sordariomycetes*, twelve species, type: *C. lignicola* Höhn., sexual morph formerly known in *Conioscyphascus*, saprobes, terrestrial, worldwide, see Crous et al. (2014c; new species, DNA), Zelski et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Réblová et al. (2016c; nomenclature), cultures and sequences are available.

Conioscyphopsis Goh & K.D. Hyde 1998, *Ascomycota* genera *incertae sedis*, one species, type: *C. australiensis* Goh & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, Australia, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coniosporium Link 1809, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, c. 20 species, type: *C. olivaceum* (Link) Tubaki, hyphomycetous, sexual morph unknown, saprobes, cosmopolitan, see Li et al. (2008a; new species), Seifert et al. (2011 morphology), Tsuneda et al. (2011a, b;

phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), cultures and sequences are available.

Coniothyrina Syd. & P. Syd. 1912 (= *Coniothyriella* Speg. 1911; = *Coniothyris* Clem. 1909 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *C. agavicola* (Speg.) Syd., coelomycetous, sexual morph unknown, saprobes, subtropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coniothyrium Corda 1840 (= *Clisosporium* Fr. 1819; = *Coniothyrinula* Petr. 1923; = *Cyclobium* C. Agardh 1821; = *Dactuliochaeta* G.L. Hartm. & J.B. Sinclair 1988; = *Monoplodia* Westend. 1859 *fide* de Gruyter et al. 2012; Species Fungorum 2017), *Coniothyriaceae*, *Pleosporales*, *Dothideomycetes*, c. 50 species, type: *C. palmarum* Corda, coelomycetous, terrestrial, pathogens, saprobes, worldwide, see de Gruyter et al. (2012; phylogeny, generic concept) Verkley & Gruyter), Crous et al. (2013b, 2016a; new species), Hyde et al. (2013; morphology), Verkley et al. (2014; phylogeny), Wijayawardene et al. (2016b; morphology, phylogeny), Thambugala et al. 2017b; new species, phylogeny), cultures and sequences are available, reference culture for type: CBS 400.71.

Coniozyma Crous 2008, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. leucospermi* (Crous & Denman) Crous, coelomycetous, terrestrial, pathogens, saprobes, South Africa, see Marincowitz et al. (2008; morphology), Wijayawardene et al. (2017; outline), holotype and ex-type: PREM 55348, CBS 114035 = STE-U 2852.

Conunctospora Udagawa & Uchiy. 1999, *Ascomycota* genera *incertae sedis*, one species, type: *C. vesiculifera* Udagawa, Uchiy. & Kamiya, hyphomycetous, sexual morph unknown, saprobes, Japan, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: CBM, Uchiyama SUM3064, cultures and sequences are unavailable.

Conlarium F. Liu & L. Cai 2012, *Sordariomycetes* genera *incertae sedis*, one species, type: *C. dupliciascosporum* F. Liu & L. Cai, sexual morph unknown, terrestrial, pathogens, saprobes, China, see Liu et al. (2012a; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type: HMAS243129, CGMCC3.14939 = GD34-5-2.

Connorsia Malloch 1974, *Pseudeurotiaceae*, *Leotiomycetes* families *incertae sedis*, one species, type: *C. rilstonei* (C. Booth) Malloch, asexual morph unknown, saprobes, British Isles, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted),

available cultures of the type: CBS 537.74, AFTOL-ID 1264.

Conoideocrella D. Johnson, G.H. Sung, Hywel-Jones & Spatafora 2009, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, type: *C. luteorostrata* (Zimm.) D. Johnson, G.H. Sung, Hywel-Jones & Spatafora, two species, asexual morph unknown, saprobes, cosmopolitan, see Johnson et al. (2009; morphology, phylogeny), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Conoplea Pers. 1797 (= *Botryoxylon* Cif. 1962; = *Streblocoalium* Chevall. 1837; = *Streptothrix* Corda 1839 *fide* Species Fungorum 2017), *Sarcosomataceae*, *Pezizales*, *Pezizomycetes*, eleven species, type: *C. sphaerica* (Pers.) Pers., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Marincowitz et al. (2008; DNA), Rivera-Orduña et al. (2011; DNA), Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Conostomatum Doweld 2013 (= *Conostoma* Bat. & J.L. Bezerra 1965 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *C. oenocarpi* (Bat. & J.L. Bezerra) Doweld, asexual morph unknown, fossil *Pteridospermae*, see Index Fungorum (2017).

Conostroma Moesz 1921 (= *Discosporella* Höhn. 1927 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, three species, type: *C. didymum* (Fautrey & Roum.) Moesz (current name: *Colpoma quercinum* (Pers.) Wallr. *fide* Species Fungorum 2017), coelomycetous, sexual morph unknown, saprobes, Asia, see Wijayawardene et al. (2012, 2017; outline), Zhao and Zhao (2012; new species, China), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Conotrempopsis Vězda 1977, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *C. weberiana* Vězda, lichenized, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Consetiella Hol.-Jech. & Mercado 1982, *Ascomycota* genera *incertae sedis*, one species, type: *C. solida* (Berk. & M.A. Curtis) Hol.-Jech. & Mercado, hyphomycetous, sexual morph unknown, saprobes, Cuba, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Constantinomyces Egidi & Onofri 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *C. virgultus* Egidi & Onofri, hyphomycetous, sexual morph unknown, rock-inhabiting, Spain, see Egidi

et al. (2014; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: CBS 117930 = TRN 79.

Constrictolumina Lücking, M.P. Nelsen & Aptroot 2015, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, nine species, type: *C. cinchonae* (Ach.) Lücking, M.P. Nelsen & Aptroot, lichenized, see Nelsen et al. (2014a; taxonomy), Aptroot and Lücking (2016; taxonomy) Lücking et al. (2017a, b; taxonomy), sequences are available.

Cookeina Kuntze 1891 (= *Boedijnopeziza* S. Ito & S. Imai, Trans. 1937; = *Peziza* sect. *Trichoscypha* Cooke 1879; = *Pilocratera* Henn. 1891 [1892]; = *Trichoscypha* (Cooke) Sacc. 1889 *fide* Species Fungorum 2017), *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, 13 species, type: *C. tricholoma* (Mont.) Kuntze, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ekanayaka et al. (2016c; revision), Tibpromma et al. (2017b; reference specimen), cultures and sequences are available.

Cookella Sacc. 1878 (= *Ascomycetella* Peck 1881; = *Phaneroascus* Baudyš 1919 *fide* Species Fungorum 2017), *Cookellaceae*, *Dothideomycetes* families *incertae sedis*, seven species, type: *C. microscopica* Sacc., asexual morph unknown, parasites, fungicolous, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Cooksonomyces H.J. Swart & D.A. Griffiths 1974, *Ascomycota* genera *incertae sedis*, one species, type: *C. banksiae* H.J. Swart & D.A. Griffiths, coelomycetous, sexual morph unknown, saprobes, Australia, see Kirk et al. (2008), however Kirk et al. (2013) did not list, cultures and sequences are unavailable.

Cophiniforma Doilom, J.K. Liu & K.D. Hyde 2012, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, three species, type: *C. eucalypti* Doilom, J.K. Liu & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, South Africa, USA, Thailand, see Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Phillips et al. (2013; taxonomy, new species, phylogeny), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline), holotype and ex-type of the type: MFLU 12-0752, MFLUCC 11-0425.

Coppinsia Lumbsch & Heibel 1998, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, one species, type: *C. minutissima* Lumbsch & Heibel, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hodkinson and Lendemer (2011b; taxonomy), Lücking et al. (2017a; classification), sequences unavailable.

Coprobolus Cain & Kimbr. 1970, *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, one species, type: *C. poculiformis* Cain & Kimbr., asexual morph unknown, coprophilous, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Copromyces N. Lundq. 1967, *Sordariaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *C. bisporus* N. Lundq., asexual morph unknown, coprophilous, terrestrial, Sweden, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Coprotiella Jeng & J.C. Krug 1976, *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *C. gongylospora* Jeng & J.C. Krug, asexual morph unknown, coprophilous, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Coprotus Korf & Kimbr. 1967 (= *Leporina* Velen. 1947 *fide* Species Fungorum 2017), *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, 29 species, type: *C. sexdecimsporus* (P. Crouan & H. Crouan) Kimbr. & Korf, asexual morph unknown, cup-fungus, saprobes or plant pathogenic, widespread, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Corallicola Volk.-Kohlm. & Kohlm. 1992, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *C. nana* Volk.-Kohlm. & Kohlm., asexual morph unknown, saprobes, marine, Belize, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures are unavailable, holotype of type: J.K. 5004 = 01594700, NYBG.

Corallocyostroma Y.N. Yu & Z.Y. Zhang 1980, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *C. oryzae* Y.N. Yu & Z.Y. Zhang, coelomycetous, saprobes or plant pathogenic, widespread, sequence data available only for *C. ornithocopreoides*, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline).

Corallomyctella Henn. 1904 (= *Corallomyces* Berk. & M.A. Curtis 1854 [1853]; = *Rhizostilbella* Wolk 1914 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, type: *C. heissenii* Henn., three species, asexual morph hyphomycetous, terrestrial, saprobes, associated with diseased plants, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Herrera et al. (2013; new

species, revision), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are available.

Corallonectria C. Herrera & P. Chaverri 2013, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. jatropheae* (Möller) C. Herrera & P. Chaverri, asexual morph hyphomycetous, terrestrial, saprobes, associated with diseased plants, cosmopolitan, see Herrera et al. (2013; new species, revision), Lombard et al. (2015b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Cordana Preuss 1851 (= *Porosphaerella* E. Müll. & Samuels 1982; = *Preussiaster* Kuntze 1891; = *Pseudobotrytis* Krzemien. & Badura 1954; = *Umbellula* E.F. Morris 1955 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Cordanaceae*, *Coniochaetales*, *Sordariomycetes*, eight species, type: *C. pauciseptata* Preuss, hyphomycetous, saprobes or plant pathogenic, widespread, sequence data available for all species, see Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2014b; phylogeny), Zelski et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), Hongsanan et al. (2017a; phylogeny, evolution), Wijayawardene et al. (2017; outline), ex-type: IMI 232041a; holotype of type: CBS H-21148.

Cordierites Mont. 1840 (= *Phyllomyces* Lloyd 1920 [1921] *fide* Species Fungorum 2017), *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *C. guianensis* Mont., asexual morph unknown, saprobes, worldwide, see Kirk et al. (2013; genus accepted), Lumbsch and Huhndorf (2010; outline), Peterson and Pfister (2010; DNA, phylogeny), cultures and sequences are available.

Cordyceps Fr. 1818, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, c. 500 species, type: *C. militaris* (L.) Fr., asexual morph unknown, on insect, widespread, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Sanjuan et al. (2015; phylogeny), Tasanathai et al. (2016; new species), cultures and sequences are available.

Coreniella Bubák & Krieg. 1912, *Ascomycota* genera *incertae sedis*, one species, type: *C. cystopodoides* Bubák & Krieg., hyphomycetous, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequence data available for unidentified species.

Coreomyces Thaxt. 1903 (= *Paracoreomyces* R.A. Poiss. 1929 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 21 species, type: *C. corisae* Thaxt., asexual morph unknown, biotrophic, worldwide, see Lee and Na (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coreomycetopsis Thaxt. 1920, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *C. oedipus* Thaxt., asexual morph unknown, biotrophic (on termites), West Indies, USA, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Corethromyces Thaxt. 1892 (= *Eucorethromyces* Thaxt. 1900; = *Rhadinomyces* Thaxt. 1893 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 82 species, type: *C. cryptobii* Thaxt., asexual morph unknown, biotrophic, West Indies, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Corethrostroma Kleb. 1933, *Ascomycota* genera *incertae sedis*, one species, type: *C. laricis* Kleb., coelomycetous, sexual morph unknown, saprobes, Europe, see Kirk et al. (2013; genus accepted), sequence data available for unidentified species.

Cornicularia (Schreb.) Ach. 1794 (= *Lichen* sect. *Cornicularia* Schreb. 1791 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *C. tristis* (Weber ex F.H. Wigg.) Hoffm., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Corniculariella P. Karst. 1884 (= *Chondropodium* Höhn. 1916; = *Collonaemella* Höhn. 1915; = *Cornularia* Sacc. 1884 *fide* Species Fungorum 2017), *Dermateaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *C. abietis* P. Karst., coelomycetous, sexual morph unknown, saprobes, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequence data available for *C. brasiliensis* but type species lack sequences.

Cornipulvina Huhndorf, A.N. Mill., F.A. Fernández & Lodge 2005, *Boloniaceae*, *Boliniales*, *Sordariomycetes*, one species, type: *C. ellipsoides* Huhndorf, A.N. Mill., F.A. Fernández & Lodge, asexual morphs unknown, saprobes, terrestrial, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Cornucopiella Höhn. 1915 (= *Clavularia* P. Karst. 1882 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *C. mirabilis* Höhn., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cornuntum Velen. 1947, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. fagineum* Velen., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; treated as a synonym of *Strossmayeria* Schulzer), cultures and sequences are unavailable.

Cornutispora Piroz. 1973, *Ascomycota* genera *incertae sedis*, five species, type: *C. limaciformis* Piroz., coelomycetous, sexual morph unknown, lichenicolous, worldwide, see Etayo (2010a; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cornutostilbe Seifert 1990, *Ascomycota* genera *incertae sedis*, five species, type: *C. medusa* Seifert, hyphomycetous, sexual morph unknown, from soil, saprobes, Indonesia, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: BO 14356.

Cornuvesica C.D. Viljoen, M.J. Wingf. & K. Jacobs 2000, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, four species, type: *C. falcata* (E.F. Wright & Cain) C.D. Viljoen, M.J. Wingf. & K. Jacobs, asexual morph chalara-like, from soil, saprobes, Europe, N. America, South Africa, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Marincowitz et al. (2015; new species, morphology, phylogeny), sequences are available for all species.

Corollospora Werderm. 1922 (= *Halosigmoidea* Nakagiri et al. 2009; *Peritrichospora* Linder 1944; = *Varicosporina* Meyers & Kohlm. 1965 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, 25 species, type: *C. maritima* Werderm., asexual morph hyphomycetous, saprobes, marine, worldwide, see Abdel-Wahab et al. (2009; new species, phylogeny), Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; diversity), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures and sequences are available.

Coronatomyces Dania García, Stchigel & Guarro 2004, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. cubensis* Dania García, Stchigel & Guarro, asexual morph unknown, saprobes, terrestrial, Cuba, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type: IMI 385316, FMR 7132.

Coronellaria (P. Karst.) P. Karst. 1870 (= *Aquarella* Velen. 1947; = *Peziza* sect. *Coronellaria* P. Karst. 1869 *fide*

Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, four species, type: *C. delitschiana* (Auersw.) P. Karst., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coronopapilla Kohlm. & Volkm.-Kohlm. 1990, *Zopfiaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *C. avellina* Kohlm. & Volkm.-Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Jones et al. (2009b, 2015; classification), Suetrong et al. (2009; morphology), Alias et al. (2010; diversity), Besitulo et al. (2010; diversity), Lee et al. (2012; checklist, Malaysia), Zhang et al. (2012e; morphology), Hyde et al. (2013; morphology, outline), Wijayawardene et al. (2014c; outline), holotype and ex-isotype: J.K. 5144a (IMS), J.K. 5144b (IMS 330805).

Coronophora Fuckel 1864 (= *Cryptosphaerella* Sacc. 1882 *fide* Species Fungorum 2017), *Coronophoraceae*, *Coronophorales*, *Sordariomycetes*, five species, type: *C. gregaria* Fuckel, asexual morph unknown, saprobes, terrestrial, Europe, see Mugambi and Huhndorf (2010; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline; phylogeny), available cultures of type: ANM1555 (*fide* Mugambi and Huhndorf 2010).

Coronophorella Höhn. 1909 (see Species Fungorum 2017 for synonyms), *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *C. chaetomoides* (Penz. & Sacc.) Höhn., asexual morph unknown, saprobes, Indonesia, see Kirk et al. (2008; treated as a synonym of *Nitschkia*), Mugambi and Huhndorf (2010; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), available cultures of type: GKM1099.

Coronoplectrum Brusse 1987, *Lecanorales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *C. namibicum* Brusse, lichenized, see Thell et al. (2012; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Corticifraga D. Hawksw. & R. Sant. 1990, *Ostropales* genera *incertae sedis*, *Lecanoromycetes*, eight species, type: *C. peltigerae* (Fuckel) D. Hawksw. & R. Sant., lichenicolous, terrestrial, widespread, asexual morph unknown, see Zhurbenko (2009; new species, review), Etayo and Sancho (2008; new species), Spribille et al. (2010; new species), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; outline), Pino-Bodas et al. (2017; phylogeny), cultures and sequences are available, holotype: ?G and isotypes K(M)-IMI and S.

Corticiruptor Wedin & Hafellner 1998, *Lecanoromycetes* genera *incertae sedis*, two species, type: *C. abeloneae* (P.M. Jørg.) Wedin & Hafellner, asexual morphs unknown, lichenicolous, terrestrial,

Europe, see Etayo and Sancho (2008; new species), Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Corticorygma M. Cáceres, Feuerstein, Aptroot & Lücking 2016, Graphidaceae, Ostropales, Lecanoromycetes, one species, type: *C. stellatum* M. Cáceres, Feuerst., Aptroot & Lücking, lichenized, see Cáceres et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Corylomyces Stchigel, M. Caldúch & Guarro 2006, Sordariales genera incertae sedis, Sordariomycetes, type: one species, *C. selenosporus* Stchigel, M. Caldúch & Guarro, asexual morph unknown, saprobes, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type culture of type: ILLS 57506, FMR 8279.

Corylophomyces R.K. Benj. 1994, Laboulbeniaceae, Laboulbeniales, Laboulbeniomycetes, 82 species, type: *C. sericoderi* (Santam.) R.K. Benj., asexual morph unknown, biotrophic, West Indies, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Corynascella Arx & Hodges 1975, Chaetomiaceae, Sordariales, Sordariomycetes, seven species, type: *C. humicola* Arx & Hodges, asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), van den Brink et al. (2012; DNA), Maharachchikumbura et al. (2015b, 2016b; outline), Wang et al. (2016c; DNA), sequences are available.

Corynecercospora V.K. Pal, Akhtar, N. Ahmad, Kamal & D.K. Agarwal 2006, Ascomycota genera incertae sedis, one species, type: *C. teraiensis* V.K. Pal, Akhtar, N. Ahmad, Kamal & D.K. Agarwal, hyphomycetous, sexual morph unknown, saprobes, terrestrial, temperate regions, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Corynecystis Brusse 1985, Lichinaceae, Lichinales, Lichinomycetes, one species, type: *C. capensis* Brusse, lichenized, South Africa, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Corynelia Ach. ex Fr. 1818 (= *Alboffia* Speg. 1898 [1899]; = *Endohormidium* Auersw. & Rabenh. 1869 fide Species Fungorum 2017), Coryneliaceae, Coryneliales, Eurotiomycetes, nine species, type species: *C. uberata* Fr., asexual morph unknown, but spermogonia often produced, terrestrial, parasitic on members of the plant family Podocarpaceae, Africa, Central and South America, Japan, Oceania, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wood et al. (2016; phylogeny, new species, proposal of the subclass Coryneliomycetidae), epitype: PREM 61207.

Coryneliella Har. & P. Karst. 1890, Ascomycota genera incertae sedis, one species, type: *C. consimilis* Har. & P. Karst., asexual morph unknown, saprobes, terrestrial, Mauritius, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Coryneliopsis Butin 1972, Coryneliaceae, Coryneliales, Eurotiomycetes, two species, type species: *C. cupulifera* Butin, asexual morph unknown, terrestrial, always associated with members of *Cytaria* Berk. which parasitize *Nothofagus* spp., Chile, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wood et al. (2016; morpho taxonomy), cultures and sequences are unavailable.

Coryneliospora Fitzp. 1942, Coryneliaceae, Coryneliales, Eurotiomycetes, two species, type: *C. fructicola* (Pat.) Fitzp., asexual morph unknown, but spermogonia often produced, terrestrial, parasitic on fruits of *Rapanea* Aubl. and *Myrsine* L., China, India, New Guinea, South Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wood et al. (2016; morpho taxonomy), cultures and sequences are unavailable.

Corynespora Güssow 1906 (= *Corynesporasca* Sivan. 1996 fide Rossman et al. 2015b), Corynesporascaceae, Pleosporales, Dothideomycetes, more than 130 species, type: *C. mazei* Güssow, hyphomycetous, sexual morph formerly known in *Corynesporasca*, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline, sexual morph), Zhang et al. (2009d; phylogeny), Ma et al. (2011c; new species), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Rossman et al. (2015b; nomenclature), sequences are available for two species, type sequence data are not available.

Corynesporella Munjal & H. S. Gill 1961, Ascomycota genera incertae sedis, ten species, type: *C. urticae* Munjal & H.S. Gill, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011 morphology), Zhang et al. (2012f; new species), Ma et al. (2012g; new species), Xia et al. (2014a; new species; key to species), cultures and sequences are unavailable.

Corynesporina Subram. 2009 (= *Corynesporina* Subram. 1994 fide Species Fungorum 2017), Ascomycota genera incertae sedis, one species, type: *C. elegans* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Singapore, see Gams et al. (2009; taxonomy), Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), holotype: MUBL 3523, cultures and sequences are unavailable.

Corynesporopsis P. M. Kirk 1981 (= *Monotretomyces* Morgan-Jones, R.C. Sinclair & Eicker 1987 fide Species Fungorum 2017), Ascomycota genera incertae sedis, 15 species, type: *C. quercicola* (Borowska) P.M. Kirk,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Castañeda-Ruiz et al. (2010d; new species), Seifert et al. (2011 morphology), Ma et al. (2010a, 2012b; new species, China), Wijayawardene et al. (2012, 2017; outline), Hernández-Restrepo et al. (2014b; new species), Xia et al. (2013a, 2014b; new species), Kirschner (2015; new species), and sequences are available but lacking for the type species.

Coryneum Nees 1816 (= *Murogenella* Goos & E.F. Morris 1965; = *Pseudovalsa* Ces. & De Not. 1863 *fide* Rossman et al. 2015a; Species Fungorum 2017), *Pseudovalsaceae*, *Diaporthales*, *Sordariomycetes*, c. 30 species, type: *C. umbonatum* Nees, sexual morph *Pseudovalsa*, pathogens, saprobes, terrestrial, temperate regions, see Wijayawardene et al. (2012, 2016b, 2017; outline, morphology, new species), Kirk et al. (2013; genus accepted), Rossman et al. (2015a; nomenclature), cultures and sequences are unavailable.

Coscinocladium Kunze 1846, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, two species, type: *C. occidentale* Kunze, lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Cosmariospora Sacc. 1880, *Ascomycota* genera *incertae sedis*, one species, type: *C. bizzozeriana* Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cosmospora Rabenh. 1862 (= *Chrysoglutin* Briosi & Farneti 1904 *fide* Species Fungorum 2017), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, c. 50 species, type: *C. coccinea* Rabenh., asexual morph acremonium-like, on fruiting bodies and stromata of other fungi or isolated from soil, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Costanetaea Bat. & J.L. Bezerra 1963, *Ascomycota* genera *incertae sedis*, one species, type: *C. caryotae* Bat. & J.L. Bezerra, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Crandallia Ellis & Sacc. 1897, *Ascomycota* genera *incertae sedis*, four species, type: *C. juncicola* Ellis & Sacc., coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, ?worldwide, see Marinowitz et al. (2008; new species), Wijayawardene et al. (2012, 2017;

outline), Rossman et al. (2016b; new combinations for pathogens, cultures and sequences are unavailable).

Craneomyces Morgan-Jones, R.C. Sinclair & Eicker 1987, *Ascomycota* genera *incertae sedis*, one species, type: *C. verrucosus* Morgan-Jones, R.C. Sinclair & Eicker, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Craspedodidymum Hol.-Jech. 1972, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, twelve species, type: *C. elatum* Hol.-Jech., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Ma et al. (2011c; new species), Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Mel'nik et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Crassicarpon Y. Marín, Stchigel, Guarro & Cano 2015, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *C. thermophilum* (Fergus & Sinden) Y. Marín, Stchigel, Guarro & Cano, asexual morph hyphomycetous, from soil, USA, see Maharachchikumbura et al. (2015b, 2016b; outline), Marin-Felix et al. (2015; taxonomy, phylogeny), cultures and sequences are available.

Crassimassarina A. Hashim. & Kaz. Tanaka 2017, *Lophiotremataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *C. macrospora* A. Hashim. & Kaz. Tanaka, asexual morph coelomycetous, saprobes, Asia, see Hashimoto et al. (2017; taxonomy, phylogeny), sequences are available.

Crassiparies M. Matsum., K. Hiray. & Kaz. Tanaka 2016, *Pleosporales* *incertae sedis*, *Dothideomycetes*, one species, type: *C. quadrisporus* M. Matsum., K. Hiray. & Kaz. Tanaka, asexual morph unknown, saprobes, terrestrial, Japan, see Li et al. (2016a; morphology, phylogeny), holotype and ex-type: HHUF 30409, MAFF 245408.

Crassoascus Checa, Barrasa & A.T. Martínez 1993, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *C. fusisporus* Checa, Barrasa & A.T. Martínez asexual morph unknown, saprobes, terrestrial, Argentina, Spain, see Lumbsch and Huhndorf (2010; outline), Catania and Romero (2012; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, treated as *Clypeosphaeriaceae*), Hongsanian et al. (2017a; phylogeny, evolution, treated *Clypeosphaeriaceae* as a synonym of *Xylariaceae*), holotype: MA-Fungi 26331, cultures and sequences are unavailable.

Crassochaeta Réblová 1999, *Chaetosphaerellaceae*, *Coronophorales*, *Sordariomycetes*, two species, type:

C. nigrita (Sacc.) Rébllová, asexual morph unknown, saprobes, terrestrial, Argentina, Spain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype: K37405, cultures and sequences are available.

Cratiria Marbach 2000, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 20 species, type: *C. lauri-cassiae* (Fée) Marbach, lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Crauatamyces Viégas 1944, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. eupatori* Viégas, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Creodiplodina Petr. 1957, *Ascomycota* genera *incertae sedis*, one species, type: *C. fusispora* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Creographa A. Massal. 1860 (see Species Fungorum 2017 for synonyms), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 30 species, type: *C. brasiliensis* A. Massal., lichenized, see Rivas Plata et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Creonecte Petr. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *C. biparasitica* Petr., coelomycetous, sexual morph unknown, fungicolous (on *Uredo*), terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Creoseptoria Petr. 1937, *Ascomycota* genera *incertae sedis*, one species, type: *C. watzlii* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Caucasus, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Creosphaeria Theiss. 1910, *Lopadostomataceae*, *Xylariales*, *Sordariomycetes*, two species, type: *C. sassafras* (Schwein.) Y.M. Ju, F. San Martín & J.D. Rogers, asexual morph unknown, saprobes, terrestrial, Argentina, Spain, see Lumbsch and Huhndorf (2010; outline), Raja et al. (2011a; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Sir et al. (2016; DNA), Wendt et al. (2017; transferred to *Lopadostomataceae*), cultures and sequences are available.

Creothyriella Bat. & C.A.A. Costa 1957, *Ascomycota* genera *incertae sedis*, one species, type: *C. sideroxyfoliae* Bat. & C.A.A. Costa, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cresponea Egea & Torrente 1993, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, 21 species; type: *C. premnea* (Ach.) Egea & Torrente, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ertz et al. (2015b, phylogeny), Lücking et al. (2017a; classification, accepted as a genus in *Opegraphaceae*), sequences are available.

Creporhaphis M.B. Aguirre 1991, *Trichosphaeriaceae*, *Trichosphaeriales*, *Sordariomycetes*, two species, type: *C. wienkampii* (J. Lahm ex Hazsl.) M.B. Aguirre, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Harada (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cribropeltis Tehon 1933, *Ascomycota* genera *incertae sedis*, one species, type: *C. citrullina* Tehon, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Criella (Sacc.) Henn. 1900 (= *Phaeorhytisma* Henn. 1899 [1900]; = *Rhytisma* sect. *Criella* Sacc. 1889 *fide* Species Fungorum 2017), *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *C. austrocaledonica* (Crié) Sacc. & P. Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Crinigera I. Schmidt 1985 (= *Crinigera* I. Schmidt 1969 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *C. maritima* I. Schmidt, asexual morph unknown, marine, temperate, on wood, saprobes, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Crinitospora B. Sutton & Alcorn 1985, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. pulchra* B. Sutton & Alcorn, coelomycetous, sexual morph unknown, saprobes, Australia, Thailand, see Kirk et al. (2013; genus accepted), Crous et al. (2014a; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), epitype and ex-epitype culture of type: CBS H-21729, CBS 138014.

Criserospaeria Speg. 1912, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. phyllostictoides* Speg., asexual morph unknown, saprobes, terrestrial, Argentina, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cristula Chenant. 1920, *Ascomycota* genera *incertae sedis*, one species, type: *C. integra* Chenant., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011) did not accept the genus, Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cristulariella Höhn. 1916 (= *Nervostroma* Narumi & Y. Harada 2006 *fide* Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *C. depraedans* (Cooke) Höhn., hyphomycetous, sexual morph *Nervostroma*, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, sexual morph), Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Cho et al. (2015; DNA, pathogens), cultures and sequences are available.

Crocellina Tehler & Ertz 2014, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *C. cinerea* (Müll. Arg.) Tehler & Ertz, lichenized, see Ertz et al. (2015b, c; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Crocicreas Fr. 1849 (= *Ascoverticillata* Kamat, Subhedar & V.G. Rao 1979 [1978]; = *Belonidium* sect. *Podobelonium* Sacc. 1889; = *Belonioscypha* Rehm 1893 [1896]; = *Belonium* sect. *Scelobelonium* Sacc. 1889; = *Conchatium* Velen. 1934; = *Davincia* Penz. & Sacc. 1902 [1901]; = *Exotrichum* Syd. & P. Syd. 1914; = *Podobelonium* (Sacc.) Penz. & Sacc. 1902 [1901]; = *Scelobelonium* (Sacc.) Höhn. 1905 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, five species, type: *C. gramineum* (Fr.) Fr., asexual morph unknown, saprobes, terrestrial, widespread (esp. north temperate), see Lumbsch and Huhndorf (2010; outline), Whitton et al. (2012b; new species), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for type species.

Crocodia Link 1833 (= *Crocodia* Link 1833; = *Cyanisticta* Gyeln. 1931; = *Diphaeosticta* Clem. 1909; = *Diphanosticta* Clem. 1909; = *Merostictina* Clem. 1909; = *Parmosticta* Nyl. 1875; = *Parmostictina* Nyl. 1875; = *Phaeosticta* Trevis. 1869; = *Phanosticta* Clem. 1909; = *Podostictina* Clem. 1909; = *Saccardoa* Trevis. 1869; = *Stictina* Nyl. 1860 *fide* Species Fungorum 2017), *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, five species; type: *C. aurata* (Ach.) Link, lichenized, see Galloway and Elix (2013; taxonomy), Moncada et al. (2013a; taxonomy),

Lücking et al. (2017a; classification), sequences are unavailable.

Crocynia (Ach.) A. Massal. [nom. cons.] 1860 (= *Lecidea* sect. *Crocynia* Ach. 1810; = *Symplocia* A. Massal. 1854 *fide* Species Fungorum 2017), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *Lecidea gossypina* (Sw.) Ach., lichenized, tropical, see Ekman et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Crotone Theiss. & Syd. 1915, *Dothideomycetes* genera *incertae sedis*, one species, type: *C. drimydis* (Lév.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, as *Dothideomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Crucellisporiopsis Nag Raj 1983, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *C. gelatinosa* Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, Venezuela, New Zealand, Zambia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014d; new species, morphology, DNA), holotype: PDD 41405, sequence data available for *C. marquesiae* P.W. Crous (holotype CBS H-21977, culture ex-type CPC 22539 = CBS 138895 *fide*).

Crucellisporium M.L. Farr 1968, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *C. selaginellae* M.L. Farr, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, Tanzania, see Marincowitz et al. (2010; new species, DNA), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for *C. umtamvunae* Marinc. et al.

Cruentotrema Rivas Plata, Papong, Lumbsch & Lücking 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *C. cruentatum* (Mont.) Rivas Plata, Lumbsch & Lücking, asexual morph unknown, lichenized, terrestrial, tropical, placement in *Graphidaceae* supported by molecular data, see Sipman et al. (2012; key), Rivas Plata et al. (2012a; taxonomy), Lumbsch et al. (2014a, b), Kalb et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Crumenella P. Karst. 1890 (= *Obscurodiscus* Raitv. 2002 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. myricae* (P. Karst.) P. Karst., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Crumenulopsis J.W. Groves 1969 (= *Crumenula* Rehm 1889 [1896]; = *Digitosporium* Gremmen 1953 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *C. pinicola* (Rebent.) J.W. Groves, asexual morph *Digitosporium*, saprobes, terrestrial, Europe, North America, see Johnston et al. (2014b; nomenclature) Pärtel et al. (2017; DNA), Wijayawardene et al. (2016b; morphology of *Digitosporium*).

Crustodiplodina Punith. 1988, *Ascomycota* genera *incertae sedis*, one species, type: *C. lichenoides* (A.L. Sm.) Punith., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Crustospathula Aptroot 1998, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *C. cartilaginea* Aptroot, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Crutarndina Parnmen, Lücking & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *C. petractoides* (P.M. Jørg. & Brodo) Parnmen, Lücking & Lumbsch, lichenized, see Parnmen et al. (2012; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Cryomyces Selbmann, de Hoog, Mazzaglia, Friedmann & Onofri 2005, *Dothideomycetes*, genera *incertae sedis*, four species, type: *C. antarcticus* Selbmann, de Hoog, Mazzaglia, Friedmann & Onofri, sexual morph unknown, rock-inhabiting, Antarctica and Italy, holotype CBS 116301 = CCFEE 534 (NCBI accession numbers: DQ028269 (ITS), DQ066713 (SSU)), taxonomic species delimitation supported by molecular data.

Additional species: *C. minteri* Selbmann, de Hoog, Mazzaglia, Friedmann & Onofri. For diagnosis see Studies in Mycology 51: 21 (2005); *C. montanus* Isola & Zucconi, and *C. funiculosus* Selbmann & de Hoog, described from the Alps (Selbmann et al. 2014). These last two taxa were invalidly published, since the holotypes are missing in the original publication [ICN (Melbourne) Art. 40]; the two species are here validated as follows:

Cryomyces montanus Isola & Zucconi sp. nov, MB819850. For diagnosis see Fungal Diversity 65(1): 177 (2014). Holotype: CBS 135221. From rock, Val de la Mare, Passo dello Stelvio, Italy; =CCFEE 5476. culture preserved in liquid nitrogen and in dried condition (NCBI accession numbers: KC315849 (ITS), GU250352 (SSU), GU250394 (LSU), GU250434 (mtSSU)), taxonomic species delimitation supported by molecular data.

Cryomyces funiculosus Selbmann & de Hoog sp. nov, MB819849. For diagnosis see Fungal Diversity 65(1):

175 (2014). Holotype: CBS 134687. From rock, Colle delle Pisse, Monte Rosa, Italy; =CCFEE 5554. culture preserved in liquid nitrogen and in dried condition (NCBI accession number: KC315848 (ITS)), taxonomic species delimitation supported by molecular data.

Cryphonectria (Sacc.) Sacc. & D. Sacc. 1905 (= *Nectria* subgen. *Cryphonectria* Sacc. 1883 *fide* Species Fungorum 2017), *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *N. parasitica* D. Sacc., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Gryzenhout et al. (2009; new species, revision), Lumbsch and Huhndorf (2010; outline), Braganca et al. (2011; new species), Maharachchikumbura et al. (2015b, 2016b; taxonomy, phylogeny, outline), cultures and sequences are available.

Cryptandromyces Thaxt. 1912 (= *Peyerimhoffiella* Maire 1916 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 13 species, type: *C. geniculatus* Thaxt., asexual morph unknown, biotrophic, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptascoma Ananthap. 1988, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. betulae* Ananthap., asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Cryptendoxyla Malloch & Cain 1970, *Cephalothecaceae*, *Sordariomycetes*, families *incertae sedis*, one species, type: *C. hypophloia* Malloch & Cain, asexual morph hyphomycetous (chalara-like), from soil, on human as pathogen, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Lysková et al. (2014; DNA, human pathogen), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), living cultures and sequence data available CBS 796.84, CBS 508.70, CBS 138.78.

Cryptphonia Frisch & G. Thor 2010, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, c. 15 species, type: *C. polillensis* (Vain.) Frisch & G. Thor, lichenized, widespread tropical, see Frisch and Thor (2010; taxonomy), Menezes et al. (2013b; taxonomy, key), Kirk et al. (2013; genus accepted), Frisch et al. (2014; systematics), Lücking et al. (2017a; classification), sequences are available but lacking for the type.

Cryptoascus Petri 1909, *Sordariomycetes* genera *incertae sedis*, two species, type: *C. oligosporus* Petri, saprobes or pathogens, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cryptoceuthospora Petr. 1921, *Ascomycota* genera *incertae sedis*, two species, type: *C. moravica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptocline Petr. 1924 (= *Gloeotrochila* Petr. 1947 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, 15 species, type: *C. effusa* Petr., coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptoclypeus A. Hashim. & Kaz. Tanaka 2017, *Lophiotremataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *C. ryukyuensis* A. Hashim. & Kaz. Tanaka, asexual morph coelomycetous, saprobes, Asia, see Hashimoto et al. (2017; taxonomy, phylogeny), sequences are available.

Cryptocoryneopsis B. Sutton 1980, *Ascomycota* genera *incertae sedis*, one species, type: *C. umbraculiformis* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, Australia, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: IMI 234264a.

Cryptocoryneum Fuckel 1870, *Cryptocoryneaceae*, *Pleosporales*, *Dothideomycetes*, 20 species, type: *C. fasciculatum* Fuckel, hyphomycetous, dictyosporium-like, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), da Silva et al. (2015; morphology), Hashimoto et al. (2016; morphology, phylogeny), holotype: PRM 155688, type species lacking cultures, seven species have sequence data.

Cryptodictyon A. Massal. 1860 (= *Amphischizonia* Mont. 1856 *fide* Species Fungorum 2017), *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, two species, type: *P. holleana* Mont. & Bosch, lichenized, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Cryptodiscus Corda 1838 (= *Diplocryptis* Clem. 1909; = *Pseudostictis* Lambotte 1887 [1888] *fide* Species Fungorum 2017), *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, nine species; type: *C. pallidus* (Pers.) Corda, saprobic rarely lichenized, see Baloch et al. (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Cryptohymenium Samuels & L.M. Kohn 1987, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *C. pycnidiphorum* Samuels & L.M. Kohn,

asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptolechia A. Massal. 1853 (= *Gyalectina* Vězda 1969 *fide* Species Fungorum 2017), *Gyalectaceae*, *Ostropales*, *Lecanoromycetes*, 13 species, type: *C. carneolutea* (Turner) A. Massal., with ascomata, lichenized (*Trentepohlia*), terrestrial (corticulous or rarely lignicolous), mainly tropical, rarely noted in the Northern Hemisphere see Mangold et al. (2008b; DNA), Lumbsch and Huhndorf (2010; outline), Kalb et al. (2012; new species), Kirk et al. (2013; genus accepted), Lücking et al. (2017a, classification), sequences are available.

Cryptoleptosphaeria Petr. 1923, *Diaporthales* genera *incertae sedis*, one species, type: *C. moravica* Petr., asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cryptomeliola S. Hughes & Piroz. 1997, *Meliolaceae*, *Meliolales*, *Sordariomycetes*, three species, type: *C. orbicularis* (Berk. & M.A. Curtis) S. Hughes & Piroz., asexual morph produced from the hyphae, saprobes, terrestrial, Cuba, Kenya, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cryptometriion Gryzenh. & M.J. Wingf. 2010, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. aestuescens* Gryzenh. & M.J. Wingf., asexual morphs coelomycetous, pathogens, terrestrial, North Sumatra, Indonesia, see Gryzenhout et al. (2010; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; taxonomy, phylogeny, outline), holotype and culture ex-type: CMW 22535.

Cryptomycella Höhn. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *C. pteridis* (Kalchbr.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptomyces Grev. 1825, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *C. wauchii* Grev., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Cryptomycetaceae*), Lantz et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; treated *Cryptomycetaceae* as a synonym of *Rhytismataceae*), cultures and sequences are available.

Cryptomycina Höhn. 1917, *Sordariomycetes*, genera *incertae sedis*, one species, type: *C. pteridis* (Rebent.) Höhn., asexual morph ? *Cryptomyces*, saprobes, terrestrial, North temperate, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cryptonectriella (Höhn.) Weese 1919, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. biparasitica* (Höhn.) Weese, see Kirk et al. (2008; treated as a synonym of *Nectriella* Nitschke ex Fuckel), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Cryptonectriopsis (Höhn.) Weese 1919, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, type: *C. biparasitica* (Höhn.) Höhn., see Kirk et al. (2008; treated as a synonym of *Phomatospora* Sacc.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Cryptoniesslia Scheuer 1993, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. setulosa* Scheuer, asexual morph unknown, saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cryptopezia Höhn. 1919, *Helotiales* genera *incertae sedis*, one species, type: *C. mirabilis* Höhn., asexual morph unknown, saprobes, terrestrial, Samoa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptophaea Van den Broeck & Ertz 2016, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *C. phaeospora* Van den Broeck & Ertz, asexual morph unknown, lichenized, terrestrial, Democratic Republic of the Congo, see Van den Broeck and Ertz (2016; morphology, phylogeny), Lücking et al. (2017a; classification), holotype and ex-type: Van den Broeck 5964 (BR).

Cryptophiale Piroz. 1968, *Ascomycota* genera *incertae sedis*, 18 species, type: *C. kakombensis* Piroz., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011 morphology, accepted as a genus in *Chaetosphaeriaceae*, *Chaetosphaerales*), Whitton et al. (2012a; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptophialoidea Kuthub. & Nawawi 1987, *Ascomycota* genera *incertae sedis*, three species, type: *C. secunda* (Kuthub. & B. Sutton) Kuthub. & Nawawi, hyphomycetous, sexual morph unknown, saprobes,

terrestrial, Cuba, Malaysia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptosphaerella Sacc. 1882 (= *Cryptosphaerella* Sacc. 1882 *fide* Species Fungorum 2017), *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, c. 15 species, type: *C. nitschkei* (Auersw.) Sacc., asexual morph unknown, saprobes, terrestrial, Cuba, Malaysia, see Lumbsch and Huhndorf (2010; outline), Mugambi and Huhndorf (2010; DNA, phylogeny), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures and sequences are available.

Cryptosphaeria Ces. & De Not. 1863 (= *Cladosphaeria* Nitschke ex Jacz. 1894; = *Cryptosphaerina* Lambotte & Fautrey ex Sacc. & P. Syd. 1902; = *Stigmatopsis* Traverso 1906 *fide* Species Fungorum 2017), *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, c. 13 species, type: *C. millepunctata* Grev., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Trouillas et al. (2015; new species, USA, DNA, Phylogeny), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures and sequences are available.

Cryptosporella Sacc. 1877 (= *Disculina* Höhn. 1916; = *Cryptospora* Tul. & C. Tul. 1863; = *Cryptospora* subgen. *Winterella* Sacc. 1883; = *Cryptosporina* Höhn. 1905; = *Hystricula* Cooke 1884; = *Ophiovalsa* Petr. 1966; = *Winterella* (Sacc.) Kuntze 1891 *fide* Mejia et al. 2008; Rossman et al. 2015a; Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, c. 26 species, type: *C. hypodermia* (Fr.) Sacc., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Mejia et al. (2008, 2011; new species, DNA, Phylogeny), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures and sequences are available.

Cryptosporium Kunze 1817 (= *Cryptomela* Sacc. 1884 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, 25 species, type: *C. atrum* Kunze, coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptostroma P.H. Greg. & S. Waller 1952, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *C. corticale* (Ellis & Everh.) P.H. Greg. & S. Waller, hyphomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cryptothecia Stir. 1876 nom. cons. (= *Herpothallon* Tobler 1937; = *Herpothallomyces* Cif. & Tomas. 1954; = *Hypochnus* Fr. ex Ehrenb. 1820; = *Myriostigma* Kremp. 1874; = *Myxotheca* Ferd. & Winge 1910 *fide* Species

Fungorum 2017), Arthoniaceae, Arthoniales, Arthoniomycetes, c. 65 species, type: *C. subnidulans* Stirt., asexual morph unknown, lichenized, epiphytic and foliicolous, widespread, pantropical to subtropical, see Ertz et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Bungartz et al. (2013a; taxonomy), Kirk et al. (2013; genus accepted), Frisch et al. (2014; phylogeny: genus *Myriostigma* reinstated for the *Cryptothecia candida* group), Jagadeesh Ram and Sinha (2016; world key), Lücking et al. (2017a; classification), sequences are available.

Cryptothele Th. Fr. 1866 (= *Cryptotheliomyces* Cif. & Tomas. 1953; = *Malmgrenia* Vain. 1939; = *Phylliscum* subgen. *Phylliscum* Nyl. 1855; = *Pyrenopsidium* (Nyl.) Forssell 1885 fide Species Fungorum 2017), Lichinaceae, Lichinales, Lichinomycetes, six species, type: *C. permiscens* (Nyl.) Th. Fr., asexual morph unknown, lichenized, saxicolous, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Cryptovalsa Ces. & De Not. ex Fuckel 1870 (= *Allescherina* Berl. 1902 fide Species Fungorum 2017), Diatrypaceae, Xylariales, Sordariomycetes, 24 species, type: *C. protracta* (Pers.) De Not., asexual morph coelomycetous, saprobes, pathogens, worldwide, see Lumbsch and Huhndorf (2010; outline; as *Sordariomycetes*, genera *incertae sedis*), Trouillas et al. (2010; pathogens), Senanayake et al. (2015; outline), Li et al. (2016a; coelomycetous asexual morph), Maharachchikumbura et al. (2016b; outline), Abdel-Wahab et al. (2017; new species), cultures and sequences are available.

Cryptumbellata Udagawa & Uchiy. 1999, Ascomycota genera *incertae sedis*, one species, type: *C. terricola* Udagawa & Uchiy., hyphomycetous, sexual morph unknown, saprobes, terrestrial, China, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ctenomyces Eidam 1880, Arthrodermataceae, Onygenales, Eurotiomycetes, one species, type: *C. serratus* Eidam, asexual morph hyphomycetous, saprobes, terrestrial, common on keratinous substrates, especially feathers, also occurring in soil and on dung, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Guarro et al. (2012; morphology), van den Brink et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), de Hoog et al. (2016a; phylogeny, outline), ex-neotype: CBS 187.61.

Ctenosporium R. Kirschner 2006, Ascomycota genera *incertae sedis*, one species, type: *C. pectinatum* R. Kirschner, hyphomycetous, sexual morph unknown,

saprobes, terrestrial, Panama, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Cubasina R.F. Castañeda 1986, Ascomycota genera *incertae sedis*, two species, type: *C. albofusca* R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: INIFAT C85/154, cultures and sequences are unavailable.

Cucujomyces Speg. 1917 (= *Stephanomyces* Speg. 1917 fide Species Fungorum 2017), Laboulbeniaceae, Laboulbeniales, Laboulbeniomycetes, 20 species, type: need typification, asexual morph unknown, biotrophic, worldwide, see Rossi and Weir (2008; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cucullosporella K.D. Hyde & E.B.G. Jones 1986 (= *Cucullospora* K.D. Hyde & E.B.G. Jones 1986 fide Species Fungorum 2017), Halosphaeriaceae, Microascales, Sordariomycetes, one species, type: *C. mangrovei* (K.D. Hyde & E.B.G. Jones) K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; DNA, phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), cultures and sequences are available.

Cucurbitodthis Petr. 1921, Cucurbitariaceae, Pleosporales, Dothideomycetes, one species, type: *C. pityophila* (J.C. Schmidt & Kunze) Petr., asexual morph unknown, saprobes, terrestrial, worldwide, see Ariyawansa et al. (2014f; accepted as a genus in Cucurbitariaceae), Wijayawardene et al. (2014c; outline), cultures and sequences are available but needs verification.

Cucurbitaria Gray 1821 (= *Crotonocarpia* Fuckel 1870 [1869–70]; = *Cucurbitariopsis* Vassilkov 1960; = *Cyathisphaera* Dumort. 1822; = *Gemmamyces* Casagr. 1969; = *Gibberidea* (Fr.) Kuntze 1898; = *Leucothyridium* Speg. 1909; = *Megaloseptoria* Naumov 1925; = *Megalospora* Naumov 1927; = *Phialospora* Raf. 1832; = *Sphaeria* b Gibberidea Fr. 1849; = *Sphaeria* ser. *Erumpentes* Fr. 1849 fide Species Fungorum 2017), Cucurbitariaceae, Pleosporales, Dothideomycetes, c. 40 species, type: *C. berberidis* (Pers.) Gray, asexual morph coelomycetous, with pyrenochaeta-like asexual morph, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Doilom et al. (2013; taxonomy, typification), Hyde et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), ex-epitype: CB1 = CBS 130007 = MFLUCC11-0384; Epitype of type: MFLU 12-0111.

Cucurbitopsis Bat. & Cif. 1957, Sordariomycetes genera *incertae sedis*, one species, type: *C. ramulorum*

Bat., Nascim. & Cif., asexual morph unknown, saprobes, pathogens, worldwide, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Cudonia Fr. 1849 (= *Leotiella* Plötzn. 1900 *fide* Species Fungorum 2017), *Cudoniaceae*, *Rhytismatales*, *Leotiomycetes*, nine species, type: *C. circinans* (Pers.) Fr., asexual morph unknown, saprobes, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for type: AFTOL-ID 353.

Cudoniella Sacc. 1889 (= *Helotium* Pers. 1801; = *Isosoma* Svrček 1989; = *Peziza* b *Helotium* (Pers.) Fr. 1822 *fide* Species Fungorum 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, 31 species, type: *C. queletii* (Fr.) Sacc., asexual morph hyphomycetous (*Trichladium*?), saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hustad and Miller (2011; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available, type lacking sequence data.

Cudoniopsis Speg. 1925, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *C. pusilla* Speg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and cultures and sequences are unavailable.

Culbersonia Essl. 2000, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, two species; type: *C. americana* Essl., lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Prieto et al. (2013; phylogeny) Prieto and Wedin (2016; phylogeny), Lücking et al. (2017a; classification), sequences are unavailable.

Culicidospora R.H. Petersen 1960, *Ascomycota* genera *incertae sedis*, two species, type: *C. aquatica* R.H. Petersen, hyphomycetous, sexual morph unknown, saprobes, aquatic, Newfoundland, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and cultures and sequences are unavailable.

Culicinomyces Couch, Romney & B. Rao 1974, *Ascomycota* genera *incertae sedis*, two species, type: *C. clavisporus* Couch, Romney & B. Rao, hyphomycetous, sexual morph unknown, on mosquitos, aquatic, USA, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequence data available for *Culicinomyces bisporalis* Sigler, Frances & C. Panter, type lacking sequence data.

Cumulospora I. Schmidt 1985, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, one species, type: *C. marina* I. Schmidt, hyphomycetous, sexual morph unknown, saprobes, marine, worldwide, see Abdel-Wahab et al. (2010; phylogeny), Seifert et al. (2011

morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cupulina Dougoud, Van Vooren & M. Vega 2015, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *C. montana* Dougoud, Van Vooren & M. Vega, asexual morph unknown, saprobes, terrestrial, Europe, see Dougoud et al. (2015; morphology, taxonomy), holotype and isotype: G00262366 (herbarium Geneva), R. Dougoud RD32.43.544.13.

Cupulomyces R.K. Benj. 1992, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *C. lasiochili* (Thaxt.) R.K. Benj., asexual morph unknown, biotrophic, Grenada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Curreya Sacc. 1883 (= *Cucurbitodothis* Petr. 1921 *fide* Species Fungorum 2017), *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *C. conorum* (Fuckel) Sacc., asexual morph coelomycetous, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Doilom et al. (2013; accepted as a genus in *Cucurbitariaceae*), Hyde et al. (2013; accepted as a genus in *Cucurbitariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), ex-holotype: S F8461.

Curucispora Matsush. 1981, *Ascomycota* genera *incertae sedis*, three species, type: *C. ponapensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Curvatispora V.V. Sarma & K.D. Hyde 2001, *Sordariomycetes* genera *incertae sedis*, one species, type: *C. singaporensis* V.V. Sarma & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Singapore, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Curvicladiella Decock & Crous 2006 (= *Curvicoladium* Decock & Crous 1998 nom. inval. Art. 53), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. cignea* (Decock & Crous) Decock & Crous, asexual morph hyphomycetous, saprobes, terrestrial, French Guiana, see Wijayawardene et al. (2012, 2017; outline), Lombard et al. (2015b; taxonomy, DNA, phylogeny), Maharachchikumbura et al. (2016b; outline), holotype and ex-type: MUCL 40269, CPC 1595 = CBS 109167.

Curvularia Boedijn 1933 (= *Curvisporium* Corbetta 1963; = *Malustela* Bat. & J.A. Lima 1960; = *Pseudocochliobolus* Tsuda, Ueyama & Nishih. 1978 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b; Species Fungorum 2017), *Pleosporaceae*, *Pleosporales*,

Dothideomycetes, more than 100 species, type: *C. lunata* (Wakker) Boedijn, sexual morph formerly known in *Pseudocochliobolus* hyphomycetous, pathogenic or saprobes on plant or humans, terrestrial, worldwide, see Seifert et al. (2011 morphology), Manamgoda et al. (2012, 2014, 2015; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Ariyawansa et al. (2015c; taxonomy, phylogeny), Rossman et al. (2015b; nomenclature), sequence data available for more than 50 species, ex-neotype: CBS 730.96.

Curvulariopsis M.B. Ellis 1961, *Ascomycota* genera *incertae sedis*, one species, type: *C. cymbisperma* (Pat.) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, Ecuador, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Cuspidatispore Shearer & Bartolata 2006, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. xiphago* Shearer & Bartolata, asexual morph unknown, saprobes, aquatic, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: ILL, Shearer A184-1, cultures and sequences are available.

Custingophora Stolk, Hennebert & Klopotek 1968, *Gondwanamycetaceae*, *Microascales*, *Sordariomycetes*, eight species, type: *C. olivacea* Stolk, Hennebert & Klopotek, sexual morph unknown, saprobes, terrestrial, worldwide, see Kolarík and Hulcr (2009; new species), Nagaraju et al. (2011a; new species), Crous et al. (2013b; new species, DNA), Dubey and Moonambeth (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for type: CBS 335.68.

Cyanoannulus Raja, J. Campb. & Shearer 2003, *Annulatasaceae*, *Annulatascales*, *Sordariomycetes*, one species, type: *C. petersenii* Raja, J. Campb. & Shearer, see Boonyuen et al. (2012; phylogeny), Cai et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: R044-1, ILL 86218, cultures and sequences are available.

Cyanochyta Höhn. 1915, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. cyanogena* (Speg.) Höhn, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cyanodermella O.E. Erikss. 1981, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *C. viridula* (Berk. & M.A. Curtis) O.E. Erikss., coelomycetous, sexual morph unknown, saprobes,

cosmopolitan, see Jaklitsch et al. (2016a; classification), van Nieuwenhuijzen et al. (2016; DNA), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Cyanodiscus E. Müll. & M.L. Farr 1971, *Saccardiaceae*, *Pezizomycotina* families *incertae sedis*, two species, type: *C. occidentalis* E. Müll. & M.L. Farr, asexual morph unknown, saprobes, terrestrial, sub tropical regions, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cyanonectria Samuels & P. Chaverri 2009, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *C. cyanostoma* (Sacc. & Flageolet) Samuels & P. Chaverri, asexual morph hyphomycetous, saprobes, terrestrial, French Guiana, see Samuels et al. (2009; morphology), Schroers et al. (2011; DNA, phylogeny), Lombard et al. (2015b; taxonomy, DNA, phylogeny), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available.

Cyanopatella Petr. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *C. iranica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Iran, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cyanophomella Höhn. 1918, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. acervalis* (Sacc.) Höhn, asexual morph coelomycetous, saprobes, Rhenogovia, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Cyanoporina Groen. 1951, *Pezizomycotina* genera *incertae sedis*, one species, type: *C. granulosa* Groen., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as in *Pyrenotrichaceae*), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences unavailable, needs a revision.

Cyanopulvis J. Fröhl. & K.D. Hyde 2000, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *C. australiensis* J. Fröhl. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as *Xylariales* genera *incertae sedis*), cultures and sequences unavailable, needs a revision.

Cyanopyrenia H. Harada 1995, *Ascomycota* genera *incertae sedis*, one species, type: *C. japonica* H. Harada, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype: HIRO, Harada 3444, cultures and sequences unavailable, needs a revision.

Cyathicula De Not. 1864 (= *Conchatium* Velen. 1934 *fide* Jaklitsch et al. 2016a), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, c. 30 species, type: *C. coronata* (Bull.) De Not., asexual morph *Glarea* (*fide* Jaklitsch et al. 2016a), saprobes on leaves, culms, herbaceous stems and wood, widespread, Kirk et al. (2013; did not list), accepted in Jaklitsch et al. (2016a), type in CUP herbarium, cultures and sequences are unavailable for the type.

Cyberlindnera Minter 2009 (= *Cyberlindnera* Minter 2009; = *Lindnera* Kurtzman, Robnett & Bas.-Powers 2008; = *Zygowilliopsis* Kudryavtsev 1960 *fide* Species Fungorum 2017), *Phaffomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, 30 species, type: *C. americana* (Wickerham) Minter, asexual reproduction is by multilateral budding and formation of pseudohyphae and true hyphae, saprophytic, one species is an opportunistic human and animal pathogen, see Minter (2009; taxonomy), cultures and sequences are available.

Cyclaneusma DiCosmo, Peredo & Minter 1983, *Marthamycetaceae*, *Rhytismatales*, *Leotiomycetes*, two species, type: *C. minus* (Butin) DiCosmo, Peredo & Minter, asexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), cultures and sequences are available.

Cyclodomus Höhn. 1909, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, three species, type: *C. umbellariae* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cyclomarsonina Petr. 1965, *Ascomycota* genera *incertae sedis*, one species, type: *C. cedrelae* (T.S. Ramakr. & K. Ramakr.) Petr., sexual morph unknown, saprobes, terrestrial, India, see coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cyclopeltella Petr. 1953, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *C. orbicularis* Petr., saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2014c; outline, reduced under *Cyclopeltis*), Hyde et al. (2013; accepted as *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), cultures and sequences are unavailable.

Cycloschizon Henn. 1902 (= *Cycloschizella* Höhn. 1919; = *Dielsiella* Henn. 1903; = *Maurodothis* Sacc., Syd. & P. Syd. ex Syd. & P. Syd. 1904 *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, c. 13 species, type: *C. brachylaenae* (Rehm) Henn., asexual morph unknown,

epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), placement in *Parmulariaceae* supported by morphology, cultures and sequences are unavailable.

Cyclostomella Pat. 1896 (= *Cocconiopsis* G. Arnaud 1918; = *Pycnostemma* Syd. 1927 *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, four species, type: *C. disciformis* Pat., epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Cyclothecea Theiss. 1914 nom. cons. prop. (= *Calopeltis* Syd. 1925; = *Hariotula* G. Arnaud 1917; = *Seynesiospora* Bat. 1960; = *Synostomella* Syd. 1927; = *Synpeltis* Syd. & P. Syd. 1917; = *Thyrosoma* Syd. & P. Syd. 1921 *fide* Species Fungorum 2017), *Dothideomycetes*, genera *incertae sedis*, nine species, type: *C. miconiae* (P. Syd.) Theiss., asexual morph unknown, saprobes, worldwide, Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; nomenclature), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Cyclothryrella Jaklitsch & Voglmayr 2016a, *Thyridariaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *C. rubronotata* (Berk. & Broome) Jaklitsch & Voglmayr, asexual morph coelomycetous, saprobes, Europe, possibly also North America, see Jaklitsch and Voglmayr (2016a; taxonomy, phylogeny), holotype of the type: K(M) 202878.

Cyclothrium Petr. 1923 (= *Coniothyriopsiella* Bender 1932; = *Coniothyriopsis* Petr. 1923; = *Kirschsteinia* Petr. 1923 *fide* Species Fungorum 2017), *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *C. ulmigenum* (Berk.) Petr., coelomycetous, ?sexual morph thyridaria-like, saprobes, terrestrial, worldwide, see Damm et al. (2008b; phylogeny), de Gruyter et al. (2012; phylogeny), Wijayawardene et al. (2014c, 2016b; outline, morphology, phylogeny), cultures and sequences are available.

Cylindrium Bonord. 1851, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, c. 50 species, type: *C. elongatum* Bonord., hyphomycetous, sexual morph unknown, on soil, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011 morphology), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; phylogeny, new species), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Cylindrocarpostylus R. Kirschner & Oberw. 1999, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *C. gregarius* (Bres.) R. Kirschner & Oberw., hyphomycetous, sexual morph unknown, in bark beetle

galleries, Europe, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cylindrocladiella Boesew. 1982 (= *Nectricladiella* Crous & C. L. Schoch 2000 *fide* Rossman et al. 2013b), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, eight species, type: *C. parva* (P.J. Anderson) Boesew., soil-borne fungi, pathogenic and/or saprobes of various plant hosts and substrates, worldwide, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2012, 2015b), Maharachchikumbura et al. (2015b, 2016b; outline), ex-epitype: CBS 114524 = ATCC 28272 = CPC 2370; holotype: ATCC 28272.

Cylindrodendrum Bonord. 1851, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *C. album* Bonord., sexual morph unknown, saprobes, pathogenic or isolated from soil, marine, fresh-water and terrestrial, worldwide, Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2012, 2015b), Maharachchikumbura et al. (2015b, 2016b; outline), ex-epitype: CBS 301.83 = ATCC 46842 = IMI 255534; isotype: CBS H-3483A = CBS H-3483B = CBS H-3483C.

Cylindrogloeum Petr. 1941, *Ascomycota* genera *incertae sedis*, one species, type: *C. arcticum* Petr., coelomycetous, sexual morph unknown, saprobes, Lapland, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cylindromyces Manohar., D.K. Agarwal & N.K. Rao 2004, *Ascomycota* genera *incertae sedis*, one species, type: *C. striatus* Manohar., D.K. Agarwal & N.K. Rao, hyphomycetous, sexual morph unknown, saprobes, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cylindrophoma (Berl. & Voglino) Höhn. 1978 (= *Macrophoma* subgen. *Cylindrophoma* Berl. & Voglino 1886 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *C. smilacina* (Desm.) Höhn., coelomycetous, sexual morph unknown, saprobes, Italy, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cylindroseptoria Quaedvl., Verkley & Crous 2013, *Dothideomycetidae*, *Dothideomycetes*, two species, type: *C. ceratoniae* Quaedvl., Verkley & Crous, saprobes, terrestrial, Spain, see Quaedvlieg et al. (2013; morphology, DNA, phylogeny), Wijayawardene et al. (2014c; outline, phylogeny), holotype and ex-type: CBS H-21301, CBS 471.69.

Cylindrosporella Höhn. 1916 (= *Actinonemella* Höhn. 1916; = *Cylindrosporella* Höhn. 1916; = *Depazea* Fr. 1818; = *Gloeosporina* Höhn. 1916; = *Ribaldia* Cif. 1954; = *Titaeosporina* Luijk 1920 [1919] *fide* Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *C. carpini* (Lib.) Höhn., saprobes, terrestrial, Spain, see Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Cylindrosympodioides Crous & M.J. Wingf. 2016, *Venturiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *C. brabejum* Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Crous et al. (2016b; new species, phylogeny), Wijayawardene et al. (2017; outline, phylogeny), holotype and ex-type of the type: CBS H-22594, CPC 25934.

Cylindrosympodium (de Hoog) W.B. Kendr. & R.F. Castañeda 1990, *Ascomycota* genera *incertae sedis*, twelve species, type: *C. variabile* (de Hoog) W.B. Kendr. & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Netherlands, see Seifert et al. (2011; morphology), Castañeda-Ruiz et al. (2012a; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: CBS-H 1634, sequence data, available, compare with *Mirandina*, *Parasypodiella*, *Solosympodiella* and *Subulispora*.

Cylindrothyrium Maire 1907, *Ascomycota* genera *incertae sedis*, one species, type: *C. subericola* Maire, coelomycetous, sexual morph unknown, saprobes, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cylindrotrichum Bonord. 1851 (= *Reticulascus* Réblová & W. Gams 2011 *fide* Maharachchikumbura et al. 2016b; Réblová et al. 2016c), *Reticulascaceae*, *Glomerellales*, *Sordariomycetes*, 14 species, type: *C. oligospermum* (Corda) Bonord., hyphomycetous, sexual morph formerly known in *Reticulascus*, Europe, see Réblová et al. (2011, 2016c; sexual morph, DNA, morphology, phylogeny, nomenclature), Seifert et al. (2011 morphology), Maharachchikumbura et al. (2015b, 2016b; outline, nomenclature), cultures available for type: CBS 561.77, CBS 570.76, CBS 557.74.

Cylindroxypium Bat. & Cif. 1963, *Ascomycota* genera *incertae sedis*, one species, type: *C. virginianum* Bat. & Cif., coelomycetous, sexual morph unknown, saprobes, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cymostachys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *C. fabispora* L. Lombard & Crous,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Lin et al. (2016b; new species), Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type: CBS H-22433, CBS 136180.

Cyniclomyces van der Walt & D.B. Scott 1971 (= *Saccharomycopsis* Guillierm. 1912 *fide* Species Fungorum 2017), *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *C. guttulatus* (C.P. Robin) van der Walt & D.B. Scott, asexual reproduction is by multilateral budding, saprophyte, stomach and intestines of rabbits and certain other herbivores, worldwide, see Lumbsch and Huhndorf (2010; outline), Kurtzman and Robnett (2013b; DNA, phylogeny), Kirk et al. (2013; genus accepted), sequences are available.

Cyphellophora G.A. de Vries 1962, *Cyphellophoraceae*, *Chaetothyriales*, *Eurotiomycetes*, 25 species, type: *C. laciniata* G.A. de Vries, hyphomycetous, sexual morph unknown, saprobes, terrestrial, southern USA, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Réblová et al. (2013; new species), Feng et al. (2014; new species), Gao et al. (2015a, b; new species), Crous et al. (2016b; new species), Madrid et al. (2016; new species), cultures and sequences are available.

Cyphellophoriella Crous & A.J. Sm. 2015, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *C. pruni* Crous & A.J. Sm., hyphomycetous, saprobes, USA, see Crous et al. (2015c; taxonomy, phylogeny), Wijayawardene et al. (2017; outline, phylogeny), holotype and ex-type: CBS H-22255, CPC 25120.

Cyrtidium Vain. 1921, *Ascomycota* genera *incertae sedis*, one species, type: *C. naevium* Vain., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline; accepted as *Dothideomycetes* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c) excluded from *Dothideomycetes*, cultures and sequences are unavailable.

Cyrtidula Minks 1876, *Ascomycota* genera *incertae sedis*, one species, type: need typification, asexual morph unknown, saprobes, Italy, see Lumbsch and Huhndorf (2010; outline; accepted as *Dothideomycetes* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c) excluded from *Dothideomycetes*, cultures and sequences are unavailable.

Cyrtopsis Vain. 1921, *Ascomycota* genera *incertae sedis*, one species, type: *C. fumosa* Vain., asexual morph unknown, saprobes, Italy, see Lumbsch and Huhndorf (2010; outline; accepted as *Dothideomycetes* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c) excluded from *Dothideomycetes*, cultures and sequences are unavailable.

Cystocoleus Thwaites 1849 (= *Chroolepus* C. Agardh 1824 1909 *fide* Species Fungorum 2017), *Cystocoleaceae*, *Capnodiales*, *Dothideomycetes*, one species; type: *C. ebeneus* (Dillwyn) Thwaites, lichenized, lichenicolous, terrestrial, see Muggia et al. (2008; taxonomy), Nelsen et al. (2009; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Hawksworth et al. (2011b; typification), Kirk et al. (2013; genus accepted), Tian et al. (2014; morphology; nomenclature), Lücking et al. (2017a; classification), sequences are available.

Cystodendron Bubák 1914, *Mollisiaceae*, *Helotiales*, *Leotiomyces*, two species, type: *C. dryophilum* (Pass.) Bubák, hyphomycetous, sexual morph unknown, saprobes, Europe, see Grünig et al. (2009; DNA), Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Selçuk et al. (2014; Turkey), cultures and sequences are available for unidentified species.

Cystodium Fée 1837, *Ascomycota* genera *incertae sedis*, one species, type: need typification, asexual morph unknown, saprobes, Italy, see Lumbsch and Huhndorf (2010; outline; accepted as *Dothideomycetes* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c) excluded from *Dothideomycetes*, cultures and sequences are unavailable.

Cystotheca Berk. & M.A. Curtis 1860 (≡ *Sphaerotheca* sect. *Cystotheca*; = *Lanomyces* Gäum. 1922 = *Sphaerotheca* auct. p.p. *fide* Braun and Cook 2012), *Erysiphaceae*, *Erysiphales*, *Leotiomyces*, nine species, type: *C. wrightii* Berk. & M.A. Curtis, sexual and hyphomycetous asexual morphs, biotrophic, obligate plant pathogens, terrestrial, Asia and North America, asexual morph: *Setoidium* (R.T.A. Cook, A.J. Inman & C. Billings) R.T.A. Cook & U. Braun, see Braun and Cook (2012; taxonomy), lectotype of type: K(M) 168934, sequences are available.

Notes: *Setoidium* was originally described as *Oidium* subgen. *Setoidium* R.T.A. Cook, A.J. Inman & C. Billings, typified by *Oidium japonicum* Syd. (Cook et al. 1997), this subgenus was subsequently raised to genus status by Braun and Cook (2012), but no binomial in *Setoidium* was given to the type species. *Oidium japonicum* is the asexual morph of *C. wrightii* (Braun and Cook 2012) and under unitary nomenclature, the genus *Setoidium* should be considered as synonym of *Cystotheca*. In addition, two names in *Setoidium* which lack a *Cystotheca* sexual state are known. The generic affinity of *S. castanopsidis* Meeboon & S. Takam. has been proven by means of molecular sequence data, justifying the reallocation of this species to *Cystotheca*. However, the combination *S. murrayae* (Hosag., U. Braun & Rabindran) U. Braun & R.T.A. Cook was only based on morphological similarity and has not yet been confirmed by sequence data, precluding a reallocation to *Cystotheca* in the current state. The

respective synonymy and a new combination are introduced as follows:

***Cystotheca wrightii* Berk. & M.A. Curtis, Proc. Amer. Acad. Arts 4:130. 1860**

= *Oidium japonicum* Syd., Mém. Herb. Boissier 4:6. 1900.

***Cystotheca castanopsisidis* (Meeboon & S. Takam.) Meeboon & S. Takam., comb. nov.**

IndexFungorum: IF553185

Basionym: *Setoidium castanopsisidis* Meeboon & S. Takam., Mycoscience 54:275. 2012.

Cystotricha Berk. & Broome 1850, *Ascomycota* genera *incertae sedis*, one species, type: *C. striola* Berk. & Broome, coelomycetous, sexual morph unknown, saprobes, terrestrial, British Isles, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cystotrichiopsis Abbas, B. Sutton & Ghaffar 2001, *Ascomycota* genera *incertae sedis*, coelomycetous, sexual morph unknown, one species, type: *C. salvadorae* Abbas, Sutton & Ghaffar, saprobes, terrestrial, Pakistan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: UCMH 759 (IMI 322594), cultures and sequences are unavailable, genus need revision.

Cytodiscula Petr. 1931, *Ascomycota* genera *incertae sedis*, one species, type: *C. carnea* Petr., coelomycetous, sexual morph unknown, saprobes, Madeira, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cytogloeum Petr. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *C. tiliae* Petr., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cytomelanconis Naumov 1951, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *C. systema-solare* Naumov, asexual morph coelomycetous, saprobes, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Cytonaema Höhn. 1914, *Ascomycota* genera *incertae sedis*, two species, type: *C. spinella* (Kalchbr.) Höhn., coelomycetous, sexual morph unknown, saprobes, Austria, South Africa, see Marinowitz et al. (2008; morphology, new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cytoplacosphaeria Petr. 1920, *Ascomycota* genera *incertae sedis*, two species, type: *C. rimos* Petr., coelomycetous, sexual morph unknown, saprobes, Europe, Hong Kong, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cytoplea Bizz. & Sacc. 1885 (= *Neopycnodothis* Tak. Kobay. 1965 *fide* Species Fungorum 2017), *Roussellaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *C. arundinacea* (Sacc.) Petr. & Syd., coelomycetous, sexual morph roussoella-like, saprobes, Europe, Hong Kong, see, Wijayawardene et al. (2012, 2014c, 2016b; outline, morphology, nomenclature), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Cytosphaera Died. 1916, *Ascomycota* genera *incertae sedis*, two species, type: *C. mangiferae* Died., coelomycetous, sexual morph unknown, saprobes, Asia, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Cytospora Ehrenb. 1818 (= *Valsa* Fr. 1825; = *Valsella* Fuckel 1870; = *Leucostoma* (Nitschke) Höhn. 1917; = *Valseutypella* Höhn. 1919; = *Leucocytospora* (Höhn.) Höhn. 1927 *fide* Rossman et al. 2015a; see Species Fungorum for additional synonyms), *Valsaceae*, *Diaporthales*, *Sordariomycetes*, c. 120 species, type: *C. chrysosperma* (Pers.) Fr., coelomycetous, sexual morph formerly known in *Valsa*, saprobes, pathogens, worldwide, see Fotouhifar et al. (2010; phylogeny), Fan et al. (2014, 2015a; morphology, pathogens, China), Liu et al. (2015a; new species), Rossman et al. (2015a; nomenclature), Yang et al. (2015b; pathogens, China), Crous et al. (2016b; new species), Maharanachchikumbura et al. (2015b, 2016b; outline, phylogeny), Norphanphoun et al. (2017, monograph, phylogeny, new species), Tibpromma et al. (2017b; new species), Thambugala et al. (2017a, new species, phylogeny), cultures and sequences are available.

Cytosporella Sacc. 1884, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, 32 species, type: *C. sycina* (Sacc.) Sacc., coelomycetous, sexual morph unknown, saprobes, temperate regions, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Fan et al. (2016a; new species), cultures and sequences are unavailable.

Cytostagonospora Bubák 1916 (= *Cytostaganis* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *C. photiniicola* Bubák, coelomycetous, sexual morph unknown, saprobes, temperate regions, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Quaedvlieg

et al. (2013; DNA, phylogeny), cultures and sequences are available for *C. martiniana* (CBS 135102), type lacking sequence data.

Cyttaria Berk. 1842, *Cyttariaceae*, *Cyttariales*, *Leotiomycetes*, type: *C. darwinii* Berk., asexual morph unknown, saprobes, south temperate regions, see Lumbsch and Huhndorf (2010; outline), Peterson and Pfister (2010; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Cyttariella Palm, *Ascomycota* genera *incertae sedis*, one species, type: *C. skottsbergii* Palm., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, cultures and sequences are unavailable.

Dacampia A. Massal. 1853 (= *Xenosphaeria* Trevis. 1860 *fide* Species Fungorum 2017), *Dacampiaceae*, *Dothideomycetes* families *incertae sedis*, 14 species, type: *D. hookeri* (Borrer) A. Massal., lichenicolous, or saprobes, terrestrial, Europe, see Halici et al. (2008, 2009a, b; new species), Brackel (2010a; new species), Hyde et al. (2013; phylogeny), Wijayawardene et al. (2014a; outline), Ertz et al. (2015a; DNA, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, available cultures for type: Hafellner 81840 (GZU), Hafellner 75980 (GZU), Hafellner 74269 (GZU), Hafellner 73897 (GZU).

Dacryoma Samuels 1988, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type *D. album* Samuels, sexual morph unknown, saprobes on palms, terrestrial, Indonesia Peru, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: BO, isotype NY.

Dactylaria Sacc. 1880 (= *Didymotrichum* Höhn. 1914; = *Diplorhinotrichum* Höhn. 1902 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, c. 100 species, type: *D. purpurella*, hyphomycetous, saprobes, pathogens, parasites, terrestrial and aquatic, widespread, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Magalhães et al. (2014), Mel'nik et al. (2013a; new species), Crous et al. (2016a; new species, phylogeny), few sequence available for this genus but not available for the type, sexual morph hypomyces-like reported for *D. mycophila*.

Dactyella Grove 1884 (= *Gangliophragma* Subram. (1978, "1977"); = *Drechsleromyces* Subram. (1978, "1977") *fide* Baral et al. 2015), *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, heterogenous, type: *D. minuta* Grove, c. 70 species, hyphomycetous, ?sexual morph *Orbilia*, saprobes, terrestrial, worldwide, see Zhang et al. (2010d; new species), Qin et al. (2011; new species), Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted),

Baral et al. (2015; genus accepted), cultures and sequences are available.

Dactyellina M. Morelet 1968 (see Baral et al. 2015 for synonyms), *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, c. 23 species, type: *D. leptospora* (Drechsler) M. Morelet, hyphomycetous, sexual morph unknown, nematode trapping, in soil, terrestrial, aquatic, Asia, Europe, see Seifert et al. (2011 morphology), Hanying et al. (2012; development), Wijayawardene et al. (2012, 2017; outline), Yu et al. (2012; biology), Kirk et al. (2013; genus accepted), Baral et al. (2015; genus accepted), cultures and sequences are available.

Dactylifera Alcorn 1927, *Ascomycota* genera *incertae sedis*, one species, type: *D. australiensis* Alcorn, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dactylina Nyl. 1860 (= *Dufoureomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *D. arctica* (Richardson) Nyl., asexual reproduction by small lateral lobules and sexual by apothecia (rarely developed), lichenized, terricolous, circumpolar, arctic-alpine Northern Hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences available for the type: AFTOL 225.

Dactylonectria L. Lombard & Crous 2014, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, ten species, type: *D. macrodidyma* (Halleen, Schroers & Crous) L. Lombard, van der Merwe & J.Z. Groenew. & Crous, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lombard et al. (2014, 2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Dactylospora Körb. 1855 (= *Kymadiscus* Kohlm. & E. Kohlm. 1971; = *Mycolecidia* P. Karst. 1888; = *Mycolecis* Clem. 1909; = *Paruephaedria* Zukal 1891; = *Pseudokarschia* Velen. 1934 *fide* Species Fungorum 2017), *Dactylosporaceae*, *Eurotiomycetes* families *incertae sedis*, c. 50 species, type: *D. floerkei* Körb., asexual morph unknown, on lichens, hepatics or saprobes, terrestrial or marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Diederich et al. (2013; phylogeny), Pang et al. (2014), Jones et al. (2015), Pino-Bodas et al. (2017; new species, phylogeny), sequence data available.

Dactylosporium Harz 1871, *Ascomycota* genera *incertae sedis*, two species, type: *D. macropus* (Corda) Harz, hyphomycetous, sexual morph unknown, saprobes,

terrestrial, Europe, Cuba, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Daldinia Ces. & De Not. 1863 (= *Annellosporium* M.L. Davey 2010; = *Hemisphaeria* Klotzsch 1843; = *Peripherostoma* Gray 1821; = *Stromatosphaeria* Grev. 1824; = *Versiomycetes* Whalley & Watling 1989 *fide* Réblová et al. 2016c; Wendt et al. 2017; see Species Fungorum 2017 for additional synonyms), *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, c. 40 species, type: *D. concentrica* (Bolton) Ces. & De Not., asexual morph hyphomycetous, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), Sir et al. (2016b; new species), Kuhnert et al. (2017; new species, revision), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available.

Dangeardiella Sacc. & P. Syd. 1899, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *D. macrospora* (J. Schröt.) Sacc. & P. Syd., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014e; transferred to *Pleosporales*, genera *incertae sedis*), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Danielozyma Kurtzman & Robnett 2014, *Metschnikowiaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *D. ontarioensis* (Kurtzman & Robnett) Kurtzman & Robnett, asexual reproduction by multilateral budding and from formation of pseudohyphae and true hyphae, saprophyte, in insect frass from various tree species, see Kurtzman and Robnett (2014; taxonomy), cultures and sequences are available.

Darkera H.S. Whitney, J. Reid & Piroz. 1975, *Phaciidaeae*, *Phaciidales*, *Leotiomycetes*, five species, type: *D. parca* H.S. Whitney, J. Reid & Piroz., asexual morph coelomycetous, pathogens, terrestrial, Europe, Cuba, see Kirk et al. (2013; genus accepted), Crous et al. (2015b; revision, morphology, DNA, phylogeny), cultures and sequences are available.

Darksidea D.G. Knapp, Kovács, J.Z. Groenew. & Crous 2015, *Lentitheciaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. alpha* D.G. Knapp, Kovács, Groenewald & Crous, endophytes, see Knapp et al. (2015; phylogeny), cultures and sequences are available.

Dasyscyphella Tranzschel 1898 (= *Lachnaster* Höhn. 1917 *fide* Jaklitsch et al. 2016a), *Lachnaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *D. cassandrae* Tranzschel, asexual morph unknown, saprobes, terrestrial, Asia, North America, see Lumbsch and Huhndorf (2010;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Dasysphaeria Speg. 1912, *Sordariomycetes* genera *incertae sedis*, one species, type: *D. andicola* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015a; outline), cultures and sequences are unavailable.

Dasydictia Speg. 1912, *Ascomycota* genera *incertae sedis*, two species, type: *D. sapindophila* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Davidgallowaya Aptroot 2007, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetidae*, one species, type: *D. cornutispora* Aptroot, asexual morph, unknown, terrestrial, New Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Davidiellomyces Crous 2017, *Cladosporiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *D. australiensis* Crous, asexual morph unknown, saprobes, Australia, see Crous et al. (2017; taxonomy), holotype: CBS H-23077, cultures and sequences are available.

Davidsoniella Z.W. de Beer, T.A. Duong & M.J. Wingf. 2014, *Ceratocystidaceae*, *Microascales*, *Sordariomycetes*, one species, type: *D. virescens* (R.W. Davidson) Z.W. de Beer, T.A. Duong & M.J. Wingf., asexual morph hyphomycetous, endophyte, terrestrial, Thailand, have sequence data, see de Beer et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Davisiella Petr. 1924, *Ascomycota* genera *incertae sedis*, two species, type: *D. elymina* (Davis) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Davisoniella Darker 1967, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, ten species, type: *D. ampla* (Davis) Darker, asexual morph unknown, saprobes, endophytes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequence available for *D. medusa* (BPI842078).

Davisoniella H.J. Swart 1988, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *D. eucalypti* H.J. Swart, coelomycetous, sexual morph teratosphaeria-like, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2014c, 2017), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Dawsicola Döbbeler 1981, *Helotiales* genera *incertae sedis*, one species, type: *D. neglecta* Döbbeler, asexual morph unknown, saprobes, endophytes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dawsomyces Döbbeler 1981, *Dothideomycetes* genera *incertae sedis*, two species, type: *D. mirabilis* Döbbeler, asexual morph unknown, saprobes, endophytes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Dawsophila Döbbeler 1981, *Dothideomycetes* genera *incertae sedis*, two species, type: *D. pygmaea* Döbbeler, asexual morph unknown, saprobes, endophytes, terrestrial, New Guinea, New Zealand, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Deakozyma Kurtzman & Robnett 2014, *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *D. indianensis* Kurtzman & Robnett, asexual reproduction by multilateral budding and from formation of pseudohyphae and true hyphae, saprophyte, rotting wood, insect eggs, Indiana, Michigan, USA, see Kurtzman and Robnett (2014; taxonomy), cultures and sequences are available.

Dearnessia Bubák 1916, *Ascomycota* genera *incertae sedis*, one species, type: *D. apocyni* Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Canada, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Debaryomyces Lodder & Kreger-van Rij 1952, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, 15 species, type: *D. hansenii* (Zopf) Lodder & Kreger-van Rij, asexual reproduction by multilateral budding and some species may also form poorly developed pseudohyphae, saprophytic, but several of the species are opportunistic pathogens of humans, soil, sea and fresh water, high osmotic foods such as cheese and sausages, bees, tree exudates, spoiled sake, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Decaisnella Fabre 1878, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, 13 species, type: *D. spectabilis* Fabre, asexual morph unknown, terrestrial and marine, worldwide, see Suetrong et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes* genera *incertae sedis*),

Jones et al. (2016; outline), sequences are available for only one species, need recollection to verify generic placement.

Decorospora Inderb., Kohlm. & Volk.-Kohlm. 2002, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. gaudefroyi* (Pat.) Inderb., Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, marine, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pleosporaceae*), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015c; phylogeny), Jones et al. (2015; outline), cultures and sequences are available.

Degelia Arv. & D.J. Galloway 1981, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 22 species, type: *D. gayana* (Mont.) Arv. & D.J. Galloway, asexual morph unknown, lichenized, epiphytic or on mossy rocks, Southern hemisphere, in humid regions, see Wedin et al. (2009b; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Spribille and Muggia (2013; phylogeny), Ekman et al. (2014; phylogeny, *Degelia* is heterogeneous and excluded the section *Amphiloma* as the genus *Pectenia*), Lücking et al. (2017a; classification), need generic revision.

Degeliella P. M. Jørg 2004, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *D. rosulata* (P.M. Jørg. & D.J. Galloway) P.M. Jørg., asexual morph unknown, lichenized, epiphytic or on mossy rocks, Gondwana distribution, Australia (Tasmania), southernmost South America, see Lumbsch and Huhndorf (2010; outline), Ekman et al. (2014; phylogeny, non-monophyly of the genus), Jorgensen and Andersen (2015; showed that the name *Degeliella* was superfluous and that *Psoromaria* (type: *P. versicolor*) should be taken up for the genus, which is still likely to be non-monophyletic), Lücking et al. (2017b; classification), sequence available for voucher Galloway 840b (BG) of the type.

Deichmannia Alstrup & D. Hawksw. 1990, *Ascomycota* genera *incertae sedis*, one species, type: *M. caespitosa* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Deightoniella S. Hughes 1952 (= *Utrechtiana* Crous & Quaedvli. 2011 *fide* Species Fungorum 2017), *Pyriculariaceae*, *Magnaportheales*, *Sordariomycetes*, 20 species, type: *D. africana* S. Hughes, hyphomycetous, sexual morph unknown, pathogens, terrestrial, worldwide, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Klaubauf et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Zhang et al. (2016d; nomenclature), cultures and sequences are available for *Deightoniella roumeguerei* (CBS 128780).

Dekkera Van der Walt 1964, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, five species, type: *D. bruxellensis* Van der Walt, saprophytic, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Delastria Tul. & C. Tul. 1843, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *D. rosea* Tul. & C. Tul., asexual morph unknown, hypogeous, Europe (southern), Northern America, see Lumbsch and Huhndorf (2010; outline), Alvarado et al. (2011a; DNA, phylogeny), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Delitschia Auersw. 1866 (= *Delitschiella* Sacc. & D. Sacc. 1905; = *Pachyspora* Kirschst. 1907 [1906] *fide* Species Fungorum 2017), *Delitschiaceae*, *Pleosporales*, *Dothideomycetes*, 52 species, type: *D. didyma* Auersw., asexual morph unknown, commonly occurs as coprophilous and saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; morphology and phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Calaça et al. (2015; new variety), holotype of the type: L 1950, cultures available for type: UME 31411.

Delortia Pat. & Gaillard 1888 (= *Hyalotrochophora* Finley & E.F. Morris 1967 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *D. palmicola* Pat. & Gaillard, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Pantropical, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Delphinella (Sacc.) Kuntze 1898 (= *Diplosphaerella* Grove 1912; = *Glonium* subgen. *Delphinella* Sacc 1891.; = *Hariotia* P. Karst. 1889; = *Pleoglonis* Clem. 1909; = *Rehmielopsis* Bubák & Kabát 1910 *fide* Species Fungorum 2017), *Dothideaceae*, *Dothideales*, *Dothideomycetes*, eight species, type: *D. strobiligena* (Desm.) Sacc. ex E. Müll. & Arx, asexual morph unknown, saprobes or pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothioraceae*), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; accepted as a genus in *Dothideaceae*), Wijayawardene et al. (2014c; outline), holotype: PC 0084688.

Delpinoella Sacc. 1899, *Sordariomycetes* genera *incertae sedis*, one species, type: *D. insignis* Sacc. & Trotter, asexual morph unknown, saprobes, terrestrial, Canada, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Delpinoina Kuntze 1891 (= *Henriquesia* Pass. & Thüm. 1879 *fide* Species Fungorum 2017), *Ascodichaenaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *D. lusitanica* (Pass. & Thüm.) Kuntze, asexual morph unknown,

saprobes, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Delpontia Penz. & Sacc. 1901, ?*Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *D. pulchella* Penz. & Sacc., asexual morph unknown, saprobes, on fern stalks, Java, see Lumbsch and Huhndorf (2010; outline: doubtful position), cultures and sequences are unavailable, classification is not evaluated phylogenetically.

Deltopyxis Baral & G. Marson 2012, *Leotiomycetes* genera *incertae sedis*, one species, type: *D. triangulispora* Baral & G. Marson, saprobes, terrestrial, Luxembourg, see Baral and Marson (2012; taxonomy), Jaklitsch et al. (2016a; classification), a sequence is available.

Dematiocladium Allegr., Aramb., Cazau & Crous 2005, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *D. celtidis* Allegr., Aramb., Cazau & Crous, saprobes, terrestrial, Argentina, China, see Wijayawardene et al. (2012, 2017; outline), Crous et al. (2014c; new species), ex-type: CBS 115994; holotype of the type: LPS 47255, isotype CBS 9921.

Dematiopleospora Wanás., Camporesi, E.B.G. Jones & K.D. Hyde, gen. nov.

Index Fungorum number: IF552175

Description: see Wanasinghe et al. (2014b)

= *Dematiopleospora* Wanás., Camporesi, E.B.G. Jones & K.D. Hyde, Cryptog Mycol 35(2): 110 (2014b), Nom. inval., Art. 42.1 (Melbourne)

Type: *Dematiopleospora mariae* Wanás., Camporesi, E.B.G. Jones & K.D. Hyde sp. nov.

Notes: Wanasinghe et al. (2014b) introduced *Dematiopleospora* but it is invalid (Art. 42.1) as the genus and type species both have same identifier IF550536. Three other species are consequently not validly published. Thus, new identifiers have been obtained and species are validly published.

Dematiopleospora alliariae Thambug., Camporesi & K.D. Hyde sp. nov.

Index Fungorum: IF552256

Description: see Hyde et al. (2016) = *Dematiopleospora alliariae* Thambug., Camporesi & K.D. Hyde, Fungal Diversity 80: 89 (2015); nom. inval., Art. 35.1

Dematiopleospora cirsii Wanás., Camporesi, E.B.G. Jones & K.D. Hyde sp. nov.

Index Fungorum number: IF552282

Description: see Hyde et al. (2016)

= *Dematiopleospora cirsii* Wanás., Camporesi, E.B.G. Jones & K.D. Hyde, Fungal Diversity 80: 89 (2015); nom. inval., Art. 35.1

Dematiopleospora luzulae Wanás., Camporesi, E.B.G. Jones & K.D. Hyde sp. nov.

Index Fungorum number: IF552283

Description: see Hyde et al. (2016)

= *Dematiopleospora luzulae* Wanas., Camporesi, E.B.G. Jones & K.D. Hyde, Fungal Diversity 75: 103 (2015); nom. inval., Art. 35.1

Dematiopleospora mariae Wanas., Camporesi, E.B.G. Jones & K.D. Hyde sp. nov.

Index Fungorum number: IF552185

Description: see Wanasinghe et al. (2014b)

= *Dematiopleospora mariae* Wanas., Camporesi, E.B.G. Jones & K.D. Hyde, Cryptog Mycol 35 (2): 110 (2014); nom. inval., Art. 35.1

Dematioscypha Svrček 1977 (= *Schizocephalum* Preuss 1852; *Haplographium* Berk. & Broome 1859 *fide* Johnston et al. 2014b), *Hyaloscypphaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *D. dematiicola* (Berk. & Broome) Svrček, asexual morph hyphomycetous, saprobes, Europe, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), Johnston et al. (2014b; nomenclature), Wijayawardene et al. (2017; outline), type in K herbarium, cultures and sequence data available for *Haplographium delicatum*.

Dencoceliopsis Korf 1971, *Rutstroemiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *D. johnstonii* (Berk.) Korf, asexual morph unknown, saprobes, terrestrial, USA, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dendriscosticta Moncada & Lücking 2013, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, five species, type: *D. wrightii* (Tuck.) Moncada & Lücking, lichenized, see Moncada et al. (2013a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Dendroclathra Voglmayr & G. Delgado 2001, *Hypocreomycetidae* genera incertae sedis, *Sordariomycetes*, one species, type: *D. caeruleofusca* Voglmayr & G. Delgado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Dendrodromus Bubák 1915, *Ascomycota* genera incertae sedis, one species, type: *D. annulatus* Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dendrographa Darb. 1895, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, six species, type: *D. leucophaea* (Tuck.) Darb., lichenized, corticolous and saxicolous, worldwide, see Lumbsch and Huhndorf (2010; outline), Ertz and Tehler (2011; phylogeny and taxonomy), Kirk et al. (2013;

genus accepted), Lücking et al. (2017a; classification), sequences are available.

Dendrophoma Sacc. 1880 (= *Amphitiarospora* Agnihothr. 1963 [1962]; = *Dendrophoma* Sacc. 1880; = *Pycnidiochaeta* Sousa da Câmara 1950 *fide* Species Fungorum 2017), *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, c. 100 species, type: *D. cytisporoides* Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2008; treated as a synonym of *Dinemasporium*), Crous et al. (2012g; reinstated the genus, DNA, phylogeny), Liu et al. (2015a; new species), Maharachchikumbura et al. (2016b; outline), epitype and ex-epitype: CBS H-12004, CBS 223.95.

Dendrosphaera Pat. 1907, *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, one species, type: *D. eberhardtii* Pat., asexual morph hyphomycetous, terrestrial, probably mycorrhizal, Asia, see Houbraken and Samson (2011; morphotaxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, needs phylogenetic assessment.

Dendrospora Ingold 1943, *Ascomycota* genera incertae sedis, ten species, type: *D. erecta* Ingold, hyphomycetous, sexual morph unknown, saprobes, terrestrial, north temperate, subtropical, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Duarte et al. (2015; DNA), cultures and sequences are available.

Dendrosporium Plakidas & Edgerton ex J.L. Crane 1972, *Ascomycota* genera incertae sedis, three species, type: *D. lobatum* Plakidas & Edgerton ex J.L. Crane, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, North and South America, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Dendryphiella Bubák & Ranoj. 1914, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, nine species, type: *D. interseminata* (Berk. & Ravenel) Bubák, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011 morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2014c, 2016b; new species, phylogeny), cultures and sequences are available.

Dendryphion Wallr. 1833 (= *Brachycladum* Corda 1838; = *Dendryphiella* Bubák & Ranoj. 1914; = *Dwayamala* Subram. 1956; = *Entomyclium* Wallr. 1833 *fide* Species Fungorum 2017), *Torulaceae*, *Pleosporales*, *Dothideomycetes*, 66 species, type: *D. comosum* Wallr., hyphomycetous, sexual morph unknown, saprobes, worldwide, see Kirk et al. (2013; genus accepted), Crous et al. (2015a; phylogeny), Su et al. (2016b; new species,

phylogeny), sequences are available for five species, needed generic placement.

Dendryphiosphaera Lunghini & Rambelli 1978, *Ascomycota* genera *incertae sedis*, four species, type: *D. taiensis* Lunghini & Rambelli, hyphomycetous, sexual morph unknown, Ivory coast, pantropical, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Deniquelata Ariyaw. & K.D. Hyde 2013, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. barringtoniae* Ariyawansa & K.D. Hyde, asexual morph unknown, pathogens, terrestrial, Thailand, see Ariyawansa et al. (2013c, 2014c; morphology, DNA, phylogeny), Hyde et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline, phylogeny), holotype and ex-type of the type: MFLU 12-0303, MFLUCC 12-0257.

Dennisiella Bat. & Cif. 1962 (= *Capnodium* subgen. *Microxiphium* Harv. ex Berk. & Desm. 1849; = *Microxiphium* (Harv. ex Berk. & Desm.) Thüm. 1879; = *Vitalia* Cif. & Bat. 1962 *fide Species Fungorum* 2017), *Coccodiniaceae*, *Chaetothyriales*, *Eurotiomycetes*, seven species, type: *D. babingtonii* (Berk.) Bat. & Cif., asexual morph microxyphium-like, saprobes, terrestrial, tropical regions, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as genera in *Coccodiniaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dennisioidiscus Svrcek 1976 (= *Trichodiscus* Kirschst. 1924 *fide Species Fungorum* 2017), *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, ten species, type: *D. prasinus* (Quél.) Svrček, asexual morph unknown, saprobes, terrestrial, Europe, Papua New Guinea, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Dennisiopsis Subram. & Chandras. 1977, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, two species, type: *D. octospora* Subram. & Chandras., asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dennisographium Rifai 1977, *Ascomycota* genera *incertae sedis*, two species, type: *D. episphaeriae* Rifai, hyphomycetous, sexual morph unknown, Java, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Denticularia Deighton 1972, *Ascomycota* genera *incertae sedis*, seven species, type: *D. modesta* (Syd.) Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dentocircinomyces R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, one species, type: *D. occidentalis* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype: INFAT C88/114, cultures and sequences are unavailable, compare with *Cryptumbellata*, *Hansfordia* and *Redbia*.

Derezia Naras. 1970, *Ascomycota* genera *incertae sedis*, one species, type: *D. lecanii* (Zimm.) Naras., hyphomycetous, sexual morph unknown, on *Lecanium viride*, India, Indonesia, Java, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Dermapteromyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *D. ctenophorus* Thaxt., asexual morph unknown, biotrophic, America (tropical), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dermateopsis Nannf. 1932, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *D. tabacina* (Cooke) Nannf., asexual morph unknown, saprobes, terrestrial, temperate regions, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Dermatiscum Nyl. 1867, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, three species, type: *D. thunbergii* (Ach.) Nyl., asexual morph is coelomycetous, lichenized, saxicolous, South Africa, Madagascar, St. Helena, see Aptroot (2008; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available for the type: AF540507 (ITS).

Dermatocarpon Eschw. 1824 (= *Capnia* Vent. 1799; = *Endocarpomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Endocarpon* subdiv. *Entosthelia* Wallr. 1831; = *Entosthelia* (Wallr.) Hue 1915; = *Epistictum* Trevis. 1869 *fide Species Fungorum* 2017), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, c. 20 species, type: *D. miniatum* (L.) W. Mann, asexual morph coelomycetous, terrestrial, lichenized, on rocks and soil, cosmopolitan, see Gueidan et al. (2009; phylogeny, outline), Prieto et al. (2012; phylogeny), Lücking et al. (2017a; classification), sequences of the type material of the type are unavailable, but sequences of non-type material are available.

Dermatodothella Viégas 1944, *Polystomellaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *D. multiseptata* Viégas, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, as *Dothideomycetes*, genera *incertae sedis*), Ariyawansa et al. (2013b; morphology, accepted as *Polystomellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Dermatodothis Racib. ex Theiss. & Syd. 1914 (= *Phaeophragmocaula* F. Stevens 1931 *fide Species Fungorum* 2017), *Dothideomycetes* genera *incertae sedis*, six species, type: *D. javanica* Racib., asexual morph unknown, saprobes, terrestrial, Asia, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Dermea Fr. 1825 (= *Bulgariastrum* Syd. & P. Syd. 1913; = *Cenangella* Sacc. 1884; *Dermatea* Fr. 1849 = *Foveostroma* DiCosmo 1978; = *Sphaeronaema* Fr. 1815 *fide* Johnston et al. 2014b; *Species Fungorum* 2017), *Dermateaceae*, *Helotiales*, *Leotiomycetes*, c. 25 species, type: *D. cerasi* (Pers.) Fr., asexual morphs formerly known in *Sphaeronaema* and *Foveostroma*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Johnston et al. (2014b; nomenclature, new comb.), cultures and sequences are available.

Dermiscillum Hafellner, Mayrhofer, & Poelt 1979, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *D. catawicense* (Willey) Hafellner & Poelt, asexual morph is coelomycetous, lichenized, saxicolous, North America, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; taxonomy), cultures and sequences are unavailable.

Descalsia A. Roldán & Honrubia 1989, *Ascomycota* genera *incertae sedis*, one species, type: *D. cruciata* A. Roldán & Honrubia, hyphomycetous, sexual morph unknown, aquatic, isolated from Spanish streams, South western Europe, Spain, Northwest Portugal, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: IMI 328087, sequences are unavailable.

Desertella Mouch. 1979, *Ascomycota* genera *incertae sedis*, two species, type: *D. globulifera* Mouch., hyphomycetous, sexual morph unknown, from soil, submerged wood in river, Egypt, USA, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures are unavailable, unverified sequences are available (KR230020-KR230041).

Deshpandiella Kamat & Ullasa 1973, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *D. jambolana* (T.S. Ramakr., Sriniv. & Sundaram) Kamat & Ullasa, asexual morph unknown, biotrophic, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Desmazierella Lib. 1829 (= *Verticiladium* Preuss 1851 *fide* Healy et al. 2016), *Chorioactidaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *D. acicola* Lib., asexual morph formerly known in *Verticiladium*, saprobes, terrestrial, India, see Pfister et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Martinović et al. (2016; DNA, phylogeographic structure), cultures and sequences are available.

Desmidiospora Thaxt. 1891, *Ascomycota* genera *incertae sedis*, three species (two fossil), type: *D. myrmecophila* Thaxt., hyphomycetous, sexual morph unknown, terrestrial, pathogen of *Camponotus* ants, North America, Africa, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Devriesia Seifert & N.L. Nick. 2004, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, 24 species, type: *D. staurophora* (W.B. Kendr.) Seifert & N.L. Nick., asexual morph hyphomycetous, saprobes, terrestrial, India, see Schoch et al. (2009; new species, phylogeny), Crous et al. (2011a, 2012c, e; new species, DNA), Frank et al. (2010; new species, DNA, phylogeny), Li et al. (2012a; new species, China), Egidi et al. (2014; new species, DNA, phylogeny), cultures and sequences are available.

Dexhowardia J.J. Taylor 1970, *Ascomycota* genera *incertae sedis*, one species, type: *D. tetraspora* J.J. Taylor, hyphomycetous, sexual morph unknown, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available.

Diaboliumbilicus I. Hino & Katum. 1955, *Ascomycota* genera *incertae sedis*, one species, type: *D. mirabilis* I. Hino & Katum., asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diachora Jul. Müll. 1893 (= *Chaetomelasmia* Danilova 1951; = *Diachorella* Höhn. 1918 *fide* Maharanachikumbura et al. 2016b; Réblová et al. 2016c), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, six species, type: *D. onobrychidis* (DC.) Jul. Müll., asexual morph *Diachorella*, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus

accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Diacrochordon Petr. 1955, *Sordariomycetes* genera *incertae sedis*, six species, type: *D. rehmii* Petr., asexual morph unknown, saprobes, terrestrial, Germany, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Diadema Shoemaker & C.E. Babc. 1989, *Diademaceae*, *Pleosporales*, *Dothideomycetes*, nine species, type: *D. tetramerum* Shoemaker & C.E. Babc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011b; taxonomy), Hyde et al. (2013; taxonomy), Ariyawansa et al. (2014d; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Diademosia Shoemaker & C.E. Babc. 1992, *Diademaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. californiana* (M.E. Barr) Shoemaker & C.E. Babc., asexual morph unknown, saprobes on herbaceous, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2015c; generic placement), Wijayawardene et al. (2014c; outline), sequences unavailable, needed generic placement.

Diademospora B.E. Soderstr. & Bååth 1979, *Ascomycota* genera *incertae sedis*, one species, type: *D. ramigera* B.E. Söderstr. & Bååth, hyphomycetous, sexual morph unknown, from forest soil, coniferous, Sweden, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), ex-type: CBS 427.76, IMI 203569, ATCC 42210.

Dialaciopsis Bat. 1959 (= *Dialaciopsis* Bat. 1959; = *Dialacium* Syd. 1930; = *Englerulella* Hansf. 1946 [1944-45] fide Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *D. landolphiae* Bat., coelomycetous, sexual morph unknown, from *Landolphia ugandensis*, Uganda, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Dialhypocrea Speg. 1919, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *D. puiggariana* Speg., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2014b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Diamantinia A.N. Mill., Læssøe & Huhndorf 2003, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *D. citrina* A.N. Mill., Læssøe & Huhndorf,

aseexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Xylariales*), holotype and ex-type: SP, Buck 26886.

Diandromyces Thaxt. 1918, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *D. chilensis* Thaxt., asexual morph unknown, biotrophic, Chile, Ecuador, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species, Ecuador), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dianesea (F. Stevens) Inácio & P.F. Cannon 2002, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. palmae* (F. Stevens) Inácio & P.F. Cannon, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), Jayasiri et al. (2016a; re-described type), sequences are unavailable.

Diaphorographis A.W. Archer & Kalb 2009, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *D. queenslandica* Kalb & A.W. Archer, asexual morph unknown, lichenized, terrestrial, tropical, see Kalb et al. (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; taxonomy), Kirk et al. (2013; genus accepted), sequences are unavailable.

Diaphoromyces Thaxt. 1926, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *D. marginatus* (Thaxt.) Thaxt., asexual morph unknown, biotrophic, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diaporthe Nitschke 1870 (= *Allantoporthe* Petr. 1921; = *Aporhytisma* Höhn. 1917; = *Chorostate* (Nitschke ex Sacc.) Traverso 1906; = *Clypeorhynchus* Kirschst. 1936; = *Dalytes* Nitschke 1867; = *Diaporthe* subgen. *Chorostate* Nitschke ex Sacc. 1882; = *Melanoporthe* Wehm. 1938; = *Phomopsis* (Sacc.) Bubák 1905; = *Septomazzantia* Theiss. & Syd. 1915 fide Udayanga et al. 2011, 2012; Rossman et al. 2015a; Maharachchikumbura et al. 2016b), *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, more than 125 species, type: *D. eres* Nitschke, asexual morph coelomycetous, endophytes, pathogens, saprobes, associate with wide host range, worldwide, see Udayanga et al. (2011, 2012, 2014a, b, 2015; review, nomenclature, phylogeny, species complexes), Gomes et al. (2013; taxonomy), Huang et al. (2013; species occur in *Citrus* sp.) Kirk et al. (2013; genus accepted), Dissanayake et al. (2015; new species, phylogeny), Fan et al. (2015b; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2015a; nomenclature), Crous et al. (2016a; taxonomy, phylogeny), Du et al. (2016; *Diaporthe* spp. in *Betula* spp.), Doilom et al. (2017, new species, phylogeny), Santos et al. (2017; new species),

Thambugala et al. (2017a, new species), Yang et al. (2017; new species), holotype and culture ex-epitype: BPI 892912, AR5193.

Diaporthella Petr. 1924, *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *D. aristata* (Fr.) Petr., asexual morph unknown, saprobes, terrestrial, Europe, North America, Eastern Asia, see Lumbsch and Huhndorf (2010; outline), Gomes et al. (2013; DNA), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Diarimella B. Sutton 1980, *Ascomycota* genera *incertae sedis*, three species, type: *D. setulosa* B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Argentina, Canada, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: IMI 103785, cultures and sequences are unavailable, genus need revision.

Diatractium Syd. & P. Syd. 1921, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *D. cordiae* (F. Stevens) Syd. & P. Syd., asexual morph unknown, pathogens, terrestrial, Central and South America, see Palmateer et al. (2008; DNA, pathogens), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available.

Diatrypasimilis J.J. Zhou & Kohlm. 2016 (= *Diatrypasimilis* J.J. Zhou & Kohlm. 2010 *fide* Species Fungorum 2017), *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *D. australiensis* J.J. Zhou & Kohlm., asexual morph unknown, saprobes, on mangroves, Central and South America, see Palmateer et al. (2008; DNA, pathogens), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline; taxonomy), Index Fungorum (2017; validate the genus), cultures and sequences are available.

Diatrype Fr. 1849 (= *Ectosphaeria* Speg. 1921; = *Libertella* Desm. 1830; = *Phaeotrype* Sacc. 1920; = *Pyrenodochium* Bonord. 1851' = *Stictosphaeria* Tul. & C. Tul. 1863; = *Valsa* Adans. 1763 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, cf. 60 species, type: *D. disciformis* (Hoffm.) Fr., asexual morph formerly known in *Libertella*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Liu et al. (2015a; new species), Maharachchikumbura et al. (2015b, 2016b; outline; taxonomy), Senanayake et al. (2015; phylogeny, outline), Réblová et al. (2016c; nomenclature), cultures and sequences are available.

Diatrypella (Ces. & De Not.) De Not. 1863 (= *Diatrype* sect. *Diatrypella* Ces. & De Not. 1863; = *Sphaerosperma* Preuss 1853 *fide* Species Fungorum 2017), *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, c. 35 species, type:

D. verruciformis (Ehrh.) Nitschke, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline; taxonomy), Mehrabi et al. (2015; new species), Senanayake et al. (2015; phylogeny, outline), Almeida et al. (2016; new species, phylogeny), Crous et al. (2016a; new species, phylogeny), Shang et al. (2017, new species, phylogeny), cultures and sequences are available.

Diatrypoidiella Manohar., Kunwar & D.K. Agarwal 2005, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *D. lignicola* Manohar., Kunwar & D.K. Agarwal, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline; taxonomy), holotype of the type: HCIO 43762, cultures and sequences are unavailable.

Dibaeis Clem. 1909 (= *Tubercularia* Weber ex F.H. Wigg. 1780; = *Verrucaria* Scop. 1777 *fide* Species Fungorum 2017), *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, 13 species, type: *D. rosea* (Pers.) Clem., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2015a; phylogeny), Lücking et al. (2017a; classification), culture available for the type: AFTOL-ID 358.

Dicephalospora Spooner 1987, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *D. calochroa* (Syd.) Spooner, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Liu et al. (2016b; new species), cultures and sequences are available.

Dichelostroma Bat. & Peres 1963, *Ascomycota* genera *incertae sedis*, one species, type: *D. powellii* Bat. & Peres, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), data inadequate, cultures and sequences are unavailable.

Dichlaena Durieu & Mont. 1849, *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, four species, type: *D. lentisci* Durieu & Mont., asexual morph hyphomycetous, terrestrial, saprobes, occurring on decaying plant material, wood and dung, Algeria, Brazil, India, Italy, Mauritania, USA, see Houbraken and Samson (2011), cultures and sequences are unavailable, needs phylogenetic assessment.

Dichlobodigitus G.P. White & Illman 1988, *Ascomycota* genera *incertae sedis*, one species, type: *D. niger* G.P. White & Illman, hyphomycetous, sexual morph unknown, on rotten wood of *Pinus contorta*, Canada, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dichosporidium Pat. 1903 (= *Dichosporium* Pat. 1899 *fide Species Fungorum* 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, eight species, type: *D. glomeratum* (Pat.) Pat., asexual morph unknown, lichenized, widespread (tropical), corticolous (occasionally saxicolous), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ertz et al. (2015c; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Dichotomophthora Mehrl. & Fitzp. ex P.N. Rao 1966, *Ascomycota* genera *incertae sedis*, four species, type: *D. portulacae* Mehrl. & Fitzp. ex P.N. Rao, hyphomycetous, sexual morph unknown, on living leaves and stems of *Portulaca oleracea*, Hawaii, USA, widespread (esp. tropical), see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Dichotomopthoropsis M.B. Ellis 1971, *Ascomycota* genera *incertae sedis*, two species, type: *D. nymphaearum* (F.V. Rand) M.B. Ellis, hyphomycetous, sexual morph unknown, plant pathogen, India, USA, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), reference strain of *D. nymphaearum* = ATCC 64770, ATCC 204261-204266, sequences are unavailable.

Dichotophora Whitton, K.D. Hyde & McKenzie 2012, *Ascomycota* genera *incertae sedis*, two species, type: *D. setiramosa* Whitton, McKenzie & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Whitton et al. (2012a; outline), Wijayawardene et al. (2012, 2017; outline), holotype: IFRD 9016, cultures and sequences are unavailable.

Diclonomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *D. eumicrophilus* Thaxt., asexual morph unknown, biotrophic, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dictyoaquaphila J.S. Monteiro, L.B. Conc., M.F.O. Marques, Gusmão & R.F. Castañeda 2016, *Ascomycota* genera *incertae sedis*, two species, type: *D. appendiculata* J.S. Monteiro, L.B. Conc., M.F.O. Marques, Gusmão & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobic, terrestrial, fresh water, Brazil, Mexico see Arias et al. (2016; taxonomy), Monteiro et al. (2016a; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Dictyoarthrinium S. Hughes 1952, *Apiosporaceae*, *Amphisphaerales*, *Sordariomycetes*, six species, type:

D. quadratum S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread (tropical), see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), Crous and Groenewald (2013; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanam et al. (2017a; phylogeny, evolution), cultures and sequences are unavailable for type species.

Dictyasterina Hansf. 1947, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. conopharyngiae* Hansf., asexual morph unknown, saprobes, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; accepted as genus in *Venturiaceae*), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dictyocatenulata Finley & E.F. Morris 1967, *Ostropomycetidae* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *D. alba* Finley & E.F. Morris, hyphomycetous, sexual morph unknown, saprobes, Karnataka, India, widespread, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), An et al. (2012; phylogenetic revision), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Dictyoceratospora Y.R. Ma & X.G. Zhang 2016, *Ascomycota* genera *incertae sedis*, three species, type: *D. cordata* Y.R. Ma & X.G. Zhang, hyphomycetous, sexual morph unknown, saprobic, terrestrial, China, see Ma et al. (2016d; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Dictyochaeta Speg. 1923 (= *Codinaeopsis* Morgan-Jones 1972; = *Dictyochaetopsis* Aramb. & Cabello 1990; = *Menisporella* Agnihothr. 1962 *fide* Seifert et al. 2011; *Species Fungorum* 2017), *Chaetosphaeriaceae*, *Chaetosphaerales*, *Sordariomycetes*, c. 80 species, type: *D. fuegiana* Speg., hyphomycetous, sexual morph unknown, saprobes, terrestrial/aquatic, widespread, see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Silva and Gusmão (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Liu et al. (2016a; new species, phylogeny), cultures and sequences are available but lacking for type species.

Dictyochirospora D'souza, Boonmee & K.D. Hyde 2016, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. rotunda* D'souza, Bhat & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, Thailand, see Boonmee et al. (2016a; phylogeny), Wang et al. (2016f; new species, phylogeny), Wijayawardene

et al. (2017; outline), holotype: MFLUCC 14-0293, cultures and sequences are available.

Dictyocoprotus J.C. Krug & R.S. Khan 1991, *Ascodesmidaceae, Pezizales, Pezizomycetes*, one species, type: *C. nipigonensis* Malloch & Cain, asexual morph unknown, coprophilous, terrestrial, Mexico, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dictyocyclus Sivan., W.H. Hsieh & Chi Y. Chen 1998, *Myriangiaceae, Myriangiales, Dothideomycetes*, one species, type: *D. hydrangeae* Sivan., W.H. Hsieh & Chi Y. Chen, asexual morph unknown, epiphytes, terrestrial, Taiwan, see Hofmann (2009; review, accepted as a genus in *Parmulariaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Parmulariaceae*), Hyde et al. (2013; review, keys, accepted as a genus in *Parmulariaceae*), Dissanayake et al. (2014; taxonomy, accepted as a genus in *Myriangiaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Myriangiaceae*), placement in *Myriangiaceae* supported by morphology, holotype: IMI 371179, cultures and sequences unavailable.

Dictyodesmium S. Hughes 1951, *Ascomycota* genera *incertae sedis*, four species, type: *D. ulmicola* (Ellis & Kellerm.) S. Hughes, hyphomycetous, sexual morph unknown, on living leaves of *Ulmus fulva*, USA, France, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dictyodochium Sivan. 1984, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. prinsepiae* Sivan., asexual morph unknown, saprobes, terrestrial, Africa, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes*, genera *incertae sedis*), data inadequate, cultures and sequences are unavailable.

Dictyodothis Theiss. & Syd. 1915, *Dothideaceae, Dothideales, Dothideomycetes*, two species, type: *D. berberidis* (Rehm) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, America, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; accepted as a genus in *Dothideaceae*), Wijayawardene et al. (2014c; outline), sequences are available.

Dictyographa Müll. Arg. 1893, *Opegraphaceae, Arthoniales, Arthoniomycetes*, two species, type: *D. arabica* Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline, treated as a synonym of *Opegrapha*), Ertz and Tehler (2011; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Dictyomeridium Aptroot, M.P. Nelsen & Lücking 2016, *Trypetheliaceae, Trypetheliales, Dothideomycetes*, seven species, type: *D. proponens* (Nyl.) Aptroot, M.P. Nelsen & Lücking, asexual morph unknown, lichenized, terrestrial, tropical, see Aptroot and Lücking (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Dictyonella Höhn. 1909 (= *Angiotheca* Syd. 1939; = *Calopeziza* Syd. & P. Syd. 1913; = *Ekmanomyces* Petr. & Cif. 1932; = *Masonia* Hansf. 1944 *fide* Species Fungorum 2017), *Saccardiaceae, Ascomycota* families *incertae sedis*, seven species, type: *D. erysiphoides* (Rehm) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), data inadequate, cultures and sequences are unavailable.

Dictyopalmispora Pinruan, Boonmee & K.D. Hyde 2016, *Dictyosporiaceae, Pleosporales, Dothideomycetes*, one species, type: *D. palmae* Pinruan & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, Thailand, see Boonmee et al. (2016a; phylogeny), Wijayawardene et al. (2017; outline), holotype: BBH 39848, cultures and sequences are available.

Dictyopeltella Bat. & I.H. Lima 1959, *Micropeltidaceae, Microthyriales, Dothideomycetes*, two species, type: *D. domingensis* (Petr. & Cif.) Bat. & I.H. Lima, asexual morph unknown, saprobes, terrestrial, tropical see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Dictyopeltis Theiss. 1913, *Dothideomycetes* genera *incertae sedis*, six species, type: *D. vulgaris* (Racib.) Theiss., asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline, as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes*, genera *incertae sedis*), data inadequate, cultures and sequences are unavailable.

Dictyophrynella Bat. & Cavalc. 1964, *Ascomycota* genera *incertae sedis*, one species, type: *D. bignoniacearum* Bat. & Cavalc., hyphomycetous, sexual morph unknown, lichenicolous, Brazil, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Dictyopolyschema M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, one species, type: *D. pirozynskii*, hyphomycetous, sexual morph unknown, saprobes on dead twigs of *Picea*, Great Britain, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), ex-type: IMI 89652, sequences are unavailable.

Dictyoporthe Petr. 1955, *Gnomoniaceae, Diaporthales, Sordariomycetes*, three species, type: *D. ahmadii* Petr., asexual morph unknown, saprobes, terrestrial, Pakistan, Taiwan, see Voglmayr and Jaklitsch (2008; DNA, transferred *Dictyoporthe acerophila* M.E. Barr to *Prostheciaceae*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are available.

Dictyostostrella U. Braun 1999, *Ascomycota* genera *incertae sedis*, one species, type: *D. caricicola* (U. Braun & Melnik) U. Braun, asexual morph hyphomycetous, terrestrial, foliicolous, Asia (Far East of Russia), North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of type: DAOM, cultures and sequences are unavailable.

Dictyospiores M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, one species, type: *D. ziziphi* M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dictyospora Brahamanage, Y.Z. Lu, Boonmee & K.D. Hyde 2017, *Tubeufiaceae, Tubeufiales, Dothideomycetes*, two species, type: *D. thailandica* Brahamanage, Y.Z. Lu, Boonmee & K.D. Hyde, asexual morph hyphomycetous, saprobes, aquatic, Asia, see Brahamanage et al. (2017; taxonomy, phylogeny), holotype of the type: MFLU 17-0500, cultures and sequences are available.

Dictyosporella Abdel-Aziz 2015, *Annulatascaeae, Annulatascales, Sordariomycetes*, one species, type: *D. aquatica* Abdel-Aziz, asexual morph unknown, saprobes, aquatic, Egypt, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type: CBS H-22127, MD1302.

Dictyosporium Corda 1836 (= *Botryosporium* Schwein. 1832 [1834]; = *Cattanea* Garov. 1875; = *Speira* Corda 1837; = *Synphragmidium* F. Strauss 1853 *fide* Species Fungorum 2017), *Dictyosporiaceae, Pleosporales, Dothideomycetes*, c. 50 species, type: *D. elegans* Corda, hyphomycetous, saprobes, terrestrial, aquatic, widespread, see Crous et al. (2009a; new species), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Liu et al. (2015a; new species), Abdel-Aziz (2016b; new species), Boonmee et al. (2016a; new species), cultures available for type species NBRC 32502.

Dictyostomiopelta Viégas 1944, *Dothideomycetes*, genera *incertae sedis*, one species, type: *D. manihotcola*

Viégas, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Dictyothyriella Spieg. 1924, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. bauhiniae* (Rehm) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, Cape Horn, see Lumbsch and Huhndorf (2010; outline, as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Dictyothyrina Theiss. 1913, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. fecunda* (Sacc.) Theiss., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Dictyothrium Theiss. 1912, *Dothideomycetes* genera *incertae sedis*, two species, type: *D. chalybeum* (Rehm) Theiss., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline, as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Diddensiella Péter, Dlauchy & Kurtzman 2012, *Trichomonascaceae, Saccharomycetales, Saccharomycetes*, three species, type: *D. caesifluorescens* G. Péter, Dlauchy & Kurtzman, asexual reproduction is by multilateral budding, saprophytic, rotten wood and forest litter, worldwide, see Péter et al. (2012; taxonomy), sequences are available.

Didonia Velen. 1934, *Helotiales* genera *incertae sedis*, *Leotiomyces*, five species, type: needs typification, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Didymascella Maire & Sacc. 1903 (= *Keithia* Sacc. 1892 *fide* Species Fungorum 2017), *Cenangiaceae, Helotiales, Leotiomyces*, five species, type: *D. oxycedri* Maire & Sacc., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available for *Didymascella thujina*.

Didymascus Sacc. 1896, *Rhytismatales* genera *incertae sedis*, *Leotiomyces*, one species, type: *D. metkinoffii* Sacc.,

asexual morph unknown, saprobes, terrestrial, Siberia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Didymella Sacc. 1880 (= *Arcangelia* Sacc. 1890; = *Dothisphaeropsis* Höhn. 1919; = *Haplotheciella* Höhn. 1918; = *Lichenosphaeria* Bornet 1873; = *Mycosphaerellopsis* Höhn. 1918 *fide* Hyde et al. 2013; Species Fungorum 2017), *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, c. 100 species, type: *D. exigua* (Niessl) Sacc., asexual morph coelomycetous, saprobes, pathogens, terrestrial, worldwide, see de Gruyter et al. (2009; phylogeny), Aveskamp et al. (2010; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Didymellaceae*, taxonomy), Wijayawardene et al. (2014c; outline), Chen et al. (2015b; phylogeny, new species, taxonomy), Liu et al. (2015a; new species), Thambugala et al. 2017b, new species, phylogeny), cultures and sequences are available.

Didymellocamarosporium Wijyaw., Camporesi, D.J. Bhat & K.D. Hyde 2016, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. tamaricis* Wijyaw., Camporesi, D.J. Bhat & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2016b, 2017; taxonomy, phylogeny, outline), holotype and ex-type: MFLU 15-3548, MFLUCC 14-0241.

Didymellopsis (Sacc.) Clem. & Shear 1931 (= *Didymella* subgen. *Didymellopsis* Sacc. 1905 *fide* Species Fungorum 2017), *Xanthopyreniaceae*, *Collemopsidiales*, *Dothideomycetes*, five species, type: *D. latitans* (Nyl.) Sacc. ex Clem. & Shear, asexual morph unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, family placed as *Ascomycota* families *incertae sedis*), Kirk et al. (2013; genus accepted), Zhurbenko et al. (2015b; new species), holotype: H-NYL 42670.

Didymobotryum Sacc. 1886, *Sordariomycetes* genera *incertae sedis*, one species, type: *D. rigidum* (Berk. & Broome) Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), cultures available for type: JCM 8837.

Didymochaetina Bat. & J.L. Bezerra 1965, *Ascomycota* genera *incertae sedis*, one species, type: *D. galactiae* Bat. & J.L. Bezerra, coelomycetous, sexual morph unknown, saprobes, terrestrial, Jamaica, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Didymochora Höhn. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *D. betulina* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Jamaica, see Wijayawardene et al. (2012, 2014c; outline, listed under

Mycosphaerellaceae and *Dothideomycetes*, genera *incertae sedis* respectively), cultures and sequences are unavailable.

Didymocoryne Sacc. & Trotter 1913, *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, type: *D. striata* (Ellis & Everh.) Sacc. & Trotter, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; reduced *D. solitaria* (Rehm) Sacc. & Trotter under *Ascocoryne albida* (Berk.) Seifert, cultures and sequences are unavailable.

Didymocrea Kowalski 1965, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. sadasivanii* (T.K.R. Reddy) Kowalski, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Didymosphaeriaceae*), Ariyawansa et al. (2014e; accepted as a genus in *Didymosphaeriaceae*, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), cultures available for type: CBS 438.65.

Didymocytidium Vain. 1921, *Dothideomycetes* genera *incertae sedis*, two species, type: need typification, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Didymocytis Vain. 1921 (= *Didymocytis* Vain. 2015; = *Diederichia* D. Hawksw. 2003 *fide* Ertz et al. 2015a; Species Fungorum 2017), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 20 species, type: *D. consimilis* Vain., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), Ertz et al. (2015a; taxonomy), cultures and sequences are unavailable.

Didymolepta Munk 1953, *Dothideomycetes* genera *incertae sedis*, two species, type: *D. winteriana* (Sacc.) Munk, asexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Didymoplella Munk 1953, *Dothideomycetes* genera *incertae sedis*, four species, type: *D. cladii* (P. Larsen & Munk) Munk, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Didymosphaeria Fuckel (1870) [1869–70] (= *Cryptodidymosphaeria* (Rehm) Höhn. 1917; = *Didymascina* Höhn. 1905; = *Didymospharella* Cooke 1889; = *Didymosphaeria* subgen. *Cryptodidymosphaeria* Rehm 1906; = *Haplovalsaria* Höhn. 1919; = *Massariellops* Curzi 1927 *fide* Species Fungorum 2017),

Didymosphaeriaceae, *Pleosporales*, *Dothideomycetes*, 25 species, type: *D. futilis* (Berk. & Broome) Rehm, asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014a, d; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type species.

Didymosporiella Traverso & Migliardi 1911, *Ascomycota* genera *incertae sedis*, type: *D. aeluropodis* Traverso & Migliardi, Kirk et al. (2008; treated as nom. dub.), Kirk et al. (2013; genus accepted).

Didymosporina Höhn. 1916 (= *Phaeomarssonia* Bubák 1915 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *D. aceris* (Lib.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2016b, 2017; outline, morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Didymostilbe Henn. 1902 (= *Putagraivam* Subram. & Bhat 1978 *fide* Species Fungorum 2017), *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, eight species, type: *D. coffeae* Henn., hyphomycetous, saprobes, terrestrial, Asia, see Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Bionectriaceae*), Lombard et al. (2016; phylogeny), cultures and sequences are available.

Didymothozetia Rangel 1915 (= *Dithozetia* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *D. mimosoensis* Rangel, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Didysimulans Tibpromma, Camporesi & K.D. Hyde 2017, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *D. mezzanensis* Tibpromma, Camporesi & K.D. Hyde, asexual morph with chlamydospores, saprobes, terrestrial, Italy, see Tibpromma et al. (2017b; ntaxonomy), holotype of the type: MFLU 15-1490, cultures and sequences are available.

Diedickea Syd. & P. Syd. 1913, *Ascomycota* genera *incertae sedis*, three species, type: *D. singularis* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diehlomyces Gilkey 1954, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *D. microsporus* (Diehl & E.B. Lamb.) Gilkey, asexual morph unknown, fungicolous,

terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, treated as *Ascomycota*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Diffractella Guarro, P.F. Cannon & Aa 1991, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *D. curvata* (Fuckel) Guarro, P.F. Cannon & Aa, asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Dicatenosporium S.M. Leão, Gusmão & R.F. Castaneda 2015, *Ascomycota* genera *incertae sedis*, one species, type: *D. polyramosum* S.M. Leão, Gusmão & R.F. Castaneda, hyphomycetous, sexual morph unknown, saprobes, Brazil, see Leão-Ferreira et al. (2015; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Digitodesmium P.M. Kirk 1981, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *D. elegans* P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, Europe, see Silvera-Simon et al. (2010; new species), Seifert et al. (2011; morphology), Boonmee et al. (2016a; phylogeny), Wijayawardene et al. (2017; outline), holotype: BBH 39848, cultures and sequences are available.

Digitodochium Tubaki & Kubono 1989, *Ascomycota* genera *incertae sedis*, one species, type: *D. rhodoleucum* Tubaki & Kubono, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Digitomyces Mercado, M. Caldúch & Gené 2003, *Ascomycota* genera *incertae sedis*, one species, type: *D. verrucosus* (Tzean & J.L. Chen) Mercado, M. Caldúch & Gené, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Digitopodium U. Braun, Heuchert & K. Schub. 2005, *Ascomycota* genera *incertae sedis*, one species, type: *D. hemileiae* (Steyaert) U. Braun, Heuchert & K. Schub., hyphomycetous, sexual morph unknown, terrestrial, fungicolous, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: BPI 426854.

Digitoramispora R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, three species, type: *D. caribensis* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes,

terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: INIFAT C89/167, cultures and sequences are unavailable, compare with *Acrodictys*, *Acrodictyopsis*, *Cheiroidea* and *Cryptocoryneopsis*.

Digitosarcinella S. Hughes 1984, *Englerulaceae*, *Dothideomycetes*, families *incertae sedis*, three species, type: *D. caseariae* S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Digitothyrea P.P. Moreno & Egea 1992, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, three species, type: *D. rotundata* (Büdel, Henssen & Wessels) P.P. Moreno & Egea, asexual morph unknown, lichenized, Africa, Central America, saxicolous, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Dilophospora Desm. 1840 (= *Lidophia* J. Walker & B. Sutton 1974 *fide* Rossman et al. 2016a), *Dothideomycetes* genera *incertae sedis*, c. two species, type: *D. graminis* Desm., coelomycetous, sexual morph formerly named in *Lidophia*, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Diluvicola K.D. Hyde, S.W. Wong & E.B.G. Jones 1998, *Annulatasceae*, *Annulatascales*, *Sordariomycetes*, type: *D. capensis* K.D. Hyde, S.W. Wong & E.B.G. Jones, asexual morph unknown, saprobes, fresh water, Brunei, see Lumbsch and Huhndorf (2010; outline, misspelled as *Diluvicola*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; accepted as a genus in *Annulatasceae*, misspelled as *Diluvicola*), cultures and sequences are unavailable.

Dimastigosporium Faurel & Schotter 1965, *Ascomycota* genera *incertae sedis*, two species, type: *D. musimonum* Faurel & Schotter, coelomycetous, sexual morph unknown, saprobes, terrestrial, Sahara, India, see Yadav and Bhat (2009a; new species, India), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dimelaena Norman 1852 (= *Beltraminia* Trevis. 1857; = *Dimelaena* sect. *Dimelaena* Norman 1853 [1852] *fide* Species Fungorum 2017), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, c. ten species, type: *D. oreina* (Ach.) Norman, asexual morph unknown, lichenized, lichenicolous, saxicolous, on siliceous or basaltic rocks, world-wide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Knudsen et al. (2013b; key), Prieto et al. (2013; phylogeny), Lücking et al. (2017a;

classification), Prieto and Wedin (2016; phylogeny; showed that *Dimelaena* is non-monophyletic as currently circumscribed), sequences are available.

Dimeriella Speg. 1908, *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *D. hirtula* Speg., asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Dimerina Theiss. 1912 (= *Chilemyces* Speg. 1910; = *Dimerinopsis* Syd. & P. Syd. 1917; = *Dimeriopsis* F. Stevens 1917; = *Pilula* Massee 1910; = *Plactogene* Theiss. 1917 [1916]; = *Stigme* Syd. & P. Syd. 1917 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, c. ten species, type: *D. strychni* (Henn.) Theiss., asexual morph unknown, saprobes, terrestrial, Sahara, India, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dimerium (Sacc. & P. Syd.) McAlpine 1903 (= *Dimerium* (Sacc. & P. Syd.) Sacc. & D. Sacc. 1905; = *Dimerosporium* subgen. *Dimerium* Sacc. & P. Syd. 1902; = *Phaeocapnodinula* Speg. 1921 *fide* Species Fungorum 2017), *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *D. pulveraceum* (Speg.) Theiss., see Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; did not include), data inadequate.

Dimeromyces Thaxt. 1896 (= *Eudimeromyces* Thaxt. 1918; = *Jeaneliomycetes* Lepesme 1945 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 114 species, type: *D. africanus* Thaxt., asexual morph unknown, biotrophic, worldwide, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species, Ecuador), Kirk et al. (2013; genus accepted), Rossi et al. (2015; new species, Ecuador), cultures and sequences are unavailable.

Dimerosporiella Speg. 1908 (= *Epinectria* Syd. & P. Syd. 1917 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, seven species, type *D. paulistana* Speg. on living leaves, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: LPS, cultures and sequences are unavailable.

Dimidiographa Ertz & Tehler 2011, *Roccellographaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *D. loandensis* (Nyl.) Ertz, Bungartz & Tehler, sexual morph with lirelliform ascomata, coelomycetous, lichenized, terrestrial, tropical, see Ertz and Tehler (2011; taxonomy, phylogeny), Lücking et al. (2017a; classification), sequences are available for three species.

Dimorphiopsis Crous 2013, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *D. brachystegiae* Crous, coelomycetous or hyphomycetous, sexual morph unknown, Zambia, see Crous et al. (2013b; morphology, DNA), Wijayawardene et al. (2014c, 2016b; outline, morphology, phylogeny), holotype and ex-type: CBS H-21430, CPC 22679, 22680 = CBS 136422.

Dimorphiseta L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *D. terrestris* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Lombard et al. (2016; taxonomy, phylogeny), holotype and ex-type: CBS H-22442, CBS 127345.

Dimorphomyces Thaxt. 1893, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 27 species, type: *Dimorphomyces denticulatus* Thaxt., asexual morph unknown, parasites, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species), cultures and sequences are unavailable.

Dimorphospora Tubaki 1958, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *D. foliicola* Tubaki, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Dimorphotricha Spooner 1987, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *D. australis* Spooner, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dinemasporium Lév. 1846, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, 35 species, type: *D. graminum* (Lib.) Lév., coelomycetous, sexual morph unknown, saprobes, worldwide, see Crous et al. (2012g, 2014c; new species, morphology, phylogeny), Hashimoto et al. (2015a; morphology, phylogeny), Liu et al. (2015a; morphology, phylogeny), Wijayawardene et al. (2016b, 2017; morphology, outline), isotype, epitype and ex-epitype: K(M) 175981, CBS H-20952, CBS 828.84.

Dingleya Trappe 1979, *Tuberaceae*, *Pezizales*, *Pezizomycetes*, six species, type: *D. verrucosa* Trappe, asexual morph unknown, pathogens, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Dioicomyces Thaxt. 1901, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 32 species, type: *D. anthici* Thaxt., asexual morph unknown, parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Bernardi et al.

(2014; Ecuador), cultures and sequences are available for unidentified species.

Diorygma Eschw. 1824, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 71 species, type: *D. hieroglyphica* (Pers.) Staiger & Kalb, asexual morph unknown, lichenized, terrestrial, tropical, see Archer (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2012; DNA, phylogeny), Rivas Plata et al. (2012a; taxonomy), Kirk et al. 2013 (genus accepted), Feuerstein et al. (2014; key), Lumbsch et al. (2014a, b; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type.

Diphymyces I.I. Tav. 1985, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 32 species, type: *D. appendiculatus* (Thaxt.) I.I. Tav., asexual morph unknown, parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Rossi and Santamaria (2010; new species), Kirk et al. (2013; genus accepted), Haelewaters et al. (2014a; new species), cultures and sequences are available for unidentified species.

Diplacella Syd. 1930, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, 32 species, type: *D. paulliniae* (Gonz. Frag. & Cif.) Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for unidentified species.

Diplocarpa Massee 1895, *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *D. curreyana* Massee, asexual morph unknown, fungicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dermateaceae*), Kirk et al. (2013; genus accepted), Pärte et al. (2017; DNA, phylogeny, treated as a genus in *Cordieritidaceae*), cultures and sequences are available, cultures available for type: KL317.

Diplocarpon F.A. Wolf 1912 (= *Entomosporium* Lév. 1857; = *Bostrichonema* Ces. 1867; = *Marssonina* Magnus 1906; = *Entomopeziza* Kleb. 1914 fide Johnston et al. 2014b), *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *D. rosae* F.A. Wolf, asexual morph *Entomosporium*; *Marssonina*, pathogens, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available.

Diplochorina Gutner 1933, *Dothideomycetes* genera incertae sedis, one species, type: *D. naumovii* Gutner, asexual morph unknown, saprobes, terrestrial, USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available for unidentified species.

Diplocadiella G. Arnaud ex M.B. Ellis 1976 (= *Diplocadiella* G. Arnaud 1954 [1953] *fide* Species Fungorum 2017), Ascomycota genera *incertae sedis*, one species, type: *D. scalaroides* G. Arnaud ex M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diplococcum Grove 1885 (= *Luxuriomyces* R.F. Castañeda 1988; = *Tretendophragmia* Subram. 1995 [1992–1993]; = *Tretendophragmia* Subram. 1994 [1991] *fide* Species Fungorum 2017), *Helminthosphaeriaceae*, *Sordariomycetidae*, families *incertae sedis*, *Sordariomycetes*, c. 25 species, type: *D. spicatum* Grove, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Shenoy et al. (2010; phylogeny, polyphyletic), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Hernández-Restrepo et al. (2012b; new species, key), Ma et al. (2012h; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Silva et al. (2014b; new species), Almeida et al. (2015, new species), Maharanachikumbura et al. (2016b; outline, accepted as a genus in *Helminthosphaeriaceae*), cultures available for type species CBS 162.47, CBS 852.73.

Diplodia Fr. 1834 (= *Cryptosphaeria* Grev. 1822; = *Diplodia* a *Eudiplodia* Sacc. 1880; = *Diplodia* b *Botryodiplodia* Sacc. 1880; = *Holcomyces* Lindau 1904 [1905] *fide* Species Fungorum 2017), *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, 23 species, type: *D. mutila* (Fr.) Mont., holomorph, asexual morph coelomycetous, endophytes, pathogens, saprobes, associated with cankers, dieback and fruit rot, worldwide, see Phillips et al. (2008, 2013; morphology, phylogeny), Liu et al. (2012b; morphology, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Yan et al. (2013; species associated with grape wine), Alves et al. (2014; morphology, phylogeny), Wijayawardene et al. (2016b; morphology, phylogeny), holotype and ex-epitype: BPI 599153 and CBS 136014,

Diplodinis Clem. 1931, Ascomycota genera *incertae sedis*, one species, type: *D. rostrupi* (Vestergr.) Clem., coelomycetous, sexual morph unknown, saprobes, terrestrial, Scandinavia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diplodinula Tassi 1902, Ascomycota genera *incertae sedis*, one species, type: need typification, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diplogelasinospora Cain 1961, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, three species, type: *D. princeps* Cain, asexual morph unknown, saprobes, terrestrial, Asia, North America, see Lumbsch and Huhndorf (2010; outline, accepted as genus in *Lasiosphaeriaceae*), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species, DNA), Maharanachikumbura et al. (2016b; outline, accepted as a genus in *Chaetomiaceae*), cultures and sequences are available.

Diplogramma Müll. Arg. 1891 (= *Diplogrammatomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *D. australiensis* Müll. Arg., lichenized, Australia, epiphytic, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; treated as a synonym of *Opegrapha*), sequences are unavailable.

Diploicia A. Massal. 1852 (= *Buellomyces* E.A. Thomas ex Cif. & Tomas. 1953 [1952]; = *Diplotomma* A. Massal. 1852 *fide* Species Fungorum 2017), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, c. twelve species, type: *D. canescens* (Dicks.) A. Massal., asexual morph unknown, lichenized, corticolous, worldwide, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Bungartz et al. (2016; generic concept, key to all species; discussed the distinction from *Diplotomma*, but showed both are distinct), Lücking et al. (2017a; classification), sequence are available for voucher MAF 8656 of the type.

Diplolaeviopsis Giralt & D. Hawksw. 1991, *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *D. ranula* Giralt & D. Hawksw., on lichens, sexual morph unknown, see Kirk et al. (2013; genus accepted), Suija et al. (2015b; position), Pärte et al. (2017; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype of type: BCC-Lich 4120.

Diplomyces Thaxt. 1895, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *D. actobianus* Thaxt., asexual morph unknown, parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Diplonaevia Sacc. 1889 (= *Asteronaevia* Petr. 1929; = *Hysteropezizella* sect. *Pyrenodiscus* (Petr.) Nannf. 1932; = *Merostictis* Clem. 1909; = *Pyrenodiscus* Petr. 1927 *fide* Species Fungorum 2017), *Calloriaceae*, *Helotiales*, *Leotiomycetes*, 23 species, type: *D. caricum* (Auersw.) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diploëspora Grove 1916, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *D. rosea* Grove, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Diploplenodomus Died. 1912 (= *Sclerochaetella* Höhn. 1917 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *D. malvae* Died. ex Died., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diplopodomyces W. Rossi & Balazuc 1977, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *D. callipodos* W. Rossi & Balazuc, asexual morph unknown, parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Santamaría et al. (2014; new species), cultures and sequences are unavailable.

Diploschistella Vain. 1926, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *D. urceolata* Vain., lichenized, pantropical, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Diploschistes Norman 1852 (= *Diploschistomyces* Werner 1976; = *Gyalecta* Eaton 1829; = *Lagerheimina* Kuntze 1891; = *Lectularia* Stirt. 1879 [1878]; = *Patellaria* Hoffm. 1789 [1790]; = *Polyschistes* J. Steiner 1898; = *Protoschistes* M. Choisy 1928; = *Urceolaria* Ach. 1803; = *Urceolariomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, c. 33 species, type: *D. scapus* (Schreb.) Norman, asexual morph coelomycetous, lichenized (young states sometimes parasitic), on rocks, soil or over mosses and other lichens, worldwide, see Lumbsch and Huhndorf (2010; outline), Fermández-Brime et al. (2013; phylogeny), Mangold et al. (2009; key, Australia), Rivas Plata et al. (2010a, 2012a; key, taxonomy), Kirk et al. (2013; genus accepted), Wang et al. (2013b; new species), Lücking et al. (2017a; classification), cultures available for the type: AFTOL-ID 98.

Diplosporonema Höhn. 1917 (= *Phragmosporonema* Moesz 1934 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *D. delastrei* (Lacroix) Höhn. ex Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diplotomma Flot. 1849 (= *Abacina* Norman 1853 [1852] *fide* Species Fungorum 2017), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 30 species, type: *D. alboatum* (Hoffm.) Flot., lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available for voucher MP5 of the type species.

Diplozythiella Died. 1916, *Ascomycota* genera *incertae sedis*, one species, type: *D. bambusina* Died., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dipodascopsis Batra & P. Millner emend. Kurtzman, Albertyn & Basehoar-Powers 2007 (= *Babjevia* Van der Walt & M.T. Sm. 1995 *fide* Kurtzman et al. 2007), *Lipomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, three species, type: *D. uninucleata* (Biggs) L.R. Batra & Millner, asexual reproduction by multilateral budding and pseudohyphae may be present, saprophytic, soil, wood, fruit fly, Canada, USA, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kurtzman and Robnett (2013b; DNA, phylogeny), sequences are available.

Dipodascus de Lagerheim 1892 (= *Endyllum* Clem. 1931; = *Fermentotrichon* E.K. Novák & Zsolt 1961; = *Galactomyces* Redhead & Malloch 1977; = *Geotrichum* Link 1809; = *Magnusiomyces* Zender 1926 [1925]; = *Oosporoidea* Sumst. 1913; = *Polymorphomyces* Coupin 1914; = *Zendera* Redhead & Malloch 1977 *fide* Species Fungorum 2017), *Dipodascaceae*, *Saccharomycetales*, *Saccharomycetes*, ten species, type: *D. albidus* Lagerh., asexual reproduction present (arthroconidia, chlamydospores may be formed), saprophytic, insects, insect galleries in pine trees, fluxes from trees, mushroom gills, cacti, worldwide, see Nagahama et al. (2008; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Dipodomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *D. monstruosus* Thaxt., asexual morph unknown, parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Diporotheca C.C. Gordon & C.G. Shaw 1961, *Diporothecaceae*, *Ascomycota* families *incertae sedis*, two species, type: *D. rhizophila* C.C. Gordon & C.G. Shaw, asexual morph unknown, saprobes, terrestrial, aquatic, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hawksworth

et al. (2016; *Diporotheca palynomorph*), holotype: WSP 48900, cultures and sequences are unavailable.

Dipyrgis Clem. 1909, *Ascomycota* genera *incertae sedis*, one species, type: data inadequate asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dirina Fr. 1825 (= *Dirinopsis* De Not. 1846 *fide* Species Fungorum 2017), *Roccellaceae*, *Arthoniales*, *Arthroniomycetes*, 24 species, type: *D. ceratoniae* (Ach.) Fr., asexual morph unknown, lichenized, saxicolous and corticolous, worldwide (coastal regions of Mediterranean and subtropical climates), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Tehler et al. (2013a; phylogeny and taxonomy), Lücking et al. (2017a; classification, but recording only 13 species), sequences are available.

Dirinaria (Tuck.) Clem. 1909 (= *Pyxine* sect. *Dirinaria* Tuck. 1877 *fide* Species Fungorum 2017), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 35 species, type: *D. picta* (Sw.) Clem. & Shear, lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available for the type.

Dirinastrum Müll.Arg. 1893 (= *Dirinastromyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Caliciaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *D. australiense* Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Bungartz et al. (2011; phylogeny), Lücking et al. (2017a; classification, treated as a genus in *Roccellaceae*), sequences are unavailable.

Diromma Ertz & Tehler 2014, *Roccellaceae*, *Arthoniales*, *Arthroniomycetes*, one species, type: *D. dirinellum* (Nyl.) Ertz & Tehler, sexual morph with apothecioid ascocarps, asexual morph unknown, lichenized, terrestrial, Mediterranean and Canary Islands, see Ertz et al. (2015c; morphology, phylogeny), Lücking et al. (2017a; classification), lectotype and ex-type strain of type: Dufour (H-NYL4756).

Discina (Fr.) Fr. 1849 (= *Discinella* P. Karst. 1891; = *Peziza* a *Discina* Fr. 1822 *fide* Species Fungorum 2017), *Discinaceae*, *Pezizales*, *Pezizomycetes*, 20 species, type: *D. perlata* (Fr.) Fr., asexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discinella Boud. 1885 (= *Pachydisca* Boud. 1885; *Geocoryne* Korf 1978 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, twelve species, type: *D. boudieri* (Quél.) Boud., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Disciotis Boud. 1885, *Morchellaceae*, *Pezizales*, *Pezizomycetes*, four species, type: *D. venosa* (Pers.) Arnould, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Methven et al. (2013; DNA, phylogeny), Carris et al. (2016; mitosporic stage; DNA, phylogeny), cultures and sequences are available.

Discocainia J. Reid & A. Funk 1966, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, type: *D. treleasei* (Sacc.) J. Reid & A. Funk, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Li et al. (2014d; new species, Japan), cultures and sequences are available.

Discocolla Prill. & Delacr. 1894, *Ascomycota* genera *incertae sedis*, one species, type: *D. pyrina* Prill. & Delacr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, north temperate, see Seifert et al. (2011), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discocurtisia Nannf. 1983, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, twelve species, type: *D. arundinariae* (Berk. & M.A. Curtis) Nannf., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discogloeum Petr. 1923, *Ascomycota* genera *incertae sedis*, one species, type: *D. veronicarum* (Ces.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discomycella Höhn. 1912, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *D. tibodensis* Höhn., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discomycetoidae Matsush. 1993, *Ascomycota* genera *incertae sedis*, one species, type: *D. aequatorialis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Discopycnothyrium Hongsanan & K.D. Hyde 2016, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, one species, type: *D. palmae* Hongsanan & K.D. Hyde, coelomycetous, sexual morph unknown, sooty moulds, Thailand, see Hongsanan et al. (2016a; taxonomy), holotype: MFLU13-0485, cultures and sequences are available.

Discorehmia Kirschst. 1936, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *D. eburnea* Kirschst.,

asexual morph unknown, saprobes, terrestrial, North America, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discosia Lib. ex Durieu & Mont. 1849 (= *Adisciso* Kaz. Tanaka, Okane & Hosoya 2011; = *Cryptostictella* Grove 1912; = *Discosia* Lib. 1837; = *Discosiospora* A.W. Ramaley 1989; = *Leptina* Bat. & Peres 1960 *fide* Wijayawardene et al. 2016b; Species Fungorum 2017), *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, c. 15 species, type: *D. strobilina* Lib., coelomycetous, sexual morphs unknown, saprobes, terrestrial, cosmopolitan, see Wijayawardene et al. (2012, 2016b, 2017; taxonomy, outline), Kirk et al. (2013; genus accepted), Li et al. (2015d; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Senanayake et al. (2015; new species, phylogeny), Tangthirasunun et al. (2015; taxonomy), Jaklitsch et al. (2016a; phylogeny), cultures and sequences are available.

Discoscellina Subram. & K.R.C. Reddy 1972, *Ascomycota* genera *incertae sedis*, one species, type: *D. longiciliata* (Agnihothr.) Subram. & K.R.C. Reddy, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discosphaerina Höhn. 1917, *Hyponectriaceae*, *Amphisphaeriales*, *Sordariomycetes*, eleven species, type: *D. discophora* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Discosporina Höhn. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *D. deplanata* (Höhn.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discotheciella Syd. & P. Syd. 1917 (= *Discothecium* Syd. & P. Syd. 1916 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *D. bakeri* (Syd. & P. Syd.) Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Discozythia Petr. 1922, *Ascomycota* genera *incertae sedis*, one species, type: *D. bakeri* (Syd. & P. Syd.) Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Disculoides Crous, Pascoe, I.J. Porter & Jacq. Edwards 2012, *Erythrogloeaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *D. eucalypti* Crous, Pascoe, I.J. Porter & Jacq. Edwards, coelomycetous, sexual morph unknown, pathogens, terrestrial, Australia, see Crous et al. (2012d, 2016a; morphology, new species, phylogeny), Maharachchikumbura et al. (2015b; classification, phylogeny), Crous et al. (2017; new species), holotype and ex-type: CBS H-20935, CPC 17650.

Dissitimirus E.G. Simmons, McGinnis & Rinaldi 1987, *Ascomycota* genera *incertae sedis*, one species, type: *D. exedrus* E.G. Simmons, McGinnis & Rinaldi, hyphomycetous, sexual morph unknown, on human, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dissoconium de Hoog, Oorschot & Hijwegen 1983, *Dissoconiaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *D. aciculare* de Hoog, Oorschot & Hijwegen, hyphomycetous, sexual morph mycosphaerella-like, saprobes, pathogens, terrestrial, worldwide, see Crous et al. (2009d; taxonomy, phylogeny), Seifert et al. (2011; morphology), Hyde et al. (2013; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2014c; outline, phylogeny), cultures and sequences are available.

Distocercospora N. Pons & B. Sutton 1988, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *D. pachyderma* (Syd. & P. Syd.) N. Pons & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, ?pathogens, terrestrial, South Africa, India, New Zealand, Philippines, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Verma and Rai (2014; morphology, new species), Braun et al. (2014b; morphology and typification), neotype S-F37683, epitype MUMH11476, duplicate: CBS H-21733; ex-epitype cultures: MUCC1716, CBS 138247.

Distolomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *D. fijianus* Thaxt., asexual morph unknown, parasites, terrestrial, Asia, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Distophragmia R.F. Castañeda, S.M. Leão & Gusmão & 2015, *Ascomycota* genera *incertae sedis*, one species, type: *D. rigidiuscula* (R.F. Castañeda) R.F. Castañeda, S.M. Leão & Gusmão, hyphomycetous, sexual morph unknown, saprobic, terrestrial, Brazil, Cuba, see Castañeda-Ruiz et al. (2015; taxonomy), Wijayawardene

et al. (2017; outline), cultures and sequences are unavailable.

Distopyrenis Aptroot 1991, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, eight species, type: *D. americana* Aptroot, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Distorimula F. San Martín, P. Lavín & Esqueda 1999, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, two species, type: *D. mexicana* F. San Martín, P. Lavín & Esqueda, asexual morph unknown, saprobes, terrestrial, Mexico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: FSM, Esqueda 3693, cultures and sequences are unavailable.

Distoseptispora K.D. Hyde, McKenzie & Maharachch. 2016, *Distoseptisporaceae*, *Diaportheomycetidae* families *incertae sedis*, *Sordariomycetes*, six species, type: *D. fluminicola* McKenzie, H.Y. Su, Z.L. Luo & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, aquatic, China, Thailand, see Hyde et al. (2016; new species), Doilom et al. (2017; species on *Tectona*), Su et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type: HKAS84006, MFLUCC 15–0417.

Distothelia Aptroot 2005, *Trypetheliaceae*, *Trypethiales*, *Dothideomycetes*, three species, type: *D. isthmospora* Aptroot, lichenized, see Lumbsch and Huhndorf (2010; outline), Aptroot and Lücking (2016; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Ditopella De Not. 1863, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *D. fusispora* De Not., asexual morph unknown, saprobes, terrestrial, Europe, see Mejía et al. (2008, 2011; DNA, phylogeny), Sogonov et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for *D. ditopa*.

Ditopellina J. Reid & C. Booth 1967, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *D. saccardoana* (Traverso & Spessa) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ditopellosis J. Reid & C. Booth 1967, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *D. clethrae* J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, North America, see Mejía et al. (2008, 2011; DNA, phylogeny), Sogonov et al. (2008; DNA, transferred *Ditopellosis racemula* (Cooke & Peck)

M.E. Barr to *Gnomoniopsis*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for unidentified *Ditopellosis* sp.

Diutina Khunnamw., Lertwatt., Jindam., Limtong & Lachance 2015, *Saccharomycetales* genera *incertae sedis*, *Saccharomycetes*, eight species, type: *D. rugosa* (Anderson) Khunnamwong, Jindamorakot, Limtong & Lachance, asexual reproduction by multilateral and pseudohyphae are produced, sexual reproduction unknown, saprophytic, but some species are common clinical isolates, rice leaf, bee, soil, soybean factory, clinical, worldwide, see Khunnamwong et al. (2015; taxonomy), sequences are available.

Diversimorbus S.F. Chen & Jol. Roux 2013, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *D. metrosideri* S.F. Chen bis & Jol. Roux, with coelomycetous asexual morph, saprobes, terrestrial, South Africa, have sequence data, see Chen et al. (2013b; morphology, phylogeny), Maharachchikumbura et al. (2015a, 2015b; outline), holotype and ex-type: PREM 60741, CMW37322.

Divinia Cif. 1955, *Ascomycota* genera *incertae sedis*, one species, type: *D. diatricha* Cif., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Dominican Republic, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dixomyces I.I. Tav. 1985, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *D. perpendicularis* (Thaxt.) I.I. Tav., asexual morph unknown, parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dolabra C. Booth & W.P. Ting 1964, *Chaetothyriomycetidae* genera *incertae sedis*, *Eurotiomycetes*, one species, type: *D. nepheliae* C. Booth & W.P. Ting, asexual morph coelomycetous (*fide* Rossman et al. 2010), saprobes, pathogens, terrestrial, Malaysia, see Lumbsch and Huhndorf (2010; outline, as *Dothideomycetes*, genera *incertae sedis*), Rossman et al. (2010, 2012; *Eurotiomycetes* genera *incertae sedis*, DNA, phylogeny, asexual morph), Kirk et al. (2013; genus accepted), Thambugala et al. (2014b; taxonomy), cultures and sequences are available.

Dolichocarpus R. Sant. 1949, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *D. chilensis* R. Sant., asexual morph unknown, lichenized, saxicolous and epiphytic, Chile and Saint Helena, see Aptroot (2008; taxonomy), Ertz and Tehler (2011; phylogenetic position), Lumbsch and Huhndorf

(2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Doliomyces Steyaert 1961 (= *Bartaliniopsis* S.S. Singh 1974 [1972] *fide Species Fungorum* 2017), *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, two species, type: *D. senegalensis* (Speg.) Steyaert, sexual morph unknown, saprobes, terrestrial, India, South America, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as in *Amphisphaeriaceae*), Wijayawardene et al. (2016b; placed in *Bartaliniaceae* (current name *Sporocadaceae*) based on morphology), cultures and sequences are unavailable.

Domingoella Petr. & Cif. 1932, *Ascomycota* genera *incertae sedis*, four species, type: *D. asterinarum* Petr. & Cif., hyphomycetous, sexual morph unknown, on fungi, terrestrial, Pantropical, Australasia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Donadinia Bellem. & Mel.-Howell 1990, *Sarcosomataceae*, *Pezizales*, *Pezizomycetes*, four species, type: *D. helvelloides* (Donadini, Berthet & Astier) Bellem. & Mel.-Howell, sexual morph unknown, saprobes, terrestrial, Europe, Asia, see Lumbsch and Huhndorf (2010; outline), Carbone et al. (2013, 2014; new species; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Doratomyces Corda 1829, *Microascaceae*, *Microasciales*, *Sordariomycetes*, two species, type: *D. neesii* Corda, sexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Sandoval-Denis et al. (2016b; phylogeny), cultures and sequences are available.

Dothichiza Lib. ex Roum. 1880 (= *Microphoma* N.F. Buchw. 1958; = *Parasclerophoma* Petr. 1924; = *Tylophoma* Kleb. 1933 *fide Species Fungorum* 2017), *Dothideomycetes* genera *incertae sedis*, 15 species, type: *D. populea* Sacc. & Briard, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c; outline, accepted as *Dothioraceae*, and *Dothideomycetes*, genera *incertae sedis* respectively), Kirk et al. (2013; genus accepted), Gonzalez-Menendez et al. (2016; DNA), Hyde et al. (2013) and Thambugala et al. (2014a) excluded from *Dothioraceae*, cultures and sequences are available.

Dothidasteroma Höhn. 1909 (= *Melanoplaca* Syd. & P. Syd. 1917 *fide Species Fungorum* 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, five species, type: *D. pterygotae* (Berk. & Broome) Höhn., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Parmulariaceae*),

Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Parmulariaceae*), Hyde et al. (2013; review, keys, excluded from *Parmulariaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Parmulariaceae*), placement in *Parmulariaceae* supported by morphology, cultures and sequences are unavailable.

Dothidasteromella Höhn. 1910 (= *Balansina* G. Arnaud 1918 [1917]; = *Dothidasteris* Clem. & Shear 1931; = *Hysteroostoma* Theiss. 1913 *fide Species Fungorum* 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *D. sepulta* (Berk. & M.A. Curtis) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hyde et al. (2013) did not accept as a genus in *Asterinaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Dothidea Fr. 1818 (= *Phragmodothis* Theiss. & Syd. 1914; = *Systremma* Theiss. & Syd. 1915 *fide Species Fungorum* 2017), *Dothideaceae*, *Dothideales*, *Dothideomycetes*, around 500 spp. (the actual number of species could be less than 20), type: *D. sambuci* (Pers.) Fr., asexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; morphology, phylogeny), Wijayawardene et al. (2014c; outline), ex-type strains of type: DAOM 231303, epitype of type: GZU 78–2002.

Dothidella Speg. 1880 (= *Pluriporus* F. Stevens & R.W. Ryan 1925; = *Polystomella* Speg. 1888 *fide Species Fungorum* 2017), *Polystomellaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *D. australis* Speg., sexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Dothideodiplodia Murashk. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *D. agropyri* Murashk., coelomycetous, sexual morph unknown, saprobes, terrestrial, USSR, see Wijayawardene et al. (2012, 2017; outline, morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dothideopsella Höhn. 1915, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. agminalis* (Sacc. & Mortier) Höhn., asexual morph unknown, saprobes, terrestrial, see Kirk et al. (2008; treated as a synonym of *Leptosphaeria*), Lumbsch and Huhndorf (2010; outline), Ariyawansa et al. (2013b; morphology), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Dothidotthia Höhn. 1918, *Dothidotthiaceae*, *Pleosporales*, *Dothideomycetes*, c. ten species, type: *D. symphoricarpi* (Rehm) Höhn., asexual morph unknown, saprobes, terrestrial, temperate, see Phillips et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, phylogeny), epitype and ex-epitype: BPI 871823, CPC 12929 = CBS 119687.

Dothiopeltis E. Müll. 1957, *Leptopeltidaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *D. arunci* E. Müll., asexual morph idriella-like, saprobes, terrestrial, Switzerland, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Leptopeltidaceae*), Granmo and Mathiassen (2013; new species), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Dothiora Fr. 1849 (= *Dothiora* subgen. *Metadothis* Sacc. 1889; = *Jaapia* Kirschst. 1938; = *Keisslerina* Petr. 1920 [1919]; = *Leptodothiora* Höhn. 1918; *Metadothis* (Sacc.) Sacc. 1892; = *Stigmea* Bonord. 1864 *fide* Species Fungorum 2017), *Dothideaceae*, *Dothideales*, *Dothideomycetes*, more than 50 species, type: *D. pyrenophora* (Fr.) Fr., asexual morph coelomycetous, hormonema-like synasexual morph, dothichiza or hormonema-like, saprobes or pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothioraceae*), Hyde et al. (2013; accepted as a genus in *Dothioraceae*), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a), Tibpromma et al. (2017b; phylogeny, asexual morph), cultures and sequences are available.

Dothiorella Sacc. 1880 (= *Botryophoma* (P. Karst.) Höhn. 1916; = *Phoma* subgen. *Botryophoma* P. Karst. 1884; = *Sclerodothiorella* Died. 1912 *fide* Species Fungorum 2017), *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, 24 species, type: *D. pyrenophora* Sacc., holomorph, asexual morph coelomycetous, endophytes, pathogens, saprobes, associated with wide range of woody hosts, worldwide distribution, see Phillips et al. (2008, 2013; morphology, phylogeny), Liu et al. (2012a, b, phylogeny), Li et al. (2014c; new species), Dissanayake et al. (2016a; morphology, phylogeny), Crous et al. (2016b; new species, phylogeny), Wijayawardene et al. (2016b; morphology, phylogeny), Doilom et al. (2017; species in *Tectona*), holotype and culture ex-epitype: IMI 63581b, CBS 115038.

Dothioropsis Riedl 1974, *Ascomycota* genera *incertae sedis*, one species, type: *D. corni* Riedl, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Dothistroma Hulbary 1941, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *D. pini* Hulbary, coelomycetous, sexual morph mycosphaerella-like, plant pathogens, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Braun et al. (2014b; morphology and typification), Barnes et al. (2016; neotypification of *Dothistroma septosporum* and epitypification of *D. pini*), epitype and ex-epitype: CBS H-12211, CMW 10951 = CBS 116487.

Dothivalsaria Petr. 1966, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. megalospora* (Auersw.) Petr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as genus in *Massariaceae*), Hyde et al. (2013; did not include as a genus in *Massariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are available.

Drechmeria W. Gams & H.-B. Jansson 1985 (= *Haptocillium* W. Gams & Zare *fide* Maharanachikumbura et al. 2015a, 2015b), *Ophiocordycipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *D. coniospora* (Drechsler) W. Gams & H.-B. Jansson, sexual morph unknown, parasitic, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline, phylogeny), Spatafora et al. (2015; new species), Zhang et al. (2016c; endoparasites of nematodes, DNA), cultures and sequences are available.

Drechslerella Subram. 1964 (= *Dactylariopsis* Mekht. 1967; = *Golovinia* Mekht. 1967; = *Orbiliaster* Dennis 1954 *fide* Baral et al. 2015), *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, two species, type: *D. acrochaeta* (Drechsler) Subram., sexual morph unknown, parasitic, terrestrial, worldwide, see Castillo et al. (2010; biocontrol), Kirk et al. (2013; genus accepted), Liu et al. (2014b; genomic studies), Baral et al. (2015; nomenclature), cultures and sequences are available.

Drepanomyces Thaxt. 1931, *Ceratomyctaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *D. malayanus* Thaxt., asexual morph unknown, parasitic, terrestrial, Sumatra, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Drepanopeziza (Kleb.) Höhn. 1917 (= *Dirimosperma* Preuss 1855 [1853]; = *Drepanopeziza* (Kleb.) Höhn. 1917; = *Gloeosporium* Desm. & Mont. 1849; = *Gloeosporidiella* Petr. 1921; = *Microsphaeropsis* Höhn. 1917; = *Pseudopeziza* subgen. *Drepanopeziza* Kleb. 1906 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *D. ribis* (Rehm ex Kleb.) Höhn., asexual

morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Drudeola Kuntze 1891 (= *Macilvainea* Nieuwl. 1916; = *Peckia* Clinton 1878 [1876] *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *D. sarraceniae* (Peck & Cooke) Kuntze, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Drumopama Subram. 1957, *Ascomycota* genera *incertae sedis*, one species, type: *D. girisa* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, Cuba, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dryosphaera Jørgen Koch & E.B.G. Jones 1989, *Ascomycota* genera *incertae sedis*, three species, type: *D. navigans* Jørg. Koch & E.B.G. Jones, asexual morph unknown, on wood associated with sand, marine, temperate and tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), cultures and sequences are unavailable.

Dualomyces Matsush. 1987, *Ascomycota* genera *incertae sedis*, two species, type: *D. taiwanensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dubitatio Speg. 1882, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. dubitationum* Speg., asexual morph coelomycetous, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Massariaceae*), Hyde et al. (2013) excluded from *Massariaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Dubujiana D.R. Reynolds & G.S. Gilbert 2005, *Dothideomycetes* genera *incertae sedis*, one species, type: *D. glandulifera* D.R. Reynolds & G.S. Gilbert, asexual morph coelomycetous, saprobes, terrestrial, Australia, see Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Duebenia Fr. 1849 (= *Briardia* Sacc. 1885 *fide* Species Fungorum 2017), *Calloriaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *D. rufa* Fr., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dufourea Ach. 1809 (see Species Fungorum 2017 for synonyms), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 23 species, type: *D. flammea* (L. f.) Ach., lichenized, see Arup et al. (2013; taxonomy), Kondratyuk et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Dumontinia L.M. Kohn 1979, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *D. tuberosa* (Bull.) L.M. Kohn, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny), cultures and sequences are available.

Duosporium K.S. Thind & Rawla 1961, *Ascomycota* genera *incertae sedis*, one species, type: *D. cyperi* K.S. Thind & Rawla, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, China, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Duplicaria Fuckel 1870, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *D. empetri* (Pers.) Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zou et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but type lacking DNA.

Duplicariella B. Erikss. 1970, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *D. phyllodoces* B. Erikss., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Lantz et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Duradens Samuels & Rogerson 1990, *Sordariomycetes* genera *incertae sedis*, one species type: *D. lignicola* Samuels & Rogerson, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Clypeosphaeriaceae*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, excluded from *Clypeosphaeriaceae* and placed in *Sordariomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Durandiella Seaver 1932 (= *Durandia* Rehm 1913 *fide* Species Fungorum 2017), *Tympanidaceae*, *Phacidiales*, *Leotiomycetes*, ten species, type: *D. fraxini* (Schwein.) Seaver, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Durella Tul. & C. Tul. 1865, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, ?five species, type: *D. compressa* (Pers.) Tul. & C. Tul., asexual morph ?coelomycetous, saprobes on wood or cones of *Pinus*, widespread in

temperate areas, see Kirk et al. (2013; genus not listed), Jaklitsch et al. (2016a; classification), cultures and sequences are available but lacking for the type species, holotype of type: PH0006280.

Durispora K.D. Hyde 1994, *Valsaceae, Diaporthales, Sordariomycetes*, two species, type: *D. elaeidicola* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Malaysia, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Durotheca Læssøe, Srikitik., Luangsa-ard & M. Stadler 2013, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *D. depressa* Læssøe & Srikitkulchai, asexual morph nodulisporium-like, saprobes, terrestrial, Thailand, see Læssøe et al. (2013; morphology, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Li et al. (2016a; new species, DNA), holotype and ex-type: BBH 18222, BBC 23016.

Dussiella Pat. 1890 (= *Echinodothis* G.F. Atk. 1894 *fide* Species Fungorum 2017), *Clavicipitaceae, Hypocreales, Sordariomycetes*, one species, type: *D. tuberiformis* (Berk. & Ravenel) Pat., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Dwayabeeja Subram. 1958, *Ascomycota* genera *incertae sedis*, three species, type: *D. sundara* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dwayaloma Subram. 1957, *Ascomycota* genera *incertae sedis*, two species, type: *D. trina* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dwayalomella Brisson, Piroz. & Pauzé 1975, *Ascomycota* genera *incertae sedis*, one species, type: *D. vaccinii* Brisson, Piroz. & Pauzé, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dwibahubeeja N. Srivast., A.K. Srivast. & Kamal 1995, *Ascomycota* genera *incertae sedis*, one species, type: *D. indica* N. Srivast., A.K. Srivast. & Kamal, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Dwibeeja Subram. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *D. sundara* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dwiroopa Subram. & Muthumary 1986, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *D. ramya* Subram. & Muthumary, coelomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Dwiroopella Subram. & Muthumary 1986, *Ascomycota* genera *incertae sedis*, one species, type: *D. sundara* Subram. & Muthumary, coelomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Dyfrolomyces K.D. Hyde, K.L. Pang, Alias, Suetrong & E.B.G. Jones 2013, *Dyfrolomycetaceae, Dyfrolomycetales, Dothideomycetes*, four species, type: *D. tiomanensis* K.L. Pang, S.A. Alias, K.D. Hyde, Suetrong & E.B.G. Jones, asexual morph unknown, saprobes, on wood in aquatic or terrestrial environments, worldwide, see Hyde et al. (2013; phylogeny, morphology, outline), Pang et al. (2013a; phylogeny, morphology), Shearer et al. (2014; review), Wijayawardene et al. (2014c; outline), Jones et al. (2015; classification, marine), Maharachchikumbura et al. (2016b; outline), holotype: MFLU 13-00063.

Dyplolabia A. Massal. 1854, *Graphidaceae, Ostropales, Lecanoromycetes*, four species, type: *D. afzelii* (Ach.) A. Massal., asexual morph unknown, lichenized, terrestrial, tropical, see Rivas Plata et al. (2012a; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kalb et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Dyrithiopsis L. Cai, Jeewon & K.D. Hyde 2003 (= *Monochaetiopsis* L. Cai et al. 2003 *fide* Maharachchikumbura et al. 2015b; Réblová et al. 2016c), *Sporocadaceae, Amphisphaerales, Sordariomycetes*, one species, type: *D. lakefuxianensis* L. Cai, Jeewon & K.D. Hyde, asexual morph formerly known in *Monochaetiopsis*, saprobes, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Bartaliniaceae*), Réblová et al. (2016c; nomenclature), Wijayawardene et al. (2012, 2017;

outline), holotype: HKU (M) 8280, cultures and sequences are available.

Dyribium M.E. Barr 1994, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type: *D. lividum* (Fr.) M.E. Barr, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Dysrhynchis Clem. 1909 (= *Balladynella* Theiss. & Syd. 1918 [1917]; = *Ceratochaete* Syd. & P. Syd. 1917; = *Dimerosporiella* Höhn. 1909; = *Dimerosporina* Höhn. 1910; = *Henningsomyces* Sacc. & D. Sacc. 1905; = *Kusanotheca* Bat. & Cif. 1963; = *Neoballadyna* Boedijn 1961; = *Neohoehnelia* Theiss. & Syd. 1918 [1917]; = *Setella* Syd. & P. Syd. 1916 *fide* Species Fungorum 2017), *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, seven species, type: *D. pulchella* (Sacc.) Clem., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Perisporiopsidaceae*), Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Ebollia Minter & Caine 1980, *Ascomycota* genera *incertae sedis*, one species, type: *E. valdiviensis* (Speg.) Minter & Caine, coelomycetous, sexual morph unknown, saprobes, terrestrial, Chile, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ebullia K.L. Pang 2015, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *E. octonae* (Kohlm.) K.L. Pang, asexual morph unknown, saprobes, marine, Brunei, Hawaiian Islands, India, Taiwan, sequence data available, see Chu et al. (2015), Maharachchikumbura et al. (2016b; outline), holotype: J.K. 4474a = 01271926, NYBG.

Echidnocybium Brusse 1987, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *E. speciosum* Brusse, asexual morph coelomycetous, lichenized, saxicolous, South Africa, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Echidnodella Theiss. & Syd. 1918, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, 34 species, type: *E. linearis* (Syd. & P. Syd.) Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, no molecular data, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Hongsanan et al. (2014c; taxonomy), Wijayawardene et al. (2014c; outline), placement in *Asterinaceae* supported by morphology, cultures and sequences are unavailable.

Echidnodes Theiss. & Syd. 1918 (= *Lembosiodothis* Höhn. 1917; = *Maurodothella* G. Arnaud 1918 [1917] *fide* Species Fungorum 2017), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, c. 17 species, type: *E. liturae* (Cooke) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Echidnodes*), Hongsanan et al. (2014c; phylogeny), cultures and sequences are unavailable for type species.

Echinoascotheca Matsush. 1995, *Phaeotrichaceae*, *Phaeotrichales*, *Dothideomycetes*, one species, type: *E. duploformis* Matsush., asexual morph unknown, saprobes, terrestrial, Pakistan, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; accepted as genus in *Phaeotrichaceae*), Hyde et al. (2013; accepted as genus in *Phaeotrichaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Echinobotryum Corda 1831, *Microascaceae*, *Microascales*, *Sordariomycetes*, one species, type: *E. atrum* Corda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Echinocatena R. Campb. & B. Sutton 1977, *Ascomycota* genera *incertae sedis*, one species, type: *E. arthrinoides* R. Campb. & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: IMI 199279, cultures and sequences are unavailable.

Echinochondrium Samson & Aa 1975, *Ascomycota* genera *incertae sedis*, one species, type: *E. pulchrum* Samson & Aa, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Echinococonidiophorum Pereira-Carv. & Dianese 2009, *Ascomycota* genera *incertae sedis*, one species, type: *E. cerradense* Pereira-Carv. & Dianese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Echinodiscus Etayo & Diederich 2000, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *E. lesdainii* (Vouaux) Etayo & Diederich, asexual morph unknown, lichenicolous, non-lichenized, Eurasia, see

Zhurbenko (2009; morphology, new species), Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable but lacking for the type species.

Echinomyces Rappaz 1987, *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *E. obesa* (Syd.) Rappaz, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Echinoplaca Fée 1825 (= *Arthotheliopsidomyces* Cif. & Tomas. 1953; = *Phallomyces* Bat. & Valle 1961; = *Pycnodermellina* Bat. & H. Maia 1957; = *Sporocybomyces* H. Maia 1967 *fide* Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, c. 40 species, type: *E. epiphylla* Fée, asexual morph unknown, lichenized, widespread (tropical), folicolous (but a few species are corticolous), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Echinosphaeria A.N. Mill. & Huhndorf 2004, *Helminthosphaeriaceae*, *Sordariomycetidae*, families *incertae sedis*, *Sordariomycetes*, seven species, type: *E. canescens* (Pers.) A.N. Mill. & Huhndorf, asexual morph unknown, saprobes, terrestrial, worldwide, see Bell and Mahoney (2008, 2008; new species), Declercq (2009; new species), Dhargalkar and Bhat (2009; new species), Bell (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miller et al. (2014; new species, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for type: SMH4666, TL5730.

Echinula Graddon 1977, *Hyaloscypheaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *E. asteriadiformis* Graddon, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Echria (N. Lundq.) Kruys, Huhndorf & A.N. Mill. 2015 (= *Arnium* sect. *Echria* N. Lundq. 1972 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, two species, type: *E. macrotheca* (P. Crouan & H. Crouan) Kruys, Huhndorf & A.N. Mill., saprobes, marine, USA, see Kruys et al. (2015; phylogeny), cultures and sequences are available.

Ecteinomyces Thaxt. 1903, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *E. trichopterophilus* Thaxt., asexual morph unknown, parasites, terrestrial, America, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2014b; the Netherlands), cultures and sequences are unavailable.

Ectendomeliola Hosag. & D.K. Agarwal 2006, *Ascomycota* genera *incertae sedis*, two species, type: *E. walsurae* Hosag. & D.K. Agarwal, asexual morph produced from the hyphae, on living leaves, terrestrial, worldwide, see Hosagoudar and Archana (2010a; new species, India), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded from *Meliolaceae*), holotype: HCIO 45237, cultures and sequences are unavailable.

Edenia M.C. González, Anaya, Glenn, Saucedo & Hanlin 2007, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *E. gomezompae* M.C. González, Anaya, Glenn, Saucedo & Hanlin, asexual morph unknown, endophytes, human pathogens, terrestrial, Pakistan, Quintana Roo, see Crous et al. (2009a; DNA, phylogeny), Hyde et al. (2013; outline, accepted as a genus in *Pleosporaceae*), Cui et al. (2013; DNA, human pathogens), Sun et al. (2013; new species, endophytic), Wijayawardene et al. (2014c; outline, accepted as a genus in *Pleosporaceae*), Ariyawansa et al. (2015c; phylogeny, excluded from *Pleosporaceae* and placed in *Phaeosphaeriaceae*), Pili et al. (2015; endophytes, Kenya, DNA), Keilor Rojas-Jimenez et al. (2016; endophytes, DNA, Costa Rica), holotype: MEXU 25346, cultures and sequences are available.

Edrudia W.P. Jord. 1980, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *E. constipans* (Nyl.) W.P. Jord., asexual morph unknown, lichenized, saxicolous, California, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Edwardiella Henssen 1986, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *E. mirabilis* Henssen, asexual morph unknown, lichenized, saxicolous, Marion Island, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Effertia Bartoli, Maggi & Persiani 1984, *Sordariaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *E. craspedoconidica* Bartoli, Maggi & Persiani, asexual morph virgariella-like, from soil, terrestrial, Ivory Coast, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: horto botanico Roma105S, cultures and sequences are unavailable.

Eiglera Hafellner 1984, *Eigleraceae*, *Acarosporales*, *Lecanoromycetes*, two species, type: *E. flava* (Hepp) Hafellner, asexual morph unknown, lichenized, saxicolous, Europe, North America, see Lumbsch and Huhndorf (2010; accepted as a genus in *Hymeneliaceae*, *Pertusariales*), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; the phylogenetic placement of *Eiglera* outside of the

Hymeneliaceae, sister to *Acarosporaceae*, needs to be confirmed, which would justify the reinstitution of the family *Eigleraceae* belonging to the *Acarosporales*, Lücking et al. (2017a; classification), sequences are available.

Eilidahlia S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2014, *Teloschistaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *E. dahlii* (Elix, S.Y. Kondr. & Kärnefelt) S.Y. Kondr., Kärnefelt, Elix, A. Thell, J. Kim, A.S. Kondr. & Hur, asexual morph unknown, lichenized, epiphytic, Australia, South Africa, see Kondratyuk et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Eiona Kohlm. 1968, *Ascomycota genera incertae sedis*, asexual morph unknown, one species, type: *E. tunicata* Kohlm., on wood often associated with sand, temperate, no sequence data, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), cultures and sequences are unavailable.

Elachopeltella Bat. & Cavalc. 1964, *Ascomycota genera incertae sedis*, two species, type: *E. sapindacearum* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Elaeodema Syd. 1922, *Ascomycota genera incertae sedis*, two species, type: *E. cinnamomi* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Elaphomyces Nees 1820 (= *Ascocleroderma* Clémencet 1932; = *Ceratogaster* Corda 1841; = *Ceraunium* Wallr. 1833; = *Hypogaeum* Pers. 1794; = *Lycoperdastrum* Haller ex Kuntze 1891; = *Phymatum* Chevall. 1826 *fide* Species Fungorum 2017), *Elaphomycetaceae*, *Elaphomycetales*, *Eurotiomycetes*, apparently polyphyletic, c. 78 species, type: *E. granulatus* Fr., terrestrial, hypogeous, forming mycorrhizas with conifers and broad-leaved trees, cosmopolitan, see Zhang and Minter (2011; new species), Castellano et al. (2011, 2012a, b, 2016; new species, phylogeny), Paz et al. (2012; new species), Buyck et al. (2016; phylogeny), cultures and sequences are available.

Elasticomyces Zucconi & Selbmann 2008, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *E. elasticus* Zucconi & Selbmann, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, Antarctica, see Selbmann et al. (2008; morphology, phylogeny), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), holotype and ex-type: CBS H-20177, CBS 122538, cultures and sequences are available.

Elattopycnis Bat. & Cavalc. 1964, *Ascomycota genera incertae sedis*, one species, type: *E. protii* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Elenkiniana S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2014, *Teloschistaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *E. gloriae* (Llimona & Werner) S. Y. Kondr., Kärnefelt, Elix, A. Thell, J. Kim, A. S. Kondratyuk et J.-S. Hur., asexual morph unknown, lichenized, saxicolous, Eurasia, see Kondratyuk et al. (2014; taxonomy), sequences are available.

Eleutherascus Arx 1971, *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, four species, type: *E. lectardii* (Nicot) Arx, asexual morph unknown, saprobes, terrestrial, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available for type: CBS 626.71.

Eleutheromycello Höhn. 1908 (= *Eleutheromyces* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Helicogoniaceae*, *Phacidiiales*, *Leotiomycetes*, one species, type: *E. mycophila* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eleutheromyces Fuckel 1870, *Helicogoniaceae*, *Phacidiiales*, *Leotiomycetes*, one species, type: *E. subulatus* (Tode) Fuckel, coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014a; DNA, phylogeny), epitype and ex-epitype: CBS H-12355, CBS113.86.

Elixia Lumbsch 1997, *Elixiaceae*, *Umbilicariales*, *Lecanoromycetes*, two species; type: *E. flexella* (Ach.) Lumbsch, lichenized, see Lumbsch and Huhndorf (2010; outline), Bendiksy and Timdal (2013; taxonomy), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are available, type specimen UPS-ACH-L-1286 (syntype).

Elletevera Deighton 1969, *Ascomycota genera incertae sedis*, one species, type: *E. parasitica* (Ellis & Everh.) Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Elliottinia L.M. Kohn 1979, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *E. kernerii* (Wettst.) L.M. Kohn, asexual morph unknown, saprobes, terrestrial,

Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny), cultures available for the type: KL402, specimen voucher TU:104529.

Ellisembia Subram. 1992, *Sordariomycetes* genera *incertae sedis*, c. 61 species, type: *E. coronata* (Fuckel) Subram., sexual reported for *E. adscendens* (Berk.) Subram., saprobes, lichenicolous (e.g. *E. lichenicola*), terrestrial, aquatic, worldwide, see Ma et al. (2008, 2010b; new species), McKenzie (2010; new species), Seifert et al. (2011; phylogeny), Delgado (2013; new species), Rajeshkumar et al. (2013; new species), Ren et al. (2013c; new species), Xia et al. (2015b; new species), Santa Izabel et al. (2013; new combinations), Su et al. (2016b; treated as synonym of *Sporidesmium*), cultures and sequences are available for few species (not from type material), but lacking for the type.

Ellisembiopsis T.S. Santa Izabel & Gusmão 2013, *Ascomycota* genera *incertae sedis*, two species, type: *E. brasiliensis* T.S. Santa Izabel & Gusmão, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, China, see Santa Izabel et al. (2013; morphology), holotype of the type: HUEFS 155102, cultures and sequences are unavailable.

Elmerinula Syd. 1934, *Dothideomycetes* genera *incertae sedis*, one species, type: *E. capnoides* Syd., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Elongatopedicellata J.F. Zhang, J.K. Liu, K.D. Hyde & Z.Y. Liu 2015, *Roussellaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *E. lignicola* J.F. Zhang, J.K. Liu, K.D. Hyde & Z.Y. Liu, asexual morph unknown, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type: MFLU 15-2717, MFLUCC 15-0642.

Elotespora R.F. Castañeda & Heredia 2010, *Ascomycota* genera *incertae sedis*, one species, type: *E. mexicana* R.F. Castañeda & Heredia, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Mexico, see Castañeda-Ruiz et al. (2010b; morphology), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: XAL CB883, cultures and sequences are unavailable, compare with *Artocarpomyces*, *Colemaniella* and *Glomerobolus*.

Elsinoë Racib. 1900 (= *Sphaceloma* de Bary 1874 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b; see Species Fungorum 2017 for additional synonyms), *Elsinoaceae*, *Myriangiales*, *Dothideomycetes*, c. 30 species (known by cultures), type: *E. canavaliae* Racib., asexual morph coelomycetous (*Sphaceloma*), pathogens, saprobes, terrestrial, Philippines, see Lumbsch and

Huhndorf (2010; outline), Chung (2011; pathogens), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Jayawardena et al. (2014; re-visit), Wijayawardene et al. (2014c; outline, nomenclature), Rossman et al. (2015b, 2016a; nomenclature, new combinations), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available.

Elytroderma Darker 1932, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *E. deformans* (Weir) Darker, asexual morph unknown, endophytes, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available for unidentified species.

Emarcea Duong, Jeewon & K.D. Hyde 2004, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, three species, type: *E. castanopsidicola* Duong, Jeewon & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, Taiwan, see Li et al. (2009b; new species, Taiwan), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Crous et al. (2015d; new species; DNA), Maharachchikumbura et al. (2015b, 2016b; outline), isotype: MRC DLA 008, HKUM 17498, CMU H224410, PDD 78748, ex-type: CBS and HKUCC 10344.

Emblemospora Jeng & J.C. Krug 1976, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *E. monotrema* Jeng & J.C. Krug, asexual morph unknown, from soil, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Embryonispora G.Z. Zhao 2014, *Ascomycota* genera *incertae sedis*, one species, type: *E. bambusicola* G.Z. Zhao, hyphomycetous, sexual morph unknown, saprobes, terrestrial, China, see Zhao et al. (2014a; taxonomy), cultures and sequences are unavailable.

Emericellopsis J.F.H. Beyma 1940 (= *Peyronellula* Malan 1952; = *Saturnomyces* Cain 1956 *fide* Species Fungorum 2017), *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, twelve species, type: *E. terricola* J.F.H. Beyma, asexual morph hyphomycetous, endophytes, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Grum-Grzhimaylo et al. (2013b; new species), Kirk et al. (2013; genus accepted), Hyde et al. (2016; new species, hyphomycetous asexual morph), sequences are available.

Emilmuelleria Arx 1986, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *E. spirotricha* (R.K. Benj.) Arx, asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Emodomelanlia Divakar & A. Crespo 2010, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *E. masonii* (Essl. & Poelt) Divakar & A. Crespo, asexual morph unknown, lichenized, saxicolous, Asian endemic (mainland China, India, Nepal and Taiwan), see Crespo et al. (2010a; phylogeny and taxonomy), Lücking et al. (2017a; classification), sequences are available.

Enantioptera Descals 1983, *Ascomycota* genera *incertae sedis*, two species, type: *E. tetra-alata* Descals, hyphomycetous, sexual morph unknown, in river foam, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Enarthromyces Thaxt. 1896, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *E. indicus* Thaxt., asexual morph unknown, parasites, saprobes, terrestrial, Asia, Africa, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Encephalographa A. Massal. 1854, *Melaspileaceae*, *Eremithallales*, *Dothideomycetes*, one species, type: *E. cerebrina* (DC.) A. Massal., asexual morph unknown, lichenized, saxicolous, Europe (southern), see Lumbsch and Huhndorf (2010; outline, doubtful position in *Arthoniomycetes*), Kirk et al. (2013; genus accepted), Ertz and Diederich (2015; phylogenetic position), Lücking et al. (2017a; classification), sequences and cultures are available; the placement of all species but *E. elisae* in *Encephalographa* should be reassessed.

Enchnoa Fr. 1849 (= *Massalongiella* Speg. 1880 *fide* Species Fungorum 2017), *Calosphaeriales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *E. lanata* (Fr.) Fr., asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Nitschiaceae*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Calosphaeriales* genera, *incertae sedis*), cultures and sequences are unavailable.

Enchylium (Ach.) Gray 1821 (see Species Fungorum 2017 for synonyms), *Collemataceae*, *Peltigerales*, *Lecanoromycetidae*, *Lecanoromycetes*, nine species, type: need typification, lichenized, see Otálora et al. (2013, 2014; taxonomy, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Encoelia (Fr.) P. Karst. 1871 (= *Cenangella* subgen. *Phaeangella* Sacc. 1889; = *Peziza* trib. *Encoelia* Fr. 1822; = *Phaeangella* (Sacc.) Massee 1895; = *Phibalis* Wallr. 1833 *fide* Species Fungorum 2017), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, 40 species, type: *E. furfuracea*

(Roth) P. Karst., asexual morph unknown, endophytes, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; morphology, ultrastructure, DNA, phylogeny), cultures and sequences are available.

Enceliopsis Nannf. 1932 (= *Neogodronia* Schläpf.-Bernh. 1969 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, four species, type: *E. rhododendri* (Ces. ex Rabenh.) Nannf., asexual morph unknown, endophytes, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Helotiaceae*), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny), cultures available for type: KL118.

Endobotrya Berk. & M.A. Curtis 1874, *Ascomycota* genera *incertae sedis*, one species, type: *E. elegans* Berk. & M.A. Curtis, coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline, morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Endobotryella Höhn. 1909, *Ascomycota* genera *incertae sedis*, one species, type: *E. oblonga* (Fuckel) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Endocalyx Berk. & Broome 1876, *Apiosporaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *E. thwaitesii* Berk. & Broome, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Capdet and Romero (2010; Argentina), Seifert et al. (2011; morphology), Wijayawardene et al. (2012; outline, accepted as *Ascomycota* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Endocarpon Hedw. 1788 (= *Dermatocarpon* W. Mann 1825; = *Endopyreniomyces* E.A. Thomas 1939; = *Leightonia* Trevis. 1853; = *Paracarpidium* Müll. Arg. 1883; = *Pyrenothamnia* Tuck. 1883; = *Pyrenothamniomyces* Cif. & Tomas. 1953; = *Rhodocarpon* Lönnr. 1858 *fide* Species Fungorum 2017), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, c. 70 species, type: *E. pusillum* Hedw., asexual morph coelomycetous, terrestrial and aquatic (maritime and freshwater), lichenized, occurring on rocks, soil and bark, cosmopolitan, see Gueidan et al. (2009; phylogeny, outline), Prieto et al. (2012; phylogeny), Joshi and Hur (2013a, b; new species), Lücking et al. (2017a; classification), sequences are available.

Endocena Cromb. 1876, *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type:

E. informis Cromb., asexual morph unknown, lichenized, terricolous, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Endococcus Nyl. 1855 (= *Discothecium* Zopf 1897; = *Pseudoendococcus* Marchand 1896; = *Sorothelia* Körb. 1865 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, 34 species, type: *E. rugulosus* (Borrer ex Leight.) Nyl., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; outline, genus accepted), Kocourková and Knudsen (2011; new species), Zhurbenko et al. (2012; new species), Jayasiri et al. (2016a; re-visited), a culture and sequence is available.

Endocolium Syd. 1937, *Ascomycota* genera *incertae sedis*, one species, type: *E. costaricanum* Syd., asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Endoconidioma Tsuneda, Hambl. & Currah 2004, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, one species, type: *E. populi* Tsuneda, Hambl. & Currah, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Endoconidiophora Münch 1907, *Ceratocystidaceae*, *Microascales*, *Sordariomycetes*, one species, type: *E. coerulescens* Münch, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012; outline, accepted as *Ascomycota* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Endoconospora Gjaerum 1971, *Ascomycota* genera *incertae sedis*, two species, type: *E. cerastii* Gjaerum, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Norway, India, see Wijayawardene et al. (2012; outline, accepted as *Ascomycota* genera *incertae sedis*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Endocoryneum Petr. 1922, ?*Didymellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *E. loculosum* (Sacc.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Norway, India, see Wijayawardene et al. (2012, 2016b; outline, accepted as *Ascomycota* genera *incertae sedis* and *Didymellaceae* respectively), Kirk et al. (2013; genus accepted), cultures and sequences are available for *E. festucae* Wijaya. et al. (MFLUCC 14–0461).

Endothiora Petr. 1929, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, one species, type: *E. sydowiana* Petr., asexual morph unknown, saprobes, terrestrial, former

USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Endogenospora R.F. Castañeda, O. Morillo & Minter 2010, *Ascomycota* genera *incertae sedis*, one species, type: *E. aspectabilis* R.F. Castañeda, O. Morillo & Minter, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Venezuela, see Castañeda-Ruiz et al., (2010b; morphology), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype: INIFAT C09/74, cultures and sequences are unavailable, compare with *Ascoconidium*, *Sporoschisma* and *Sporoschismopsis*.

Endohyalina Marbach 2000, *Physciaceae*, *Lecanorales*, *Lecanoromycetidae*, type: *E. rappii* (Imshaug ex R.C. Harris) Marbach, ten species, three varieties, asexual morph unknown, lichenized, terrestrial, corticolous, lignicolous, sometimes obligately lichenicolous, Australia, tropical to subtropical, see Giralt et al. (2010; taxonomy, Elix and Kantvilas (2015; new records, new taxa), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Endomelanconiopsis E.I. Rojas & Samuels 2008, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, two species, type: *E. endophytica* E.I. Rojas & Samuels, coelomycetous, saprobes, terrestrial, South America, see Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Phillips et al. (2013; taxonomy, new species, phylogeny), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Endomelanconium Petr. 1940, *Ascomycota* genera *incertae sedis*, four species, type: *E. pini* (Corda) Petr., coelomycetous, sexual morph unknown, endophytes, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2016b, 2017; morphology; outline), cultures and sequences are unavailable.

Endomeliola S. Hughes & Piroz. 1994, *Meliolaceae*, *Meliolales*, *Sordariomycetes*, one species, type: *E. dingleyae* S. Hughes & Piroz., asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: DAOM 167533, cultures and sequences are available for PDD:98304 (unpublished).

Endomyces Reess 1870, *Saccharomycetales* genera *incertae sedis*, *Saccharomycetes*, four species, type: *E. decipiens* (Tul. & C. Tul.) Reess, asexual reproduction is by pseudohyphae or septate hyphae or blastoconidia, parasite of mushrooms, grow on various mushrooms,

worldwide, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Endophoma Tsuneda & M.L. Davey 2011, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *E. elongata* Tsuneda & M.L. Davey, coelomycetous, sexual morph unknown, saprobes (soil of bat cave), terrestrial, Canada, see Tsuneda et al. (2011a; taxonomy, DNA, phylogeny), Chen et al. (2015b; phoma-like taxa in *Didymellaceae*), holotype and ex-type: UAMH 11216.

Endophragmiella B. Sutton 1973, *Helminthosphaeriaceae*, *Sordariales*, *Sordariomycetes*, c. 70 species, type: *E. pallescens* B. Sutton, hyphomycetous, Manitoba, see Seifert et al. (2011; morphology), Ma et al. (2012c; new species), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Endophragmiopsis M.B. Ellis 1966, *Ascomycota* genera *incertae sedis*, one species, type: *E. pirozynskii* M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Endoplacodium Petr. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *E. nigrescens* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Iran, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Endoramularia Petr. 1923 (= *Endocladis* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *E. ulmi* Petr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Endoscypha Syd. 1924, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *E. perforans* Syd., asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available PDD:102231 (unpublished).

Endosporoideus W.H. Ho, Yanna, K.D. Hyde & Goh 2005, *Ascomycota* genera *incertae sedis*, one species, type: *E. pedicellatus* W.H. Ho, Yanna, K.D. Hyde & Goh, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Hong Kong, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), cultures and sequences are unavailable.

Endostilbum Malençon 1964, *Ascomycota* genera *incertae sedis*, one species, type: *E. cerasi* (Bourdot & Galzin) Malençon, hyphomycetous, sexual morph

unknown, saprobes, terrestrial, Europe, Morocco, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), cultures and sequences are unavailable.

Endothia Fr. 1849 (= *Calopactis* Syd. & P. Syd. *fide* Rossman et al. 2015a), *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *E. gyrosa* (Schwein.) Fr., asexual morphs unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2015a; nomenclature), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Endotryblidium Petr. 1959, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *E. insculptum* (Cooke) Petr., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), cultures and sequences are unavailable.

Endoxyla Fuckel 1871 (= *Eutypopsis* P. Karst. 1878; = *Lentomitella* Höhn. 1906 [1905]; = *Linostomella* Petr. 1925 *fide* Species Fungorum 2017), *Boloniaceae*, *Boloniiales*, *Sordariomycetes*, ten species, type: *E. operculata* (Nitschke) Sacc., asexual morphs unknown, saprobes, terrestrial, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Untereiner et al. (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Endoxylina Romell 1892 (= *Pogonospora* Petr. 1957 [1956] *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, seven species, type: *E. stellulata* Romell, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Vasilyeva (2010; new species; Russia), Kirk et al. (2013; genus accepted), Untereiner et al. (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Endozythia Petr. 1959, *Ascomycota* genera *incertae sedis*, one species, type: *E. moravica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, former Yugoslavia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Enerthidium Syd. 1939, *Ascomycota* genera *incertae sedis*, one species, type: *E. canarii* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Wijayawardene et al. (2012, 2017; outline, morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Engelhardtia A. Funk 1973, *Ascomycota* genera *incertae sedis*, one species, type: *E. alba* A. Funk, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Englerodothis Theiss. & Syd. 1915 (= *Dothophaeis* Clem. 1931 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, three species, type: *E. kilimandscharica* (Henn.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Parmulariaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Parmulariaceae*), Hyde et al. (2013; review, keys, excluded from *Parmulariaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences unavailable.

Engleromyces Henn. 1900 (≡ *Stromne* Clem. 1909; = *Cerillum* Clem. 1931; *Colletomanginia* Har. & Pat. 1906 *fide* Species Fungorum 2017; Wendt et al. 2017, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *E. goetzei* Henn., asexual morph unknown, saprobes, terrestrial, Africa, Asia, see Lumbsch and Huhndorf (2010; outline), Whalley et al. (2010; new species, China), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Englerula Henn. 1904 (= *Anatexis* Syd. 1928 *fide* Species Fungorum 2017), *Englerulaceae*, *Dothideomycetes* families *incertae sedis*, six species, type: *E. macarangae* Henn., asexual morph coelomycetous, saprobes, epiphytic, terrestrial, Africa, India, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Englerulaceae*), Dai et al. (2014c; morphology), cultures and sequences are unavailable.

Engyodontium de Hoog 1978, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *E. parvisporum* (Petch) de Hoog, hyphomycetous, sexual morph unknown, saprobes, on insects and spiders, cosmopolitan, see Seifert et al. (2011; morphology), Jaklitsch et al. (2016a; classification), Maharachchikumbura et al. (2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Enigmotrema Lücking 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *E. rubrum* Lücking, lichenized, see Sipman et al. (2012; taxonomy), Lücking et al. (2013, 2016b; phylogeny, classification), Lücking (2014; taxonomy), sequences are available.

Enridescalsia R.F. Castañeda & Guarro 1998, *Ascomycota* genera *incertae sedis*, one species, type: *E. speciosa* R.F. Castañeda & Guarro, hyphomycetous, sexual morph unknown, saprobes, aquatic (fresh water), Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: INIFAT C95/96, cultures and sequences are unavailable, compare with *Xylomyces* and *Gonatobotrys*.

Enterocarpus Locq.-Lin. 1977, *Microascaceae*, *Microascales*, *Sordariomycetes*, two species, type: *E. uniporus* Locq.-Lin., asexual morph unknown, coprophilous, terrestrial, Sahara, Italy, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Enterodictyon Müll. Arg. 1892 (= *Enterodictyonomyces* Cif. & Tomas. 1953; = *Enterodictyum* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *E. indicum* Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Enterographa Fée 1824 (= *Arthonaria* Fr. 1825; = *Chiodectionomyces* Cif. & Tomas. 1953; = *Platygramma* Leight. 1854; = *Stigmatidium* G. Mey. 1825 *fide* Species Fungorum 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, ±60 species, type: *E. crassa* (DC.) Fée, sexual morph with punctiform or rounded to lirelliform ascocarps, lichenized, rarely lichenicolous, terrestrial, world-wide with a preference for (sub)tropical regions, see Ertz and Tehler (2011; phylogeny), Lumbsch and Huhndorf (2010; outline), Aptroot et al. (2013a; new species), Kirk et al. (2013; genus accepted), Ertz et al. (2015c, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Enthallopycnidium F. Stevens 1925, *Ascomycota* genera *incertae sedis*, one species, type: *E. gouldiae* F. Stevens, coelomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Entoderma Hanula, Andreadis & M. Blackw. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *E. colletosporium* Hanula, Andreadis & M. Blackw., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Entodesmium Riess 1854, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, eight species, type: *E. rude* Riess, sexual morph unknown, saprobes, terrestrial, temperate, see Schoch et al. (2009; DNA,

phylogeny), Hyde et al. (2013; outline, accepted as a genus in *Phaeosphaeriaceae*), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*, phylogeny), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015a; new species, Russia, DNA, phylogeny), Tibpromma et al. (2017b; new species), cultures available for the type: CBS 650.86.

Entoleuca Syd. 1922, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, three species, type: *E. callimorpha* Syd., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Entonaema Möller 1901, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, six species, type: *E. liquescens* Möller, asexual morph unknown, terrestrial, saprobes, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available.

Eoaleurina Korf & W.Y. Zhuang 1986, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *E. foliicola* (Henn.) Korf & W.Y. Zhuang, asexual morph unknown, saprobes, terrestrial, Central and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eopyrenula R.C. Harris 1973, *Ascomycota* genera *incertae sedis*, three species, type: *E. leucoplaca* (Wallr.) R.C. Harris, sexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; outline, accepted as a genus in *Dacampiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as a genus in *Dacampiaceae*), cultures and sequences are unavailable.

Eosphaeria Höhn. 1917 (= *Herminia* R. Hilber 1979 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *E. uliginosa* (Kunze) Höhn., asexual morph hyphomycetous, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Eoterfezia G.F. Atk. 1902, *Eoterfeziaceae*, *Ascomycota* families *incertae sedis*, type: *E. parasitica* G.F. Atk., six species, asexual morph unknown, terrestrial, saprobes, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Epaphroconidia Calat. & V. Atienza 1995, *Ascomycota* genera *incertae sedis*, one species, type: *E. hawksworthii* Calat. & V. Atienza, coelomycetous, sexual morph unknown, lichenicolous, terrestrial, Spain, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ephebe Fr. 1825 (= *Ephebeia* Nyl. 1875; = *Ephebomyces* Cif. & Tomas. 1953; = *Spilonematopsis* A.E. Dahl 1950 *fide* Species Fungorum 2017), *Lichinaceae*, *Lichinales*, *Lichinomycetidae*, 13 species, type: *E. lanata* (L.) Vain., asexual morph unknown, lichenized, terrestrial, corticolous, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Ephelidium C.W. Dodge & E.D. Rudolph 1955, *Ascomycota* genera *incertae sedis*, one species, type: *E. heardense* C.W. Dodge & E.D. Rudolph, coelomycetous, sexual morph unknown, saprobes, terrestrial, Antarctica, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Epibelonium E. Müll. 1963, *Saccardiaceae*, *Ascomycota* families *incertae sedis*, one species, type: *E. gaeumannii* E. Müll., asexual morph unknown, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Epibryon Döbbeler 1978, *Epibryaceae*, *Chaetothyriales*, *Eurotiomycetes*, c. 30 species, type: *E. plagiochilae* (Gonz. Frag.) Döbbeler, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; DNA, symbioses), Hyde et al. (2013; outline, accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Gueidan et al. (2014a; phylogeny), cultures and sequences are available.

Epichloë (Fr.) Tul. & C. Tul. 1865 (= *Cordyceps* subgen. *Epichloë* Fr. = *Neotyphodium* Glenn, C.W. Bacon & Hanlin 1996 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, six species, type: *E. typhina* (Pers.) Tul. & C. Tul., sexual morph unknown, saprobes, endophytes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Young et al. (2014; endophytes), Hettiarachchige et al. (2015; endophytes), Maharanachikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are available.

Epicladonia D. Hawksw. 1981, *Leotiomycetes* genera *incertae sedis*, four species, type: *E. sandstedei* (Zopf) D. Hawksw., on lichens, sexual morph unknown, see Kirk et al. (2013; genus accepted), Pino-Bodas et al. (2017; new

species, phylogeny), cultures and sequences are unavailable, topotype material of type: B (Krypt. Exs. Vindob. No. 1330b).

Epiclinium Fr. 1849, *Ascomycota* genera *incertae sedis*, two species, type: *E. pezizoideum* (Schwein.) Fr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Epicoccospora Budathoki & S.K. Singh 1995, *Ascomycota* genera *incertae sedis*, two species, type: *E. nepalensis* Budathoki & S.K. Singh, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Nepal, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Epicoccum Link 1816 (= *Cerebella* Ces. 1851; = *Clathrococcum* Höhn. 1911; = *Malacharia* Fée 1843; = *Paratrichaegum* Faurel & Schotter 1966 [1965]; = *Thyrococcum* R.E. Buchanan 1911 *fide* Species Fungorum 2017), *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, c. ten species, type: *E. nigrum* Link, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Aveskamp et al. (2010, 2010; DNA, new species, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c; outline, accepted as a genus in *Didymellaceae*), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), Kirk et al. (2013; genus accepted), Chen et al. (2015b; taxonomy, new species, DNA, phylogeny), Thambugala et al. 2017b, new species, phylogeny), ex-type cultures available for type: CBS 173.73.

Epicoleosporium Videira & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *E. ramularioides* Videira, H.D. Shin & Crous, hyphomycetous, sexual morph unknown, plant pathogenic, see Videira et al. (2016; morphology, phylogeny), holotype KUS-F19603, isotype CBS H-22542, culture ex-type: CBS 141103 = CPC 10672.

Epicrea Petr. 1950, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, six species, type: *E. insignis* Petr., asexual morph unknown, saprobes, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Epidermophyton Sabour. 1907 (= *Epidermomyces* Loeffler 1983 *fide* Species Fungorum 2017), *Arthrodermataceae*, *Onygenales*, *Eurotiomycetes*, one species: *E. floccosum* (Harz) Langeron & Miloch., hyphomycetous, terrestrial, on human skin, one of the main etiological agents of dermatophytosis worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), neotype strain of the type: =CBS 230.76, sequences are available from de Hoog et al. (2017).

Epigloeoa Zukal 1889 (= *Epigloeomyces* Cif. & Tomas. 1957; = *Vorarlbergia* Grummann 1969 *fide* Species Fungorum 2017), *Epigloeaceae*, *Ascomycota* families *incertae sedis*, 13 species, type: *E. bactrospora* Zukal, on algae, Europe, Antarctica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Epilichen Clem. 1909, *Rhizocarpaceae*, *Rhizocarpales*, *Lecanoromycetes*, three species; type: *E. scabrosus* (Ach.) Clem., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable, type specimen UPS-ACH-L-130 (isotype),

Epinephroma Zhurb. 2012, *Ascomycota* genera *incertae sedis*, two species, type: *E. kamchatica* Zhurb. & Stepanch., coelomycetous, lichenicolous, terrestrial, Russia, see Zhurbenko et al. (2012; morphology), cultures and sequences are unavailable.

Epiphloea Trevis 1880 (= *Amphidium* Nyl. 1876 [1877]; = *Latelia* Zahlbr. 1926 *fide* Species Fungorum 2017), *Heppiaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *E. terrena* (Nyl.) Trevis., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline). Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Episclerotium L.M. Kohn 1984, *Helotiales* genera *incertae sedis*, *Leotiomyces*, two species, type: *E. sclerotiorum* (Rostr.) L.M. Kohn, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Episphaerella Petr. 1924, *Pseudoperisporiaceae*, *Dothideomycetes*, families *incertae sedis*, six species, type: *E. manihotis* (Henn.) Petr., asexual morph unknown, saprobes, terrestrial, USA, Africa, China, see Lumbsch and Huhndorf (2010; outline), Zhao and Zhao (2012; new species, China), Hyde et al. (2013; outline, accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as a genus in *Pseudoperisporiaceae*), cultures and sequences are available for *Episphaerella dodonaeae* PDD:104466 (unpublished).

Episporogoniella U. Braun 1994, *Ascomycota* genera *incertae sedis*, one species, type: *E. muscigena* (Speg.) U. Braun, hyphomycetous, sexual morph unknown, terrestrial, muscicolous, South America, see Seifert et al. (2011), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of type: LPS 12764, cultures and sequences are unavailable.

Epistigme Syd. 1924, *Ascomycota* genera *incertae sedis*, two species, type: *E. nidulans* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Epithamnolia Zhurb. 2012, *Ascomycota* genera *incertae sedis*, one species, type: *E. karatyginii* Zhurb., sexual morph unknown, lichenicolous, terrestrial, Canada, see Zhurbenko (2011; morphology), cultures and sequences are unavailable.

Epithyrium (Sacc.) Trotter 1931 (= *Biatoridina* Schczedr. 1964; = *Coniothyrium* subgen. *Epithyrium* Sacc. 1892 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *E. resinae* (Sacc. & Berl.) Trotter, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Wijayawardene et al. (2012, 2016b, 2017; outline, morphology), Kirk et al. (2013; did not list), cultures and sequences are unavailable.

Equisetcola Abdelsalam, Tibpromma, Wanasa. & K.D. Hyde 2016, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *E. fusispora* Abdelsalam, Tibpromma, Wanasinghe & K.D. Hyde, asexual morph unknown, saprobes, Italy, see Abd-Elsalam et al. (2016; taxonomy), holotype: MFLU 14-0702, cultures and sequences are available.

Eremascus Eidam 1877, *Eremascaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *E. albus* Eidam, asexual morph absent in the type species, but hyphomycetous in *E. terrestris* Asthana & Mahmud (Asthana & Mahmud 1944), terrestrial, saprobes, usually xerophilic, occurring on substrates with low water and high carbohydrate content, such as dried fruits, honey, jams, mustard and others, also reported from soil, Argentina, Australia, India, Western Europe, see Gueidan et al. (2008; phylogeny), Pitt and Hocking (2009; morphology, food spoilage), Wynns (2015; phylogeny, proposal of a segregate genus), material of the type species is lost and neotypification is necessary.

Eremastrella S. Vogel 1955, *Lecideaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *E. tobleri* S. Vogel, asexual morph unknown, lichenized, terrestrial, see Ekman and Blaaid (2011; molecular sequence), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017b; classification), sequences are available.

Eremiomyces Trappe & Kagan-Zur 2005, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *E. echinulatus* (Trappe & Marasas) Trappe & Kagan-Zur, asexual morph unknown, saprobes, from soil, terrestrial, South Africa, Spain, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Pezizaceae*), Alvarado et al. (2011b; new species, Spain, DNA, placed in *Tuberaceae*), cultures and sequences are available.

Eremomyces Malloch & Cain 1971 (= *Pithoascina* Valmaseda, A.T. Martínez & Barrasa 1987 *fide* Species Fungorum 2017), *Eremomycetaceae*, *Dothideomycetes* genera *incertae sedis*, two species, type: *E. bilateralis* Malloch & Cain, asexual morph hyphomycetous, saprobes, terrestrial, North America, Netherlands, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for type.

Eremothecella Syd. & P. Syd. 1917 (= *Nematidia* Stirz. 1879 [1878] *fide* Species Fungorum 2017), *Arthoniaceae*, *Arthoniales*, *Arthoniomycetidae*, type: *E. calamicola* Syd. & P. Syd., eleven species, asexual morph unknown, lichenized, terrestrial, mainly tropical, see Lumbsch and Huhndorf (2010; outline). Kirk et al. (2013; genus accepted), Cáceres and Aptroot (2014; taxonomy), Lücking et al. (2017a; classification), molecular and culture data unavailable.

Eremothecium Borzi emend. Kurtzman 1995 (= *Ashbya* Guillerm. 1928; = *Crebrothecium* Routien 1949; = *Spermophthora* S.F. Ashby & W. Nowell 1926 *fide* Species Fungorum 2017), *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, six species, type: *E. cymbalariae* Borzí, asexual reproduction is by budding cells are absent or present, and when present, budding is multilateral on a narrow base, enteroarthric conidia are infrequently produced by one species, pseudohyphae and true hyphae are generally present, plant pathogens, cotton, hazel nuts, tomatoes, mustard, insects, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Erikssonia Penz. & Sacc. 1898 (= *Paidania* Racib. 1909; = *Periaster* Theiss. & Syd. 1917 [1916]; = *Rinia* Penz. & Sacc. 1902 [1901] *fide* Species Fungorum 2017), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, five species, type: *E. pulchella* Penz. & Sacc., sexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Erikssonopsis M. Morelet 1971 (= *Grovesiella* B. Erikss. 1970 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *E. ericae* (Fr.) M. Morelet, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Erinacellus T. Sprib., Muggia & Tønsberg 2014, *Peltigerales* genera *incertae sedis*, *Lecanoromycetes*, two species, type: *E. dendroides* (Henssen) T. Sprib., Tønsberg & Muggia, lichenized, see Spribille et al. (2014b;

taxonomy), Lücking et al. (2017a; classification), sequences are available.

Eriocercospora Deighton 1969, *Dothideomycetes* genera *incertae sedis*, two species, type: *E. balladynae* (Hansf.) Deighton, hyphomycetous, sexual morph unknown, saprobes, mycoparasites, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eriocercosporella Rak. Kumar, A.N. Rai & Kamal ex U. Braun 1998, *Ascomycota* genera *incertae sedis*, one species, type: *E. indica* Rak. Kumar, A.N. Rai & Kamal ex U. Braun, hyphomycetous, sexual morph unknown, terrestrial, foliicolous, plant pathogenic, Asia, see Seifert et al. (2011), Wijayawardene et al. (2012, 2014c; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: K(M) IMI 302747.

Erioderma Fée 1825 (= *Malmella* C.W. Dodge 1933 *fide* Species Fungorum 2017), *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 40 species, type: *E. polycarpum* Fée, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Eriomycopsis Speg. 1911, *Ascomycota* genera *incertae sedis*, 13 species, type: *E. bomplandi* Speg., hyphomycetous, sexual morph unknown, fungicolous, terrestrial, worldwide, see Seifert et al. (2011), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eriopezia (Sacc.) Rehm 1892 (= *Tapesia* subgen. *Eriopezia* Sacc. 1889; = *Tapesiella* Velen. 1939 *fide* Species Fungorum 2017), *Arachnopezizaceae*, *Helotiales*, *Leotiomycetes*, 30 species, type: *E. caesia* (Pers.) Rehm, asexual morph unknown, saprobes, terrestrial, Europe, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Erioscyphella Kirschst. 1938, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, c. ten species, type: *E. longispora* (P. Karst.) Kirschst., sexual morph unknown, saprobes, cosmopolitan, see Kirk et al. (2013; treated as a synonym of *Lachnum* Retz.), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Eriosphaeria Sacc. 1875 (= *Cantharosphaeria* Thaxt. 1920; = *Eriosphaerella* Höhn. 1906; = *Malacosphaeria* Syd. 1924; = *Penzigina* Kuntze 1891 *fide* Species Fungorum 2017), *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, six species, type: *E. vermicularia* (Nees) Sacc., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura

et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Eriospora Berk. & Broome 1850, *Ascomycota* genera *incertae sedis*, one species, type: *E. leucostoma* Berk. & Broome, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, South America, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Eriosporella Höhn. 1916 (= *Pseuderiospora* Keissl. 1924 *fide* Species Fungorum 2017), *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *E. calami* (Niessl) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Thailand, see Wijayawardene et al. (2012, 2014c; outline), Kirk et al. (2013; genus accepted), Dai et al. (2014e; new species, DNA, phylogeny), cultures and sequences are available for *E. bambusicola* D.Q. Dai et al. (ex-type MFLUCC 11-0436).

Eriosporopsis Petr. 1947, *Ascomycota* genera *incertae sedis*, one species, type: *E. albida* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eriothyrium Speg. 1887, *Ascomycota* genera *incertae sedis*, one species, type: *E. dubiosum* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Erispora Pat. 1922, *Ascomycota* genera *incertae sedis*, one species, type: *E. parasitica* Pat., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, holotype is lost.

Ernakulamia Subram. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *E. cochinensis* (Subram.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South East Asia, see Capdet and Romero (2010; Argentina), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Erysiphe R. Hedw. ex DC. 1805: Fr. 1829 emend. Braun & Takamatsu 2000 (see Species Fungorum 2017 for synonyms), *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, 378 species in *Erysiphe* and 72 species in *Pseudoidium*, type: *E. polygoni* DC., hyphomycetous asexual morphs, biotrophic, obligate plant pathogens, terrestrial, worldwide, see Braun and Cook (2012; taxonomy), Takamatsu et al. (2015a, b; phylogeny), lectotype of type: G 00122108.

Erythrocarpon Zukal 1886, *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, six species, type:

E. microstomum Zukal, asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Erythrodecon G. Thor 1991, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *E. granulatum* (Mont.) G. Thor, lichenized, see Lumbsch and Huhndorf (2010; outline), Ertz and Tehler (2011, phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Erythrogloeum Petr. 1953, *Erythrogloeaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *E. hymenaeae* Gonz. Frag. & Cif. ex Petr., coelomycetous, sexual morph unknown, biotrophic, terrestrial, Costa Rica, South and Central America, see Crous et al. (2012d; DNA, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), epitype and ex-epitype of the type: CBS H-20937, CPC 18819 = CBS 132185.

Erythromada Huhndorf, A.N. Mill., F.A. Fernández & Lodge 2005, *Chaetosphaeriales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *E. lanciospora* Huhndorf, A.N. Mill., F.A. Fernández & Lodge, asexual morphs unknown, saprobes, terrestrial, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Eschatogonia Trevis. 1853, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, seven species; type: *E. montagnei* Trevis., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), isotype of the type: UPS-L-129880, sequences are unavailable.

Escovopsioides H.C. Evans & J.O. Augustin 2012, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *E. nivea* H.C. Evans & J.O. Augustin, asexual morph hyphomycetous, on nest of leaf-cutting ants, terrestrial, Brazil, see Augustin et al. (2012; morphology, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), holotype IMI 501181; isotype DOA627-VIC 31754, cultures and sequences are available.

Escovopsis J.J. Muchovej & Della Lucia 1990 (= *Phialocladus* Kreisel 1972 *fide Species Fungorum* 2017), *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *E. weberi* J.J. Muchovej & Della Lucia, hyphomycetous, sexual morph unknown, on nest of attine ants, terrestrial, worldwide, see Augustin et al. (2012; morphology, DNA, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Masiulionis et al. (2015; new species), Meirelles

et al. (2015; new species), cultures and sequences are available.

Esdipatilia Phadke 1981, *Ascomycota* genera *incertae sedis*, one species, type: *E. indica* Phadke, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: AMH4616, cultures and sequences are unavailable.

Esfandiariomyces Ershad 1985 (= *Esfandiaria* Petr. 1955 *fide Species Fungorum* 2017), *Sordariomycetes*, genera *incertae sedis*, one species, type: *E. insignis* (Petr.) Ershad, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Esslingeriana Hale & M.J. Lai 1980, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *E. idahoensis* (Essl.) Hale & M.J. Lai, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Esteya J.Y. Liou, J.Y. Shih & Tzean 1999, *Ascomycota* genera *incertae sedis*, one species, type: *E. vermicola* J.Y. Liou, J.Y. Shih & Tzean, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Taiwan, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Etayoia Diederich & Ertz 2014, *Phaeococcymycetaceae*, *Lichenostigmatales*, *Arthoniomycetes*, one species, type: *E. trypethelii* (Flakus & Kukwa) Diederich & Ertz, both sexual and asexual morphs stromatic, lichenicolous, non-lichenized, on corticolous lichens, worldwide, see Ertz et al. (2014b; morphology, phylogeny), cultures and sequences are available.

Etheiophora Kohlm. & Volk.-Kohlm. 1989, *Etheiophoraceae*, *Torpedosporales*, *Sordariomycetes*, three species, type: *E. bijubata* Kohlm. & Volk.-Kohlm., saprobes, marine, Atlantic Ocean, Pacific Ocean, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for type.

Euacanthe Theiss. 1917, *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, two species, type: *E. usambarensis* (Rehm) Theiss., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Mugambi and Huhndorf (2010; DNA), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for *E. foveolata* (GKM1221 *fide* Mugambi and Huhndorf 2010), treated as a

synonym of *Acanthonitschkea* Speg. in Species Fungorum (2017).

Euantennaria Speg. 1918 (= *Aithalomyces* Woron. 1926; = *Ophiocapnodium* Speg. 1918; = *Phaeocapnias* Cif. & Bat. 1963; = *Skoteinospora* Bat. 1962 *fide* Species Fungorum 2017), *Euantennariaceae*, *Dothideomycetes* families *incertae sedis*, six species, type: *E. tropicicola* Speg., saprobes, parasitic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Eubelonis Höhn. 1926, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *E. albosanguinea* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eucalyptostroma Crous & M.J. Wingf. 2016, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *E. eucalypti* Crous & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type of the type: CBS H-22871, CPC 28764.

Eucantharomyces Thaxt. 1895, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 26 species, type: *E. atrani* Thaxt., asexual morph unknown, parasites, saprobes, terrestrial, worldwide, see Rossi and Ponziani (2008; new species), Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Eucasperhaeria Crous 2007, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *E. capensis* Crous, asexual morph coelomycetous, saprobes, terrestrial, South Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Azeem et al. (2015; DNA), Crous et al. (2016a; new species, phylogeny), holotype and the ex-type strains of the type: CBS-H19764, CPC 12796 = CBS 120028.

Euceramia Bat. & Cif. 1962, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *E. palmicola* Bat. & Cif., asexual morph unknown, epiphytic, epiphylloous, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Euceratomyces Thaxt. 1931, *Euceratomyctaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *E. terrestris* Thaxt., asexual morph unknown, parasites, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species), cultures and sequences are unavailable.

Eudarluca Speg. 1908 (= *Cryptocrea* Petr. 1937; = *Parabotryon* Syd. 1926; = *Xenodimerium* Petr. 1947 *fide*

Species Fungorum 2017), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, c. five species, type: *E. australis* Speg., asexual morph unknown, hyperparasitic associated with rust species (but not necessarily parasitic), terrestrial, worldwide, all species have sequence data, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; phylogeny), Wijayawardene et al. (2014c; outline), isotype of the type: S-F11420, cultures and sequences are available.

Eudimeriolum Speg. 1912 (= *Astiothyrium* Bat. 1964; = *Ontostheca* Bat. 1963 *fide* Species Fungorum 2017), *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, eight species, type: *E. elegans* Speg., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Euepixylon Füisting 1867 (= *Fuckelia* (Nitschke ex Sacc.) Cooke 1869 *fide* Species Fungorum 2017), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *E. udum* (Pers.) Læssøe & Spooner, asexual morph unknown, saprobes, terrestrial, Europe, Eastern Russia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Eugeniella Lücking, Sérus. & Kalb 2008, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, nine species, type: *E. psychotriae* (Müll. Arg.) Lücking, Sérus. & Kalb, lichenized, see Lücking (2008; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Euhaplomyces Thaxt. 1901, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *E. aencyphori* Thaxt., asexual morph unknown, parasites, saprobes, terrestrial, worldwide, see Rossi and Ponziani (2008; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eumela Syd. 1925, *Pseudoperisporiaceae*, *Dothideomycetes*, families *incertae sedis*, one species, type: *E. chiococcae* Syd., asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Eumonoicomycetes Thaxt. 1901, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two or three species, type: *E. californicus* Thaxt., asexual morph unknown, parasites, terrestrial, worldwide, see Rossi and

Ponziani (2008; new species), Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Euopsis Nyl. 1875, *Harpidiaceae*, *Lichinales*, *Lichenomycetes*, two species, type: *E. granatina* (Sommerf.) Nyl., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017b; classification), sequences are available.

Eupelte Syd. 1924 (= *Maurodothina* G. Arnaud ex Piroz. & Shoemaker 1970; = *Pirozynskia* Subram. 1972 *fide* Rossman et al. 2016a), *Dothideomycetes* genera *incertae sedis*, four species, type: *E. amicta* Syd., asexual morph (=sporidesmium-like, *fide* Sivanesan 1984), epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, treated as a doubtful genus in *Asterinales*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), Rossman et al. (2016a; nomenclature), holotype of type: IMI 240625, cultures and sequences are unavailable.

Eupenidiella Quaedvl. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *E. venezuelensis* (Crous & U. Braun) Quaedvl. & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Venezuela, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; nomenclature), holotype, ex-type strain of the type: CBS H-19934, CBS 106.75.

Euphoriomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 14 species, type: *E. bilateralis* Thaxt., asexual morph unknown, parasites, terrestrial, worldwide, see Rossi and Ponziani (2008; new species), Lumbsch and Huhndorf (2010; outline), Lee and Na (2011; new species), cultures and sequences are unavailable.

Eupropolella Höhn. 1917, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *E. vaccinii* (Rehm) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Eurychora Fuckel 1870 (= *Discomycopsis* Müll. Arg. 1893 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *E. sedi* (Link) Fuckel, asexual morph placosphaeria-like, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Eurytheca De Seynes 1878, *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, one species, type: *E. monspeliensis* De Seynes, asexual morph unknown,

saprobes, terrestrial, Trinidad, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Dissanayake et al. (2014; morphology), Wijayawardene et al. (2014c; outline), holotype of the type: F11411.

Eusynaptomyces Thaxt. 1931, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, five species, type: *E. borealis* Thaxt., asexual morph unknown, parasites, terrestrial, worldwide, see Rossi and Carolina Proaño Castro (2009; compare with *Rhachomyces*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Euteratosphaeria Quaedvl. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *E. verrucosiafricana* (Crous & M.J. Wingf.) Quaedvl. & Crous, asexual morph unknown, pathogens, terrestrial, Indonesia, have sequence data, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; outline), holotype of the type: CBS H-19705, ex-type strain of type: CBS 118496.

Eutiarosporella Crous 2015, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, seven species, type: *E. tritici* (B. Sutton & Marasas) Crous, coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, cosmopolitan, see Crous et al. (2015b; taxonomy, phylogeny), Thynne et al. (2015; new species, pathogens), Li et al. (2016a; new species, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: PREM 60866, CMW 38423.

Eutypa Tul. & C. Tul. 1863, *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, c. 35 epithets, type: *E. lata* (Pers.) Tul. & C. Tul., asexual morph coelomycetous, saprobes, pathogenic, terrestrial, widely distribute, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; morphology, phylogeny, new species), Dai et al. (2017; morphology, phylogeny), Almeida et al. (2016; new species), cultures and sequence are available, but lacking for type material, need generic revision.

Eutypella (Nitschke) Sacc. 1875 (= *Peroneutypella* Berl. 1902; = *Pseudotrype* Henn. 1900; = *Scoptria* Nitschke 1867; = *Tilakomyces* Sathe & Vaidya 1979; = *Valsa* subgen. *Eutypella* Nitschke 1870 *fide* Species Fungorum 2017), *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *E. cerviculata* (Fr.) Sacc., asexual morph unknown, saprobes, pathology, terrestrial, worldwide, see Carmarán et al. (2009; Argentina), Lumbsch and Huhndorf (2010; outline), Trouillas et al. (2010, 2011; new species, Australia), Fournier and Lechat (2011; new species), Isaka et al. (2011; endophytes, active compounds), Chacón et al. (2013; Panama, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b;

outline), Almeida et al. (2016; new species, Brazil), Thambugala et al. (2017a, phylogeny, new species), cultures and sequences are available.

Euzodiomyces Thaxt. 1900, *Euceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *E. lathrobii* Thaxt., asexual morph unknown, parasites, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Evanidomus Caball. 1941, *Ascomycota* genera *incertae sedis*, one species, type: *E. valentinus* Caball., coelomycetous, sexual morph unknown, saprobes, terrestrial, Spain, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Everhartia Sacc. & Ellis 1882, *Ascomycota* genera *incertae sedis*, three species, type: *E. hymenuloides* Sacc. & Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Evernia Ach. 1809 (= *Everniomyces* Cif. & Tomas. 1953; = *Hologymnia* Nyl. 1900 *fide Species Fungorum* 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, ten species, type: *E. prunastri* (L.) Ach., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Everniicola D. Hawksw. 1982, *Ascomycota* genera *incertae sedis*, one species, type: *E. flexispora* D. Hawksw., coelomycetous, sexual morph unknown, on lichens, terrestrial, Europe and North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype material of type: *E.*

Everniopsis Nyl. 1860 (= *Hendrickxia* P.A. Duvign. 1942 *fide Species Fungorum* 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, one species; type: *E. trulla* (Ach.) Nyl., lichenized, Central and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Eversia J.L. Crane & Schokn. 1977, *Ascomycota* genera *incertae sedis*, one species, type: *E. subopaca* (Cooke & Ellis) J.L. Crane & Schokn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Evlachovaea B.A. Borisov & Tarasov 1999, *Ascomycota* genera *incertae sedis*, one species, type: *E. kintrischica* B.A. Borisov & Tarasov, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Russia, see

Wijayawardene et al. (2012, 2017; outline), Humber et al. (2013; DNA), Kirk et al. (2013; genus accepted), Tibpromma et al. (2017b; new species), cultures and sequences are unavailable.

Exarmidium P. Karst. 1873 (= *Clypeothecium* Petr. 1922; = *Trematostoma* (Sacc.) Shear 1942; = *Zignoëlla* subgen. *Trematostoma* Sacc. 1883 *fide Species Fungorum* 2017), *Hypocreaceae*, *Amphisphaeriales*, *Sordariomycetes*, nine species, type: *E. hysteriforme* P. Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Excipularia Sacc. 1884, *Ascomycota* genera *incertae sedis*, two species, type: *E. fusispora* (Berk. & Broome) Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Russia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Excipulariopsis P.M. Kirk & Spooner 1982 (= *Kentingia* Sivan. & W.H. Hsieh 1989 *fide Wijayawardene et al. 2014c*; Rossman et al. 2015b), *Dothideomycetes*, genera *incertae sedis*, two species, type: *E. narsapurensis* (Subram.) Spooner & P.M. Kirk, hyphomycetous, sexual morph formerly known in *Kentingia*, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline; sexual morph), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline, nomenclature), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), cultures and sequences are unavailable.

Exiliseptum R.C. Harris 1986, *Dothideomycetes*, genera *incertae sedis*, two species, type: *E. ocellatum* (Müll. Arg.) R.C. Harris, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Trypetheliaceae*), Hyde et al. (2013; excluded from *Trypetheliaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Exochalara W. Gams & Hol.-Jech. 1976, *Leotiomycetes* genera *incertae sedis*, two species, type: *E. longissima* (Grove) W. Gams & Hol.-Jech., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, Europe, see Réblová et al. (2011; DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available: CBS 393.82 and CBS 980.73.

Exophiala J.W. Carmich. 1966 (= *Foxia* Castell. 1908; = *Melanclonus* Calandron 1953; = *Wangiella* McGinnis 1977 *fide Species Fungorum* 2017), *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, polyphyletic, (the core of

the genus belongs in *Herpotrichiellaceae*, but some species are related to other families of *Chaetothyriales*, such as *Chaetothyriaceae* and *Trichocomiaceae*), c. 41 species, type: *E. salmonis* J.W. Carmich., predominantly terrestrial, although some species occur in freshwater or sea environments, mostly saprobes associated with soil and plant material, but also occurring as endophytes or as opportunistic pathogens in a wide spectrum of animal hosts, including humans, certain species are common in harsh habitats such as rock surfaces or areas contaminated with aromatic hydrocarbons, cosmopolitan, see Zeng and de Hoog (2008; diagnostics of selected species), Crous and Groenewald (2011; new species), Gueidan et al. (2008, 2014a; phylogeny), de Hoog et al. (2011; DNA), Seyedmousavi et al. (2013; pathogenicity to various animals), Zupančič et al. (2016; species in indoor environments), Isola et al. (2013, 2016; species associated with polluted habitats and rock surfaces), Maciá-Vicente et al. (2016; new endophytic species), Madrid et al. (2016; phylogeny, some species from soil and river sediments). ex-type: CBS 157.67.

Exophoma Weedon 1926, *Ascomycota* genera *incertae sedis*, one species, type: *E. magnoliae* Weedon, coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Sutton (1977; nomenclature), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Tibpromma et al. (2017b; new species), cultures and sequences are unavailable.

Exormatostoma Gray 1821, *Diaporthales* genera *incertae sedis*, Sordariomycetes, type: *E. nebulosa* (Pers.) Gray, see Kirk et al. (2008; treated as nom. dub.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Exosporella Höhn. 1912, *Ascomycota* genera *incertae sedis*, one species, type: *E. symploci* Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Java, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), cultures and sequences are unavailable.

Exosporiella P. Karst. 1892 (= *Anomalemma* Sivan. 1983; = *Phanerocorynella* Höhn. 1923 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b; Species Fungorum 2017), *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *E. fungorum* (Fr.) P. Karst., sexual morph *Anomalemma* Sivan., saprobes or fungicolous, terrestrial, commonly occur on *Corticium* species in Europe, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2012, 2014c, 2016a, b, c; outline, nomenclature), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), Tian et al. (2015; morphology; nomenclature, phylogeny), cultures and

sequences are unavailable for both sexual and asexual morphs.

Exosporium Link 1809 (= *Cephaloedium* Kunze 1828; = *Cuspidosporium* Cif. 1955 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *E. tiliæ* Link, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Java, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), cultures and sequences are unavailable.

Exosporodiella Ganie, Azam & A.H. Wani 2012, *Ascomycota* genera *incertae sedis*, one species, type: *Exosporodiella phoenicis* Ganie, Azam & A.H. Wani, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Ganie et al. (2012; morphology), holotype of the type: HCIO 50058, cultures and sequences are unavailable.

Exserohilum K.J. Leonard & Suggs 1974 (= *Setosphaeria* K.J. Leonard & Suggs 1974 *fide* Wijayawardene et al. 2014c), *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, c. 30 species, type: *E. turicum* (Pass.) K.J. Leonard & Suggs, hyphomycetous, sexual morph *Setosphaeria*, on grasses, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline, nomenclature), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2015c; phylogeny), Rossman et al. (2015b; nomenclature), cultures and sequences are available.

Exserticlava S. Hughes 1978, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, seven species, type: *E. vasiformis* (Matsush.) S. Hughes, hyphomycetous, sexual morph chaetosphaeria-like, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Exrawettsteinina M.E. Barr 1972, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *E. minuta* M.E. Barr, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, accepted as a genus in *Pleosporaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015c; morphology, taxonomy), cultures and sequences are unavailable.

Extremus Quaedvl. & Crous 2014, *Extremaceae*, *Capnodiales*, *Dothideomycetes*, hyphomycetous, two species, sexual morph unknown, rock-inhabiting, terrestrial, Antarctica, Spain, have sequence data, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), holotype of the type: CBS 118292, ex-type strain of type: CBS 118292.

Extrusothecium Matsush. 1996, *Dothideomycetes* genera *incertae sedis*, one species, type: *E. caffrum* Matsush., asexual morph unknown, saprobes, terrestrial, see Kirk et al. (2008; treated as a synonym of *Leptosphaerulina*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Fabrella Kirschst. 1941, *Cenangiaceae*, *Helotiales*, *Leotiomyces*, one species, type: *F. tsugae* (Farl.) Kirschst., asexual morph unknown, saprobes, pathogens, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Fairmaniella Petr. & Syd. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *F. leprosa* (Fairm.) Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2016b; outline, morphology, taxonomic notes), Kirk et al. (2013; genus accepted), holotype of the type: IMI 57994, cultures and sequences are unavailable.

Falciformispora K.D. Hyde 1992, *Trematosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *F. lignatilis* K.D. Hyde, asexual morph unknown, saprobes on mangrove wood or pathogenic on human, Mexico, Senegal, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, accepted as a genus in *Trematosphaeriaceae*), Kirk et al. (2013; genus accepted), Ahmed et al. (2014a; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), holotype of type species BRIP 16972, cultures and sequences are available.

Falcocladium S.F. Silveira, Alfenas, Crous & M.J. Wingf. 1994, *Falcocladiaceae*, *Falcocladiales*, *Sordariomycetes*, four species, type: *F. multivesiculatum* S.F. Silveira, Alfenas, Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see da Cruz and Gusmão (2009; diversity), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Jones et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: PREM 51541, cultures and sequences are available.

Fanniomycetes T. Majewski 1972, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *F. ceratophorus* (Whisler) T. Majewski, asexual morph unknown, parasites, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Fantasmomyces DongHyeon Lee, Marinc., Z.W. de Beer & M.J. Wingf. 2016, *Sordariomycetes* genera *incertae sedis*, one species, type: *F. hyalinus* DongHyeon Lee, Marinc., Z.W. de Beer & M.J. Wingf., asexual morph

hyphomycetous, saprobes, terrestrial, South Africa, see Crous et al. (2016a; taxonomy, phylogeny), holotype and ex-type of the type: PREM 61340, CBS 139038.

Farasanispora Abdel-Wahab, Bahkali & E.B.G. Jones 2016, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *F. avicenniae* Abdel-Wahab, Bahkali & E.B.G. Jones, asexual morph unknown, saprobes, marine, Saudi Arabia, see Li et al. (2016a; taxonomy, phylogeny), holotype and ex-type of the type: CBS H-22559, MF 1207.

Farnoldia Hertel 1983, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, six species, type: *F. jurana* (Schaer.) Hertel, lichenized, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Farriolla Norman 1884 (= *Farriollomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *F. distans* Norman, asexual morph unknown, saprobes, terrestrial, Norway, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Farrowia D. Hawksw. 1975, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, four species, type: *F. longicollis* (Krzemien. & Badura) D. Hawksw., asexual morph botyrotrichum-like, on cellulosic substrata, Europe, Asia and North America, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, lectotype of type: BPI-A 1212.

Fasciatispora K.D. Hyde 1991, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, eight species, type: *F. nypae* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Liu et al. (2015a; designate reference specimen and deposited cultures), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), reference specimen and available cultures for type: MFLU 15-0042, MFLUCC 11-0382.

Fassia Dennis 1964, *Sordariomycetes* genera *incertae sedis*, eight species, type: *F. scabrosa* Dennis, asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Sordariomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Faurellina Locq.-Lin. 1975, *Chadefaudiellaceae*, *Microascales*, *Sordariomycetes*, four species, type: *F. fimigena* Locq.-Lin., asexual morph arthrophagous-like,

coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Réblová et al. (2011; DNA), Rossman et al. (2012; DNA), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are available *F. indica* (CBS 301.78), *F. elongata*.

Fauriea S.Y. Kondr., L. Lökö & J.-S. Hur 2016, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *F. orientochinensis* S.Y. Kondr., X.Y. Wang & Hur, lichenized, see Kondratyuk et al. (2016b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Favostroma B. Sutton & E.M. Davison 1983, *Ascomycota* genera *incertae sedis*, one species, type: *F. crypticum* B. Sutton & E.M. Davison, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 266341, cultures and sequences are unavailable, genus need revision.

Felipes Frisch & G. Thor 2014, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *F. leucopellaeus* (Ach.) Frisch & G. Thor, lichenized, see Frisch et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Felisbertia Viégas 1944, *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *F. melastomacearum* (Speg.) Viégas, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Fellhanera Vězda 1986 (= *Lobaca* Vězda 1983 *fide* Species Fungorum 2017), *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, 97 species, type: *F. fuscatula* (Müll. Arg.) Vězda, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2013b; new species), Lücking et al. (2017a; classification), sequences are available.

Fellhaneropsis Sérus. & Coppins 1996, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, seven species, type: *F. myrtillicola* (Erichsen) Sérus. & Coppins, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Feltgeniomycetes Dieder. 1990, *Ascomycota* genera *incertae sedis*, five species, type: *F. luxemburgensis* Dieder., hyphomycetous, lichenicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of type: Diederich 8477 (LG: holotype; IMI 335016, UPS, herb. Diederich: isotypes), cultures and sequences are unavailable.

Fenestella Tul. & C. Tul. 1863, *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, c. four species, type: *F. princeps* Tul. & C. Tul., asexual morph unknown, saprobes or biotrophic or hemibiotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Fenestellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Phookamsak and Hyde (2015; taxonomy), Wanasinghe et al. (2017b; new species, phylogeny, placement), cultures and sequences are available.

Fenestroconidia Calatayud & Etayo 1999, *Ascomycota* genera *incertae sedis*, one species, type: *F. caloplacae* Calat. & Etayo, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, Spain, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: VAB (lich.) 15301, cultures and sequences are unavailable.

Ferrarisia Sacc. 1919 (= *Microthyriolum* Speg. 1919 [1918] *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, ten species, type: *F. philippina* Sacc., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), placement in *Parmulariaceae* is supported by morphology, cultures and sequences are unavailable.

Ferraroa Lücking, Sérus. & Vězda 2005, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *F. hyalina* (Lücking) Lücking, Sérus. & Vězda, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Fibrillithecis Frisch 2006, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 16 species, type: *F. vernicosa* (Zahlbr.) Frisch, asexual morph unknown, lichenized fungi, terrestrial, tropical, placement in *Graphidaceae* supported by molecular data, see Rivas Plata et al. (2010a, 2012a; key, taxonomy), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Filariomycetes Shanor 1952 (= *Labiduromyces* Ishik. 1941 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *F. forcipulae* Shanor, asexual morph unknown, parasites, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Filicupula Y.J. Yao & Spooner 1996, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, two species, type: *F. suboperculata* (Döbbeler & P. James) Y.J. Yao & Spooner, asexual morph unknown, parasites, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk

et al. (2013; genus accepted), cultures and sequences are unavailable.

Filiella Videira & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *F. pastinacae* (P. Karst.) Videira & Crous, hyphomycetous, sexual morph unknown, plant pathogenic, see Videira et al. (2016; morphology, phylogeny), cultures and sequences are available, neotype is in HAL.

Filosporella Nawawi 1976 (= *Coeloanguillospora* Dyko & B. Sutton 1978 [1977]; = *Rogersia* Shearer & J.L. Crane 1976 *fide* Species Fungorum 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *F. aquatica* Nawawi, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Baschien et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Filsoniana S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetidae*, *Lecanoromycetes*, five species, type: *F. australiensis* (S.Y. Kondr., Kärnefelt & Filson) S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur, lichenized, see Kondratyuk et al. (2016b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Fimetariella N. Lundq. 1964, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *F. rabenhorstii* (Niessl) N. Lundq., asexual morph unknown, saprobes, terrestrial, Europe, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Finkia Vain. 1929, *Lichinaceae*, *Lichinales*, *Lichinomycetes*, one species, type: *F. portoricensis* Vain., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Fischerula Mattir. 1928, *Morchellaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *F. macrospora* Mattir., asexual morph unknown, ectomycorrhizal, saprobes, terrestrial, Italy, USA, see Lumbsch and Huhndorf (2010; outline, accepted as *Morchellaceae*), Trappe et al. (2010b; DNA, phylogeny), Alvarado et al. (2011a; DNA, phylogeny, accepted as a genus in *Helvellaceae*), Healy et al. (2013; DNA, phylogeny, ectomycorrhizal), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; accepted as a genus in *Morchellaceae*), cultures and sequences are available.

Fissurina Fée 1824, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, over 150 species, type: *F. dumastii* Fée, asexual morph unknown, lichenized, terrestrial,

mostly tropical, see Archer (2009; key Australia), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2014, 2016a, b), Kirk et al. (2013; genus accepted), sequences are available.

Fissuroma Jian K. Liu, Phook., E.B.G. Jones & K.D. Hyde 2011, *Aigialaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *F. maculans* (Rehm) J.K. Liu, E.B.G. Jones & K.D. Hyde, asexual morph unknown, saprobes, Japan, Thailand, see Liu et al. (2011b; morphology, DNA, phylogeny), Hyde et al. (2013; accepted as a genus in *Aigialaceae*), Wijayawardene et al. (2014c; outline), Phookamsak et al. (2015b; taxonomy), holotype and ex-type of the type: MFLU 11-1143, MFLUCC 10-0886.

Fitzpatrickella Benny, Samuelson & Kimbr. 1985, *Coryneliaceae*, *Coryneliales*, *Eurotiomycetes*, one species: *F. operculata* Benny, Samuelson & Kimbr., asexual morph unknown, terrestrial, parasitic on fruits of *Drimys confertifolia* Phil., Juan Fernández Island, Chile, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wood et al. (2016; morphotaxonomy), cultures and sequences are unavailable, needs phylogenetic assessment.

Flabellocladia Nawawi 1985, *Ascomycota* genera *incertae sedis*, four species, type: *F. gigantea*, hyphomycetous, from foam, aquatic, worldwide, Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Fiúza and Gusmão (2013; morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Flabellospora Alas. 1968, *Ascomycota* genera *incertae sedis*, six species, type: *F. crassa*, hyphomycetous, on submerged plant material, aquatic, worldwide, see Patil et al. (2010; morphology), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Flagellosphaeria Aptroot 1995, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type: *F. polytrichospora* (Lucas & Sousa de Câmara) Aptroot, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015a, b; outline), cultures and sequences are unavailable.

Flagellospora Ingold 1942, *Leotiaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *F. curvula* Ingold, hyphomycetous, saprobes, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Baschien et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Lombard et al. (2014; phylogeny), cultures and sequences are available.

Flakea O.E. Erikss. 1992, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *F. papillata* O.E. Erikss., lichenized, see Lumbsch and

Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Flammeascoma Phook. & K.D. Hyde 2015, *Anteagloniaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *F. bambusae* Phook. & K.D. Hyde, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2015a; taxonomy, new species), Liu et al. (2015a; taxonomy), holotype and ex-type of the type: MFLU 10-0061, MFLUCC 10-0551.

Flammispora Pinruan, Sakay., K.D. Hyde & E.B.G. Jones 2004, *Savoryellales*, genera *incertae sedis*, *Sordariomycetes*, two species, type: *F. bioteca* Pinruan, Sakay., K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, Thailand, USA, see Lumbsch and Huhndorf (2010; outline, as a genus in *Hypocreomycetidae*, genera *incertae sedis*), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type of the type: BBH WAH134, BCC13368.

Flammocladia Crous, L. Lombard & R.K. Schumach. 2015, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *F. aceris* Crous, L. Lombard & R.K. Schumach., asexual morph coelomycetous, saprobes, Germany, see Crous et al. (2015c; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22263, CPC 24422.

Flavobathelium Lücking, Aptroot & G. Thor 1997, *Strigulaceae*, *Strigulales*, *Dothideomycetes*, one species, type: *F. epiphyllum* Lücking, Aptroot & G. Thor, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2011b; DNA, phylogeny), Hyde et al. (2013; accepted as a genus in Strigulaceae), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Lücking et al. (2017a; classification), cultures available for the type: MPN67 (*fide* Nelsen et al. 2011a, b).

Flavocetraria Kärnefelt & A. Thell 1994, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *F. cucullata* (Bellardi) Kärnefelt & A. Thell, asexual morph, unknown, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Flavomyces D.G. Knapp, Kovács, J.Z. Groenew. & Crous 2015, *Massarinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *F. fulophazii* D.G. Knapp, Kovács, J.Z. Groenew. & Crous, endophytes, see Knapp et al. (2015; phylogeny), cultures and sequences are available.

Flavoparmelia Hale 1986, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 40 species, type: *F. caperata* (L.) Hale, asexual morph unknown, lichenized, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Del-Prado et al. (review,

phylogeny), Lücking et al. (2017a; classification), sequences are available.

Flavoplaca Arup, Søchting & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 27 species, type: *F. citrina* (Hoffm.) Arup, Frödén & Søchting, asexual morph unknown, lichenized, terrestrial, widely distributed, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Flavopunctelia (Krog) Hale 1984 (= *Punctelia* subgen. *Flavopunctelia* Krog 1982 *fide* Species Fungorum 2017), seven species, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, type: *F. flaventior* (Stirt.) Hale, asexual morph unknown, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny, molecular), Divakar et al. (2015; evolutionary study of family), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Flegographa A. Massal. 1860 (see Species Fungorum 2017 for synonyms), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *F. leprieurii* (Mont.) A. Massal., lichenized, see Rivas Plata et al. (2013; taxonomy, reinstated the genus), Lücking et al. (2017a; classification), sequences are available.

Floricola Kohlm. & Volk.-Kohlm. 2000, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *F. striata* Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, cosmopolitan, see Thambugala et al. (2015b; taxonomy), Ariyawansa et al. (2015a; new species), Jaklitsch et al. (2016c; treated as a synonym of *Teichospora*, but in here we do not agree with them), Wijayawardene et al. (2016b; taxonomy), cultures and sequences are available.

Flosculomyces B. Sutton 1978, *Ascomycota* genera *incertae sedis*, two species, type: *F. floridaensis* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, Ivory Coast, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: IMI 215981c, cultures and sequences are unavailable, genus need revision.

Fluctua Marbach 2000, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *F. megapotamica* (Malme) Marbach, lichenized, see Lücking et al. (2017a; classification), sequences are unavailable.

Fluminicola S.W. Wong, K.D. Hyde & E.B.G. Jones 1999, *Papulosaceae*, *Diaportheomycetidae* families *incertae sedis*, *Sordariomycetes*, one species, type: *F. bipolaris* S.W. Wong, K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, Philippines, see Lumbsch and Huhndorf (2010; outline, as a genus in *Annulatasaceae*), Kirk et al. (2013; genus accepted), Réblová et al. (2013; placed in *Papulosaceae*),

Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Papulosaceae*), holotype of the type: HKU (M) 3127.

Fluviatispora K.D. Hyde 1994, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, three species, type: *F. tunicata* K.D. Hyde, saprobes, aquatic, Brunei, Papua New Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016a, b, c), cultures and sequences are unavailable, holotype of type: BRIP 21391.

Fodinomyces Vázquez-Campos & Neilan 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *F. uranophilus* X. Vázquez-Campos, hyphomycetous, sexual morph unknown, from soil, terrestrial, Australia, see Vazquéz-Campos et al. (2014; morphology, DNA, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS 136962.

Foliocryphia Cheew. & Crous 2009, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *F. eucalypti* Cheew. & Crous, coelomycetous, sexual morph unknown, pathogens, terrestrial, Australia, see Cheewangkoon et al. (2009; morphology, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type cultures of type: CBS H-20299, ex-type CPC 12494 = CBS 124779.

Follmannia C.W. Dodge 1967 (see Species Fungorum 2017 for synonyms), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *F. rufa* C.W. Dodge, lichenized, see Arup et al. (2013; reinstated the genus), Lücking et al. (2017a; classification), sequences are available.

Follmanniella Peine & B. Werner 1995, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *F. scutellata* Peine & B. Werner, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Fonsecaea Negroni 1936, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, eight species, type: *F. pedrosoi* (Brumpt) Negroni, hyphomycetous, mostly terrestrial, in soil and on plant material, but also reported from aquatic environments, most species are pathogenic to animals, ranging from crustaceans to humans, America, Asia, Europe and Africa, see Vicente et al. (2008, 2012; environmental reservoirs, new species), Najafzadeh et al. (2010, 2011a, b; phylogeny, new species, molecular epidemiology), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Seyedmousavi et al. (2013; pathogenicity to animals), de Azevedo et al. (2015; new species), ex-neotype: CBS 271.37.

Fontanospora Dyko 1978, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *F. eccentrica* (R.H. Petersen) Dyko, hyphomycetous, sexual morph unknown, saprobes, aquatic, USA, see Baschien et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Forliomyces Phukhams., Camporesi & K.D. Hyde 2016, *Sporormiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *F. uniseptata* Phukhams., Camporesi & K.D. Hyde, coelomycetous, saprobes, terrestrial, Italy, see Phukhamsakda et al. (2016a; taxonomy, phylogeny), Wijayawardene et al. (2016b, 2017; taxonomy, outline), holotype and ex-type of the type: MFLU 16-0031, MFLUCC 15-0765.

Fouragea Trevis. 1880 (see Species Fungorum 2017 for synonyms), *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *F. filicina* (Mont.) Trevis., lichenized, see Frisch et al. (2014; reinstated), Lücking et al. (2017a; classification), sequences are available.

Foveostroma DiCosmo 1978 (= *Micropera* Lév. 1846; = *Micula* Duby 1858 fide Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, six species, type: *F. drupacearum* (Lév.) DiCosmo, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Fracchiaea Sacc. 1873, *Nitschiaceae*, *Coronophorales*, *Sordariomycetes*, three species, type: *F. heterogenea* Sacc., asexual morph unknown, lichenicolous, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Mugambi and Huhndorf (2010; DNA), Vasilyeva et al. (2010; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for *Fracchiaea broomeana* (SMH347 fide Mugambi and Huhndorf 2010).

Fragosphaeria Shear 1923, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, four species, type: *F. purpurea* Shear, asexual morph hyphomycetous, from soil, pathogens (human), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not accept), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for type: CBS 133.34.

Franwilsia S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *F. bastowii* (S.Y. Kondr. & Kärnefelt) S.Y. Kondr., Kärnefelt, Elix, A. Thell, J. Kim, A.S. Kondr. & Hur; lichenized, see Kondratyuk et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Fremineavia Nieuwl. 1916 (= *Titania* Berl. 1890 *fide* Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *F. berkeleyi* (Berl.) Nieuwl., asexual morph unknown, saprobes, North America, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Fremitomyces P.F. Cannon & H.C. Evans 1999, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *F. punctatus* P.F. Cannon & H.C. Evans, sexual morph unknown, saprobes, terrestrial, East Africa, Seychelles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Friedmanniomycetes Onofri 1999, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *F. endolithicus* Onofri, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Antarctica, see Schoch et al. (2009; phylogeny), Selbmann et al. (2008; DNA, review, phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Teratosphaeriaceae*), Kirk et al. (2013; genus accepted), Egidi et al. (2014; phylogeny), Ertz et al. (2014b; phylogeny), Quaedvlieg et al. (2014; phylogeny), cultures available for the epitype: CBS 119423=CCFEE 5208.

Frigidispora K.D. Hyde & Goh 1999, *Ascomycota* genera *incertae sedis*, six species, type: *F. colnensis* K.D. Hyde & Goh, coelomycetous, sexual morph unknown, saprobes, aquatic, British Isles, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Frigidopyrenia Grube 2005, *Ascomycota* genera *incertae sedis*, one species, type: *F. bryospila* (Nyl.) Grube, lichenized, subarctic, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Xanthopyreniaceae*), Lücking et al. (2017a; classification), sequences are unavailable.

Frondicola K.D. Hyde 1992, *Hypocretriaceae*, *Amphisphaerales*, *Sordariomycetes*, nine species, type: *F. tuniticuspis* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Brunei, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Frondisphaeria K.D. Hyde 1996, *Dothideomycetes* genera *incertae sedis*, two species, type: *F. palmicola* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, misspelled as *Frondisphaera*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

accepted), Maharachchikumbura et al. (2015b, 2016b; outline, misspelled as *Frondisphaera*), Daranagama et al. (2016d; taxonomy), cultures and sequences are unavailable.

Frondispora K.D. Hyde 1993, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type: *F. bicalcarata* (Ces.) K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Frutidella Kalb 1994, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *F. caesioatra* (Schaer.) Kalb, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Fujimyces Minter & Caine 1980, *Ascomycota* genera *incertae sedis*, two species, type: *F. oödes* (Bayl. Ell.) Minter & Caine, coelomycetous, sexual morph unknown, saprobes, terrestrial, British Isles, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI35344, cultures and sequences are unavailable.

Fulgidea Bendiksby & Timdal 2013, *Umbilicariaceae*, *Umbilicariales*, *Lecanoromycetes*, two species, type: *F. oligospora* (Timdal) Bendiksby & Timdal, lichenized, see Bendiksby and Timdal (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available. holotype of type species: O-L-767,

Fuligomyces Morgan-Jones & Kamal 1984, *Ascomycota* genera *incertae sedis*, two species, type: *F. macrosporus*, asexual morph hyphomycetous, saprobes (symptoms on leaves similar to sooty mould), terrestrial, India, (Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of type: AUA, IMI 248960, GPU (KA 69), cultures and sequences are unavailable.

Fulvia Cif. 1954 (= *Berteromyces* Cif. 1954; = *Cercodeuterospora* Curzi 1932; = *Ragnhildiana* Solheim 1931 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *F. fulva* (Cooke) Cif., hyphomycetous, sexual morph unknown, on *Lycopersicon esculentum*, plant pathogenic, see Videira et al. (2016; morphology, phylogeny), holotype BPI 426698, for epitypification see Videira et al. (2017).

Fulvocentrum E.B.G. Jones & Abdel-Wahab 2014, *Juncigenaceae*, *Torpedosporales*, *Sordariomycetes*, two species, type: *F. aegyptiaca* (Abdel-Wahab, El-Shar. & E.B.G. Jones) E.B.G. Jones & Abdel-Wahab, asexual morph undermined, marine, tropical/subtropical, see Jones et al. (2014, 2015; morphology, phylogeny, outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Fulvoflamma Crous 2006, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *F. eucalypti* Crous, coelomycetous, sexual morph unknown, saprobes, endophytes, terrestrial, Spain, see Wijayawardene et al. (2012, 2017; outline), holotype and ex-type strains of the type: CBS H-18045, 1243 = CBS 118549.

Fulvophyton Ertz & Tehler 2011, *Roccellographaceae*, *Arthoniales*, *Arthoniomycetes*, nine species, type: *F. stalactinum* (Nyl.) Ertz & Tehler, sexual morph with punctiform or shortly lirellate ascocarps, lichenized, terrestrial, tropical, see Ertz and Tehler (2011; morphology, phylogeny), Tehler et al. (2013a; one new combination), Lücking et al. (2017a; classification), sequences are available.

Fumagopsis Speg. 1911 (= *Tricornispora* Bonar 1967 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *F. triglifoides*, hyphomycetous, sexual morph unknown, saprobes or plant pathogenic on plant leaves, terrestrial, known from Argentina and India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of type: in LPS, cultures and sequences are unavailable.

Fumiglobus D.R. Reynolds & G.S. Gilbert 2006, *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, 15 species, type: *F. ficiinus* (Bat., Nascim. & Cif.) D.R. Reynolds & G.S. Gilbert, coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Wijayawardene et al. (2014c; outline), Bose et al. (2014; DNA, new species), cultures and sequences are available.

Funbolia Crous & Seifert 2011, *Dothideomycetes* genera *incertae sedis*, one species, type: *F. dimorpha* Crous & Seifert, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Crous et al. (2011a; morphology, DNA), Wijayawardene et al. (2012, 2014c; outline), holotype and ex-type strains of the type: CBS H-20577, CPC 14170.

Funiliomyces Aptroot 2004, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type: *F. biseptatus* Aptroot, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available for the type: AY772015 (LSU) (fide Aptroot 2004).

Furcaspora Bonar 1965, *Ascomycota* genera *incertae sedis*, two species, type: *F. pinicola* Bonar, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI35344, ex-type cultures and sequences are available for *Furcaspora eucalypti* Crous & Verkley (CPC 12556 = CBS 119111).

Fusamen (Sacc.) P. Karst. 1890 (= *Calogloeum* Syd. 1924; = *Fusarium* subgen. *Fusamen* Sacc. 1886; = *Platycarpium* P. Karst. 1905 [1905-1906] *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *F. fungicola* P. Karst., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Fusariella Sacc. 1884 (= *Kurssanovia* Pidopl. 1948 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypoocreales*, *Sordariomycetes*, 19 species, type: *F. atrovirens*, asexual morph hyphomycetous, on plants, leaf litter, dung and in soil, terrestrial, worldwide, Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Lin et al. (2016a; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Fusarium Link 1809 (= *Gibberella* Sacc. 1877 *fide* Maharachchikumbura et al. 2016b; see Lombard et al. 2015b and Species Fungorum 2017 for additional synonyms), *Nectriaceae*, *Hypoocreales*, *Sordariomycetes*, c. 120 species, type: *F. roseum* Link, hyphomycetous, sexual morph *Gibberella*, terrestrial, pathogens, worldwide, see Seifert et al. (2011; morphology), Zhu et al. (2013; pathogens), Wijayawardene et al. (2012, 2017; outline), Lombard et al. (2015b; phylogeny, taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Fuscidea V. Wirth & Vězda 1972 (= *Biatorinella* Deschâtres & Werner 1974; = *Fuscidea* V. Wirth & Vězda 1972 *fide* Species Fungorum 2017), *Fuscideaceae*, *Umbilicariales*, *Lecanoromycetes*, 35 species, type: *F. aggregata* (Flot.) V. Wirth & Vězda, lichenized, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; taxonomy), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Fuscoderma (D.J. Galloway & P.M. Jørg.) P.M. Jørg. & D.J. Galloway 1989 (= *Leioderma* subgen. *Fuscoderma* D.J. Galloway & P.M. Jørg. 1987 *fide* Species Fungorum 2017), *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, five species, type: *F. applanatum* (D.J. Galloway & P.M. Jørg.) P.M. Jørg. & D.J. Galloway, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Fuscolachnum J.H. Haines 1989, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *F. pteridis* (Alb. & Schwein.) J.H. Haines, asexual morph unknown, saprobes, epiphytes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Fuscopannaria P.M. Jørg. [nom. cons. prop.] 1994 (= *Kroswia* P.M. Jørg. 2002; = *Moelleropsis* Gyeln. 1939 *fide* Jørgensen et al. 2013; Species Fungorum 2017), *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 58 species, type: *F. leucosticta* (Tuck.) P.M. Jørg., lichenized, see Ekman et al. (2014; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jørgensen et al. (2013; *Fuscopannaria* over *Moelleropsis*), Lücking et al. (2017a; classification), sequences are available.

Fuscophialis B. Sutton 1977, *Ascomycota* genera *incertae sedis*, four species, type: *F. brasiliensis* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, Cuba, Mauritius, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, genus need revision.

Fuscosclera Hern.-Restr et al. 2017, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *F. lignicola* Hern.-Restr., J. Mena & Gene, hyphomycetous, sexual morph unknown, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Fuscosporella J. Yang, Bhat & K.D. Hyde 2016, *Fuscosporellaceae*, *Fuscosporellales*, *Sordariomycetes*, one species, type: *F. pyriformis* J. Yang, Bhat & K.D. Hyde, hyphomycetous, saprobes, Thailand, see Yang et al. (2016c; taxonomy, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Fuscostagonospora Kaz. Tanaka & K. Hiray. 2015, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *F. sasae* Kaz. Tanaka & K. Hiray., coelomycetous, saprobes, terrestrial, Japan, see Tanaka et al. (2015; morphology, phylogeny), Wijayawardene et al. (2016b; taxonomy), holotype and ex-type of the type: HHUF 29106, CBS 139687.

Fusichalara S. Hughes & Nag Raj 1973, *Ascomycota* genera *incertae sedis*, two species, type: *F. dimorphospora* S. Hughes & Nag Raj, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012; outline, treated as a genus in *Chaetosphaeriaceae*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; excluded from *Chaetosphaeriaceae* and placed in *Ascomycota* genera *incertae sedis*), cultures and sequences are unavailable.

Fusicladiella Höhn. 1919, *Ascomycota* genera *incertae sedis*, five species, type: *F. aronici* (Sacc.) Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology) and Wijayawardene et al. (2012; treated as *Mycosphaerellaceae*), Kirk et al. (2013; genus accepted),

Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Fusicladium Bonord. 1851 (see Species Fungorum 2017 for synonyms), *Sympoventuriaceae*, *Venturiales*, *Dothideomycetes*, c. 75 species, type: *P. virescens* Bonord., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Braun (2009; new species), Seifert et al. (2011; morphology), Crous et al. (2016b; new species, phylogeny), Wijayawardene et al. (2017; outline), cultures are available.

Fusiconidium J.F. Li, Phookamsak & K.D. Hyde 2017, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. aquaticus* Z.L. Luo, Baht & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Italy, China, see Li et al. (2017a; taxonomy, phylogeny), holotype of the type: MFLU 16-2139, cultures and sequences are available.

Fusicolla Bonord. 1851 (= *Fusarium* subgen. *Leptosporium* Sacc. 1886; = *Leptosporium* (Sacc.) Höhn. 1923 *fide* Species Fungorum 2017), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, eight species, type: *F. betae* Bonord., hyphomycetous, holomorph, diverse substrates and hosts, terrestrial and aquatic, worldwide, see Gräfenhan et al. (2011; DNA, phylogeny) Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available.

Fusoidiella Videira & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *F. depressa* (Berk. & Broome) Videira & Crous, hyphomycetous, sexual morph mycosphaerella-like, plant pathogenic, see Videira et al. (2016; morphology, phylogeny), ex-type strain unavailable, authentic specimen (herb. K(M) 29181, on *Angelica sylvestris*, Great Britain).

Fusoidispora Vijaykr., Jeewon & K.D. Hyde 2005, *Annulatasaceae*, *Annulatascales*, *Sordariomycetes*, one species, type: *F. aquatica* Vijaykr., Jeewon & K.D. Hyde, asexual morph unknown, saprobes, aquatic, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: HKU (M) 17484, cultures and sequences are available.

Fusticeps Webster & R.A. Davey 1980, *Ascomycota* genera *incertae sedis*, five species, type: *F. bullatus* Webster & R.A. Davey, hyphomycetous, sexual morph unknown, on plant litter in water, aquatic, Tropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Chuaseeharonnachai et al. (2014; key), holotype of type: IMI 245198, cultures and sequences are unavailable.

Gabura Adans., *Arctomiaceae*, *Arctomiales*, *Lecanoromycetes*, one species, type: *G. fascicularis* (L.)

P.M. Jørg.), see Jørgensen (2014: taxonomy), Lücking et al. (2017b: classification), sequences are available.

Gaeumannella Petr. 1952, *Ascomycota* genera *incertae sedis*, one species, type: *G. singularis* Petr., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gaeumannomyces Arx & D.L. Olivier 1952 (= *Harpophora* W. Gams 2000; = *Raphidophora* Ces. & De Not. 1863; = *Raphidospora* Fr. 1849 *fide* Zhang et al. 2016d; Species Fungorum 2017), *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, c. 13 species, type: *G. graminis* (Sacc.) Arx & D.L. Olivier, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011c; phylogeny), Kirk et al. (2013; genus accepted), Zhang et al. (2015a, b; phylogeny), Luo et al. (2015a; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Gaillardiella Pat. 1895, *Bertiaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *G. pezizoides* Pat., asexual morph unknown, saprobes, terrestrial, South America, see Mugambi and Huhndorf (2010; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures available for the type: GK1245.

Galactomyces Redhead & Malloch 1977, ?*Dipodascaceae*, *Saccharomycetales*, *Saccharomycetes*, five species, type: *G. geotrichum* (E.E. Butler & L.J. Petersen) Redhead & Malloch, asexual reproduction is present, saprophytic, soil, rotted fruit and other plant parts, tree exudates, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Galeaticarpa A. Hashim. & Kaz. Tanaka 2017, *Lophiotremataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *G. aomoriensis* A. Hashim. & Kaz. Tanaka, asexual morph coelomycetous, saprobes, Asia, see Hashimoto et al. (2017; taxonomy, phylogeny), sequences are available.

Galeoscypha Svrček & J. Moravec 1989, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *G. pileiformis* (Svrček) Svrček & J. Moravec, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Galiella Nannf. & Korf 1957, *Sarcosomataceae*, *Pezizales*, *Pezizomycetes*, eight species, type: *G. rufa* (Schwein.) Nannf. & Korf, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Galicola Tibpromma, Camporesi & K.D. Hyde 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *Gallicola pseudophaeosphaeria* Tibpromma, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Ariyawansa et al. (2015a; taxonomy, phylogeny), cultures available for type: MFLUCC 14-0524.

Gampsonema Nag Raj 1975, *Ascomycota* genera *incertae sedis*, one species, type: *G. exile* (Tassi) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gamsia M. Morelet 1969, *Microascaceae*, *Microascales*, *Sordariomycetes*, three species, type: *G. dimera* (W. Gams) M. Morelet, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Sandoval-Denis et al. (2016b; DNA, phylogeny), cultures and sequences are available for *G. simplex* (CBS 546.69) and *G. aggregata* (CBS 251.69).

Gamsylella M. Scholler, Hagedorn & A. Rubner 1999, *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, type: *G. arcuata* (Scheuer & J. Webster) M. Scholler, Hagedorn & A. Rubner, cf. six species, hyphomycetous, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Gangliophora Subram. 1992, *Ascomycota* genera *incertae sedis*, one species, type: *G. casuarinae* (Matsush.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Taiwan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gangliostilbe Subram. & Vittal 1976, *Ascomycota* genera *incertae sedis*, one species, type: *G. indica* Subram. & Vittal, hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Ma et al. (2014a, b; new species, China), cultures and sequences are unavailable.

Garnaudia Borowska 1977, *Ascomycota* genera *incertae sedis*, three species, type: *G. elegans* Borowska, hyphomycetous, synanamorph selenosporella-like, on wood, bark and dead leaves, terrestrial, Cuba, Poland, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted),

holotype of type: WA 20592, cultures and sequences are unavailable.

Gassicurtia Féé 1825 (see Species Fungorum 2017 for synonyms), *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 27 species, type: *G. coccinea* Féé, lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Gaubaea Petr. 1942, *Ascomycota* genera *incertae sedis*, two species, type: *G. insignis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, central Asia, see Wijayawardene et al. (2012, 2016b, 2017; outline, morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Geastrumia Bat. 1960 (= *Benekea* Bat. & J.L. Bezerra 1960; = *Shomea* Bat. & J.L. Bezerra 1964 [1963] *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *G. polystigmatis* Bat. & M.L. Farr, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Geejayessia Schroers, Gräfenhan & Seifert 2011, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, c. five species, type: *G. cicanticum* (Berk.) Schroers (current name: *Fusarium cicanticum* (Berk.) O'Donnell & Geiser *fide* Geiser et al. 2013), sexual morph unknown, saprobes, pathogens, terrestrial, cosmopolitan, see Geiser et al. 2013; treated the type species as a synonym of *Fusarium cicanticum* (Berk.), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Geisleria Nitschke 1861 (see Species Fungorum 2017 for synonyms), *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *G. sychnogonoides* Nitschke, lichenized, see Aptroot et al. (2015b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Gelasinospora Dowding 1933, *Sordariaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *G. tetrasperma* Dowding, asexual morph unknown, saprobes, terrestrial, Ivory Coast, see Lumbsch and Huhndorf (2010; outline), Nygren et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available for type: ATCC 11345.

Gelatinipulvinella Hosoya & Y. Otani 1995 (= *Aureohyphozyma* Hosoya & Y. Otani *fide* Johnston et al. 2014b), *Helicogoniaceae*, *Phacidiales*, *Leotiomycetes*, two species, type: *G. astraeicola* Hosoya & Y. Otani, asexual morph hyphomycetous, saprobe, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), holotype of the type: TNS-F-52070.

Gelatinocrinis Matsush. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *G. roseus* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gelatinomyces N. Sanoamuang, W. Jitjak, S. Rodtong & A.J.S. Whalley 2013, *Leotiomycetes* genera *incertae sedis*, one species, type: *G. siamensis* N. Sanoamuang, W. Jitjak, S. Rodtong & A.J.S. Whalley, asexual morph hyphomycetous, saprobes, terrestrial, Thailand, see Sanoamuang et al. (2013; morphology, DNA, phylogeny), Jaklitsch et al. (2016a; classification, treated as a synonym of *Myriodiscus*), holotype and ex-type cultures of the type: K(M) 188078, CBS 135071, close to *Leotiomycetes* in phylogenetic analyses.

Gelatinopsis Rambold & Triebel 1990 (= *Micropyxis* Seeler 1943 *fide* Species Fungorum 2017), *Helicogoniaceae*, *Phacidiales*, *Leotiomycetes*, cf. eight species, type: *G. geoglossi* (Ellis & Everh.) Rambold & Triebel, asexual morph unknown, saprobes, lichenicolous, terrestrial, worldwide, see Navarro-Rosinés et al. (2008; new species), Kocourková and Knudsen (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gelatinopycnis Dyko & B. Sutton 1979, *Ascomycota* genera *incertae sedis*, one species, type: *G. hartigiana* (Allesch.) Dyko & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gelatinosporium Peck 1873 (= *Gelatinosporis* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *G. betulinum* Peck, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gelineostroma H.J. Swart 1988, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *G. athrotaxis* H.J. Swart, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gelinginia Alstrup & D. Hawksw. 1990, *Helicogoniaceae*, *Phacidiales*, *Leotiomycetes*, one species, type: *G. associata* (Th.Fr.) Alstrup & D. Hawksw., asexual morph unknown, on lichens, terrestrial, Arctic, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Odontotremataceae*), Kirk et al. (2013; genus accepted), Suija et al. (2014; phylogeny),

Jaklitsch et al. (2016a, b; position), holotype of type: UPS, cultures and sequences are unavailable.

Geminispora Pat. 1893 (= *Diplosporis* Clem. 1909 *fide* Species Fungorum 2017), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *G. mimosae* Pat., sexual morph unknown, saprobes, terrestrial, East Africa, Seychelles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Geminoarcus K. Ando 1993, *Ascomycota* genera *incertae sedis*, three species, type: *G. pachysporus* K. Ando, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gemmaspora D. Hawksw. & Halici 2007, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *G. lecanorae* (Werner) D. Hawksw. & Halici, lichenicolous, Syria, Turkey and Uzbekistan, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; position), holotype of type: BC and epitype ERC-MGH 02274, cultures and sequences are unavailable.

Gemmulina Descals & Marvanová 1999, *Ascomycota* genera *incertae sedis*, one species, type: *G. botryosa* (Descals) Descals & Marvanová, asexual morph hyphomycetous, has a yeast-like phase, from foam in streams, aquatic, Canada and United Kingdom, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of type: IMI 254853, cultures and sequences are unavailable.

Genea Vittad. 1831 (= *Genabea* Tul. & C. Tul. 1845; = *Hydnocaryon* Wallr. 1833; = *Myrmecocystis* Harkn. 1899; = *Pseudogenea* Bucholtz 1901 *fide* Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *G. verrucosa* Vittad., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Alvarado et al. (2014; new species, phylogeny), Alvarado et al. (2016; revisit), Kaounas et al. (2016; new species), cultures and sequences are available.

Geniculospora Sv. Nilsson ex Marvanová & Sv. Nilsson 1971 (= *Geniculospora* Sv. Nilsson 1964 *fide* Species Fungorum 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *G. inflata* (Ingold) Sv. Nilsson ex Marvanová & Sv. Nilsson, sexual morph unknown, saprobes, aquatic, worldwide, see Wijayawardene et al. (2012, 2017; outline), Duarte et al. (2012; DNA), Kirk et al. (2013; did not list), cultures and sequences are available.

Geodina Denison 1965, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *G. guanacastensis* Denison, asexual morph unknown, saprobe, terrestrial, Costa Rica, Canary Islands, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Geoglossum Pers. 1794 (= *Cibalocoryne* Hazsl. 1881; = *Corynetes* Hazsl. 1881; = *Gloeoglossum* E.J. Durand 1908; = *Microglossum* Sacc. 1884; = *Thuemenidium* Kuntze 1891 *fide* Species Fungorum 2017), *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, two species, type: *G. glabrum* Pers., asexual morph unknown, saprobe, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Arauzo and Iglesias (2014; new species), Loizides et al. (2015; new species), Crous et al. (2016b; new species), cultures and sequences are available.

Geomycetes Traaen 1914, *Myxotrichaceae*, *Leotiomycetes* families *incertae sedis*, c. ten species, type: *G. auratus* Traaen, hyphomycetous, sexual morph unknown, saprobes, in soil, on many other substrata, humans and other animals, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017), Chen et al. (2017b; new species), cultures and sequences are available.

Geopora Harkn. 1885 (= *Peziza* subgen. *Sepultaria* Cooke 1879; = *Pseudohydnotrya* E. Fisch 1897; = *Sepultaria* (Cooke) P. Karst. 1884 *fide* Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, c. 20 species, type: *G. cooperi* Harkn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Frank et al. (2011; new species), Kirk et al. (2013; genus accepted), Flores-Rentería et al. (2014; new species), cultures and sequences are available.

Geopyxis (Pers.) Sacc. 1889, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *G. carbonaria* (Alb. & Schwein.) Sacc., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wang et al. (2016c; taxonomy), cultures and sequences are available.

Geosmithia Pitt 1979, *Hypocreales* genera *incertae sedis*, six species, type: *G. lavendula* (Raper & Fennell) Pitt, sexual morph unknown, saprobes, terrestrial, Ecuador, see Kolařík et al. (2011, 2015b; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wu et al. (2013b; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Geotrichum Link 1809 (= *Endylium* Clem. 1931; = *Fermentotrichon* E.K. Novák & Zsolt 1961; = *Galactomyces* Redhead & Malloch 1977; = *Geotrichum* Link 1809; = *Magnusiomyces* Zender 1926 [1925]; = *Oosporoidea* Sumst. 1913; = *Polymorphomyces* Coupin

1914; = *Zendera* Redhead & Malloch 1977 *fide* Species Fungorum 2017), *?Dipodascaceae*, *Saccharomycetales*, *Saccharomycetes*, three species, type: *G. candidum* Link, asexual morph present, saprophytic, soil, rotted fruit and other plant parts, tree exudates, worldwide, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gesasha Abdel-Wahab & Nagahama 2011, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, three species, type: *G. peditatus* Abdel-Wahab & Nagahama, saprobes, marine, Japan, asexual morph unknown, see Abdel-Wahab and Nagahama (2011; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype of the type: IMI 397958.

Gibbago E.G. Simmons 1986, *Dothideomycetes* genera *incertae sedis*, one species, type: *G. trianthemae* E.G. Simmons, sexual morph unknown, pathogens, terrestrial, USA, see Wijayawardene et al. (2012, 2014c; outline), Akhtar et al. (2013; pathogens, Pakistan), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Gibbera Fr. 1825 (= *Maireella* Syd. ex Maire 1908; = *Montagnina* Höhn. 1910; = *Pseudothia* Henn. 1900; = *Winteromyces* Speg. 1912 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, c. 28 species, type: *G. vaccinii* (Sowerby) Fr., sexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Hyde et al. (2013; excluded from *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are available.

Gibberidea Fuckel 1870 (= *Gibberinula* Kuntze 1898 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *G. visci* Fuckel, sexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Gibbosporina Elvebakk, Hong & P.M. Jørg. 2016, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 13 species, type: *G. boninensis* (Kurok.) Elvebakk & P.M. Jørg., lichenized, see Elvebakk et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Gibellia Sacc. 1885, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *G. dothideoides* Sacc. & Berl., asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Gibellina Pass. 1886 (= *Gibellia* Pass. 1886 *fide* Species Fungorum 2017), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, three species, type: *G. cerealis* (Pass.) Pass., asexual morph unknown, pathogens, terrestrial, Asia, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for type: MF-C22704 (unpublished).

Gibellula Cavara 1894 (= *Synsterigmatocystis* Costantin 1888 *fide* Species Fungorum 2017), *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *G. pulchra* Cavara, hyphomycetous, sexual morph unknown, parasitic, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Han et al. (2013; new species, China), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for *Gibellula leiopus*.

Gibellulopsis Bat. & H. Maia 1959, *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*, two species, type: *G. piscis* Bat. & H. Maia, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Han et al. (2013; new species, China), Kirk et al. (2013; genus accepted), Hirooka et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Gigantospora B.S. Lu & K.D. Hyde 2003, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *G. gigaspora* B.S. Lu & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Gilkeya M.E. Sm., Trappe & Rizzo 2007, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *G. compacta* (Harkn.) M.E. Sm. & Trappe, asexual morph unknown, biotrophic, terrestrial, worldwide, see Gladdish et al. (2010; DNA), Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Gilletiella Sacc. & P. Syd. 1899 (= *Epibotrys* Theiss. & Syd. 1915; = *Heterochlamys* Pat. 1895; = *Melanochlamys* Syd. & P. Syd. 1914 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *G. chusqueae* (Pat.) Sacc. & P. Syd., asexual morph coelomycetous (ascochyta-like), saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Gillotia Sacc. & Trotter 1913, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type:

G. orbicularis (Syd. & P. Syd.) Sacc. & Trotter, asexual morph asteromella-like, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mycosphaerellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Gilmaniella G.L. Barron 1964 (= *Adhogamina* Subram. & Lodha 1964 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, nine species, type: *G. humicola* G.L. Barron, hyphomycetous, from soil, dung, litter and bamboo, terrestrial, worldwide, see Dubey et al. (2011; new species, India), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Goh et al. (2013; new species, Malaysia), Kirk et al. (2013; genus accepted), ex-type strain of type: ATCC 16013 = O.A.C. 10076. = CBS 220.65 = IFO 9235, Holotype of type: O.A.C. 10076 (dried culture).

Gintarasia Kraichak, Lücking & Lumbsch 2013, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *G. megalophtalma* (Müll. Arg.) Kraichak, Lücking & Lumbsch, lichenized, terrestrial, tropical, see Kraichak et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Giulia Tassi 1904, *Ascomycota* genera *incertae sedis*, one species, type: *G. tenuis* (Sacc.) Tassi ex Sacc. & D. Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Glaphyriopsis B. Sutton & Pascoe 1987, *Ascomycota* genera *incertae sedis*, two species, type: *G. brunnea* B. Sutton & Pascoe, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Glarea Bills & Paláez 1999, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *G. lozoyensis* Bills & Peláez, sexual morph unknown, saprobes, terrestrial, Spain, see Wijayawardene et al. (2012; outline, listed as a genus in *Dermateaceae*), Chen et al. (2013d; DNA, phylogeny, accepted as a genus in *Helotiaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are available for type: ATCC 20868.

Glaucotrema Rivas Plata & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *G. glaucophaeum* (Kremp.) Rivas Plata & Lumbsch, lichenized, see Rivas Plata et al. (2012b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Glaxoa P.F. Cannon 1997, *Pleosporales* genera *incertae sedis*, one species, type: *G. pellucida* P.F. Cannon, asexual morph unknown, saprobes, terrestrial, UK, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in

Tubeufiaceae), Boonmee et al. (2014a; accepted as genus in *Pleosporales* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, cultures and sequences are unavailable.

Glaziella Berk. 1880, *Glaziellaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *G. vesiculos* Berk., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Glioannellodochium Matsush. 1989, *Ascomycota* genera *incertae sedis*, one species, type: *G. austriale* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: Matsushima, MFC-8A230, cultures and sequences are unavailable.

Glioblastocladium Matsush. 1989, *Ascomycota* genera *incertae sedis*, one species, type: *G. austriale* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: Matsushima, MFC-8A329, cultures and sequences are unavailable.

Gliocephalis Matr. 1899, *Pyxidiophoraceae*, *Pyxidiophorales*, *Laboulbeniomycetes*, two species, type: *G. hyalina* Matr., asexual morph unknown, parasitic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Gliocephalotrichum J.J. Ellis & Hesselt. 1962, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, 13 species, type: *G. bulbilium* J.J. Ellis & Hesselt., hyphomycetous, sexual morph unknown, in soil, associated with post-harvest fruit spoilage of several important tropical fruit crops, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2014, 2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: BPI 414619, ex-type of the type: CBS 242.62 = ATCC 22228 = IFO 9325 = IMI 096357 = MUCL 18575 = NRRL 2899 = QM 9007.

Gliocladopsis S.B. Saksena 1954 (= *Acontiopsis* Negru 1961; = *Tetracytum* Vanderw. 1945 *fide Species Fungorum* 2017), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, ten species, type *G. sagariensis* S.B. Saksena., soil-borne fungi, worldwide, see Seifert et al. (2011; morphology), Lombard and Crous (2012), Wijayawardene et al. (2012, 2017; outline), Lombard et al. (2015b; phylogeny), holotype of the type: CBS 199.55, ex-type of the type: CBS 199.55.

Gliomastix Guég. 1905 (= *Basitorula* G. Arnaud 1954 [1953]; = *Haplotrichella* G. Arnaud 1954 [1953]; = *Nematomyces* Faurel & Schotter 1966 [1965]; = *Torulina* Sacc. & D. Sacc. 1906; = *Torulopsis* Oudem. 1903 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, c. 25 species, type: *G. chartarum* (Cooke) Guég., hyphomycetous, sexual morph unknown, saprobes, in soil, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Summerbell et al. (2011; revision, phylogeny), Whitton et al. (2012a; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Globoea Bat. & H. Maia 1962, *Dothideomycetes* genera *incertae sedis*, one species, type: *G. toddaliae* Bat. & H. Maia, asexual morph unknown, saprobes, terrestrial, Uganda, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Globobotrys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *G. sanseviericola* (Crous & M.J. Wingf.) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Tanzania, see Lombard et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: CBS H-22220, CBS 138872.

Globoconidiopsis G.F. Sepúlveda, Pereira-Carv. & Dianese 2009, *Ascomycota* genera *incertae sedis*, one species, type: *G. cerradensis* G.F. Sepúlveda, Pereira-Carv. & Dianese, hyphomycetous, sexual morph unknown, superficial on leaf trichomes, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of type: UB Mycol. Col. 9926, cultures and sequences are unavailable.

Globoconidium G.F. Sepúlveda, Pereira-Carv. & Dianese 2009, *Ascomycota* genera *incertae sedis*, one species, type: *G. cerradense* G.F. Sepúlveda, Pereira-Carv. & Dianese, hyphomycetous, sexual morph unknown, superficial on leaf trichomes, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of type: UB Mycol. Col. 8387, cultures and sequences are unavailable.

Globonectria Etayo 2002, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *G. cochenensis* Etayo, asexual morph unknown, lichenicolous, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), holotype of the type: COL Etayo 15859, sequences are unavailable.

Globosphaeria D. Hawksw. 1990, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *G. jamesii* D. Hawksw., lichenicolous, terrestrial, Australia (Tasmania), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted),

Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: NSW LI4339, sequences are unavailable.

Globulina Speg. 1889, *Dothideomycetes* genera *incertae sedis*, one species, type: *G. toddaliae* Bat. & H. Maia, asexual morph unknown, saprobes, terrestrial, Uganda, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Gloeandromyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, five species (est. spp. 5+), type: *G. streblae* (Peyr.) Thaxt., asexual morph unknown, biotrophic, hosts are *Streblidae* (Diptera: Hippoboscidae), neotropics, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2015b, 2017a, b; DNA, review, taxonomy), ribosomal sequence data available but not for the type.

Gloeodiscus Dennis 1961, *Dothideomycetes* genera *incertae sedis*, one species, type: *G. nigrorufus* (Berk.) Dennis, asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline), Ariyawansa et al. (2013b; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Gloeohepnia Gyeln. 1935, *Gloeohepniaceae*, *Lichinales*, *Lichinomycetes*, five species, type: *G. turgida* (Ach.) Gyeln., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Gloeopeziza Zukal 1891, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, c. eight species, type: *G. rehmii* Zukal, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gloeosporidiella Petr. 1921, *Ascomycota* genera *incertae sedis*, one species, type: *G. ribis* (Lib.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gloeosporidina Petr. 1921, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *G. moravica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Gloeosporiella Cavara 1892, *Ascomycota* genera *incertae sedis*, one species, type: *G. rosicola* Cavara, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology),

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gloeotinia M. Wilson, Noble & E.G. Gray 1954 (= *Endoconidium* Prill. & Delacr. *fide* Johnston et al. 2014b), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *G. temulenta* (Prill. & Delacr.) M. Wilson, Noble & E.G. Gray, asexual morph *Endoconidium*, saprobe, terrestrial, India, Canary Islands, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available for unidentified species.

Gloiosphaera Höhn. 1902 (= *Scopula* G. Arnaud 1952 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *G. globuligera* Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Glomerilla Norman 1869, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *G. subtilis* Norman, lichenized, terrestrial, Norway, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Glomerobolus Kohlm. & Volk.-Kohlm. 1996, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *G. gelineus* Kohlm. & Volk.-Kohlm., saprobes, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMS, Kohlmeyer 5555, cultures and sequences are available.

Glomerulophoron Frisch, Ertz & G. Thor 2015, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *G. mauritiae* Frisch, Ertz & G. Thor, lichenized, see Frisch et al. (2015; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Gloniella Sacc. 1883, *Hysteriaceae*, *Hysterialales*, *Dothideomycetes*, twelve species, type: *G. sardoa* Sacc. & Traverso, asexual morph unknown, saprobes, terrestrial/aquatic, cosmopolitan, see Boehm et al. (2009a, b; overview of genus) Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Hysteriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Gloniopsis De Not. 1847, *Hysteriaceae*, *Hysterialales*, *Dothideomycetes*, seven species, type: *G. decipiens* De Not., asexual morph coelomycetous (aposporeia-like *fide* Lohman 1933a), saprobes, terrestrial, see Boehm et al. (2009a, b; overview of genus), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted),

Wijayawardene et al. (2014c; outline), sequence data indicate this genus is polyphyletic.

Glonium Muhl. ex Fr. 1823 (= *Glonium* Muhl. 1813; = *Solenarium* Spreng. 1827 *fide* Species Fungorum 2017), *Gloniaceae*, *Mytilinidiales*, *Dothideomycetes*, c. 13 species, type: *G. stellatum* Muhl., asexual morph asteromella-like, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Gloniaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Glutinium Fr. 1849 (= *Malacodermis* Bubák & Kabát 1912 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *G. exasperans* Fr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Glutinoglossum Hustad, A.N. Mill, Dentinger & P.F. Cannon 2013, *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, two species, type: *G. glutinosum* (Pers.) Hustad, A.N. Mill., Dentinger & P.F. Cannon, asexual morph unknown, from soil, terrestrial, worldwide, see Hustad et al. (2013; morphology, taxonomy, DNA, phylogeny), Kirk et al. (2013; genus accepted), holotype of the type: 910.261-767 (L 0110938 [Persoon Herb.], cultures and sequences are available: ANM 2231.

Glyphis Ach. 1814 (= *Glyphomyces* Cif. & Tomas. 1953; = *Gymnotrema* Nyl. 1858 *fide* Species Fungorum 2017), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, seven species, type: *G. cicatricosa* Ach., lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Glyphium Nitschke ex F. Lehmann 1886, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, three species, type: *G. dolabriiforme* (Wallr.) F. Lehmann., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Boehm et al. (2015; phylogeny, taxonomy), cultures and sequences are available.

Glypholecia Nyl. 1853 (= *Glypholeciomyces* Cif. & Tomas. 1953; = *Laureriella* Hepp 1867 *fide* Species Fungorum 2017), *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, one species, type: *G. candidissima* Nyl., lichenized, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Glyhopeltis Brusse 1985, *Psoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *G. eburina* Brusse, lichenized, Mediterranean Europe, South Africa, Brazil,

see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Glyphopsis Aptroot 2014, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *G. aurantioidisca* Aptroot, lichenized, see Aptroot (2014; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Gnomonia Ces. & De Not. 1863 (= *Melanopelta* Kirschst. 1939; = *Rehmiella* G. Winter 1883; = *Rostrocoronophora* Munk 1953 *fide* Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, c. 70 species, type: *G. vulgaris* Ces. & De Not., asexual morph unknown, saprobes, terrestrial, Europe, see Mejía et al. (2008; DNA, phylogeny), Sogonov et al. (2008; new species, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available but lacking for the type.

Gnomoniella Sacc. 1881, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, c. 70 species, type: *G. tubaeformis* (Tode) Sacc., asexual morph unknown, saprobes, terrestrial, Europe, see Sogonov et al. (2008; DNA, phylogeny), Singh et al. (2009; new species, India), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available but lacking for the type.

Gnomoniopsis Berl. 1893, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, c. 20 species, type: *G. chamaemori* (Fr.) Berl., asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Sogonov et al. (2008; reinstated the genus), Lumbsch and Huhndorf (2010; outline), Walker et al. (2010; new species, revision), Visentin et al. (2012; new species, pathogens), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Tibpromma et al. (2017b; reference specimen), cultures and sequences are available.

Godronia Moug. & Lév. 1846 (= *Cenangiella* Lambotte 1887; = *Cenangium* trib. *Scleroderris* Fr. 1822; = *Crumenula* De Not. 1864; *Scleroderris* (Fr.) Bonord. 1851 = *Sphaeronaema* Fr. *fide* Johnston et al. 2014b; Species Fungorum 2017), *Godroniaceae*, *Helotiales*, *Leotiomycetes*, c. 30 species, type: *G. muehlenbeckii* Moug. & Lév., asexual morph unknown, saprobes, terrestrial, worldwide, see de Gruyter et al. (2009; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available but lacking for type.

Godroniopsis Diehl & E.K. Cash 1929 (= *Sphaeronaema* Fr., *fide* Johnston et al. 2014b), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type:

G. quernea (Schwein.) Diehl & E.K. Cash, asexual morph ?*Sphaeronaema*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are unavailable.

Golovinomyces (U. Braun) Heluta 1988 (= *Euoidium* Y.L. Paul & J.N. Kapoor 1985; = *Erysiphe* sect. *Golovinomyces* U. Braun 1978 [1977] *fide* Rossman et al. 2016a; Species Fungorum 2017), *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, c. 63 species, type: *G. cichoracearum* (DC.) Heluta, asexual morphs hyphomycetous, biotrophic, obligate plant pathogens, terrestrial, worldwide, see Cunningham et al. (2010; phylogeny), Takamatsu et al. (2013a, b; phylogeny), Braun and Cook (2012; taxonomy), Rossman et al. (2016a; nomenclature), lectotype of type: G 00298360.

Gomphillus Nyl. 1855 (= *Baeopodium* Trevis. 1857; = *Mycetodium* A. Massal. 1856 *fide* Species Fungorum 2017), *Gomphillaceae*, *Ostrospores*, *Lecanoromycetes*, six species, type: *G. calycioides* (Delise ex Duby) Nyl., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Gonatobotrys Corda 1839 (see Species Fungorum 2017 for synonyms), *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, type: *G. simplex* Corda., hyphomycetous, sexual morph unknown, fungicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable, Index Fungorum (2017) listed Gonatobotrys as a synonym of *Melanospora* Corda.

Gonatobotryum Sacc. 1880 (= *Christiaster* Kuntze 1891; = *Gonatorrhodis* Clem. & Shear 1931; = *Gonatorrhodiella* Thaxt. 1891; = *Thomiella* C.W. Dodge 1935 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, four species, type: *G. fuscum* (Sacc.) Sacc., hyphomycetous, sexual morph unknown, fungicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for *G. apiculatum* (unpublished) but lacking for type.

Gonatophragmium Deighton 1969, *Acrospermaceae*, *Acrospermales*, *Dothideomycetes*, 15 species, type: *G. mori* (Sawada) Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014d; new species, DNA), Berger et al. (2015; new species, lichenicolous, Austria), cultures and sequences are available.

Gonatorrhodum Corda 1839, Ascomycota genera *incertae sedis*, one species, type: *G. speciosum* Corda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gondwania Søchting, Frödén & Arup 2013, Teloschistaceae, Teloschistales, Lecanoromycetes, four species, type: *G. cibrosa* (Hue) Søchting, Frödén & Arup, lichenized, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Goniopila Marvanová & Descals 1985, Ascomycota genera *incertae sedis*, one species, type: *G. monticola* (Dyko) Marvanová & Descals, hyphomycetous, sexual morph unknown, saprobes, aquatic, USA, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for type: CCM F-11686.

Goosia B. Song 2003, Englerulaceae, Dothideomycetes families *incertae sedis*, one species, type: *G. melastomatis* B. Song, asexual morph unknown, parasitic, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), Dai et al. (2014c; morphology), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Goosiella Morgan-Jones, Kamal & R.K. Verma 1986, Ascomycota genera *incertae sedis*, one species, type: *G. helicospora* Morgan-Jones, Kamal & R.K. Verma, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Goosiomycetes N.K. Rao & Manohar. 1989, Ascomycota genera *incertae sedis*, one species, type: *G. digitatus* N.K. Rao & Manohar., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Dubey and Neelima (2014; new species), cultures and sequences are unavailable.

Gordonomyces Crous & Marinc. 2011, Pleosporales, genera *incertae sedis*, Dothideomycetes, one species, type: *G. mucovaginatus* Crous & Marinc., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Crous et al. (2011d; morphology, DNA), Wijayawardene et al. (2012, 2014c, 2016a, b, c; outline, morphology, phylogeny), holotype and ex-type strains of the type: PREM 59592, CMW 22212.

Gorgadesia Tav. 1964, Roccellaceae, Arthoniales, Arthoniomycetes, one species, type: *G. mira* Tav.,

lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Gorgomyces M. Gönczöl & Révay 1985, Leotiomycetes genera *incertae sedis*, two species, type: *G. hungaricus* J. Gönczöl & Révay, hyphomycetous, sexual morph unknown, nematophagous, aquatic, Hungary, Spain, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Baschien et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), ex-type culture of the type: CCM F-12696.

Gorgoniceps (P. Karst.) P. Karst. 1871 (= Peziza sect. Gorgoniceps P. Karst. 1869 fide Species Fungorum 2017), Helotiidae genera *incertae sedis*, Leotiomycetes, three species, type: *G. aridula* (P. Karst.) P. Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gossypiothallon Aptroot 2014, Arthoniales genera *incertae sedis*, Arthoniomycetes, one species, type: *G. appendisporum* Aptroot, lichenized, see Aptroot (2014; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Govindua Bat. & H. Maia 1960, Dothideomycetes, genera *incertae sedis*, one species, type: *G. shoreae* Bat. & H. Maia, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Microthyriaceae), Wu et al. (2011; transferred to Asterinaceae), Hyde et al. (2013) and Hongsanan et al. (2014c) did not accept as Asterinaceae, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as a genus in Dothideomycetes genera *incertae sedis*), cultures and sequences are unavailable.

Gowardia Halonen, Myllys, Velmala & Hyvärinen 2009, Parmeliaceae, Lecanorales, Lecanoromycetes, two species, type: *G. nigricans* (Ach.) Halonen, Myllys, Velmala & Hyvärinen, lichenized, see Myllys et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Gracilistilbella Seifert 2000, Bionectriaceae, Hypocreales, Sordariomycetes, four species, type: *G. clavulata* (Mont.) Seifert, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Graddonia Dennis 1955, Helotiaceae, Helotiales, Leotiomycetes, seven species, type: *G. coracina* (Bres.) Dennis, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Dermateaceae), Hustad and Miller (2011; DNA, phylogeny), Raja et al. (2011a; DNA,

phylogeny), Kirk et al. (2013; genus accepted), cultures available for type: ILLS60491.

Graddonidiscus Raity. & R. Galán 1992, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *G. coruscatus* (Graddon) Raity. & R. Galán, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Grahamiella Spooner 1981, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *G. dryadis* (Nannf. ex L. Holm) Spooner, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Grallomyces F. Stevens 1918 (= *Ophiopodium* G. Arnaud 1954 [1953]; = *Phialetea* Bat. & Nascim. 1960 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, one species, type: *G. portoricensis* F. Stevens, hyphomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Grandibotrys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *G. pseudotheobromae* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Nepal, see Lombard et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: CBS H-22443, CBS 136170.

Grandigallia M.E. Barr, Hanlin, Cedeño, Parra & R. Hern. 1987, *Shiraiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *G. dictyospora* M.E. Barr, Hanlin, Cedeño, Parra & R. Hern., asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), Ariyawansa et al. (2013b; accepted as a genus in *Shiraiaceae*), Hyde et al. (2013; accepted as a genus in *Shiraiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Granulomanus de Hoog & Samson 1978, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *G. aranearium* (Petch) de Hoog & Samson, hyphomycetous, sexual morph unknown, parasitic on Arachnida, terrestrial, British Isles, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for *Gibellula leiopus*.

Granulopyrenis Aptroot 1991, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, six species, type:

G. antillensis Aptroot, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Graphidastra (Redinger) G. Thor 1991 (= *Dirina* sect. *Graphidastra* Redinger 1936 *fide Species Fungorum* 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, four species, type: *G. multiformis* (Mont. & Bosch) G. Thor, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Graphiopsis Trail 1889, *Cladosporiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *G. chlorocephala* (Fresen.) Trail, hyphomycetous, terrestrial, Asia, Europe, North America, New Zealand, see Braun et al. (2008; nomenclature), Crous et al. (2009d; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), neotype and ex-neotype strain of the type: HAL 1924 F; CBS 121523 = CPC 11969.

Graphiothecium Fuckel 1870 (= *Fresenia* Fuckel 1866 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, six species, type: *G. fresenii* Fuckel, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Graphis Adans. 1763 (see *Species Fungorum* 2017 for synonyms), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, c. 500 species, type: *G. scripta* (L.) Ach., asexual morph unknown, lichenized, terrestrial, mostly tropical, see Lücking et al. (2008a, 2009, 2016b; key, Costa Rica, classification), Archer (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; taxonomy), Kirk et al. (2013; genus accepted), Barcenas Peña et al. (2014; key), Kraichak et al. (2015; taxonomy), sequences are available.

Graphium Corda 1837, *Graphiaceae*, *Microascales*, *Sordariomycetes*, three species, type: *G. penicilliodes* Corda, hyphomycetous, sexual morph unknown, saprobes or from soil terrestrial, worldwide, see Cruywagen et al. (2010; new species, South Africa), Paciura et al. (2010b; new species, China), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Kolařík et al. (2015b; new species; Costa Rica), Wu et al. (2015; new species, China), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type strain of the type: JCM 1049.

Graphostroma Piroz. 1974, *Graphostromataceae*, *Xylariales*, *Sordariomycetes*, one species, type: *G. platystoma* (Schwein.) Piroz., asexual morph hyphomycetous (nodulisporium-like), saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010;

outline), Kirk et al. (2013; genus accepted), Koukol et al. (2014; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures available for the type: AFTOL-ID 1249.

Graphyllum Clem. 1901, *Hysteriales* genera *incertae sedis*, eleven species, type: *G. chloës* Clem., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Diademaceae*), Hyde et al. (2013) did not accept as *Diademaceae*, Kirk et al. (2013; genus accepted), Almeida et al. (2014b; DNA, new species, phylogenetic analyses showed that *G. caracolinense* D.A.C. Almeida et al. belongs to *Hysteriales*), Wijayawardene et al. (2014c; accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are available only for *G. caracolinense*.

Greeneria Scribn. & Viala 1887 (= *Frankia* Brunch. 1886; = *Frankiella* Speschnew 1900 *fide* Species Fungorum 2017), *Diaporthales* genera *incertae sedis*, three species, type: *G. fuliginea* Scribn. & Viala, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy, phylogeny), Tangthirasunun et al. (2014a; DNA, phylogeny), cultures and sequence available for *G. saprophytica* (MFLUCC 12-0298 *fide* Tangthirasunun et al. 2014a, b).

Gregarithecum Kaz. Tanaka & K. Hiray. 2015, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *G. curvisporum* Kaz. Tanaka & K. Hiray., asexual morph unknown, saprobes, terrestrial, Japan, see Tanaka et al. (2015; morphology, phylogeny), holotype and ex-type of the type: HHUF 30134, CBS 139688.

Gregatothecium L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *G. humicola* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Papua New Guinea, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22445, CBS 205.96.

Gregrella Lumbsch 2005, *Arctomiaceae*, *Arctomiales*, *Lecanoromycetes*, one species, type: *G. humida* (Kullh.) Lumbsch, asexual morph coelomycetous, lichenized, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Miądlikowska et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Gremmenia Korf 1962, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *G. pini-cembrae* (Rehm) Crous, asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Crous et al. (2014b; DNA, phylogeny), cultures and sequences are available.

Gremmeniella M. Morelet 1969 (= *Brunchorstia* Erikss 1891 *fide* Johnston et al. 2014b), *Godroniaceae*, *Helotiales*,

Leotiomycetes, three species, type: *G. abietina* (Lagerb.) M. Morelet, asexual morph *Brunchorstia*, pathogens, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Laflamme and Smerlis (2012; DNA, new species, Canada), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available.

Griggsia F. Stevens & Dalbey 1919, *Dothideomycetes* genera *incertae sedis*, one species, type: *G. cyathea* F. Stevens & Dalbey, asexual morph unknown, saprobes, terrestrial, West Indies, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014e; taxonomy), cultures and sequences are unavailable.

Grimmicola Döbbeler & Hertel 1983, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *G. parasiticus* Döbbeler & Hertel, asexual morph unknown, saprobes, terrestrial, Marion Island, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Griphosphaerioma Höhn. 1918 (= *Labridella* Brenckle 1929 *fide* Maharachchikumbura et al. 2015b; Wijayawardene et al. 2016b), *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, two species, type: *G. symphoricarpi* (Ellis & Everh.) Höhn. ex Petr., asexual morphs *Labridella*, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, nomenclature), Wijayawardene et al. (2016b; taxonomy, nomenclature), Rossman et al. (2016a) proposed to protect *Labridella* over *Griphosphaerioma*, cultures and sequences are unavailable.

Groenewaldozyma Kurtzman 2016, *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, three species, type: *G. auringiensis* (Santa María) Kurtzman, saprophytic, worldwide, see Kurtzman (2016; taxonomy), cultures and sequences are available.

Groenhiella Jorg., E.B.G. Jones & S.T. Moss 1983, *Nitschiaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *G. bivestia* Jørg. Koch, E.B.G. Jones & S.T. Moss, asexual morph unknown, marine, on wood often associated with sand, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline) cultures and sequences are unavailable.

Grosmannia Goid. 1936, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, one species, type: *G. penicillata* (Grosmann) Goid., holomorph, asexual morph hyphomycetous, associated with bark beetles, terrestrial, worldwide, see Masuya et al. (2009; new

species), Lumbsch and Huhndorf (2010; outline), Duong et al. (2012; new species, phylogeny), Linnakoski et al. (2012, 2016; new species, DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures available for the type: CMW43738.

Groveolopsis Boedijn 1951 (= *Septoriopsis* Höhn. 1920 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, six species, type: *G. pandani* (Höhn.) Boedijn, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Grovesia Dennis 1960, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *G. pulchella* Dennis, asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Grovesiella M. Morelet 1969, *Tympanidaceae*, *Phacidiales*, *Leotiomycetes*, one species, type: *G. abieticola* (Zeller & Goodd.) M. Morelet & Gremmen, asexual morph unknown, saprobes, terrestrial, USA, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available for the type: ZT 91110.

Grovesinia M.N. Cline, J.L. Crane & S.D. Cline 1983 (= *Hinomyces* Narumi & Y. Harada 2006 *fide* Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *G. pyramidalis* M.N. Cline, J.L. Crane & S.D. Cline, asexual morph *Hinomyces*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature).

Guanomyces M.C. González, Hanlin & Ulloa 2000, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *G. polythrix* M.C. González, Hanlin & Ulloa, asexual morph unknown, saprobes, terrestrial, Mexico, Trinidad, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available (see Gonzalez et al. 2000).

Gudelia Henssen 1995, *Gloeoheppiaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *G. mexicana* Henssen, lichenized, Mexico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Guedea Rambelli & Bartoli 1978, *Ascomycota* genera *incertae sedis*, six species, type: *G. sacra* Rambelli & Bartoli, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Guelichia Spieg. 1886, *Ascomycota* genera *incertae sedis*, six species, type: *G. paradoxa* Spieg., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012; outline, accepted as a genus in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b) and Wijayawardene et al. (2014c) did not accept as a genus in *Tubeufiaceae*, cultures and sequences are unavailable.

Guestia G.J.D. Sm. & K.D. Hyde 2001, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *G. gonetropospora* G.J.D. Sm. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Guilliermondia Boud. 1904, *Sordariaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *G. saccoboloides* Boud., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Guttulispora Thambug., Qing Tian & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *G. crataegi* Qing Tian, Thambug., Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Thambugala et al. (2015b; morphology, phylogeny), holotype of type: MFLU 14–0584, ex-type strains of type: MFLUCC 13–0442.

Gyalecta Ach. 1808 (= *Gyalectomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Phialopsis* Körb. 1855; = *Secoliga* Norman 1853 [1852]; = *Volvaria* DC. 1805 *fide* Species Fungorum 2017), *Gyalectaceae*, *Ostropales*, *Lecanoromycetes*, 50 species, type: *G. cupularis* (Hedw.) Schaer., lichenized, terrestrial (corticulous, saxicolous, terricolous), mainly temperate, see Baloch et al. (2010, 2013; taxonomy), Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; DNA), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; DNA), Aptroot and Moon (2015; new species), Resl et al. (2015; DNA), Lücking et al. (2017a; classification), sequences are available but lacking for the type.

Gyalectaria Lumbsch, Kalb & Schmitt 2010, *Coccotremataceae*, *Pertusariales*, *Lecanoromycetes*, three species, type: *G. jamesii* (Kantvilas) I. Schmitt, Kalb & Lumbsch, lichenized, terrestrial (corticulous), Chile, New Guinea, North America, Tasmania, from lowland to subalpine, see Schmitt et al. (2010; taxonomy), Resl

et al. (2015; DNA), Lücking et al. (2017a; classification), sequences are available.

Gyalectidium Müll. Arg. 1881 (= *Gonothecis* Clem. 1909; = *Gonothecium* (Vain.) Clem. & Shear 1931; = *Lecidea* sect. *Gonothecium* Vain. 1890; = *Lopadiopsidomyces* Cif. & Tomas. 1953; = *Lopadiopsis* Vain. 1896; = *Tauromyces* Cavalc. & A.A. Silva 1972 *fide* Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, 52 species, type: *G. filicinum* Müll. Arg., lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Gyalidea Lettau ex Vězda [nom. cons.] 1966 (= *Aglaothecium* Groenh. 1962; = *Cappellettia* Tomas. & Cif. 1952 *fide* Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, 50 species, type: *G. lecideopsis* (A. Massal.) Lettau ex Vězda, lichenized, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Gyalideopsis Vězda [nom. cons.] 1972, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, 95 species, type: *G. peruviana* G. Merr. ex Vězda, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Gyalolechia A. Massal. 1852, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 30 species, type: *G. aurea* (Schaer.) A. Massal., lichenized, see Arup et al. (2013; taxonomy), Kondratyuk et al. (2014; taxonomy), Jaklitsch et al. (2016a; outline), Lücking et al. (2017a; classification), sequences are available.

Gymnascella Peck 1884 (= *Pseudoarachniotus* Kuehn 1957 *fide* Species Fungorum 2017), *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, 14 species, type: *G. aurantiaca* Peck, asexual morphs unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Gymnoascoideus G.F. Orr, K. Roy & G.R. Ghosh 1977, *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *G. petalosporus* G.F. Orr, K. Roy & G.R. Ghosh, asexual morphs unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Gymnoascus Baran. 1872 (= *Macronodus* G.F. Orr 1977; = *Tripedotrichum* G.F. Orr & Kuehn 1964 *fide* Species Fungorum 2017), *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, eight species, type: *G. reessii* Baran., asexual morphs unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are available.

Gymnoderma Nyl. 1860 (= *Gymnodermatomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *G. coccocarpum* Nyl., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Gymnographopsis C.W. Dodge 1967, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *G. chilena* C.W. Dodge, lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rivas Plata et al. (2012a; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Gymnohydnotrya B.C. Zhang & Minter 1989, *Discinaceae*, *Pezizales*, *Pezizomycetes*, three species, type: *G. australiana* B.C. Zhang & Minter, asexual morphs unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Gymnostellatospora Udagawa, Uchiy. & Kamiya 1993, *Myxotrichaceae*, *Leotiomycetes*, families *incertae sedis*, two species, type: *G. japonica* Udagawa, Uchiy. & Kamiya, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype of the type: BF 24290, cultures available for type: UAMH 9239 (*fide* Rice and Currah 2006).

Gymnoxyphium Cif., Bat. & I.J. Araújo 1963, *Ascomycota* genera *incertae sedis*, six species, type: *G. malum* Bat., Nascim. & Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gynonectria Döbbeler 2012, *Hypocreales* genera *incertae sedis*, six species, type: *G. intraspora* Döbbeler, asexual morph unknown, saprobes, terrestrial, South America, see Döbbeler (2012; outline), cultures and sequences are unavailable.

Gyoerffyella Kol 1928 (= *Ingoldia* R.H. Petersen 1962 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, nine species, type: *G. tatraica* Kol, hyphomycetous, sexual morph unknown, saprobes, pathogens, saprobes, aquatic, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Baschien et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures unavailable for the type, ex-type strains available for *G. entomobryooides* (CBS268.63 *fide* Baschien et al. 2013), *G. tricapillata* (CBS 451.64 *fide* Baschien et al. 2013).

Gypsoplaea Timdal 1990, *Gypsoplacaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *G. macrophylla* (Zahlbr.) Timdal, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Gyrocollema Vain. 1929 (= *Ginzbergerella* Zahlbr. 1931 *fide* Species Fungorum 2017), *Lichinaceae*, *Lichinales*, *Lichinomycetes*, two species, type: *G. scyphuliferum* Vain., with ascocarps, lichenized (Nostoc), terrestrial (saxicolous), North and Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Gyrographa Ertz & Tehler 2015, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *G. gyrocarpa* (Flot.) Ertz & Tehler, sexual morph with knot-like or clustered-gyrose to lirelliform ascocarps, asexual morph coelomycetous, lichenized, terrestrial, worldwide, see Ertz et al. (2015b, c; taxonomy, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Gyromitra Fr. 1849 (see Species Fungorum 2017 for synonyms), *Discinaceae*, *Pezizales*, *Pezizomycetes*, 20 species, type: *G. esculenta* (Pers.) Fr., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Methven et al. (2013; DNA, phylogeny), cultures available for the type: AFTOL-ID 5017.

Gyronactis Ertz & Tehler 2015, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *G. asiatica* Ertz & Tehler, sexual morph with apothecial ascocarps, asexual morph coelomycetous, lichenized, terrestrial, tropical, see Ertz et al. (2015b, c; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Gyrophthorus Hafellner & Sancho 1990, *Ascomycota* genera *incertae sedis*, two species, type: *G. perforans* Hafellner & Sancho, asexual morph unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gyrostroma Naumov 1914 (= *Kaskaskia* Born & J.L. Crane 1972 *fide* Species Fungorum 2017), *Diaporthales* genera *incertae sedis*, one species, type: *G. sinuosum* Naumov, sexual morph unknown, saprobes, terrestrial, Europe, see Hirooka et al. (2012; morphology, accepted as a genus in *Diaporthales*, genera *incertae sedis*), Wijayawardene et al. (2012; accepted as a genus in *Hypocreales*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Diaporthales*, genera *incertae*

sedis), cultures and sequences are unavailable for the type: thus *Gyrostroma s. str.* uncertain.

Gyrothrix Corda 1842 (= *Campsotrichum* sect. *Gyrothrix* Corda 1839; = *Peglonia* Goid. 1937 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, c. 22 species, type: *G. podosperma* (Corda) Rabenh, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Rambelli (2011; morphology), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Gyrotrema Frisch 2006, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *G. sinuosum* (Sipman) Frisch, with ascocarps, lichenized (green algae), terrestrial (corticicolous), Central and South America see Rivas Plata et al. (2008; outline), Kalb (2009; outline), Lumbsch and Huhndorf (2010; outline), Lumbsch et al. (2011; outline); Cáceres et al. (2014; DNA), Kraichak et al. (2014; DNA), Lücking et al. (2017a; classification), sequences are available but lacking for type.

Hadroneema Syd. & P. Syd. 1909, *Ascomycota* genera *incertae sedis*, four species, type: *H. orbiculare* Syd. & P. Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, China, UK see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hadrospora Boise 1989, *Trematosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *H. fallax* (Mouton) Boise, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Phaeosphaeriaceae*), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; accepted as a genus in *Trematosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hadrosporium Syd. 1938, *Ascomycota* genera *incertae sedis*, two species, type: *H. fraserianum* Syd., hyphomycetous, saprobes, terrestrial, Australia, New Zealand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hadrotrichum Fuckel 1865 (= *Microbasidium* Bubák & Ranj. 1914 *fide* Species Fungorum 2017), *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *H. phragmitis* Fuckel, hyphomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline),

cultures and sequences are available for *H. pyrenaicum* (YMJ 320 fide Hsieh et al. 2005).

Haematomma A. Massal. 1852 (= *Lepadolemma* Trevis. fide Species Fungorum 2017), *Haematommataceae*, *Lecanorales*, *Lecanoromycetes*, c. 40 species, type: *H. vulgare* A. Massal., with ascomata, lichenized (*Pseudotreobuxia*), terrestrial (corticulous, lignicolous or saxicolous), cosmopolitan, but most diverse in the tropics, see Brodo et al. (2008; outline and chemistry), Ekman et al. (2008; DNA), Kalb et al. (2008; DNA), Lumbsch et al. (2008; DNA), Stocker-Wörgötter et al. (2009; chemistry, mycobionts), Lumbsch and Huhndorf (2010; outline), Miądlikowska et al. (2014a; DNA), Lücking et al. (2017a; classification), sequences are available but lacking for the type.

Haematomyxa Sacc. 1884, *Ascomycota genera incertae sedis*, two species, type: *H. vinosa* (Cooke & Ellis) Sacc., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hafellia Kalb, H. Mayrhofer & Scheid. 1986, *Physciaceae*, *Teloschistales*, *Lecanoromycetes*, 22 species, type: *H. leptoclinoides* (Nyl.) Scheid. & H. Mayrhofer, with ascomata, lichenized, (green algae), terrestrial (corticulous, lignicolous, rare saxicolous), tropical, subtropical and temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2015f; new species), sequences available but lacking for the type species.

Hagleromyces Sousa, Morais, Lachance & Rosa 2014, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *H. auroranensis* Sousa, asexual reproduction by multilateral budding, saprophytic, from water tanks (phytotelmata) of *Bromelia karatas* in the cerrado ecosystem of Aurora do Tocantins, Brazil, see Sousa et al. (2014; taxonomy), cultures and sequences are available.

Haiyangia K.L. Pang & E.B.G. Jones 2008, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *H. salina* (Meyers) K.L. Pang & E.B.G. Jones, asexual morph unknown, saprobes, marine, worldwide, see Pang et al. (2008b; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Halazon Abdel-Aziz, Abdel-Wahab & Nagah. 2010, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, two species, type: *H. meliae* Abdel-Aziz, Abdel-Wahab & Nagah., coelomycetous, sexual morph unknown, saprobes, marine, worldwide, see Abdel-Wahab et al. (2010; morphology, phylogeny), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al.

(2015b, 2016b; outline), holotype and ex-type of the type: IMI 397964, MF819.

Halbania Racib. 1889, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, three species, type: *H. cyathearum* Racib., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, keys), Wijayawardene et al. (2014c; outline), placement in *Asterinaceae* supported by morphology, holotype of type: ZT Myc 1488, cultures and sequences are unavailable.

Halecania M. Mayrhofer 1987, *Catillariaceae*, *Lecanorales*, *Lecanoromycetes*, 22 species, type: *H. alpivaga* (Th. Fr.) M. Mayrhofer, with ascomata, lichenized (green algae), terrestrial (saxicolous, corticolous, lichenicolous or growing on mosses and plant debris), cosmopolitan, see Ekman et al. (2008; DNA), van den Boom (2009; new species), Andreev (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lendemer and Hodkinson (2013; taxonomy), Aptroot and Moon (2015; new species), Kondratyuk (2015; new species), Lücking et al. (2017a; classification), sequences are available for type: AY756368 (LSU).

Halegrapha Rivas Plata & Lücking 2011, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, eight species, type: *H. chimaera* Rivas Plata & Lücking, asexual morph unknown, lichenized, terrestrial, tropical, Lücking et al. (2011; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Halenospora E.B.G. Jones 2009, *Leotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *H. varia* (Anastasiou) E.B.G. Jones, hyphomycetous, sexual morph unknown, saprobe, marine, worldwide, see Jones et al. (2009b, 2015; outline, phylogeny), Abdel-Wahab et al. (2010; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Haleomyces D. Hawksw. & Essl. 1993, ?*Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *H. oropogonicola* D. Hawksw. & Essl., lichenicolous, terrestrial, Mexico, asexual morph unknown, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; outline), holotype of type: K(M)-IMI 3551875.

Haligena Kohlm. 1961, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *H. elaterophora* Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al.

(2015b, 2016b; outline), cultures and sequences are available, holotype of type: IMI 297765.

Halodiatripe Dayarathne & K.D. Hyde 2016, *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, three species, type: *H. salinicola* Dayarathne & K.D. Hyde, saprobes, found on decaying wood at mangrove stands, Thailand, see Dayarathne et al. (2016b; taxonomy, phylogeny), holotype of type: MFLU 15-0179.

Halographis Kohlm. & Volkm.-Kohlm. 1988, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *H. runica* Kohlm. & Volkm.-Kohlm., endolithic lichenoid from corals and snails, marine, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Lücking et al. (2017a; classification), sequences unavailable.

Haloguignardia Cribb & J. W. Cribb 1956, *Lulworthiaceae*, *Lulorthiales*, *Sordariomycetes*, five species, type: *H. decidua* Cribb & J.W. Cribb, asexual morph unknown, endophytes or parasites, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Halojulella Suetrong, K.D. Hyde & E.B.G. Jones 2013, *Halojullaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *H. avicenniae* (Borse) Suetrong, K.D. Hyde & E.B.G. Jones, asexual morph coelomycetous, saprobes, on woody substrates in mangrove habitats, worldwide, sequence data available, see Ariyawansa et al. (2013a, 2014a, c; phylogeny, morphology), Hyde et al. (2013; phylogeny, morphology, outline), Abdel-Wahab et al. (2014; diversity), Gnavi et al. (2014, phylogeny), Wijayawardene et al. (2014c; outline), Jones et al. (2015; classification, marine), Liu et al. (2015a; phylogeny), holotype of type: IMI 304217, cultures and sequences are available.

Halokirschsteiniothelia Boonmee & K.D. Hyde 2012, *Dothideomycetes* genera *incertae sedis*, one species, type: *H. maritima* (Linder) Boonmee & K.D. Hyde, asexual morph unknown, saprobes, marine, USA, see Boonmee et al. (2012; taxonomy, phylogeny; treated as a genus in *Mytilinidiaceae*), Hyde et al. (2013; not accepted as a genus in *Mytilinidiaceae*), Wijayawardene et al. (2014c; listed as *Dothideomycetes* genera *incertae sedis*), cultures available for the type: CBS 221.60.

Halomassarina (Kohlm. & Volkm.-Kohlm.) Suetrong, Sakay., E.B.G. Jones, Kohlm., Volkm.-Kohlm. & C.L. Schoch 2009, *Trematosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *H. thalassiae* (Kohlm. & Volkm.-Kohlm.) Suetrong, Sakay., E.B.G. Jones, Kohlm., Volkm.-Kohlm. & C.L. Schoch, asexual morph unknown, saprobes, marine, worldwide, see Suetrong et al. (2009; phylogeny, morphology), Schoch

et al. (2009; phylogeny), Alias et al. (2010; diversity), Lumbsch and Huhndorf (2010; outline), Zainuddin et al. (2010; antimicrobial activities), Sakayaroj et al. (2011b; diversity), Boonmee et al. (2012; phylogeny), Dai et al. (2012; phylogeny), Lee et al. (2012; checklist), Jones et al. (2012; phylogeny), Zhang et al. (2012e; morphology, phylogeny), Borse et al. (2013; checklist), Hyde et al. (2013; morphology, phylogeny), Ahmed et al. (2014a; phylogeny), Gnavi et al. (2014; phylogeny), Shearer et al. (2014; phylogeny), Wijayawardene et al. (2014c; outline), Singtripop et al. (2015; phylogeny), Sarma (2015; key), holotype and ex-isotype strain of type: J.K. 4804a, J.K.4804b.

Halonectria E.B.G. Jones 1965, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *H. milfordensis* E.B.G. Jones, asexual morph unknown, marine, on wood, mostly temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Haloplaca Arup, Søchting & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *H. britannica* (R. Sant.) Arup, Frödén & Søchting, lichenized (with green algae), saxicolous, corticolous, lignicolous, lichenicolous, terrestrial, United Kingdom, Greece, Morocco, Turkey, see Vondrák et al. (2012; DNA as *Caloplaca suaesae* s. l.), Arup et al. (2013, DNA), Lücking et al. (2017a; classification), sequences are available for all species KC179292 (ITS) (*fide* Arup et al. 2013).

Halorosellinia Whalley, E.B.G. Jones, K.D. Hyde & Laessøe, 2000, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *H. oceanica* (S. Schatz) Whalley, E.B.G. Jones, K.D. Hyde & Laessøe, asexual morph unknown, tropical, on wood especially mangrove substrates, sequence data for both species, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; morphology, phylogeny), Jones et al. (2015; outline), cultures and sequences are available.

Halosarpheia Kohlm. & E. Kohlm. 1977, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, polyphyletic, four species, type: *H. fibrosa* Kohlm. & E. Kohlm., asexual morph unknown, marine, freshwater, saprobes on herbaceous material, wood, widespread, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jones et al. (2015; outline), holotype of the type: NY J.K. 3726a, cultures available for type: AFTOL-ID 409, CBS 197.60.

Halosphaeria Linder 1944, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, c. 20 species, type:

H. appendiculata Linder, asexual morph unknown, saprobes, marine, temperate, subtropics, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), sequence are available.

Halosphaeriopsis T.W. Johnson 1958 (= *Culcitalna* Meyers & R.T. Moore *fide* Maharachchikumbura et al. 2016b), *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *H. mediosetigera* (Cribb & J.W. Cribb) T.W. Johnson, asexual morph *Trichocladium acharospora*, saprobes, marine, temperate, subtropics, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), sequence are available.

Halothia Kohlm. 1963, *Halothiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *H. posidoniae* (Durieu & Mont.) Kohlm., asexual morph coelomycetous, saprobes, marine habitats, Mediterranean, see Suetrong et al. (2009; phylogeny, morphology), Jones et al. (2009b; classification), Zhang et al. (2012e, 2013d; morphology, phylogeny), Hyde et al. (2013; morphology, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Jones et al. (2015; classification).

Halysiomyces E.G. Simmons 1981, *Ascomycota* genera *incertae sedis*, one species, type: *H. saxatilis* E.G. Simmons, hyphomycetous, epiphyte, free living, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hamatocanthoscypha Svrček 1977 (= *Debaryoscyphus* Arendh. & R. Sharma 1986; = *Uncinia* Velen. 1934; = *Unciniella* K. Holm & L. Holm 1977 *fide* Species Fungorum 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *H. laricionis* (Velen.) Svrček, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), Jaklitsch et al. (2016a; classification), cultures available for the type: TNS-F13530.

Hamigera Stolk & Samson 1971 (= *Merimbla* Pitt 1979 *fide* Houbraken and Samson 2011), *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, two species, type: *H. avellanea* Stolk & Samson, asexual morph formerly named in *Merimbla*, saprobes, terrestrial, temperate, see Peterson et al. (2010; phylogeny), Houbraken and Samson (2011; phylogeny, nomenclature), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Hanliniomyces Raja & Shearer 2008, *Sordariomycetes* genera *incertae sedis*, one species, type: *H. hyaloapicalis*

Raja & Shearer, asexual morph unknown, freshwater, northern and southern Florida, see Raja and Shearer (2008; taxonomy), Raja et al. (2009b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: R037-3, ILL 40116, cultures and sequences are unavailable.

Hanseniaspora Zikes 1912 (= *Kloeckera* Janke 1928; see Species Fungorum 2017 for additional synonyms), *Saccharomycodaceae*, *Saccharomycetales*, *Saccharomycetes*, 15 species, type: *H. valbyensis* Klöcker, asexual reproduction is by bipolar budding on a broad base (bud-fission), saprophytic, in soil, fresh water, sea water, rotting plants, fruit, fruit juices, fruit flies, grape must, beer, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Hansfordia S. Hughes 1951, *Ascomycota* genera *incertae sedis*, seven species, type: *H. ovalispora* S. Hughes, hyphomycetous, saprobes, terrestrial, marine, China, UK, Spain, and India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hansfordiella S. Hughes 1951, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, eight species, type: *H. asterinearum* S. Hughes, hyphomycetous, saprobes, on fungi, terrestrial, wide spread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hansfordiellopsis Deighton 1960, *Dothideomycetes* genera *incertae sedis*, one species, type: *H. aburiensis* Deighton, hyphomycetous, sexual morph unknown, saprobes, lichenicolous, terrestrial, Africa, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: IMI 4778, cultures and sequences are unavailable.

Hansfordiopeltis Bat. & C.A.A. Costa 1956, *Ascomycota* genera *incertae sedis*, five species, type: *H. congoensis* Bat. & C.A.A. Costa, coelomycetous, saprobes, terrestrial, Congo and Brazil, see Wijayawardene et al. (2012, 2017), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hansfordiopeltopsis M.L. Farr 1986, *Ascomycota* genera *incertae sedis*, one species, type: *H. amazonensis* M.L. Farr, coelomycetous, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hansfordiopsis Bat. 1959, *Dothideomycetes* genera *incertae sedis*, one species, type: *H. phaeospora* (Hansf.) Bat., asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline,

accepted as a genus in *Micropeltidaceae*), Hyde et al. (2013; did not accept as a genus *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, *Dothideomycetes* genera *incertae sedis*), holotype of the type: PDD 6788 (as *Microthyriella phaeospora* Hansf.), cultures and sequences are unavailable.

Hapalocystis Auersw. ex Fuckel 1863, *Sydiomycetaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *H. berkeleyi* Auersw. ex Fuckel, asexual morphs coelomycetous (phoma-like), saprobes, terrestrial, Europe, North America, see de Silva et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Liu et al. (2015a; DNA, phylogeny), Maharachchikumbura et al. (2016b; outline, phylogeny), Senanayake et al. (2017, placement), cultures available for the type: AR 3851.

Haplosphaeria Syd. 1908, *Ascomycota* genera *incertae sedis*, one species, type: *H. deformans* (Syd.) Syd., coelomycetous, pathogens, terrestrial, Germany, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Haploriopsis Oudem. 1903, *Ascomycota* genera *incertae sedis*, two species, type: *H. fagicola* Oudem., hyphomycetous, saprobes, terrestrial, India, Holland, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Haplobasidion Erikss. 1889, *Ascomycota* genera *incertae sedis*, three species, type: *H. thalictri* Erikss., hyphomycetous, saprobes, pathogens, terrestrial, Sweden, Asia, Malaysia, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Haplodina Zahlbr. 1930, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, three species, need typification, lichenized, terrestrial (corticicolous, saxicolous), China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Haplographium Berk. & Broome 1859, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, 15 species, type: *H. delicatum* Berk. & Broome, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for type species FMR 11585, TNS-F17834.

Hapolepis Syd. 1925, *Ascomycota* genera *incertae sedis*, three species, type: *H. polyadelpha* Syd., coelomycetous, saprobes, terrestrial, Sri Lanka, Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Hapoloma Trevis. 1857, *Lecanoromycetes* genus *incertae sedis*, one species, type: *H. fraudulentum* Trevis., lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not accept), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Hapomyces Thaxt. 1893, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *H. californicus* Thaxt., asexual morph unknown, biotrophic, North America, Europe, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species, Ecuador), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Haplopeltheca Bat., J.L. Bezerra & Cavalc. 1963 (= *Phaeostomiopeltis* Bat. & Cavalc. 1963 *fide Species Fungorum* 2017), *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *H. hyleensis* Bat., J.L. Bezerra & Cavalc., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Haplophyse Theiss. 1916, *Rhytidomatales* genera *incertae sedis*, *Leotiomycetes*, ten species, type: *H. oahuensis* Theiss., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hapsidascus Kohlm. & Volk.-Kohlm. 1991, *Sordariomycetes* genera *incertae sedis*, one species, type: *H. hadrus* Kohlm. & Volk.-Kohlm., asexual morph unknown, marine, on mangrove wood and salt marsh sedge, tropical, see Jones et al. (2009b, 2015; marine, outline), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Hapsidospora Malloch & Cain 1970, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *H. irregularis* Malloch & Cain, asexual morph hyphomycetous (acremonium-like), saprobes, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: ATCC 22087.

Haptocara Drechsler 1975, *Ascomycota* genera *incertae sedis*, one species, type: *H. latirostrum* Drechsler, hyphomycetous, saprobes, insect-trapping, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Haptospora G.L. Barron 1991, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *H. appendiculata* G.L. Barron, hyphomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, New Zealand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Haradamyces Masuya, Kusunoki, Kosaka & Aikawa 2009, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *H. foliicola* Masuya, Kusunoki, Kosaka & Aikawa, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: TFM FPH7930, cultures available for the type: MAFF 411026.

Harea Sacc. & P. Syd. 1913, *Ascomycota* genera *incertae sedis*, one species, type: *H. japonica* Sacc. & P. Syd., asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type of the type: DAOM 167533, MAFF 411026.

Harknessia Cooke 1881 (= *Caudosporella* Höhn. 1914; = *Mastigonetron* Kleb. 1914 *fide* Species Fungorum 2017), *Harknessiaceae*, *Diaporthales*, *Sordariomycetes*, c. 50 species, type: *H. eucalypti* Cooke, holomorph with wuestneia-like sexual morphs, pathogens, saprobes, terrestrial, worldwide, have sequence data, see Crous et al. (2012f; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Wijayawardene et al. (2012, 2016b, 2017; morphology, phylogeny), Rossman et al. (2016a; new combination), cultures and sequences are available.

Harknessiella Sacc. 1889, *Dothideomycetes* genera *incertae sedis*, one species, type: *H. purpurea* (W. Phillips & Harkn.) Sacc., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Harmoniella V.N. Boriss. 1981, *Ascomycota* genera *incertae sedis*, two species, type: *H. chrysocephalus* V.N. Boriss, hyphomycetous, saprobes, terrestrial, Ukraine, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Madrid et al. (2016; new species), cultures and sequences are unavailable.

Harpidium Körb. 1855, *Harpidiaceae*, *Lichinales*, *Lichenomycetes*, three species, type: *H. rutilans* Körb., with ascocata and conidia, lichenized (with green algae), terrestrial (saxicolous), Eurasian mountains, North America, South Africa, see Lumbsch and Huhndorf (2010; outline), Lumbsch et al. (2011; new species), Kirk et al. (2013; genus accepted), Lücking et al. (2017b; classification), sequences are available.

Harpographium Sacc. 1880, *Ascomycota* genera *incertae sedis*, five species, type: *H. fasciculatum* Sacc., hyphomycetous, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Harposporium Lohde 1874 (= *Podocrella* Seaver 1928; = *Atricordyceps* Samuels 1983; = *Polyrhina* Sorokin 1876 *fide* Quandt et al. 2014; Maharachchikumbura et al. 2015b), *Ophiocordycipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *H. anguillulae* Lohde, hyphomycetous, sexual morph unknown, parasitic on nematodes, terrestrial, temperate, see Kuo et al. (2008; taxonomy, DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015a, b; outline, phylogeny), Spatafora et al. (2015; new species), cultures available for the type: ARSEF 5593.

Harpostroma Höhn. 1928, *Ascomycota* genera *incertae sedis*, one species, type: *H. notha* (Sacc.) Höhn., coelomycetous, sexual morph unknown, on cheese and fungi, terrestrial, Europe, North America, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Valsaceae*), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Harzia Costantin 1888 (= *Chlorospora* Speg. 1891; = *Eidamia* Lindau 1904 [1907] *fide* Species Fungorum 2017), *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *H. acremonioides* (Harz) Costantin, hyphomycetous, sexual morph unknown, saprobes, foliicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2013b; new species, DNA), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015a, b; outline, phylogeny), ex-type cultures available for *H. cameroonensis* Crous & Jol. Roux (CPC 22065, 22066 = CBS 136420).

Hassea Zahlbr. 1902, *Dothideomycetes* genera *incertae sedis*, one species, type: *H. bacillosa* (Nyl.) Zahlbr., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Havispora K. L. Pang & Vrijmoed 2008, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *H. longyearbyensis* K.L. Pang & Vrijmoed, asexual morph unknown, saprobes, marine, Arctic, see Pang et al. (2008a, 2011a; taxonomy), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: BBH 18329, cultures and sequences are available.

Hawksworthiana U. Braun 1988, *Ascomycota* genera *incertae sedis*, one species, type: *H. peltigericola* (D. Hawksw.) U. Braun, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, UK, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hazslinszkyia Körb. 1861, *Asterinales* genera *incertae sedis*, *Dothideomycetes*, four species, type: *H. gibberulosa* (Ach.) Körb., saprobes, Europe, Asia, see Ertz and Diederich (2015; taxonomy, phylogeny), needs further studies to characterize the genus and establish its taxonomic status, sequences are available.

Hegermila Raity. 1995, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *H. andina* (Pat.) Raity., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Heimiadora Nicot 1960, *Ascomycota* genera *incertae sedis*, one species, type: *H. verticillata* Nicot, hyphomycetous, saprobes, terrestrial, Thailand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Heiomasia Nelsen, Lücking & Rivas Plata 2010, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *H. sipmanii* (Aptroot, Lücking & Rivas Plata) Nelsen, Lücking & Rivas Plata, lichenized, see Nelsen et al. (2010; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Heleiosa Kohlm., Volkm.-Kohlm. & O.E. Erikss. 1996, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *H. barbatula* Kohlm., Volkm.-Kohlm. & O.E. Erikss., asexual morph unknown, tropical, marine on maritime sedge, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Jones et al. (2015; outline), cultures and sequences are unavailable.

Helensiella (R.F. Castañeda, Heredia & R.M. Arias) Minter, R.F. Castañeda & Heredia 2015 (= *Digitella* R.F. Castañeda, Heredia & R.M. Arias 2013 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *H. rigidophora* (R.F. Castañeda, Heredia &

R.M. Arias) Minter, R.F. Castañeda & Heredia, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Mexico, see Index Fungorum (2017), holotype: XAL CB1692, cultures and sequences are unavailable, compare with *Ceratosporella* and *Dendryphion*.

Heleococcum C.A. Jørg. 1922, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *H. aurantiacum* C.A. Jørg., isolated from soil, marine or freshwater, Asia and Europe see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type of the type: CBS 201.35, slides of holotype of the type: C, NY (few remaining ascomata were broken).

Helhonia B. Sutton 1980 (= *Ceuthosporella* Höhn. 1929; = *Septochora* Höhn. 1917 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *H. rhamnigena* (Fautrey) B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequence are unavailable.

Heliastrum Petr. 1931, *Sordariomycetes* genera *incertae sedis*, one species, type: *H. concinnum* Petr., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Helicangiospora Boonmee, Bhat & K.D. Hyde 2014, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, type: *H. lignicola* Boonmee, Bhat & K.D. Hyde, asexual morph hyphomycetous, saprobes, terrestrial, temperate, see Boonmee et al. (2014b; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), holotype and ex-type cultures of the type: MFLU11-0137, MFLUCC 11-0378 = BCC 52029 = ICMP 20069.

Helicascus Kohlm. 1969, *Morosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *H. kanaloanus* Kohlm., asexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Amnuaykanjanasin et al. (2009; DNA), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Zhang et al. (2013a, 2014c; new species, phylogeny), Wijayawardene et al. (2014c; outline), Luo et al. (2016c; new species), cultures available for the type: F1R2B2 (*fide* Amnuaykanjanasin et al. 2009).

Helicobolomyces Matzer 1995, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *H. lichenicola* Matzer, with conidiomata, lichenized (with green algae) or lichenicolous, terrestrial, tropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helicocentralis Sri-indr., Chuaseehar., Boonyuen, K. Yamag., Suetrong & C.K.M. Tsui 2015, *Leotiomycetes* genera *incertae sedis*, one species, type: *H. hyalina* Sri-indr., Chuaseehar., Boonyuen, K. Yamag., Suetrong & C.K.M. Tsui, sexual morph unknown, saprobes, freshwater, Thailand, see Sri-indrasutdh et al. (2015; morphology, phylogeny), cultures and sequences are available, holotype of type: BBH27889.

Helicodendron Peyronel 1918 (= *Belospora* Clem. 1909; *Ciboriella* Seaver 1951; *Hymenoscypha* (Fr.) W. Phillips 1887; *Lambertellinia* Korf & Lizoň 1994; = *Peziza* trib. *Hymenoscypha* Fr. 1822; *Poculopsis* Kirschst. 1935; *Septatium* Velen. 1934 *fide* Jaklitsch et al. 2016; Species Fungorum 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *H. paradoxum* Peyr., hyphomycetous, sexual morph unknown, saprobes, aero-aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Chuaseeharonnachai et al. (2013; diversity, Thailand), Kirk et al. (2013; genus accepted), Panzer et al. (2015; DNA, phylogeny), Sri-indrasutdh et al. (2015; DNA, phylogeny), cultures available for the type: KF767.

Helicodochium S. Monteiro, R.F. Castañeda, A.C. Cruz & Gusmão 2014, *Ascomycota* genera *incertae sedis*, one species, type: *H. amazonicum* J.S. Monteiro, R.F. Castañeda, A.C. Cruz & Gusmão, hyphomycetous, sexual morph unknown, lignicolous, terrestrial, Brazil, see Monteiro et al. (2014b; taxonomy), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Helicofilia Matsush. 1983, *Ascomycota* genera *incertae sedis*, one species, type: *H. madrasensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, UK, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: MFC-11630, cultures and sequences are unavailable.

Helicogermislita Lodha & D. Hawksw. 1983, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *H. celastri* (S.-V. Kale & S.V.S. Kale) Lodha & D. Hawksw., asexual morphs unknown, on twigs, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2013; genus accepted), Jaklitsch et al. (2016a; position), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: HCIO 29374.

Helicogonium W.L. White 1942 (= *Myriogonium* Cain 1948 *fide* Jaklitsch et al. 2016a), *Helicogoniaceae*, *Phacidiales*, *Leotiomycetes*, 19 species, type: *H. jacksonii* W.L. White, asexual reproduction is present or absent, and when present, produce phialides directly on the hyphae, or form small phialide-bearing penicilli, mycoparasitic,

Europe, North America, see Baral and Kutorga (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Helicogoosia Hol.-Jech. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *H. paradoxa* Hol.-Jech., hyphomycetous, saprobes, terrestrial, Czechoslovakia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helicoma Corda 1837 (= *Helicocoryne* Corda 1854; = *Helicopsis* P. Karst. 1889; = *Lituaria* Riess 1853 *fide* Species Fungorum 2017), *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, type: *H. muelleri* Corda, hyphomycetous, saprobes, terrestrial, worldwide, see Boonmee et al. (2011, 2014b; taxonomy, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), available cultures for the type: CBS 964.69.

Helicomycetes Link 1809, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, c. 15, type: *H. roseus* Link, hyphomycetous, saprobes, aero-aquatic, worldwide, see Boonmee et al. (2011, 2014b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), Wijayawardene et al. (2014c; outline), Luo et al. (2017, sexual morph), available cultures for the type: AFTOL-ID 1613, CBS 283.51.

Helicoön Morgan 1892, *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, c. 20 species, type: *H. sessile* Morgan, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Helicorhoidion S. Hughes 1958, *Ascomycota* genera *incertae sedis*, six species, type: *H. botryoideum* (Cooke) S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, Asia, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helicosingula P.S. van Wyk, Marasas, Baard & Knox-Dav. 1985, *Ascomycota* genera *incertae sedis*, one species, type: *H. leucadendri* P.S. van Wyk, Marasas, Baard & Knox-Dav., hyphomycetous, pathogen, terrestrial, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helicosporium Nees 1816 (= *Helicotrichum* Nees & T. Nees 1818 *fide* Species Fungorum 2017), *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, c. 15, type: *H. vegetum* Nees, hyphomycetous, saprobes, terrestrial, worldwide, see

Cruz et al. (2009; new species, Brazil), Dharkar et al. (2010; new species, India), Boonmee et al. (2011, 2014b; taxonomy, phylogeny), Seifert et al. (2011; morphology), Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Brahamanage et al. (2017, new species, phylogeny), available cultures for the type: BCC 3332.

Helicothyrium I. Hino & Katum. 1961, *Ascomycota* genera *incertae sedis*, one species, type: *H. ryukyuense* I. Hino & Katum., coelomycetous, pathogen, Japan, see Wijayawardene et al. (2012, 2017, outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helicoubisia Lunghini & Rambelli 1979, *Ascomycota* genera *incertae sedis*, one species, type: *H. coronata* Lunghini & Rambelli, hyphomycetous, saprobe, terrestrial, Ivory Coast, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helioccephala V. Rao, K.A. Reddy & de Hoog 1984 (= *Holubovaniella* R.F. Castañeda 1985 *fide* Heredia et al. 2011), *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, seven species, type: *H. proliferans* V. Rao, K.A. Reddy & de Hoog, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Heredia et al. (2011; genus revised), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Mel'nik et al. (2013b; new species), cultures and sequences are available, compare with *Holubovaniella*, *Gonatobotrys* and *Uncispora*.

Heliscella Marvanová 1980, *Ascomycota* genera *incertae sedis*, two species, type: *H. stellata* (Ingold & V.J. Cox) Marvanová, hyphomycetous, saprobes, terrestrial, former Czechoslovakia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Heliscina Marvanová 1980, *Ascomycota* genera, *incertae sedis*, two species, type: *H. campanulata* Marvanová, hyphomycetous, saprobes, aquatic, former Czechoslovakia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helminthascus Tranzschel 1898 (= *Helminthascus* Tranzschel 1899 *fide* Species Fungorum 2017), *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *H. arachnophorus* Tranzschel, asexual morphs unknown, on insects, terrestrial, former USSR, see

Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Helminthocarpon Fée 1837, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *H. leprevostii* Fée, lichenized, terrestrial (corticulous, saxicolous), tropical, see Kirk et al. (2013; genus accepted), Bungartz et al. (2013a; outline), Lücking et al. (2017a; classification), sequences are available.

Helminthopeltis Sousa da Câmara 1950, *Dothideomycetes*, genera *incertae sedis*, one species, type: *H. almeidaeana* Sousa da Câmara, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, *Microthyriaceae*), Hyde et al. (2013) did not accept as a genus in *Microthyriaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Helminthosphaeria Fuckel 1870 (= *Litschaueria* Petr. 1923 *fide* Species Fungorum 2017), *Helminthosphaeriaceae*, *Chaetosphaerales*, *Sordariomycetes*, c. 20 species, type: *H. clavariarum* (Desm.) Fuckel, asexual morph diplococcum-like, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miller et al. (2014; new species, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Bell and Mahoney (2016; new species), culture available for the type: SMH4609.

Helminthosporiella Hern.-Restr., G.A. Sarria & Crous 2016, *Massarinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *H. stilbacea* (Moreau) Hern.-Restr., G.A. Sarria & Crous, hyphomycetous, saprobes, terrestrial, Colombia, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Helminthosporiomyces G.F. Sepúlveda, Pereira-Carv. & Dianese 2009, *Ascomycota* genera *incertae sedis*, one species, type: *H. cerradensis* G.F. Sepúlveda, Pereira-Carv. & Dianese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Pereira-Carvalho et al. (2009b; taxonomy), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Helminthosporium Link 1809 (= *Macroön* Corda 1833; = *Mydonotrichum* Corda 1831; = *Oxysporium* Lév. 1863 *fide* Species Fungorum 2017), *Dothideomycetes*, genera *incertae sedis*, c. 40 species, type: *H. velutinum* Link, hyphomycetous, sexual morph unknown, saprobes, lignicolous, terrestrial, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012;

outline, listed under *Massarinaceae*), Hyde et al. (2013) did not accept as a genus in *Massarinaceae*, Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes*, genera *incertae sedis*), Zhu et al. (2016; DNA, new species), cultures and reference specimen available for the type: MFLUCC 15-0243, HKAS 84015 (*fide* Zhu et al. 2016).

Helocarpon Fr. 1860 (= *Hastifera* D. Hawksw. & Poelt 1986; *Helocarpon* Th. Fr. 1860; = *Stereocauliscum* Nyl. 1865 *fide* Species Fungorum 2017), *Helocarpaceae*, *Lecanoromycetes* families *incertae sedis*, three species; type: *H. crassipes* Th. Fr., lichenized, see Lumbsch and Huhndorf (2010; outline), Miadlikowska et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Helochora Sherwood 1979, *Ascomycota* genera *incertae sedis*, one species, type: *H. hypertropha* Sherwood, asexual morph unknown, saprobes, terrestrial, India, UK, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Mesnieraceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013) Wijayawardene et al. (2014c) did not regard as a *Dothideomycetes*, cultures and sequences are unavailable.

Helodiomyces F. Picard 1913, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *H. elegans* F. Picard, asexual morph unknown, biotrophic, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Helotiella Sacc. 1884, *Leotiomycetes* genera *incertae sedis*, ?one species, type: *H. citri* (Penz.) Sacc., asexual morph unknown, saprobes, Europe, see Kirk et al. (2008; listed as nom. dub.), Lumbsch and Huhndorf (2010; outline, treated as *Helotiales*, genera *incertae sedis*), Jaklitsch et al. (2016a; classification).

Helvella L. 1753 (see Species Fungorum 2017 for synonyms), *Helvellaceae*, *Pezizales*, *Pezizomycetes*, c. 60 species, type: *H. mitra* L., asexual morph unknown, saprobes, ?from soil, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hwang et al. (2015; DNA, phylogeny), Zhao et al. (2016b; 2017, DNA, phylogeny, new species), Tibpromma et al. (2017b; new species), cultures and sequences are available but lacking for the type.

Hemibeltrania Piroz. 1963, *Ascomycota* genera *incertae sedis*, eleven species, type: *H. cinnamomi* (Deighton) Piroz., hyphomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are

available, needs generic revision as have different opinions.

Hemicorynespora M.B. Ellis 1972, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, twelve species, type: *H. deightonii* M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Whitton et al. (2012a; new species), Ma et al. (2012b; new species), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available but lacking for the type species.

Hemicorynesporella Subram. 1992, *Ascomycota* genera *incertae sedis*, one species, type: *H. guadalcanalensis* (Matsush.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Guadalcanal, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hemidothis Syd. & P. Syd. 1916, *Ascomycota* genera *incertae sedis*, one species, type: *H. miconiae* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hemiglossum Pat. 1890, *Helotiales* genera *incertae sedis*, two species, type: *H. yunnanense* Pat., asexual morph unknown, saprobes, terrestrial, China, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hemigrapha (Müll. Arg.) R. Sant. ex D. Hawksw. 1975 (= *Melanographa* sect. *Hemigrapha* Müll. Arg. 1882 *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, eight species, type: *H. asteriscus* (Müll. Arg.) R. Sant. ex D. Hawksw., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Wijayawardene et al. (2014c; outline), placement in *Parmulariaceae* supported by morphology, cultures and sequences are unavailable.

Hemileucoglossum S. Arauzo 2014, *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, four species, type: *H. littorale* (Rostr.) S. Arauzo, asexual morph unknown, saprobes, terrestrial, China, Japan, see Arauzo and Iglesias (2014; taxonomy, phylogeny), cultures available for the type: C: 35673.

Hemimyriangium J. Reid & Piroz. 1966, *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, one species, type: *H. betulae* J. Reid & Piroz., asexual morph unknown, saprobes, terrestrial, Sweden, see Lumbsch and

Huhndorf (2010; outline), Dissanayake et al. (2014; morphology), the genus was not listed in Kirk et al. (2013).

Hemisphaericaspora Hui, Ren, Chen, Li, Zhang & Niu 2014, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *H. nanyangensis* F.L. Hui, Y.C. Ren, L. Chen, Z.H. Li, L. Zhang, Q.H. Niu, asexual reproduction is by multilateral budding, saprophytic, frass of beetle larvae, South Africa, China, see Hui et al. (2014; taxonomy), sequences are available.

Hemisphaeropsis Petr. 1947, *Ascomycota* genera *incertae sedis*, one species, type: *H. magnoliae* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hemisynnema Subram. 1995 [1992], *Ascomycota* genera *incertae sedis*, one species, type: *H. malaysianum* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not genus accepted), cultures and sequences are unavailable.

Hemithecium Trevis. 1853, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, c. 50 species, type: *H. balbisii* (Fée) Trevis., asexual morph unknown, lichenized, terrestrial, tropical, see Archer (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rivas Plata et al. (2012a; taxonomy), Lücking et al. (2014, 2016; classification), sequences are available.

Hendersoniella (Sacc.) Tassi 1900 (= *Hendersonia* subgen. *Hendersoniella* Sacc. 1884; = *Hendersoniella* (Sacc.) Sacc. 1902 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *H. epixyla* Tassi, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hendersonina E.J. Butler 1913, *Ascomycota* genera *incertae sedis*, one species, type: *H. sacchari* E.J. Butler, coelomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hendersoniopsis Höhn. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *H. thelebola* (Sacc.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hendersonula Speg. 1880, *Ascomycota* genera *incertae sedis*, c. 20, species, type: *H. australis* Speg., coelomycetous, sexual morph unknown, saprobes or plant pathogenic, worldwide, see Wijayawardene et al.

(2012, 2016b, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hendersonulina Tassi 1902, *Ascomycota* genera *incertae sedis*, one species, type: need typification, coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Henfella Halici, D. Hawksw., Z. Kocak & M. Kocak. 2016, *Ascomycota* genera *incertae sedis*, one species, type: *H. muriformis* Halici, D. Hawksw., Z. Kocak & M. Kocak, sexual morph unknown, lichenicolous, cultures and sequences are unavailable, holotype of the type: ERC-MGH 1.231.

Henicospora P.M. Kirk & B. Sutton 1980, *Ascomycota* genera *incertae sedis*, six species, type: *H. minor* P.M. Kirk & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Monteiro et al. (2014a; Brazil), cultures and sequences are unavailable.

Henningsiella Rehm 1895, *Schizothyriaceae*, *Dothideomycetes*, families *incertae sedis*, two species, type: *H. quitensis* Rehm, asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013, accepted as *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Henrica B. de Lesd. 1921, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, four species, type: *H. ramulosa* B. de Lesd., lichenized, Europe, North America, see Savić and Tibell (2008b; taxonomy, revision), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Hepnia Nägeli 1854 (see Species Fungorum 2017 for synonyms), *Lichinaceae*, *Lichinales*, *Lichinomycetes*, seven species, type: *H. adglutinata* A. Massal., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Heppsora D.D. Awasthi & K. Singh 1977, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *H. indica* D.D. Awasthi & Kr.P. Singh, lichenized, India, Middle East, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Heptameria Rehm & Thüm. 1879 (= *Verlotia* Fabre 1879 [1878] *fide* Species Fungorum 2017), *Dothideomycetes*, genera *incertae sedis*, two species, type: *H. elegans* Rehm & Thüm., asexual morph

unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Heracleicola Tibpromma, Camporesi & K.D. Hyde 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *H. premilcurensis* Tibpromma, Camporesi & K.D. Hyde, saprobes, Italy, see Ariyawansa et al. (2015a; morphology, phylogeny), cultures available for type: MFLUCC 14-0518.

Herbampulla Scheuer & Nograsek 1993, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *H. crassirostris* Scheuer & Nograsek, asexual morph unknown, saprobes, terrestrial, Austria, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Hercospora Fr. 1825, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *H. tiliae* (Pers.) Tul. & C. Tul., asexual morph unknown, saprobes, terrestrial, Austria, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), holotype and ex-type of the type: BPI 748440, AR3526.

Hermatomyces Speg. 1911, *Hermatomycetaceae*, *Pleosporales*, *Dothideomycetes*, 15 species, type: *H. tucumanensis* Speg., saprobes, terrestrial, Africa, Asia, Caribbean, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Doilom et al. (2017; new species, phylogeny), Hyde et al. (2016; new species), Tibpromma et al. (2016a, 2017b; new species), Hashimoto et al. (2017; new family and new species), cultures and sequences are available.

Herpomyces Thaxt. 1902, *Herpomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 25 species, type: *H. chaetophilus* Thaxt., asexual morph unknown, biotrophic, dioecious, hosts are Blattodea, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2015b; DNA), sequences are available for type: (SSU, ITS, LSU), GenBank accession number for ITS of *Herpomyces chaetophilus*: KT800039.

Herposira Syd. 1938, *Ascomycota* genera *incertae sedis*, one species, type: *H. velutina* Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Herpothallon Tobler 1937, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, 46 species, type: *H. sanguineum* (Vain.) Tobler, lichenized, see Lumbsch and Huhndorf (2010;

outline), Lücking et al. (2017a; classification), sequences are available.

Herpotrichia Fuckel 1868 (= *Enchnosphaeria* Fuckel 1870 [1869–70]; = *Khekia* Petr. 1921; = *Sordariella* J.N. Kapoor, S.P. Lal & Bahl 1975 [1974] *fide* Species Fungorum 2017), *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, 101 species, type: *H. herpotrichoides* (Fuckel) P.F. Cannon, asexual morph coelomycetous, parasitic or saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2012, 2014c, 2017; outline; phylogeny, coelomycetous taxa), Kirk et al. (2013; genus accepted), Tian et al. (2015; morphology, nomenclature, phylogeny), cultures and sequences are available but lacking for the type.

Herreromyces R.F. Castañeda & W.B. Kendr. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *H. catenosporus* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype: INIFAT C91/47, cultures and sequences are unavailable, compare with *Exochalara*, *Chloridium* and *Monilochaetes*.

Herteliana P. James 1980, *Squamariaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *H. taylorii* (Salwey) P. James, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017b; classification), sequences are available.

Hertelidea Printzen & Kantvilas 2004, *Stereocaulaceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *H. botryosa* (Fr.) Printzen & Kantvilas, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Hertella Henssen 1985, *Placynthiaceae*, *Peltigerales*, *Lecanoromycetes*, three species, type: *H. subantarctica* Henssen, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Hesperomyces Thaxt. 1891, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eleven species, type: *H. virescens* Thaxt., no asexual state, biotrophic, hosts are *Biphyllidae*, *Coccinellidae*, *Mycetophagidae* (*Coleoptera*), and *Anthocoridae* (*Hemiptera*: *Heteroptera*), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2015b, 2016; DNA), sequences are available for the type: (SSU, ITS, LSU), GenBank accession numbers for ITS of *Hesperomyces virescens*: JN127409, KF192910–KF192920, KT800043–KT800048, KU574864

Heterocarpon Müll. Arg. 1885, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type:

H. ochroleucum (Tuck.) Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Heterocephalum Thaxt. 1903, *Ascomycota* genera *incertae sedis*, two species, type: *H. aurantiacum* Thaxt., hyphomycetous, sexual morph unknown, coprophilous, saprobes, terrestrial, worldwide, see Puangsombat et al. (2010; Thailand), Seifert et al. (2011; morphology), Silva et al. (2011; Amazon forest), Cort (2012; distribution), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and Sequences are unavailable, neotype of type: CBS H-15402.

Heteronrium Petr. 1949, *Dothideomycetes* genera *incertae sedis*, 20 species, type: *H. citharexylī* Petr., hyphomycetous, sexual morph unknown, saprobes, endophytes, terrestrial, worldwide, see Ruiz et al. (2008; new species), Zhang et al. (2010c; new species), Seifert et al. (2011; morphology), Ma et al. (2012f; new species), Ren et al. (2012b; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Xia et al. (2013b; new species), Lahlali et al. (2014; resistance to clubroot), sequences are available for most of the species.

Heterocyphelium Vain. 1927, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *H. leucampyx* (Tuck.) Vain., lichenized, see Lumbsch and Huhndorf (2010; outline), Tibell and Frisch (2010; Africa), Kirk et al. (2013; genus accepted), Pandit (2015; India), Lücking et al. (2017a; classification), sequences are available.

Heterodermia Trevis. 1868 (= *Chaudhuria* Zahlbr. 1932; = *Pseudophyscia* Müll. Arg. 1894; = *Pseudophyscia* Müll. Arg. 1893 *fide* Species Fungorum 2017), *Physciaceae*, *Caliciales*, *Lecanoromycetes*, 90 species, type: *H. speciosa* (Wulff) Trevis., lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Heteropichloë E. Tanaka, C. Tanaka, Gafur & Tsuda 2002, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, type: *H. bambusae* (Pat.) E. Tanaka, C. Tanaka, Gafur & Tsuda, asexual morphs unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, nomenclature), cultures and sequences are available

Heteromyces Müll. Arg. 1889, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *H. rubescens* Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences unavailable.

Heterophoma Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *H. sylvatica* (Sacc.) Q. Chen & L. Cai, coelomycetous, saprobes, ?pathogens, terrestrial, cosmopolitan, see Chen et al. (2015b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), neotype and ex-neotype of the type: HMAS 246700, CBS 874.97.

Heteroplacidium Breuss 1996, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, twelve species, type: *H. imbricatum* (Nyl.) Breuss, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Heteroseptata E.F. Morris 1972, *Ascomycota* genera *incertae sedis*, one species, type: *H. bacillospora* E.F. Morris, sexual morph unknown, saprobes, terrestrial, Costa Rica, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Heterosphaeria Grev. 1824 (= *Heteropatella* Fuckel *fide* Johnston et al. 2014b), *Heterosphaeriaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *H. patella* (Tode) Grev., asexual morph hyphomycetous, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature; new combination), cultures and sequences are unavailable.

Heterosphaeriopsis Hafellner 1979, *Dothideomycetes* genera *incertae sedis*, one species, type: *H. fulvodisca* (Pat.) Hafellner, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of type: (FH, Sweden), cultures and sequences are unavailable.

Heterosporicola Crous, *nom. nov.* MycoBank MB821707

≡ *Phoma* sect. *Heterospora* Boerema, Gruyter & Noordel., Persoonia 16(3): 336. 1997.

≡ *Heterospora* (Boerema, Gruyter & Noordel.) Gruyter, Aveskamp & Verleyen, Stud. Mycol. 75: 18. 2012 [2013]. nom. illeg., [ICN (Melbourne) Art. 53.1], non *Heterospora* P. Kuckuck 1895 (*Tilopteridaceae*)

Leptosphaeriaceae, *Pleosporales*, *Dothideomycetes*, two species, coelomycetous, sexual morph unknown, saprobic or pathogenic on stems and leaves of *Chenopodium* spp., Argentina, Belgium, Peru, The Netherlands, see de Gruyter et al. (2012; morphology, phylogeny).

Holotype: Belgium, Beverloo, from leaves of *Chenopodium suecicum* (album) and *Chenopodium urbicum* (*Chenopodiaceae*), no date, G.D. Westendorp, Herb. Crypt. (Ed. Beyaert-Feys), No. 959. BR, holotype of *Phyllosticta chenopodii* Westend. ex herb. G.D.

Westendorp. Epitype (designated by de Gruyter et al. 2012): Netherlands, Baarn, from leaf spots in *Chenopodium album*, 3 Jul. 1968, H.A. van der Aa, CBS H-16386; culture ex-epitype, CBS 448.68.

Heterosporicola chenopodii (Westend.) Crous, **comb. nov.** MycoBank: MB821708

≡ *Phyllosticta chenopodii* Westend., Bull. Acad. R. Sci. Belg., Cl. Sci., sér. 2 2(7): 567. 1857.

≡ *Heterospora chenopodii* (Westend.) Gruyter, Aveskamp & Verkley, Stud. Mycol. 75: 18. 2012 [2013].

≡ *Septoria westendorpii* G. Winter, Hedwigia 26: 26. 1887. nom. inval., [ICN (Melbourne) Art. 36.1(b)],

≡ *Phoma variospora* Aa & Kesteren, Persoonia 10: 268. 1979. nom. illeg., [ICN (Melbourne) Art. 53.1],

≡ *Phoma heteromorphospora* Aa & Kesteren, Persoonia 10: 542. 1980.

Heterosporicola dimorphospora (Speg.) Crous, **comb. nov.** MycoBank: MB821709

≡ *Phyllosticta dimorphospora* Speg., Anal. Mus. nac. B. Aires, Ser. 3 13: 334. 1911.

≡ *Phoma dimorphospora* (Speg.) Aa & Kesteren, Persoonia 10(2): 269. 1979.

≡ *Heterospora dimorphospora* (Speg.) Gruyter, Aveskamp & Verkley, Stud. Mycol. 75: 18. 2012 [2013].

= *Stagonospora chenopodii* Peck, Rep. (Annual) New York State Mus. Nat. Hist. 40: 60. 1887.

Notes: The genus *Heterospora* (Boerema, Gruyter & Noordel.) Gruyter, Verkley & Crous 2012 is illegitimate, being a later homonym of the algal genus *Heterospora* P. Kuckuck 1895. In order to replace this name, the new name *Heterosporicola* nom. nov. is provided for this genus. Species of the genus are known plant pathogens of *Chenopodiaceae* in South America and Europe (de Gruyter et al. 2012).

Heuflera Bail 1860, *Ascomycota* genera *incertae sedis*, type: *H. betulae* Bail, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Heufleridium Müll. Arg. 1883, *Pyrenulales* genera *incertae sedis*, *Eurotiomycetes*, two species, type: *H. pentagastricum* (Müll. Arg.) Müll. Arg., asexual morphs unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hexacladium D.L. Olivier 1983, *Ascomycota* genera *incertae sedis*, one species, type: *H. corynephorum* D.L. Olivier, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hexagonella F. Stevens & Guba ex F. Stevens 1925, *Schizothyriaceae*, *Dothideomycetes*, families *incertae sedis*, two species, type: *H. peleae* F. Stevens & Guba ex F. Stevens, asexual morph unknown, saprobes, terrestrial, Hawaii, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013, accepted as *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Heydenia Fresen. 1852 (= *Riccoa* Cavara 1903; = *Rupinia* Speg. & Roum. 1879 *fide* Species Fungorum 2017), *Pezizales* genera *incertae sedis*, *Pezizomycetes*, four species, type: *H. alpina* Fresen., hyphomycetous, sexual morph unknown, saprobes, terrestrial, north temperate, see Seifert et al. (2011; morphology), Leuchtmann and Clemenccon (2012; DNA, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), cultures available for the type: 0732.

Heyderia Link 1833, *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *H. abietis* (Fr.) Link, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Hemiphacidiaceae*), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny, accepted as a genus in *Cenangiaceae*), cultures available for the type: KL20.

Hidakaea I. Hino & Katum. 1955, *Dothideomycetes* genera *incertae sedis*, two species, type: *H. tumidula* I. Hino & Katum, asexual morph unknown, saprobes, terrestrial, Hawaii, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Wu et al. (2010; placed in *Hypocreales* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes* genera *incertae sedis*), cultures and sequences are unavailable, generic revision is essential as it has different opinions.

Hiemsia Svrček 1969, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, two species, type: *H. pseudoampezzana* (Svrček) Svrček, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hilberina Huhndorf & A.N. Mill. 2004, *Helminthosphaeriaceae*, *Chaetosphaerales*, *Sordariomycetes*, c. 20 species, type: *H. caudata* (Fuckel) Huhndorf & A.N. Mill., asexual morph diplococcium-like, saprobes, terrestrial, worldwide, see Declercq (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miller et al. (2014; new species, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Bell and Mahoney (2016; new species), culture available for the type: SMH1542 (*fide* Miller et al. 2014).

Himantermia I.M. Lamb 1964 (= *Nimisia* Kärnefelt & A. Thell 1993 *fide Species Fungorum* 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *H. lugubris* (Hue) I.M. Lamb, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Hinoa Hara & I. Hino 1961 (= *Haraella* Hara & I. Hino 1955 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, two species, type: *H. sasae* (Hara & I. Hino) Hara & I. Hino, ?coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hippocrepedea Sérus. 1997, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *H. nigra* Sérus., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Hirudinaria Ces. 1856 (= *Hippocrepidium* Sacc. 1874 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, two species, type: *H. mespili* Ces., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hispidicarpomyces Nakagiri 1993, *Hispidicarpomycetaceae*, *Spathulosporales*, *Sordariomycetes*, one species (probably more than this), type: *H. galaxauricola* Nakagiri, spermatiophores or asexual morph hyphomycetous, biotrophic on marine red alga, Japan, see Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharanachikumbura et al. (2016b; outline, the family accepted as in *Spathulosporales*), cultures and sequences are unavailable, holotype of the type: IFO H-12135 and paratype IFO H-12136, needs generic revision since have different opinions.

Hispidoconidioma Tsuneda & M.L. Davey 2010, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *H. alpinum* Tsuneda & M.L. Davey, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Tsuneda et al. (2010; taxonomy), Wijayawardene et al. (2012, 2014c; outline), holotype and ex-type of the type: UAMH 11010, L2-2.

Histeridomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, six species, type: *H. ramosus* Thaxt., asexual morph unknown, biotrophic, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Histoplasma Darling 1906, *Ajellomycetaceae*, *Onygenales*, *Eurotiomycetes*, seven species, type: *H. capsulatum* Darling, hyphomycetous, sexual morph unknown, on humans and animals, bat guano, terrestrial, Africa, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Hobsoniopsis D. Hawksw. 2001, *Ascomycota* genera *incertae sedis*, one species, type: *H. santessonii* (Lowen & D. Hawksw., sexual morphs unknown, lichenicolous, Europe, terrestrial, see Seifert et al. (2011; morphology), cultures and sequences are available, holotype of type at UPS.

Hobus Jaklitsch & Voglmayr 2016, *Thyridariaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *H. wogradensis* Jaklitsch & Voglmayr, asexual morph unknown, saprobes, Central Europe, see Jaklitsch and Voglmayr (2016; taxonomy, phylogeny), holotype of the type: WU 36874.

Hoehneliella Bres. & Sacc. 1902 (= *Klebahnopycnis* Kirschst. 1939 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, two species, type: *H. perplexa* Bres. & Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Austria, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hoffmannoscypha Stielow, Göker & Klenk 2012 [2013], *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *H. pellita* (Sacc.) Stielow, Göker & Klenk, asexual morph unknown, saprobes, terrestrial, Europe, see Stielow et al. (2013; taxonomy, phylogeny), epitype of the type: M-0156529 (holotype of *Peziza pellita*), cultures and sequences are available.

Holmiella Petrini, Samuels & E. Müll. 1979 (= *Caldesia* Rehm 1890 [1896]; = *Eutryblidiella* Höhn 1918 *fide Species Fungorum* 2017), *Patellariaceae*, *Patellariales*, *Dothideomycetes*, two species, type: *H. sabina* (De Not.) Petrini, Samuels & E. Müll., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), cultures and sequences are unavailable.

Holocryphia Gryzenh. & M.J. Wingf. 2006, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *H. eucalypti* (M. Venter & M.J. Wingf.) Gryzenh. & M.J. Wingf., asexual morph unknown, pathogens, terrestrial, Australia, New Zealand, South Africa, see Lumbsch and Huhndorf (2010; outline), Chen

et al. (2013b; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: PREM 56211, cultures and sequences are available.

Holubovaea Mercado 1983, Ascomycota genera incertae sedis, two species, type: *H. roystoneicola* Mercado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Holwaya Sacc. 1889 (= *Crinula* Fr. 1821; = *Ditiola* Schulzer 1860; = *Jugglerandia* Lloyd 1923 fide Johnston et al. 2014b; Species Fungorum 2017), Tympanidaceae, Phacidiales, Leotiomycetes, two species, type: *H. ophiobolus* (Ellis) Sacc., asexual morph *Crinula*, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Johnston et al. (2014b; nomenclature), living cultures and sequences are available for sexual morph.

Homaromyces R.K. Benj. 1955, Laboulbeniaceae, Laboulbeniales, Laboulbeniomycetes, one species, type: *H. epieri* R.K. Benj., asexual morph unknown, biotrophic, USA, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Homortomyces Crous & M.J. Wingf. 2012, Homortomycetaceae, Dothideomycetes families incertae sedis, type: *H. combreti* Crous & M.J. Wingf., two species, saprobes, South Africa, Italy, see Crous et al. (2012b; morphology, phylogeny), Wijayawardene et al. (2014c, 2014e, 2016b; morphology, phylogeny), Thambugala et al. (2017a; phylogeny), ex-epitype strains of type: CPC 19800; epitype CBS H-21049.

Homostegia Fuckel 1870 [1869–70] (= *Trematosphaeris* Clem. & Shear 1931 fide Species Fungorum 2017), Dothideomycetes, genera incertae sedis, five species, type: *H. adusta* Fuckel, asexual morph unknown, saprobes, terrestrial, Cuba, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Homothecium A. Massal. 1853 (= *Lecidocollema* Vain. 1890 fide Species Fungorum 2017), Pannariaceae, Peltigerales, Lecanoromycetes, five species, type: *H. opulentum* (Mont.) A. Massal., lichenized, see Kirk et al. (2013; genus accepted), Ekman et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Hongkongmyces C.C.C. Tsang, J.F.W. Chan, N.J. Trendell-Sm., A.H.Y. Ngan, I.W.H. Ling, K.Y. Yuen, S.K.P. Lau & P.C.Y. Wood 2014, Lindgomycetaceae, Pleosporales, Dothideomycetes, two species, type: *H. pedis* C.C.C. Tsang et al., asexual morph unknown, human

pathogen, Hong Kong, see Tsang et al. (2014b; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), holotype and ex-type strain of the HKU35.

Hoornsmania Crous 2007, Cladosporiaceae, Capnodiales, Dothideomycetes, one species, type: *H. pyrina* Crous, hyphomycetous, sexual morph unknown, hyperparasitic on *Neonectria ditissima*, terrestrial, France, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), holotype of the type: CBS H-19769, cultures and sequences are unavailable.

Hormiactis Preuss 1851 (= *Hormiactina* Bubák 1916 fide Species Fungorum 2017), Ascomycota genera incertae sedis, five species, type: *H. alba* Preuss, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Sri Lanka, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hormiokrypsis Bat. & Nascim. 1957, Ascomycota genera incertae sedis, one species, type: *H. libocedri* Bat. & Nascim., coelomycetous, sexual morph unknown, sooty mould on plants, terrestrial, Australasia, North America, Oceania, see Seifert et al. (2011; morphology, accepted as a genus in Metacapnodiaceae), Wijayawardene et al. (2012, 2014c; outline, accepted as a genus in Metacapnodiaceae), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Hormiscioides M. Blackw. & Kimbr. 1979, Ascomycota genera incertae sedis, one species, type: *H. filamentosus* M. Blackw. & Kimbr., hyphomycetous, sexual morph unknown, on insects, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hormisciomyces Bat. & Nascim. 1957, Euantennariaceae, Dothideomycetes families incertae sedis, eight species, type: *H. prepusus* Bat. & Nascim., sooty mould, terrestrial, Australasia, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hormocephalum Syd. 1939, Ascomycota genera incertae sedis, five species, type: *H. ecuadorense* Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Sri Lanka, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hormocladium Höhn. 1923 (see Species Fungorum 2017 for synonyms), Ascomycota genera incertae sedis,

one species, type: *H. kaki* (Hori & Yoshino) Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hormoconis Arx & G.A. de Vries 1973, *Amorphothecaceae*, *Leotiomycetes*, families *incertae sedis*, two species, type: *H. resinae* (Lindau) Arx & G.A. de Vries, hyphomycetous, sexual morph unknown, saprobes on resins, hydrocarbons, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available.

Hormographis Guarro, Punsola & Arx 1986, *Ascomycota* genera *incertae sedis*, one species, type: *H. ramirezzii* Guarro, Punsola & Arx, hyphomycetous, sexual morph unknown, from soil, terrestrial, Spain, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hormonema Lagerb. & Melin 1927, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, six species, type: *H. dematiodoides* Lagerb. & Melin, saprobes, human pathogen, terrestrial, Spain, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), Crous et al. (2015d; new species, DNA), ?polyphyletic, needs generic revision and epitypification.

Hormosphaeria Lév. 1863, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *H. tessellata* Lév., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Hortaea Nishim. & Miyaji 1984, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *H. werneckii* (Horta) Nishim. & Miyaji, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; new species, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c; outline, phylogeny), Hyde et al. (2013; accepted as a genus in *Teratosphaeriaceae*), Egidi et al. (2014; phylogeny), Quaedvlieg et al. (2014; phylogeny), cultures available for the type: CBS 117931=TRN122 (*fide* Egidi et al. 2014).

Hosseusia Gyeln. 1940, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, one species, type: *H. gertrudiana* Gyeln., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Houjia G.Y. Sun & Crous 2010, *Phaeothecoidiellaceae*, *Dothideomycetes* family, *incertae sedis*, two species,

hyphomycetous, sexual morph unknown, saprobes, ?pathogens, terrestrial, USA, China, see Yang et al. (2010; morphology, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Hongsanan et al. (2017b; phylogeny, family placement), holotype and ex-type strains of the type: CBS H-20325, CPC 16114.

Huangshania O.E. Erikss. 1992, *Triblidiaceae*, *Triblidiales*, *Leotiomycetes*, two species, type: *H. verrucosa* O.E. Erikss., asexual morph unknown, saprobes, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Huea C.W. Dodge & G.E. Baker 1938 (= *Carbonea* (Hertel) Hertel 1983 *fide* Lücking et al. (2017a; see Species Fungorum 2017 for additional synonyms), *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, 20 species, type: *H. flava* C.W. Dodge & G.E. Baker, lichenized, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Teloschistaceae*), Lücking et al. (2017a; classification), sequences are available.

Hueidea Kantvilas & P.M. McCarthy 2003, *Fuscideaceae*, *Umbilicariales*, *Lecanoromycetes*, one species, type: *H. australiensis* Kantvilas & P.M. McCarthy, lichenized, see Lücking et al. (2017a; classification), sequences are unavailable.

Hughesinia J.C. Lindq. & Gamundí 1970, *Ascomycota* genera *incertae sedis*, one species, type: *H. chusqueae* J.C. Lindq. & Gamundí, hyphomycetous, sexual morph unknown, from soil, terrestrial, Chile, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Humaria Fuckel 1870 (= *Lachnea* Boud. 1885; = *Leucopezis* Clem. 1909; = *Mycolachnea* Maire 1937 *fide* Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, c. 16 species, type: *H. hemisphaerica* (F.H. Wigg.) Fuckel, asexual morph unknown, saprobes, terrestrial, north temperate, see Palmer et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Gaitnieks et al. (2016; DNA), Klavina et al. (2016; DNA, mycorrhization), cultures and sequences are available.

Humicola Traaen 1914 (= *Melanogone* Wollenw. & H. Richt. 1934 *fide* Species Fungorum 2017), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *H. fuscoatra* Traaen, hyphomycetous, sexual morph unknown, saprobes, from soil, terrestrial, worldwide, see Jiang and Zhang (2010; new species, China), Wu and Zhang (2013; new species), Kirk et al. (2013; genus accepted), Geng and Zhang (2014; new species), Zhang and Zhang (2014; new species, China), Maharachchikumbura et al. (2015b, 2016b; outline),

cultures available for the type: NRRL 22980, DAOM 35882.

Humicolopsis Cabral & S. Marchand 1976, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *H. cephalosporioides* Cabral & Marchand, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Argentina, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Huneckia S.Y. Kondr., Elix, Kärnefelt, A. Thell & Hur 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *H. pollinii* (A. Massal.) S.Y. Kondr., Kärnefelt, A. Thell, Elix, J. Kim, A.S. Kondr. & Hur, lichenized, see Kondratyuk et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Huntiella Z.W. de Beer, T.A. Duong & M.J. Wingf. 2014, *Ceratocystidaceae*, *Microascales*, *Sordariomycetes*, 17 species, type: *H. moniliformis* (Hedg.) Z.W. de Beer, T.A. Duong & M.J. Wingf., asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see De Beer et al. (2014; phylogeny), Maharachchikumbura et al. (2015; phylogeny, outline), sequences are available.

Hyalacrotes (Korf & L.M. Kohn) Raity. 1991 (= *Urceolella* subgen. *Hyalacrotes* Korf & L.M. Kohn 1980 *fide* Species Fungorum 2017), *Calloriaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *H. hamulata* (Rehm) Raity., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalobelemnospora Matsush. 1993, *Ascomycota* genera *incertae sedis*, one species, type: *H. amazonica* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Peru, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalocamposporium Révay & J. Gönczöl 2007, *Ascomycota* genera *incertae sedis*, four species, type: *H. longiflagellatum* Révay & J. Gönczöl, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: BP 98968, cultures and sequences are unavailable.

Halocephalotrichum Nagaraju, Kunwar, Sureshk. & Manohar. 2011, *Ascomycota* genera *incertae sedis*, one species, type: *H. indicum* Nagaraju, Kunwar, Sureshk. & Manohar., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, holotype of the type: HCIO 50144, cultures and sequences are unavailable.

Hyalocladium Moustafa 1977 [1976], *Ascomycota* genera *incertae sedis*, one species, type: *H. moubasherii* Moustafa, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Kuwait, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalocrea Syd. & P. Syd. 1917 (= *Poeltia* Petr. 1972 [1971]; = *Poeltiella* Petr. 1974 [1972] *fide* Species Fungorum 2017), *Dothideomycetes*, genera *incertae sedis*, four species, type: *H. epimyces* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hyalocylindrophora J.L. Crane & Dumont 1978, *Ascomycota* genera *incertae sedis*, three species, type: *H. rosea* (Petch) Réblová & W. Gams, hyphomycetous, sexual morph unknown, on cheese and fungi, terrestrial, Europe, North America, see Réblová et al. (2011; reinstated the genus), Seifert et al. (2011; morphology), Maharachchikumbura et al. (2015b, 2016b; outline, treated as a genus in *Australiascaceae*), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Hyalodendriella Crous 2007, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *H. betulae* Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Netherlands, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype and ex-type strains of the type: CBS-H 19895, CBS 261.82.

Hyaloderma Speg. 1884 (= *Mycophaga* F. Stevens 1924 *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, one species, type: *H. imperspicuum* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Hyalodermella Speg. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *H. gardeniae* (Keissl.) Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalodictyum Woron. 1916, *Ascomycota* genera *incertae sedis*, one species, type: *H. colchicum* Woron., coelomycetous, sexual morph unknown, saprobes, terrestrial, former USSR, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalohelicomina T. Yokoy. 1974, *Ascomycota* genera *incertae sedis*, one species, type: *H. deutiae* (T. Yokoy.)

T. Yokoy., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalomeliolina F. Stevens 1924, *Perisporiopsidaceae*, *Dothideomycetes*, families *incertae sedis*, two species, type: *H. guianensis* F. Stevens, asexual morph unknown, saprobes, terrestrial, central and south America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Perisporiopsidaceae*), cultures and sequences are unavailable.

Hyalopeziza Fuckel 1870 (= *Hyalotricha* Dennis 1949; = *Pseudolachnea* Velen. 1934 *fide* Species Fungorum 2017), *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, c. 15 species, type: *H. patula* (Pers.) Fuckel, asexual morph phialophora-like, saprobes, ?lichenicolous, terrestrial, worldwide, see Huhtinen et al. (2008; new species, Alaska), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), type in G and S herbarium, cultures and sequences are available but lacking for the type.

Hyalopleiochaeta R.F. Castañeda, Guarro & Cano 1996, *Ascomycota* genera *incertae sedis*, one species, type: *H. delicata* R.F. Castañeda, Guarro & Cano, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype: INIFAT C94/140, cultures and sequences are unavailable, compare with *Campylosporium*.

Hyalorbilia Baral & G. Marson 2001 (= *Dwyaangam* Subram. 1978 *fide* Baral et al. 2015), *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, 13 species, type: *H. berberidis* (Velen.) Baral, asexual morph ?hyphomycetous, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available but lacking for type species.

Hyalorhinocladiella H.P. Upadhyay & W.B. Kendr. 1975, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, one species, type: *H. minuta-bicolor* (R.W. Davidson) H.P. Upadhyay & W.B. Kendr., hyphomycetous, saprobes, worldwide, see Lu et al. (2009; DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Hyalorostratum Raja & Shearer 2010, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type:

H. brunneisporum Raja & Shearer, freshwater, Alaska and New Hampshire see Raja et al. (2010a; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: A573-1 ILL 40792.

Hyaloscolecostroma Bat. & J. Oliveira 1967, *Dothideomycetes* genera *incertae sedis*, one species, type: *H. rondonense* Bat. & J. Oliveira, asexual morph unknown, lichenicolous, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; outline, genus accepted), Wijayawardene et al. (2014c; outline), culture and sequence are unavailable.

Hyaloscypha Boud. 1885 (= *Eupezizella* Höhn. 1926; = *Orbiliopsis* Höhn. 1926; = *Parorbiliopsis* Spooner & Dennis 1986 [1985]; = *Pseudaegerita* J.L. Crane & Schokn. 1981; = *Truncicola* Velen. 1934 *fide* Jaklitsch et al. 2016a; Species Fungorum 2017), *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, c. 45 species, type: *H. vitreola* (P. Karst.) Boud., asexual morph hyphomycetous in *H. albohyalina* var. *spiralis* and *H. albohyalina* var. *monodictys*, saprobes, terrestrial, worldwide, see Baral et al. (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), cultures and sequences are available for the type: M220 (*fide* Baral et al. 2009).

Hyaloseta A.W. Ramaley 2001, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *H. nolinae* A.W. Ramaley, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type culture for the type: CBS 109837.

Hyalosphaera F. Stevens 1917 (= *Dexteria* F. Stevens 1917; = *Hyalotaxis* Syd. 1925 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, four species, type: *H. miconiae* F. Stevens, asexual morph unknown, saprobes, terrestrial, North and Central America, see Alves et al. (2010; new species, Costa Rica), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequence data available for unidentified *Hyalosphaera* species.

Hyalosynnema Matsush. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *H. multiseptatum* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, Taiwan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalothelos Speg. 1908, *Dothideomycetes*, genera *incertae sedis*, one species, type: *H. dimerosperma* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Alves et al. (2010; new species, Costa Rica), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Elsinoaceae*), Li et al. (2011; excluded from *Elsinoaceae*), Jayawardena et al. (2014; excluded from

Elsinoaceae), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequence data available for unidentified *Hyalosphaera* species.

Hyalothyridium Tassi 1900 (= *Hyalothyris* Clem. 1909 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, one species, type: *H. viburnicola* Tassi, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyalotiella Papendorf 1967, *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, four species, type: *H. transvalensis* Papendorf, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), Li et al. (2015d; new species); Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; outline), sequence data and cultures available but lacking for the type.

Hyalotopsis Punith. 1970 (= *Ellurema* Nag Raj & W.B. Kendr. *fide Réblová et al. 2016c*), *Sordariomycetes* genera *incertae sedis*, one species, type: *H. subramanianii* (Agnihothr. & Luke) Punith., coelomycetous, sexual morph formerly known in *Ellurema*, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), Réblová et al. (2016c; nomenclature), cultures and sequences are unavailable.

Hydea K.L. Pang & E.B.G. Jones 2010, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, one species, type: *H. pygmaea* (Kohlm.) K.L. Pang & E.B.G. Jones, hyphomycetous, sexual morph unknown, saprobes, marine, pantropical, see Abdel-Wahab et al. (2010; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Hydnobolites Tul. & C. Tul. 1843, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *H. cerebriformis* Tul. & C. Tul., asexual morph unknown, hypogeous, terrestrial, North America, Europe, see Lumbsch and Huhndorf (2010; outline), Healy et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures available for type: PA9 (*fide* Klavina et al. 2016).

Hydnocystis Tul. & C. Tul. 1845 (= *Protogeneia* Kobayasi 1964 [1963] *fide Species Fungorum* 2017), *Pezizales* genera *incertae sedis*, *Pezizomycetes*, two species, type: *H. arenaria* Tul. & C. Tul., asexual morph unknown, hypogeous, terrestrial, Japan, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available (unpublished) but lacking for the type.

Hydnotrya Berk. & Broome 1846 (= *Geoporella* Soehner 1951; = *Gyrocratera* Henn. 1899 *fide Species Fungorum* 2017), *Discinaceae*, *Pezizales*, *Pezizomycetes*, c. 15 species, type: *H. tulasnei* (Berk.) Berk. & Broome, asexual morph unknown, hypogeous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Li et al. (2013; DNA, phylogeny, new species), Methven et al. (2013; DNA, phylogeny), cultures available for the type: BB70 (766) (*fide* Stielow et al. 2010).

Hydnotryopsis Gilkey 1916, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *H. setchellii* Gilkey, asexual morph unknown, hypogeous, terrestrial, North America, see Brock et al. (2009; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available for the type: RBG Kew K(M)99871.

Hydraeomyces Thaxt. 1896, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *H. halipli* (Thaxt.) Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hydrocina Scheuer 1991, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *H. chaetocladia* Scheuer, asexual morph unknown, saprobes, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline), Baschien et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available for the type: HME4375, CCM F-10890 (*fide* Baschien et al. 2013).

Hydromelitis A. Ferrer, A.N. Mill., C. Sarmiento & Shearer 2012, *Sordariomycetidae* genera *incertae sedis*, one species, *H. pulchella* A. Ferrer, A.N. Mill., Sarmiento & Shearer, freshwater, submerged substrates, Costa Rica, see Ferrer et al. (2012; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: AF284-2, ILL 40117.

Hydrometrospora J. Gönczöl & Révay 1985, *Ascomycota* genera *incertae sedis*, one species, type: *H. symmetrica* J. Gönczöl & Révay, hyphomycetous, sexual morph unknown, saprobes, aquatic, Hungary, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, Holotype and cultures of type species unavailable.

Hydronectria Kirschst. 1925, *Sordariomycetes* genera *incertae sedis*, one species, type: *H. kriegeriana* Kirschst., asexual morph unknown, algicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are

unavailable, cultures and sequences of type species are unavailable.

Hydrophilomyces Thaxt. 1908, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, twelve species, type: *H. rhynchophorus* (Thaxt.) Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hydropisphaera Dumort. 1822 (= *Hyphonectria* (Sacc.) Petch 1937; = *Nectria* subgen. *Hyphonectria* Sacc. 1883; = *Neohenningsia* Koord. 1907; = *Neuronectria* Munk 1957; = *Perrotiella* Naumov 1916 *fide Species Fungorum* 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, 25 species, type: *H. peziza* (Tode) Dumort., asexual morph unknown, terrestrial, aquatic, worldwide, Lechat et al. (2010; asexual morph), Luo and Zhuang (2010; new species), cultures and sequences are available.

Hydropunctaria C. Keller, Gueidan & Thüs 2009, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, eight species, type: *H. maura* (Wahlenb.) C. Keller, Gueidan & Thüs, marine, lichenized, Asia, Europe, Australia, South America, see Gueidan et al. (2009; description, phylogeny), Roux et al. (2011; new species), Lücking et al. (2017a; classification), cultures and sequences are available, but not from type material.

Hymenelia Kremp. 1852 (= *Pinacisca* A. Massal. 1854 *fide Species Fungorum* 2017), *Hymeneliaceae*, *Hymenellales*, *Lecanoromycetes*, 25 species, type: *H. prevostii* (Duby) Kremp., lichenized, worldwide, see Wirth et al. (2011; Germany), Himelbrant et al. (2013; Russia), Aptroot and Moon (2014; Korea), Kantvilas (2014b; new species), Hansen (2015; Greenland), Lücking et al. 2016 (classification), cultures and sequences are available but lacking for type species.

Hymenella Fr. 1822 (= *Hymenula* Fr. 1825 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, eleven species, type: *H. arundinis* Fr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Hyde et al. (2011; checklist), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hymenopeltis Bat. 1959, *Ascomycota* genera *incertae sedis*, three species, type: *H. erythroxyl* Bat. & A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, Holotype and ex-type of type species unavailable.

Hymenobia Nyl. 1854 (= *Hymenobiella* Triebel 1989 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, one species, type: *H. insidiosa* Nyl., asexual morph

unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, Holotype or ex-type of type species unavailable.

Hymenobolus Durieu & Mont. 1845, *Helotiaceae* genera *incertae sedis*, *Leotiomycetes*, three species, type: *H. agaves* Durieu & Mont., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hymenopsis Sacc. 1886 (= *Amerosporium* subgen. *Phaeopolynema* (Speg.) Petr. & Syd. 1937; = *Chaetodiscula* Bubák & Kabát 1910; = *Phaeodochium* M.L. Farr 1968; = *Phaeopolynema* Speg. 1912; = *Schoenbornia* Bubák 1906 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, 13 species, type: *H. trochiloides* (Sacc.) Sacc., sexual morph unknown, saprobes, fungicolous, terrestrial, worldwide, see Schmidt et al. (2010; extrolites), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, holotype and ex-type of type species unavailable.

Hymenoscyphus Gray 1821 (= *Belospora* Clem. 1909; = *Ciboriella* Seaver 1951; = *Hymenoscypha* (Fr.) W. Phillips 1887; = *Peziza* trib. *Hymenoscypha* Fr. 1822; = *Septatium* Velen. 1934 *fide Species Fungorum* 2017), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, c. 170 species, type: *H. fructigenus* (Bull.) Gray, asexual morph ?hyphomycetous, pathogens, saprobes, terrestrial, North and South America, see Lumbsch and Huhndorf (2010; outline), Queloz et al. (2011; cryptic species), Kirk et al. (2013; genus accepted), Baral et al. (2014; nomenclature), Gross et al. (2014; pathogens), Zheng and Zhuang (2014; new species, China), cultures and sequences are available.

Hymenorrendiella P.R. Johnst., Baral & R. Galán 2014, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, nine species, type: *H. eucalypti* (Berk.) P.R. Johnst., Baral & R. Galán, asexual morph unknown, saprobes, terrestrial, Europe, see Johnston et al. (2014a; taxonomy, DNA, phylogeny), cultures and sequences are available.

Hyperphyscia Müll. Arg. 1894 (= *Dimelaena* sect. *Tetramelaena* Trevis. 1868; = *Physcia* sect. *Tetramelaena* (Trevis.) Zahlbr. 1803; = *Physciopsis* M. Choisy 1950; = *Tetramelaena* (Trevis.) C.W. Dodge 1971 *fide Species Fungorum* 2017), *Physciaceae*, *Caliciales*, *Lecanoromycetes*, nine species, type: *H. synthalea* (C. Knight) Müll. Arg., lichenized, worldwide, see Esslinger et al. (2012; new species), Lücking et al. (2017a; classification), cultures and sequences are available.

Hypochlaena Cif. 1962, *Ascomycota* genera *incertae sedis*, one species, type: *H. caespitosa* Cif., sterile

mycelium, sexual morph unknown, saprobes, terrestrial, Dominican Republic, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypoconis Egidi & Quaedvli. 2014, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *H. sterilis* Egidi & Quaedvli., hyphomycetous, sexual morph unknown, rock-inhabiting fungus, see Egidi et al. (2014; taxonomy), Wijayawardene et al. (2017; outline), cultures available for the type: CBS 118321 =TRN278.

Hyphodiscosia Lodha & K.R.C. Reddy 1974, *Ascomycota* genera *incertae sedis*, five species, type: *H. jaipurensis* Lodha & K.R.C. Reddy, hyphomycetous, saprobes, pathogen, terrestrial, Argentina, Australia, Brazil, Cuba, Czechoslovakia, India, Japan, Mexico, Nigeria, Poland, Russia, Taiwan, Thailand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species, Holotype and ex-type of type species RUBL 730.

Hyphodiscosiodes Matsush. 1993, *Ascomycota* genera *incertae sedis*, one species, type: *H. tambopataensis* Matsush., hyphomycetous, saprobes, terrestrial, Peru, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, Holotype and ex-type of type species MFC 1P-449.

Hyphodiscus Kirschst. 1907 (= *Catenulifera* Hosoya 2002; = *Incrupilella* Svrček 1986; = *Tanglella* Höhn. 1918 fide Johnston et al. 2014b; Species Fungorum 2017), *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, c. 20 species, type: *H. gregarius* Kirschst., asexual morph *Catenulifera* (hyphomycetous), saprobes, north temperate, see Huhtinen et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Hosoya et al. (2011; DNA, phylogeny, new species), Pärte and Pöldmaa (2011; new species), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature; new combinations), cultures and sequences are available but lacking for the type species.

Hyphopeziza J.G. Han, Hosoya & H.D. Shin 2014, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *H. pygmaea* (Mouton) J.G. Han, Hosoya & H.D. Shin, asexual morph unknown, saprobes, terrestrial, worldwide, see Han et al. (2014; taxonomy), cultures and sequences are unavailable, holotype and available cultures of the type species Mouton 473 (BR), KUS F51564.

Hyhopichia von Arx & van der Walt 1976, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, nine species, type: *H. burtonii* (Boidin, Pignal, Lehodey, Vey & Abadie) von Arx & van der Walt,

aseexual reproduction is by multilateral budding, saprophytic, on pollen, alfalfa silage, grain, various fruits, human skin, sputum, decaying insect infested wood, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Hypopolynema Nag Raj 1977, *Ascomycota* genera *incertae sedis*, six species, type: *H. tropicale* Nag Raj, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, Brazil, Colombia, India, USA, UK, see Pinho et al. (2010; morphology, new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Hyposcypha Velen. 1934, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *H. loricata* Velen., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hyposoma Syd. 1924, *Metacapnodiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *H. hypoxylloides* Syd., see Kirk et al. (2008; nom. conf.).

Hypostereum Pat. 1892, *Ascomycota* genera *incertae sedis*, one species, type: *H. pendulum* Pat., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypothyrium B. Sutton & Pascoe 1989, *Ascomycota* genera *incertae sedis*, one species, type: *H. pallidum* B. Sutton & Pascoe, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypobryon Döbbeler 1983, *Dothideomycetes*, genera *incertae sedis*, seven species, type: *H. poeltii* (Döbbeler) Döbbeler, asexual morph unknown, on hepatic, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hypocenomyce M. Choisy 1951, *Ophioparmaceae*, *Umbilicariales*, *Lecanoromycetes*, four species, type: *H. scalaris* (Ach. ex Lilj.) M. Choisy, lichenized, worldwide, see Björk (2010; USA), Bendiksby and Timdal (2013, phylogeny, taxonomy), Stepanchikova et al. (2013; Russia), Miadlikowska et al. (2014a; Poland), Lücking et al. (2017a; classification), cultures and sequences are available, holotype and ex-type of type species unavailable.

Hypocline Syd. 1939, *Ascomycota* genera *incertae sedis*, one species, type: *H. penniseti* Syd.,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, Dominican Republic, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypocopra (Fr.) Kickx f. 1867 (= *Coprolepa* Fuckel 1870; = *Coprolepa* Fuckel 1870 [1869–70]; = *Massaria* subgen. *Hypocopra* Fr. 1849 *fide* Species Fungorum 2017; Wendt et al. 2017), Xylariaceae, Xylariales, Sordariomycetes, c. 35 species, type: *H. merdaria* (Fr.) J.J. Kickx, asexual morph unknown, terrestrial, saprobes, mainly coprophilous, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jayanetti et al. (2014; applications), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type species.

Hypocrella Sacc. 1878 (= *Aschersonia* Mont. *fide* Maharachchikumbura et al. 2015b, 2016b; Rossman et al. 2016a), Clavicipitaceae, Hypocreales, Sordariomycetes, 43 species, type: *H. discoidea* (Berk. & Broome) Sacc., asexual morph *Aschersonia*, saprobes, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, nomenclature), Rossman et al. (2016a; nomenclature), Tibpromma et al. (2017b; new species), cultures and sequences are available.

Hypocreodendron Henn. 1897 (= *Discoxylaria* J.C. Lindq. & J.E. Wright 1964; = *Poroniopsis* Speg. 1922 *fide* Rébllová et al. 2016c; Species Fungorum 2017), Xylariaceae, Xylariales, Sordariomycetes, one species, type: *H. sanguineum* Henn., hyphomycetous, sexual morph formerly known in *Discoxylaria*, saprobes, Argentina, Mexico, see Lumbsch and Huhndorf (2010; outline), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Stadler et al. (2013), Rébllová et al. (2016c; nomenclature), Daranagama et al. (2017; morphology, taxonomy, key, sexual-aseexual link), sequence data available for one species *D. myrmecophila*.

Hypocreopsis P. Karst. 1873 (= *Dozya* P. Karst. 1873 *fide* Species Fungorum 2017), Hypocreaceae, Hypocreales, Sordariomycetes, three species, type: *H. riccioidea*, (Bolton) P. Karst., asexual morph hyphomycetous, saprobes and/or pathogens, terrestrial, Atlantic coast of Europe, Eastern USA, South-Eastern Australia, New Zealand, see Lumbsch and Huhndorf (2010; outline), Grundy et al. (2012; molecular analysis), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Hypoderma De Not. 1847 (= *Epidermella* Tehon 1935; = *Hypoderopsis* Kuntze 1898; = *Locelliderma* Tehon

1935; = *Schizochorella* Höhn. 1918 *fide* Species Fungorum 2017), Rhytismataceae, Rhytismatales, Leotiomycetes, one species, type: *H. rubi* (Pers.) DC., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Lantieri et al. (2011; phylogeny, new species), Lantz et al. (2011; DNA, phylogeny), Wang et al. (2013d; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available, lectotype 'Hb. Pers. mis Chaill.

Hypodermella Tubeuf 1895, Rhytismataceae, Rhytismatales, Leotiomycetes, three species, type: *H. laricis* Tubeuf, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypodermellina Höhn. 1917, Rhytismataceae, Rhytismatales, Leotiomycetes, one species, type: *H. ruborum* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypodermina Höhn. 1916 (= *Daedala* Hazsl. 1887; = *Mazzantiella* Höhn. 1918 *fide* Species Fungorum 2017), Ascomycota genera incertae sedis, one species, type: *H. nervisequa* (Link) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypoftavia Marbach 2000, Caliciaceae, Caliciales, Lecanoromycetes, three species, type: *H. velloziae* (Kalb) Marbach, lichenized, Brazil, Thailand, Chile, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Hypogloeum Petr. 1923, Ascomycota genera incertae sedis, one species, type: *H. euonymi* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hypogymnia (Nyl.) Nyl. 1896 (= *Ceratophyllum* M. Choisy 1951; = *Parmelia* subgen. *Hypogymnia* Nyl. 1881 *fide* Species Fungorum 2017), Parmeliaceae, Lecanorales, Lecanoromycetes, 90 species, type: *H. physodes* (L.) Nyl., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Miądlikowska et al. (2011, phylogeny), Kirk et al. (2013; genus accepted), McCune and Wang (2014, inventory), Lücking et al. (2017a; classification), cultures and sequences are available.

Hypohelion P.R. Johnst. 1990 (= *Leptostroma* Fr. 1815 *fide* Johnston et al. 2014b), *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, three species, type: *H. scirpinum* (DC.) P.R. Johnst., asexual morph *Leptostroma*, saprobes, north temperate, see Lumbsch and Huhndorf (2010; outline), Lantz et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Wang et al. (2014b; DNA, phylogeny), cultures and sequences are available, cultures available for type: Lantz 394 (UPS) (*fide* Lantz et al. 2011).

Hypomyces (Fr.) Tul. 1870 (= *Cladobotryum* Nees 1816; = *Sepedonium* Link 1809; = *Stephanoma* Wallr. 1833 *fide* Jaklitsch et al. 2016a; see Species Fungorum 2017 for additional synonyms), *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, more than 150 species, type: *H. lactifluorum* (Schwein.) Tul. & C. Tul., asexual morph hyphomycetous, fungicolous, worldwide, see Jaklitsch et al. (2008; new species), Lumbsch and Huhndorf (2010; outline), Pöldmaa (2011; new species, red pigment), Seifert et al. (2011; morphology of asexual morphs), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2012, 2017; outline), Gea et al. (2016; pathogenicity of *Cladobotryum*), Jaklitsch et al. (2016a; classification, nomenclature), sequence data available, holotype of type: CBS H-19806 (as *Sphaeria lactifluorum* Schw., Syn. Fung. 1190-44 Herb.), ex-type strains are unavailable for type species.

Hyponectria Sacc. 1878 (= *Anisostomula* Höhn. 1919 *fide* Species Fungorum 2017), *Hyponectriaceae*, *Amphisphaerales*, *Sordariomycetes*, c. 30 species, type: *H. buxi* (DC.) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhao and Zhao (2012; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), DNA sequences and cultures available (*fide* Winka and Eriksson 2000), type specimen of type: CBS H-4954.

Hypophloeda K.D. Hyde & E.B.G. Jones 1989, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *H. rhizospora* K.D. Hyde & E.B.G. Jones, asexual morph unknown, marine, mangrove roots, Brunei, Seychelles, see Jones et al. (2009b; key, description), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Hypospilina (Sacc.) Traverso 1913 (= *Hyospila* subdiv. *Hypospilina* Sacc. 1883 *fide* Species Fungorum 2017), *Valsaceae*, *Diaporthales*, *Sordariomycetes*, four species,

type: *H. bifrons* (DC.) Traverso, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Hypotarzetta Donadini 1985, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *H. insignis* (Berthet & Riousset) Donadini, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Hypotrachyna (Vain.) Hale 1974 (= *Parmelia* sect. *Hypotrachyna* Vain. 1890 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 260 species, type: *H. brasiliiana* (Nyl.) Hale, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Flakus et al. (2012; new species), Divakar et al. (2013a; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Hypotrachynicola Etayo 2002, *Ascomycota* genera *incertae sedis*, one species, type: *H. rubra* Etayo, asexual morph unknown, lichenicolous, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable, holotype of the type: COL Etayo 16515.

Hypoxylon Bull. 1791 (= *Nodulisporium* Preuss 1849 *fide* Réblová et al. 2016c; see Wendt et al. 2017 for additional synonyms), *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, c. 140 species, type: *H. fragiforme* (Pers.) J. Kickx f., asexual morph formerly known in *Nodulisporium*, terrestrial, saprobes, mainly coprophilous, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), Sir et al. (2016a; new species), Dai et al. (2017; new species), Wendt et al. (2017; phylogeny, resurrected *Hypoxylaceae*), cultures and sequences are available, needs generic revision as the genus comprises more than 1000 species epithets (Index Fungorum 2017).

Hypsostroma Huhndorf 1992, *Hypsostromataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *H. saxicola* Huhndorf, asexual morph unknown, saprobes, terrestrial, worldwide, see Mugambi and Huhndorf (2009a; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Hypsostromataceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available, cultures available for the type: SMH 5005, holotype of type species NY, Harris 26462.

Hysteridium P. Karst. 1905, *Ascomycota* genera *incertae sedis*, one species, type: *H. phragmitis* P. Karst., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hysterium Pers. 1791 (= *Pragmoparopsis* Höhn. 1917 *fide* Species Fungorum 2017), *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, 14 species, type: *H. pulicare* (Lightf.: Fr.) Pers., asexual morph unknown, saprobes, terrestrial, worldwide, see Boehm et al. (2009a, b, overview of genus), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Tibpromma et al. (2017b; phylogeny), cultures and sequences are available, cultures available for type species CBS 123377, holotype of type: CBS H-5484.

Hysterothecium E. Boehm & C.L. Schoch 2010, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, four species, type: *H. mori* (Schwein.) E.W.A. Boehm & C.L. Schoch, asexual morph coelomycetous (apospshaeria-like), saprobes, terrestrial, worldwide, see Boehm et al. (2009a, b; overview), Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), Tibpromma et al. (2017b; phylogeny), cultures and sequences are available, holotype and ex-type of type: BPI878733, CBS 123336.

Hysterocarpina H. Zogg 1949, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, one species, type: *H. paulistae* H. Zogg, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hysterodictum D.A.C. Almeida, Gusmão & A.N. Mill. 2014, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, one species, type: *H. partisporum* D.A.C. Almeida, Gusmão & A.N. Mill, asexual morph unknown, saprobes, terrestrial, Brazil, see Almeida et al. (2014b; taxonomy), cultures and sequences are available, holotype of type: HUEFS:42865

Hysterodiscula Petr. 1942, *Ascomycota* genera *incertae sedis*, one species, type: *H. kalmiae* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hysterogonium Rehm ex Lindau 1896, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, two species, type: *H. ovatum* (Cooke) Lindau, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hysterographium Corda 1842, *Pleosporomycetidae* genera *incertae sedis*, *Dothideomycetes*, two species, type: *H. elongatum* (Wahlenb.) Corda, asexual morph unknown, saprobes, terrestrial, worldwide, see Boehm et al. (2009a, b; taxonomy, DNA, phylogeny), Mugambi and Huhndorf (2009b; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; key to distinguish from other histeriaceous genera), Kirk et al. (2013; genus accepted), Logesh et al. (2013; new species, India), cultures and sequences are available.

Hysteronaevia Nannf. 1984, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, twelve species, type: *H. holoschoenii* (De Not.) Nannf., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hysteropatella Rehm 1890, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, three species, type: *H. prostii* (Duby) Rehm, asexual morph unknown, saprobes, terrestrial, Switzerland, see Boehm et al. (2009a; phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as genus in *Hysteriaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013) did not accept as a genus in *Hysteriaceae*, Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), cultures and sequences are available, cultures available for the type species H.B. 9934b

Hysteropeltella Petr. 1923, *Schizophyriaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *H. moravica* Petr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as *Dothideomycetes*, genera *incertae sedis*), Ariyawansa et al. (2013b; transferred to *Schizophyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hysteropezizella Höhn. 1917, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, 19 species, type: *H. subvelata* (Rehm) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hysteropsis Rehm 1887, *Dothideomycetes* genera *incertae sedis*, four species, type: *H. culmigena* Rehm, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hysteropycnis Hiltizer 1929, *Ascomycota* genera *incertae sedis*, one species, type: need typification, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Hysterostegiella Höhn. 1917 (= *Hysteropezizella* sect. *Hysterostegiella* (Höhn.) Nannf. 1932; = *Stegopeziza* Höhn. 1917 *fide* Species Fungorum 2017), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, ten species, type: *H. fenestrata* (Roberge ex Desm.) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype and ex-type of type species unavailable.

Hysterostomella Speg. 1885 (= *Hysterostomina* Theiss. & Syd. 1915 *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, 40 species, type: *H. guaranitica* Speg., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Hyweljonesia R.G. Shivas, Y.P. Tan, Marney & Abell 2016, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *H. queenslandica* R.G. Shivas, Y.P. Tan, Marney & Abell, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type of the type: BRIP 61322b.

Ialomitzia Gruia 1964, *Ascomycota genera incertae sedis*, one species, type: *I. cavernicola* Gruia, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Icmadophila Trevis. [nom. cons.] 1852 (= *Cystolobis* Clem. 1909; = *Glossodium* Nyl. 1855; = *Icmadophilomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Knightiella* Müll. Arg. 1886; = *Phycodiscis* Clem. 1909; = *Thelidea* Hue 1902 [1901]; = *Tupia* L. Marchand 1830 *fide* Species Fungorum 2017), *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, eight species, type: *I. aeruginosa* (Scop.) Trevis., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Idiocercus B. Sutton 1967, *Ascomycota genera incertae sedis*, two species, type: *I. pirozynskii* B. Sutton, coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, Australia, Tanzania, Sierra Leone, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type species IMI 107154a.

Idiomyces Thaxt. 1893, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *I. peyritschii* Thaxt., asexual morph unknown, biotrophic, terrestrial, Europe, Japan, see Lumbsch and Huhndorf

(2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Idriella P.E. Nelson & S. Wilh. 1956 (= *Chloridiella* G. Arnaud 1954 [1953] *fide* Species Fungorum 2017), *Microdochiaeae*, *Xylariales*, *Sordariomycetes*, c. 30 species, type: *I. lunata* P.E. Nelson & S. Wilh., hyphomycetous, sexual morphs unknown, in soil, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2016b; DNA, phylogeny), cultures and sequences are available.

Idriellopsis Hern.-Restr. & Crous 2015, *Xylariales*, genera *incertae sedis*, *Sordariomycetes*, one species, type: *I. uncinospora* (R.F. Castañeda & W.B. Kendr.) Hern.-Restr. & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Hernández-Restrepo et al. (2016b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Igneocumulus A.W. Ramaley 2003, *Ascomycota genera incertae sedis*, ten species, type: *I. yuccae* A.W. Ramaley, asexual morph hyphomycetous (lecythophora-like), saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species UC 1713843.

Ijuhya Starbäck 1899 (= *Lepidonectria* Speg. 1910; = *Mollisia* subgen. *Peristomialis* W. Phillips 1887 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, 14 species, type: *I. vitrea* Starbäck, asexual morph acremonium-like, saprobes, terrestrial, worldwide, see Lechat and Courtecuisse (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Lechat and Fournier (2017; new species), cultures and sequences are available.

Illosporiopsis D. Hawksw. 2001, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *I. christiansenii* (B.L. Brady & D. Hawksw.) D. Hawksw., lichenicolous, Europe, terrestrial, see Seifert et al. (2011; genus accepted), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: K(M)-IMI 226836.

Illosporium Mart. 1817 (= *Lepra* Willd. 1787 *fide* Species Fungorum 2017), *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, 18 species, type: *I. carneum* Fr., asexual morph hyphomycetous, lichenicolous, USA and Europe, see Pindi (2012; marine), Kossowska and Szczepańska (2013; Poland), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ilyomyces F. Picard 1917, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *I. mairei* F. Picard, asexual morph unknown, biotrophic, terrestrial, Europe, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ilyonectria P. Chaverri & C. Salgado 2011, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, 22 species, type: *I. radicicola* (Gerlach & L. Nilsson) P. Chaverri & Salgado, root pathogenic or saprobes, cosmopolitan, see Chaverri et al. (2011; phylogeny), Cabral et al. (2012; new species), Lombard et al. (2015b; phylogeny), cultures and sequences are available, ex-type of type species CBS 264.65.

Ilytheomyces Thaxt. 1917, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 15 species, type: *I. elegans* Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Imia Trappe & Kovács 2008, *Morchellaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *I. gigantea* (S. Imai) Trappe & Kovács, asexual morph unknown, ectomycorrhizal, saprobes, terrestrial, North American, East Asia, see Kovács et al. (2008; DNA, taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), cultures and sequences are available, Holotype and ex-type of type species unavailable.

Imazekia Tak. Kobay. & Y. Kawabe 1992, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *I. ryukyuensis* Tak. Kobay. & Y. Kawabe, asexual morph unknown, from living leaves, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable Holotype of type species TFM, FPH-7163.

Imicles Shoemaker & Hambl. 2001, *Ascomycota* genera *incertae sedis*, six species, type: *I. leptospora* (Sacc. & Roum.) Shoemaker & Hambl., hyphomycetous, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, compare with *Sporidesmium*.

Immersaria Rambold & Pietschm. 1989, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, eight species; type: *I. athroocarpa* (Ach.) Rambold & Pietschm., lichenized, Asia, Africa, Europe, Australia, see Lumbsch and Huhndorf (2010; outline), Valadbeigi et al. (2011b; new species), Kirk et al. (2013; genus accepted), Pandit (2014; India), Gasparyan et al. (2016; Armenia), Lücking et al. (2017a; classification), cultures and sequences are available.

Immersidiscosia Kaz. Tanaka, Okane & Hosoya 2011, *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *I. eucalypti* (Pat.) Kaz. Tanaka, Okane & Hosoya, sexual morph unknown, saprobes, terrestrial, France, Japan, Tunisia, see Tanaka et al. (2011a; morphology, phylogeny), Jaklitsch et al. (2016a; phylogeny), cultures and sequences are available, holotype of the type species in FH.

Immersiella A.N. Mill. & Huhndorf 2004, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, two species, type: *I. immersa* (P. Karst.) A.N. Mill. & Huhndorf, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type species SMH2589, SMH4104, holotype and ex-type of type species unavailable.

Immersiporthe S.F. Chen, M.J. Wingf. & Jol. Roux [2012] 2013, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *I. knoxdaviesiana* S.F. Chen, M.J. Wingf. & Jol. Roux, coelomycetous, sexual morph unknown, pathogens, terrestrial, South Africa, see Chen et al. (2013e; taxonomy, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype and the ex-type of type: PREM 60738, CMW37314 =CBS132862.

Immersisphaeria Jaklitsch 2007, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *I. eichleriana* (Bres.) Jaklitsch, asexual morph unknown, pathogens, terrestrial, South Africa, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Daranagama et al. (2016d; accepted in *Xylariales*), cultures and sequences are unavailable.

Immotthia M.E. Barr 1987, *Dothideomycetes* genera *incertae sedis*, one species, type: *I. hypoxylon* (Ellis & Everh.) M.E. Barr, asexual morph coniothyrium-like, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline, misspelled as *Immotthia*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Imshaugia F.C. Mey. 1985, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, seven species, type: *I. aleurites* (Ach.) S.L.F. Mey., lichenized, boreal and temperate regions of the Northern Hemisphere and from cool-montane regions in the Southern Hemisphere, see Hauck et al. (2009; acid tolerance), Lumbsch and Huhndorf (2010; outline), Thell et al. (2012; description, phylogeny), Kirk et al. (2013; genus accepted), Kharpukhayeva and Mukhortova (2016; Baikal), Lücking et al. (2017a; classification), sequences are available.

Inaequalispora L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *I. prestonii* (M.C. Tulloch) L. Lombard & Crous, hyphomycetous, sexual morph unknown, from forest soil, terrestrial, Malaysia, see Lombard et al. (2016; taxonomy, phylogeny), holotype and ex-isotype of the type: CBS H-7392, CBS 175.73.

Incertomyces Egidi & Zucconi 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *I. perditus* Egidi & Zucconi, hyphomycetous, sexual morph unknown, rock-inhabiting fungus, Italy, see Egidi et al. (2014; taxonomy), Wijayawardene et al. (2014c, 2017; outline), type: CBS 136105 =CCFEE 5385, cultures and sequences are available.

Incrucipulum Baral 1985, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *I. ciliare* (Schrad.) Baral, asexual morph unknown, saprobes, terrestrial, temperate, see Hosoya et al. (2010; new species, Japan, DNA), Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Incrupila Raity. 1970, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, ten species, type: *I. aspidii* (Lib.) Raity., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Induratia Samuels, E. Müll. & Petrini 1987, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *I. apiospora* Samuels, E. Müll. & Petrini, asexual morph nodulosporium-like, saprobes, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2013), Daranagama et al. (2017; morphology, taxonomy, key), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), sequences are unavailable.

Inesiosporium R.F. Castañeda & W. Gams 1997, *Ascomycota* genera *incertae sedis*, two species, type: *I. longispirale* (R.F. Castañeda) R.F. Castañeda & W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; valid genus), Argôlo Magalhães et al. (2014; Bahia), Castañeda-Ruiz et al. (2016a; distribution), holotype: INIFAT C84/107, cultures and sequences are unavailable, compare with *Ceratophorum* and *Zalerion*.

Inflatispora Yin. Zhang, J. Fourn. & K.D. Hyde 2011, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *I. pseudostromatica* Ying Zhang, J. Fourn. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, France and Thailand, see Zhang et al. (2011e; taxonomy, phylogeny), Tibpromma et al. (2017b; taxonomy, phylogeny), cultures and sequences are available, holotype and ex-type of type: IFRD 8989, CBS 123110.

Infundibulomyces Plaingam, Somrith. & E.B.G. Jones 2003, *Chaetosphaeriaceae*, *Chaetosphaerales*, *Sordariomycetes*, two species, type: *I. cupulata* Plaingam, Somrith. & E.B.G. Jones, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available, epitype and ex-epitype of the type: SFC 1615, BCC 13400.

Infundichalara Réblová & W. Gams 2011, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *I. microchona* (W. Gams) Réblová & W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Netherlands, Czech Republic, see Réblová and Gams (2011; morphology, phylogeny), Koukol (2012; morphology, new species), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, holotype and ex-type strains of the type: CBS-H 19895, CBS 261.82.

Ingaderia Darb. 1897 (= *Darbshirella* Zahlbr. 1898; = *Dictyographa* Darb. 1897 fide Species Fungorum 2017), *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *I. pulcherrima* Darb., lichenized, Chile, see Lumbsch and Huhndorf (2010; outline), Ertz and Tehler (2011; phylogeny), Notov (2014; outline), Lücking et al. (2017a; classification), sequences are available.

Ingvariella Guderley & Lumbsch 1997, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *I. bispora* (Bagl.) Guderley & Lumbsch, lichenized, arid and semi-arid area, see Lumbsch and Huhndorf (2010; outline), Fernández-Brime et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Inifatiella R.F. Castañeda, *Ascomycota* genera *incertae sedis*, one species, type: *I. nectandrae* R.F. Castañeda, coelomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Inocyclus Theiss. & Syd. 1915 (= *Aspidothaea* Syd. 1927; = *Ellimonia* Syd. 1930; = *Fraserula* Syd. 1938 fide Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, eight species, type: *I. psychotriae* (Syd. & P. Syd.) Theiss. & Syd, asexual morph unknown, epiphytes, pathogens, biotrophic, terrestrial, worldwide, see Hofmann (2009; review, placed in *Parmulariaceae*), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; review, keys), Wijayawardene et al. (2014c; outline), Guatimosim et al. (2015; phylogeny), cultures and sequences are available.

Inoderma (Ach.) Gray 1821 (= *Phaeothrombis* Clem. 1909; = *Verrucaria* sect. *Inoderma* (Ach.) W. Mann 1825;

= *Werneria* Zschacke ex Werner 1934; = *Werneromyces* Werner 1976 *fide Species Fungorum* 2017, Arthoniaceae, Arthoniales, Arthoniomycetes, three species, type: *I. byssaceum* (Weigel) Gray, lichenized, temperate Europe, North America, Eastern Siberia, Japan, and eastern Africa (Uganda), see Frisch et al. (2015; phylogeny, key, description), Lücking et al. (2017a; classification), sequences are available.

Intercalarispora J.L. Crane & Schokn. 1983, *Ascomycota* genera *incertae sedis*, one species (probably more described as synasexual morphs of *Helicoma*, *Monodictys* or *Xenosporium*), type: *I. nigra* J.L. Crane & Schokn., hyphomycetous, sexual morph unknown (probably *Tubeufia*), saprobes, on rotten submerged wood, America and Africa, see Hyde et al. (2011; checklist), Kirk et al. (2013; valid genus), cultures and sequences are unavailable, holotype of type: ILLS 43047.

Intralichen D. Hawksw. & M.S. Cole 2002, *Ascomycota* genera *incertae sedis*, four species, type: *I. christiansenii* (D. Hawksw.) D. Hawksw. & M.S. Cole, sexual morph unknown, lichenicolous, North temperate, see Seifert et al. (2011; genus accepted), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Involutopyrenium Breuss 1996, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, nine species, type: *I. waltheri* (Kremp.) Breuss, lichenized, Europe, North America, Asia, see Prieto et al. (2010a; key, description), Prieto et al. (2012; phylogeny), Breuss (2016; nomenclature), Lücking et al. (2017a; classification), sequences are available, holotype of type species unavailable.

Involutroscypha Raitv. 2002, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, ten species, type: *I. involucrata* (B. Erikss.) Raitv., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type species TAA165831.

Iodophanus Korf 1967, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, 16 species, type: *I. carneus* (Pers.) Korf, asexual morph unknown, coprophilous, terrestrial, worldwide, see Schmitt et al. (2008; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, type material of type species CBS H-6320.

Iodosphaeria Samuels, E. Müll. & Petrini 1987, *Iodosphaeriaceae*, *Xylariales*, *Sordariomycetes*, three species, type: *I. phyllophila* (Mouton) Samuels, E. Müll. & Petrini, asexual morph hyphomycetous (selenosporella-like and ceratosporium-like), saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura

et al. (2015b, 2016b; outline), cultures and sequences are available, holotype of the type: PDD 47872.

Iodowynnea Medel, Guzmán & S. Chacón 1996, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *I. auriformis* (Pat. ex Le Gal) Medel, Guzmán & S. Chacón, asexual morph unknown, coprophilous, terrestrial, worldwide, see Schmitt et al. (2008; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Ionaspis Th.Fr. 1871 (= *Aphragmia* Trevis. 1880; = *Durietzia* Gyeln. 1935 *fide Species Fungorum* 2017), *Hymeneliaceae*, *Hymeneliales*, *Lecanoromycetes*, seven species, type: *I. chrysophana* (Körb.) Th. Fr., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Ionophragmium Peres 1961, *Ascomycota* genera *incertae sedis*, one species, type: *I. notatum* Peres & L. Xavier, coelomycetous, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, compare with *Asterostomella*, holotype of type: IMUR 20676.

Ioplaca Poelt 1977, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *I. sphalerata* Poelt, lichenized, Himalayas, see Lumbsch and Huhndorf (2010; outline), Arup et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); cultures and sequences are available.

Irianiella Petr. 1949, *Sordariomycetes* genera *incertae sedis*, one species, type: *I. rechingeri* Petr., asexual morph unknown, saprobes, terrestrial, Iran, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Irenopsis F. Stevens 1927 (= *Meliola* subgen. *Irenopsis* (F. Stevens) Cif. 1938 *fide Species Fungorum* 2017), *Meliolaceae*, *Meliolales*, *Sordariomycetes*, 75 species, type: *I. tortuosa* (G. Winter) F. Stevens, asexual morph produced from the hyphae, saprobes, pathogens, terrestrial, worldwide, see Hosagoudar and Archana (2010b; new species, India), Hosagoudar and Sabeena (2010; new species, India), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pinho et al. (2013; new species, Brazil, DNA, phylogeny), Hosagoudar et al. (2014; new species, Andaman Islands), Hongsanan et al. (2015d; new species, taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), Rodríguez et al. (2015; new species), cultures and sequences are available, holotype and ex-type of type species unavailable.

Iridinea Velen. 1934, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, two species, type: need typification,

asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, data inadequate.

Irpicomycetes Deighton 1969, *Ascomycota* genera *incertae sedis*, three species, type: *I. schiffnerulae* Deighton, hyphomycetous, sexual morph unknown, on leaves or twigs, Malaysia and Japan, see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species IMI 54929b.

Irydonia Racib. 1900, *Rhytismatales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *I. filicis* Racib., asexual morph unknown, saprobes, Java, see Lumbsch and Huhndorf (2010; outline), holotype and ex-type of type species unavailable.

Isalonactis Ertz, Tehler, Eb. Fisch., Killmann, Razafin. & Sérus. 2014, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *I. madagascariensis* Ertz, Tehler, Eb. Fisch., Killmann, Razafin. & Sérus., sexual morph with apothecoid ascomata, lichenized, terrestrial, Madagascar, see Ertz et al. (2014c; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available, lectotype and ex-type strain of type: Ertz 13024.

Isaria Pers. 1794 (= *Alphitomyces* Reissek 1856; = *Amphichorda* Fr. 1825; = *Coremiopsis* Sizova & Suprun 1957; = *Beejasamuha* Subram. & Chandrash. 1977 *fide* Species Fungorum 2017), *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, c. 13 species, type: *I. farinosa* (Holmsk.) Fr., hyphomycetous, entomogenous, worldwide, see Kim et al. (2010; thermotolerance), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, epitype and ex-type of type: CBS-H6586, CBS 111113.

Isariella Henn. 1909, *Ascomycota* genera *incertae sedis*, two species, type: *I. auerswaldiae* Henn., hyphomycetous, saprobes, terrestrial, Brazil, Dominican Republic, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), compare with *Atractilina*, sequences are unavailable, holotype of type species S-F40445.

Ischnostroma Syd. & P. Syd. 1914, *Ascomycota* genera *incertae sedis*, one species, type: *I. merrillii* Syd. & P. Syd., coelomycetous, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), compare with *Trachythriolum*, sequences are unavailable.

Isia D. Hawksw. & Manohar. 1978, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *I. neocaldoniensis* (C. Moreau) D. Hawksw. &

Manohar, on plant material, Asia and Pacific, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Isomunkia Theiss. & Syd. 1915, *Dothideomycetes* genera *incertae sedis*, one species, type: *I. pulvinula* (Pat.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, Ecuador, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Isothea Fr. 1849 (= *Didymella* Speg. 1879 *fide* Species Fungorum 2017), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *I. rhytismaoides* (Bab.) Fr., asexual morph unknown, from living leaves, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), DNA sequences are unavailable.

Isthmoconidium Etayo & Berger 2013, *Ascomycota* genera *incertae sedis*, one species, type: *I. bacidiicola* Etayo & F. Berger, coelomycetous, sexual morph unknown, parasymbiont on *Bacidia*, terrestrial, Caribbean, see Etayo and Berger (2013; morphology, distinction of similar lichenicolous genera), sequences are unavailable, holotype of the type species LI, F. Berger 22323 & S. La Greca.

Isthmolongispora Matsush. 1971, *Ascomycota* genera *incertae sedis*, eleven species, type: *I. intermedia* Matsush., hyphomycetous, saprobes, aquatic and terrestrial, Argentina, Australia, Brazil, China, Cuba, Hungary, Japan, Malaysia, Papua New Guinea, Peru, Portugal, Scotland, Taiwan, United Kingdom, Venezuela and West Indies, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), compare with *Isthmophragmospora* and *Wiesneriomycetes*, cultures and sequences are available, holotype of type: MFC-3052.

Isthmophragmospora Kuthub. & Nawawi 1992, *Ascomycota* genera *incertae sedis*, two species, type: *I. verruculosa* Kuthub. & Nawawi, hyphomycetous, sexual morph unknown, saprobes, on submerged rotten angiosperms twigs, on submerged decaying leaves in a black water stream, Malaysia and Brazil, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Monteiro et al. (2014a; new species), sequences are unavailable, holotype of type: IMI 344358.

Isthmospora F. Stevens 1918 (= *Carlosia* G. Arnaud 1954 [1953] *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, three species, type: *I. spinosa* F. Stevens, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Pantropical, see Seifert et al. (2011; morphology, accepted as a genus in *Microthyriaceae*),

Wijayawardene et al. (2012, 2014c; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Isthmosporella Shearer & J.L. Crane 1999, *Dothideomycetes* genera *incertae sedis*, one species, type: *I. pulchra* Shearer & J.L. Crane, asexual morph unknown, saprobes, fresh water, USA, see Lumbsch and Huhndorf (2010; outline, placed in *Phaeosphaeriaceae*), Zhang et al. (2012e; placed in *Pleosporales*), Hyde et al. (2013; accepted in *Phaeosphaeriaceae*), Phookamsak et al. (2014c; excluded from *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes*), sequences are unavailable, holotype of type: ILLS 53086.

Isthmotricladia Matsush. 1971, *Ascomycota* genera *incertae sedis*, three species, type: *I. laeensis* Matsush., hyphomycetous, saprobes, aquatic and terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), compare with *Articulospora*, holotype of type: MFC-2803, cultures are available.

Italiomyces Senan., Camporesi & K.D. Hyde 2017, *Sydiwellaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *I. centaureii* Senan., Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Europe, see Senanayake et al. (2017; taxonomy, phylogeny), holotype of the type: MFLU 16–2866, cultures and sequences are available.

Ityorhoptrum P.M. Kirk 1986, *Ascomycota* genera *incertae sedis*, four species, type: *I. verruculosum* (M.B. Ellis) P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Whitton et al. (2012a; new species), Wijayawardene et al. (2012, 2017; outline), Mel'nik et al. (2014; key species), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ivanpisutia S.Y. Kondr., L. Lokös & Hur 2015, *Lecanorales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *I. oxneri* S.Y. Kondr., Lökös & Hur, lichenized, Russia, see Kondratyuk et al. (2015e; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type: KoLRI 020233.

Iwilsoniella E.B.G. Jones 1991, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *I. rotunda* E.B.G. Jones, asexual morph unknown, marine on wood, Great Britain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: IMI 271838.

Iyengarina Subram. 1958, *Ascomycota* genera *incertae sedis*, three species, type: *I. elegans* Subram., hyphomycetous, sexual morph unknown, saprobes, lignicolous, India, Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of type: MUBL 2015 (Madras, India) (?IMI 113126).

Jackrogersella L. Wendt, Kuhnert & M. Stadler 2017, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, seven species, type: *J. multififormis* (Fr. Fr.) L. Wendt, Kuhnert & M. Stadler, four species, asexual morph hyphomycetous, terrestrial, saprobes, cosmopolitan, see Wendt et al. (2017; taxonomy, phylogeny), cultures and sequences are available.

Jacobsonia Boedijn 1935, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *J. glauca* Boedijn, asexual morph unknown, saprobes, terrestrial, Sumatra, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Jaffuela Speg. 1921, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *J. chilensis* Speg., asexual morph unknown, saprobes, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothioraceae*), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; transferred to *Pseudoperisporiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Jafnea Korf 1960, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, four species, type: *J. fusicarpa* (W.R. Gerard) Korf, saprobes, terrestrial, north temperate, Australia, Tasmania, see Lumbsch and Huhndorf (2010; outline), Gates and van Vooren (2016; new species), cultures and sequences are available.

Jahniella Petr. 1921, *Ascomycota* genera *incertae sedis*, three species, type: *J. bohemica* Petr., coelomycetous, saprobe, pathogen, terrestrial, Czechoslovakia, France, Germany, Russia, Ukraine, see Campbell et al. (2007; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), compare with *Septoria*, sequences are unavailable, holotype of type: IMI 204129.

Jahnula Kirschst. 1936, *Aliquandostipitaceae*, *Jahnulales*, *Dothideomycetes*, 16 species, type: *J. aquatica* (Plöttner & Kirschst.) Kirschst., asexual morph unknown, saprobes, aquatic, worldwide, see Raja et al. (2008a; new species), Sivichai and Boonyuen (2010; new species), Sivichai et al. (2011; asexual morph), Jones et al. (2012; phylogeny), Fournier et al. (2015; new species), cultures and sequences are available, GenBank Number: NR_120015 (*Jahnula bipileata*), holotype and ex-type of type species unavailable.

Jalapriya D'souza, Su, Luo & K.D. Hyde 2016, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *J. pulchra* D'souza, Su, Luo & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, aquatic, cosmopolitan, see Boonmee et al. (2016a; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Jamesiella Lücking, Sérus. & Vězda 2005, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *J. anastomosans* (P. James & Vězda) Lücking, Sérus. & Vězda, lichenized, cosmopolitan, see Hauck (2009, Europe), Lumbsch and Huhndorf (2010; outline), Motiejūnaitė (2011; Lithuania), Yatsyna and Motiejūnaitė (2015; Belarus), Lücking et al. (2017a; classification), sequences are unavailable, holotype of type species unavailable.

Janannfeldtia Subram. & Sekar 1993, *Nitschkiaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *J. karwarensis* Subram. & Sekar, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable, isotype of the type: IMI 324682.

Janetia M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, 22 species, type: *J. euphorbiae* M.B. Ellis, hyphomycetous, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), compare with *Sporidesmium* and *Stanjehughesia*, cultures and sequences are available but lacking for the type species, holotype of type: IMI 163941.

Japewia Tønsberg 1990, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *J. tornoënsis* (Nyl.) Tønsberg, lichenized, cool temperate Northern Hemisphere, see Czarnota (2009; Carpathians), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Allen and Lendemer (2015; key), Lücking et al. (2017a; classification), cultures and sequences are available.

Japewiella Printzen 1999, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *J. carrollii* (Coppins & P. James) Printzen, lichenized, humid temperate and subtropical regions, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Allen and Lendemer (2015; new species, key), Lücking et al. (2017a; classification), sequences are unavailable.

Jarmania Kantvilas 1996, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *J. tristis* Kantvilas, lichenized, Australia, New Zealand, Kerguelen Islands, see Øvstedral and Gremmen (2009; Kerguelen Islands), Lumbsch and Huhndorf (2010; outline), Ludwig and

Kantvilas (2015; New Zealand), Lücking et al. (2017a; classification), sequences are unavailable.

Jarxia D. Hawksw. 1989, *Naetrocymbaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *J. thelenula* (Müll. Arg.) D. Hawksw., on twigs, terrestrial, Caribbean, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Naetrocymbaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Jaklitsch et al. (2016a; position), sequences are unavailable, holotype of type species G.

Jasonhuria S.Y. Kondr., L. Lököö & S.O. Oh 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *J. bogilana* (Y. Joshi & Hur) S.Y. Kondr., L. Lököö, J. Kim, A.S. Kondr. & S.O. Oh, lichenized, South Korea, see Kondratyuk et al. (2015b; genera proposed), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species KoLRI, Hur 041679.

Jattaea Berl. 1900 (= *Phragmocalosphaeria* Petr. 1923; = *Wegelina* Berl. 1900 *fide* Species Fungorum 2017), *Calosphaeriaceae*, *Calosphaerales*, *Sordariomycetes*, twelve species, type: *J. algeriensis* Berl., asexual morph hyphomycetous (phialophora-like), from living leaves, terrestrial, worldwide, see Marincowitz et al. (2008; new species), Lumbsch and Huhndorf (2010; outline), Réblová et al. (2011; taxonomy, revision, new species), Kirk et al. (2013; genus accepted), Réblová et al. (2015b; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Abdel-Wahab et al. (2017, new species), cultures and sequences are available, verified culture for type species STE-U 6201 (*fide* Damm et al. 2008b).

Javaria Boise 1984, *Astrospheariellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *J. samuelsii* Boise, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; morphology, phylogeny), Kirk et al. (2013; genus accepted), Tian et al. (2015; morphology, nomenclature, phylogeny), Phookamsak et al. (2015b; phylogeny), sequences are unavailable.

Javonarxia Subram. ex Subram. 1995 (= *Javonarxia* Subram. 1992 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *J. triseptata* Subram. ex Subram., hyphomycetous, sexual morph unknown, saprobes, on twigs, Malaysia, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of type: MUBL 3124.

Jayarambhatia J. Pratibha 2013, *Ascomycota* genera *incertae sedis*, one species, type: *J. mangrovei* J. Pratibha, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Pratibha (2013; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, holotype of the type: HCIO 51502.

Jenmania W. Wächt. 1897 (= *Jenmaniomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Lichenaceae*, *Lichinales*, *Lichenomycetes*, two species, type: *J. goebelii* W. Wächt., lichenized, British Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species unknown.

Jerainum Nawawi & Kuthub. 1992, *Ascomycota* genera *incertae sedis*, one species, type: *J. triquetrum* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, on submerged rotten twigs, Malaysia, see Hyde et al. (2011; checklist), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of type species KLU 300009.

Jobellisia M.E. Barr 1993, *Jobellisiaceae*, *Calosphaeriales*, *Sordariomycetes*, five species, type: *J. luteola* (Ellis & Everh.) M.E. Barr, asexual morph unknown, saprobes, terrestrial, fresh water, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Untereiner et al. (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, GenBank Number: NR_138381 (*Jobellisia guangdongensis*), holotype of the type: J.P. Morgan 1109.

Joergensenia Passo, Stenroos & Calvelo 2008, *Lecanorales* genera *incertae sedis*, *Lecanomycetes*, one species, type: *J. cephalodina* (Zahlbr.) Passo, S. Stenroos & Calvelo, lichenized, Chile, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type: UPS L-78746.

Johanssonia Sacc. 1889 (= *Johansoniella* Bat., J.L. Bezerra & Cavalc. 1966 *fide* Species Fungorum 2017), *Saccardiaceae*, *Ascomycota* families *incertae sedis*, c. twelve species, type: *J. setosa* (G. Winter) Sacc., asexual morph unknown, saprobes, terrestrial, widespread (tropical), see Crous et al. (2010a; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable for type species.

Johnalcornia Y.P. Tan & R.G. Shivas 2014, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *J. aberrans* (Alcorn) Y.P. Tan & R.G. Shivas, asexual morph hyphomycetous, associated with leaf spots, terrestrial, Australia, see Tan et al. (2014;

morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type: BRIP 16281, isotype: IMI 335210, ex-isotype: CBS 510.91.

Josefpoeltia S.Y. Kondr. & Kärnefelt 1997, *Teloschistaceae*, *Teloschistales*, *Lecanomycetes*, three species, type: *J. boliviensis* S.Y. Kondr. & Kärnefelt, lichenized, Peru, Chile, see Gaya et al. (2008; phylogeny), Arup et al. (2013, taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species MIN (2732).

Jubispora B. Sutton & H.J. Swart 1986, *Ascomycota* genera *incertae sedis*, one species, type: *J. acaciae* B. Sutton & H.J. Swart, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Wijayawardene et al. (2012, 2016b, 2017; morphology, outline), sequences are unavailable, holotype of the type: IMI 167364.

Juglanconis Voglmayr & Jaklitsch 2017, *Juglanconidaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *J. juglandina* (Kunze) Voglmayr & Jaklitsch, asexual morph coelomycetous, saprobes, Europe, see Voglmayr et al. (2017; taxonomy), holotype: PRM 935803, cultures and sequences are available.

Jugulospora N. Lundq. 1972 (= *Coscinospora* Mirza 1963 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, two species, type: *J. rotula* (Cooke) N. Lundq., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, cultures available for the type species ATCC 38359, type material of type species CBS H-13717.

Julella Fabre 1879 (= *Catharinia* (Sacc.) Sacc. 1895; = *Hyalospora* Nieuwl. 1916; = *Peltosphaeria* Berl. 1888; = *Pleospora* subgen. *Catharinia* Sacc. 1883; = *Polyblastiopsis* Zahlbr. 1903 *fide* Species Fungorum 2017), *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, c. 20 species, type: *J. buxi* Fabre, asexual morph unknown, saprobes, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2011b; phylogeny), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014e; phylogeny), cultures and sequences are available, holotype of type: S F599.

Jumillera J.D. Rogers, Y.M. Ju & F. San Martín 1997, *Lopadostomataceae*, *Xylariales*, *Sordariomycetes*, eight species, type: *J. mexicana* J.D. Rogers, Y.M. Ju & F. San Martín, asexual morph libertella-like, with geniculosporium-like synanamorph, saprobes, Rio Grande do Sul, Hawaiian Island, Mexico, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2013), Daranagama et al. (2017 morphology, taxonomy, key), sequences are unavailable,

placement in *Lopadostomataceae* needs to be confirmed by molecular data.

Juncaceicola Tennakoon, Camporesi, Phook. & K.D. Hyde 2016, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, eight species, type: *J. luzulae* Tennakoon, Camporesi, Phook. & K.D. Hyde, asexual morph unknown, saprobes, Italy, see Tennakoon et al. (2016a; taxonomy, phylogeny), Hyde et al. (2016; new species, phylogeny), holotype of the type: HKAS 93700, cultures and sequences are available.

Juncigena Kohlm., Volk.-Kohlm. & O.E. Erikss. 1997 (= *Moheitospora* Abdel-Wahab et al. 2010 *fide* Réblová et al. 2016c), *Juncigenaceae*, *Torpedosporales*, *Sordariomycetes*, two species, type: *J. adarca* Kohlm., Volk.-Kohlm. & O.E. Erikss., asexual morph formerly known in *Moheitospora*, saprobes, marine, widespread, see Abdel-Wahab et al. (2010, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Junctospora Minter & Hol.-Jech. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *J. pulchra* Minter & Hol.-Jech., hyphomycetous, sexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species IMI 243573.

Junewangia W.A. Baker & Morgan-Jones 2002, *Ascomycota* genera *incertae sedis*, four species, type: *J. sphaerospora* W.A. Baker & Morgan-Jones, sexual morph unknown, saprobes, terrestrial, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, holotype of type species AUA.

Kabatia Bubák. 1904 (= *Colletotrichella* Höhn. 1916; = *Pseudogloeosporium* Jacz. 1917 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, ten species, type: *K. latemarensis* Bubák, coelomycetous, pathogen, terrestrial, Armenia, Canada, China, Czechoslovakia, Germany, Italy, Japan, Poland, Russia, Spain, Scotland, Switzerland, Turkey, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), compare with *Leptothyrium*, sequences are unavailable, holotype of type: IMI 213594.

Kabatiella Bubák 1907 (= *Exobasidiopsis* Karak. 1922 *fide* Species Fungorum 2017), *Saccotheciaceae*, *Dothideales*, *Dothideomycetes*, 18 species, type: *K. microsticta* Bubák, hyphomycetous, sexual morph unknown, saprobes, pathogen, terrestrial, worldwide, see Simon et al. (2009; DNA), Seifert et al. (2011; morphology), Bai et al. (2012; pathogens), Bills et al. (2012; new species), Wijayawardene et al.

(2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; genus accepted), cultures and sequences are available.

Kabatina R. Schneid. & Arx 1966, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, five species, type: *K. thujae* R. Schneid. & Arx, hyphomycetous, sexual morph unknown, saprobes, pathogen, terrestrial, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; accepted as a genus in *Dothideaceae*), cultures and sequences are available.

Kacosphaeria Speg. 1887, *Calosphaeriaceae*, *Calosphaeriales*, *Sordariomycetes*, one species, type: *K. antarctica* Speg., asexual morph unknown, from living leaves, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type species unavailable.

Kaernefelia S.Y. Kondr., Elix, A. Thell & Hur 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *K. kaernefeltii* (S.Y. Kondr., Elix & A. Thell) S.Y. Kondr., A. Thell, Elix, J. Kim, A.S. Kondr. & J.-S. Hur, lichenized, Australia, see Kondratyuk et al. (2013a; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available.

Kainomyces Thaxt. 1901, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *K. isomali* Thaxt., asexual morph unknown, biotrophic, terrestrial, Western Africa, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kalaallia Alstrup & D. Hawksw. 1990, *Dacampiaceae*, *Dothideomycetes* families *incertae sedis*, one genus, type: *K. reactiva* Alstrup & D. Hawksw., on lichens, Europe, see Etayo and Sancho (2008; synonym of *Opegrapha*, *Roccellaceae*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, holotype of type: C.

Kalharituber Trappe & Kagan-Zur 2005, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *K. pfeilii* (Henn.) Trappe & Kagan-Zur, mycorrhizal, terrestrial, see Adeleke and Dames (2014; interaction with bacteria), Trappe et al. (2008; desert truffles), cultures and sequences are available, holotype and ex-type of type species unavailable.

Kalamarospora G. Delgado 2010 [2011], *Ascomycota* genera *incertae sedis*, one species, type: *K. multiflagellata* G. Delgado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

cultures and sequences are unavailable, holotype of the type: BPI 879811A.

Kalapuya M.J. Trappe, Trappe & Bonito 2010, *Morchellaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *K. brunnea* M.J. Trappe, Trappe & Bonito, asexual morph unknown, epigeous, saprobes, mycorrhizal, terrestrial, Italy, USA, see Trappe et al. (2010a; DNA, phylogeny), Jaklitsch et al. (2016a; accepted as a genus in *Morchellaceae*), cultures and sequences are available.

Kalbiana Henssen 1988, *Verrucariales* genera *incertae sedis*, *Eurotiomycetes*, one species, type: *K. brasiliensis* Henssen, lichenized, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Kalbographa Lücking 2007, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *K. caracasana* (Müll. Arg.) Lücking, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Kalb et al. (2009; Thailand), Plata et al. (2012a; placed in family), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, holotype of type species unavailable.

Kalchbrenneriella Diederich & M. S. Christ. 2002, *Ascomycota* genera *incertae sedis*, one species, type: *K. cyanescens* (Kalchbr.) Diederich & M.S. Christ., hyphomycetous, sexual morph unknown, lichenicolous, non-lichenized, terrestrial, Europe, North America and Papua New Guinea, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), sequences are unavailable.

Kaleidosporium Van Warmelo & B. Sutton 1981, *Ascomycota* genera *incertae sedis*, one species, type: *K. fenestratum* (Ellis & Everh.) Van Warmelo & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2016b; morphology), cultures and sequences are unavailable.

Kallichroma Kohlm. & Volk.-Kohlm. 1993, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, type: *K. tethys* (Kohlm. & E. Kohlm.) Kohlm. & Volk.-Kohlm., four species, asexual morph unknown, saprobes, marine, tropical regions of the Atlantic, Indian and Pacific Oceans and east coast of Australia, see Dethoup and Manoch (2009; Thailand), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Abdel-Wahab et al. (2016; new species, phylogeny), cultures and sequences are available.

Kallistoskypha Pfister, Agnello, Lantieri & LoBuglio 2013, *Caloscypthaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *K. incarnata* (Duvernoy & Maire) Pfister, Agnello, Lantieri & LoBuglio, on soil associated with

Eucalyptus, Mediterranean region, see Pfister et al. (2013; proposed genera), sequences are unavailable.

Kalmusia Niessl 1872 (= *Dendrothyrium* Verkley et al. 2013; = *Diapleella* Munk 1953 *fide* Wijayawardene et al. 2014c; Species Fungorum 2017), *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 15 species, type: *K. ebuli* Niessl, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genera in *Didymosphaeriaceae*), Ariyawansa et al. (2014e; taxonomy, phylogeny), Wijayawardene et al. (2014c, 2016b; phylogeny, taxonomy, outline, nomenclature), Zhang et al. (2014d; neotyped *Kalmusia ebuli*), Liu et al. (2015a; new species, Italy), Tanaka et al. (2015; phylogeny), cultures and sequences are available, neotype and ex-neotype culture for type: BJFC 200201, CBS 123120.

Kalmusibambusa Phook., Tennakoon, Thambug. & K.D. Hyde 2017, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *K. triseptata* Phook., Tennakoon, & K.D. Hyde, asexual morph unknown, saprobes, Thailand, see Thambugala et al. (2017b; taxonomy, phylogeny), holotype of the type: MFLU 12-2473.

Kamalomyces R.K. Verma, N. Sharma & Soni 2008, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, three species, type: *K. indicus* R.K. Verma, N. Sharma & Soni, asexual morph unknown, saprobes, India, see Boonmee et al. (2014b; taxonomy), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Kamatella Anahosur 1969, *Ascomycota* genera *incertae sedis*, one species, type: *K. indica* Anahosur, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, India, see Wijayawardene et al. (2012, 2016b, 2017; outline, morphology), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of type species unavailable.

Kameshwaramyces Kamal, R.K. Verma & Morgan-Jones 1986, *Ascomycota* genera *incertae sedis*, two species, type: *K. globosus* Kamal, R.K. Verma & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Verma et al. (2008; new species, India), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Kananascus Nag Raj 1984, *Trichosphaeriaceae*, *Trichosphaeriales*, *Sordariomycetes*, three species, type: *K. verrucisporus* Nag Raj, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Kantvilasia P.M. McCarthy, Elix & Sérus. 2000, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *K. hians* P.M. McCarthy, Elix & Sérus., lichenized, Australia, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable, holotype of type species HO 445410.

Kapooria J. Reid & C. Booth 1989, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *K. musarum* (J.N. Kapoor) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type species unknown.

Karschia Körb. 1865, *Asterinales* genera *incertae sedis*, *Dothideomycetes*, four species, type: *K. talcophila* (Ach.) Körb., asexual morph unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Muggia et al. (2013; phylogeny), Thambugala et al. (2014b; transferred to *Lichenotheliaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Lichenotheliaceae*), Ertz and Diederich (2015; phylogeny, accepted as *Asterinales*), cultures and sequences are available.

Karstenella Harmaja 1969, *Karstenellaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *K. vernalis* Harmaja, saprobes, terrestrial, Europe (Finland) and North America (New Mexico), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pfister et al. (2013; genus accepted), cultures and sequences are available, holotype of type species unavailable.

Karstenia Fr. 1885 (= *Chailletia* P. Karst. 1871 *fide* Species Fungorum 2017), *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, ten species, type: *K. sorbina* (P. Karst.) P. Karst., asexual morph unknown, saprobes, Europe, USA, see Lumbsch and Huhndorf (2010; outline), Baral (2015; new combination), sequences are unavailable, holotype of type species unavailable.

Karsteniomyces D. Hawksw. 1980, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *K. peltigerae* (P. Karst.) D. Hawksw., asexual macroconidial morphs of *Scutula* Tul., lichenicolous, widespread, see Hyde et al. (2011; checklist), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of type species H-Karst 2041.

Karstenula Speg. 1879 (= *Staurosphaeria* Rabenh. 1858 *fide* Species Fungorum 2017), *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 16 species, type: *K. rhodostoma* (Alb. & Schwein.) Speg., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch

and Huhndorf (2010; outline), Zhang et al. (2012e; accepted in *Didymosphaeriaceae*), Hyde et al. (2013; accepted in *Didymosphaeriaceae*), Ariyawansa et al. (2014e; taxonomy, accepted in *Didymosphaeriaceae*, phylogeny), Wijayawardene et al. (2014c; outline), Tanaka et al. (2015; DNA, phylogeny), holotype of the type species PH 01048835 (holotype of *Sphaeria rhodostoma*), cultures available for type species CBS 690.94, CBS 691.94.

Kashiwadia S.Y. Kondr., L. Lökö & Hur 2014, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *K. orientalis* (Kashiw.) SY Kondr., L. Lökö et J.-S. Hur, lichenized, eastern Asia, see Lücking et al. (2017a; classification), cultures and sequences are available.

Kastanostachys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *K. aterrima* (Fuckel) L. Lombard & Crous, asexual morph hyphomycetous, from forest soil, terrestrial, Czech Republic, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), epitype and ex-epitype of the type: PRM 934970, CBS 101310.

Kathistes Malloch & M. Blackw. 1990, *Kathistaceae*, *Ophiostomatales*, *Sordariomycetes*, type: *K. calyculata* Malloch & M. Blackw., three species, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences and cultures are available (unpublished), holotype of type species TRTC 51145.

Katumotoa Kaz. Tanaka & Y. Harada 2005, *Lentitheciaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *K. bambusicola* Kaz. Tanaka & Y. Harada, asexual morph unknown, saprobes, terrestrial, Japan, see Tanaka et al. (2009, 2015; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type species HHUF 28661, JCM 13131.

Kazachstania Zubkova 1971, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, 35 species, type: *K. viticola* Zubkova, asexual reproduction is by multilateral budding, saprophytic, but a few species are opportunistic pathogens of mammals, in soil, maize silage, waste water, soft drinks, fermented foods, tree fluxes, faeces, infected mammals, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Keinstirschia J. Reid & C. Booth 1989, *Diaporthales* *incertae sedis*, *Sordariomycetes*, one species, type: *K. megalosperma* (Kirschst.) J. Reid & C. Booth, terrestrial, saprobes, Germany, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type species (see MBT#133497).

Keissleriella Höhn. 1919 (= *Chaetopyrena* Sacc. 1883; = *Chaetopyrenis* Clem. & Shear 1931; = *Coenosphaeria* Munk 1953; = *Zopfinula* Kirschst. 1939 *fide* Species Fungorum 2017), *Lentitheciaeae*, *Pleosporales*, *Dothideomycetes*, c. 34, type: *K. aesculi* (Höhn.) Höhn., asexual morph dendrophoma-like, saprobes, terrestrial, worldwide, Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; accepted as a genus in *Lentitheciaeae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013; outline, accepted as a genus in *Lentitheciaeae*), Crous et al. (2014c; new species, The Netherlands), Ariyawansa et al. (2015a; new species), Liu et al. (2015a; new species, Italy), Singtripop et al. (2015; new species, phylogeny), Tanaka et al. (2015; new species, phylogeny), Tibpromma et al. (2017b; new species), cultures and sequences are available.

Keissleriomycetes D. Hawksw. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *K. sansteadeans* (Keissl.) D. Hawksw., lichenicolous, Europe, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species W 1921/185.

Kellermania Ellis & Everh. 1885 (= *Planstromella* A.W. Ramaley 1993 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b), *Planstromellaceae*, *Botryosphaeriales*, *Dothideomycetes*, c. 20 species, type: *K. yuccigena* Ellis & Everh., coelomycetous, sexual morph *Planstromella*, saprobes, terrestrial, worldwide, see Minnis et al. (2012; taxonomy, phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline, nomenclature), Hyde et al. (2013; taxonomy), Monkai et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), cultures and sequences are available.

Kendrickomyces B. Sutton, V.G. Rao & Mhaskar 1976, *Ascomycota* genera *incertae sedis*, two species, type: *K. indicus* B. Sutton, V.G. Rao & Mhaskar, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of the type species IMI 190465.

Kensinia J. Reid & C. Booth 1989, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *K. umbrina* (Jenkins) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Kephartia R.C. Harris & Lendemer 2013, *Ascomycota* genera *incertae sedis*, two species, type: *K. crystalligera* R.C. Harris & Lendemer, asexual morph unknown, lichenized, terrestrial, USA, see Lendemer et al. (2013;

taxonomy), sequences are unavailable, holotype of type: NY(26883).

Keratosphaera H.B.P. Upadhyay 1964, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, six species, type: *K. batistae* H.B.P. Upadhyay, asexual morph unknown, lichenicolous, terrestrial, worldwide, see Farkas (2010; new species), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Kernia Nieuwl. 1916 (= *Magnusia* Sacc. 1878 *fide* Species Fungorum 2017), *Microascaceae*, *Microascales*, *Sordariomycetes*, five species, type: *K. nitida* (Sacc.) Nieuwl., asexual morph unknown, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype material of type: CBS H-13743.

Kerniomycetes Toro 1939, *Schizophyriaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *K. costi* Toro, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Schizophyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Ketubakia Kamat, Varghese & V.G. Rao 1987, *Ascomycota* genera *incertae sedis*, one species, type: *K. indica* Kamat, Varghese & V.G. Rao, hyphomycetous, sexual morph unknown, saprobes, lignicolous, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Kiehlia Viégas 1944 (= *Lateropeltis* Shanor 1946 *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *K. obscura* Viégas, asexual morph unknown, epiphytes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type species IACM.

Kiliophora Kuthub. & Nawawi 1993 (= *Danaëa* Caneva & Rambelli 1981 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, three species, type: *K. fusispora* Kuthub. & Nawawi, hyphomycetous, sexual morph unknown, saprobes, foliicolous, Ivory Coast, Malaysia and Vietnam, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hidayat et al. (2014; phylogeny, placed in *Amphisphaeriaceae*), Rajeshkumar et al. (2016; family

needs revision), cultures and sequences are available, holotype of type: IMI 355816.

Kimbropezia Korf & W.Y. Zhuang 1991, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *K. campestris* Korf & W.Y. Zhuang, saprobes, terrestrial, Canary Island, cultures and sequences are available, holotype of type: CUP-MM 2762.

Kionocephala P.M. Kirk 1986, *Ascomycota* genera *incertae sedis*, one species, type: *K. catenulata* (M.B. Ellis) P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, lignicolous, UK, Russia, Cuba, see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Kionochaeta P.M. Kirk & B. Sutton 1986 (= *Phalicorona* Subram. 1993; = *Phialocorona* Subram. 1995 [1992–1993] *fide* Species Fungorum 2017), *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, seven species, type: *K. ramifera* (Matsush.) P.M. Kirk & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type species, holotype of the type: IMI188783.

Kirschsteiniothelia D. Hawksw. 1985 (= *Dendryphiopsis* S. Hughes 1953 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b), *Kirschsteiniotheliaceae*, *Kirschsteiniotheliales*, *Dothideomycetes*, 19 species, type: *K. aethiops* (Berk. & Curtis) D. Hawksw., asexual morph formerly known in *Dendryphiopsis*, on wood, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes* genera *incertae sedis*), Boonmee et al. (2012; phylogeny), Hyde et al. (2013; accepted as a genus in *Kirschsteiniotheliaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Rossman et al. (2015b; nomenclature), Jaklitsch et al. (2016a; classification), Su et al. (2016b; new species, phylogeny), cultures and sequences are available, Hernández-Restrepo et al. (2017; new species, phylogeny), holotype of type species K(M), genus remains polyphyletic.

Kirstenboschia Quaedvl., Verkley & Crous 2013, *Ascomycota* genera *incertae sedis*, one species, type: *K. diospyri* Quaedvl., Verkley & Crous, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Quaedvlieg et al. (2013; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type: CBS H-21326, CBS 134911.

Kitesporella Jheng & K. L. Pang 2012, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one

species, type: *K. keelungensis* J.S. Jheng & K.L. Pang, asexual morph unknown, saprobes, marine, Taiwan, see Pang and Jheng (2012a; placement in *Halosphaeriaceae*), Maharachchikumbura et al. (2015b, 2016b; genus accepted), sequence are unavailable, holotype of type: BBH30190.

Klasterskyia Petr. 1940, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, one species, type: *K. acuum* (Mouton) Petr., asexual morph hyalorhinocladiella-like, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Kleidiomyces Thaxt. 1908, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *K. furcillatus* (Thaxt.) Thaxt., asexual morph unknown, biotrophic, terrestrial, America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kluyveromyces van der Walt 1956 (= *Dekkeromyces* Santa María & C. Sánchez-Pinto 1970; = *Dekkeromyces* Wick. & K.A. Burton 1956; = *Fabospora* Kudryavtsev 1960; = *Zygodabospora* Kudryavtsev 1960; = *Zygorenomyspora* Krassiln. 1954 *fide* Species Fungorum 2017), *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, eight species, type: *K. marxianus* (E.C. Hansen) van der Walt, asexual reproduction is by multilateral budding, saprophytic, sea water, *Drosophila* spp., mushrooms, tree fluxes, cacti, dairy products, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Kmetia Bres. & Sacc. 1902, *Ascomycota* genera *incertae sedis*, one species, type: *K. exigua* Bres. & Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kmetiopsis Bat. & Peres 1960, *Ascomycota* genera *incertae sedis*, one species, type: *K. hymenaeae* Bat. & Peres, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Knemiothyrium Bat. & J.L. Bezerra 1960, *Ascomycota* genera *incertae sedis*, one species, type: *K. weldenii* Bat. & J.L. Bezerra, coelomycetous, sexual morph unknown, saprobes, terrestrial, Jamaica, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Knightiella Müll. Arg. 1886, *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type: *K. splachnirruma* Müll. Arg., see Lücking et al. (2017b; classification), sequences are available.

Knox Daviesia M.J. Wingf., P.S. van Wyk & Marasas 1988 (= *Gondwanamyces* G.J. Marais & M.J. Wingf. 1998 *fide* Réblová et al. 2016c), *Microascaceae*, *Microascales*, *Sordariomycetes*, two species, type: *K. proteae* M.J. Wingf., P.S. van Wyk & Marasas, hyphomycetous, sexual morph unknown, coprophilous, terrestrial, South Africa, see Lumbsch and Huhndorf (2010; outline), de Beer et al. (2013; new species), Kirk et al. (2013; genus accepted), Aylward et al. (2014, 2015; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures and sequences are available, holotype of the type: PREM48928, cultures and sequences are available.

Knufia L.J. Hutchison & Unter. 1996, *Trichomeriaceae*, *Chaetothyriales*, *Eurotiomycetes*, 13 species, type: *K. cryptophialidica* L.J. Hutchinson & Unter., hyphomycetous, mostly terrestrial, associated with soil, bark, rock surfaces and human skin, America, Asia and Europe, see Li et al. (2008a; human infection, as “*Coniosporium*”), Tsuneda et al. (2011b; outline), Seifert et al. (2011; morphology), Untereiner et al. (2011; phylogeny, sexual morph), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2013b; new species), Réblová et al. (2013; phylogeny, new combination), holotype and ex-isotype strain of the type: TRTC 51492, DAOM 216555.

Kochiella Sakayaroj, K.L. Pang & E.B.G. Jones 2011, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *K. crispa* (Kohlm.) Sakay., K.L. Pang & E.B.G. Jones, asexual morph unknown, saprobes, marine, Denmark, Hawaii, Japan, Liberia, Martinique, Taiwan, see Sakayaroj et al. (2011a; morphology, phylogeny), Jones et al. (2015; outline, phylogeny), cultures and sequences are available.

Kockiozyma Jindam., Yukphan, & Y. Yamada 2012, *Lipomyctaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *K. suomiensis* (M.T. Sm., Van der Walt & Y. Yamada) Jindam., Yukphan & Y. Yamada, saprophytic, see Jindamorakot et al. (2012; taxonomy).

Kodamaea Y. Yamada, T. Suzuki, Matsuda & Mikata emend. Rosa, Lachance, Starmer, Barker, Bowles & Schlag-Edler 1999, ?*Metschnikowiacaeae*, *Saccharomycetales*, *Saccharomycetes*, seven species, type: *K. ohmeri* (Etchells & T.A. Bell) Y. Yamada, Suzuki, M. Matsuda & Mikata, saprophytic, on flowers of morning glory, (*Hibiscus* spp.) and other plants including cacti, various beetles, on *Drosophila*, pickle brines, see Lumbsch and Huhndorf (2010; outline), Freitas et al.

(2013; DNA), Kirk et al. (2013; genus accepted), sequences are available.

Kodonospora K. Ando 1993, *Ascomycota* genera *incertae sedis*, one species, type: *K. tetracolumnaris* K. Ando, hyphomycetous, sexual morph unknown, saprobes, on leaves, Japan, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), sequences are unavailable, holotype of type: TNS F-176499.

Koerbereria A. Massal. 1854 (= *Tingiopsisidium* Werner 1939 *fide* Species Fungorum 2017), *Koerberiaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *K. biformis* A. Massal., lichenized, Europe, Asia, USA, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Placynthiaceae*), Kirk et al. (2013; genus accepted), Spribile and Muggia (2013, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species unavailable.

Koerberiella Stein 1879 (= *Lecanorella* Frey 1926; = *Perspicinora* Riedl 1990 *fide* Species Fungorum 2017), *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, two species, type: *K. wimmeriana* (Körb.) Stein, lichenized, Europe, North America, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Schmull et al. (2011; phylogeny), Lücking et al. (2016a; classification), cultures and sequences are available, holotype of type species unavailable.

Kohlmeyeriella E. B. G. Jones, R. G. Johnson & S. T. Moss 1983, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, two species, type: *K. tubulata* (Kohlm.) E.B.G. Jones, R.G. Johnson & S.T. Moss, asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; genus accepted), Jones et al. (2015; outline, phylogeny), cultures and sequences are available.

Kohlmeyeriopsis S. Klaubauf, M.H. Lebrun & P.W. Crous 2014, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *K. medullaris* (Kohlm., Volkm.-Kohlm. & O.E. Erikss.) Klaubauf, Lebrun & Crous, holomorph, asexual morph hyphomycetous, saprobes, marine, USA, see Abdel-Wahab et al. (2014; phylogeny), Klaubauf et al. (2014; phylogeny), Jones et al. (2015, outline, phylogeny), Maharachchikumbura et al. (2015b, 2016b; genus accepted), cultures and sequences are available, ex-type culture of type species unavailable, holotype of type species J.K. 5528 = 01315596-99 = 01315600-05, NYBG.

Kohninia Holst-Jensen, Vrålstad & T. Schumach. 2004, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *K. linnaeicola* Holst-Jensen, Vrålstad & T. Schumach., asexual morph unknown, saprobes, terrestrial,

Norway, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; phylogeny), cultures and sequences are available.

Kolletes Kohlm. & Volkm.-Kohlm. 2005, *Ascomycota* genera *incertae sedis*, one species, type: *K. undulatus* Kohlm. & Volkm.-Kohlm., hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; not accepted as valid genus), cultures and cultures and sequences are unavailable, holotype of the type: IMS, Kohlmeyer 5728.

Komagataella Y. Yamada, Matsuda, Maeda & Mikata 1995, *?Pichiaceae, Saccharomycetales, Saccharomycetes*, six species, type: *K. pastoris* (Guilliermond) Y. Yamada, M. Matsuda, K. Maeda & Mikata, asexual reproduction is by multilateral budding, saprophytic, fluxes from various trees and from rotted wood, worldwide, see Kurtzman (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Kompsoscypha Pfister 1989, *Sarcoscyphaceae, Pezizales, Pezizomycetes*, four species, type: *K. chudei* (Pat. ex Le Gal) Pfister, saprobes, on leaves, fruits and twigs, tropical and subtropical, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Konenia Hara 1913, *Sordariomycetes* genera *incertae sedis*, two species, type: *K. bambusae* Hara, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Konradia Racib. 1900 (= *Xylariopsis* F.L. Tai 1934 *fide* Species Fungorum 2017), *Clavicipitaceae, Hypocreales, Sordariomycetes*, two species, type: *K. bambusina* Racib., asexual morph unknown, saprobe, terrestrial, Java, Indonesia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Kontospora A. Roldán, Honrubia & Marvanová 1990, *Ascomycota* genera *incertae sedis*, one species, type: *K. halophila* A. Roldán, Honrubia & Marvanová, hyphomycetous, sexual morph unknown, saprobes, aquatic, Spain, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Koorschalamia Subram. 1953, *Trichosphaeriaceae, Trichosphaeriales, Sordariomycetes*, nine species, type: *K. madreeya* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread (pan-tropical), see Allegrucci et al. (2011; key to species), Hyde et al. (2011; outline), Seifert et al. (2011; morphology),

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable for type species.

Koorschalamella Chona, Munjal & J.N. Kapoor 1958, *Ascomycota* genera *incertae sedis*, one species, type: *K. oryzae* Chona, Munjal & J.N. Kapoor, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Koordersiella Höhn. 1909 (= *Hansfordiellopsis* Deighton 1960; *Ascohansfordiellopsis* D. Hawksw. 1979; = *Phanerococcus* Theiss. & P. Syd. 1918 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, three species, type: *K. javanica* Höhn., asexual morph unknown, lichenicolous, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Koralionastes Kohlm. & Volkm.-Kohlm. 1987, *Koralionastaceae, Koralionastetales, Sordariomycetes*, five species, type: *K. ovalis* Kohlm. & Volkm.-Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Campbell et al. (2009a; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2015b, 2016b; genus accepted), holotype: J.K. 4965a, cultures and sequences are available.

Korfia J. Reid & Cain 1963, *Cenangiaceae, Helotiales, Leotiomycetes*, one species, type: *K. tsugae* (E.K. Cash & R.W. Davidson) J. Reid & Cain, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, holotype of type species unavailable.

Korfiella D.C. Pant & V. P. Tewari 1970, *Sarcosomataceae, Pezizales, Pezizomycetes*, one species, type: *K. karnika* D.C. Pant & V.P. Tewari, saprobes, on conifer needles, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Korfiomyces Iturr. & D. Hawksw. 2004, *Lecanoromycetes* genera *incertae sedis*, one species, type: *K. gelatinosus* Iturr. & D. Hawksw. (as “gelatinosum”), asexual morph unknown, on resin and wood, Venezuela, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, holotype of type: VEN Iturriaga 615.

Korunomyces Hodges & F.A. Ferreira 1981, *Ascomycota* genera *incertae sedis*, three species, type: *K. terminaliae* Hodges & F.A. Ferreira, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, ex-type of type: CBS 224.80.

Kostermansinda Rifai 1968, *Ascomycota* genera *incertae sedis*, three species, type: *K. magna* (Boedijn) Rifai, hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kostermansindiopsis R.F. Castañeda 1986, *Ascomycota* genera *incertae sedis*, one species, type: *K. theobromae* R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kotlabaea Svrcek 1969, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, twelve species, type: *K. deformis* (P. Karst.) Svrcek, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lindemann et al. (2015; phylogeny), cultures and sequences are available.

Kramabeeja G.V. Rao & K.A. Reddy 1981, *Ascomycota* genera *incertae sedis*, one species, type: *K. shrungashakha* G.V. Rao & K.A. Reddy, hyphomycetous, sexual morph unknown, saprobes, on dead rachis of *Areca*, India, see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kramasamuha Subram. & Vittal 1973, *Ascomycota* genera *incertae sedis*, one species, type: *K. sibika* Subram. & Vittal, hyphomycetous, sexual morph unknown, saprobes, on leaves, widespread, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kraurogymnogarpa Udagawa & Uchiy. 1999, *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *K. lenticulispora* Udagawa & Uchiy., asexual morphs unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Sigler et al. (2013; DNA), cultures and sequences are available.

Kravtzevia Schwarzman 1961, *Sordariomycetes* genera *incertae sedis*, three species, type: *K. halimodendri* Schwarzman, asexual morph unknown, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf

(2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Kregervanrija Kurtzman 2006, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, three species, type: *K. fluxuum* (Phaff & E.P. Knapp) Kurtzman, asexual reproduction multilateral budding, saprophytic, on tree fluxes, apples, apple juice, wine and other fermented beverages, pickle brines, Europe and North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Kreiseliella U. Braun 1991, *Ascomycota* genera *incertae sedis*, one species, type: *K. typhae* (Vasyag.) U. Braun, coelomycetous, terrestrial, foliicolous, leaf-spotting, Asia (Kazakhstan), see Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kretzschmaria Fr. 1849 (= *Ustulina* Tul. & C. Tul. 1863; = *Ascostroma* Bonord. 1851; = *Rhopalopsis* Cooke 1883 *fide* Wendt et al. 2017), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, more than 50 species, type: *K. clavus* (Fr.) Sacc., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2013), Daranagama et al. (2017 morphology, taxonomy, key), sequences are available but lacking for the type species.

Kretzschmariella Viégas 1944, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *K. culmorum* (Cooke) Y.-M. Ju & J. D. Rogers, asexual morph present, saprobes, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2013), Daranagama et al. (2017 morphology, taxonomy, key), sequences are unavailable.

Kriegeriella Höhn. 1918, *Dothideomycetes* genera *incertae sedis*, four species, type: *K. mirabilis* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline, placed in *Pleosporaceae*), Wu et al. (2011; classification), Zhang et al. (2012e; placed in *Microthyriaceae*), Hyde et al. (2013; classification), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015c; did not accept as in *Pleosporaceae*), cultures and sequences are unavailable, generic revision is essential

Krishnamyces Hosag. 2003, *Ascomycota* genera *incertae sedis*, one species, type: *K. indica* Hosag., ?hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Krogia Timdal 2002, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *K. coralloides*

Timdal, lichenized, Asia, Africa, see Timdal (2009; new species), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type species O L21909.

Kruphaiomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *K. trochoidei* Thaxt., asexual morph unknown, biotrophic, terrestrial, Fiji, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kuehniella G.F. Orr 1976, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *K. racovitzae* (Lagarde) G.F. Orr, asexual morph unknown, saprobes, terrestrial, Northern America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Kullhemia P. Karst. 1878, *Dothideomycetes* genera *incertae sedis*, three species, type: *K. moriformis* (Ach.) P. Karst., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Kumanasamuha P. Rag. Rao & D. Rao 1964, *Ascomycota* genera *incertae sedis*, four species, type: *K. sundara* P. Rag. Rao & D. Rao, hyphomycetous, sexual morph unknown, saprobes, lignicolous, widespread, see Nagao et al. (2009; new species), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kundalinella Hande & Subhedar 2001, *Ascomycota* genera *incertae sedis*, one species, type: *K. raoella* Hande & Subhedar, asexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Kuraishia Y. Yamada, Maeda & Mikata 1994, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, seven species, type: *K. capsulata* (Wickerham) Y. Yamada, K. Maeda & Mikata, asexual reproduction is by multilateral budding, saprophytic, on insect frass, rotted wood, soil, mushroom, worldwide, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Kurssanovia Kravtzev 1955, *Sordariomycetes* genera *incertae sedis*, three species, type: *K. vassiljevskii* Kravtzev, asexual morph unknown, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not accept), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Kurtzmaniella Lachance & Starmer 2008, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, twelve species, type: *K. cleridarum* Lachance & Starmer, in

asexual reproduction cells produced by asexual reproduction are ovoid to ellipsoid and form by multilateral budding, saprophytic, one species is an opportunistic human pathogen, cactus flowers, mushrooms, tree fluxes, beetles, rotted wood, insect frass, apple cider, rotted fruit, olives, soil, seawater, human faeces, sputum, worldwide, see Lumbsch and Huhndorf (2010; outline), Lachance et al. (2013b; DNA), sequences are available.

Kusanobotrys Henn. 1904 (= *Chaetobotrys* Clem. 1931 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, three species, type: *K. bambusae* Henn., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Kylindria DiCosmo, S.M. Berch & W.B. Kendr. 1983, *Reticulascaceae*, *Glomerellales*, *Sordariomycetes*, nine species, type: *K. triseptata* (Matsush.) DiCosmo, S.M. Berch & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Zhang et al. (2010e; new species, China), Réblová et al. (2011; DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available but lacking for type species.

Kyphomyces I.I. Tav. 1985, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 14 species, type: *K. ansatus* (Thaxt.) I.I. Tav., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Kyphophora B. Sutton 1991, *Ascomycota* genera *incertae sedis*, one species, type: *K. avicenniae* B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Laboulbenia Mont. & C.P. Robin 1853 (= *Ceratiomyces* Thaxt. 1901; = *Eumisomyces* Speg. 1912; = *Laboulbeniella* Speg. 1912; = *Schizolaboulbenia* Middelh. 1957; = *Thaxteria* Giard 1892 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, c. 625 species, type: *L. rougetii* Mont. & C.P. Robin, asexual morph unknown, biotrophic, hosts are diverse, including *Acari*, *Blattodea*, *Coleoptera* (*Carabidae*, *Chrysomelidae*, *Curculionidae*, *Erotylidae*, *Gyrinidae*, *Staphylinidae*, and others), *Diptera*, *Hemiptera*, *Hymenoptera* (*Formicidae*), and *Orthoptera* (*Gryllotalpidae*), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2015b; DNA), Rossi et al. (2016a; morphology), Wang et al. (2016b; morphology), sequences are available but lacking for the type species.

Laboulbeniopsis Thaxt. 1920, *Laboulbeniomycetes* genera *incertae sedis*, one species, type: *L. termitarius* Thaxt., asexual morph unknown, parasitic on termites, terrestrial, North America, Caribbean, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Labrocarpon Pérez-Ortega & Etayo 2010, *Asterinales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *L. canariense* (D. Hawksw.) Etayo & Pérez-Ortega, asexual morph unknown, lichenicolous on *Pertusaria* species, parasymbiont, Canary Islands, Madeira, mainland Spain and Portugal, Italy and Greece, see Pérez-Ortega and Etayo (2010; genus proposed), Ertz and Diederich (2015; phylogeny), cultures and sequences are available, holotype of type: IMI 259313.

Laburnicola Wanas., Camporesi, E.B.G. Jones & K.D. Hyde 2016, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *L. muriformis* Wanas., Camporesi, E.B.G. Jones & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Europe, Russia, see Wanasinghe et al. (2016a; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of type: MFLU 16-0230, MFLUCC 16-0290.

Labyrinthia Malcolm, Elix & Owe-Larsson 1995, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *L. implexa* Malcolm, Elix & Owe-Larss., lichenized, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type species CHR 470232.

Labyrinthomyces Boedijn 1939, *Tuberaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *L. steenisii* Boedijn, truffles, ectomycorrhizal, terrestrial, Australia, see Trappe et al. (2008; desert truffles), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Lacazia Taborda, V.A. Taborda & McGinnis 1999, *Onygenales*, genera *incertae sedis*, *Eurotiomycetes*, two species, type: *L. loboi* Taborda, V.A. Taborda & McGinnis, asexual morph unknown, on humans and animals, terrestrial, neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Lacellina Sacc. 1913, *Ascomycota* genera *incertae sedis*, three species, type: *L. libyca* Sacc. & Trotter, hyphomycetous, sexual morph unknown, on plants, widespread (pantropical), see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lacellinopsis Subram. 1953, *Ascomycota* genera *incertae sedis*, three species, type: *L. sacchari* Subram.,

hyphomycetous, sexual morph unknown, saprobes, on leaves (grasses) and litter, widespread (pantropical), see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lachancea Kurtzman 2003, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, ten species, type: *L. thermotolerans* (Filippov) Kurtzman, asexual reproduction is by multilateral budding, saprophytic, tree fluxes, rotted wood, alpechin, fruit, fruit flies, spoiled beverages, soil, seawater near mangroves, worldwide, see Lumbsch and Huhndorf (2010; outline), sequences are available.

Lachnellula P. Karst. 1884, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, c. 40 species, type: *L. chrysophthalma* (Pers.) P. Karst., asexual morph unknown, saprobes, pathogen, terrestrial, widespread (temperate), Lumbsch and Huhndorf (2010; outline), Whitton et al. (2012b; morphology), Kirk et al. (2013; genus accepted), Schumacher et al. (2015; pathogens), Jaklitsch et al. (2016a; classification), cultures and sequences are available but lacks for type species, needs generic revision.

Lachnodochium Marchal 1895, *Ascomycota* genera *incertae sedis*, one species, type: *L. candidum* Marchal, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lachnum Retz. 1769 (see Species Fungorum 2017 for synonyms), *Lachnaceae*, *Helotiales*, *Leotiomycetes*, c. 50 species, type: *L. agaricinum* Retz., asexual morph unknown, saprobes on ferns, bamboo culms, herbs, herbaceous stems, cones, cupules, wood or leaves of palms, coniferous or broadleaved trees, cosmopolitan, see Nagao (2008; Japan), Hosoya et al. (2010; phylogeny, cultures), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Laciniocladium Petri 1917, *Ascomycota* genera *incertae sedis*, one species, type: *L. castaneae* Petri, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lacrymospora Aptroot 1991, *Requienellaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *L. parasitica* Aptroot, asexual morph unknown, lichenicolous, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable,

holotype of the type: herb. Aptroot, Aptroot and Hensen 12613.

Lacunospora Cailleux 1969, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, two species, type: *L. stercoraria* Cailleux, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Laetinaevia Nannf. 1932 (= *Myridium* Clem. 1909 *fide* Species Fungorum 2017), *Calloriaceae*, *Helotiales*, *Leotiomycetes*, c. 15 species, type: *L. lapponica* (Nannf.) Nannf., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Baral and Rama (2015; DNA), cultures and sequences are available but lacking for type species.

Laeviomyces D. Hawksw. 1981 (= *Spilomium* Nyl. 1858 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, four species, type: *L. pertusariicola* (Nyl.) D. Hawksw., sexual morphs unknown, lichenicolous, terricolous, widespread, see Kirk et al. (2013; genus accepted), Zhurbenko and Otte (2012; genus accepted, illustrations), cultures and sequences are unavailable, holotype of type species H-Nyl 23576.

Lagenulopsis Fitzp. 1942, *Coryneliaceae*, *Coryneliales*, *Eurotiomycetes*, one species, type: *L. bispora* Fitzp., asexual morph unknown, saprobes, terrestrial, Northern America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wood et al. (2016; phylogeny, taxonomy), cultures and sequences are available.

Lahmia Körb. 1861 (= *Parkerella* A. Funk 1976 *fide* Species Fungorum 2017), *Lahmiaceae*, *Lahmiales*, *Leotiomycetes*, one species, type: *L. kunzei* Körb., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, holotype of type species unavailable.

Lahmiomyces Cif. & Tomas. 1953, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *L. piceae* Cif. & Tomas., asexual morph unknown, saprobes, terrestrial, Italy, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification, treated as a genus in *Patellariaceae*), cultures and sequence are unavailable.

Lambdasporium Matsush. 1971, *Ascomycota* genera *incertae sedis*, three species, type: *L. wauense* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lambertella Höhn. 1918 (= *Coprotinia* Whetzel 1944; = *Phaeociboria* Höhn. 1918; = *Phaeodiscus* L.R. Batra 1968 *fide* Species Fungorum 2017), *Rutstroemiaceae*, *Helotiales*, *Leotiomycetes*, c. six species, type: *L. corni-maris* Höhn., asexual morph helicodendron-like, terrestrial, aquatic, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline), Zhao et al. (2013b; Japan), Kirk et al. (2013; genus accepted), Galán et al. (2015; phylogeny), sequence data available for ten species, holotype and ex-type of type species unavailable.

Lambiella Hertel 1984 (= *Mosigia* Fr. 1845 *fide* Species Fungorum 2017), *Xylographaceae*, *Trapeliales*, *Lecanoromycetes*, twelve species, type: *L. psephota* (Tuck.) Hertel, lichenized, see Spribile et al. (2014a; phylogeny), Resl et al. (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Labminonia Sérus. & Diederich 2005, *Ascomycota* genera *incertae sedis*, one species, type: *L. strigulae* (Elenkin & Woron.) Sérus. & Diederich, hyphomycetous, sexual morph unknown, lichenicolous, non-lichenized, on foliicolous lichens, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Lambro Racib. 1900 (= *Stegophora* Syd. & P. Syd. 1916 *fide* Senanayake et al. 2017), *Sydiowilliaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *L. insignis* Racib., asexual morphs unknown, saprobes, terrestrial, Indonesia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Senanayake et al. (2017, placement), cultures and sequence are unavailable.

Lamproconium (Grove) Grove 1937 (= *Melanconium* sect. *Lamproconium* Grove 1918 *fide* Species Fungorum 2017), *Lamproconiaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *L. desmazieri* (Berk. & Broome) Grove, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), Norphanphon et al. (2016; taxonomy, phylogeny), cultures and sequences are available, holotype of type: CBS H-13785.

Lamprospora De Not. 1864 (= *Moravecia* Benkert, Caillet & Moyne 1987; = *Barlaea* Sacc. 1889; = *Barlaeina* Sacc. & P. Syd. 1899; = *Crouania* Fuckel 1870 [1869–70] *fide* Jaklitsch et al. 2016a; Species Fungorum 2017), *Pezizales* genera *incertae sedis*, *Pezizomycetes*, c. 60 species, type: *L. miniata* (P. Crouan & H. Crouan) De Not., parasitic on mosses, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus

accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available, holotype of the type: CBS H-14139.

Lanatosphaera Matzer 1996, *Dothideomycetes* genera *incertae sedis*, two species, type: *L. porinicola* Matzer, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of the type species GZU, Matzer 1422 & Pelzmann.

Lanceispora Nakagiri, Okane, Tad. Ito & Katum. 1997, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *L. amphibia* Nakagiri, Okane, Tad. Ito & Katum., asexual morph unknown, endophytic, saprophytic, subtropics, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Amphisphaeriaceae*), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available, holotype and ex-type of the type: IFO H-12218, NBRC 32918.

Lanspora K.D. Hyde & E.B.G. Jones 1986, *Phomatosporaceae*, *Phomatosporales*, *Sordariomycetes*, one species, type: *L. coronata* K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, marine, Seychelles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2016; taxonomy, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Lanzia Sacc. 1884, *Rutstroemiaceae*, *Helotiales*, *Leotiomycetes*, 55 species, type: *L. flavorufa* (Sacc.) Sacc., asexual morph unknown, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Johnston and Park (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), cultures and sequences are available but lacking for the type species.

Laocoön J.C. David 1997, *Ascomycota* genera *incertae sedis*, one species, type: *L. paradoxus* (Syd. & P. Syd.) J.C. David, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c; outline, accepted as a genus in *Mycosphaerellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline, transferred to *Ascomycota* genera *incertae sedis*), cultures and sequences are unavailable.

Lapidomyces de Hoog & Stielow 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *L. hispanicus* de Hoog & Stielow, hyphomycetous, sexual morph unknown, rock-inhabiting fungus, Spain, see Egidi et al. (2014; taxonomy, phylogeny), Quaedvlieg et al. (2014; phylogeny),

Wijayawardene et al. (2017; outline), type: CBS 118355 = TRN500, cultures and sequences available.

Lappodochium Matsush. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *L. langeniforme* Matsush., hyphomycetous, sexual morph unknown, on soil, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Laquearia Fr. 1849, *Rhytismatales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *L. sphaeralis* (Fr.) Fr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lareunionomyces Crous & M.J. Wingf. 2016, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *L. syzygii* Crous & M.J. Wingf., hyphomycetous, saprobes, terrestrial, France, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type species CBS H-22623, CPC 26531.

Larissia Raiv. 2008, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *L. pyrolae* Raiv., asexual morph unknown, saprobes, terrestrial, Russia, see Raitviiir (2008; taxonomy), the genus was not listed in any recent outlines or checklists, cultures and sequences are unavailable, holotype of type species TAA-182408.

Lasallia Mérat 1821 (= *Macrodictya* A. Massal. 1852; = *Umbilicariomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Umbilicariaceae*, *Umbilicariales*, *Lecanoromycetes*, 17 species, type: *L. pustulata* (L.) Mérat, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Lasiobelonium Ellis & Everh. 1897 (= *Lasiobelonis* Clem. & Shear 1931; = *Sponheimeria* Kirschst. 1941 *fide* Species Fungorum 2017), *Lachnaceae*, *Helotiales*, *Leotiomycetes*, c. 20 species, type: *L. subflavidum* Ellis & Everh., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for type species.

Lasiobertia Sivan. 1978, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *L. africana* Sivan., asexual morph unknown, saprobes, terrestrial, Ghana, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Daranagama et al. (2016d; accepted in *Xylariales*), cultures and sequences are available but lack for the type species, holotype of the type: IMI 38816.

Lasiobolidium Malloch & Cain 1971, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, seven species, type: *L. spirale* Malloch & Cain, asexual morph unknown, saprobes, on dung, North America, see Doveri et al. (2013; key, Italy), Hansen et al. (2013; polyphyletic genus), cultures and sequences are available.

Lasiobolus Sacc. 1884 (= *Peziza* subgen. *Pyronemella* Vido 1879; = *Pyronemella* (Vido) Sacc. 1889 *fide* Species Fungorum 2017), *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, eleven species, type: *L. pilosus* (Fr.) Sacc., coprophilous, sexual morph unknown, saprobes, on herbivore dung, worldwide, see Güngör et al. (2008; proposed family), Chang and Wang (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, holotype of type: CBS H-13792.

Lasiobotrys Kunze 1823, *Venturiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *L. lonicerae* (Fr.) Kunze, asexual morph unknown, saprobes, terrestrial, ulocladium-like, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Hyde et al. (2013) did not accept as a genus in *Venturiaceae*, Zhang et al. (2011d; accepted as a genus in *Venturiales* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lasiodiplodia Ellis & Everh. 1896 (= *Combodia* Fr. 1849; = *Macrophomella* Died. 1916; = *Macrophomopsis* N.E. Stevens & Baechler 1926; = *Nemadiplodia* Zambett. 1955 [1954]; = *Striодiplodia* Zambett. 1955 [1954]; = *Traversoa* Sacc., Syd. & P. Syd. 1913 *fide* Species Fungorum 2017), *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, at least 31 species, type: *L. theobromae* (Pat.) Griffon & Maubl., asexual morph coelomycetous, saprobes, pathogens, terrestrial, worldwide, see Pavlic et al. (2008; new species), Phillips et al. (2008, 2013; taxonomy, phylogeny), Abdollahzadeh et al. (2010; new species), Liu et al. (2012a, b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Yan et al. (2013; species associated with grape wine), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c, 2016b; outline, taxonomy, phylogeny), Linaldeddu et al. (2015; new species), Dissanayake et al. (2016b; taxonomy), Tennakoon et al. (2016c; sexual morph of *L. pseudotheobromae*, phylogeny), Doilom et al. (2017; species in *Tectona*), neotype and ex-neotype of the type: CBS H-21411, CBS 164.96 (*fide* Phillips et al. 2013).

Lasiodiplodiella Zambett 1955, *Ascomycota* genera *incertae sedis*, three species, type: *L. triflorae* (B.B. Higgins) Zambett., coelomycetes, sexual morph unknown, widespread, see Hyde et al. (2011; checklist), Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable, holotype of type species unavailable.

Lasioloma R. Sant. 1952, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, nine species, type: *L. arachnoideum* (Kremp.) R. Sant., lichenized, Asia, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Lasiomollisia Raitv. & Vesterh. 2006, *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *L. phalaridis* Raitv. & Vesterh., asexual morph unknown, saprobes, terrestrial, Sweden, cultures and sequences are unavailable, holotype of the type: TAA (M), Hanson 2003-147.

Lasionectria (Sacc.) Cooke 1884 (= *Nectria* subgen. *Lasionectria* Sacc. 1883 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, twelve species, type: *L. mantuana* (Sacc.) Cooke, asexual morph acremonium-like, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lechat and Fournier (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Lasiospaeria Ces. & De Not. 1863 (= *Hormosperma* Penz. & Sacc. 1897; = *Lasiella* Quél. 1875; = *Lasiosordariella* Chenant. 1919 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, 72 species, type: *L. ovina* (Pers.) Ces. & De Not., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Raja et al. (2011a; DNA), Kirk et al. (2013; genus accepted), Miller et al. (2014; phylogeny), Kruys et al. (2015; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Bell and Mahoney (2016; new species, phylogeny), cultures available for the type species ANM 1295 (Raja et al. 2011a, b), holotype of type: CBS H-13806.

Lasiosphaeriella Sivan. 1975, *Chaetosphaeriales* genera *incertae sedis*, *Sordariomycetes*, twelve species, type: *L. dennisii* Sivan., asexual morph acremonium-like, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline, accepted as *Sordariomycetidae*, genera *incertae sedis*), Huhndorf and Miller (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Lasiospaeriopsis D. Hawksw. & Sivan. 1980, *Nitschiaceae*, *Coronophorales*, *Sordariomycetes*, seven species, type: *L. salisburyi* D. Hawksw. & Sivan, asexual morph unknown, lichenicolous, widespread, terrestrial, see Lumbsch & Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: K(M)-IMI 80468.

Lasiosphaeris Clem. 1909 (= *Lasiadelphia* Réblová & W. Gams 2011 *fide* Maharachchikumbura et al. 2015b, 2016b; Réblová et al. 2016c), *Sordariales* genera *incertae sedis*, *Sordariomycetes*, three species, type: *L. hispida* (Tode) Clem., asexual morph formerly known in *Lasiadelphia*, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miller et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures available for the type: TL6019, SMH 3336 (Miller and Huhndorf 2004).

Lasiostemma Theiss., Syd. & P. Syd. 1917 (= *Chaetyllis* Clem. 1931; = *Epiplaca* Kleb. 1918; = *Lasiostemella* Petr. 1950; = *Neodimerium* Petr. 1950; = *Pseudoperisporium* Clem. & Shear 1931; = *Pseudoperisporium* Toro 1926 *fide* Species Fungorum 2017), *Pseudoperisporiaceae*, *Dothideomycetes*, families *incertae sedis*, five species, type: *L. melioloides* (Berk. & Ravenel) Theiss., Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Mapook et al. (2016b; transferred to *Pseudoperisporiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lasiostictella Sherwood 1986, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *L. bambusae* Sherwood, asexual morph unknown, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lasiothyrium Syd. & P. Syd. 1913, *Ascomycota* genera *incertae sedis*, one species, type: *L. cycloschizon* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lasmenia Speg. 1886, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, five species, type: *L. balansae* Speg., coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, South America, Australia, Hong Kong, Philippines, see Serrato-Diaz et al. (2011; DNA), Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Lasmeniella Petr. & Syd. 1927, *Ascomycota* genera *incertae sedis*, 13 species, type: *L. guaranitica* (Speg.) Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lasseria Dennis 1960, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *L. chrysophthalma* Dennis, asexual morph unknown, saprobes, terrestrial, Argentina, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), living cultures and sequences are unavailable.

Latericonis G.V. Rao, K.A. Reddy & de Hoog 1984, *Ascomycota* genera *incertae sedis*, one species, type: *L. obscura* V. Rao, K.A. Reddy & de Hoog, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lateriramulosa Matsush. 1971, *Ascomycota* genera *incertae sedis*, five species, type: *L. uni-inflata* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Laterispora Uecker, W.A. Ayers & P.B. Adams 1982, *Ascomycota* genera *incertae sedis*, one species, type: *L. brevirama* Uecker, W.A. Ayers & P.B. Adams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lathagrium (Ach.) Gray 1821 (see Species Fungorum 2017 for synonyms), *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, ten species, type: ?*L. nigrescens* (Huds.) Gray, lichenized, North America, Europe, Russia, see Otálora et al. (2013, 2014; phylogeny, review), Lücking et al. (2017a; classification), cultures and sequences are available.

Lathraeodiscus Dissing & Sivertsen 1989, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *L. arcticus* Dissing & Sivertsen, on soil with mosses with *Octospora*, *Lamprospora*, *Geopora*, high arctic, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species C, isotype TRH.

Latorua Crous 2015, *Latoruaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. caligans* (Bat. & H.P. Upadhyay) Crous, hyphomycetous, saprobes, Brazil, see Crous et al. (2015a; taxonomy, phylogeny), Wijayawardene et al. (2017; outline, phylogeny), holotype and ex-type of the type: CBS H-22267, CBS 576.65.

Lauderlindsaya J.C. David & D. Hawksw. 1989 (= *Lenormandia* Delise 1841; = *Normandinomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017),

Verrucariaceae, *Verrucariales*, *Eurotiomycetes*, three species, type: *L. borrerei* (Tul.) J.C. David & D. Hawksw., some species lichenicolous, some species lichenized, Europe, Japan, Papua New Guinea, see Lumbsch and Huhndorf (2010; outline), Muggia et al. (2010a; phylogeny), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, lectotype of type species K-Borrer.

Lauriomycetes R.F. Castañeda 1990, *Lauriomycetaceae*, *Lauriomycetales*, *Leotiomycetes*, five species, type: *L. pulcher* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Crous et al. (2009a; DNA, phylogeny), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Liu et al. (2015a; new species, Thailand, accepted as *Ascomycota incertae sedis*), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Lautitia S. Schatz 1984, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. danica* (Berl.) S. Schatz, asexual morph unknown, parasites, marine, Denmark, France, UK, USA, see Jones et al. (2009b; review), cultures and sequences are unavailable, holotype and ex-type unavailable.

Lautospora K.D. Hyde & E.B.G. Jones 1989, *Lautosporaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *L. gigantea* K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, marine, Australia, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jones et al. (2015; outline, phylogeny), sequences are available, holotype of the type species IMI 327273.

Lawalrea Diederich 1990, *Ascomycota* genera *incertae sedis*, one species, type: *L. lecanorae* Diederich, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: LG (66/14656).

Lazarenkoia Zerova 1938, *Dothideomycetes* genera *incertae sedis*, three species, type: *L. selaginellae* Zerova, asexual morph unknown, saprobes, terrestrial, Siberia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lazarenkoella S.Y. Kondr., Kärnefelt, A. Thell, Elix & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *L. zoroasteriorum* S.Y. Kondr. & M. Haji Moniri ex S.Y. Kondr., Kärnefelt, A. Thell, Elix, J. Kim, A.S. Kondr. & Hur, lichenized, Iran,

Uzbekistan, see Kondratyuk et al. (2015c; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species KW-L.

Lazuardia Rifai 1988, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *L. lobata* (Berk. & M.A. Curtis) Rifai, asexual morph unknown, hypogeous, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for type: AAU 43756.

Lecanactis Körb. [nom. cons.] 1855 (= *Arthoniactis* (Vain.) Clem. 1909; = *Bacidiactis* M. Choisy 1931; = *Lecanactiomycetes* Cif. & Tomas. 1953; = *Lecanactis* subgen. *Arthoniactis* Vain. 1901; = *Pyrenotea* Fr. 1821 *fide* Species Fungorum 2017), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, 25 species, type: *L. abietina* (Ehrh. ex Ach.) Körb., lichenized, worldwide, see Ertz et al. (2015b, c; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Lecania A. Massal. 1853 (= *Adermatis* Clem. 1909; = *Aipospila* Trevis. 1857; = *Bayrhofferia* Trevis. 1857; = *Dimerospora* Th. Fr. 1860; = *Dyslecanis* Clem. 1909; = *Lecaniella* Jatta 1889; = *Lecaniomyces* E.A. Thomas 1939; = *Polyozosia* A. Massal. 1855 *fide* Species Fungorum 2017), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, 50 species, type: *L. fuscella* (Schaer.) A. Massal., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), van den Boom (2016; new species), cultures and sequences are available.

Lecanicillium W. Gams & Zare 2001, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, 26 species, type: *L. lecanii* (Zimm.) Zare & W. Gams, hyphomycetous, entomogenous, fungicolous and in soil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, neotype and ex-neotype of type: IMI 331550, CBS 102067.

Lecanidiella Sherwood 1986, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *L. contortae* Sherwood, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), holotype of the type: CBS, Aptroot 38022, cultures and sequences are unavailable.

Lecaniocola Brain 1923, *Ascomycota* genera *incertae sedis*, one species, type: *L. parasitica* (Linder) Brain, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and

sequences are unavailable, holotype of type species unavailable.

Lecanographa Egea & Torrente 1994, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, 37 species, type: *L. lyncea* (Sm.) Egea & Torrente, sexual morph with rounded to lirelliform ascomata, lichenized, rarely lichenicolous, terrestrial, rather world-wide with a preference for coastal habitats with a Mediterranean or Desert type climate, see Ertz and Tehler (2011; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Lecanora Ach. 1809 (see Species Fungorum 2017 for synonyms), *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, 550 species, type: *L. allophana* Nyl., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Pérez-Ortega et al. (2010; phylogeny), Ranković et al. (2011; Antioxidant, antimicrobial and anticancer activity), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Lecanosticta Syd. 1922 (= *Eruptio* M.E. Barr 1996 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, seven species, type: *L. pini* Syd., coelomycetous, sexual morph formerly named in *Eruptio*, pathogens, see Quaedvlieg et al. (2012; morphology, phylogeny), Wijayawardene et al. (2014c, 2016b; taxonomy, phylogeny), cultures and sequences are available, epitype and ex-epitype of type: CBS H-21113, CBS 133791.

Lecanostictopsis B. Sutton & Crous 1997, *Ascomycota* genera *incertae sedis*, four species, type: *L. kamatii* (Ullasa) B. Sutton & Crous, coelomycetous, sexual morph unknown, saprobes, terrestrial, France, India, South Africa, see Wijayawardene et al. (2012, 2017; outline, morphology), cultures and sequences are unavailable, holotype of the type: IMI 147817.

Lecidea Ach. 1803 (= *Cladopycnidium* H. Magn. 1940; = *Diplotheca* (Zahlbr.) Räsänen 1943; = *Lecidea* sect. *Diplotheca* Zahlbr. 1907; = *Nothoporpidia* Hertel 1984; = *Sporoacania* A. Massal. 1855; = *Stenhammara* A. Massal. ex Zahlbr. 1924; = *Stereonema* Kütz. 1836; = *Zosterodiscus* Hertel 1984 *fide* Species Fungorum 2017), *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, 100 species, type: *L. fuscoatra* (L.) Ach., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Lecidella Körb. 1855 (= *Diplophragmia* Vain. 1934; = *Lecidellomyces* E.A. Thomas 1939; = *Lecideola* A. Massal. 1861 *fide* Species Fungorum 2017), *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, 80 species, type:

L. viridans (Flot.) Körb., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kantvilas and Elix (2013; review), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species unavailable.

Lecideopsella Höhn. 1909 (= *Gymnopeltis* F. Stevens 1924; = *Plectomyriangium* C. Moreau & M. Moreau 1959 *fide* Species Fungorum 2017), *Schizothyriaceae*, *Dothideomycetes*, families *incertae sedis*, c. eleven species, type: *L. gelatinosa* Höhn., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Phookamsak et al. (2016; new species, taxonomy), cultures and sequences are unavailable.

Lecidoma Gotth. Schneid. & Hertel 1981 (= *Lecidea* subdiv. *Lepidoma* Ach. 1803; = *Lepidoma* (Ach.) Gray 1821 *fide* Species Fungorum 2017), *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *L. demissum* (Rutstr.) Gotth. Schneid. & Hertel, lichenized, clearings of alpine grasslands and in places with prolonged snow cover, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Lecidopyrenopsis Vain. 1907, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *L. corticola* Vain., lichenized, Thailand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Leciophysma Th. Fr. 1865, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, three species, type: *L. finmarkicum* Th. Fr., asexual morph unknown, lichenized, terrestrial, Siberia, see Wedin et al. (2009b; DNA), Lumbsch and Huhndorf (2010; outline), Øvstedral and Lewis Smith (2011; new species, Antarctica), Kirk et al. (2013; genus accepted), Ekman et al. (2014; new species), cultures and sequences are available.

Lecophagus M.W. Dick 1999, *Ascomycota* genera *incertae sedis*, two species, type: *L. fasciculatus* M.W. Dick, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Lectera P.F. Cannon 2012, *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*, two species, type: *L. colletotrichoides* (J.E. Chilton) P.F. Cannon, coelomycetous, sexual morph unknown, pathogens,

terrestrial, worldwide, see Cannon et al. (2012; taxonomy, phylogeny), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available, lectotype and cultures of the type species ISC 217496, CBS 109728.

Lecythiomyces Doweld 2013 (= *Lecythium* Zukal 1893 *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, three species, type: *L. aeruginea* (Zukal) Doweld, asexual morph unknown, saprobes, terrestrial, Austria, see Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Lecythothecium Réblová & Winka 2001, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, four species, type: *L. duriligni* Réblová & Winka, asexual morph hyphomycetous (sporidesmium-like), terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and ex-type of the type: PRM 842977, CBS 101317.

Leeina Petr. 1923, *Ascomycota* genera *incertae sedis*, one species, type: *L. philippinensis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leifidium Wedin 1993, *Sphaerophoraceae*, *Lecanorales*, *Lecanomycetes*, one species, type: *L. tenerum* (Laurer) Wedin, asexual morph unknown, lichenized, terrestrial, Australia, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Leightonella Henssen 1965, *Pannariaceae*, *Peltigerales*, *Lecanomycetes*, one species, type: *L. zeylanensis* (Leight.) Henssen, lichenized, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Weerakoon et al. (2016, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Leightonomyces D. Hawksw. & B. Sutton 1977, *Ascomycota* genera *incertae sedis*, two species, type: *L. phillipsii* (Berk. & Leight.) D. Hawksw. & B. Sutton, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, Europe, North America, see Seifert et al. (2011; genus accepted), Wijayawardene et al. (2012, 2017; outline), Pérez-Vargas et al. (2013; morphology, new species), cultures and sequences are unavailable, holotype of the type: TFC Lich 12742.

Leimonia R.C. Harris 2009, *Pilocarpaceae*, *Lecanorales*, *Lecanomycetes*, one species, type: *L. erratica* (Körb.) R.C. Harris & Lendemer, asexual

morph unknown, lichenized, terrestrial, North America, see Harris (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are available.

Leioderma Nyl. 1888, *Pannariaceae*, *Peltigerales*, *Lecanomycetes*, one species, type: *L. pycnophorum* Nyl., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ekman et al. (2014; DNA, phylogeny), Lücking et al. (2017a; classification), sequences and cultures are available.

Leiorreuma Eschw. 1824, *Graphidaceae*, *Ostropales*, *Lecanomycetes*, 15 species, type: *L. hepaticum* Eschw., asexual morph unknown, lichenized, terrestrial, North America, see Lendemer and Knudsen (2008; new species), Moon et al. (2008; new species), Dubey et al. (2010; new species), Joshi et al. (2013a; Taiwan), Kirk et al. (2013; genus accepted), McDonald et al. (2013; DNA), Lücking et al. (2014, 2016a, b; taxonomy), Poengsungnoen et al. (2014; new species), Wang et al. (2015c; China), sequences and cultures are available.

Leiosphaerella Höhn. 1919 (= *Pseudomassariella* Petr. 1955 *fide* Species Fungorum 2017), *Xylariales* genera *incertae sedis*, *Sordariomycetes*, nine species, type: *L. praeclara* (Rehm) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Jaklitsch and Voglmayr (2012; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, isotype of the type: W11860, epitype and ex-epitype of the type species WU 31314, CBS 125586.

Leiothecium Samson & Mouch. 1975, *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, two species, type: *L. ellipsoideum* Samson & Mouch., saprobes, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Houbraken and Samson (2011; DNA), Kirk et al. (2013; genus accepted), Marin-Felix et al. (2014; new species, phylogeny), cultures and sequences are available.

Lemalis Fr. 1825 (= *Catinula* Lév. 1848; = *Patellina* Speg. 1880 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *L. alismatis* (Pers.) Fr., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable, holotype of type species unavailable.

Lembosia Lév. 1845 (= *Heraldoa* Bat. 1959; = *Lembosidium* Speg. 1921; = *Lembosiellina* Bat. & H. Maia 1960; = *Micrographa* Müll. Arg. 1890; = *Micrographomyces* Cif. & Tomas. 1953; = *Morenoella* Speg. 1885 *fide* Species Fungorum 2017), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, 257 species, type: *L. tenella* Lév., asexual morph unknown, epiphytes, biotrophic,

terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in Asterinaceae), Hongsanan et al. (2014c; taxonomy, DNA, keys), Wijayawardene et al. (2014c; outline), Guatimosim et al. (2015; phylogeny, DNA), Tibpromma et al. (2017b, new species), most species do not have molecular data.

Lembosiella Sacc. 1891, *Aulographaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *L. polyspora* (Pat.) Sacc., asexual morph unknown, saprobes, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Microthyriaceae), Hyde et al. (2013; accepted as a genus in Aulographaceae), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; accepted as a genus in Aulographaceae), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lembosina Theiss. 1913, *Aulographaceae*, *Dothideomycetes*, families *incertae sedis*, 21 species, type: *L. aulographoides* (E. Bommer, M. Rousseau & Sacc.) Theiss., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Asterinaceae), Hyde et al. (2013; accepted as a genus in Aulographaceae), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; accepted as a genus in Aulographaceae), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lembosiopeltis Bat. & J.L. Bezerra 1967, *Dothideomycetes* genera *incertae sedis*, two species, type: *L. nectandrae* Bat. & J.L. Bezerra, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Lembosiopsis Theiss. 1918, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *L. andromedae* (Tracy & Earle) Theiss., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Asterinaceae), Hyde et al. (2013; accepted as a genus in Aulographaceae), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; transferred to *Mycosphaerellaceae* from Asterinaceae), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lembuncula Cif. 1954, *Ascomycota* genera *incertae sedis*, one species, type: *L. pothoidei* Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, Santo Domingo, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lemkea Morgan-Jones & R.C. Sinclair 1983, *Ascomycota* genera *incertae sedis*, one species, type:

L. sphaerospora Morgan-Jones & R.C. Sinclair, hyphomycetous, sexual morph unknown, saprobes, lignicolous, North America, South Africa, Spain, Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lemmopsis (Vain.) Zahlbr. 1906 (= *Leptogium* sect. *Lemmopsis* Vain. 1890 *fide Species Fungorum* 2017), *Lichenaceae*, *Lichinales*, *Lichinomycetes*, three species, type: *L. arnoldiana* (Hepp) Zahlbr., asexual morph unknown, lichenized, terrestrial, Europe, see van den Boom and Brand (2008; Netherlands), Lumbsch and Huhndorf (2010; outline), Motiejūnaitė (2012; Lithuania), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Lemonniera De Wild. 1894, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, seven species, type: *L. aquatica* De Wild., hyphomycetous, sexual morph unknown, saprobes, aquatic, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Lempholemma Körb. 1855 (= *Arnoldia* A. Massal. 1856; = *Physma* sect. *Plectopsora* (A. Massal.) Zahlbr. 1906; = *Plectopsora* A. Massal. 1860 [1859-1860]; = *Schizoma* Nyl. ex Cromb. 1894; = *Synalissina* Nyl. 1885 [1885-86] *fide Species Fungorum* 2017), *Lichenaceae*, *Lichinales*, *Lichinomycetes*, 35 species, type: *L. compactum* (Wallr.) Körb., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Prieto et al. (2012; DNA, phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available but lack for the type species.

Lentithecium K.D. Hyde, J. Fourn. & Ying Zhang 2009, *Lentitheciaceae*, *Pleosporales*, *Dothideomycetes*, ten species, type: *L. fluviatile* (Aptroot & Van Ryck.) K.D. Hyde, asexual morph coelomycetous, saprobes, aquatic (freshwater or marine), Europe, Japan, Africa, see, Zhang et al. (2009d; taxonomy, phylogeny), Hyde et al. (2013, 2016; taxonomy, phylogeny), Wijayawardene et al. (2014c, 2017; outline), Su et al. (2016c; new species), Tibpromma et al. (2017b; taxonomy, phylogeny), cultures and sequence are available, holotype of type species GENT (509).

Lentomitella Höhn. 1906 (= *Eutypopsis* P. Karst. 1878; = *Linostomella* Petr. 1925 *fide Species Fungorum* 2017), *Sordariomycetidae*, genera *incertae sedis*, seven species, type: *L. vestita* (Sacc.) Höhn., saprobes, terrestrial, temperate, see Huhndorf et al. (2008; new species, DNA, phylogeny), Marinowitz et al. (2008; new species, South Africa), Raja and Shearer (2009a, b; compare with *Hanliniomyces*), Réblová and Štěpánek (2009;

phylogeny), Lumbsch and Huhndorf (2010; outline), Rébllová et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), sequence data and cultures available but lacking for the type species.

Leohumicola N.L. Nick. 2005, *Leotiomycetes* genera *incertae sedis*, seven species, type: *L. verrucosa* N.L. Nick., Hambl. & Seifert, saprobes, from soil, terrestrial, worldwide, see Nguyen and Seifert (2008; new species, phylogeny), Chen et al. (2009; new records to China), Wijayawardene et al. (2012, 2017; outline), Takashima et al. (2014; symbiotic associations with *Lycopodiaceae* roots), holotype and ex-type of the type: DAOM 226889, DAOM226889.

Leotia Pers. 1797 (= *Cucullaria* Corda 1842; = *Fungodaster* Haller ex Kuntze 1891; = *Hygromitra* Nees 1816 [1816-17]; = *Leotia* Pers. 1801 *fide* Species Fungorum 2017), *Leotiaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *L. lubrica* (Scop.) Pers., asexual morph unknown, lichenized, terrestrial, worldwide, see Palmer et al. (2008; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, type specimen of type species CBS H-13841.

Lepidocollema Vain. 1890 (see Species Fungorum 2017 for synonyms), *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *L. carassense* Vain., lichenized, Brazil, Solomon Islands, see Lumbsch and Huhndorf (2010; outline), Ekman et al. (2014; new species), Lücking et al. (2017a; classification), cultures and sequences are available.

Lepidopterella Shearer & J.L. Crane 1980, *Argynnaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *L. palustris* Shearer & J.L. Crane, asexual morph unknown, saprobes, aquatic, USA, see Shearer et al. (2009; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Argynnaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Peter et al. (2016; DNA, comparison), cultures available for the type species CBS 459.81.

Lepidosphaeria Parg.-Leduc 1970, *Testudinaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *L. nicotiae* Parg.-Leduc, asexual morph unknown, saprobes, aquatic, USA, see Kruys and Wedin (2009; DNA), Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Testudinaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available, type specimen and cultures available for type species CBS H-7282, CBS 559.71 (*fide* Kruys and Wedin 2009).

Lepidotia Boud. 1885, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *L. hispida* (Quél.)

Boud., asexual morph unknown, saprobes, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list the genus), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Lepisticola W. Gams 2009 (= *Harziella* Costantin & Matr. 1899 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *L. capitata* (Costantin & Matr.) W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, Europe, see Gams et al. (2009; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of type species unavailable.

Leprantha Dufour ex Körb. 1855 (see Species Fungorum 2017 for synonyms), *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: need typification, lichenized, see Lücking et al. (2017a; classification), sequences are unavailable.

Lepraria Ach. 1803 (= *Amphiloma* Nyl. 1855; = *Conia* Vent. 1799; = *Epinycitis* Wallr. 1831; = *Pulina* Adans. 1763 *fide* Species Fungorum 2017), *Stereocaulaceae*, *Lecanorales*, *Lecanoromycetes*, 74 species, type: *L. incana* (L.) Ach., sterile, terrestrial (various substrata), cosmopolitan, see Nelsen et al. (2008; phylogeny), Lendemer (2010a, 2011, 2013; new species, DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Flakus and Kukwa (2011; new species), Flakus et al. (2011; new species), Bungartz et al. (2013b; new species), Elix (2013; new species), Lendemer and Hodkinson (2013; DNA, phylogeny), Miądlikowska et al. (2014a, DNA), sequences are available.

Leprieuria Læssøe, J.D. Rogers & Whalley 1989, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *L. bacillum* (Mont.) Læssøe, J.D. Rogers & Whalley, asexual morph geniculosporium-like, saprobes, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stadler et al. (2013), Daranagama et al. (2017 morphology, taxonomy, key), sequences are unavailable.

Leprieurinella Bat. & H. Maia 1961, *Ascomycota* genera *incertae sedis*, one species, type: *L. ingae* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Laprocaulon Nyl. 1879, *Laprocaulaceae*, *Laprocaulales*, *Lecanoromycetes*, 17 species, type: *L. nanum* Nyl., coelomycetous, lichenized, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lendemer and Hodkinson (2013; new species), Lücking et al. (2017a; classification), cultures and sequences are available.

Leprocollema Vain. 1890, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, three species, type: *L. americanum* Vain., lichenized, Europe, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2016a, b; classification), cultures and sequences are unavailable.

Leproplaca (Nyl.) Nyl. 1888 (see Species Fungorum 2017 for synonyms), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, six species, type: *L. xantholyta* (Nyl.) Nyl., lichenized, Europe, see Arup et al. (2013; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Leptascospora Speg. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *L. uredinis* (Racib.) Speg., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsan et al. (2015d; excluded the genus from *Meliolaceae*), cultures and sequences are unavailable, holotype of type species unavailable.

Lepteutypa Petr. 1923, *Amphisphaeriaceae*, *Amphisphaerales*, *Sordariomycetes*, c. 20 species, type: *L. fuckelii* (Nitschke) Petr., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016b; new species, phylogeny), cultures and sequences are unavailable.

Leptochidium M. Choisy 1952, *Massalongiaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *L. albociliatum* (Desm.) M. Choisy, lichenized, Europe, North America, see Lücking et al. (2017a; classification), cultures and sequences are available.

Leptochlamys Died. 1921, *Ascomycota* genera *incertae sedis*, one species, type: *L. scapicola* (P. Karst.) Died., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leptocucurthis Aptroot 1998, *Dacampiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *L. quadrata* Aptroot, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in *Dacampiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of the type species CBS, Aptroot 37234.

Leptodermella Höhn. 1915, *Ascomycota* genera *incertae sedis*, one species, type: *L. incarnata* (Bres.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leptodiscella Papendorf 1969, *Dothideomycetes* genera *incertae sedis*, four species, type: *L. africana* (Papendorf) Papendorf, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, Spain, South Africa, see Madrid et al. (2011b; new species, DNA), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures for the type: CBS 400.65, type specimen of type species CBS H-720.

Leptodontidium de Hoog 1979 (= *Leptodontium* de Hoog 1977 *fide* Species Fungorum 2017), *Leptodontidiaceae*, *Helotiales*, *Leotiomyces*, eleven species, type: *L. elatius* (F. Mangenot) de Hoog, sexual morph unknown, saprobes, terrestrial, Europe, North America, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), cultures and sequences are available, holotype and cultures for the type species CBS H-7283, CBS 329.53 (Sogonov et al. 2008).

Leptogidium Nyl. 1873 (see Species Fungorum 2017 for synonyms), *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, three species, type: *L. dendriscum* (Nyl.) Nyl., lichenized, Europe, North America, see Muggia et al. (2011b; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Leptogium (Ach.) Gray 1821 (see Species Fungorum 2017 for synonyms), *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, 70 species, type: *L. lacerum* (Sw.) Gray, lichenized, North America, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species unavailable.

Leptographium Lagerb. & Melin 1927, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, 68 species, type: *L. lundbergii* Lagerb. & Melin, asexual morph unknown, hyphomycetous, saprobes, worldwide, see Zhou et al. (2008; new species), Seifert et al. (2011; morphology), Duong et al. (2012; new species), Wijayawardene et al. (2012, 2017; outline), Paciura et al. (2010a; new species), de Beer et al. (2013; new species), Kirk et al. (2013; genus accepted), Huang and Chen (2014; new species), Romón et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Yin et al. (2015; new species), cultures and sequences are available.

Leptoguignardia E. Müll. 1955, *Phyllostictaceae*, *Botryosphaerales*, *Dothideomycetes*, three species, type: *L. onobrychidis* E. Müll., asexual morph unknown,

saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Botryosphaeriaceae*), Liu et al. (2012b; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c, 2017; outline, accepted as a genus in *Phyllostictaceae*), cultures and sequences are unavailable, type specimen of type species CBS H-13858.

Leptokalpion Brumm. 1977, *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, two species, type: *L. albicans* Brumm., asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leptomassaria Petr. 1914, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *L. simplex* (Nitschke ex G.H. Otth) Petr., asexual morph unknown, saprobes, Europe and Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Daranagama et al. (2017) morphology, taxonomy, key), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), sequences are unavailable, placement in *Xylariaceae* needs to be confirmed by molecular data.

Leptomelanconium Petr. 1923, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *L. asperulum* (Moesz) Petr., coelomycetous, sexual morph teratosphaeria-like, saprobes, terrestrial, north temperate, see Crous et al. (2009f; transferred *L. australiense* to *Teratosphaeria*), Zhao and Zhao (2012; new species), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2016b; outline, taxonomy, accepted as a genus in *Teratosphaeriaceae*), cultures and sequences are unavailable.

Leptomeliola Höhn. 1919 (= *Phaeophragmeriella* Hansf. 1944; = *Stevensula* Speg. 1921 *fide Species Fungorum* 2017), *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, nine species, type: *L. hyalospora* (Lév.) Höhn., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; symbiosis with bryophytes, DNA), Silvério et al. (2011; taxonomy, South America), Hyde et al. (2013; accepted in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type species.

Leptopeltis Höhn. 1917 (= *Dothithyrella* Höhn. 1918; = *Leptopeltella* Höhn. 1917; = *Leptopeltina* Petr. 1947; = *Leptopeltinella* Petr. 1951; = *Leptopeltopsis* Petr. 1947; = *Moesziella* Petr. 1927; = *Moeszopeltis* Petr. 1947; = *Opegraphellomyces* Cif. & Tomas. 1953 *fide Species Fungorum* 2017), *Leptopeltidaceae*, *Dothideomycetes* genera *incertae sedis*, one species, type: *L. filicina* (Lib.) Höhn., asexual morph leptothyrium-like, saprobes,

terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, accepted as a genus in *Leptopeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, available specimen of type: CBS H-13860.

Leptoperidia Rappaz 1987 (= *Paraeutypa* Subram. & Ananthap. 1988 *fide Species Fungorum* 2017), *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *L. macropunctata* (Rehm) Rappaz, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, available specimen of type: CBS H-6460.

Leptophyllosticta I.E. Brezhnev 1939, *Ascomycota* genera *incertae sedis*, two species, type: *L. lychnitis* I.E. Brezhnev, coelomycetous, sexual morph unknown, saprobes, terrestrial, former USSR, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leptorhaphis Körb. 1855 (= *Campylacia* A. Massal. ex Beltr. 1858; = *Endophis* Norman 1852; = *Leptomychoraphis* Cif. & Tomas. 1953; = *Leptorhaphiomycetes* Cif. & Tomas. 1953; = *Microtheliomyces* Cif. & Tomas. 1953; = *Mycoleptorhaphis* Cif. & Tomas. 1953 *fide Species Fungorum* 2017), *Naetrocymbaceae*, *Dothideomycetes* genera *incertae sedis*, 15 species, type: *L. oxyspora* (Nyl.) Körb., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, accepted in *Naetrocymbaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Leptosacca Syd. 1928, *Sordariomycetes* genera *incertae sedis*, two species, type: *L. lumae* Syd., asexual morph unknown, saprobes, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type species.

Leptosillia Höhn. 1928, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *L. notha* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Leptosphaerella Speg. 1909, *Sordariomycetes* genera *incertae sedis*, seven species, type: *L. fagaricola* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline),

Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Leptosphaeria Ces. & De Not. 1863 (see Species Fungorum 2017 for synonyms), *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 148 species, type: *L. doliolum* (Pers.) Ces. & De Not., asexual morph coelomycetous, saprobes or parasitic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012a, b; taxonomy, phylogeny), Hyde et al. (2013, 2016; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c, 2017; outline), Ariyawansa et al. (2015b; taxonomy, phylogeny), Dayarathne et al. (2015; new species), cultures and sequence are available, available specimen for type: CBS H-13880.

Leptosphaerulina McAlpine 1902 (= *Pseudoplea* Höhn. 1918; = *Scleropella* Höhn. 1918 fide Species Fungorum 2017), *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, 30 species, type: *L. australis* McAlpine, asexual morph hyphomycetous (pithomyces-like), saprobes, terrestrial, worldwide, see de Gruyter et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Crous et al. (2011d; DNA, phylogeny, on *Proteaceae*), Zhang et al. (2012e; DNA, phylogeny), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2013; new species with asexual morphs), Wijayawardene et al. (2014c; outline), Chen et al. (2015b; phylogeny), cultures and sequences are available, available specimen and cultures for type: CBS H-13980, CBS 311.51.

Leptospora Rabenh. 1857, *Dothideomycetes* genera *incertae sedis*, 15 species, type: *L. porphyrogona* (Tode) Rabenh., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, accepted in *Naetrocybaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Hyde et al. (2016; new species), cultures and sequences are available, holotype of type species unavailable.

Leptosporella Penz. & Sacc. 1897, *Chaetosphaerales* genera *incertae sedis*, *Sordariomycetes*, twelve species, type: *L. gregaria* Penz. & Sacc., asexual morph unknown, saprobes, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Huhndorf and Miller (2011; morphology, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Dai et al. (2017; morphology, phylogeny, new species), cultures and sequences are available but lacking for type material.

Leptostromella (Sacc.) Sacc. 1884 (= *Discostromella* Petr. 1924; = *Leptostroma* subgen. *Leptostromella* Sacc. 1882; = *Leptostromella* subgen. *Leptostromella* (Sacc.)

Sacc. 1884 fide Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *L. septorioides* Sacc. & Roum., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leptothryrella Sacc. 1885 (= *Asterinula* Ellis & Everh. 1889 fide Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, ten species, type: *L. mougeotiana* Sacc. & Roum., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species unavailable.

Leptothryrina Höhn. 1915, *Ascomycota* genera *incertae sedis*, one species, type: *L. rubi* (Duby) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Switzerland, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species unavailable.

Leptothyrium Kunze 1823 (= *Manginulopsis* Bat. & Peres 1963; = *Myxodiscus* Höhn. 1906; = *Rhabdothyrella* Höhn. 1917; = *Rhabdothyrium* Höhn. 1915 fide Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *L. lunariae* Kunze, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leptotrichila P. Karst. 1871 (= *Dibelonis* Clem. 1909; = *Dibelonis* Clem. & Shear 1931; = *Ephelina* Sacc. 1889; = *Fabrea* Sacc. 1881; = *Placopezizia* Höhn. = *Sporonema* Desm. 1874 fide Johnston et al. 2014b; Species Fungorum 2017), *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, 15 species, type: *L. radians* (Desm.) P. Karst., asexual morphs formerly named in *Sporonema*, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequence are unavailable.

Leptoxypium Speg. 1918 (= *Astragoxyphium* Bat., Nascim. & Cif. 1963; = *Caldariomyces* Woron. 1926; = *Megaloxypium* Cif., Bat. & Nascim. 1956 fide Species Fungorum 2017), *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *L. graminum* (Pat.) Speg., asexual morph unknown, sooty moulds, terrestrial, cosmopolitan, see Cheewangkoon et al. (2009; DNA, phylogeny), Chomnunti et al. (2011; taxonomy, new species), Crous et al. (2011a; new species), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Yang et al. (2014a; DNA, new species), Hongsan et al. (2016c, evolution), Kajale

et al. (2015; associated with insects), Park et al. (2015; pathogens), cultures and sequences are available.

Letendrea Sacc. 1880, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. eurotioides* Sacc. asexual morph unknown, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2014e; taxonomy, DNA, phylogeny), Hyde et al. (2013; accepted in *Didymosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures available for the type species CBS 212.31 (*fide* Kodsueb et al. 2016).

Letendraeopsis K.F. Rodrigues & Samuels 1994, *Dothideomycetes* genera *incertae sedis*, one species, type: *L. palmarum* K.F. Rodrigues & Samuels, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Tubeufiaceae*), Hyde et al. (2013; not accepted in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; not listed in *Tubeufiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Letharia (Th. Fr.) Zahlbr. 1892 (= *Chlorea* Nyl. 1855; = *Evernia* subdiv. *Letharia* Th. Fr. 1871; = *Nylanderaria* Kuntze 1891; = *Rhytidocaulon* Nyl. ex Elenkin 1916 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *L. vulpina* (L.) Hue, asexual morph unknown, lichenized, terrestrial, worldwide, see McCune and Altermann (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Lethariella (Motyka) Krog 1976 (= *Usnea* subgen. *Lethariella* Motyka 1936 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, ten species, type: *L. intricata* (Moris) Krog, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Letrouitia Hafellner & Bellem. 1982, *Letrouitiaceae*, *Teloschistales*, *Lecanoromycetes*, 16 species, type: *L. domingensis* (Pers.) Hafellner & Bellem., asexual morph unknown, lichenized, terrestrial, worldwide, see Elix and Kondratyuk (2008; new species), Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; DNA, phylogeny), Joshi et al. (2013b; Vietnam), Kirk et al. (2013; genus accepted), cultures available for type species AFTOL-ID 102.

Lettavia D. Hawksw. & R. Sant. 1990, *Fuscideaeaceae*, *Umbilicariales*, *Lecanoromycetes*, three species, type: *L. cladoniicola* D. Hawksw. & R. Sant., asexual morph unknown, on lichens, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures

and sequences not available, holotype of type species B 7700.

Leucangium Quél. 1883, *Morchellaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *L. ophthalmosporum* Quél., asexual morph unknown, saprobes, hypogeous, worldwide, see Lumbsch and Huhndorf (2010; outline), Sbissi et al. (2010; phylogeny, comparison), Trappe et al. (2010b; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Leucoconiella Bat., H. Maia & Peres 1960, *Ascomycota* genera *incertae sedis*, type species *L. paraguayensis* Bat., H. Maia & Peres, asexual morph unknown, saprobes, terrestrial, Paraguay, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leucoconis Theiss. & Syd. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *L. erysiphina* (Syd. & P. Syd.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leucodecton A. Massal. 1860, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, type: *L. punctatum* (Ach.) A. Massal., asexual morph unknown, lichenized, terrestrial, pantropical, see Mangold et al. (2009; key, Australia), Rivas Plata et al. (2010a; key), Sipman et al. (2012; key, Costa Rica), Joshi et al. (2010b, 2012; Korea, India), Lumbsch and Huhndorf (2010; outline), Lücking and Breuss (2012; new combination), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type species.

Leucodermia Kalb 2015, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, ten species, type: *L. leucomelos* (L.) Kalb, lichenized, see Mongkolsuk et al. (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species GB, Arvidsson & Arvidsson 4158 (as *Heterodermia fertilis* Moberg).

Leucodiaporthe Lar. N. Vassiljeva 2007, *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *L. acerina* M.E. Barr & Lar. N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: VLA P-2941.

Leucodochium Syd. & P. Syd. 1917, *Ascomycota* genera *incertae sedis*, type: *L. pipturi* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leucoglossum S. Imai 1942, *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, two species, type: *L. durandii* (Teng) S. Imai, asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline, accepted as *Leotiomycetes*), Arauzo and Iglesias (2014; new species), Fedosova and Kovalenko (2015; DNA, phylogeny, Russia), cultures and sequences are available but lacking for the type species.

Leuconeurospora Malloch & Cain 1970, *Pseudeurotiaceae*, *Leotiomycetes* families *incertae sedis*, three species, type: *L. pulcherrima* (G. Winter) Malloch & Cain, asexual morph unknown, saprobes, terrestrial, Europe, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, type specimen and cultures available for type species CBS H-14009, CBS 343.76.

Leucoscypha Boud. 1885, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, two species, type: *L. leucotricha* (Alb. & Schwein.) Boud., asexual morph unknown, hypogeous, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Leucosphaerina Arx 1987 (= *Leucosphaera* Arx, Mukerji & N. Singh 1978 *fide Species Fungorum* 2017), *Hypocreales* genera *incertae sedis*, *Hypocreales*, *Sordariomycetes*, two species, type: *L. indica* (Arx, Mukerji & N. Singh) Arx, asexual morph hyphomycetous, terrestrial, saprobes, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Bionectriaceae*), Giraldo et al. (2012; phylogeny), cultures and sequences are available, cultures available for type: CBS 123.78.

Leucothecium Arx & Samson 1973, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, three species, type: *L. emdenii* Arx & Samson, asexual morph unknown, saprobes, on marine and river sediments, Spain, see Gueidan et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline, accepted in *Onygenaceae*), Kirk et al. (2013; genus accepted), Giraldo et al. (2014a; DNA, phylogeny), cultures and sequences are available, available specimen and cultures for the type: CBS H-7299 CBS 576.73.

Leucovibrissa (A. Sánchez) Korf 1990 (= *Vibrissa* sect. *Leucovibrissa* A. Sánchez 1966 *fide Species Fungorum* 2017), *Vibrissaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *L. obconica* (Kanouse) Korf, asexual morph unknown, from fine roots of *Pinus*, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leuiisinea Matsush. 1985, [as '*Leuiisinea*'], *Ascomycota* genera *incertae sedis*, two species, type: *L. bambusicola* Matsush., hyphomycetous, sexual morph unknown, saprobes, lignicolous, Australia, Peru and Taiwan, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Leveillella Theiss. & Syd. 1915 (= *Parasterinella* Speg. 1924 *fide Species Fungorum* 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *L. drimydis* (Lév.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, Chile, see Hofmann (2009; review, accepted in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, excluded from *Asterinaceae*), Wijayawardene et al. (2014c; outline, accepted in *Dothideomycetes*), cultures and sequences are unavailable.

Leveillina Theiss. & Syd. 1915, *Dothideomycetes* genera *incertae sedis*, two species, type: *L. arduinae* (Kalchbr. & Cooke) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, Africa, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Leveillula G. Arnaud 1921, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, 40 species, type: *L. taurica* (Lév.) G. Arnaud (≡ *Erysiphe taurica* Lév.), hyphomycetous asexual morphs, obligate plant pathogens, terrestrial, worldwide, see Khodaparast et al. (2012, 2016; phylogeny), Braun & Cook (2012; taxonomy), sequences are available, lectotype of type: G 00298357.

Libartania Nag Raj 1979, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *L. laserpitii* (Bres.) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Liberomyces Pažoutová, M. Kolařík & Kubátová 2012, *Sordariomycetes* genera *incertae sedis*, two species, type: *L. salicophilus* Pažoutová, M. Kolařík & Kubátová, coelomycetous, sexual morph unknown, endophytes, terrestrial, Czech Republic, see Pažoutová et al. (2012; morphology, DNA), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures and sequences are available, holotype and ex-type of the type species PRM 921954, CCF 4020.

Libertasomyces Crous & Roets 2016, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *L. myopori* Crous & Roets, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, New Zealand, see Crous et al. (2016a, b; new species,

phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22617, CPC 27354.

Libertiella Speg. & Roum. 1880 (= *Nicholsoniella* Kuntze 1891; = *Shecutia* Nieuwl. 1916 *fide* Species Fungorum 2017), Ascomycota genera incertae sedis, six species, type: *L. malmedyensis* Speg. & Roum., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Motiejūnaitė et al. (2011; record from Lithuania), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), current generic concept is uncertain, cultures and sequences are unavailable.

Lichenobactrium Diederich & Etayo 1995, Ascomycota genera incertae sedis, one species, type: *L. pertusariae* Diederich & Etayo, hyphomycetous, sexual morph unknown, lichenicolous, non-lichenized, terrestrial, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type species LG, Diederich 9219.

Lichenobarya Etayo, Diederich & Lawrey 2015, Hypocreaceae, Hypocreales, Sordariomycetes, one species, type: *L. usneae* (Etayo) Etayo, Diederich & Lawrey, asexual morphs unknown, lichenicolous, terrestrial, Europe, see Lawrey et al. (2015; taxonomy, phylogeny), sequences are available.

Lichenochora Hafellner 1989 (= *Laestadia* [unranked] *Paralaestadia* Sacc. & D. Sacc. 1905; = *Paralaestadia* (Sacc. & D. Sacc.) Vain. 1921 *fide* Species Fungorum 2017), Phyllachoraceae, Phyllachorales, Sordariomycetes, five species, type: *L. thallina* (Cooke) Hafellner, asexual morph unknown, lichenicolous, terrestrial, worldwide, see Etayo and Navarro-Rosines (2008; new species), Zhurbenko (2008; new species, Italy), Halici et al. (2009b; new species), Lumbsch and Huhndorf (2010; outline), Etayo (2011; new species), Hafellner and Zimmermann (2012; new species), Etayo et al. (2013b; new species), Kirk et al. (2013; genus accepted), Zhurbenko (2013; new species), Lendemer et al. (2016; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: K.

Lichenoconium Petr. & Syd. 1927, Lichenoconiaceae, Lichenoconiales, Dothideomycetes, five species, type: *L. lichenicola* (P. Karst.) Petr. & Syd., coelomycetous, sexual morph unknown, lichenicolous, terrestrial, worldwide, see Lawrey et al. (2011; new species), Wijayawardene et al. (2012, 2014c, 2016b; outline, taxonomy), Hyde et al. (2013; taxonomy), cultures and sequences are available but lacking for the type species.

Lichenodiplis Dyko & D. Hawksw. 1979, Chaetothyriales genera incertae sedis, Eurotiomycetes, 13 species, type: *L. lecanorae* (Vouaux) Dyko & D. Hawksw.,

coelomycetous, sexual morph unknown, lichenicolous, terrestrial, worldwide, see Atienza et al. (2009; morphology, new species), Knudsen and Kocourkova (2009; morphology, new species), Kirk et al. (2013; genus accepted), Pérez-Vargas et al. (2013; new species), Muggia et al. (2015b; phylogeny), Wijayawardene et al. (2016b; outline, morphology, phylogeny), cultures and sequences are available, lectotype of type species ANGUC.

Lichenohendersonia Calatayud & Etayo 2001, Ascomycota genera incertae sedis, three species, type: *L. squamarinae* Calat. & Etayo, sexual morph unknown, parasymbiont on different lichens, terrestrial, Northern Hemisphere, see Wijayawardene et al. (2016b; morphology), sequence data and cultures unavailable, holotype of the type: IMI 363556.

Lichenopeltella Höhn. 1919 (= *Didymopyrostroma* Bat. & Cavalc. 1964 [1962]; = *Micropeltopsis* Vain. 1921; = *Microthyris* Clem. 1931; = *Trichothyrina* (Petr.) Petr. 1950; = *Trichothryium* subgen. *Trichothyrina* Petr. 1940 *fide* Species Fungorum 2017), Trichothryiaceae, Dothideomycetes, families incertae sedis, c. 31 species, type: *L. maculans* (Zopf) Höhn., asexual morph unknown, saprobes, lichenicolous, terrestrial, worldwide, see Pérez-Ortega and Spribille (2009; new species), Etayo (2010a, b; new species), Lumbsch and Huhndorf (2010; outline, accepted as a genus in Asterinaceae), Zhurbenko (2010a; new species), Brackel (2010b, 2011b; new species), Flakus and Kukwa (2012; new records in Bolivia), van den Boom and Giralt (2012; records in Portugal), Hyde et al. (2013; accepted as a genus in Aulographaceae), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014a; transferred to Mycosphaerellaceae from Asterinaceae), Wijayawardene et al. (2014c; outline), Zhurbenko and Pino-Bodas (2015; new species, New Zealand), cultures and sequences are unavailable.

Lichenopeziza Zukal 1884, Ascomycota genera incertae sedis, one species, type: *L. bryophile* Zukal, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), current generic concept is uncertain, cultures and sequences are unavailable.

Lichenopuccinia D. Hawksw. & Hafellner 1984, Ascomycota genera incertae sedis, one species, type: *L. poeltii* D. Hawksw. & Hafellner, sexual morphs unknown, terrestrial, lichenicolous, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species K(M)-IMI 281393.

Lichenopyrenis Calat., M.J. Sanz & Aptroot 2001, Pleomassariaceae, Pleosporales, Dothideomycetes, one species, type: *L. galligena* Calat., M.J. Sanz & Aptroot, asexual morph unknown, lichenicolous, terrestrial, Spain,

Canada, Japan, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; accepted as a genus in *Pleomassariaceae*), Hyde et al. (2013; accepted as a genus in *Pleomassariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of the type species MA (Lich.) 12715.

Lichenostella Etayo & Calat. 1999, *Ascomycota* genera *incertae sedis*, one species, type: *L. hypotrachynae* Etayo & Calat., hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lichenosticta Zopf 1898, *Lecanorales* genera *incertae sedis*, one species, type: *L. podeticola* Zopf, coelomycetous, sexual morph unknown, lichenicolous, terrestrial, north temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Pino-Bodas et al. (2017; new species, phylogeny), current generic concept is uncertain, cultures and sequences are unavailable.

Lichenostigma Hafellner 1983 (= *Diederimyces* Etayo 1995; = *Phaeosporobolus* D. Hawksw. & Hafellner 1986 *fide* Ertz et al. 2014b; Species Fungorum 2017), *Phaeococcomyctaceae*, *Lichenostigmatales*, *Arthoniomycetes*, type: *L. maureri* Hafellner, asexual morph hyphomycetous, lichenicolous, terrestrial, worldwide, see Pérez-Ortega and Calatayud (2009; new species, Spain), Fernández-Brime et al. (2010; new species), Knudsen and Kocourkova (2010a, b; new species), Lumbsch and Huhndorf (2010; outline), Valadbeigi and Brackel (2011; new species), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2013b; new species), Ertz et al. (2014b; new species; phylogeny), cultures and sequences are available, current generic concept is uncertain, cultures available for type species Diederich 1730, Diederich 17337, Diederich 17326, Diederich 17337.

Lichenothelia D. Hawksw. 1981 (= *Anzia* Garov. 1868 *fide* Species Fungorum 2017), *Lichenotheliaceae*, *Lichenotheliales*, *Dothideomycetes*, 29 species, type: *L. scopolaria* (Nyl.) D. Hawksw., asexual morphs pycnidial or hyphomycetous, on rock or lichenicolous, terrestrial, widespread, some species may belong in *Lichenostigma* Hafellner, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; genus accepted), Muggia et al. (2015a; phylogeny), Jaklitsch et al. (2016a; position), cultures and sequences are available, isotypes of type species BM and K(M) [Anzi, Lich. Langob. 35].

Lichenoverruculina Etayo 2011, *Sordariomycetes* genera *incertae sedis*, one species, type: *L. sigmatospora* (Speg.) Etayo & Sharuddin, asexual morph unknown,

lichenicolous on *Heterodermia*, terrestrial, South America, see Etayo and Sharuddin (2011; genus proposed), cultures and sequences are unavailable, holotype of the type species J. Puiggari 124 (LPS-1715).

Lichina C. Agardh [nom. cons.] 1817 (= *Lichinomyces* E.A. Thomas ex Cif. & Tomas. 1939; = *Neolichina* Gyeln. 1940; = *Pygmaea* Stackh. 1809; = *Thamnidium* Tuck. ex Schwend. 1869 *fide* Species Fungorum 2017), *Lichenaceae*, *Lichinales*, *Lichinomycetes*, nine species, type: *L. pygmaea* (Lightf.) C. Agardh, lichenized, marine, west-European coast, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Lichinella Nyl. 1873 (= *Gonohymenia* J. Steiner 1902; = *Rechingeria* Servít 1931 *fide* Species Fungorum 2017), *Lichenaceae*, *Lichinales*, *Lichinomycetes*, 30 species, type: *L. stipatula* Nyl., lichenized, North America, Europe, Asia, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Lichinodium Nyl. 1875, *Lichenaceae*, *Lichinales*, *Lichinomycetes*, four species, type: *L. sirosiphoidicum* Nyl., lichenized, North America, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Licopolia Sacc., Syd. & P. Syd. 1900, *Dothideomycetes* genera *incertae sedis*, two species, type: *L. franciscana* Sacc., Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Brazil, Kenya, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lignincola Höhnk 1955, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, four species, type: *L. laevis* Höhnk, asexual morph unknown, saprobes, marine, worldwide, see Jones et al. (2009b, 2015; classification), Alias et al. (2010; diversity), Abdel-Wahab (2011a; diversity), Liu et al. (2011a; morphology), Pang et al. (2011b, 2013b; monograph, phylogeny), Sakayaroj et al. (2011a, b; phylogeny, diversity), Lee et al. (2012; checklist), Pang and Jheng (2012a; checklist), Alsheikh-Hussain et al. (2014, review), Overy et al. (2014; natural product), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, neotype of type species J.K. 4459a.

Ligninsphaeria J.F. Zhang, J.K. Liu, K.D. Hyde & Z.Y. Liu 2016, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *L. jonesii* J.F. Zhang, J.K. Liu, K.D. Hyde & Z.Y. Liu, saprobes, Thailand, see Zhang et al. (2016b; taxonomy, phylogeny), holotype: MFLU 15-2718, cultures and sequences are available.

Lignoscripta B.D. Ryan 2004, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, one species, type: *L. atroalba* B.D. Ryan & T.H. Nash, lichenized, Pacific Northwest, see Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; taxonomy), Lücking et al. (2017a, b; classification), cultures and sequences are unavailable, holotype of type species ASU (9092).

Lignosphaeria Boonmee, Thambug. & K.D. Hyde 2015, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *L. thailandica* Boonmee, Thambug. & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Thailand, see Thambugala et al. (2015b; morphology, phylogeny), holotype and ex-type of type: MFLU 11–0135, MFLUCC 11–0376.

Lilicoa Sherwood 1977, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *L. palicoureae* (Seaver & Whetzel) Sherwood, saprobes, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Limaciniaseta D.R. Reynolds 1988, *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *L. californica* D.R. Reynolds, asexual morph unknown, saprobes, terrestrial, USA, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Limaciopsis J.M. Mend. 1925, *Dothideomycetes* genera *incertae sedis*, type: *L. rollandiae* J.M. Mend., data inadequate, see Kirk et al. (2008; treated as nom. dub.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Limacinula Höhn. 1907 (= *Deslandesia* Bat. 1962; = *Limacinia* subgen. *Limacinula* Sacc. & D. Sacc. 1905; = *Limacinula* (Sacc. & D. Sacc.) Höhn. 1909; = *Naetrocymbe* Bat. & Cif. 1963; = *Shanoriella* Bat. & Cif. 1962; = *Teichospora* subgen. *Teichosporina* G. Arnaud 1911; = *Teichosporina* (G. Arnaud) Cif. & Bat. 1962 *fide* Species Fungorum 2017), *Coccodiniaceae*, *Chaetothyriales*, *Eurotiomycetes*, c. 16 species, type: *L. samoensis* Höhn., aerial surface of living plants, epifoliar fungus, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Limacospora Jørg. Koch & E.B.G. Jones 1995, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *L. sundica* (Jørg. Koch & E.B.G. Jones) Jørg. Koch & E.B.G. Jones, asexual morph unknown, saprobes, marine, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Limboria Ach. 1815, *Pezizomycetes* genera incertae sedis, type: *L. constellata* Ach., see Lücking et al. (2017b; nomenclature, typification), sequences are unavailable.

Limnaiomyces Thaxt. 1900, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *L. tropisterni* Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Limongia Jindam., Am-in, Yukphan & Y. Yamada 2011, *Lipomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *L. smithiae* (Van der Walt, M.J. Wingf. & Y. Yamada) Jindam., Am-in, Yukphan & Y. Yamada, saprophytic, see Jindamorakot et al. (2011; taxonomy), cultures and sequences are unavailable.

Lindauella Rehm 1900, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *L. pyrenocarpoidea* Rehm, asexual morph unknown, from living leaves, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Lindgomyces K. Hiray., Kaz. Tanaka & Shearer 2010, *Lindgomycetaceae*, *Pleosporales*, *Dothideomycetes*, 13 species, type species: *L. ingoldianus* (Shearer & K.D. Hyde) K. Hiray., Kaz. Tanaka & Shearer, asexual morph unknown, saprobes, aquatic, China, Japan, USA, see Hirayama et al. (2010; morphology, phylogeny), Raja et al. (2011b, 2013a; morphology, phylogeny), Zhang et al. (2014e; morphology, phylogeny), Crous et al. (2015e; morphology, phylogeny), Li et al. (2016a; morphology, phylogeny), Raja et al. (2017; taxonomy), cultures and sequences are available, holotype and ex-type of the type species ILLS 52289, ATCC 200398.

Lindquistomyces Aramb., E. Müll. & Gamundí 1982, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type: *L. antarcticus* (Speg.) Aramb., E. Müll. & Gamundí (Art. 33.3; Invalid), asexual morph unknown, saprobes, terrestrial, Argentina, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lindra I. M. Wilson 1956, *Lulworthiaceae*, *Lulorthiales*, *Sordariomycetes*, five species, type: *L. inflata* I.M. Wilson, asexual morph hyphomycetous, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2016b; outline), sequences are available, holotype of type: IMI 62909, needs generic revision as it is polyphyletic, type species needs to be sequenced.

Lineolata Kohlm. & Volk.-Kohlm. 1990, *Dothideomycetes* genera *incertae sedis*, one species, type:

L. rhizophorae (Kohlm. & E. Kohlm.) Kohlm. & Volkmar-Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Suetrong et al. (2009; phylogeny, morphology), Jones et al. (2009b, 2015; classification), Alias et al. (2010; diversity), Sakayaroj et al. (2011b; diversity), Zhang et al. (2012e; phylogeny, morphology), Abdel-Wahab et al. (2014; diversity), Wijayawardene et al. (2014c; outline), holotype of type: J.K. 2390a.

Lineostroma H.J. Swart 1988, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. banksiae* (Cooke) H.J. Swart, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Jayasiri et al. (2016a; classification), cultures and sequences are unavailable, available specimen for type: CBS H-5512.

Linkosia A. Hern.-Gut. & B. Sutton 1997, *Ascomycota* genera *incertae sedis*, ten species, type: *L. coccothrinacis* (A. Hern. Gut. & J. Mena) A. Hern. Gut. & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Ma et al. (2011a; morphology), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Almeida et al. (2014c; morphology, new species), Delgado (2014; morphology), sequence data available, holotype of the type species HAC, HACCM 206.

Linocarpon Syd. & P. Syd. 1917 (= *Micronectriopsis* Höhn. 1918 *fide Species Fungorum* 2017), *Xylariales* genera *incertae sedis*, Sordariomycetes, 42 species, type: *L. pandani* Syd. & P. Syd., asexual morph phialophora-like, saprobes, pathogen, endophytes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures available for type: HKUM16278, HKUM16279.

Linochora Höhn. 1910 (= *Xanthopsora* Speg. 1922 *fide Species Fungorum* 2017), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, five species, type: *L. leptospermi* Höhn., coelomycetous, sexual morph unknown, from living leaves, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type species unknown.

Linochorella Syd. & P. Syd. 1912, *Ascomycota* genera *incertae sedis*, one species, type: *L. striiformis* Syd. & P. Syd., coelomycetous, sexual morph unknown, lichenicolous, terrestrial, South Africa, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Linodochium Höhn. 1909, *Ascomycota* genera *incertae sedis*, two species, type: *L. hyalinum* (Lib.) Höhn., hyphomycetous, sexual morph unknown, saprobes,

terrestrial, Europe, North America, see Seifert et al. (2011; morphology, accepted as a genus in *Helotiaceae*), Wijayawardene et al. (2017; outline), Kirk et al. (2013; did not list), cultures and sequences are unavailable, .

Linopeltis I. Hino & Katum. 1961 (= *Linopeltis* I. Hino & Katum. 1960 *fide Species Fungorum* 2017), *Schizothyriaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *L. ryukyuensis* I. Hino & Katum., asexual morph unknown, saprobes, terrestrial, Hawaii, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013, accepted as a genus in *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lipomyces Lodder & Kreger-van Rij 1952 (= *Myxozyma* van der Walt, Weijman & von Arx 1981; see *Species Fungorum* 2017 for additional synonyms), *Lipomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, 20 species, type: *L. starkeyi* Lodder & Kreger, asexual reproduction is by multilateral budding, saprophytic, predominantly from soil, but not uncommon in tree frass from bark beetles, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Yamazaki and Kawasaki (2014; new species, DNA), sequences are available.

Lirellodisca Aptroot 1998, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *L. pyrenulispora* Aptroot, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), cultures and sequences are unavailable, holotype of the type species CBS, Aptroot 38022.

Lirula Darker 1967, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, three species, type: *L. nervisequa* (DC.) Darker, asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Lantz et al. (2011; DNA, phylogeny), Fan et al. (2012b; new species, China, DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Listeromyces Penz. & Sacc. 1901 (= *Cylomyces* Clem. 1931 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, one species, type: *L. insignis* Penz. & Sacc., hyphomycetous, sexual morph unknown, saprobes, lignicolous, Asia, Oceania, see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lithoglypha Brusse 1988, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, one species, type:

L. aggregata Brusse, lichenized, South Africa, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Lithographia Nyl. 1857 (= *Haplographa* Anzi 1860; = *Lithographomyces* Cif. & Tomas. 1953; = *Mycoplacographa* Reinke 1895; = *Placographa* Th. Fr. 1860 *fide* Species Fungorum 2017), *Xylographaceae*, *Trapeliales*, *Lecanoromycetes*, ten species, type: *L. petraea* (Nyl.) Nyl., lichenized, North and South America, Australia, see Resl et al. (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Lithogyalideopsis Lücking, Sérus. & Vězda 2005, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *L. poeltii* (Vězda) Lücking, Sérus. & Vězda, lichenized, see Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Lithophila Selbmann & Isola. For diagnosis see Fungal Diversity 76: 88 (2016). *Trichomeriaceae*, *Chaetothyriales*, *Eurotiomycetes*, rock-inhabiting fungus, monospecific; sexual morph unknown; type: *L. guttulata* Selbmann & Isola. For diagnosis see Fungal Diversity 76: 90 (2016). Holotype: CBS 139723. From marble stone, Cortile della Pigna (Vatican City State); =CCFEE 5907 (NCBI accession numbers: KP791773 (ITS), KR781061 (LSU)), taxonomic species delimitation supported by molecular data. *Lithophila* Selbmann & Isola (2016), introduced as a genus of *Trichomeriaceae*, is a later homonym of *Lithophila* Sw., Prodri, 14, 1788, a name previously and validly published for a genus of *Amaranthaceae*.

Notes: *Lithophila* Selbmann & Isola is illegitimate and unavailable for use [ICN (Melbourne) Art. 53.1]; it is here renamed ***Lithohyppha*** Selbmann & Isola **nom. nov.**, MB819853. For diagnosis see Fungal Diversity 76: 88 (2016). The type species is here renamed ***Lithohyppha guttulata*** (Selbmann & Isola) Selbmann & Isola **comb. nov.**, MB819854 basionym: *Lithophila guttulata* Selbmann & Isola, Fungal Diversity 76: 90 (2015).

Lithopythium Bornet & Flahault 1891, *Ascomycota* genera *incertae sedis*, three species, type: *L. gangliiforme* Bornet & Flahault, asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lithothelium Müll. Arg. 1885 (= *Plagiocarpa* R.C. Harris 1973; = *Pleurotrema* Müll. Arg. 1885; = *Pleurotrematomyces* Cif. & Tomas. 1957; = *Trichotrema* Clem. 1909 *fide* Species Fungorum 2017), *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, eight species, type: *L. cubanum* Müll. Arg., asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Valadbeigi and Sipman (2010;

Iran), Weerakoon et al. (2012a; DNA, phylogeny), Kirk et al. (2013; genus accepted), Gueidan et al. (2016; phylogeny), cultures and sequences are available but lacking for the type species.

Livia Velen. 1947, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *L. aurea* Velen., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Lizonia (Ces. & De Not.) De Not. 1863 (= *Cucurbitaria* subgen. *Lizonia* Ces. & De Not. 1863; = *Pseudolizonia* Pirotta 1889 *fide* Species Fungorum 2017), *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *L. empirigonia* (Auersw.) De Not., asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; DNA), Hyde et al. (2013; accepted in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, type specimen of type species CBS H-14093.

Limonaea Egea & Torrente 1991, *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, four species, type: *L. occulta* Egea & Torrente, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Arthoniales*), Ertz and Tehler (2011; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available, holotype of type: MUB 13619.

Llimoniella Hafellner & Nav.-Ros. 1993, *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *L. scabridula* (Müll. Arg.) Nav.-Ros. & Hafellner, asexual morph unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Suija et al. (2015b; phylogeny), Pärte et al. (2017; DNA, phylogeny), cultures and sequences are available.

Lobaria (Schreb.) Hoffm. 1796 (= *Lichen* sect. *Lobaria* Schreb. 1791; = *Lobariomyces* E.A. Thomas 1939; = *Pulmonaria* Hoffm. 1789 [1790]; = *Reticularia* Baumg. 1790; = *Ricasoliomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Stictomyces* E.A. Thomas ex Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, 60 species, type: *L. pulmonaria* (L.) Hoffm., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Moncada et al. (2013a; classification), Miadlikowska et al. (2014a; phylogeny), Cornejo and Scheidegger (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Lobariella Yoshim. 2002, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, 35 species, type: *L. crenulata* (Hook.) Yoshim., lichenized, neo-tropical distribution, see

Lumbsch and Huhndorf (2010; outline, accepted as a synonym of *Lobaria*), Moncada et al. (2013a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Lobarina Nyl. ex Cromb. 1894, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, 15 species, type: *L. scrobiculata* (Scop.) P. Gaertn., lichenized, Northern Hemisphere and in oceanic areas of Africa, Australia, New Zealand and South America, see Moncada et al. (2013a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Lobatopedis P.M. Kirk 1979, *Ascomycota* genera *incertae sedis*, five species, type: *L. foliicola* P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, on leaf litter, widespread, see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Heredia et al. (2014; new species), cultures and sequences are unavailable, holotype of type species IMI 222919.

Lobothallia (Clauzade & Cl. Roux) Hafellner 1991 (= *Aspicilia* subgen. *Lobothallia* Clauzade & Cl. Roux 1984; = *Protoplacodium* Motyka 1995 *fide* Species Fungorum 2017), *Megasporaceae*, *Pertusariales*, *Lecanoromycetes*, nine species, type: *L. alphoplaca* (Wahlenb.) Hafellner, lichenized, Asia, Europe, North America, see Lücking et al. (2017a; classification), cultures and sequences are available.

Lockeria K. D. Hyde 1994, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *L. palmicola* K.D. Hyde, asexual morph unknown, freshwater, Australia, Florida, see Raja and Shearer (2008; new species), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: BRIP 21334.

Loculistroma F. Patt. & Charles 1910, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *L. bambusae* F. Patt., Charles & Veihmeyer, asexual morph hyphomycetous, terrestrial, saprobes, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: BPI 632466.

Loculohypoxylon M.E. Barr 1976, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. grandineum* (Berk. & Ravenel) M.E. Barr, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Teichosporaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Loculotuber Trappe, Parladé & I.F. Alvarez 1993 (see Species Fungorum 2017 for synonyms), *Pezizales* genera *incertae sedis*, *Pezizomycetes*, two species, type:

L. gennadii (Chatin) Trappe, Parladé & I.F. Alvarez, asexual morph unknown, saprobes, hypogeous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lodderomyces van der Walt 1971, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, c. two species, type: *L. elongisporus* (Recca & Mrak) van der Walt, asexual reproduction is by multilateral budding, saprophytic, but some species are common human and animal pathogens, clinical isolates, soil, plant debris, beetles, water, yeast cake, worldwide, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Loekoesia S.Y. Kondr., S.O. Oh & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *L. austrocoreana* (S.Y. Kondr., L. Lököös & J.S. Hur) S.Y. Kondr., J. Kim, A.S. Kondr., S.O. Oh & J.S. Hur, lichenized, South Korea, see Kondratyuk et al. (2015d; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species KoLRI 015504.

Loflammia Vězda 1986, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *L. flammnea* (Müll. Arg.) Vězda, lichenized, Australia, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Loftamniopsis Lücking & Kalb 2000, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *L. brasiliensis* Lücking & Kalb, lichenized, Brazil, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type species Kalb, Kalb & Kalb 27140.

Logilia Vězda 1986, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *L. gilva* (Müll. Arg.) Vězda, lichenized, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Lohwagia Petr. 1942, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, three species, type: *L. intermedia* (Speg.) Petr., asexual morph unknown, from living leaves, terrestrial, central and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Lojkania Rehm 1905 (= *Sydowina* Petr. 1923 *fide* Species Fungorum 2017), *Testudinaceae*, *Pleosporales*, *Dothideomycetes*, ten species, type: *L. hungarica* Rehm, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Fenestellaceae*), Kirk et al.

(2013; genus accepted), Wijayawardene et al. (2014c; outline), Phookamsak and Hyde (2015; taxonomy, placed in *Testudinaceae*), cultures and sequences are available but lacking for the type species.

Lolia Abdel-Aziz & Abdel-Wahab 2011, *Lindgomycetaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *L. aquatica* Abdel-Aziz & Abdel-Wahab, coelomycetous, saprobes, aquatic, Egypt, see Abdel-Aziz and Abdel-Wahab (2010; taxonomy, phylogeny), Hyde et al. (2013; accepted in *Lindgomycetaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2012, 2014c; outline), Abdel-Aziz (2016a; new species, phylogeny), cultures and sequences are available, holotype and ex-type of the type species IMI 398675, MF644.

Loliomyces Maire 1937, *Ascomycota* genera *incertae sedis*, one species, type: *L. temulenti* Maire, hyphomycetous, saprobes, Morocco, see Seifert et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lollipoparia Inderb. 2001, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *L. minuta* Inderb., asexual morph unknown, saprobes, terrestrial, Thailand, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Lomaantha Subram. 1954, *Ascomycota* genera *incertae sedis*, three species, type: *L. pooga* Subram., hyphomycetous, sexual morph unknown, saprobes, dead wood and culms, India and China, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Ma et al. (2011c; new species, revision), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lomachashaka Subram. 1956, *Ascomycota* genera *incertae sedis*, one species, type: *L. kera* Subram., hyphomycetous, saprobes, Western Africa, India, see Yadav and Bhat (2009b; new species, India), Seifert et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Longicollum Zelski, F.R. Barbosa, Raja, A.N. Mill. & Shearer 2011, *Annulatasaceae*, *Annulatascales*, *Sordariomycetes*, one species, type: *L. biappendiculatum* Zelski, F.R. Barbosa, Raja, A.N. Mill. & Shearer, asexual morph unknown, saprobes, aquatic, USA, South America, see Zelski et al. (2011a; taxonomy), cultures and sequences are unavailable, holotype of the type: ILL 40794.

Longiostiolum Doilom, Ariyaw. & K.D. Hyde 2016, *Massarinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. tectonae* Doilom, Ariyaw. & K.D. Hyde, saprobes, terrestrial, Thailand, see Li et al. (2016a; taxonomy, phylogeny), cultures and sequences are available, ex-type strains of type: MFLUCC 12-0562,

MKT 078, ICMP 21259, holotype of type species MFLU 15-3532.

Longipedicellata H. Zhang, K.D. Hyde & J.K. Liu 2016, *Longipedicellataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. aptrootii* (K.D. Hyde & S.W. Wong) H. Zhang, K.D. Hyde & J.K. Liu, asexual morph hyphomycetous, saprobes, Asia, see Zhang et al. (2016a; taxonomy, phylogeny), Phukhamsakda et al. (2016b; evolution), cultures and sequences are available.

Lopadium Körb. [nom. cons.] 1855 (= *Heterothecium* Mont. 1852; = *Lopadiomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Lopadiaceae*, *Lecideales*, *Lecanoromycetes*, five species, type: *L. pezizoideum* (Ach.) Körb., lichenized, worldwide, see Ekman et al. (2008; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Lopadostoma (Nitschke) Traverso 1906 (= *Anthostoma* subgen. *Lopadostoma* Nitschke 1867 *fide* Species Fungorum 2017), *Lopadostomataceae*, *Xylariales*, *Sordariomycetes*, twelve species, type: *L. turgidum* (Pers.) Traverso, asexual morph libertella-like, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2014a), Senanayake et al. (2015), Maharachchikumbura et al. (2015b, 2016b; outline), Daranagama et al. (2016c; morphology, phylogeny), cultures and sequence are available but lacking for the type species.

Lopezaria Kalb & Hafellner 1990, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *L. versicolor* (Fée) Kalb & Hafellner, asexual morph unknown, lichenized, Morocco, see Seifert et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for type species AFTOL-ID 108.

Lophiella Sacc. 1878, *Dothideomycetes* genera *incertae sedis*, one species, type: *L. cristata* (Pers.) Sacc., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Lophiostomataceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, Index Fungorum (2016) treated type species as a synonym of *Lophiostoma angustilabrum* (Berk. & Broome) Cooke, holotype of type material unavailable.

Lophiohelichrysum Dayarathne, Camporesi & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *L. helichrysi* Dayarathne, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 15-0175, MFLUCC 15-0701.

Lophionema Sacc. 1883, *Dothideomycetes* genera *incertae sedis*, nine species, type: *L. vermisporum* (Ellis) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Lophiostomataceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lophiopoacea Ariyaw., Thambug., Camporesi & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *L. arundinis* Ariyaw., Thambugala, Camporesi & K.D. Hyde, asexual morph unknown, saprobes on *Poaceae*, terrestrial, Italy, Japan, see Thambugala et al. (2015b; morphology, phylogeny), cultures and sequenced are available, holotype and ex-type of type species MFLU 14-0590, MFLUCC 11-0463.

Lophiosphaerella Hara 1948, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *L. euryae* (Syd.) Hara, asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Li et al. (2014b; transferred to *Mycosphaerellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Lophistoma Ces. & De Not. 1863 (see Species Fungorum 2017 for synonyms), *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, c. 100 species, type: *L. macrostomum* (Tode) Ces. & De Not., asexual morph coelomycetous, only known for *L. semilibereum*, saprobes, terrestrial, aquatic, worldwide, sequence data available for four species, see Lumbsch and Huhndorf (2010; outline), Hirayama and Tanaka (2011; phylogeny), Hyde et al. (2013; outline), Kirk et al. (2013; genus accepted), Hirayama et al. (2014; new species), Wijayawardene et al. (2014c; outline), Thambugala et al. (2015b; phylogeny, review), cultures and sequences are available, epitype of the type: IFRD 2005.

Lophiotrema Sacc. 1878, *Lophiotremataceae*, *Pleosporales*, *Dothideomycetes*, 15 species, type: *L. nucula* (Fr.) Sacc., asexual morph coelomycetous, saprobes, terrestrial habitats, worldwide, see Hirayama and Tanaka (2011; taxonomy, phylogeny), Zhang et al. (2012e; taxonomy, phylogeny), Hyde et al. (2013, 2016; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Tibpromma et al. (2017b; new species), cultures and sequences are available, holotype of type: Scler. suec. n. 230.

Lophium Fr. 1818 (= *Lophidium* P. Karst 1873; = *Scolecostroma* Bat. & Peres 1960 *fide* Species Fungorum 2017), *Mytilinidiaceae*, *Mytilinidiales*, *Dothideomycetes*, five species, type: *L. mytilinum* (Pers.) Fr., asexual morph papulaspora-like, saprobes, terrestrial, Europe, America,

see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mytilinidiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Hernández-Restrepo et al. (2016a; new species), cultures and sequences are available.

Lophodermella Höhn. 1917, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, nine species, type: *L. sulcigena* (Link) Höhn. asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Lophodermium Chevall. 1826 (= *Lophomerum* Quell. & Magasi 1966 *fide* Jaklitsch et al. 2016a; see Species Fungorum 2017 for additional synonyms), *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, c. 145 species, type: *L. arundinaceum* (Schrad.) Chevall., asexual morph unknown, pathogens, saprobes, endophytes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Lophophacidium Lagerb. 1949 (= *Neonaumovia* Schwarzman 1959 *fide* Species Fungorum 2017), *Phaciaceae*, *Phaciidales*, *Leotiomycetes*, five species, type: *L. hyperboreum* Lagerb., asexual morph unknown, pathogens, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lafalmme et al. (2015; pathogens, phylogeny), cultures and sequences are available but lacking for the type species.

Lophotrichus R.K. Benj. 1949, *Microascaceae*, *Microascales*, *Sordariomycetes*, one species, type: *L. ampullus* R.K. Benj., asexual morph hyphomycetous (humicola-like), saprobes, human pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Eghrari et al. (2015; human pathogens, DNA), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, available specimen and cultures for the type: CBS H-14124, UAMH 11809.

Loramycetes W. Weston 1929, *Loramycetaceae*, *Helotioidales*, *Leotiomycetes*, two species, type: *L. juncicola* W. Weston, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; phylogeny), Tanney et al. (2016; DNA, phylogeny), cultures and sequences are available.

Loranitschka Lar. N. Vassiljeva 1990, *Nitschkiaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *L. viticola* Lar. N. Vassiljeva, asexual morph unknown, saprobes, terrestrial, Asia, see Vasilyeva et al. (2009, 2010, 2013; taxonomy, distribution), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus

accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of the type: VLA, Vasilyeva.

Loratospora Kohlm. & Volkm. 1993, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *L. aestuarii* Kohlm. & Volkm.-Kohlm., asexual morph unknown, saprobes, terrestrial, USA, Italy, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; revision), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015a; phylogeny), cultures and sequences are available, holotype and cultures available for the type: IMS J.K. 5505, JK 5535B.

Loricella Velen. 1934, *Callotiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *L. juncina* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Lotinia Pérez-Butrón, Fern.-Vic. & P. Alvarado 2015, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *L. verna* Pérez-Butrón, Fern.-Vic. & P. Alvarado, asexual morph unknown, saprobes, Spain, see Crous et al. (2015e; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Loxospora A. Massal. 1852, *Sarrameanaceae*, *Sarrameanales*, *Lecanoromycetes*, 13 species, type: *L. elatina* (Ach.) A. Massal., lichenized, Australia, Asia, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Loxosporopsis Henssen 1995, *Pertusariaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type: *L. corallifera* Brodo, Henssen & Imshaug, asexual morph unknown, lichenized, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Resl et al. (2015; DNA, phylogeny), cultures and sequences are available, holotype of type: herb. Henssen, Henssen 13604; isotypes CANL, FH, GZU, UPS, US.

Lucidascocarpa A. Ferrer, Raja & Shearer 2008, *Dothideomycetes* genera *incertae sedis*, one species, type: *L. pulchella* A. Ferrer, Raja & Shearer, asexual morph unknown, submerged wood from freshwater, Panama, Ecuador, see Ferrer et al. (2008; genus proposed), Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: ILL 40558 (AF178-2).

Luciotrichus R. Galán & Raity. 1995, *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *L. lasioboloides* R. Galán & Raity., asexual morph

unknown, saprobes, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: RG [sic] 6808.

Ludwigomyces Kirschst. 1939, *Ascomycota* genera *incertae sedis*, one species, type: *L. parasiticus* Kirschst., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lueckingia Aptroot & Umaña 2006, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *L. polyspora* Aptroot & Umaña, asexual morph unknown, lichenized, terrestrial, tropical, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type species INB, Aptroot 60206.

Lulwoana Kohlm., Volkm.-Kohlm., J. Campb., Spatafora & Gräfenhan 2005, *Lulworthiaceae*, *Lulorthiales*, *Sordariomycetes*, one species, type: *L. uniseptata* (Nakagiri) Kohlm., Volkm.-Kohlm., J. Campb., Spatafora & Gräfenhan, asexual morph unknown, endophytic, saprobes, marine, Japan, see Lumbsch and Huhndorf (2010; outline), Jones et al. (2015; outline of marine genera), Maharachchikumbura et al. (2015b, 2016b; outline), Panzer et al. (2015; DNA, distribution), Torta et al. (2015; endophytes, DNA), cultures and sequences are available.

Lulwoidea Kohlm., Volkm.-Kohlm., J. Campbell, Spatafora & Gräfenhan 2005, *Lulworthiaceae*, *Lulorthiales*, *Sordariomycetes*, one species, type: *L. lignoarenaria* (Jørg. Koch & E. B. G. Jones) Kohlm., Volkm.-Kohlm., J. Campb., Spatafora & Gräfenhan, asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Jones et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: Herb. J. Koch 398, slides 1–5 (CP) cultures and sequences available.

Lulworthia G.K. Sutherl. 1915 (= *Halophiobolus* Linder 1944 *fide* Species Fungorum 2017), *Lulworthiaceae*, *Lulorthiales*, *Sordariomycetes*, eleven species, type: *L. fucicola* G.K. Sutherl., asexual morph hyphomycetous, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), ex-neotype of type: ATCC 64288.

Lunatiannulus Daranag., Camporesi & K.D. Hyde 2015, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *L. irregularis* Daranag., Camporesi & K.D. Hyde, asexual morph libertella-like, saprobes, terrestrial, Italy, Europe, see Daranagama et al. (2015a; morphology, phylogeny), Maharachchikumbura et al. (2016b; morphology, phylogeny), holotype and ex-type strains of type species MFLU 14-0236, MFLUCC 14-0014.

Lunulospora Ingold 1942, *Sordariales* genera *incertae sedis*, two species, type: *L. curvula* Ingold, hyphomycetous, sexual morph unknown, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Luteocirrus C.F. Crane, B.L. Shearer & T.I. Burgess 2013, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *L. shearrii* C.F. Crane, B.L. Shearer & T.I. Burgess, coelomycetous, sexual morph unknown, pathogens, terrestrial, Australia, see Crane and Burgess (2013; taxonomy, phylogeny), Norphanphoun et al. (2016; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and the ex-type cultures of the type species PERTH 08439362, CBS 130776 = WAC 13425.

Luttrellia Shearer 1978, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, four species, type: *L. estuarina* Shearer, asexual morph unknown, saprobes, marine, fresh water, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Luzfridiella R.F. Castañeda & W.B. Kendr. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *L. insignis* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Lylea Morgan-Jones 1975, *Ascomycota* genera *incertae sedis*, six species, type: *L. catenulata* Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, Europe, see Karandikar and Singh (2010; new species, India), McKenzie (2009; new species, New Zealand), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Xia et al. (2014a, b; new species, China), cultures and sequences are unavailable.

Lyromma Bat. & H. Maia 1965 (= *Anconomyces* Cavalc. & A.A. Silva 1972; = *Lyrommotheca* Bat. & T. Herrera 1967 *fide* Species Fungorum 2017), *Lyrommataceae*, *Chaetothyriales*, *Eurotiomycetes*, seven species, type: *L. nectandrae* Bat. & H. Maia, coelomycetous, sexual morph unknown, lichenicolous, lichenized, worldwide, see Lücking (2008; new species, taxonomy), Wijayawardene et al. (2012; outline, accepted as *Pyrenulales*), Kirk et al. (2013; genus accepted), Flakus and Farkas (2013; new species, Bolivia), Farkas and Flakus (2015; South Africa), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Lysotheca Cif. 1962, *Ascomycota* genera *incertae sedis*, six species, type: *L. suprastromatica* Cif., coelomycetous, sexual morph unknown, saprobes, Dominican Republic, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Maasoglossum K.S. Thind & R. Sharma 1985, *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, one species, type: *M. verrucisporum* K.S. Thind & R. Sharma, asexual morph unknown, saprobes, terrestrial, Asia, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as *Leotiomycetes*), Kirk et al. (2013; genus accepted), Hustad and Miller (2015; revision, phylogeny), cultures and sequences are available, cultures available for the type: CUP-IN-000606.

Macgarvieomyces Klaubauf, Lebrun & Crous 2014, *Pyriculariaceae*, *Magnaportheales*, *Sordariomycetes*, two species, type: *M. borealis* (de Hoog & Oorschot) Klaubauf, Lebrun & Crous, hyphomycetous, pathogens, terrestrial, UK, Netherlands, see Klaubauf et al. (2014; phylogeny, morphology, asexual morph), Wijayawardene et al. (2017; outline), cultures and sequences available, ex-type strain of type species CBS 461.65.

Mackenziella Yanna & K.D. Hyde 2009 (= *Mackenziea* Yanna & K.D. Hyde 2002 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *M. livistonae* (Yanna & K.D. Hyde) Yanna & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Gams et al. (2009; nomenclature), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: HKU (M) 13304.

Macowaniella Doidge 1921, *Dothideomycetes* genera *incertae sedis*, two species, type: *M. congesta* (G. Winter) Doidge, epiphytes, terrestrial, see Hofmann (2009; review, accepted in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, treated as a doubtful genus in *Asterinales*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are unavailable.

Macroallantina Speer 1987, *Ascomycota* genera *incertae sedis*, one species, type: *M. caespitosa* Petr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Macroconia (Wollenw.) Gräfenhan, Seifert & Schroers 2011 (= *Nectria* sect. *Macroconia* Wollenw. 1926 *fide* Species Fungorum 2017), *Nectriaceae*, *Hypoocreales*, *Sordariomycetes*, five species, type: *M. leptosphaeriae* (Niessl) Gräfenhan & Schroers, hyphomycetous, terrestrial,

mostly fungicolous, Northern hemisphere, see Gräfenhan et al. (2011; phylogeny), Lombard et al. (2015b; phylogeny), cultures and sequences are available, lectotype of type: K(M) 165805.

Macroderma Höhn. 1917, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *M. curtisii* (Berk. & Ravenel) Höhn., asexual morph unknown, saprobes, terrestrial, North America, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Macrodiaporthe Petr. 1920 (= *Calospora* Nitschke ex Fuckel 1870 [1869-70] *fide* Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *M. occulta* (Fuckel) Petr., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Macrodiploidia Sacc. 1884, *Ascomycota* genera *incertae sedis*, two species, type: *M. curreyi* (Sacc. & Roum.) Sacc., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Macrodiplodiopsis Petr. 1922 (= *Macrodiplois* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Macrodiplodiopsidaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *M. desmazieri* (Mont.) Petr., asexual morph coelomycetous, saprobes, Europe, see Wijayawardene et al. (2012, 2014b, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), Crous et al. (2015a; taxonomy, sexual morph, typification), Tanaka et al. (2015; phylogeny), cultures and sequences are available, lectotype: PC 0142158, epitype and ex-epitype: CBS H-22269, CBS 140062.

Macrographa Etayo 2008, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *M. antarctica* Etayo, asexual morph unknown, terrestrial, parasympiont, or parasite on *Nephroma antarcticum*, South America, see Etayo and Sancho (2008; morphology), Hyde et al. (2013; accepted as *Trichothyriaceae*, but without molecular data), cultures and sequences are unavailable, holotype of the type species MAF, Etayo 23269.

Macrohilum H.J. Swart 1988, *Macrohilaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *M. eucalypti* H.J. Swart, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2016b; outline, morphology, phylogeny), Crous et al. (2015a; morphology, phylogeny),

cultures and sequences are available, epitype and ex-epitype of the type: CBS H-22279, CBS 140063.

Macrophomina Petr. 1923, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, three species, type: *M. phaseolina* (Tassi) Goid., coelomycetous, sexual morph unknown, saprobes, pathogens, in soil, terrestrial, cosmopolitan, see Phillips et al. (2008, 2013; taxonomy, phylogeny), Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c, 2016b; outline, taxonomy, phylogeny), Dissanayake et al. (2016b; taxonomy), cultures and sequences are available.

Macrorhabdus Tomaszewski, Logan, Snowden, Kurtzman & Phalen 2003, *Saccharomycetales* genera *incertae sedis*, *Saccharomycetes*, one species, type: *M. ornithogaster* Tomasz., asexual reproduction is by fission, typically infects the stomach area of budgerigars (*Melopsittacus undulatus*) and other birds causing a wasting disease, on Budgerigars and other birds, worldwide, sequences are available.

Macroskytea Etayo, Flakus, Suija & Kukwa 2015, *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, one species, type species: *M. parmotrematis* Etayo, Flakus & Kukwa, asexual morph unknown, lichenicolous, terrestrial, Bolivia, see Etayo et al. (2015; DNA, description), DNA sequences available, cultures unavailable, holotype of type species in UGDA.

Macrovalsaria Petr. 1962, *Dothideomycetes* genera *incertae sedis*, two species, type: *M. leonensis* (Deighton) Petr., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in *Botryosphaeriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted in *Botryosphaeriaceae*), Doilom et al. (2017; DNA, phylogeny), cultures and sequences are available but lacking for the type species.

Macroventuria Aa 1971, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *M. anomochaeta* Aa, asexual morph unknown, saprobes, ?pathogens, terrestrial, cosmopolitan, see Aveskamp et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Chen et al. (2015b; taxonomy, phylogeny), cultures and sequences are available.

Maculatifrondes K.D. Hyde 1996, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *M. aequatoriensis* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline),

cultures and sequences are unavailable, holotype of type: BRIP 23240.

Maculatipalma J. Fröhl. & K.D. Hyde 1995, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *M. frondicola* J. Fröhl. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: BRIP 21402.

Madurella Brumpt 1905 (= *Indiella* Brumpt 1906; = *Rubromadurella* Talice 1935 *fide* Species Fungorum 2017), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, nine species, type: *M. mycetomatis* (Laveran) Brumpt, hyphomycetous, sexual morph unknown, saprobes, human (causal agent of black-grain mycetoma) and animal pathogens, worldwide, see Fahala et al. (2010; pathogen, treatments), Seifert et al. (2011; morphology), de Hoog et al. (2012, 2013; new species, Sudan, Sumatera), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Irinyi et al. (2015; barcode), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, cultures available for the type species CBS 247.48.

Magmopsis Nyl. 1875 (see Species Fungorum 2017 for synonyms), *Arthopyreniaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. pertenella* Nyl., lichenized, see Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Magnaportheiopsis J. Luo & N. Zhang 2013, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, six species, type: *M. poae* (Landsch. & N. Jacks.) J. Luo & N. Zhang, asexual morph hyphomycetous, pathogens, terrestrial, worldwide, see Luo and Zhang (2013; taxonomy, phylogeny), Klaubauf et al. (2014; new species, phylogeny), Luo et al. (2014; new species, phylogeny), Crous et al. (2015e; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Khemmuk et al. (2016; phylogeny), holotype and cultures available for the type species DAR 59044, M47.

Magnibotryascoma Thambug. & K.D. Hyde 2015, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. uniseriatum* (Mugambi, A.N. Mill. & Huhndorf) Thambug. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, USA, see Thambugala et al. (2015b; phylogeny), Liu et al. (2017, phylogeny, placement), cultures and sequences are available, holotype of type species ANM 909.

Magnicamarosporium Kaz. Tanaka & K. Hiray. 2015, *Sulcatisporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type species: *M. iriomotense* Kaz. Tanaka & K. Hiray., coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Tanaka et al. (2015; morphology,

phylogeny), Wijayawardene et al. (2016b, 2017; morphology, phylogeny, outline), Phukhamsakda et al. (2017; new species, phylogeny), cultures and sequences are available, holotype and ex-type: HHUF 30125, CBS 139696.

Magnisphaera J. Campb., J.L. Anderson & Shearer 2003 (= *Matsusphaeria* K.L. Pang & E.B.G. Jones *fide* Maharachchikumbura et al. 2015b, 2016b), *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *M. spartinae* (E.B.G. Jones) J. Campb., J.L. Anderson & Shearer, asexual morph unknown, saprobes, marine, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Réblová et al. (2017; DNA, phylogeny), cultures and sequences are available.

Magnohelicospora R.F. Castañeda, Hern.-Restr., Gené & Guarro 2013, *Venturiaceae*, *Venturiales*, *Dothideomycetes*, two species, type: *M. iberica* R.F. Castañeda, Hern.-Restr., Gené & Guarro, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Portugal, see Castañeda-Ruiz et al. (2013c; taxonomy), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype of the type: HAL (Fungi) 2447, cultures and sequences are unavailable.

Magnusiomyces Zender 1977 (= *Saprochaete* Coker & Shanor ex D.T.S. Wagner & Dawes 1970; see Index Fungorum 2017 for additional synonyms), *Dipodascaceae*, *Saccharomycetales*, *Saccharomycetes*, eight species, type: *M. ludwigii* Zender, asexual reproduction consists of true hyphae that branch at acute angles with acuminate apices, and that disarticulate into arthroconidia, saprophytic, but clinical isolates are not uncommon, soil, tree fluxes, waste water, insect tunnels in trees, worldwide, cultures and sequences are available.

Mahabalella B. Sutton & S.D. Patil 1966, *Ascomycota* genera *incertae sedis*, two species, type: *M. acutisetosa* B. Sutton & S.D. Patil, hyphomycetous, with acremonium-like synasexual morph, sexual morph unknown, saprobes, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Maheşhwaramyces Hosag. 2009, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. pachygones* Hosag., Archana & Dan, asexual morph unknown, living on leaves, India, see Hosagoudar et al. (2009; taxonomy), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Maireella Syd. ex Maire 1908, *Dothideomycetes* genera *incertae sedis*, type: *M. maculans* Syd. ex Maire, data inadequate, see Kirk et al. (2008; treated as a synonym of

Gibbera, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Majewskia Y.B. Lee & K. Sugiy. 1986, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *M. japonica* Y.B. Lee & K. Sugiy., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Malacaria Syd. 1930, *Dothideomycetes* genera *incertae sedis*, two species, type: *M. meliolicola* Syd., asexual morph unknown, saprobes, pantropical, see Lumbsch and Huhndorf (2010; outline, accepted in *Tubeufiaceae*), Hyde et al. (2013; not accepted in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; not listed in *Tubeufiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Malaysiasca Crous & M.J. Wingf. 2016, *Glomerellales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *M. phaii* Crous & M.J. Wingf., asexual morph coelomycetous, saprobes, terrestrial, Malaysia, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type: CBS H-22616, CPC 27548.

Malbranchea Sacc. 1882 (= *Thermoidium* Miehe 1907 *fide Species Fungorum* 2017), *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, 22 species, type: *M. pulchella* Sacc. & Penz., hyphomycetous, sexual morph unknown, in soil, dung, keratin, cellulose, man and other substrata, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Malcolmella Vězda 1997, *Teloschistales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *M. cinereovirens* Vězda, lichenized, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available, holotype of type: CHR (2757).

Mallochia Arx & Samson 1986, *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, four species, type: *M. echinulata* (B.G. Dutta & G.R. Ghosh) Arx & Samson, asexual morph unknown, saprobes, palaeotropical, see Lumbsch and Huhndorf (2010; outline, accepted in *Onygenaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, available specimen for type: CBS H-6703.

Malneomyces Starbäck 1899, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. pulchellus* Starbäck, asexual morph unknown,

terrestrial, saprobes, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Malmidea Kalb, Rivas Plata & Lumbsch 2011, *Malmideaceae*, *Lecanorales*, *Lecanoromycetes*, 50 species, type: *M. piperis* (Spreng.) Kalb, Rivas Plata & Lumbsch, ascomata apotheciate, conidiomata pycnidial, lichen-forming, mostly on bark, pantropical, see Kalb et al. (2012; phylogeny), Miądlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Malmographina M. Cáceres, Rivas Plata & Lücking 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *M. plicosa* (C.F.W. Meissn.) M. Cáceres, Rivas Plata & Lücking, asexual morph unknown, lichenized, terrestrial, pantropical, see Cáceres et al. (2012a; taxonomy), Lumbsch et al. (2014a, b; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Malthomyces K.D. Hyde & P.F. Cannon 1999, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *M. calamigena* (Berk. & Broome) K.D. Hyde & P.F. Cannon, asexual morph unknown, saprobes, terrestrial, India, Sri Lanka, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Mamiania Ces. & De Not. 1863 (= *Kubinia* Schulzer 1866 *fide Species Fungorum* 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *M. fimbriata* (Pers.) Ces. & De Not., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Mamianiella Höhn. 1917, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, three species, type: *M. coryli* (Batsch) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Sogonov et al. (2008; accepted as *Diaporthales*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Gnomoniaceae*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as *Diaporthales*), cultures and sequences are available, available specimen and cultures for the type: CBS H-14211, BPI 877578.

Mamillisphaeria K.D. Hyde, S.W. Wong & E.B.G. Jones 1996, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. dimorphospora* K.D. Hyde, S.W. Wong & E.B.G. Jones, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and

Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Melanommataceae*), Wijayawardene et al. (2014c; outline), Tian et al. (2015; accepted as a genus in *Melanommataceae*), holotype of the type: BRIP 22967, cultures and sequences are unavailable.

Mammaria Ces. ex Rabenh. 1854 (= *Pseudocercophora* Subram. & Sekar 1986 *fide* Réblová et al. 2016c), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, two species, type: *M. echinobotryoides* Ces., sexual morph unknown, saprobes, terrestrial, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, available specimen for type: CBS H-14212.

Manglicola Kohlm. & E. Kohlm. 1971, *Mangicolaceae*, *Jahnulales*, *Dothideomycetes*, two species, type: *M. guatemalensis* Kohlm. & E. Kohlm., asexual morph unknown, saprobes, marine, Brunei, Guatemala, Thailand, see Jones et al. (2009b, 2015; classification), Suetrong et al. (2010, 2011a; morphology, phylogeny), Jones et al. (2012; morphology, phylogeny), Hyde et al. (2013; morphology, phylogeny), Shearer et al. (2014; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available, holotype and ex-isotype strain of type species J.K. Nos. 2700 (NY), J.K. 2700a (IMS).

Mangoldia Lücking, Parnmen & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *M. australiana* Lücking, Parnmen & Lumbsch, asexual morph unknown, lichenized, terrestrial, Australia, see Lücking et al. (2012, 2014; taxonomy), Lumbsch et al. (2014a, b; phylogeny), sequences are available, holotype: CANB, Mangold 27zB.

Mangrovispora K.D. Hyde & Nakagiri 1991, *Sordariomycetes* genera *incertae sedis*, two species, type: *M. pemphii* K.D. Hyde & Nakagiri, asexual morph unknown, saprobes, marine, Australia, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; accepted as *Phyllachorales*), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available (unpublished), holotype of the type: BRIP 17112.

Manoharachariella Bagyan., N.K. Rao & Kunwar 2009, *Tubeufiaceae*, *Tubeufiales*, *Dothidiomycetes*, four species, type: *M. lignicola* Bagyan., N.K. Rao & Kunwar, sexual morph unknown, saprobes, terrestrial, Asia and Turkey, see Bagyanarayana et al. (2009; established genus), Rajeshkumar and Singh (2012; new species), Selçuk and Ekici (2014; new species), Doilom et al. (2017; new species), cultures and sequences are available for *M. tectonae*, holotype of type: IMI 296857.

Manokwaria K.D. Hyde 1993, *Xylariomycitidae* genera *incertae sedis*, *Sordariomycetes*, one species, type: *M. notabilis* K.D. Hyde, asexual morph unknown, saprobes, fresh water, Indonesia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: BRIP 20854.

Mapletonia B. Sutton 1991, *Ascomycota* genera *incertae sedis*, one species, type: *M. falcata* B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type species IMI 263431a.

Marasasiomyces Crous 2015, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, one species, type: *M. karo* (B. Sutton & Marasas) Crous, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Crous et al. (2015b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-isotype of the type: PREM 44967, CBS 118718.

Marcelaria Aptroot, Nelsen & Parnmen 2013 (= *Buscalionia* Cengia Sambo 1940), *Trypetheliaceae*, *Trypethliales*, *Dothideomycetes*, three species, type: *M. purpurina* (Nyl.) Aptroot, Nelsen & Parnmen, asexual morph unknown, lichenized, terrestrial, worldwide, see Aptroot et al. (2013b; taxonomy), Nelsen et al. (2014b; DNA, phylogeny), Luangsuphabool et al. (2016; DNA, phylogeny), Aptroot and Nepi (2017; nomenclature), cultures and sequences are available, holotype and cultures available for the type species 236 (PC), MPN323A (*fide* Nelsen et al. 2014).

Marcelleina Brumm., Korf & Rifai 1967, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *M. persoonii* (P. Crouan & H. Crouan) Brumm., asexual morph unknown, hypogeous, terrestrial, worldwide, see Lantieri and Pfister (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rubio et al. (2011; new species, Spain), cultures and sequences are available, cultures available for type: TL-5696.

Marchantiana S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *M. occidentalis* (Elix, S. Y. Kondr. & Kärnefelt) S.Y. Kondr., Kärnefelt, Elix, A. Thell, J. Kim, A.S. Kondr. & Hur, asexual morph unknown, lichenized, terrestrial, Australia, see Kondratyuk et al. (2014, 2015b; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available, holotype and available cultures of the type: CANB Elix 32479, SK981 and SK982.

Margaritispora Ingold 1942, *Helotiales* genera *incertae sedis*, one species, type: *M. aquatica* Ingold, hyphomycetous, sexual morph unknown, saprobes,

aquatic, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Mariannaea G. Arnaud ex Samson 1974 (= *Mariannaea* G. Arnaud 1952 *fide* Species Fungorum 2017), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *M. elegans* (Corda) Samson, asexual morph hyphomycetous, terrestrial, saprobes and in soil, Asia, see Gräfenhan et al. (2011; phylogeny), Lombard et al. (2015b; phylogeny), cultures and sequences are available, holotype of type species in herb. Corda (as *Penicillium elegans*).

Marielliotia Shoemaker 1999, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *M. biseptata* (Sacc. & Roum.) Shoemaker, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology, accepted AY004786 (DAOM 208987) as the ITS barcode), Wijayawardene et al. (2012, 2014c, 2017; outline), Ariyawansa et al. (2015c; accepted as a genus in *Phaeosphaeriaceae*), cultures and sequences are available.

Marinokulati E.B.G. Jones & K.L. Pang 2014, *Juncigenaceae*, *Torpedosporales*, *Sordariomycetes*, one species, type: *M. chaetosa* (Kohlm.) E.B.G. Jones & K.L. Pang, saprobes, marine, temperate, asexual morph unknown, see Jones et al. (2014, 2015; morphology, phylogeny; outline), cultures and sequences are available.

Marinosphaera K.D. Hyde 1989, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *M. mangrovei* K.D. Hyde, asexual morph unknown, saprobes, marine, terrestrial, Brunei, see Jones et al. (2009b, 2015; phylogeny, outline), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of the type: IMI 325412.

Marinospora A.R. Caval 1966 (= *Ceriosporella* A.R. Caval. 1966 *fide* Species Fungorum 2017), *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *M. calyptata* (Kohlm.) A.R. Caval., asexual morph unknown, saprobes, marine, Argentina, Belgium, Denmark, France, Germany, Scotland, Iceland, Spain, USA, Canada, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), cultures and sequences are available.

Marisolaris Jørgen Koch & E.B.G. Jones 1989, *Sordariomycetes* genera *incertae sedis*, one species, type: *M. ansata* Jørg. Koch & E.B.G. Jones, asexual morph unknown, marine, temperate, on wood associated with sand, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Ascomycota* genera *incertae sedis*), Kirk et al.

(2013; genus accepted), Jones et al. (2015; outline), no sequence data.

Maronea A. Massal. 1856 (= *Maroneomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Fuscideaceae*, *Umbilicariales*, *Lecanoromycetes*, 13 species, type: *M. berica* A. Massal., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; phylogeny), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Maronella M. Steiger 1959 (= *Myrioblastus* Trevis. 1857; = *Piccollia* A. Massal. 1856 *fide* Species Fungorum 2017), *Lecanoromycetes* genera *incertae sedis*, one species, type: *M. laricina* M. Steiner, lichenized, temperate northern hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Marthamyces Minter 2003, *Marthamycetaceae*, *Rhytismatales*, *Leotiomycetes*, nine species, type: *M. emarginatus* (Cooke & Massee) Minter, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lantz et al. (2011; DNA, phylogeny), cultures and sequences are available.

Martinellisia V.G. Rao & Varghese 1977, *Ascomycota* genera *incertae sedis*, one species, type: *M. indica* V.G. Rao & Varghese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Martininia Dumont & Korf 1970 (= *Martinia* Whetzel 1942 *fide* Species Fungorum 2017), *Sclerotiniaceae*, *Helotiidales*, *Leotiomycetes*, one species, type: *M. panamaensis* Dumont & Korf, asexual morph unknown, saprobes, terrestrial, America, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype and ex-type of the type: CUP 27040, CBS 207.47.

Martiniozyma Kurtzman 2015, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *M. abiesophila* (Kurtzman) Kurtzman, asexual reproduction is by multilateral budding, saprophytic, water, soil, fruit, sap of red fir, worldwide, see Kurtzman (2015; taxonomy), cultures and sequences are available.

Martinjahnsia S.Y. Kondr., Fedorenko, S. Stenoos, Kärnefelt, Elix, J.S. Hur & A. Thell 2012, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, ten species, type: *M. resendei* (Poelt & C. Tav.) S.Y. Kondr., Fedorenko, S. Stenoos, Kärnefelt, Elix, J.S. Hur & A. Thell, asexual morph unknown, lichenized, terrestrial, worldwide, see

Fedorenko et al. (2012; taxonomy), cultures and sequences are unavailable.

Massalongia Körb. 1855 (= *Massalongomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Massalongiaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *M. cariosa* (Dicks.) Körb., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Massalongina Bubák 1916, *Ascomycota* genera *incertae sedis*, one species, type: *M. aquilina* (C. Massal.) Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Massaria De Not. 1844 (= *Aglaospora* De Not. 1844; = *Bathystomum* Füsing 1868; = *Phaeomassaria* Speg. 1880 *fide* Species Fungorum 2017), *Massariaceae*, *Pleosporales*, *Dothideomycetes*, 31 species, type: *M. inquinans* (Tode) De Not., asexual morph coelomycetous, saprobes or weak parasitic, terrestrial, northern temperate climate zone, see Lumbsch and Huhndorf (2010; outline), Voglmayr and Jaklitsch (2011; taxonomy, phylogeny), Hyde et al. (2013, 2016; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequence data available for 17 species, holotype of type species WU 30526.

Massarina Sacc. 1883 (= *Abaphospora* Corda 1939; = *Bertia* subgen. *Bertiella* Sacc. 1882; = *Massarinula* Géneau 1894; = *Oraniella* Speg. 1909; = *Parasphaeria* Syd. 1924; = *Pseudodiaporthe* Speg. 1909 *fide* Species Fungorum 2017), *Massarinaceae*, *Pleosporales*, *Dothideomycetes*, c. 100 species, type: *M. eburnea* (Tul. & C. Tul.) Sacc., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Zhang et al. (2009a, b; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Tanaka et al. (2015; taxonomy, phylogeny), cultures and sequences are available.

Massariola Füsing 1868, *Dothideomycetes* genera *incertae sedis*, two species, type: need typification, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Massariosphaeria (E. Müll.) Crivelli 1983 (= *Leptosphaeria* subgen. *Massariosphaeria* E. Müll. 1950 *fide* Species Fungorum 2017), *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, 25 species, type: *M. phaeospora* (E. Müll.) Crivelli, asexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf

(2010; outline, accepted as a genus in *Lophiostomataceae*), Tanaka et al. (2011b; new species, Pakistan), Hyde et al. (2013; excluded from *Lophiostomataceae*) Kirk et al. (2013; genus accepted), Thambugala et al. (2015b; excluded from *Lophiostomataceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Pleosporales*), cultures available for the type: CBS 611.86 (*fide* Schoch et al. 2009).

Massariothea Syd. 1939 (= *Apocoryneum* B. Sutton 1975 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, eight species, type: *M. themiae* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Massariovalsa Sacc. 1882 (= *Melanconiopsis* Ellis & Everh. *fide* Rossman et al. 2015a), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *M. sudans* (Berk. & M.A. Curtis) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongisan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable, available specimen for type: CBS H-14272.

Masseea Sacc. 1889, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, ?four species, type: *M. quisquiliarum* (Berk. & M.A. Curtis) Sacc., asexual morph unknown, saprobes, terrestrial, America, Asia see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mastigosporella Höhn. 1914 (= *Wuestneiopsis* J. Reid & Dowsett 1990 *fide* Rossman et al. 2015a), *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *M. hyalina* (Ellis & Everh.) Höhn., asexual morph coelomycetous, pathogens, saprobes, terrestrial, worldwide, see Crous et al. (2013b; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2015a; nomenclature), Wijayawardene et al. (2017; outline), cultures and sequences are available but lack for the type.

Mastigosporium Riess 1852 (= *Amastigis* Clem. & Shear 1931; = *Amastigosporium* Bondartsev 1921 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, four species, type: *M. album* Riess, coelomycetous, sexual morph unknown, pathogens, terrestrial, temperate, see Kirk et al. (2013; genus accepted), Crous et al. (2014a; taxonomy, epitypification, phylogeny), cultures and sequences are available but lacking for the type species.

Mastodia Hook.f. & Harv. 1847 (= *Dermatomeris* Reinsch 1890; = *Kohlmeyera* S. Schatz 1980; = *Leptogiopsis* Nyl. 1884 *fide* Species Fungorum 2017), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, five species, type: *M. tessellata* (Hook. f. & Harv.) Hook. f. & Harv., lichenized, Southern hemisphere, especially Antarctica, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Mastodiaceae*), Pérez-Ortega et al. (2010; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Matsushimaea Subram. 1978, *Ascomycota* genera *incertae sedis*, three species, type: *M. fasciculata* (Matsush.) Subram., sexual morph unknown, saprobes, terrestrial, Asia, North America, Africa, see Hyde et al. (2011; checklist), Rambelli (2011; Mediterranean), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Matsushimaomyces Rahul Sharma & Rohit Sharma 2015, *Latoruaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. bohaniensis* Rahul Sharma, Rohit Sharma & Crous, hyphomycetous, keratinophilic, from soil, India, see Sharma et al. (2015a; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Matsushimiella R.F. Castañeda & Heredia 2001, *Ascomycota* genera *incertae sedis*, two species, type: *M. queenslandica* (Matsush.) R.F. Castañeda & Heredia, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Monteiro et al. (2015a; new species, Brazil), holotype of type: MFC-8A063, cultures and sequences are unavailable, compare with *Minimelanolocus*, *Nigrolentilocus*, *Pseudospiropes* and *Spiropes*.

Matsushimomyces V.G. Rao & Varghese 1979, *Ascomycota* genera *incertae sedis*, two species, type: *M. indicus* V.G. Rao & Varghese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Matsusporium E.B.G. Jones & K.L. Pang 2010, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, one species, type: *M. tropicale* (Kohlm.) E.B.G. Jones & K.L. Pang, hyphomycetous, sexual morph unknown, saprobes, marine, pantropical, see Abdel-Wahab et al. (2010; phylogeny), cultures and sequences are available, holotype of type: IMI 325412.

Mattirolella S. Colla 1929, *Kathistaceae*, *Ophiostomatales*, *Sordariomycetes*, one species, type: *M. silvestrii* S. Colla, saprobes, Africa, see

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharanachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Mattirolia Berl. & Bres. 1889, *Thyridiaceae*, *Sordariomycetes* families *incertae sedis*, one species, type: *M. roseovirens* Berl. & Bres., saprobes, Africa, see Lumbsch and Huhndorf (2010; outline), Checa et al. (2013; new species), Kirk et al. (2013; genus accepted), Maharanachchikumbura et al. (2015b, 2016b; outline), sequences and cultures are unavailable, epitype of type: WU 32154.

Mattirolomyces E. Fisch. 1938, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *M. terfezioides* (Mattirol.) E. Fisch., asexual morph unknown, hypogeous, terrestrial, worldwide, see Trappe et al. (2010b; DNA), Kovács et al. (2011b; DNA, new combination), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Maublancia G. Arnaud 1918 (= *Caudellopeltis* Bat. & H. Maia 1960 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *M. myrtacearum* G. Arnaud, asexual morph unknown, saprobes, terrestrial, South America, South Africa, see Lumbsch and Huhndorf (2010; outline, accepted in *Microthyriaceae*), Wu et al. (2011; excluded from *Microthyriaceae*), Hyde et al. (2013; excluded from *Microthyriaceae*), Wijayawardene et al. (2014c; accepted in *Dothideomycetes*), cultures and sequences are unavailable, available specimen of type: CBS H-14276.

Mauginiella Cavara 1925, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. scaettiae* Cavara, asexual morph unknown, saprobes, pathogens, terrestrial, South America, South Africa, see Seifert et al. (2011; morphology), Abdullah et al. (2015; pathogens), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Mauritiana Poonyth, K.D. Hyde, Aptroot & Peeraly 2000, *Halothiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. rhizophorae* Poonyth, K.D. Hyde, Aptroot & Peeraly, asexual morph unknown, saprobes, marine, see Alias and Jones (2009; monograph), Suetrong et al. (2009; phylogeny, morphology), Jones et al. (2009b, 2015; outline, marine), Zainuddin et al. (2010; antimicrobial activities), Lee et al. (2012; checklist, Malaysia), Zhang et al. (2012e; phylogeny, morphology), Hyde et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available, holotype of type: HKU(M) 10219.

Mawsonia C.W. Dodge 1948, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *M. harrissonii* C.W. Dodge, lichenized, Brazil, Antarctica, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Mazaediothecium Aptroot 1991, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, three species, type: *M. rubiginosum* Aptroot, asexual morph unknown, lichenized, terrestrial, Papua New Guinea, Costa Rica, Malaysia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type: Aptroot (17464).

Mazosia A. Massal. 1854, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, 27 species, type: *M. rotula* (Mont.) A. Massal., sexual morph with apothecoid ascocarps, asexual morph coelomycetous, lichenized, rarely lichenicolous, terrestrial, tropical to subtropical regions, see Lücking (2008a, b; morphology), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ertz et al. (2015b, c; phylogeny), Frisch et al. (2014; phylogeny), Lücking et al. (2017a; classification), sequence data available for a few species.

Mazzantia Mont. 1855 (= *Mazzantiella* Höhn. 1925 *fide* Rossman et al. 2015a), *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *M. galii* (Fr.) Mont., asexual morph coelomycetous, saprobes, terrestrial, Europe, America, see Sogonov et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for the type: AR4658 (*fide* Udayanga et al. 2014a, b).

Mebaria J. Reid & C. Booth 1989, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *M. thujina* (Nag Raj & DiCosmo) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Medeolaria Thaxt. 1922, *Medeolariaceae*, *Medeolariales*, *Leotiomycetes*, one species, type: *M. farlowii* Thaxt., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Medusamyces G.L. Barron & Szijarto 1990, *Ascomycota* genera *incertae sedis*, two species, type: *M. lunulosporus* G.L. Barron & Szijarto, hyphomycetous, sexual morph unknown, on rotifers, aquatic, Canada, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: OAC 10845.

Megacapitula J.L. Chen & Tzean 1993, *Ascomycota* genera *incertae sedis*, two species, type: *M. villosa* J.L. Chen & Tzean, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Prabhugaonkar and Bhat (2011; record in India, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype and cultures available for the type: PPH17, GUFCC 15515.

Megalaria Hafellner 1984, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, 29 species, type: *M. grossa* (Pers. ex Nyl.) Hafellner, lichenized, worldwide, see Ekman et al. (2008; Ancestral state reconstructions), Schmull et al. (2011; phylogeny), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Megaloblastenia Sipman 1983, *Megalosporaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *M. flavidatra* (Nyl.) Sipman, lichenized, Australia, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Megalodochium Deighton 1960, *Ascomycota* genera *incertae sedis*, four species, type: *M. palmicola* Deighton, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, China, India, Sierra Leone (tropical Africa), see Tiwari and Rai (2009; morphology), Wijayawardene et al. (2012, 2017; outline), Zhao and Zhao (2012; morphology), Crous et al. (2015d; compare with *Melanconium*), cultures and sequences are unavailable.

Megalohyppha A. Ferrer & Shearer 2007, *Aliquandostipitaceae*, *Jahnulales*, *Dothideomycetes*, one species, type: *M. aqua-dulces* A. Ferrer & Shearer, freshwater, Panama, Thailand, see Lumbsch and Huhndorf (2010; outline), Suetrong et al. (2011a; phylogeny), Hyde et al. (2013; accepted as a genus in *Aliquandostipitaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available, holotype of type: AF005 ILL.

Megaloseptoria Naumov 1925 (see Species Fungorum 2017 for synonyms), *Ascomycota* genera *incertae sedis*, one species, type: *M. mirabilis* Naumov, coelomycetous, sexual morph unknown, saprobes, terrestrial, Russia, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Megalospora Meyen 1843 (= *Bombyliospora* De Not. 1852; = *Bombyliosporomyces* Cif. & Tomas. 1953; = *Byssophragmia* M. Choisy 1931; = *Dumoulinia* Stein 1883; = *Heterothecium* Flot. 1850 *fide* Species Fungorum 2017), *Megalosporaceae*, *Teloschistales*, *Lecanoromycetes*, 35 species, type: *M. sulphurata* Meyen, lichenized, tropical and warm temperate zones, see Lumbsch and Huhndorf (2010; outline), Kantvilas and Lumbsch (2012; phylogeny),

Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Megalotremis Aptroot 1991, *Monoblastiaceae*, *Monoblastiales*, *Dothideomycetes*, twelve species, type: *M. verrucosa* (Makhija & Patw.) Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, India, Costa Rica, Sri Lanka, see Nelsen et al. (2009, 2011a, b; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures available for type: MPN104 (*fide* Nelsen et al. 2011a, b).

Megaspora (Clauzade & Cl. Roux) Hafellner & V. Wirth 1987 (= *Aspicilia* subgen. *Megaspora* Clauzade & Cl. Roux 1984 *fide* Species Fungorum 2017), *Megasporaceae*, *Pertusariales*, *Lecanoromycetes*, three species, type: *M. verrucosa* (Ach.) Hafellner & V. Wirth, lichenized, Austria, Iran, Armenia, see Lumbsch and Huhndorf (2010; outline), Valadbeigi et al. (2011a; new species), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Zakeri et al. (2016; new species), cultures and sequences are available.

Meionomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, six species, type: *M. asteni* Thaxt., asexual morph unknown, biotrophic, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Melanamphora Lafl. 1976, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *M. spinifera* (Wallr.) Lafl., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Melanconiella Sacc. 1882, *Melanconiellaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *M. spodiaea* (Tul. & C. Tul.) Sacc., asexual morph discosporina-like and melanconium-like, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Voglmayr et al. (2012; taxonomy, phylogeny, asexual morph, typification), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available, cultures available for type species SPOD1, SPOD, MVS, MSH (*fide* Voglmayr et al. 2012).

Melanconis Tul. & C. Tul. 1863 (= *Melanconium* Link 1809 *fide* Rossman et al. 2015a), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, c. 100 species, type:

M. stilbostoma (Fr.) Tul. & C. Tul., asexual morph formerly known in *Melanconium*, saprobes, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Fan et al. (2016b; species associated with *Betula*), Wijayawardene et al. (2016b, 2017; taxonomy of *Melanconium*, outline), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are available.

Melanelia Essl. 1978, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *M. stygia* (L.) Essl., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Melanelisia O. Blanco, A. Crespo, Divakar, Essl., D. Hawksw. & Lumbsch 2004, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, eight species, type: *M. glabra* (Schaer.) O. Blanco, A. Crespo, Divakar, Essl., D. Hawksw. & Lumbsch (syn. *Parmelia olivacea* f. *glabra* Schaer.), lichenized, widespread, terrestrial, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; position), Lücking et al. (2017a; classification), cultures and sequences are available, lectotype of type species G.

Melanocamarosporium Wijyaw., Camporesi, D.J. Bhat & K.D. Hyde 2016, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. galicola* Wijyaw., Camporesi, D.J. Bhat & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2016b, 2017; taxonomy, phylogeny, outline), holotype and ex-type of the type: MFLU 15-3451, MFLUCC 13-0545.

Melanocarpus Arx 1975, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, three species, type: *M. albomyces* (Cooney & R. Emers.) Arx, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted in *Sordariales*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), van den Brink et al. (2015; thermophilic nature), cultures and sequences are available.

Melanocephala S. Hughes 1979, *Ascomycota* genera incertae sedis, five species, type: *M. cupulifera* S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Melanoctona Qing Tian, Doilom & K.D. Hyde 2016, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *M. tectonae* Qing Tian, Doilom & K.D. Hyde, asexual morph hyphomycetous, Thailand, see Tian

et al. (2016b; taxonomy), cultures and sequences are available.

Melanodothis R.H. Arnold 1972 (= *Hyalodothis* Pat. & Har. 1893 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *M. caricis* R.H. Arnold, asexual morph ramularia-like, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Mycosphaerellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Melanographa Müll. Arg. 1882 (= *Mycomelaspilea* Reinke 1895 *fide* Ertz and Diederich 2015), *Asterinales* genera *incertae sedis*, *Dothideomycetes*, type: *M. tribuloides* (Tuck.) Müll. Arg., lichenicolous, cosmopolitan, see Ertz and Diederich (2015; taxonomy, phylogeny), sequences are available.

Melanographium Sacc. 1913 (= *Pseudocamptoum* Gonz. Frag. & Cif. 1925; = *Sporostachys* Sacc. 1919 *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, nine species, type: *M. spleniosporum* Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Melanohalea O. Blanco, A. Crespo, Divakar, Essl., D. Hawksw. & Lumbsch 2004, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 19 species, type: *M. exasperata* (De Not.) O. Blanco, A. Crespo, Divakar, Essl., D. Hawksw. & Lumbsch (syn. *Parmelia exasperata* De Not.), lichenized, widespread, terrestrial, see Jaklitsch et al. (2016a; position), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type species RO.

Melanolecia Hertel 1981, *Hymeneliaceae*, *Hymeniales*, *Lecanoromycetes*, eight species, type: *M. transitoria* (Arnold) Hertel, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available, cultures available for the type: Meltra167.

Melanomma Nitschke ex Fuckel 1870 (= *Moriolopis* Norman ex Keissl. 1927 *fide* Species Fungorum 2017), *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, c. 30 species, type: *M. pulvis-pyrius* (Pers.) Fuckel, asexual morph unknown, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Melanommataceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Tian et al. (2015; taxonomy and phylogeny), Tibpromma et al. (2017b; new data), cultures available for type: CBS 371.75, CBS 109.77, CBS 124080, CBS 371.75.

Melanopeziza Velen. 1939, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *M. artemisiae* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, dried culture is available PRM 152097.

Melanophloea P. James & Vězda 1971, *Ascomycota* genera *incertae sedis*, one species, type: *M. pacifica* P. James & Vězda, asexual morph unknown, lichenized, terrestrial, tropical South-East Asia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Thelocarpaceae*), Aptroot and Schumm (2012b; evolution), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Melanophoma Papendorf & J.W. du Toit 1967, *Ascomycota* genera *incertae sedis*, one species, type: *M. karroo* Papendorf & G.W. du Toit, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Wijayawardene et al. (2012, 2017; outline), cultures are available ex-type CBS 589.66, ATCC 18417, PRE43876.

Melanops Nitschke ex Fuckel 1870, *Melanopsaceae*, *Botryosphaeriales*, *Dothideomycetes*, one species, type: *M. tulasnei* Fuckel, asexual morph coelomycetous, saprobes, terrestrial, cosmopolitan, see Phillips and Alves (2009; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Slippers et al. (2013; phylogeny, introduced of a new family *Melanopsaceae*), Wijayawardene et al. (2014c; outline, phylogeny), epitype and ex-epitype of the type: LISE 95179, CBS 116805.

Melanopsamma Niessl 1876, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, c. 40 species, type: *M. pomiformis* (Pers.) Sacc., asexual morph custingophora-like, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Niessliaceae*), Kirk et al. (2013; genus accepted), Wang et al. (2015e; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Lombard et al. (2016; phylogeny), cultures available for type: CBS 114119, CBS 325.50, CBS 114119, CBS 101322.

Melanormia Körb. 1865, *Leotiomycetes* genera *incertae sedis*, one species, type: *M. velutina* Körb., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Melanospora Corda 1836 (= *Gonatobotrys* Corda 1839 *fide* Réblová et al. 2016c; see Species Fungorum 2017 for additional synonyms), *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, 35 species, type: *M. zamiae* Corda, asexual morphs formerly named in *Gonatobotrys*, saprobes, from soil, fungicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010;

outline), Fan et al. (2012a; species from China), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures and sequences are available.

Melanotopelia Lumbsch & Mangold 2008, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *M. toensbergii* (Vězda & Kantvilas) Lumbsch & Mangold, asexual morph unknown, lichenized, terrestrial, worldwide, see Mangold et al. (2008a; taxonomy), Séruaux et al. (2009a; new species, Rwanda), Lumbsch and Huhndorf (2010; outline), Lumbsch et al. (2010a, 2014a; new species, Falkland Island, accepted in *Graphidaceae*), Rivas Plata et al. (2012; accepted in *Graphidaceae*), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type species, holotype of the type species BG, Tønsberg 25545 (holotype of *Topeliopsis toensbergii* Kantvilas & Vězda).

Melanotrema Frisch 2006, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, eleven species, type: *M. platystomum* (Mont.) Frisch, asexual morph unknown, lichenized, terrestrial, worldwide, see Mangold et al. (2009; new species, Australia), Lumbsch and Huhndorf (2010; outline), Papong et al. (2010; new species), Rivas Plata et al. (2013; DNA, phylogeny), Kraichak et al. (2014; new species), Sipman (2014; new species), Lumbsch et al. (2014b; accepted in *Graphidaceae*), Lücking et al. (2017a; classification), cultures and sequences are available, cultures available for the type: DNA3492.

Melanotrigonum Réblová 2015, *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, one species, type: *M. ovale* Réblová, asexual morph hyphomycetous, saprobes, terrestrial, Czech Republic, see Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2017; taxonomy), holotype and ex-type: PRM 933852, CBS 138743.

Melarthonitis Frisch & G. Thor 2014, *Chrysotrichaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *M. piceae* Frisch & G. Thor, asexual morph unknown, saprobes, terrestrial, see Frisch et al. (2014; taxonomy), Lücking et al. (2017a; classification), holotype of the type: UPS, Thor 25995, sequences are available.

Melaspilea Nyl. 1857 (= *Eremithallus* Lücking, Lumbsch & L. Umaña 2008 *fide* Ertz and Diederich 2015; see Species Fungorum 2017 for additional synonyms), *Melaspileaceae*, *Eremithallales*, *Dothideomycetes*, 163 species, type: *M. arthonioides* (A. Massal.) Nyl., asexual morph unknown, saprobes, marine and terrestrial, worldwide, see Jones et al. (2009b, 2015; classification), Pérez-Ortega and Etayo (2010; morphology), Zainuddin et al. (2010; antimicrobial activities), Zhurbenko and Zhdanov (2013; morphology), Flakus et al. (2014; morphology), Ertz and Diederich

(2015; phylogeny), Perlmutter et al. (2015; morphology), Lücking et al. (2017a; classification), sequences are available but lacking for the type.

Melaspilella (P. Karst.) Vain. 1921 (= *Banhegyia* L. Zeller & Tóth 1969 *fide* Ertz and Diederich 2015), *Asterinales* genera incertae sedis, *Dothideomycetes*, type: *M. proximella* (Nyl.) Ertz & Diederich, saprobes, Europe, see Ertz and Diederich (2015; taxonomy, phylogeny, designate the lectotype), sequences are available.

Melaspileopsis (Müll. Arg.) Ertz & Diederich 2015, *Melaspileaceae*, *Eremithallales*, *Dothideomycetes*, one species, type: *M. diplasiospora* (Nyl.) Ertz & Diederich, sexual morph with lirelliform ascocarps, saprobes, terrestrial, tropical and subtropical, see Ertz and Diederich (2015; morphology, phylogeny), Ertz et al. (2016b; phylogeny), sequences are available, lectotype of the type: Lindig 846 (PC0146198).

Melastiza Boud. 1885, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 14 species, type: *M. miniata* (Fuckel) Boud., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Melchioria Penz. & Sacc 1897 (= *Exomassarinula* Teng 1940; = *Gaeumannia* Petr. 1950 *fide* Species Fungorum 2017), *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, six species, type: *M. leucomelaena* Penz. & Sacc., asexual morph unknown, terrestrial, saprobes, Indonesia and South America, cultures and sequences are unavailable.

Melinomyces Hambl. & Sigler 2005, *Hyaloscyphaceae*, *Helotiidae*, *Leotiomycetes*, three species, type: *M. variabilis* Hambl. & Sigler, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are available, holotype of the type: UAMH 8861.

Meliola Fr. 1825 (= *Arberia* Nieuwl. 1916; = *Asteridium* (Sacc.) Speg. ex Sacc. 1891; = *Asterina* subgen. *Asteridium* Sacc. 1882; = *Chaetomeliola* (Cif.) Bat., H. Maia & M.L. Farr 1962; = *Meliola* subgen. *Chaetomeliola* Cif. 1938; = *Laeviomeliola* Bat. 1960; = *Meliothecium* Mussat 1901; = *Myxothecium* Kunze 1829 *fide* Hongsanan et al. 2015d; Species Fungorum 2017), *Meliolaceae*, *Meliolales*, *Sordariomycetes*, c. 1300, type: *M. nidulans* (Schwein.) Cooke, asexual morph produced from the hyphae, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d, 2016c; taxonomy, evolution), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Meliolaster Höhn. 1918 (= *Meliolinopsis* Beeli 1920; = *Patouillardina* G. Arnaud 1917 *fide* Species Fungorum

2017), Asterinaceae, Asterinales, Dothideomycetes, three species, type: *M. clavisporus* (Pat.) Höhn., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in Asterinaceae), Wijayawardene et al. (2014c; outline), Hongsanan et al. (2014c; taxonomy, keys), cultures and sequences unavailable, holotype of type: FH 258798.

Meliolina Syd. & P. Syd. 1914, Meliolinaceae, Dothideomycetes, families incertae sedis, c. 40 species, type: *M. cladotricha* (Lév.) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; listed the family under excluded, doubtful and poorly known families), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Melioliphila Speg. 1921 (= *Subiculicola* Speg. 1921 fide Species Fungorum 2017), Dothideomycetes genera incertae sedis, seven species, type: *M. graminicola* (F. Stevens) Speg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted in Tubeufiaceae), Hyde et al. (2013; excluded from Tubeufiaceae), Boonmee et al. (2014b; excluded from Tubeufiaceae), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Melliosporiella Höhn. 1919, Marthamycetaceae, Rhytismatales, Leotiomycetes, two species, type: *M. pulchella* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), isotype of the type: CUP- D-12032, sequences are unavailable.

Melliosporium Corda 1838 (= *Pleistictis* Rehm 1882 fide Species Fungorum 2017), Marthamycetaceae, Rhytismatales, Leotiomycetes, ?eleven species, type: *M. versicolor* Corda, asexual morph unknown, saprobes, terrestrial, America, Europe, New Zealand, see Lumbsch and Huhndorf (2010; outline, accepted as Rhytismatales, genera incertae sedis), Lantz et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Melnikia Wijyaw., Goonas., D.J. Bhat & K.D. Hyde, 2016, Phaeosphaeriaceae, Pleosporales, Dothideomycetes, one species, type: *M. anthoxanthii* Wijyaw., Goonas., Camporesi, D.J. Bhat & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2016b, 2017; taxonomy, phylogeny, outline), holotype and ex-type of the type: MFLU 15-2657, MFLUCC 14-1010.

Melnikomyces Crous & U. Braun 2014, Chaetothyriales genera incertae sedis, Eurotiomycetes, one species, type:

M. vietnamensis Crous & U. Braun, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Vietnam, see Crous et al. (2014c; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type strain of type: CBS H-21715, CPC 23554.

Meloderma Darker 1967, Rhytismataceae, Rhytismatales, Leotiomycetes, nine species, type: *M. desmazieri* (Duby) Darker, asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lantz et al. (2011; DNA, phylogeny), cultures and sequences are available.

Melogramma Fr. 1849 (= *Dothiora* Fr. 1836 fide Species Fungorum 2017), Melogrammataceae, Amphisphaerales, Sordariomycetes, three species, type: *M. campylosporum* Fr., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline, accepted in Melanconidaceae), Jaklitsch and Voglmayr. (2012; phylogeny, accepted in Melogrammataceae), Kirk et al. (2013; genus accepted), Senanayake et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2016b; outline, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Melomastia Nitschke ex Sacc. 1875, Sordariomycetes genera incertae sedis, four species, type: *M. friesii* Nitschke, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Melophia Sacc. 1884, Ascomycota genera incertae sedis, four species, type: *M. ophiospora* (Lév.) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Memnoniella Höhn. 1923 (= *Fuckelina* Sacc. 1875; = *Gliobotrys* Höhn. 1902; = *Hyalobotrys* Pidopl. 1948; = *Hyalostachybotrys* Sriniv. 1958; = *Memnoniella* Höhn. 1923 [1924]; = *Ornatispora* K.D. Hyde, Goh, Joanne E. Taylor & J. Fröh. 1999; = *Synsporium* Preuss 1849 fide Species Fungorum 2017), Stachybotryaceae, Hypocreales, Sordariomycetes, 13 species, type: *M. echinata* (Riv.) Galloway hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011; synonym of *Stachybotrys*), Wang et al. (2015e; synonym of *Stachybotrys*), Lombard et al. (2016; reinstated the genus, phylogeny), Lin et al. (2016b; new species), Wijayawardene et al. (2017; outline), cultures and sequences are available, epitype and ex-epitype culture of type species CBS H-22447, CBS 216.32.

Mendogia Racib. 1900 (= *Pleistostomella* Syd. & P. Syd. 1917; = *Uleopeltis* Henn. 1904 fide Species Fungorum

2017), *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, four species, type: *M. bambusina* Racib., asexual morph unknown, epiphytic, terrestrial, Indonesia, Philippine, Thailand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Schizophyriaceae*), Wijayawardene et al. (2014c; outline), Dai et al. (2017, description, new species), cultures and sequences are unavailable.

Mendoziopeltis Bat. 1959, *Dothideomycetes* genera *incertae sedis*, four species, type: *M. byrsinimae* Bat. & Nascim. ex Bat., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as *Micropeltidaceae*), Hyde et al. (2013; excluded from *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus *Dothideomycetes*), cultures and sequences are unavailable.

Menegazzia A. Massal. 1854, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 70 species, type: *M. terebrata* (Hoffm.) A. Massal., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Menidochium R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, one species, type: *M. formosum* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Menispora Pers. 1822 (= *Ciliofusa* Clem. & Shear 1931; = *Ciliofusarium* Rostr. 1892; = *Eriomene* (Sacc.) Clem. & Shear 1931; = *Eriomenella* Peyronel 1918; = *Erionema* Maire 1906; = *Menispora* subgen. *Eriomene* Sacc. 1886; = *Zignoëlla* Sacc. 1878 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, 14 species, type: *M. glauca* (Link) Pers., hyphomycetous, sexual morph formerly named in *Zignoëlla*, saprobes, terrestrial, temperate, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures and sequences are unavailable.

Menisporopsis S. Hughes 1952 (= *Menisporopascus* Matsush. 2003 *fide* Réblová et al. 2016c), *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, nine species, type: *M. theobromae* S. Hughes, hyphomycetous, sexual morph formerly named in *Menisporopascus*, saprobes, terrestrial, temperate, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted),

Maharachchikumbura et al. (2015b, 2016b; outline), Liu et al. (2016a; phylogeny), Réblová et al. (2016c; nomenclature), cultures and sequences are available, available specimen of type: CBS H-19592.

Mercadomyces J. Mena 1988, *Ascomycota* genera *incertae sedis*, one species, type: *M. camaguevensis* J. Mena, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Meridianelia Kantvilas & Lumbsch 2009, *Elixiaceae*, *Umbilicariales*, *Lecanoromycetes*, one species, type: *M. maccarthyana* Kantvilas & Lumbsch, lichenized, corticolous, Tasmania, see Kantvilas and Lumbsch (2009; taxonomy), Bendiksby and Timdal (2013; phylogeny), Miądlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Merismatium Zopf 1898 (= *Heterophracta* (Sacc. & D. Sacc.) Theiss. & Syd. 1918; = *Leptosphaerulina* subgen. *Heterophracta* Sacc. & D. Sacc. 1905; = *Phaeocyrtis* Vain. 1921 *fide* Species Fungorum 2017), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, eleven species, type: *M. lopadii* (Anzi) Zopf, asexual morph unknown, lichenicolous, terrestrial, Europe, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Yazici and Aslan (2016; Asia, Turkey), cultures and sequences are unavailable.

Merismella Syd. 1927, *Ascomycota* genera *incertae sedis*, six species, type: *M. concinna* Syd., sexual morph *Chaetothyrium* Speg., epiphytic, terrestrial, Brazil, Greece, India, Panama, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Meristemomyces Isola & Onofri 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *M. frigidus* Isola & Onofri, hyphomycetous, sexual morph unknown, rock and plant inhabiting fungus, Italy, America, see Egidi et al. (2014; taxonomy), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2017; outline), Crous et al. (2016b; new species), type: CBS 136109 = CCFFEE 5508, cultures and sequences available.

Merodontis Clem. 1909 (= *Davincia* subgen. *Davinciella* Sacc. & D. Sacc. 1906; = *Davinciella* (Sacc. & D. Sacc.) Trotter 1928 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *M. tenella* (Penz. & Sacc.) Clem., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Merugia Rogerson & Samuels 1990, *Sordariomycetidae* genera *incertae sedis*, seven species, type: *M. palicoureae* Rogerson & Samuels, saprobes, terrestrial, Guyana, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of the type: NY, Samuels 6552.

Mesniera Sacc. & P. Syd. 1902, *Mesnieraceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *M. rottlerae* (Racib.) Sacc. & P. Syd., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, available specimen of type: CBS H-4212.

Metacapnodium Speg. 1918 (= *Antennularia* Rchb. 1828; = *Capnocybe* S. Hughes 1966; = *Capnophialophora* S. Hughes 1966; *Capnosporium* S. Hughes 1976; = *Ophiocapnocoma* Bat. & Cif. 1963; = *Sthughesia* M.E. Barr 1987 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Metacapnodiaceae*, *Capnodiales*, *Dothideomycetes*, 13 species, type: *M. juniperi* (W. Phillips & Plowr.) Speg., asexual morph unknown, sooty mould on living plants, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hughes et al. (2012; morphology), Hyde et al. (2013; accepted as a genus in *Metacapnodiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Metacoleroa Petr. 1927, *Venturiaceae*, *Venturiales*, *Dothideomycetes*, one species, type: *M. dickiei* (Berk. & Broome) Petr., asexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2008; treated as a synonym of *Venturia* Sacc.), Lumbsch and Huhndorf (2010; outline), Raja et al. (2011a; DNA), Zhang et al. (2011d; morphology, phylogeny), Hyde et al. (2013; accepted as in *Venturiaceae*, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Metacordyceps G.H. Sung, J.M. Sung, Hywel-Jones & Spatafora 2007, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, 16 species, type: *M. taiii* (Z.Q. Liang & A.Y. Liu) G.H. Sung, J.M. Sung, Hywel-Jones & Spatafora, asexual morph hyphomycetous, terrestrial, entomogenous and in soil and dung, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kepler et al. (2012a; sexual morph), Maharachchikumbura et al. (2015b, 2016b; outline), Wen et al. (2015; new species), cultures and sequences are available.

Metadiplodia Syd. 1937, *Ascomycota* genera *incertae sedis*, c. 40 species, type: *M. eucalypti* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Metadothella Henn. 1904, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *M. stellata* Henn., asexual morph hyphomycetous, terrestrial, Peru, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; phylogeny), cultures and sequences are unavailable, holotype and ex-type strain not available.

Metahyphopichia Sipiczki & Pfleigler 2016, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *M. laotica* Sipiczki, Pfleigler, Safar, Morais, & Rosa, dimorphic, asexual morph unknown, on plants, terrestrial, Southeast Asia, South America, see Sipiczki et al. (2016; taxonomy), cultures available for type: CBS 13022, CCY 092-001-001, NCAIM Y.02126, sequences are available.

Metamelanea Henssen 1989, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, three species, type: *M. umbonata* Henssen, lichenized, asexual morph unknown, terrestrial, Europe, North America, see Kirk et al. (2013; genus accepted), Prieto et al. (2015; records in Sweden), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Metameris Theiss. & Syd. 1915 (= *Scirrhodothis* Theiss. & Syd. 1915; = *Scirrhophragma* Theiss. & Syd. 1915 *fide* Species Fungorum 2017), *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, three species, type: *M. japonica* (Syd. & P. Syd.) Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (accepted as a genus in *Phaeosphaeriaceae*), Phookamsak et al. (2014c; placed in *Botryosphaeriaceae*), Wijayawardene et al. (2014c; outline, accepted in *Botryosphaeriaceae*), cultures and sequences are unavailable.

Metapezizella Petr. 1968, *Leotiomycetes* genera *incertae sedis*, one species, type: *M. phyllachorivora* Petr., asexual morph unknown, saprobes, terrestrial, Mexico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Metapochonia Kepler, S.A. Rehner & Humber 2014, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *M. suchlasporia* (W. Gams & Dackman) Kepler, Rehner & Humber, asexual morph hyphomycetous, terrestrial, nematogenous and in soil, rotifers and plant roots, cosmopolitan, see Kepler et al. (2014; taxonomy), cultures and sequences are available, Holotype and ex-type of type species in Suecia, C. Dackman No. 10 (1983); CBS 251.83.

Metarhiziopsis D.W. Li, R.S. Cowles, Vossbrinck 2008, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. microspora* D.W. Li, R.S. Cowles & Vossbrinck, hyphomycetous, sexual morph unknown, terrestrial, entomogenous, USA, see Li et al. (2008b; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, holotype and ex-type of type: BPI 878276 = UAMH 10901.

Metarhizium Sorokin 1879 (= *Chromostylium* Giard 1889 *fide* Species Fungorum 2017), *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, 24 species, type: *M. anisopliae* (Metschn.) Sorokin, hyphomycetous, sexual morph unknown, terrestrial, entomogenous and in soil, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kepler et al. (2014; phylogeny), Chen et al. (2017c; new species, phylogeny), cultures and sequences are available.

Metathyriella Syd. 1927 (= *Ciferriotheca* Bat. & I.H. Lima 1959; = *Stomiopeltina* Bat. 1963 *fide* Species Fungorum 2017), *Schizophyriaceae*, *Dothideomycetes* families *incertae sedis*, three species, type: *M. roupalae* Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013, accepted as a genus in *Schizophyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Phookamsak et al. (2016; taxonomy), cultures and sequences are unavailable.

Metazythia Petr. 1950, *Ascomycota* genera *incertae sedis*, one species, type: *M. caespitosa* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Metazythiopsis M. Morelet 1988, *Ascomycota* genera *incertae sedis*, one species, type: *M. halepensis* M. Morelet, coelomycetous, sexual morph unknown, saprobes, terrestrial, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Metschnikowia Kamienski 1899 (= *Anthomyces* Grüss 1918; = *Chlamydozyma* Wick. 1964; = *Metschnikowiella* Henckel 1913; = *Monospora* Metschn. 1884; = *Monosporella* Keilin 1920; = *Nectaromyces* Syd. & P. Syd. 1919 [1918] *fide* Species Fungorum 2017), *Metschnikowiaceae*, *Saccharomycetales*, *Saccharomycetes*, c. 100 species (including c. 50 *Candida* species in the *Metschnikowia* clade), type: *M. bicuspidata* (Metschnikoff) T. Kamienski, asexual reproduction is by multilateral budding, saprophytic, on beetles, fruit flies, rotted plants, berries, water, seawater, flowers of various plant species, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Crous et al. (2015d; new species, phylogeny), asexual species in this clade are classified in the genus *Candida*, but will be transferred to *Metschnikowia* as new combinations, sequences are available.

Metulocladosporiella Crous, Schroers, J.Z. Groenew., U. Braun & K. Schub. 2006, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, two species, type: *M. musae* (E.W. Mason) Crous, Schroers, J.Z. Groenew., U. Braun & K. Schub., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Neotropics, see Seifert et al. (2011; morphology), Xu et al. (2011; phylogeny), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Metus D.J. Galloway & P. James 1987, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *M. conglomeratus* (F. Wilson) D.J. Galloway & P. James, asexual morph unknown, lichenized, terrestrial, Australasia, see Lumbsch et al. (2010b; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; DNA, phylogeny), Lücking et al. (2017a, classification), cultures and sequences are available, cultures available for the type: AFTOL-ID 4945.

Meyerozyma Kurtzman & M. Suzuki 2010, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, five species, type: *M. guilliermondii* (Wickerham) Kurtzman & M. Suzuki, asexual reproduction is by multilateral budding, saprophytic, but *M. guilliermondii* is a common clinical species, atmosphere, soil, water, sugar cane and other plants, plant debris, fruit, mammals, worldwide, see Kurtzman and Suzuki (2010; taxonomy), Bautista-Rosales et al. (2013; bio control agent of *Colletotrichum* sp.), Coda et al. (2013; antifungal properties), sequences are available.

Micarea Fr. [nom. cons.] 1825 (= *Hastifera* D. Hawksw. & Poelt 1986; = *Helocarpon* Th. Fr. 1860; = *Stereocauliscum* Nyl. 1865 *fide* Species Fungorum 2017), *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, 100 species, type: *M. prasina* Fr., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Micareopsis R.C. Harris & Lendemer 2013, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *M. irriguata* R.C. Harris & Lendemer, asexual morph unknown, lichenized, terrestrial, USA, see Lendemer et al. (2013; taxonomy), cultures and sequences are unavailable, holotype of the type species NY, Buck 49935.

Micraspis Darker 1963, *Tympanidaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *M. acicola* Darker, asexual morph unknown, saprobes, North America, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk

et al. (2013; genus accepted), Index Fungorum (2015; new family), cultures and sequences are unavailable.

Microascus Zukal 1886 (= *Fairmania* Sacc. 1906; = *Nephrospora* Loubière 1923; = *Peristomium* Lechmere 1912 *fide* Species Fungorum 2017), *Microascaceae*, *Microasciales*, *Sordariomycetes*, 51 species, type: *M. longirostris* Zukal, asexual morph hyphomycetous, coprophilous, on human, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Sandoval-Denis et al. (2016a; new species, phylogeny), Crous et al. (2016b; new species), Jagielski et al. (2016; new species, phylogeny), neotype and ex-neotype of type: CBS H-14440, MUCL 9058.

Microblastosporon Cif., 1930, *Ascomycota* genera *incertae sedis*, one species, type: *M. botryoideum* (Chaborski) Cif., hyphomycetous, saprobes or plant pathogenic, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microcalicium Vain. 1927 (= *Coniocybopsis* Vain. 1927; = *Strongylopsis* Vain. 1927 *fide* Species Fungorum 2017), *Microcaliciaceae*, *Ascomycota* families *Incatae sedis*, four species, type: *M. disseminatum* (Ach.) Vain., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Prieto et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), voucher for the type species (sequence data also deposited under this number) Wedin 6353 (Prieto et al. 2012).

Microcallis Syd. 1926 (= *Barnettia* Bat. & J.L. Bezerra 1962; = *Fraseria* Bat. 1962; = *Hansfordina* Bat. 1962; = *Kanousea* Bat. & Cif. 1962; = *Microcalliomycetes* Bat. & Cif. 1962; = *Microcalliopsis* Bat. & Cif. 1962; = *Neopeltis* Petr. 1949; = *Recifea* Bat. & Cif. 1962 *fide* Species Fungorum 2017), *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, ten species, type: *M. phoebe* Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microcera Desm. 1848, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, nine species, type: *M. coccophila* Desm., asexual morph hyphomycetous, terrestrial, entomogenous, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Gräfenhan et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, lectotype of type: K(M) 165807.

Microclava F. Stevens 1917 (= *Ommatospora* Bat. & Cavalc. 1964 *fide* Species Fungorum 2017), *Ascomycota*

genera *incertae sedis*, six species, type: *M. miconiae* F. Stevens, hyphomycetous, saprobes or plant pathogenic, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microcyclella Theiss. 1914, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. nervisequia* (Höhn.) Theiss., asexual morph unknown, saprobes, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Microcyclephaeria Bat. 1958, *Hypocreales*, genera *incertae sedis*, one species, type: *M. palmicola* (Syd.) Bat. & H. Maia ex Bat., asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariomycetes* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Microcyclospora J. Frank, Schroers & Crous 2010, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, seven species, type: *M. pomicola* J. Frank, B. Oertel, Schroers & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Canada, Iran, see Frank et al. (2010; taxonomy, phylogeny), Arzanlou and Bakhshi (2011; new species), Seifert et al. (2011; morphology), Crous et al. (2012c; new species, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Microcyclosporella J. Frank, Schroers & Crous 2010, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *M. mali* J. Frank, Schroers & Crous, hyphomycetous, sexual morph unknown, plant pathogenic, epiphytic, Slovenia, see Frank et al. (2010; morphology, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, holotype and ex-type of type: CBS H-20413; CBS 126136.

Microcyclus Sacc., Syd. & P. Syd. 1904 (= *Diplochora* Syd. & P. Syd. 1913; = *Diplochorella* Syd. & P. Syd. 1913; = *Lizoniella* Henn. ex Sacc. & D. Sacc. 1905; = *Melanopsammopsis* Stahel 1915; = *Pseudosphaerella* Höhn. 1911 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, c. 15 species, type: *M. angolensis* Sacc., Syd. & P. Syd., asexual morph unknown, pathogen, saprobes, endophytes, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable for type species.

Microdiplodia Allesch. 1901 (= *Microbotryodiplodia* Sousa da Câmara 1951; = *Syndiplodia* Peyronel 1915 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae*

sedis, 30 species, type: *M. conigena* Allesch., sexual morph unknown, saprobes, terrestrial, Europe, Canada, Iran, see Wijayawardene et al. (2016b; outline, taxonomic notes and treated as doubtful), cultures and sequences are available, need generic revision and epitypification.

Microdiscula Höhn. 1915, *Ascomycota* genera *incertae sedis*, two species, type: *M. rubicola* (Bres.) Höhn., coelomycetous, sexual morph unknown, saprobes or plant pathogenic, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted); cultures and sequences are unavailable.

Microdiscus Sacc. 1916 (= *Brachyascus* Syd. & P. Syd. 1917; = *Microascus* Sacc. 1916 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *M. americanus* (Sacc.) Trotter, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microdochella M. Hern.-Restr. & Crous 2016, *Orbiliiales*, genera *incertae sedis*, *Orbiliomycetes*, one species, type: *M. fusariooides* (D.C. Harris) M. Hern.-Restr. & Crous, hyphomycetous, sexual morph unknown, fungicolous, see Hernández-Restrepo et al. (2016a, b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: IMI 281715, CBS 741.83.

Microdochium Syd. & P. Syd. 1924 (= *Gerlachia* W. Gams & E. Müll. 1980; = *Griphosphaerella* Petr. 1927; = *Monographella* Petr. 1924 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Microdochiaeaceae*, *Xylariales*, *Sordariomycetes*, 25 species, type: *M. phragmitis* Syd., hyphomycetous, sexual morph formerly named in *Monographella*, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Hernández-Restrepo et al. (2016b; phylogeny, introduced *Microdochiaeaceae*), Réblová et al. (2016c; nomenclature), holotype and epitype of the type: K-IMI 193888, CBS H-22135, ex-epitype culture CBS 285.71.

Microdothella Syd. & P. Syd. 1914 (= *Phaeopolystomella* Bat. & H. Maia 1960 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, two species, type: *M. culmicola* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, North America, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Microdothiorella C.A.A. Costa & Sousa da Câmara 1955, *Ascomycota* genera *incertae sedis*, one species, type: *M. sphaeralceae* M.E.A. Costa & Sousa da Câmara, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Portugal, see Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microeurotium Ghatak 1936, *Pezizales* genera *incertae sedis*, *Pezizomyces*, two species, type: *M. albidum* Ghatak, asexual morph unknown, saprobes or coprophilous, terrestrial, UK, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microglossum Gillet 1879 (= *Geoglossum* subgen. *Leptoglossum* Cooke 1879; = *Helote* Hazsl. 1881; = *Leptoglossum* (Cooke) Sacc. 1884; = *Leptoglossum* sect. *Xanthoglossum* Sacc. 1889; = *Microglossum* Gillet 1908; = *Ochroglossum* S. Imai 1955; = *Xanthoglossum* (Sacc.) Kuntze 1891 *fide* Species Fungorum 2017), *Leotiaceae*, *Helotiales*, *Leotiomycetes*, ten species, type: *M. olivaceum* (Pers.) Gillet, asexual morph unknown, saprobes, on soil, terrestrial, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kučera et al. (2014; new species, phylogeny), Pärte et al. (2017; phylogeny), cultures and sequences are available for the type species SAV 9902 (Kučera et al. 2014) and KL220.

Microhendersonula Dias & Sousa da Câmara 1952, *Ascomycota* genera *incertae sedis*, one species, type: *M. cestri* Dias & Sousa da Câmara, coelomycetous, saprobes or plant pathogenic, Portugal, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microhilum H.Y. Yip & A.C. Rath 1989, *Cordycitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. oncoperae* H.Y. Yip & A.C. Rath, hyphomycetous, terrestrial, entomogenous, Australia, see Hyde et al. (2011; checklist), Wijayawardene et al. (2012; checklist), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; *Sordariomycetes*), cultures and sequences are available, holotype of type species DATH 64D6.

Microidium (To-anun & S. Takam.) To-anun & S. Takam. 2012 (= *Oidium* subgen. *Microidium* To-anun & S. Takam. 2005 *fide* Species Fungorum 2017), *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, four species, type: *M. phyllanthi* (J.M. Yen) To-anun & S. Takam. (≡ *Oidium phyllanthi* J.M. Yen), hyphomycetous, sexual morphs unknown, biotrophic, obligate plant pathogens, terrestrial, Asia, South America, see Braun and Cook (2012; taxonomy), Tam et al. (2015; Vietnam), sequences are available.

Micromastia Speg. 1909, *Ascomycota* genera *incertae sedis*, two species, type: *M. trigonospora* Speg., asexual morph unknown, saprobes, on soil, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Micronectria Speg. 1885, *Hyponectriaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *M. guaranitica* Speg., asexual morph unknown, saprobes, see Kirk et al. (2008; treated as nom. dub.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Micronematobotrys Xiang Sun & L.D. Guo 2010, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 14 species, type: *M. verrucosus* Xiang Sun & L.D. Guo, hyphomycetous, sexual morph unknown, saprobes, terrestrial, China, see Sun and Guo (2010; taxonomy, phylogeny), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, holotype and cultures available for type species HMAS 186049; E13-2151.

Micropeltis Mont. 1842 (= *Dictyothyriella* Rehm 1914; = *Hormopeltis* Speg. 1912; = *Micropeltella* Syd. & P. Syd. 1913; = *Micropeltidium* Speg. 1919 [1918]; = *Ophiopeltis* J.V. Almeida & Sousa da Câmara 1903; = *Parapeltella* Speg. 1919 [1918]; = *Scolecopeltella* Speg. 1921; = *Scolecopeptopsis* Höhn. 1909; = *Theciopeltis* F. Stevens & Manter 1925 *fide* Species Fungorum 2017), *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, c. 110 species, type: *M. applanata* Mont., asexual morph unknown, epiphytes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; outline), Hongsanan et al. (2015a; new species), cultures and sequences are unavailable.

Micoperella Höhn. 1909, *Ascomycota* genera *incertae sedis*, one species, type: *M. quercus* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Micopeziza Fuckel 1870 (= *Actinoscypha* P. Karst. 1888; = *Calloriella* Höhn. 1918; = *Crustomollisia* Svrček 1987 [1986]; = *Niesella* Höhn. 1919 [1918] *fide* Jaklitsch et al. 2016a; Species Fungorum 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, twelve species, type: *M. cornea* (Berk. & Broome) Nannf., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Micropodia Boud. 1885, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *M. pteridina* (Nyl.) Boud., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Micropustulomyces R.W. Barreto 1995, *Ascomycota* genera *incertae sedis*, one species, type: *M. mucilaginosus* R.W. Barreto, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microscypha Syd. & P. Syd. 1919 (= *Mollisiaster* Kirschst. 1939 *fide* Species Fungorum 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *M. grisella* (Rehm) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Europe, South Africa, see Huhtinen et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Microsomyses Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *M. psammoechi* Thaxt., asexual morph unknown, biotrophic, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microsphaeropsis Höhn. 1917 (= *Chaetosphaeropsis* Curzi & Barbaini 1927; = *Coniothyriopsis* Speg. 1911; = *Cryptophaeella* Höhn. 1917; = *Microsporella* Höhn. 1918; = *Phyllostictella* Tassi 1901; = *Sclerothyrium* Höhn. 1918 *fide* Species Fungorum 2017), *Microsphaeropsidaceae*, *Pleosporales*, *Dothideomycetes*, 37 species, type: *M. olivacea* (Bonord.) Höhn., coelomycetous, sexual morph unknown, saprobes, pathogens of both plants and animals, lichenicolous, terrestrial, worldwide, see Etayo and Sancho (2008; new species), Marinowitz et al. (2008; new species), Pethybridge et al. (2008; new species, Australia), de Gruyter et al. (2009; DNA, phylogeny), Etayo and Yazici (2009; new species, lichenicolous), Alves et al. (2010; new species, Costa Rica), Crous et al. (2011d; new species, pathogens, *Proteaceae*), Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), Kirk et al. (2013; genus accepted), Brackel (2014; new species), Chen et al. (2015b; taxonomy, phylogeny), cultures available for the type: CBS 116669, CBS 442.83.

Microsporum Gruby 1843 (= *Closteroleurosporia* Grigoraki 1924; = *Closterosporia* Grigoraki 1924; = *Sabouraudites* M. Ota & Langeron 1923; = *Thallomicrosporon* Benedek 1964 *fide* Species Fungorum 2017), *Arthrodermataceae*, *Onygenales*, *Eurotiomycetes*, c. 17 species, type: *M. audouinii* Gruby, hyphomycetous, sexual morph unknown, on humans and other mammals, causing microspores, terrestrial, worldwide, see Seifert et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), Hubka et al. (2014b; new species), cultures and sequences are available, culture available for the type: CBS 280.63.

Microstoma Bernstein 1852 (= *Anthopeziza* Wettst. 1885; = *Articulariella* Höhn. 1909; = *Articulis* Clem. & Shear 1931 *fide* Species Fungorum 2017), *Sarcoscyphaceae, Pezizales, Pezizomycetes*, one species, type: *M. hiemale* Bernstein, asexual morph unknown, saprobe, terrestrial, Costa Rica, Canary Islands, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microtheliopsis Müll. Arg. 1890 (= *Chaetomonodorus* Bat. & H. Maia 1961; = *Micropyrenula* Vain. 1921; = *Microtheliopsidomyces* Cif. & Tomas. 1953; = *Scutomyces* J.L. Bezerra & Cavalc. 1972 *fide* Species Fungorum 2017), *Microtheliopsidaceae, Dothideomycetes* families *incertae sedis*, four species, type: *M. uleana* Müll. Arg., asexual morph unknown, saprobe, terrestrial, Central and South America, Australasia, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Microthia Gryzenh. & M.J. Wingf. 2006, *Cryphonectriaceae, Diaporthales, Sordariomycetes*, two species, type: *M. havanensis* (Bruner) Gryzenh. & M.J. Wingf., asexual morph unknown, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and cultures available for the type: BPI 614275; CMW 10879 = CBS 115758, PREM 57522.

Microthyrium Desm. 1841 (= *Calothyriella* Höhn. 1917; = *Microthyrina* Bat. 1960; = *Phragmothyrium* Höhn. 1912 *fide* Species Fungorum 2017), *Microthyriaceae, Microthyriales, Dothideomycetes*, c. 60 species, type: *M. microscopicum* Desm., asexual morph unknown, epiphytes, terrestrial, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Wu et al. (2011, 2014a; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Microthyriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Microtyle Speg. 1919, *Ascomycota* genera *incertae sedis*, one species, type: *M. bergii* Speg., asexual morph coelomycetous, saprobes or plant pathogenic, South America, See Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and cultures and sequences are unavailable.

Microxiphium (Harv. ex Berk. & Desm.) Thüm. 1879 (= *Dennisiella* Bat. & Cif. 1962 *fide* Rossman et al. 2016a), *Coccidiiniaceae, Chaetothyriales, Eurotiomycetes*, two species, type: *M. footii* (Harv. ex Berk. & Desm.) Thüm., hyphomycetous, sexual morph formerly known in *Dennisiella*, sooty moulds, Europe, North America, South America, see Seifert et al. (2011; morphology),

Wijayawardene et al. (2017; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Microxyphiella Speg. 1918, *Ascomycota* genera *incertae sedis*, 15 species, type: *M. fuligo* (Berk. & Desm.) Speg., coelomycetous, saprobes or plant pathogenic, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Microxyphiopsis Bat. 1963, *Ascomycota* genera *incertae sedis*, two species, type: *M. byrsonimiae* Bat., coelomycetous, saprobes or plant pathogenic, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Micularia Boedijn 1961, *Myriangiaceae, Myriangiales, Dothideomycetes*, one species, type: *M. merremiae* Boedijn, asexual morph unknown, saprobes, parasitic, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline, accepted genus as in *Elsinoaceae*), Hyde et al. (2013; accepted in *Myriangiaceae*), Dissanayake et al. (2014; accepted in *Myriangiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Middlehovenomyces Kurtzman & Robnett 2014, *?Trichomonascaceae, Saccharomycetales, Saccharomycetes*, two species, type: *M. tepae* (J. Grinbergs) C.P. Kurtzman & C.J. Robnett, asexual reproduction is by multilateral budding and from formation of pseudohyphae, saprophytes, rotted wood, Chile, see Kurtzman and Robnett (2014; taxonomy), sequences are available.

Midotiopsis Henn. 1902, *Cordieritidaceae, Helotiales, Leotiomycetes*, two species, type: *M. bambusicola* Henn., asexual morph unknown, saprobes, terrestrial, America (tropical), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Miladina Svrček 1972 (= *Actinospora* Descals et al. 1999 *fide* Healy et al. 2016), *Pyronemataceae, Pezizales, Pezizomycetes*, two species, type: *M. lecithina* (Cooke) Svrček, asexual morph formerly named in *Actinospora*, saprobes, hypogeous, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), sequences are available for the type species from voucher KH.04.22 (Hansen et al. 2013).

Millerburtonia Cif. 1951, *Gnomoniaceae, Diaporthales, Sordariomycetes*, one species, type: *M. oyedaeae* Cif., coelomycetes, sexual morph unknown, saprobes, terrestrial, Venezuela, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Millerozyma Kurtzman & M. Suzuki 2010, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, five species, type: *M. farinosa* (Lindner) Kurtzman & M. Suzuki, asexual reproduction is by multilateral budding, saprophytic, on beetle larvae, food contaminants, dung, worldwide, see Kurtzman and Suzuki (2010; taxonomy), sequences are available.

Milosporium D. Hawksw. 1975 (= *Spilomium* Nyl. 1856 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, three species, type: *M. graphideorum* (Nyl.) D. Hawksw., lichenicolous, terrestrial, sexual morph unknown, Europe, see Seifert et al. (2011; genus accepted), Kirk et al. (2013; genus accepted), Crous et al. (2015a; phylogeny), cultures and sequences are available, conserved type and epitype of type species BM (Herb. Lich. Paris 72), CBS H-22271.

Miltidea Stirt. 1898, *Miltideaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type: *L. rubricatula* Stirt., lichenized, Southern Hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Mimeomyces Thaxt. 1912, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 16 species, type: *M. decipiens* Thaxt., asexual morph unknown, parasites, worldwide, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species), Kirk et al. (2013; genus accepted), sequences and culture unavailable.

Mindoia Petr. 1949, *Ascomycota* genera *incertae sedis*, two species, type: *M. ingae* Petr., coelomycetous, saprobes or plant pathogenic, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Miniancora Marvanová & Bärl. 1989, *Leotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *M. allisonensis* Marvanová and Bärl., hyphomycetous, sexual morph unknown, saprobes, freshwater, Canada, see Wijayawardene et al. (2012, 2017; outline), sequence is available, ex-type strain CCM F-30487.

Minimelanolocus R.F. Castañeda & Heredia 2001, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, 26 species, type: *M. navicularis* (R.F. Castañeda) R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, aquatic, China, Portugal, Mexico, Cuba, see Ma et al. (2011a, b; China; new species), Hernández-Restrepo et al. (2012a; Portugal), Heredia et al. (2014; Mexico), Liu et al. (2015a, c; phylogeny), Hyde et al. (2016; new species), Tian et al. (2016b; new species), sequence available but lacking for the type species, holotype of the type: INIFAT C86/157, compare with *Pseudospiropes*, *Spiropes* and *Sympodioplanus*.

Minimidochium B. Sutton 1970 (= *Neottiosporella* Höhn. ex Graniti 1951 *fide* Seifert et al. 2011), *Ascomycota* genera *incertae sedis*, eight species, type: *M. setosum* B. Sutton, hyphomycetous, saprobes or plant pathogenic, widespread (tropical), see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 103360c.

Minksia Müll. Arg. 1882 (= *Cyrtographa* Müll. Arg. 1894 [1893] *fide* Species Fungorum 2017), *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, six species, type: *M. caesiella* Müll. Arg., lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Mintera Inácio & P.F. Cannon 2003, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *M. reticulata* (Starbäck) Inácio & P.F. Cannon, asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, Isotype of type species S F20503.

Minteriella Heredia, R.F. Castañeda & R.M. Arias 2013, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. cenotigena* Heredia, R.F. Castañeda & R.M. Arias, hyphomycetous, sexual morph unknown, saprobes, aquatic, Mexico, see Heredia et al. (2013; morphology), holotype of the type: XAL CB897, cultures and sequences are unavailable, compare with *Cylindrosympodium*, *Polyscytalum* and *Sympodioplanus*.

Minutiella Crous 2015, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, one species, type: *M. tardicola* (Damm & Crous) Crous, coelomycetous, sexual morph unknown, in necrotic wood of *Prunus armeniaca*, pathogens, terrestrial, South Africa, see Crous et al. (2015d; phylogeny), cultures and sequences are available, holotype and ex-type culture of type: CBS H-20000, CBS 121757.

Minutisphaera Shearer, A.N. Mill. & A. Ferrer 2011, *Minutisphaeraceae*, *Minutisphaerales*, *Dothidiomycetes*, four species, type: *M. fimbriatispora* Shearer, A.N. Mill. & A. Ferrer, freshwater, USA, Japan, see Ferrer et al. (2011; taxonomy), Raja et al. (2013b, 2015; phylogeny, new species), cultures and sequences are available, holotype of type: A242-8 (ILL).

Minutoexcipula V. Atienza & D. Hawksw. 1994, *Ascomycota* genera *incertae sedis*, seven species, type: *M. tuckerae* V. Atienza & D. Hawksw., coelomycetous, sexual morph unknown, terrestrial, lichenicolous, Europe, North America, see Atienza et al. (2009; generic concept),

Kirk et al. (2013; genus accepted), Wijayawardene et al. (2016b; morphology), cultures and sequences are unavailable, holotype of type: K(M)-IMI 352335.

Minutophoma D. Hawksw. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *M. chrysophthalmae* D. Hawksw., sexual morph unknown, lichenicolous, Europe, terrestrial, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Mirandina G. Arnaud ex Matsush. 1975 (= *Chaetonectrioides* Matsush. 1996; = *Mirandina* G. Arnaud ex Matsush. 1971; = *Mirandina* G. Arnaud 1952 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, eleven species, type: *M. corticola* G. Arnaud ex Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial and aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Ma et al. (2015; morphology), Fiúza et al. (2016; morphology), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Mirannulata Huhndorf, F.A. Fernández, A.N. Mill. & Lodge 2003, *Sordariomycetidae* genera *incertae sedis*, four species, type: *M. samuelsii* Huhndorf, F.A. Fernández, A.N. Mill. & Lodge, asexual morph unknown, saprobes, terrestrial, neotropics, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and cultures for the type species F (Huhndorf 1880); SMH 1880.

Miricatena Punith. & Spooner 2011, *Ascomycota* genera *incertae sedis*, two species, type: *M. prunicola* Punith. & Spooner, hyphomycetous, sexual morph unknown, pathogens, terrestrial, UK, see Punithalingam and Spooner (2011b; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Mirimyces Nag Raj 1993, *Ascomycota* genera *incertae sedis*, one species, type: *M. pulcher* Nag Raj, coelomycetous, saprobes or plant pathogenic, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: DAOM 215253.

Miriiquidica Hertel & Rambold 1987, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, 30 species, type: *M. complanata* (Körb.) Hertel & Rambold (*Lecanora complanata* Körb.), lichenized fungi, saxicolous, lichenicolous, worldwide temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Singh et al. (2013a; phylogeny), Miadlikowska et al. (2014a; phylogenetic placement), Lücking et al. (2017a; classification), sequence are available.

Mischoblastia A. Massal. 1852, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, three species, type: *M. oxydata* A. Massal., lichenized, see Gaya et al. (2012; phylogeny), Prieto et al. (2013; phylogeny), Prieto and Wedin (2016; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Misgomyces Thaxt. 1900, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *M. dyschirii* Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Misturatosphaeria Mugambi & Huhndorf 2009, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *M. aurantiacinotata* Mugambi & Huhndorf, asexual morph unknown, saprobes, Italy, see Mugambi and Huhndorf (2009a; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Lophiostomataceae*), Wijayawardene et al. (2014c; accepted as a genus in *Lophiostomataceae*), Jaklitsch et al. (2016c; treated as a synonym of *Teichospora* but in here we do not agree with them), cultures and sequences are available.

Mitopeltis Speg. 1921, *Dothideomycetes* genera *incertae sedis*, two species, type: *M. chilensis* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted in *Micropeltidaceae*), Hyde et al. (2013; excluded from *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Mitrula Fr. 1821, *Mitrulaceae*, *Helotiales*, *Leotiomycetes*, eight species, type: *M. paludosa* Fr., asexual morph hyphomycetous, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Mitrulinia Spooner 1987, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *M. ushuaiae* (Rehm) Spooner, asexual morph unknown, saprobes, terrestrial, southern hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures or sequence are unavailable.

Miuraea Hara 1948 (= *Hyalodictys* Subram. 1962; = *Rhopaloconidium* Petr. 1952 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *M. degenerans* (Syd. & P. Syd.) Hara, hyphomycetous, sexual morph mycosphaerella-like, plant pathogenic, see Crous et al. (2012a; phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, holotype of type: M. Miura 572 (S-F1753).

Mixtoconidium Etayo 1995, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, two species, type:

M. insidens (Vouaux) Etayo & van den Boom asexual morph related Niebla and sexual morph related to California, parasymbiont on *Ramalina*, terrestrial, Canary Islands, see Hyde et al. (2011; checklist), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), van den Boom and Etayo (2017; classification) cultures and sequences are unavailable, holotype of the type: IMI Etayo 3470.

Mixtura O.E. Erikss. & J.Z. Yue 1990, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. saginata* (Syd.) O.E. Erikss. & J.Z. Yue, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Phaeosphaeriaceae*), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; placed in *Didymellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequence are unavailable.

Miyakeomyces Hara 1913, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. bambusae* Hara, asexual morph hyphomycetous, terrestrial, saprobes, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; phylogeny), cultures and sequence are unavailable.

Miyoshiella Kawam. 1929 (see Species Fungorum 2017 for synonyms), *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *M. fusispora* Kawam., asexual morph unknown, terrestrial, saprobes, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; not accepted as valid genus), Index Fungorum (2016; synonym of *Chaetosphaeria*), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequence are unavailable.

Mniaecia Boud. 1885; *Helotiaceae*, *Helotiales*, *Leotiomycetidae*, *Leotiomycetes*, three species, type: *M. jungermanniae* (Fr.) Boud., asexual morph unknown, bryosymbiont/parasite, terrestrial, Europe, America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Czarnota and Hernik (2013; lichenized), cultures are available, two species have sequence data.

Moana Kohlm. & Volkm.-Kohlm. 1989, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *M. turbinulata* Kohlm. & Volkm.-Kohlm., asexual morph unknown, marine, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Mobergia H. Mayrhofer & Sheard 1992, *Physciaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *M. calculiformis* (W.A. Weber) H. Mayrhofer & Sheard, asexual morph unknown, terrestrial, saprobes, temperate,

see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Moelleriella Bres. 1896, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, 25 species, type: *M. sulphurea* Bres., asexual morph hyphomycetous or coelomycetous, terrestrial, entomogenous, tropical South America and French Guiana, see Chaverri et al. (2008; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Li et al. (2016a; new species, coelomycetous), Tibpromma et al. (2017b; new species), cultures and sequences are available, neotype of type: CUP 067787, ex-type strain P.C. 756.

Moleospora Abdel-Wahab, Abdel-Aziz & Nagah. 2010, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, one species, type: *M. maritima* Abdel-Wahab, Abdel-Aziz & Nagah., hyphomycetous, sexual morph unknown, saprobes, marine, Egypt, Saudi Arabia, see Abdel-Wahab et al. (2010, phylogeny), Jones et al. (2015; phylogeny, outline), Maharachchikumbura et al. (2015b, 2016b; phylogeny, outline), cultures and sequences are available.

Mollerella G. Winter 1886 (= *Agyrona* Höhn. 1909; = *Capnodiopsis* Henn. 1902; = *Elachophyllum* Petr. 1931; = *Elenkinella* Woron. 1922; = *Nostocothecea* Starbäck 1899 fide Species Fungorum 2017), *Elsinoaceae*, *Myriangiales*, *Dothideomycetes*, four species, type: *M. mirabilis* G. Winter, asexual morph unknown, saprobes, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jayawardene et al. (2014; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Mollicamarops Lar.N. Vassiljeva 2007, *Boliniaceae*, *Boliniales*, *Sordariomycetes*, one species, type: *M. stellata* Lar.N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, north temperate zone, see Huhndorf and Miller (2008; morphology), Lumbsch and Huhndorf (2010; outline), Untereiner et al. (2013; accepted in *Boliniaceae*), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: VLA P-2942.

Mollisia (Fr.) P. Karst. 1871 (= *Aleuriella* (P. Karst.) P. Karst. 1871; = *?Atrocybe* Velen. 1947; = *Belonopsis* (Sacc.) Rehm 1891; = *Crustula* Velen. 1934; = *Cystodendron* Bubák 1914; = *Dibeloniella* Nannf. 1932; = *Haglundia* Nannf. 1932; *Mollisiella* Boud. 1885; *Peziza* sect. *Aleuriella* P. Karst. 1869; *Peziza* trib. *Mollisia* Fr. 1822; = *Syntexis* Theiss. 1916; = *Tapesia* (Pers.) Fuckel 1870 fide Jaklitsch et al. 2016a; Species Fungorum 2017), *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, c. 130 species, type: *M. cinerea* (Batsch) P. Karst., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Tanney et al. (2016; DNA), cultures and

sequences are available, available specimen and cultures for the type: CBS H-14530, AFTOL-ID 76.

Mollisina Höhn. ex Weese 1926 (= *Dendrotrichoscypha* Svrček 1977 *fide Species Fungorum* 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, eleven species, type: *M. rubi* (Rehm) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, America, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), cultures and sequences are available.

Mollisinopsis Arendh. & R. Sharma 1984, *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *M. filicis* Arendh. & R. Sharma, asexual morph unknown, saprobes, terrestrial, Bhutan, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of the type: PAN 17819.

Mollisiopsis Rehm 1908, *?Helotiales genera incertae sedis*, *Leotiomycetes*, seven species, type: *M. subcinerea* Rehm, asexual morph unknown, saprobes, terrestrial, America, Europe, see Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification; treated as a synonym of *Pyrenopeziza* Fuckel), cultures and sequences are unavailable.

Monandromyces R.K. Benj. 1999, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eleven species, type: *M. microveliae* (Thaxt.) R.K. Benj., asexual morph unknown, biotrophic, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Monascella Guarro & Arx 1986, *Orygenaceae*, *Orygenales*, *Eurotiomycetes*, one species, type: *M. botryosa* Guarro & Arx, asexual morph unknown, saprobes, Spain, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for the type: CBS 233.85.

Monascostroma Höhn. 1918, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, c. five species, type: *M. innumerousum* (Desm.) Höhn., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Kirk et al. (2008; treated as a synonym of *Hendersonia* Berk.), Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), cultures and sequences are available.

Monascus Tiegh. 1884 (= *Allescheria* Sacc. & P. Syd. 1899; = *Basipetospora* G.T. Cole & W.B. Kendr. 1968; = *Backusia* Thirum., M.D. Whitehead & P.N. Mathur 1965 [1964]; = *Eurotiella* Lindau 1900; = *Eurotiopsis* Costantin ex Laborde 1897; = *Physomyces* Harz 1890; *fide* Rossman et al. 2016a), *Monascaceae*, *Eurotiales*, *Eurotiomycetes*, 14 species, type: *Monascus ruber* Tiegh., asexual morph

unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016a; nomenclature), cultures and sequences are available, available specimen for type: CBS H-14541.

Monerolechia Trevis. 1857, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, five species, type: *M. bayrhofferi* (Schaer.) Trevis., lichenized, Asia, Australia, Europe, see Gaya et al. (2012; phylogeny), Prieto et al. (2013; phylogeny), Prieto and Wedin (2016; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Monilia Honey 1928 (= *Monilia* Bonord. 1851 *fide* Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, c. 30 species, type: *M. fructicola* (G. Winter) Honey, asexual morph formerly named in *Monilia*, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Zhu et al. (2016; pathogens, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Monilochaetes Halst. ex Harter 1916 (= *Australiasca* Sivan. & Alcorn 2002; = *Dischloridium* B. Sutton 1977 [1976]; = *Hyalocylindrophora* J.L. Crane & Dumont 1978 *fide* Réblová et al. 2016c; *Species Fungorum* 2017), *Australiascaceae*, *Glomerellales*, *Sordariomycetes*, two species, type: *M. infuscans* Harter, hyphomycetous, sexual morph formerly named in *Australiasca*, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Réblová et al. (2011, 2016c; taxonomy, phylogeny, nomenclature), Maharanachikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available, available specimen and cultures for the type: CBS H-1013, CBS 869.96.

Monoblastia Riddle 1923, *Monoblastiaceae*, *Monoblastiales*, *Dothideomycetes*, eleven species, type: *M. palmicola* Riddle, asexual morph unknown, lichenized, Spain, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Monoblastiopsis R.C. Harris & C.A. Morse 2008, *Dothideomycetes genera incertae sedis*, two species, type: *M. konzana* R.C. Harris & C.A. Morse, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: KANU 371963.

Monochaetia (Sacc.) Allesch. 1902 (= *Lennisia* Nieuwl. 1916; = *Pestalotia* subgen. *Monochaetia* Sacc. 1884 *fide* *Species Fungorum* 2017), *Xylariomycitidae* genera *insertae*

sedis, *Sordariomycetes*, 28 species, type: *M. monocheata* (Desm.) Allesch., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; phylogeny), de Silva et al. (2017, new species, phylogeny), cultures and sequences are available, available specimen and cultures for the type: CBS H-14561; CBS 115004, CBS 191.82.

Monochaetiella E. Castell. 1943, *Ascomycota* genera *incertae sedis*, three species, type: *M. hyparrheniae* E. Castell., asexual morph coelomycetous, saprobes or plant pathogenic, widespread, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Monochaetellopsis B. Sutton & DiCosmo 1977 (= *Hypotheca* Tommerup 1970 *fide* Johnston et al. 2014b), *Helotiales* genera *incertae sedis*, *Leotiomyces*, two species, type: *M. themedae* (M. Kandasw. & Sundaram) B. Sutton & DiCosmo, coelomycetous, sexual morph *Hypotheca*, saprobes, terrestrial, India, Australia, Tanzania, see Wijayawardene et al. (2012, 2017; outline), Johnston et al. (2014b; nomenclature), cultures and sequences are unavailable, holotype of the type species is not designated, other specimens IM I78370a, 151159a, 62200.

Monochaetinula Muthumary, Abbas & B. Sutton 1996, *Ascomycota* genera *incertae sedis*, six species, type: *M. terminaliae* (Bat. & J.L. Bezerra) Muthumary, Abbas & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Monocillium S.B. Saksena 1955, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, c. 15 species, type: *M. indicum* S.B. Saksena, asexual morph unknown, saprobes, on fungi and lichens, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Monodia Breton & Faurel 1970, *Ascomycota* genera *incertae sedis*, two species, type: *M. elegans* Breton & Faurel, coelomycetous, coprophilous and saprobes, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Monodictys S. Hughes 1958, *Dothideomycetes* genera *incertae sedis*, c. 50 species, type: *M. putredinis* (Wallr.) S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology, treated as heterogenous),

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Monodidymaria U. Braun 1994, *Ascomycota* genera *incertae sedis*, five species, type: *M. canadensis* (Ellis & Everh.) U. Braun (\equiv *Ramularia canadensis* Ellis & Everh.), asexual morphs hyphomycetous, terrestrial, plant pathogenic, leaf-spotting, Asia, Europe, North and South America, see Hyde et al. (2013; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequence are unavailable, lectotype of type: NY 1293230.

Monodisma Alcorn 1975, *Ascomycota* genera *incertae sedis*, one species, type: *M. fragilis* Alcorn, hyphomycetous, sexual morph unknown, pathogens, terrestrial, widespread, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, cultures available for type: IMI 73184, IMI 41189.

Monoicomycetes Thaxt. 1900, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 49 species, type: *M. homalotae* Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species, Ecuador), Kirk et al. (2013; genus accepted) sequences are available only for MT004.

Monorhizina Theiss. & Syd. 1916, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *M. filicina* (Berk. & Broome) Theiss. & Syd., terrestrial, on leaves, Sri Lanka, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Monosporascus Pollack & Uecker 1974 (= *Bitrimonospora* Sivan., Talde & Tilak 1974 *fide* Species Fungorum 2017), *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *M. cannonballus* Pollack & Uecker, asexual morph unknown, thermophilic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted in *Sordariomycetes*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, available specimen and cultures for the type: BPI 71820; ATCC 26931 (unpublished), FMR6682.

Monostichella Höhn. 1916, *Ascomycota* genera *incertae sedis*, 15 species, type: *M. robergei* (Desm.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Monotosporella S. Hughes 1958, *Savoryellaceae*, *Savoryellales*, *Sordariomycetes*, probably polyphyletic, c. 10 species, type: *M. albida* G. Weber, Spaaij & Oberw., asexual morph unknown, saprobes, Asia, Australia, see Seifert et al. (2011; morphology, treated as *Pleosporales*),

Kirk et al. (2013; genus did not list), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Montagnella Speg. 1881, *Dothideomycetes* genera *incertae sedis*, nine species, type: *M. curumamuel* Speg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*), cultures and sequences are unavailable.

Montagnula Berl. 1896, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 30 species, type: *M. infernalis* (Niessl) Berl., asexual morph unknown, saprobes, terrestrial, worldwide, see Zhang et al. (2012e; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Hyde et al. (2014; taxonomy, phylogeny), Ariyawansa et al. (2014a, e; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), Wanasinghe et al. (2016a; phylogeny), Tennakoon et al. (2016b; taxonomy, phylogeny), cultures and sequences are available.

Montanelia Divakar, A. Crespo, Wedin & Essl. 2012, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *M. panniformis* (Nyl.) Divakar, A. Crespo, Wedin & Essl., asexual morph unknown, lichenized, terrestrial, worldwide, see Divakar et al. (2012; taxonomy), Leavitt et al. (2015; morphology, phylogeny), Szczepańska et al. (2015; distribution in Poland and Europe), cultures and sequences are available.

Monticola Selbmann & Egidi 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *M. elongata* Selbmann & Egidi, hyphomycetous, sexual morph unknown, rock-inhabiting, Italy, see Egidi et al. (2014; taxonomy), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2014c, 2017; outline), type: CBS 136206 = CCFEE 5394, cultures and sequences available.

Moorella P. Rag. Rao & D. Rao 1964, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, one species, type: *M. speciosa* P. Rag. Rao & D. Rao, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Morakotiella (C.A. Farrant & E.B.G. Jones) Sakay. 2005, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *M. salina* (C.A. Farrant & E.B.G. Jones) Sakay., asexual morph unknown, saprobes, marine, USA, see Lumbsch and Huhndorf (2010; outline), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b;

outline), cultures and sequences are available, holotype of the type: IMI 297765.

Moralesia Urries 1956, *Ascomycota* genera *incertae sedis*, one species, type: *M. arundinis* Urries, sexual morph unknown, saprobes, terrestrial, Canary Islands, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Morchella Dill. ex Pers. 1794 (= *Costantinella* Matr. 1892 *fide* Healy et al. 2016; see Species Fungorum 2017 for additional synonyms), *Morchellaceae*, *Pezizales*, *Pezizomycetes*, 40 species, type: *M. esculenta* (L.) Pers., asexual morph unknown, saprobes, terrestrial, temperate, see Isiloglu et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Clowez (2012; new species, France), Kirk et al. (2013; genus accepted), Elliott et al. (2014; new species, New South Wales), Loizides et al. (2016; new species, Cyprus), cultures and sequences are available, epitype of type: PhC10.

Morenoina Theiss. 1913 (= *Aulographella* Höhn. 1917 *fide* Species Fungorum 2017), *Aulographaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, 25 species, type: *M. antarctica* (Speg.) Theiss., asexual morph coelomycetous (acarella-like), epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted in *Asterinaceae*), Hongsan et al. (2014c; taxonomy, accepted in *Aulographaceae*), Wijayawardene et al. (2014c; outline, accepted in *Aulographaceae*), Tibpromma et al. (2017b; new species, phylogeny), cultures and sequences are available, epitype of type: IMI 349680.

Morinia Berl. & Bres. 1889 (= *Rinomia* Nieuwl. 1916 *fide* Species Fungorum 2017), *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, two species, type: *M. pestalozziioides* Berl. & Bres., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), cultures and sequences are available.

Moriola Norman 1872, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 15 species, type: *M. descensa* Norman, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Moriolomyces Cif. & Tomas. 1953, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. descensae* Cif. & Tomas., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Moristroma A.I. Romero & Samuels 1991, *Chaetothyriomycetidae*, genera *incertae sedis*,

Eurotiomycetes, four species, type: *M. polysporum* A.I. Romero & Samuels, asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Moromyces Abdel-Wahab, K.L. Pang, Nagah., Abdel-Aziz & E.B.G. Jones 2010, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, one species, type: *M. varius* (Chatmala & Somrith.) Abdel-Wahab, K.L. Pang, Nagah., Abdel-Aziz & E.B.G. Jones, hyphomycetous, sexual morph unknown, saprobes, marine, Egypt, Japan and Thailand, see Abdel-Wahab et al. (2010; phylogeny), Wijayawardene et al. (2012, 2017; outline), Jones et al. (2015; phylogeny, outline), Maharachchikumbura et al. (2015b, 2016b; phylogeny, outline), cultures and sequences are available, holotype of the type: BBH IT152.

Morosphaeria Suetrong, Sakay., E.B.G. Jones & C.L. Schoch 2009, *Morosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *M. velatispora* (K.D. Hyde & Borse) Suetrong, Sakay., E.B.G. Jones & C.L. Schoch, asexual morph unknown, saprobes, on wood in mangrove habitats, worldwide, see Schoch et al. (2009; phylogeny), Suetrong et al. (2009, 2011a, b; phylogeny, morphology), Lumbsch and Huhndorf (2010; outline), Zainuddin et al. (2010; antimicrobial activity), Pang et al. (2011b; monograph), Sakayaroj et al. (2011b; diversity), Boonmee et al. (2012; phylogeny), Zhang et al. (2012e; phylogeny, morphology), Hyde et al. (2013; phylogeny, morphology, outline), Sarma and Raghukumar, (2013; diversity), Ariyawansa et al. (2014a, e; phylogeny), Shearer et al. (2014; phylogeny, freshwater), Wijayawardene et al. (2014c; outline), Jones et al. (2015; outline, marine), Liu et al. (2015a; phylogeny), Tanaka et al. (2015, phylogeny), Singtripop et al. (2015; phylogeny), sequence data available for two species, holotype of type species IMI297770.

Morrisographium M. Morelet 1968 (= *Phragmographium* E.F. Morris 1966 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, eight species, type: *M. pilosum* M. Morelet, hyphomycetous, sexual morph unknown, saprobes, terrestrial, freshwater, widespread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Moserella Pöder & Scheuer 1994, *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *M. radicicola* Pöder & Scheuer, asexual morph unknown, on roots, terrestrial, Austria, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), sequences are unavailable, holotype/paratype? of the type species IB 88/310.

Mouliniea C.P. Robin 1853, *Ascomycota* genera *incertae sedis*, accepted as a genus in *Pezizomycotina* Kirk et al. (2008) and Kirk et al. (2013) genus accepted. However, previous outlines i.e. Lumbsch and Huhndorf (2010) or Wijayawardene et al. (2012, 2017) did not list as a genus in *Pezizomycotina*.

Moutoniella Penz. & Sacc. 1902, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *M. polita* Penz. & Sacc., asexual morph unknown, saprobes, terrestrial, India, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mucispora J. Yang, Bhat & K.D. Hyde 2016, *Fuscosporellaceae*, *Fuscosporellales*, *Sordariomycetes*, one species, type: *M. pseudoseptata* J. Yang, Bhat & K.D. Hyde, hyphomycetous, saprobes, Thailand, see Yang et al. (2016c; taxonomy, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Muciturbo P.H.B. Talbot 1989, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, three species, type: *M. reticulatus* P.H.B. Talbot, asexual morph unknown, saprobes, terrestrial, Australia, see Kirk et al. (2008; genus accepted), Lumbsch and Huhndorf (2010; genus not accepted), Healy et al. (2013; DNA), Kirk et al. (2013; genus not accepted), Index Fungorum (2016; genus accepted), cultures and sequences are available.

Mucoharknessia Crous, R.M. Sánchez & Bianchin. 2015, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, two species, type: *M. cortaderiae* Crous, R.M. Sánchez & Bianchin, coelomycetous, sexual morph unknown, saprobes, terrestrial, Argentina, Italy, see Crous et al. (2015b; taxonomy, phylogeny), Li et al. (2016a; new species, phylogeny), Wijayawardene et al. (2017; outline), isotype and ex-isotype of the type: CBS H-21853, CBS 131032.

Mucomassaria Petr. & Cif. 1932, *Mucomassariaceae*, *Ascomycota* genera *incertae sedis*, one species, type: *M. maxima* Petr. & Cif., asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mucomycosphaerella Quaedvl. & Crous 2014, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *M. eurypotami* (Kohlm., Volkm.-Kohlm. & O.E. Erikss.) Quaedvl. & Crous, asexual morph unknown, terrestrial, USA, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), cultures and sequences available, holotype and ex-type of type: JK5586; JK5586J.

Muellerella Hepp ex Müll. Arg. 1862 (= *Muellerellomyces* Cif. & Tomas. 1953; = *Pleothelis* M. Choisy 1949; = *Polycarpella* Theiss. & Syd. 1918; =

Sychnogonia Körb. 1855; = *Thelidiola* C.W. Dodge 1968 *fide Species Fungorum* 2017), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 18 species, type: *M. polyspora* Hepp ex Müll. Arg., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Etayo and Sancho (2008; new species, Magallanes) Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Berger and LaGreca (2014; new species, Bermuda), Muggia et al. (2015b; DNA), cultures and sequences are available but lacking for the type species.

Muellerites L. Holm 1968, *Dothidotthiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. juniperi* (E. Müll. & Arx) L. Holm, asexual morph unknown, saprobes, terrestrial, temperate, Switzerland, see Lumbsch and Huhndorf (2010; outline, accepted in *Dothideomycetes*), Hyde et al. (2013; outline, phylogeny), Kirk et al. (2013; genus accepted), Dai et al. (2014a; transferred to *Dothidotthiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type species.

Muelleromyces Kamat & Anahosur 1968, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *M. indicus* Kamat & Anahosur, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Muiaria Thaxt. 1914, *Laboulbeniales* genera *incertae sedis*, one species, type: *M. gracilis* Thaxt., hyphomycetous, sexual morph unknown, on insects, terrestrial, Africa, Borneo, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Muiogone Thaxt. 1914, *Ascomycota* genera *incertae sedis*, two species, type: *M. chromopteri* Thaxt., hyphomycetous, sexual morph unknown, pathogen, terrestrial, West Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Muirella R. Sprague 1959, *Ascomycota* genera *incertae sedis*, one species, type: *M. alasensis* R. Sprague, hyphomycetous, sexual morph unknown, pathogens, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Mulderomyces Crous, Jacq. Edwards & P.W.J. Taylor 2016, *Ostropales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *M. natalis* Crous, Jacq. Edwards & P.W.J. Taylor, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017;

outline), holotype and ex-type of the type: CBS H-22609, CPC 25519.

Multilocularia Phook., Ariyaw. & K.D. Hyde 2016, *Parabambusicolaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. bambusae* Phook., Ariyaw. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Li et al. (2016a; taxonomy, phylogeny), holotype and ex-isotype of the type: MFLU 11-0216, MFLUCC 11-0180.

Multiseptospora Phook. & K.D. Hyde 2015, *Parabambusicolaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *M. thailandica* Phook. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Liu et al. (2015a; taxonomy, phylogeny), Li et al. (2016a; new species), holotype and ex-type of the type: MFLU 11-0219, MFLUCC 11-0183.

Multisporascus Marinc., M.J. Wingf. & Crous 2008, *Ascomycota* genera *incertae sedis*, one species, type: *M. proteae* Marinc., M.J. Wingf. & Crous, asexual morph unknown, saprobes, terrestrial, South Africa, see Marincowitz et al. (2008; morphology), cultures and sequences are unavailable, holotype of the type: PREM 59465.

Munkia Speg. 1886 (= *Aschersoniopsis* Henn. 1902; = *Pycnostroma* Clem. 1909; = *Shropshiria* F. Stevens 1927 *fide Species Fungorum* 2017), *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *M. martyris* Speg., asexual morph hyphomycetous, terrestrial, saprobes, South America, see Hyde et al. (2011; checklist), Wijayawardene et al. (2012, 2017; outline), Maharanachikumbura et al. (2015b, 2016b; phylogeny), cultures and sequences are unavailable.

Munkiella Speg. 1885 (= *Apotrabutia* Petr. 1929; = *Coscinopeltis* Speg. 1909 *fide Species Fungorum* 2017), *Polystomellaceae*, *Dothideomycetes* families *incertae sedis*, three species, type: *M. caa-guazu* Speg., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Munkovalsaria Aptroot 1995, *Dacampiaceae*, *Dothideomycetes* families *incertae sedis*, three species, type: *M. donacina* (Niessl) Aptroot, asexual morph hyphomycetous (lecythophora-like), saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline), Kirk et al. (2013; genus accepted), Pitt et al. (2014; from grapevine, DNA), Wijayawardene et al. (2014c; outline), cultures and sequences are available, type specimen and cultures available for the type: CBS H-14398; HVV01.

Muraeriatata Huhndorf, Greif, Mugambi & A.N. Mill. 2008, *Magnaportheaceae*, *Magnaporthales*,

Sordariomycetes, one species, type: *M. collapsa* Huhndorf, Greif, Mugambi & A.N. Mill., asexual morph unknown, saprobes, terrestrial, Austria, see Huhndorf et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available, holotype and available cultures for the type: GKM1084, SMH4553.

Murangium Seaver 1951, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *M. sequoiae* (Plowr. ex W. Phillips) Seaver, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted in *Patellariaceae*), cultures and sequences are unavailable.

Muricopeltis Viégas 1944, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. brasiliensis* Viégas, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Micropeltidaceae*), Hyde et al. (2013; did not accept in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Muriformistrickeria Q. Tian, Wanasi., Camporesi & K.D. Hyde 2015, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. rubi* Q. Tian, Wanasi., Camporesi & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Italy, see Tian et al. (2015; morphology, nomenclature, phylogeny), cultures and sequences are available for type species, holotype and ex-type strains of type: MFLU 15-1409; MFLUCC 15-0681 = KUMCC15-0079.

Murilentithecium Wanasi., Camporesi, E.B.G. Jones & K.D. Hyde 2014, *Lentitheciaeae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. clematis* Wanasi., Camporesi, E.B.G. Jones & K.D. Hyde, asexual morph camarosporium-like, saprobes, terrestrial, Europe, see Wanasinghe et al. (2014a; morphology, phylogeny), cultures and sequences are available, holotype and ex-type strains of type species MFLU 14-0334; MFLUCC 14-0562 = MUCL 55405.

Muriphaeosphaeria Phukhams., Bulgakov & K.D. Hyde 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *M. galatellae* Phukhams., Bulgakov & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Russia, see Phukhamsakda et al. (2015; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: MFLU 15-0087, cultures and sequences are available.

Muripulchra Z.L. Luo, H.Y. Su & K.D. Hyde 2017, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, one species,

type: *M. aquatica* Z.L. Luo, H.Y. Su & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, see Luo et al. (2017; taxonomy, phylogeny), holotype of the type: HKAS 83980, cultures and sequences are available.

Muripulchra Z.L. Luo, H.Y. Su & K.D. Hyde 2017, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, one species, type: *M. aquatica* Z.L. Luo, H.Y. Su & K.D. Hyde, sporodochial hyphomycete, sexual morph unknown, saprobes, aquatic, China, see Luo et al. (2017; morphology, phylogeny), cultures and sequences are available, type HKAS 83980.

Murispora Y. Zhang bis, J. Fourn. & K.D. Hyde 2009, *Amniculicolaceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *M. rubicunda* (Niessl) Y. Zhang ter, J. Fourn. & K.D. Hyde, asexual morph phoma-like, saprobes, terrestrial and aquatic, Europe, significantly substrate stained purple, see Zhang et al. (2009a, b, 2012e; morphology, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Amniculicolaceae*), Wijayawardene et al. (2014c; outline), Wanasinghe et al. (2015; morphology, phylogeny), cultures and sequences are available.

Muroia I. Hino & Katum. 1958, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. nipponica* I. Hino & Katum., asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Lophiostomataceae*), Hyde et al. (2013; did not accept in *Lophiostomataceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Muscia Gizhitsk. 1929, *Helotiaceae* genera *incertae sedis*, *Leotiomycetes*, one species, type: *M. catharineae* Gizhitsk., asexual morph unknown, on living stalk of *Catharinea* sp., terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Muscicola Velen. 1934, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *M. dubia* (Velen.) Svrček, asexual morph unknown, saprobes, terrestrial, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, lectotype PRM 148165.

Muscodor Worapong, Strobel & W.M. Hess 2001, *Sordariomycetes* genera *incertae sedis*, 19 species, type: *M. albus* Worapong, Strobel & W.M. Hess, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Meshram et al. (2017; new species, phylogeny), Wendt et al. (2017; accepted as a genus in *Sordariomycetes* genera *incertae sedis*), cultures and sequences are available.

Musicillium Zare & W. Gams 2007, *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*,

one species, type: *M. theobromae* (Turconi) Zare & W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available, neotype (dry culture) and ex-type of the type: Herb. CBS 19864, CBS 968.72.

Muyocopron Speg. 1881 (= *Ellisiodothis* Theiss. 1914; = *Haplopeltis* Theiss. 1914; = *Peltella* Syd. & P. Syd. 1917; = *Peltopsis* Bat. 1960 *fide* Species Fungorum 2017), *Muyocopronaceae*, *Muyocpronales*, *Dothideomycetes*, five species, type: *M. correntinum* Speg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted in *Microthyriaceae*), Hyde et al. (2013; introduced *Muyocopronaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Mapook et al. (2016c; taxonomy, phylogeny), Tibpromma et al. (2016d; new species), cultures and sequences are available, but lacking for the type.

Mycelephas R.F. Castañeda 2009 (= *Arnoldiella* R.F. Castañeda 1984 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *M. robustus* (R.F. Castañeda) R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Myceliophthora Costantin 1892 (= *Corynascus* Arx 1973 *fide* Maharachchikumbura et al. 2015b, 2016b), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *M. lutea* Costantin, sexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, nomenclature), van den Brink et al. (2015; DNA, phylogeny), cultures and sequences are available for the type species CBS 146.50, CBS 145.77 (van den Brink et al. 2015), available specimen for type: CBS H-10236.

Mycerema Bat., J.L. Bezerra & Cavalc. 1963 (see Species Fungorum 2017 for synonyms), *Schizothyriaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *M. vochysiacearum* Bat., J.L. Bezerra & Cavalc., asexual morph plenotrichiaus-like, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Phookamsak et al. (2016; new species, taxonomy), cultures and sequences are unavailable.

Mycoarachis Malloch & Cain 1970, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *M. inversa* Malloch & Cain, asexual morph

hyphomycetous, terrestrial, dung, Africa and North America, see Hirooka et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Bionectriaceae*), Maharachchikumbura et al. (2015b, 2016b; phylogeny), cultures and sequences are available, holotype and ex-type of type: TRTC 662166f; CBS 517.70.

Mycoarctium K.P. Jain & Cain 1973, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, ?two species, type: *M. ciliatum* K.P. Jain & Cain, asexual morph unknown, coprophilous, terrestrial, Australia, Canary Islands, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), sequences are unavailable, holotype and ex-type of the type species? CBS 857.73, ATCC 26246, IMI 179697, TRTC 46385.

Mycoarthris Marvanová & P.J. Fisher 2002, *Hyaloscypthaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *M. corallina* Marvanová & P.J. Fisher, hyphomycetous, sexual morph unknown, saprobes, aquatic, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available, holotype of the type: species IMI 385400.

Mycobilimbia Rehm 1890 (= *Probilimbia* Vain. 1899; = *Weitenwebera* Opiz 1857 *fide* Species Fungorum 2017), *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, five species, type: *M. obscurata* (Sommerf.) Rehm, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Joshi and Hur (2012; Korea), Kirk et al. (2013; genus accepted), Miądkowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type species.

Mycoblastus Norman [nom. cons.] 1853 (= *Megalospora* A. Massal. 1852; = *Mycoblastomyces* Cif. & Tomas. 1953; = *Oedemocarpus* Trevis. 1857 *fide* Species Fungorum 2017), *Tephromelataceae*, *Lecanorales*, *Lecanoromycetes*, ten species, type: *M. sanguinarius* (L.) Norman, lichenized, worldwide, see Spribille et al. (2011; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Mycocalicium Vain. 1890 (= *Sphinctrinella* Nádv. 1942 *fide* Species Fungorum 2017), *Mycocaliciaceae*, *Mycocaliciales*, *Eurotiomycetes*, twelve species, type: *M. subtile* (Pers.) Szatala, asexual morph phialophora-like, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Mycocentrodochium K. Matsush. & Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *M. curvisporum* K. Matsush. & Matsush., asexual morph hyphomycetous, Japan, saprobes or plant pathogenic, see

Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: Matsushima Fungus Collection, Kobe, 5M406.

Mycocentrospora Deighton 1972 (= *Ansatospora* A.G. Newhall 1944; = *Centrospora* Neerg. 1942 *fide* Species Fungorum 2017), *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, four species, type: *M. acerina* (R. Hartig) Deighton (≡ *Cercospora acerina* R. Hartig), hyphomycetous, terrestrial, plant pathogenic (leaves, roots) and saprobes (isolated from soil), Asia, Australia, Europe, North America, New Zealand, see Seifert et al. (2011; phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, neotype of type: Herb. Mycol. Oecon. 743 (B).

Mycochaetophora Hara & Ogawa 1931, *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *M. japonica* Hara & Ogawa, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Kobayashi et al. (2009; morphology, phylogeny), Seifert et al. (2011; phylogeny), Wijayawardene et al. (2012, 2017; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are available but lacking for the type species.

Mycochlamys S. Marchand & Cabral 1976, *Ascomycota* genera *incertae sedis*, one species, type: *M. macrospora* S. Marchand & Cabral, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Argentina, see Wijayawardene et al. (2012, 2017; outline), cultures available for type: CBS 639.76, BAFC 2663, IMI 185605, F277901.

Mycocitrus A. Möller 1901 (= *Shiraiella* Hara 1914 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *M. aurantium* Möller, asexual morph hyphomycetous, terrestrial, plant pathogen, South America, Japan, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; phylogeny), cultures and sequences are unavailable, epitype of type: BPI 631727.

Mycoclelandia Trappe & G.W. Beaton 1984 (= *Clelandia* Trappe 1979 *fide* Species Fungorum 2017), *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *M. arenacea* (Trappe) Trappe & G.W. Beaton, asexual morph unknown, hypogeous, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Trappe et al. (2010b; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for the type: OSC:131132.

Mycocryptospora J. Reid & C. Booth 1987, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. anthostiomoides* (Rehm) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, Germany, see

Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Mycodidymella C.Z. Wei, Y. Harada & Katum. 1998, *Dothideomycetes* genera *incertae sedis*, one species, type: *M. aesculi* C.Z. Wei, Y. Harada & Katum., asexual morph unknown, saprobes, terrestrial, Germany, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: HHUF 22892.

Mycodiella Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *M. eucalypti* Crous, asexual morph unknown, saprobes, terrestrial, Australia, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type: CBS H-22885, CPC 29226.

Mycöenterolobium Goos 1970, *Ascomycota* genera *incertae sedis*, one species, type: *M. platysporum* Goos, sexual morph unknown, saprobes, terrestrial, Hawaii, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Karandikar et al. (2015; morphology), cultures and sequences are unavailable.

Mycofalcella Marvanová, Om-Kalth. & J. Webster 1993, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *M. calcarata* Marvanová, Om-Kalth. & J. Webster, hyphomycetous, sexual morph unknown, saprobes, freshwater, British Isles (Great Britain), Pakistan, see Baschien et al. (2013; phylogeny), Benucci et al. (2016; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype and culture available for type species IMI 351879; CCM F-10289.

Mycogalopsis Gjurašin 1925, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *M. retinospora* Gjurašin, asexual morph unknown, saprobes or plant pathogenic, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycoglaena Höhn. 1909, *Dothideomycetes* genera *incertae sedis*, 16 species, type: *M. subcoeruleascens* (Nyl.) Höhn., asexual morph unknown, saprobes and plant pathogens, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycogone Link 1809 (= *Coccosporella* P. Karst. 1893 [1893-1894]; = *Mycobanche* Pers. 1818 *fide* Species Fungorum 2017), *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, 29 species, type: *M. rosea* Link, hyphomycetous, widespread, mycoparasites (mushroom), saprobes or plant pathogenic, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

Rossman et al. (2013b; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for the type: CBS 132.97, CBS 260.39, CBS 488.67, CBS 527.80, CBS 563.78A, CBS 563.78B, available specimen of type: CBS H-14849, CBS H-14850.

Mycohypallage B. Sutton 1963, *Ascomycota* genera *incertae sedis*, three species, type: *M. congesta* (Berk. & Broome) B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Africa, South Asia, Russia, see Marincowitz et al. (2010; morphology, new species), Wijayawardene et al. (2016b; morphology), cultures and sequences are unavailable, holotype of the type: IM I84983.

Mycoleptodiscus Ostaz. 1968, *Magnaportheaceae*, *Magnaporthales*, *Sordariomycetes*, c. eleven species, type: *M. terrestris* (Gerd.) Ostaz., sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Whitton et al. (2012a; new species), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Mycomalus Möller 1901, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. bambusinus* Möller, asexual morph unknown, saprobes, terrestrial, Brazil, see Tanaka et al. (2008; compare with *Ustilaginoidea*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Mycomediospora G.C. Carroll & Munk 1964, *Ascomycota* genera *incertae sedis*, one species, type: *M. flava* (Rick) Carroll & Munk, asexual morph unknown, saprobes, terrestrial, Brazil, see Tanaka et al. (2008; compare with *Ustilaginoidea*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Mycomelanea Velen. 1947, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *M. ludmilae* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline, misspelled as *Mycomelaena*), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as a synonym of *Coccozymes* De Not.), cultures and sequences are unavailable.

Mycomicrothelia Keissl. 1936, *Trypetheliaceae*, *Trypethiales*, *Dothideomycetes*, three species, type: *M. macularis* (Hampe ex A. Massal.) Keissl., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2011b; DNA, phylogeny), Hyde et al. (2013; accepted as a

genus in *Trypetheliaceae*), Wijayawardene et al. (2014c; outline), cultures available but lacking for the type species.

Myconenesia Kirschst. 1936, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *M. formosa* (Kirschst.) Kirschst., asexual morph unknown, saprobes, see Kirk et al. (2008; treated as a synonym of *Anthostomella* Sacc.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a distinct genus), cultures and sequences are unavailable.

Mycopappus Redhead & G.P. White 1985, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *M. alni* (Dearn. & Barthol.) Redhead & G.P. White, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Lee et al. (2013b; pathogens, DNA), Phookamsak et al. (2014c; phylogeny), Tian et al. (2015; accepted as a genus in *Melanommataceae*), cultures and sequences are available.

Mycopara Bat. & J.L. Bezerra 1960, *Ascomycota* genera *incertae sedis*, one species, type: *M. shawii* Bat. & J.L. Bezerra, coelomycetous, sexual morph unknown, on insects, terrestrial, Africa, Borneo, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycoporon Boise 1987, *Dothideomycetes* genera *incertae sedis*, two species, type: *M. anensis* Boise, asexual morph unknown, saprobes, terrestrial, Germany, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type.

Mycophilomyces Crous & M.J. Wingf. 2016, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. periconiae* Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type of the type: CBS H-22878, CPC 27558.

Mycophyrias Kohlm. & Volkm.-Kohlm. 1998, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *M. ascophylli* (Cotton) Kohlm. & Volkm.-Kohlm., saprobes, marine, cosmopolitan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Verrucariaceae*), Toxopeus et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Jaklitsch et al. (2016a; classification, accepted as a genus in *Verrucariaceae*), cultures and sequences are available.

Mycoporellum Müll. Arg. 1884 (= *Keisslerellum* Werner 1944 *fide* Species Fungorum 2017),

Dothideomycetes genera *incertae sedis*, seven species, type: *M. sparsellum* (Nyl.) Müll. Arg., asexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Mycoporis Clem. 1909, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *M. perexigua* (Müll. Arg.) Clem., asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideaceae*), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; transferred to *Mycosphaerellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Mycoporopsis Müll. Arg. 1885, *Dothideomycetes* genera *incertae sedis*, six species, type: *M. sorenocarpa* (C. Knight) Müll. Arg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Mycoporum Flot. ex Nyl. [nom. cons.] 1855 (= *Bottaria* A. Massal. 1856; = *Dictyothyrium* Grove 1932; = *Mycoporum* G. Mey. 1825; = *Pseudomycoporon* Marchand 1896 *fide* Species Fungorum 2017), *Mycoporaceae*, *Pleosporales*, *Dothideomycetes*, 25 species, type: *M. elabens* (A. Massal.) Flot. ex Nyl., asexual morph unknown, non-lichenized or facultatively lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mycoporaceae*), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Mycorrhynchidium Malloch & Cain 1971, *Pyxidiophoraceae*, *Pyxidiophorales*, *Laboulbeniomycetes*, one species, type: *M. saccatum* Malloch & Cain, asexual morph unknown, coprophilous, terrestrial, Kenya, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycosphaerangium Verkley 1999, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *M. tetrasporum* (Ellis) Verkley, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycosphaerelloides Videira & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *M. madeirae* (Crous & Denman) Videira & Crous, asexual morph unknown, plant pathogens, see Videira et al. (2016; morphology, phylogeny), holotype

and ex-type of type species CBS H-9898; CBS 112895 = CPC 3745.

Mycospraguea U. Braun 1993, *Ascomycota* genera *incertae sedis*, one species, type: *M. campicola* (Rogerson & R. Sprague) U. Braun, coelomycetous, terrestrial, plant pathogenic, leaf-spotting, North America, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, lectotype and isotypes of type: WSP 42832; NY 256526, 1028516.

Mycosticta Höhn. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *M. ovalis* (Pass.) Höhn., coelomycetous, sexual morph unknown, on insects, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycosylva M.C. Tulloch 1973, *Ascomycota* genera *incertae sedis*, three species, type: *M. clarkii* M.C. Tulloch, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Great Britain, Finland, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures for type: CBS 608.73, ATCC 32970, IMI 163345, type specimen of type species CBS H-14862.

Mycosymbioces J.L. Frank 2014, *Leotiomycetes* genera *incertae sedis*, one species, type: *M. mycenophila* J.L. Frank [as ‘mycenaphila’], asexual morph unknown, saprobes, terrestrial, USA, see Index Fungorum (2017; taxonomy), holotype of the type: JLF2627 (OSC148294), sequences are available.

Mycothyridium Petr. 1962 (= *Xylosphaeria* G.H. Otth 1869 [1868] *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, 30 species, type: *Xylosphaeria elliptica* G.H. Otth, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Mycotodea Kirschst. 1936, *Ascomycota* genera *incertae sedis*, 14 species, type: *M. heufleri* (Niessl) Kirschst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycousteria M.L. Farr 1986 (= *Drummondia* Bat. & H. Maia 1963; = *Usteria* Bat. & H. Maia 1962 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *M. morulinae* (Bat. & H. Maia) M.L. Farr, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Mycovellosiella Rangel 1917 (= *Vellosiella* Rangel 1915; = *Walkeromyces* Thaung 1976 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *M. cajani* Henn., hyphomycetous, sexual morph unknown, plant pathogenic, see Videira et al. (2017; morphology, phylogeny, and designation of neotype), cultures and sequences are available.

Mycowinteria Sherwood 1986 (= *Trematosphaeria* subgen. *Winteria* Rehm 1881; = *Winteria* (Rehm) Sacc. 1883 *fide* Species Fungorum 2017), *Protothelenellaceae*, *Lecanoromycetes* families *incertae sedis*, three species, type: *M. anodonta* (Nyl.) Sherwood & Boise, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Myelochroa (Asahina) Elix & Hale 1987 (= *Parmelia* subsect. *Myelochroa* Asahina 1952 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 30 species, type: *M. aurulenta* (Tuck.) Elix & Hale, lichenized, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Myeloconis P.M. McCarthy & Elix 1996, *Porinaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *M. fecunda* P.M. McCarthy & Elix, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Nelsen et al. (2014a; DNA, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, cultures available for the type: MPN3C, MPN759, holotype of type: E, Coppins 5131.

Myelorrhiza Verdon & Elix 1986, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *M. antrea* Verdon & Elix, asexual morph unknown, lichenized, terrestrial, Australia, see Lumbsch et al. (2010b; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures are available but lacking for the type species.

Myelosperma Syd. & P. Syd. 1915, *Myelospermataceae*, *Xylariales*, *Sordariomycetes*, four species, type: *M. tumidum* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Myiocoprula Petr. 1955, *Ascomycota* genera *incertae sedis*, two species, type: *M. gregaria* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Mychoidea Printzen, T. Sprib. & Tønsberg 2008, *Lecanorales* genera *incertae sedis*, *Lecanoromycetes*, four species, type: *M. rufofusca* (Anzi) Printzen, T. Sprib. & Tønsberg, asexual morph unknown, lichenized, terrestrial, worldwide, see Printzen et al. (2008; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Myotisia Kubátová, M. Kolařík & Hubka 2017, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *M. cremea* Kubátová, M. Kolařík & Hubka, asexual morph unknown, on bat dropping, Czech Republic, see Crous et al. (2017; taxonomy), holotype: PRM 935803, cultures and sequences are available.

Myriangiella Zimm. 1902, *Schizothyriaceae*, *Dothideomycetes* families *incertae sedis*, five species, type: *M. orbicularis* Zimm., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Schizothyriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Myriangiopsis Henn. 1902 (= *Ascomycetella* Sacc. 1889 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, two species, type: *M. sulfurea* (G. Winter) Henn., asexual morph unknown, saprobes, terrestrial, Mexico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type species.

Myriangium Mont. & Berk. 1845 (= *Phymatodiscus* Speg. 1919 [1918]; = *Phymatosphaeria* Pass. 1875; = *Pyrenotheca* Pat. 1886 *fide* Species Fungorum 2017), *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, one species, type: *M. duriae* Mont. & Berk., asexual morph unknown, saprobes, terrestrial, Algeria, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), Dissanayake et al. (2014; morphology), Wijayawardene et al. (2014c; outline), cultures and sequences are available, holotype of the type species NL 8414.

Myriellina Höhn. 1915, *Ascomycota* genera *incertae sedis*, two species, type: *M. cydoniae* (Desm.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Papua New Guinea, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: Pl. crypt. Fr. nr. 79.

Myriochapsa M. Cáceres, Lücking & Lumbsch 2013, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *M. psoromica* (M. Cáceres, Santos de Jesus & Santos Vieira) M. Cáceres, Lücking & Lumbsch, asexual

morph unknown, lichenized, terrestrial, worldwide, see Parnmen et al. (2013; taxonomy), Lumbsch et al. (2014a, b; accepted in *Graphidaceae*), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of the type: ASE, Cáceres 7595 (holotype of *Chapsa psoromica* M. Cáceres, L.S. de Jesus & T.S. Vieira), cultures available for the type: CHAR106 (*fide* Lumbsch et al. 2014a, b).

Myriconium Syd. & P. Syd. 1912, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, c. ten species, type: *M. scirpi* Syd. & P. Syd., hyphomycetous, sexual morph unknown, in soil, saprobes, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Myriodiscus Boedijn 1935 (= *Ascotremellopsis* Teng & S.H. Ou ex S.H. Ou 1936 *fide* Species Fungorum 2017), *Tymanidaceae*, *Phacidiales*, *Leotiomycetes*, one species, type: *M. sparassoides* Boedijn, asexual morph unknown, saprobes on bamboo culms, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myriodontium Samson & Polon. 1978, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *M. keratinophilum* Samson & Polon., sexual morph unknown, human pathogen, saprobes, Spain, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Man et al. (2015; DNA), cultures and sequences are available, available specimen and cultures for the type: CBS H-7385; S7-P-3-16 (*fide* Man et al. 2015).

Myriogenospora G.F. Atk. 1894, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. paspali* G.F. Atk., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Myriolecis Clem. 1909, *Lecanoraceae*, *Lecanorales*, *Lecanomycetes*, 31 species, type: *M. sambuci* (Pers.) Clem., lichenized, worldwide, see Śliwa et al. (2012; phylogeny), Zhao et al. (2016c; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Myrionora R.C. Harris 1988, *Lecanoraceae*, *Lecanorales*, *Lecanomycetes*, one species, type: *M. albidula* (Willey) R.C. Harris, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Myriophacidium Sherwood 1974, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, six species, type:

M. aphyophyllicum Sherwood, asexual morph unknown, saprobes, terrestrial, worldwide, see Hou and Piepenbring (2009a, b; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of the type: CUP 53225, isotype of the type: OSC 34178.

Myriosclerotinia N.F. Buchw. 1947 (= *Myrioconium* Syd. & P. Syd. 1912 *fide* Jaklitsch et al. 2016a), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, ten species, type: *M. scirpicola* (Rehm) N.F. Buchw., asexual morph unknown, on roots, terrestrial, America, Europe, see Lumbsch and Huhndorf (2010; outline), Andrew et al. (2012; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Myriospora Nägeli ex Uloth 1861 (= *Myriospora* Nägeli 1853 *fide* Species Fungorum 2017), *Acarosporaceae*, *Acarosporales*, *Lecanomycetes*, nine species, type: *M. rufescens* (Ach.) Nägeli, lichenized, Europe, USA, Africa, see Arcadia and Knudsen (2012; phylogeny), Gueidan et al. (2014c; phylogeny), Westberg et al. (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Myriostigma Kremp. 1874, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, seven species, type: *M. candidum* Kremp., lichenized, see Frisch et al. (2014; phylogeny), Ertz et al. (2015b, new combination), Jagadeesh Ram and Sinha (2016; identification key), Lücking et al. (2017a; classification), sequences are available.

Myriostigmella G. Arnaud 1952 (= *Myriostigma* G. Arnaud 1925 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *M. guatteriae* (G. Arnaud) G. Arnaud, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type species.

Myriotrema Fée 1825 (= *Coscinedia* A. Massal. 1860 [1859-1860] *fide* Species Fungorum 2017), *Graphidaceae*, *Ostropales*, *Lecanomycetes*, 54 species, type: *M. olivaceum* Fée, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Mangold et al. (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Sipman et al. (2012; key, Costa Rica), Rivas Plata et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2014, 2017a; phylogeny, classification), cultures and sequences are available.

Myrmaeciella Lindau 1897 (= *Myrmaecium* Sacc. 1880 *fide* Species Fungorum 2017), *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. endoleuca* (Sacc.) Lindau, asexual morph unknown, saprobes or fungicolous, terrestrial, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Jaklitsch and Voglmayr (2011; taxonomic notes),

Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type species.

Myrmaecium Nitschke ex Fuckel 1870, *Valsariaceae*, *Valsariales*, *Dothideomycetes*, two species, type: *M. rubricosum* (Fr.) Fuckel asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2015; taxonomy, and phylogeny), cultures and sequences are available, lectotype of the type: L9118162, cultures available for the type: CBS 139067.

Myrmecomycetes Jouvenaz & Kimbr. 1991, *Ascomycota* genera *incertae sedis*, two species, type: *M. annellisiae* Jouvenaz & Kimbr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, Argentina, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of type: FLAS 55591.

Myrmecridium Arzanlou, W. Gams & Crous 2007, *Myrmecidiaceae*, *Myrmecridiales*, *Sordariomycetes*, eleven species, type: *M. schulzeri* (Sacc.) Arzanlou, W. Gams & Crous, sexual morph unknown, saprobes, terrestrial, worldwide, see Crous et al. (2012e, 2015d, 2016a; new species, phylogeny), Jie et al. (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline, accepted in *Hypocreomycetidae*), Réblová et al. (2016a; new species, phylogeny), Tibpromma et al. (2017b; new species), cultures and sequences are available for the type: CBS 134.68 = ATCC 16310.

Myrotheciastrum Abbas & B. Sutton 1988, *Ascomycota* genera *incertae sedis*, one species, type: *M. salvadorae* Abbas & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Pakistan, see Wijayawardene et al. (2012, 2016b, 2017; morphology, outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: IMI 139985i.

Myrothecium Tode 1790 (= *Godroniella* P. Karst. 1884; = *Myrotheciella* Speg. 1911; = *Myxormia* Berk. & Broome 1850; = *Starkeyomyces* Agnihothr. 1956 *fide* Species Fungorum 2017), *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, 83 species, type: *M. inundatum* Tode, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014c; notes), Maharachchikumbura et al. (2015a, b; outline), Chen et al. (2016, b, c; review, phylogeny), Dai et al. (2017; morphology, phylogeny, new species), Lombard et al. (2016; phylogeny), Tibpromma et al. (2017b; new

species), cultures and sequences are available but lacking for type, epitype of type: CBS H-14898.

Myrtapenidiella Quaedvly. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *M. tenuiramis* (Crous & Summerell) Quaedvly. & Crous, hyphomycetous, sexual morph unknown, pathogens, saprobes, terrestrial, Australia, Thailand, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; outline), Crous et al. (2015d, 2017; new species, phylogeny), sequence data available, holotype and ex-type of the type species CBS H-20253; CPC 13692.

Mystrosporiella Munjal & Kulsh. 1969, *Ascomycota* genera *incertae sedis*, two species, type: *M. litseae* Munjal & Kulsh., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myrtacremonium Crous 2017, *Niesliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *M. eucalypti* Crous, hyphomycetous, sexual morph unknown, saprobes, Australia, see Crous et al. (2017; taxonomy), holotype: CBS H-23123, cultures and sequences are available.

Mytilinidion Duby 1861 (= *Murashkinskija* Petr. 1928 *fide* Species Fungorum 2017), *Mytilinidiaceae*, *Mytilinidiales*, *Dothideomycetes*, twelve species, type: *M. aggregatum* (DC.) Duby, asexual morph septonema-like, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mytilinidiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Mytilodiscus Kropp & S.E. Carp. 1984, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *M. alnicola* Kropp & S.E. Carp., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of the type: OSC 41,244.

Mytilostoma P. Karst. 1879, *Dothideomycetes* genera *incertae sedis*, two species, type: *M. deflectens* P. Karst., asexual morph unknown, saprobes, terrestrial, Finland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Myxocephala G. Weber, Spaaij & Oberw. 1989, *Sordariomycetes* genera *incertae sedis*, one species, type: *M. setosa* (Berk. & M.A. Curtis) S. Hughes, hyphomycetous, sexual morph unknown, saprobes, Germany, see Kirk et al. (2008, 2013; genus accepted), cultures and sequences are available.

Myxocyclus Riess 1852, *Pleomassariaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *M. confluens* Riess, coelomycetous, sexual morph *Splanchnonema sensu lato*, saprobes, terrestrial, USA, Canada, Japan, see Pastirčáková and Pastirčák (2010; morphology), Wijayawardene et al. (2016b, 2017; morphology, outline), cultures and sequences are unavailable, holotype of the type species IMI 180178.

Myxoparaphysella Caball. 1941, *Ascomycota* genera *incertae sedis*, two species, type: *M. cocoës* Caball., coelomycetous, sexual morph unknown, saprobes, terrestrial, Spain, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myxophora Döbbeler & Poelt 1978, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, seven species, type: *M. amerospora* Döbbeler & Poelt, asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Myxospora L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *M. masonii* (M.C. Tulloch) L. Lombard & Crous, hyphomycetous, sexual morph unknown, from forest soil, terrestrial, cosmopolitan, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), ex-type of the type: CBS 174.73.

Myxosporella Sacc. 1881, *Ascomycota* genera *incertae sedis*, one species, type: *M. miniata* Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myxosporidiella Negru 1960, *Ascomycota* genera *incertae sedis*, one species, type: *M. fritillariae* Negru, coelomycetous, sexual morph unknown, saprobes, terrestrial, Romania, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myxostomellina Syd. 1931, *Ascomycota* genera *incertae sedis*, one species, type: *M. apoensis* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myxothyriopsis Bat. & A.F. Vital 1956 (= *Plenotrichella* Bat. & A.F. Vital 1959; = *Stigmopeltopsis* Peres 1961 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *M. astronifolia*

Bat. & A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myxothyrium Bubák & Kabát 1915, *Ascomycota* genera *incertae sedis*, one species, type: *M. leptideum* (Fr.) Bubák & Kabát, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Myxotrichum Kunze 1823 (= *Actinospira* Corda 1854; = *Campsotrichum* Ehrenb. 1818; = *Eidamella* Matr. & Dassonv. 1901; = *Myxotrichella* (Sacc.) Sacc. 1899; = *Myxotrichum* subgen. *Myxotrichella* Sacc. 1892; = *Oncidium* Nees 1823; = *Toxotrichum* G.F. Orr & Kuehn 1964 *fide* Species Fungorum 2017), *Myxotrichaceae*, *Leotiomycetes* families *incertae sedis*, twelve species, type: *M. chartarum* Kunze, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Nadsonia Sydow 1912 (= *Schizoblastosporion* Cif. 1930; = *Guilliermondia* Nadson & Konok. 1911 *fide* Species Fungorum 2017), *?Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, four species, type: *N. fulvescens* (Nadson & Konokotina) Syd., asexual reproduction is by budding, saprophytic, on exudates of various trees, soil, peat, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kurtzman and Robnett (2014; DNA, taxonomy), sequences are available.

Nadsoniella Issatsch. 1914, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *N. nigra* Issatsch., hyphomycetous, sexual morph unknown, saprobes, marine, USSR, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Nadvornikia Tibell 1984 (= *Stephanophoron* Nádv. 1942 *fide* Species Fungorum 2017), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *N. hawaiiensis* (Tuck.) Tibell, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; DNA, phylogeny), Lumbsch et al. (2014a, b; accepted in *Graphidaceae*), Lücking et al. (2017a; classification), cultures and sequences are available.

Naemacyclus Fuckel 1874 (= *Lasiostictis* Sacc. & Berl. ex Sacc. 1889; = *Lasiostictis* Sacc. & Berl. 1885 [1884-1885] *fide* Species Fungorum 2017), *Marthamycetaceae*, *Rhyismatales*, *Leotiomycetes*, six species, type: *N. pinastri* Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Helotiaceae*), Kirk et al. (2013; genus

accepted), cultures and sequences are available but lacking for the type species.

Naemosphaera P. Karst. 1888 (= *Naemosphaera* (Sacc.) Sacc. 1892; = *Sphaeronaema* subgen. *Naemosphaera* Sacc. 1884 *fide* Species Fungorum 2017), Ascomycota genera incertae sedis, one species, type: *N. rufa* P. Karst., coelomycetous, sexual morph unknown, saprobes, marine, Finland, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Naemosphaerella Höhn. 1923, Ascomycota genera incertae sedis, two species, type: *N. ceratophora* (Speg.) Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, marine, Brazil, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Naetrocymbae Körb. 1865 (= *Santessoniolichen* Tomas. & Cif. 1952; = *Santessoniomycetes* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Naetrocymbaceae*, *Dothideomycetes* families incertae sedis, six species, type: *N. fuliginea* Körb., asexual morph unknown, epiphytes, terrestrial, worldwide, see Alstrup et al. (2009; new species, USA), Knudsen and Lendemer (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Minter et al. (2015; new species), cultures and sequences are unavailable.

Naevala B. Hein 1976 (= *Naevia* Fr. 1849; = *Ocellariella* Petr. 1947 *fide* Species Fungorum 2017), *Calloriaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *N. minutissima* (Auersw.) B. Hein, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for the type: CBS 115920, CBS 115003 (Verkley and van Kempen unpub.), available specimen for type: CBS H-15904.

Naeviella Clem. 1909, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *N. paradoxa* (Rehm) Clem., asexual morph unknown, saprobes, terrestrial, Europe, see Chlebicki (2009b; endophytes, Poland), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Naeviopsis B. Hein 1976, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *N. epilobii* (P. Karst.) B. Hein, asexual morph unknown, saprobes, terrestrial, Europe, see Raitvii (2008; Magadan), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nagrajia R.F. Castañeda & W.B. Kendr. 1991, Ascomycota genera incertae sedis, one species, type: *N. havanensis* R.F. Castañeda, coelomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: INIFAT C91/10-3.

Nagrajomyces Melnik 1984, Ascomycota genera incertae sedis, one species, type: *N. dictyosporus* Melnik, coelomycetous, sexual morph unknown, saprobes, terrestrial, Russia, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Naïs Kohlm. 1962, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *N. inornata* Kohlm., asexual morph unknown, marine, freshwater, on wood, worldwide, see Sakayaroj et al. (2011a; phylogeny), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Nakaseomyces Kurtzman 2003, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, seven species, type: *N. delphensis* (van der Walt & Tscheuschner) Kurtzman, asexual reproduction is by multilateral budding, saprophytic, but several of the species are important human pathogens, on exudate of oaks, figs, human pathogens, worldwide, see Lumbsch and Huhndorf (2010; outline), Limtong et al. (2011; asexual morph as *Candida* sp.), sequences are available.

Nakataea Hara 1939 (= *Magnaporthe* R.A. Krause & R.K. Webster 1972 *fide* Zhang et al. 2016d), *Magnaporthaceae*, *Magnaporthales*, *Sordariomycetes*, c. ten species, type: *N. oryzae* (Catt.) J. Luo & N. Zhang, sexual morph formerly known in *Magnaporthe*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, sexual morph), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Nakatopsis Whitton, McKenzie & K.D. Hyde 2001, Ascomycota genera incertae sedis, two species, type: *N. malaysiana* Whitton, McKenzie & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: HKU (M) 5093.

Nakazawaea Y. Yamada, Maeda & Mikata 1994, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, ten species, type: *N. holstii* (Wickerham) Y. Yamada, K. Maeda & Mikata, asexual reproduction is by multilateral budding, saprophytic, on beetle larvae tunnels in spruce and pine trees, alpechin, cider, grapes, sauerkraut, silage, mastitis milk, embalmed human corpse, human faeces, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kaewwichian and

Limtong (2014; new species, DNA), Kurtzman and Robnett (2014; taxonomy), sequences are available.

Nalanthamala Subram. 1956 (= *Rubrinectria* Rossman & Samuels 1999 *fide* Rossman et al. 2013b), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, six species, type: *N. madreeya* Subram., sexual morph formerly known in *Rubrinectria*, saprobes, plant pathogens, terrestrial, worldwide, see Ligoxigakis et al. (2013; Greece), Hong et al. (2013; selective medium), Severn-Ellis et al. (2012; South Africa), Li et al. (2013; indoor, phylogeny), Persoh et al. (2011; EcoChip), Rossman et al. (2013b; acceptance of genus), Su et al. (2013; secondary metabolite), Summerbell et al. (2011; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, but lacking for type.

Nannfeldtia Petr. 1947, *Leptopeltidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *N. atra* Petr., asexual morph unknown, saprobes, terrestrial, Austria, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Nannizziopsis Currah 1985, *Nannizziopsidaceae*, *Orygenales*, *Eurotiomycetes*, one species, type: *N. vriesii* (Apinis) Currah, asexual morph hyphomycetous, saprobes, animal pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stchigel et al. (2013; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), type specimen and ex-type of the type: CBS H-6714; IMI 149994.

Nanomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *N. fijianus* Thaxt., asexual morph unknown, biotrophic, terrestrial, Pacific, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nanoschema B. Sutton 1980, *Ascomycota* genera *incertae sedis*, one species, type: *N. elaeocarpi* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, British Solomon Islands, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type species IMI 187436.

Nanoscypha Denison 1972, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, ten species, type: *N. tetraspora* (Seaver) Denison, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Nanostictis M.S. Christ. 1954, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *N. peltigerae* M.S. Christ., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Etayo and Sancho (2008; new species),

Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Naothrysium Bat. 1960, *Ascomycota* genera *incertae sedis*, one species, type: *N. splendidum* Bat. & J.L. Bezerra, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Narasimhella Thirum. & P.N. Mathur 1966, *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *N. poonensis* Thirum. & P.N. Mathur, asexual morph unknown, from bee hive, Venezuela, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), sequences and culture unavailable.

Natantiella Réblová 2009, *Sordariomycetes* genera *incertae sedis*, one species, type: *N. ligneola* (Berk. & Broome) Réblová, asexual morph unknown, saprobes, terrestrial, Europe, see Réblová and Štěpánek (2009; morphology), Maharachchikumbura et al. (2016b; outline), holotype and culture of the type species K 84426 (holotype of *Sphaeria ligneola*); CBS 123470.

Natantispora J. Campb., J.L. Anderson & Shearer 2003, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, three species, type: *N. retorquens* (Shearer & J.L. Crane) J. Campb., J.L. Anderson & Shearer, marine, freshwater, see Lumbsch and Huhndorf (2010; outline), Jones et al. (2015; outline, phylogeny), Liu et al. (2015a; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Natarajania Pratibha & Bhat 2006, *Stilboporaceae*, *Diaporthales* *incertae sedis*, *Sordariomycetes*, one species, type: *N. indica* Pratibha & Bhat, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Shenoy et al. (2010; phylogeny), Seifert et al. (2011; morphology), Crous et al. (2012f; phylogeny, familial position unresolved), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Natipusilla A. Ferrer, A.N. Mill. & Shearer 2011, *Natipusillaceae*, *Natipusillales*, *Dothideomycetes*, four species, type: *N. decorospora* A. Ferrer, A.N. Mill. & Shearer, saprobes, on submerged wood, Central and South America, see Raja et al. (2012; placed in *Natipusillaceae*), Hyde et al. (2013; accepted as a genus in *Natipusillaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available, holotype of type: AF236-1 (ILL 86249).

Naumovela Kravtzev 1955, *Sordariomycetes* genera *incertae sedis*, two species, type: need typification, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Naumovozyma Kurtzman 2008 (= *Naumovia* Kurtzman 2003 *fide* Species Fungorum 2017), *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *N. dairenensis* (H. Naganiishi) Kurtzman, asexual reproduction is by multilateral budding, saprophytic, soil, spoiled milk, ensiled maize, fermented vegetables, dried fruit, cecum of baboon, worldwide, see Lumbsch and Huhndorf (2010; outline), sequences are available.

Nautosphaeria E.B.G. Jones 1964, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *N. cristaminuta* E.B.G. Jones, asexual morph unknown, marine, on submerged wood, temperate, tropical, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015; outline), cultures and sequences are available.

Navicella Fabre 1879, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *N. julii* Fabre, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014f; transferred to *Melanommataceae*), Wijayawardene et al. (2014c; outline), Tian et al. (2015; accepted as a genus in *Melanommataceae*), cultures and sequences are unavailable.

Nawavia Marvanová 1980, *Chaetosphaerales* genera *incertae sedis*, type: *N. filiformis* (Nawawi) Marvanová, hyphomycetous, sexual morph unknown, saprobes, aquatic, worldwide, see Crous et al. (2009b; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Goh et al. (2014a; new species), cultures and sequences are available but lacking for the type.

Nebularia P.M. Jørg. 2014, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *N. incrassata* (P.M. Jørg.) P.M. Jørg., lichenized, Ecuador, see Ekman et al. (2014; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available.

Nectria (Fr.) Fr. 1849 (= *Tubercularia* Tode 1790 *fide* Rossman et al. 2013b; see Species Fungorum 2017 for additional synonyms), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, around 1000 species, type: *N. cinnabarina* (Tode) Fr., saprobes, pathogens, worldwide, see Lumbsch and Huhndorf (2010; outline), Hirooka et al. (2011; monograph), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Lombard et al. (2015b; phylogeny), cultures and sequences are available, epitype and ex-epitype of the type: BPI 879981; CBS 125165 = AR 4477 = CLL 7152.

Nectriella Nitschke ex Fuckel 1870 (= *Notariella* Speg. 1909; = *Stylolectriella* Höhn. 1915 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, 74 species, type: *N. fuckelii* Nitschke

ex Fuckel, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Nectriopsis Maire 1911 (= *Dasyphthora* Clem. 1909; = *Peloronectriella* Yoshim. Doi 1968 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, c. 70 species, type: *N. violacea* (J.C. Schmidt ex Fr.) Maire, asexual morph acremonium-like, saprobes, fungicolous, lichenicolous, terrestrial, worldwide, see Etayo (2010a; new species), Lumbsch and Huhndorf (2010; outline), Luo and Zhuang (2010; new species), Brackel and Zimmermann (2012; new species), Whitton et al. (2012b; new species), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Kondratyuk et al. (2016a; new species), Zare and Gams (2016; phylogeny), cultures and sequences are available.

Neelakesa Udaian & Hosag. 1992, *Pseudeurotiaceae*, *Leotiomycetes* families *incertae sedis*, three species, type: *N. lignicola* Udaian & V.S. Hosag., saprobes, submerged wood, freshwater, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nemania Gray 1821 (= *Geniculosporium* Chesters & Greenh. 1964; = *Gamosphaera* Dumort. 1822; = *Geniculosynnema* Okane & Nakagiri 2007 *fide* Réblová et al. 2016c; Wendt et al. 2017), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, 68 species, type: *N. serpens* (Pers.) Gray, asexual morph formerly named in *Geniculosporium*, saprobes or endophytes, terrestrial or marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Nematococomyces C.L. Hou, M. Piepenbr. & Oberw. 2004, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, nine species, type: *N. rhododendri* C.L. Hou, M. Piepenbr. & Oberw., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Tian et al. (2013; new species, China, phylogeny), cultures and sequences are available, holotype and culture available for type: HMAS (112); voucher HOU 469A.

Nematogonum Desm. 1834 (= *Botryocladium* Preuss 1851 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *N. aurantiacum* Desm., hyphomycetous, sexual morph unknown, mycoparasites, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nematographium Goid. 1935, *Ascomycota* genera *incertae sedis*, one species, type: *N. hippotrichoides* (Lindau) Goid., hyphomycetous, sexual morph unknown,

saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, type specimen of type species B herb. Lindau.

Nematostoma Syd. & P. Syd. 1914 (= *Ceratosperma* Speg. 1918; = *Dimeriellopsis* F. Stevens 1927 *fide* Species Fungorum 2017), *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, 13 species, type: *N. artemisiae* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Nematothecium Syd. & P. Syd. 1912 (= *Ophiogene* Petr. 1931 *fide* Species Fungorum 2017), *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *N. vinosum* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Nemozythiella Höhn. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *N. lonicerae* (Died.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neocanthostigma Boonmee, Bhat & K.D. Hyde 2014, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, three species, type: *N. fusiforme* Boonmee, Bhat & K.D. Hyde, asexual morph hyphomycetous, saprobes, terrestrial, worldwide, Boonmee et al. (2014b; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), Lu et al. (2017b, new species, phylogeny), YZ cultures and sequences are available, holotype and available ex-type cultures for the type: MFLU 11-1146; MFLUCC 11-0510 = BCC 52383 = ICMP 20073.

Neoalpakesa Punith. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *N. poae* Punith., coelomycetous, sexual morph unknown, saprobes, terrestrial, UK, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: IMI 252188.

Neoanthostomella D.Q. Dai & K.D. Hyde 2016, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *N. pseudostromatica* D.Q. Dai & K.D. Hyde, lichenicolous, asexual morph unknown, saprobes, terrestrial, Thailand, see Dai et al. (2017; phylogeny, morphology, sexual morph), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), holotype

and ex-type of type species MFLU 15-1190; MFLUCC 11-0610.

Neoquastroma Wanás., E.B.G. Jones & K.D. Hyde 2017, *Parabambusicolaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. guttulatum* Wanás. E.B.G. Jones & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Wanasinghe et al. (2017a; taxonomy, phylogeny), holotype of the type: MFLU 16-2524, cultures and sequences are available.

Neorachnotheca Ulfig, Cano & Guarro 1997, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *N. keratinophila* Ulfig, Cano & Guarro, asexual morph unknown, saprobes, on marine and river sediments, Spain, see Lumbsch and Huhndorf et al. (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: IMI 351982.

Nearbucula B. Sutton 1983 (= *Arbuscula* Bat. & Peres 1965; = *Arbusculidium* B. Sutton 1982 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *N. eugeniae* (Bat. & Peres) B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, specimens available for the type: IMI 159330, 192110, 198635, 199104, 227705a, 227705b, 296847.

Neoascochyta Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *N. exitialis* (Morini) Q. Chen & L. Cai, coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Chen et al. (2015b; taxonomy, phylogeny), Crous et al. (2016a, b; new species, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Neascotaiwania Hern.-Restr. et al. 2017, *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, two species, type: *A. terrestris* Hern.-Restr., J. Mena & Guarro, asexual morph hyphomycetous, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Neoastrophaeriella J.K. Liu, E.B.G. Jones & K.D. Hyde 2011, *Aigialaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. krabiensis* J.K. Liu, E.B.G. Jones & K.D. Hyde, asexual morph unknown, saprobes, Thailand, see Liu et al. (2011b; morphology, DNA, phylogeny), Hyde et al. (2013; accepted as a genus in *Aigialaceae*), Wijayawardene et al. (2014c; outline), holotype: MFLU 11-1148.

Neobambusicola Crous & M.J. Wingf. 2014, *Sulcatisporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. strelitziae* Crous & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Crous et al. (2014d; morphology, DNA), Tanaka et al. (2015; phylogeny),

cultures and sequences are available, holotype and ex-type of the type species CBS H-22000; CPC 24182 = CBS 138869.

Neobarclaya Kuntze 1898 (= *Barklayella* Sacc. 1892; = *Neobarclaya* Sacc. 1899 *fide* Species Fungorum 2017), Ascomycota genera *incertae sedis*, one species, type: *N. primaria* (Ellis & Everh.) Kuntze, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neobarya Lowen 1986 (= *Barya* Fuckel 1864; = *Baryella* Rauschert 1988 *fide* Species Fungorum 2017), Clavicipitaceae, Hypocreales, Sordariomycetes, ten species, type: *N. parasitica* (Fuckel) Lowen, asexual morph acremonium-like, mycoparasites, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lawrey et al. (2015; lichenicolous, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), sequence available for type species; GenBank Number: KP899626.

Neobroomella Petr. 1947, Xylariomycitidae genera *insertae sedis*, Sordariomycetes, one species, type: *N. ciliata* Petr., asexual morph unknown, saprobes, terrestrial, Syria, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Senanayake et al. (2015; accepted in Phlogocyliaceae), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Neobulgaria Petr. 1921 (= *Evulla* Kavina 1939 *fide* Species Fungorum 2017), Gelatinodiscaceae, Helotiales, Leotiomycetes, eleven species, type: *N. pura* (Pers.) Petr., asexual morph hyphomycetous (phialophora-like), saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Johnston et al. (2010; new species, New Zealand, asexual morph, DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available, available specimen and cultures for type: CBS H-15111; AFTOL-ID 1259 (Schoch unpub.).

Neocamarosporium Crous & M.J. Wingf. 2014, Pleosporaceae, Pleosporales, Dothideomycetes, four species, type: *N. goegapense* Crous & M.J. Wingf., coelomycetous, saprobes, terrestrial, worldwide, see Crous et al. (2014d; taxonomy, phylogeny), Wijayawardene et al. (2016b; outline, taxonomy), Ariyawansa et al. (2015c; phylogeny, new combination), cultures and sequences are available, holotype and ex-type strains of the type species CBS H-21722; CPC 23676.

Neocarpenteles Udagawa & Uchiy. 2002, Trichocomaceae, Eurotiales, Eurotiomycetes, one species, type: *N. acanthosporum* (Udagawa & Takada) Udagawa & Uchiy., asexual morph unknown, saprobes, terrestrial,

temperate, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available, holotype and cultures available for the type: NHL 22462; CBS 447.65, CBS 558.71.

Neocatapyrenium H. Harada 1993, Verrucariaceae, Verrucariales, Eurotiomycetes, four species, type: *N. cladonioideum* (Vain.) H. Harada, asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available but lacking for the type species, holotype of type species TUR (142).

Neocatenulostroma Quaedvl. & Crous 2014, Teratosphaeriaceae, Capnodiales, Dothideomycetes, three species, type: *N. microsporum* (Joanne E. Taylor & Crous) Quaedvl. & Crous, hyphomycetous, pathogens, saprobes, terrestrial, Sweden, Germany, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, nomenclature), holotype and ex-type of the type species PREM 56207a; CPC 1960.

Neoceratosperma Crous & Cheew. 2014, Mycosphaerellaceae, Capnodiales, Dothideomycetes, four species, type: *N. eucalypti* Crous & Cheew., hyphomycetous, sexual morph mycosphaerella-like, pathogens or associated with leaf spots, Brazil, China, Thailand, see Crous et al. (2014c; morphology, phylogeny), Wijayawardene et al. (2014c, 2017; outline), Guatimosim et al. (2016a, b; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type species CBS H-21712; CPC 23465.

Neocercospora M. Bakhshi, Arzanlou, Babai-ahari & Crous 2015, Mycosphaerellaceae, Capnodiales, Dothideomycetes, one species, type: *N. ammicola* M. Bakhshi, Arzanlou, Babai-ahari & Crous, hyphomycetous, sexual morph unknown, plant pathogenic, see Bakhshi et al. (2015a, b; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of type: IRAN 16461 F; CCTU 1186 = CBS 136450.

Neochaetosphaerella Lar. N. Vassiljeva, S.L. Stephenson & Chernyshev 2012, Nitschkiaceae, Coronophorales, Sordariomycetes, one species, type: *N. thaxteriospora* Lar. N. Vassiljeva, S.L. Stephenson & Chernyshev, asexual morphs unknown, saprobes, terrestrial, southern USA, see Vasilyeva et al. (2012; morphology), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: VLA P-2489.

Neochaetospora B. Sutton & Sankaran 1991 (= *Chaetospora* Faurel & Schotter 1965 [1964] *fide* Species Fungorum 2017), Ascomycota genera *incertae sedis*, one species, type: *N. quezelii* (Faurel & Schotter) B. Sutton &

Sankaran, coelomycetous, sexual morph unknown, coprophilous, terrestrial, Sahara, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neocladophialophora Crous & R.K. Schumach. 2014, *Eurotiomycetes* genera *incertae sedis*, one species, type: *N. quercina* Crous & R.K. Schumach., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Germany, see Crous et al. (2014c; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of type: CBS H-22003; CPC 24426.

Neoclaviceps J.F. White, Bills, S.C. Alderman & Spatafora 2001, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *N. monostipa* J.F. White, Bills, S.C. Alderman & Spatafora, asexual morph unknown, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, cultures available for the type species INBio 6-141, holotype: RUTPP INBio 6-141.

Neococcoomyces Y.R. Lin, C.T. Xiang & Z.Z. Li 1999, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, three species, type: *N. rhododendri* Y.R. Lin, C.T. Xiang & Z.Z. Li, asexual morph unknown, saprobes, terrestrial, China, see Hou and Piepenbring (2009a, b; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: AAU [Anhui] AAUFP 67.

Neocoleroa Petr. 1934, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, six species, type: *N. sibirica* Petr., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Farkas (2010; new species), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Neocordana Hern.-Restr. & Crous 2015, *Pyriulariaceae*, *Magnaportheales*, *Sordariomycetes*, five species, type: *N. musae* (Zimm.) M. Hern.-Rest. & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Hernández-Restrepo et al. (2015; taxonomy, phylogeny), Crous et al. (2016a, b; new species, phylogeny), Crous et al. (2017; new species), Wijayawardene et al. (2017; outline), neotype and ex-neotype of the type: CBS H-22054, CPC 18127.

Neocordyceps Kobayasi 1984, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *N. kohyanensis* Kobayasi, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Neocosmospora E.F. Sm. 1899, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, 25 species, type: *N. vasinfecta* E.F. Sm., asexual morph hyphomycetous (acremonium-like), saprobes, pathogens on human, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Nalim et al. (2011; new species), Hirooka et al. (2012; new species), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; new species, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Neocrinula Crous 2016, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *N. xanthorrhoeae* Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Crous et al. (2016a; new species, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22859, CPC 29474.

Neocryptospora Petr. 1959, *Sordariomycetes* genera *incertae sedis*, one species, type: *N. rickii* Petr., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Neocucurbitaria Wanás, E.B.G. Jones & K.D. Hyde 2017, *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *N. unguis-hominis* (Punith. & M.P. English) Wanás, E.B.G. Jones & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Thailand, see Wanasinghe et al. (2017b; taxonomy, phylogeny), holotype of the type: MFLU 16-2710, cultures and sequences are available.

Neocudoniella S. Imai 1941, *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *N. jezoensis* (S. Imai) S. Imai, asexual morph unknown, saprobes, terrestrial, Japan, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Neocurreya Thambug. & K.D. Hyde 2015, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. austroafricana* (Marinc., M.J. Wingf. & Crous) Thambugala & K.D. Hyde, asexual morph unknown, saprobes, cosmopolitan, see Thambugala et al. (2015b; taxonomy), Jaklitsch et al. (2016c; treated as a synonym of *Teichospora*), Wijayawardene et al. (2016b; taxonomy), cultures and sequences are available.

Neocylindroseptoria Thambug. & K.D. Hyde 2014, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, one species, type: *N. pistaciae* (Quaedvl., Verkley & Crous) Thambug. & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Spain, see Thambugala et al. (2014a; phylogeny), cultures and sequences are available, holotype and ex-type of type: CBS H-21301; CBS 471.69.

Neodactylaria Guevara-Suarez, Deanna A. Sutton, Wiederhold & Gené 2017, *Dothideomycetes* genera *incertae sedis*, one species, type: *N. obpyriformis* Guevara-Suarez, Deanna A. Sutton, Wiederhold & Gené, hyphomycetous, sexual morph unknown, from human bronchoalveolar lavage, USA, see Crous et al. (2017; taxonomy), holotype: CBS H-23131, cultures and sequences are available.

Neodasyscypha Suková & Spooner 2005 (= *Dasysscypha* Fuckel 1870 [1869-70]; = *Neodasyscypha* Spooner 1987 *fide* Species Fungorum 2017), *Lachnaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *N. cerina* (Pers.) Spooner, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Whitton et al. (2012b; on *Pandanus*), Jaklitsch et al. (2016a; classification), cultures and sequences are available, available specimen of type: CBS H-15118.

Neodeightonia C. Booth 1970, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, six species, type: *N. subglobosa* Booth, asexual morph coelomycetous, saprobes, pathogens, in soil, terrestrial, cosmopolitan, see Phillips et al. (2008, 2013; taxonomy, phylogeny), Liu et al. (2012; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c, 2016b; outline, taxonomy, phylogeny), Dissanayake et al. (2016b; taxonomy), Dai et al. (2017, new species, phylogeny), Konta et al. (2016a; new species), holotype and ex-type of the type: IMI 57769(f), CBS 448.91.

Neodeightoniella Crous & W.J. Swart 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *N. phragmiticola* Crous & W.J. Swart, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Crous et al. (2013b; morphology), cultures and sequences are available, holotype and ex-type of type: CBS H-21427; CPC 22059.

Neodevriesia Quaedvl. & Crous 2014, *Neodevriasiaceae*, *Capnodiales*, *Dothideomycetes*, 17 species, type: *N. hilliana* (Crous & U. Braun) Quaedvl. & Crous, hyphomycetous, with teratosphaeria-like sexual morph, saprobes, terrestrial, worldwide, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), Crous et al. (2015d; morphology, phylogeny, sexual morph), cultures and sequences are available, holotype and ex-type of the type: CBS H-20340; CPC 15382.

Neodidymella Phook., R.H. Perera & K.D. Hyde 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *N. thailandicum* Phook. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type of the type: MFLU11-0176, MFLUCC 11-0140.

Neodidymeliopsis Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *N. cannabis* (G. Winter) Q. Chen & L. Cai, asexual morph coelomycetous, saprobes, ?pathogens, terrestrial, cosmopolitan, see Chen et al. (2015b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), ex-type of the type: CBS 121.75.

Neodiplodina Petr. 1954, *Ascomycota* genera *incertae sedis*, one species, type: *N. verruciformis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neoeriomyopsis Crous & M.J. Wingf. 2015, *Sordariomycetes*, genera *incertae sedis*, one species, type: *N. aristata* (B. Sutton & Hodges) Crous & M.J. Wingf., coelomycetous, saprobes, Brazil, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2017; outline, phylogeny), epitype and ex-epitype of the type: CBS H-22243, CPC 25050.

Neoerysiphe U. Braun 1999, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, c. 15 species, type species: *N. galeopsidis* (DC.) U. Braun, sexual and hyphomycetous asexual morphs, obligate plant pathogens, terrestrial, almost worldwide, asexual morph: *Striatoidium* (R.T.A. Cook, A.J. Inman & C. Billings) R.T.A. Cook & U. Braun, see Takamatsu (2008a; phylogeny), Heluta et al. (2010; phylogeny), Braun and Cook (2012; taxonomy), DNA sequences available, lectotype of type: G 00122104.

Notes: *Striatoidium* was originally described as *Oidium* subgen. *Striatoidium* R.T.A. Cook, A.J. Inman & C. Billings, typified by *Oidium leucas-javanicae* Sawada, which is a synonym of *Oidium lamii* Rabenh. according to Braun and Cook (2012). This subgenus was subsequently raised to genus status by Braun and Cook (2012), but no binomial in *Striatoidium* was given to the type species. In addition, four species in *Striatoidium* which lack a *Neoerysiphe* sexual morph are known, i.e. *S. aloysiae* (S. Wolcan, S. Takam. & Havryl.) S. Wolcan, S. Takam. & Havryl., *S. baccharidis* (Havryl. & U. Braun) U. Braun & Havryl., *S. jaborosae* M.G. Cabrera & U. Braun, and *S. maquii* (Havryl. & U. Braun) U. Braun & Havryl. *Oidium lamii* is the asexual morph of *N. galeopsidis* (Braun and Cook 2012) and under unitary nomenclature, the genus *Striatoidium* should be considered a synonym of *Neoerysiphe*. *Oidium leonuri-sibiricae* Sawada is also a synonym of *O. lamii* (Braun and Cook 2012), therefore these names and *O. leucas-javanicae* need to be synonymized with *N. galeopsidis*. In addition, molecular phylogenetic studies have revealed the placement of *S. aloysiae*, *S. baccharidis*, and *S. maquii* in *Neoerysiphe*, indicating that these names need to be reallocated accordingly. The phylogenetic placement of *S. jaborosae*

is unknown and therefore no new combination can be proposed for this taxon in the current state.

***Neoerysiphe galeopsidis* (DC.) U. Braun,
Schlechtendalia 3:50. 1999**

- = *Oidium lamii* Rabenh., Flora 36:204. 1853.
- = *Oidium leucas-javanicae* Sawada, Bull. Dept. Agric. Gov. Res. Inst. Formosa 24:22. 1927
- = *Oidium leonuri-sibiricae* Sawada, loc. cit.:24

Additional synonyms of *N. galeopsidis* are listed in Braun and Cook (2012).

***Neoerysiphe aloysiae* (S. Wolcan, S. Takam. & Havryl.) S. Takam., comb. nov.**

IndexFungorum number: IF553242

Basionym: *Oidium aloysiae* S. Wolcan, S. Takam. & Havryl., in Takamatsu et al. Mycol. Res. 112:644, 2008.

≡ *Striatoidium aloysiae* (S. Wolcan, S. Takam. & Havryl.) S. Wolcan, S. Takam. & Havryl., in U. Braun & R.T.A. Cook, Taxonomic Manual of the *Erysiphales* (Powdery Mildews): 347. 2012.

***Neoerysiphe baccharidis* (Havryl. & U. Braun) S. Takam., comb. nov.**

IndexFungorum number: IF553243

Basionym: *Oidium baccharidis* Havryl. & U. Braun, Nova Hedw. 66(3–4):515. 1998.

≡ *Striatoidium baccharidis* (Havryl. & U. Braun) U. Braun & Havryl., Taxonomic Manual of the *Erysiphales* (Powdery Mildews):348. 2012.

***Neoerysiphe maquii* (Havryl. & U. Braun) S. Takam., comb. nov.**

IndexFungorum number: IF553244

Basionym: *Oidium maquii* Havryl. & U. Braun, Nova Hedw. 66(3–4):518. 1998.

≡ *Striatoidium maquii* (Havryl. & U. Braun) U. Braun & Havryl., Taxonomic Manual of the *Erysiphales* (Powdery Mildews):349. 2012.

***Neofabraea* H.S. Jacks. 1913 (= *Allantozythia* Höhn. 1924; = *Phlyctema* Desm. 1847 fide Johnston et al. 2014b), *Dermateaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *N. malicorticis* H.S. Jacks., asexual morph hyphomycetous, saprobes, pathogens, terrestrial, worldwide, see Raitivir (2008; Magadan), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature, new combination), Crous et al. (2015e; new species), de Jong et al. (2015; phylogeny), Chen et al. (2016b; new species), cultures and sequences are available, cultures available for the type: DAOM 227090, neotype of type: CBS H-22219.**

***Neoflageoletia* J. Reid & C. Booth 1966, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *N. bambusina* (Syd.) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al.**

(2015b, 2016b; outline), cultures and sequences are unavailable.

***Neofracchiaea* Teng 1938, *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *N. callista* (Berk. & M.A. Curtis) Teng, asexual morph unknown, saprobes, see Mugambi and Huhndorf (2010; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.**

***Neofuckelia* Zeller & Goodd. 1935, *Ascomycota* genera *incertae sedis*, one species, type: *N. pinicola* (Zeller & Goodd.) Zeller & Goodd., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.**

***Neofusicoccum* Crous, Slippers & A.J.L. Phillips 2006, *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, c. 29 species, type: *N. parvum* (Pennycook & Samuels) Crous, Slippers & A.J.L. Phillips, coelomycetous (with dichomera-like synasexual morphs), saprobes, pathogens, terrestrial, cosmopolitan, see Phillips et al. (2008, 2013; taxonomy, phylogeny), Pavlic et al. (2009; new species), Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Slippers et al. (2013; phylogeny), Yan et al. (2013; species associated with grape wine), Wijayawardene et al. (2014c, 2017; outline, taxonomy, phylogeny), Dissanayake et al. (2016b; taxonomy), holotype and ex-type of the type: PDD 45438, ATCC 58191.**

***Neogaeumannomyces* D.Q. Dai & K.D. Hyde 2015, *Magnaporthaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *N. bambusicola* D.Q. Dai & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Liu et al. (2015a; phylogeny, taxonomy), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available, holotype and ex-type of type: MFLU 14–0822; MFLUCC 11–0390.**

***Neogymnomycetes* G.F. Orr 1970, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *N. demonbreunii* (Ajello & S.L. Cheng) G.F. Orr, asexual morph unknown, saprobes, Spain, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available, available specimen and cultures for the type: CBS H-7197; CBS 233.85.**

***Neohaplomyces* R.K. Benj. 1955, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *N. medonalis* R.K. Benj., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.**

Neohelicomyces Z.L. Luo, D.J. Bhat & K.D. Hyde 2017, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, two species, type: *N. aquaticus* Z.L. Luo, Baht & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, see Luo et al. (2017; taxonomy, phylogeny), holotype of the type: MFLU 16-2543, cultures and sequences are available.

Neohendersonia Petr. 1921, *Neohendersoniaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *N. piriformis* (G.H. Otth) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neoheppia Zahlbr. 1909, *Peltulaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *N. brasiliensis* Zahlbr., asexual morph unknown, saprobes, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neoheteroceras Nag Raj 1993 (= *Heteroceras* Sacc. 1915 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *N. flageoletii* (Sacc.) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, France, Japan, see Yonezawa and Tanaka (2008; morphology, new species), Wijayawardene et al. (2012, 2016b, 2017; morphology, outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: PAD 940.

Neohortaea Quaedvli. & Crous 2014, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *N. acidophila* (Hölker, Bend, Pracht, Tetsch, Tob. Müll., M. Höfer & de Hoog) Quaedvli. & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Germany, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), cultures and sequences are available, ex-type strain of type: CBS 113389.

Neohypodiscus J.D. Rogers, Y.M. Ju & Læssøe 1994 (= *Hypodiscus* Lloyd 1923 *fide* Species Fungorum 2017), *Boliniaceae*, *Boliniales*, *Sordariomycetes*, three species, type: *N. rickii* (Lloyd) J.D. Rogers, Y.M. Ju & Læssøe, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Senanayake et al. (2015; accepted in *Xylariomycetidae*), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Neoidriella Hern.-Restr. & Crous 2015, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *N. desertorum* (Nicot & Mouch.) Hern.-Restr. & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Egypt, see Hernández-Restrepo et al. (2016b; taxonomy, phylogeny), Maharachchikumbura et al.

(2015b, 2016b; outline), cultures and sequences are available.

Neojohnstonia B. Sutton 1983 (= *Johnstonia* M.B. Ellis 1971 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *N. colocasiae* (M.B. Ellis) B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type was not designated but several specimens are at IMI and CBS (CBS H-4370).

Neokalmusia Kaz. Tanaka, Ariyaw. & K.D. Hyde 2014, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *N. brevispora* (Nagas. & Y. Otani) Kaz. Tanaka, Ariyaw. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Japan, Thailand, see Ariyawansa et al. (2014e; taxonomy, phylogeny), Dai et al. (2015a, 2017; taxonomy, phylogeny), Tanaka et al. (2015; taxonomy, phylogeny), Thambugala et al. (2017b, phylogeny, new species), cultures and sequences are available.

Neolamya Theiss. & Syd. 1918 (= *Lamyella* Berl. 1899 *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, two species, type: *N. peltigerae* (Mont.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, Brazil, see Kocourková (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Neolecta Speg. 1881, *Neolectaceae*, *Neolectales*, *Neolectomycetes*, three species, type: *N. flavovirescens* Speg., asexual morph unknown, saprobes, terrestrial, Brazil, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available but lacking for the type species.

Neoleptosphaeria Ariyawansa & K.D. Hyde 2015, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. rubefaciens* (Togliani) Ariyawansa & K.D. Hyde, asexual morph unknown, saprobes, Europe, see Ariyawansa et al. (2015b; taxonomy), Wanasinghe et al. (2016c; new species), cultures and sequences are available.

Neoligniella Naumov 1951, *Ascomycota* genera *incertae sedis*, four species, type: *N. immersa* Naumov, coelomycetous, sexual morph unknown, saprobes, terrestrial, former USSR, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neolinocarpon K.D. Hyde 1992, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, ten species, type: *N. globosicarpum* K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; treated as a genus in *Sordariomycetes*), Maharachchikumbura et al.

(2015b, 2016b; outline), cultures and sequences are available.

Neolophiostoma S. Boonmee & K.D. Hyde 2015, *Halothiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. pigmentatum* Boonmee & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 10-0062, MFLUCC 10-0129.

Neomassaria Mapook, Camporesi & K.D. Hyde 2016, *Massariaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. fabacearum* Mapook, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Hyde et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 16-1875, MFLUCC 14-1117.

Neomassariosphaeria Y. Zhang bis, J. Fourn. & K.D. Hyde 2009, *Lindgomycetaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. typhicola* (P. Karst.) Y. Zhang ter, J. Fourn. & K.D. Hyde, asexual morph unknown, saprobes, aquatic, cosmopolitan, see Zhang et al. (2009d; taxonomy, accepted as a genus in *Amniculicolaceae*), Hyde et al. (2013; accepted as a genus in *Amniculicolaceae*), Ariyawansa et al. (2015a; accepted as a genus in *Lindgomycetaceae*), Jones et al. (2015; outline, accepted as a genus in *Amniculicolaceae*), cultures and sequences are unavailable.

Neomelanconium Petr. 1940, *Ascomycota* genera *incertae sedis*, three species, type: *N. gelatosporum* (Zimm.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, former USSR, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy, new species), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neomicrosphaeropsis Thambug., Camporesi & K.D. Hyde 2016, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *N. italica* Thambug., Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Thambugala et al. (2017a, b; taxonomy, phylogeny), Hyde et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 16-0674, MFLUCC 16-0284.

Neomunkia Petr. 1947, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *N. sydowii* Petr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; phylogeny), cultures and sequences are available.

Neomycosphaerella Crous 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *N. pseudopentameridis* Crous, asexual morph unknown, foliicolous, South Africa, see Crous et al. (2013b;

morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), cultures and sequences are available, holotype and ex-type of type: CBS H-21416; CPC 21127.

Neomyrothecium L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *N. humicola* L. Lombard & Crous, hyphomycetous, sexual morph unknown, from soil, terrestrial, Papua New Guinea, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), holotype and ex-type of the type: CBS H-22456, CBS 310.96.

Neonectria Wollenw. 1917 (= *Heliscus* Sacc. 1880; = *Cylindrocarpon* Wollenw. 1913 *fide* Rossman et al. 2013b, 2016a; see Species Fungorum 2017 for additional synonyms), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, 28 species, type *N. ramulariae* Wollenw., on bark, sometimes causing cankers, cosmopolitan, mostly in temperate regions, see Chaverri et al. (2011; phylogeny), Rossman et al. (2013b; *Neonectria* protected against *Cylindrocarpon*), Lombard et al. (2015b; synonymised *Heliscus* under *Neonectria*), Maharachchikumbura et al. (2015b, 2016b; phylogeny), Rossman et al. (2016a; nomenclature), cultures and sequences are available, cultures available for type: CBS 151.29 =IMI 113894 =MUCL 28083), holotype of the type species B.

Neooccultibambusa Doilom & K.D. Hyde 2016, *Occultibambusaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. chiangraiensis* Doilom & K.D. Hyde, saprobes, terrestrial, see Jayasiri et al. (2016, b, c; new species), Doilom et al. (2017; phylogeny), cultures and sequences are available, holotype and ex-type strains of type: MFLU 15-3531; MFLUCC 12-0559, MKT 075, ICMP 21165.

Neophiosphaerella Kaz. Tanaka & K. Hiray. 2015, *Lentiteliaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. sasicola* (Nagas. & Y. Otani) Kaz. Tanaka & K. Hiray., asexual morph unknown, saprobes, terrestrial, Japan, see Tanaka et al. (2015; morphology, phylogeny), cultures and sequences are available, holotype, epitype and ex-epitype of the type: TMI 3176, HHUF 29443; CBS 120247.

Neovularia U. Braun 1992, *Ascomycota* genera *incertae sedis*, six species, type: *N. nomuriana* (Sacc.) U. Braun (≡ *Tuberculina nomuriana* Sacc.), asexual morphs hyphomycetous, terrestrial, plant pathogenic, leaf-spotting, Asia, Europe, North America, see Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: Nomura (PAD).

Neoparodia Petr. & Cif. 1932, *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *N. ekmani* Petr. & Cif., asexual morph chuppia-like or sarcinella-like, saprobes, terrestrial, West Indies, see Lumbsch and Huhndorf (2010; outline), Hyde et al.

(2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Neopeckia Sacc. 1883 (= *Didymotrichia* Berl. 1893 *fide* Species Fungorum 2017), *Dothideomycetes* genera incertae sedis, one species, type: *N. coulteri* (Peck) Sacc., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but unavailable for type species.

Neopeltella Petr. 1950, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, three species, type: *N. aequatoriensis* Petr., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline, accepted in *Schizothyriaceae*), Hyde et al. (2013, accepted in *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Schizothyriaceae*), Phookamsak et al. (2016; placed in *Micropeltidaceae*), cultures and sequences are unavailable.

Neopeltis Syd. 1937, *Ascomycota* genera incertae sedis, three species, type: *N. brenesii* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, central America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neopenidiella Quaedvl. & Crous 2014, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *N. nectandrae* (Crous, U. Braun & R.F. Castañeda) Quaedvl. & Crous, hyphomycetous, sexual morph unknown, foliicolous, terrestrial, Cuba, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), cultures and sequences are available, holotype and ex-type culture of type: INIFAT C87/45; CBS 734.87.

Neopericonia Kamal, A.N. Rai & Morgan-Jones 1983, *Ascomycota* genera incertae sedis, three species, type: *N. indica* Kamal, A.N. Rai & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neapestalotiopsis Maharachch., K.D. Hyde & Crous 2014, *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, 27 species, type: *N. protearum* (Crous & L. Swart) Maharachch., K.D. Hyde & Crous, coelomycetous, sexual morph unknown, saprobes, pathogens on plants, worldwide, see Maharachchikumbura et al. (2014b, 2016c; phylogeny, morphology), Jayawardena et al. (2016a; new species), Wijayawardene et al. (2016b; outline, taxonomy), sequence data available for all species, holotype and ex-type of the type: PREM 56186; CBS 114178.

Neopetromyces Frisvad & Samson 2000, *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, one species, type: *N. muricatus* (Udagawa, Uchiy. & Kamiya) Frisvad & Samson, asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available, holotype and ex-type strain of type species CBM BF-42515; NRRL 35674.

Neophacidium Petr. 1950, *Rhytidomycetidae*, *Rhytidomycetales*, *Leotiomycetes*, two species, type: *N. macrocarpum* (Pat.) Petr., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neophaeococcomyces Crous & M.J. Wingf. 2015, *Strelitzianaceae*, *Chaetothyriales*, *Eurotiomycetes*, two species, type: *N. aloes* (Crous & M.J. Wingf.) Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, South Africa, Switzerland, see Crous et al. (2015e; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-21441, CPC 21873.

Neophaeocryptopus Wanas., Camporesi, E.B.G. Jones & K.D. Hyde 2016, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, one species, type: *N. cytisi* Wanas., Camporesi, E.B.G. Jones & K.D. Hyde, asexual morph coelomycetous, sexual morph bitunicate ascomycetous, saprobes, terrestrial, Europe, see Li et al. (2016a; taxonomy, phylogeny), holotype and ex-type strain of type: MFLU 15-3542; MFLUCC 14-0970.

Neophaeomoniella Rooney-Latham & Crous 2015, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, three species, type: *N. eucalypti* Rooney-Latham & Crous, coelomycetous, sexual morph unknown, on living leaves, endophytic, terrestrial, USA, South Korea, South Africa, see Crous et al. (2015d; morphology, phylogeny), cultures and sequences are available, holotype and ex-type culture of type: CBS H-22249; CPC 25161.

Neophaeosphaeria M.P.S. Câmara, M.E. Palm & A.W. Ramaley 2003, *Neophaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *N. filamentosa* (Ellis & Everh.) M.P.S. Câmara, M.E. Palm & A.W. Ramaley, asexual morph coelomycetous, saprobes or pathogenic, terrestrial, Mexico, USA, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Leptosphaeriaceae*), Crous et al. (2013b; taxonomy), Ariyawansa et al. (2015a; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), sequence data available for all species.

Neophaethocoidea Quaedvl. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *N. proteae* (Crous) Quaedvl. & Crous, hyphomycetous, saprobes, terrestrial, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; outline), cultures and

sequences are available, holotype and ex-type of the type: CBS H-20092; CPC 2828.

Neophoma Petr. & Syd. 1927, *Ascomycota* genera *incertae sedis*, two species, type: *N. graminella* (Sacc.) Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America., see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neophyllis F. Wilson 1891 (= *Phyllis* F. Wilson 1889 *fide* Species Fungorum 2017), *Sphaerophoraceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *N. melacarpa* F. Wilson, asexual morph unknown, lichenized, terrestrial, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available for the type species, holotype of type species available.

Neophysalospora Crous & M.J. Wingf. 2014, *Clypeophysalosporaceae*, *Amphisphaerales*, *Sordariomycetes*, one species, type: *N. eucalypti* Crous & M.J. Wingf., asexual morph coelomycetous, saprobes, terrestrial, Mozambique, South Africa, see Crous et al. (2014d; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), Giraldo et al. (2017; phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Neoplaconema B. Sutton 1977, *Ascomycota* genera *incertae sedis*, two species, type: *N. napelli* (Maire & Sacc.) B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, Germany, see Wijayawardene et al. (2012, 2017; outline), sequences are available, holotype of the type species was not designated (IMI 24787).

Neoplatysporoides Crous & M.J. Wingf. 2015, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. aloicola* Crous & M.J. Wingf., asexual morph coelomycetous, saprobes, Tanzania, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2017; outline, phylogeny), holotype and ex-type of the type: CBS H-22230, CPC 24435.

Neopodoconis Rifai 2008, *Ascomycota* genera *incertae sedis*, three species, type: *N. ampullacea* (Petch) Rifai, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Indonesia, see Rifai (2008; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, holotype of type species available.

Neopseudocercospora Crous 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *N. terminaliae* Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Zambia,

type, see Crous et al. (2013b; morphology), Braun et al. (2014b; morphology, keys), cultures and sequences are available, holotype and ex-type of the type: CBS H- 21431; CPC 22685.

Neopseudocercospora Videira & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *N. capsellae* (Ellis & Everh.) Videira & Crous, hyphomycetous, sexual morph mycosphaerella-like, plant pathogenic on *Brassica* species, USA, see Videira et al. (2016; morphology, phylogeny), cultures and sequences are available, holotype and isotype of type species NY 883641; BPI 399944.

Neopseudolachnella A. Hashim. & Kaz. Tanaka 2015, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, three species, type: *N. acutispora* A. Hashim., Sat. Hatak. & Kaz. Tanaka, coelomycetous, sexual morph unknown, saprobes, Japan, see Hashimoto et al. (2015b; morphology, phylogeny), sequences are available, holotype and ex-type strain of type: HHUF 29727; MAFF 244358.

Neopsoromopsis Gyeln. 1940, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *N. argentina* Gyeln., asexual morph unknown, lichenized, terrestrial, Argentina, see Lumbsch and Huhndorf (2010; outline), Thell et al. (2012; review), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Neopyricularia Klaubauf, Lebrun & Crous 2014, *Pyriculariaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *N. commelinicola* (M.J. Park & H.D. Shin) Klaubauf, Lebrun & Crous, asexual morph unknown, saprobes, terrestrial, South Korea, see Klaubauf et al. (2014; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and available cultures for the type: KUS (F) 22838; CBS 128308 =KACC 43081.

Neoramichloridium Phook., Thambug. & K.D. Hyde 2017, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *N. bambusicola* Thambug., Phook. & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, Thailand, see Thambugala et al. (2017a, 2017b; taxonomy, phylogeny), holotype of the type: MFLU 16-2278.

Neoramularia U. Braun 1991, *Ascomycota* genera *incertae sedis*, nine species, type: *N. eurotiae* (Gamalizk.) U. Braun, hyphomycetous, sexual morph unknown, terrestrial, plant pathogenic, leaf-spotting, Asia, Europe, North America, see Hyde et al. (2011; checklist), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: LE 41968.

Neorehmia Höhn. 1902, *Trichosphaeriaceae*, *Trichosphaeriales*, *Sordariomycetes*, two species, type: *N. ceratophora* Höhn., asexual morph tritirachium-like, saprobes, Austria, see Lumbsch and Huhndorf (2010;

outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Neorhamphoria Boonmee, E. Hüseyin & F. Selçuk 2016, *Tubeufiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *N. garethjonesii* Boonmee, E. Hüseyin & F. Selçuk, asexual morph unknown, saprobes, Turkey, see Boonmee et al. (2016b; taxonomy), holotype: KE 201, cultures and sequences are available.

Neoroussoella J.K. Liu, Phook. & K.D. Hyde 2014, *Roussellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. bambusae* Phook., J.K. Liu & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Asia, see Liu et al. (2014a; taxonomy, phylogeny), Hyde et al. (2016; outline), cultures and sequences are available.

Neosartorya Malloch & Cain 1973 (= *Hemisartorya* J.N. Rai & H.J. Chowdhery 1976 [1975]; = *Saitoa* Rajendran & Muthappa 1980 *fide* Species Fungorum 2017), *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, 15 species, type: *N. fischeri* (Wehmer) Malloch & Cain, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, available specimen of type: CBS H-15123.

Neoscolecobasidium Crous 2016, *Microthyriales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *N. agapanthi* Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type of the type: CBS H-22870, CPC 28778.

Neoscytalidium Crous & Slippers 2006, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, two species, type: *N. dimidiatum* (Penz.) Crous & Slippers, coelomycetous, sexual morph unknown, saprobes, pathogen, tropical and sub-tropical regions of Europe, Africa, Asia, North and South America, see Liu et al. (2012b; phylogeny), Phillips et al. (2013; morphology, phylogeny, new comb. for type species), Huang et al. (2016a; new species), cultures and sequences are available, isotype and ex-isotype strain of the type species CBS H-7745; CBS 145.78.

Neoseptoria Quaedvl., Verkley & Crous 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *N. caricis* Quaedvl., Verkley & Crous, coelomycetous, sexual morph unknown, saprobes, terrestrial, Netherlands, see Quaedvlieg et al. (2013; morphology, phylogeny), sequence data available, holotype and ex-type of the type: CBS H-21293; CBS 135097.

Neoseptorioides Crous, J. Edwards & I.G. Pascoe 2015, *Saccharataceae*, *Botryosphaeriales*, *Dothideomycetes*, two species, type: *N. eucalypti* Crous, J. Edwards & I.G. Pascoe, coelomycetous, sexual morph unknown, saprobes, Australia, see Crous et al. (2015e; taxonomy, phylogeny),

Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22395, CPC 25529.

Neosergipea M. Cáceres, Ertz & Aptroot 2016, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *N. aurata* (M. Cáceres, Ertz & Aptroot) Lücking, M. Gut. & Moncada, lichenized, terrestrial, tropical, see Lücking et al. (2016a, 2017b; nomenclature), cultures and sequences are available, holotype of type species ISE, Cáceres & Jesus 12539 (as *Sergipea aurata* M. Cáceres, Ertz & Aptroot).

Neosetophoma Gruyter, Aveskamp & Verkley 2010, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *N. samararum* (Desm.) Gruyter, Aveskamp & Verkley, asexual morph coelomycetous, saprobes and plant endophytes, terrestrial, aquatic, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Li et al. (2015c; phylogeny), Phookamsak et al. (2014c; phylogeny), Tibpromma et al. (2017b; new species), Thambugala et al. (2017b, new species, phylogeny), cultures and sequences are available, holotype and ex-type strains of type: CBS H-20319; CBS 138.96 =PD 82/653.

Neospegazzinia Petr. & Syd. 1936, *Ascomycota* genera *incertae sedis*, two species, type: *N. pulchra* Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neosphaerellopsis Crous & Trakun. 2014, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. thailandica* Crous & Trakun., on uredinio rust sori on leaves of *Bothriochloa bladhii*, terrestrial, Thailand, see Trakunyingcharoen et al. (2014; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type: CBS H-21847; CPC 21659.

Neosporidesmium Mercado & J. Mena 1988, *Ascomycota* genera *incertae sedis*, two species, type: *N. maestrense* Mercado & J. Mena, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Ma et al. (2011b; new species), Mel'nik and Braun (2013; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Prasher and Verma (2014; new species, India), Mel'nik et al. (2016; new species), cultures and sequences are unavailable.

Neostagonospora Quaedvl., Verkley & Crous 2013, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. caricis* Quaedvl., Verkley & Crous, coelomycetous, sexual morph unknown, saprobes, terrestrial, Netherlands, South Africa, see Quaedvlieg et al. (2013; morphology, phylogeny), Phookamsak et al. (2014c; phylogeny), Thambugala et al. (2017b, new species, phylogeny), cultures and sequences are available,

holotype and ex-type of the type: CBS H-21306; CBS 135092.

Neostrelitziana Crous & M.J. Wingf. 2015, *Chaetothyriales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *N. acacigena* Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, Malaysia, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22232, CPC 24873.

Neosulcatispora Crous & M.J. Wingf. 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. agave* Crous & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, France, see Crous et al. (2015e, 2016b; new species, phylogeny), Wijayawardene et al. (2016b, 2017; outline, phylogeny), holotype and ex-type of the type: CBS H-22404, CPC 26407.

Neotapesia E. Müll. & Hüttner 1963, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *N. graddonii* E. Müll. & Hüttner, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Neotestudina Segretain & Destombes 1961 (= *Pseudodelitschia* J.N. Kapoor, Bahl & S.P. Lal 1976 [1975]; = *Pseudophaeotrichum* Arx, E. Müll. & C. Stoll 1969 *fide* Species Fungorum 2017), *Testudinaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *N. rosatii* Segretain & Destombes, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Testudinaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are available, available specimen and cultures for the type: CBS H-200; CBS 690.82.

Neothyridaria Petr. 1934, *Sordariomycetes* genera *incertae sedis*, one species, type: *N. moravica* Petr., asexual morph unknown, saprobes, terrestrial, Europe, see Kocourková (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Neothyronectria Crous & Thangavel 2016, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *N. sophorae* Crous & Thangavel, coelomycetous, sexual morph unknown, saprobes, terrestrial, New Zealand, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type of the type: CBS H-22880, CPC 29690.

Neotorula Ariyaw., Z.L. Luo & K.D. Hyde 2016, *Torulaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *N. aquatica* Z.L. Luo & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, aquatic, worldwide, see Hyde et al. (2016; new species), Su et al. (2016c;

taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Neotrematosphaeria Thambug., Kaz. Tanaka & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *N. biappendiculata* (Kaz. Tanaka, Y. Harada & M.E. Barr) Thambug., Kaz. Tanaka & K.D. Hyde, asexual morph pleurophomopsis-like, saprobes, freshwater, Japan, see Thambugala et al. (2015b; morphology, phylogeny), sequence data available, holotype and ex-type of type species HHUF 28026; MAFF 239297.

Neotrimmatostroma Quaedv. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *N. excentricum* (B. Sutton & Ganapathi) Quaedv. & Crous, with hyphomycetous asexual morph, foliicolous, terrestrial, Australia, New Zealand, see Quaedvlieg et al. (2014; morphology, phylogeny), Crous et al. (2016b; new species, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype of sexual morph of the type: CBS H-19829, ex-type strain of the type: CPC 13092.

Neotrotteria Sacc. 1921, *Nitschiaceae*, *Coronophorales*, *Sordariomycetes*, two species, type: *A. pulchella* Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Mugambi and Huhndorf (2010; DNA), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Neotruncatella Hyang B. Lee & T.T.T. Nguyen 2016, *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, one species, type: *N. endophytica* Hyang B. Lee, P.M. Kirk, K.D. Hyde, Maharachch., & T.T.T. Nguyen, coelomycetous, sexual morph unknown, endophytes, terrestrial, Korea, see Hyde et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 15-0399, EMLAS5-1.

Neottiella Cooke Sacc. 1889 (= *Neottiopeziz* Clem. 1903; = *Peziza* subgen. *Neottiella* Cooke 1879; = *Peziza* subgen. *Scypharia* Quél. 1886; = *Scypharia* (Quél.) Quél. 1886 *fide* Species Fungorum 2017), *Pezizales* genera *incertae sedis*, *Pezizomycetes*, 13 species, type: *N. albocincta* (Berk. & M.A. Curtis) Sacc., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Neottiospora Desm. 1843 (= *Samukuta* Subram. & K. Ramakr. 1957; = *Spilomyces* Petr. & Syd. 1927 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *N. caricum* Desm., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Neottiosporina Subram. 1961 (= *Sadasivanella* Agnihothr. 1964 *fide* Species Fungorum 2017), *Dothideomycetes*, genera *incertae sedis*, eleven species, type: *N. apoda* (Speg.) Subram., sexual morph unknown, saprobes, terrestrial, Asia, Australia, South America, see de Gruyter et al. (2009; phylogeny), Wijayawardene et al. (2012, 2016b, 2017; outline), Trakunyingcharoen et al. (2014; phylogeny), cultures and sequences are available for one species (*N. paspali*).

Neotyphodium Glenn, C.W. Bacon & Hanlin 1996, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *N. coenophialum* (Morgan-Jones & W. Gams) Glenn, C.W. Bacon & Hanlin, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, available specimen and culture for type: CBS H-3511; CBS 494.82.

Neournula Paden & Tylutki 1969, *Chorioactidaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *N. nordmanensis* Paden & Tylutki, asexual morph unknown, saprobes, terrestrial, USA, Europe, North Africa, see Pfister et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), cultures and sequences are available but lacking for the type species, holotype of type: WSP 56281.

Neoventuria Syd. & P. Syd. 1919 (= *Venturiella* Speg. 1909 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, two species, type: *N. argentinensis* (Speg.) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Neozythia Petr. 1958 (= *Sclerozythia* Petr. 1955 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *N. nectrioides* (Petr.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Iran, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Nephroma Ach. 1809 (= *Dermatodea* Vent. 1799; = *Nephromomyces* E.A. Thomas 1939; = *Nephromiomyces* E.A. Thomas ex Cif. & Tomas. 1953; = *Nephromium* Nyl. 1860; = *Opisteria* (Ach.) Vain. 1909; = *Ornatinephroma* Gyeln. 1934; = *Peltidea* subdiv. *Opisteria* Ach. 1803 *fide* Species Fungorum 2017), *Nephromataceae*, *Peltigerales*, *Lecanoromycetes*, 36 species, type: *N. polaris* (Ach.) Ach., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al.

(2017a; classification), cultures and sequences are available.

Nephromopsis Müll. Arg. 1891 (= *Cetrariopsis* Kurok. 1980; = *Flavocetrariella* D.D. Awasthi 2007; = *Kaernefeltia* Thell & Goward 1996; = *Masonhalea* Kärnefelt 1977; = *Tuckermanella* Essl. 2003; = *Tuckermanopsis* Gyeln. 1933 *fide* Divakar et al. 2017; Lücking et al. 2017a), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 24 species, type: *N. stracheyi* (C. Bab.) Müll. Arg., asexual morph unknown, lichenized, terrestrial, Argentina, see Thell et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Randlane et al. (2013; new species), Lücking et al. (2017a; classification), Divakar et al. (2017; phylogeny), sequences are available.

Neptunella K.L. Pang & E.B.G. Jones 2003, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *N. longirostris* (Cribb & J.W. Cribb) K.L. Pang & E.B.G. Jones, asexual morph unknown, saprobes, marine, Australia, see Lumbsch and Huhndorf (2010; outline), Jones et al. (2015; review), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Nereiospora E.B.G. Jones, R.G. Johnson & S.T. Moss 1983, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *N. comata* (Kohlm.) E.B.G. Jones, R.G. Johnson & S.T. Moss, asexual morph hyphomycetous, on wood, marine, worldwide, see Sakayaroj et al. (2011a; phylogeny), Jones et al. (2015; outline), Maharachchikumbura et al. (2015a, b; outline), cultures and sequences are available.

Neta Shearer & J.L. Crane 1971, *Ascomycota* genera *incertae sedis*, two species, type: *N. patuxentica* Shearer & J.L. Crane, hyphomycetous, sexual morph unknown, saprobes, terrestrial and aquatic, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: ILLS 34520.

Neurospora Shear & B.O. Dodge 1927 (= *Anixiella* Saito & Minoura 1948; = *Chrysonilia* Arx 1981 *fide* Réblová et al. 2016c), *Sordariaceae*, *Sordariales*, *Sordariomycetes*, 52 species, type: *N. sitophila* Shear & B.O. Dodge, asexual morph formerly named in *Chrysonilia*, saprobes, terrestrial, worldwide, see Villalta et al. (2009; new species), Lumbsch and Huhndorf (2010; outline), Nygren et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available for type species.

Nevesia P.M. Jørg, L. Lindblom, Wedin & S. Ekman 2014, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, one species, type: *N. sampaiana* (Tavares) P.M. Jørg., L. Lindblom, Wedin & S. Ekman, lichenized, terrestrial, worldwide, see Ekman et al. (2014; taxonomy), Lücking

et al. (2017a; classification), cultures and sequences are available.

Nidulispora Nawawi & Kuthub. 1990, *Ascomycota* genera *incertae sedis*, one species, type: *N. quadrifida* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, aquatic, Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Niebla Rundel & Bowler 1978 (see Species Fungorum 2017 for synonyms), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, 22 species, type: *N. homalea* (Ach.) Rundel & Bowler, lichenized, New World, see Lumbsch and Huhndorf (2010; outline), Sérusiaux et al. (2010; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Niesslia Auersw. 1869 (= *Lohwagiella* Petr. 1970 [1969]; = *Nitschka* Nitschke ex Fuckel 1870 [1869-70] *fide* Species Fungorum 2017), *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, 42 species, type: *N. chaetomium* (Corda) Auersw., asexual morph unknown, lichenicolous, parasitizing members of *Peltigerales*, terrestrial, Australia, Asia and Europe, see Etayo and Sancho (2008; three new species), Motiejūnaitė and Czyżewska (2008; Poland), Zhurbenko and Dillman (2010; Alaska), Etayo et al. (2013a; new species), Himelbrant et al. (2014; Russia), Judith et al. (2015; phylogeny), Zhurbenko and Pino-Bodas (2015; New Zealand), sequence data available for *Niesslia exilis*.

Nigrocornus Ryley & Langdon 2003, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *N. scleroticus* (Pat.) Ryley, asexual morph contains simple conidiophores, endophytes, terrestrial, Africa, SE Asia, Australia, see Cheplick and Faeth (2009; ecology), Wijayawardene et al. (2012, 2017; outline), Pańka et al. (2013; Poland), Maharachchikumbura et al. (2015b, 2016b; outline), DNA sequence data available (as *Balansia sclerotica*), culture data unavailable, neotype of the type species FH (Farlow Herbarium).

Nigrograna Gruyter, Verkley & Crous 2012, *Nigrogranaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *N. mackinnonii* (Borelli) Gruyter, Verkley & Crous, asexual morph coelomycetous, saprobes, Europe, see de Gruyter et al. (2012; taxonomy), Jaklitsch and Voglmayr (2016; taxonomy, phylogeny), Tibpromma et al. (2017b; new species), holotype of the type: WU 36874.

Nigrolentilocus R.F. Castañeda & Heredia 2001, *Ascomycota* genera *incertae sedis*, five species, type: *N. africanus* (B. Sutton) R.F. Castañeda & Heredia, hyphomycetous, sexual morph unknown, saprobes, terrestrial & aquatic, Malawi, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), holotype of type: IMI 347112a, cultures and sequences

are unavailable, compare with *Matsushimaella*, *Minimelanolocus*, *Pseudospirodes*, *Spiropes* and *Sympodioplanus*.

Nigromacula Etayo 2002, *Ascomycota* genera *incertae sedis*, one species, type: *N. hypotrichynae* Etayo, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, South America, Europe, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: COL Etayo 16185.

Nigromammilla K.D. Hyde & J. Fröhl. 2003, *Sordariomycetes* genera *incertae sedis*, one species, type: *N. calami* K.D. Hyde & J. Fröhl. [as 'Nigramammilla calami'], asexual morph unknown, saprobes, terrestrial, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: HKU (M) JF212.

Nigropuncta D. Hawksw. 1981, *Ascomycota* genera *incertae sedis*, two species, type: *N. rugulosa* D. Hawksw., coelomycetous, lichenized, lichenicolous, Europe, terrestrial, sexual morph unknown, see Wijayawardene et al. (2012, 2016b, 2017; morphology, outline), Kirk et al. (2013; genus accepted), cultures and sequences unavailable, holotype of type: K(M)-IMI 241409.

Nigrosabulum Malloch & Cain 1970, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, twelve species, type: *N. globosum* Malloch & Cain, asexual morph acremonium-like, saprobes, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted in *Bionectriaceae*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; accepted in *Hypocreales*), cultures and sequences are available, holotype of type species TRTC 43288.

Nigrospora Zimm. 1902 (= *Basisporium* Molliard 1902; = *Dichotomella* Sacc. 1914; = *Khuskia* H.J. Huds. 1963; = *Phaeoconis* Clem. 1909 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Sordariomycetes*, genera *incertae sedis*, four species, type: *N. panici* Zimm., hyphomycetous, sexual morph formerly named in *Khuskia*, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures and sequences are available.

Nigrovothelium Lücking, M.P. Nelsen & Aptroot 2016, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, two species, type: *N. tropicum* (Ach.) Lücking, M.P. Nelsen & Aptroot, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Aptroot and Lücking (2016; monograph), Lücking et al. (2017a, c; phylogeny, classification), cultures and sequences are available,

lectotype and iso-lectotype of type: H-ACH 707A; BM-ACH.

Nimbospora Jørg. Koch 1982, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *N. effusa* Jørg. Koch, asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; phylogeny), Chu et al. (2015; phylogeny), sequences are available, holotype of type species J. Koch 387, Herbarium CP, Denmark.

Nimisiostella Calat., Barreno & O.E. Erikss. 1997, *Lecanoromycetes* genera *incertae sedis*, one species, type: *N. lichenicola* Calat., Barreno & O.E. Erikss., lichenized, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species VAB (Lich.) 7546.

Nipholepis Syd. 1935, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *N. filicina* Syd., lichenized, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nipicola K.D. Hyde 1992, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *N. carbospora* K.D. Hyde, asexual morph unknown, saprobes, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Daranagama et al. (2017) morphology, taxonomy, phylogeny, key), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Nipponoparmelia (Kurok.) K.H. Moon, Y. Ohmura & Kashiw. 2010 (= *Parmelia* subgen. *Nipponoparmelia* Kurok. 1994 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *N. laevior* (Nyl.) K.H. Moon, Y. Ohmura & Kashiw., asexual morph unknown, lichenized, terrestrial, Argentina, see Crespo et al. (2010b; taxonomy, phylogeny), Kondratyuk et al. (2013d; new species), Divakar et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Niptera Fr. 1849 (= *Nimbomollisia* Nannf. 1983 *fide* Jaklitsch et al. 2016a), *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, c. ten species, type: *N. lacustris* (Fr.) Fr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Whitton et al. (2012b; morphology), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable for type species.

Nipterella Starbäck ex Dennis 1962, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *N. duplex* (Starbäck) Starbäck & Dennis, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch

and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nitidochapsa Parnmen, Lücking & Lumbsch 2013, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *N. leprieurii* (Mont.) Parnmen, Lücking & Lumbsch, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Parnmen et al. (2013; genus proposed), Lücking et al. (2017a; classification), cultures and sequences are available.

Nitschchia G.H. Otth ex P. Karst. 1873 (see Species Fungorum 2017 for synonyms), *Nitschkiaceae*, *Coronophorales*, *Sordariomycetes*, 27 species, type: *N. cupularis* (Pers.) P. Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type, available specimen of type: CBS H-4803.

Nitschkiopsis Nannf. & R. Sant. 1975, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *N. stictarum* Nannf. & R. Sant., asexual morph unknown, saprobes, see Kirk et al. (2008; treated as a synonym of *Niesslia* Auersw.), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a distinct genus), cultures and sequences are unavailable.

Ndobryoria Common & Brodo 1995, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *N. abbreviata* (Müll. Arg.) Common & Brodo, asexual morph unknown, lichenized, terrestrial, North America, Greenland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Divakar et al. (2015; DNA, phylogeny), cultures and sequences are available, voucher specimen of type: St. Clair 52291 (BRY-C).

Nodulosphaeria Rabenh. 1858 (= *Leptosphaeria* subgen. *Pocosphaeria* Sacc. 1883; = *Pocosphaeria* (Sacc.) Berl. 1892 *fide* Species Fungorum 2017), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, c. 52 species, type: *N. hirta* Rabenh., asexual morph coelomycetous, saprobes, terrestrial, aquatic, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; taxonomy), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline, phylogeny), Ariyawansa et al. (2015a; new species, Italy), Li et al. (2015c; new species, phylogeny, asexual morph), Hyde et al. (2016; new species, Italy), Mapook et al. (2016a; new species, revision), Tibpromma et al. (2017b; new species), cultures and sequences are available.

Nohea Kohlm. & Volkmar-Kohlm. 1991, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, three

species, type: *N. umiumi* Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Abdel-Wahab (2011b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; phylogeny), cultures and sequences are available, holotype of type: J.K. 4847 =01348233, NYBG.

Nomuraea Maubl. 1903, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *N. prasina* Maubl., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Noosia Crous, R.G. Shivas & McTaggart 2011, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *N. banksiae* Crous, R.G. Shivas & McTaggart, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Crous et al. (2011a; morphology, DNA), Wijayawardene et al. (2012, 2014c; outline), holotype and ex-type of the type: CBS H-20587; CPC 17282 =CBS 129526.

Normandina Nyl. 1855, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, three species, type: *N. jungermanniae* (Nyl.) Nyl., asexual morph unknown, lichenized, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Ascomycota*), Kirk et al. (2013; genus accepted), Muggia et al. (2010a; phylogeny), Frisch and Ohmura (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Normanogalla Diederich 2014, *Ostropales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *N. cibiformis* (Norman) Diederich, asexual morph is unknown, lichenicolous, terrestrial, Greenland and Scandinavia, see Diederich et al. (2014; taxonomy), lectotype: O, Norman s.n., sequences are unavailable.

Norrlinia Theiss. & Syd. 1918, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *N. peltigericola* (Nyl.) Theiss. & Syd., asexual morph unknown, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for the type species.

Nosophloea Fr. 1849, *Ascomycota* genera *incertae sedis*, three species, type: need typification, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nothojafnea Rifai 1968, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, two species, type: *N. cryptotricha* Rifai, asexual morph unknown, saprobes, terrestrial, Australia,

South America, see Lumbsch and Huhndorf (2010; outline, treated as *Pyronemataceae*), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Nothomitra Maas Geest. 1964, *Geoglossomycetes* genera *incertae sedis*, three species, type: *N. cinnamomea* Maas Geest., asexual morph unknown, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Hustad and Miller (2011; taxonomy), Kirk et al. (2013; genus accepted), Hustad et al. (2013; taxonomy, phylogeny), cultures and sequences are available.

Nothophaeidium J. Reid & Cain 1962, *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *N. abietinellum* (Dearn.) J. Reid & Cain, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: DAOM herb. J. Dearnness 5748.

Nothophoma Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, six species, type: *N. infossa* (Ellis & Everh.) Q. Chen & L. Cai, coelomycetous, saprobes, terrestrial, cosmopolitan, see Chen et al. (2015b; taxonomy, phylogeny), Crous et al. (2016b; new species, phylogeny), Tibpromma et al. (2017b; new data), Wijayawardene et al. (2017; outline), neotype and ex-neotype of the type: CBS H-20145, CBS 123395.

Nothorhytisma Minter, P.F. Cannon, A.I. Romero & Peredo 1998, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, three species, type: *N. nahuelitae* Minter, P.F. Cannon, A.I. Romero & Peredo, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: BAFC.

Nothospora Peyronel 1913, *Ascomycota* genera *incertae sedis*, one species, type: *N. ambigua* Peyronel, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Notocladonia S. Hammer 2003, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *N. cochleata* (Müll. Arg.) S. Hammer, asexual morph unknown, lichenized, terrestrial, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Notolecidea Hertel 1984, *Lecanoromycetes* genera *incertae sedis*, one species, type: *N. subcontinua* (Nyl.) Hertel, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al.

(2017a; classification), cultures and sequences are available.

Notoparmelia A. Crespo, Ferencova & Divakar 2014, *Parmeliaceae, Lecanorales, Lecanoromycetes*, 16 species, type: *N. signifera* (Nyl.) A. Crespo, Ferencova & Divakar, lichenized, Australasia, see Ferencova et al. (2014; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Novomicrothelia Aptroot, M.P. Nelsen & Lücking 2016, *Trypetheliaceae, Trypetheliales, Dothideomycetes*, one species, type: *N. oleosa* (Aptroot) Aptroot, M.P. Nelsen & Lücking, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Aptroot and Lücking (2016; monograph), Lücking et al. (2017a; phylogeny), cultures and sequences are available, holotype of type: F, ISE.

Novozymia W.P. Wu 2005, *Ascomycota* genera *incertae sedis*, one species, type: *N. elegans* W.P. Wu, hyphomycetous, sexual morph unknown, saprobes, terrestrial, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nummospora E. Müll. & Shoemaker 1964, *Ascomycota* genera *incertae sedis*, one species, type: *N. hexamera* E. Müll. & Shoemaker, coelomycetous, sexual morph unknown, saprobes, terrestrial, Switzerland, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nusia Subram. 1995, *Ascomycota* genera *incertae sedis*, two species, type: *N. scheeleae* Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nyctalospora E.F. Morris 1972, *Ascomycota* genera *incertae sedis*, one species, type: *N. compacta* E.F. Morris, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Costa Rica, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nycteromyces Thaxt. 1917 (= *Streblomyces* Thaxt. 1920 fide Species Fungorum 2017), *Laboulbeniaceae, Laboulbeniales, Laboulbeniomycetes*, one species, type: *N. streblidinus* Thaxt., asexual morph unknown, biotrophic, only known from *Strebla wiedemannii*, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nymandomyces Henn. 1899 (= *Synglonium* Penz. & Sacc. 1898 [1897] fide Species Fungorum 2017), *Rhytismatales* genera *incertae sedis*, *Leotiomycetes*, one

species, type: *N. aceris-laurini* Henn., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Nypaelia K.D. Hyde & B. Sutton 1992, *Ascomycota* genera *incertae sedis*, two species, type: *N. frondicola* K.D. Hyde & B. Sutton, coelomycetous, sexual morph unknown, saprobes, marine, on intertidal fronds of *Nypa fruticans*, Brunei, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: IMI 348629.

Nyungwea Sérus., Eb. Fischer & Killmann 2006, *Opegraphaceae, Arthoniales, Arthoniomycetes*, one species, type: *N. pallida* Sérus., Eb. Fischer & Killmann, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Frisch et al. (2014; phylogeny), sequences are available.

Obconicum Velen. 1939, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: need typification, asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Obeliospora Nawawi & Kuthub. 1990, *Ascomycota* genera *incertae sedis*, three species, type: *O. basispira* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Obolarina Pouzar 1986, *Graphostromataceae, Xylariales, Sordariomycetes*, two species, type: *O. dryophila* (Tul. & C. Tul.) Pouzar, asexual morph hyphomycetous (rhinocladiella-like), saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Pažoutová et al. (2010; DNA, phylogeny), Mirabolfathy et al. (2013; new species, Iran), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; transferred to *Graphostromataceae*), cultures available for the type: H76.

Obryzum Wallr. 1825, *Obryzaceae, Ascomycota* families *incertae sedis*, two species, type: *O. corniculatum* (Hoffm.) Wallr., asexual morph unidentified, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline). This name is to be considered a synonym of *Scytinium* in *Collemataceae*, since it was originally based on *Collema corniculatum* Hoffm., and can only be retained in the sense of a lichenicolous fungus by conservation (see Lücking et al. 2017b).

Obstipipilus B. Sutton 1968, *Ascomycota* genera *incertae sedis*, one species, type: *O. malabaricus* (T.S. Ramakr. & K. Ramakr.) B. Sutton, coelomycetous, sexual morph unknown, pathogens, terrestrial, India, see

Wijayawardene et al. (2012, 2016b; outline, morphology), cultures and sequences are unavailable, holotype of the type: IMI 110502.

Obstipispora R.C. Sinclair & Morgan-Jones., 1979, *Ascomycota* genera *incertae sedis*, one species, type: *O. chewaclensis* R.C. Sinclair & Morgan-Jones, sexual morph unknown, submerged decaying leaves, Alabama, see Wijayawardene et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: BPI.

Obtectodiscus E. Müll., Petrini & Samuels 1980, *Loramycetaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *O. aquaticus* E. Müll., Petrini & Samuels, asexual morph unknown, saprobes, terrestrial, Switzerland, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ocala Raja & Shearer 2009, *Natipusillales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *O. scalariformis* Raja & Shearer, submerged wood, Florida, see Raja et al. (2009a, 2015; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Phaeosphaeriaceae*), Shearer et al. (2009; phylogeny), Wijayawardene et al. (2014c; accepted as a genus in *Dothideomycetes*), cultures and sequences are available, holotype of type: F121-1, ILL.

Occultibambusa D.Q. Dai & K.D. Hyde 2016, *Occultibambusaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *O. bambusae* D.Q. Dai & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Thailand, see Dai et al. (2017; phylogeny, morphology, sexual and asexual morphs), Zhang et al. (2017b; new species, phylogeny), cultures and sequences are available, holotype and ex-type cultures of type: MFLU 15-1212; MFLUCC 13-0855 =CBS 139960 =MUCL 55882.

Occultitheca J.D. Rogers & Y.M. Ju 2003, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *O. costaricensis* J.D. Rogers & Y.M. Ju, asexual morph unknown, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable, holotype of type: INB Rogers.

Occultocarpon L.C. Mejía & Zhu L. Yang 2012, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *O. ailaoshanense* L.C. Mejía & Zhu L. Yang, asexual morph unknown, saprobes, terrestrial, China, see Mejía et al. (2012; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and ex-type cultures of the type: BPI 879253, LCM524.01.

Oceanitis Kohlm. 1977 (= *Ascocalsum* J. Campb., J.L. Anderson & Shearer 2003 *fide* Species Fungorum 2017),

Halosphaeriaceae, *Microascales*, *Sordariomycetes*, four species, type: *O. scuticella* Kohlm., asexual morph unknown, saprobes, freshwater or marine, worldwide, see Dupont et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequence data available for all species, holotype of type: J.K. 2971a, not found in NYBG.

Ocellomma Ertz & Tehler 2015, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *O. picconianum* (Bagl.) Ertz & Tehler, sexual morph with apothecioid ascocata, lichenized, terrestrial, Mediterranean, see Ertz et al. (2014b; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available, lectotype and ex-type strain of type: Ertz 17143; 17132.

Ocellularia G. Mey. 1825 (see Species Fungorum 2017 for synonyms), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 400 species, type: *O. obturata* (Ach.) Spreng., asexual morph unknown, lichenized fungi, terrestrial, tropical, see Mangold et al. (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; phylogeny), Sipman et al. (2012; key, Costa Rica), Kirk et al. (2013; genus accepted), Lücking (2014; key), Lücking et al. (2014, 2017a; classification, phylogeny), cultures and sequences are available.

Ochotrichobolus Kimbr. & Korf 1983, *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *O. polysporus* Kimbr. & Korf, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: CUP 59844.

Ochrocladosporium Crous & U. Braun 2007, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, three species, type: *O. elatum* (Harz) Crous & U. Braun, hyphomycetous, terrestrial, Africa, Europe, see Hyde et al. (2011; outline), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Ochroconis de Hoog & Arx 1974, *Sympoventuriaceae*, *Venturiales*, *Dothideomycetes*, 25 species, type: *O. constricta* (E.V. Abbott) de Hoog & Arx, asexual morph hyphomycetous, saprobes, opportunistic pathogens on human and animals, terrestrial, worldwide, see Ge et al. (2012; human pathogen), Giraldo et al. (2014b; new species), Machouart et al. (2014; phylogeny), Samerpitak et al. (2014, 2015; human and animal pathogens, phylogeny), Wijayawardene et al. (2014c, 2017; outline), Crous et al. (2015e, 2016a; new species, phylogeny), cultures and sequences are available, available specimen and cultures for type: CBS H-15203; CBS 211.53, CBS 202.27.

Ochrolechia A. Massal. 1852 (= *Korkir* Adans. 1763 *fide* Species Fungorum 2017), *Ochrolechiaceae*, *Pertusariales*, *Lecanoromycetes*, 60 species, type:

O. tartarea (L.) A. Massal., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Aptroot and Aslan (2012; new records), Kirk et al. (2013; genus accepted), Ertz et al. (2016a; taxonomy), sequence are available for the type.

Ochroconectria Rossman & Samuels 1999, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *O. calami* (Henn. & E. Nyman) Rossman & Samuels, asexual morph unknown, saprobes, terrestrial, Asia, South America, see Guu et al. (2010; Taiwan), Pinruan et al. (2010; Thailand), Li et al. (2016a; new species), cultures and sequences are available, holotype and culture data unavailable.

Ocostaspora E.B.G. Jones, R.G. Johnson & S.T. Moss 1983, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *O. apilongissima* E.B.G. Jones, R.G. Johnson & S.T. Moss, asexual morph unknown, saprobes, marine, Wales, Denmark, temperate, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; DNA, phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ocotomyces H.C. Evans & Minter 1985 (= *Uyucamyces* H.C. Evans & Minter *fide* Johnston et al. 2014b), *Leotiomycetes* genera *incertae sedis*, one species, type: *O. parasiticus* H.C. Evans & Minter, asexual morph formerly named in *Uyucamyces*, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, holotype of type: IMI 286340.

Octopodotus Kohlm. & Volk.-Kohlm. 2003, *Ascomycota* genera *incertae sedis*, one species, type: *O. stupendous* Kohlm. & Volk.-Kohlm., coelomycetous, sexual morph unknown, saprobes, marine, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Octospora Hedw. 1789 (= *Humaria* (Fr.) Boud. 1885; = *Humarina* Seaver 1927; = *Leucoloma* Fuckel 1870 [1869–70]; = *Peziza* trib. *Humaria* Fr. 1822; = *Pseudocenangium* A. Knapp 1924 *fide* Species Fungorum 2017), *Pezizales* genera *incertae sedis*, *Pezizales*, c. 60 species, type: *O. leucoloma* Hedw., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Octosporella Döbbeler 1980, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, nine species, type: *O. jungermanniarum* (P. Crouan & H. Crouan) Döbbeler, asexual morph unknown, on mosses and liverworts,

terrestrial, Europe, Venezuela, see Lumbsch and Huhndorf (2010; outline), Stenroos et al. (2010; DNA), Döbbeler (2011; new species, Costa Rica and French Guiana), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Octosporopsis U. Lindem. & M. Vega 2014, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, nine species, type: *O. nicolai* U. Lindemann, M. Vega & T. Richter, asexual morph unknown, saprobes, terrestrial, Europe, see Lindemann et al. (2014; taxonomy, phylogeny), sequences are available, voucher specimen for type: MV20130205-1.

Oculimacula Crous & W. Gams 2003 (= *Helgardia* Crous & W. Gams 2003 *fide* Johnston et al. 2014b), *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *O. yellundae* (Wallwork & Spooner) Crous & W. Gams, asexual morph formerly named in *Helgardia*, saprobes, pathogens, terrestrial, Germany, Australia, France, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, holotype of the type species unavailable.

Odontodictyospora Mercado 1984, *Ascomycota* genera *incertae sedis*, one species, type: *O. comasii* Mercado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Odontotrema Nyl. 1858, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, c. seven species, type: *O. phacidoides* Nyl., asexual morph unknown, lichenized, temperate, see Baloch et al. (2010; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Zhurbenko (2010a; new species), sequences are available.

Oedocephalum Preuss 1851 (= *Amblyosporiopsis* Fairm. 1922; = *Sporocephalum* G. Arnaud 1952 *fide* Species Fungorum 2017), *Pezizaceae*, *Pezizales*, *Pezizomycetes*, c. 23 species, type: *O. elegans* Preuss, hyphomycetous, sexual morph unknown, Nematode trapping, in soil, terrestrial, aquatic, Asia, Europe, see Davison et al. (2009; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Oedohysterium E.W.A. Boehm & C.L. Schoch 2009, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, three species, type: *O. insidens* (Schwein.) E. Boehm & C.L. Schoch, saprobes, terrestrial, North and South America, Europe, South Africa, China, see Boehm et al. (2009a, b; generic review), Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Oedothaea Syd. 1930, *Ascomycota genera incertae sedis*, one species, type: *O. vismiae* Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Oevstedalia Ertz & Diederich 2004, *Ascomycota genera incertae sedis*, one species, type: *O. antarctica* (C.W. Dodge) Ertz & Diederich, sexual morph perithecioid, asexual morph represented by true ascocnidia, lichenized, on wood, rock, detritus and whale bones, Antarctica, probably endemic, first lichenized ascomycete known to produce true ascocnidia, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of the type: Follmann 14173 (FH-Dodge).

Ogataea Y. Yamada, K. Maeda & Mikata 1994, *Pichiaceae*, *Saccharomycetales*, *Saccharomyces*, 40 species, type: *O. minuta* (Wickerham) Y. Yamada, K. Maeda & Mikata, asexual reproduction is by multilateral budding, saprophytic, soil, flowers, tree leaves, tree sap, rotted wood, insect frass in trees and other plants, animal faeces, fruit of osage orange, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Ohleria Fuckel 1868, *Ohleriaceae*, *Pleosporales*, *Dothideomycetes*, 13 species, type: *O. modesta* Fuckel, asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Tian et al. (2015; morphology, nomenclature, phylogeny), Jaklitsch and Voglmayr (2016; taxonomy, phylogeny), cultures and sequences are unavailable, holotype of type species Rhenani exsicc. 2173 present in G.

Ohleriella Earle 1902, *Delitschiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *O. neomexicana* Earle, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Delitschiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Oidiiodendron Robak 1932, *Myxotrichaceae*, *Leotiomycetes* families *incertae sedis*, two species, type: *O. fuscum* Robak, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for type: CBS 238.31, MUCL 1057.

Ojibwaya B. Sutton 1973, *Ascomycota genera incertae sedis*, one species, type: *O. perpulchra* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Belarus, Canada, Malawi, see Seifert et al.

(2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: IMI 132708.

Okeanomyces K.L. Pang & E.B.G. Jones 2004, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *O. cucullatus* (Kohlm.) K.L. Pang & E.B.G. Jones, asexual morph hyphomycetous, saprobes, marine, worldwide, Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), sequence data available, holotype of type: J.K. 1686 =00966750, NYBG.

Olegblumia S.Y. Kondr., L. Lököö & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *O. demissa* Flot. ex S.Y. Kondr., L. Lököö, J. Kim, A.S. Kondr., S.O. Oh & Hur, lichenized, Europe and North America, see Kondratyuk et al. (2015d; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Oleoguttula Selbmann & de Hoog 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *O. mirabilis* Selbmann & de Hoog, hyphomycetous, sexual morph unknown, rock-inhabiting fungus, Antarctica, see Egidi et al. (2014; taxonomy), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2014c, 2017; outline), type: CBS 136102 =CCFEE 5523, cultures and sequences available.

Oletheriostrigula Huhndorf & R.C. Harris 1996, *Strigulaceae*, *Strigulales*, *Dothideomycetes*, one species, type: *O. papulosa* (Durieu & Mont.) Huhndorf & R.C. Harris, asexual morph unknown, lichenized, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), Lücking et al. (2017a; classification), cultures and sequence are unavailable.

Olla Velen. 1934 (= *Pseudolla* Velen. 1934 *fide* Species Fungorum 2017), *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, twelve species, type: *O. ulmariae* Velen., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Olpitrichum G.F. Atk. 1894, *Ascomycota genera incertae sedis*, three species, type: *O. carpophilum* G.F. Atk., hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Ombrophila Fr. 1849 (= *Haplocybe* Clem. 1909; = *Kubickia* Svrček 1957 *fide* Species Fungorum 2017), *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, eleven species, type: *O. violacea* (Hedw.) Fr., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus

accepted), Osmundson et al. (2013; DNA), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Omega B. Sutton & Minter 1988, *Ascomycota* genera *incertae sedis*, one species, type: *O. coenobiticum* B. Sutton & Minter, coelomycetous, sexual morph unknown, saprobes, terrestrial, Greece, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type species IMI 326014.

Ommatomyces Kohlm., Volkm.-Kohlm. & O.E. Erikss. 1995, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, three species, type: *O. coronatus* Kohlm., Volkm.-Kohlm. & O.E. Erikss., asexual morph unknown, saprobes, marine or terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMS JK5509.

Omnidemputus P.F. Cannon & Alcorn 1994, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *O. affinis* P.F. Cannon & Alcorn, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of the type: BRIP 17195.

Omphalodium Meyen & Flot. 1843, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, four species, type: *O. pisacomense* Meyen & Flot., asexual morph unknown, lichenized, terrestrial, North and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Divakar et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Omphalora T.H. Nash & Hafellner 1990, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *O. arizonica* (Tuck. ex Willey) T.H. Nash & Hafellner, asexual morph unknown, lichenized, terrestrial, North America, see Crespo et al. (2010b; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Omphalospora Theiss. & Syd. 1915 (= *Placosphaerella* Pat. 1897; = *Plectosphaerella* Kirschst. 1938; = *Plectosphaerina* Kirschst. 1938 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, two species, type: *O. stellariae* (Lib.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Oncocladium Wallr. 1833 (= *Actinodendron* G.F. Orr & Kuehn 1963 *fide* Species Fungorum 2017), *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, one

species, type: *O. flavum* Wallr., hyphomycetous, sexual morph unknown, soil borne, Europe, Canada, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and cultures are unavailable.

Oncopodiella G. Arnaud ex Rifai 1965, *Dothideomycetes* genera *incertae sedis*, 14 species, type: *O. trigonella* (Sacc.) Rifai, hyphomycetous, soil borne, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), cultures and sequences are unavailable, cultures available for the type: CBS 126413, MUCL 52643.

Oncopodium Sacc. 1904, *Ascomycota* genera *incertae sedis*, twelve species, type: *O. antoniae* Sacc. & D. Sacc., hyphomycetous, soil borne, Europe and North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Oncospora Kalchbr. 1880, *Ascomycota* genera *incertae sedis*, eight species, type: *O. bullata* Kalchbr. & Cooke, coelomycetous, widespread, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Oncosporella P. Karst. 1887, *Ascomycota* genera *incertae sedis*, one species, type: *O. punctiformis* P. Karst. 1887, coelomycetous, Finland, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Oncostroma Bat. & Marasas 1966, *Ascomycota* genera *incertae sedis*, one species, type: *O. toddiae* Bat. & Marasas, hyphomycetous, South Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ondiniella E.B.G. Jones, R.G. Johnson & S.T. Moss 1984, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *O. torquata* (Kohlm.) E.B.G. Jones, R.G. Johnson & S.T. Moss, asexual morph unknown, saprobes on wood, marine, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Onychocola Sigler 1990, *Arachnomycetaceae*, *Arachnomycetales*, *Eurotiomycetes*, five species, type: *O. canadensis* Sigler, human pathogens, widespread, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, UTMB 3250.

Onychophora W. Gams, P.J. Fisher & J. Webster 1984, *Ascomycota* genera *incertae sedis*, one species, type:

O. coprophila W. Gams, P.J. Fisher & J. Webster, hyphomycetous, sexual morph unknown, coprophilous (rabbit dung), terrestrial, UK, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: CBS 1740.

Onygena Pers. 1800 (= *Piligena* Schumach. 1803; = *Sphaeropus* Paulet 1808 *fide* Species Fungorum 2017), *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *O. equina* (Willd.) Pers., asexual morph unknown, on feathers, bones, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Huang et al. (2015b; DNA, protein degrading taxa), cultures and sequences are available, cultures available for the type species (Huang et al. 2013), type specimen of type species CBS H-15271.

Onygenopsis Henn. 1910, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *O. engleriana* Henn., asexual morph unknown, saprobes, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Oomyces Berk. & Broome 1851 (= *Coscinaria* Ellis & Everh. 1886 *fide* Species Fungorum 2017), *Acrospermaceae*, *Acrospermales*, *Dothideomycetes*, ten species, type: *O. carneoalbus* (Lib.) Berk. & Broome, asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable, need generic revision.

Osporidium Stautz 1931, *Saccharomycetes* genera *incertae sedis*, two species, type: *O. margaritiferum* Stautz, saprobes, terrestrial, Europe and North America, Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for type species CBS 2531, ATCC 10676, CCRC 21536, DSM 70351, IFO 1208, NRRL Y-1519, VKM Y-816.

Oothyrium Syd. 1939, *Ascomycota* genera *incertae sedis*, one species, type: *O. butyrospermi* Syd., coelomycetous, sexual morph unknown, Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Opegrapha Ach. 1809 (see Species Fungorum 2017 for synonyms), *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, but strongly polyphyletic, 300 species, type: *O. vulgata* (Ach.) Ach., asexual morph unknown, lichenized, saprobes, lichenicolous, terrestrial, worldwide, see Ertz et al. (2009; DNA), Lumbsch and Huhndorf (2010; outline), Ertz and Tehler (2011; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Ophidiomyces Sigler, Hambl. & Paré 2013, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species,

type: *O. ophioditicola* (Guarro, Deanna A. Sutton, Wickes & Rajeev) Sigler, Hambleton & Paré, asexual morph unknown, pathogens on snakes, Northern America, see Sigler et al. (2013; taxonomy, phylogeny), Allender et al. (2015; pathogens), cultures and sequences are available, cultures available for the type species UAMH 10717 (Sigler et al. 2013).

Ophiobolus Riess 1854 (= *Leptosphaeriopsis* Berl. 1892 *fide* Species Fungorum 2017), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *O. disseminans* Riess, asexual morph coniothyrium-like; phoma-like, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; taxonomy), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Ophioceras Sacc. 1883, *Ophioceraceae*, *Magnaportheales*, *Sordariomycetes*, 39 species, type: *O. dolichostomum* (Berk. & M.A. Curtis) Sacc., saprobes, aquatic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hu et al. (2012b; new species), Kirk et al. (2013; genus accepted), Klaubauf et al. (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ophiociliomyces Bat. & I.H. Lima 1955, *Ascomycota* genera *incertae sedis*, one species, type: *O. bauhiniae* Bat. & I.H. Lima, asexual morph unknown, from living leaves, tropical, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded the genus from *Meliolaceae*), cultures and sequences are unavailable.

Ophiocordyceps Petch 1931 (= *Cordycepioideus* Stifler 1941 *fide* Index Fungorum 2017), *Ophiocordycipitaceae*, *Hypocreales*, *Sordariomycetes*, c. 200 species, type: *O. blattae* (Petch) Petch, asexual morph acremonium-like, on insects, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Evans et al. (2011; new species), Kirk et al. (2013; genus accepted), Wen et al. (2013, 2014, 2016a, b; new species, phylogeny), Araújo et al. (2015; new species), Ban et al. (2015; new species), Kobmoo et al. (2015; new species), Sanjuan et al. (2015; new species), Spatafora et al. (2015; new species, nomenclature), Maharachchikumbura et al. (2015b, 2016b; outline), Crous et al. (2016a; new species), cultures and sequences are available.

Ophioleira Kohlm. & Volk.-Kohlm. 1988, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *O. monosemeia* Kohlm. & Volk.-Kohlm., marine, Caribbean, see Lumbsch and Huhndorf (2010; outline), Jones et al. (2015; outline, marine),

Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences available.

Ophiodiaporthe Y.M. Ju, H.M. Hsieh, C.H. Fu, C.Y. Chen & T.T. Chang 2013, Diaporthaceae, Diaporthales, Sordariomycetes, one species, type: *O. cyatheae* Y.M. Ju, H.-M. Hsieh, C.H. Fu, C.Y. Chen & T.T. Chang, asexual morph coelomycetous, pathogens, terrestrial, Taiwan, see Fu et al. (2013; phylogeny, taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype and ex-type strain of type: YMJ1364; CRC 34961.

Ophiodothella (Henn.) Höhn. 1910 (= *Microphiodothis* Speg. 1919 [1918]; = *Ophiodothis* subgen. *Ophiodothella* Henn. 1904; = *Rhopographina* Theiss. & Syd. 1915; = *Scolecodothis* Theiss. & Syd. 1914 *fide* Species Fungorum 2017), Phyllachoraceae, Phyllachorales, Sordariomycetes, 31 species, type: *O. atromaculans* (Henn.) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hanlin and González (2013; new species), Fu et al. (2013; DNA), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ophiognomonia (Sacc.) Speg. 1896 (= *Gnomonia* subgen. *Ophiognomonia* Sacc. 1882; = *Ophiognomonia* (Sacc.) Sacc. 1899 *fide* Species Fungorum 2017), Gnomoniaceae, Diaporthales, Sordariomycetes, 46 species, type: *O. melanostyla* (DC.) Sacc., asexual morph unknown, endophytes, pathogens, saprobes, primarily temperate and rarely subtropics, worldwide, see Lumbsch and Huhndorf (2010; outline), Walker et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Ophioirenina Sawada & W. Yamam. 1959, Dothideomycetes genera incertae sedis, one species, type: *O. theae* Sawada & W. Yamam., asexual morph unknown, from leaves, terrestrial, India, Taiwan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Meliolaceae), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; outline, excluded the genus from Meliolaceae and placed in Dothideomycetes), cultures and sequences are unavailable.

Ophiomassaria Jacz. 1894, Sordariomycetes genera incertae sedis, one species, type: *O. selenospora* (G.H. Otth) Jacz., asexual morph unknown, saprobes or endophytes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ophiomeliola Starbäck 1899, Sordariomycetes genera incertae sedis, four species, type: *O. lindmanii* Starbäck,

aseexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in Perisporiopsidaceae), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Hongsanan et al. (2015d; accepted as a genus in Sordariomycetes), cultures and sequences are unavailable.

Ophionectria Sacc. 1878 (= *Antipodium* Piroz. 1974 *fide* Rossman et al. 2016a), Nectriaceae, Hypocreales, Sordariomycetes, two species, type: *O. trichospora* (Berk. & Broome) Sacc., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are available, type specimen and cultures available for the type: CBS H-6331; CBS 109876.

Ophioparma Norman 1852 (= *Haematommato myces* Cif. & Tomas. 1953; = *Ophioparma* Norman 1853 *fide* Species Fungorum 2017), Ophioparmaceae, Umbilicariales, Lecanoromycetes, two species, type: *O. ventosa* (L.) Norman, asexual morph unknown, saprobes, terrestrial, tropical, see Khare et al. (2010; taxonomy, India), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Le Pogam et al. (2016; taxonomy), Lücking et al. (2017a; classification), cultures available for the type: AFTOL-ID 1694.

Ophioparodia Petr. & Cif. 1932, Perisporiopsidaceae, Dothideomycetes families incertae sedis, one species, type: *O. pulchra* Petr. & Cif., asexual morph unknown, saprobes, terrestrial, West Indies, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in Perisporiopsidaceae), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Ophiorosellinia J.D. Rogers, A. Hidalgo, F.A. Fernández & Huhndorf 2004, Xylariales genera incertae sedis, Sordariomycetes, one species, type: *O. costaricensis* J.D. Rogers, A. Hidalgo, F.A. Fernández & Huhndorf, asexual morph unknown, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in Xylariales genera incertae sedis), cultures and sequences are unavailable, holotype of the type: USJ72848.

Ophiosira Petr. 1955, Ascomycota genera incertae sedis, one species, type: *O. austriaca* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ophisphaerella Speg. 1909, Phaeosphaeriaceae, Pleosporales, Dothideomycetes, ten species, type: *O. graminicola* Speg., asexual morph wojnowicia-like,

saprobes, terrestrial, aquatic, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; taxonomy), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline, phylogeny), Ariyawansa et al. (2015a; new species, phylogeny), cultures and sequences are available.

Ophiostoma Syd. & P. Syd. 1919 (= *Europhium* A.K. Parker 1957; = *Linostoma* Höhn. 1918 *fide* Species Fungorum 2017), *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, c. 230 species, type: *O. piliferum* (Fr.) Syd. & P. Syd., saprobes, worldwide, see Linnakoski et al. (2008, 2010; new species), Kamgan et al. (2010, 2012; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Romón et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Oplothecium Syd. & P. Syd. 1923, *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, two species, type: *O. arecae* Syd. & P. Syd., asexual morph unknown, saprobes, Austria, see Kirk et al. (2008; treated as a synonym of *Neorehmia*), Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Orbicula Cooke 1871 (= *Anixia* H. Hoffm. 1862; = *Mycogala* Rostaf. ex Sacc. 1884 *fide* Species Fungorum 2017), *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *O. cyclospora* Cooke, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Orbilia Fr. 1836 (= *Dicranidion* Harkn. 1885; = *Pedilospora* Höhn. 1902 *fide* Baral et al. 2015; see Species Fungorum 2017 for additional synonyms), *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, c. 75 species, type: *O. leucostigma* (Fr.) Fr., asexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Zhang et al. (2009f; new species, China), Lumbsch and Huhndorf (2010; outline), Su et al. (2011b; new species, China), Quijada et al. (2013; new species, Spain), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Orbiliopsis (Sacc. & D. Sacc.) Syd. & P. Syd. 1924 (= *Orbilia* subgen. *Orbiliopsis* Sacc. & D. Sacc. 1906 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *O. coleosporioides* (Sacc.) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available but lacking for the type species.

Orbimyces Linder 1944, *Lulworthiaceae*, *Lulwothiales*, *Sordariomycetes*, one species, type: *O. spectabilis* Linder, sexual morph unknown, hyphomycetous, marine, see Jones et al. (2008; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; phylogeny, outline), cultures and sequences are available, holotype: CBS H-15318.

Orbiocrella D. Johnson, G.H. Sung, Hywel-Jones & Spatafora 2009, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *O. petchii* (Hywel-Jones) D. Johnson, G.H. Sung, Hywel-Jones & Spatafora, asexual morphs unknown, on insects, terrestrial, Thailand, see Johnson et al. (2009; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, nomenclature), cultures and sequences are available.

Orcadia G.K. Sutherl. 1914, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *O. ascophylli* G.K. Sutherl., asexual morph unknown, marine, on brown algae, UK, Norway, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Pezizales*), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), cultures and sequences are unavailable.

Orceolina Hertel 1970 (= *Urceolina* Tuck. 1875 *fide* Species Fungorum 2017), *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, two species, type: *O. kerguelensis* (Tuck.) Hertel, asexual morph unknown, lichenized, terrestrial, subantarctica islands, see Lumbsch and Huhndorf (2010; outline), Lumbsch et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Kantvilas et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Orcularia (Malme) Kalb & Giralt 2011, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, four species, type: *O. insperata* (Nyl.) Kalb & Giralt, asexual morph present (in the original description, authors mentioned that genus has filiform conidia), lichenized, terrestrial, worldwide, see Kalb and Giralt (2011; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Orientophila Arup, Søchting & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, four species, type: *O. loekoesii* (S.Y. Kondr. & J.S. Hur) Arup, Søchting & Frödén, asexual morph unknown, lichenized, terrestrial, worldwide, see Arup et al. (2013; taxonomy, phylogeny), Kondratyuk et al. (2014, 2016a, b; new species, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Ormomyces I.I. Tav. 1985, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *O. clivinae* (Thaxt.) I.I. Tav., asexual morph unknown, biotrophic, terrestrial, Africa, Indonesia, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Oropogon Th. Fr. 1861 (= *Atestia* Trevis. 1861 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 30 species, type: *O. loxensis* (Fée) Th. Fr., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Leavitt et al. (2012; diversity in Neotropical), Kirk et al. (2013; genus accepted), Divakar et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, voucher specimen for the type: Nelson 4197A (F).

Orphanocoela Nag Raj 1989, *Ascomycota* genera *incertae sedis*, three species, type: *O. calamagrostidis* (H.C. Greene) Nag Raj, coelomycetous, sexual morph unknown, pathogens, terrestrial, USA, Africa, Colombia, Mexico, Nicaragua, Mexico, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; morphology), cultures and sequences are unavailable, holotype of the type species unavailable.

Orphniospora Körb. 1874 (= *Orphniosporomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Elixiaceae*, *Umbilicariales*, *Lecanoromycetes*, three species, type: *O. groenlandica* Körb., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Biložit and Mayrhofer (2014; new records in Vorarlberg, Australia), cultures and sequences are unavailable.

Orphnodactylis Malloch & A. Mallik 1998, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, 31 species, type: *O. kalmiae* Malloch & A. Mallik, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Orromyces B. Sur & G.R. Ghosh 1987, *Gymnoascaceae*, *Onygenales*, *Eurotiomycetes*, four species, type: *O. spiralis* B. Sur & G.R. Ghosh, asexual morph chrysosporium-like, saprobes, from soil, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Orthobellus A.A. Silva & Cavalc. 1973, *Dothideomycetes* genera *incertae sedis*, three species, type: *O. leguminosarum* A.A. Silva & Cavalc., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Schizothyriaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Schizothyriaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Schizothyriaceae*), Phookamsak et al. (2016; taxonomy, transferred to *Dothideomycetes*

genera *incertae sedis*), cultures and sequences are unavailable.

Oscarbrefeldia Holterm. 1898, *Saccharomycetes* genera *incertae sedis*, one species, type: *O. pellucida* Holterm., information unavailable, see Kirk et al. (2008, 2013; genus accepted in *Saccharomycetes*) cultures and sequences are unavailable.

Osoriomyces Terada 1981, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *O. rhizophorus* Terada, asexual morph unknown, biotrophic, terrestrial, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ostracoderma Fr. 1825, *Ascomycota* genera *incertae sedis*, three species, type: *O. pulvinatum* Fr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Ostracodermidium Mukerji 1973, *Ascomycota* genera *incertae sedis*, one species, type: *O. serpentinum* (J.N. Rai & J.P. Tewari) Mukerji, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ostrechnion Duby 1861 (= *Ostreion* Sacc. 1883 *fide* Species Fungorum 2017), *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, three species, type: *O. americanum* Duby, asexual morph unknown, saprobes, terrestrial, Atlantic, China, see Boehm et al. (2009a, b; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Ostreola Darker 1963, *Mytilinidiaceae*, *Mytilinidiales*, *Dothideomycetes*, twelve species, type: *O. consociata* Darker, asexual morph septonema-like, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mytilinidiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Ostropa Fr. 1825, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *O. barbara* (Fr.) Nannf., asexual morph unknown, lichenized, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), available specimen and cultures for the type: CBS H-15322; AFTOL-ID 77.

Ostropella (Sacc.) Höhn. 1918, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, five species, type: *O. albocincta* (Berk. & M.A. Curtis) Höhn., asexual morph unknown, saprobes, worldwide, see Lumbsch and

Huhndorf (2010; outline, treated as *Pleosporales*, genera *incertae sedis*), Hyde et al. (2013; accepted as a genus in *Platystomaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Oswaldina Rangel 1921, *Ascomycota* genera *incertae sedis*, one species, type: *O. icarthyensis* Rangel, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Phyllachoraceae*), Wijayawardene et al. (2017; not accepted in *Phyllachoraceae*), cultures and sequences are unavailable.

Otidea (Pers.) Bonord. 1851 (= *Cochlearia* (Cooke) Lambotte 1887 [1888]; = *Flavoscypha* Harmaja 1974; = *Otidea* subgen. *Pseudotis* Boud. 1885; = *Peziza* †† *Otidea* Pers. 1822; = *Peziza* sect. *Cochleate* Fr. 1822; = *Peziza* subgen. *Cochlearia* Cooke 1879; = *Pseudotis* (Boud.) Boud. 1907 *fide* Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 40 species, type: *O. onotica* (Pers.) Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Carbone and Van Vooren (2011; new species), Kirk et al. (2013; genus accepted), Olariaga et al. (2015; taxonomy, monograph), cultures and sequences are available, epitype and cultures are available for the type species KH.10.284 (S - epitype); KH.10.284.

Otthia Nitschke ex Fuckel 1870 (= *Otthia* subgen. *Otthiella* Sacc. 1882; = *Otthiella* (Sacc.) Sacc. & D. Sacc. 1905 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, eleven species, type: *O. spiraeae* (Fuckel) Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available, cultures available for the type: HKUCC 10896 (unpublished).

Otwaya G.W. Beaton 1978, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, twelve species, type: *O. verruculospora* G.W. Beaton, asexual morph unknown, saprobes, or endophytes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as a synonym of *Aeruginoscyphus* Dougoud), cultures and sequences are unavailable.

Ovadendron Sigler & J.W. Carmich. 1976, *Ascomycota* genera *incertae sedis*, one species, type: *O. sulphureo-ochraceum* (J.F.H. Beyma) Sigler & J.W. Carmich., hyphomycetous, sexual morph unknown, human pathogen, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Oviacomia Y.J. Yao & Spooner 1996, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *O. paludosum* (Dennis) Y.J. Yao & Spooner, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Oviculispora Etayo 2010, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *O. parmeliae* Etayo, asexual morph unknown, lichenicolous, terrestrial, tropical, see Etayo (2010a; morphology), cultures and sequences are available.

Ovulinia F.A. Weiss 1940 (= *Ovulitis* N.F. Buchw 1970 *fide* Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, nine species, type: *O. azaleae* F.A. Weiss, asexual morph formerly named in *Ovulitis*, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are unavailable.

Oxnerella S.Y. Kondr., Lókös & Hur 2014, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *O. safavidiorum* S.Y. Kondr., Zarei-Darki, Lókös & Hur, lichenized, see Resl et al. (2016; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type: KW-L 70296.

Oxodeora K.D. Hyde & P.F. Cannon 1999, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *O. petrakii* (Cif.) K.D. Hyde & P.F. Cannon, asexual morph unknown, on living palm leaves, terrestrial, Caribbean, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Oxydothis Penz. & Sacc. 1898 (= *Merrilliopeltis* Henn. 1908; = *Plagiogynion* Schrantz 1962; = *Plagiothecium* Schrantz 1961 *fide* Species Fungorum 2017), *Oxydothidaceae*, *Amphisphaerales*, *Sordariomycetes*, c. 35 species, type: *O. grisea* Penz. & Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Liu et al. (2015a; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Konta et al. (2016b; phylogeny, introduced *Oxydothidaceae*), cultures and sequences are available but lacking for the type species.

Paathramaya Subram. 1956, *Ascomycota* genera *incertae sedis*, five species, type: *P. sundara* Subram., hyphomycetous, sexual morph unknown, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype: CBS H-194, need generic revision.

Pachnolepia A. Massal. 1855, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. pruinata* (Pers.) Frisch & G. Thor, lichenized, see Frisch et al. (2014; reinstated), Lücking et al. (2017a; classification), sequences are available, need generic revision.

Pachyascus Poelt & Hertel 1968, *Pachyascaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *P. lapponicus* Poelt & Hertel, asexual morph unknown, lichenized, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Pachycladina Marvanová 1987, *Ascomycota* genera *incertae sedis*, three species, type: *P. mutabilis* Marvanová, hyphomycetous, sexual morph unknown, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Pachydisca Boud. 1885, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. guernisacii* (P. Crouan & H. Crouan) Boud., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as a synonym of *Discinella* Boud.), cultures and sequences are available but lacking for the type species.

Pachyella Boud. 1907 (= *Gorodkoviella* Vassilkov 1969; = *Peltidium* Kalchbr. 1862; = *Pulvinaria* Velen. 1934 *fide* Species Fungorum 2017), *Pezizaceae*, *Pezizales*, *Pezizomycetes*, 19 species, type: *P. barlaeana* (Bres.) Boud., asexual morph unknown, saprobes, terrestrial, worldwide, see Healy et al. (2009b; species in USA and Mexico), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Pachypatella Theiss. & Syd. 1915 (= *Apoa* Syd. 1931 *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. alsophilae* (Racib.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, Brazil, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype and isotype of type species KRAM?: S F9863.

Pachypeltis Søchting, Arup & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, four species, type: *P. castellana* (Räsänen) Søchting, Frödén & Arup, asexual morph unknown, lichenized, terrestrial, Europe, see Arup et al. (2013; taxonomy), Lücking et al.

(2017a; classification), cultures and sequences are available.

Pachyphloides Zobel 1854 (= *Cryptica* R. Hesse 1884; = *Pachyphloeus* Tul. & C. Tul. 1845 *fide* Jaklitsch et al. 2016a; Species Fungorum 2017), *Pezizaceae*, *Pezizales*, *Pezizomycetes*, 19 species, type: *P. ligericus* Zobel, asexual morph unknown, saprobes, saprobes, terrestrial, worldwide, see Doweld (2013; new combinations), Healy et al. (2015; new species, DNA), cultures and sequences are available.

Pachyphytis R.C. Harris & Ladd 2007, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *P. ozarkana* R.C. Harris & Ladd, asexual morph unknown, on dolomite, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available, holotype and cultures are available for the type species Pacoza51; NY, Harris 44707.

Pachysacca Syd. 1930, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. eucalypti* Syd., asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline, accepted as genus in *Dothideaceae*), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; excluded from *Dothideaceae* and placed in *Teratosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pachysolen Boidin & Adzet 1957, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *P. tannophilus* Boidin & Adzet, asexual reproduction is by multilateral budding, saprophytic, extracts of chestnut (*Castanea vesca*) and mimosa (*Acacia mollissima*) for use in leather tanning, France, Germany, Spain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Pachythrium G. Arnaud ex Spooner & P.M. Kirk 1990, *Trichothyriaceae*, *Dothideomycetes*, families *incertae sedis*, one species, type: *P. parasiticum* (Fabre) G. Arnaud ex Spooner & P.M. Kirk, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; accepted as a genus in *Trichothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pachytrype Berl. ex M.E. Barr, J.D. Rogers & Y.M. Ju 1993, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type species: *P. princeps* (Penz. & Sacc.) M.E. Barr, J.D. Rogers & Y.M. Ju, asexual morph cytospora-like, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Paecilomyces Bainier 1907 (= *Byssochlamys* Westling 1909 = *Corollium* Sopp 1912; = *Graphidium* Lindau 1909; = *Spicariopsis* R. Heim 1939 *fide* Rossman et al. 2016a; Species Fungorum 2017), *Thermoascaceae*, *Eurotiales*, *Eurotiomycetes*, c. 50 species, type: *P. variotii* Bainier, hyphomycetous, sexual morph formerly known in *Byssochlamys*, widespread, see Liang et al. (2009b; taxonomy), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016a; nomenclature), cultures available for type species CBS 102.74, CBS 110431.

Palawania Syd. & P. Syd. 1914, *Palawaniaceae*, *Dothideomycetes* families, *incertae sedis*, two species, type: *P. grandis* (Niessl) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Mapook et al. (2016d; new species, phylogeny, place in *Palawaniaceae*), cultures and sequences are available.

Palawaniella Doidge 1921 (= *Byliana* Dippen. 1930 *fide* Species Fungorum 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, eight species, type: *P. eucleae* Doidge, asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: PREM 12375.

Palicella Rodr. Flakus & Printzen 2014, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *P. glaucopa* (Hook. f. & Taylor) Rodr. Flakus & Printzen, lichenized, Argentina, Canada, Europe, see Flakus and Printzen (2014; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Paliphora Sivan. & B. Sutton 1985, *Ascomycota* genera *incertae sedis*, six species, type: *P. aurea* Sivan. & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Gusmão et al. (2008; morphology, new species, distribution), Goh et al. (2014b; morphology, new species), Wijayawardene et al. (2017; outline), sequences are available, holotype: IMI 263424g.

Pallidocercospora Crous 2012, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, eight species, type: *P. heimii* (Crous) Crous, hyphomycetous, sexual morph reported, pathogenic, terrestrial, worldwide, see Crous et al. (2012a; morphology; phylogeny), cultures and sequences are available, holotype: PREM 51749 (sexual

morph), holotype of the type: PREM 51748 (aseexual morph), ex-type strain of type: CPC 760.

Pallidogramme Staiger, Kalb & Lücking 2008 (= *Hemithecium* subgen. *Leucogramma* Staiger 2002 *fide* Species Fungorum 2017), *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, twelve species, type: *P. chrysenteron* (Mont.) Staiger, Kalb & Lücking, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Archer (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2014, 2017b; phylogeny, classification), cultures and sequences are available, holotype of type species PC (Leprieur 23).

Palmiascoma Phook. & K.D. Hyde 2015, *Bambusicolaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. gregariascomum* Phook. & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Thailand, see Liu et al. (2015a; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 11–0211, MFLUCC 11–0175.

Palmicola K.D. Hyde 1993, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *P. archontophoenicis* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Australia, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type species, holotype of the type: BRIP 20438.

Palmaria K.D. Hyde, J. Fröhl. & Joanne E. Taylor, **nom. nov.**

= *Palmomyces* K.D. Hyde, J. Fröhl. & Joanne E. Taylor 1998, *Sydiowia* 50(1): 59 (1998) (nom. illegit., Art. 53.1); non *Palmomyces* Maire 1922

Index Fungorum number: IF553238

Xylariaceae, *Xylariales*, *Sordariomycetes*, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, treated as *Clypeosphaeriaceae*), Hongsan et al. (2017a, b; phylogeny, evolution, treated *Clypeosphaeriaceae* as a synonym of *Xylariaceae*), holotype: HKU (M) 1485, cultures and sequences are unavailable.

Type: *Palmaria montaneus* (K.D. Hyde, J. Fröhl. & Joanne E. Taylor) K.D. Hyde, J. Fröhl. & Joanne E. Taylor, **comb. nov.**

Index Fungorum number: IF553239

= *Palmomyces montaneus* K.D. Hyde, J. Fröhl. & Joanne E. Taylor, *Sydiowia* 50(1): 60 (1998)

Holotype: HKU (M) 1485

See Hyde et al. (1998) for the illustrations and description

Notes: The genus *Palmomyces* K.D. Hyde, J. Fröhl. & Joanne E. Taylor (Hyde et al. 1998) is illegitimate, being a later homonym of *Palmomyces* Maire 1922. Thus, the new name *Palmaria* K.D. Hyde, J. Fröhl. & Joanne E. Taylor, nom. nov. is introduced to replace this illegitimate name.

Pandanicola K.D. Hyde 1994, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *P. calocarpa* (Syd. & P. Syd.) K.D. Hyde, asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Pannaria Delise ex Bory 1828 (= *Amphinomium* Nyl. 1888; = *Lepidogium* Clem. & Shear 1931; = *Lepidoleptogium* A.L. Sm. 1922; = *Pannaria* subgen. *Lepidoleptogium* (A.L. Sm.) P.M. Jørg. 2004; = *Trachyderma* Norman 1853 [1852] *fide* Species Fungorum 2017), *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 60 species, type: *P. rubiginosa* (Thunb.) Delise, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Lumbsch et al. (2011; new species), Passo and Calvelo (2011; new species), Elvebakken (2012; new species), Jørgensen and Gjerde (2012; new species), Kirk et al. (2013; genus accepted), Magain and Sérusiaux (2014; phylogeny), cultures and sequences are available.

Pannoparmelia (Müll. Arg.) Darb. 1912 (= *Anzia* sect. *Pannoparmelia* Müll. Arg. 1889 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *P. angustata* (Pers.) Zahlb., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Thell et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Panorbis J. Campb., J.L. Anderson & Shearer 2003, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, three species, type: *P. viscosus* (I. Schmidt) J. Campb., J.L. Anderson & Shearer, on debris near sea, Germany, USA, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Pantospora Cif. 1938 (= *Dictyocephala* A.G. Medeiros 1962 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. guazumae* Cif., hyphomycetous, sexual morph unknown, plant pathogenic, terrestrial, see Minnis et al. (2011; morphology, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), cultures and sequences are available, epitype and ex-type of type: BPI 880778; CBS 130299.

Paoayensis Cababela, Jeewon & K.D. Hyde 2007, *Sordariomycetes* genera *incertae sedis*, one species, type: *P. lignicola* Cababela, Jeewon & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type species HKU(M) 17516, holotype of type: HKUCC 10927.

Papilionospora V.G. Rao & B. Sutton 1976, *Ascomycota* genera *incertae sedis*, one species, type: *P. aspergilloides* (Speg.) V. Rao & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Myanmar, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Papilionovela Aptroot 1997, *Hypocreaceae*, *Amphisphaeriales*, *Sordariomycetes*, 17 species, type: *P. albothallina* Aptroot, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable, holotype of type: CBS Aptroot 33014.

Pappimyces B. Sutton & Hodges 1975, *Ascomycota* genera *incertae sedis*, one species, type: *P. hastatus* B. Sutton & Hodges, hyphomycetous, sexual morph unknown, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Papulaspora Preuss 1851 (= *Baryeidamia* H. Karst. 1888; = *Helicosporangium* H. Karst. 1865 *fide* Species Fungorum 2017), *Coronophorales* genera *incertae sedis*, *Sordariomycetes*, 24 species, type: *P. sepedonioides* Preuss, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Ahmed et al. (2016; DNA, phylogeny), cultures and sequences are available, type specimen for type species B.

Papulosa Kohlm. & Volk.-Kohlm. 1993, *Papulosaceae*, *Diaportheomycetidae* families *incertae sedis*, *Sordariomycetes*, one species, type: *P. amerospora* Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and cultures available for the type species IMS J.K. 5491; AFTOL-ID 748.

Paraoria R.K. Verma & Kamal 1987, Ascomycota genera *incertae sedis*, one species, type: *P. himalayana* R.K. Verma & Kamal, coelomycetous, sexual morph unknown, Nepal, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 294092.

Paraarthrocladium Matsush. 1993, Ascomycota genera *incertae sedis*, one species, type: *P. amazonense* Matsush, hyphomycetous, sexual morph unknown, Peru, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: Matsushima Fungus Collection Kobe OP-146, need generic revision.

Parabagliettoa Gueidan & Cl. Roux 2009, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, three species, type: *P. dufourii* (DC.) Gueidan & Cl. Roux, asexual morph unknown, lichenicolous, terrestrial, Europe, Asia, see Diederich and Schultz (2009), Gueidan et al. (2009; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available, available specimen and cultures for the type: G; AFTOL-ID 2254.

Parabambusicola Kaz. Tanaka & K. Hiray. 2015, *Parabambusicolaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. bambusina* (Teng) Kaz. Tanaka & K. Hiray., asexual morph unknown, saprobes, terrestrial, Japan, see Tanaka et al. (2015; morphology, phylogeny), cultures and sequences are available.

Parablastocatena Y.D. Zhang & X.G. Zhang 2012, Ascomycota genera *incertae sedis*, one species, type: *P. tetraceriae* Y.D. Zhang & X.G. Zhang, hyphomycetous, sexual morph unknown, saprobes, terrestrial, China, see Zhang et al. (2012h; taxonomy), Wijayawardene et al. (2017; outline), holotype of the type: HSAUP H3357, cultures and sequences are unavailable.

Paraboeremia Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *P. selaginellae* (Sacc.) Q. Chen & L. Cai, coelomycetous, saprobes, ?pathogens, terrestrial, cosmopolitan, see Chen et al. (2015b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), neotype and ex-neotype of the type: HMAS 246693, CBS 122.93.

Paracainiella Lar. N. Vassiljeva 1983, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type: *P. dryadis* Lar. N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, arctic north-eastern Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable, holotype of type species VLA P-2943.

Paracamarosporium Wijyaw. & K.D. Hyde 2014, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five

species, type: *P. psoraleae* (Crous & M.J. Wingf.) Wijyaw. & K.D. Hyde, asexual morph coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Africa, USA, see Hyde et al. (2013; accepted in *Didymosphaeriaceae*), Wijayawardene et al. (2014c, d, 2016b; outline, phylogeny), Crous et al. (2015c; new species, phylogeny), Thambugala et al. (2017a, b; new species), cultures and sequences are available, holotype and ex-type cultures available for the type: CBS H-21440; CPC 21632.

Paracapsulospora Konta. & K. D. Hyde 2016, *Sordariomycetidae*, genera *incertae sedis*, one species, type: *P. metroxyli* Konta. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Hyde et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 15-0704, MFLUCC 15-0250.

Paraceratocladium R.F. Castañeda 1987, Ascomycota genera *incertae sedis*, six species, type: *P. silvestre* R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Asia, Caribbean, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paracercospora Deighton 1979, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *P. egenula* (Syd.) Deighton, hyphomycetous, sexual morph unknown, plant pathogenic, terrestrial, see Crous et al. (2012a; morphology; phylogeny), Braun et al. (2013; morphology), cultures and sequences are available, reference specimen and culture of type: CBS H-20836; CPC 12537, holotype and isotype of type species PRE 25999; IMI 89597, genus requires epitypification.

Parachionomyces Thaung 1979, Ascomycota genera *incertae sedis*, one species, type: *P. acroconidiellinae* Thaung, asexual morph hyphomycetous, sexual morph unknown, Myanmar, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Parachnopeziza Korf 1978, *Arachnopezizaceae*, *Helotiales*, *Leotiomycetes*, eight species, type: *P. miniopsis* (Ellis) Korf, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paracladophialophora Crous 2016, *Chaetothyriales* genera *incertae sedis*, *Eurotiomycetes*, one species, type: *P. carceris* Crous & Roets, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Crous et al. (2016a; new species, phylogeny), holotype and ex-type of the type: CBS H-22865, CPC 27596.

Paracoccidioides F.P. Almeida 1930 (= *Lutziomyces* O.M. Fonseca 1939 *fide* Species Fungorum 2017), *Ajellomycetaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *P. brasiliensis* (Splend.) F.P. Almeida, hyphomycetous, sexual morph unknown, human pathogen, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), de Melo Teixeira et al. (2014; clinical control), cultures and sequences are available.

Paracollema Otálora & Wedin 2014, *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *P. italicum* (B. de Lesd.) Otálora, P.M. Jørg. & Wedin, asexual morph coelomycetous, lichenized, terrestrial, Southern Europe, Northern Africa, see Otálora et al. (2014; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Paraconiothyrium Verkley 2004, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *P. estuarinum* Verkley & M. da Silva, asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Damm et al. (2008c; taxonomy, phylogeny), de Gruyter et al. (2012; new species), Wijayawardene et al. (2012, 2014c, 2016b; outline, phylogeny, taxonomy), Hyde et al. (2013; accepted in *Didymosphaeriaceae*), Verkley et al. (2014; new species, phylogeny), Ariyawansa et al. (2015a; new species, phylogeny), cultures and sequences are available, holotype and cultures available for the type species CBS H-10528, CCT6596 =CBS 109850.

Paracostantinella Subram. & Sudha 1989 (= *Xenobotrytis* R.F. Castañeda & W.B. Kendr. 1990 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *P. pulchra* Subram. & Sudha, hyphomycetous, sexual morph unknown, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Paracremonium L. Lombard & Crous 2015, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *P. inflatum* L. Lombard & Crous, asexual morph unknown, pathogens, saprobes, terrestrial, tropical, see Lombard et al. (2015b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Lynch et al. (2016; new species, phylogeny), Wijayawardene et al. (2017; outline), holotype and cultures of the type: CBS H-21946; CBS 110348.

Paracryptophiale Kuthub. & Nawawi 1994, *Ascomycota* genera *incertae sedis*, two species, type: *P. kamaruddinii* Kuthub. & Nawawi, hyphomycetous, sexual morph unknown, Malaysia, China, see Seifert et al. (2011; morphology), Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: KLU 300030.

Paracylindrocarpone Crous, Roets & L. Lombard 2016, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. aloicola* Crous, Roets & L. Lombard., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22613, CPC 27362.

Paracytospora Petr. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *P. salicis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paradendryphiella Woudenberg & Crous 2013, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. salina* (G.K. Sutherl.) Woudenberg & Crous, hyphomycetous, sexual morph unknown, saprobes, France, Europe, see Woudenberg et al. (2013; morphology, phylogeny), Jones et al. (2015; outline, phylogeny), cultures and sequences are available, holotype of the type species and culture ex-type were not mentioned.

Paradendryphiopsis M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, five species, type: *P. cambreensis* M.B. Ellis, hyphomycetous, sexual morph unknown, Europe, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type species IMI 6151c, need generic revision.

Paradictyoarthrinium Matsush. 1996, *Paradictyoarthrinaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. diffractum* Matsush., hyphomycetous, sexual morph unknown, Transvaal, Thailand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Isaka et al. (2015; DNA, secondary metabolites), Liu et al. (2015a; new species, Thailand, phylogeny), Doilom et al. (2017, description), cultures and sequences are available, holotype and ex-type available for the type: Matsushima Fungus Collection Kobe 5A135; MFC-5A135.

Paradidymobotryum C.J.K. Wang & B. Sutton 1984, *Ascomycota* genera *incertae sedis*, one species, type: *P. oblongum* C.J.K. Wang & B. Sutton, asexual morph hyphomycetous, sexual morph unknown, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Paradiplococcum Hern.-Restr. 2017, *Castanediellaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *X. lusitanica* Hern.-Restr., J. Mena & Guarro, hyphomycetous, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Paradiplodia Speg. ex Trotter 1931, *Ascomycota* genera *incertae sedis*, six species, type: *P. aurantiorum* (Speg.) Clem. & Shear, coelomycetous, sexual morph unknown, widespread, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Paradischloridium Bhat & B. Sutton 1985, *Ascomycota* genera *incertae sedis*, one species, type: *P. ychaffrei* Bhat & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, New Zealand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paradiscula Petr. 1941, *Ascomycota* genera *incertae sedis*, one species, type: *P. spuria* (Vestergr.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paradoxa Mattir. 1935, *Tuberaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *P. monospora* Mattir., asexual morph unknown, saprobes, hypogeous, terrestrial, Italy, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list genus), cultures and sequences are available, need generic revision.

Paradoxomyces Matzer 1996, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. nymanii* Matzer, sexual morph with lirellate ascomata, lichenicolous, terrestrial, Indonesia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, holotype of type: Nyman 87b (UPS).

Paraepicoccum Matsush. 1993, *Ascomycota* genera *incertae sedis*, one species, type: *P. amazonense* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Peru, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2017a; description, molecular data), cultures and sequences are unavailable.

Parafulvia Kamal, A.N. Rai & Morgan-Jones 1983, *Ascomycota* genera *incertae sedis*, one species, type: *P. indica* Kamal, A.N. Rai & Morgan-Jones., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Parafuscosporella J. Yang & K.D. Hyde 2016, *Fuscosporellaceae*, *Fuscosporellales*, *Sordariomycetes*, one species, type: *P. moniliformis* J. Yang, Bhat & K.D. Hyde, hyphomycetous, saprobes, Thailand, see Boonyuen et al. (2016; new species), Yang et al. (2016c; taxonomy, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Paragnomonia Senan. & K.D. Hyde 2017, *Sydiowellaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *P. fragariae* (Kleb.) Senan. & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Europe, see Senanayake et al. (2017; taxonomy, phylogeny), cultures and sequences are available.

Parahaplotrichum W.A. Baker & Partr. 2001, *Ascomycota* genera *incertae sedis*, one species, type: *P. idahoense* W.A. Baker & Partr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paraharknessia Matsush. 2003, *Ascomycota* genera *incertae sedis*, one species, type: *P. ellipsoidea* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parahyalotopsis Nag Raj 1976, *Ascomycota* genera *incertae sedis*, one species, type: *P. borassi* (Thaung) Nag Raj., coelomycetous, sexual morph unknown, saprobes, terrestrial, Myanmar, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paraidriella Hern.-Restr. & Crous 2015, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. jambosae* (R.F. Castañeda & W.B. Kendr.) Hern.-Restr. & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Hernández-Restrepo et al. (2016b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Paraingaderia Ertz & Tehler 2011, *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. placodioidea* Ertz & Tehler, sexual morph with punctiform to shortly lirelliform ascomata in groups (synascomata), lichenized, terrestrial, Socotra island, see Ertz and Tehler (2011; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available, holotype of type: Tehler 9344 (S).

Parainoa Resl & T. Sprib. 2014, *Baeomycetaceae*, *Baeomycetales*, *Lecanoromycetes*, one species, type:

P. subconcolor (Anzi) Resl & T. Sprib., lichenized, see Resl et al. (2015; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available.

Paralecanographa Ertz & Tehler 2011, *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. grumulosa* (Dufour) Ertz & Tehler, lichenicolous (becoming sometimes lichenized), terrestrial, mainly (sub-)tropical, Mediterranean, see Ertz and Tehler (2011; morphology, phylogeny), sequences are available.

Paralecia Brackel, Greiner, Peršoh & Rambold 2015, *Squamariaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *P. pratorum* Brackel, Greiner, Peršoh & Rambold, asexual morph unknown, lichenized, terrestrial, worldwide, see Liu et al. (2015a; taxonomy, phylogeny), cultures and sequences are available, holotype of the type: M-0045925.

Paraleptosphaeria Gruyter, Verkley & Crous 2012, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *P. nitschkei* (Rehm ex G. Winter) Gruyter, Aveskamp & Verkley, asexual morph coelomycetous, saprobes, terrestrial, worldwide, see de Gruyter et al. (2012; morphology, phylogeny), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015b; phylogeny), Tibpromma et al. (2017b; new species), cultures and sequences are available, epitype and ex-epitype of the type: CBS H-20822; CBS 306.51.

Paralethariicola Calatayud, Etayo & Diederich 2001, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *P. aspiciliae* Calat., Etayo & Diederich, asexual morph unknown, lichenicolous, non-lichenized, Italy and Spain, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable, holotype of the type species MA (Fungi) 49914.

Paraliomyces Kohlm. 1959, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. lentifer* Kohlm., asexual morph unknown, saprobes, marine, worldwide, see Suetrong et al. (2009; phylogeny, morphology), Jones et al. (2009b, 2015; classification), Pang and Jheng (2012a; checklist), Zhang et al. (2012e; morphology), Wijayawardene et al. (2014c; outline), cultures and sequences are available, type specimen of type: CBS H-5921.

Paramassariothea Subram. & Muthumary 1979, *Ascomycota* genera *incertae sedis*, one species, type: *P. falcata* Subram. & Muthumary., coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parameliola Hongsanan, Peršoh & K.D. Hyde 2016, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *P. dimocarpi* Hongsanan & K.D. Hyde,

coelomycetous, sexual morph unknown, saprobes, terrestrial, Saudi Arabia, see Li et al. (2016a; taxonomy, phylogeny), holotype of the type: MFLU 15-0045.

Paramenisporopsis Matsush. 2003, *Ascomycota* genera *incertae sedis*, one species, type: *P. undulosetulata* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Paramicrodochium Hern.-Restr. & Crous 2015, *Microdochiaeae*, *Xylariales*, *Sordariomycetes*, one species, type: *P. gracile* (Mouch. & Samson) Hern.-Restr. & Crous, hyphomycetous, sexual morphs unknown, on rabbit dung, terrestrial, worldwide, see Hernández-Restrepo et al. (2016b; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Paramicrothyrium H.X. Wu & K.D. Hyde 2011, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *P. chinensis* H.X. Wu & K.D. Hyde, saprobes, terrestrial, Philippines, see Wu et al. (2011; taxonomy), Hyde et al. (2013; accepted as a genus in *Microthyriaceae*), cultures and sequences are available, holotype and ex-type of the type: IFRD 8987; IFRDCC2258.

Paramphisphaeria F.A. Fernández, J.D. Rogers, Y.M. Ju, Huhndorf & Umaña 2004, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. costaricensis* F.A. Fernández, J.D. Rogers, Y.M. Ju, Huhndorf & L. Umaña, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Senanayake et al. (2015; accepted as a genus in *Xylariaceae*), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable, holotype of the type: INB 3466410.

Paramycoleptodiscus Crous & M.J. Wingf. 2016, *Acrospermales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *P. albizziae* Crous & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22615, CPC 27552.

Paramycosphaerella Crous & Jol. Roux 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 15 species, type: *P. brachystegia* Crous & Jol. Roux, asexual morph hyphomycetous (zasmidium-like), foliicolous, terrestrial, worldwide, see Crous et al. (2013b; morphology, phylogeny), Quaedvlieg et al. (2014; phylogeny), Guatimosim et al. (2016a, b; phylogeny, new comb.), cultures and sequences are available, holotype and ex-type culture of type: CBS H-21445; CPC 21136.

Paramyrothecium L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, twelve species, type: *P. roridum* (Tode) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, from soil, terrestrial, cosmopolitan, see Lombard et al. (2016; taxonomy, phylogeny), epitype and ex-epitype of the type: CBS H-22438, CBS 357.89.

Paranannizziopsis Sigler, Hambl. & Paré 2013, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, four species, type: *P. australasiensis* Sigler, Hambleton & Paré, asexual morph hyphomycetous, animal pathogens, USA, see Sigler et al. (2013; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype of the type: UAMH 11645, cultures available for the type: UAMH 10439, UAMH 11719.

Paranectria Sacc. 1878 (= *Ciliomyces* Höhn. 1906 *fide* Species Fungorum 2017), *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *P. affinis* (Grev.) Sacc., asexual morph unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Paraniesslia K.M. Tsui, K.D. Hyde & Hodgkiss 2001, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. tuberculata* K.M. Tsui, K.D. Hyde & Hodgkiss, asexual morph unknown, saprobes, terrestrial, Hong Kong, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype: HKU (M) 4647.

Parapenidiella Crous & Summerell 2012, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *P. tasmaniensis* (Crous & M.J. Wingf.) Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Crous et al. (2012c; morphology; phylogeny), cultures and sequences are available, holotype of the type: PREM 55340.

Parapericonia M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, one species, type: *P. angusii* M.B. Ellis., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa and Australia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parapericoniella U. Braun, Heuchert & K. Schub. 2005, *Ascomycota* genera *incertae sedis*, one species, type: *P. asterinae* (Deighton) U. Braun, Heuchert & K. Schub., hyphomycetous, terrestrial, fungicolous, Africa, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of type: (K(M) IMI 11851b).

Paraphaeoisaria de Hoog & Morgan-Jones 1978, *Ascomycota* genera *incertae sedis*, one species, type: *P. alabamensis* de Hoog & Morgan-Jones., hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paraphaeomoniella Crous 2015, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, one species, type: *P. capensis* Crous & A.R. Wood, coelomycetous, sexual morph unknown, on living leaves, endophytic, terrestrial, South Africa, see Crous et al. (2015d; taxonomy, phylogeny), cultures and sequences are available, holotype and ex-type culture of type: CBS H-20159; CPC 15416.

Paraphaeosphaeria O.E. Erikss. 1967, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 33 species, type: *P. michotii* (Westend.) O.E. Erikss., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014a; taxonomy, phylogeny), Verkley et al. (2014; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), Liu et al. (2015a; taxonomy, phylogeny), Tanaka et al. (2015; taxonomy, phylogeny), Tibpromma et al. (2017b; new species), Thambugala et al. (2017b, phylogeny, new species), Crous et al. (2017; new species), sequence data available for 12 species, holotype for type species.

Paraphoma Morgan-Jones & J.F. White 1983, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *P. radicina* (McAlpine) Morgan-Jones & J.F. White, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see de Gruyter et al. (2012; phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Quaedvlieg et al. (2013; new species, phylogeny), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*), cultures and sequences are available.

Parapithomyces Thaung 1976, *Ascomycota* genera *incertae sedis*, one species, type: *P. brideliae* Thaung., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Myanmar and Australia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parapleurotheciopsis P.M. Kirk 1982, *Beltraniaceae*, *Amphisphaerales*, *Sordariomycetes*, one species, type: *P. inaequiseptata* (Matsush.) P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, terrestrial, British Isles, Japan, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanan et al. (2017a; phylogeny, evolution), DNA sequence available, available specimen and cultures for the type: IMI 234741; MUCL 41089.

Paraporpidia Rambold & Pietschm. 1989, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *P. aboriginum* Rambold, asexual morph unknown, lichenized, on soil and boulders, terrestrial, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parapyrenis Aptroot 1991, *Requienellaceae*, *Xylariales*, *Sordariomycetes*, eight species, type: *P. aurora* (Zahlbr.) Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parapyricularia M.B. Ellis 1972, *Ascomycota* genera *incertae sedis*, one species, type: *P. musae* M.B. Ellis & Peregrine., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pararobillarda Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *P. caffra* Matsush., coelomycetous, sexual morph unknown, saprobes, terrestrial, Cape Province, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Parasarcopodium Mel'nik, S.J. Lee & Crous 2004, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *P. ceratocaryi* Mel'nik, S.J. Lee & Crous, hyphomycetous, saprobes, terrestrial, South Africa, see Lombard et al. (2016; phylogeny), Tibpromma et al. (2016c; holomorph), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype and ex-type culture of type: PREM 57603; STE-U 5494.

Parasedosporium Gilgado, Gené, Cano & Guarro 2007, *Microascaceae*, *Microascales*, *Sordariomycetes*, two species, type: *P. tectonae* (C. Booth) Gilgado, Gené, Cano & Guarro, asexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available, available specimen and ex-type strain of the type species CBS H-7195; CBS 127.84.

Paraschismatomma Ertz & Tehler 2011, *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. ochroleucum* (Zahlbr.) K. Knudsen, Ertz & Tehler, sexual morph with apothecoid ascocarps, lichenized, terrestrial, Californian coast and Baja

California, see Ertz and Tehler (2011; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Parascutellinia Svrček 1975, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, three species, type: *P. violacea* (Velen.) Svrček, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type species.

Parasiphula Kantvilas & Grube 2006, *Coccotremataceae*, *Pertusariales*, *Lecanoromycetes*, seven species, type: *P. fragilis* (Hook. f. & Taylor) Kantvilas & Grube, asexual morph unknown, lichenized, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Paraspphaeropsis Petr. 1953, *Ascomycota* genera *incertae sedis*, one species, type: *P. shearii* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Hawaii, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parastagonospora Quaedv., Verkley & Crous 2013, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *P. nodorum* (Berk.) Quaedv., Verkley & Crous, coelomycetous, pathogenic, saprobes, terrestrial, Germany, Netherlands, Denmark, see Quaedvlieg et al. (2013; taxonomy, phylogeny), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*), Wijayawardene et al. (2014c, 2017; outline), Crous et al. (2016a; new species, phylogeny), Li et al. (2015c, 2016a; new species), Thambugala et al. (2017b, new species, phylogeny), culture of type: CBS 110109 (no ex-type cultures).

Parastenella J.C. David 1991 (= *Stenellopsis* Morgan-Jones 1980 *fide Species Fungorum* 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *P. magnoliae* (Weedon) J.C. David, coelomycetous, sexual morph unknown, saprobes, associated with leaf spots, USA, Brazil (Rondônia), Taiwan, see Dornelo-Silva et al. (2008; new species), Wijayawardene et al. (2012, 2017; outline), Kirschner and Chen (2010; new species), cultures and sequences are available but lacking for the type species.

Parasterinopsis Bat. 1960, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, three species, type: *P. sersalisiae* (Hansf.) Bat., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, keys), Hongsanan et al. (2014c; taxonomy, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, isotype of type: URM 11923.

Parastigmataea Doidge 1921 (= *Hypostigme* Syd. 1925 *fide Species Fungorum* 2017), *Polystomellaceae*, *Dothideomycetes* families *incertae sedis*, three species, type: *P. nervisita* Doidge, sexual morph unknown, widespread (tropical), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Polystomellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable for type species.

Parastigmatellina Bat. & C.A.A. Costa 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. asiatica* Bat. & C.A.A. Costa, coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines and Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parasypodiella Ponnappa 1975, *Parasypodiellaceae*, *Coronophorales*, *Sordariomycetes*, seven species, type: *P. laxa* (Subram. & Vittal) Ponnappa, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Cheewangkoon et al. (2009; new species, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), cultures and sequences are available, specimens available for the type: IMI 50645, IMI 173095h, cultures available for the type: CBS 102698.

Parateratosphaeria Quaedvly. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *P. bellula* (Crous & M.J. Wingf.) Quaedvly. & Crous, asexual morph unknown, pathogens, terrestrial, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), Crous et al. (2017; new species), cultures and sequences are available, epitype and ex-epitype strain of the type: CBS H-20094; CPC 1821.

Paratetraploa M.K.M. Wong & K.D. Hyde 2002, *Ascomycota* genera *incertae sedis*, one species, type: *P. exappendiculata* M.K.M. Wong, Goh & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, aquatic, Hong Kong, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type species HKU (M) 12646.

Parathyridaria Jaklitsch & Voglmayr 2016, *Thyridariaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *P. ramulicola* Jaklitsch, Fourn. & Voglmayr, asexual morph coelomycetous, saprobes, France, Germany, see Jaklitsch and Voglmayr (2016; taxonomy, phylogeny), Tibpromma et al. (2017b; new species), holotype of the type: WU 36867.

Paratomenticola M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, one species, type: *P. lanceolata* (Cooke)

M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Paratopeliopsis Mercado-Díaz, Lücking & Parnmen 2014, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *P. caraibica* Mercado-Díaz, Lücking & Parnmen, asexual morph unknown, lichenized, terrestrial, worldwide, see Mercado-Díaz et al. (2014; taxonomy), Lumbsch et al. (2014a, b; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type: Mercado-Díaz 836.

Paratricharia Lücking 1997, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *P. paradoxa* (Lücking) Lücking, asexual morph unknown, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: UPS (114).

Paratricharina Van Vooren, U. Lindemann, M. Vega, Ribes, Illescas & Matočec 2015, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *P. poiraultii* (Boud.) Van Vooren, U. Lindemann, M. Vega, Ribes, Illescas & Matočec, asexual morph unknown, saprobes, terrestrial, Europe, see sequences are available, voucher specimen of the type: U.L. 178-14.

Paratrichoconis Deighton & Piroz. 1972, *Ascomycota* genera *incertae sedis*, four species, type: *P. chinensis* (Hansf.) Deighton & Piroz., hyphomycetous, sexual morph unknown, saprobes or fungicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paratrichophaea Trigaux 1985, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, five species, type: *P. macrocystis* Trigaux, asexual morph unknown, from soil, saprobes, terrestrial, Europe, North America, see Benkert (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available but lacking for the type species.

Paraulocladium R.F. Castañeda 1986, *Ascomycota* genera *incertae sedis*, two species, type: *P. angulosporum* R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parauncinula S. Takam. & U. Braun 2005, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, two species, type: *P. septata* (E.S. Salmon) S. Takam. & U. Braun, biotrophic, obligate plant pathogens, terrestrial, Asia (China, Japan), see Lumbsch and Huhndorf (2010;

outline), Braun and Cook (2012; taxonomy), sequences are available, lectotype of type: TNS-F-214605.

Paravalsa Ananthap. 1990, *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *P. indica* Ananthap., asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Parawiesneriomycetes Crous & M.J. Wingf. 2016, *Tubeufiales*, *Wiesneriomycetales*, *Dothideomycetes*, one species, type: *P. syzygii* Crous & M.J. Wingf., hyphomycetous, saprobes, terrestrial, France, see Crous et al. (2016b; taxonomy, phylogeny), Bezerra et al. (2017a; phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22630, CPC 26528.

Parberya C.A. Pearce & K.D. Hyde 2001, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *P. kosciuskoia* C.A. Pearce & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: DAR 28023.

Parencelia Petr. 1950, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, four species, type: *P. andina* Petr., asexual morph unknown, saprobes, terrestrial, Central America, West Indies, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parenglerula Höhn. 1910, *Englerulaceae*, *Dothideomycetes* families *incertae sedis*, nine species, type: *P. macowaniana* (Thüm.) Höhn., asexual morph unknown, epiphytic, terrestrial, tropical region, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Dai et al. (2014c; morphology), cultures and sequences are unavailable.

Pareutypella Y.M. Ju & J.D. Rogers 1996, *Sordariomycetes* genera *incertae sedis*, one species, type: *P. sulcata* Y.M. Ju & J.D. Rogers, asexual morph unknown, saprobes, terrestrial, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Parmelia Ach. [nom. cons.] 1803 (= *Aspidelia* Stirt. 1900 [1899]; = *Lichen* L. 1753 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 50 species, type: *P. saxatilis* (L.) Ach., lichenized, worldwide, see Crespo et al. (2010a, b; phylogeny), Kirk et al. (2013; genus accepted), Lumbsch and Huhndorf (2010; outline), Divakar et al. (2012, 2013a, b; India, phylogeny), Ferencova et al. (2014; new species), Lücking et al.

(2017a; classification), cultures and sequences are available.

Parmelia Müll. Arg. 1862, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 90 species, type: *P. triptophylla* (Ach.) Müll. Arg., lichenized (with cyanobacteria), terrestrial (saxicolous, corticolous or lignicolous), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ekman et al. (2014; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Parmelina Hale 1974 (= *Parmeliomyces* E.A. Thomas ex Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, seven species, type: *P. tiliacea* (Hoffm.) Hale, lichenized (with green algae), terrestrial (saxicolous, corticolous or lignicolous), Northern Hemisphere with a center of distribution in western North America and Europe, see Crespo et al. (2010a, b; phylogeny), Lumbsch and Huhndorf (2010; outline), Núñez-Zapata et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), sequence data available for six species, including the generic type.

Parmelinella Elix & Hale 1987, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, nine species, type: *P. wallichiana* (Taylor) Elix & Hale, lichenized (with green algae), terrestrial (saxicolous, corticolous or lignicolous), tropical, see Crespo et al. (2010a, b; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Benatti (2014; current status), Lücking et al. (2017a; classification), sequence data available only for the type species.

Parmeliopsis (Nyl.) Nyl. 1866 (= *Foraminella* S.L.F. Mey. 1882; = *Parmelia* subgen. *Parmeliopsis* Nyl. 1861 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, eight species, type: *P. ambigua* (Wulfen) Nyl., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; DNA, phylogeny), Lücking et al. (2017a; classification), Mark et al. (2016; DNA, phylogeny), cultures available for the type: AFTOL-ID 199.

Parmostictina Nyl. 1875 (see Species Fungorum 2017 for synonyms), *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, 15 species, type: *S. hirsuta* Mont., lichenized,?, rejected against *Pseudocyclphellaria* Vain. 1890, see Moncada et al. (2013a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Parmotrema A. Massal. 1860, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 300 species, type: *P. perforatum* (Jacq.) A. Massal., lichenized, terrestrial (corticolous, saxicolous, terricolous or lignicolous), worldwide, but with the center of distribution in tropical regions, see Crespo et al. (2010a, b; phylogeny), Lumbsch

and Huhndorf (2010; outline), Del-Prado et al. (2011; *Parmotrema reticulatum*), Lendemer and Hodkinson (2012b; phylogeny), Divakar et al. (2013b; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Parmotremopsis Elix & Hale 1987, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *P. antillensis* (Nyl.) Elix & Hale, asexual morph unknown, lichenized, terrestrial, Central America, West Indies, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parmularia Lév. 1846 (= *Clypeum* Massee 1896; = *Pycnographa* Müll. Arg. 1890; = *Schneepia* Speg. 1885 *fide Species Fungorum* 2017), *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, 31 species, type: *P. styracis* Lév., asexual morph unknown, pathogens, epiphytes, terrestrial, Brazil, Paraguay, Uruguay, see Inácio and Cannon (2008; morphology), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Guatimosim et al. (2015; phylogeny), cultures and sequences are available but lacking for type material, holotype and epitype of type: PC; VIC 42447.

Parmulariella Henn. 1904, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. vernoniae* Henn., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, isotype of type: K, NY.

Parmulariopsella Sivan. 1970, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. burseracearum* Sivan., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: IMI 34052.

Parmulariopsis Petr. 1954, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. pulchella* Petr., asexual morph unknown, epiphytes, terrestrial, Malaysia, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, isotype of type: IMI 330018.

Parmulina Theiss. & Syd. 1914, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, six species, type: *P. exsculpta* (Berk.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and

sequences are unavailable, neotype of type: K Herb. C.E. Broome 593.

Parodiella Speg. 1880 (= *Diplodiopsis* Henn. 1904; = *Pyrenophaetina* Syd. & P. Syd. 1916 *fide Species Fungorum* 2017), *Parodiellaceae*, *Dothideomycetes* families *incertae sedis*, four species, type: *P. perisporioides* (Berk. & M.A. Curtis) Speg., asexual morph unknown, biotrophic, terrestrial, America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parodiellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, available specimen of type: CBS H-15496.

Parodiellina Henn. ex G. Arnaud 1918, *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. manaosensis* (Henn.) G. Arnaud, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Paropodia Cif. & Bat. 1956, *Dothideomycetes* genera *incertae sedis*, type: *P. intermedia* Cif. & Bat., data inadequate, see Kirk et al. (2008; treated as a synonym of *Triposporiopsis*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Parorbiliopsis Spooner & Dennis 1986 (= *Eupezizella* Höhn. 1926; = *Orbiliopsis* Höhn. 1926; = *Parorbiliopsis* Spooner & Dennis 1986 [1985]; = *Truncicola* Velen. 1934 *fide Species Fungorum* 2017), *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *P. minuta* Spooner & Dennis, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Parvacoccum R.S. Hunt & A. Funk 1988, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *P. pini* R.S. Hunt & A. Funk, asexual morph unknown, saprobes, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: DAVFP 23419.

Parvomyces Santam. 1995, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. merophysiae* Santam., asexual morph unknown, biotrophic, terrestrial, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paroplaca Arup, Søchting & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, five species, type: *P. tirolensis* (Zahlbr.) Arup, Søchting & Frödén, asexual morph unknown, lichenized, terrestrial,

worldwide, see Arup et al. (2013; taxonomy, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, culture available for the type: U1289, U1290.

Parvothecium L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. terrestre* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, from soil, terrestrial, Brazil, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), epitype and ex-epitype of the type: CBS H-4380, CBS 198.89.

Paschelkiella Sherwood 1987, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *P. pini* (Romell) Sherwood, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Paspalomyces Linder 1933, *Ascomycota* genera *incertae sedis*, one species, type: *P. aureus* Linder, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Passalora Fr. 1849, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, c. 250 species, type: *P. bacilligera* Fr. & Mont., hyphomycetous, sexual morph unknown, plant pathogens, terrestrial, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Phengsintham et al. (2013; species in Laos), Moreno-Rico et al. (2014; new species), Braun et al. 2015; new species), Rossman et al. (2016a; new combination), cultures and sequences are available.

Passeriniella Berl. 1890 (= *Santiella* Tassi 1900 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, seven species, type: *P. dichroa* (Pass.) Berl., asexual morph unknown, saprobes, marine habitat, see Suetrong et al. (2009; phylogeny, morphology), Zhang et al. (2009b, 2012e; phylogeny), Jones et al. (2009b, 2015; outline, marine), Alias et al. (2010; diversity), Besitudo et al. (2010; diversity), Lee et al. (2012; checklist), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Passerinula Sacc. 1875 (= *Spegazzinula* Sacc. 1833 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, one species, type: *P. candida* Sacc., asexual morph unknown, saprobes or fungicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Patellaria Fr. 1822 (= *Lecanidion* Endl. 1830 *fide* Species Fungorum 2017), *Patellariaceae*, *Patellariales*,

Dothideomycetes, twelve species, type: *P. atrata* (Hedw.) Fr., asexual morph unknown, saprobes, terrestrial, temperate, see Schoch et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in *Patellariaceae*, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; taxonomy), cultures and sequences are available, cultures available for the type: CBS 958.97, BCC 28876, holotype of type: CBS H-17437.

Patellariopsis Dennis 1964, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, five species, type: *P. clavispora* (Berk. & Broome) Dennis, asexual morph unknown, saprobes, terrestrial, Europe, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Patescospora Abdel-Wahab & El-Sharouny 2002, *Aliquandostipitaceae*, *Jahnulales*, *Dothideomycetes*, one species, type: *P. separans* Abdel-Wahab & El-Shar., asexual morph unknown, saprobes, freshwater, Egypt, see Suetrong et al. (2010; phylogeny), cultures and sequences are available, holotype of the type: IMI 386405, ?

Patinella Sacc. 1875 (= *Odontoschizon* Syd. & P. Syd. 1914 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *P. hyalophaea* Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures are available for the type (unpublished).

Patinellaria P. Karst. 1885, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. sanguinea* (Pers.) P. Karst., asexual morph unknown, saprobes, terrestrial, Europe, Mexico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Patriciomyces D. Hawksw. 2001, *Ascomycota* genera *incertae sedis*, one species, type: *P. valentinianus* D. Hawksw., hyphomycetous, sexual morph unknown, lichenicolous, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures are sequences unavailable.

Pauahia F. Stevens 1925, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. sideroxyli* F. Stevens, asexual morph unknown, saprobes, terrestrial, Hawaii, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded from *Meliolaceae* and placed in *Dothideomycetes*), cultures and sequences are unavailable.

Paucispora Thambug., Kaz. Tanaka & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. quadrispora* (K. Hiray. & Kaz. Tanaka)

Thambug., Kaz. Tanaka & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Japan, see Thambugala et al. (2015b; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of type: HHUF 27321; MAFF 239455.

Paucithecium Lloyd 1923, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. rickii* Lloyd, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Paulia Fée 1836 (= *Pasithoe* M.J. Decne. 1840 *fide* Species Fungorum 2017), *Lichenaceae*, *Lichinales*, *Lichenomycetes*, ten species, type: *P. pullata* Fée, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available but lack for the type.

Paulkirkia Wijayaw., Wanasa., Tangthirasunun, Camporesi & K.D. Hyde 2016, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. arundinis* Wijayaw., Wanasa., Tangthirasunun, Camporesi & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2016b, 2017; taxonomy, phylogeny, outline), holotype and ex-type of the type: MFLU 13-0315, MFLUCC 12-0328.

Paurocotylis Berk. 1855, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *P. pila* Berk., asexual morph unknown, from soil, hypogeous, terrestrial, Europe, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wang et al. (2016c; DNA, phylogeny), cultures and sequences are available, voucher specimen of the type: Trappe 12583 (OSC).

Payosphaeria H.Y.M. Leung 1990, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. minuta* H.Y.M. Leung, asexual morph unknown, marine, on wood, mangrove, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: IMI 327472.

Peccania A. Massal. ex Arnold 1858 (= *Corinophoros* A. Massal. 1856; = *Peccaniomyces* Cif. & Tomas. 1953; = *Peccaniopsis* M. Choisy 1949; = *Pleoconis* Clem. 1909 *fide* Species Fungorum 2017), *Lichenaceae*, *Lichinales*, *Lichenomycetes*, eight species, type: *P. coralloides* (A. Massal.) A. Massal., asexual morph unknown, lichenized, terrestrial, Europe, North America, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Pectenia P.M. Jørg., L. Lindblom, Wedin & S. Ekman 2014, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, four species, type: *P. plumbea* (Lightf.) P.M. Jørg., L. Lindblom, Wedin & S. Ekman, lichenized, terrestrial, Europe, Africa, mostly along the Atlantic coast, see Ekman et al. (2014; taxonomy), cultures and sequences are available.

Pectinotrichum Varsavsky & G.F. Orr 1971, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *P. llanense* Varsavsky & G.F. Orr, asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence available for the type, available specimen of type: CBS H-7460.

Pedumispora K.D. Hyde & E.B.G. Jones 1992, *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *Pedumispora rhizophorae* K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, mangroves, marine, Republic of Seychelles, Guam, Thailand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Klaysuban et al. (2014; DNA, phylogeny), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), sequence data available, holotype of type: BRIP 19201.

Peethambara Subram. & Bhat 1978, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. sundara* Subram. & Bhat, asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Lombard et al. (2016; phylogeny), cultures and sequences are available, ex-type culture of type: CBS 646.77.

Peethasthabeeja P. Rag. Rao 1981, *Ascomycota* genera *incertae sedis*, one species, type: *P. krishna* P. Rag. Rao, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pellionella (Sacc.) Sacc. 1902 (= *Diplodiella* subgen. *Pellionella* Sacc. 1899; = *Pellionella* (Sacc.) Petch 1924 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *P. deformans* Penz. & Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Java, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pellucida Dulym., Sivan., P.F. Cannon & Peckally 2001, *Hypocreaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *P. pendulina* Dulym., Sivan., P.F. Cannon & Peckally, asexual morph unknown, saprobes, terrestrial,

Mauritius, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: Mauritius (Dulymamode P6).

Peloronectria Möller 1901, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. vinosa* Möller, asexual morph unknown, South America, see Kirk et al. (2008; treated as nom. dub.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Peltaster Syd. & P. Syd. 1917, *Dothideomycetes* genera *incertae sedis*, seven species, type: *P. hedyotidis* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes or pathogens, terrestrial, Philippines, South America, Slovenia, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Medjedović et al. (2014; new species, phylogeny), cultures and sequences are available but lacking for the type.

Peltasterella Bat. & H. Maia 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. ocoteae* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peltasterinostroma Punith. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *P. rubi* Punith., coelomycetous, sexual morph unknown, saprobes, terrestrial, British Isles, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peltasteropsis Bat. & H. Maia 1959, *Ascomycota* genera *incertae sedis*, seven species, type: *P. discreta* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peltigera Willd. [nom. cons.] 1787 (see Species Fungorum 2017 for synonyms), *Peltigeraceae*, *Peltigerales*, *Lecanoromycetes*, 100 species, type: *P. canina* (L.) Willd., lichenized (green algae or, and cyanobacteria), terrestrial (saxicolous, terricolous, corticolous or lignicolous) or aquatic, worldwide, see Sérusiaux et al. (2009b; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014b; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available.

Peltigeromyces Möller 1901 (= *Peltophoromyces* Clem. 1909 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *P. microsporus* Möller, asexual morph unknown,

saprobes, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärte et al. (2017; phylogeny), cultures and sequences are available for unidentified species.

Peltistroma Henn. 1904, *Ascomycota* genera *incertae sedis*, one species, type: *P. juruanum* Henn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, lectotype of type: K (Mycotheca Brasiliensis no. 93).

Peltistromella Höhn. 1907, *Ascomycota* genera *incertae sedis*, one species, type: *P. brasiliensis* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peltosoma Syd. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *P. freycinetiae* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peltostromellina Bat. & A.F. Vital 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. atra* Bat. & A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, Sri Lanka, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peltostromopsis Bat. & A.F. Vital 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. artocarpi* Bat. & A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, Sri Lanka, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peltula Nyl. 1853, *Peltulaceae*, *Lichinales*, *Lichenomycetes*, 32 species, type: *P. radicata* Nyl., asexual morph unknown, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), McDonald et al. (2013; DNA, phylogeny), Marques et al. (2013; new species, Portugal), sequences are available.

Peltularia R. Sant. 1944, *Coccocarpiaceae*, *Peltigerales*, *Lecanoromycetes*, three species, type: *P. gyrophoroides* (Räsänen) R. Sant., asexual morph unknown, lichenized, terrestrial, South America, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pemphidium Mont. 1840, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, 17 species, type: *P. nitidum* Mont., coelomycetous, common on palms, terrestrial, South America, Australia, Indonesia, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, type specimen of type: CBS H-6336.

Pendulispora M.B. Ellis 1961, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. venezuelanica* M.B. Ellis, sexual morph unknown, hyphomycetous, on dead branches, Venezuela, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Penicillifer Emden 1968 (= *Viridispora* Samuels & Rossman 1999 *fide* Maharachchikumbura et al. 2015b, 2016b; Lombard et al. 2016; Rossman et al. 2016a), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, seven species, type: *P. pulcher* Emden, asexual morph hyphomycetous, saprobes, from soil, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Lombard et al. (2016; phylogeny), Rossman et al. (2016a; nomenclature), cultures and sequences are available, type specimen and culture available for type: CBS H-7461; CBS 560.67.

Penicilliopsis Solms 1887 (= *Pseudocordyceps* Hauman 1936; = *Sarophorum* Syd. & P. Syd. 1917 *fide* Species Fungorum 2017), *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, three species, type: *P. clavariiformis* Solms, asexual morph unknown, saprobes, terrestrial, worldwide, see Houbraken and Samson (2011; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Trichocomaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are available, lectotype and culture available for type: Hauman (BR); CBS 761.68.

Penicillium Link 1809 (see Species Fungorum 2017 for synonyms), *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, c. 300 species, type: *P. crustaceum* Link, hyphomycetous, pathogens, saprobes, widespread (common moulds), see Houbraken and Samson (2011; morphology, phylogeny), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Zhou et al. (2016; new species, polluted soil), Tibpromma et al. (2017b; new species), cultures and sequences available.

Penidiella Crous & U. Braun 2007, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, six species, type: *P. columbiana* Crous & U. Braun, hyphomycetous, sexual morph teratosphaeria-like, pathogens, associated with leaf spots, terrestrial, worldwide, see Crous et al. (2009e; morphology, phylogeny), Crous and Groenewald (2011; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, outline), cultures and sequences are

available, epitype and ex-epitype strain of the type: CBS H-9937; CBS 486.80.

Penidiellopsis Sandoval-Denis, Gené, Deanna A. Sutton & Guarro 2016, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. radicularis* Sandoval-Denis, Gené, Deanna A. Sutton & Guarro, hyphomycetous, human pathogen, terrestrial, USA, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22389, CBS 140695.

Pentagenella Darb. 1897 (= *Roccella* sect. *Pentagenella* (Darb.) Follmann 1998 *fde* Species Fungorum 2017), *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *P. fragillima* Darb., lichenized, terrestrial, neotropical, see Lumbsch and Huhndorf (2010; outline), Ertz and Tehler (2011; new species, phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Penzigomyces Subram. 1992, *Ascomycota* genera *incertae sedis*, 13 species, type: *P. nodipes* (Penz. & Sacc.) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Perelegamycetes R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, one species, type: *P. parviechinulatus* W.B. Kendr. & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Perennicordyceps Matočec & I. Kušan 2014, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *P. prolifica* (Kobayasi) Matočec & I. Kušan, asexual morph unknown, South America, see Matočec et al. (2014; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Periamphispore J.C. Krug 1989, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *P. phacelodes* J.C. Krug, asexual morph unknown, on dung, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariales*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Periconia Tode 1791 (= *Cephalotrichum* Berk. ex Sacc. 1886; = *Harpocephalum* G.F. Atk. 1897; = *Pachytrichum* Syd. 1925; = *Sporocybe* Fr. 1825; = *Sporodum* Corda 1837; = *Trichocephalum* Costantin 1888 *fide* Species Fungorum 2017), *Periconiaceae*, *Pleosporales*, *Dothideomycetes*, c. 45 species, type: *P. lichenoides* Tode, hyphomycetous,

terrestrial, aquatic, worldwide, see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Jones et al. (2015; outline, marine), Tanaka et al. (2015; phylogeny), Thambugala et al. (2017b, new species, phylogeny), sequence data available.

Periconiella Sacc. 1885 (= *Acrocladium* Petr. 1949; = *Acrodesmis* Syd. 1926 *fide* Species Fungorum 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 70 species, type: *P. velutina* (G. Winter) Sacc., hyphomycetous, sexual morph unknown, plant pathogens, terrestrial, see cultures and sequences are available, Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), epitype and ex-epitype of type: CBS H-15612; CBS 101948.

Peridiothelia D. Hawksw. 1985, *Pleomassariaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *P. fuliguncta* (Norman) D. Hawksw., asexual morph unknown, on bark, Europe, terrestrial, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pleomassariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, lectotype of type: O.

Periglandula U. Steiner, Leistner & Leuchtm. 2011, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *P. ipomoeae* U. Steiner, E. Leistner & Schardl, asexual morph unknown, on living leaves, terrestrial, Ecuador, Panama, see Steiner et al. (2011; taxonomy, outline), Beaulieu et al. (2015; DNA), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Perigrapha Hafellner 1996, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, four species, type: *P. superveniens* (Nyl.) Hafellner, lichenicolous, terrestrial, Asia, Europe and North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Perischizon Syd. & P. Syd. 1914, *Dothideomycetes* genera *incertae sedis*, three species, type: *P. oleifolium* (Kalchbr. & Cooke) Syd. & P. Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted in *Parmulariaceae*), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; excluded from *Parmulariaceae*), Wijayawardene et al. (2014c; outline, accepted in *Dothideomycetes*), cultures and sequences unavailable, Holotype of type: K (P. MacOwan 7).

Perisporiopsis Henn. 1904 (see Species Fungorum 2017 for synonyms), *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, 17 species, type: *P. struthanthi* Henn., asexual morph unknown, saprobes, endophytes, terrestrial, tropical, see Chaverri and Gazis (2010, 2011; new species, endophytes), Kawada

et al. (2010; antitumour compounds), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Peristomialis (W. Phillips) Boud. 1907, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, c. five species, type: *P. berkeleyi* Boud., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Perizomella Syd. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *P. inquinans* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Central America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peroneutypa Berl. 1902, *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, 47 species, type: *P. bellula* (Desm.) Berl., asexual morph unknown, saprobes, pathogens, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Dai et al. (2017; morphology, phylogeny), Shang et al. (2017, new species, phylogeny), cultures and sequences are available but lacking for type.

Peroschaeta Bat. & A.F. Vital 1957, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. artocarpi* Bat. & A.F. Vital, asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Perrotia Boud. 1901, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, 19 species, type: *P. flammea* (Alb. & Schwein.) Boud., asexual morph unknown, saprobes, terrestrial, worldwide, see Monaghan et al. (2009; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny), sequence available for the type.

Persicospora P.F. Cannon & D. Hawksw. 1982, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *P. moreauii* P.F. Cannon & D. Hawksw., fungicolous, widespread, terricolous, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; position), cultures and sequences are available, holotype of type: K(M)-IMI 238745.

Perthomyces Crous 2017, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *P. podocarpi* Crous, coelomycetous, sexual morph unknown, saprobes, Australia, see Crous et al. (2017; taxonomy), holotype: CBS H-23122, cultures and sequences are available.

Pertusaria DC. [nom. cons.] 1805, *Pertusariaceae*, *Pertusariales*, *Lecanoromycetes*, 500 species, type: *P. communis* DC., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Perusta Egidi & Sielow 2014, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *P. inaequalis* Egidi & Sielow, hyphomycetous, sexual morph unknown, rock-inhabiting fungus, Spain, see Egidi et al. (2014; taxonomy), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2014c, 2017; outline), type: CBS 118271 = TRN272 cultures and sequences available.

Pesotum J.L. Crane & Schokn. 1973 (= *Phialographium* H.P. Upadhyay & WB. Kendr. 1974 *fide* Seifert et al. 2011), *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, seven species, type: *P. ulmi* (M.B. Schwarz) J.L. Crane & Schokn., hyphomycetous, saprobes, pathogens, worldwide, see Kamgan Nkuekam et al. (2008; new species), Grobbelaar et al. (2009; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Pestalopezia Seaver 1942, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *P. brunneopruinosa* (Zeller) Seaver, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pestalotiopsis Steyaert 1949 (= *Discosiosis* Edward et al. 1972; = *Pestalosphaeria* M.E. Barr 1975; = *Robillarda* Castagne 1845 *fide* Réblová et al. 2016c; Species Fungorum 2017), *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, c. 300 species, type: *P. maculans* (Corda) Nag Raj, coelomycetous, sexual morph formerly named in *Pestalosphaeria*, saprobes or pathogens on plants, worldwide, see Maharanachikumbura et al. (2011, 2012, 2013, 2014a, b, 2016a, c; phylogeny, morphology), Zhang et al. (2012j, 2013f; species in ornamental plants), Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Réblová et al. (2016c; nomenclature), Chen et al. (2017a; new species), 56 species have ex-type sequences.

Pestalozziella Sacc. & Ellis ex Sacc. 1882, *Ascomycota* genera *incertae sedis*, four species, type: *P. inquinans* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Petchiomycetes E. Fisch. & Mattir. 1938, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *P. thwaitesii* (Berk. & Broome) E. Fisch. & Mattir., asexual morph unknown, saprobes, terrestrial, North America, Sri Lanka, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peterozyma Kurtzman & Robnett 2010, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *P. toletana* (Socias, C. Ramírez & Peláez) Kurtzman & Robnett, asexual reproduction is by multilateral budding, saprophytic, tanning fluid, insect frass from trees and tree bark, Europe, North America, see Kurtzman and Suzuki (2010; taxonomy), sequences are available.

Petractis Fr. 1845, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, six species, type: *P. exanthematica* (Sm.) Fr., asexual morph unknown, lichenized, terrestrial, worldwide, see Orange (2009a; new species, DNA), Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Petrakia Syd. & P. Syd. 1913 (= *Echinospodium* Woron. 1913 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, three species, type: *P. echinata* (Peglion) Syd. & P. Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Asia, see Wijayawardene et al. (2012, 2017; outline), Butin et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), Li et al. (2016a; phylogeny, taxonomy), cultures available for the type: OH 080921.4.

Petrakiella Syd. 1924, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *P. insignis* Syd., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Petrakina Cif. 1932, *Dothideomycetes* genera *incertae sedis*, two species, type: *P. mirabilis* Cif., asexual morph unknown, epiphytes, terrestrial, see Hofmann (2009; review, accepted as a genus in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hongsan et al. (2014c; taxonomy, keys, accepted as a genus in *Dothideomycetes*), Wijayawardene et al. (2014c; outline, accepted in *Dothideomycetes*), cultures and sequences unavailable, syntype of type: NY 00936794.

Petrakiopeltis Bat., A.F. Vital & Cif. 1958, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. byrsionimae* Bat., A.F. Vital & Cif., asexual morph unknown, epiphytes, terrestrial, Guyana, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in

Microthyriaceae), Wu et al. (2011; not accepted as *Microthyriaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013; not accepted as a genus in *Microthyriaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*), cultures and sequences are unavailable.

Petrakiopsis Subram. & K.R.C. Reddy 1968, *Ascomycota* genera *incertae sedis*, one species, type: *P. elegans* Subram. & K.R.C. Reddy, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Petriella Curzi 1930, *Microascaceae*, *Microascales*, *Sordariomycetes*, two species, type: *P. asymmetrica* Curzi, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available, available specimen and cultures for the type: CBS H-7548; CBS 238.38.

Petrophila de Hoog & Quaedvlyl. 2014, *Extremaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. incerta* de Hoog & Quaedvlyl., hyphomycetous, sexual morph unknown, rock-inhabiting fungus, Spain, see Egidi et al. (2014; taxonomy), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2017; outline), type: CBS 118608 = TRN139b, cultures and sequences available.

Peyritschia Thaxt. 1890, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 47 species, type: *P. curvata* Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peyronelia Cif. & Gonz. Frag. 1927, *Ascomycota* genera *incertae sedis*, six or seven species (*fide* Seifert et al. 2011) but Index Fungorum (2017) lists only four epithets, type: *P. sirodesmiooides* Cif. & Gonz. Frag., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Peyronellaea Goid. ex Togliani emend. Aveskamp, Gruyter & Verkley 2010, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, c. 20 species, type: *P. glomerata* (Corda) Goid. ex Togliani, asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Aveskamp et al. (2009, 2010; generic amendment, treated as a distinct genus), Wijayawardene et al. (2012, 2014c, 2017; outline), cultures and sequences are available.

Pezicula Tul. & C. Tul. 1865 (= *Cryptosporiopsis* Bubák & Kabát 1912; = *Lagynodella* Petr. 1922; = *Ocellaria* (Tul. & C. Tul.) P. Karst. 1871 *fide* Johnston et al. 2014b; Jaklitsch et al. 2016a), *Dermateaceae*, *Helotiales*, *Leotiomycetes*, c. 50 species, type: *P. carpinea* (Pers.) Tul. ex Fuckel, asexual morph *Cryptosporiopsis*, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Chen et al. (2016b; new species), Ekanayaka et al. (2016a; new species, Thailand), epitype and cultures available for the type: CBS H-17476; CBS 923.96, CBS 921.96.

Peziza Dill. ex Fr. 1822 (see Species Fungorum 2017 for synonyms), *Pezizaceae*, *Pezizales*, *Pezizomycetes*, more than 100 species, type: *P. vesiculosa* Bull., asexual morph unknown, hypogeous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Ashraf et al. (2012; Pakistan, ectomycorrhizae), Barseghyan and Solomon (2011; Israel). Kirk et al. (2013; genus accepted), cultures and sequences are available, available specimen and cultures for type: CBS H-15930; AFTOL-ID 507.

Pezolepis Syd. 1925, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *P. denigrata* Syd., asexual morph unknown, saprobes, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pezoloma Clem. 1909 (= *Ciliatula* Velen. 1922; = *Pseudodiscinella* Dennis 1956 [1955]; = *Sphagnicola* Velen. 1934 *fide* Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, 14 species, type: *P. denigrata* Clem., asexual morph unknown, saprobes, terrestrial, temperate, see Huhtinen et al. (2010; new species, Sweden), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Bruzone et al. (2016; DNA), cultures and sequences are available but lacking for the type.

Pezomela Syd. 1928, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. saxe gotha ae* Syd., asexual morph unknown, saprobes, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärte et al. (2017; phylogeny), cultures and sequences are available for unidentified species.

Phacellium Bonord. 1860, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, 27 species, type: *P. in honestum* Bonord., hyphomycetous, saprobes, pathogens, terrestrial, Asia, Europe, North America, South America, see Crous et al. (2009d; DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Phaciella P. Karst. 1884, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. salicina* P. Karst., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014d; new species, treated as in *Ostropales*), cultures and sequences are available but lacking for the type, needs generic revision since have different opinions.

Phacidina Höhn. 1917, *Leptopeltidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. gracilis* (Niessl) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Leptopeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phaciopycnis Potebnia 1912 (= *Potebniamyces* Smerlis 1962, = *Discosporiopsis* Petr. 1921 *fide* Johnston et al. 2014b), *Phaciaceae*, *Phacidiales*, *Leotiomycetes*, five species, type: *P. malorum* Potebnia, asexual morph coelomycetous, saprobes, pathogens terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014b; phylogeny), Johnston et al. (2014b; nomenclature), Díaz et al. (2015; pathogens, DNA), cultures and sequences are available but lacking for the type.

Phacidium Fr. 1815 (= *Ceuthospora* Grev. 1826; = *Basilocula* Bubák 1914; = *Coccobolus* Wallr. 1833; = *Melanostroma* Corda 1829; = *Phaciostroma* Höhn. 1917; *Siroplaconema* Petr. 1922 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Phaciaceae*, *Phacidiales*, *Leotiomycetes*, 270 species, type: *P. lacerum* Fr., asexual morph formerly named in *Ceuthospora*, saprobes, plant pathogens, terrestrial, worldwide, see Barbeito et al. (2013; pathogenicity), Petäistö et al. (2013; pathogenicity), Crous et al. (2014b, 2015a, b; phylogeny, new species), Johnston et al. (2014b; nomenclature), Suija et al. (2015b; phylogeny), Wiseman et al. (2016; new disease), cultures and sequences are available, epitype and ex-epitype of type: CBS H-10302; CBS 761.73.

Phacographa Hafellner 2009, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *P. glaucomaria* (Nyl.) Hafellner, sexual morph with rounded ascospores, lichenicolous, terrestrial, Northern Hemisphere, see Hafellner (2009; morphology), Frisch et al. (2014; phylogeny), sequences are available, lectotype of the type: Nylander (H-Nyl. 11019).

Phacopsis Tul. 1852 (= *Nesolechia* A. Massal. 1856 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 22 species, type: *P. vulpina* Tul., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline),

Zhurbenko (2010b; new species), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Phacostroma Petr. 1955, *Ascomycota* genera *incertae sedis*, one species, type: *P. hypodermium* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phacostromella Petr. 1955, *Ascomycota* genera *incertae sedis*, one species, type: *P. coronata* (Fuckel) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Germany, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phacothecium Trevis. 1856, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. varium* (Tul.) Trevis., lichenicolous, terrestrial, Northern Hemisphere, see Hafellner (2009; morphology), sequences are unavailable, epitype of the type: J. Hafellner 47633 & A. Hafellner.

Phaeangellina Dennis 1955 (= *Diehlia* Petr. 1951 *fide* Species Fungorum 2017), *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. empetri* (W. Phillips) Dennis, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeangium Pat. 1894, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *P. lefebvrei* Pat., asexual morph unknown, saprobes, terrestrial, North Africa, Arabian Peninsula, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Zitouni-Haouar et al. (2015; DNA), cultures available for the type: BMBH4 (Zitouni-Haouar et al. 2015).

Phaeoacremonium W. Gams, Crous & M.J. Wingf. 1996 (= *Romelia* Berl. 1900; = *Togninia* Berl. 1900 *fide* Réblová et al. 2016c), *Togniniaceae*, *Togniniales*, *Sordariomycetes*, 50 species, type: *P. parasiticum* (Ajello, Georg & C.J.K. Wang) W. Gams, Crous & M.J. Wingf., hyphomycetous, with acremonium-like synasexual morph, sexual morph formerly named in *Togninia*, pathogens, saprobes, terrestrial, cosmopolitan, see Damm et al. (2008a; phylogeny, species associated with *Prunus* wood, new species) Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2016a; new species, phylogeny), Dai et al. (2017; sexual morph), holotype and ex-type of the type: CBS H-22886, CPC 27637.

Phaeoblastophora Partr. & Morgan-Jones 2002, *Ascomycota* genera *incertae sedis*, two species, type: *P. resinae* (Fr.) Partr. & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Seifert et al. (2011; morphology),

Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Phaeobotryon Theiss. & Syd. 1915, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, four species, type: *P. cercidis* (Cooke) Theiss. & Syd., asexual morph coelomycetous, saprobes, pathogens, terrestrial, cosmopolitan, see Phillips et al. (2008, 2013; taxonomy, phylogeny), Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c, 2016b; outline, taxonomy, phylogeny), Dissanayake et al. (2016b; taxonomy), cultures and sequences are available.

Phaeocalicium A.F.W. Schmidt 1970, *Mycocaliciaceae*, *Mycocaliciales*, *Eurotiomycetes*, eleven species, type: *P. praecedens* (Nyl.) A.F.W. Schmidt, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Tibell et al. (2014; new species), Tuovila et al. (2014; DNA), Telfer et al. (2015; DNA, phylogeny), sequences are available, voucher specimen of the type: Tuovila 09-240 (TUR).

Phaeocandellabrum R.F. Castañeda, Gusmão, Guarro & Iturr. 2009, *Ascomycota* genera *incertae sedis*, three species, type: *P. elegans* (R.F. Castañeda) R.F. Castañeda, Heredia, Saikawa, hyphomycetous, sexual morph unknown, saprobes, terrestrial, neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeocercospora Crous 2012, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. colophospermi* Crous, hyphomycetous, pathogens, South Africa, see Crous et al. (2012a; morphology; phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), sequence data available, holotype and ex-type of the type: CBS H-20966; CPC 19813.

Phaeochora Höhn. 1909, *Phaeochoraceae*, *Phyllachorales*, *Sordariomycetes*, five species, type: *P. chamaeropis* (Cooke) Höhn., saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phaeochorella Theiss. & Syd. 1915, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, six species, type: *P. parinarii* (Henn.) Theiss. & Syd., from living leaves, South Africa, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, available specimen of type: CBS H-15956.

Phaeochoropsis K.D. Hyde & P.F. Cannon 1999, *Phaeochoraceae*, *Phyllachorales*, *Sordariomycetes*, four species, type: *P. neowashingtoniae* (Shear) K.D. Hyde & P.F. Cannon, saprobes, North and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phaeococcoomyces de Hoog 1979 (= *Melanocryptococcus* Della Torre & Cif. 1964; = *Nigrococcus* E.K. Novák & Zsolt 1961; = *Phaeococcus* de Hoog 1977 fide Species Fungorum 2017), *Phaeococcoomycetaceae*, *Lichenostigmatales*, *Arthoniomycetes*, eight species, type: *P. nigricans* (M.A. Rich & A.M. Stern) de Hoog, hyphomycetous, sexual morph unknown, on human, saprobes, Asia, Europe, North America, South America, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species), Moreno-Rico et al. (2014; new species), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Phaeocryptopus Naumov 1915 (= *Adelopus* Theiss. 1918 [1917]; = *Cryptopus* Theiss. 1914 fide Species Fungorum 2017), *Dothideaceae*, *Dothideales*, *Dothideomycetes*, eight species, type: *P. abietis* Naumov, asexual morph unknown, growing on conifer needles, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; review), cultures and sequences are available, holotype of type: F67760.

Phaeocyrtidula Vain. 1921, *Dothideomycetes* genera *incertae sedis*, two species, type: *P. pinea* (Nyl.) Vain., saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phaeocytostroma Petr. 1921, *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *P. istrica* Petr., coelomycetous, sexual morph unknown, pathogens, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; phylogeny, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Phaeodactylium Agnihothr. 1968, *Ascomycota* genera *incertae sedis*, five species, type: *P. venkatesanum* Agnihothr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Castañeda-Ruiz et al. (2009c, 2013c; new species), Seifert et al. (2011; morphology), Zhang et al. (2011b; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available (unpublished).

Phaeodimeriella Speg. 1908, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, 15 species, type:

P. occulta (Racib.) Speg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Mapook et al. (2016b; taxonomy, phylogeny), cultures and sequences are available, but lacking for the type.

Phaeodiscula Cub. 1891, *Ascomycota* genera *incertae sedis*, one species, type: *P. celottii* Cub., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeodomus Höhn. 1909, *Ascomycota* genera *incertae sedis*, three species, type: *P. lauracearum* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeodothis Syd. & P. Syd. 1904, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *P. tricuspidis* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in *Didymosphaeriaceae*), Ariyawansa et al. (2014e; accepted in *Didymosphaeriaceae*), Wijayawardene et al. (2014c; outline), Verkley et al. (2014; new species, phylogeny), sequences and cultures available only for *P. winteri* but lacking for the type, available specimen for type: CBS H-4793.

Phaeofabraea Rehm 1909, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. miconiae* Rehm, asexual morph unknown, saprobes, terrestrial, South America, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeographis Müll. Arg. 1882, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, c. 180 species, type: *P. dendritica* (Ach.) Müll. Arg., asexual morph unknown, lichenized, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; classification), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Phaeographopsis Sipman 2007, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *P. indica* (Patw. & Nagarkar) Sipman & Aptroot, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2013; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Phaeohelotium Kanouse 1935, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, 41 species, type: *P. flavum* Kanouse,

aseexual morph unknown, saprobes, terrestrial, worldwide, see Hengstmengel (2009; outline), Kaya (2009; Turkey), Lumbsch and Huhndorf (2010; outline), Baral et al. (2013a; new combination), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Phaeohiratsukaea Udagawa & Iwatsu 1990, *Ascomycota* genera *incertae sedis*, one species, type: *P. expansa* Udagawa & Iwatsu, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: National Institute of Hygienic Sciences, Tokyo, NHL2968.

Phaeoidiomyces Dorn.-Silva & Dianese 2004, *Ascomycota* genera *incertae sedis*, one species, type: *P. qualeae* Dorn.-Silva & Dianese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: UB (col. micol.) 9851.

Phaeoisaria Höhn. 1909 (= *Graphiopsis* Bainier 1907; = *Hansfordiula* E.F. Morris 1963; = *Spiculostilbella* E.F. Morris 1963 *fide* Species Fungorum 2017), *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, 20 species, type: *P. bambusae* Höhn., hyphomycetous, sexual morph unknown, pathogens, terrestrial, worldwide, see Seifert et al. (2011; morphology), Mel'nik (2012; new species, Vietnam), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Cheng et al. (2014; new species, China), Crous et al. (2015c; new species), Liu et al. (2015a; new species), Maharanachikumbura et al. (2015b, 2016b; outline), Réblová et al. (2017; phylogeny), cultures and sequences are available but lacking for the type.

Phaeolabrella Speg. 1912, *Ascomycota* genera *incertae sedis*, one species, type: *P. eryngicola* Speg., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeomoniella Crous & W. Gams 2000, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, one species, type: *P. chlamydospora* (W. Gams, Crous, M.J. Wingf. & Mugnai) Crous & W. Gams, hyphomycetous, sexual morph unknown, with coelomycetous synasexual morph, pathogens, saprobes, endophytic, terrestrial, Italy, see Chen et al. (2015a; phylogeny), Crous et al. (2015d; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype of type: CBS H-229.95

as *Phaeoacremonium chlamydosporum* W. Gams et al., reference culture of type: CBS 229.95.

Phaeomonilia R.F. Castañeda, Heredia & R.M. Arias 2007, *Ascomycota genera incertae sedis*, four species, type: *P. pleiomorpha* R.F. Castañeda, Heredia & R.M. Arias, hyphomycetous, sexual morph unknown, saprobes, terrestrial and aquatic, Mexico, see Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Ma et al. (2015; new species), Yang et al. (2016a; new species), Wijayawardene et al. (2017; outline), holotype of the type: XAL CB746, cultures and sequences are unavailable, compare with *Botryomonilia*, *Pseudaegerita*, *Staheliella* and *Oidiodendron*.

Phaeomonostichella Keissl. ex Petr. 1941, *Ascomycota genera incertae sedis*, one species, type: *P. symploci* (Keissl.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeonectriella R.A. Eaton & E.B.G. Jones 1971, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *P. lignicola* R.A. Eaton & E.B.G. Jones, asexual morph unknown, saprobes on wood, marine, temperate, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Phaeopeltosphaeria Berl. & Peglion 1892, *Dothideomycetes genera incertae sedis*, two species, type: *P. caudata* Berl. & Peglion, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phaeophacidium Henn. & Lind 1897, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, four species, type: *P. escalloniae* Henn. & Lindau, asexual morph unknown, saprobes, causing leaf spots, terrestrial, Asia, Australia, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeophleospora Rangel 1916, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, eleven species, type: *P. eugeniae* Rangel, coelomycetous, with mycosphaerella-like sexual morphs, plant pathogens, see Wijayawardene et al. (2012, 2017; outline), Quaedvlieg et al. (2014; morphology; phylogeny), Crous et al. (2015d, 2016b; new species), cultures and sequences are available, neotype of type: IMI 372655.

Phaeophloeosporella Crous & B. Sutton 1997, *Ascomycota genera incertae sedis*, one species, type: *P. ekebergiae* (Syd.) Crous & B. Sutton, coelomycetous, sexual morph unknown, Wijayawardene et al. (2012, 2017;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeophyscia Moberg 1977, *Physciaceae*, *Teloschistales*, *Lecanoromycetes*, 66 species, type: *P. orbicularis* (Neck.) Moberg, foliose lichen, terrestrial, worldwide, see Shukla & Upadhyay (2009; bioindicator), Lendemer (2009; record), Brackel (2011b; Iceland), Arslan et al. (2011; Turkey), Shukla et al. (2012; physiology), Pindaru et al. (2013; chemical composition), Lücking et al. (2017a; classification), cultures and sequences are available for most of the species including type.

Phaeopoacea Thambug., Dissan. & K.D. Hyde 2017, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. festucae* Dissan. & K.D. Hyde, asexual morph coelomycetous, saprobes, Europe, see Thambugala et al. (2017b; taxonomy, phylogeny), holotype of the type: MFLU 17-0121.

Phaeopyxis Rambold & Triebel 1990, *Helotiales genera incertae sedis*, *Leotiomycetes*, five species, type: *P. punctum* (A. Massal.) Rambold, lichenicolous, terrestrial, Europe, Asia, North and South America, Australia, see Diederich et al. (2009; Europe), Motiejūnaitė et al. (2011; Lithuania), Matwiejuk and Bohdan (2011; Poland), Himelbrant et al. (2013; Russia), Suija et al. (2015b; phylogeny), Pino-Bodas et al. (2017; treated as *Ostropomycetidae*), sequences are available.

Phaeoramularia Munt.-Cvetk. 1960, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, more than 100 species names, but genus appears one species, type: *P. gomphrenicola* (Speg.) Munt.-Cvetk., hyphomycetous, plant pathogens, see Videira et al. (2017; morphology, phylogeny), cultures and sequences are available, holotype of type: LPS 914, isotypes of type: BPI 436740, BPI 722393; K (M) IMI 7706 (slide), MICH 15302.

Phaeorrhiza H. Mayrhofer & Poelt 1979, *Physciaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *P. nimbosa* (Fr.) H. Mayrhofer & Poelt, asexual morph unknown, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Resl et al. (2016; phylogeny), cultures and sequences are available, cultures available for the type: T805.

Phaeosaccardinula Henn. 1905, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, 16 species, type: *P. diospyricola* Henn., asexual morph unknown, epiphytes, sooty moulds, terrestrial, China, Thailand, see Chomnunti et al. (2012b; 2014; morphology; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Yang et al. (2014b; morphology; phylogeny), cultures and sequences are available for some species, available specimen of type: CBS H-4260.

Phaeoschizotrichum C.R. Silva, Gusmão & R.F. Castañeda 2015, *Ascomycota* genera *incertae sedis*, one species, type: *P. ramosum* Silva, Gusmão & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Silva et al. (2015b; morphology), cultures and sequences are unavailable, holotype of the type: HUEFS 210423, compare with *Passalora*, *Pseudocercospora* and *Schizotrichum*.

Phaeosclera Sigler, Tsuneda & J.W. Carmich. 1981, *Dothideomycetes* genera *incertae sedis*, two species, type: *P. dematoides* Sigler, Tsuneda & J.W. Carmich., hyphomycetous, sexual morph unknown, terrestrial, saprobes, worldwide, see Raja et al. (2011a; DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, isotype and cultures available for the type: CBS H-7557; CBS 157.81.

Phaeosclerotinia Hori 1916, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. nipponica* Hori, asexual morph hyphomycetous, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeoscypha Spooner 1984, *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. cladii* (Nag Raj & W.B. Kendr.) Spooner, asexual morph unknown, saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Phaeoseptum Ying Zhang, J. Fourn. & K.D. Hyde 2013, *Halothiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. aquaticum* Ying Zhang, J. Fourn. & K.D. Hyde, asexual morph unknown, saprobes, freshwater, France, see Zhang et al. (2013d; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), sequence data available for the type, holotype and ex-type strains of type: IFRD 8986; CBS 123113.

Phaeosperma Nitschke ex Fuckel 1870, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. dryophilum* Nitschke ex G.H. Otth, asexual morph unknown, saprobes, terrestrial, Switzerland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, epitype and ex-epitype strains of type: WU 32035; CBS 133213, LG21.

Phaeosphaeria I. Miyake 1909, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 90 species, type: *P. oryzae* I. Miyake, asexual morph coelomycetous, saprobes or pathogens, terrestrial, worldwide, see Zhang et al. (2012e; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus

accepted), Phookamsak et al. (2014c; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), Tibpromma et al. (2017b; new species), sequence data available for more than 10 species, need adequate sequence data, ex-epitype strains of type: CBS 110110; lectotype of type: S-F9572.

Phaeosphaeriopsis M.P.S. Câmara, M.E. Palm & A.W. Ramaley 2003, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, ten species, type: *P. glaucopunctata* (Grev.) M.P.S. Câmara, M.E. Palm & A.W. Ramaley, asexual morph unknown coelomycetous, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013 accepted as a genus in *Phaeosphaeriaceae*), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*), Wijayawardene et al. (2014c, 2017; outline), Thambugala et al. (2014c; new species, phylogeny), Crous et al. (2016a; new species, phylogeny), Tibpromma et al. (2017b; new species), cultures and sequences are available.

Phaeospora Hepp 1867, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 17 species, type: *P. rimosicola* (Leight. ex Mudd) Hepp, asexual morph unknown, lichenicolous, terrestrial, worldwide, see Etayo (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), van den Boom and Etayo (2014; new species), cultures and sequences are unavailable for the type.

Phaeosporis Clem. 1909, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *P. melasperma* (Nyl.) Clem., asexual morph unknown, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phaeosporobolus D. Hawksw. & Hafellner 1986, six species, *Ascomycota* genera *incertae sedis*, type: *P. usneae* D. Hawksw. & Hafellner, hyphomycetous, sexual morph unknown, lichenicolous, widespread, see Seifert et al. (2011; morphology), Ertz et al. (2015; phylogeny), cultures and sequences are available, holotype of type: K(M)-IMI 281394.

Phaeostagonospora A.W. Ramaley 1997, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. nolinae* A.W. Ramaley, coelomycetous, sexual morph paraphaeosphaeria-like, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Quaedvlieg et al. (2013; accepted phaeostagonospora-like asexual morphs in *Paraphaeosphaeria*), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*), cultures and sequences are unavailable.

Phaeostalagmus W. Gams 1976, *Ascomycota* genera *incertae sedis*, seven species, type: *P. cyclosporus* (Grove) W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology, treated as *Chaetosphaeriaceae*), Wijayawardene et al. (2012, 2017; outline, accepted as a genus in *Ascomycota*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, listed in *Chaetosphaeriaceae*), cultures and sequences are unavailable, available specimen of type: CBS H-19245.

Phaeostigme Syd. & P. Syd. 1917, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, c. 20 species, type: *P. picea* (Berk. & M.A. Curtis) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phaeostilbelloides Armando, Z.M. Chaves & Dianese 2015, *Ascomycota* genera *incertae sedis*, one species, type: *P. velloziae* Armando, Z.M. Chaves & Dianese, hyphomycetous, sexual morph unknown, on living leaves, terrestrial, Brazil, see de Souza Armando et al. (2015; morphology), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, holotype of the type: UBMC 18404, compare with *Myrothecium* and *Phaeostilbella*.

Phaeotheca Sigler et al. 1981 *Capnodiales*, genera *incertae sedis*, *Dothideomycetes*, four species, type: *P. fissurella* Sigler, Tsuneda & J.W. Carmich., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2016b; new species, phylogeny), Type: CBS 520.89, cultures are available.

Phaeothecoidea Crous 2007, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *P. eucalypti* Crous & Summerell, hyphomycetous, sexual morph unknown, pathogens, saprobes, Australia, see Crous et al. (2009e; morphology, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017 outline), cultures and sequences are available, holotype and ex-type culture of type: CBS-H 19836, CPC 12918.

Phaeothecodiella Batzer & Crous 2010, *Phaeothecodiellaceae*, *Dothideomycetes* family, *incertae sedis*, two species, type: *P. missouriensis* Batzer & Crous, hyphomycetous, sexual morph unknown, pathogens, saprobes, USA, see Yang et al. (2010; morphology, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Hongsanan et al. (2017b; phylogeny, family placement), cultures and sequences are available, holotype and ex-type culture of type: CBS H-20322, CPC 16116.

Phaeothyriolum Syd. 1938, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. eucalyptinum* Syd., asexual morph unknown, saprobes, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phaeothyrium Petr. 1947, *Ascomycota* genera *incertae sedis*, one species, type: *P. pulchellum* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeotomaselia Katum. 1981, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. ruwenzorensis* Katum., asexual morph unknown, saprobes, terrestrial, Uganda, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phaeotrichoconis Subram. 1956, *Ascomycota* genera *incertae sedis*, two species, type: *P. crotalariae* (M.A. Salam & P.N. Rao) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaeotrichosphaeria Sivan. 1983, *Sordariomycetes* genera *incertae sedis*, two species, type: *P. indica* Sivan. & N.D. Sharma, asexual morph unknown, pathogens, terrestrial, India, UK, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phaeotrichum Cain & M.E. Barr 1956, *Phaeotrichaceae*, *Phaeotrichales*, *Dothideomycetes*, two species, type: *P. hystricinum* Cain & M.E. Barr, asexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; accepted as a genus in *Phaeotrichaceae*), Hyde et al. (2013; accepted as a genus in *Phaeotrichaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type.

Phaeoxyphiella Bat. & Cif. 1963, *Ascomycota* genera *incertae sedis*, seven species, type: *P. morototoni* Bat. & Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Phaeoxyphium Bat. & Cif. 1963, *Ascomycota* genera *incertae sedis*, type: *P. dubium* (Sacc.) Bat. & Cif., see Kirk et al. (2008; treated as nom. dub.), Kirk et al. (2013; genus accepted).

Phaffomyces Y. Yamada 1997, *Phaffomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, five species, type: *P. opuntiae* (Starmer, Phaff, M. Miranda, M.W. Miller & J.S.F. Barker) Y. Yamada, asexual reproduction is by multilateral budding, saprophytic, necrotic tissue of the columnar cactus (*Cephalocereus rojenii*), the prickly pear cactus (*Opuntia inermis*) and the senita cactus (*Lophocereus schottii*), Australia, USA, islands of the Caribbean, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Phalacrichomycetes R.K. Benj. 1992, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *P. normalis* R.K. Benj., asexual morph unknown, biotrophic, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phalangispora Nawawi & J. Webster 1982, *Ascomycota* genera *incertae sedis*, three species, type: *P. constricta* Nawawi & J. Webster, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available (unpublished).

Phanerococcus Cif. 1954, *Ascomycota* genera *incertae sedis*, two species, type: *P. rapaneae* Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, Santo Domingo, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaneromyces Speg. & Har. 1889, *Phaneromycetaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *P. macrosporus* (Boud.) Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phaulomyces Thaxt. 1931 (= *Adelomyces* Taxt. 1931 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 14 species, type: *P. corylophodis* Thaxt., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phellostroma Syd. & P. Syd. 1914, *Ascomycota* genera *incertae sedis*, one species, type: *P. hypoxyloides* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phialemoniopsis Perdomo, Dania García, Gené, Cano & Guarro 2013, *Sordariomycetes* genera *incertae sedis*, six species, type: *P. ocularis* (Gené & Guarro) Perdomo, Dania García, Gené & Guarro, asexual morph unknown, pathogens on human, endophytes, terrestrial, America,

China, see Perdomo et al. (2013; taxonomy, phylogeny), Desoubeaux et al. (2014; pathogens, phylogeny), Tsang et al. (2014a; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Su et al. (2016d; new species), cultures and sequences are available, cultures available for the type: FMR 6632.

Phialemonium W. Gams & McGinnis 1983, *Cephalothecaceae*, *Sordariomycetes* families *incertae sedis*, six species, type: *P. obovatum* W. Gams & McGinnis, hyphomycetous, sexual morph unknown, saprobes, human pathogen, terrestrial, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Perdomo et al. (2013; new species, phylogeny), Crous et al. (2015d; new species), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures and sequences are available, available specimen and cultures for the type: CBS H-1665; CBS 279.76.

Phialoarthrobotryum Matsush. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *P. triseptatum* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, available specimen for type: CBS H-7561.

Phialoascus Redhead & Malloch 1977, ?*Saccharomycetales* genera *incertae sedis*, ?*Saccharomycetes* one species, type: *P. borealis* Redhead & Malloch, asexual reproduction unknown, mycoparasitic on the mushroom *Cortinarius huronensis*, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phialocephala W.B. Kendr. 1961 (= *Phaeomollisia* T.N. Sieber & Grünig 2009 *fide* Johnston et al. 2014b), *Dermateaceae*, *Helotiiales*, *Leotiomycetes*, 27 species, type: *P. dimorphospora* W.B. Kendr., hyphomycetous, sexual morph formerly named in *Phaeomollisia*, saprobes, terrestrial, aquatic, worldwide, see Grünig et al. (2009; new species), Wang et al. (2009b; new species, Missouri), Lumbsch and Huhndorf (2010; outline), Day et al. (2012; new species), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures available for the type: CBS 300.62.

Phialogeniculata Matsush. 1971, *Ascomycota* genera *incertae sedis*, one species, type: *P. guadalcanalensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Guadeloupe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phialomyces P.C. Misra & P.H.B. Talbot 1964, *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, three species, type: *P. macrosporus* P.C. Misra & P.H.B. Talbot, asexual

morph unknown, saprobes, terrestrial, worldwide, see Houbraken and Samson (2011; taxonomy, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, available specimen and culture for type: CBS H-7563; CBS 430.64.

Phialophaeoisaria Matsush. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *P. spongiosa* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: Matsushima Fungus Collection, Kobe, MFC-2J770.

Phialophora Medlar 1915, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, nine species, type: *P. verrucosa* Medlar, hyphomycetous, mostly saprobes, terrestrial, isolated from soil, decaying plant material and other substrates, also including important clinically-relevant fungi, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Rébllová et al. (2013; reallocation of selected taxa), Madrid et al. (2016; reallocation), Li et al. (2017b; outline, new species), cultures and sequences are available, ex-epitype of the type: CBS 140325.

Phialophorophoma Linder 1944, *Pleosporales* genera *incertae sedis*, one species, type: *P. litoralis* Linder, coelomycetous, sexual morph unknown, marine, saprobes, Europe, USA, see Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), Wijayawardene et al. (2017; outline), cultures and sequences are available, available specimen of type: CBS H-17521.

Phialosporostilbe Mercado & J. Mena 1985, *Ascomycota* genera *incertae sedis*, five species, type: *P. turbinata* Mercado & J. Mena, hyphomycetous, sexual morph unknown, saprobes, in soil, terrestrial, Asia, Caribbean, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available (unpublished).

Phialostele Deighton 1969, *Ascomycota* genera *incertae sedis*, one species, type: *P. scytopetali* Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phialotubus R.Y. Roy & Leelav. 1966, *Ascomycota* genera *incertae sedis*, one species, type: *P. microsporus* R.Y. Roy & Leelav., hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al.

(2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, available specimen of type: CBS H-7598.

Phillipsia Berk. 1881, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, 17 species, type: *P. domingensis* Berk., asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Romero et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for the type: DHP 7169.

Phillipsiella Cooke 1878, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, twelve species, type: *P. atra* Cooke, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Saccardiaceae*), Kirk et al. (2013; genus accepted), da Silva et al. (2016; phylogeny), cultures and sequences are available.

Philobryon Döbbeler 1988, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. anuliferum* Döbbeler, on Bryophyta, Papua New Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Philonectria Hara 1914, *Dothideomycetes* genera *incertae sedis*, three species, type: *P. variabilis* Hara, saprobes, Japan, Uganda, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phloeopeccania J. Steiner 1902, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, four species, type: *P. pulvinulina* J. Steiner, asexual morph unknown, lichenized, terrestrial, Asia, Africa, North America, see Schultz et al. (2009; Namibia, South Africa), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Candan and Schultz (2015; new species, taxonomy), Lücking et al. (2017a; classification), sequence data available for one species.

Phloeoospora Wallr. 1833, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, c. 150 species but need a revision, type: *P. ulmi* (Fr.) Wallr., coelomycetous, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2013; phylogeny), cultures and sequences are unavailable.

Phloeosporina Höhn. 1924, *Ascomycota* genera *incertae sedis*, one species, type: *P. minor* (Ellis & Kellerm.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phlogicylindrium Crous, Summerb. & Summerell 2006, *Phlogicylindriaceae, Amphisphaerales, Sordariomycetes*, three species, type: *P. eucalypti* Crous, Summerb. & Summerell, asexual morph unknown, saprobes, all occurring on *Eucalyptus* species, Australia, see Wijayawardene et al. (2012, 2017; outline), Senanayake et al. (2015; phylogeny, morphology), Crous et al. (2016a; new species, phylogeny), sequence data available for all species, holotype and ex-type of the type: CBS H-19749; CBS 120080.

Phlyctaeniella Petr. 1922, *Ascomycota genera incertae sedis*, two species, type: *P. polonica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Australia, see Tiwari et al. (2012; new species, India), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phlytema Desm. 1847, *Dermateaceae, Helotiales, Leotiomycetes*, c. 30, type: *P. vagabunda* Desm., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2013; new species), Johnston et al. (2014b; treated as a synonym of *Neofabraea*), Chen et al. (2016b; resurrected the genus), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available.

Phlyctis (Wallr.) Flot., 1850, *Phlyctidaceae, Lecanorales, Lecanoromycetes*, 20 species, type: *P. agelaea* (Ach.) Flot., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Fernandez-Brime et al. (2011; phylogeny), Kelly et al. (2011; DNA barcoding), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Weerakoon et al. (2016; new species and records), sequence data available.

Phoebus R.C. Harris & Ladd, 2007, *Arthoniales genera incertae sedis, Arthoniomycetes*, one species, type: *P. hydrophobius* R.C. Harris & Ladd, asexual morph unknown, lichenized, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Phoma Sacc. 1880, *Didymellaceae, Pleosporales, Dothideomycetes*, c. 100 species, type: *P. herbarum* Westend., coelomycetous, sexual morph didymella-like, saprobes, pathogens, terrestrial, worldwide, see de Gruyter et al. (2009, 2010, 2012; taxonomy, phylogeny), Aveskamp et al. (2010; phylogeny), Woudenberg et al. (2009; sexual morph), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Didymellaceae*), Kirk et al. (2013; genus accepted), Chen et al. (2015b; phylogeny), cultures and sequences are available.

Phomachora Petr. & Syd. 1925, *Ascomycota genera incertae sedis*, two species, type: *P. lucida* (Berk. & M.A.

Curtis) Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, America, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phomachorella Petr. 1947, *Ascomycota genera incertae sedis*, one species, type: *P. parinarii* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phomatodes Q. Chen & L. Cai 2015, *Didymellaceae, Pleosporales, Dothideomycetes*, c. 20 species, type: *P. aubrietiae* (Moesz) Q. Chen & L. Cai, coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Chen et al. (2015b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Phomatospora Sacc., 1875, *Phomatosporaceae, Phomatosporales, Sordariomycetes*, c. 100 species, type: *P. berkeleyi* Sacc., saprobes on submerged wood or decaying twigs, widespread, asexual morph sporothrix-like reported from culture, see Raja and Shearer (2008; overview), Lumbsch and Huhndorf (2010; outline), Jones et al. (2015; marine species, outline), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2016; phylogeny), sequences are available for four species but lacking for the type, available specimen of type: CBS H-5533.

Phomatosporella Tak. Kobay. & K. Sasaki 1982, *Ascomycota genera incertae sedis*, one species, type: *P. albomaculans* Tak. Kobay. & K. Sasaki, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, accepted in as a genus in *Sordariomycetes*), cultures and sequences are unavailable.

Phomyces Clem. 1931, *Ascomycota genera incertae sedis*, one species, type: *P. meliolicola* (Speg.) Clem., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phragmaspidium Bat. 1960, *Dothideomycetes genera incertae sedis*, three species, type: *P. corruscans* (Rehm) Bat., asexual morph unknown, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phragmeriella Hansf. 1946, *Pseudoperisporiaceae, Dothideomycetes families incertae sedis*, one species,

type: *P. ireninae* Hansf., asexual morph unknown, saprobes, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phragmitensis K.M. Wong, Poon & K.D. Hyde 1998, *Hyponectriaceae*, *Amphisphaerales*, *Sordariomycetes*, two species, type: *P. marina* M.K.M. Wong, Poon & K.D. Hyde, asexual morph unknown, intertidal, in salt marsh, on *Phragmites*, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable, holotype of type: HKU (M) 5151.

Phragmiticola Sherwood 1987, *Marthamycetaceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *P. rhopalospermum* (Kirschst.) Sherwood, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phragmocamarosporium Wijayaw., Yong Wang bis & K.D. Hyde 2015, *Lentitheciaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. platani* Wijayaw., Yong Wang bis & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, Germany, China, see Wijayawardene et al. (2015, 2016b, 2017; taxonomy, phylogeny), holotype: HGUP NNW G0928-1, cultures and sequences are unavailable.

Phragmocapnia Theiss. & Syd. 1918 (= *Conidiocarpus* Woron. 1927 *fide* Wijayawardene et al. 2014c), *Capnodiales*, *Dothideomycetes*, five species, type: *P. beite* (Syd., P. Syd. & E.J. Butler) Theiss. & Syd., asexual morph unknown, tropical, see Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2011, 2014; accepted as a genus in *Capnodiales*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Phragmocephala E.W. Mason & S. Hughes 1951, *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, nine species, type: *P. cookei* E.W. Mason & S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2017; phylogeny), cultures and sequences are available but lacking for the type.

Phragmoconidium G.F. Sepúlveda, Pereira-Carv. & Dianese 2009, *Ascomycota* genera *incertae sedis*, two species, type: *P. cerradense* G.F. Sepúlveda, Pereira-Carv. & Dianese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al.

(2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phragmodiaporthe Wehm. 1941, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, five species, type: *P. caryae* (Peck) Wehm., asexual morph unknown, saprobes, North America, China, see Vasilyeva et al. (2008; new species, China), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Phragmodiscus Hansf. 1947, *Sordariomycetes* genera *incertae sedis*, two species, type: *P. arundinariae* Hansf., asexual morph unknown, saprobes, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Phragmogibbera Samuels & Rogerson 1990, *Dothideomycetes* genera *incertae sedis*, three species, type: *P. xylariicola* Samuels & Rogerson, saprobes, Venezuela, see Lumbsch and Huhndorf (2010; outline, accepted in *Venturiaceae*), Hyde et al. (2013; not accepted in *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: VEN.

Phragmopeltis Henn. 1904, *Ascomycota* genera *incertae sedis*, five species, type: *P. siparunae* Henn., coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phragmoporthe Petr. 1934, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *P. ploettneriana* (Henn.) Petr., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Sogonov et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Phragmoscutella Woron. & Abramov 1927, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. abchasica* Woron. & Abramov, saprobes, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phragmospathula Subram. & N.G. Nair 1966, *Ascomycota* genera *incertae sedis*, one species, type: *P. phoenicis* Subram. & N.G. Nair, hyphomycetous,

sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phragmospathulella J. Mena & Mercado 1986, *Ascomycota* genera *incertae sedis*, one species, type: *P. matsushimaiae* J. Mena & Mercado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phragmosperma Theiss. & Syd. (1917) [1916], *Dothideomycetes* genera *incertae sedis*, type: *P. marattiae* (Henn.) Theiss. & Syd., data inadequate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable,

Phragmotrichum Kunze 1823, *Ascomycota* genera *incertae sedis*, four species, type: *P. chailletii* Kunze, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Asia, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, available specimen for type: CBS H-16912.

Phthora d'Hérelle 1909, *Ascomycota* genera *incertae sedis*, one species, type: *P. vastatrix* d'Hérelle, asexual morph unknown, saprobes, terrestrial, Guatemala, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phurmomyces Thaxt. 1931, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. obtusus* Thaxt., asexual morph unknown, saprobes, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phycomelaina Kohlm., 1968, *Phyllachorales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. laminariae* (Rostr.) Kohlm., asexual morph unknown, on marine alga *Laminaria*, temperate, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Phyllachoraceae*), Kirk et al. (2013; genus accepted), Jones et al. (2015; review), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Phyllachorales*), cultures and sequences are unavailable.

Phycarella Döbbeler 1980, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. scytonematis* Döbbeler, parasites, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Phylacia Lév. 1845 (= *Henningsina* Möller 1901; = *Leveillea* Fr. 1849 *fide* Wendt et al. 2017), *Hypoxylaceae*,

Xylariales, *Sordariomycetes*, eight species, type: *P. globosa* Lév., asexual morph unknown, terrestrial, saprobes, mainly coprophilous, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available.

Phyllachora Nitschke ex Fuckel 1870, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, c. 1000 species (need revision), type: *P. graminis* (Pers.) Fuckel, saprobes, North and South America, see Lumbsch and Huhndorf (2010; outline), Santos et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), DNA sequence available for the voucher UB21307 of the type, available specimen of type: CBS H-5212.

Phyllachorella Syd. 1914, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, one species, type: *P. micheliae* Syd., asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, available specimen of type: CBS H-4052.

Phyllactinia Lév. 1851, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, 111 species, type: *P. guttata* (Wallr. : Fr.) Lév., asexual morph hyphomycetous, biotrophic, obligate plant pathogens, terrestrial, worldwide, see Takamatsu et al. (2008b, 2016; phylogeny), Lumbsch and Huhndorf (2010; outline), Braun and Cook (2012; taxonomy), Kirk et al. (2013; genus accepted), sequences are available, lectotype of type: G 00122109.

Phylleutypa Petr. 1934, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *P. dioscoreae* (Wakef.) Petr., saprobes, Africa, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phylliscidiopsis Sambo 1937, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *P. abissinica* Sambo., asexual morph unknown, lichenized, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Phylliscidium Forssell 1885, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *P. monophyllum* (Kremp.) Forssell, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline) Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Phyllisciella Henssen 1984, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, three species, type: *P. marionensis* Henssen, asexual morph unknown, lichenized, terrestrial, Africa, see de Lange et al. (2012; conservation), Lumbsch and Huhndorf (2010; outline) Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Phylliscum Nyl. 1855, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, eight species, type: *P. demangeonii* (Moug. & Mont.) Nyl., asexual morph unknown, lichenized, terrestrial, worldwide, see Bajpai et al. (2009; biodeterioration), Sheik et al. (2009, 2016; taxonomy), Lumbsch and Huhndorf (2010; outline), Anderson and Neily (2012, Nova Scotia), Prieto et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Goni et al. (2014; Jammu), Shukla et al. (2014b; diversity), Lücking et al. (2017a; classification), sequences are available.

Phyllobaeis Kalb & Gierl 1993, *Baeomycetaceae*, *Baeomycetales*, *Lecanoromycetes*, six species, type: *P. imbricata* (Hook.) Kalb & Gierl, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Cao et al. (2013; new species, phylogeny), Kirk et al. (2013; genus accepted), Flakus et al. (2015; taxonomy), Mittermeier et al. (2015; metabolic profiling), Lücking et al. (2017a; classification), sequences are available.

Phyllobathelium (Müll. Arg.) Müll. Arg. 1890, *Strigulaceae*, *Strigulales*, *Dothideomycetes*, five species, type: *P. epiphyllum* (Müll. Arg.) Müll. Arg., asexual morph unknown, lichenized fungi, terrestrial, tropical, see Nelsen et al. (2009, 2011b; description, phylogeny), Lücking (2008; partial monograph), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Lücking et al. (2017a; classification), cultures and sequences are available.

Phylloblastia Vain. 1921, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, twelve species, type: *P. dolichospora* Vain., asexual morph unknown, lichenicolous, terrestrial, pantropical, see Flakus and Lücking (2008; new species, Bolivia), Lücking (2008; new species), Lumbsch and Huhndorf (2010; outline), McCarthy (2010; new species), Kirk et al. (2013; genus accepted), McCarthy and Stajsic (2013; new species), van den Broeck et al. (2014; new species), Lücking et al. (2017a; classification), cultures and sequences are unavailable for the type.

Phyllocelis Syd. 1925, *Sordariomycetes* genera *incertae sedis*, two species, type: *P. oyedaeae* Syd., saprobes, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phyllocratera Sérusiaux & Aptroot 1997, *Strigulaceae*, *Strigulales*, *Dothideomycetes*, two species, type: *P. papuana* Sérusiaux & Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted in *Strigulaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: LG (Lambinon 87.427).

Phyllocrea Höhn. 1918, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *P. quitensis* (Pat.) Höhn., from living leaves, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phylloedium Fr. 1825, *Ascomycota* genera *incertae sedis*, one species, type: *P. epiphylla* Fr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phyllogalyidea Lücking & Aptroot 2008, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *P. epiphylla* (Vězda) Lücking & Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lücking (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Phyllohendersonia Tassi 1902, *Ascomycota* genera *incertae sedis*, 25 species, type: *P. mahoniae* Unamuno, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phyllopezis Petr. 1949, *Leotiomycetes* genera *incertae sedis*, one species, type: *P. andina* Petr., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Phylloporthe Syd. 1925, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *P. vernoniae* Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Phyllopsora Müll. Arg. 1894, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, 70 species, type: *P. breviuscula* (Nyl.) Müll. Arg. (*Lecidea breviuscula* Nyl.), lichenized, corticolous, tropical, extending into

oceanic temperate, see Timdal (2008; Peru), Lücking et al. (2017a; classification), sequence data available.

Phyllosticta Pers. 1818 (= *Guignardia* Viala & Ravaz 1892 *fide* Wijayawardene et al. 2014c), *Phyllostictaceae*, *Botryosphaerales*, *Dothideomycetes*, c. 100 species, type: *P. convallariae* Pers., coelomycetous, sexual morph formerly named in *Guignardia*, saprobes, pathogens, endophytes, terrestrial, worldwide, see Wulandari et al. (2009; pathogens), Glienke et al. (2011; taxonomy), Wikee et al. (2011, 2013a, b; review), Su and Cai (2012; phylogeny), Hyde et al. (2013; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline, phylogeny), Wu et al. (2014c; species in banana), cultures and sequences are available, available specimen of type: CBS H-16994.

Phymatotrichopsis Hennebert 1973, *Rhizinaceae*, *Pezizales*, *Pezizomycetes*, 17 species, type: *P. omnivora* (Duggar) Hennebert, hyphomycetous, sexual morph unknown, pathogens, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Physalidiella Rulamort 1990, *Ascomycota* genera *incertae sedis*, three species, type: *P. elegans* (Mosca) Rulamort, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, Europe, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Physalidiopsis R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, one species, type: *P. carbonis* (R.F. Castañeda) R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Physalospora Niessl 1876, *Hypocretriaceae*, *Amphisphaerales*, *Sordariomycetes*, 37 species, type: *P. alpestris* Niessl, asexual morph unknown, terrestrial, aquatic, saprobes and endophytic, widespread, see Raja and Shearer (2008; new species, freshwater), Polashock et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available for three species.

Physalosporopsis Bat. & H. Maia 1955, *Dothideomycetes* genera *incertae sedis*, 17 species, type: *P. rhizophoricola* Bat. & H. Maia, asexual morph unknown, pathogens, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus

accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Physcia (Schreb.) Michaux 1803, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, 80 species, type: *P. tenella* (Scop.) DC., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny), Prieto et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2015g; new species), Prieto and Wedin (2016; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Physcidia Tuck 1862, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, ten species, type: *P. wrightii* (Tuck.) Tuck., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Aptroot and Caceres (2014; key), Dong et al. (2015; new records), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Physciella Essl. 1986, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, four species, type: *P. chloantha* (Ach.) Essl., lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny), Prieto et al. (2013; phylogeny), Prieto and Wedin (2016; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Physconia Poelt 1965, *Physciaceae*, *Lecanorales*, *Lecanoromycetes*, 25 species, type: *P. pulverulenta* (Schreb.) Poelt, asexual morph unknown, lichenized, terrestrial, worldwide, see Lendemer et al. (2009; taxonomy), Begum and HariKrishna (2010; air pollution), Björk (2010; new record), Esslinger and Dillman (2010; new record), Loppi and Nascimbene (2010; bioindicators), Stamenković and Cvijan (2010; air pollution), Stamenković et al. (2010; bioindicators), Lisowska (2011; air pollution), Lumbsch and Huhndorf (2010; outline), Tsurykau et al. (2011; checklist), Yatsyna (2011; taxonomy), Gaya et al. (2012; sequence), Jury (2013; list), Kirk et al. (2013; genus accepted), Bilovitz (2014; biodiversity), Goni et al. (2014; checklist), Karakoti et al. (2014; taxonomy), Paoli et al. (2015; air pollution), Yavuz et al. (2015; epiphytes diversity), Lücking et al. (2017a; classification), Motiejūnaitė et al. (2016; list), sequences are available for few species.

Physma A. Massal. 1854, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 51 species, type: *P. boryanum* (Pers.) A. Massal., asexual morph unknown, lichenized, terrestrial, worldwide, see Wedin et al. (2009b; phylogeny), Diederich et al. (2010b; lectotypification), Otálora et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline), Din et al. (2010; chemoprofile), Jayalal et al. (2013a; Vietnam), Kirk et al. (2013; genus accepted), Kerr (2014; checklist), Magain and Sérusiaux (2014; partial sequence), Ekman et al. (2014; transferred to *Pannariaceae*), Aptroot and

John (2015; taxonomy), Lücking et al. (2017a; classification), Ohmura et al. (2016; Japan), Rangsiruji et al. (2016; phylogeny), sequences are available but lacking for type.

Physmatomyces Rehm 1900, *Leotiomycetes* genera *incertae sedis*, one species, type: *P. compacti* E.A. Thomas ex Cif. & Tomas., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Picardella I.I. Tav. 1985, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *P. endogaea* (F. Picard) I.I. Tav., asexual morph unknown, biotrophic, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Piccolia A. Massal. 1856, *Lecanoromycetes* genera *incertae sedis*, seven species, type: *P. crocea* (Spreng.) A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Piceomphale Svrček 1957, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. bulgarioides* (P. Karst.) Svrček, asexual morph unknown, saprobes, cosmopolitan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Rutstroemiaceae*), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Pichia E.C. Hansen 1904, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, 35 species, type: *P. membranifaciens* (E.C. Hansen) E.C. Hansen, asexual reproduction is by multilateral budding, saprophytic, but a few strains are of clinical origin, insect frass from various trees, cacti, rotted fruit, spoiled fruit juice, soil, tanning liquor, sputum, cecum of a cow, human and animal faeces, various clinical sources (*P. kudriavzevii* = *Candida krusei*), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), a few asexual species of this clade presently remain in *Candida*, but will be transferred to *Pichia* as new combinations, cultures and sequences are available.

Picoa Vittad. 1831, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, three species, type: *P. juniperi* Vittad., asexual morph unknown, saprobes, hypogeous, terrestrial, worldwide, see Guecin et al. (2010; Turkey), Lumbsch and Huhndorf (2010; outline), Sbissi et al. (2010; DNA, phylogeny), Ammarellou et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Pidoplichkoviella Kiril. 1975, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. terricola* Kiril., asexual morph unknown, from soil, terrestrial, British Islesraine, see Lumbsch and Huhndorf

(2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Piedraia Fonseca & Leão 1928, *Piedraiaeae*, *Capnodiales*, *Dothideomycetes*, two species, type: *P. hortae* Fonseca & Leão, asexual morph unknown, human pathogen, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Piggotia Berk. & Broome 1851, *Ascomycota* genera *incertae sedis*, three species, type: *P. astroidea* (Berk.) Berk. & Broome, coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pileomyces K.L. Pang & Jheng 2012, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *P. formosanus* K.L. Pang & J.S. Jheng, asexual morph unknown, saprobes, marine, Taiwan, see Pang and Jheng (2012b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype of type: BBH 30192, BIOTEC Bangkok Herbarium.

Pilgeriella Henn. 1900, *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *P. perisporioides* Henn., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pilidium Kunze 1823 (= *Discohainesia* Nannf. 1932; *Hainesia* Ellis & Sacc. 1884; *Sclerotiopsis* Speg. 1882 fide Johnston et al. 2014b), *Chaetomellaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *P. acerinum* (Alb. & Schwein.) Kunze, coelomycetous, sexual morph formerly named in *Discohainesia*, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2013b, 2015e; new species), Johnston et al. (2014b; nomenclature, new combination), cultures and sequences are available, available specimen and cultures for the type: CBS H-17074; CBS 736.68.

Pilophorus Th. Fr., 1857, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, 38 species, type: *P. robustus* Th. Fr., asexual morph unknown, lichenized, terrestrial, worldwide, see Wang et al. (2010c, 2011a, b; new species, taxonomy), Lim et al. (2010; pyrosequencing), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequence data available.

Pinatubo J.B. Manandhar & Mew 1996, *Ascomycota* genera *incertae sedis*, one species, type: *P. oryzae* J.B. Manandhar & Mew, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: ?CALP [as 'UPLB'] 10284.

Piperivora Siboe, P.M. Kirk & P.F. Cannon 1999, *Ascomycota* genera *incertae sedis*, one species, type: *P. nodosa* Siboe, P.M. Kirk & P.F. Cannon, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 376523.

Piricauda Bubák 1914, *Ascomycota* genera *incertae sedis*, eight species, type: *P. uleana* (Sacc. & P. Syd.) Bubák, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Whitton et al. (2012a; morphology), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Piricaudilium Hol.-Jech. 1988, *Ascomycota* genera *incertae sedis*, two species, type: *P. lobatum* Hol.-Jech., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, available specimen for type: CBS H-4357.

Piricaudiopsis J. Mena & Mercado 1987, *Ascomycota* genera *incertae sedis*, one species, type: *P. elegans* J. Mena & Mercado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Zhang et al. (2014b; transferred five species to new genus *Xiuguozhangia*), cultures and sequences are unavailable.

Pirispora Faurel & Schotter 1966, *Ascomycota* genera *incertae sedis*, one species, type: *P. trisetosa* Faurel & Schotter, coelomycetous, sexual morph unknown, saprobes, terrestrial, Algeria, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pirostomella Sacc. 1914, *Ascomycota* genera *incertae sedis*, two species, type: *P. raimundi* Sacc., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pirottaea Sacc. 1878, *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, 28 species, type: *P. veneta* Sacc. & Speg., asexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline), Chlebicka and Konvalinkova (2010; new species), Kirk et al. (2013; genus accepted), cultures available but lacking for the type.

Pirozynskia Subram. 1972, *Dothideomycetes* genera *incertae sedis*, two species, type: *P. farriae* Subram., sexual morph maurodothina-like, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Asterinaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*), cultures and sequences unavailable.

Pirozynskiella S. Hughes 2007, *Asterinaceae*, *Asterinales*, *Dothidomycetes*, three species, type: *P. solaninum* (Sacc. & P. Syd.) S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2017; new species, phylogeny), cultures and sequences are available.

Pisomyxa Corda 1837, *Eurotiomycetidae* genera *incertae sedis*, one species, type: *P. rhacodiooides* Corda, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available but lacking for the type.

Pisorisporium Réblová & J. Fourn. 2015, *Pisorisporiaceae*, *Pisorisporiales*, *Sordariomycetes*, one species, type: *P. cymbiforme* Réblová & J. Fourn., asexual morph unknown, from submerged branch, aquatic, France, Belgium, see Réblová et al. (2015a; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and ex-type cultures available for the type: PRM 924377; CBS 138884.

Pithomyces Berk. & Broome 1873, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, c. 38 species, type: *P. flavus* Berk. & Broome, hyphomycetous, sexual morph didymella-like, saprobes, human and plant pathogens, terrestrial, worldwide, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Irinyi et al. (2015; DNA barcodes for clinically important species), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Pithosira Petr. 1949, *Ascomycota* genera *incertae sedis*, two species, type: *P. sydowii* Petr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Ecuador, Caribbean, see Seifert et al. (2011; morphology, accepted as a genus in *Venturiaceae*), Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pithya Fuckel 1870, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, five species, type: *P. cupressina* (Batsch) Fuckel, asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Pithyella Boud. 1885, *?Helotiaceae*, *Helotiales*, *Leotiomycetes*, eight species, type: *P. hypnina* (Quél.) Boud., asexual morph unknown, saprobes, on bryophytes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Pittostroma Kowalski & T.N. Sieber 1992, *Ascomycota* genera *incertae sedis*, one species, type: *P. abietinum* Kowalski & T.N. Sieber, coelomycetous, sexual morph unknown, saprobes, terrestrial, Poland, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: ZT.

Placella Syd. 1938, *Ascomycota* genera *incertae sedis*, one species, type: *P. fraseriana* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Placiopsis Beltr. 1858, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 20 species, type: *P. grappae* Beltr. (=*P. cinerascens*), asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Prieto et al. (2010b; 2010c; species delimitation), Kirk et al. (2013; genus accepted), sequences are available for the type.

Placidium A. Massal. 1855, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 28 species, type: *P. michelii* A. Massal., asexual morph unknown, lichenicolous, lichenized, terrestrial, worldwide, see Gueidan et al. (2009; new species), Lumbsch and Huhndorf (2010; outline), Prieto et al. (2012; new species), Kirk et al. (2013; genus accepted), sequences are available for the voucher M. Prieto 702 of the type.

Placoasterella Sacc. ex Theiss. & Syd. 1915, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, four species, type: *P. schweinfurthii* (Henn.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted in *Asterinaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, keys, transferred to *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Placoasterina Toro 1930, *Dothideomycetes* genera *incertae sedis*, three species, type: *P. antioquensis* Toro, asexual morph unknown, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Placocarpus Trevis. 1860, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, five species, type: *P. schaeferi* (Fr.) Breuss, asexual morph unknown, lichenized, terrestrial, worldwide, see Knudsen et al. (2009a; new species, USA), Moon and Aptroot (2009; new species, South Korea), Lumbsch and Huhndorf (2010; outline), Prieto et al. (2010a; new species), Roux and Gueidan (2011; new species, France), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Placocrea Syd. 1939, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *P. pulchella* Syd., asexual morph unknown, saprobes, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Placodiplodia Bubák 1916, *Ascomycota* genera *incertae sedis*, two species, type: *P. copelandii* Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Placodothis Syd. 1928, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. petrakii* Syd., asexual morph unknown, saprobes, terrestrial, West Indies, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Placolecis Trevis 1857, *Catillariaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *P. balaniana* (Fr.) Trevis., asexual morph unknown, lichenized, terrestrial, Asia, Europe, see Lumbsch and Huhndorf (2010; outline), Rizzi et al. (2011; Sardinia), Tufan-Çetin and Sümbül (2011; taxonomy), Kirk et al. (2013; genus accepted), Aptroot and Moon (2014; Korea), Sipman and Raus (2015; Greece), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Placomaronea Räsänen 1944, *Candeliariaceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *P. candelarioides* Räsänen, asexual morph unknown, lichenized, terrestrial, America, Africa, see Westberg et al. (2009; monograph), Lumbsch and Huhndorf (2010; outline), Westberg and Sohrabi (2012; taxonomy), Kirk et al. (2013; genus accepted), Flakus et al. (2015; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Placomelan Cif. 1962, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. dipteridis* Cif., sexual morph dothidasteroma-like, epiphytes, terrestrial, see Hofmann (2009; review, accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2012, 2014c; outline), cultures and sequences are unavailable.

Placonema (Sacc. & D. Sacc.) Petr. 1921, *Ascomycota* genera *incertae sedis*, three species, type: *P. bambusacearum* (Sacc., Syd. & P. Syd.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Placonemina Petr. 1921, *Ascomycota* genera *incertae sedis*, one species, type: *P. dothideooides* (Mont.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Placopsis (Nyl.) Linds. 1866, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, 60 species, type: *P. gelida* (L.) Linds., asexual morph unknown, lichenized, terrestrial, worldwide, see Brackel and Berger (2010; *Polycoccum* host), Galloway (2010, 2013; taxonomy), Singh et al. (2010; ecology), de los Ríos et al. (2011; ecophysiology), Lumbsch and Huhndorf (2010; outline), Lumbsch et al. (2011; taxonomy, new species), Øvstdal and Lewis Smith (2011; new species), Schmull et al. (2011; phylogeny), Raggio et al. (2012; ecophysiology), Kukwa et al. (2012; new records), Gao and Ren (2013; new records), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Schneider et al. (2016; phylogeny), sequences are available.

Placopyrenium Breuss 1987, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 22 species, type: *P. bucekkii* (Nádv. & Servít) Breuss, asexual morph unknown, lichenized, terrestrial, worldwide, see Ertz et al. (2008; Europe), Gueidan et al. (2009; phylogeny), Breuss (2009; description, key), Orange (2009b; partial sequence), Lumbsch and Huhndorf (2010; outline), Aptroot et al. (2010; biodiversity), Prieto et al. (2010a, 2012; genus considered polyphyletic, phylogeny), Breuss (2011; Spain), Miller et al. (2011a, b; USA), Aptroot and Yazici (2012; new species), Kirk et al. (2013; genus accepted), Valadbeigi (2014; Iran), Sipman and Raus (2015; taxonomy), Lücking et al. (2017a; classification), Urbanavichus and Ismailov (2016; new records), sequences are available.

Placosoma Syd. 1924, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *P. nothopanacis* Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Asterinaceae*), Lumbsch

and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hongsanan et al. (2014c; taxonomy, keys, transferred to *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: PDD 1003.

Placosphaerina Maire 1917, *Ascomycota* genera *incertae sedis*, one species, type: *P. myriospora* (Pat.) Maire, coelomycetous, sexual morph unknown, saprobes, terrestrial, Algeria, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Placostromella Petr. 1947, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *P. macrospora* Petr., asexual morph unknown, saprobes, terrestrial, China, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes*), Kirk et al. (2013; genus accepted), Thambugala et al. (2014b; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Placothea Syd. 1931, *Ascomycota* genera *incertae sedis*, one species, type: *P. philippensis* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Placothelium Müll. Arg. 1893, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *P. staurothelioides* Müll. Arg., asexual morph unknown, lichenized, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Placothyrium Bubák 1916, *Ascomycota* genera *incertae sedis*, one species, type: *P. athyrum* Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Placynthiella Elenkin 1909, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, seven species, type: *P. perfurarea* (Nyl.) Gyeln., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Voytsekhovich et al. (2011; photobiont), Joshi et al. (2011b; new records from South Korea), Kirk et al. (2013; genus accepted), Kantvilas et al. (2015; phylogeny), cultures and sequences are available but lacking for the type.

Placynthiopsis Zahlbr. 1932, *Placynthiaceae*, *Peltigerales*, *Lecanoromycetes*, one species, type: *P. africana* Zahlbr., lichenized, terrestrial, central Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al.

(2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Placynthium (Ach.) Gray. 1821, *Placynthiaceae*, *Peltigerales*, *Lecanoromycetes*, 25 species, type: *P. nigrum* (Huds.) Gray, asexual morph unknown, lichenized, terrestrial, worldwide, see Øvstedral et al. (2009; new species), Wedin et al. (2009b; phylogeny), Schmull et al. (2011; phylogeny) Lumbsch and Huhndorf (2010; outline), Urbanavichus et al. (2011; new records), Wirth et al. (2011; checklist), Yatsyna (2011; new record), Aptroot and Iqbal (2012; checklist), Hansen (2012; new records, 2014; taxonomy), Aslan (2013; new records), Kirk et al. (2013; genus accepted), McCarthy and Kantvilas (2014; new species), Miądlikowska et al. (2014a; phylogeny, systematics), Aptroot et al. (2015d; new records), Malíček et al. (2015a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Plagiascoma Réblová & J. Fourn. 2015, *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, one species, type: *P. frondosum* Réblová & J. Fourn., asexual morph hyphomycetous, saprobes, terrestrial, France, see Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2017; taxonomy), holotype and ex-type: PRM 933854, CBS 139031.

Plagiophiale Petr. 1955, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *P. eucarpa* (P. Karst.) Petr., asexual morph unknown, saprobes, North America, Sweden, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Plagiosphaera Petr. 1941, *Sordariomycetes* genera *incertae sedis*, five species, type: *P. moravica* (Petr.) Petr., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Plagiostigme Syd. 1925, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *P. couraliae* Syd., asexual morph unknown, saprobes, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Plagiostigmella Petr. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *P. clypeata* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plagiostoma Fuckel 1870 (= *Diplodina* Westend. 1857; = *Septomyxa* Sacc. 1884; = *Cryptodiaporthe* Petrak 1921 fide Rossman et al. 2015a), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, 50 species, type: *P. euphorbiae* (Fuckel) Fuckel, asexual morph coelomycetous, saprobes, terrestrial, tropical, see Sogonov et al. (2008; new species, phylogeny), Lumbsch and Huhndorf (2010; outline), Mejía et al. (2011; new species, phylogeny), Kirk et al. (2013; genus accepted), Walker et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, available specimen and cultures or the type: CBS H-17115; CBS 817.79.

Plagiostromella Höhn. 1917, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. pleurostoma* Höhn., asexual morph unknown, saprobes, India, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Planistroma A.W. Ramaley 1991, *Planistromellaceae*, *Botryosphaeriales*, *Dothideomycetes*, two species, type: *P. yuccigenum* A.W. Ramaley, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Minnis et al. (2012; treated as a synonym of *Kellermania*), Kirk et al. (2013; genus accepted), Monkai et al. (2013; treated as a distinct genus), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Plasia Sherwood 1981, *Ascomycota* genera *incertae sedis*, one species, type: *P. ramicola* (Cooke & Massee) Sherwood, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Platismatia W.L. Culb. & C.F. Culb. 1968, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, ten species, type: *P. glauca* (L.) W.L. Culb. & C.F. Culb., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Thell et al. (2012; phylogeny, accepted as a genus in *Parmeliaceae*), Kirk et al. (2013; genus accepted), Mitrovic et al. (2014; antimicrobial properties), cultures available for the type: AFTOL-ID 201.

Platychora Petr. 1925, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. ulmi* (J. Schröt.) Petr., asexual morph coelomycetous, lichenized, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013, 2016; accepted as a genus in *Didymellaceae*, coelomycetous), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Platygramme Fée 1874, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 30 species, type: *P. caesiopruinosa* (Fée) Fée, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2014, 2017a; phylogeny, classification), cultures and sequences are available.

Platypeltella Petr. 1929, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, three species, type: *P. smilacis* Petr., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Microthyriaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hongsan et al. (2014c; keys, accepted as a genus in *Asterinaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Asterinaceae*), Wu et al. (2014b; taxonomy, transferred to *Asterinaceae*), cultures and sequences are unavailable, holotype of type: W 257.

Platysporoides (Wehm.) Shoemaker & C.E. Babc., Can. J. Bot. 1992, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, twelve species, type: *P. chartarum* (Fuckel) Shoemaker & C.E. Babc., asexual morph unknown, saprobes, terrestrial, California, Europe and India, See Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015c; taxonomy), cultures and sequences are unavailable.

Platystomum Trevis. 1877, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, c. 20 species, type: *P. compressum* (Pers.) Trevis., asexual morph unknown, saprobes, worldwide, see Kirk et al. (2008; treated as a synonym of *Lophiostoma* Ces. & De Not.), Lumbsch and Huhndorf (2010; outline, treated as *Pleosporales*, genera *incertae sedis*), Hyde et al. (2013; accepted as a genus in *Platystomaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Platystomaceae*), Thambugala et al. (2015b; accepted as a genus in *Lophiostomataceae*), Tibpromma et al. (2017b; new species), cultures and sequences are available.

Platytheclium Staiger 2002, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 25 species, type: *P. grammatis* (Fée) Staiger, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Neuwirth et al. (2017; key), cultures and sequences are available.

Platytrachelon Réblová 2013, *Sordariomycetes* genera *incertae sedis*, one species, type: *P. abietis* (Réblová) Réblová, asexual morph unknown, saprobes, Czech Republic, Ukraine, see Réblová (2013; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b;

outline, phylogeny), cultures and sequences are available, holotype and cultures available for the type: PRM 842974; CBS 125235.

Plectania Fuckel 1870, *Sarcosomataceae*, *Pezizales*, *Pezizomycetes*, c. 20 species, type: *P. melastoma* (Sowerby) Fuckel, asexual morph conoplea-like, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Plectocarpon Fée 1825, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, 37 species, type: *P. pseudosticta* (Fée) Fée, sexual morph with mainly multilocular-lirelliform stromatic ascocarps, lichenicolous (only one lichenized), terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Frisch et al. (2014; phylogeny), Lücking et al. (2017a; classification), sequences are available but lacking for the type.

Plectomyces Thaxt. 1931, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. gracilis* Thaxt., asexual morph unknown, saprobes, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plectonaemella Höhn. 1915, *Ascomycota* genera *incertae sedis*, one species, type: *P. fuckeliana* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plectopeltis Syd. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *P. egenula* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Central America, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plectophomopsis Petr. 1922, *Ascomycota* genera *incertae sedis*, one species, type: *P. rivularis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plectosira Petr. 1929, *Ascomycota* genera *incertae sedis*, one species, type: *P. adeana* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plectosphaera Theiss. 1916, *Clypeophsalosporaceae*, *Amphisphaerales*, *Sordariomycetes*, 38 species epithets in Index Fungorum (2017), type: *P. bersamae* (Lingelsh.) Theiss., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2008; treated as a

synonym of *Phyllachora* Nitschke ex Fuckel), Maharachchikumbura et al. (2015b, 2016b; outline), Giraldo et al. (2017; phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Plectosphaerella Kleb. 1929 (= *Plectosporium* M.E. Palm et al. 1995 *fide* Maharachchikumbura et al. 2015b, 2016b; Réblová et al. 2016c), *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*, 13 species, type: *P. cucumeris* Kleb., asexual morph formerly known in *Plectosporium*, pathogens, nematophagous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Carlucci et al. (2012; new species, phylogeny), Liu et al. (2013a; new species), Kirk et al. (2013; genus accepted), Crous et al. (2015d; new species), Maharachchikumbura et al. (2015b, 2016b outline, phylogeny), Réblová et al. (2016c; nomenclature), cultures and sequences are available but lacking for the type.

Plectronidiopsis Nag Raj 1979, *Ascomycota* genera *incertae sedis*, one species, type: *P. chilensis* (Speg.) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, Chile, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plectronidium Nag Raj 1977, *Ascomycota* genera *incertae sedis*, four species, type: *P. sinense* Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plectrothrix Shear 1902, *Ascomycota* genera *incertae sedis*, one species, type: *P. globosa* Shear, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleiochaeta (Sacc.) S. Hughes 1951, *Pleosporineae*, *Pleosporales*, *Dothideomycetes*, four species, type: *P. setosa* (Kirchn.) S. Hughes, hyphomycetous, sexual morph unknown, pathogens, saprobes, terrestrial, North America, see Garibaldi et al. (2012; pathogens, DNA), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Gur and Frenkel (2015; pathogens, DNA), cultures and sequences are available, cultures are available, holotype: CBS H-17161.

Pleiopatella Rehm 1908, *Ostropomycetidae* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *P. harperi* Rehm, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Pleiotomellina Bat., J.L. Bezerra & H. Maia 1964, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. pernambucensis* Bat., J.L. Bezerra & Cavalc., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Plejobolus (E. Bommer, M. Rousseau & Sacc.) O.E. Erikss. 1967, *Dothideomycetes* genera *incertae sedis*, type: *P. arenarius* (E. Bommer, M. Rousseau & Sacc.) O.E. Erikss., data inadequate, see Kirk et al. (2008; treated as a synonym of *Robergea*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Plenocatenulis Bat. & Cif. 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. magnoliae* Bat. & Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plenodomus Preuss 1851, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 18 species, type: *P. lingam* (Tode: Fr.) Höhn., asexual morph coelomycetous, pathogens, saprobes, terrestrial, worldwide, see de Gruyter et al. (2012; taxonomy, phylogeny, treated *Plenodomus rabenhorstii* Preuss as a nom. Dub. and reduced under *Plenodomus lingam*), Hyde et al. (2013; accepted as a genus in *Leptosphaeriaceae*), Wijayawardene et al. (2014c, 2017; outline), Ariyawansa et al. (2015b; phylogeny), cultures and sequences are available, cultures available for the type: CBS 260.94 (de Gruyter et al. 2013).

Plenophysa Syd. & P. Syd. 1920, *Ascomycota* genera *incertae sedis*, one species, type: *P. mirabilis* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, Sri Lanka, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plenotrichaius Bat. & Valle 1961, *Ascomycota* genera *incertae sedis*, one species, type: *P. swartziae* Bat. & Valle, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plenotrichopsis Bat. 1961, *Ascomycota* genera *incertae sedis*, one species, type: *P. coffeae* Bat., H. Maia & C.T. Vasconc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plenotrichum Syd. 1927, *Ascomycota* genera *incertae sedis*, two species, type: *P. mirabile* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Central

America, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plenozythia Syd. & P. Syd. 1916, *Ascomycota* genera *incertae sedis*, two species, type: *P. euphorbiae* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleochaeta Sacc. & Speg. 1881 emend. Kimbrough & Korf 1963, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, five species, type: *P. lynckii* (Speg.) Speg., sexual and hyphomycetous asexual morphs, biotrophic, obligate plant pathogens, terrestrial, Asia, North and South America, see Lumbsch and Huhndorf (2010; outline), Lee et al. (2011; phylogeny, Korea), Kirk et al. (2013; genus accepted), Takamatsu (2013a, b; phylogeny, phenotypic evolution), sequences are available but lacking for the type, holotype of type: LPS.

Pleocouturea G. Arnaud 1911, *Ascomycota* genera *incertae sedis*, two species, type: *P. castagnei* G. Arnaud, coelomycetous, sexual morph unknown, saprobes, terrestrial, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleocryptospora J. Reid & C. Booth 1969, *Sordariomycetes* genera *incertae sedis*, two species, type: *P. bambusae* (Speg.) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Pleogibberella Sacc. ex Berl. & Voglino 1886, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *P. calamia* (Cooke) Berl. & Voglino, asexual morph unknown, saprobe, saprobes, terrestrial, India, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pleomeliola (Sacc.) Sacc. 1899, *Ascomycota* genera *incertae sedis*, one species, type: *P. fenestrata* (Cooke & Ellis) Sacc., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded from *Meliolaceae*), cultures and sequences are unavailable.

Pleomelogramma Speg. 1909, *Herpotrichiellaceae*, *Chaetothyriales*, *Chaetothyriomycetes*, one species, type: *P. argentinensis* Speg., asexual morph unknown, saprobes, South America, see Lumbsch and Huhndorf (2010;

outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleomerium Speg. 1918, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. fuscoviridescens* (Rehm) Speg., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded from *Meliolaceae* and placed in *Dothideomycetes*), cultures and sequences are unavailable.

Pleomonodictys Hern.-Restr., J. Mena & Gene 2017, *Pleomonodictyaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. descalsii* Hern.-Restr., J. Mena & Gene, hyphomycetous, sexual morph unknown, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny, designate lectotype and epitype), sequences are available.

Pleophragmia Fuckel 1870, *Sporormiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. leporum* Fuckel, asexual morph unknown, saprobes, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Hyde et al. (2013; accepted as *Sporormiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pleospidium Körb. 1855, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, four species, type: *P. flavum* Körb., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Knudsen (2011; new species), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Pleoscutula Vouaux 1913, *Helotiiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *P. arsenii* Vouaux, asexual morph unknown, lichenicolous, terrestrial, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available but lacking for the type.

Pleoseptum A.W. Ramaley & M.E. Barr 1995, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. yuccaesedum* A.W. Ramaley & M.E. Barr, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Phaeosphaeriaceae*), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; excluded from *Phaeosphaeriaceae* and placed in *Pleosporaceae*), Wijayawardene et al. (2014c; accepted as a genus in *Pleosporaceae*), cultures and sequences are unavailable, holotype of the type: BPI 802381.

Pleosphaerellula Naumov & Czerepan. 1952, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *P. cornicola* Naumov & Czerepan., asexual

morph unknown, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as *Pleosporales*), cultures and sequences are unavailable.

Pleosphaeria Speg. 1881, *Sordariomycetes* genera *incertae sedis*, 15 species, type: *P. australis* Speg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pleostigma Kirschst. 1939, *Dothideomycetes* genera *incertae sedis*, type: data inadequate, see Kirk et al. (2008; treated as a synonym of *Teichosporella*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Pleotrichiella Sivan. 1984, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. australiensis* Sivan., asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pleotrichocladium Hern.-Restr., R.F. Castaneda & Gene 2017, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. opacum* (Corda) Hern.-Restr., R.F. Castaneda & Gen, hyphomycetous, sexual morph unknown, saprobes, from soil, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Plesiospora Drechsler 1971, *Ascomycota* genera *incertae sedis*, one species, type: *P. globosa* Drechsler, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleuroascus Massee & E.S. Salmon 1901, *Pseudeurotiaceae*, *Leotiomycetes* families *incertae sedis*, eleven species, type: *P. nicholsonii* Massee & E.S. Salmon, saprobes, from soil, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Crous et al. (2013b; new species), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 345, available specimen for type: CBS H-7657.

Pleurocatena G. Arnaud ex Aramb., Gamundí, W. Gams & G.R.W. Arnold 2007, *Pyxidiophoraceae*, *Pyxidiophorales*, *Laboulbeniomycetes*, three species, hyphomycetous, sexual morph unknown, on nematodes, leaf litter, Europe, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Pleuroceras Riess 1854, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, twelve species, type: *P. ciliatum* Riess, asexual morph unknown, saprobes, terrestrial, north temperate, see Mejía et al. (2008; DNA, phylogeny), Sogonov et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Pleurocytospora Petr. 1923, *Ascomycota* genera *incertae sedis*, three species, type: *P. vestita* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleurodesmospora Samson, W. Gams & H.C. Evans 1979, *Ascomycota* genera *incertae sedis*, one species, type: *P. coccorum* (Petch) Samson, W. Gams & H.C. Evans, hyphomycetous, sexual morph unknown, parasites, on insects (*Aleyrodes*, aphids, leaf hoppers), mites (*Tydeus*, *Paratetranychus*) and fungi (*Meliola*), Africa, Asia, Central America, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleurodiscula Höhn. 1926, *Ascomycota* genera *incertae sedis*, one species, type: *P. neglecta* (Desm.) Höhn, coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleurodomus Petr. 1934, *Ascomycota* genera *incertae sedis*, one species, type: *P. humuli* Petr., coelomycetous, sexual morph unknown, saprobes, on decaying stems of *Humuli lupuli*, Siberia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleuropedium Marvanová & S.H. Iqbal., 1973, *Ascomycota* genera *incertae sedis*, three species, type: *P. tricladoides* Marvanová & S.H. Iqbal, sexual morph unknown, on submerged dead *Juncus effusus* axes and from foam in mountain streams, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: BRNU 445.796

Pleurophoma Höhn. 1914, *Lentitheciaceae*, *Pleosporales*, *Dothideomycetes*, nine species, type: *P. pleurospora* (Sacc.) Höhn., sexual morph unknown, saprobes, terrestrial, UK, see Wijayawardene et al. (2012, 2017; outline), de Gruyter et al. (2012; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Crous et al. (2015d, 2016a; new species, phylogeny), Tibpromma et al.

(2017b; new species), cultures and sequences are available, cultures available for the type: CBS 130329.

Pleurophomopsis Petr. 1924, Ascomycota genera incertae sedis, seven species, type: *P. salicicola* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Pleurophragmium Costantin 1888, *Sordariomycetes* genera incertae sedis, 29 species, type: *P. bicolor* Costantin, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widely distributed, see Réblová (2009; phylogeny), D'Souza and Bhat (2012; key), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015; outline), Dai et al. (2017; morphology, phylogeny), cultures and sequences are available but lacking for type.

Pleuroplaconema Petr. 1923, Ascomycota genera incertae sedis, two species, type: *P. sambuci* Petr., coelomycetous, sexual morph unknown, endophytes, Europe and Asia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleuroplacosphaeria Syd. 1928, Ascomycota genera incertae sedis, one species, type: *P. negeriana* Syd., coelomycetous, sexual morph unknown, saprobes, Chile, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleurosticta Petr. 1931, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *P. lichenicola* Petr., asexual morph unknown, lichenized, terrestrial, Asia, Europe, Africa, see Lumbsch and Huhndorf (2010; outline), Crespo et al. (2011; generic concept), Hawksworth et al. (2011a; checklist), Schmull et al. (2011; sequence), Kirk et al. (2013; genus accepted), Millot et al. (2014; cancer), Favero-Longo et al. (2015; saxicolous community), Papazi et al. (2015; symbiosis), Kaźmierska and Kossowska (2016; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available.

Pleurostoma Tul. & C. Tul. 1863 (= *Pleurostomophora* Vijaykr. et al. 2004 fide Maharachchikumbura et al. (2015b, 2016b; Réblová et al. 2016c), *Pleurostomataceae*, *Calosphaeriales*, *Sordariomycetes*, seven species, type: *P. candollei* Tul. & C. Tul., asexual morph hyphomycetous, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2015b, 2016c; new species, nomenclature), cultures and sequences are available but lacking for the type.

Pleurostromella Petr. 1922, *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, 15 species, type: *P. ulmicola* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Phookamsak and Hyde (2015; accepted as in *Fenestellaceae*), Wanasinghe et al. (2017b; treated *Fenestellaceae* as a synonym of *Cucurbitariaceae*) cultures and sequences are unavailable.

Pleurotheciella Réblová, Seifert & J. Fourn. 2012, *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, nine species, type: *P. rivularia* Réblová, Seifert & J. Fourn., asexual morph hyphomycetous, saprobes, terrestrial, Europe, Canada, see Réblová et al. (2012; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and ex-type cultures of the type: PRM 899852; CBS 125238.

Pleurotheciopsis B. Sutton 1973, Ascomycota genera incertae sedis, seven species, type: *P. pusilla* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, on litter (*Calophyllum*, *Castanea*, *Rubus*) and bank (*Betula*, *Ilex*, *Salix*), Caribbean, Europe, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: IMI 168990.

Pleurothecium Höhn. 1919 (= *Carpoligna* F.A. Fernández & Huhndorf 1999 fide Réblová et al. 2016c), *Pleurotheciaceae*, *Pleurotheciales*, *Sordariomycetes*, nine species, type: *P. recurvatum* (Morgan) Höhn., hyphomycetous, sexual morph formerly named in *Carpoligna*, saprobes, terrestrial, Europe, Canada, see Wu and Zhang (2009; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Réblová et al. (2012, 2016c; taxonomy, phylogeny, nomenclature), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 138686.

Pleurothyriella Petr. & Syd. 1925, Ascomycota genera incertae sedis, one species, type: *P. pinastri* (Oudem.) Petr. & Syd., coelomycetous, sexual morph unknown, saprobes, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pleurovularia R. Kirschner & U. Braun 2002, Ascomycota genera incertae sedis, one species, type: *P. polliniae* (Henn.) R. Kirschner & U. Braun (≡ *Ovularia polliniae* Henn.), hyphomycetous, terrestrial, plant pathogens, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline),

cultures and sequences are unavailable, holotype of type: Yoshinaga 25 (B).

Pliariona A. Massal. 1860, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *P. montagnei* (Bosch) A. Massal., asexual morph unknown, lichenized, terrestrial, tropical, see Rivas Plata et al. (2013; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Plicaria Fuckel 1870, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, ten species, type: *P. trachycarpa* (Curr.) Boud., asexual morph unknown, hypogeous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for the type: DAOM 195830, genus needs revision since it is polyphyletic.

Plicariella (Sacc.) Rehm 1894, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *P. radula* (Berk. & Broome) Rehm, asexual morph unknown, saprobes, see Kirk et al. (2013; did not list), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Plochmopeltis Theiss. 1914, *Schizothyriaceae*, *Dothideomycetes* families *incertae sedis*, five species, type: *P. intricata* (Ellis & G. Martin) Theiss., asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013, accepted as a genus in *Schizothyriaceae*), Wijayawardene et al. (2014c; outline), Phookamsak et al. (2016; taxonomy), cultures and sequences are unavailable.

Ploettnera Henn. 1900, *Calloriaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *P. caeruleoviridis* (Rehm) Henn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ploioderma Darker 1967 (= *Cryocaligula* Minter 1986 *fide* Johnston et al. 2014b), *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, seven species, type: *P. hedgcockii* (Dearn.) Darker, asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Shi et al. (2010; new species, China), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available but lacking for the type.

Plokamidomyces Bat., C.A.A. Costa & Cif. 1957, *Euantennariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *P. colensoi* Bat., C.A.A. Costa & Cif., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Oceania, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plowrightia Sacc. 1883, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, more than 50 species, type: *P. ribesia* (Pers.) Sacc., asexual morph coelomycetous, parasitic or saprobes on leaves, twigs and wood in terrestrial habitats, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014b; phylogeny), Wijayawardene et al. (2014c; outline), sequence data available for two species, holotype of type: cannot be located.

Plurisperma Sivan. 1970, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *P. dalbergiae* Sivan., asexual morph unknown, saprobes, terrestrial, Pakistan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Plurispermopsis Pereira-Carv., Inácio & Dianese 2010, *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. cerradensis* Pereira-Carvalho, Inácio & Dianese, asexual morph unknown, saprobes, terrestrial, Brazil, see Pereira-Carvalho et al. (2010; taxonomy), cultures and sequences are unavailable.

Pneumocystis Delanoë & Delanoë 1912, *Pneumocystidaceae*, *Pneumocystidales*, *Pneumocystidomycetes*, five species, type: *P. carinii* P. Delanoë & Delanoë, asexual reproduction occurs by binary fission, pathogens, on mammals (frequent cause of pneumonia in HIV/AIDS patients), worldwide, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Poaceascoma Phook. & K.D. Hyde 2015, *Lentitheciaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. helicoides* Phook. & K.D. Hyde, asexual morph unknown, saprobes, Asia, see Phookamsak et al. (2015a; taxonomy), Luo et al. (2016b; new species), cultures and sequences are available.

Poaceicola W.J. Li, Camporesi, D.J. Bhat & K.D. Hyde 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *P. arundinis* W.J. Li, Camporesi, D.J. Bhat & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Italy, see Li et al. (2015c; taxonomy, phylogeny), Wijayawardene et al. (2016b; taxonomy, phylogeny), Hyde et al. (2016; taxonomy, phylogeny), Thambugala et al. (2017b, new species, phylogeny), holotype of the type: MFLU 15-0702.

Pochonia Bat. & O.M. Fonseca 1965, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *P. humicola* Bat. & O.M. Fonseca, asexual morph unknown, saprobes, from soil, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Nonaka et al. (2013c; new species, Japan), Huang et al. (2015a; new species, Tibet, China), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type, type specimen of type: CBS H-7885.

Pocillopycnis Dyko & B. Sutton 1979, *Ascomycota* genera *incertae sedis*, one species, type: *P. umensis* (Bubák & Vleugel) Dyko & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Sweden, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pocillum De Not. 1864, *Vibrissaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. cesatii* (Mont.) De Not., asexual morph unknown, saprobes on fruits and leaves, Europe and America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), DNA sequences and cultures unavailable.

Poculinia Spooner 1987, *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. nothofagi* (Rodway) Spooner, asexual morph unknown, saprobes, terrestrial, Tasmania, see Lumbsch and Huhndorf (2010; outline, listed as *Poculina*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Podocapsa Tiegh. 1887, *Ascomycota* genera *incertae sedis*, one species, type: *P. palmata* Tiegh., asexual morph unknown, saprobes, terrestrial, Europe, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pododimeria E. Müll. 1959, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, four species, type: *P. gallica* E. Müll., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Podonectria Petch 1921, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, nine species, type: *P. coccicola* (Ellis & Everh.) Petch, asexual morph unknown, associated with scale insects, widespread in tropical regions, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Tubeufiaceae*), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Podophacidium Niessl 1868, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *P. terrestre* Niessl, asexual morph unknown, saprobes, terrestrial, terrestrial, Europe, North America see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Podoplaconema Petr. 1921, *Ascomycota* genera *incertae sedis*, one species, type: *P. melaenum* (Fr.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2014c; genus accepted, in *Dothideaceae*), Hyde et al. (2013; genus accepted, in *Dothideaceae*), Kirk et al. (2013;

genus accepted), Wijayawardene et al. (2017; listed genus as *Ascomycota*), cultures and sequences are unavailable, isosyntype of type: B (also distributed in Scleromyceti Suecicae no. 200).

Podosordaria Ellis & Holw. 1897 (= *Pedisordaria* Clem. 1909 *fide* Wendt et al. 2017), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *P. mexicana* Ellis & Holw., asexual morph hyphomycetous, saprobes, coprophilous, terrestrial, worldwide, see Hsieh et al. (2010; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; phylogeny), sequences are available for the voucher 176 (WSP) of the type.

Podosphaera Kunze 1823 emend. Braun & Takamatsu 2000, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, 95 species, type: *P. myrtillina* (Schub. : Fr.) Kunze, sexual and hyphomycetous asexual morphs, biotrophic, obligate plant pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Takamatsu et al. (2010; phylogeny), Braun and Cook (2012; taxonomy), Kirk et al. (2013; genus accepted), sequences are available, neotype of type: Fungi Saxon. Exs. 72b. (HAL).

Podospora Ces. 1856, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, c. 100 species, type: *P. fimiseda* (Ces. & De Not.) Niessl, asexual morph consists of dark celled phialides such as *Cladorrhinum*, *Phialophora*, *Rhizoctonia*, *Ramophialophora*, saprobes on dung of animals, worldwide, see Lumbsch and Huhndorf (2010; outline), Chang et al. (2010; phylogeny), Mungai et al. (2012c; from dung), Kirk et al. (2013; genus accepted), Kruys et al. (2015; phylogeny), Melo et al. (2015, taxonomy), about 24 species have sequence data, cultures available for the type: CBS 482.64.

Podosporiella Ellis & Everh. 1895, *Ascomycota* genera *incertae sedis*, four species, type: *P. humilis* Ellis & Everh., hyphomycetous, sexual morph unknown, saprobes, on leaves (*Garrya*, *Faurea*), Africa, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Podosporiopsis Jian Ma, X.G. Zhang & R.F. Castañeda 2016, *Ascomycota* genera *incertae sedis*, two species, type: *P. sinensis* Jian Ma, X.G. Zhang & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobic, terrestrial, China, see Ma et al. (2016b; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Podosporium Schwein. 1832, *Ascomycota* genera *incertae sedis*, 67 species, type: *P. rigidum* Schwein, hyphomycetous, sexual morph unknown, saprobes, on dead stems and litter (*Ampelopsis*, *Bambusa*, *Rhus* and others),

terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Podostictina Clem. 1909, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, six species, type: *P. endochrysoides* (Müll. Arg.) Clem., lichenized, see Moncada et al. (2013a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Podotara Malcolm & Vězda 1996, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *P. pilophoriformis* Malcolm & Vězda, asexual morph unknown, lichenized, terrestrial, New Zealand, see Hsieh et al. (2010; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available, holotype of type: CHR 486638.

Poeltaria Hertel 1984, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, six species, type: *P. turgescens* (Körb.) Hertel, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Fryday and Hertel (2014; new species, Antarctica, Tasmania), Lücking et al. (2017a; classification), cultures and sequences are available for an unidentified species but lacking for the type.

Poeltidea Hertel & Hafellner 1984, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, three species, type: *P. perusta* (Nyl.) Hertel & Hafellner, asexual morph unknown, lichenized, terrestrial, Kerguelen, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Fryday and Hertel (2014; new species, New Zealand), Lücking et al. (2017a; classification), cultures and sequences are unavailable, lectotype of type: BM.

Poeltinula Hafellner 1984, *Rhizocarpaceae*, *Rhizocarpales*, *Lecanoromycetes*, two species, type: *P. cerebrina* (DC.) Hafellner, asexual morph unknown, lichenized, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Poetschia Körb. 1861, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, four species, type: *P. buellioides* Körb., asexual morph unknown, saprobes, terrestrial, Europe, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; taxonomy), cultures and sequences are unavailable.

Poikilosperma Bat. & J.L. Bezerra 1961, *Ascomycota* genera *incertae sedis*, one species, type: *P. scutellare* Bat. & J.L. Bezerra, coelomycetous, sexual morph unknown, saprobes, on leaves of *Brosimum*, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Polaroscyphus Huhtinen 1987, *Hyaloscypheaceae*, *Helotiidales*, *Leotiomycetes*, one species, type: *P. spetsbergianus* Huhtinen, asexual morph unknown, saprobes, terrestrial, Svalbard, Norway, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polistroma Clemente 1807, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *P. fernandezii* Clemente, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lumbsch et al. (2014a, b; taxonomy), Lücking et al. (2017a; classification), sequences and cultures are unavailable.

Polyancora Voglmayr & Yule 2006, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *P. globosa* Voglmayr & Yule, sexual morph unknown, saprobes, submerged leaves, Peninsular Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharanachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and ex-type strains of type: WU 26489; CBS 118182.

Polyandromyces Thaxt. 1920, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. coptosomatis* Thaxt., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences available for the type: KT800020, KT800035.

Polyascomyces Thaxt. 1900, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. trichophyiae* Thaxt., asexual morph unknown, saprobes, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polyblastia A. Massal. 1852, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 120 species, type: *P. cupularis* A. Massal., asexual morph unknown, saprobes, terrestrial, worldwide, see Savić et al. (2008; phylogeny), Lumbsch and Huhndorf (2010; outline), Savić and Tibell (2012; northern Europe, adjacent Arctic), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Polyblastidium Kalb 2015, *Physciaceae*, *Caliciales*, *Lecanoromycetes*, 18 species, type: *P. japonicum* (M. Satô) Kalb, lichenized, worldwide, see Mongkolsuk et al. (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Polybulbophiale Goh & K.D. Hyde 1998, *Ascomycota* genera *incertae sedis*, one species, type: *P. palmicola* Goh & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, on palm petiole (*Licuala*), Brunei, see Seifert

et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: HKU (M) 4717.

Polycauliona Hue 1908, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 25 species, type: *P. regalis* (Vain.) Hue, lichenized, see Arup et al. (2013; phylogeny), Kondratyuk et al. (2014; phylogeny), Jaklitsch et al. (2016a; notes), Lücking et al. (2017a; classification), cultures and sequences are available.

Polycephalomyces Kobayasi 1941, *Ophiocordycepitaceae*, *Hypocreales*, *Sordariomycetes*, twelve species, type: *P. formosus* Kobayasi, hyphomycetous, on insects, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kepler et al. (2013; new species), Kirk et al. (2013; genus accepted), Matočec et al. (2014; revision), Quandt et al. (2014; nomenclature), cultures and sequences are available.

Polychaetella Speg. 1918, *Ascomycota* genera *incertae sedis*, three species, type: *P. schweinitzii* (Berk. & Desm.) Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polychidium (Ach.) Gray 1821, *Massalongiaceae*, *Peltigerales*, *Lecanoromycetes*, c. ten species, type: *P. muscicola* (Sw.) Gray, asexual morph unknown, lichenized, terrestrial, worldwide, see Wedin et al. (2009b; phylogeny), Hansen (2009; Iceland), Lumbsch and Huhndorf (2010; outline), Muggia et al. (2011b; phylogeny, systematics), Anderson and Neily (2012; Nova Scotia), Jayalal et al. (2012a; new records), Kirk et al. (2013; genus accepted), Weerakoon and Aptroot (2016; checklist), Lücking et al. (2017a; classification), sequences are available.

Polycladium Ingold. 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. equiseti* Ingold, sexual morph unknown, found on dead internodes of *Equisetum fluviatile*, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 73532.

Polyclypeolina Bat. & I.H. Lima 1959, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. brideliae* (Hansf.) Bat., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review, accepted as a genus in *Aulographaceae*), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Aulographaceae*), Hyde et al. (2013; review, excluded from *Aulographaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*), cultures and sequences are unavailable, holotype of type: K(M) 177972.

Polycoccum Saut. ex Körb. 1865, *Polycoccaceae*, *Trypetheliales*, *Dothideomycetes*, 66 species, type: *P. sauteri* Körb., asexual morph unknown, saprobes, lichenicolous, terrestrial, worldwide, see Etayo (2010; new species), Brackel and Berger (2010; new species), Lumbsch and Huhndorf (2010; outline), Zhurbenko and Dillman (2010; new species), van den Boom (2012; new species), Hyde et al. (2013; outline, accepted as a genus in *Dacampiaceae*), Halici et al. (2013; new species, Turkey), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Ertz et al. (2015a; phylogeny, introduced *Polycoccaceae*), Hafellner (2015; new species), cultures and sequences are available but lacking for the type.

Polycyclina Theiss. & Syd. 1915, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, one species, type: *P. rhytidmoides* (Speg.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Polycyclinopsis Bat., A.F. Vital & I.H. Lima 1958, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. solani* Bat., A.F. Vital & I.H. Lima, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as a genus in *Dothideomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Polycyclus Höhn. 1909, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, two species, type: *P. andinus* (Pat.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, São Paulo, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, keys), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Polydesmia Boud. 1885, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, seven species, type: *P. pruinosa* (Berk. & Broome) Boud., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Polydesmus Mont. 1845, *Ascomycota* genera *incertae sedis*, 14 species, type: *P. elegans* Durieu & Mont., hyphomycetous, sexual morph unknown, saprobes, terrestrial, American, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Polydiscidium Wakef. 1934, *Leotiomycetes* genera *incertae sedis*, seven species, type: *P. martynii* Wakef., asexual morph unknown, saprobes, Africa, Guyana, worldwide, see Lumbsch and Huhndorf (2010; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polydiscina Syd. 1930, *Leotiomycetes* genera *incertae sedis*, one species, type: *P. boliviiana* Syd., asexual morph unknown, saprobes, terrestrial, Bolivia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polyetron Bat. & Peres 1963, *Ascomycota* genera *incertae sedis*, one species, type: *P. myrtaceicola* Bat. & Peres, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polylobatispora Matsush. 1996, *Ascomycota* genera *incertae sedis*, two species, type: *P. deltoidea* Matsush., coelomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Moro et al. (2015; new species, key), cultures and sequences are unavailable, holotype of the type: Matsushima Fungus Collection, Kobe, 5T195.

Polymeridium (Müll. Arg.) R.C. Harris 1980, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, 50 species, type: *P. contendens* (Nyl.) R.C. Harris, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Aptroot and Cáceres (2014; key), Aptroot and Lücking (2016; monograph), Lücking et al. (2017a; classification), cultures and sequences are available.

Polypedia Bat. & Peres 1959, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *P. leopoldinensis* Bat. & Peres, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Micropeltidaceae*), Hyde et al. (2013; not accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; not accepted as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polyphialoseptoria Quaedvl., R.W. Barreto, Verkley & Crous 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. terminaliae* Quaedvl., R.W. Barreto, Verkley & Crous, coelomycetous, sexual morph unknown, pathogens, terrestrial, Brazil, see Quaedvlieg et al. (2013; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type: CBS H-21298; CBS 135106.

Polyplosphaeria Kaz. Tanaka & K. Hiray. 2009, *Tetraplosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. fusca* Kaz. Tanaka & K. Hiray., asexual morph hyphomycetous, saprobes, terrestrial, Japan,

Thailand, see Tanaka et al. (2009; morphology, phylogeny), Li et al. (2016a; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type: HHUF 29399; JCM 13175.

Polypyrenula D. Hawksw. 1985, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, one species, type: *P. sexlocularis* (Müll. Arg.) D. Hawksw., asexual morph unknown, on bark, doubtfully lichenized, terrestrial, Caribbean, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type: G.

Polyrhizon Theiss., Syd. & P. Syd. 1914, *Dothideomycetes* genera *incertae sedis*, two species, type: *P. terminaliae* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, India, South Africa, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Polyrostrata T.P. Devi & N. Mathur 2009, *Ascomycota* genera *incertae sedis*, two species, type: *P. indica* T.P. Devi & N. Mathur, coelomycetous, sexual morph unknown, pathogens, terrestrial, India, see Mathur et al. (2009; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, holotype of the type: HCIO 46892.

Polyschema H.P. Upadhyay 1966, *Latoruaceae*, *Pleosporales*, *Dothideomycetes*, 21 species, type: *P. terricola* H.P. Upadhyay 1966, hyphomycetous, sexual morph not unknown, saprobes, one species from human clinical sample, terrestrial, cosmopolitan, see Mota et al. (2008; new species, key), Shenoy et al. (2010; monophyly within *Pleosporales*), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2015d; placement in *Latoruaceae*), ex-type of the type: CBS 301.65.

Polyscytalum Riess 1853, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, 13 species, type: *P. fecundissimum* Riess, hyphomycetous, saprobes, pathogens, terrestrial, worldwide, see Cheewangkoon et al. (2009; new species, Portugal), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), Senanayake et al. (2015; phylogeny), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available, available specimen and cultures for the type: CBS H-6049; CBS 100506.

Polysporella Woron. 1916, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. woronowii* Woron., see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list).

Polysporidiella Petr. 1960, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. iranica* Petr., asexual morph unknown, saprobes, terrestrial, Iran, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Polysporina Vězda [nom. cons.] 1978, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, ten species, type: *P. simplex* (Taylor) Vězda, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Polystigma DC. 1815 (= *Polystigmina* Sacc. 1884, *Rhodoseptoria* Naumov 1913 *fide* Maharachchikumbura et al. 2015b, 2016b; Réblová et al. 2016c), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, 25 species, type: *P. rubrum* (Pers.) DC., pathogens, worldwide, see Suzuki et al. (2008; pathogens, Japan), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Habibi and Banihashemi (2016; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Polystomellina Bat. & A.F. Vital 1958, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. didymopanacis* Bat. & A.F. Vital, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; accepted as a genus in *Dothideomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Polystomellomyces Bat. 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. atheniensis* Bat., coelomycetous, sexual morph unknown, saprobes, terrestrial, Greece, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Polystomellopsis F. Stevens 1924 [1923], *Dothideomycetes* genera *incertae sedis*, one species, type: *P. mirabilis* F. Stevens, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Polystratorictus Matsush. 1993, *Ascomycota* genera *incertae sedis*, two species, type: *P. fusarioideus* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Peru, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: Matsushima Fungus Collection, Kobe, OP-353.

Polythrincium Kunze 1817 (= *Cymadothea* F.A. Wolf 1935 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *P. trifolii* Kunze, hyphomycetous, sexual morph formerly known in *Cymadothea*, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline, sexual morph), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; nomenclature), Rossman et al. (2015b; nomenclature), Tibpromma et al. (2017b; phylogeny), cultures and sequences are available.

Polytolypa J.A. Scott & Malloch 1993, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *P. hystricis* J.A. Scott & Malloch, asexual morph unknown, coprophilous, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hirooka et al. (2016; phylogeny), cultures and sequences are available, cultures available for the type: UAMH 7299, holotype of type: TRTC 51349.

Polytretophora Mercado 1983, *Ascomycota* genera *incertae sedis*, one species, type: *P. calcarata* Mercado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pontogeneia Kohlm. 1975, *Koralionastetaceae*, *Koralionastetales*, *Sordariomycetes*, eight species, type: *P. padinae* Kohlm., asexual morph unknown, saprobes, marine, worldwide, Campbell et al. (2009a; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequence data available for one species, holotype of type: J.K. 3541a = 00985934, NYBG.

Pontoporeia Kohlm. 1963, *Halothiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. biturbinata* (Durieu & Mont.) Kohlm., asexual morph coelomycetous, saprobes, marine, Mediterranean Sea, see Jones et al. (2009b, 2015; classification), Suetrong et al. (2009; phylogeny, morphology), Zhang et al. (2012e, 2013c; morphology, phylogeny), Hyde et al. (2013; morphology, phylogeny), Wijayawardene et al. (2014c; phylogeny, nomenclature), cultures and sequences are available.

Populocrescentia Wanás., E.B.G. Jones & K.D. Hyde 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. forlicesenensis* Wanás., Camporesi, E.B.G. Jones & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Europe, Russia, see Ariyawansa et al. (2015a; taxonomy, phylogeny), cultures and sequences are available, holotype and ex-type strains of type: MFLU 15-1504; MFLUCC 14-0651.

Porina Ach. [nom. cons.] 1883, *Porinaceae*, *Ostropales*, *Lecanoromycetes*, 140 species, type: *P. nucula* Ach.,

lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Porobeltraniella Gusmão 2004, *Beltraniaceae*, *Amphisphaerales*, *Sordariomycetes*, one species, type: *P. porosa* (Piroz. & S.D. Patil) Gusmão, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Rajeshkumar et al. (2016; phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available, available specimen and cultures available for the type: IMI 234741; NFCC 3994 (Rajeshkumar et al. 2016).

Porocercospora Amaradasa, Amundsen, Madrid & Crous 2014, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. seminalis* (Ellis & Everh.) Amaradasa, Amundsen, H. Madrid & Crous, hyphomycetous, sexual morph unknown, pathogens, USA, see Amaradasa et al. (2014; morphology, phylogeny), cultures and sequences are available, epitype and ex-epitype culture of type: CBS H-21149, CBS 134906.

Poroocladium Descals., 1976, *Ascomycota* genera *incertae sedis*, one species, type: *P. aquaticum* Descals, hyphomycetous, sexual morph unknown, saprobes, terrestrial, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 199900.

Porocyphus Körb. 1855, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, eight species, type: *P. coccodes* Flot. ex Körb., asexual morph unknown, lichenized, terrestrial, worldwide, see Högnabba et al. (2009a, b; evolution of photobiont), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type.

Poroleprieuria M.C. González, Hanlin, Ulloa & Elv. Aguirre 2004, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. rogersii* M.C. González, Hanlin, Ulloa & Elv. Aguirre, asexual morph hyphomycetous, saprobes, terrestrial, Mexico, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; phylogeny), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable, holotype of the type: MEXU 23019.

Poronia Willd. 1787, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *P. gleditschii* Willd., asexual morph unknown, saprobes, coprophilous,

terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hembrom et al. (2013; new species, India), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; phylogeny), cultures and sequences are available but lacking for the type.

Poropeltis Henn. 1904, *Ascomycota* genera, one species, type: *P. davillae* Henn., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Porophilomyces U. Braun 2000, *Ascomycota* genera *incertae sedis*, one species, type: *P. poricola* (Bonar) U. Braun, hyphomycetous, terrestrial, fungicolous, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype and isotype of type: UC 1272179; UC 568840.

Porophoromyces Thaxt. 1926, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. tmesiphori* Thaxt., asexual morph unknown, biotrophic, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Porosubramaniana Hol.-Jech. 1985, *Ascomycota* genera *incertae sedis*, two species, type: *P. moniliformis* Hol.-Jech., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Porpidia Körb. 1855, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, 50 species, type: *P. trullisata* (Kremp.) Körb., lichenized, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Schmull et al. (2011; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Porpidinia Timdal 2010, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *P. tumidula* (Sm.) Timdal (*Lichen tumidulus* Sm.), lichenized fungi, saxicolous, Europe, North Africa, see Timdal (2010; phylogenetic placement based on anatomical characters), Lücking et al. (2017b; classification), sequences are unavailable, holotype of type: Smith s.n., LINN.

Psora Hoffm. 1796, *Psoraceae*, *Lecanorales*, *Lecanoromycetes*, c. 35 species, type: *P. decipiens* (Hedw.) Hoffm. (*Lichen decipiens* Hedw.), lichenized fungi, saxicolous, terricolous, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Psorinia Gotth. Schneid. 1980, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, c. 15 species, type:

P. conglomerata (Ach.) Gotth. Schneid. (*Lecidea conglomerata* Ach.), lichenized, saxicolous, temperate northern hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Porrectotheca Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *P. divaricata* Matsush., coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: Matsushima Fungus Collection, Kobe, 5H436.

Potamomyces K.D. Hyde 1995, *Ascomycota* genera *incertae sedis*, one species, type: *P. armatisporus* K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: BRIP 21490.

Potridiscus Döbbeler & Triebel 2000, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. polymorphus* Döbbeler & Triebel, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: M Rambold 3269.

Potriphila Döbbeler 1996, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *P. navicularis* Döbbeler, asexual morph unknown, saprobes, terrestrial, Austria, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: M (Döbbeler 6759).

Praelongicaulis E.B.G. Jones, Abdel-Wahab & K.L. Pang 2015, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *P. kandeliae* (Abdel-Wahab & E.B.G. Jones) E.B.G. Jones, Abdel-Wahab & K.L. Pang, asexual morph unknown, saprobes, marine, Hong Kong, see Sakayaroj et al. (2011a; phylogeny), Jones et al. (2015, phylogeny, nomenclature), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and ex-type of the type: IMI 379307; CY 1492.

Pragmopora A. Massal 1855 (= *Pragmopycnis* B. Sutton & A. Funk 1975 *fide* Johnston et al. 2014b), *Tympanidaceae*, *Phacidiales*, *Leotiomycetes*, seven species, type: *P. amphibola* A. Massal, asexual morph formerly known in *Pragmopycnis*, saprobes, common in temperate northern hemisphere, see Kirk et al. (2013; genus accepted), Quijada et al. (2015; phylogeny), lectotype in Royal Botanic Gardens (K), cultures and sequences are unavailable.

Prataprajella Hosag. 1992, *Meliolaceae*, *Meliolales*, *Sordariomycetes*, two species, type: *P. turpinicola*

(Hosag.) Hosag., asexual morph unknown, saprobes, terrestrial, India, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; treated as a doubtful genus in *Meliolaceae*), cultures and sequences are unavailable.

Premilcurensis Tibpromma, Camporesi & K.D. Hyde 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. senencionis* Tibpromma, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, Italy, see Tibpromma et al. (2015; taxonomy), holotype: MFLU 14-0678, cultures and sequences are available.

Preussia Fuckel 1867 (= *Sporormiella* Ellis & Everh. 1892; = *Spororminula* Arx & Aa 1987 *fide* Hyde et al. 2013), *Sporomiaceae*, *Pleosporales*, *Dothideomycetes*, 49 species, type: *P. funiculata* (Preuss) Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Chang and Wang (2009; new species), Kruys and Wedin (2009; new species), Schoch et al. (2009; phylogeny), Asgari and Zare (2010; new species), Lumbsch and Huhndorf (2010; outline), Mungai et al. (2012d; *Sporormiella* from dung), Hyde et al. (2013; accepted as a genus in *Sporomiaceae*, phylogeny), Kirk et al. (2013; genus accepted), Mapperson et al. (2014; endophytes), Wijayawardene et al. (2014c; outline), Kruys (2015; new species, Sweden), cultures and sequences are available, type specimen and cultures available for the type: CBS H-17339; CBS 659.74.

Priceomyces Kurtzman & M. Suzuki 2010, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, nine species, type: *P. haplophilus* (Shifrine & Phaff) M. Suzuki, asexual reproduction is by multilateral budding, saprophytic, in tree fluxes, spoiled wine, alpechin, soil, insect frass in various trees, worldwide, see Kurtzman and Suzuki (2010; taxonomy), Crous et al. (2016b; new species), cultures and sequences are available.

Prillieuxina G. Arnaud 1918 (= *Leprieurina* G. Arnaud 1918 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015a), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, 55 species, type: *P. winteriana* (Pazschke) G. Arnaud, asexual morph unknown, parasymbiont on *Pertusaria*, terrestrial, tropical, see Hosagoudar and Chandraprabha (2009b; new species), Lumbsch and Huhndorf (2010; outline), Hosagoudar (2012; new species, India), Hosagoudar and Divya (2013; new species, India), Hyde et al. (2013; accepted as a genus in *Asterinaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Gautam (2015; new species), Guatimosim et al. (2015; phylogeny), Rossman et al. (2015b; nomenclature), cultures and sequences are available but lacking for the type.

Pringsheimia Schulzer 1866, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, nine species, type: *P. rosarum* Schulzer, asexual morph unknown, parasitic

or saprobes, terrestrial habitats, worldwide, see Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; review), Wijayawardene et al. (2014c; outline), sequence data available for one species, but no ex-type sequence data are available.

Proboscispora Punith. 1984, *Ascomycota* genera *incertae sedis*, one species, type: *P. manihotis* Punith., coelomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Proliferodiscus J.H. Haines & Dumont 1983, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, eight species, type: *P. inspersus* (Berk. & M.A. Curtis) J.H. Haines & Dumont, asexual morph unknown, saprobes, terrestrial, worldwide, see Hofton et al. (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Proliferosphaera T.P. Devi 2012, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. capsici* T.P. Devi, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Devi et al. (2012; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are available, cultures available for the type: PP-2011 isolate 6494, holotype of type: ITCC 6494.

Prolixandromyces R.K. Benj. 1970, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eight species, type: *P. veliae* R.K. Benj., asexual morph unknown, parasitic on Veliidae bugs, semiaquatic, Africa, Asia, Central America, Europe, see Weir (2008; review), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pfliegler et al. (2016b; taxonomy), cultures unavailable, sequences are available but lacking for the type.

Pronectria Clem. 1931, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, 41 species, type: *P. lichenicola* (Ces.) Clem., asexual morph hyphomycetous, parasymbiont to parasites on lichens, more rarely algae, worldwide, see Etayo and Sancho (2008; morphology, keys), López et al. (2009; new species), Brackel (2014; morphology), Maharachchikumbura et al. (2015b, 2016b; outline), sequence data only for the type *P. robergei* as *Illosporium carneum* (ATCC 201590), lectotype of type: PC, Roberge.

Propolidium Sacc. 1884, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, two species, type: *P. glaucum* (Ellis) Sacc., asexual morph unknown, saprobes, terrestrial, Africa, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Propolina Sacc. 1884, *Marthamycetaceae*, *Rhytismatales*, *Leotiomycetes*, one species, type:

P. cervina Sacc., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Propolis (Fr.) Corda 1838, *Marthamycetaceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *P. farinosa* (Pers.) Fr., asexual morph unknown, saprobes, terrestrial, Asia, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Proprioscypha Spooner 1987, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. corticicola* Spooner, asexual morph unknown, saprobes, pathogens, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Prosopidicola Crous & C.L. Lennox 2004, *Prosopidicolaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *P. mexicana* Crous & C.L. Lennox, coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Wijayawardene et al. (2012, 2017; outline), Crous et al. (2016b; new species, phylogeny), cultures are available.

Prosthemium Kunze 1817 (= *Pleomassaria* Speg. 1880 *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b), *Pleomassariaceae*, *Pleosporales*, *Dothideomycetes*, c. seven species, type: *P. betulinum* Kunze, coelomycetous, sexual morph formerly known in *Pleomassaria*, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline, sexual morph), Tanaka et al. (2010; taxonomy, phylogeny), Wijayawardene et al. (2012, 2014c, 2016b; outline, taxonomy, nomenclature), Zhang et al. (2012e; accepted as a genus in *Pleomassariaceae*, phylogeny), Hyde et al. (2013; accepted as a genus in *Pleomassariaceae*), Rossman et al. (2015b; nomenclature), Ariyawansa et al. (2016b; new species), cultures available for the type: CBS 279.74.

Prostratus Sivan., W.H. Hsieh & Chi Y. Chen 1993, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *P. cyclobalanopsidis* Sivan., W.H. Hsieh & Chi Y. Chen, asexual morph unknown, saprobes, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable, holotype of type: IMI 354625.

Protoblastenia (Zahlbr.) J. Steiner 1911, *Psoraceae*, *Lecanorales*, *Lecanoromycetes*, 30 species, type: *P. rupestris* (Scop.) J. Steiner, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures available for the type: Prorup415.

Protocrea Petch 1937, Hypocreaceae, Hypocreales, Sordariomycetes, six species, type: *P. farinosa* (Berk. & Broome) Petch, asexual morph acremonium-like, saprobes, terrestrial, worldwide, see Jaklitsch et al. (2008; new species, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, epitype and cultures available for the type: BPI 747356; CBS 121554.

Protocreopsis Yoshim. Doi 1977, Bionectriaceae, Hypocreales, Sordariomycetes, twelve species, type: *P. zingibericola* Yoshim. Doi, asexual morph acremonium-like, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lechat and Fournier (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Protocucurbitaria Naumov 1951, Sordariomycetes genera incertae sedis, six species, type: *P. ribicola* Naumov, asexual morph unknown, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Protomicarea Hafellner 2001, Pilocarpaceae, Lecanorales, Lecanoromycetes, one species, type: *P. limosa* (Ach.) Hafellner, asexual morph unknown, lichenized, terrestrial, Europe, Turkey, see Lumbsch and Huhndorf (2010; outline), Kinalioglu and Aptroot (2011; new records in Turkey), Schmull et al. (2011; phylogeny), cultures available for the type: Obermayer 07939 (Schmull et al. 2011).

Protomyces Unger 1833, Protomycetaceae, Taphriniales, Taphrinomycetes, ten species, type: *P. macrosporus* Unger, asexual morph unknown, biotrophic on plants, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available, available specimen of type: CBS H-17359.

Protomycopsis Magnus 1905, Protomycetaceae, Taphriniales, Taphrinomycetes, ten species, type: *P. leucanthemi* P. Magnus, asexual morph unknown, biotrophic on plants, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Protopannaria (Gyeln.) P.M. Jørg. & S. Ekman 2000, Pannariaceae, Peltigerales, Lecanoromycetes, five species, type: *P. pezizoides* (Weber) P.M. Jørg. & S. Ekman, lichenized, Southern Hemisphere, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are available.

Protoparmelia M. Choisy 1929, Parmeliaceae, Lecanorales, Lecanoromycetes, 35 species, type: *P. badia* (Hoffm.) Hafellner, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Papong et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Protoparmeliopsis Choisy 1929, Lecanoraceae, Lecanorales, Lecanoromycetes, 20 species, type: *P. muralis* (Schreb.) M. Choisy, lichenized, see Kondratyuk et al. (2015f; taxonomy), Zhao et al. (2016c; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Protoroccella Follmann 2001, Roccellaceae, Arthoniales, Arthoniomycetes, two species, type: *P. minima* (R. Sant.) Follmann ex Follmann, lichenized, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Protoscypha Syd. 1925, Protoscyphaceae, Dothideomycetes families incertae sedis, one species, type: *P. pulla* Syd., asexual morph unknown, biotrophic, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Protostegia Cooke 1880, Mycosphaerellaceae, Capnodiales, Dothideomycetes, one species, type: *P. eucleae* Kalchbr. & Cooke, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Kirk et al. (2013; genus accepted), Crous et al. (2015a; phylogeny, epitypification), Wijayawardene et al. (2017; outline), epitype and ex-epitype of the type: PREM 60879; CPC 23549.

Protostegiomycetes Bat. & A.F. Vital 1955, Ascomycota genera incertae sedis, one species, type: *P. lembosiae* Bat. & A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Protostroma Bat. 1957, Ascomycota genera incertae sedis, one species, type: *P. hyphaenes* Bat., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Protothelenella Räsänen 1943, Protothelenellaceae, Ostropales, Lecanoromycetes, eleven species, type: *P. reducta* (Th. Fr.) Räsänen, lichenized, Asia, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Resl et al. (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Protothyrium G. Arnaud 1917, Parmulariaceae, Dothideomycetes families incertae sedis, two species,

type: *P. salvadorae* (Cooke) G. Arnaud, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, isotype of type: IMI 158625.

Protounguicularia Raity. & R. Galán 1986, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. brevicapitata* Raity. & R. Galán, asexual morph unknown, saprobes, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Protousnea (Motyka) Krog 1976, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, eight species, type: *P. magellanica* (Mont.) Krog, lichenized, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Protoventuria Berl. & Sacc. 1887 (= *Ramalia* Bat. 1957 *fide* Rossman et al. 2016a), *Venturiaceae*, *Venturiales*, *Dothideomycetes*, c. five species, type: *P. rosae* (De Not.) Berl. & Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011d; morphology, phylogeny), Hyde et al. (2013; accepted as in *Venturiaceae*, phylogeny), Wijayawardene et al. (2014c; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are available.

Proxipyricularia Klaubauf, Lebrun & Crous 2014, *Pyriculariaceae*, *Magnaportheales*, *Sordariomycetes*, two species, type: *P. zingiberis* (Y. Nisik.) Klaubauf, Lebrun & Crous, hyphomycetous, sexual morph unknown, pathogens, terrestrial, Japan, see Klaubauf et al. (2014; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Wijayawardene et al. (2017; outline), available cultures for the type: CBS 133594.

Psammina Sacc. & M. Rousseau ex E. Bommer & M. Rousseau 1891, *Ascomycota* genera *incertae sedis*, eight species, type: *P. bommeriae* Sacc. & M. Rousseau, hyphomycetous, sexual morph unknown, saprobes, lichenicolous, terrestrial, Africa, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: CBS H-17363.

Psathyrophlyctis Brusse 1987, *Phlyctidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *P. serpentaria* Brusse, asexual morph unknown, lichenized, terrestrial, South Africa, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Pselaphidomyces Speg. 1917, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. pselapti* Speg. asexual morph unknown, biotrophic, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudaleuria Lusk 1987, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *P. quinaultiana* Lusk, asexual morph unknown, from soil, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Xu et al. (2012; soil fungi), Kirk et al. (2013; genus accepted), Nouhra et al. (2013; DNA, ectomycorrhizal), sequences are available.

Pseudallescheria Negroni & I. Fisch. 1944, *Microascaceae*, *Microascales*, *Sordariomycetes*, nine species, type: *P. shearii* Negroni & I. Fisch., asexual morph unknown, saprobes, human pathogens, terrestrial, worldwide, see Gilgado et al. (2010; new species, human pathogen), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available but lacking for the type.

Pseudarctomia Gyeln. 1939, *Lichinaceae*, *Lichinales*, *Lichinomycetes*, one species, type: *P. microleuca* (Nyl.) Gyeln., asexual morph unknown, lichenized, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Pseudascozonius Brumm. 1985, *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, one species, type: *P. racemosporus* Brumm., asexual morph unknown, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudephebe M. Choisy 1930, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *P. pubescens* (L.) M. Choisy, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Wang and McCune (2010; China), Thell et al. (2012; phylogeny, accepted as a genus in *Parmeliaceae*), Kirk et al. (2013; genus accepted), Mitrovic et al. (2014; antimicrobial properties), Boluda et al. (2016; cryptic species), sequences are available.

Pseudeurotium J.F.H. Beyma 1937, *Pseudeurotiaceae*, *Leotiomycetes* families *incertae sedis*, six species, type: *P. zonatum* J.F.H. Beyma, asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Minnis and Lindner (2013; new species), cultures and sequences are available.

Pseudevernia Zopf 1903, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, four species, type: *P. furfuracea* (L.) Zopf, lichenized, North America, Europe, see Lumbsch

and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Pseudoacremonium Crous 2014, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. sacchari* Crous, hyphomycetous, sexual morph unknown, Vietnam, see Crous et al. (2014c; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), holotype and ex-type CBS H-21704, CPC 2315.

Pseudoacrodixys W.A. Baker & Morgan-Jones 2003, *Ascomycota* genera *incertae sedis*, 14 species, type: *P. eickeri* (Morgan-Jones) W.A. Baker & Morgan-Jones, sexual morph unknown, saprobes, terrestrial, see Castañeda-Ruiz et al. (2010a, 2016b; new species), Zhang et al. (2011g; new species), Dubey and Pandey (2012; new species), Fiua et al. (2014; new species), Ma et al. (2015; new species), two species have sequence data.

Pseudoamauroascus Cano, M. Solé & Guarro 2002, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *P. australiensis* Cano, M. Solé & Guarro, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), sequences are available for the type (unpublished), holotype of type: IMI 370014.

Pseudoanguillospora S.H. Iqbal 1974, *Ascomycota* genera *incertae sedis*, two species, type: *P. stricta* S.H. Iqbal, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudoannulatascus Z.L. Luo, Maharachch. & K.D. Hyde 2015, *Annulatascaeae*, *Annulatascales*, *Sordariomycetes*, one species, type: *P. batriisporus* (K.D. Hyde) Z.L. Luo, Maharachch. & K.D. Hyde, asexual morph unknown, aquatic, Australia, Thailand, see Luo et al. (2015b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: BRIP 21481, cultures and sequences are available.

Pseudoanthostomella Daranag., Camporesi & K.D. Hyde 2016, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. pini-nigrae* Daranag., Camporesi & K.D. Hyde, asexual morph unknown, terrestrial, saprobes, Italy, see Daranagama et al. (2016d; taxonomy), Wendt et al. (2017; placed in *Xylariales*), cultures and sequences are available.

Pseudoaristastoma Suj. Singh 1979, *Ascomycota* genera *incertae sedis*, one species, type: *P. indicum* Suj. Singh, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudoascochyta Valenzuela-Lopez, Stchigel, Cano-Canals, Guarro & Cano 2016, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. pratensis* Valenzuela-Lopez, Cano-Canals, Stchigel, Guarro & Cano, coelomycetous, sexual morph unknown, saprobes, terrestrial, Spain, see Crous et al. (2016a; taxonomy, phylogeny), holotype and ex-type of the type: CBS H-22735, FMR 14524.

Pseudoascotaiwania J. Yang, Bhat & K.D. Hyde 2016 *Fuscosporellaceae*, *Fuscosporellales*, *Sordariomycetes*, one species, type: *P. persoonii* (Fallah, J.L. Crane & Shearer) J. Yang, Bhat & K.D. Hyde, hyphomycetous, Manitoba, see Yang et al. (2016c; taxonomy, phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available.

Pseudoasperisporium U. Braun 2000, *Ascomycota* genera *incertae sedis*, three species, type: *P. tupae* (Speg.) U. Braun, hyphomycetous, terrestrial, plant pathogens, leaf-spotting, Asia, South America, see Seifert (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of type: LPS 959.

Pseudoasteromassaria M. Matsum. & Kaz. Tanaka 2015, *Latoruaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. fagi* M. Matsum. & Kaz. Tanaka, asexual morph coelomycetous (stagonospora-like), saprobes, terrestrial, Japan, see Ariyawansa et al. (2015a; morphology, phylogeny), Tibpromma et al. (2017b; new data), cultures and sequences are available, holotype and ex-type of the type: HHUF 30471; MAFF 245221.

Pseudoastrospphaeriella Phook., Z.L. Luo & K.D. Hyde 2015, *Pseudoastrospphaeriellaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. thailandensis* Phook., Z.L. Luo & K.D. Hyde, asexual morph coelomycetous, saprobes, Asia, see Phookamsak et al. (2015b; taxonomy, phylogeny), cultures and sequences are available.

Pseudoaurantiascoma Thambug. & K.D. Hyde 2015, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. kenyense* (Mugambi & Huhndorf) Thambugala & K.D. Hyde, asexual morph unknown, saprobes, cosmopolitan, see Thambugala et al. (2015b; taxonomy), Jaklitsch et al. (2016c; treated as a synonym of *Teichospora* but in here we do not agree with them), cultures and sequences are available.

Pseudobaeomyces M. Satô 1940, *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type: *P. insignis* (Zahlbr.) M. Satô, asexual morph unknown, lichenized, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudobasidiospora Dyko & B. Sutton 1978, *Ascomycota* genera *incertae sedis*, one species, type:

P. caroliniana Dyko & B. Sutton, coelomycetous, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sexual morph unknown, cultures and sequences are unavailable, holotype of type: IMI 208458.

Pseudobeltrania Henn. 1902, *Beltraniaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *P. cedrelae* Henn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2015e; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Rajeshkumar et al. (2016; phylogeny), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available but lacking for the type.

Pseudocalopadina Lücking 1999, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *P. mira* Lücking, lichenized, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type: M.

Pseudocamarosporium Wijyaw. & K.D. Hyde 2014, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, ten species, type: *P. propinquum* (Sacc.) Wijyaw., Camporesi & K.D. Hyde, coelomycetous, sexual morph unknown, has camarosporium-like and coniothyrium-like morphologies, saprobes, Europe, see Wijayawardene et al. (2014c, d, 2016b, c; morphology, phylogeny), Crous et al. (2015c; new species, phylogeny, coniothyrium-like species), Thambugala et al. (2017a, phylogeny, new species), sequence data available, holotype and ex-type strains of type: MFLU 14-0090; MFLUCC 14-0192 = ICMP 20203.

Pseudocanalisporium R.F. Castañeda & W.B. Kendr. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *P. circumfecundum* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudocatenomycopsis Crous & L.A. Shuttlew. 2013, *Coronophorales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. rothmanniae* Crous & L.A. Shuttlew., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Zambia, see Crous et al. (2013b; morphology, phylogeny), Maharachchikumbura et al. (2015b, 2016b; phylogeny, outline), holotype and ex-type of the type: CBS H- 21432; CBS 22733.

Pseudocenangium P. Karst. 1886, *Ascomycota* genera *incertae sedis*, one species, type: *P. pinastri* P. Karst., coelomycetous, sexual morph unknown, saprobes,

terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudocercospora Speg. 1910, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, c. 1500 accepted species (Mycobank), type: *P. vitis* (Lév.) Speg., hyphomycetous, sexual morph mycosphaerella-like, plant pathogens, endophytes or saprobes, cosmopolitan, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Phengsintham et al. (2013; species in Laos), Crous et al. (2012a; phylogeny), Bakhshi et al. (2014; phylogeny), Shivas et al. (2015; new species), sequences are available, epitype of type: CBS HAL (MBT 102230).

Pseudocercosporaella Deighton 1973, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 150 species, type: *P. ipomoeae* Sawada ex Deighton, hyphomycetous, sexual morph mycosphaerella-like, plant pathogens, terrestrial, five species confirmed based on DNA, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Frank et al. (2010; morphology, phylogeny), Rossman et al. (2016b; new combination), epitype and ex-epitype of type: CBS H-20409; CBS 125685.

Pseudocercosporidium Deighton 1973, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *P. venezuelanum* (Syd.) Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Pseudochapsa Parnmen, Lücking & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 17 species, type: *P. dilatata* (Müll. Arg.) Parnmen, Lücking & Lumbsch, asexual morph unknown, lichenized, terrestrial, tropical, see Parnmen et al. (2012; genus proposed), cultures and sequences are available.

Pseudochuppia Kamal, A.N. Rai & Morgan-Jones 1984, *Ascomycota* genera *incertae sedis*, one species, type: *P. sarcinispore* Kamal, A.N. Rai & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudociboria Kanouse 1944, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. umbrina* Kanouse, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudoclathrosphaerina Voglmayr 1997, Ascomycota genera *incertae sedis*, one species, type: *P. evamariae* Voglmayr, hyphomycetous, sexual morph unknown, saprobes, from submerged deciduous leaves, USA, New Zealand, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable (unpublished), holotype of the type: WU 17015.

Pseudocoleodictyospora Doilom & K.D. Hyde 2016, *Pseudocoleodictyosporaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *P. tectonae* Doilom, D.J. Bhat & K.D. Hyde, saprobes, terrestrial, see Doilom et al. (2017; phylogeny), cultures and sequences are available, holotype and ex-holotype type strains of type: MFLU 15-3527; MFLUCC 12-0385, MKT 058, ICMP 21160.

Pseudocoleophoma Kaz. Tanaka & K. Hiray. 2015, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. calamagrostidis* Kaz. Tanaka & K. Hiray., asexual morph coelomycetous, saprobes, terrestrial, Japan, see Tanaka et al. (2015; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of the type: HHUF 30450; CBS 139700.

Pseudoconium Petr. 1969, Ascomycota genera *incertae sedis*, one species, type: *P. pallescens* (Bäumler) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available (unpublished).

Pseudocosmospora C. Herrera & P. Chaverri 2013, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *P. eutypellae* C. Herrera & P. Chaverri, asexual morph hyphomycetous (acremonium-like to verticillium-like), saprobes, terrestrial, worldwide, see Herrera et al. (2013; taxonomy, phylogeny), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype and ex-type cultures of the type: BPI 884165; CBS 133966 = A.R. 4562.

Pseudocryptoclypeus A. Hashim. & Kaz. Tanaka 2017, *Lophiotremataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. yakushimensis* A. Hashim. & Kaz. Tanaka, asexual morph coelomycetous, saprobes, Asia, see Hashimoto et al. (2017; taxonomy, phylogeny), sequences are available.

Pseudocryptospora J. Reid & C. Booth 1969, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. polylepidis* (E. Müll.) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al.

(2015b, 2016b; outline), cultures and sequences are unavailable.

Pseudocyphellaria Vain. [nom. cons.] 1890 (= *Crocodia* Link 1833; = *Cyanisticta* Gyeln. 1931; = *Diphaeosticta* Clem. 1909; = *Diphanosticta* Clem. 1909; = *Merostictina* Clem. 1909); = *Parmosticta* Nyl. 1875; = *Parmostictina* Nyl. 1875; = *Phaeosticta* Trevis. 1869; *Phanosticta* Clem. 1909; *Podostictina* Clem. 1909; = *Saccardoa* Trevis. 1869; = *Stictina* Nyl. 1860), *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, 100 species, type: *Ps. curvata* (Ach.) Vain., lichenized, South America, Asia, see Lumbsch and Huhndorf (2010; outline), Quilhot et al. (2010; taxonomy), Miller et al. (2011a, b; taxonomy), Pandit and Sharma (2012; new record), Fritis et al. (2013; biological activity), Kirk et al. (2013; genus accepted), Moncada et al. (2014b; taxonomy, phylogeny), Yoshiaki and Ohmura (2014; transplant experiment), Beese et al. (2015; ecology), Lücking et al. (2017a; classification), Messuti et al. (2016; sequence), sequences are available but lacking for type.

Pseudocytoplacosphaeria Punith. & Spooner 2002, Ascomycota genera *incertae sedis*, one species, type: *P. conicola* Punith. & Spooner, coelomycetous, sexual morph unknown, saprobes, British Isles, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: K(M) 85582.

Pseudocytospora Petr. 1923, Ascomycota genera *incertae sedis*, one species, type: *P. allantospora* Petr., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudodichomera Höhn. 1918, Ascomycota genera *incertae sedis*, three species, type: *P. varia* (Pers.) Höhn., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudodictyosporium Matsush. 1971, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *P. wauense* Matsush., hyphomycetous, sexual morph unknown, saprobes, Asia, South America, see Menkis and Vasaitis (2011; DNA), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Kirschner et al. (2013; new species, Taiwan), Hyde et al. (2016; new species, Thailand), Boonmee et al. (2016a; phylogeny), cultures and sequences are available, type specimen and cultures available for the type: CBS H-5642; KRP88-6 (Menkis and Vasaitis 2011).

Pseudodidymaria U. Braun 1993, Ascomycota genera *incertae sedis*, three species, type: *P. wyethiae* (Ellis & Everh.) U. Braun, hyphomycetous, sexual morph unknown,

terrestrial, foliicolous, leaf-spotting, Asia, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, lectotype of type: NY 1087025.

Pseudodidymella C.Z. Wei, Y. Harada & Katum. 1997 (= *Pycnopleiospora* C.Z. Wei et al. *fide* Wijayawardene et al. 2014c; Rossman et al. 2015b), *Dothideomycetes* genera *incertae sedis*, one species, type: *P. fagi* C.Z. Wei, Y. Harada & Katum., asexual morph formerly known in *Pycnopleiospora*, saprobes, Japan, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2012, 2014c, 2017; outline, nomenclature), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), cultures and sequences are unavailable, holotype of type: HHUF 22903.

Pseudodidymosphaeria Thambug. & K.D. Hyde 2015, *Massarinaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. spartii* Thambug., Camporesi & K.D. Hyde, asexual morph coelomycetous, saprobes, Europe, see Pavlic et al. (2015a; morphology, phylogeny), Li et al. (2016a; new species), Tibpromma et al. (2017b; new data), sequence data available, reference specimen and ex-type of type: MFLU 14-0578; MFLUCC 13-0273.

Pseudodinemasperium A. Hashim. & Kaz. Tanaka 2015, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *P. fabiformis* A. Hashim., G. Sato & Kaz. Tanaka, coelomycetous, sexual morph unknown, saprobes, Japan, Malaysia, see Hashimoto et al. (2015b; morphology, phylogeny), Crous et al. (2015c; phylogeny), holotype and ex-type of the type: HHUF 29716; MAFF 244361.

Pseudodiplococcum Hern.-Restr., J. Mena & Gene 2017, *Xenospadicoidaceae*, *Xenospadicoidales*, *Sordariomycetes*, one species, type: *X. ibericum* Hern.-Restr., J. Mena & Guarro, hyphomycetous, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Pseudodiplodia (P. Karst.) Sacc. 1884, *Ascomycota* genera *incertae sedis*, 45 species, type: *P. lignaria* (P. Karst.) Sacc., coelomycetous, sexual morph unknown, saprobes, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), culture and sequences are available for an unidentified species.

Pseudodiscula Laubert 1911, *Ascomycota* genera *incertae sedis*, two species, type: *P. endogenospora* Laubert, coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudodiscus Arx & E. Müll. 1959, *Saccardiaceae*, *Ascomycota* families *incertae sedis*, one species, type: *P. nootkatensis* (Terrier) Arx & E. Müll., asexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, available specimen of type: CBS H-17396.

Pseudoecteinomyces W. Rossi 1977, *Euceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *P. zuphiicola* (Speg.) W. Rossi, asexual morph unknown, biotrophic, Africa, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudoepicoccum M.B. Ellis 1971, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *P. cocos* (F. Stevens) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudofuscophialis Sivan. & H.S. Chang 1995, *Ascomycota* genera *incertae sedis*, one species, type: *P. lignicola* Sivan. & H.S. Chang, hyphomycetous, sexual morph unknown, saprobes, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 356633.

Pseudofusicoccum Mohali, Slippers & M.J. Wingf. 2006, *Pseudofusicoccumaceae*, *Botryosphaeraiales*, *Dothideomycetes*, seven species, type: *P. stromaticum* (Mohali, Slippers & M.J. Wingf.) Mohali, Slippers & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, endophytes, terrestrial, cosmopolitan, see Pavlic et al. (2008; new species), Phillips et al. (2008, 2013; taxonomy, phylogeny), Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c, 2017; outline), Doilom et al. (2015; species in *Tectona*), Dissanayake et al. (2016b; taxonomy), Yang et al. (2017; phylogeny), holotype and ex-type of the type: PREM 58237, CMW 13366.

Pseudogaster Höhn. 1907, *Ascomycota* genera *incertae sedis*, one species, type: *P. singularis* Höhn., hyphomycetous, sexual morph unknown, saprobes, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudogibellula Samson & H.C. Evans 1973, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one

species, type: *P. formicarum* (Mains) Samson & H.C. Evans, hyphomycetous, sexual morph unknown, on insects, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Pseudogliophragma Phadke & V.G. Rao 1980, *Wiesneriomycetales*, *Tubeufiales*, *Dothideomycetes*, one species, type: *P. indicum* [as India] Phadke & V.G. Rao, hyphomycetous, sexual morph unknown, saprobes, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Pratibha et al. (2015; phylogeny, typification), cultures and sequences are available, ex-type of the type: MTCC 11985.

Pseudographiella E.F. Morris 1966, *Ascomycota* genera *incertae sedis*, one species, type: *P. variiseptata* E.F. Morris., hyphomycetous, sexual morph unknown, saprobes, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudographis Nyl. 1855, *Triblidiaeae*, *Triblidiales*, *Leotiomycetes*, three species, type: *P. elatina* (Ach.) Nyl., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, available specimen of type: CBS H-17399.

Pseudogymnoascus Raillo 1929, *Myxotrichaceae*, *Leotiomycetes* families *incertae sedis*, ten species, type: *P. roseus* Raillo, asexual morph unknown, animal pathogens, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Minnis and Lindner (2013; new species), Hoyt et al. (2015; control disease), cultures and sequences are available.

Pseudohalonectria Minoura & T. Muroi 1978, *Pseudohalonectriaceae*, *Magnaporthales*, *Sordariomycetes*, eleven species, type: *P. lignicola* Minoura & T. Muroi, asexual morph unknown, saprobes, terrestrial, aquatic, worldwide, see Lumbsch and Huhndorf (2010; outline), Hu et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Perera et al. (2016c; new species, phylogeny), type specimen and cultures available for the type: CBS H-19515; SMH2440 IFRDCC 3110.

Pseudohelicomyces Garnica & E. Valenz. 2000, *Ascomycota* genera *incertae sedis*, one species, type: *P. albus* Garnica & E. Valenz., hyphomycetous, sexual morph unknown, saprobes, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus

accepted), cultures and sequences are unavailable, holotype of type: AH 19575.

Pseudohelotium Fuckel 1870, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, c. 50 species, type: *P. pineti* (Batsch) Fuckel, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudohendersonia Crous & M.E. Palm 1999, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. proteae* Crous & M.E. Palm, coelomycetous, sexual morph unknown, saprobes, associated with leaf spots, USA, Italy, see Wijayawardene et al. (2016b; morphology, phylogeny), cultures and sequences are available, holotype of type: BPI 1107849.

Pseudohepatica P.M. Jørg. 1993, *Ascomycota* genera *incertae sedis*, one species, type: *P. pachyderma* P.M. Jørg., asexual morph unknown, lichenized, Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: US, Steyermark 93298.

Pseudoheppia Zahlbr. 1903, *Heppiaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *P. schuleri* Zahlbr., asexual morph unknown, lichenized, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudohypocrea Yoshim. Doi 1972, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. citrinella* (Ellis) Yoshim. Doi, asexual morph acremonium-like, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pseudoidriella Crous & R.G. Shivas 2011, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. syzygii* Crous & R.G. Shivas, hyphomycetous, sexual morph unknown, saprobe, associated with insect damaged leaves, terrestrial, Australia, see Crous et al. (2011c; morphology, DNA, phylogeny), Wijayawardene et al. (2012, 2017; outline), holotype and ex-type culture of type: CBS H-20758; CBS 131307.

Pseudolachnea Ranoj. 1910, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, five species, type: *P. bubakii* Ranoj., coelomycetous, sexual morph unknown, terrestrial, saprobes, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hashimoto et al. (2015b; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Pseudolachnella Teng. 1936, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, 18 species, type: *P. scolecospora* (Teng & C.I. Chen) Teng, coelomycetous, sexual morph unknown, saprobes, worldwide, see Hashimoto et al. (2015b; morphology, phylogeny), Li et al. (2016a; new species), cultures and sequences are available, holotype of the type: “Ling 138”.

Pseudolachnum Velen. 1934, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. lateritium* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudolecanactis Zahlbr. 1907, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. filicicola* Zahlbr., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ertz et al. (2015b, c; phylogenetic study), Lücking et al. (2017a; classification), sequences are unavailable.

Pseudolembosia Theiss. 1913, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *P. geographica* (Massee) Theiss., asexual morph unknown, saprobes, terrestrial, Australia, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of type: NY.

Pseudoleptogium Müll.Arg. 1885, *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, one species, type: *P. diffractum* (Kremp.) Müll. Arg., lichenized, see Otálora et al. (2013, 2014; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Pseudolignincola Chatmala & E.B.G. Jones 2006, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *P. siamensis* Chatmala & E.B.G. Jones, asexual morph unknown, on wood, marine, tropical, see Sakayaroj et al. (2011a; phylogeny), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Pseudolophiostoma Thambug., Kaz. Tanaka & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. vitigenum* (Kaz. Tanaka & Y. Harada) Thambug., Kaz. Tanaka & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Japan, see Thambugala et al. (2015b; morphology, phylogeny), cultures and sequences are available, holotype and ex-type of type: HHUF 26930; MAFF 239459.

Pseudolophiotrema A. Hashim. & Kaz. Tanaka 2017, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *P. elymicola* A. Hashim. & Kaz. Tanaka,

aseexual morph unknown, saprobes, Europe, see Hashimoto et al. (2017; taxonomy, phylogeny), sequences are available.

Pseudomassaria Jacz. 1894, *Pseudomassariaceae*, *Amphisphaeriales*, *Sordariomycetes*, c. 20 species, type: *P. chondrospora* (Ces.) Jacz., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Jaklitsch and Voglmayr (2012; phylogeny), Senanayake et al. (2015; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for type: MFLUCC 15-0545.

Pseudomassariosphaeria Phukhams., Ariyaw., Campor. & K.D. Hyde 2015, *Amniculicolaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. bromicola* Phukhams., Ariyaw., Camporesi & K.D. Hyde, asexual morph unknown, saprobes, aquatic and terrestrial, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type culture of type: MFLU 15–1403, MFLUCC 15–0031.

Pseudombrophila Boud. 1885, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, 28 species, type: *P. pedrottii* Boud., asexual morph unknown, coprophilous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Pseudomeliola Speg. 1889, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *P. brasiliensis* Speg., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pseudomeria G.L. Barron 1980, *Ascomycota* genera *incertae sedis*, one species, type: *P. mucosa* G.L. Barron, hyphomycetous, sexual morph unknown, on nematodes, terrestrial, Canada, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Pseudomicrodochium B. Sutton 1975, *Ascomycota* genera *incertae sedis*, nine species, type: *P. aciculare* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Réblová et al. (2013; compare with *Cyphellophora*), cultures and sequences are unavailable, holotype of type: IMI 169711.

Pseudomisturatosphaeria Thambug. & K.D. Hyde 2015, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. cruciformis* (Mugambi & Huhndorf)

Thambugala & K.D. Hyde, asexual morph unknown, saprobes, cosmopolitan, see Thambugala et al. (2015b; taxonomy), Jaklitsch et al. (2016c; treated as a synonym of *Teichospora* but in here we do not agree with them), cultures and sequences are available.

Pseudomitrula Gamundí 1980, *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *P. horakii* Gamundí, asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudomonodictys Doilom, Ariyaw., D.J. Bhat & K.D. Hyde 2015, *Macrodiplodiopsidaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. tectonae* Doilom, Ariyaw., D.J. Bhat & K.D. Hyde, saprobes, terrestrial, see Ariyawansa et al. (2015a; phylogeny), Doilom et al. (2017; phylogeny), cultures and sequences are available, holotype and ex-holotype type strains of type: MFLU 15-1413; MFLUCC 12-0552.

Pseudomorfea Punith. 1981, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. coffeae* Punith., asexual morph unknown, saprobes, terrestrial, India, Argentina, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pseudonaevia Dennis & Spooner 1993, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *P. caricina* Dennis & Spooner, asexual morph unknown, saprobes, terrestrial, British Isles, see Raatviir (2008; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: K.

Pseudonectria Seaver 1909, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, c. 10 species, type: *P. buxi* (DC.) Seifert, Gräfenhan & Schroers., hyphomycetous, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures available for type: AFTOL-ID 191, AR2716, CBS 114049, CBS 122566, CBS 114049, AR 2714.

Pseudonectriella Petr. 1959, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. ahmadii* Petr., asexual morph unknown, saprobes, terrestrial, South Africa, see Kirk et al. (2008; treated as a synonym of *Catabotrys* Theiss. & Syd.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; not listed), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Pseudoneottiospora Faurel & Schotter 1965, *Ascomycota* genera *incertae sedis*, two species, type: *P. cunicularia* Faurel & Schotter, coelomycetous, sexual morph unknown, saprobes, Algeria, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Pseudoneurospora Dania García, Stchigel & Guarro 2004, *Sordariaceae*, *Sordariales*, *Sordariomycetes*, two species, type: *P. amorphoporcata* (Udagawa) Dania García, Stchigel & Guarro, asexual morph unknown, saprobes, from soil, terrestrial, Thailand, Spain, see Lumbsch and Huhndorf (2010; outline), Nygren et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available but lacking for the type.

Pseudoniptera Velen. 1947, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, 25 species, type: *P. quercina* Velen., asexual name unknown, saprobes, terrestrial, ?Europe, see Kirk et al. (2008; treated as a synonym of *Hymenoscyphus* Gray), Lumbsch and Huhndorf (2010; outline, accepted as in *Dermateaceae*), Jaklitsch et al. (2016a; classification, placed in *Helotiaceae*), cultures and sequences are unavailable.

Pseudonitschka Coppins & S.Y. Kondr. 1995, *Dacampiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. parmotrematis* Coppins & S.Y. Kondr., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes*), Hyde et al. (2013; outline, accepted as a genus in *Dacampiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of the type: BM, Alston 5836.

Pseudopannaria (B.de Lesd.) Zahlbr. 1924, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *P. marci* (B. de Lesd.) Zahlbr., lichenized, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Pseudoparmelia Lynge 1914, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 15 species, type: *P. cyphellata* Lynge, lichenized, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Buaruang et al. (2015; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Pseudoparodia Theiss. & Syd. 1917, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *P. pseudopeziza* (Pat.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in

Patellariaceae), cultures and sequences are unavailable, holotype of type: FH 7735.

Pseudoparodiella F. Stevens 1927, *Venturiaceae*, *Venturiales*, *Dothideomycetes*, one species, type: *P. vernoniae* F. Stevens, asexual morph unknown, saprobes, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011d; morphology, phylogeny), Hyde et al. (2013; accepted as in *Venturiaceae*, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pseudopassalora Crous 2011, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *P. gouriqua* Crous, hyphomycetous, sexual morph unknown, saprobes, South Africa, see Crous et al. (2011d; morphology, phylogeny), Wijayawardene et al. (2014c, 2017; outline), epitype and ex-epitype culture of type: CBS H-20682, CBS 101954.

Pseudopatellina Höhn. 1908, *Ascomycota* genera *incertae sedis*, one species, type: *P. conigena* (Niessl) Höhn., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, type specimen of type: CBS H-17414.

Pseudopaulia M. Schultz 2002, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *P. tessellata* M. Schultz, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type: B (Schultz 14042a).

Pseudopeltis L. Holm & K. Holm 1978, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. filicum* L. Holm & K. Holm, asexual morph unknown, saprobes, terrestrial, Sweden, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudopeltistroma Katum. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *P. kyusyuense* Katum., coelomycetous, sexual morph unknown, saprobes, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudopeltula Henssen 1995, *Gloeoheppiaceae*, *Lichinales*, *Lichenomycetes*, four species, type: *P. myriocarpa* Henssen, lichenized, Mexico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type: COLO.

Pseudopenidiella Crous & Koukol 2012, *Ascomycota* genera *incertae sedis*, one species, type: *P. piceae* Crous & Koukol, hyphomycetous, saprobes, Czech Republic, see Crous et al. (2012e; morphology; phylogeny), Wijayawardene et al. (2012, 2017; outline), cultures and

sequences are available, holotype of the type: CBS H-20964, ex-type strain of the type: CPC 19969.

Pseudoperitheca Elenkin 1922, *Ascomycota* genera *incertae sedis*, one species, type: *P. murmanica* Elenkin, asexual morph unknown, saprobes, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudopestalotiopsis Maharanachch., K.D. Hyde & Crous 2014, *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, seven species, type: *P. theae* (Sawada) Maharanachch., K.D. Hyde & Crous, coelomycetous, sexual morph unknown, saprobes or pathogens on leaves, China, India, Indonesia, Malaysia, Thailand, see Maharanachchikumbura et al. (2014b, 2016a, b; morphology, phylogeny), Wijayawardene et al. (2016b; outline, morphology), all species have sequence data, epitype and ex-epitype: MFLU 12-0116, MFLUCC 12-0055.

Pseudopetrakia M.B. Ellis 1971, *Ascomycota* genera *incertae sedis*, two species, type: *P. kambakkamensis* (Subram.) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, Asia, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudopezicula Korf 1986, *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *P. tetraspora* Korf, Pearson & Zhuang, asexual morph *Phialophora*-like, pathogen of *Vitaceae*, Europe and North America, see Kirk et al. (2013; genus accepted), holotype of the type: CUP-061765, cultures and sequences are unavailable.

Pseudopeziza Fuckel 1870, *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *P. trifoli* (Biv.) Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available (unpublished) but lacking for the type.

Pseudophacidium P. Karst. 1885, *Phaciaceae*, *Phacidiiales*, *Leotiomycetes*, five species, type: *P. ledii* (Alb. & Schwein.) P. Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Ascodichaenaceae*), Crous et al. (2014b; phylogeny), cultures and sequences are available for the type: CBS 377.59.

Pseudophaeomoniella Nigro, Antelmi & Crous 2015, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, one species, type: *P. oleae* Nigro, Antelmi & Crous, coelomycetous, sexual morph unknown, on living leaves, pathogens, terrestrial, Italy, see Crous et al. (2015d; morphology, phylogeny), cultures and sequences are

available, holotype and ex-type culture of type: CBS H-22250, FV84 = CBS 139191.

Pseudophaeosphaeria Jayasiri, Camporesi & K.D. Hyde 2016, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. rubi* Jayasiri, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Hyde et al. (2016; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 15-1400, MFLUCC 14-0259.

Pseudophloeospora Crous & R.G. Shivas 2010, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *P. syzygii* Crous & M.J. Wingf., coelomycetous, associated with leaf spots, terrestrial, Australia, see Crous et al. (2010c, 2016b; new species, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H 20493, CPC 17051.

Pseudophloeosporella U. Braun 1993, *Ascomycota* genera *incertae sedis*, one species, type: *P. dioscoreae* (Miyabe & S. Ito) U. Braun, coelomycetous, sexual morph unknown, terrestrial, plant pathogens, leaf-spotting, Asia, see Hong et al. (2010; Korea), Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: Miyabe & Hanzawa 9 (SAPA).

Pseudophragmotrichum W.P. Wu, B. Sutton & Gange 1998, *Ascomycota* genera *incertae sedis*, one species, type: *P. cubense* W.P. Wu, B. Sutton & Gange, coelomycetous, sexual morph unknown, saprobes, Cuba, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 372456.

Pseudopithomyces Ariyaw. & K.D. Hyde 2015, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *P. chartarum* (Berk. & M.A. Curtis) J.F. Li, Ariyaw. & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, from human, cosmopolitan, see Ariyawansa et al. (2015a; taxonomy, phylogeny), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available.

Pseudopithyella Seaver 1928, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *P. minuscula* (Boud. & Torrend) Seaver, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Romero et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), cultures available for the type: AFTOL-ID 69 (unpublished).

Pseudoplagiostoma Cheew., M.J. Wingf. & Crous 2010, *Pseudoplagiostomataceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *P. eucaalypti* Cheew., M.J. Wingf. & Crous, asexual morph coelomycetous, saprobes, terrestrial, tropical, see Cheewangkoon et al. (2010; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), holotype

and ex-type cultures of the type: CBS H-20303, CPC 13341=CBS 124807.

Pseudoplatystomum Thambug. & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. scabridisporum* (Abdel-Wahab & E.B.G. Jones) Thambug. & K.D. Hyde, asexual morph unknown, saprobes, on marine driftwood, Australia, see Thambugala et al. (2015b; taxonomy), cultures and sequences are available, holotype of type: IMI 379750.

Pseudoplectania Fuckel 1870, *Sarcosomataceae*, *Pezizales*, *Pezizomycetes*, ten species, type: *P. nigrella* (Pers.) Fuckel, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Romero et al. (2012; phylogeny), Holec and Kříž (2013; morphology), Carbone (2013; taxonomy), Kirk et al. (2013; genus accepted), Glejdura et al. (2015; new species), cultures available for the type: KL BK-4914, neotype: KL BK-4914.

Pseudopleospora Petr. 1920, *Dothideomycetes* genera *incertae sedis*, three species, type: *P. ruthenica* Petr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pseudopolystigmina Murashk. 1928, *Ascomycota* genera *incertae sedis*, two species, type: *P. spiraeicola* Murashk., coelomycetous, sexual morph unknown, saprobes, Russia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudoproboscispora Punith. 1999, *Annulatascaeae*, *Annulatascales*, *Sordariomycetes*, two species, type: *P. aquatica* (S.W. Wong & K.D. Hyde) Punith., asexual morph unknown, saprobes, aquatic, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype of type: CC63 (HKU(M) 2792).

Pseudopyrenidium Nav.-Ros., Zhurb. & Cl. Roux 2010, *Adelococcaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *P. tartaricola* (Linds.) Nav.-Ros., Zhurb. & Cl. Roux, coelomycetous, sexual morph unknown, lichenicolous, Asia, USA, see Navarro-Rosinés et al. (2010; taxonomy), cultures and sequences are unavailable.

Pseudopyrenula Müll. Arg. 1883, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, 20 species, type: *P. diluta* (Fée) Müll. Arg., asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Aptroot and Lücking (2016; monograph), Lücking et al. (2017a; classification), cultures and sequences are available.

Pseudopyricularia Klaubauf, Lebrun & Crous 2014, Pyriculariaceae, Magnaportheales, Sordariomycetes, one species, type: *P. kyllingae* Klaubauf, Lebrun & Crous, sexual morph unknown, pathogens, terrestrial, worldwide, see Klaubauf et al. (2014; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), holotype and ex-type cultures for the type: CBS H-21840, CBS 133595.

Pseudoramichloridium Cheew. & Crous 2009, Extremaceae, Capnodiales, Dothideomycetes, one species, type: *P. henryi* Cheew. & Crous, hyphomycetous, sexual morph unknown, saprobes, pathogens, Australia, see Cheewangkoon et al. (2010; taxonomy, phylogeny), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2014c, 2017; outline), holotype and ex-type cultures for the type: CBS H-20293, CPC 13121 = CBS 124775.

Pseudosplanchnonema Chethana & K.D. Hyde 2015, Massarinaceae, Pleosporales, Dothideomycetes, one species, type: *P. phorcioides* (I. Miyake) Chethana, Camporesi & K.D. Hyde, asexual morph coelomycetous, saprobes, Italy, see Chethana et al. (2015; taxonomy), holotype: MFLU 14-0929, cultures and sequences are available.

Pseudoramonia Kantvilas & Vězda 2000, Graphidaceae, Ostropales, Lecanoromycetes, three species, type: *P. stipitata* (Vězda & Hertel) Kantvilas & Vězda, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; key), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Pseudoramularia Matsush. 1983, Ascomycota genera incertae sedis, two species, type: *P. ponapensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, Uganda, Pacific islands, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudorbilia Y. Zhang bis, Z.F. Yu, Baral & K.Q. Zhang 2007, Orbiliaceae, Orbiliales, Orbiliomycetes, one species, type: *P. bipolaris* Y. Zhang bis, Z.F. Yu, Baral & K.Q. Zhang, saprobes, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable, holotype of type: XBL17 (M H.B. 8310).

Pseudorhizina Jacz. 1913, Discinaceae, Pezizales, Pezizomycetes, three species, type: *P. korshinskii* Jacz., asexual morph unknown, saprobes, terrestrial, North America, see Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Pseudorhizopogon Kobayasi 1983, Ascomycota genera incertae sedis, one species, type: *P. sudaе* Kobayasi, coelomycetous, sexual morph unknown, saprobes, Japan,

see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudorhynchia Höhn. 1909, Niessliaceae, Hypocreales, Sordariomycetes, one species, type: *P. polyrrhyncha* (Penz. & Sacc.) Höhn., asexual morph unknown, saprobes, terrestrial, see Kirk et al. (2008; treated as a synonym of *Trichosphaeria* Fuckel), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; not listed), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Pseudorhytisma Juel 1894, Rhytismataceae, Rhytismatales, Leotiomycetes, one species, type: *P. bistortae* (DC.) Juel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, type specimen of type: CBS H-17420.

Pseudorobillarda M. Morelet 1968, Dothideomycetes genera incertae sedis, twelve species, type: *P. phragmitis* (Cunnell) M. Morelet, coelomycetous, sexual morph unknown, saprobes, marine, terrestrial, worldwide, see Suetrong et al. (2009; phylogeny, morphology), Rungindamai et al. (2012; morphology, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Moreno-Pérez et al. (2014; antagonist), Tangthirasunun et al. (2014b; new species, Thailand), Crous et al. (2015a; phylogeny), Jones et al. (2015; outline), available specimen and cultures for the type: CBS 842.84, CBS H-17422.

Pseudosaccharomyces Briosi & Farneti 1906, Ascomycota genera incertae sedis, 20 species, type: *P. citri* Briosi & Farneti, hyphomycetous, sexual morph unknown, saprobes, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudosagedia (Müll. Arg.) Choisy 1949, Porinaceae, Ostropales, Lecanoromycetes, 80 species, type: *P. cerasi* (Schrad.) M. Choisy, lichenized, see Lücking et al. (2017a; classification), cultures and sequences are available.

Pseudosarcosoma M. Carbone, Agnello & P. Alvarado 2013, Chorioactidaceae, Pezizales, Pezizomycetes, two species, type: *P. latahense* (Paden & Tylutki) M. Carbone, Agnello & P. Alvarado, asexual morph unknown, saprobes, terrestrial, USA, see Carbone et al. (2013; taxonomy), sequences are available of the voucher TUR-A 195801 of the type:, holotype of the type: WTU-F-33051.

Pseudoschismatomma Ertz & Tehler 2014, Roccellaceae, Arthoniales, Arthoniomycetes, one species, type: *P. rufescens* (Pers.) Ertz & Tehler, sexual morph with lirelliform ascomata, lichenized, terrestrial, Europe, Macaronesia, North America, Asia, Australia, see Ertz

et al. (2014b; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Pseudoschizothyra Punith. 1980, *Ascomycota* genera *incertae sedis*, one species, type: *P. thaungii* Punith., coelomycetous, sexual morph unknown, saprobes, Myanmar, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudoscopulariopsis Sandoval-Denis, Gené & Guarro 2015, *Microascaceae*, *Microascales*, *Sordariomycetes*, one species, type: *P. schumacheri* (E.C. Hansen) M. Sandoval-Denis, J. Gené & J. Guarro, asexual morph hyphomycetous, saprobes, terrestrial, North America, see Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Pseudoscypha Duby 1861, *Hysteriaceae*, *Histeriales*, *Dothideomycetes*, three species, type: *P. abietis* J. Reid & Piroz., asexual unknown, terrestrial, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Pseudoseptoria Speg. 1911, *Saccotheciaceae*, *Dothideales*, *Dothideomycetes*, eight species, type: *P. donacicola* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2013; accepted as a genus in *Dothioraceae*, phylogeny), Thambugala et al. (2015c; phylogeny), cultures and sequences are available but lacking for the type.

Pseudosigmoidea K. Ando & N. Nakam. 2000, *Ascomycota* genera *incertae sedis*, one species, type: *P. cranei* K. Ando & N. Nakam., hyphomycetous, sexual morph unknown, saprobes, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Diene et al. (2013; new species), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Pseudospiromastix Guarro, Stchigel & Cano 2014, *Spiromastigaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *P. tentaculata* (Guarro, Gené & De Vroey) Guarro, asexual morph unknown, from soil, terrestrial, Somalia, see Rizzo et al. (2014; morphology; phylogeny), Hirooka et al. (2016; phylogeny), holotype and ex-type strains of type: IMI 351264; RV 24167 = CBS 184.92 = FMR 3842.

Pseudosporidesmium K.D. Hyde & McKenzie 2016, *Xylariomycetidae* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. knawiae* (Crous) K.D. Hyde & McKenzie, hyphomycetous, sexual morph unknown, saprobes, KwaZulu-Natal, see Su et al. (2016b; phylogeny), Wijayawardene et al. (2017; outline),

holotype of the type: CBS H-20158 (as *Sporidesmium knawiae* Crous).

Pseudostegia Bubák 1906, *Ascomycota* genera *incertae sedis*, one species, type: *P. nubilosa* Bubák, coelomycetous, sexual morph unknown, saprobes, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudostigmidium Etayo 2008, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *P. nephromiarium* (Linds.) Etayo, asexual morph unknown, lichenicolous, South America, see Etayo and Sancho (2008; morphology, keys), Hyde et al. (2013; accepted as a genus in *Mycosphaerellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of the type: E, J.D. Hooker.

Pseudostrickeria Q. Tian, Wanás., Camporesi & K.D. Hyde 2015, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *P. muriformis* Wanás., Q. Tian, Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy, see Tian et al. (2015; taxonomy), cultures and sequences are available for type, holotype and ex-type strains of type: MFLU 15-1407; MFLUCC 13-0764 = KUMCC15-0062.

Pseudosydowia Thambug. & K.D. Hyde 2014, *Saccotheciaceae*, *Dothideales*, *Dothideomycetes*, one species, type: *P. eucalypti* (Verwoerd & du Plessis) Thambug. & K.D. Hyde, asexual morphs unknown, saprobes, terrestrial, South Africa, see Thambugala et al. (2015c; phylogeny), cultures and sequences are available, holotype of type: PREM 46423.

Pseudotaeniolina J.L. Crane & Schokn. 1986, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *P. convolvuli* (Esfand.) J.L. Crane & Schokn., hyphomycetous, sexual morph unknown, on litter, rock-inhabiting, cosmopolitan, see Egidi et al. (2014; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Pseudoteratosphaeria Quaedv. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, six species, type: *P. perpendicularis* (Crous & M.J. Wingf.) Quaedv. & Crous, asexual morph unknown, foliicolous, pathogens, saprobes, Brazil, Colombia, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), cultures and sequences are available, holotype and ex-type culture of type: CBS H-19691, CBS 118367.

Pseudotetraploa Kaz. Tanaka & K. Hiray. 2009, *Tetraplosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *P. curviappendiculata* (Sat. Hatak., Kaz. Tanaka & Y. Harada) Kaz. Tanaka & K. Hiray., hyphomycetous, saprobes, terrestrial, Indonesia, Japan, see Tanaka et al. (2009; phylogeny), Seifert et al. (2011;

morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, holotype and ex-type of the type: HHUF 28582, JCM 12852.

Pseudothelomma M. Prieto & Wedin 2016, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, two species, type: *P. ocellatum* (Körb.) M. Prieto and Wedin, lichenized, see Prieto and Wedin (2016; genus proposed, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, voucher specimen of type: Tehler 8063 (S).

Pseudothiella Petr. 1928 (= *Pseudothiopsis* Petr. 1928 *fide* Réblová et al. 2016c), *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, six species, type: *P. hirtellae* (Henn.) Petr., asexual morph formerly named in *Pseudothiopsis*, from living leaves, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures and sequences are unavailable.

Pseudothysis Theiss. & Syd. 1914, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. machaerii* (Rehm) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pseudothyrium Höhn. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *P. polygonati* (Tassi) Höhn., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudotopeliopsis Parnmen, Lücking & Lumbsch 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *P. laceratula* (Müll. Arg.) Parnmen, Lücking & Lumbsch, asexual morph unknown, lichenized, terrestrial, tropical, see Parnmen et al. (2012; genus proposed, phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available, lectotype of type: Sayer s.n. – G (Australia).

Pseudotorula Subram. 1958, *Ascomycota* genera *incertae sedis*, three species, type: *P. heterospora* Subram., hyphomycetous, sexual morph unknown, saprobes, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudotracylla B. Sutton & Hodges 1976, *Ascomycota* genera *incertae sedis*, three species, type: *P. dentata* B. Sutton & Hodges, coelomycetous, sexual morph unknown, saprobes, Brazil, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudotrichia Kirschst. 1939, *Pleomassariaceae*, *Pleosporales*, *Dothideomycetes*, eight species, type: *P. stromatophila* Kirschst., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Melanommataceae*), Hyde et al. (2013; accepted as a genus in *Melanommataceae*), Wijayawardene et al. (2014c; outline), Liu et al. (2015a; new species, Thailand), Tian et al. (2015; phylogeny, accepted as a genus in *Pleomassariaceae*), cultures and sequences are available.

Pseudotrichococonis W.A. Baker & Morgan-Jones 2001, *Ascomycota* genera *incertae sedis*, one species, type: *P. echinophila* (C. Massal.) W.A. Baker & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudotripoconidium Z.F. Yu & K.Q. Zhang 2011, *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, one species, type: *P. sinense* Z.F. Yu & K.Q. Zhang, saprobes, terrestrial, China, see Yu et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: YMF 1.01843.

Pseudotrochila Höhn. 1917, *Rhytismatales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *P. rhododendri* (Racib.) Höhn., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudotryblidium Rehm 1890, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. neesii* (Flot.) Rehm, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pseudotulostoma O.K. Mill. & T.W. Henkel 2001, *Elaphomycetaceae*, *Eurotiales*, *Eurotiomycetes*, two species, type: *P. volvatum* O.K. Mill. & T.W. Henkel, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Smith et al. (2011; DNA), cultures and sequences are available.

Pseudovalsaria Spooner 1986, *Boliniaceae*, *Boliniales*, *Sordariomycetes*, four species, type: *P. foedans* (P. Karst.) Spooner, asexual morphs unknown, saprobes, terrestrial, Europe, China, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Clypeosphaeriaceae*), Untereiner et al. (2013; accepted as a genus in *Boliniaceae*, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available but lacking for the type.

Pseudovalsella Höhn. 1918, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type:

P. thelebola (Fr.) Höhn., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Pseudoveronaea Crous & Batzer 2012, *Dissoconiaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *P. ellipsoidea* Batzer & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Li et al. (2012a; morphology; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available, holotype and ex-type of the type: CBS H-20927, CBS 132086.

Pseudovirgaria H.D. Shin, U. Braun, Arzanlou & Crous 2007, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *P. hyperparasitica* H.D. Shin, U. Braun, Arzanlou & Crous, asexual morph hyphomycetous, fungicolous, Asia, Europe, see Braun et al. (2011; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype and ex-type strains of type: HAL 2054 F; CBS 121739–121741.

Pseudoxylomyces Kaz. Tanaka & K. Hiray. 2015, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *P. elegans* (Goh, W.H. Ho, K.D. Hyde & K.M. Tsui) Kaz. Tanaka & K. Hiray., sexual morph unknown, saprobes, aquatic, worldwide, see Tanaka et al. (2015; morphology, phylogeny), cultures and sequences are available, holotype of the type: HKU (M) 2777.

Pseudoyuconia Lar. N. Vassiljeva 1983, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. thalictri* (G. Winter) Lar. N. Vassiljeva, asexual morph unknown, saprobe, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; accepted as a genus in *Pleosporaceae*), Wijayawardene et al. (2014c; outline), Ariyawansa et al. (2015c; accepted as a genus in *Pleosporaceae*), cultures and sequences are unavailable.

Pseudozythia Höhn. 1902, *Ascomycota* genera *incertae sedis*, one species, type: *P. pusilla* Höhn., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Psilachnum Höhn. 1926, *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, 28 species, type: *P. lateritioalbum* (P. Karst.) Höhn., asexual morph unknown, saprobes, terrestrial, worldwide, see Han et al. (2009, 2014; new species, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Psilocistella Svrček 1977, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, ten species, type: *P. obsoleta* (Velen.) Svrček, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Psiloglonium Höhn. 1918, *Hysteriaceae*, *Hysteriales*, *Dothideomycetes*, c. 15 species, type: *P. lineare* (Fr.) Petr., asexual morph unknown, saprobes, pathogens, terrestrial, cosmopolitan, see Kirk et al. (2008; treated as a synonym of *Glonium* Muhl. ex Fr.), Boehm et al. (2009a, b; overview of genus), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Hysteriaceae*), Wijayawardene et al. (2014c; outline), Li et al. (2016a; new species), Rossman et al. (2016b; new combination, pathogen), sequences are available.

Psilolechia A. Massal. 1860, *Psilolechiaceae*, *Lecanorales*, *Lecanoromycetes*, four species, type: *P. lucida* (Ach.) M. Choisy, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Psiloparmelia Hale 1989, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 13 species, type: *P. distincta* (Nyl.) Hale, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Psilopezia Berk. 1847, *Rhizinaceae*, *Pezizales*, *Pezizomycetes*, five species, type: *P. nummularia* Berk., asexual morph unknown, pathogens, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Pezizales*), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Psilophana Syd. 1939, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. andina* Syd., asexual morph unknown, saprobes, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Psilosphaeria Cooke 1879, *Ascomycota* genera *incertae sedis*, one species, type: need typification, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Psilothecium Clem. 1903, *Leotiomycetes* genera *incertae sedis*, one species, type: *P. incurvum* Clem., asexual morph unknown, saprobes, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Helotiales*), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Psora Hoffm. [nom. cons.] 1796, *Psoraceae*, *Lecanorales*, *Lecanoromycetes*, 35 species, type: *L. decipiens* (Hedw.) Ach., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Psoroglaena Müll. Arg. 1891, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 17 species, type: *P. cubensis* Müll. Arg., lichenized, see Muggia et al. (2010a; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), McCarthy and Kantvilas (2013a; new species), Yatsyna (2014; new records), Himelbrant et al. (2015; new records), Käffer et al. (2015; new records), Kubiak et al. (2016; biodiversity), Lücking et al. (2017a; classification), Weerakoon and Aptroot (2016; new records), cultures and sequences are available.

Psorama Michaux 1803, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, 70 species, type: *P. hypnorum* (Vahl) Gray, lichenized, worldwide, see Ekman et al. (2014; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Psoromaria Nyl. ex Nyl., *Pannariaceae*, *Peltigerales*, *Lecanoromycetes* (= *Degelia* P.M.Jørg. [nom. illeg.]), two species, type: *P. subdescendens* (Nyl.) Nyl., see Jørgensen and Andersen (2015; nomenclature), Lücking et al. (2017b; classification).

Psoromella Gyeln. 1940, *Lecanorales* genera *incertae sedis*, *Lecanoromycetes*, one species, type: *P. pampana* Gyeln., lichenized, see Lumbsch and Huhndorf (2010; outline), Thell et al. (2012; overview), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Psoromidium Stirt. 1877, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *P. wellingtonii* Stirt., lichenized, southern hemisphere, see Lumbsch and Huhndorf (2010; outline), Rubio et al. (2013; taxonomy), Jørgensen and Andersen (2015; genus re-evaluation, sequence), Lücking et al. (2017a; classification), cultures and sequences are available.

Psoronactis Ertz & Tehler 2015, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *P. dilleniana* (Ach.) Ertz & Tehler, sexual morph with apothecioid ascocarps, asexual morph coelomycetous, lichenized, terrestrial, Euro-Siberian, see Ertz et al. (2015b, c; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Psorophorus Elvebakk & S.G. Hong 2010, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *P. pholidotus* Elvebakk & S.G. Hong, asexual morph unknown, lichenized, terrestrial, South America, see Elvebakk et al. (2010; new genus, sequence), sequences are available.

Psorotheciopsis Rehm 1900, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, seven species, type: *P. decipiens* Rehm, lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Psorotichia A. Massal. 1855, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, 50 species, type: *P. murorum* A. Massal., lichenized, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Psorula Gotth. Schneid. 1980, *Psoraceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *P. rufonigra* (Tuck.) Gotth. Schneid., lichenized, North America, Europe, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Pteridosperma J.C. Krug & Jeng 1979, *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *P. foveolatum* (Udagawa & Y. Horie) J.C. Krug & Jeng, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pteridospora Penz. & Sacc. 1897, *Astrospheariellaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *P. javanica* Penz. & Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as *Dothideomycetes*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014b, 2015b; phylogeny, epitypifying), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes*), available cultures for the type: MFLUCC 11-0195.

Pteromyces E. Bommer, M. Rousseau & Sacc. 1906, *Helotiaceae* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. ambiguus* E. Bommer, M. Rousseau & Sacc. ex Sacc., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pteromycula P.F. Cannon 1997, *Ascomycota* genera *incertae sedis*, one species, type: *P. clypeiformis* P.F. Cannon, asexual morph unknown, saprobes, terrestrial, Great Britain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: IMI 362154a.

Pterosporidium W.H. Ho & K.D. Hyde 1996, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *P. rhizophorae* (Vizioli) W.H. Ho & K.D. Hyde, from living leaves, worldwide, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pterulopsis Wakef. & Hansf. 1943, *Ascomycota* genera *incertae sedis*, one species, type: *P. dummeri* Wakef. & Hansf., hyphomycetous, sexual morph unknown, saprobes, Uganda, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pterygiopsis Vain. 1890, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, 17 species, type: *P. atra* Vain., lichenized, North America, South America, Europe, Australia, Africa, see Eichler et al. (2010; list), Schultz and Moon (2011; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Pterygosporopsis P.M. Kirk 1983, *Ascomycota* genera *incertae sedis*, two species, type: *P. fragilis* P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ptychographa Nyl. 1874, *Xylographaceae*, *Trapeliales*, *Lecanoromycetes*, one species, type: *P. xylographoides* Nyl., lichenized, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Spribille et al. (2014a; sequence), Resl et al. (2015; sequence, partial sequence), Lücking et al. (2017a; classification), cultures and sequences are available.

Pubigera Baral, Gminder & Svrček 2014 (= *Pubigera* Baral, Gminder & Svrček 1995; Nom. inval., Art. 40.6 (Melbourne) fide Species Fungorum 2017), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *P. subvillosula* (Rehm) Baral, Gminder & Svrček, sexual morph unknown, saprobes, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Pucciniospora Speg. 1886, *Ascomycota* genera *incertae sedis*, one species, type: *P. chusqueae* Speg., coelomycetous, sexual morph unknown, saprobes, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pulchromyces Hennebert 1973, *Ascomycota* genera *incertae sedis*, one species, type: *P. fimicola* (V. Dring) Hennebert, coelomycetous, sexual morph unknown, coprophilous, Ghana, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available, cultures available for the type: ATCC 18595 (unpublished).

Pulmosphaeria Joanne E. Taylor, K.D. Hyde & E.B.G. Jones 1996, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. archontophoenicus* Joanne E. Taylor, K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Daranagama et al. (2016d; accepted in *Xylariales*), cultures and sequences are unavailable.

Pulvinaria Bonord. 1851, *Sordariomycetes* genera *incertae sedis*, one species, type: *P. bohemica* Velen., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pulvinella A.W. Ramaley 2001, *Ascomycota* genera *incertae sedis*, one species, type: *P. nolinae* A.W. Ramaley, hyphomycetous, sexual morph unknown, coprophilous, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: UC 1512945.

Pulvinodecton Henssen & G. Thor 1998, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *P. kurzii* (Kremp.) Henssen & G. Thor, lichenized, Seychelles, Asia, Australia, see Jaklitsch et al. (2016a; phylogeny), Lücking et al. (2017a; classification), sequences are unavailable.

Pulvinula Boud. 1885, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, 28 species, type: *P. convexella* (P. Karst.) Pfister, asexual morph unknown, coprophilous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), sequences are available, voucher specimen of the type: KH.03.19 (FH).

Pumilus Viala & Marsais 1934, *Sordariomycetes* genera *incertae sedis*, one species, type: *P. medullae* Viala & Marsais, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Punctelia Krog 1982, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 45 species, type: *P. borreri* (Sm.) Krog, lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Punctillina Toro 1934, *Ascomycota* genera *incertae sedis*, one species, type: *P. solteroii* Toro, coelomycetous, sexual morph unknown, saprobes, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Punctillum Petr. & Syd. 1924, *Dothideomycetes* genera *incertae sedis*, two species, type: *P. hepaticarum* (Cooke) Petr. & Syd., asexual morph unknown, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Punctonora Aptroot 1997, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *P. nigropulvinata* Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: B, Aptroot 33356.

Pureke P.R. Johnst. 1991, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, seven species, type: *P. zelandicum* P.R. Johnst., asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type, holotype of type: PDD 45591.

Purpureocillium Luangsa-ard, Hywel-Jones, Houbraken & Samson 2011, *Ophiocordycipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *P. lilacinum* (Thom) Luangsa-ard, Houbraken, Hywel-Jones & Samson, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Luangsa-ard et al. (2011; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), Spatafora et al. (2015; new combinations), Hyde et al. (2016; new species), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Pustulipora P.F. Cannon 1982, *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *P. corticola* P.F. Cannon, asexual morph unknown, saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Pustulomyces D.Q. Dai, D.J. Bhat & K.D. Hyde 2014, *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *P. bambusicola* D.Q. Dai, D.J. Bhat & K.D. Hyde, asexual morph coelomycetous, sexual morph unknown, saprobes on dead bamboo culms, terrestrial, only known in Thailand, see Dai et al. (2014d; phylogeny, morphology), Wijayawardene et al. (2016b; morphology, phylogeny), sequence data available, holotype and ex-type strain of type: MFLU 13-0369; MFLUCC 11-0436.

Puttea S. Stenroos & Huhtinen 2009, *Lecanorales* genera *incertae sedis*, *Lecanoromycetes*, three species, type: *P. margaritella* (Hulting) S. Stenroos & Huhtinen, lichenized, Eastern Europe, America, Russia, see Stenroos et al. (2009, 2010; taxonomy, DNA), Lumbsch and

Huhndorf (2010; outline), Davydov and Printzen (2012; taxonomy), Czarnota and Hernik (2013; taxonomy, new report), Dillman et al. (2014; new combination), Lücking et al. (2017a; classification), cultures and sequences are available.

Puttemansia Henn. 1902, *Paranectriellaceae*, *Dothideomycetes* families *incertae sedis*, 19 species, type: *P. lanosa* Henn., asexual morph hyphomycetous, parasitic on *Nectandra*, worldwide, see Hyde et al. (2013; accepted as a genus in *Paranectriellaceae*), Wijayawardene et al. (2014c; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnidioarxiella Punith. & N.D. Sharma 1980, *Ascomycota* genera *incertae sedis*, one species, type: *P. indica* Punith. & N.D. Sharma, coelomycetous, sexual morph unknown, saprobes, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnidiopeltis Bat. & C.A.A. Costa 1959, *Ascomycota* genera *incertae sedis*, one species, type: *P. smilacina* Bat. & C.A.A. Costa, coelomycetous, sexual morph unknown, saprobes, USA, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnis Bref. 1881, *Ascomycota* genera *incertae sedis*, one species, type: *P. sclerotivora* Bref., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnocarpon Theiss. 1913, *Dothideomycetes* genera *incertae sedis*, two species, type: *P. magnificum* (Syd., P. Syd. & E.J. Butler) Theiss., asexual morph unknown, saprobes, terrestrial, India, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pycnodactylus Bat., A.A. Silva & Cavalc. 1967, *Ascomycota* genera *incertae sedis*, one species, type: *P. maranhensis* Bat., A.A. Silva & Cavalc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnodallia Kohlm. & Volk.-Kohlm. 2001, *Ascomycota* genera *incertae sedis*, one species, type: *P. dupla* Kohlm. & Volk.-Kohlm., coelomycetous, sexual morph unknown, saprobes, marine, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; marine genera), cultures and sequences are unavailable, holotype of type: IMS Kohlmeyer & Volkmann-Kohlmeyer 5650.

Pycnoderma Syd. & P. Syd. 1914, *Cookellaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. bambusinum* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Cookellaceae*), Hongsanan et al. (2014a; accepted as a genus in *Cookellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pycnoharknessia Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *P. pakistanica* Matsush., coelomycetous, sexual morph unknown, saprobes, terrestrial, Pakistan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: Matsushima Fungus Collection, Kobe, IPX295.

Pycnomma Syd. 1924, *Ascomycota* genera *incertae sedis*, one species, type: *P. canariense* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Canary Islands, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnomoreletia Rulamort 1990, *Ascomycota* genera *incertae sedis*, two species, type: *P. aristata* (J. Ahmad) Rulamort, coelomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Pakistan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnopeziza W.L. White & Whetzel 1938 (= *Acarosporium* Bubák & Vleugel ex Bubák 1911; = *Chaetalysis* Peyronel 1922; = *Cilirosira* Syd. 1942 fide Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *P. sympodialis* W.L. White & Whetzel, asexual morphs formerly known in *Acarosporium*, *Chaetalysis*, *Cilirosira*, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Pärtel et al. (2017; DNA, phylogeny), cultures and sequences are available.

Pycnopsammina Diederich & Etayo 1995, *Ascomycota* genera *incertae sedis*, one species, type: *P. lobariae* Diederich & Etayo, coelomycetous, sexual morph unknown, lichenicolous, non-lichenized, terrestrial, France, Spain, UK, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, holotype of the type: LG (Diederich 9697).

Pycnora Hafellner 2001, *Pycnoraceae*, *Candeliariales*, *Lecanoromycetes*, three species, type: *P. xanthococca* (Sommerf.) Hafellner, lichenized, Europe, Asia, North America, see Bendiksby and Timdal (2013; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Pycnoseynesia Kuntze 1898, *Ascomycota* genera *incertae sedis*, one species, type: *P. caronae* (Pass.)

Kuntze, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnothelia Duf. 1821, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *P. papillaria* (Ehrh.) L.M. Dufour, lichenized, see Coste and Dufrêne (2009; checklist), Crisan and Ardelean (2010; ecology), Ketner-Oostra et al. (2010; outline), Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; sequence), Zarabska and Rosadzinski (2011; new records), Grochowski (2012; ecology), Kirk et al. (2013; genus accepted), McCune and Rosentreter (2014; new record), Miadlikowska et al. (2014a; sequence), Lücking et al. (2017a; classification), Şenkardeşler et al. (2016; taxonomy), cultures and sequences are available.

Pycnothera N.D. Sharma & G.P. Agarwal 1974, *Ascomycota* genera *incertae sedis*, two species, type: *P. cordiae* G.P. Agarwal & N.D. Sharma, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnothyriella Bat. 1952, *Ascomycota* genera *incertae sedis*, two species, type: *P. discreta* Bat., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnothyrium Died. 1913, *Ascomycota* genera *incertae sedis*, six species, type: *P. litigiosum* (Desm.) Died., Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pycnotrema Rivas Plata & Lücking 2012, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *P. pycnoporellum* (Nyl.) Rivas Plata & Lücking, lichenized, terrestrial, tropical, see Rivas Plata et al. (2012a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Pyramidospora Sv. Nilsson., 1962, *Ascomycota* genera *incertae sedis*, seven species, type: *P. casuarinae* Sv. Nilsson, hyphomycetous, sexual morph unknown, on submerged decaying branches of *Casuarina*, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pyrenidium Nyl. 1865, *Dacampiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *P. actinellum* Nyl., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline,

accepted as a genus in *Dacampiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pyrenobotrys Theiss. & Syd. 1914, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. conferta* (Fr.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, America, Scandinavia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Zhang et al. (2011d; did not accept as a genus in *Venturiaceae*), Hyde et al. (2013; not accepted as a genus in *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pyrenocarpon Trevis. 1855, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *P. flotowianum* (Hepp) Trevis., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Medeiros et al. (2014; ecology and new record), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Pyrenochaeta De Not. 1849, *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *P. nobilis* De Not., coelomycetous, saprobes, plant endophytes, terrestrial, aquatic, worldwide, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), de Gruyter et al. (2010, 2013; phylogeny), Doilom et al. (2013; phylogeny), cultures and sequences are available, holotype and ex-type strains of type: CBS H-500; CBS 407.76.

Pyrenochaetopsis Gruyter, Aveskamp & Verkley 2010, *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *P. leptospora* (Sacc. & Briard) Gruyter, Aveskamp & Verkley, coelomycetous, saprobes, plant endophytes, terrestrial, aquatic, worldwide, see de Gruyter et al. (2010, 2012; phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline) Kirk et al. (2013; genus accepted), Doilom et al. (2013; phylogeny), Papizadeh et al. (2017; phylogeny), cultures and sequences are available, holotype and ex-type strains of type: CBS H-20313; CBS 101635= PD 71/1027.

Pyrenochium Link 1833, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. pini* Link, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pyrenocollema Reinke 1895, *Ascomycota* genera *incertae sedis*, 19 species, type: *P. tremelloides* Reinke, lichenized, marine, terrestrial, near sea shores, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Pérez-Ortega et al. (2016; phylogeny), cultures and sequences are available, holotype of type: H9505110.

Pyrenocyclus Petr. 1955, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. ambiguus* Petr., asexual morph unknown, saprobes, terrestrial, Hawaii, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pyrenodesmia A. Massal. 1853, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, six species, type: *P. chalybaea* (Fr.) A. Massal., lichenized, Europe, North America, China, see Arup et al. (2013; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Pyrenographa Aptroot 1991, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, three species, type: *P. xylographoides* Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of the type: CBG, Streimann 37582.

Pyrenomyxa Morgan 1895 (= *Pulveria* Malloch & Rogerson 1977 *fide* Wendt et al. 2017), *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, eight species, type: *P. globosa* Lév., asexual morph unknown, terrestrial, saprobes, mainly coprophilous, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available.

Pyrenomyxa Morgan 1895, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, three species, type: *P. invocans* Morgan, asexual morph unknown, saprobes, North America, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Pyrenopeziza Fuckel 1870 (= *Cylindrosporium* Grev. 1822 *fide* Johnston et al. 2014b), *Ploettnerulaceae*, *Helotiidae*, *Leotiomycetes*, three species, type: *P. chailletii* (Pers.) Fuckel, asexual morph formerly named in *Cylindrosporium*, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Carter et al. (2013; DNA, pathogens), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are available but lacking for the type.

Pyrenophora Fr. 1849, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, 16 species, type: *P. phaeocomes* (Rebent.) Fr., asexual morph hyphomycetous, saprobes or pathogens (phytopathogens) in terrestrial habitats, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012e; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Pleosporaceae*), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2014c, 2015c;

taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Pyrenopolyporus Lloyd 1917, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, five species, type: *P. hunteri* Lloyd, asexual morph hyphomycetous, terrestrial, saprobes, worldwide, see Wendt et al. (2017; resurrected the genus, placed in *Hypoxylaceae*), cultures and sequences are available.

Pyrenopsis Nyl. [nom. cons.] 1858, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, 40 species, type: *P. fuscatula* Nyl., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are available.

Pyrenostigme Syd. 1926, *Dothideomycetes* genera *incertae sedis*, one species, type: *P. siparunae* Syd., asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Pyrenothrix Riddle 1917, *Pyrenotrichaceae*, *Chaetothyriales*, *Eurotiomycetes*, two species, type: *P. nigra* Riddle, lichenized, asexual morph unknown, saprobes, terrestrial, North America, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available.

Pyrenowilmsia R.C. Harris & Aptroot 1991, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, one species, type: *P. ferruginea* (Müll. Arg.) R.C. Harris & Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pyrenula Ach. 1814, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, 175 species, type: *P. nitida* (Weig.) Ach., asexual morph unknown, lichenized, terrestrial, mostly tropical, see Lumbsch and Huhndorf (2010; outline), Aptroot (2012; key), Weerakoon et al. (2012a; phylogeny), Aptroot et al. (2013c; new species), Kirk et al. (2013; genus accepted), Gueidan et al. (2016; phylogeny), Lücking et al. (2017a; classification), sequences are available for the voucher F 5929 of the type.

Pyrenyllum Clem. 1909, *Ascomycota* genera *incertae sedis*, two species, type: *P. analeptum* (Ach.) Clem., coelomycetous, sexual morph unknown, lichenized, terrestrial, distribution unknown, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pyrgidium Nyl. 1867, *Sphinctrinaceae*, *Mycocaliciales*, *Eurotiomycetes*, twelve species, type: *P. bengaliense* Nyl., asexual morph unknown, saprobes, terrestrial, worldwide,

see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pyrgillus Nyl. 1857, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, four species, type: *P. americanus* Nyl., asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Weerakoon et al. (2012a; phylogeny), Gueidan et al. (2016; phylogeny), Lücking et al. (2017a; classification), cultures and sequences are available but lacking for the type.

Pyrgostroma Petr. 1951, *Ascomycota* genera *incertae sedis*, two species, type: *P. shearii* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pyricularia (Sacc.) Sacc. 1886, *Pyriculariaceae*, *Magnaporthales*, *Sordariomycetes*, 83 species, type: *P. grisea* (Sacc.) Sacc., sexual morph unknown, plant pathogens, saprobes, terrestrial, worldwide, see Consolo et al. (2008; pathotype diversity in Argentina), Hayashi and Fukuta (2009; Blast differentiation), Park and Shin (2009; new species), McKenzie et al. (2010; new species), Suzuki et al. (2010; fungicide resistance), Zhang et al. (2010a; spot disease in banana), Kohli et al. (2011; pathology), Netam et al. (2011; plant extract for controlling rice blast), Debona et al. (2012; pathological biochemistry), Yang et al. (2012; early detection of rice blast), Alexandrova et al. (2013; new species), Crous et al. (2013b, 2015a, 2016a; new species), Choi et al. (2013a; relationship of pathogenicity and phylogeny), Sousa et al. (2013; phytopathology), Klaubauf et al. (2014; phylogeny, new species), Murata et al. (2014; phylogeny), Castroagudín et al. (2016; new species), sequence data available for most of the species, including type, epitype and ex-epitype of the type: CBS H-22280, CBS 138707.

Pyriculariopsis M.B. Ellis 1971, *Magnaporthaceae*, *Magnaporthales*, *Sordariomycetes*, six species, type: *P. parasitica* (Sacc. & Berl.) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Thongkantha et al. (2009; DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Silva et al. (2015b; new species, Brazil), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Pyrimiascoma Daranag., Camporesi & K.D. Hyde 2015, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *P. trilobatum* Daranag., Camporesi & K.D. Hyde, asexual morph hyphomycetous, saprobes, terrestrial, Italy, Europe, see Daranagama et al. (2015a; morphology, phylogeny), Maharachchikumbura et al.

(2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are available, holotype and ex-type strains of type: MFLU 14-0231; MFLUCC 14-0012.

Pyrigemmula D. Magyar & Shoemaker 2011, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *P. aurantiaca* D. Magyar & R. Shoemaker, hyphomycetous, sexual morph unknown, terrestrial, saprobes, worldwide, see Magyar et al. (2011; taxonomy), Crous et al. (2012g; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available, holotype and ex-type cultures of the type: BP 101176, CBS 126743.

Pyriponomyces Cavalc. 1972, *Ascomycota* genera *incertae sedis*, one species, type: *P. maranhensis* Cavalc., coelomycetous, sexual morph unknown, lichenized, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Pyronema Carus 1834, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, two species, type: *P. Marianum* Carus, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Traeger et al. (2013; evolution), sequences are available but lacking for the type.

Pyropyxis Egger 1984, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *P. rubra* (Peck) Egger, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), sequences are available for the voucher K. Egger 323 (DAOM) of the type.

Pyrrhospora Körb. 1855, *Lecanoraceae*, *Lecanorales*, *Lecanomycetes*, one species, type: *P. quernea* (Dicks.) Körb., lichenized, see Rotter (2010; checklist), Lumbsch and Huhndorf (2010; outline), Perlmutter et al. (2012; checklist), Kirk et al. (2013; genus accepted), Singh et al. (2015; sequence), Urbanavichus and Urbanavichene (2015; checklist), Gasparyan and Sipman (2016; new records), Lücking et al. (2017a; classification), cultures and sequences are available.

Pyxidiophora Bref. & Tavel 1891, *Pyxidiophoraceae*, *Pyxidiophorales*, *Laboulbeniomycetes*, 17 species, type: *P. nyctalidis* Bref. & Tavel, asexual morph hyphomycetous, mainly coprophilous, terrestrial, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Pyxine Fr. 1825, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 75 species, type: *P. sorediata* (Ach.) Mont., lichenized, see Rout et al. (2010; ecology), Jungbluth and Marcelli (2011; taxonomy), Jungbluth et al. (2011; new species and new records), Lumbsch and Huhndorf (2010; list), Bajpai and Upadhyay (2012; ecology), Gaya et al. (2012; sequence), Mongkolsuk et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Nayaka et al. (2013; new species), Prieto et al. (2013; taxonomy, phylogeny), Shukla et al. (2013; bioaccumulation), Karakoti et al. (2014; ecology), Aptroot et al. (2014e; key, taxonomy), Pandit (2014; ecology), Bajpai et al. (2015; genotoxicity), Aptroot (2016; new records), Prieto and Wedin (2016; taxonomy), Lücking et al. (2017a; classification), cultures and sequences are available.

Quadraceae Lunghini, Pinzari & Zucconi 1996, *Ascomycota* genera *incertae sedis*, three species, type: *Q. mediterranea* Lunghini, Pinzari & Zucconi, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, see Seifert et al. (2011; morphology), Zhang et al. (2012g; new species), Wijayawardene et al. (2012, 2017; outline), Barbosa et al. (2013a, b; new species), Kirk et al. (2013; genus accepted), holotype: IMI 367816, cultures and sequences are unavailable.

Quadicladium Nawawi & Kuthub. 1989, *Ascomycota* genera *incertae sedis*, one species, type: *Q. aquaticum* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, aquatic, Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 327415, cultures and sequences are unavailable.

Quadricrura Kaz. Tanaka, K. Hiray. & Sat. Hatak. 2009, *Tetraplosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *Q. septentrionalis* Kaz. Tanaka, K. Hirayama & Sat. Hatak., sexual morph unknown, saprobes, terrestrial, Japan, see Tanaka et al. (2009; morphology, phylogeny), Hyde et al. (2013; accepted as a genus in *Tetraplosphaeriaceae*), Wijayawardene et al. (2014c; outline), holotype and ex-type of the type: HHUF 28782, CBS 125430, cultures and sequences are available.

Quasiconcha M.E. Barr & M. Blackw. 1981, *Mytilinidiaceae*, *Mytilinidiales*, *Dothideomycetes*, one species, type: *Q. reticulata* M.E. Barr & M. Blackw., asexual morph hyphomycetous, mainly coprophilous, terrestrial, USA, see Boehm et al. (2009a, b; phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mytilinidiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Quasidiscus B. Sutton 1991, *Ascomycota* genera *incertae sedis*, one species, type: *Q. simmondsii* B. Sutton, coelomycetous, sexual morph unknown, saprobes, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 263349c, cultures and sequences are unavailable.

Quasiphloeoospora B. Sutton, Crous & Shamoun 1996, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *Q. saximontanensis* (Deighton) B. Sutton, Crous & Shamoun, sexual morph unknown, one species, foliicolous, pathogens, North America, see Seifert et al. (2011; morphology), Braun et al. (2013; Cercosporoid fungi), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c, 2017; outline), holotype of type: IMI 98069, cultures and sequences are unavailable.

Quaternaria Tul. & C. Tul. 1863, *Diatrypaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *Q. persoonii* Tul. & C. Tul., asexual morph unknown, saprobes, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Queenslandia Bat. & H. Maia 1959, *Ascomycota* genera *incertae sedis*, five species, two varieties, type: *Q. pulchella* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Queenslandipendiella Quaedv. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *Q. kurandae* (Crous & J.K. Stone) Quaedv. & Crous, hyphomycetous, sexual morph unknown, pathogens, saprobes, Australia, see Quaedvlieg et al. (2014; morphology, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type culture of type: CBS H-19932, CPC 13333.

Queirozia Viégas & Cardoso 1944, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, one species, type: *Q. turbinata* Viégas & Cardoso, sexual and hyphomycetous asexual morph, biotrophic, obligate plant pathogens, terrestrial, South America, see Braun and Cook 2012 (taxonomy), Kirk et al. (2013; genus accepted), holotype of the type: IACM 26560, sequences are available.

Quezelia Faurel & Schotter 1965, *Ascomycota* genera *incertae sedis*, one species, type: *Q. hoggariana* Faurel & Schotter, coelomycetous, sexual morph unknown, coprophilous, Sahara, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Quintaria Kohlm. & Volk.-Kohlm. 1991, *Dothideomycetes* genera *incertae sedis*, four species,

type: *Q. lignatilis* (Kohlm.) Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, freshwater or marine, worldwide, see Schoch et al. (2009; phylogeny), Suetrong et al. (2009; placement in *Testudinaceae*), Hirayama et al. (2010), Wijayawardene et al. (2014c; accepted as a genus in *Dothideomycetes*), cultures and sequences are available, holotype of type: J.K.4365a = 01097442, NYBG.

Rachicladosporium Crous, U. Braun & C.F. Hill 2007, *Cladosporiaceae*, *Capnodiales*, *Dothideomycetes*, twelve species, type: *R. luciliae* Crous, U. Braun & C.F. Hill, asexual morphs hyphomycetous, terrestrial, Antarctica, Europe, New Zealand, North America, see Crous et al. (2009d; phylogeny), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2014c, 2017; outline) Egidi et al. (2014; rock-inhabiting species), Selbmann et al. (2015; distributional records of Antarctic fungi), cultures and sequences are available, holotype and ex-type strains of the type: CBS H-19891; CBS 121620 = CPC 11407.

Rachidicola K.D. Hyde & J. Fröhl. 1995, *Hypocreaceae*, *Amphisphaerales*, *Sordariomycetes*, one species, type: *R. palmae* K.D. Hyde & J. Fröhl., asexual morph unknown, saprobes, terrestrial, Hong Kong, see Lumbsch and Huhndorf (2010; outline, misspelled as *Rhachidicola*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, misspelled as *Rhachidicola*), holotype of the type: BRIP 22762, cultures and sequences are unavailable.

Raciborskiomyces Siemaszko 1925, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, c. four species, type: *R. polonicus* Siemaszko, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Racodium Fr. [nom. cons.] 1829, *Racodiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *R. rupestre* Pers., lichenized, see Muggia et al. (2008; phylogeny), Nelsen et al. (2009; phylogeny), Hawksworth et al. (2011b; typification), Lücking et al. (2017a; classification); sequences are available, holotype of type: UPS:BOT:L-024182.

Racoleus R. Sant. & D. Hawksw. 2011, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *R. trichoporus* R. Sant. & D. Hawksw., filamentous lichen, tropical, Africa, Asia, and South America, sexual morph unknown, see Hawksworth et al. (2011b; morphology, taxonomy), Jaklitsch et al. (2016a; position), Lücking et al. (2017a; classification), cultures and sequences are unavailable, holotype of type: UPS:BOT:L-638266.

Radiatispora Matsush. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *R. yunnanensis*

Matsush., coelomycetous, sexual morph unknown, coprophilous, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: MFC-0Y237.

Radotinea Velen. 1934, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *R. caudata* Velen., asexual morph unknown, saprobes, former Czechoslovakia, see Kirk et al. (2008; mentioned as *Helotiales*, genera *incertae sedis*), Lumbsch and Huhndorf (2010; did not accept as genus in *Helotiales*), Kirk et al. (2013; genus accepted), data inadequate, cultures and sequences are unavailable.

Radulidium Arzanlou, W. Gams & Crous 2007, *Dothideomycetes* genera *incertae sedis*, three species, type: *R. subulatum* (de Hoog) Arzanlou, W. Gams & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Czech Republic, USA, see Wu et al. (2013c; new species, morphology), Wijayawardene et al. (2017; outline), ex-type culture of type: CBS 405.76.

Raesaeenia D. Hawksw. C. Boluda & H. Lindgren 2015, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *R. huuskonenii* (Räsänen) D. Hawksw. C. Boluda & H. Lindgren, asexual morph unknown, on lichens, Europe, terrestrial, see Divakar et al. (2015; phylogeny), cultures and sequences are unavailable, isotype of type: K(M)-IMI 209424.

Raesaeeniana S.Y. Kondr., Kärnefelt, A. Thell, Elix & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *R. maulensis* (S.Y. Kondr. & J.-S. Hur) S.Y. Kondr., Elix, Kärnefelt & A. Thell, lichenized, see Kondratyuk et al. (2015c; classification), Lücking et al. (2017a; classification), sequences are available.

Raffaelea Arx & Hennebert 1965, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, 31 species, type: *R. ambrosiae* Arx & Hennebert, sexual morph unknown, symbionts of Ambrosia beetle, plant pathogen, terrestrial, worldwide, see Harrington et al. (2008, new species), Mayfield et al. (2008; laurel wilt on avocado), Kim et al. (2009a, c; effect of Silver nanoparticles, new species), Kolarík and Hulcr (2009; *Scolytodes unipunctatus* mycoxiota), Smith et al. (2009; laurel wilt on Camphor), Harrington et al. (2010, 2011; phylogeny, new species, origin of *Raffaelea lauricola*), Matsuda et al. (2010; phylogeny), Seo et al. (2012; phylogeny), Dreaden et al. (2014a, b; multilocus PCR assays, phylogeny), Jeyaprakash et al. (2014; molecular diagnostics), Musvuugwa et al. (2015; sexual state), Simmons et al. (2016; new species), cultures and sequences are available, holotype and ex type of the type: CBS H-7710, CBS 185.64.

Raizadenia S.L. Srivast. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *R. garhwalensis* S.L.

Srivast., hyphomycetous, sexual morph unknown, saprobes, India, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Chomnunti et al. (2014; morphology), cultures and sequences are unavailable.

Ramakrishnanella Kamat & Ullasa ex Ullasa 1970, *Ascomycota* genera *incertae sedis*, one species, type: *R. indica* Ullasa, coelomycetous, sexual morph unknown, saprobes, India, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ramalea Nyl. 1866, *Lecanorales* genera *incertae sedis*, *Lecanoromycetes*, four species; type: *R. tribulosa* Nyl.; lichenized; see Lücking et al. (2017a; classification); sequences are available.

Ramalina Ach. 1809, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, c. 230 species, type: *R. fraxinea* (L.) Ach., lichenized, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Moncada et al. (2014a; taxonomy), Lücking et al. (2017a; classification), sequences are available for few species.

Ramalodium Nyl. 1879, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, five species, type: *R. succulentum* Nyl., lichenized, see Wedin et al. (2009b; molecular), Hill et al. (2011; new record), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), de Lange and Blanchon (2014; endemic record), Ekman et al. (2014; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Ramboldia Kantvilas & Elix 1994, *Ramboldiaceae*, *Lecanorales*, *Lecanoromycetes*, 33 species, two varieties, type: *R. stuartii* (Hampe) Kantvilas & Elix, lichenized, see Kalb et al. (2009; new species), Upreti et al. (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirika et al. (2012; new record), Kuznetsova et al. (2012; new record), Mayrhofer et al. (2012; new record), Gumboski (2014; new records), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; phylogeny), Resl et al. (2015; partial sequence), Singh et al. (2015; molecular), Kantvilas (2016; new species, taxonomy), Zhao et al. (2016c; classification), Lücking et al. (2017a; classification), sequences are available for few species.

Rangea Brumm. 1992, *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, one species, type: *R. annulispore* Brumm., asexual morph unknown, saprobes, terrestrial, Netherlands, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ramicapitulum Whitton, K.D. Hyde & McKenzie 2012, *Ascomycota* genera *incertae sedis*, one species, type:

R. chinense Whitton, K.D. Hyde & McKenzie, hyphomycetous, sexual morph unknown, saprobes, Hong Kong, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ramicephala Voglmayr & G. Delgado 2002, Ascomycota genera incertae sedis, one species, type: *R. sphaerospora* Voglmayr & G. Delgado, hyphomycetous, sexual morph unknown, saprobes, aquatic, Cuba, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: WU 22035, cultures and sequences are unavailable.

Ramiclitoridium Stahel ex de Hoog 1977, *Dissconiaceae*, *Capnodiales*, *Dothideomycetes*, 35 species, type: *R. apiculatum* (J.H. Mill., Giddens & A.A. Foster) de Hoog, hyphomycetous, sexual morph unknown, saprobes, from soil terrestrial, USA, see Braun et al. (2009; new species), Crous et al. (2009d, e, f, 2014c; phylogeny, new species), Kirschner and Chen (2010; new species), Shivas et al. (2010; new species), Li et al. (2012a; new species; phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), cultures and sequences are available, ex-type cultures of the type: QM 7716, MUCL 7991, MUCL 15753, JCM 6972, IMI 100716, CBS 156.59, ATCC 13211 (fide Videira et al. 2016).

Ramimonilia Stielow & Quaedvli. 2014, *Capnodiales* genera incertae sedis, *Dothideomycetes*, one species, type: *R. apicalis* Stielow. & Quaedvli., hyphomycetous, sexual morph unknown, rock-inhabiting, Spain, see Egidi et al. (2014; morphology, phylogeny), Wijayawardene et al. (2014c, 2017; outline), type: CBS 118327 = TRN437, cultures and sequences available.

Ramoconidiifera B. Sutton, Carmarán & A.I. Romero 1996, Ascomycota genera incertae sedis, three species, type: *R. verrucosa* B. Sutton, Carmarán & A.I. Romero, hyphomycetous, sexual morph unknown, saprobes, South America, see Seifert et al. (2011; morphology), Whitton et al. (2012a; new species, morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype BAFC 33.275.

Ramonia Stizenb. 1862, *Gyalectaceae*, *Ostropales*, *Lecanoromycetes*, 24 species, type: *R. valenzueliana* (Mont.) Stizenb., lichenized, cosmopolitan, see Lendemer et al. (2009; new species), Knudsen and Kocourková (2010a, b; checklist), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Stepanchikova et al. (2013; new records), Aptroot et al. (2015c; new species), Lücking et al. (2017a; classification), sequences are available.

Ramopenidiella Crous & R.G. Shivas 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *R. cycadicola* Crous & R.G. Shivas,

hyphomycetous, sexual morph unknown, saprobe, associated with leaf spots, terrestrial, Australia, see Crous et al. (2014c; morphology, phylogeny), Wijayawardene et al. (2017; outline), culture and sequence are available, holotype and ex-type culture of type: CBS 137973, CBS H-21684.

Ramophialophora M. Caldúch, Stchigel, Gené & Guarro 2004, *Sordariales*, genera incertae sedis, *Sordariomycetes*, two species, type: *R. vesiculospora* M. Caldúch, Stchigel, Gené & Guarro, hyphomycetous, sexual morph unknown, in soil, terrestrial, Europe, see Madrid et al. (2010a; new species, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Ramsbottomia W.D. Buckley 1924, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, two species, type: *R. lamprosporoidea* W.D. Buckley, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Ramularia Unger 1833 (= *Mycosphaerella* Johanson 1884 fide Wijayawardene et al. 2014c; Rossman et al. 2015b), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 1047 species, of which 100 species are presently recognized based on DNA, type: *R. pusilla* Unger, hyphomycetous, sexual morph formerly named in *Mycosphaerella*, plant pathogens, terrestrial, mostly foliicolous, see Crous et al. (2009e, f, 2011e; taxonomy, phylogeny, new species), Kirschner (2009; monograph), Bakhshi et al. (2012; checklist), Braun (2013a, b; monograph), Bruckart et al. (2014; characterization and evaluation, biological control), Ruszkiewicz-Michalska and Połec (2014; new species, records), Videira et al. (2016; morphology, phylogeny), Park and Shin (2016; new species, taxonomy), Rossman et al. (2016b; new combination), cultures and sequences are available for most of the species, epitype and ex-epitype culture of type: CBS H-22527, CBS 124973.

Ramulariopsis Speg. 1910, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *R. cnidoscoli* Speg., hyphomycetous, with mycosphaerella-like sexual morph (only known for *R. gossypii*), plant pathogens, associated with leaf spots, Africa, Asia, Australia, Europe, North and South America, Oceania, see Seifert et al. (2011), Pirnia et al. (2012; taxonomy), Wijayawardene et al. (2012, 2014, 2017; outline), Videira et al. (2016, phylogeny), sequence data only available for the non-type: *R. gossypii* and *R. pseudoglycines*, lectotype of type: LPS 12851.

Ramulispora Miura 1920, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, twelve species, type:

R. andropogonis Miura, hyphomycetous, sexual morph unknown, plant pathogens, on Poaceae, see Crous et al. (2009d, f; review, phylogeny, taxonomy), Wijayawardene et al. (2012, 2017; outline), Videira et al. (2017; morphology, phylogeny), cultures and sequences are available.

Ramusculicola Thambug. & K.D. Hyde 2015, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *R. hailandica* Thambugala & K.D. Hyde, asexual morph unknown, saprobes, cosmopolitan, see Thambugala et al. (2015b; taxonomy), Jaklitsch et al. (2016c; treated as a synonym of *Teichospora* but in here we do not agree with them), cultures and sequences are available.

Rasamonia Houbraken & Frisvad 2011, *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, twelve species, type: *R. emersonii* (Stolk) Houbraken & Frisvad, thermotolerant, saprobes, human pathogen, terrestrial, worldwide, see Houbraken et al. (2012; description, morphology, phylogeny), Doyon et al. (2013; pulmonary and graft infection), Giraud et al. (2013; pathogenesis), Houbraken et al. (2013; antifungal susceptibility), Salgüero et al. (2013; infection in dog), Su and Cai (2013; new species), Tanney and Seifert (2013; new species), Steinmann et al. (2014; real time PCR, clinical diagnosis), Matos et al. (2015; isolation from cystic fibrosis patient), Crous et al. (2016b; new species, phylogeny), Gudmundsson et al. (2016; extrolite profiling), Ishiwada et al. (2016; mixed infection), Jurjevic et al. (2016; new species), Mouhajir et al. (2016; rep-PCR, clinical diagnosis), Steinmann et al. (2016; isavuconazole drug response), cultures and sequences are available for all species, holotype and ex-type of type: CBS H-7817; CBS 393.64.

Ranulospora Senan., Camporesi & K.D. Hyde 2017, *Sydiwiellaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *R. alnii* Senan., Camporesi & K.D. Hyde, asexual morphs unknown, saprobes, terrestrial, Europe, see Senanayake et al. (2017; taxonomy, phylogeny), holotype of the type: MFLU 16-2868, cultures and sequences are available.

Rasutoria M.E. Barr 1987, *Euantennariaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *R. abietis* (Dearn.) M.E. Barr, asexual morph unknown, plant pathogens, see Crous et al. (2009d, e, f; morphology, phylogeny), Wijayawardene et al. (2012, 2014c; outline), GenBank: GU371809, holotype of type: BPI 691065, cultures are unavailable.

Rattania Prabhug. & Bhat 2009, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *R. setulifera* Prabhug. & Bhat, hyphomycetous, sexual morph unknown, Endophytic, Terrestrial, India, see Shenoy et al. (2010; phylogeny), Seifert et al. (2011;

morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available, ex-type culture: GUFCC15501 (MTCC 9698).

Readeriella Syd. & P. Syd. 1908, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, five species, type: *R. mirabilis* Syd. & P. Syd., coelomycetous, sexual morph unknown, with *Cibiessia* synasexual morph, saprobes, pathogens, terrestrial, cosmopolitan, see Crous et al. (2009c, d, e, f, 2016a, 2016b; new species, phylogeny), Wijayawardene et al. (2012, 2014c, 2016b, 2017; outline, taxonomy, phylogeny), Quaedvlieg et al. (2014; new species, phylogeny), cultures and sequences are available.

Readerielliopsis Crous & C. Decock 2015, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *R. fuscoporiae* Crous & C. Decock, coelomycetes, sexual morph unknown, saprobes, French Guiana, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2016b, 2017; outline, phylogeny), holotype and ex-type of the type: CBS H-22229, CPC 24637.

Rebentischia P. Karst. 1869, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, 16 species, type: *R. pomiformis* P. Karst., saprobes on decaying wood, widespread, asexual morph unknown, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rechingeriella Petr. 1940, *Zopfiaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *R. insignis* Petr., asexual morph unknown, saprobes, terrestrial, Asia, Europe, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Hyde et al. (2013; morphology, outline), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Reconditella Matzer & Hafellner 1990, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *R. physconiarum* Matzer & Hafellner, asexual morph unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rectifusarium L. Lombard, Crous & W. Gams 2015, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *R. ventricosum* (Appel & Wollenw.) L. Lombard & Crous, asexual morph hyphomycetous, saprobes, terrestrial, Germany, see Lombard et al. (2015b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), holotype and ex-type cultures of the type: CBS H-21948, CBS 430.91 = NRRL 25729.

Recurvomyces Selbmann & de Hoog 2008, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*,

hyphomycetous, one species, type: *R. mirabilis* Selbmann & de Hoog, sexual morph unknown, rock-inhabiting, Antarctica, Europe, see Selbmann et al. (2008; morphology; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Quaedvlieg et al. (2014; phylogeny), Selbmann et al. (2015; distributional pattern), holotype and ex-type strain of the type: CBS H-20178, CBS 119434, cultures and sequences available.

Redbia Deighton & Piroz. 1972, *Ascomycota* genera *incertae sedis*, two species, type: *R. pucciniicola* Deighton & Piroz., hyphomycetous, sexual morph unknown, fungicolous, tropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Reddellomyces Trappe, Castellano & Malajczuk 1992, *Tuberaceae*, *Pezizales*, *Pezizomycetes*, four species, type: *R. westraliensis* (G.W. Beaton & Malajczuk) Trappe, Castellano & Malajczuk, asexual morph unknown, saprobes, saprobes, terrestrial, Australasia, Mediterranean, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species (unpublished).

Redheadia Y. Suto & Suyama 2005, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *R. quercus* Y. Suto & Suyama, asexual morph unknown, saprobes, Honshu, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Redingeria Frisch 2006, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, twelve species, type: *R. leiostoma* (Nyl.) Frisch, asexual morph unknown, lichenized, terrestrial, tropical, placement in *Graphidaceae* supported by molecular data, see Lumbsch and Huhndorf (2010; outline), Cáceres et al. (2012b; new combinations), Rivas Plata et al. (2012a, b; molecular, classification), Kirk et al. (2013; genus accepted), van den Broeck et al. (2014; new species), Lücking et al. (2017a; classification), sequences are available.

Redonia C.W. Dodge 1973, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, two species, type: *R. chilena* C.W. Dodge, lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are unavailable.

Redonographa Lücking, Tehler & Lumbsch 2013, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, five species, type: *R. chilensis* (Zahlbr.) Lücking & Tehler, asexual morph unknown, lichenized, terrestrial, tropical, see Lücking et al. (2013, 2014, 2017a; phylogeny, new species, classification), sequences are available.

Refractohilum D. Hawksw. 1977, *Ascomycota* genera *incertae sedis*, six species, type: *R. galligenum* D. Hawksw., hyphomycetous, sexual morph unknown, lichenicolous, fungicolous, widespread, terrestrial, see Seifert et al. (2011; genus accepted), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: K(M)-IMI 212682.

Regiocrella P. Chaverri & K.T. Hodge 2006, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *R. camerunensis* P. Chaverri & H.C. Evans, asexual morph unknown, mainly coprophilous, terrestrial, Cameroon, China, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Maharanachikumbura et al. (2015b, 2016b; outline), holotype of the type: CUP 67512, cultures and sequences are available.

Rehmiodothis Theiss. & Syd. 1914, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *R. osbeckiae* (Berk. & Broome) Theiss. & Syd., from living leaves, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rehmiomycella E. Müll. 1962, *Sordariomycetes* genera *incertae sedis*, one species, type: *R. phoradendri* (Rehm) E. Müll., saprobes, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Reichlingia Diederich & Scheid. 1996, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, four species, type: *R. leopoldii* Diederich & Scheid., two species known from the sexual stage, the generic type only known from the asexual stage with loose, sporodochia-like conidiomata, lichenized, on saxicolous and corticolous, Europe and Africa (Uganda), see Motiejūnaitė (2011; taxonomy), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Frisch et al. (2014), Lücking et al. (2017a; classification), holotype of the type: STU Wirth 23450, sequences are available.

Reimnitzia Kalb 2001, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *R. santensis* (Tuck.) Kalb, asexual morph unknown, lichenized, terrestrial, tropical, see Rivas Plata et al. (2012a; phylogeny), see Lumbsch and Huhndorf (2010; outline), Kraichak et al. (2014; phylogeny), Lücking et al. (2017a; phylogeny, classification), sequences are available.

Relicina (Hale & Kurok.) Hale, 1974, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 60 species, type: *R. eumorpha* (Hepp) Hale, asexual morph, undetermined, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Benatti (2011; taxonomy), Kirk et al.

(2013; genus accepted), Sethy et al. (2012; taxonomy), Crespo et al. (2010b; phylogeny), de Paz et al (2011; phylogeny), Buaruang et al. (2015; lineage), Kirka et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Remersonia Samson & Seifert 1997, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *R. thermophila* (Fergus) Seifert & Samson, hyphomycetous, sexual morph unknown, in soil, terrestrial, Asia, Australasia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Remispora Barghoon & Linder 1944, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, six species, type: *R. maritima* Linder, asexual morph unknown, saprobes, marine, worldwide, see Jones et al. (2009b, 2015; outline), Lumbsch and Huhndorf (2010; outline), Abdel-Wahab (2011b; phylogeny), Sakayaroj et al. (2011a; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are available.

Remototrachyna Divakar & A. Crespo 2009, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 20 species, type: *R. flexilis* (Kurok.) Divakar & A. Crespo, asexual morph, undetermined, terrestrial, see Flakus et al. (2012; taxonomy), Crespo et al. (2010b; phylogeny), de Paz et al. (2011; origin and diversification), Divakar et al. (2012; data partial sequencing), Thell et al. (2012; review of the family Parmeliaceae), Masson et al. (2015; sequencing data), Lücking et al. (2017a; classification), cultures and sequences are available.

Renispora Sigler & J.W. Carmich. 1979, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, two species, type: *R. flavissima* Sigler, P.K. Gaur, Lichtw. & J.W. Carmich., asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pitt et al. (2013; partial sequence), cultures available for the type: ATCC 38503.

Repetoblastiella R.F. Castañeda, Minter & M. Stadler 2010, *Ascomycota* genera *incertae sedis*, one species, type: *R. olivacea* R.F. Castañeda, Minter & M. Stadler, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Repetophragma Subram. 1992, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, 29 species, type: *R. biseptatum* (M.B. Ellis) Subram., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Marinowitz et al. (2008; new species), Silvera-Simón et al. (2009; new species), Zhang et al. (2009c; phylogeny),

Castañeda-Ruiz et al. (2011, 2013b; new species, revision), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Ma et al. (2014a, b; new species), cultures and sequences are available but lacking for the type.

Requienella Fabre 1883, *Requienellaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *R. seminuda* (Pers.) Boise, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; phylogeny), Lücking et al. (2017a; classification), cultures available for the type: RS12, RS13, sequences are available.

Resendea Bat. 1961, *Dothideomycetes* genera *incertae sedis*, two species, type: *R. paraguayensis* Bat., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Retiarius D.L. Olivier 1978, *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, two species, type: *R. superficiaris* D.L. Olivier, hyphomycetous, sexual morph unknown, on pollen, South Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Reticulosphaeria Sivan. & Bahekar 1982, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *R. indica* Sivan. & Bahekar, asexual morph unknown, on stems of *Achyranthes aspera*, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 254813, cultures and sequences are unavailable.

Retroa P.F. Cannon 1991, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *R. dimorphandrae* (F. Stevens) P.F. Cannon, from living leaves, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Retroconis de Hoog & Bat. Vegte 1989, *Ascomycota* genera *incertae sedis*, one species, type: *R. fusiformis* (S.M. Reddy & Bilgrami) de Hoog & Bat. Vegte, hyphomycetous, sexual morph unknown, saprobes, on human skin, terrestrial, Africa, Asia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 452.71.

Retrostium Nakagiri & Tad. Ito 1997, *Spathulopsporaceae*, *Spathulopsporales*, *Sordariomycetes*, one species, type: *R. amphiroeae* Nakagiri & Tad. Ito, spermatiophores or asexual morph hyphomycetous,

biotrophic on marine red alga, Japan, holotype of the type: IFO H-12210, cultures and sequences are unavailable

Rhabdo clema Syd. 1939, *Ascomycota* genera *incertae sedis*, two species, type: *R. chusqueae* Syd., coelomycetous, sexual morph unknown, saprobes, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhabdo cline Syd. 1922 (= *Meria* Vuill. 1896; *Hartigiella* Syd. & P. Syd. 1900, *Rhabdogloeum* Syd. 1922, *fide* Johnston et al. 2014b), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *R. pseudotsugae* Syd., asexual morph hyphomycetous, saprobes, pathogens, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Morgenstern et al. (2013; pathogens), Johnston et al. (2014b; nomenclature), Sweet et al. (2015; pathogens, DNA), cultures available for the type.

Rhabdodiscus Vain. 1921, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 42 species, type: *R. auberianus* (Mont.) Vain., asexual morph unknown, lichenized, terrestrial, tropical, see Rivas Plata et al. (2012a; phylogeny), Cáceres et al. (2014; taxonomy), Papong et al. (2014; taxonomy), Peláez et al. (2014; new combination), Lücking et al. (2017a; classification), Ohmura et al. (2016; taxonomy), sequences are available.

Rhabdogloeopsis Petr. 1925, *Ascomycota* genera *incertae sedis*, two species, type: *R. balsameae* (Davis) Petr., coelomycetous, sexual morph unknown, saprobes, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhabdomyces Balbiani 1889, *Ascomycota* genera *incertae sedis*, one species, type: *R. lobjoyi* Balbiani., coelomycetous, sexual morph unknown, saprobes, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhabdopsora Müll. Arg. 1888, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, two species, type: *R. polymorpha* Müll. Arg., lichenized; see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Rhabdospora (Durieu & Mont.) Sacc. 1884, *Dothideomycetes* genera *incertae sedis*, c. 150 species, type: *S. oleandri* Durieu & Mont., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhabdostromella Höhn. 1915, *Ascomycota* genera *incertae sedis*, one species, type: *R. rubi* Höhn., coelomycetous, sexual morph unknown, saprobes, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhabdostromina Died. 1921, *Ascomycota* genera *incertae sedis*, three species, type: *R. empetri* (Rostr.) Died., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Punithalingam and Spooner (2011a; reinstated the genus, morphology, new comb.), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Rhachomyces Thaxt. 1895 (= *Acanthomyces* Tixt. 1892 *fide* Species Fungorum 2017), *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, c. 80 species, type: *R. lasiophorus* (Thaxt.) Thaxt., asexual morph unknown, biotrophic, worldwide, see Rossi and Proano (2009; new species), Santamaría and Faille (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2015b; DNA, review), sequences are available.

Rhagadolobiopsis Guatim. & R.W. Barreto 2014, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes* families *incertae sedis*, two species, type: *R. thelypteris* Guatimosim & R.W. Barreto, asexual morph unknown, saprobes, terrestrial, Brazil, see Guatimosim et al. (2014; taxonomy), holotype of the type: VIC 31939, cultures and sequences are available.

Rhagadolobium Henn. & Lindau 1897, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, *Dothideomycetes*, seven species, type: *R. hemitheliae* Henn. & Lindau, asexual morph unknown, epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Inácio et al. (2012; taxonomy), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rhagadostoma Körb. 1865, *Nitschiaceae*, *Coronophorales*, *Sordariomycetes*, six species, type: *R. corrugatum* Körb., asexual morph unknown, saprobes, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rhagadostomella Etayo 2002, *Nitschiaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *R. gregaria* Etayo, asexual morph unknown, lichenicolous on *Sticta*, terrestrial, South America, see Maharachchikumbura et al. (2015b, 2016b; outline), need revision, holotype of the type: COL, Etayo 15943, cultures and sequences are unavailable.

Rhamphoria Niessl 1876, *Annulatasaceae*, *Annulatascales*, *Sordariomycetes*, 14 species, type: *R. delicatula* Niessl, asexual morph hyphomycetous (phaeoisaria-like), saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Rhamphosphaeria Kirschst. 1936, *Sordariomycetes* genera *incertae sedis*, one species, type: *R. capillata* Kirschst., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rhaphidicyrtis Vain. 1921, *Pyrenulales incertae sedis*, *Eurotiomycetes*, one species, type: *R. trichosporella* (Nyl.) Vain., asexual morph unknown, lichenized, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Rhexoacrodictys W.A. Baker & Morgan-Jones 2002, *Ascomycota* genera *incertae sedis*, five species, type: *R. erecta* (Ellis & Everh.) W.A. Baker & Morgan-Jones, sexual morph unknown, saprobes, terrestrial, see Seifert et al. (2011; morphology), Zhao et al. (2011a; outline), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Rhexoampullifera P.M. Kirk 1982, *Pyrenulales incertae sedis*, *Eurotiomycetes*, three species, type: *R. fagi* (M.B. Ellis) P.M. Kirk & C.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, terrestrial, British Isles, Brazil, Czech Republic, see Seifert et al. (2011; morphology), Koukol (2012; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhexocercosporidium U. Braun 1994, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *R. carotae* (Årvoll) U. Braun, hyphomycetous, terrestrial, soil-borne pathogens, root-inhabiting and on foliage, Asia, Europe, North America, see Nekoduka et al. (2010, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), ex-type strains of type: ATCC 18513 = CBS 418.65 = DAOM 116857.

Rhexodenticula W.A. Baker & Morgan-Jones 2001, *Cordanaceae*, *Coniochaetales*, *Sordariomycetes*, four species, type: *R. cylindrospora* (R.F. Castañeda, Saikawa & Hennebert) W.A. Baker & Morgan-Jones, saprobes, saprobes, cosmopolitan, Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Klaubauf et al. (2014; taxonomy, phylogeny), Crous et al. (2016a, b; new species, phylogeny), cultures and sequences are available.

Rhexophiale Th. Fr. 1860, *Sagiolechiaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *R. coronata* Th. Fr., lichenized, see Baloch et al. (2010; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Rhexoprolifer Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *R. variabilis* Matsush., hyphomycetous, sexual morph unknown, saprobes, France, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhexosporium Udagawa & Furuya 1977, *Sordariales incertae sedis*, *Sordariomycetes*, one species, type: *R. terrestre* Udagawa & Furuya, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rhexothecium Samson & Mouch. 1975, *Eremomycetaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *R. globosum* Samson & Mouch, asexual morph hyphomycetous (trichosporiella-like), saprobes, terrestrial, Egypt, Kenya, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Eremomycetaceae*), Kirk et al. (2013; genus accepted), Giraldo et al. (2014a; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type.

Rhinocladiella Nannf. 1934, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, c. 13 species, type: *R. atrovirens* Nannf., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Wu et al. (2013c; new species), Hernández-Restrepo et al. (2016a; new species), Madrid et al. (2016; new species, phylogeny), cultures and sequences are available.

Rhinocladium Sacc. & Marchal 1885, *Microascaceae*, *Microascales*, *Sordariomycetes*, nine species, type: *R. coprogenum* Sacc. & Marchal ex Marchal, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Lackner et al. (2012; DNA), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are available but lacking for the type.

Rhinotrichella G. Arnaud ex de Hoog 1977, *Ascomycota* genera *incertae sedis*, four species, type: *R. globulifera* G. Arnaud ex de Hoog, hyphomycetous, sexual morph unknown, saprobes, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhipidiomyces Thaxt. 1926, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *R. acriti* Thaxt., asexual morph unknown, biotrophic, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Rhipidocarpon (Theiss.) Theiss. & Syd. 1915, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *R. javanicum* (Pat.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, Java, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rhipidocephalum Trail 1888, *Ascomycota* genera *incertae sedis*, two species, type: *R. abietis* Trail, hyphomycetous, sexual morph unknown, saprobes, British Isles, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhizina Fr. 1815, *Rhizinaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *R. undulata* Fr., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Palomares-Rius et al. (2014; DNA), cultures available for the type: AFTOL-ID 918.

Rhizoblepharia Rifai 1968, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, two species, type: *R. jugispora* Rifai, asexual morph unknown, saprobes, terrestrial, Australia, Jamaica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhizocarpon Ramond ex DC. 1805, *Rhizocarpaceae*, *Lecanorales*, *Lecanoromycetes*, 325 species, 71 subspecies, 197 varieties, 188 forma species, type: *R. geographicum* (L.) DC. morph, undetermined, terrestrial, Europe, Middle East, South Korea, Turkey, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Bradwell (2010; applied studies), Matthews and Trenbirth (2011; lichenometry), Sánchez et al. (2014a; physiology), Joshi et al. (2010a; rare lichens), Aptroot and Aslan (2011; rare lichens), Lücking et al. (2017a; classification), sequences are available.

Rhizocladosporium Crous & U. Braun 2007, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *R. argillaceum* (Minoura) Crous & U. Braun asexual morph hyphomycetous, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), holotype and ex-type strains of the type: Tubaki 4262 (IFO); CBS 241.67 = IFO 7055.

Rhizodermea Verkley & Zijlstra 2010, *Dermateaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *R. veluwensis* Verkley & Zijlstra, produced only chlamydospores, sexual morph unknown, from roots, terrestrial, Europe, see

Verkley and Hofland-Zijlstra (2010; taxonomy), holotype of the type: CBS 110605 (deposited as culture).

Rhizodiscina Hafellner 1979, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *R. lignyota* (Fr.) Hafellner, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), cultures and sequences are unavailable.

Rhizogene Syd. & P. Syd. 1921, *Dothideomycetes* genera *incertae sedis*, one species, type: *R. symphoricarpi* (Syd. & P. Syd.) Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Zhang et al. (2011d; provisionally accepted as a genus in *Venturiaceae*), Hyde et al. (2013; excluded the genus from *Venturiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rhizolecia Hertel 1984, *Lecideaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *R. hybrida* (Zahlbr.) Hertel, asexual morph, undetermined, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hertel (2014; taxonomy revision), cultures and sequences are unavailable.

Rhizomyces Thaxt. 1896, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, ten species, type: *R. ctenophorus* Thaxt., asexual morph unknown, biotrophic, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Rhizophila K.D. Hyde & E.B.G. Jones 1989, *Sordariomycetes* genera *incertae sedis*, one species, type: *R. marina* K.D. Hyde & E.B.G. Jones, asexual morph unknown, saprobes, marine, Seychelles, Brunei, Thailand, North Sumatra, Bali, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: IMI 325411, cultures and sequences are unavailable.

Rhizoplaeca Zopf 1905, *Lecanoraceae*, *Lecanoromycetes*, *Lecanorales*, eleven species, five sub species, seven varieties, type: *R. opaca* (Ach.) Zopf., asexual morph unknown, lichenized, terrestrial, see Sinha and Jagadeesh Ram (2011; taxonomy), Leavitt et al. (2011a; species delimitation, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Goni et al. (2014; checklist), Kondratyuk et al. (2014; molecular phylogeny), Zhao et al. (2016c; generic concepts, taxonomy), Lücking et al. (2017a; classification), sequences are available.

Rhizopodomycetes Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, seven species, type: *R. merragatae* Thaxt., asexual morph unknown, biotrophic, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Rhizosphaera L. Mangin & Har. 1907, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, six species, type: *R. abietis* L. Mangin & Har., sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c; outline), Thambugala et al. (2014a; accepted as a genus in *Dothideaceae*, phylogeny), Kirk et al. (2013; genus accepted), Azeem et al. (2015; DNA, fungi on conifer), cultures and sequences are available but lacking for the type.

Rhizosphaerina B. Sutton 1986, *Ascomycota* genera *incertae sedis*, two species, type: *R. variabilis* B. Sutton, coelomycetous, sexual morph unknown, saprobes, Australia, South East Asia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhizotexis Theiss. & Syd. 1917, *Dothideomycetes* genera *incertae sedis*, one species, type: *R. bauhiniarum* (Henn.) Theiss. & Syd., asexual morph unknown, epiphytes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Dai et al. (2014a; taxonomy), cultures and sequences are unavailable.

Rhizothyrium Naumov 1915 (= *Rhizocalyx* Petr. 1928; *Bactrexcipula* Höhn. 1918 *fide* Johnston et al. 2014b), *Helotiales* genera *incertae sedis*, *Leotiomyces*, one species, type: *R. abietis* Naumov, asexual morph coelomycetous, saprobes, terrestrial, Russia, Chile, Siberia, see Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are unavailable.

Rhodesia Grove 1937, *Ascomycota* genera *incertae sedis*, two species, type: *R. subiecta* (Roberge ex Desm.) Grove, coelomycetous, sexual morph unknown, saprobes, Germany, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhodesiopsis B. Sutton & R. Campb. 1979, *Ascomycota* genera *incertae sedis*, two species, type: *R. gelatinosa* B. Sutton & R. Campb., coelomycetous, sexual morph unknown, saprobes, British Isles, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhodopeziza Hohmeyer & J. Moravec 1995, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *R. tuberculata* (Gamundí) J. Moravec & Hohmeyer, asexual morph unknown, saprobes, terrestrial, South

America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhodoscypha Dissing & Sivertsen 1983, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *R. ovilla* (Peck) Dissing & Sivertsen, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), cultures available for the type: HMAS97505.

Rhodosticta Woron. 1911, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *R. caraganae* Woron., coelomycetous, sexual morph unknown, saprobes, Asia, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhodotarzetta Dissing & Sivertsen 1983, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *R. rosea* (Rea) Dissing & Sivertsen, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Rhodothallus Bat. & Cif. 1959, *Ascomycota* genera *incertae sedis*, two species, type: *R. caseariaefolius* Bat., Cif. & C.A.A. Costa ex Bat. & Cif., coelomycetous, sexual morph unknown, saprobes, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhodoveronica Arzanlou, W. Gams & Crous, 2007, *Sordariomycetes* genera *incertae sedis*, type: *R. varioseptata* Arzanlou, W. Gams & Crous, hyphomycetous, sexual morph unknown, one species, saprobes, terrestrial, Germany, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype and ex-type culture of type: CBS-H 19932, CBS 431.88.

Rhombostilbella Zimm. 1902, *Ascomycota* genera *incertae sedis*, two species, type: *R. rosae* Zimm., hyphomycetous, sexual morph unknown, fungicolous, Asia, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhopalocladium Schroers, Samuels & W. Gams 1999, *Ascomycota* genera *incertae sedis*, two species, type: *R. myxophilum* Schroers, Samuels & W. Gams, hyphomycetous, sexual morph unknown, fungicolous, Asia, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhopalostroma D. Hawksw. 1977, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, eleven species, type: *R. indicum* D. Hawksw. & Muthappa, on wood and bark, tropical, Africa and Asia, asexual morph nodulisporium-like, see Lumbsch and Huhndorf (2010; outline), Stadler et al. (2010a, 2013; phylogeny), Kirk et al. (2013; genus accepted), Daranagama et al. (2014; phylogeny), Daranagama et al. (2015b; includes description of type), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available but lacking for the type: holotype of type: K(M)-IMI 190605.

Rhopographella (Henn.) Sacc. & Trotter 1913, *Sordariomycetes* genera *incertae sedis*, one species, type: *R. gaduae* (Henn.) Sacc. & Trotter, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rhopographus Nitschke ex Fuckel 1870, *Dothideomycetes* genera *incertae sedis*, six species, type: *R. filicinus* (Fr.) Nitschke ex Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rhymbocarpus Zopf 1896, *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, 17 species, type: *R. punctiformis* Zopf, asexual morph unknown, lichenicolous, terrestrial, Europe, see Diederich et al. (2010b; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Suija et al. (2015b; phylogeny), cultures and sequences are available but lacking for the type.

Rhynchodiplodia Briosi & Farneti 1904, *Ascomycota* genera *incertae sedis*, one species, type: *R. citri* Briosi & Farneti, coelomycetous, sexual morph unknown, saprobes, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhynchosomeliola Speg. 1884, *Rhynchostomataceae*, *Eurotiomycetes* families *incertae sedis*, three species, type: *R. pulchella* Speg., asexual morph unknown, lichenicolous, terrestrial, South America, Australasia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Rhynchophoromyces Thaxt. 1908, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eight species, type: *R. rostratus* (Thaxt.) Thaxt., asexual morph unknown, biotrophic on Coleoptera (Hydrophilidae), worldwide, see

Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhynchoseptoria Unamuno 1940, *Ascomycota* genera *incertae sedis*, one species, type: *R. caballeroana* Unamuno, coelomycetous, sexual morph unknown, saprobes, Morocco, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhynchosphaeria (Sacc.) Berl. 1890, *Sordariomycetes* genera *incertae sedis*, one species, type: *R. dusenii* Henn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rhynchosporina Arx 1957, *Ascomycota* genera *incertae sedis*, two species, type: *R. meinersii* (R. Sprague) Arx, hyphomycetous, sexual morph unknown, terrestrial, plant pathogens, leaf spotting, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rhynchosporium Heinzen ex A.B. Frank 1897, *Ploettnerulaceae*, *Helotiales*, *Leotiomycetes*, five species, type: *R. graminicola* Heinzen ex A.B. Frank, hyphomycetous, terrestrial, plant pathogens, leaf spotting, worldwide, see Seifert et al. (2011; morphology), Zaffarano et al. (2011, taxonomy, phylogeny), King et al. (2013, 2014, genetics, taxonomy), Torriani et al. (2014, genetics), Braun (2016, nomenclature of the type), Jaklitsch et al. (2016a; classification), ex-epitype: RS99CH6-C1a.

Rhynchostoma P. Karst. 1870 (= *Rhynchostoma* P. Karst. 1873 *fide* Species Fungorum 2017), *Rhynchostomataceae*, *Eurotiomycetes* families *incertae sedis*, c. 10 species, type: *R. minutum* P. Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see Marincowitz et al. (2008; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Rhytidenglerula Höhn. 1918, *Englerulaceae*, *Dothideomycetes* families *incertae sedis*, eleven species type: *R. carnea* (Ellis & G. Martin) Höhn., asexual morph coelomycetous, saprobes, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Dai et al. (2014c; morphology), cultures and sequences are unavailable, type specimen need recollect and sequence.

Rhytidhysteron Speg. 1881 (= *Brunaudia* (Sacc.) 1898; = *Eutryblidiella* (Rehm) Höhn., 1959; = *Rhytidopeziza* Speg. 1885; = *Triblidium* subgen. *Brunaudia* Sacc. 1883; = *Tryblidiella* Sacc. 1883; = *Tryblidiella* sect. *Eutryblidiella* Rehm 1881 *fide* Species Fungorum 2017), *Hysteriaceae*,

Hysteriales, *Dothideomycetes*, 19 species, type: *R. brasiliense* Speg, diplodia- and aposphaeria-like, saprobes, terrestrial, worldwide, see Boehm et al. (2009a, b; phylogeny), Murillo et al. (2009; morphology, chemistry, phylogeny), Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, key, phylogeny), Almeida et al. (2014a; diversity), Chacón et al. (2014; diversity), Mahajan et al. (2014; pathogenicity), Mishra et al. (2014a; pathogenicity), Pudhom and Teerawatananond (2014; chemistry), Yacharoen et al. (2015; revision, classification, morphology), Chokpaiboon et al. (2016; chemistry), Doilom et al. (2017; morphology, phylogeny), Thambugala et al. (2016; morphology, phylogeny), cultures and sequences are available, holotype of type: LPS 1189.

RhytidIELLA Zalasky 1968, *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *R. moriformis* Zalasky, asexual morph coelomycetous (phaeoseptoria-like), pathogen, terrestrial, see Lumbsch and Huhndorf (2010; outline), Doilom et al. (2013; accepted as a genus in *Cucurbitariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Rhytidospora Jeng & Cain 1977, *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, five species, type: *R. tetraspora* Jeng & Cain, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rhytisma Fr. 1819 (= *Melasmia* Lév. 1846 *fide* Johnston et al. 2014b), *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, 127 species, type: *R. acerinum* (Pers.) Fr., asexual morph unknown, saprobes, terrestrial, worldwide, see Suto (2009; new species, Japan), Wang et al. (2009a; new species), Hou et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Masumoto et al. (2014, 2015; new species, Japan, phylogeny), sequences are available for voucher Hou 203 (AAUF).

Ricasolia De Not. 1846, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, 15 species, type: *R. amplissima* (Scop.) De Not., lichenized; see Moncada et al. (2013a; taxonomy), Cornejo and Scheidegger (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Richonia Boud. 1885, *Zopfiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *R. variospora* Boud., asexual morph unknown, France, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Zopfiaceae*), Kirk et al. (2013; genus accepted),

Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rickia Cavara 1899, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 161 species, type: *R. wasmannii* Cavara, asexual morph unknown, biotrophic hosts are Acari (*Chelicera*), *Diplopoda* (*Myriapoda*), Formicidae (*Hexapoda*: *Hymenoptera*), and several other groups of insects (*Hexapoda*: *Blattodea*, *Coleoptera*, *Orthoptera*), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Haelewaters et al. (2015a; review of *R. wasmannii*), Santamaría et al. (2016; most recent species descriptions), Pflieger et al. (2016a; host specificity), sequences are available.

Rikatlia P.F. Cannon 1993, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *R. lungusaensis* (Henn.) P.F. Cannon, from living leaves, Eastern America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rileya A. Funk 1979, *Ascomycota* genera *incertae sedis*, one species, type: *R. piceae* A. Funk, coelomycetous, sexual morph unknown, saprobes, Canada, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rimaconus Huhndorf, F.A. Fernández, Joanne E. Taylor & K.D. Hyde 2001, *Chaetosphaeriales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *R. jamaicensis* (Seaver) Huhndorf, F.A. Fernández, Joanne E. Taylor & K.D. Hyde, hyphomycetous, sexual morph unknown, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariomycetidae*, genera *incertae sedis*), Huhndorf and Miller (2011; phylogeny), Maharanachikumbura et al. (2015b, 2016b; outline), cultures available for voucher SMH 5212 of the type.

Rimora (Kohlm. & Vittal) Kohlm., Volkm.-Kohlm., Suetrong, Sakay. & E.B.G. Jones 2009, *Aigialaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *R. mangrovei* (Kohlm. & Vittal) Kohlm., Volkm.-Kohlm., Suetrong, Sakay. & E.B.G. Jones, asexual morph unknown, saprobes, marine, Belize, India, Thailand, see Suetrong et al. (2009; morphology, phylogeny), Hyde et al. (2013; accepted as a genus in *Aigialaceae*), Wijayawardene et al. (2014c; outline, accepted as *Aigialaceae*), Jones et al. (2015; outline, marine), Phookamsak et al. (2015b; taxonomy), sequences are available.

Rimula Velen. 1934, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *R. faginea* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010;

outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as *Patellariaceae*), cultures and sequences are unavailable.

Rimularia Nyl., 1868, *Trapeliaceae*, *Baeomycetales*, *Lecanoromycetes*, 40 species, type, *R. limborina* Nyl., asexual morph, undetermined, terrestrial, terricolous, See Lumbsch and Huhndorf (2010; outline), Øvstedral (2012; taxonomy), Kirk et al. (2013; genus accepted), Kantvilas (2014a; taxonomy), Spribille et al. (2014a; molecular systematics), Resl et al. (2015; phylogenetic studies), Lücking et al. (2017a; classification), cultures and sequences are available.

Rinaldiella D. A. Sutton, Y. Marín, Guarro & E.H. Thomps. 2014, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *R. pentagonospora* D.A. Sutton, Y. Marín, Guarro & E.H. Thomps., asexual morph unknown, human pathogens, Georgia, see Crous et al. (2014c; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Rinodina (Ach.) Gray, 1821, *Physciaceae*, *Lecanorales*, *Lecanoromycetes*, 692 species, type: *R. sophodes* (Ach.) A. Massal., asexual morph unknown, terrestrial, corticolous, muscicolous, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Giralt et al. (2009; new species), Giralt and Klaus (2010; taxonomy), Nadyeina et al. (2010; sequencing and taxonomy), Sheard (2010; taxonomy), Kinalioglu and Aptroot (2011; new records), Sheard et al. (2011; new species), Gaya et al. (2012; comprehensive phylogenetic studies), Uperti and Patel (2012; environmental studies, ecology), Lücking et al. (2017a; classification), Resl et al. (2016, sequencing data, unpublished), sequences are available.

Rinodinella H. Mayrhofer & Poelt, 1978, *Physciaceae*, *Lecanorales*, *Lecanoromycetes*, six species, four varieties, type: *R. controversa* (A. Massal.) H. Mayrhofer & Poelt, asexual morph unknown, terrestrial, see Elix (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; taxonomy), Gasparyan and Sipman (2013; new records), Kirk et al. (2013; genus accepted). Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Riomycetes A. Ferrer, A.N. Mill., C. Sarmiento et Shearer 2012, *Sordariomycetes* genera *incertae sedis*, one species, type: *R. rotundus* A. Ferrer, A.N. Mill., Costa Rica, see Ferrer et al. (2012; taxonomy), cultures and sequences are available.

Rivilata Kohlm., Volkm.-Kohlm. & O.E. Erikss. 1998, *Dothideomycetes* genera *incertae sedis*, one species, type: *R. ius* Kohlm., Volkm.-Kohlm. & O.E. Erikss., asexual morph unknown, saprobes, USA, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rivulicola K.D. Hyde 1997, *Sordariomycetes* genera *incertae sedis*, three species, type: *R. incrustata* K.D. Hyde, saprobes, found on decaying woody substrates in freshwater habitats, Queensland, Florida, Hong Kong, see Raja et al. (2009a; morphology), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: BRIP 23340.

Rizalia Syd. & P. Syd. 1914, *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, two species, type: *R. fasciculata* Syd. & P. Syd, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Robakia Petr. 1952, *Ascomycota* genera *incertae sedis*, one species, type: *R. arctica* Petr., coelomycetous, sexual morph unknown, saprobes, Norway, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Robergea Desm., 1847, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, 13 species, type: *R. unica* Desm., asexual morph, undetermined, terrestrial, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Robillarda Sacc. 1880, *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, eight species, type: *R. sessilis* (Sacc.) Sacc., sexual morph unknown, saprobes, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2015a; phylogeny, epitypification, new species), Maharachchikumbura et al. (2015b, 2016b; outline), epitype and ex-epitype of the type: CBS H-22273, CBS114312.

Robillardiella S. Takim. 1943, *Dothideomycetes* genera *incertae sedis*, type: data inadequate, see Kirk et al. (2008; treated as nom. dub.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Robincola Velen. 1947, *Lecanoromycetes* genera *incertae sedis*, one species, type: *R. gregaria* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as *Lecanoromycetes*), cultures and sequences are unavailable.

Roburnia Velen. 1947, *Lecanoromycetes* genera *incertae sedis*, one species, type: *R. karschioides* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch

et al. (2016a; classification, treated as *Lecanoromycetes*), cultures and sequences are unavailable.

Roccella DC. 1805, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, 24 species, type: *R. fuciformis* (L.) DC., asexual morph unknown, lichenized, terrestrial, Africa, Asia, see Lumbsch and Huhndorf (2010; outline), Tehler et al. (2009b, 2010; evolution, molecular taxonomy), Aptroot and Schumm (2011; worldwide key), Bates et al. (2011; ecology), Tehler (2011, 2012; taxonomy), Kirk et al. (2013; genus accepted), Ertz et al. (2014b; taxonomy), Kambar et al. (2014; antimicrobial), Notov (2014; taxonomy), Shukla et al. (2014a; natural dyes), Fiorentino (2015; notes), Lücking et al. (2017a; classification, but wrongly citing 40 species), sequences are available.

Roccellina Darb. 1898, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, 35 species, type: *R. condensata* Darb., asexual morph unknown, lichenized, terrestrial, see Kocourková et al. (2010; taxonomy), Lumbsch and Huhndorf (2010; outline), Rotter (2010; list), Ertz and Tehler (2011; phylogeny), Kirk et al. (2013; genus accepted), de Lange (2014; taxonomy), Ertz et al. (2015b, c; phylogeny), Lücking et al. (2017ab; classification), sequences are available.

Roccellinastrum Follmann 1968, *Micareaceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *R. spongoideum* Follmann, asexual morph unknown, lichenized, terrestrial, Chile, Tasmania, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), Lücking et al. (2017a; classification); cultures and sequences are unavailable.

Roccellographa J. Steiner, 1902, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, four species, type: *R. cretacea* J. Steiner, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Rodacea W. Rossi & Santam. 2012, *Teratomyceteae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *R. salgadoi* W. Rossi & Santam., asexual morph unknown, biotrophic only host known is *Eucatops curtus* Salgado (Coleoptera, Cholevidae), Ecuador, see Rossi and Santamaría (2012; taxonomy, morphology).

Rodentomyces Doveri, Pecchia, Sarrocco & Vannacci 2010, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *R. reticulatus* Doveri, Pecchia, Sarrocco & Vannacci, asexual morph unknown, saprobes, terrestrial, Italy, see Doveri et al. (2010; taxonomy, phylogeny), Lombard et al. (2015b; taxonomy, phylogeny, doubt the placement in *Nectriaceae* since low bootstrap values in phylogenetic analyses), Maharachchikumbura et al. (2015b, 2016b; outline), Index Fungorum (2016; fixed

the nomenclature issue), holotype and ex-type cultures of the type: CBS H-21948, CBS 128675.

Rodwayella Spooner 1986 (= *Remleria* Raitv. 2004 *fide* Jaklitsch et al. 2016a), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *R. sessilis* (Rodway) Spooner, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Roesleria Thüm. & Pass. 1877, *Roesleriaceae*, *Helotiales*, *Leotiomycetes*, c. 50 species, type: *R. subterranea* (Weinm.) Redhead, asexual morph unknown, saprobes, terrestrial, Europe, see Kirchmair et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Roeslerina Redhead 1985, *Ascomycota* genera *incertae sedis*, two species, type: *R. radicella* Redhead, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available for the type (unpublished).

Rogelia Döbbeler 1999, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *R. nectrioidea* Döbbeler, asexual morph unknown, saprobes, terrestrial, Tierra del Fuego, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rogeroosiella A. Hern.-Gut. & J. Mena 1996, *Ascomycota* genera *incertae sedis*, one species, type: *R. roystoneicola* A. Hern. Gut. & J. Mena, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rogersonia Samuels & Lodge 1996, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *R. striolata* Samuels & Lodge, asexual morph unknown, saprobes, terrestrial, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Rolfidium Moberg, 1986, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, two species, *R. peltatum* Moberg, asexual morph, undetermined, terrestrial, East Africa, Sri Lanka, Mascarenes, see Ekman et al. (2008; evolution of the ascus), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); Molecular and culture data unavailable.

Rolueckia Papong, Thammath. & Boonpr. 2008, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, three species; type: *R. conspersa* (Stirt.) Papong, Thammath. &

Boonpr., lichenized, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; classification), Lücking et al. (2017a; classification), sequences are unavailable.

Romellina Petr. 1955, *Sordariomycetes* genera *incertae sedis*, one species, type: *R. variabilis* Petr., asexual morph unknown, on insects, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Romjularia Timdal 2007, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *R. lurida* (Ach.) Timdal, lichenized, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Rommelaarsia Baral & Haelew. 2015, *Hyaloscrophaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *R. flavovirens* Baral, Tanchaud & Romm., asexual morph hyphomycetous, saprobes, terrestrial, see Baral and Haelewaters (2015; morphology, DNA, phylogeny), Guatimosim et al. (2016b; phylogeny), sequences are available.

Ronnigeria Petr. 1947, *Leptopeltidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *R. arctica* (Oudem.) Petr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, accepted as a genus in *Leptopeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Ropalospora A. Massal. 1860, *Ropalosporaceae*, *Umbilicariales*, *Lecanoromycetes*, eight species and three forma species, type: *R. caffra* A. Massal., asexual morph, unknown, terrestrial, Europe, Japan, see Hauck et al. (2009; biochemical studies), Löhmus and Löhmus (2010; conservation studies), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lendemer (2012a, b; floristic studies), Lücking et al. (2017a; classification), sequences re available.

Rosasphaeria Jaklitsch & Voglmayr 2012, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *R. moravica* (Petr.) Jaklitsch & Voglmayr, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), epitype and ex-epitype of the type: WU 31334, LMM = CBS124270.

Roscoepoundia Kuntze 1898, *Ascomycota* genera *incertae sedis*, one species, type: *R. croceola* (Sacc.) Kuntze, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rosellinia De Not. 1844 (= *Amphisphaerella* Henn. 1902; = *Byssitheca* Bonord. 1864; = *Dematophora* R. Hartig 1883; = *Vrikshopama* D. Rao & P. Rag. Rao *fide*

Réblová et al. 2016c; Wendt et al. 2017), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *R. aquila* (Fr.) Ces. & De Not., asexual morph unknown, saprobes, terrestrial, Italy, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Petrini and Petrini (2012; taxonomy, monograph), Maharachchikumbura et al. (2015b, 2016b; outline), Réblová et al. (2016c; phylogeny), Su et al. (2016a; new species), Crous et al. (2017; new species), Tibpromma et al. (2017b; new species), cultures and sequences are available.

Roselliniella Vain. 1921, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, 15 species, type: *R. nephromatis* (P. Crouan & H. Crouan) Matzer & Hafellner, asexual morph unknown, saprobes, lichenicolous, terrestrial, worldwide, see Zhurbenko et al. (2009; new species), Hawksworth et al. (2010; new species, phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariales*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), van den Boom (2015; new species), cultures and sequences are available but lacking for the type.

Rosellinomyces Matzer & Hafellner 1990, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *R. trichotheliorum* Matzer & Hafellner, asexual morph unknown, lichenicolous, terrestrial, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Roselliniopsis Matzer & Hafellner 1990, *Sordariales*, genera *incertae sedis*, *Sordariomycetes*, nine species, type: *R. groedensis* (Zopf) Matzer & Hafellner, asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kaur and Gehlot (2011; new species, India), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2013b; new species, South Korea), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Rosellinula R. Sant. 1986, *Dothideomycetes* genera *incertae sedis*, one species, type: *R. haplospora* (Th. Fr. & Almq.) R. Sant., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rosenscheldia Speg. 1885, *Dothideomycetes* genera *incertae sedis*, one species, type: *R. paraguaya* Speg., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Rosenscheldiella Theiss. & Syd. 1915, *Dothideomycetes* genera *incertae sedis*, one species, type: *R. styracis* (Henn.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Sultan et al. (2011; new species, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), cultures and sequences are available only for *R. korthalsellae* Sultan, P.R. Johnst., D.C. Park & A.W. Robertson.

Roseodiscus Baral 2006, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, six species, type: *R. rhodoleucus* (Fr.) Baral, asexual morph unknown, from soil, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Hyaloscypheaceae*), Wieschollek et al. (2011; new species), Kirk et al. (2013; genus accepted), Zheng and Zhuang (2013; new species, Chi), Baral and Haelewaters (2015; DNA, phylogeny), cultures and sequences are available.

Rossiomyces R.K. Benj. 2001, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *R. falcatus* (T. Majewski) R.K. Benj., asexual morph unknown, biotrophic on Coleoptera in fungi (*Cryptophagus* species, *Cryptophagidae*), Europe, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Rossmania Lar. N. Vassiljeva 2001, *Sydiomycetidae*, *Diaporthales*, *Sordariomycetes*, one species, type: *R. ukurunduensis* Lar. N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, temperate north-eastern Asia, see Cheewangkoon et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline), Kruys and Castlebury (2012; phylogeny), Voglmayr and Jaklitsch (2014; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Senanayake et al. (2017; placement), holotype of type: VLA P-2944.

Rostania Trevis. 1880, *Collemataceae*, *Peltigerales*, *Lecanoromycetes*, eight species, type: *R. quadrata* (J. Lahm ex Körb.) Trevis., lichenized, see Otálora et al. (2013, 2014; classification), Tagirdzhanova et al. (2014; new report, Russia), Jørgensen and Palice (2015; new combination), McMullin (2015; checklist), Stehn et al. (2015; list), Lücking et al. (2017a; classification), culture and sequences are unavailable.

Rostraureum Gryzenh. & M.J. Wingf. 2005, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *R. tropicale* Gryzenh. & M.J. Wingf., asexual morph unknown, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline),

holotype and ex-type cultures of type: PREM 57519, CMW 9972.

Rstrohypoxylon J. Fourn. & M. Stadler 2010, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, eight species, type: *R. terebratum* J. Fourn. & M. Stadler, asexual morph hyphomycetous (sporothrix-like to virgariella-like), terrestrial, saprobes, Thailand, see Fournier et al. (2010; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available.

Rostrupiella Jørg. Koch, K. L. Pang & E. B. G. Jones 2007, *Lulworthiaceae*, *Lulworthiales*, *Sordariomycetes*, one species, type: *R. danica* Jørg. Koch, K.L. Pang & E.B.G. Jones, asexual morph unknown, saprobes, marine, Denmark, Norway, see Jones et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2015b, 2016b; review), sequences are available, holotype of type: JK 539.

Rosulomyces S. Marchand & Cabral 1976, *Ascomycota* genera *incertae sedis*, one species, type: *R. arthrosporioides* Marchand & Cabral, hyphomycetous, sexual morph unknown, from soil, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rota Bat., Cif. & Nascim. 1959, *Ascomycota* genera *incertae sedis*, one species, type: *R. pulchra* Bat., Cif. & Nascim., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rotiferophthora G.L. Barron 1991, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, 27 species, type: *R. globospora* G.L. Barron, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Roumegueria (Sacc.) Henn. 1908, *Dothideomycetes* genera *incertae sedis*, one species, type: *R. goudotii* (Lév.) Sacc. ex Clem. & Shear, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Roumegueriella Speg. 1880, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *R. muricospora* Speg., asexual morph gliocladium-like, coprophilous, saprobes, terrestrial, temperate, see Hirooka et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted),

Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Roussoella Sacc. 1888, *Roussellaceae*, *Pleosporales*, *Dothideomycetes*, 34 species, type: *R. nitidula* Sacc. & Paol., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ahmed et al. (2014b; new species, opportunistic pathogens), Crous et al. (2014d, 2015a, 2016b; new species), Liu et al. (2014a; new species, phylogeny), Ariyawansa et al. (2015a; new species), Wijayawardene et al. (2014c, 2016b; taxonomy), Dai et al. (2017; new species, phylogeny), Thambugala et al. (2017b, new species, phylogeny), cultures available for the type: MFLUCC 11-0182, MFLUCC 11-0634.

Roussoëllopsis I. Hino & Katum. 1965, *Roussellaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *R. japonica* (I. Hino & Katum.) I. Hino & Katum., asexual morph coelomycetous, saprobes, terrestrial, Asia, see Kirk et al. (2013; genus accepted), Liu et al. (2014a; new species, phylogeny), Phookamsak et al. (2014b; phylogeny, epitypifying *R. macrospora*), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type.

Rubikia H.C. Evans & Minter 1985, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *R. samsonii* H.C. Evans & Minter, coelomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available.

Rubrotricha Lücking, Sérus. & Vězda 2005, *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *R. helminthospora* (R. Sant.) Lücking, Sérus. & Vězda, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Rufoplaca Arup, Søchting & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, six species, type: *R. subpallida* (H. Magn.) Arup, Søchting & Frödén, asexual morph unknown, lichenized, terrestrial, worldwide, see Gaya et al. (2012; phylogeny), Arup et al. (2013; taxonomy, phylogeny), Diederich et al. (2014; new records), Halıcı et al. (2014b; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Ruggieria Cif. & Montemart. 1958, *Ascomycota* genera *incertae sedis*, one species, type: *R. glaucescens* (Trab.) Cif. & Montemart., coelomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Rugonectria P. Chaverri & Samuels 2011, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *R. rugulosa* (Pat. & Gaillard) Samuels, P. Chaverri & C. Salgado, hyphomycetous asexual morph (cylindrocarpon-like), saprobes, pathogens, terrestrial, pantropical, Japan, Indonesia, see Chaverri et al. (2011; morphology, phylogeny), Lombard et al. (2015b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 126565.

Ruhlandiella Henn. 1903, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, four species, type: *R. berolinensis* Henn., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Rubio et al. (2010; new species), Lantieri et al. (2012; new species, Italy), Kirk et al. (2013; genus accepted), sequence are available for the voucher Mycoflora of Macaronesia 1230 (C) of the type.

Ruptoseptoria Quaedv., Verkley & Crous 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *R. unedonis* (Roberge ex Desm.) Quaedv., Verkley & Crous, coelomycetous, sexual morph unknown, pathogens, terrestrial, France, see Quaedvlieg et al. (2013; morphology, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-14645, CBS 355.86.

Rusavskia S.Y. Kondr. & Kärnefelt 2003, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 17 species, type: *R. elegans* (Link) S.Y. Kondr. & Kärnefelt, lichenized, see Arup et al. (2013; taxonomy), Kondratyuk et al. (2013c, 2015; new species), Ahti et al. (2015; new combinations), Halıcı (2015; new record), Kaźmierska and Kossowska (2016; taxonomy), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Rutola J.L. Crane & Schokn. 1978, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *R. graminis* (Desm.) J.L. Crane & Schokn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Crous et al. (2015a; typification, phylogeny), isotype of the type: L0054599, cultures and sequences are unavailable.

Rutstroemia P. Karst. 1871 (= *Poculum* Velen. 1934 *fide* Jaklitsch et al. 2016a), *Rutstroemiaceae*, *Helotiales*, *Leotiomycetes*, c. 111 species, type: *R. firma* (Pers.) P. Karst., asexual morph unknown, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; phylogeny), Dougoud (2015; new species), Pärtel et al. (2017; phylogeny), cultures available for the type: KL292.

Ruwenzoria J. Fourn., M. Stadler, Læssøe & Decock 2010, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, eight

species, type: *R. pseudoannulata* J. Fourn., M. Stadler, Laessøe & Decock, asexual morph hyphomycetous (sporothrix- or nodulisporium-like), terrestrial, saprobes, Democratic Republic of Congo, see Stadler et al. (2010c; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in Hypoxylaceae), holotype of the type: MUCL 51394, cultures and sequences are available.

Ruzenia Hilber 2002, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *R. spermoides* (Hoffm.) O. Hilber, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Miller et al. (2014; new species, DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), culture available for the type: SMH4655.

Saagaromyces K.L. Pang & E.B.G. Jones 2003, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, four species, type: *S. ratnagiriensis* (S.D. Patil & Borse) K.L. Pang & E.B.G. Jones, asexual morph unknown, saprobes, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Liu et al. (2015a; new species), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Sabahriopsis Crous & M.J. Wingf. 2015, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *S. eucalypti* Crous & M.J. Wingf., hyphomycetous, saprobes, Malaysia, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2017; outline, phylogeny), holotype and ex-type of the type: CBS H-22235, CPC 24957.

Sabalicola K.D. Hyde 1995, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *S. sabalensioides* (Ellis & G. Martin) K.D. Hyde, asexual morph unknown, terrestrial, saprobes, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype: MUCL 51394, cultures and sequences are unavailable.

Sablicola E.B.G. Jones, K.L. Pang & Vrijmoed 2004, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *S. chinensis* E.B.G. Jones, K.L. Pang & Vrijmoed, asexual morph unknown, on beached driftwood, Guangdong-China, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype of the type: IMI 389302.

Sabuloglossum Hustad, A.N. Mill, Dentinger & P.F. Cannon 2013, *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, one species, type: *S. arenarium* (Rostr.) Hustad, A.N. Mill., Dentinger & P.F. Cannon, asexual morph unknown, saprobes, terrestrial, worldwide, see Hustad et al. (2013; taxonomy, phylogeny), holotype of

the type: C, no. C-F-70804, sequences are available, NCBI RefSeq: NR_120040, NG_042605.

Saccardia Cooke 1878, *Saccardiaceae*, *Ascomycota* families *incertae sedis*, three species, type: *S. quercina* Cooke, asexual morph unknown, saprobes, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Saccardinula Speg. 1885, *Trichopeltinaceae*, *Dothideomycetes* families *incertae sedis*, eleven species, type: *S. guaranitica* Speg., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Elsinoaceae*), Kirk et al. (2013; genus accepted), Jayawardena et al. (2014; excluded from *Elsinoaceae* and placed in *Trichopeltinaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Saccardoëlla Speg. 1879, *Sordariomycetes* genera *incertae sedis*, 18 species, type: *S. montellica* Speg., asexual morph unknown, terrestrial, saprobes, aquatic, widespread, see Suetrong et al. (2009; molecular), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline).

Saccharata Denman & Crous 2004, *Saccharataceae*, *Botryosphaeriales*, *Dothideomycetes*, 19 species, type: *S. proteae* (Wakef.) Denman & Crous, asexual morph coelomycetous, saprobes, associated with leaf spots, terrestrial, South Africa, see Crous et al. (2008, 2016a, b; new species, morphology, phylogeny), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c, 2017; outline), holotype and ex-type culture of type: K(M) 49510 as *Phyllachora proteae* Wakef., CBS 115206.

Saccharomyces Meyen 1870, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, seven species, type: *S. cerevisiae* Meyen ex E.C. Hansen, asexual reproduction is by multilateral budding, saprophytic, but some strains are of clinical origin, in beer, wine and other alcoholic beverages, grain-based fermented foods, occasionally in soil, tree bark, fruit flies, occasional clinical isolates, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), *Saccharomyces cerevisiae* is a model organism for studies of genetics and molecular biology and was the first eukaryote to have its genome sequence determined.

Saccharomycodes E.C. Hansen 1904, *Saccharomycodaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *S. ludwigii* (E.C. Hansen) E.C. Hansen, asexual reproduction is by bipolar budding on a broad base (bud-fission) and are lemon-shaped (apiculate), saprophytic, in tree exudates, soil, spoiled alcoholic beverages, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus

accepted), Wang et al. (2015a; classification), cultures and sequences are available.

Saccharomyopsis Schiönnung 1903, *Saccharomycetidaeae*, *Saccharomycetales*, *Saccharomycetes*, 15 species, type: *S. capsularis* Schiönnung, asexual reproduction is by multilateral budding and production of arthroconidia by some species, saprophytic, on hawthorne fruit, grapes and other fruit, soil of fruit orchards, starchy foods, beetles, tree exudates, tanning fluids, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Saccobolus Boud. 1869, *Ascobolaceae*, *Pezizales*, *Pezizomycetes*, about 27 species, type: *S. kervernii* (P. Crouan & H. Crouan) Boud., asexual morph hyphomycetous, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Ramos et al. (2011; molecular studies), Mungai et al. (2012b; ecology, taxonomy), Kirk et al. (2013; genus accepted), Doveri (2014; taxonomy, keys), Melo et al. (2014; ecology, taxonomy, keys), evolutionary lineages of *Pezizales* has been confirmed and refined by ascus septal ultrastructure, number of nuclei in spores and molecular phylogenetics.

Saccothecium Fr. 1836, *Saccotheciaceae*, *Dothideales*, *Dothideomycetes*, two species, type: *S. sepincola* (Fr.) Fr., asexual morph hyphomycetous, terrestrial, Italy, saprobes, terrestrial, Italy, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Li et al. (2016a; new species), cultures available for the type: CBS 278.32.

Sadasivania Subram. 1957, *Ascomycota* genera *incertae sedis*, three species, type: *S. girisa* Subram., hyphomycetous, sexual morph unknown, saprobes, Asia, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sagedia Ach. 1809, *Megasporaceae*, *Pertusariales*, *Lecanoromycetes*, three species, type: *S. zonata* Ach., lichenized, see Nordin et al. (2010; reinstated the genus), Lücking et al. (2017a; classification), sequences are available.

Sagediopsis Sacc. ex Vain. 1921, *Adelococcaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *S. tartarina* (Nyl.) Vain., asexual morph unknown, lichenicolous, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sagema Poelt & Grube 1993, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *S. potentillae* Poelt & Grube, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus

accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Sagenidiopsis R.W. Rogers & Hafellner 1987, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, six species; type: *S. merrotsyi* R.W. Rogers & Hafellner, lichenized, Africa, Asia, see Lumbsch and Huhndorf (2010; outline), Zambare and Christopher (2012; biopharmaceutical potential), Kirk et al. (2013; genus accepted), Ram (2014; new species - Andaman), Ertz et al. (2015b, c; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Sagenomella W. Gams 1978, *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, 13 species, type: *S. diversispora* (J.F.H. Beyma) W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Hsieh et al. (2009; new species), Houbraken and Samson (2011; phylogeny), Samson et al. (2011; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 399.69, CBS 398.69.

Sageria A. Funk 1975, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *S. tsugae* A. Funk, sexual morph unknown, saprobes, Canada, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Sagiolechia A. Massal. 1854, *Sagiolechiaceae*, *Ostropales*, *Lecanoromycetes*, nine species, three varieties, three forma species, type: *S. protuberans* (Ach.) A. Massal., asexual morph unknown, lichenized, terrestrial, Europe, see Baloch et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Gomphillaceae*), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), cultures available for the type: AFTOL-ID 4896.

Saitoella S. Goto, Sugiyama, Hamamoto & Komagata 1987, *Protomycetaceae*, *Taphriniales*, *Taphrinomycetes*, two species, type: *S. complicata* Goto, Sugiyama, Hamamoto & Komagata, asexual reproduction is mainly by multipolar enteroblastic budding, saprophyte, soil, insect frass, Bhutan, western USA, see Kirk et al. (2013; genus accepted), cultures and sequences are available.

Sakireeta Subram. & K. Ramakr. 1957, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, one species, type: *S. madreeya* Subram. & K. Ramakr., coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, India, see Crous et al. (2015b; reinstated the genus, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Salsuginaea K.D. Hyde 1991, *Salsuginaceae*, *Pleosporales*, *Dothideomycetes*, one species, type:

S. ramicola K.D. Hyde, asexual morph unknown, saprobes, marine, worldwide, see Suetrong et al. (2009; phylogeny, morphology), Jones et al. (2009b, 2015; classification), Alias et al. (2010; diversity), Zainuddin et al. (2010; antimicrobial activity), Zhang et al. (2012e; phylogeny, morphology), Hyde et al. (2013; morphology, phylogeny), Wijayawardene et al. (2014c; outline), holotype of type: BRIP 17102, cultures and sequences are available.

Samarospora Rostr. 1892, *Eurotiomycetidae* genera *incertae sedis*, one species, type: *S. potamogetonis* Rostr., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sambucina Velen. 1947, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *S. aculeata* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Samuelsia P. Chaverri & K.T. Hodge 2008, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *S. rufobrunnea* Chaverri & K.T. Hodge, asexual morph coelomycetous (aschersonia-like), saprobes, on insects, terrestrial, Bolivia, Thailand, see Chaverri et al. (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Mongkolsamrit et al. (2011; new species), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: CUP 067858, cultures and sequences are available.

Sandersoniomyces R.K. Benj. 1968, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. divaricatus* R.K. Benj., asexual morph unknown, biotrophic on Coleoptera, Staphylinidae (*Quedius* sp.), America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sanguinotrema Lücking 2015, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *S. rightii* (Taylor) Lücking, asexual morph unknown, lichenized, terrestrial, tropical, see Lücking et al. (2017a; phylogeny, classification), cultures and sequences are unavailable.

Santessonnia Hale & Vobis 1978, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, ten species, type: *S. namibensis* Hale & Vobis, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Physciaceae*), Gaya et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Sapucchaka K. Ramakr. 1956, *Dothideomycetes* genera *incertae sedis*, two species, type: *S. madreeya* K. Ramakr., asexual morph unknown, saprobes, terrestrial, India, see

Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Sarcinella Sacc. 1880 (= *Schiffnerula* Höhn. 1909; = *Questieriella* G. Arnaud ex S. Hughes 1983 *fide* Rossman et al. 2016a), *Englerulaceae*, *Dothideomycetes* genera *incertae sedis*, 71 species type: *S. heterospora* Sacc., schiffnerulaceous fungi, *Questieriella*, *Digitosarcinella* and *Mitteriella* synanamorphs of *Schiffnerula*, saprobes, plant pathogens, terrestrial, worldwide. see Hosagoudar and Riju (2009, 2011; morphology), Hosagoudar et al. (2010b, 2011b; new species), Hosagoudar et al. (2010b, 2011a, b; Schiffnerulaceous fungi), Soni et al. (2011; neem disease) Singh and Singh (2012; new species), Bhise et al. (2014; new species), Dai et al. (2014c; family description), Rossman et al. (2016a; nomenclature), sequences are unavailable.

Sarcinomyces Lindner 1898, *Eurotiomycetes* genera *incertae sedis*, five species, type: *S. crustaceus* Lindner, sexual morph unknown, saprobes, plant and animal pathogen, terrestrial, worldwide, see Ruibal et al. (2008, 2009; phylogeny, ecology), Tsuneda et al. (2011b; phylogeny), Muggia et al. (2013; lichen connection), Nagano et al. (2014; deep sea), sequences are available.

Sarcinosporon D.S. King & S.C. Jong 1975, *Ascomycota* genera *incertae sedis*, one species, type: *S. inkin* (Oho) D.S. King & S.C. Jong, hyphomycetous, sexual morph unknown, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sarcoexcipula Etayo 2008, *Ascomycota* genera *incertae sedis*, one species, type: *S. pannariae* Etayo, asexual morph unknown, parasymbiont on *Pannaria*, terrestrial, South America, see Etayo and Sancho (2008; morphology, distinction of similar lichenicolous genera), holotype of the type: MAF Etayo 23100, cultures and sequences are unavailable.

Sarcographa Fée 1921, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, c. 40 species, type: *S. cinchonarum* Fée, asexual morph unknown, lichenized, terrestrial, tropical, see Rivas Plata et al. (2012a; classification), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted). Lücking et al. (2017a; classification); sequences are available.

Sarcographina Müll. Arg. 1887, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, eight species, type: *S. cyclospora* Müll. Arg., asexual morph unknown, lichenized, terrestrial, tropical, see Rivas Plata et al. (2012a; classification), Kirk et al. (2013; genus accepted),

Kraichak et al. (2014; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Sarcogyne Flot., 1851, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, 88 species, 38 varieties, two subspecies, 32 forma species, type: *S. corrugata* Flot., asexual morph unknown, terrestrial, terricolous, saxicolous, crustose, squamulose, America, Australia, Japan, Turkey, see Knudsen et al. (2009b, 2013a; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Knudsen and Kocourková (2012; neotypification), McCarthy and Kantvilas (2013b; new species), Tokizawa et al. (2015; new species), Westberg et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Sarcoleotia S. Ito & S. Imai 1934, *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, one species, type: *S. nigra* S. Ito & S. Imai, asexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; phylogeny, accepted as a genus in *Rutstroemiaceae*), Hustad et al. (2013; taxonomy, phylogeny), sequences are available but lacking for type.

Sarcomyces Massee 1891, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *S. vinosus* Massee, asexual morph unknown, saprobes, America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as a synonym of *Haematomyxa* Sacc. in *Patellariales*), cultures and sequences are unavailable.

Sarconiptera Raiv. 2003, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *S. vinacea* Raiv., asexual morph unknown, saprobes, terrestrial, Greenland, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2016a; classification), holotype of the type: TAA 137803, cultures and sequences are unavailable.

Sarcophoma Höhn. 1916, *Ascomycota* genera *incertae sedis*, three species, type: *S. endogenospora* Höhn., asexual morph unknown, saprobes, widespread, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sarcopodium Ehrenb. 1818 (= *Lanatonectria* Samuels & Rossman 1999; *Actinostilbe* Petch 1925 *fide* Rossman et al. 2016a), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, 14 species, type: *S. circinatum* Ehrenb., hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Lombard et al. (2015b; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), cultures available for the type: CBS 587.92.

Sarcopyrenia Nyl. 1858, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. gibba* Nyl., asexual

morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Sarcosagium A. Massal. 1856, *Thelocarpaceae*, *Thelocarpales*, *Ascomycota* orders *incertae sedis*, one species, type: *S. biatorellum* A. Massal., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Sarcoscypha (Fr.) Boud. 1885 (= *Mollardiomyces* Paden 1984 *fide* Healy et al. 2016), *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, c. 30 species, type: *S. coccinea* (Jacq.) Sacc., asexual morph formerly named in *Mollardiomyces*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Romero et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Wang et al. (2016e; new species), cultures available for the type: AFTOL-ID 930 (unpublished).

Sarcosoma Casp. 1891, *Sarcosomataceae*, *Pezizales*, *Pezizomycetes*, five species, type: *S. globosum* (Schmidel) Casp., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; DNA), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Sarcosphaera Auersw. 1869, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *S. macrocalyx* (Riess) Auersw., asexual morph unknown, hypogeous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Romero et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Sarcostroma Cooke 1871, *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, 28 species, type: *S. berkeleyi* Cooke, asexual morph coelomycetous, sexual morph unknown, saprobes, endophytes, foliar pathogens in tropical and temperate regions, see Maharachchikumbura et al. (2015b, 2016b; outline), Norphanphoun et al. (2015; phylogeny), need generic revision.

Sarcotrichila Höhn. 1917, *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *S. alpina* (Fuckel) Höhn., asexual morph unknown, saprobes, terrestrial, North America, Europe, see Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available but lacking for the type.

Sarcoxylon Cooke 1883, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *S. punctatum* (Jungh.) Cooke, asexual morph unknown, terrestrial, saprobes, Asia, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline),

Senanayake et al. (2015; DNA, phylogeny), cultures and sequences are available.

Sardinella Linaldeddu, A. Alves & A.J.L. Phillips 2016, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, nine species, type: *S. urbana* Linaldeddu, A. Alves & A.J.L. Phillips, coelomycetous, sexual morph unknown, pathogens, terrestrial, Italy, see Linaldeddu et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: LISE 96308, CBS 141580 = BL179.

Sarea Fr. 1825, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, two species, type: *S. difformis* (Fr.) Fr., asexual morph unknown, on resins, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; reinstated the genus), Kirk et al. (2013; genus accepted), Sanz-Ros et al. (2015; DNA, phylogeny), Burneviča et al. (2016; DNA), Lücking et al. (2017a; classification), cultures and sequences are available.

Sarimanas M. Matsum., K. Hiray. & Kaz. Tanaka 2015, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *S. shirakamiense* Matsumura, K. Hiray. & Kaz. Tanaka, asexual morph unknown, saprobes, terrestrial, Japan, see Liu et al. (2015a; morphology, phylogeny), Tian et al. (2015; phylogeny), holotype and ex-type of the type: HHUF 30454, MAFF 244768.

Sarocladium W. Gams & D. Hawksw. 1976, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, eight species, type: *S. oryzae* (Sawada) W. Gams & D. Hawksw., hyphomycetous, sexual morph unknown, on plants, isolated from soil, human tissues, widespread, terrestrial, see Seifert et al. (2011; genus accepted), Summerbell et al. (2011; phylogeny and circumscription), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; position), cultures and sequences are available, holotype of type: TAI.

Sarrameana Vězda & P. James 1973, *Sarrameanaceae*, *Sarrameanales*, *Lecanoromycetes*, three species, type: *S. paradoxa* Vězda & P. James, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sartorya Vuill. 1927, *Sordariomycetes*, genera *incertae sedis*, one species, type: *S. fumigata* Vuill., see Kirk et al. (2008; treated as nom. dub.), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline).

Satchmopsis B. Sutton & Hodges 1975, *Ascomycota* genera *incertae sedis*, one species, type: *S. brasiliensis* B. Sutton & Hodges, coelomycetous, sexual morph unknown,

saprobes, terrestrial, widespread, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Sativumoides S.C. Ren, Jian Ma & X.G. Zhang 2012, *Ascomycota* genera *incertae sedis*, one species, type: *S. punicae* S.C. Ren, Jian Ma & X.G. Zhang, hyphomycetous, saprobes, terrestrial, China, see Ren et al. (2012a; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Saturnispora Liu & Kurtzman 1991, *Pichiaceae*, *Saccharomycetales*, *Saccharomycetes*, 20 species, type: *S. disporea* (Dekker) Z.W. Liu & Kurtzman, asexual reproduction is by multilateral budding, saprophytic, in plant rhizosphere in salt marsh, tree exudates, grapes, fruit flies, sauerkraut, soil, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), many species in this genus represent asexual taxa originally classified in the genus *Candida*, but have been transferred to the genus *Saturnispora* as new combinations, cultures and sequences are available.

Savoronala Ertz, Eb. Fisch., Killmann, Razafindr. & Sérus. 2013, *Malmideaceae*, *Lecanorales*, *Lecanoromycetes*, one species; type: *S. madagascariensis* Ertz, Eb. Fisch., Killmann, Razafindr. & Sérus., lichenized, see Ertz et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Savoryella E.B.G. Jones & R.A. Eaton. 1969, *Savoryellaceae*, *Savoryellales*, *Sordariomycetes*, ten species, type: *S. lignicola* E.B.G. Jones & R.A. Eaton, asexual morph unknown, saprobes, freshwater and marine, Australia, Brunei, China, Hong Kong, India, Japan, Malaysia, Mauritius, Mexico, Philippines, South Africa, Thailand, Taiwan, UK, see Dethoup and Manoch (2009; diversity), Kurniawati et al. (2010; diversity), Alias et al. (2010; diversity), Boonyuen et al. (2011; outline, phylogeny, morphology, classification), Awaluddin et al. (2015; diversity), Jones et al. (2016; monograph, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available, holotype of type: IMI 129784.

Savulescua Petr. 1959, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, type: *S. insignis* Petr., asexual morph unknown, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are unavailable.

Sawadaea Miyabe 1914, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, ten species, type: *S. bicornis* (Wallr. : Fr.) Homma, sexual and hyphomycetous asexual morphs, biotrophic, obligate plant pathogens, terrestrial, worldwide, see Takamatsu et al. (2008c; phylogeny and taxonomy of *S. nankinensis*), Lumbsch and Huhndorf

(2010; outline), Braun and Cook (2012; taxonomy), Kirk et al. (2013; genus accepted), sequences are available.

Saxomyces Selbmann & Isola, For diagnosis see Fungal Diversity 65(1): 174 2014, *Dothideomycetes*, genera *incertae sedis*, two species, hyphomycetous, type: *Saxomyces alpinus* Zucconi & Selbmann, sexual morph unknown, rock-inhabiting fungus. From granite, Val de La Mare, Passo Stelvio, Alps, Italy; CCFEE 5470. culture preserved in liquid nitrogen and in dried condition (NCBI accession numbers: KC315861 (SSU), KC315872 (LSU), KC315883 (mtSSU)), taxonomic species delimitation supported by molecular data.

Notes: The genus *Saxomyces* was invalidly published because the type: *S. alpinus* was invalidly published [ICN (Melbourne) Art. 40.6].

The genus and the type are here validated as follows:

Saxomyces Selbmann & Isola gen. nov., MB819855. For diagnosis see Fungal Diversity 65(1): 174 (2014).

Type: *Saxomyces alpinus* Zucconi & Selbmann sp. nov., MB819851. For diagnosis see Fungal Diversity 65(1): 174 (2014). Holotype: CBS 135222. From granite, Val de La Mare, Passo Stelvio, Alps, Italy; =CCFEE 5470, cultures and sequences available.

Additional species: *Saxomyces penninicus* Zucconi & Onofri. For diagnosis see Fungal Diversity 65(1): 175 (2014). From sandstone, Punta Indren, Monte Rosa, Alps, Italy; CCFEE 5495. culture preserved in dried condition (NCBI accession numbers: KC315864 (SSU), KC315875 (LSU), KC315886 (mtSSU)), taxonomic species delimitation supported by molecular data. The species was invalidly published because the type: *S. alpinus* was invalidly published [ICN (Melbourne) Art. 40]. It is here validated as follows:

Saxomyces penninicus Zucconi & Onofri sp. nov., MB819852. For diagnosis see Fungal Diversity 65(1): 175 (2014). From rock, Punta Indren, Monte Rosa, Alps, Italy. Holotype: CBS H-22053 (exsiccatum) = CCFEE 5495.

Saxophila Selbmann & de Hoog 2016, *Extremaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *S. tyrrhenica* Selbmann & de Hoog, hyphomycetous, sexual morph unknown, rock-inhabiting, Italy, Spain, see Isola et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), type: CBS 139725 = CCFEE 5935, cultures and sequences available.

Suttoniella S. Ahmad 1961, *Ascomycota* genera *incertae sedis*, three species, type: *S. gaubae* (Petr.) S. Ahmad, coelomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, tropical, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Scalenomyces I.I. Tav. 1985, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. endogaeus* (F. Picard) I.I. Tav., asexual morph unknown, parasites, Sardinia, see Lumbsch and Huhndorf (2010; outline), Rossi (2010; new species), Kirk et al. (2013; genus accepted), sequences and culture unavailable.

Scaphidiomyces Thaxt. 1912, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, five species, type: *S. baeocerae* Thaxt., asexual morph unknown, biotrophic on Coleoptera (Scaphidiinae), South America, West Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scaphidium Clem. 1901, *Ascomycota* genera *incertae sedis*, one species, type: *S. boutelouae* Clem., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available.

Scedosporium Sacc. ex Castell. & Chalm. 1919, *Microascaceae*, *Microascales*, *Sordariomycetes*, seven species, type: *S. apiospermum* Sacc. ex Castell. & Chalm., hyphomycetous, sexual morph unknown, saprobes, human pathogen, terrestrial, worldwide, see Gilgado et al. (2008; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lackner et al. (2014; new species), Vandepitte et al. (2014; genome study), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Crous et al. (2016a; taxonomy, phylogeny), cultures available for the type: IHEM 14462.

Scelophoromyces Thaxt. 1912, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. osorianus* Thaxt., asexual morph unknown, biotrophic on Coleoptera (Staphylinidae, *Osorius* sp.), South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scepastocarpus Santam. 2004, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. peritheciiformis* Santam., asexual morph unknown, biotrophic on aquatic Coleoptera (Hydrophilidae, *Hydrochus* sp.), Spain, see Lumbsch and Huhndorf (2010; outline), related to *Zodiomyces* Thaxt., cultures and sequences are unavailable.

Sceptrifera Deighton 1965, *Ascomycota* genera *incertae sedis*, one species, type: *S. pulchra* Deighton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schadonia Körb. 1859, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *S. alpina* Körb., asexual morph unknown, lichenized, terrestrial, see Ekman et al. (2008; molecular), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), culture and sequences are unavailable for type.

Schaereria Th. Fr. 1855, *Schaereriaceae*, *Sarrameanales*, *Lecanoromycetes*, 16 species, type: *S. lugubris* (A. Massal.) Körb., lichenized, Africa, Europe, Asia, see Kukwa and Flakus (2009; taxonomy), Lewis Smith (2009; taxonomy), Spribile et al. (2009; taxonomy, new species), Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; taxonomy), Schmull et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Resl et al. (2015; phylogeny), Tarasova et al. (2016; ecology), Lücking et al. (2017a; classification), sequences are available.

Scharifia Petr. 1955, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. procumbens* Petr., asexual morph unknown, saprobes, terrestrial, Iran, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Scheffersomyces Kurtzman & M. Suzuki 2010, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, ten species, type: *S. stipitis* (Pignal) Kurtzman & M. Suzuki, asexual reproduction is by multilateral budding, saprophytic, in beetles and beetle larvae in various trees, rotted wood, one species (*S. spartinae*) was isolated from oyster grass (*Spartina alterniflora*) marshes, worldwide, see Kurtzman and Suzuki (2010; taxonomy), *S. stipitis* (formerly *Pichia stipitis*) and *S. shehatae* (formerly *Candida*) are important for biomass utilization because of their ability to ferment D-xylose, cultures and sequences are available.

SchenckIELLA Henn. 1893 (= *Allothyrium* Syd. 1939 *fide* Hongsanan et al. 2014c), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, one species, type: *S. marcgraviae* Henn., asexual morph unknown, epiphytes, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline; accepted as a genus in *Saccardiaceae*), Hongsanan et al. (2014c; taxonomy, keys, nomenclature, accepted as a genus in *Asterinaceae*), Wijayawardene et al. (2014c; outline), placement in *Asterinaceae* supported by morphology, cultures and sequences are unavailable.

Schiffnerula Höhn. 1909, *Englerulaceae*, *Dothideomycetes* families *incertae sedis*, 98 species, type: *S. mirabilis* Höhn. 1909, asexual morph hyphomycetous, epiphytes, pathogens, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Dai et al. (2014c; morphology), cultures

and sequences are unavailable, type specimen need recollect and sequence.

Schismatomma Flot. & Körb. ex A. Massal. 1852, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, ten species, type: *S. pericleum* (Ach.) Branth & Rostr., Asia, North America, Europe, see Hauck and Wirth (2010; ecology), Kocourková et al. (2010; taxonomy), Lumbsch and Huhndorf (2010; outline), Ertz and Tehler (2011; phylogeny), Stepanchikova et al. (2011; new records), Kirk et al. (2013; genus accepted), Tehler et al. (2013a; phylogeny), Lendemer and Harris (2014; taxonomy), Himelbrant et al. (2016; new records), London (2016; new record), Motiejūnaitė et al. (2016; new records), Lücking et al. (2017a; classification), sequences are available.

Schistophoron Stirt. 1876, *Graphidaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *S. tenue* Stirt., Asia, see Tehler et al. (2009a; systematic study), Lumbsch and Huhndorf (2010; outline), Singh and Swarnalatha (2011; new species), Rivas Plata et al. (2012a; classification), Lücking et al. (2013; molecular phylogeny), Prieto et al. (2012; classification), Kirk et al. (2013; genus accepted), Weerakoon and Aptroot (2016; new species from Sri Lanka), sequences are available.

Schizochora Syd. & P. Syd. 1913, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *S. elmeri* Syd. & P. Syd., from living leaves, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Schizodiscus Brusse 1888, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *S. afroalpinus* Brusse, subpolar region, see Lumbsch and Huhndorf (2010; outline), Hertel (2014; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Schizopelte Th. Fr. 1875, *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, four species, type: *S. californica* Th. Fr., lichenized, see Lumbsch and Huhndorf (2010; outline), Ertz, and Tehler (2011; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Schizosaccharomyces Lindner 1893, *Schizosaccharomycetaceae*, *Schizosaccharomycetales*, *Schizosaccharomycetes*, four species, type: *S. pombe* Lindner, asexual reproduction is by fission, saprophytic, fruit wines and fruit, honey, dried figs, sugar cane molasses, tree fluxes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Schizothecium Corda 1838, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *S. fimicola* Corda, asexual morph unknown, on dung,

terrestrial, Spain, see Chang et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariales* genera *incertae sedis*), Mungai et al. (2012f; from dung), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Schizothyra Bat. & C.A.A. Costa 1957, *Ascomycota* genera *incertae sedis*, one species, type: *S. minuta* Bat. & C.A.A. Costa, coelomycetous, sexual morph unknown, saprobes, terrestrial, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schizothyrella Thüm. 1880, *Ascomycota* genera *incertae sedis*, one species, type: *S. quercina* (Lib.) Thüm., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schizothyrioma Höhn. 1917, *Dermateaceae*, *Helotiaceae*, *Leotiomycetes*, one species, type: *S. ptarmicae* (Desm.) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schizothyrium Desm. 1849 (= *Zygomphiala* E.W. Mason 1945 *fide* Rossman et al. 2015b, *Schizothyriaceae*, *Dothideomycetes* families *incertae sedis*, five species, type: *S. acerinum* Desm., asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013, accepted as a genus in *Schizothyriaceae*), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2014c; outline), Hongsanan et al. (2016c; evolution), Phookamsak et al. (2016; taxonomy), cultures and sequences are available.

Schizothyropsis Bat. & A.F. Vital 1960, *Ascomycota* genera *incertae sedis*, one species, type: *S. citri* Bat. & A.F. Vital., coelomycetous, sexual morph unknown, saprobes, terrestrial, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schizotrema Mangold & Lumbsch 2009, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 13 species, type: *S. zebrinum* Mangold, asexual morph unknown, lichenized, terrestrial, tropical, see Kalb et al. (2009; new report), Rivas Plata et al. (2010a; key), Sipman et al. (2012; key), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Schizotrichum McAlpine 1903, *Ascomycota* genera *incertae sedis*, one species, type: *S. lobeliae* McAlpine, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Seifert et al. (2011;

morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schizoxylon Pers. 1811, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *S. sepincola* Pers., Asia, see Lumbsch and Huhndorf (2010; outline), Muggia et al. (2011a; partial sequence), Kirk et al. (2013; genus accepted), Konoreva et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Schnablia Sacc. & P. Syd. 1899, *Leotiomycetes* genera *incertae sedis*, one species, type: *B. schnablianum* Rehm, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Schrakia Hafellner 1979, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, two species, type: *S. crassula* (Starbäck) Hafellner, asexual morph unknown, saprobes, South and North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schroeteria G. Winter 1881, *Ascomycota* genera *incertae sedis*, one species, type: *S. delastrina* (Tul. & C. Tul.) G. Winter, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schwanniomyces Klöcker emend. M. Suzuki & Kurtzman 2010, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, eight species, type: *S. occidentalis* Klöcker, asexual reproduction is by multilateral budding, saprophytic, soil, tree exudates, cucumber brines, fruit flies, human faeces, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kurtzman and Suzuki (2010; revision), cultures and sequences are available.

Schwarzmannia Pisareva 1968, *Ascomycota* genera *incertae sedis*, one species, type: *S. goebeliae* Pisareva, coelomycetous, sexual morph unknown, saprobes, terrestrial, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Schweinitziella Speg. 1888, *Trichosphaeriaceae*, *Trichosphaeriales*, *Sordariomycetes*, three species, type: *S. styracum* Speg., asexual morph unknown, fungicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Scirrhia Nitschke ex Fuckel 1870, *Dothideomycetes* genera *incertae sedis*, ten species, type: *S. rimosa* (Alb. & Schwein.) Fuckel, asexual morph unknown, saprobes,

pathogens, terrestrial, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Crous et al. (2011b; phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes* genera *incertae sedis*), cultures and sequences are available but lacking for the type.

Scirrhophoma Petr. 1941, *Ascomycota* genera *incertae sedis*, one species, type: *S. dothideoides* (Mont.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sclerococcum Fr. 1825, *Dactylosporaceae*, *Eurotiomycetes* families *incertae sedis*, 18 species, type: *S. sphaerale* (Ach.) Fr., hyphomycetous, sexual morph unknown, saprobes, lichenicolous, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Scleroconidioma Tsuneda, Currah & Thormann 2000, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. sphagnicola* Tsuneda, Currah & Thormann, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), cultures available for the type: UAMH 9731.

Sclerocrana Samuels & L.M. Kohn 1987, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, four species, type: *S. atra* Samuels & L.M. Kohn, asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sclerographiopsis Deighton 1973, *Ascomycota* genera *incertae sedis*, one species, type: *S. dalbergiae* Deighton, hyphomycetous, sexual morph unknown, saprobes, Africa, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sclerographium Berk. 1854, *Ascomycota* genera *incertae sedis*, three species, type: *S. aterrimum* Berk., hyphomycetous, sexual morph unknown, saprobes, Africa, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scleromeris Syd. 1926, *Ascomycota* genera *incertae sedis*, three species, type: *S. guazumae* Syd., coelomycetous, sexual morph unknown, saprobes, Central America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scleromitrla S. Imai 1941, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *S. shiraiana* (Henn.) S. Imai, asexual morph unknown, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; phylogeny, strains of type were grouped in *Rutstroemiaceae* and *Sclerotiniaceae*).

Scleropezicula Verkley 1999 (= *Cryptosympodula* Verkley 1999 *fide* Johnston et al. 2014b), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *S. alnicola* (J.W. Groves) Verkley, asexual morph formerly known in *Cryptosympodula*, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Sclerophoma Höhn. 1909, *Ascomycota* genera *incertae sedis*, 30 species, type: needs typification, coelomycetous, sexual morph unknown, saprobes, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sclerophora Chevall. 1826, *Coniocybaceae*, *Coniocybales*, *Coniocybomycetes*, six species; type: *S. furfuracea* (L.) Chevall., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Suija et al. (2016; phylogeny), sequences are available.

Sclerophyton Eschw. 1824, *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, 14 species, (probably less, needs revision), type: *S. elegans* Eschw., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Ertz. and Tehler (2011; phylogeny, molecular), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available

Scleropycnis Syd. & P. Syd. 1911, *Ascomycota* genera *incertae sedis*, two species, type: *S. abietina* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scleropyrenium H. Harada 1993, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, two species, type: *S. japonicum* H. Harada, asexual morph unknown, lichenized, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Prieto et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Sclerostagonospora Höhn. 1917, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, eight species, type: *S. heraclei* (Sacc.) Höhn., coelomycetous, sexual morph

unknown, saprobes, terrestrial, cosmopolitan, see Crous et al. (2011a, 2016b; new species, phylogeny), Quaedvlieg et al. (2013; new species, phylogeny), Wijayawardene et al. (2016b, 2017; taxonomy, outline, phylogeny), cultures are available.

Sclerotinia Fuckel 1870, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, 15 species, type: *S. libertiana* Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; phylogeny), cultures and sequences are available.

Sclerotium Tode 1790, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, c. 100 species (probably more than this), type: *S. complanatum* Tode, hyphomycetous, sexual morph unknown, see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable for type.

Sclerozythia Petch 1937, *Ascomycota* genera *incertae sedis*, one species, type: *S. brassicae* Petch, coelomycetous, sexual morph unknown, saprobes, British Isles, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scolecobasidiella M.B. Ellis 1971, *Ascomycota* genera *incertae sedis*, two species, type: *S. avellanea* (Sappa & Mosca) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, Africa, Caribbean, see Shenoy et al. (2010; DNA), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available.

Scolecobasidium E.V. Abbott 1927, *Sympoventuriaceae*, *Venturiales*, *Dothideomycetes*, 64 species, type: *S. terreum* E.V. Abbott, sexual stage unknown, animal pathogens, saprobes, terrestrial, aquatic, worldwide, see Hamada and Niichiro (2008; bathroom), Jiang and Zhang (2008; new species), Wu and Zhang 2008; China, soil), Hamayun et al. (2009; growth enhancement in plant), Abe and Hamada (2011; detergent rich environment), Punithalingam and Spooner (2011a; new species), Ge et al (2012; human pathogen), Muthukrishnan et al. (2012; alkali tolerance in scrub forest), D'Souza and Bhat (2013; litter and endophyte), Hao et al. (2013; new species), Ren et al. (2013a; new species), Samerpitak et al. (2014; phylogeny, taxonomy, classification), Garzoli et al. (2015; marine), Hou et al. (2015; corals), Oliveira et al. (2016a; drinking water, Brazil), Pundhir et al. (2016; pneumonia), Shi et al. (2016; human pathogen), sequences are available, ex type of type: CBS 203.27.

Scolecobeltrania Iturr., R.F. Castañeda & R. Fernández 2013, *Ascomycota* genera *incertae sedis*, one species, type: *S. vermispora* Iturr., R.F. Castañeda & R. Fernández, hyphomycetous, sexual morph unknown, saprobes,

terrestrial, Venezuela, see Iturriaga et al. (2013; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Scolecobonaria Bat. 1962, *Dothideomycetes* genera *incertae sedis*, two species, type: *S. filiformis* (W. Yamam.) Bat., asexual morph unknown, saprobes, terrestrial, Taiwan, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Scolecodochium K. Matsush. & Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *S. malayense* K. Matsush. & Matsush., hyphomycetous, sexual morph unknown, saprobes, Malaysia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scolecopeltidium F. Stevens & Manter 1925, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, c. 80 species, type: *S. salacensis* (Racib.) F. Stevens & Manter, asexual morph unknown, epiphytes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Wu et al. (2011; taxonomy), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Scolecosporiella Petr. 1921, *Ascomycota* genera *incertae sedis*, six species, type: *S. typhae* (Oudem.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scolecostigmina U. Braun 1998, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 25 species, type: *S. mangiferae* (Koord.) U. Braun & Mouch., hyphomycetous, plant pathogens, leaf spotting, Africa, Asia, Australia, North and South America, Oceania, West Indies, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Crous et al. (2012a; phylogeny), Braun et al. (2014b; taxonomy), cultures available for the type: CBS 125467, holotype of type species: Indonesia, Java, Prov. Kedu, Purworedjo, on *Mangifera indica*, 21 Sep. 1905, S.H. Koorders, no. 6, ser. 5 (unknown).

Scolecotheca Søchting & B. Sutton 1997, *Ascomycota* genera *incertae sedis*, one species, type: *S. cornuta* Søchting & B. Sutton, coelomycetous, sexual morph unknown, saprobes, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scolecoxyphium Cif. & Bat. 1956, *Dothideomycetes* genera *incertae sedis*, four species, type: *S. fraserae* Cif. &

Bat., coelomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scolecozythia Curzi 1927, *Ascomycota genera incertae sedis*, one species, type: *S. valsivora* Curzi, coelomycetous, sexual morph unknown, saprobes, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scoliciosporum A. Massal. 1852, *Scoliciosporaceae*, *Lecanorales*, *Lecanoromycetes*, 15 species; type: *S. holomelaeum* (Flörke) A. Massal., lichenized, see Spribille et al. (2009; new species), Lumbsch and Huhndorf (2010; outline), Kukwa et al. (2012; new records: Poland), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Scolicosporium Lib. ex Roum. 1880, *?Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 13 species, type: *S. fagi* Lib. ex Roum., coelomycetous, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011; treated as hyphomycetous), Wijayawardene et al. (2012, 2013b, 2016b; outline, treated as coelomycetous, phylogeny, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Scoliocarpon Nyl. 1858, *Sordariomycetes genera incertae sedis*, one species, type: *S. pupula* (Tuck.) Nyl., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Scolionema Theiss. & Syd. 1918, *Perisporiopsidaceae*, *Dothideomycetes families incertae sedis*, one species, type: *S. palmarum* (Kunze & Fr.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Perisporiopsidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Scoliotidium Bat. & Cavalc. 1963, *Ascomycota genera incertae sedis*, one species, type: *S. lauracearum* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scopaphoma Dearn. & House 1925, *Ascomycota genera incertae sedis*, one species, type: *S. corioli* Dearn. & House, coelomycetous, sexual morph unknown, saprobes, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk

et al. (2013; genus accepted), cultures and sequences are unavailable.

Scopinella Lév. 1846, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, nine species, type: *S. pleiospora* (J. Schröt.) Sacc., asexual morph unknown, saprobes, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Hypocreales*, genera incertae sedis), Kirk et al. (2013; genus accepted), Sánchez et al. (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Scopulariella Gjaerum 1971, *Ascomycota genera incertae sedis*, one species, type: *S. vaccinii* Gjaerum, hyphomycetous, sexual morph unknown, saprobes, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scopulariopsis Bainier 1907, *Microascaceae*, *Microascales*, *Sordariomycetes*, 32 species, type: *S. brevicaulis* (Sacc.) Bainier, hyphomycetous, sexual morph unknown, from soil, saprobes, human pathogen, terrestrial, worldwide, see Seifert et al. (2011; morphology), Iwen et al. (2012; human pathogen, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Pan et al. (2014; new species, China), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Grum-Grzhimaylo et al. (2016; DNA), cultures available for the type: UAMH 10915.

Scoriadopsis J.M. Mend. 1930, *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *S. miconiae* J.M. Mend., asexual morph unknown, epiphytes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2011; taxonomy), Hyde et al. (2013; accepted as a genus in *Capnodiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Scorias Fr. 1832, *Capnodiaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *S. spongiosa* (Schwein.) Fr., asexual morph unknown, epiphytes, terrestrial, tropical, see Crous et al. (2009d; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2011; taxonomy), Hyde et al. (2013; accepted as a genus in *Capnodiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Scortechinia Sacc. 1891, *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, eight species, type: *S. culcitella* (Berk. & Ravenel) Speg., asexual morph unknown, saprobes, cosmopolitan, see Mugambi and Huhndorf (2010; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Scortechiniella Arx & E. Müll. 1954, *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *S. similis* (Bres.) Arx & E. Müll., asexual morph unknown, saprobes, see Kirk et al. (2008; treated as a synonym of *Nitschkia*), Mugambi and Huhndorf (2010; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Scortechiniellopsis Sivan. 1974, *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *S. leonensis* Sivan., asexual morph unknown, saprobes, cosmopolitan, see Mugambi and Huhndorf (2010; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Scothelius Bat., J.L. Bezerra & Cavalc. 1965, *Ascomycota* genera *incertae sedis*, one species, type: *S. guianensis* Bat., J.L. Bezerra & Cavalc., coelomycetous, sexual morph unknown, saprobes, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scotiosphaeria Sivan. 1977, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. endoxyliniae* Sivan., asexual morph unknown, saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Sculptolumina Marbach 2000, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, three species, type: *S. japonica* (Tuck.) Marbach, asexual morph unknown, lichenized, terrestrial, see Giralt et al. (2009; taxonomy), Joshi et al. (2010a; new record, taxonomy), Lumbsch and Huhndorf (2010; outline), Giralt et al. (2014; taxonomy), Kondratyuk (2015; new records), Lücking et al. (2017a; classification), sequences are unavailable.

Scutaria Söchting, Arup & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *S. andina* (Räsänen) Söchting, Frödén & Arup., lichenized, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Scutellinia (Cooke) Lambotte 1887, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 75 species, type: *S. scutellata* (L.) Lambotte, asexual morph unknown, from soil, saprobes, terrestrial, Europe, see Zhuang and Yang (2008; new species), Han et al. (2010a; new species, Korea), Lumbsch and Huhndorf (2010; outline), Zhuang (2010; new species), Choi et al. (2013b; new species), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Zhuang (2013; new species, key to species in

China), cultures and sequences are available but lacking for the type.

Scutisporus Ando & Tubaki 1985, *Ascomycota* genera *incertae sedis*, one species, type: *S. brunneus* K. Ando & Tubaki, hyphomycetous, saprobes, terrestrial, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Fiua et al. (2015a; taxonomy), holotype of type: TKB-F-5093, cultures and sequences are unavailable, compare with *Biflagellosporella*.

Scutobelonium Graddon 1984, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *S. amorilens* Graddon, asexual morph unknown, saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Scutomollisia Nannf. 1976, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, 14 species, type: *S. punctum* (Rehm) Nannf., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scutopeltis Bat. & H. Maia 1957, *Ascomycota* genera *incertae sedis*, two species, type: *S. alafae* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, Ghana, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scutopycnis Bat. 1957, *Ascomycota* genera *incertae sedis*, two species, type: *S. osmanthi* Bat., coelomycetous, sexual morph unknown, saprobes, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scutula Tul. 1852, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, 41 species, two subspecies, three varieties, four forma species, type: *S. wallrothii* Tul., asexual morph unknown, lichenicolous, see Egan (2009; recent literature), Etayo and Triebel (2010; new records), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Pino-Bodas et al. (2017; new species, phylogeny), cultures and sequences are unavailable.

Scutulopsis Velen. 1934, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *S. pinacea* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Scyphospora L.A. Kantsch. 1928, *Apiosporaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *S. phyllostachydis* L.A. Kantsch., coelomycetous, sexual morph apiospora-like, saprobes, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; taxonomy,

outline), Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanan et al. (2017a; phylogeny, evolution), probably a synonym of *Arthrinium* Kunze, cultures and sequences are unavailable.

Scytalidium Pesante 1957, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, 23 species, type: *S. lignicola* Pesante, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Russia, Chile, Siberia, see Kang et al. (2010; new species), Wu and Zhang (2010; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Gautam et al. (2015; new species, India), ex-type culture available for the type: UAMH 1502, needs generic revision as genus is heterogenous (fide Seifert et al. 2011).

Scytinium (Ach.) Gray 2014, *Collemataceae*, *Lecanorales*, *Lecanoromycetes*, 48 species; type: *S. palmatum* (Huds.) Gray, lichenized, see Otálora et al. (2013, 2014; taxonomy, new combination), Diederich et al. (2014; new records: France), Gasparian and Sipman (2016; new records, Armenia), Lücking et al. (2017a; classification), sequences unavailable.

Seaverinia Whetzel 1945 (= *Verrucobotrys* Hennebert 1973 *fide* Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *Verrucobotrys geranii* (Seaver) Hennebert, asexual morph formerly named in *Verrucobotrys*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are unavailable.

Sedecimiella K. L. Pang, Alias & E. B. G. Jones 2010, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. taiwanensis* K.L. Pang, Alias & E.B.G. Jones, asexual morph unknown, saprobes, marine, Hong Kong, Taiwan, see Pang et al. (2010a; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: BBH 26390.

Sedelnikovaea S.Y. Kondr., M.H. Jeong & Hur 2015, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *S. baicalensis* (Zahlbr.) S.Y. Kondr., M.H. Jeong & Hur, asexual morph unknown, lichenized, see Kondratyuk et al. (2015b; taxonomy), Lücking et al. (2017a; classification), sequences are available but lacking for type.

Segestria Fr. 1825, *Porinaceae*, *Ostropales*, *Lecanoromycetes*, 70 species, type: *S. lectissima* Fr., lichenized, see Lücking (2008; taxonomy), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification); sequences are available but lacking for type.

Seifertia Partr. & Morgan-Jones 2002, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *S. azaleae* (Peck) Partr. & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), Li et al. (2016b; new species, China), cultures available for the type: DAOM 239135.

Seimatospriopsis B. Sutton, Ghaffer & Abbas 1972, *Ascomycota* genera *incertae sedis*, two species, type: *S. salvadorae* B. Sutton, Ghaffer & Abbas, coelomycetous, sexual morph unknown, saprobes, Pakistan, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Seimatosporium Corda 1833 (= *Discostroma* Clem. 1909 *fide* Réblová et al. 2016c), *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, 75 species, type: *S. rosae* Corda, asexual morph coelomycetous, saprobes, ?pathogens, worldwide, see Barber et al. (2011; new species, phylogeny), Tanaka et al. (2011a; phylogeny), Wijayawardene et al. (2012, 2016a, b, 2017; outline, taxonomy, new species), Crous et al. (2014c; new species), Ariyawansa et al. (2015a; new species), Norphanphoun et al. (2015; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Senanayake et al. (2015; outline, phylogeny), Réblová et al. (2016c; nomenclature), Rossman et al. (2016b; new combination), Goonasekara et al. (2016; new species), Perera et al. (2016a; sexual morph of *S. cornii*), cultures available for the type: MFLUCC 14–0621.

Seiridium Nees 1816 (= *Blogiascospora* Shoemaker et al. 1966 *fide* Réblová et al. 2016c), *Sporocadaceae*, *Amphisphaeriales*, *Sordariomycetes*, 35 species, type: *S. marginatum* Nees, asexual morph coelomycetous, saprobes, ?pathogens, worldwide, see Wijayawardene et al. (2012, 2016b; outline, taxonomy, new species), Crous et al. (2012c, 2014c; new species), Senanayake et al. (2015; outline, phylogeny), Maharachchikumbura et al. (2015a, b, 2016a; outline, phylogeny), Jaklitsch et al. (2016a; phylogeny), Réblová et al. (2016c; nomenclature), cultures available for the type: SEI1.

Seirophora Poelt 1983, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, twelve species, type: *S. magara* (Kremp.) Poelt, lichenized, see Lumbsch and Huhndorf (2010; outline), Moniri and Sipman (2011; new record), Arup et al. (2013; phylogeny), Benesperi et al. (2013; ecology), Kirk et al. (2013; genus accepted), Zhurbenko (2013; new record), Giordani et al. (2015; ecology), Kondratyuk et al. (2014, 2015a, b; taxonomy, phylogeny), Moniri et al. (2014; new record, Iran), Lücking et al. (2017a; classification), sequences are available.

Selenaspora R. Heim & Le Gal 1936, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, two species, type: *S. batava* R. Heim & Le Gal, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted).

accepted), cultures and sequences are unavailable for the type.

Selenodriella R.F. Castañeda & W.B. Kendr. 1990, *Microdochiaeae*, *Xylariales*, *Sordariomycetes*, seven species, type: *S. fertilis* (Piroz. & Hodges) R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Hernández-Restrepo et al. (2016a; new species, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Selenophoma Maire 1907, *Saccotheciaeae*, *Dothideales*, *Dothideomycetes*, 13 species, type: *S. catananches* Maire, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Cheewangkoon et al. (2009; new species, phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2015c; phylogeny), cultures and sequences are available but lacking for the type.

Selenosira Petr. 1957, *Ascomycota* genera *incertae sedis*, one species, type: *S. unica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Selenosporella G. Arnaud ex MacGarvie 1968, *Ascomycota* genera *incertae sedis*, twelve species, type: *S. curvispora* G. Arnaud ex MacGarvie, hyphomycetous, saprobes, on decaying leaves, terrestrial, worldwide, Castañeda-Ruiz et al. (2009a; key and new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Fiúza et al. (2015b; new species), holotype of type: IMI 129890, cultures and sequences are unavailable.

Selenosporopsis R.F. Castañeda & W.B. Kendr. 1991, *Ascomycota* genera *incertae sedis*, one species, type: *S. elegans* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Selenozyma Yarrow 1977, *Saccharomycetes* genera *incertae sedis*, two species, type: *S. intestinalis* (Krassiln.) Yarrow, yeast, sexual morph unknown, saprobes, terrestrial, Netherlands, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable for the type.

Selinia P. Karst. 1876, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *S. pulchra* (G. Winter)

Sacc., asexual morph tubicularia-like, coprophilous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Chaverri et al. (2011; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures are available.

Semidelitschia Cain & Luck-Allen 1969, *Delitschiaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *S. agasmatica* Cain & Luck-Allen, asexual morph unknown, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Delitschiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Semifissispora H.J. Swart 1982, *Massarinaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *S. fusiformis* H.J. Swart, asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes*, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Crous et al. (2015e; DNA, new species, phylogeny), cultures and sequences are available but lacking for the type.

Semigyalecta Vain. 1921, *Gylectaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *S. paradoxa* Vain., lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Semisphaeria K. Holm & L. Holm 1991, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. sigmundii* K. Holm & L. Holm, asexual morph unknown, saprobes, terrestrial, Norway, Sweden, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Sepedonium Link 1809, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, 13 species, type: *S. mycophilum* (Pers.) Nees, hyphomycetous, on fungi, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Septocyta Petr. 1927, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, five species, type: *S. ramealis* (Roberge ex Desm.) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Aleksić et al. (2011; use in disease control, DNA), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Septocytella Syd. 1929, *Ascomycota* genera *incertae sedis*, one species, type: *S. bambusina* Syd.,

coelomycetous, sexual morph unknown, saprobes, China, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Septofusidium W. Gams 1971, *Tilachlidiaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *S. elegantulum* (Pidopl.) W. Gams, hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Septogloewum Sacc. 1880, *Ascomycota* genera *incertae sedis*, two species, type: *S. carthusianum* (Sacc.) Sacc., coelomycetous, sexual morph unknown, saprobes, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Septomyrothecium Matsush. 1971, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *S. uniseptatum* Matsush., asexual morph unknown, saprobes, terrestrial, Asia, Caribbean, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Liu et al. (2015a; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Lombard et al. (2016; phylogeny), cultures available for the type: CBS 100966.

Septomyxella (Höhn.) Höhn. 1923, *Ascomycota* genera *incertae sedis*, one species record, type: *M. spaethianum* Allesch., coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Septonema Corda 1837, *Mytilinidiaceae*, *Mytilinidiales*, *Dothideomycetes*, c. 15 species, type: *S. secedens* Corda, sexual stage unknown, terrestrial, saprobes, lichenized, Asia, Europe, Africa, North America, see Jankowiak and Rossa (2008; associations with *Pityogenes*), Wang et al. (2008a; fallen leaves), Koukol (2010; new species), Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Wiewióra (2012; seed health), Chigoziri and Ekefan (2013; seed borne fungi), Álvarez et al. (2015; ophiostomatoid fungus), sequences are unavailable.

Septopatella Petr. 1925, *Ascomycota* genera *incertae sedis*, one species, type: *S. septata* (Jaap) Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Septoria Sacc. 1884, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, c. 1000 (200 are presently recognized based on DNA), type: *S. cytisi* Desm., coelomycetous, sexual morph mycosphaerella-like, of which, plant pathogens, terrestrial, worldwide, see Quaedvlieg et al. (2011, 2013; morphology, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Verkley et al. (2013; morphology, phylogeny), cultures and sequences are available.

Septoriella Oudem. 1889 (= *Wojnowicia* Sacc. 1892 fide Crous et al. 2015a), *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 19 species, type: *S. phragmitis* Oudem., coelomycetous, sexual morph unknown, saprobes, worldwide, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2013a, 2014c, 2017; outline, taxonomy), Crous et al. (2014c, 2015a; new species, phylogeny), Li et al. (2015c; new species), Thambugala et al. (2017b, new species, phylogeny), epitype and ex-epitype cultures of the type: CBS H-22281, CPC 24118 = CBS 140065.

Septorioides Quaedvli., Verkley & Crous 2013, *Septorioideaceae*, *Botryosphaeriales*, *Dothideomycetes*, one species, type: *S. pini-thunbergii* (S. Kaneko) Quaedvli., Verkley & Crous, coelomycetous, sexual morph unknown, pathogens, terrestrial, Japan, see Quaedvlieg et al. (2013; morphology, phylogeny), cultures and sequences are available.

Septosporiopsis W.A. Baker & Morgan-Jones 2009, *Ascomycota* genera *incertae sedis*, one species, type: *S. elaeidis* (J.M. Yen & Sulmont) W.A. Baker & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, on palm litter, Africa, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Septosporium Corda 1831, *Ascomycota* genera *incertae sedis*, 28 species, type: *S. atrum* Corda, hyphomycetous, sexual morph unknown, saprobes, on wood (*Acer*, *Carpinus*, *Sorbus*, and many others) and litter (*Bambusa*), cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Septotinia Whetzel ex J.W. Groves & M.E. Elliott 1961 (= *Septotis* N.F. Buchw. ex Arx 1970 fide Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *S. podophyllina* Whetzel, asexual morph formerly known in *Septotis*, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures and sequences are unavailable.

Septotrapelia Aptroot & Chaves 2007, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, four species, seven

species, type: *S. glauca* Aptroot & Chaves, lichenized, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Aptroot and Iqbal (2012; new record, Pakistan), Bungartz et al. (2013b; taxonomy), Aptroot and Moon (2014; new species, Korea), Lücking et al. (2017a; classification), holotype of the type: INB, Aptroot 60426, cultures and sequences are unavailable

Septotrullula Höhn. 1902, *Ascomycota genera incertae sedis*, four species, type: *S. bacilligera* Höhn., hyphomycetous, sexual morph unknown, saprobes, on bark (*Betula*, *Fagus*, *Quercus*), Australasia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Serenomyces Petr. 1952, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, five species, type: *S. shearii* Petr., saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Phaeochoraceae*), Kirk et al. (2013; genus accepted), Elliott and Des Jardin (2014; species associated with palms in southeastern USA), Maharachchikumbura et al. (2015b, 2016b; outline), cultures are available for unidentified species (unpublished).

Seriascoma Phook., D.Q. Dai & K.D. Hyde 2016, *Occultibambusaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. didymospora* Phook., D.Q. Dai, Karunar. & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Thailand, see Dai et al. (2017; phylogeny, morphology, sexual and asexual morphs), ex-type culture of type: MFLUCC 11–0179.

Servazziella J. Reid & C. Booth 1987, *Sordariomycetes genera incertae sedis*, five species, type: *S. longispora* (Servazzi) J. Reid & C. Booth, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Servitia M.S. Christ. & Alstrup 2001, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, species, type: *S. inconspicuum* (Lynge) M.S. Christ. & Alstrup, asexual morph unknown, lichenized, terrestrial, Greenland, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Sessiospora D. Hawksw. 1979, *Ascomycota genera incertae sedis*, one species, type: *S. bicolor* D. Hawksw., sexual morph unknown, lichenicolous on foliicolous lichens, Asia, terrestrial, see Seifert et al. (2011; genus accepted), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: K(M)-IMI 41011c.

Setameliola D.R. Reynolds 2010, *Meliolaceae*, *Meliolales*, *Sordariomycetes*, 26 species, type: *S. argentina* (Speg.) D.R. Reynolds, asexual morph unknown, saprobes, terrestrial, Singapore, see Reynolds (2010; taxonomy), cultures and sequences are unavailable.

Setiferotheca Matsush. 1995, *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *S. nipponica* Matsush., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Setocampanula Sivan. & W.H. Hsieh 1989, *Trichosphaeriaceae*, *Trichosphaeriales*, *Sordariomycetes*, one species, type: *S. taiwanensis* Sivan. & W.H. Hsieh, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences unavailable.

Setolibertella Punith. & Spooner 1999, *Ascomycota genera incertae sedis*, one species, type: *S. solani* Punith. & Spooner, sexual morph unknown, coelomycetous, saprobes, on dead stems of *Solanum mauritanium*, Azores, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: KM 58669, cultures and sequences are unavailable.

Setomelanomma M. Morelet 1980, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. holmii* M. Morelet, asexual morph coelomycetous, saprobes or pathogens, terrestrial, cosmopolitan, see Zhang et al. (2012e; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Phaeosphaeriaceae*), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Setophaeosphaeria Crous & Y. Zhang ter 2014, *Pleosporales genera incertae sedis*, *Dothideomycetes*, three species, type: *S. hemerocallidis* Crous & Y. Zhang ter, with coelomycetous asexual morph, saprobes, ?pathogens, terrestrial, China, see Crous et al. (2014c; morphology, phylogeny), Wijayawardene et al. (2014c; outline), holotype and ex-type culture of type: CBS H-21720, CPC 23645.

Setophiale Matsush. 1995, *Ascomycota genera incertae sedis*, one species, type: *S. unisetulata* Matsush., sexual morph unknown, hyphomycetous, saprobes, on palm petioles, Peru, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: Matsushima Fungus Collection, Kobe, MFC-4P766, cultures and sequences are unavailable.

Setophoma Gruyter, Aveskamp & Verkley 2010, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, asexual morph coelomycetous, five species, type: *S. terrestris* (H.N. Hansen) Gruyter, Aveskamp &

Verkley, saprobes, ?pathogens, terrestrial, Brazil, North America, see de Gruyter et al. (2010; morphology, phylogeny), Quaedvlieg et al. (2013; morphology, phylogeny), Crous et al. (2014c, 2016b; morphology, phylogeny, sexual morph), Phookamsak et al. (2014a; sexual morph, phylogeny), Wijayawardene et al. (2014c; outline), Thambugala et al. (2017b, new species, phylogeny), lectotype and ex-lectotype culture of type: CBS H-20311, CBS 335.29.

Setoseptoria Quaedvly., Verkley & Crous 2013, *Lentiteliaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. phragmitis* Quaedvly., Verkley & Crous, coelomycetous, sexual morph unknown, pathogens, terrestrial, Hong Kong, see Quaedvlieg et al. (2013; morphology, phylogeny), Wijayawardene et al. (2014c; outline), holotype of the type: CBS H-21324, culture ex-type CBS 114802.

Setosporella Mustafa & Abdul-Wahid 1989, *Ascomycota* genera *incertae sedis*, one species, type: *S. mahmoudii* Moustafa & Abdul-Wahid, sexual morph unknown, hyphomycetous, saprobes, isolated from soil, Egypt, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 326932, cultures and sequences are unavailable.

Setosynnema D.E. Shaw & B. Sutton 1985, *Ascomycota* genera *incertae sedis*, one species, type: *S. isthmosporum* D.E. Shaw & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Seuriatiopsis Woron. 1934, *Seuriatiaceae*, *Ascomycota* families *incertae sedis*, one species, type: *S. epiphylla* Woron., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Seychellomyces Matsush. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *S. hexagonus* Matsush., sexual morph unknown, hyphomycetous, saprobes, isolated from soil, Seychelles, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Seynesia Sacc. 1883, *Cainiaceae*, *Xylariales*, *Sordariomycetes*, two species, type: *S. nobilis* (Welw. & Curr.) Sacc., tropical, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable for type.

Seynesiella G. Arnaud 1918, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, c. 80 species, type:

S. juniperi (Desm.) G. Arnaud, asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; taxonomy), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Seynesiopeltis F. Stevens & R.W. Ryan 1925, *Seynesiopeltidaceae*, *Dothideomycetes*, *incertae sedis*, one species, type: *S. tetraplasandrae* F. Stevens & R.W. Ryan, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Seynesiopsis Henn. 1904, *Ascomycota* genera *incertae sedis*, one species, type: *S. rionegrensis* Henn., sexual morph unknown, coelomycetous, saprobes, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Shackletonia Søchting, Frödén & Arup 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, five species, type: *S. hertelii* (Søchting, Øvstvedal & Sancho) Søchting, Frödén & Arup, asexual morph unknown, lichenized, terrestrial, see Arup et al. (2013; taxonomy), Kondratyuk et al. (2014; phylogeny), Garrido-Benavent et al. (2016; new species), Lücking et al. (2017a; classification), sequences are available.

Shanarella R.K. Benj. 1956, *Arthrodermataceae*, *Onygenales*, *Eurotiomycetes*, 17 species, type: *S. spirotricha* R.K. Benj., a sexual morph unknown, on humans and other mammals, causing microsporoses, terrestrial, USA, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), culture available for the type: CBS 305.56.

Shawiella Hansf. 1957, *Ascomycota* genera *incertae sedis*, one species, type: *S. grevilleae* Hansf., sexual morph unknown, coelomycetous, saprobes, on leaves of *Grevillea robusta*, New Guinea, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Shearia Petr. 1924, *Ascomycota* genera *incertae sedis*, two species, type: *S. magnoliae* (Shear) Petr., coelomycetous, sexual morph unknown, saprobes, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sheariella Petr. 1952, *Ascomycota* genera *incertae sedis*, one species, type: *S. carbonacea* Petr., sexual morph unknown, coelomycetous, saprobes, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sheathnema Dubey & Moonnambeth 2014, *Ascomycota* genera *incertae sedis*, two species, type: *S. indicum* Dubey & Moonnambeth, hyphomycetous, sexual morph unknown, saprobes, epiphytes, India, see Dubey and Moonnambeth (2014a; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Shimizuomyces Kobayasi 1981, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *S. paradoxus* Kobayasi, asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Sung et al. (2010, 2011; culture characters of the type, applications), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: EFCC 6279.

Shiraiia Henn. 1900, *Shiraiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. bambusicola* Henn., asexual morph coelomycetous, parasitic, terrestrial, China and Japan, see Lumbsch and Huhndorf (2010; outline), Liu et al. (2013b; morphology, nomenclature, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; 2017; outline), cultures and sequences are available, ex-epitype strains of type: GZAAS2.0708 = CGMCC 3.15200 = MFLUCC 13-0408; holotype of type: Shirai (B!), epitype of type: GZAAS K-2002.

Shivomyces Hosag. 2004, *Ascomycota* genera *incertae sedis*, two species, type: *S. ligustri* Hosag. & Kamar., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Shrungabeeja V.G. Rao & K.A. Reddy 1981, *Tetrapsphaeriaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *S. vadrajensis* V.G. Rao & K.A. Reddy, hyphomycetous, saprobes, on bamboo (*Bambusa*) and other plants (*Begonia*, *Melicope*), Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Ariyawansa et al. (2015a; new species, phylogeny), cultures and sequences are unavailable.

Sibirina G.R.W. Arnold 1970, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *S. fungicola* G.R.W. Arnold, hyphomycetous, sexual morphs unknown, mycoparasites, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available.

Siemaszkoa I.I. Tav. & T. Majewski 1976, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, seven species, type: *S. flexa* (T. Majewski) I.I. Tav. & T. Majewski, asexual morph unknown, biotrophic on

Coleoptera (Ptinidae), Europe, Southern America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sigarispora Thambug. & K.D. Hyde 2015, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *S. ravennica* (Tibpromma, Camporesi & K.D. Hyde) Thambug. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Italy and Japan, six species, see Thambugala et al. (2015b; morphology, phylogeny), Li et al. (2016a; new species), Tibpromma et al. (2017b; new species), holotype of type: MFLU 14-0692, ex-type strains of type: MFLUCC 14-0005.

Sigleria Hirooka, Tanney & Seifert 2015, *Spiromastigaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *S. carmichaelii* Hirooka, Tanney & Seifert, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Vietnam, see Hirooka et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype of type: DAOM 571575.

Sigmatomyces Sacc. & P. Syd. 1913, *Ascomycota* genera *incertae sedis*, one species, type: *S. bakeri* Sacc., Syd. & P. Syd., sexual morph unknown, hyphomycetous, saprobes, Philippines, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sigmoidea J.L. Crane 1968, *Phaeotrichaceae*, *Phaetrichales*, *Dothideomycetes*, type: *S. prolifera* (R.H. Petersen) J.L. Crane, hyphomycetous, freshwater, on wood and leaves, worldwide, Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available. Marine *Sigmoidea* (= *Halosigmoidea*) species group in *Corollospora*, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, see Jones et al. (2009b; morphology, phylogeny), Jones et al. (2015) and Réblová et al. (2016c).

Sigridia Tehler 1993, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, six species, type: *S. californica* (Tuck.) Tehler, lichenized (only one lichenicolous), terrestrial, tropical to subtropical with a preference for coastal habitats with a Mediterranean climate, see Ertz and Tehler (2011; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ertz et al. (2015b, c; morphology, phylogeny), Lücking et al. (2017a; classification), lectotype and ex-type strain of type: Bolander 129 (FH).

Sillia P. Karst. 1873, *Sydiellaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *S. ferruginea* (Pers.) P. Karst., asexual morphs unknown, saprobes, terrestrial, Europe, America, see Lumbsch and Huhndorf (2010; outline), Kruys and Castlebury (2012; phylogeny),

Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2017; placement), Tibpromma et al. (2017b; new species), cultures available for the type: CBS 126567.

Simonyella J. Steiner 1902, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *S. variegata* J. Steiner, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification, treated as *Roccellaceae*), sequences are available.

Simplicillium W. Gams & Zare 2001, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, nine species, type: *S. lanosonivum* (J.F.H. Beyma) Zare & W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, Europe, North America, see Seifert et al. (2011; morphology), Liu and Cai (2012; new species, China), Wijayawardene et al. (2012, 2017; outline), Nonaka et al. (2013b; new species, Japan), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Sinodidymella J.Z. Yue & O.E. Erikss. 1985, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes* families *incertae sedis*, five species, type: *S. verrucosa* (Petr.) J.Z. Yue & O.E. Erikss., see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Sinotrichium Juan Wang, S.Q. Xu & Strongman 2010, *Ascomycota* genera *incertae sedis*, one species, type: *S. chironomidarum* Juan Wang, S.Q. Xu & Strongman, asexual morph unknown, from the hindgut of a chironomid larva, China, see Wang et al. (2010b; taxonomy), holotype of the type: XIN-4, cultures and sequences are unavailable.

Siphula Fr. [nom. cons.] 1824, *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, 26 species; type: *S. ceratites* (Wahlenb.) Fr., lichenized, see Valadbeigi and Sipman (2010; new records), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Resl et al. (2015; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Siphulastrum Müll.Arg. 1889, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, four species; type: *S. triste* Müll. Arg., lichenized see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ekman et al. (2014; phylogeny), Jørgensen and Andersen (2015; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Siphulella Kantvilas, Elix & P. James 1992, *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type: *S. coralloidea* Kantvilas, Elix & P. James, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Sipmania Egea & Torrente 1994, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *S. peltata* Egea & Torrente, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Sipmaniella Kalb 2009, *Megalosporaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *S. sulphureofusca* (Fée) Kalb, lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Sirastachys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, seven species, type: *S. phaeospora* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22460, CBS 100155.

Sirenophila Søchting, Arup & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, seven species, type: *S. gintarasii* (S.Y. Kondr. & Kärnefelt) Arup, Frödén & Søchting, lichenized, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), Søchting et al. (2016; new species, southern hemisphere), sequences are available.

Sirexipula Bubák 1907, *Ascomycota* genera *incertae sedis*, one species, type: *S. kabatiana* Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirococcus Preuss 1855, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, three species, type: *S. strobilinus* Preuss, coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Crous et al. (2012f, 2016a, b; phylogeny, new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Sirocyphis Clem. 1909, *Ascomycota* genera *incertae sedis*, one species, type: *S. kabatiana* Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirogloea Petr. 1923, *Ascomycota* genera *incertae sedis*, one species, type: *S. euonymi* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Siroligniella Naumov 1926, *Ascomycota* genera *incertae sedis*, one species, type: *S. salicicola* Naumov, coelomycetous, sexual morph unknown, saprobes, terrestrial, former USSR, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirophoma Höhn. 1917, *Ascomycota* genera *incertae sedis*, three species, type: *S. singularis* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Siroplacodium Petr. 1940, *Ascomycota* genera *incertae sedis*, six species, type: *S. atrum* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, China, see Wijayawardene et al. (2012, 2017; outline), Zhao and Zhao (2012; new species, China), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Siropleura Petr. 1934, *Ascomycota* genera *incertae sedis*, one species, type: *S. pulchella* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Siroscyphellina Petr. 1923, *Ascomycota* genera *incertae sedis*, two species, type: *S. arundinacea* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Austria, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirosperma Syd. & P. Syd. 1917, *Ascomycota* genera *incertae sedis*, two species, type: *S. hypocrella* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Papua New Guinea, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirospphaera Syd. & P. Syd. 1913, *Ascomycota* genera *incertae sedis*, two species, type: *S. botryosa* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirosporium Bubák & Serebrian. 1912, *?Dothideomycetes* genera *incertae sedis*, c. 26 species, type: *S. antenniforme* (Berk. & M.A. Curtis) Bubák & Serebrian., hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are available.

Silosporonaemella Naumov 1951, *Ascomycota* genera *incertae sedis*, one species, type: *S. strobilina* Naumov, coelomycetous, sexual morph unknown, saprobes, terrestrial, Russia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirothecium P. Karst. 1887, *Ascomycota* genera *incertae sedis*, three species, type: *S. saepiarium* P. Karst., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirothyriella Höhn. 1910, *Ascomycota* genera *incertae sedis*, two species, type: *S. pinastri* Höhn. ex Sacc., coelomycetous, saprobes, terrestrial, Europe, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Sirothyrium Syd. & P. Syd. 1916, *Ascomycota* genera *incertae sedis*, one species, type: *S. taxi* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirozythia Höhn. 1904, *Ascomycota* genera *incertae sedis*, two species, type: *S. rosea* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sirozythiella Höhn. 1909, *Ascomycota* genera *incertae sedis*, one species, type: *S. sydowiana* (Sacc.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sitochora H.B.P. Upadhyay 1964, *Ascomycota* genera *incertae sedis*, one species, type: *S. ellipsospora* H.B.P. Upadhyay, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sivanesania W.H. Hsieh & Chi Y. Chen 1996, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, one species, type: *S. rubi* W.H. Hsieh & Chi Y. Chen, asexual morph unknown, saprobes, terrestrial, Taiwan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Sivanesaniella Gawande & D.K. Agarwal 2004, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. prunicola* Gawande & D.K. Agarwal, asexual morph

unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013; did not accept as a genus in *Venturiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Skottsbergiella Petr. 1927, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *S. diaporthoides* Petr., asexual morph unknown, saprobes, terrestrial, Chile, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Skyathea Spooner & Dennis 1986, ?*Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *S. hederae* Spooner & Dennis, asexual morph unknown, saprobe of wood, Skye Island, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype of the type: K 32267, cultures and sequences are unavailable.

Skyttea Sherwood, D. Hawksw. & Coppins 1981, *Cordieritidaceae*, *Leotiomycetes* families *incertae sedis*, 29 species, type: *S. nitschkei* (Körb.) Sherwood, D. Hawksw. & Coppins lichenicolous, widespread, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Suija et al. (2014, 2015b; phylogeny), Jaklitsch et al. (2016a; position), sequences are available, holotype of type: L 910.195-559.

Skyttella D. Hawksw. & R. Sant. 1988, ?*Cordieritidaceae*, *Leotiomycetes* families *incertae sedis*, one species, type: *S. muelleri* (Willey) D. Hawksw. & R. Sant., on lichens, terrestrial, Europe and North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; position), sequences are available, holotype of type: FH.

Slimacomycetes Minter 1986, *Ascomycota* genera *incertae sedis*, two species, type: *S. monosporus* (W.B. Kendr.) Minter, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Slopeiomyces Klaubauf, Lebrun & Crous 2014, *Magnaportheaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *S. cylindrosporus* (D. Hornby, Slope, Gutter. & Sivan.) Klaubauf, Lebrun & Crous, asexual morph hyphomycetous, saprobes, terrestrial, UK, see Klaubauf et al. (2014; morphology, phylogeny), Maharachchikumbura et al. (2015b; outline), ex-type culture of type: CBS 609.75.

Smaragdiniseta L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one

species, type: *S. bisetosa* (V.G. Rao & de Hoog) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), ex-type of the type: CBS 459.82.

Smardaea Svrček 1969, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, eight species, type: *S. amethystina* (W. Phillips) Svrček, asexual morph unknown, saprobes, terrestrial, north temperate, see Van Vooren (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Smarodsia Raity. & Vimba 2006, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *S. stollii* Raity. & Vimba, asexual morph unknown, from soil, terrestrial, Latvia, see Lumbsch and Huhndorf (2010; outline), holotype of the type: RIG 6493, cultures and sequences are unavailable.

Smeringomyces Thaxt. 1908, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *S. anomalus* (Thaxt.) Thaxt., asexual morph unknown, biotrophic on Coleoptera (Staphilinidae), USA, Europe, Japan, Trinidad, Sumatra, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sodiomyces Grum-Grzhimaylo, Debets & Bilanenko 2013, *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*, two species, type: *S. alkalinus* (Bilanenko & M. Ivanova) A.A. Grum-Grzhimaylo, A.J.M. Debets & E.N. Bilanenko, asexual morph unknown, saprobes, terrestrial, worldwide, see Grum-Grzhimaylo et al. (2013a; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Solanella Vaňha 1910, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, one species, type: *S. rosea* Vaňha, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification, treated as *Pezizales*), cultures and sequences are unavailable.

Soleella Darker 1967, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, seven species, type: *S. striiformis* (Darker) Darker, asexual morph unknown, saprobes, terrestrial, North America, China, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Solenopezia Sacc. 1889, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *S. solenia* Sacc., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline),

Kirk et al. (2013; genus accepted), cultures and sequences are unavailable for type.

Solenopsora A. Massal. 1855, *Catillariaceae*, *Rhizocarpales*, *Lecanoromycetes*, 20 species, type: *S. requienii* A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Van Den Boom (2010; new records, Spain), Farkas et al. (2011; distribution Hungary), Kinalioglu and Aptroot (2011; new records, Turkey), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), Yazici and Aslan (2016; new species, Turkey), sequences are available.

Solicorynespora R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, 27 species, type: *S. zapatensis* R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see McKenzie (2010; new species), Seifert et al. (2011; morphology), Ma et al. (2012d, 2016a; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2014a; new species), cultures and sequences are unavailable.

Solitaria Arup, Søchting & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *S. chrysophthalma* (Degelius) Arup, Søchting & Frödén, asexual morph unknown, lichenized, terrestrial, see Arup et al. (2013; taxonomy), Gaya et al. (2015; phylogeny), Lücking et al. (2017a; classification); sequences are

Soloacrospora W.B. Kendr. & R.F. Castañeda 1991, *Ascomycota* genera *incertae sedis*, two species, type: *S. flagellisetosa* W.B. Kendr. & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Soloacrosporiella Crous & M.J. Wingf. 2015, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. acaciae* Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, Malaysia, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22223, CPC 24871.

Solorina Ach. 1808, *Peltigeraceae*, *Peltigerales*, *Lecanoromycetes*, ten species, type: *S. crocea* (L.) Ach., lichenized, Asia, Europe, North America, see Gagunashvili et al. (2009; molecular), Björk (2010; new record), Lumbsch and Huhndorf (2010; outline), Wedin et al. (2011; molecular), Grube et al. (2012; symbiosis), Lisov et al. (2012; ecology), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; molecular), Centeno et al. (2016; metabolic profiling), Magain et al. (2016;

molecular), Lücking et al. (2017a; classification), sequences are available.

Solorinaria (Vain.) Gyeln., 1935, *Lichinaceae*, *Lichinales*, *Lichinomycetes*, one species, type: *S. fuscata* (Vain.) Gyeln., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Aptroot and Iqbal (2012; checklist), Kirk et al. (2013; genus accepted), Brackel (2013; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Solosympodiella Matsush. 1971, *Ascomycota* genera *incertae sedis*, eight species, type: *S. clavata* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Solotermilospora Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *S. caffra* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: Matsushima Fungus Collection, Kobe, 5A089, cultures and sequences are unavailable.

Sonderhenia H.J. Swart & J. Walker 1988, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *S. eucalyptorum* (Hansf.) H.J. Swart & J. Walker coelomycetous, sexual morph mycosphaerella-like, two species, plant pathogens on *Eucalyptus*, terrestrial, see Crous et al. (2013a; morphology, phylogeny), Videira et al. (2017; typification), Wijayawardene et al. (2017; outline), holotype DAR 45719, isotype IMI 280474 (sexual morph), holotype WARI 2007 (aseexual morph); representative cultures CBS 120220 = CPC 12553, CPC 12554–12555.

Soosiella Hujslová & M. Kolařík 2014, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *S. minima* Hujslová & M. Kolařík, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Czech Republic, see Hujslová et al. (2014; phylogeny), holotype and ex-type of the type: PRM 922619, CBS 136257.

Sordaria Ces. & De Not. 1863, *Sordariaceae*, *Sordariales*, *Sordariomycetes*, twelve species, type: *S. fimicola* (Roberge ex Desm.) Ces. & De Not., asexual morph unknown, terrestrial, saprobes, coprophilous, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; phylogeny), cultures and sequences are available.

Sorocybe Fr. 1849, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, three species, type: *S. resinae* (Fr.) Fr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology, genus accepted), Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; did not accept genus), cultures and sequences are available.

Sorokina Sacc. 1892, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, six species, type: *S. microspora* (Berk.) Sacc., asexual morph unknown, saprobes, terrestrial, pantropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sorokinella J. Fröhl. & K.D. Hyde 2000, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, type: *S. appendiculosa* J. Fröhl. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Hong Kong, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Sowerbyella Nannf. 1938, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, eight species, type: *S. radiculata* (Sowerby) Nannf., asexual morph unknown, saprobes, terrestrial, Europe, China, see Zhuang (2009; new species), Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), sequences are available.

Spadicoides S. Hughes 1958, *Sordariomycetes* genera *incertae sedis*, c. 40 species, type: *S. bina* (Corda) S. Hughes 1958, hyphomycetous, sexual morph unknown, phialophora-like and selenosporella-like synanamorphs, saprobes, terrestrial and aquatic, cosmopolitan, see Shenoy et al. (2010; polyphyly, affinities with *Dothideomycetes* and *Sordariomycetes*), Seifert et al. (2011; morphology), Ma et al. (2012a, 2012b, 2014a, new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Heredia et al. (2015, new species), sequences are available but lacking for the type.

Sparria Ertz & Tehler 2011, *Opegraphaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *S. cerebriformis* (Egea & Torrente) Ertz & Tehler, lichenized, see Californian coast, Baja California and Europe, see Ertz and Tehler (2011; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Sparticola Phukhams., Ariyaw., Camporesi & K.D. Hyde 2016, *Sporomiaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *S. junci* Phukhams., Camporesi & K.D. Hyde asexual morph hyphomycetous, saprobes, terrestrial, Italy, see Phukhamsakda et al. (2016a; taxonomy, phylogeny), Thambugala et al. (2017b; new species, phylogeny), Wijayawardene et al. (2017; outline), Karunarathna et al. (2017; new species, phylogeny), holotype and ex-type of the type: MFLU 15-1405, MFLUCC 15-0030.

Spataporthe Bronson, Klymiuk, Stockey & Tomescu 2013, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *S. taylorii* Bronson, Klymiuk, Stockey &

Tomescu, asexual morph unknown, fossil fungi, see Bronson et al. (2013; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Spathaspora N.H. Nguyen, S.O. Suh & M. Blackw. 2006, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, ten species, type: *S. passalidarum* N.H. Nguyen, S.O. Suh & M. Blackw., asexual reproduction is by multilateral budding, saprophytic, associated with wood boring insect larvae, possibly worldwide, see Lumbsch and Huhndorf (2010; outline), *S. passalidarum* and other species in the clade are noted for fermentation of D-xylose, which is important for fermentation of biomass to fuel ethanol, many species in this genus represent asexual taxa originally classified in the genus *Candida*, but which have been transferred to the genus *Spathaspora* as new combinations, cultures and sequences are available.

Spathularia Pers. 1797, *Cudoniaceae*, *Rhytismatales*, *Leotiomycetes*, twelve species, type: *S. flava* Pers., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ge et al. (2014; DNA, phylogeny), cultures and sequences are available.

Spathulospora A.R. Caval. & T.W. Johnson 1965, *Spathulosporaceae*, *Spathulosporales*, *Sordariomycetes*, five species, type: *S. phycophila* A.R. Caval. & T.W. Johnson, spermatiophores or asexual morph hyphomycetous, biotrophic on marine red alga, temperate to Antarctic sea of Southern hemisphere, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: (R.E. Norris, 3892).

Speerschneidera Trevis. 1861, *Leprocaulaceae*, *Leprocaulales*, *Lecanoromycetes*, one species, type: *S. euploca* (Tuck.) Trevis., lichenized, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Lecanorales, genera *incertae sedis*), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); sequences are available.

Spegazzinia Sacc. 1880, *Apiosporaceae*, *Amphisphaeriales*, *Sordariomycetes*, nine species, type: *S. ornata* Sacc., hyphomycetous, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Whitton et al. (2012a; morphology with two types of conidia), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanan et al. (2017a; phylogeny, evolution), Thambugala et al. (2017b, phylogeny, new species), cultures and sequences are available.

Speiropsis Tubaki 1958, *Dothideomycetes* genera *incertae sedis*, nine species, type: *S. pedatospora* Tubaki,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, North and South America, see Prihatini et al. (2008; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Pratibha et al. (2016; DNA, phylogeny), cultures available for the type: GUFCC 18008.

Spencermartinsia A.J.L. Phillips, A. Alves & Crous 2008, *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, nine species, type: *S. viticola* (A.J.L. Phillips & J. Luque) A.J.L. Phillips, A. Alves & Crous, asexual morph coelomycetous, saprobes, weakly pathogens, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Liu et al. (2012b; phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Phillips et al. (2013; phylogeny), Abdollahzadeh et al. (2014; new species), Slippers et al. (2014; phylogeny), Wijayawardene et al. (2014c, 2016b; outline, taxonomy), Dissanayake et al. (2016b; taxonomy), Yang et al. (2016d; reduced *Spencermartinsia* to synonymy under *Dothiorella* but here we treat *Spencermartinsia* as a distinct genus pending further studies), holotype and ex-type of the type: LISE 95177, CBS 117009.

Spencermartiniella Péter, Dlauchy, Tornai-Lehoczki, M. Suzuki & Kurtzman 2011, *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, four species, type: *S. europaea* G. Péter, Dlauchy, Tornai-Leh., M. Suzuki & Kurtzman, cell division is by multilateral budding, saprophytic, rotted wood, Europe, South America, see Péter et al. (2011; taxonomy), cultures and sequences are available.

Spermatoloncha Speg. 1908, *Ascomycota* genera *incertae sedis*, one species, type: *S. maticola* Speg., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spermochaetella Cif. 1954, *Ascomycota* genera *incertae sedis*, one species, type: *S. tecumae* (Chardón & Toro) Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spermopora R. Sprague 1948, *Ascomycota* genera *incertae sedis*, nine species, type: *S. subulata* (R. Sprague) R. Sprague, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spermosporella Deighton 1969, *Ascomycota* genera *incertae sedis*, nine species, type: *S. aggregata* Deighton,

hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Asia, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spermotrichum Kuntze 1898, *Ascomycota* genera *incertae sedis*, one species, type: *S. pulchellum* (Speg.) Kuntze, coelomycetous, sexual morph unknown, saprobes, terrestrial, Paraguay, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphacelia Lév. 1827, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, seven species, type: *S. segetum* Lév., hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Pažoutová et al. (2008; DNA, phylogeny), Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Sphaerellopsis Cooke 1883, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *S. quercuum* Cooke, asexual morph coelomycetous, pathogens, lichenicolous, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c, 2017; outline), Trakunyingcharoen et al. (2014; new species, phylogeny), Crous et al. (2016a, b; new species, phylogeny), cultures and sequences are available but lacking for the type.

Sphaerellothecium Zopf 1897, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, type: *S. araneosum* (Rehm) Zopf, c. 30 species, asexual morph unknown, terrestrial, on lichens, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mycosphaerellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), van den Boom (2016; new species), cultures and sequences are unavailable.

Sphaerostromella Bubák 1916, *Ascomycota* genera *incertae sedis*, one species, type: *S. pteridina* (Sacc. & Roum.) Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphaerothyrium Bubák 1916, *Ascomycota* genera *incertae sedis*, two species, type: *S. filicinum* Bubák, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphaerocordyceps Kobayasi 1981, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *S. palustris* (Berk.) Kobayasi, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: CUP 067858, cultures and sequences are unavailable.

Sphaerodes Clem. 1909, *Coronophorales* genera *incertae sedis*, *Sordariomycetes*, 13 species, type: *S. episphaerium* (W. Phillips & Plowr.) Clem., asexual morph unknown, saprobes, terrestrial, worldwide, see Vujanovic and Goh (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Sphaerodothella C.A. Pearce & K.D. Hyde 2001, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *S. danthoniae* (McAlpine) C.A. Pearce & K.D. Hyde, saprobes, Australia, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Sphaerodothis (Sacc. & P. Syd.) Shear 1909, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, 20 species, type: *S. arengae* (Racib.) Shear ex Theiss. & Syd., saprobes, South East Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Sphaerognomiella Naumov & Kusnezowa 1952, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. cornicola* Naumov & Kusnezowa, asexual morph unknown, saprobes, terrestrial, former USSR, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Sphaerographium Sacc. 1884, *Chaetomellaceae*, *Helotiales*, *Leotiomycetes*, twelve species, type: *S. squarrosum* Riess, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Minnis et al. (2010; new species, DNA), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; accepted as valid genus), cultures available but lacking for the type.

Sphaeromma H.B.P. Upadhyay 1964, *Ascomycota* genera *incertae sedis*, two species, type: *S. mazosiae* H.B.P. Upadhyay, coelomycetous, sexual morph unknown, lichenicolous, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphaeronaemella P. Karst. 1884 (= *Gabarnaudia* Samson & W. gams 1974 *fide* Rébllová et al. 2016c), *Microascales* genera *incertae sedis*, *Sordariomycetes*, ten species, type: *S. helvillae* (P. Karst.) P. Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see

Lumbsch and Huhndorf (2010; outline), de Beer et al. (2013; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Rébllová et al. (2016c; nomenclature), Wijayawardene et al. (2017; outline), sequences are available.

Sphaerophoma Petr. 1924, *Ascomycota* genera *incertae sedis*, two species, type: *S. brencklei* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphaerophoropsis Vain. 1890, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *S. stereocauloides* Vain., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Sphaerophorus Pers. 1794, *Sphaerophoraceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *S. coralloides* Pers., asexual morph unknown, fruticose, terrestrial, saxicolous, see Wedin et al. (2009a; taxonomy), Hua-Jie et al. (2010; metal studies), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), McMullin and Arsenault (2016; new records), Root et al. (2014; climate indicator), Chamizo et al. (2016, spatial distribution), Lücking et al. (2017a; classification), sequences are available.

Sphaeropsis Sacc. 1880 (= *Phaeobotryosphaeria* Speg. 1908 *fide* Phillips et al. 2013; Wijayawardene et al. 2014c; Rossman et al. 2015b), *Botryosphaeriaceae*, *Botryosphaerales*, *Dothideomycetes*, five species, type: *S. visci* (Alb. & Schwein.) Sacc., coelomycetous, sexual morph formerly known in *Phaeobotryosphaeria*, saprobes, endophytes, terrestrial, cosmopolitan, see Phillips et al. (2008, 2013; taxonomy, phylogeny), Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Slippers et al. (2013; phylogeny), Wijayawardene et al. (2014c, 2016b; outline, taxonomy, phylogeny, nomenclature), Doilom et al. (2016; species in *Tectona*), Rossman et al. (2015b; nomenclature), Dissanayake et al. (2016b; taxonomy), neotype and ex-neotype of the type: CWU (MYC) AS 2271 (MBT176099), CBS 122527.

Sphaerosoma Klotzsch 1839, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, three species, type: *S. fuscescens* Klotzsch, asexual morph unknown, hypogeous, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphaerospora (Svrček) Svrček & Kubička 1961, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, three species, type: *S. brunnea* (Alb. & Schwein.) Svrček & Kubička, asexual morph hyphomycetous, hypogeous, terrestrial,

worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Sánchez et al. (2014b; asexual reproduction, DNA), cultures and sequences are available for the type.

Sphaerosporium Schwein. 1832, *Onygenales* genera *incertae sedis*, two species, type: *S. lignatile* Schwein., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North and South America, see Seifert et al. (2011; morphology), Ropars et al. (2012; DNA, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Sphaerostilbella (Henn.) Sacc. & D. Sacc. 1905 (= *Gliocladium* Corda 1840 *fide* Rossman et al. 2013b; Maharachchikumbura et al. 2016b), *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *S. lutea* (Henn.) Sacc. & D. Sacc., asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Sphaerozone Zobel 1854, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, four species, type: *S. tulasnei* Zobel, asexual morph unknown, hypogeous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphaerulina Sacc. 1878, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, 66 species, type: *S. myriadea* (DC.) Sacc., asexual morph coelomycetous, asexual morph septoria-like, saprobes or plant pathogens, terrestrial, freshwater or marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Crous et al. (2011e; taxonomy), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Mycosphaerellaceae*), Verkley et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Sphaerulomyces Marvanová. 1977, *Ascomycota* genera *incertae sedis*, one species, type: *S. coralloides* Marvanová, hyphomycetous, saprobes, Czechoslovakia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sphaleromyces Thaxt. 1894, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *S. lathrobii* Thaxt., asexual morph unknown, biotrophic on Coleoptera (Staphylinidae), Europe, America, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spheconisca (Norman) Norman 1876, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, twenty species, type: *S. hypocrita* Norman, asexual morph unknown,

lichenized, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Sphinctrina Fr. 1825, *Sphinctrinaceae*, *Mycocaliciales*, *Eurotiomycetes*, ten species, type: *S. turbinata* (Pers.) De Not., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Muniz et al. (2013; new species), Tibell et al. (2014; new species, DNA), cultures available for the type: AFTOL-ID 1721.

Sphinctrinopsis Woron. 1927, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *S. pertusariae* Woron., lichenized, see Gaya et al. (2012; taxonomy), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are unavailable.

Spicellum Nicot & Roquebert 1976, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *S. roseum* Nicot & Roquebert, hyphomycetous, sexual morph unknown, saprobes, terrestrial, France, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Spilodochium Syd. 1927, *Ascomycota* genera *incertae sedis*, four species, type: *S. vernoniae* Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, Neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spilonema Bornet 1856, *Coccocarpiaceae*, *Peltigerales*, *Lecanoromycetes*, five species, type: *S. paradoxum* Bornet, lichenized, see Lumbsch and Huhndorf (2010; outline), Anderson and Neily (2012; new records), Kirk et al. (2013; genus accepted), Medeiros et al. (2014; ecology), Spribille et al. (2014b; evolution), McMullin et al. (2015; new records), Prieto et al. (2015; new record), Lücking et al. (2017a; classification), sequences are available.

Spilopodia Boud. 1885, *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, four species, type: *S. nervisequa* (Pers.) Boud., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Spilopodiella E. Müll. 1889, *Drepanopezizaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *S. arxii* E. Müll., asexual morph unknown, saprobes, terrestrial, Switzerland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spinulosphaeria Sivan. 1974, *Chaetosphaerellaceae*, *Coronophorales*, *Sordariomycetes*, 14 species, type:

S. thaxteri (Pat.) Sivan., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Mugambi and Huhndorf (2010; new species, phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline, phylogeny), cultures and sequences are available but lacking for the type.

Spinulospora Deighton 1973, *Ascomycota* genera *incertae sedis*, four species, type: *S. pucciniiphila* Deighton, hyphomycetous, sexual morph unknown, fungicolous, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spiralum J.L. Mulder 1975, *Ascomycota* genera *incertae sedis*, two species, type: *S. helicosporum* J.L. Mulder, hyphomycetous, sexual morph unknown, fungicolous, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spirodecospora B.S. Lu, K. D. Hyde & W. H. Ho 1998, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, two species, type: *S. bambusicola* B.S. Lu et al., asexual morph unknown, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Spirographa Zahlbr. 1903 (= *Pleospilis* Clem. 1909 *fide* Jaklitsch et al. 2016a), *Ostropomycetidae* genera *incertae sedis*, *Lecanoromycetes*, four species, type: *S. spiralis* (Müll. Arg.) Zahlbr., asexual morph unknown, saprobes, lichenicolous, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Helotiales*), Jaklitsch et al. (2016a; classification), Flakus and Kukwa (2012; new species), cultures and sequences are unavailable.

Spiromastigoides Doweld 2013, *Spiromastigaceae*, *Onygenales*, *Eurotiomycetes*, 17 species, type: *S. warcupii* (Kuehn & G.F. Orr) Doweld, asexual morph unknown, ?saprobes, from soil, terrestrial, Australia, see Index Fungorum (2013), culture available for the type: AFTOL-ID 430.

Spiromastix Kuehn & G.F. Orr 1962, *Spiromastigaceae*, *Onygenales*, *Eurotiomycetes*, eleven species, type: *S. warcupii* Kuehn & G.F. Orr, asexual morph hyphomycetous, from dung, soil, animal, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Doweld (2013; nomenclatural), Hirooka et al. (2016; new species), holotype and ex-type of the type: NRRL A-10679, CBS 576.63.

Spiroplana Voglmayr, M.J. Park & H.D. Shin 2011, *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, two

species, type: *S. centripetala* Voglmayr, M.J. Park & H.D. Shin, hyphomycetous, sexual morph unknown, parasitic, terrestrial, Korea, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype and ex-type of the type: KUS-F22135, KACC42611.

Spirosphaera Beverw. 1953, *Helotiales* genera *incertae sedis*, nine species, type: *S. floriformis* Beverw, hyphomycetous, sexual morph unknown, on decaying birch leaves in water, see Seifert et al. (2011; morphology), Voglmayr et al. (2011; taxonomy), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Splanchnonema Corda 1829, *Pleomassariaceae*, *Pleosporales*, *Dothideomycetes*, 37 species, type: *S. pustulatum* Corda, asexual morph unknown, saprobes, terrestrial, Europe, North America, China, India, see Lumbsch and Huhndorf (2010; outline, sexual morph), Zhang et al. (2012e; accepted as a genus in *Pleomassariaceae*, phylogeny), Zhao and Zhao (2012; new species, China), Hyde et al. (2013; accepted as a genus in *Pleomassariaceae*, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type.

Splanchospora Lar.N. Vassiljeva 1998, *Ascomycota* genera *incertae sedis*, one species, type: *S. ampullacea* (Pers.) Lar.N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, north temperate zone, see Honda et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spondylocladiella Linder 1934, *Ascomycota* genera *incertae sedis*, two species, type: *S. botrytoides* Linder, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spondylocladiopsis M.B. Ellis 1963, *Ascomycota* genera *incertae sedis*, two species, type: *S. cupulicola* M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, India, see Pratibha et al. (2009; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Spooneromyces T. Schumach. & J. Moravec 1989, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, five species, type: *S. laeticolor* (P. Karst.) T. Schumach. & J. Moravec, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Hansen et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Van Vooren (2014; new species), cultures and sequences are available.

Sporastatia A. Massal. 1854, *Sporastatiaceae*, *Rhizocarpales*, *Lecanoromycetes*, four species, type: *S. testudinea* (Ach.) A. Massal., lichenized, saxicolous, worldwide temperate, see Lumbsch and Huhndorf (2010; outline), Bendiksby and Timdal (2013; phylogenetic placement), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogenetic placement), Lücking et al. (2017a; classification), sequences are available.

Sporendocladia G. Arnaud ex Nag Raj & W.B. Kendr. 1975, *Microascales* genera *incertae sedis*, *Sordariomycetes*, seven species, type: *S. castaneae* G. Arnaud ex Nag Raj & W.B. Kendr., asexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Kiyuna et al. (2012; DNA), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Sporendonema Desm. 1827, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, nine species, type: *S. casei* Desm., coelomycetous, sexual morph unknown, on cheese and fungi, terrestrial, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Sporaphplus H.B.P. Upadhyay 1964, *Ascomycota* genera *incertae sedis*, one species, type: *S. rondoniensis* H.B.P. Upadhyay, asexual morph unknown, lichenized, terrestrial, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Sporidesmajora Batzer & Crous 2010, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. pennsylvaniensis* Batzer & Crous, hyphomycetous, sexual morph unknown, one species, saprobes,? pathogens, terrestrial, USA, see Yang et al. (2010; morphology, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type culture of type: CBS H-20326, CPC 16112.

Sporidesmiella P.M. Kirk 1982, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 32 species, type: *S. claviformis* P.M. Kirk, hyphomycetous, sexual morph unknown, saprobes, on dung, terrestrial, worldwide, see Braun and Heuchert (2010; new species), Seifert et al. (2011; morphology), Ma et al. (2012e; new species, China), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Monteiro et al. (2014a; new species, Brazil), Heredia et al. (2015; new species, Mexico), Tian et al. (2015; excluded *Melanommataceae* from and placed in *Didymosphaeriaceae*), Ma (2016; new species) holotype of the type: IMI 246982, cultures and sequences are available but lacking for the type.

Sporidesmina Subram. & Bhat 1989, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. malabarica* Subram. & Bhat, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: NN43118.

Sporidesmioides J.F. Li, Phook. & K.D. Hyde 2016, *Torulaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. thailandica* J.F. Li, Phook. & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Thailand, see Li et al. (2016c; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: MFLU14-0827, cultures and sequences are available.

Sporidesmiopsis Subram. & Bhat 1989, *Ascomycota* genera *incertae sedis*, one species, type: *S. malabarica* Subram. & Bhat, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, South America, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline, treated as *Ascomycota* genera *incertae sedis*), cultures and sequences are unavailable.

Sporidesmiopsis Subram. & Bhat, Kavaka 1989, *Ascomycota* genera *incertae sedis*, three species, type: *S. malabarica* Subram. & Bhat, hyphomycetous, sexual morph unknown, saprobes, Asia, Caribbean, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; did not list), cultures and sequences are unavailable, need generic revision.

Sporidesmium Link 1809 (= *Podoconis* Boedijn 1933 *fide* Species Fungorum 2017), *Sporidesmiaceae*, *Diaportheomycetidae* families *incertae sedis*, *Sordariomycetes*, around 330 species, type: *S. atrum* Link, hyphomycetous, saprobes, mycoparasites, terrestrial and aquatic, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2016; new species, phylogeny), Mena-Portales et al. (2016; new species), Su et al. (2016b; phylogeny of *Sporidesmiaceae*), cultures and sequences are available.

Sporoctomorpha J.V. Almeida & Sousa da Câmara 1903, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. magnoliae* J.V. Almeida & Sousa da Câmara, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Sporodictyon A. Massal. 1852, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, five species, 38 species,

type: *S. schaeerianum* A. Massal., lichenized, see Dillman et al. (2014; new records, Alaska), Lücking et al. (2017a; classification); sequences are available.

Sporodophoron Frisch, Y. Ohmura, Ertz & G. Thor 2015, Arthoniaceae, Arthoniales, Arthoniomycetes, four species, type: *S. gossypinum* Frisch, Y. Ohmura & G. Thor, lichenized, see Frisch et al. (2015), Lücking et al. (2017a; classification), sequences are available.

Sporoglena Sacc., 1894, Ascomycota genera *incertae sedis*, one species, type: *S. velutina* Sacc., hyphomycetous, sexual morph unknown, Papua New Guinea, see Seifert et al. (2011; mentioned as a doubtful genus), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Sporomega Corda 1842, Rhytismataceae, Rhytismatales, Leotiomycetes, seven species, type: *S. degenerans* (Fr.) Corda, asexual morph unknown, saprobes, terrestrial, North America, Europe, see Lumbsch and Huhndorf (2010; outline), Lantz et al. (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), sequences are available.

Sporopachydermia Rodrigues de Miranda 1978, Saccharomycetales genera *incertae sedis*, *Saccharomycetes*, five species, type: *S. lactativora* Rodrigues de Miranda, asexual reproduction is by multilateral budding, saprophytic, in cacti, seawater, tree exudates, waste lagoon, clinical, Europe, North America, Antarctica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Sporophagomyces K. Pöldmaa & Samuels 1999, Hypocreaceae, Hypocreales, Sordariomycetes, three species, type: *S. chrysostomus* (Berk. & Broome) K. Pöldmaa & Samuels., asexual morph unknown, fungicolous, terrestrial, worldwide, see Jaklitsch et al. (2008; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Sporophiala P. Rag. Rao 1970, Ascomycota genera *incertae sedis*, three species, type: *S. prolifica* P. Rag. Rao, hyphomycetous, sexual morph unknown, Asia, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Sporopodiopsis Sérusiaux 1997, Pilocarpaceae, Lecanorales, Lecanoromycetes, two species, type: *S. sipmanii* Sérus., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Sporopodium Mont. 1851, Pilocarpaceae, Lecanorales, Lecanoromycetes, 24 species, type: *S. leprieurii* Mont., lichenized, see Lumbsch and Huhndorf (2010; outline), Weerakoon et al. (2012a; taxonomy), Kirk et al. (2013; genus accepted), Rashmi and Rajkumar (2015; new report), Lücking et al. (2017a; classification), sequences are available.

Sporormia De Not. 1845, Sporormiaceae, Pleosporales, Dothideomycetes, 29 species, type: *S. fimetaria* De Not., asexual morph unknown, saprobes on dung, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; accepted genus), Hyde et al. (2013; phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Sporoschisma Berk. & Broome 1847 (= *Melanochaeta* E. Müll., Harr & Sulmont 1969 *fide* Maharachchikumbura et al. 2016b; Réblová et al. 2016c), Chaetosphaeriaceae, Chaetosphaeriales, Sordariomycetes, type: *M. hemipsila* (Berk. & Broome) E. Müll., Harr & Sulmont., hyphomycetous, sexual morphs formerly named in *Melanochaeta*, saprobes, terrestrial (wood), Europe, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Réblová et al. (2016c; nomenclature), Luo et al. (2016a; new species), Yang et al. (2016b; new species, phylogeny), cultures available for type: SMH 3251, SMH 2125, AR4690, GKM1247, GKM1112.

Sporoschismopsis Hol.-Jech. & Hennebert 1972 (= *Porosphaerellopsis* Samuels & E. Müll. 1982 *fide* Réblová et al. 2016c), Reticulascaceae, Glomerellales, Sordariomycetes, eight species, type: *S. moravica* Hol.-Jech. & Hennebert, asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Zhao and Zhao (2012; new species, China), Kirk et al. (2013; genus accepted), Réblová (2014; new species, holomorph, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Réblová et al. (2016c; nomenclature), cultures and sequences are available but lacking for the type.

Sporostigma Grube 2001, Arthoniales genera *incertae sedis*, Arthoniomycetes, one species, type: *S. melaspora* (Tuck.) Grube, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Sporothrix Hektoen & C.F. Perkins 1901, Ophiostomataceae, Ophiostomatales, Sordariomycetes, c. 51 species, type: *S. schenckii* Hektoen & C.F. Perkins, terrestrial, worldwide, including saprobes associated with plant material and soil, and important clinically-relevant species causing sporotrichosis in humans and other vertebrates, see Madrid et al. (2009, 2010b; geographical

distribution, new species), Musvuugwa et al. (2015; new species), Rodrigues et al. (2015a, 2016; molecular diagnostics of pathogenic species, new species), de Beer et al. (2016; phylogeny, outline, new combinations), de Errasti et al. (2016, new species).

Sporotretophora Whitton, K.D. Hyde & McKenzie 2012, Ascomycota genera *incertae sedis*, one species, type: *S. punctata* Whitton, K.D. Hyde & McKenzie, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Whitton et al. (2012a; taxonomy), cultures and sequences are unavailable.

Spumatoria Massee & E.S. Salmon 1901, Ophiostomataceae, Ophiostomatales, Sordariomycetes, one species, type: *S. longicollis* Massee & E.S. Salmon, asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Squamaria Poelt, 1958, Squamarinaceae, Lecanorales, Lecanoromycetes, species, sub species, varieties, forma species, type: *S. gypsacea* (Sm.) Poelt, asexual morph, undetermined, terricolous, saxicolous, widespread, see Dietrich and Buerger-Meyer (2010; taxonomy, rare occurrences in surroundings), Lumbsch and Huhndorf (2010; outline), Miralles-Mellado et al. (2011; indicator species), Jorge-Villar et al. (2011; Raman spectroscopy study), Bevan et al. (2012; transplant experiments), Cansaran-Duman and Halici (2012; biochemistry and antimicrobial study), Kirk et al. (2013; genus accepted), Bastida et al. (2014; ecology), Martellos et al. (2014; ecology, species distribution) Delgado Baquerizo et al. (2015, chemistry), Lange and Belnap (2016; soil crust studies), Lücking et al. (2017b; classification); sequences are available.

Squamella S. Hammer 2001, Cladoniaceae, Lecanorales, Lecanoromycetes, one species, type: *S. spumosa* S. Hammer, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Squamotubera Henn. 1903, Xylariaceae, Xylariales, Sordariomycetes, one species, type: *S. leratii* Henn., see Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; genus accepted).

Squamulea Arup, Søchting & Frödén, 2013, Teloschistaceae, Teloschistales, Lecanoromycetidae, five species, type: *S. subsoluta* (Nylander) Arup, Søchting & Frödén, asexual morph unknown, lichenized, terrestrial, Antarctica, see Arup et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2014, 2016; revised taxonomy), Søchting et al. (2014a; phylogeny species), Lücking et al. (2017a; classification), sequences are available.

Stachybotryella Ellis & Barthol. 1902, Ascomycota genera *incertae sedis*, three species, type: *S. repens* Ellis & Barthol., hyphomycetous, sexual morph unknown, America, see Seifert et al. (2011; mentioned as a doubtful genus), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Stachybotryna Tubaki & T. Yokoy. 1971, Ascomycota genera *incertae sedis*, five species, type: *S. columaris* Tubaki & T. Yokoy., hyphomycetous, sexual morph unknown, saprobes, ?pathogens, terrestrial, Asia, Caribbean, Europe, see Seifert et al. (2011; morphology), Heredia et al. (2012; new species, Mexico), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), culture available for an unidentified species.

Stachybotrys Corda 1837, Stachybotryaceae, Hypocreales, Sordariomycetes, c. 60 species, type: *S. chartarum* (Ehrenb.) S. Hughes, hyphomycetous, sexual morph available, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species), Wang et al. (2015e; new species, revision, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Lombard et al. (2016; phylogeny, revision, sexual morph), cultures and sequences are available.

Stachylidium Link 1809, Plectosphaerellaceae, Glomerellales, Sordariomycetes, two species, type: *S. bicolor* Link, hyphomycetous, sexual morph unknown, pathogens, nematophagous, terrestrial, worldwide, see Réblová et al. (2011; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures sequences are available for the type: DAOM 226658.

Stagonopatella Petr. 1927, Ascomycota genera *incertae sedis*, one species, type: *S. aeruginosa* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stagonopsis Sacc. 1884, Ascomycota genera *incertae sedis*, four species, type: *S. pallida* (Berk. & M.A. Curtis) Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stagonospora (Sacc.) Sacc. 1884, Massarinaceae, Pleosporales, Dothideomycetes, 220 species, type: *S. paludosa* (Sacc. & Speg.) Sacc., coelomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Wijayawardene et al. (2012, 2014c, 2017;

outline), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2013; new species, phylogeny), Crous et al. (2014c, d, 2015a; DNA, new species, phylogeny), Tanaka et al. (2015; new species, phylogeny), Thambugala et al. (2017b, phylogeny, new species), neotype and ex-neotype of the type: CBS H-21317, S601 = CBS 135088.

Stagonosporina Tassi 1902, Ascomycota genera *incertae sedis*, one species, type: *S. sparti* Tassi, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stagonosporopsis Died. 1912, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, 22 species, type: *S. hortensis* (Sacc. & Malbr.) Petr., coelomycetous, with sexual morph, saprobes, ?pathogens, terrestrial, cosmopolitan, see Vaghefi et al. (2012; species associated with ray blight disease of Asteraceae), Chen et al. (2015b; taxonomy, phylogeny), Hyde et al. (2016; sexual morph), Tibpromma et al. (2017b; new species), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Stagonostromella Petr. & Syd. 1927, Ascomycota genera *incertae sedis*, one species, type: *S. sparti* Tassi, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Staheliella Emden 1974, Ascomycota genera *incertae sedis*, two species, type: *S. nodosa* Emden, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Staibia Bat. & Peres 1966, *Leptopeltidaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *S. connari* Bat. & Peres, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, accepted as a genus in *Leptopeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Stalagmochaetia Cif. & Bat. 1963, Ascomycota genera *incertae sedis*, two species, type: *S. basilisca* Bat., Cif. & Nascim., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stamnaria Fuckel 1870 (= *Titaeospora* Bubák 1916 *fide* Johnston et al. 2014b), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, five species, type: *S. persoonii* (Moug.) Fuckel, asexual morph formerly known in *Titaeospora*, saprobes, terrestrial, worldwide, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Baral and Haelewaters (2015; DNA), Wijayawardene et al. (2017; outline), cultures and sequences are available but lacking for the type.

Stanhughesia Constant. 1989, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, four species, type: *S. linnaeae* Constant., sexual morph unknown, epiphytes, epiphyllous, hyphomycetous, terrestrial, Sweden, South Africa, see Seifert et al. (2011; morphology), Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stanimwardia B. Sutton 1971, *Extremaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *S. breviuscula* B. Sutton, coelomycetous, sexual morph unknown, saprobes, pathogens, Australia, see Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2014c, 2017; outline, taxonomy), cultures and sequences are available but lacking for the type.

Stanjehughesia Subram. 1992 (= *Umbrinosphaeria* Réblová 1999 *fide* Réblová et al. 2016c), *Sordariomycetes* genera *incertae sedis*, 16 species, type: *S. hormisciooides* (Corda) Subram., hyphomycetous, sexual morph formerly named in *Umbrinosphaeria*, saprobes, terrestrial, worldwide, see Delgado (2008; new species, USA), Marincowitz et al. (2008; new species), Ma et al. (2011c; new species, China), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), et al. (2016c; nomenclature), cultures and sequences are available but lacking for the type.

Stanjemonium W. Gams, O'Donnell, Schroers & M. Chr. 1999, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *S. grisellum* W. Gams, Schroers & M. Chr., hyphomycetous, sexual morph unknown, from soil, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 655.79.

Staphylotrichum J. Mey. & Nicot 1957, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, four species, type: *S. coccosporum* J.A. Mey. & Nicot, sexual morph unknown, saprobes, from soil, terrestrial, Africa, Asia, Europe, see Seifert et al. (2011; morphology), Nonaka et al. (2012; new species, Japan phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for the type.

Starbaeckia Rehm ex Starbäck 1899, *Phaciidaeae*, *Phacidiiales*, *Leotiomycetes*, one species, type:

S. pseudotryblidiooides Rehm & Starbäck, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Starmeria Y. Yamada, Higashi, Ando & Mikata 1997, *Phaffomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, ten species, type: *S. amethionina* (Starmer, Phaff, M. Miranda & M.W. Miller) Y. Yamada, Higashi, S. Ando & Mikata, cell division is by multilateral budding, saprophytic, in cacti, tunnels of wood boring insects, slime fluxes of trees, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Starmerella Rosa & Lachance 1998, *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, 25 species, type: *S. bombicola* C.A. Rosa & Lachance, cell division is by multilateral budding, saprophytic, in honey, fruit juice concentrate, flowers, various beetles, bees, worldwide, see Kurtzman et al. (2010; phylogeny, revision), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), most of the species in this genus represent asexual forms originally classified in the genus *Candida* thus, they will be transferred to *Starmerella* as new combinations, cultures and sequences are available.

Staurolemma Körb. 1867, *Pannariaceae*, *Peltigerales*, *Lichenomycetes*, three species, type: *S. dalmaticum* Körb., asexual morph unknown, lichenized, terrestrial, see Wedin et al. (2009b; taxonomy), Jørgensen (2010; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Bendiksby et al. (2014; long-distance dispersal), Dixon et al. (2014; mapping), Ekman et al. (2014; taxonomy), Otálora et al. (2013, 2014; revised generic classification), Tapper et al. (2014; endemics on granite), Boutabia et al. (2015; floristic), Paoli et al. (2015; indicator species), Lücking et al. (2017a; classification), sequences are available.

Stauronema (Sacc.) Syd., P. Syd. & E.J. Butler 1916, *Ascomycota* genera *incertae sedis*, four species, type: *S. cruciferum* (Ellis) Syd., P. Syd. & E.J. Butler, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stauronematopsis Abbas, B. Sutton & Ghaffar 2002, *Ascomycota* genera *incertae sedis*, one species, type: *S. sojae* (Uecker & Kulik) Abbas, B. Sutton & Ghaffar, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Staurophoma Höhn. 1907, *Ascomycota* genera *incertae sedis*, one species, type: *S. panici* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Hong Kong, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Staurothele Norman 1853, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 40 species, type: *S. clopima* (Wahlb.) Th. Fr., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Gueidan et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Stearophora L. Mangin & Viala 1905, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. radicicola* L. Mangin & Viala, asexual morph unknown, saprobes, terrestrial, North Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Steganopycnis Syd. & P. Syd. 1916, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. oncospermatis* Syd. & P. Syd., see Lumbsch and Huhndorf (2010; outline), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Sordariomycetes* genera *incertae sedis*), the genus was not accepted as a genus in Kirk et al. (2008, 2013).

Stegasphaeria Syd. & P. Syd. 1916, *Mesnieraceae*, *Dothideomycetes* families *incertae sedis*, five species, type: *S. pavonina* Syd. & P. Syd., see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Mesnieraceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Stegobolus Mont. 1845, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 15 species, type: *S. berkeleyanus* Mont., asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Sipman et al. (2012; key, Costa Rica), Rivas Plata et al. (2012a; phylogeny), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); sequences are available.

Stegonsporiopsis Van Warmelo & B. Sutton 1981, *Ascomycota* genera *incertae sedis*, one species, type: *S. cenangiooides* (Ellis & Rothr.) Van Warmelo & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Hong Kong, South America, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stegonsporium Corda 1827, *Stilboporaceae*, *Diaporthales*, *Sordariomycetes*, eight species, type: *S. pyriforme* (Hoffm.) Corda, asexual morph coelomycetous, saprobes, terrestrial, worldwide, see

Voglmayr and Jaklitsch (2008, 2014; phylogeny, taxonomy, holomorph), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2016b, 2017; taxonomy, outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Stegophorella Petr. 1947, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. lagerheimii* Petr., asexual morph unknown, saprobes, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Stegothyrium Höhn. 1918, *Dothideomycetes* genera *incertae sedis*, two species, type: *S. denudans* (Rehm) Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wu et al. (2011) and Hyde et al. (2013) did not accept as *Microthyriaceae*, Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Steinera Zahlbr. 1906, *Koerberiaceae*, *Peltigerales*, *Lecanoromycetes*, four species, type: *S. molybdoplaca* (Nyl.) Zahlbr., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Spribille and Muggia (2013; evolutionary), Lücking et al. (2017a; classification), sequences are available.

Steineropsis T. Sprib. & Muggia 2010, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, one species, type: *S. alaskana* T. Sprib. & Muggia, lichenized, Europe, Africa, mostly along the Atlantic coast, see Spribille et al. (2010; taxonomy), Spribille and Muggia (2013; phylogenetics), Schneider et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Steinia Körb., 1873, *Aphanopsidaceae*, *Pezizomycotina* families *incertae sedis*, three species, type: *S. luridescens* Körb., asexual morph unknown, lichenized, terrestrial, European distribution, see Lumbsch and Huhndorf (2010; outline), Kärnefelt et al. (2012; systematic, biodiversity), Printzen et al. (2012; partial sequencing), Kirk et al. (2013; genus accepted), Czarnota and Hernik (2014; ecology), Yatsyna and Motiejūnaitė (2015; new contributions), Kondratyuk et al. (2015b; molecular phylogeny), Lücking et al. (2017a; classification); sequences are available.

Stellarangia Frödén, Arup & Søchting 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *S. elegantissima* (Nyl.) Frödén, Arup & Søchting, lichenized, see Arup et al. (2013; taxonomy), Gaya et al. (2015; adaptations, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Stellatospora Tad. Ito & Nakagiri 1994, *Sordariaceae*, *Sordariales*, *Sordariomycetes*, one species, type: *S. terricola* Tad. Ito & Nakagiri asexual morph unknown, saprophytic, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: IFO H-12166, ex-type cultures of type: NBRC 32597.

Stellifraga Alstrup & Olech 1993, *Ascomycota* genera *incertae sedis*, one species, type: *S. cladoniicola* Alstrup & Olech, asexual morph unknown, saprobes, terrestrial, Spitsbergen, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stellomyces Morgan-Jones, R.C. Sinclair & Eicker 1987, *Ascomycota* genera *incertae sedis*, two species, type: *S. suidafricanus* Morgan-Jones, R.C. Sinclair & Eicker, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stellopeltis Bat. & A.F. Vital 1959, *Ascomycota* genera *incertae sedis*, two species, type: *S. philodendricola* (Bat.) Bat. & A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stellosetifera Matsush. 1996, *Sordariomycetes* genera *incertae sedis*, one species, type: *S. malaysiana* Matsush., asexual morph unknown, saprobes, terrestrial, Malaysia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Stellospora Alcorn & B. Sutton 1984, *Ascomycota* genera *incertae sedis*, two species, type: *S. appendiculellae* Alcorn & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stellothyriella Bat. & Cif. 1959, *Ascomycota* genera *incertae sedis*, two species, type: *S. graminis* (Wehm.) Bat. & Cif., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stemmatomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type: *S. anoplischii* (Thaxt.) Thaxt., asexual morph unknown, biotrophic on Coleoptera (Elateridae), USA, see Lumbsch

and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stemphylium Wallr. (= *Pleospora* Rabenh. ex Ces. & De Not. 1863 *fide* Wijayawardene et al. 2014c), *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, 56 species, type: *S. botryosum* Wallr., hyphomycetous, sexual morph formerly known in *Pleospora*, saprobes or pathogens in terrestrial habitats, worldwide, see Seifert et al. (2011; morphology), Zhang et al. (2012e; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Ariyawansa et al. (2015c; taxonomy, phylogeny), Rossman et al. (2015b; nomenclature), Crous et al. (2016b; new species, phylogeny), sequences are available.

Stenella Syd. 1930, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, c. 140 species, type: *S. araguata* Syd., hyphomycetous, saprobes, pathogens, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2014; phylogeny), cultures and sequences are available.

Stenhammarella Hertel 1967, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *S. turgida* (Ach.) Hertel, asexual morph unknown, lichenized, terrestrial, Europe, Central Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures available for the type: Stetur3861.

Stenocarpella Syd. & P. Syd. 1917, *Diaporthaceae*, *Diaporthales*, *Sordariomycetes*, two species, type: *S. zeae* Syd. & P. Syd., coelomycetous, sexual morph unknown, pathogens, terrestrial, worldwide, see Wicklow et al. (2011; bioactive metabolites), Wijayawardene et al. (2012, 2016b; phylogeny, taxonomy), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for the type: NRRL13609.

Stenocephalopsis Chamuris & C.J.K. Wang 1998, *Ascomycota* genera *incertae sedis*, one species, type: *S. subalutacea* (Peck) Chamuris & C.J.K. Wang, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stenocladiella Marvanová & Descals 1987, *Ascomycota* genera *incertae sedis*, one species, type: *S. neglecta* (Marvanová & Descals) Marvanová & Descals, hyphomycetous, sexual morph unknown, saprobes, aquatic, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stenocybe Nyl. ex Körb. 1855, *Mycocaliciaceae*, *Mycocaliciales*, *Eurotiomycetes*, 14 species, type: *S. byssacea* (Fr.) Nyl. ex Körb., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Stenospora Deighton 1969, *Ascomycota* genera *incertae sedis*, one species, type: *S. uredinicola* Deighton, hyphomycetous, sexual morph unknown, fungicolous, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stephanocyclos Hertel 1983, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *S. henssenianus* Hertel, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Schmull et al. (2011; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Stephanonectria Schroers & Samuels 1999, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *S. keithii* (Berk. & Broome) Schroers & Samuels, asexual morph unknown, saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Stephanotheca Syd. & P. Syd. 1914, *Dothideomycetes* genera *incertae sedis*, four species, type: *S. micromera* Syd. & P. Syd., asexual morph unknown, foliar epiphytes, terrestrial, India, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jayawardene et al. (2014; morphology), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable, holotype of the species S-F6581.

Stephembruneria R.F. Castañeda 1988, *Ascomycota* genera *incertae sedis*, one species, type: *S. elegans* R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stephensia Tul. & C. Tul. 1845 (= *Densocarpa* Gilkey 1954; = *Elderia* McLennan 1961 *fide* Species Fungorum 2017), *Pezizales* genera *incertae sedis*, *Pezizomycetes*, seven species, type: *S. bombycinia* (Vittad.) Tul., asexual morph unknown, hypogeous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wang et al. (2016c; DNA, phylogeny), sequences are available.

Stereocaulon Hoffm. 1795. *Stereocaulaceae*, *Lecanorales*, *Lecanoromycetes*, 140 species, type:

S. paschale (L.) Hoffm., asexual morph unknown, lichenized, terrestrial, worldwide, see Khare et al. (2009; taxonomy), Morita et al. (2009; antimitotic study), Seo et al. (2009b; bioactivity study), Wedin et al. (2009b; molecular), Fryday (2010; taxonomy), Hafellner et al. (2010; biodiversity), Huang (2010; environment), Lumbsch and Huhndorf (2010; outline), Melechin (2010; taxonomy), Goel et al. (2011; antifungal study), Hansen (2011; taxonomy), Ismed et al. (2012; chemistry), Bhattacharai et al. (2013; bioactivity study), Kim et al. (2013; antimicrobial study), Kirk et al. (2013; genus accepted), Rincon-Espitia and Mateus (2013; taxonomy), Singh et al. (2013b; physiology), Högnabba et al. (2014; phylogeny), Meunier et al. (2014; environment), Nayaka and Saxena (2014; physiology), Oset and Kukwa (2014; typification), Ranković et al. (2014; bioactivity), Oset (2015; taxonomy), Doering et al. (2016; taxonomy), Kossowska and Szczepańska (2016; taxonomy), Lücking et al. (2017a; classification), Shukla et al. (2016; amino acids), sequences are available.

Stereocrea Syd. & P. Syd. 1917, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *S. schizostachyi* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, paleotropics, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), holotype of the type: CUP 067858, cultures and sequences are unavailable.

Stereosphaeria Kirschst. 1939, *Sordariomycetes* genera *incertae sedis*, two species, type: *S. phloeophila* Kirschst., widespread in temperate regions, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a), one sequence is available.

Sterigmatobotrys Oudem. 1886, *Pleurotheciaeae*, *Pleurotheciales*, *Sordariomycetes*, three species, type: *S. elata* (Sacc.) Oudem., hyphomycetous, sexual morph unknown, saprobes, terrestrial, north temperate, see and Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures available but lacking for the type.

Stevensonula Petr. 1952, *Ascomycota* genera *incertae sedis*, one species, type: *S. ciliata* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, New Caledonia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stichomyces Thaxt. 1901, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *S. conosomatis* Thaxt., asexual morph unknown, biotrophic on Coleoptera (Staphylinidae), Europe, Africa, USA,

Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sticta (Schreb.) Ach. 1803, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, c. 200 species, type: *S. sylvatica* (Huds.) Ach., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Moncada and Lücking (2012; new species), Moncada et al. (2013b; new species), Lücking et al. (2017a; classification), cultures and sequences are available.

Stictis Pers. 1800, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, four species, type: *S. radiata* (L.) Pers., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures available for type: AFTOL-ID 398, MW6493, P164, F907E267.

Stictographa Mudd 1861, *Asterinales* genera *incertae sedis*, *Dothideomycetes*, one species, type: *S. lentiginosa* (Lyell ex Leight.) Mudd, lichenicolous, cosmopolitan, see Ertz and Diederich (2015; taxonomy, phylogeny), sequences are available.

Stictopatella Höhn. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *S. euonymi* (Desm.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stictosepta Petr. 1964, *Ascomycota* genera *incertae sedis*, one species, type: *S. cupularis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stictospora Cif., 1957, *Stictidaceae*, *Ostropales*, *Ostropomycetes*, one species, type: *S. oleae* (De Not.) Cif., asexual morph unknown, lichenized, terrestrial, see Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stigmatellina Bat. & H. Maia 1960, *Ascomycota* genera *incertae sedis*, one species, type: *S. epilobii* Bat. & H. Maia, coelomycetous, sexual morph unknown, saprobes, terrestrial, Belgium, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stigmatochroma Marbach, 2000, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, nine species, type: *S. epimarta* (Nyl.) Marbach, asexual morph unknown, lichenized, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Cavalcante (2012; comparative studies), Gaya et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Giralt et al. (2014; new species), Logesh et al. (2015; floristics), Mohabe et al. (2015; new species), Prieto and Wedin

(2016; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Stigmatodiscus Voglmayr & Jaklitsch 2016, *Stigmatodiscaceae*, *Stigmatodiscales*, *Dothideomycetes*, two species, type: *S. enigmaticus* Voglmayr & Jaklitsch, asexual morph coelomycetous, saprobes, Europe, see Voglmayr et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype: WU 35914, cultures and sequences are available.

Stigmatodothis Syd. & P. Syd. 1914, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. palawanensis* Syd. & P. Syd., asexual morph unknown, saprobes, terrestrial, Philippines, Australia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Micropeltidaceae*), Hyde et al. (2013; did not recognize this genus as in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline) cultures and sequences are unavailable.

Stigmatomyces H. Karst. 1869, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 145 species, type: *S. muscae* (Peyr.) H. Karst., asexual morph unknown, biotrophic on Diptera, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossi and Leonardi (2012, most recent species descriptions Africa), sequences are available but lacking for the type.

Stigmatophragmia Tehon & G.L. Stout 1929, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. sassafrasicola* Tehon & G.L. Stout, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Micropeltidaceae*), Hyde et al. (2013; did not recognize this genus as in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Stigmatula (Sacc.) Syd. & P. Syd. 1901, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, ten species, type: *S. sutherlandiae* (Kalchbr. & Cooke) Syd. & P. Syd., saprobes, temperate, see Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Stigmea Fr. 1836, *Ascomycota* genera *incertae sedis*, one species, type: *S. submaculans* Mont., saprobes, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stigmella Lév. 1842, *Ascomycota* genera *incertae sedis*, 28 species, type: *S. dryina* (Corda) Lév., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b; outline, taxonomy), Kirk et al. (2013; genus accepted), Wang et al. (2013e; new species, China), cultures and sequences are unavailable.

Stigmidiump Trevis. 1860, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *S. schaeferi* (A. Massal.) Trevis., lichenicolous or lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Stigmochora Theiss. & Syd. 1914, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, twelve species, type: *S. controversa* (Starbäck) Theiss. & Syd., saprobes, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Stigmopeltis Syd. 1927, *Ascomycota* genera *incertae sedis*, two species, type: *S. roupaleae* Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Central America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stilbella Lindau 1900, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *S. fimetaria* (Pers.) Lindau, hyphomycetous, sexual morph unknown, from soil and dung, saprobes, worldwide, see Mazzaferro et al. (2010; secondary metabolites), Seifert et al. (2011; morphology), Summerbell et al. (2011; phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Stilbellula Boedijn 1951, *Ascomycota* genera *incertae sedis*, one species, type: *S. pallida* Boedijn, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stilbocrea Pat. 1900, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *S. dussii* Pat., asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Sung et al. (2008; DNA), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Stilbodendron Syd. & P. Syd. 1916, *Ascomycota* genera *incertae sedis*, one species, type: *S. camerunense* Syd. & P. Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stilbophyoxylon Henn. 1902, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, ten species, type: *S. moelleri* Henn., asexual morph unknown, terrestrial, saprobes, subtropical, see Hsieh et al. (2010; DNA, phylogeny), Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Stilbopeziza Speg. 1908, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *S. yerbae* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stilbophoma Petr. 1942, *Ascomycota* genera *incertae sedis*, one species, type: *S. microspora* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, India, Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stilbospora Pers. 1794 (= *Prosthecium* Fresen. 1852 *fide* Rossman et al. 2015a; Voglmayr and Jaklitsch 2014), *Stilbosporaceae*, *Diaporthales*, *Sordariomycetes*, 20 species, type: *S. macroisperma* Pers., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Voglmayr and Jaklitsch (2008, 2014; phylogeny, taxonomy, holomorph), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2016b, 2017; taxonomy, outline, phylogeny), cultures and sequences are available.

Stioclettia Dennis 1975, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. luzulina* Dennis, asexual morph unknown, saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Stirtonia A.L. Sm. 1926, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, 25 species, type: *S. obvallata* A.L. Sm., lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are available.

Stirtoniella D.J. Galloway, Hafellner & Elix 2005, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *S. kelica* (Stirt.) D.J. Galloway, Hafellner & Elix, asexual morph unknown, lichenized, terrestrial, Australasia, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Stomatogene Theiss. 1917, *Perisporiopsidaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *S. agaves* (Ellis & Everh.) Theiss., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Stomatogenella Petr. 1955, *Sordariomycetes* genera *incertae sedis*, two species, type: *S. mirabilis* Petr.,

asexual morph unknown, saprobes, terrestrial, North Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Stomiopeltis Theiss. 1914, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, 25 species, type: *S. aspersa* (Berk.) Theiss., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for type.

Stomiopeltopsis Bat. & Cavalc. 1963, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, two species, type: *S. linacearum* Bat. & Cavalc., asexual morph unknown, saprobes, terrestrial, Brazil, New Mexico, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Stomiotheca Bat. 1959, *Micropeltidaceae*, *Microthyriales*, *Dothideomycetes*, two species, type: *S. amazonensis* Bat., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Wu et al. (2011; new combination), Hyde et al. (2013; accepted as a genus in *Micropeltidaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Stoufferia Kovács & Trappe 2011, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, four species, type: *S. longii* (Gilkey) Kovács & Trappe, asexual morph unknown, hypogeous, terrestrial, Argentina, USA, see Kovács et al. (2011b; taxonomy, phylogeny), sequences and cultures are unavailable.

Strangospora Körb. 1860, *Strangosporaceae*, *Pezizomycotina* families *incertae sedis*, nine species, type: *S. pinicola* (A. Massal.) Körb., asexual morph unknown, lichenized, terrestrial distribution, Lumbsch and Huhndorf (2010; outline), Stepanchikova et al (2011; new records), Kirk et al. (2013; genus accepted), Miądlikowska et al. (2014a; phylogeny), Rogers et al. (2015; lichen monitoring), Lücking et al. (2017a; classification), Urbanavichus and Ismailov (2016; new records), sequences are available.

Strasseria Bres. & Sacc. 1902, *Ascomycota* genera *incertae sedis*, one species, type: *S. carpophila* Bres. & Sacc., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Strasseriopsis B. Sutton & Tak. Kobay. 1970, *Ascomycota* genera *incertae sedis*, one species, type:

S. tsugae (Tak. Kobay.) B. Sutton & Tak. Kobay., coelomycetous, sexual morph unknown, saprobes, terrestrial, Japan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stratiphoromyces Goh & K.D. Hyde 1998, Ascomycota genera *incertae sedis*, one species, type: *S. brunneisporus* Goh & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stratisporaella Hafellner 1979, *Patellariaceae*, *Patellariales*, *Dothideomycetes*, one species, type: *S. episemoides* (Nyl.) Hafellner, asexual morph unknown, lichenicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), cultures and sequences are unavailable.

Strattonia Cif. 1954, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, nine species, type: *S. tetraspora* (R. Stratton) Cif., asexual morph unknown, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariales* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Streimannia G. Thor 1991, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, one species, type: *S. varieseptata* G. Thor., lichenized, terrestrial, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); sequences unavailable.

Streimanniella S.Y. Kondr., Kärnfeldt, A. Thell, Elix & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, four species, type: *S. michelagoensis* (Elix, S.Y. Kondr. & Kärnfeldt) S.Y. Kondr., Kärnfeldt, A. Thell, Elix, J. Kim, A.S. Kondr. & Hur, lichenized, see Kondratyuk et al. (2015c; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Strelitziana Arzanlou & Crous 2006, *Strelitzianaceae*, *Chaetothyriales*, *Eurotiomycetes*, seven species, type: *S. africana* Arzanlou & Crous, hyphomycetous, sexual morph unknown, saprobes, cosmopolitan, see Zhang et al. (2009b; new species), Crous et al. (2010b, 2015e; new species, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Streptotinia Whetzel 1945 (= *Streptobotrys* Hennebert 1973 *fide* Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *S. arisaematis* Whetzel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Striabotrys L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, six species, type: *S. eucylindrospora* (D.W. Li) L. Lombard & Crous, asexual morph hyphomycetous, saprobes, from soil, terrestrial, cosmopolitan, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), ex-type of the type: MUCL 9483.

Striaticonidium L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *S. cinctum* (Corda) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), epitype and ex-epitype of the type: CBS H-22471, CBS 932.69.

Striatodecospora D.Q. Zhou, K.D. Hyde & B.S. Lu 2000, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. bambusae* D.Q. Zhou, K.D. Hyde & B.S. Lu., asexual morph unknown, on dead bamboo culm, see Lumbsch and Huhndorf (2010; outline), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are unavailable.

Striatosphaeria Samuels & E. Müll. 1979, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *S. codinaeaphora* Samuels & E. Müll., asexual morph unknown, saprobes, Central and South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available for the type.

Strickeria Körb. 1865, *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, ten species, type: *S. kochii* Körb., asexual morph coelomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Sordariales* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; taxonomy, phylogeny, holomorph), cultures available for the type: C149, C138.

Strigopodia Bat. 1957, *Euantennariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *S. piceae* Bat., asexual morph unknown, saprobes,

terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; accepted as a genus in *Euantennariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Strigula Fr. 1823, *Strigulaceae*, *Strigulales*, *Dothideomycetes*, one species, type: *S. smaragdula* Fr., asexual morph unknown, lichenized, foliicolous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Nelsen et al. (2011b; DNA, phylogeny), Hyde et al. (2013; accepted as a genus in *Strigulaceae*), Jayalal et al. (2013b; DNA, South Korea), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Lücking et al. (2017a; classification), sequences are available.

Striosphaeropsis Verkley & Aa 1997, *Ascomycota* genera *incertae sedis*, one species, type: *S. mirabilis* Verkley & Aa, coelomycetous, sexual morph unknown, saprobes, terrestrial, Papua New Guinea, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Strobiloscypha N.S. Weber & Denison 1995, *Chorioactidaceae*, *Pezizales*, *Pezizomycetes*, two species, type: *S. keliae* N.S. Weber & Denison, asexual morph unknown, saprobes, terrestrial, Montenegro, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Perić et al. (2013; new species, DNA, phylogeny), sequences are available.

Stromaster Höhn. 1930, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, twelve species, type: *S. tuberculatus* (Pat.) Höhn., saprobes, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Stromatella Henssen 1989, *Lichinaceae*, *Lichinales*, *Lichinomycetes*, one species, type: *S. bermudana* (Riddle) Henssen, lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Stromatinia (Boud.) Prill. 1907, *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *S. rapulum* (Bull.) Boud., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence available for the type: Z73801 (ITS).

Stromatocrea W.B. Cooke 1952, *Ascomycota* genera *incertae sedis*, one species, type: *S. cerebriformis* W.B. Cooke, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura

et al. (2015b, 2016b; outline, treated as a genus in *Bionectriaceae* or *Hypocreaceae*), cultures and sequences are unavailable.

Stromatographium Höhn. 1907 (= *Fluviostroma* Samuels & E. Müll. 1980; = *Stromatostilbella* Samuels & E. Müll. 1980 *fide* et al. 2016c; Species Fungorum 2017), *Trichosphaeriaceae*, *Trichosphaeriales*, *Sordariomycetes*, one species, type: *S. stromaticum* (Berk.) Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), et al. (2016c; nomenclature), cultures and sequences are unavailable.

Stromatonectria Jaklitsch & Voglmayr 2011, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *S. caraganae* (Höhn.) Jaklitsch & Voglmayr, asexual morph coelomycetous, saprobes, terrestrial, Austria, see Jaklitsch and Voglmayr (2011; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; phylogeny), ex-epitype of the type: CBS 127387.

Stromatoneurospora S.C. Jong & E.E. Davis. 1973, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. phoenix* S.C. Jong & E.E. Davis., tropical, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Senanayake et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), a sequence is available.

Stromatopogon Zahlbr. 1897, *Ascomycota* genera *incertae sedis*, three species, type: *S. baldwinii* Zahlbr., coelomycetous, sexual morph unknown, lichenicolous, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stromatopycnis A.F. Vital 1956, *Ascomycota* genera *incertae sedis*, one species, type: *S. rosetum* A.F. Vital, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stromatoseptoria Quaedvl., Verkley & Crous 2013, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *S. castaneicola* (Desm.) Quaedvl., Verkley & Crous, coelomycetous, sexual morph unknown, pathogens, terrestrial, Austria, France, Netherlands, one species, see Quaedvlieg et al. (2013; morphology, phylogeny), holotype PC0084574, reference specimens: CBS H-21200, culture CBS 102322; requires epitypification.

Stromatostysanus Höhn. 1919, *Ascomycota* genera *incertae sedis*, three species, type: *S. caprifoliorum* (Desm.) Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Stromatothecia D.E. Shaw & D. Hawksw. 1971, *Odontotemataceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *S. nothofagi* D.E. Shaw & D. Hawksw, on bark, Papua New Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, holotype of type: K(M)-IMI 151852a.

Stromatothelium Trevis. 1861, *Pyrenulales* genera *incertae sedis*, *Eurotiomycetes*, four species, need typification, hyphomycetous, sexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Strongylothallus Bat. & Cif. 1959, *Ascomycota* genera *incertae sedis*, one species, type: *S. scutatus* Bat., Cif. & C.A.A. Costa, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Strossmayeria Schulzer 1881 (= *Pseudospiropes* M.B. Ellis 1971 *fide* Johnston et al. 2014b), *Helotiales* genera *incertae sedis*, *Leotiomycetes*, 16 species, type: *S. rackii* Schulzer, asexual morph hyphomycetous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hustad and Miller (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Johnston et al. (2014b) nomenclature), Wijayawardene et al. (2017; outline), cultures and sequences are available but lacking for the type.

Stuartella Fabre 1879, *Dothideomycetes* genera *incertae sedis*, three species, type: *S. formosa* Fabre, asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Stygiomyces Coppins & S.Y. Kondr. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *S. galliformis* Coppins & S.Y. Kondr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Tasmania, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: HO 34310, cultures and sequences are unavailable.

Stylaspergillus B. Sutton, Alcorn & P.J. Fisher 1982, *Ascomycota* genera *incertae sedis*, one species, type: *S. laxus* B. Sutton, Alcorn & P.J. Fisher, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Stylobothis Arx & E. Müll. 1975, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, two species, type: *S. puccinoides* (DC.) Arx & E. Müll., asexual morph unknown, saprobes, terrestrial, Europe, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; accepted as a genus in *Dothideaceae*), Wijayawardene et al. (2014c; outline), sequences are available.

Stylolectria Höhn. 1915, *Nectriaceae*, *Hypoocreales*, *Sordariomycetes*, five species, type: *S. applanata* Höhn., sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Gräfenhan et al. (2011; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Subbaromyces Hesselt. 1953, *Ophiostomataceae*, *Ophiostomatales*, *Sordariomycetes*, two species, type: *S. splendens* Hesselt., asexual morph unknown, saprobes, North America, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available.

Suberoteratosphaeria Quaedvl. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *S. suberosa* (Crous, F.A. Ferreira, Alfenas & M.J. Wingf.) Quaedvl. & Crous, asexual morph unknown, pathogens, Australia, Brazil, Uruguay, see Quaedvlieg et al. (2014; morphology, phylogeny), holotype and ex-type culture of the type: PREM 51082, CPC 515.

Subglobosporium Doilom & K.D. Hyde 2016, *Pseudocoleodictyosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. tectonae* Doilom & K.D. Hyde, saprobes, terrestrial, see Doilom et al. (2017; taxonomy), ex-type: MFLUCC 12-0393, MKT 066, ICMP 21258; holotype of type: MFLU 15-3560.

Subhysteropycnis Wedin & Hafellner 1998, *Ascomycota* genera *incertae sedis*, one species, type: *S. maculiformans* Wedin & Hafellner, lichenized, lichenicolous, terrestrial, restricted distribution, see Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), Zhurbenko and Tugi (2013; records), Wijayawardene et al. (2017; outline), sequences are unavailable.

Submersisphaeria K.D. Hyde 1996, *Annulatasaceae*, *Annulatascales*, *Sordariomycetes*, five species, type: *S. aquatica* K.D. Hyde, asexual morph unknown, terrestrial, saprobes, aquatic, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: BRIP 22743, one putative strain is available of *S. aquatica*.

Subplenodomus Gruyter, Aveskamp & Verkley 2012, *Leptosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *S. violicola* (P. Syd.) Gruyter, Aveskamp & Verkley coelomycetous, saprobes, terrestrial, worldwide, see de Gruyter et al. (2012; morphology; phylogeny), Wijayawardene et al. (2014c, 2017; outline), Ariyawansa et al. (2015b; phylogeny), Tibpromma et al. (2017b; new species), cultures and sequences are available.

Subramaniomyces Varghese & V.G. Rao 1980, *Beltraniaceae*, *Amphisphaeriales*, *Sordariomycetes*, three species, type: *S. indicus* Varghese & V.G. Rao, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Hongsanan et al. (2017a; phylogeny, evolution), cultures and sequences are available but lacking for the type.

Subramaniula Arx 1985, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, four species, type: *S. thielavioides* (Arx, Mukerji & N. Singh) Arx, asexual morph unknown, saprobes, opportunistic pathogen on human, terrestrial, India, South Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Ahmed et al. (2016; new species, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 560.84, CBS 122.78.

Subsessila C.G. Lin & K.D. Hyde 2017, *Beltraniaceae*, *Amphisphaeriales*, *Sordariomycetes*, one species, type: *S. turbinata* C.G. Lin & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Thailand, see Lin et al. (2017; taxonomy, phylogeny), holotype of the type: MFLU 15-3271, cultures and sequences are available.

Subulispora Tubaki 1971, *Ascomycota* genera *incertae sedis*, eight species, type: *S. procurvata* Tubaki, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, tropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Sugiyamaella Kurtzman & Robnett 2007, *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, 20 species, type: *S. smithiae* (Giménez-Jurado) Kurtzman & Robnett, asexual reproduction is by multilateral budding and through formation of blastoconidia, saprophytic, insect frass in trees, soil, worldwide, Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Sugiyamaemyces I.I. Tav. & Balazuc 1989, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. oroussetii* I.I. Tav. & Balazuc, asexual morph unknown, biotrophic, only known from *Clidicus formicarius* (Coleoptera: Scydmaenidae), Malaysian Borneo, a second species on *Clidicus laticeps*

(from India) is suggested but not formally described because of lack of material, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Suhomyces M. Blackwell & Kurtzman 2016, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, 24 species, type: *S. tanzawaensis* (Nakase & Itoh) M. Blackwell & Kurtzman, asexual reproduction is by multilateral budding, saprophytic, soil, insects, moss, gum exudates of black cherry, maize kernels, rotted mushroom, worldwide, see Kurtzman et al. (2016; taxonomy, phylogeny), sequences are available.

Sulcaria Bystrek 1971, *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, five species, four variety, type: *S. sulcata* (Lév.) Bystrek, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2012c; chemistry), Kirk et al. (2013; genus accepted), Myllys et al. (2014; taxonomy), Hisayoshi et al. (2015; molecular), Lücking et al. (2017a; classification); sequences are available.

Sulcatispora Kaz. Tanaka & K. Hiray. 2015, *Sulcatisporaceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *S. acerina* Kaz. Tanaka & K. Hiray., asexual morph coelomycetous, saprobes, terrestrial, Japan, see Tanaka et al. (2015; taxonomy, phylogeny), Wijayawardene et al. (2016b, 2017; morphology, phylogeny), holotype and ex-type of the type: HHUF 30449, CBS 139703.

Sulcatistroma A.W. Ramaley 2005, *Calosphaeriaceae*, *Calosphaeriales*, *Sordariomycetes*, one species, type: *S. nolinae* A.W. Ramaley, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Sulcopyrenula H. Harada 1999, *Pyrenulaceae*, *Pyrenulales*, *Eurotiomycetes*, four species, type: *S. staurospora* (Tuck.) H. Harada, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Aptroot (2012; new species), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Sulcosporium Phook. & K.D. Hyde 2015, *Halothiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. thailandicum* Phook. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2015a; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 11-0243, MFLUCC 12-0004.

Sungaiicola Fryar & K.D. Hyde 2004, *Sordariomycetes* genera *incertae sedis*, two species, type: *S. bactrodesmiella* Fryar & K.D. Hyde, asexual morph unknown, saprobes, marine, Brunei, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b;

outline), holotype of the type: HKU (M) 15201, cultures and sequences are unavailable.

Superstratomyces van Nieuwenhuijzen, Miadl. & Samson 2016, *Superstratomycetaceae*, *Superstratomycetales*, *Dothideomycetes* three species, type: *S. albomucosus* van Nieuwenhuijzen & Samson, coelomycetous, sexual morph unknown, see van Nieuwenhuijzen et al. (2016; taxonomy), cultures and sequences are available.

Surculiseries Okane, Nakagiri & Tad. Ito 2001, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *S. rugispora* Okane, Nakagiri & Tad., sexual morph unknown, endophytic on mangroves, saprophytic, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: IFO H-12236, ex-type cultures of type: NBRC 33167.

Suttonina H.C. Evans 1984, *Ascomycota* genera *incertae sedis*, one species, type: *S. guatemalitica* H.C. Evans, coelomycetous, sexual morph unknown, saprobes, terrestrial, Guatemala, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Suttonomyces Wijyaw., Camporesi & K.D. Hyde 2015, *Massarinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *S. clematis* Wijyaw., Camporesi & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, Italy, see Tanaka et al. (2015; phylogeny), Wijayawardene et al. (2015, 2016b; taxonomy, phylogeny), holotype and ex-type of the type: HHUF 28661, JCM 13131.

Swampomyces Kohlm. & Volk.-Kohlm. 1987, *Etheiophoraceae*, *Torpedosporales*, *Sordariomycetes*, two species, type: *S. armeniacus* Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, mangroves, Australia, Belize, Brunei, Egypt, North Sumatra, Papua New Guinea, Philippines, Saudi Arabia, South Africa, Thailand, see Besitulo et al. (2010; diversity), Sakayaroj et al. (2011b; diversity), Abdel-Wahab et al. (2014; diversity), Jones et al. (2014, 2015; outline, phylogeny, morphology, re-classification), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Syamithabeeja Subram. & Natarajan 1976, *Ascomycota* genera *incertae sedis*, one species, type: *S. girisa* Subram. & Natarajan, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sydowia Bres., 1895, *Dothideaceae*, *Dothideales*, *Dothideomycetes*, 19 species, type: *S. gregaria* Bres., asexual morph coelomycetous, parasitic or saprobes on

conifer plants or branches and wood, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Thambugala et al. (2014a; taxonomy), Wijayawardene et al. (2014c; outline), holotype of type: S-F6473, cultures and sequences are available.

Sydoiella Petr. 1923, *Sydoiellaceae*, *Diaporthales*, *Sordariomycetes*, four species, type: *S. fenestrans* (Duby) Petr., asexual morphs unknown, saprobes, terrestrial, Europe, North America, see Marinowitz et al. (2008; new species), Lumbsch and Huhndorf (2010; outline), Kruys and Castlebury (2012; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2017, new species, placement), cultures available for the type: CBS 125530.

Sylvacollaea Cif. 1963, *Ascomycota* genera *incertae sedis*, one species, type: *S. termitaria* Cif., coelomycetous, sexual morph unknown, on termites, terrestrial, Asia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Symbiotaphrina Kühlw. & Jurzitz ex W. Gams & Arx 1980, *Xylonomycetaceae*, *Xylonomycetales*, *Xylonomycetes*, two species, type: *S. buchneri* Gräbner ex W. Gams & Arx, coelomycetous, sexual morph unknown, on termites, terrestrial, Asia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Gazis et al. (2016; phylogeny), cultures and sequences are unavailable.

Sympheophyma Speg. 1912, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *S. subtropicale* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Sympaster Theiss. & Syd. 1915, *Dothideomycetes* genera *incertae sedis*, one species, type: *S. gesneraceae* (Henn.) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, South Africa, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Sympyosirinia E.A. Ellis 1956 (= *Sympyosirella* Seifert 2009 *fide* Johnston et al. 2014b), *Helotiaceae*, *Helotiales*, *Leotiomycetes*, six species, type: *S. galii* E.A. Ellis, asexual morph formerly known in *Sympyosirella*, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature),

Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Sympylos Bat. & Cavalc. 1967, *Ascomycota* genera *incertae sedis*, one species, type: *S. sapindaceifolii* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Symplectromyces Thaxt. 1908, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. vulgaris* (Thaxt.) Thaxt., asexual morph unknown, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Sympodiella W.B. Kendr. 1958, *Ascomycota* genera *incertae sedis*, one species, type: *S. acicola* W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Cuba, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are available (unpublished).

Sympodiocladium Descals 1982, *Ascomycota* genera *incertae sedis*, one species, type: *S. frondosum* Descals, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Sympodioclathra Voglmayr 1997, *Ascomycota* genera *incertae sedis*, one species, type: *S. globosa* Voglmayr, hyphomycetous, sexual morph unknown, saprobes, aquatic, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), holotype of the type: WU 17019, cultures and sequences are unavailable.

Sympodioplanus R.C. Sinclair & Boshoff 1997, *Ascomycota* genera *incertae sedis*, one species, type: *S. capensis* R.C. Sinclair & Boshoff, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), holotype: PREM 55315, cultures and sequences are unavailable.

Sympodomycetes R.K. Benj. 1973, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. pentacellularis* R.K. Benj., asexual morph unknown, biotrophic on Coleoptera (Tenebrionoideae), Papua-New Guinea, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Sympoventuria Crous & Seifert 2007, *Sympoventuriaceae*, *Venturiales*, *Dothideomycetes*, one species, type: *S. capensis* Crous & Seifert, asexual morph unknown, worldwide, see Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011d; phylogeny), Hyde et al.

(2013; accepted as a genus in *Sympoventuriaceae*), Wijayawardene et al. (2014c; outline), sequences are available.

Synalissa Fr. 1825, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, 39 species, type: *S. ramulosa* (Bernh.) Körb., asexual morph unknown, lichenized, terrestrial, see Dietrich and Buergi-Meyer (2010; taxonomy), Eichler et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Schultz and Moon (2011; taxonomy), Paukov and Teptina (2012; new record), de la Rosa et al. (2013; ecology), Kirk et al. (2013; genus accepted), Stordeur et al. (2015; new record), Lücking et al. (2017a; classification), sequences unavailable.

Synandromyces Thaxt. 1912, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, ten species, type: *S. telephani* Thaxt., no asexual state, biotrophic, hosts are Biphyllidae, Cryptophagidae, Cucujidae, Nitidulidae, and Tenebrionidae (Coleoptera: Cucujiformia), widespread but not in Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Synaptomyces Thaxt. 1912, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, type: *S. argentinus* Thaxt., no asexual state, biotrophic, only known from *Hydrochara* sp. [as *Hydrocharis*] (Coleoptera, Hydrophilidae), Argentina, two more species are suggested but not formally described, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Synaptopsora Cain 1957, *Helminthosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, four species, type: *S. petrakii* Cain, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Miller et al. (2014; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), sequences are available but lacking for the type.

Synarthonia Müll. Arg. 1891, *Arthoniaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *S. bicolor* Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Menezes et al. (2013a; new species), Joseph and Sinha (2015; taxonomy), Lücking et al. (2017a; classification, treated as *Arthoniales* genera *incertae sedis*), sequences are unavailable.

Synarthothelium Sparrius 2009, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, two species, type: *S. sipmanianum* Sparrius, lichenized, terrestrial, Venezuela, see Sparrius (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), holotype B Sipman 26697, cultures and sequences are unavailable.

Syncarpella Theiss. & Syd. 1915 (= *Syntholus* A.W. Ramaley & M.E. Barr 1997; Art. 59), *Cucurbitariaceae*, *Pleosporales*, *Dothideomycetes*, c. six species, type:

S. tumefaciens (Ellis & Harkn.) Theiss. & Syd., asexual morph coelomycetous, saprobes, terrestrial, cosmopolitan, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Cucurbitariaceae*), Doilom et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), in here we propose to consider *Syntholus* as a synonym of *Syncarpella*, ex-holotype type strains of type: NY, 1677.

Syncesia Taylor 1938 (= *Feigeana* Mies, Lumbsch & Tehler 1995 *fide* Ertz and Tehler 2011), *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, 27 species, type: *S. myrticola* (Fée) Tehler, terrestrial, see Lumbsch and Huhndorf (2010; outline), Ertz and Tehler (2011; phylogeny), Joshi et al. (2011a; new species), Kirk et al. (2013; genus accepted), Joseph et al. (2013; new record), Menezes et al. (2013a; new species), Ertz et al. (2010, 2015b, c; phylogeny), Cáceres and Aptroot (2016; inventory), Lücking et al. (2017a; classification), sequences are available.

Synchaetomella Decock & Seifert 2005, *Chaetomellaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *S. lunatospora* Decock, G. Delgado & Seifert, hyphomycetous, saprobes, terrestrial, Brazil, Singapore, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Crous et al. (2012c; taxonomy), Fiúza et al. (2015c; taxonomy), sequences are available for two species, compare with *Chaetomella*, *Sphaerographium*, holotype of type: MUCL 43804.

Synchronoblastia Uecker & F.L. Caruso 1988, *Ascomycota* genera *incertae sedis*, one species, type: *S. crypta* Uecker & F.L. Caruso, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Syncladium Rabenh. 1859, *Ascomycota* genera *incertae sedis*, one species, type: *S. nietneri* Rabenh., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Synnemacrodictys W.A. Baker & Morgan-Jones 2009, *Ascomycota* genera *incertae sedis*, one species, type: *S. stilboidea* (J. Mena & Mercado) W.A. Baker & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Synnemadiella Crous & M.J. Wingf. 2016, *Xylariales*, genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. eucalypti* Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, with subverticillium-like synasexual

morph, saprobes, terrestrial, Malaysia, see Crous et al. (2016a; new species, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22886, CPC 27637.

Synnemapestalooides T. Handa & Y. Harada 2004, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *S. rhododendri* T. Handa & Y. Harada, coelomycetes, sexual morph unknown, leaves and twigs of *Rhododendron brachycarpum*, Japan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype and ex-type of the type: HHUF26527, MAFF239201.

Synnemaseimatooides K. Matsush. & Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *S. nipponica* K. Matsush. & Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Synnematomyces Kobayasi 1981, *Ascomycota* genera *incertae sedis*, one species, type: *S. capitatus* Kobayasi, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available (unpublished).

Synnemellisia N.K. Rao, Manohar. & Goos 1989, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *S. hyalospora* N.K. Rao, Manohar. & Goos, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2016a; new species, phylogeny), cultures and sequences are available.

Synostomina andina Petr. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *S. andina* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Synsphaeria Bonord. 1851, *Sordariomycetes*, genera *incertae sedis*, one species, type: *S. fumigata* Vuill., asexual morph unknown, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline).

Syphosphaera Dumort. 1822, *Ascomycota* genera *incertae sedis*, one species, type: *S. coronata* Dumort., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Syrropeltis Bat., J.L. Bezerra & Matta 1964, *Dothideomycetes* genera *incertae sedis*, one species, type:

S. xylopiae Bat., J.L. Bezerra & Matta, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Sympastospora P.F. Cannon & D. Hawksw. 1982, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *S. parasitica* (Tul.) P.F. Cannon & D. Hawksw., fungicolous, widespread, terrestrial, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in Hypocreaceae), Kirk et al. (2013; genus accepted), cultures and sequences are available, holotype of type: ? PC.

Systemmopsis Petr. 1923, *Ascomycota* genera *incertae sedis*, one species, type: *S. ribesia* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Szczawinskia A. Funk, 1984, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *S. tsugae* A. Funk, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Øvstdal and Gremmen (2010; new species), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Taeniarella S. Hughes 1958, *Kirschsteiniotheliaceae*, *Kirschsteiniothliales*, *Dothideomycetes*, c. 40 species, type: *T. exilis* (P. Karst.) S. Hughes, sexual morph unknown, saprophytic, endophytic, lichenicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Ertz et al. (2016b; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Taeniolina M.B. Ellis 1976, *Ascomycota* genera *incertae sedis*, six species, type: *T. centaurii* (Fuckel) M.B. Ellis, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Zhang et al. (2012i; new species, China), Kirk et al. (2013; genus accepted), Prasher and Verma (2014; new species, India), cultures and sequences are unavailable.

Tainospaeria F.A. Fernández & Huhndorf 2005, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, one species, type: *T. crassiparies* F.A. Fernández & Huhndorf, asexual morph unknown, saprobes, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Liu et al. (2016a; new species, phylogeny), Lu et al. (2016; new species), Perera et al. (2016b; new records in Thailand), sequence available for the type: AF466056 (beta tubulin).

Taiwanascus Sivan. & H.S. Chang 1997, *Niesliaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *T. tetrasporus* Sivan. & H.S. Chang asexual morph unknown, saprobes, lignicolous, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rajeshkumar and Rossman (2013; new species), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: IMI 364835, cultures and sequences are unavailable.

Takamatsuella U. Braun & A. Shi 2012, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, one species, type: *T. circinata* (Cooke & Peck) U. Braun & A. Shi, asexual morph unknown, biotrophic, obligate plant pathogens, terrestrial, North America, see Braun and Cook (2012; taxonomy), sequences are available, lectotype of type: NYS-F-762.

Talaromyces C.R. Benj. 1955 (= *Erythrogymnotheca* Yaguchi & Udagawa; = *Paratalaromyces* Matsush.; = *Penicillium* Link subgenus *Biverticillium* Dierckx; = *Penicillium* Link subgenus *Biverticillata-Symmetrica* Thom; = *Sagenoma* Stolk & G.F. Orr *fide* Samson et al. 2011), *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, c. 114 species, type: *T. flavus* (Klöcker) Stolk & Samson, terrestrial, but sometimes also isolated from aquatic habitats, associated with diverse substrates, including living or decaying plant material, soil, compost, dung, house dust, foodstuffs, leather, arthropods, rodents, air, cosmopolitan, the genus includes some clinically-relevant species, of which the most important is *Talaromyces marneffei* (Segretain, Capponi & Sureau) Samson, Yilmaz, Frisvad & Seifert (= *Penicillium marneffei* Segretain, Capponi & Sureau), a thermally dimorphic fungus which commonly infects immunocompromised patients in southeast Asia, see Proksa (2010; secondary metabolites), Samson et al. (2011; phylogeny, new combinations, epitypification of the type:), Houbraken and Samson (2011; phylogeny), Guarro et al. (2012; soil borne species), Yilmaz et al. (2012, 2014, 2016a, 2016b; phylogeny, revision, new species), Frisvad et al. (2013; new species), Sang et al. (2013; new species), Visagie et al. (2015; new species), Chen et al. (2016a; new species), Wang et al. (2016a; new species), ex-epitype strain of the type: = CBS 310.38.

Talekpea Lunghini & Rambelli 1979, *Ascomycota* genera *incertae sedis*, one species, type: *T. foeticia* Lunghini & Rambelli, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Ivory Coast, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Talpapellis Alstrup & M.S. Cole 1998, *Ascomycota* genera *incertae sedis*, two species, type: *T. peltigerae* Alstrup & M.S. Cole, hyphomycetous, sexual morph unknown, lichenicolous, Canada, see Kirk et al. (2008;

genus accepted), Heuchert et al. (2014; taxonomic revision), Zhurbenko et al. (2015c; new species), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, holotype of type: M.S. Cole 1632 (C).

Tamaricicola Thambug., E. Camporesi & K.D. Hyde 2016, *Pleosporaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *T. muriformis* Thambug., Camporesi & K.D. Hyde, asexual morph coelomycetous, saprobes, terrestrial, Italy, see Thambugala et al. (2017a; taxonomy, phylogeny), holotype and ex-type of the type: MFLU 16-0676, MFLUCC 15-0488.

Tamhinispora Rajeshkumar & Rahul Sharma 2013, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, one species, type: *T. indica* Rajeshkumar & Rahul Sharma, hyphomycetous, sexual morph unknown, saprobes, see Rajeshkumar and Sharma (2013; morphology), Boonmee et al. (2014b; placed in *Tubeufiales*, phylogeny, key), holotype and ex-type of the type: AMH 9555, NFCCI 2924.

Tamsiniella S.W. Wong, K.D. Hyde, W.H. Ho & S.J. Stanley 1998, *Sordariomycetes* genera *incertae sedis*, two species, type: *T. labiosa* S.W. Wong, K.D. Hyde, W.H. Ho & S.J. Stanley, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: HKU (M) 2276, cultures and sequences are unavailable.

Tandonea M.D. Mehrotra 1991, *Ascomycota* genera *incertae sedis*, one species, type: *T. dargentiana* M.D. Mehrotra, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 269926, cultures and sequences are unavailable.

Tangerinosporium L. Lombard & Crous 2016, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *T. thaliticola* L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, UK, see Lombard et al. (2016; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: CBS H-22472, CBS 317.61.

Tania Egea, Torrente & Sipman 1995, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, two species, type: *T. lanosa* Egea, Torrente & Sipman, lichenized, terrestrial, Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), Lücking et al. (2017a; classification), sequences unavailable.

Tapellaria Müll. Arg. 1898, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, 22 species, type: *T. herpetospora* Müll. Arg., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010;

outline), Lumbsch et al. (2011; new species), Messuti et al. (2011; new species), Kirk et al. (2013; genus accepted), Farkas (2014; taxonomy), van den Broeck et al. (2014; taxonomy), Lücking et al. (2017a; classification), McCarthy and Elix (2016; new species), sequences are unavailable.

Tapellariopsis Lücking 1999, *Pilocarpaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *T. octomera* Lücking, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable, The name as published in 1999 was deemed invalid in *Index Fungorum* because of supposed lack of a Latin diagnosis; however, all required elements are present in the protologue and the name is valid.

Taphridium Lagerh. & Juel 1902, *Protomycetaceae*, *Taphrinales*, *Taphrinomycetes*, two species, type: *T. umbelliferarum* (Rostr.) Lagerh. & Juel, asexual morph unknown, ?saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Taphrina Fr. 1815, *Taphrinaceae*, *Taphrinales*, *Taphrinomycetes*, ca. 100 species, type: *T. aurea* (Pers.) Fr., asexual morph unknown, biotrophic, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Selbmann et al. (2014; new species, taxonomy), cultures and sequences are available for some species.

Four new combinations into *Taphrina* for names in *Lalaria* were invalidly published by Selbmann et al. (2014), since the pagination of the whole publication on which the basionyms appeared, rather than the individual pages, were cited [ICN (Melbourne) Art. 41.5 Note 1].

The four new combinations are here validated as follows:

Taphrina arrabidae (Á. Fonseca, J. Inácio & M.G. Rodrigues) S. Onofri & L. Zucconi, **comb. nov.** Index Fungorum 553576. Basionym: *Lalaria arrabidae* Á. Fonseca, J. Inácio & M.G. Rodrigues (FEMS Yeast Research 4:551, 2004).

Taphrina inositophila (Á. Fonseca, J. Inácio & M.G. Rodrigues) A. Yurkov & P. Buzzini, **comb. nov.** Index Fungorum 553577. Basionym: *Lalaria inositophila* Á. Fonseca, J. Inácio & M.G. Rodrigues (FEMS Yeast Research 4:552, 2004).

Taphrina kurtzmanii (Á. Fonseca, J. Inácio & M.G. Rodrigues) B. Turchetti & C. Cecchini, **comb. nov.** Index Fungorum 553578. Basionym: *Lalaria kurtzmanii* Á. Fonseca, J. Inácio & M.G. Rodrigues (FEMS Yeast Research 4:553, 2004).

Taphrina veronaerambelli (Á. Fonseca, J. Inácio & M.G. Rodrigues) L. Selbmann & C. Cecchini, **comb. nov.** Index Fungorum 553579. Basionym: *Lalaria*

veronaerambellii Á. Fonseca, J. Inácio & M.G. Rodrigues (FEMS Yeast Research 4:553, 2004).

Taphrophila Scheuer 1988, *Dothideomycetes* genera *incertae sedis*, four species, type: *T. cornu-capreoli* Scheuer, asexual morph unknown, on rotten leaves, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tarasginia S.Y. Kondr., Kärnefelt, A. Thell, Elix & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *T. whinrayi* (S.Y. Kondr. & Kärnefelt) S.Y. Kondr., Kärnefelt, A. Thell, Elix, J. Kim, A.S. Kondr. & Hur, lichenized, see Kondratyuk et al. (2015c; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Tarbertia Dennis 1974, *Arthoniales* genera *incertae sedis*, one species, type: *T. juncina* Dennis, ?lichenized or saprobes, terrestrial, British Isles, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Tarsodisporus Bat. & A.A. Silva 1965, *Ascomycota* genera *incertae sedis*, one species, type: *T. paraensis* Bat. & A.A. Silva, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tarzetta (Cooke) Lambotte 1887, *Pezizales* genera *incertae sedis*, *Pezizomycetes*, nine species, type: *P. tarzetta* Cooke, asexual morph unknown, hypogeous, terrestrial, worldwide, see Hansen et al. (2013; phylogeny), Olariaga et al. (2015; taxonomy), culture and sequences are available but lacking for the type.

Tasmidella Kantvilas, Hafellner & Elix 1999, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *T. variabilis* Kantvilas, Hafellner & Elix, lichenized, see Ekman et al. (2008; taxonomy), Lendemer (2010b; taxonomy), Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; phylogeny), Lumbsch et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences unavailable.

Tassiloa S.Y. Kondr., Kärnefelt, A. Thell, Elix & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *T. digitaurea* (Søgaard, Søchting & Sancho) S.Y. Kondr., Kärnefelt, A. Thell, Elix, J. Kim, A.S. Kondr. & Hur, lichenized, see Kondratyuk et al. 2015e, Lücking et al. (2017a; classification), sequences are available.

Tatraea Svrček 1993, ?*Helotiaceae*, *Helotiales*, *Leotiomycetes*, two species, type: *T. dumbirensis* (Velen.) Svrček, asexual morph unknown, saprobe of wood,

widespread (temperate), see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype of the type: PRM 148876 (Svrček 1993), cultures and sequences are unavailable.

Tavaresiella T. Majewski 1980, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *T. hebri* T. Majewski, asexual morph unknown, biotrophic, hosts are Hebridae (Hemiptera: Heteroptera), Africa, Asia, and Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tayloriellina S.Y. Kondr., Kärnefelt., A. Thell, Elix & Hur 2016, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *T. erythrosticta* (Taylor) S.Y. Kondr., Kärnefelt., A. Thell, Elix, J. Kim, A.S. Kondr. & Hur, lichenized, see Kondratyuk et al. (2016a; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Tectacervulus A.W. Ramaley 1992, *Ascomycota* genera *incertae sedis*, one species, type: *T. mahoniae* A.W. Ramaley, coelomycetous, sexual morph unknown, pathogens, on leaf spots, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: BPI 1107358, cultures and sequences are unavailable.

Teichospora Fuckel 1870, *Teichosporaceae*, *Pleosporales*, *Dothideomycetes*, 35 species, type: *T. trabicola* Fuckel, asexual morph coelomycetous, terrestrial, saprobes, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Teichosporaceae*), Wijayawardene et al. (2014c; outline), Crous et al. (2016a; new species, phylogeny), Jaklitsch et al. (2016c; new species, phylogeny), Liu et al. (2017, placement), cultures available for type: C160, C134, C141.

Teichosporella (Sacc.) P. Karst. 1895, *Dothideomycetes* genera *incertae sedis*, 26 species, type: *T. dura* Sacc., asexual morph unknown, saprobes, terrestrial, America, Germania, Germany, Italy, temperate, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Telimena Racib. 1900, *Telimenaceae*, *Phyllachorales*, *Sordariomycetes*, c. 16 species, type: *T. erythrinae* Racib., from living leaves, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline, accepted as a genus in *Phyllachoraceae*), Mardones et al. (2017; phylogeny), cultures and sequences are unavailable.

Telimenella Petr. 1940, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, two species, type: *T. persica* Petr., from living leaves, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al.

(2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Telimenochora Sivan. 1987, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *T. abortiva* (F. Stevens) Sivan., from living leaves, Mexico, Puerto Rico, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Teliocleipeum Viégas 1962, *Ascomycota* genera *incertae sedis*, one species, type: *T. lavrense* Viégas, saprobes, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Telogalla Nik. Hoffm. & Hafellner 2000, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *T. olivieri* (Vouaux) Nik. Hoffm. & Hafellner, asexual morph unknown, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Teloschistes Norman 1853, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 20 species, type: *T. flavicans* (Sw.) Norman, asexual morph unknown, lichenized, terrestrial, worldwide, see Muggia et al. (2008; taxonomy, phylogeny), Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012, 2015; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures available for the type: AFTOL-ID 315.

Teloschistopsis Frödén, Søchting & Arup 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *T. chrysocarpoides* (Vainio) Frödén, Arup & Søchting, asexual morph unknown, lichenized, terrestrial, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Temerariomyces B. Sutton 1993, *Ascomycota* genera *incertae sedis*, one species, type: *T. acutulus* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Malawi, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 347181, cultures and sequences are unavailable.

Temperantia K. Hansen, Healy & Kovács 2011, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, nine species, type: *T. tiffanyae* (Healy) K. Hansen, Healy & Kovács, asexual morph unknown, hypogeous, terrestrial, USA, see Kovács et al. (2011b; taxonomy, phylogeny), holotype of the type: ISC, Healy 231, sequence are available.

Tengomyces 1999, *Helminthosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, four species, type: *T. indicus* (Varghese & V.G. Rao), asexual morph unknown, saprobes, terrestrial, India, China, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al.

(2015b, 2016b; outline), cultures and sequences are unavailable.

Tenuiappendicula Senan., Camporesi & K.D. Hyde 2017, *Sydotiellaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *T. alnicola* Senan., Camporesi & K.D. Hyde, asexual morphs coelomycetous, saprobes, terrestrial, Europe, see Senanayake et al. (2017; taxonomy, phylogeny), holotype of the type: MFLU 16–1265A, cultures and sequences are available.

Tenuimurus Senan., Camporesi & K.D. Hyde 2016, *Phomatosporaceae*, *Phomatosporales*, *Sordariomycetes*, one species, type: *T. clematis* Senan., Camporesi & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, temperate, Italy, see Senanayake et al. (2016; taxonomy, phylogeny), cultures and sequences are available.

Tephromela M. Choisy 1929, *Tephromelataceae*, *Lecanorales*, *Lecanoromycetes*, 30 species, type: *T. atra* (Huds.) Hafellner, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Muggia et al. (2010b, 2014; morphology, phylogeny, taxonomy), Pérez-Vargas et al. (2010a; new species), Fryday (2011; new species), Spribille et al. (2011; molecular), Elix (2012; new species), Kirk et al. (2013; genus accepted), Wang et al. (2013b; taxonomy), Muggia et al. (2014; taxonomy), Kantvilas (2015; taxonomy), Szczepańska (2015; new records), Cestaro et al. (2016; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Teracosphaeria & Seifert 2007, *Sordariomycetes* genera *incertae sedis*, two species, type: *T. petroica* & Seifert, asexual morph unknown, saprobes, terrestrial, New Zealand, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Annulatasaceae* and *Sordariomycetidae*, genera *incertae sedis*), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: PDD 81436, cultures and sequences are available.

Teratomyces Thaxt. 1893, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, nine species, type: *T. mirificus* Thaxt., asexual morph unknown, biotrophic, hosts are Staphylininae (Coleoptera: Staphylinidae), widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossi (2010; most recent species description), Haelewaters et al. (2015c; synonymy of *T. brevicaulis* with *T. actobii*), cultures and sequences are unavailable.

Teratoschaeta Bat. & O.M. Fonseca 1967, *Dothideomycetes* genera *incertae sedis*, two species, type: *T. rondoniensis* Bat. & O.M. Fonseca, asexual morph unknown, lichenicolous, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Teratosperma Syd. & P. Syd. 1909, *Ascomycota* genera *incertae sedis*, one species, type: *T. singulare* Syd. & P. Syd., saprobes, Africa, North America, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Teratosphaeria Syd. & P. Syd. 1912 (= *Colletogloeopsis* Crous & M.J. Wingf. 1997; = *Kirramyces* J. Walker et al. 1992 *fide* Crous et al. 2009f; Wijayawardene et al. 2014c; Rossman et al. 2015b), *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, c. 90 species, type: *T. fibrillosa* Syd. & P. Syd., asexual morph coelomycetous (formerly known in *Colletogloeopsis*, *Kirramyces*), saprobes, pathogens, terrestrial, cosmopolitan, see Crous et al. (2009c, d, e, f, 2011e; taxonomy, phylogeny), Andjic et al. (2010, 2016; new species, Australia, cryptic species), Lumbsch and Huhndorf (2010; outline), Yamazaki et al. (2010; acidophilic), Hunter et al. (2011; diseases), Taole et al. (2012; *T. suttonii* complex), Taylor et al. (2012; pathogens), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, nomenclature), Quaedvlieg et al. (2014; phylogeny), cultures and sequences are available.

Teratosphaericola Quaedv. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *T. pseudoafricana* (Crous & T.A. Cout.) Quaedv. & Crous, asexual morph unknown, pathogens, Zambia, see Quaedvlieg et al. (2014; morphology, phylogeny), holotype and ex-type culture of the type: PREM 54973, CPC 1229.

Teratosphaeriopsis Quaedv. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *T. pseudoafricana* Quaedv. & Crous asexual morph unknown, one species, pathogens, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), holotype and ex-type culture of the type: CBS H-21137, CBS 111171.

Terfezia (Tul. & C. Tul.) Tul. & C. Tul. 1851, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, 21 species, type: *T. arenaria* (Moris) Trappe, asexual morph unknown, hypogeous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kovács et al. (2011a; new species), Bordallo et al. (2012, 2013, 2015; new species), Kirk et al. (2013; genus accepted), sequences are available.

Termitaria Thaxt. 1920, *Ascomycota* genera *incertae sedis*, three species, type: *T. snyderi* Thaxt., hyphomycetous, sexual morph unknown, America, Sardinia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Termitariopsis M. Blackw., Samson & Kimbr. 1980, *Ascomycota* genera *incertae sedis*, three species, type:

T. cavernosa M. Blackw., Samson & Kimbr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North and Central America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Terriera B. Erikss. 1970, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, 26 species, type: *T. cladophila* (Lév.) B. Erikss., asexual morph unknown, saprobes, terrestrial, China, Europe, Thailand, see Lumbsch and Huhndorf (2010; outline), Lantz et al. (2011; DNA, phylogeny), Chen et al. (2012; new species, China), Kirk et al. (2013; genus accepted), Zhou et al. (2013; new species, China), Wu et al. (2015; new species), Hyde et al. (2016; new species, Thailand), sequences are available.

Testudina Bizz. 1885, *Testudinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *T. terrestris* Bizz., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Testudinaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Testudomyces Cano, M. Solé & Guarro 2002, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *T. verrucosus* M. Solé, Cano & Guarro, asexual morph unknown, saprobes, terrestrial, Spain, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences is available for the type (unpublished).

Tetrabrachium Nawawi & Kuthub. 1987, *Ascomycota* genera *incertae sedis*, three species, type: *T. elegans* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tetrabrunneospora Dyko 1978, *Ascomycota* genera *incertae sedis*, two species, type: *T. ellisii* Dyko, hyphomycetous, sexual morph unknown, saprobes, aquatic, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tetrachaetum Ingold 1942, *Leotiomycetes* genera *incertae sedis*, three species, type: *T. elegans* Ingold, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available (unpublished).

Tetracladium De Wild. 1893, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, eight species, type:

T. marchalianum De Wild., hyphomycetous, sexual morph unknown, saprobes, aquatic, terrestrial, worldwide, see Sati et al. (2009; new species), Letourneau et al. (2010; DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CCM F-11001.

Tetracoccosprium Szabó 1905, *Ascomycota* genera *incertae sedis*, three species, type: *T. paxianum* Szabó, hyphomycetous, sexual morph unknown, on dung and soil, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tetracrium Henn., 1902, *Dothideomycetes* genera *incertae sedis*, nine species, type: *T. aurantii* Henn., hyphomycetous, sexual morph unknown, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tetramelas Norman 1853, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, 16 species, type: need typification, lichenized, see Gaya et al. (2012; phylogeny), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; taxonomy), sequences are available.

Tetrameronycha Speg. ex W. Rossi & M. Blackw. 1990, *Ascomycota* genera *incertae sedis*, one species, type: *T. bonaerensis* (Speg.) W. Rossi & M. Blackw., hyphomycetous, sexual morph unknown, on insects, terrestrial, Africa, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tetranaciella Kohlm. & Volkm.-Kohlm. 2001, *Ascomycota* genera *incertae sedis*, one species, type: *T. papillata* Kohlm. & Volkm.-Kohlm., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tetranacrium H.J. Huds. & B. Sutton 1964, *Ascomycota* genera *incertae sedis*, one species, type: *T. gramineum* H.J. Huds. & B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, West Indies, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tetrandromyces Thaxt. 1912, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, six species, type: *T. brachidae* Thaxt., no asexual state, biotrophic, hosts are Anthicidae and Staphylinidae (Coleoptera), widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al.

(2013; genus accepted), cultures and sequences are unavailable.

Tetrapisispora Ueda-Nishimura & Mikata emend. Kurtzman 2003, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, seven species, type: *T. phaffii* (van der Walt) Ueda-Nishimura & Mikata, asexual reproduction is by multilateral budding, saprophytic, in soil, Japanese blueberry, mangrove trees, cockroach, food processing plant, worldwide, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Tetraploa Berk. & Broome 1850 (= *Tetraplosphaeria* Kaz. Tanaka & K. Hiray fide Wijayawardene et al. 2014c; Rossman et al. 2015b), *Tetraplosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, 16 species, type: *T. aristata* Berk. & Broome, asexual morph coelomycetous, saprobes, terrestrial or aquatic, worldwide, see Tanaka et al. (2009; phylogeny, morphology), Wijayawardene et al. (2014c; nomenclature), Rossman et al. (2015b; nomenclature), holotype of the type: IMI 17841, cultures and sequences are available.

Tetrasporium S. Hughes 1951, *Ascomycota* genera *incertae sedis*, two species, type: *T. asterinearum* S. Hughes, hyphomycetous, sexual morph unknown, fungicolous, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tettigomyces Thaxt. 1915, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, 16 species, type: *T. gryllotalpae* Thaxt., asexual morph unknown, biotrophic, hosts are *Gryllotalpa* species and *Scapteriscus* sp. (*Orthoptera: Gryllotalpidae*), widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Teunomyces Kurtzman & M. Blackwell 2016, *Debaromyctaceae*, *Saccharomycetales*, *Saccharomycetes*, twelve species, type: *T. kruisii* (Kocková-Kratochvílová & Ondrušová) Kurtzman & M. Blackwell, asexual reproduction is by multilateral budding, saprophytic, in insects, often the guts of insects, worldwide, see Kurtzman et al. (2016; taxonomy), cultures and sequences are available.

Teuva Sohrabi & S. Leavitt 2013, *Megasporaceae*, *Pertusariales*, *Lecanoromycetes*, three species, type: *T. uxorius* (Werner) Sohrabi, V. J. Rico & S. Leavitt, lichenized, see Sohrabi et al. (2013b; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Texosporium Nádv. ex Tibell & Hofsten 1968, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, two species, type: *T. sancti-jacobi* (Tuck.) Nádv. ex Tibell & Hofsten, asexual morph unknown, lichenized, terrestrial, see

Lumbsch and Huhndorf (2010; outline), Kirby et al. (2011; ecology), Kirk et al. (2013; genus accepted), Root et al. (2011, 2012, 2013; ecology), Rosentreter et al. (2014; ecology), Lücking et al. (2017a; classification), Prieto and Wedin (2016; ecology), sequences are available.

Textotheca Matsush. 1996, *Ascomycota* genera *incertae sedis*, one species, type: *T. caffra* Matsush., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thailandiomyces Pinruan, Sakay., K.D. Hyde & E.B.G. Jones 2008, *Tirisporellaceae*, *Tirisporellales*, *Sordariomycetes*, one species, type: *T. bisetulosus* Pinruan, Sakay., Hyde & E.B.G. Jones, saprobes, aquatic, see Lumbsch and Huhndorf (2010; outline), Jones et al (2015; classification, phylogeny, marine), Maharachchikumbura et al. (2015b, 2016b; classification, phylogeny), Suetrong et al. (2015; phylogeny), holotype and cultures available for the type: BBH, Pinruan, Wah 110, BCC18.

Thalassoascus Ollivier 1926, *Dothideomycetes* genera *incertae sedis*, three species, type: *T. tregoubovii* Ollivier, asexual morph unknown, parasites, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Jones et al (2015; classification, marine), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Thalassogena Kohlm. & Volk.-Kohlm. 1987, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *T. sphaerica* Kohlm. & Volk.-Kohlm., asexual morph unknown, saprobes, marine, worldwide, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of type: J.K. 5044a = 01390521, NYBG.

Thalespora Chatmala & E.B.G. Jones 2006, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *T. appendiculata* Chatmala & E.B.G. Jones, asexual morph unknown, marine, on wood, tropical, see Sakayaroj et al. (2011a; phylogeny), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Thallinocarpon E. Dahl 1950, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, two species, type: *T. pulvinatum* E. Dahl, lichenized, see Lücking et al. (2017b; classification), sequences are unavailable.

Thalloloma Trevis. 1853, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 19 species, type: *T. anguinum* (Mont.) Trevis., asexual morph unknown, lichenized, terrestrial, tropical, placement in *Graphidaceae* supported by molecular data, see Rivas Plata et al. (2012a; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013;

genus accepted), Lücking et al. (2017a; classification), sequences are available.

Thallomyces H.J. Swart 1975, *Parmulariaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *T. oritis* (Hansf.) H.J. Swart, asexual morph unknown, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Thamnochrolechia Aptroot & Sipman 1991, *Pertusariaceae*, *Pertusariales*, *Lecanoromycetes*, one species, type: *T. verticillata* Aptroot & Sipman, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available, holotype of the type: B, Aptroot 18622.

Thamnogalla D. Hawksw. 1980, *Cordieritidaceae*, *Leotiomycetes* families *incertae sedis*, one species, type: *T. crombiei* (Mudd) D. Hawksw., lichenicolous, Europe and North America, terrestrial, see Kirk et al. (2013; genus accepted), Suija et al. (2015b; phylogeny), Jaklitsch et al. (2016a; classification), cultures and sequences are available, lectotype of type: BM.

Thamnolecania (Vain.) Gyeln., 1933, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, eight species, type: *L. brialmontii* Vain., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Thamnolia Ach. ex Schaer. 1850, *Icmadophilaceae*, *Pertusariales*, *Lecanoromycetes* four species, type: *T. vermicularis* (Sw.) Schaer., asexual morph unknown, lichenized, terrestrial, see Nelsen. and Gargas (2009; molecular), Diederich et al. (2010b; taxonomy), Manojlović et al. (2010a, b; HPLC analysis, bioactivity), Schmull et al. (2011; molecular), Lumbsch and Huhndorf (2010; outline), Hansen (2013; taxonomy), Kirk et al. (2013; genus accepted), Knight et al. (2013; chemotaxonomy), Árnason (2014; chemistry), Lücking et al. (2017a; classification), sequences are available.

Thamnomyces Ehrenb. 1820, *Hypoxylaceae*, *Xylariales*, *Sordariomycetes*, five species, type: *T. chamissonis* Ehrenb., asexual morph hyphomycetous, terrestrial, saprobes, tropical, see Lumbsch and Huhndorf (2010; outline), Stadler et al. (2010b; new species), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available for unidentified species.

Thaptospora B. Sutton & Pascoe 1987, *Ascomycota* genera *incertae sedis*, one species, type: *T. atrobrunnea* B.

Sutton & Pascoe, coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thaumasiomyces Thaxt. 1931, *Ceratomyctaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, three species, type: *T. scabellularius* Thaxt., no asexual state, biotrophic, hosts are Hydrophilidae (Coleoptera), Africa and Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thaxteria Sacc. 1891, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, nine species, type: *T. kunkelii* Giard, asexual morph unknown, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Thaxteriella Petr., 1924, *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, nine species, type: *T. corticola* Petr., saprobes on decaying wood, Puerto Rico, Thailand, asexual morph hyphomycetous, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; taxonomy), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Thaxteriellopsis Sivan., Panwar & S.J. Kaur 1976 [1977], *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, three species, type: *T. lignicola* Sivan., Panwar & S.J. Kaur, saprobes on decaying wood, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; taxonomy), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Thecaria Fée 1824, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *T. quassitcola* Fée, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2012a; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Thecographa A. Massal. 1860, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *T. ceramia* A. Massal., asexual morph unknown, lichenized fungi, terrestrial, tropical, see Rivas Plata et al. (2012a; taxonomy), Lücking et al. (2017a; classification), sequences are unavailable.

Thecotheus Boud. 1869, *Ascobolaceae*, *Pezizales*, *Pezizomycetes*, 20 species, type: *T. pelletieri* (P. Crouan & H. Crouan) Boud., asexual morph unknown, coprophilous, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kušan et al. (2015a; new species), cultures and sequences are available but lacking for the type.

Thegdonia B. Sutton 1973, *Helotiales* genera *incertae sedis*, one species, type: *T. ligustrina* (Boerema) B. Sutton, hyphomycetous, sexual morph unknown, terrestrial, plant pathogens, leaf-spotting, Asia, Europe, North America, see Crous et al. (2009a, 2012a, phylogeny), Seifert et al. (2011), Wijayawardene et al. (2017; outline), ex-type culture: CBS 148.59, holotype of type: the Netherlands, Eefde, on *Ligustrum ovalifolium*, 23 Mar. 1959, G.H. Boerema (L).

Theissenia Maubl. 1914, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, four species, type: *T. pyrenocrata* (Theiss.) Maubl., asexual morph hyphomycetous, terrestrial, saprobes, pantropical, see Hsieh et al. (2010; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), cultures and sequences are available.

Thelebolus Tode 1790 (= *Lasiothelebolus* Kimbr. & Luck-Allen 1974; = *Pezizella* P. Karst., 1873; = *Pezizula* P. Karst. 1871; = *Ryparobius* Boud. 1869; = *Telebolus* Lindau 1905 fide Species Fungorum 2017), *Thelebolaceae*, *Thelebolales*, *Leotiomycetes*, c. 13 species, type: *T. stercoreus* Tode, asexual morph unknown, saprobes, from soil, on dung, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Mukhopadhyay et al. (2014; extracellular enzymes), Singh et al. (2014; extracellular enzymes), Crous et al. (2015d; new species), cultures available for the type: CBS 709.69, CBS 717.69 (unpublished).

Thelenella Nyl. 1855, *Thelenellaceae*, *Ostropales*, *Lecanoromycetes*, 30 species, type: *T. modesta* (Nyl.) Nyl., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kinalioğlu and Aptroot (2011; new records of Turkey), Kirk et al. (2013; genus accepted), Aptroot et al. (2014a; new species), Cáceres and Aptroot (2016; new species), Nelsen et al. 2016, Lücking et al. (2017a; classification), sequences are available.

Thelenidia Nyl. 1886, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. monospora* Nyl., asexual morph unknown, lichenized, terrestrial, Greenland, Switzerland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline, accepted as *Dothideomycetes*), Lücking et al. (2017a; classification), sequences are unavailable.

Theliella Fink 1933, *Sordariomycetes* genera *incertae sedis*, two species, type: *T. blastenica* Fink ex J. Hedrick, asexual morph unknown, lichenicolous, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Thelidiopsis Vain. 1921, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, one species, type: *T. robinsonii* Vain., asexual morph unknown, lichenized, terrestrial, Europe, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Thelidium A. Massal. 1855, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, c. 100 species, type: *T. amylaceum* A. Massal., asexual morph unknown, lichenized, terrestrial, fresh water, Europe, Asia, see Holger et al. (2008; fresh water species), Lumbsch and Huhndorf (2010; outline), Pykälä et al. (2012; species in Russia), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Thelignya A. Massal. 1856, *Lichenaceae*, *Lichinales*, *Lichenomycetes*, three species, type: *T. fuliginea* (Ach.) A. Massal., asexual morph unknown, lichenized, terrestrial, see Jørgensen and Palice (2010; taxonomy), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kristinsson et al. (2014; taxonomy), Lücking et al. (2017a; classification), sequences unavailable.

Thelliana S.Y. Kondr., Kärnefelt, Elix & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species; type: *T. pseudokiamae* S.Y. Kondr., Kärnefelt, Elix & Hur; lichenized; see Kondratyuk et al. (2015c; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Thelocarpella Nav.-Ros. & Cl. Roux, 1999, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, one species, type: *T. gordensis* Nav.-Ros. & Cl. Roux, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Tonouchi (2010; characterization), Knudsen et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Gueidan et al. (2014c; taxonomy), Westberg et al. (2015; phylogeny), Roux et al. (2016; taxonomy), Lücking et al. (2017a; classification); sequences are available.

Thelocarpon Nyl. 1853, *Thelocarpaceae*, *Thelocarpales*, *Ascomycota* orders *incertae sedis*, 27 species, type: *T. laureri* (Flot.) Nyl., asexual morph unknown, lichenized or non-lichenized, terrestrial, worldwide, see Moon and Aptroot (2009; new species, South Korea), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Flakus and Kukwa (2014; new species), Cáceres and Aptroot (2016; new species, Brazil), van den Boom (2016; new species), Jaklitsch et al. (2016a; classification), Lücking et al. (2017a; classification), sequences are available.

Thelomma A. Massal. 1860, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, seven species; type: *T. mammosum* (Hepp) A. Massal., lichenized, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; phylogeny),

Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Thelonectria P. Chaverri & C. Salgado 2011, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, 43 species, type: *T. discophora* (Mont.) P. Chaverri & C. Salgado, asexual morph cylindrocarpon-like, saprobes, terrestrial, worldwide, see Chaverri et al. (2011; taxonomy, phylogeny), Salgado-Salazar et al. (2012, 2014, 2016; new species, phylogeny), Zeng and Zhuang (2013b; new species, China), Maharachchikumbura et al. (2015b, 2016b; outline), Lombard et al. (2016; phylogeny), cultures available for the type: CBS 125153.

Thelopsis Nyl. 1855, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, nine species, type: *T. rubella* Nyl., asexual morph unknown, lichenized, terrestrial, see Giordani et al. (2009; new records), Øvstedral and Gremmen (2009; new record), Aptroot et al. (2010; biodiversity), Potenza et al. (2010; new records), Lumbsch and Huhndorf (2010; outline), Dymytrova et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Aptroot et al. (2014c; world key), Mies (2015; new records), Stehn et al. (2015; list), Lücking et al. (2017a; classification), sequences unavailable.

Thelotrema Ach. 1803, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 106 species, type: *T. lepadinum* (Ach.) Ach., asexual morph unknown, lichenized, terrestrial, mostly tropical, see Mangold et al. (2009; key, Australia), Lumbsch and Huhndorf (2010; outline), Rivas Plata et al. (2010a, 2012a; key), Sipman et al. (2012; key, Costa Rica), Weerakoon et al. (2012b; new species), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Themisia Velen. 1939, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, eight species, type: *T. nuda* Velen., asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thermoascus Miehe 1907, *Thermoascaceae*, *Eurotiales*, *Eurotiomycetes*, five species, type: *T. aurantiacus* Miehe, asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as in *Trichocomaceae*), Houbraken and Samson (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Thermothelomyces Y. Marín, Stchigel, Guarro & Cano 2015, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, four species, type: *T. thermophila* (Apinis) Y. Marín, Stchigel, Guarro & Cano, see Marín-Felix et al. (2014; taxonomy), Maharachchikumbura et al. (2016b; outline), cultures and sequences are available.

Thermitis Fr. 1825, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, two species, type: *T. velutina* (Ach.) Flot., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Urbanavichus and Urbanavichene (2011; new records), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Thermutopsis Henssen 1990, *Lichinaceae*, *Lichinales*, *Lichenomycetes*, one species, type: *T. jamesii* Henssen, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); sequences are unavailable.

Therrya Sacc. 1882, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, seven species, type: *T. gallica* Sacc. & Penz., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Solheim et al. (2013; DNA, phylogeny), cultures and sequences are available but lacking for the type.

Thielavia Zopf 1876, *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, 31 species, type: *T. basicola* Zopf, asexual morph unknown, saprobes, saprobes, terrestrial, worldwide, see Kusari et al. (2009; endophytes, secondary metabolites), Lumbsch and Huhndorf (2010; outline), Berka et al. (2011; genomic analysis), Kirk et al. (2013; genus accepted), Van den Brink et al. (2015; DNA, thermophilic properties), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 229.82.

Thielaviopsis Went 1893, *Ceratocystidaceae*, *Microascales*, *Sordariomycetes*, six species, type: *T. ethacetica* Went, asexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see de Beer et al. (2014; new species), Mbenoun et al. (2014, 2016; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), de Melo et al. (2016; pathogens), Rossman et al. (2016b; new combination), cultures available for the type: CMW37775.

Thindia Korf & Waraitch 1971, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *T. cupressii* Korf & Waraitch, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thindiomyces Arendh. & R. Sharma 1983, *Hyaloscyphaceae*, *Helotiidae*, *Leotiomycetes*, one species, type: *T. epiphyllus* Arendh. & Sharma, asexual morph unknown, saprobes of leaves, Bhutan, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), holotype: PAN 17589, cultures and sequences are unavailable.

Thirumalacharia Rathaiah 1981, *Ascomycota* genera *incertae sedis*, one species, type: *T. curcumae* Rathaiah, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tholomyces Matsush. 2003, *Ascomycota* genera *incertae sedis*, one species, type: *T. flagellifer* Matsush., coelomycetous, sexual morph unknown, saprobes, terrestrial, Honshu, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tholurna Norman 1861, *Caliciaceae*, *Caliciales*, *Lecanoromycetes*, one species, type: *T. dissimilis* Norman, asexual morph unknown, lichenized, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Gaya et al. (2012; molecular), Allen and Lendemer (2013; molecular), Kirk et al. (2013; genus accepted), Prieto et al. (2013; phylogeny), Beese et al. (2015; ecology), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Thoracella Oudem. 1901, *Ascomycota* genera *incertae sedis*, one species, type: *T. ledi* Oudem., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thozetella Kuntze 1891, *Chaetosphaeriaceae*, *Chaetosphaerales*, *Sordariomycetes*, 17 species, type: *T. nivea* (Berk.) Kuntze, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Whitton et al. (2012a; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Perera et al. (2016b; new species), cultures and sequences are available for unidentified species.

Thrauste Theiss. 1916, *Englerulaceae*, *Dothideomycetes* families *incertae sedis*, three species, type: *T. medinillae* (Racib.) Theiss., sexual morph hyphomycetous, epiphytes, saprobes, terrestrial, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Dai et al. (2014c; morphology), cultures and sequences are unavailable.

Thrinacospora Petr. 1948, *Ascomycota* genera *incertae sedis*, one species, type: *T. insignis* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Ecuador, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thripomyces Speg. 1915, *Ceratomycetaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, two species, type:

T. italicus Speg., asexual morph unknown, biotrophic, hosts are *Hydraena* species (Coleoptera: Hydraenidae), Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thrombium Wallr. 1831, *Thrombiaceae, Ostropales, Lecanoromycetes*, five species, type: *T. epigaeum* (Pers.) Wallr., Russia, America, see Nadyeina (2009; taxonomy), Kukwa (2010; list), Lumbsch and Huhndorf (2010; outline), Stepanchikova et al. (2011; new records), Kirk et al. (2013; genus accepted), Resl et al. (2015), Lücking et al. (2017a; classification), sequences are available.

Thryptospora Petr. 1947, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. singularis* Petr., asexual morph unknown, saprobes, terrestrial, Syria, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Thuemenella Penz. & Sacc. 1898 (= *Chromocreopsis* Seaver 1910 fide Wendt et al. 2017), *Hypoxylaceae, Xylariales, Sordariomycetes*, one species, type: *T. javanica* Penz. & Sacc., asexual morph hyphomycetous, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Hypoxylaceae*), cultures and sequences are available.

Thyrea A. Massal. 1856, *Lichinaceae, Lichinales, Lichenomycetes*, 13 species, type: *T. plectospora* A. Massal., asexual morph unknown, lichenized, terrestrial, South Korea, Iran, Turkey, USA, see Björk (2010; new records), Lumbsch and Huhndorf (2010; outline), Schultz and Moon (2011; taxonomy), Kirk et al. (2013; genus accepted), Candan and Schultz (2015; new records), Ahti et al. (2016; checklist), Lücking et al. (2017a; classification); sequences are available.

Thyridaria Sacc. 1875, *Thyridariaceae, Pleosporales, Dothideomycetes*, 52 species, type: *T. incrustans* Sacc., asexual morph unknown, saprobes, terrestrial, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology; phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Thyridella (Sacc.) Sacc. 1895, *Sordariomycetes* genera *incertae sedis*, three species, type: *T. colliculus* Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Thyridium Nitschke 1867, *Thyridiaceae, Diaportheomycetidae* families *incertae sedis*,

Sordariomycetes, c. 20 species, type: *T. vestitum* (Fr.) Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), sequence are available.

Thyrinula Petr. & Syd. 1924, *Ascomycota* genera *incertae sedis*, one species, type: *T. eucalypti* (Cooke & Massee) H.J. Swart, coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, Australia, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Thyriodictyella Cif. 1962, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. microsperma* (Syd. & P. Syd.) Cif., asexual morph unknown, saprobes, terrestrial, Dominican Republic, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Micropeltidaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Thyriopsis Theiss. & Syd. 1915 (= *Dothiclypeolum* Höhn. 1916 fide Species Fungorum 2017), *Aulographaceae, Dothideomycetes* families *incertae sedis*, one species, type: *T. halepensis* (Cooke) Theiss. & Syd., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hyde et al. (2013; accepted as a genus in *Aulographaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; taxonomy, accepted as a genus in *Aulographaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Thyriostromella Bat. & C.A.A. Costa 1959, *Ascomycota* genera *incertae sedis*, one species, type: *T. calophylli* (Sacc.) Bat. & C.A.A. Costa, coelomycetous, sexual morph unknown, saprobes, terrestrial, Malaysia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thronectria Sacc. 1875 (= *Pleonectria* Sacc. 1876; = *Chilonectria* Sacc. 1878; = *Nectria* subg. *Aponectria* Sacc. 1878; ≡ *Aponectria* (Sacc.) Sacc. 1883; = *Mattirolia* Berl. & Bres. 1889; = *Scoleconectria* Seaver 1909; = *Thronectroidea* Seaver 1909 fide Jaklitsch and Voglmayr 2014), *Nectriaceae, Hypocreales, Sordariomycetes*, c. 40 species, type: *T. patavina* Sacc., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch and Voglmayr (2014; revisited, phylogeny), Lombard et al. (2015a; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Thyrostroma Höhn. 1911 (= *Thyrococcum* (Sacc.) Sacc. 1913; = *Thyrococcum* Höhn. 1907; = *Thyrostromella* Syd.

1924; = *Wilsonomyces* Adask., J.M. Ogawa & E.E. Butler 1990 *fide Species Fungorum* 2017), *Dothidotthiaceae*, *Pleosporales*, *Dothideomycetes*, c. 45 species, type: *T. compactum* (Sacc.) Höhn., hyphomycetous, sexual morph unknown, saprobes, pathogens, terrestrial, worldwide, see Phillips et al. (2008; accepted as genus in *Dothidotthiaceae*), Seifert et al. (2011; morphology), Hyde et al. (2013; accepted as genus in *Dothidotthiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Crous et al. (2016b; phylogeny), cultures available but lacking for the type.

Thyrostromella Höhn. 1919, *Ascomycota* genera *incertae sedis*, three species, type: *T. myriana* (Desm.) Höhn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australia, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thyrotheca Kirschst. 1944, *Sordariomycetes* genera *incertae sedis*, three species, type: *T. nyssae* (Berk. & M.A. Curtis) Kirschst., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Thrysidiella Höhn. 1909 (= *Thrysidiella* Höhn. 1905 *fide Species Fungorum* 2017), *Ascomycota* genera *incertae sedis*, two species, type: *T. lignicola* (Höhn.) Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Thysidina Höhn. 1905, *Ascomycota* genera *incertae sedis*, one species, type: *T. carneominiata* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Thysanorea Arzanlou, W. Gams & Crous 2007, *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *T. papuana* (Aptroot) Arzanlou, W. Gams & Crous, hyphomycetous, sexual morph unknown, saprobes, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirschner (2016; taxonomy, phylogeny), holotype and cultures available for the type: CBS, Aptroot 36647, CBS 212.96.

Thysanothecium Mont. & Berk. 1846, *Cladoniaceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *T. hookeri* Mont. & Berk., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Tiarospora Sacc. & Marchal 1885, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *T. westendorpii* Sacc. & Marchal, coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Phookamsak et al. (2014c; accepted as a genus in *Phaeosphaeriaceae*), cultures and sequences are unavailable.

Tiarosporella Höhn. 1919 (= *Tiarosporella* Höhn. 1924 *fide Species Fungorum* 2017), *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, c. five species, type: *S. visci* (Alb. & Schwein.) Sacc., coelomycetous, saprobes, endophytes, terrestrial, cosmopolitan, see Phillips et al. (2008, 2013; taxonomy, phylogeny), Liu et al. (2012b; taxonomy, phylogeny), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Slippers et al. (2013; phylogeny), Thambugala et al. (2014d; sexual morph, phylogeny), Crous et al. (2015b; phylogeny), Wijayawardene et al. (2014c, 2017 outline), Dissanayake et al. (2016b; taxonomy), cultures and sequences are available.

Tiarosporellivora Punith. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *T. caricina* Punith., coelomycetous, sexual morph unknown, lichenicolous, terrestrial, Germany, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tibellia Vezda & Hafellner 1992, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *T. dimerelloides* Vezda & Hafellner, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Ticogloea G. Weber, Spaaij & W. Gams 1994, *Ascomycota* genera *incertae sedis*, one species, type: *T. guttulata* G. Weber, Spaaij & W. Gams, hyphomycetous, sexual morph unknown, saprobes, ?endophytes, terrestrial, Europe, Central America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Ticomyces Toro 1952 (= *Dontuzia* L.D. Gómez 1973; = *Tonduzia* F. Stevens 1927 *fide Species Fungorum* 2017), *Sordariomycetes* genera *incertae sedis*, one species, type: *T. psychotriae* (F. Stevens) Toro, asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsan et al. (2015d; excluded the genus from *Meliolaceae* and placed in *Sordariomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Ticonectria Döbbeler 1998, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *T. perianthii* Döbbeler, asexual morph unknown, on liverworts, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Ticosynnema R.F. Castañeda, Granados & Mardones 2013, *Ascomycota* genera *incertae sedis*, one species, type: *T. carrranzae* R.F. Castañeda, Granados & Mardones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Costa Rica, see Castañeda-Ruiz et al. (2013a; taxonomy), Wijayawardene et al. (2017; outline), holotype of the type: HAL 2454F, cultures and sequences are unavailable.

Tilachlidium Preuss 1851, *Tilachlidiaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *T. pinnatum* Preuss, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; phylogeny, new family), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Tilakidium Vaidya, C.D. Naik & Rathod 1986, *Hypocreales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *T. indicum* Vaidya, C.D. Naik & Rathod, asexual morph unknown, saprobes, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Tilakiella Srinivas. 1973, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. indica* Srinivas., asexual morph unknown, saprobes, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Timdalia Hafellner 2001, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, one species, type: *T. intricata* (H. Magn.) Hafellner, lichenized, see Lumbsch and Huhndorf (2010; outline), Westberg et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available.

Tingoldiago K. Hiray. & Kaz. Tanaka 2010, *Lentitheciaeae*, *Pleosporales*, *Dothideomycetes*, one species, type: *T. graminicola* K. Hiray. & Kaz. Tanaka, saprobes, Japan, see Hirayama et al. (2010; taxonomy, phylogeny), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Tinhaudeus K.L. Pang, S.Y. Guo & E.B.G. Jones 2015, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *T. formosanus* K.L. Pang, S.Y. Guo & E.B.G. Jones, asexual morph unknown, saprobes, marine,

Taiwan, see Ariyawansa et al. (2015a; morphology, phylogeny), Maharachchikumbura et al. (2016b; phylogeny, outline), holotype and ex-type strains of type: F28727, BCRC FU30496.

Tirispora E.B.G. Jones & Vrijmoed 1994, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, two species, type: *T. unicaudata* E.B.G. Jones & Vrijmoed, asexual morph unknown, two species, marine, on mangrove wood, tropical, see Lumbsch and Huhndorf (2010; outline), Sakayaroj et al. (2011a; phylogeny), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Tirisporella E.B.G. Jones, K.D. Hyde & Alias, 1996, *Tirisporellaceae*, *Tirisporellales*, *Sordariomycetes*, one species, type: *T. beccariana* (Ces.) E.B.G. Jones, K.D. Hyde & Alias asexual morph phialophora-like, on petioles of mangrove palm *Nypa*, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Tirmania Chatin 1892, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, three species, type: *T. nivea* (Desf.) Trappe, asexual morph unknown, hypogeous, terrestrial, worldwide, see Al-Laith (2010; Antioxidant components and antioxidant/antiradical activities), Lumbsch and Huhndorf (2010; outline), Dib-Bellahouel and Fortas (2011; antibacterial activity), Kirk et al. (2013; genus accepted), Osmundson et al. (2013; DNA), sequences are available.

Titaea Sacc. 1876 (= *Aorate* Syd. 1929; = *Maxillispora* Höhn. 1914; = *Monogrammia* F. Stevens 1917 fide Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, 23 species, type: *T. callispora* Sacc., hyphomycetous, see Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Titaeopsis B. Sutton & Deighton 1984, *Ascomycota* genera *incertae sedis*, one species, type: *T. ugandae* (Hansf.) B. Sutton & Deighton, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Titaeospora Bubák 1916, *Ascomycota* genera *incertae sedis*, two species, type: *T. detospora* (Sacc.) Bubák, coelomycetous, sexual morph unknown, lichenicolous, terrestrial, temperate, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Toensbergia Bendiksby & Timdal 2013, *Sporastatiaceae*, *Rhizocarpales*, *Lecanoromycetes*, one

species, type: *T. leucococca* (R. Sant.) Bendiksby & Timdal, lichenized, corticolous, temperate northern hemisphere, see Bendiksby and Timdal (2013; phylogenetic placement), Lücking et al. (2017a; classification), sequences are available, type specimen UPS- L-86993 (holotype).

Togniniella Réblová, L. Mostert, W. Gams & Crous 2004 (= *Phaeocrella* Réblová et al. 2004 *fide* Réblová et al. 2016c), *Calosphaeriaceae*, *Calosphaeriales*, *Sordariomycetes*, twelve species, type: *T. acerosa* Réblová, L. Mostert, W. Gams & Crous, asexual morph formerly named in *Phaeocrella*, from living leaves, terrestrial, New Zealand, see Damm et al. (2008b; DNA, morphology of ascii), Lumbsch and Huhndorf (2010; outline), Réblová (2011; phylogeny), Réblová et al. (2016c; phylogeny), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: CBS 113726.

Tolypocladium W. Gams 1971 (= see Species Fungorum 2017 for synonyms), *Elaphocordyceps* G.H. Sung & Spatafora 2007 *fide* Maharachchikumbura et al. 2016b), *Ophiocordycipitaceae*, *Hypocreales*, *Sordariomycetes*, c. 15 species, type: *T. inflatum* W. Gams, hyphomycetous, sexual morph formerly known in *Elaphocordyceps*, on soil, rotifers and mosquito larvae, terrestrial, cosmopolitan, see Sato et al. (2010a, b; new species), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Tomasellia A. Massal. 1856, *Naetrocymbaceae*, *Dothideomycetes* families *incertae sedis*, 50 species, type: *T. arthonioides* (A. Massal.) A. Massal., asexual morph unknown, epiphytes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Naetrocymbaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tomenticola Deighton 1969, *Ascomycota* genera *incertae sedis*, two species, type: *T. trematis* Deighton, hyphomycetous, sexual morph unknown, on living leaves, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Tomeoa I. Hino 1954, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. viridicollculosa* I. Hino, asexual morph unknown, saprobes, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Tompetchia Subram. 1985, *Ascomycota* genera *incertae sedis*, one species, type: *T. webberi* (H.S. Fawc.) Subram., hyphomycetous, sexual morph unknown, on insects, terrestrial, Australasia, North and South America, see

Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Toninia A. Massal. [nom. cons.] 1852 (see Species Fungorum 2017 for synonyms), *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, species, 85 species, type: *T. squalida* (Ach.) A. Massal., asexual morph unknown, lichenized, terrestrial, worldwide, see Moniri et al. (2009; taxonomy), Lumbsch and Huhndorf (2010; outline), Farkas et al. (2011; ecology), Schmull et al. (2011; phylogeny), Urbanavichus et al. (2011; new records), Bendiksby and Timdal (2013; phylogeny), Kirk et al. (2013; genus accepted), Zhurbenko and Zhdanov (2013; taxonomy), Maier et al. (2014; ecology), Joshi et al. (2015a; new records), Sezer et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Toniniopsis Frey 1926, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *T. obscura* Frey, lichenized, see Lumbsch and Huhndorf (2010; outline), Lücking et al. (2017a; classification), sequences are unavailable.

Topelia P.M. Jørg. & Vezda, 1984 (= *Clathroporinopsis* M. Choisy 1929 *fide* Species Fungorum 2017), *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, six species, type: *T. rosea* (Servít) P.M. Jørg. & Vezda, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Aptroot et al. (2014c; checklist), Lücking et al. (2017a; classification), sequences are unavailable.

Topeliopsis Kantvilas & Vezda 2000, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, 20 species, type: *T. muscicola* Kantvilas & Vezda, asexual morph unknown, lichenized, terrestrial, see Messuti et al. (2010; new species), Lumbsch and Huhndorf (2010; outline), Sipman et al. (2012; new combinations), Mangold et al. 2009 (key Australia), Rivas Plata et al. (2010a, 2012a; taxonomy), Lücking et al. (2014; new species), Lumbsch et al. (2014a, b; phylogeny), Weerakoon et al. (2015; new species), Lücking et al. (2017a; classification), sequences are available.

Toriella (Kohlm.) Sakay., K.L. Pang & E.B.G. Jones 2011, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *T. tubulifera* (Kohlm.) Sakay., K.L. Pang & E.B.G. Jones, asexual morph unknown, saprobes, marine, Canada, Denmark, England, Germany, Iceland, Norway, Sweden, USA, Wales, see Sakayaroj et al. (2011a; morphology, phylogeny), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), holotype and ex-type of the type: BCC33511, BCC33512, BCC33513.

Tornabea Østh. 1980 (= *Tornabenia* Trevis. 1853; = *Tornabeniopsis* Follmann 1980 *fide* Species Fungorum 2017), *Physciaceae*, *Caliciales*, *Lecanoromycetes*, one

species, type: *T. montagnei* Trevis., asexual morph unknown, lichenized, terrestrial, Iran, Pakistan, Kayseri-Adamna, see Lumbsch and Huhndorf (2010; outline), Moniri (2010; taxonomy), Sohrabi et al. (2010b; taxonomy), Aptroot and Iqbal (2012; checklist), Gaya et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), Parsa et al. (2013; taxonomy), Halici et al. (2014a; taxonomy), Nattah et al. (2014, taxonomy, inventory), Lücking et al. (2017a; classification), Prieto and Wedin (2016; phylogeny), sequences are available.

Toroa Syd. 1926, *Pseudoperisporiaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *T. dimerosporioides* (Speg.) Syd., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Pseudoperisporiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Torpedospora Meyers 1957 (= *Glomerulispora* Abdel-Wahab & Nagah. *fide* Réblová et al. 2016c), *Torpedosporaceae*, *Torpedosporales*, *Sordariomycetes*, five species, type: *T. radiata* Meyers, asexual morph formerly named in *Glomerulispora*, saprobes, marine, worldwide, see Abdel-Wahab et al. (2010; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Réblová et al. (2016c; nomenclature).

Torrendiella Boud. & Torrend 1911 (= *Zoellneria* Velen. 1934 *fide* Species Fungorum 2017), *Rutstroemiaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *T. ciliata* Boud., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; taxonomy, phylogeny), Jaklitsch et al. (2016a; classification), cultures available for the type: F132996.

Torrentispora K.D. Hyde, W.H. Ho, E.B.G. Jones, K.M. Tsui & S.W. Wong 2000, *Annulatasccaceae*, *Annulatascales*, *Sordariomycetes*, one species, type: *T. fibrosa* K.D. Hyde, Wai H. Ho, E.B.G. Jones, K.M. Tsui & S.W. Wong, asexual morph unknown, saprobes, aquatic, Hong Kong, Brunei, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Barbosa et al. (2013a; new species, Costa Rica), Maharachchikumbura et al. (2016b; outline), holotype of the type: HKU (M) 4519, cultures and sequences are available.

Torrubiella Boud. 1885, *Cordycipitaceae*, *Hypocreales*, *Sordariomycetes*, 71 species, type: *T. aranicida* Boud., asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Johnson et al. (2009; taxonomy, DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Sato et al. (2010a; new species), Kirk et al. (2013; genus

accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Tortispora Lachance & Kurtzman 2013, *Trigonopsidaceae*, *Saccharomycetales*, *Saccharomycetes*, eight species, type: *T. ganteri* Lachance & Kurtzman, asexual growth is by multilateral budding, saprophytic, in rotted cacti, Costa Rica, Hawaii, USA, Argentina, Brazil, Mexico, see Lachance et al. (2013a; taxonomy), cultures and sequences are available.

Tortilispora Senan. & K.D. Hyde 2017, *Sydiowelliaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *T. aurantiaca* (Wehm.) Senan. & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Europe, see Senanayake et al. (2017; taxonomy, phylogeny), holotype of the type: MICH 13964, cultures and sequences are available.

Tortulomyces Lar. N. Vassiljeva, S.L. Stephenson, Chernyshev & K.D. Hyde 2013, *Nitschkiaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *T. thailandicus* Lar. N. Vassiljeva, S.L. Stephenson, A.V. Chernyshev & K.D. Hyde, asexual morphs unknown, saprobes, terrestrial, south-eastern Asia, see Maharachchikumbura et al. (2016b; outline), holotype of type: MFLU 11-1163, cultures and sequences are unavailable.

Torula Pers. 1795 (= *Hormiscium* Kunze 1817; = *Satwalekera* D. Rao, V.G. Rao & P.Rag. Rao 1970 [1969]; = *Taeniola* Bonord. 1851; = *Tetraclonium* Kunze ex Link 1824 *fide* Species Fungorum 2017), *Torulaceae*, *Pleosporales*, *Dothideomycetes*, eleven species, type: *T. herbarum* (Pers.) Link, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Crous et al. (2015a, 2016a; typification, new species, phylogeny), Su et al. (2016b; phylogeny), Li et al. (2016d; phylogeny, new species), neotype and epitype of the type: CBS H-22275, CPC 24114.

Torulaspora Lindner 1904 (= *Azymomyces* E.K. Novák & Zsolt 1961; = *Debaryomyces* Klöcker 1909; = *Zymodebaryomyces* E.K. Novák & Zsolt 1961 *fide* Species Fungorum 2017), *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, six species, type: *T. delbrueckii* (Lindner) Lindner, asexual reproduction is by multilateral budding, saprophytic, soil, plant material, Indonesian ragi, grapes, tree bark, mangrove trees, spoiled fruit juice, dough, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Torulopsisella Bender 1932 (= *Torulopsis* Speg. 1918 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *T. fumaginea* (Speg.) Bender, hyphomycetous, sexual morph unknown, on leaves (sooty moulds), terrestrial, Australasia, South America, see Seifert

et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Tothia Bat. 1960, *Venturiaceae*, *Venturiales*, *Dothideomycetes*, two species, type: *T. fuscella* (Sacc.) Bat., asexual morph unknown, saprobes, terrestrial, Austria, Niederösterreich, Hungary and Italy, see Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; morphology; phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline; phylogeny), Liu et al. (2015a; morphology, phylogeny), cultures and sequences are available for all species.

Tovariella Syd. 1930, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *T. pittieriana* Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Towyspora Wanas., E.B.G. Jones & K.D. Hyde 2016, *Lentitheciaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *T. aestuari* Wanas., E.B.G. Jones & K.D. Hyde, coelomycetous, sexual morph unknown, saprobes, terrestrial, UK, see Li et al. (2016a; taxonomy, phylogeny), holotype and ex-isotype of the type: MFLU 15–3543, MFLUCC 15–1274.

Toxicocladosporium Crous & U. Braun 2007, *Cladosporiaceae*, *Capnodiales*, *Dothideomycetes*, 14 species, type: *T. irritans* Crous & U. Braun, asexual morphs hyphomycetous, terrestrial, Africa, Asia, Australia, South America, see Seifert et al. (2011; morphology), Hyde et al. (2013; accepted as a genus in *Cladosporiaceae*), Wijayawardene et al. (2012, 2014c, 2017; outline), Crous et al. (2016b; new species, phylogeny), Bezerra et al. (2017b; new species), holotype and ex-type strains of the type: CBS-H 19892, CBS 185.58.

Toxosporiella B. Sutton 1986, *Ascomycota* genera *incertae sedis*, one species, type: *T. bactrodesmioides* B. Sutton, coelomycetous, sexual morph unknown, on leaves (sooty moulds), terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Toxosporiopsis B. Sutton & Sellar 1966, *Ascomycota* genera *incertae sedis*, one species, type: *T. capitata* B. Sutton & Sellar, coelomycetous, sexual morph unknown, on leaves (sooty moulds), terrestrial, Australia, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trabutia Sacc. & Roum. 1881, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *T. abortiva* (F. Stevens) Sivan., from living leaves, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus not listed), Maharachchikumbura

et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Tracylla (Sacc.) Tassi 1904 (= *Leptothyrium* subgen. *Tracylla* Sacc. 1895; = *Mitrorhizopeltis* Bat. & Cavalc. (1964) [1963] *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *T. spartinae* (Peck) Tassi, coelomycetous, saprobes, terrestrial, Brazil, Dominican Republic, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trailia G.K. Sutherl. 1915, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, three species, type: *T. ascophylli* G.K. Sutherl., asexual morph unknown, terrestrial, one marine on brown seaweeds, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Trapelia M. Choisy [nom. cons.] 1929 (= *Discocera* A.L. Sm. & Ramsb. 1918) [1917]; = *Trapelina* Motyka 1996 *fide* Species Fungorum 2017), *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, 13 species, type: *T. coarctata* (Turner ex Sm.) M. Choisy, asexual morph unknown, lichenized, terrestrial, Macaronesia, Korea, Australia, Romania, Poland, Italy, Iraq, India, see Czarnota and Kukwa (2009; taxonomy), Baloch et al. (2010; phylogeny), Çobanoğlu et al. (2010; new records), Hodkinson (2010; biodiversity), Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; taxonomy), Tsurykau and Khramchankova (2011; checklist), Aptroot and Schumm (2012a; new species and new records), Łubek (2012; ecology), Rajakaruna et al. (2012; new record), Wieczorek and Łysko (2012; taxonomy), Joshi et al. (2013a; new records), Kirk et al. (2013; genus accepted), Vondrák and Liška (2013; checklist), Ertz et al. (2014a; new species), Giralt et al. (2014; taxonomy), Brodo and Lendemer (2015; new species), Goni and Sharma (2015; new records), Kantvilas et al. (2015; new species), Resl et al. (2015; phylogeny), Telfer et al. (2015; biodiversity inventory, DNA barcoding), Aziz and Qadir (2016; new records), Schneider et al. (2016; phylogeny and partial sequence), Lücking et al. (2017a; classification), sequences are available.

Trapeliopsis Hertel & Gotth. Schneid. 1980, *Trapeliaceae*, *Trapeliales*, *Lecanoromycetes*, 20 species, type: *T. wallrothii* (Flörke ex Spreng.) Hertel & Gotth. Schneid., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hodkinson and Lendemer (2011b; taxonomy), Aptroot and Schumm (2012a; new species), Elix (2013; *Trapeliopsis granulosa* var. *australis*, new species), Kirk et al. (2013;

genus accepted), Sanz-Resl et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification), sequences are available for the type: CP956.

Traponora Aptroot 1997, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, five species, type: *T. asterella* Aptroot, asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Tremateia Kohlm., Volk.-Kohlm. & O.E. Erikss. 1995, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *T. halophila* Kohlm., Volk.-Kohlm. & O.E. Erikss., asexual morph phoma-like., saprobes, marine, Atlantic Coast, see Suetrong et al. (2009; phylogeny, morphology), Jones et al. (2009b, 2015; classification), Ariyawansa et al. (2014e; accepted as a genus in *Didymosphaeriaceae*), Wijayawardene et al. (2014c; outline), holotype of type: J.K. 5522 (IMS).

Trematophoma Petr. 1924, *Ascomycota* genera *incertae sedis*, two species, type: *T. lignicola* Petr., coelomycetous, saprobes, terrestrial, Brazil, Dominican Republic, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available for unidentified species.

Trematosphaeria Fuckel 1870 (= *Botanamphora* Nograsek & Scheuer 1990 *fide* Species Fungorum 2017), *Trematosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, c. 20 species, type: *T. pertusa* Fuckel, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Pleosporales*, genera *incertae sedis*), Suetrong et al. (2011b; phylogeny, new family), Hyde et al. (2013; accepted as a genus in *Trematosphaeriaceae*, phylogeny), Kirk et al. (2013; genus accepted), Zhang et al. (2012e; phylogeny), Ahmed et al. (2014a; new species), Wijayawardene et al. (2014c; outline), Zhou et al. (2014; new species), Tanaka et al. (2015; phylogeny), cultures and sequences are available.

Trematosphaeriopsis Elenkin 1901, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. parmeliana* Jacz. & Elenkin, lichenicolous, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Dai et al. (2014a; taxonomy), Wijayawardene et al. (2014c; outline), sequences are unavailable.

Trematovalsa Jacobesco 1906, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *T. matruchotii* Jacobesco, asexual morph unknown, saprobes, terrestrial, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Tremellidium Petr. 1927 (= *Kordyanella* Höhn. 1904 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae*

sedis, one species, type: *T. piskorzii* Petr., coelomycetous, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tremolecia M. Choisy 1953, *Hymeneliaceae*, *Lecanorales*, *Lecanoromycetes*, six species, type: *T. dicksonii* (Ach.) M. Choisy, asexual morph unknown, lichenized, terrestrial, Australia, Asia, North America, see Hansen (2009, 2014, 2015; taxonomy), Lumbsch and Huhndorf (2010; outline), Aptroot and Iqbal (2012; checklist), Kirk et al. (2013; genus accepted), Kantvilas (2014b; taxonomy, new species), McCune and Rosentreter (2014; new records), Miadlikowska et al. (2014a; molecular), Lücking et al. (2017a; classification); sequences are available.

Tremotylium Nyl. 1868 (= *Phanotylium* Clem. 1909 *fide* Species Fungorum 2017), *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, type: *T. angolense* Nyl., one species, lichenized, terrestrial see Makhija and Patwardhan (1995; systematics), Lücking et al. (2017b; classification), sequences are unavailable.

Trenomyces Chatton & F. Picard 1909, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eleven species, type: *T. histophtorus* Chatton & F. Picard, asexual morph unknown, biotrophic, dioecious, hosts are Mallophaga (Arthropoda: Phthiraptera), widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence are unavailable.

Tretocephala Subram. 1995 (= *Tretocephala* Subram. 1992 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *T. decidua* Subram., hyphomycetous, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tretolytea Cantillo, R.F. Castañeda & Gusmão 2015, *Ascomycota* genera *incertae sedis*, one species, type: *T. pleiomorpha* T. Cantillo, R.F. Castañeda & Gusmão, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Cantillo-Pérez et al. (2016; morphology), holotype of the type: HUEFS 211337, cultures and sequences are unavailable, compare with *Capnophialophora*, *Capnosporium Heteroconium*, *Lylea*, *Taeniola* and *Taeniolina*.

Tretospeira Piroz. 1972, *Ascomycota* genera *incertae sedis*, one species, type: *T. ugandensis* (Hansf.) Piroz., hyphomycetous, saprobes, terrestrial, Africa, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tretovularia Deighton 1984, *Ascomycota* genera *incertae sedis*, one species, type: *T. villiana* (Magnus) Deighton, hyphomycetous, saprobes, terrestrial, Europe,

see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Treibiomycetes Höhn. 1909 (= *Chaetomeris* Clem. 1931; = *Marceloa* Bat. & Peres 1962; = *Mycostevensonia* Bat. & Cif. 1962; = *Paropsis* Speg. 1918 *fide* Species Fungorum 2017), *Chaethyriaceae*, *Chaethyriales*, *Eurotiomycetes*, seven species, type: *T. pulcherrimus* Höhn., asexual morph unknown, epiphytes, sooty moulds, terrestrial, America, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Triacutus G.L. Barron & Tzean 1981, *Sordariomycetes* genera *incertae sedis*, 18 species, type: *T. subcuticularis* G.L. Barron & Tzean, hyphomycetous, saprobes, from soil, on immunocompromised patients, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Edathodu et al. (2013; human pathogens *Triadelphia pulvinata*, DNA, treated as *Hypocreomycetidae incertae sedis*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable for the type, genus could be polyphyletic.

Triadelphia Shearer & J.L. Crane 1971, *Microascales* genera *incertae sedis*, *Sordariomycetes*, 18 species, type: *T. heterospora* Shearer & J.L. Crane, hyphomycetous, sexual morph unknown, saprobes, human pathogen, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Triainomyces W. Rossi & A. Weir 1998, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, *T. hollowayanus* W. Rossi & A. Weir, asexual morph unknown, biotrophic, known from *Procyliosoma tuberculatum* (Diplopoda: Chilognatha: Oniscomorpha: Sphaerotheriidae), New Zealand, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable, needs revision.

Triangularia Boedijn 1934 (= *Trigonia* J.F.H. Beyma 1933 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, six species, type: *T. bambusae* (J.F.H. Beyma) Boedijn, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Triblidium Rebent. 1805 (= *Blitridium* De Not. (1864) [1863]; = *Phaciopsis* Hazsl. 1873; = *Triblidium* subgen. *Triblidium* Rebent. 1805 *fide* Species Fungorum 2017), *Triblidaceae*, *Triblidiales*, *Leotiomycetes*, six species, type: *T. caliciforme* Rebent., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and

Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch and Voglmayr (2016; classification), cultures and sequences are unavailable.

Tribolospora D.A. Reid 1966, *Ascomycota* genera *incertae sedis*, one species, type: *T. sycopsisidis* D.A. Reid, coelomycetous, sexual morph unknown, saprobes, on leaves of *Sycopsis dunnii*, New Guinea, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tribulatia Joanne E. Taylor, K.D. Hyde & E.B.G. Jones 2003, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *T. appendicospora* Joanne E. Taylor, K.D. Hyde & E.B.G. Jones, saprobes, Queensland, see Lumbsch and Huhndorf (2010; outline), Maharanachikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Tricellula Beverw. 1954, *Ascomycota* genera *incertae sedis*, one species, type: *T. inaequalis* Beverw., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology, accepted as a genus in *Orbiliaceae*), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Triceromyces T. Majewski 1980, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, eight species, type: *T. balazucii* T. Majewski, no asexual state, biotrophic, monoecious, dioecious, and trioeccious, hosts are Hebridae, Hydrometridae, and Veliidae (Hemiptera: Heteroptera), worldwide, see Santamaría (2008; morphology, review), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Tricharia Fée 1825 (= *Microxyphiomyces* Bat. 1961; = *Psathyromyces* Bat. & Peres (1964) [1963] *fide* Species Fungorum 2017), *Gomphillaceae*, *Ostropales*, *Lecanoromycetes*, 30 species, type: *T. melanothrix* Fée, asexual morph unknown, lichenized, terrestrial, India, Brazil, Colombia, see Lumbsch and Huhndorf (2010; outline), Lumbsch et al. (2011; new species), Messuti et al. (2011; new species), Mateus et al. (2012; taxonomy), Kirk et al. (2013; genus accepted), Rashmi and Rajkumar (2015; taxonomy, new record), Sanders and Lücking (2015; new species), Sanders and de los Rios (2016; morphotaxonomy), Gupta and Sinha (2016; new report), Lücking et al. (2017a; classification), van den Boom and Sipman (2016; new records), sequences are available.

Tricharina Eckblad 1968 (= *Tricharia* Boud. 1885 *fide* Species Fungorum 2017), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 15 species, type: *T. gilva* (Boud. ex Cooke) Eckblad, asexual morph unknown, hypogeous, terrestrial, temperate, see Benkert (2010; new species), Lumbsch and Huhndorf (2010; outline), Hansen (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted),

Stielow et al. (2013; phylogeny), Kušan et al. (2015b; new species), Lindemann and Böhning (2016; new species), sequences are available for voucher DED 7356 (SFSU) of the type: (*fide* Hansen et al. 2013).

Trichasterina G. Arnaud 1918 (= *Doguetia* Bat. & J.A. Lima 1960; = *Trichosia* Bat. & R. Garnier 1960 *fide* Species Fungorum 2017), *Asterinaceae*, *Asterinales*, *Dothideomycetes*, nine species, type: *T. styracis* (Theiss.) G. Arnaud, asexual morph unknown, foliar epiphytes, terrestrial, tropical, three species, cultures and sequences are unavailable, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Asterinaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014c; taxonomy, accepted as a genus in *Asterinaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichobolbus Bat. 1964, *Ascomycota* genera *incertae sedis*, one species, type: *T. palmae* Bat. & Cavalc., coelomycetous, sexual morph unknown, saprobes, on leaves of Palmae, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichobolus (Sacc.) Kimbr. & Cain 1967, *Ascodesmidaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *T. zukalii* (Heimerl) Kimbr., asexual morph unknown, saprobes, from soil, on dung, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Doveri (2012; new species, coprophilous, Italy), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Trichobotrys Penz. & Sacc. 1902, *Ascomycota* genera *incertae sedis*, four species, type: *T. pannosa* Penz. & Sacc., hyphomycetous, saprobes, from soil, terrestrial, Africa, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichocladium Harz 1871 (= *Harziella* Kuntze 1891 *fide* Species Fungorum 2017), *Chaetomiaceae*, *Sordariales*, *Sordariomycetes*, 32 species, type: *T. asperum* Harz, hyphomycetous, sexual morph unknown, terrestrial, aquatic, saprobes, worldwide, see Abdel-Wahab (2011b; phylogeny), Seifert et al. (2011; morphology), Abdel-Wahab and Bahkali (2012; phylogeny), Wijayawardene et al. (2012, 2017; outline), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), polyphyletic.

Trichocoma Jungh. 1838 (= *Trichoskytale* Corda 1842 *fide* Species Fungorum 2017), *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, two species, type: *T. paradoxa* Jungh., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Houbraken and Samson (2011; phylogeny), Kirk et al. (2013; genus

accepted), cultures available for the type: CBS 247.57, CBS 788.83 (*fide* Houbraken and Samson 2011).

Trichoconis Clem. 1909 (= *Appelia* (Sacc. & D. Sacc.) Trotter 1931; = *Pyricularia* sect. *Appelia* Sacc. & D. Sacc. 1906 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, 20 species, type: *T. caudata* (Appel & Strunk) Clem., hyphomycetous, saprobes, terrestrial, pantropic, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Brackel (2014; new species), cultures and sequences are available but lacking for the type.

Trichodelitschia Munk 1953, *Phaeotrichaceae*, *Phaeotrichales*, *Dothideomycetes*, four species, type: *T. bisporula* (P. Crouan & H. Crouan) Munk, asexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; outline, phylogeny), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures available for the type: CBS 262.69.

Trichoderma Pers. 1794 (= *Hypocrea* Fr. 1825; = *Aphysiostroma* Barrasa, A.T. Martínez & G. Moreno 1986 *fide* Rossman et al. 2013b; Jaklitsch and Voglmayr 2015), *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, c. 100 species, type: *T. viride* Pers., hyphomycetous, sexual morph *Hypocrea*, in soil, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2013b; nomenclature), Jaklitsch and Voglmayr (2015; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Trichodiscula Vouaux 1910, *Ascomycota* genera *incertae sedis*, one species, type: *T. lesdainii* Vouaux, coelomycetous, saprobes, terrestrial, France, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichodochium Syd. 1927, *Ascomycota* genera *incertae sedis*, one species, type: *T. disseminatum* Syd., hyphomycetous, saprobes, terrestrial, Asia, Central and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichodothella Petr. 1946, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. blumeri* Petr. ex S. Blumer, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Zhang et al. (2011d; morphology, accepted as a genus in *Venturiaceae*), Hyde et al. (2013) did not accept as a

genus in Venturiaceae, Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichodothis Theiss. & Syd. 1914, *Dothideomycetes* genera *incertae sedis*, three species, type: *T. comata* (Berk. & Ravenel) Theiss. & Syd., asexual morph unknown, saprobes, North America, Africa, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Venturiaceae*), Zhang et al. (2011d; provisionally included in *Venturiaceae*), Hyde et al. (2013) did not accept as a genus in *Venturiaceae*, Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichoglossum Boud. 1885, *Geoglossaceae*, *Geoglossales*, *Geoglossomycetes*, 20 species, type: *T. hirsutum* (Pers.) Boud., asexual morph unknown, on soil, saprobes, terrestrial, temperate, see Schoch et al. (2009; phylogeny, accepted as a genus in *Rutstroemiaceae*), Kučera et al. (2010; new records in Slovakia), Lumbsch and Huhndorf (2010; outline), Akata and Kaya (2013; new records in Turkey), Hustad et al. (2013; DNA, phylogeny), sequences are available for the type.

Trichohelotium Killerm. 1935, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, two species, need typification, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichomaris Hibbits, G.C. Hughes & Sparks 1981, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *T. invadens* Hibbits, G.C. Hughes & Sparks, on tanner crab, marine, Alaska, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Trichomatoclava G.F. Sepúlveda, Pereira-Carv. & Dianese 2009, *Ascomycota* genera *incertae sedis*, one species, type: *T. cerradensis* G.F. Sepúlveda, Pereira-Carv. & Dianese, hyphomycetous, saprobes, terrestrial, South America, see Pereira-Carvalho et al. (2009b; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of type: UB (Mycol. Col.) 19442, cultures and sequences are unavailable.

Trichomatomyces Dorn.-Silva & Dianese 2004, *Ascomycota* genera *incertae sedis*, one species, type: *T. byrsiniae* (Bat. & Peres) Dorn.-Silva & Dianese, hyphomycetous, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Trichomatosphaera Pereira-Carv., G.F. Sepúlveda & Dianese 2009, *Ascomycota* genera *incertae sedis*, one

species, type: *T. cerradensis* Pereira-Carv., G.F. Sepúlveda & Dianese, hyphomycetous, saprobes, terrestrial, South America, see Pereira-Carvalho et al. (2009b; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: UB (Mycol. Col.) 19597, cultures and sequences are unavailable.

Trichomerium Speg. 1918, *Trichomeriaceae*, *Chaetothyriales*, *Eurotiomycetes*, 32 species, type: *T. coffeicola* (Puttemans) Speg., asexual morph hyphomycetous, possibly *Tripospermum* Speg., epiphytes or saprobes, sooty moulds, terrestrial, Australia, Brazil, Thailand, see Lumbsch and Huhndorf (2010; outline), Chomnunti et al. (2012a; morphology; phylogeny), Liu et al. (2015a; morphology, phylogeny), Hongsanan et al. (2016b; asexual morph of *T. gloeosporum*), Hyde et al. (2016; morphology, phylogeny), cultures and sequences are available but lacking for type.

Trichometasphaeria Munk 1953, *Dothideomycetes* genera *incertae sedis*, eight species, type: *T. dianthi* (Rostr.) Munk, asexual morph coelomycetous, saprobes, North America, see Chlebicki (2009a; new species), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichonectria Kirschst. 1907, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, 19 species, type: *T. aculeata* Kirschst., asexual morph unknown, saprobes, lichenicolous, terrestrial, Austria, see Etayo and Sancho (2008; new species), Lumbsch and Huhndorf (2010; outline), Brackel (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Farkas and Flakus (2016; new species), cultures and sequences are available but lacking for the type.

Trichopeltella Höhn. 1910 (= *Raciborskiella* Speg. 1919 [1918] fide Species Fungorum 2017), *Trichopeltinaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *T. montana* (Racib.) Höhn., asexual morph unknown, epiphytes, Java, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014a; accepted as a genus in *Trichopeltinaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichopelttheca Bat., C.A.A. Costa & Cif. 1958, *Trichopeltinaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *T. asiatica* Bat., C.A.A. Costa & Cif., asexual morph unknown, epiphytes, Asia, Pacific, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014a; accepted as a genus in *Trichopeltinaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichopeltina Theiss. 1914 (= *Trichopeltina* Theiss., Centbl. Bakt. ParasitKde, Abt. II 39: 630 (1914) fide Species Fungorum 2017), *Trichopeltinaceae*,

Dothideomycetes families *incertae sedis*, two species, type: *T. labecula* (Mont.) Theiss., asexual morph unknown, epiphytes, South America, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014a; accepted as a genus in *Trichopeltinaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichopeltospora Bat. & Cif. 1958, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, two species, type: *T. pipericola* Bat., Cif. & C.A.A. Costa, asexual morph unknown, epiphytes, terrestrial, worldwide, no molecular data, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; taxonomy), Hongsanan et al. (2014c; keys), Wijayawardene et al. (2014c; outline), holotype of type: URM 2492, cultures and sequences are unavailable.

Trichopeltulum Speg. 1889 (= *Trichopeltium* Clem. 1909 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *T. pulchellum* Speg., coelomycetes, sexual morph unknown, saprobes, terrestrial, Hawaii, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichopeltum Bat., Cif. & C.A.A. Costa (1958) [1957], ?*Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, type: *T. hawaiiense* Bat. & C.A.A. Costa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wu et al. (2011; did not accept the placement).

Trichopeziza Fuckel 1870, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, 30 species, type: *T. sulphurea* (Pers.) Fuckel, asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), cultures available for the type: KUS-F52218.

Trichophaea Boud. 1885 (= *Dichobotrys* Hennebert 1973 *fide* Healy et al. 2016), *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, 32 species, type: *T. woolhopeia* (Cooke & W. Phillips) Boud., asexual morph unknown, hypogeous, saprobes, terrestrial, worldwide, see Benkert (2010; new species), Lumbsch and Huhndorf (2010; outline), Rubini et al. (2011; DNA, phylogeny), Hansen (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), Stielow et al. (2013; phylogeny), Perić et al. (2014; new species), sequences are available.

Trichophaeopsis Korf & Erb 1972, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, three species, type: *T. bicuspis* (Boud.) Korf & Erb, asexual morph unknown, hypogeous, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available.

Trichophyma Rehm 1904, *Arthoniales* genera *incertae sedis*, two species, type: *T. bunchosiae* Rehm, asexual morph unknown, lichenicolous, terrestrial, tropical, see

Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype of the type: E. Ule 1120 (S), cultures and sequences are unavailable.

Trichophyton Malmsten 1848 (see Species Fungorum 2017 for synonyms), *Arthrodermataceae*, *Onygenales*, *Eurotiomycetes*, 32 species, type: *T. tonsurans* Malmsten, hyphomycetous, sexual morph unknown, on humans and other animal, saprobes, on soil, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Hubka et al. (2014c; new species), Lorch et al. (2014; new species), Hiruma et al. (2016; DNA, mating types), culture and sequences are available for the type.

Trichoseptoria Cavara 1892 (= *Macroseptoria* Petr. 1923 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *T. alpei* Cavara, coelomycetes, sexual morph unknown, saprobes, terrestrial, Europe, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichospermella Speg. 1912, *Sordariomycetes* genera *incertae sedis*, one species, type: *T. pulchella* Speg., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Trichosphaerella E. Bommer, M. Rousseau & Sacc. 1891 (= *Bresadolella* Höhn. 1903; = *Larseniella* Munk 1942; = *Neorehmia* Höhn. 1902; = *Oplotheciosis* Bat. & Cif. 1963; = *Oplothecium* Syd. & P. Syd. 1923; = *Trichohleria* Sacc. 1908 *fide* Species Fungorum 2017), *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *T. decipiens* E. Bommer, M. Rousseau & Sacc., asexual morph unknown, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Whitton et al. (2012b; morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Lendemer et al. (2016; new species), cultures and sequences are unavailable for type.

Trichosphaeria Fuckel 1870 (= *Bakeromyces* Syd. & P. Syd. 1917; = *Pseudorhynchia* Höhn. 1909 *fide* Species Fungorum 2017), *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, c. 20 new species, type: *T. pilosa* (Pers.) Fuckel, saprobes, terrestrial, widespread, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for type: 089319.

Trichosphaeropsis Bat. & Nascim. 1960, *Sordariomycetes* genera *incertae sedis*, one species, type: *T. crescentiae* Bat. & Nascim., asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted),

Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Trichosporiella Kamyschko 1960, *Ascomycota* genera *incertae sedis*, two species, type: *T. hyalina* Kamyschko, hyphomycetous, sexual morph unknown, saprobes, oh human, on soil, terrestrial, Asia, Europe, see Seifert et al. (2011; morphology, accepted as a genus in *Sordariales*), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequence available for unidentified species.

Trichosporodochium Dorn.-Silva & Dianese 2004, *Ascomycota* genera *incertae sedis*, two species, type: *T. cerradensis* Dorn.-Silva & Dianese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trichothallus F. Stevens 1925 (= *Plokamidomyces* Bat. et al. 1958; *Trichopelttheca* Bat. et al. 1958 *fide* Rossman et al. 2016a), *Euantennariaceae*, *Dothideomycetes* families *incertae sedis*, two species, type: *T. hawaiiensis* F. Stevens, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, Oceania, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Trichothecium Link 1809 (= *Cephalothecium* Corda 1838; = *Hyphelia* Fr. 1825; = *Hyphoderma* Fr. 1849 = *Trichodermia* Hoffm. 1795; = *Trichotheciopsis* J.M. Yen 1979 *fide* Species Fungorum 2017), *Hypocreales* genera *incertae sedis*, nine species, type: *T. roseum* (Pers.) Link, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Summerbell et al. (2011; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Trichothelium Müll. Arg. 1885 (= *Actiniopsis* Starbäck 1899; = *Asteropeltis* Henn. 1904; = *Ophiodictyon* Sacc. & P. Syd. 1902; = *Stereochlamydomyces* Cif. & Tomas. 1953; = *Stereochlamys* Müll. Arg. 1885; = *Trichotheliomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Porinaceae*, *Ostropales*, *Lecanoromycetes*, 40 species, type: *T. epiphyllum* Müll. Arg., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Trichothyriella Theiss. 1914, *Dothideomycetes* genera *incertae sedis*, one species, type: *T. quercigena* (Berk. ex Cooke) Theiss., asexual morph unknown, terrestrial, South

East Asia, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Wu et al. (2011; treated as a doubtful genus), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichothyrinula Petr. 1950, *Trichopeltinaceae*, *Dothideomycetes* families *incertae sedis*, one species, type: *T. labecula* (Mont.) Theiss., asexual morph unknown, epiphytes, Ecuador, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; morphology), Kirk et al. (2013; genus accepted), Hongsanan et al. (2014a; accepted as a genus in *Trichopeltinaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Trichothyrium Speg. 1889 (= *Trichopeltis* Speg. 1889; = *Trichopeltopsis* Höhn. 1909; = *Trichopeltula* Theiss. 1914 *fide* Species Fungorum 2017), *Trichothyriaceae*, *Dothideomycetes* families *incertae sedis*, twelve species, type: *T. sarciniferum* Speg., asexual morph unknown, saprobes, terrestrial, subtropical, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Hyde et al. (2013; accepted as a genus in *Trichothyriaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Tricladiella K. Ando & Tubaki 1984, *Ascomycota* genera *incertae sedis*, one species, type: *T. pluvialis* K. Ando & Tubaki, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tricladiopsis Descals 1982, *Ascomycota* genera *incertae sedis*, two species, type: *T. foliosa* Descals, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, Europe, Central and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tricladiospora Nawawi & Kuthub. 1988, *Ascomycota* genera *incertae sedis*, three species, type: *T. stricta* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tricladium Ingold 1942, *Helotiaceae*, *Helotiales*, *Leotiomycetes*, 25 species, type: *T. splendens* Ingold, hyphomycetous, saprobes, aquatic, worldwide, see Campbell et al. (2009b; evolutionary relationships), Seifert et al. (2011, polyphyletic), Gulis et al. (2012; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Baschien et al. (2013;

molecular phylogeny), sequences are available for 15 species, compare with *Varicosporium*, holotype of type: CBS 234.46, MUCL 28105.

Trifurcospora K. Ando & Tubaki 1988, *Ascomycota* genera *incertae sedis*, two species, type: *T. irregularis* (Matsush.) K. Ando & Tubaki, hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trigonopsis Schachner emend. Kurtzman & Robnett 2007, *Trigonopsidaceae*, *Saccharomycetales*, *Saccharomycetes*, four species, type: *T. variabilis* Schachner, asexual reproduction is by multilateral budding, saprophytic, grape must, wine, beer, worldwide, cultures and sequences are available.

Trigonosporium Tassi 1900, *Ascomycota* genera *incertae sedis*, two species, type: *T. australiense* Tassi, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trimmatostruma Corda 1837, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, c. 30 species type: *T. salicis* Corda, sexual morph unknown, saprobes, terrestrial, aquatic, halophilic, worldwide, see Vaupotíč et al. (2008; physiology), Bartosik et al. (2010; biodeterioration), Diederich et al. (2010a; new species), El-Said et al. (2010; Egypt), Mansour (2010; Libya), Sepcic et al. (2010; metabolite profiling), Shenoy et al. (2010; phylogeny), Mohammadi (2011; spectrofluorometry), Ruiz-González et al. (2013; isolation protocol), Sarma and Raghukumar (2013; India), Gessler et al. (2014; review), Gunde-Cimerman and Zalar (2014; solar salterns), Heuchert and Braun (2014; new species), cultures and sequences are available.

Trimmatothele Norman ex Zahlbr. 1903, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, three species, type: *T. perquisita* (Norman) Norman ex Zahlbr., lichenized, see Lücking et al. (2017a; classification), sequences are available.

Trimmatothelopsis Zschacke 1934, *Acarosporaceae*, *Acarosporales*, *Lecanoromycetes*, one species, type: *T. versipellis* (Nyl.) Zschacke, lichenized, see Gueidan et al. (2014c; taxonomy), Roux et al. (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Trinacrium Riess 1852, *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, ten species, type: *T. subtile* Riess, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trinathotrema Lücking, Rivas Plata & Mangold 2011, *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, three species, type: *T. stictideum* (Nyl.) Lücking, R. Miranda & Kalb, asexual morph unknown, lichenized, terrestrial, Bolivia, Canary Iceland, see Boom and Ertz (2012; new species, new records), Flakus et al. (2013; taxonomy), Lücking et al. (2017a; classification); sequences are available.

Trinosporium Crous & Decock 2012, *Xylonomycetaceae*, *Xylonomycetales*, *Xylonomycetes*, *T. guianense* Crous & C. Decock, coelomycetous, sexual morph unknown, as a contaminant, see Crous et al. (2012e; taxonomy), Gazis et al. (2016; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Triplosphaeria Kaz. Tanaka & K. Hiray. 2009, *Triplosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *T. maxima* Kaz. Tanaka & K. Hirayama, asexual morph hyphomycetous, saprobes, terrestrial, Japan, see Tanaka et al. (2009; morphology, phylogeny), Lumbsch and Huhndorf (2010; outline), Wijayawardene et al. (2014c; outline), holotype and ex-type of the type: HHUF29330, JCM 13172.

Tripoconidium Subram. 1978, *Ascomycota* genera *incertae sedis*, one species, type: *T. aphanopagum* (Drechsler) Subram., hyphomycetous, sexual morph unknown, on nematodes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tripospermum Speg. 1918 (= *Pentaposporium* Bat. 1957 *fide* Species Fungorum 2017), *Neodevriesiaceae*, *Capnodiales*, *Dothideomycetes*, 27 species, type: *T. acerinum* (P. Syd.) P. Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Rocha et al. (2010; new species), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Quaedvlieg et al. (2014; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Tripospora Sacc. 1886 (= *Tripocorynelia* Kuntze 1898 *fide* Species Fungorum 2017), *Coryneliaceae*, *Coryneliales*, *Eurotiomycetes*, four species, type: *T. cookei* Sacc., asexual morph unknown, saprobes, South Africa, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Triposporina Höhn. 1912, *Ascomycota* genera *incertae sedis*, two species, type: *T. uredinicola* Höhn., hyphomycetous, sexual morph unknown, fungicolous, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Triposporium Corda 1837, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, 41 species, type: *T. elegans* Corda, hyphomycetous, sexual morph unknown, fungicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c; outline, accepted as a genus in *Asterinaceae*), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species, phylogeny, accepted as a *Helotiales*), Wijayawardene et al. (2017; outline), cultures and sequences are available but lacking for the type, taxonomic placement is based on sequence of non-type.

Tripterospora Subram. & Lodha 1968 (= *Cleistobombardia* J.H. Mirza 1968 *fide Species Fungorum* 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, six species, type: *T. coprophila* Subram. & Lodha, asexual morph unknown, saprobes, coprophilous, terrestrial, Asia, Africa, see Doveri (2010; new species and two varieties), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Triramulispora Matsush. 1975, *Ascomycota* genera *incertae sedis*, three species, type: *T. gracilis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Triscelophorus Ingold 1943, *Ascomycota* genera *incertae sedis*, eight species, type: *T. monosporus* Ingold, hyphomycetous, saprobes, aquatic, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Fiúza et al. (2015a; morphology), cultures and sequences are available.

Triscelosporium Nawawi & Kuthub. 1987, *Ascomycota* genera *incertae sedis*, one species, type: *T. verrucosum* Nawawi & Kuthub., hyphomycetous, sexual morph unknown, saprobes, aquatic, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: IMI 313480, cultures and sequences are unavailable.

Tristratiperidium Daranag., Camporesi & K.D. Hyde 2015, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *T. microsporum* Daranag., Camporesi & K.D. Hyde, asexual morph hyphomycetous, saprobes, terrestrial, Italy, Europe, one species, see Daranagama et al. (2016b; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), ex-type strains of type: MFLUCC 15-0413; holotype MFLU 15-0656.

Trisulcosporium H.J. Huds. & B. Sutton 1964, *Ascomycota* genera *incertae sedis*, one species, type: *T. acerinum* H.J. Huds. & B. Sutton, hyphomycetous,

sexual morph unknown, saprobes, aquatic, Africa, Asia, Europe, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trizodia Laukka 2010, *Ascomycota* genera *incertae sedis*, one species, type: *T. acrobia* Laukka, lichenized, terrestrial, Finland, see Stenroos et al. (2010; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Trochila Fr. 1849 (= *Pyrenotrochila* Höhn. 1917 *fide Species Fungorum* 2017), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, 15 species, type: *T. craterium* (DC.) Fr., asexual morph unknown, saprobes, terrestrial, temperate, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dermateaceae*), Kirk et al. (2013; genus accepted), Pärtel et al. (2017; DNA, phylogeny, accepted as a genus in *Cenangiaceae*), cultures available for the type: KL333.

Trochoideomyces Thaxt. 1931, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, one species, *T. gracilicaulis* Thaxt., asexual morph unknown, biotrophic, only known from *Trochoideus desjardinsii* (Coleoptera: Endomychidae), Malaysian Borneo, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trochophora R.T. Moore 1955 (= *Helicostilbe* Linder 1929 *fide Species Fungorum* 2017), *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *T. simplex* (Petch) R.T. Moore, hyphomycetous, sexual morph unknown, plant pathogens on *Daphniphyllum* shrubs and trees in Asia, terrestrial, see Crous et al. (2009a; morphology, phylogeny), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype NY00945981, for epitype see Videira et al. (2017).

Troglomyces S. Colla 1932, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, six species, type: *T. manfredii* S. Colla, asexual morph unknown, biotrophic on millipedes (Blaniulidae & Julidae), mostly Europe, Madeira, Tenerife, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Santamaría et al. (2014 morphology), Enghoff and Santamaría (2015, morphology, taxonomy, key, specificity and transmission), cultures and sequences are unavailable.

Tromeropsis Sherwood 1981, *Ascomycota* genera *incertae sedis*, one species, type: *T. microtheca* (P. Karst.) Sherwood, asexual morph unknown, saprobes, terrestrial, Finland, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Troposporella P. Karst. 1892 (= *Slimacomycetes* Minter 1986 *fide Species Fungorum 2017*), *Dothideomycetes* genera *incertae sedis*, four species, type: *T. fumosa* P. Karst., hyphomycetous, sexual morph unknown, saprobes, aquatic, Australasia, Europe, North and South America, see Tsui and Berbee (2010; new species, DNA, phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: MUCL 15695.

Troposporium Harkn. 1884, *Ascomycota* genera *incertae sedis*, one species, type: *T. album* Harkn., hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Troposporopsis Whitton, McKenzie & K.D. Hyde 1999, *Ascomycota* genera *incertae sedis*, two species, type: *T. atroapicis* Whitton, McKenzie & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Australasia, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Trullula Ces. 1852 (= *Cesatia* Rabenh. 1850; = *Epidochium* subgen. *Hormodochium* Sacc. 1886; = *Hormodochis* Clem. 1909; = *Hormodochium* (Sacc.) Höhn. 1911 *fide Species Fungorum 2017*), *Leotiomycetes* genera *incertae sedis*, 15 species, type: *T. oreoselini* Ces., coelomycetous, sexual morph unknown, saprobes, terrestrial, north temperate, see Wijayawardene et al. (2012, 2016b; outline, taxonomy, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type, taxonomic placement is based on sequences of non-type.

Truncatella Steyaert 1949 (= *Ahmadinula* Petr. 1953; = *Chailletia* Fuckel 1864; = *Pseudopestalotia* Elenkin & Ohl 1912 *fide Species Fungorum 2017*), *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, eleven species, type: *T. truncata* (Lév.) Steyaert, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2016b, 2017; outline, morphology), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2015; new species, phylogeny), Tibpromma et al. (2017b; new data), cultures and sequences are available.

Tryblidaria (Sacc.) Rehm 1903 (= *Blitridium* subgen. *Tryblidaria* Sacc. 1889 *fide Species Fungorum 2017*), *Patellariaceae*, *Patellariales*, *Dothideomycetes*, nine species, type: *T. breutelii* Rehm, asexual morph unknown, saprobes, terrestrial, subtropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted)

as a genus in *Patellariaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Yacharoen et al. (2015; accepted as a genus in *Patellariaceae*), cultures and sequences are unavailable.

Tryblidiopsis P. Karst. 1871 (= *Biatorellina* Henn. 1903; = *Tryblidis* Clem. 1909 *fide Species Fungorum 2017*), *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, three species, type: *T. pinastri* (Pers.) P. Karst., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; listed as *Tribliopsis*), Kirk et al. (2013; genus accepted), Wang et al. (2014a; DNA, phylogeny), cultures and sequences are available for the type: AFTOL-ID 1319.

Tryblidiopycnis Höhn. 1918, *Ascomycota* genera *incertae sedis*, one species, type: *T. pinastri* Höhn., coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tryblis Clem. 1931, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *T. arnoldii* (Rehm) Clem., asexual morph unknown, ?saprobes, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable

Trypetheliopsis Asah. 1937, *Monoblastiaceae*, *Monoblastiales*, *Dothideomycetes*, six species, type: *T. boninensis* Asah., asexual morph unknown, lichenized, terrestrial, tropical, new synonym *Musaespora* Aptroot & Sipman, see Kashiwadani et al. (2009; taxonomy), Nelsen et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Trypethelium Spreng. [nom. cons.] 1804 (= *Bathelium* Ach. 1803; = *Holstiella* Henn. 1895; = *Leightonia* Trevis. 1861; = *Phyllothelium* Trevis. 1861; = *Porophora* Zenker 1827; = *Trypetheliomyces* Cif. & Tomas. 1953 *fide Species Fungorum 2017*), *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, 16 species, type: *T. eluteriae* Spreng., asexual morph unknown, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Aptroot and Lücking (2016; taxonomy, monograph), Lücking et al. (2017a; classification); sequences are available.

Tryssglobulus B. Sutton & Pascoe 1987, *Ascomycota* genera *incertae sedis*, one species, type: *T. aspergilloides* B. Sutton & Pascoe, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tubakia B. Sutton 1973 (= *Actinopelti* Sacc. 1913 *fide Species Fungorum 2017*), *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, eight species, type: *T. japonica*

(Sacc.) B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; treated as coelomycetous), Yun and Rossman (2011; new species), Harrington et al. (2012; new species), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Braun et al. (2014a; new species), cultures and sequences are available but lacking for the type.

Tubakiella (Tubaki) Sakay., K.L. Pang & E.B.G. Jones 2011, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *T. galerita* (Tubaki) Sakay., K.L. Pang & E.B.G. Jones, asexual morph unknown, saprobes, marine, France, Denmark, Germany, Japan, Taiwan, USA, Yugoslavia, see Sakayaroj et al. (2011a; morphology, phylogeny), Jones et al. (2015; outline), Maharachchikumbura et al. (2015, b; outline, phylogeny), holotype and ex-type of the type: BCC33500, PP5577.

Tuber P. Micheli ex F.H. Wigg. 1780 (= *?Loculotuber* Trappe, Parladé & I.F. Alvarez 1993 *fide* Jaklitsch et al. 2016a; see Species Fungorum 2017 for additional synonyms), *Tuberaceae*, *Pezizales*, *Pezizomycetes*, 135 species, type: *T. aestivum* Vittad., asexual morph unknown, saprobes, saprobes, hypogeous, terrestrial, worldwide, see Bonito et al. (2010a, b; new species), Culleré et al. (2010; aromatic compounds), Lumbsch and Huhndorf (2010; outline), Benucci et al. (2011; ectomycorrhizal), Fan et al. (2013; new species), Kirk et al. (2013; genus accepted), Li et al. (2014a; new species, China), Su et al. (2014; new species), Guevara-Guerrero et al. (2015; new species), cultures and sequences are available.

Tuberculispora Deighton & Piroz. 1972, *Ascomycota* genera *incertae sedis*, one species, type: *T. jamaicensis* Deighton & Piroz., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tubeufia Penz. & Sacc. 1898 (= *Linobolus* Syd. & P. Syd. 1917 *fide* Species Fungorum 2017), *Tubeufiaceae*, *Tubeufiales*, *Dothideomycetes*, 29 species, type: *T. javanica* Penz. & Sacc., saprobes on decaying wood or sheaths of bamboo, widespread, asexual morph hyphomycetous, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Boonmee et al. (2014b; taxonomy, phylogeny), Wijayawardene et al. (2014c, 2017; outline), Dai et al. (2017, new species, phylogeny), Doilom et al. (2017, new species, phylogeny), Lu et al. (2017a, new species, phylogeny), Luo et al. (2017, phylogeny, new species), cultures and sequences are available, ex-type strains of type: MFLUCC12-0545; epitype of type: MFLU13-0371.

Tubolachnum Velen. 1934, *Lachnaceae*, *Helotiales*, *Leotiomycetes*, two species, type: need typification, asexual morph unknown, saprobes, terrestrial, former Czechoslovakia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tulipispora Révay & J. Gönczöl 2009, *Ascomycota* genera *incertae sedis*, one species, type: *T. ingoldii* Révay & J. Gönczöl, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: BP 99766, cultures and sequences are unavailable.

Tumidispora Hongsanan & K.D. Hyde 2015, *Microthyriaceae*, *Microthyriales*, *Dothideomycetes*, one species, type: *T. shoreae* Hongsanan & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Thailand, see Ariyawansa et al. (2015a; taxonomy, phylogeny), Hongsanan et al. (2016c; evolution), holotype and ex-type of the type: MFLU 15-1391, MFLUCC 12-0409.

Tumularia Descals & Marvanová 1987, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *T. tuberculata* (J. Gönczöl) Descals & Marvanová, hyphomycetous, sexual morph unknown, saprobes, aquatic, Europe, North and South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; morphology, phylogeny), Hyde et al. (2013; accepted as a genus in *Lophiostomataceae*), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Tunicago B. Sutton & Pollack 1977, *Ascomycota* genera *incertae sedis*, two species, type: *T. uniolae* B. Sutton & Pollack, coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, Australia, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tunicatispora K.D. Hyde 1990, *Halosphaeriaceae*, *Microascales*, *Sordariomycetes*, one species, type: *T. australiensis* K.D. Hyde, saprobes, marine, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Tunstallia Agnihothr. 1961, *Sordariomycetes* genera *incertae sedis*, one species, type: *T. aculeata* (Petch) Agnihothr., asexual morph unknown, saprobes, pathogens, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Turgidosculum Kohlm. & E. Kohlm. 1972, *Ascomycota* genera *incertae sedis*, two species, type: *T. ulvae* (M. Reed)

Kohlm. & E. Kohlm., asexual morph unknown, symbiotic, marine, worldwide, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Turturconchata J.L. Chen, T.L. Huang & Tzean 1999, *Ascomycota* genera *incertae sedis*, two species, type: *T. reticulata* J.L. Chen, T.L. Huang & Tzean, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Tylophorella Vain. 1890 (= *Tylophorellomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *T. polyspora* Vain., lichenized, terrestrial, Bolivia, New guinea, see Lumbsch et al. (2009b; molecular), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), McCarthy (2014; new record), Lücking et al. (2017a; classification), sequences are unavailable.

Typhorion Nyl. ex Stizenb. 1862 (= *Ditylis* Clem. 1909; = *Sporodochiolichen* Aptroot & Sipman 2011 *fide* Diederich et al. 2013; Species Fungorum 2017), *Arthoniaceae*, *Arthoniales*, *Lecanoromycetes*, eight species, type: *T. protrudens* Nyl., lichenized, terrestrial, mainly tropical, see Lumbsch et al. (2009b; molecular), Lumbsch and Huhndorf (2010; outline), Aptroot et al. (2010; biodiversity), Ertz et al. (2011; molecular), Reddy et al. (2011; new record), Sethy et al. (2012; new record), Kirk et al. (2013; genus accepted), Lendemer et al. (2013; new species), Lendemer and Harris (2014; taxonomy), Pérez-Pérez et al. (2015; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Tylotballia P. James & H. Kilias 1981, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, three species, type: *T. biformigera* (Leight.) P. James & H. Kilias, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Elix and Kantvilas (2013; new records), Kirk et al. (2013; genus accepted), Øvstedral and Gremmen (2014; new combination), Lücking et al. (2017a; classification), sequences are available.

Tympanis Tode 1790 (= *Sirodothis* Clem. 1909; *Pleurophomella* Höhn 1914 *fide* Johnston et al. 2014b), *Tympanidaceae*, *Phacidiales*, *Leotiomycetes*, 27 species, type: *T. saligna* Tode, asexual morphs formerly known in *Sirodothis* and *Pleurophomella*, weak parasite of gymno- and angiosperm trees commonly associated with cankers, worldwide, common in temperate areas, Kirk et al. (2013; genus accepted), Quijada et al. (2015; phylogeny), cultures and sequences are available but lacking for the type.

Tympanopsis Starbäck 1894, *Scortechiniaceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *T. euomphala* (Berk. & M.A. Curtis) Starbäck, asexual morph unknown, saprobes, cosmopolitan, see Mugambi and Huhndorf (2010; phylogeny), Vasilyeva et al. (2012; new species), Maharachchikumbura et al. (2015b, 2016b; outline), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Tympansporitum W. Gams 1974, *Ascomycota* genera *incertae sedis*, one species, type: *T. parasiticum* W. Gams, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Typhulochaeta S. Ito & Hara 1915, *Erysiphaceae*, *Erysiphales*, *Leotiomycetes*, four species, type: *T. japonica* S. Ito & Hara, biotrophic, obligate plant pathogens, terrestrial, Asia, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are available.

Tyrannicordyceps Kepler & Spatafora 2012, *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, five species, type: *T. fraticida* (Tanda & Kobayasi) Kepler & Spatafora, asexual morph unknown, saprobes, terrestrial, cosmopolitan, see Kepler et al. (2012b; taxonomy), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Tyrannosorus Unter. & Malloch 1995, *Venturiaceae*, *Venturiales*, *Dothideomycetes*, one species, type: *T. pinicola* (Petrini & P.J. Fisher) Unter. & Malloch, asexual morph unknown, saprobes, terrestrial, Canada, Pakistan, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideomycetes* genera *incertae sedis*), Zhang et al. (2011d; accepted as in *Venturiaceae*), Hyde et al. (2013; accepted as in *Venturiaceae*, phylogeny), Wijayawardene et al. (2014c; outline), cultures available for the type: AFTOL-ID 1235.

Uberispora Piroz. & Hodges 1973, *Ascomycota* genera *incertae sedis*, four species, type: *U. simplex* (Ichinoe) Piroz. & Hodges, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, Asia, Caribbean, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Jian et al. (2014; new species), Ma et al. (2014; new species, China), cultures and sequences are unavailable.

Ubrizsya Negru 1965, *Ascomycota* genera *incertae sedis*, one species, type: *U. oxycoeci* Negru & R. Sandor, coelomycetous, sexual morph unknown, saprobes, terrestrial, Rumania, see Wijayawardene et al.

(2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Uleodothis Theiss. & Syd. 1915 (= *Didothis* Clem. 1931 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, four species, type: *U. balansiana* (Sacc., Roum. & Berl.) Theiss. & Syd., saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Uleomyces Henn. 1895 (= *Dictyomollisia* Rehm 1909; = *Hyalocurreya* Theiss. & Syd. 1915; = *Kusanoa* Henn. 1900; = *Kusanoopsis* F. Stevens & Weedon 1923; = *Micromyriangium* Petr. 1929; = *Myriangina* (Henn.) Höhn. 1909; = *Myrianginella* F. Stevens & Weedon 1923; = *Myriangium* sect. *Myriangina* Henn. 1902 *fide* Species Fungorum 2017), *Cookellaceae*, *Myriangiales*, *Dothideomycetes* families *incertae sedis*, four species, type: *U. parasiticus* Henn., asexual morph unknown, on rusts and ascomycetes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Cookellaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Uleoporthe Petr. 1941, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *U. orbiculata* (Syd. & P. Syd.) Petr., asexual morphs unknown, saprobes, terrestrial, America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Senanayake et al. (2017, new species, placement), cultures and sequences are unavailable.

Uleothyrium Petr. 1929, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, two species, type: *U. amazonicum* Petr., asexual morph coelomycetous (= septothyrella-like, *fide* Hofmann 2009), epiphytes, terrestrial, worldwide, see Hofmann (2009; review), Lumbsch and Huhndorf (2010; outline), Wu et al. (2011; taxonomy), Hongsanan et al. (2014c; keys), Wijayawardene et al. (2014c; outline), placement in *Asterinaceae* supported by morphology, cultures and sequences are unavailable, holotype of type: W 10746.

Ulocoryphus Michaelides, L. Hunter & W.B. Kendr. 1982, *Ascomycota* genera *incertae sedis*, one species, type: *U. mastigophorus* Michaelides, L. Hunter & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Ulospora D. Hawksw., Malloch & Sivan. 1979, *Testudinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *U. bilgramii* (D. Hawksw., C. Booth & Morgan-Jones) D. Hawksw., Malloch & Sivan., marine

sediments and plant material, Asia, terrestrial and marine, see Zhang et al. (2008b; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures available, holotype of type: K(M)-IMI 12081.

Ulurua Trappe, Claridge & Kovács 2010, *Pezizaceae*, *Pezizales*, *Pezizomycetes*, one species, type: *U. nonparaphysata* Trappe & Claridge, asexual morph unknown, hypogeous, terrestrial, Northern Territory, see Trappe et al. (2010b; taxonomy, phylogeny), isotype of the type: OSC 131126, sequences are available.

Umbellidion B. Sutton & Hodges 1975, *Ascomycota* genera *incertae sedis*, one species, type: *U. radulans* B. Sutton & Hodges, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Umbilicaria Hoffm [nom. cons.] 1789 (see Species Fungorum 2017 for synonyms), *Umbilicariaceae*, *Umbilicariales*, *Lecanoromycetes*, 70 species, type: *U. exasperata* Hoffm., asexual morph unknown, lichenized, terrestrial, worldwide, see Hájek et al. (2009; physiological exp.), Hestmark (2009, 2010, 2014a, b, 2016; taxonomy, new records), Krzewicka and Wilk (2009; new species), Krzewicka et al. (2009; taxonomy), Lee et al. (2009a; bioactivity study), Luo et al. (2009c; antioxidant study), Ranković et al. (2009; bioactivity study), Seo et al. (2009a; bioactivity study), Hestmark (2010; taxonomy), Hestmark et al. (2010, 2014a, b; phylogeny), Kim et al. (2010; bioactivity study), Krzewicka (2010; combination), Lumbsch and Huhndorf (2010; outline), Manojlović et al. (2011; antimicrobial study), Harańczyk et al. (2012; experimental study), Jupa et al. (2012; physiology), Kosanić et al. (2012; bioactivity study), McCune and Curtis (2012; taxonomy), Bendiksby and Timdal 2013, Buçukoglu et al. (2013; antioxidant study), Singh and Ravindra (2013; environmental study), Kirk et al. (2013; genus accepted), Park et al. (2013; mutagenic study), Hestmark (2014a, b; taxonomy), McCune et al. (2014; new species), Tibell (2014; taxonomy), Wang et al. (2014d; antithrombotic study), Cao et al. (2015; molecular), Du et al. (2015; bioactivity study), Bogdanović et al. (2016; floristic), Davydov and Ohmura (2016; taxonomy), Hestmark (2016; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Umbilithecium Etayo 2002, *Lecanoromycetes* genera *incertae sedis*, one species, type: *U. pseudocyphellariae* Etayo, asexual morph unknown, parasymbiont on macrolichens of *Pseudocyphellaria*, terrestrial, South America, see Etayo and Sancho (2008; morphology, distinction of similar lichenicolous genera), needs

revision, holotype of the type: MAF Etayo 23090, cultures and sequences are unavailable.

Umthunziomyces Crous & M.J. Wingf. 2016, *Planistromellaceae*, *Botryosphaerales*, *Dothideomycetes*, one species, type: *U. hagahagensis* Crous & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Crous et al. (2016a; taxonomy, phylogeny), holotype and ex-type of the type: CBS H-22873, CPC 29917.

Umushamyces Etayo 2002, *Scoliciosporaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *U. kuturnum* Etayo, asexual morph unknown, parasymbiont on *Coccotrema*, terrestrial, South America, see Etayo and Sancho (2008; morphology, distinction of similar lichenicolous genera), need revision, holotype of the type: MAF Etayo 22460, cultures and sequences are unavailable.

Uncinocarpus Sigler & G.F. Orr 1976 (= *Brunneospora* Guarro & Punsola 1987 *fide* Species Fungorum 2017), *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, three species, type: *U. reesii* Sigler & G.F. Orr, asexual morph unknown, saprobes, terrestrial, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are available for the type (unpublished).

Uncispora R.C. Sinclair & Morgan-Jones 1979, *Chaetothyriales* genera *incertae sedis*, *Eurotiomycetes*, three species, type: *U. harroldiae* R.C. Sinclair & Morgan-Jones, hyphomycetous, sexual morph unknown, saprobes, terrestrial, China, North America, see Seifert et al. (2011; morphology), Yang et al. (2011a; new species, China, DNA), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Li et al. (2015a; new species; DNA), cultures and sequences are available but lacking for the type, generic placement is based on no-type.

Underwoodia Peck 1890 (= *Geomorium* Speg. 1922 *fide* Species Fungorum 2017), ?*Helvellaceae*, *Pezizales*, *Pezizomycetes*, three species, type: *U. columnaris* Peck, asexual morph unknown, saprobes, from soil, terrestrial, worldwide, see Bonito et al. (2013; DNA, phylogeny, stated that the genus is polyphyletic), Kirk et al. (2013; genus accepted), sequences are available for the type: U42658 (SSU).

Unguicularia Höhn. 1905 (= *Catinella* Kirschst. 1924; = *Chytrella* Kirschst. 1941; = *Mycopandora* Velen. 1947; = *Phalothrix* Clem. 1909 *fide* Species Fungorum 2017), *Hyaloscypheaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *U. unguiculata* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available (unpublished).

Unguiculariella K.S. Thind & R. Sharma 1990, *Hyaloscypheaceae*, *Helotiales*, *Leotiomycetes*, one species,

type: *U. bhutanica* K.S. Thind & R. Sharma, asexual morph unknown, saprobes, terrestrial, Bhutan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Unguiculariopsis Rehm 1909 (= *Encoeliella* Höhn. 1910; = *Deltosperma* W.Y. Zhuang 1988; = *Mollisia* subgen. *Mollisiella* W. Phillips 1887; *Mollisiella* (W. Phillips) Massee 1895 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Cordieritidaceae*, *Helotiales*, *Leotiomycetes*, 25 species, type: *U. ilicincola* (Berk. & Broome) Rehm, asexual morph formerly known in *Deltosperma*, fungicolous and lichenicolous, terrestrial, worldwide, see Etayo and Sancho (2008; new species), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Helotiaceae*), Brackel (2011a; new species), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), Pärtel et al. (2017; DNA, phylogeny), Wijayawardene et al. (2017; outline, taxonomy), cultures and sequences are available but lacking for the type.

Unguiculella Höhn. 1906 (= *Globulina* Velen. 1934 *fide* Species Fungorum 2017), *Hyaloscypheaceae*, *Helotiales*, *Leotiomycetes*, 17 species, type: *U. falcipila* Höhn., asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Uniseta Ciccar. 1948, *Ascomycota* genera *incertae sedis*, one species, type: *U. flagellifera* (Ellis & Everh.) Ciccar., coelomycetous, sexual morph unknown, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Unisetosphaeria Pinnoi, E.B.G. Jones, McKenzie & K.D. Hyde 2003, *Trichosphaeriaceae*, *Trichosphaerales*, *Sordariomycetes*, one species, type: *S. styracum* Speg., asexual morph unknown, saprobes, terrestrial, Thailand, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Urceolella Boud. 1885 (= *Pilatia* Velen. 1934; = *Urceolella* subgen. *Urceolella* Boud. 1885 *fide* Species Fungorum 2017), *Hyaloscypheaceae*, *Helotiales*, *Leotiomycetes*, 24 species, type: *U. crispula* (P. Karst.) Boud., asexual morph unknown, saprobes, terrestrial, worldwide, see Baral et al. (2009; new species), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Han et al. (2014; DNA, phylogeny), sequences are available for voucher TNS-F18014 of the type.

Uredinophila Rossman 1987, *Dothideomycetes* genera *incertae sedis*, two species, type: *U. tropicalis* (Speg.) Rossman, asexual morph unknown, parasitic on *Pucciniales* on leaves of *Dryopteris tetragona*,

Venezuela, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Urnula Fr. 1849, *Sarcosomataceae*, *Pezizales*, *Pezizomycetes*, ten species, type: *U. craterium* (Schwein.) Fr., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Pfister et al. (2008; phylogeny), Lumbsch and Huhndorf (2010; outline), Carbone et al. (2013; new species, DNA, phylogeny), Kirk et al. (2013; genus accepted), Wang and Huang (2015; new species, DNA, phylogeny), sequences are available for voucher DHP 04-511 (FH) of the type.

Urohendersonia Speg. 1902 (= *Uroconis* Clem. 1902 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, five species, type: *U. platensis* Speg., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, South America, India, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Urohendersoniella Petr. 1955, *Ascomycota* genera *incertae sedis*, one species, type: *U. mastigospora* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Australia, see Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Uropolystigma Maubl. 1920, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *U. atrostestaceum* Maubl., in living leaves, Brazil, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Urosporella G.F. Atk. 1897 (= *Fabreola* Kuntze 1891; = *Sauvageautia* Har. 1892; = *Urospora* Fabre 1879 [1878] *fide* Species Fungorum 2017), *Sordariomycetes* genera *incertae sedis*, two species, type: *U. americana* G.F. Atk., cosmopolitan, Lumbsch and Huhndorf (2010; outline, treated as *Amphisphaeriaceae*), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Urosporellopsis W.H. Hsieh, Chi Y. Chen & Sivan. 1994, *Xylariomycitidae* genera *insertae sedis*, *Sordariomycetes*, one species, type: *U. taiwanensis* W.H. Hsieh, Chi Y. Chen & Sivan., asexual morph unknown, saprobes, Taiwan, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype IMI 354629, cultures and sequences are unavailable.

Ursicollum Gryzenh. & M.J. Wingf. 2006, *Cryphonectriaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *U. fallax* Gryzenh. & M.J. Wingf., coelomycetous, sexual morph unknown, pathogens, saprobes, terrestrial, USA, see Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al.

(2015b, 2016b; outline), holotype and ex-type cultures of type: PREM 58840, CMW 18119 = CBS 118663.

Urupe Viégas 1944, *Sordariomycetes* genera *incertae sedis*, one species, type: *U. guaduae* Viégas, asexual morph unknown, saprobes, terrestrial, Central America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded from *Meliolaceae* and placed in *Sordariomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Usnea Dill. ex Adans. 1763 (= *Alectoria* Link 1833; = *Eumitria* Stirt. 1881; = *Neuropogon* Nees & Flot. 1835 [1834]; = *Usnomyces* E.A. Thomas ex Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, c. 350 species, type: *U. florida* (L.) Weber ex Wigg., asexual morph unknown, lichenized, terrestrial, worldwide, see Verma et al. (2008; antioxidant and hepatoprotective activity), Lumbsch and Huhndorf (2010; outline), Çobanoğlu (2011, new records), Kelly et al. (2011; DNA barcoding), Li et al. (2011, Kirk et al. (2013; accepted as a valid genus), Truong et al. (2013; molecular, species delimitation), Ohmura (2014; New to Asia), Shukla et al. (2014b; taxonomy, India), Çobanoğlu et al. (2016; antimicrobial activity), Lücking et al. (2017a; classification), Mark et al. (2016; circumscribed species, phylogeny), sequences are available.

Usnochroma Søchting, Arup & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *U. carphineum* (Fr.) Søchting, Arup & Frödén, asexual morph unknown, lichenized, terrestrial, France, South Africa, see Arup et al. (2013; taxonomy, phylogeny), Gaya et al. (2015; DNA), Lücking et al. (2017a; classification), sequences are available. cultures available for the type: Ccar201 (*fide* Gaya et al. 2015).

Ustilaginoidea Bref. 1895 (= *Dubiomycetes* Lloyd 1895; = *Villosiclava* E. Tanaka & C. Tanaka 2009 *fide* Rossman et al. 2016a), *Clavicipitaceae*, *Hypocreales*, *Sordariomycetes*, six species, type: *U. oryzae* (Pat.) Bref., hyphomycetous, sexual morph unknown, saprobes, terrestrial, subtropical, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are available.

Utriascus Réblová 2003, *Sordariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *U. gabretiae* Réblová, asexual morph unknown, lichenicolous, terrestrial, Czech Republic, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), cultures and sequences are unavailable.

Uvarispora Goos & Piroz. 1975, *Ascomycota* genera *incertae sedis*, one species, type: *U. lignicola* Goos & Piroz., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Central America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Uwebraunia Crous & M.J. Wingf. 1996, *Dissconiaceae*, *Capnodiales*, *Dothideomycetes*, seven species, type: *U. juvenis* Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, worldwide, see Seifert et al. (2011; morphology), Li et al. (2012a; phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Uwemyces Hern.-Restr., G.A. Sarria & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *U. elaeidis* Hern.-Restr., G.A. Sarria & Crous, hyphomycetous, associated with leaf spots, terrestrial, Australia, see Crous et al. (2016b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), culture and sequences are available.

Vagicola Chethana & K.D. Hyde 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *V. vagans* (Niessl) O. Erikss., Chethana & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, worldwide, see Ariyawansa et al. (2015a; taxonomy), Jayasiri et al. (2015; new species), Thambugala et al. (2017b, new species, phylogeny), holotype of type species UPS:BOT:F-169873, cultures and sequences are available for the type species.

Vaginatispore K.D. Hyde 1995, *Lophiostomataceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *V. aquatica* K.D. Hyde, asexual morph unknown, saprobic, fresh water or terrestrial, Hong Kong, Japan, Thailand, all species have sequence data, see Kirk et al. (2013; genus accepted), Thambugala et al. (2015b; morphology, phylogeny, accepted as a genus in *Lophiostomataceae*), Wanasinghe et al. (2016b; morphology, phylogeny), Tibpromma et al. (2017b; new data), ex-type strains of type: MFLUCC 11-0083 = IFRDCC 2462; holotype MFLU 11-1115.

Vagnia D. Hawksw. & Miadl. 1997, *Ascomycota* genera *incertae sedis*, one species, type: *V. cirriformia* D. Hawksw. & Miadl., lichenicolous, Europe, terrestrial, see Kirk et al. (2013; genus accepted), culture and sequences are unavailable, holotype of type: K(M)-IMI 371358.

Vahliella P.M. Jørg. 2008, *Vahliellaceae*, *Peltigerales*, *Lecanoromycetes*, eight species, type: *V. leucophaea* (Vahl) P.M. Jørg., asexual morph unknown, lichenized, terrestrial, Europe, see Jørgensen (2008; taxonomy) Lumbsch and Huhndorf (2010; outline), Wedin et al. (2011; new species), Wedin et al. (2011; phylogeny),

Perez-Vargas et al. (2014; new species), Lücking et al. (2017a; classification), cultures and sequences are available, sequences are available.

Vainionora Kalb 1991, *Lecanoraceae*, *Lecanorales*, *Lecanoromycetes*, nine species, type: *V. pallidostraminea* (Vain.) Kalb, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Papong and Lumbsch (2011; taxonomy), Menezes et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Papong et al. (2013; molecular), Lücking et al. (2017a; classification), sequences are unavailable.

Valdensia Peyronel 1923 (= *Asterobolus* Redhead & P.W. Perrin 1972; = *Valdensinia* Peyronel 1953 *fide* Johnston et al. 2014b), *Sclerotiniaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *V. heterodoxa* Peyronel, asexual morph formerly known in *Valdensinia*, saprobes, terrestrial, Europe, Japan, see Lumbsch and Huhndorf (2010; outline), Nekoduka et al. (2012; pathogens, Japan), Kirk et al. (2013; genus accepted), cultures available for the type: HBI0401.

Valetoniella Höhn. 1909, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *V. crucipila* Höhn., asexual morph unknown, saprobes, or fungicolous, terrestrial, tropics, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Valetoniellopsis Samuels & M.E. Barr 1998, *Niessliaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *V. laxa* Samuels & M.E. Barr, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 191.97.

Valsalnicola D.M. Walker & Rossman 2012, *Diaporthales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *V. oxystoma* (Rehm) D. Walker & Rossman, asexual morph dimorphic, pathogens, saprobes, terrestrial, Europe, America, see Crous et al. (2012c; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), lectotype of the type: BPI 884138, cultures available for the type: AR4833.

Valsaria Ces. & De Not. 1863, *Valsariaceae*, *Valsariales*, *Dothideomycetes*, 140 epithets, type: *V. insitiva* (Tode) Ces. & De Not. 1863, asexual morph coelomycetous, saprobes, terrestrial, widely distributed, see Lumbsch and Huhndorf (2010; outline), Jaklitsch et al. (2015; sexual and asexual morphs, morphology, phylogeny), epitype of the type: WU 33462,

Vamsapriya Gawas & Bhat 2006, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, four species, type: *V. indica* Gawas & Bhat, hyphomycetous, saprobes, terrestrial, India,

Thailand, see Pratibha and Bhat (2008, morphology), Dai et al. (2014b; asexual morph, morphology, phylogeny, 2017; sexual morph, morphology, phylogeny), ex-epitype culture of type: MFLUCC 12–0544 = ICMP 20734.

Vanakripa Bhat, W.B. Kendr. & Nag Raj 1993, *Ascomycota* genera *incertae sedis*, nine species, type: *V. gigaspora* Bhat, W.B. Kendr. & Nag Raj, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Brazil, see Hu et al. (2010a; new species, China), Mel'nik (2011; new species, Vietnam), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Leão-Ferreira et al. (2013; new species; Brazil), cultures and sequences are unavailable.

Vanbeverwijkia Agnihothr. 1961, *Ascomycota* genera *incertae sedis*, one species, type: *V. spirospora* Agnihothr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Assam, USA, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Vanderwaltozyma Kurtzman 2003, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *V. polyspora* (van der Walt) Kurtzman, asexual reproduction is by multilateral budding, saprophytic, soil, cactus fruit, tanning fluids, worldwide, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Vanderystiella Henn. 1908 (= *Deightonia* Petr. 1947 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *V. leopoldvilleana* Henn., coelomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Vanterpoolia A. Funk 1982, *Ascomycota* genera *incertae sedis*, one species, type: *V. tsugae* A. Funk, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Vargamycetes Tóth 1980, *Amniculicolaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *V. aquaticus* (Dudka) Tóth, hyphomycetous, sexual morph unknown, saprobes, aquatic, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny, designate lectotype and epitype), sequences are available.

Varicellaria Nyl. 1858 (= *Varicellariomyces* Cif. & Tomas. 1953 *fide* Species Fungorum 2017), *Ochrolechiaceae*, *Pertusariales*, *Lecanoromycetes*, seven species, type: *V. microsticta* Nyl., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010;

outline), Schmitt et al. (2012; circumscription), Kirk et al. (2013; genus accepted), Ren and Li (2013; new record), Miądlikowska et al. (2014a; DNA), Lücking et al. (2017a; classification), cultures and sequences are unavailable for type.

Varicosporium W. Kegel 1906, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, eleven species, type: *V. elodeae* W. Kegel, hyphomycetous, sexual morph unknown, in soil, saprobes, terrestrial, aquatic, worldwide, see Campbell et al. (2009b; phylogeny, polyphyletic), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 541.92.

Variocladium Descals & Marvanová 1998, *Mollisiaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *V. rangiferinum* (Descals) Descals & Marvanová, hyphomycetous, sexual morph unknown, saprobes, terrestrial, aquatic, temperate, see Campbell et al. (2009b; phylogeny), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 508.71 (unpublished).

Variospora Arup, Søchting & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, twelve species, type: *V. velana* (A. Massal.) Arup, Søchting & Frödén, asexual morph unknown, lichenized, terrestrial, Europe, North Africa, see Arup et al. (2013; taxonomy, phylogeny), Kondratyuk et al. (2014; DNA), Gaya et al. (2015; DNA, phylogeny), Lücking et al. (2017a; classification); cultures available for the type: Cavel192, sequences are available.

Vasudevella Chona, Munjal & Bajaj 1957, *Ascomycota* genera *incertae sedis*, one species, type: *V. sporoboli* Chona, Munjal & Bajaj, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Velloziomyces Armando, Z.M. Chaves & Dianese 2015, *Ascomycota* genera *incertae sedis*, one species, type: *V. ramosiconidialis* Armando, Z.M. Chaves & Dianese, hyphomycetous, sexual morph unknown, on living leaves, terrestrial, Brazil, see de Souza Armando et al. (2015; morphology), holotype of the type: UBMC 8849, cultures and sequences are unavailable, compare with *Dictyodochium*, *Minimidochium* and *Iyengarina*.

Velutaria Fuckel 1870 (= *Schweinitzia* Massee 1895 *fide* Species Fungorum 2017), *Pezizellaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *V. griseovitellina* (Fuckel) Fuckel, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al.

(2016a; classification), cultures and sequences are unavailable.

Velutarina Korf 1971 (= *Velutarina* Korf 1953 *fide* Species Fungorum 2017), *Cenangiaceae*, *Helotiales*, *Leotiomycetes*, three species, type: *V. rufo-olivacea* (Alb. & Schwein.) Korf, asexual morph unknown, saprobes, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Perić et al. (2015; new species), Jaklitsch et al. (2016a; classification), Pärte et al. (2017; DNA, phylogeny), cultures and sequences are available.

Velutipila D. Hawksw. 1987, *Ascomycota* genera *incertae sedis*, one species, type: *V. poeltii* D. Hawksw., algicolous, terrestrial, Europe, hyphomycetous, sexual morph unknown, see Seifert et al. (2011; genus accepted), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: K(M)-IMI 309488.

Ventrographium H.P. Upadhyay, Cavalc. & A.A. Silva 1986, *Ascomycota* genera *incertae sedis*, one species, type: *V. nervicola* H.P. Upadhyay, Cavalc. & A.A. Silva, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Venturia Sacc. 1882 (= *Fusicladium* Bonord. 1851; = *Pollaccia* E. Bald. & Cif. 1937; = *Pseudocladosporium* U. Braun 1998 *fide* Rossman et al. 2015b; see Species Fungorum 2017 for additional synonyms), *Venturiaceae*, *Venturiales*, *Dothideomycetes*, c. 60 species, type: *V. inaequalis* (Cooke) G. Winter, asexual morph hyphomycetous, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Zhang et al. (2011d; phylogeny), Hyde et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Rossman et al. (2015b; nomenclature), Shen et al. (2016; new species), cultures and sequences are available.

Venturiocistella Raitv. 1979, *Hyaloscyphaceae*, *Helotiales*, *Leotiomycetes*, seven species, type: *V. ventrioides* (Sacc. & Roum.) Raitv., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Hosoya et al. (2011; taxonomy, DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Venustocephala Matsush. 1995, *Ascomycota* genera *incertae sedis*, two species, type: *V. aequatorialis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Neotropics, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Venustosynnema R.F. Castañeda & W.B. Kendr. 1990, *Ascomycota* genera *incertae sedis*, three species, type: *V. ciliatum* (R.F. Castañeda, G.R.W. Arnold & A.G. Guerra) R.F. Castañeda & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Caribbean, Taiwan, see Castañeda-Ruiz et al. (2009b; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Kirschner and Liu (2014; new species, Taiwan), cultures and sequences are unavailable.

Veracruzomyces Mercado, Guarro, Heredia & J. Mena 2002, *Ascomycota* genera *incertae sedis*, one species, type: *V. obclavatus* Mercado, Guarro, Heredia & J. Mena, hyphomycetous, sexual morph unknown, fungicolous, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Veramycella G. Delgado 2009, *Ascomycota* genera *incertae sedis*, one species, type: *V. bispora* G. Delgado, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Veramyces Matsush. 1993, *Ascomycota* genera *incertae sedis*, one species, type: *V. manuensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Verdipulvinus A.W. Ramaley 1999, *Ascomycota* genera *incertae sedis*, one species, type: *V. nolinae* A.W. Ramaley, hyphomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Vermiconia Egidi & Onofri 2014, *Extremaceae*, *Capnodiales*, *Dothideomycetes*, four species, type: *V. foris* Egidi & Onofri, hyphomycetous, sexual morph unknown, rock-inhabiting fungus, Spain, Italy, see Egidi et al. (2014; taxonomy), Quaedvlieg et al. (2014; phylogeny), Isola et al. (2016; new species, phylogeny), Wijayawardene et al. (2017; outline), type: CBS 136106 = CCFFE 5459, cultures and sequences available.

Vermiculariopsiella Bender 1932 (= *Oramasia* Urries 1956 [1955]; = *Singera* Bat. & J.L. Bezerra 1960; = *Vermiculariopsis* Höhn. 1918 *fide* Species Fungorum 2017), *Vermiculariopsiellaceae*, *Vermiculariopsiellales*, *Sordariomycetes*, 19 species, type: *V. immersa* (Desm.) Bender, hyphomycetous, sexual morph unknown, saprobes, worldwide, see Dhargalkar and Bhat (2009; new species), Seifert et al. (2011; morphology), Wijayawardene et al.

(2012, 2017; outline), Hernández-Restrepo et al. (2012a; new species), Kirk et al. (2013; genus accepted), Dubey and Moonambeth (2014b; new species), Crous et al. (2016b; new species, phylogeny), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), cultures and sequence available but lack for the type, generic placement is based on non-type.

Vermispora Deighton & Piroz. 1972, *Orbiliaceae*, *Orbiliales*, *Orbiliomycetes*, seven species, type: *V. grandispora* Deighton & Piroz., hyphomycetous, sexual morph unknown, fungicolous, nematode cysts, in soil, terrestrial, Africa, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available but lacking for the type.

Veronaea Cif. & Montemart. 1957 (= *Sympodina* Subram. & Lodha 1964 *fide* Species Fungorum 2017), *Herpotrichiellaceae*, *Chaetothyriales*, *Eurotiomycetes*, 21 species, type: *V. botryosa* Cif. & Montemart., hyphomycetous, sexual morph unknown, saprobes fungicolous, on dung, on soil, worldwide, see Soares and Barreto (2008; new species), Chlebicki (2009a; new species), Pan et al. (2009; new species, China), de Hoog et al. (2011; DNA), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 254.57.

Veronaella Subram. & K.R.C. Reddy 1975, *Ascomycota* genera *incertae sedis*, one species, type: *V. indica* Subram. & K.R.C. Reddy, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Veronaeopsis Arzanlou & Crous 2007, *Sympoventuriaceae*, *Venturiales*, *Dothideomycetes*, one species, type: *V. simplex* (Papendorf) Arzanlou & Crous, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011; morphology), Zhang et al. (2011d; accepted as a genus in *Sympoventuriaceae*), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in *Sympoventuriaceae*), Samerpitak et al. (2014; phylogeny), holotype and ex-type of the type: CBS H-7810, CBS 588.66 = IMI 203547.

Veronaia Benedek 1961, *Eurotiomycetidae* genera *incertae sedis*, *Eurotiomycetes*, two species, type: *V. castellanii* Benedek, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Veronidia Negru 1964, *Ascomycota* genera *incertae sedis*, one species, type: *V. oxycocci* Negru,

coelomycetous, sexual morph unknown, saprobes, terrestrial, Rumania, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Verpa Sw. 1815 (= *Monka* Adans. 1763; = *Ptychoverpa* Boud. 1907; = *Relhanum* Gray 1821 *fide* Species Fungorum 2017), *Morchellaceae*, *Pezizales*, *Pezizomycetes*, five species, type: *V. conica* (O.F. Müll.) Sw., asexual morph unknown, saprobes, terrestrial, north temperate, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Osmundson et al. (2013; DNA), cultures available for type: AFTOL-ID 74 (unpublished).

Verrucaria Schrad. [nom. cons.] 1794 (see Species Fungorum 2017 for synonyms), *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, 300 species, type: *V. rupestris* Schrad., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Verrucariella S. Ahmad 1967, *Ascomycota* genera *incertae sedis*, one species, type: *V. capparicola* S. Ahmad, coelomycetous, sexual morph unknown, saprobes, terrestrial, Pakistan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Verrucocladosporium K. Schub., Aptroot & Crous 2007, *Cladosporiaceae*, *Capnodiales*, *Dothideomycetes*, two species, type: *V. dirinae* K. Schub., Aptroot & Crous, hyphomycetous, sexual morph unknown, lichenicolous, terrestrial, temperate, see Braun et al. (2009; new species), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2014c, 2017; outline), Zhurbenko et al. (2015c; new species) holotype and ex-type culture: CBS H-19883, CBS 112794.

Verrucoconiothyrium Crous 2015, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *V. nitidae* (Crous & Denman) Crous, coelomycetous, sexual morph unknown, saprobes, Germany, see Crous et al. (2015c; taxonomy, phylogeny), Wijayawardene et al. (2016b, 2017; outline, phylogeny), holotype of the type species PREM 55346.

Verruconis Samerp., H.J. Choi, Van den Ende, Horré & de Hoog 2013, *Sympoventuriaceae*, *Venturiales*, *Dothideomycetes*, one species, type: *V. gallopava* (W.B. Cooke) Samerpitak & de Hoog, hyphomycetous, sexual morph unknown, saprobes, human pathogen, terrestrial, South Africa, see Giraldo et al. (2014b; phylogeny), Samerpitak et al. (2014, 2016; phylogeny, barcoding), Seyedmousavi et al. (2014; antifungal susceptibility), Wijayawardene et al. (2014c, 2017; outline), Geltner et al. (2015; Necrotizing mycosis), ex-type cultures of the type: CBS 437.64= MUCL 6683=IFM 52605.

Verrucophragmia Crous, M.J. Wingf. & W.B. Kendr. 1994, *Ascomycota* genera *incertae sedis*, one species, type: *V. splendens* Crous, M.J. Wingf. & W.B. Kendr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: PREM 51441, cultures and sequences are unavailable.

Verrucoplaca S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type: *V. verruculifera* (Vain.) S.Y. Kondr., Kärnefelt, Elix, A. Thell, J. Kim, M.-H. Jeong, N.N. Yu, A.S. Kondr. & Hur, asexual morph unknown, lichenized, terrestrial, Europe, see Kondratyuk et al. (2014; taxonomy, DNA), Gaya et al. (2015; DNA, phylogeny), cultures and sequences are unavailable.

Verrucostoma Hirooka, Tak. Kobay. & P. Chaverri 2010, *Bionectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *V. freycinetiae* Hirooka, Tak. Kobay. & P. Chaverri, asexual morph acremonium-like, saprobes, terrestrial, tropical, see Hirooka et al. (2010; taxonomy), Crous et al. (2015d; new specie), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: BPI 878948, cultures and sequences are available.

Verruculina Kohlm. & Volk. Kohlm. 1990, *Testudinaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *V. enalia* (Kohlm.) Kohlm. & Volk. Kohlm., asexual morph unknown, saprobes, marine, Liberia, see Suetrong et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Didymosphaeriaceae*), Hyde et al. (2013; accepted as a genus in *Testudinaceae*), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline, marine), cultures available for the type: BCC 18402.

Verruculopsis Gueidan, Nav.-Ros. & Cl. Roux 2007, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, six species, type: *V. poeltiana* (Clauzade & Cl. Roux) Gueidan, Nav.-Ros. & Cl. Roux, asexual morph unknown, lichenized, lichenicolous, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Krzewicka (2012; new species, Poland), Kukwa and Pérez-Ortega (2010; phylogeny), Lücking et al. (2017a; classification); sequences are available, cultures available for the type: AFTOL-ID 2298.

Versicolorisporium Sat. Hatake., Kaz. Tanaka & Y. Harada 2008, *Occultibambusaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *V. triseptatum* Sat. Hatake., Kaz. Tanaka & Y. Harada, coelomycetous, saprobes, terrestrial, Asia, see Wijayawardene et al. (2012, 2016b, 2017; outline, morphology), Dai et al. (2017; phylogeny, morphology, sexual and asexual morphs), ex-type culture of type: MFLUCC 11-0179.

Verticicola K.D. Hyde, Ranghoo & S.W. Wong 2000, *Annulatasaceae*, *Annulatascales*, *Sordariomycetidae*, two species, type: *V. caudatus* K.D. Hyde, S.W. Wong & Ranghoo, asexual morph unknown, saprobes, fresh water, terrestrial, Philippines, Hong Kong, Costa Rica, see Lumbsch and Huhndorf (2010; outline), Barbosa et al. (2013a; new species, Costa Rica), Maharachchikumbura et al. (2016b; outline), holotype of the type: HKU (M) 3108, cultures and sequences are available but lacking for the type.

Verticicladus Matsush. 1993, *Ascomycota* genera *incertae sedis*, three species, type: *V. amazonensis* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, aquatic, Africa, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Guo et al. (2015; new species), cultures and sequences are unavailable.

Verticillium Nees 1816 (= *Verticillium* sect. *Verticillium* Nees 1816 [1816-17] *fide* Species Fungorum 2017), *Plectosphaerellaceae*, *Glomerellales*, *Sordariomycetes*, c. 20 species (in strict sense), type: *V. dahliae* Kleb., hyphomycetous, Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for type species ATCC 16535.

Verticimonosporium Matsush. 1971, *Hypocreaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *V. diffractum* Matsush., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, North and South America, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures available for the type: NRRL 26545.

Vesicidiella Crous & M.J. Wingf. 1994, *Ascomycota* genera *incertae sedis*, two species, type: *V. capitata* (B. Sutton) Crous & M.J. Wingf., hyphomycetous, sexual morph unknown, saprobes, Australia, see Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), holotype of type: IMI163277a.

Vesiculohyphomyces Armando, Pereira-Carv. & Dianese 2009, *Ascomycota* genera *incertae sedis*, one species, type: *V. cerradensis* Armando, Pereira-Carv. & Dianese, hyphomycetous, sexual morph unknown, saprobes, terrestrial, South America, see Pereira-Carvalho et al. (2009b; taxonomy), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), holotype of the type: UB (Mycol. Col.) 14595, cultures and sequences are unavailable.

Vestergrenia Rehm 1901 (= *Catacaumella* Theiss. & Syd. 1915; = *Coscinopeltella* Chardón 1930; = *Guignardiella* Sacc. & P. Syd. 1902; = *Haplodothella*

Werderm. 1923; = *Pediascus* Chardón & Toro 1934; = *Whetzelia* Chardón & Toro 1934 *fide* Species Fungorum 2017), *Botryosphaeriaceae*, *Botryosphaeriales*, *Dothideomycetes*, 13 species, type: *V. nervisequia* Rehm, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Dothideaceae*), Kirk et al. (2013; genus accepted), Hyde et al. (2013; accepted as a genus in *Botryosphaeriaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Vestergrenopsis Gyeln. 1940, *Koerberiaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *V. elaeina* (Wahlenb. ex Ach.) Gyeln., asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Spribille and Muggia (2013; new species, DNA, phylogeny), sequences are available for voucher Spribille 24525 (KLGO) of the type, Lücking et al. (2017a; classification), sequences are available.

Vestigium Piroz. & Shoemaker 1972, *Ascomycota* genera *incertae sedis*, two species, type: *V. felicis* Piroz. & Shoemaker, coelomycetous, sexual morph unknown, saprobes, aquatic, cosmopolitan, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Shoemaker et al. (2013; new species), cultures and sequences are available.

Vezdaea Tscherm.-Woess & Poelt 1976 (= *Vezdaea* Poelt & Döbbeler 1975 *fide* Species Fungorum 2017), *Vezdaeaceae*, *Vezdaeales*, *Ascomycota* orders *incertae sedis*, twelve species, type: *V. aestivalis* (Ohlert) Tscherm.-Woess & Poelt, asexual morph unknown, lichenized, saprobes, Europe, see Czarnota and Kukwa (2009; Poland), Lumbsch et al. (2009a; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification) cultures and sequences are available.

Vialaea Sacc. 1896 (= *Boydia* A.L. Sm. 1919 [1918] *fide* Species Fungorum 2017), *Vialaeaceae*, *Amphisphaeriales*, *Sordariomycetes*, three species, type: *V. insculpta* (Fr.) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Senanayake et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures available for the type: AFTOL-ID:1302.

Vibrissea Fr. 1822 (= *Anavirga* B. Sutton 1975; = *Apostemidium* (P. Karst.) P. Karst. 1871; = *Apostemium* P. Karst. 1870 [1871]; = *Leptosporium* Bonord. 1857; = *Ophiogloea* Clem. 1903; = *Peziza* sect. *Apostemidium* P. Karst. 1869 *fide* Johnston et al. 2014b; Species Fungorum 2017), *Vibrisseaceae*, *Helotiales*, *Leotiomycetes*, 34 species, type: *V. truncorum* (Alb. & Schwein.) Fr., asexual morph formerly known in *Anavirga*, saprobes,

aquatic or semiaquatic, worldwide, see Grünig et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Hustad and Miller (2011; DNA, phylogeny), Raja et al. (2011a; DNA, phylogeny), Kirk et al. (2013; genus accepted), Johnston et al. (2014b; nomenclature), cultures available for the type: CBS 143.92.

Viegasella Inácio & P.F. Cannon 2003, *Parmulariaceae*, *Dothideomycetes* genera *incertae sedis*, one species, type: *V. pulchella* (Speg.) Inácio & P.F. Cannon, asexual morph unknown, saprobes, terrestrial, Brazil, see Lumbsch and Huhndorf (2010; outline, listed as *Viegasiella*), Hyde et al. (2013; accepted as a genus in *Parmulariaceae*), Wijayawardene et al. (2014c; outline), Tian et al. (2016a; taxonomy), holotype of type: LPS, cultures and sequences are unavailable.

Vigneronia Ertz 2015, *Roccellaceae*, *Arthoniales*, *Arthoniomycetes*, three species, type: *V. spieri* (Aptroot & Sparrius) Ertz & Bungartz, sexual morph with lirelliform ascocarps, lichenized, terrestrial, neotropical, see Ertz et al. (2015b, c; morphology, phylogeny), Lücking et al. (2017a; classification), sequences are available, holotype and ex-type of the type: Aptroot 65014 (CDS31595).

Vikalpa D'souza, Boonmee, Bhat & K.D. Hyde 2016, *Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*, three species, type: *V. australiense* (B. Sutton) D'souza, Boonmee & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, Thailand, China, see Boonmee et al. (2016a; phylogeny), Wijayawardene et al. (2017; outline), holotype: HKAS 83979, cultures and sequences are available.

Villophora Søchting, Arup & Frödén 2013, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, two species, type: *V. isidioclada* (Zahlbr.) Søchting, Frödén & Arup, asexual morph unknown, lichenized, saxicolous or epiphytic, terrestrial, southern hemisphere, see Arup et al. (2013; taxonomy, phylogeny), Kondratyuk et al. (2015c; new species, DNA, phylogeny), Lücking et al. (2017a; classification); sequence available for the type: KC179325 (ITS) (*fide* Arup et al. 2013).

Violella T. Sprib. 2011, *Tephromelataceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *V. fucata* (Stirt.) T. Sprib., lichenized, see Spribille et al. (2011; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Virgaria Nees 1816 (= *Ascovirgaria* J.D. Rogers & Y.M. Ju 2002 *fide* Réblová et al. 2016c), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, 16 species, type: *V. nigra* (Link) Nees, sexual morph formerly known in *Ascovirgaria*, saprobes, terrestrial, Africa, Australia, Europe, Japan, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Nonaka et al. (2013a; new species, Japan, DNA), Maharachchikumbura et al.

(2015b, 2016b; outline), Réblová et al. (2016c; nomenclature), cultures available for the type: NBRC 9453.

Virgariella S. Hughes 1953, *Ascomycota* genera *incertae sedis*, eleven species, type: *V. globigera* (Sacc. & Ellis) S. Hughes, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, Caribbean, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Virgatospora Finley 1967, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *V. echinofibrosa* Finley, hyphomycetous, sexual morph unknown, saprobes, terrestrial, pantropical, see Seifert et al. (2011; morphology), Lombard et al. (2016; phylogeny), Wijayawardene et al. (2017; outline), culture available for the type: CBS.

Virgella Darker 1967, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, one species, type: *V. robusta* (Tubeuf) Darker, asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Viridiannula Etayo 2002, *Ascomycota* genera *incertae sedis*, one species, type: *V. pertusariae* Etayo, coelomycetous, sexual morph unknown, parasymbiont, terrestrial, Spain, see Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable, needs revision, holotype of the type: hb. Etayo, Etayo 14496.

Viridothelium Lücking, M.P. Nelsen & Aptroot 2016, *Trypetheliaceae*, *Trypetheliales*, *Dothideomycetes*, ten species, type: *V. virens* (Tuck. ex Michener) Lücking, M.P. Nelsen & Aptroot, asexual morph unknown, lichenized fungi, terrestrial, tropical, see Aptroot and Lücking (2016; taxonomy), Lücking et al. (2017a; c; classification), sequences are available.

Vishnumyces Hosag. 2010, *Dothideomycetes* genera *incertae sedis*, one species, type: *V. otonephelii* Hosag. & Harish, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Asterinaceae*), Hosagoudar and Harish (2010; taxonomy, accepted as a genus in *Asterinaceae*), Hyde et al. (2013) and Hongsanan et al. (2014c) did not treat as a genus in *Asterinaceae*, Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Vismaya V.V. Sarma & K.D. Hyde 2001, *Diaporthales* genera *incertae sedis*, one species, type: *V. chaturbeeja* V.V. Sarma & K.D. Hyde, asexual morph unknown, saprobes, terrestrial, Hong Kong, see Lumbsch and Huhndorf (2010; outline, listed as *Vismya*), Maharachchikumbura et al. (2015b, 2016b; outline),

lectotype of the type: HKU (M) 12457, cultures and sequences are unavailable.

Vitreostroma P.F. Cannon 1991, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *V. desmodii* (Henn.) P.F. Cannon, in living leaves, montane paleotropics, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Vittalia Gawas & Bhat 2007, *Ascomycota* genera *incertae sedis*, one species, type: *V. indica* Gawas & Bhat, hyphomycetous, sexual morph unknown, saprobes, terrestrial, India, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Vittatispora P. Chaudhary, J. Campb., D. Hawksw. & K.N. Sastry 2006, *Ceratostomataceae*, *Coronophorales*, *Sordariomycetes*, one species, type: *V. coorgii* P. Chaudhary, J. Campb., D. Hawksw. & K.N. Sastry, asexual morph unknown, saprobes, terrestrial, India, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures available for the type: BICC 7817.

Vivantia J.D. Rogers, Y.M. Ju & Cand. 1996, *Graphostromataceae*, *Xylariales*, *Sordariomycetes*, one species, type: *V. guadalupensis* J.D. Rogers, Y.M. Ju & Cand., asexual morph hyphomycetous, terrestrial, saprobes, Guadeloupe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; transferred to *Graphostromataceae*), cultures and sequences are unavailable.

Vizella Sacc. 1883 (= *Entopeltis* Höhn. 1910; = *Haplopyrenula* Müll. Arg. 1883; = *Haplopyrenulomyces* Cif. & Tomas. 1953; = *Haplospora* Räsänen 1943; = *Hypocelis* Petr. 1929; = *Mycerema* Bat., J.L. Bezerra & Cavalc. 1963; = *Phaeaspis* Clem. & Shear 1931; = *Phaeopeltis* Petch 1919; = *Stigmatopeltis* Dodge 1927 fide Species Fungorum 2017), *Vizzellaceae*, *Dothideomycetes* genera *incertae sedis*, eleven species, type: *V. conferta* (Cooke) Sacc., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Vizzellaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Vizelopsis Bat., J.L. Bezerra & T.T. Barros 1969, *Asterinaceae*, *Asterinales*, *Dothideomycetes*, one species, type: *V. grevilleae* Bat., J.L. Bezerra & T.T. Barros, asexual morph unknown, foliar epiphytes, terrestrial, New Caledonia, see Lumbsch and Huhndorf (2010; outline, accepted as *Dothideomycetes* genera *incertae sedis*), Kirk et al. (2013; genus accepted), Dai et al. (2014a; accepted as

a genus in *Asterinaceae* based on morphology), Hongsanan et al. (2014c; taxonomy, accepted as a genus in *Asterinaceae*), Wijayawardene et al. (2014c; 2017; outline), cultures and sequences are unavailable.

Vladracula P.F. Cannon, Minter & Kamal 1986, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, two species, type: *V. annuliformis* (Syd., P. Syd. & E.J. Butler) P.F. Cannon, Minter & Kamal, asexual morph unknown, saprobes, terrestrial, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Vleugelia J. Reid & C. Booth 1969, *Sordariomycetes* genera *incertae sedis*, one species, type: *V. betulina* (Bubák & Vleugel) J. Reid & C. Booth, asexual morph unknown, saprobes, terrestrial, Sweden, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Volkartia Maire 1907, *Protomycetaceae*, *Taphrinales*, *Taphrinomycetes*, one species, type: *V. rhaetica* (Volkart) Maire, asexual reproduction is unknown, plant parasite, parasitic on species of *Asteraceae*, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Volutella Fr. 1832 (= *Chaetodochium* Höhn. 1932; = *Psilonia* Fr. 1825; = *Thysanopyxis* Rabenh. ex Bonord. 1864; = *Volutellonectria* J. Luo & W.Y. Zhuang *fide* Rossman et al. 2016a; Species Fungorum 2017), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, c. 25 species, type: *V. ciliata* (Alb. & Schwein.) Fr., hyphomycetous, sexual morph formerly known in *Volutellonectria*, saprobes, terrestrial, widespread, see Seifert et al. (2011; morphology), Gräfenhan et al. (2011, morphology, phylogeny), Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2016b; outline), Rossman et al. (2016a; nomenclature), cultures available for type: WBS025, YIMPH30011, GJS 85.205, DAOM 226718, CBS 483.61.

Vonarxella Bat., J.L. Bezerra & Peres 1965, *Saccardiaceae*, *Ascomycota* families *incertae sedis*, one species, type: *V. dipterygis* Bat., J.L. Bezerra & Peres, asexual morph unknown, saprobes, Brazil, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Vonarxia Bat. 1960, *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, one species, type: *V. anacardii* Bat. & J.L. Bezerra, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Crous et al. (2009a; DNA, phylogeny, treated as hyphomycetous-i.e. with sporodochial conidioma), Seifert et al. (2011; treated as coelomycetous), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Vouauxiella Petr. & Syd. 1927 (= *Alysia* Cavalc. & A.A. Silva 1972 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, five species, type: *V. verrucosa* (Vouaux) Petr. & Syd., coelomycetous, sexual morph unknown, lichenicolous, terrestrial, Europe, South America, see Wijayawardene et al. (2012, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Vrystaatia Quaedv., W.J. Swart, Verkley & Crous 2013, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *V. aloicola* Quaedv., Verkley, W.J. Swart & Crous coelomycetous, sexual morph unknown, pathogenic, terrestrial, South Africa, see Quaedvlieg et al. (2013; morphology, phylogeny), holotype and ex-type strains of the type: CBS H-21313; CBS 135107.

Wadeana Coppins & P. James 1978, *Ascomycota* genera *incertae sedis*, two species, type: *W. dendrographa* (Nyl.) Coppins & P. James, asexual morph unknown, lichenized, terrestrial, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); sequences unavailable.

Wahlenbergiella Gueidan & Thüs 2009, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, three species, type: *W. mucosa* (Wahlenb.) Gueidan & Thüs, asexual morph unknown, lichenized, terrestrial, Europe, see Gueidan et al. (2009, 2011; taxonomy), Lumbsch and Huhndorf (2010; outline), Shivarov and Stoykov (2013; new records, Bulgaria), Higgins et al. (2015; factors in distributing), Lücking et al. (2017a; classification), sequences are available, cultures available for the type: AFTOL-ID 2264.

Waihonghopes Yanna & K.D. Hyde 2002, *Ascomycota* genera *incertae sedis*, one species, type: *W. australiensis* Yanna & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable.

Wallrothiella Sacc. 1882 (= *Zignoina* Cooke 1885 *fide* Species Fungorum 2017), *Amplistromataceae*, *Amplistromatales*, *Sordariomycetes*, ten species, type: *W. congregata* (Wallr.) Sacc., asexual morph unknown, saprobes, terrestrial, worldwide, see Huhndorf et al. (2009; DNA, phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures available for the type: ANM81.

Waltonia Saho 1970, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *W. pinicola* Saho, asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable.

Warcupia Paden & J.V. Cameron 1972, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, three species, type: *W. terrestris* Paden & J.V. Cameron, asexual morph unknown, hypogeous, saprobes, terrestrial, Canada, see Lumbsch and Huhndorf (2010; outline), Barua et al. (2012; environmental adaptations), Hansen (2013; DNA, phylogeny), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 891.69.

Warcupiella Subram. 1972 (= *Sporophormis* Malloch & Cain 1973 [1972] *fide* Species Fungorum 2017), *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, three species, type: *W. spinulosa* (Warcup) Subram., asexual morph unknown, saprobes, terrestrial, worldwide, see Houbraken and Samson (2011; taxonomy, phylogeny), Kirk et al. (2013; genus accepted), culture available for type: CBS 430.64.

Wardinella Bat. & Peres 1960, *Ascomycota* genera *incertae sedis*, one species, type: *W. jaboatonensis* Bat. & Peres, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Wardomyces F.T. Brooks & Hansf. 1923, *Microascaceae*, *Microascales*, *Sordariomycetes*, ten species, type: *W. anomalus* F.T. Brooks & Hansf., hyphomycetous, saprobes, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Sandoval-Denis et al. (2016b; new species, phylogeny), cultures and sequences are available.

Wardomycopsis Udagawa & Furuya 1978, *Microascaceae*, *Microascales*, *Sordariomycetes*, four species, type: *W. inopinata* Udagawa & Furuya, hyphomycetous, saprobes, Asia, Europe, North America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline), Sandoval-Denis et al. (2016b; new species, phylogeny), cultures and sequences are available.

Wawea Henssen & Kantvilas 1985, *Arctomiaceae*, *Arctomiales*, *Lecanoromycetes*, one species, type: *W. fruticulosa* Henssen & Kantvilas, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Wawelia Namysł. 1908, *Xylariaceae*, *Xylariales*, *Sordariomycetes*, three species, type: *W. regia* Namysł., asexual morph unknown, terrestrial, coprophilous, Europe, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al.

(2015b, 2016b; outline), cultures and sequences are unavailable.

Waydora B. Sutton 1976 (= *Pulvinaria* Rodway 1918 [1917] *fide* Species Fungorum 2017), *Valsaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *W. typica* (Rodway) B. Sutton, coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures available for voucher PDD 103894 of the type: (unpublished).

Waynea Moberg 1990, *Ramalinaceae*, *Lecanorales*, *Lecanoromycetes*, seven species, type: *W. californica* Moberg, lichenized, see Lücking et al. (2017a; classification), sequences are available.

Websteromyces W.A. Baker & Partr. 2000, *Ascomycota* genera *incertae sedis*, one species, type: *W. verruculosus* W.A. Baker & Partr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), holotype of the type: AUA, Webster, 11 Jul. 1991, cultures and sequences are unavailable.

Weddellomyces D. Hawksw. 1986, *Dacampiaceae*, *Dothideomycetes* families *incertae sedis* 14 species, type: *W. epicallopisma* (Weddell) D. Hawksw., see Lumbsch & Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), sequences are unavailable, isotype of type: H-Nyl 391.

Wegea Aptroot & Tibell 1997, *Arthoniales* genera *incertae sedis*, *Arthoniomycetes*, one species, type: *W. tylophorelloides* Aptroot & Tibell, lichenized, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequences are unavailable.

Wehmeyera J. Reid & C. Booth 1989, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *W. acerina* (Wehm.) J. Reid & C. Booth, asexual morph unknown, saprobes, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongsanan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are unavailable.

Wentiomycetes Koord. 1907 (= *Acarothallium* Syd. 1937; = *Calochaetis* Syd. 1935; = *Capnodinula* Speg. 1918; = *Chaetostigme* Syd. & P. Syd. 1917; = *Dichaetis* Clem. 1931; = *Neocoleroa* Petr. 1934 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, c. 50 species, type: *W. javanicus* Koord., Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Wenyingia Zheng Wang & Pfister 2001, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, one species, type: *W. sichuanensis* Zheng Wang & Pfister, asexual

morph unknown, saprobes, terrestrial, China, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are unavailable.

Westea H.J. Swart 1988, *Dothideomycetes* genera *incertae sedis*, one species, type: *W. banksiae* H.J. Swart, saprobes, terrestrial, Australia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Westerdykella Stolk 1955, *Sporormiaceae*, *Pleosporales*, *Dothideomycetes*, 49 species, type: *W. ornata* Stolk, asexual morph unknown, saprobes, terrestrial, worldwide, see Kruys and Wedin (2009; new species), Schoch et al. (2009; phylogeny), Asgari and Zare (2010; new species), Lumbsch and Huhndorf (2010; outline), Ebead et al. (2012; new species), Armentano et al. (2013; pathogens of dogs), de Gruyter et al. (2012; new species, phylogeny), Hyde et al. (2013; accepted as a genus in *Sporormiaceae*, phylogeny), Wijayawardene et al. (2014c; outline), Kirk et al. (2013; genus accepted), cultures available for the type: CBS 379.55.

Wetmoreana Arup, Söchting & Frödén 2013 (= *Fulgogasparrea* S.Y.Kondr., N.-H.Jeong, Kärnefelt, Elix, A.Thell & Hur 2013), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *W. texana* (Wetmore & Kärnefelt) Arup, Frödén & Söchting, lichenized, terrestrial, Northern Hemisphere, see Arup et al. (2013; description), Kondratyuk et al. (2013; synonym), Jaklitsch et al. (2016a; classification), sequences are available; this name is deemed invalid in *Index Fungorum* because "... *The identifier cited in the protologue, 'MB 02011', was not issued by a recognized repository ...*"; however, the erroneous number is a typographic error for 'MB 202011' as apparent from the context and it is unclear whether this renders the name invalid (e.g. ICN Art. 33.2).

Wettsteinina Höhn. 1907 (= *Pseudosphaeria* Höhn. 1907 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, 29 species, type: *W. gigantospora* (Rehm) Höhn., saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type.

Weufia Bhat & B. Sutton 1985, *Ascomycota* genera *incertae sedis*, one species, type: *W. tewoldei* Bhat & B. Sutton, hyphomycetous, sexual morph unknown, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Whalleya J.D. Rogers, Y.M. Ju & F. San Martín 1997, *Lopadostomataceae*, *Xylariales*, *Sordariomycetes*, two species, type: *W. microplaca* (Berk. & M.A. Curtis) J.D.

Rogers, Y.M. Ju & F. San Martín, asexual morph unknown, terrestrial, saprobes, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; transferred to *Lopadostomataceae*), cultures and sequences are available.

Wickerhamia Soneda 1960, *Metschnikowiaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *W. fluorescens* (Soneda) Soneda, saprobes, Japan, France, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Wickerhamiella van der Walt 1973, *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, 25 species, type: *W. domercqiae* van der Walt & Liebenberg, asexual reproduction is by multilateral budding on either a narrow or a broad base, saprophytic, on nitidulid beetles, flowers of morning glory (*Hibiscus* spp.) and cacti, wine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Wickerhamomyces Kurtzman et al. 2008, *Phaffomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, 20 species, type: *W. canadensis* (Wickerham) Kurtzman, Robnett & Basehoar-Powers, asexual reproduction is by multilateral budding, saprophytic, but several species are infrequent clinical isolates, in tree sap fluxes, insect tunnels in trees, leaf litter, berries, soil, tanned leather, soy sauce, blood, lung, eye, worldwide, see de García et al. (2010; new species, DNA), Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Wicklowia Raja, A. Ferrer & Shearer 2010, *Wicklowiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *W. aquatica* Raja, A. Ferrer & Shearer, asexual morph unknown, saprobes, aquatic, USA, see Raja et al. (2010b; taxonomy, phylogeny), holotype of the type: ILL 40790.

Wiesneriomycetes Koord. 1907 (= *Aschizotrichum* Rieff 1962; = *Chaetopeltis* Sacc. 1898; = *Chaetosira* Clem. 1931; = *Gamonaemella* Fairm. 1922; = *Gamospora* Sacc. 1885; = *Setodochium* Bat. & Cif. 1957; = *Tassia* Syd. & P. Syd. 1919 *fide* Species Fungorum 2017), *Wiesneriomycetaceae*, *Tubeufiales*, *Dothideomycetes*, two species, type: *W. javanicus* Koord., hyphomycetous, saprobe, terrestrial, widespread, see Hyde et al. (2011; outline), Kirk et al. (2013; genus accepted), Suetrong et al. (2014; morphology, phylogeny), Bezerra et al. (2017a; phylogeny), Liu et al. (2017, placement), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Wilcoxina Chin S. Yang & Korf 1985, *Pyronemataceae*, *Pezizales*, *Pezizomycetes*, three species, type: *W. mikolae* (Chin S. Yang & H.E. Wilcox) Chin S. Yang & Korf,

asexual morph unknown, hypogeous, saprobes, terrestrial, Europe, North America, see Siemens and Zwiazek (2008; mycorrhizal with *Hebeloma crustuliniforme*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures are available for the type but unpublished, sequence available for the type: DQ220468 (LSU).

Willeya Müll. Arg. 1883, *Verrucariaceae*, *Verrucariales*, *Eurotiomycetes*, eleven species, type: *W. diffractella* (Nyl.) Müll. Arg., lichenized, see Gueidan et al. (2014a; taxonomy), Gueidan and Lendemer (2015; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Wirthiotrema Rivas Plata, Kalb, Frisch & Lumbsch 2010, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, seven species, type: *W. glaucopallens* (Nyl.) Rivas Plata & Kalb, asexual morph unknown, lichenized, terrestrial, tropical, see Rivas Plata et al. (2010a; classification), Sipman et al. (2012; key), Lücking et al. (2017a; classification), sequences are available.

Wojnowiciella Crous, Hern.-Restr. & M.J. Wingf. 2015, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, seven species, type: *W. eucalypti* Crous, Hern.-Restr. & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Crous et al. (2015a, d, 2016a, b; new species, phylogeny), Hernández-Restrepo et al. (2016a; new species, new species, phylogeny), Wijayawardene et al. (2016b, 2017; taxonomy, outline, phylogeny), cultures are available.

Wolfina Seaver ex Eckblad 1968 (= *Wolfina* Seaver 1937 *fide* Species Fungorum 2017), *Chorioactidaceae*, *Pezizales*, *Pezizomycetes*, three species, type: *W. aurantiopsis* (Ellis) Seaver, asexual morph unknown, saprobes, terrestrial, USA, China, see Lumbsch and Huhndorf (2010; outline), Hustad and Miller (2011; DNA, phylogeny), Kirk et al. (2013; genus accepted), sequences are available.

Wolkia Ramsb. 1915 (= *Protascus* Wolk 1913 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *W. decolorans* (Wolk) Ramsb., asexual morph unknown, saprobes, terrestrial, Java, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Wongia Khemmuk, Geering & R.G. Shivas 2016, *Papulosaceae*, *Diaportheomycetidae* families *incertae sedis*, *Sordariomycetes*, two species, type: *W. garrettii* (P. Wong & M.L. Dickinson) Khemmuk, Geering & R.G. Shivas, asexual morph unknown, pathogen, Australia, see Khemmuk et al. (2016; taxonomy, phylogeny), holotype: DAR 76937, cultures and sequences are available.

Woodiella Sacc. & P. Syd. 1899, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type:

W. natalensis Sacc. & P. Syd., asexual morph unknown, saprobes, terrestrial, South Africa, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Woswasia Jaklitsch, Réblová & Voglmayr 2012, *Sordariomycetidae* genera *incertae sedis*, *Sordariomycetes*, one species, type: *W. atropurpurea* Jaklitsch, Réblová & Voglmayr, asexual morph formed in culture only (penicilliate heads with terminally arranged, sympodially proliferating conidiogenous cells producing simple ellipsoid to ovoid, hyaline, aseptate conidia holoblastically), myotrophic saprobes, terrestrial, Italy, see Jaklitsch et al. (2013; original description, phylogeny), Réblová et al. (2014; close relationship with *Xylochrysis*), Maharachchikumbura et al. (2015b, 2016b; outline), sequence are available, holotype and ex-type of type: WU 32007, CBS 133167.

Wuestneia Auersw. ex Fuckel 1864, *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, 20 species, type: *W. aurea* (Fuckel) Auersw., forms pycnidial asexual morphs harknessia-like, saprobes, terrestrial, worldwide, see Crous et al. (2012f; placement in new family *Harknessiaceae*), Wijayawardene et al. (2012, 2017; outline), Maharachchikumbura et al. (2015b, 2016b; outline, as *Melanconidaceae*), Hongisan et al. (2017a; evolution, treated *Melanconidaceae* as a synonym of *Gnomoniaceae*), cultures and sequences are available.

Wynnea Berk. & M.A. Curtis 1867, *Sarcoscyphaceae*, *Pezizales*, *Pezizomycetes*, seven species and two variants, type: *W. gigantea* Berk. & M.A. Curtis, saprobes, terrestrial, Asia, North and Central America, see Lumbsch and Huhndorf (2010; outline), Romero et al. (2012; phylogeny), Kirk et al. (2013; genus accepted), sequences are available

Wynnella Boud. 1885 (= *Midotis* Fr. 1828 *fide* Species Fungorum 2017), *Helvellaceae*, *Pezizales*, *Pezizomycetes*, three species, type: *W. auricula* (Schaeff.) Boud., saprobes, terrestrial, Europe, Asia, North America, see Kirk et al. (2013; genus accepted), Landeros et al. (2015; phylogeny), Zhao et al. (2015, 2016a; new species, phylogeny), sequences are available.

Xalcooa Kraichak, Lücking & Lumbsch 2013, *Graphidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *X. ocellata* (Fr.) Kraichak, Lücking & Lumbsch, asexual morph unknown, lichenized fungi, terrestrial, temperate, see Kraichak et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Xanthocarpia A. Massal. & De Not. 1853, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 13 species, type: *X. ochracea* (Schaer.) A. Massal. & De Not., lichenized, see Arup et al. (2013; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Xanthodactylon P.A. Duvign 1941, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, seven species, type: *X. perforatum* P.A. Duvign, asexual morph unknown, lichenized, terrestrial, see Fedorenko et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Nyati et al. (2013; phylogeny), sequences are unavailable for type species.

Xanthomendoza S.Y. Kondr. & Kärnefelt 1997, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, 20 species, type: *X. mendozae* (Räsänen) S.Y. Kondr. & Kärnefelt, asexual morph unknown, lichenized, terrestrial, see Fedorenko et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Lamit et al. (2011; ecology), Messuti et al. (2011; new species), Moniri et al. (2011; new record), Riddell et al. (2011; air pollution), Bomble (2012; taxonomy), Arup et al. (2013; taxonomy), Hauck et al. (2013a; new records), Kirk et al. (2013; genus accepted), Leavitt et al. (2013; taxonomy), Lindblom and Søchting (2013; genetic diversity of photobiont), Nyati et al. (2013; phylogeny), Diederich et al. (2014; new combination), Kondratyuk et al. (2014; phylogeny), Lindblom (2014; taxonomy), Valadbeigi (2014; floristic), Crittenden et al. (2015; ecology), Gaya et al. (2012, 2015; molecular), Holt et al. (2015; ecology), McMullin et al. (2015; new records), Miralles et al. (2015; biostrategy), Suija et al. (2015c; new record), Tripp (2015; inventory), Jaklitsch et al. (2016a; classification), Lücking et al. (2017a; classification), sequences are available.

Xanthoparmelia (Vain.) Hale 1974 (= *Almbornia* Essl. 1981; = *Namakwa* Hale 1988; = *Parmelia* sect. *Xanthoparmelia* Vain. 1890; = *Xanthomaculina* Hale 1985 *fide* Species Fungorum 2017), *Parmeliaceae*, *Lecanorales*, *Lecanoromycetes*, 820 species, type: *X. conspersa* (Ehrh. ex Ach.) Hale, asexual morph unknown, lichenized, terrestrial, worldwide, see Elix and Kantvilas (2009; new species), Tuzen et al. (2009; biosorptive study), Dailey et al. (2010; chemical study), Lumbsch and Huhndorf (2010; outline), Pérez-Vargas et al. (2010b; new species), Yazici et al. (2010; new records), Hawksworth et al. (2011a; checklist), Hodkinson and Lendemer (2011a; molecular), Leavitt and Clair (2011; ecology), Leavitt et al. (2011b, c; taxonomy and molecular), Biazrov (2012; physiology), de Paz et al. (2012; molecular taxonomy), Favero-Longo and Piervittori (2012; transplant expt.), Molnár et al. (2012; molecular), Kirk et al. (2013; genus accepted), Rizzi and Giordani (2013; ecology), Goni et al. (2014; checklist), Jayalal et al. (2014; new species, taxonomy), Cao et al. (2015; molecular), Lücking et al. (2017a; classification), Kanigowski et al. (2016; taxonomy), sequences are available.

Xanthopeltis R. Sant. 1949, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, one species, type:

X. rupicola R. Sant., lichenized, see Lumbsch and Huhndorf (2010; outline), Arup et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are available.

Xanthopsorella Kalb & Hafellner 1984 (= *Xanthopsora* Gotth. Schneid. & W.A. Weber 1980 [1979] *fide* Species Fungorum 2017), *Catillariaceae*, *Lecanorales*, *Lecanoromycetes*, one species, type: *X. texana* (W.A. Weber) Kalb & Hafellner, asexual morph unknown, lichenized, Mexico, USA, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification); COLO 448, COLO (holotype), cultures and sequences are unavailable, need generic revision.

Xanthopsoroma Elvebakk & S.G. Hong 2010, *Pannariaceae*, *Peltigerales*, *Lecanoromycetes*, two, species, type: *X. contextum* Elvebakk & S.G. Hong, asexual morph unknown, lichenized, terrestrial, see Elvebakk et al. (2010; molecular), sequences are available for type.

Xanthoria (Fr.) Th. Fr. 1861 (see Species Fungorum 2017 for synonyms), *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, ten species, type: *X. parietina* (L.) Beltr., asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Sari and Tuzen (2010; biosorption), Karagöz et al. (2011; new species), Pisani et al. (2011; physiology), Beck and Mayr (2012; physiology), Demiray et al. (2012; biomonitoring), Arup et al. (2013; taxonomy), Kirk et al. (2013; genus accepted), Kondratyuk et al. (2013c; new species), Nyati et al. (2013; phylogeny), Motiejunaite and Grochowski (2014; new records), Joshi et al. (2015a; checklist), Kaya et al. (2015; natural product), Anar et al. (2016; antigenotoxic and Antioxidant study), Lücking et al. (2017a; classification), sequences are available.

Xanthoriicola D. Hawksw. 1973, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *X. physciae* (Kalchbr.) D. Hawksw., sexual morph unknown, lichenicolous, mainly in Europe, Seifert et al. (2011; morphology), Ruibal et al. (2011; sequences), Ertz et al. (2013; phylogeny), Kirk et al. (2013; genus accepted), Pérez-Ortega et al. (2015; position), sequences are available, isotype of type: S.

Xanthothecium Arx & Samson 1973, *Onygenaceae*, *Onygenales*, *Eurotiomycetes*, one species, type: *X. peruvianum* (Cain) Arx & Samson, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), sequence available for the type: AB053453.

Xenidiocercus Nag Raj 1993, *Ascomycota* genera *incertae sedis*, one species, type: *X. macarangae* (T.S. Ramakr.) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, West Africa, see

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xenoacremonium L. Lombard & Crous 2015, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *X. recifei* (Leão & Lôbo) L. Lombard & Crous, hyphomycetous, sexual morph unknown, saprobes, Brazil, New Zealand, see Lombard et al. (2015b; morphology, phylogeny), Wijayawardene et al. (2017; outline), ex-type of type: CBS 137.3.

Xenocamarosporium Crous & M.J. Wingf. 2015, *Didymosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *X. acaciae* Crous & M.J. Wingf., coelomycetous, sexual morph unknown, saprobes, Malaysia, see Crous et al. (2015d; taxonomy, phylogeny), Wijayawardene et al. (2016b, 2017; outline, phylogeny), holotype and ex-type of the type: CBS H-22224, CPC 24755.

Xenochalara M.J. Wingf. & Crous 2000, *Dermateaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *X. juniperi* M.J. Wingf. & Crous, hyphomycetous, sexual morph unknown, saprobes, The Netherlands, see Seifert et al. (2011; morphology), Wijayawardene et al. (2017; outline), holotype and ex-type of type: PREM 26210, CBS 670.75.

Xenochora Petr. 1948, *Ascomycota* genera *incertae sedis*, one species, type: *X. sydowii* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Ecuador, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable, need generic revision.

Xenoconiothyrium Crous & Marinc. 2011, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *X. catenatum* Crous & Marinc., coelomycetous, saprobes, South Africa, see Crous et al. (2011c; morphology, phylogeny), Wijayawardene et al. (2016b, 2017; morphology, phylogeny, outline), holotype and ex-type: PREM 59486, CMW 22113.

Xenocylindrocladium Decock, Hennebert & Crous 1997 (= *Xenocalonectria* Crous & C.L. Schoch 2000 *fide* Rossman et al. 2016a), *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, three species, type: *X. serpens* Decock, Hennebert & Crous, hyphomycetous, sexual morph formerly known in *Xenocalonectria*, saprobes, terrestrial, South America, see Seifert et al. (2011; morphology), Lombard et al. (2015b; phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Rossman et al. (2016a; nomenclature), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Xenocylindrosporium Crous & Verkley 2009, *Phaeomoniellaceae*, *Phaeomoniellales*, *Eurotiomycetes*, one species, type: *X. kirstenboschense* Crous & Verkley, hyphomycetous, pathogens, South Africa, see Crous and

Verkley (2009; morphology, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type culture of type: CBS H-20346, CPC 16311.

Xenodidymella Q. Chen & L. Cai 2015, *Didymellaceae*, *Pleosporales*, *Dothideomycetes*, four species, type: *X. applanata* (Niessl) Q. Chen & L. Cai, coelomycetous, saprobes, ?pathogens, terrestrial, cosmopolitan, see Chen et al. (2015b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), epitype and ex-epitype of the type: HMAS 246688, CBS 195.36.

Xenodium Syd. 1935 (= *Xenodiella* Syd. 1935 *fide* Rossman et al. 2016a), *Sordariomycetes* genera *incertae sedis*, two species, type: *X. petrakii* Syd., asexual morph formerly known in *Xenodiella*, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Jayawardena et al. (2014) excluded from *Elsinoaceae* and placed in *Sordariomycetes*, Rossman et al. (2016a; nomenclature), cultures and sequences are available.

Xenodomus Petr. 1922 (= *Bactropycnis* Höhn. 1920; = *Coleonaema* Höhn. 1924; = *Rhabdostromellina* Höhn. 1917; = *Xenodomus* Petr. 1922 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *X. sydowii* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, North America, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Xenogliocladiopsis Crous & W.B. Kendr. 1994, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *X. eucalyptorum* Crous & W.B. Kendr., coelomycetous, saprobes, Australia, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Lombard et al. (2015b; morphology, phylogeny), epitype and ex-epitype of type: CBS H-21952, CBS 138758.

Xenoheteroconium Bhat, W.B. Kendr. & Nag Raj 1993, *Ascomycota* genera *incertae sedis*, one species, type: *X. bicolor* Bhat, W.B. Kendr. & Nag Raj, hyphomycetous, sexual morph unknown, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Xenolecia Hertel 1984, *Lecideaceae*, *Lecideales*, *Lecanoromycetes*, one species, type: *X. spadicomma* (Nyl.) Hertel, lichenized, see Lumbsch and Huhndorf (2010; outline), Schmull et al. (2011; taxonomy), Kirk et al. (2013; genus accepted), Lücking et al. (2017a; classification), sequences are unavailable.

Xenoleptographium Marinc., T.A. Duong, Z.W. de Beer & M.J. Wingf. 2015, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, one species, type: *X. phialoconidium* Marinc., T.A. Duong, Z.W. de Beer & M.J. Wingf.,

hyphomycetous, sexual morph unknown, saprobes, Indonesia, see Crous et al. (2015e; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type of the type: PREM 61244, CBS 134694.

Xenolophium Syd. 1925 (= *Schizostoma* Ces. & De Not. ex Sacc. 1878 *fide* Species Fungorum 2017), *Pleosporales* genera *incertae sedis*, *Dothideomycetes*, c. five species, type: *X. leve* Syd., asexual morph unknown, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline, treated as *Pleosporales*, genera *incertae sedis*), Hyde et al. (2013; accepted as a genus in *Platystomaceae*), Wijayawardene et al. (2014c; outline, accepted as a genus in *Platystomaceae*), cultures and sequences are available.

Xenomeris Syd. 1924, *Dothideomycetes* genera *incertae sedis*, eleven species, type: *X. pruni* Syd., asexual morph hormonema-like, Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available but lacking for the type, need generic revision.

Xenomycosphaerella Quaedvl. & Crous 2014, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *X. elongata* (Crous & M.J. Wingf.) Quaedvl. & Crous, asexual morph unknown, pathogens, Brazil, Venezuela, see Quaedvlieg et al. (2014; morphology, phylogeny), Guatimosim et al. (2016; morphology, phylogeny), holotype and ex-type culture of type: CBS-H 19824, CPC 13378.

Xenomyxa Syd. 1939, *Ascomycota* genera *incertae sedis*, one species, type: *X. disseminata* Syd., asexual morph unknown, saprobes, terrestrial, Ecuador, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Xenonectriella Weese 1919, *Nectriaceae*, *Hypocreales*, *Sordariomycetes*, twelve species, type: *X. lutescens* (Arnold) Weese, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Etayo and van den Boom (2013; new species, Dominican Republic), Kirk et al. (2013; genus accepted), Brackel et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Xenopeltis Syd. & P. Syd. 1919, *Ascomycota* genera *incertae sedis*, one species, type: *X. philippinensis* Syd. & P. Syd., coelomycetous, sexual morph unknown, saprobes, terrestrial, Philippines, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Xenopenidiella Quaedvl. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *X. rigidophora* (Crous, R.F. Castañeda & U. Braun) Quaedvl. & Crous, hyphomycetous, saprobes, Cuba, see Quaedvlieg et al. (2014; morphology, phylogeny), holotype and ex-type culture of type: CBS H-19938, CBS 314.95.

Xenophacidiella Crous 2011, *Dothideomycetes* genera *incertae sedis*, one species, type: *X. pseudocatenata* Crous, hyphomycetous, sexual morph unknown, saprobes, South Africa, terrestrial, see Crous and Groenewald (2011; morphology, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type culture of type: CBS H-20536, CPC 18472.

Xenophaeosphaeria Crous & M.J. Wingf. 2014, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *X. grewiae* Crous & M.J. Wingf., asexual morph unknown, Tanzania, terrestrial, see Crous et al. (2014d; morphology, phylogeny), holotype and ex-type culture of type: CBS H-21999, CPC 24398.

Xenoplaca Petr. 1949, *Ascomycota* genera *incertae sedis*, one species, type: *X. aequatoriensis* Petr., hyphomycetous, sexual morph unknown, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Xenopolyscytalum Crous 2010, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, one species, type: *X. pinea* Crous, hyphomycetous, sexual morph unknown, The Netherlands, terrestrial, see Crous and Groenewald (2010; morphology, phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type culture of type: CBS-H 20488, CPC 14225.

Xenopyricularia Klaubauf, Lebrun & Crous 2014, *Pyriculariaceae*, *Magnaportheales*, *Sordariomycetes*, one species, type: *X. zizaniicola* (Hashioka) Klaubauf, Lebrun & Crous, hyphomycetous, sexual morph unknown, pathogens, terrestrial, worldwide, see Klaubauf et al. (2014; taxonomy, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), Wijayawardene et al. (2017; outline), neotype and ex-neotype cultures for the type: CBS H-21846, CBS133593 = MAFF 240219.

Xenoramularia Videira, H.D. Shin & Crous 2016, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, three species, type: *X. polygonicola* Videira, H.D. Shin & Crous hyphomycetous, sexual morph unknown, plant pathogens, see Videira et al. (2016; morphology, phylogeny), holotype KUS-F19688, isotype CBS H-22541, culture ex-type CBS 141102 = CPC 10852.

Xenoseptoria Quaedvl., H.D. Shin, Verkley & Crous 2013, *Phaeosphaeriaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *X. neosaccardoi* Quaedvl., H.D. Shin,

Verkley & Crous, coelomycetous, sexual morph unknown, pathogenic, terrestrial, South Korea, see Quaedvlieg et al. (2013; morphology, phylogeny), holotype and ex-type strains of the type: CBS H-21312; CBS 128665.

Xenosonderhenia Crous 2012, *Capnodiales* genera *incertae sedis*, *Dothideomycetes*, two species, type: *X. syzygii* Crous, coelomycetous, saprobes, terrestrial, worldwide, see de Gruyter et al. (2012; morphology; phylogeny), Wijayawardene et al. (2017; outline), holotype and ex-type strain of the type: CBS H-20968; CPC 19790.

Xenospadicoides Hern.-Restr., J. Mena & Gene 2017, *Xenospadicoidaceae*, *Xenospadicoidales*, *Sordariomycetes*, one species, type: *X. atra* (Corda) Hern.-Restr., J. Mena & Gene, hyphomycetous, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Xenopodium Penz. & Sacc. 1902 (= *Xenoporella* Höhn. 1923 [1924] *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, 18 species, type: *X. mirabile* Penz. & Sacc., hyphomycetous, sexual morph unknown, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Xenostigme Syd. 1930, *Sordariomycetes* genera *incertae sedis*, one species, type: *X. trichophila* Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Meliolaceae*), Kirk et al. (2013; genus accepted), Hongsanan et al. (2015d; excluded from *Meliolaceae* and placed in *Sordariomycetes* genera *incertae sedis*), cultures and sequences are unavailable.

Xenostigmella Crous 1998, *Melanommataceae*, *Pleosporales*, *Dothideomycetes*, two species, type: *X. zilleri* (A. Funk) Crous, hyphomycetous, saprobes, terrestrial, Canada, British Columbia, see Crous et al. (2009a; morphology, phylogeny), Wijayawardene et al. (2012, 2014c, 2017; outline; phylogeny), Kirk et al. (2013; genus accepted), Tian et al. (2015; morphology, nomenclature, phylogeny), cultures and sequences are available, ex-type strains of type: CBS 115685 = CPC 4010; holotype of type: DAVFP 23272.

Xenostomella Syd. 1930 (= *Plochmothea* Syd. 1939 *fide* Species Fungorum 2017), *Dothideomycetes* genera *incertae sedis*, two species, type: *X. tovarensis* Syd., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline, accepted as a genus in *Microthyriaceae*), Kirk et al. (2013; genus accepted), Wu et al. (2011) and Hyde et al. (2013) did not accept as *Microthyriaceae*, Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Xenostroma Höhn. 1915, *Ascomycota* genera *incertae sedis*, one species, type: *X. caespitosum* (Fuckel) Höhn.,

coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, need generic revision.

Xenoteratosphaeria Quaedvl. & Crous 2014, *Teratosphaeriaceae*, *Capnodiales*, *Dothideomycetes*, one species, type: *X. jonkershoekensis* (P.S. van Wyk, Marasas & Knox-Dav.) Quaedvl. & Crous, asexual morph unknown, pathogens, South Africa, see Quaedvlieg et al. (2014; morphology, phylogeny), epitype and ex-epitype culture of type: CBS H-20095, CBS 122897.

Xenothecium Höhn. 1919, *Hypocretriaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *X. jodophilum* Höhn., asexual morph unknown, saprobes, terrestrial, South America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), holotype of the type: BRIP 22762, cultures and sequences are unavailable.

Xenotypa Petr. 1955 (= *Radulum* Fr. 1825; = *Tylodon* Banker 1902 *fide* Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, one species, type: *X. aterrima* (Fr.) Petr., asexual morph unknown, saprobes, terrestrial, Europe, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Xenus Kohlm. & Volk. Kohlm., 1992, *Pyrenulales* genera *incertae sedis*, *Eurotiomycetes*, one species, type: *X. lithophylli* Kohlm. & Volk. Kohlm., asexual morph unknown, saprobes, lichenized, marine, on coral, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jones et al. (2015; outline), Lücking et al. (2017a; classification), cultures and sequences are unavailable.

Xepicula Nag Raj 1993, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, four species, type: *X. leucotricha* (Peck) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Kirk et al. (2013; genus accepted), Lombard et al. (2016; new species, phylogeny), Wijayawardene et al. (2016b, 2017; outline, taxonomy), ex-type of the type: CBS 256.57 = MUCL 9860, need generic revision.

Xepiculopsis Nag Raj 1993, *Stachybotryaceae*, *Hypocreales*, *Sordariomycetes*, two species, type: *X. perpulchra* Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Lombard et al. (2016; taxonomic notes), cultures are unavailable.

Xerochrysium Pitt 2013, *Trichocomaceae*, *Eurotiales*, *Eurotiomycetes*, two species, type: *X. dermatitidis* (A. Agostini) Pitt, hyphomycetous, sexual morph unknown,

saprobes, terrestrial, Europe, Australia, see Pitt et al. (2013; taxonomy, phylogeny), lectotype and ex-lectotype of the type: CBS 132.31, IMI 96729=UAMH 802.

Xeroconium D. Hawksw. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *X. borealis* (P. Karst.) D. Hawksw., on wood, Europe, terrestrial, sexual morph unknown, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, holotype of type: H-Karst 1493.

Xerombrophila Baral 2013, *Gelatinodiscaceae*, *Helotiales*, *Leotiomycetes*, one species, type: *X. crystallifera* Baral, G. Marson & Unter., asexual morph hyphomycetous, saprobes, terrestrial, Europe, see Baral et al. (2013b; taxonomy, phylogeny), Wijayawardene et al. (2017; outline), ex-type cultures of the type: CBS 128289.

Xeromedulla Korf & W.Y. Zhuang 1987, *Helotiales* genera *incertae sedis*, *Leotiomycetes*, three species, type: *X. leptospora* W.Y. Zhuang & Korf, asexual morph hyphomycetous, saprobes, terrestrial, China, Philippines, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xeromyces L.R. Fraser 1954 (= *Fraseriella* Cif. & A.M. Corte *fide* Rossman et al. 2016a), *Aspergillaceae*, *Eurotiales*, *Eurotiomycetes*, one species: *X. bisporus* L.R. Fraser, type: *Xeromyces bisporus* L.R. Fraser asexual morph hyphomycetous (*Fraseriella bispora* Cif. & A.M. Corte *fide* Petterson et al. 2011 and Seifert et al. 2011), terrestrial, strongly xerophilic, isolated from dry, sugar-rich foodstuffs, cosmopolitan, see Seifert et al. (2011; morphology), Petterson et al. (2011; phylogeny, morphology), Under unitary nomenclature (Hawksworth et al. 2011a), the names *Xeromyces* and *Fraseriella* compete for use. Since the former genus name is more widely used and older than the latter, we consider *Fraseriella* to represent a synonym of *Xeromyces*. The synonymy is formally proposed herein as follows:

Xeromyces bisporus L.R. Fraser, Proc. Linn. Soc. N.S.W. 78:245. 1954.

≡ *Monascus bisporus* (L.R. Fraser) Arx, Gen. Fungi Spor. Pure Cult.: 84. 1970.

= *Fraseriella bispora* Cif. & A.M. Corte, in Corte Atti Ist. Bot. Univ. Lab. crittog. Pavia Ser. 4 14:109. 1957.

Xerotrema Sherwood & Coppins 1980, *Odontotremataceae*, *Ostropales*, *Lecanoromycetes*, two species, type: *X. megalospora* Sherwood & Coppins, asexual morph unknown, ?saprobes, terrestrial, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xiambola Minter & Hol.-Jech. 1981, *Ascomycota* genera *incertae sedis*, one species, type: *X. mirabilis*

Minter & Hol.-Jech., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xiphomyces Syd. & P. Syd. 1916, *Ascomycota* genera *incertae sedis*, two species, type: *X. sacchari* Syd. & P. Syd., hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xiuguo Zhangia K. Zhang, R.F. Castañeda, Jian Ma & L.G. Ma 2014, *Ascomycota* genera *incertae sedis*, five species, type: *X. rosae* (K. Zhang & X.G. Zhang) K. Zhang & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Zhang et al. (2014b; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Xyladictyochaeta Hern.-Restr., R.F. Castaneda & Gene 2017, *Castanediellaceae*, *Xylariales*, *Sordariomycetes*, one species, type: *X. lusitanica* Hern.-Restr., J. Mena & Guarro, hyphomycetous, saprobes, Europe, see Hernández-Restrepo et al. (2017; taxonomy, phylogeny), sequences are available.

Xylaria Hill ex Schrank 1789 (= *Moelleroclavus* Henn. 1902; = *Xylocoremium* J.D. Rogers 1984 *fide* Rébllová et al. 2016c; see Species Fungorum 2017 for additional synonyms), *Xylariaceae*, *Xylariales*, *Sordariomycetes*, c. 350 species, type: *X. hypoxylon* (L.) Grev., asexual morph hyphomycetous, terrestrial, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Fournier et al. (2011; new species), Rogers and Ju (2012; new species), Ma et al. (2012a, 2013; new species), Kirk et al. (2013; genus accepted), Gu et al. (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Senanayake et al. (2015; outline, phylogeny), Rébllová et al. (2016c; new species; nomenclature), cultures and sequences are available.

Xyleborus R.C. Harris & Ladd 2007, *Stereocaulaceae*, *Lecanorales*, *Lecanoromycetes*, two species, type: *X. sporodochifer* R.C. Harris & Ladd, asexual morph unknown, lichenized, terrestrial, see Lumbsch and Huhndorf (2010; outline), Lendemer et al. (2013; list), Lendemer and Harris (2015; new species), Lücking et al. (2017a; classification), sequences unavailable.

Xylobotryum Pat. 1895 (= *Melanobotrys* 1926 [1925]; = *Trachyxyloaria* 1901; = *Xyloceras* A.L. Sm. 1901 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, two species, type: *X. andinum* Pat., asexual morph unknown, saprobes, terrestrial, tropical, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylochia B. Sutton 1983, Ascomycota genera *incertae sedis*, one species, type: *X. stegonsporoides* B. Sutton, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylochora Arx & E. Müll. 1954, Xylariomycitidae genera *insertae sedis*, Sordariomycetes, three species, type: *X. craticola* (H. Wegelin) Arx & E. Müll., asexual morph unknown, saprobes, Sweden, Switzerland and India, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylocrea Möller 1901, Xylariales genera *incertae sedis*, Sordariomycetes, one species, type: *X. piriformis* Möller, asexual morph unknown, saprobes, see Kirk et al. (2008; treated as a synonym of *Sarcoxylo* Cooke), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Maharachchikumbura et al. (2015b, 2016b; outline, accepted as a distinct genus), Wendt et al. (2017; accepted as a genus in Xylariales genera *incertae sedis*), cultures and sequences are unavailable.

Xyloglyphis Clem. 1909, Ascomycota genera *incertae sedis*, one species, type: *X. striola* (Fr.) Clem., coelomycetous, sexual morph unknown, saprobes, terrestrial, Europe, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylogramma Wallr. 1833 (= *Globuligera* (Sacc.) Höhn. 1918; = *Patellea* (Fr.) Sacc. 1884; = *Patinella* subgen. *Globuligera* Sacc. 1889; = *Peziza* trib. *Patellea* Fr. 1822; = *Scutularia* P. Karst. 1885 [1881-1885]; = *Sphaeropeziella* P. Karst. (1885) [1881-1885] fide Species Fungorum 2017), Helotiaceae, Helotiales, Leotiomycetes, 18 species, type: *X. sticticum* (Fr.) Wallr., asexual morph unknown, saprobes, terrestrial, China, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylographa (Fr.) Fr., 1836 (= *Spiloma* Ach. 1803; = *Spilonematopsis* Vain. 1909; = *Stictis* subgen. *Xylographa* Fr. 1822; = *Xylographomyces* Cif. & Tomas. 1953 fide Species Fungorum 2017), Xylographaceae, Trapeliales, Lecanoromycetes, 20 species, type: need typification, see Heininger and Spribille (2009; new species), Wetmore (2009; floristic study), Zhdanov (2009; new record), Agnello et al. (2010; ecology), Bilovitz et al. (2010; biodiversity), Brodo (2010; new record), Lumbsch and Huhndorf (2010; outline), Hansen (2012; new record), Mayrhofer et al. (2012; new records), Bennett (2013; review), Hauck et al. (2013b; new record), Kirk et al. (2013; genus accepted), Mayrhofer et al. (2013; new record), Svensson (2013; biodiversity), Svensson et al.

(2013, 2016; ecology, biodiversity), Tønsberg (2013; new species), Urbanavichene et al. (2013, new record), Abbas et al. (2014; new records), Goni et al. (2014; new record), Spribille et al. (2014a; systematic, new species), Hämäläinen et al. (2015; biodiversity), Malíček et al. (2015b; new record), Resl et al. (2015; phylogeny), Tarasova et al. (2015, 2016; new record), Węgrzyn et al. (2015; new record), Himelbrant et al. (2016; new record), Kharpukhæva and Mukhortova (2016; ecology), Motiejūnaitė et al. (2016; new record), Lücking et al. (2017a; classification), sequences are available.

Xylohypha (Fr.) E.W. Mason 1960 (= *Hyphelia* sect. *Xylohypha* Fr. 1849; = *Xylohypha* (Fr.) E.W. Mason 1953 fide Species Fungorum 2017), Ascomycota genera *incertae sedis*, six species, type: *X. nigrescens* (Pers.) E.W. Mason, hyphomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylohypopsis W.A. Baker & Partr. 2000, Ascomycota genera *incertae sedis*, two species, type: *X. curta* (Corda) W.A. Baker & Partr., hyphomycetous, sexual morph unknown, saprobes, terrestrial, South Africa, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylomelasma Réblová 2006, Sordariomycetidae genera *incertae sedis*, Sordariomycetes, three species, type: *X. sordida* Réblová, asexual morph hyphomycetous, saprobes, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Vasilyeva and Stephenson (2014; new species), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), Réblová et al. (2016a; DNA, phylogeny), cultures are available for the type: CBS 131683.

Xylomyces Goos, R.D. Brooks & Lamore 1977, Aliquandostipitaceae, Jahnulales, Dothideomycetes, ten species, type: *X. chlamydosporus* Goos, R.D. Brooks & Lamore, hyphomycetous, sexual morph unknown, aquatic (freshwater/marine), see Shearer et al. (2009; phylogeny), Sivichai and Boonyuen (2010; association with *Jahnula*), Seifert et al. (2011; morphology), Hu et al. (2010a; diversity), Barbosa and Gusmão (2011; diversity), Suetrong et al. (2011a; phylogeny), Sivichai et al. (2011; taxonomy), Zhang et al. (2011a; checklist), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2012, 2014c, 2017; outline), Hyde et al. (2013; accepted as a genus in Aliquandostipitaceae), Borse et al. (2014; diversity), da Silva et al. (2015; diversity), Jones et al. (2015; marine genera), Oliveira et al. (2015; diversity), cultures available for the type: SS2917.

Xylona Gazis & P. Chaverri 2012, *Xylonomycetaceae*, *Xylonomycetales*, *Xylonomycetes*, *T. heveae* Gazis & P. Chaverri, coelomycetous, sexual morph unknown, endophytes, see Gazis et al. (2016; phylogeny), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Xylopezia Höhn. 1917, *Dothideomycetes* genera *incertae sedis*, type: data inadequate, see Kirk et al. (2008; treated as a synonym of *Exarmidium*), Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; did not list), Wijayawardene et al. 2014c; outline), sequences are unavailable.

Xylopsora Bendiksby & Timdal 2013, *Umbilicariaceae*, *Umbilicariales*, *Lecanoromycetes*, two species, type: *X. friesii* (Ach.) Bendiksby & Timdal, lichenized, lignicolous, corticolous, often on charred substrata, temperate northern hemisphere, Australia, see Bendiksby and Timdal (2013; phylogenetic placement), Lücking et al. (2017a; classification), type specimen H-ACH-436A sequences are available.

Xyloschistes Vain. ex Zahlbr. 1903 (= *Xyloschistomyces* Cif. & Tomas. 1903 *fide Species Fungorum* 2017), *Stictidaceae*, *Ostropales*, *Lecanoromycetes*, one species, type: *X. platytropa* (Nyl.) Vain. ex Zahlbr., lichenized, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Miadlikowska et al. (2014a; phylogeny), Lücking et al. (2017a; classification), sequences are available.

Xyloschizon Syd. 1922, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, two species, type: *X. weirianum* Syd., asexual morph unknown, saprobes, terrestrial, North America, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Xylotumulus J.D. Rogers, Y.M. Ju & Hemmes 2006, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *X. gibbisporus* J.D. Rogers, Y.M. Ju & Hemmes, asexual morph unknown, saprobes, terrestrial, Hawaii, see Lumbsch and Huhndorf (2010; outline), Rogers and Ju (2012; Hawaii fungi), Chang et al. (2014; secondary compounds), Maharachchikumbura et al. (2015b, 2016b; outline), Wendt et al. (2017; accepted as a genus in *Xylariales* genera *incertae sedis*), holotype: BISH, Rogers, 2 Aug. 2005, cultures available for the type: NBRC 9453, ATCC MYA-4109.

Yalomycetes Nag Raj 1993, *Ascomycota* genera *incertae sedis*, six species, type: *Y. pulcher* Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, worldwide, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Yamadazyma Billon-Grand 1989, *Debaryomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, 20 species, type:

Y. philogaea (van der Walt & Johannsen) Billon-Grand, asexual reproduction is by multilateral budding, saprophytic, in tree exudates and insect frass, insect guts, soil, cactus rots, seawater, soy sauce, infrequent clinical isolates, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), many species in this genus represent asexual taxa originally classified in the genus *Candida*, thus they will be transferred to *Yamadazyma* as new combinations, cultures and sequences are available.

Yarrowia van der Walt & von Arx 1980, *Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, 13 species, type: *Y. lipolytica* (Wickerham, Kurtzman & Herman) van der Walt & von Arx, asexual reproduction is by multilateral budding, saprophytic, but clinical isolates are not uncommon, on wet milled maize, refrigerated meats, soil, clinical, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), some species in this genus represent asexual taxa originally classified in the genus *Candida* thus they will be transferred to *Yarrowia* as new combinations, cultures and sequences are available.

Yarrumia D.J. Galloway 2015, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, two species, type: *Y. coronata* D.J. Galloway, lichenized, see Galloway (2015; taxonomy), Lücking et al. (2017a; classification); sequences are available.

Yatesula Syd. & P. Syd. 1917 (= *Ciferriusia* Bat. 1962 *fide Species Fungorum* 2017), *Chaetothyriaceae*, *Chaetothyriales*, *Eurotiomycetes*, two species, type: *Y. calami* Syd. & P. Syd., asexual morph unknown, epiphytic, sooty moulds, terrestrial, America, Philippines, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Yinmingella Goh, K.M. Tsui & K.D. Hyde 1999, *Ascomycota* genera *incertae sedis*, six species, type: *Y. mitriformis* Goh, K.M. Tsui & K.D. Hyde, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, South America, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Yoshimuria S.Y. Kondr., Kärnefelt, Elix, A. Thell & Hur 2014, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, three species, type: *Y. spodoplaca* (Nyl.) S.Y. Kondr., Kärnefelt, Elix, A. Thell, J. Kim, A.S. Kondr. & Hur, lichenized, see Kondratyuk et al. (2014a; taxonomy), Lücking et al. (2017a; classification), cultures available for the type: SK725, SK724.

Yoshimuriella Moncada & Lücking 2013, *Lobariaceae*, *Peltigerales*, *Lecanoromycetes*, eight species, type: *Y. fendleri* (Tuck. & Mont.) Moncada & Lücking, lichenized, see Moncada et al. (2013a; taxonomy),

Lücking et al. (2017a; classification), sequences are available.

Yoshinagaia Henn. 1904 (= *Japonia* Höhn. 1909; = *Yoshinagamyces* Hara 1912; = *Monoloculia* Hara 1927 *fide* Rossman et al. 2016a), *Dothideomycetes* genera *incertae sedis*, two species, type: *Y. quercus* Henn., asexual morph unknown, saprobes, terrestrial, Japan, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Rossman et al. (2016a; nomenclature), cultures and sequences are unavailable.

Yoshinagella Höhn. 1913, *Dothideomycetes* genera *incertae sedis*, four species, type: *Y. japonica* Höhn, asexual morph unknown, saprobes and parasitic, reported from Japan and Hawaii, see Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), Dai et al. (2014a; phylogeny, *Dothideomycetes*), cultures and sequences are unavailable.

Ypsilina J. Webster, Descals & Marvanová 1999, *Ascomycota* genera *incertae sedis*, one species, type: *Y. graminea* (Ingold, P.J. McDougall & Dann) Descals, J. Webster & Marvanová, hyphomycetous, sexual morph unknown, aquatic, see Seena et al. (2010; DNA barcoding of fungi), Hyde et al. (2011; checklist), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), sequences are available:

Ypsilomyces D.A.C. Almeida & Gusmão 2014, *Ascomycota* genera *incertae sedis*, one species, type: *Y. elegans* D.A.C. Almeida & Gusmão, hyphomycetous, sexual morph unknown, one species, saprobes, terrestrial, Brazil, see Almeida and Gusmão (2014; morphology), holotype of the type: HUEFS 155081, cultures and sequences are unavailable, compare with *Biflagellospora*, *Diademospora*, *Diplocladiella*, *Iyengarina*, *Vanterpoolia*, *Weufia* and *Ypsilina*.

Ypsilonilia Lév. 1846 (= *Acanthothecium* Speg. 1889; = *Psalidosperma* Syd. & P. Syd. 1914 *fide* Species Fungorum 2017), *Sordariales* genera *incertae sedis*, five species, type: *Y. cuspidata* Lév., coelomycetous, sexual morph of unknown, saprobes, aquatic, see Hyde et al. (2011; outline), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Yuccamyces Gour, Dyko & B. Sutton 1979, *Ascomycota* genera *incertae sedis*, seven species, type: *Y. purpureus* Gour, Dyko & B. Sutton, hyphomycetous, sexual morph unknown, saprobes, Canada and Cuba, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Yuea O.E. Erikss. 2003, *Xylariales* genera *incertae sedis*, *Sordariomycetes*, one species, type: *Y. chusqueicola*

O.E. Erikss., asexual morph unknown, terrestrial, saprobes, Chile, see Lumbsch and Huhndorf (2010; outline), Maharachchikumbura et al. (2015b, 2016b; outline, phylogeny), cultures and sequences are unavailable.

Yueomyces Q.M. Wang, L. Wang, M. Groenew. & T. Boekhout 2015, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, one species, type: *Y. sinensis* (J.Z. Yue) Q.M. Wang, L. Wang, M. Groenew. & T. Boekhout, asexual reproduction is by bipolar budding, saprophytic, Soil, China, see Wang et al. (2015a; taxonomy), cultures and sequences are available.

Zahlbrucknerella Herre 1912 (= *Lecanephebe* Frey 1929; = *Leptopterygium* Zahlbr. 1930; = *Zahlbrucknera* Herre 1910 *fide* Species Fungorum 2017), *Lichenaceae*, *Lichinales*, *Lichinomycetes*, eight species, type: *Z. calcarea* (Herre) Herre, asexual morph unknown, lichenized, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zakatoshia B. Sutton 1973, *Ascomycota* genera *incertae sedis*, two species, type: *Z. hirschiopori* B. Sutton, hyphomycetous, sexual morph unknown, fungicolous, terrestrial, worldwide, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zalerion R.T. Moore & Meyers 1962, polyphyletic, six species, type: *Z. nepura* R.T. Moore & Meyers, hyphomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), marine species referred to *Lulworthiaceae*, *Lulorthiales*, *Sordariomycetes* and *Leotiaceae*, *Helotiales*, *Leotiomycetes* (see Jones et al. 2009, 2015; outline, phylogeny), cultures and sequences available. Terrestrial species need re-evaluation and need to be reassigned, cultures and sequences are available.

Zanclospora S. Hughes & W.B. Kendr. 1965, *Chaetosphaeriaceae*, *Chaetosphaeriales*, *Sordariomycetes*, ten species, type: *Z. novae-zelandiae* S. Hughes & W. B. Kendr., hyphomycetous, sexual morph unknown, saprobes, tropical and subtropical distribution with only two records in temperate region, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Almeida et al. (2013; new species), Kirk et al. (2013; genus accepted), Hernández-Restrepo et al. (2017; taxonomy, phylogeny), cultures and sequences are available.

Zasmidium Fr. 1849, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, c. 70 species names, type: *Z. cellare* (Pers.) Fr., hyphomycetous, sexual morph mycosphaerella-like, five species confirmed based on

DNA, plant pathogens, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Braun et al. (2013; morphology), Phengsintham et al. (2013; species in Laos), Videira et al. (2017; morphology, phylogeny), cultures and sequences are available.

Zebraspora McKenzie 1991, *Ascomycota* genera *incertae sedis*, two species, type: *Z. bicolor* McKenzie, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, Pacific Islands, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zelandiocoela Nag Raj 1993, *Ascomycota* genera *incertae sedis*, one species, type: *Z. ambigua* (Nag Raj & W.B. Kendr.) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, New Zealand, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zeloasperisporium R.F. Castañeda 1996 (= *Neomicrothyrium* Boonmee, H.X. Wu & K.D. Hyde 2014 *fide* Hongsanan et al. 2015b), *Zeloasperisporiaceae*, *Zeloasperisporiales*, *Dothideomycetes*, seven species, type: *Z. hyphopodioides* R.F. Castañeda, hyphomycetous, sexual morph unknown, in air, saprobes, Asia, Australasia, Caribbean, see Cheewangkoon et al. (2009; new species, phylogeny), Seifert et al. (2011; morphology), Crous et al. (2015e; new species, phylogeny), Hongsanan et al. (2015b, 2016c; new species, phylogeny, evolution), Wijayawardene et al. (2017; outline), cultures and sequences are available.

Zelodactylaria A.C. Cruz, Gusmão & R.F. Castañeda 2012, *Ascomycota* genera *incertae sedis*, one species, type: *Z. verticillata* A.C. Cruz, Gusmão & R.F. Castañeda, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Cruz et al. (2012; taxonomy), Wijayawardene et al. (2017; outline), cultures and sequences are unavailable.

Zelopelta B. Sutton & R.D. Gaur 1984 (= *Cladosporothrygium* Katum. 1984 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, one species, type: *Z. thrinacospora* B. Sutton & R.D. Gaur, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zelosatchmopsis Nag Raj 1991, *Ascomycota* genera *incertae sedis*, one species, type: *Z. sacciformis* (R.F. Castañeda) Nag Raj & R.F. Castañeda, coelomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zernya Petr. 1947, *Ascomycota* genera *incertae sedis*, one species, type: *Z. clypeata* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Brazil, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zeroviella S.Y. Kondr. & Hur 2015, *Teloschistaceae*, *Teloschistales*, *Lecanoromycetes*, eight species, type: *Z. papillifera* (Vain.) S.Y. Kondr. & Hur, lichenized, see Kondratyuk et al. (2015d; taxonomy), Lücking et al. (2017a; classification), sequences are available.

Zetesimomyces Nag Raj 1988, *Ascomycota* genera *incertae sedis*, one species, type: *Z. setibicolor* (R.F. Castañeda) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, Cuba, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zetiasplozna Nag Raj 1993 (= *Pestalotia* subgen. *Pestalozzina* Sacc. 1884; = *Pestalozzina* (Sacc.) Sacc. 1895 *fide* Species Fungorum 2017), *Sporocadaceae*, *Amphisphaerales*, *Sordariomycetes*, six species, type: *Z. unicolor* (Berk. & M.A. Curtis) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, cosmopolitan, see Nicoletti et al. (2010; endophytes), Wijayawardene et al. (2012, 2016b, 2017; outline, taxonomy), Kirk et al. (2013; genus accepted), Crous et al. (2014c; new species), Senanayake et al. (2015; outline), Jaklitsch et al. (2016a; accepted as a genus in *Sporocadaceae*), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available.

Zeugandromyces Thaxt. 1912, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *Z. australis* Thaxt., asexual morph unknown, biotrophic on Coleoptera (Staphylinidae, Paederini), America, Philippines, Sumatra, Sri Lanka, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zeus Minter & Diam. 1987, *Rhytismataceae*, *Rhytismatales*, *Leotiomycetes*, two species, type: *Z. olympius* Minter & Diam., asexual morph unknown, saprobes, terrestrial, Greece, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zevadia J.C. David & D. Hawksw. 1995, *Ascomycota* genera *incertae sedis*, one species, type: *Z. peroccidentalis* J.C. David & D. Hawksw., lichenicolous, Europe, sexual morph unknown, see Seifert et al. (2011; genus accepted), Kirk et al. (2013; genus accepted), no cultures or sequences are available, holotype of type: K(M)-IMI 401823.

Zilingia Petr. 1934, *Ascomycota* genera *incertae sedis*, one species, type: *Z. sibirica* Petr., coelomycetous, sexual morph unknown, saprobes, terrestrial, Siberia, see

Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zimmermanniella Henn. 1902, *Phyllachoraceae*, *Phyllachorales*, *Sordariomycetes*, one species, type: *Z. trispora* Henn., from living leaves, South East Asia, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Zinzipegasa Nag Raj 1993, *Ascomycota* genera *incertae sedis*, one species, type: *Z. argentinensis* (Speg.) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, Argentina, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zodiomyces Thaxt. 1891, *Laboulbeniaceae*, *Laboulbeniales*, *Laboulbeniomycetes*, four species, type: *Z. vorticellarius* Thaxt., asexual morph unknown, biotrophic on aquatic Coleoptera (Hydrophilidae), worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Rossi et al. (2016b; most recent species description), ribosomal sequence are available.

Zoggium Lar.N. Vassiljeva 2001, *Mytilinidiaceae*, *Mytilinidiales*, *Dothideomycetes*, one species, type: *Z. mayorii* (H. Zogg) Lar.N. Vassiljeva, asexual morphs unknown, saprobes, terrestrial, north temperate, Boehm et al. (2009a, b; accepted as in *Mytilinidiaceae*), Hyde et al. (2013; accepted as in *Mytilinidiaceae*), Wijayawardene et al. (2014c; outline), cultures and sequences are unavailable.

Zopfia Rabenh. 1874 (= *Pontoporeia* Kohlm. 1963 *fide* Species Fungorum 2017), *Zopfiaceae*, *Pleosporales*, *Dothideomycetes*, five species, type: *Z. rhizophila* Rabenh., asexual morph unknown, saprobes, terrestrial, worldwide, see Schoch et al. (2009; phylogeny), Lumbsch and Huhndorf (2010; outline), Hyde et al. (2013; accepted as a genus in *Zopfiaceae*), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2014c; outline), cultures and sequences are available.

Zopfiella G. Winter 1884 (= *Ariefia* Jacz. 1922; = *Tripterospora* Cain 1956 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, 30 species, type: *Z. tabulata* (Zopf) G. Winter, asexual morph unknown, coprophilous, saprobes, terrestrial, marine, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Zopfiofoveola D. Hawksw. 1979, *Zopfiaceae*, *Pleosporales*, *Dothideomycetes*, one species, type: *Z. punctata* (D. Hawksw. & C. Booth) D. Hawksw., on earthworm excrements, Sweden, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Jaklitsch et al. (2016a; classification), cultures and sequences are unavailable, holotype of type: (microscope slide) UPS.

Zopheromyces B. Sutton & Hodges 1977, *Ascomycota* genera *incertae sedis*, one species, type: *Z. fasciatus* B. Sutton & Hodges, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Australasia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Zugazaea Korf, Iturr. & Lizoň 1998, *Helotiales* genera *incertae sedis*, *Leotiomyces*, one species, type: *Z. agyrioides* Korf, Iturr. & Lizoň, asexual morph unknown, saprobes, Canary Islands, Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), holotype of the type: CUP-MM 2844, cultures and sequences are unavailable.

Zukaliopsis Henn. 1904, *Myriangiaceae*, *Myriangiales*, *Dothideomycetes*, one species, type: *Z. amazonica* Henn., asexual morph unknown, parasitic, terrestrial, Peru, isotype of the type: S, F5604, see Dissanayake et al. (2014; morphology), Wijayawardene et al. (2014c; outline).

Zunura Nag Raj 1993, *Ascomycota* genera *incertae sedis*, one species, type: *Z. appendiculata* (B. Sutton) Nag Raj, coelomycetous, sexual morph unknown, saprobes, terrestrial, India, see Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable.

Table 2 Nom. dub. genera in Ascomycota (based on Kirk et al. 2008) which were not included in recent literature

Generic name	Authors	Year	Comments
<i>Asteronema</i>	Trevis.	1845	
<i>Blastotrichum</i>	Corda	1838	
<i>Borinquenia</i>	F. Stevens	1917	
<i>Botryodiplodia</i>	Sacc.	1884	
<i>Byssocladium</i>	Link	1816	
<i>Calothyriopeltis</i>	F. Stevens & R.W. Ryan	1925	
<i>Cellulosporium</i>	Peck	1879	
<i>Chaetobasidiella</i>	Höhn.	1918	
<i>Chmelia</i>	Svob.-Pol.	1966	
<i>Coccosporium</i>	Corda	1831	
<i>Coniothecium</i>	Corda	1833	
<i>Cyphellopsus</i>	Fayod	1889	
<i>Dinemasporiella</i>	Spieg.	1911	
<i>Diplodiella</i>	(P. Karst.) Sacc.	1884	
<i>Endaematus</i>	Raf.	1814	
<i>Halterophora</i>	Endl.	1836	
<i>Hypha</i>	Pers.	1822	
<i>Hyphasma</i>	Rebent. ex Nocca & Balb.	1821	
<i>Leptodothis</i>	Theiss. & Syd.	1914	
<i>Leucosporium</i>	Corda	1833	Seifert et al. (2011) also treated as nom. dub.
<i>Levieuxia</i>	Fr.	1849	
<i>Maurya</i>	Pat.	1898	Nom. illegit., Art. 53.1 (Index Fungorum 2017)
<i>Microstelium</i>	Pat.	1899	
<i>Myxonyphe</i>	Rchb.	1841	
<i>Neottiosporella</i>	Höhn. ex Falck	1923	Seifert et al. (2011) treated as a synonym of <i>Minimidochium</i> B. Sutton 1969
<i>Phaeoantenariella</i>	Cavalc.	1969	
<i>Piptostoma</i>	Berk. & Broome	1873	
<i>Piptostomum</i>	Lév.	1845	
<i>Rhinotrichum</i>	Corda	1837	
<i>Scopularia</i>	Preuss	1851	Nom. illegit., Art. 53.1 (Index Fungorum 2017)
<i>Spermatodium</i>	Fée ex Trevis.	1860	
<i>Stemmaria</i>	Preuss	1851	
<i>Variolaria</i>	Pers.	1794	Nom. illegit., Art. 53.1; Nom. rejic., see Arts 14.6, Ex. 5 and 14.7 (Index Fungorum 2017)
<i>Voglinoana</i>	Kuntze	1891	
<i>Weinmannodora</i>	Fr.	1849	
<i>Xylodon</i>	(Pers.) Gray	1821	Nom. rejic., see Arts 14.6, Ex. 5 and 14.7 (Index Fungorum 2017)

Zwackhia Körb. 1855, *Lecanographaceae*, *Arthoniales*, *Arthoniomycetes*, six species, type: *Z. viridis* (Ach.) Poetsch & Schied., lichenized, see Ertz and Tehler (2011; taxonomy) Diederich et al. (2012a; taxonomy), Lücking et al. (2017a; classification); sequences are available.

Zwackhiomacromyces Etayo & van den Boom 2014, ?*Xanthopyrenaceae*, *Collemopsidiales*, *Dothideomycetes*, one species, type: *Z. constrictocarpus* Etayo & Van den

Boom asexual morph unknown, parasymbiont on *Megalospora*, terrestrial, Mediterranean Europe, see van den Boom and Etayo (2014; morphology, distinction of similar lichenicolous genera), holotype of the type: P. & B. van den Boom 27046 (BCN), cultures and sequences are unavailable.

Zwackhiomyces Grube & Hafellner 1990 (= *Zwackhiomyces* Grube & Triebel 1989 nom. inval. *fide*

Species Fungorum 2017), *Xanthopyreniaceae*, *Collemopsidiales*, *Dothideomycetes*, 35 species, type: *Z. coepulonus* (Norman) Grube & R. Sant., asexual morph unknown, lichenicolous, terrestrial, worldwide, see Brackel (2008; new species), Diederich and Schultz (2009; new species), Halici and Candan (2009; new species), Roux (2009; new species), van den Boom (2010; new species), Lumbsch and Huhndorf (2010; outline, family placed as *Ascomycota* families *incertae sedis*), Kocakaya et al. (2011; new species), Pérez-Ortega et al. (2011a, 2016; new species, phylogeny), Kirk et al. (2013; genus accepted), Boom and Etayo (2014; new species), holotype of type: O-F72536, cultures available for the type: RO31.

Zygoascus M. T. Smith 1986, ?*Trichomonascaceae*, *Saccharomycetales*, *Saccharomycetes*, six species, type: *Z. hellenicus* M.T. Smith, asexual reproduction is by multilateral budding, saprophytic, but there are some clinical isolates, in soft drinks, fruit juices, insect tunnels, cactus fruits, soil, tanning fluids, clinical isolates, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Zygopleurage Boedijn 1962, *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, type: *Z. zygospora* (Speg.) Boedijn, saprobes, coprophilous, asexual morph phialophora-like, terrestrial, worldwide, see Richardson (2008, taxonomy), Chang et al. (2010, phylogeny), Mungai et al. (2011; taxonomy), Kruys et al. (2015, phylogeny).

Zygosaccharomyces B.T.P. Barker 1901 (= *Zygosaccharis* Clem. & Shear 1931; = *Zygosaccharomyces* Nishiw. 1929 *fide* Species Fungorum 2017), *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, eleven species, type: *Z. rouxii* (Boutroux) Yarrow, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), cultures and sequences are available.

Zygospermella Cain 1935 (= *Zygospermum* Cain 1934 *fide* Species Fungorum 2017), *Lasiosphaeriaceae*, *Sordariales*, *Sordariomycetes*, six species, type: *Z. setosa* (Cain) Cain, asexual morph unknown, saprobes, terrestrial, worldwide, see Lumbsch and Huhndorf (2010; outline), Kirk et al. (2013; genus accepted), Kruys et al. (2015; DNA, phylogeny), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are available but lacking for the type.

Zygosporium Mont. 1842 (= *Pimina* Grove 1888; = *Urobasidium* Giesenh. 1893; = *Urophiala* Vuill. 1909 *fide* Species Fungorum 2017), *Ascomycota* genera *incertae sedis*, 16 species, type: *Z. oscheoides* Mont.,

hyphomycetous, saprobe, terrestrial, widespread, see Hyde et al. (2011; outline), Seifert et al. (2011; morphology), Kirk et al. (2013; genus accepted), Wijayawardene et al. (2012, 2017; outline), cultures and sequences are unavailable for type species.

Zygotulaspora Kurtzman 2003, *Saccharomycetaceae*, *Saccharomycetales*, *Saccharomycetes*, two species, type: *Z. mrakii* (Capriotti) Kurtzman, asexual reproduction is by multilateral budding, saprophytic, fruit drinks, fruit flies, grape must, silage, worldwide, see Lumbsch and Huhndorf (2010; outline), cultures and sequences are available.

Zymoseptoria Quaedvl. & Crous 2011, *Mycosphaerellaceae*, *Capnodiales*, *Dothideomycetes*, seven species, type: *Z. tritici* (Desm.) Quaedvl. & Crous, coelomycetous, sexual morph mycosphaerella-like, plant pathogens on Poaceae, worldwide, see Quaedvlieg et al. (2011, 2014; morphology, phylogeny), epitype and ex-epitype culture of type: CBS H-20545, IPO 323 = CBS 115943.

Zythia Fr. 1825 (= *Zythia* Fr. 1849 *fide* Species Fungorum 2017), *Gnomoniaceae*, *Diaporthales*, *Sordariomycetes*, type: *Z. resinae* (Ehrenb.) P. Karst. 1890, coelomycetous, sexual morph unknown, endophytic, saprobes, widespread, see Ding et al. (2010; antimicrobial activity), Hyde et al. (2011; checklist), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), Maharachchikumbura et al. (2015b, 2016b; outline), cultures and sequences are unavailable.

Zyziphora B. Sutton 1981, *Ascomycota* genera *incertae sedis*, one species, type: *Z. appendiculata* (B. Sutton) Nag Raj, hyphomycetous, sexual morph unknown, saprobes, terrestrial, Asia, see Seifert et al. (2011; morphology), Wijayawardene et al. (2012, 2017; outline), Kirk et al. (2013; genus accepted), cultures and sequences are unavailable, needs generic revision.

Doubtful genera

In Table 2, we listed the genera in *Ascomycota* which have been listed as nom. dub. in Kirk et al. 2008 and have not been included in recent literature.

Acknowledgements Nalin Wijayawardene would like to thank Lechat Christian, Yuanpin Xiao, Danushka Sandaruwan, Paul Mungai, Huang Zhang, Ishani Goonasekara, Chada Norphanphon, Ishara Manawasingha, Rajesh Jeewon, Thilini Chethana and Hasini Ekanayaka for their assistances and suggestions. We would like to thank Mark Stadler for his help to check names in *Xylariales*. Nalin Wijayawardene and Pedro Crous thank Ulrike Damm for her comments and suggestions for validating several names. K.D. Hyde thanks The Chinese Academy of Sciences, for the award of Visiting Professorship for Senior International Scientists at Kunming Institute of Botany. K.D. Hyde and Monika C. Dayaratne would like to thank the Thailand Research Fund (TRF) grant no RSA5980068 entitled

Biodiversity, phylogeny and role of fungal endophytes on above parts of *Rhizophora apiculata* and *Nypa fruticans*, National Research Council of Thailand (NRCT) entitled Diseases of mangrove trees and maintenance of good forestry practice (Grant number: 60201000201) and Mae Fah Luang University grant “Biodiversity, phylogeny and role of fungal endophytes of Pandanaceae” (Grant number: 592010200112). Hugo Madrid was funded by Comisión Nacional de Investigación Científica y Tecnológica (CONICYT), Fondo Nacional de Desarrollo Científico y Tecnológico (FONDECYT), Chile, project no. 11140562.” Rafael F. Castañeda-Ruiz is grateful to the Organización Superior de Dirección Empresarial, Grupo Agrícola, (OSDE) from the Cuban Ministry of Agriculture and “Programa de Salud Animal y Vegetal”, project P13ILH003033. Dong Qin Dai would like to thank the Key Laboratory of Yunnan Province Universities of the Diversity and Ecological Adaptive Evolution for Animals and plants on Yun-Gui Plateau for the support. Ka-Lai Pang thanks Ministry of Science and Technology, Taiwan for financial support (105-2621-B-019 -002-). Guo Zhu Zhao was funded by the National Natural Science Foundation of China (No. 31570019). Mingkwan Doilom acknowledges the Royal Golden Jubilee Ph.D. Program (PHD./0072/2553 in 4.S.M.F./53/A.2. K. Tanaka would like to thank the Japan Society for the Promotion of Science (JSPS; 26291084 and 16K07474). Walter P. Pflieger was supported through the ÚNKP-16-4-IV New National Excellence Program of the Hungarian Ministry of Human Capacities. Samantha C. Karunaratna thanks Yunnan Provincial Department of Human Resources and Social Security funded postdoctoral project (number 179122) for supporting his postdoctoral research study. The authors extend their appreciation to the International Scientific Partnership Program ISPP at King Saud University for funding this research work through ISPP#0089. KC Rajeshkumar thanks SERB, DST, Government of India for providing financial support under the project YSS/2015/001590 and Dr. K. M. Paknikar, Director, ARI for providing the facility. Mats Wedin thanks the Swedish Research Council, grants VR 621-2012-3990 and VR 2016-03589. Alan JL Phillips acknowledges the support from Biosystems and Integrative Sciences Institute (BioISI, FCT/UID/ Multi/04046/2013). L. Selbmann, L. Zucconi and S. Onofri thank the Italian National Program for Antarctic Researches (PNRA) for the financial support. The Italian National Antarctic Museum (MNA) is acknowledged for supporting the Mycological Section and the Culture Collection of Fungi from Extreme Environments (CCFEE).

References

- Abbas A, Mamut R, Abdoula A, Ababakri G (2014) A newly recorded lichen genus *Xylographa* (Fr.) Fr. from China. *Plant Divers Resour* 36:578–580
- Abdel-Aziz FA (2016a) The genus *Lolia* from freshwater habitats in Egypt with one new species. *Phytotaxa* 267:279–288
- Abdel-Aziz FA (2016b) Two new cheiroporous asexual taxa (*Dictyosporiaceae*, *Pleosporales*, *Dothideomycetes*) from freshwater habitats in Egypt. *Mycosphere* 7:448–457
- Abdel-Aziz FA, Abdel-Wahab MA (2010) *Lolia aquatica* gen. et sp. nov. (*Lindgomycetaceae*, *Pleosporales*), a new coelomycete from freshwater habitats in Egypt. *Mycotaxon* 114:33–42
- Abd-Elsalam KA, Tibpromma S, Wanasinghe DN, Camporesi E et al (2016) *Equiseticola* gen. nov. (*Phaeosphaeriaceae*), from *Equisetum* sp. in Italy. *Phytotaxa* 284:169–180
- Abdel-Wahab MA (2011a) Lignicolous marine fungi from Yokosuka, Japan. *Bot Mar* 54:209–221
- Abdel-Wahab MA (2011b) Marine fungi from Sarushima Island, Japan, with a phylogenetic evaluation of the genus *Naufragella*. *Mycotaxon* 115:443–456
- Abdel-Wahab MA, Nagahama T (2011) *Gesasha* (*Halosphaeriaceae*, *Ascomycota*), a new genus from the Gesashi mangroves in Japan. *Nova Hedw* 92:497–512
- Abdel-Wahab MA, Bahkali AH (2012) Taxonomy of filamentous anamorphic marine fungi: morphology and molecular evidence. In: Pang KL (ed) *Marine fungi and fungal-like organisms labyrinthomycetes and oomycetes*. De Gruyter series: marine and freshwater botany. Walter De Gruyter GMBH, Germany, pp 59–82
- Abdel-Wahab MA, Nagahama T, Abdel-Aziz FA (2009) Two new *Corollospora* species and one anamorph based on morphological and molecular data. *Mycoscience* 50:147–155
- Abdel-Wahab MA, Pang KL, Nagahama T, Abdel-Aziz FA, Jones EBG (2010) Phylogenetic evaluation of anamorphic species of *Cirrenalia* and *Cumulospora* with the description of eight new genera and four new species. *Mycol Prog* 9:537–558
- Abdel-Wahab MA, Hodhod MS, Bahkali AHA, Jones EBG (2014) Marine fungi of Saudi Arabia. *Bot Mar* 27:323–335
- Abdel-Wahab MA, Bahkali AH, Jones EBG, Elgorban AM et al (2016) Two new species of *Kallichroma* (*Bionectriaceae*, *Hypocreales*) from Saudi Arabian mangroves. *Phytotaxa* 260:66–74
- Abdel-Wahab MA, Dayarathne MC, Suetrong S, Guo SY et al (2017) New saprobic marine fungi and a new combination. *Bot Mar*. doi:10.1515/bot-2016-0118
- Abdollahzadeh J, Goltepah EM, Javadi A, Shams-Bakhsh M et al (2009) *Barriopsis iraniana* and *Phaeobotryon cupressi*: two new species of the *Botryosphaeriaceae* from trees in Iran. *Persoonia* 23:1–8
- Abdollahzadeh J, Javadi A, Goltepah EM, Zare R, Phillips AJ (2010) Phylogeny and morphology of four new species of *Lasiodiplodia* from Iran. *Persoonia* 25:1–10
- Abdollahzadeh J, Zare R, Phillips AJ (2013) Phylogeny and taxonomy of *Botryosphaeria* and *Neofusicoccum* species in Iran, with description of *Botryosphaeria scharifii* sp. nov. *Mycologia* 105:210–220
- Abdollahzadeh J, Javadi A, Zare R, Phillips AJ (2014) A phylogenetic study of *Dothiorella* and *Spencermartinsia* species associated with woody plants in Iran, New Zealand, Portugal and Spain. *Persoonia* 32:1–12
- Abdullah SK, Asensio L, Monfort E, Gomez-Vidal S et al (2015) Occurrence in elx, se spain of inflorescence rot disease of date palms caused by *Mauginiella scaettiae*. *J Phytopathol* 153:417–422
- Abe N, Hamada N (2011) Molecular characterization and surfactant utilization of *Scolecobasidium* isolates from detergent-rich indoor environments. *Biocontrol Sci* 16:139–147
- Acharya K, Dutta AK, Pradhan P (2011) *Bipolaris sorokiniana* (Sacc.) Shoem.: the most destructive wheat fungal pathogen in the warmer areas. *Australian J Crop Sci* 5:1064
- Adamčík S, Cai L, Chakraborty D, Chen XH et al (2015) Fungal biodiversity profiles 1–10. *Cryptogam Mycol* 36:121–166
- Adeleke R, Dames JF (2014) *Kalaharituber pfeilii* and associated bacterial interactions. *South Afr J Bot* 90:68–73
- Aggarwal R, Das S, Jahani M, Singh D (2008) Histopathology of spot blotch disease of wheat caused by *Bipolaris sorokiniana* (Teleomorph: *Cochliobolus sativus*). *Acta Phytopathol et Entomol Hung* 43:23–30
- Aggarwal R, Singh VB, Gurjar MS, Gupta S, Srinivas P (2009) Intraspecific variations in Indian isolates of *Bipolaris sorokiniana* infecting wheat based on morphological, pathogenic and molecular characters. *Indian Phytopathol* 62:449–460
- Aggarwal R, Singh VB, Shukla R, Gurjar MS et al (2010) URP-based DNA fingerprinting of *Bipolaris sorokiniana* isolates causing spot blotch of wheat. *J Phytopathol* 158:210–216

- Aggarwal R, Gupta S, Banerjee S, Singh VB (2011a) Development of a SCAR marker for detection of *Bipolaris sorokiniana* causing spot blotch of wheat. *Can J Microbiol* 57:934–942
- Aggarwal R, Purwar S, Kharbikar L, Gupta S (2011b) Induction of a wheat β-1, 3-glucanase gene during the defense response to *Bipolaris sorokiniana*. *Acta Phytopathol et Entomol Hung* 46:39–47
- Agnello G, Catinon M, Ayrault S, Asta J et al (2010) A comparative study of atmospheric deposits and lichen populations in a protected alpine area in the Grenoble region (France). *Int J Environ Health* 4(2–3):235–249
- Agnihothrudu V (1962) Notes on fungi from North East India X. *Chaetospermum camelliae* sp. nov. on tea (*Camellia sinensis* (L.) O. Kuntze). *Mycopathologia et Mycologia Applicata* 16: 113–116
- Ahmadvour A, Donyadoost-Chelan M, Heidarian Z, Javan-Nikkhah M (2011) New species of *Bipolaris* and *Curvularia* on grass species in Iran. *Rostaniha* 12:39–49
- Ahmadvour A, Heidarian Z, Donyadoost-Chelan M, Javan-Nikkhah M, Tsukiboshi T (2012) A new species of *Bipolaris* from Iran. *Mycotaxon* 120:301–307
- Ahmed S, Imdad SS, Jamil A (2012) Comparative study for the kinetics of extracellular xylanases from *Trichoderma harzianum* and *Chaetomium thermophilum*. *Electron J Biotechnol* 15:3. doi:10.2225/vol15-issue3-fulltext-2
- Ahmed SA, de Sande WWJV, Stevens DA, Fahal A et al (2014a) Revision of agents of black-grain eumycetoma in the order *Pleosporales*. *Persoonia* 33:141–154
- Ahmed SA, Stevens DA, van de Sande WWJ, Meis JF, de Hoog GS (2014b) *Roussoella percutanea*, a novel opportunistic pathogen causing subcutaneous mycoses. *Med Mycol* 52:689–698
- Ahmed SA, Desbois N, Quist D, Miossec C et al (2015) Phaeohyphomycosis caused by a novel species: *Pseudochaetosphaeronema martinelli*. *J Clin Microbiol* 53:2927–2934
- Ahmed SA, Khan Z, Wang XW, Moussa TA et al (2016) Chaetomium-like fungi causing opportunistic infections in humans: a possible role for extremotolerance. *Fungal Divers* 76:11–26
- Ahti T, Stenroos S (2013) *Cladonia*. In: Ahti T, Stenroos S, Moberg R (eds) Nordic Lichen Flora, vol 5. Nordic Lichen Society, Uppsala, pp 8–87
- Ahti TE, Kondratyuk SY, Kärnefelt I, Thell A (2015) Nomenclatural corrections and notes on some taxa in the *Teloschistaceae* (lichenized ascomycetes). *Gr Scr* 27(1–2):37–41
- Ahti T, Mayrhofer H, Schultz M, Tehler A et al (2016) First supplement to the lichen checklist of South Africa: original research. *Bothalia-Afr Biodivers Conserv* 46:1–8
- Akata I, Kaya A (2013) Two earth-tongue genera new for Turkey. *Mycotaxon* 125:87–90
- Akhtar KP, Sarwar N, Saleem K, Ali S (2013) *Gibbago trianthemae* causes *Trianthema portulacastrum* (horse purslane) blight in Pakistan. *Australas Plant Dis Notes* 8(1):109–110
- Al Izquierdo-San, Sierra S, Rodriguez-Gutierrez I, Castro-Santiuste S et al (2013) A new species of *Scutellinia* from Mexico. *Mycotaxon* 125:258
- Alamouti SM, Tsui CK, Breuil C (2009) Multigene phylogeny of filamentous ambrosia fungi associated with ambrosia and bark beetles. *Mycol Res* 113:822–835
- Aleksić M, Stepanović M, Duduk B, Rekanović E (2011) *Septocyta ruborum*: occurrence and possibility of disease control in Serbia. In: X international Rubus and Ribes symposium, vol 946, pp 277–281
- Alexandrova AV, Braun U, Mel'nik VA (2013) *Pyricularia contorta* sp. nov.—a new species from Vietnam. *Schlechtendalia* 25:73–76
- Ali H, Ahmad W, Haq T (2009) Decolorization and degradation of malachite green by *Aspergillus flavus* and *Alternaria solani*. *Afr J Biotechnol* 8
- Alias AS, Jones EBG (2009) Marine fungi from mangrove of Malaysia: IOES monograph series no. 8. Institute of Ocean and Earth Sciences, University of Malaya, Malaysia.
- Alias AS, Zainuddin N, Jones EBG (2010) Biodiversity of marine fungi in Malaysian mangroves. *Bot Mar* 53:545–554
- Al-Laith AA (2010) Antioxidant components and antioxidant/antiradical activities of desert truffle (*Tirmania nivea*) from various Middle Eastern origins. *J Food Compos Anal* 23:15–22
- Allegrucci N, Eliades L, Cabello M, Arambarri A (2011) New species *Koorchaloma* and *Ciliochorella* from xeric forests in Argentina. *Mycotaxon* 115:175–181
- Allen JL, Lendemer JC (2013) Uncovering chemical variability: molecular data reveal the identity of a sterile crustose lichen from the Yukon and affirm an expanded circumscription for *Buellia griseovirens*. *N Am Fungi* 8:1–14
- Allen JL, Lendemer JC (2015) *Japewiella dollypartoniana*, a New Widespread Lichen in the Appalachian Mountains of Eastern North America. *Castanea* 80:59–65
- Allender MC, Raudabaugh DB, Gleason FH, Miller AN (2015) The natural history, ecology, and epidemiology of *Ophidiomyces ophiodiicola* and its potential impact on free-ranging snake populations. *Fungal Ecol* 17:187–196
- Almaguer M, Rojas TI, Dobal V, Batista A, Aira MJ (2013) Effect of temperature on growth and germination of conidia in *Curvularia* and *Bipolaris* species isolated from the air. *Aerobiologia* 29:13–20
- Almeida DAC, Gusmão LFP (2014) *Ypsilomyces*, a new thallic genus of conidial fungi from the semi-arid Caatinga biome of Brazil. *Mycotaxon* 129:181–186
- Almeida DAC, Cruz ACR, Marques MFO, Gusmão LFP (2013) Conidial fungi from the semi-arid Caatinga biome of Brazil. New and interesting *Zanclospora* species. *Mycosphere* 4:684–692
- Almeida DA, Gusmão LF, Miller AN (2014a) Brazilian Semi-Arid Ascomycetes I: New and interesting records of hysteriaceous ascomycetes. *Mycosphere* 5:379–391
- Almeida DAC, Gusmão LFP, Miller AN (2014b) A new genus and three new species of hysteriaceous ascomycetes from the semiarid region of Brazil. *Phytotaxa* 176:298–308
- Almeida DAC, Miller AN, Gusmão LFP (2014c) New species and combinations of conidial fungi from the semi-arid Caatinga biome of Brazil. *Nova Hedw* 98:431–447
- Almeida DA, Santa Izabel TD, Gusmão LF, Castañeda-Ruiz RF (2015) A new species of *Diplococcum* from the Brazilian semi-arid region. *Mycotaxon* 130:495–498
- Almeida DAC, Gusmão LFP, Miller AN (2016) Taxonomy and molecular phylogeny of *Diatripaceae* (Ascomycota, Xylariales) species from the Brazilian semi-arid region, including four new species. *Mycol Prog* 15:1–27
- Almeida DAC, Gusmão LFP, Miller AN (2017) A new species of *Bertiella* (Melanommataceae) from Brazil and a key to accepted species. *Mycosphere* 8:392–8396
- Al-Nasrawi HG, Hughes AR (2012) Fungal diversity associated with salt marsh plants *Spartina alterniflora* and *Juncus roemerianus* in Florida. *Jordan J Biol Sci* 5:247–254
- Alsheikh-Hussain A, Altenaiji E, Yousef LF (2014) Fungal cellulases from mangroves forests- a short review. *J Biochem Technol* 5:765–774
- Alstrup V, Kocourkova J, Kukwa M, Motiejunaite J et al (2009) The lichens and lichenicolous fungi of south Greenland. *Folia Cryptogam Estonica* 46:1–24
- Aluoch AM, Obonyo MA, Okun DO, Akinyi A et al (2015) Morphological diversity of *Ascobolus* and *Pilobolus* fungi from wild herbivore dung in Nairobi National Park, Kenya. *J Microbiol Res* 5:134–141
- Alvarado P, Moreno G, Manjón JL, Gelpí C et al (2011a) First molecular data on *Delastria rosea*, *Fischerula macrospora* and

- Hydnocystis piligera*. Boletín de la Sociedad Micológica de Madrid 35:75–81
- Alvarado P, Moreno G, Manjón JL, Sanz MA (2011b) *Eremiomyces magnisporus* (*Pezizales*), a new species from central Spain. Mycotaxon 118:103–111
- Alvarado P, Cabero J, Moreno G, Bratek Z et al (2014) Species diversity of *Genea* (*Ascomycota*, *Pezizales*) in Europe. Ascomycete.org 6:41–51
- Alvarado P, Cabero J, Moreno G, Bratek Z et al (2016) Phylogenetic overview of the genus *Genea* (*Pezizales*, *Ascomycota*) with an emphasis on European taxa. Mycologia 108(2):441–456
- Álvarez G, Fernández M, Diez JJ (2015) Ophiostomoid fungi associated with declined *Pinus pinaster* stands in Spain. For Syst 24:6
- Alvarez LV, Groenewald JZ, Crous PW (2016) Revising the *Schizopharmaceae*: *Coniella* and its synonyms *Pilidiella* and *Schizopharne*. Stud Mycol 85:1–34
- Alves JL, Barreto RW, Pereira OL, Soares DJ (2010) Additions to the mycoflora of the invasive weed *Miconia calvescens* (*Melastomataceae*). Mycologia 102:69–82
- Alves JL, Woudenberg JHC, Duarte LL, Crous PW, Barreto RW (2013) Reappraisal of the genus *Alternariaster* (*Dothideomycetes*). Persoonia 31:77–85
- Alves A, Linaldeddu BT, Deidda A, Scana B, Phillips AJL (2014) The complex of *Diplodia* species associated with *Fraxinus* and some other woody hosts in Italy and Portugal. Fungal Divers 67:143–156
- Aly AH, Edrada-Ebel R, Indriani ID, Wray V et al (2008) Cytotoxic metabolites from the fungal endophyte *Alternaria* sp. and their subsequent detection in its host plant *Polygonum senegalense*. J Nat Prod 71:972–980
- Amaradasa BS, Amundsen K (2014) First report of *Curvularia inaequalis* and *Bipolaris spicifera* causing leaf blight of buffalograss in Nebraska. Am Phytopathol Soc 98:279
- Ammarellou A, Smith ME, Tajick MA, Trappe JM (2011) The phylogenetic placement of *Picoa*, with a first report on *Picoa lefebvrei* (Pat.) Maire (= *Phaeangium lefebvrei*) from Iran. Int J Environ Res 5:509–514
- Amnuaykanjanasin A, Phonghanpot S, Sengpanich N, Cheevadhanarak S et al (2009) Insect-specific polyketide synthases (PKSs), potential PKS-nonribosomal peptide synthetase hybrids, and novel PKS clades in tropical fungi. Appl Environ Microbiol 75:3721–3732
- An KD, Degawa Y, Fujihara E, Mikawa T et al (2012) Molecular phylogenetic analyses based on the nuclear rRNA genes and the intron–exon structures of the nuSSU rRNA gene in *Dictyocatenulata alba* (anamorphic Ascomycota). Fungal Biol 116:1134–1145
- Anand T, Bhaskaran R, Raguchander T, Samiyappan R et al (2009) Defence responses of chilli fruits to *Colletotrichum capsici* and *Alternaria alternata*. Biol Plant 53:553–559
- Anar M, Aslan A, Agar G, Ozgenclı I (2016) Antigenotoxic and antioxidant activity of lichens *Anaptychia ciliaris*, *Bryoria fuscescens*, *Parmotrema chinensa* and *Xanthoria candelaria*: An *in vitro* study. Med Aromat Plants, 5(233). ISSN: 2167-0412
- Anderson F, Neily TH (2012) New and noteworthy macrolichens records for Nova Scotia. Evansia 29:1–3
- Andersen B, Dongo A, Pryor BM (2008) Secondary metabolite profiling of *Alternaria dauci*, *A. porri*, *A. solani*, and *A. tomatophila*. Mycol Res 112:241–250
- Andersen B, Sørensen JL, Nielsen KF, van den Ende BG, de Hoog S (2009) A polyphasic approach to the taxonomy of the *Alternaria infectioria* species-group. Fung Gen Biol 46:642–656
- Andjic V, Whyte G, Hardy G, Burgess T (2010) New *Teratosphaeria* species occurring on eucalypts in Australia. Fungal Divers 43:27–38
- Andjic V, Maxwell A, Hardy GE, Burgess TI (2016) New cryptic species of *Teratosphaeria* on Eucalyptus in Australia. IMA Fungus 7:253–263
- Ando Y, Masuya H, Motohashi K, Linnakoski R, Yamaoka Y (2016) Phylogenetic relationship of Japanese isolates belonging to the *Grosmannia piceiperda* complex (*Ophiostomatales*). Mycoscience 57:123–135
- Andreev M (2010) *Halecania santessonii*, a new lichenicolous lichen from Russia. Lichenologist 42:249–252
- Andrew M, Peever TL, Pryor BM (2009) An expanded multilocus phylogeny does not resolve morphological species within the small-spored *Alternaria* species complex. Mycologia 101:95–109
- Andrew M, Barua R, Short SM, Kohn LM (2012) Evidence for a common toolbox based on necrotrophy in a fungal lineage spanning necrotrophs, biotrophs, endophytes, host generalists and specialists. PLoS ONE 7:e29943
- Antoniazzi N, Deschamps C, Bach EE (2008) Effect of xanthan gum and allicin as elicitors against *Bipolaris sorokiniana* on barley in field experiments. J Plant Dis Prot 115:104–107
- Aoki N, Moriyama H, Kodama M, Arie T et al (2009) A novel mycovirus associated with four double-stranded RNAs affects host fungal growth in *Alternaria alternata*. Virus Res 140:179–187
- Aptroot A (2008) Lichens of St Helena and Ascension Island. Bot J Linn Soc 158:147–171
- Aptroot A (2012) A world key to the species of *Anthracothecium* and *Pyrenula*. Lichenologist 44:5–53
- Aptroot A (2014) Two new genera of *Arthoniales* from New Caledonia and the Solomon Islands, with the description of eight further species. Bryologist 117:282–289
- Aptroot A (2016) Preliminary checklist of the lichens of Madagascar, with two new thiotremoid *Graphidaceae* and 131 new records. Willdenowia 46:349–365
- Aptroot A, Aslan A (2011) *Lecanora wrightiana* and *Rhizocarpon inimicum*, rare lichens new to Turkey and the Middle East. Mycotaxon 117:145–148
- Aptroot A, Aslan A (2012) *Candelariella*, *Ochrolechia*, *Physcia*, and *Xanthoria* species new to Turkey. Mycotaxon 119:149–156
- Aptroot A, Cáceres MES (2013) Pyrenocarpous lichens (except *Trypetheliaceae*) in Rondônia. Lichenologist 45:763–785
- Aptroot A, Cáceres MES (2014) A refined species concept in the tropical lichen genus *Polymeridium* (*Trypetheliaceae*) doubles the number of known species, with a worldwide key to the species. Nova Hedw 98:1–29
- Aptroot A, Cáceres MES (2014) A key to the corticolous microfoliose, foliose and related crustose lichens from Rondônia, Brazil, with the description of four new species. Lichenologist 46:783–799
- Aptroot A, Iqbal SH (2012) Annotated checklist of the lichens of Pakistan, with reports of new records. Herzogia 25:211–229
- Aptroot A, John V (2015) An historical lichen collection from New Caledonia. Herzogia 28:307–321
- Aptroot A, Lücking R (2016) A revisionary synopsis of the *Trypetheliaceae*. (*Ascomycota*: *Trypetheliales*). Lichenologist 48:763–982
- Aptroot A, Moon KH (2014) 114 new reports of microlichens from Korea, including the description of five new species, show that the microlichen flora is predominantly Eurasian. Herzogia 27:347–365
- Aptroot A, Moon KH (2015) New lichen records from Korea, with the description of the lichenicolous *Halecania parasitica*. Herzogia 28:193–203
- Aptroot A, Nepi C (2017) (2492) Proposal to conserve the name *Marcelaria* against *Buscalionia* (*Trypetheliaceae*, lichenized *Ascomycota*). Taxon 66:200–201

- Aptroot A, Schumm F (2011) Fruticose *Roccellaceae*: An anatomical-microscopical atlas and guide with a worldwide key and further notes on some crustose *Roccellaceae* or similar lichens. BoD—Books on Demand. GmbH, Norderstedt 374 pp
- Aptroot A, Schumm F (2012a) A new terricolous *Trapelia* and a new *Trapeliopsis* (*Trapeliaceae, Baeomycetales*) from Macaronesia. *Lichenologist* 44:449–456
- Aptroot A, Schumm F (2012b) The genus *Melanophloea*, an example of convergent evolution towards polypory. *Lichenologist* 44:501–509
- Aptroot A, Sipman HJ (2011) *Sporodochiolichen*, a new genus of tropical hyphomycetous lichens. *Lichenologist* 43:357–362
- Aptroot A, Yazici K (2012) A new *Placopyrenium* (*Verrucariaceae*) from Turkey. *Lichenologist* 44:739–741
- Aptroot A, Sparrius LB, LaGreca S, Bungartz F (2008) *Angiactis*, a new crustose lichen genus in the family *Roccellaceae* with species from Bermuda, the Galapagos islands, and Australia. *Bryologist* 111:510–516
- Aptroot A, Seaward MR, Sparrius LB (2010) Biodiversity and ecology of lichens. *Bibliotheca Lichenologica* 99:1–439
- Aptroot A, Ertz D, Lima EL, de Jesus KA et al (2013a) Two new species of *Roccellaceae* (*Ascomycota: Arthoniales*) from Brazil, with the description of the new genus *Sergipea*. *Lichenologist* 45:627–634
- Aptroot A, Nelsen MP, Parnmen S (2013b) *Marcelaria*, a new genus for the *Laurera purpurina* group in the *Trypetheliaceae* (*Ascomycota: Dothideomycetes*). *Glalia* 5:1–14
- Aptroot A, Sipman HJM, Cáceres MES (2013c) Twenty-one new species of *Pyrenula* from South America, with notes on overmature ascospores. *Lichenologist* 45:169–198
- Aptroot A, Andrade DS, Mendonça C, de Lima EL, Cáceres MES (2014a) Ten new species of corticolous pyrenocarpous lichens from NE Brazil. *Phytotaxa* 197:197–206
- Aptroot A, de Oliveira MC, Ferraro LI, Cáceres MES (2014b) A world key to species of the genera *Topelia* and *Thelopsis* (*Stictidaceae*), with the description of three new species from Brazil and Argentina. *Lichenologist* 46:801–807
- Aptroot A, Ferraro LI, Cáceres MES (2014c) New pyrenocarpous lichens from NE Argentina. *Lichenologist* 46:95–102
- Aptroot A, Jungbluth P, Cáceres MES (2014d) A world key to the species of *Pyxine* with lichenanthone, with a new species from Brazil. *Lichenologist* 46:669–672
- Aptroot A, Parnmen S, Lücking R, Baloch E et al (2014e) Molecular phylogeny resolves a taxonomic misunderstanding and places *Geisleria* close to *Absconditella* s. str. (*Ostropales: Stictidaceae*). *Lichenologist* 46:115–128
- Aptroot A, Ertz D, Silva JR, Grube M, Cáceres MES (2015a) The phylogenetic position of *Coniarthonia*, with a key to the species and new information about the polyphyly of *Cryptothecia* and *Stirtonia* and the transfer of *Cryptothecia miniata* to *Myriostigma* (*Arthoniaceae*, lichenized ascomycetes). *Phytotaxa* 218:128–136
- Aptroot A, Parnmen S, Lücking R, Baloch E et al (2015b) Molecular phylogeny resolves taxonomic misunderstanding and places *Geisleria* close to *Absconditella* s. str. (*Ostropales: Stictidaceae*). *Lichenologist* 46:115–128
- Aptroot A, Sobreira P, Silva Da, Cáceres MES (2015c) A remarkable new *Ramonia* (*Gyalectaceae*) from Brazil, with a key to the species. *Lichenologist* 47:21–29
- Aptroot A, Yazici K, Košuthová A (2015d) Three *Placynthiaceae* reported new to Asia from limestone in Turkey, with the description of the apothecia of *Placynthium posterulum*. *Herzogia* 28:288–292
- Araújo JP, Evans HC, Geiser DM, Mackay WP, Hughes DP (2015) Unravelling the diversity behind the *Ophiocordyceps unilateralis* (*Ophiocordycipitaceae*) complex: three new species of zombie-ant fungi from the Brazilian Amazon. *Phytotaxa* 220:224–238
- Arauzo S, Iglesias P (2014) La familia *Geoglossaceae* en la península Ibérica y la Macaronesia. *Errötari* 11:166–259
- Arcadia L, Knudsen K (2012) The name *Myriospora* is available for the *Acarospora smaragdula* group. *Opusc Phil* 11:19–25
- Archer AW (2009) *Graphidaceae*. *Fl. Australia* 57:84–194
- Arias MS, Suárez J, Cazapal-Monteiro CF, Francisco I et al (2013) *Trematodes* enhance the development of the nematode-trapping fungus *Arthrobotrys (Duddingtonia) flagrans*. *Fungal Biol* 117:540–544
- Arias RM, Heredia G, Castañeda-Ruiz RF (2016) Two new species of *Bactrodesmium* and *Dictyoaquaphila* from Mexico. *Mycotaxon* 131:291–295
- Ariyawansa HA, Jones EBG, Suetrong S, Alias AS et al (2013a) *Halojellaceae* a new family of the order *Pleosporales*. *Phytotaxa* 130:14–24
- Ariyawansa HA, Kang JC, Alias SA, Chukeatirote E et al (2013b) Towards a natural classification of *Dothideomycetes*: The genera *Dermatodothella*, *Dothideopsella*, *Grandigallia*, *Hysteropeltella* and *Gloeodiscus* (*Dothideomycetes incertae sedis*). *Phytotaxa* 147:35–47
- Ariyawansa HA, Maharachchikumbura SSN, Karunaratne SC, Chukeatirote E et al (2013c) *Deniquelata barringtoniae* gen. et sp. nov., associated with leaf spots of *Barringtonia asiatica*. *Phytotaxa* 105:11–20
- Ariyawansa HA, Camporesi E, Thambugala KM, Mapook A et al (2014a) Confusion surrounding *Didymosphaeria* phylogenetic and morphological evidence suggest *Didymosphaeriaceae* is not a distinct family. *Phytotaxa* 176:102–119
- Ariyawansa HA, Hawksworth DL, Hyde KD, Jones EBG et al (2014b) Epitypification and neotypification: guidelines with appropriate and inappropriate examples. *Fungal Divers* 69:57–91
- Ariyawansa HA, Kang JC, Alias SA, Chukeatirote E, Hyde KD (2014c) *Pyrenophora*. *Mycosphere* 5:351–362
- Ariyawansa HA, Phookamsak R, Tibpromma S, Kang JC, Hyde KD (2014d) A molecular and morphological reassessment of *Diademaceae*. *Sci World J*:1–11
- Ariyawansa HA, Tanaka K, Thambugala KM, Phookamsak R et al (2014e) A molecular phylogenetic reappraisal of the *Didymosphaeriaceae* (*Montagnulaceae*). *Fungal Divers* 68:69–104
- Ariyawansa HA, Thambugala KM, Kang JC, Alias SA et al (2014f) Towards a natural classification of *Dothideomycetes* 2: The genera *Cucurbitothis*, *Heterosphaeropsis*, *Hyalosphaera*, *Navicella* and *Pleiostomellina* (*Dothideomycetes incertae sedis*). *Phytotaxa* 176:7–17
- Ariyawansa HA, Hyde KD, Jayasiri SC, Buyck B et al (2015a) Fungal diversity notes 111–252 taxonomic and phylogenetic contributions to fungal taxa. *Fungal Divers* 75:27–274
- Ariyawansa HA, Phukhamsakda C, Thambugala KM, Bulgakov TS et al (2015b) Revision and phylogeny of *Leptosphaeriaceae*. *Fungal Divers* 74:19–51
- Ariyawansa HA, Thambugala KM, Manamgoda DS, Jayawardena R et al (2015c) Towards a natural classification and backbone tree for *Pleosporaceae*. *Fungal Divers* 71:85–139
- Ariyawansa HA, Hyde KD, Liu JK, Wu SP et al (2016a) Additions to Karst Fungi 1: *Botryosphaeria minutispermatia* sp. nov., from Guizhou Province, China. *Phytotaxa* 275:35–44
- Ariyawansa HA, Hyde KD, Tanaka K, Maharachchikumbura SSN et al (2016b) Additions to Karst Fungi 3: *Prosthemium sinense* sp nov., from Guizhou Province, China. *Phytotaxa* 284:281–291
- Ariyawansa HA, Hyde KD, Thambugala KM, Maharachchikumbura SSN et al (2016c) Additions to Karst Fungi 2: *Alpestrisphaeria jonesii* from Guizhou Province, China. *Phytotaxa* 277:255–265

- Armentano RA, Cooke KL, Wickes BL (2013) Disseminated mycotic infection caused by *Westerdykella* species in a German Shepherd Dog. J Am Vet Med Assoc 242:381–387
- Árnason H (2014) Isolation and structure determination of a heteroglycan from whiteworm lichen (*Thamnolia subuliformis*). PhD Thesis, University of Iceland
- Arslan B, Öztürk S, Oran S (2011) *Lecanora*, *Phaeophyscia* and *Rinodina* species new to Turkey. Mycotaxon 116:49–52
- Arunpanichlert J, Rukachaisirikul V, Tadpatch K, Phongpaichit S et al (2012) A dimeric chromanone and a phthalide: Metabolites from the seagrass-derived fungus *Bipolaris* sp. PSU-ES64. Phytochem Lett 5:604–608
- Arup U, Søchting U, Frödén P (2013) A new taxonomy of the family *Teloschistaceae*. Nordic J Bot 31:16–83
- Arzanlou M, Bakhshi M (2011) *Microcyclospora rumicis*, a new species on *Rumex crispus* from Iran. Mycotaxon 118:181–186
- Arzanlou M, Ghasemi S, Baradarani Bagheri M (2016a) *Collophora hispanica*, a new pathogen and potential threat to the almond industry in Iran. J Phytopathol 164:833–839
- Arzanlou M, Samadi R, Frisvad JC, Houbraken J, Ghosta Y (2016b) Two novel *Aspergillus* species from hypersaline soils of the National Park of Lake Urmia, Iran. Mycol Prog 15:1081–1092
- Asad S, Iftikhar S, Munir A, Ahmad I (2009) Characterization of *Bipolaris sorokiniana* isolated from different agro-ecological zones of wheat production in Pakistan. Pak J Bot 41:301–308
- Asam S, Lichtenegger M, Liu Y, Rychlik M (2012) Content of the *Alternaria* mycotoxin tenuazonic acid in food commodities determined by a stable isotope dilution assay. Mycotoxin Res 28:9–15
- Asgari B, Zare R (2010) Two new species of *Preussia* from Iran. Nova Hedw 90:533–548
- Asgari B, Zare R (2011) A contribution to the taxonomy of the genus *Coniocephala* (*Xylariales*). Mycol Prog 10:189–206
- Asgari B, Zare R, Zamanizadeh HR, Rezaee S (2014) *Aspergillus osmophilus* sp. nov. and a new teleomorph for *A. proliferans*. Mycoscience 55:53–62
- Ashraf T, Hanif M, Khalid AN (2012) *Peziza michelii* and its ectomycorrhizae with *Alnus nitida* (*Betulaceae*) from Pakistan. Mycotaxon 120:181–188
- Aslan A (2013) New *Lecanora*, *Lecidea*, *Melaspilea*, *Placynthium*, and *Verrucaria* records for Turkey and Asia. Mycotaxon 123:321–326
- Atienza V, Pérez-Ortega S, Etayo J (2009) Two new conidial lichenicolous fungi from Spain indicate the distinction of *Lichenodiplis* and *Minutoexcipula*. Lichenologist 41:223–229
- Augustin JO, Elliot SL, Nascimento RJ, Mizubuti ESG, Barreto RW, Evans HC (2012) Novelties in the taxonomy and ecology of mycoparasites associated with nests of *Acromyrmex* leaf-cutting ants. PLoS ONE 8:1–17
- Avenot H, Morgan DP, Michailides TJ (2008) Resistance to pyraclostrobin, boscalid and multiple resistance to Pristine® (pyraclostrobin+ boscalid) fungicide in *Alternaria alternata* causing *Alternaria* late blight of pistachios in California. Plant Pathol 57:135–140
- Avenot H, Sellam A, Michailides T (2009) Characterization of mutations in the membrane-anchored subunits AaSDHC and AaSDHD of succinate dehydrogenase from *Alternaria alternata* isolates conferring field resistance to the fungicide boscalid. Plant Pathol 58:1134–1143
- Aveskamp MM, Verkley GJM, de Gruyter J, Murace MA et al (2009) DNA phylogeny reveals polyphyly of *Phoma* section *Peyronellaea* and multiple taxonomic novelties. Mycologia 101:359–378
- Aveskamp MM, De Gruyter J, Woudenberg JH, Verkley GJ, Crous PW (2010) Highlights of the *Didymellaceae*: a polyphasic approach to characterise *Phoma* and related pleosporalean genera. Stud Mycol 65:1–60
- Awaluddin HH, Nor NAM, Nor HM, Sharuddin SS et al (2015) Biodiversity of marine lignicolous fungi from mangroves of Sulu Sea. Malay J Med Sci 34:43–57
- Aylward J, Dreyer LL, Steenkamp ET, Wingfield MJ, Roets F (2014) Development of polymorphic microsatellite markers for the genetic characterisation of *Knox daviesiae proteae* (*Ascomycota: Microascales*) using ISSR-PCR and pyrosequencing. Mycol Prog 13:439–444
- Aylward J, Dreyer LL, Steenkamp ET, Wingfield MJ, Roets F (2015) *Knox daviesiae proteae* is not the only *Knox daviesiae*-symbiont of *Protea repens*. IMA Fungus 6:471–476
- Azcarate MP, Patriarca A, Terminello L, Pinto VF (2008) *Alternaria* toxins in wheat during the 2004 to 2005 Argentinean harvest. J Food Prot 71:1262–1265
- Azeem M, Terenius O, Rajarao GK, Nagahama K et al (2015) Chemodiversity and biodiversity of fungi associated with the pine weevil *Hylobius abietis*. Fungal Biol 119:738–746
- Aziz FH, Qadir SB (2016) Common and new records of Lichens from Iraqi Kurdistan Region, Iraq. Bull Iraq Nat Hist Mus 14:51–68
- Back CG, Kim YH, Jo WS, Chung H, Jung HY (2010) Cobweb disease on *Agaricus bisporus* caused by *Cladobotryum mycophilum* in Korea. J Gen Plant Pathol 76:232–235
- Badali H, Gueidan C, Najafzadeh MJ, Bonifaz A et al (2008) Biodiversity of the genus *Cladophialophora*. Stud Mycol 61:175–191
- Badali H, Carvalho VO, Vicente V, Attili-Angelis D et al (2009) *Cladophialophora saturnica* sp. nov., a new opportunistic species of *Chaetothyriales* revealed using molecular data. Med Mycol 47:55–66
- Badali H, Prenafeta-Boldú FX, Guarro J, Klaassen CH et al (2011) *Cladophialophora psammophila*, a novel species of *Chaetothyriales* with a potential use in the bioremediation of volatile aromatic hydrocarbons. Fungal Biol 115:1019–1029
- Bagyanarayana G, Rao NK, Kunwar IK (2009) *Manoharachariella*, a new dematiaceous hyphomycetous genus from India. Mycotaxon 109:301–305
- Bai QR, Han S, Xie YY, Dong R, Gao J, Li Y (2012) First report of daylily leaf streak caused by *Kabatiella microsticta* in China. Plant Dis 10:1579
- Bajpai R, Upreti DK (2012) Accumulation and toxic effect of arsenic and other heavy metals in a contaminated area of West Bengal, India, in the lichen *Pyxine cocoës* (Sw.) Nyl. Ecotoxicol Environ Saf 83:63–70
- Bajpai R, Upreti DK, Dwivedi SK, Nayaka S (2009) Lichen as quantitative biomonitor of atmospheric heavy metals deposition in Central India. J Atmos Chem 63:235–246
- Bajpai R, Shukla V, Singh N, Rana TS, Upreti DK (2015) Physiological and genetic effects of chromium (+VI) on toxitolent lichen species, *Pyxine cocoës*. Environ Sci Pollut Res 22:3727–3738
- Bakhshi M, Arzanlou M, Babai-Ahari A (2012) Comprehensive check list of Cercosporoid fungi from Iran. Plant Pathol Quar 2:44–55
- Bakhshi M, Arzanlou M, Babai-Ahari A, Groenewald JZ et al (2014) Multi-gene analysis of *Pseudocercospora* spp. from Iran. Phytotaxa 184(5):245–264
- Bakhshi M, Arzanlou M, Babai-Ahari A, Groenewald JG, Crous PW (2015a) Is morphology in *Cercospora* a reliable reflection of generic affinity? Phytotaxa 213:22–34
- Bakhshi M, Arzanlou M, Babai-Ahari A, Groenewald JZ et al (2015b) Application of the consolidated species concept to *Cercospora* spp. from Iran. Persoonia 34:65

- Baloch E, Gilenstam G, Wedin M (2009) Phylogeny and classification of *Cryptodiscus*, with taxonomic synopsis of the Swedish species. *Fungal Divers* 38:51–68
- Baloch E, Lücking R, Lumbsch HT, Wedin M (2010) Major clades and phylogenetic relationships between lichenized and non-lichenized lineages in *Ostropales* (Ascomycota: *Lecanoromycetes*). *Taxon* 59:1483–1494
- Baloch E, Gilenstam G, Wedin M (2013) The relationships of *Odontotrema* (*Odontotremataceae*) and the resurrected *Sphaeropezia* (*Stictidaceae*)—new combinations and three new *Sphaeropezia* species. *Mycologia* 105:384–397
- Ban S, Sakane T, Nakagiri A (2015) Three new species of *Ophiocordyceps* and overview of anamorph types in the genus and the family *Ophiocordyceptaceae*. *Mycol Prog* 14:1017
- Bansal H, Srivastava S, Chaurasia A, Jabalia NA (2014) Comparative Study of Antifreeze Proteins from *Antarctomyces psychrotrophicus* and *Typhula ishikariensis* using computational tools and servers. *VIVECHAN Int J Res* 5:21–28
- Baral HO (2015) Nomenclatural novelties. *Index Fungorum*, No. 255, pp 1–3
- Baral HO, Kutorga E (2010) *Helicogonium fusisporum* sp. nov., an intrahymenial parasite in *Orbilia eucalypti*. *Mycotaxon* 113:331–336
- Baral HO, Marson G (2012) *Deltopyxis triangulispora* gen. et sp. nov., a polysporous Tromeropsis-like discomycete of unclear relationships. *Andrias* 19:176
- Baral HO, Haelewaters D (2015) *Rommelaarsia flavovirens* gen. et sp. nov. (*Helotiales*), a new discomycete on *Equisetum* with a peculiar asexual state. *Ascomycete.org* 7:321–330
- Baral HO, Rämä T (2015) Morphological update on *Calycina marina* (*Pezizellaceae*, *Helotiales*, *Leotiomycetes*), a new combination for *Laetinaevia marina*. *Bot Mar* 58:523–534
- Baral HO, De Sloover JR, Huhtinen S, Laukka T, Stenroos S (2009) An emendation of the genus *Hyaloscypha* to include *Fuscoscypha* (*Hyaloscypheaceae*, *Helotiales*, *Ascomycotina*). *Karstenia* 49:1–17
- Baral HO, Garcia G, Bogale M, O’Hara MJ, Untereiner WA (2012) *Colipila*, a new genus in the Helotiales. *Mycol Prog* 11:201–214
- Baral HO, Galán R, Platas G, Tena R (2013a) *Phaeohelotium undulatum* comb. nov. and *Phaeoh. succineoguttulatum* sp. nov., two segregates of the *Discinella terrestris* aggregate found under Eucalyptus in Spain: taxonomy, molecular biology, ecology and distribution. *Mycosistema* 32:386–428
- Baral HO, Marson G, Bogale M, Untereiner WA (2013b) *Xerombrophila crystallifera*, a new genus and species in the Helotiales. *Mycol Prog* 12:475–488
- Baral HO, Queloz VK, Hosoya TS (2014) *Hymenoscyphus fraxineus*, the correct scientific name for the fungus causing ash dieback in Europe. *IMA Fungus* 5:79–80
- Baral HO, Weber E, Gams W, Stadler M et al (2015) Recommendations about Genera to be protected or suppressed in the *Orbiliaceae* (*Orbiliomycetes*). *Mycol Prog*. doi:[10.1007/s11557-017-1300-6](https://doi.org/10.1007/s11557-017-1300-6)
- Barbeito I, Brücker RL, Rixen C, Bebi P (2013) Snow fungi—induced mortality of *Pinus cembra* at the alpine treeline: evidence from plantations. *Arct Antarct Alp Res* 45:455–470
- Barber PA, Crous PW, Groenewald JZ, Pascoe IG, Keane P (2011) Reassessing *Vermisporium* (*Amphisphaeriaceae*), a genus of foliar pathogens of *Eucalypts*. *Persoonia* 27:90–118
- Barbosa FR, Gusmão LFP (2011) Conidial fungi from semi-arid Caatinga Biome of Brazil. Rare freshwater hyphomycetes and other new records. *Mycosphere* 2:475–485
- Barbosa FR, Gusmão LF, Raja HA, Shearer CA (2013a) New species and new records of freshwater ascomycetes from Brazil and Costa Rica. *Mycologia* 105:335–343
- Barbosa FR, Raja HA, Shearer CA, Gusmão LFP (2013b) Some freshwater fungi from the Brazilian semi-arid region, including two new species of hyphomycetes. *Cryptog Mycol* 34:243–258
- Barbosa MA, Costa PM, Malosso E, Castañeda-Ruiz RF (2016) *Codinaea leomaiae* sp. nov. from the Brazilian Atlantic Forest. *Mycotaxon* 131:423–428
- Barcenas Peña A, Lücking R, Miranda-González R, Herrera-Campos MA (2014) Three new species of *Graphis* (Ascomycota: *Ostropales*: *Graphidaceae*) from Mexico, with updates to taxonomic key entries for 41 species described between 2009 and 2013. *Lichenologist* 46:69–82
- Barnes I, Van der Nest A, Mullett MS, Crous PW et al (2016) Neotypification of *Dothistroma septosporum* and epitypification of *D. pini*, causal agents of *Dothistroma* needle blight of pine. *For Pathol* 46:388–407
- Barreto RW, Johnston PR, Crous PW, Evans HC (2012) A new species of the lenticel fungal genus *Claviradulomyces* (*Ostropales*) from the Brazilian Atlantic forest tree *Xylopia sericea* (*Annonaceae*). *IMA Fungus* 3:135–141
- Barseghyan GS, Solomon P (2011) The genus *Peziza* Dill. ex Fr. (*Pezizales*, *Ascomycota*) in Israel. *Ascomycete.org* 2(4):39–50
- Bartkowiak ME, Bennett JP (2016) Floristic study of lichens in Portage County, Wisconsin. *Evansia*. doi:[10.1639/0747-9859-32.4.176](https://doi.org/10.1639/0747-9859-32.4.176)
- Bartosik M, Źakowska Z, Cedzińska K, Rožniakowski K (2010) Biodeterioration of optical glass induced by lubricants used in optical instruments technology. *Polskie Towarzystwo Mikrobiologów Pol Soc Microbiol* 59:295–300
- Bartra J, Belmonte J, Torres-Rodriguez JM, Cistero-Bahima A (2008) Sensitization to *Alternaria* in patients with respiratory allergy. *Front Biosci* 14:3372–3379
- Barua BS, Suzuki A, Pham HN, Inatomi S (2012) Adaptation of ammonia fungi to urea enrichment environment. *J Agric Technol* 8:173–189
- Baschien C, Tsui CKM, Gulis V, Szewzyk U, Marvanová L (2013) The molecular phylogeny of aquatic hyphomycetes with affinity to the *Leotiomycetes*. *Fungal Biol* 117:660–672
- Bashir G, Hussain W, Rizvi A (2009) *Bipolaris hawaiiensis* keratomycosis and endophthalmitis. *Mycopathologia* 167:51–53
- Bashya LM, Chand R, Kushwaha C, Sen D et al (2010) Association of melanin content with conidiogenesis in *Bipolaris sorokiniana* of barley (*Hordeum vulgare* L.). *World J Microbiol Biotechnol* 26:309–316
- Bashya LB, Chand R, Prasad L, Joshi A (2011) Partial resistance components for the management of spot blotch pathogen *Bipolaris sorokiniana* of barley (*Hordeum vulgare* L.). *Acta Phytopathol et Entomol Hung* 46:49–57
- Bastida F, Jehmlich N, Ondoño S, von Bergen M et al (2014) Characterization of the microbial community in biological soil crusts dominated by *Fulgensia desertorum* (Tomin) Poelt and *Squamaria cartilaginea* (With.) P. James and in the underlying soil. *Soil Biol Biochem* 76:70–79
- Bates ST, Bungartz F, Lücking R, Herrera-Campos MA, Zambrano A (2011) Biomonitoring, ecology and systematics of lichens. *Acta Bot Hung* 53:442–443
- Baturo-Ciesniewska A (2011) Genetic variability and pathogenicity among polish isolates of *Bipolaris sorokiniana* from spring barley. *J Plant Pathol* 93:291–302
- Bautista-Rosales PU, Calderon-Santoyo M, Servín-Villegas R, Ochoa-Álvarez NA, Ragazzo-Sánchez JA (2013) Action mechanisms of the yeast *Meyerozyma caribbica* for the control of the phytopathogen *Colletotrichum gloeosporioides* in mangoes. *Biol Control* 65:293–301
- Beaulieu WT, Panaccione DG, Ryan KL, Kaonongbua W, Clay K (2015) Phylogenetic and chemotypic diversity of *Periglandula*

- species in eight new morning glory hosts (*Convolvulaceae*). *Mycologia* 107:667–678
- Beck A, Mayr C (2012) Nitrogen and carbon isotope variability in the green-algal lichen *Xanthoria parietina* and their implications on mycobiont–photobiont interactions. *Ecol Evol* 2:3132–3144
- Beese WJ, Sandford J, Symon M, Major S (2015) Distribution and abundance of *Pseudocyphellaria rainierensis* (old-growth specklebelly lichen) on Vancouver Island and a portion of the Central Mainland Coast of British Columbia. *Evansia* 32:136–153
- Begoude ADB, Gryzenhout M, Wingfield MJ, Roux J (2010) *Aurifilum*, a new fungal genus in the *Cryphonectriaceae* from *Terminalia* species in Cameroon. *Antonie Van Leeuwenhoek* 98:263–278
- Begum A, HariKrishna S (2010) Monitoring Air pollution using lichens species in South Bangalore, Karnataka. *Int J ChemTech Res* 2:255–260
- Bell A (2010) *Echinospaeria cincinnati*, a new species from Rimutaka Forest park near Wellington New Zealand. *Austral Mycol* 29:24–26
- Bell A, Mahoney DP (2008) *Echinospaeria medusa*, a new species from New Zealand, with notes on related species. *Austral Mycol* 27:141–148
- Bell A, Mahoney DP (2016) Field studies in the Lasiosphaeriaceae and *Helminthosphaeriaceae sensu lato*. *CBS Biodivers Ser* 15:3–124
- Benatti MN (2011a) A simple clearing technique to aid in the recognition of cilia and rhizinae structure in the *Parmeliaceae*. *Opusc Phil* 9:21–25
- Benatti MN (2013) A review of the genus *Bulbothrix* Hale: the isidiate, sorediate, and pustulate species with medullary salazinic acid. *Mycosphere* 4:1–30
- Benatti MN (2014) An update on the genus *Parmelinella* Elix & Hale (Parmeliaceae, lichenized ascomycetes). *Mycosphere* 5:770–789
- Bendiksby M, Timdal E (2013) Molecular phylogenetics and taxonomy of *Hypocenomyces sensu lato* (Ascomycota: Lecanoromycetes): extreme polyphyly and morphological/ecological convergence. *Taxon* 62:940–956
- Bencsik O, Papp T, Berta M, Zana A et al (2014) Ophiobolin A from *Bipolaris oryzae* perturbs motility and membrane integrities of porcine sperm and induces cell death on mammalian somatic cell lines. *Toxins* 6:2857–2871
- Bendiksby M, Haugan R, Spribile T, Timdal E (2015) Molecular phylogenetics and taxonomy of the *Calvitimela aglaea* complex (Tephromelataceae, Lecanorales). *Mycologia* 107:1172–1183
- Bendiksby M, Mazzoni S, Jørgensen MH, Halvorsen R, Holien H (2014) Combining genetic analyses of archived specimens with distribution modelling to explain the anomalous distribution of the rare lichen *Staurolemma omphalariooides*: long-distance dispersal or vicariance? *J Biogeogr* 41:2020–2031
- Benesperi R, Lastrucci L, Nascimbene J (2013) Human disturbance threats the red-listed macrolichen *Seirophora villosa* (Ach.) Frödén in coastal Juniperus habitats: Evidence from Western Peninsular Italy. *Environ Manag* 52:939–945
- Benkert D (2010) Seltene und kritische *Pezizales*-Funde (Ascomycota) aus der Bundesrepublik Deutschland. *Zeitschrift für Mykologie* 76:27–58
- Bennett JP (2013) The lichens and allied fungi of Great Smoky Mountains National Park. *Bryologist* 116:218–220
- Bensch K, Braun U, Groenewald JZ, Crous PW (2012) The genus *Cladosporium*. *Stud Mycol* 72:1–401
- Bensch K, Groenewald JZ, Braun U, Dijksterhuis J et al (2015) Common but different: the expanding realm of *Cladosporium*. *Stud Mycol* 82:23–74
- Benucci GM, Raggi L, Albertini E, Grebenc T et al (2011) Ectomycorrhizal communities in a productive *Tuber aestivum* Vittad. orchard: composition, host influence and species replacement. *FEMS Microbiol Ecol* 76:170–184
- Benucci GMN, Lefevre C, Bonit G (2016) Characterizing root-associated fungal communities and soils of Douglas-fir (*Pseudotsuga menziesii*) stands that naturally produce Oregon white truffles (*Tuber oregonense* and *Tuber gibbosum*). *Mycorrhiza* 26:367–376
- Ben-Ze'ev IS, Assouline I, Levy E, Elkind G (2011) First report of *Bipolaris caktivora* causing fruit blotch and stem rot of dragon fruit (*Pitaya*) in Israel. *Phytoparasitica* 39:195–197
- Berger F, LaGreca S (2014) Contributions to the lichen flora of bermuda-part 1. *Evansia* 31:41–68
- Berger F, Braun U, Heuchert B (2015) *Gonatophragmium lichenophilum* sp. nov.—a new lichenicolous hyphomycete from Austria. *Mycobiotika* 5:7–13
- Berka RM, Grigoriev IV, Ottilar R, Salamov A et al (2011) Comparative genomic analysis of the thermophilic biomass-degrading fungi *Myceliophthora thermophila* and *Thielavia terrestris*. *Nat Biotechnol* 10:922–927
- Berkeley MJ (1876) Notices of North American fungi. *Grevillea* 4:105–106
- Bernardi M, Barragán A, Rossi W (2014) New records of *Laboulbeniales* (Fungi: Ascomycota) from Ecuador and other countries. *Webbia* 69:281–289
- Besitulo A, Moslem MA, Hyde KD (2010) Occurrence and distribution of fungi in a mangrove forest on Siargao Island, Philippines. *Bot Mar* 53:535–543
- Bevan J, Alexander R, Lazaro-Suau R (2012) Transplant experiments to examine the habitat exclusivity of lichen dominated soil crust communities in the El Cautivo badlands, SE Spain. In: EGU General Assembly Conference Abstracts 14:11040
- Bezerra JD, Oliveira RJ, Paiva LM, Silva GA et al (2017a) *Bezerromycetales* and *Wiesneriomycetales* ord. nov. (class Dothideomycetes), with two novel genera to accommodate endophytic fungi from Brazilian cactus. *Mycol Prog* 4:297–309
- Bezerra JDP, Sandoval-Denis M, Paiva LM, Silva GA et al (2017b) New endophytic *Toxicocladosporium* species from cacti in Brazil, and description of *Neocladosporium* gen. nov. *IMA Fungus* 8:77–97
- Bhattarai HD, Kim T, Oh H, Yim JH (2013) A new pseudodepsidone from the Antarctic lichen *Stereocalon alpinum* and its antioxidant, antibacterial activity. *J Antibiot* 66:559
- Bhise MR, Patil CR, Salunkhe CB (2014) Three new black mildew fungi from Mahabaleshwar, Maharashtra, India. *Int J Life Sci* 2:304–310
- Biazrov LG (2012) Stable nitrogen isotopes (d15N) in thalli of arid vagrant lichen *Xanthoparmelia camtschadalensis* across an altitudinal gradient of the Khangai Plateau of Mongolia. *Contemp Probl Ecol* 5:200–207
- Bills GF, Menéndez VG, Platas G (2012) *Kabatiella bupleuri* sp. nov. (Dothideales), a pleomorphic epiphyte and endophyte of the Mediterranean plant *Bupleurum gibraltarium* (Apiaceae). *Mycologia* 104:962–973
- Bilovitz PO (2014) The importance of old deciduous trees and wooden fences for lichen diversity—an example from the Teichalm area (Eastern Alps). *Herzogia* 27:199
- Bilovitz PO, Mayrhofer H (2014) Lichenized and lichenicolous fungi from the valley 'Ochsental' (Eastern Alps, Vorarlberg, Austria). *Fritschiana: Veröffentlichungen aus dem Herbarium des Instituts für Botanik der Karl-Franzens-Universität Graz (GZU)* 78:47
- Bilovitz PO, Stešević D, Mayrhofer H (2010) Epiphytic lichens and lichenicolous fungi from the northern part of Montenegro. *Herzogia* 23:249
- Björk CR (2010) Additions to the Lichen Flora of Washington State, United States.I. *Evansia* 27:18–20

- Boehm EWA, Mugambi G, Miller AN, Huhndorf S et al (2009a) A molecular phylogenetic reappraisal of the *Hysteriaceae*, *Mytilinidiaceae* and *Gloniaceae* (*Pleosporomycetidae*, *Dothideomycetes*) with keys to world species. *Stud Mycol* 64:49–83
- Boehm EWA, Schoch CL, Spatafora JW (2009b) On the evolution of the *Hysteriaceae* and *Mytilinidiaceae* (*Pleosporomycetidae*, *Dothideomycetes*, *Ascomycota*) using four nuclear genes. *Mycol Res* 113:461–479
- Boehm EWA, Marson G, Mathiassen GH, Schoch CL (2015) An overview of the genus *Glyphium* and its phylogenetic placement in *Patellariales*. *Mycologia* 107:607–618
- Bogdanović G, Trajković S, Micić T (2016) Flora lichen western mountains Vranje place. *J Process Manag New Technol* 4:69
- Boitano S, Flynn AN, Sherwood CL, Schulz SM, Hoffman J, Gruzinova I, Daines MO (2011) *Alternaria alternata* serine proteases induce lung inflammation and airway epithelial cell activation via PAR2. *Am J Physiol Lung Cell Mol Physiol* 300:L605–L614
- Boluda CG, Hawksworth DL, Divakar PK, Crespo A, Rico VJ (2016) Microchemical and molecular investigations reveal *Pseudephebe* species as cryptic with an environmentally modified morphology. *Lichenologist* 48:527–543
- Bomble FW (2012) *Candelaria pacifica* and *Xanthomendoza borealis* in the region of Aachen, North Rhine-Westphalia—new to Germany. *Veröff. Bochumer Bot Ver* 4:1–8
- Bonito GM, Gryganskyi AP, Trappe JM, Vilgalys R (2010a) A global meta-analysis of *Tuber* ITS rDNA sequences: species diversity, host associations and long-distance dispersal. *Mol Ecol* 19:4994–5008
- Bonito G, Trappe JM, Rawlinson P, Vilgalys R (2010b) Improved resolution of major clades within *Tuber* and taxonomy of species within the *Tuber gibbosum* complex. *Mycologia* 102:1042–1057
- Bonito G, Smith ME, Nowak M, Healy RA et al (2013) Historical biogeography and diversification of truffles in the *Tuberaceae* and their newly identified southern hemisphere sister lineage. *PLoS ONE* 8:e52765
- Boom PVD, Ertz D (2012) Lichens and lichenicolous fungi from El Hierro (Canary Islands), a survey, including five new species. *Cryptog Mycol* 33:59–97
- Boonmee S, Zhang Y, Chomnunti P, Chukeatirote E et al (2011) Revision of lignicolous *Tubeufiaceae* based on morphological reexamination and phylogenetic analysis. *Fungal Divers* 51:63–102
- Boonmee S, Ko Ko TW, Chukeatirote E, Hyde KD et al (2012) Two new *Kirschsteiniothelia* species with *Dendryphiopsis* anamorphs cluster in *Kirschsteiniotheliaceae* fam. nov. *Mycologia* 104:698–714
- Boonmee S, Bhat JD, Maharachchikumbura SS, Hyde KD (2014a) *Clavatispora thailandica* gen. et sp. nov., a novel taxon of *Venturiiales* (*Dothideomycetes*) from Thailand. *Phytotaxa* 176:92–101
- Boonmee S, Rossman AY, Liu JK, Li WJ et al (2014b) *Tubeufiales*, ord. nov., integrating sexual and asexual generic names. *Fungal Divers* 68:239–298
- Boonmee S, D'souza MJ, Luo Z, Pinruan U et al (2016a) *Dictyosporiaceae* fam. nov. *Fungal Divers* 80:457–482
- Boonmee S, Ekanayaka AH, Hyde KD, Hüseyin E et al (2016b) Phylogenetic placement of *Neorhamphoria garethjonesii* gen. et sp. nov. (*Tubeufiales*, genus, *incertae sedis*). *Mycosphere* 7:1447–1457
- Boonyuen N, Chuaseeharonnachai C, Suetrong S, Veera VS et al (2011) *Savoryellales* (*Hypocreomycetidae*, *Sordariomycetes*): a novel lineage of aquatic ascomycetes inferred from multiple-gene phylogenies of the genera *Ascotaiwania*, *Ascothailandia*, and *Savoryella*. *Mycologia* 103:1351–1371
- Boonyuen N, Sri-Indrasutdh V, Suetrong S, Sivichai S, Jones EBG (2012) *Annulatascus aquatorba* sp. nov., a lignicolous freshwater ascomycete from Sirindhorn Peat Swamp Forest, Narathiwat, Thailand. *Mycologia* 104:746–757
- Boonyuen N, Chuaseeharonnachai C, Suetrong S, Sujinda S et al (2016) *Parafuscosporella garethii* sp. nov. (*Fuscosporellales*) from a community northern forest, in Thailand. *Mycosphere* 7:1265–1272
- Bordallo JJ, Rodriguez AN, Honrubia MA, Morte AS (2012) *Terfezia canariensis* sp. nov. una nueva especie de trufa encontrada en las Islas Canarias. *Cantarella* 56:1–8
- Bordallo JJ, Rodríguez A, Muñoz-Mohedano JM, Suz LM et al (2013) Five new *Terfezia* species from the Iberian Peninsula. *Mycotaxon* 124:189–208
- Bordallo JJ, Rodríguez A, Kounas V, Camello F et al (2015) Two new *Terfezia* species from Southern Europe. *Phytotaxa* 230:239–249
- Borse BD, Borse KN, Pawar NS, Tuwar AR (2013) Marine fungi from India-XII. A revised checklist. *Ind J Geo-Marine Sci* 42:110–119
- Borse BD, Jagdale PE, Patil SY (2014) Freshwater higher fungi from Pune district (MS, India) I: the mitosporic genus *Xylomyces*. *BIOINFOLET-A Quart J Life Sci* 11(3a):730–734
- Bose T, Reynolds DR, Berbee ML (2014) Common, unsightly and until now undescribed: *Fumiglobus pieridicola* sp. nov., a sooty mold infesting *Pieris japonica* from western North America. *Mycologia* 106:746–756
- Boutabia L, Telailia S, de Belair G (2015) Corticolous lichen flora on *Quercus suber* L. in the wetlands of El Kala national park (North-Eastern Algeria). *Adv Environ Biol* 9:360–373
- Boyce RD, Deziel PJ, Otley CC, Wilhelm MP et al (2010) *Phaeohyphomycosis* due to *Alternaria* species in transplant recipients. *Transpl Infect Dis* 12:242–250
- Brackel WV (2008) *Zwackhiomyces echinulatus* sp. nov. and other lichenicolous fungi from Sicily, Italy. *Herzogia* 21:181–198
- Brackel WV (2010a) *Dacampia cyrtellae*, a new name for *Dacampia lecaniae* Brackel. *Herzogia* 23:315–316
- Brackel WV (2010b) Some lichenicolous fungi and lichens from Iceland, including *Lichenopeltella uncialicola* sp. nov. *Herzogia* 23:93–109
- Brackel WV (2011a) Lichenicolous fungi and lichens from Puglia and Basilicata (southern Italy). *Herzogia* 24:65–101
- Brackel WV (2011b) *Lichenopeltella rangiferinae* sp. nov. and some other lichenicolous fungi from Iceland. *Acta Bot Islandica* 15:51–60
- Brackel WV (2013) Miscellaneous records of lichenicolous fungi from the Italian Alps. *Herzogia* 26:141–157
- Brackel WV (2014) Kommentierter Katalog der flechtenbewohnenden Pilze Bayerns. *Biblioth Lichenol* 109:1–476
- Brackel WV, Berger F (2010) Gall-inducing species of *Polycoccum* (*Ascomycota*) on the lichen genus *Placopsis*. *Herzogia* 23:195–204
- Brackel WV, Zimmermann DG (2012) A new species of *Nectriopsis* on *Cladonia* from Nordrhein-Westfalen (Germany). *Herzogia* 25:231–234
- Bradwell T (2010) Studies on the growth of *Rhizocarpon geographicum* in NW Scotland, and some implications for lichenometry. *Geogr Annal Series A Phys Geogr* 92:41–52
- Braga FR, Carvalho RO, Araújo JM, Silva AR et al (2009) Predatory activity of the fungi *Duddingtonia flagrans*, *Monacrosporium thaumasium*, *Monacrosporium sinense* and *Arthrobotrys robusta* on *Angiostrongylus vasorum* first-stage larvae. *J Helminthol* 83:303–308
- Braga FR, Silva AR, Carvalho RO, Araújo JV et al (2010) *In vitro* predatory activity of the fungi *Duddingtonia flagrans*, *Monacrosporium thaumasium*, *Monacrosporium sinense* and

- Arthrobotrys robusta* on *Ancylostoma ceylanicum* third-stage larvae. *Vet Microbiol* 146:183–186
- Braga FR, Carvalho RO, Silva AR, Araújo JV et al (2014) Predatory capability of the nematophagous fungus *Arthrobotrys robusta* preserved in silica gel on infecting larvae of *Haemonchus contortus*. *Trop Anim Health Prod* 46:571–574
- Braganca H, Rigling D, Diogo E, Capelo J et al (2011) *Cryphonectria nateriae*: a new species in the *Cryphonectria-Endothia* complex and diagnostic molecular markers based on microsatellite-primed PCR. *Fungal Biol* 115:852–861
- Brahamanage RS, Lu YZ, Bhat DJ, Wanasinghe DN (2017) Phylogenetic investigations on freshwater fungi in *Tubeufiaceae* (*Tubeufiales*) reveals the new genus *Dictyospora* and new species *Chlamydotubeufia aquatica* and *Helicosporium flavum*. *Mycosphere* 8:917–933
- Braun U (1995) A monograph of *Cercosporella*, *Ramularia* and allied genera (phytopathogenic hyphomycetes), vol 1. IHW-Verlag, Eching, p 333
- Braun U (1998) A monograph of *Cercosporella*, *Ramularia* and allied genera (phytopathogenic hyphomycetes), vol 2. IHW-Verlag, Eching, p 493
- Braun U (2009) New species, combinations and records of hyphomycetes. *Schlechtendalia* 19:63–71
- Braun U (2013a) 2210–2232) Proposals to conserve the teleomorph-typified name *Blumeria* against the anamorph-typified name *Oidium* and twenty-two teleomorph-typified powdery mildew species names against competing anamorph-typified names (*Ascomycota: Erysiphaceae*). *Taxon* 62:1262–1331
- Braun U (2013b) A monograph of *Cercosporella*, *Ramularia* and allied genera (phytopathogenic hyphomycetes). Additions to host range and distribution. *Schlechtendalia* 1:41–43
- Braun U (2016) *Rhynchosporium graminicola* revisited and reinstated. *Schlechtendalia* 30:39–40
- Braun U, Cook RTA (2012) Taxonomic Manual of the *Erysiphales* (Powdery Mildews). CBS Biodivers Series 11:1–707
- Braun U, Crous P (2016) (2415) Proposal to conserve the name *Cercospora* (*Ascomycota: Mycosphaerellaceae*) with a conserved type. *Taxon* 65:185–185
- Braun U, Heuchert B (2010) *Sporidesmiella lichenophila* sp. nov. a new lichenicolous hyphomycete. *Herzogia* 23:69–74
- Braun U, Urtiaga R (2013) New species and new records of cercosporoid hyphomycetes from Cuba and Venezuela (Part 2). *Mycosphere* 4:176–216
- Braun U, Crous PW, Schubert K (2008) Taxonomic revision of the genus *Cladosporium* s. lat.8. Reintroduction of *Graphiopsis* (= *Dichocladosporium*) with further reassessments of cladosporoid hyphomycetes. *Mycotaxon* 103:207–216
- Braun U, Heuchert B, Diederich P (2009) Two new and another interesting lichenicolous hyphomycete. *Herzogia* 22:165–171
- Braun U, Crous PW, Groenewald JZ, Scheuer C (2011) *Pseudovirgaria*, a fungicolous hyphomycete genus. *IMA Fungus* 2:65–69
- Braun U, Nakashima C, Crous PW (2013) Cercosporoid fungi (*Mycosphaerellaceae*) 1. Species on other fungi, *Pteridophyta* and *Gymnospermae*. *IMA Fungus* 4:265–345
- Braun U, Bien S, Hantsch L, Heuchert B (2014a) *Tubakia chinensis* sp. nov. and a key to the species of the genus *Tubakia*. *Schlechtendalia* 28:23–28
- Braun U, Crous PW, Nakashima C (2014b) cercosporoid fungi (*Mycosphaerellaceae*) 2. Species on monocots (*Acoraceae* to *Xyridaceae*, excluding *Poaceae*). *IMA Fungus* 5:203–390
- Braun U, Crous PW, Nakashima C (2015) cercosporoid fungi (*Mycosphaerellaceae*) 4. Species on dicots (*Acanthaceae* to *Amaranthaceae*). *IMA Fungus* 6:373–469
- Breuss O (2009) A synopsis of the lichen genus *Placopyrenium* (*Verrucariaceae*), with descriptions of new taxa and a key to all species. *Biblioth Lichenol* 99:93–112
- Breuss O (2011) Notable records of pyrenocarpous lichens (*Verrucariaceae*) from Spain. *Herzogia* 24:375–378
- Breuss O (2016) *Involucropyrenium romeum* is the correct name for *I. squamulosum* (lichenised *Ascomycota*, *Verrucariaceae*). *Herzogia* 29:196–197
- Brock PM, Döring H, Bidartondo MI (2009) How to know the unknown fungi: the role of a herbarium. *New Phytol* 181:719–724
- Brodo IM (2010) The lichens and lichenicolous fungi of Haida Gwaii (Queen Charlotte Islands), British Columbia, Canada. 5. A new species of *Lecanora* from shoreline rocks. *Botany* 88:352–358
- Brodo IM, Lendemer JC (2015) A revision of the saxicolous, esorediate species of *Ainoa* and *Trapelia* (*Baeomycetaceae* and *Trapeliaceae*, lichenized *Ascomycota*) in North America, with the description of two new species. *Bryologist* 118:385–399
- Brodo IM, Culberson WL, Culberson CF (2008) *Haematomma* (*Lecanoraceae*) in North and Central America, including the West Indies. *Bryologist* 111:363–423
- Bronson AW, Klymiuk AA, Stockey RA, Tomescu AM (2013) A perithecial sordariomycete (*Ascomycota, Diaporthales*) from the lower cretaceous of Vancouver Island, British Columbia, Canada. *Int J Plant Sci* 174(3):278–292
- Brown EM, McTaggart LR, Zhang SX, Low DE et al (2013) Phylogenetic analysis reveals a cryptic species *Blastomyces gilchristii*, sp. nov. within the human pathogenic species *Blastomyces dermatitidis*. *PLoS ONE* 8:e59237
- Bruckart WL, Eskandari FM, Berner DK (2014) Characterization and evaluation of *Ramularia crupinae*, a candidate for biological control, and of its host, two varieties of *Crupina vulgaris* in the United States. *Biol Control* 71:40–48
- Bruzone MC, Fehrer J, Fontenla SB, Vohník M (2016) First record of *Rhizoscyphus ericae* in Southern Hemisphere's *Ericaceae*. *Mycorrhiza* 25:1–7
- Buaruang K, Scharnagl K, Divakar PK, Leavitt SD et al (2015) Molecular data support *Pseudoparmelia* as a distinct lineage related to *Relicina* and *Relicinopsis* (*Ascomycota, Lecanorales*). *Lichenologist* 47:43–49
- Buğukoglu TZ, Albayrak S, Halıcı MG, Tay T (2013) Antimicrobial and antioxidant activities of extracts and lichen acids obtained from some *Umbilicaria* species from Central Anatolia, Turkey. *J Food Process Preserv* 37:1103–1110
- Büdel B (2011) Lichens of the Namib Desert-A guide to their identification. *Herzogia* 24:167–167
- Bungartz FR, Grube J, Elix JA, Heininger C, Mayrhofer H (2011) A taxonomic revision of the *Buellia subalbula*-group in the Southern Hemisphere using fluorescence microscopy. *Biblioth Lichenol* 106:21–39
- Bungartz F, Dután-Patiño VL, Elix JA (2013a) The lichen genera *Cryptothecia*, *Herpothallon* and *Helminthocarpon* (*Arthoniales*) in the Galapagos Islands, Ecuador. *Lichenologist* 45:739–762
- Bungartz F, Hillmann G, Kalb K, Elix JA (2013b) Leprose and leproid lichens of the Galapagos, with a particular focus on *Lepraria* (*Stereocaulaceae*) and *Septotrapelia* (*Pilocarpaceae*). *Phytotaxa* 150:1–28
- Bungartz F, Elix JA, Kalb K, Giralt M (2016) New and overlooked species from the Galapagos Islands: the generic concept of *Diploicia* reassessed. *Lichenologist* 48:489–515
- Burgos MRG, Katimbang MLB, Paz MD, Beligan GA et al (2013) Genotypic variability and aggressiveness of *Bipolaris oryzae* in the Philippines. *Eur J Plant Pathol* 137:415–429

- Burneviča N, Jansons Ā, Zaļuma A, Kļaviņa D et al (2016) Fungi inhabiting bark stripping wounds made by large game on stems of *Picea abies* (L.) Karst. in Latvia. *Baltic For* 22:2–7
- Butin H, Holdenrieder O, Sieber TN (2013) The complete life cycle of *Petrakia echinata*. *Mycol Prog* 12:427–435
- Buyck B, Hosaka K, Masi S, Hofstetter V (2016) Molecular analyses of first collections of *Elaphomycetes* Nees (*Elaphomycetaceae*, *Eurotiales*, Ascomycota) from Africa and Madagascar indicate that the current concept of *Elaphomycetes* is polyphyletic. *Cryptog Mycol* 37:3–14
- Cabral A, Groenewald JZ, Rego C, Oliveira H, Crous PW (2012) *Cylindrocarpon* root rot: multi-gene analysis reveals novel species within the *Ilyonectria radicicola* species complex. *Mycol Prog* 11:655–688
- Cáceres MES, Aptroot A (2016) First inventory of lichens from the Brazilian Amazon in Amapá State. *Bryologist* 119:250–265
- Cáceres MES, da Silva Plata ER, Lücking R (2012a) *Malmographina*, a new genus for *Graphina malmei* (Ascomycota: Ostropales: Graphidaceae). *Lichenologist* 44:115–120
- Cáceres MES, Santos Viera T, Santos De Jesus L, Lücking R (2012b) New and interesting lichens from the Caxiuanã National Forest in the Brazilian Amazon. *Lichenologist* 44:807–812
- Cáceres MES, Aptroot A, Parnmen S, Lücking R (2014) Remarkable diversity of the lichen family *Graphidaceae* in the Amazon rain forest of Rondônia, Brazil. *Phytotaxa* 189:87–136
- Cai L, Guo XY, Hyde KD (2008) Morphological and molecular characterisation of a new anamorphic genus *Cheirosporium*, from freshwater in China. *Persoonia* 20:53–58
- Cai L, Hu DM, Liu F, Hyde KD, Jones EBG (2014) The molecular phylogeny of freshwater *Sordariomycetes* and discomycetes. *Freshwater Fungi and Fungal-like Organisms*. De Gruyter, Berlin, pp 47–72
- Calça FJS, Delmont M, Xavier-Santos S (2015) *Delitschia gigaspora* var. *pescanii*: a new variety of coprophilous fungus from Brazil. *Mycosphere* 6:122–126
- Calado ML, Carvalho L, Pang KL, Barata M (2015) Diversity and ecological characterization of sporulating higher filamentous marine fungi associated with *Spartina maritima* (Curtis) Fernald in two Portuguese salt marshes. *Microb Ecol* 70:612–633
- Campbell J, Ferrer A, Raja HA, Sivachai S et al (2007) Phylogenetic relationships among taxa in the *Jahnulales* inferred from 18S and 28S nuclear ribosomal DNA sequences. *Botany* 85:873–882
- Campbell J, Inderbitzin P, Kohlmeyer J, Volkmann-Kohlmeyer B (2009a) *Koralionastetales*, a new order of marine Ascomycota in the Sordariomycetes. *Mycol Res* 113:373–380
- Campbell J, Marvanová L, Gulis V (2009b) Evolutionary relationships between aquatic anamorphs and teleomorphs: *Trichladium* and *Varicosporium*. *Mycol Res* 113:1322–1334
- Candan M, Schultz M (2015) New and additional records of cyanolichens from Turkey. *Herzogia* 28:359–369
- Cannon PF, Damm U, Johnston PR, Weir BS (2012) *Colletotrichum*—current status and future directions. *Stud Mycol* 73:181–213
- Cansaran-Duman D, Halici MG (2012) *Squamaria lentigera* türlerinde usnik asit konsantrasyonunun antimikrobiyal aktivitesi. *Turk Hijyen ve Deneysel Biyoloji Dergisi*
- Cantillo-Pérez T, Gusmão LF, Castañeda-Ruiz RF (2016) *Tretolytea*, a new genus from the Brazilian semiarid region. *Mycotaxon* 130:977–981
- Cao S, Wei X, Zhou Q, Wei J (2013) *Phyllobaeis crustacea* sp. nov. from China. *Mycotaxon* 126:31–36
- Cao S, Zhang F, Liu C, Hao Z et al (2015) Distribution patterns of haplotypes for symbionts from *Umbilicaria esculenta* and *U. muehlenbergii* reflect the importance of reproductive strategy in shaping population genetic structure. *BMC Microbiol* 15:212
- Capdet M, Romero AI (2010) Fungi from palms in Argentina. 1. *Mycotaxon* 112:339–355
- Carbone M (2013) A type study of *Pseudoplectania stygia* (Pezizales). *Ascomycete* 5:33–38
- Carbone M, Van Vooren N (2011) *Otidea fuckelii*, una nuova specie pubblicata per chiarire le differenti interpretazioni di *Otidea leporina*. *Rivista di Micologia* 52:313–330
- Carbone M, Agnello C, Alvarado P (2013) Phylogenetic studies in the family *Sarcosomataceae* (Ascomycota, Pezizales). *Ascomycete* 5:1–2
- Carbone M, Agnello C, Alvarado P, Krom I (2014) *Donadinia sibirica* (Ascomycota, Pezizales) a new species from Russia. *Mycol Montenegrina* 17:53–64
- Carissimi M, Giraudo MS, Germani JC, Van Der Sand ST (2010) Antifungal activity of *Bacillus* sp. E164 against *Bipolaris sorokiniana*. *Biociências* (On-line) 17
- Carlucci A, Raimondo M, Santos J, Phillips AJL (2012) *Plectosphaerella* species associated with root and collar rots of horticultural crops in southern Italy. *Persoonia* 28:34–48
- Carmarán CC, Pildain MB, Vassiljeva LN (2009) The family *Diatrypaceae* (Ascomycota) in Argentina: new species and new records. *Nova Hedw* 88:521–530
- Carmarán CC, Berretta M, Martínez S, Barrera V et al (2015) Species diversity of *Cladorrhinum* in Argentina and description of a new species, *Cladorrhinum australe*. *Mycol Prog* 14:94
- Carmo LT, Monteiro JS, Gusmão LF, Sotão HM et al (2014) *Anabahusakala*, a new genus from the Brazilian Amazon rainforest. *Mycotaxon* 127:11–15
- Carneiro de Almeida DA, Miller AN, Pascholati Gusmão LF (2014) New species and combinations of conidial fungi from the semi-arid Caatinga biome of Brazil. *Nova Hedw* 98:431–447
- Carris LM, Peever T, McCotter SW (2016) Mitosporic stages of *Disciotis*, *Gyromitra* and *Morchella* in the inland Pacific Northwest USA. *Mycologia* 107:729–744
- Carter HE, Cools HJ, West JS, Shaw MW, Fraaije BA (2013) Detection and molecular characterisation of *Pyrenopeziza brassicae* isolates resistant to methyl benzimidazole carbamates. *Pest Manag Sci* 69:1040–1048
- Carvalho RO, Braga FR, Araújo JV (2011) Viability and nematophagous activity of the freeze-dried fungus *Arthrobotrys robusta* against *Ancylostoma* spp. infective larvae in dogs. *Vet Parasitol* 176:236–239
- Casaregola S, Jacques N, Louis-Mondesir C, Coton M, Coton E (2013) *Citeromyces nyonsensis* sp. nov., a novel yeast species isolated from black olive brine. *Int J Syst Evol Microbiol* 63:3086–3090
- Castañeda-Ruiz RF, Guerrero B, Adamo GM, Morillo O et al (2009a) A new species of *Selenosporella* and two microfungi recorded from a cloud forest in Mérida, Venezuela. *Mycotaxon* 109:63–74
- Castañeda-Ruiz RF, Gusmão LFP, de Moraes Júnior VO et al (2009b) Two setose anamorphic fungi: *Ampullicephala* gen. nov. and *Venustosynnema grandiae*. *Mycotaxon* 109:275–288
- Castañeda-Ruiz RF, Iturriaga T, Minter DW, Heredia G et al (2009c) Two new anamorphic fungi and some microfungi recorded from 'El Ávila', Venezuela. *Mycotaxon* 107:225–237
- Castañeda-Ruiz RF, Heredia G, Arias RM, Stadler M et al (2010a) *Anaselenosporella sylvatica* gen. & sp. nov. and *Pseudoacrodictys aquatica* sp. nov., two new anamorphic fungi from Mexico. *Mycotaxon* 112:65–74
- Castañeda-Ruiz RF, Heredia G, Arias RM, Becerra C et al (2010b) *Elotespora*, an enigmatic anamorphic fungus from Tabasco, Mexico. *Mycotaxon* 111:197–203
- Castañeda-Ruiz RF, Morillo O, Tovar B, Hernández Z et al (2010c) *Endogenospora*, a new genus of anamorphic fungi from Venezuela. *Mycotaxon* 112:75–82
- Castañeda-Ruiz RF, Silvera-Simón C, Gené J, Guarro J et al (2010d) A new species of *Corynesporopsis* from Portugal. *Mycotaxon* 114:407–415

- Castañeda-Ruiz RF, Heredia G, Arias RM, McKenzie EHC et al (2011) A new species and re-disposed taxa in *Repetophragma*. *Mycosphere* 2:273–289
- Castañeda-Ruiz RF, Heredia G, Arias RM, Stadler M et al (2012a) Two new fungi from Mexico: *Anaseptoidium* gen. nov. and *Cylindrosympodium sosae* sp. nov. *Mycotaxon* 119:141–148
- Castañeda-Ruiz RF, Hernández-Restrepo M, Gené J, Guarro J et al (2012b) Two new microfungi from Portugal: *Magnohelicospora iberica* gen. & sp. nov. and *Phaeodactylium stadleri* sp. nov. *Mycotaxon* 121:171–179
- Castañeda-Ruiz RF, Granados MM, Mardones M, Staler M et al (2013a) A microfungus from Costa Rica: *Ticosynnema* gen. nov. *Mycotaxon* 122:255–259
- Castañeda-Ruiz RF, Hernández-Restrepo M, Gené J, Granados M, Guarro J (2013b) Two new species of *Repetophragma* from the Iberian Peninsula. *Mycotaxon* 125:209–215
- Castañeda-Ruiz RF, Hernández-Restrepo M, Gené J, Guarro J et al (2013c) Two new microfungi from Portugal: *Magnohelicospora iberica* gen. & sp. nov. and *Phaeodactylium stadleri* sp. nov. *Mycotaxon* 121:171–179
- Castañeda-Ruiz RF, Leão-Ferreira SM, Gusmão LF (2015) *Distostrophmia*, a new genus of microfungi to accommodate *Endophragmiella rigidiuscula*. *Mycotaxon* 130:499–503
- Castañeda-Ruiz RF, Heredia G, Gusmão LFP, Li DW (2016a) Fungal diversity of Central and South America. In biology of microfungi. Springer, New York, pp 197–217
- Castañeda-Ruiz RF, Ma YR, Xia JW, Zhang XG (2016b) Two new species of *Pseudoacrodiclys* from Hainan, China. *Nova Hedw* 102:69–75
- Castellano MA, Trappe JM, Vernes K (2011) Australian species of *Elaphomycetes* (*Elaphomycetaceae*, *Eurotiales*, *Ascomycota*). *Aust Syst Bot* 24:32–57
- Castellano MA, Beever RE, Trappe JM (2012a) Sequestrate fungi of New Zealand: *Elaphomycetes* (*Ascomycota*, *Eurotiales*, *Elaphomycetaceae*). *NZ J Bot* 50:423–433
- Castellano MA, Guevara Guerrero G, García Jiménez J, Trappe JM (2012b) *Elaphomycetes appalachensis* and *E. verruculosus* sp. nov. (*Ascomycota*, *Eurotiales*, *Elaphomycetaceae*) from Eastern North America. *Rev Mex Micol* 35:17–22
- Castellano MA, Dentinger BTM, Séne O, Elliott TF et al (2016) New species of *Elaphomycetes* (*Elaphomycetaceae*, *Eurotiales*, *Ascomycota*) from tropical rainforests of Cameroon and Guyana. *IMA Fungus* 7:59–73
- Castillo JD, Lawrence KS, Kloepper JW, Van Santen E (2010) Evaluation of *Drechslerella dactyloides*, *Drechslerella brochopaga*, and *Paecilomyces lilacinus* for biocontrol of *Rotylenchulus reniformis*. *Nematropica* 40:71–86
- Castroagudín VL, Moreira SI, Pereira DAS, Moreira SS et al (2016) *Pyricularia graminis-tritici*, a new *Pyricularia* species causing wheat blast. *Persoonia* 37:199–216
- Catal M, Jordan SA, Butterworth SC, Schilder AM (2007) Detection of *Eutypa lata* and *Eutypella vitis* in grapevine by nested multiplex polymerase chain reaction. *Phytopathology* 97:737–747
- Catania MD, Romero A (2012) *Crassoascus monocaudatus* and *Iodosphaeria podocarpi*, two new species on *Podocarpus parlatorei* from "Las Yungas", Argentina. *Mycosphere* 3:37–44
- Catania MD, Romero AI, Huhndorf SM, Miller AN (2011) A new species and new records of *Cercophora* from Argentina. *Mycologia* 103:1372–1383
- Caalcante JG (2012) Fatores associados à estrutura da comunidade de liquens corticólicos crostosos em duas áreas de caatinga no estado de alagoas. PhD thesis, Universidade Federal de Sergipe
- Centeno DC, Hell AF, Braga MR, del Campo EM, Casano LM (2016) Contrasting strategies used by lichen microalgae to cope with desiccation-rehydration stress revealed by metabolite profiling and cell wall analysis. *Environ Microbiol* 18:1546–1560
- Cestaro L, Tønsberg T, Muggia L (2016) Phylogenetic data and chemical traits characterize a new species in the lichen genus *Tephromela*. *Herzogia* 29:383–402
- Chacón S, Dörge D, Weisenborn J, Piepenbring M (2013) A new species and a new record of *Diatrypaceae* from Panama. *Mycologia* 105:681–688
- Chacón S, Tapia F, Esqueda M (2014) New records of *Dothideomycetes* from Mexico. *Mycotaxon* 128:145–157
- Chakraborty S, Basak B, Dutta S, Bhunia B, Dey A (2013) Decolorization and biodegradation of congo red dye by a novel white rot fungus *Alternaria alternata* CMERI F6. *Bioresour Technol* 147:662–666
- Chamizo S, Rodríguez-Caballero E, Roncero B, Román JR, Cantón Y (2016) April. Factors controlling spatial distribution patterns of biocrusts in a heterogeneous and topographically complex semiarid area. In EGU General Assembly Conference Abstracts 18:7937
- Chand R, Sen D, Prasad KD, Singh AK et al (2008) Screening for disease resistance in barley cultivars against *Bipolaris sorokiniana* using callus culture method. *Indian J Exp Biol* 46:249
- Chand R, Kumar M, Kushwaha C, Shah K, Joshi AK (2014) Role of melanin in release of extracellular enzymes and selection of aggressive isolates of *Bipolaris sorokiniana* in barley. *Curr Microbiol* 69:202–211
- Chang JH, Wang YZ (2009) The genera *Sporormia* and *Preussia* (*Sporomiaceae*, *Pleosporales*) in Taiwan. *Nova Hedw* 88:245–254
- Chang JH, Kao HW, Wang YZ (2010) Molecular phylogeny of *Cercophora*, *Podospora*, *Schizothecium* (*Lasiosphaeriaceae*, *Pyrenomycetes*). *Taiwania* 55:110–116
- Chang YC, Lu CK, Chiang YR, Wang GJ et al (2014) Diterpene glycosides and polyketides from *Xylotumulus gibbisporus*. *J Nat Prod* 77:751–757
- Chaverri P, Gazis RO (2010) *Perisporiopsis lateritia*, a new species on decaying leaves of *Hevea* spp. from the Amazon basin in Peru. *Mycotaxon* 113:163–169
- Chaverri P, Gazis RO (2011) Linking ex planta fungi with their endophytic stages: *Perisporiopsis*, a common leaf litter and soil fungus, is a frequent endophyte of *Hevea* spp. and other plants. *Fung Ecol* 4:94–102
- Chaverri P, Liu M, Hodge KT (2008) A monograph of the entomopathogenic genera *Hypocrella*, *Moelleriella*, and *Samuelsiagen*. nov. (*Ascomycota*, *Hypocreales*, *Clavicipitaceae*), and their aschersonia-like anamorphs in the Neotropics. *Stud Mycol* 60:1–66
- Chaverri P, Salgado C, Hirooka Y, Rossman AY, Samuels GJ (2011) Delimitation of *Neonectria* and *Cylindrocarpon* (*Nectriaceae*, *Hypocreales*, *Ascomycota*) and related genera with Cylindrocarpon-like anamorphs. *Stud Mycol* 68:57–78
- Checa J, Arenal F, Blanco N, Rogers JD (2008) *Coniolariella hispanica* sp. nov. and other additions to *Coniolariella*. *Mycol Res* 112:795–801
- Checa J, Blanco MN, Moreno G (2013) Contributions to the family *Thyridiaceae*. New data on *Sphaeria mutabilis*. *Mycotaxon* 125:149–164
- Cheewangkoon R, Groenewald JZ, Summerell BA, Hyde KD et al (2009) *Myrtaceae*, a cache of fungal biodiversity. *Persoonia* 23:55–85
- Cheewangkoon R, Groenewald JZ, Verkley HJM, Hyde KD et al (2010) Re-evaluation of *Cryptosporiopsis eucalypti* and *Cryptosporiopsis*-like species occurring on Eucalyptus leaves. *Fungal Divers* 44:89–105

- Cheewangkoon R, Groenewald JZ, Hyde KD, To-anun C, Crous PW (2012) Chocolate Spot disease of *Eucalyptus*. *Micol Prog* 11:61–69
- Chen J, Meng ZX, Chen XM, Lv YL et al (2009) *Leohumicola*, a genus new to China. *Mycotaxon* 108:337–340
- Chen S, Gryzenhout M, Roux J, Xie Y et al (2011) Novel species of *Celoporthae* from *Eucalyptus* and *Syzygium* trees in China and Indonesia. *Mycologia* 103:1384–1410
- Chen JL, Lin YR, Hou CL, Wang SJ (2012) Species of *Rhytismataceae* on *Camellia* spp. from the Chinese mainland. *Mycotaxon* 118:219–230
- Chen S, Van Wyk M, Roux J, Wingfield MJ et al (2013a) Taxonomy and pathogenicity of *Ceratocystis* species on *Eucalyptus* trees in South China, including *C. chiaueucensis* sp. nov. *Fungal Divers* 58:267–279
- Chen SF, Wingfield MJ, Roux J (2013b) *Diversimorbus metrosiderotis* gen. et sp. nov. and three new species of *Holocryphia* (*Cryphonectriaceae*) associated with cankers on native *Metrosideros angustifoliata* trees in South Africa. *Fungal Biol* 117:289–310
- Chen YL, Gao Y, Zhang KQ, Zou CG (2013c) Autophagy is required for trap formation in the nematode-trapping fungus *Arthrobotrys oligospora*. *Environ Microbiol Rep* 5:511–517
- Chen L, Yue Q, Zhang X, Xiang M et al (2013d) Genomics-driven discovery of the pneumocandin biosynthetic gene cluster in the fungus *Glarea lozoyensis*. *BMC Genom* 14:339
- Chen SF, Wingfield MJ, Roets F, Roux J (2013e) A serious canker disease caused by *Immersiporthe knoxdaviesiana* gen. et sp. nov. (*Cryphonectriaceae*) on native *Rapanea melanophloeos* in South Africa. *Plant Pathol* 62:667–678
- Chen H, Jiang XJ, Yu Z, Duns GJ et al (2014) A novel mitovirus isolated from *Buergerella spartinae*. *Virilogica Sinica* 29:257–260
- Chen KH, Miadlikowska J, Molnár K, Arnold AE et al (2015a) Phylogenetic analyses of euotiomyctetous endophytes reveal their close affinities to *Chaetothyriales*, *Eurotiales*, and a new order—*Phaeomoniellales*. *Mol Phyl Evol* 85:117–130
- Chen Q, Jiang JR, Zhang GZ, Cai L, Crous PW (2015b) Resolving the *Phoma enigma*. *Stud Mycol* 82:137–217
- Chen AJ, Sun BD, Houbraken J, Frisvad JC et al (2016a) New *Talaromyces* species from indoor environments in China. *Stud Mycol* 84:119–144
- Chen C, Verkley GJ, Sun G, Groenewald JZ, Crous PW (2016b) Redefining common endophytes and plant pathogens in *Neofabrea*, *Pezicula*, and related genera. *Fungal Biol* 120:1291–1322
- Chen Y, Ran SF, Dai DQ, Wang Y et al (2016c) Mycosphere essays 2. *Myrothecium*. *Mycosphere* 7:64–80
- Chen YY, Maharachchikumbura SSN, Liu JK, Hyde KD et al (2017a) Fungi from Asian Karst formations I. *Pestalotiopsis photinicola* sp. nov., causing leaf spots of *Photinia serrulata*. *Mycosphere* 8:103–110
- Chen WH, Zeng GP, Luo Y, Liang ZQ, Han YF (2017b) Morphological traits and molecular analysis for *Geomycetes fujianensis* sp. nov. from China. *Mycosphere* 8:38–43
- Chen WH, Han YF, Liang JD, Liang ZQ, Jin DC (2017c) *Metarhizium dendrolimatis*, a novel *Metarhizium* species parasitic on *Dendrolimus* sp. larvae. *Mycosphere* 8:37
- Cheng XL, Li W, Zhang TY (2014) A new species of *Phaeoisaria* from intertidal marine sediment collected in Weihai, China. *Mycotaxon* 127:17–24
- Cheng XL, Li W, Cai L (2015) Molecular phylogeny of *Ascotricha*, including two new marine algae-associated species. *Mycologia* 27:14–210
- Cheplick GP, Faeth S (2009) Ecology and evolution of the grass-endophyte symbiosis. Oxford University Press, Oxford
- Chethana T, Liu M, Ariyawansa HA, Konta S et al (2015) Splanchnonema-like species in *Pleosporales*: introducing *Pseudosplanchnonema* gen. nov. in *Massarinaceae*. *Phytotaxa* 231:133–144
- Chhikara S, Chaudhury D, Dhankher OP, Jaiwal PK (2012) Combined expression of a barley class II chitinase and type I ribosome inactivating protein in transgenic *Brassica juncea* provides protection against *Alternaria brassicae*. *Plant Cell Tissue Organ Cult* 108:83–89
- Chigoziri E, Ekefan EJ (2013) Seed-borne fungi of Chilli pepper (*Capsicum frutescens*) from pepper producing areas of Benue state, Nigeria. *Agric Biol J Am* 4:370–374
- Chilvers MI, Rogers JD, Dugan FM, Stewart JE et al (2009) *Didymella pisi* sp. nov., the teleomorph of *Ascochyta pisi*. *Micol Res* 113:391–400
- Chlebicki A (2009a) Fungi on higher plants of the upper limit of the alpine zone: new species from Tian Shan. *Mycotaxon* 110:443–450
- Chlebicki A (2009b) Some endophytes of *Juncus trifidus* from Tatra Mts. in Poland. *Acta Mycologica* 44:11–17
- Chlebicka M, Konvalinkova T (2010) *Pirottaea lychnidis* comb. nov. from the Bohemian Switzerland National Park, Czech Republic. *Czech Mycol* 62:19–32
- Cho Y, Kim KH, La Rota M, Scott D et al (2009) Identification of novel virulence factors associated with signal transduction pathways in *Alternaria brassicicola*. *Mol Microbiol* 72:1316–1333
- Cho Y, Srivastava A, Ohm RA, Lawrence CB et al (2012) Transcription factor Amr1 induces melanin biosynthesis and suppresses virulence in *Alternaria brassicicola*. *PLoS Pathog* 8:1002974
- Cho SE, Park JH, Choi JK, Shin HD (2015) First report of zonate leaf spot of *Glycine max* caused by *Cristulariella moricola* in Korea. *J Phytopathol* 163:1019–1022
- Choi JS, Graser Y, Walther G, Peano A et al (2012) *Microsporum mirabile* and its teleomorph *Arthroderma mirabile*, a new dermatophyte species in the *M. cookei* clade. *Med Mycol* 50:161–169
- Choi J, Park SY, Kim BR, Roh JH et al (2013a) Comparative analysis of pathogenicity and phylogenetic relationship in *Magnaporthe grisea* species complex. *PLoS ONE* 8:57196
- Choi YJ, Shin HD, Han JG, Pfister DH (2013b) *Scutellinia (Pezizales)* in Korea, with a new species and eight new records. *Nova Hedw* 97:457–476
- Choi IY, Kim BS, Park JH, Cho SE, Shin HD (2014) First report of black stem caused by *Botryosporium longibrachiatum* on statice in Korea. *Plant Dis* 98:1431
- Chokpaiboon S, Choodej S, Boonyuen N, Teerawatananond T, Pudhom K (2016) Highly oxygenated chromones from mangrove-derived endophytic fungus rhytidhysteron rufulum. *Phytochemistry* 122:172–177
- Chomnunti P, Schoch CL, Aguirre-Hudson B, Ko Ko TW et al (2011) Capnodiaeae. *Fungal Divers* 51:103–134
- Chomnunti P, Bhat DJ, Jones EB, Chukeatirote E et al (2012a) *Trichocomeriaceae*, a new sooty mould family of *Chaetothyriales*. *Fungal Divers* 56:63–76
- Chomnunti P, Ko TW, Chukeatirote E, Hyde KD, Cai L et al (2012b) Phylogeny of *Chaetothyriaceae* in northern Thailand including three new species. *Mycologia* 104:382–395
- Chomnunti P, Hongsanan S, Hudson BA, Tian Q et al (2014) The sooty moulds. *Fungal Divers* 66:1–36
- Chowdhary A, Randhawa HS, Singh V, Khan ZU et al (2011) *Bipolaris hawaiiensis* as etiologic agent of allergic bronchopulmonary mycosis: first case in a paediatric patient. *Med Mycol* 49:760–765

- Chruszcz M, Chapman MD, Osinski T, Solberg R et al (2012) *Alternaria alternata* allergen Alt a 1: a unique β-barrel protein dimer found exclusively in fungi. *J Allergy Clin Immunol* 130:241–247
- Chu Y, Alias SA, Iddi R, Guo SY, Pang KL (2015) A molecular reappraisal of *Nimbospora* (*Halosphaeriaceae*, *Microascales*) and a new genus *Ebullia* for *N. octonae*. *Mycoscience* 56:34–41
- Chuaseeharonnachai C, Yamaguchi K, Sri-Indrasutdhia V, Somrithipol S et al (2013) Diversity of aero-aquatic hyphomycetes from six streams in Doi Inthanon and Khao Yai tropical forests, Thailand. *Cryptog Mycol* 34:183–197
- Chuaseeharonnachai C, Somrithipol S, Boonyuen N (2014) A new species of *Fusticeps* from Thailand. *Mycosphere* 5:313–317
- Chung KR (2011) *Elsinoë fawcettii* and *Elsinoë australis*: the fungal pathogens causing citrus scab. *Mol Plant Pathol* 12:123–135
- Chungu D, Gryzenhout M, Muimba-Kankolongo A, Wingfield MJ, Roux J (2010) Taxonomy and pathogenicity of two novel *Chrysoporthe* species from *Eucalyptus grandis* and *Syzygium guineense* in Zambia. *Mycol Prog* 9:379–393
- Clowez P (2012) Les morilles, une nouvelle approche mondiale du genre *Morchella*. *Bulletin de la Société Mycologique de France* 126:199–376
- Çobanoğlu G (2011) Additional and new lichen records for the province of Giresun. *Fen Bilimleri Dergisi* 23:83–88
- Çobanoğlu G, Yavuz M, Costache I, Radu I (2010) Additional and new lichen records from Cozia National Park, Romania. *Mycotaxon* 114:193–196
- Çobanoğlu G, Sesal C, Açıkgöz B, Karaltı İ (2016) Evaluation of antimicrobial activity of the lichens *Physcia aipolia*, *Xanthoria parietina*, *Usnea florida*, *Usnea subfloridana* and *Melanohalea exasperata*. *Mod Phytomorphol* 10:19–24
- Coda R, Rizzello CG, Di Cagno R, Trani A et al (2013) Antifungal activity of *Meyerozyma guilliermondii*: identification of active compounds synthesized during dough fermentation and their effect on long-term storage of wheat bread. *Food Microbiol* 33:243–251
- Consolo VF, Cordo CA, Salerno GL (2008) DNA fingerprint and pathotype diversity of *Pyricularia oryzae* populations from Argentina. *Australas Plant Pathol* 37:357–364
- Cook RTA, Inman AJ, Billings C (1997) Identification and classification of powdery mildew anamorphs using light and scanning electron microscopy and host range data. *Micol Res* 101:975–1002
- Coppins BJ (2008) *Ameliella*, a new genus of lichen-forming fungi from north-west Europe and western Canada. *Lichenologist* 40:387–397
- Cornejo C, Scheidegger C (2015) Multi-gene phylogeny of the genus *Lobaria*: evidence of species-pair and allopatric cryptic speciation in East Asia. *Am J Bot* 102:2058–2073
- Cort C (2012) A phylogenetic analysis of *Heterocephalum*: Using molecular tools to reassess the taxonomic placement of an unusual and infrequently collected fungus. Honors Theses. Paper 7
- Costa PM, Barbosa MA, Malosso E, Castañeda-Ruiz RF (2016) *Anaverticladus uncinatus* gen. & sp. nov. from decaying leaves from Brazil. *Mycotaxon* 131:687–691
- Coste C, Dufrêne P (2009) Reasoned Inventory of the lichens and lichenicolous fungi of the Cascades of Mortain (department of Manche, 50). Société Tarnaise de Sciences Naturelles, pp 57–80
- Cota BB, Rosa LH, Caligorne RB, Raballo ALT et al (2008) Altenusin, a biphenyl isolated from the endophytic fungus *Alternaria* sp., inhibits trypanothione reductase from *Trypanosoma cruzi*. *FEMS Microbiol Lett* 285:177–182
- Crane C, Burgess TI (2013) *Luteocirrhush shearii* gen. sp. nov. (*Diaporthales*, *Cryphonectriaceae*) pathogenic to *Proteaceae* in the south Western Australian Floristic Region. *IMA Fungus* 4:111–122
- Craven KD, Véliz H, Cho Y, Lawrence CB, Mitchell TK (2008) Anastomosis is required for virulence of the fungal necrotroph *Alternaria brassicicola*. *Eukaryot Cell* 7:675–683
- Crespo A, Ferencova Z, Pérez-Ortega S, Elix JA, Divakar PK (2010a) *Austroparmelina*, a new Australian lineage in parmelioid lichens (*Parmeliaceae*, *Ascomycota*). *Syst Biodivers* 8:209–221
- Crespo A, Kauff F, Divakar PK, del Prado R et al (2010b) Phylogenetic generic classification of parmelioid lichens (*Parmeliaceae*, *Ascomycota*) based on molecular, morphological and chemical evidence. *Taxon* 59:1735–1753
- Crespo A, Divakar PK, Hawksworth DL (2011) Generic concepts in parmelioid lichens, and the phylogenetic value of characters used in their circumscription. *Lichenologist* 43:511–535
- Crisan F, Ardelean I (2010) The analysis of ecological behaviour of the macrolichens from Rosia Montana (Romania). *Studia Universitatis Vasile Goldis Seria Stiintele Vietii (Life Sciences Series)* 20
- Crittenden PD, Scrimgeour CM, Minnallina G, Sutton MA et al (2015) Lichen response to ammonia deposition defines the footprint of a penguin rookery. *Biogeochemistry* 122:295–311
- Crous PW, Verkley GJM (2009) *Xenocylindrosporium*. *Fungal Planet* 44
- Crous PW, Groenewald JZ (2010) Fungal planet 55. *Xenopolyscytalum pinea* gen. et sp. nov. *Persoonia* 25:130–131
- Crous PW, Groenewald JZ (2011) Why everlastings dont last. *Persoonia* 26:70–84
- Crous PW, Groenewald JZ (2013) A phylogenetic re-evaluation of *Arthrinium*. *IMA Fungus* 4:133–154
- Crous PW, Wood AR, Okada G, Groenewald JZ (2008) Follicolous microfungi occurring on *Encephalartos*. *Persoonia* 21:135–146
- Crous PW, Braun U, Wingfield MJ, Wood AR et al (2009a) Phylogeny and taxonomy of obscure genera of microfungi. *Persoonia* 22:139–161
- Crous PW, Groenewald JZ, Lee SS (2009b) Fungal planet 41. *Nawawia malaysiana*. *Persoonia* 23:194–195
- Crous PW, Groenewald JZ, Summerell BA, Wingfield BD, Wingfield MJ (2009c) Co-occurring species of *Teratosphaeria* on *Eucalyptus*. *Persoonia* 22:38–48
- Crous PW, Schoch CL, Hyde KD, Wood AR et al (2009d) Phylogenetic lineages in the *Capnodiales*. *Stud Mycol* 64:17–47
- Crous PW, Summerell BA, Carnegie AJ, Wingfield MJ, Groenewald JZ (2009e) Novel species of *Mycosphaerellaceae* and *Teratosphaeriaceae*. *Persoonia* 23:119–146
- Crous PW, Summerell BA, Carnegie AJ, Wingfield MJ et al (2009f) Unravelling *Mycosphaerella*: do you believe in genera? *Persoonia* 23:99–118
- Crous PW, Barreto RW, Alfenas AC, Alfenas RF, Groenewald JZ (2010a) what is *Johansonia*? *IMA Fungus* 1:117–122
- Crous PW, Groenewald JZ, Shin HD (2010b) Fungal planet 56. *Strelitziana albiziae* sp. nov. *Persoonia* 25:132–133
- Crous PW, Groenewald JZ, Shivas RG (2010c) Fungal planet 60. *Pseudophloeospora eucalypti* gen. et sp. nov. *Persoonia* 25:140–141
- Crous PW, Groenewald JZ, Shivas RG, Edwards J et al (2011a) Fungal Planet description sheets:69–91. *Persoonia* 26:108–156
- Crous PW, Minnis AM, Pereira OL, Alfenas AC et al (2011b) What is *Scirrhia*? *IMA Fungus* 2:127–133
- Crous PW, Summerell BA, Shivas RG, Romberg M et al (2011c) Fungal Planet description sheets: 92–106. *Persoonia* 27:130–162
- Crous PW, Summerell BA, Swart L, Denman S et al (2011d) Fungal pathogens of *Proteaceae*. *Persoonia* 27:20–45
- Crous PW, Tanaka K, Summerell BA, Groenewald JZ (2011e) Additions to the *Mycosphaerella* complex. *IMA Fungus* 2:49–64

- Crous PW, Braun U, Hunter GC, Wingfield MJ et al (2012a) Phylogenetic lineages in *Pseudocercospora*. Stud Mycol 75:37–114
- Crous PW, Groenewald JZ, Lombard L, Wingfield MJ (2012b) *Homortomyces* gen. nov., a new dothidealean pycnidial fungus from the Cradle of Humankind. IMA Fungus 3:109–115
- Crous PW, Shivas RG, Wingfield MJ, Summerell BA et al (2012c) Fungal Planet description sheets 128–153
- Crous PW, Summerell BA, Alfenas AC, Edwards J et al (2012d) Genera of diaporthalean coelomycetes associated with leaf spots of tree hosts. Persoonia 28:66–75
- Crous PW, Summerell BA, Shivas RG, Burgess TI et al (2012e) Fungal planet description sheets: 107–127. Persoonia 28:138–182
- Crous PW, Summerell BA, Shivas RG, Carnegie AJ et al (2012f) A re-appraisal of *Harknessia* (*Diaporthales*), and the introduction of *Harknessiaceae* fam. nov. Persoonia 28:59–65
- Crous PW, Verkley GJM, Christensen M, Groenewald JZ (2012g) How important are conidial appendages? Persoonia 28:126–137
- Crous PW, Quaedvlieg W, Sarpkaya K, Can C, Erkiliç A (2013a) Septoria-like pathogens causing leaf and fruit spot of pistachio. IMA Fungus 4:187–199
- Crous PW, Wingfield MJ, Guarro J, Cheewangkoon R et al (2013b) Fungal Planet description sheets: 154–213. Persoonia 31:188–296
- Crous PW, Giraldo A, Hawksworth DL, Robert V et al (2014a) The Genera of Fungi: fixing the application of type species of generic names. IMA Fungus 5:141–160
- Crous PW, Quaedvlieg W, Hansen K, Hawksworth DL, Groenewald JZ (2014b) *Phacidium* and *Ceuthospora* (*Phaciidaeae*) are congeneric: taxonomic and nomenclatural implications. IMA Fungus 5:173–193
- Crous PW, Shivas RG, Quaedvlieg W, van Der Bank M et al (2014c) Fungal planet description sheets: 214–280. Persoonia 32:184–306
- Crous PW, Wingfield MJ, Schumacher RK, Summerell BA et al (2014d) Fungal planet description sheets: 281–319. Persoonia 33:212–289
- Crous PW, Carris LM, Giraldo A, Groenewald JZ et al (2015a) The Genera of Fungi: fixing the application of the type species of generic names – G 2: *Allantophomopsis*, *Latorua*, *Macrodiploidiopsis*, *Macrohilum*, *Milospium*, *Protostegia*, *Pyricularia*, *Robillarda*, *Rotula*, *Septoriella*, *Torula*, and *Wojnowicia*. IMA Fungus 6:163–198
- Crous PW, Muller MM, Sánchez RM, Giordano L et al (2015b) Resolving *Tiarosporella* spp. allied to *Botryosphaeriaceae* and *Phaciidaeae*. Phytotaxa 202:73–93
- Crous PW, Schumacher RK, Wingfield MJ, Lombard L et al (2015c) Fungal Systematics and Evolution: FUSE 1. Sydowia 67:81–118
- Crous PW, Wingfield MJ, Guarro J, Hernandez-Restrepo M et al (2015d) Fungal Planet description sheets: 320–370. Persoonia 34:167–266
- Crous PW, Wingfield MJ, Le Roux JJ, Richardson DM et al (2015e) Fungal Planet description sheets: 371–399. Persoonia 35:264–327
- Crous PW, Wingfield MJ, Burgess TI, Hardy GS et al (2016a) Fungal planet description sheets: 469–557. Persoonia 37:218–403
- Crous PW, Wingfield MJ, Richardson DM, Leroux JJ et al (2016b) Fungal planet description sheets: 400–468. Persoonia 36:316–458
- Crous PW, Wingfield MJ, Burgess TI, Hardy GS et al (2017) Fungal planet description sheets: 558–624. Persoonia 38:240–384
- Cruywagen EM, de Beer ZW, Roux J, Wingfield MJ (2010) Three new *Graphium* species from baobab trees in South Africa and Madagascar. Persoonia 25:61–71
- Cruz ACR, dos Santos SIT, Leão-Ferreira SM, Gusmão LFP (2009) Conidial fungi from the semi-arid Caatinga biome of Brazil. New species and new records of *Helicosporium*. Mycotaxon 110:53–64
- Cruz ACR, Gusmão LF, Castañeda-Ruiz RF, Stadler M, Minter DW (2012) *Zelodactylaria*, an interesting new genus from semi-arid northeast Brazil. Mycotaxon 119:241–248
- Cui Y, Jia H, He D, Yu H et al (2013) Characterization of *Edenia gomezpompae* isolated from a patient with keratitis. Mycopathologia 176:75–81
- Culleré L, Ferreira V, Chevreau B, Venturini ME et al (2010) Characterisation of aroma active compounds in black truffles (*Tuber melanosporum*) and summer truffles (*Tuber aestivum*) by gas chromatography–olfactometry. Food Chem 122:300–306
- Cunnington JH, Lawrie AC, Pascoe IG (2010) Genetic characterization of the *Golovinomyces cichoracearum* complex in Australia. Plant Pathol 59:158–164
- Czarnota P (2009) *Japewia tornensis* and further localities of *J. subaurifera* found in the Carpathians. Acta Mycologica 44:59–264
- Czarnota P, Guzow-Krzeminska B (2012) ITS rDNA data confirm a delimitation of *Bacidina arnoldiana* and *B. sulphurella* and support a description of a new species within the genus *Bacidina*. Lichenologist 44:743–755
- Czarnota P, Hernik E (2013) *Mniaecia jungermanniae* and *Puttea margaritella* (lichenized Ascomycota) found in Poland. Acta Societatis Botanicorum Poloniae 82
- Czarnota P, Hernik E (2014) Some peltigericolous microlichens from southern Poland. Acta Botanica Croatica 73:159–170
- Czarnota P, Kukwa M (2009) Contribution to the knowledge of some poorly known lichens in Poland. III. *Trapelia corticola* and the genus *Vezdaea*. Folia Cryptog Estonica 46:25–31
- D'Souza MA, Bhat DJ (2013) Occurrence of microfungi as litter colonizers and endophytes in varied plant species from the Western Ghats forests, Goa, India. Mycosphere 4:567–582
- da Cruz ACR, Gusmão LFP (2009) Conidial fungi from caatinga: Lignicolous species. Acta Bot Bras 23:1133–1144
- da Cunha KC, Sutton DA, Fothergill AW, Cano J et al (2012) Diversity of *Bipolaris* species in clinical samples in the United States and their antifungal susceptibility profiles. J Clin Microbiol 50:4061–4066
- da Silva SS, Gusmão LF (2015) Some facultative-aquatic conidial fungi from the Caatinga biome II: new records for Brazil and Americas. Braz J Bot 38:323–331
- da Silva SS, Gusmão LFP, Castañeda-Ruiz RF (2015) *Cryptocoryneum parvulum*, a new species on *Araucaria angustigolia* (Brazilian pine). Mycotaxon 130:465–469
- da Silva M, Pinho DB, Pereira OL, Fernandes FM, Barreto RW (2016) Naming potentially endangered parasites: foliicolous mycobiota of *Dimorphandra wilsonii*, a highly threatened Brazilian tree species. PloS ONE 11:e0147895
- Dai DQ, Bhat DJ, Liu J, Chukeatirote E et al (2012) *Bambusicola*, a new genus from bamboo with asexual and sexual morphs. Cryptog Mycol 33:363–379
- Dai DQ, Bahkali AH, Bhat DJ, Xiao YP et al (2014a) Towards a natural classification of *Dothideomycetes* 3: The genera *Muellerites*, *Trematosphaeriopsis*, *Vizzellopsis* and *Yoshinagella* (*Dothideomycetes incertae sedis*). Phytotaxa 176:18–27
- Dai DQ, Bahkali AH, Li QR, Bhat DJ et al (2014b) *Vamsapriya* (*Xylariaceae*) re-described, with two new species and molecular sequence data. Cryptog Mycol 35:339–357
- Dai DQ, Boonmee S, Tian Q, Xiao YP et al (2014c) *Englerulaceae* (*Dothideomycetes*). Phytotaxa 176:139–155
- Dai DQ, Wijayawardene NN, Bhat DJ, Chukeatirote E et al (2014d) *Pustulomyces* gen. nov. accommodated in *Diaporthaceae*,

- Diaporthales*, as revealed by morphology and molecular analyses. *Cryptog Mycol* 35:63–72
- Dai DQ, Wijayawardene NN, Bhat DJ, Chukeatirote E et al (2014e) The phylogenetic placement of *Eriosporella bambusicola* sp. nov. in *Capnodiales*. *Cryptog Mycol* 35:41–49
- Dai DQ, Bahkali AH, Ariyawansa HA, Li WJ et al (2015a) *Neokalmusia didymospora* (*Didymosphaeriaceae*), a new species from bamboo. *Syndowia* 68:17–25
- Dai DQ, Bahkali AH, Li WJ, Bhat DJ et al (2015b) *Bambusicola loculata* sp. nov. (*Bambusicolaceae*) from bamboo. *Phytotaxa* 213(2):122–130
- Dai DQ, Jiang HB, Tang LZ, Bhat DJ (2016) Two new species of *Arthrinium* (*Apiosporaceae, Xylariales*) associated with bamboo from Yunnan, China. *Mycosphere* 7:1332–1345
- Dai DQ, Phookamsak R, Wijayawardene NN, Li WJ et al (2017) Bambusicolous fungi. *Fungal Divers* 82:1–105
- Dailey R, Siemion R, Raisbeck M, Jesse C (2010) Analysis of salazinic, norstictic, and usnic acids in *Xanthoparmelia chlorochroa* by ultra-performance liquid chromatography/tandem mass spectrometry. *J AOAC Int* 93:1137–1142
- Dallagnol LJ, Rodrigues FA, DaMatta FM, Mielli MV, Pereira SC (2011a) Deficiency in silicon uptake affects cytological, physiological, and biochemical events in the rice-*Bipolaris oryzae* interaction. *Phytopathology* 101:92–104
- Dallagnol LJ, Rodrigues FA, Martins SC, Cavatte PC, DaMatta FM (2011b) Alterations on rice leaf physiology during infection by *Bipolaris oryzae*. *Australas Plant Pathol* 40:360–365
- Dallagnol LJ, Rodrigues FA, Chaves ARM, Vale FXR, DaMatta FM (2013) Photosynthesis and sugar concentration are impaired by the defective active silicon uptake in rice plants infected with *Bipolaris oryzae*. *Plant Pathol* 62:120–129
- Damm U, Mostert L, Crous PW, Fourie PH (2008a) Novel species associated with necrotic wood of *Prunus* trees. *Persoonia* 20:87–102
- Damm U, Crous PW, Fourie PH (2008b) A fissitunicate ascus mechanism in the *Calosphaeriaceae*, and novel species of *Jattaea* and *Calosphaeria* on *Prunus* wood. *Persoonia* 20:39–52
- Damm U, Verkley GJM, Crous PW, Fourie PH et al (2008c) *Paraconiothyrium* species on stone fruit trees and other woody hosts, including *P. variabile* and *P. africanum* spp. nov. *Persoonia* 20:9–17
- Damm U, Baroncelli R, Cai L, Kubo Y et al (2010a) *Colletotrichum*: species, ecology and interactions. *IMA Fungus* 1(2):161–165
- Damm U, Fourie PH, Crous PW (2010b) *Coniochaeta* (*Lecythophora*), *Collophora* gen. nov. and *Phaeomoniella* species associated with wood necroses of *Prunus* trees. *Persoonia* 24:60–80
- Damm U, Cannon PF, Woudenberg JH, Crous PW (2012a) The *Colletotrichum acutatum* species complex. *Stud Mycol* 73:37–113
- Damm U, Cannon PF, Woudenberg JH, Johnston PR et al (2012b) The *Colletotrichum boninense* species complex. *Stud Mycol* 73:1–36
- Damm U, Cannon PF, Liu F, Barreto RW et al (2013) The *Colletotrichum orbiculare* species complex: Important pathogens of field crops and weeds. *Fungal Divers* 61:29–59
- Daranagama DA, Liu XZ, Chamuyang S, Stadler M, Hyde KD (2014) A multiple gene genealogy reveals phylogenetic placement of *Rhopalostroma lekae*. *Phytotaxa* 186:177–187
- Daranagama DA, Camporesi E, Tian Q, Liu XZ et al (2015a) *Anthostomella* is polyphyletic comprising several genera in *Xylariaceae*. *Fungal Divers* 73:203–238
- Daranagama DA, Liu XZ, Chamuyang S, Stadler M et al (2015b) *Rhopalostroma brevistipitatum* sp. nov. from Thailand with an extended generic description for *Rhopalostroma*. *Phytotaxa* 227:229–242
- Daranagama DA, Jones EBG, Liu XZ, To-anun C et al (2016a) Mycosphere Essays 13 – Do xylariaceous macromycetes make up most of the *Xylariomycetidae*? *Mycosphere* 7:582–601
- Daranagama DA, Camporesi E, Liu XZ, Bhat DJ et al (2016b) *Tristratiperidium microsporum* gen. et sp. nov. (*Xylariales*) on dead leaves of *Arundo plinii*. *Mycol Prog* 15:8
- Daranagama DA, Camporesi E, Liu XZ, Chamuyang S et al (2016c) *Lopadostoma fagi* (*Lopadostomataceae*) on *Fagus sylvatica* from Italy. *Stud Fungi* 1:80–89
- Daranagama DA, Liu XZ, Chamuyang S, Stadler M et al (2016d) Towards a natural classification of *Sordariomycetes*: The genera *Frondisphaeria*, *Immersisphaeria*, *Lasiobertia*, *Pulmosphaeria* and *Yuea* (*Sordariomycetes incertae sedis*). *Phytotaxa* 258:153–163
- Daranagama DA, Hyde KD, Sir EB, Thambugala KM et al. (2017) Towards a natural classification and backbone tree for *Graphostromataceae*, *Hypoxylaceae*, *Lopadostomataceae* and *Xylariaceae* (in prep)
- Davey ML, Currah RS (2009) *Atradidymella muscivora* gen. et sp. nov. (*Pleosporales*) and its anamorph *Phoma muscivora* sp. nov.: a new pleomorphic pathogen of boreal bryophytes. *Am J Bot* 96:1281–1288
- Davison EM, Davison PJN, Tan YP, Shivas RG (2009) Fungal Planet 31. *Oedocephalum adhaerens*. *Persoonia* 22:166–167
- Davydov EA, Printzen C (2012) Rare and noteworthy boreal lichens from the Altai Mountains (South Siberia, Russia). *Bryologist* 115:61–73
- Davydov EA, Ohmura Y (2016) Lectotypification of the name *Umbilicaria esculenta*. *Mycotaxon* 131:407–412
- Day MJ, Hall JC, Currah RS (2012) Phialide arrangement and character evolution in the helotialean anamorph genera *Cadophora* and *Phialocephala*. *Mycologia* 104:371–378
- Dayarathne MC, Phookamsak R, Ariyawansa HA, Jones EBG et al (2015) Phylogenetic and morphological appraisal of *Leptosphaeria italica* sp. nov. (*Leptosphaeriaceae, Pleosporales*) from Italy. *Mycosphere* 6:634–642
- Dayarathne MC, Maharachchikumbura SSN, Phookamsak R, Fryar S et al (2016a) Morpho-molecular characterization and epitypification of *Annulatascus velatisporus*. *Mycosphere* 7:1389–1398
- Dayarathne MC, Phookamsak R, Hyde KD, Manawasinghe IS et al (2016b) *Halodiatripe*, a novel diatrypaceous genus from mangroves with *H. salinicola* and *H. avicenniae* spp. nov. *Mycosphere* 7:612–627
- de Azevedo CMPS, Gomes RR, Vicente VA, Santos DWCL et al (2015) *Fonsecaea pugnacius*, a novel agent of disseminated chromoblastomycosis. *J Clin Microbiol* 53:2674–2685
- de Beer ZW, Seifert KA, Wingfield MJ (2013) Nomenclator for ophiostomatoid genera and species in the *Ophiostomatales* and *Microasciales*. *The Ophiostomatoid Fungi* 12:245–322
- de Beer ZW, Duong TA, Barnes I, Wingfield BD, Wingfield MJ (2014) Redefining *Ceratocystis* and allied genera. *Stud Mycol* 79:187–219
- de Beer ZW, Duong TA, Wingfield MJ (2016) The divorce of *Sporothrix* and *Ophiostoma*: solution to a problematic relationship. *Stud Mycol* 83:165–191
- de Errasti A, de Beer ZW, Coetzee MPA, Roux J et al (2016) Three new species of *Ophiostomatales* from *Nothofagus* in Patagonia. *Mycol Prog* 15:17
- de García V, Brizzio S, Libkind D, Rosa CA, van Broock M (2010) *Wickerhamomyces patagonicus* sp. nov., an ascomycetous yeast species from Patagonia, Argentina. *Int J Syst Evol Microbiol* 60:1693–1696
- de Gruyter J, Aveskamp MM, Woudenberg JH, Verkley GJ et al (2009) Molecular phylogeny of *Phoma* and allied anamorph

- genera: towards a reclassification of the *Phoma* complex. *Mycol Res* 113:508–519
- de Gruyter J, Woudenberg JH, Aveskamp MM, Verkley GJ et al (2010) Systematic reappraisal of species in *Phoma* section *Paraphoma*, *Pyrenophaeta* and *Pleurophoma*. *Mycologia* 102:1066–1081
- de Gruyter J, Woudenberg JHC, Aveskamp MM, Verkley GJM et al (2012) Redisposition of *Phoma*-like anamorphs in *Pleosporales*. *Stud Mycol* 75:1–36
- de Hoog GS, Vicente VA, Najafzadeh MJ, Harrak MJ et al (2011) Waterborne *Exophiala* species causing disease in cold-blooded animals. *Persoonia* 27:46–72
- de Hoog GS, van Diepeningen AD, Mahgoub E-S, van de Sande WWJ (2012) New species of *Madurella*, causative agents of black-grain mycetoma. *J Clin Microbiol* 50:988–994
- de Hoog GS, Ahmed SA, Najafzadeh MJ, Sutton DA et al (2013) Phylogenetic findings suggest possible new habitat and routes of infection of human eumycetoma. *PLoS Negl Trop Dis* 7:e2229
- de Hoog GS, Dukik K, Monod M, Packeu A et al. (2016a) Toward a novel multilocus phylogenetic taxonomy for the dermatophytes. *Mycopathologia* 1–27
- de Hoog GS, Redhead SA, Feng P, Jiang Y et al (2016b) (2465–2466) Proposals to conserve *Blastomyces* Gilchrist & W.R. Stokes against *Blastomyces* Costantin & Rolland and *Ajellomycetaceae* against *Paracoccidioidaceae* (Ascomycota: Onygenales). *Taxon* 65:1167–1169
- de Hoog GS, Dukik K, Monod M, Packeu A et al (2017) Toward a novel multilocus phylogenetic taxonomy for the dermatophytes. *Mycopathologia* 182(1–2):5–31
- de Jong SN, Levesque CA, Verkley GJM, Abeln ECA et al (2015) Phylogenetic relationships among *Neofabrea* species causing tree cankers and bull's-eye rot of apple based on DNA sequencing of ITS nuclear rDNA, mitochondrial rDNA, and the beta-tubulin gene. *Mycol Res* 105:658–669
- de la Rosa JPM, Porcel MC, Warke PA (2013) Mapping stone surface temperature fluctuations: Implications for lichen distribution and biomodification on historic stone surfaces. *J Cult Herit* 14:346–353
- de Lange PJ (2014) The flora of Egeria Rock, Northern Kermadec Island Group. *Wellington Bot Soc Bull* 55:41–49
- de Lange PJ, Blanchon D (2014) The hunt for *Ramalodium dumosum* (*Pannariaceae*, lichenised ascomycetes)-an enigmatic lichen endemic to the southern Waitakere Ranges
- de Lange PJ, Galloway DJ, Blanchon DJ, Knight A et al (2012) Conservation status of New Zealand lichens. *NZ J Bot* 50:303–363
- de los Ríos A, Raggio J, Pérez-Ortega S, Vivas M et al (2011) Anatomical, morphological and ecophysiological strategies in *Placopsis pycnotheca* (lichenized fungi, Ascomycota) allowing rapid colonization of recently deglaciated soils. *Flora-Morphol Distrib Funct Ecol Plants* 206:857–864
- de Melo Teixeira M, Theodoro RC, de Oliveira FF, Machado GC et al (2014) *Paracoccidioides lutzii* sp. nov. biological and clinical implications. *Med Mycol* 52:19–28
- de Melo MP, da Silva Matos K, Júnior JE et al (2016) *Thielaviopsis musarum* causes postharvest crown and fruit rot of banana in Northeastern Brazil. *Trop Plant Pathol* 41:258–263
- de Moura Nascimento EJ, Van Der Sand ST (2008) Restriction analysis of the amplified ribosomal DNA spacers ITS1 and ITS2 of *Bipolaris sorokiniana* isolates. *World J Microbiol Biotechnol* 24:647–652
- de Paz AG, Cubas P, Divakar PK, Lumbsch HT, Crespo A (2011) Origin and Diversification of major clades in parmeliod lichens (*Parmeliaceae*, Ascomycota) during the Paleogene inferred by Bayesian analysis. *PLoS ONE* 6:e28161
- de Paz GA, Cubas P, Crespo A, Elix JA, Lumbsch HT (2012) Transoceanic dispersal and subsequent diversification on separate continents shaped diversity of the *Xanthoparmelia pulilla* group (Ascomycota). *Plos ONE* 7:e39683
- de Silva H, Castlebury LA, Green S, Stone JK (2009) Characterisation and phylogenetic relationships of *Anisogramma virgultorum* and *A. anomala* in the *Diaporthales* (Ascomycota). *Mycol Res* 113:73–81
- de Silva NI, Phookamsak R, Maharachchikumbura SSN, Thambugala KM et al (2017) *Monochaetia ilexae* sp. nov. (*Pestalotiopsidaceae*) from Yunnan Province in China. *Phytotaxa* 291:123–132
- de Souza Armando EA, Chaves ZM, Dianese JC (2015) *Phaeostilbelloides* and *Velloziomyces*—new dematiaceous genera from the Brazilian Cerrado. *Mycotaxon* 130:757–767
- Debona D, Rodrigues FA, Rios JA, Nascimento KJT (2012) Biochemical changes in the leaves of wheat plants infected by *Pyricularia oryzae*. *Phytopathology* 102:1121–1129
- Declercq B (2009) Omertent *Lasiosphaeria* s.l. *Sterbeekia* 28:35–41
- Delgado G (2008) South Florida microfungi: a new species of *Stanjeughesia* (hyphomycetes) from Sabal palm. *Mycotaxon* 103:229–234
- Delgado G (2013) South Florida microfungi: a new species of *Ellisembia* (hyphomycetes) with new records from the USA. *Mycotaxon* 123:445–450
- Delgado G (2014) South Florida microfungi: *Linkosia longirostrata*, a new hyphomycete on paurotis palm. *Mycotaxon* 129:41–46
- Delgado Baquerizo M, Gallardo A, Covelo F et al (2015) Differences in thallus chemistry are related to species-specific effects of biocrust forming lichens on soil nutrients and microbial communities. *Funct Ecol* 29:1087–1098
- Dellavalle PD, Cabrera A, Alem D, Larrañaga P et al (2011) Antifungal activity of medicinal plant extracts against phytopathogenic fungus *Alternaria* spp. *Chil J Agric Res* 71:231
- Del-Prado R, Divakar PK, Crespo A (2011) Using genetic distances in addition to ITS molecular phylogeny to identify potential species in the *Parmotrema reticulatum* complex: a case study. *Lichenologist* 43:569–583
- Del-Prado R, Blanco O, Lumbsch HT, Divakar PK et al (2013) Molecular phylogeny and historical biogeography of the lichen-forming fungal genus *Flavoparmelia* (Ascomycota: Parmeliaceae). *Taxon* 62:928–939
- Demiray AD, Yolcubal I, Akyol NH, Çobanoğlu G (2012) Biomonitoring of airborne metals using the lichen *Xanthoria parietina* in Kocaeli Province, Turkey. *Ecol Ind* 18:632–643
- Desoubeaux G, García D, Baily E, Augereau O et al (2014) Subcutaneous phaeohyphomycosis due to *Phialemoniopsis ocularis* successfully treated by voriconazole. *Med Mycol Case Rep* 5:4–8
- Dethoup T, Manoch L (2009) Diversity of marine fungi in eastern Thailand. *Kasetsart J Nat Sci* 43:100–106
- Devi TP, Mathur N, Kamil D, Prabhakaran N (2012) *Proliferosphaera*, a new genus from India. *Indian Phytopathol* 65:395–398
- Dey P, Banerjee J, Maiti MK (2011) Comparative lipid profiling of two endophytic fungal isolates—*Colletotrichum* sp. and *Alternaria* sp. having potential utilities as biodiesel feedstock. *Biores Technol* 102:5815–5823
- Dhargalkar S, Bhat DJ (2009) *Echinospaera pteridis* sp. nov. and its *Vermiculariopsiella* anamorph. *Mycotaxon* 108:115–122
- Dharkar N, Subhedar A, Hande D, Shahezad MA (2010) Two new fungal species from Vidarbha, India. *J Mycol Plant Pathol* 40:235–237
- Díaz GA, Zoffoli JP, Lolas M, Blanco A et al (2015) Occurrence of *Phacidopycnis washingtonensis* causing speck rot on stored pink lady apple fruit in Chile. *Plant Dis* 100:211

- Dib-Bellahouel S, Fortas Z (2011) Antibacterial activity of various fractions of ethyl acetate extract from the desert truffle, *Tirmania pinoyi*, preliminarily analyzed by gas chromatography–mass spectrometry (GC–MS). *Afr J Biotechnol* 10:9694–9699
- Diederich PA (2013) Redescription of *Sporodochiolichen flavus* (lichenized sporodochial Ascomycetes). *Bull Soc Nat Luxemb* 114:55
- Diederich P, Schultz M (2009) *Zwackhiomyces namibiensis*, a new lichenicolous ascomycete (*Xanthopyreniaceae*) on *Psorotrichia* from Namibia. *Herzogia* 22:73–176
- Diederich P, Palice Z, Ertz D (2008) *Cheiromycina ananas* is a synonym of *Dictyocatenulata alba*, a widespread, lichenized, synnematosus hyphomycete herewith reported as new for Europe. *Sauteria* 15:205–214
- Diederich P, Ertz D, Van den Broeck D, van den Boom P et al (2009) New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. XII. *Bull Soc Nat Luxemb* 110:75–92
- Diederich P, Braun U, Heuchert B, Ertz D (2010a) Four new lichen-associated *Trimmatostroma* species (hyphomycetes). *Bull Soc Nat Luxemb* 111:47–55
- Dietrich M, Buergi-Meyer K (2010) Interesting saxicolous lichens at Tossen near Schimbrigbad in the UNESCO Biosphere Entlebuch (Canton of Lucerne, Central Switzerland). *Herzogia* 23:75–84
- Diederich P, Ertz D, Etayo J (2010b) An enlarged concept of *Llimoniella* (lichenicolous Helotiales), with a revised key to the species and notes on related genera. *Lichenologist* 42:253–269
- Diederich P, Ertz D, Eichler M, Cezanne R et al (2012a) New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. XIV. *Bull Soc Nat Luxemb* 113:95–115
- Diederich P, Lawrey JD, Sikaroodi M, van den Boom P, Ertz D (2012b) *Briancoppinsia*, a new coelomycetous genus of *Arthoniaceae* (*Arthoniales*) for the lichenicolous *Phoma cytospora*, with a key to this and similar taxa. *Fungal Divers* 52:1–12
- Diederich P, Ertz D, Lawrey JD, Sikaroodi M, Untereiner WA (2013) Molecular data place the hyphomycetous lichenicolous genus *Sclerococcum* close to *Dactylospora* (*Eurotiomycetes*) and *S. parmeliae* in *Cladophialophora* (*Chaetothyriales*). *Fungal Divers* 58:61–72
- Diederich P, Ertz D, Eichler M, Cezanne R et al (2014) New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. XV. *Bull Soc Nat Luxemb* 115:157–165
- Diene O, Wang W, Narisawa K (2013) *Pseudosigmoidea ibarakiensis* sp. nov., a dark septate endophytic fungus from a cedar forest in Ibaraki, Japan. *Microbes Environ* 28:381–387
- Dillman KL, Ahti T, Björk CR, Clerc P et al (2012) New records, range extensions and nomenclatural innovations for lichens and lichenicolous fungi from Alaska, USA. *Herzogia* 25(2):177–210
- Din LB, Zakaria Z, Samsudin MW, Elix JA (2010) Chemical profile of compounds from lichens of Bukit Larut, Peninsular Malaysia. *Sains Malays* 39:901–908
- Ding T, Jiang T, Zhou J, Xu L, Gao ZM (2010) Evaluation of antimicrobial activity of endophytic fungi from *Camptotheca acuminata* (Nyssaceae). *Genet Mol Res* 9:2104–2112
- Dissanayake AJ, Jayawardena RS, Boonmee S, Thambugala KM et al (2014) The status of *Myriangiaceae* (*Dothideomycetes*). *Phytotaxa* 176:219–237
- Dissanayake AJ, Liu M, Zhang W, Chen Z et al (2015) Morphological and molecular characterisation of *Diaporthe* species associated with grapevine trunk disease in China. *Fungal Biol* 19:283–294
- Dissanayake AJ, Camporesi E, Hyde KD, Phillips AJL et al (2016a) *Dothiorella* species associated with woody hosts in Italy. *Mycosphere* 7:51–63
- Dissanayake AJ, Phillips AJL, Li XH, Hyde KD (2016b) *Botryosphaeriaceae*: current status of genera and species. *Mycosphere* 7:1001–1073
- Dissanayake AJ, Camporesi E, Hyde KD, Zhang W et al (2017) Molecular phylogenetic analysis reveals seven new *Diaporthe* species from Italy. *Mycosphere* 8:853–877
- Divakar PK, Lumbsch HT, Ferencova Z, Del Prado R, Crespo A (2010) *Remototrichyna*, a newly recognized tropical lineage of lichens in the *Hypotrichyna* clade (*Parmeliaceae, Ascomycota*), originated in the Indian subcontinent. *Am J Bot* 97:579–590
- Divakar PK, Del-Prado R, Lumbsch HT, Wedin M et al (2012) Diversification of the newly recognized lichen-forming fungal lineage *Montanelia* (*Parmeliaceae, Ascomycota*) and its relation to key geological and climatic events. *Am J Bot* 99:2014–2026
- Divakar PK, Crespo A, Núñez-Zapata Flakus A et al (2013a) A molecular perspective on generic concepts in the *Hypotrichyna* clade (*Parmeliaceae, Ascomycota*). *Phytotaxa* 132:21–38
- Divakar PK, Kauff F, Crespo A, Leavitt SD, Lumbsch HT (2013b) Understanding phenotypical character evolution in parmeloid lichenized fungi (*Parmeliaceae, Ascomycota*). *PLoS ONE* 8:e83115
- Divakar PK, Crespo A, Wedin M, Leavitt SD et al (2015) Evolution of complex symbiotic relationships in a morphologically derived family of lichen-forming fungi. *New Phytol* 208:1217–1226
- Divakar PK, Crespo A, Kraichak E, Leavitt SD et al (2017) Using a temporal phylogenetic method to harmonize family-and genus-level classification in the largest clade of lichen-forming fungi. *Fungal Divers* 84:101–117
- Dixon AP, Faber-Langendoen D, Josse C, Morrison J, Loucks CJ (2014) Distribution mapping of world grassland types. *J Biogeogr* 41:2003–2019
- Döbbeler P (2010) New species and records of *Bryocentria*—a hypocrealean genus of bryophilous ascomycetes. *Karstenia* 50:11–23
- Döbbeler P (2011) Two new hepaticolous species of the genus *Octosporella* (*Pezizales*). *Herzogia* 24:357–365
- Döbbeler P (2012) *Gynonectria intraspora* gen. et sp. nov. (*Hypocreales*), a perianthicolous phyllosphere fungus with a novel ascospore type. *Mycol Prog* 11:473–481
- Doering J, Deduke C, Piercy-Normore MD (2016) Variation in lichen species assemblages and secondary metabolites surrounding *Stereocaulon* species in the boreal forest of Northwestern Manitoba. *Evansia* 33:70–82
- Doherty TA, Khorram N, Chang JE, Kim HK et al (2012) STAT6 regulates natural helper cell proliferation during lung inflammation initiated by *Alternaria*. *Am J Physiol* 303:L577–L588
- Doilom M, Liu JK, Jaklitsch WM, Ariyawansa H et al (2013) An outline of the family *Cucurbitariaceae*. *Sydowia* 65:167–192
- Doilom M, Shuttleworth LA, Roux J, Chukeatirote E et al (2014) *Barriopsis tectonae* sp. nov. a new species of *Botryosphaeriaceae* from *Tectona grandis* (teak) in Thailand. *Phytotaxa* 176:81–91
- Doilom M, Taylor JE, Bhat DJ, Chukeatirote E et al (2016) Checklist of fungi on teak. *Mycosphere* 7:656–678
- Doilom M, Dissanayake AJ, Wanasinghe AN, Boonmee S et al (2017) Microfungi on *Tectona grandis* (teak) in northern Thailand. *Fungal Divers* 82:107–182
- Dong N, Liu X, Lu Y, Du L et al (2010) Overexpression of TaPIEP1, a pathogen-induced ERF gene of wheat, confers host-enhanced resistance to fungal pathogen *Bipolaris sorokiniana*. *Funct Integr Genomics* 10:215–226
- Dong L, Park JS, Woo JJ (2015) A-34: contribution to the Lichen Flora of Vietnam, with ten new record species. *교학회소식* 27:62

- Dornelo-Silva D, Pereira-Carvalho RC, Dianese JC (2008) New *Stenella* and *Parastenella* species from the Brazilian cerrado. *Mycologia* 99:753–764
- Dougoud R (2012) *Aeruginoscyphus*, un nouveau genre pour *Peziza sericea* (*Helotiales*, *Hyaloscyphaceae*). Ascomycete 4:1–4
- Dougoud R (2015) *Rutstroemia alnobetulae* sp. nov. (*Helotiales*, *Rutstroemiaceae*), une espèce nouvelle des aulnes verts. Ascomycete 7:336–340
- Dougoud R, Van Vooren N, Vega M (2015) *Cupulina montana* (*Pezizales*, *Pyronemataceae*), un nouveau genre et une nouvelle espèce des montagnes d'Europe. Ascomycete 7:39–44
- Doveri F (2008) Pagine di micologia. Associazione Micologica Bresadola Centro Studi Micologici 65–66:110
- Doveri F (2010) Une nouvelle variété de *Tripterospora* sur crottin une occasion de recombiner le pyrenomycète cleistothelial *Cercophora heterospora* dans le genre *Tripterospora*. Bull Mycol Bot Dauphiné-Savoie 50:49–55
- Doveri F (2011) Addition to “*Fungi Fimicoli Italici*”: An update on the occurrence of coprophilous Basidiomycetes and Ascomycetes in Italy with new records and descriptions. *Mycosphere* 2:331–427
- Doveri F (2012) Coprophilous discomycetes from the *Tuscan archipelago* (Italy). Description of two rare species and a new *Trichobolus*. *Mycosphere* 3:503–522
- Doweld G (2013) Nomenclatural novelties. Index Fungorum 31:1
- Doveri F (2014) An update on the genera *Ascobolus* and *Saccobolus* with keys and descriptions of three Coprophilous species, new to Italy. *Mycosphere* 5:86–135
- Doveri F, Sarrocco S (2013) *Sporormiella octomegaspora*, a new hairy species with eight-celled ascospores from Spain. *Mycotaxon* 123:129–140
- Doveri F, Peccia S, Sarrocco S, Minnoci A, Vannacci G (2010) *Rodentomyces*, a new hypocrealean genus from Italy. *Fungal Divers* 42:57–69
- Doveri F, Sarrocco S, Vannacci G (2013) Studies on three rare coprophilous plectomycetes from Italy. *Mycotaxon* 124:279–300
- Doyon JB, Sutton DA, Theodore P, Dhillon G et al (2013) *Rasamonia argillacea* pulmonary and aortic graft infection in an immune-competent patient. *J Clin Microbiol* 51:719–722
- Dreaden TJ, Davis JM, De Beer ZW, Ploetz RC et al (2014a) Phylogeny of ambrosia beetle symbionts in the genus *Raffaelea*. *Fungal Biol* 118:970–978
- Dreaden TJ, Davis JM, Harmon CL, Ploetz RC et al (2014b) Development of multilocus PCR assays for *Raffaelea lauricola*, causal agent of laurel wilt disease. *Plant Dis* 98:379–383
- D’Souza MA, Bhat DJ (2012) A new species of *Pleurophragmium* from India. *Mycotaxon* 119:477–482
- Du YQ, Liu Y, Wang JH (2015) Polysaccharides from *Umbilicaria esculenta* cultivated in Huangshan Mountain and immunomodulatory activity. *Int J Biol Macromol* 72:1272–1276
- Du Z, Fan XL, Hyde KD, Yang Q et al (2016) Phylogeny and morphology reveal two new species of *Diaporthe* from *Betula* spp. in China. *Phytotaxa* 269:90–102
- Duarte S, Seena S, Barlocher F, Cassio F, Pascoal C (2012) Preliminary insights into the phylogeography of six aquatic hyphomycete species. *PLoS ONE* 7:E45289
- Duarte S, Batista D, Bärlocher F, Cássio F, Pascoal C (2015) Some new DNA barcodes of aquatic hyphomycete species. *Mycoscience* 56:102–108
- Dubey R, Moonambeth AM (2013) *Custingophora ratnagiriensis* sp.nov.—a novel species of *Custingophora* from Konkan, Maharashtra, India. *NEBIO Int J Environ Biodivers* 4:31–32
- Dubey R, Neelima A (2014) *Goosiomycetes bambusicola*—a new cheiroporous anamorphic species from Western Ghats, India. *Curr Res Environ Appl Mycol* 4:211–216
- Dubey R, Moonambeth AM (2014a) *Sheathnema indicum* gen. et sp. nov. a new sooty mold fungus from northern Western Ghats, India. *J Threat Taxa* 6:6549–6555
- Dubey R, Moonambeth AM (2014b) *Vermiculariopsiella papaye* sp. nov.—a new species of *Vermiculasporiella* from Western Ghats, India. *Int J Sci Res* 3:35–36
- Dubey R, Pandey AK (2011) *Bispora aeglei* sp. nov.—a new follicolous hyphomycete from India. *J Mycopathol Res* 49:357–359
- Dubey R, Pandey AK (2012) *Pseudoacrodictys steviae*—a new generic and species record for India. *J Mycol Pl Pathol* 42:251–253
- Dubey U, Upadhyay DK, Nayaka S (2010) A new species of *Leiorreuma* Eschw. from India. *Lichenologist* 42:711–713
- Dubey R, Pandey AK, Manoharachary C (2011a) *Beltramono*—a hitherto undescribed Hyphomycetes genus. *J Mycopathol Res* 49:321–324
- Dubey RK, Rai AN, Shrivastava S, Verma NK (2011b) Two new records of hyphomycetes from India. *Indian Phytopathol* 64:390–391
- Duong TA, de Beer ZW, Wingfield BD, Wingfield MJ (2012) Phylogeny and taxonomy of species in the *Grosmannia serpens* complex. *Mycosphere* 104:715–732
- Dupont J, Magnin S, Rousseau F, Zbinden M et al (2009) Molecular and ultrastructural characterization of two ascomycetes found on sunken wood off Vanuatu Islands in the deep Pacific Ocean. *Mycol Res* 112:1351–1364
- Durkin SR, Henderson T, Raju R, Ellis D (2008) Successful treatment of phaeohyphomycotic keratitis caused by *Bipolaris australiensis*. *Clin Exp Ophthalmol* 36:697–699
- Dyer ZA, Wright RS, Rong IH, Jacobs A (2008) Back pain associated with endobronchial mucus impaction due to *Bipolaris australiensis* colonization representing atypical allergic bronchopulmonary mycosis. *Med Mycol* 46:589–594
- Dymytrova LV, Breuss O, Kondratyuk SY (2011) *Agonimia borysthenica*, a new lichen species (*Verrucariales*) from Ukraine. *Österr Z Pilzkd* 20:25–28
- Dymytrova L, Nadyrina O, Naumovych A, Keller C, Scheidegger C (2013) Primeval beech forests of Ukrainian Carpathians are sanctuaries for rare and endangered epiphytic lichens. *Herzogia* 26:73–89
- Ebead GA, Overy DP, Berrue F, Kerr RG (2012) *Westerdykella reniformis* sp. nov., producing the antibiotic metabolites melinacidin IV and chetracin B. *IMA Fungus* 3:189–201
- Edathodu J, Al-Abdely HM, AlThawadi S, Wickes BL et al (2013) Invasive fungal infection due to *Triadelphia pulvinata* in a patient with acute myeloid leukemia. *J Clin Microbiol* 51:3426–3429
- Egan RS (2009) Recent literature on lichens-212. *Bryologist* 112:239–256
- Eghrari AO, Gibas C, Watkins T, Vahedi S et al (2015) First human case of fungal keratitis caused by a putatively novel species of *Lophotrichus*. *J Clin Microbiol* 53:3063–3067
- Eghtedari M (2015) Asymptomatic fungal cyst of conjunctiva caused by *Bipolaris spicifera*. *Iran J Med Sci* 31:56–58
- Egidi E, De Hoog GS, Isola D, Onofri S et al (2014) Phylogeny and taxonomy of meristematic rock-inhabiting black fungi in the *Dothideomycetes* based on multi-locus phylogenies. *Fungal Divers* 65:127–165
- Eichler M, Cezanne R, Diederich P, Ertz D et al (2010) New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. XIII. *Bull Soc Nat Luxemb* 111:33–46
- Eisa M, Chand R, Joshi AK (2013) Biochemical and histochemical traits: a promising way to screen resistance against spot blotch

- (*Bipolaris sorokiniana*) of wheat. Eur J Plant Pathol 137:805–820
- Ekanayaka AH, Daranagama DA, Ariyawansa HA, Jones EBG et al (2016a) *Pezicula chiangraiensis* sp. nov. from Thailand. Mycotaxon 131:739–748
- Ekanayaka AH, Dissanayake AJ, Jayasiri SC, To-anun C et al (2016b) *Aplosporella thailandica*; a novel species revealing the sexual-aseexual connection in *Aplosporellaceae* (*Botryosphaerales*). Mycosphere 7:440–447
- Ekanayaka AH, Hyde KD, Zhao Q (2016c) The genus *Cookeina*. Mycosphere 7:1399–1413
- Ekanayaka AH, Zhao Q, Jones EBG, Pu ED et al (2016d) Two novel *Acervus* species extend their distribution within Yunnan, China. Phytotaxa 283:74–82
- Ekman S, Andersen HL, Wedin M (2008) The limitations of ancestral state reconstruction and the evolution of the ascus in the *Lecanorales* (lichenized *Ascomycota*). Syst Biol 57:141–156
- Ekman S, Blaalid R (2011) The devil in the details: interactions between the branch-length prior and likelihood model affect node support and branch lengths in the phylogeny of the *Psoraceae*. Syst Biol 60:541–561
- Ekman S, Svensson M (2014) *Brianaria* (*Psoraceae*), a new genus to accommodate the *Micarea sylvicola* group. Lichenologist 46:285–294
- Ekman S, Wedin M, Lindblom L, Jørgensen PM (2014) Extended phylogeny and a revised generic classification of the *Pannariaceae* (*Peltigerales*, *Ascomycota*). Lichenologist 46:627–656
- El Mhadri M, Benkirane R, Touhami AO, Douira A (2009) *Citrullus lanatus*, a new host of *Bipolaris spicifera* in Morocco. Phytopathol Meditarr 48:291–293
- Elix JA (2009) New saxicolous species and new records of *Buellia* sens. lat. and *Rinodinella* (*Ascomycota*, *Physciaceae*) in Australia. Australas Lichenol 65:10–19
- Elix JA (2012) Four new species and a new record of *Tephromela* (lichenized *Ascomycota*) from Australia. Australas Lichenol 71:3–11
- Elix JA (2013) New crustose lichen taxa (lichenized *Ascomycota*) from Australia. Australas Lichenol 73:45–54
- Elix JA, Kantvilas G (2009) Further new species of *Xanthoparmelia* (*Parmeliaceae*, *Ascomycota*) from Tasmania. Biblioth Lichenol 99:123–132
- Elix JA, Kantvilas G (2013) New taxa and new records of *Amandinea* (*Physciaceae*, *Ascomycota*) in Australia. Australas Lichenol 72:3–19
- Elix JA, Kantvilas G (2014) New species and new records of the lichen genus *Baculifera* (*Physciaceae*, *Ascomycota*) in Australia. Australas Lichenol 75:28–37
- Elix JA, Kantvilas G (2015) New taxa and new records of crustose lichens in the family *Physciaceae* (*Ascomycota*) in Australia. Australas Lichenol 76:16–23
- Elix JA, Kondratyuk SY (2008) Two new species of *Letrouitia* (*Letrouitiaceae*, *Ascomycota*) from Australia. Australas Lichenol 62:16–19
- Elix JA, Mayrhofer H (2016) Two new species of *Buellia* sens. lat. (*Ascomycota*, *Physciaceae*) from New Zealand with 1-septate ascospores. Australas Lichenol 79:10–15
- Elliott ML, Des Jardin EA (2014) *Serenomyces* associated with palms in southeastern USA: isolation, culture storage and genetic variation. Mycologia 106:698–707
- Elliott TF, Bouger NL, O'Donnell K, Trappe JM (2014) *Morchella australiana* sp. nov., an apparent Australian endemic from New South Wales and Victoria. Mycologia 106:113–118
- El-Said AH, Sohair TH, El-Hadi AG (2010) Studies on the mycoflora associated with sugarcane factory waste and pollution of River Nile in upper Egypt. J Yeast Fungal Res 1:188–196
- Elvebak A (2012) *Pannaria howeana* and *Pannaria streimannii*, two related new lichen species endemic to Lord Howe Island, Australia. Lichenologist 44:457–463
- Elvebak A, Robertsen EH, Park CH, Hong SG (2010) *Psorophorus* and *Xanthopsoroma*, two new genera for yellow-green, corticolous and squamulose lichen species, previously in *Psoroma*. Lichenologist (Lond) 42:563–585
- Elvebak A, Hong SG, Park CH, Robertsen EH et al (2016) *Gibbosporina*, a new genus for foliose and tripartite, Palaeotropi. *Pannariaceae* species previously assigned to *Psoroma*. Lichenol 48(1):13–52
- Endoh R, Suzuki M, Benno Y (2008) *Ambrosiozyma kamigamensis* sp. nov. and *A. neoplatypodis* sp. nov., two new ascomycetous yeasts from ambrosia beetle galleries. Antonie Van Leeuwenhoek 94:365–376
- Enghoff H, Santamaría S (2015) Infectious intimacy and contaminated caves—three new species of ectoparasitic fungi (*Ascomycota*: *Laboulbeniales*) from blaniulid millipedes (Diplopoda: Julida) and inferences about their transmittal mechanisms. Org Divers Evol 15:249–263
- Ertz D, Diederich P (2015) Dismantling *Melaspileaceae*: a first phylogenetic study of *Buellia*, *Hemigrapha*, *Karschia*, *Labrocarpon* and *Melaspilea*. Fungal Divers 71:41–164
- Ertz D, Tehler A (2011) The phylogeny of *Arthoniales* (*Pezizomycotina*) inferred from nucLSU and RPB2 sequences. Fungal Divers 49:47–71
- Ertz D, Diederich P, Brand AM, van den Boom P, Séruviaux E (2008) New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. XI. Bull Soc Nat Luxemb 109:35–51
- Ertz D, Miadlikowska J, Lutzoni F, Dessein S et al (2009) Towards a new classification of the *Arthoniales* (*Ascomycota*) based on a three-gene phylogeny focussing on the genus *Opegrapha*. Mycol Res 113:141–152
- Ertz D, Killmann D, Razafindrahaja T, Séruviaux E, Fischer E (2010) Two new species of *Syncesia* (*Arthoniales*, *Roccellaceae*) from Africa. Lichenologist 42:43–49
- Ertz D, Bungartz F, Diederich P, Tibell L (2011) Molecular and morphological data place *Blarneya* in *Tylophoron* (*Arthoniaceae*). Lichenologist 43:345–356
- Ertz D, Fischer E, Killmann D, Razafindrahaja T, Séruviaux E (2013) *Savoronala*, a new genus of *Malmideaceae* (*Lecanorales*) from Madagascar with stipes producing sporodochia. Mycol Prog 12:645–656
- Ertz D, Aptroot A, Van de Vijver B, Sliwa L et al (2014a) Lichens from the Utsteinen Nunatak (Sør Rondane Mountains, Antarctica), with the description of one new species and the establishment of permanent plots. Phytotaxa 191:99–114
- Ertz D, Lawrey JD, Common RS, Diederich P (2014b) Molecular data resolve a new order of *Arthoniomycetes* sister to the primarily lichenized *Arthoniales* and composed of black yeasts, lichenicolous and rock-inhabiting species. Fungal Divers 66:113–137
- Ertz D, Tehler A, Fischer E, Killmann D et al (2014c) *Isalonactis*, a new genus of *Roccellaceae* (*Arthoniales*), from southern Madagascar. Lichenologist 46:159–167
- Ertz D, Diederich P, Lawrey JD, Berger F et al (2015a) Phylogenetic insights resolve *Dacampiaceae* (*Pleosporales*) as polyphyletic: *Didymocyrtis* (*Pleosporales*, *Phaeosphaeriaceae*) with Phoma-like anamorphs resurrected and segregated from *Polycoccum* (*Trypethiales*, *Polycoccaceae* fam. nov.). Fungal Divers 74:53–89
- Ertz D, Flakus A, Oset M, Sipman HJ, Kukwa M (2015b) A first assessment of lichenized *Arthoniales* in Bolivia with descriptions of two new species. Phytotaxa 217:1–25

- Ertz D, Tehler A, Irestedt M, Frisch A et al (2015c) A large-scale phylogenetic revision of *Roccellaceae* (*Arthoniales*) reveals eight new genera. *Fungal Divers* 70:31–53
- Ertz D, Fryday A, Schmitt I, Charrier M et al (2016a) *Ochrolechia kerguelensis* sp. nov. from the Southern Hemisphere and *O. antarctica* reinstated from the synonymy of *O. parella*. *Phytotaxa* 280:129–140
- Ertz D, Heuchert B, Braun U, Freebury CE et al (2016b) Contribution to the phylogeny and taxonomy of the genus *Taeniopeltis*, with a focus on lichenicolous taxa. *Fungal Biol* 120:1416–1447
- Escuredo O, Seijo MC, Fernández-González M, Iglesias I (2011) Effects of meteorological factors on the levels of *Alternaria* spores on a potato crop. *Int J Biometeorol* 55:243–252
- Esslinger TL, Dillman KL (2010) *Physconia grumosa* in North America. *Bryologist* 113:77–80
- Esslinger TL, Morse CA, Leavitt SD (2012) A new North American species of *Hyperphyscia* (*Physciaceae*). *Bryologist* 115:31–41
- Etayo J (2010a) Hongos liquenícolas de Perú. Homenaje a Rolf Santesson. *Bull Soc Linn Provence* 61:83–128
- Etayo J (2010b) Euskal Herriko Unibertsitatea/Universidad del País Vasco Guineana 16:501
- Etayo J (2010c) Lichenicolous fungi from the western Pyrenees. V. Three new ascomycetes. *Opuscula Phil* 8:131–139
- Etayo J (2011) *Lichenochora hyperphysiae*, a new lichenicolous fungus found in Mexico and Spain. *Biblioth Lichenol* 106:54
- Etayo JA, Berger FR (2009) *Carnegieispora rimeliae*, a new genus of lichenicolous fungus from the Azores. *Opuscula Phil* 7:17–20
- Etayo J, Fr Berger (2013) *Isthmoconidium bacidiicola*, a new genus and species of lichenicolous hyphomycetes on Bacidia from Bermuda. *Opuscula Phil* 13:1–5
- Etayo J, Navarro-Rosines P (2008) A combination and three new species of *Lichenochora* (*Phyllachorales*, lichenicolous *Ascomycota*), and additional reports for the genus. *Revta Catal Micol* 30:27–44
- Etayo J, Sancho LG (2008) Hongos liquenícolas del sur de Sudamérica, especialmente de Isla Navarino (Chile). *Biblioth Lichenol* 98:1–302
- Etayo J, Sharuddin SS (2011) *Verruculina* and *Licheno verruculina* nom. nov. *Herzogia* 24:379–380
- Etayo J, Triebel D (2010) New and interesting lichenicolous fungi at the Botanische Staatssammlung München. *Lichenologist* 42:231–240
- Etayo J, Yazici K (2009) *Microsphaeropsis caloplacae* sp. nov. on *Caloplaca persica* in Turkey. *Mycotaxon* 107:297–302
- Etayo JA, van den Boom PI (2013) A first checklist of lichenicolous fungi from the Dominican Republic, including the description of a new species of *Xenonectriella*. *Opuscula Phil* 12:142–150
- Etayo J, Flakus A, Kukwa M (2013a) *Niesslia echinoides* (*Niessliaceae*, *Ascomycota*), a new lichenicolous fungus on Erioderma from Bolivia. *Lichenologist* 45:21–24
- Etayo J, Flakus A, Kukwa M, Rodriguez Flakus P (2013b) *Lichenochora tertia* (*Phyllachorales*): the third species of the genus growing on *Xanthoria elegans*. *Mycotaxon* 123:9–13
- Etayo J, Flakus A, Suija A, Kukwa M (2015) *Macroskyttea parmotrematis* gen. et sp. nov. (*Helotiaceae*, *Leotiomycetes*, *Ascomycota*), a new lichenicolous fungus from Bolivia. *Phytotaxa* 224:247–257
- Evans HC, Elliot S, Hughes DP (2011) Hidden diversity behind the zombie-ant fungus *Ophiocordyceps unilateralis*: four new species described from carpenter ants in Minas Gerais, Brazil. *PLoS ONE* 6:e17024
- Evans HC, Johnston PR, Park D, Barreto RW, Soares DR (2010) *Claviradulomyces*, a new genus of *Odontotremataceae* from West African rainforest. *Fungal Biol* 114:41–48
- Evidente A, Punzo B, Andolfi A, Berestetskiy A, Motta A (2009) Alternethanolins A and B, polycyclic ethanones produced by *Alternaria sonchi*, potential mycoherbicides for Sonchus arvensis biocontrol. *J Agric Food Chem* 57:6656–6660
- Fahala AH, Rahmana IA, El-Hassana AM, Abdel ME et al (2010) The safety and efficacy of itraconazole for the treatment of patients with eumycetoma due to *Madurella mycetomatis*. *Trans R Soc Trop Med Hyg* 105:127–132
- Falbo MK, Soccol VT, Sandini IE, Vicente VA et al (2013) Isolation and characterization of the nematophagous fungus *Arthrobotrys conoides*. *Parasitol Res* 112:177–185
- Fan L, Hou C, Cannon PF, Li Y (2012a) A new species of *Melanospora* on truffles from China. *Mycologia* 104:1433–1442
- Fan L, Lin YR, Wang S, Hou CL (2012b) Two species of *Lirula* on *Abies* from Yunnan, southwest China. *Mycol Prog* 11:279–286
- Fan L, Cao JZ, Hou CL (2013) *Tuber subglobosum* and *T. wenchuanense*—two new species with spino-reticulate ascospores. *Mycotaxon* 123:95–101
- Fan XL, Liang YM, Ma R, Tian CM (2014) Morphological and phylogenetic studies of *Cytospora* (*Valsaceae*, *Diaporthales*) isolates from Chinese scholar tree, with description of a new species. *Mycoscience* 55:252–259
- Fan XL, Hyde KD, Yang Q, Liang Y et al (2015a) *Cytospora* species associated with canker disease of three anti-desertification plants in northwestern China. *Phytotaxa* 197:227–244
- Fan XL, Hyde KD, Udayanga D, Wu XY et al (2015b) *Diaporthe rostrata*, a novel ascomycete from *Juglans mandshurica* associated with walnut dieback. *Mycol Prog* 14:82–90
- Fan XL, Du Z, Hyde KD, Liang YM et al (2016a) *Cryptosporella platyphylla*, a new species associated with *Betula platyphylla* in China. *Phytotaxa* 253:285–292
- Fan XL, Du Z, Liang Y, Tian C (2016b) *Melanconis* (*Melanconidaceae*) associated with *Betula* spp. in China. *Mycol Prog* 15:40–49
- Fangyan Z, Danyang Q, Changsheng L (2013) Studies on control effects of different density of 5 fungicides against *Agaricodochium camellia*. *Guangdong For Sci Technol* 2:005
- Farkas E (2010) Notes and schedae to Lichenes delicati exsiccati editae in memoriam Antonín Vezda (1920–2008), fasc. 1. *Acta Bot Hung* 52:331–340
- Farkas E (2014) Bibliography of foliicolous lichenized fungi, 1952–2013. *Acta Bot Hung* 56:33–68
- Farkas E (2015) Names of Bacidia s. l. in current use for foliicolous lichens—an annotated nomenclatural study. *Acta Bot Hung* 57:51–70
- Farkas E, Flakus A (2015) Rare or Overlooked? Two Species of *Lyromma* (*Lyrommataceae*, Lichenized *Ascomycota*) are New for Africa. *Herzogia* 28:204–211
- Farkas E, Flakus A (2016) *Trichonectria calopadiicola* sp. nov. (*Hypocreales*, *Ascomycota*): the second species of the family Bionectriaceae parasitic on foliicolous lichens discovered in Tanzania. *Phytotaxa* 278:281–286
- Farkas E, Veres K, Czúcz B (2011) Statistical analysis of substrate investigations of tericolous *Toninia* species in Hungary. *Acta Bot Hung* 53:313–320
- Fatima F, Bajpai P, Pathak N, Singh S et al (2015) Antimicrobial and immunomodulatory efficacy of extracellularly synthesized silver and gold nanoparticles by a novel phosphate solubilizing fungus *Bipolaris tetramera*. *BMC Microbiol* 15:1
- Favero-Longo SE, Piervittori R (2012) Cultivation of isidia and transplantation of adult thalli of *Xanthoparmelia tinctina* in an abandoned asbestos mine. *Lichenologist* 44:840–844
- Favero-Longo SE, Matteucci E, Morando M, Rolfo F et al (2015) Metals and secondary metabolites in saxicolous lichen communities on ultramafic and non-ultramafic rocks of the western Italian Alps. *Aust J Bot* 63:276–291

- Fawzi EM, Khalil AA, Afifi AF (2009) Antifungal effect of some plant extracts on *Alternaria alternata* and *Fusarium oxysporum*. Afr J Biotechnol 8:2598–2603
- Fedorenko N, Stenroos S, Thell A, Kärnefelt I, Kondratyuk S (2009) A phylogenetic analysis of xanthorioid lichens (*Teloschistaceae*, *Ascomycota*) based on ITS and mtSSU sequences. Biblioth Lichenol 100:49–84
- Fedorenko NM, Stenroos S, Thell A, Kärnefelt I et al (2012) Molecular phylogeny of xanthorioid lichens (*Teloschistaceae*, *Ascomycota*), with notes on their morphology. Systematics, biodiversity and ecology of lichens. Biblioth Lichenol 108:45–64
- Fedosova AG, Kovalenko AE (2015) Studies on the Geoglossoid fungi of Russia: the genus *Leucoglossum*. Mycol Prog 14:26
- Felle HH, Herrmann A, Schäfer P, Hückelhoven R, Kogel KH (2008) Interactive signal transfer between host and pathogen during successful infection of barley leaves by *Blumeria graminis* and *Bipolaris sorokiniana*. J Plant Physiol 165:52–59
- Feng PY, de Hoog GS, Najafzadeh MJ, Gerrits van den Ende AHG et al (2013) *Cladophialophora abundans*, a novel species of *Chaetothyriales* isolated from the natural environment. Mycol Prog 13:381–391
- Feng PY, Lu Q, Najafzadeh MJ, van den Ende AG et al (2014) *Cyphellophora* and its relatives in *Phialophora*: biodiversity and possible role in human infection. Fungal Divers 65:17–45
- Ferencova Z, Cubas P, Divakar PK, Molina MC, Crespo A (2014) *Notoparmelia*, a new genus of *Parmeliaceae* (*Ascomycota*) based on overlooked reproductive anatomical features, phylogeny and distribution pattern. Lichenologist 46:51–67
- Fernandes MDRV, Pfenning LH, Costa-Neto CMD, Heinrich TA et al (2009) Biological activities of the fermentation extract of the endophytic fungus *Alternaria alternata* isolated from *Coffea arabica* L. Braz J Pharm Sci 45:677–685
- Fernández-Brime S, Llimona X, Navarro-Rosinés P (2010) *Lichenostigma rupicolae* (*Lichenotheliaceae*), a new lichenicolous species growing on *Pertusaria rupicola*. Lichenologist 42:242–247
- Fernandez-Brime S, Llimona X, Molnar K, Stenroos S et al (2011) Expansion of the Stictidaceae by the addition of the saxicolous lichen-forming genus *Ingvariella*. Mycologia 103:755–763. doi:10.3852/10-287
- Fermández-Brime S, Llimona X, Lutzoni F, Gaya E (2013) Phylogenetic study of *Diploschistes* (lichen-forming *Ascomycota*: *Ostropales*: *Graphidaceae*), based on morphological, chemical, and molecular data. Taxon 62:267–280
- Ferrer A, Raja HA, Shearer CA (2008) *Lucidascocarpa pulchella*, a new ascomycete genus and species from freshwater habitats in the American tropics. Mycologia 100:642–646
- Ferrer A, Miller AN, Shearer CA (2011) *Minutisphaera* and *Natipusilla*: two new genera of freshwater *Dothideomycetes*. Mycologia 103:411–423
- Ferrer A, Miller AN, Sarmiento C, Shearer CA (2012) Three new genera representing novel lineages of *Sordariomycetidae* (*Sordariomycetes*, *Ascomycota*) from tropical freshwater habitats in Costa Rican. Mycologia 104:865–879
- Feuerstein SH, Cunha-Dias IPR, Aptroot A, Eliasaro S, Cáceres MES (2014) Three new *Diorygma* (*Graphidaceae*) species from Brazil, with a revised world key. Lichenologist 46:753–761
- Filippova NV (2012) Discomycetes from plant, leave and sphagnum litter in ombrotrophic bog (West Siberia). Environ Dyn Global Climate Change 3:5
- Fiorentino J (2015) Clarification regarding old records of *Roccella* in the Maltese Islands. Mycosphere 6:673–680
- Fischer M, Schneider P, Kraus C, Molnar M et al (2017) Grapevine trunk disease in German viticulture: occurrence of lesser known fungi and first report of *Phaeoacremonium viticola* and *P. fraxinopennsylvanicum*. VITIS—J Grapevine Res 55:145–156
- Fluza PO, Gusmão LFP (2013) Ingoldian fungi from the semi-arid Caatinga biome of Brazil. Mycosphere 4:1133–1150
- Fluza PO, Gusmão LF, Cruz AC, Castañeda-Ruiz RF (2014) Conidial fungi from the semiarid Caatinga biome of Brazil: a new species of *Pseudoacrodiclys*. Mycotaxon 127:33–37
- Fluza PO, Gusmão LFP, Castañeda-Ruiz RF (2015a) Conidial fungi from the semiarid Caatinga biome of Brazil: a new species of *Selenosporella* from submerged leaves. Mycotaxon 130:601–605
- Fluza PO, Gusmão LFP, Castañeda-Ruiz RF (2015b) *Synchaetomella aquatica* sp. nov. from submerged leaves from Brazil. Mycotaxon 130:1135–1139
- Fluza PO, Ottoni-Boldrini BMP, Monteiro JS, Catena NR et al (2015c) First records of Ingoldian fungi from the Brazilian Amazon. Braz J Bot 38:615–621
- Fluza PO, Monteiro JS, Gusmão LFP, Castañeda-Ruiz RF (2016) *Mirandina uncinata* sp. nov. from submerged leaves from Brazil. Mycotaxon 131:141–144
- Flakus A, Farkas E (2013) A contribution to the taxonomy of *Lyromma* (*Lyrommataceae*, lichenized *Ascomycota*) with a species key. Mycotaxon 124:127–141
- Flakus A, Kukwa M (2011) *Lepraria maderensis* Kukwa & Flakus, a new lichen species containing gyrophoric and lecanoric acids. Nova Hedwig 92:95–99
- Flakus AD, Kukwa MA (2012) New records of lichenicolous fungi from Bolivia. Opusc Phil 11:36–48
- Flakus A, Kukwa M (2014) The first squamulose *Thelecarpon* species (*Thelecarpaceae*, *Ascomycota*) discovered in the biological soil crusts in the Bolivian Andes. Phytotaxa 175:281–286
- Flakus A, Lücking R (2008) New species and additional records of folicolous lichenized fungi from Bolivia. Lichenologist 40:423–436
- Flakus PR, Printzen C (2014) *Palicella*, a new genus of lichenized fungi and its phylogenetic position within *Lecanoraceae*. Lichenol 46:535–552
- Flakus A, Elix JA, Rodriguez P, Kukwa M (2011) New species and records of *Lepraria* (*Stereocaulaceae*, lichenized *Ascomycota*) from South America. Lichenologist 43:57–66
- Flakus A, Rodriguez Saavedra P, Kukwa M (2012) A new species and new combinations and records of *Hypotrachyna* and *Remototrichyna* from Bolivia. Mycotaxon 119:157–166
- Flakus A, Sipman HJ, Bach K, Flakus PR et al (2013) Contribution to the knowledge of the lichen biota of Bolivia. 5. Pol Bot J 58:697–733
- Flakus A, Etayo J, Kukwa M (2014) *Melaspilea tucumana*, a new gall-forming lichenicolous fungus from the tropical Andes in Bolivia. Lichenologist 46:657–662
- Flakus A, Sipman HJ, Rodriguez Flakus P, Jabłońska A et al (2015) Contribution to the knowledge of the lichen biota of Bolivia. 7. Pol Bot J 60:81–98
- Fleck SC, Burkhardt B, Pfeiffer E, Metzler M (2012) *Alternaria* toxins: altertoxin II is a much stronger mutagen and DNA strand breaking mycotoxin than alternariol and its methyl ether in cultured mammalian cells. Toxicol Lett 214:27–32
- Flores-Rentería L, Lau MK, Lamit LJ, Gehring CA (2014) An elusive ectomycorrhizal fungus reveals itself: a new species of *Geopora* (*Pyronemataceae*) associated with *Pinus edulis*. Mycologia 106:553–563
- Flors V, Ton J, Van Doorn R, Jakab G et al (2008) Interplay between JA, SA and ABA signalling during basal and induced resistance against *Pseudomonas syringae* and *Alternaria brassicicola*. Plant J 54:81–92
- Fotouhifar KB, Hedjaroude GA, Leuchtmann A (2010) ITS rDNA phylogeny of Iranian strains of *Cytospora* and associated teleomorphs. Mycologia 102:1369–1382

- Fournier J, Lechat C (2011) *Eutypella phaeospora*, a new species on *Chenopodiaceae*. Mycotaxon 118:441–446
- Fournier J, Stadler M, Hyde KD, Duong ML (2010) The new genus *Rostrohypoxylon* and two new *Annulohypoxylon* species from Northern Thailand. Fungal Divers 40:23–36
- Fournier J, Flessa F, Persöhl D, Stadler M (2011) Three new *Xylaria* species from southwestern Europe. Mycol Prog 10:33–52
- Fournier J, Raja HA, Shearer CA (2015) Freshwater ascomycetes: *Jahnula purpurea* (*Jahnulales, Dothideomycetes*), a new species on submerged wood from Martinique Island, Lesser Antilles. MycoKeys 9:29–36
- Frank J, Crous PW, Groenewald JZ, Oertel B et al (2010) *Microcyclospora* and *Microcycloporella*: novel genera accommodating epiphytic fungi causing sooty blotch on apple. Persoonia 24:93–105
- Frank JL, Southworth D, Reha C (2011) From mycorrhiza to fruiting body: a new species of *Geopora* on *Cercocarpus ledifolius* (*Rosaceae*). Mycologia 103:1194–1200
- Freire ES, Campos VP, Pinho RSC, Oliveira DF et al (2012) Volatile substances produced by *Fusarium oxysporum* from coffee rhizosphere and other microbes affect *Meloidogyne incognita* and *Arthrobotrys conoides*. J Nematol 44:321
- Freitas LF, Barriga EJ, Barahona PP, Lachance MA, Rosa CA (2013) *Kodamaea transpacifica* fa. sp. nov., a yeast species isolated from ephemeral flowers and insects in the Galápagos Islands and Malaysia: further evidence for ancient human transpacific contacts. Int J Syst Evol Microbiol 63:4324–4329
- Friebes G (2012) A key to the non-lichenicolous species of the genus *Capronia* (*Herpotrichiellaceae*). Ascomycete 4:55–64
- Friebes G, Wendelin I (2014) Über einige seltene und interessante Ascomyceten-Funde vom Reinerkogel (Graz, Steiermark, Österreich). Joannea Bot 11:5–33
- Frisch A, Ohmura Y (2015) The phylogenetic position of *Normandina simodensis* (*Verrucariaceae*, lichenized Ascomycota). Bull Natl Mus Natl Sci 41:1–7
- Frisch A, Thor G (2010) *Cryphtonia*, a new genus of byssoid Arthoniaceae (lichenised Ascomycota). Mycol Prog 9:281–303
- Frisch A, Thor G, Ertz D, Grube M (2014) The Arthonialean challenge: restructuring Arthoniaceae. Taxon 63:727–744
- Frisch A, Ohmura Y, Ertz D, Thor G (2015) Inoderma and related genera in Arthoniaceae with elevated white pruinose pycnidia or sporodochia. Lichenol 47:233–256
- Frisvad JC, Yilmaz N, Thrane U, Rasmussen KB, Samson RA (2013) *Talaromyces atroroseus*, a new species efficiently producing industrially relevant red pigments. PLoS ONE 8:e84102
- Fritis MC, Lagos CR, Sobralo NQ, Venegas IM et al (2013) Depsides and triterpenes in *Pseudocyphellaria coriifolia* (lichens) and biological activity against *Trypanosoma cruzi*. Nat Prod Res 27:1607–1610
- Fryday AM (2010) A brief lichen foray in the Mount Washington alpine zone—including *Clauroxia chalybeoides*, *Porina norrlinii* and *Stereocaulon leucophaeopsis* new to North America. Opusc Phil 8:1–10
- Fryday AM (2011) New species and combinations in *Calvitimela* and *Tephromela* from the southern subpolar region. Lichenologist 43:225
- Fryday AM, Hertel HA (2014) Contribution to the family *Lecideaceae* s. lat. (*Lecanoromycetidae* inc. sed., lichenized Ascomycota) in the southern subpolar region; including eight new species and some revised generic circumscriptions. Lichenologist 46:389–412
- Fryday AM, Tønsberg T (2015) *Ameliella*—a bryicolous lichen genus rediscovered in North America after 50 years. Evansia 32:171–175
- Fryday AM, Printzen C, Ekman S (2014) *Bryobilimbia*, a new generic name for *Lecidea hypnorum* and closely related species. Lichenologist 46:25–37
- Fu CH, Hsieh HM, Chen CY, Chang TT et al (2013) *Ophiodiaporthe cyatheae* gen. et sp. nov., a diaporthalean pathogen causing a devastating wilt disease of *Cyathea lepifera* in Taiwan. Mycologia 105:861–872
- Fuijano N, Fusilli S, Incorvaia C (2012) A role for measurement of nasal IgE antibodies in diagnosis of *Alternaria*-induced rhinitis in children. Allergol Immunopathol 40:71–74
- Funck JA, Bec A, Perrière F, Felten V, Danger M (2015) Aquatic hyphomycetes: a potential source of polyunsaturated fatty acids in detritus-based stream food webs. Fungal Ecol 13:205–210
- Gagunashvili AN, Davíðsson SP, Jónsson ZO, Andrésson ÓS (2009) Cloning and heterologous transcription of a polyketide synthase gene from the lichen *Solorina crocea*. Mycol Res 113:354–363
- Gaitnieks T, Klavina D, Muiznieks I, Pennanen T et al (2016) Impact of *Heterobasidion* root-rot on fine root morphology and associated fungi in *Picea abies* stands on peat soils. Mycorrhiza 26:465–473
- Galán R, Checa J, Blanco MN, Platas G et al (2015) Taxonomic position of the genus *Bicornispora* and the appearance of a new species *Bicornispora seditiosa*. Mycologia 107:793–807
- Galloway DJ (2010) Additions to the *Placopsis* mycobiota (*Trapeliaceae, Ascomycota*) of southern South America, with notes on new records (including *Aspiciliopsis macrophthalma*), and a revised regional key to the species. Lichenologist 42:727–737
- Galloway DJ (2013) The lichen genera *Aspiciliopsis*, and *Placopsis* (*Trapeliaceae: Ascomycota*) in New Zealand. Phytotaxa 120:1–194
- Galloway DJ (2015) Contributions to a history of New Zealand lichenology 5*. James Murray (1923–1961). Phytotaxa 198(1):1–67
- Galloway DJ, Elix JA (2013) Reinstatement of *Crocodia* link (*Lobariaceae: Ascomycota*) for five species formerly included in *Pseudocyphellaria* Vain. Australas Lichenol 72:32–42
- Gams W, Seifert KA, Morgan-Jones G (2009) New and validated hyphomycete taxa to resolve nomenclatural and taxonomic issues. Mycotaxon 110:89–108
- Ganie AA, Azam MF, Wani AH (2012) *Exosporodiella* gen. nov.—a new coelomycetous fungus on leaves of *Phoenix* from India. Arch Phytopathol Plant Prot 45(13):1582–1585
- Gao TL, Ren Q (2013) New records of *Ochrolechia* and *Placopsis* from the Hengduan Mountains, China. Mycotaxon 122:461–466
- Gao SS, Li XM, Wang BG (2009) Perylene derivatives produced by *Alternaria alternata*, an endophytic fungus isolated from *Laurencia* species. Nat Prod Commun 4:1477–1480
- Gao JM, Xia JW, Ma Y, Li Z, Zhang XG (2015a) *Blastophragma chongqingense* sp. nov. and a new record of *Bahusutrabeja angularis* from southern China. Mycotaxon 130:821–825
- Gao L, Ma Y, Zhao W, Wei Z et al (2015b) Three new species of *Cyphellophora* (*Chaetothyriales*) associated with sooty blotch and flyspeck. PLoS ONE 10:e0136857
- Garibaldi A, Bertetti D, Poli A, Gullino ML (2012) First report of leaf spot of garden lupin (*Lupinus polyphyllus*) caused by *Pleiochaeta setosa* in Italy. Plant Dis 96:909
- Garrido-Benavent I, Pérez-Ortega S (2015) Unravelling the diversity of European *Caliciopsis* (*Coryneliaceae, Ascomycota*): *Caliciopsis valentina* sp. nov. and *C. beckhausii* comb. nov., with a worldwide key to *Caliciopsis*. Mycol Prog 14:1–11
- Garrido-Benavent I, Söchting U, de los Ríos Murillo A, Pérez-Ortega S (2016) *Shackletonia cryodesertorum* (*Teloschistaceae*). Mycol Prog 15:743–754
- Garzoli L, Gnavi G, Tamme F, Tosi S et al (2015) Sink or swim: updated knowledge on marine fungi associated with wood

- substrates in the Mediterranean Sea and hints about their potential to remediate hydrocarbons. *Prog Oceanogr* 137:140–148
- Gasparyan A, Sipman HJ (2013) New lichen records from Armenia. *Mycotaxon* 123:491–492
- Gasparyan A, Sipman HJ (2016) The epiphytic lichenized fungi in Armenia: diversity and conservation. *Phytotaxa* 281:1–68
- Gates G, Van Vooren N (2016) *Jafnea pallida* (Pezizales), a new name for a cup fungus rediscovered in Tasmania. *Ascomycete* 8:101–104
- Gautam AK (2015) *Prillieuxina aeglicola* sp. nov. (ascomycota), a new black mildew fungus from Himachal Pradesh, India. *Curr Res Environ Appl Mycol* 5:70–73
- Gautam AK, Avasthi S, Bhadauria R (2015) A new species of *Scytalidium* from Himachal Pradesh, India. *J New Biol Rep* 4:7–10
- Gaya E, Navarro-Rosinés P, Llimona X, Hladun N, Lutzoni F (2008) Phylogenetic reassessment of the *Teloschistaceae* (lichen-forming Ascomycota, Lecanoromycetes). *Mycol Res* 112:528–546
- Gaya E, Högnabba F, Holguín Á, Molnar K et al (2012) Implementing a cumulative supermatrix approach for a comprehensive phylogenetic study of the *Teloschistales* (Pezizomycotina, Ascomycota). *Mol Phyl Evol* 63:374–387
- Gaya E, Fernández-Brime S, Vargas R, Lachlan RF et al (2015) The adaptive radiation of lichen-forming *Teloschistaceae* is associated with sun screening pigments and a bark-to-rock substrate shift. *PNAS* 112:11600–11605
- Gayatri Mambiar R, Raveendran K (2009) Eco-biodiversity of marine mycoflora of gulf of Mannar Biosphere Reserve (India). *Ecol Environ Conserv* 15:739–743
- Gazis R, Miadlikowska J, Lutzoni F, Arnold AE, Chaverri P (2012) Culture-based study of endophytes associated with rubber trees in Peru reveals a new class of Pezizomycotina: Xylonomycetes. *Mol Phyl Evol* 65:294–304
- Gazis R, Kuo A, Riley R, LaButti K et al (2016) The genome of *Xylona heveae* provides a window into fungal endophytism. *Fungal Biol* 120:26–42
- Ge YP, Lv GX, Shen YN, Li M et al (2012) First report of subcutaneous phaeohyphomycosis caused by *Ochroconis tshawytschae* in an immunocompetent patient. *Med Mycol* 50:637–640
- Ge ZW, Yang ZL, Pfister DH, Carbone M et al (2014) Multigene molecular phylogeny and biogeographic diversification of the earth tongue fungi in the genera *Cudonia* and *Spathularia* (Rhytismatales, Ascomycota). *PLoS ONE* 9:e103457
- Gea FJ, Navarro MJ, Suz LM (2016) First report of *Cladobotryum mycophilum* causing cobweb on cultivated king oyster mushroom in Spain. *Plant Dis* 100:2334
- Geiser DM, Aoki T, Bacon CW, Baker SE et al (2013) One fungus, one name: defining the genus *Fusarium* in a scientifically robust way that preserves longstanding use. *Phytopathology* 103(5):400–408
- Geltner C, Sorschag S, Willinger B, Jaritz T et al (2015) Necrotizing mycosis due to *Verruconis gallopava* in an immunocompetent patient. *Infection* 43:743–746
- Geng YH, Zhang TY (2014) Two new species of *Humicola* from soil. *Mycotaxon* 33:800–805
- Geng RM, Fu Y, Zhang WM, Zhang JP, Yu LQ (2008) Efficacy and safety of *Bipolaris sorokiniana* and *Bipolaris coicis* for the control of *Echinochloa crus-galli* in paddy field. *Chin J Rice Sci* 3:015
- Gessler NN, Egorova AS, Belozerskaya TA (2014) Melanin pigments of fungi under extreme environmental conditions (review). *Appl Biochem Microbiol* 50:105–113
- Ghazvini H, Tekauz A (2008) Host-pathogen interactions among barley genotypes and *Bipolaris sorokiniana* isolates. *Plant Dis* 92:225–233
- Gilbert OL, Hawksworth DL (2009) *Clauroxia* D. Hawksw (1988). In: Smith CW, Aptroot A, Coppins BJ, Fletcher A, Gilbert OL, James PW, Wolseley PA (eds) The lichens of Great Britain and Ireland, 2nd edn. British Lichen Society, London, pp 338–339
- Gilgado F, Cano J, Gené J, Sutton DA, Guarro J (2008) Molecular and phenotypic data supporting distinct species statuses for *Scedosporium apiospermum* and *Pseudallescheria boydii* and the proposed new species *Scedosporium dehoogii*. *J Clin Microbiol* 46:766–771
- Gilgado F, Gene J, Cano JF, Guarro J (2010) Heterothallism in *Scedosporium apiospermum* and description of its teleomorph *Pseudallescheria apiosperma* sp. nov. *Med Mycol* 48:122–128
- Giordani P, Benesperi R, Rizzi G, Brunialti G (2009) New records for lichen regional floras of Italy. *Webbia* 64:153–158
- Giordani P, Benesperi R, Mariotti MG (2015) Local dispersal dynamics determine the occupied niche of the red-listed lichen *Seirophora villosa* (Ach.) Frödén in a Mediterranean Juniperus shrubland. *Fungal Ecol* 13:77–82
- Giraldo A, Gené J, Cano J, de Hoog GS, Guarro J (2012) Two new species of *Acremonium* from Spanish soils. *Mycologia* 8:11–402
- Giraldo A, Gené J, Sutton DA, Madrid H et al (2014a) Phylogenetic circumscription of *Arthrographis* (Eremomycetaceae, Dothideomycetes). *Persoonia* 32:102–114
- Giraldo A, Sutton DA, Samerpitak K, de Hoog GS et al (2014b) Occurrence of *Ochroconis* and *Verruconis* species in clinical specimens from the United States. *J Clin Microbiol* 52:4189–4201
- Giraldo A, Gené J, Cano J, de Hoog S et al (2014c) *Acremonium* with catenate elongate conidia: phylogeny of *Acremonium fusidioides* and related species. *Mycologia* 106:328–338
- Giraldo A, Crous PW, Schumacher RK, Cheewangkoon R et al (2017) The genera of fungi-G3: *Aleurocystis*, *Blastacervulus*, *Clypeophysalospora*, *Licrostroma*, *Neohendersonia* and *Spumatoria*. *Mycol Prog* 16:1–24
- Giralt M (2009) *Rinodina etayoi*, a new saxicolous lichen species from the Canary Islands. *Lichenologist* 41:141–145
- Giralt M, Klaus KALB (2010) *Rinodina fuscoisidiata*, a new muscicolous, isidiate species from Venezuela. *Lichenologist* 42:73–76
- Giralt M, Klaus KALB, Mayrhofer H (2009) *Rinodina brasiliensis*, a new corticolous isidiate species, and closely related taxa. *Lichenologist* 41:179–187
- Giralt M, van den Boom PPG, Elix JA (2010) *Endohyalina*, the genus in the Physciaceae to accommodate the species of the *Rinodina ericina*-group. *Mycol Prog* 9:37–48
- Giralt M, Van den Boom PP, Mayrhofer H, Elix JA (2014) Three new species of Crustose Physciaceae from Guatemala, with notes on some additional species. *Phytotaxa* 164:79–90
- Giraud S, Favenne L, Bougnoux ME, Bouchara JP (2013) *Rasamsonia argillacea* species complex: taxonomy, pathogenesis and clinical relevance. *Future Microbiol* 8:967–978
- Giri DK, Sims WP, Suran R, Cooper JJ, Gavrilov BK, Mansell J (2011) Cerebral and renal phaeohyphomycosis in a dog infected with *Bipolaris* species. *Vet Pathol Online* 48:754–757
- Gladdish S, Frank JL, Southworth D (2010) The serpentine syndrome belowground: hypogeous fungi and ectomycorrhizas associated with conifers. *Can J For Res* 40:1671–1679
- Glejdura S, Kučera V, Lizoň P, Kunca V (2015) *Pseudoplectania lignicola* sp. nov. described from central Europe. *Mycotaxon* 130:1–10
- Glienke C, Pereira OL, Stringari D, Fabris J, Kava-Cordeiro V, Galli-Terasawa L, Cunningham J, Shivas RG, Groenewald JZ, Crous PW (2011) Endophytic and pathogenic *Phyllosticta*

- species, with reference to those associated with Citrus Black Spot. Persoonia 26:47–56
- Gnavi G, Ercole E, Panno L, Vizzini A, Varese GC (2014) *Dothideomycetes* and *Leotiomycetes* sterile mycelia isolated from the Italian seagrass *Posidonia oceanica* based on rDNA data. SpringerPlus 3:508
- Goel M, Sharma PK, Dureja P, Rani A, Uniyal PL (2011) Antifungal activity of extracts of the lichens *Parmelia reticulata*, *Ramalina roesleri*, *Usnea longissima* and *Stereocaulon himalayense*. Arch Phytopathol Plant Prot 44:1300–1311
- Goh TK, Lee LL, Teo KC (2013) A new coprophilous *Gilmaniella* species from Malaysia. Mycotaxon 125:234–240
- Goh TK, Lau WY, Teo KC (2014a) A new species of *Nawawia* from Malaysia, with a synopsis of the genus. Mycotaxon 129:109–118
- Goh TK, Lau WY, Teo KC (2014b) *Paliphora curviapicis* sp. nov. from Malaysia, and a synopsis of the genus. Mycotaxon 127:145–153
- Goldmann L, Weir A (2012) Position specificity in *Chitonomycetes* (*Ascomycota*, *Laboulbeniomycetes*) on *Laccophilus* (Coleoptera, Dytiscidae): a molecular approach resolves a century-old debate. Mycologia 104:1143–1158
- Gomathinayagam S, Rekha M, Murugan SS, Jagessar JC (2010) The biological control of paddy disease brown spot (*Bipolaris oryzae*) by using *Trichoderma viride* in vitro condition. J Biopestic 3:93–95
- Gomes RR, Glienke C, Videira SIR, Lombard L et al (2013) *Diaporthe*: a genus of endophytic, saprobic and plant pathogenic fungi. Persoonia 31:1–41
- Goni R, Sharma N (2015) Additions to lichen flora of Jammu and Kashmir, India. Trop Plant Res 2:78–81
- Goni R, Raina AK, Magotra R, Sharma N (2014) Lichen flora of Jammu and Kashmir State, India: an updated checklist. Trop Plant Res 2:64–71
- Gonzalez-Menendez V, Perez-Bonilla M, Martin J, Munoz F et al (2016) Multicomponent analysis of the differential induction of secondary metabolite profiles in fungal endophytes. Molecules 21:234. doi:10.3390/molecules21020234
- Goonasekara ID, Maharachchikumbura SS, Wijayawardene NN, Phookamsak R et al (2016) *Seimatosporium quercina* sp. nov. (*Discosphaeriaceae*) on *Quercus robur* from Germany. Phytotaxa 255:240–248
- Gräfenhan T, Schroers H-J, Nirenberg HI, Seifert KA (2011) An overview of the taxonomy, phylogeny, and typification of nectriaceous fungi in *Cosmospora*, *Acremonium*, *Fusarium*, *Stilbella*, and *Volutella*. Stud Mycol 68:79–113
- Gramaje D, Agustí-Brisach C, Pérez-Sierra A, Moralejo E et al (2012) Fungal trunk pathogens associated with wood decay of almond trees on Mallorca (Spain). Persoonia 28:1–13
- Granados M, Castañeda-Ruiz RF, Castro O, Minter DW, Kendrick B (2014) Microfungi from Costa Rica. A new species and a new combination in *Codinaea*. Mycotaxon 127:115–120
- Granmo A, Mathiassen G (2013) A new species of *Dothiopeltis* from Norway. Sydowia 65(2):343–348
- Greif MD, Stchigel AM, Miller AN, Huhndorf SM (2009) A re-evaluation of genus *Chaetomidium* based on molecular and morphological characters. Mycologia 101:554–564
- Grinn-Gofroń A, Rapiejko P (2009) Occurrence of *Cladosporium* spp. and *Alternaria* spp. spores in Western, Northern and Central-Eastern Poland in 2004–2006 and relation to some meteorological factors. Atmos Res 93:747–758
- Grinn-Gofroń A, Strzelczak A (2008) Artificial neural network models of relationships between *Alternaria* spores and meteorological factors in Szczecin (Poland). Int J Biometeorol 52:859–868
- Grobelaar JW, Aghayeva DN, de Beer ZW, Bloomer P et al (2009) Delimitation of *Ophiostoma quercus* and its synonyms using multiple gene phylogenies. Mycol Prog 8:221–236
- Grochowski P (2012) Secondary succession with the participation of protected species of lichens in the charred areas of the forest inspectorate of Lubsko. W: Lipnicki L. (red) Lichen protection—protected lichen species, Sonar Literacki, Gorzów Wielkopolski, pp 205–212
- Groenewald M, Linde CC, Groenewald JZ, Crous PW (2008) Indirect evidence for sexual reproduction in *Cercospora beticola* populations from sugar beet. Plant Pathol 57:25–32
- Groenewald JZ, Nakashima C, Nishikawa J, Shin HD et al (2013) Species concepts in *Cercospora*: spotting the weeds among the roses. Stud Mycol 75:115–170
- Gross A, Holdenrieder O, Pautasso M, Queloz V, Sieber TN (2014) *Hymenoscyphus pseudoalbidus*, the causal agent of European ash dieback. Mol Plant Pathol 15:5–21
- Grube M, Köberl M, Lackner S, Berg C, Berg G (2012) Host-parasite interaction and microbiome response: effects of fungal infections on the bacterial community of the Alpine lichen *Solorina crocea*. FEMS Microbiol Ecol 82:472–481
- Grum-Grzhimaylo AA, Debets AJ, van Diepeningen AD, Georgieva ML, Bilanenko EN (2013a) *Sodiomyces alkalinus*, a new holomorphic alkaliphilic ascomycete within the *Plectosphaerellaceae*. Persoonia 31:147–158
- Grum-Grzhimaylo AA, Georgieva ML, Debets AJM, Bilanenko EN (2013b) Are alkalitolerant fungi of the *Emericellopsis* lineage (*Bionectriaceae*) of marine origin? IMA Fungus 4:213–228
- Grum-Grzhimaylo AA, Georgieva ML, Bondarenko SA, Debets AJ, Bilanenko EN (2016) On the diversity of fungi from soda soils. Fungal Divers 76:27–74
- Grundy KC, Woodward S, Genney DR, Taylor AF (2012) A molecular approach to explore the extent of the threatened fungus *Hypocreopsis rhododendri* within wood. Fungal Biol 116:354–362
- Grünig CR, Queloz V, Duò A, Sieber TN (2009) Phylogeny of *Phaeomollisia piceae* gen. sp. nov. a dark, septate, conifer-needle endophyte and its relationships to *Phialocephala* and *Acephala*. Mycol Res 113:207–221
- Gryzenhout M, Wingfield BD, Wingfield MJ (2009) Taxonomy, phylogeny, and ecology of bark-inhabiting and tree-pathogenic fungi in the *Cryphonectriaceae*. American Phytopathological Society (APS Press), Saint Paul
- Gryzenhout M, Tarigan M, Clegg PA, Wingfield MJ (2010) *Cryptometriion aestuans* gen. [et] sp. nov. (*Cryphonectriaceae*) pathogenic to Eucalyptus in Indonesia. Australas Plant Pathol 39:161–169
- Gu W (2009) Bioactive metabolites from *Alternaria brassicicola* ML-P08, an endophytic fungus residing in *Malus halliana*. World J Microbiol Biotechnol 25:1677–1683
- Guang-yu X, Bi-fa S, Ke-ming C, Si-ai F (2008) Study on the control of *Agaricodochium camelliae*. J Hebei Agric Sci 9:17
- Guarro J, Gené J, Stchigel A, Figueras MJ (2012) Atlas of soil ascomycetes. CBS Biodiversity Series 10, Utrecht
- Guatimosim E, Pinto HJ, Barreto RW, Prado J (2014) *Rhagadolobiopsis*, a new genus of *Parmulariaceae* from Brazil with a description of the ontogeny of its ascocarps. Mycologia 106:276–281
- Guatimosim E, Firmino A, Bezerra J, Pereira O et al (2015) Towards a phylogenetic reappraisal of *Parmulariaceae* and *Asterinaceae* (*Dothideomycetes*). Persoonia 35:230–241
- Guatimosim E, Schwartsburd PB, Barreto RW, Crous PW (2016a) Novel fungi from an old niche: cercosporoid and related sexual morphs on ferns. Persoonia 37:106–141

- Guatimosim E, Schwartsburd PB, Crous PW, Barreto RW (2016b) Novel fungi from an ancient niche: lachnoid and chalara-like fungi on ferns. *Mycol Prog* 15:1239–1267
- Gudmundsson M, Hansson H, Karkehabadi S, Larsson A et al (2016) Structural and functional studies of the glycoside hydrolase family 3 β-glucosidase Cel3A from the moderately thermophilic fungus *Rasamsonia emersonii*. *Acta Crystallogr D* 72:860–870
- Guecin F, Kaya A, Soylu MK, Uzun Y (2010) Picoa Vittad A new truffle genus record for Turkey. *Biol Divers Conserv* 3:23–25
- Gueidan C, Lendemer JC (2015) Molecular data confirm morphological and ecological plasticity within the North-American endemic lichen *Willeya diffractella* (*Verrucariaceae*). *Syst Bot* 40(2):369–375
- Gueidan C, Villaseñor CR, de Hoog GS, Gorushina AA et al (2008) A rock-inhabiting ancestor for mutualistic and pathogen-rich fungal lineages. *Stud Mycol* 61:111–119
- Gueidan C, Savić S, Thüs H, Roux C et al (2009) Generic classification of the *Verrucariaceae* (*Ascomycota*) based on molecular and morphological evidence: recent progress and remaining challenges. *Taxon* 58:184–208
- Gueidan C, Thüs H, Pérez-Ortega S (2011) Phylogenetic position of the brown algae-associated lichenized fungus *Verrucaria tavaresiae* (*Verrucariaceae*). *Bryologist* 114:563–569
- Gueidan C, Aptroot A, Cáceres MES, Badali H, Stenroos S (2014a) A reappraisal of orders and families within the subclass *Chaetothyriomycetidae* (*Eurotiomycetes*, *Ascomycota*). *Mycol Prog* 13:1027–1039
- Gueidan C, Monnat JY, Navarro-Rosines P, Roux C (2014b) *Trimmatothelopsis versipellis*: discovery of several localities in the department of Finistere (France), phylogenetic placement and taxonomic consequences. *Bull Soc Linn Provence* 65:47–65
- Gueidan C, Truong VD, Lu NT (2014c) Phylogeny and taxonomy of *Staurothele* (*Verrucariaceae*, lichenized ascomycetes) from the karst of northern Vietnam. *Lichenologist* 46:515–533
- Gueidan C, Aptroot A, Cáceres MES (2016) Molecular phylogeny of the tropical family *Pyrenulaceae*: contribution from dried specimens and FTA card samples. *Mycol Prog* 15:1–21
- Guevara-Guerrero G, Bonito G, Cázares-González E, Healy RA, Trappe J (2015) Novel *Tuber* spp. (*Tuberaceae*, *Pezizales*) in the *Puberulum* Group from North American. *Ascomycets* 7:367–374
- Guinea J, Sandoval-Denis M, Escrivano P, Peláez T et al (2015) *Aspergillus citrinoterreus*, a new species of the section Terrei isolated from samples of patients with nonhematological predisposing conditions. *J Clin Microb* 53:611–617
- Gulder TA, Hong H, Correa J, Egereva E et al (2012) Isolation, structure elucidation and total synthesis of lajollamide A from the marine fungus *Asteromyces cruciatus*. *Mar Drugs* 10:2912–2935
- Gulis V, Baschien C, Marvanová L (2012) Two new *Trichladium* species from streams in Alaska. *Mycologia* 104:1510–1516
- Gumboski EL (2014) Contribution to the knowledge of lichenized mycota from Southern Brazil. I-Species of *Ramboldia* (*Ascomycota*: *Lecanoraceae*). *Acta Biol Catarinense* 1(2):60–66
- Gunde-Cimerman N, Zalar P (2014) Extremely halotolerant and halophilic fungi inhabit brine in solar salterns around the globe. *Food Technol Biotechnol* 52:170
- Guo MT, Qiao M, Li JY, Wang W, Yu ZF (2015) *Verticilicladus hainanensis*, a new aquatic hyphomycete. *Mycotaxon* 130:275–278
- Gupta P, Sinha GP (2016) A first note on foliicolous lichens of Assam, India. *J Threat Taxa* 8:9014–9023
- Gupta V, Shamas N, Razdan VK, Sharma BC et al (2013) Foliar application of fungicides for the management of brown spot disease in rice (*Oryza sativa* L.) caused by *Bipolaris oryzae*. *Afr J Agric Res* 8:3303–3309
- Gur L, Frenkel O (2015) First report of leaf spot on blue lupin (*Lupinus pilosus*) caused by *Pleiochaeta setosa* in Israel. *Plant Dis* 100:525
- Gusmão LF, Leao-Ferreira SM, Marques MF, de Almeida DA (2008) New species and records of *Paliphora* from the Brazilian semi-arid region. *Mycologia* 100:306–309
- Guu JR, Ju YM, Hsieh HJ (2010) Bionectriaceous fungi collected from forests in Taiwan. *Bot Stud* 51:61–74
- Guzow-Krzemińska B, Halda JP, Czarnota P (2012) A new *Agonimia* from Europe with a flabelliform thallus. *Lichenologist* 44:55–66
- Habibi A, Banihashemi Z (2016) Mating system and role of pycnidiospores in biology of *Polystigma amygdalinum*, the causal agent of almond red leaf blotch. *Phytopathol Mediterr* 55:98
- Habibi A, Banihashemi Z, Mostowfizadeh-Ghalmfarsa R (2015) Phylogenetic analysis of *Polystigma* and its relationship to *Phyllachorales*. *Phytopathol Mediterr* 54:45
- Hadizadeh I, Peivastegan B, Hamzehzarghani H (2009) Antifungal activity of essential oils from some medicinal plants of Iran against *Alternaria alternata*. *Am J Appl Sci* 6:857–861
- Haelewaters D, De Kesel A (2013) A new species of *Cantharomyces* (*Laboulbeniales*, *Ascomycota*) from the Netherlands. *Mycotaxon* 123:467–472
- Haelewaters D, Yaakop S (2014) New and interesting *Laboulbeniales* from southern and southeastern Asia. *Mycotaxon* 129:454–493
- Haelewaters D, Schiltzhoven M, Pfister DH (2014a) On *Diphymyces* (*Laboulbeniales*, *Ascomycota*) in Malaysian Borneo. *Plant Ecol Evol* 147:93–100
- Haelewaters D, Vorst O, De Kesel A (2014b) New and interesting *Laboulbeniales* (Fungi, Ascomycota) from the Netherlands. *Nova Hedwig* 98:113–125
- Haelewaters D, Boer P, Noordijk J (2015a) Studies of *Laboulbeniales* (Fungi, Ascomycota) on Myrmica ants: *Rickia wasmannii* in the Netherlands. *J Hymenopt Res* 44:39
- Haelewaters D, Gorczak M, Pflieler W, Tertally A et al (2015b) Bringing *Laboulbeniales* into the 21st century: enhanced techniques for extraction and PCR amplification of DNA from minute ectoparasitic fungi. *IMA Fungus* 6:363–372
- Haelewaters D, Zhao SY, De Kesel A, Handlin RE et al (2015c) *Laboulbeniales* (Ascomycota) of the Boston Harbor Islands I: species parasitizing Coccinellidae and Staphylinidae, with comments on typification. *Northeast Nat* 22:459–477
- Haelewaters D, Minnaar IA, Clusella-Trullas S (2016) First finding of the parasitic fungus *Hesperomyces virescens* (*Laboulbeniales*) on native and invasive ladybirds (Coleoptera, Coccinellidae) in South Africa. *Parasite* 23:5
- Haelewaters D, Pflieler WP, Szentiványi T, Földvári M et al (2017a) Parasites of parasites of bats: *Laboulbeniales* (Fungi: Ascomycota) on bat flies (Diptera: Nycteribiidae) in central Europe. *Parasites Vectors* 10:96
- Haelewaters D, Verhaeghen SJC, Ríos González TA, Bernal Vega JA et al (2017b) New and interesting *Laboulbeniales* from Panama and neighboring areas. *Nova Hedwig*. doi:[10.1127/nova_hedwigia/2017/0410](https://doi.org/10.1127/nova_hedwigia/2017/0410)
- Hafellner J (2009) *Phacothecium* resurrected and the new genus *Phacographa* (Arthoniales) proposed. *Biblioth Lichenol* 100:85–121
- Hafellner J (2015) Distributional and other data for some species of *Didymocytis* (*Dothideomycetes*, *Pleosporales*, *Phaeosphaeriaceae*), including their Phoma-type anamorphs. *Fritschiana* 80:43–88
- Hafellner J, Zimmermann E (2012) A lichenicolous species of *Pleospora* (Ascomycota) and a key to the fungi invading *Physcia* species. *Herzogia* 25:47–59
- Hafellner J, Herzog G, Mayrhofer H (2008) Zur Diversität von lichenisierten und lichenicolosen Pilzen in den Ennstaler Alpen

- (Osterreich: Steiermark, Oberösterreich). Mitt Naturw Ver Steierm 137:131–204
- Hafellner J, Karnefelt I, Wirth V (2010) Diversity and ecology of lichens in polar and mountain ecosystems. Nova Hedwig 91:3–4
- Hajek J, Vaczi P, Bartak M (2009) Photosynthetic electron transport at low temperatures in the green algal foliose lichens *Lasallia pustulata* and *Umbilicaria hirsuta* affected by manipulated levels of ribitol. Photosynthetica 47:199–205
- Halici MG (2015) New records of crustose *Teloschistaceae* and lichenicolous fungi from Turkey. Mycotaxon 130:769–773
- Halici MG, Candan M (2009) New lichenicolous fungi from Turkey. Nova Hedwig 88:483–490
- Halici MG, Ozdemir Turk A, Candan M (2008) *Dacampia cladoniicola* sp. nov. (*Ascomycota*, *Dacampiaceae*) on *Cladonia* sp. from Turkey. Mycotaxon 103:53–57
- Halici MG, Etayo J, Candan M (2009a) Two new lichenicolous species of *Dacampia* on *Teloschistaceae*. Mycotaxon 109:393–398
- Halici MG, Knudsen K, Candan M (2009b) *Lichenochora atrans* (*Phyllachoraceae*), a new lichenicolous species on *Psora decipiens* from Turkey. Mycotaxon 109:239–243
- Halici MG, Knudsen K, Candan M, Turk AO (2009c) A new species of *Polycoccum* (*Dothideales*, *Dacampiaceae*) from Turkey. Nova Hedwig 89:431–436
- Halici M, Hawksworth DL, Candan M, Ozdemir Turk A (2010) A new lichenicolous species of *Capronia* (*Ascomycota*, *Herpotrichiellaceae*), with a key to the known lichenicolous species of the genus. Fungal Divers 40:37–40
- Halici MG, Akgil HE, Öztürk C, Kılıç E (2013) *Polycoccum anatolicum* sp. nov. on *Lepraria incana* and a key to *Polycoccum* species known from Turkey. Mycotaxon 124:45–50
- Halici M, Kocakaya M, Kiriş Z (2014a) Lichenized and lichenicolous fungi of Bakırdağı (Kayseri, Adana). Acta Bot Hung 56:319–332
- Halici MG, Vondrak J, Demirel R, Ceylan A, Candan M (2014b) *Teloschistaceae* (lichenized Ascomycetes) in Turkey II. Some poorly known taxa. Supported by molecular data. Nova Hedwig 98:449–458
- Halonen P, Mylllys L, Velmala S, Hyvarinen H (2009) *Gowardia* (*Parmeliaceae*)—a new alectorioid lichen genus with two species. Bryologist 112:138–146
- Hamada N, Niichiro ABE (2008) Characteristics of recent fungal contamination in bathrooms. Seikatsu Eisei J Urban Living Health Assoc 52:98–106
- Hämäläinen A, Kouki J, Lohmus P (2015) Potential biodiversity impacts of forest biofuel harvest: lichen assemblages on stumps and slash of Scots pine. Can J For Res 45:1239–1247
- Hamayun M, Khan SA, Kim HY, Chaudhary MF et al (2009) Gibberellin production and plant growth enhancement by newly isolated strain of *Scolecobasidium tshawytschae*. J Microbiol Biotechnol 19:560–565
- Han HB, Yang XY, Yang LR, Wang R et al (2008) Determination of contents of crude protein and amino acids in *Arthrobotrys oligospora*. Chin Vet Sci 7:17
- Han JG, Park MJ, Shin HD (2009) *Psilachnum staphyleae*, a new member of foliicolous *Hyaloscyphaceae* from Korea. Mycotaxon 110:219–224
- Han JG, Choi YJ, Pfister DH, Shin HD (2010a) *Scutellinia jejuensis* (*Pezizales*), a new species from Korea. Mycotaxon 112:47–53
- Han Q, Huang L, Buchenauer H, Wang C, Kang Z (2010b) Cytological study of wheat spike infection by *Bipolaris sorokiniana*. J Phytopathol 158:22–29
- Han JG, Hosoya T, Shin HD (2012) *Amicodisca castaneae* sp. nov. (*Hyaloscyphaceae*, *Helotiales*) on Japanese chestnut bur. Mycotaxon 118:89–94
- Han YF, Chen WH, Zou X, Liang ZQ (2013) *Gibellula curvispora*, a new species of *Gibellula*. Mycosistema 32:778
- Han JG, Hosoya T, Sung GH, Shin HD (2014) Phylogenetic reassessment of *Hyaloscyphaceae sensu lato* (*Helotiales*, *Leotiomycetes*) based on multigene analyses. Fungal Biol 118:150–167
- Hanlin RT, González MC (2013) *Ophiiodothella angustissima* comb. nov., a new name for *Acerviclypeatus poriformans* and *O. vaccinii*. Mycotaxon 123:327–334
- Hansen ES (2009) A contribution to the lichen flora of Iceland. Folia Cryptogam Estonica 46:45–54
- Hansen ES (2011) A contribution to lichen flora of the Kobbefjord area, West Greenland. Bot Lith 17:151–156
- Hansen ES (2012) Notes on some new and interesting Greenland lichens IX. Gr Scr 24:55–59
- Hansen ES (2013) Lichens from Peary Land, North Greenland. Folia Cryptogam Estonica 50:3–11
- Hansen ES (2014) Contribution to the lichen flora of South East Greenland. I. The Kangerlussuaq area. Bot Lith 20:153–158
- Hansen ES (2015) Contribution to the lichen flora of South East Greenland. III. The coastal area between 63 and 65 N. Bot Lith 21:119–124
- Hansen K, Perry BA, Dranginis AW, Pfister DH (2013) A phylogeny of the highly diverse cup-fungus family *Pyrenopeltaceae* (*Pezizomycetes*, *Ascomycota*) clarifies relationships and evolution of selected life history traits. Mol Phylogen Evol 67:311–335
- Hanying Y, Wei X, Xingxi G (2012) Development of a transformation method for the nematophagous fungus *Dactylellina cionopaga*. Afr J Biotech 11:3370–3378
- Hao L, Chen C, Zhang R, Zhu M et al (2013) A new species of *Scolecobasidium* associated with the sooty blotch and flyspeck complex on banana from China. Mycol Prog 12:489–495
- Harada H (2013) *Agonimia deguchii* (lichenized *Ascomycota*, *Verrucariaceae*), a new saxicolous species from central Japan. Hikobia 16:307–310
- Harada H (2014) *Cresporaphis chibaensis* sp. nov. (lichenized *Ascomycota*, *Trichosphaeriaceae*) from Chiba-ken, central Japan. Lichenology 12:31–36
- Haranczyk H, Nowak P, Bacior M, Lisowska M et al (2012) Bound water freezing in Antarctic *Umbilicaria aprina* from Schirmacher Oasis. Antarct Sci 24:342–352
- Harimoto Y, Tanaka T, Kodama M, Yamamoto M et al (2008) Multiple copies of AMT2 are prerequisite for the apple pathotype of *Alternaria alternata* to produce enough AM-toxin for expressing pathogenicity. J Gen Plant Pathol 74:222–229
- Harrington TC, Fraedrich SW, Aghayeva DN (2008) *Raffaelea lauricola*, a new ambrosia beetle symbiont and pathogen on the Lauraceae. Mycotaxon 104:399–404
- Harrington TC, Aghayeva DN, Fraedrich SW (2010) New combinations in *Raffaelea*, *Ambrosiella*, and *Hyalarhinocladiella*, and four new species from the redbay ambrosia beetle, *Xyleborus glabratus*. Mycotaxon 111:337–361
- Harrington TC, Yun HY, Lu SS, Goto H et al (2011) Isolations from the redbay ambrosia beetle, *Xyleborus glabratus*, confirm that the laurel wilt pathogen, *Raffaelea lauricola*, originated in Asia. Mycologia 103:1028–1036
- Harrington TC, McNew D, Yun HY (2012) Bur oak blight, a new disease on *Quercus macrocarpa* caused by *Tubakia iowensis* sp. nov. Mycologia 104:79–92
- Harrington TC, McNew D, Mayers C, Fraedrich SW, Reed SE (2014) *Ambrosiella roeperi* sp. nov. is the mycangial symbiont of the granulate ambrosia beetle, *Xylodendrus crassiusculus*. Mycologia 106:835–845
- Harris RC (2009) Four novel lichen taxa in the lichen biota of eastern North America. Opusc Phil 6:149–156

- Harris LB, Rogers SO (2011) Evolution of small putative group I introns in the SSU rRNA gene locus of *Phialophora* species. *BMC Res Notes* 4:258
- Hashimoto A, Sato G, Matsuda T, Hirayama K et al (2015a) Molecular taxonomy of *Dinemasporium* and its allied genera. *Mycoscience* 56:86–101
- Hashimoto A, Sato G, Matsuda T, Matsumura M et al (2015b) Taxonomic revision of *Pseudolachnea* and *Pseudolachnella* and establishment of *Neopseudolachnella* and *Pseudodinemasperium* gen. nov. *Mycologia* 107:383–408
- Hashimoto A, Matsumura M, Hirayama K, Yonezawa H, Tanaka K (2016) Taxonomy and phylogeny of *Cryptocoryneum* (*Pleosporales*, *Dothideomycetes*). *Mycol Prog* 15:45
- Hashimoto A, Matsumura M, Hirayama K, Tanaka K (2017) Revision of *Lophiotremataceae* (*Pleosporales*, *Dothideomycetes*): *Aquasubmersaceae*, *Cryptocoryneaceae*, and *Hermatomycetaceae* fam. nov. *Persoonia* 39:51–73
- Hauck M, Wirth V (2010) Preference of lichens for shady habitats is correlated with intolerance to high nitrogen levels. *Lichenologist* 42:475–484
- Hauck M, Jurgens SR, Huneck S, Leuschner C (2009) High acidity tolerance in lichens with fumarprotocetraric, perlatolic or thamnolic acids is correlated with low pK_{a1} values of these lichen substances. *Environ Pollut* 157:2776–2780
- Hauck M, Tonsberg T, Mayrhofer H, Breuss O (2013a) Lichen-forming and lichenicolous fungi new to Kazakhstan. *Herzogia* 26:103–116
- Hauck M, Tonsberg T, Mayrhofer H, de Bruyn U et al (2013b) New records of lichen species from western Mongolia. *Folia Cryptogam Estonica* 50:13–22
- Hawksworth DL (2015) Proposals to clarify and enhance the naming of fungi under the International Code of Nomenclature for algae, fungi, and plants. *IMA Fungus* 6(1):199–205
- Hawksworth DL (2016) A synopsis of the tropical pleomorphic lichenicolous genus *Koordersiella* (syn. *Hansfordiellopsis*) under a unified nomenclature. *J Fungal Res* 13:227–232
- Hawksworth DL, Etayo J (2011) *Calongea*, *Calongia*, and *Calongomyces* nom. nov. *Lichenologist* 43:275–276
- Hawksworth DL, Millanes AM, Wedin M (2010) *Roselliniella* revealed as an overlooked genus of *Hypocreales*, with the description of a second species on parmelioid lichens. *Persoonia* 24:12–17
- Hawksworth DL, Divakar PK, Crespo A, Teuvo AHTI (2011a) The checklist of parmelioid and similar lichens in Europe and some adjacent territories: additions and corrections. *Lichenologist* 43:639–645
- Hawksworth DL, Santesson R, Tibell L (2011b) *Racoleus*, a new genus of sterile filamentous lichen-forming fungi from the tropics, with observations on the nomenclature and typification of *Cystocoleus* and *Racodium*. *IMA Fungus* 2:71–79
- Hawksworth DL, Aguirre-Hudson B, Ainsworth AM (2014) *Sphinctrina tigillaris*, an overlooked species of *Chaenothecopsis* growing on *Perenniporia meridionalis*, a polypore new to the UK. *Lichenologist* 46:729–735
- Hawksworth DL, van Geel B, Wiltshire PWJ (2016) The enigma of the *Diporotheca* palynomorph. *Rev Palaeobot Palynol* 235:94–98
- Hayashi N, Fukuta Y (2009) Proposal for a new international system of differentiating races of blast (*Pyricularia oryzae* Cavara) by using LTH monogenic lines in rice (*Oryza sativa* L.). *JIRCAS Work Rep* 63:11–15
- Healy RA, Bonito G, Guevara G (2009a) The truffle genus *Pachyphloeus* in the US and Mexico: phylogenetic analysis and a new species. *Mycotaxon* 107:61–71
- Healy RA, Bonito G, Trappe JM (2009b) *Calongea*, a new genus of truffles in the *Pezizaceae* (*Pezizales*). *Anales Jard Bot Madrid* 66:25–32
- Healy RA, Smith ME, Bonito GM, Pfister DH et al (2013) High diversity and widespread occurrence of mitotic spore mats in ectomycorrhizal *Pezizales*. *Mol Ecol* 22:1717–1732
- Healy R, Hobart C, Tocci GLG, Conde IP, Smith ME (2015) Fun with the discomycetes: revisiting collections of Korf's anamorphic *Pezizales* and Thaxter's New England truffles leads to a connection between forms and the description of two new truffle species: *Pachyphloeus pfisteri* and *P. nemoralis*. *Ascomycete Org* 7:357–366
- Healy R, Pfister DH, Rossman AY, Marvanova L, Hansen K (2016) Competing sexual-aseexual generic names of *Pezizomycetes* and recommendations for use. *IMA Fungus* 7:285–288
- Hedayati MT, Arabzadehmoghadam A, Hajheydari Z (2009) Specific IgE against *Alternaria alternata* in atopic dermatitis and asthma patients. *Eur Rev Med Pharmacol Sci* 13:187–191
- Heininger C, Spribille T (2009) The sorediate species of *Xylographa* in Austria (*Baeomycetales*, lichenized Ascomycetes). *Herzogia* 22:129–134
- Heluta VP, Takamatsu S, Harada M, Voytyuk S (2010) Molecular phylogeny and taxonomy of Eurasian *Neoerysiphe* species infecting Asteraceae and Geranium. *Persoonia* 24:81–92
- Hembrom ME, Parihar A, Das K (2013) A new species of *Poronia* from India. *Curr Res Environ Appl Mycol* 3:182–185
- Hengstmengel J (2009) Notes on *Hymenoscyphus*-3: on the nomenclature of *Hymenoscyphus subcarneus* (Ascomycota, Helotiales). *Mycotaxon* 107:267–276
- Heredia G, Castañeda-Ruiz RF, AriasMota RM, Hernández Becerra CI et al (2011) A new species of *Heliocephala* from Mexico with an assessment of the systematic positions of the anamorph genera *Heliocephala* and *Holubovaniella*. *Mycologia* 103:631–640
- Heredia G, Arias RM, Castaneda-Ruiz RF, Gamboa-Angulo M (2012) New species of *Hughesinia* and *Stachybotryna* and new records of anamorphic fungi from the Yucatan Peninsula, Mexico. *Mycol Prog* 11:927–935
- Heredia G, Castañeda-Ruiz RF, Arias RM, Gamboa-Angulo M, De la Rosa S (2013) *Minteriella cenotigena* anam. gen. & sp. nov. from submerged plant material in Mexico. *Mycol Prog* 12:271–275
- Heredia G, Arias RM, Castañeda Ruiz RF, Minter DW (2014) New species of *Lobatopedis* and *Minimelanolocus* (anamorphic fungi) from a Mexican cloud forest. *Nova Hedwig* 98:31–40
- Heredia G, Gamboa-Angulo M, Arias RM, Castañeda-Ruiz RF (2015) Two new species of *Spadicoides* and *Sporidesmiella* from Yucatán, Mexico. *Mycotaxon* 130:511–516
- Hernández-Gutiérrez A (2013) New or rare fungi from eastern Amazonia. 1. *Circinoconiopsis amazonica* gen. and sp. nov. *Mycotaxon* 123:107–111
- Hernández-Restrepo M, Castañeda-Ruiz RF, Gené J, Guarro J, Minter DW, Stadler M (2012a) Microfungi from Portugal: *Minimelanolocus manifestus* sp. nov. and *Vermiculariopsis pediculata* comb. nov. *Mycotaxon* 122:135–143
- Hernández-Restrepo M, Silvera-Simón C, Mena-Portales J, Mercado-Sierra A et al (2012b) Three new species and a new record of *Diplococcum* from plant debris in Spain. *Mycol Prog* 11:191–199
- Hernández-Restrepo M, Mena-Portales J, Gene J, Cano J, Guarro J (2013) New *Bactrodesmiastrum* and *Bactrodesmium* from decaying wood in Spain. *Mycologia* 105:172–180
- Hernández-Restrepo M, Castañeda-Ruiz RF, Gene J, Silvera-Simón C et al (2014a) Two new species of *Solicorynespora* from Spain. *Mycol Prog* 13:157–164

- Hernández-Restrepo M, Gené J, Castañeda-Ruiz RF, Kirk PM, Guarro J (2014b) A new species of *Corynesporopsis* from Spain. *Mycotaxon* 127:155–160
- Hernández-Restrepo M, Groenewald JZ, Crous PW (2015) *Neocordana* gen. nov., the causal organism of Cordana leaf spot of banana. *Phytotaxa* 205:229–238
- Hernández-Restrepo M, Schumacher RK, Wingfield MJ, Ahmad I et al (2016a) Fungal systematics and evolution: FUSE 2. *Sydowia* 68:193–230
- Hernández-Restrepo M, Groenewald JZ, Crous PW (2016b) Taxonomic and phylogenetic re-evaluation of *Microdochium*, *Monographella* and *Idriella*. *Persoonia* 36:57–82
- Hernández-Restrepo M, Gené J, Castañeda-Ruiz RF, Mena-Portales J et al (2017) Phylogeny of saprobic microfungi from Southern Europe. *Stud Mycol* 86:53–97
- Herrera CS, Rossman AY, Samuels GJ, Lechat C, Chaverri P (2013) Revision of the genus *Corallomyctella* with *Corallonectria* gen. nov. for *C. jatropheae* (Nectriaceae, Hypocreales). *Mycosistema* 32:518–544
- Hertel H (2014) A contribution to the family *Lecideaceae* s. lat. (*Lecanoromycetidae* inc. sed., lichenized Ascomycota) in the southern subpolar region; including eight new species and some revised generic circumscriptions. *Lichenologist* 46:389–412
- Hesami S, Khodaparast SA, Zare R (2011) New reports on *Cercospora* and *Cercospora*-like fungi from Guilan Province, Iran. *Iran J Plant Pathol* 47(4):131
- Hestmark G (2009) New observations and records for *Umbilicaria* (Umbilicariaceae) in Bolivia. *Bryologist* 112:833–838
- Hestmark G (2010) Typification of the Andean taxa of *Umbilicaria* described by William Nylander. *Mycotaxon* 111:51
- Hestmark G (2014) Lectotypification of the name *Umbilicaria nylanderiana* (Umbilicariaceae). *Taxon* 63:914–917
- Hestmark G (2016) The lichen genus *Umbilicaria* in Ecuador. *Nordic J Bot* 34:257–268
- Hestmark G, Miadlikovska J, Kauff F, Fraker E et al (2010) Single origin and subsequent diversification of Central Andean endemic *Umbilicaria* species. *Mycologia* 103:45–56
- Hestmark G, Lendemer J, Lumbsch HT, Schmull M et al (2014) A multigene phylogenetic synthesis for the class *Lecanoromycetes* (Ascomycota): 1307 fungi representing 1139 infrageneric taxa, 317 genera and 66 families. *Mol Phylogen Evol* 79:132–168
- Hettiarachchige IK, Ekanayake PN, Mann RC, Guthridge KM et al (2015) Phylogenomics of asexual *Epichloe* fungal endophytes forming associations with perennial ryegrass. *BMC Evol Biol* 15:72
- Heuchert B, Braun U (2014) Two new lichen-associated *Trimmatostroma* species. *Herzogia* 27:227–236
- Heuchert B, Zhurbenko MP, Braun U (2014) Reassessment of the lichenicolous hyphomycete genus *Talpapellis*. *Herzogia* 27:83–92
- Heuchert B, Sancho LG, Braun U, Hawksworth DL (2015) *Bicoloromyces kyffinensis*: a new genus and species of lichen-inhabiting conidial fungi from 83°46'S. *Antarct Sci* 27:421–428
- Hidayat I, Harahap I, Rahayu G (2014) The first phylogenetic study of *Kiliophora* (Fungi, Anamorphic Xylariales). *Mycosphere* 5:78–84
- Higgins NF, Connan S, Stengel DB (2015) Factors influencing the distribution of coastal lichens *Hydropunctaria maura* and *Wahlenbergiella mucosa*. *Mar Ecol* 36:1400–1414
- Hill DJ, Jorgensen PM, Aptroot A (2011) *Ramalodium fecundissimum* Henssen discovered in New Guinea. *Lichenologist* 43:175–177
- Himelbrant DE, Motiejunaitė J, Pykala J, Schiefelbein U, Stepanchikova IS (2013) New records of lichens and allied fungi from the Leningrad Region, Russia. IV. *Folia Cryptogam Est* 50:23–31
- Himelbrant DE, Motiejunaitė J, Stepanchikova IS, Tagirdzhanova GM (2014) New records of lichens and allied fungi from the Leningrad Region, Russia. V. *Folia Cryptogam Est* 51:49–55
- Himelbrant D, Stepanchikova I, Motiejunaite J, Vondrak J et al (2015) New records of lichens and allied fungi from the Leningrad Region, Russia. VI. *Folia Cryptogam Est* 52:21–28
- Himelbrant DE, Stepanchikova IS, Motiejunaitė J, Gagarina LV, Dyomina AV (2016) New records of lichens and allied fungi from the Leningrad Region, Russia. VII. *Folia Cryptogam Est* 53:25–34
- Hipolito E, Faria E, Alves AF, De Hoog GS et al (2009) Alternaria infectoria brain abscess in a child with chronic granulomatous disease. *Eur J Clin Microbiol Infect Dis* 28:377–380
- Hirayama K, Tanaka K (2011) Taxonomic revision of *Lophiostoma* and *Lophiotrema* based on re-evaluation of morphological characters and molecular analyses. *Mycoscience* 52:401–412
- Hirayama K, Tanaka K, Raja HA, Miller AN, Shearer CA (2010) A molecular phylogenetic assessment of *Massarina ingoldiana* sensu lato. *Mycologia* 102:729–746
- Hirayama K, Hashimoto A, Tanaka K (2014) A new species, *Lophiostoma versicolor*, from Japan (Pleosporales, Dothideomycetes). *Mycosphere* 5:411–417
- Hirooka Y, Kobayashi T, Ono T, Rossman AY, Chaverri P (2010) *Verrucostoma*, a new genus in the Bionectriaceae from the Bonin Islands, Japan. *Mycologia* 102:418–429
- Hirooka Y, Rossman AY, Chaverri P (2011) A morphological and phylogenetic revision of the *Nectria cinnabarina* complex. *Stud Mycol* 68:35–56
- Hirooka Y, Rossman AY, Samuels GJ, Lechat C, Chaverri P (2012) A monograph of *Allantonectria*, *Nectria*, and *Pleonectria* (Nectriaceae, Hypocreales, Ascomycota) and their pycnidial, sporodochial, and synnematous anamorphs. *Stud Mycol* 71:1–210
- Hirooka Y, Kawaradani M, Sato T (2014) Description of *Gibellulopsis chrysanthemi* sp. nov. from leaves of garland chrysanthemum. *Mycol Prog* 13:13–19
- Hirooka Y, Tanney JB, Nguyen HD, Seifert KA (2016) Xerotolerant fungi in house dust: taxonomy of *Spiromastix*, *Pseudospromastix* and *Sigleria* gen. nov. in Spiromastigaceae (Oxygenales, Eurotiomycetes). *Mycologia* 108:135–156
- Hiruma J, Okubo M, Kano R, Kumagawa M et al (2016) Mating type gene (MAT) and itraconazole susceptibility of *Trichophyton tonsurans* strains isolated in Japan. *Mycopathologia* 181:441–444
- Hisayoshi T, Shinomura M, Yokokawa K, Kuze I et al (2015) Inhibition of the DNA polymerase and RNase H activities of HIV-1 reverse transcriptase and HIV-1 replication by *Brasenia schreberi* (Junsai) and *Petasites japonicus* (Fuki) components. *J Nat Med* 69:432–440
- Hladki AI, Romero AI (2009) Taxonomic and nomenclatural aspects of *Hypoxylon* taxa from southern South America proposed by Spegazzini. *Mycologia* 101:733–744
- Hodhod MS, Abdel-Wahab MA, Bahkali AH, Hyde KD (2012) *Amarenographium solium* sp. nov. from Yanbu mangroves in the Kingdom of Saudi Arabia. *Cryptog Mycol* 33:285–294
- Hodkinson BP (2010) A first assessment of lichen diversity for one of North America's biodiversity hotspots' in the southern Appalachians of Virginia. *Castanea* 75:126–133
- Hodkinson BP, Lendemer JC (2011a) Molecular analyses reveal semi-cryptic species in *Xanthoparmelia tasmanica*. *Biblioth Lichenol* 106:108–119
- Hodkinson BP, Lendemer JC (2011b) The orders of *Ostropomycetidae* (Lecanoromycetes, Ascomycota): recognition

- of *Sarrameanales* and *Trapeliales* with a request to retain Pertusariales over Agyriales. *Phytologia* 93:407–412
- Hodkinson BP, Lendemer JC (2013) Next-generation sequencing reveals sterile crustose lichen phylogeny. *Mycosphere* 4:1028–1039
- Hofmann TA (2009) Plant parasitic Asterinaceae and Microthyriaceae from the Neotropics (Panama). Johann Wolfgang Goethe-University, Frankfurt
- Hofton TH, Baral HO, Homble K (2009) The ascomycete *Proliferodiscus tricolor* (Sowerby: Fr.) Baral comb. nov., recollected in Scandinavia after 60 years. *Agarica* 28:33–42
- Högnabba F, Stenroos S, Thell A (2009a) Evolution of cyanobacterial symbioses in *Ascomycota*. *Biodiversity and Ecology of lichens. Liber Amicorum Harrie Sipman. Biblioth Lichenol* 99:163–184
- Högnabba F, Stenroos S, Thell A (2009b) Phylogenetic relationship and evolution of photobiont associations in the *Lobariaceae* (*Peltigerales*, *Lecanoromycetes*, *Ascomycota*). *Biblioth Lichenol* 100:157–187
- Högnabba F, Raquel PB, Nordin A, Myllys L, Stenroos S (2014) Phylogenetic position of the crustose *Stereocaulon* species. *Lichenologist* 46:103–114
- Holec J, Kriz M (2013) Current occurrence of *Pseudoplectania melaena* (Fungi, Ascomycota) in the Boubínský Prales National Nature Reserve. *Silva Gabreta* 19:73–80
- Holger TH, Nascimbene J (2008) Contributions toward a new taxonomy of Central European freshwater species of the lichen genus *Thelidium* (*Verrucariales*, *Ascomycota*). *Lichenologist* 40:499–521
- Holt EA, Zemp N, Van Orman M, Perry J et al (2015) Macrolichen substrate selection: Patterns among aspen, non-aspen hardwood, and conifer-dominated forests in the Wasatch Mountains, Utah. *Bryologist* 118:357–366
- Honda K, Kamiyama M, Hirayama K, Menik VA (2012) Taxonomic revision of pleomassariaceous fungi based on phylogenetic analyses of SSU and LSU nrDNA. In: The 55th annual meeting of the mycological society of Japan (<https://www.jstage.jst.go.jp/article/>)
- Hong SK, Kim WG, Lee YK, Choi HW et al (2010) Leaf spot of yam caused by *Pseudophloeosporella dioscoreae* in Korea. *Mycobiology* 38:78–80
- Hong CF, Hsieh HY, Chen CT, Huang HC (2013) Development of a semiselective medium for detection of *Nalanthamala psidii*, causal agent of wilt of guava. *Plant Dis* 97:1132–1136
- Hongsanan S, Bahkali AH, McKenzie EHC, Chukeatirote EHC, Hyde KD (2014a) *Trichopeltinaceae* (*Dothideomycetes*), an earlier name for Brefeldiellaceae, with a new species of *Trichopeltina*. *Phytotaxa* 176:270–282
- Hongsanan S, Chomnunti P, Crous PW, Chukeatirote E, Hyde KD (2014b) Introducing *Chaetothyriothecium*, a new genus of Microthyriales. *Phytotaxa* 161:157–164
- Hongsanan S, Li YM, Liu JK, Hofmann T et al (2014c) Revision of genera in Asterinales. *Fungal Divers* 68:1–68
- Hongsanan S, Hyde KD, Bahkali AH, Camporesi E et al (2015a) Fungal biodiversity profiles 11–20. *Cryptogam Mycol* 36:355–380
- Hongsanan S, Tian Q, Bahkali AH, Yang JB et al (2015b) *Zelotasperisporiales* ord. nov., and two new species of *Zelotasperporium*. *Cryptogam Mycol* 36:301–317
- Hongsanan S, Tian Q, Hyde KD, Chomnunti P (2015c) Two new species of sooty moulds, *Capnodium coffeicola* and *Conidiocarpus plumeriae* in Capodiaceae. *Mycosphere* 6:814–824
- Hongsanan S, Tian Q, Persoh D, Zeng XY et al (2015d) *Meliolales*. *Fungal Divers* 74:91–141
- Hongsanan S, Bahkali AH, Chomnunti P, Liu JK et al (2016a) *Discopycnothyrium palmae* gen. & sp. nov. (*Asterinaceae*). *Mycotaxon* 131(4):859–869
- Hongsanan S, Tian Q, Hyde KD, Hu DM (2016b) The asexual morph of *Trichomerium gloeosporum*. *Mycosphere* 7:1473–1479
- Hongsanan S, Sánchez-Ramírez S, Crous PW, Ariyawansa HA et al (2016c) The evolution of fungal epiphytes. *Mycosphere* 7:1690–1712
- Hongsanan S, Maharanachikumbura SS, Hyde KD, Samarakoon MC et al (2017a) An updated phylogeny of Sordariomycetes based on phylogenetic and molecular clock evidence. *Fungal Divers* 84:25–41
- Hongsanan S, Zhao RL, Hyde KD (2017b) A new species of *Chaetothyrina* on branches of mango, and introducing *Phaeothecoidiellaceae* fam. nov. *Mycosphere* 8:137–146
- Horsfield A, Wicks T, Davies K, Wilson D, Paton S (2010) Effect of fungicide use strategies on the control of early blight (*Alternaria solani*) and potato yield. *Australas Plant Pathol* 39:368–375
- Hosagoudar VB (2012) Asterinales of India. *Mycosphere* 3:617–852
- Hosagoudar VB (2013) My contribution to the fungal knowledge of India. *J Threat Taxa* 5:4129–4348
- Hosagoudar VB, Archana GR (2009) Host range of meliolaceous fungi in India. *J Threat ned Taxa* 1:269–282
- Hosagoudar VB, Archana GR (2010a) A new species of the genus *Ectendomeliola* (*Meliolaceae*) from Kerala, India. *J Threat Taxa* 2:1092–1095
- Hosagoudar VB, Archana GR (2010b) *Meliolaceae* of Kerala, India—XXXI. *J Threat Taxa* 2:889–891
- Hosagoudar VB, Chandraprabha A (2009a) *Bramhamyces*, a new anomorphic genus from India. *Indian J Sci Technol* 2:17
- Hosagoudar VB, Chandraprabha A (2009b) Two new *Prillieuxina* species from Kerala, India. *Indian J Sci Technol* 2:18
- Hosagoudar VB, Riju MC (2009) Studies on foliicolous fungi-XXXI: the genus *Schiffnerula* and *Sarcinella*. *Indian J Sci Technol* 2:7–8
- Hosagoudar VB, Harish M (2010) Vishnumyces a new genus of the family Asterinaceae. *Indian Phytopathol* 63:85–86
- Hosagoudar VB, Sabeena A (2010) Five new *Meliolaceae* members from Kerala, India. *Biosci Discov* 2:117–121
- Hosagoudar VB, Riju MC (2011) *Sarcinella tamarindi* sp. nov. from Kerala, India. *Mycosphere* 2:157–160
- Hosagoudar VB, Robin PJ (2011) Five new black mildews from the Western Ghats of Peninsular India. *Biosci Discov* 2:264–267
- Hosagoudar V, Divya B (2013) Additions to black mildews. *IJPBAS* 2:430–438
- Hosagoudar VB, Archana GR, Dan M (2009) *Maheshwaramyces*, a new genus of the family Lembosiaceae. *Indian J Sci Technol* 2:12–13
- Hosagoudar VB, Sabeena A, Riju M (2010a) *Bheemamyces*, a new genus of the family Asterinaceae (Ascomycetes). *J Threat Taxa* 2:1323–1324
- Hosagoudar VB, Sabeena A, Riju MC (2010b) *Sarcinella limoniae* sp. nov. from Kerala, India. *Indian Phytopathol* 63:236–237
- Hosagoudar VB, Thomas J, Agarwal DK (2010c) Additions to *Meliolales* of India. *Nelumbo* 52:1–8
- Hosagoudar VB, Thimmaiah CJ, Jayashankara M (2011a) Schiffnerulaceous fungi of Kodagu, Karnataka, India. *J Threat Taxa* 3:2268–2271
- Hosagoudar VB, Thomas J, Agarwal DK (2011b) Two new schiffnerulaceous fungi from Kerala, India. *J Yeast Fungal Res* 2:85–87
- Hosagoudar VB, Sabeena A, Divya B (2013) CEPF Western Ghats special series: new fungi from Kerala, India. *J Threat Taxa* 5:4805–4807
- Hosagoudar VB, Mathew SP, Babu D (2014) Foliicolous fungi of Andaman Islands. *J Threat Taxa* 6:5447–5463

- Hosoya T, Sasagawa R, Hosaka K, Gi-Ho S et al (2010) Molecular phylogenetic studies of *Lachnum* and its allies based on the Japanese material. *Mycoscience* 51:170–181
- Hosoya T, Han JG, Sung GH, Hirayama Y et al (2011) Molecular phylogenetic assessment of the genus *Hyphodiscus* with description of *Hyphodiscus hyaloscopoides* sp. nov. *Micol Prog* 10:239–248
- Hou CL, Piepenbring M (2009a) Two new *Rhytismatales* on *Rhododendron* from China. *Mycologia* 101:383–389
- Hou CL, Piepenbring M (2009b) New species and new records of *Rhytismatales* from Panama. *Mycologia* 101:565–572
- Hou CL, Trampe T, Piepenbring M (2010) A new species of *Rhytisma* causes tar spot on *Comarostaphylis arbutooides* (Ericaceae) in Panama. *Mycopathologia* 169:225–229
- Hou XM, Xu RF, Gu YC, Wang CY, Shao CL (2015) Biological and chemical diversity of coral- derived microorganisms. *Curr Med Chem* 22:3707–3762
- Hou LW, Liu F, Duan WJ, Cai L (2016) *Colletotrichum aracearum* and *C. camelliae-japonicae*, two holomorphic new species from China and Japan. *Mycosphere* 7:1111–1123
- Houbraken J, Samson R (2011) Phylogeny of *Penicillium* and segregation of *Trichocomaceae* into three families. *Stud Mycol* 70:1–51
- Houbraken J, Spierenburg H, Frisvad JC (2012) *Rasamonia*, a new genus comprising thermotolerant and thermophilic *Talaromyces* and *Geosmithia* species. *Antonie Van Leeuwenhoek* 101:403–421
- Houbraken J, Giraud S, Meijer M, Bertout S et al (2013) Taxonomy and antifungal susceptibility of clinically important *Rasamonia* species. *J Clin Microbiol* 51:22–30
- Hoyt JR, Cheng TL, Langwig KE, Hee MM et al (2015) Bacteria isolated from bats inhibit the growth of *Pseudogymnoascus destructans*, the causative agent of white-nose syndrome. *PLoS ONE* 10:e0121329
- Hsieh HM, Ju YM, Hsueh PR, Lin HY et al (2009) Fungal keratitis caused by a new filamentous hyphomycete *Sagenomella keratitidis*. *Bot Stud* 50:331–335
- Hsieh HM, Lin CR, Fang MJ, Rogers JD et al (2010) Phylogenetic status of *Xylaria* subgenus *Pseudoxylaria* among taxa of the subfamily *Xylarioideae* (*Xylariaceae*) and phylogeny of the taxa involved in the subfamily. *Mol Phylogen Evol* 54:957–969
<http://www.speciesfungorum.org/Names/SynSpecies.asp?RecordID=504433>
- Hu D, Cai L, Chen H, Bahkali AH, Hyde KD (2010a) Fungal diversity on submerged wood in a tropical stream and an artificial lake. *Biodivers Conserv* 19:3799–3808
- Hu DM, Cai L, Chen H, Bahkali AH, Hyde KD (2010b) Four new freshwater fungi associated with submerged wood from Southwest Asia. *Sydotwia* 62:191–203
- Hu DM, Cai L, Bahkali AH, Hyde KD (2012a) Two new freshwater species of *Annulatasaceae* from China. *Mycotaxon* 120:81–88
- Hu DM, Chen H, Cai L, Bahkali AH, Hyde KD (2012b) *Aquapeziza*: a new genus from freshwater and its morphological and phylogenetic relationships to *Pezizaceae*. *Mycologia* 104:540–546
- Hu DM, Cai L, Hyde KD (2012c) Three new Ascomycetes from freshwater in China. *Mycologia* 104:1478–1489
- Hu J, Chen C, Peever T, Dang H et al (2012d) Genomic characterization of the conditionally dispensable chromosome in *Alternaria arborescens* provides evidence for horizontal gene transfer. *BMC Genomics* 13:1
- Hua-Jie L, Zhen C, QingFeng W (2010) Differences in enrichment capability of Co, Cr, Pb, Cu among five Antarctic lichens. *Mycosistema* 29:719–725
- Hua H, Luo H, Bai Y, Wang K et al (2014) A thermostable glucoamylase from *Bispora* sp. MEY-1 with stability over a broad pH range and significant starch hydrolysis capacity. *PLoS ONE* 9:e113581
- Huang MR (2010) Altitudinal patterns of *Stereocaulon* (lichenized Ascomycota) in China. *Acta Oecol* 36:173–178
- Huang YT, Chen CY (2014) *Leptographium globosum* sp. nov., a new species with globose conidia. *Mycol Prog* 13:841–848
- Huang CH, Pan JH, Chen B, Yu M et al (2011) Three bianthraquinone derivatives from the mangrove endophytic fungus *Alternaria* sp. ZJ-9-B from the South China Sea. *Mar Drugs* 9:832–843
- Huang F, Hou X, Dewdney MM, Fu Y et al (2013) *Diaporthe* species occurring on citrus in China. *Fungal Divers* 61(1):237–250
- Huang G, Guo L, Liu N (2014) *Xylaria byttneriae* sp. nov. from Yunnan Province in China. *Mycosistema* 33:567–570
- Huang H, Wang M, Zhang L, Yue HM et al (2015a) *Pochonia cordycepsociata*, a new species associated with Chinese cordyceps in Tibet, China. *Phytotaxa* 208:278–286
- Huang Y, Busk PK, Herbst FA, Lange L (2015b) Genome and secretome analyses provide insights into keratin decomposition by novel proteases from the non-pathogenic fungus *Onygena corvine*. *Appl Microbiol Biotechnol* 99:9635–9649
- Huang SK, Tangthirasunun N, Phillips AJ, Dai DQ et al (2016a) Morphology and phylogeny of *Neoscytalidium orchidacearum* sp. nov. (Botryosphaeriaceae). *Mycobiology* 44:79–84
- Huang HP, Ma S, Huang JH, Zheng JL, Yi KX (2016b) Whole genome re-sequencing and transcriptome analysis of the Stylosanthes Anthracnose pathogen *Colletotrichum gloeosporioides* reveal its characteristics. *Mycosphere* 7:1124–1146
- Hubka V, Dobiasova S, Lyskova P, Mallatova N et al (2013a) *Auxarthron ostraviense* sp. nov., and *A. umbrinum* associated with non-dermatophytic onychomycosis. *Med Mycol* 51:614–624
- Hubka V, Kolarik M, Kubatova A, Peterson SW (2013b) Taxonomic revision of *Eurotium* and transfer of species to *Aspergillus*. *Mycologia* 105:912–937
- Hubka V, Peterson SW, Frisvad JC, Yaguchi T et al (2013c) *Aspergillus waksmanii* sp. nov. and *Aspergillus marvanovae* sp. nov., two closely related species in section Fumigati. *Int J Syst Evol Microbiol* 63:783–789
- Hubka V, Cmokova A, Skorepova M, Mikula P, Kolarik M (2014) *Trichophyton onychocola* sp. nov. isolated from human nail. *Med Mycol* 52:285–292
- Hubka V, Dobiasova S, Dobias R, Kolarik M (2014a) *Microsporum aenigmaticum* sp. nov. from *M. gypseum* complex, isolated as a cause of tinea corporis. *Med Mycol* 52:387–396
- Hubka V, Reblova M, Rehulka J, Selbmann L et al (2014b) *Bradymyces* gen. nov. (*Chaetothyriales, Trichomeriaceae*), a new ascomycete genus accommodating poorly differentiated melanized fungi. *Antonie Van Leeuwenhoek* 106:979–992
- Hubka V, Novakova A, Kolarik M, Jurjevic Z, Peterson SW (2015) Revision of *Aspergillus* section Flavipes: seven new species and proposal of section Jani sect. nov. *Mycologia* 107:169–208
- Hubka V, Novakova A, Peterson SW, Frisvad JC et al (2016a) A reappraisal of *Aspergillus* section Nidulantes with descriptions of two new sterigmatocystin-producing species. *Plant Syst Evol* 302:1267–1299
- Hubka V, Novakova A, Samson RA, Houbraken J et al (2016b) *Aspergillus europaeus* sp. nov., a widely distributed soil-borne species related to *A. wentii* (section Cremei). *Plant Syst Evol* 302:641–650
- Hughes SJ, Seifert KA (2012) Taxonomic and nomenclatural notes on sooty mould names based on species mixtures: *Hormiscium handelii* and *Torula lechleriana*. *Mycoscience* 53:17–24
- Hughes SJ, Atkinson TJ, Seifert KA (2012) New Zealand fungi 37: two new species of the sooty mould genus *Metacapnodium* with dictyoseptate ascospores. *NZ J Bot* 50:381–387

- Huhndorf SM, Miller AN (2008) A new species of *Camarops* and phylogenetic analysis of related taxa in the Boliniaceae. *N Am Fungi* 3:231–239
- Huhndorf SM, Miller AN (2011) A molecular re-appraisal of taxa in the *Sordariomycetidae* and a new species of *Rimaconus* from New Zealand. *Stud Mycol* 68:203–210
- Huhndorf SM, Greif M, Mugambi GK, Miller AN (2008) Two new genera in the *Magnaportheaceae*, a new addition to *Ceratosphaeria* and two new species of *Lentomitella*. *Mycologia* 100:940–955
- Huhndorf SM, Miller AN, Greif M, Samuels GJ (2009) *Amplistroma* gen. nov. and its relation to *Wallrothiella*, two genera with globose ascospores and acrodontium-like anamorphs. *Mycologia* 101:904–919
- Huhtinen S, Hawksworth DL, Ihlen PG (2008) Observations of two glassy-haired lichenicolous discomycetes. *Lichenologist* 40:549–557
- Huhtinen S, Laukka T, Dobbeler P, Stenroos S (2010) Six novelties to European bryosymbiotic discomycetes. *Nova Hedw* 90:413–431
- Hui F, Ren Y, Chen L, Li Y et al (2014) Molecular phylogenetic analysis reveals the new genus *Hemisphaericaspora* of the family Debaryomycetaceae. *PLoS ONE* 9:e103737
- Hujsllová M, Kubatova A, Kostovcik M, Kolarik M (2013) *Acidiella bohemica* gen. et sp. nov. and *Acidomyces* spp. (*Teratosphaeriaceae*), the indigenous inhabitants of extremely acidic soils in Europe. *Fungal Divers* 58:33–45
- Hujsllová M, Kubatova A, Kostovcik M, Blanchette RA, de Beer ZW (2014) Three new genera of fungi from extremely acidic soils. *Mycol Progr* 13:819–831
- Humber RA, Rocha LF, Inglis PW, Kipnis A, Luz C (2013) Morphology and molecular taxonomy of Evlachovaea-like fungi, and the status of this unusual conidial genus. *Fungal Biol* 117:1–12
- Hunter GC, Crous PW, Carnegie AJ, Burgess TI, Wingfield MJ (2011) *Mycosphaerella* and *Teratosphaeria* diseases of *Eucalyptus*; easily confused and with serious consequences. *Fungal Divers* 50:145
- Hüseyin E, Selcuk F, Churakov BP (2014) A new species of *Berkleasmium* from Ulyanovsk, Russia. *Mycosphere* 5:462–466
- Hustad VP, Miller AN (2011) Phylogenetic placement of four genera within the *Leotiomycetes*. *N Am Fungi* 6:1–13
- Hustad VP, Miller AN (2015) *Maasoglossum*, a basal genus in *Geoglossomycetes*. *Mycoscience* 56:572–579
- Hustad VP, Miller AN, Moingeon JM, Priou JP (2011) Inclusion of *Nothomitra* in *Geoglossomycetes*. *Mycosphere* 2:646–654
- Hustad VP, Miller AN, Dentinger BT, Cannon PF (2013) Generic circumscriptions in *Geoglossomycetes*. *Persoonia* 31:101–111
- Hwang J, Zhao Q, Yang ZL, Wang Z, Townsend JP (2015) Solving the ecological puzzle of mycorrhizal associations using data from annotated collections and environmental samples—an example of saddle fungi. *Environ Microbiol Rep* 7:658–667
- Hyde KD, Fröhlich J, Taylor JE (1998) Fungi from palms. XXXVI. Reflections on unitunicate ascomycetes with apiospores. *Sydowia* 50:21–80
- Hyde KD, Cai L, Cannon PF, Crouch JA, Crous PW et al (2009) *Colletotrichum*—names in current use. *Fungal Divers* 39:147–182
- Hyde KD, McKenzie EHC, Koko TW (2011) Towards incorporating anamorphic fungi in a natural classification—checklist and notes for 2010. *Mycosphere* 2:1–88
- Hyde KD, Jones EBG, Liu JK, Ariyawansa H et al (2013) Families of *Dothideomycetes*. *Fungal Divers* 63:1–313
- Hyde KD, Nilsson RH, Alias SA, Ariyawansa HA et al (2014) One stop shop: backbones trees for important pytopathogenic genera: I. *Fungal Divers* 67:21–125
- Hyde KD, Hongsanan S, Jeewon R, Bhat DJ et al (2016) Fungal diversity notes 367–490: taxonomic and phylogenetic contributions to fungal taxa. *Fungal Divers* 80:1–270
- Hyde KD, Maharachchikumbura SS, Hongsanan S, Samarakoon MC et al (2017) The ranking of fungi: a tribute to David L. Hawksworth on his 70th birthday. *Fungal Divers* 84:1–23
- Iftikhar S, Asad S, Mujeeb-Kazi A, Ahmad I (2009a) In vitro screening of synthetic hexaploid wheat lines against *Bipolaris sorokiniana* in Pakistan. *Pak J Bot* 41:1989–2001
- Iftikhar S, Asad S, Munir A, Sultan A, Ahmad I (2009b) Hosts of *Bipolaris sorokiniana*, the major pathogen of spot blotch of wheat in Pakistan. *Pak J Bot* 41:1433–1436
- Imazaki A, Tanaka A, Harimoto Y, Yamamoto M et al (2010) Contribution of peroxisomes to secondary metabolism and pathogenicity in the fungal plant pathogen *Alternaria alternata*. *Eukaryotic cell* 9:682–694
- Imoulan A, Wu HJ, Lu WL, Li Y et al (2016) *Beauveria medogensis* sp. nov., a new fungus of the entomopathogenic genus from China. *J Invertebr Pathol* 139:74–81
- Inácio CA, Cannon PF (2008) The genera of the ‘*Parmulariaceae*’. CBS Fungal Biodiversity Centre, Utrecht
- Inácio CA, Araúz K, Piepenbring M (2012) A new genus of *Parmulariaceae* from Panama. *Mycol Prog* 11:1–6
- Index Fungorum (2013) <http://www.indexfungorum.org/names/NAMESRecord.asp?RecordID=550246>
- Index Fungorum (2016) <http://www.indexfungorum.org>. Accessed in December 2016
- Inuma T, Khodaparast SA, Takamatsu S (2007) Multilocus phylogenetic analyses within *Blumeria graminis*, a powdery mildew fungus of cereals. *Molecular Phylogenetics and Evolution* 44:741–751
- Irinyi L, Serena C, Garcia-Hermoso D, Arabatzis M et al (2015) International Society of Human and Animal Mycology (ISHAM)-ITS reference DNA barcoding database—the quality controlled standard tool for routine identification of human and animal pathogenic fungi. *Med Mycol* 53:313–337
- Isaka M, Yangchum A, Intamas S, Kocharin K et al (2009) Aigialomycins and related polyketide metabolites from the mangrove fungus *Aigialus parvus* BCC 5311. *Tetrahedron* 65:4396–4403
- Isaka M, Palasarn S, Prathumpai W, Laksanacharoen P (2011) Pimarane diterpenes from the endophytic fungus *Eutypella* sp. BCC 13199. *Chem Pharm Bull (Tokyo)* 59:1157–1159
- Isaka M, Chinthanom P, Rachtaewee P, Srichomthong K et al (2015) Cytotoxic hydroanthraquinones from the mangrove-derived fungus *Paradicthyothrinium diffractum* BCC 8704. *Journal of Antibiotics* 68:334–338
- Ishihara A, Nakao T, Mashimo Y, Murai M et al (2011) Probing the role of tryptophan-derived secondary metabolism in defense responses against *Bipolaris oryzae* infection in rice leaves by a suicide substrate of tryptophan decarboxylase. *Phytochemistry* 72:7–13
- Ishiwada N, Takeshita K, Yaguchi T, Nagasawa K et al (2016) The first case of invasive mixed-mold infections due to *Emericella nidulans* var. *echinulata* and *Rasamonia piperina* in a patient with chronic granulomatous disease. *Mycopathologia* 181:305–309
- Isiloglu M, Alli H, Spooner BM, Solak MH (2010) *Morchella anatolica* (Ascomycota), a new species from southwestern Anatolia, Turkey. *Mycologia* 102:455–458
- Ismed F, Lohézic-Le Dévéhat F, Delalande O, Sinbandhit S et al (2012) Lobarin from the Sumatran lichen, *Stereocaulon halei*. *Fitoterapia* 83:1693–1698
- Isola D, Selbmann L, de Hoog GS, Fenice M et al (2013) Isolation and screening of black fungi as degraders of volatile aromatic hydrocarbons. *Mycopathologia* 175:369–379
- Isola D, Zucconi L, Onofri S, Caneva G et al (2016) Extremotolerant rock inhabiting black fungi from Italian monumental sites. *Fungal Divers* 76:75–96

- Iturriaga T, Fernandez R, Castaneda-Ruiz RF, Minter DW, da Cruz AC (2013) *Scolecobeltrania*, an interesting new microfungus from Venezuela. *Mycotaxon* 124:143–148
- Iwen PC, Schutte SD, Florescu DF, Noel-Hurst RK, Sigler L (2012) Invasive *Scopulariopsis brevicaulis* infection in an immunocompromised patient and review of prior cases caused by *Scopulariopsis* and *Microascus* species. *Med Mycol* 50:561–569
- Jagadeesh Ram TAM, Sinha GP (2016) A world key to *Cryptothecia* and *Myriostigma* (Arthoniaceae), with new species and new records from the Andaman and Nicobar Islands, India. *Phytotaxa* 266:103–114
- Jagielski T, Sandoval-Denis M, Yu J, Yao L et al (2016) Molecular taxonomy of scopulariopsis-like fungi with description of new clinical and environmental species. *Fungal Biol* 120:586–602
- Jaklitsch WM, Voglmayr H (2011) *Stromatonectria* gen. nov. and notes on *Myrmeciciella*. *Mycologia* 103:431–440
- Jaklitsch WM, Voglmayr H (2012) Phylogenetic relationships of five genera of Xylariales and *Rosaspheeria* gen. nov. (*Hypocreales*). *Fungal Divers* 52:75–98
- Jaklitsch WM, Voglmayr H (2015) Biodiversity of *Trichoderma* (*Hypocreaceae*) in Southern Europe and Macaronesia. *Stud Mycol* 80:1–87
- Jaklitsch WM, Voglmayr H (2016) Hidden diversity in *Thyridaria* and a new circumscription of the *Thyridariaceae*. *Stud Mycol* 85:35–64
- Jaklitsch WM, Voglmayr H (2017) Three former taxa of *Cucurbitaria* and considerations on *Petrakia* in the *Melanommataceae*. *Sydowia* 69:81–95
- Jaklitsch W, Poldmaa K, Samuels GJ (2008) Reconsideration of *Protocrea* (*Hypocreales*, *Hypocreaceae*). *Mycologia* 100:962–984
- Jaklitsch WM, Reblova M, Voglmayr H (2013) Molecular systematics of *Woswasia atropurpurea* gen. et sp. nov. (*Sordariomycetidae*), a fungicolous ascomycete with globose ascospores and holoblastic conidiogenesis. *Mycologia* 105:476–485
- Jaklitsch WM, Fournier J, Rogers JD, Voglmayr H (2014a) Phylogenetic and taxonomic revision of *Lopadostoma*. *Persoonia* 32:52–82
- Jaklitsch WM, Lechat C, Voglmayr H (2014b) The rise and fall of *Sarawakus* (*Hypocreaceae*, *Ascomycota*). *Mycologia* 106:133–144
- Jaklitsch WM, Fournier J, Dai DQ, Hyde KD, Voglmayr H (2015) *Valsaria* and the *Valsariales*. *Fungal Divers* 73:159–202
- Jaklitsch WM, Voglmayr H (2014) Persistent hamathelial threads in the *Nectriaceae*, *Hypocreales*: *Thyronectria* revisited and reinstated. *Persoonia* 33:182–211
- Jaklitsch W, Baral HO, Lücking R, Lumbsch HT (2016a) *Ascomycota*. In: Frey W (ed) Syllabus of Plant Families, 23rd edn. Borntraeger Science Publishers, Stuttgart
- Jaklitsch WM, Gardiennet A, Voglmayr H (2016b) Resolution of morphology-based taxonomic delusions: *Acrocordiella*, *Basiseptospora*, *Blogiascospora*, *Clypeosphaeria*, *Hymenoplella*, *Lepteutypa*, *Pseudapiospora*, *Requienella*, *Seiridium* and *Strickeria*. *Persoonia* 37:82–105
- Jaklitsch WM, Olariaga I, Voglmayr H (2016c) *Teichospora* and the *Teichosporaceae*. *Mycol Prog* 15:1–20
- Jamshidnejad V, Sahebani N Etebarian H (2013) Potential biocontrol activity of *Arthrobotrys oligospora* and *Trichoderma harzianum* BI against *Meloidogyne javanica* on tomato in the greenhouse and laboratory studies. *Archives of Phytopathology and Plant Protection* 46:1632–1640
- Jankowiak R, Rossa R (2008) Associations between *Pityogenes bidentatus* and fungi in young managed Scots pine stands in Poland. *Forest Pathol* 38:169–177
- Janni M, Sella L, Favaron F, Blechl AE et al (2008) The expression of a bean PGIP in transgenic wheat confers increased resistance to the fungal pathogen *Bipolaris sorokiniana*. *Mol Plant-Microbe Interact* 21:171–177
- Jayalal U, Aptroot A, Hur JS (2012a) The lichen genus *Polychidium* new to South Korea. *Mycobiology* 40:252–254
- Jayalal U, Wolseley P, Gueidan C, Aptroot A et al (2012b) *Anzia mahaelyensis* and *Anzia flavotenuis*, two new lichen species from Sri Lanka. *Lichenologist* 44:381–389
- Jayalal U, Aptroot A, Nguyen TT, Dzung NA et al (2013a) Further additions to the macrolichen mycota of Vietnam. *Mycotaxon* 124:51–59
- Jayalal U, Oh SO, Lücking R, Joshi S et al (2013b) Contributions to the foliicolous lichens flora of South Korea. *Mycobiology* 41:202–209
- Jayalal U, Divakar PK, Joshi S, Oh SO et al (2014) Overview of *Xanthoparmelia* taxa from South Korea including the description of two new species (*Parmeliaceae*, *Ascomycota*). *Phytotaxa* 181:96–109
- Jayanetti DR, Yue Q, Bills GF, Gloer JB (2014) Hypocoprins A-C: new sesquiterpenoids from the coprophilous fungus *Hypocopra rostrata*. *J Nat Prod* 78:396–401
- Jayasiri SC, Wanasinghe DN, Ariyawansa HA, Jones EB et al (2015) Two novel species of *Vagicola* (*Phaeosphaeriaceae*) from Italy. *Mycosphere* 6:716–728
- Jayasiri SC, Ariyawansa HA, Jones EBG, Kang JC (2016a) Towards a natural classification of *Dothideomycetes*: 8. The genera *Cocconia*, *Dianesea*, *Endococcus* and *Lineostroma*. *Phytotaxa* 255:66–74
- Jayasiri SC, Hyde KD, Jeewon R, Bhat JD et al (2016b) *Neoccultibambusa jonesii*, a novel taxon within *Occultibambusaceae*. *Mycosphere* 7:1458–1472
- Jayasiri SC, Jones EBG, Kang JC, Promputtha I et al (2016c) A new species of genus *Anteaglonia* (*Anteagloniaceae*, *Pleosporales*) with its asexual morph. *Phytotaxa* 263:233–244
- Jayawardena RS, Ariyawansa HA, Singtripop C, Li YM et al (2014) A re-assessment of *Elsinoaceae* (*Myriangiales*, *Dothideomycetes*). *Phytotaxa* 176:120–138
- Jayawardena RS, Zhang W, Liu M, Maharachchikumbura SSN et al (2015) Identification and characterization of *Pestalotiopsis*-like fungi related to grapevine diseases in China. *Fungal Biol* 119:348–361
- Jayawardena RS, Liu M, Maharachchikumbura SSN, Zhang W et al (2016a) *Neopestalotiopsis vitis* sp. nov. causing grapevine leaf spot in China. *Phytotaxa* 258:63–74
- Jayawardena RS, Hyde KD, Damm U, Cai L et al (2016b) Notes on currently accepted species of *Colletotrichum*. *Mycosphere* 7:1192–1260
- Jayawardena RS, Hyde KD, Jeewon R, Li XH et al (2016c) Mycosphere essay 6: Why is it important to correctly name *Colletotrichum* species? *Mycosphere* 7:1076–1092
- Jayawardena RS, Huang JK, Jin BC, Yan JY et al (2016d) An account of *Colletotrichum* species associated with strawberry anthracnose in China based on morphology and molecular data. *Mycosphere* 7:1147–1163
- Jayawardena RS, Li XH, Liu M, Zhang W, Yan JY (2016e) Mycosphere Essay 16: *Colletotrichum*: Biological control, bio-catalyst, secondary metabolites and toxins. *Mycosphere* 7:1164–1176
- Jeyaprakash A, Davison DA, Schubert TS (2014) Molecular detection of the laurel wilt fungus, *Raffaelea lauricola*. *Plant Dis* 98:559–564

- Jia GJ, Lin YR, Hou CL (2011) A new species of *Cocomyces* (*Rhytismatales*, Ascomycota) from Mt Huangshan, China. *Mycotaxon* 118:231–235
- Jia C, Zhang L, Liu L, Wang J, Li C, Wang Q (2013) Multiple phytohormone signalling pathways modulate susceptibility of tomato plants to *Alternaria alternata* f. sp. *lycopersici*. *J Exp Bot* 64:637–650
- Jiang YL, Zhang TY (2008) New species of *Bipolaris*, *Scolecobasidium* and *Torula* from soil. *Mycotaxon* 104:135–140
- Jiang YL, Zhang TY (2010) Two new species of *Humicola* from soil. *Mycotaxon* 111:117–120
- Jiang YL, Xu JJ, Wu YM, Zhang YL et al (2011) Studies on *Cephalotrichum* from soils in China twelve new species and two new combinations. *Mycotaxon* 117:207–225
- Jiao JIA, Xing JH, Dong JG, Han JM, Liu JS (2011) Functional analysis of MYB73 of *Arabidopsis thaliana* against *Bipolaris oryzae*. *Agric Sci China* 10:721–727
- Jie CY, Zhou QX, Zhao WAS, Jiang YL et al (2013) A new *Myrmecidium* species from Guizhou, China. *Mycotaxon* 124:1–8
- Jindamorakot S, Am-in S, Yukphan P, Yamada Y (2011) *Limtongia* gen. nov. for *Zygozyma smithiae* (Lipomycetaceae). *Ann Microbiol* 61:689–693
- Jindamorakot S, Yukphan P, Yamada Y (2012) *Kockiozyma* gen. nov., for *Zygozyma suomiensis*: the phylogeny of the Lipomycetaceous yeasts. *Ann Microbiol* 62:1831–1840
- Johnson D, Sung GH, Hywel-Jones NL, Luangsa-ard JJ et al (2009) Systematics and evolution of the genus *Torrubiella* (Hypocreales, Ascomycota). *Mycol Res* 113:279–289
- Johnston PR, Park D, Manning MA (2010) *Neobulgaria alba* sp. nov. and its Phialophora-like anamorph in native forests and kiwifruit orchards in New Zealand. *Mycotaxon* 113:385–396
- Johnston PR, Park D (2013) The phylogenetic position of *Lanzia berggrenii* and its sister species. *Mycosistema* 32:366–385
- Johnston PR, Steel JB, Park D, Ludwig LR (2013) A bryophilous member of the *Leotiomycetes* from New Zealand, *Bryoclavicular campylopi* gen. et sp. nov. *NZ J Bot* 51:321–327
- Johnston PR, Park D, Baral HO, Galán R et al (2014a) The phylogenetic relationships of *Torreidiella* and *Hymenotorrendiella* gen. nov. within the *Leotiomycetes*. *Phytotaxa* 177:1–25
- Johnston PR, Seifert KA, Stone JK, Rossman AY, Marvanova L (2014b) Recommendations on generic names competing for use in *Leotiomycetes* (Ascomycota). *IMA Fungus* 5:91–120
- Jones EG, Klaysuban A, Pang KL (2008) Ribosomal DNA phylogeny of marine anamorphic fungi: *Cumulospora varia*, *Dendryphiella* species and *Orbimyces spectabilis*. *Raffles Bull Zool* 19(suppl):11–18
- Jones EBG, Sakayaroj J, Suetrong S, Somrithipol S, Pang KL (2009a) Classification of marine Ascomycota, anamorphic taxa and Basidiomycota. *Fungal Divers* 35:1–187
- Jones EBG, Zuccaro A, Nakagiri A, Mitchell JL, Pang KL (2009b) Phylogenetic relationships of the genus *Sigmoidea* and a new genus *Halosigmoidea* gen. nov. *Botany* 52:349–359
- Jones EBG, Hyde KD, Pang KL, Suetrong S (2012) Phylogeny of the Dothideomycetes and other classes of marine Ascomycota. In: Jones EBG, Pang KL (eds) Marine Fungi and Fungal-like Organisms. De Gruyter, Berlin, pp 17–34
- Jones EBG, Suetrong S, Cheng WH, Rungjindamai N et al (2014) An additional fungal lineage in the Hypocreomycetidae (Falcocladiaceae) and the taxonomic revaluation of *Chaetosphaeria chaetosa* and *Swampomyces* species, based on morphology, ecology and phylogeny. *Cryptog Mycol* 35:119–138
- Jones EBG, Suetrong S, Sakayaroj J, Bahkali AH et al (2015) Classification of marine Ascomycota, Basidiomycota, Blastocladiomycota and Chytridiomycota. *Fungal Divers* 73:1–72
- Jones EBG, To-anun C, Suetrong S, Boonyuen N (2016) Mycosphere Essays 12. Progress in the classification of the water-cooling tower ascomycete *Savoryella* and a tribute to John Savory: a review. *Mycosphere* 7:570–581
- Jørgensen PM (2008) *Vahliella*, a new lichen genus. *Lichenologist* 40:221–225
- Jørgensen PM (2010) New discoveries in the lichen genus *Staurolemma* Korber. *Nova Hedw* 90:153–159
- Jørgensen PM (2014) Taxonomy and nomenclature of *Collema fasciculare* (L.) G.H. Weber. *Lichenologist* 46:594
- Jørgensen PM, Nordin A (2009) Lichens in Scandinavia known mainly from Norwegian type-specimens. *Graphis Scripta* 21:1–20
- Jørgensen PM, Palice Z (2010) Additions to the lichen family Pannariaceae in Ecuador. *Nordic J Bot* 28:623–628
- Jørgensen PM, Gjerde I (2012) Notes on some pannariaceous lichens from New Caledonia. *Cryptog Mycol* 33:3–9
- Jørgensen PM, Palice Z (2015) Additions to the cyanolichens of the Andes. *Herzogia* 28:285–287
- Jorgensen PM, Andersen HL (2015) The lichen genus *Psoromidium* (Pannariaceae) re-evaluated, with nomenclatural notes on *Degelia* and *Psoromaria*. *Lichenologist* 47:343–348
- Jorgensen PM, Ekman S, Wedin M (2013) Proposal to conserve the name *Fuscopannaria* against *Moelleropsis* (lichenized Ascomycota). *Taxon* 62:629–639
- Jorge-Villar SE, Miralles I, Ferrón CC, Hernandez V (2011) Raman spectroscopy study of lichens using three spectrometers under different experimental conditions: analyses of the results with relevance for extraplanetary exploration. *Anal Methods* 3:2783–2791
- Joseph S, Sinha GP (2015) Contributions to the genus *Synarthonia* (lichenized Ascomycota, Arthoniaceae). *Lichenologist* 47:123–130
- Joseph S, Sinha GP, Ramachandran VS (2013) A new record of *Syncesia* (Lichenized ascomycota) from India with additional taxonomic characters. *Taiwania* 58:300–304
- Joshi S, Hur JS (2012) A New Record of the genus *Mycobilimbia* (Ramalinaceae) from South Korea. *Mycobiology* 40:91–93
- Joshi Y, Koh YJ, Hur JS (2010a) Three new records of lichen genus *Rhizocarpon* from South Korea. *Mycobiology* 38:219–221
- Joshi Y, Wang XY, Nguyen TT, Koh YJ, Hur JS (2010b) Notes on the existence of *Leucodection desquamescens* (Thelotremonoid Graphidaceae) in South Korea. *Mycobiology* 38:149–152
- Joshi S, Upreti DK, Divakar PK (2011a) A new species of lichen genus *Syncesia* (Roccellaceae) from India. *Bryologist* 114:215–219
- Joshi Y, Nguyen TT, Lokos L, Koh YJ, Hur JS (2011b) Two new records of the lichen genus *Placynthiella elenkin* in South Korea. *Mycobiology* 39:54–56
- Joshi S, Upreti DK, Haridas Biju (2012) Nomenclatural notes on the lichen genera *Leucodection* and *Myriotrema* (Graphidaceae) in India. *Mycotaxon* 122:467–482
- Joshi S, Jayalal U, Lokos L, Park JS et al (2013a) *Leiorreuma exaltatum* and *Trapelia coarctata*, New to Korean Lichen Flora. *Mycobiology* 41:56–58
- Joshi S, Nguyen TT, Dzung NA, Jayalal U et al (2013b) New records of corticolous lichens from Vietnam. *Mycotaxon* 123:479–489
- Joshi Y, Upadhyay S, Shukla S, Nayaka S, Rawal RS (2015a) New records and an updated checklist of lichenicolous fungi from India. *Mycosphere* 6:195–200
- Joshi Y, Gagarina L, Halda JP, Oh SO, Hur JS (2015b) A new species and a new record of the lichen genus *Coenogonium* (Ostropales: Coenogoniaceae) from South Korea, with a world-wide key to crustose *Coenogonium* having prothalli. *Mycosphere* 6:667–672

- Josphine A, Senthilkumar G, Madhanraj P, Panneerselvam A (2011) Diversity of fungi from drift wood of muthupet mangroves. Asian J Pharm Ana 1:53–55
- Joubert A, Bataille-Simoneau N, Campion C, Guillemette T et al (2011a) Cell wall integrity and high osmolarity glycerol pathways are required for adaptation of *Alternaria brassicicola* to cell wall stress caused by brassicaceous indolic phytoalexins. Cell Microbiol 13:62–80
- Joubert A, Simoneau P, Campion C, Bataillé-Simoneau N et al (2011b) Impact of the unfolded protein response on the pathogenicity of the necrotrophic fungus *Alternaria brassicicola*. Mol Microbiol 79:1305–1324
- Judith C, Rossman AY, Kennedy AH, Cáceres O, Piepenbring M (2015) *Microchrysosphaera graminicola*, an enigmatic new genus and species in the Hypocreales from Panama. Mycol Prog 14:1–12
- Jungbluth P, Marcelli MP (2011) The *Pyxine pungens* complex in São Paulo State, Brazil. Bryologist 114:166–177
- Jungbluth P, Marcelli MP, Kalb K (2011) A new species and a new record of *Pyxine* (Physciaceae) with norstictic acid from São Paulo State, Brazil. Mycotaxon 115:435–442
- Jupa R, Hajek J, Hazdrova J, Bartak M (2012) Interspecific differences in photosynthetic efficiency and spectral reflectance in two *Umbilicaria* species from Svalbard during controlled desiccation. Czech Polar Rep 2:31–41
- Jurjevic Z, Kubatova A, Kolarik M, Hubka V (2015) Taxonomy of *Aspergillus* section *Petersonii* sect. nov. encompassing indoor and soil-borne species with predominant tropical distribution. Plant Syst Evol 301:2441–2462
- Jurjevic Z, Hubka V, Peterson S (2016) *Rasamsonia columbiensis*. Persoonia 36:404–405
- Jury SL (2013) Lichens collected during the fifth “Iter Mediterraneum” in Morocco, 8–27 June, 1992. Bocconeia 26:145–149
- Kaewwichian R, Limtong S (2014) *Nakazawaea siamensis* sp. nov., a yeast species isolated from phylloplane. Int J Syst Evol Microbiol 64:266–270
- Kaffer MI, Koch NM, Aptroot A, de M A, Suzana M (2015) New records of corticolous lichens for South America and Brazil. Plant Ecol Evol 148:111–118
- Kajale SC, Sonawane MS, Sharma R, Shouche YS (2015) *Leptoxiphyllum kurandae*-new record of insect gut associated sooty mould fungus from India. Mycosphere 6:133–138
- Kajitani Y, Masuya H (2011) *Ceratocystis ficicola* sp. nov., a causal fungus of fig canker in Japan. Mycoscience 52:349–353
- Kalb K (2009) New taxa and new records of thelotremonoid *Graphidaceae*. Herzogia 22:17–42
- Kalb K, Giralt M (2011) Orcularia, a segregate from the lichen genera *Buellia* and *Rinodina* (*Lecanoromycetes*, *Caliciaceae*). Phytotaxa 38:53–60
- Kalb K, Staiger B, Elix JA, Lange U, Lumbsch HT (2008) A new circumscription of the genus *Ramboldia* (*Lecanoraceae*, *Ascomycota*) based on morphological and molecular evidence. Nova Hedw 86:23–42
- Kalb K, Archer AW, Sutjaritturakan J, Boonpragob K (2009) New or otherwise interesting lichens V. Biblioth Lichenol 99:225–246
- Kalb K, Buaruang K, Mongkolsuk P, Boonpragob K (2012) New or otherwise interesting lichens. VI, including a lichenicolous fungus. Phytotaxa 42:35–47
- Kalb K, Polyiam W, Rivas PE, Bawingan PA et al (2016) ‘Missing links’ alive? Novel taxa represent morphological transitions between distinctive phenotypes among extant *Graphidaceae* (lichenized *Ascomycota*: *Ostropales*). Phytotaxa 268:110–122
- Kalko GV (2011) New for Russia and rare species of anamorphic fungi on conifers from saint petersburg suburbs. Mikol Fitopatol 45:356–361
- Kamal MM, Mia MAT (2009) Diversity and pathogenicity of the rice brown spot pathogen, *Bipolaris oryzae* (Breda de Haan) Shoem. in Bangladesh assessed by genetic fingerprint analysis. Bangladesh J Bot 38:119–125
- Kambar Y, Vivek MN, Manasa M, Vinayaka KS, Mallikarjun N, Kekuda PTR (2014) Antimicrobial activity of *Leptogium burnetiae*, *Ramalina hossei*, *Roccella montagnei* and *Heterodermia diademata*. Int J Pharm Phytoparm Res 4:164–168
- Kamgan NG, Jacobs K, Beer ZW, de Wingfield MJ, Roux J (2008) *Pesotum australi* [sic] sp. nov. and *Ophiostoma quercus* associated with *Acacia mearnsii* trees in Australia and Uganda, respectively. Australas Pl Path 37:406–416
- Kamgan NG, Solheim H, Beer ZW, de Grobbelaar JW et al (2010) *Ophiostoma* species, including *Ophiostoma borealis* sp. nov., infecting wounds of native broad-leaved trees in Norway. Cryptog Mycol 31:285–303
- Kamgan NG, Beer ZW, de Wingfield MJ, Roux J (2012) A diverse assemblage of *Ophiostoma* species, including two new taxa on eucalypt trees in South Africa. Mycol Prog 11:526
- Kang HJ, Sigler L, Lee J, Gibas CF et al (2010) *Xylogone ganoderophthora* sp. nov., an ascomycetous pathogen causing yellow rot on cultivated mushroom *Ganoderma lucidum* in Korea. Mycologia 102:1167–1184
- Kanigowski P, Flakus A, Oset M, Kowalewska A et al (2016) The lichen family *Parmeliaceae* in Poland. *Xanthoparmelia* species containing usnic acid. Herzogia 29:108–119
- Kantvilas G (2012) *Cameronia* (lichenized Ascomycetes), a remarkable new alpine genus from Tasmania. Lichenologist 44:91–100
- Kantvilas G (2014a) New species of the genus *Rimularia* Nyl. from Tasmania. Lichenologist 46:349–364
- Kantvilas G (2014b) The lichen family *Hymeneliaceae* in Tasmania, with the description of a new species. Kanunnah 7:127–140
- Kantvilas G (2015) Observations on the genus *Tephromela* (lichenised Ascomycetes) in Tasmania, with the description of a new species. Herzogia 28:430–444
- Kantvilas G (2016) Further additions to the lichen genus *Ramboldia* (*Lecanoraceae*) from Australia. Muelleria 34:103–109
- Kantvilas G, Divakar PK (2013a) New combinations in *Austroparmelina* (*Parmeliaceae*). - Australasian Lichenology 73:6–7
- Kantvilas G, Elix JA (2013) The lichen genus *Lecidella* (*Lecanoraceae*), with special reference to the Tasmanian species. Muelleria 31:31–47
- Kantvilas G, Lumbsch HT (2009) *Meridianelia*, a new genus in the *Elixiaceae* (*Ascomycota*) from Tasmania. Lichenologist 41:261–270
- Kantvilas G, Lumbsch HT (2012) Reappraisal of the genera of *Megalosporaceae* (*Teloschistales*, *Ascomycota*). Aust Syst Bot 25:210–216
- Kantvilas G, Leavitt SD, Elix JA, Lumbsch HT (2015) Additions to the genus *Trapelia* (*Trapeliaceae*: lichenised Ascomycetes). Aust Syst Bot 27:395–402
- Kaounas V, Agnello C, Alvarado P (2016) *Genea cephalonicae* sp. nov. (*Ascomycota*, *Pezizales*), a new hypogeous species from Greece. Ascomycete org 8:105–110
- Karagöz Y, Aslan A, Aptroot A (2011) *Diplotomma*, *Lecanora*, and *Xanthoria* lichen species new to Turkey. Mycotaxon 115:115–119
- Karakehian JM, LoBuglio KF, Pfister DH (2014) Placement of the genus *Angelina* within *Rhytismatales* and observations of *Angelina rufescens*. Mycologia 106:154–162
- Karakoti N, Bajpai R, Upreti DK, Mishra GK et al (2014) Effect of metal content on chlorophyll fluorescence and chlorophyll

- degradation in lichen *Pyxine cocoes* (Sw.) Nyl.: a case study from Uttar Pradesh, India. Environ Earth Sci 71:2177–2183
- Karandikar KG, Singh SK (2010) *Lylea indica*: a new hyphomycete species from India. Mycotaxon 112:257–260
- Karandikar KG, Singh PN, Singh PN (2015) *Mycoenterolobium flabelliforme*: a new anamorphic fungus from India. Plant Pathol Quar 5:49–51
- Karaoglanidis GS, Luo Y, Michailides TJ (2011) Competitive ability and fitness of *Alternaria alternata* isolates resistant to QoI fungicides. Plant Dis 95:178–182
- Kärnefelt I, Seaward MR, Thell A (2012) Systematics, biodiversity and ecology of lichens. Acta Bot Hung 54:3–4
- Kasahara K, Miyamoto T, Fujimoto T, Oguri H et al (2010) *Solanapyrone synthase*, a possible Diels-Alderase and iterative type I polyketide synthase encoded in a biosynthetic gene cluster from *Alternaria solani*. Chem Bio Chem 11:1245–1252
- Karunaratna A, Phookamsak R, Wanasinghe DN, Wijayawardene NN et al (2017) Taxonomy and phylogeny of *Sparticola muriformis* sp. nov. on decaying grass. Mycosphere 8:603–614
- Kashiwadani H, Aptroot A, Moon KH (2009) Pyrenocarpous lichens of Japan, with the resurrection of the genus *Trypetheliopsis*. Biblioth Lichenol 99:247–258
- Kaur S, Gehlot P (2011) New pyrenomycetous genus from India. J Mycol Plant Pathol 41:487–488
- Kawada M, Someno T, Inoue H, Ohba SI et al (2010) a new antitumor compound against human prostate cancer cells, produced by *Perisporiopsis melioloides* Mer-f16716. J Antibiot 63:319–323
- Kawahigashi H, Kasuga S, Ando T, Kanamori H et al (2011) Positional cloning of ds1, the target leaf spot resistance gene against *Bipolaris sorghicola* in sorghum. Theor Appl Genet 123:131–142
- Kaya A (2009) Macrofungi of Huzurlu high plateau (Gaziantep-Turkey). Turkish J Bot 33:429–437
- Kaya M, Halıcı MG, Duman F, Erdogan S et al (2015) Characterisation of α -chitin extracted from a lichenised fungus species *Xanthoria parietina*. Nat Prod Res 29:1280–1284
- Każmierska E, Kossowska M (2016) Contribution to the lichen biota of the Stawy Milickie nature reserve and its adjacent area (Lower Silesia, southwestern Poland). Histry 51:1075
- Keča N, Milenovic I, Keca L (2015) Mycological complex of poplars in Serbia. J For Sci 61:169–174
- Keilor Rojas-Jimenez K, Hernandez M, Blanco J, Vargas LD et al (2016) Richness of cultivable endophytic fungi along an altitudinal gradient in wet forests of Costa Rica. Fungal Ecol 20:124–131
- Kekuda PTR, Akarsh S, Nawaz AN, Ranjitha MC et al (2016) In vitro antifungal activity of some plants against *Bipolaris sorokiniana* (Sacc.) Shoem. Int J Curr Microbiol Applied Sci 5:331–337
- Kelly LJ, Hollingsworth PM, Coppins BJ, Ellis CJ et al (2011) DNA barcoding of lichenized fungi demonstrates high identification success in a floristic context. New Phytol 191:288–300
- Kepler RM, Sung GH, Ban S, Nakagiri A et al (2012a) New teleomorph combinations in the entomopathogenic genus *Metacordyceps*. Mycologia 104:182–197
- Kepler RM, Sung GH, Harada Y, Tanaka K et al (2012b) Host jumping onto close relatives and across kingdoms by *Tyrannicordyceps* (Clavicipitaceae) gen. nov. and *Ustilaginoides* (Clavicipitaceae). Am J Bot 99:552–561
- Kepler R, Ban S, Nakagiri A, Bischoff J et al (2013) The phylogenetic placement of hypocrealean insect pathogens in the genus *Polycephalomyces*: an application of One Fungus One Name. Fungal Biol 117:611–622
- Kepler RM, Humber RA, Bischoff JF, Rehner SA (2014) Clarification of generic and species boundaries for Metarhizium and related fungi through multigene phylogenetics. Mycologia 106:811–829
- Kerr AM (2014) Catalogue of the Lichens of the Mariana Islands ex Elix & McCarthy's (2008) Checklist of Pacific Island Lichens.
- Ketner-Oostra R, Sparrius LB, Sýkora KV (2010) Development of lichen-rich communities. Inland drift sand landscapes, Paleoecol Landsc Ecol, pp 235–254
- Khare R, Upreti DK, Nayaka S, Gupta RK (2009) Diversity of Soil lichens in India. Soil Microflora. Daya Publishing House, New Delhi, pp 64–75
- Khare R, Rai H, Upreti DK, Dubey U et al (2010) Occurrence of rare, crustose lichen, *Ophioparma ventosa* (*Ophioparmaceae*, lichenized Ascomycota), from alpine regions of Arunachal Pradesh, India. Feddes Repertorium 121:184–188
- Kharpukhaeva TM, Mukhortova LV (2016) Dynamics of interaction between lichens and fallen deadwood in forest ecosystems of the eastern Baikal region. Contemp Probl Ecol 9:125–139
- Khemmuk W, Geering ADW, Shivas RG (2016) *Wongia* gen. nov. (*Papulosaceae*, *Sordariomycetes*), a new generic name for two root-infecting fungi from Australia. IMA Fungus 7:247–252
- Khodaparast SA, Takamatsu S, Harada M, Abbasi M, Samadi S (2012) Additional rDNA ITS sequences and its phylogenetic consequences for the genus *Leveillula* with emphasis on conidium morphology. Mycol Prog 11:741–752
- Khodaparast SA, Takamatsu S, Shadloo A, Damadi M et al (2016) Notes on the genus *Leveillula* (*Erysiphaceae*): a new unrecorded species and notes on *Leveillula* infecting *Ficus*, *Cucurbita* and *Tropaeolum* in Iran. Phytotaxa 260:267–275
- Khunnamwong P, Lertwattanasakul N, Jindamorakot S, Limtong S et al (2015) Description of *Diutina* gen. nov., *Diutina siamensis* sp. nov., and reassignment of *Candida catenulata*, *Candida mesorugosa*, *Candida neorugosa*, *Candida pseudorugosa*, *Candida ranongensis*, *Candida rugosa* and *Candida scorzettiae* to the genus *Diutina*. Int J Syst Evol Microbiol 65:4701–4709
- Kihara J, Moriwaki A, Tanaka N, Tanaka C et al (2008) Characterization of the BMR1 gene encoding a transcription factor for melanin biosynthesis genes in the phytopathogenic fungus *Bipolaris oryzae*. FEMS Microbiol Lett 281:221–227
- Kihara J, Tanaka N, Ueno M, Arase S (2009) Cloning and expression analysis of two opsins-like genes in the phytopathogenic fungus *Bipolaris oryzae*. FEMS Microbiol Lett 295:289–294
- Kilic M, Altintas DU, Yilmaz M, Kendirli SG et al (2010) The effects of meteorological factors and *Alternaria* spore concentrations on children sensitised to *Alternaria*. Allergologia et Immunopathol 38:122–128
- Kim KH, Choi YJ, Seo ST, Shin HD (2009a) *Raffaelea quercus-mongolicae* sp. nov. associated with *Platypus koryoensis* on oak in Korea. Mycotaxon 110:189–197
- Kim SW, Kim KS, Lamsal K, Kim YJ et al (2009b) An in vitro study of the antifungal effect of silver nanoparticles on oak wilt pathogen *Raffaelea* sp. J Microbiol Biotechnol 19:760–764
- Kim MY, Sohn JH, Ahn JS, Oh H (2009c) Alternaramide, a cyclic depsipeptide from the marine-derived fungus *Alternaria* sp. SF-5016. J Nat Prod 72:2065–2068
- Kim HS, Kim JY, Lee HK, Kim MS et al (2010) Dendritic cell activation by glucan isolated from *Umbilicaria esculenta*. Immune Network 10:188–197
- Kim MK, Park H, Oh TJ (2013) Antimicrobial properties of the bacterial associates of the Arctic lichen *Stereocaulon* sp. Afr J Microbiol Res 7:3651–3657
- Kinalioglu K, Aptroot A (2011) *Carbonea*, *Gregorella*, *Porpidia*, *Protomicarea*, *Rinodina*, *Solenopsora*, and *Thelenella* lichen species new to Turkey. Mycotaxon 115:125–129
- King KM, West JS, Brunner PC, Dyer PS, Fitt BDL (2013) Evolutionary relationships between *Rhynchosporium lolii* sp. nov. and other *Rhynchosporium* species on grasses. PLoS ONE 8:e72536

- King KM, West JS, Fitt BDL, Dyer PS (2014) Differences in MAT gene distribution and expression between *Rhynchosporium* species on grasses. *Plant Pathol* 64:344–354
- Kirby BM, Barnard D, Tuffin IM, Cowan DA (2011) Ecological distribution of microorganisms in terrestrial, psychrophilic habitats. *Extremophiles Handbook*. Springer, Tokyo, pp 839–863
- Kirchmair M, Neuhauser S, Buzina W, Huber L (2008) The taxonomic position of *Roesleria subterranea*. *Mycol Res* 112:1210–1219
- Kirika P, Mugambia G, Lücking R, Lumbsch HT (2012) New records of lichen-forming fungi from Kenya. *J East Afr Nat Hist* 101:73–98
- Kirika PM, Leavitt SD, Divakar PK, Crespo A et al (2015) The monotypic genus *Bulborrhizina* belongs to *Bulbothrix* sensu lato (Parmeliaceae, Ascomycota). *Bryologist* 118:164–169
- Kirika PM, Divakar PK, Crespo A, Leavitt SD et al (2016) Polyphyly of the genus *Canoparmelia*—uncovering incongruences between phenotype-based classification and molecular phylogeny within lichenized Ascomycota (Parmeliaceae). *Phytotaxa* 289:36–48
- Kirk P, Cannon PF, Minter DW, Stalpers JA (2008) Ainsworth & Bisby's Dictionary of the Fungi, 10th edn. CAB International, Wallingford
- Kirk PM, Stalpers JA, Braun U, Crous PW et al (2013) A without-prejudice list of generic names of fungi for protection under the International Code of Nomenclature for algae, fungi, and plants. *IMA Fungus* 4:381–443
- Kirschner R (2009) *Cercosporaella* and *Ramularia*. *Mycologia* 101:110–119
- Kirschner R (2015) Phylogenetic placement of a new species of *Corynesporopsis* from dead acacia wood indicates occurrence of tretic conidiogenesis within Xylariales. *Phytotaxa* 192:24–34
- Kirschner R (2016) Revision of the morphology and biogeography of *Thysanorea papuana*. *Mycosphere* 7:820–827
- Kirschner R, Chen CJ (2010) Two new species of Ramichloridium-like hyphomycetes from senescent leaves of night-scented lily (*Alocasia odora*) in Taiwan. *Fungal Divers* 40:41–50
- Kirschner R, Liu LC (2014) Mycosphaerellaceous fungi and new species of *Venustosynnema* and *Zasmidium* on ferns and fern allies in Taiwan. *Phytotaxa* 176:309–323
- Kirschner R, Pang KL, Jones EBG (2013) Two cheiroporous hyphomycetes reassessed based on morphological and molecular examination. *Mycol Prog* 12:29–36
- Kiyuna T, An KD, Kigawa R, Sano C et al (2011) Molecular assessment of fungi in “black spots” that deface murals in the Takamatsuzuka and Kitara Tumuli in Japan: *Acremonium* sect. *Gliomastix* including *Acremonium tumulicola* sp. nov. and *Acremonium felinum* comb. nov. *Mycoscience* 52:1–7
- Kiyuna T, An KD, Kigawa R, Sano C et al (2012) Bristle-like fungal colonizers on the stone walls of the Kitara and Takamatsuzuka Tumuli are identified as *Kendrickiella phycomyces*. *Mycoscience* 53:446–459
- Kjer J, Wray V, Edrada-Ebel R, Ebel R et al (2009) Xanalteric acids I and II and related phenolic compounds from an endophytic *Alternaria* sp. isolated from the mangrove plant Sonneratia alba. *J Nat Prod* 72:2053–2057
- Klaubauf S, Tharreau D, Fournier E, Groenewald JZ et al (2014) Resolving the polyphyletic nature of *Pyricularia* (Pyriculariaceae). *Stud Mycol* 79:85–120
- Klavina D, Muiznieks I, Gaitnieks T, Lazzins A et al (2016) Fine root development and mycorrhization in Norway spruce stands a year after fertilization with potassium sulphate and wood ash. *J For Sci* 62:17–23
- Klaysuban A, Sakayaroj J, Jones EBG (2014) An additional marine fungal lineage in the Diatrypaceae, Xylariales: *Pedumispora rhizophorae*. *Bot Mar* 57:413–420
- Kleczewski NM, Flory SL (2010) Leaf blight disease on the invasive grass *Microstegium vimineum* caused by a *Bipolaris* sp. *Plant Dis* 94:807–811
- Knapp DG, Kovacs GM, Zajta E, Groenewald JZ, Crous PW (2015) Dark septate endophytic pleosporalean genera from semiarid areas. *Persoonia* 35:87–100
- Knight NL, Platz GJ, Lehmensiek A, Sutherland MW (2010) An investigation of genetic variation among Australian isolates of *Bipolaris sorokiniana* from different cereal tissues and comparison of their abilities to cause spot blotch on barley. *Aust Plant Pathol* 39:207–216
- Knight A, Bannister JM, Ludwig LR, Malcolm WM, Orlovich DA (2013) Rediscovery of pycnidia in *Thamnolia vermicularis*: implications for chemotype occurrence and distribution. *Lichenologist* 45:397–411
- Knudsen K (2011) A preliminary study of *Pleopsidium stenosporum* (Stizenb. ex Hasse) K. Knudsen. *Opusc Phil* 9:77–83
- Knudsen K, Kocourkova J (2009) Lichens, lichenicolous and allied fungi of the Santa Monica mountains, Part 4: additions and corrections to the annotated checklist. *Opusc Phil* 7:29–48
- Knudsen K, Kocourkova J (2010a) A new *Lichenostigma* species (genus *incertae sedis*) from southern California. *Bryologist* 113:229–234
- Knudsen K, Kocourkova J (2010b) Lichens, lichenicolous and allied fungi of the Santa Monica Mountains, Part 5: additions and corrections to the Annotated Checklist. *Opusc Phil* 8:83–100
- Knudsen K, Kocourkova J (2012) Lichenological notes 5: Neotypification of *Sarcogyne magnussonii* (Acarosporaceae). *Mycotaxon* 121:139–145
- Knudsen K, Lendemer JC (2009) *Naetrocymbe herrei* (Pleosporales: Ascomycetes), a new lichenized saxicolous species from the coast of central California, USA. *Opusc Phil* 6:59–64
- Knudsen K, Elix JA, Reeb V (2008) A preliminary study of the genera *Acarospora* and *Pleopsidium* in South America. *Opusc Phil* 5:1–22
- Knudsen K, Breuss O, Kocourkova J (2009a) A new species of *Placocarpus* (Verrucariaceae) from southern California. *Lichenologist* 41:627–630
- Knudsen K, Gokhan Halici M, Kocakaya M (2009b) *Sarcogyne magnispora* (Acarosporaceae), a new species in the nivea group from Turkey. *Mycotaxon* 107:413–417
- Knudsen K, Lendemer JC, Harris RC (2011) Lichens and lichenicolous fungi—no. 15: miscellaneous notes on species from eastern North America. *Opusc Phil* 9:45–75
- Knudsen K, Flakus A, Kukwa M (2012) A contribution to the study of Acarosporaceae in South America. *Lichenologist* 44:253–262
- Knudsen K, Kocourkova J, McCune B (2013a) *Sarcogyne mitziae* (Acarosporaceae), a new species from biotic soil crusts in western North America. *Bryologist* 116:122–126
- Knudsen K, Sheard JW, Kocourkova J, Mayrhofer H (2013b) A new lichenicolous lichen from Europe and western North America in the genus *Dimelaena* (Physciaceae). *Bryologist* 116:257–262
- Kobayashi H, Sano A, Aragane N, Fukuoka M et al (2008) Disseminated infection by *Bipolaris spicifera* in an immunocompetent subject. *Med Mycol* 46:361–365
- Kobayashi T, Iijima K, Radhakrishnan S, Mehta V et al (2009) Asthma-related environmental fungus, *Alternaria*, activates dendritic cells and produces potent Th2 adjuvant activity. *J Immunol* 182:2502–2510
- Kobmoo N, Mongkolsamrit S, Wutikhun T, Tasanathai K et al (2015) New species of *Ophiocordyceps unilateralis*, an ubiquitous pathogen of ants from Thailand. *Fung Biol* 119:44–52
- Kocakaya M, Halici MG, Aksoy A (2011) *Zwackhiomyces turcicus* sp. nov. (Ascomycota, Xanthopyreniaceae) from Turkey. *Mycotaxon* 116:329–333

- Kocharin K, Supothina S, Prathumpai W (2013) Hypothemycin production and its derivatives diversifying of *Aigialus parvus* BCC 5311 influenced by cultural condition. *Adv Biosci Biotechnol* 4:1049–1056
- Kocourková J (2009) Observations on the genus *Neolamya*, with the description of the new species *N. xanthoparmeliae* (*Ascomycota*, genera *incertae sedis*). *Opusc Phil* 6:137–148
- Kocourková J, Hawksworth DL (2008) *Acaroconium punctiforme* gen. sp. nov., a new lichenicolous coelomycete on *Acarospora* species and *Sarcogyne regularis*. *Lichenologist* 40:105–109
- Kocourková J, Knudsen K (2009) *Gelatinopsis acarosporicola* (Helotiaceae), a new lichenicolous fungus on *Acarospora socialis* from western North America. *Bryologist* 112:363–367
- Kocourková J, Knudsen K (2011) *Endococcus thelomatis*, a new lichenicolous fungus from southern California. *Bibliotheca Lichenol* 106:173–178
- Kocourková J, Fryday AM, Knudsen K, Lendemer JC (2010) Studies in lichens and lichenicolous fungi: more notes on taxa from North America 6. *Mycotaxon* 111:423–429
- Kohli MM, Mehta YR, Guzman E, Viedma LD, Cubilla LE (2011) *Pyricularia* blast—a threat to wheat cultivation. *Czech J Genet Plant Breed* 47:S130–S134
- Kolarík M, Hulcr J (2009) Mycobiota associated with the ambrosia beetle *Scolytodes unipunctatus* Blandford (Coleoptera: Curculionidae, Scolytinae). *Mycol Res* 113:44–60
- Kolarík M, Freeland E, Utley C, Tisserat N (2011) *Geosmithia morbida* sp. nov., a new phytopathogenic species living in symbiosis with the walnut twig beetle (*Pityophthorus juglandis*) on *Juglans* in USA. *Mycologia* 103:325–332
- Kolarík M, Hujslová M, Vazquéz-Campos X (2015a) Acidotolerant genus *Fodinomyces* (*Ascomycota*: *Capnodiales*) is a synonym of *Acidiella*. *Czech Mycol* 67:37–38
- Kolarík M, Hulcr J, Kirkendall LR (2015b) New species of *Geosmithia* and *Graphium* associated with ambrosia beetles in Costa Rica. *Czech Mycol* 67:29–35
- Kondratyuk SY (2015) *Agonimia blumii* sp. n. (*Verrucariales*, lichen-forming *Ascomycota*), a new taxon from Eastern Asia. *Ukr Bot J* 72:246–251
- Kondratyuk S, Jeong MH, Yu NH et al (2013a) Four new genera of teloschistoid lichens (*Teloschistaceae*, *Ascomycota*) based on molecular phylogeny. *Acta Bot Hung* 55:251–274
- Kondratyuk S, Lókös L, Tschabanenko S, Haji Moniri M et al (2013b) New and noteworthy lichen-forming and lichenicolous fungi. *Acta Bot Hung* 55:275–349
- Kondratyuk S, Yatsyna A, Lókös L, Galanina I et al (2013c) Three new *Xanthoria* and *Rusavskia* species (*Teloschistaceae*, *Ascomycota*) from Europe. *Acta Bot Hung* 55:351–365
- Kondratyuk SY, Tschabanenko SI, Elix JA, Oh S et al (2013d) *Nipponoparmelia perplicata* sp. nov. (*Parmeliaceae*, *Ascomycota*) from eastern Asia. *Mycotaxon* 126:37–44
- Kondratyuk S, Kärnefelt I, Thell A, Elix J et al (2014) A revised taxonomy for the subfamily *Caloplacoideae* (*Teloschistaceae*, *Ascomycota*) based on molecular phylogeny. *Acta Bot Hung* 56:141–178
- Kondratyuk S, Lókös L, Kim JA, Kondratyuk AS et al (2015a) New members of the *Pertusariales* (*Ascomycota*) proved by combined phylogenetic analysis. *Stud Bot Hung* 46:95–110
- Kondratyuk SY, Jeong MH, Galanina IA, Yakovchenko LS et al (2015b) Molecular phylogeny of placiodioid lichen-forming fungi reveal a new genus, *Sedelnikovaea*. *Mycotaxon* 129:269–282
- Kondratyuk SY, Kärnefelt I, Thell A, Elix JA et al (2015c) *Brownlielloideae*, a new subfamily in the *Teloschistaceae* (*Lecanoromycetes*, *Ascomycota*). *Acta Bot Hung* 57:321–343
- Kondratyuk SY, Kim JA, Yu NH, Jeong MH et al (2015d) *Zeroviella*, a new genus of xanthorioid lichens (*Teloschistaceae*, *Ascomycota*) proved by three gene phylogeny. *Ukr Bot J* 72:574–584
- Kondratyuk SY, Lókös L, Farkas E, Oh SO, Hur JS (2015e) New and noteworthy lichen-forming and lichenicolous fungi 2. *Acta Bot Hung* 57:77–141
- Kondratyuk SY, Lókös L, Farkas E, Oh SO, Hur JS (2015f) New and noteworthy lichen-forming and lichenicolous fungi 3. *Acta Bot Hung* 57:345–382
- Kondratyuk SY, Lókös L, Kapetz NV, Pleskach LY et al (2015g) *Physcia ucrainica* sp. nova (*Physciaceae*, *Ascomycota*) from the Crimean Peninsula, proved by molecular phylogeny. *Acta Bot Hung* 57:143–163
- Kondratyuk SY, Lókös L, Kim JA, Kondratyuk AS et al (2015h) Three new monotypic genera of the caloplacoid lichens (*Teloschistaceae*, Lichen-Forming Ascomycetes). *Mycobiology* 43:195–202
- Kondratyuk SY, Lókös L, Halda JP, Haji Moniri M et al (2016a) New and noteworthy lichen-forming and lichenicolous fungi 4. *Acta Bot Hung* 58:75–136
- Kondratyuk SY, Lókös L, Kim JA, Kondratyuk AS et al (2016b) A new genus of the lecanoroid caloplacoid lichens (*Teloschistaceae*, lichen-forming ascomycetes). *Acta Bot Hung* 58:303–318
- Konoreva LA, Chesnokov SV, Davydov EA (2016) *Stictis* and *Schizoxylon* (*Stictidaceae*, *Ostropales*) in Russia. *Herzogia* 29:706–711
- Kosanić M, Ranković B, Stanojković T (2012) Antioxidant, antimicrobial, and anticancer activity of 3 *Umbilicaria* species. *J Food Sci* 77:T20–T25
- Kossowska M, Szczepańska K (2016) Current distribution and ecology of lichens in a biodiversity hotspot in the Mały Śnieżny Kocioł glacial cirque, Sudetes, Poland. *Herzogia* 29:120–136
- Konta S, Hongsan S, Phillips AJL, Jones EBG et al (2016a) *Botryosphaeriaceae* from palms in Thailand II - two new species of *Neodeightonia*, *N. rattanica* and *N. rattanicola* from *Calamus* (rattan palm). *Mycosphere* 7:950–961
- Konta S, Hongsan S, Tibpromma S, Thongbai B et al (2016b) An advance in the endophyte story: *Oxydorhidaceae* fam. nov. with six new species of *Oxydorthis*. *Mycosphere* 7:1425–1446
- Konta S, Phillips AJ, Bahkali AH, Jones EB et al (2016c) *Botryosphaeriaceae* from palms in Thailand-*Barriopsis archontophoenicis* sp. nov. from *Archontophoenix alexandrae*. *Mycosphere* 7:921–932
- Kossowska M, Szczepańska K (2013) New records of lichenicolous fungi from SW Poland. *Pol Bot J* 58(2):735–739
- Koukol O (2010) Revision of “*Septonema ochraceum*” revealed three new species of *Venturiaceae* and *Herpotrichiellaceae*. *Micol Prog* 9:369–378
- Koukol O (2011) New species of *Chalara* occupying coniferous needles. *Fungal Divers* 49:75–91
- Koukol O (2012) A new species of *Infundichalara* from pine litter. *Mycotaxon* 120:343–352
- Koukol O, Kolárová Z (2010) *Bactrodesmium gabretiae* (anamorphic *Helotiaceae*), a new sporodochial species described from spruce needles. *Nova Hedw* 91:243–248
- Koukol O, Kelnarova I, Černý K (2014) Recent observations of sooty bark disease of sycamore maple in Prague (Czech Republic) and the phylogenetic placement of *Cryptostroma corticale*. *For Pathol* 45:21–27
- Kovács GM, Trappe Amalsheikh JM, Boka K, Elliott TF (2008) *Imaia*, a new truffle genus to accommodate *Terfezia gigantea*. *Mycologia* 100:930–939
- Kovács GM, Balázsz TK, Calonge FD, Martín MP (2011a) The diversity of *Terfezia* desert truffles: new species and a highly

- variable species complex with intrasporocarpic nrDNA ITS heterogeneity. *Mycologia* 103:841–853
- Kovács GM, Trappe JM, Alsheikh AM, Hansen K et al (2011b) *Terfezia* disappears from the American truffle mycota as two new genera and *Mattirolomyces* species emerge. *Mycologia* 103:831–840
- Kowalski T, Holdenrieder O (2009) The teleomorph of *Chalara fraxinea*, the causal agent of ash dieback. *For Pathol* 39:304–308
- Kowalski M, Hausner G, Piercey-Normore MD (2011) Bioactivity of secondary metabolites and thallus extracts from lichen fungi. *Mycoscience* 52:413–418
- Kowalsky T, Krai W, Bednarz B (2016) Fungi on stems and twigs in initial and advanced stages of dieback of European ash (*Fraxinus excelsior*) in Poland. *Eur J For Res* 135:565–579
- Kraichak E, Parnmen S, Lücking R, Lumbsch HT (2013) *Gintarasia* and *Xalocoa*, two new genera to accommodate temperate to subtropical species in the predominantly tropical *Graphidaceae* (*Ostropales, Ascomycota*). *Aust Syst Bot* 26:466–474
- Kraichak E, Parnmen S, Lücking R, Plata ER et al (2014) Revisiting the phylogeny of Ocellulariae, the second largest tribe within *Graphidaceae* (lichenized *Ascomycota: Ostropales*). *Phytotaxa* 189:52–81
- Kraichak E, Lücking R, Aptroot A, Beck A et al (2015) Hidden diversity in the morphologically variable script lichen (*Graphis scripta*) complex (*Ascomycota, Ostropales, Graphidaceae*). *Org Divers Evol* 15:447–458
- Kristiina M, Lauri S, Andres S, Thell A, Randlane T (2012) Testing morphology-based delimitation of *Vulpicida juniperinus* and *V. tubulosus* (*Parmeliaceae*) using three molecular markers. *Lichenologist* 44:757–772
- Kristinsson H, Heiðmarsson S, Hansen ES (2014) Lichens from Iceland in the collection of Svanhildur Svane. *Bot Lith* 20:14–18
- Kruys Å (2015) New species of Preussia with 8-celled ascospores (*Sporormiaceae, Pleosporales, Ascomycota*). *Phytotaxa* 234:143–150
- Kruys Å, Wedin M (2009) Phylogenetic relationships and an assessment of traditionally used taxonomic characters in the *Sporormiaceae* (*Pleosporales, Dothideomycetes, Ascomycota*), utilising multi-gene phylogenies. *Syst Biodivers* 7:465–478
- Kruys Å, Castlebury LA (2012) Molecular phylogeny of *Sydiowiellaceae*—resolving the position of *Cainiella*. *Mycologia* 104:419–426
- Kruys A, Huhndorf S, Miller A (2015) Coprophilous contributions to the phylogeny of *Lasiosphaeriaceae* and allied taxa within *Sordariales* (*Ascomycota, Fungi*). *Fungal Divers* 70:101–113
- Krzewicka B (2010) *Umbilicaria rhizinata* comb. nov. (lichenized *Ascomycota*). *Lichenologist* 42:491–493
- Krzewicka B (2012) A revision of *Verrucaria* sl (*Verrucariaceae*) in Poland. *Pol Bot Stud* 27:143
- Krzewicka B, Wilk K (2009) *Umbilicaria isidiosa* (lichenized *Ascomycota*), a new species from Bolivia. *Mycotaxon* 109:165–170
- Krzewicka B, García MA, Johansen SD, Sancho LG et al (2009) Morphological and nuclear ribosomal DNA data support distinguishing two new species of *Umbilicaria* (*Umbilicariaceae, Ascomycota*) from Europe. *Lichenologist* 41:631–648
- Kuan CS, Yew SM, Toh YF, Chan CL, et al. (2015) Dissecting the fungal biology of *Bipolaris papendorffii*: from phylogenetic to comparative genomic analysis. *DNA Research*, p dsv007
- Kubiak D, Guzow-Krzemińska B (2014) Herbarium record of *Vulpicida tubulosus* (Lichenized *Ascomycota*) from locality beyond its known distribution range. *Pol J Environ Stud* 23:577–583
- Kubiak D, Osyczka P, Rola K (2016) Spontaneous restoration of epiphytic lichen biota in managed forests planted on habitats typical for temperate deciduous forest. *Biodivers Conserv* 25:1937–1954
- Kučera V, Lizoň P, Kautmanová I (2010) Geoglossoid fungi in Slovakia II. *Trichoglossum octopartitum*, a new species for the country. *Czech Mycol* 62:13–18
- Kučera V, Lizoň P, Tomšovský M, Kučera J et al (2014) Re-evaluation of the morphological variability of *Microglossum viride* and *M. griseoviride* sp. nov. *Mycologia* 106:282–290
- Kuhnert E, Surup F, Sir EB, Lambert C et al (2015) Lenormandins A–G, new azaphilones from *Hypoxyton lenormandii* and *Hypoxyton jaklitschii* sp. nov., recognised by chemotaxonomic data. *Fungal Divers* 71:165–184
- Kuhnert E, Sir EB, Lambert C, Hyde KD et al (2017) Phylogenetic and chemotaxonomic resolution of the genus *Annulohypoxylon* (*Xylariaceae*) including four new species. *Fungal Divers*. doi:10.1007/s13225-016-0377-6
- Kukwa M (2010) Additions to the list of lichens and lichenicolous fungi of Gdanskie Pomerania. *Fragm Florist Geobot Pol* 17:196–199
- Kukwa M, Flakus A (2009) New or interesting records of lichenicolous fungi from Poland VII. Species mainly from Tatra Mountains. *Herzogia* 22:191–211
- Kukwa M, Pérez-Ortega S (2010) A second species of *Botryolepraria* from the Neotropics and the phylogenetic placement of the genus within *Ascomycota*. *Mycol Prog* 9:345–351
- Kukwa M, Łubek A, Szymczyk R, Zalewska A (2012) Seven lichen species new to Poland. *Mycotaxon* 120:105–118
- Kukwa M, Schiefelbein U, Flakus A (2013) A contribution to the lichen family *Graphidaceae* (*Ostropales, Ascomycota*) of Bolivia. *Herzogia* 26:231–252
- Kumar JV, Kumari BR, Sujatha G, Castano E (2008a) Production of plants resistant to *Alternaria carthami* via organogenesis and somatic embryogenesis of safflower cv NARI-6 treated with fungal culture filtrates. *Plant Cell, Tissue Organ Cult* 93:85–96
- Kumar V, Haldar S, Pandey KK, Singh RP et al (2008b) Cultural, morphological, pathogenic and molecular variability amongst tomato isolates of *Alternaria solani* in India. *World J Microbiol Biotechnol* 24:1003–1009
- Kumar U, Joshi AK, Kumar S, Chand R, Röder MS (2009) Mapping of resistance to spot blotch disease caused by *Bipolaris sorokiniana* in spring wheat. *Theor Appl Genet* 118:783–792
- Kumar U, Joshi AK, Kumar S, Chand R, Röder MS (2010) Quantitative trait loci for resistance to spot blotch caused by *Bipolaris sorokiniana* in wheat (*T. aestivum* L.) lines ‘Ning 8201’ and ‘Chiryu 3’. *Mol Breed* 26:477–491
- Kumar P, Anshu V, Kumar S (2011) Morpho-pathological and molecular characterization of *Bipolaris oryzae* in rice (*Oryza sativa*). *J Phytopathol* 159:51–56
- Kuna P, Kaczmarek J, Kupeczyk M (2011) Efficacy and safety of immunotherapy for allergies to *Alternaria alternata* in children. *J Allergy Clin Immunol* 127:502–508
- Kuo CC, Chen P, Chen CY (2008) Harposporium from Taiwan. *Fung Sci* 23:35–41
- Kurniawati E, Zhang H, Chukeatirote E, Sulistyowati L et al (2010) Diversity of freshwater ascomycetes in freshwater bodies at Amphoe Mae Chan, Chiang Rai. *Cryptog Mycol* 31:323–331
- Kurtzman CP (2009) Biotechnological strains of *Komagataella (Pichia) pastoris* are *Komagataellaphaffii* as determined from multigene sequence analysis. *J Ind Microbiol Biotechnol* 36:1435–1438
- Kurtzman CP (2012) *Citeromyces hawaiiensis* sp. nov., an ascosporic yeast associated with *Myoporum sandwicense*. *Int J Syst Evol Microbiol* 62:1215–1219
- Kurtzman CP (2015) Description of *Martiniozyma* gen. nov. and transfer of seven *Candida* species to *Saturnispora* as new combinations. *Antonie van Leeuwenhoek* 108:803–809

- Kurtzman CP (2016) Description of *Groenewaldozyma* gen. nov. for placement of *Candida auringiensis*, *Candida salmanticensis* and *Candida tartarivorans*. Antonie van Leeuwenhoek 109:1041–1045
- Kurtzman CP, Robnett CJ (2013a) *Alloascoidea hylecoeti* gen. nov., comb. nov., *Alloascoidea africana* comb. nov., *Ascoidea tarda* sp. nov., and *Nadsonia starkeyi-henicci* comb. nov., new members of the *Saccharomycotina* (Ascomycota). FEMS Yeast Res 13:423–432
- Kurtzman CP, Robnett CJ (2013b) Relationships among genera of the *Saccharomycotina* (Ascomycota) from multigene phylogenetic analysis of type species. FEMS Yeast Res 13:23–33
- Kurtzman CP, Robnett CJ (2014) Three new anascoporic genera of the Saccharomycotina: *Danielozyma* gen. nov., *Deakozyma* gen. nov. and *Middlehovenomyces* gen. nov. Antonie van Leeuwenhoek 105:933–942
- Kurtzman CP, Suzuki M (2010) Phylogenetic analysis of ascomycete yeasts that form coenzyme Q-9 and the proposal of the new genera *Babjeviella*, *Meyeromyza*, *Milleromyza*, *Priceomyces*, and *Scheffersomyces*. Mycoscience 51:2–14
- Kurtzman CP, Robnett CJ, Basehoar-Powers E (2008) Phylogenetic relationships among species of *Pichia*, *Issatchenkia* and *Williopsis* determined from multigene sequence analysis, and the proposal of *Barnetozyma* gen. nov., *Lindnera* gen. nov. and *Wickerhamomyces* gen. nov. FEMS Yeast Res 8:939–954
- Kurtzman CP, Price NP, Ray KJ, Kuo TM (2010) Production of sophorolipid biosurfactants by multiple species of the *Starmerella* (*Candida*) *bombicola* yeast clade. FEMS Microbiol Lett 311:140–146
- Kurtzman CP, Robnett CJ, Blackwell M (2016) Description of *Teunomyces* gen. nov. for the *Candida kruisii* clade, *Suhomyces* gen. nov. for the *Candida tanzawaensis* clade and *Suhomyces kilbournensis* sp. nov. FEMS Yeast Res 16:fow041
- Kušan I, Matičec N, Mešić A, Tkalcčec Z (2015a) A new species of *Thecotheus* from Croatia with a key to the known species with apiculate spores. Sydowia 67:51–63
- Kušan I, Matičec N, Mešić A, Tkalcčec Z (2015b) *Tricharina topfisia*—a new species from Croatia, with a revision of *T. japonica* (Pyronemataceae, Pezizales). Phytotaxa 221:35–47
- Kusari S, Zühlke S, Košuth J, Čellárová E, Spiteller M (2009) Light-independent metabolomics of endophytic *Thielavia subthermophila* provides insight into microbial hypericin biosynthesis. J Nat Prod 72:1825–1835
- Kuznetsova ES, Motiejūnaitė J, Stepanchikova IS, Himelbrant DE, Czarnota P (2012) New records of lichens and allied fungi from the Leningrad Region, Russia. III. Folia Cryptogamica Estonica 49:31–37
- Lackner M, Klaassen CH, Meis JF, van den Ende AG et al (2012) Molecular identification tools for sibling species of *Scedosporium* and *Pseudallescheria*. Med Mycol 50:497–508
- Lachance MA, Kurtzman CP (2013) The yeast genus *Tortispora* gen. nov., description of *Tortispora ganteri* sp. nov., *Tortispora mauiana* sp. nov., *Tortispora agaves* sp. nov., *Tortispora sangerardensis* sp. nov., *Tortispora cuajiniquilana* sp. nov., *Tortispora starmeri* sp. nov. and *Tortispora phaffii* sp. nov., reassignment of *Candida caseinolytica* to *Tortispora caseinolytica* fa. comb. nov., emendation of *Botryozyma*, and assignment of *Botryozyma*, *Tortispora* gen. nov. and *Trigonopsis* to the family *Trigonopsidaceae* fam. nov. Int J Syst Evol Microbiol 63:3104–3114
- Lachance MA, Perri AM, Farahbakhsh AS, Starmer WT (2013) Genetic structure of *Kurtzmaniella cleridarum*, a cactus flower beetle yeast of the Sonoran and Mojave Deserts: speciation by distance? FEMS Yeast Res 13:674–861
- Lackner M, de Hoog GS, Yang L, Moreno LF et al (2014) Proposed nomenclature for *Pseudallescheria*, *Scedosporium* and related genera. Fungal Divers 67:1–10
- Læssøe T, Srikitkulchai P, Luangsa-ard JJD, Stadler M (2013) *Theissenia* reconsidered, including molecular phylogeny of the type species *T. pyrenocrata* and a new genus *Durotheca* (Xylariaceae, Ascomycota). IMA Fungus 4:57–69
- Laflamme G, Smerlis E (2012) *Gremmeniella balsamea* sp. nov. on balsam fir in Canada. N Am Fungi 16(7):1–4
- Lafalmme G, Broders KD, Cote C, Munck I et al (2015) Priority of *Lophophacidium* over *Canavirgella*: taxonomic status of *Lophophacidium dooksi* and *Canavirgella banfieldii*, causal agents of a white pine needle disease. Mycologia 107:45–75
- Lahlali R, McGregor L, Song T, Gossen BD et al (2014) *Heteroconium chaetospira* induces resistance to clubroot via upregulation of host genes involved in jasmonic acid, ethylene, and auxin biosynthesis. PLOS ONE 9:e94144
- Lamit LJ, Bowker MA, Holeski LM, Næsborg RR et al (2011) Genetically-based trait variation within a foundation tree species influences a dominant bark lichen. Fung Ecol 4:103–109
- Landeros F, Iturriaga T, Rodríguez A, Vargas-Amado G, Guzmán-Dávalos L (2015) Advances in the phylogeny of *Helvella* (Fungi: Ascomycota), inferred from nuclear ribosomal LSU sequences and morphological data. Revista Mexicana de Biodiversidad 86:856–871
- Lange OL, Belnap J (2016) How Biological soil crusts became recognized as a functional unit: a selective history. In: Biological soil crusts: an organizing principle in drylands. Springer International Publishing, pp 15–33
- Lantieri A, Pfister DH (2010) A new species of *Marcelleina* from Italy. Mycotaxon 111:465–469
- Lantieri A, Johnston PR, Park D, Lantz H, Merdadi G (2011) *Hypoderma siculum* sp. nov. from Italy. Mycotaxon 118:393–401
- Lantieri A, Smith ME, Pfister DH (2012) A new species of *Ruhlandiella* (Pezizaceae) from Italy. Mycol Prog 11:509–513
- Lantz H, Johnston PR, Park D, Minter DW (2011) Molecular phylogeny reveals a core clade of *Rhytismatales*. Mycologia 103:57–74
- Lapmak K, Lumyong S, Thongkuntha S, Wongputtisin P, Sardsud U (2010) L-asparaginase production by Bipolaris sp. BR438 isolated from brown rice in Thailand. Chiang Mai J Sci 37:160–164
- Latha P, Anand T, Ragupathi N, Prakasam V, Samiyappan R (2009) Antimicrobial activity of plant extracts and induction of systemic resistance in tomato plants by mixtures of PGPR strains and Zimmu leaf extract against *Alternaria solani*. Biol Control 50:85–93
- Laundon JR (2012) Proposal to conserve *Chrysotrichia*, nom. cons., against an additional name, *Alysphaeria* (lichenised Ascomycota). Taxon 61:1321
- Lawrence CB, Mitchell TK, Craven KD, Cho YR, Cramer RA, Kim KH (2008) At death's door: *Alternaria* pathogenicity mechanisms. Plant Pathol J 24:101–111
- Lawrence DP, Gannibal PB, Peever TL, Pryor BM (2013) The sections of *Alternaria*: formalizing species-group concepts. Mycologia 105:530–546
- Lawrey JD, Diederich P, Nelsen MP, Sikaroodi M et al (2011) The obligately lichenicolous genus *Lichenoconium* represents a novel lineage in the Dothideomycetes. Fungal Biol 115:176–187
- Lawrey JD, Marson G, Etayo J, Dal-Forno M et al (2015) Molecular data support establishment of a new genus for the lichenicolous species *Neobarya usneae* (Hypocreales). Bryologist 118:83–92
- Le Pogam P, Le Lamer AC, Siva B, Legouin B et al (2016) Minor pyranonaphthoquinones from the apothecia of the lichen *Ophioparma ventosa*. J Nat Prod 79:1005–1011

- Leão-Ferreira SM, Pascholati Gusmão LF, Castañeda Ruiz RF (2013) Conidial fungi from the semi-arid Caatinga biome of Brazil. Three new species and new records. *Nova Hedw* 96:479–494
- Leão-Ferreira SM, Gusmão LF, de Almeida DA, Castañeda-Ruiz RF (2015) *Digicatenosporium polyramosum*, a new hyphomycete from Brazil. *Mycotaxon* 130:479–482
- Leavitt SD, Clair LLS (2011) Estimating *Xanthoparmelia* (Parmeliaceae) population density in subalpine communities in southern Utah, USA using two distance methods, with implications for assessing community composition. *Bryologist* 114:625–636
- Leavitt SD, Fankhauser JD, Leavitt DH, Porter LD et al (2011a) Complex patterns of speciation in cosmopolitan “rock posy” lichens—discovering and delimiting cryptic fungal species in the lichen-forming *Rhizoplaca melanophthalma* species-complex (Lecanoraceae, Ascomycota). *Mol Phylogenet Evol* 59:587–602
- Leavitt SD, Johnson LA, Goward T, Clair LLS (2011b) Species delimitation in taxonomically difficult lichen-forming fungi: an example from morphologically and chemically diverse *Xanthoparmelia* (Parmeliaceae) in North America. *Mol Phylogenet Evol* 60:317–332
- Leavitt SD, Johnson LA, St Clair LL (2011c) Species delimitation and evolution in morphologically and chemically diverse communities of the lichen-forming genus *Xanthoparmelia* (Parmeliaceae, Ascomycota) in western North America. *Am J Bot* 98:175–188
- Leavitt SD, Esslinger TL, Lumbsch HT (2012) Neogene-dominated diversification in neotropical diversification in neotropical (Parmeliaceae). *Am J Bot* 99:1764–1777
- Leavitt SD, Lumbsch HT, St Clair LL (2013) Contrasting demographic histories of two species in the lichen-forming fungal genus *Xanthomendoza* (Teloschistaceae, Ascomycota). *Bryologist* 116:337–349
- Leavitt SD, Divakar PK, Ohmura Y, Wang LS et al (2015) Who's getting around? Assessing species diversity and phylogeography in the widely distributed lichen-forming fungal genus *Montanelia* (Parmeliaceae, Ascomycota). *Mol Phylogenet Evol* 90:85–96
- Lechat C, Courtecuisse R (2010) A new species of *Ijuhya*, *I. antillana*, from the French West Indies. *Mycotaxon* 113:443–447
- Lechat C, Fournier J (2013) Two new species of *Lasionectria* (Bionectriaceae, Hypocreales) from Guadeloupe and Martinique (French West Indies). *Mycotaxon* 121:275–280
- Lechat C, Fournier J (2017) Four new species of *Ijuhya* (Bionectriaceae) from Belgium, metropolitan France and French Guiana. *Ascomycete.org* 9:1–18
- Lechat C, Farr DF, Hirooka Y, Minnis AM, Rossman AY (2010) A new species of *Hydropsphaera*, *H. bambusicola*, is the sexual state of *Gliomastix fusigera*. *Mycotaxon* 111:95–102
- Lee YB, Na YH (2009) A new species of the genus *Coreomyces* (Laboulbeniales, Ascomycotina) collected from the island of Java, Indonesia. *Mycobiology* 37:300–301
- Lee YB, Na YH (2011) A new species of the genus *Euphoriomycetes* (Laboulbeniales Ascomycotina) collected in Korea. *Mycobiology* 39:59
- Lee JS, Min GH, Lee JS (2009a) Nutritional and physicochemical characteristics of the antidementia acetylcholinesterase-inhibiting methanol extracts from *Umbilicaria esculenta*. *Mycobiology* 37:203–206
- Lee SB, Go YS, Bae HJ, Park JH et al (2009b) Disruption of glycosylphosphatidylinositol-anchored lipid transfer protein gene altered cuticular lipid composition, increased plastoglobules, and enhanced susceptibility to infection by the fungal pathogen *Alternaria brassicicola*. *Plant Physiol* 150:42–54
- Lee SJ, Strobel GA, Eisenman K, Geary B et al (2009c) *Aurosphaeria*, a novel coelomycetous genus. *Mycotaxon* 107:463–472
- Lee SK, Lee SH, Cho SE, Park JH, Shin HD (2011) Confirmation of the Presence of *Pleochaeta shiriana* associated with powdery mildew of *Celtis sinensis* in Korea. *Res Plant Dis* 17:391–395
- Lee SS, Alias SA, Jones EBG, Zainuddin N, Chan HT (2012) Checklist of fungi of Malaysia. Research Pamphlet No.132. Swan Printing Sdn. Bhd., Seri Kembangan, Malaysia
- Lee LW, Wang GJ, Lin MH, Ju YM et al (2013a) Isolation and characterization of sesquiterpenes from *Arecophila saccharicola* YMJ96022401 with NO production inhibitory activity. *Phytochemistry* 85:129–136
- Lee SC, Han KS, Park JH, Cho SE, Shin HD (2013b) First report of frosty mildew caused by *Mycopappus alni* on Asian pear in Korea. *J Phytopathol* 161:866–869
- Lee S, Tamayo-Castillo G, Pang C, Clardy J et al (2016) Diketopiperazines from Costa Rican endolichenic fungus *Colpoma* sp. CR1465A. *Bioorg Med Chem Lett* 26:2438–2441
- Lendemer JC (2009) Another record of the rare and endangered species *Phaeophyscia leana*. *Evansia* 26:142–143
- Lendemer JC (2010a) *Lepraria larrainiana* (Stereocaulaceae, Lichenized Ascomycetes), a new species from Central Chile. *Gayana Bot* 67:238–241
- Lendemer JC (2010b) Notes on *Lepraria* s.l. (Lecanoromycetes, Ascomycota) in North America: New species, new reports, and preliminary keys. *Brittonia* 62:267–292
- Lendemer JC (2010c) Reassessment of the genus *Catillochroma* (lichenized Ascomycota, Ramalinaceae). *Lichenologist* 42:587–600
- Lendemer JC (2011) A taxonomic revision of the North American species of *Lepraria* s.l. that produce divaricatic acid, with notes on the type species of the genus *L. incana*. *Mycologia* 103:1216–1229
- Lendemer JC (2012a) Contributions to the Lichen Flora of Pennsylvania.—Further New and Interesting Reports of Lichens and Lichenicolous Fungi, Part 2
- Lendemer JC (2012b) Perspectives on chemotaxonomy: molecular data confirm the existence of two morphologically distinct species within a chemically defined *Lepraria caesiella* (Stereocaulaceae). *Castanea* 77:89–105
- Lendemer JC (2013) Molecular phylogenetic evidence corroborates morphology but not chemistry in the *Lepraria neglecta* group (Stereocaulaceae). *Mem N Y Bot Garden* 108:127–153
- Lendemer JC, Knudsen K (2008) Studies in lichens and lichenicolous fungi: further notes on North American taxa. *Mycotaxon* 103:75–86
- Lendemer JC, Hodkinson BP (2012a) *Chirleja buckii*, a new genus and species of lichenized-fungi from Tierra del Fuego, southern South America. *N Z J Bot* 50:449–456
- Lendemer JC, Hodkinson BP (2012b) Recognition of the *Parmelia crozalsiana* group as the genus *Crespoa*. *N Am Fungi* 7:1–5
- Lendemer JC, Hodkinson BP (2013) A radical shift in the taxonomy of *Lepraria* s.l.: molecular and morphological studies shed new light on the evolution of asexuality and lichen growth form diversification. *Mycologia* 105:994–1018
- Lendemer JC, Harris RC (2014) Studies in lichens and lichenicolous fungi—No. 19: further notes on species from the coastal plain of southeastern North America. *Opusc Phil* 13:155–176
- Lendemer JC, Harris RC (2015) *Xyleborus nigricans*, a second species for the previously monospecific genus newly found in the Mid-Atlantic Coastal Plain of North America. *Bryologist* 118:284–292
- Lendemer JC, Kocourková J, Knudsen K (2009) Studies in lichen and lichenicolous fungi: more notes on taxa from North America. *Mycotaxon* 108:491–497

- Lendemer JC, Harris RC, Tripp EA (2013) The lichens and allied fungi of Great Smoky Mountains National Park. Mem N Y Bot Gard 104:1–152
- Lendemer JC, Harris RC, Ruiz AM (2016) A review of the lichens of the dare regional biodiversity hotspot in the mid-atlantic coastal plain of North Carolina, eastern North America. Castanea 81:1–77
- Leonardo L, Mazzilli BP, Damatto SR, Saiki M et al (2011) Assessment of atmospheric pollution in the vicinity of a tin and lead industry using lichen species *Canoparmelia texana*. J Environ Radioact 102:906–910
- Letourneau A, Seena S, Marvanová L, Bärlocher F (2010) Potential use of barcoding to identify aquatic hyphomycetes. Fungal Divers 40:51–64
- Leuchtmann A, Clemenccon H (2012) The taxonomic position of the genus *Heydenia* (*Pyronemataceae*, *Pezizales*) based on molecular and morphological data. Mycol Prog 11:699–710
- Lewis Smith RI (2009) Further additions to the lichen flora of Antarctica and South Georgia. Nova Hedw 88:157–168
- Li DM, de Hoog GS, Lindhardt Saunte DM, Gerrits van den Ende AHG et al (2008a) *Coniosporium epidermidis* sp. nov., a new species from human skin. Stud Mycol 61:131–136
- Li DW, Cowles R, Vossbrinck C (2008b) *Metarhiziosis microspora* gen. et sp. nov. associated with the elongate hemlock scale. Mycologia 100:460–466
- Li J, Yang J, Liang L, Zhang KQ (2008c) Taxonomic revision of the nematode-trapping fungus *Arthrobotrys multisequendaria*. J Microbiol 46:513–518
- Li S, Yu Z, Zhang Y, Qiao M et al (2009a) *Arthrobotrys nonseptata*, a new anamorph from an *Orbilia* species. Mycotaxon 109:247–254
- Li XF, Lu T, Lu BS, Hyde KD (2009b) *Emarcea rostrispora* nom. nov. (*Xylariaceae*, *Ascomycota*) on wood from Taiwan. Mycosistema 28:151–153
- Li YM, Haixia WU, Hang C, Hyde KD (2011) Morphological studies in Dothideomycetes: *Elsinoe* (*Elsinoaceae*), *Butleria* and three excluded genera. Mycotaxon 115:507–520
- Li HY, Sun GY, Zhai XR, Batzer JC et al (2012a) *Dissocloniaceae* associated with sooty blotch and flyspeck on fruits in China and the United States. Persoonia 28:113–125
- Li WH, Xiao YX, Wang CS, Zhang R et al (2012b) A new species of *Devriesia* causing sooty blotch and flyspeck on rubber tree in China. Mycol Prog 12:733–738
- Li DW, Zhao G, Yang C, Jalsrai A, Kerin B (2013) Four noteworthy hyphomycetes from indoor environments. Mycotaxon 125:111–121
- Li SH, Zheng LY, Liu CY, Wang Y et al (2014a) Two new truffles species, *Tuber alboumbilicum* and *Tuber pseudobrumeale* from China. Mycol Prog 13:1157–1163
- Li WJ, Bhat JD, Hyde KD, Wang Y (2014b) Towards a natural classification of Dothideomycetes 4: The genera *Bryopelta*, *Bryorella*, *Bryosphaeria*, *Lophiosphaera* and *Maireella* (*Dothideomycetes incertae sedis*). Phytotaxa 176:28–41
- Li WJ, Liu JK, Bhat DJ, Camporesi E et al (2014c) Introducing the novel species, *Dothiorella synphoricarpicola*, from snowberry in Italy. Cryptogam Mycol 35(3):257–270
- Li ZJ, Cao N, Chen HF, Taylor JE, Hou CL (2014d) New species and new records of Rhytismataceae from Japan. Mycol Prog 13:951–958
- Li JY, Qiao M, Peng J, Qian WY et al (2015a) *Uncispora hainanensis* sp. nov. isolated from decayed leaves. Mycotaxon 129:473–476
- Li Q, Chen YX, Zheng CT, Lu DD, Lin YR (2015b) *Coccozymes neolitseaie* sp. nov. from Zhangjiajie, China. Mycotaxon 130:289–293
- Li WJ, Bhat DJ, Camporesi E, Tian Q, Wijayawardene NN et al (2015c) New asexual morph taxa in *Phaeosphaeriaceae*. Mycosphere 6:681–708
- Li WJ, Maharachchikumbura SSN, Li QR, Bhat DJ et al (2015d) Epitypification of *Broomella vitaliae* and introduction of a novel species of *Hyalotiella*. Cryptog Mycol 36:93–108
- Li GJ, Hyde KD, Zhao RL, Hongsanan S et al (2016a) Fungal diversity notes 253–366: taxonomic and phylogenetic contributions to fungal taxa. Fungal Divers 78:1–237
- Li J, Phookamsak R, Mapook A, Boonmee S, Bhat JD et al (2016b) *Seifertia shangrilaensis* sp. nov. (*Melanommataceae*), a new species from Southwest China. Phytotaxa 273:34–42
- Li JF, Bhat DJ, Phookamsak R, Mapook A, Lumyong S, Hyde KD (2016c) *Sporidesmioides thailandica* gen. et sp. nov. (*Dothideomycetes*) from northern Thailand. Mycol Prog 15:1169–1178
- Li JF, Phookamsak R, Jeewon R, Bhat DJ et al (2016d) Molecular taxonomy and morphological characterization reveal new species and new host records of *Torula* species (*Torulaceae*, *Pleosporales*). Mycol Prog 16:447–461
- Li JF, Jeewon R, Luo Z, Phookamsak R, Bhat DJ et al (2017a) Morphological characterization and DNA based taxonomy of *Fusiconidium* gen. nov. with two novel taxa within *Melanommataceae* (*Pleosporales*). Phytotaxa 308:206–218
- Li Y, Xiao J, de Hoog GS, Wang X et al (2017b) Biodiversity and human-pathogenicity of *Phialophora verrucosa* and relatives in Chaetothyriales. Persoonia 38:1–19
- Liang J, Han Y, Du W, Liang Z, Li Z (2009a) *Chrysosporium linfenense*: a new *Chrysosporium* species with keratinolytic activity. Mycotaxon 110:65–71
- Liang ZQ, Han YF, Chu HL, Fox RT (2009b) Studies on the genus *Paecilomyces* in China V. *Taifanglania* gen. nov. for some monophialidic species. Fungal Divers 31:34–69
- Liang L, Wu H, Liu Z, Shen R et al (2013) Proteomic and transcriptional analyses of *Arthrobotrys oligospora* cell wall related proteins reveal complexity of fungal virulence against nematodes. Appl Microbiol Biotechnol 97:8683–8692
- Liang L, Shen R, Mo Y, Yang J et al (2015) A proposed adhesin AoMad1 helps nematode-trapping fungus *Arthrobotrys oligospora* recognizing host signals for life-style switching. Fung Gen Biol 81:172–181
- Ligoxigakis EK, Papaioannou IA, Markakis EA, Typas MA (2013) First report of pink rot of *Phoenix* and *Washingtonia* species caused by *Nalanthamala vermoesenii* in Greece. Plant Dis 97:285
- Lim YW, Kim BK, Kim C, Jung HS et al (2010) Assessment of soil fungal communities using pyrosequencing. J Microbiol 48:284–289
- Lima NB, de Batista MV, De Moraes Jr MA, Maria A et al (2013) Five *Colletotrichum* species are responsible for mango anthracnose in northeastern Brazil. Fungal Divers 61(1):75–88
- Limtong S, Jindamorakot S, Am-In S, Kaewwichian R et al (2011) *Candida uthaihanina* sp. nov., an anamorphic yeast species in *Nakaseomyces* clade isolated in Thailand. Antonie van Leeuwenhoek 99:865–871
- Lin CH, Chung KR (2010) Specialized and shared functions of the histidine kinase-and HOG1 MAP kinase-mediated signaling pathways in *Alternaria alternata*, a filamentous fungal pathogen of citrus. Fung Gen Biol 47:818–827
- Lin CH, Yang SL, Chung KR (2009) The YAP1 homolog-mediated oxidative stress tolerance is crucial for pathogenicity of the necrotrophic fungus *Alternaria alternata* in citrus. Mol Plant Microbe Interact 22:942–952
- Lin CH, Yang SL, Wang NY, Chung KR (2010) The FUS3 MAPK signaling pathway of the citrus pathogen *Alternaria alternata* functions independently or cooperatively with the fungal redox-responsive AP1 regulator for diverse developmental, physiological and pathogenic processes. Fung Gen Biol 47:381–391

- Lin CH, Yang SL, Chung KR (2011) Cellular responses required for oxidative stress tolerance, colonization, and lesion formation by the necrotrophic fungus *Alternaria alternata* in citrus. *Curr Microbiol* 62:807–815
- Lin CG, Chen Y, McKenzie EH, Bhat DJ et al (2016a) The genus *Fusariella*. *Mycol Prog* 12:1313–1326
- Lin CG, McKenzie EH, Bhat DJ, Ran SF et al (2016b) Stachybotrys-like taxa from karst areas and a checklist of stachybotrys-like species from Thailand. *Mycosphere* 7:1273–1291
- Lin CG, Dai DQ, Bhat DJ, Hyde KD et al (2017) *Subsessila turbinata* gen. et. sp. nov. (*Beltraniaceae*), a Beltrania-like fungus from Thailand. *Mycol Prog* 4:393–401
- Linaldeddu BT, Deidda A, Scana B, Franceschini A et al (2015) Diversity of *Botryosphaeriaceae* species associated with grapevine and other woody hosts in Italy, Algeria and Tunisia, with descriptions of *Lasiodiplodia exigua* and *Lasiodiplodia mediterranea* sp. nov. *Fungal Divers* 71:201–214
- Linaldeddu BT, Alves A, Phillips AJL (2016) *Sardiniella urbana* gen. et sp. nov., a new member of the *Botryosphaeriaceae* isolated from declining *Celtis australis* trees in Sardinian streetscapes. *Mycosphere* 7:893–905
- Lindblom L (2014) *Xanthomendoza poeltii* is a synonym of *X. oregana* (*Teloschistaceae*, lichen-forming ascomycetes). *Lichenologist* 46:829–832
- Lindblom L, Söchting U (2013) Genetic diversity of the photobiont of the bipolar lichen-forming ascomycete *Xanthomendoza borealis*. *Herzogia* 26:307–322
- Lindemann U, Böhning T (2016) *Tricharina glabra* (Pezizales)—eine neue Art in einer schwierigen Gattung. *Z Mykol* 82:449–458
- Lindemann U, Vega M, Richter T, Alvarado P (2014) *Octosporopsis nicolai*—a mysterious member of the *Pyronemataceae*. *Z Mykol* 80:565–592
- Lindemann U, Vega M, Alvarado P (2015) Revision der Gattung *Kotlabaea*: *K. deformis*, *K. delectans* und *K. Benkertii*. *Dtsch Ges Mykol* 81:373–402
- Linnakoski R, de Beer ZW, Rousi M, Niemelä P et al (2008) Fungi, including *Ophiostoma karelicum* sp. nov., associated with *Scolytus ratzeburgii* infesting birch in Finland and Russia. *Mycol Res* 112:1475–1488
- Linnakoski R, de Beer ZW, Ahtiainen J, Sidorov E et al (2010) *Ophiostoma* spp. associated with pine- and spruce-infesting bark beetles in Finland and Russia. *Persoonia* 25:72–93
- Linnakoski R, de Beer ZW, Duong TA, Niemelä P et al (2012) *Grosmannia* and *Leptographium* spp. associated with conifer-infesting bark beetles in Finland and Russia, including *Leptographium taigensis* sp. nov. *Antonie van Leeuwenhoek* 102:375–399
- Linnakoski R, Mahilainen S, Harrington A, Vanhanen H et al (2016) Seasonal succession of fungi associated with *Ips typographus* beetles and their phoretic mites in an outbreak region of Finland. *PLoS ONE* 11:e0155622
- Lisov A, Zavarzina A, Zavarzin A, Demin V, Leontievsky A (2012) Dimeric and monomeric laccases of soil-stabilizing lichen *Solorina crocea*: purification, properties and reactions with humic acids. *Soil Biol Biochem* 45:161–167
- Lisowska M (2011) Lichen recolonisation in an urban-industrial area of southern Poland as a result of air quality improvement. *Environ Monit Assess* 179:177–190
- Liu F, Cai L (2012) Morphological and molecular characterization of a novel species of *Simplicillium* from China. *Cryptogam Mycol* 33:137–144
- Liu JK, Jones EBG, Chukeatirote E, Bahkali AH, Hyde KD (2011a) *Lignincola conchicola* from palms with a key to the species of *Lignincola*. *Mycotaxon* 117:343–349
- Liu JK, Phookamsak R, Jones EBG, Zhang Y et al (2011b) *Astrospheeriaella* is polyphyletic, with species in *Fissuroma* gen. nov., and *Neoastrospheeriaella* gen. nov. *Fungal Divers* 51:135–154
- Liu F, Hu DM, Cai L (2012a) *Conlarium duplumascospora* gen. et. sp. nov. and *Jobellisia guangdongensis* sp. nov. from freshwater habitats in China. *Mycologia* 104:1178–1186
- Liu JK, Phookamsak R, Doilom M, Wikee S et al (2012b) Towards a natural classification of *Botryosphaerales*. *Fungal Divers* 57:149–210
- Liu TT, Hu DM, Liu F, Cai L (2013a) Polyphasic characterization of *Plectosphaerella oligotrophica*, a new oligotrophic species from China. *Mycoscience* 54:387–393
- Liu YX, Hyde KD, Ariyawansa HA, Li WJ et al (2013b) *Shiraiaceae*, new family of *Pleosporales* (*Dothideomycetes*, *Ascomycota*). *Phytotaxa* 103:51–60
- Liu JK, Phookamsak R, Dai DQ, Tanaka K et al (2014a) *Rousselaceae*, a new pleosporalean family to accommodate the genera *Neoroussella* gen. nov., *Roussella* and *Rousselopsis*. *Phytotaxa* 181:1–33
- Liu K, Zhang W, Lai Y, Xiang M et al (2014b) *Drechslerella stenobrocha* genome illustrates the mechanism of constricting rings and the origin of nematode predation in fungi. *Bmc Genom* 15:114
- Liu S, Su H, Su X, Zhang F et al (2014c) *Arthrobotrys xiangyunensis*, a novel nematode-trapping taxon from a hot-spring in Yunnan Province, China. *Phytotaxa* 174:89–96
- Liu JK, Hyde KD, Jones EBG, Ariyawansa HA et al (2015a) Fungal diversity notes 1–110: taxonomic and phylogenetic contributions to fungal species. *Fungal Divers* 72:1–197
- Liu Q, Wang Y, Luo H, Wang L et al (2015b) Isolation of a novel cold-active family 11 Xylanase from the filamentous fungus *Bispora antennata* and deletion of its N-terminal amino acids on thermostability. *Appl Biochem Biotechnol* 175:925–936
- Liu JK, Yang J, Maharachchikumbura SS, McKenzie EH et al (2016a) Novel chaetosphaeriaceous hyphomycetes from aquatic habitats. *Mycol Prog* 10:1157–1167
- Liu XX, Zhuang WY, Zeng ZQ, Zhao P (2016b) Newly discovered sclerotiniaceous fungi from China. *Nova Hedwig* 102:347–357
- Liu JK, Hyde KD, Jeewon R, Phillips AJL et al (2017) Ranking higher taxa using divergence times: a case study in Dothideomycetes. *Fungal Divers* 84:75–99
- Logesh AR, Kalaiselvam M, Upadhyay DK (2013) New species of non-lichenized ascomata from India. *Mycosphere* 4:152–154
- Logesh AR, Chinlampianga M, Shukla AC, Upadhyay DK (2015) Studies on lichens of *Mizoram*, northeast India. *Proc Natl Acad Sci USA* 31:1–13
- Logrieco A, Moretti A, Solfrizzo M (2009) *Alternaria* toxins and plant diseases: an overview of origin, occurrence and risks. *World Mycotoxin J* 2:129–140
- Löhmus A, Löhmus P (2010) Epiphyte communities on the trunks of retention trees stabilise in 5 years after timber harvesting, but remain threatened due to tree loss. *Biol Conserv* 143:891–898
- Loizides M, Carbone M, Alvarado P (2015) *Geoglossum dunense* (*Ascomycota*, *Geoglossales*): a new species from the Mediterranean islands of Cyprus and Malta. *Mycol Prog* 14:41
- Loizides M, Bellanger JM, Clowez P, Richard F, Moreau PA (2016) Combined phylogenetic and morphological studies of true morels (*Pezizales*, *Ascomycota*) reveal Cyprus as the new hotspot of *Morchella* diversity. *Mycol Prog* 15:1–28
- Lombard L, Crous PW (2012) Phylogeny and taxonomy of the genus *Gliocladiopsis*. *Persoonia* 28:25–33
- Lombard L, Crous PW, Wingfield BD, Wingfield MJ (2010a) Phylogeny and systematics of the genus *Calonectria*. *Stud Mycol* 66:31–69

- Lombard L, Crous PW, Wingfield BD, Wingfield MJ (2010b) Species concepts in *Calonectria* (*Cylindrocladum*). Stud Mycol 66:1–14
- Lombard L, Shivas RG, To-Anun C, Crous PW (2012) Phylogeny and taxonomy of the genus *Cylindrocladiella*. Mycol Prog 11:835–868
- Lombard L, van der Merwe NA, Groenewald JZ, Crous PW (2014) Lineages in *Nectriaceae*: re-evaluating the generic status of *Ilyonectria* and allied genera. Phytopathol Mediterr 53:515–532
- Lombard L, Chen SF, Mou X, Zhou XD et al (2015a) New species, hyper-diversity and potential importance of *Calonectria* spp. from *Eucalyptus* in South China. Stud Mycol 80:151–188
- Lombard L, van der Merwe NA, Groenewald JZ, Crous PW (2015b) Generic concepts in *Nectriaceae*. Stud Mycol 80:189–245
- Lombard L, Houbraeken J, Decock C, Samson RA, Meijer M et al (2016) Generic hyper-diversity in *Stachybotriaceae*. Persoonia 36:156–246
- London LS (2016) *Schismatomma galactinum* (lichenized ascomycota), a lichen species new to India. Curr Sci 110:770
- López de Silanes ME, Etayo J, Paz-Bermúdez G (2009) *Pronectria pilosa* (*Hypocreaceae*) sp. nov. and other lichenicolous fungi found on Collemataceae in the Iberian Peninsula. Bryologist 112:101–108
- Loppi S, Nascimbene J (2010) Monitoring H2S air pollution caused by the industrial exploitation of geothermal energy: the pitfall of using lichens as bioindicators. Environ Pollut 158:2635–2639
- Lorch JM, Minnis AM, Meteyer CU, Redell JA et al (2014) The fungus *Trichophyton redellii* sp. nov. causes skin infections that resemble white-nose syndrome of hibernating bats. J Wildl Dis 51:36–47
- Lou J, Fu L, Peng Y, Zhou L (2013) Metabolites from *Alternaria* fungi and their bioactivities. Molecules 18:5891–5935
- Lu M, Zhou XD, De Beer ZW, Wingfield MJ, Sun JH (2009) Ophiostomatoid fungi associated with the invasive pine-infesting bark beetle, *Dendroctonus valens*, in China. Fungal Divers 38:133–145
- Lu YZ, Liu JK, Hyde KD, Bhat DJ et al (2016) *Brunneodinemasporium jonesii* and *Tainosphaeria jonesii* spp. nov. (*Chaetosphaeriaceae*, *Chaetosphaerales*) from southern China. Mycosphere 7:1323–1332
- Lu YZ, Boonmee S, Dai DQ, Liu JK et al (2017a) Four new species of *Tubeufia* (*Tubeufiaceae*, *Tubeufiales*) from Thailand. Mycol Prog 16:403–417
- Lu YZ, Boonmee S, Liu JK, Hyde KD et al (2017b) Novel *Neoacanthostigma* species from aquatic habitats. Cryptogam Mycol 38:169–190
- Luangsa-ard JJ, Houbraeken J, van Doorn T et al (2011) *Purpureocillium*, a new genus for the medically important *Paecilomyces lilacinus*. FEMS Microbiol Lett 321:141–149
- Luangsuphabool T, Piapukiew J, Parnmen S, Nelsen MP et al (2016) Diversity of the *Trypethelium eluteriae* group in Thailand (Ascomycota, Trypetheliaceae). Lichenologist 48:53–60
- Łubek A (2012) The lichen biota of Skałki Piekło pod Niekłaniem “Nature Reserve-current state and changes in species composition over the past 100 years. Pol J Nat Sci 27:135–150
- Lücking R (2008) Foliicolous lichenized fungi. Flora Neotrop Monogr 103:1–866
- Lücking R (2014) A key to species of the *Ocellularia papillata*, *perforata* and *terebrata* morphodemes. Glalia 6:1–35
- Lücking R, Breuss O (2012) A new species of *Thelotrema*, a new combination, *Leucodecon isidioides*, and a key to thelotremoid lichens of Macaronesia (lichenised Ascomycota: Graphidaceae). Österr Z Pilzk 21:127–133
- Lücking R, Moncada B (2017) Dismantling *Marchandiophalina* into *Agonimia* (Verrucariaceae) and *Lawreymyces* gen. nov. (Corticiciaceae): setting a precedent to the formal recognition of thousands of voucherless fungi based on type sequences. Fungal Divers 84:119–138
- Lücking R, Chaves JL, Sipman HJM, Umana L, Aptroot A (2008a) A first assessment of the Ticolichen biodiversity inventory in Costa Rica: the genus *Graphis*, with notes on the genus *Hemithecium* (Ascomycota: Ostropales: Graphidaceae). Fieldiana Bot New Ser 46:1–140
- Lücking R, Lumbsch HT, Di Stefano JF, Lizano D et al (2008b) *Eremithallus costaricensis* (Ascomycota: Lichinomycetes: Eremothallales), a new fungal lineage with a novel lichen symbiotic lifestyle discovered in an urban relict forest in Costa Rica. Symbiosis 46:161–170
- Lücking R, Archer AW, Aptroot A (2009) A world-wide key to the genus *Graphis* (Ostropales: Graphidaceae). Lichenologist 41:363–452
- Lücking R, Rivas Plata E, Kalb K, Common R et al (2011) *Halegrapha* (Ascomycota: Graphidaceae), an enigmatic new genus of tropical lichenized fungi dedicated to Mason E. Hale. Jr. Lichenologist 43:331–344
- Lücking R, Parnmen S, Lumbsch HT (2012) *Mangoldia*, a new lichen genus in the family Graphidaceae (Ascomycota: Ostropales). Phytotaxa 69:1–5
- Lücking R, Tehler A, Bungartz F, Rivas Plata E, Lumbsch HT (2013) Journey from the West: did tropical Graphidaceae (lichenized Ascomycota: Ostropales) evolve from a saxicolous ancestor along the American Pacific coast. Am J Bot 100:844–856
- Lücking R, Johnston MK, Aptroot A, Kraichak E et al (2014) One hundred and seventy-five new species of Graphidaceae: closing the gap or a drop in the bucket? Phytotaxa 189:7–38
- Lücking R, Gutiérrez MC, Moncada B (2016a) *Neosergipea*, a new name for the lichen fungus *Sergipea*, with an updated phylogeny and notes on the genus *Dichosporidium* (lichenized Ascomycota: Arthoniales: Roccellaceae). Lichenologist 48:269–273
- Lücking R, Nelsen MP, Aptroot A, Barillas de Klee R et al (2016b) A phylogenetic framework for reassessing generic concepts and species delimitation in the lichenized family Trypetheliaceae (Ascomycota: Dothideomycetes). Lichenologist 48:739–762
- Lücking R, Hodkinson BP, Leavitt SD (2017a) The 2016 classification of lichenized fungi in the Ascomycota and Basidiomycota—approaching one thousand genera. Bryologist 119:361–416
- Lücking R, Hodkinson BP, Leavitt SD (2017b) Corrections and amendments to the 2016 classification of lichenized fungi in the Ascomycota and Basidiomycota. Bryologist 120:58–69
- Lumbsch HT, Huhndorf SM (2010) Outline of Ascomycota 2009. Myconet 14:1–64
- Lumbsch HT, Nelsen MP, Lücking R (2008) The phylogenetic position of *Haematommataceae* (Lecanorales, Ascomycota), with notes on secondary chemistry and species delimitation. Nova Hedwig 86:105–114
- Lumbsch HT, Zimmermann DG, Schmitt I (2009a) Phylogenetic position of ephemeral lichens in Thelocarpaceae and Vezdaeaceae (Ascomycota). Biblioth Lichenol 100:389–398
- Lumbsch HT, Lücking R, Tibell L (2009b) Molecular data place Typholoron as an additional calicoid genus in the Arthoniales (Ascomycota). Biblioth Lichenol 99:285–296
- Lumbsch HT, Divakar PK, Messuti MI, Mangold A, Lücking R (2010a) A survey of thelotremoid lichens (Ascomycota: Ostropales) in subantarctic regions excluding Tasmania. Lichenologist 42:203–224
- Lumbsch HT, Parnmen S, Rangsiruji A, Elix JA (2010b) Phenotypic disparity and adaptive radiation in the genus *Cladonia* (Lecanorales, Ascomycota). Aust Syst Bot 23:239–247
- Lumbsch HT, Ahti T, Altermann S, Amo De Paz G et al (2011) One hundred new species of lichenized fungi: a signature of undiscovered global diversity. Phytotaxa 18:1–127

- Lumbsch T, Kantvilas G, Parmen S (2012) Molecular data support placement of *Cameronia* in Ostropomycetidae (*Lecanoromycetes, Ascomycota*). MycoKeys 30:31–44
- Lumbsch HT, Kraichak E, Parmen S, Rivas Plata E et al (2014a) New higher taxa in the lichen family Graphidaceae (lichenized Ascomycota: Ostropales) based on a three-gene skeleton phylogeny. Phytotaxa 189:89–151
- Lumbsch HT, Parmen S, Kraichak E, Papong KB, Lücking R (2014b) High frequency of character transformations is phylogenetically structured within the lichenized fungal family Graphidaceae (Ascomycota: Ostropales). Syst Biodivers 12:271–291
- Luo J, Zhang N (2013) *Magnaporthiopsis*, a new genus in Magnaportheaceae (Ascomycota). Mycologia 105:1019–1029
- Luo H, Li J, Yang J, Wang H et al (2009a) A thermophilic and acid stable family-10 xylanase from the acidophilic fungus *Bispora* sp. MEY-1. Extremophiles 13:849–857
- Luo H, Wang Y, Wang H, Yang J et al (2009b) A novel highly acidic beta-mannanase from the acidophilic fungus *Bispora* sp. MEY-1: gene cloning and overexpression in *Pichia pastoris*. App Microbiol Biotechol 82:453–461
- Luo H, Yamamoto Y, Kim JA, Jung JS et al (2009c) Lecanoric acid, a secondary lichen substance with antioxidant properties from *Umbilicaria antarctica* in maritime Antarctica (King George Island). Polar Biol 32:1033–1040
- Luo J, Zhuang WY (2010) New species and new Chinese records of Bionectriaceae (Hypocreales, Ascomycota). Mycol Prog 9:17–25
- Luo J, Walsh E, Zhang N (2014) Four new species in Magnaportheaceae from grass roots in New Jersey Pine Barrens. Mycologia 106:580–588
- Luo J, Walsh E, Zhang N (2015a) Toward monophyletic generic concepts in Magnaportheales: species with *Harpophora* asexual states. Mycologia 107:641–664
- Luo ZL, Maharachchikumbur SSN, Liu X, Chen L et al (2015b) *Annulatascus saprophyticus* sp. nov. and *Pseudoannulatascus* gen. nov. to accommodate *Annulatascus biatriisporus* (*Annulatascales, Sordariomycetes*) from Thailand. Phytotaxa 239:174–182
- Luo ZL, Bao DF, Bhat DJ, Yang J et al (2016a) *Sporoschisma* from submerged wood in Yunnan, China. Mycol Prog 15:1145–1155
- Luo ZL, Bahkali AH, Liu XY, Phookamsak R et al (2016b) *Poaceascoma aquaticum* sp. nov. (Lentitheciaceae), a new species from submerged bamboo in freshwater. Phytotaxa 253:71–80
- Luo ZL, Yang J, Liu JK, Su HY et al (2016c) Two new species of *Helicascus* (Morosphaeriaceae) from submerged wood in northern Thailand. Phytotaxa 270:182–190
- Luo ZL, Bhat J, Jeewon R, Boonmee S et al (2017) Molecular phylogeny and morphological characterization of asexual fungi (Tubuliflora) from freshwater habitats in Yunnan, China. Cryptogam Mycol 38:1–28
- Lynch SC, Twizeyimana M, Mayorquin JS, Wang DH et al (2016) Identification, pathogenicity and abundance of *Paracremonium pembeum* sp. nov. and *Graphium euwallaceae* sp. nov.—two newly discovered mycangial associates of the polyphagous shot hole borer (*Euwallacea* sp.) in California. Mycologia 108:313–329
- Lysková P, Hubka V, Navrátilová P, Kolařík M et al (2014) Two cases of fungal infection of nails with participation of *Cryptendoxyla hypophloia*. Klin Mikrobiol Infekc Lek 20:23–26
- Ma J (2016) *Sporidesmiella guangdongensis* and *S. jiangxiensis* spp. nov. on dead branches from southern China. Sydowia 68:113–118
- Ma J, Zhang K, Zhang XG (2008) Two new *Ellisembia* species from Hainan China. Mycotaxon 104:141–145
- Ma J, Ren SC, Ma LG, Zhang YD, Zhang XG (2010a) New records of *Corynesporopsis* from China. Mycotaxon 114:423–428
- Ma J, Zhang YD, Ma LG, Ren SC, Zhang XG (2010b) Taxonomic studies of *Ellisembia* from Hainan China. Mycotaxon 114:417–421
- Ma J, Ma LG, Zhang YD, Zhang XG (2011a) Three new hyphomycetes from southern China. Mycotaxon 117:247–253
- Ma J, Wang Y, Ma LG, Zhang YD, Castañeda-Ruiz RF, Zhang XG (2011b) Three new species of *Neosporidesmium* from Hainan China. Mycol Prog 10:157–162
- Ma LG, Ma J, Zhang YD, Zhang XG (2011c) *Craspedodidymum* and *Corynespora* spp. nov. and a new anamorph recorded from southern China. Mycotaxon 117:351–358
- Ma HX, Vasilyeva L, Li Y (2012a) The genus *Xylaria* in the south of China—3. *X. atroglobosa* sp. nov. Mycotaxon 119:381–384
- Ma J, Ma LG, Zhang YD, Castañeda-Ruiz RF, Zhang XG (2012b) New species and record of *Corynesporopsis* and *Hemicorynespora* from southern China. Nova Hedwig 95:233–241
- Ma J, Ma LG, Zhang YD, Castañeda-Ruiz RF, Zhang XG (2012c) New species or records of *Endophragmiella* and *Heteroconium* from southern China. Cryptogam Mycol 33:127–135
- Ma J, Ma LG, Zhang YD, Xia JW, Zhang XG (2012d) New species and a new record of *Solicorynespora* from southern China. Mycotaxon 119:95–102
- Ma J, Zhang YD, Ma LG, Castañeda-Ruiz RF, Zhang XG (2012e) Three new species of *Sporidesmiella* from southern China. Mycoscience 53:187–193
- Ma LG, Ma J, Zhang YD, Castañeda-Ruiz RF, Zhang XG (2012f) New species and records of *Heteroconium* (anamorphic fungi) from southern China. Mycoscience 53:466–470
- Ma LG, Ma J, Zhang YD, Zhang XG (2012g) A new *Corynesporella* species and two first records from China. Mycotaxon 119:83–88
- Ma LG, Ma J, Zhang YD, Zhang XG (2012h) *Spadicoides camelliae* and *Diplococcum livistona*, two new hyphomycetes on dead branches from Fujian Province, China. Mycoscience 53:25–30
- Ma HX, Vasilyeva L, Li Y (2013) *Xylaria* in southern China—8. *X. bannaensis* sp. nov. and *X. brunneovinosa* new to the mainland. Mycotaxon 125:251–256
- Ma LG, Xia JW, Ma YR, Castañeda-Ruiz RF, Zhang XG (2014a) Two new species of *Spadicoides* and *Gangliostilbe* from southern China. Mycol Prog 13:547–552
- Ma J, Zhang XG, Castañeda-Ruiz RF (2014b) New species of *Acrodictys* and *Repetophragma* from dead branches in China. Mycotaxon 127:129–134
- Ma J, Ji-Wen X, Rafael F, Castañeda-Ruiz Xiu-Guo Z (2014) *Antinakataea setulosa* gen. et sp. nov. and *Uberispora formosa* sp. nov. from southern China. Mycologia 13:753–758
- Ma YR, Xia JW, Zhang XG, Castañeda-Ruiz RF (2015) New species of *Phaeomonilia* and *Mirandina* from dead branches in China. Mycotaxon 130:775–781
- Ma J, Xia JW, Zhang XG, Luo YQ, Castañeda-Ruiz RF (2016a) *Solicorynespora* species associated with dead branches of subtropical forests in southern China. Cryptogam Mycol 37:37–44
- Ma J, Zhang XG, Castañeda-Ruiz RF (2016b) *Podosporiopsis*, a new genus of synnematous hyphomycetes from China. Mycotaxon 131:773–780
- Ma YR, Xia JW, Gao JM, Li Z, Zhang XG (2016c) *Anacaccumisporium*, a new genus based on morphology and molecular analyses from Hainan, China. Cryptogam Mycol 37:45–59
- Ma YR, Xia JW, Gao JM, Li Z, Zhang XG (2016d) *Dictyoceratospora*, gen. nov., with the description of two new species collected from Hainan, China. Sydowia 68:57–61

- Machado AR, Pinho DB, Silva M, Pereira OL (2012) First report of leaf spot disease caused by *Cercosporaella pfaffiae* on Brazilian ginseng (*Pfaffia glomerata*) in Brazil. *Plant Dis* 96:1702
- Machouart M, Samerpitak K, de Hoog GS, Gueidan C (2014) A multigene phylogeny reveals that *Ochroconis* belongs to the family *Sympoventuriaceae* (*Venturiales, Dothideomycetes*). *Fungal Divers* 65:77–88
- Macía-Vicente JG, Glynnou K, Piepenbring M (2016) A new species of *Exophiala* associated with roots. *Mycol Prog* 15:18
- Madrid H, Cano J, Gené J, Bonifaz A et al (2009) *Sporothrix globosa*, a pathogenic fungus with widespread geographical distribution. *Rev Iberoam Micol* 26:218–222
- Madrid H, Cano JF, Stchigel A, Gené J, Guarro J (2010a) *Ramophialophora humicola* and *Fibulochlamys chilensis*, two new microfungi from soil. *Mycologia* 102:605–612
- Madrid H, Gené J, Cano J, Silvera C et al (2010b) *Sporothrix brunneoviolacea* and *S. dimorphospora*, two new members of the *Ophiostoma stenoceras*-*Sporothrix schenckii* complex. *Mycologia* 102:1193–1203
- Madrid H, Cano J, Gené J, Guarro J (2011a) Two new species of *Cladorrhinum*. *Mycologia* 103:795–805
- Madrid H, Gené J, Cano J, Guarro J (2011b) A new species of *Leptodiscella* from Spanish soil. *Mycol Prog* 11:535–541
- Madrid H, Hernández-Restrepo M, Gené J, Cano J et al (2016a) New and interesting chaetothyrialean fungi from Spain. *Mycol Prog* 15:1179–1201
- Madrid H, Torres D, Silva V (2016b) A new species of *Harmoniella* from central Chile. *Mycotaxon* 131:535–539
- Magain N, Séruisiaux E (2012) A further new species in the lichen genus *Arctomia*: *A. borbonica* from Réunion (Mascarene archipelago). *MycoKeys* 4:9–21
- Magain N, Séruisiaux E (2014) Do photobiont switch and cephalodia emancipation act as evolutionary drivers in the lichen symbiosis? A case study in the *Pannariaceae* (*Peltigerales*). *PLoS ONE* 9:e89876
- Magain N, Séruisiaux E (2015) Dismantling the treasured flagship lichen *Sticta fuliginosa* (*Peltigerales*) into four species in Western Europe. *Mycol Prog* 14:97
- Magain N, Miadlikowska J, Goffinet B, Séruisiaux E, Lutzoni F (2016) Macroevolution of specificity in cyanolichens of the genus *Peltigera* section polydactylon (*Lecanoromycetes, Ascomycota*). *Syst Biol.* doi:10.1093/sysbio/syw065
- Magalhães DMA, Luz EDMN, Magalhães AF, Dos Santos MVO et al (2014) Anamorphic fungi of the Atlantic forest of southern Bahia: new records and *Dactylaria pseudomanifesta* sp. nov. *Mycotaxon* 128:185–194
- Magyar D, Shoemaker RA, Bobovs J, Crous PW, Groenewald JZ (2011) *Pyrigemmula*, a novel hyphomycete genus on grapevine and tree bark. *Mycol Prog* 10:307–314
- Mahajan VK, Sharma V, Prabha N, Thakur K et al (2014) A rare case of subcutaneous phaeohyphomycosis caused by a *rhytidhysteron* species: a clinico-therapeutic experience. *Int J Derm* 53:1485–1489
- Maharachchikumbura SSN, Guo LD, Chukeatirote E, Bahkali AH et al (2011) *Pestalotiopsis*—morphology, phylogeny, biochemistry and diversity. *Fungal Divers* 50:167–187
- Maharachchikumbura SSN, Guo LD, Cai L, Chukeatirote E et al (2012) A multi-locus backbone tree for *Pestalotiopsis*, with a polyphasic characterization of 14 new species. *Fungal Divers* 56:95–129
- Maharachchikumbura SSN, Guo LD, Chukeatirote E, McKenzie EH et al (2013) A destructive new disease of *Syzygium samarangense* in Thailand caused by the new species *Pestalotiopsis samarangensis*. *Trop Plant Pathol* 38:227–235
- Maharachchikumbura SSN, Guo LD, Chukeatirote E, Hyde KD (2014a) Improving the backbone tree for the genus *Pestalotiopsis*; addition of *P. steyaertii* and *P. magna* sp. nov. *Mycol Prog* 13:617–624
- Maharachchikumbura SSN, Hyde KD, Groenewald JZ, Xu J et al (2014b) *Pestalotiopsis* revisited. *Stud Mycol* 79:121–186
- Maharachchikumbura SSN, Camporesi E, Liu ZY, Hyde KD (2015a) *Seiridium venetum* redescribed, and *S. camelliae*, a new species from *Camellia reticulata* in China. *Mycol Prog* 14:85–93
- Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EH et al (2015b) Towards a natural classification and backbone tree for *Sordariomycetes*. *Fungal Divers* 72:199–301
- Maharachchikumbura SSN, Guo LD, Liu ZY, Hyde KD (2016a) *Pseudopestalotiopsis ignota* and *Ps. camelliae* spp. nov. associated with grey blight disease of tea in China. *Mycol Prog* 15:22
- Maharachchikumbura SSN, Hyde KD, Jones EBG, McKenzie EH et al (2016b) Families of *Sordariomycetes*. *Fungal Divers* 79:1–317
- Maharachchikumbura SSN, Larignon P, Hyde KD, Al-Sadi AM, Liu YZ (2016c) Characterization of *Neopestalotiopsis*, *Pestalotiopsis* and *Truncatella* species associated with grapevine trunk diseases in France. *Phytopathol Mediterr* 55:380–390
- Mahmoud RS, Narisawa K (2013) A new fungal endophyte, *Scolecobasidium humicola*, promotes tomato growth under organic nitrogen conditions. *PLoS ONE* 8:e78746
- Maier S, Schmidt TS, Zheng L, Peer T, Wagner V, Grube M (2014) Analyses of dryland biological soil crusts highlight lichens as an important regulator of microbial communities. *Biodiversity and conservation* 23:1735–1755
- Makhija U, Patwardhan PG (1995) Nomenclatural notes on some species of *Arthothelium* (lichenized Ascomycotina). *Trop Bryol* 10:205–213
- Malíček J, Bouda F, Liška J, Palice Z, Peksa O (2015a) Contribution to the lichen biota of the Romanian Carpathians. *Herzogia* 28:713–735
- Malíček J, Palice Z, Vondrák J (2015b) New lichen records and rediscoveries from the Czech Republic and Slovakia. *Herzogia* 27:257–284
- Magain A, Roychowdhury R, Tah J (2013) *Alternaria* pathogenicity and its strategic controls. *Res J Biol* 1:1–9
- Man B, Wang H, Xiang X, Wang R et al (2015) Phylogenetic diversity of culturable fungi in the Heshang Cave, central China. *Front Microbiol* 6:1158
- Manamgoda DS, Cai L, McKenzie EH, Crous PW et al (2012) A phylogenetic and taxonomic re-evaluation of the *Bipolaris-Cochliobolus-Curvularia* complex. *Fungal Divers* 56:131–144
- Manamgoda DS, Udayanga D, Cai L, Chukeatirote E et al (2013) Endophytic *Colletotrichum* from tropical grasses with a new species *C. endophytica*. *Fungal Divers* 61(1):107–115
- Manamgoda DS, Rossman AY, Castlebury LA, Crous PW et al (2014) The genus *Bipolaris*. *Stud Mycol* 79:221–288
- Manamgoda DS, Rossman AY, Castlebury LA, Chukeatirote E et al (2015) A taxonomic and phylogenetic re-appraisal of the genus *Curvularia* (*Pleosporaceae*): human and plant pathogens. *Phytotaxa* 212:175–198
- Mangold A, Elix JA, Lumbsch HT (2009) *Thelotremaeaceae*. *Flora Aust* 57:653–659
- Mangold A, Martin MP, Kalb K, Lücking R, Lumbsch HT (2008a) Molecular data shows that *Topeliopsis* (*Ascomycota, Thelotremaeaceae*) is polyphyletic. *Lichenologist* 40:39–46
- Mangold A, Martin MP, Luecking R, Lumbsch HT (2008b) Molecular phylogeny suggests synonymy of *Thelotremaeaceae* within *Graphidaceae* (*Ascomycota: Ostropales*). *Taxon* 57:476–486
- Manoharachary C, Bagyanarayana G, Rao NK, Kunwar IK (2009) *Chasakopama*, a new dematiaceous hyphomycetous genus from India. *Mycotaxon* 110:459–463

- Manoharachary C, Kunwar IK, Kumar GS, Reddy BS (2010) Two new species of *Actinocladium* Ehrenb. from India. Indian Phytopathol 63:83–84
- Manojlović NT, Vasiljević P, Juskovic M, Najman S et al (2010a) Screening of *Thamnolia vermicularis* var. *subuliformis* for antimicrobial, antioxidant and cytotoxic activities. Planta Med 76:442
- Manojlović NT, Vasiljević P, Jusković M, Janković SNS (2010b) HPLC analysis and cytotoxic potential of extracts from the lichen, *Thamnolia vermicularis* var. *subuliformis*. J Med Plants Res 4:817–823
- Manojlović NT, Vasiljević PJ, Maskovic PZ, Juskovic M, Bogdanovic-Dusanovic G (2011) Chemical composition, antioxidant, and antimicrobial activities of lichen *Umbilicaria cylindrica* (L.) Delise (*Umbilicariaceae*). Evid Complement Altern Med. doi:10.1155/2012/452431
- Mansour AM (2010) Contribution to knowledge of some soil fungi in eastern region in Libya. J Prod Dev 15:395–404
- Mapook A, Boonmee S, Ariyawansa HA, Tibpromma S et al (2016a) Taxonomic and phylogenetic placement of *Nodulosphaeria*. Mycol Prog 15:1–5
- Mapook A, Boonmee S, Liu JK, Jones EBG et al (2016b) Taxonomic and phylogenetic placement of *Phaeodimeriella* (*Pseudoperisporiaceae*, *Pleosporales*). Cryptogam Mycol 37:157–176
- Mapook A, Hyde KD, Dai DQ, Li J et al (2016c) *Muyocopronales*, ord. nov., (*Dothideomycetes*, *Ascomycota*) and a reappraisal of *Muyocopron* species from northern Thailand. Phytotaxa 265:225–237
- Mapook A, Hyde KD, Hongsanan S, Phukhamsakda C et al (2016d) Palawaniaceae fam. nov., a new family (*Dothideomycetes*, *Ascomycota*) to accommodate *Palawania* species and their evolutionary time estimates. Mycosphere 7:1732–1745
- Mapperson RR, Kotiw M, Davis TA, Dearnaley JDW (2014) The diversity and antimicrobial activity of *Preussia* sp. Endophytes isolated from Australian dry rainforests. Curr Microbiol 68:30–37
- Mardones M, Trampe-Jaschik T, Oster S, Elliott M et al (2017) Phylogeny of the order Phyllachorales (*Ascomycota*, *Sordariomycetes*): among and within order relationships based on five molecular loci. Persoonia 39:74–90
- Marincowitz S, Crous PW, Groenewald JZ, Wingfield MJ (2008) Microfungi occurring on the *Proteaceae* in the fynbos. CBS Diversity Ser. 7: [iii], Utrecht: Centraalbureau voor Schimmelcultures
- Marincowitz S, Gryzenhout M, Wingfield MJ (2010) New and rare coelomycetes with appendage-bearing conidia from Pondoland, South Africa. Mycotaxon 111:309–322
- Marincowitz S, Duong TA, de Beer ZW, Wingfield MJ (2015) *Cornuvesica*: a little known mycophilic genus with a unique biology and unexpected new species. Fungal Biol 119:615–630
- Marin-Felix Y, Cano-Lira JF, Guarro J, Stchigel AM (2014) *Leiothecium cristatum* sp. nov. and *Aspergillus posadasensis* sp. nov., two species of *Eurotiales* from rainforest soils in South America. Int J Syst Evol Micro 64:2871–2877
- Marin-Felix Y, Stchigel AM, Miller AN et al (2015) A re-evaluation of the genus *Myceliophthora* (Sordariales, Ascomycota): its segregation into four genera and description of *Corynascus fumimontanus* sp. nov. Mycologia 107:619–632
- Marin-Felix Y, Groenewald JZ, Cai L, Chen Q et al (2017) Genera of phytopathogenic fungi: GOPHY 1. Stud Mycol 86:99–216
- Mark K, Saag L, Saag A, Thell A, Randlane T (2012) Testing morphology-based delimitation of *Vulpicida juniperinus* and *V. tubulosus* (*Parmeliaceae*) using three molecular markers. Lichenologist 44:757–772
- Mark K, Cornejo C, Keller C, Fluck D, Scheidegger C (2016) Barcoding lichen-forming fungi using 454 pyrosequencing is challenged by artifactual and biological sequence variation. Genome 59:685–704
- Marques J, Schultz M, Graciela PB (2013) A *Peltula* Nyl. diversity hotspot in north-east Portugal, with one species new to science and three species new to mainland Europe. Lichenologist 45:483–496
- Martellos S, Attorre F, Farcomeni A, Francesconi F et al (2014) Species distribution models backing taxa delimitation: the case of the lichen *Squamaria cartilaginea* in Italy. Flora Morphol Distrib Funct Ecol Plants 209:698–703
- Martinez ADS, Franzener G, Stangerlin JR (2010) Damages caused by *Bipolaris maydis* in *Panicum maximum* cv. Tanzânia. Semin Ciências Agrár Lond 31:863–870
- Martinović T, Koukol O, Hirose D (2016) Distinct phylogeographic structure recognized within *Desmazierella acicola*. Mycologia 108:20–30
- Marvanová L, Laichmanová M (2014) *Campylospora leptosoma* sp. nov. and characteristics of *Campylospora* spp., based on morphology and on ITS sequences. Mycosphere 5:245–261
- Masiulionis VE, Cabello M, Seifert KA, Rodrigues A, Pagnocca FC (2015) *Escovopsis trichodermoides* sp. nov., isolated from a nest of the lower attine ant *Mycocepurus goeldii*. Antonie van Leeuwenhoek 107:731–740
- Masson D, Divakar PK, Serusiaux E (2015) *Hypotrachyna penduliloba* and *Remototrachyna pandani*, two new species in the hyperdiverse lichen family Parmeliaceae from Reunion in the Mascarene archipelago. Mycol Prog 14:22
- Masumoto S, Tojo M, Uchida M, Imura S (2014) *Rhytisma polaris*: morphological and molecular characterization of a new species from Spitsbergen Island, Norway. Mycol Prog 13:181–188
- Masumoto S, Tojo M, Uchida M, Imura S (2015) Morphological and molecular characterization of *Rhytisma filamentosum* sp. nov. from Nagano Prefecture, Japan. Mycol Prog 14:1–6
- Masuya H, Yamaoka Y, Wingfield MJ (2009) Ophiostomatoid fungi and their associations with bark beetles in Japan. CBS Biodivers Ser 96:275–293
- Mateus N, Aguirre J, Luecking R (2012) Contributions to the foliicolous lichen biota of Chocó (Colombia). Caldasia 34:25–32
- Mathur N, Singh O, Maheswari CU (2009) *Polyrostrata*, a new genus of Sphaeropsidaceae from India. Indian Phytopathol 62:233–236
- Matočec N, Kušan I, Ozimec R (2014) The genus *Polycephalomyces* (*Hypocreales*) in the frame of monitoring *Veterica* cave (Croatia) with a new segregate genus *Perennicordyceps*. Ascomycete.org 6:125–133
- Matos T, Cerar T, Praprotnik M, Krivec U, Pirš M (2015) First recovery of *Rasamsonia argillacea* species complex isolated in adolescent patient with cystic fibrosis in Slovenia—case report and review of literature. Mycoses 58:506–510
- Matsuda Y, Kimura K, Ito SI (2010) Genetic characterization of *Raffaelea quercivora* isolates collected from areas of oak wilt in Japan. Mycoscience 51:310–316
- Matsuwaki Y, Wada K, White T, Moriyama H, Kita H (2012) *Alternaria* fungus induces the production of GM-CSF, interleukin-6 and interleukin-8 and calcium signaling in human airway epithelium through protease-activated receptor 2. Int Arch Allergy Immunol 158(Suppl. 1):19–29
- Matthews JA, Trenbirth HE (2011) Growth rate of a very large crustose lichen (*Rhizocarpon subgenus*) and its implications for lichenometry. Geogr Ann Ser A Phys Geogr 93:27–39
- Matusinsky P, Frei P, Mikolasova R, Svacinova I et al (2010) Species-specific detection of *Bipolaris sorokiniana* from wheat and barley tissues. Crop Prot 29:1325–1330
- Matwiejuk A, Bohdan A (2011) New Records of *Plectocarpon lichenum* from Bialowieza Forest (Poland). Herzogia 24:381–383
- Mayers CG, McNew DL, Harrington TC, Roeper RA et al (2015) Three genera in the *Ceratostidaceae* are the respective

- symbionts of three independent lineages of ambrosia beetles with large, complex mycangia. *Fungal Biol* 119:1075–1092
- Mayfield AE III, Smith JA, Hughes M, Dreden TJ (2008) First report of laurel wilt disease caused by a *Raffaelea* sp. on avocado in Florida. *Plant Dis* 92:976–1976
- Mayrhofer H, Malíček J, Rohrer A, Svoboda D, Biložit PO (2012) New or otherwise interesting lichenized and lichenicolous fungi from Macedonia. *Fritschiana* (Graz) 71:1–35
- Mayrhofer H, Drescher A, Stešević D, Biložit PO (2013) Lichenized fungi of a chestnut grove in Livari (Rumija, Montenegro). *Acta Bot Croat* 72:337–346
- Mazzaferrro L, Piñuel L, Minig M, Breccia JD (2010) Extracellular monoenzyme deglycosylation system of 7-O-linked flavonoid β-rutinosides and its disaccharide transglycosylation activity from *Stilbella fimetaria*. *Arch Microbiol* 192:383–393
- Mbenoun M, De Beer ZW, Wingfield MJ, Wingfield BD, Roux J (2014) Reconsidering species boundaries in the *Ceratocystis paradoxica* complex, including a new species from oil palm and cacao in Cameroon. *Mycologia* 106:757–784
- Mbenoun M, Wingfield MJ, Begoude Boyogueno AD et al (2016) Diversity and pathogenicity of the *Ceratocystidaceae* associated with cacao agroforests in Cameroon. *Plant Pathol* 65:64–78
- McCarthy PM (2010) New and interesting foliicolous lichens from Australia. *Australas Lichenol* 66:4–15
- McCarthy PM (2014) Additional lichen records from Australia 77. *Verrucariaceae*. *Australas Lichenol* 75:3–5
- McCarthy PM, Kantvilas G (2013a) *Psoroglaena halmaturina* sp. nov. (lichenised Ascomycota, Verrucariaceae) from Kangaroo Island, South Australia. *J Adel Bot Gard* 1:1–4
- McCarthy PM, Kantvilas G (2013b) Two new species of *Sarcogyne* (lichenised Ascomycota: Acarosporaceae) from central and southern Australia. *J Adel Bot Gard* 26:15–21
- McCarthy PM, Stajsic V (2013) *Phylloblastia blechnicola* (Ascomycota, Verrucariaceae), a new leaf-inhabiting lichen from southern Victoria, Australia. *Muelleria* 31:49–52
- McCarthy PM, Kantvilas G (2014) *Placynthium australiense* sp. nov. (lichenised Ascomycota, Placynthiaceae) from South Australia. *J Adel Bot Gard* 27:1–5
- McCarthy P, Elix J (2016) Five new lichen species (Ascomycota) from south-eastern Australia. *Telopea* 19:137–151
- McCune B, Altermann S (2009) *Letharia gracilis* (Parmeliaceae), a new species from California and Oregon. *Bryologist* 112:375–378
- McCune B, Stone J (2009) *Leightoniomyces phillipsii*, a synnematosus soil-dwelling hyphomycete new to North America. *N Am Fungi* 4:1–4
- McCune B, Curtis MJ (2012) *Umbilicaria semitensis* (lichenized fungi: Umbilicariaceae) resurrected. *Bryologist* 115:255–264
- McCune B, Rosentreter R (2014) New lichen records from Oregon to Alaska in North America. *Evensia* 31:1–7
- McCune B, Wang LS (2014) The lichen genus Hypogymnia in southwest China. *Mycosphere* 5:27–76
- McCune B, Di Meglio J, Curtis MJ (2014) An unusual ascospore shape and a new species, *Umbilicaria nodulospora* (Umbilicariaceae), from California and Oregon. *Bryologist* 117:170–178
- McDonald TR, Gaya E, Lutzoni F (2013) Twenty-five cultures of lichenizing fungi available for experimental studies on symbiotic systems. *Symbiosis* 59:165–171
- McGrann GR, Stavrinides A, Russell J, Corbitt MM et al (2014) A trade off between mlo resistance to powdery mildew and increased susceptibility of barley to a newly important disease, *Ramularia* leaf spot. *J Exp Bot* 65:1025–1037
- McKenzie EHC (2008) Two new dictyosporous hyphomycetes on Pandanaceae. *Mycotaxon* 104:23–28
- McKenzie EHC (2009) A new species of *Lylea* (hyphomycetes) on *Rhopalostylis* (Arecaceae) in New Zealand. *Mycotaxon* 109:39–42
- McKenzie EHC (2010) Three new phragmosporous hyphomycetes on *Ripogonum* from an ‘ecological island’ in New Zealand. *Mycotaxon* 111:183–196
- McKenzie EHC, Park D, Bellgard SE, Johnston PR (2010) A new species of *Pyricularia* (hyphomycetes) on *Cortaderia* (Poaceae) in New Zealand. *Mycosphere* 1:223–228
- McMullin RT (2015) The Lichens of Prince Edward Island, Canada: a second checklist, with species ranked for conservation status. *Rhodora* 117:454–484
- McMullin RT, Arsenault A (2016) The Calicioids of Newfoundland, Canada. *Opusc Philolichenum* 15:92–104
- McMullin RT, Thompson ID, Newmaster SG (2013) Lichen conservation in heavily managed boreal forests. *Conserv Biol* 27:1020–1030
- McMullin RT, Maloles JR, Newmaster SG (2015) New and interesting lichens from Ontario, Canada-II. *Opusc Philolichenum* 14:93–108
- McMullin RT, Lendemer JC, Braid HE, Newmaster SG (2016) Molecular insights into the lichen genus *Alectoria* (Parmeliaceae) in North America. *Botany* 94:165–175
- McNeill J, Turland NJ, Monro A, Lepschi BJ (2011) XVIII International Botanical Congress: preliminary mail vote and report of Congress action on nomen clature proposals. *Taxon* 60:1507–1520
- Medeiros ID, Fryday AM, Rajakaruna N (2014) Additional lichen records and mineralogical data from metal-contaminated sites in Maine. *Rhodora* 116:323–347
- Medeiros ID, Kraichak E, Lücking R, Mangold A, Lumbsch HT (2017) Assembling a taxonomic monograph of tribe *Wirthiotremataceae* (lichenized Ascomycota: Ostropales: Graphidaceae). *Fieldiana Life Earth Sci* 9:1–31
- Medjedović A, Frank J, Schroers HJ, Oertel B et al (2014) *Peltaster cerophilus* is a new species of the apple sooty blotch complex from Europe. *Mycologia* 106:525–536
- Meena N, Mishra VK, Baranwal DK, Singh AK et al (2014) Genetic evaluation of spring wheat (*Triticum aestivum* L.) recombinant inbred lines for Spot Blotch (*Bipolaris sorokiniana*) resistance and yield components under natural conditions for south Asia. *J Agric Sci Technol* 16:1429–1440
- Meena PD, Awasthi RP, Chattopadhyay C, Kolte SJ, Kumar A (2016) *Alternaria* blight: a chronic disease in rapeseed–mustard. *Journal of Oilseed Brassica* 1:1–11
- Mehrabi M, Hemmati R, Vasilyeva LN, Trouillas FP (2015) A new species and a new record of *Diatrypaceae* from Iran. *Mycosphere* 6:60–68
- Meirelles LA, Montoya QV, Solomon SE, Rodrigues A (2015) New light on the systematics of fungi associated with attine ant gardens and the description of *Escovopsis krieselii* sp. nov. *PLoS ONE* 10:e0112067
- Mejia LC, Castlebury LA, Rossman AY, Sogonov MV, White JF (2008) Phylogenetic placement and taxonomic review of the genus *Cryptospora* and its synonyms *Ophiovalsa* and *Winterella* (Gnomoniaceae, Diaporthales). *Mycol Res* 112:23–35
- Mejia LC, Rossman AY, Castlebury LC, White JF (2011) New species, phylogeny, host-associations, and geographic distribution of genus *Cryptospora* (Gnomoniaceae, Diaporthales). *Mycologia* 103:379–399
- Mejia LC, Rossman AY, Castlebury LA, Yang ZL, White JF Jr (2012) *Occultocarpon*, a new monotypic genus of Gnomoniaceae on *Alnus nepalensis* from China. *Fungal Divers* 52:99–105

- Mel'nik VA (2011) Anamorphic fungi of Vietnam. I. Mikologiya i Fitopatologiya 45:323–331
- Mel'nik VA (2012) *Phaeoisaria vietnamensis* sp. nov. and *P. clematidis* (hyphomycetes) from Vietnam. Mycosphere 3:957–960
- Mel'nik VA, Braun U (2013) *Atractilina alinae* sp. nov. and *Neosporidesmium vietnamense* sp. nov.—two new synnematosus hyphomycetes from Vietnam. Mycobiota 3:1–9
- Mel'nik VA, Crous PW (2014) *Braunomyces dictyosporus* gen. sp. nov. from vietnam. IMA Fungus 5:1–5
- Mel'nik VA, Braun U, Alexandrova V (2013a) *Dactylaria mucoglobifera* sp. nov.—a new species from Vietnam. Schlechtendalia 25:49–52
- Mel'nik VA, Castañeda-Ruiz RF, Granados M (2013b) A new species of *Heliocephala* from Vietnam. Mycotaxon 123:281–284
- Mel'nik VA, Alexandrova AV, Braun U (2014) Two new species and new records of hyphomycetes from Vietnam. Mycosphere 5:591–600
- Mel'nik VA, Popov ES, Braun U (2016) *Neosporidesmium subramanianii* sp. nov. from Vietnam. Mycosphere 7:148–153
- Melechin AV (2010) *Stereocaulon leucophaeopsis* and *S. tornense* new to Russia from the Murmansk region. Gr Scr 22:63–64
- Melo RFR, Miller AN, Santiago ALCMA, Maia LC (2014) The genera *Ascobolus* and *Saccobolus* (Ascobolaceae, Pezizales) in Brazil. Mycosphere 5:790–804
- Melo RFR, Miller AN, Maia LC (2015) The genus *Podospora* (Lasiosphaeriaceae, Sordariales) in Brazil. Mycosphere 6:201–215
- Méndez-Mayboca F, Checa J, Esqueda M, Chacón S (2010) New records of loculoascomycetes from natural protected areas in Sonora, Mexico. Mycotaxon 111:19–30
- Mena-Portales J, Hernández-Restrepo M, Guarro J, Minter DW et al (2016) New species of *Penzigomyces*, *Sporidesmium* and *Stanjehughesia* from plant debris in Spain. Nova Hedwig 103:359–371
- Menezes A, Leite A, Otsuka A, Jesus L, Cáceres MES (2011) New records of crustose and microfoliaceous corticolous lichens in Caatinga vegetation of the semi-arid region in Alagoas. Acta Bot Bras 25:885–889
- Menezes AA, Lima EL, Xavier-Leite AB, Maia LC et al (2013a) New species of *Arthoniales* from NE Brazil. Lichenologist 45:611–617
- Menezes AA, Xavier-Leite AB, De Jesus KA, Aptroot A, Cáceres MES (2013b) Two new *Cryptohnia* species and a new *Syncessia* from Chapada do Araripe, Ceará, NE Brazil (Ascomycota: Arthoniales), with a key to *Cryptohnia*. Lichenologist 45:657–664
- Menkis A, Vasaitis R (2011) Fungi in roots of nursery grown *Pinus sylvestris*: ectomycorrhizal colonisation, genetic diversity and spatial distribution. Microb Ecol 61:52–63
- Menkis A, Urbina H, James TY, Rosling A (2014) *Archaeorhizomyces borealis* sp. nov. and a sequence-based classification of related soil fungal species. Fungal Biol 118:943–955
- Mercado-Díaz JA, Lücking R, Parnmen S (2014) Two new genera and twelve new species of *Graphidaceae* from Puerto Rico suggest a high degree of endemism. Phytotaxa 189:186–203
- Messuti MI, Codesal PL, Mangold A, Luecking R, Lumbsch HT (2010) New or interesting *Chapsa* and *Topeliopsis* species (Ascomycota: Ostropales) from Argentina. Lichenologist 42:191–195
- Messuti MI, Lumbsch HT, Luecking R, Altermann S (2011) One hundred new species of lichenized fungi: a signature of undiscovered global diversity. Phytotaxa 18:1–127
- Messuti MI, Passo A, Scervino JM, Vidal-Russell R (2016) The species pair *Pseudocyphellaria pilosella*–*piloselloides* (lichenized Ascomycota: Lobariaceae) is a single species. Lichenologist 48:141–146
- Methven AS, Zelski SE, Miller AN (2013) A molecular phylogenetic assessment of the genus *Gyromitra* in North America. Mycologia 105:1306–1314
- Meunier JD, Kirman S, Strasberg D, Grauby O, Dussouillez P (2014) Incipient weathering by *Stereocaulon vulcani* at Réunion volcanic island. Chem Geol 382:123–131
- Meshram V, Kapoor N, Chopra G, Saxena S (2017) *Muscodor camphora*, a new record from *Cinnamomum camphora*. Mycosphere 8:568–582
- Miądlikowska J, Schoch CL, Kageyama SA, Molnar K et al (2011) *Hypogymnia* phylogeny, including *Cavernularia*, reveals biogeographic structure. Bryologist 114:392–400
- Miądlikowska J, Kauff F, Högnabba F, Oliver JC et al (2014a) A multigene phylogenetic synthesis for the class *Lecanoromycetes* (Ascomycota): 1307 fungi representing 1139 infrageneric taxa, 317 genera and 66 families. Mol Phylogenet Evol 79:132–168
- Miądlikowska J, Richardson D, Magain N, Ball B et al (2014b) Phylogenetic placement, species delimitation, and cyanobiont identity of endangered aquatic *Peltigera* species (lichen-forming Ascomycota, Lecanoromycetes). Am J Bot 101:1141–1156
- Mies BA (2015) New or interesting lichens from the Limousin region (Western Massif Central, France). Herzogia 28:473–483
- Miller JE, McCune B, Kofranek D, Villella J et al (2011a) Lichens from the South Slough and Horsfall Dunes on the southern Oregon coast. Evansia 28:92–99
- Miller JE, Rossman A, Rosentreter R, Ponzetti J (2011b) Lichen ecology and diversity of a sagebrush steppe in Oregon: 1977 to the present. N Am Fungi 6:1–14
- Miller AN, Huhndorf SM, Fournier J (2014) Phylogenetic relationships of five uncommon species of *Lasiosphaeria* and three new species in the *Helminthosphaeriaceae* (Sordariomycetes). Mycologia 106:505–524
- Millot M, Delebassée S, Liagre B, Vignaud L et al (2014) Screening of lichen extracts on HT-29 human colon–cancer cells. Plant Med 80:PIN5
- Minckler D, Small KW, Walsh TJ (2014) Clinical and pathologic features of *Bipolaris endophthalmitis* after intravitreal triamcinolone. JAMA Ophthalmol 132:630–632
- Minnis AM, Lindner DL (2013) Phylogenetic evaluation of *Geomyces* and allies reveals no close relatives of *Pseudogymnoascus destructans*, comb. nov., in bat hibernacula of eastern North America. Fungal Biol 117:638–649
- Minnis AM, Rossman AY, Farr DF, Olsen RT (2010) Fungal planet 51. *Sphaerographium nysicola* sp. nov. Persoonia 25:122–123
- Minnis AM, Kennedy AH, Grenier DB, Rehner SA, Bischoff JF (2011) *Asperisporium* and *Pantospora* (Mycosphaerellaceae): epitypifications and phylogenetic placement. Persoonia 27:1–8
- Minnis AM, Kennedy AH, Grenier DB, Palm ME, Rossman AY (2012) Phylogeny and taxonomic revision of the Planistromellaceae including its coelomycetous anamorphs: contributions towards a monograph of the genus *Kellermania*. Persoonia 29:11–28
- Minter DW (2009) Cyberlindnera, a replacement name for *Lindnera* Kurtzman et al., nom. illegit. Mycotaxon 110:473–476
- Minter DW, Aguirre-Hudson B, Cannon PF (2015) *Naetrocybme nitescens*. [Descriptions of Fungi and Bacteria]. IMI Descriptions of Fungi and Bacteria No. 206 pp. Sheet 2059 ref. 10
- Mirabolafathy M, Ju YM, Hsieh HM, Rogers JD (2013) *Obolarina persica* sp. nov., associated with dying *Quercus* in Iran. Mycoscience 54:315–320
- Miralles I, Edwards HG, Domingo F, Jorge-Villar SE (2015) Lichens around the world: a comprehensive study of lichen survival

- biostrategies detected by Raman spectroscopy. *Anal Methods* 7:6856–6868
- Miralles-Mellado I, Cantón Y, Solé-Benet A (2011) Two-dimensional porosity of crusted silty soils: indicators of soil quality in semiarid rangelands? *Soil Sci Soc Am J* 75:1330–1342
- Mishra AK, Mishra A, Kehri HK, Sharma B, Pandey AK (2009) Inhibitory activity of Indian spice plant *Cinnamomum zeylanicum* extracts against *Alternaria solani* and *Curvularia lunata*, the pathogenic dematiaceous moulds. *Ann Clin Microbiol Antimicrob* 8:1
- Mishra K, Das S, Goyal S, Gupta C et al (2014a) Subcutaneous mycoses caused by *Rhytidhysterion* species in an immunocompetent patient. *Med Mycol Case Rep* 5:32–34
- Mishra S, Singh BR, Singh A, Keswani C et al (2014b) Biofabricated silver nanoparticles act as a strong fungicide against *Bipolaris sorokiniana* causing spot blotch disease in wheat. *PLoS ONE* 9:e97881
- Mitrovic T, Stamenkovic S, Cvetkovic V, Radulovic N et al (2014) *Platismatia glauca* and *Pseudevernia furfuracea* lichens as sources of antioxidant, antimicrobial and antibiofilm agents. *Excli J* 13:938
- Mittermeier VK, Schmitt N, Volk LP, Suárez JP et al (2015) Metabolic profiling of alpine and ecuadorian lichens. *Molecules* 20:18047–18065
- Miyamoto Y, Masunaka A, Tsuge T, Yamamoto M et al (2008) Functional analysis of a multicopy host-selective ACT-toxin biosynthesis gene in the tangerine pathotype of *Alternaria alternata* using RNA silencing. *Mol Plant Microbe Interact* 21:1591–1599
- Miyamoto Y, Ishii Y, Honda A, Masunaka A et al (2009) Function of genes encoding acyl-CoA synthetase and enoyl-CoA hydratase for host-selective ACT-toxin biosynthesis in the tangerine pathotype of *Alternaria alternata*. *Phytopath* 99:369–377
- Mizuno H, Kawahigashi H, Kawahara Y, Kanamori H et al (2012) Global transcriptome analysis reveals distinct expression among duplicated genes during *sorghum*–*Bipolaris sorghicola* interaction. *BMC Plant Biol* 12:1
- Mohabe S, Nayaka S, Reddy AM (2015) *Stigmatochroma microspora* (Physciaceae), a new species from India. *J New Biol Rep* 4:128
- Mohammadi P (2011) Spectrofluorometry as a tool to evaluate biocide efficiency on deteriorating rock fungi. *Iran J Biol* 24:818–825
- Molnár K, Lőkös L, Schrett-Major Á, Farkas E (2012) Molecular genetic analysis of *Xanthoparmelia pulvinaris* (Ascomycota, Lecanorales, Parmeliaceae). *Acta Bot Hung* 54:125–130
- Monaghan MT, Wild R, Elliot M, Fujisawa T et al (2009) Accelerated species inventory on Madagascar using coalescent-based models of species delineation. *Syst Biol* 58:298–311
- Moncada B, Lücking R (2012) Ten new species of *Sticta* and counting: Colombia as a hot spot for unrecognized diversification in a conspicuous macrolichen genus. *Phytotaxa* 74:1–29
- Moncada B, Lücking R, Betancourt-Macuase L (2013a) Phylogeny of the *Lobariaceae* (lichenized Ascomycota: Peltigerales), with a reappraisal of the genus *Lobariella*. *Lichenologist* 45:203–263
- Moncada B, Luecking R, Coca LF (2013b) Six new apotheciate species of *Sticta* (lichenized Ascomycota: Lobariaceae) from the Colombian Andes. *Lichenologist* 45:635–656
- Moncada B, Lücking R, Suárez A (2014a) Molecular phylogeny of the genus *Sticta* (lichenized Ascomycota: Lobariaceae) in Colombia. *Fungal Divers* 64:205–231
- Moncada B, Reidy B, Lücking R (2014b) A phylogenetic revision of Hawaiian *Pseudocyclphellaria* sensu lato (lichenized Ascomycota: Lobariaceae) reveals eight new species and a high degree of inferred endemism. *Bryologist* 117:119–160
- Mongkolsamrit S, Luangsa-ard JJ, Hywel-Jones NL (2011) *Samuelsia mundiveteris* sp. nov. from Thailand. *Mycologia* 103:921–927
- Mongkolsuk P, Meesim S, Poengsungnoen V, Kalb K (2012) The lichen family *Physciaceae* in Thailand—I. The genus *Pyxine*. *Phytotaxa* 59:32–54
- Mongkolsuk P, Meesim S, Poengsungnoen V, Buaruang K et al (2015) The lichen family *Physciaceae* in Thailand—II. Contributions to the genus *Heterodermia* sensu lato. *Phytotaxa* 235:1–66
- Moniri MH (2010) A Small Lichen Collection from Darkesh, NE Iran. *J Environ Sci Eng* 4:36
- Moniri MH, Sipman HJ (2011) Lichens from three mountain sites in Khorasan provinces, Iran, including four species new to Iran. *Cryptogam Mycol* 32:145–150
- Moniri MH, Sipman HJ, Schultz M (2014) New records of lichenized and lichenicolous fungi from Northeastern Iran. *Herzogia* 27:367–376
- Moniri MH, Soltani A, Kamyabi S (2009) Some lichens from Kashmar, NE Iran. *J Appl Nat Sci* 1:286–290
- Monkai J, Liu JK, Boonmee S, Chomnunti P et al (2013) *Planistromellaceae* (Botryosphaeriales). *Cryptogam Mycol* 34:45–77
- Monteiro JS, Gusmão LFP (2014) Two new species of *Ceratosporella* (anamorphic fungi) from Brazilian Amazon forest. *Nova Hedw* 98:481–490
- Monteiro JS, Carmo LT, Fiúza PO, Ottoni BM et al (2014a) New species of microfungi from Brazilian Amazon rainforests. *Mycotaxon* 127:81–87
- Monteiro JS, Castañeda-Ruiz RF, Gusmão LFP (2014b) *Helicodochium*, a new microfungus from submerged wood from Brazil. *Mycotaxon* 127:5–9
- Monteiro JS, Gusmão LF, Castañeda-Ruiz RF (2014c) Two new microfungi from Brazilian Amazon Forest: *Atrogeniculata submersa* and *Nigrolentilocus amazonicus*. *Mycotaxon* 127:39–45
- Monteiro JS, Gusmão LF, Castañeda-Ruiz RF (2015a) A new species of *Matsushimaella* from submerged leaves in the Brazilian Amazon Forest. *Mycotaxon* 130:311–314
- Monteiro JS, Gusmão LF, Castañeda-Ruiz RF (2015b) *Brachycephala exotica*, a new hyphomycete from Brazil. *Mycotaxon* 130:489–493
- Monteiro JS, Conceição LB, Marques MF, Gusmão LF, Castañeda-Ruiz RF (2016a) *Dictyoaquaphila appendiculata* gen. & sp. nov. from submerged wood from Brazil. *Mycotaxon* 131:177–183
- Monteiro JS, Gusmão LF, Castañeda-Ruiz RF (2016b) *Anacoronospora diversiseptata* gen. & sp. nov. from Brazil. *Mycotaxon* 131:185–192
- Moon KH, Aptroot A (2009) Pyrenocarpous lichens in Korea. *Biblthca. Lichenol* 99:297–314
- Moon KH, Nakanishi S, Kashiwadani H (2008) Notes on species of *Graphidaceae* (Ascomycota) from eastern Asia with three new species. *Mem Natl Sci Musuem Tokyo* 45:85–91
- Moreno G, Alvarado P, Manjón JL (2012a) Phylogenetic affiliation of *Choriomyces magnusii* and *C. venosus* (*Tuberaceae* Ascomycota) from Spain. *Mycol Prog* 11:463–471
- Moreno G, Lizárraga M, Esqueda M, Galán R, Alvarado P (2012b) New records of little-known species of *Carbomyces* (*Carbomycetaceae*, Ascomycota). *Mycotaxon* 120:89–98
- Moreno-Pérez P, Gamboa-Angulo M, Heredia G, Canto-Canché B et al (2014) Antagonistic properties of micromycetes isolated from sinkholes of the Yucatán Peninsula against fungal phytopathogens. *Rev Mex Mic* 40:27–36
- Moreno-Rico O, Groenewald JZ, Crous PW (2014) Follicolous fungi from *Arctostaphylos pungens* in Mexico. *IMA Fungus* 5:7–15

- Morgenstern K, Döring M, Krabel D (2013) Rhabdocline needle cast-investigations on various Douglas fir tissue types. *Eur J Plant Pathol* 137:495–504
- Morin L, Shivas RG, Piper MC, Tan YP (2010) *Austropleospora osteospermi* gen. et sp. nov. and its host specificity and distribution on *Chrysanthemoides monilifera* ssp. *rotundata* in Australia. *Fungal Divers* 40:65–74
- Morita H, Tsuchiya T, Kishibe K, Noya S et al (2009) Antimitotic activity of lobaric acid and a new benzofuran, sakisacaulon A from *Stereocaulon sasakii*. *Bioorg Med Chem Lett* 19:3679–3681
- Moro LB, Delgado G, Schoenlein-Crusius IH (2015) *Polylobatispora setulosa*, a new freshwater hyphomycete from Ilhabela, São Paulo state, Brazil. *Mycosphere* 6:13–18
- Mostafanezhad H, Sahebani N, Nourinejad Zarghani S (2014) Control of root-knot nematode (*Meloidogyne javanica*) with combination of *Arthrobotrys oligospora* and salicylic acid and study of some plant defense responses. *Biocontrol Sci Technol* 24:203–215
- Mota RMA, Abarca GH, Ruíz RFC, Hernández CIB (2008) Two new species of *Polyschema* and *Vanakripa* and other microfungi recorded from mangrove in Veracruz, Mexico. *Mycotaxon* 106:29–40
- Motiejūnaitė J (2011) Lichens and allied fungi from Kamanos state strict nature reserve (northern Lithuania). *Bot Lith* 17:109–116
- Motiejūnaitė J (2012) New data on distribution of *Lemmopsis pelodes* in Lithuania. *Bot Lith* 18:166–168
- Motiejūnaitė J, Czyżewska K (2008) Additions to the biota of lichens and lichenicolous fungi of Poland with note on *Lecania prasinoides* in eastern and central Europe. *Polish Bot J* 53:155–162
- Motiejūnaitė J, Grochowski P (2014) Miscellaneous new records of lichens and lichenicolous fungi. *Herzogia* 27:193–198
- Motiejūnaitė J, von Brackel W, Stoncius D, Preikša Ž (2011) Contribution to the Lithuanian flora of lichens and allied fungi III. *Bot Lith* 17:39–46
- Motiejūnaitė J, Kukwa M, Löhmus P, Markovskaja S et al (2013) Contribution to the Lithuanian flora of lichens and allied fungi IV. *Bot Lith* 19:3–7
- Motiejūnaitė J, Cheskakov SV, Czarnota P, Gagarina LV et al (2016) Ninety-one species of lichens and allied fungi new to Latvia with a list of additional records from Kurzeme. *Herzogia* 29:143–163
- Motlagh MRS, Kaviani B (2008) Characterization of new *Bipolaris* spp.: the causal agent of rice brown spot disease in the north of Iran. *Int J Agri Biol* 10:638–642
- Motlagh MRS, Anvari M (2010) Genetic variation in a population of *Bipolaris oryzae* based on RAPD-PCR in north of Iran. *Afr J Biotech* 9:800–5804
- Mouhajir A, Matray O, Giraud S, Mély L et al (2016) Long-term *Rasamonia argillacea* complex species colonization revealed by rep-PCR in cystic fibrosis patients. *J Clin Microbiol* 54(11):2804–2812
- Moya-Elizondo EA, Jacobsen BJ, Hogg AC, Dyer AT (2011) Population dynamics between *Fusarium pseudograminearum* and *Bipolaris sorokiniana* in wheat stems using real-time qPCR. *Plant Dis* 95:1089–1098
- Mugambi GK, Huhndorf SM (2009a) Molecular phylogenetics of *Pleosporales*: *Melanommataceae* and *Lophiostomataceae* recircumscribed (*Plesporomycetidae*, *Dothideomycetes*, *Ascomycota*). *Stud Mycol* 64:103–121
- Mugambi GK, Huhndorf SM (2009b) Parallel evolution of hysterothelial ascomata in ascolocularous fungi (*Ascomycota*, *Fungi*). *Syst Biodiv* 7:453–464
- Mugambi GK, Huhndorf SM (2010) Multigene phylogeny of the *Coronophorales*: morphology and new species in the order. *Mycologia* 102:185–210
- Mugambi GK, Huhndorf SM, Rogers JD (2009) Two new *xylariaceous* species from Kenya. *Mycotaxon* 108:499–504
- Muggia L, Hafellner J, Wirtz N, Hawksworth DL, Grube M (2008) The sterile microfilamentous lichenized fungi *Cystocoleus ebeneus* and *Racodium rupestre* are relatives of plant pathogens and clinically important dothidealean fungi. *Mycol Res* 112:50–56
- Muggia L, Gueidan C, Grube M (2010a) Phylogenetic placement of some morphologically unusual members of *Verrucariales*. *Mycologia* 102:835–846
- Muggia L, Zellnig G, Rabensteiner J, Grube M (2010b) Morphological and phylogenetic study of algal partners associated with the lichen-forming fungus *Tephromela atra* from the Mediterranean region. *Symbiosis* 51:149–160
- Muggia L, Baloch E, Stabentheiner E, Grube M, Wedin M (2011a) Photobiont association and genetic diversity of the optionally lichenized fungus *Schizoxylon albescens* FEMS Microbial Ecology 75:255–272
- Muggia L, Nelson P, Wheeler T, Yakovchenko, LS et al (2011b) Convergent evolution of a symbiotic duet: the case of the lichen genus *Polychidium* (*Peltigerales*, *Ascomycota*). *Am J Bot* 98:1647–1656
- Muggia L, Gueidan C, Knudsen K, Perlmutter G, Grube M (2013) The lichen connections of black fungi. *Mycopath* 175:523–535
- Muggia L, Pérez-Ortega S, Fryday A, Spribile T, Grube M (2014) Global assessment of genetic variation and phenotypic plasticity in the lichen-forming species *Tephromela atra*. *Fungal Divers* 64:233–251
- Muggia L, Kocourková J, Knudsen K (2015a) Disentangling the complex of *Lichenothelia* species from rock communities in the desert. *Mycologia* 107:1233–1253
- Muggia L, Kopun T, Ertz D (2015b) Phylogenetic placement of the lichenicolous, anamorphic genus *Lichenodiopsis* and its connection to *Muellerella*-like teleomorphs. *Fungal Biol* 119:1115–1128
- Mukherjee AK, Carp MJ, Zuchman R, Ziv T et al (2010) Proteomics of the response of *Arabidopsis thaliana* to infection with *Alternaria brassicicola*. *J Proteom* 73:709–720
- Mukhopadhyay SK, Chatterjee S, Gauri SS, Das SS et al (2014) Isolation and characterization of extracellular polysaccharide Thelebolan produced by a newly isolated psychrophilic Antarctic fungus *Thelebolus*. *Carbohydr Polym* 104:204–212
- Müller ME, Korn U (2013) *Alternaria* mycotoxins in wheat—A 10 years survey in the Northeast of Germany. *Food Control* 34:191–197
- Mungai PG, Hyde KD, Cai L, Njogu JG, Chukeatirote K (2011) Coprophilous ascomycetes of northern Thailand. *Curr Res Environ Appl Mycol* 1:135–159
- Mungai PG, Chukeatirote E, Njogu JG, Hyde KD (2012a) Coprophilous ascomycetes in Kenya: *Chaetomium* species from wildlife dung. *Curr Res Environ Appl Mycol* 2:113–128
- Mungai PG, Chukeatirote E, Njogu JG, Hyde KD (2012b) Coprophilous ascomycetes in Kenya: *Saccobolus* species from wildlife dung. *Mycosphere* 3:111–129
- Mungai PG, Chukeatirote E, Njogu JG, Hyde KD (2012c) Studies of coprophilous ascomycetes in Kenya. *Podospora* species from wildlife dung. *Mycosphere* 3:978–981
- Mungai PG, Njogu JG, Chukeatirote E, Hyde KD (2012d) Coprophilous ascomycetes in Kenya: *Sporormiella* from wildlife dung. *Mycology* 3:234–251
- Mungai PG, Njogu JG, Chukeatirote E, Hyde KD (2012e) Studies of coprophilous ascomycetes in Kenya—*Ascobolus* species from wildlife dung. *Curr Res Environ Appl Mycol* 2:1–16
- Mungai PG, Njogu JG, Chukeatirote E, Hyde KD (2012f) Studies of coprophilous ascomycetes in Kenya: Coprophilous

- Schizothecium* from wildlife dung. Curr Res Environ Appl Mycol 2:84–97
- Muniz D, Llop E, Hladun NL (2013) *Sphinctrina paramerae*, a new Mediterranean lichenicolous species with non-septate spores. Lichenologist 45:137–143
- Munzenberger B, Bubner B, Wollecke J, Sieber TN, Bauer R (2009) The ectomycorrhizal morphotype *Pinirhiza sclerotia* is formed by *Acephala macrosclerotorum* sp. nov., a close relative of *Phialocephala fortinii*. Mycorrhiza 19:481–492
- Murata N, Aoki T, Kusaba M, Tosa Y, Chuma I (2014) Various species of *Pyricularia* constitute a robust clade distinct from *Magnaporthe salvinii* and its relatives in *Magnaporthaceae*. J Gen Plant Pathol 80:66–72
- Murillo C, Albertazzi FJ, Carranza J, Lumbsch HT, Tamayo G (2009) Molecular data indicate that *Rhytidhysteron rufulum* (ascomycetes, *Patellariales*) in Costa Rica consists of four distinct lineages corroborated by morphological and chemical characters. Mycol Res 113:405–416
- Musvuugwa T, Dreyer LL, De Beer WZ, Oberlander K et al (2015) New species of *Ophiostomatales* from Scolytinae and Platypodinae beetles in the Cape Floristic Region, including the discovery of the sexual state of Raffaelea. Antonie van Leeuwenhoek 108:933–950
- Muthukrishnan S, Sanjayan KP, Jahir HK (2012) Species composition, seasonal changes and community ordination of alkalotolerant micro fungal diversity in a natural scrub jungle ecosystem of Tamil Nadu, India. Mycosphere 3:92–109
- MycoBank (2016) <http://www.mycobank.org>. Accessed Dec 2016
- Mylllys L, Velmala S, Holien H, Halonen P et al (2011) Phylogeny of the genus *Bryoria*. Lichenologist 43:617–638
- Mylllys L, Velmala S, Lindgren H, Glavich D et al (2014) Taxonomic delimitation of the genera *Bryoria* and *Sulcaria*, with a new combination *Sulcaria spiralisfera* introduced. Lichenologist 46:737–752
- Nadyeina O (2009) The lichen-forming and lichenicolous fungi of the Donets Upland (Ukraine). Mycol Balt 6:37–53
- Nadyeina O, Grube M, Mayrhofer H (2010) A contribution to the taxonomy of the genus *Rinodina* (*Physciaceae*, lichenized Ascomycotina) using combined ITS and mtSSU rDNA data. Lichenologist 42:521–531
- Nagahama T, Abdel-Wahab MA, Nogi Y, Miyazaki M et al (2008) *Dipodascus tetrasporous* sp. nov., an ascosporogenous yeast isolated from deep-sea sediments in the Japan Trench. Int J Syst Evol Microbiol 58:1040–1046
- Nagao H (2008) Discomycetes on decayed tree fern. (3) *Lachnum lanariceps* and *Lachnum oncospermaum* new to Japan. Mycoscience 49:403–406
- Nagao H, Kurogi SI, Kiyota E, Sasatomi K (2009) *Kumanasamuha geaster* sp. nov., an anamorph of *Chorioactis geaster* from Japan. Mycologia 101:871–877
- Nagano Y, Nagahama T, Abe F (2014) Cold-adapted yeasts in deep-sea environments. Cold-adapted yeasts. Springer, Berlin Heidelberg, pp 149–171
- Nagaraju D, Kunwar IK, Suresh Kumar G, Manoharachary C (2011a) *Custingophora lignicola* sp. nov. and *Chaetopsina indica* sp. nov. from India. J Mycol Plant Pathol 41:9
- Nagaraju D, Kunwar IK, Sureshkumar G, Manoharachary C (2011b) A new synnematosus hyphomycetous fungus—*Bhadradriella* gen. nov. from India. J Mycol Plant Pathol 41:238
- Najafzadeh MJ, Sun J, Vicente V, Xi L et al (2010) *Fonsecaea nubica* sp. nov., a new agent of human chromoblastomycosis revealed using molecular data. Med Mycol 48:800–806
- Najafzadeh MJ, Sun J, Vicente VA, Klaassen CHW et al (2011a) Molecular epidemiology of *Fonsecaea* species. Emerg Infect Dis 17:464–469
- Najafzadeh MJ, Vicente VA, Sun J, Meis JF, de Hoog GS (2011b) *Fonsecaea multimorphosa* sp. nov., a new species of *Chaetothyriales* isolated from a feline cerebral abscess. Fungal Biol 115:1066–1076
- Nalim FA, Samuels GJ, Wijesundara RL, Geiser DM (2011) New species from the *Fusarium solani* species complex derived from perithecia and soil in the old world tropics. Mycologia 103:1302–1330
- Nallathambi P, Umamaheswari C, Thakore BBL, More TA (2009) Post-harvest management of ber (*Ziziphus mauritiana* Lamk) fruit rot (*Alternaria alternata* Fr. Keissler) using *Trichoderma* species, fungicides and their combinations. Crop Prot 28:525–532
- Nambiar GR, Raveendran K (2009a) Marine mycoflora in backwater ecosystem of kerala, India. J Environ Biol 30:647–649
- Nambiar GR, Raveendran K (2009b) Manglicolous marine fungi on *Avicennia* and *Rhizophora* along Kerala Coast (India). Middle East J Sci Res 4:48–51
- Nambiar GR, Raveendran K, Jaleel CA (2009) Marine mycoflora of south India with special emphasis on lignicolous marine fungi. Front Biol China 4:436–441
- Nascimento MM, Selbmann L, Sharifyna S, Al-Hatmi AM et al (2016) *Arthrocladium*, an unexpected human opportunist in *Trichocomaceae* (*Chaetothyriales*). Fungal Biol 120:207–218
- Nattah I, Ajaj A, Chliyeh M, Touhami AO, Benkirane R, Douira A (2014) Contribution to the inventory of lichens encountered in the reserve of Sidi Bouhaba (Morocco). Int J Pure App Biosci 2:103–107
- Naumann TA, Wicklow DT, Kendra DF (2009) Maize seed chitinase is modified by a protein secreted by *Bipolaris zeicola*. Physiol Mol Plant Pathol 74:134–141
- Navarrete F, Abreo E, Martínez S, Bettucci L, Sandra LU (2011) Pathogenicity and molecular detection of Uruguayan isolates of *Greeneria uvicola* and *Cadophora luteo-olivacea* associated with grapevine trunk diseases. Phytopathol Mediterr 50:166–175
- Navarro-Rosinés P, Hladun NL, Llimona X (2008) *Gelatinopsis heppiae* sp. nov. (Leotiales, Ascomycota) un hongo liquenícola sobre *Heppia desprauxii* de Aragón (NE de España). Rev Catalana Micol 30:57–64
- Navarro-Rosinés P, Zhurbenko M, Roux C (2010) *Pseudopyrenidium* un nouveau genre pour inclure *Weddellomyces tartaricola* (champignon lichenique non lichenisé, Ascomycota). Bull Mens Soc Linn Paris 61:129–140
- Nayaka S, Saxena P (2014) Physiological responses and ecological success of lichen *Stereocaulon foliolosum* and moss *Racomitrium subsecundum* growing in same habitat in Himalaya. Indian J Fund Appl Life Sci 4:167–179
- Nayaka S, Upadhyay DK, Ponmurgan P, Ayyappadasan G (2013) Two new species of saxicolous *Pyxine* with yellow medulla from southern India. Lichenologist 45:3–8
- Nekoduka S, Tanaka K, Harada Y, Sano T (2010) Phylogenetic affinity of *Mycochaetophora gentianae*, the causal fungus of brown leaf spot on gentian (*Gentiana triflora*), to *Pseudocercospora*-like hyphomycetes in Helotiaceae. Mycoscience 51:123–133
- Nekoduka S, Kanematsu S, Tanaka K, Harada Y, Sano T (2012) *Valdensia* leaf blight of highbush blueberry caused by *Valdensinia heterodoxa*, a new fungal disease in Japan. J Gen Plant Pathol 78:151–159
- Nelsen MP, Gargas A (2009) Assessing clonality and chemotype monophly in *Thamnolia* (*Icmadophilaceae*). Bryologist 112:42–53
- Nelsen MP, Lumbsch HT, Lücking R, Elix JA (2008) Further evidence for the polyphyly of *Lepraria* (*Lecanorales: Stereocaulaceae*). Nova Hedwig 87:361–371

- Nelsen MP, Lücking R, Grube M, Mbatchou JS et al (2009) Unravelling the phylogenetic relationships of lichenised fungi in *Dothideomyceta*. Stud Mycol 64:135–144
- Nelsen MP, Lücking R, Plata ER, Mbatchou JS (2010) *Heiomasia*, a new genus in the lichen-forming family *Graphidaceae* (Ascomycota: Lecanoromycetes: Ostropales) with disjunct distribution in Southeastern North America and Southeast Asia. Bryologist 113:742–751
- Nelsen MP, Chavez N, Sackett-Hermann E, Thell A et al (2011a) The cetrarioid core group revisited (Lecanorales: Parmeliaceae). Lichenologist 43:537–551
- Nelsen MP, Lücking R, Mbatchou JS, Andrew CJ et al (2011b) New insights into relationships of lichen-forming *Dothideomycetes*. Fungal Divers 51:155–162
- Nelsen MP, Lücking R, Andrew CJ, Plata ER et al (2012) Dismantling *Herpothallon*: *Herpothallon antillarum* (Arthoniomycetes: Arthoniaceae) is a member of the genus *Diorygma* (Lecanoromycetes: Graphidaceae). Bryologist 115:313–321
- Nelsen MP, Thell A, Leavitt SD, Hampton-Miller CJ et al (2013) A reappraisal of *Masonhalea* (Parmeliaceae, Lecanorales) based on molecular and morphological data. Lichenologist 45:729–738
- Nelsen MP, Lücking R, Andrew CJ, Aptroot A et al (2014a) Molecular phylogeny reveals the true colors of *Myeloconidaceae* (Ascomycota: Ostropales). Aust Syst Bot 27:38–47
- Nelsen MP, Lücking R, Aptroot A, Andrew CJ et al (2014b) Elucidating phylogenetic relationships and genus-level classification within the fungal family *Trypetheliaceae* (Dothideomycetes: Ascomycota). Taxon 63:974–992
- Nelsen MP, Lücking R, Cáceres MES, Aptroot A (2016) Assessing the phylogenetic placement and redundancy of *Aspidotheliaceae* (Ascomycota), an orphaned family of lichenforming fungi. Syst Biodivers 15:63–73
- Netam RS, Bahadur AN, Tiwari U, Tiwari RKS (2011) Efficacy of plant extracts for the control of (*Pyricularia grisea*) blast of rice under field condition of bastar, Chhattisgarh. Res J Agric Sci 2:269–271
- Neuwirth G, Aptroot A, Stocker-Woergoetter E (2017) *Platythecium seychellense*, a new species in the family *Graphidaceae* (lichenized Ascomycota: Ostropales) from the Seychelles and a world key to the genus. Lichenologist 49:85–91
- Nguanhom J, Cheewangkoon R, Groenewald JZ, Braun U et al (2015) Taxonomy and phylogeny of *Cercospora* spp. from Northern Thailand. Phytotaxa 233:27–48
- Nguyen HDT, Seifert KA (2008) Description and DNA barcoding of three new species of *Leohumicola* from South Africa and the United States. Persoonia 21:57–69
- Nicoletti R, Wolczanska A, Carella A (2010) Occurrence of *Zetiasplozna unicolor* as an endophyte of myrtle (*Myrtus communis*) in Italy. Petria 20:581
- Nilsson RH, Hyde KD, Pawłowska J, Ryberg M et al (2014) Improving ITS sequence data for identification of plant pathogenic fungi. Fungal Divers 67:11–19
- Niu XM, Zhang KQ (2011) *Arthrobotrys oligospora*: a model organism for understanding the interaction between fungi and nematodes. Mycology 2(2):59–78
- Nizam S, Verma S, Singh K, Aggarwal R et al (2012) High reliability transformation of the wheat pathogen *Bipolaris sorokiniana* using *Agrobacterium tumefaciens*. J Microbiol Methods 88:386–392
- Nkuekam GK, Wingfield MJ, Mohammed C, Carnegie AJ et al (2012) *Ceratocystis* species, including two new species associated with nitidulid beetles, on eucalypts in Australia. Antonie Van 101:217–241
- Nonaka K, Miyazaki H, Iwatsuki M, Shiomi K et al (2012) *Staphylocutrichum boninense*, a new hyphomycete (Chaetomiaceae) from soils in the Bonin Islands, Japan. Mycoscience 53:312–318
- Nonaka K, Ishii T, Shiomi K, Ōmura S, Masuma R (2013a) *Virgaria boninensis*, a new hyphomycete (Xylariaceae) from soils in the Bonin Islands, Japan. Mycoscience 54:394–399
- Nonaka K, Kaifuchi S, Ōmura S, Masuma R (2013b) Five new *Simplicillium* species (Cordycipitaceae) from soils in Tokyo, Japan. Mycoscience 54:42–53
- Nonaka K, Kaifuchi S, Omura S, Masuma R (2013c) Three new *Pochonia* taxa (Clavicipitaceae) from soils in Japan. Mycologia 6:12–32
- Nordin A, Savić S, Tibell L (2010) Phylogeny and taxonomy of *Aspicilia* and *Megasporaceae*. Mycologia 102:1339–1349
- Norphanphoun C, Maharachchikumbura SSN, Daranagama A, Bulgakov TS et al (2015) Towards a backbone tree for *Seimatosporium*, with *S. physocarpi* sp. nov. Mycosphere 6:385–400
- Norphanphoun C, Hongsanan S, Doilom M, Bhat DJ et al (2016) *Lamproconiaceae* fam. nov. to accommodate *Lamproconium desmazieri*. Phytotaxa 270:89–102
- Norphanphoun C, Doilom M, Daranagama DA, Phookamsak R et al (2017) Revisiting the genus *Cytospora* and allied species. Mycosphere 8:51–97
- Norvell LL (2010) Report of the nomenclature committee for fungi: 15. Taxon 59:291–293
- Noser J, Schneider P, Rother M, Schmutz H (2011) Determination of six *Alternaria* toxins with UPLC-MS/MS and their occurrence in tomatoes and tomato products from the Swiss market. Mycotoxin research 27:265–271
- Notov AA (2014) Fruticose lichens: structural diversity, taxonomic characteristics and evolution. Wulfenia 21:21–31
- Nouhra E, Urcelay C, Longo S, Tedersoo L (2013) Ectomycorrhizal fungal communities associated to *Nothofagus* species in Northern Patagonia. Mycorrhiza 23:487–496
- Nowicki M, Nowakowska M, Niezgoda A, Kozik E (2012) *Alternaria* black spot of crucifers: symptoms, importance of disease and perspectives of resistance breeding. Veg Crops Res Bull 76:5–19
- Ntui VO, Thirukkumaran G, Azadi P, Khan RS et al (2010) Stable integration and expression of wasabi defensin gene in “Egusi” melon (*Colocynthis citrullus* L.) confers resistance to *Fusarium* wilt and *Alternaria* leaf spot. Plant Cell Rep 29:943–954
- Núñez-Zapata J, Divakar PK, Del-Prado R, Cubas P et al (2011) Conundrums in species concepts: the discovery of a new cryptic species segregated from *Parmelina tiliacea* (Ascomycota: Parmeliaceae). Lichenologist 43:603–616
- Nyati S, Bhattacharya D, Werth S, Honegger R (2013) Phylogenetic analysis of LSU and SSU rDNA group I introns of lichen photobionts associated with the genera *Xanthoria* and *Xanthomendoza* (Teloschistaceae, lichenized ascomycetes). J Phycol 49:1154–1166
- Nygren K, Strandberg R, Wallberg A, Nabholz B et al (2011) A comprehensive phylogeny of *Neurospora* reveals a link between reproductive mode and molecular evolution in fungi. Mol Phylogen Evol 59:649–663
- O’Connell RJ, Thon MR, Hacquard S, Amyotte SG et al (2012) Lifestyle transitions in plant pathogenic *Colletotrichum* fungi deciphered by genome and transcriptome analyses. Nat Genet 44:1060–1065
- Ohmura Y, Yakovchenko L, Zhurbenko MP (2014) *Carbonea vitellinaria* new to Japan, with a key to lichenicolous fungi growing on species of *Candelariella*. Mycosphere 5:607–611
- Ohmura Y, Frisch A, Lendemer JC, Nakanishi M (2016) Lichens of Mikura-jima Island, Izu Islands, Central Japan. Bull Natl Mus Natl Sci Ser 42:1–18

- Olariaga I, Van Vooren N, Carbone M, Hansen K (2015) A monograph of *Otidea* (*Pyronemataceae*, *Pezizomycetes*). *Persoonia* 35:166–229
- Oliveira RJ, Lima TE, Cunha IB, Coimbra VR et al (2014) *Corniculariella brasiliensis*, a new species of coelomycetes in the rhizosphere of *Caesalpinia echinata* (*Fabaceae*, *Caesalpinioideae*) in Brazil. *Phytotaxa* 178:197–204
- Oliveira MS, Malosso E, Barbosa MA, Araújo MA, Castañeda-Ruiz RF (2015) *Xylomyces acerosisporus* sp. nov. from submerged leaves from Brazil. *Mycotaxon* 130:875–878
- Oliveira H, Santos C, Paterson RRM, Gusmão NB, Lima N (2016a) Fungi from a groundwater-fed drinking water supply system in Brazil. *Int J Environ Res Publ Health* 13:304
- Oliveira MS, Malosso E, Castañeda-Ruiz RF (2016b) A new species and a new combination in *Codinaea* from Brazil. *Mycotaxon* 130:1045–1049
- Orange A (2009a) A new species of *Petractis* (*Ostropales s. lat.*, lichenized *Ascomycota*) from Wales. *Lichenologist* 41:213–321
- Orange A (2009b) Two parasitic species of *Placopyrenium* (*Verrucariaceae*) from freshwater habitats in North-west Europe. *Lichenologist* 41:131–139
- Oset M (2015) The lichen genus *Stereocaulon* (Schreb.) Hoffm. in Poland—a taxonomic and ecological study. *Monogr Bot* 104:1–81
- Oset M, Kukwa M (2014) Typification of two *Stereocaulon* names (*Lecanorales*, *Ascomycota*). *Phytotaxa* 181:188–191
- Osmundson TW, Robert VA, Schoch CL, Baker LJ et al (2013) Filling gaps in biodiversity knowledge for macrofungi: contributions and assessment of an herbarium collection DNA barcode sequencing project. *PLoS ONE* 8:e62419
- Osono T, Masuya H (2012) Endophytic fungi associated with leaves of *Betulaceae* in Japan. *Can J Microbiol* 58:507–515
- Ostry V (2008) *Alternaria* mycotoxins: an overview of chemical characterization, producers, toxicity, analysis and occurrence in foodstuffs. *World Mycotoxin J* 1:175–188
- Otalora MAG, Wedin M (2013) *Collema fasciculare* belongs in *Arctomiaceae*. *Lichenologist* 45:295–304
- Otalora MA, Aragón G, Molina MC, Martínez I, Lutzoni F (2010) Disentangling the *Collema-Leptogium* complex through a molecular phylogenetic study of the *Collemataceae* (*Peltigerales*, lichen-forming *Ascomycota*). *Mycologia* 102:279–290
- Otalora MA, Aragón G, Martínez I, Wedin M (2013) Cardinal characters on a slippery slope—A re-evaluation of phylogeny, character evolution, and evolutionary rates in the jelly lichens (*Collemataceae* s. str) Mol. Phylogenetic Evol 68:185–198
- Otalora MA, Jørgensen PM, Wedin M (2014) A revised generic classification of the jelly lichens, *Collemataceae*. *Fungal Divers* 64:275–293
- Ouda SM (2014) Antifungal activity of silver and copper nanoparticles on two plant pathogens, *Alternaria alternata* and *Botrytis cinerea*. *Res J Microbiol* 9:34
- Overy DP, Baymane P, Kerr RG, Bills GF (2014) An assessment of natural product discovery from marine (*sensu strictu*) and marine-derived fungi. *Mycology* 5:145–167
- Øvstedal DO (2012) New species, combinations and records of lichenized fungi from the Falkland Islands (Islas Malvinas). *Lichenologist* 44:483–500
- Øvstedal DO, Gremmen NJM (2009) Additional lichen records from Subantarctica 1. The Kerguelen Islands. *Australas Lichenol* 64:3–9
- Øvstedal DG, Gremmen NJM (2010) New lichen species from Tristan da Cunha and Gough Island. *Folia Cryptogam Estonica* 47:43–49
- Øvstedal DO, Lewis Smith RI (2011) Four additional lichens from the Antarctic and South Georgia, including a new *Leciophysma* species. *Folia Cryptogam Estonica* 48:65–68
- Øvstedal DO, Gremmen NJ (2014) Additional lichen records from Subantarctica III. Marion Island. *Australas Lichenol* 74:2–7
- Øvstedal D, Tønsberg T, Elvebakk A (2009) The lichen flora of Svalbard. *Sommerfeltia* 33:3–393
- Paciura D, de Beer ZW, Jacobs K, Zhou XD et al (2010a) Eight new *Leptographium* species associated with tree-infesting bark beetles in China. *Persoonia* 25:94–108
- Paciura D, Zhou XD, De Beer ZW, Jacobs K, Ye H (2010b) Characterisation of synnematous bark beetle-associated fungi from China, including *Graphium carbonarium* sp. nov. *Fungal Divers* 40:75–88
- Palmateer AJ, Perez JM, Cating RA, Ploetz RC (2008) First report of tar spot on orange geiger (*Cordia sebestena*) caused by *Diatractium cordianum* in the US. *Plant Dis* 92:1250
- Palmer JM, Lindner DL, Volk TJ (2008) Ectomycorrhizal characterization of an American chestnut (*Castanea dentata*)-dominated community in Western Wisconsin. *Mycorrhiza* 19:27–36
- Palomares-Rius JE, Hirooka Y, Tsai IJ, Masuya H et al (2014) Distribution and evolution of glycoside hydrolase family 45 cellulases in nematodes and fungi. *BMC Evol Biol* 14:69
- Pan HQ, Zhang TY, Kong JH (2009) Notes on soil dematiaceous hyphomycetes from the Yellow River source area, China I. *Mycosistema* 28:14–19
- Pan HQ, Wang HF, Jiang YL, Zhang TY (2014) *Scopulariopsis*: three new species and a key to species from soils in China. *Mycosistema* 33:1–6
- Pandey SP, Sharma S, Chand R, Shahi P, Joshi AK (2008) Clonal variability and its relevance in generation of new pathotypes in the spot blotch pathogen, *Bipolaris sorokiniana*. *Current Microbiol* 56:33–41
- Pandit GS (2014) Lichens of the Mahabaleshwar Panchgani Ecosensitive zone (MPESZ), Maharashtra, India. *J Threatened Taxa* 6:5784–5791
- Pandit G (2015) Review of lichens of the high level Ferricretes and Mesas of the North Western Ghats, India. *Current Res Environ Appl Mycol* 5:180–195
- Pandit GS, Sharma BO (2012) New records in the lichen family *Lobariaceae* from the Western Ghats of India. *Mycosphere* 3:430–435
- Pandit RJ, Bhatt VD, Mukhopadhyaya PN, Joshi CG, Kunjadia AP (2014) Biochemical and molecular characterization of protease from *Arthrobotrys conoides* and *Duddingtonia flagrans*. *Int J Adv Biotechnol Res* 5:552–561
- Pane C, Celano G, Villecco D, Zaccardelli M (2012) Control of *Botrytis cinerea*, *Alternaria alternata* and *Pyrenopeziza lycopersici* on tomato with whey compost-tea applications. *Crop Prot* 38:80–86
- Pang KL, Jheng JS (2012a) Checklist of marine fungi of Taiwan with a description of *Kitesporella keelungensis* gen. et sp. nov. *Bot Mar* 55:459–466
- Pang KL, Jheng JS (2012b) *Pileomyces formosanus* gen. et sp. nov. (*Halosphaeriaceae*, *Ascomycota*) from a rocky shore of Taiwan. *Bot Stud* 53:535–539
- Pang KL, Chiang MWL, Vrijmoed LLP (2008a) *Havispora longyearbyensis* gen. et sp. nov.: an arctic marine fungus from Svalbard, Norway. *Mycologia* 100:291–295
- Pang KL, Jones EBG, Vrijmoed LLP (2008b) Autecology of *Antennosporea* (*Ascomycota*, *Fungi*) and its phylogeny. *Raff Bull Zool* 19:1–10
- Pang KL, Alias SA, Chiang MWL, Vrijmoed LLP, Jones EBG (2010a) *Sedecimiella taiwanensis* gen. et sp. nov., a marine

- mangrove fungus in the Hypocreales (*Hypocreomycetidae*, *Ascomycota*). Bot Mar 53:493–498
- Pang KL, Sharuddin SS, Alias SA, Nor NAM, Awaluddin HH (2010b) Diversity and abundance of lignicolous marine fungi from the east and west coasts of Peninsular Malaysia and Sabah (Borneo Island). Bot Mar 53:515–523
- Pang KL, Chow RKK, Chan CW, Vrijmoed LLP (2011a) Diversity and physiology of marine lignicolous fungi in Arctic waters: a preliminary account. Polar Res 30:1–5
- Pang KL, Jheng JS, Jones EBG (2011b) Marine mangrove fungi of Taiwan. National Taiwan Ocean University Press, Keelung
- Pang KL, Hyde KD, Alias SA, Suetrong S et al (2013a) *Dyfrolomycetaceae*, a new family in the *Dothideomycetes*. Cryptogam Mycol 34:223–232
- Pang KL, Vrijmoed LLP, Jones EBG (2013b) Genetic variation within the cosmopolitan aquatic fungus *Lignincola laevis* (*Microascales*, *Ascomycota*). Org Divers Evol 13:301–309
- Pang KL, Guo SY, Alias SA, Hafellner J, Jones EBG (2014) A new species of marine *Dactylospora* and its phylogenetic affinities within the *Eurotiomycetes*, *Ascomycota*. Bot Mar 57:315–321
- Pańska D, Jeske M, Troczyński M (2013) Occurrence of *Neotyphodium* and *Epichloë* fungi in meadow fescue and red fescue in Poland and screening of endophyte isolates as potential biological control agents. Acta Sci Pol Hortorum Cultus 12:67–83
- Panzer K, Yilmaz P, Weiß M, Reich L et al (2015) Identification of habitat-specific biomes of aquatic fungal communities using a comprehensive nearly full-length 18S rRNA dataset enriched with contextual data. PLoS ONE 10:e0134377
- Paoli L, Munzi S, Gutová A, Senko D et al (2015) Lichens as suitable indicators of the biological effects of atmospheric pollutants around a municipal solid waste incinerator (S Italy). Ecol Indic 52:362–370
- Papazi A, Kastanaki E, Pirintsos S, Kotzabasis K (2015) Lichen symbiosis: nature's high yielding machines for induced hydrogen production. PloS ONE 10:e0121325
- Papizadeh M, Soudi MR, Amini L, Wijayawardene NN, Hyde KD (2017) *Pyrenophaetopsis tabarestanensis* (*Cucurbitariaceae*, *Pleosporales*), a new species isolated from rice farms in north Iran. Phytotaxa 291:15–28
- Papong K, Lumbsch HT (2011) A taxonomic survey of *Lecanora* *sensu stricto* in Thailand (*Lecanoraceae*; *Ascomycota*). Lichenologist 43:299–320
- Papong K, Lücking R, Thammathaworn A, Boonpragob K (2009a) Four new taxa of *Chroodiscus* (thelotretemoid *Graphidaceae*) from Southeast Asia. Bryologist 112:152–163
- Papong K, Mangold A, Corush J, Lücking R, Lumbsch HT (2009b) Phylogenetic position of the foliicolous genus *Chroodiscus* (*Ostropales*, *Ascomycota*) inferred from nuclear and mitochondrial ribosomal DNA sequences. Fungal Divers 38:147–153
- Papong K, Boonpragob K, Mangold A, Divakar PK et al (2010) Thelotretemoid lichen species recently described from Thailand: a re-evaluation. Lichenologist 42:131–137
- Papong K, Kantvilas G, Lumbsch HT (2011) Morphological and molecular evidence places *Maronina* into synonymy with *Protoparmelia* (*Ascomycota*: *Lecanorales*). Lichenol 43:561–567
- Papong K, Boonpragob K, Parnmen S, Lumbsch HT (2013) Molecular phylogenetic studies on tropical species of *Lecanora* *sensu stricto* (*Lecanoraceae*, *Ascomycota*). Nova Hedw 96:1–13
- Papong KB, Luecking R, Kraichak E, Parnmen S et al (2014) Twenty-three new species in the lichen family *Graphidaceae* from New Caledonia (*Ostropales*, *Ascomycota*). Phytotaxa 189:204–231
- Parada RY, Sakuno E, Mori N, Oka K et al (2008) *Alternaria brassicae* produces a host-specific protein toxin from germinating spores on host leaves. Phytopathology 98:458–463
- Park MJ, Shin HD (2009) A new species of *Pyricularia* on *Commelina communis*. Mycotaxon 108:449–456
- Park JH, Shin HD (2016) *Ramularia hydrangeicola* sp. nov. with distinctive traits on *Hydrangea serrata* f. *acuminata* in Korea. Mycotaxon 131:95–102
- Park MS, Romanoski CE, Pryor BM (2008) A re-examination of the phylogenetic relationship between the causal agents of carrot black rot, *Alternaria radicina* and *A. carotinctulae*. Mycologia 100:511–527
- Park MJ, Choi YJ, Hong SB, Shin HD (2010) Genetic variability and mycohost association of *Ampelomyces quisqualis* isolates inferred from phylogenetic analyses of ITS rDNA and actin gene sequences. Fungal Biol 114:235–247
- Park SY, Jeong MH, Wang HY, Kim JA et al (2013) *Agrobacterium tumefaciens*-mediated transformation of the lichen fungus, *Umbilicaria muehlenbergii*. PloS ONE 8(12):e83896
- Park JH, Cho SE, Hong SH, Choi IY, Shin HD (2015) Sooty mould on *hibiscus rosa-sinensis* caused by *Leptoxyphium kurandae* is associated with extrafloral nectaries. J Phytopathol 163:1027–1030
- Parnmen S, Rangsiruji A, Mongkolsuk P, Boonpragob K et al (2010) Morphological disparity in *Cladoniaceae*: the foliose genus *Heterodea* evolved from fruticose *Cladia* species (*Lecanorales*, lichenized *Ascomycota*). Taxon 59:841–849
- Parnmen S, Lücking R, Lumbsch HT (2012) Phylogenetic classification at generic level in the absence of distinct phylogenetic patterns of phenotypical variation: a case study in *Graphidaceae* (*Ascomycota*). PLoS ONE 7:e51392
- Parnmen S, Cáceres ME, Lücking R, Lumbsch HT (2013) *Myriochapsa* and *Nitidochapsa*, two new genera in *Graphidaceae* (*Ascomycota*: *Ostropales*) for chroodiscoid species in the *Ocellularia* clade. Bryologist 116:127–133
- Parsa SG, Moniri MH, Asoodeh A (2013) A study on *Tornabea scutellifera* (Lichenized Ascomycete, *Lecanorales*) in northeastern Iran. J App Nat Sci 5:148–152
- Pärte K, Pöldmaa K (2011) A new species of *Hyphodiscus* (*Helotiiales*) on *Stereum*. Mycotaxon 115:11–17
- Pärte K, Baral HO, Tamm H, Pöldmaa K (2017) Evidence for the polyphyly of *Encoelia* and *Encoelioidae* with reconsideration of respective families in *Leotiomycetes*. Fungal Divers 82:183–219
- Pasche JS, Gudmestad NC (2008) Prevalence, competitive fitness and impact of the F129L mutation in *Alternaria solani* from the United States. Crop Prot 27:427–435
- Passo A, Calvelo S (2011) *Pannaria byssoidaea* (*Pannariaceae*), a new squamulose species from southern South America. Bryologist 114:756–763
- Pastirčáková K, Pastirčák M (2010) Two important ascomycetes and their anamorphs on twigs of *Betula pendula* Slovakia. Pol Bot J 55:373–380
- Pastor FJ, Guarro J (2008) *Alternaria* infections: laboratory diagnosis and relevant clinical features. Clin Microbiol Infect 14:734–746
- Patil SY, Patil VR, Nemade LC (2010) Biodiversity of freshwater mitosporic fungi from Dhule District (MS), India. J Ecobiotech 2:25–28
- Paukov AG, Teptina AY (2012) New records of lichens from Middle Urals, Russia. Folia Cryptogam Estonica 49:39–43
- Pavlic D, Wingfield MJ, Barber P, Slippers B et al (2008) Seven new species of the *Botryosphaeriaceae* from baobab and other native trees in Western Australia. Mycologia 100:851–866
- Pavlic D, Slippers B, Coutinho TA, Wingfield MJ (2009) Molecular and phenotypic characterization of three phylogenetic species discovered within the *Neofusicoccum parvum*/N. *ribis* complex. Mycologia 101:636–647

- Paz A, Lavoise C, Barrio L, Richard F, Moreau PA (2012) Propuesta de dos nuevas especies del género *Elaphomyces*, dos primeras citas para la Península Ibérica y una clave de identificación de las especies del género para Europa. *Bol Micol FAMCAL* 7:85–104
- Pažoutová S, Kolařík M, Odvody GN, Frederickson DE et al (2008) A new species complex including *Claviceps fusiformis* and *Claviceps hirtella*. *Fungal Divers* 31:95–110
- Pažoutová S, Srutka P, Holusa J, Chudickova M et al (2010) The phylogenetic position of *Obolarina dryophila* (*Xylariales*). *Mycol Prog* 9:501–507
- Pažoutová S, Odvody GN, Frederickson DE, Chudíčková M et al (2011) New *Claviceps* species from warm-season grasses. *Fungal Divers* 49:145–165
- Pažoutová S, Šrůtka P, Holuša J, Chudíčková M et al (2012) *Liberomyces* gen. nov. with two new species of endophytic coelomycetes from broad leaf trees. *Mycologia* 104:198–210
- Pažoutová S, Pešicová K, Chudíčková M, Šrůtka P, Kolařík M (2015) Delimitation of cryptic species inside *Claviceps purpurea*. *Fungal Biol* 119:7–26
- Pedras MSC, Chumala PB, Jin W, Islam MS, Hauck DW (2009) The phytopathogenic fungus *Alternaria brassicicola*: phytotoxin production and phytoalexin elicitation. *Phytochemistry* 70:394–402
- Peláez F, Collado J, Platas G, Overy DP et al (2011) Phylogeny and intercontinental distribution of the pneumocandin-producing anamorphic fungus *Glarea lozoyensis*. *Mycology* 2:1–17
- Peláez RN, Moncada B, Lücking R (2014) High diversity of *Ocellularia* (*Ascomycota: Graphidaceae*) in the Colombian Llanos, including two species new to science. *Phytotaxa* 189:245–254
- Peng L, Yang Y, Hyde KD, Bahkali AH et al (2012) *Colletotrichum* species on Citrus leaves in Guizhou and Yunnan provinces, China. *Cryptogam Mycol* 33:267–283
- Perdomo H, García D, Gené J, Cano J et al (2013) *Phialemoniopsis*, a new genus of *Sordariomycetes*, and new species of *Phialemonium* and *Lecythophora*. *Mycologia* 105:398–421
- Pereira J, Rogers JD, Bezerra JL (2010) New *Annulohypoxylon* species from Brazil. *Mycologia* 102:248–252
- Pereira-Carvalho RC, Dornelo-Silva D, Inácio CA, Dianese JC (2009a) *Chaetothyriomyces*: a new genus in family *Chaetothyriaceae*. *Mycotaxon* 107:483–488
- Pereira-Carvalho RC, Sepúlveda-Chavera GF, Armando EAS, Inácio CA, Dianese JC (2009b) An overlooked source of fungal diversity: novel hyphomycete genera on trichomes of cerrado plants. *Mycol Res* 113:261–274
- Pereira-Carvalho RD, Inácio CA, Dianese JC (2010) *Plurispermiopsis*: a new capnodiaceous genus from the Brazilian Cerrado. *Mycologia* 102:1163–1166
- Perello A, Moreno M, Sisterna M (2008) Alternaria infectoria species-group associated with black point of wheat in Argentina. *Plant Pathol* 57:379–379
- Perera RH, Maharachchikumbura SS, Bahkali AH, Camporesi E et al (2016a) Sexual morph of *Seimatosporium cornii* found on *Cornus sanguinea* in Italy. *Phytotaxa* 257:51–60
- Perera RH, Maharachchikumbura SSN, Bhat JD, Al-Sadi AM et al (2016b) New species of *Thozetella* and *Chaetosphaeria* and new records of *Chaetosphaeria* and *Tainosphaeria* from Thailand. *Mycosphere* 7:1301–1321
- Perera RH, Maharachchikumbura SS, Ariyawansa H, Bahkali AH et al (2016c) Two new *Pseudohalonectria* species on beech cupules (*Fagus sylvatica*) and a new genus to accommodate *P. suthepensis*. *Phytotaxa* 278:115–131
- Pérez-Ortega S, Calatayud V (2009) *Lichenostigma epirupestre*, a new lichenicolous species on *Pertusariaceae* from Spain. *Mycotaxon* 107:189–195
- Pérez-Ortega S, Etayo J (2010) *Labrocarpon* gen. nov. for *Melaspilea canariensis*, with the description of *Buellia protoparmeliopsis* sp. nov. from South America. *Lichenologist* 42:271–276
- Pérez-Ortega S, Spribille T (2009) Three new species of *Lichenopeltella* (*Microthyriaceae, Ascomycota*) from Northwest North America. *Nova Hedw* 89:219–228
- Pérez-Ortega S, de los Ríos A, Crespo A, Sancho LG (2010) Symbiotic lifestyle and phylogenetic relationships of the bionts of *Mastodia tessellata* (*Ascomycota, incertae sedis*). *Am J Bot* 97:738–752
- Pérez-Ortega S, Etayo J, Elix JA, Crespo A (2011a) A new species of *Zwackhiomyces* (*Xanthopyreniaceae, Ascomycota*) growing on *Austroparmelina* from Australia. *Nova Hedw* 93:395–400
- Pérez-Ortega S, Suija A, de los Ríos A (2011b) The connection between *Abrothallus* and its anamorph state *Vouauxiomycetes* established by denaturing gradient gel electrophoresis (DGGE). *Lichenologist* 43:277–279
- Pérez-Ortega S, Suija A, Crespo A, de los Ríos A (2014) Lichenicolous fungi of the genus *Abrothallus* (*Dothideomycetes: Abrothallales ordo nov.*) are sister to the predominantly aquatic *Jahnulales*. *Fungal Divers* 64:295–304
- Pérez-Ortega S, Garrido-Benavent I, de Los Ríos A (2015) *Austrostigmidium*, a new austral genus of lichenicolous fungi close to rock-inhabiting meristematic fungi in *Teratosphaeriaceae*. *Lichenologist* 47:143–156
- Pérez-Ortega S, Garrido-Benavent I, Grube M, Olmo R, de los Ríos A (2016) Hidden diversity of marine borderline lichens and a new order of fungi: *Collemopsidiales* (*Dothideomyceta*). *Fungal Divers* 80:285–300
- Pérez-Pérez RE, Castillo-Campos G, da Silva Cáceres ME (2015) Diversity of corticolous lichens in cloud forest remnants in La Cortadura, Coatepec, Veracruz, México in relation to phorophytes and habitat fragmentation. *Cryptogam Mycol* 36:79–92
- Pérez-Vargas I, Etayo J, Hernández-padrón C (2013) New species of lichenicolous fungi from the Canary Islands. *Phytotaxa* 99:58–64
- Pérez-Vargas I, Padrón CH, Pérez de Paz PL, Elix JA (2010a) *Tephromela follmannii* (lichenized *Ascomycota*), a new species from the Canary Islands. *Mycotaxon* 112:9–14
- Pérez-Vargas I, Padrón CH, Pérez de Paz PL, Elix JA (2010b) *Xanthoparmelia teydea*, a new brown *Xanthoparmelia* (*Parmeliaceae*) from the Canary Islands. *Bryologist* 113:51–54
- Perez-Vargas I, Hernández-Padrón C, De Paz PP, Van Den PP et al (2014) A new species in the lichen genus *Vahliella* from the Canary Islands, including a key to *Vahliellaceae, Pannariaceae*, and *Coccocarpiaceae* in Macaronesia. *Phytotaxa* 167:183–188
- Perić B, Lobuglio KF, Pfister DH (2013) The genus *Strobiloscypha*: a new species and an unresolved phylogenetic placement. *Mycol Montenegrina* 16:7–22
- Perić B, Van Vooren N, Healy R, Lazarević J (2014) Une trichophaea rare récoltée en france et au monténégró: *T. flavobrunnea* comb. nov. (*Pezizales*). *Mycol Montenegrina* 17:65–87
- Perić B, Baral HO, Pärtel K (2015) *Cenangiopsis raghavanii* and *C. junipericola* spp. nov. (*Cenangiaceae, Helotiales*) collected in Montenegro, with redescription of a recent collection of *C. quericola*. *Mycol Montenegrina* 18:7–40
- Perlmutter GB (2013) Lichen biotic inventory of turnipseed preserve, North Carolina, USA. *Evansia* 30:1–10
- Perlmutter GP, Lendemer JC, Guccion JC, Harris RC et al (2012) A provisional survey of lichen diversity in south-central South Carolina, USA, from the 19th Tuckerman lichen workshop. *Opusc Phil* 11:104–119
- Perlmutter GB, Tucker SC, Plata ER, Cerc P et al (2015) *Melaspilea demissa* (Tuck.) Zahlbr. (lichenized *Ascomycota*) in eastern North America with a key to North American species of *Melaspilea* s. lat. *Lichenologist* 47:167–182

- Peršoh D, Weig AR, Rambold G (2011) A transcriptome—targeting ecochip for assessing functional mycodiversity. *Microarrays* 1:25–41
- Persson M, Rasmussen M, Falk A, Dixielius C (2008) Barley mutants with enhanced level of resistance to Swedish isolates of *Bipolaris sorokiniana*, causal agent of spot blotch. *Plant Breeding* 127:639–643
- Persson M, Falk A, Dixielius C (2009) Studies on the mechanism of resistance to *bipolaris sorokiniana* in the barley lesion mimic mutant bst1. *Mol Plant Pathol* 10:587–598
- Petäistö RL, Lilja A, Hantula J (2013) Artificial infection and development of snow mold fungus (*Phacidium infestans*) in container-grown Norway spruce seedlings. *Balt For* 19:31–38
- Péter G, Dlauchy D, Tornai-Lehoczki J, Suzuki M, Kurtzman CP (2011) *Spencermartiniella europaea* gen. nov., sp. nov., a new member of the family *Trichomonascaceae*. *Int J Syst Evol Microbiol* 61:993–1000
- Péter G, Dlauchy D, Price NP, Kurtzman CP (2012) *Diddensiella caesifluorescens* gen. nov., sp. nov., a riboflavin-producing yeast species of the family *Trichomonascaceae*. *Int J Syst Evol Microbiol* 62:3081–3087
- Peter M, Kohler A, Ohm RA, Kuo A et al (2016) Ectomycorrhizal ecology is imprinted in the genome of the dominant symbiotic fungus *Cenococcum geophilum*. *Nat Commun* 7:12662. doi:[10.1038/ncomms12662](https://doi.org/10.1038/ncomms12662)
- Peterson SW (2008) Phylogenetic analysis of *Aspergillus* species using DNA sequences from four loci. *Mycologia* 100:205–226
- Peterson KR, Pfister DH (2010) Phylogeny of *Cytaria* inferred from nuclear and mitochondrial sequence and morphological data. *Mycologia* 102:1398–1416
- Peterson SW, Jurjevic Z, Bills GF, Stchigel AM et al (2010) Genus *Hamigera*, six new species and multilocus DNA sequence based phylogeny. *Mycologia* 102:847–864
- Peterson SW, Manitchoptisit P, Leathers TD (2013) *Aureobasidium thailandense* sp. nov. isolated from leaves and wooden surfaces. *Int J Syst Evol Microb* 63:790–795
- Pethybridge SJ, Jones SJ, Shivas RG, Hay FS, Wilson CR, Groom T (2008) Tan spot: a new disease of pyrethrum caused by *Microsphaeropsis tanaceti* sp. nov. *Plant Path* 57:1058–1065
- Petrini LE, Petrini O (2012) *Rosellinia* species (*Xylariaceae*) from South and Central America—an annotated list. *Kurtziana* 37:127–139
- Petterson OV, Leong SL, Lantz H, Rice T et al (2011) Phylogeny and intraspecific variation of the extreme xerophile, *Xeromyces bisporus*. *Fungal Biol* 115:1100–1111
- Pfister DH, Slater C, Hansen K (2008) *Chorioactidaceae*: a new family in the *Pezizales* (*Ascomycota*) with four genera. *Mycol Res* 112:513–527
- Pfister DH, Agnello C, Lantieri A, LoBuglio KF (2013) The *Caloscyphaceae* (*Pezizomycetes*, *Ascomycota*), with a new genus. *Mycol Prog* 12:667–674
- Pfliegler WP, Báthori F, Haelewaters D, Tartally A (2016a) Studies of *Laboulbeniales* on Myrmica ants (III): myrmecophilous arthropods as alternative hosts of *Rickia wasmannii*. *Parasite*. doi:[10.1051/parasite/2016060](https://doi.org/10.1051/parasite/2016060)
- Pfliegler WP, Fekete J, Báthori F, Tartally A (2016b) First Central European record of the fungus *Prolixandomyces triandrus* Santam., 1988 (*Ascomycota*: *Laboulbeniales*), a parasite of Veliid bugs (*Heteroptera*: *Veliidae*), with notes on its biology and DNA-barcoding. *Aquat Insects* 37:215–223
- Phengsintham P, Braun U, McKenzie EH, Chukeatirote E et al (2013) Monograph of cercospoid fungi from Thailand. *Plant Pathol Quar* 3:67–138
- Phillips AJ, Alves A, Pennycook SR, Johnston PR et al (2008) Resolving the phylogenetic and taxonomic status of dark-spored teleomorph genera in the *Botryosphaeriaceae*. *Persoonia* 21:29–55
- Phillips AJ, Alves A, Abdollahzadeh J, Slippers B et al (2013) The *Botryosphaeriaceae*: genera and species known from culture. *Stud Mycol* 76:51–167
- Phookamsak R, Hyde KD (2015) *Fenestellaceae*. *Mycosphere* 6:402–413
- Phookamsak R, Liu JK, Chukeatirote E, McKenzie EHC et al (2013) Convergent evolution: Phylogeny and morphology of *Leptosphaerulina saccharicola* sp. nov. and *Pleosphaerulina oryzae* and relationships with *Pithomyces*. *Cryptogam Mycol* 34:303–319
- Phookamsak R, Liu JK, Manamgoda DS, Chukeatirote E et al (2014a) The sexual state of *Setophoma*. *Phytotaxa* 176:260–269
- Phookamsak R, Liu JK, Manamgoda DS, Wanasinghe DN et al (2014b) Epitypification of two bambusicolous fungi from Thailand. *Cryptogam Mycol* 35:239–256
- Phookamsak R, Liu JK, McKenzie EH, Manamgoda DS et al (2014c) Revision of *Phaeosphaeriaceae*. *Fungal Divers* 68:159–238
- Phookamsak R, Manamgoda DS, Li WJ, Dai DQ et al (2015a) *Poaceascoma helicoides* gen et sp. nov., a new genus with scolecospores in *Lentitheciaceae*. *Cryptogam Mycol* 36:225–236
- Phookamsak R, Norphanphon C, Tanaka K, Dai DQ et al (2015b) *Astrosphaeriella*-like species; introducing *Astrosphaeriellaceae* and *Pseudoastrospphaeriellaceae* fam. nov. and *Astrosphaeriellopsis*, gen. nov. *Fungal Divers* 74:143–197
- Phookamsak R, Boonmee S, Norphanphon C, Wanasinghe DN et al (2016) *Schizophyriaceae*. *Mycosphere* 7:154–189
- Phukhamsakda C, Ariyawansa HA, Phookamsak R, Chomnunti P et al (2015) *Muriphaeosphaeria galatellae* gen. et sp. nov. in *Phaeosphaeriaceae* (*Pleosporales*). *Phytotaxa* 227:55–65
- Phukhamsakda C, Ariyawansa HA, Phillips AJ, Wanasinghe DN et al (2016a) Additions to *Sporormiaceae*: Introducing two novel genera, *Sparticola* and *Forliomyces*, from *Spartium*. *Cryptogam Mycol* 37:75–97
- Phukhamsakda C, Hongsanan S, Ryberg M, Ariyawansa HA et al (2016b) The evolution of *Massarinea* with *Longipedicellataceae* fam. nov. *Mycosphere* 7:1713–1731
- Phukhamsakda C, Bhat DJ, Hongsanan S, Tibpromma S et al (2017) *Magnicamarosporium diospyricola* sp. nov. (*Sulcatisporaceae*) from Thailand. *Mycosphere* 8:512–520
- Phuwapraisirisan P, Rangsan J, Siripong P, Tip-pyang S (2009) New antitumour fungal metabolites from *Alternaria porri*. *Nat Product Res* 23(12):1063–1071
- Pili NN, Franca SC, Kyndt T, Makumba BA et al (2015) Analysis of fungal endophytes associated with rice roots from irrigated and upland ecosystems in Kenya. *Plant Soil* 405:371–380
- Pindaru DM, Tanase C, Olariu RI, Arsene C (2013) Chemical composition and ions concentration in *Xanthoria parietina* and *Phaeophyscia orbicularis* lichenised fungi species from Iasi, North-Eastern Romania. *Revista de Chimie* 64:807–814
- Pindi PK (2012) Diversity of fungi at various depths of marine water. *Res Biotechnol* 3(4):62–66
- Pinho DB, Firmino AL, Pereira OL (2010) *Hyphoplynema ingae* sp. nov., associated with leaf spot disease on *Inga edulis* in Brazil. *Mycotaxon* 114:55–59
- Pinho DB, Honorato J, Nicoli A, Hora BT et al (2012) Phylogenetic placement of the genus *Anhelia* and the description of *A. nectandrae* sp. nov. *Mycologia* 104:1291–1298
- Pinho DB, Firmino AL, Ferreira-Junior WG, Pereira OL (2013) New *Meliolaceae* from the Brazilian Atlantic Forest 2: species on host families *Annonaceae*, *Cecropiaceae*, *Meliaceae*, *Piperaceae*, *Rubiaceae*, *Rutaceae* and *Tiliaceae*. *Mycologia* 105:697–711
- Pino-Bodas R, Zhurbenko MP, Stenroos S (2017) Phylogenetic placement within *Lecanoromycetes* of lichenicolous fungi

- associated with *Cladonia* and some other genera. Persoonia 39:91–117
- Pinnoi A, Phongpaichit P, Jeewon R, Tang AM et al (2010) Phylogenetic relationships of *Astrocytis eleiodoxae* sp. nov. (*Xylariaceae*). Mycosphere 1:1–9
- Pino-Bodas R, Martin MP, Burgaz AR (2012) *Cladonia subturgida* and *C. iberica* (*Cladoniaceae*) form a single, morphologically and chemically polymorphic species. Mycol Prog 11:269–278
- Pinruan U, Rungjindamai N, Sakayaroj J, Lumyong S et al (2010) *Baipadisphaeria* gen. nov., a freshwater ascomycete (*Hypocreales*, *Sordariomycetes*) from decaying palm leaves in Thailand. Mycosphere 1:53–63
- Pirnia M, Zare R, Zamanizadeh HR, Khodaparast A (2012) New records of cercosporoid hyphomycetes from Iran. Mycotaxon 120:157–169
- Pisani T, Munzi S, Paoli L, Baćkor M, Loppi S (2011) Physiological effects of arsenic in the lichen *Xanthoria parietina* (L.) Th. Fr. Chemosphere 82:963–969
- Pitt JI, Hocking AD (2009) Fungi and food spoilage. Springer, New York
- Pitt JI, Lantz H, Pettersson OV, Leong SL (2013) *Xerochrysium* gen. nov. and *Bettsia*, genera encompassing xerophilic species of *Chrysosporium*. IMA Fungus 4:229–241
- Pitt WM, Urbez-Torres JR, Trouillas FP (2014) *Munkovalsaria donacina* from grapevines and Desert Ash in Australia. Mycosphere 5:656–661
- Plattner A, Kim JJ, Reid J, Hausner G et al (2009) Resolving taxonomic and phylogenetic incongruence within species *Ceratocystiopsis minuta*. Mycologia 101:878–887
- Plaza CM, de Torres LE, Lücking RK, Vizcaya M, Medina GE (2014) Antioxidant activity, total phenols and flavonoids of lichens from Venezuelan Andes. J Pharm Pharm Res 2:138–147
- Poengsungnoen V, Manoch L, Mongkolsuk P, Kalb K (2014) New species of *Graphidaceae* from Loei Province, Thailand. Phytotaxa 189:255–267
- Pogány M, von Rad U, Grün S, Dongó A et al (2009) Dual roles of reactive oxygen species and NADPH oxidase RBOHD in an *Arabidopsis-Alternaria* pathosystem. Plant Physiol 151:1459–1475
- Polashock JJ, Caruso FL, Oudemans PV, McManus PS, Crouch JA (2009) The North American cranberry fruit rot fungal community: a systematic overview using morphological and phylogenetic affinities. Plant Pathol 58:1116–1127
- Pöldmaa K (2011) Tropical species of *Cladobotryum* and *Hypomyces* producing red pigments. Stud Mycol 68:1–34
- Poloni A, Pessi IS, Frazzon APG, van Der Sand ST (2009a) Morphology, physiology, and virulence of *Bipolaris sorokiniana* isolates. Curr Microbiol 59:267–273
- Poloni A, Pessi IS, Frazzon APG, van Der Sand ST (2009b) Vegetative incompatibility among monoconidial isolates of *Bipolaris sorokiniana*. Curr Microbiol 58:153–158
- Pose G, Patriarca A, Kyanko V, Pardo A, Pinto VF (2009) Effect of water activity and temperature on growth of *Alternaria alternata* on a synthetic tomato medium. Int J Food Microbiol 135:60–63
- Potenza G, Fascetti S, Ravera S, Puntillo D (2010) Lichens from sandy dune habitats on the Ionian Coast (Basilicata, southern Italy). Cryptog Mycol 31:59–65
- Prabhugaonkar A, Bhat DJ (2011) New record of *Megacapitula villosa* and *Paradicthyarthrinium diffractum* from India. Mycosphere 2:463–467
- Prasher IB, Verma RK (2014) *Taeniolina echinata* sp. nov. from North India. Kavaka 43:11–13
- Prasher IB, Verma RK (2015) *Neosporidesmium appendiculatus* sp. nov. from North-Western India. Mycol Prog 14:87–93
- Pratibha J (2013) *Jayarambhatia rhizophorae* gen. et sp. nov., an asexually reproducing fungus from Goa, India. Mycotaxon 125:139–144
- Pratibha J, Bhat DJ (2008) New and unusual hyphomycetes from Mahabaleshwar, India. Mycotaxon 105:423–431
- Pratibha J, Prabhugaonkar A (2015) Multi-gene phylogeny of *Pithomyces* with the sexual morph of *P. flavus* Berk. & Broome. Phytotaxa 218:84–90
- Pratibha J, Raghukumar S, Bhat DJ (2009) New species of *Digitoramispora* and *Spondylocladiopsis* from the forests of Western Ghats, India. Mycotaxon 107:383–390
- Pratibha J, Amandeep K, Shenoy BD, Bhat DJ (2010) *Caliciopsis indica* sp. nov. from India. Mycosphere 1:65–72
- Pratibha J, Bhat DJ, Raghukumar S (2011) Four anamorphic fungi (with two new species) from forests of Western Ghats, India. Mycotaxon 117:269–278
- Pratibha J, Prabhugaonkar A, Hyde KD, Bhat DJ (2014) Phylogenetic placement of *Bahuandhika*, *Cancellidium* and *Pseudoepicoccum* (asexual Ascomycota). Phytotaxa 176:68–80
- Pratibha J, Nguyen HD, Mel'nik VA, Bhat DJ et al (2015) Lectotyping, epitypification, and molecular phylogeny of the synnematous hyphomycete *Pseudogliophragma indicum*, the second genus in the Wiesneriomycetaceae. Mycoscience 56:387–395
- Pratibha J, Bhat DJ, Prabhugaonkar A (2016) Molecular phylogeny of *Speiropsis pedatospora*. Mycosphere 7:679–686
- Prelle A, Spadaro D, Garibaldi A, Gullino ML (2013) A new method for detection of five *Alternaria* toxins in food matrices based on LC-APCI-MS. Food Chem 140:161–167
- Prieto M, Wedin M (2016) Phylogeny, taxonomy and diversification events in the *Caliciaceae*. Fungal Divers 82:221–238
- Prieto M, Aragon G, Martinez I (2010a) The genus *Catapyrenium s. lat.* (*Verrucariaceae*) in the Iberian Peninsula and the Balearic Islands. Lichenologist 42:637–684
- Prieto M, Martinez I, Aragon G (2010b) The genus *Placiopsis* in the Iberian Peninsula. Mycotaxon 114:463–472
- Prieto M, Martinez I, Aragón G, Otálora MA, Lutzoni F (2010c) Phylogenetic study of *Catapyrenium s. str.* (*Verrucariaceae*, lichen-forming Ascomycota) and related genus *Placiopsis*. Mycologia 102:291–304
- Prieto M, Martínez I, Aragón G, Gueidan C, Lutzoni F (2012) Molecular phylogeny of *Heteroplacidium*, *Placidium*, and related catapyrenioid genera (*Verrucariaceae*, lichen-forming Ascomycota). Am J Bot 99:23–35
- Prieto M, Baloch E, Tehler M, Wedin M (2013) Mazaedium evolution in the *Ascomycota* (Fungi) and the classification of mazaediata groups of unclear relationship. Cladistics 29:296–308
- Prieto M, Westberg M, Schultz M (2015) New records of *Lichenomycetes* in Sweden and the Nordic countries. Herzogia 28:142–152
- Prihatini R, Boonyuen N, Sivichai S (2008) Phylogenetic evidence that two submerged-habitat fungal species, *Speiropsis pedatospora* and *Xylomyces chlamydosporus*, belong to the order *Jahnulales* insertae sedis *Dothideomycetes*. Microbiol Indonesia 2:136–140
- Printzen C (2010) Lichen systematics: the role of morphological and molecular data to reconstruct phylogenetic relationships. Prog Botany 71:233–275
- Printzen C (2014) A molecular phylogeny of the lichen genus *Biatora* including some morphologically similar species. Lichenologist 46:441–453
- Printzen C, Spribille T, Tønsberg T (2008) *Myochroidea*, a new genus of corticolous, crustose lichens to accommodate the *Lecidea leprosula* group. Lichenologist 40:195–207

- Printzen C, Cezanne R, Eichler M, Lumbsch HT (2012) The genera *Aphanopsis* and *Steinia* represent basal lineages within Leotiomyceta. *Biblioth Lichenol* 108:177–186
- Proksa B (2010) *Talaromyces flavus* and its metabolites. *Chem Pap* 64:696–714
- Puangsombat P, Sangwanit U, Marod D (2010) Diversity of soil fungi in different land use types in Tha Kum-Huai Raeng Forest Reserve, Trat Province. *Nat Sci* 44:1162–1175
- Pudhom K, Teerawatananond T (2014) Rhytidenes A–F, spirobisnaphthalenes from *Rhytidhysteron* sp. AS21B, an endophytic fungus. *J Nat Prod* 77:1962–1966
- Pundhir P, Tuda C, Vincentelli C, Morlote D et al (2016) *Scolecobasidium granulomatous* pneumonia and abscess—an emerging opportunistic fungal pathogen: a case report. *Int J STD AIDS* 0956462416646688
- Punithalingam E, Spooner BM (2011a) A new fungicolous *Scolecobasidium* (hyphomycetes) and *Caducirostrum* gen. nov. (coelomycetes) from leaf litter in the UK and Italy. *Kew Bull* 66:309–324
- Punithalingam E, Spooner BM (2011b) *Miricatena prunicola* (Hyphomycetes), a new genus and species causing leaf spots of *Prunus serotina* in the UK. *Kew Bull* 66:637–642
- Pykälä J, Mylllys L (2016) Three new species of Atla from calcareous rocks (*Verrucariaceae*, lichenized Ascomycota). *Lichenologist* 48:111–120
- Pykälä J, Stepanchikova IS, Himelbrant DE, Kuznetsova ES, Alexeeva NM (2012) The lichen genera *Thelidium* and *Verrucaria* in the Leningrad Region (Russia). *Folia Cryptogam Estonica* 49:45–57
- Qiao M, Zhang Y, Li SF, Baral HO et al (2012) *Orbilia blumenaviiensis* and its *Arthrobotrys* anamorph. *Micol Prog* 11:255–262
- Qin L, Qiao M, Yang Y, Yang GZ et al (2011) *Dactylella pseudobrevistipitata*, a new species from China. *Ann Microbiol* 61:591–595
- Qu CR, Yin GH, Zhao GZ, Cui BK et al (2014) Three new species of *Berkleasmium* (hyphomycetes) from China. *Nova Hedw* 98:151–161
- Quaedvlieg W, Kema GH, Groenewald JZ, Verkley GJ et al (2011) *Zymoseptoria* gen. nov.: a new genus to accommodate *Septoria*-like species occurring on graminicolous hosts. *Persoonia* 26:57–69
- Quaedvlieg W, Groenewald JZ, de Jesús Yáñez-Morales M et al (2012) DNA barcoding of *Mycosphaerella* species of quarantine importance to Europe. *Persoonia* 29:101–115
- Quaedvlieg W, Verkley GJ, Shin HD, Barreto RW et al (2013) Sizing up *Septoria*. *Stud Mycol* 75:307–390
- Quaedvlieg W, Binder M, Groenewald JZ, Summerell BA et al (2014) Introducing the consolidated species concept to resolve species in the *Teratosphaeriaceae*. *Persoonia* 33:1–40
- Quandt CA, Kepler RM, Gams W, Araújo JP et al (2014) Phylogenetic-based nomenclatural proposals for *Ophiocordycepsitaceae* (*Hypocreales*) with new combinations in *Tolyphocladium*. *IMA Fungus* 5:121–134
- Queloz V, Grünig CR, Berndt R, Kowalski T et al (2011) Cryptic speciation in *Hymenoscyphus albidos*. *For Pathol* 41:133–142
- Quijada L, Baral HO, Beltrán-Tejera E (2013) New species of *Orbilia* (*Orbiliales*) from arid ecosystems of the Canary Islands (Spain). *Nova Hedw* 96:237–248
- Quijada L, Baral HO, Jaén-Molina R, Weiss M et al (2015) Exploring the phylogenetic and morphological relationships of disregarded genera in *Helotiales* s.l. XVII Congress of European Mycologist (21–25 September, Madeira, Portugal)
- Quilhot W, Cuellar M, Diaz R, Riquelme F, Rubio C (2010) Preliminary study of the lichen flora of Isla Mocha, southern Chile. *Gayana Botánica* 67:206–212
- Raggio J, Pintado A, Ascaso C, De La Torre R et al (2011) Whole lichen thalli survive exposure to space conditions: results of Lithopanspermia experiment with *Aspicilia fruticulosa*. *Astrobiology* 11:281–292
- Raggio J, Green TGA, Crittenden PD, Pintado A et al (2012) Comparative ecophysiology of three *Placopsis* species, pioneer lichens in recently exposed Chilean glacial forelands. *Symbiosis* 56:55–66
- Raitivir A (2008) The *Helotiales* of the Magadan and Chukotka areas of the Russian Arctic. – *Sommerfeltia* 31:179–190
- Raja HA, Shearer CA (2008) Freshwater ascomycetes: new and noteworthy species from aquatic habitats in Florida. *Mycologia* 100:467–489
- Raja HA, Carter A, Platt HW, Shearer CA (2008a) Freshwater ascomycetes: *Jahnula apiospora* (*Jahnulales, Dothideomycetes*), a new species from Prince Edward Island, Canada. *Mycoscience* 49:326–328
- Raja HA, Miller AN, Shearer CA (2008b) Freshwater ascomycetes: *Aquaporiterium pinicola*, a new genus and species of *Helotiales* (*Leotiomycetes*) from Florida. *Mycologia* 100:141–148
- Raja HA, Ferrer A, Shearer CA (2009a) Freshwater ascomycetes: a new genus, *Ocala scalariformis* gen. et sp nov., and two new species, *Ayria nubispora* sp nov and *Rivulicola cygnea* sp nov. *Fungal Divers* 34:79–86
- Raja HA, Schmit JP, Shearer CA (2009b) Latitudinal, habitat and substrate distribution patterns of freshwater ascomycetes in the Florida Peninsula. *Biodivers Conserv* 18:419–455
- Raja H, Miller A, Shearer CA (2010a) Freshwater ascomycetes: *Hyalorostratum brunneisporum*, a new genus and species in the *Diaporthales* (*Sordariomycetidae, Sordariomycetes*) from North America. *Mycosphere* 1:275–288
- Raja HA, Ferrer A, Shearer CA, Miller AN (2010b) Freshwater ascomycetes: *Wicklowia aquatica*, a new genus and species in the *Pleosporales* from Florida and Costa Rica. *Mycoscience* 51:208–214
- Raja HA, Violi HA, Shearer CA (2010c) Freshwater ascomycetes: *Alascospora evergladensis*, a new genus and species from the Florida Everglades. *Mycologia* 102:33–38
- Raja HA, Schoch C, Hustad V, Shearer C, Miller A (2011a) Testing the phylogenetic utility of MCM7 in the *Ascomycota*. *MycoKeys* 1:63–94
- Raja HA, Tanaka K, Hirayama K, Miller A, Shearer CA (2011b) Freshwater Ascomycetes: two new species of *Lindgomycetes* (*Lindgomycetaceae, Pleosporales, Dothideomycetes*) from Japan and USA. *Mycologia* 103:1421–1432
- Raja HA, Miller AN, Shearer CA (2012) Freshwater ascomycetes: *Natipusillaceae*, a new family of tropical fungi, including *Natipusilla bellaspore* sp. nov. from the Peruvian Amazon. *Mycologia* 104:569–573
- Raja HA, Oberlies NH, El-Elimat T, Miller AN et al (2013a) *Lindgomycetes angustiascus*, (*Lindgomycetaceae, Pleosporales, Dothideomycetes*), a new lignicolous species from freshwater habitats in the USA. *Mycoscience* 54:353–361
- Raja HA, Oberlies NH, Figueroa M, Tanaka K et al (2013b) Freshwater ascomycetes: *Minutisphaera* (*Dothideomycetes*) revisited, including one new species from Japan. *Mycologia* 105:959–976
- Raja HA, Oberlies NH, Elimat Tamam-El, Miller AN et al (2013c) *Lindgomycetes angustiascus* (*Lindgomycetaceae, Pleosporales, Dothideomycetes*), a new lignicolous species from freshwater habitats in the USA. *Mycoscience* 54:353–361
- Raja HA, El-Elimat T, Oberlies NH, Shearer CA et al (2015) *Minutisphaerales* (*Dothideomycetes, Ascomycota*): a new order of freshwater ascomycetes including a new family, *Minutisphaeraceae*, and two new species from North Carolina, USA. *Mycologia* 107:845–862

- Raja HA, Paguigan ND, Fournier J, Oberlies NH (2017) Additions to *Lindgomyces* (*Lindgomycetaceae*, *Pleosporales*, *Dothideomycetes*), including two new species occurring on submerged wood from North Carolina, USA, with notes on secondary metabolite profiles. *Mycol Prog* 16:535–552
- Rajakaruna N, Knudsen K, O'dell R, Nathaniel P et al (2012) Investigation of the importance of rock chemistry for saxicolous lichen communities of the New Idria serpentinite mass, San Benito County, California, USA. *Lichenologist* 44:695–714
- Rajeshkumar KC, Singh SK (2012) *Manoharachariella indica* sp. nov. from the Western Ghats, India. *Mycotaxon* 120:43–48
- Rajeshkumar KC, Rossman AY (2013) *Taiwanascus samuelsii* sp. nov., an addition to *Niessliaceae* from the Western Ghats, Kerala, India. *IMA Fungus* 4:1–4
- Rajeshkumar KC, Sharma R (2013) *Tamhinispora* a new genus belongs to family *Tubeufiaceae* from the Western Ghats, India based on morphology and phylogenetic analysis. *Mycosphere* 4:165–174
- Rajeshkumar KC, Kajale SC, Sutar SA, Singh SK (2013) *Ellisembia karadkensis* sp. nov. from southern Western Ghats, India. *Mycotaxon* 121:181–186
- Rajeshkumar KC, Crous PW, Groenewald JZ, Seifert KA (2016) Resolving the phylogenetic placement of *Porobeltraniella* and allied genera in the *Beltraniaceae*. *Mycol Prog* 15:1119–1136
- Ram TJ (2014) New species and new records in *Roccellaceae* (*Arthoniales*) from the Andaman Islands, India. *Phytotaxa* 177:155–162
- Rämä T, Mathiassen GH, Kauserud H (2014) Marine fungi new to Norway, with an outlook to the overall diversity. *Agarica* 34:35–54
- Rambelli A (2011) Some dematiaceous hyphomycetes from Mediterranean maquis litters. *Flora Mediterranea* 21:5–204
- Ramos AM, Cinto IE, Tadic LF, Ranalli ME (2011) AFLP molecular characterizations of some *Saccobolus* species (Ascomycota, Pezizales). *Mycotaxon* 116:355–364
- Randlane T, Saag A, Thell A, Ahti T (2013) Third world list of cetrarioid lichens—in a new databased form, with amended phylogenetic and type information. *Cryptog Mycol* 34:79–94
- Randriamanantany ZA, Annesi-Maesano I, Moreau D, Raherison C et al (2010) *Alternaria* sensitization and allergic rhinitis with or without asthma in the French Six Cities study. *Allergy* 65:368–375
- Rangsiruij A, Boonpragob K, Mongkolsuk P, Sodamuk M et al (2016) Diversity and phylogenetic survey of cyanobacterial lichens (Collematinae, Ascomycota) in mangrove forests of eastern Thailand. *Bryologist* 119:123–130
- Ranjbar-Bahadori S, Razzaghi-Abyaneh M, Bayat M, Eslami A et al (2010) Studies on the effect of temperature, incubation time and in vivo gut passage on survival and nematophagous activity *Arthrobotrys oligospora* var. *oligospora* and *A. cladodes* var. *macroides*. *Glob Vet* 4:112–117
- Ranković B, Mišić M, Sukdolak S (2009) Antimicrobial activity of extracts of the lichens *Cladonia furcata*, *Parmelia caperata*, *Parmelia pertusa*, *Hypogymnia physodes* and *Umbilicaria polyphylla*. *Biologija* 64:53–58
- Ranković BR, Kosančić MM, Stanojković TP (2011) Antioxidant, antimicrobial and anticancer activity of the lichens *Cladonia furcata*, *Lecanora atra* and *Lecanora muralis*. *BMC Complement Altern Med* 11:97–105
- Ranković BV, Kosančić MA, Stanojković TA (2014) *Stereocaulon paschale* lichen as antioxidant, antimicrobial and anticancer agent. *Farmacia* 62:306–317
- Rashmi S, Rajkumar HG (2015) First report of foliicolous lichen biota in South Karnataka-India. *Int J Curr Microbiol App Sci* 4:250–256
- Réblová M (2008) *Bellojisia* a new sordariaceous genus for *Jobellisia rhynchostoma* and a description of *Jobellisiaceae* fam. nov. *Mycologia* 100:893–901
- Réblová M (2011) New insights into the systematics and phylogeny of the genus *Jattaea* and similar fungi of the *Calosphaerales*. *Fungal Divers* 49:167–198
- Réblová M (2013) Two taxonomic novelties in the *Sordariomycetidae*: *Ceratolenta caudata* gen. et sp. nov. and *Platytrachelon abietis* gen. et comb. nov. for *Ceratosphaeria abietis*. *Mycologia* 105:462–475
- Réblová M (2014) *Sporoschismopsis angustata* sp. nov., a new holomorph species in the *Retiulascaceae* (*Glomerellales*), and a reappraisal of *Sporoschismopsis*. *Mycol Prog* 13:671–681
- Réblová M, Štěpánek V (2009) New fungal genera, *Tectonidula* gen. nov. for Calosphaeria-like fungi with holoblastic-denticulate conidiogenesis and *Natantiella* gen. nov. for three species segregated from *Ceratostomella*. *Mycol Res* 113:991–1002
- Réblová M, Gams W (2011) The new hyphomycete genera *Brachyalara* and *Infundichalara*, the similar *Exochalara* and species of 'Phialophora' sect. *Catenulatae*' (Leotiomycetes). *Fungal Divers* 46:67–86
- Réblová M, Seifert KA (2011) Discovery of the teleomorph of the hyphomycete, *Sterigmatobotrys macrocarpa*, and epitypification of the genus to holomorphic status. *Stud Mycol* 68:193–202
- Réblová M, Seifert KA (2012) *Cirrosporium novae-zelandiae*, an enigmatic coelomycete with meristem arthroconidia, with ancestors in the *Eurotiomycetes*. *Mycologia* 104:1315–1324
- Réblová M, Fournier J, Hyde KD (2010) *Achroceratosphaeria*, a new ascomycete genus in the *Sordariomycetes*, and re-evaluation of *Ceratosphaeria incolorata*. *Fungal Divers* 43:75–84
- Réblová M, Gams W, Seifert KA (2011) *Monilochaetes* and allied genera of the *Glomerellales*, and a reconsideration of families in the *Microascales*. *Stud Mycol* 68:163–191
- Réblová M, Seifert KA, Fournier J, Štěpánek V (2012) Phylogenetic classification of *Pleurothecium* and *Pleurotheciella* gen. nov. and its dactylaria-like anamorph (*Sordariomycetes*) based on nuclear ribosomal and protein-coding genes. *Mycologia* 104:1299–1314
- Réblová M, Untereiner WA, Réblová K (2013) Novel evolutionary lineages revealed in the *Chaetothyriales* (Fungi) based on multigene phylogenetic analyses and comparison of ITS secondary structures. *PloS ONE* 8:e.0063547
- Réblová M, Štěpánek V, Schumacher RK (2014) *Xylochrysis lucidagen*. et sp. nov., a new lignicolous ascomycete (*Sordariomycetidae*) with holoblastic conidiogenesis. *Mycologia* 106:564–572
- Réblová M, Fournier J, Štěpánek V (2015a) *Pisorisporiales*, a new order of aquatic and terrestrial fungi for *Achroceratosphaeria* and *Pisorisporium* gen. nov. in the *Sordariomycetes*. *Persoonia* 34:40–49
- Réblová M, Jaklitsch W, Réblová K, Štěpánek V (2015b) Phylogenetic reconstruction of the *Calosphaerales* and *Togniniales* using five genes and predicted RNA secondary structures of ITS, and *Flabellascus tenuirostris* gen. et sp. nov. *Plos ONE* 10:e.0144616
- Réblová M, Réblová K, Štěpánek V (2015c) Molecular systematics of *Barbatosphaeria* (*Sordariomycetes*): multigene phylogeny and secondary ITS structure. *Persoonia* 35:21–38
- Réblová M, Fournier J, Štěpánek V (2016a) Two new lineages of aquatic ascomycetes: *Atractospora* gen. nov. and *Rubellisphaeria* gen. et sp. nov., and a sexual morph of *Myrmecridium montsegurinum* sp. nov. *Mycol Prog* 15:1–18
- Réblová M, Hubka V, Thureborn O, Lundberg J et al (2016b) From the tunnels into the treetops: new lineages of black yeasts from biofilm in the Stockholm metro system and their relatives among

- ant-associated fungi in the *Chaetothyriales*. PLoS ONE 11:e0163396
- Réblová M, Miller AN, Rossman AY, Seifert KA et al (2016c) Recommendations for competing sexual—asexually typified generic names in *Sordariomycetes* (except *Diaporthales*, *Hypocreales*, and *Magnaportheales*). IMA Fungus 7:131–153
- Réblová M, Seifert KA, Fournier J, Štěpánek V (2017) Newly recognised lineages of perithecial ascomycetes: the new orders *Conioscyphales* and *Pleurotheciales*. Persoonia 37:57–81
- Recio M, del Mar Trigo M, Docampo S, Melgar M et al (2012) Analysis of the predicting variables for daily and weekly fluctuations of two airborne fungal spores: *Alternaria* and *Cladosporium*. Int J Biometeorol 56:983–991
- Reddy MA, Nayaka S, Shankar PC, Reddy SR, Rao BR (2011) New distributional records and checklist of lichens for Andhra Pradesh, India. Indian Forester 137:1371–1376
- Redhead SA, Demoulin V, Hawksworth DL, Seifert KA, Turland NJ (2014) Fungal nomenclature at IMC10: Report of the Nomenclature Sessions. IMA Fungus 5:449–462
- Rehner SA, Minnis AM, Sung GH, Luangsa-ard JJ et al (2011) Phylogeny and systematics of the anamorphic, entomopathogenic genus *Beauveria*. Mycologia 103:1055–1073
- Ren Q, Li S (2013) New records of crustose lichens from China-1. Mycotaxon 125:65–67
- Ren F, Zhuang WY (2014) A new species of the genus *Chlorencoelia* (*Helotiales*) from China. Mycoscience 55:227–230
- Ren SC, Ma J, Ma LG, Zhang YD, Zhang XG (2012a) *Sativumoides* and *Cladosporiopsis*, two new genera of hyphomycetes from China. Mycol Prog 11:443–448
- Ren SC, Ma J, Zhang XG (2012b) Two new *Heteroconium* species and two other forest microfungi newly recorded from China. Mycotaxon 119:361–367
- Ren J, Jie CY, Zhou QX, Li XH et al (2013a) Molecular and morphological data reveal two new species of *Scolecobasidium*. Mycoscience 54:420–425
- Ren J, Zeng C, Jiang YL, Hyde KD, Wang Y (2013b) A new Thai species of *Astrosphaeriella* (*Dothideomycetes*, *Ascomycota*) from submerged wood in freshwater. Sydowia 65:33–43
- Ren SC, Ma J, Zhang XG (2013c) Two new *Ellisembia* species from Hainan and Yunnan, China. Mycotaxon 122:83–87
- Resl P, Schneider K, Westberg M, Printzen C et al (2015) Diagnostics for a troubled backbone: testing topological hypotheses of trapezioid lichenized fungi in a large-scale phylogeny of *Ostropomyctidae* (*Lecanoromycetes*). Fungal Divers 73:239–258
- Resl P, Mayrhofer H, Clayden SR, Spribille T et al (2016) Morphological, chemical and species delimitation analyses provide new taxonomic insights into two groups of *Rinodina*. Lichenologist 48:469–488
- Reynolds DR (2010) Epifoliar fungi of Singapore. Gdns' Bull Singapore 61:401–435
- Richardson MJ (2008) Records of coprophilous fungi from the Lesser Antilles and Puerto Rico. Carrib J Sci 44:206–214
- Riddell J, Jovan S, Padgett P, Sweat K (2011) Tracking lichen community composition changes due to declining air quality over the last century: the Nash legacy in Southern California. Biblio Lichenol 106:263–277
- Rifai MA (2008) Another note on *Podoconis megasperma* Boedijn (Hyphomycetes). Reinwardtia 12:277–279
- Rikkinen J, Tuovila H, Beimforde C, Seyfullah L et al (2014) *Chaenothecopsis neocaledonica* sp. nov.: the first resinicolous mycocalicioid fungus from an araucarian conifer. Phytotaxa 173:49–60
- Rincon-Espitia A, Mateus N (2013) Morphological and anatomical characterization of the genus *Stereocaulon hoffmann* (Lichenized ascomycetes) in Colombia. Caldasia 35:241–260
- Rivas Plata E, Lücking R (2012) High diversity of *Graphidaceae* (lichenized *Ascomycota*: *Ostropales*) in Amazonian Perú. Fungal Divers 58:13–32
- Rivas Plata ET, Lücking R, Lumbsch HT, Chaves JL (2008) *Gyrotrema wirthii*, a new species of *Gyrotrema* (*Ascomycota*: *Ostropales*: *Graphidaceae*) from Costa Rica. Sauteria 15:417–420
- Rivas Plata E, Lücking R, Sipman HJM, Mangold A et al (2010a) A world-wide key to the thelotremoid *Graphidaceae*, excluding the *Ocellularia*-*Myriotrema*-*Stegobolus* clade. Lichenologist 42:139–185
- Rivas Plata E, Kalb K, Frisch A (2010b) *Wirthiotrema*: a new genus for the *Thelotrema glaucopallens* group (*Ascomycota*: *Ostropales*: thelotremoid *Graphidaceae*). Lichenologist 42:187–202
- Rivas Plata E, Lücking R, Lumbsch HT (2012a) A new classification for the family *Graphidaceae* (*Ascomycota*: *Lecanoromycetes*: *Ostropales*). Fungal Divers 52:107–121
- Rivas Plata E, Lücking R, Lumbsch HT (2012b) Molecular phylogeny and systematics of the *Ocellularia* clade (*Ascomycota*: *Ostropales*: *Graphidaceae*). Taxon 61:1161–1179
- Rivas Plata E, Parmen S, Staiger B, Mangold A et al (2013) A molecular phylogeny of *Graphidaceae* (*Ascomycota*, *Lecanoromycetes*, *Ostropales*) including 428 species. Myco Keys 6:55–94
- Rivera-Orduña FN, Suarez-Sánchez RA, Flores-Bustamante ZR, Gracida-Rodríguez JN, Flores-Cotera LB (2011) Diversity of endophytic fungi of *Taxus globosa* (Mexican yew). Fungal Divers 47:65–74
- Rizzi G, Giordan P (2013) The ecology of the lichen genus *Xanthoparmelia* in Italy: an investigation throughout spatial scales. Plant Biosyst 147:33–39
- Rizzi G, Incerti G, Ginaldi F, Kodnik D et al (2011) A contribution to the lichen flora of Sardinia. Mycotaxon 115:535
- Rizzo L, Sutton DA, Wiederhold NP, Thompson EH, Friedman R et al (2014) Isolation and characterisation of the fungus *Spiromastix asexualis* sp. nov. from discospondylitis in a German Shepherd dog, and review of *Spiromastix* with the proposal of the new order *Spiromastixales* (*Ascomycota*). Mycoses 57:419–428
- Robène-Soustrade I, Jouen E, Pastou D, Payet-Hoarau M et al (2015) Description and phylogenetic placement of *Beauveria hoplocheli* sp. nov. used in the biological control of the sugarcane white grub, *Hoplochelus marginalis*, on Reunion Island. Mycologia 107:1221–1232
- Rocha FB, Barreto RW, Bezerra JL, Neto JAAM (2010) Foliar mycobiota of *Coussapoa floccosa*, a highly threatened tree of the Brazilian Atlantic forest. Mycologia 102:1240–1252
- Rodriguez Flakus PR, Printzen C (2014) Molecular evidence for the occurrence of the lichen genus *Biatora* (*Lecanorales*, *Ascomycota*) in the Southern Hemisphere. Phytotaxa 172:271–279
- Rodrigues AM, de Hoog GS, De Camargo ZP (2015) Molecular diagnosis of pathogenic *Sporothrix* species. PLoS Negl Tropical Dis 9:e0004190
- Rodríguez JD, Kirschner R, Piepenbring M (2015) New species and new records of *Meliolaceae* from Panama. Fungal Divers 70:73–84
- Rodrigues AM, Cruz-Choappa R, Fernandes GF, de Hoog GS, de Camargo ZP (2016) *Sporothrix chilensis* sp. nov. (*Ascomycota*: *Ophiostomatales*), a soil-borne agent of human sporotrichosis with mild pathogenic potential to mammals. Fungal Biol 120:246–264
- Rogers JD, Ju YM (2012) The *Xylariaceae* of the Hawaiian Islands. North Am Fungi 7:1–35

- Rogers PC, Rogers RW, Hedrich AE, Moss PT (2015) Lichen monitoring delineates biodiversity on a Great Barrier Reef Coral Cay. *Forests* 6:1557–1575
- Rojas EI, Herre EA, Mejía LC, Arnold AE, Chaverri P, Samuels GJ (2008) *Endomelanconiopsis*, a new anamorph genus in the *Botryosphaeriales*. *Mycologia* 100:760–775
- Romero AI, Robledo G, LoBuglio KF, Pfister DH (2012) *Rickiella edulis* and its phylogenetic relationships within *Sarcoscyphaceae*. *Kurtziana* 37:79–89
- Romón P, De Beer ZW, Fernández M, Diez J, Wingfield BD, Wingfield MJ (2014) Ophiostomatoid fungi including two new fungal species associated with pine root-feeding beetles in Northern Spain. *Antonie Van Leeuwenhoek* 106:1167–1184
- Root HT, McCune B (2012) Regional patterns of biological soil crust lichen species composition related to vegetation, soils, and climate in Oregon, USA. *J Arid Env* 79:93–100
- Root HT, Miller JE, McCune B (2011) Biotic soil crust lichen diversity and conservation in shrub-steppe habitats of Oregon and Washington. *Bryologist* 114:796–812
- Root HT, McCune M, McCune B (2013) Wind farm potential is higher in prime habitat for uncommon soil crust lichens. *Ecol Process* 2:10. doi:[10.1186/2192-1709-2-10](https://doi.org/10.1186/2192-1709-2-10)
- Root HT, McCune B, Jovan S (2014) Lichen communities and species indicate climate thresholds in southeast and south-central Alaska, USA. *Bryologist* 117:241–252
- Ropars J, Cruaud C, Lacoste S, Dupont J (2012) A taxonomic and ecological overview of cheese fungi. *Int J Food Microbiol* 155:199–210
- Rosentreter R, Rai H, Upreti DK (2014) Distribution ecology of soil crust lichens in India: a comparative assessment with global patterns. *Terricolous lichens in India*. Springer, New York, pp 21–31
- Rosenzweig N, Olaya G, Atallah ZK, Cleere S et al (2008) Monitoring and tracking changes in sensitivity to azoxystrobin fungicide in *Alternaria solani* in Wisconsin. *Plant Dis* 92:555–560
- Rosling A, Cox F, Cruz-Martinez K, Ihrmark K et al (2011) *Archaeorhizomycetes*: unearthing an ancient class of ubiquitous soil fungi. *Science* 333:876–879
- Rosow L, Jiang JX, Deuel T, Lechpammer M et al (2011) Cerebral phaeohyphomycosis caused by *Bipolaris spicifera* after heart transplantation. *Transp Infect Dis* 13:419–423
- Rossi W (2010) New *Laboulbeniales* (Ascomycota) parasitic on Staphylinidae from Ecuador. *Mycol Prog* 9:407–415
- Rossi W, Bergonzo E (2008) New and interesting *Laboulbeniales* from Brazil. *Aliso* 26:1–8
- Rossi W, Ponziani PR (2008) New *Laboulbeniales* (Ascomycotina) from French Guyana. *Cryptog Mycol* 29:3–9
- Rossi W, Santamaría S (2008) New *Laboulbeniales* parasitic on endogeal ground beetles. *Mycologia* 100:636–641
- Rossi W, Santamaría S (2008) *Cesariella*, a new genus of *Laboulbeniales*. *Mycol Res* 112:917–920
- Rossi W, Weir A (2008) New Species of *Cucujomyces* (*Laboulbeniales*) on Chilean Leiodidae. *Aliso* 26:9–14
- Rossi W, Proano Castro AC (2009) New species of *Rhachomyces* from Ecuador, one of which is dimorphic. *Mycologia* 101:674–680
- Rossi W, Santamaría S (2010) New species of *Diphymyces* (*Laboulbeniales*, Ascomycota). *Mycol Prog* 9:597–601
- Rossi W, Weir A (2011) New species of *Stigmatomyces* from Asia. *Mycologia* 103:131–134
- Rossi W, Leonardi M (2012) New species of *Stigmatomyces* (*Laboulbeniomycetes*) from Sierra Leone. *Plant Biosyst* 147:79–83
- Rossi W, Santamaría S (2012) *Rodoucea*, a new genus of the *Laboulbeniales*. *Mycologia* 104:785–788
- Rossi W, Santamaría S (2015) New species of *Aporomyces*. *Turkish J Bot* 39:354–358
- Rossi W, Bernardi M, Torres JA (2015) New species of *Dimeromyces* from Ecuador. *Mycol Prog* 14:5. doi:[10.1007/s11557-015-1026-2](https://doi.org/10.1007/s11557-015-1026-2)
- Rossi W, Bernardi M, Torres JA (2016a) New species of *Laboulbenia* parasitic on leaf beetles. *Mycol Prog* 15:1–7
- Rossi W, Haelewaters D, Pfister DH (2016b) Fireworks under the microscope: a spectacular new species of *Zodiomyces* from the Thaxter collection. *Mycologia* 108:709–715
- Rossman AY, Schoch CL, Farr DF, Nishijima K et al (2010) *Dolabra nepheliae* on rambutan and lychee represents a novel lineage of phytopathogenic *Eurotiomycetes*. *Mycoscience* 51:300–309
- Rossman AY, Melgar J, Walker D, Gonzales A, Ramirez T, Rivera J (2012) First report of *Dolabra nepheliae* associated with corky bark disease of rambutan and pulasan in Honduras. *Plant Dis* 96:765
- Rossman AY, Manamgoda DS, Hyde KD (2013a) (2233) Proposal to conserve the name *Bipolaris* against *Cochliobolus* (Ascomycota: Pleosporales: Pleosporaceae). *Taxon* 62:1331–1332
- Rossman AY, Seifert KA, Samuels GJ, Minnis AM et al (2013b) Genera in *Bionectriaceae*, *Hypocreaceae*, and *Nectriaceae* (Hypocreales) proposed for acceptance or rejection. *IMA Fungus* 4:41–51
- Rossman AY, Adams GC, Cannon PF, Castlebury LA et al (2015a) Recommendations of generic names in *Diaporthales* competing for protection or use. *IMA Fungus* 6:145–154
- Rossman AY, Crous PW, Hyde KD, Hawksworth DL et al (2015b) Recommended names for pleomorphic genera in *Dothideomycetes*. *IMA Fungus* 6:507–523
- Rossman AY, Allen WC, Braun U, Castlebury LA et al (2016a) Overlooked competing asexual and sexually typified generic names of *Ascomycota* with recommendations for their use or protection. *IMA Fungus* 7:285–304
- Rossman AY, Allen WC, Castlebury LA (2016b) New combinations of plant-associated fungi resulting from the change to one name for fungi. *IMA Fungus* 7:1–7
- Rossmann B, Muller H, Smalla K, Mpiira S et al (2012) Banana-associated microbial communities in Uganda are highly diverse but dominated by enterobacteriaceae. *Appl Environ Microbiol* 78:4933–4941
- Rotter M (2010) A preliminary list of the lichen flora of the San Francisco Presidio. *Bull California Lichen Soc* 17:23–26
- Rout J, Singha AB, Upreti DK (2010) Pigment profile and chlorophyll degradation of *Pyxine cocoes* lichen: a comparative study of the different degree of disturbance in Cachar District, Assam. *Assam Uni J Sci Technol* 5:85–88
- Roux C (2009) *Naetrocymbae saxicola*, likeno kun *Trentepohlia*. *Bull Soc Linn Provence* 60:127–142
- Roux C, Gueidan C (2011) Du novaj specioj de Verrucariaceae el Pyrénées-Orientales (Farncio): *Placocarpus melanophthalmosus* sp. nov. kaj *Placopyrenium breussii* sp. nov. *Bull de la Soc Linn Provence* 14:163–176
- Roux C, Gueidan C, Navarro-Rosinés P, Monnat JY (2016) Encore Trimmatothelopsis! *Bull Soc Linn Provence* 67:159
- Rubini A, Belfiori B, Passeri V, Falini LB et al (2011) The AD-type ectomycorrhizas, one of the most common morphotypes present in truffle fields, result from fungi belonging to the *Trichophaea woolhopeia* species complex. *Mycorrhiza* 21:17–25
- Rubio E, Tena R, Ormad J, Suárez A (2010) *Ruhlandiella reticulata* comb. nov. y *Ruhlandiella truncata* comb. nov. (Ascomycota, Pezizales) Nuevas combinaciones para dos raras especies semihipogea, eucaliptícolas y pirófilas de origen austral: *Muciturbus reticulatus* y *Muciturbus truncatus*. *Revista Catalana de Micología* 32:23–30

- Rubio E, Tabarés M, Martínez MA (2011) *Marcelleina parvispora* (Ascomycota, Pezizales), a new *Marcelleina* species from Catalonia (Spain). Rev Catalana de Micol 32:31–35
- Rubio C, Saavedra M, Cuéllar M, Díaz R, Quilhot W (2013) Epiphytic lichens of Conguillío National Park, southern Chile/Líquenes epífitos en el Parque Nacional Conguillío, sur de Chile. Gayana. Botanica 70:66
- Ruibal C, Platas G, Bills GF (2008) High diversity and morphological convergence among melanised fungi from rock formations in the Central Mountain System of Spain. Persoonia 21:93–110
- Ruibal C, Gueidan C, Selbmann L, Gorbushina AA et al (2009) Phylogeny of rock-inhabiting fungi related to *Dothideomycetes*. Stud Mycol 64:123–133
- Ruibal R, Millanes AM, Hawksworth DL (2011) Molecular phylogenetic studies on the lichenicolous *Xanthoriicola physciæ* reveal Antarctic rock-inhabiting fungi and *Piedraia* species among closest relatives in the *Teratosphaeriaceae*. IMA Fungus 2:97–103
- Ruiz RFC, Iturriaga T, Abarca GH, Minter DW et al (2008) Notes on *Heteroconium* and a new species from Venezuela. Mycotaxon 105:175–184
- Ruiz-González MX, Lauth J, Leroy C, Jauneau A et al (2013) An efficient protocol for isolating melanised chaetothyrialean anamorphic fungi associated with plant-ants. J Basic Microbiol 53:98–100
- Rungindamai N, Sakayaroj S, Somrithipol S, Plaingam N, Jones EBG (2012) Phylogeny of the appendaged coelomycete genera: *Pseudorobillarda*, *Robillarda*, and *Xepiculopsis* based on nuclear ribosomal DNA sequences. Cryptog Mycol 33:319–332
- Ruszkievicz-Michalska M, Połec E (2014) Additions to *Ramularia* species (hyphomycetes) in Poland. Mycotaxon 127:63–72
- Saag A, Randlane T, Saag L, Thell A, Ahti T (2013) Third world list of cetrarioid lichens, a databased tool for documentation of nomenclatural data—lessons learned. Taxon 62:591–603
- Saag L, Mark K, Saag A, Randlane T (2014) Species delimitation in the lichenized fungal genus *Vulpicida* (Parmeliaceae, Ascomycota) using gene concatenation and coalescent-based species tree approaches. American J Bot 101:2169–2182
- Sabirov A, Hamilton RG, Jacobs JB, Hillman DE et al (2008) Role of local immunoglobulin E specific for *Alternaria alternata* in the pathogenesis of nasal polyposis. Laryngoscope 118:4–9
- Saccente M, Woods GL (2010) Clinical and laboratory update on *Blastomycosis*. Clin Microbiol Rev 23:367–381
- Saha S, Sarkar J, Chattopadhyay D, Patra S et al (2010) Production of silver nanoparticles by a phytopathogenic fungus *Bipolaris nodulosa* and its antimicrobial activity. Dig J Nanomater Biostruct 5:887–895
- Saha D, Fetzner R, Burkhardt B, Podlech J et al (2012) Identification of a polyketide synthase required for alternariol (AOH) and alternariol-9-methyl ether (AME) formation in *Alternaria alternata*. PLoS One 7:e40564
- Sakayaroj J, Pang KL, Jones EBG (2011a) Multi-gene phylogeny of the *Halosphaeriaceae*: its ordinal status, relationships between genera and morphological character evolution. Fungal Divers 46:87–109
- Sakayaroj J, Supaphon O, Jones EBG, Phongpaichit S (2011b) Diversity of higher marine fungi at Hat Khanom-Mu Ko Thale Tai National Park, Southern Thailand. Songklanakarin J Sci Technol 33:15–22
- Salgado-Salazar C, Rossman A, Samuels GJ, Capdet M, Chaverri P (2012) Multigene phylogenetic analyses of the *Thelonectria coronata* and *T. veuillotiana* species complexes. Mycologia 104:1325–1350
- Salgado-Salazar C, Rossman AY, Samuels GJ, Hirooka Y et al (2014) Phylogeny and taxonomic revision of *Thelonectria discophora* (Ascomycota, Hypocreales, Nectriaceae) species complex. Fungal Divers 70:1–29
- Salgado-Salazar C, Rossman AY, Chaverri P (2016) The genus *Thelonectria* (Nectriaceae, Hypocreales, Ascomycota) and closely related species with cylindrocarpon-like asexual states. Fungal Divers 80:411–455
- Salgüero R, Borman AM, Herrtage M, Benckebroun G et al (2013) *Rasamsonia argillacea* mycosis in a dog: first case in Europe. Vet Rec Case Rep 1:101373
- Samarakoon MC, Hyde KD, Promputtha I, Hongsanan S et al (2016) Evolution of *Xylariomycetidae* (Ascomycota: Sordariomycetes). Mycosphere 7:1746–1761
- Samerpitak K, Van der Linde E, Choi HJ, van den Ende AG et al (2014) Taxonomy of *Ochroconis*, genus including opportunistic pathogens on humans and animals. Fungal Divers 65:89–126
- Samerpitak K, Gerrits van den Ende AHG, Menken SBJ, de Hoog GS (2015) Three new species of the genus *Ochroconis*. Mycopath 180:7–1817
- Samerpitak K, van den Ende BH, Stielow JB, Menken SB et al (2016) Barcoding and species recognition of opportunistic pathogens in *Ochroconis* and *Verruconis*. Fungal Biol 120:219–230
- Samón-Legrá E, Lavandera DE, González MC (2014) A preliminary checklist of marine fungi from Zapata Swamp Biosphere Reserve, South of Cuba. Middle-East J Sci Res 21:1771–1775
- Samson RA, Yilmaz N, Houbraken J, Spierenburg H et al (2011) Phylogeny and nomenclature of the genus *Talaromyces* and taxa accommodated in *Penicillium* subgenus *Biverticillium*. Stud Mycol 70:159–183
- Samson RA, Visagie CM, Houbraken J, Hong SB et al (2014) Phylogeny, identification and nomenclature of the genus *Aspergillus*. Stud Mycol 78:141–173
- Samuels GJ, Lu BS, Chaverri P, Candoussau F et al (2009) *Cyanonectria*, a new genus for *Nectria cyanostoma* and its *Fusarium* anamorph. Mycol Prog 8:49–58
- Sánchez RM, Giordano L, Anderson FE, Bianchinotti M (2013) A new species of *Scopinella* from pampas grass in Argentina. Mycotaxon 122:265–270
- Sánchez FJ, Meeßen J, del Carmen Ruiz M, Leopoldo G et al (2014a) UV-C tolerance of symbiotic *Trebouxia* sp. in the space-tested lichen species *Rhizocarpon geographicum* and *Circinaria gyrosa*: role of the hydration state and cortex/screening substances. Int J Astrobiol 13:1–18
- Sánchez S, Gómez E, Martín M, De Miguel AM et al (2014b) Experiments on the life cycle and factors affecting reproduction of *Sphaerosporella brunnea* provide evidence for rapid asexual propagation by conidiospores and for homothallism in an ectomycorrhizal competitor of cultivated truffle species. Fung Ecol 8:59–65
- Sanders WB, de los Ríos A (2016) Structure of foliicolous thalli of the *Gomphillaceae* in a south-western Florida lichen community. Lichenologist 48:293–303
- Sanders WB, Lücking R (2015) Three new species of foliicolous *Gomphillaceae* (lichen-forming ascomycetes) from southern Florida. Bryologist 118:170–177
- Sanders WB, de los Ríos A (2016) Structure of foliicolous thalli of the *Gomphillaceae* in a south-western Florida lichen community. Lichenologist 48:293–303
- Sanderson NA, Hawksworth DL, Aptroot A (2009) *Melaspilea* Nyl. In: Smith CW, Aptroot A, Coppins BJ, Fletcher A et al. (eds). The lichens of Great Britain and Ireland. British Lichen Society, Department of Botany. The Natural History Museum pp 576–579
- Sandoval-Denis M, Gené J, Sutton DA, Wiederhold NP, Guarro J (2015a) *Acrophialophora*, a poorly known fungus with clinical significance. J Clin Microbiol 53:1549–1555

- Sandoval-Denis M, Sutton DA, Martin-Vicente A, Cano-Lira JF, Wiederhold N, Guarro J, Gené J (2015b) *Cladosporium* species recovered from clinical samples in the United States. *J Clin Microbiol*. doi:10.1128/JCM.01482-15
- Sandoval-Denis M, Gené J, Sutton DA, Cano-Lira JF et al (2016a) Redefining *Microascus*, *Scopulariopsis* and allied genera. *Persoonia* 36:1–36
- Sandoval-Denis M, Guarro J, Cano-Lira JF, Sutton DA et al (2016b) Phylogeny and taxonomic revision of *Microascaceae* with emphasis on synnematous fungi. *Stud Mycol* 83:193–233
- Sandoval-Leiva P, Carmarán CC, Park D, Romero AI, Johnston PR (2014) Vibrissaceous fungi from the southern hemisphere, including *Chlorovibrissea chilensis* (*Helotiales*, *incertae sedis*) sp. nov. *Mycologia* 106:1159–1167
- Sang H, An T, Kim CS, Shin GS et al (2013) Two novel *Talaromyces* species isolated from medicinal crops in Korea. *J Microbiol* 51:704–708
- Sanjuan TI, Franco-Molano AE, Kepler RM, Spatafora JW et al (2015) Five new species of entomopathogenic fungi from the Amazon and evolution of neotropical *Ophiocordyceps*. *Fungal Biol* 119:901–916
- Sanoamuang N, Jitjak W, Rodtong S, Whalley A (2013) *Gelatinomyces siamensis* gen. sp. nov. (*Ascomycota*, *Leotiomycetes incertae sedis*) on bamboo in Thailand. *IMA Fungus* 4:71–87
- Santa-Izabel TS, Cruz ACR, Gusmão LFP (2013) Conidial fungi from the semi-arid Caatinga biome of Brazil. *Ellisembiopsis* gen. nov., new variety of *Sporidesmiella* and some notes on *Sporidesmium* complex. *Mycosphere* 4:156–163
- Santa Izabel TD, Almeida DA, Monteiro JS, Marques MF et al (2015) *Anaexserticlavula caatingae*, a new conidial fungus from the semi-arid Caatinga biome of Brazil. *Mycotaxon* 130:445–449
- Santamaría S (2008) *Laboulbeniales* on semiaquatic *Heteroptera*. A new species of *Triceromyces* (*Ascomycota*, *Laboulbeniales*) on *Microvelia* (*Heteroptera*, *Veliidae*) from Spain. *Alico* 26:15–21
- Santamaría S, Faillé A (2009) New species of *Laboulbenia* and *Rhachomyces* (*Laboulbeniales*, *Ascomycota*), some of them polymorphic, parasitic on termiticolous ground beetles from tropical Africa. *Nova Hedw* 89:97–120
- Santamaría S, Enghoff H, Reboleira ASPS (2014) *Laboulbeniales* on millipedes: *Diplopodomyces* and *Troglobomyces*. *Mycologia* 106:1027–1038
- Santamaría S, Enghoff H, Reboleira ASPS (2016) Hidden biodiversity revealed by collections-based research—*Laboulbeniales* in millipedes: genus *Rickia*. *Phytotaxa* 243:101–127
- Santos MD, Fonseca ME, Boiteux LS, Dianese JC (2012) First approach to molecular phylogeny within the genus *Phyllachora*. *Phytopath* 102:105
- Santos L, Phillips AJL, Crous P, Alves A (2017) Diaporthe species on Rosaceae with descriptions of *D. pyracanthae* sp. nov. and *D. malorum* sp. nov. *Mycosphere* 8:485–511
- Sanz-Ros AV, Müller MM, San Martin R, Diez JJ (2015) Fungal endophytic communities on twigs of fast and slow growing Scots pine (*Pinus sylvestris* L.) in Northern Spain. *Fungal Biol* 119:870–883
- Saravanakumar D, Ciavarella A, Spadaro D, Garibaldi A, Gullino ML (2008) *Metschnikowia pulcherrima* strain MACH1 outcompetes *Botrytis cinerea*, *Alternaria alternata* and *Penicillium expansum* in apples through iron depletion. *Postharvest Biol Technol* 49:121–128
- Sari A, Tuzen M (2010) Biosorption of As (III) and As (V) from aqueous solution by lichen (*Xanthoria parietina*) biomass. *Sep Sci Technol* 45:463–471
- Sarkar J, Ray S, Chattopadhyay D, Laskar A, Acharya K (2012) Mycogenesis of gold nanoparticles using a phytopathogen *Alternaria alternata*. *Bioprocess Biosyst Eng* 35:637–643
- Sarma VV (2015) A dichotomous key to genera of marine fungi recorded on *Rhizophora* spp. *Indian J Geo-Marine Sci* 44:1–7
- Sarma VV, Raghukumar S (2013) Mangicolous fungi from Chorao mangroves, Goa, West coast of India: diversity and frequency of occurrence. *Nova Hedw* 97:533–542
- Sati SC, Arya P, Belwal M (2009) *Tetracladium nainitalense* sp. nov., a root endophyte from Kumaun Himalaya. *India. Mycologia* 101:692–695
- Sato H, Ban S, Masuya H, Hosoya T (2010a) Reassessment of type specimens of *Cordyceps* and its allies described by Dr Yosio Kobayashi preserved in the mycological herbarium of the National Museum of Nature and Science (TNS). Part 1: the genus *Torrubilla*. *Mycoscience* 51:154–161
- Sato H, Ban S, Masuya H, Hosoya T (2010b) Reassessment of type specimens of *Cordyceps* and its allies described by Dr Yosio Kobayashi, preserved in the National Museum of Nature and Science. Part 2: *Cordyceps* (*Elaphocordyceps*) on *Elaphomyces*. *Mycoscience* 51:387–390
- Savić S, Tibell L (2008a) *Atla*, a new genus in the *Verrucariaceae* (*Verrucariales*). *Lichenologist* 40:269–282
- Savić S, Tibell L (2008b) The lichen genus *Henrica* (*Verrucariaceae*, *Eurotiomycetes*) in Northern Europe. *Nordic J Bot* 26:237–247
- Savić S, Tibell L (2012) Polyblastia in northern Europe and the adjacent Arctic. *Uppsala: Acta Universitatis Upsaliensis* 36:1–69
- Savić S, Tibell L, Gueidan C, Lutzoni F (2008) Molecular phylogeny and systematics of *Polyblastia* (*Verrucariaceae*, *Eurotiomycetes*) and allied genera. *Mycol Res* 112:1307–1318
- Sbissi I, Neffati M, Boudabous A, Murat C, Gtari M (2010) Phylogenetic affiliation of the desert truffles *Picoa juniperi* and *Picoa lefebvrei*. *Antonie van Leeuwenhoek* 98:429–436
- Schmechel D, Green BJ, Blachere FM, Janotka E, Beechold DH (2008) Analytical bias of cross-reactive polyclonal antibodies for environmental immunoassays of *Alternaria alternata*. *J Allergy Clin Immunol* 121:763–768
- Schmidt V (2015) Fungal infections in reptiles—an emerging problem. *J Exotic Pet Med* 24:267–275
- Schmidt LE, Deyrup ST, Baltrusaitis J, Swenson DC et al (2010) Hymenopsins A and B and a macrophorin analogue from a fungicolous *Hymenopsis* sp. *J Nat Prod* 73:404–408
- Schmidt V, Plenz B, Pfaff M, Pees M (2012) Disseminated systemic mycosis in *Veiled chameleons* (*Chamaeleo calyptratus*) caused by *Chamaeleomyces granulomatis*. *Vet Microbiol* 161:145–152
- Schmitt I, Partida-Martinez LP, Winkler R, Voigt K et al (2008) Evolution of host resistance in a toxin-producing bacterial-fungal alliance. *ISME J* 2:632–641
- Schmitt I, Fankhauser JD, Sweeney K, Spribille T et al (2010) *Gyalectoid pertusaria* species form a sister-clade to *Coccotrema* (*Ostropomyctidae*, *Ascomycota*) and comprise the new lichen genus *Gyalectaria*. *Mycology* 1:75–83
- Schmitt I, Otte J, Parmen S, Sadowska-Deś A et al (2012) A new circumscription of the genus *Varicellaria* (*Pertusariales*, *Ascomycota*). *MycoKeys* 4:23
- Schmull M, Miadlikowska J, Pelzer M, Stocker-Worgötter E et al (2011) Phylogenetic affiliations of members of the heterogeneous lichen-forming fungi of the genus *Lecidea* sensu *Zahlbrückner* (*Lecanoromycetes*, *Ascomycota*). *Mycologia* 103:983–1003
- Schneider K, Resl P, Westberg M, Spribille T (2015) A new, highly effective primer pair to exclude algae when amplifying nuclear large ribosomal subunit (LSU) DNA from lichens. *Lichenologist* 47:269–275

- Schneider K, Resl P, Spribille T (2016) Escape from the cryptic species trap: lichen evolution on both sides of a cyanobacterial acquisition event. *Mol Ecol* 25:3453–3468
- Schoch CL, Crous PW, Groenewald JZ, Boehm EWA et al (2009) A class-wide phylogenetic assessment of *Dothideomycetes*. *Stud Mycol* 64:1–15
- Schroers HJ, Gräfenhan T, Nirenberg HI, Seifert KA (2011) A revision of *Cyanonectria* and *Geejayessia* gen. nov., and related species with *Fusarium*-like anamorphs. *Stud Mycol* 68:115–138
- Schultz M, Moon KH (2011) Notes on taxonomy and distribution of some critical cyanobacterial lichens from South Korea. *Nova Hedw* 92:479–486
- Schultz M, Zedda L, Rambold G (2009) New records of lichen taxa from Namibia and South Africa. *Biblioth Lichenol* 99:315–334
- Schumacher J, Reinbold J, Aikaterini D (2015) The *Lachnellula* canker of *Abies alba*—a new disease phenomenon in the German Black Forest. *Jahrbuch der Baumpflege Yearbook of Arboriculture*, pp 45–51
- Scott JA, Ewaze JO, Summerbell RC, Arocha-Rosete Y et al (2016) Multilocus DNA sequencing of the whiskey fungus reveals a continental-scale speciation pattern. *Persoonia* 37:13–20
- Seena S, Pascoal C, Marvanova L, Cassio F (2010) DNA barcoding of fungi: a case study using ITS sequences for identifying aquatic hyphomycete species. *Fungal Divers* 44:77–87
- Seena S, Duarte S, Pascoal C, Cassio F (2012) Intraspecific variation of the aquatic fungus *Articulospora tetracladia*: an ubiquitous perspective. *PLoS One* 7:e35884
- Seephueak P, Phongpaichit S, Hyde KD, Petcharat V (2011a) Diversity of saprobic fungi on decaying rubber logs (*Hevea brasiliensis*). *Sydowia* 63:249–282
- Seephueak P, Phongpaichit S, Hyde KD, Petcharat V (2011b) Diversity of saprobic fungi on decaying branch litter of the rubber tree (*Hevea brasiliensis*). *Mycosphere* 2:307–330
- Seifert KA, Morgan-Jones G, Gams W, Kendrick B (2011) The genera of Hyphomycetes. CBS Biodiversity Series 9, CBS-KNAW Fungal Biodiversity Centre, Utrecht
- Selbmann L, De Hoog GS, Zucconi L, Isola D et al (2008) Drought meets acid: three new genera in a dothidealean clade of extremotolerant fungi. *Stud Mycol* 61:1–20
- Selbmann L, Isola D, Egidi E, Zucconi L et al (2014) Mountain tips as reservoirs for new rock-fungal entities: *Saxomyces* gen. nov. and four new species from the Alps. *Fungal Divers* 6:167–182
- Selbmann L, Zucconi L, Isola D, Onofri S (2015) Rock black fungi: excellence in the extremes, from the Antarctic to space. *Curr Genetics* 61:335–345
- Selçuk F, Ekici K (2014a) A new species of *Manoharachariella* (hyphomycetes) from Central Anatolia, Turkey. *Mycosphere* 5:419–423
- Selçuk F, Ekici K (2014b) A new species of *Manoharachariella* (hyphomycetes) from Central Anatolia, Turkey. *Mycosphere* 5:419–423
- Selçuk F, Hüseyin E, Şahin A, Cebeci CC (2014) Hyphomycetous fungi in several forest ecosystems of Black Sea provinces of Turkey. *Mycosphere* 5:334–344
- Selva SB (2013) The calicioid lichens and fungi of the Acadian Forest Ecoregion of northeastern North America, I. New species and range extensions. *Bryologist* 116:248–256
- Senanayake IC, Maharachchikumbura SS, Mortimer PE, Xu J et al (2014) Phylogenetic studies on *Vialaeaceae* reveals a novel species, *Vialaea mangiferae*. *Sydowia* 66:203–216
- Senanayake IC, Maharachchikumbura SSN, Hyde KD, Bhat JD et al (2015) Towards unraveling relationships in *Xylariomycetidae* (*Sordariomycetes*). *Fungal Divers* 73:73–144
- Senanayake IC, Al-Sadi AM, Bhat DJ, Camporesi E et al (2016) *Phomatosporales* ord. nov. and *Phomatosporaceae* fam. nov., to accommodate *Lanspora*, *Phomatospora* and *Tenuimurus*, gen. nov. *Mycosphere* 7:628–641
- Senanayake IC, Maharachchikumbura SSN, Jeewon R, Promputtha I et al (2017) Morphophylogenetic study of *Sydiwiellaceae* reveals several new genera. *Mycosphere* 8:172–217
- Şenkardeşler A, Duman DC, Lököş L, Ahti T (2016) *Cladonia trapezuntica* (*Cladoniaceae*, lichenized Ascomycota): a robust morphotype of *Pycnothelia papillaria*, a taxonomic study with conservational survey. *Turkish J Botany* 40:104–111
- Seo C, Choi YH, Ahn JS, Yim JH et al (2009a) PTP1B inhibitory effects of tridepside and related metabolites isolated from the Antarctic lichen *Umbilicaria antarctica*. *J Enzyme Inhibition Med Chem* 24:1133–1137
- Seo C, Sohn JH, Ahn JS, Yim JH et al (2009b) Protein tyrosine phosphatase 1B inhibitory effects of depsidone and pseudodepsidone metabolites from the Antarctic lichen *Stereocaulon alpinum*. *Bioorganic Med Chem Lett* 19:2801–2803
- Seo MY, Matsuda Y, Nakashima C, Ito SI (2012) Taxonomic reevaluation of *Raffaelea quercivora* isolates collected from mass mortality of oak trees in Japan. *Mycoscience* 53:211–219
- Sepcik K, Zalar P, Gunde-Cimerman N (2010) Low water activity induces the production of bioactive metabolites in halophilic and halotolerant fungi. *Marine Drugs* 9:43–58
- Serrato-Diaz LM, Rivera-Vargas LI, Goenaga R, Verkley GJM, FrenchMonar RD (2011) First report of a *Lasmenia* sp. causing rachis necrosis, flower abortion, fruit rot, and leaf spots on rambutan in Puerto Rico. *Plant Dis* 95:1313
- Sert HB, Sümbül H, Sterflinger K (2011) Two new species of *Capnobotryella* from historical monuments. *Mycol Prog* 10:333–339
- Sert HB, Wuczkowski M, Sterflinger K (2012) *Capnobotryella isiloglui*, a new rock-inhabiting fungus from Austria. *Turkish J Botany* 36:401–407
- Sérusiaux E, Brand AM, Fischer E, Killmann D et al (2009a) A new species of *Melanotopelia* (*Graphidaceae*) from Africa. *Lichenologist* 41:243–247
- Sérusiaux E, Goffinet B, Miadlikowska J, Vitikainen O (2009b) Taxonomy, phylogeny and biogeography of the lichen genus *Peltigera* in Papua New Guinea. *Fungal Divers* 38:185–224
- Sérusiaux E, Van den Boom P, Ertz D (2010) A two-gene phylogeny shows the lichen genus *Niebla* (*Lecanorales*) is endemic to the new world and does not occur in Macaronesia nor in the Mediterranean Basin. *Fungal Biol* 114:528–537
- Sethy PP, Pandit GS, Sharma BO (2012) Lichens on mangrove plants in Andaman Islands, India. *Mycosphere* 3:476–484
- Severn-Ellis A, Schoeman MH, Willemse S, Sippel A et al. (2012) Towards guava wilt disease resistance in South Africa. In III International Symposium on Guava and other Myrtaceae, pp 73–79
- Seyedmousavi S, Guillot J, de Hoog GS (2013) *Phaeohyphomycoses*, emerging opportunistic diseases in animals. *Clin Microbiol Rev* 26:19–34
- Seyedmousavi S, Samerpitak K, Rijs AJ, Melchers WJ et al (2014) Antifungal susceptibility patterns of opportunistic fungi in the genera *Verruconis* and *Ochroconis*. *Antimicrobial Agents Hemotherapy* 58:3285–3292
- Sezer O, Özgiş K, Yaylaci ÖK, Öztürk D et al (2016) Contributions to lichen biota of Günyüzü (Eskişehir). *J Applied Biol Sci* 10:26–32
- Shabana YM, Abdel-Fattah GM, Ismail AE, Rashad YM (2008) Control of brown spot pathogen of rice (*Bipolaris oryzae*) using some phenolic antioxidants. *Brazilian J Microbiol* 39:438–444
- Shali A, Ghasemi S, Ahmadian G, Ranjbar G et al (2010) *Bacillus pumilus* SG2 chitinases induced and regulated by chitin, show

- inhibitory activity against *Fusarium graminearum* and *Bipolaris sorokiniana*. *Phytoparasitica* 38:141–147
- Shang QJ, Hyde KD, Phookamsak R, Doilom M et al (2017) *Diatrypella tectonae* and *Peroneutypa mackenziei* spp. nov. (*Diatrypaceae*) from northern Thailand. *Mycol Prog* 16:463–476
- Sharma BO, Khadikar P (2011) Two additional new species of *Carbacanthographis* from India. *Lichenologist* 43:293–297
- Sharma BO, Makija U, Khadikar P (2010) Two species of *Carbacanthographis* from India. *Lichenologist* 42:391–395
- Sharma R, Gräser Y, Singh SK (2011) *Auxarthronopsis*, a new genus of *Onygenales* isolated from the vicinity of Bandhavgarh National Park, India. *IMA fungus* 4:89–102
- Sharma R, Sharma R, Crous PW (2015a) *Matsushimamycetes*, a new genus of keratinophilic fungi from soil in central India. *IMA Fungus* 6:337–343
- Sharma G, Pinnaka AK, Shenoy BD (2015b) Resolving the *Colletotrichum siamense* species complex using ApMat marker. *Fungal Divers* 71:247–264
- Sharma G, Shenoy BD (2016) *Colletotrichum* systematics: past, present and prospects. *Mycosphere* 7:1093–1102
- Sheard JW (2010) The lichen genus *Rinodina* (Ach.) gray (*Lecanoromycetidae, Physciaceae*) in North America, North of Mexico. NRC Research Press, Ottawa
- Sheard JW, Knudsen K, Mayrhofer H, Morse CA (2011) Three new species of *Rinodina* (*Physciaceae*) and a new record from North America. *Bryologist* 114:453–465
- Shearer BL, Crane CE (2014) Host range of the stem canker pathogen *Luteocirrhus shearii* mainly limited to Banksia. *Australas Plant Pathol* 43:257–266
- Shearer CA, Raja HA, Miller AN, Nelson P et al (2009) The molecular phylogeny of freshwater *Dothideomycetes*. *Stud Mycol* 64:145–153
- Shearer CA, Pang KL, Suetrong S, Raja HA (2014) Phylogeny of the *Dothideomycetes* and other classes of freshwater fissiliticate *Ascomycota*. In: Jones EBG, Hyde KD, Pang KL (eds) Freshwater fungi and fungal-like organisms. De Gruyter, Berlin, pp 25–45
- Sheik MA, Raina AK, Upreti DK (2009) Lichen flora of Surinsar-Mansar wildlife sanctuary. *J Appl Nat Sci* 1(1):79–81
- Sheikh MA, Raina AK, Upreti DK (2016) Lichens—enshrouding the bare rocks in pappad area of Samba district, Jammu and Kashmir. *Int J Curr Res Rev* 8:29–32
- Shen W, Mao H, Huang Q, Dong J (2015) Benzenediol lactones: a class of fungal metabolites with diverse structural features and biological activities. *European J Med Chem* 97:747–777
- Shen M, Zhang JQ, Zhang Y (2016) *Venturia* species form sooty mold-like colonies on leaves of *Salix*: introducing *Venturia fuliginosa* sp. nov. *Mycosphere* 7:1292–1300
- Shenoy BD, Jeewon R, Wang H, Amandeep K et al (2010) Sequence data reveals phylogenetic affinities of fungal anamorphs *Bahusutrabeeja*, *Diplococcum*, *Natarajania*, *Paliphora*, *Polyschema*, *Rattania* and *Spadicoides*. *Fungal Divers* 44:161–169
- Shi GK, Luo JT, Hou CL (2010) Species of the *Rhytismataceae* on *Pinus armandii*. *Mycosistema* 29:159–163
- Shi H, Qian Z, Wang X, Liu D et al (2015) The genus *Letrovittia* (*Letrovitiaceae*: Lichenized Ascomycota) new to Cambodia. *Mycobiology* 43:163–165
- Shi D, Lu G, Mei H, de Hoog GS et al (2016) Subcutaneous infection by *Ochroconis mirabilis* in an immunocompetent patient. *Med Mycol Case Rep* 11:44–47
- Shirouzu T, Hirose D, Tokumasu S, To-Anun C, Maekawa N (2010) Host affinity and phylogenetic position of a new anamorphic fungus *Beltraniella botryospora* from living and fallen leaves of evergreen oaks. *Fungal Divers* 43:85–92
- Shivarov VV, Stoykov DY (2013) New records of pyrenocarpous lichenized fungi from Bulgaria. *Mycotaxon* 121:133–138
- Shivas RG, Grice KRE, Young AJ (2010) *Ramichloridium* spp. on *Musa* in northern Queensland: introducing *Ramichloridium ducassei* sp. nov. on leaf streaks of Ducasse banana. *Australas Pl Path* 40:61–65
- Shivas RG, Marney TS, Tan YP, McTaggart AR (2015) Novel species of *Cercospora* and *Pseudocercospora* (*Capnodiales, Mycosphaerellaceae*) from Australia. *Fungal Biol* 119:362–369
- Shoemaker RA, Malloch D, Hambleton S, Liu M (2013) A new *Vestigium* on *Abies balsamea*. *N Am Fungi* 8:1–6
- Shrestha G, St Clair LL (2009) The lichen flora of southwestern Colorado. *Evansia* 26:102–123
- Shukla V, Upreti DK (2009) Polycyclic aromatic hydrocarbon (PAH) accumulation in lichen, *Phaeophyscia hispidula* of DehraDun City, Garhwal Himalayas. *Environ Monit Assess* 149:1–7
- Shukla V, Upreti DK, Patel DK (2012) Physiological attributes of lichen, *Phaeophyscia hispidula* in heavy metal rich sites of Dehra Dun, India. *J Environ Biol* 33:1051
- Shukla V, Patel DK, Upreti DK, Yunus M, Prasad S (2013) A comparison of metallic contents in lichen *Pyxine subcineraria*, its substratum and soil. *Int J Environ Sci Technol* 10:37–46
- Shukla P, Upreti DK, Nayaka S, Tiwari P (2014a) Natural dyes from Himalayan lichens. *Indian J Tradit Knowl* 13:195–201
- Shukla V, Upreti DK, Bajpai R (2014b) Lichen diversity in different lichenogeographical regions of India. In: lichens to biomonitor the environment. Springer, India. pp 61–96
- Shukla V, Kumari R, Patel DK, Upreti DK (2016) Characterization of the diversity of mycosporine-like amino acids in lichens from high altitude region of Himalaya. *Amino acids* 48:129–136
- Siegel D, Rasenko T, Koch M, Nehls I (2009) Determination of the *Alternaria* mycotoxin tenuazonic acid in cereals by high-performance liquid chromatography-electrospray ionization ion-trap multistage mass spectrometry after derivatization with 2, 4-dinitrophenylhydrazine. *J Chromatogr A* 1216:4582–4588
- Siegel D, Merkel S, Koch M, Nehls I (2010) Quantification of the *Alternaria* mycotoxin tenuazonic acid in beer. *Food Chem* 120:902–906
- Siemens JA, Zwiazek JJ (2008) Root hydraulic properties and growth of balsam poplar (*Populus balsamifera*) mycorrhizal with *Hebeloma crustuliniforme* and *Wilcoxina mikolae* var. *mikolae*. *Mycorrhiza* 18:393–401
- Sigler L, Gibas CF, Kokotovic B, Bertelsen MF (2010a) Disseminated mycosis in veiled chameleons (*Chamaeleo calyptratus*) caused by *Chamaeleomyces granulomatis*, a new fungus related to *Paecilomyces viridis*. *J Clin Microbiol* 48:3182–3192
- Sigler L, Sutton DA, Gibas CFC, Summerbell RC et al (2010b) *Phialosimplex*, a new anamorphic genus associated with infections in dogs and having phylogenetic affinity to the *Trichocomaceae*. *Med Mycol* 48:335–345
- Sigler L, Hambleton S, Paré JA (2013) Molecular characterization of reptile pathogens currently known as members of the *Chrysosporium* anamorph of *Nannizziopsis vriesii* complex and relationship with some human-associated isolates. *J Clin Microbiol* 51:3338–3357
- Silva SS, Gusmão LF (2013) Conidial fungi from the semi-arid Caatinga Biome of Brazil. A new species of *Dictyochaeta*. *Mycosphere* 4:701–705
- Silva BF, Carrijo-Mauad JR, Braga FR, Campos AK et al (2010) Efficacy of *Duddingtonia flagrans* and *Arthrobotrys robusta* in controlling sheep parasitic gastroenteritis. *Parasitol Res* 106:1343–1350
- Silva GB, Lustosa DC, Nechet KL, Conceicao CS et al (2011) Diversity of plant pathogenic fungi associated with native Amazon forest species. *Phytopathology* 101:166–167

- Silva MD, Castañeda-Ruiz RF, Pereira OL, Barreto RW (2012) *Alveariospora*, a new anamorphic genus from trichomes of *Dimorphandra mollis* in Brazil. Mycotaxon 119:109–116
- Silva CR, Castañeda-Ruiz RF, Gusmão P, Fernando L (2014a) *Anacraspedodidymum*, a new genus from submerged wood in Brazil. Mycotaxon 128:11–15
- Silva SS, Cruz AC, Gusmão LF, Castañeda-Ruiz RF (2014b) *Diplococcum variegatum*, a new conidial fungus from the semi-arid Caatinga biome of Brazil. Mycotaxon 127:59–62
- Silva SS, Gusmão LFP, Castañeda-Ruiz RF (2015a) Conidial fungi on *Araucaria angustifolia*: a new species of *Trichoconis* and new records from Brazil. Mycotaxon 130:1051–1059
- Silva CR, Gusmão LFP, Castañeda-Ruiz RF (2015b) Two new conidial fungi from Chapada Diamantina, Brazil. Mycotaxon 130:437–443
- Silva SS, Gusmão LF, Castañeda-Ruiz RF (2016) *Arthromoniliphora araucariae* gen. & sp. nov. from Brazilian pine. Mycotaxon 131:821–826
- Silvera-Simón C, Mena-Portales J, Gene J, Cano JF, Guarro J (2009) *Repetophragma calongei* [sic] sp. nov. and other interesting dematiaceous hyphomycetes from the north of Spain. An Jard Bot Madr 66(S1):33–39
- Silvera-Simón C, Mercado-Sierra A, Mena-Portales J, Gene J, Guarro J (2010) Two new species of *Digitodesmium* from Euskadi (Spain). Cryptog Mycol 31:211–218
- Silvério ML, Cavalcanti MA, Bezerra JL (2011) First record of *Leptomeliola uvariae* for South America. Mycotaxon 116:1–5
- Simmons DR, de Beer ZW, Huang YT, Bateman C et al (2016) New *Raffaelea* species (*Ophiostomatales*) from the USA and Taiwan associated with ambrosia beetles and plant hosts. IMA Fungus 7:265–273
- Simon UK, Groenewald JZ, Crous PW (2009) *Cymadothea trifolii*, an obligate biotrophic leaf parasite of *Trifolium*, belongs to *Mycosphaerellaceae* as shown by nuclear ribosomal DNA analyses. Persoonia 22:49–55
- Singh KP, Swarnalatha G (2011) A new species of *Schistophoron* from India. Lichenologist 43:209–212
- Singh SK, Singh PN (2012) A New species of *Sarcinella* from India. Indian Phytopath 58:341–342
- Singh SM, Ravindra R (2013) Impact of climate change on Lichen and Moss Communities in Ny-Ålesund, Arctic: some preliminary observations. Earth system processes and disaster management. Springer, Heidelberg, pp 93–100
- Singh SK, Gaikwad VP, Waingankar VM (2009) A new endophytic ascomycete from healthy leaves of *Pongamia pinnata* Merr. Indian Phytopath 62:124–125
- Singh SM, Pereira N, Ravindra R (2010) Adaptive mechanisms for stress tolerance in Antarctic plants. Curr Sci 99:334–340
- Singh UB, Sahu A, Singh RK, Singh DP et al (2012) Evaluation of biocontrol potential of *Arthrobotrys oligospora* against *Meloidogyne graminicola* and *Rhizoctonia solani* in Rice (*Oryza sativa* L.). Biol Control 60:262–270
- Singh G, Divakar PK, Dal Grande F, Otte J et al (2013a) The sister-group relationships of the largest family of lichenized fungi, *Parmeliaceae* (*Lecanorales*, *Ascomycota*). Fungal Biol 117:715–721
- Singh R, Ranjan S, Nayaka S, Pathre UV, Shirke PA (2013b) Functional characteristics of a fruticose type of lichen, *Stereocaulon foliolosum* Nyl. in response to light and water stress. Acta Physiologiae Plantarum 35:1605–1615
- Singh UB, Sahu A, Sahu N, Singh BP, Singh RK et al (2013c) Can endophytic *Arthrobotrys oligospora* modulate accumulation of defence related biomolecules and induced systemic resistance in tomato (*Lycopersicon esculentum* Mill.) against root knot disease caused by *Meloidogyne incognita*. Appl Soil Ecol 63:45–56
- Singh SM, Singh PN, Singh SK, Sharma PK (2014) Pigment, fatty acid and extracellular enzyme analysis of a fungal strain *Thelebolus microsporus* from Larsemann Hills, Antarctica. Polar Rec 50:31–36
- Singh G, Dal Grande F, Divakar PK, Otte J et al (2015) Coalescent-based species delimitation approach uncovers high cryptic diversity in the cosmopolitan lichen-forming fungal genus *Protoparmelia* (*Lecanorales*, *Ascomycota*). PLoS One 10:e0124625
- Singtripop C, Hongsanan SC, Camporesi E, Ariyawansa HA et al (2015) *Keissleriella dactyliidis*, sp. nov., from *Dactylis glomerata* and its phylogenetic placement. Sci Asia 41:295–304
- Singtripop C, Hongsanan S, Li J, De Silva NI et al (2016) *Chaetothyridia mangiferae* sp. nov., a new species of *Chaetothyridia*. Phytotaxa 255:21–33
- Sinha R, Chattopadhyay S (2011) Changes in the leaf proteome profile of *Mentha arvensis* in response to *Alternaria alternata* infection. J Proteom 74:327–336
- Sinha GP, Jagadeesh Ram TAM (2011) Lichen diversity in Sikkim. Biodiversity of Sikkim. Exploring and Conserving a Global Hotspot, Gangtok: Sikkim:Information and Public Relations Department, pp 13–28
- Sipiczki M, Pfiegl WP, Safar SWB, Morais PB, Rosa CA (2016) *Metahyphopichia laotica* gen. nov., sp. nov., a novel polymorphic yeast related to *Hyphopichia*. Int J Syst Evol Microbiol 66:2550–2557
- Sipman HJM (2014) New species of *Graphidaceae* from the Neotropics and Southeast Asia. Phytotaxa 189:289–311
- Sipman HJ, Raus T (2015) Lichens and lichenicolous fungi from the island of Chios (Aegean Sea, Greece). Herzogia 28:496–519
- Sipman HJM, Lücking R, Aptroot A, Kalb K et al (2012) A first assessment of the *Ticolichen* biodiversity inventory in Costa Rica and adjacent areas: the thelotremoid *Graphidaceae* (*Ascomycota*: *Ostropales*). Phytotaxa 55:1–214
- Sir EB, Kuhnert E, Lambert C, Hladki AI et al (2016a) New species and reports of *Hypoxyylon* from Argentina recognized by a polyphasic approach. Mycol Prog. doi:10.1007/s11557-016-1182-z
- Sir EB, Lambert C, Wendt L, Hladki AI, Romero AI, Stadler M (2016b) A new species of *Daldinia* (*Xylariaceae*) from the Argentine subtropical montane forest. Mycosphere 7:1378–1388
- Sivakumar T (2013) A review on biodiversity of marine and mangrove fungi. Int J Curr Res Acad Rev 1:26–44
- Sivanesan A (1984) The bitunicate ascomycetes and their anamorphs. J Cramer, Vaduz
- Sivichai S, Boonyuen N (2010) *Jahnula morakotii* sp. nov. and *J. appendiculata* from a peat swamp in Thailand. Mycotaxon 112:475–481
- Sivichai S, Sri-Indrasutdi V, Jones E (2011) *Jahnula aquatica* and its anamorph *Xylomyces chlamydosporus* on submerged wood in Thailand. Mycotaxon 116:137–142
- Six DL, Stone WD, de Beer ZW, Woolfolk SW (2009) *Ambrosiella beaveri* sp. nov., associated with an exotic ambrosia beetle, *Xylosandrus multilatus* (Coleoptera: Curculionidae, Scolytinae), in Mississippi, USA. Antonie van Leeuwenhoek 96:17–29
- Skjøth CA, Sommer J, Frederiksen L, Gosewinkel Karlson U (2012) Crop harvest in Denmark and Central Europe contributes to the local load of airborne *Alternaria* spore concentrations in Copenhagen. Atmos Chem Phys 12:11107–11123
- Slippers B, Boissin E, Phillips AJL, Groenewald JZ, Wingfield MJ et al (2013) Phylogenetic lineages in the *Botryosphaeraiales*: a systematic and evolutionary framework. Stud Mycol 76:31–49
- Small KW, Chan CK, Silva-Garcia R, Walsh TJ (2014) Onset of an outbreak of *Bipolaris hawaiiensis* fungal endophthalmitis after intravitreal injections of triamcinolone. Ophthalmology 121:952–958

- Smith JA, Mount L, Mayfield AE III, Bates CA et al (2009) First report of laurel wilt disease caused by *Raffaelea lauricola* on camphor in Florida and Georgia. *Plant Dis* 93:198–198
- Smith ME, Henkel TW, Catherine Aime M, Premier AK, Vilgalys R (2011) Ectomycorrhizal fungal diversity and community structure on three co-occurring leguminous canopy tree species in a Neotropical rainforest. *New Phytol* 192:699–712
- Snelgrove RJ, Gregory LG, Peiró T, Akthar S et al (2014) *Alternaria*-derived serine protease activity drives IL-33-mediated asthma exacerbations. *J Allergy Clin Immunol* 134:583–592
- Soares DJ, Barreto RW (2008) Fungal pathogens of the invasive riparian weed *Hedychium coronarium* from Brazil and their potential for biological control. *Fungal Divers* 28:85–96
- Sobreira PN, Aptroot A, da Silva Cáceres ME (2015) A world key to species of the genus *Bactrospora* (*Roccellaceae*) with a new species from Brazil. *Lichenologist* 47:131–136
- Søchting U, Garrido-Benavent I, Seppelt R, Castello M et al (2014a) *Charcotiana* and *Amundsenia*, two new genera in *Teloschistaceae* (lichenized Ascomycota, subfamily *Xanthorioideae*) hosting two new species from continental Antarctica, and *Austroplaca frigida*, a new name for a continental Antarctic species. *Lichenologist* 46:763–782
- Søchting U, Søgaard MZ, Elix JA, Arup U et al (2014b) *Catenarina* (*Teloschistaceae*, Ascomycota), a new Southern Hemisphere genus with 7-chlorocatenarin. *Lichenologist* 46:175–187
- Sogonov MV, Castlebury LA, Rossman AY, Mejía LC, White JF Jr (2008) Leaf-inhabiting genera of the *Gnomoniaceae*, *Diaporthales*. *Stud Mycol* 62:1–79
- Sohrabi M, Myllys L, Stenroos S (2010a) Successful DNA sequencing of a 75 year-old herbarium specimen of *Aspicilia aschabadiensis* (J. Steiner) Mereschk. *Lichenologist* 42:626–628
- Sohrabi M, Sipman H, Tohranegar Z, Nejadsattari T (2010b) A contribution to the lichenized mycota of Zanjan province Iran. *Iranian J Botany* 16:125–129
- Sohrabi M, Leavitt SD, Halici MG, Shrestha G, Stenroos S (2013a) *Teuvoa*, a new lichen genus in *Megasporaceae* (Ascomycota: Pertusariales), including *Teuvoa junipericola* sp. nov. *Lichenologist* 45:347–360
- Sohrabi M, Stenroos S, Myllys L, Søchting U et al (2013b) Phylogeny and taxonomy of the “manna lichens”. *Mycol Prog* 12:231–269
- Solheim H, Torp TB, Hietala AM (2013) Characterization of the ascomycetes *Therrya fuckelii* and *T. pini* fruiting on Scots pine branches in Nordic countries. *Mycol Prog* 12:37–44
- Song W, Ma X, Tan H, Zhou J (2011) Abscisic acid enhances resistance to *Alternaria solani* in tomato seedlings. *Plant Physiol Biochem* 49:693–700
- Soni K, Pyasi A, Tiwari P, Verma R (2011) Black Mildew of neem seedlings: a new disease record from India. *Indian J Trop Biodiv* 19:133–136
- Sousa RS, Rodrigues FA, Schurt DA, Souza NF, Cruz MFA (2013) Cytological aspects of the infection process of *Pyricularia oryzae* on leaves of wheat plants supplied with silicon. *Trop Plant Pathol* 38:472–477
- Sousa FM, Morais PB, Lachance MA, Rosa CA (2014) *Hagleromyces* gen. nov., a yeast genus in the *Saccharomycetaceae*, and description of *Hagleromyces aurorensis* sp. nov., isolated from water tanks of bromeliads. *Int J Syst Evol Microbiol* 64:2915–2918
- Sparrius LB (2009) *Synarthrothelium*, a new genus in the Arthoniaceae with a thalline exciple, similar to *Synarthronia*. *Biblioth Lichenol* 99:373–382
- Sparrius L, Elix JA, Archer AW (2010) *Astrographa*, a new genus in the *Roccellaceae* with three species from Australia. *Australasian Lichenol* 67:23–28
- Sparrius L, Elix JA, Archer AW (2013) Index Fungorum 27:1
- Spatafora JW, Quandt CA, Kepler RM, Sung GH et al (2015) New 1F1N species combinations in *Ophiocordycipitaceae* (*Hypocreales*). *IMA Fungus* 6:357–362
- Species Fungorum (2017) <http://www.speciesfungorum.org/Names/SynSpecies.asp?RecordID=323547>
- Spooren M (2014) A new species of *Bloxamia* from freshwater in the Netherlands. *Mycosphere* 5:346–349
- Spribille T, Muggia L (2013) Expanded taxon sampling disentangles evolutionary relationships and reveals a new family in *Peltigerales* (*Lecanoromycetidae*, Ascomycota). *Fungal Divers* 58:171–184
- Spribille T, Björk CR, Ekman S, Elix JA et al (2009) Contributions to an epiphytic lichen flora of northwest North America: I. Eight new species from British Columbia inland rain forests. *Bryologist* 112:109–137
- Spribille T, Pérez-Ortega S, Tønsberg T, Schirokauer D (2010) Lichens and lichenicolous fungi of the Klondike Gold Rush National Historic Park, Alaska, in a global biodiversity context. *Bryologist* 113:439–515
- Spribille T, Goffinet B, Klug B, Muggia L et al (2011) Molecular support for the recognition of the *Mycoblastus fucatus* group as the new genus *Violella* (*Tephromelataceae*, *Lecanorales*). *Lichenologist* 43:445–466
- Spribille T, Resl P, Ahti T, Perez-Ortega S et al (2014a) Molecular systematics of the wood-inhabiting, lichen-forming genus *Xylographa* (*Baeomycetales*, *Ostropomycetidae*) with eight new species. *Symb Bot Ups* 37:1–87
- Spribille T, Tønsberg T, Stabentheiner E, Muggia L (2014b) Reassessing evolutionary relationships in the filamentous cyanolichen genus *Spilonema* (*Peltigerales*, *Lecanoromycetes*). *Lichenologist* 46:373–388
- Sri-Indrasutdh V, Boonyuen N, Suetrong S, Chuaseeharonnachai C et al (2010) Wood-inhabiting freshwater fungi from Thailand: *Ascothailandia grenadoidia* gen. et sp. nov., *Canalisporium grenadoidia* sp. nov. with a key to *Canalisporium* species (Sordariomycetes, Ascomycota). *Mycoscience* 51:411–420
- Sri-Indrasutdh V, Tsui CKM, Chuaseeharonnachai C, Yamaguchi K et al (2015) *Helicocentralis hyalina* gen. et sp. nov., an aero-aquatic helicosporous fungus (*Leotiomycetes*, *Ascomycota*) in Thailand. *Mycol Prog* 14:81–93
- Stadler M, Fournier J, Gardt S, Persöhl D (2010a) The phylogenetic position of *Rhopalostroma* as inferred from a polythetic approach. *Persoonia* 25:11–21
- Stadler M, Fournier J, Læssøe T, Chlebicki A et al (2010b) Chemotaxonomic and phylogenetic studies of *Thamnomyces* (*Xylariaceae*). *Mycoscience* 51:189–207
- Stadler M, Fournier J, Læssøe T, Decock C, Persöhl D, Rambold G (2010c) *Ruwenzoria*, a new genus of the *Xylariaceae* from central Africa. *Mycol Prog* 9:169–179
- Stadler M, Kuhnert E, Persöhl D, Fournier J (2013) The *Xylariaceae* as model example for a unified nomenclature following the “One fungus—one name” (1F1N) concept. *Mycology* 4:5–21
- Stadler M, Læssøe T, Fournier J, Decock C et al (2014) A polyphasic taxonomy of *Daldinia* (*Xylariaceae*). *Stud Mycol* 77:1–43
- Stamenković S, Cvijan M (2010) Determination of air pollution zones in knjaževac (south eastern Serbia) by using epiphytic lichens. *Biotechnol Biotechnol Equip* 24:278–283
- Stamenković S, Cvijan M, Arandjelović M (2010) Lichens as bioindicators of air quality in Dimitrovgrad (South-Eastern Serbia). *Archives of Biological Sciences (Serbia)*. *Arch Biol Sci* 62:643–648
- Stchigel AM, Sutton DA, Cano-Lira JF, Cabañas FJ et al (2013) Phylogeny of *Chrysosporia* infecting reptiles: proposal of the new family *Nannizziopsiaceae* and five new species. *Persoonia* 31:86–100

- Stehn SE, Walton JK, Nelson PR, Hampton-Miller CJ, Roland CA (2015) A lichen species list for Denali National Park and Preserve, Alaska, with comments on several new and noteworthy records. *Evansia* 32:195–215
- Steiner U, Leibner S, Schardl CL, Leuchtmann A, Leistner E (2011) *Periglandula*, a new fungal genus within the *Clavicipitaceae* and its association with *Convolvulaceae*. *Mycologia* 103:1133–1145
- Steinmann J, Giraud S, Schmidt D, Sedlacek L et al (2014) Validation of a novel real-time PCR for detecting *Rasamsonia argillacea* species complex in respiratory secretions from cystic fibrosis patients. *New Microbes New Infect* 2:72–78
- Steinmann J, Dittmer S, Houbraken J, Buer J, Rath PM (2016) In vitro activity of isavuconazole against *Rasamsonia* species. *Antimicrob Agents Chemother* 60:6890–6891
- Stenroos S, Huhtinen S, Lesonen A, Palice Z, Printzen C (2009) *Puttea*, gen. nov., erected for the enigmatic lichen *Lecidea margaritella*. *Bryologist* 112:544–557
- Stenroos S, Laukka T, Huhtinen S, Döbbeler P et al (2010) Multiple origins of symbioses between ascomycetes and bryophytes suggested by a five-gene phylogeny. *Cladistics* 26:281–300
- Stepalska D, Wótek J (2009) Intra-diurnal periodicity of fungal spore concentrations (*Alternaria*, *Botrytis*, *Cladosporium*, *Didymella*, *Ganoderma*) in Cracow, Poland. *Aerobiologia* 25:333–340
- Stepanchikova IS, Himelbrant DE, Kukwa M, Kuznetsova ES (2011) New records of lichens and allied fungi from the Leningrad Region, Russia. II. *Folia Cryptogam Est* 48:85–94
- Stepanchikova IS, Gagarina LV, Kataeva OA (2013) New and rare lichens and allied fungi from the Novgorod Region, Russia. *Folia Cryptogam Estonica* 50:49–55
- Stielow B, Hensel G, Strobelt D, Makonde HM et al (2013) *Hoffmannoscyphe*, a novel genus of brightly coloured, cupulate *Pyronemataceae* closely related to *Tricharina* and *Geopora*. *Mycol Prog* 12:675–686
- Stocker-Wörgötter E, Hager A, Elix JA (2009) Intraspecific chemical variation within the crustose lichen genus *Haematomma*: anthraquinone production in selected cultured mycobionts as a response to stress and nutrient supply. *Phytochem Rev* 8:561–569
- Stordeur R, Cezanne R, Eichler M, Heinrich D et al (2015) First records and noteworthy lichens and lichenicolous fungi from Saxony-Anhalt and the western part of the Harz National Park (Lower Saxony). *Herzogia* 28:654–678
- Su YY, Cai L (2012) Polyphasic characterisation of three new *Phyllosticta* spp. *Persoonia* 28:76–84
- Su YY, Cai L (2013) *Rasamsonia composticola*, a new thermophilic species isolated from compost in Yunnan, China. *Mycol Prog* 12:213–221. Su HY, Liu S, Li Y, Cao Y et al. (2011a) *Arthrobotrys latispora*, a new nematode-trapping fungus from southwest China. *Mycotaxon* 117:29–36
- Su HY, Zhang Y, Baral HO, Yang XY et al (2011) Four new species of *Orbiliaceae* from Yunnan, China. *Mycol Prog* 10:373–381
- Su CM, Wang SW, Lee TH, Tzeng WP et al (2013) *Trichodermin* induces cell apoptosis through mitochondrial dysfunction and endoplasmic reticulum stress in human chondrosarcoma cells. *Toxicol Appl Pharmacol* 272:335–344
- Su KM, Xiong WP, Wang Y, Li SH, Xie R, Baima D (2014) *Tuber boumiense*, a new truffle species from Tibet, China. *Mycotaxon* 126:127–132
- Su H, Li QR, Kang JC, Wen TC et al (2016a) *Rosellinia convexa* sp. nov. (*Xylariales*, *Pezizomycotina*) from China. *Mycoscience* 57:164–170
- Su HY, Hyde KD, Maharachchikumbura SSN, Ariyawansa HA et al (2016b) The families *Distoseptisporaceae* fam. nov., *Kirschsteiniotheliaceae*, *Sporormiaceae* and *Torulaceae*, with new species from freshwater in Yunnan Province, China. *Fungal Divers* 80:375–409
- Su HY, Luo ZL, Liu XY, Su XJ et al (2016c) *Lentithecium cangshanense* sp. nov. (*Lentitheciaceae*) from freshwater habitats in Yunnan Province, China. *Phytotaxa* 267:61–69
- Su L, Deng H, Niu YC (2016d) *Phialemoniopsis endophytica* sp. nov., a new species of endophytic fungi from *Luffa cylindrica* in Henan, China. *Mycol Prog* 15:1–7
- Suetrong S, Schoch CL, Spatafora JW, Kohlmeyer J et al (2009) Molecular systematics of the marine *Dothideomycetes*. *Stud Mycol* 64:155–173
- Suetrong S, Sakayaroj J, Phongpaichit S, Jones EBG (2010) Morphological and molecular characteristics of a poorly known marine ascomycete, *Manglicola guatemalensis* (*Jahnulales*: *Pezizomycotina*; *Dothideomycetes*, *Incertae sedis*): new lineage of marine ascomycetes. *Mycologia* 102:83–92
- Suetrong S, Boonyuen N, Pang KL, Ueapattanakit J et al (2011a) A taxonomic revision and phylogenetic reconstruction of the *Jahnulales* (*Dothideomycetes*), and the new family *Manglicolaceae*. *Fungal Divers* 51:163–188
- Suetrong S, Hyde KD, Zhang Y, Bahkali AB, Jones EBG (2011b) *Trematosphaeriaceae* fam. nov. (*Dothideomycetes*, *Ascomycota*). *Cryptog Mycol* 32:343–358
- Suetrong S, Rungjindamai N, Sommai S, Rung-Areerate P et al (2014) *Wiesneromyces* a new lineage of *Dothideomycetes* (*Ascomycota*) basal to *Tubeufiales*. *Phytotaxa* 176:283–297
- Suetrong S, Klaysuban K, Sakayaroj J, Preedanon S et al (2015) *Tirisporellaceae*, a new family in the order *Diaporthales* (*Sordariomycetes*, *Ascomycota*). *Cryptog Mycol* 36:1–12
- Sugui JA, Peterson SW, Figat A, Hansen B et al (2014) Genetic relatedness versus biological compatibility between *Aspergillus fumigatus* and its sister species. *J Clin Microbiol* 52:3707–3721
- Suija A, de los Rios A, Pérez-Ortega S (2015a) A molecular reappraisal of *Abrothallus* species growing on lichens of the order *Peltigerales*. *Phytotaxa* 195:201–226
- Suija A, Ertz D, Lawrey JD, Diederich P (2015b) Multiple origin of the lichenicolous life habit in *Helotiaceae*, based on nuclear ribosomal sequences. *Fungal Divers* 70:55–72
- Suija A, Oja E, Suu A, Marmor L (2015c) New Estonian records: lichenized and lichenicolous fungi. *Folia Cryptogam Est* 52:129–130
- Suija A, Suu A, Lõhmus P (2016) Substrate specificity corresponds to distinct phylogenetic lineages: the case of *Chaenotheca brunneola*. *Herzogia* 29:355–363
- Sultan A, Johnston PR, Park D, Robertson AW (2011) Two new pathogenic ascomycetes in *Guignardia* and *Rosenschiediella* on New Zealand's pygmy mistletoes (*Korthalsella*: *Viscaceae*). *Stud Mycol* 68:237–247
- Summerbell RC, Gueidan C, Schroers HJ, de Hoog GS et al (2011) *Acremonium* phylogenetic overview and revision of *Gliomastix*, *Sarocladium*, and *Trichothecium*. *Stud Mycol* 68:139–162
- Sun X, Guo LD (2010) *Micronematobotrys*, a new genus and its phylogenetic placement based on rDNA sequence analyses. *Mycol Prog* 9:567–574
- Sun BD, Chen AJ, Gao WW, Zhou YG et al (2013) A new endophytic ascomycete associated with the medicinal plant, *Achyranthes bidentata* Blume (*Amaranthaceae*). *Pakistan J Bot* 45:319–323
- Sun JZ, Dong CH, Liu XZ, Liu JK, Hyde KD (2016) *Calcarisporium cordycipiticola* sp. nov., an important fungal pathogen of *Cordyceps militaris*. *Phytotaxa* 268:135–144
- Sun JZ, Hyde KD, Liu XZ, Zhao Q, Maharachchikumbura SSN et al (2017) *Calcarisporium xylariicola* sp. nov. and introduction of *Calcarisporiaceae* fam. nov. in *Hypocreales*. *Mycol Prog* 16:433–445
- Sundin R, Thor G, Frisch A (2012) A literature review of *Arthonia s. lat.* *Biblioth Lichenol* 108:257–290
- Sung GH, Poinar GO, Spatafora JW (2008) The oldest fossil evidence of animal parasitism by fungi supports a Cretaceous

- diversification of fungal–arthropod symbioses. Mol Phylogenet Evol 49:495–502
- Sung GH, Shrestha B, Park KB, Sung JM (2010) Cultural characteristics of *Shimizuomyces paradoxus* collected from Korea. Mycobiology 38:189–194
- Sung GH, Shrestha B, Park KB, Han SK, Sung JM (2011) Enhancing effect of *shimizuomyces paradoxus* on seed germination and seedling growth of canola, plant growth of cucumber, and harvest of tomato. Mycobiology 39:7–11
- Suresh Kumar G, Manoharachary C, Kunwar IK (2009) *Bhadradriomyces*, a new dematiaceous hyphomycetous genus from India. J Mycol Plant Pathol 39:238–240
- Suto Y (2009) Three ascomycetes on leaves of evergreen Ilex trees from Japan: *Rhytisma ilicis-integrale* sp. nov., *R. ilicis-latifoliae*, and *R. ilicis-pedunculosae* sp. nov. Mycoscience 50:357–368
- Sutton BC (1977) Coelomycetes VI. Nomenclature of generic names proposed for Coelomycetes. Mycol Pap 141:1–253
- Sutton BC (1980) The Coelomycetes: Fungi imperfecti with pycnidia, acervuli, and stromata. Commonwealth Mycological Institute, Kew
- Suzuki Y, Hatakeyama S, Harada Y, Tanaka K (2008) *Polystigma fulvum*, a red leaf blotch pathogen on leaves of *Prunus* spp., has the *Polystigmella pallescens* anamorph/andromorph. Mycoscience 49:395–398
- Suzuki F, Yamaguchi J, Koba A, Nakajima T, Arai M (2010) Changes in fungicide resistance frequency and population structure of *Pyricularia oryzae* after discontinuance of MBI-D fungicides. Plant Disease 94:329–334
- Svensson M (2013) Occurrence patterns of dead wood and wood-dependent lichens in managed boreal forest landscapes. Acta Univ Agric Suec 84:1652–6880
- Svensson M, Dahlberg A, Ranius T, Thor G (2013) Occurrence patterns of lichens on stumps in young managed forests. PLoS One 8:p.e62825
- Svensson M, Johansson V, Dahlberg A, Frisch A et al (2016) The relative importance of stand and dead wood types for wood-dependent lichens in managed boreal forests. Fung Ecol 20:166–174
- Swe A, Jeewon R, Pointing SB, Hyde KD (2008) Taxonomy and molecular phylogeny of *Arthrobotrys mangrovispora*, a new marine nematode-trapping fungal species. Bot Mar 51:331–338
- Sweet M, Bulling M, Cerrano C (2015) A novel sponge disease caused by a consortium of micro-organisms. Coral Reefs 34:871–883
- Szczepańska K (2015) New records of rare lichenicolous and lichen-forming fungi from volcanic rocks in SW Poland. Acta Mycol 50:1056–1064
- Szczepańska K, Pruchniewicz D, Sołtysiak J, Kossowska M (2015) Lichen-forming fungi of the genus montanelia in Poland and their potential distribution in central Europe. Herzogia 28:697–712
- Tabar AI, Lizaso MT, García BE, Gómez B et al (2008) Double-blind, placebo-controlled study of *Alternaria alternata* immunotherapy: Clinical efficacy and safety. Pediatr Allergy Immunol 19:67–75
- Tagirdzhanova GM, Kataeva OA, Stepanchikova IS (2014) New lichen records from the Novgorod Region, Russia. Folia Cryptogam Estonica 51:103–107
- Taj-Aldeen SJ, Al-Ansari N, El Shafei S, Meis JF et al (2009) Molecular identification and susceptibility of trichosporon species isolated from clinical specimens in Qatar: Isolation of *Trichosporon dohaense* Taj-Aldeen, Meis & Boekhout sp. nov. J Clin Microbiol 47:1791–1799
- Takamatsu S (2013a) Origin and evolution of the powdery mildews (*Ascomycota, Erysiphales*). Mycoscience 54:75–86
- Takamatsu S (2013b) Molecular phylogeny reveals phenotypic evolution of powdery mildews (*Erysiphales, Ascomycota*). J General Plant Pathol 79:218–226
- Takamatsu S (2014) Molecular phylogeny reveals phenotypic evolution of powdery mildews (*Erysiphales, Ascomycota*). J General Plant Pathol 79:218–226
- Takamatsu S, Havrylenko M, Wolcan SM, Matsuda S, Niinomi S (2008a) Molecular phylogeny and evolution of the genus *Neoerysiphe* (*Erysiphaceae, Ascomycota*). Mycol Res 112:639–649
- Takamatsu S, Inagaki M, Niinomi S, Khodaparast SAM et al (2008b) Comprehensive molecular phylogenetic analysis and evolution of the genus *Phyllactinia* (*Ascomycota: Erysiphales*) and its allied genera. Mycol Res 112:299–315
- Takamatsu S, Ito T, Yamamoto H, Braun U (2008c) *Sawadaea nankinensis* comb. nov.: a powdery mildew fungus of *Acer buergeriana*. Mycoscience 49:161–167
- Takamatsu S, Niinomi S, Harada M, Havrylenko M (2010) Molecular phylogenetic analyses reveal a close coevolutionary relationship between *Podosphaera* (*Erysiphales: Erysiphaceae*) and its rosaceous hosts. Persoonia 24:38–48
- Takamatsu S, Matsuda S, Grigaliūnaitė B (2013) Comprehensive phylogenetic analysis of the genus *Golovinomyces* (*Ascomycota: Erysiphales*) reveals close evolutionary relationships with host plants. Mycologia 105:1135–1152
- Takamatsu S, Ito H, Shiroya Y, Kiss L, Heluta V (2015a) First comprehensive phylogenetic analysis of the genus *Erysiphe* (*Erysiphales, Erysiphaceae*) I. The *Microsphaera* lineage. Mycologia 107:475–489
- Takamatsu S, Kiss L, Heluta V (2015b) First comprehensive phylogenetic analysis of the genus *Erysiphe* (*Erysiphales, Erysiphaceae*) II: the *Uncinula* lineage. Mycologia 107:903–914
- Takamatsu S, Siahuan SAS, Moreno-Rico O, Cabrera de Alvarez MG, Braun U (2016) Early evolution of endoparasitic group in powdery mildews: molecular phylogeny suggests missing link between *Phyllactinia* and *Leveillula*. Mycologia 108:837–850
- Takashima Y, Narisawa K, Hidayat I, Rahayu G (2014) First report on fungal symbionts of *Lycopodiaceae* root from Mount Gede Pangrango National Park Indonesia. J Dev Sustain Agric 9:81–88
- Tam LT, Hoai HT, Thao LP, Huong NT et al (2015) First report of *Microdium phyllanthi* causing powdery mildew on chamber bitter in Vietnam. New Dis Rep 32:32
- Tan YP, Madrid H, Crous PW, Shivas RG (2014) *Johnalcornia* gen. et. comb. nov., and nine new combinations in *Curvularia* based on molecular phylogenetic analysis. Austral Plant Pathol 43:589–603
- Tanaka E (2009) Specific in situ visualization of the pathogenic endophytic fungus *Aciculosporium* take, the cause of witches' broom in bamboo. Appl Environ Microbiol 75:4829–4834
- Tanaka E, Ashizawa T, Sonoda R, Tanaka C (2008) *Vilosclavula virens* gen. nov., comb. nov., the teleomorph of *Ustilaginoidea virens*, the causal agent of rice false smut. Mycotaxon 106:491–501
- Tanaka K, Hirayama K, Yonezawa H, Hatakeyama S et al (2009) Molecular taxonomy of bambusicolous fungi: *Tetrapsphaeriaceae*, a new pleosporalean family with Tetrapsphaea-like anamorphs. Stud Mycol 64:175–209
- Tanaka K, Mel'nik VA, Kamiyama M, Hirayama K, Shirouzu T (2010) Molecular phylogeny of two coelomycetous fungal genera with stellate conidia, *Prosthemium* and *Asterosporium*, on Fagales trees. Botany 88:1057–1071
- Tanaka K, Endo M, Hirayama K, Okane I et al (2011a) Phylogeny of *Discosia* and *Seimatosprium*, and introduction of *Adisciso* and *Immersidiscosia* genera nova. Persoonia 26:85–98

- Tanaka K, Hirayama K, Iqbal SH (2011b) *Massariosphaeria websteri* sp. nov. and several members of the *Pleosporales* noteworthy to Pakistan. *Mycol Balc* 7:77–85
- Tanaka K, Hirayama K, Yonezawa H, Sato G et al (2015) Revision of the *Massarinea* (Pleosporales, Dothideomycetes). *Stud Mycol* 82:75–136
- Tangthirasunun N, Philippe SP, Bhat DJ, Maharanachikumbura SSN, Hyde KD (2014a) *Greeneria saprophytica* sp. nov. on dead leaves of *Syzygium cumini* from Chiang Rai, Thailand. *Phytotaxa* 184:275–282
- Tangthirasunun N, Silar P, Bhat DJ, Chukeatirote E et al (2014b) Morphology and phylogeny of *Pseudorobillarda eucalypti* sp. nov., from Thailand. *Phytotaxa* 176:251–259
- Tangthirasunun N, Philippe SP, Bhat DJ, Maharanachikumbura SSN et al (2015) Morphology and phylogeny of two appended genera of coelomycetes: *Ciliochorella* and *Discosia*. *Sydowia* 67:217–226
- Tanney JB, Seifert KA (2013) *Rasamonia pulvericola* sp. nov., isolated from house dust. *IMA Fungus* 4:205–212
- Tanney JB, Douglas B, Seifert KA (2016) Sexual and asexual states of some endophytic *Phialocephala* species of *Picea*. *Mycologia* 108:255–280
- Taole MM, Burgess TI, Gryzenhout M, Wingfield BD, Wingfield MJ (2012) DNA sequence incongruence and inconsistent morphology obscure species boundaries in the *Teratosphaeria suttonii* species complex. *Mycoscience* 53:270–283
- Tapper SL, Byrne M, Yates CJ, Keppel G et al (2014) Prolonged isolation and persistence of a common endemic on granite outcrops in both mesic and semi-arid environments in southwestern Australia. *J Biogeogr* 41:2032–2044
- Tarasova V, Androsova V, Sonina A, Ahti T (2015) The lichens from the City of Petrozavodsk in the Herbarium of the Botanical Museum, University of Helsinki. *Folia Cryptogam Est* 52:41–50
- Tarasova VN, Sonina AV, Androsova VI, Stepanchikova IS (2016) The lichens of forest rocky communities of the hill *Muroigora* (Arkhangelsk Region, Northwest Russia). *Folia Cryptogam Est* 53:111–121
- Tarigan M, Roux J, Van Wyk M, Tjahjono B, Wingfield MJ (2011) A new wilt and die-back disease of *Acacia mangium* associated with *Ceratocystis manginecans* and *C. acaciivora* sp. nov. in Indonesia. *South Afr J Bot* 77:292–304
- Tarnowski TLB, Palmateer AJ, Crane JH (2013) First report of fruit rot on *Hylocereus undatus* caused by *Bipolaris cactivora* in South Florida. *Austral Plant Pathol* 42:437–440
- Tasanathai K, Thanakitpipattana D, Noisripoon W, Khonsanit A et al (2016) Two new *Cordyceps* species from a community forest in Thailand. *Mycol Prog* 15:1–8
- Taylor K, Andjic V, Barber PA, Hardy GE, Burgess TI (2012) New species of *Teratosphaeria* associated with leaf diseases on *Corymbia calophylla* (Marri). *Mycol Prog* 11:159–169
- Tehler A (2011) *Roccella*, the Sonoran species reviewed. *Biblioth Lichenol* 106:309–318
- Tehler A (2012) Fruticose *Roccellaceae*. *Bryologist* 115:210–214
- Tehler A, Baloch E, Tibell L, Wedin M (2009a) The systematic position of *Schistophoron*. *Biblioth Lichenol* 99:383–392
- Tehler A, Irestedt M, Bungartz F, Wedin M (2009b) Evolution and reproduction modes in the *Roccella galapagoensis* aggregate (*Roccellaceae*, Arthoniales). *Taxon* 58:438–456
- Tehler A, Irestedt M, Wedin M, Ertz D (2010) The Old World *Roccella* species outside Europe and Macaronesia: taxonomy, evolution and phylogeny System. *Biodiversity* 8:223–246
- Tehler A, Ertz D, Irestedt M (2013a) The genus *Dirina* (Roccellaceae, Arthoniales) revisited. *Lichenologist* 45:427–476
- Tehler A, Irestedt M, Ertz D (2013b) *Austroroccella*, a new fruticose genus in the family *Roccellaceae*. *Bryologist* 116:162–168
- Telfer AC, Young MR, Quinn J, Perez K et al (2015) Biodiversity inventories in high gear: DNA barcoding facilitates a rapid biotic survey of a temperate nature reserve. *Biodivers Data* 3:e6313
- Templeton GE (2016) *Alternaria* toxins related to pathogenesis in plants. *Microbial Toxins* 8:169–192
- Tennakoon DS, Hyde KD, Phookamsak R, Wanasinghe DN et al (2016a) Taxonomy and phylogeny of *Juncaceicola* gen. nov. (Phaeosphaeriaceae, Pleosporinae, Pleosporales). *Cryptogam Mycol* 37:135–156
- Tennakoon DS, Hyde KD, Wanasinghe DN, Bahkali AH et al (2016b) Taxonomy and phylogenetic appraisal of *Montagnula jonesii* sp. nov. (Didymosphaeriaceae, Pleosporales) from Italy. *Mycosphere* 7:1346–1356
- Tennakoon DS, Phillips AJL, Phookamsak R, Ariyawansa HA et al (2016c) Sexual morph of *Lasiodiplodia pseudotheobromae* (Botryosphaeriaceae, Botryosphaerales, Dothideomycetes) from China. *Mycosphere* 7:990–1000
- Teran CG, Downes K, Medows M (2014) Fatal *Bipolaris spicifera* infection in an immunosuppressed child. *BMJ Case Reports*:p.bcr2013009703
- Thambugala KM, Ariyawansa HA, Li YM, Boonmee S et al (2014a) *Dothideales*. *Fungal Divers* 68:105–158
- Thambugala KM, Ariyawansa HA, Liu ZY, Chukeatirote E, Hyde KD (2014b) Towards a natural classification of dothideomycetes 6: the genera *Dolabra*, *Placostromella*, *Pleosphaerellula*, *Polysporidiella* and *Pseudotrichia* (Dothideomycetes incertae sedis). *Phytotaxa* 176:55–67
- Thambugala KM, Camporesi E, Ariyawansa HA, Phookamsak R et al (2014c) Phylogeny and morphology of *Phaeosphaeriopsis triseptata* sp. nov., and *Phaeosphaeriopsis glaucopunctata*. *Phytotaxa* 176:238–250
- Thambugala KM, Daranagama DA, Camporesi E, Singtripop C et al (2014d) Multi-locus phylogeny reveals the sexual state of *Tiarosporella* in *Botryosphaeriaceae*. *Cryptogam Mycol* 35:359–367
- Thambugala KM, Singtripop C, Chunfang Y, McKenzie EHC et al (2014e) Towards a natural classification of Dothideomycetes 7: The genera *Allosoma*, *Austroplespora*, *Dangeardiella*, *Griggsia* and *Karschia* (Dothideomycetes incertae sedis). *Phytotaxa* 181:34–46
- Thambugala KM, Fang YC, Camporesi E, Bahkali AH et al (2015a) *Pseudodidymosphaeria* gen. nov. in *Massarinaceae*. *Phytotaxa* 231:271–282
- Thambugala KM, Hyde KD, Tanaka K, Tian Q et al (2015b) Towards a natural classification and backbone tree for *Lophiostomataceae*, *Floricolaceae*, and *Amorosiaceae* fam. nov. *Fungal Divers* 74:199–266
- Thambugala KM, Hyde KD, Eungwanichayapant PD, Romero AI, Liu ZY (2016) Additions to the genus *Rhytidhysteron* in *Hysteriaceae*. *Cryptog Mycol* 37:99–116
- Thambugala KM, Daranagama DA, Phillips AJ, Bulgakov TS et al (2017a) Microfungi on *Tamarix*. *Fungal Divers* 82:239–306
- Thambugala KM, Wanasinghe DN, Phillips AJL, Camporesi E et al (2017b) Mycosphere notes 1–50: Grass (Poaceae) inhabiting Dothideomycetes. *Mycosphere* 8:697–796
- Thell A, Högnabba F, Elix JA, Feuerer T et al (2009) Phylogeny of the cetrarioid core (Parmeliaceae) based on five genetic markers. *Lichenologist* 41:489–511
- Thell A, Crespo A, Divakar PK, Kärnefelt I et al (2012) A review of the lichen family *Parmeliaceae*—history, phylogeny and current taxonomy. *Nordic J Bot* 30:641–664
- Thongkantha S, Jeewon R, Vijaykrishna D, Lumyong S et al (2009) Molecular phylogeny of *Magnaporthe* (Sordariomycetes) with a new species *Ophioceras chiangdaoense* from *Dracaena loureiroi* in Thailand. *Fungal Divers* 34:157–173

- Thüs H, Muggia L, Pérez-Ortega S, Favero-Longo SE et al (2011) Revisiting photobiont diversity in the lichen family *Verrucariaceae* (*Ascomycota*). *Eur J Phycol* 46:399–415
- Thynne E, McDonald MC, Evans M, Wallwork H et al (2015) Reclassification of the causal agent of white grain disorder on wheat as three separate species of *Eutiarosporella*. *Australas Plant Pathol* 44:527–539
- Tian HZ, Yang Z, Wang S, Hou CL, Piepenbring M (2013) A new species and phylogenetic data for *Nematococomyces*. *Botany* 91:592–596
- Tian Q, Chomnunti P, Bhat JD, Alias SA et al (2014) Towards a natural classification of *Dothideomycetes* 5: The genera *Ascostratum*, *Chaetoscutula*, *Ceratocarpia*, *Cystocoleus*, and *Colensoniella* (*Dothideomycetes incertae sedis*). *Phytotaxa* 176:42–54
- Tian Q, Liu JK, Hyde KD, Wanasinghe DN et al (2015) Phylogenetic relationships and morphological reappraisal of *Melanommataceae* (*Pleosporales*). *Fungal Divers* 74:267–324
- Tian Q, Hongsan S, Dai DQ, Alias SA et al (2016a) Towards a natural classification of *Dothideomycetes*: clarification of *Aldona*, *Aldonata* and *Viegasella* (*Parmulariaceae*). *Mycosphere* 7:511–524
- Tian Q, Doilom M, Luo ZL, Chomnunti P et al (2016b) Introducing *Melanoctona tectonae* gen. et sp. nov. and *Minimelanolocus yunnanensis* sp. nov. (*Herpotrichiellaceae*, *Chaetothyriales*). *Cryptog Mycol* 37:477–492
- Tibell L (2014) *Umbilicaria crustulosa* on the Swedish West Coast. *Graphis Scripta* 26:51–53
- Tibell L, Frisch A (2010) New data on crustose mazaediate lichens from tropical Africa. *Biblioth Lichenol* 104:323–332
- Tibell S, Tibell L (2015) Two new species of *Atla* (*Verrucariaceae*). *Lichenologist* 47:93–98
- Tibell L, Frisch A, Thor G (2014) Additions to the calicioid flora of Japan and Korea, with the descriptions of two new species. *Ann Bot Fenn* 51:189–194
- Tibpromma S, Promputtha I, Phookamsak R, Boonmee S et al (2015) Phylogeny and morphology of *Premilcurensis* gen. nov. (*Pleosporales*) from stems of *Senecio* in Italy. *Phytotaxa* 236:40–52
- Tibpromma S, Bhat J, Doilom M, Lumyong S et al (2016a) Three new *Hermatomyces* species (*Lophiotremataceae*) on *Pandanus odorifer* from Southern Thailand. *Phytotaxa* 275:127–139
- Tibpromma S, Boonmee S, Wijayawardene NN, Maharachchikumbura SSN et al (2016b) The holomorph of *Parasarcopodium* (*Stachybotryaceae*), introducing *P. pandanicola* sp. nov. on *Pandanus* sp. *Phytotaxa* 266(4):250–260
- Tibpromma S, Liu JK, Promputtha I, Camporesi E et al (2016c) *Ophiostomulans tanaceti* gen. et sp. nov. (*Phaeosphaeriaceae*) on *Tanacetum* sp. (*Asteraceae*) from Italy. *Mycol Prog* 15:46–57
- Tibpromma S, McKenzie EHC, Karunaratna SC, Xu J, Hyde KD et al (2016d) *Muyocopron garethjonesii* sp. nov. (*Muyocopronales*, *Dothideomycetes*) on *Pandanus* sp. *Mycosphere* 7:1480–1489
- Tibpromma S, Wijayawardene NN, Manamgoda DS, Boonmee S et al (2016e) *Camarosporium arezzoensis* on *Cytisus* sp., an addition to sexual state of *Camarosporium sensu stricto*. *Saudi J Bio Sci* 23:1–8
- Tibpromma S, Daranagama DA, Boonmee S, Promputtha I, Nontachaiyapoom S, Hyde KD (2017a) *Anthostomelloides krabiensis* gen. et sp. nov. (*Xylariaceae*) from *Pandanus odorifer* (*Pandanaceae*). *Turk J Bot* 41:107–116
- Tibpromma S, Hyde KD, Jeewon R, Maharachchikumbura SSN et al (2017b) Fungal diversity notes 491–602: taxonomic and phylogenetic contributions to fungal taxa. *Fungal Divers* 83:1–261
- Timdal E (2008) Studies on *Eschatogonia* (*Ramalinaceae*) in Peru. *Lichenologist* 40:31–38
- Timdal E (2009) *Krogia antillarum* (*Ramalinaceae*), a new lichen species from the West Indies. *Bryol* 112:387–389
- Timdal E (2010) *Porpidinia* (*Porpidiaceae*), a new genus for *Toninia tumidula*. *Bibl Lichenol* 104:333–337
- Tiwari V, Rai AN (2009) New species of *Megalodochium* and *Monochaetia* from India. *Indian Phytopathol* 62:268–272
- Tiwari CK, Jagrati P, Verma RK (2012) A new and rare species of *Phlyctaeniella* from central India. *Mycosphere* 3:450–453
- To-Anun C, Hidayat I, Meeboon J (2010) *Cercospora cristellae*, a new cercosporoid fungus associated with weed *Cristella parasitica* from northern Thailand. *J Agric Technol* 6:331–339
- To-Anun C, Hidayat I, Meeboon J (2011) Genus *Cercospora* in Thailand: taxonomy and phylogeny (with a dichotomous key to species). *Plant Pathol Quar* 1:11–87
- Tokizawa M, Ohmura Y, Moon KH, Takeshit S (2015) *Sarcogyne endopetrophila* (*Acarosporaceae*, Lichenized *Ascomycota*), a New Species from Japan Shokubutsu Kenkyu Zasshi 90:46–51
- Tomaso-Peterson M, Balbalian CJ (2010) First report of *Bipolaris oryzae* causing leaf spot of switchgrass in Mississippi. *Plant Disease* 94:643
- Tonouchi A (2010) Isolation and characterization of a novel facultative anaerobic filamentous fungus from Japanese rice field soil. *Int J Microbiol* 2009:571383
- Tønsberg T (2013) *Xylographa soralifera* (lichenized *Ascomycetes*) new to Switzerland. *Opera Bot* 119:1–61
- Torriani SFF, Penselin D, Knogge W, Felder M et al (2014) Comparative analysis of mitochondrial genomes from closely related *Rhynchosporium* species reveals extensive intron invasion. *Fung Gen Biol* 62:34–42
- Torta L, Lo Piccolo S, Piazza G, Burruano S et al (2015) *Lulwoana* sp., a dark septate endophyte in roots of *Posidonia oceanica* (L.) Delile. *Plant Biology* 17:505–511
- Toxopeus J, Kozera CJ, O'Leary SJ, Garbary DJ (2011) A reclassification of *Mycophyrias ascophylli* (*Ascomycota*) based on nuclear large ribosomal subunit DNA sequences. *Bot Mar* 54:325–334
- Traeger S, Altegoer F, Freitag M, Gabaldon T, Kempken F, Kumar A, Marcket-Houben M, Pöggeler S, Stajich JE, Nowrouzian M (2013) The genome and development-dependent transcriptomes of *Pyronema confluens*: a window into fungal evolution. *PLoS Genet* 9:e1003820
- Trakunyingcharoen T, Lombard L, Groenewald JZ, Cheewangkoon R et al (2014) Mycoparasitic species of *Sphaerellopsis*, and allied lichenicolous and other genera. *IMA Fungus* 5:391–414
- Trakunyingcharoen T, Lombard L, Groenewald JZ, Cheewangkoon R et al (2015) Caulicolous *Botryosphaerales* from Thailand. *Persoonia* 34:87–99
- Trappe JM, Claridge AW, Arora D, Smit WA (2008) Desert truffles of the African Kalahari: ecology, ethnomycology, and taxonomy. *Econ Bot* 62:521
- Trappe MJ, Trappe JM, Bonito GM (2010a) *Kalapuya brunnea* gen. & sp. nov. and its relationship to the other sequestrate genera in *Morchellaceae*. *Mycologia* 102:1058–1065
- Trappe JM, Kovács GM, Claridge AW (2010b) Comparative taxonomy of desert truffles of the Australian outback and the African Kalahari. *Mycol Prog* 9:131–143
- Travadon R, Lawrence DP, Rooney-Latham S, Gubler WD et al (2015) *Cadophora* species associated with wood-decay of grapevine in North America. *Fungal Biol* 119:53–66
- Tripp EA (2015) Lichen inventory of white rocks open space (city of Boulder, Colorado). *Western North Am Nat* 75:301–310
- Tripp EA, Lendemer JC, Harris RC (2010) Resolving the genus *Graphina* Müll. Arg. in North America: new species, new

- combinations, and treatments for *Acanthothecis*, *Carbacanthographis*, and *Diorgyma*. *Lichenologist* 42:55–71
- Trouillas FP, Sosnowski MR, Gubler WD (2010) Two new species of *Diatrypaceae* from coastal wattle in Coorong National Park, South Australia. *Mycosphere* 1:183–188
- Trouillas FP, Pitt WM, Sosnowski MR, Huang R et al (2011) Taxonomy and DNA phylogeny of *Diatrypaceae* associated with *Vitis vinifera* and other woody plants in Australia. *Fungal Divers* 49:203–223
- Trouillas FP, Hand FP, Inderbitzin P, Gubler WD (2015) The genus *Cryptosphaeria* in the western United States: taxonomy, multi-locus phylogeny and a new species, *C. multicontinentalis*. *Mycologia* 107:1304–1313
- Truong C, Divakar PK, Yahr R, Crespo A, Clerc P (2013) Testing the use of ITS rDNA and protein-coding genes in the generic and species delimitation of the lichen genus *Usnea* (*Parmeliaceae*, *Ascomycota*). *Mol Phylogen Evol* 68:357–372
- Tsang CC, Chan JF, Ip PP, Ngan AH et al (2014a) Subcutaneous phaeohyphomycotic nodule due to *Phialemoniopsis hongkongensis* sp. nov. *J Clin Microbiol* 52:3280–3289
- Tsang CH, Chan JFW, Trendell-Smith NJ, Ngan AHY et al (2014b) Subcutaneous phaeohyphomycosis in a patient with Ig G4-related sclerosing disease caused by a novel ascomycete, *Hongkongmyces pedis* gen. et sp. nov.: first report of human infection associated with the family *Lindgomycetaceae*. *Med Mycol* 52:736–747
- Tsang CC, Hui TWS, Lee KC, Chen JHK et al (2016) Genetic diversity of *Aspergillus* species isolated from onychomycosis and *Aspergillus hongkongensis* sp. nov., with implications to antifungal susceptibility testing. *Diagn Microbiol Infect Dis* 84:125–134
- Tsuge T, Harimoto Y, Akimitsu K, Ohtani K et al (2013) Host-selective toxins produced by the plant pathogenic fungus *Alternaria alternata*. *FEMS Microbiol Rev* 37:44–66
- Tsui CK, Berbee ML (2010) Transfer of two *Helicoma* species to *Troposporella* based on molecular and morphological data. *Mycoscience* 51:144–148
- Tsuneda A, Davey ML, Tsuneda I, Currah RS (2010) Two new dothideomycetous endoconidial genera from declining larch. *Botany* 88:471–487
- Tsuneda A, Davey ML, Tsuneda I, Hudgins A, Currah RS (2011a) *Endophoma*, a new didymellaceous endoconidial genus from bat-cave soil. *Mycologia* 103:1146–1155
- Tsuneda A, Hambleton S, Currah RS (2011b) The anamorph genus *Knufia* and its phylogenetically allied species in *Coniosporium*, *Sarcinomyces* and *Phaeococomyces*. *Botany* 89:523–536
- Tsurykau A, Khramchankova V (2011) Lichens from gomel region: a provisional checklist. *Bot Lith* 17:157–163
- Tu EY (2009) *Alternaria* keratitis: clinical presentation and resolution with topical fluconazole or intrastromal voriconazole and topical caspofungin. *Cornea* 28:116–119
- Tufan-Çetin Ö, Sümbül H (2011) Lichens of the Köprülü Canyon National Park in Turkey. *Mycotaxon* 115:536
- Tuovila HA, Larsson P (2011) Three resinicolous North American species of Mycocaliciaceae in Europe with a re-evaluation of *Chaenothecopsis oregana* Rikkinen. *Karstenia* 51:37–49
- Tuovila H, Rikkinen J, Huhtinen S (2012) Nomenclatural corrections in calicioid fungi. *Kastenia* 52:73–74
- Tuovila H, Davey ML, Yan L, Huhtinen S, Rikkinen J (2014) New resinicolous *Chaenothecopsis* species from China. *Mycologia* 106:989–1003
- Tuzen M, Sari A, Mendil D, Soylak M (2009) Biosorptive removal of mercury (II) from aqueous solution using lichen (*Xanthoparmelia conspersa*) biomass: kinetic and equilibrium studies. *J Hazard Mater* 169:263–270
- Twaroch TE, Focke M, Fleischmann K, Balic N et al (2012) Carrier-bound Alt a 1 peptides without allergenic activity for vaccination against *Alternaria alternata* allergy. *Clin Exp Allergy* 42:966–975
- Udayanga D, Liu XZ, McKenzie EHC, Chukeatirote E et al (2011) The genus *Phomopsis*: biology, applications, species concepts and names of common phytopathogens. *Fungal Divers* 50:189–225
- Udayanga D, Liu X, Crous PW, McKenzie EH et al (2012) A multi-locus phylogenetic evaluation of *Diaporthe* (*Phomopsis*). *Fungal Divers* 56:157–171
- Udayanga D, Manamgoda DS, Liu X, Chukeatirote E et al (2013) What are the common anthracnose pathogens of tropical fruits? *Fungal Divers* 61(1):165–179
- Udayanga D, Castlebury LA, Rossman A, Chukeatirote Ekachai, Hyde KD (2014a) Insights into the genus *Diaporthe*: Phylogenetic species delimitation in the *D. eres* species complex. *Fungal Divers* 67:203–229
- Udayanga D, Castlebury LA, Rossman AY, Hyde KD (2014b) Species limits in *Diaporthe*: molecular re-assessment of *D. citri*, *D. cytospora*, *D. foeniculina* and *D. rufidis*. *Persoonia* 32:83–101
- Udayanga D, Castlebury LA, Rossman AY, Chukeatirote E et al (2015) The *Diaporthe* sojae species complex: phylogenetic reassessment of pathogens associated with soybean, cucurbits and other field crops. *Fungal Biol* 119:383–407
- Untereiner WA, Gueidan C, Orr MJ, Diederich P (2011) The phylogenetic position of the lichenicolous ascomycete *Capronia peltigerae*. *Fungal Divers* 49:225–233
- Untereiner WA, Bogale M, Carter A, Hanson SÅ et al (2013) Molecular phylogeny of *Boliniales* (*Sordariomycetes*) with an assessment of the systematics of *Apiorhynchostoma*, *Endoxyla* and *Psseudovalsaria*. *Mycologia* 105:564–588
- Uppadhyay V, Ingle KK, Trivedi S, Upadhyay V, Upadhyay DK (2016) Diversity and distribution of lichens from the monuments of Gwalior division, Madhya Pradesh with special reference to rock porosity and lichen growth. *Trop Plant Res* 3:384–389
- Upadhyay DK, Patel DK (2012) *Rinodina sophodes* (Ach.) Massal.: a bioaccumulator of polycyclic aromatic hydrocarbons (PAHs) in Kanpur City, India. *Environ Monit Assess* 184:229–238
- Upadhyay DK, Nayaka S, Joshi Y (2009) *Ramboldia amarkantakana* (*Lecanoraceae*, *Ascomycota*), a new lichen species from India. *Mycotaxon* 107:239–242
- Urbanavichene I, Urbanavichus G, Mežaka A, Palice Z (2013) New records of lichens and lichenicolous fungi from the Southern Ural Mountains, Russia. II. *Folia Cryptogam Estonica* 50:73–80
- Urbanavichus G, Ismailov A (2016) New records of lichens and lichenicolous fungi from Dagestan, Russia. *Folia Cryptogam Estonica* 53:65–69
- Urbanavichus G, Urbanavichene I (2011) New records of lichens and lichenicolous fungi from the Ural Mountains, Russia. *Folia Cryptogam Estonica* 48:119–124
- Urbanavichus G, Urbanavichene I (2015) New records of lichens and lichenicolous fungi from the NW Caucasus (Russia). *Herzogia* 28:185–192
- Urbanavichus G, Gabibova A, Ismailov A (2011) New records of lichens and lichenicolous fungi for Russia and the Caucasus. *Turkish J Bot* 35:291–297
- Usui E, Takashima Y, Narisawa K (2016) *Cladophialophora inabaensis* sp. nov., a new species among the dark septate endophytes from a secondary forest in Tottori, Japan. *Microbes Environ* 31:357–360
- Vaghefi N, Pethybridge SJ, Ford R, Nicolas ME et al (2012) *Stagonosporopsis* spp. associated with ray blight disease of Asteraceae. *Australas Plant Pathol* 41:675–686

- Valadbeigi T (2014) Lichen flora of the Ilam province, South West Iran. *Mycotaxon* 128:248–255
- Valadbeigi T, Sipman HJ (2010) New records of lichens and lichenicolous fungi from Iran and their biogeographical significance. *Mycotaxon* 113:191–194
- Valadbeigi T, von Brackel W (2011) Two new species of *Lichenostigma* (*Lichenotheliaceae*) from Iran. *Willdenowia* 41:191–195
- Valadbeigi T, Nordin A, Tibell L (2011a) *Megaspora rimisorediata* (*Pertusariales*, *Megasporaceae*), a new sorediate species from Iran and its affinities with *Aspicilia sensu lato*. *Lichenologist* 43:285–291
- Valadbeigi T, Sipman HJ, Rambold G (2011b) The genus *Immersaria* (*Lecideaceae*) in Iran, including *I. iranica* sp. nov. *Lichenologist* 43:203–208
- van den Boom PPG (2009) New *Halecania* species (*Catillariaceae*) from Europe and South America. *Bryologist* 112:827–832
- van den Boom PPG (2010) New or interesting lichens and lichenicolous fungi of Gran Canaria (Canary Islands, Spain). *Willdenowia* 40:359–367
- van den Boom PP (2012) Additions and notes to the checklist of lichens and lichenicolous fungi of Cape Verde. *Österr Z für Pilzkd* 21:5–16
- van den Boom PPG (2015) Lichens and lichenicolous fungi from graveyards of the area of Eindhoven (the Netherlands), with the description of two new species. *Ann des Naturhistorischen Mus Wien Ser B* 117:245–276
- van den Boom PPG (2016) Lichens and lichenicolous fungi of the azores (portugal), collected on são miguel and terceira with the descriptions of seven new species. *Acta Bot Hung* 58:199–222
- van den Boom PPG, Brand AM (2008) New records of lichens, lichenicolous and allied fungi from Belgium and The Netherlands. *Sauteria* 15:95–101
- van den Boom PP, Giralt M (2012) Checklist and three new species of lichens and lichenicolous fungi of the Algarve (Portugal). *Sydowia* 64:149–207
- van den Boom PI, Etayo JA (2014) New records of lichenicolous fungi and lichenicolous lichens from the Iberian Peninsula, with the description of four new species and one new genus. *Opusc Phil* 13:44–79
- van den Boom PP, Sipman HJ (2016) Foliicolous lichens from Suriname and Guyana: new records and three new species. *Folia Cryptogam Estonica* 53:101–110
- van den Boom PPG, Etayo J (2017) Further interesting lichens and lichenicolous fungi from Fuerteventura, Canary Islands (Spain), with three new species and notes on *Mixtoconidium*. *Ascomycete org* 9:124–134
- van den Brink J, Samson RA, Hagen F, Boekhout T, de Vries RP (2012) Phylogeny of the industrial relevant, thermophilic genera *Myceliophthora* and *Corynascus*. *Fungal Divers* 52:197–207
- van den Brink J, Facun K, De Vries M, Stielow JB (2015) Thermophilic growth and enzymatic thermostability are polyphyletic traits within *Chaetomiaceae*. *Fungal Biol* 119:1255–1266
- van Broeck D, Ertz D (2016) *Cryptophaea*, a new genus of byssoid *Arthoniaceae* (lichenized *Ascomycota*) and its phylogenetic position. *Phytotaxa* 261:168–176
- van Broeck D, Luecking R, Damien ER (2014) The foliicolous lichen biota of the Democratic Republic of the Congo, with the description of six new species. *Lichenologist* 46:141–158
- van Nieuwenhuijzen EJ, Miadlikowska JM, Houbraken JA, Adan OC et al (2016) Wood staining fungi revealed taxonomic novelties in *Pezizomyctina*: new order *Superstratomycetales* and new species *Cyanodermella oleoligni*. *Stud Mycol* 85:107–124
- van Vooren N (2009) Note sur *Smardaea ovalispora* comb. nov. (*Pezizales*). *Bull Mycol Bot Dauphiné-Savoie* 49(192):3–38
- van Vooren N (2014) Type-revision of *Lachnea velenovskyi*, a new species in the genus *Spooneromyces* (*Pezizales*). *Ascomycete.org* 6:89–92
- van Wyk M, Wingfield BD, Clegg PA, Wingfield MJ (2009) *Ceratocystis larium* sp. nov., a new species from *Styrax benzoin* wounds associated with incense harvesting in Indonesia. *Persoonia* 22:75–82
- van Wyk M, Wingfield BD, Marin M, Wingfield MJ (2010) New *Ceratocystis* species infecting coffee, cacao, citrus and native trees in Colombia. *Fungal Divers* 40:103–117
- van Wyk M, Wingfield BD, Wingfield MJ (2011) Four new *Ceratocystis* spp. associated with wounds on *Eucalyptus*, *Schizolobium* and *Terminalia* trees in Ecuador. *Fungal Divers* 46:111–131
- Vandepitte P, Gharnawi S, Rechenmann M, Iltis A et al (2014) Draft genome sequence of the pathogenic fungus *Scedosporium apiospermum*. *Genome Announc* 2:e00988
- Varga J, Samson RA, Frisvad JC (2010a) Polyphasic taxonomy of *Aspergillus* section Sparsi. *IMA Fungus* 1:187–195
- Varga J, Samson RA, Frisvad JC (2010b) *Aspergillus* sect. *Aeni* sect. nov., a new section of the genus for *A. karnatakaensis* sp. nov. and some allied fungi. *IMA Fungus* 1:197–205
- Varga J, Frisvad JC, Samson RA (2011) Two new aflatoxin producing species, and an overview of *Aspergillus* section Flavi. *Stud Mycol* 69:57–80
- Vasilyeva LN (2010) New species of pyrenomycetes from eastern Russia: *Endoxylina rufula* (*Diatrypaceae*). *Mycosphere* 1:163–166
- Vasilyeva LN, Stephenson SL (2010) Biogeographical patterns in pyrenomycetous fungi and their taxonomy. 1. The Grayan disjunction. *Mycotaxon* 114:281–303
- Vasilyeva LN, Stephenson SL (2014) Notes on pyrenomycetous fungi in the Mountain Lake area of southwestern Virginia. *Mycosphere* 5:218–227
- Vasilyeva LN, Rossman AY, Farr DF (2008) [2007] New species of the Diaporthales from eastern Asia and eastern North America. *Mycologia* 99:916–923
- Vasilyeva L, Li Y, Stephenson S (2009) Some pyrenomycetous fungi new to China. *Mycotaxon* 109:415–428
- Vasilyeva L, Chernyshev A, Stephenson SL (2010) Pyrenomycetes of the Russian Far East 4: family *Nitschiaceae* (*Coronophorales*, *Ascomycota*). *Mycologia* 102:233–247
- Vasilyeva LN, Chernyshev AV, Stephenson SL (2012) *Neochaetosphaerella thaxteriospora* gen. et sp. nov. and *Tympnopopsis texensis* sp. nov. (*Coronophorales*, *Ascomycota*) from Texas, USA. *Fungal Divers* 52:191–196
- Vasilyeva LN, Chernyshev AV, Stephenson SL, Hyde KD (2013) *Tortulomyces thailandicus* gen. et sp. nov. and *Nitschki siamensis* sp. nov. (*Coronophorales*, *Ascomycota*) from northern Thailand. *Mycoscience* 54:110–115
- Vasilyeva LN, Ma HX, Stephenson SL (2016) Biogeography and taxonomy of pyrenomycetous fungi 3. The area around the Sea of Japan. *Mycotaxon* 126:1–14
- Vaupotić T, Veranic P, Petrović U, Gunde-Cimerman N, Plemenitaš A (2008) HMG-CoA reductase is regulated by environmental salinity and its activity is essential for halotolerance in halophilic fungi. *Stud Mycol* 61:61–66
- Vazquéz-Campos X, Kinsela AS, Waite TD, Collins RN, Neilan BA (2014) *Fodinomyces uranophilus* gen. nov. sp. nov. and *Coniochaeta fodinicola* sp.nov., two uranium mine-inhabiting *Ascomycota* fungi from northern Australia. *Mycologia* 106:1073–1089
- Véliz H, Glassbrook NJ, Daub ME (2008) Mannitol biosynthesis is required for plant pathogenicity by *Alternaria alternata*. *FEMS Microbiol Lett* 285:122–129

- Velmala S, Myllys L, Goward T, Holien H, Halonen P (2014) Taxonomy of *Bryoria* section *Implexae* (*Parmeliaceae*, *Lecanoromycetes*) in North America and Europe, based on chemical, morphological and molecular data. *Ann Bot Fenn* 51:345–371
- Verkley GJM, Hofland-Zijlstra JD (2010) *Rhizodermea veluwensis* gen. et sp. nov. *Persoonia* 24:130–131
- Verkley GJM, Quaedvlieg W, Shin HD, Crous PW (2013) A new approach to species delimitation in *Septoria*. *Stud Mycol* 75:213–305
- Verkley GJ, Dukik K, Renfurm R, Göker M, Stielow JB (2014) Novel genera and species of coniothyrium-like fungi in *Montagnulaceae* (*Ascomycota*). *Persoonia* 32:25–51
- Verma NK, Rai AN (2014) *Distocercospora indica*, a new dematiaceous hyphomycete from central India. *Mycotaxon* 127:97–101
- Verma RK, Sharma N, Soni KK (2008) Forest Fungi of Central India Lucknow. International Book Distributing Co., Lucknow
- Vermeulen M, Gryzenhout M, Wingfield MJ, Roux J (2013) Species delineation in the tree pathogen genus *Celoporthes* (*Cryphonectriaceae*) in southern Africa. *Mycologia* 105:297–311
- Vicente VA, Attili-Angelis D, Pie MR, Queiroz-Telles F et al (2008) Environmental isolation of black yeast-like fungi involved in human infection. *Stud Mycol* 61:137–144
- Vicente VA, Orélis-Ribeiro R, Najafzadeh MJ, Sun J et al (2012) Black yeast-like fungi associated with lethargic crab disease (LCD) in the mangrove-land crab, *Ucides cordatus* (Ocyopodidae). *Vet Microbiol* 158:109–122
- Videira SIR, Groenewald JZ, Braun U, Shin HD, Crous PW (2016) All that glitters is not Ramularia. *Stud Mycol* 83:49–163
- Videira SIR, Groenewald JZ, Nakashima C, Barreto R et al. (2017) *Mycophaerellaceae*—chaos or clarity? *Stud Mycol* (in prep)
- Vieira WA, Michereff SJ, de Morais Jr MA, Hyde KD et al (2014) Endophytic species of *Colletotrichum* associated with mango in northeastern Brazil. *Fungal Divers* 67:181–202
- Villalta CF, Jacobson DJ, Taylor JW (2009) Three new phylogenetic and biological *Neurospora* species: *N. hispaniola*, *N. metzenbergii* and *N. perkinsii*. *Mycologia* 101:777–789
- Visagie CM, Yilmaz N, Frisvad JC, Houbraken J et al (2015) Five new *Talaromyces* species with ampulliform-like phialides and globose rough walled conidia resembling *T. verruculosus*. *Mycoscience* 56:486–502
- Visagie CM, Yilmaz N, Renaud JB, Sumarah MW et al (2017) A survey of xerophilic *Aspergillus* from indoor environments, including descriptions of two new section *Aspergillus* species producing eurotium-like sexual states. *MycoKeys* 19:1–30
- Visentin I, Gentile S, Valentino D, Gonthier P et al (2012) *Gnomoniopsis castanea* sp. nov. (*Gnomoniaceae*, *Diaporthales*) as the causal agent of nut rot in sweet chestnut. *J Plant Pathol* 94:411–419
- Vitoria NS, Cavalcanti MAQ, Hyde KD, Bezerra JL (2011) *Arecomyces* new to Brazil, including *A. attaleae* sp. nov. *Cryptogam Mycol* 32:103–108
- Vitoria NS, Cavalcanti MA, Hyde KD, Bezerra JL (2012) *Brunneiapiospore brasiliensis* sp. nov. (*Clypeosphaeriaceae*) on palms from Brazil. *Nova Hedw* 94:245–250
- Voglmayr H, Jaklitsch WM (2008) *Prostheciaceae* species with *Stegonsporium* anamorphs on Acer. *Mycol Res* 112:885–905
- Voglmayr H, Jaklitsch WM (2011) Molecular data reveal high host specificity in the phylogenetically isolated genus *Massaria* (*Ascomycota*, *Massariaceae*). *Fungal Divers* 46:133–170
- Voglmayr H, Jaklitsch WM (2014) *Stilboporaceae* resurrected: generic reclassification and speciation. *Persoonia* 33:61–82
- Voglmayr H, Park MJ, Shin HD (2011) *Spiroplana centripeta* gen. & sp. nov., a leaf parasite of *Philadelphus* and *Deutzia* with a remarkable aeroaquatic conidium morphology. *Mycotaxon* 116:203–216
- Voglmayr H, Rossman AY, Castlebury LA, Jaklitsch WM (2012) Multigene phylogeny and taxonomy of the genus *Melanconiella* (*Diaporthales*). *Fungal Divers* 57:1–44
- Voglmayr H, Gardiennet A, Jaklitsch WM (2016) *Asterodiscus* and *Stigmatodiscus*, two new apothecial dothideomycete genera and the new order *Stigmatodiscales*. *Fungal Divers* 80:271–284
- Voglmayr H, Castlebury LA, Jaklitsch WM (2017) *Juglanconis* gen. nov. on *Juglandaceae*, and the new family *Juglanconidaceae* (*Diaporthales*). *Persoonia* 38:136–155
- von Arx JA, Müller E (1954) Die Gattungen der amerosporen Pyrenomyceten. Beiträge zur Kryptogamenflora der Schweiz 11:1–434
- von Brackel W, Etayo J, Lechat C (2012) Notes on three lichenicolous species of *Acremonium* including two new species. *Mycosphere* 3:854–862
- Vondrák J, Liška J (2013) Lichens and lichenicolous fungi from the Retezat Mts and overlooked records for the checklist of Romanian lichens. *Herzogia* 26:293–305
- Vondrák J, Khodosovtsev A, Soun J, Vondráková O (2012) Two new European species from the heterogeneous *Caloplaca holocarpa* group (*Teloschistaceae*). *Lichenologist* 44:73–89
- Vondrák J, Frolov I, Davydov EA, Urbanavichene I et al (2016) The extensive geographical range of several species of *Teloschistaceae*: evidence from Russia. *Lichenologist* 48:171–189
- Voitsekovich A, Dymytrova L, Nadyeina O (2011) Photobiont composition of some taxa of the genera *Micarea* and *Placynthiella* (*Lecanoromycetes*, lichenized *Ascomycota*) from Ukraine. *Folia Cryptogam Estonica* 48:135–148
- Vu AL, Dee MM, Gwinn KD, Ownley BH (2011) First report of spot blotch and common root rot caused by *Bipolaris sorokiniana* on switchgrass in Tennessee. *Plant Dis* 95:1195
- Vu AL, Dee MM, Gualandi RJ Jr, Huff S et al (2015) First report of leaf spot caused by *Bipolaris spicifera* on switchgrass in the United States. *Phytopathology* 105:1146–1154
- Vujanovic V, Goh YK (2009) *Sphaerodes mycoparasitica* sp. nov., a new biotrophic mycoparasite on *Fusarium avenaceum*, *F. graminearum* and *F. oxysporum*. *Mycol Res* 113:1172–1180
- Walker DM, Castlebury LA, Rossman AY, Sogonov MV et al (2010) Systematics of genus *Gnomoniopsis* (*Gnomoniaceae*, *Diaporthales*) based on a three gene phylogeny, host associations and morphology. *Mycologia* 102:1479–1496
- Walker DM, Castlebury LA, Rossman AY, Mejia LC et al (2012) Phylogeny and taxonomy of *Ophiognomonia* (*Gnomoniaceae*, *Diaporthales*), including twenty-five new species in this highly diverse genus. *Fungal Divers* 57:85–147
- Walker DM, Lawrence BR, Wooten JA, Rossman AY et al (2014) Five new species of the highly diverse genus *Plagiostoma* (*Gnomoniaceae*, *Diaporthales*) from Japan. *Mycol Prog* 13:1057–1067
- Walsh E, Luo J, Zhang N (2014) *Acidomelania panicicola* gen. et sp. nov. from switchgrass roots in acidic New Jersey pine barrens. *Mycologia* 106:856–864
- Walsh E, Luo J, Naik A, Preteroti T, Zhang N (2015) *Barrenia*, a new genus associated with roots of switchgrass and pine in the oligotrophic pine barrens. *Fungal Biol* 119:1216–1225
- Wanasinghe DN, Jones EBG, Camporesi E, Boonmee S et al (2014a) An exciting novel member of *Lentiteliaceae* in Italy from *Clematis vitalba*. *Cryptogam Mycol* 35(4):323–337
- Wanasinghe DN, Jones EBG, Camporesi E, Boonmee S et al (2014b) *Dematiopleospora mariae* gen. sp. nov., from *Ononis Spinosa* in Italy. *Cryptog Mycol* 35:105–117
- Wanasinghe DN, Jones EBG, Camporesi E, Mortimer PE et al (2015) The genus *Murispora*. *Cryptog Mycol* 36:419–448

- Wanasinghe DN, Jones EBG, Camporesi E, Dissanayake AJ et al (2016a) Taxonomy and phylogeny of *Laburnicola* gen. nov. and *Paramassariosphaeria* gen. nov. (*Didymosphaeriaceae*, *Massarinea* sp., *Pleosporales*). *Fungal Biol* 120:1354–1373
- Wanasinghe DN, Jones EBG, Dissanayake AJ, Hyde KD (2016b) Saprobic *Dothideomycetes* in Thailand: *Vaginatispora appendiculata* sp. nov. (*Lophiostomataceae*) introduced based on morphological and molecular data. *Stud Fungi* 1:56–68
- Wanasinghe DN, Camporesi E, Hu DM (2016c) *Neoleptosphaeria jonesii* sp. nov., a novel saprobic sexual species, in *Leptosphaeriaceae*. *Mycosphere* 7:1368–1377
- Wanasinghe DN, Hyde KD, Konta S, To-Anun C et al (2017a) Saprobic *Dothideomycetes* in Thailand: *Neoaquastroma* gen. nov. (*Parabambusicolaceae*) introduced based on morphological and molecular data. *Phytotaxa* 302:133–144
- Wanasinghe DN, Phookamsak R, Jeewon R, Li WJ et al (2017b) Fenestellaceae with descriptions of new *Fenestella* species and *Neocucurbitaria* gen. nov. *Mycosphere* 8:397–414
- Wang LS, McCune B (2010) Contributions to the lichen flora of the Hengduan mountains, China 1. Genus *Pseudephebe* (lichenized Ascomycota, *Parmeliaceae*). *Mycotaxon* 113:431–437
- Wang YZ, Huang CL (2015) A new species of *Urnula* (*Sarcosomataceae*, *Pezizales*) from Taiwan. *Nova Hedw* 101:49–55
- Wang HK, Hyde KD, Soytong K, Lin FC (2008a) Fungal diversity on fallen leaves of *Ficus* in northern Thailand. *J Zhejiang Univ Sci B* 9:835–841
- Wang R, Yang XY, Han HB (2008b) Influence of amino acids on traps forming of nematode-trapping fungus-*Arthrobotrys oligospora*. *Heilongjiang Anim Sci Vet Med* 7:3
- Wang Y, Bao Y, Shen D, Feng W et al (2008c) Biocontrol of *Alternaria alternata* on cherry tomato fruit by use of marine yeast *Rhodosporidium paludigenum* Fell & Tallman. *Int J Food Microbiol* 123:234–239
- Wang MM, Jin LT, Jiang CX, Hou CL (2009a) *Rhytisma huangshanense* sp. nov. described from morphological and molecular data. *Mycotaxon* 108:73–82
- Wang W, McGhee D, Gibas CFC, Tsuneda A, Currah RS (2009b) *Phialocephala urceolata*, sp. nov., from a commercial, water-soluble heparin solution. *Mycologia* 101:136–141
- Wang W, Wang Y, Tao H, Peng X et al (2009c) Cerebrosides of the halotolerant fungus *Alternaria raphani* isolated from a sea salt field. *J Nat Prod* 72:1695–1698
- Wang Y, Yu T, Li Y, Cai D et al (2009d) Postharvest biocontrol of *Alternaria alternata* in Chinese winter jujube by *Rhodosporidium paludigenum*. *J Appl Microbiol* 107:1492–1498
- Wang H, Luo H, Li J, Bai Y et al (2010a) An alpha-galactosidase from an acidophilic *Bispora* sp. MEY-1 strain acts synergistically with beta-mannanase. *Bioresour Technol* 101:8376–8382
- Wang J, Xu S, Strongman DB (2010b) Two new *Harpellales* inhabiting the digestive tracts of midge larvae and other trichomycetes from Tianshan Mountains, China. *Mycologia* 102:135–141
- Wang XY, Joshi Y, Hur JS, Oh SO, Wang LS (2010c) Taxonomic studies on the lichen flora of southwestern China (1). *Pilophorus yunnanensis* sp. nov. (*Cladoniaceae*). *Bryologist* 113:345–349
- Wang XY, Joshi Y, Oh SO, Hur JS, Wang LS (2011a) *Pilophorus fruticosus* (*Cladoniaceae*), a new species from south-western China. *Lichenologist* 43:137–140
- Wang XY, Wang LS, Kawakami H, Hur SJ (2011b) The genus *Pilophorus* (Lichenized Ascomycota, *Cladoniaceae*) from China. *菌物学报* 30:889–894
- Wang YL, Abdulla A, Abbas A (2012) *Anzina*, a lichen genus new to China. *Arid Zone Res* 1:029
- Wang J, Wang R, Yang XY (2013a) Efficacy of an *Arthrobotrys oligospora* N mutant in nematode-trapping larvae after passage through the digestive tract of sheep. *Vet Microbiol* 161:359–361
- Wang LS, Wang XY, Lumbsch HT (2013b) Eight lecanoroid lichen species new to China. *Cryptog Mycol* 34:343–348
- Wang QX, Yang JL, Qi QY, Bao L et al (2013c) 3-Anhydro-6-hydroxy-ophiobolin A, a new sesquiterpene inhibiting the growth of methicillin-resistant *Staphylococcus aureus* and inducing the cell death by apoptosis on K562, from the phytopathogenic fungus *Bipolaris oryzae*. *Bioorg Med Chem Lett* 23(12):3547–3550
- Wang S, Taylor J, Hou CL (2013d) Species of *Rhytismatales* on *Berberis* from China. *Mycol Prog* 12:629–635
- Wang Y, Chen XR, Yang CD (2013e) A new *Stigmella* species associated with *Lycium* leaf spots in northwestern China. *Mycotaxon* 122:69–72
- Wang S, Cannon P, Li ZJ, Hou CL (2014a) Multigene phylogenetic analysis detects cryptic species of *Tryblidiopsis* in China. *Mycologia* 106:95–104
- Wang S, Kirschner R, Wang YW, Hou CL (2014b) Hypohelion from China. *Mycol Prog* 13:781–789
- Wang XY, Gu HL, Li Q, Wang LH, Lin YR (2014c) A new species of *Coccomyces* with dimorphic paraphyses. *Mycotaxon* 129:297–302
- Wang Y, Shao J, Yao S, Zhang S et al (2014d) Study on the antithrombotic activity of *Umbilicaria esculenta* polysaccharide. *Carbohydr Polym* 105:231–236
- Wang L, Groenewald M, Wang QM, Boekhout T (2015a) Reclassification of *Saccharomyces sinensis*, Proposal of *Yueomyces sinensis* gen. nov., comb. nov. within *Saccharomycetaceae* (*Saccharomycetales*, *Saccharomycotina*). *PLoS ONE* 10:e0136987
- Wang X, Gao Q, Bao J (2015b) Transcriptional analysis of *Amorphotheca resiniae* ZN1 on biological degradation of furfural and 5-hydroxymethylfurfural derived from lignocellulose pretreatment. *Biotechnol Biofuels* 8:1
- Wang XH, Xu L, Jia ZF (2015c) The lichen genus *Leiorreuma* in China. *Mycotaxon* 130:247–251
- Wang XY, Goffinet B, Liu D, Liang MM et al (2015d) Taxonomic study of the genus *Anzia* (*Lecanorales*, lichenized Ascomycota) from Hengduan Mountains, China. *Lichenologist* 47:99–115
- Wang Y, Hyde KD, McKenzie EH, Jiang YL et al (2015e) Overview of *Stachybotrys* (*Memnoniella*) and current species status. *Fungal Divers* 71:17–83
- Wang LS, Liu D, Shi HX, Zhang YY et al (2015f) *Alectoria spinosa*, a new lichen species from Hengduan Mountains, China. *Mycosphere* 6:159–164
- Wang QM, Zhang YH, Wang B, Wang L (2016a) *Talaromyces neofusisporus* and *T. qii*, two new species of section *Talaromyces* isolated from plant leaves in Tibet, China. *Sci Rep* 6:18622
- Wang TW, Haelewaters D, Pfister DH (2016b) Long-hidden in Thaxter's treasure trove: *Laboulbenia camerunensis* sp. nov. parasitic on African Curculionidae. *Mycotaxon* 131:613–619
- Wang X, Huhtinen S, Hansen K (2016c) Multilocus phylogenetic and coalescent-based methods reveal dilemma in generic limits, cryptic species, and a prevalent intercontinental disjunct distribution in *Geopyxis* (*Pyronemataceae* sl., *Pezizomycetes*). *Mycologia* 19:16–20
- Wang XW, Houben J, Groenewald JZ, Meijer M et al (2016d) Diversity and taxonomy of *Chaetomium* and chaetomium-like fungi from indoor environments. *Stud Mycol* 84:145–224
- Wang YZ, Huang CL, Wei JL (2016e) Two new species of *Sarcoscypha* (*Sarcoscyphaceae*, *Pezizales*) from Taiwan. *Phytotaxa* 245:169–177

- Wang RX, Luo ZL, Hyde KD, Bhat DJ et al (2016f) New species and records of *Dictyochirospora* from submerged wood in northwestern Yunnan, China. *Mycosphere* 7:1357–1367
- Waxman KD, Bergstrom GC (2015) First report of a leaf spot caused by *Bipolaris oryzae* on switchgrass in New York. *Curr Res Fungal Pathog Assoc Rice* 8:482–501
- Wedén C, Larsson S, Burman R, Backlund A (2009) The edible truffle *Choiromyces venosus* and its use in Sweden. *Acta Bot Yunnanica* 31:94–96
- Wedin M, Högnabba F, Goward T (2009a) A new species of *Sphaerophorus*, and a key to the family *Sphaerophoraceae* in western North America. *Bryologist* 112:368–374
- Wedin M, Wiklund E, Jørgensen PM, Ekman S (2009b) Slippery when wet: phylogeny and character evolution in the gelatinous cyanobacterial lichens (*Peltigerales*, Ascomycetes). *Mol Phylogenetics Evol* 53:862–871
- Wedin M, Jørgensen PM, Ekman S (2011) *Vahliellaceae*, a new family of cyanobacterial lichens (*Peltigerales*, Ascomycetes). *Lichenologist* 43:67–72
- Weerakoon G, Aptroot A (2016) Nine new lichen species and 64 new records from Sri Lanka. *Phytotaxa* 280:152–162
- Weerakoon G, Aptroot A, Lumbsch HT, Wolseley PA et al (2012a) New molecular data on *Pyrenulaceae* from Sri Lanka reveal two well-supported groups within this family. *Lichenologist* 44:639–647
- Weerakoon G, Wijeyaratne SC, Wolseley PA, Plata ER et al (2012b) Six new species of *Graphidaceae* from Sri Lanka. *Bryologist* 115:74–83
- Weerakoon G, Jayalal U, Wijesundara S, Karunaratne V, Lücking R (2015) Six new *Graphidaceae* (lichenized Ascomycota: Ostropales) from Horton Plains National Park, Sri Lanka. *Nova Hedw* 101:77–88
- Weerakoon G, Wolseley PA, Arachchige O, da Silva E et al (2016) Eight new lichen species and 88 new records from Sri Lanka. *Bryologist* 119:131–142
- Węgrzyn M, Wietrzyk P, Adamska E, Nicia P (2015) New records of driftwood lichens in the Kaffiøya Plain (NW Spitsbergen, Svalbard). *Pol Polar Res* 36:189–195
- Wei LX, Zhang HX, Tan JL, Chu YS et al (2011) Arthrobotrisins A-C, oligosporons from the nematode-trapping fungus *Arthrobotrys oligospora*. *J Nat Prod* 74:1526–1530
- Weir A (2008) The genus *Prolixandromyces* (*Laboulbeniales*) in the Old World. *Alico* 26:29–35
- Weir BS, Johnston PR, Damm U (2012) The *Colletotrichum gloeosporioides* species complex. *Stud Mycol* 73:115–180
- Weisenborn JL, Kirschner R, Piepenbring M (2010) A new darkly pigmented and keratinolytic species of *Acremonium* (Hyphomycetes) with relationship to the *Plectosphaerellaceae* from human skin and nail lesions in Panama. *Nova Hedwig* 90:457–468
- Wen TC, Zhu RC, Kang JC, Huang MH et al (2013) *Ophiocordyceps xuefengensis* sp. nov. from larvae of *Phassus nodus* (Hepialidae) in Hunan Province, southern China. *Phytotaxa* 123:41–50
- Wen TC, Xiao YP, Li WJ, Kang JC, Hyde KD (2014) Systematic analyses of *Ophiocordyceps ramosissimum* sp. nov., a new species from a larvae of Hepialidae in China. *Phytotaxa* 161:227–234
- Wen T, Zha LS, Xiao YP, Wang Q, Kang JC, Hyde KD (2015) *Metacordyceps shibinensis* sp. nov. from larvae of Lepidoptera in Guizhou Province, southwest China. *Phytotaxa* 226:51–62
- Wen TC, Xiao YP, Zha LS, Hyde KD, Kang JC (2016a) Multigene phylogeny and morphology reveal a new species, *Ophiocordyceps tettigonia*, from Guizhou Province, China. *Phytotaxa* 280:141–151
- Wen TC, Wei DP, Long FY, Zeng XY, Kang JC (2016b) Multigene phylogeny and HPLC analysis reveal fake *Ophiocordyceps sinensis* in markets. *Mycosphere* 7:853–867
- Wendt L, Sir EB, Kuhnert E, Heitkämper S et al (2017) Resurrection and emendation of the *Hypoxylaceae*, recognised from a multigene phylogeny of the *Xylariales*. *Mycol Prog*. doi:10.1007/s11557-017-1311-3
- Westberg M, Sohrabi M (2012) A conspectus of the lichen genus *Candelariella* (*Candelariaceae*, Ascomycota) in Southwest Asia with emphasis on Iran. *Nova Hedw* 95:531–546
- Westberg M, Frödén P, Wedin M (2009) A monograph of the genus *Placomaronea* (Ascomycota, Candelariales). *Lichenologist* 41:513–527
- Westberg M, Millanes AM, Knudsen K, Wedin M (2015) Phylogeny of the *Acarosporaceae* (Lecanoromycetes, Ascomycota, Fungi) and the evolution of carbonized ascomata. *Fungal Divers* 70:145–158
- Wetmore CM (2009) Lichens collected in the Big Horn Mountains, Wyoming. *Evansia* 26:10–14
- Whalley MA, Khalil AM, Wei TZ, Yao YJ, Whalley AJ (2010) A new species of *Engleromyces* from China, a second species in the genus. *Mycotaxon* 112:317–323
- Whitton SR, Mckenzie EHC, Hyde KD (2012a) Anamorphic fungi associated with *Pandanaceae*. In: Hyde KD, Whitton SR, Mckenzie EHC (eds) *Fungi associated with Pandanaceae*. Springer, Dordrecht
- Whitton SR, Mckenzie EHC, Hyde KD (2012b) Teleomorphic microfungi associated with *Pandanaceae*. In: Whitton SR, Mckenzie EH, Hyde KD (eds) *Fungi associated with Pandanaceae*. Springer, Dordrecht, pp 23–124
- Wicklow DT, Rogers KD, Dowd PF, Gloer JB (2011) Bioactive metabolites from *Stenocarpella maydis*, a stalk and ear rot pathogen of maize. *Fungal Biol* 115:133–142
- Wieczorek A, Łysko A (2012) Lichen biota of the Wolin Island (Poland). *Biodivers Res Conserv* 25:83–89
- Wieschollek D, Hellemann S, Baral HO, Richter T (2011) *Roseodiscus formosus* spec. nov. ein Bryophiler Pionier mit falschem Namen. *Z für Mykol* 77:161–174
- Wiewióra B (2012) The effect of seed health of perennial ryegrass (*Lolium perenne* L.) on germination capacity. *Plant Breed Seed Sc* 65:51–62
- Wight WD, Kim KH, Lawrence CB, Walton JD (2009) Biosynthesis and role in virulence of the histone deacetylase inhibitor depudecin from *Alternaria brassicicola*. *Mol Plant-Microbe Interact* 22:1258–1267
- Wijayawardene DNN, Mckenzie EHC, Hyde KD (2012) Towards incorporating anamorphic fungi in a natural classification—checklist and notes for 2011. *Mycosphere* 3:157–228
- Wijayawardene DNN, Song Y, Bhat DJ, Mckenzie EHC et al (2013a) *Wojnowicia viburni* sp. nov. from China and its phylogenetic placement. *Sydowia* 65(1):181–190
- Wijayawardene NN, Camporesi E, Song Y, Dai DQ et al (2013b) Multigene analyses reveal taxonomic placement of *Scolicosporium minkevicii* in Phaeosphaeriaceae (Pleosporales). *Cryptogam Mycol* 34(4):357–366
- Wijayawardene NN, Bhat DJ, Hyde KD, Camporesi E et al (2014a) *Camarosporium sensu stricto* in Pleosporinae, Pleosporales with two new species. *Phytotaxa* 183:16–26
- Wijayawardene NN, Camporesi E, Bhat DJ, Song Y et al (2014b) *Macrodiplodiopsis* in Lophiostomataceae, Pleosporales. *Phytotaxa* 176:192–200
- Wijayawardene NN, Crous PW, Kirk PM, Hawksworth DL et al (2014c) Naming and outline of *Dothideomycetes*—2014 including proposals for the protection or suppression of generic names. *Fungal Divers* 69:1–55

- Wijayawardene NN, Hyde KD, Bhat DJ, Camporesi E et al (2014d) Camarosporium-like species are polyphyletic in *Pleosporales*; introducing *Paracamarosporium* and *Pseudocamarosporium* gen. nov. in *Montagnulaceae*. Cryptog Mycol 35:177–219
- Wijayawardene NN, Hyde KD, Camporesi E, Bhat DJ et al (2014e) *Homortomyces tamaricis* sp. nov. and convergent evolution of *Homortomyces* and *Stilbospora*. Phytotaxa 176:156–163
- Wijayawardene NN, Hyde KD, Bhat DJ, Goonasekara ID, Nadeeshan D et al (2015) Additions to brown spored coelomycetous taxa in *Massarinaceae*, *Pleosporales*: Introducing *Phragmocamarosporium* gen. nov. and *Suttonomyces* gen. nov. Cryptog Mycol 36:213–224
- Wijayawardene NN, Goonasekara ID, Camporesi E, Wang Y, An YL (2016a) Two new *Seimatosprium* species from Italy. Mycosphere 7:204–213
- Wijayawardene NN, Hyde KD, Wanasinghe DN, Papizadeh M et al (2016b) Taxonomy and phylogeny of dematiaceous coelomycetes. Fungal Divers 77:1–316
- Wijayawardene NN, Tibpromma S, Hyde KD, An YL et al (2016c) Additions to *Pseudocamarosporium*; two new species from Italy. Stud Fungi 1:1–10
- Wijayawardene NN, Hyde KD, Tibpromma S, Wanasinghe DN et al. (2017) Towards incorporating asexual fungi in a natural classification: checklist and notes 2012–2016. Mycosphere (accepted)
- Wikee S, Udayanga D, Crous PW, Chukeatirote E et al (2011) *Phyllosticta*—an overview of current status of species recognition. Fungal Divers 51:43–61
- Wikee S, Lombard L, Crous PW, Nakashima C et al (2013a) *Phyllosticta capitalensis*, a widespread endophyte of plants. Fungal Divers 60:91–105
- Wikee S, Lombard L, Nakashima C, Motohashi K et al (2013b) A phylogenetic re-evaluation of *Phyllosticta* (*Botryosphaeriales*). Stud Mycol 76:1–29
- Wirth V, Hauck M, Brackel WV, Cezanne R et al (2011) Checklist of lichens and lichenicolous fungi in Germany. Georg August Univ Göttingen 2:19
- Wiseman MS, Kim YK, Dugan FM, Rogers JD, Xiao CL (2016) A new postharvest fruit rot in apple and pear caused by *Phacidium lacerum*. Plant Dis 100:32–39
- Wolf J, O'Neill NR, Rogers CA, Muilenberg ML, Ziska LH (2010) Elevated atmospheric carbon dioxide concentrations amplify *Alternaria alternata* sporulation and total antigen production. Environ Health Perspect 118:1223
- Wong SS, Ngan AH, Riggs CM, Teng JL et al (2011) Brittle tail syndrome is an emerging infection in horses caused by a keratinolytic fungus *Equicapillimyces hongkongensis* gen. nov., sp. nov. Vet Microbiol 155:399–408
- Wongsawas M, Wang HK, Hyde KD, Lin FC (2009) Two new hyphomycetes from submerged wood collected in China. Sydowia 61:345–351
- Wood AR, Damm U, van der Linde EJ, Groenewald JZ et al (2016) Finding the missing link: resolving the *Coryneliomycetidae* within *Eurotiomycetes*. Persoonia 37:37–56
- Woudenberg JH, Aveskamp MM, De Gruyter J, Spiers AG, Crous PW (2009) Multiple *Didymella* teleomorphs are linked to the *Phoma clematidina* morphotype. Persoonia 22:56–62
- Woudenberg JHC, Groenewald JZ, Binder M, Crous PW (2013) *Alternaria* redefined. Stud Mycol 75:171–212
- Woudenberg JHC, Truter M, Groenewald JZ, Crous PW (2014) Large-spored *Alternaria* pathogens in section *Porri* disentangled. Stud Mycol 79:1–47
- Wrzosek M, Dubiel G, Gorczak M, Pawłowska J et al (2016) New insights on the phylogeny and biology of the fungal ant pathogen *Aegeritella*. J Invertebr Pathol 133:1–7
- Wu YM, Zhang TY (2008) Soil dematiaceous hyphomycetes from Dahingganling Mountain Range, Inner Mongolia, China I. Mycosistema 27:5–15
- Wu YM, Zhang TY (2009) New species of *Phialosporostilbe* and *Pleurothecium* from soil. Mycotaxon 110:1–4
- Wu YM, Zhang TY (2010) Three new species of *Scytalidium* from soil. Mycotaxon 114:205–210
- Wu YM, Zhang TY (2013) New species of *Humicola* and *Endophragmiella* from China. Mycotaxon 121:147–151
- Wu HX, Schoch CL, Boonmee S, Bahkali AH et al (2011) A reappraisal of *Microthyriaceae*. Fungal Divers 51:189–248
- Wu DK, Zhang CP, Zhu CY, Wang YL et al (2013a) Metabolites from carnivorous fungus *Arthrobotrys entomopaga* and their functional roles in fungal predatory ability. J Agric Food Chem 61:4108–4113
- Wu YM, Xu JJ, Wang HF, Zhang TY (2013b) *Geosmithia tibetensis* sp. nov. and new *Gibellulopsis* and *Scopulariopsis* records from Qinghai-Tibet. Mycotaxon 125:59–64
- Wu YM, Xu JJ, Wang HF, Zhang TY (2013c) *Radulidium xigazense* sp. nov., *Rhinocladiella tibetensis* sp. nov., and three new records of *Ramichloridium* from China. Mycotaxon 125:123–130
- Wu HX, Li Y, Ariyawansa HA, Li W et al (2014a) A new species of *Microthyrium* from Yunnan, China. Phytotaxa 176:213–218
- Wu HX, Tian Q, Li WJ, Hyde KD (2014b) A reappraisal of *Microthyriaceae*. Phytotaxa 176:201–212
- Wu SP, Liu YX, Yuan J, Wang Y et al (2014c) *Phyllosticta* species from banana (*Musa sp.*) in Chongqing and Guizhou Provinces, China. Phytotaxa 188:135–144
- Wu Y, Wang SJ, Meng YQ, Tang YP, Lin YR (2015) *Terriera fici* sp. nov. on *Ficus vesculosa* from Hainan Province, China. Mycotaxon 130:1111–1116
- Wu YM, Xu JJ, Wang HF, Zhang TY (2013d) *Radulidium xigazense* sp. nov., *Rhinocladiella tibetensis* sp. nov., and three new records of *Ramichloridium* from China. Mycotaxon 125:123–130
- Wulandari NF, To-Anun C, Hyde KD, Duong LM et al (2009) *Phyllosticta citriasiiana* sp. nov., the cause of Citrus tan spot of *Citrus maxima* in Asia. Fungal Divers 34:23–39
- Wynns AA (2015) Convergent evolution of highly reduced fruiting bodies in *Pezizomycotina* suggests key adaptations to the bee habitat. BMC Evol Biol 15:145
- Xia JW, Ma LG, Ma YR, Castañeda-Ruiz RF, Zhang XG (2013a) *Corynesporopsis curvularioides* sp. nov. and new records of microfungi from southern China. Cryptog Mycol 34:281–288
- Xia JW, Ren SC, Ma LG, Zhang XG (2013b) *Heteroconium bannaense* sp. nov. and a new record of the genus from China. Mycotaxon 121:413–417
- Xia JW, Ma LG, Castañeda-Ruiz RF, Zhang XG (2014a) *Minimelanolocus bicolorata* sp. nov., *Paradendryphiopsis elegans* sp. nov. and *Corynesporella bannaense* sp. nov. from southern China. Mycoscience 55:299–307
- Xia JW, Ma YR, Zhang XG (2014b) New species of *Corynesporopsis* and *Lylea* from China. Sydowia 66:241–248
- Xia JW, Ma YR, Gao JM, Li Z, Zhang XG (2015a) *Codinaea jianfenglingensis* sp. nov. and new records from southern China. Mycotaxon 130:835–841
- Xia JW, Ma YR, Zhang XG (2015b) *Anungitea guangxiensis* and *Ellisembia longchiensis*, two new species from southern China. Mycotaxon 130:41–46
- Xia JW, Ma YR, Gao JM, Zhang XG, Li Z (2016a) A new species of *Blastophragma* and three new records of other dematiaceous hyphomycetes from southern China. Nova Hedw 102:223–232
- Xia JW, Ma YR, Gao JM, Zhang XG, Li Z (2016b) *Atrosynnema*, a new hyphomycete from dead branches in China. Mycotaxon 131:287–290

- Xie J, Strobel GA, Mends MT, Hilmer J et al (2013) *Collophora aceris*, a novel antimycotic producing endophyte associated with douglas maple. *Microb Ecol* 66:784–795
- Xu D, Ondeyka J, Harris GH, Zink D et al (2011) Isolation, structure, and biological activities of fellutamides C and D from an undescribed *Metulocladosporiella* (*Chaetothyriales*) using the genome-wide *Candida albicans* fitness test. *J Nat Prod* 74:1721–1730
- Xu L, Ravnskov S, Larsen J, Nilsson RH, Nicolaisen M (2012) Soil fungal community structure along a soil health gradient in pea fields examined using deep amplicon sequencing. *Soil Biol Biochem* 46:26–32
- Xu X, Zhang Y, Meng Q, Meng K et al (2013) overexpression of a fungal β-mannanase from *Bispora* sp. MEY-1 in maize seeds and enzyme characterization. *PLoS ONE* 8:e56146
- Xu C, Wang CH, Ju LL, Zhang R et al (2015) Multiple locus genealogies and phenotypic characters reappraise the causal agents of apple ring rot in China. *Fungal Divers* 71:215–231
- Yacharoen S, Tian Q, Chomnunti P, Boonmee S et al (2015) *Patellariaceae* revisited. *Mycosphere* 6:290–326
- Yadav S, Bhat DJ (2009a) *Dimastigosporium yanense*, a new coprophilous fungus from the forests of Western Ghats in Karnataka State, India. *Mycotaxon* 107:397–403
- Yadav S, Bhat DJ (2009b) *Lomachashaka gomaya*, a new sporodochial hyphomycete from India. *Mycotaxon* 110:357–361
- Yago JI, Lin CH, Chung KR (2011) The SLT2 mitogen-activated protein kinase-mediated signalling pathway governs conidiation, morphogenesis, fungal virulence and production of toxin and melanin in the tangerine pathotype of *Alternaria alternata*. *Mol Plant Pathol* 12:653–665
- Yamazaki A, Kawasaki H (2014) *Lipomyces chichibuensis* sp. nov., isolated in Japan, and reidentification of the type strains of *Lipomyces kononenkoae* and *Lipomyces spencermartinsiae*. *Int J Syst Evol Microbiol* 64:2566–2572
- Yamazaki A, Toyama K, Nakagiri A (2010) A new acidophilic fungus *Teratosphaeria acidotherma* (*Capnodiales, Ascomycota*) from a hot spring. *Mycoscience* 51:443–455
- Yan JY, Xie Y, Zhang W, Wang Y et al (2013) Species of *Botryosphaeriaceae* involved in grapevine dieback in China. *Fungal Divers* 61:221–236
- Yan JY, Jayawardena MMRS, Goonasekara ID, Wang Y et al (2015) Diverse species of *Colletotrichum* associated with grapevine anthracnose in China. *Fungal Divers* 71:233–246
- Yang SL, Chung KR (2012) The NADPH oxidase-mediated production of hydrogen peroxide (H₂O₂) and resistance to oxidative stress in the necrotrophic pathogen *Alternaria alternata* of citrus. *Mol Plant Pathol* 13:900–914
- Yang SL, Lin CH, Chung KR (2009) Coordinate control of oxidative stress tolerance, vegetative growth, and fungal pathogenicity via the AP1 pathway in the rough lemon pathotype of *Alternaria alternata*. *Physiol Mol Plant Pathol* 74:100–110
- Yang HL, Sun GY, Batzer JC, Crous PW et al (2010) Novel fungal genera and species associated with the sooty blotch and flyspeck complex on apple in China and the United States. *Persoonia* 24:29–37
- Yang GZ, Lu J, Yu Z, Zhang K, Qiao M (2011a) *Uncispora sinensis*, a new species from China. *Mycotaxon* 116:171–174
- Yang J, Wang L, Ji X, Feng Y, Li X et al (2011b) Genomic and proteomic analyses of the fungus *Arthrobotrys oligospora* provide insights into nematode-trap formation. *PLoS Pathog* 7:e1002179
- Yang Y, Chai R, He Y (2012) Early detection of rice blast (*Pyricularia*) at seedling stage in Nipponbare rice variety using near-infrared hyper-spectral image. *Afr J Biotech* 11:6809–6817
- Yang J, Yu Y, Li J, Zhu W et al (2013) Characterization and functional analyses of the chitinase-encoding genes in the nematode-trapping fungus *Arthrobotrys oligospora*. *Arch Microbiol* 195:453–462
- Yang H, Ariyawansa HA, Wu HX, Hyde KD (2014a) The genus *Leptoxypium* (*Capnodiaceae*) from China. *Phytotaxa* 176:174–183
- Yang H, Chomnunti P, Ariyawansa HA, Wu HX et al (2014b) The genus *Phaeosaccardinula* (*Chaetothyriales*) from Yunnan, China, introducing two new species. *Chiang Mai J Sci* 41:873–884
- Yang J, Maharachchikumbura SSN, Bhat DJ, Bahkali AH et al. (2015a) *Aquapteridospora lignicola* gen. et sp. nov., a new hyphomycetous taxon (*Sordariomycetes*) from wood submerged in a freshwater stream. *Cryptog Mycol* 36(4):469–478
- Yang Q, Fan XL, Crous PW, Liang YM, Tian CM (2015b) *Cytospora* from *Ulmus pumila* in northern China. *Mycol Prog* 14:74
- Yang C, Wang J, Xia J, Ma Y et al (2016a) *Phaeomonilia nanningensis* sp. nov. and a new *Craspedodidymum* record from southern China. *Mycotaxon* 131:547–551
- Yang J, Liu JK, Bhat DJ, Hyde KD et al (2016b) New species of *Sporoschisma* (*Chaetosphaeriaceae*) from aquatic habitats in Thailand. *Phytotaxa* 289:147–157
- Yang J, Maharachchikumbura SSN, Bhat DJ, Hyde KD et al (2016c) *Fuscosporellales*, a new order of aquatic and terrestrial *Hypocreomycetidae* (*Sordariomycetes*). *Cryptog Mycol* 37:449–475
- Yang T, Groenewald JZ, Cheewangkoon R, Jami F et al (2016d) Families, genera, and species of *Botryosphaerales*. *Fungal Biol* 121:322–346
- Yang Q, Fan XL, Du Z, Tian CM (2017) *Diaporthe juglandicola* sp. nov. (*Diaporthales, Ascomycetes*), evidenced by morphological characters and phylogenetic analysis. *Mycosphere* 8:817–826
- Yatsyna A (2011) The first contribution to lichens, lichenicolous and allied fungi from Braslav Lakes National Park (NW Belarus). *Botanica Lithuanica* 17:177–184
- Yatsyna A (2014) Lichens from Manor Parks in Minsk Region (Belarus). *Botanica Lithuanica* 20:159–168
- Yatsyna A, Motiejūnaitė J (2015) New and noteworthy lichens To Belarus. *Botanica Lithuanica* 21:57–63
- Yavuz Y, Özdemir Türk A, Böcük H (2015) Epiphytic lichen diversity of the Sündiken mountains (Eskişehir, Turkey). *Herzogia* 28:753–761
- Yazawa T, Kawahigashi H, Matsumoto T, Mizuno H (2013) Simultaneous transcriptome analysis of Sorghum and *Bipolaris sorghicola* by using RNA-seq in combination with de novo transcriptome assembly. *PLoS ONE* 8:e62460
- Yazici K, Aslan A (2016) *Merismatium*, *Porpidia* and *Protoparmelia* spp. new for Turkey and Asia. *Mycotaxon* 131:337–343
- Yazici K, Elix JA, Aslan A (2010) Some parmelioid lichens new to Turkey and Asia. *Mycotaxon* 111:489–494
- Yeshitela K, Fischer E, Killmann D, Serusiaux E (2009) *Aspidothelium hirsutum* (*Thelenellaceae*) and *Caprettia goderei* (*Monoblastiaceae*), two new species of foliicolous lichens from Ethiopia and Kenya. *Bryologist* 112:850–855
- Yilmaz N, Houbraken J, Hoekstra ES, Frisvad JC et al (2012) Delimitation and characterisation of *Talaromyces purpurogenus* and related species. *Persoonia* 29:39–54
- Yilmaz N, Visagie CM, Houbraken J, Frisvad JC, Samson RA (2014) Polyphasic taxonomy of the genus *Talaromyces*. *Stud Mycol* 78:175–341
- Yilmaz N, López-Quintero CA, Vasco-Palacios AM, Frisvad JC et al (2016a) Four novel *Talaromyces* species isolated from leaf litter from Colombian Amazon rain forests. *Mycol Prog* 15:1041–1056
- Yilmaz N, Visagie CM, Frisvad JC, Houbraken J et al (2016b) Taxonomic re-evaluation of species in *Talaromyces* section Islandici, using a polyphasic approach. *Persoonia* 36:37–56

- Yin M, Duong TA, Wingfield MJ, Zhou X, de Beer ZW (2015) Taxonomy and phylogeny of the *Leptographium procerum* complex, including *Leptographium sinense* sp. nov. and *Leptographium longiconidiophorum* sp. nov. *Antonie Van Leeuwenhoek* 107:547–563
- Yonezawa H, Tanaka K (2008) A second species of *Neoheteroceras* and additional characters of the genus. *Mycoscience* 49:152–154
- Yoshiaki KON, Ohmura Y (2014) Regeneration of *Pseudocyphellaria aurata* transplanted on a tree in Japan. *Lichenologist* 46:833–836
- Young CA, Charlton ND, Takach JE, Swoboda GA et al (2014) Characterization of *Epichloë coenophiala* within the US: are all tall fescue endophytes created equal? *Front Chem* 2:95
- Yu ZF, Qiao M, Zhang Y, Qin L, Zhang KQ (2011) *Pseudotriopconidium*, a new anamorph genus connected to *Orbilia*. *Mycologia* 103:164–173
- Yu H, Duan J, Wang B, Jiang X (2012) The function of snodprot in the cerato-platanin family from *Dactyellina cionopaga* in nematophagous fungi. *Biosci Biotechnol Biochem* 76:1835–1842
- Yuan L, Zhao PJ, Ma J, Li GH, Shen YM (2008) Tricycloalternarenes A-E: five new mixed terpenoids from the endophytic fungal strain *Alternaria alternata* Ly83. *Helv Chim Acta* 91:1588–1594
- Yuan ZL, Lin FC, Zhang CL, Kubicek CP (2010) A new species of *Harpophora* (*Magnaportheaceae*) recovered from healthy wild rice (*Oryza granulata*) roots, representing a novel member of a beneficial dark septate endophyte. *FEMS Microbiol Lett* 307:94–101
- Yun HY, Rossman AY (2011) *Tubakia seoraksanensis*, a new species from Korea. *Mycotaxon* 115:369–373
- Zaffarano PL, McDonald BA, Linde CC (2008) Rapid speciation following recent host shifts in the plant pathogenic fungus *Rhynchosporium*. *Evolution* 62:1418–1436
- Zaffarano PL, McDonald BA, Linde CC (2011) Two new species of *Rhynchosporium*. *Mycologia* 103(1):195–202
- Zainuddin N, Alias SA, Lee CW, Ebel N et al (2010) Antimicrobial activities of marine fungi from Malaysia. *Bot Mar* 53:507–513
- Zakeri Z, Gasparyan A, Aptroot A (2016) A new corticolous *Megaspora* (*Megasporaceae*) species from Armenia. *Willdenowia* 46:245–251
- Zambare VP, Christopher LP (2012) Biopharmaceutical potential of lichens. *Pharm Biology* 50:778–798
- Zarabska D, Rosadzinski S (2011) New records of the lichen species *Pycnothelia papillaria* in Poland in the context of threats to species. *Roczniki Akademii Rolniczej w Poznaniu Botanika-Steciana* 15:149–152
- Zare R, Gams W (2016) More white verticillium-like anamorphs with erect conidiophores. *Mycol Prog* 15:993–1030
- Zare R, Asgaria B, Gams W (2010) The species of *Coniolariella* (*Xylariales*). *Mycologia* 102:1383–1388
- Zelski SE, Raja HA, Miller AN, Babosa FR et al (2011a) *Longicollum biappendiculatum* gen. et sp. nov., a new freshwater ascomycete from the Neotropics. *Mycosphere* 2:539–545
- Zelski SE, Raja HA, Miller AN, Shearer CA (2011b) *Chaetorostrum quincemilensis*, gen. et sp. nov., a new freshwater ascomycete and its Taeniola-like anamorph from Peru. *Mycosphere* 5:593–600
- Zelski SE, Raja HA, Miller AN, Shearer CA (2014) Conioscypha peruviana sp. nov., its phylogenetic placement based on 28S rRNA gene, and a report of *Conioscypha gracilis* from Peru. *Mycoscience* 56:319–325
- Zeng JS, de Hoog GS (2008) *Exophiala spinifera* and its allies: diagnostics from morphology to DNA barcoding. *Med Mycol* 46:193–208
- Zeng XY, Wen TC, Chomnunti P, Liu JK et al (2016) *Ceramothyrium longivolcaniforme* sp. nov., a new species of *Chaetothyriaceae* from northern Thailand. *Phytotaxa* 267:51–60
- Zeng ZQ, Zhuang WY (2013a) A new combination of *Allantonectria* (*Hypocreales*). *Mycosistema* 32:294
- Zeng ZQ, Zhuang WY (2013b) Four new taxa of *Ilyonectria* and *Thelonectria* (*Nectriaceae*) revealed by morphology and combined ITS and β-tubulin sequence data. *Phytotaxa* 85:15–25
- Zhang JZ, Li MJ (2009) A new species of *Bipolaris* from the halophyte *Sesuvium portulacastrum* in Guangdong Province, China. *Mycotaxon* 109:289–300
- Zhang BC, Minter DW (2011) *Elaphomyces spinoreticulatus* sp. nov., with notes on Canadian species of *Elaphomyces*. *Can J Bot* 67:909–914
- Zhang YL, Zhang TY (2014) Notes on soil dematiaceous hyphomycetes from Guangxi Province, China. *Mycosistema* 33:793–799
- Zhang Y, Fournier J, Pointing SB, Hyde KD (2008a) Are *Melanomma pulvis-pyrius* and *Trematosphaeria pertusa* congeneric? *Fungal Divers* 33:47–60
- Zhang Y, Jeewon R, Fournier J, Hyde KD (2008b) Multi-gene phylogeny and morphotaxonomy of *Amniculicola lignicola*: a novel freshwater fungus from France and its relationships to the *Pleosporales*. *Mycol Res* 112:1186–1194
- Zhang K, Ma J, Ma LG, Zhang XG (2009a) A new species of *Berkleasmium* from Chongqing, China. *Mycotaxon* 108:5–7
- Zhang R, Yang HL, Sun GY, Li HY et al (2009b) *Strelitziana mali*, a new species causing sooty blotch on apple fruit. *Mycotaxon* 110:477–485
- Zhang Y, Fournier J, Crous PW, Pointing SB, Hyde KD (2009c) Phylogenetic and morphological assessment of two new species of *Amniculicola* and their allies (*Pleosporales*). *Persoonia* 23:48–54
- Zhang Y, Schoch CL, Fournier J, Crous PW et al (2009d) Multi-locus phylogeny of *Pleosporales*: a taxonomic, ecological and evolutionary re-evaluation. *Stud Mycol* 64:85–102
- Zhang Y, Wang HK, Fournier J, Crous PW et al (2009e) Towards a phylogenetic clarification of *Lophiostoma/Massarina* and morphologically similar genera in the *Pleosporales*. *Fungal Divers* 38:225–251
- Zhang Y, Yu ZF, Baral HO, Mo MH, Zhang KQ (2009f) New species and records of *Orbilia* (*Orbiliaceae, Ascomycota*) from China. *Fungal Divers* 36:141–153
- Zhang H, Zhang X, Qi YX, Pu JJ et al (2010a) A new leaf spot disease of banana caused by *Pyricularia angulata*. *J Tropical Org* 2:12
- Zhang Y, Qiao M, Weber E, Baral HO et al (2010b) *Arthrobotrys scaphoides* from China and Europe with a phylogenetic analysis including the type strain. *Mycotaxon* 111:291–300
- Zhang YD, Ma J, Ma LG, Zhang XG (2010c) A new species of *Heteroconium* from Fujian, China. *Mycotaxon* 114:315–318
- Zhang YD, Ma J, Ma LG, Zhang XG (2010d) Taxonomic studies of *Dactyella* from Fujian, China. *Mycotaxon* 114:259–261
- Zhang YD, Ma J, Ma LG, Zhang XG (2010e) Two new species of *Kylinaria* from Fujian, China. *Mycotaxon* 114:367–371
- Zhang H, Jones EBG, Zhou D, Bahkali AH, Hyde KD (2011a) Checklist of freshwater fungi in Thailand. *Cryptog Mycol* 32:199–217
- Zhang M, Ko KT W, Hyde KD (2011b) Towards a monograph of *Dothideomycetes*: Studies on *Diademaceae*. *Cryptog Mycol* 32:115–126
- Zhang N, Zhao S, Shen Q (2011c) A six-gene phylogeny reveals the evolution of mode of infection in the rice blast fungus and allied species. *Mycologia* 103:1267–1276
- Zhang Y, Crous PW, Schoch CL, Bahkali AHKD et al (2011d) A molecular, morphological and ecological re-appraisal of *Venturiales*—a new order of *Dothideomycetes*. *Fungal Divers* 51:249–277

- Zhang Y, Fournier J, Bahkali AH, Hyde KD (2011e) Inflatispora, a novel lignicolous genus of Pleosporales from France. *Sydowia* 63:287–295
- Zhang Y, Yu ZF, Xu J, Zhang KQ (2011f) Divergence and dispersal of the nematode-trapping fungus *Arthrobotrys oligospora* from China. *Environ Microbiol Rep* 3:763–773
- Zhang YD, Ma J, Ma LG, Castañeda-Ruiz RF, Zhang XG (2011g) New species and record of *Pseudoacrodiclys* from China. *Mycol Prog* 10:261–265
- Zhang YD, Ma J, Ma LG, Castañeda-Ruiz RF, Zhang XG (2011h) New species of *Phaeodactylium* and *Neosporidesmium* from China. *Sydowia* 63:125–130
- Zhang H, Hyde KD, McKenzie EHC, Bahkali AH, Zhou D (2012a) Sequence data reveals phylogenetic affinities of *Acrocylindma aquatica* sp. nov., *Aquasubmersa mircensis* gen. et sp. nov. and *Clovesomyces aquaticus* (Freshwater coelomycetes). *Cryptog Mycol* 33:333–346
- Zhang HX, Tan JL, Wei LX, Wang YL, Zhang CP et al (2012b) Morphology regulatory metabolites from *Arthrobotrys oligospora*. *J Nat Prod* 75:1419–1423
- Zhang HX, Wang LS, Du ZZ, Hua Y (2012c) Comparative analyses on chemical composition of *Sulcaria sulcata* and *Sulcaria sulcata* f. *vulpinoidea*. *J Southwest For Univ* 2:25
- Zhang W, Liu BY, Wu YM, Zhang TY (2012d) A preliminary report of dematiaceous hyphomycetes from soil in Liaoning Province, China. *Mycosistema* 31:1–7
- Zhang Y, Crous PW, Schuh CL, Hyde KD (2012e) *Pleosporales*. *Fungal Divers* 53:1–221
- Zhang YD, Ma J, Ma LG, Castañeda-Ruiz RF, Zhang XG (2012f) A new species of *Corynesporaella* and two new records from southern China. *Cryptog Mycol* 33:99–104
- Zhang YD, Ma J, Ma LG, Castañeda-Ruiz RF, Zhang XG (2012g) A new species of *Quadraceae* and new records of other dematiaceous hyphomycetes from southern China. *Nova Hedw* 94:405–411
- Zhang YD, Ma J, Ma LG, Zhang XG (2012h) *Parablastocatena tetraceriae* gen. et sp. nov. and *Corynesporaella licualiae* sp. nov. from Hainan, China. *Mycoscience* 53:381–385
- Zhang YD, Ma J, Ma LG, Zhang XG (2012i) Two new species of *Taeniolina* from southern China. *Mycol Prog* 11:71–74
- Zhang YM, Maharachchikumbura SSN, McKenzie EH, Hyde KD (2012j) A novel species of *Pestalotiopsis* causing leaf spots of *Trachycarpus fortunei*. *Cryptogam, Mycol* 33(3):311–318
- Zhang Z, Liu X, Wang X, Zhou M et al (2012k) An R2R3 MYB transcription factor in wheat, TaPIMP1, mediates host resistance to *Bipolaris sorokiniana* and drought stresses through regulation of defense-and stress-related genes. *New Phytol* 196:1155–1170
- Zhang H, Hyde KD, Abdel-Wahab MA, Abdel-Aziz FA, Ariyawansa HA et al (2013a) A modern concept for *Helicascus* with a Pleurohomopsis-like asexual state. *Sydowia* 65:147–166
- Zhang Q, Guo WJ, Fu CL, Ma S, Zhu MQ (2013b) Chemical constituents from an endophyte, *Cercosporaella* sp. *Chem Nat Compd* 49:117–118
- Zhang Y, Fournier J, Bahkali AH, Hyde KD (2013c) *Misturatosphaeria mariae* sp. nov. from France, a first record of *Misturatosphaeria* in Europe. *Mycoscience* 54(2):106–109
- Zhang Y, Fournier J, Phookamsak R, Bahkali AH, Hyde KD (2013d) *Halothiaceae* fam. nov (*Pleosporales*) accommodates the new genus *Phaeoseptum* and several other aquatic genera. *Mycologia* 105:603–609
- Zhang Y, Qiao M, Xu J, Cao Y et al (2013e) Genetic diversity and recombination in natural populations of the nematode-trapping fungus *Arthrobotrys oligospora* from China. *Ecol Evol* 3:312–325
- Zhang YM, Maharachchikumbura SSN, Tian Q, Hyde KD (2013f) *Pestalotiopsis* species on ornamental plants in Yunnan Province, China. *Sydowia* 65:113–128
- Zhang H, Hyde KD, Zhao Y, McKenzie EHC, Zhou DQ (2014a) Freshwater ascomycetes: *Lophiostoma vaginatispora* comb. nov. (*Dothideomycetes*, *Pleosporales*, *Lophiostomaceae*) based on morphological and molecular data. *Phytotaxa* 176:184–191
- Zhang K, Ma LG, Ma J, Castañeda-Ruiz RF (2014b) *Xiuguozhangia*, a new genus of microfungi to accommodate five *Piricaudiopsis* species. *Mycotaxon* 128:131–135
- Zhang Y, Yang Y, Zhang XD, Cui BK et al (2014c) *Helicascus gallicus* sp. nov., a new freshwater pleosporalean ascomycete from France. *Phytotaxa* 183:183–192
- Zhang Y, Zhang J, Wang Z, Fournier J et al (2014d) Neotypification and phylogeny of *Kalmusia*. *Phytotaxa* 176:164–173
- Zhang Y, Zhang X, Fournier J, Chen J, Hyde KD (2014e) *Lindgomycetes griseosporus*, a new aquatic ascomycete from Europe including new records. *Mycoscience* 55:43–48
- Zhang J, Zhou Y, Dou Z, Fournier J, Zhang Y (2015a) *Helicascus unilocularis* sp. nov., a new freshwater pleosporalean ascomycete from the Caribbean area. *Mycol Prog* 14:47
- Zhang Y, Liu F, Wu W, Cai L (2015b) A phylogenetic assessment and taxonomic revision of the thermotolerant hyphomycete genera *Acrophialophora* and *Taifanglania*. *Mycologia* 107:768–779
- Zhang H, Dong W, Hyde KD, Bahkali AH et al (2016a) Molecular data shows *Didymella aptrootii* is a new genus in *Bambusicolaceae*. *Phytotaxa* 247:99–108
- Zhang JF, Liu JK, Hyde KD, Liu YX et al (2016b) *Ligninsphaeria jonesii* gen. et sp. nov., a remarkable bamboo inhabiting ascomycete. *Phytotaxa* 247:109–117
- Zhang L, Zhou Z, Guo Q, Fokkens L et al (2016c) Insights into adaptations to a near-obligate nematode endoparasitic lifestyle from the finished genome of *Drechmeria coniospora*. *Sci Rep* 6:23122
- Zhang N, Luo J, Rossman AY, Aoki T et al (2016d) Generic names in Magnaportheales. *IMA Fungus* 7:155–159
- Zhang X, Xu G, Tang H, Li Y, Liu C (2017a) Development of loop-mediated isothermal amplification (lamp) assay for the rapid detection of *Alternaria alternata*. *J AOAC Int* 100:99–103
- Zhang JF, Liu JK, Hyde KD, Yang W, Liu ZY (2017b) Fungi from Asian Karst formations II. Two new species of *Occultibambusa* (*Occultibambusaceae*, *Dothideomycetes*) from karst landforms of China. *Mycosphere* 8:550–559
- Zhao GC, Zhao RL (2012) The higher microfungi from forests of Yunnan Province. Yunnan Science and Technology Press, Kunming, p 564
- Zhao GZ, Liu XZ, Xie XM, Cao AX (2009) Saprobic dematiaceous hyphomycetes from Shennongjia region, China. *Nova Hedw* 88:217–227
- Zhao GZ, Cao AX, Zhang TY, Liu XZ (2011a) *Acrodictys* (Hyphomycetes) and allied genera from China. *Mycol Prog* 10:67–83
- Zhao P, Luo J, Zhuang WY (2011b) Practice towards DNA barcoding of the nectriaceous fungi. *Fungal Divers* 46:183–191
- Zhao GZ, Yu P, Liu XZ (2013a) *Cancellidium* and *Canalisporium* (hyphomycetes) from China. *Nova Hedw* 96:221–236
- Zhao YI, Hosoya T, Shirouzu T, Kakishima M, Yamaoka Y (2013b) *Lambertella pyrolae* (Rutstroemiaceae, Ascomycota), a new species from Japan. *Phytotaxa* 136:54–60
- Zhao KN, Yin GH, Zhao GZ, Cao AX (2014a) *Embryonispora*, a new genus of hyphomycetes from China. *Mycotaxon* 126:77–81
- Zhao X, Wang Y, Zhao Y, Huang Y et al (2014b) Malate synthase gene AoMIs in the nematode-trapping fungus *Arthrobotrys*

- oligospora* contributes to conidiation, trap formation, and pathogenicity. *Appl Microbiol Biotechnol* 98:2555–2563
- Zhao Q, Tolgor B, Zhao YC, Yang ZL, Hyde KD (2015) Species diversity within the *Helvella crispa* group (*Ascomycota: Helvellaceae*) in China. *Phytotaxa* 239:130–142
- Zhao Q, Brooks S, Zhao YC, Yang ZL et al (2016a) Morphology and phylogenetic position of *Wynnella subalpina* sp. nov. (*Helvellaceae*) from western China. *Phytotaxa* 270:41–48
- Zhao Q, Zhao Y, Yang Z, Hyde KD (2016b) Species diversity within the *Helvella crispa* group (*Ascomycota: Helvellaceae*) in China. *Phytotaxa* 239(2):130–142
- Zhao X, Leavitt SD, Zhao ZT, Zhang LL et al (2016c) Towards a revised generic classification of lecanoroid lichens (*Lecanoraceae, Ascomycota*) based on molecular, morphological and chemical evidence. *Fungal Divers* 78:293–304
- Zhao Q, Brooks S, Zhao YC, Hyde KD (2017) Taxonomy and multi-gene phylogeny of the *Helvella atra* complex (*Helvellaceae*) in southwest China. *Mycosphere* 8:577–586
- Zhdanov I (2009) Interesting records of lichens on the coast of Baydaratskaya Bay (Yamal-Nenets Autonomous District, Russia). *Folia Cryptogamica Estonica* 46:79–82
- Zheng CJ, Shao CL, Guo ZY, Chen JF et al (2012) Bioactive hydroanthraquinones and anthraquinone dimers from a soft coral-derived *Alternaria* sp. fungus. *J Nat Prod* 75:189–197
- Zheng HD, Zhuang WY (2013) A new species of *Roseodiscus* (*Ascomycota, Fungi*) from tropical China. *Phytotaxa* 105:51–57
- Zheng HD, Zhuang WY (2014) *Hymenoscyphus albidooides* sp. nov. and *H. pseudoalbidus* from China. *Mycol Prog* 13:625–638
- Zhou XD, Jacobs K, Kirisits T, Chhetri DB, Wingfield MJ (2008) *Leptographium bhutanense* sp. nov., associated with the root collar weevil *Hylobitellus chenkupidorjii* on *Pinus wallichiana* in Bhutan. *Persoonia* 21:1–8
- Zhou F, Wang XY, Zhang L, Lin YR (2013) *Terriera angularis* sp. nov. on *Ilicium simonsii* from China. *Mycotaxon* 122:355–359
- Zhou Y, Gong G, Zhang S, Liu N et al (2014) A new species of the genus *Trematosphaeria* from China. *Mycol Prog* 13:33–43
- Zhou QX, Houbraken J, Li QL, Xu Y et al (2016) Diversity of *Penicillium* species isolated from heavy metal polluted soil in Guizhou Province, China. *Phytotaxa* 273:167–174
- Zhu XF, Pan Y, Chen LJ, Duan YY, Wang YY (2012) First report of leaf spot of soybean caused by *Aristostoma guttulosum* in China. *Plant Dis* 96:1694
- Zhu L, Chen GQ, Zhao XL, Deng CL et al (2013) *Fusarium* spp. are responsible for Shoot Canker of kumquat in China. *J Phytopathol* 161:59–62
- Zhu XQ, Niu CW, Chen XY, Guo LY (2016) *Monilinia* species associated with brown rot of cultivated apple and pear fruit in China. *Plant Dis* 100:2240–2250
- Zhuang WY (2009) The genus *Sowerbyella* (*Pezizales*) in China. *Mycotaxon* 109:233–237
- Zhuang WY (2010) Taxonomic assessment of some pyronemataceous fungi from China. *Mycotaxon* 112:31–46
- Zhuang WY (2013) The genus *Scutellinia* (*Pyronemataceae*) from China with a key to the known species of the country. *Mycosistema* 32:432
- Zhuang WY, Yang ZL (2008) [2007] Some pezizalean fungi from alpine areas of southwestern China. *Mycologia Montenegrina* 10:235–249
- Zhuang WY, Luo J, Zhao P (2011) Two new species of *Acervus* (*Pezizales*) with a key to species of the genus. *Mycologia* 103:400–406
- Zhurbenko MP (2008) *Lichenochora thorii* (*Phyllachorales*), a new lichenicolous ascomycete from Europe. *Nova Hedw* 87:509–512
- Zhurbenko MP (2009) Lichenicolous fungi and some lichens from the Holarctic. *Opuscula Phil* 6:87–120
- Zhurbenko MP (2010a) Lichenicolous fungi and lichens growing on *Stereocaulon* from the Holarctic, with a key to the known species. *Opuscula Phil* 8:9–39
- Zhurbenko MP (2010b) New and interesting lichenicolous fungi from Eurasia, II. *Mycosphere* 1:213–222
- Zhurbenko MP (2011) Lichenicolous fungi growing on *Thamnolia*, mainly from the Holarctic, with a worldwide key to the known species. *Lichenologist* 44:147–177
- Zhurbenko MP (2013) Lichenicolous fungi and some allied lichens from the Canadian Arctic. *Opuscula Phil* 12:180–197
- Zhurbenko M, Dillman K (2010) *Polycoccum hymenicolola* comb. nov. (*Dacampiaceae*) and other interesting lichenicolous fungi from southeastern Alaska. *Bryologist* 113:260–266
- Zhurbenko MP, Braun (2012) *Cecidiomyces*, a new subantarctic lichenicolous hypomycete genus. *Lichenologist* 44:801–806
- Zhurbenko MP, Otte V (2012) Lichenicolous fungi from the Caucasus: new records and a first synopsis. *Herzogia* 25:235–244
- Zhurbenko MP, Braun U (2013) *Ameroconium cladoniae* gen. et sp. nov. and *Phoma psoromatis* sp. nov., new anamorphic lichenicolous fungi from the Holarctic. *Lichenologist* 45:583–591
- Zhurbenko MP, Tugi EW (2013) Some noteworthy lichenicolous fungi from Khabarovsk Territory of Russia. *Mycosphere* 4:46–51
- Zhurbenko MP, Zhdanov IS (2013) *Melaspilea galligena* sp. nov. and some other lichenicolous fungi from Russia. *Folia Cryptog Estonica* Fasc 50:89–99
- Zhurbenko MP, Pino-Bodas R (2015) New lichenicolous fungi growing on *Cladonia* in New Zealand. *Lichenologist* 4:395–402
- Zhurbenko MP, Kukwa M, Oset M (2009) *Roselliniella stereocaulorum* (*Sordariales, Ascomycota*), a new lichenicolous fungus from the Holarctic. *Mycotaxon* 109:323–328
- Zhurbenko MP, Himelbrant DE, Kuznetsova ES, Stepanchikova IS (2012) Lichenicolous fungi from the Kamchatka Peninsula, Russia. *Bryologist* 115:295–312
- Zhurbenko MP, Braun U, Heuchert B, Kobzeva AA (2015a) Three new lichenicolous hyphomycetes from Asia. *Herzogia* 28:584–598
- Zhurbenko MP, Etayo J, Fedrowitz K, Thor G (2015b) Miscellaneous reports of lichenicolous fungi from Argentina including the new species *Didymelopsis nephromatis*. *Opuscula Phil* 14:82–89
- Zhurbenko MP, Heuchert B, Braun U (2015c) *Talpapellis solorinae* sp. nov. and an updated key to the species of *Talpapellis* and *Verrucocladosporium*. *Phytotaxa* 234:191–194
- Zitouni-Haouar FE, Alvarado P, Sbissi I, Boudabous A et al (2015) Contrasted genetic diversity, relevance of climate and host plants, and comments on the taxonomic problems of the genus *Picoa* (*Pyronemataceae, Pezizales*). *PLoS ONE* 10:e0138513
- Zou S, Li Q, Kong L (2011) Additional gene data and increased sampling give new insights into the phylogenetic relationships of *Neogastropoda*, within the caenogastropod phylogenetic framework. *Mol Phylogenetic Evol* 61:425–435
- Zupančič J, Novak Babič M, Zalar P, Gunde-Cimerman N (2016) The black yeast *Exophiala dermatitidis* and other selected opportunistic human pathogens spread from dishwashers to kitchens. *PLoS ONE* 11:e0148166