

Associations of Participation in Organized Sports and Physical Activity in Preschool Children: A Cross-Sectional Study

Chu Chen (✉ chen.chu@ki.se)

Karolinska Institutet <https://orcid.org/0000-0001-6350-3797>

Fanny Sellberg

Karolinska Institutet

Viktor H. Ahlqvist

Karolinska Institutet

Martin Neovius

Karolinska Institutet Institute of Environmental Medicine

Filip Christiansen

Karolinska Institutet

Daniel Berglind

Karolinska Institutet

Research article

Keywords: Exercise, sedentary, sports, preschool, accelerometer

Posted Date: April 29th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-24897/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on July 2nd, 2020. See the published version at <https://doi.org/10.1186/s12887-020-02222-6>.

Abstract

Background Participation in organized sports is associated with higher physical activity (PA) levels in school-aged-children. Yet, little is known about PA determinants in preschool-aged-children. We examined associations between organized sports participation and preschoolers' daily PA.

Methods The study comprised 290 3–5 years old children and PA was measured for 1 week via accelerometers. Organized sports participation was parent-reported and preschool time was teacher-recorded. Linear mixed models, nested on preschool level, were used to examine associations between organized sports participation and children's PA outside preschool time, during preschool time and throughout the day.

Results In total, 146 children (50.3%) participated in organized sports at least 1 hour/week. Participation in organized sports was associated with 6.0 more minutes of moderate-to-vigorous PA (MVPA) (95% CI: 0.6, 11.3) throughout the day and 5.7 more minutes of MVPA (95% CI: 1.6, 9.7) outside preschool time after adjustment. There was no association between organized sports participation and PA during preschool time.

Conclusions This is the first study to show positive associations between organized sports participation and preschoolers' PA levels outside preschool time and throughout the day. In addition, findings from this study do not support PA compensation. Therefore, targeting organized sports may be successful in improving PA, even among preschoolers.

Introduction

There are well-established associations between total physical activity (PA), moderate to vigorous PA (MVPA) (1), steps (2) and both short- and long-term health benefits in children, while more conflicting findings have been reported for objectively measured sedentary time (ST) (3). Beyond the immediate early life (0–4 years) health benefits associated with regular PA, physically active children also have a tendency to continue being physically active across their lifespan (4). Therefore, a major concern is the low levels of PA among young children (5). According to a review on preschoolers' objectively measured physical activity level, preschoolers spend 2–41% of their day in MVPA, 4–33% in light PA (LPA), and 34–94% sedentary (6). Moreover, Swedish data with objectively measured PA, show that preschool aged children's PA levels are low, especially outside preschool hours and on weekend days (7).

Organized sports, which is defined as *“PA that is directed by adult or youth leaders and involves rules and formal practice and competition”* (8), have been associated with reduced risk of morbidity. In adolescents, participation in organized sports is associated with higher levels of PA (9), decreased risk for cardiovascular disease (10), improved mental health and social adjustment (11). Thus, participation in organized sports is a potential strategy to improve adolescents' physical and mental health by increasing their PA (9–11). Moreover, childhood exposure to organized sports may be influential on participation in

later years (12). However, little is known about the association between organized sports participation and PA levels in preschool aged children (13).

There are studies showing a PA synergy, where participation in exercise/PA increases PA at other times of the day (14, 15). However, there is a conflicting “activity-stat” hypothesis, which proposes that children compensate for increased PA in during part of the day (e.g. during organized sports participation) by decreasing their PA at another time (e.g. during preschool time); thus, maintaining a fixed level of total PA (16). This hypothesis is supported by review data indicating that approximately 63% of exercise/PA intervention studies, that specifically examined this hypothesis in children, have reported PA compensation (17). If preschool children maintain a fixed level of total PA despite engaging in organized sports, any interventions targeting organized sports may be ineffective in achieving health benefits associated with PA. The aforementioned review study also calls for more evidence to examine the validity of the hypothesis (17). In preschool children, the compensatory behavior was not observed in studies comparing days with and without teacher-led structured physical activity (18), comparing between locations (childcare center and home) (19), and when increasing outdoor time (20). Yet, to the best of our knowledge, no previous study has investigated the association between participation in organized sports and preschool children’s activity levels in different segments of the day.

Based on these knowledge gaps, the aims of the current study were to examine associations between participation in organized sports and objectively measured PA and ST: (i) outside preschool time, (ii) during preschool time and (iii) throughout the day in a sample of Swedish 3–5 years old children.

Methods

Study setting, design, and population

Within the Södermalm district of Stockholm Sweden, a convenience sample of 30 out of the total 51 municipal preschools were invited to participate in the current cross-sectional observational study. Out of 30 preschools invited, 27 preschools chose to participate, including a total of 1178 children. At the participating preschools, all children between 3–5 years of age were invited to participate. Out of the 1178 total children, 405 (23%) children and their parents consented to participation in the study. In Sweden, children aged below 6 attend preschools every weekday (i.e. 5/7 in a week). The study was approved by the Stockholm Ethical Review Board (Dnr: 2018/890 – 31/2), and informed consent was obtained from participating children’s parents and preschool teachers. The fieldwork measurements, including body measures of children, 7 days of accelerometer measures of PA in children and parental questionnaires, were carried out during September to November 2018.

Participation in organized sports (exposure)

Parents to participating children filled in a questionnaire with the question “Does your child participate in any kind of organized sports?”. The answer options were (i) no participation, (ii) 1–2 hours organized sports/week, (iii) 3 hours organized sports/week or (iv) 4 or more hours organized sports/week.

Organized sports are generally accepted as structured leisure time activities in non-profit organization (21). Therefore, the participation in organized sports assessed here was explicitly about sports participation outside preschool hours. Due to the low participation in organized sports of more than 2 hours/week (10%), we classified participation in organized sports into a dichotomous variable: (i) No organized sports and (ii) at least one-hour participation in organized sports/week.

Physical activity and sedentary time outside preschool time, during preschool time and throughout the day (outcomes)

The outcome measures, PA and ST outside preschool time, during preschool time and throughout the day, were measured via the triaxial Actigraph GT3X + accelerometer, which has been widely used to assess PA and ST in pediatric research (22). We consulted best practices for wear protocol and analysis (22) and as such decided our procedure to be the following: All children were instructed to wear accelerometer, at right hip for 7 consecutive days, during all waking hours children that had worn the accelerometer for at least 3 days with 10 or more wear-time hours per day were included in the analytical dataset (22). Non-wear time was defined as 60 or more consecutive minutes with zero counts, allowing up to 2 minutes of interruptions with non-zero counts (23) after adaptation for a potentially less compliant sample (preschool children) (24). We analyzed vector magnitude (V_m) activity counts ($V_m = \sqrt{X^2 + Y^2 + Z^2}$) in 60 s epochs and MVPA, light physical activity (LPA) and ST were calculated based on cut-offs developed specifically for V_m activity counts in 4-year old children (25): ST was calculated as any minute of less than 820 counts per minute (cpm), LPA as 820–3907 cpm and MVPA as ≥ 3908 cpm. Steps were determined using the manufacturer's step algorithm, using the normal filter setting.

During the 7 days of accelerometer measures in children, preschool staff recorded time arrival to and leaving the preschool on daily basis for each participating child. This data was thereafter matched, on daily level, to time-stamped accelerometer data which enabled us to calculate PA and ST before, after and during preschool. PA and ST before and after preschool time were then combined with PA and ST during the weekend to calculate PA and ST outside preschool time. PA and ST throughout the day was considered as all wear-time hours during the day. The mean daily PA and ST outside preschool time, during preschool time and throughout the day were calculated on the individual level and then matched with data on organized sports participation.

Covariates

Anthropometry and family characteristics were also documented. Weight and height of participating children were measured using validated scales and stadiometers, respectively (scale: VB2-200-EC, Vetek AB, Väddö, Sweden, stadiometer: Seca 213, Seca, Chino, CA, USA). We used an age and sex specific international body mass index (BMI) classification by Cole et al (26) to classify children as normal weight, overweight or obese. Parents filled out a questionnaire on demographical variables such as number of siblings within the family and highest parental education level, categorized into elementary school, upper secondary school and university education.

Statistical analysis

Appropriate measures of variability and central tendency, mean and standard deviation (SD) for normally distributed variables or median and interquartile range (IQR) for variables with skewed distribution, are presented for various background characteristics and PA outcomes of the total study population and stratified by participation in organized sports.

We used linear mixed models, nested on preschool level, to examine associations between participation in organized sports with children's daily levels of MVPA, LPA, steps and ST outside preschool time, during preschool time and throughout the day. Separate models were fitted for each activity intensity. In addition, we performed robust multilevel mixed-effects Poisson regression, nested on preschool level, to examine the relative risk of meeting the current World Health Organization (WHO) PA guidelines for children under five (27), of at least 60 minutes of daily MVPA, in organized sports participation with no participation in organized sports as reference level. All models were adjusted for confounders, factors that potentially influences both exposure (participation in organized sports) and outcomes, selected based on causal reasoning (28). The selected confounders were age of the children, sex, overweight/obesity status (29), accelerometer wear-time (30), number of siblings (31) and parental education (32). We calculated all p-values for both Poisson and linear mixed models using Wald tests, testing the coefficient of interest being equal to zero. To test for sex-specific associations, we performed a Wald test of the coefficient of a product interaction between sex and organized sports participation and stratified our main analysis.

In sensitivity analyses, we compared the descriptive characteristics between the analytical dataset (n = 290) and excluded observations (n = 104).

Raw accelerometer data was processed in Actilife version 6.13.3 and all statistical analysis were conducted in Stata version 14.2.

Results

In total, 404 children aged 3–5 years participated (Fig. 1). Among participants, 114 children were excluded with 10 children disqualified due to invalid accelerometer data and 104 children excluded due to missing value. Of the children with missing values, we first excluded 30 children who had missing data on organized sports participation, secondly the 33 children who had missing data on preschool arrival and departure time, thirdly the 33 children who did not have weekend accelerometer data to determine PA outside preschool time, and finally 8 children who had missing co-variate data (weight, height, number of siblings and parental education). Consequently, the final analytical sample comprised 290 children with valid accelerometer data, data on organized sports participation, time outside/at preschool and all co-variate variables.

Table 1 displays an overview of descriptive characteristics and children's levels of PA (outcomes) by participation in organized sports (exposure) in categories of no participation in organized sports versus participation in at least 1 hour of organized sports/week. In total, 146 children (50.3%) participated in

organized sports at least 1 hour/week. Of those children who participated in organized sports, 131 (90%) participated 1–2 hours/week, 12 (8%) 3 hours/week and 3 (2%) 4 or more hours/week.

Figure 2 illustrates associations between participation in organized sports and MVPA, LPA, steps and ST outside preschool time, during preschool time throughout the whole day in adjusted analyses. Children who participated in organized sports took 693 more steps (95% CI: 331, 1056; $p < 0.001$) outside preschool time and 658 more steps (95% CI: 237, 1079; $p < 0.001$) throughout the whole day compared with children who did not participate in organized sports (Additional file 1, Additional Table 1). Similarly, children participating in organized sports spent 5.7 more minutes in MVPA (95% CI: 1.6, 9.7; $p = 0.03$) outside preschool time and 6.0 more minutes in MVPA (95% CI: 0.6, 11.3; $p = 0.01$) throughout the whole day. There were no associations between participation in organized sports and steps ($p = 0.20$) or MVPA ($p = 0.18$) during preschool time. Although the interaction between sex and organized sports participation was not statistically significant ($P > 0.05$), associations between participation in organized sports and all PA indicators outside preschool time, during preschool time and throughout the whole day were stronger in boys compared to girls (Additional file 1, additional Table 1).

Table 1

Descriptive characteristics in relation to participation in organized sports Abbreviation: SD = standard deviation

Participation in organized sports			
	No (n = 144)	Yes (n = 146) At least 1 h per week	Total (n = 290)
Descriptive characteristics			
Girls, n (%)	67 (46.5%)	64 (43.8%)	131 (45.2%)
Age, mean (SD)	4.4 (0.8)	5.0 (0.6)	4.7 (0.8)
Overweight/obese, n (%)	11 (7.6%)	16 (11.0%)	27 (9.3%)
Number of siblings,	1.0 (0.8)	1.1 (0.7)	1.1 (0.8)
0, n (%)	33 (22.9%)	22 (15.7%)	55 (19.0%)
1, n (%)	91 (63.2%)	88 (60.3%)	179 (61.7%)
2, n (%)	16 (11.1%)	33 (22.7%)	49 (16.9%)
≥3, n (%)	4 (2.8%)	3 (2.3%)	7 (2.4%)
Parental education,			
Elementary school, n (%)	2 (1.4%)	3 (2.0%)	5 (1.7)
Secondary school, n (%)	29 (20.1%)	21 (14.4%)	50 (17.3%)
University, n (%)	113 (78.5%)	122 (83.6%)	235 (81.0%)
Physical activity outside preschool time, mean (SD)			
Moderate-to-vigorous physical activity (min)	31.3 (14.6)	40.8 (18.8)	36.1 (17.5)
Light physical activity (min)	220.5 (37.2)	225.5 (41.5)	223.1 (39.5)
Steps (counts)	5510 (1427)	6627 (1663)	6072 (1646)
Sedentary time (min)	270.3 (61.1)	266.0 (72.3)	268.2 (66.9)
Wear-time (min)	522.2 (68.2)	532.4 (71.1)	527.3 (69.7)
Physical activity during preschool time, mean (SD)			

Participation in organized sports			
Moderate-to-vigorous physical activity (min)	32.3 (20.1)	40.6 (20.5)	36.5 (20.7)
Light physical activity (min)	247.0 (49.5)	240.9 (40.1)	243.9 (45.1)
Steps (counts)	6684 (2185)	7161 (2108)	6924 (2156)
Sedentary time (min)	172.1 (43.0)	161.5 (40.2)	166.7 (41.9)
Wear-time (min)	451.5 (66.2)	443.1 (50.7)	447.3 (59.0)
Physical activity whole day, mean (SD)			
Moderate-to-vigorous physical activity (min)	47.1 (21.3)	60.6 (25.3)	53.9 (24.3)
≥ 60 min/day, n (%)	38 (26.4%)	64 (43.8%)	102 (35.2%)
Light physical activity (min)	350.9 (43.4)	355.3 (42.1)	353.1 (42.7)
Steps (counts)	9053 (2061)	10397 (2134)	9730 (2200)
Sedentary time (min)	355.2 (64.7)	342.6 (66.3)	348.8 (65.7)
Wear-time (min)	753.2 (55.5)	758.5 (56.8)	755.9 (56.1)

Additional Table 2 in Additional file 1 illustrates associations between participation in organized sports and meeting the PA recommendation, of 60 or more minutes of daily MVPA, in crude and adjusted analyses. In adjusted analysis, participation in organized sports was associated with a 20% higher risk of meeting the daily PA recommendation (RR 1.2, 95% CI: 0.8, 1.9). However, this association was not statistically significant ($p > 0.05$).

Additional Table S3 in Additional file 1 provides an overview of descriptive characteristics and child PA indicators throughout the whole day between the analytical dataset ($n = 290$) and excluded observations ($n = 104$). On average, participants in the analytical dataset and those excluded from analyses were similar with regards to descriptive characteristics and PA throughout the day.

Discussion

The current study was the first to examine associations between participation in organized sports and objectively measured PA and ST outside preschool time, during preschool time and throughout the day in

preschool-aged children. Our findings showed that approximately 50% of children participated in organized sports for 1 or more hours/week. Participation in organized sports was positively associated with children's MVPA and steps outside preschool time and throughout the whole day. There was no association between participation in organized sports and PA during preschool time. Thus, findings from the current study do not support a PA compensation in preschool children participating in organized sports.

Studies in school aged children and adolescents have shown a positive association between participation in organized sports and children's accelerometer measured PA throughout the day (33–35). Previous studies have also shown a significant interaction between organized sports participation and sex on children's PA (34–36). Results from the current study showing that organized sports participation was associated with 6.0 minutes (10%) more MVPA throughout the day, and potentially a greater association in boys compared with girls, are in line with those presented in above-mentioned studies. The activity-stat hypothesis is supported by several exercise intervention studies in older children (school aged), showing a compensation in PA (17). In contrast, results from the current study show that preschoolers who participate in organized sports are equally active as those who do not participate in organized sports during preschool time and more active both outside preschool time and throughout the whole day.

Given the low and decreasing levels of PA among young children (5, 37), it is important to find effective strategies to increase young children's PA. Meta-analyses data of trials examining the effectiveness of interventions to increase PA in children show minor treatment effects; adding up to approximately 4 minutes of additional walking or running per day (38). Although the results from this cross-sectional study cannot be compared to results from that of interventions, the 6.0 minutes (10%) more MVPA associated with participation in organized sports is a practically significant difference which may warrant further intervention studies of organized sports in preschool children. This finding, in combination with the positive effect participation in organized sports may have on the tracking of PA from childhood to adulthood (4), supports early life participation in organized sports as a strategy to achieve higher levels of PA throughout childhood, adolescence and adulthood. The higher PA level, potentially resulted from participation in organized sports, can further benefit a wide range of health indicator in preschool children comprising bone strength, motor development, fitness and psychosocial health (1). However, the observational and cross-sectional nature of the current study can only provide evidence of a positive association between organized sports participation and PA throughout the day in preschoolers. A randomized controlled trial in preschool aged children is warranted to examine if the associations observed in the current study hold.

A strength of this study was the accelerometer measured PA which limit several biases associated with self-reported measures, e.g. social desirability and recall difficulties. Second, the time-stamped accelerometer PA data and detailed daily data on time at and outside preschool enabled us to assess the association between organized sport participation and PA levels during preschool time, outside preschool time and throughout the day. Third, in the aspect of obtaining organized sports participation information

and objectively measured PA data, the current study included a fairly large number of children in a previous unstudied age group.

Limitations of our study include that information on organized sports participation relied on parental self-reporting, which has several inherent biases and we did not take type of sport performed into account. Second, the observational and cross-sectional nature of the study preclude any inference about causality, and we cannot rule out that our results may be explained by unmeasured confounders or reverse causality, i.e. children who are more active chose to participate in organized sports and not vice versa. Third, although accelerometry is considered as a desirable measurement of PA among preschool aged children in free-living conditions, hip-worn accelerometers are unable to detect some types of PA, e.g. cycling, swimming or PA involving upper body movements (39). Forth, by using the normal filter to process accelerometer data we may have underestimated children's number of steps taken per day. Fifth, the sample of preschool children was selected based on acceptance of the invitation from the invited children and parents which could potentially influence the generalizability of the study. Finally, the study population was homogenous in terms of socio-economic aspects and urbanity. This is of importance as participation in organized sports may be associated with household finances due to high costs for many of these activities. Consequently, our results have limited generalizability to rural and lower socioeconomic areas.

Conclusions

To the best of our knowledge, the current study is the first to show, in preschool children, a positive association between participation in organized sports and objectively measured moderate-to-vigorous physical activity and steps outside preschool time and throughout the whole day. There was no association between participation in organized sports and physical activity during preschool time. Thus, results from the current study do not support PA compensation among preschool aged children. Targeting organized sports may be successful in improving PA, even among very young children (3–5 years old). This could contribute to forming interest and habit of sport participation from early age and throughout childhood (4) with a wide range of benefits such as improving fitness, motor development and psychosocial health (1).

List Of Abbreviations

IQR Interquartile range

LPA Light physical activity

MVPA Moderate to vigorous physical activity

PA Physical activity

SD Standard deviation

ST Sedentary time

WHO World health organization

Declarations

Ethics approval and consent to participate

The study was approved by the Stockholm Ethical Review Board (Dnr: 2018/890-31/2) and written informed consent was obtained from participating children's parents and preschool teachers.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was partly funded by Lindhés Advokatbyrå and by Södermalm municipality. The funding parties had no partaking in study design, data analyses/interpretation or writing of the current article.

Authors' contributions

CC: data analyses, data collection/management, manuscript editing, FS: data analyses, manuscript editing, VA: data analyses and manuscript editing, MN: data interpretation, manuscript editing, FC: data collection, manuscript editing, DB: study design, data analyses/interpretation, manuscript writing. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank all fund providers, participating preschools, parents and children who made this study possible.

References

1. Carson V, Lee EY, Hewitt L, Jennings C, Hunter S, Kuzik N, et al. Systematic review of the relationships between physical activity and health indicators in the early years (0–4 years). BMC Public Health.

- 2017;17(Suppl 5):854.
2. Tudor-Locke C, Craig CL, Beets MW, Belton S, Cardon GM, Duncan S, et al. How Many Steps/Day are Enough? for Children and Adolescents. *Int J Behav Nutr Phy.* 2011;8.
 3. Cliff DP, Hesketh KD, Vella SA, Hinkley T, Tsiros MD, Ridgers ND, et al. Objectively measured sedentary behaviour and health and development in children and adolescents: systematic review and meta-analysis. *Obes Rev.* 2016;17(4):330–44.
 4. Jones RA, Hinkley T, Okely AD, Salmon J. Tracking Physical Activity and Sedentary Behavior in Childhood: A Systematic Review. *Am J Prev Med.* 2013;44(6):651–8.
 5. Bornstein DB, Beets MW, Byun W, McIver K. Accelerometer-derived physical activity levels of preschoolers: a meta-analysis. *Journal of Science Medicine in Sport.* 2011;14(6):504–11.
 6. Hnatiuk JA, Salmon J, Hinkley T, Okely AD, Trost S. A review of preschool children's physical activity and sedentary time using objective measures. *Am J Prev Med.* 2014;47(4):487–97.
 7. Berglind D, Tynelius P. Objectively measured physical activity patterns, sedentary time and parent-reported screen-time across the day in four-year-old Swedish children. *BMC Public Health.* 2017;18(1):69.
 8. Logan K, Cuff S, Council On Sports M. Fitness. Organized Sports for Children, Preadolescents, and Adolescents. *Pediatrics.* 2019.
 9. Mandic S, Bengoechea EG, Stevens E, de la Barra SL, Skidmore P. Getting kids active by participating in sport and doing it more often: focusing on what matters. *Int J Behav Nutr Phys Act.* 2012;9:86.
 10. Hebert JJ, Klakk H, Moller NC, Grontved A, Andersen LB, Wedderkopp N. The Prospective Association of Organized Sports Participation With Cardiovascular Disease Risk in Children (the CHAMPS Study-DK). *Mayo Clin Proc.* 2017;92(1):57–65.
 11. Eime RM, Young JA, Harvey JT, Charity MJ, Payne WR. A systematic review of the psychological and social benefits of participation in sport for children and adolescents: informing development of a conceptual model of health through sport. *Int J Behav Nutr Phys Act.* 2013;10:98.
 12. Allender S, Cowburn G, Foster C. Understanding participation in sport and physical activity among children and adults: a review of qualitative studies. *Health Educ Res.* 2006;21(6):826–35.
 13. De Craemer M, De Decker E, De Bourdeaudhuij I, Vereecken C, Deforche B, Manios Y, et al. Correlates of energy balance-related behaviours in preschool children: a systematic review. *Obes Rev.* 2012;13(s1):13–28.
 14. Goodman A, Mackett RL, Paskins J. Activity compensation and activity synergy in British 8–13 year olds. *Prev Med.* 2011;53(4–5):293–8.
 15. Long MW, Sobol AM, Cradock AL, Subramanian SV, Blendon RJ, Gortmaker SL. School-Day and Overall Physical Activity Among Youth. *Am J Prev Med.* 2013;45(2):150–7.
 16. Rowland TW. The biological basis of physical activity. *Med Sci Sport Exer.* 1998;30(3):392–9.
 17. Gomersall SR, Rowlands AV, English C, Maher C, Olds TS. The ActivityStat hypothesis: the concept, the evidence and the methodologies. *Sports medicine.* 2013;43(2):135–49.

18. Van Cauwenberghe E, De Craemer M, De Decker E, De Bourdeaudhuij I, Cardon G. The impact of a teacher-led structured physical activity session on preschoolers' sedentary and physical activity levels. *Journal of Science Medicine in Sport*. 2013;16(5):422–6.
19. Hesketh KR, Griffin SJ, van Sluijs EMF. UK Preschool-aged children's physical activity levels in childcare and at home: a cross-sectional exploration. *Int J Behav Nutr Phy*. 2015;12(1):123.
20. Alhassan S, Sirard JR, Robinson TN. The effects of increasing outdoor play time on physical activity in Latino preschool children. *International Journal of Pediatric Obesity*. 2007;2(3):153–8.
21. Støckel JT, Strandbu Å, Solenes O, Jørgensen P, Fransson K. Sport for children and youth in the Scandinavian countries. *Sport in Society*. 2010;13(4):625–42.
22. Migueles JH, Cadenas-Sanchez C, Ekelund U, Delisle Nystrom C, Mora-Gonzalez J, Lof M, et al. Accelerometer Data Collection and Processing Criteria to Assess Physical Activity and Other Outcomes: A Systematic Review and Practical Considerations. *Sports medicine*. 2017.
23. Choi L, Liu Z, Matthews CE, Buchowski MS. Validation of accelerometer wear and nonwear time classification algorithm. *Med Sci Sport Exer*. 2011;43(2):357.
24. Knaier R, Höchsmann C, Infanger D, Hinrichs T, Schmidt-Trucksäss A. Validation of automatic wear-time detection algorithms in a free-living setting of wrist-worn and hip-worn ActiGraph GT3X. *BMC Public Health*. 2019;19(1):244-.
25. Butte NF, Wong WW, Lee JS, Adolph AL, Puyau MR, Zakeri IF. Prediction of energy expenditure and physical activity in preschoolers. *Med Sci Sports Exerc*. 2014;46(6):1216–26.
26. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320(7244):1240–3.
27. WHO. WHO guidelines on physical activity, sedentary behaviour and sleep for children under 5 years of age. In: WHO, editor. <https://apps.who.int/iris/bitstream/handle/10665/311664/9789241550536-eng.pdf?sequence=1&isAllowed=y>: WHO; 2019.
28. Williamson EJ, Aitken Z, Lawrie J, Dharmage SC, Burgess JA, Forbes AB. Introduction to causal diagrams for confounder selection. *Respirology*. 2014;19(3):303–11.
29. Dolinsky DH, Brouwer RJN, Evenson KR, Siega-Riz AM, Østbye T. Correlates of sedentary time and physical activity among preschool-aged children. *Prev Chronic Dis*. 2011;8(6):A131-A.
30. Oliver M, Schofield GM, Kolt GS. Physical activity in preschoolers: understanding prevalence and measurement issues. *Sports medicine*. 2007;37(12):1045–70.
31. Kracht CL, Sisson SB. Sibling influence on children's objectively measured physical activity: a meta-analysis and systematic review. *BMJ open sport exercise medicine*. 2018;4(1):e000405-e.
32. Wijtzes AI, Jansen W, Bouthoorn SH, Pot N, Hofman A, Jaddoe VWV, et al. Social inequalities in young children's sports participation and outdoor play. *Int J Behav Nutr Phys Act*. 2014;11:155-.
33. Hebert JJ, Moller NC, Andersen LB, Wedderkopp N. Organized Sport Participation Is Associated with Higher Levels of Overall Health-Related Physical Activity in Children (CHAMPS Study-DK). *PloS one*. 2015;10(8).

34. Marques A, Ekelund U, Sardinha LB. Associations between organized sports participation and objectively measured physical activity, sedentary time and weight status in youth. *Journal of Science Medicine in Sport*. 2016;19(2):154–7.
35. Lagestad P, Mikalsen H, Ingulfsvann LS, Lyngstad I, Sandvik C. Associations of Participation in Organized Sport and Self-Organized Physical Activity in Relation to Physical Activity Level Among Adolescents. *Frontiers in public health*. 2019;7.
36. Telford RM, Telford RD, Cochrane T, Cunningham RB, Olive LS, Davey R. The influence of sport club participation on physical activity, fitness and body fat during childhood and adolescence: The LOOK Longitudinal Study. *J Sci Med Sport*. 2016;19(5):400–6.
37. Nyström CD, Larsson C, Ehrenblad B, Eneroth H, Eriksson U, Friberg M, et al. Results from Sweden's 2016 report card on physical activity for children and youth. *Journal of physical activity health*. 2016;13(s2):284-S90.
38. Metcalf B, Henley W, Wilkin T. Effectiveness of intervention on physical activity of children: systematic review and meta-analysis of controlled trials with objectively measured outcomes (EarlyBird 54). *BMJ*. 2012;345:e5888.
39. Cliff DP, Reilly JJ, Okely AD. Methodological considerations in using accelerometers to assess habitual physical activity in children aged 0–5 years. *J Sci Med Sport*. 2009;12(5):557–67.

Figures




Figure 1

Flow chart of participants Description of participants included in the analytical data set (n=290) and excluded participants (n=114)




Figure 2

Associations between participation in organized sports and moderate-to-vigorous physical activity, steps, light physical activity and sedentary time outside preschool time, during preschool time and throughout the whole day Adjusted for sex, age, overweight/obesity status, accelerometer wear-time, parental education and number of siblings Square: not participating in organized sports Triangle: participating in organized sports

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)