

The mosses (Bryophyta) of Capitán Prat Province, Aisén Region, southern Chile

Juan Larraín¹

¹ Instituto de Biología, Facultad de Ciencias, Pontificia Universidad Católica de Valparaíso, Campus Curauma, Av. Universidad 330, Curauma, Valparaíso, Chile

Corresponding author: Juan Larraín (musgoschiloe@gmail.com)

Academic editor: Matt von Konrat | Received 12 May 2016 | Accepted 25 July 2016 | Published 16 August 2016

Citation: Larraín J (2016) The mosses (Bryophyta) of Capitán Prat Province, Aisén Region, southern Chile. PhytoKeys 68: 91–116. doi: 10.3897/phytokeys.68.9181

Abstract

The bryophytes of Capitán Prat province have remained one of the least explored in Chile. The eventual construction of several dams on the rivers Baker and Pascua required prospection of all groups of organisms including bryophytes, work that was facilitated by the recent construction of vehicular roads that now offer easy access to previously almost unaccessible locations. The results of intense bryophyte collecting during the austral summer of 2007 are here presented. A total of 260 moss taxa are reported for the province, corresponding to 256 species and four infraspecific taxa, of which 211 are new records for the province, 54 are new for Aisén Region, and two are new records for continental Chile (*Pohlia longicollis* (Hedw.) Lindb. and *Rigodium toxarium* var. *robustum* (Broth.) Zomlefer). Twelve species extend their known distribution ranges to the north, whereas 49 extend them to the south.

Keywords

Biodiversity, checklist, range extensions, new records, southern South America

Introduction

The Aisén Region is divided into four administrative provinces, including Capitán Prat as the southernmost province of the Region, bordering by the south with Última Esperanza Province in the Magallanes Region. It is of particular botanical interest due to: (i) the diversity of climates, ranging from the per-humid temperate rainforests along the coast line and the western archipelago, to the extremely dry steppe in the eastern border with Argentina, and (ii) it is the southernmost ice-free area of continental Chile before the beginning of the massive Southern Patagonian Ice Field that partially isolates it from the neighboring Magallanes Region.

Although the Aisén Region has received considerable exploration by bryologists, and there is even a moss flora already published (Seki 1974, published under the incorrect title of “Provincia de Aisén”), the southernmost province of the region has received very few visits by moss collectors (Fig. 1A). Surprisingly, to date, only two expeditions have collected mosses in the province, i.e., those by Carl Skottsberg at Cañal Baker (Caleta Hale, Caleta Connor and Puerto Cueri-Cueri) in the austral winter of 1908, and at Lago O’Higgins in the summer of 1909 (Skottsberg 1916), and those by Arturo Donat at Lago O’Higgins on 1933, as part of the Argentinian expedition “Gaea” (Donat 1936a, 1936b). The first bryophytes reported for the province were 15 liverwort species (Stephani 1911), based on Skottsberg’s collections of 1908. The main scope of Skottsberg’s work was investigating vascular plants, yet Skottsberg (1916) reported 19 moss species, 38 liverworts, and some lichens. Skottsberg’s collections where further studied by Cardot and Brotherus (1923), who reported five new records for the localities mentioned above. Later, Donat reported 12 additional moss species (Donat 1936a), including six new to science (although one of them in the Argentinian side of Lago O’Higgins, locally named there Lago San Martín), and in a second work describing the flora of the western shore of O’Higgins lake (Donat 1936b), he added five more species, making a total of 40 moss species known for the province by 1936. In his moss flora of Aisén Region, Seki (1974) did not report a single locality that belonged to Capitán Prat Province. More recently, Bell et al. (2007) described a new genus of mosses, *Ombronesus* N.E.Bell, N.Pedersen & A.E.Newton, based on collections gathered by the Raleigh International expedition of 2003 in Katalalixar National Reserve, near Caleta Tortel. The floristic results of the latter expedition have not yet been published.

Unfortunately, in his checklist of Chilean mosses, Müller (2009a) reports most of the species listed by Cardot and Brotherus (1923) and all those reported by Donat (1936a) as belonging to Aisén Province, and did not mention the works by Skottsberg (1916) or by Donat (1936b). The result is that Müller (2009a) reports only seven taxa for Capitán Prat Province in his checklist. This partial information led Cuvertino et al. (2012) to report a new record for the province, *Distichium capillaceum* (Hedw.) Bruch & Schimp., although this taxon was already reported 100 years ago by Skottsberg (1916) from the NE shore of Lago O’Higgins. With the descrip-




Figure 1. Map of Aisén Region showing the four provinces (AI = Aisén; CO = Coihaique; GC = General Carrera; CP = Capitán Prat) and the itinerary of bryological expeditions. **A** Expeditions that collected mosses previous to this study: 1. Charles Darwin, 1834; 2. Per Dusén, 1896–97; 3. Carl Skottsberg, 1908–09; 4. Federico Reichert and Cristóbal Hicken, 1921; 5. Heikki Rovainen, 1929; 6. Arturo Donat, 1933; 7. Gerhard Schwabe, 1939–41; 8. Rolf Santesson, 1940–41; 9. Tarow Seki, 1967; 10. Hironori Deguchi, 1981 **B** Explored area during the 2007 expedition where collections reported here were made. Inset: map of Chile showing in red the location of Aisén Region. (dotted line = regional border; continuous line = provincial limit).

tion of the new species *Ombronesus stuvensis* (Bell et al. 2007), *Ulota billbuckii* Garilletti, Mazimpaka & F.Lara and *U. streptodon* Garilletti, Mazimpaka & F.Lara (Garilletti et al. 2012), and *U. larrainii* Garilletti, Mazimpaka & F.Lara (Garilletti et al. 2015), and the new reports of *Lepyrodon patagonicus* (Cardot & Broth.) B.H.Allen (Allen 1999), *Brachythecium subpilosum* (Hook.f. & Wilson) A.Jaeger (Cuvertino et al. 2012), *Hedwigia ciliata* var. *nivalis* Hampe (Ellis et al. 2014), *Philonotis esquelensis* Matteri (Jimenez et al. 2014), and *P. brevifolia* Herzog (Jimenez et al. 2016), the moss taxa known for the province rises to 49. This number is in contrast with the 250 moss taxa known from the neighboring Última Esperanza Province, or with the 311 taxa reported for Aisén Province (Müller 2009a).

Methods

Study site

Capitan Prat Province lies in the southern portion of Aisén Region (XI) in southern Chile, spanning between lat. 46°57'-49°09'S and long. 71°51'-75°36'W, with a total area of 37200 km². The province is crossed by three major river basins from north to south (the rivers Baker, Bravo and Pascua), with Baker river the largest river in Chile in terms of the volume of water. The province shows a remarkable heterogeneity of landscapes and climates (Fig. 2), ranging from extremely rainy areas in the western archipelago with rainfall measured up to 4266 mm a year at San Pedro station, to 191 mm in the steppe habitats in the easternmost areas, measured at Chile Chico, in adjacent General Carrera Province (di Castri and Hajek 1976), the closest available weather station in the easternmost side. Temperatures are mild, with a mean annual temperature measured at 8.2 °C at San Pedro and 11.5 °C at Chile Chico (di Castri and Hajek 1976). The geography is very rough, with snow-capped mountains dominating the landscape, Mt. San Lorenzo being the highest peak in the province, at 3706 m a.s.l. The province is flanked by the Northern and Southern Patagonian Ice Fields, and to the west it sinks into the Pacific Ocean in a large number of islands, fjords and channels, many of them still barely explored. It is one of the least populated provinces of Chile with a little more than 3000 people as measured by the 2012 census, and with more than twice the area of the whole Metropolitan Region of Chile, where more than 7 million people live.

In terms of the vegetation that dominates the landscape, there is a clearly marked gradient from east to west, mostly determined by the extreme variation in rainfall as explained above. In the easternmost parts of the province the vegetation fits into Gajardo's (1984) concept of the "Patagonian Steppe of Aisén", dominated by low shrubs and grasses, where the dominant species are *Baccharis patagonica* Hook. & Arn., *Stipa neaei* Nees ex Steud., *Festuca simpliciuscula* (Hack.) E.B.Alexeev, *Mulinum spinosum* Pers., *Adesmia longipes* Phil., and *Azorella incisa* Wedd., among others. This dry vegetation can be found in Estancia Chacabuco, around Lago Cochrane, and in Tamango National Reserve.

Descending in altitude, and towards the west, the vegetation is dominated by deciduous forests and shrublands, inserted in what Gajardo (1984) calls "Deciduous Forest fo Aisén". This vegetation is very heterogeneous depending on the degree of disturbance of the land and in the local variations in soils, exposition, slopes and drainage. The dominant tree species are *Nothofagus pumilio* (Poepp. & Endl.) Krasser, *N. betuloides* (Mirb.) Oerst., *N. antarctica* (G.Forst.) Oerst., and *Embothrium coccineum* J.R.Forst. & G.Forst. In the drier places shrublands can be found, dominated mostly by *Gaultheria mucronata* Hook. & Arn., *Chiliotrichum diffusum* (G.Forst.) Kuntze, *Berberis microphylla* G.Forst. and *Escallonia serrata* Sm. In the areas with stronger grazing impact, and where the forest has been cut, fields of *Taraxacum officinale* F.H.Wigg.,




Figure 2. Different habitats of Capitán Prat. **A** Steppe at Estancia Chacabuco **B** *Nothofagus dombeyi*–*N. antarctica*–*Pilgerodendron uviferum* forest at Lago Vargas **C** *Nothofagus dombeyi*–*Drimys winteri*–*Podocarpus nubigenus* forest at Río Bravo **D** Peatland at Lago Leal.

Holcus lanatus L. and *Dactylis glomerata* L. are common. The mosaic formed by these plant formations is frequent at mid altitude and is where most of the farming activity of the province is developed, mostly in the middle part of Río Baker.

Moving further towards the west, the vegetation changes into what Gajardo (1984) classifies as the region of “Evergreen Forests and Peatlands”. Several different kinds of forests coexist here, and there is a marked gradient too in its composition that relates with the available rainfall. The less humid forests are dominated by *Nothofagus dombeyi* (Mirb.) Oerst. and *N. nitida* (Phil.) Krasser, whereas the most humid are dominated by *Podocarpus nubigenus* Lindl., *Drimys winteri* J.R.Forst. & G.Forst., *Weinmannia trichosperma* Cav., *Raukaua laetevirens* (Gay) Frodin, and several Myrtaceae species. Areas with little drainage are dominated by *Pilgerodendron uviferum* (D.Don) Florin, *Tepualia stipularis* (Hook. & Arn.) Griseb., and *Sphagnum magellanicum* Brid. peatlands. Moving even further to the west the vegetation is dominated by cushion plants and shrubs like *Astelia pumila* (G.Forst.) Gaudich., *Donatia fascicularis* J.R.Forst. & G.Forst., *Oreobolus obtusangulus* Gaudich., *Empetrum rubrum* Vahl ex Willd. and *Lepidothamnus fonckii* Phil. (Gajardo 1984), which alternates with very humid evergreen forests in the ravines and protected places. This vegetation can be found along the coastline at Caleta Tortel, Puerto Yungay, and around Lago Quetru.

Data collection

Two field trips were made during the Austral summer of 2007, making up a total of 13 full days of collecting. During these expeditions a total of 1283 bryophyte collections were made throughout the area between the localities of Puerto Bertrand and Villa O'Higgins, most of them adjacent to the main vehicular roads (Route 7, the road to Tortel, and the road to Villa O'Higgins) spanning the whole basin of Baker River and some sites along the Bravo and Pascua River basins (Fig. 1B), distributed in 56 collecting sites. Of this total, 1091 collections corresponded to mosses, with the remainder being liverworts and hornworts. Additionally, 74 recent moss collections from this same area kept at the Universidad de Concepción herbarium (CONC) were examined (Suppl. material 1).

The collecting localities were arbitrarily chosen, attempting to include the largest number of different floristic associations and landscapes, both in the driest and the wettest areas of the province and throughout all the gradients in between, from the coast line to about 700 m a.s.l.

The specimens were identified with the monographs or taxonomic revisions currently available for each group, and comparing with reference material kept at CONC, NY, and F herbaria. Additionally, many colleagues helped with the identification of difficult material, or for groups where there are no revisions available: María Jesús Cano and Mayte Gallego (Pottiaceae), Ricardo Garilletti (*Ulota*), Paco Lara (*Orthotrichum*), Lars Hedenäs (Amblystegiaceae), John Spence (*Bryum*), Barbara Murray (*Andreaea*), Guillermo Suárez (*Pohlia*), Soledad Jimenez (*Philonotis*) and Barbara Andreas (*Blindia*). All vouchers are deposited in the Universidad de Concepción herbarium (CONC).

The species list is given below in alphabetical order, indicating the new records for Chile (***)¹, for Aisén Region (**) and for Capitán Prat Province (*). A systematic arrangement of the taxa, including habitat information, altitude where taxa were found, frequency in the studied area, global and Chilean distribution, and a list of studied specimens, is presented in the Suppl. material 1.

For the analysis of the distribution patterns, the many different patterns observed were reduced into five major categories for simplification: (1) Wide distribution (WD), meaning cosmopolitan or subcosmopolitan species present in both hemispheres and in both tropical and temperate areas of the planet; (2) Bipolar, meaning species distributed in the temperate areas of both hemispheres with eventual isolated populations at high altitudes in the tropics; (3) Austral, meaning species with various distribution patterns along the Southern Hemisphere - this is the most heterogeneous group because it includes both species with very wide circumsubantarctic distributions to narrow “sub-endemics” maybe recorded from a few populations in southern South America and South Africa, for example; (4) Andean, meaning species mostly distributed throughout the Andes but incidentally extending into Antarctica, Africa or some subantarctic Islands; and (5) Endemic, meaning species only known from southern South America and eventually extending into Juan Fernández or the Falkland and South Georgia Islands.

Results

A total of 260 moss taxa belonging to 256 species and 4 infraspecific taxa, in 42 botanical families are reported for Capitán Prat Province (Suppl. material 1). The most diverse families were the Orthotrichaceae with 33 taxa, the Pottiaceae with 24 and the Bryaceae with 20 taxa. The most species rich genera were *Bryum* with 20 species, *Ulota* with 15 and *Syntrichia* with 11 species. Two taxa are reported for continental Chile for the first time, 54 for Aisén Region, and 211 are for the first time reported for Capitán Prat Province, with 12 taxa extending their known distribution limits to the north, and 49 to the south.

- **Achrophyllum anomalum* (Schwägr.) H.Rob. var. *anomalum*
- Achrophyllum anomalum* var. *pallidum* (Cardot & Broth.) S.He
- **Achrophyllum haesselianum* (Matteri) Matteri
- **Achrophyllum magellanicum* (Besch.) Matteri (Figure 3G)
- Acrocladium auriculatum* (Mont.) Mitt. (Figure 3C)
- ***Acroschisma wilsonii* (Hook.f.) A.Jaeger
- **Amphidium tortuosum* (Hornschr.) Cufod.
- **Ancistrodes genuflexa* (Müll.Hal.) Crosby
- Andreaea alpina* Hedw.
- Andreaea fuegiana* (Cardot) S.W.Greene
- Andreaea regularis* Müll.Hal.
- Andreaea vaginalis* Herzog
- Arbusculohypopterygium arbuscula* (Brid.) M.Stech, T.Pfeiff. & W.Frey
- **Bartramia ithyphyloides* Schimp. ex Müll.Hal.
- Bartramia mossmaniana* Müll.Hal.
- Bartramia patens* Brid.
- **Bartramia robusta* Hook.f. & Wilson
- **Bartramia stricta* Brid.
- **Blindia connecta* (Hook.f. & Wilson) Müll.Hal.
- Blindia magellanica* Schimp.
- Brachytheciastrum paradoxum* (Hook.f. & Wilson) Ignatov & Huttunen
- **Brachythecium albicans* (Hedw.) Schimp.
- **Brachythecium austroglaresum* (Müll.Hal.) Paris
- **Brachythecium austrosalebrosum* (Müll.Hal.) Paris
- Brachythecium subpilosum* (Hook.f. & Wilson) A.Jaeger
- **Brachythecium subplicatum* (Hampe) A.Jaeger
- **Breutelia angustiretis* E.B.Bartram
- **Breutelia aureola* (Besch. ex Müll.Hal.) Besch.
- **Breutelia dumosa* Mitt.
- Breutelia integrifolia* (Taylor) A.Jaeger
- **Breutelia plicata* Mitt.
- **Breutelia subplicata* Broth.

- ***Bryum algovicum* Sendtn. ex Müll.Hal.
***Bryum archangelicum* Bruch & Schimp.
**Bryum australe* Hampe
***Bryum billarderi* Schwägr. (Figure 4E)
**Bryum caespiticium* Hedw.
***Bryum canariense* Brid.
**Bryum capillare* Hedw.
Bryum clavatum (Schimp.) Müll.Hal.
**Bryum coronatum* Schwägr.
Bryum crassinervium Lorentz
***Bryum dichotomum* Hedw.
Bryum donati Thér.
Bryum gemmatum Müll.Hal.
**Bryum laevigatum* Hook.f. & Wilson
**Bryum macrophyllum* Cardot & Broth.
**Bryum perlimbatum* Cardot
***Bryum platyphyllum* (Schwägr.) Müll.Hal.
Bryum pseudotriquetrum (Hedw.) Schwägr.
***Bryum puconense* Herzog & Thér.
**Bryum subapiculatum* Hampe
**Calyptopogon mnioides* (Schwägr.) Broth.
**Camptodontium cryptodon* (Mont.) Reimers
***Campylium stellatum* (Hedw.) C.E.O.Jensen
**Campyloporidium euchlorum* (Mont.) Matteri
**Campyloporidium medium* (Duby) Giese & J.-P.Frahm
**Campylopus acuminatus* Mitt.
**Campylopus clavatus* (R.Br.) Wilson
**Campylopus incrassatus* Müll.Hal.
**Campylopus introflexus* (Hedw.) Brid.
Campylopus purpureocaulis Dusén
**Campylopus pyriformis* (Schultz) Brid.
**Campylopus vesticaulis* Mitt.
**Catagonium nitens* (Brid.) Cardot
**Ceratodon purpureus* (Hedw.) Brid. subsp. *purpureus*
**Ceratodon purpureus* subsp. *convolutus* (Reichardt) Burley
Chorisodontium aciphyllum (Hook.f. & Wilson) Broth.
**Chorisodontium dicranellatum* (Dusén) Roiv.
**Chorisodontium spegazzinii* (Müll.Hal.) Roiv.
***Chrysoblastella chilensis* (Mont.) Reimers
**Conostomum pentastichum* (Brid.) Lindb.
Cratoneuropsis relaxa (Hook.f. & Wilson) M.Fleisch. ex Broth. subsp. *minor* (Hook.f. & Wilson) Ochyra
**Cryptothallia consimilis* Mont.

- **Cryphaeophilum molle* (Dusén) M.Fleisch.
- **Daltonia gracilis* Mitt.
- **Daltonia trachydonta* Mitt.
- **Dendroligotrichum dendroides* (Hedw.) Brid.
- Dendroligotrichum squamosum* (Hook.f. & Wilson) Broth. ex Cardot
- **Dicranella campylophylla* (Taylor) A.Jaeger
- **Dicranella hookeri* (Müll.Hal.) Cardot
- ***Dicranella pseudorufescens* Cardot & Broth.
- Dicranoloma billardieri* (Brid.) Paris
- **Dicranoloma chilense* (De Not.) Ochyra & Matteri
- ***Dicranoloma imponens* (Mont.) Renaud
- **Dicranoloma menziesii* (Hook.f. & Wilson) Paris
- **Dicranoloma perremotifolium* (Dusén) Broth.
- Dicranoloma robustum* (Hook.f. & Wilson) Paris
- **Didymodon andreaeoides* Cardot & Broth.
- **Didymodon australasiae* (Hook. & Grev.) R.H.Zander
- **Didymodon fuscus* (Müll.Hal.) J.A.Jiménez & M.J.Cano
- Distichium capillaceum* (Hedw.) Bruch & Schimp.
- **Distichophyllum dicksonii* (Hook. & Grev.) Mitt.
- **Ditrichum cylindricarpum* (Müll.Hal.) F.Muell.
- **Ditrichum difficile* (Duby) M.Fleisch.
- **Ditrichum ditrichoideum* (Cardot) Ochyra
- **Ditrichum heteromallum* (Hedw.) E.Britton
- ***Drepanocladus aduncus* (Hedw.) Warnst.
- **Drepanocladus longifolius* (Mitt.) Broth. ex Paris
- **Drepanocladus polygamus* (Schimp.) Hedenäs
- **Dryptodon austrofunalis* (Müll.Hal.) Ochyra & Żarnowiec
- **Dryptodon humilis* (Mitt.) Ochyra & Żarnowiec
- **Dryptodon navicularis* (Herzog) Ochyra & Żarnowiec
- Dryptodon trichophyllus* (Grev.) Brid.
- **Encalypta ciliata* Hedw.
- ***Encalypta rhaftocarpa* Schwägr.
- **Eriodon conostomus* Mont.
- **Eucamptodon perichaetialis* (Mont.) Mont.
- **Eurhynchiella acanthophylla* (Mont.) M.Fleisch.
- **Eustichia longirostris* (Brid.) Brid
- ***Fabronia jamesonii* Taylor
- **Fissidens curvatus* Hornsch.
- **Fissidens rigidulus* Hook.f. & Wilson
- **Funaria hygrometrica* Hedw.
- **Glyphothecium sciurooides* (Hook.) Hampe
- **Gymnostomum calcareum* Nees & Hornsch.
- ***Hebantia rigida* (Lorentz) G.L.Merr.

- Hedwigia ciliata* Hedw. var. *nivalis* Hampe (Figure 4D)
***Hennediella antarctica* (Ångström) Ochyra & Matteri
**Hennediella arenae* (Besch.) R.H.Zander
***Hennediella heimii* (Hedw.) R.H.Zander
**Hymenodontopsis mnioides* (Hook.) N.E.Bell, A.E.Newton & D.Quandt (Figure 4H)
Hymenoloma crispulum (Hedw.) Ochyra
Hymenoloma turpe (Cardot) Cardot & Broth.
**Hymenostylium recurvirostrum* (Hedw.) Dixon
**Hypnodendron microstictum* Mitt. ex A.Jaeger & Sauerb.
**Hypnum chrysogaster* Müll.Hal.
**Hypnum cupressiforme* Hedw. var. *cupressiforme*
***Hypnum cupressiforme* var. *filiforme* Brid.
**Hypnum cupressiforme* var. *mossmanianum* (Müll.Hal.) Ando
**Hypnum skottsbergii* Ando
**Hypopterygium didictyon* Müll.Hal.
**Kiaeria pumila* (Mitt.) Ochyra
**Leptobryum pyriforme* (Hedw.) Wilson
***Leptodontium longicaule* Mitt. var. *microruncinatum* (Dusén) R.H.Zander
***Leptostomum menziesii* R.Br.
Leptotheca gaudichaudii Schwäger.
**Lepyrodon hexastichus* (Mont.) Wijk & Margad.
Lepyrodon lagurus (Hook.) Mitt.
Lepyrodon patagonicus (Cardot & Broth.) B.H.Allen
**Lepyrodon tomentosus* (Hook.) Mitt.
***Looseria orbiculata* (Thér.) D.Quandt, Huttunen, Tangney & M.Stech
**Lopidium concinnum* (Hook.) Wilson
***Macromitrium pertriste* Müll.Hal.
***Mahua enervis* W.R.Buck
**Matteria gracillima* (Besch.) Goffinet
**Matteria papillosoala* (Thér.) Goffinet
**Neckera scabridens* Müll.Hal.
**Notoligotrichum minimum* (Cardot) G.L.Sm. (Figure 4A)
**Notoligotrichum trichodon* (Hook.f. & Wilson) G.L.Sm.
**Oligotrichum austroaligerum* G.L.Sm.
Ombronesus stuvensis N.E.Bell, N.Pedersen & A.E.Newton
***Orthodontium lineare* Schwägr.
**Orthotrichum assimile* Müll.Hal.
**Orthotrichum brotheri* Dusén ex Lewinsky
**Orthotrichum elegantulum* Schimp. ex Mitt.
***Orthotrichum hortense* Bosw.
**Orthotrichum incanum* Müll.Hal.
**Orthotrichum ludificans* Lewinsky
***Orthotrichum pariatum* Mitt.

- **Orthotrichum rupestre* Schleich. ex Schwägr.
- **Pararhacocarpus patagonicus* (Broth.) J.-P.Frahm
- Philonotis brevifolia* Herzog
- Philonotis esquelensis* Matteri
- **Philonotis krausei* (Müll.Hal.) Broth.
- ***Philonotis polymorpha* (Müll.Hal.) Kindb.
- ***Philonotis scabrifolia* (Hook.f. & Wilson) Braithw.
- Philonotis vagans* (Hook.f. & Wilson) Mitt.
- ***Pilopogon schilleri* Herzog & Thér.
- ***Plagiothecium lucidum* (Hook.f. & Wilson) Paris
- **Platyneuron praealtum* (Mitt.) Ochyra & Bednarek-Ochyra (Figure 3H)
- Pohlia cruda* (Hedw.) Lindb.
- ****Pohlia longicollis* (Hedw.) Lindb.
- **Pohlia nutans* (Hedw.) Lindb. (Figure 3A)
- **Pohlia wahlenbergii* (F.Weber & D.Mohr) A.L.Andrews (Figure 4G)
- **Polytrichadelphus magellanicus* (Hedw.) Mitt. (Figure 4B)
- ***Polytrichastrum alpinum* (Hedw.) G.L.Sm.
- **Polytrichastrum longisetum* (Sw. ex Brid.) G.L.Sm.
- **Polytrichum juniperinum* Hedw.
- Polytrichum piliferum* Hedw. (Figure 4C)
- ***Polytrichum strictum* Menzies ex Brid.
- **Porotrichum arbusculans* (Müll.Hal.) Ochyra
- **Pseudocrossidium crinitum* (Schultz) R.H.Zander
- Ptychomniella ptychocarpon* (Schwägr.) W.R.Buck, C.J.Cox, A.J.Shaw & Goffinet
- **Ptychomnion cygnisetum* (Müll.Hal.) Kindb.
- **Racomitrium didymum* (Mont.) Lorentz (Figure 3B)
- Racomitrium geronticum* Müll.Hal. (Figure 3F)
- **Racomitrium heterostichoides* Cardot
- Racomitrium laevigatum* A.Jaeger
- **Racomitrium lamprocarpum* (Müll.Hal.) A.Jaeger (Figure 4F)
- **Racomitrium orthotrichaceum* (Müll.Hal.) Paris
- **Racomitrium pachydictyon* Cardot
- **Racomitrium rupestre* (Hook.f. & Wilson) Wilson & Hook.f.
- **Racomitrium subcrispipilum* (Müll.Hal.) A.Jaeger
- **Rhacocarpus purpurascens* (Brid.) Paris
- ***Rhaphidorrhynchium amoenum* (Hedw.) M.Fleisch.
- **Rhaphidorrhynchium callidum* (Mont.) Broth. (Figure 3D)
- **Rhaphidorrhynchium dendroligotrichum* (Dusén) Broth.
- **Rigodium adpressum* Zomlefer
- **Rigodium brachypodium* (Müll.Hal.) Paris
- **Rigodium pseudothuidium* Dusén
- **Rigodium tamarix* Müll.Hal.
- **Rigodium toxarium* (Schwägr.) A.Jaeger var. *toxarium*




Figure 3. Mosses of Capitán Prat Province. **A** *Pohlia nutans* **B** *Racomitrium didymum* **C** *Acrocladium auriculatum* **D** *Rhaphidorrhynchium callidum* **E** *Tetraplodon fuegianus* **F** *Racomitrium geronicum* **G** *Achrophyllum magellanicum* **H** *Platyneuron praealtum*.



Figure 4. Mosses of Capitán Prat Province. **A** *Notoligotrichum minuum* **B** *Polytrichadelphus magellanicus* **C** *Polytrichum piliferum* **D** *Hedwigia ciliata* var. *nivalis* **E** *Bryum billardieri* **F** *Racomitrium lamprocarpum* **G** *Pohlia wahlenbergii* **H** *Hymenodontopsis mnioides*.

- ****Rigodium toxarion* var. *robustum* (Broth.) Zomlefer
**Sanionia uncinata* (Hedw.) Loeske
**Sauloma tenella* (Hook.f. & Wilson) Mitt.
Schistidium andinum (Mitt.) Herzog
***Sciuro-hypnum plumosum* (Hedw.) Ignatov & Huttunen
***Scorpidium revolvens* (Sw.) Rubers
Scouleria patagonica (Mitt.) A.Jaeger
**Sematophyllum scorpiurus* (Mont.) Mitt.
**Sphagnum falcatulum* Besch.
Sphagnum fimbriatum Wilson
Sphagnum magellanicum Brid.
Straminergon stramineum (Dicks. ex Brid.) Hedenäs
Syntrichia anderssonii (Ångström) R.H.Zander
***Syntrichia costesii* (Thér.) R.H.Zander
***Syntrichia epilosa* (Broth. ex Dusén) R.H.Zander
***Syntrichia fragilis* (Taylor) Ochyra
***Syntrichia glacialis* (Kunze ex Müll.Hal.) R.H.Zander
**Syntrichia lithophila* (Dusén) Ochyra & R.H.Zander
**Syntrichia magellanica* (Mont.) R.H.Zander
**Syntrichia princeps* (De Not.) Mitt.
**Syntrichia pseudorobusta* (Dusén) R.H.Zander
***Syntrichia ruralis* (Hedw.) F.Weber & D.Mohr
***Syntrichia scabrella* (Dusén) R.H.Zander
**Tayloria dubyi* Broth.
***Tayloria magellanica* (Brid.) Mitt.
***Tayloria stenophysata* (Herzog) A.K.Kop.
**Tetraplodon fuegianus* Besch. (Figure 3E)
***Tortella knightii* (Mitt.) Broth.
***Tortula atrovirens* (Sm.) Lindb.
Ulota billbuckii Garilleti, Mazimpaka & F.Lara
***Ulota carinata* Mitt.
**Ulota fuegiana* Mitt.
**Ulota germana* (Mont.) Mitt.
Ulota larrainii Garilleti, Mazimpaka & F.Lara
**Ulota luteola* (Hook.f. & Wilson) Wijk & Marg.
***Ulota macrocalycina* Mitt.
**Ulota macrodonzia* Dixon & Malta
**Ulota magellanica* (Mont.) A.Jaeger
***Ulota phyllantha* Brid.
***Ulota pusilla* Malta
***Ulota pycnophylla* Dusén ex Malta
**Ulota pygmaeothecia* (Müll.Hal.) Kindb.
**Ulota rufula* (Mitt.) A.Jaeger

- Ulota streptodon* Garilleti, Mazimpaka & F.Lara
- **Vittia pachyloma* (Mont.) Ochyra
- **Warnstorffia exannulata* (Schimp.) Loeske
- ***Warnstorffia fluitans* (Hedw.) Loeske
- **Warnstorffia fontinialopsis* (Müll.Hal.) Ochyra
- **Weymouthia cochlearifolia* (Schwägr.) Dixon
- **Weymouthia mollis* (Hedw.) Broth.
- ***Zygodon hookeri* Hampe var. *hookeri*
- **Zygodon hookeri* var. *leptobolax* (Müll.Hal.) Calabrese
- **Zygodon magellanicus* Dusén ex Malta
- **Zygodon papillatus* Mont.
- **Zygodon pentastichus* (Mont.) Müll.Hal.
- **Zygodon pichinchensis* (Taylor) Mitt.
- Zygodon reinwardtii* (Hornschr.) A.Braun

From the analysis of the five simplified distribution patterns, 38.08% (n=99) of the taxa are actually endemics of southern South America, followed by the “austral” taxa with 30.77% (n=80). The rest of the taxa were found to be either of “wide distribution” (13.85%, n=36), “bipolar” (9.62%, n=25), or “Andean” (7.69%, n=20).

Discussion

The known diversity of Capitán Prat Province was incremented from 49 to 260 taxa, representing a 531% increment. This is remarkable and demonstrates the very little attention this area has historically received in terms of its bryophyte diversity. The current number of species reported for the province makes sense with the numbers known from the adjacent provinces of Aisén (311) and Última Esperanza (250). The number of moss taxa known from adjacent General Carrera province [27 in Müller's (2009a) checklist] is still very low and certainly needs further study, although General Carrera province lacks the most humid habitats found in the coastal lowland rainforests, that in this study yielded a large number of taxa not found anywhere else.

Three taxa were described as new based on the collections gathered during this study in Capitán Prat (*Ulota billbuckii*, *U. larraini* and *U. streptodon*), and four others were found to be new records for Chile (*Hedwigia ciliata* var. *nivalis*, *Philonotis esquelensis*, *Pohlia longicollis* and *Rigodium toxarium* var. *robustum*), the latter previously known to be an endemic of the Juan Fernández Islands. The number of new species and new records for Chile might increment even more after study of the material that still remains unidentified.

The high level of endemic taxa in southern South America temperate rainforests has already been reported in the literature for neighboring areas (i.e., Seki 1974, Villagrán et al. 2003, Larraín 2005). This is also the principal distribution pattern in Capitán Prat Province, with almost 40% of all the taxa known for the area being endemics.

It is also interesting that the moss flora of these Austral forests is way more related to distant Austral lands as New Zealand, Australia, or some remote subantarctic islands rather than to neighboring Neotropical countries like Perú or Bolivia: only 7.69% of the taxa studied has a continuous distribution along the Andes into the northern Tropics. This is explained by the condition of “biogeographic island” of southern Chile forests, isolated from the rest of the continent by the very high peaks of central and northern Chile, and extremely arid deserts both to the north (Atacama) and to the east (Patagonian steppe).

Several species were found farther south from their previously known distribution ranges. This is interesting as it would appear to suggest that many of these taxa are apparently not able to disperse south of the Southern Patagonia Ice Field. This ice field would act as a natural barrier to the dispersion of Valdivian rainforest endemics such as *Ancistrodes genuflexa*, *Cryphaea consimilis*, *Cryphaeophilum molle*, *Eriodon conostomus*, *Eurhynchiella acanthophylla*, *Rhaphidorrhynchium dendroligotrichum*, *Rigidium tamarix*, and *Weymouthia cochlearifolia*, among others, preventing their colonization of Magallanes Region. This might be due also to less collecting effort in the more wet areas of Magallanes, because of the logistical difficulties to access the most western islands, where rainfall is similar to what can be found in the wettest areas of Capitán Prat.

It is important to mention that not a single locality above 700 m was visited. Although most of the area above this altitude is covered in ice and snow all year round, it would be interesting to visit higher altitude spots and glaciers, since several of the species found by Donat around the glaciers of Lago O’Higgins were not found during this expedition. Covering these sites would maybe increase even more the number of moss taxa known from Capitán Prat Province.

Acknowledgements

This work was made as a consultancy for Hidroaysén hydroelectric project within the framework of the baseline study prior to the environmental-impact study as required by the Chilean authority. I am specially acknowledged to the staff at the botany department of Universidad de Concepción, and to the many colleagues that helped with the determination of difficult groups of mosses: María Jesús Cano, Mayte Gallego, Ricardo Garilleti, Paco Lara, Lars Hedenäs, John Spence, Barbara Murray, Guille Suárez, Sol Jimenez and Barbara Andreas. Reinaldo Vargas and Götz Palfner are acknowledged for company during the fieldwork. Thanks also to Alicia Marticorena, Alejandra Jiménez, Ernesto Teneb and Roberto Rodríguez for making their collections from the area available for my study. I acknowledge the staff at CONAF office in Cochrane for allowing me to collect inside Tamango National Reserve. Finally I would like to thank the staff at CONC, F and NY herbaria for providing working space and full access to their bryophyte collections.

References

- Allen BH (1987) A revision of the Dicnemonaceae (Musci). *Journal of the Hattori Botanical Laboratory* 62: 1–100.
- Allen BH (1999) A revision of the moss genus *Lepyrodon* (Leucodontales, Lepyrodontaceae). *Bryobrothera* 5: 23–48.
- Allen BH (2002) Moss Flora of Central America, Part 2. Encalyptaceae-Orthotrichaceae. Monographs in Systematic Botany from the Missouri Botanical Garden 90. Missouri Botanical Garden Press, Saint Louis, 1–699.
- Ando H (1972a) Distribution and speciation in the genus *Hypnum* in the circum-Pacific region. *Journal of the Hattori Botanical Laboratory* 35: 68–98.
- Ando H (1972b) A phytogeographical account of *Hypnum cypresiforme* Hedw. in Japan. *Bulletin of the Yokohama Phytosociological Society of Japan* 16: 339–348.
- Ando H (1982) *Hypnum* in Australasia and the Southern Pacific. *Journal of the Hattori Botanical Laboratory* 52: 93–106.
- Ando H, Matteri CM (1982) Patagonian bryophytes 7. *Hypnum revolutum* (Mitt.) Lindb. new to South America, with notes on the *Hypnum* species known from the Transecta zone, S Argentina. *Lindbergia* 8(1): 60–64.
- Andreas B (2013) A revision of *Blindia* (Seligeriaceae) from southern South America. *The Bryologist* 116(3): 263–280. doi: 10.1639/0007-2745-116.3.263
- Arts T (2001) The moss genus *Leptobryum* and the identity of *Pohlia integra*. *Journal of Bryology* 23(4): 325–330. doi: 10.1179/jbr.2001.23.4.325
- Bartlett JK, Frahm J-P (1983) Notes on *Campylopus* and *Chorisodontium* from New Zealand. *Journal of Bryology* 12(3): 365–382. doi: 10.1179/jbr.1983.12.3.365
- Bartlett JK, Vitt DH (1986) A survey of species in the genus *Blindia* (Bryopsida, Seligeriaceae). *New Zealand Journal of Botany* 24(2): 203–246. doi: 10.1080/0028825X.1986.10412674
- Bednarek-Ochyra H (2006) A monograph of the moss genus *Codriophorus* P. Beauv. (Grimmiaceae). W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków, 1–276.
- Bednarek-Ochyra H (2011) *Bucklandiella striatipila* (Cardot) Bednarek-Ochyra & Ochyra [in Tristan da Cunha]. In: Ellis TL (Ed.) New national and regional bryophyte records, 28. *Journal of Bryology* 33(3): 239–240. doi: 10.1179/1743282011Y.0000000022
- Bednarek-Ochyra H (2014) Taxonomic status of two subantarctic species of *Philonotis* (Bartramiaceae, Musci). *Cryptogamie, Bryologie* 35(4): 387–395. doi: 10.7872/cryb.v35.iss4.2014.387
- Bednarek-Ochyra H, Ochyra R (2009) *Bucklandiella striatipila* (Cardot) Bednarek-Ochyra & Ochyra [in Îles Crozet]. In: Blockeel TL (Ed.) New national and regional bryophyte records, 22. *Journal of Bryology* 31(3): 202. doi: 10.1179/037366809X12469790518367
- Bednarek-Ochyra H, Ochyra R (2010) *Bucklandiella striatipila* (Cardot) Bednarek-Ochyra & Ochyra [in Îles Kerguelen]. In: Ellis TL (Ed.) New national and regional bryophyte records, 25. *Journal of Bryology* 32(4): 311–312. doi: 10.1179/jbr.2010.32.4.311
- Beever J, Allison KW, Child J (1992) The Mosses of New Zealand. Second edition. University of Otago Press, Dunedin, 1–214.

- Beever J, Malcolm B, Malcolm N (2002) The moss genus *Fissidens* in New Zealand. An illustrated key. Micro-Optics Press, Nelson, 1–99.
- Bell BG (1974) A synoptic flora of South Georgian mosses: V. *Willia* and *Racomitrium*. British Antarctic Survey Bulletin 38: 73–101.
- Bell NE, Pedersen N, Newton A (2007) *Ombronesus stuvensis*, a new genus and species of the Ptychomniaceae (Bryophyta) from south west Chile. Taxon 56(3): 887–896. doi: 10.2307/25065870
- Biasuso AB (2007) The genus *Hedwigia* (Hedwigiaceae, Bryophyta) in Argentina. Lindbergia 32(1): 5–17. <http://www.jstor.org/stable/20150227>
- Blockeel TL (1990) The genus *Hennediella* Par.: a note on the affinities of *Tortula brevis* Whitehouse & Newton and *T. stanfordensis* Steere. Journal of Bryology 16(2): 187–192. doi: 10.1179/jbr.1990.16.2.187
- Buck WR (1983) A synopsis of the South American taxa of *Fabronia* (Fabroniaceae). Brittonia 35(3): 248–254. doi: 10.2307/2806024
- Buck WR, Ireland RR (1989) Plagiotheciaceae. Flora Neotropica Monographs 50: 1–22.
- Burley JS, Pritchard NM (1990) Revision of the genus *Ceratodon* (Bryophyta). Harvard Papers in Botany 2: 17–76.
- Calabrese GM (2006) A taxonomic revision of *Zygodon* (Orthotrichaceae) in southern South America. The Bryologist 109(4): 453–509. doi: 10.1639/0007-2745(2006)109[453:ATR OZO]2.0.CO;2
- Calabrese GM, Matteri CM (1999) Los musgos (Bryophyta) de Puerto Blest y alrededores (Parque Nacional Nahuel Huapi, Argentina) I. Lista comentada de las especies. Revista del Museo Argentino de Ciencias Naturales, nueva serie, 1(1): 23–48.
- Cano MJ (2008) Taxonomic revision of *Hennediella* Paris (Pottiaceae, Bryophyta). Bryophytorum Bibliotheca 64. J. Cramer, Berlin, 1–142.
- Cano MJ, Gallego MT (2008) The genus *Tortula* (Pottiaceae, Bryophyta) in South America. Botanical Journal of the Linnean Society 156: 173–220. doi: 10.1111/j.1095-8339.2007.00739.x
- Cardot J (1908) La flora bryologique des Terres magellaniques, de la Géorgie du Sud et de l'Antarctide. Wissenschaftliche Ergebnisse der Schwedischen Südpolar-Expedition 1901–1903 4(8): 1–298, + 11 plates.
- Cardot J (1916) Note sur des Mousses de Kerguelen. Bulletin du Muséum d'Histoire Naturelle 22: 336–341.
- Cardot J, Brotherus VF (1923) Botanische Ergebnisse der Schwedischen Expedition nach Patagonien und dem Feuerlande 1907–1909. X. Les Mousses. Kungl. Svenska Vetenskapsakademiens Handlingar 63(10): 1–73.
- Churchill SP (1985) The systematics and biogeography of *Scouleria* Hook. (Musci: Scouleriaceae). Lindbergia 11(1): 59–71. <http://www.jstor.org/stable/20149542>
- Churchill SP, Buck WR (1982) A taxonomic investigation of *Leptotheca* (Rhizogoniaceae). Brittonia 34(1): 1–11. doi: 10.2307/2806391
- Churchill SP, Linares EL (1995) Prodromus Bryologiae Novo-Granatensis, Introducción a la flora de musgos de Colombia. Instituto de Ciencias Naturales - Museo de Historia Natural, Biblioteca “José Jerónimo Triana”, No. 12, Bogotá, 1–924.

- Crum H (1984) Sphagnopsida, Sphagnaceae. North American Flora. Series II, 11: i-ii, 1–180.
- Crum HA, Anderson LE (1981) Mosses of Eastern North America (2 vols.). Columbia University Press, New York, 1–1328.
- Cuvertino J, Ardiles V, Osorio F, Romero X (2012) New records and additions to the Chilean bryophyte flora. Ciencia e Investigación Agraria 39(2): 245–254. doi: 10.4067/S0718-16202012000200001
- Dalton PJ, Seppelt RD, Buchanan AM (1991) Annotated checklist of the moss flora of Tasmania. In: Banks MR, Smith SJ, Orchard AE, Kantvilas G (Eds) Aspects of Tasmanian Botany. A tribute to Winifred Curtis. Royal Society of Tasmania, Hobart, 15–32.
- di Castri F, Hajek ER (1976) Bioclimatología de Chile. Editorial Universidad Católica de Chile, Santiago, 1–160.
- Dixon HN (1926) Studies in the bryology of New Zealand, with special reference to the herbarium of Robert Brown. New Zealand Institute Bulletin 3(4): 153–238. doi: 10.5962/bhl.title.25898
- Donat A (1936a) Resultados fitogeográficos de la expedición patagónica “Gaea” 1933. I. Algunas briófitas nuevas de la Cordillera Central Patagónica. Revista Sudamericana de Botánica 3(3): 61–68.
- Donat A (1936b) Bericht über die pflanzengeographischen Ergebnisse der Inlandeisexpedition der Argentinischen Geographischen Gesellschaft “Gaea” 1933. Berichten Deutschen Botanischen Gesellschaft 54(1): 27–46.
- Dusén P (1906) Beiträge zur Bryologie der Magellanslander von Westpatagonien und Sudchile. V. Arkiv för Botanik utgivet av K. Svenska Vetenskaps-Akademien 6(10): 1–32 + 6 plates.
- Eckel PM (1997) *Tortella knightii*, an Australasian species new to the South American continent in Tierra del Fuego. Bryologist 100(2): 210–211. <http://www.jstor.org/stable/3244052>
- Ellis LT, Aleffi M, Tacchi R, Alegro A, Alonso M, Asthana AK, Sahu V, Biasoso AB, Callaghan DA, Ezer T, Kara R, Seyli T, Garilleti R, Gil-López MJ, Gwynne-Evans D, Hedderson TA, Kiebacher T, Larraín J, Long D, Lüth M, Malcolm B, Mamontov YS, Newsham KK, Nobis M, Nowak A, Ochyra R, Pawlikowski P, Plášek V, Čihal L, Potemkin AD, Puche F, Rios D, Gallego MT, Guerra J, Sawicki J, Schäfer-Verwimp A, Segarra-Moragues JG, Šegota V, Sofronova EV, Štefanuť S, Szűcs P, Bidló A, Papp B, Szurdoki E, Tan BC, Váňa J, Vigalondo B, Draper I, Lara F, Yoon Y-J, Sun B-Y, Nishimura N (2014) New national and regional bryophyte records, 41. Journal of Bryology 36(4): 306–324. doi: 10.1179/1743282014Y.0000000123
- Essl F, Steinbauer K, Dullinger S, Mang T, Moser D (2014) Little, but increasing evidence of impacts by alien bryophytes. Biological Invasions 16(5): 1175–1184. doi: 10.1007/s10530-013-0572-2
- Frahm J-P (1983) A monograph of *Pilopogon* Brid. Lindbergia 9(2): 99–116. <http://www.jstor.org/stable/20149474>
- Frahm J-P (1984) *Campylopus* Brid. 1. Subgenus *Thysanomitrium* (Schwaegr.) Kindb. emend. J.-P. Frahm. Nova Hedwigia 39: 585–621.
- Frahm J-P (1988) The subantarctic and Southern Hemispheric species of *Campylopus* (Dicranaceae), with contributions to the origin and speciation of the genus. Journal of the Hattori Botanical Laboratory 64: 367–387.

- Frahm J-P (1990) A short survey of the *Campylopus* flora of New Caledonia. Cryptogamie: Bryologie, Lichénologie 11(4): 369–375.
- Frahm J-P (1991) Dicranaceae: Campylopodioidae, Paraleucobryoideae. Flora Neotropica Monographs 54: 1–237.
- Frahm J-P (1994) A new synopsis of the *Campylopus* species from Australia. Journal of Bryology 18(2): 311–327. doi: 10.1179/jbr.1994.18.2.311
- Frahm J-P (1996) Revision der gattung *Rhacocarpus* Lindb. (Musc). Cryptogamie: Bryologie, Lichénologie 17: 39–65.
- Frahm J-P, Klöcker T, Schmidt R, Schöter C (2000) Revision der Gattung *Amphidium* (Musc), Dicranaceae. Tropical Bryology 18: 173–184.
- Fransén S (2004a) Taxonomic revision of extra-Neotropical *Bartramia* section *Vaginella* C. Müll. Lindbergia 29(2): 73–107. <http://www.jstor.org/stable/20150159>
- Fransén S (2004b) A taxonomic revision of *Bartramia* Hedw. section *Bartramia*. Lindbergia 29(3): 113–122. <http://www.jstor.org/stable/20150166>
- Gajardo R (1984) La vegetación natural de Chile. Clasificación y distribución geográfica. Editorial Universitaria, Santiago, 1–165.
- Gallego MT (2005) A taxonomic study of the genus *Syntrichia* Brid. (Pottiaceae, Musci) in the Mediterranean region and Macaronesia. Journal of the Hattori Botanical Laboratory 98: 47–122.
- Gallego MT, Cano MJ, Ros RM, Guerra J (2002) An overview of *Syntrichia ruralis* complex (Pottiaceae, Musci) in the Mediterranean region and neighbouring areas. Botanical Journal of the Linnaean Society 138: 209–224. doi: 10.1046/j.1095-8339.2002.138002209.x
- Garilletti R, Mazimpaka V, Lara F (2012) New *Ulota* species with multicellular spores from southern South America. The Bryologist 115(4): 585–600. doi: 10.1639/0007-2745-115.4.585
- Garilletti R, Mazimpaka V, Lara F (2015) *Ulota larrainii* (Orthotrichoideae, Orthotrichaceae, Bryophyta) a new species from Chile, with comments on the worldwide diversification of the genus. Phytotaxa 217(2): 133–144. doi: 10.11646/phytotaxa.217.2.3
- Giese M, Frahm J-P (1985 [1986]) A revision of *Campylopodium* (C. Müll.) Besch. Lindbergia 11(2-3): 125–133. <http://www.jstor.org/stable/20149553>
- Goffinet B, Shaw AJ, Cox CJ (2010) Phylogenetic inferences in the dung-moss family Splachnaceae from analyses of cpDNA sequence data and implications for the evolution of entomophily. American Journal of Botany 91(5): 748–759. doi: 10.3732/ajb.91.5.748
- Gradstein SR, Churchill SP, Sallazar-Allen N (2001) Guide to the Bryophytes of Tropical America. Memoirs of The New York Botanical Garden 86: 1–577.
- Hassel K, Söderström L (2005) The expansion of the alien mosses *Orthodontium lineare* and *Campylopus introflexus* in Britain and continental Europe. Journal of the Hattori Botanical Laboratory 97: 183–193.
- Hattaway RA (1984) A monograph of the Ptychomniaceae (Bryopsida). PhD Thesis, University Ann Arbor, Michigan, USA.
- He S (1998) A checklist of the mosses of Chile. Journal of the Hattori Botanical Laboratory 85: 103–189.
- Hedenäs L (2003) Amblystegiaceae (Musc). Flora Neotropica 89. Organization for Flora Neotropica by The New York Botanical Garden, New York, 1–108.

- Horton DG (1983) A revision of the Encalyptaceae (Musci), with particular reference to the North American taxa. Part II. *Journal of the Hattori Botanical Laboratory* 54: 353–532.
- Hyvönen J (1987) A synopsis of the genus *Leptostomum* (Leptostomataceae, Musci). *Acta Botanica Fennica* 24: 63–72. <http://www.jstor.org/stable/23725591>
- Hyvönen J (1991) *Chorisodontium* (Dicranaceae, Musci) in southern South America. *Annales Botanici Fennici* 28: 247–258.
- Ireland RR, Bellolio G (2002) The mosses of Easter Island. *Tropical Bryology* 21: 11–19.
- Ireland RR, Robinson H, Bellolio G (2006) On the distribution and morphology of *Hebantia rigida* (Bryopsida, Polytrichaceae). *Journal of the Hattori Botanical Laboratory* 100: 201–209.
- Jiménez JA, Cano MJ (2006) Two new combinations in *Didymodon* (Pottiaceae) from South America. *The Bryologist* 109: 391–397. doi: 10.1639/0007-2745(2006)109[391:tncidp]2.0.co;2
- Jimenez MS, Suárez GM, Schiavone MM (2014) Fruiting material of *Philonotis esquelensis* (Bartramiaceae, Bryophyta) discovered in Chile. *Boletín de la Sociedad Argentina de Botánica* 49(4): 463–471. <http://ref.scielo.org/c95dz2>
- Jimenez MS, Suárez GM, Larraín J (2016) Rediscovery and lectotypification of *Philonotis brevifolia* Herzog (Bartramiaceae, Bryophyta), a neglected species from Chile. *Cryptogamie, Bryologie* 37(2): 113–118. doi: 10.7872/cryb/v37.iss2.2016.113
- Jofre J, Massardo F, Rozzi R, Goffinet B, Marino P, Raguso RA, Navarro NP (2010) Fenología de *Tayloria dubyi* (Splachnaceae) en las turberas de la Reserva de Biosfera Cabo de Hornos. *Revista Chilena de Historia Natural* 83: 195–206. doi: 10.4067/S0716-078X2010000100012
- Jofre J, Goffinet B, Marino P, Raguso RA, Nihei SS, Massardo F, Rozzi R (2011) First evidence of insect attraction by a Southern Hemisphere Splachnaceae: the case of *Tayloria dubyi* Broth. in the Reserve Biosphere Cape Horn, Chile. *Nova Hedwigia* 92(3-4): 317–326. doi: 10.1127/0029-5035/2011/0092-0317
- Karlin E, Boles SB, Shaw AJ (2008) Systematics of *Sphagnum* section *Sphagnum* in New Zealand: A microsatellite-based analysis. *New Zealand Journal of Botany* 46(2): 105–118. doi: 10.1080/00288250809509758
- Karlin E, Buck WR, Seppelt RD, Boles SB, Shaw AJ (2013) The double allopolyploid *Sphagnum x falcatum* (Sphagnaceae) in Tierra del Fuego, a Holantarctic perspective. *Journal of Bryology* 35(3): 157–172. doi: 10.1179/1743282013Y.0000000066
- Klazenga N (2003) A revision of the Australasian species of *Dicranoloma* (Bryophyta, Dicranaceae). *Australian Systematic Botany* 16(4): 427–471. doi: 10.1071/SB02032
- Koponen A (1982) The family Splachnaceae in Australasia and the Pacific. *Journal of the Hattori Botanical Laboratory* 52: 87–91.
- Krijger JD (2002) Hypopterygiaceae of the World. *Blumea, Supplement* 13: 1–388.
- Kühnemann O, Gonçalves Carralves MF (1976) Sobre *Cryptaeophilum molle* (Dus.) Fleisch. Meteoriaceae-Briofitas. *Boletín de la Sociedad Argentina de Botánica* 17(3-4): 247–251.
- Larraín J (2005) Musgos de la Cordillera de la Costa de Valdivia, Osorno y Llanquihue: consideraciones ecológicas y lista de especies. In: Smith-Ramirez C, Armesto JJ, Valdovinos C (Eds) *Historia, Diversidad y Ecología de los Bosques Costeros de Chile*. Editorial Universitaria, Santiago, Chile, 159–177.

- Larraín J (2007) Adiciones a la flora de musgos de la Isla Grande de Chiloé, Chile. *Gayana Botánica* 64(1): 7–23. doi: 10.4067/S0717-66432007000100002
- Larraín J (2012) Filogenia del género *Racomitrium* Brid. (Bryophyta, Grimmiaceae) y taxonomía de las especies latinoamericanas. PhD Thesis, Universidad de Concepción, Chile.
- Larraín J, Suárez G, Bednarek-Ochyra H, Ochyra R (2010) The rediscovery of *Dicranella circinata* Herzog, a rare endemic moss from southern Chile, with comments on other southern South American species of *Dicranella*. *Nova Hedwigia* 91(3-4): 361–376. doi: 10.1127/0029-5035/2010/0091-0361
- Lewinsky J (1984) *Orthotrichum* Hedw. in South America 1. Introduction and taxonomic revision of taxa with immersed stomata. *Lindbergia* 10(2): 65–94. <http://www.jstor.org/stable/20149513>
- Lewinsky J (1987) *Orthotrichum* Hedw. (Orthotrichaceae) in South America 2. Taxonomic revision of taxa with superficial stomata. *Memoirs of the New York Botanical Garden* 45: 326–370.
- Lightowers PJ (1986) Taxonomy and distribution of the subantarctic species of *Tortula*. *Journal of Bryology* 14(2): 281–295. doi: 10.1179/jbr.1986.14.2.281
- Lin SH (1984) A taxonomic revision of Phyllogoniaceae (Bryopsida). Part II. *Journal of Taiwan Museum* 37: 1–54.
- Malta N (1927) Die *Ulota*-Arten Sud-Amerikas. *Acta Horti Botanici Universitatis Latviensis* 2: 165–208.
- Matteri CM (1968) Las especies de “*Philonotis* (Bartramiaceae)” del sur de Argentina. *Revista del Museo Argentino Ciencias Naturales “Bernardino Rivadavia” e Instituto Nacional de Investigación de las Ciencias Naturales Botánica* 3(4): 185–234.
- Matteri CM (1972) Las Hookeriaceae (Musci) Andino-Patagónicas II. *Revista del Museo Argentino Ciencias Naturales “Bernardino Rivadavia” e Instituto Nacional de Investigación de las Ciencias Naturales Botánica* 4(2): 243–280.
- Matteri CM (1973) El género *Breutelia* (Bartramiaceae, Musci) en la región Andino-Patagónica. *Revista del Museo Argentino Ciencias Naturales “Bernardino Rivadavia” e Instituto Nacional de Investigación de las Ciencias Naturales Botánica* 4(4): 321–360.
- Matteri CM (1975) Las Hookeriaceae (Musci) Andino-Patagónicas, I. *Nova Hedwigia* 26: 649–724.
- Matteri CM (1982) Patagonian Bryophytes 6. Fruiting *Sarconeurum glaciale* (C. Muell.) Card. et Bryhn newly found in southern Patagonia. *Lindbergia* 8(2): 105–109. <http://www.jstor.org/stable/20149428>
- Matteri CM (1985) Bryophyta, Musci: Bartramiaceae. *Flora Criptogámica de Tierra del Fuego* 14(7): 1–62.
- Matteri CM (2000) Sobre la identidad de *Orthotrichum inclinatum*, *O. incanum* y *O. crenato-erosum* (Musci, Orthotrichaceae). *Revista del Museo Argentino de Ciencias Naturales*, n.s. 2(1): 17–21.
- Matteri CM (2003) Los musgos (Bryophyta) de Argentina. *Tropical Bryology* 24: 33–100.
- Matteri CM, Calabrese GM (1999) Taxonomic notes and distribution of *Matteria papillosum* and *Microdus euchlorus*, from northern Patagonia (Argentina), with a new combination in *Campylopodium*. *Annales Botanici Fennici* 36: 43–49. <http://www.jstor.org/stable/23726646>

- Matteri CM, Schiavone MM (2002) Catálogo de los musgos (Bryophyta) de la Región Fueguina en Argentina y Chile. Revista del Museo Argentino de Ciencias Naturales, n.s. 4(2): 111–138.
- McFarland KD (1988) Revision of *Brachythecium* (Muscii) for Mexico, Central America, South America, Antarctica and circum-subantarctic Islands. PhD Thesis, University of Tennessee, Knoxville, USA.
- Menzel M (1992) Preliminary checklist of the mosses of Peru (Studies on Peruvian Bryophytes IV). Journal of the Hattori Botanical Laboratory 71: 175–254.
- Mitten W (1859) Descriptions of some new species of Musci from New Zealand and other parts of the Southern Hemisphere, together with an enumeration of the species collected in Tasmania by William Archer, arranged upon the plan proposed in the *Musci Indiae Orientalis* (1859). Journal of the Proceedings of the Linnean Society 4: 64–100.
- Müller C (1849) Synopsis Muscorum Frondosorum omnium hucusque cognitorum. Pars prima. Sumpibus Alb. Foerstener, Berolini, 1–812.
- Müller C (1885) Bryologia fuegiana. Flora 68: 391–429.
- Müller C (1898) Symbolae ad bryologiam Australiae II. Hedwigia 37: 76–171.
- Müller F (2009a) An updated checklist of the mosses of Chile. Archive for Bryology 58: 1–124.
- Müller F (2009b) New records and new synonyms for the Chilean moss flora. Tropical Bryology 30: 77–84.
- Müller F, Pursell RA (2003) The genus *Fissidens* (Muscii, Fissidentaceae) in Chile. Journal of the Hattori Botanical Laboratory 93: 117–139.
- Muñoz J (1999) A revision of *Grimmia* (Muscii, Grimmiaceae) in the Americas. 1: Latin America. Annals of the Missouri Botanical Garden 86(1): 118–191. doi: 10.2307/2666219
- Muñoz J, Pando F (2000) A world synopsis of the genus *Grimmia*. Monographs in Systematic Botany from the Missouri Botanical Garden 83: (i–viii) + 1–113.
- Murray B (2006) Andreaeaceae. Flora of Australia 51: 108–123.
- O’Shea BJ (2006) Checklist of the mosses of sub-Saharan Africa (version 5, 12/06). Tropical Bryology Research Reports 6: 1–252.
- Ochi H (1970) A revision of the subfamily Bryoideae in Australia, Tasmania, New Zealand and adjacent islands. The Journal of the Faculty of Education, Tottori University, Natural Sciences 21: 7–67.
- Ochi H (1980) A revision of the Neotropical Bryoideae, Musci (First part). Journal of the Faculty of Education, Tottori University, Natural Science 29(2): 49–154.
- Ochi H (1982) A revision of the Bryoideae (Muscii) in southern South America. Journal of the Faculty of Education, Tottori University, Natural Science 31: 11–47.
- Ochi H (1990) Bryaceous mosses as the source of distributional records from the Andes region. Bulletin of the National Science Museum, (Tokyo) Series B, Botany 16(2): 61–71.
- Ochyra R (1999) Antipodal mosses: X. The status of *Dichodontium brasiliense* and *D. opacifolium* (Dicranaceae). Fragmenta Floristica et Geobotanica 44: 227–232.
- Ochyra R, Bednarek-Ochyra H (1997) The type specimen of *Leptotrichum prealtum* (Muscii, Ditrichaceae). Fragmenta Floristica et Geobotanica 42: 567–570.
- Ochyra R, Lewis Smith RI, Bednarek-Ochyra H (2008) The Illustrated Moss Flora of Antarctica. Cambridge University Press, Cambridge, 1–685.

- Ochyra R, Matteri CM (2001) Bryophyta, Musci: Amblystegiaceae. Flora Criptogámica de Tierra del Fuego 14(10): 1–96.
- Ochyra R, Newton ME (1985) The taxonomy and distribution of *Dicranella cardotii* (R.Br. ter.) Dix., an addition to the moss flora of Antarctica. Lindbergia 11(2-3): 94–98. <http://www.jstor.org/stable/20149550>
- Ochyra R, Plášek V (2016) *Racomitrium patagonicum* Bednarek-Ochyra & Ochyra [in Chile]. In: Ellis TL (Ed.) New national and regional bryophyte records, 46. Journal of Bryology 38(1): 54. doi: 10.1080/03736687.2015.1123344
- Ochyra R, Van Rooy J (2013) Distribution of *Bucklandiella lamprocarpa* (Grimmiaceae, Musci) in South Africa. Cryptogamie, Bryologie 34(3): 359–366. doi: 10.7872/cryb.v34.iss3.2013.359
- Ochyra R, Zander RH (2007) Czy *Tortula lithophila* jest konspecyficzna z *Sarconeurum glaciale* (Bryopsida: Pottiaceae)? Fragmenta Floristica et Geobotanica Polonica 14: 209–212.
- Peralta DF, Yano O (2010) On the occurrence of *Notoligotrichum minimum* (Cardot) G. L. Smith and *Oligotrichum canaliculatum* (J. D. Hook. & Arnott) Mitt. (Polytrichaceae) in Brazil. Tropical Bryology 32: 103–107.
- Ramsay HP (2012) Australian Mosses Online. 1. Sematophyllaceae: *Rhaphidorrhynchium*. http://www.anbg.gov.au/abrs/Mosses_online/01_Sematophyllaceae/Semat_Rhaphidorrhynchium.pdf
- Rao P (2001) Taxonomic studies on *Cryphaea* (Cryphaeaceae, Bryopsida). 3. Revision of European, African, Australian and Oceanian, and American species. Bryobrothera 7: 37–111.
- Reimers H (1936) Über die Laubmoosgattungen *Bryopoteria* Thér., *Camptodontium* Dus., *Hymenoloma* Dus., *Verrucidens* Card. und ihre systematische Stellung. Notizblatt des Botanischen Gartens und Museums zu Berlin-Dahlem 13(116): 36–52. doi: 10.2307/3994913
- Robinson H (1975) The mosses of Juan Fernández Islands. Smithsonian Contributions to Botany 27: 1–88. http://www.sil.si.edu/smithsoniancontributions/Botany/pdf_hi/sctb-0027.pdf
- Sainsbury GOK (1955) A Handbook of the New Zealand mosses. Royal Society of New Zealand, Bulletin No. 5, Wellington, New Zealand, 1–490.
- Sastre-de Jesús I (1987) A revision of the Neckeraceae Schimp. and the Thamnobryaceae Marg. & Dur. in the Neotropics. PhD Thesis, City University of New York, USA.
- Schiavone MM (1993) Bryophyta Musci: Polytrichinales. Flora Criptogámica de Tierra del Fuego 14(12): 1–61.
- Seki T (1974 [1976]) A moss flora of Provincia de Aisén, Chile. Journal of Science of the Hiroshima University, Series B 15: 9–101.
- Seppelt RD (1978) Studies on the bryoflora of Macquarie Island II. *Ulota phyllantha* Brid. New Zealand Journal of Botany 16(1): 21–23. doi: 10.1080/0028825X.1978.10429654
- Seppelt RD (1982) A monographic revision of the genus *Ditrichum* (Musci: Ditrichaceae). I. Australian and New Zealand species. Journal of the Hattori Botanical Laboratory 51: 99–150.
- Skottsberg C (1916) Botanische Ergebnisse der Schwedischen Expedition nach Patagonien und dem Feuerlande 1907–1909. V. Die Vegetationsverhältnisse längs der Cordillera de los Andes S. von 41° S. Br. Ein Beitrag zur Kenntnis der Vegetation in Chiloé, West-Patagonien, dem Andinen Patagonien und Feuerland. Kungl. Svenska Vetenskapsakademiens Handlingar,

- n.s. 56(5): 1–366 + 23 plates. <http://www.biodiversitylibrary.org/item/130979#page/681/mode/1up>
- Smith AJE (2004) The Moss Flora of Britain and England, 2nd edition. Cambridge University Press, Cambridge, 1–1012. doi: 10.1017/CBO9780511541858.004
- Stech M, Frey W (2010) A morpho-molecular classification of the mosses (Bryophyta). *Nova Hedwigia* 86(1-2): 1–21. doi: 10.1127/0029-5035/2008/0086-0001
- Stech M, Pfeiffer T, Frey W (2008) Phylogenetic relationships within *Dendroligotrichum* and *Polytrichadelphus* (Polytrichaceae, Polytrichopsida) in the Palaeoaustral region. *Studies in austral temperate rainforest bryophytes 32. Australian Systematic Botany* 21: 39–49. doi: 10.1071/SB07005
- Stephani F (1911) Botanische Ergebnisse der schwedischen Expedition nach Patagonien und dem Feuerlande 1907–1909. II. Die Lebermoose. *Kungl. Svenska Vetenskapsakademiens Handlingar*, n.s. 46(9): 1–92.
- Streimann H, Curnow J (1989) Catalogue of the mosses of Australia and its external territories. *Australian Flora and Fauna Series number 10. Australian Government Publishing Service, Canberra*, 1–479.
- Suárez GM, Schiavone MM (2009) El género *Pilopogon* (Bryophyta, Dicranaceae) en la Argentina. *Darwiniana* 47(2): 309–314. <http://www.jstor.org/stable/23230190>
- Suárez GM, Schiavone MM (2011) *Pohlia* Hedw. section *Pohlia* (Bryaceae) in Central and South America. *Nova Hedwigia* 92(3-4): 453–477. doi: 10.1127/0029-5035/2011/0092-0453
- Tangney R, Huttunen S, Stech M, Quandt D (2010) Review of the systematic position of the pleurocarpous moss genus *Acrocladium* Mitten. *Tropical Bryology* 31: 164–170.
- Thériot I (1934 [1935]) Contribution à la flore bryologique du Chili (11^e article). *Revue Bryologique et Lichénologique* 7(3-4): 167–183.
- Touw A (1971) A taxonomic revision of the Hypnodendraceae (Musci). *Blumea* 19(2): 211–354.
- TROPICOS (2016) Tropicos.org. Missouri Botanical Garden. <http://www.tropicos.org/Name/35188759> [27 February 2016]
- Villagrán C, Barrera E, Cuvertino J, García N (2003) Musgos de la Isla Grande de Chiloé, X Región, Chile: Lista de especies y rasgos fitogeográficos. *Boletín del Museo Nacional de Historia Natural, Chile* 52: 17–44.
- Zander RH (1972) Revision of the genus *Leptodontium* (Musci) in the New World. *The Bryologist* 75(3): 213–280. doi: 10.2307/3241464
- Zander RH (1977) The Tribe Pleuroveisieae (Pottiaceae, Musci) in Middle America. *The Bryologist* 80(2): 233–269. doi: 10.2307/3242469
- Zander RH (1993) Genera of the Pottiaceae: mosses of harsh environments. *Bulletin of the Buffalo Society of Natural Sciences* 32, Buffalo, New York, 1–378.
- Zander RH (2007) *Didymodon*. In: Flora of North America Editorial Committee (Eds) *Flora of North America* Vol. 27, Bryophytes, Mosses, Part I. Oxford University Press, New York, 539–561.
- Zander RH, Toren D, Eckel PM (2007) *Gymnostomum aeruginosum*, *G. calcareum* and *G. viridulum* (Pottiaceae, Bryopsida) in California. *Journal of Bryology* 29: 27–32. doi: 10.1179/174328207X160568
- Zomlefer W (1993) A revision of *Rigodium* (Musci: Rigodiaceae). *The Bryologist* 96(1): 1–72. doi: 10.2307/3243321

Supplementary material I**Systematic list of taxa and additional information**

Authors: Juan Larraín

Data type: specimens data

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.