

Original Research Article

Proximate Analysis and Phytochemical Screening of *Triclisia subcordata* Oliv Leaf

Abstract

Proximate analysis and phytochemical screening of *Triclisia subcordata* Oliv leaf were investigated using standard analytical methods. Results of the proximate analysis showed contents for moisture ($79.28 \pm 2.32\%$), Ash ($1.91 \pm 0.21\%$), crude fibre ($1.83 \pm 0.13\%$), lipid ($1.81 \pm 0.16\%$), crude protein ($8.32 \pm 1.2\%$) and carbohydrate ($6.91 \pm 0.74\%$). Phytochemical screening gave tannins ($0.01 \pm 0.00\%$), saponins ($5.81 \pm 0.23\%$), alkaloids ($2.32 \pm 0.13\%$), flavonoids ($6.01 \pm 0.36\%$), cyanogenic glycosides (0.20 ± 0.00 mg/kg) and phenol ($0.03 \pm 0.00\%$). The study therefore concludes that *Triclisia subcordata* Oliv leaf has high contents for moisture, flavonoids and saponins and low lipid, cyanogenic glycosides, tannins and phenols.

Comment [emma chis1]: There should be a space between the standard deviation and %

Keywords: *Triclisia subcordata* oliv; Proximate profile; Phytochemicals; Traditional medicine.

Comment [emma chis2]: Use proximate analysis instead of proximate profile

Comment [emma chis3]: Traditional medicine is not mentioned in the abstract. please use words in the abstract, such as analytical methods

1.0 Introduction

Phytochemicals are secondary metabolites produced by plants. These products are biologically active, naturally occurring chemicals in various parts of a plant, providing health benefits for humans further than those attributed to macronutrients and micronutrients. These compounds have been linked to human health by contributing to protection against degenerative diseases.

Comment [emma chis4]: There should be a citation here

Triclisia subcordata Oliv (Menispermaceae) is a medicinal plant traditionally used for the treatment of various diseases in West Africa. The Menispermaceae is a temperate to tropical family of around 70 genera (including *Triclisia*) and 450 species of dicotyledenous small tropical flowering vines that are short in pedunculate clusters with twining stems, an appressed pubescent fruits and a few herbs, shrubs and trees. Leaves are reticulate, alternate and simple, but may be palmately veined and often lobed [1]. *T. subcordata* Oliv is commonly called

“Alugboran” in Yoruba language, “ogwu-aju” in Nsukka (which means antidizziness), “Ezize” in Ohafia (which means benign skin growth), “ike mbekwu” in Umuoji in Eastern part of Nigeria (which is gotten from the shape of the leaf) [2]. The stem is used as a rough fibre (tietie) [3].

Comment [emma chis5]: The highlighted part should come immediately after the first paragraph

It is a nature plant of west tropical Africa including Nigeria, Ghana, Ivory Coast, Sierra Leone, Senegal and Togo [4]. The plant frequently serves as rope used for tying purposes; the importance of the species however rest with their medicinal application including the use of root extract for the treatment of snake bite, ulcer, diarrhea, malaria pyorrhea, swelling of extremities, anemia joint pains, cancer, rheumatic pains and hypertension in Nigeria [2, 3].

Evidenced-based studies revealed that *Triclisia subcordata* has antiulcer, antihistamine, antimicrobial, anticancer, antioxidant and antidiabetic activities [5, 6, 7, 8]. The cytotoxicity Effects and Apoptosis Induction in ovarian and cancer cells by Bisbenzylisoquinoline and Isochondodendrine and 2'-norcocculine alkaloids from *Triclisia subcordata* was assayed by Uche *et al.* [9] [10].




Plate 1: photograph showing *Triclisia subcordata* Oliv leaf (arrowed)

Medicinally, several hypotheses suggest that a decoction of the root is drunk to treat fever and malaria. Also, root decoction is also taken as an emmenagogue and abortifacient (drugs or chemicals that induce abortion). The root pulp is rubbed in or root sap is rubbed into scarifications as a treatment for rheumatism, arthritis, anaemia and sleeping sickness. The leaf or a root juice is mixed with salt in palm wine and taken as treatment for coughs and bronchial disorders and as well function as sedative on the heart and to help wash palpitations. The leaf sap has a soothing effect on coughs. A decoction of the leaves and twigs is drunk, or leaf pulp is rubbed in, to treat oedema of the legs. Furthermore, it was hypothesized that a decoction of the leaves and stem are used as a nasal or ocular instillation and as a purgative or bathe against epilepsy and as treatment for stomach ache respectively. The stem bark is powdered and applied topically to syphilitic sores and leprosy while the bark pulp is used as a purgative. A methanolic leaf extract has shown significant anti-ulcer effects [11]. This paper is aimed at analyzing the proximate content and phytochemical screening of *Triclisia subcordata* Oliv leaf.

Comment [emma chis6]: taken

Comment [emma chis7]: The root pulp or root sap is rubbed into

Comment [emma chis8]: There must be a reason why you are analysing the proximate and photochemical constituents, possibly, due to it's acclaimed ethnomedicinal uses by African natives and it must reflect in the aim

2.0 Materials and Methods

2.1. Sample Collection

Fresh *Triclisia subcordata* Oliv leaf was obtained from University of Port Harcourt campus environment, Choba, Rivers State, Nigeria. The leaves were pulverized using a mechanical grinder to obtain a smooth mixture.

2.2. Experimental Design

After grinding of the leaf, approximately 500g was weighed using electronic weighing balance and used for proximate and phytochemical screening using standard analytical methods.

Comment [emma chis9]: 500 g

2.3. Proximate Analysis

Determination of Moisture content [12]

A dry clean petridish was placed in an oven at 80°C for about 30 minutes, cooled in a desiccator and weighed as (w). 5g of the samples was added to the petridish and weighed as (b). The petridish and its content were placed in an oven adjusted to 70°C. After 5 hours, the petridish containing the sample was removed and quickly transferred to a desiccator for cooling. The petridish was put back into the oven and adjusted to 105°C for another 5 hours after which it was removed, put in desiccators for cooling. This process was repeated and weighed until a constant weight (c) was obtained.

Comment [emma chis10]: 5 g

The % moisture content was determined as follows;

$$\% \text{ moisture content} = \frac{b-c}{b-w} \times 100$$

Where W = weight of moisture can, b = weight of petridish + sample, c= weight of petridish + sample after drying.

Determination of ash content [12]

An empty crucible was first ignited in a muffle furnace for 1minute and allowed to cool in desiccators containing silica gel. 5g of the sample was accurately weighed into the preheated dish. The weight of the porcelain dish and the samples were noted. Afterwards, the dish was heated with a Bunsen burner in a fume cupboard until smoking ceases and later transferred into a muffle furnace at 550-570°C for about 18-24hours to burn off all organic matter. After ashing, the crucible was removed from the furnace and placed in desiccator to cool at room temperature and weighed. The percentage ash content of the sample was calculated thus;

Comment [emma chis11]: 5 g

Comment [emma chis12]: 18-24 hours

$$\% \text{ Ash} = \frac{\text{weight of ash}}{\text{weight of sample}} \times 100$$

$$= \frac{W_3 - W_1}{W_2 - W_1} \times 100$$

Where; W_1 = weight of empty crucible, W_2 = weight of crucible + sample before ashing

W_3 = weight of crucible + sample after ashing

Determination of lipid [12]

A 5g of the sample was weighed into a thimble and was extracted with petroleum ether until it siphons using the Soxhlet extraction method. The lipid was exhaustively extracted using petroleum ether at 40 – 60°C for 6hrs. The sample in the thimble was removed and dried in air at 50°C for 5 minutes, cooled in a desiccator and weighed. The % lipid content was calculated as follows;

$$\begin{aligned}\% \text{ Lipid} &= \frac{\text{weight of sample (extracted fat)}}{\text{weight of sample}} \times 100 \\ &= \frac{W_2 - W_1}{W} \times 100\end{aligned}$$

Where; W_1 = weight of empty thimble, W_2 = weight of thimble + sample W = weight of sample

Determination of crude fiber [12]

A 2g of the defatted sample was weighed into conical flask and 200 mLs of 1.25% of boiling sulphuric acid was added within a minute. The content of the flask was filtered through a buchner funnel prepared with wet 12.5cm filter paper. The sample was washed back into the original flask with 200mLs of 1.25% NaOH, and boiled for 30minutes. All insoluble matter was transferred to the crucible and treated till the sample was free from acid. The sample was again washed in a muffle furnace at 550°C/hour. The crucible was then cooled in desiccator and reweighed.

$$\% \text{ Crude fiber} = \frac{W_2 - W_1}{W} \times 100$$

Where; W = weight of sample, W_1 = weight of crucible + sample

W_2 = weight of crucible+ filter paper after ashing.

Determination of Crude protein [12]

Comment [emma chis13]: 5 g not 5g

Comment [emma chis14]: 1.25 %
Effect the correction all through. There should be a space between the values and the unit, expect for degree Celsius

Comment [emma chis15]: 12.5 cm

A 1g of the sample was weighed and transferred into Khedahl flask. Few chips of antibumping granules, 4g of digestion catalyst and 20mLs of concentrated sulphuric acid were added at a 40°C angle with a retort stand on an electro thermal heater. The flask was gently heated for frothing to occur and subside, and then heat was increased to about 250°C. The digestion was carried out within 2-6 hours by which time the entire sample was digested completely. The digest was cooled to room temperature and diluted to 100mLs with distilled water. For distillation, 20mLs aliquot of the digest was transferred into a round bottomed flask. This flask was connected to a Liebig condenser through a monoarm steel head (Adaptor). The liebig condenser was connected to a receiver flask through a receiver adapter. 10mLs of 2% boric acid and two drops of double indicator were pipetted into the distillation flask. 30mLs of 40% sodium hydroxide was injected into the distillation flask through a cork with the aid of a syringe. The flask was heated for 10minutes to digest the content. The distillate was collected in the boric acid and then titrated with 0.1M HCL. The vol. of HCl added was recorded as the titre value.

The % Crude protein was calculated thus;

$$\% \text{ Crude protein} = \% \text{ Nitrogen} \times 6.25$$

$$\% \text{ Nitrogen} = \frac{\text{titre value} \times 1.4 \times 100 \times 10}{1000 \times \text{wt of sample} \times \text{aliquot digest}}$$

Where, 1.4 = N₂ equivalent to 0.1NHCl used in titration

100 = Total volume of digest

Determination of total carbohydrate [12]

The total carbohydrate content of the sample was estimated as the Nitrogen free extract (NFE).

The arithmetic different methods involve adding the total percentage value of crude volume.

$$\text{Total CHO} = 100 - (\% \text{fibre} + \% \text{protein} + \% \text{Moisture} + \% \text{ash} + \% \text{fats})$$

2.4. Phytochemical Screening

Preparation of the leaf extracts

Ten grams (10g) of the leaf sample was soaked in 100mL of water in a beaker and left for about 8hours. The solution obtained was filtered using filter paper, and the filtrate used for phytochemical screening.

Quantitative determination of phytochemicals

Determination of Saponins [12]

A 10g of the ground sample was measured into a conical flash and 100mL of 20% ethanol added to it. The suspension was heated over hot bath at 55°C for 12hourrs with continuous stirring using a magnetic stirrer. The mixture was filtered and the residue re-extracted with another 200mL of 2% aqueous ethanol. The combined extract was reduced to 40mL of the original size over a water bath at about 55°C. The purification process was repeated two more times. A 4g of Sodium chloride (NaCl) was added to adjust the pH meter. The solution was shaken with 60mL and 30mL portions of n-butanol extract and later washed twice with 10mL of aqueous NaCl. The remaining solution was evaporated to dryness in water bath. After evaporation, the sample was dried in the oven to a constant weight.

Comment [emma chis16]: 12 hours

The saponin content was calculated in g/100g as;

$$\% \text{ saponins} = \frac{\text{weight of residue}}{\text{weight of sample}} \times 100$$

$$\% \text{ saponins} = \frac{W_2 - W_1}{W} \times 100$$

Where; W = weight of sample, W₁ = weight of evaporating dish

W₂ = weight of evaporating dish + saponin content after drying

Determination of Alkaloids [12]

A 2g of freshly crushed sample was weighed and dispensed into 100 mL of 10 % acetic acid. The mixture was shaken and allowed to stand for 4 hours before filtration. The mixture was filtered to remove all debris and evaporated to ¼ of the original volume. A 1% concentrated ammonium hydroxide (NH₄OH conc) was added drop wise to precipitate the alkaloids. It was filtered with well weighed ppt. in the filter paper, oven dried at 60°C for 30 minutes and then reweighed.

The alkaloid content was calculated as;

$$\% \text{ Alkaloid} = \frac{\text{weight of residue}}{\text{weight of sample}} \times 100$$

$$\% \text{ Alkaloid} = \frac{W_2 - W_1}{W} \times 100$$

Where; W = weight of sample, W₁ = weight of empty filter paper

W₂ = weight of filter paper of ppt.

Determination of Flavonoids [13]

The method used for determination of flavonoids was that of Bohn and Kocipai (1994). 5g of the blended sample was extracted repeatedly and separated with 50mL of 40% aqueous methanol at room temperature. The solution was shaken for homogeneity and left to stand for about 4hours and later filtered into a weighed beaker. The filtrate was later transferred into a crucible and evaporated to dryness over a water bath, then dried in an electric oven to a constant weight.

Comment [emma chis17]: 4 hours

The flavonoid content was expressed in percentage as follows;

$$\% \text{ flavonoid} = \frac{\text{weight of residue}}{\text{weigh tof sample}} \times 100$$

Determination of cyanogenic glycoside [12]

One gram (1g) of sample was weighed into a 250mL round bottomed flask. 200mL distilled water was added and allowed to stand for 2hours (for autolysis to occur). Full distillation was then carried out and 150-170mLs of distillate was collected in a 250mL conical flask containing

2mLs of 2.5% NaOH. Anti-foaming agent (tannic acid) was added before distillation. To 100mL of the distillate containing cyanogenic glycoside, 8mL of 6N NH₄OH and 2mLs of 5% potassium iodide (KI) was added, mixed and titrated with 0.02M silver nitrate (AgNO₃) using a micro burette against a black background. Permanent turbidity indicated end point.

Cyanogenic glycoside in the sample was calculated thus;

$$\text{cyanogenic glycosides (mg/kg)} = \frac{\text{Titre value (mL)} \times 1.0\text{g} \times \text{extract vol. (mL)} \times 100}{\text{Aliquot vol. (mL)} \times \text{weight of sample (g)}}$$

Determination of tannin [12]

A 0.5g of the sample was weighed into 100mL plastic bottle. 50mLs of distilled water was added and shaken for 1hour in a mechanical shaker. Then 5mLs of the filtrate was pipette out into a tube and mixed with 3mLs of 0.1M Iron (III) chloride (FeCl₃) in 0.1N hydrochloric acid (HCl) and 0.008M potassium ferrocyanide K₄[Fe(CN)₆]. The absorbance was measured in spectrophotometer at 720 nm wavelength within 10 minutes. A blank sample was prepared, and the colour also developed and read at same wavelength. A standard was prepared using tannic acid to get 100 ppm and measured using the formula below.

$$\text{Tannin} = \frac{A_n}{A_s} \times C \times 100 / W \times v_f / v_g$$

Comment [emma chis18]: Use equation editor

Where; A_n = Absorbance of test sample, A_s = Absorbance of standard solution, C = Concentration of standard solution, W = Weight of sample used, V_f = Total volume of extract, V_g = volume of extract analyzed.

Determination of phenol [12]

A 0.1g of sample was weighed into a 250mL round bottomed flask. 100 ml distilled water was added, boiled for 30 minutes and the volume made up to 100mL. A 2.5mL aliquot is measured into a conical flask and 5mL of 0.1M NaOH added into it. The solution was heated to 50°C and

allowed to cool, then 2.5mL of 0.005 iodine added. The flask was cork with a foil and allowed to stand at room temperature for 2hourrs. 0.5ml of concentrated HCl was added and titrated to pale yellow with 0.1M Na₂S₂O₃ with the addition of 3 drops of starch indicator. The solution is titrated again with 0.1M Na₂S₂O₃. A blank is formed using 2.5mL distilled water and absorbances read off.

2.5. Method of Data Analysis

All data collected were subjected to Students t-test for statistical analysis. All data were represented in mean \pm standard deviation (M \pm S.D). Level of significance was determined at a confidence level of determination ($p \leq 0.05$).

3.0 Results

Table 1: Proximate analysis of *Triclisia subcordata* Oliv leaf

Proximate content	<i>Triclisia subcordata</i> (%)
Moisture	79.28 \pm 2.32
Ash	1.91 \pm 0.21
Lipid	1.81 \pm 0.16
Crude protein	8.32 \pm 1.2
Crude fibre	1.83 \pm 0.13
Carbohydrate	6.91 \pm 0.74

Comment [emma chis19]: space

Comment [emma chis20]: SD not in uniform with others in the table

Values are Mean \pm SD of triplicate determinations. n=3.

Table 2: Phytochemical screening of *Triclisia subcordata* Oliv leaf

Phytochemical content	<i>Triclisia subcordata</i> Oliv leaf
Tannins (%)	0.01±0.00
Saponins (%)	5.81±0.23
Alkaloids (%)	2.32±0.13
Flavonoids (%)	6.01±0.36
HCN (mg/kg)	0.20±0.00
Phenol (%)	0.03±0.00

Values are Mean ± SD of triplicate determinations. n=3.

4.0 Discussion

The proximate content of *Triclisia subcordata* oliv leaf showed appreciable amount of the parameters as expressed in Table 1. From this study, the moisture content was very high (79.28%) when compared with findings of [14] which imply that this leaf cannot be stored for long and is prone to microbial attack.

Comment [emma chis21]: Recast

Ash contents express the mineral index contents in biological mass. This study revealed relatively lower ash in *Triclisia subcordata* oliv leaf (1.91%) than in fluted pumpkin leaf

Comment [emma chis22]: 1.91 %
Correct it throughout the manuscript

(9.68%), bitter leaf, (15.86%) and *Moringa oleifera* (15.09%) leaves [15]. This implies that *Triclisia subcordata* oliv leaves are not good source of mineral elements (Table 1).

Lipid in food absorbs and retains the flavour thereby enhancing the palatability [16]. The lipid content recorded in this study (1.81%) is far lower than values obtained by [14] which implies consumption of *T. subcordata* will not enhance palatability.

Proteins are necessary for the build-up of tissues and substances like hormones and enzymes. The values of protein obtained in this study (8.32%) are low when compared with [14], 24% in *Amaranthus vividis* [17], 20.72% in *Moringa oleifera* [18], 21.0% in *Lasianthera africana* and 15.0% in *Heinsia crinata* [15, 19] which makes it insufficient source of protein.

The crude fibre contents of *Triclisia subcordata* oliv leaf (1.83%) is very low when compared with those of *Lasianthera africana* (15.3 – 18.1% dry mass) [20] and *Heinsia crinata* (13 – 15% dry mass) [19]. Although intake of dietary fibre can lower the serum cholesterol level, risk of coronary heart disease, hypertension, diabetes, obesity, constipation and cardiovascular diseases, colon and breast cancer [21, 22]. The major problem associated with nutrition of vegetables by human is the high fibre content which can cause intestinal irritation and lower nutrient bioavailability [15, 23]. Intake of fibre can stimulate peristaltic movement, weakening of hunger, increase stool bulk and reduce serum levels of cholesterol [24, 25].

The carbohydrate content recorded was 6.91. Carbohydrates are pivotal nutrients needed for a balanced diet [26]. Generally, the low contents of protein, lipid and carbohydrate in this leaf imply that it is not a good source of energy.

Table 2 revealed the phytochemical screening of *Triclisia subcordata* oliv leaf. The study showed substantial amounts of flavonoids, saponins and alkaloids and small amounts of phenol,

Comment [emma chis23]: space

tannins and cyanogenic glycosides. When compared with the study of [27], tannins, saponins, alkaloids and phenols were lower while flavonoid content was high.

Tannins are complex moiety with wide pharmacological activities and are produced by majority of plants as protective substance. Tannin has astringent property, hastens the healing of wounds and inflamed mucous membrane and has been used since past as tanning agents. Tannin has received considerable attention in the fields of nutrition, health and medicine, largely due to their physiological activity, such as antioxidant, antimicrobial and anti-inflammatory properties [28, 29].

Saponins protect against hyperglycaemia, hypercholesterolaemia, hypertension [30], have antibiotic properties and antiinflammatory property and aid healing [31].

Alkaloids have been reported to be powerful pain relievers, exert an anti-pyretic, antihypertensive, antifungal, anti-inflammatory, anti-fibrogenic effect [32], stimulating, anaesthetic action [33] and inhibiting activity against most bacteria [34].

The flavonoids show antioxidant activity and have strong anti-cancer activity [35]. Flavonoids exhibit their antioxidative properties through several mechanisms, such as scavenging of free radicals, chelation of metal ions, such as iron and copper, inhibition of hydrolytic and oxidative enzymes inhibit antibiotics resistant microbes and also act as anti-inflammatory agent [36].

Conclusion

It can be concluded that *Triclisia subcordata* oliv leaf has high moisture, flavonoids and saponins with low lipid, cyanogenic glycosides, tannins and phenols. The proximate analysis does not support their nutritional relevance as they are seen in smaller quantity when compared to other plants (*Telfairia occidentalis* etc) but the phytochemical screening backs their usage in

traditional medicine research. Again the proximate composition could provide the micronutrients needed but not the macronutrients.

References

- [1] Ezimah, A.C., Ajugwo, A.O., Osuala, F.N. & Aghatise, K. (2013). Comparative Studies of EDTA, Sodium Citrate and Aqueous Extract of *Triclisia Dictyophylla*. *Scholars Academic Journal of Biosciences*, 1(2), 55-58.
- [2] Dalziel, J.M. (1937). The Useful Plants of West Tropical Africa, The Crown Agents for the Colonies, London. PP. xii + 612pp
- [3] Irvine, F.R. (1961). Woody plants of Ghana, Oxford University Press, London. Pp. xcv + 868
- [4] Trease, G. & Evans S.M. (1993). Medicinal Plants and Traditional Medicines in Africa, 2nd Ed., Spectrum Books. Pp. 35 – 53.
- [5] Asuzu, I.U. & Anaga, A.O. (1995). The anti-ulcer effect of methanolic extract of *Triclisia subcordata* O. leaves in rats. *Journal of Herbs, Spices and Medicinal Plants*, 3(3), 45-53.
- [6] Abo, K.A., Lawal, I.O. & Ogunkanmi, A. (2011). Evaluation of extracts of *Triclisia subcordata* Oliv and *Heinsia crinita* (Afz) G. Taylor for antimicrobial activity against some clinical bacterial isolates and fungi. *African Journal of Pharmacy and Pharmacology*, 5(2), 125-131.
- [7] Ayoola, M.D., Akinwunmi, K.F. & Agboola, O.B. (2017). Antidiabetic and antioxidant activities of *Entandrophragma cylindricum* and *Triclisia subcordata*. *Nigerian Journal of Natural Products and Medicine*, 21(1), 24-31.
- [8] Li, W. (2016). Cytotoxicity effects and apoptosis induction by cycleanine and tetrandrine. *Planta Medica: Natural Products and Medicinal Plant Research*, 30(9), 1533-1539.
- [9] Uche, F.I., Drijfhout, F.P., McCullagh, J., Richardson, A. & Li, W.W. (2016). Cytotoxicity effects and apoptosis induction by bisbenzylisoquinoline alkaloids from *Triclisia subcordata*. *Phytotherapy Research*, 30(9), 1533-1539.
- [10] Uche, F.I., Abed, M.N., Abdullah, M.I., Drijfhout, F.P., McCullagh, J., Claridge, T.W., Richardson, A. & Li, W. (2017). Isochondodendrine and 2'-norcocsuline: additional

alkaloids from *Triclisia subcordata* induce cytotoxicity and apoptosis in ovarian cancer cell lines. *Royal Society of Chemistry Advances*, **7**(70), 44154-44161.

- [11] Tropical Plants Database, Ken Fern. tropical.theferns.info.
/tropical.theferns.info/viewtropical.php?id=Triclisia+subcordata/ Retrieved 2021-05-01.
(<http://www.prota.org>)
- [12] AOAC. (1990). Association of Official Analytical Chemists. Method of analysis
Washington DC, U.S., 15, 1250-1255.
- [13] Bohn, C. & Kocipai, G. (1994). Method of analysis. Washington DC, U.S., 8, 1233-1246.
- [14] Akwukwaegbu, P.I., Peters, D.E., & Wegwu, M.O. (2016). Proximate analysis and
phytochemical screening of fluted pumpkin (*Telfairia occidentalis*) pod. *American
Journal of Food, Nutrition and Health*, **1**(1), 1-6.
- [15] Effiong, G. S., Ogban, P. I., Ibia, T. O., & Adam, A. A. (2009). Evaluation of nutrient
supplying potentials of Fluted Pumpkin (*Telfairia occidentalis*, Hook, F.) and Okra
(*Abelmoschus esculentus*) (L.) Moench. *Academic Journal of Plant Sciences*, **2**(3), 209–
214.
- [16] Fagbohun, E.D., Egbebi, A.O. & Lawal, O.U. (2012). Phytochemical screening, proximate
analysis and in-vitro antimicrobial activities of methanolic extract of *Cnidoscolus
aconitifolius* leaves. *International Journal of Pharmaceutical Science Review and
Research*, **13**, 28-33.
- [17] Sena, L. P., vander Jagt, D. J., Rivera, C., Tsin, A. T. C., Muhammadu, I., Muhamadou, O.,
Milson, M., Pastosyn, A. & Glew, R. H. (1998). Analysis of Nutritional components of
eight famine foods of the republic of Niger. *Plant Foods for Human Nutrition*, **52**, 17-30.
- [18] Lockeett, C. T., Calvert, C. C. & Grivetti, L. E. (2000). Energy and micronutrient
composition of dietary and medicinal wild plants consumed during drought. Study of
rural Fulani, Northeastern Nigeria. *International Journal of Food Science and Nutrition*,
51, 195- 208.

- [19] Udosen, E. O., Udok, U. E. & Unuigue, O. S. (1998). The comparison of the Nutrient composition of *Lasianthera africana* and *Heinsia crinata*. *Journal of Food Biochemistry*, **23**, 571-576.
- [20] Isong, E. U. & Idiong, U. I. (1997). Comparative studies on the nutritional and toxic composition of three varieties of *Lesianthera africana*. *Plant Foods for Human Nutrition*, **51**:79-84.
- [21] Aberoumand, A. & Deokule, S.S. (2009). Studies on nutritional values of some wild edible plants from Iran and India. *Pakistan Journal of Nutrition*, **8**, 26-31.
- [22] Chukwu, E.C., Osuocha, K.U. & Uhegbu, F.O. (2018). Nutrient Composition and Selected Biochemical Effects of *Cnidoscopus aconitifolius* Leaf Extracts in Male Albino Rats. *Journal of Forensic Research*, **9**(1), 409.
- [23] Iwuji, S.C., Nwafor, A., Ekwurugwu, J., Ejeta, K. & Akpan, U. (2013). Comparative characterization of the phytomedicinal constituents of *Cnidoscopus aconitifolius* leaf extracts. *American Journal of PharmTech Research*, **3**, 780-784.
- [24] Umar, K.J., Hassan, L.G., Dangoggo, S.M. & Ladan, M.J. (2007). Nutritional composition of water spinach (*Ipomea aquatica* Forsk) leaves. *Journal of Applied Science*, **7**, 803-809.
- [25] Ramula, P. & Rao, P.U. (2003). Dietary fibre content of fruits and leafy vegetables. *Nutrition News*, **24**, (3) 1-6.
- [26] Emebu, P.K. & Anyika, J.U. (2011). Proximate and mineral composition of Kale (*Brassica oleracea*) grown in Delta State, Nigeria. *Pakistan Journal of Nutrition*, **10**, 190-194.
- [27] Okpara, F.N. & Akwukwaegbu, P.I. (2020). Effect of aqueous leaf extract of *Cnidoscopus aconitifolius* on lipid profile and haematology of carbon tetrachloride treated rats. *Acta Chemica Iasi*, **28**(2), 237-256.doi: 10.2478/achi-2020-0017.
- [28] Asquith, T.N. & Butter, L.G. (1986). Interaction of condensed tannin with selected proteins. *Phytochemistry*, **25**(7), 1591-1593.

- [29] Killedar, S.G. & More, H.N. (2010). Estimation of tannins in different parts of *Memecylon umbellatum* Burm. *Journal of Pharmacy Research*, **3**(3), 554-556.
- [30] Trease, G.E. & Evans, W.C. (1985). *Pharmacognosy*, 14th edition, London: W.B. Sanders company. Pp 23-29.
- [31] Krishnaiah, D., Devi, T., Bono, A. & Sarbatly, R. (2009). Studies on phytochemical constituents of six alaysian medical plants. *Journal of Medicinal Plants Research*, **3**(2), 67-72.
- [32] Awoyinka, O.A., Balogun, I.O. & Ogunnowo, A.A. (2007). Phytochemical screening and in-vitro bioactivity of *Cnidioscolus aconitifolius* (Euphorbiaceae). *Journal of Medicinal Plants Research*, **1**(3), 63-65.
- [33] Edeoga, H.O. & Enata, D.O. (2001). Alkaloids, tannins and saponin content of some medicinal plants. *Journal of Medicinal Aromatic Plant Science*, **23**, 344-349.
- [34] Al-Bayati, F.A. & Sulaiman, K.D. (2008). In-vitro antimicrobial activity of *Salvadora persica* L. extracts against some isolated oral pathogens in Iraq. *Turkish Journal of Biology*, **32**(1), 57-62.
- [35] Salah, N., Miller, N.J., Paganga, G., Tijburg, L., Bolwell, G.P., Rice, E. & Evans, C. (1995). Polyphenolic flavonols as scavenger of aqueous phase radicals and chain breaking antioxidants. *Archives of Biochemistry and Biophysics*, **322**(2), 339-346.
- [36] Kessler, R.C., Berglund, P., Demler, O., Jin, R. & Koretz, D. (2003). The epidemiology of major depressive disorder: results from the National

Comorbidity Survey Replication (NCS-R). *Journal of American Medical Association*, **289**, 3095-3105.

UNDER PEER REVIEW