

## Original Research Article

### Acute and Sub-acute Toxicity Assessment of *Euphorbia lateriflora* (Schum and Thonn) in Wistar Albino Rats

#### Abstract

**Aims:** The toxicity of ethanol whole plant extract of *Euphorbia lateriflora* was assessed in Albino Wistar rats.

**Methodology:** The LD<sub>50</sub> was at single 5000 mg/kg body weight, the sub-acute dosage of the extract was administered orally for 7 days at 250 and 500 mg/kg b.w.t per day and the effect of the extract on liver, kidney, and haematological parameters were assessed and recorded during these periods

**Results:** The result of the oral acute toxicity study at single high dose of 5000 mg/kg/bwt shows that the LD<sub>50</sub> of the extract is greater than 5000 mg/kg/bwt. After 28days of oral administration, 500mg/kg/bwt of the extract caused a significant ( $p < 0.05$ ) decrease in the packed cell volume. At 500 mg/kg/bwt, the extract caused a significant ( $p < 0.05$ ) increased in ALP, total protein, bilirubin and albumin and decrease in GGT, serum electrolytes ( $\text{Na}^+$ ,  $\text{K}^+$  and  $\text{Cl}^-$ ). Histopathological analysis revealed the expansion of partial spaces by fibrous spaces in the liver and thickening of the glomerular basement of the kidney in the group fed with 500mg/kg/b.w.t of extracts

**Conclusion:** In conclusion, the dose and time-dependent selective organ toxicity effect of this extract suggested that the extract might be relatively unsafe for consumption at especially at high concentrations.

**Keywords:** Toxicity; Kidney; Liver; Haematology; LD<sub>50</sub>.

#### Introduction

Medicinal plants had a long dated history in the management of several human diseases. They play important role in the development of potent therapeutic agents and have contributed significantly toward the development of modern medicine [1]. The therapeutic properties attributed to medicinal plants are linked to phytochemicals such as terpenoids, saponins, phenolic compounds and alkaloids [2, 3]. Several studies have shown that the bioactivities of medicinal

plants include anti-malaria, anti-microbial, anti-inflammatory, antioxidant, and anti-diabetes properties [3, 13, 29].

The major advantages of medicinal plants over the conventional drugs seem to be their perceived efficacy, low cost and the low incidence of serious adverse effects [4, 28]. This claim has however, been challenged as some medicinal plants have demonstrated some toxicological effects in both human and animals [5]. Therefore, it should be stressed that before the use of traditional medicinal plants, the safety should be ascertained [6]. Furthermore, it is necessary for medicinal plants to be evaluated for safety or toxicity and necessary recommendations be made regarding their use [7].

*Euphorbia lateriflora* (Schum and Thonn) also known in Yoruba as “Enu opiri” and “Fidda sartse” in Hausa. It is a shrub with smooth- gracious and erect branches [8]. The latex of the plant is used for the treatment of ringworm, and in a dilute aqueous solution as purgative [8, 9]. The leaves are also useful in the treatment of dermatoses [8, 9]. Several studies have revealed the pharmacological potential of this plant. According to Obi *et al.* [10], *E. Lateriflora* extract exhibited antiviral activity in human carcinoma cell line. However, for proper recommendation, and public awareness on the safety of the plant extracts, the acute oral toxicity effects of the ethanol extracts of the whole plant on Albino rats was investigated.

## **2.0 Material and methods**

### **2.1 Plants collection**

Whole plant of *E. lateriflora* was obtained within Ogbomosho North LGA, Oyo State Nigeria through the help of a traditional healer. Ogbomoso North lies between latitude 8.156697 and longitude 4.264409. The plant was identified and authenticated by Prof. A.T.J Ogunkunle of the

Department of Pure and Applied Biology, Ladoke Akintola University of Technology, Oyo State Nigeria. The plant sample was deposited in the University Herbarium.

## **2.2 Preparation of plant extract**

The ethanol extraction of the plant sample was carried out by using modified method of Mbaka *et al.* [11]. Briefly, 400g of the pulverized plant sample was loaded in a soxhlet extractor in batches for 5h each and subjected to extraction with ethanol. After extraction, mixtures were concentrated at 48 °C using rotary evaporator and the concentrate was kept in a refrigerator (4 °C) until use.

## **2.3 Experimental animals**

Thirty male albino rats (Wistar stock) weighing between 150-200 g were obtained from the Department of Anatomy Animal House, Ladoke Akintola University of Technology Ogbomosho Oyo State, Nigeria. They were acclimatized for a period of two weeks and kept under the normal 12 h light/dark cycle. The animals were allowed access to food and water *ad libitum* throughout the study period. The animal feed was specially prepared from chick Grower's mash (Feed Mill Company Ogbomosho Oyo State, Nigeria). Animals were handled following the standard protocol on animal care and use.

## **2.4 Experimental design**

### **2.4.1 Determination of LD<sub>50</sub>**

Ten (10) rats were used for this study. After 7 days acclimatization, the rats were fasted over night but allowed access to water prior to dosing. The up and down method by Dixon, [12] was used. The method involves testing the rats, one at a time in a staircase method i.e. limit test

which involves either starting with the lowest dose (2000mg/kg/bwt) and increasing the dosage if no death or toxic effect is noticed or starting with the highest dose (5000mg/kg/bwt) and reducing the dosage if death occurs at the highest dose. The animals were weighed and dosed with a limit dose of 5000mg/kg body weight of the freshly prepared ethanol whole plant extract of *Euphorbia lateriflora*. The rats were monitored for 72 hours for any change in behaviour.

#### **2.4.2 Sub-Acute toxicity test**

The sub-acute toxicity study was performed for the ethanol whole plant extracts of *Euphorbia lateriflora*. Rats were randomly separated into 3 groups (5 rats per group) and were orally administered with 1/10 and 1/20 LD<sub>50</sub> respectively for *Euphorbia lateriflora*. At the end of 7 days of twice administration daily, all surviving rats were sacrificed by decapitation and the internal organs examined.

- Group A= control rats given 1ml distilled water (no ethanolic plant extract)
- Group B= rats given 1/10 of the LD<sub>50</sub> of ethanolic plant extract. (500mg/kg)
- Group C= rats given 1/20 of the LD<sub>50</sub> of ethanol whole plant extract (250mg/kg).

#### **2.5. Collection of Blood samples**

Animals were anaesthetized with chloroform and decapitated. Blood samples were then collected into heparinized and non heparinized tubes. Those in heparinized tubes was used for haematological evaluation while the non heparinized were centrifuged and serum was analysed for biochemical parameters.

#### **2.6 Haematological and serum biochemical Parameters**

Haematological parameters assayed included red blood cell (RBC) and white blood cell (WBC) counts platelets, haematocrit, and haemoglobin (Hb) estimation. Erythrocyte indices (mean corpuscular volume (MCV), mean corpuscular haemoglobin concentration (MCHC), and mean

corpuscular haemoglobin (MCH)) was determined from values obtained from red blood cell (RBC) count, Hb concentration, and packed cell volume (PCV) values. All haematological parameters were analysed using the automated method with automatic analyser “Haematology Auto-analyser sysmex KX-21N”. The biochemical analysis evaluated includes the serum activity of aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), gamma glutamyl transferase, creatinine, urea, and total protein level. Electrolytes such as sodium, chloride and Potassium were also assayed.

## **2.7 Histopathology**

The histological procedures were done according to the method of Avwioro, [14]. The kidney and liver were removed and fixed into 10% buffered formalin in labeled bottles. Dewax in Xylene for 15 mins, Taken through Absolute Alcohol, 95 % and 70 % Alcohol. Rinsed the section in water, Stain in Harris haematoxylin for 5 mins, Rinse in water, Differentiate in 1% acid alcohol briefly, Rinse in water, Blue under running tap water for 10 mins, Counterstain with 1% aqueous Eosin for 2 min, Rinse in water, Dehydrate in ascending grades of alcohol and Clear in xylene and Mount in DPX.

## **2.8 Statistical analysis**

All data were calculated and expressed as mean  $\pm$  SEM using Graph Pad Prism Software and were considered significant at ( $p < 0.05$ ).

## **3.0 Results**

### **3.1 LD<sub>50</sub> Determination**

Animals treated with 5000mg/kg/bwt of the ethanolic extract of *E. lateriflora* showed no changes in behaviour and no apparent toxicity symptoms after 72hrs post treatment. The rats survived the

highest dose, hence, it was concluded that the LD<sub>50</sub> of the plants extracts is greater than 5000mg/kg/bwt.

### 3.2 Sub-Acute toxicity tests

#### Effect of whole plant extract of *E. lateriflora* on haematological parameters of rats

Table 1 showed the effects of the administration of *Euphorbia lateriflora* on the haematological parameters of rats. Administration of *Euphorbia lateriflora* whole plant extract at 250 mg/kg/b.w.t showed no significant (p<0.05) effect on the WBC, RBC, HGB, Platelet, Neutrophil and Lymphocyte but significantly (p<0.05) reduced the packed cell volume (HCT) when 500 mg/kg/b.w.t of the extract was administered in rat.

**Table 1: Toxic effects of *Euphorbia lateriflora* administration on Haematological indices**

Haematological	Control	<i>Euphorbia lateriflora</i> (mg/kg)	
Parameters		250	500
WBC 10 <sup>3</sup> /NL	10.2±0.45	10.5±0.90	11.3±1.56
NEUTROPHILS %	25±4.38	27.4±8.10	35.8±6.28
LYMPHOCYTE %	74±6.62	70.5±8.84	64.0±9.37
RBC ×10 <sup>6</sup> /NL	0.09±0.01	0.09±0.05	0.12±0.10
HGB g/dl	13.0±0.37	13.1±0.51	13.8±0.78
HCT %	39.4±1.15	38.3±0.95	35.4±0.54**
MCV	95.0±0.48	94.2±0.74	90.0±0.57
MCHC g/dl	450.7±87.62	445.1±69.30	440.0±57.16
PLATELET × 10 <sup>3</sup> /NL	783±51.71	574±76.39	680 ±120.83
PCT L/L	0.60±0.19	0.57±0.14	0.59±0.30

Values were expressed as mean ± SEM and considered significant at p-value <0.05. \*significant difference when compared with the Control group. #significant difference when the 250mg group is compared with 500mg dosage group.

### 3.3 Effect of *E. lateriflora* on biochemical parameters of rats

The results presented in Table 2 showed the effects of *Euphorbia lateriflora* administration at 250mg/kg/bwt and 500mg/kg/bwt on serum liver markers (ALT, AST, ALP and GGT). The administration of *Euphorbia lateriflora* had no significant ( $p<0.05$ ) effects on the serum activity of ALT and AST at both doses but a significant ( $p<0.05$ ) increase was observed in serum ALP activity when the extracts treated groups were compared with the control. A significant ( $p<0.05$ ) decrease was observed in serum GGT activity in the 250 and 500mg/kg/bwt treated groups when compared with the control. No significant ( $p<0.05$ ) difference was observed in serum GGT activity when the 250mg/kg/bwt dosage group was compared with 500mg/kg/bwt dosage group.

**Table 2: Serum Liver markers in *Euphorbia lateriflora* and *Terminalia Ivorensis* toxicity**

Parameters	Control	<i>Euphorbia lateriflora</i> (mg/kg)	
		250	500
ALT(U/L)	7.74±0.22	6.9±0.27	7.16±0.15
AST(U/L)	6.86±0.23	7.2±0.2	6.2±0.52
ALP(U/L)	36.28±0.29	42.6±0.29***	51.48±0.27***#
GGT(U/L)	40.57±0.25	39.1±0.12***	39±0.13***

Values were expressed as mean ± SEM and considered significant at P value <0.05. \*significant difference when compared with the Control group. #significant difference when the 250mg group is compared with 500mg dosage group. n=5

### 3.4 Effect of *Euphorbia lateriflora* on Electrolyte concentration

In table 3, treatment of rats with 250mg/kg/bwt of *Euphorbia lateriflora* had no significant ( $p<0.05$ ) effects on the serum sodium, potassium and calcium concentration when compared with

the control, but treatment with 500mg/kg/bwt significantly ( $p<0.05$ ) decreased the serum sodium and calcium concentration. No significant ( $p<0.05$ ) difference was observed between the serum potassium concentration of the control rats and rats treated with 500mg/kg/bwt of *Euphorbia lateriflora* while a significant ( $p<0.05$ ) decrease was observed in the serum sodium concentration of rats treated with 500mg/kg/bwt of *Euphorbia lateriflora* when compared with the group treated with 250mg/kg/bwt. In addition, *Euphorbia lateriflora* at 250mg/kg/bwt and 500mg/kg/bwt showed no significant effects on the serum urea and creatinine concentration.

**Table 3: Electrolytes concentration in *Euphorbia lateriflora* and *Terminalia Ivorensis* toxicity**

Parameters	Control	<i>Euphorbia lateriflora</i> (mg/kg)	
		250	500
<b>Sodium(<math>N_a^+</math>)</b>	166.95 $\pm$ 0.03	166.08 $\pm$ 0.21	160.82 $\pm$ 3.67**#
<b>Potassium(<math>k^+</math>)</b>	4.61 $\pm$ 0.05	3.33 $\pm$ 0.18	2.08 $\pm$ 0.12
<b>Chloride(<math>Cl^-</math>)</b>	51.44 $\pm$ 0.19	48.8 $\pm$ 0.36	45.34 $\pm$ 0.14*
<b>Urea (mg/dl)</b>	56.3 $\pm$ 0.19	56.18 $\pm$ 0.21	57.86 $\pm$ 0.11
<b>Creatinine (mg/dl)</b>	4.9 $\pm$ 0.13	4.1 $\pm$ 0.15	3.22 $\pm$ 0.12

Values were expressed as mean  $\pm$  SEM and considered significant at P value  $<0.05$ . \*significant difference when compared with the Control group. #significant difference when the 250mg group is compared with 500mg dosage group. n=5

The effects of *Euphorbia lateriflora* on the serum total protein, bilirubin and albumin level are showed in table 4. Significant ( $p<0.05$ ) increase was observed in the serum total protein and albumin level of rats treated with 500mg/kg/bwt of *Euphorbia lateriflora* when compared with the group treated with 250mg/kg/bwt and the control group. No significant ( $p<0.05$ ) difference




was observed in the total protein level of the control rats and the rats treated with 250mg/kg/bwt of *Euphorbia lateriflora*, while a significant ( $p<0.05$ ) increase was observed in the serum albumin level of rats treated with 250mg of *Euphorbia lateriflora* when compared with rats in the control group. No significant ( $p<0.05$ ) difference was observed in the serum bilirubin level of rats treated with *Euphorbia lateriflora* when compared with the control rats.

**Table 4: Effect of *Euphorbia lateriflora* on Protein concentration**

Parameters	Control	<i>Euphorbia lateriflora</i> (mg/kg)	
		250	500
<b>Total protein (g/dl)</b>	68.68±0.32	69.88±1.08	77.56±1.29***#
<b>Total bilirubin (mg/dl)</b>	2.86±0.10	3.16±0.09	3.84±0.13
<b>Albumin (g/dl)</b>	39.86±0.10	46.70±0.27***	50.32±0.22***#


Values were expressed as mean  $\pm$  SEM and considered significant at P value  $<0.05$ . \*significant difference when compared with the Control group. #significant difference when the 250mg group is compared with 500mg dosage group. n=5

### 3.5 Histopathological Studies




Shown renal tissue with preserved architecture, No abnormality seen, the glomeruli and tubules appear normal, No inflammatory cells and no cellular atypia seen.

**(A) Figure: 1.0: Photomicrograph of kidney Section of normal rat. MG X100, H&E stain**




Shown renal parenchyma with preserved architecture, No cellular atypia , both glomeruli and tubules appear normal .

**Figure: 2.0: Photomicrograph of kidney section of rat treated with 250 mg/kg of *E. lateriflora*. MG X100, H&E stain**




Glomerular basement thickening. No cellular atypia, dense eosinophilic substances distort through renal parenchyma

**Figure: 3.0: Photomicrograph of kidney section of rat treated with 500 mg/kg of *E. lateriflora*. MG X100, H&E stain**




Showed liver parenchyma with preserved architecture

**Figure: 4.0: Photomicrograph of liver section of normal rat. MG X100, H&E stain**



Shown liver with parenchyma, central veins (CV) and partial vasculature appears normal

**Figure: 5.0: Photomicrograph of liver section of rat treated with 250 mg/kg of *E. lateriflora*. MG X100, H&E stain**



Showed liver with parenchyma, there is expansion of partial spaces by fibrous spaces. Central veins and partial vasculature appears normal. No cellular atypia

**Figure: 6.0: Photomicrograph of liver section of rat treated with 500 mg/kg of *E. lateriflora*.  
MG X100, H&E stain**

#### **4.0 Discussion**

Determination of LD<sub>50</sub> is usually the initial step in the evaluation of the toxicity of a substance [7]. The LD<sub>50</sub> provides information on the dose of the test substance that is lethal to the test subject. In this study, 5000mg/kg/bwt of *E. lateriflora* had no lethal effect on the rats. Data from acute toxicity study may provide initial information on the likely toxic effect of a substance and help to arrive at a dose of a new compound (s) [6]. Liver, being the organ responsible for the metabolism of most xenobiotics is particularly susceptible to chemical injury [15] while the kidney functions in getting rid of the body waste materials that are either ingested or produced by the detoxification process of the liver. Hence, they are predisposed to damages upon accumulation of these toxic metabolites or chemicals [16]. In this study, we investigated the toxic effects of *E. lateriflora* on the liver and kidney of rat.

Biochemical indices monitored in the serum that includes electrolytes and other secretory substances of the liver and kidney can be used as markers for assessing the functional capacities of the organs [7]. Alteration in these parameters indicates impairment in the normal functions of the organs. The various biochemical parameters assessed in this study are useful indices for evaluating toxicity of plant extracts in animals [17-19].

The result of this study showed that the ethanol extracts of *E. lateriflora* have no hepatotoxic effects in rats. Increase in serum activity of AST, ALT, ALP and GGT is usually an indication of damage to the liver cells. These enzymes are mainly localized in the liver and are released into circulation upon damage to the hepatic cells. Thus, the changes in the activity and

concentration of these enzymes could reflect the extent of hepatotoxicity. Ethanolic whole plant extracts of *E. lateriflora* caused no alteration in some of these serum biomarkers, thus might not be toxic at the tested dose.

The liver synthesizes most serum protein [20]. The parenchymal cells are responsible for the synthesis of albumin, fibrinogen and other coagulation factors and most of globulin b and a [21]. The quantitative changes in some of these serum protein could have negative effects on the normal functioning of a cell and may also reflects impairment in the secretory functions of the liver [6].

The result of this study further affirmed the non-hepatotoxic nature of *E. lateriflora* as it produced no changes in the serum total protein and bilirubin level at 250mg/kg/bwt but increase total protein and albumin when administered at 500mg/kg/bwt. Nutritional status and disease state are known to affect the serum albumin level thereby affecting the blood osmotic pressure [7]. In this study, the increased observed in albumin level might be a reflection of increased appetite in the rats.

Bilirubin is derived from haemoglobin upon degradation of RBC. It is a metabolic breakdown product of heme and it is one of the most commonly used liver function test [22]. Abnormal increase in serum bilirubin is indicative of liver impairment. The observation in this study suggested that *E. lateriflora* had no toxicological effects at 250mg/kg/bwt but might be slightly toxic at 500mg/kg/bwt.

Evaluation of haematological parameters can be used to determine the extent of deleterious effects of *E. lateriflora* extracts on the blood, it can also be used to explain blood related functions of a plant extract or its product [18]. Changes in red blood cell, mean corpuscular volume, mean corpuscular Hb and platelets could be used to assess erythropoiesis, morphology

and osmotic fragility of the RBC [23]. The extracts of *E. lateriflora* caused no significant changes in the haematological parameters thus might be non-toxic at the tested dose.

Furthermore, renal function indices are usually needed to assess the normal functioning of the different parts of the nephron [24]. In addition, serum electrolyte, uric acid, urea and creatinine concentration could give an insight into the effects of plant extract on the tubular and glomerular functions of the kidney [7]. The result in table showed that the administration of *E. laterifolia* is non-toxic to the kidney. Glomerular filtration process removes creatinine from plasma and then excretes it in the urine without reabsorption by the tubules to any significant extent. In addition, when the plasma level increases above normal, creatinine could also be excreted through the tubules. The implication is that the serum creatinine level in renal disease generally does not increase until the renal function is substantially impaired [25]. However, high serum urea and creatinine level may be an indication of renal failure [15]. Normal renal function depends on a normal filtration rate. In this study, *E. laterifolia* had no toxic effect on the renal function at both doses. The result of these findings is consistent with earlier findings of Usman *et al.* [9] on *E. laterifolia*.

The microscopic examination of the kidney and liver of rats treated with 250mg/kg/bwt and 500mg/kg/bwt of *E. lateriflora* supported the results of the biochemical analysis. In the liver, the architecture of the parenchyma was preserved; the central veins and Partial vasculature appear normal, no cellular atypia, no inflammatory exudates. The entire tissue appears normal and no cellular atypia. Generally, any damage to the parenchyma liver cells result in the elevation of both transaminases in the blood [26]. Hence, the insignificant changes in serum ALT and AST activity observed in *E. lateriflora* treated group.



The renal histopathology study also revealed preserved kidney architecture. Both the glomeruli and tubules appear normal, no inflammatory cells and no cellular atypia in the control and *E. lateriflora* treated group. Rise in urea and creatinine level is only observed when a marked damage to functional nephron occur [27]. Therefore, the observation in this study might be attributed to the preservation of the renal and liver integrity.

## Conclusion

This study showed that the extract of *E. lateriflora* is relatively safe for consumption when consumed at very low dose. Therefore, medicinal plant could be maximized for providing relief to various disease; however, the public must be informed on the need to exercise caution while using medicinal plants to meet their health care need.

## Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the authors.

## References

1. Obaseki OE, Adesegun OI, Anyasor, GN, Abebawo OO. "Evaluation Of The Anti-Inflammatory Properties of the Hexane Extract Of *Hydrocotyle Bonariensis* Comm. Ex Lam. Leaves". *African Journal of Biotechnology*. 2016; 15(49): 2759-2771
2. Iwalewa EO, Mc-Gaw LJ, Naidoo V, Eloff JN. "Inflammation: The Foundation of Diseases and Disorders. A Review Of Phytochemicals Of South African Origin Used to Treat Pain and Inflammatory Conditions". *African Journal of Biotechnology*. 2007; 6: 2868-2885.
3. Hodzic Z, Pasalic H, Memisevic A, Srabovic M, Saletovic M, Poljakovic M. "The Influence Of Total Phenols Content On Antioxidant Capacity In The Whole Grain Extract. *European Journal of Science Research*. 2009;28 (3):471–7.
4. Sangita C, Priyanka C, Protapaditya, D, Sanjib B. "Evaluation Of In Vitro Anti-Inflammatory activity Of Coffee Against The Denaturation Of Protein". *Asian Pacific Journal of Tropical Biomed*. 2002; 2(1):S178-S180.



5. Ertekin V, Selimoglu MA, Altinkaynak SA. "Combination of Unusual Presentations Of Datura Stramonium Intoxication In a Child: Rhabdomyolysis and Fulminant Hepatitis". *Journal of Emergency Med.* 2005; 28: 227-228.
6. Ping KY, Darah I, Chen Y, Sreeramanan S, Sasidharan S. "Acute And Subchronic Toxicity Study of *Euphorbia hirta* Methanol Extract in Rats". *Journal of Biomed Research International.* 2013; 1-14
7. Ukwuani AN, Abubaka G, Hassan SW, Agaie BM. "Toxicological Studies of Hydromethanolic Leaves Extract of *Grewia Crenata*". *International Journal of Pharmaceutical Sciences and Drug Research.* 2012; 4(4): 245-249.
8. Hambali MJ. African Traditional Medicine. A Case Study of Hausa Medicinal Plants and Therapy. Zaria: Gaskiya Cooperation Ltd. 1990. P. 78.
9. Usman MM, Sule MS, Gwarzo MY. Toxicological studies of aqueous root extract of *Euphorbia lateriflora* (Schum and Thonn) in rats. *Journal of Medicinal Plants Studies.* 2014; 2(2): 58-62
10. Obi RK, Iroagba II, Ojiako OA. Viracidal Potential of Some Edible Nigerian Vegetables. *Afric J Biotech* 2006; 5(19):1785–1788.
11. Mbaka GO, Adeyemi OO, Anunobi CC. "Anti-Hyperglycaemic Effects of Ethanol Leaf Extract of *Sphenocentrumjollyanum* in Normal and Alloxan-Induced Diabetic Rabbits". *Global Journal of Pharmacology.* 2008; (3):46-51.
12. Dixon WJ. Staircase Bioassay: The Up and Down Method. *Neurosci. Biobehav. Rev.* 1991; 15:47-50
13. Olorunnisola OS, Adetutu A, Popoola RB, Owoade AO, Adegbola P, Adesina BT. Nephroprotective effect of ethanolic leaf extract of *Thaumatococcus danielli* (benth.) in streptozotocin induced diabetic rats. *Functional Foods in Health and Disease.* 2017; 7(12): 923-935
14. Avwioro OG. Histochemistry and Tissue Pathology, Principles and Techniques. Claverian um Press, Nigeria. 2010.
15. Hodgson E, Mailman RB, Chambers JE. Macmillan Dictionary of Toxicology. London: *The Macmillan Press*; 31, 62, 89, 100, 164, 186–218 and 322. 1988
16. Arthur CG, John EH. Textbook Of Medical Physiology. Edition 10, Philadelphia: W.B. Saunders. 2000; 279–281.
17. Yakubu MT, Bilbis LS, Lawal M, Akanji MA. "Evaluation Of Selected Parameters Of Rat Liver And Kidney Function Following Repeated Administration Of Yohimbine". *Journal of Biochemistry.* 2003; 15: 50-56.
18. Yakubu MT, Akanji MA, Anda T, Oladiji. "Hematological Evaluation In Male Albino Rats Following Chronic Administration Of Aqueous Extract Offadogia Agrestisstem," *In pharmacognosy Magazine.* 2007; 3:34–38.
19. Yakubu MT, Akanji MA, Oladiji AT. "Alterations In Serum Lipid Profile Of Male Rats By Oral Administration Of Aqueous Extract Of Fadogia Argrestis Stem". *Research Journal of Medicinal Plant.* 2008; 2: 66-73.
20. Rosalki SB, McIntyre N. Biochemical Investigations in the management of liver disease. Oxford Textbook of Clinical Hepatology, 2<sup>nd</sup> Edition. New York; Oxford University Press, 1999; 503-521
21. Thapa BR, Anuj W. "Liver Function Tests and their Interpretation". *Indian Journal of Pediatrics.* 2007; 74 (7): 663- 671.

22. Friedman SF, Martin P, Munoz JS. "Laboratory Evaluation of the Patient with Liver Disease". *A Textbook of Liver Disease. Saunders Publication, Philadelphia*, 2003; Pp. 661- 09.
23. Guyton AC, Hall JE. *Textbook of Medical Physiology. W.B Saunder, Philadelphia, PA, USA*. 2000
24. Abolaji AO, Adebayo AH, Odesanmi OS. Effect of Ethanolic Extract of *Parinari polyandra* (Rosaceae) on Serum Lipid Profile and Some Electrolytes in Pregnant Rabbits. *Research Journal of Medicinal Plant*. 2007; **1**: 121- 127.
25. Faulkner WR, King JW. Renal Function. In: W.T. Norbert (ed). *Fundamentals of Clinical Chemistry*. USA: WB Sounder Company. 1982; 975-978, 994 – 995.
26. Slichter SJ. "Relationship between Platelet Count and Bleeding Risk in Thrombocytopenic Patients Transfusion". *Medicine Reviews*. 2004; 18(3):153–167.
27. Lameire N, Van Biesen, Vanholder R. "Acute Renal Failure," *The Lancet*. 2005; 365(9457): 417–430.
28. Pari L, Murugavel P. Protective effect of  $\alpha$ -lipoic acid against chloroquine-induced hepatotoxicity in rats, *Journal of Applied Toxicology*. 2004; 24(1): 21–26.
29. Akinkurolere RO, Adedire CO, Odeyemi OO, Raji J, Owoeye JA. Bioefficacy of Extracts of some Indigenous Nigerian Plants on the developmental stages of mosquito (*Anopheles gambiae*). *Jordan Journal of Biological Sciences*. 2011; 4(4): 237-242