

Proximate Analysis and Phytochemical Screening of *Triclisia subcordata* Oliv Leaf

Okpara, F.N.^a; Nwaichi, E.O.*^a and Akaninwor, J.O.^a

^a Department of Biochemistry, University of Port Harcourt, Choba

Abstract

Proximate analysis and phytochemical screening of *Triclisia subcordata* Oliv leaf were investigated using standard analytical methods. Results of the proximate analysis showed contents for moisture (79.28 ± 2.32 %), Ash (1.91 ± 0.21 %), crude fibre (1.83 ± 0.13 %), lipid (1.81 ± 0.16 %), crude protein (8.32 ± 1.20 %) and carbohydrate (6.91 ± 0.74 %). Phytochemical screening showed the presence of all six metabolites undertaken but at different concentrations. Saponin, flavonoids were highly present, alkaloids were moderately while phenol, tannin and cyanogenic glycoside were slightly present. Tannins gave (0.01 ± 0.00 %), saponins (5.81 ± 0.23 %), alkaloids (2.32 ± 0.13 %), flavonoids (6.01 ± 0.36 %), cyanogenic glycosides (0.20 ± 0.00 mg/kg) and phenol (0.03 ± 0.00 %). The study therefore submits that *Triclisia subcordata* Oliv leaf has high contents of moisture, flavonoids and saponins, moderate alkaloids content and low contents of lipid, cyanogenic glycosides, tannins and phenols. This study supports the medicinal use of the plant, and unveils its potential as a source of micronutrients.

Keywords: *Triclisia subcordata* oliv; Proximate analysis; Phytochemical screening; Analytical methods; Traditional medicine.

1.0 Introduction

Phytochemicals are secondary metabolites produced by plants. These products are biologically active, naturally occurring chemicals in various parts of a plant, providing health benefits for humans further than those attributed to macronutrients and micronutrients. These compounds have been linked to human health by contributing to protection against degenerative diseases [1].

Triclisia subcordata Oliv (Menispermaceae) is a medicinal plant traditionally used for the treatment of various diseases in West Africa [2]. *T. subcordata* Oliv is commonly called “Alugboran” in Yoruba language, “ogwu-aju” in Nsukka (which means antidizziness), “Ezize” in Ohafia (which means benign skin growth), “ike mbekwu” in Umuoji in Eastern part of Nigeria (which is gotten from the shape of the leaf) [3]. The Menispermaceae is a temperate to tropical family of around 70 genera (including *Triclisia*) and 450 species of dicotyledenous small tropical flowering vines that are short in pedunculate clusters with twining stems, an appressed pubescent fruits and a few herbs, shrubs and trees. Leaves are reticulate, alternate and simple, but may be palmately veined and often lobed [2]. The stem is used as a rough fibre (tietie) [4].

It is a nature plant of west tropical Africa including Nigeria, Ghana, Ivory Coast, Sierra Leone, Senegal and Togo [5]. The plant frequently serves as rope used for tying purposes; the importance of the species however rest with their medicinal application including the use of root extract for the treatment of snake bite, ulcer, diarrhea, malaria pyorrhea, swelling of extremities, anemia joint pains, cancer, rheumatic pains and hypertension in Nigeria [3, 4].

Evidenced-based studies revealed that *Triclisia subcordata* has antiulcer, antihistamine, antimicrobial, anticancer, antioxidant and antidiabetic activities [6, 7, 8, 9]. The cytotoxicity Effects and Apoptosis Induction in ovarian and cancer cells by Bisbenzylisoquinoline and Isochondodendrine and 2'-norcocsuline alkaloids from *Triclisia subcordata* was assayed by Uche *et al.* [10] [11].



Plate 1: photograph showing *Triclisia subcordata* Oliv leaf

Medicinally, several hypotheses suggest that a decoction of the root is taken to treat fever and malaria. Also, root decoction is also taken as an emmenagogue and abortifacient (drugs or chemicals that induce abortion). The root pulp is rubbed in scarifications as a treatment for rheumatism, arthritis, anaemia and sleeping sickness. The leaf or a root juice is mixed with salt in palm wine and taken as treatment for coughs and bronchial disorders and as well function as sedative on the heart and to help wash palpitations. The leaf sap has a soothing effect on coughs. A decoction of the leaves and twigs is drunk, or leaf pulp is rubbed in, to treat oedema of the

legs. Furthermore, it was hypothesized that a decoction of the leaves and stem are used as a nasal or ocular instillation and as a purgative or bathe against epilepsy and as treatment for stomach ache respectively. The stem bark is powdered and applied topically to syphilitic sores and leprosy while the bark pulp is used as a purgative. A methanolic leaf extract has shown significant anti-ulcer effects [12]. Due to its acclaimed pharmacological importance by African natives, this paper is aimed at analyzing the proximate content and phytochemical screening of *Triclisia subcordata* Oliv leaf.

2.0 Materials and Methods

2.1. Sample Collection

Fresh *Triclisia subcordata* Oliv leaf was obtained from University of Port Harcourt campus environment, Choba, Rivers State, Nigeria. The plant was identified by a taxonomist (Dr. Nwosu) of the department of plant science and biotechnology University of Port Harcourt. The leaves were pulverized using a mechanical grinder to obtain a smooth mixture.

2.2. Experimental Design

After grinding of the leaf, approximately 50 g was weighed using electronic weighing balance and used for proximate and phytochemical screening using standard analytical methods.

2.3. Proximate Analysis

Determination of Moisture content [13]

A dry clean petridish was placed in an oven at 80°C for about 30 minutes, cooled in a desiccator and weighed as (w). 5 g of the samples was added to the petridish and weighed as (b). The petridish and its content were placed in an oven adjusted to 70°C. After 5 hours, the petridish containing the sample was removed and quickly transferred to a desiccator for cooling. The petridish was put back into the oven and adjusted to 105°C for another 5 hours after which it was

removed, put in desiccators for cooling. This process was repeated and weighed until a constant weight (c) was obtained.

The % moisture content was determined as follows;

$$\% \text{ moisture content} = \frac{b-c}{b-w} \times 100$$

Where W = weight of moisture can, b = weight of petridish + sample, c= weight of petridish + sample after drying.

Determination of ash content [13]

An empty crucible was first ignited in a muffle furnace for 1 minute and allowed to cool in desiccators containing silica gel. 5 g of the sample was accurately weighed into the preheated dish. The weight of the porcelain dish and the samples were noted. Afterwards, the dish was heated with a Bunsen burner in a fume cupboard until smoking ceases and later transferred into a muffle furnace at 550-570°C for about 18-24 hours to burn off all organic matter. After ashing, the crucible was removed from the furnace and placed in desiccator to cool at room temperature and weighed. The percentage ash content of the sample was calculated thus;

$$\begin{aligned} \% \text{ Ash} &= \frac{\text{weight of ash}}{\text{weight of sample}} \times 100 \\ &= \frac{W_3 - W_1}{W_2 - W_1} \times 100 \end{aligned}$$

Where; W_1 = weight of empty crucible, W_2 = weight of crucible + sample before ashing

W_3 = weight of crucible + sample after ashing

Determination of lipid [13]

A 5 g of the sample was weighed into a thimble and was extracted with petroleum ether until it siphons using the Soxhlet extraction method. The lipid was exhaustively extracted using petroleum ether at 40 – 60°C for 6 hours. The sample in the thimble was removed and dried in

air at 50°C for 5 minutes, cooled in a desiccator and weighed. The % lipid content was calculated as follows;

$$\begin{aligned}\% \text{ Lipid} &= \frac{\text{weight of sample (extracted fat)}}{\text{weight of sample}} \times 100 \\ &= \frac{W_2 - W_1}{W} \times 100\end{aligned}$$

Where; W_1 = weight of empty thimble, W_2 = weight of thimble + sample W = weight of sample

Determination of crude fiber [13]

A 2 g of the defatted sample was weighed into conical flask and 200 mLs of 1.25 % of boiling sulphuric acid was added within a minute. The content of the flask was filtered through a buchner funnel prepared with wet 12.5 cm filter paper. The sample was washed back into the original flask with 200 mLs of 1.25 % NaOH, and boiled for 30 minutes. All insoluble matter was transferred to the crucible and treated till the sample was free from acid. The sample was again washed in a muffle furnace at 550°C/hour. The crucible was then cooled in desiccator and reweighed.

$$\% \text{ Crude fiber} = \frac{W_2 - W_1}{W} \times 100$$

Where; W = weight of sample, W_1 = weight of crucible + sample

W_2 = weight of crucible+ filter paper after ashing.

Determination of Crude protein [13]

A 1 g of the sample was weighed and transferred into Kjeldahl flask. Few chips of antibumping granules, 4 g of digestion catalyst and 20 mLs of concentrated sulphuric acid were added at a 40°C angle with a retort stand on an electro thermal heater. The flask was gently heated for frothing to occur and subside, and then heat was increased to about 250°C. The digestion was carried out within 2-6 hours by which time the entire sample was digested completely. The digest

was cooled to room temperature and diluted to 100 mLs with distilled water. For distillation, 20 mLs aliquot of the digest was transferred into a round bottomed flask. This flask was connected to a Liebig condenser through a monoarm steel head (Adaptor). The liebig condenser was connected to a receiver flask through a receiver adapter. 10 mLs of 2 % boric acid and two drops of double indicator were pipetted into the distillation flask. 30 mLs of 40 % sodium hydroxide was injected into the distillation flask through a cork with the aid of a syringe. The flask was heated for 10 minutes to digest the content. The distillate was collected in the boric acid and then titrated with 0.1M HCL. The volume of HCl added was recorded as the titre value.

The % Crude protein was calculated thus;

$$\% \text{ Crude protein} = \% \text{ Nitrogen} \times 6.25$$

$$\% \text{ Nitrogen} = \frac{\text{titre value} \times 1.4 \times 100 \times 10}{1000 \times \text{wt of sample} \times \text{aliquot digest}}$$

Where, 1.4 = N₂ equivalent to 0.1NHCl used in titration

100 = Total volume of digest

Determination of total carbohydrate [13]

The total carbohydrate content of the sample was estimated as the Nitrogen free extract (NFE).

The arithmetic different methods involve adding the total percentage value of crude volume.

$$\text{Total CHO} = 100 - (\% \text{ fibre} + \% \text{ protein} + \% \text{ Moisture} + \% \text{ ash} + \% \text{ fats})$$

2.4. Phytochemical Screening

Preparation of the leaf extracts

Ten grams (10 g) of the leaf sample was soaked in 100 mL of water in a beaker and left for about 8 hours. The solution obtained was filtered using filter paper, and the filtrate used for phytochemical screening.

Qualitative determination of phytochemicals

Screening for presence of secondary metabolites (qualitative) were performed following standard methods by AOAC (1990) [13].

Quantitative determination of phytochemicals

Determination of Saponins [13]

A 10 g of the ground sample was measured into a conical flash and 100 mL of 20 % ethanol added to it. The suspension was heated over hot bath at 55°C for 12 hours with continuous stirring using a magnetic stirrer. The mixture was filtered and the residue re-extracted with another 200 mL of 2 % aqueous ethanol. The combined extract was reduced to 40 mL of the original size over a water bath at about 55°C. The purification process was repeated two more times. A 4 g of Sodium chloride (NaCl) was added to adjust the pH meter. The solution was shaken with 60 mL and 30 mL portions of n-butanol extract and later washed twice with 10 mL of aqueous NaCl. The remaining solution was evaporated to dryness in water bath. After evaporation, the sample was dried in the oven to a constant weight.

The saponin content was calculated in g/100g as;

$$\% \text{ saponins} = \frac{\text{weight of residue}}{\text{weight of sample}} \times 100$$

$$\% \text{ saponins} = \frac{W_2 - W_1}{W} \times 100$$

Where; W = weight of sample, W_1 = weight of evaporating dish

W_2 = weight of evaporating dish + saponin content after drying

Determination of Alkaloids [13]

A 2 g of freshly crushed sample was weighed and dispensed into 100 mL of 10 % acetic acid. The mixture was shaken and allowed to stand for 4 hours before filtration. The mixture was

filtered to remove all debris and evaporated to $\frac{1}{4}$ of the original volume. A 1 % concentrated ammonium hydroxide (NH_4OH concentration) was added drop wise to precipitate the alkaloids. It was filtered with well weighed precipitate in the filter paper, oven dried at 60°C for 30 minutes and then reweighed.

The alkaloid content was calculated as;

$$\% \text{ Alkaloid} = \frac{\text{weight of residue}}{\text{weight of sample}} \times 100$$

$$\% \text{ Alkaloid} = \frac{W_2 - W_1}{W} \times 100$$

Where; W = weight of sample, W_1 = weight of empty filter paper

W_2 = weight of filter paper of precipitate

Determination of Flavonoids [14]

The method used for determination of flavonoids was that of Bohn and Kocipai (1994). 5 g of the blended sample was extracted repeatedly and separated with 50 mL of 40 % aqueous methanol at room temperature. The solution was shaken for homogeneity and left to stand for about 4 hours and later filtered into a weighed beaker. The filtrate was later transferred into a crucible and evaporated to dryness over a water bath, then dried in an electric oven to a constant weight.

The flavonoid content was expressed in percentage as follows;

$$\% \text{ flavonoid} = \frac{\text{weight of residue}}{\text{weigh tof sample}} \times 100$$

Determination of cyanogenic glycoside [13]

One gram (1 g) of sample was weighed into a 250 mL round bottomed flask. 200 mL distilled water was added and allowed to stand for 2 hours (for autolysis to occur). Full distillation was then carried out and 150-170 mLs of distillate was collected in a 250 mL conical flask containing 2 mLs of 2.5 % NaOH . Anti-foaming agent (tannic acid) was added before distillation. To 100

mL of the distillate containing cyanogenic glycoside, 8 mL of 6N NH₄OH and 2 mLs of 5 % potassium iodide (KI) was added, mixed and titrated with 0.02 M silver nitrate (AgNO₃) using a micro burette against a black background. Permanent turbidity indicated end point.

Cyanogenic glycoside in the sample was calculated thus;

$$\text{cyanogenic glycosides (mg/kg)} = \frac{\text{Titre value (mL)} \times 1.0 \text{ g} \times \text{extract volume (mL)} \times 100}{\text{Aliquot volume (mL)} \times \text{weight of sample (g)}}$$

Determination of tannin [13]

A 0.5 g of the sample was weighed into 100 mL plastic bottle. 50 mLs of distilled water was added and shaken for 1 hour in a mechanical shaker. Then 5 mLs of the filtrate was pipette out into a tube and mixed with 3 mLs of 0.1 M Iron (III) chloride (FeCl₃) in 0.1N hydrochloric acid (HCl) and 0.008 M potassium ferrocyanide K₄[Fe(CN)₆]. The absorbance was measured in spectrophotometer at 720 nm wavelength within 10 minutes. A blank sample was prepared, and the colour also developed and read at same wavelength. A standard was prepared using tannic acid to get 100 ppm and measured using the formula below.

$$\text{Tannin} = \frac{A_n}{A_s} \times C \times \frac{100}{W} \times \frac{v_f}{v_g}$$

Where; A_n = Absorbance of test sample, A_s = Absorbance of standard solution, C = Concentration of standard solution, W = Weight of sample used, V_f = Total volume of extract, V_g = volume of extract analyzed.

Determination of phenol [13]

A 0.1 g of sample was weighed into a 250 mL round bottomed flask. 100 ml distilled water was added, boiled for 30 minutes and the volume made up to 100 mL. A 2.5 mL aliquot is measured into a conical flask and 5 mL of 0.1M NaOH added into it. The solution was heated to 50°C and allowed to cool, then 2.5 mL of 0.005 Iodine added. The flask was cork with a foil and allowed

to stand at room temperature for 2 hours. 0.5 ml of concentrated HCl was added and titrated to pale yellow with 0.1 M Na₂S₂O₃ with the addition of 3 drops of starch indicator. The solution is titrated again with 0.1 M Na₂S₂O₃. A blank is formed using 2.5 mL distilled water and absorbances read off.

2.5. Method of Data Analysis

All data collected were subjected to Students t-test for statistical analysis. All data were represented in mean \pm standard deviation (M \pm S.D). Level of significance was determined at a confidence level of determination ($p \leq 0.05$).

3.0 Results

Table 1: Proximate analysis of *Triclisia subcordata* Oliv leaf

Proximate content	<i>Triclisia subcordata</i> (%)
Moisture	79.28 \pm 2.32
Ash	1.91 \pm 0.21
Lipid	1.81 \pm 0.16
Crude protein	8.32 \pm 1.20
Crude fibre	1.83 \pm 0.13
Carbohydrate	6.91 \pm 0.74

Values are Mean \pm SD of triplicate determinations. n=3.

Table 2: Qualitative phytochemical screening of *Triclisia subcordata* Oliv leaf

Phytochemical content	<i>Triclisia subcordata</i> Oliv leaf
Tannins (%)	+
Saponins (%)	+++
Alkaloids (%)	++
Flavonoids (%)	+++
HCN (mg/kg)	+
Phenol (%)	+

Key: + = slightly present; ++ = moderately present; +++ = highly present

Table 3: Phytochemical screening of *Triclisia subcordata* Oliv leaf

Phytochemical content	<i>Triclisia subcordata</i> Oliv leaf
Tannins (%)	0.01±0.00
Saponins (%)	5.81±0.23
Alkaloids (%)	2.32±0.13
Flavonoids (%)	6.01±0.36
HCN (mg/kg)	0.20±0.00
Phenol (%)	0.03±0.00

Values are Mean ± SD of triplicate determinations. n=3.

4.0 Discussion

The proximate content of *Triclisia subcordata* oliv leaf showed appreciable amount of the parameters as expressed in Table 1. From this study, the moisture content was very high (79.28 %) when compared with findings of [15] on fluted pumpkin pod which imply that this leaf may not be stored for long and can be prone to microbial attack.

Ash contents express the mineral index contents in biological mass. This study revealed relatively lower ash in *Triclisia subcordata* oliv leaf (1.91 %) than in fluted pumpkin leaf (9.68 %), bitter leaf, (15.86 %) and *Moringa oleifera* (15.09 %) leaves [16]. This implies that *Triclisia subcordata* oliv leaves are not good source of mineral elements (Table 1).

Lipid in food absorbs and retains the flavour thereby enhancing the palatability [17]. The lipid content recorded in this study (1.81 %) is far lower than values obtained by [15] which implies consumption of *T. subcordata* will not enhance pallatability.

Proteins are necessary for the build-up of tissues and substances like hormones and enzymes. The values of protein obtained in this study (8.32 %) are low when compared with [15], 24 % in *Amaranthus vividis* [18], 20.72 % in *Moringa oleifera* [19], 21.0 % in *Lasianthera africana* and 15.0 % in *Heinsia crinata* [16, 20] which makes it insufficient source of protein.

The crude fibre contents of *Triclisia subcordata* oliv leaf (1.83 %) is very low when compared with those of *Lasianthera africana* (15.3 – 18.1 % dry mass) [21] and *Heinsia crinata* (13 – 15 % dry mass) [20]. Although intake of dietary fibre can lower the serum cholesterol level, risk of coronary heart disease, hypertension, diabetes, obesity, constipation and cardiovascular diseases, colon and breast cancer [22, 23]. The major problem associated with nutrition of vegetables by human is the high fibre content which can cause intestinal irritation and lower nutrient

bioavailability [16, 24]. Intake of fibre can stimulate peristaltic movement, weakening of hunger, increase stool bulk and reduce serum levels of cholesterol [25, 26].

The carbohydrate content recorded was 6.91 %. Carbohydrates are pivotal nutrients needed for a balanced diet [27]. Generally, the low contents of protein, lipid and carbohydrate in this leaf imply that it is not a good source of energy.

Table 2 revealed that saponin, flavonoids were highly present, alkaloids were moderately present while cyanogenic glycosides, tannins and phenols were slightly present in *Triclisia subcordata* oliv leaf.

Table 3 revealed the phytochemical screening of *Triclisia subcordata* oliv leaf. The study showed substantial amounts of flavonoids, saponins and alkaloids and small amounts of phenol, tannins and cyanogenic glycosides. When compared with the study of [28], tannins, saponins, alkaloids and phenols were lower while flavonoid content was high.

Tannins are complex moiety with wide pharmacological activities and are produced by majority of plants as protective substance. Tannin has astringent property, hastens the healing of wounds and inflamed mucous membrane and has been used since past as tanning agents. Tannin has received considerable attention in the fields of nutrition, health and medicine, largely due to their physiological activity, such as antioxidant, antimicrobial and anti-inflammatory properties [29, 30].

Saponins protect against hyperglycaemia, hypercholesterolaemia, hypertension [31], have antibiotic properties and antiinflammatory property and aid healing [32].

Alkaloids have been reported to be powerful pain relievers, exert an anti-pyretic, antihypertensive, antifungal, anti-inflammatory, anti-fibrogenic effect [33], stimulating, anaesthetic action [34] and inhibiting activity against most bacteria [35].

The flavonoids show antioxidant activity and have strong anti-cancer activity [36]. Flavonoids exhibit their antioxidative properties through several mechanisms, such as scavenging of free radicals, chelation of metal ions, such as iron and copper, inhibition of hydrolytic and oxidative enzymes inhibit antibiotics resistant microbes and also act as anti-inflammatory agent [37].

Conclusion

It can be concluded that *Triclisia subcordata* oliv leaf has high moisture (which show their low shelf-life), alkaloid, flavonoids and saponins (pharmacological relevance) with low lipid, cyanogenic glycoside, tannins and phenols. The low cyanogenic glycoside, tannin and phenol implies that the plant is non-poisonous and inhibits inflammatory enzymes. Generally, the proximate analysis does not support their nutritional relevance as macronutrient supplement (as they are seen in smaller quantity when compared to other plants (*Telfairia occidentalis* etc)) but provide the micronutrients required for proper body functioning. The low concentration of lipid obtained makes it suitable and healthy for people on low fat diet. However, the phytochemical screening backs their usage in traditional medicine research and will be helpful in the coping of different diseases.

References

- [1] Dandjesso, C., Klotee, J.R., Dougnon, T.V., Segbo, J.M., Gbaguidi, F., Fah, L., Fanou, B., Loko, F. & Dramane, K. (2012). Phytochemistry and hemostatic properties of some medicinal plant sold as anti-hemorrhagic in Cotonou markets (Benin). *Indian Journal of Science and Technology*, **5**(8):3105-3109.
- [2] Ezimah, A.C., Ajugwo, A.O., Osuala, F.N. & Aghatise, K. (2013). Comparative Studies of EDTA, Sodium Citrate and Aqueous Extract of *Triclisia Dictyophylla*. *Scholars Academic Journal of Biosciences*, **1**(2), 55-58.
- [3] Dalziel, J.M. (1937). The Useful Plants of West Tropical Africa, The Crown Agents for the Colonies, London. PP. xii + 612pp
- [4] Irvine, F.R. (1961). Woody plants of Ghana, Oxford University Press, London. Pp. xcv + 868
- [5] Trease, G. & Evans S.M. (1993). Medicinal Plants and Traditional Medicines in Africa, 2nd Ed., Spectrum Books. Pp. 35 – 53.
- [6] Asuzu, I.U. & Anaga, A.O. (1995). The anti-ulcer effect of methanolic extract of *Triclisia subcordata* O. leaves in rats. *Journal of Herbs, Spices and Medicinal Plants*, **3**(3), 45-53.
- [7] Abo, K.A., Lawal, I.O. & Ogunkanmi, A. (2011). Evaluation of extracts of *Triclisia subcordata* Oliv and *Heinsia crinita* (Afz) G. Taylor for antimicrobial activity against some clinical bacterial isolates and fungi. *African Journal of Pharmacy and Pharmacology*, **5**(2), 125-131.
- [8] Ayoola, M.D., Akinwunmi, K.F. & Agboola, O.B. (2017). Antidiabetic and antioxidant activities of *Entandrophragma cylindricum* and *Triclisia subcordata*. *Nigerian Journal of Natural Products and Medicine*, **21**(1), 24-31.
- [9] Li, W. (2016). Cytotoxicity effects and apoptosis induction by cycleanine and tetrandrine. *Planta Medica: Natural Products and Medicinal Plant Research*, **30**(9), 1533-1539.
- [10] Uche, F.I., Drijfhout, F.P., McCullagh, J., Richardson, A. & Li, W.W. (2016). Cytotoxicity effects and apoptosis induction by bisbenzylisoquinoline alkaloids from *Triclisia subcordata*. *Phytotherapy Research*, **30**(9), 1533-1539.
- [11] Uche, F.I., Abed, M.N., Abdullah, M.I., Drijfhout, F.P., McCullagh, J., Claridge, T.W., Richardson, A. & Li, W. (2017). Isochondodendrine and 2'-norcocsuline: additional

- alkaloids from *Triclisia subcordata* induce cytotoxicity and apoptosis in ovarian cancer cell lines. *Royal Society of Chemistry Advances*, **7**(70), 44154-44161.
- [12] Tropical Plants Database, Ken Fern. tropical.theferns.info.
/tropical.theferns.info/viewtropical.php?id=Triclisia+subcordata/ Retrieved 2021-05-01.
(<http://www.prota.org>)
- [13] AOAC. (1990). Association of Official Analytical Chemists. Method of analysis
Washington DC, U.S., 15, 1250-1255.
- [14] Bohn, C. & Kocipai, G. (1994). Method of analysis. Washington DC, U.S., 8, 1233-1246.
- [15] Akwukwaegbu, P.I., Peters, D.E., & Wegwu, M.O. (2016). Proximate analysis and phytochemical screening of fluted pumpkin (*Telfairia occidentalis*) pod. *American Journal of Food, Nutrition and Health*, **1**(1), 1-6.
- [16] Effiong, G. S., Ogban, P. I., Ibia, T. O., & Adam, A. A. (2009). Evaluation of nutrient supplying potentials of Fluted Pumpkin (*Telfairia occidentalis*, Hook, F.) and Okra (*Abelmoschus esculentus*) (L.) Moench. *Academic Journal of Plant Sciences*, **2**(3), 209–214.
- [17] Fagbohun, E.D., Egbebi, A.O. & Lawal, O.U. (2012). Phytochemical screening, proximate analysis and in-vitro antimicrobial activities of methanolic extract of *Cnidioscolus aconitifolius* leaves. *International Journal of Pharmaceutical Science Review and Research*, **13**, 28-33.
- [18] Sena, L. P., vander Jagt, D. J., Rivera, C., Tsin, A. T. C., Muhammadu, I., Muhamadou, O., Milson, M., Pastosyn, A. & Glew, R. H. (1998). Analysis of Nutritional components of eight famine foods of the republic of Niger. *Plant Foods for Human Nutrition*, **52**, 17-30.
- [19] Lockeett, C. T., Calvert, C. C. & Grivetti, L. E. (2000). Energy and micronutrient composition of dietary and medicinal wild plants consumed during drought. Study of rural Fulani, Northeastern Nigeria. *International Journal of Food Science and Nutrition*, **51**, 195- 208.
- [20] Udosen, E. O., Udok, U. E. & Unuigue, O. S. (1998). The comparison of the Nutrient composition of *Lasianthera africana* and *Heinsia crinata*. *Journal of Food Biochemistry*, **23**, 571-576.

- [21] Isong, E. U. & Idiong, U. I. (1997). Comparative studies on the nutritional and toxic composition of three varieties of *Lesianthera africana*. *Plant Foods for Human Nutrition*, **51**:79-84.
- [22] Aberoumand, A. & Deokule, S.S. (2009). Studies on nutritional values of some wild edible plants from Iran and India. *Pakistan Journal of Nutrition*, **8**, 26-31.
- [23] Chukwu, E.C., Osuocha, K.U. & Uhegbu, F.O. (2018). Nutrient Composition and Selected Biochemical Effects of *Cnidoscolus aconitifolius* Leaf Extracts in Male Albino Rats. *Journal of Forensic Research*, **9**(1), 409.
- [24] Iwuji, S.C., Nwafor, A., Egwurugwu, J., Ejeta, K. & Akpan, U. (2013). Comparative characterization of the phytomedicinal constituents of *Cnidoscolus aconitifolius* leaf extracts. *American Journal of PharmTech Research*, **3**, 780-784.
- [25] Umar, K.J., Hassan, L.G., Dangoggo, S.M. & Ladan, M.J. (2007). Nutritional composition of water spinach (*Ipomea aquatica* Forsk) leaves. *Journal of Applied Science*, **7**, 803-809.
- [26] Ramula, P. & Rao, P.U. (2003). Dietary fibre content of fruits and leafy vegetables. *Nutrition News*, **24**, (3) 1-6.
- [27] Emebu, P.K. & Anyika, J.U. (2011). Proximate and mineral composition of Kale (*Brassica oleracea*) grown in Delta State, Nigeria. *Pakistan Journal of Nutrition*, **10**, 190-194.
- [28] Okpara, F.N. & Akwukwaegbu, P.I. (2020). Effect of aqueous leaf extract of *Cnidoscolus aconitifolius* on lipid profile and haematology of carbon tetrachloride treated rats. *Acta Chemica Iasi*, **28**(2), 237-256.doi: 10.2478/achi-2020-0017.
- [29] Asquith, T.N. & Butter, L.G. (1986). Interaction of condensed tannin with selected proteins. *Phytochemistry*, **25**(7), 1591-1593.
- [30] Killedar, S.G. & More, H.N. (2010). Estimation of tannins in different parts of *Memecylon umbellatum* Burm. *Journal of Pharmacy Research*, **3**(3), 554-556.

- [31] Trease, G.E. & Evans, W.C. (1985). *Pharmacognosy*, 14th edition, London: W.B. Sanders company. Pp 23-29.
- [32] Krishnaiah, D., Devi, T., Bono, A. & Sarbatly, R. (2009). Studies on phytochemical constituents of six alaysian medical plants. *Journal of Medicinal Plants Research*, **3**(2), 67-72.
- [33] Awoyinka, O.A., Balogun, I.O. & Ogunnowo, A.A. (2007). Phytochemical screening and in-vitro bioactivity of *Cnidoscolus aconitifolius* (Euphorbiaceae). *Journal of Medicinal Plants Research*, **1**(3), 63-65.
- [34] Edeoga, H.O. & Enata, D.O. (2001). Alkaloids, tannins and saponin content of some medicinal plants. *Journal of Medicinal Aromatic Plant Science*, **23**, 344-349.
- [35] Al-Bayati, F.A. & Sulaiman, K.D. (2008). In-vitro antimicrobial activity of *Salvadora persica* L. extracts against some isolated oral pathogens in Iraq. *Turkish Journal of Biology*, **32**(1), 57-62.
- [36] Salah, N., Miller, N.J., Paganga, G., Tijburg, L., Bolwell, G.P., Rice, E. & Evans, C. (1995). Polyphenolic flavonols as scavenger of aqueous phase radicals and chain breaking antioxidants. *Archives of Biochemistry and Biophysics*, **322**(2), 339-346.
- [37] Kessler, R.C., Berglund, P., Demler, O., Jin, R. & Koretz, D. (2003). The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *Journal of American Medical Association*, **289**, 3095-3105.