

Lopped ster

Exfoliating Bark

Stem showing prickles



**Bipinnate** Leaves

Fruits

## Location :

Indopiptadenia is monotypic genus occurring in India and Nepal region. Only a few trees of Indopiptadenia oudhensis (Brandis) Brenan are surviving today on Indo-Nepal border in Balrampur district of Uttar Pradesh. This tree was first locally discovered in 1871 by Mr. Richard Thompson and was found to be heavily lopped for fodder. Currently the exact location is in Siriya nala in between Barahwa range and Tulsipur range.

Vernacular / Local Name : Genti, Gainti (India), Haathi Paula (Nepali).

## **Classification :**

Kingdom	:	Plantae	Division	:	Magnoliophyta
Class	:	Magnoliopsida	Order	:	Fabales
Family	:	Fabaceae (Leguminosae)	Subfamily	:	Mimosoideae

GOI notification S.O.998 (E) dated 15th April 2009 Under section 38 of the Biological Diversity Act 2002, the Government of India has declared this species to be on the verge of extinction and has prohibited and  $regulated \, the \, collection \, of \, this \, plant.$ 

## •Uttar Pradesh State Biodiversity Board•

East Wing, III Floor, A-Block, PICUP Bhawan, Vibhuti Khand, Gomti Nagar, Lucknow-226 010, U.P., INDIA Phone : +91 522 4006746, +91 2306491, E-mail: upstatebiodiversityboard@gmail.com Website: www.upsbdb.org

